

A manual of the dissection of the human body / by Luther Holden ; with notes and additions by Erskine Mason.

Contributors

Holden, Luther, 1815-1905.

Mason, Erskine, 1837-1882

Francis A. Countway Library of Medicine

Publication/Creation

1868

Persistent URL

<https://wellcomecollection.org/works/mdshtj5g>


License and attribution

This material has been provided by This material has been provided by the Francis A. Countway Library of Medicine, through the Medical Heritage Library. The original may be consulted at the Francis A. Countway Library of Medicine, Harvard Medical School. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>




A 3.A.1868.4

Harvard University
Library of
The Medical School
and
The School of Public Health




The Gift of





To
John P. S. Jackson M.D.

With the
Compliments of the
Editor ^{and} Publisher



Digitized by the Internet Archive
in 2011 with funding from
Open Knowledge Commons and Harvard Medical School

☞ The Only Book in the Market up to the Times. ☞

THE FOURTH EDITION NOW READY.

A PRACTICAL TREATISE

ON THE

Diseases of the Sexual Organs in Women

By F. W. VON SCANZONI,

Professor of Midwifery and Diseases of Females in the University of Wurzburg; Counsellor to his Majesty, the King of Bavaria; Chevalier of many Orders.

TRANSLATED FROM THE FRENCH OF DRS. H. DOR AND A. SOCIN.

And Annotated with the Approval of the Author.

BY AUGUSTUS K. GARDNER, A. M., M. D.,

Professor of Clinical Midwifery and the Diseases of Women, in the N. Y. Medical College, Author of "The Causes and Curative Treatment of Sterility," Editor of "Tyler Smith's Lectures on Obstetrics," etc.

WITH UPWARD OF SIXTY ILLUSTRATIONS.

It is almost unnecessary to speak of the merits of the above work; Scanzoni has a world-wide reputation, and on the Continent of Europe he is considered as standing at the head of Obstetrical Science, being Professor of Midwifery in the leading College in Germany, and from his great ability and well known experience, it makes this book the authority on this subject. The American Editor, who is himself noted among the principal Obstetricians of the metropolis of New-York, thus speaks of this book:

"Had I fully appreciated the amount of labor involved in the translation and editing of an American edition of this work, I should not have ventured on so great a responsibility. Having, however, once commenced, I found myself unconsciously sustained by the interest the volume excited in my mind, and the further my task progressed, the stronger grew the conviction that I should be doing a good deed in presenting to the profession of my country this treatise of the celebrated teacher of Wurzburg—a work which, on its subject, I think is unequalled in German, French or English Languages.

"In the etiology, pathology and therapeutics of female diseases, with all the improvements which have been realized during the last twenty years, this volume is exceedingly rich; while in its arrangement it is so methodical that it must constitute one of the text-books for students and one of the most reliable aids to the busy practitioner.

"The Paris translation, of which the present is an English version, was subjected to the revision of the author. The valuable notes which the French translators have added are retained with the initials of the writers, while those of myself are enclosed in brackets [], and have been incorporated in the text on account of their practical character.

In its publication no expense has been spared. In typography, paper and illustrations, as now issued, it contains nearly 800 pages, Royal Octavo, handsomely bound in cloth, with beveled boards, and retails at \$5 per copy.

☞ Copies of the above book sent, postage paid, on receipt of price. Address

ROBERT M. DE WITT, Publisher,

13 Frankfort St., N. Y.

Copies of this book was sent to every Medical Hospital by the United States Government for the use of the Surgeons.

SEVENTH AMERICAN EDITION.

Cellular Pathology,

AS BASED UPON

PHYSIOLOGICAL AND PATHOLOGICAL HISTOLOGY TWENTY LECTURES,

Delivered in the Pathological Institute of Berlin, during the months of February, March and April, 1858.

By RUDOLPH VIRCHOW,

Public Professor in Ordinary of Pathological Anatomy, General Pathology and Therapeutics in the University of Berlin; Director of the Pathological Institute, and Physician to the Charity Hospital, etc., etc.

Translated from the Second Edition of the original, by FRANK CHANCE, B. A., M. B., Licentiate of the Royal College of Physicians; Physician to the Blenheim Free Dispensary and Infirmary. With Notes and numerous Emendations, principally from MS. Notes of the Author, and Illustrated by 144 Engravings on Wood.

The Publisher would call the attention of the Trade to the above very valuable Medical Book now first reprinted from the English Translation of the Third German Edition.

The above Lectures were delivered in the year 1858, in the new Pathological Institute of the University of Berlin, in the presence of a large number of the first Medical Men in that city.

The importance of the subject, the new ideas advanced, and the well-known reputation of the Lecturer, compelled the publication of these Lectures in Book form, and it is now considered the Standard Work throughout Germany, on this subject. We would herewith submit the translator's Preface:—

"Professor Virchow and his works are so well known wherever the science of Medicine is studied, that I think it quite unnecessary to give any account of them here.

"When I arrived in Berlin, in March, 1858, these Lectures were in the course of delivery, and I was present at a few of the concluding ones. Subsequently, whilst attending the lectures, classes, and post-mortem examinations (from 700 to 800 bodies are examined annually in the Institute) which are held in the Pathological Institute, by Professor Virchow, I had ample opportunities for seeing practical illustrations of most of the doctrines advocated in this book. It was natural, therefore, that I should feel a desire to translate these lectures, the more especially as I had every reason to suppose that the views put forward in them still remained unknown—in consequence, no doubt, of their German dress—to a large proportion of the English Medical Public, although they had already, many of them several years previously, appeared in Professor Virchow's larger works.

"The translation will in many instances be found to differ somewhat from the original, for numerous additions, subtractions and substitutions have been made, many of them at the suggestions of the author, many at my own, but all with the author's sanction.

"A few Notes will be found, especially in the latter Lectures. Of these some are literal, some free translations of, or are based upon answers I received from Professor Virchow to questions I had put to him, while others were made entirely at his own suggestion, and are literal translations of his words. In all cases, however, the Notes have been submitted to the author, and approved by him.

"An Index too, I thought might be of service, and I have therefore added a tolerably full one.

"I cannot sufficiently thank Professor Virchow for the very great trouble—a trouble of which nobody but myself can have any idea—which he has taken in revising this translation, nor for the courtesy and kindness with which he has replied to the very numerous questions—many of them put for my own private information—which I have plagued him with. He has written me fully fifty letters, most of them very long ones; and when I reflect that he daily passes eight or nine hours at the Charity, that he reads all the more important German, French and English Medical Works which appear, and is besides constantly engaged in publishing something fresh, I can scarcely conceive how he has managed to find time to write these letters, of which a large proportion reached me by return of post.

In its publication no expense has been spared. In typography, paper or illustrations: as now issued, it contains nearly 800 pages, Royal Octavo, handsomely bound in cloth, with beveled boards, and retailed at \$5 per copy.

Copies of the above book will be sent postage paid, on receipt of price. Address,

ROB'T M. DE WITT, Publisher,

13 Frankfort Street, N. Y.

The Modern Practice of Midwifery.

THIRD AMERICAN EDITION NOW READY.

A Course of Lectures on Obstetrics

By WILLIAM TYLER SMITH, M. D.,

Physician, Accoucheur and Lecturer on Midwifery and Diseases of Females in St. Mary's Hospital Medical School; Member of the Royal College of Physicians; Vice-President of the Medical Society, London; Honorary Fellow of the Obstetrical Society of Dublin, &c.

WITH AN INTRODUCTORY LECTURE ON THE

HISTORY OF THE ART OF MIDWIFERY, AND COPIOUS PRACTICAL ANNOTATIONS.

BY AUGUSTUS K. GARDNER, A. M., M. D.,

Late Instructor on Obstetrics in the N. Y. Preparatory School of Medicine; Author of "The Causes and Curative Treatment of Sterility," etc.

ILLUSTRATED BY ENGRAVINGS.

The above Course of Lectures on Obstetrics, lately published in the columns of the London Lancet has been carefully revised by the author, and by the addition of two new chapters and frequent amplifications, and some score of new engravings, has been arranged to form one of Churchill's Celebrated Manuals for Students. The Publisher has the pleasure to announce, that in accordance with the general high estimate of their value and the frequently expressed desire that they might be made available to the profession, he made arrangements with the distinguished practical accoucheur and obstetric student, Augustus K. Gardner, A. M., M. D., Fellow of the New-York Academy of Medicine; N. Y. Pathological Society; Author of the "Causes and Curative Treatment of Sterility," &c., &c., to revise these Lectures, and properly arrange them for the use of the American Student, who to secure such a result, has interspersed copious annotations throughout the work, and added a Preliminary Lecture on the History of the Art of Midwifery, and the Os and Cervix Uteri in its Obstetric Relations. In its publication no expense has been spared in typography, paper or illustration. Numerous wood cuts have been added by the editor illustrative of practical points; thus rendering it the most profusely illustrated work on Obstetrics ever published, and entitling it in every respect to be considered as the Modern Practice of Midwifery. As now issued it occupies nearly 800 pages royal octavo, handsomely bound in cloth, bevelled boards.

EXTRACTS FROM LEADING AMERICAN MEDICAL JOURNALS.

From the SAVANNAH JOURNAL OF MEDICINE.

"It is especially adapted for polishing up rusty practitioners and keeping them well posted up in the discourses of their contemporaries."

From the MEDICAL AND SURGICAL REPORTER.

"The original lectures though previously quite soluble, have been much improved by the additions of Mr. Gardner. * * * In many points he has materially improved the original by additions, which though small, would have been much merited by the reader."

From the MAINE MEDICAL AND SURGICAL REPORTER.

"We must admit, that we have never before seen so comprehensive a text-book, on the art and science of Obstetrics. It is a matter for congratulation with the whole profession, that we have so valuable a work."

From the CINCINNATI LANCET AND OBSERVER.

"There is so much virtue in most of these Lectures that the work will commend itself largely to the profession."

From the PENINSULAR AND INDEPENDENT MED. JOUR.

"It is a very complete compendium of the knowledge extant."

From the BOSTON MEDICAL AND SURGICAL JOURNAL.

"Dr. Smith's style is pleasant, his English good, his mode of expression clear and elegant, and his work as a whole, very readable, as well as practically instructive."

From D. F. CONDIE, Am. Jour. Med. Sciences.

"The name, Dr. Smith, is a sufficient guarantee for the general accuracy of the theoretical teaching and practical directions of this manual. * * * The Editor has fulfilled his duties, on the whole, in an acceptable manner. The preliminary lecture on the "History of the Art of Midwifery" is, to say the least of it, interesting and instructive, while his practical annotations, together with his lecture upon the Os and Cervix Uteri demanded in cases of rigidity of the latter parts are, in the main, sound and judicious."

From Prof. C. T. ELLIOT, N. Y. Jour. of Medicine.

"We are of those who are only too happy to see English books reproduced in this country, when, as in this case, the additions are such as to justify the use of the annotators named on the title page. * * * We can confidently recommend alike the book and the annotations to the study of the student and the library of the professional man."

From A. T. WOODWARD, M. D., Prof. of Obstetrics, &c. in Castleton Med. College

"I deem it, all in all, the very best Text-book I have ever had the pleasure to peruse, and worthy of the author and editor who have contributed so much to the profession."

It will be sent by mail or express, free of cost, for \$5 per copy. Orders should be addressed to

ROBT M. DE WITT, Publisher,

13 Frankfort Street, N. Y.

STERILITY: ITS CAUSES AND CURATIVE TREATMENT.

With a Preliminary Statement of the Physiology of Generation.

COLORER LITHOGRAPHS AND NUMEROUS WOOD CUT ILLUSTRATIONS.

By AUGUSTUS K. GARDNER, A. M., M. D.,

Permanent Member of the American Medical Association, Fellow of the New York Academy of Medicine, &c., &c.

This is one of the ablest and most valuable Medical Works ever published on a subject of vital importance to all those who have heretofore supposed STERILITY to be incurable.

It is clearly and concisely written, and illustrated in a manner to make it easy of comprehension to readers of all classes.

We give a portion of the

CONTENTS.

I.—PHYSIOLOGY OF GENERATION.

In what reproduction consists—Fecundation in plants and fish—Stated times of impregnation in animals and women—That the woman who menstruates is capable of being fecundated—Identity of fecundation in all animal life—Theory of producing sexes at will—Proportion of male to female births—Description of the ovum—Male life principle—The female organs of reproduction—Their functions—Manner of menstruation—The human egg—Anatomy of egg—Man in his first estate—Semen—Its composition—Development of Spermatozoa—Manner of union of the male and female life giving principle—Action of uterus in congress—Bischoff's and others' views—Distinctive duties of the divisions of the germ vesicle—Growth of development ceases and the growth of augmentation commences.

II.—PATHOLOGY OF STERILITY.

Fecundation—How insured—Circumstances preventing—Affections of the vagina—impotence of male from malformations, disease, &c—Effects of various vaginal discharges—Various appearance of hymen—Constriction of, and strictures in the vagina—Prolapsus, malpositions, flexions and versions of uterus—Affections of os and cervix uteri—Polyphi of uterus—Value of the speculum in diagnosis—Diseases of the os and cervix, the analogue of various skin diseases—A scrofulous diathesis having a marked tendency to uterine disease, also to fecundation—Symptoms of uterine disease—Character of vaginal discharges—Cause of death of spermatozoa—Falling of the womb

Leucorrhœa an evidence of a strumous diathesis—Speculum os uteri; Effects of inordinate sexual intercourse; Diseases of cavity of the uterus; Uterus ossified; Metritis; Diseases of the fallopian tubes; Stricture; Rupture; Obliteration; Disease of fimbriated extremity; Polyphi; Disease of ovary; Tumors in and around; Hernia of, absence of, atrophy; Rheumatism and neuralgia of uterus and appendages; Influence of fatness as causing sterility; "Free martins;" the female of male and female twins, barren; Mental manifestations of the diseased; Absence of sexual appetites from over stimulation, venereal excesses, masturbation, &c.

III.—THERAPEUTICS OF STERILITY.

Treatment and cure sometimes synonymous; Requisites for fecundation; Operations on Hymen; Strictures of vagina treated; Sensitiveness of vagina from urethral tumors relieved by local anesthetics; Removal of urethral tumors by knife, ligature, cautery; Eversion of the uterus treated by Simpson's uterine supporter; Polyphi; Various modes of treatment; excision, forsyon, caustics curette; Little danger of hemorrhage from; Removal by Chaissegagnac's ecraseur; By galvanism; Hydatids; How removed; New form of speculum; Ulcerations treated by local injections of cold water; Infusions of tan-

nin; Caustic applications; alteratives; Sexual intercourse not necessarily interrupted. Issues upon os uteri, Leeches and their danger; Stricture of cervix, and fissures of os; Dysmenorrhœa; Instrument for division of strictures; Sponge Tents; Catarrh of the womb; Treatment by injections; Dangers of; Obliteration of the cavity of the uterus; Probing fallopian tube; Treatment for affections of the ovary; excessive fat; Torpor of uterine system; Application of electricity; Syphilitic taint; Sobriety of body and mind recommended.

Read the following from some of the most eminent members of the Profession:

Dr. JOHN W. FRANCIS, New York, writes to the author: "You have chosen a subject of great intricacy, but have manifested by your labors an earnestness in its investigation commensurate with its practical importance to the happiness and well-being of the social state. Your anatomical and pathological expositions must have been the result of close observation, cautious reflection, and professional devotion."

Dr. R. S. KISSAM, New York, writes: "Its simple and perspicuous style will commend it both to the professional and general reader, thus conveying to the afflicted the knowledge that sterility is not always an incurable condition."

Prof. J. HUMPHREYS STORER, Boston, writes: "I have read it with no little interest, and would recommend it to the Medical Student as a valuable compend, exhibiting great familiarity on the part of the author with his subject, which is rendered perfectly intelligible to all by the clearness and simplicity of manner with which it is treated."

"Many practical suggestions, particularly with regard to the treatment of the derangements spoken of, are distributed throughout the work, which cannot but be of importance to the members of the profession."

Prof. J. V. P. QUACKENBUSH, Albany, writes: "I am very much pleased with the work, and consider it a valuable addition to my library. In novelty, literary ability, typographical appearance, and beauty of illustration, it compares favorably with the best European works, and the profession demands and requires such publications."

Prof. THOS. G. PRIDLEAU, Charleston, S. C., says: "It should have a place in every library with Tyler Smith, Meigs, &c."

THE MEDICAL PRESS speaks of the book in the most flattering terms.

"It is just such a book as the Profession wants on this generally ignored subject." [*Philadelphia Medical and Surgical Journal*].

"Dr. Gardner has made the Diseases of Females an especial object of attention, and this book proves that he has profited by his opportunities. It is an excellent work, which we recommend to all practitioners, young and old." [*N. H. Journal of Medicine, Manchester*].

Price \$3; for which amount we will mail a copy, pre-paid, to any distance within 3,000 miles.

R. M. DE WITT, Publisher, 13 Frankfort St., N. Y.

R. T. Jackson.

A MANUAL
OF THE
DISSECTION OF THE HUMAN BODY.

By LUTHER HOLDEN, F.R.C.S.,
ASSISTANT SURGEON OF, AND LECTURER ON ANATOMY AT, ST. BARTHOLOMEW'S HOSPITAL,
LONDON.

With Notes and Additions

By ERSKINE MASON, M.D.,
DEMONSTRATOR OF ANATOMY AT THE COLLEGE OF PHYSICIANS AND SURGEONS, AND
SURGEON TO THE CHARITY HOSPITAL, NEW YORK.

Illustrated with Numerous Wood Engravings.

NEW YORK:
ROBERT M. DE WITT, PUBLISHER,
NO. 13 FRANKFORT STREET.

[1868.]

HARVARD UNIVERSITY
SCHOOL OF MEDICINE AND PUBLIC HEALTH

LIBRARY

Transferred from Mus. of Comp. Zool.

7 MAY 1947

A 3.A.1868.4

Entered, according to Act of Congress, in the year 1868, by
ROBERT M. DE WITT,
In the Clerk's Office of the District Court of the United States for the Southern District of
New York.

Electrotyped by
SMITH & MCDUGAL,
84 Beekman St., N.Y.

THE NEW YORK PRINTING COMPANY,
81, 83, and 85 Centre Street,
NEW YORK.

R. T. Jackson.

P R E F A C E .

IN preparing this edition of Holden's Manual of Anatomy, it has been the aim of the editor to place within reach of the American student the best guide, as he believes, that has yet appeared for use in the dissecting room: the work being neither too concise nor diffuse, whether viewed as a text-book for the student in Practical Anatomy, or for the Physician or Surgeon wishing to refresh his memory upon anatomical subjects. The illustrations are so clear and elaborate, and at the same time so true to nature; the language plain, and descriptions so perfect, that with such a guide, the difficulties which beset the beginner in his studies are in a great measure removed. Such additions have been made, both as regards the anomalies of the vessels and muscles, as, in the opinion of the editor, the student should be familiarized with while pursuing his studies, together with the measurement and weight of organs, and occasional reference made to some operation which it is well for him to practise upon the cadaver. These additions will be found enclosed within brackets—[]. But little originality is claimed in the preparation of this work; the additions being almost entirely compilations from standard works, arranged so as simply to be within reach of the student while pursuing his anatomical studies. The author-

ities from whence these have been taken are for the most part from Sharpey and Quain, Powers on the Arteries, Cruveilhier, and Todd's Cyclopædia of Anatomy and Physiology. The classification of the muscles according to their uses, after Dr. Barclay, together with the grouping of the vessels and nerves as given by Ellis in his Demonstrations of Anatomy, are also added, believing they may be of service for ready reference.

113 WEST 44TH STREET, }
February, 1868. }

A MANUAL

OF THE

DISSECTION OF THE HUMAN BODY.

THE DISSECTION OF THE NECK.

MAKE an incision through the skin, down the middle of the neck from the jaw to the sternum ; a second along the clavicle to the acromion ; a third along the base of the jaw as far as the mastoid process. Reflect the skin, and expose the cutaneous muscle called the *platysma myoides*. Between the platysma and the skin is a layer of adipose tissue, sometimes called the "superficial fascia." It varies in thickness in different subjects, but is generally more abundant at the upper part of the neck, especially in corpulent individuals, in whom it occasions a double chin.

Platysma Myoides.—The *platysma myoides* is the cutaneous muscle of the neck. It arises in the subcutaneous tissue over the pectoralis major and deltoid muscles ; thence proceeding over the clavicle and the side of the neck, its fibres become more closely aggregated, and terminate thus :—The anterior cross those of the opposite platysma, immediately below the symphysis of the jaw, and are lost in the skin of the chin ; the middle are attached along the base of the jaw : the posterior cross the masseter muscle, and terminate partly in the subcutaneous tissue of the cheek, partly in the muscles at the corner of the mouth.

The platysma forms a strong muscular defence for the neck. It is also a muscle of expression.* It is supplied with nerves by the cervical plexus, and by the cervical branch of the facial nerve.

Reflect the platysma. Beneath it lies the general investment of the neck, called the "*cervical fascia*." Upon this fascia we trace the superficial branches of the cervical plexus of nerves, the external jugular vein, and a smaller vein in front called the anterior jugular. These superficial veins are so variable in size and course, that a general description only is applicable.

External Jugular Vein.—The external jugular vein is formed within the substance of the parotid gland, by the junction of the temporal and internal maxillary veins. After receiving the transverse facial and posterior auricular veins, it appears at the lower border of the gland, crosses obliquely over the sterno-mastoid muscle (fig. 1), and joins the subclavian vein. A line drawn from the angle of the jaw to the middle of the clavicle would indicate its course. To trace the vein, during life, press upon it just above the clavicle; but do not be surprised if you fail to find it; it is sometimes wanting.

Near the angle of the jaw the external jugular vein communicates by a large branch with the internal jugular: hence bleeding from the external jugular relieves congestion of the brain.

* If the entire muscle be permanently contracted, it may occasion wry-neck, though distortion from such a cause is an exceedingly rare occurrence. A case in point is related by Mr. Gooch (*Chirurg. Works*), in which a complete cure was effected, after the failure of all ordinary means of relief, by the division of the platysma a little below the jaw.

The platysma myoides belongs to a class of muscles called "*cutaneous*," from their office of moving the skin. There are not many in man, except upon the neck and face, and there is a little one (*palmaris brevis*) in the palm of the hand. To understand their use thoroughly we must refer to the lower orders of animals, in whom they fulfill very important functions, by moving not only the skin, but also its appendages. For instance, by muscles of this kind the hedgehog, porcupine, and animals of that family can roll themselves up and erect their quills: we are all familiar with the broad "*panniculus carnosus*" on the sides of herbivorous quadrupeds, which enables them to twitch their skins, and thus rid themselves from insects. In birds, too, these cutaneous muscles are extremely numerous, each feather having appropriate muscles to move it.

Before its termination the external jugular vein generally receives the supra-scapular, posterior scapular, and other unnamed veins: a disposition very unfavorable for the surgeon,

Fig. 1.




DIAGRAM OF THE SUPERFICIAL NERVES AND VEINS OF THE NECK.

because there is a *confluence of veins immediately over the subclavian artery* in the place where it is usually tied.

In opening the external jugular vein, remember that the puncture should be made in the direction of the fibres of the sterno-mastoid muscle. The fibres of the platysma, divided obliquely, will be less likely to contract over the orifice.

Anterior Jugular Vein.—The anterior jugular vein is situated more in the middle of the neck, and is much smaller than the external jugular. It commences by small branches below the chin, and runs down the front of the neck, nearly to the sternum: it then curves outwards, beneath the sterno-mastoid muscle, and opens into the external jugular, or perhaps the subclavian vein. We commonly meet with two anterior jugular veins, one on either side; immediately above the sternum they communicate by a transverse branch.

The size of the anterior jugular vein is inversely proportion-

ate to that of the external jugular. When the external jugular is small, or terminates in the internal jugular, then the anterior jugular becomes an important supplemental vein, and attains considerable size. It is not uncommon to find it a quarter of an inch in diameter, and I have seen it nearly half an inch. These varieties should be remembered in tracheotomy.

Superficial absorbent glands are sometimes found near the cutaneous veins of the neck. They are small, and escape observation unless enlarged by disease. One or two are situated over the sterno-mastoid muscle; others about the mesial line.

Cutaneous Nerves of the Neck.—The cutaneous nerves of the neck are superficial branches of the cervical plexus: the plexus itself cannot at present be seen. It lies under the sterno-mastoid muscle, close to the transverse processes of the four upper cervical vertebræ, and is formed by the communications of the anterior divisions of the four upper cervical nerves. The cutaneous branches of the plexus emerge beneath the posterior border of the sterno-mastoid, and take different directions. We divide and name them thus: (fig. 1)

Cutaneous branches of the cervical plexus.	{	Ascending branches . . .	{ Auriculo-parotidean.
		Transverse branch . . .	{ Occipitalis minor.
		Descending branch . . .	{ Superficialis colli.
			{ Sternal.
			{ Clavicular.
			{ Acromial.

The *auriculo-parotidean n.* comes from the second and third cervical nerves, and ascends obliquely over the sterno-mastoid muscle, near the external jugular vein, towards the parotid gland. Near the gland it divides into two principal branches, of which the anterior is distributed to the skin over the gland and the side of the cheek; the posterior ascends to the cartilage of the ear, and ramifies chiefly upon its occipital surface. Other filaments communicate in the substance of the parotid gland with branches of the portio-dura, or facial nerve.

The *n. occipitalis minor* comes from the second cervical nerve. It runs near the posterior border of the sterno-mastoid muscle to the occiput, where it supplies the back of the scalp. Beneath the sterno-mastoid this nerve commonly forms a loop which embraces the nervus accessorius, and sends a branch to it.

The *transverse branch*, called the *n. superficialis colli*, comes

from the second cervical nerve. It passes forwards over the sterno-mastoid muscle, and supplies the front of the neck. Some of its filaments ascend towards the jaw, and join the cervical branch of the facial nerve.

The *descending* branch is derived from the fourth cervical nerve. It subdivides into three branches, which cross over the clavicle, and supply the skin of the front of the chest and shoulder. Of these, one, called the *sternal*, supplies the skin over the upper part of the sternum: another, the *clavicular*, passes over the middle of the clavicle, and is distributed to the skin over the pectoral muscle, the mammary gland, and the nipple; the third, named *acromial*, crosses over the acromion to supply the skin of the shoulder.

Reviewing these cutaneous branches of the cervical plexus, we find that they have a very wide distribution: for they supply the skin covering the following parts, viz: the ear, the back of the scalp, the parotid gland, the front and side of the neck, the upper and front part of the chest and shoulder.

Cervical Branch of the Facial Nerve.—Look for this branch beneath the fascia near the angle of the jaw. It leaves the parotid gland, and divides into filaments which curve forwards below the jaw; some of these join the transverse branch of the cervical plexus; others supply the platysma.

Cervical Fascia.—Now turn your attention to the membranous investment, called "*cervical fascia*," which encloses the several structures of the neck. In some subjects this fascia is very thin; in others, with strong muscles, it is proportionably dense and resisting. It is always relatively stronger in particular situations, for the more effective protection of the parts beneath; for instance, in front of the trachea, in the fossa above the clavicle, and below the angle of the jaw. It not only covers the soft parts of the neck collectively, but, by its inflections, forms separate sheaths for the muscles, vessels, and glands. It isolates them, and keeps them in their proper relative position. A lengthened description of its numerous layers would be not only extremely tedious, but unintelligible, without considerable knowledge of the anatomy of the neck. I propose, therefore, to give only a general outline of the fascia, and to describe,—first, its attachments at the upper part

of the neck; secondly, its attachments at the lower part; and lastly, its intermediate attachments.

At the upper part of the neck, the cervical fascia is attached along the base of the jaw, and especially to the angle, from which it extends backward to the styloid process and forms what is called the "stylo-maxillary ligament." It is also attached to the mastoid process, and the superior curved line of the occipital bone.

At the lower part of the neck, the fascia presents two clearly defined layers, a superficial and a deep one. The superficial layer is attached to the *front* of the sternum, and along the clavicle. The deep layer is attached to the *back* of the sternum and along the edge of the first rib: from thence we trace this deep layer *under* the clavicle along with the axillary vessels and nerves into the axilla. The interval between these two layers is especially manifest immediately above the sternum; here it contains more or less fat and one or two small absorbent glands.

The intermediate attachments of the fascia are to the os hyoides, to the transverse and the spinous processes of the cervical vertebræ.

A correct knowledge of the attachments of the principal layers of the cervical fascia is essential to a right understanding of the course which pus takes when it forms in the neck. For instance, suppose the pus to be formed at the lower part of the neck. If it be seated immediately under the superficial layer (which is attached to the clavicle), it may burrow beneath the clavicle into the axilla. But if it be seated beneath the deep layer (which is attached to the first rib), then it becomes a much more serious affair, since the pus may readily travel through the loose tissue by the side of the pharynx, and make its way into the chest, where it will probably burrow down the anterior or the posterior mediastinum, and produce the most disastrous results by bursting into the trachea or the œsophagus.

Besides forming sheaths for the several structures of the neck, there are other purposes to which the cervical fascia is subservient. The firm attachment of its layers to the sternum, the first rib, and the clavicle, forms a fibrous barrier at the

upper opening of the chest, which supports the soft parts, and prevents their yielding to the pressure of the atmosphere during inspiration. Dr. Allan Burns* first pointed out this important function of the cervical fascia, and has recorded a case exemplifying the results of its destruction by disease.

[In the case of a gentleman who suffered from an abscess at the root of the neck, which destroyed the fascia and tissues of this part, the trachea and the anterior innominata were found, after recovery, to be merely covered by a very thin pellicle of polished skin. An insufficient protection against the pressure of the external air upon the trachea, which so diminished its calibre as to render his respiration laborious.]

Moreover, the great veins at the root of the neck, namely, the internal jugular, subclavian, and innominate, are so closely united by means of the cervical fascia to the adjacent bones and muscles, that when divided they gape. They are, as the French express it, "*canalisées*," and are therefore better able to resist the pressure of the atmosphere, which tends to render them flaccid and impervious during inspiration. But this anatomical disposition of the great veins renders them more liable to the entrance of air when wounded. The danger of this occurrence is well known. Many deaths are recorded, resulting from the sudden entrance of air into the veins during operations about the neck, or even the axilla.

Sterno-cleido-mastoideus.—The sterno-cleido-mastoideus *arises* by a flat tendon from the upper part of the sternum, and by fleshy fibres, from the sternal third of the clavicle. It is *inserted* into the mastoid process, and about the outer half of the superior semicircular ridge of the occipital bone.

Observe that the sternal origin of the muscle is at first separated from the clavicular by a slight interval ; subsequently the sternal fibres gradually overlap the clavicular.

[The interval at the origin of this muscle is sometimes wanting. Its sternal origin may be very broad, arising from more than a third of the clavicle. The lower extremity may reach as low as the rectus abdominis, or the ensiform cartilage of the sternum.]

The muscle is confined by its strong sheath of fascia, in such

* Surgical Anatomy of the Head and Neck.

a manner that it forms a slight curve, with the convexity forwards. Observe especially that its front border overlaps the common carotid artery; along this border we make the incision in the operation of tying the vessel.

Action of Sterno-mastoid.—When both sterno-mastoidei act simultaneously, they draw the head and neck forwards, and are therefore especially concerned in raising the head from the recumbent position. When one sterno-mastoid acts singly, it turns the head obliquely towards the opposite shoulder; in this action it co-operates with the splenius of the other side.*

[We should observe that the anterior jugular vein is liable to be wounded in the division of this muscle, as it passes directly behind it, and runs for a short distance along the upper and posterior part of the clavicle.]

On emergency, the sterno-mastoid acts as a muscle of inspiration, by raising the sternum, its fixed point being, in this case, at the head. The sterno-mastoid is supplied with nerves by the n. accessorius and by a branch of the cervical plexus. It has three nutrient arteries: an upper mastoid, a middle, and a lower.

The upper mastoid, a branch of the occipital artery, enters the muscle with the n. accessorius; the middle mastoid is a branch of the superior thyroid; the lower mastoid is a branch of the supra scapular.

Triangles of the Neck.—Anatomists avail themselves of the sterno-mastoid muscle, to divide the neck on each side into two great triangles, an anterior and a posterior (fig. 2). The base of the anterior triangle is formed by the jaw, its sides by the mesial line and the front border of the sterno-mastoid. The posterior has the clavicle for the base, while the sides are defined by the hind border of the sterno-mastoid, and the free border of the trapezius.

The omo-hyoid muscle, which crosses the neck under the sterno-mastoid, subdivides these great primary triangles into


* The single action of the muscle is well seen when it becomes rigid and causes a wry neck. Other means of relief failing, the division of the muscle near its origin is sometimes beneficial in curing the distortion. In deciding as to the propriety of this operation, we should be careful to examine the condition of the other muscles, lest, after having divided the sterno-mastoid, we should be disappointed in removing the deformity.

four smaller ones (1, 2, 3, 4, fig. 2), of unequal size; an anterior superior, an anterior inferior, a posterior superior, and a posterior inferior. The direction of the omo-hyoid muscle renders their boundaries at once obvious.

[The anterior superior is sometimes called the superior carotid, or omo-mastoid; the anterior inferior, the inferior carotid, or omo-tracheal; the posterior superior, the sub-occipital, and the posterior inferior, the sub-clavian.]

Nervus Accessorius.—The upper part of the sterno-mastoid is traversed obliquely by a large nerve called the spinal accessory.* This nerve, one of the three divisions of the eighth pair of cerebral nerves, arises from the cervical portion of the spinal cord by a series of filaments from the same tract which gives origin to the anterior or motor roots of all the

Fig. 2.



TRIANGLES OF THE NECK.

spinal nerves. It ascends through the foramen magnum into the skull, leaves the skull through the foramen lacerum posterius, runs behind the internal jugular vein, passes obliquely through the upper third of the sterno-mastoid, and crosses the posterior triangle of the neck to the trapezius, to which it is distributed.

* Nervus accessorius of Willis (Willis, nervor. descr. cap. xxiii; Opera omnia, Amst. 1682).

The *nervus accessorius* supplies the sterno-mastoid, and, after leaving the muscle, is joined by a branch from the second or third cervical nerve.

To find the *nervus accessorius* before it enters the sterno-mastoid, dissect deep near the angle of the jaw, and you will see it emerging beneath the posterior belly of the digastricus, close to the transverse process of the atlas (fig. 2). The upper mastoid artery, a branch of the occipital, runs into the muscle with the nerve.

Depressor Muscles of the Os-hyoides and Larynx.—Let us now examine the flat muscles in front of the neck, which pull down the larynx; namely, the sterno-hyoid, sterno-thyroid, and omo-hyoid.* Remove the fascia which covers them; disturb them as little as possible, and take care of the nerves (branches of the *descendens noni*), which enter their outer borders. At this stage of the dissection, reflect the sternal portion of the sterno-mastoid from its origin: you will thus obtain a better view of your work.

Sterno-hyoid.—The sterno-hyoid *arises* from the back part of the sternum and sterno-clavicular ligament, and is *inserted* into the body of the os-hyoides. This is the most superficial of the muscles in front of the neck. Notice that opposite the cricoid cartilage and the two or three upper rings of the trachea, it is in contact with its fellow of the opposite side. We cut in the mesial line between these muscles in laryngotomy.

Sterno-thyroid.—The sterno-thyroid *arises* from the back part of the sternum and the cartilage of the first rib, and is *inserted* into the oblique ridge on the thyroid cartilage.

This muscle is situated immediately under, and is much broader than, the sterno-hyoid. It covers the thyroid gland, and for some distance above the sternum is in contact with its fellow of the other side; the result of which is, that the trachea is completely covered by muscular fibres.

[These muscles may be double, and arise from the same


* The sterno-hyoid and sterno-thyroid muscles often present slight transverse tendinous lines. These tendinous intersections are quite rudimentary in man; but in some animals with long necks, *e. g.* the giraffe, they are so developed that each depressor muscle is composed of alternations of muscle and tendon.

osseous points. We have seen the sterno-thyroids joined together by transverse fibres.]

Omo-hyoid.—The omo-hyoid *arises* from the notch in the scapula and the ligament over it; and is *inserted* into the body of the os-hyoides near the great cornu.

This muscle is digastric; that is, it consists of two fleshy portions connected by a tendon. Pay attention to its direction (fig. 3). From the scapula it comes nearly horizontally forwards across the lower part of the neck, and passes beneath the sterno-mastoid, *over* the sheath of the great vessels of the

Fig. 3.



CENTRAL LINE OF THE NECK.—COURSE AND RELATIONS OF COMMON CAROTID A.

neck; then, changing its direction, it ascends nearly vertically close to the outer border of the sterno-hyoid. Observe, therefore, that the muscle does not proceed straight from origin to insertion, but that it forms an obtuse angle beneath the sterno-mastoid. Now the intermediate tendon is situated at the angle, and it is maintained in its angular position by a broad layer of the cervical fascia, which comes up to it from the clavicle. What is the object of this peculiar direction of the omo-hyoid? I think it is to keep tense that part of the cervical fascia which covers the apex of the lung, and thus to resist atmospheric pressure.

[The size and direction of this muscle often vary. The tendinous intersection may be altogether or partially wanting; the posterior belly may be wanting; the muscles arising from the centre of the clavicle, or the anterior belly, be absent, it terminating in the fascia over the artery. We have met with this anomaly several times.]

Action of the Depressor Muscles.—The sterno-hyoid, sterno-thyroid, and omo-hyoid muscles, co-operate in fixing the larynx and os-hyoides, *e. g.* in sucking, or they depress the larynx after it has been raised in deglutition. Again, they depress it in the utterance of grave notes. That the larynx is raised or depressed according to the height of the note may be ascertained by placing the finger upon it while we go through the gamut.

Now these depressor muscles are all supplied with nerves (fig. 3, p. 11) by the “*descendens noni*” (a branch of the ninth, or hypoglossal), and by the “*communicantes noni*” (branches of the cervical plexus). The *descendens noni* sends a separate branch to each belly of the omo-hyoid. They are supplied with blood by the superior thyroid artery.

Thyro-hyoid.—The thyro-hyoid *arises* from the oblique line on the ala of the thyroid cartilage, and is *inserted* into the body and half the great cornu of the os-hyoides.

This muscle is a continuation of the sterno-thyroid. It is supplied by a special branch from the hypoglossal nerve. It covers the thyro-hyoid membrane, and the superior laryngeal nerve and artery as they enter the larynx.

We are now about to examine the course and relations of the common carotid artery. But before we do so, you must

particularly notice the strong layer of fascia which lies under the sterno-mastoid, and forms the under part of its sheath. It is attached to the angle of the jaw, thence descends over and protects the great vessels of the neck, and is firmly connected to the clavicle and first rib. This fascia prevents matter from coming to the surface, when suppuration takes place by the side of the pharynx.

Remove this fascia, taking care not to remove with it the *descendens noni* and *communicantes noni* nerves, which cross the sheath of the common carotid. Dissect out any absorbent glands which lie about the sheaths of the great vessels.

Course and Relations of the Common Carotid Artery.—

The common carotid arises on the right side from the *arteria innominata*; on the left, from the arch of the aorta. It ascends in front of the bodies of the cervical vertebræ, by the side of the trachea, thyroid gland, and larynx, as high as the upper border of the thyroid cartilage, and then divides into the external and internal carotid. Thus a line drawn from the sternal end of the clavicle to the angle of the jaw will nearly indicate its course. It is contained in a sheath of cervical fascia. In the same sheath are the internal jugular vein and the pneumogastric nerve. The vein lies on the outer side of, and parallel with the artery: the nerve lies behind and between the artery and vein. Behind the sheath is the sympathetic nerve. Lastly, along the vertebral column the sheath lies successively upon the *longus colli* and the *rectus capitis anticus major* muscles.

[Frequently well marked septa are found within the sheath, dividing it into three separate compartments for the vein, artery and nerve.]

At the lower part of the neck the carotid artery is deeply seated:—it is covered by the sternal portion of the sterno-mastoid, the sterno-hyoid and thyroid muscles, and about two inches above the clavicle it is crossed by the *omo-hyoid*. Above this point the artery becomes more superficial, and is covered by the *platysma*, cervical fascia, and only slightly overlapped by the sterno-mastoid. Lying immediately upon the sheath of the artery we find the *descendens noni* and *communicantes noni* nerves. Above all, observe that the sheath is crossed by three veins, namely, the facial, the supe-

rior, and inferior thyroid, which empty themselves into the internal jugular. This is the general rule, and we draw especial attention to it, because these veins are so liable to be overlooked and injured in the operation of tying the carotid.

It is plain that it is easier to tie the common carotid above the omo-hyoid than below it. We make an incision (three inches long) down the inner border of the sterno-mastoid; we cut through the platysma and cervical fascia, draw aside the overlapping edge of the sterno-mastoid, and expose the sheath of the vessel. We then make a small opening in the sheath large enough to admit the aneurismal needle, and tie the vessel, taking care not to include the pneumogastric or descendens noni nerves in the ligature.

[To tie the artery, the head should be well thrown back, and when the operation is to be above the omo-hyoid, an incision should be made a little below the angle of the jaw to the cricoid cartilage. When below this muscle, we should commence on a level with the cricoid cartilage. The sterno-mastoid must be well drawn to the outside before the vessel is observed; the descendens noni here runs on the tracheal side, and the recurrent laryngeal nerve behind it. In either operation the needle should be passed from without, inwards.]

In what respects the Left Carotid differs from the Right.—In the first part of its course the left carotid differs from the right in the following particulars:—

1. It arises from the arch of the aorta, is therefore longer and deeper seated than the right, and is covered by the first bone of the sternum.
2. It is crossed by the left vena innominata.
3. It is in close relation with the œsophagus.
4. It is in close relation with the left recurrent nerve.
5. It is in close relation with the left thoracic duct.

Division of the Common Carotid.—The common carotid at its division is often a little bulbous. This dilatation is in some instances so marked, that during life I have seen it mistaken for an incipient aneurism. It is necessary to be aware that the carotid sometimes divides much lower than usual. Several times I have seen the division as low as the level of the cricoid cartilage.

[The common carotid may be absent—the external and inter-

nal arising separately from the arch. Its bifurcation has been observed both above the angle of the jaw, and as low down as the sixth cervical vertebra. It may also give off the inferior thyroid, superior laryngeal, pharyngeal ascendens, superior thyroid and right vertebral.]

Internal Jugular Vein.—The internal jugular vein returns the blood from the brain. Leaving the skull at the foramen jugulare, the vein descends the outer side of the carotid, but in the sheath with it, and joins the subclavian vein at nearly a right angle to form the vena innominata. In its course down the neck it receives the pharyngeal, occipital, lingual, facial, superior and inferior thyroid veins.

Previous to its termination the internal jugular advances slightly to meet the subclavian vein, so that it lies on a plane a little anterior to the carotid. Besides which, the direction of the vein being perpendicular, and that of the artery oblique, there is necessarily a small interval between them at the lower part of the neck, more *especially on the right side*. In this interval we find the pneumogastric nerve, and, deeper, the vertebral artery. Here, too, on the left side, we look for the thoracic duct.

Descendens noni.—The descendens noni (page 11), a branch of the hypoglossal, runs down obliquely over the sheath of the carotid to supply the depressor muscles of the os-hyoides. Trace the nerve upwards, and you will see that it leaves the hypoglossal where this nerve curves round the occipital, or perhaps the mastoid artery. For a short distance the descendens noni lies within the carotid sheath; but about the level of the os-hyoides it comes through the sheath, and crosses obliquely the carotid, from the outer to the inner side. Now the descendens noni is reinforced by one or more nerves termed communicantes noni (derived from the second and third cervical nerves). These communicating branches descend on the outer side of the internal jugular vein, and form generally two loops in front of the carotid sheath. From these loops the nerves proceed to the depressor muscles of the larynx.

In some subjects the descendens noni seems to be wanting; but look carefully and you will probably discover it concealed *within* the carotid sheath: in such case the reinforcing loops

from the cervical nerves will be found behind the internal jugular vein.

Let us now examine the thyroid body, and in order to expose it, let us reflect the sterno-hyoid and thyroid muscles from their insertions, so that they can be replaced if necessary. Afterwards we will examine the absorbent glands of the neck, and then survey the objects in the central line of the neck, from the jaw to the sternum.

Thyroid body.—This very vascular gland-like substance lies over the front and sides of the upper part of the trachea, and extends upwards on each side of the larynx. It consists of two lateral lobes, connected a little below the cricoid cartilage by a transverse portion called the "isthmus." Each lobe is conical, with the base opposite the sixth or seventh ring of the trachea, and the apex by the side of the thyroid cartilage. Its anterior surface is covered by the sterno-hyoid and thyroid muscles; its deep surface clasps the sides of the trachea and larynx, and usually extends so far backwards as to be in contact with the pharynx. Its external border overlaps, in most cases partially, but sometimes completely, the common carotid artery, particularly on the right side; and there are instances in which the lobe is deeply grooved by the vessel.

The isthmus lies over the 2nd, 3rd, and 4th rings of the trachea. This portion of the organ varies much in its dimensions. In some instances there is no transverse portion. This corresponds with the normal disposition in most of the lower orders of mammalia; but in man it is a failure in the union of the two halves by which the organ is originally developed. Generally, the vertical measurement of the isthmus is about one inch. Between its upper border and the cricoid cartilage is a space about four or five lines in extent, where the trachea is free; this space therefore is the more preferable situation for tracheotomy. But the vertical measurement of this isthmus is sometimes of very considerable length. I have seen it covering the trachea almost down to the sternum.*

* From the upper part of the isthmus, or from the adjacent border of either lobe, most commonly the left, a conical prolongation of the thyroid body, called "the pyramid," frequently ascends in front of the crico-thyroid mem-

The thyroid body is closely connected, by fibrous tissue, to the sides of the trachea and cricoid cartilage. This explains why it rises and falls with the larynx in deglutition.

Taken as a whole, the thyroid varies in size in different individuals and at different periods of life. It is relatively larger in the child than the adult, in the female than the male. In old age it diminishes in size, becomes firmer, and occasionally contains earthy matter.

But by far the most notable considerations in respect to the thyroid body are the number, the large size, and the free innosculation of its blood-vessels. In fact, it appears to be composed of a tissue of arteries and veins. The superior thyroid arteries come from the external carotid, and enter the front surface of the apex of each lobe; the inferior thyroid come from the subclavian, and enter the under surface of the base. An artery, called the middle thyroid, is observed in some subjects; it comes from the *arteria innominata*, or the arch of the aorta, and ascends directly in front of the trachea to the isthmus.

Its veins are equally large, and form a plexus upon it. The superior and inferior thyroid veins cross the common carotid, and open into the internal jugular. The middle thyroid veins descend over the front of the trachea, communicate freely with each other, and terminate in the left *vena innominata*. When you perform tracheotomy, bear in mind the size of these middle thyroid veins, and the possible existence of a middle thyroid artery.

Its nerves are furnished by the laryngeal branch of the pneumogastric and the cervical ganglia of the sympathetic. They accompany the arteries.

Structure of the Thyroid Body.—The thyroid body is

brane, as high as the "*promontorium Adami*," and is connected to the body of the *os-hyoides* by fibrous tissue. In some subjects we may observe a few muscular fibres passing from the *os-hyoides* to the pyramid. This constitutes the "*levator glandulæ thyroideæ*" (see preparation in Museum of St. Barth. Hosp., Patholog. Series, No. 14,) of some anatomists. There are instances in which the pyramid is double; and lastly, I have seen a considerable portion of this thyroid substance lying over the crico-thyroid membrane, completely isolated from the rest of the organ. These varieties deserve the notice of the surgeon, because any one portion of this structure may become enlarged, independent of the rest, and occasion a bronchocele.

invested by a thin covering of condensed areolar tissue, which penetrates into it and divides it into lobes. It consists of a multitude of vesicles which do not communicate with each other, and vary in size. Some of them may be recognized with the naked eye; but the greater number require the aid of the microscope. In hypertrophy of the gland we sometimes see them as large as a horse-bean, or even larger. They contain a glairy, transparent fluid, full of nucleated cells. The arteries ramify most minutely upon their walls. Of its function nothing is with certainty known. The presumption is that it is concerned in the elaboration of the blood.

An enlargement of the thyroid body is termed a "Bronchocele." If the relation of its lobes to the trachea and œsophagus be properly understood, it is easy to predicate the consequences which may result from their enlargement. The nature and severity of the symptoms will to a certain extent be determined by the part of the organ affected. For instance, if the isthmus be enlarged, difficulty in breathing will probably be the prominent symptom; and an enlargement of the left lobe is more likely to produce a difficulty in swallowing, on account of the inclination of the œsophagus towards the left side.

An instance is related by Allen Burns in which the isthmus was placed between the trachea and the œsophagus. It must be obvious that enlargement of a part so situated would occasion most fearful difficulty in swallowing. I have seen two cases in which the lateral lobes projected so far inwards, that they completely embraced the back of the œsophagus.

[The normal weight of this body may be stated as from one to two ounces: in disease it has been known to weigh a pound and a half; its transverse measurement three inches, and its vertical about two inches. This body has been successfully removed for enlargement by Dr. Voss of this city; and the superior and inferior thyroid arteries have been ligated by Dr. Thomas M. Markoe for bronchocele, with the result of greatly diminishing the size of the tumor.]

Small absorbent glands are observed about the thyroid body, especially in front of the trachea; one is often situated over the crico-thyroid membrane. These glands, if enlarged by disease, might be mistaken for a small bronchocele.

Deep Cervical Absorbent Glands.—In the areolar tissue which surrounds the great vessels of the neck, we meet with a series of absorbent glands, called the deep cervical. They form an uninterrupted chain (whence their name *glandulæ concatenatæ*), from the base of the skull, along the side of the neck, to the clavicle, beneath which they are continuous with the thoracic and the axillary glands. Notice that some of these glands lie anterior to the common carotid artery, others, between it and the spine. This disposition explains the well-known fact, that, when these glands are enlarged, the great vessels and nerves of the neck are liable to become imbedded in their substance.

The glands are particularly numerous near the division of the common carotid, by the side of the pharynx, and the posterior belly of the digastricus. The absorbents connected with them come from all parts of the head and neck. These vessels unite, to form, on both sides of the neck, one or more absorbent trunks, called the jugular. On the left side this jugular trunk joins the thoracic duct, or opens by a separate orifice into the left subclavian vein: on the right it always opens into the subclavian vein.

The contiguity of the glands to the great vessels and nerves of the neck explains the symptoms produced by their enlargement. The tumor may be so situated as to be raised and depressed by the pulsation of the carotid, and thus simulate an aneurism. A careful examination, however, will distinguish between a real and an apparent pulsation. By grasping the tumor we become sensible that the rising and falling does not depend upon any variation of its magnitude, but upon the impulse derived from the artery; consequently, if we lift the tumor from the vessel, all feeling of pulsation ceases.

Survey of the Central Line of the Neck.—Study well the parts in the central line of the neck (fig. 3, p. 11). Beginning at the chin, we observe the insertions of the digastric muscles. Below these is the junction or “raphé,” of the mylo-hyoid muscles. Then comes the os-hyoides. Below the os-hyoides comes, after a short interval, the pomum Adami, or notch of the thyroid cartilage. Below the thyroid cartilage is the cricoid. These two cartilages are connected by the crico-thyroid membrane. Below the cricoid c. is the trachea. This is

crossed by the isthmus of the thyroid body, and lower down it is covered by the middle thyroid veins.

Now the chief surgical interest lies just above and below the cricoid cartilage. We can feel this cartilage very plainly in the living subject. In laryngotomy, the crico-thyroid membrane is divided transversely. Remember that the membrane should be divided *close* to the edge of the cricoid c., for two reasons:—1. In order to be far from the vocal cords. 2. To avoid the crico-thyroid artery which crosses the middle of the membrane. If more room be required, the cricoid cartilage should be divided longitudinally.

In tracheotomy, we open the trachea by a perpendicular incision, either above the isthmus of the thyroid body, or below it. The operation above the isthmus, if there be space enough for the introduction of the canula, is the easier and safer of the two; for here the trachea is nearer to the surface, and no large blood-vessels are, generally speaking, in the way. The space available measures from a quarter to half an inch; and the isthmus is not so firmly adherent to the trachea as to prevent its being drawn downwards for a short distance. However, it is fair to state, that in one case out of every eight or ten, there is *no* available space.

Tracheotomy below the isthmus, is neither an easy nor a safe operation, for many reasons. 1. The trachea recedes from the surface as it descends, so that just above the sternum it is nearly an inch and a half from the skin. 2. The large middle thyroid veins are in the way. 3. A middle thyroid artery may run up in front of the trachea, direct from the arteria innominata. 4. The arteria innominata itself lies sometimes upon the trachea higher than usual, and may, therefore, be in danger. 5. The left vena innominata in some cases crosses the trachea above the edge of the sternum instead of below it. The celebrated French surgeon Bécларd used to relate in his lectures the following occurrence:—A student had fallen into the Seine, and was nearly drowned. As he was recovering very gradually, some kind friends attempted to accelerate the process by making an opening into the trachea. In so doing they unhappily wounded the vena innominata. Blood poured into the trachea, and the result was instantly fatal.

Some surgeons advocate the practice of dividing the isth-

mus vertically, in order to get at the trachea beneath it. Let such surgeons remember that an artery, which would bleed freely if divided, generally runs along the upper edge of the isthmus. The artery is a branch of the superior thyroid, and inosculates with a corresponding branch on the opposite side.

Whoever pays attention to this subject in the dissecting-room will soon be convinced of the fact, that not only large veins but large arteries occasionally cross the crico-thyroid membrane as well as the trachea. Every winter session convinces me more and more of the necessity of cutting *cautiously* down to, and fairly exposing the air tube, before we venture to open it.

[When an incision is made through the crico-thyroid membrane, it is termed laryngotomy; when through the cricoid cartilage and first ring of the trachea, laryngo-tracheotomy: below the isthmus of the thyroid body, tracheotomy. The first of these operations is evidently the most readily performed, and should be selected when tracheotomy is not absolutely indicated. In the child, tracheotomy is exceedingly difficult at times, owing to the trachea being smaller and more deeply placed, and more movable, than in the adult. We also have the thymus body in the infant, ascending a variable distance.]

Dissection of the Submaxillary Region, or the Digastric Triangle.—When the platysma and the cervical fascia have been removed from their attachment to the jaw, the most conspicuous object is the submaxillary gland. Observe that the fascia forms for it a complete case. Beneath the jaw we find several absorbent glands, of which some lie superficial to the salivary gland, others beneath it. These glands receive the absorbents of the face, the tonsils, and the tongue.


A little dissection will expose a muscle called the digastricus, consisting of two distinct portions connected by a tendon. They form, with the body of the jaw, a triangle, of which we propose to examine the contents. And first of the digastric muscle itself.

Digastricus.—The digastricus *arises* from the digastric fossa of temporal bone, and is *inserted* close to the symphysis of the lower jaw.

Raise the submaxillary gland to see the intermediate tendon of

the digastricus, the angle which it forms, and how it is fastened by aponeurosis to the os-hyoides : observe also that this aponeurosis is connected in the mesial line with its fellow of the opposite side, so that a fibrous expansion occupies the interval between the anterior portions of the digastrici.

Fig. 4.



DIGASTRIC TRIANGLE AND CONTENTS.

The chief action of the digastricus is to depress the lower jaw. But if the lower jaw be fixed, then the muscle raises the os-hyoides, as in deglutition.

The posterior belly of the digastricus is supplied by a nerve from the facial ; the anterior belly by a branch from the mylo-hyoidean nerve (which comes from the third division of the fifth pair).

Stylo-hyoideus.—The stylo-hyoideus *arises* from the styloid process of the temporal bone, and is *inserted* into the body of the os-hyoides.

This muscle runs close along the posterior belly of the digastricus. Most frequently the digastric tendon runs through the substance of it. Its nerve comes from the facial. Its action is to raise and draw back the os-hyoides.

The objects which we have to examine in the digastric triangle are twelve in number, as follows:—

- | | |
|-----------------------------------|---|
| 1. Submaxillary gland. | 7. Stylo-maxillary ligament. |
| 2. Facial vein. | 8. Part of the parotid gland. |
| 3. Facial artery. | 9. Part of the external carotid artery. |
| 4. Submental artery. | 10. Mylo-hyoideus muscle. |
| 5. Mylo-hyoidean nerve. | 11. Hypoglossal nerve. |
| 6. Submaxillary absorbent glands. | 12. Part of the hyo-glossus muscle. |

Submaxillary Salivary Gland.—In the ordinary position of the head, the submaxillary gland is partially concealed by the jaw, but when the head falls back the gland is more exposed. It is about the size of a chestnut, and is divided into several lobes. Its upper margin is covered by the body of the jaw; its lower margin overlaps the side of the os-hyoides. Its cutaneous surface is flat, but the lobes on its deep surface are irregular, and often continuous with those of the sub-lingual gland. By raising the gland we find that it lies upon the mylo-hyoideus, the hyo-glossus, the tendon of the digastricus, and a portion of the hypoglossal nerve, seen above the tendon. Mark these relations well, because they are of importance, as we shall presently explain, in tying the lingual artery.

The duct of the gland (Wharton's duct*) passes from its under surface, runs forward under the mylo-hyoideus, and opens into the floor of the mouth, by the side of the frenum linguæ. In length it is about two inches; its dimensions are not equal throughout, for it is dilated about the middle, and contracted at the orifice. Saliva, collected in the dilated portion, is sometimes spirted to a considerable distance out of the narrow orifice, in consequence of the sudden contraction of the neighboring muscles.

An abnormal dilatation of the submaxillary duct is called a "ranula." It makes a tumor with semi-transparent walls, per-

* Thom. Wharton, *Adenographia, seu glandularum totius corporis descriptio*. 12mo. Amstel. 1659.

ceptible beneath the tongue. In some cases, however, what appears to be a "ranula" is in reality a cyst, formed in the loose areolar tissue under the tongue, or an enlargement of one of the small bursæ which are normally placed here.

Facial vein.—The facial vein does not accompany the facial artery. It leaves the face at the anterior edge of the masseter m., and then runs *over* the submaxillary gland, the digastricus and stylo-hyoideus and the carotid artery, to join the internal jugular. This is the rule,—but there are frequent exceptions. The principal point to remember is, that the vein runs superficial to the gland, and that we must be cautious in opening abscesses under the jaw.

Facial artery.—The facial artery is the third branch of the external carotid. It runs tortuously under the hypoglossal nerve, the posterior belly of the digastricus and stylo-hyoideus, and underneath or through the substance of the submaxillary gland to the face, where it appears at the anterior border of the masseter. Below the jaw the facial gives off the four following branches :—

1. The *ascending palatine* runs up between the stylo-glossus and the stylo-pharyngeus m. to the soft palate and tonsils, and inosculates with the *descending palatine*, a branch of the internal maxillary.

2. The *tonsillar* runs up between the internal pterygoid and the stylo-glossus m. to the tonsil.

3. *Glandular* branches to the submaxillary gland.

4. The *submental* runs forward upon the mylo-hyoideus m., turns over the chin, and inosculates with the terminal branches of the inferior dental. It supplies the mylo-hyoideus, the anterior belly of the digastricus, and the sublingual absorbent glands.

[The facial often arises together with the lingual. If it comes off above its normal position, it resumes its usual course by passing down beneath the angle of the jaw. It may end simply as a submental artery, supplying only a portion of the face. The deficiency is then made up by branches of the transverse facial or internal maxillary, or by an increase in the size of the nasal branches of the ophthalmic. The ascending palatine may cease before it reaches the soft palate, its place being supplied by the pharyngeal artery. The tonsillar

branch is also found wanting at times. The submental branch may come from the lingual.]

Mylo-hyoidean nerve.—Look for the mylo-hyoidean nerve near the submental artery. The nerve comes from the inferior dental (before its entrance into the dental foramen), and running along a groove on the inner side of the jaw, advances between the bone and the internal pterygoid m., to supply the mylo-hyoideus and the anterior belly of the digastricus.

The submaxillary absorbent glands receive the absorbent vessels of the face and the tongue. We often see them enlarged in cancer of the tongue and cancer of the lower lip. It is worth remembering that there are absorbent glands in the mesial line below the chin.

Mylo-hyoideus.—The mylo-hyoideus *arises* from the mylo-hyoid ridge of the lower jaw, and is *inserted* into the body of the os-hyoides. Between this and the symphysis of the jaw, it joins its fellow in a mesial tendinous line, termed a "*raphé*."

Observe the posterior fibres only of the muscle are inserted into the os-hyoides; the anterior fibres successively decrease in length, and meet their fellow in the mesial line. Thus the muscles of the opposite sides form a muscular floor for the mouth. It is supplied with nerves by the mylo-hyoid branch of the dental, with blood by the submental a.

The muscles of opposite sides conjointly elevate the os-hyoides and the floor of the mouth,—as in deglutition.

Stylo-maxillary ligament.—This is simply a deep layer of the cervical fascia, extending from the angle of the jaw to the styloid process. It is a broad sheet of fascia, and separates the submaxillary gland from the parotid. It is continuous with the fascia covering the pharynx; this gives it a surgical interest, because it prevents accumulations of matter formed near the tonsils and upper part of the pharynx from coming to the surface.

The remaining objects seen in the submaxillary triangle, namely, the parotid gland, the external carotid, the hypoglossal nerve, and the hyo-glossus muscle will be described presently when they can be better seen. I wish now to call your attention to a little piece of surgical anatomy, which

will enable you readily to find and tie the lingual artery. It is this :

Suppose a horizontal incision to be made along the body and greater cornu of the os-hyoides through the skin, the platysma, and the cervical fascia, you will come at once upon the lower edge of the submaxillary gland. Lift up the gland, which can be easily done, and underneath it you will observe that the tendon of the digastricus makes two sides of a triangle, of which the base is formed by the hypoglossal nerve crossing the hyo-glossus m. Within this little triangle, cut through the fibres of the hyo-glossus, and under them is the lingual artery. The first time you perform this operation on the dead subject, you will probably miss the artery and cut (through the middle constrictor) into the pharynx.

Pursuing the dissection, reflect the anterior belly of the digastricus from its insertion into the jaw; reflect the mylo-hyoideus from the mesial line and the os-hyoides, and turn it over the body of the jaw. Thus we get at the parts which lie under the mylo-hyoideus. To obtain a full view of them, however, it is necessary to saw through the symphysis of the jaw, so as to be able to separate the two halves of it, and draw the tongue out of the mouth. This done, we have to make out the following objects, represented in fig. 5., p. 27.

- | | |
|-----------------------|---------------------------|
| 1. Genio-hyoideus. | 6. Sublingual gland. |
| 2. Hyo-glossus. | 7. Hypoglossal nerve. |
| 3. Stylo-glossus. | 8. Gustatory nerve. |
| 4. Genio-hyo-glossus. | 9. Submaxillary ganglion. |
| 5. Submaxillary duct. | 10. Lingual artery. |

Genio-hyoideus.—The genio-hyoideus *arises* from the symphysis of the jaw, and is *inserted* into the front of the body of the os-hyoides.


This round muscle is situated in the mesial line, and parallel to its fellow. Its nerve comes from the hypoglossal, and its blood from the lingual artery. Its action is to draw the os-hyoides forwards and upwards.

Hyo-glossus.—The hyo-glossus *arises* from the body and great cornu of the os-hyoides, and is *inserted* into the posterior two-thirds of the side of the tongue.

This is a square and flat muscle; its fibres ascend nearly

perpendicularly from origin to insertion. Its nerve comes from the hypoglossal and its blood from the lingual artery. Its action (with that of its fellow) is to depress the tongue. Observe the several objects which lie *upon* the hyo-glossus, namely, the hypoglossal and gustatory nerves, the submaxillary ganglion, the duct of the submaxillary gland, and the

Fig. 5.



MUSCLES, VESSELS, AND NERVES OF THE TONGUE.

sublingual gland. *Beneath* the hyo-glossus muscle lie the lingual artery, part of the middle constrictor of the pharynx, and part of the genio-hyo-glossus.

Genio-hyo-glossus.—The genio-hyo-glossus *arises* by a tendon from the tubercle behind the symphysis of the jaw, and is *inserted* into the os-hyoides and the tongue from base to tip.

This is the largest and most important of the muscles of the tongue. It is fan-shaped, with the apex attached to the symphysis. From thence its fibres radiate into the entire length of the tongue. It derives its nerves from the hypoglossal and its blood from the lingual artery. Its action is various. The posterior fibres, by raising the os-hyoides and

drawing forwards the base of the tongue, protrude the tongue out of the mouth; the anterior fibres draw the tongue back again. When every part of the muscle acts, it draws down the whole tongue, and is therefore one of the chief muscles concerned in suction.

Stylo-glossus.—The stylo-glossus *arises* from the apex of the styloid process and the stylo-maxillary ligament, and is *inserted* along the side of the tongue.

This is a long and slender muscle. It runs outside the hyo-glossus nearly to the tip of the tongue. Its nerve comes from the hypoglossal. Its action is to retract the tongue.

Hypoglossal nerve.—The hypoglossal or ninth cerebral nerve, is the *motor* nerve of the muscles of the tongue. It arises (by several roots) from the front of the medulla oblongata between the pyramid and the olive. It leaves the skull through the anterior condyloid foramen, and runs down obliquely between the internal carotid and the internal jugular vein. Just below the posterior belly of the digastricus, the nerve curves forward over the occipital, both carotid, and the facial arteries; it then crosses the hyo-glossus muscle, and divides into branches which supply all the muscles of the tongue, namely, the stylo-glossus, hyo-glossus, genio-hyo-glossus, lingualis, and the genio-hyoideus.

As it curves round the occipital artery, the hypoglossal n. sends the “*descendens noni*” to the depressors of the os-hyoideus, (page 11.) Besides this, it sends a separate nerve to the thyro-hyoideus. While crossing the hyo-glossus, it communicates with the gustatory nerve. (Fig 5.)

The hypoglossal at its origin is purely a *motor* nerve. But soon after it leaves the skull, it receives communications from the first cervical nerve. These communications are important physiologically for two reasons: 1. They account for the hypoglossal nerve containing sensitive fibres: 2. They contribute the greater part of the filaments of the “*descendens noni*.”

Sublingual Salivary Gland.—The sublingual gland lies immediately beneath the mucous membrane of the floor of the mouth. Its shape is oblong, with the long axis, (about an inch and a half) directed from before backwards. It rests upon the upper surface of the mylo-hyoid muscle, and to-

wards the mesial line it is in contact with the hyo-glossus and the genio-hyo-glossus.

The ducts of the sublingual gland (ducts of Rivinus*) vary in number from seven to twelve. They terminate by minute openings behind the orifice of the submaxillary duct, along a ridge upon the floor of the mouth. The ducts of some of the lobes terminate in the submaxillary duct.

The duct of the submaxillary gland may now be traced across the hyo-glossus, and under the gustatory nerve to the floor of the mouth.

Gustatory Nerve.—This nerve is a branch of the inferior maxillary, or third division of the fifth pair. It descends between the ramus of the jaw and the internal pterygoid muscle, and comes forward along the upper part of the hyo-glossus, crossing at an acute angle over the duct of the submaxillary gland. Having reached the under part of the tongue, the nerve divides into filaments which supply the papillæ on its anterior three-fourths.

Submaxillary Ganglion.—Look at the lower border of the gustatory nerve before it crosses the submaxillary duct, and you will find a small ganglion, about the size of a pin's head. Like the other ganglia in connection with the branches of the fifth pair, it receives nerve-filaments of three different kinds, viz., motor, sensitive, and sympathetic. Its motor root is the chorda tympani, which, though apparently a branch of the gustatory, is in point of fact derived from the facial nerve (a nerve of motion). Its sensitive roots proceed from the gustatory; and its connection with the sympathetic system of nerves is established by a branch which comes from the "nervi molles" round the facial artery. Thus provided with nerves from these sources, the ganglion supplies the submaxillary and sublingual glands and their ducts.

Lingual Artery.—The lingual artery is generally the second branch of the external carotid. Curving slightly upwards from its origin, the artery soon runs forwards beneath the hyo-glossus muscle, parallel to the os-hyoides. At the anterior edge of the hyo-glossus it ascends to the under-surface of the tongue, and is continued forward to the apex of the

* Aug. Quirin. Rivinus, de Dyspepsia. Lips. 1678.

tongue under the name of "ranine." The curves thus made by the artery are obviously for the purpose of allowing the elongation of the tongue. Observe that, under the hyoglossus, the artery lies upon the middle constrictor of the pharynx; under the tongue, it lies between the hyoglossus and the lingualis m. Its branches are as follow:—

1. The *hyoid*, a small artery which runs along the hyoid bone, supplying the muscles.
2. The *dorsales linguæ*, two or more, which run up under the hyoglossus to the back of the tongue.
3. The *sublingual*, which arises near the anterior border of the hyoglossus, and supplies the sublingual gland. This artery generally gives off the little artery of the "frænum linguæ," which is sometimes wounded in the operation of cutting the frænum in children who are tongue-tied, especially when we neglect the practical rule of pointing the scissors downwards and backwards.

[**Varieties of the Lingual Artery.**—The lingual artery at times arises in common with the branch above, *i. e.*, the facial. Sometimes this artery is joined with the superior thyroid. The hyoid branch is sometimes absent, or very small; then we find the hyoid branch of the superior thyroid, larger than usual. The sublingual is sometimes given off from the facial, and passes through the mylo-hyoid muscle. The lingual again at times gives off the submental and ascending palatine.]

The *lingual vein* runs over the hyoglossus, and empties itself into the internal jugular.

I have already mentioned (p. 26) the best place for finding and tying the lingual artery. Of course, the rule laid down is trustworthy, only when the artery runs its normal course. I have known an instance in which a good anatomist failed in an attempt to tie the lingual artery, because the vessel arose from the facial behind the submaxillary gland, and then passed through the mylo-hyoideus to reach the tongue.

The next object of study should be the course and relations of the external carotid artery, and its branches in the neck. In preparing a view of them, you will observe, that nearly all the veins lie in front of their corresponding arteries.

Course and Relations of the External Carotid Artery.—The external carotid arises from the common carotid about

the level of the upper border of the thyroid cartilage. It ascends beneath the hypoglossal nerve, the posterior belly of the digastricus and stylo-hyoideus, enters the parotid gland, and terminates near the neck of the jaw by dividing into the temporal and internal maxillary arteries.

Notice the relative position which the external and internal carotids bear to each other. The external lies at first nearer to the side of the pharynx than the internal. But the external soon changes its position and crosses obliquely in front of the internal to reach the space between the angle of the jaw and the mastoid process. The internal carotid, bear in mind, ascends perpendicularly by the *side of the pharynx* to the base of the skull.

The external is separated from the internal carotid by the stylo-glossus and stylo-pharyngeus muscles, the glosso-pharyngeal nerve, and the stylo-hyoid ligament.

[The external carotid varies in length according to the point of division of the common carotid. It has also been known to arise as an independent branch from the arch.]

The external carotid gives off the following branches:—

- | | |
|--------------------------|------------------------------|
| 1. The superior thyroid. | 5. The posterior auris. |
| 2. The lingual. | 6. The internal maxillary. |
| 3. The facial. | 7. The temporal. |
| 4. The occipital. | 8. The ascending pharyngeal. |

Superior Thyroid Artery.—The superior thyroid is the first branch of the external carotid. It runs beneath the omohyoid, sterno-hyoid, and sterno-thyroid muscles to the upper and front surface of the thyroid body, in which it terminates. Its branches are the four following:—

1. The *hyoid*, a small muscular branch, runs horizontally inwards below the greater cornu of the os-hyoides.

2. The *superior laryngeal* branch, accompanied by the superior laryngeal nerve, runs beneath the thyro-hyoid muscle, perforates the thyro-hyoid membrane, (sometimes the thyroid cartilage,) and supplies the muscles and the mucous membrane of the larynx.

3. The *sterno-mastoid*, a small branch variable as to origin and size, descends over the sheath of the carotid artery to the mastoid muscle.

4. The *crico-thyroid*, an artery of great interest in reference to the operation of laryngotomy, crosses the crico-thyroid membrane, and communicates with a corresponding branch on the opposite side (fig 3. p. 11). One or two small branches pass through the membrane to the interior of the larynx. It is important to know that the crico-thyroid artery often varies in direction and size. In most cases it is small, and runs across the centre of the membrane; we should therefore be least likely to wound it in laryngotomy, by dividing the membrane close to the cricoid cartilage. But it is by no means unfrequent to find this artery of considerable size, taking an oblique or a perpendicular direction in front of the membrane, and finally distributed to one of the lobes of the thyroid body. I have even seen several instances in which the membrane was crossed by the main trunk of the superior thyroid. These facts should establish the practical rule in laryngotomy, not to make an opening into the larynx until we have fairly exposed the parts, and ascertained whether any large artery lies in the way.

Among the many arterial inosculations about the thyroid body are two which deserve notice; the one is formed between the two superior thyroid arteries along the upper border of the isthmus, the other takes place along the back part of the lateral lobe between the superior and inferior thyroid.

[The superior thyroid sometimes arises from the common carotid directly, or from a branch with the lingual. We have seen two thyroid arteries on the same side. It may again be small, and terminate in the mastoid muscle and larynx. The hyoid branch may be absent. The laryngeal may arise from the external or from the common carotid. This branch has been seen passing through a foramen in the thyroid cartilage.]

The *superior thyroid vein* crosses the sheath of the common carotid, and joins the internal jugular.

The superior laryngeal nerve, mentioned as accompanying the superior laryngeal artery, comes from the ganglion of the pneumogastric. It descends behind both carotid arteries, enters the larynx through the thyro-hyoid membrane, and supplies the mucous membrane of the larynx with common sensa-

tion. It is this nerve which presides over the exquisite sensibility of the "rima glottidis."

Before it enters the larynx, the nerve sends the "external laryngeal branch" to supply the crico-thyroid muscle.

Lingual Artery.—The lingual artery and its branches have been described (p. 29).

Facial Artery.—The facial artery and its branches below the jaw have been described (p. 24).

Occipital Artery.—The occipital artery is the fourth branch of the external carotid. It runs backwards along the lower border of the digastricus towards the mastoid process. It then passes under all the muscles inserted into the mastoid process, namely, the sterno-mastoid, the splenius capitis and the trachelo-mastoid. Arrived at the back of the head, the artery runs superficial to the complexus and divides into wide-spreading branches for the supply of the scalp. Observe that in the first part of its course, the occipital artery crosses the internal jugular vein, and is itself crossed by the hypo-glossal nerve. It sends off the three following branches:—

1. The *sterno-mastoid*, which enters the muscle with the nervus accessorius.
2. The *posterior meningeal* ascends along with the internal jugular vein, and enters the cranium to supply the dura mater.
3. The *princeps cervicis*, which we shall see better hereafter, runs down the back of the neck under the complexus, supplies this muscle and the semi-spinalis colli.

The *occipital vein* usually terminates in the internal jugular.

[The occipital artery may arise from the internal carotid or even from the vertebral, and has been seen coming from the pharyngeal ascendens. It may pass over the trachelo-mastoid muscle. The posterior auricular, and pharyngeal, have been seen to take their origin from this vessel.]

Posterior Auricular Artery.—The posterior auricular artery is the fifth branch of the external carotid. It arises above the digastric muscle, and runs, under cover of the parotid gland, to the furrow between the cartilage of the ear and the mastoid process. It supplies the back of the scalp and the cartilage of the ear. Its only named branch is the *stylo-mastoid*, a very constant little artery, which runs through the stylo-mastoid foramen to the tympanum of the ear.

[The posterior auricular may arise from a trunk common to it and the occipital, and may give off the transverse facial.]

Posterior Auricular Nerve.—We find the posterior auricular nerve close to the artery of the same name. It is the first branch of the facial after its exit from the stylo-mastoid foramen. It runs behind the ear and sends motor filaments to the little muscles which move the ear, and to the posterior belly of the occipito-frontalis. This is what we should expect, considering that the facial nerve supplies all the muscles of expression.

Ascending Pharyngeal Artery.—This artery arises near the division of the common carotid. It ascends towards the base of the skull, by the side of the pharynx, the upper part of which it supplies, as well as the Eustachian tube and the tonsils.

[The ascending pharyngeal may come off from the internal carotid, occipital, facial, or superior thyroid; also two, and even three, instead of a single artery, have been met with.]

The examination of the two remaining branches of the external carotid, namely, the internal maxillary and the temporal, must for the present be postponed. Meantime let us notice the cervical plexus and its branches, and then turn our attention to the posterior triangle of the neck.

Cervical Plexus of Nerves.—This plexus is formed by the anterior branches of the four upper cervical nerves. It consists of a series of loop-like communications which take place between these nerves, close to the transverse processes of the four upper cervical vertebræ. The plexus is situated behind the internal jugular vein, and lies in front of the scalenus anticus and the levator anguli scapulæ muscles.

The plexus gives off *superficial* and *deep* branches. The superficial branches are very extensive; these you have already traced, (p. 4.) The *deep* branches are distributed to eleven muscles, as follow :—

Muscles supplied by the Cervical Plexus.—The sternomastoid, the trapezius, the levator anguli scapulæ, the rectus capitis anticus major and minor, the rectus capitis lateralis, the longus colli, the sterno-hyoid, sterno-thyroid, and omo-hyoid, and lastly, the diaphragm.

The cervical plexus communicates with the hypoglossal

nerve, the *nervus accessorius*, and the superior cervical ganglion of the sympathetic. But the dissection of all this requires much time and care, and is of no practical value.

Posterior Triangle of the Neck.—The boundaries of the posterior triangle of the neck are, the sterno-mastoid, the trapezius, and the clavicle. This triangle is divided by the omo-hyoid into two lesser triangles, an upper and a lower.

When the platysma and cervical fascia are cleared away, you will find in the upper triangle (posterior superior) the following muscles proceeding from above (fig. 2, p. 9),—the *splenius capitis*, the *levator anguli scapulæ* and the *scaleni*. You also find the descending branches of the cervical plexus, the *nervus accessorius*, the posterior scapular a. (*transversalis colli*), and its branch the *superficialis colli*, which chiefly supplies the trapezius.

The posterior inferior triangle is by far the more important of the two, because it contains the subclavian vessels and the brachial plexus of nerves. To this triangle let us now address ourselves.

ANATOMY OF THE SUPRA-CLAVICULAR REGION.

Anatomy of the Supra-clavicular Region.—The arm should be placed by the side of the body, that the clavicle and shoulder may not be raised above their natural position. The space we propose to examine is bounded by the clavicle, the outer border of the sterno-mastoid, and the posterior portion of the omo-hyoid muscle. The area of the triangle thus formed will vary in proportion to the obliquity of the omo-hyoid muscle, and the extent to which the sterno-mastoid is attached to the clavicle: the trapezius must also be taken into the account, for in some instances it comes so far forwards as almost to meet the sterno-mastoid. The depth at which the vessels and nerves contained in this space are situated depends not only upon the degree to which the clavicle arches forwards, but varies, as one may ascertain in one's own person, with the elevation and depression of the shoulder.

Immediately beneath the skin covering this region we find the platysma myoides, some cutaneous branches of the cervical plexus, and a layer of fascia which binds down the omohyoid muscle to the clavicle. Beneath this is a deeper layer of fascia, which covers the subclavian vessels and the brachial plexus of nerves, and descends with them beneath the clavicle into the axilla. Between these two layers we meet with more or less fat and areolar tissue, and absorbent glands continuous with those in the axilla. It will be easily understood how a collection of matter, originating in the axilla, may ascend in front of the vessels and point above the clavicle, or, *vice versâ*, matter formed in the neck may travel under the clavicle and point in the axilla.

Near the border of the sterno-mastoid muscle the external jugular vein passes through both layers of the fascia, and terminates in the subclavian: but before its termination it is commonly joined by the supra-scapular, the posterior scapular, and other unnamed veins proceeding from the surrounding muscles; so that we have in this particular situation a *confluence of veins*, which are under any circumstances troublesome, and, when large or distended, exceedingly embarrassing.

The fascia and the glands should be carefully removed. The clavicle should be sawn through the middle, and the sternal half raised with the sterno-mastoid attached to it, so that the bone can be replaced from time to time in order to see its bearings to the parts beneath.

Behind and nearly parallel with the clavicle is the supra-scapular (*transversalis humeri*) artery, a branch of the thyroid axis. A little higher is the *transversalis colli*, or posterior scapular (commonly a branch of the thyroid axis) which crosses the lower part of the neck towards the posterior superior angle of the scapula. But both these arteries, the last particularly, are very irregular in respect to their origin. The scalene muscles descend from the transverse processes of the cervical vertebræ to the first and the second ribs. The phrenic nerve is observed upon the surface of the *scalenus anticus*. The subclavian artery comes into view behind the outer border of the *scalenus anticus*, and continues its course over the surface of the first rib. The subclavian vein also

lies upon the rib, but in front of the scalenus anticus. The large nerves constituting the brachial plexus emerge between the scalene muscles, higher than the subclavian artery.

The scalene muscles, so called from their resemblance to a scalene triangle, extend from the transverse processes of the cervical vertebræ to the first and second ribs. They may be considered as intercostal muscles, since the transverse processes of the cervical vertebræ are but rudimentary ribs. Anatomists describe them as three separate muscles—an anterior, a middle, and a posterior; the anterior and middle are attached to the first rib, the posterior to the second. In plan and purpose these three muscles form but one.

Scalenus Anticus.—The scalenus anticus *arises* from the anterior tubercles of the transverse processes of the third, fourth, fifth, and sixth cervical vertebræ, and is *inserted* by a flat tendon into the tubercle on the *inner* border of the first rib.

Scalenus Medius.—The scalenus medius *arises* from the posterior tubercles of the transverse processes of the six lower cervical vertebræ, and is *inserted* into the first rib behind the scalenus anticus.

Scalenus Posticus.—The scalenus posticus *arises* from the posterior tubercles of the transverse processes of the two or three lower cervical vertebræ, and is *inserted* into the second rib between the tubercle and angle.

The scalene muscles are important agents in raising the thorax, in a deep inspiration. Take a deep breath, and you can easily feel them contracting in your own person. They can bend forwards the cervical portion of the spine, if their lower attachment be the fixed point, as in rising from the recumbent position.

The scalenus anticus is just one of those muscles about which we ought to know well all that lies in front of it, and all that lies behind it. In front of it, then, you see the phrenic nerve, the subclavian vein, the supra-scapular and posterior scapular arteries. Behind it you see the subclavian artery and the five nerves which form the brachial plexus.

Make your finger familiar with the feel of the tubercle on the first rib, to which the scalenus anticus is attached. This tubercle is the best guide to the subclavian artery, for it

guides you to the *outer edge* of the scalenus anticus, where you are to look for the vessel. Is the scalenus anticus entirely concealed from view by the sterno-mastoid or not? This will depend upon the breadth of the clavicular attachment of the sterno-mastoid. As a general rule, it may be said, that the scalene muscle is concealed by the sterno-mastoid, and that consequently, in tying the subclavian artery, it will greatly facilitate the operation, if the clavicular origin of the muscle be partially divided.

Phrenic Nerve.—The phrenic nerve runs perpendicularly upon the scalenus anticus. Trace it upwards, and you will find that it arises from the third, fourth, and fifth cervical nerves; but chiefly from the fourth. It enters the chest between the subclavian artery and vein, close to the inner border of the scalenus, and continues its course between the pericardium and the pleura, in front of the root of the lung, to the diaphragm, which muscle it supplies.

When the spinal cord is injured above the fourth cervical vertebra, the origin of the phrenic is implicated, and therefore the diaphragm, as well as all the other muscles of inspiration, are paralysed. Death is the immediate result.*

COURSE AND RELATIONS OF THE SUBCLAVIAN ARTERIES.


The left subclavian artery differs from the right, not only in its origin but in the relations of the first part of its course. Let us therefore examine the right subclavian first, and then consider the differences which exist between it and the left.

Right Subclavian Artery.—The right subclavian artery is one of the two great branches into which the *arteria innominata* divides behind the sterno-clavicular joint. It runs out-

* The phrenic nerve is frequently joined by a filament from that branch of the brachial plexus which supplies the subclavius muscle. It is important to be aware that cases sometimes occur in which this seemingly insignificant filament is a branch of considerable size, and forms the greater portion of the phrenic itself. I have met with many instances in which this accessory branch was larger than the regular trunk; in all of them it crossed over the

wards behind the scalenus anticus, and then inclines downwards over the first rib, at the outer border of which it takes the name of "axillary." The artery describes a slight curve,

Fig. 6.



of which the greatest convexity is between the scalene muscles. The height to which the arch ascends varies. Gener-

subclavian artery in the third part of its course, and would probably have been injured in the operation of tying this vessel. That such an accident has actually happened is recorded by Bransby Cooper in his surgical lectures. With his usual candor he speaks of having injured this accessory branch of the phrenic in tying the subclavian artery. The patient had incessant spasm of the diaphragm till he died.

ally, it rises higher in women than in men, on the right side than on the left.

To study its relations more precisely, we divide the course of the subclavian artery into three parts:—1. the part which intervenes between the trachea and the scalene muscle: 2. the part which lies behind the scalene muscle: 3. the part which intervenes between the outer border of the scalenus, and the outer border of the first rib.

The *first* portion of the artery is covered by the sternal end of the clavicle, the sterno-mastoid, sterno-hyoid, and sterno-thyroid muscles, and a layer of fascia. It is covered by the confluence of the internal jugular and subclavian, and also by the vertebral veins. It is crossed by the pneumogastric and phrenic nerves, and some cardiac filaments of the sympathetic. Behind the artery are the recurrent branch of the pneumogastric nerve, the sympathetic nerve, the pleura, and the apex of the lung.

In the *second* part of its course, the artery lies between the scalene muscles. It is covered by the clavicular origin of the sterno-mastoid, the cervical fascia, and the scalenus anticus, which separates it from the subclavian vein.

In the *third* part of its course, the artery lies upon the surface of the first rib. Here it is covered only by the platysma, and two layers of cervical fascia: and, what is of much more consequence, it is here crossed by the external jugular and (often) the supra and posterior scapular veins; so that we have a confluence of large veins in front of the artery. The subclavian vein is situated below the artery, but on a plane anterior to it. Above the artery are the trunk nerves of the brachial plexus. One of these nerves (the conjoined fifth and sixth cervical), runs so nearly parallel with the artery, and *on a plane anterior* to it, that many a surgeon has mistaken the nerve for the artery, in the operation of tying it. I once heard a hospital surgeon of great experience say, that he had seen the subclavian tied four times during life, and that in three out of the four cases, the nerve in question had been tied instead of the artery.

Left Subclavian Artery.—The left subclavian is the last of the three great branches which arise from the arch of the aorta. It ascends nearly vertically out of the chest, and then

arches in front of the apex of the lung, so as to reach the inner border of the scalenus anticus, behind which it runs over the first rib.

In the first part of its course, the left subclavian lies deep in the chest near the spine. It is covered on its left side by the pleura. On its right side are the thoracic duct and the œsophagus: the pneumogastric, phrenic, and sympathetic nerves run nearly parallel with it: the left vena innominata crosses in front of it.

When the left subclavian reaches the level of the upper part of the chest, it arches, like the right, over the apex of the lung, and has similar relations; namely, in front, it is covered by the sternal end of the clavicle, the sterno-mastoid, sterno-hyoid, and sterno-thyroid muscles, and by the confluence of the internal jugular and subclavian veins, and also by the vertebral vein: behind it, are the apex of the lung and the pleura.

Behind the scalenus anticus, and on the surface of the first rib, the relations of the left subclavian are similar to those of the right (page 40).

The left subclavian, then, differs from the right only in the first part of its course. Now what are these differences?

1. The left subclavian comes direct from the arch of the aorta, and is therefore longer, deeper in the chest, and more vertical than the right, which comes from the arteria innominata.

2. The left subclavian is in close relation with the œsophagus and the thoracic duct: the right is not.

3. The left subclavian is crossed by the left vena innominata.

4. The left subclavian has the phrenic, pneumogastric, and sympathetic nerves nearly parallel with it; on the right side, these nerves cross the artery at right angles.

5. The left subclavian is not embraced by the recurrent laryngeal nerve, like the right subclavian.

[The origin of the right subclavian varies according to bifurcation of the innominate. The right subclavian arises at times directly from the aortic arch, being the first, second, or third in order; the most frequent anomaly is to find it the last of these branches, in which case it springs from the upper or posterior part of the arch; it also has been known to come from the descending aorta. Its usual course is to cross

obliquely to the right side, in front of the vertebra, and behind the trachea; very rarely has it been known to pass between the œsophagus and trachea. The left has been seen to spring from the ductus arteriosus. We have also noticed this vessel running in front, instead of behind, the scalenus. The vein may run with the artery, behind the muscle. The first portion of the left subclavian was tied by J. K. Rodgers, at New York Hospital, in 1845. The first successful case of ligature of the artery, in the third portion of its course, was that by Dr. W. Post, of New York, in 1817.]

The thoracic duct bears an important relation to the left subclavian artery. It ascends from the chest between the œsophagus and the artery, forms an arch with the convexity upwards behind the internal jugular vein, and terminates in the subclavian vein at its junction with the jugular. The duct is so thin and transparent that it easily escapes observation: you will most readily find it by searching with the handle of the scalpel in the loose tissue on the inner edge of the scalenus anticus, in front of the vertebral vein.

Before we trace the branches of the subclavian artery, let us consider some points relating to the operation of tying it.

To tie the artery in the first part of its course, namely, on the inner edge of the scalenus anticus muscle, is an operation of great difficulty and danger, even with the parts in a normal position. The great depth at which the artery is placed, the size and close proximity of its numerous branches, the large veins by which it is covered, its connexion with the pneumogastric, recurrent laryngeal, phrenic, and sympathetic nerves, and above all its close contiguity with the pleura, form a combination of circumstances so formidable that one cannot be surprised the operation has never been performed with a favorable result.

In the second part of its course, namely, between the scalene muscles, the artery is more accessible. It would be necessary to divide the clavicular origin of the sterno-mastoid, the cervical fascia, and the scalenus anticus muscle, in order to reach the vessel; the phrenic nerve and the subclavian vein would be the chief objects exposed to injury. This operation was performed first and with success by Dupuytren in the year 1819. More recently it has been performed by Dr. Warren,

of Boston. The patient recovered, though the pleura was wounded.*

But in the last part of its course, that is, on the outer side of the scalenus, the artery may be tied with comparative facility. The incision should be made from three to four inches in length, parallel with the upper border of the clavicle. We divide the platysma, some of the supra-clavicular nerves, and the cervical fascia. The external jugular vein must be drawn to the outer side, or divided and tied at both ends. With the finger and the handle of the scalpel we then make our way down to the outer edge of the scalenus anticus muscle, behind which the artery will be found lying upon the first rib. Remember the tubercle on the inner edge of the rib which indicates the insertion of the scalenus: this tubercle is the best guide to the artery. It will be necessary to divide a layer of fascia which immediately covers the vessel before the needle can be introduced around it. Mr. Ramsden of St. Bartholomew's Hospital was the first who tied the subclavian in the third part of its course, in the year 1809; since that time the operation has been repeatedly performed and often with a favorable result.

In the hands of a surgeon possessed of a practical knowledge of anatomy the operation is easy, provided all circumstances be favorable; but circumstances are often very unfavorable. It often happens that the aneurismal or other tumor, on account of which the operation is performed, raises the clavicle beyond its natural level, and so disturbs the parts, that to expose the artery and place a ligature around it becomes exceedingly difficult. Under such circumstances one cannot be surprised that even distinguished anatomists have committed mistakes. Sir Astley Cooper† failed in one instance. Dupuytren perforated the artery with the point of the needle, and included one of the nerves in the ligature: fatal hæmorrhage was the result.‡ I was present at an operation in which the large nerve (a branch of the brachial plexus) which runs parallel with and on a plane anterior to the artery was mis-

* Med. Chirurg. Trans., vol. xxix. p. 25.

† London Medical Review, vol. ii. p. 300.

‡ Edinburgh Med. and Surg. Journal, vol. xvi. 1820.

taken for it and tied; the surgeon being deceived by the pulsation communicated to the nerve.

Branches of the Subclavian Artery.—The branches of the subclavian extend so far, that in the present dissection we can trace them only for a short distance. They are four in number:—

1. The vertebral.
2. The thyroid axis, which gives off inferior thyroid, supra-scapular, posterior-scapular, and cervicalis ascendens.
3. The internal mammary.
4. The superior intercostal, which gives off the deep cervical.

The rule is, that these branches arise from the subclavian in the first part of its course. The most frequent variety is, that the posterior scapular (*transversalis colli*) arises from the subclavian in the third part of its course.*

Vertebral Artery.—This, the first and largest branch, arises from the upper part of the subclavian. For a short distance it lies in the interval between the *scalenus anticus* and the *longus colli*. Here it enters the foramen in the transverse process of the sixth cervical vertebra, and ascends through the foramina in the transverse processes of the succeeding vertebræ. In the interval between the axis and the atlas, recollect that the artery makes a sigmoid curve, in order that it may not be stretched in the rotation of the head. After passing through the foramen of the atlas, the vessel curves backwards along the groove upon its posterior arch, enters the skull through the foramen magnum, and unites with its fellow near the lower border of the "*pons Varolii*," to form the basilar artery.

The vertebral artery is accompanied by slender nerves from the inferior cervical ganglion of the sympathetic. These nerves communicate with the spinal nerves forming the brachial plexus.

Being intended for the brain, the vertebral gives off no branches in the neck, except a few small ones to the deeply-

* With reference to the origin of the posterior scapular (*transversalis colli*) artery, I made special observations during the winter session of 1858-59. I found that this artery was given off most frequently, not by the thyroid axis, but by the subclavian, in the *third* part of its course. Under these circumstances the *superficialis colli a.* generally came from the thyroid axis.

seated muscles; it furnishes, however, to the spinal cord and its membranes small arteries which pass through the inter-vertebral foramina.

[The right vertebral at times comes from the common carotid, while the left may be a branch from the aorta. They have been found to arise from two roots; these unite before, or soon after, entering the vertebral canal. Instead of entering the transverse process of the sixth vertebra, it may pass through the foramen of the fifth, fourth, third, or even the second vertebra. Their respective size often differs, the left being the larger.]

The *vertebral vein* does not come out of the skull with the vertebral artery, but is formed by small branches from the muscles near the foramen magnum. It descends in front of the artery through the foramina in the transverse processes, and joins the brachio-cephalic vein. It receives the veins from the cervical portion of the spinal cord. In some subjects it communicates with the lateral sinus by a branch through the posterior condyloid foramen.

Thyroid Axis.—The *thyroid axis* arises from the subclavian near the inner edge of the scalenus anticus, and after a course of a quarter of an inch divides into four branches which take different directions; namely the inferior thyroid, the posterior scapular, the supra-scapular, and the ascending cervical.

[The thyroid axis may be absent; the vessels coming off separately from the subclavian. The axis has in two cases been known to come off on the outer side of the scalenus muscle.]

The *inferior thyroid* artery ascends tortuously behind the sheath of the common carotid artery and the sympathetic nerve, to the deep surface of the thyroid body, in which it communicates freely with the superior thyroid. It gives small branches to the trachea, the oesophagus, and the larynx.

The *ascending cervical* often arises from the inferior thyroid. It ascends in the interval between the scalenus anticus and the rectus capitis anticus major, and terminates in small branches, some of which supply these muscles; others enter the inter-vertebral foramina and supply the spinal cord.

The *supra-scapular* artery (*transversalis humeri*) runs outwards over the scalenus anticus, then directly *beneath* and

parallel with the clavicle towards the notch of the scapula. Here it is joined by the supra-scapular nerve, and divides into branches, some of which ramify above, others below, the spine of this bone. It inosculates freely in the infra-spinous fossa with the "dorsalis scapulæ," a branch of the subscapular artery.

The *posterior scapular* (transversalis colli) artery, though generally described as a branch of the thyroid axis, more frequently arises from the subclavian in the last part of its course. It runs tortuously across the side of the neck (higher than the supra-scapular), over the scalene muscles and the great nerves of the brachial plexus (sometimes between them), and disappears beneath the trapezius and the levator anguli scapulæ to reach the posterior angle of the scapula. It then runs beneath the rhomboid muscles, in which it is lost. In the space between the sterno-mastoid and trapezius, the posterior scapular gives off the "*superficialis colli*." This vessel proceeds tortuously across the posterior triangle of the neck to the under surface of the trapezius, which, with the levator anguli scapulæ, it principally supplies.

You will very often find that the *superficialis colli* comes direct from the thyroid axis, while the posterior scapular comes from the last part of the subclavian.

The veins corresponding to the supra-scapular and posterior scapular arteries terminate in the external jugular, sometimes in the subclavian. The inferior thyroid vein crosses in front of the common carotid artery, and joins the internal jugular.

Internal Mammary.—This artery arises from the subclavian opposite to the thyroid axis. It enters the chest behind the subclavian vein, and descends behind the cartilages of the ribs, about half an inch from the sternum. Its further progress will be examined in the dissection of the chest. The corresponding vein most frequently terminates in the vena innominata.

Superior Intercostal.—This artery is given off by the subclavian behind the scalenus anticus, so that you must divide the muscle to see it. It descends into the chest, passing over the necks of the first and second ribs, and furnishes the arteries of the two upper intercostal spaces. It usually inos-

culates with the first intercostal branch of the aorta. The corresponding vein terminates on the right side in the vena-azygos; on the left in the vena innominata.

[The superior intercostal is not unfrequently absent.]

Deep Cervical.—This artery arises from the superior intercostal, seldom direct from the subclavian. It goes to the back of the neck between the first rib and the transverse process of the last cervical vertebra, and ascends between the complexus and the semispinalis colli, both of which it supplies. It sometimes inosculates with the “princeps cervicis,” a branch of the occipital a.

If you wish to test your knowledge of the branches of the subclavian artery, reflect upon the answer to the following question: If

the artery were tied in the *first* part of its course before it gives off any branches, how would the arm be supplied with blood? The answer is, by five collateral channels, as follow: 1. By the communications between the superior and inferior thyroid: 2. By the communications between the two vertebral: 3. By the communications between the internal mammary on the one hand, and the intercostals and the epigastric on the other: 4. By the communications between the thoracic branches of the axillary, and the intercostal branches of the aorta: 5. By the communications between the superior inter-

Fig. 7.




DIAGRAM TO SHOW THE INOSCULATIONS OF THE SUBCLAVIAN ARTERY.

costal and the aortic intercostal. These inosculations are shown in the diagram, page 47.

Again, if the subclavian were tied in the *third* part of its course, the circulation would be carried on by the communications: 1. Between the supra-scapular and dorsalis scapulæ (branch of subscapular): 2. Between the posterior scapular and the subscapular: 3. Between the long and short thoracic on the one hand, and the internal mammary and aortic intercostals on the other.

Subclavian Vein.—The subclavian vein does not form an arch like the artery, but proceeds in a nearly straight line over the first rib to join the internal jugular. Throughout its whole course the vein is situated on a plane anterior to and a little lower than the artery, from which it is separated by the scalenus anticus, the phrenic and pneumogastric nerves. It has a valve just before its junction with the internal jugular. It receives the external jugular, anterior jugular, supra-scapular, and posterior scapular veins.

Brachial Plexus of Nerves.—The large nerves forming the plexus which supplies the upper extremity are the anterior divisions of the four lower cervical nerves and the first dorsal. Emerging from the inter-vertebral foramina they appear between the anterior and middle scalene muscles, and pass with the subclavian artery into the axilla. To this bundle of nerves the name "plexus" is given, on account of their mutual communications. The plexus at its root is wide, and situated higher than the subclavian artery, and nearly on the same plane; but as the plexus descends beneath the clavicle its component nerves converge, and in the axilla, completely surround the artery.

Remember that the plexus is crossed superficially by the omo-hyoid muscle, and by the supra-scapular and posterior scapular arteries.

The arrangement of the nerves in the formation of the plexus is usually thus:—The fifth and sixth cervical unite to form a single cord; the eighth cervical and the first dorsal unite to form another; the seventh cervical runs for some distance alone and then divides so as to unite with the other two cords.

[We think the arrangement is more frequently for the fifth,

sixth, and seventh to unite, and form a single cord, (outer cord) while the eighth and first dorsal unite to make the other cord (inner cord). These cords then each give off a branch, which uniting, give rise to the third, or posterior cord.]

The plexus gives off above the clavicle the following nerves:—

a. *Nerve to the subclavius muscle*.—This proceeds from the fifth and sixth cervical, and crosses the subclavian artery in the third part of its course. It frequently sends a filament which passes in front of the subclavian vein to join the phrenic.

b. *Nerves to the scalene and the longus colli muscles*.

c. *Nerve to the rhomboid muscles*.—This arises from the fifth and sixth cervical nerves, and accompanies the posterior scapular artery beneath the levator anguli scapulæ, which, as well as the rhomboid muscles, it supplies.

d. The *supra-scapular* nerve arises from the fifth and sixth cervical, runs to the notch in the scapula, where it meets with the corresponding artery, and terminates in the supra-spinatus and infra-spinatus.

e. The *nerve* (called *external respiratory* by Sir C. Bell) to the *serratus magnus*, arises from the fifth and sixth cervical, appears between the middle and posterior scalene muscles, then descends behind the plexus and the subclavian vessels to the outer surface of the serratus magnus, to which it is exclusively distributed.

f. *Nerves* (called *anterior thoracic*) to the *pectoral muscles*.—Generally two from the fifth and sixth cervical; they pass beneath the clavicle in front of the axillary artery.

g. A branch forming one of the roots of the phrenic, arises from the fifth cervical nerve.

It only remains to be observed that the upper chord of the brachial plexus receives a branch from the lower chord of the cervical plexus, and that each of its component nerves communicate by slender filaments with the sympathetic system.

Below the clavicle the plexus divides into branches for the supply of the arm; namely, the subscapular (three in number, to the subscapularis, the latissimus dorsi, and teres major), the circumflex (to the deltoid), the median, the radial, the ulnar, the external cutaneous, the internal cutaneous, and the lesser internal cutaneous (nerve of Wrisberg).

DISSECTION OF THE FACE.

Much practice is required to make a good dissection of the face. The muscles of expression are numerous and complicated; they are interwoven with the subcutaneous tissue and closely united to the skin; their fibres are often pale and indistinct. The face is amply supplied with motor and sensitive nerves, of which the ramifications extend far and wide. Therefore you must not be discouraged, if in a first attempt you fail to make a satisfactory display of the parts we propose to examine.

Make the cheeks tense by filling the mouth with horse-hair, and stitch the lips together. Make an incision down the medial line of the face, and another from the condyle of the jaw to the angle of the mouth.

Recollect that the nerve which supplies all the muscles of the face is the "*portio dura*," or facial division of the seventh cerebral nerve. It emerges from the stylo-mastoid foramen, and divides into branches, which pass through the parotid gland, forming a plexus termed the "*pes anserinus*."

The sensitive nerves of the face are supplied by the three divisions of the fifth cerebral nerve: namely, the supra-orbital branch, the infra-orbital, and the mental; each being named after the foramen in the bone through which it emerges. No other nerve takes any share in conferring sensation upon the face except the auriculo-parotidean branch of the cervical plexus (p. 4), which supplies the skin covering the parotid gland and part of the cheek.

The skin covering the cartilages of the nose is supplied by a small nerve called the *naso-lobular*. It is the terminal branch of the nasal division of the ophthalmic nerve. It appears between the bone and the cartilage.

Musculus Risorius (Santorini).—In the dissection of the neck we saw that some of the fibres of the platysma pass over the angle of the jaw and are prolonged as far as the angle of the mouth, where they intermingle with the orbicularis oris and other muscles in this situation. This facial part of the platysma is termed "*musculus risorius*." It produces the smile, not of good humor, but of derision.

It is convenient to arrange the muscles of the face under three heads; appertaining respectively to the mouth, the nose, the eyebrows and lids. It matters little which group be first selected. Begin with those of the mouth.

The muscles of the mouth are arranged thus: there is an orbicular or sphincter muscle surrounding the lips; from this, as from a common centre, muscles diverge and are fixed into the surrounding bones. They are named elevators, depressors, &c., according to their respective action.

Orbicularis Oris.—This muscle, nearly an inch in breadth, surrounds the mouth. Its size and thickness in different individuals, produce the variety in the prominence of the lips. Observe that its fibres do not surround the mouth in one unbroken series, but that those of the upper and lower lip decussate at the angles of the mouth, and intermingle with the fibres of the buccinator and other muscles which converge from different parts of the face. The cutaneous surface of the muscle is intimately connected with the lips and the surrounding skin; the deep surface is separated from the mucous membrane by the labial glands and the coronary vessels.

The “orbicularis” is the antagonist of all the muscles which move the lips. Upon a nice balance of their opposite actions depends the play of the mouth with which every one is familiar.*

Depressor Anguli Oris.—This muscle *arises* from the oblique line of the lower jaw below the foramen mentale, and is *inserted* into the angle of the mouth, intermingling with the zygomatic muscles.

It is an important muscle in the expression of sorrowful emotions. We see its action remarkably well when children cry.

Depressor Labii Inferioris, or Quadratus Menti.—This muscle *arises* from the oblique line of the lower jaw, and is *inserted* into the lower lip. It covers the vessels and nerves which come through the foramen mentale. It must therefore be divided in the operation of cutting through the nerve in cases of neuralgia.

* In strong muscular lips the upper part of the orbicularis sends a small subcutaneous slip of muscle from each side along the septum nasi nearly to the apex. The interval between the two slips corresponds to the furrow which leads from the nose to the lip. This is the “naso-labialis” or “depressor septi narium” of some anatomists, *e. g.* Haller, Albinus.

Levator Menti.—This muscle *arises* from the lower jaw below the incisor teeth, and is inserted into the skin of the chin. To see it properly you should evert the lower lip and remove the mucous membrane on either side of the frænum. There are two of them, one for each side. Their action is well shown when we shave.

Zygomaticus Major and Minor.—The Z. major *arises* from the outer surface of the malar bone, and is *inserted* into the angle of the mouth, joining the depressor anguli oris. The Z. minor *arises* from the outer surface of the malar bone, and is *inserted* into the angle of the mouth, joining the levator labii superioris. The action of the zygomatici is seen in laughing.

[The zygomaticus minor is frequently absent, and, when present, may terminate before reaching the angle of the mouth.]

Before we examine the orbicularis palpebrarum we must notice the tendo oculi. To make the tendon more apparent the tarsal cartilages should be drawn outwards.

Tendo Oculi.—This tendon is about two lines* in length, and is readily felt beneath the skin at the inner angle of the eye by drawing the eyelids outwards. It is fixed to the nasal process of the superior maxillary bone, passes outwards, and divides into two portions, one of which is attached to the upper, the other to the lower tarsal cartilage. The tendon crosses the lachrymal sac a little above the centre, and furnishes a tendinous expansion which covers the sac and is attached to the margin of the bony groove. To see all this expansion we must reflect that portion of the orbicularis palpebrarum which covers the sac.

In puncturing the lachrymal sac the knife is introduced below the tendon, in a direction downwards, outwards, and a little backwards. We have to divide the skin, a few fibres of the orbicularis, and the fibrous expansion from the tendo oculi. The angular artery and vein are situated on the inner side of the incision.

Orbicularis Oculi.—This muscle *arises* from the tendo oculi, and from the inner margin of the orbit, for some distance *above* and *below* this tendon. It is *inserted* into the skin of the brow, temple, and cheek.

* A line is the twelfth part of an inch.

This broad sphincter muscle surrounds the margin of the orbit and eyelids. Its circumferential fibres take a wide sweep, and mingle, on the forehead, with the occipito frontalis, on the cheek, with the zygomaticus minor, and the levator labii superioris.

The fibres which belong to the eyelids (*orbicularis palpebrarum*), are thin and pale. They arise from the tendo oculi, and form, over each eyelid, a series of elliptical curves which meet at the external canthus of the lids. The degree of their curvature becomes less as they approach the margin of the lids, so that some fibres proceed close to the lashes. This was first pointed out by Riolanus,* and described as "*musculus ciliaris*."†

You will observe that no fat is ever found on the eyelids: nothing intervenes between the skin and the muscle but a quantity of loose areolar tissue, that there may be no impediment to the free play of the lids.

The orbicular muscle not only shuts the eye but protects it. For instance, suppose the eye to be threatened by a blow, the muscle suddenly contracts, presses the eye back into the orbit, and contracts the skin of the brow and cheek so as to form a soft cushion in front of the orbit. The cushion itself may be severely bruised, as is seen in a "black eye;" but, how rarely the globe itself is injured! When the eye is simply closed, as in winking, the palpebral portion of the muscle contracts. Observe this movement attentively, and you will see that the lids are drawn slightly inwards as well as closed. Now this inward motion is to direct the tears towards the inner angle of the eyelids, where they are absorbed by the lachrymal ducts.

Since the orbicular muscle is supplied by the facial nerve, it is affected in facial palsy, and the patient cannot shut the eye.

Tensor Tarsi.—This muscle has been particularly described by Horner,‡ an American anatomist. It should be considered not as a distinct muscle, but as a portion of the orbicularis.

* *Anthropologia*, lib. v. cap. 10.

† Strictly speaking the ciliary muscle arises from the two little divisions of the tendo oculi, and is inserted, at the external canthus, into the fibrous tissue which unites the two tarsal cartilages

‡ *Philadelphia Journal*, Nov. 1824. But this muscle was accurately described by Rosenmüller in his *Handbuch der Anatomie*. Leipzig, 1819.

The best way to see Horner's muscle, is to cut perpendicularly through the middle of the upper and lower lids, and to evert the inner halves towards the nose. After removing the mucous membrane you will see this little muscle arising from the ridge on the lachrymal bone. It passes outwards and divides into two portions, which are inserted into the upper and lower tarsal cartilages, close to the orifices of the lachrymal ducts. It is probable that the tensor tarsi draws backwards the open mouths of these ducts, so that they may absorb the tears at the inner angle of the eye.

Corrugator Supercilii.—This *arises* from the superciliary ridge of the frontal bone and is *inserted* into the under surface of the orbicularis oculi. It lies concealed beneath the orbicularis, and is the proper muscle of frowning.

Pyramidalis Nasi.—This is situated on the bridge of the nose, on each side of the mesial line, and is regarded as a prolongation of the occipito-frontalis. It mingles with the fibres of the triangularis nasi, and its action is to elevate the skin of the nose; for instance, in the expression of surprise.

The present being a fitting opportunity to examine the structure of the eyelids, let us postpone for the present the dissection of the remaining muscles of the face.

STRUCTURE OF THE EYELIDS.—The eyelids are composed of different tissues arranged in successive strata one beneath the other. There are—1, the skin; 2, the orbicularis palpebrarum; 3, the expanded tendon of the levator palpebræ (in the upper lid only); 4, the tarsal cartilage; 5, Meibomian glands; 6, mucous membrane. These structures are separated by areolar tissue, which for very good reasons never contains fat.

The *skin* of the eyelids is remarkably smooth and delicate. It is abundantly supplied with sensitive nerves by branches of the fifth pair, namely, by the supra-orbital, supra-trochlear, infra-trochlear, lachrymal, and infra-orbital nerves.

The *orbicularis palpebrarum* has been already described (p. 53). It is supplied with nerves by the facial.

The *levator palpebræ* arises from the back of the orbit, gradually becomes broader, and terminates in a thin aponeurosis, which unites with the broad tarsal ligament, and is lost on the outer surface of the tarsal cartilage.


Tarsal Cartilages.—These are plates of fibro-cartilage

which support and give figure to the eyelids. There is one for each lid, and they are connected at the angles (commisures or canthi) of the lids through the medium of fibrous tissue. One can best examine them by everting the lids. Each cartilage resembles its lid in form. The upper is the larger of the two, is broad in the middle, and gradually becomes narrower at either end. The lower is nearly of uniform breadth throughout. Both are thicker on the nasal than the temporal side. They are connected to the margin of the orbit, and maintained in position by the *tendo oculi*, and by what are called the *broad tarsal ligaments*: these ligaments are continuations from the periosteum of the orbit to the tarsal cartilages. There are two of them, termed upper and lower, and proceeding to each cartilage respectively. When an abscess forms, which is not uncommon after erysipelas, in the areolar tissue of the lids, these ligaments prevent the matter from making its way into the orbit.

The ciliary margin is the thickest part of the tarsal cartilages. It is generally stated that the inner edge of each is sloped or beveled off; and that, consequently, when the lids are closed, there is formed, with the globe of the eye, a triangular channel. This channel is said to conduct the tears to the puncta lacrymalia. According to my observation, this triangular channel does not exist: for when the lids are closed, their margins are in such accurate apposition, that not the slightest interstice can be discovered between them.

The *puncta lacrymalia* are two pin-hole apertures, easily discovered at the inner angles of the lids. They are the orifices of the canals which convey the tears to the lachrymal sac. Observe that their orifices are directed backwards.

In the introduction of probes for the purpose of opening the contracted puncta, or of slitting up the lachrymal duct, it is necessary to know the exact direction of these tubes. (*See diagram.*) By passing a bristle into one of them, we find that it does not run straight from the punctum to the sac, but that



it proceeds for a short distance perpendicularly, and then, dilating into a small pouch, makes a sharp bend inwards to the lachrymal sac. In the majority of cases, the tubes open into the sac by a common orifice. When, from any cause, the tears are secreted in greater quantity than usual, they overflow and trickle down the cheek.

The *eyelashes* (cilia) are planted in rows along the edge of the tarsal cartilages. The bulbs of the lashes are situated between the tarsal cartilage and the fibres of the ciliary muscle. They are supplied with blood by the palpebral branches of the ophthalmic which run parallel and close to the free borders of the lids under the ciliary muscle.

Meibomian Glands.—These sebaceous glands, so called after the anatomist* who first described them, are situated on the under surface of each of the tarsal cartilages. In the upper lid there are from thirty to forty of them, not so many in the lower. On everting the lid they are seen running in longitudinal rows in grooves of the cartilage. Examined with the microscope, one sees that they consist of a central tube, round the sides of which are a number of openings leading to short cæcal dilatations. The orifices of these glands are situated on the free margin of the lid behind the lashes. Their use is to secrete an unctuous substance, which prevents the lids from sticking together.

Caruncula Lacrymalis.—This name is given to a small reddish body situated at the inner corner of the eye. It is composed of an aggregation of sebaceous glands covered by mucous membrane. In some instances minute hairs grow upon it. What is the use of this caruncula? Probably to support the inner junction of the eyelids. When the caruncle is diminished in size by disease, the puncta lacrymalia become displaced, and the tears are apt to run down the cheek.

The conjunctival coat of the eyelid will be described with the anatomy of the eye. We merely observe at present, that it is more vascular than the conjunctival coat of the eye, and that it presents a number of minute papillæ, which, when enlarged and aggregated by inflammation, give rise to the disease called "granular lids."

* H. Meibom, De vasis palpebrarum novis, Helmstedt, 1666.

Such, in outline, is the structure of the eyelids. Their use is best described by Socrates, who, in answer to the question whether animals were made by chance or design, replies:—"Think you not that it looks like the work of *prescience*, because the sight is delicate, to have guarded it with eyelids, which open when we want to see, but shut when we go to sleep; to have fenced those lids with eyelashes, which, like a sieve, strain the dusty wind, and hinder it from hurting the eyes; and over the eyes to have placed eyebrows, as eaves, to carry off the sweat of the brow from disturbing the sight?"*

After this digression, let us proceed with the dissection of the muscles of the face, and raise the lower circumference of the orbicularis oculi, in order to see the origin of the elevators of the upper lip and the nose.

Levator Labii Superioris Alæque Nasi.—This *arises* from the superior maxillary bone near its orbital margin, and is *inserted* into the side of the nose and upper lip. It acts chiefly in expressing the smile of derision, and the scornful affections of the mind. Its habitual use occasions corresponding permanent folds in the skin, which are to a certain extent, indicative of the feelings and passions.

Levator Labii Superioris.—This *arises* from the margin of the orbit above the infra-orbital foramen, and is *inserted* into the upper lip. It is nearly an inch in breadth at its origin, and it covers the infra-orbital vessels and nerves.

Levator Anguli Oris.—This *arises* from the canine fossa of the superior maxillary bone, below the infra-orbital foramen, and is *inserted* into the angle of the mouth.

This muscle lies beneath, and is covered by the levator labii superioris.

Now examine two muscles belonging to the nose—the triangularis or compressor nasi, and the depressor labii superioris.

Compressor Nasi.—This *arises* from the superior maxillary bone, near the ala of the nose, and is *inserted* into the dorsum

* Xenophon's Memorabilia, b. 1. c. iv.: Οὐ δοκεῖ σοι καὶ τόδε προνοίας ἔργον εἰκέναι, τὸ, ἐπεὶ ἀσθενὴς μὲν ἐστὶν ἡ ὄψις, βλεφάροις αὐτὴν θυρῶσαι, αἱ, ὅταν μὲν αὐτῇ χρῆσθαι τι δέη, ἀναπετάννυνται, ἐν δὲ τῷ ὕπνῳ συγκλείεται; ὥς δ' ἂν μηδὲ ἀνεμοὶ βλάπτωσιν, ἡθμὸν βλεφαρίδας ἐμφύσαι· ὀφρύσι τε ἀπογείσῃαι τὰ ὑπὲρ τῶν ὀμμάτων ὥς μὴδ' ὁ ἐκ τῆς κεφαλῇ; ἰδρῶς κακουργῇ.

of the nose with its fellow of the opposite side. The origin of this muscle is concealed by the levator labii superioris alæque nasi.

Depressor Labii Superioris.—This *arises* from the superior maxillary bone, above the second incisor tooth, and is *inserted* into the septum and ala of the nose. It is situated between the mucous membrane and the muscular structure of the upper lip. To expose it, the upper lip must be everted and the mucous membrane removed.

Besides the muscles above described, we find, in connection with the cartilages of the ala of the nose, pale muscular fibres, in which no definite arrangement can be traced. They constitute the "*dilatator narium anterior*" and the "*dilatator narium posterior*" of some anatomists. By acting upon the cartilages of the nose, these little muscles contribute in some degree to express the condition of the mind. Some of them dilate the nostrils; for instance in dyspnœa: others contract them, as in smelling, when we sniff up the air towards the olfactory nerve. Though apparently feeble, they are strong enough to answer their purpose, from the facility with which the cartilages of the nose are moved upon each other.

Buccinator.—The buccinator *arises* from the alveolar border of upper and lower jaw, near the last two molars, and from the pterygo-maxillary ligament. It is *inserted* into the angle of the mouth and the muscular structure of the lips.

The buccinator is the great muscle of the cheek. It forms, with the superior constrictor of the pharynx, a continuous muscular wall for the side of the mouth and pharynx. The bond of connection between the buccinator and the superior constrictor, is the "so called" pterygo-maxillary ligament. Now this ligament (*see diagram*), extends from the hamular process vertically to the lower jaw near the last molar. It is wrong to call it a ligament, for it is simply a fibrous intersection between the two muscles.


The duct of the parotid gland passes obliquely through the buccinator into the mouth, near the second molar of the upper jaw.

The chief use of the buccinator is to keep the food between the teeth during mastication. It can also widen the mouth. Its power of expelling air from the mouth, as in whistling

or playing on a wind instrument, has given rise to its peculiar name. It is supplied by the facial nerve, and is, therefore, affected in facial paralysis.

Buccal Fascia.—The buccinator muscle is covered by a

Fig. 9.



MUSCLES OF THE PHARYNX.

layer of fibrous tissue, which adheres closely to its surface, and is attached to the alveolar border of the upper and lower jaw. This structure is thin over the anterior part of the muscle, but more dense behind, where it is continuous with the aponeurosis of the pharynx. It is called the "bucco-pharyngeal" fascia, since it supports and strengthens the muscular walls of these cavities. In consequence of the resistance of

this fascia abscesses do not readily burst into the mouth or the pharynx.

Molar Glands.—The molar glands, not more than three or four in number, situated immediately outside the posterior part of the buccinator, are about the size of a split pea, and their secretion is conveyed to the mouth by separate ducts near the last molar teeth.

Between the buccinator and the masseter there is, in almost all subjects, an accumulation of fat. It is found, beneath the zygoma especially, in large round masses, and may be turned out with the handle of the scalpel. It fills up the zygomatic fossa, and being soft and elastic, does not present the least obstacle to the free movements of the jaw. Its absorption in emaciated individuals occasions the sinking of the cheek.

Facial Artery.—The facial (external maxillary) artery is the third branch of the external carotid. It runs tortuously beneath the hypoglossal nerve, the posterior belly of the digastricus, and the stylo-hyoideus, next *through* the substance of the submaxillary gland, and mounts over the base of the jaw at the *anterior edge of the masseter* muscle. Up to this point we traced it in the dissection of the neck (page 24). It now ascends tortuously near the corner of the mouth and the ala of the nose, towards the inner angle of the eye, where, much diminished in size, it inosculates with the terminal branch of the ophthalmic, a branch of the internal carotid. In the first part of its course on the face, the facial artery is covered only by the platysma; above the corner of the mouth it is crossed by a few fibres of the orbicularis oris and the zygomatici; still higher it is covered by some of the fibres of the elevators of the upper lip and nose. It lies successively upon the buccinator, levator anguli oris, and levator labii superioris muscles. In its course along the face it gives off the following branches:—

The *inferior coronary* artery comes off near the angle of the mouth, runs tortuously along the lower lip, between the mucous membrane and the orbicularis muscle, and inosculates with its fellow.

The *superior coronary* proceeds along the upper lip close to the mucous membrane, and inosculates with its fellow; thus a complete arterial circle is formed round the mouth. These

arteries can be felt in one's own person pulsating on the inner side of the lip, not far from the free border. From this circle numerous branches pass off to the papillæ of the lips, and the labial glands. The superior coronary gives off a small branch (artery of the septum) which ascends along the septum to the apex of the nose.

The *lateral artery of the nose*, a branch of considerable size, arises opposite the ala nasi, and ramifies upon the external surface of the nose.

The *angular artery*, which may be regarded as the termination of the facial, inosculates on the inner side of the tendo oculi with the ophthalmic.

The facial artery supplies numerous branches to the muscles of the face, and inosculates with the transversalis faciei, infra-orbital, and mental arteries.

The facial artery and its branches are surrounded by a minute plexus of nerves (*nervi molles*), scarcely visible to the naked eye. They are derived from the superior cervical ganglion of the sympathetic, and exert a powerful influence over the contraction and dilatation of the capillary vessels, and thus occasion those sudden changes in the countenance indicative of certain mental emotions: *e. g.*, blushing or sudden paleness.*

The *facial vein* does not run with the artery, but takes a straight course from the inner angle of the eye to the anterior edge of the masseter. In this course it descends beneath the zygomatic muscles, over the termination of the parotid duct, and at the anterior border of the masseter passes over the jaw, behind the facial artery, and joins the internal jugular vein.

The facial vein is a continuation of the frontal, which descends over the forehead, and, after receiving the supra-orbital, takes the name of angular at the corner of the eye. It communicates with the ophthalmic vein, receives the veins of the eyelids, the external parts of the nose, the coronary veins, and others from the muscles of the face. Near the angle of the mouth it is often increased in size by a large vein which

* M. Bernard and Brown Séquard have proved by experiment that if the branches of the sympathetic, which accompany the facial artery, be divided, the capillary vessels of the face, being deprived of their contractile power, become immediately distended with blood, and the temperature of the face is raised.

comes from a venous plexus deeply seated behind the superior maxillary bone.

Arteria Transversalis Faciei.—This artery arises from the external carotid, or the temporal in the substance of the parotid gland. It runs forwards across the masseter above the parotid duct, and is distributed to the *glandula socia parotidis*, but chiefly to the masseter itself.

The transverse facial artery is seldom of large size, except when it supplies those parts which usually receive blood from the facial. I have seen it as large as a goose-quill, furnishing the coronary and the nasal arteries; the facial itself not being larger than a sewing thread.

Now direct your attention to the parotid gland; its boundaries, its deep relations, the course of its duct, and the objects contained within the gland itself.

Parotid Gland.—The parotid salivary gland occupies the space between the ramus of the jaw and the mastoid process. It is bounded above by the zygoma, below by the sterno-mastoid and digastric muscles; behind, by the meatus auditorius and the mastoid process; in front, it lies over the ascending ramus of the jaw, and is prolonged for some distance over the masseter. It is separated from the submaxillary gland by the stylo-maxillary ligament: sometimes the two glands are directly continuous.

The superficial structure of the gland is flat, and covered by a strong layer of fascia, a continuation of the cervical fascia. It not only envelopes the whole gland in a common sheath, but sends down numerous partitions which form a frame-work for the individual lobes. The density of this sheath explains the pain caused by inflammation of the gland, the tardiness with which abscesses within it make their way to the surface, and the propriety of an early opening.

The deep surface of the gland is irregular, and moulded upon the subjacent parts. Thus, it extends inwards between the neck of the jaw and the internal lateral ligament; it reaches the styloid process and the muscles attached to it, and sometimes penetrates deep enough to be in contact with the internal jugular vein.

That portion of the gland which lies on the masseter muscle is called the "*glandula socia parotidis*." It varies in size in

different cases. The greater part of it is situated above the parotid duct, into which it pours its secretion by one or two smaller ducts.

The *duct* of the parotid gland (ductus Stenonis*), is thick and strong. In this respect it differs much from the duct of the submaxillary gland, which is less exposed to injury. It runs forwards over the masseter muscle, perforates the buccinator obliquely, and opens into the mouth opposite the second molar tooth of the upper jaw. Near its termination it is crossed by the zygomaticus major and the facial vein. After perforating the buccinator, the duct passes for a short distance between the muscle and the mucous membrane. Its orifice is small and contracted when compared with the diameter of the rest of the duct, which will admit a crow-quill: it is not easily found in the mouth, because it is concealed by a fold of mucous membrane.

Since it is desirable, in operations about the face, to avoid injuring the parotid duct, it is well to know that the precise direction of the duct corresponds with a line drawn from the tip of the lobe of the ear to a point midway between the nose and the upper lip.

Now cut into the substance of the parotid, to see the objects contained in its interior, which are as follow, proceeding in the order of their depth from the surface:—

1. Two or more small absorbent glands.
2. The “pes anserinus,” or the primary branches of the facial nerve.
3. The external jugular vein formed by the junction of the internal maxillary and the temporal veins.
4. The external carotid, which, after giving many branches to the gland, divides opposite the neck of the jaw, into the internal maxillary and temporal arteries.†

* Nic. Steno, De glandulis oris, &c. Lugd. Bat. 1661.

† Reviewing the intricate and deep connections of the parotid gland, one cannot but conclude that it is almost, if not quite impracticable, to remove it entirely during life. If this conclusion be correct, even in the normal condition of the gland, what must it be when the gland is enlarged by disease? John Bell, however, relates a case in which he was induced to attempt the extirpation of a diseased parotid (J. Bell's Principles of Surgery, vol. iii. p. 262). Other surgeons, too, of more modern date, have attempted the same thing. It

[This gland was first extirpated in this country by Dr. J. Warren, of Boston, in 1798; by Dr. McClellan, of Franklin County, Pa., in 1805; also successfully removed by Dr. S. White, of Hudson, N. Y., in 1808. It has also been removed by Drs. Mott, Parker, and other surgeons of this country. It was not, however, generally regarded feasible until Dr. George McClellan, in 1826, showed that the difficulties attendant on its removal were less in a diseased than in a healthy condition.]

The absorbent glands about the parotid deserve notice, because they are liable to become enlarged, and thus to simulate disease of the parotid itself. An absorbent gland lies close to the root of the zygoma, just in front of the cartilage of the ear: this gland is sometimes affected in disease of the external tunics of the eye; for instance, in purulent ophthalmia.

The easiest way to display the plexus of nerves (*pes anserinus*), formed by the branches of the facial in the parotid gland, is to find one of the larger branches, say one of the malar, on the face, and then to trace this into the substance of the gland, as a clue to the others.

Portio Dura or Facial Nerve.—This is the motor nerve of the face. It supplies all the muscles of expression except those which move the eyes. It arises immediately below the “pons Varolii,” from the lateral tract of the medulla oblongata. The nerve enters the meatus auditorius internus, traverses a canal (“*aqueductus Fallopii*”) in the petrous portion of the temporal bone, and leaves the skull at the stylo-mastoid foramen. Its course and connections in the temporal bone will be investigated hereafter: at present we are interested in the facial part of the nerve.

Having emerged from the stylo-mastoid foramen, the nerve enters the parotid gland, and soon divides into two primary branches, named, from their distribution, the *temporo-facial* and the *cervico-facial*. These primary branches cross over the external carotid, and the external jugular vein; they then form, by their communications within the substance of the parotid, the plexus to which the term “*pes anserinus*” has been ap-

is not unlikely that they have mistaken a tumor in the substance of the parotid for disease of the parotid itself.

plied, from its resemblance to the skeleton of a goose's foot. You ought to trace this plexus and the direction of its ramifications carefully; for you may be called upon to remove tumors formed in the interior of the parotid; you may find the nerves spread over the tumor, and you must dissect between the nerves without injuring them.

Close to the stylo-mastoid foramen the facial nerve sends off its *posterior auricular* branch (p. 32), which supplies the muscles of the ear and the posterior portion of the occipito-frontalis; also a branch to the posterior belly of the digastricus, and another to the stylo-hyoideus.

The *temporo-facial* division crosses the external carotid and the neck of the jaw, receives two or more communications from the temporo-auricular (branch of the fifth), and subdivides into temporal, malar, and infra-orbital branches.

The *temporal* branches ascend over the zygoma, supply the frontalis, the attrahens, and attollens aurem, and communicate with filaments of the supra-orbital nerve.

The *malar* branches cross the zygoma, supply the orbicularis oculi, and communicate with filaments of the lachrymal nerve.

The *infra-orbital* branches proceed forwards beneath the zygomatici over the masseter, and supply the elevators of the upper lip and the muscles of the nose. One large filament accompanies the parotid duct, and supplies the buccinator. Beneath the levator labii superioris we find a free communication with the infra-orbital branches of the second division of the fifth nerve.

The *cervico-facial* division, joined by filaments from the auriculo-parotidean (branch of cervical plexus), descends towards the angle of the jaw, and subdivides into supra and infra-maxillary branches.

The *supra-maxillary* branches advance over the masseter and facial artery, and run under the depressor muscles of the lower lip, all of which they supply. Some of the filaments communicate with the mental branch of the dental nerve.

The *infra-maxillary* or cervical branches, one or more in number, were dissected with the neck (page 5). They arch forwards below the jaw, covered by the platysma, and com-

municate with the *superficialis colli*, a branch of the cervical plexus.

Respecting the function of the facial nerve, it is necessary to remember that though at its origin it is purely a motor nerve, yet after leaving the stylo-mastoid foramen it becomes a compound nerve, in consequence of the filaments which it receives from the temporo-auricular branch of the fifth, and from the auriculo-parotidean branch of the cervical plexus. These communications explain the pain which is often felt in facial paralysis along the track of the facial nerves.

Sensitive Nerves of the Face.—These are the supra-orbital, the infra-orbital, and the mental, all branches of the fifth pair.

The *supra-orbital* nerve is a branch of the first division of the fifth pair. It leaves the orbit through the supra-orbital notch, and is at first covered by the orbicularis, and occipito-frontalis. But it presently divides into wide-spreading branches, which supply the skin of the forehead and scalp. The supra-orbital artery is a branch of the ophthalmic.

The *infra-orbital* nerve is the terminal branch of the superior maxillary or second division of the fifth nerve. It emerges with its artery from the infra-orbital foramen, covered by the levator labii superioris. The nerve immediately divides into several branches, of which some, ascending beneath the orbicularis, supply the lower eyelid; others pass inwards to supply the nose; by far the greater number descend into the upper lip, and eventually terminate in lashes of filaments, which endow the papillæ of the lip with exquisite sensibility.

The *infra-orbital* artery is the terminal branch of the internal maxillary; it supplies the muscles and skin, and inosculates with branches of the facial.

The *mental* nerve is a branch of the inferior maxillary or third division of the fifth nerve. It emerges from the mental foramen in the lower jaw, in a direction upwards and backwards, beneath the depressor labii inferioris. It soon divides into a number of branches, some of which supply the skin of the chin, but the greater number ascend to the lower lip, in the papillæ of which they terminate.

The *mental* artery is a branch of the inferior dental. It supplies the gums and the chin, and inosculates with the submental and inferior coronary arteries.

MUSCLES OF MASTICATION.—TEMPORAL AND SPHENO-MAXILLARY REGIONS.

In the dissection we are about to commence, I advise you to examine the parts in the following order :—

- | | |
|---|--|
| 1. Superficial arteries and nerves of the temple. | 5. Pterygoid muscles. |
| 2. Masseter muscle. | 6. Internal maxillary artery and branches. |
| 3. Temporal aponeurosis. | 7. Inferior maxillary nerve and branches. |
| 4. Temporal muscle. | |

Reflect the skin of the temple in any way most convenient. Under the skin you will come upon a layer of tough fibro-cellular tissue, which is continuous, above, with the aponeurosis of the scalp, below, with the fascia covering the masseter and the parotid gland. In this tissue are contained the superficial temporal vessels and nerves, of which let us trace the ramifications.

Temporal Artery.—This is one of the terminal branches of the external carotid. Arising in the substance of the parotid gland near the neck of the jaw, it passes *over* the root of the zygoma close to the meatus auditorius, ascends for about an inch and a half upon the temporal fascia, and then divides into an anterior and a posterior branch. Above the zygoma it is superficial, and is only covered by the *attrahens aurem* muscle. It gives off the following branches :—

- a. Several small branches to the parotid gland.
- b. Sometimes the *transversalis faciei* (p. 62.)
- c. The *anterior auricular* branches ramify upon the pinna of the ear, inosculating with the posterior auricula.

Of the two branches into which the temporal divides, the *anterior* is usually chosen for arteriotomy. It ascends tortuously towards the external angle of the frontal bone, from which it is distant generally about an inch. Its ramifications extend over the forehead, supplying the orbicularis and frontal muscles, and inosculate with the other arteries of the scalp. The *posterior* runs towards the back of the head, and inosculates freely with the occipital and posterior auricular arteries.

Temporo-auricular Nerve.—This nerve supplies the tem-

ples and side of the head with common sensibility. It arises from the third division of the fifth pair by two roots (between which the arteria meningeae media runs). From its origin it proceeds outwards between the neck of the jaw and the internal lateral ligament, to the root of the zygoma, *over* which it runs with the temporal artery. Its ramifications correspond with those of the artery.

Near the condyle of the jaw the temporo-auricular nerve sends branches to the upper division of the facial nerve, endowing it with common sensibility. Above the zygoma it gives filaments (auricular) to the pinna of the ear.

Lastly, in the subcutaneous tissue of the temple we find the temporal branches of the facial nerve, which supply the frontalis, the attrahens, and attollens aurem.

Masseter Muscle.—This muscle *arises* from the lower edge of the zygoma, and is *inserted* into the outer side of the ramus and coronoid process of the jaw. The masseter is composed of a superficial and a deep layer of fibres which cross each other like the letter X. The superficial fibres, constituting the principal part of the muscle, arise from the anterior two-thirds of the zygoma by a strong tendon which occupies the front border of the muscle, and sends aponeurotic partitions into its substance. The fibres incline backwards, and are inserted into the angle and part of the ramus of the jaw. The deep fibres arise from the posterior part of the zygoma, incline forwards, and are inserted into the upper half of the ramus and the coronoid process. Besides these, a few fibres, taking origin from the inner surface of the zygoma, are inserted into the coronoid process and the tendon of the temporal muscle. Its *action* is to masticate the food: it closes the jaw with great force.

The following objects lie upon the masseter: 1. Glandula sociæ parotidis, and parotid duct; 2. Transversalis faciei artery; 3. Facial artery and vein; 4. Branches of the facial nerve.

Reflect the masseter from its origin. Observe the direction of the superficial and deep fibres, and the tendinous partitions of which the object is to augment the power of the muscle by increasing its extent of origin. The masseteric nerve and artery enter the under surface of the muscle through

the sigmoid notch of the jaw; the artery comes from the internal maxillary, the nerve from the motor division of the inferior maxillary.

Temporal Aponeurosis.—This is the strong shining membrane which covers the temporal muscle. It is attached, above, to the temporal ridge, below, to the zygoma. Its chief use is to give additional origin to the temporal muscle. It increases in thickness from above downwards, and near the zygoma divides into two layers, which are attached to the outer and inner surfaces of the zygoma. These layers are separated by a small quantity of fat. The density of this aponeurosis explains why abscesses in the temporal fossa rarely point outwards; the matter makes its way beneath the zygoma, into the mouth.

Reflect the aponeurosis from the temporal muscle. Notice that the aponeurosis is separated from the muscle near the zygoma by fat. The absorption of this fat, and the wasting of the muscle, occasion the falling in of the temple in the aged and those emaciated by disease.

Temporal Muscle.—This muscle *arises* from the temporal fossa and the temporal aponeurosis. It is *inserted* into the inner surface and anterior border of the coronoid process.

The fibres of this muscle converge from their wide origin, pass under the zygomatic arch, and terminate upon their tendon, the outer surface of which is partially concealed by the insertion of those fibres which come from the temporal aponeurosis; remove them, and you will see how admirably this tendon is adapted to the muscle, and radiates into its substance like the ribs of a fan. Thus the whole force of the muscle is collected into one focus.*


Spheno-maxillary Region.—Remove with a small saw the zygomatic arch; this will expose the coronoid process of the jaw, the insertion of the temporal muscle, and the loose fat which surrounds it. Now saw through the coronoid process in a direction downwards and forwards, so as to include the

* The power possessed by the muscles which close the jaw is quite astonishing. Haller (Elem. Phys. lib. xi. sect. 2, § 26) mentions that one Thomas Topham lifted with his lower jaw a table six feet in length, with 50 lbs. attached to its lower end. It is not uncommon to see mountebanks balance on their chins a ladder so heavy that they can hardly lift it.

insertion of the temporal muscle; turn it upwards with the muscle without injuring the subjacent vessels and nerves. Then ask your demonstrator how much of the neck and ramus of the jaw should be removed in order to obtain a good view of the spheno-maxillary region.

In the spheno-maxillary region we have to examine the two pterygoid muscles, the internal maxillary artery, and the inferior maxillary nerve.

Fig 10.



PTERYGOID MUSCLES AND INTERNAL MAXILLARY ARTERY.

External Pterygoid.—This muscle *arises* by two heads, one from the external pterygoid plate, the other from the great wing of the sphenoid. It is *inserted* into the neck of the jaw, and slightly into the inter-articular fibro-cartilage of the joint of the jaw.

The object of the insertion of some of its fibres into the inter-articular cartilage is, that the cartilage may follow the condyle in all its movements. When the jaw is dislocated, it is chiefly by the action of this muscle, which pulls the condyle

into the zygomatic fossa; the inter-articular cartilage is dislocated with the condyle.

Internal Pterygoid.—This muscle *arises* from the inner surface of the external pterygoid plate, and tuberosity of the palate bone. It is *inserted* into the inner side of the angle of the jaw.

Notice particularly the direction of the fibres of the pterygoid muscles, else you cannot understand their action upon the jaw. The fibres of the external pterygoid run horizontally outwards and backwards from their origin; the fibres of the internal pterygoid run downwards and backwards from their origin. In structure, these muscles are similar to the masseter; that is, they are intersected by tendinous septa for the purpose of giving origin to muscular fibre.

The action of the pterygoid muscle is to produce the lateral movement of the jaw essential to the proper mastication of the food. Consequently we find them enormously developed in all ruminants, and comparatively feeble in carnivorous animals.

Disarticulate the condyle of the jaw with its fibro-cartilage from the glenoid cavity, and turn it forwards with the external pterygoid, so that the condyle can be replaced if desirable. A little dissection will bring into view the internal maxillary artery.

Internal Maxillary Artery.—This is one of the terminal branches into which the external carotid divides opposite the neck of the jaw. It runs horizontally forwards between the neck of the jaw and the internal lateral ligament; it then runs tortuously in some cases above, in others beneath the external pterygoid, enters the spheno-maxillary fossa between the two heads of the external pterygoid, and terminates by dividing into several branches.


We divide the course of this artery into three stages. In the first stage, the artery lies between the neck of the jaw and the internal lateral ligament; in the second stage, it lies either over or under the external pterygoid; in the third stage, it lies in the spheno-maxillary fossa.

BRANCHES OF THE INTERNAL MAXILLARY ARTERY IN THE THREE STAGES OF ITS COURSE.

BRANCHES IN THE FIRST STAGE.	BRANCHES IN THE SECOND STAGE.	BRANCHES IN THE THIRD STAGE.
Tympanic.	Six to the muscles of mastication, namely:	Superior dental.
Meningea media.	Masseteric.	Infra-orbital.
Meningea parva.	Anterior and posterior temporal.	Descending palatine.
Inferior dental.	External and internal pterygoid.	Vidian.
	Buccal.	Nasal.
		Pterygo-palatine.

The *tympanic* artery passes through the glenoid fissure to the tympanum. This small branch varies as to its origin.

The *middle meningeal* artery ascends between the two roots of the temporo-auricular nerve through the foramen spinosum



into the cranium, where it ramifies between the dura mater and the bones.

The *meningea parva* ascends through the foramen ovale into the skull, and supplies chiefly the ganglion of the fifth pair. It often comes from the *meningea media*.

The *inferior dental* artery descends behind the neck of the jaw to the dental foramen, which it enters in company with

the dental nerve. It then proceeds through a canal in the diploe to the symphysis, where it minutely inosculates with its fellow. In this canal, which runs beneath the roots of all the teeth, the artery gives branches which ascend through the little apertures in the fangs, and ramify upon the pulp in their interior. Opposite the foramen mentale arises the mental branch already described (p. 66). Before entering the jaw the dental artery furnishes a small branch which accompanies the nerve proceeding to the mylo-hyoid muscle.

The *masseteric* branch passes through the sigmoid notch of the jaw to the under surface of the masseter, with the masseteric nerve.

The anterior and posterior *temporal* arteries ascend to supply the temporal muscle, ramifying close to the bone, one near the front, the other near the posterior border of the muscle.

The *pterygoid* branches supply the internal and external pterygoid muscles.

The *buccal* branch runs forwards with the buccal nerve to the buccinator.

The *superior-dental* branch runs along the back part of the superior maxillary bone, and sends small arteries through the foramina in the bone to supply the pulps of the molar and bicuspid teeth. It also supplies the gums.

The *infra-orbital* branch passes through the spheno-maxillary fissure, then runs forwards along the infra-orbital canal with the superior maxillary nerve, and emerges upon the face at the infra-orbital foramen. While in the infra-orbital canal the artery sends branches downwards through little canals in the bone to supply the incisor and canine teeth.

The *descending palatine*, a branch of considerable size, runs down the posterior palatine canal with the palatine nerve, and then along the hard palate, in which it is lost. It supplies the gums, the glands, and mucous membrane of this part, and furnishes branches to the soft palate.

The *Vidian*, an insignificant branch, runs backwards through the Vidian canal with the Vidian nerve, and is lost upon the Eustachian tube.

The *nasal* branch enters the nose through the spheno-palatine foramen in company with the nasal nerve from Meckel's

(spheno-palatine) ganglion, and ramifies upon the spongy bones and the septum narium.

The *pterygo-palatine* is a small but constant branch which runs backwards through the pterygo-palatine canal, and ramifies upon the upper part of the pharynx and the Eustachian tube.

[The internal maxillary is very constant as regards its place of origin. It has been found arising from the facial. When it passes beneath the external pterygoid it is found lodged in a depression in the posterior margin of the external pterygoid plate. Where the internal carotid has been absent, two branches from the internal maxillary have been found entering the foramen rotundum and ovale to supply its place.]

The *internal-maxillary vein* is formed by the veins corresponding to the branches of the artery. It joins the temporal vein in the substance of the parotid gland.

Inferior Maxillary Nerve and Branches.—This great nerve is the largest of the three divisions of the fifth cerebral nerve. It differs from the other two divisions, *i. e.* the ophthalmic and the superior maxillary, in that it contains motor as well as sensitive filaments; the motor being furnished by the small non-ganglionic root of the fifth nerve. Thus much of its physiology it is necessary to know, in order to understand the extensive distribution of the nerve; for we shall find that the motor portion supplies all the muscles concerned in the movements of the lower jaw;* whilst the other portion confers only common sensibility upon the parts to which it is distributed.

The nerve, then, composed of sensitive and motor filaments, emerges from the skull through the foramen ovale as a single thick trunk, under the name of the inferior maxillary. It lies directly external to the Eustachian tube: it is covered by the external pterygoid muscle, which should be turned on one side to expose it. Immediately after its exit from the skull, the nerve divides into its several branches; some, destined for the muscles, contain motor as well as sensitive filaments; others are purely sensitive.

* Excepting the genio-hyoidei muscles, which are supplied by the hypoglossal nerve.

BRANCHES OF THE INFERIOR MAXILLARY NERVE.

BRANCHES SENSITIVE.

Temporo-auricular.
Inferior dental.
Gustatory.
Buccal.

BRANCHES MOTOR.

To temporal muscle.
— masseter.
— external pterygoid.
— internal pterygoid.
— tensor palati.
— mylo-hyoideus.
— anterior belly of digastricus.


The branches to the *temporal* muscle pass outwards close to the great wing of the sphenoid bone, and ascend in company with the temporal arteries to the muscle.

The branch to the *masseter* runs outwards above the external pterygoid, through the sigmoid notch of the jaw.

The branch to the external pterygoid comes apparently from the buccal nerve in its passage through this muscle.

The branch to the *internal pterygoid* and *tensor palati* muscles is rather difficult to find. It proceeds from the inner side of

Fig 12



PLAN OF THE BRANCHES OF THE INFERIOR MAXILLARY NERVE.

the main trunk, descends between the internal pterygoid and the tensor palati, and supplies both.

The *buccal* branch passes either above or between the fibres of the external pterygoid to the buccinator, where it spreads out into filaments, which supply the skin, the mucous membrane and glands of the cheek, with common sensibility. The *motor* power of the buccinator, remember, is derived from the facial nerve. This is proved not only by careful dissection, but by the fact that the muscle loses its power of contraction when the facial nerve is paralyzed; and that it retains its power of contraction when the muscles of mastication are paralyzed.

The *temporo-auricular* branch arises by two roots which embrace the middle meningeal artery before it enters the skull. The nerve runs outward behind the neck of the jaw, ascends over the root of the zygoma with the superficial temporal artery, and ramifies with this artery upon the temple and side of the head, supplying the skin. While behind the condyle it sends filaments to the skin lining the meatus auditoris. Lastly, branches proceed from this nerve to the upper division of the facial, and endow it with common sensibility.

The *inferior dental* branch descends between the ramus of the jaw and the internal lateral ligament of the jaw to the dental foramen, which it enters with the dental artery. It then runs in the canal in the diploe of the jaw below the fangs of all the teeth. It furnishes filaments which ascend through the canals in the fangs of the teeth to the pulp in their interior. Opposite the foramen mentale the mental branch is given off. Observe that the same nerve which supplies the teeth supplies the gums; hence the sympathy between these parts.

The *mylo-hyoid* branch, apparently originating from the dental, and so described, is derived from the motor root of the fifth, and may, with careful dissection, be traced to it. It leaves the dental nerve near the foramen in the jaw, and runs in a groove on the inner side of the ramus to the lower surface of the mylo-hyoid, which muscle, together with the anterior portion of the digastricus, it supplies (p. 25.)

The *gustatory* branch descends first between the ramus of the jaw and the internal pterygoid muscle, then for a short

distance between the jaw and the superior constrictor of the pharynx. In this situation it lies close to the mucous membrane of the mouth near the last molar tooth of the lower jaw. The gustatory n. then runs along the hyo-glossus muscle to the tongue, and is distributed to the papillæ of the tip and sides of that organ.

The gustatory nerve is joined by the chorda tympani (a branch of the facial nerve), which emerges from the glenoid fissure, and eventually runs to the submaxillary ganglion.

Internal Lateral Ligament of the Jaw.—We have now a good opportunity of examining this so-called ligament. By some anatomists it is not recognised as a ligament, and with reason, because it is placed at such a distance from the joint. It proceeds from the spinous process of the sphenoid bone, and is attached to the jaw on the inner side of the foramen dentale. Between this ligament and the neck of the jaw we find the internal maxillary artery and vein, the temporo-auricular nerve, the middle meningeal artery, the dental nerve and artery, and a portion of the parotid gland.

At this stage of the dissection you will be able to trace the course and relations of the internal carotid artery. But before you do this, examine the several objects which intervene between the external and internal carotid. These objects are, 1. The stylo-glossus, 2. The stylo-pharyngeus, 3. The glosso-pharyngeal nerve, 4. The stylo-hyoid ligament.

Stylo-glossus.—This *arises* from the styloid process near the apex, and the stylo-maxillary ligament, and is *inserted* into the side of the tongue, external to the hyo-glossus. Its action is to retract the tongue. Its nerve comes from the hypoglossal.

Stylo-pharyngeus.—This *arises* from the styloid process near the base, and is *inserted* into the posterior edge of the thyroid cartilage. It descends along the side of the pharynx between the superior and the middle constrictor. Its nerve comes from the pharyngeal plexus. Its *action* is to raise the larynx with the pharynx in deglutition.

Between the stylo-glossus and the stylo-pharyngeus, and nearly parallel with both, is the stylo-hyoid ligament. It extends from the apex of the styloid process to the lesser cornu of the os-hyoides. It is often more or less ossified.

The *ascending palatine artery*, a branch of the external maxillary or facial (p. 24), runs up between the stylo-glossus and stylo-pharyngeus, and divides into branches which supply these muscles, the palate, the side of the pharynx, and the tonsils.

Glosso-pharyngeal Nerve.—The best guide to the glosso-pharyngeal nerve is the lower border of the stylo-pharyngeus muscle. This nerve is one of the divisions of the eighth pair. It arises by four or five filaments from the medulla oblongata in the groove between the olive and the restiform tract. At its origin it is purely a sensitive nerve. Leaving the skull through the foramen lacerum posterius, it descends in front of the internal carotid artery, and then proceeds along the lower border of the stylo-pharyngeus. But now it curves forwards over the stylo-pharyngeus (p. 27) and disappears beneath the hyo-glossus, where it divides into its terminal branches for the supply of the mucous membrane of the pharynx, the back of the tongue, and the tonsils.

According to the present state of our knowledge, the glosso-pharyngeal is, at its origin, purely a sensitive nerve. But soon after its exit from the skull it receives communications from the facial and the nervus accessorius, so that it soon becomes a compound nerve, *i. e.* composed of both sensitive and motor filaments. Moreover, at the base of the skull it is provided with a ganglion, called the ganglion of "Andersch." The minute branches given off by this ganglion will be noticed in their proper place; at present we are more concerned with what is called the pharyngeal plexus of nerves.

Pharyngeal Plexus.—By the side of the middle constrictor of the pharynx, we find an intricate interlacement of nerves, constituting the plexus which presides over deglutition. Its dissection requires much time and care, and a pharynx prepared exclusively for the purpose. The nerves which enter into its composition are derived from the glosso-pharyngeal, the pneumogastric, the spinal accessory, and the sympathetic. Consequently it possesses nerves of three different kinds—ganglionic, sensitive, and motor. Its minute ramifications supply the pharynx, the back of the tongue, and tonsils.

Course and Relations of the Internal Carotid Artery.—The internal carotid artery, proceeding from the division of

the common carotid, ascends to the base of the skull, *by the side of the pharynx* close to the transverse processes of the three upper cervical vertebræ. It enters the skull through the carotid canal in the temporal bone, runs very tortuously by the side of the body of the sphenoid, and terminates in branches which supply the orbit and the brain. In the first part of its course, it is situated immediately *outside* the external carotid, near the inner border of the sterno-mastoid muscle. But it soon gets beneath the external carotid, and lies deeply seated by the side of the pharynx. In its course it is crossed successively by the hypoglossal nerve and the occipital artery; still higher, it is crossed obliquely by the styloid process, by the styloid-glossus and pharyngeus muscles, by the glosso-pharyngeal nerve, and the stylo-hyoid ligament, all of which intervene between it and the external carotid.

The internal jugular vein runs along the outer side of the artery. Behind the artery are the pneumogastric nerve and the superior cervical ganglion of the sympathetic. The rectus capitis anticus major separates it from the cervical vertebræ. But after all, the most important relation of the artery in a surgical point of view is, that it ascends close by the *side of the pharynx* and *tonsils*. In opening an abscess, therefore, near the tonsils, or at the back of the pharynx, be careful to introduce the knife with its point inwards towards the mesial line: observe this caution the more, because, in some subjects, the internal carotid makes a curve, or even a complete curl upon itself, in its ascent near the pharynx. In such cases the least deviation of the instrument in an outward direction would injure the vessel; an occurrence of which instances have been recorded.

[When the internal carotid comes off from the arch, it is found nearer to the trachea than the external, which passes forwards over the former vessel at the larynx. Instead of being straight, it may present many curves in its course; a few cases are recorded where this vessel was wanting.]

Ascending Pharyngeal Artery.—This artery generally arises from the angle of the common carotid, or from the commencement of the external carotid; it ascends by the pharynx, towards the base of the skull. It gives off branches which supply the pharynx, the tonsils, the Eustachian tube,

and the muscles in front of the spine. A very constant branch runs down with the levator palati, above the superior constrictor of the pharynx, and supplies the soft palate.

[Two ascending pharyngeal arteries are occasionally met with.]

Pneumogastric Nerve.—The pneumogastric nerve is one of the three divisions of the eighth pair of cerebral nerves. It arises from the medulla oblongata by a series of roots along the groove between the olive and the restiform tract. It passes out of the skull with the nervus accessorius through the foramen jugulare. Very soon a ganglion is observed in its substance analogous to the posterior or sensitive roots of all the spinal nerves, and it is joined by a branch from its companion the nervus accessorius.

Thus, the pneumogastric, probably only sensitive at its origin, becomes in consequence of this reinforcement from the nervus accessorius a compound nerve, and in all respects analogous to a spinal nerve.

Leaving the skull, then, at the foramen jugulare, the nerve descends in front of the cervical vertebræ, lying successively upon the rectus capitis anticus major and the longus colli muscles. In the upper part of the neck it is situated behind the internal carotid artery; in the lower, between and behind the common carotid and the internal jugular vein. It enters the chest, on the right side crossing in front of the subclavian artery nearly at a right angle; on the left running nearly parallel with it.

In their course through the chest, the pneumogastric nerves have not similar relations. The right nerve descends by the side of the trachea and then behind the right bronchus to the posterior part of the œsophagus. The left nerve crosses in front of the arch of the aorta, and then passes behind the left bronchus to the anterior part of the œsophagus. Both nerves accompany the œsophagus into the abdomen, and are eventually distributed to the stomach.

In their long course from the medulla oblongata to the abdomen, the pneumogastric nerves supply branches to most important organs: namely, to the pharynx, the larynx, the heart, the lungs, the œsophagus, and the stomach.

Within the foramen jugulare, a small ganglion (Arnold's

ganglion) is formed upon the pneumogastric nerve. This ganglion will be described hereafter. But soon after leaving the skull, the pneumogastric nerve swells considerably, and forms a second ganglion (inferior ganglion) of a reddish-grey color.

This ganglion occupies about an inch of the nerve. It is united to the hypoglossal nerve, from which it receives filaments. It also receives filaments from the first and second spinal nerves, and from the superior cervical ganglion of the sympathetic. The branches of this ganglion are:—

1. The *pharyngeal*, which descend either in front of or behind the internal carotid to join the pharyngeal plexus (p. 78).

2. The *superior laryngeal*, which descends behind the external carotid to the interval between the os-hyoides and the thyroid cartilage, where it enters the larynx through the thyrohyoid membrane, and is distributed to the mucous membrane of the larynx. Remember that this superior laryngeal gives off the “external laryngeal” which supplies the crico-thyroid muscle (p. 59).

3. The *cardiac*, two in number, which descend behind the sheath of the carotid to the cardiac plexus. They vary much as to the situation of their origin,—proceeding either from the ganglion, or from the main trunk of the nerve in its course down the neck. On their passage to the heart, the right cardiac nerves run behind the arch of the aorta, the left in front of it, to the great cardiac ganglion. This cardiac ganglion is situated between the arch of the aorta and the trachea; and from it the nerves of the heart are given off.

The *inferior laryngeal* or *recurrent* branch of the pneumogastric turns on the right side under the subclavian artery (p. 40), and runs up to the larynx between the trachea and the bodies of the cervical vertebræ; on the left side, it turns under the arch of the aorta, and ascends to the larynx between the trachea and the œsophagus. On both sides it enters the larynx beneath the lower border of the inferior constrictor.

The remaining branches of the pneumogastric, to the lungs, the œsophagus and stomach will be examined in the dissection of the chest.

Sympathetic Nerve.—We now pass on to examine the cervical ganglia of the sympathetic system of nerves. Speaking in general terms of this system, it may be said that it consists

of a series of ganglia arranged on either side of the spine, from the first cervical to the last sacral vertebra. The successive ganglia of the same side are connected by intermediate nerves, so as to form a continuous cord on each side of the spine; this constitutes what is called the trunk of the sympathetic system. This system of nerves is connected with all the spinal nerves. Its upper or cephalic extremity penetrates into the cranium through the carotid canal, surrounds the internal carotid artery, communicates with the third, fourth, fifth, and sixth cranial nerves, and is said to join its fellow of the opposite side upon the anterior communicating artery.* Its sacral extremity joins its fellow by means of a little "ganglion impar," situated in the mesial line, upon the last sacral vertebra.

The sympathetic system of nerves presides over the functions of those organs which are withdrawn from our control. Thus it regulates the circulation of the blood, respiration, digestion, and secretion; all of which processes go on without our being conscious of them.

Cervical Ganglia of Sympathetic.—In the cervical portion of the sympathetic are three ganglia, named, from their position,—superior, middle, and inferior.

The *superior cervical ganglion*, the largest of the three, is situated near the base of the skull, upon the rectus capitis anticus major behind the internal carotid artery. It is of a reddish-grey color like other ganglia, of an elongated oval shape, varying in length from one to two inches. It may facilitate the description of its several branches if we divide them into—1st, those which are presumed to connect it with other nerves; and 2ndly, those which originate from it.

It is then connected by branches as follows:—

- a. With each of the four upper spinal nerves.
- b. With the hypoglossal, pneumogastric, and glosso-pharyngeal nerves.
- c. With the third, fourth, fifth, and sixth cerebral nerves (in the cavernous sinus).
- d. With the several ganglia of the sympathetic system about the head and neck; namely, the ophthalmic, sphenopalatine, otic, and submaxillary.

* Here, it is said, is situated the little ganglion of Ribes. I have never seen it.

The branches which it distributes are—

e. Nerves to the heart.—One or more descend behind the sheath of the carotid, and entering the chest, join the cardiac plexus.

f. Nerves to the pharynx.—These join the pharyngeal plexus.

g. Nerves to the blood-vessels.—These nerves, named on account of their delicacy “*nervi molles*,” ramify around the external carotid artery and its branches.

The *middle cervical ganglion* is something less than a barley-corn in size. It is situated behind the carotid sheath, about the fifth or sixth cervical vertebra, near the inferior thyroid artery. It receives branches from the fifth and sixth spinal nerves, and gives off—

a. Branches to the thyroid body.—These accompany the inferior thyroid artery.

b. Branch to the heart.—This usually descends in front of the subclavian artery into the chest, and joins the cardiac plexus.

In cases where the middle cervical ganglion is absent, the preceding nerves are supplied by the sympathetic cord connecting the superior and inferior ganglia.

The *inferior cervical ganglion* is of considerable size, and is situated in the interval between the transverse process of the seventh cervical vertebra and the first rib, immediately behind the vertebral artery. It receives branches from the seventh and eighth spinal nerves, and others which, descending from the fourth, fifth, and sixth, through the foramina in the transverse processes of the vertebræ, form a plexus around the vertebral artery.

The only branch which it gives off is—

a. Inferior cardiac nerve.—This communicates with the recurrent laryngeal, and joins the cardiac plexus beneath the arch of the aorta.

DISSECTION OF THE CHEST.

Before we examine the several organs contained in the chest, we ought to have some knowledge of its frame work. The true ribs with their cartilages describe a series of arcs, in-

creasing in length from above downwards, and form, with the spine and the sternum, a barrel of a conical shape, broader in the lateral than in the antero-posterior direction. The lower aperture or base of the cavity is closed in the recent state by a muscle called the diaphragm, which forms a partition between the chest and the abdomen. The partition is not flat, but arched, so that it constitutes a vaulted floor for the chest, and moreover by its capability of alternately falling and rising can increase or diminish the capaciousness of the chest. The spaces between the ribs are filled by the intercostal muscles. In each intercostal space there are two layers of these muscles, arranged like the letter X. The fibres of the outer layer run obliquely from above downwards, and forwards; those of the inner layer in the reverse direction.

The upper aperture of the chest gives passage to the trachea, the oesophagus, the great vessels of the neck and the arms, and also to certain muscles and nerves; the interspaces between these parts being occupied by a dense fibro-cellular tissue.

Such, in outline, is the frame-work of the cavity, closed on all sides, which contains the heart and lungs. Observe that its walls are made up of different structures, bone, cartilage, and muscle, admirably disposed to fulfil two important conditions: by their solidity and elasticity they protect the important organs contained in them; by their alternate dilatation and contraction they act as mechanical powers of respiration. For they can increase the cavity of the chest in three directions: in height, by the descent of the diaphragm; in width, by the turning outwards of the ribs; and in depth, by the elevation of the sternum.

The chest of the female differs from that of the male in the following points:—Its general capacity is less; the sternum is shorter; the upper opening is larger in proportion to the lower; the upper ribs are more movable, and therefore permit a greater enlargement of the chest at its upper part, in adaptation to the condition of the abdomen during pregnancy.

In the dissection of the chest let us take the parts in the following order:—

1. *Triangularis sterni*, with the internal mammary artery.

2. Anterior mediastinum.
3. Right and left brachio-cephalic veins and superior vena cava.
4. Course and relations of the arch of the aorta.
5. The three great branches of the arch.
6. Course of the phrenic nerves.
7. Posterior mediastinum and its contents; namely, the aorta, the thoracic duct, the vena azygos, the œsophagus, and pneumogastric nerves.
8. Sympathetic nerve.
9. Intercostal muscles, vessels, and nerves.
10. Cardiac plexus of nerves.
11. Position and relations of the heart.
12. Pericardium.
13. Position and form of the lungs.
14. Pleura.

In the first place you must make a window into the chest, by carefully removing the sternum and the cartilages of all the true ribs.* Cut through the cartilages of the ribs, close to the bone of the rib, and then raise the sternum with its attached cartilages; in doing this, take care not to wound the pleura which is closely connected with the cartilages. On one side remove the internal mammary artery, on the other leave it.

On the under surface of the sternum and cartilages of the ribs thus removed, is a muscle named "*triangularis sterni*."

Triangularis Sterni.—This muscle *arises* from the ensiform cartilage and the lower part of the sternum, and is *inserted* by digitations into the cartilages of the true ribs from the second to the sixth; its fibres ascend outwards to their insertion; its action therefore is to depress the costal cartilages. It is one of the muscles which act on emergency in expiration. Its nerve comes from the intercostal nerves: its arteries from the internal mammary.

Internal Mammary Artery.—This is a branch of the subclavian. It runs perpendicularly about half an inch from the sternum between the cartilages of the ribs and the *triangularis sterni*, enters the wall of the abdomen behind the rectus abdominis, and finally inosculates with the epigastric (a branch of the external iliac). Its branches are as follow:—

* Those who are more proficient in dissection should not remove the whole of the sternum, but leave, say a quarter of an inch of its upper part with the first rib attached to it. This little portion serves as a valuable landmark, although it obstructs, to a certain extent, the view of the subjacent vessels

a. Arteria comes nervi phrenici.—A very slender artery which accompanies the phrenic nerve to the diaphragm.

b. Mediastinal and thymic.—These branches supply the cellular tissue of the anterior mediastinum, the pericardium, and the triangularis sterni muscle. The *thymic* are only visible in childhood, and disappear with the thymus gland.

c. Intercostal.—One at least, and often two, for each intercostal space. They inosculate with the intercostal arteries from the aorta.

d. Perforating arteries, which pass through the intercostal spaces, and supply the pectoral muscle and skin of the chest. In the female they are of large size, for the supply of the mammary gland.

e. The intercosto-phrenic branch runs outwards behind the cartilages of the false ribs, and terminates near the last intercostal space. It supplies small arteries to the diaphragm, to the sixth, seventh, and sometimes the eighth intercostal spaces.

There are several *absorbent glands* in the neighborhood of the internal mammary artery. They receive the absorbents from the inner portion of the mammary gland, from the diaphragm, and the wall of the abdomen. In disease of the inner portion of the mamma, these glands may enlarge, without any enlargement of those in the axilla.

Mediastina.—The “mediastina” are the spaces which the two pleural sacs leave between them in the antero-posterior plane of the chest. There is an anterior, and posterior mediastinum. To put these spaces in the simplest light, let us imagine the heart and lungs to be removed from the chest, and the two pleural sacs to be left in it by themselves. The two sacs, if inflated, would then appear like two bladders, in contact only in the middle, as shown by the dotted outlines in the annexed scheme (fig. 13). The interval marked *a*, behind the sternum, would represent the anterior mediastinum: the interval *b*, the posterior mediastinum. But now let us introduce the heart and lungs again, between the two pleural sacs; these must give way to make room for them. Therefore, with the heart and lungs interposed, the pleural sacs appear as shown in the diagram, fig. 14, which represents a transverse section through the chest. Here, you see that the heart and

the lungs become invested by the pleural sacs ; or, to speak more technically, the sacs are reflected over the heart and the

lungs, on each side respectively. But the anterior and posterior mediastina remain unaltered.

Looking at the chest in front, the anterior mediastinum appears as shown in the cut (p. 97). It is not precisely longitudinal in its direction, but inclines slightly towards the left, owing to the position of the heart. Its area varies : thus it is extremely narrow

in the middle where the edges of the lungs nearly meet : but it is wider above and below, where the lungs diverge.

Fig. 13.




Fig. 14.




DIAGRAM OF THE REFLECTIONS OF THE PLEURAL SACS IN DOTTED LINES.

What parts are contained in the anterior mediastinum ?—The remains of the thymus gland, the origins of the sterno-hyoid, sterno-thyroid, and triangularis sterni muscles, the left brachio-cephalic vein (which crosses behind the first bone of the sternum), and a few absorbent glands.

The *posterior mediastinum* (fig. 14), contains the œsophagus, the two pneumogastric nerves, the aorta, the thoracic duct, and the vena azygos.

Before we examine the brachio-cephalic veins, we must remove a layer of the cervical fascia which descends from the neck over these veins and is lost upon the pericardium. The coats of all the great veins are closely connected to this fascia; and its chief use appears to be to keep these veins always wide and open for the free return of the blood to the heart.

Brachio-cephalic Veins.—The right and left brachio-cephalic (innominate) veins are formed, near the sternal end of the clavicle, by the confluence of the internal jugular and subclavian. They differ in their course and relations, and must, therefore, be described separately.

The *left brachio-cephalic vein* passes behind the first bone of the sternum from the left toward the right side, to join the vena cava superior. It is about two inches and a half in length, and its direction inclines a little downwards. It crosses over the trachea and three primary branches of the arch of the aorta (p. 39). We are reminded of this fact in some cases of aneurism of these vessels; for what happens? The vein becomes compressed between the aneurism and the sternum; hence the swelling and venous congestion of the parts from which it returns the blood; namely, of the left arm and left side of the neck. Observe, moreover, that the upper border of the vein lies not far from the upper border of the sternum; in some instances it lies even higher, and I have seen it crossing in front of the trachea a full inch above the sternum. Remember this in the performance of tracheotomy.

The *right brachio-cephalic vein* descends nearly vertically to join the superior vena cava. It is about one and a half inch in length, and is situated about one inch from the mesial line of the sternum. On its left side runs the corresponding *arteria innominata*; on its right side is the pleura. Between the vein and the pleura is the phrenic nerve.

The veins which empty themselves into the left and right brachio-cephalic, are as follow :—

The RIGHT B. C. Vein receives:—

The vertebral,
The deep cervical,
The internal mammary.

The LEFT B. C. Vein receives:—

The vertebral,
The deep cervical,
The internal mammary,
The middle thyroid,
The superior intercostal,
The pericardiac.

Vena Cava Superior.—This is one of the great channels through which the impure blood returns into the right auricle. It is formed by the confluence of the right and left brachiocephalic veins, which unite at nearly a right angle opposite the first intercostal space on the right border of the sternum; that is, about the level of the highest point of the arch of the aorta (p. 39). The vena cava descends vertically with a slight inclination backwards, to the upper part of the right auricle. It is from two and a half to three inches long. The lower half of it is covered by the pericardium; you must, therefore, open this bag in order to see that the serous layer of the pericardium is reflected over the front and sides of the vein. In respect to its relations, notice that the vein lies in front of the right bronchus and the right pulmonary vessels, and that it is overlapped by the ascending aorta. In the upper half of its course, that is, above the pericardium, it is covered on its right side by the pleura; on this side, in contact with it, descends the phrenic nerve.

Before it is covered by the pericardium, the vena cava receives the vena azygos, which comes in arching over the right bronchus.

[The two innominate veins which unite to form the superior cava, have been seen to open separately into the right auricle. Then the vein of the right side takes the ordinary course of the vena cava; the left vein sends a branch to the right, then passes to the left side of the heart, and ends on the back of the right auricle with the coronary vein. This is a natural condition in the earlier period of the foetus, and is constant in birds and in some mammalia.]

Course and Relations of the Arch of the Aorta.—The aorta is the great trunk from which all the arteries of the body carrying red blood are derived. It arises from the upper and back part of the left ventricle of the heart. Its

origin is on the left side of the sternum, about the level of the third intercostal space. It ascends forwards and to the right as high as the first intercostal space on the right side; it then curves backwards toward the left side of the body of the second dorsal vertebra, and, turning downwards over the third, completes the arch. The direction of the arch, therefore, is from the sternum to the spine, and rather oblique from the right towards the left side. For convenience of description, we divide it into an ascending, a transverse, and a descending portion.

Ascending portion.—To see this portion of the aorta you must open the pericardium. You then observe that this part of the artery is covered all round by the serous layer of the pericardium, except where it is in contact with the pulmonary artery. Now what are its relations? Its commencement is covered by the pulmonary artery, and overlapped by the appendix of the right auricle. On its right side, but on a posterior plane, descends the superior vena cava; on its left, is the division of the pulmonary artery; behind it, is a part of the auricle, the right pulmonary artery, and the right bronchus.

Transverse portion.—This portion of the aorta crosses in front of the trachea a little above its bifurcation. Near its summit runs the left vena innominata; within its concavity are the left bronchus, the bifurcation of the pulmonary artery, and the remains of the ductus arteriosus.

The transverse part of the arch is crossed by the left phrenic and the left pneumogastric nerves; and the recurrent branch of the left pneumogastric ascends behind it to the larynx.

Descending portion.—This part of the arch lies upon the left side of the body of the third dorsal vertebra. On its right side are the œsophagus and thoracic duct; on its left is the pleura.

What parts are contained within the arch of the aorta?—The left bronchus, the right pulmonary artery, the left recurrent nerve, the remains of the ductus arteriosus, and the great cardiac ganglion.

The arch of the aorta presents partial dilatations in certain situations. One of these, called the sinus or bulge of the aorta, is observed on the right side of the arch about the

junction of the ascending with the transverse portion; just where one might expect such a dilatation from the impulse of the blood. It is little marked in the infant, but increases with age. Three other dilatations (the sinuses of Valsalva), one corresponding to each of the valves at the commencement of the aorta, will be examined hereafter.

[The aorta sometimes arches over the root of the right lung instead of the left, and then continues down the right side of the vertebra. In such a case, we find a transposition of the viscera. Sometimes after having taken this abnormal course, it passes back to its normal position on the left side within the thorax.]

Relations of the Arch of the Aorta to the Sternum.—These relations vary according to the size of the heart, the obliquity of the ribs, and the general development of the chest. In a well-formed adult the ascending aorta is, at the most prominent part of its bulge, about half of an inch behind the first bone of the sternum. The highest part of the arch is about one inch below the upper edge of the sternum.*

From the upper part of the arch arise three great arteries for the head, neck, and upper limbs; namely, the brachio-cephalic or innominate artery, the left carotid, and the left subclavian.

Brachio-cephalic, or Innominate Artery.—This arises from the commencement of the transverse part of the arch. It ascends obliquely toward the right, and after a course of about one inch and a half, divides behind the right sternoclavicular joint into two arteries of nearly equal size—the right carotid, and the right subclavian.

[The length of the innominate sometimes exceeds two inches, at others it is an inch or less. It has usually no branches, though a thyroid and thymic branch have been

* The relations of the arch of the aorta to the sternum vary even in adults, more especially if there be any hypertrophy of the heart. As an instance among many, we may mention that of a young female who died of phthisis. The position of the aortic valves was opposite the middle of the sternum on a level with the middle of the second costal articulation. The highest part of the arch was on a level with the upper border of the sternum; the arteria innominata was situated entirely in front of the trachea; and the left vena innominata crossed the trachea so much above the sternum that it would have been directly exposed to injury in tracheotomy.

seen coming off, as well as the left common carotid. Its place of bifurcation is interesting in a surgical view, as upon this depends the success of ligating the vessel. This vessel has been tied eleven times. First by Dr. Valentine Mott at the New York Hospital, in 1818. The only successful case was the last one recorded. The operation was performed in New Orleans, May 15, 1864, by Dr. A. W. Smith, at the Charity Hospital. In this instance, the common carotid was also tied, and subsequently the vertebral, owing to secondary hemorrhage.]

What are the relations of the innominate artery? It ascends obliquely in front of the trachea; the right vena innominata descends on its right side; the left vena innominata crosses in front of it: parallel and close to the artery are the slender cardiac nerves.*

With the anatomy of the parts before you, it is easy to understand that an aneurism of the innominate artery might be distinguished from an aneurism of the aorta. 1. By a pulsation in the neck between the sterno-mastoid muscles, *i. e.* in the fossa above the sternum; 2. By occasional dyspnoea owing to pressure on the trachea; 3. By venous congestion in the *left* arm; 4. By the aneurismal thrill being confined to the *right* arm.


Left Carotid Artery.—This artery arises from the arch of the aorta close to the arteria innominata. It ascends obliquely behind the first bone of the sternum to the neck. In the first part of its course it lies upon the trachea, but it soon passes to the left side of the trachea, and then lies for a short distance upon the œsophagus. In the rest of its course it resembles the right carotid (p. 13).

Left Subclavian Artery.—This is the third branch of the arch. It ascends out of the chest to the inner border of the first rib, and then curves outwards behind the scalenus anticus. In the first part of its course it is deeply seated, and is covered on its left side by the pleura. Close to its right side

* In some cases the innominate artery ascends for a short distance above the clavicle before it divides, lying close to the right of the trachea. We have already alluded to the fact that it occasionally gives off a middle thyroid artery (p. 17), which ascends in front of the trachea to the thyroid body, and is therefore directly in the way in tracheotomy.

is the œsophagus ; between it and the œsophagus we find the thoracic duct. Like the other primary branches of the arch, it is crossed by the left brachio-cephalic vein. The upper part of its course, where the vessel passes in front of the apex of the lung, has been described with the anatomy of the neck (p. 39).

Fig. 15.



Course of the Phrenic Nerves through the Chest.—The phrenic nerve comes from the third, fourth, and fifth cervical nerves. It descends over the scalenus anticus, and enters the chest between the subclavian vein and artery. It then descends in front of the root of the lung between the pleura

and the pericardium to the diaphragm (fig. 15) which muscle it supplies.

In what respects do the phrenic nerves differ from each other in their course?—The right phrenic runs along the outer side of the innominate vein and superior vena cava; the left crosses in front of the transverse part of the arch of the aorta; besides which, the left is rather longer than the right, because it has to go round the point of the heart.*

On the upper surface of the diaphragm the phrenic divides into a number of branches, which are distributed, some to the greater, others to the lesser crus of the muscles.†

Having studied these dry anatomical details, consider for a moment what kind of symptoms are likely to be produced by an aneurism of the arch of the aorta, or any of the primary branches. A glance at the important parts in the neighborhood will answer the question. The effects will vary according to the part of the artery which is the seat of the aneurism, and according to the volume, the form, and the position of the tumor. One can understand that compression of the vena cava superior, or either of the brachio-cephalic veins, would occasion swelling and congestion of the parts from which it returns the blood; that compression of the trachea or one of the bronchi might occasion dyspnoea, and thus simulate disease of the larynx‡; that compression of the œsophagus would give rise to symptoms of stricture. Nor must we forget the immediate vicinity of the thoracic duct and the recurrent nerve,§ and the effects which would be produced by their compression. Can one, then, be surprised that a disease which may give rise to so many different symptoms should be a fertile source of fallacy in diagnosis?

* In the upper part of the chest, the phrenic is sometimes joined by a branch from the brachial plexus, and less frequently by a branch from the descendens noni.


† Does the phrenic ever send filaments to the pericardium? In some cases it unquestionably does; hence possibly the pain in the arm and shoulder in some diseases of the heart.

‡ In the Museum of Guy's Hospital there is a preparation, No. 1487, in which laryngotomy was performed under the circumstances described in the text.

§ See Med. Gaz., Dec. 22nd, 1843. A case in which loss of voice was produced by the pressure of an aneurismal tumor upon the left recurrent nerve.

You can easily see how aneurisms of the aorta prove fatal, by bursting into the contiguous tubes or cavities; for instance, into the trachea, the œsophagus, the pleura, or the pericardium. You will see too, why an aneurism of the first part of the arch is so much more dangerous than elsewhere. The reason is, that in this part of its course the aorta is covered only by a thin layer of serous membrane: now if an aneu-

Fig. 16.



RELATIVE POSITION OF THE HEART AND ITS VALVES WITH REGARD TO THE WALLS OF THE CHEST.

The valves are denoted by curved lines. The *aortic valves* are opposite the third intercostal space on the left side, close to the sternum. The *pulmonary valves* are just above the aortic, opposite the junction of the third rib with the sternum. The *mitral valves* are opposite the third intercostal space, about one inch to the left of the sternum. The *tricuspid valves* lie behind the middle of the sternum, about the level of the fourth rib. *Aortic murmurs*, as shown by the arrow, are propagated up the aorta; *mitral murmurs*, as shown by the arrow, are propagated towards the apex of the heart.

rism take place here, the coats of the vessel soon become distended, give way, and allow the blood to escape into the pericardium; an occurrence which is speedily fatal, because the pericardium being filled with blood prevents the heart from acting.

Position and Form of the Heart.—The heart is situated obliquely in the chest, between the lungs. Its base, *i. e.*, the part by which it is attached, and from which its great vessels proceed, is directed upwards toward the right shoulder; its apex points downwards and to the left, between the fifth and sixth ribs. It is maintained in its position by a membranous bag termed the “pericardium;” which is lined by a polished membrane to facilitate its movements. It is supported by the tendinous centre of the diaphragm.


Cut open the pericardium, and observe that the heart is conical in form, and convex everywhere except upon its lower surface, which is flat, and rests upon the tendinous centre of the diaphragm. But this general observation is not sufficient. You must notice the several objects exposed, when the pericardium is thus laid open; they are as follow:—1. Part of the right ventricle; 2. Part of the left ventricle; 3. Part of the right auricle with its appendix overlapping the root of the aorta; 4. The appendix of the left auricle overlapping the root of the pulmonary artery; 5. The aorta; 6. The pulmonary artery; 7. The vena cava superior.

The heart then is placed behind the lower half of the sternum, occupies more of the left than the right half of the chest, and rests upon the central tendon of the diaphragm, which is nearly on a plane with the lowest part of the fifth rib. At each contraction the apex of the heart may be felt beating between the fifth and sixth ribs, about two inches below the nipple, and one inch on its sternal side.

Præcordial Region.—The “*præcordial region*” is the outline of the heart traced upon the front wall of the chest. We ought to be able to define this region with something like precision. We should know how much of this region is covered and separated from the wall of the chest by intervening lung (fig. 17. p. 97). Here then is the rule:—Let the middle of the fifth costal cartilage be the centre of a circle two inches in diameter: this circle will define well enough for

all practical purposes, that part of the præcordial region which is naturally less resonant to percussion than the rest ; for here the heart is uncovered except by pericardium and loose cellular tissue, and is very near the wall of the chest. In the rest of the præcordial region the heart is covered and separated from the chest by the intervening lung.

Fig. 17.



FORM OF THE LUNGS, AND THE EXTENT TO WHICH THEY OVERLAP THE HEART AND ITS VALVES.

Where must we put the stethoscope when we listen to the valves of the heart? For practical purposes it is enough to remember that the mouth of an ordinary-sized stethoscope will cover a portion of them all, if it be placed a little to the left of the mesial line of the sternum opposite the third intercostal space (fig. 16, p. 95). They are all covered by a thin portion of lung; for this reason we ask a patient to stop breathing while we listen to his heart.

The position of the heart alters a little with the position of the body. Of this any one may convince himself by leaning alternately forwards and backwards, by lying on this side and on that, placing at the same time the hand upon the præcordial region. He will find that he can, in a slight degree, alter the place and the extent of the impulse of the heart; and the reason is obvious. Inspiration and expiration also alter the position of the heart. In inspiration the heart descends with the tendinous centre of the diaphragm about half an inch.

Pericardium.—The pericardium is the membranous bag which encloses the heart and the large vessels at its base. It is attached below to the tendinous centre of the diaphragm; above, it is lost on the great vessels of the heart, and is connected with the cervical fascia. On each side it is covered by the pleura. In front of it is the anterior mediastinum; behind it, the posterior. Of all the objects in the posterior mediastinum, that which is nearest to the pericardium is the œsophagus. It is worth remembering that the œsophagus is in close contact with the back of the pericardium for nearly two inches; this fact accounts for what one sometimes observes in cases of pericarditis where there is much effusion; I mean pain and difficulty in swallowing.

The pericardium is what is called a "*fibro serous*" membrane. Its fibrous layer, which constitutes the chief strength of it, is external. This layer, as just now stated, is attached below to the central tendon of the diaphragm. Above it forms eight sheaths for the vessels at the base of the heart; namely, one for the vena cava superior, four for the pulmonary veins, two for the pulmonary arteries, and one for the aorta. The *serous* layer forms a shut sac. It lines the fibrous layer, and is reflected over the great vessels and the heart. The best way to see where the serous layer is reflected over the vessels, is to distend the pericardium with air. Thus you will find that this layer is reflected over the aorta as high as the origin of the arteria innominata. It is reflected over the vena cava superior, after the entrance of the vena azygos.*

* Those who choose to follow the reflections of the serous layer of the pericardium, will find that it covers the great vessels to an extent greater than is generally imagined; though in truth the extent is not precisely similar in all bodies. The aorta and pulmonary artery are wrapped in a complete sheath

How much larger is the pericardium than the heart? In the healthy state the capacity of the pericardium pretty nearly corresponds to the size of the heart when distended to its utmost. The healthy pericardium, with the heart *in situ*, may be made to hold, in the adult, about ten ounces of fluid. The pericardium is not extensible. When an aneurism bursts into it, death is caused, not by the loss of blood, but by compression of the heart in consequence of the inextensibility of the pericardium.

[The pericardium has been known to be entirely absent.]

The pericardium derives its blood chiefly from the internal mammary arteries. Hence the direct relief afforded by local abstraction of blood from the præcordial region.

Position and Form of the Lungs.—The lungs are situated in the chest, one on each side of the heart. Each fits accurately into the cavity which contains it. Each, therefore, is conical in form; the base rests on the diaphragm; the apex projects in the neck a little above the sternal end of the clavicle. Its outer surface is adapted to the ribs; its inner surface is excavated to make room for the heart. But the best way to see the shape of the lungs is to inject the trachea with wax, which is tantamount to taking a cast of each thoracic cavity. In such a preparation, besides the general convexities and concavities alluded to, you would find in the right lung a little indentation for the right brachio-cephalic vein, in the left an indentation for the arch of the aorta and the left subclavian artery.

Each lung is divided into an upper and a lower lobe by a deep fissure, which commences behind about three inches

two inches in length, so that these vessels are covered all round by the serous layer, except where they are in contact. Indeed you can pass your finger behind them both, through a foramen bounded in front, by the two great vessels themselves, behind, by the upper part of the auricles, and above by the right pulmonary artery. Again, the back of the aorta, where it lies on the auricles, is covered by the serous pericardium. The superior cava is covered all round, except behind where it crosses the right pulmonary artery. What little there is of the inferior cava within the pericardium is also covered all round. The left pulmonary veins are covered nearly all round; the right less so. Behind the auricles, chiefly the left, the serous layer extends upwards in the form of a pouch, rising above their upper border, so as to be loosely connected to the left bronchus.

from the apex, and proceeds obliquely downwards to the front a little lower than the fifth costal cartilage. The upper lobe of the right lung is divided by a second fissure, which slices off a triangular portion, called the "middle lobe."

The dimensions of the right lung are greater than those of the left in all directions except the vertical; the reason of this exception is the greater elevation of the diaphragm on the right side, by the bulk of the liver. On an average, the right lung is to the left, in point of size, as 11 to 10.

To understand rightly the shape and play of the lungs in inspiration and expiration, cut away the intercostal muscles and the pleura, without wounding the lungs, and then distend the lungs by blowing into the trachea with bellows. By this proceeding you will learn much that is worth knowing. Thus, you will ascertain how far the front edge of the lung overlaps the pericardium; how low the lung descends between the ribs and the diaphragm on the side and at the back of the chest. You will see the great gap in the left lung for the point of the heart. By making the lung expand and contract, you will observe how it slides along the pleural lining of the chest. This sliding takes place in health without the least noise. But when the naturally smooth and polished surface of the pleura becomes roughened by inflammatory deposit, a sound of greater or less distinctness (friction sound) is occasioned by the rubbing of the opposite surfaces.

The practical result of this investigation should be to enable us to trace upon a living chest the outline of the lungs, that we may know what parts are naturally resonant on percussion.

Commencing, then, from above, (fig. 17, p. 97), we find that the apex of the lung rises into the neck from half an inch to an inch above the sternal end of the clavicle. This part of the lung mounts up behind the subclavian artery and the anterior scalene muscle, and deserves especial attention, because it is, more than any other, obnoxious to tubercular disease. From the sternal end of the clavicles the lungs converge towards the mesial line, where their edges almost meet opposite the junction of the second rib. There is little or no lung behind the first bone of the sternum.

From the level of the second costal cartilage to the level of

the fourth, the inner margins of each lung run parallel and almost close behind the middle of the sternum; consequently they overlap the great vessels at the root of the heart.

Below the level of the fourth costal cartilage the margins of the lungs diverge from each other, but not in equal degree. The *left* presents the notch for the heart, and curves pretty nearly in the course of the fourth costal cartilage; at the lower part of its curve it projects more or less over the apex of the heart. The *right* descends almost perpendicularly behind the sternum as low as the attachment of the ensiform cartilage, and then, turning outwards, corresponds with the direction of the sixth costal cartilage. Hypertrophy of the heart, or effusion into the pericardium, will not only raise above the ordinary level the point where the lungs diverge, but also increase their divergence; hence the greater dulness on percussion.

Laterally, the margin of the lung comes down as low as the eighth rib; posteriorly, as low as the tenth rib.

Pleura.—As the lungs are continually gliding to and fro, within the chest, they are provided with a serous membrane to facilitate their motion. This membrane is termed the pleura. There is one for each lung. Each pleura forms a completely closed sac, and is disposed like all other serous sacs, that is, one part of the sac lines the containing cavity, while the other is reflected over the contained organ. Its several parts are named after the surface to which they adhere: that which lines the ribs is called “pleura costalis;” that which forms the mediastinum, “pleura mediastinalis;” that which covers the lung, “pleura pulmonalis.” Unlike the peritoneum, the pleura forms no folds except a small one called “ligamentum latum pulmonis” which extends from the root of the lung to the diaphragm.

If asked to divide the reflections of the plural sac, (fig. 14, p. 87), I should say, that it lines the ribs and part of the sternum; from the sternum it is reflected backwards over the pericardium, from thence it passes over the front of the root of the lung, and so on over the entire lung itself, to the back part of its root, whence it is reflected over the sides of the vertebræ, and thus reaches the ribs again.

[The late Dr. C. E. Isaacs in a paper read before the New

York Academy of Medicine, April 1st, 1867, on the Extent of the Pleura above the Clavicle, states, that in one hundred cases, the pleura rose two inches and upwards above the clavicle, in fourteen cases on the right side, in four on the left, and on both sides in five—in all, twenty-three cases—thus giving, in one hundred, the proportion of nearly one in four. In referring to the low extent of the pleura as occasionally met with, he says, that there were only five cases out of one hundred in which the pleura did not rise above the superior margin of the clavicle. In one case the pleura on the right side was of the usual height, but on the left it only rose a little above the first rib. The sternal edge of the pleura on the right side extended across and to the left of the median line, in eleven cases to the distance of about an inch. The average width of the pleura in these eleven cases, was four inches. This fact is important in reference to the surgical relations of the vessels, and in removal of tumors from this region of the neck; as one in every nine presented the peculiarity of great width of the pleura. These facts are also of interest to the physician, as the apex of the lungs corresponds with the concavity of the pleural sac above the clavicle. Dr. Isaacs has also noticed that the pleura rose highest in those having long necks. We should bear in mind that the first portion of the subclavian always rests on the pleura, while the second portion of the vessel is always in contact with this membrane; and when the pleura rises high in the neck above the clavicle, it carries with it the third portion of the artery, though usually the vessel in this portion of its course does not rest upon the pleural sac. This paper goes on to state, that in twenty-three cases the subclavian artery was high in the neck, and in many of them, the artery in its third portion rested upon the pleura for a considerable extent. 1. In one case in which the pleura rose very high in the neck, one inch and a quarter of the right carotid, three-fourths of an inch of the left carotid, and one inch and a quarter of the third portion of the right subclavian, extending from the outer margin of the scalenus anticus to the upper edge of the first rib, rested upon the pleura. 2. The third portion of the right subclavian, to the extent of one inch was in contact with the pleura. 3. The third portion of the right subclavian to the extent of one inch and five-eighths,

rested upon the pleura. 4. The third portion of the right subclavian was in contact with the pleura to the extent of one inch and a quarter. 5. The third portion of the left subclavian to the extent of one inch and five-eighths, rested upon the pleura. The top of this pleural cavity does not always present a regular or dome-like form. Sometimes it forms cul-de-sacs, which extend upwards, like the fingers of a glove, into the recesses of the neck. These may also be prolonged laterally from the right pleura across the median line to the left side, and pass upwards either between the bodies of the cervical vertebræ and the œsophagus, or between the œsophagus and the trachea.]

The spaces called "anterior" and "posterior" mediastina formed by the pleuræ, I have already described, p. 87.

In health the internal surface of the pleura is smooth, polished, and lubricated by moisture sufficient to facilitate the sliding of the lung. When the surface is thickened and roughened by inflammation, the moving lung produces a "friction" sound. When the pleural sac is distended by serum, it constitutes hydro-thorax; when by pus, "empyema."

Take an opportunity of introducing your hand into the pleural sac, and ascertain that the reflection of the pleura on to the diaphragm corresponds with an imaginary line commencing at the lower part of the sternum, and sloping along the cartilages of the successive ribs down to the lower border of the last rib. Now suppose a musket ball were to lodge in the pleural sac, it would fall upon the dome of the diaphragm, and roll down to the lowest part of the pleural cavity. The place, therefore, to extract it would be in the back, at the eleventh intercostal space. This operation has been done during life with success.

Posterior Mediastinum and its Contents.—The posterior mediastinum (p. 88) is formed by the reflection of the pleural sac on each side, from the root of the lung to the sides of the bodies of the dorsal vertebræ. It is bounded in front by the pericardium. To obtain a view of it, draw the right lung out, and fasten it to the left side. This mediastinum contains the aorta; in front of the aorta, the œsophagus, with the pneumogastric nerves; on the right of the aorta is the vena azygos; between this vein and the aorta is the thoracic duct. To ex-

pose all these we have only to remove the pleura, and a layer of tolerably firm fascia which lines the chest outside the pleura.

Thoracic Aorta.—We have already traced the arch of the aorta to the left side of the body of the third dorsal vertebra. From this point the aorta descends on the left side of the

Fig. 18.




DIAGRAM TO SHOW THE COURSE OF THE VENA AZYGOS AND THE THORACIC DUCT.

spine, gradually approaching towards the mesial line. Opposite the last dorsal vertebra it passes between the crura of the diaphragm and enters the abdomen. Its left side is covered by the pleura: on its right side run the vena azygos and thoracic duct; in front of it, and nearer to the mesial line, is the œsophagus. Its branches will be described presently.

Vena Azygos.—This vein commences in the abdomen by small branches from one of the lumbar veins, and generally communicates with the renal, or even the vena cava itself. This, indeed, is the main point about the origin of the vena azygos, that it communicates directly or indirectly with the vena cava inferior. It enters the chest through the aortic opening of the diaphragm, and ascends on the right side of the aorta through the posterior mediastinum. When the vein reaches the level of the third dorsal vertebra,

it arches over the right bronchus, and terminates in the superior vena cava just before this vessel is covered by pericardium. In its course it receives nine or ten of the lower

intercostal, the spinal veins, the œsophageal, and commonly the right bronchial veins.

[The azygos major has been found to terminate in the subclavian, though rarely.]

But another vein, called "*vena azygos minor*," runs up the left side of the spine. This vein commences in the abdomen by small branches communicating with the inferior cava, and ascends on the left side of the aorta. On a level with the sixth or seventh dorsal vertebra, it passes beneath the aorta and joins the azygos major. It receives five or six of the lower intercostal veins of the left side. None of these veins are provided with valves.

What is the purpose of the "azygos" vein? The answer is, it is supplemental to the vena cava.

Thoracic Duct.—The thoracic duct (fig. 18) is the canal through which the contents of the lacteal vessels from the intestines and the absorbent vessels from the lower limbs are conveyed into the blood. These vessels converge to a general receptacle, termed *receptaculum chyli*, situated in front of the body of the second lumbar vertebra. From this receptacle, the duct passes through the aortic opening of the diaphragm into the chest, and runs up the posterior mediastinum along the right side of the aorta. Near the *third* dorsal vertebra, it passes under the œsophagus, and ascends on the left side of this tube between it and the left subclavian artery as high as the level of the seventh cervical vertebra, where it describes a curve with the convexity upwards, and opens into the back part of the confluence of the left internal jugular and subclavian veins. The orifice of the duct is guarded by two valves, which permit fluid to pass from the duct into the vein, but not vice versâ. Valves too, disposed like those in the venous system, are placed at short intervals along the duct, so that its contents can only pass one way. The duct is not much larger than a crow-quill: its walls are thin and transparent.*

Œsophagus.—The "œsophagus" is that part of the ali-

* It is right to state that the thoracic duct varies in size in different individuals. I have seen it of all sizes, intermediate between a crow-quill and a goose-quill. It may divide in its course into two branches, which subsequently reunite; instead of one, there may be several terminal orifices. Instances

mentary canal which conveys the food from the pharynx to the stomach. It commences about the fifth cervical vertebra, that is, nearly opposite the cricoid cartilage, runs down the posterior mediastinum in front of the descending aorta, and passes through a special opening in the diaphragm to the stomach. It is from nine to ten inches long. Its course is not exactly vertical; in the neck, it lies to the left of the trachea; in the chest, *i. e.* about the fourth dorsal vertebra, it inclines towards the right side, to make way for the aorta; but again inclines to the left before it perforates the diaphragm.

Observe that the œsophagus, before it passes through the diaphragm, lies in close contact with the pericardium for nearly two inches: this fact accounts for the pain which is sometimes experienced in cases of pericarditis, during the passage of the food.

The œsophagus is supplied with blood by the inferior thyroid, the œsophageal branches of the aorta, and the coronaria ventriculi. It is supplied with nerves by the pneumogastric and the sympathetic.

The œsophagus is composed of two coats, a muscular and a mucous; and these are very loosely connected together by firm areolar tissue, which some anatomists describe as a distinct coat. The muscular coat consists of an outer longitudinal, and an inner circular layer of fibres. Both are of the non-striped variety. The longitudinal layer is particularly strong, and arranged all round the œsophagus so as to support the circular. The mucous membrane is of a pale color, and considerable thickness, and in the contracted state of the œsophagus is arranged in longitudinal folds. It is protected by a very thick layer of scaly epithelium. In the submucous tissue are many small glands, especially towards the lower end of the œsophagus.

Course and Branches of the Pneumogastric Nerves.—

The *right* pneumogastric nerve (fig. 6) enters the chest between the subclavian artery and vein, descends by the side of the trachea, then behind the right bronchus to the posterior surface of the œsophagus, upon which it divides into branches,

have been observed in which the duct has terminated on the right instead of on the left side (Fleischman, *Leichenöffnungen*, 1815). It has been seen to terminate in the vena azygos (Müller's Archives, 1834).

which form a plexus (posterior œsophageal) upon the tube. The *left* pneumogastric descends into the chest between the left subclavian and the left carotid arteries, crosses in front of the arch of the aorta, then passes behind the left bronchus to the anterior surface of the œsophagus, upon which it also forms a plexus (anterior œsophageal). The branches of the pneumogastric nerve in the chest are as follow :—

a. The inferior laryngeal or recurrent.—This nerve on the right side turns under the subclavian artery, on the left under the arch of the aorta, and ascends to the larynx. It supplies with motor power all the muscles which act upon the rima glottidis except the crico-thyroid (supplied by the superior laryngeal nerve).

b. Cardiac branches.—These are very small, and join the cardiac plexus.

c. Pulmonary branches.—These accompany the bronchial tubes. The greater number are seen behind the root of the lung, forming the posterior pulmonary plexus. A few, forming the anterior pulmonary plexus, proceed over the front of the lung's root. Both these plexus are joined by filaments from the sympathetic system. But the nerves of the lungs are very small, and cannot be traced far into their substance.*

d. Œsophageal plexus.—We have already mentioned that below the root of the lung each pneumogastric nerve is subdivided so as to form an interlacement of nerves round the œsophagus (plexus gulæ). From this plexus numerous filaments supply the coats of the tube ; but the majority of them are collected into two or more nerves, which pass with the œsophagus through the diaphragm for the supply of the stomach.

Having examined the contents of the posterior mediastinum from the right side, now do so from the left. Turn the left lung out of its cavity, and fasten it towards the right side. After removing the pleura, we see the descending thoracic aorta, the pneumogastric nerve crossing the arch and sending the recurrent branch through it ; also the first part of the course of the left subclavian, covered externally by pleura. Trace the pneumogastric nerve behind the left bronchus to the œsophagus, and dissect the œsophageal plexus on this

* Upon this subject, see the beautiful plates of Scarpa.

side. Lastly, notice the lesser vena azygos, which crosses under the aorta about the sixth or seventh dorsal vertebra, and joins the vena azygos major.

Thoracic Portion of the Sympathetic.—There are generally twelve ganglia in the thoracic portion of the sympathetic, one over the head of each

rib. But often, you find only ten ganglia, because one or two of them have fused together. The first thoracic ganglion is the largest.

Each ganglion receives branches from the corresponding spinal nerve. The nerves proceeding from the ganglia supply the thoracic and part of the abdominal viscera. They are as follow:— (see the diagram).

a. Minute nerves from the first and second ganglia to the *cardiac plexus*.

b. Minute nerves from the third and fourth ganglia to the *pulmonary plexus*.

c. The *greater splanchnic nerve*. This is generally formed by branches from the sixth to the tenth ganglion. The branches descend obliquely along the sides of the bodies of the vertebræ, and unite into a single nerve, which passes through the corresponding crust of the diaphragm, and joins the




DIAGRAM OF THE THORACIC PORTION OF THE SYMPATHETIC.

semilunar ganglion of the abdomen.

d. The *lesser splanchnic nerve*.—This is commonly formed by branches from the eleventh and twelfth ganglia. It passes through the crust of the diaphragm to the renal plexus.*

* In a few instances we have traced a minute filament from one of the ganglia into the body of the vertebra. According to a celebrated French anatomist (Cruveilhier), each vertebra receives one.

Intercostal Muscles.—The intercostal muscles occupy the intervals between the ribs. Between each rib there are two layers of muscles which cross each other like the letter X. The external layer runs obliquely from behind, forwards, like the external oblique muscle of the abdomen. The internal layer runs from before backwards, like the internal oblique. Observe that a few fibres of the inner layer pass over one or even two ribs, and terminate upon a rib lower down.

But neither of these layers extend all the way between the sternum and the spine : for the outer layer, beginning at the spine, ceases at the cartilages of the ribs ; while the inner, commencing at the sternum, ceases at the angles of the ribs.

The intercostal muscles present a curious intermixture of tendinous and fleshy fibres ; and they are covered both inside and outside the chest by a glistening fascia, to give greater protection to the intercostal spaces.

The external intercostal muscles elevate the ribs, and are therefore muscles of inspiration. The internal intercostal muscles depress the ribs, and are therefore muscles of expiration.

Intercostal Arteries.—There are twelve intercostal arteries on each side. The two upper are supplied by the intercostal branch of the subclavian ; the remaining ten are supplied by the aorta : and since this vessel lies rather on the left side of the spine, the right intercostal arteries are longer than the left. The upper intercostal arteries from the aorta ascend obliquely to reach their intercostal spaces ; the lower run more transversely. Having reached the intercostal space, each artery divides into an *anterior* and a *posterior* branch.

The *anterior* branch, in direction and size, appears to be the continuation of the common trunk. At first it runs *along the middle of the intercostal space*, and is separated from the cavity of the chest by the pleura and intercostal fascia. Here, therefore, it is liable to be injured by a wound in the back. But near the angle of the rib it passes between the intercostal muscles, and occupies the groove in the lower border of the rib. Here, too, it gives off a branch, which runs for some distance along the upper border of the rib below, and is lost in the muscles. In some cases I have seen this branch as

large as the intercostal itself, and situated so as to be directly exposed to injury in the operation of tapping the chest.

In its course along the intercostal space, each artery sends branches to the intercostal muscles and the ribs. About midway between the sternum and the spine, each gives off a small branch, which accompanies the lateral cutaneous branch of the intercostal nerve. The continued trunk, gradually decreasing in size, becomes very small towards the anterior part of its space, and is placed more in the middle of it. Those of

Fig. 20.




DIAGRAM OF A SPINAL NERVE.

the true intercostal spaces inosculate with branches of the internal mammary: those of the false are lost between the layers of the abdominal muscles.

The *posterior* branch passes backwards between the transverse processes of the vertebræ, to the muscles and skin of the back. Each sends an artery through the intervertebral foramen to the spinal cord and its membranes.

The intercostal veins terminate in the vena azygos.

Intercostal Nerves.—These are the anterior divisions of the spinal nerves. Remember that each spinal nerve arises from the spinal cord by two roots—an anterior or motor root, a posterior or sensitive root. The sensitive root has a ganglion upon it. The two roots unite in the intervertebral foramen, and form a *compound* nerve. After passing through the foramen, this nerve then divides into an anterior and a posterior branch. The anterior branch proceeds along the intercostal space, in company with and immediately below the corresponding arteries. In its course it supplies the intercostal muscles; also a lateral cutaneous and an anterior cutaneous branch.

Observe that the first dorsal nerve, after furnishing a nerve to the first intercostal space, ascends perpendicularly over the neck of the first rib, and contributes to form the brachial plexus.

Intercostal absorbent glands.—These are situated near the heads of the ribs ; there are some between the layers of the intercostal muscles. They are of small size, and send their absorbent vessels into the thoracic duct. I have seen these intercostal glands enlarged and diseased in phthisis.

Bronchial and Œsophageal Arteries.—Small arteries, arising from the concavity of the arch of the aorta, accompany the bronchial tube into the substance of the lung. Their distribution and office will be considered with the anatomy of the lung. Other arteries proceed from the front of the descending aorta to supply the Œsophagus.

Having finished the posterior mediastinum, replace the lung, and turn your attention once more to the great vessels at the root of the heart.

Pulmonary Artery.—This vessel carries the impure blood from the heart to the lungs. It proceeds from the right ventricle, crosses obliquely in front of the root of the aorta, and on the left side of that vessel divides into two branches, one for each lung. The right branch passes through the arch of the aorta to the right lung : the left is easily followed to its corresponding lung by removing its investing layer of pericardium.

Draw towards the left side the first part of the arch of the aorta, and dissect the pericardium from the great vessels at the base of the heart. Thus a good view will be obtained of the trachea and its bifurcation into the two bronchi. Below the division of the trachea the right pulmonary artery is seen passing in front of the right bronchus. The superior vena cava is seen descending in front of, and nearly at right angles to, the right pulmonary artery. The vena azygos is also seen arching over the right bronchus and terminating in the vena cava. Notice, especially, a number of absorbent glands called "*bronchial*," at the angle of bifurcation of the trachea. The situation of these glands in the midst of so many tubes explains the variety of symptoms which may be produced by their enlargement.

Cardiac Plexus of Nerves.—The nerves of the heart are derived from the pneumogastric and from the cervical ganglia of the sympathetic. A general description of them will suffice. The cardiac nerves of the right side descend chiefly

behind the arch of the aorta; those of the right chiefly in front of the arch. The nerves from both sides, however, converge towards the posterior part of the arch of the aorta. Here they form a plexus called the "*great cardiac*," and in it we recognise two or more ganglia. The precise situation of this plexus is under the arch of the aorta above the bifurcation of the pulmonary artery.

From this plexus the nerves proceed, in company with the

Fig 2




DIAGRAM SHOWING THE CONSTITUENTS OF THE ROOT OF EACH LUNG, AND THEIR RELATIVE POSITION: ALSO THE POSITION OF THE VALVES OF THE HEART. THE ARROWS INDICATE THE DIRECTION IN WHICH AORTIC AND MITRAL MURMURS ARE PROPAGATED.

coronary arteries, to the heart. Those which accompany the anterior coronary artery form the "*anterior coronary plexus*." The "*posterior coronary plexus*" proceeds with its artery to the posterior part of the heart.

But it is not an easy matter to trace the nerves into the substance of the heart. For this purpose a horse's heart is the best, and previous maceration in water is desirable. The nerves in the substance of the heart are peculiar in this respect; that they present minute ganglia in their course, which are presumed to preside over the rhythmical contractions of the heart.

Constituents of the Root of each Lung.—Draw aside the margin of the right lung; divide the superior vena cava above the vena azygos, and turn down the lower part. Remove the layer of pericardium which covers the pulmonary veins, and the constituent parts of the root of the right lung will be exposed. They lie from before backwards in the following order:—In front are the two pulmonary veins; behind the veins are the subdivisions of the pulmonary artery; and behind the artery are the divisions of the bronchus. From above downwards they are disposed thus:—On the right side we find—1st, the bronchus; 2nd, the artery; 3rd, the veins. On the left, we find:—1st, the artery; 2nd, the bronchus; 3rd, the veins, as shown in fig. 21.

DISSECTION OF THE HEART.

It is presumed that you are familiar with the general form of the heart, described at page 95. You should notice the longitudinal grooves on the upper and lower surfaces of the heart, indicating the divisions of the ventricles; and the circular groove near the base, indicating the separation between the ventricles and auricles. These grooves are occupied by the coronary vessels, and by more or less fat.

The heart is a double organ; that is, it is composed of two hearts, a right and a left, separated by a septum. Each consists of an auricle and a ventricle, which communicate by a wide orifice: the right heart propels the blood through the lungs, and is called the *pulmonic*; the left propels the blood through the body, and is called the *systemic*. True, these two hearts are not placed apart, because important advantages result from their union; by being enclosed in a single bag they occupy less room in the chest; and the action of their corres-

ponding cavities being precisely synchronous, their fibres mutually intermixing, contribute to their mutual support. We shall examine the cavities of the heart in the order in which the blood circulates through them.

Right Auricle.—This is situated at the right side of the base of the heart, and forms a quadrangular cavity between the two venæ cavæ, from which it receives the blood. From its front a small pouch projects towards the left, and overlaps the root of the aorta: this part is termed the appendix of the auricle, and resembles a dog's ear in shape: *unde nomen*.

To see the interior, make a horizontal incision through the anterior wall, from the apex of the appendage transversely

Fig. 22.




DIAGRAM OF THE INTERIOR OF THE RIGHT AURICLE.

across the cavity; from this make another upwards at right angles into the superior cava. Observe that the interior is lined by a polished membrane, and that it is everywhere flat, except in the appendix, where the muscular fibres are collected into bundles, called, from their resemblance to the teeth of a comb, "*musculi pectinati*." They radiate from the auricles to the edges of the auriculo-ventricular opening. Observe the openings of the two venæ cavæ: they are not directly opposite to each other: the superior is situated on a plane rather in front of the inferior, that the streams of blood may not meet. The inferior cava, after passing through the tendinous centre of the diaphragm, makes a slight curve to the left be-

fore it opens into the auricle: and the reason of this is that the stream of its blood may be directed towards the auriculo-ventricular opening. The orifice of each vena cava is nearly circular, and is surrounded by a few muscular fibres continuous with those of the auricle.

[Between the openings of the vena cava, the wall of the auricle projects a little, forming a slight elevation in some hearts, called the tubercle of Lower.]

The left wall of the auricle is formed by the "*septum auricularum*." Upon this septum, above the orifice of the vena cava inferior, is an oval depression (*fossa ovalis*), bounded by a prominent border (*annulus ovalis*), (fig. 22). This depression marks the position of the opening (*foramen ovale*), through which the auricles communicated in foetal life. After birth this opening closes; but if the closure is imperfect, the stream of dark blood in the right auricle mixes with the florid blood in the left, and occasions what is called "*morbis cæruleus*."

Extending from the anterior margin of the vena cava inferior to the anterior border of the fossa ovalis, is seen in young subjects a thin fold of the lining membrane of the heart: it is the remains of what was, in foetal life, the Eustachian* valve. The direction of this valve in the foetus is such that it directs the current of blood from the inferior cava towards the foramen ovale. It is a valve of considerable size in the foetus, and contains a few muscular fibres; but after birth it gradually disappears, being no longer required. To the left of this—that is, between the remains of the Eustachian valve and the auriculo-ventricular opening—is the orifice of the coronary vein; it is covered by a semicircular valve, called "*valvula Thebesii*." Here and there upon the posterior wall of the auricle may be observed minute openings called "*foramina Thebesii*": they are the orifices of small veins returning blood from the substance of the heart. Lastly, to the left and rather in front of the orifice of the vena cava inferior is the auriculo-ventricular opening. It is oval in form, and admits the passage of three fingers.

Right Ventricle.—This forms the right border and about two-thirds of the upper surface of the heart. To examine

* Eustachius, Libell. de vena sine pari.

its interior, a triangular flap should be raised from its anterior wall. The apex of this flap should be below ; one cut along the right edge of the ventricle, the other along the line of the ventricular septum. We observe that the wall of the ventricle is much thicker than that of the auricle. The cavity of the ventricle is conical, with its base upwards and to the right. Its walls present bands of muscular fibres (*columnæ carneæ*) of various length and thickness, which cross each other in every direction ; this muscular network is generally filled with coagulated blood. Of these *columnæ carneæ* there are three kinds :—one stands out in high relief from the ventricle : another is attached to the ventricle by their extremities only ; a third, and by far the most important set, called “*musculi papillares*” is fixed by one extremity to the ventricle, while the other extremity gives attachment to the fine tendinous cords (*cordæ tendineæ*) which regulate the action of the tricuspid valve. The number of these *musculi papillares* is equal to the number of the chief divisions of the valve ; consequently there are three in the right, and two in the left ventricle. Of those in the right ventricle, one proceeds from the septum.

There are two openings in the right ventricle. One, the *auriculo-ventricular*, through which the blood passes from the auricle, is placed at the base of the ventricle. It is surrounded by a ring of fibrous tissue, to which is attached the tricuspid valve. From the upper and front part of the ventricle, a smooth passage (“*infundibulum*”) leads to the other opening—that of the pulmonary artery. It is situated to the left and in front of the *auriculo-ventricular*, and about three-fourths of an inch higher.

Tricuspid Valve.—This is situated at the right *auriculo-ventricular* opening. Like all the valves of the heart, it is formed by a fold of the lining membrane “*endocardium*” of the heart, strengthened by fibrous tissue. The base of the valve is attached to the tendinous ring round the opening. Its free border presents three principal triangular flaps ; and besides these, intermediate flaps of smaller size. Of the three principal flaps, the largest is so placed that when not in action it partially covers the orifice of the pulmonary artery.

Observe the arrangement of the tendinous cords which regu-

late the action of the valve. In the first place they are all attached to the ventricular surface of the valve. Secondly, the tendinous cords proceeding from a given papillary muscle, are attached to the adjacent halves of two of the larger flaps, and to a smaller intermediate one; consequently, when the ventricle contracts, and the papillary muscle also, the adjacent borders of the flaps will be approximated. Thirdly, to insure the strength of every part of the valve, the tendinous cords are inserted at three different points of it in straight lines; accordingly, they are divisible into three sets. Those of the first are attached to the base of the valve; those of the second to the middle of it; those of the third to its loose border.*

Pulmonary or Semilunar Valves.—These are three membranous folds situated at the orifice of the pulmonary artery. Their convex borders are attached to the fibrous ring at the root of the artery; their free edges present a festooned border, in the centre of which is a little cartilaginous body called the “nodulus,” or “*corpus Arantii*.” The use of these little bodies is very obvious. Since the valves are semilunar, when they fall together they would not exactly close the artery; there would be a space of a triangular form left between them in the centre, just as there is when we put the thumb, fore, and middle fingers together. The space is filled by the little bits of cartilage, so that the septum becomes complete.

The valves are composed of a fold of the “*endocardium*,” or lining membrane of the heart. But since this would not

* The best mode of showing the action of the valve is to introduce a glass tube into the pulmonary artery, and then to pour water through it into the ventricle until the cavity is quite distended. By gently squeezing the ventricle in the hand, so as artificially to imitate its natural contraction, the tricuspid valve will flap back like a flood-gate, and close the auriculo-ventricular opening. In this way one can understand how, when the ventricle contracts, the blood catches the margin of the valve, and by its pressure gives it the proper distension and figure requisite to block up the aperture into the auricle. It is obvious that the tendinous cords will prevent the valve from being pushed too far back into the auricle; and this purpose is assisted by the papillary muscles, which nicely adjust the degree of tension of the cords at a time when they would otherwise be too much slackened by the contraction of the ventricle.

of itself be sufficiently strong, we find between the folds a thin layer of fibrous tissue, which is prolonged from the tendinous ring at the orifice of the artery. Observe, however, that this layer of fibrous tissue reaches the free edge of the valve at three points only; namely, at the centre, or corpus Arantii, and at each extremity. Between these points it stops short, and leaves a crescent-shaped portion of the valve thinner than the rest, and consisting simply of endocardium. This crescent-shaped portion (called the "*lunula*") is not wholly without fibrous tissue, for a thin tendinous cord runs along its free edge, to give it additional strength to resist the impulse of the blood; just as sails are strengthened by cords along their edges, that they may not be torn by the wind. Behind each of the valves, the artery bulges, and forms three slight dilatations called the "*sinuses of Valsalva*." These, we shall presently see, are more marked at the orifice of the aorta.

The action of these valves is obvious. During the contraction of the ventricle, the valves lie against the side of the artery, and offer no impediment to the current of blood; during its dilatation, the elasticity of the distended artery would force back the column of blood, but that the valves, being caught by the reflux blood, bag, and fall together so as to close the tube. The greater the pressure, the more accurate is the closure. The coats of the artery are very elastic and yielding, while the valve, as well as the circumference to which it is attached, is quite unyielding; consequently, when the artery is distended by the impulse of the blood, its wall is removed from the contact of the free margin of the valves, and these are more readily caught by the retrograde motion of the blood. The force of the reflux is sustained by the tendinous part of the valves, and also by the muscular wall of the ventricle, as shown by Mr. Savory. According to Haller, the valves are capable of sustaining a weight of sixty-three pounds before they give way. The thinner portions (the *lunule*) become placed so as to lie side by side, each one with that of the adjacent valve. All this may be seen by filling the artery with water.

Left Auricle.—This is situated at the left side, and posterior part of the base of the heart. It is somewhat quadrilateral, and receives the pulmonary veins, two on either side,

which return the purified blood from the lungs. From its upper and left side the auricular appendage projects towards the right, curling over the root of the pulmonary artery. The auricle should be opened by a horizontal incision proceeding from one pulmonary vein to another: from this a second should be made into the appendix. Its interior is smooth and flat, excepting in the appendix, which contains the "*musculi pectinati*." Notice the openings of the four pulmonary veins. Upon the septum between the auricles is a depression indicating the remains of the foramen ovale. At the lower and front part of the auricle is the *auriculo-ventricular opening*. It is oval, with its long axis nearly transverse, and in the adult admits the passage of two fingers.

Left Ventricle.—This occupies the left border, and forms the apex of the heart. One third of it only is seen on the upper surface. To examine the interior, raise a triangular flap from its front wall, with the apex of the flap below. Then observe that it is about three times as thick as that of the right ventricle, and that this thickness gradually diminishes towards the apex. The interior of the left ventricle is so much like that of the right, that one need not describe it in detail. Observe that the *auriculo-ventricular valve* consists of only two principal flaps: hence its name "*mitral*," or "*bicuspid*." The largest of these flaps is placed between the aortic and auricular orifices. There are only two "*musculi papillares*;" one attached to the anterior, the other to the posterior surface of the ventricle. They are thicker, and their "*cordæ tendineæ*" stronger than those of the right ventricle, but their plan is precisely similar. From the upper and back part of the ventricle, a smooth passage leads to the orifice of the aorta. The orifice is placed rather in front and to the right side of the auriculo-ventricular opening; but the two orifices are close together, and only separated by the larger flap of the mitral valve. The aortic orifice is guarded by three semilunar valves, of which the arrangement, structure, and mode of action are precisely like those of the pulmonary artery. Their framework is proportionately stronger, consistently with the greater strength of the left ventricle and the greater impulse of the blood. In the "*sinuses of Valsalva*" are observed the orifices of the coronary arteries.

At the openings between the auricles and ventricles, and also at the commencement of the aorta and pulmonary artery, we find fibrous rings. These rings serve as fixed points for the attachment of the muscular fibres of the heart.

How the Great Arteries are attached to the Ventricles.—

The fibrous rings at the arterial orifices present three festoons, with concavities directed upwards. These festoons give attachment to the middle coat of the artery above, to the muscular fibres of the heart below, and internally, to the tendinous fibres of the valves. The vessels are also connected to the heart by the serous layer of the pericardium, and by a continuation of the lining membrane (endocardium) of the ventricle.

Heart, how Supplied with Blood.—The heart is supplied with blood by the two coronary arteries, a right or posterior, and a left or anterior. Both arise from the aorta just above the semilunar valves, and at such a distance as always to admit the passage of blood: both run in the furrows between the ventricles and auricles: both are accompanied by branches of the coronary vein, and by the cardiac nerves.

The anterior or left coronary artery arises from the left side of the aorta. It appears on the left of the pulmonary artery, and then runs down the inter-ventricle furrow on the anterior surface of the heart to the apex, where it inosculates with the posterior coronary artery. In this course, its principal branch is that which turns to the left, along the furrow between the left ventricle and auricle, and then communicates at the back of the heart with a branch of the posterior coronary.

The posterior or right coronary artery arises from the right side of the aorta. It turns to the right, along the furrow between the right ventricle and auricle, to the back of the heart, where it inosculates with the horizontal branch of the left coronary. Besides this, it sends a branch down the inter-ventricular furrow at the back of the heart to the apex, where it communicates with the left coronary.

Thus the coronary arteries form two circles about the heart: the one, horizontal, runs round the base of the heart, in the furrow between the auricles and ventricles. The other, perpendicular, runs in the furrow between the ventricles.

The veins corresponding to the coronary arteries terminate

in a single trunk, which opens into the right auricle between the remains of the Eustachian valve and the auriculo-ventricular opening. The orifice of the vein is guarded by a valve, to prevent regurgitation of blood.

Arrangement of the Muscular Fibres of the Heart.—Most of the fibres are attached by both extremities to the fibrous rings of the heart. The fibres of the auricles are distinct from those of the ventricles. They consist of a superficial layer common to both cavities, and a deeper layer proper to each. The superficial fibres run transversely across the auricles; some pass into the septum. The deeper fibres run in circles chiefly round the auricular appendages and the entrance of the great veins, upon which a few may be traced for a short distance.

Of the ventricular fibres, some are common to both ventricles, others proper to each. The septum is formed principally by the fibres of the left. All the fibres take a more or less spiral course from the base towards the apex of the heart, where they coil round, pass into the interior of the ventricle, and form either “*carneæ columnæ*” or “*musculi papillares*.”*

Reduced to their simplest expression, the ventricles consist of two muscular sacs, enclosed in a third equally muscular. The same may be said of the auricles.

Weight.—The average weight of the heart is from ten to twelve ounces in the male, and from eight to ten in the female; but much depends upon the size and condition of the body generally. We have seen the heart so reduced in size by diabetic disease, that it weighed only five and a half ounces. As a general rule, it may be stated, that the heart gradually increases in length, breadth, and thickness, from childhood to age.†

Thickness of the Cavities.—The average thickness of the right auricle is about one line; that of the left, one and a half. Of course, the measure must not be taken during the “*rigor mortis*.”

The average thickness of the right ventricle at its thickest part—*i. e.*, the base—is about two lines. The average thick-


* For further information upon this subject, see the article in Todd's Cyclopædia.

† Consult Bizot, *Mém. de la Soc. Méd. d'Obser. de Paris*, tom. i. 1836.

ness of the left ventricle at its thickest part—*i. e.*, the middle—is about half an inch. In the female the average is less.

[The usual length of the heart from base to apex, we find to be from five and a half to five and three-quarters inches; about four inches belonging to the ventricles. The circumfer-

Fig. 23.



SCHEME OF THE FŒTAL CIRCULATION.

ence at the auriculo-ventricular groove is from eight to nine inches; in breadth, three to four inches; and in thickness, about two inches. The average thickness of the septum between the ventricles is about five lines; it being thicker in the middle. The circumference of the aortic opening measures

about two and three-quarters inches ; of the pulmonary, three ; of the left auriculo-ventricular, four ; and the right, four and a half inches. In the foetus and infant, the heart is placed vertically, instead of oblique, as in the adult. This organ is subject to many congenital malformations, apparently due to an arrest of development. A majority of these are incompatible with a long life.]

Peculiarities of Foetal Heart.—The heart of the foetus differs from that of the adult in the following points :—1. The Eustachian valve is well developed, in order to guide the current of blood from the vena cava inferior into the foramen ovale. 2. The foramen ovale is widely open. 3. The right and left pulmonary arteries are very much contracted, so as to admit very little blood to the lungs. 4. The ductus arteriosus (from the pulmonary artery to the aorta) is widely open. 5. The right and left ventricles are of equal thickness, because they have equal work to do.

Circulation of the Blood in the Foetus.—The umbilical vein, fig. 23, bringing pure blood from the placenta, enters at the umbilicus, and goes to the under surface of the liver, where it divides into many branches, which join the hepatic circulation. One of these branches, termed the "*ductus venosus*," goes straight into the inferior vena cava. From the inferior vena cava the blood enters the right auricle, and this stream (directed by the Eustachian valve) flows through the foramen ovale into the left auricle. From the left auricle it runs into the left ventricle, and thence through the aorta, into the great vessels of the head and the upper limbs, which are in this way supplied by almost pure blood.

From the head and upper limbs, the blood returns (impure) through the superior vena cava into the right auricle, and flows into the right ventricle. From the right ventricle it passes through the pulmonary artery, and the "*ductus arteriosus*," into the third part of the arch of the aorta ; only a very small quantity of it goes to the lungs. From the aorta the blood is conveyed through the umbilical arteries to the placenta, where it is purified.

STRUCTURE OF THE LUNGS.

The lungs are very vascular, spongy organs, in which the blood is purified by exposure to atmospheric air. Their situation and shape have been already described (p. 99). Our present concern is with their general structure.

The lungs are composed of cartilaginous and membranous tubes, of which the successive subdivisions convey the air into closely-packed, minute cells, called the air vesicles; of the ramifications of the pulmonary artery and veins; of the bronchial vessels concerned in their nutrition; of lymphatics and nerves. These component parts are united by areolar tissue, and covered externally by pleura. The point at which they respectively pass in and out is called the "root" of the lung.

The lungs are the lightest organs in the body, and swim in water. When entirely deprived of air, and reduced to their own substance, they sink. This is observed in certain pathological conditions,—for instance, when one lung is compressed, by effusion, into the chest, or rendered solid by inflammation.

Contractility of the Lung.—If an opening be made into the chest, the lung, which was in contact with the ribs, immediately recedes from them, and, provided there be not adhesions, gradually contracts. Again, if the lungs be artificially inflated, either in or out of the chest, we observe that, if left to themselves, they spontaneously expel a part of the air. This constant disposition to contract, in the living and the dead lung, is owing to the elastic tissue in the bronchial tubes and the air-cells; but more especially to a layer of delicate elastic tissue on the surface of the lung, which has been described by some anatomists as a distinct coat, under the name of the second or inner layer of the pleura.*

Colour.—The lungs are of a livid red or violet colour; they often present a mixture of tints, giving them a marble-like appearance. This is not the natural colour of the organ, since it is produced in the act of dying. It depends upon the

* In some animals, the seal especially, the elasticity of this tissue is very strongly marked.

stagnation of the venous blood, which the right ventricle still impels into the lungs, though respiration is failing. The tint varies in particular situations in proportion to the accumulation of the blood, and is always the deepest at the back of the lung. But the colour of the proper tissue of the lung, apart from the blood which it contains, is pale and light grey. This colour is seldom seen except in the lungs of infants who have never breathed, or after death from profuse hæmorrhage. It was particularly observed by the French surgeons in the lungs of individuals who were executed by the guillotine.*

Upon or near the surface of the lung, we observe black spots, which do not depend upon the blood, since they are seen in the palest lungs. They vary in number and size, and increase with age. The source of these discolorations is not exactly known. Some consider them as deposits of minute particles of carbon which have been inhaled with the air; others, as masses of black pigment cells.

Trachea.—This is a partly membranous, partly cartilaginous, tube, which proceeds from the larynx opposite the fifth cervical vertebra, and divides about the third dorsal vertebra into two tubes, called the right and left bronchi, one for each lung. Its length is from four to four and a half inches. Its diameter is about eight or ten lines in the adult; but it varies, according to the age of the individual, and the natural volume of the lungs. It is kept constantly open by a series of cartilaginous rings, from sixteen to twenty in number, which extend round two-thirds of its circumference. At the posterior part of the tube the rings are deficient. This deficiency is for the purpose of allowing the trachea to expand or contract; and the membranous part of the tube is provided with transverse muscular fibres, which can approximate the ends of the rings.

Bronchi.—The two bronchi differ in length, direction, and diameter. The *right* is shorter than the left, and passes more horizontally to the root of its lung. It is larger in all its diameters than the left; hence, foreign bodies which have accidentally dropped into the trachea are more likely to be carried into the right bronchus by the stream of the air. The *left*

* Bichat, Anat. descript. tom. iv. p. 12.

bronchus is about two inches in length, and descends more obliquely to its lung than the right.

At the root of the lung, each bronchus divides into two branches, an upper and a lower, corresponding to the lobes of the lung; on the right side, the lower branch sends a small division to the third lobe. The tubes diverge through the lung, and divide into branches, successively smaller and smaller, until they lead to the air-cells. These ramifications do not communicate with each other; they are like the branches of a tree; hence, when a bronchial tube is obstructed, all supply of air is cut off from the cells to which it leads.

The several tissues, cartilaginous, fibrous, muscular, mucous, and glandular, which collectively compose the air-passages, are not present in equal proportions throughout all their ramifications, but each is placed in greater or less amount where it is required. The cartilaginous rings, so necessary to keep the larger bronchi permanently open, become in the smaller tubes fewer and less regular in form. As the subdivisions of the tubes multiply, the cartilages consist of small pieces placed here and there,—they become less and less firm, and finally disappear altogether. The air-passages, when no longer traceable by the naked eye, are entirely membranous.

The *cartilages* of the trachea vary in number from sixteen to twenty: those of the right bronchus, from six to eight; those of the left, from nine to twelve. They form about two-thirds of a circle, and resemble a horse-shoe in form. The cartilage at the bifurcation of the trachea is shaped like the letter V; its angle projects into the centre of the main tube, and its sides belong one to each bronchus.

The cartilages are connected, and covered on their outer and inner surfaces by a tough membrane, consisting of fibrous and yellow elastic tissue. This membrane is attached to the circumference of the cricoid cartilage, and is continued through the whole extent of the trachea and the bronchial tubes. Posteriorly, where the cartilages are deficient, it maintains the integrity of the tube; in this situation, it consists of parallel and closely arranged longitudinal fibres, which are seated immediately beneath the mucous membrane, and raise it into folds. The elasticity of this structure admits of the elongation and contraction of the trachea.

Trachial Glands.—On the outer surface of the membranous part of the trachea and bronchi are small glands, of which the minute excretory ducts open into the tube. In health, their secretion is clear, not tenacious, and just sufficient to lubricate the air-passages. In bronchitis, they are the sources of the abundant viscid expectoration.

Muscular Fibre.—After removing the glands and the fibrous membrane from the back of the trachea, we expose a thin stratum of non-striped muscular fibres, which extend transversely between the cartilages. By their contraction, they approximate the ends of the cartilages, and diminish the calibre of the tube.

The *mucous membrane* lining the air-passages is a continuation from that of the larynx. Its colour in the natural state is nearly white, but in catarrhal affections it becomes bright red, in consequence of accumulation of blood in the capillary vessels. Its surface is lined by a layer of epithelium of the ciliated kind, of which the vibratal movement is directed in such a way as to favour the expectoration of the mucus. The mucous membrane ceases at the commencement of the air-cells: in them there is no ciliated epithelium.


[The mucous membrane should be described rather as becoming thinner as it passes along the bronchial tubes; and then continued into the cells, where it is transparent,—the ciliated columnæ epithelium, becoming changed to the pavement variety in the air-cells.]

Lobules of the Lung.—The surface of the healthy lung is mapped out by faint white lines, which circumscribe spaces of various form and size. These spaces indicate the lobules of the lung. Each lobule is a lung in miniature. Whoever understands the structure of a single lobule, understands the structure of the entire lung. The lobules are connected by fine areolar tissue, called “interlobular,” which is everywhere soft and elastic, to allow the free expansion of the organ. The cells of this tissue have no communication with the air-vesicles unless the latter be ruptured by excessive straining, and then this intermediate tissue becomes inflated with air, and gives rise to “interlobular emphysema.” When infiltrated with serum, it constitutes “anasarca” of the lungs.

Each lobule receives a small bronchial tube, which subdi-


vides into smaller branches. Thus reduced in size, the walls of the tubes no longer present traces of cartilaginous tissue, but are composed of a delicate elastic membrane, upon which the capillaries ramify in a very minute network.* Each tube finally leads into an irregular passage (intercellular passage), from which proceed on all sides numerous dilatations: these are the air-cells (fig. 24). The air-cells themselves present a

Fig. 24.



ULTIMATE AIR-CELLS OF THE LUNG (FROM
KOLLIKER).

Fig. 25.



MAGNIFIED TWENTY-
FIVE TIMES.

number of shallow depressions, separated by somewhat prominent partitions, so that their interior has a honeycomb appearance, as shown in fig. 25. The purpose of this is to increase the extent of surface upon which the capillaries may ramify. The structure of the minute air-cell of the human lung is in all respects similar to the large respiratory sac of the reptile.

Pulmonary Vessels.—The branches of the pulmonary artery subdivide with the bronchial tubes. Their ultimate ramifications spread out in such profusion over the air-cells, that a successfully injected lung appears a mass of the finest network of blood-vessels. This network is so close, that the interstices are even narrower than the vessels, which are, on

* In phthisis, the expectoration contains some of the debris of this elastic framework of the air vesicles; they can be seen under the microscope. This is not a bad test of the character of the sputa.

an average, about $\frac{1}{3000}$ th of an inch in diameter. The blood and air are not in actual contact. Nothing, however, intervenes but the wall of the cell and the capillary vessel, which are such delicate structures that they oppose no obstacle to the free interchange of gases by which the blood is purified. This purification is effected by the taking in of oxygen, and the elimination of carbonic acid. The most complete purification takes place in the single layer of capillaries between the folds of membrane projecting into the cell; for in this situation, both sides of these vessels are exposed to the action of the air. The blood, circulating in steady streams through this capillary plexus, returns through the pulmonary veins. These, at first extremely minute, gradually coalesce into larger and larger branches, which accompany those of the artery, and finally emerge from the root of the lung by two large trunks, which carry the pure florid blood to the left auricle of the heart.

From this outline of the anatomy of the lung, we see that the organ is so constructed as, in a given space to allow the largest possible quantity of impure blood to be brought in communication with the largest possible quantity of atmospheric air. It is difficult to conceive how any apparatus could be better adapted to the object in view. A stratum of blood, of great superficial extent, is exposed to an equal stratum of air, and these strata of contiguous fluids are contained in the interior of an organ so small as to lie within the compass of the chest.

Bronchial Vessels.—These are small arteries, two or more in number for each lung. They arise from the concavity of the arch of the aorta, and enter the lung in company with the divisions of the bronchi. They are the proper nutritive vessels of the organ. The old anatomists called them the "*vasa privata pulmonum*," to distinguish them from the "*vasa publica pulmonum*"—namely, the pulmonary arteries. The former provide "*pro existentia privata pulmonum*;" the latter "*pro bono publico totius organismi*." The bronchial vessels are distributed in various ways; some of their branches supply the coats of the air-passages and the large blood vessels; others, the interlobular tissue; and a few reach the surface of the lung, and ramify beneath the pleura. The right *bronchial*

veins terminate in the vena azygos ; the left, in the superior intercostal vein.

The *nerves* of the lung are derived from the pneumogastric and the sympathetic. They enter with the bronchial tubes, forming a plexus in front and behind them (anterior and posterior pulmonary plexus).

The *absorbents* of the lung form a network upon its surface and in the interlobular spaces. They all pass through the bronchial glands. Of these, the larger are situated about the bronchi near the root of the lung, particularly under the bifurcation of the trachea.

DISSECTION OF THE PHARYNX.

To obtain a view of the pharynx, cut through the trachea, œsophagus, and the great vessels of the neck, and then separate them from the bodies of the cervical vertebræ, to which they are loosely connected. The head should be removed at the first vertebra ; thus the pharynx and larynx will be left attached to the base of the skull. Horse-hair should be introduced through the mouth and the œsophagus to distend the pharynx.

[Another mode of preparing the parts for dissection, is, to divide the trachea and œsophagus at the top of the sternum, turn them up over the face, then separate the skull into two halves by a section of the saw carried behind the styloid process, and directed inwards and a little forwards. The occiput and the vertebral column will constitute one of the pieces, while the other will contain the face, tongue, trachea, and pharynx.]

The term "pharynx" is applied to that part of the alimentary canal which receives the food after it has been masticated, and propels it downwards into the œsophagus. It is a funnel-formed, muscular bag. Its upper part is attached to the basilar process of the occipital bone ; from thence it extends perpendicularly as low as the cricoid cartilage, where the continuation of it takes the name of "œsophagus." The bag is connected behind to the bodies of the cervical ver-

tebræ by loose cellular tissue which never contains fat. Remember that, in abscesses at the back of the pharynx, the matter is seated in this tissue. *Parallel with, and close to its sides, run the internal carotid arteries.* Its dimensions are not equal throughout. Its breadth at the upper part is just equal to that of the posterior openings of the nose; for here it is only required to transmit the air inspired through the nose; but it becomes much wider in the situation where it transmits the food—that is, at the back of the mouth: thence it gradually contracts to the œsophagus. The pharynx, therefore, may be compared to a funnel communicating in front, by wide apertures, with the nose, the mouth, and the larynx; while the œsophagus represents the tube leading from its lower end. The upper part of the funnel forms a cul-de-sac at the basilar process of the occipital bone. At this part there is, on each side, the opening of a narrow canal, called the “Eustachian tube,” through which air passes to the tympanum of the ear.*

Before we can examine the muscles of the pharynx, we must remove a layer of condensed cellular membrane, called by some anatomists the “*pharyngeal fascia.*” It is only a layer of cervical fascia behind the pharynx, and must not be confounded with the proper pharyngeal “aponeurosis,” which intervenes between its muscular and mucous walls.

At the back of the pharynx, near the base of the skull, we find some *absorbent glands.* They sometimes enlarge, and form a perceptible tumor in the pharynx.


In removing the fascia from the pharyngeal muscles, notice that a number of veins ramify and communicate in all directions. They constitute the “*pharyngeal venous plexus,*” and terminate in the internal jugular.

Constrictor Muscles of the Pharynx.—They are three in

* Observe that the pharynx conducts to the œsophagus by a gradual contraction of its channel. This transition, however, is in some cases sufficiently abrupt to detain a foreign body, such as a morsel of food more bulky than usual, at the top of the œsophagus. If such a substance become firmly impacted in this situation, one can readily understand that it will not only prevent the descent of food into the stomach, but that it may occasion, by its pressure on the trachea, alarming symptoms of suffocation. Supposing that the obstacle can neither be removed by the forceps, nor pushed into the stomach by the probang, it may then become necessary to extract it by making an incision into the œsophagus on the left side of the neck.

number, and arranged so that they overlap each other,—*i. e.* the inferior overlaps the middle, and the middle the superior (fig. 26). They have the same attachments on both sides of the body; and the fibres from the right and left meet together, and are inserted in the mesial line, the insertion being marked by a white longitudinal line called the “raphé.”

Fig. 26



MUSCLES OF THE PHARYNX.

The *inferior constrictor* arises from the side of the cricoid and thyroid cartilages. Its fibres expand over the lower part of the pharynx. The superior fibres ascend: the middle run transversely; the inferior descend, and are identified with the oesophagus. Beneath its lower border the recurrent laryngeal nerve enters the larynx.

The *middle constrictor* arises from the upper edge of the greater cornu of the os-hyoides, from its lesser cornu, and part of the stylo-hyoid ligament. Its fibres take different directions, so that, with those of the opposite muscle, they form a lozenge. The lower angle of the lozenge is covered by the inferior constrictor; the upper angle ascends nearly to the basilar process of the occiput, and terminates upon the pharyngeal aponeurosis. The external surface of the muscle is covered at its origin by the hyo-glossus.

The *superior constrictor* arises from the hamular process of the sphenoid bone; from the pterygo-maxillary ligament (which connects it with the buccinator); from the mylo-hyoid ridge of the lower jaw, and from the side of the tongue. The fibres pass inward to the mesial line: some of them are inserted through the medium of the pharyngeal aponeurosis into the basilar process.

The upper border of the superior constrictor presents on either side a free semilunar edge with its concavity upward, so that between it, and the base of the skull, a space is left in which the muscle is deficient (fig. 26). Here the pharynx is strengthened and walled in by its own aponeurosis. The space is called the "*sinus of Morgagni*;" and in it, with a little dissection, we get at the muscles which raise and tighten the soft palate; *i. e.* the levator palati, the tensor palati. The Eustachian tube opens into the pharynx just here. Observe that the fibres of the stylo-pharyngeus pass in between the superior and middle constrictors, and expand upon the side of the pharynx; some of them mingle with those of the constrictors, but most of them are inserted into the posterior margin of the thyroid cartilage.

Pharyngeal Membrane or Aponeurosis.—The pharyngeal aponeurosis, intervenes between the muscles and the mucous membrane of the pharynx. It is attached to the basilar process of the occiput, and to the points of the petrous portions of the temporal bones. It maintains the strength and integrity of the pharynx at its upper part, where the muscular fibres are deficient; but it gradually diminishes in thickness as it descends, and is finally lost on the œsophagus. Notice the number of mucous glands upon this aponeurosis, especially near the base of the skull and the Eustachian tube.

These glands sometimes enlarge and occasion deafness from pressure on the tube.

Openings into the Pharynx.—Now lay open the pharynx by a longitudinal incision, and observe the seven openings into it (fig. 27): 1. The posterior openings of the nares. 2. On either side of them, near the lower turbinated bones, are the openings of the Eustachian tubes: below the nares is the soft palate, with the uvula. 3. Below the soft palate is the communication with the mouth, called the “isthmus faucium.” On either side of this are two folds of mucous membrane, constituting the anterior and posterior half-arches of the palate; between them are the tonsils. Below the isthmus faucium is the epiglottis, which is connected to the base of the tongue by three folds of mucous membrane. 4. Below the epiglottis is the aperture of the larynx. 5. Lastly, is the opening into the œsophagus.


All these parts are lined by mucous membrane, common to the entire tract of the respiratory passages and the alimentary canal. But this membrane presents characteristic differences in the different parts of these channels, according as they are intended as passages for air or for food. The mucous membrane of the pharynx above the velum palati, being intended to transmit air only, is very delicate in its texture, and lined by ciliated epithelium like the rest of the air-passages. But opposite the fauces, the mucous membrane in every respect resembles that of the mouth, and is provided with squamous epithelium. At the back of the larynx the membrane is corrugated into folds, to allow the expansion of the pharynx during the passage of the food.

The membrane is lubricated by a plentiful secretion from the numerous mucous glands which are situated in the sub-mucous tissue throughout the whole extent of the pharynx, more particularly in the neighborhood of the Eustachian tubes.

Posterior openings of the Nose.—Notice the size of these two oval openings, that you may know how large to make the plug, when requisite in hæmorrhage from the nose. They are barely one inch in the long, half an inch in the short diameter. They are bounded above by the body of the sphenoid bone, externally by its pterygoid plate, below by the horizontal por-

tion of the palate bone: they are separated from each other by the vomer. If you remove the mucous membrane from the posterior part of the roof of the nose and the top of the pharynx, you will find beneath it much fibrous tissue. This is the reason why polypi, growing from these parts, are generally of a fibrous nature.

Fig. 27.



VIEW OF THE PHARYNX LAID OPEN FROM BEHIND.

Isthmus Faucium.—This name is given to the opening by which the mouth communicates with the pharynx. Look into the throat of a living person, and observe that this aperture is bounded, above by the soft palate and uvula, below by the

root of the tongue, and on either side by the arches of the palate, enclosing the tonsils between them.

Soft Palate.—This movable prolongation of the roof of the mouth is attached to the posterior border of the hard palate. It constitutes an imperfect partition between the mouth and the posterior nares. Its upper or nasal surface is convex, and continuous with the floor of the nose; its lower surface is concave, in adaptation to the back of the tongue, and is marked in the middle by a white line, indicating its original formation by two lateral halves. When the soft palate is at rest, it hangs obliquely downwards and backwards; but in swallowing, it is raised to the horizontal position by the levatores palati, comes into opposition with the back of the pharynx, and thus prevents the food from returning through the nose.

If you make a perpendicular section through the soft palate, you will see that the great bulk of it is made up of mucous glands, which lie thick on its under surface, to lubricate the passage of the food. Above these glands we come upon the aponeurosis of the palate; still higher, upon the fibres of the levator palati; and lastly, upon the nasal mucous membrane. The soft palate is supplied with blood by the descending palatine branch of the internal maxillary, and the ascending palatine branch of the external maxillary. Its nerves are derived from the second division of the fifth pair.

Uvula.—The uvula projects from the middle of the soft palate, and gives the free edge of it the appearance of a double arch. It contains a number of mucous glands and a little muscle, the “azygos uvulae.” Its length varies in different individuals, and in the same person at different times, according to the state of its muscle. It occasionally becomes permanently elongated, and causes considerable irritation, a tickle in the throat, and harassing cough. When you have to remove a portion of it, you should cut off only the redundant mucous membrane.

Arches of the Palate.—The soft palate is continued into the tongue and pharynx by two folds of mucous membrane on each side, enclosing muscular fibres. These are the anterior and posterior half-arches or pillars of the palate. The anterior arch describes a curve from the base uvula, to the side of

the tongue. It is well seen when the tongue is put out. The posterior arch, commencing at the side of the uvula, curves along the free margin of the palate, and terminates on the side of the pharynx. The posterior arches, when the tongue is depressed, can be seen through the span of the anterior. The tonsils are situated on each side between the arches. Now, what are the uses of these arches? They are chiefly concerned in the mechanism of deglutition. The anterior, (enclosing the "palato glossi" muscles,) contract so as to prevent the food from coming back into the mouth: the posterior (enclosing the "palato pharyngei" muscles,) contract like side curtains, and co-operate in preventing the food from passing into the nose. True, in vomiting, food does sometimes escape through the nostrils, but one cannot wonder at this, considering the violence with which it is driven into the pharynx.

Muscles of the Soft Palate.—The muscles of the soft palate lie immediately under the mucous membrane. There are four pairs; namely, the "levator palati," the "circumflexi or tensores palati," the "palato glossi," and "palato pharyngei:" the "azygos uvulæ" is single.

Levator Palati.—This muscle *arises* from the petrous portion of the temporal bone, and from the cartilage of the Eustachian tube. Its fibres spread out, and are inserted along the upper surface of the soft palate (fig. 27). Its action is to raise the palate, so as to make it horizontal in deglutition.

Circumflexus or Tensor Palati.—This muscle is situated between the internal pterygoid and the internal pterygoid plate of the sphenoid bone. It *arises* from the scaphoid fossa, and from the Eustachian tube. Thence it descends perpendicularly, ends in a tendon which turns round the hamular process, and expands into a broad aponeurosis, which is connected to its fellow of the opposite side, and is also attached to the horizontal plate of the palate bone. It gives strength to the soft palate. A synovial membrane facilitates the play of the tendon round the hamular process. Its *action* is to draw down and tighten the soft palate.

Azygos or Levator Uvulæ.—This consists of one or sometimes two thin bundles of muscular fibre, which arise from

the aponeurosis of the palate, and descend along the uvula, nearly down to its extremity.

Palato-glossus and Palato-pharyngeus.—These muscles are contained within the arches of the soft palate. The palato-glossus, within the interior arch, proceeds from the anterior surface of the soft palate to the side of the tongue, and is lost in the stylo-glossus muscle. The palato-pharyngeus, within the posterior arch, proceeds from the posterior border of the palate to the side of the pharynx and mixes with the fibres of the constrictor and the stylo-pharyngeus.

[These muscles, and the relation they bear to surrounding parts, should be particularly noticed, as the surgeon may have to divide them in the operation of staphyloraphy. Sir William Ferguson has shown that in cleft-palate, the fissure is separated by the levatores-palati and palato-pharyngei muscles; which offer an impediment to the parts healing after they are brought together. He recommends the division of these muscles as a necessary proceeding. The flaps are thus relaxed, and readily retained in position. The pillars should be divided just below the tonsils, with a pair of blunt-pointed scissors. To divide the levator-palati, Mr. Pollock puts the flap upon the stretch, and passes a double-edged knife through the soft palate, on the inner side of the hamular process, and above the line of the levator-palati. The handle is now raised and depressed, a sweeping cut being made along the posterior surface of the soft palate.]

Tonsils.—The tonsils consist of an aggregation of muciparous glands. They are situated at the entrance of the fauces, between the arches of the palate. Their use is to lubricate the fauces during the passage of the food. The orifices of their ducts, visible on the surface, give the tonsil an appearance like the shell of an almond. Hence, as well as from their oval figure, they are called "*amygdalæ*." These openings lead into small cells in the substance of the tonsil, lined by mucous membrane; and into these, numerous smaller ducts pour their secretion. The fluid, viscid and transparent in the healthy state, is apt to become white and opaque in inflammatory affections of the tonsils, and occasionally accumulates in these superficial cells, giving rise to the deceptive appearance of a small ulcer, or even a slough in the part.

Concerning the relations of the tonsil, remember that it lies close to the inner side of the internal carotid artery. It is only separated from this great vessel by the superior constrictor and the aponeurosis of the pharynx. Therefore, in removing a portion of the tonsil, or in opening an abscess near it; the point of the instrument should never be directed outwards, but *inwards* towards the mesial line.* The tonsils are supplied with blood by the tonsillar branch of the external maxillary artery.

Eustachian Tube.—This canal conveys air from the pharynx to the tympanum of the ear. Its orifice is situated opposite the back part of the inferior spongy bone. The direction of the tube from the pharynx is upwards, backwards, and outwards; it is an inch and a half long. The narrowest part is about the middle, and here its walls are in contact. Near the tympanum its walls are osseous, but towards the pharynx they are composed of fibro-cartilage and fibrous membrane. The cartilaginous end projects between the origins of the levator and the tensor palati, and gives attachment to some of their fibres. It is situated at the base of the skull, in the furrow between the petrous portion of the temporal and the great wing of the sphenoid bone. It adheres closely to the bony furrow, as well as to the fibro-cartilage filling up the "foramen lacerum medium." The orifice is not trumpet-shaped, as usually described, but an elliptical slit about half an inch long and nearly perpendicular. The fibro-cartilage bounds it only on the inner and the upper part of the circumference; the integrity of the canal externally is maintained by tough fibrous membrane.

The Eustachian tube is lined by a continuation of the mucous membrane of the pharynx, and covered by ciliated epithelium. Hence, inflammatory affections of the throat or tonsils are liable to be attended with deafness, from temporary obstruction of the tube.

Mucous glands surround the orifice of the tube like a second tonsil. They are similar in nature and function to the

* Cases are related by Portal and Béclard, in which the carotid artery was punctured in opening an abscess into the tonsil. The result was immediately fatal hæmorrhage.

glands beneath the mucous membrane of the mouth, the palate, and the pharynx.

[The passing of the Eustachian catheter may now be practiced. To do this, keep the point of the catheter upon the floor of the nostril, its concavity upwards, and inclined against the septum of the nose; pass it backwards until it reaches the soft palate, which in the patient is readily made known by his attempt to swallow. At this time turn the point of the catheter outwards, rotating it a little, then by a slight movement forwards and backwards, it passes into the tube. That it has entered the tube is known by its steadiness, as well as by the sensation of the patient.]

The *hard palate*, formed by the superior maxillary and palate bones, serves as a fulcrum for the tongue in the act of tasting, in mastication, in deglutition, and in the articulation of sounds. The tissue covering the bones is thick and close in its texture, and firmly united to the asperities on the bones. But it is not everywhere of equal thickness. Along the raphé in the mesial line, it is much thinner than at the sides, this is probably the reason why the hard palate is in this situation more prone to be perforated by syphilitic disease.

A thick layer of glands (*glandulae palatinae*), are arranged in rows on either side of the hard palate. These glands become more numerous and larger as we approach the soft palate. Their orifices are visible to the naked eye. The mucous membrane has a very thick epithelial coat, which gives the white colour to the palate. The descending palatine branch of the internal maxillary artery, and the palatine nerves from the superior maxillary, may be traced along each side of the roof of the mouth. The ramifications of these arteries and nerves supply the soft as well as the hard palate.

Mechanism of Deglutition.—With the anatomy of the parts fresh in your mind, consider for a moment the mechanism of deglutition. The food duly masticated, is collected into a mass upon the back of the tongue; the lower jaw is closed to give a fixed point for the action of the muscles which raise the os-hyoides and larynx; the food is then carried back into the pharynx by the pressure of the tongue against the palate, at the same time that the pharynx is elevated

and expanded to receive it.* Having reached the pharynx, the food is prevented from ascending into the nasal passages by the approximation of the posterior palatine arches, and the elevation of the soft palate which thus forms a temporary roof to the pharynx; it is prevented from returning into the mouth by the pressure of the retracted tongue, and the contraction of the anterior palatine arches: it cannot enter the larynx, because the "rima glottidis" is closed and protected by the falling of the epiglottis: consequently, being forcibly compressed by the constrictors of the pharynx, the food passes into the œsophagus.†

The food passes with different degrees of rapidity through the different parts of its course; but most rapidly through the pharynx. The necessity of this is obvious, when we reflect that the air tube must be closed while the food passes over it, and that the closure produces a temporary interruption to respiration. The progress of the food through the œsophagus is slow and gradual.

DISSECTION OF THE LARYNX.

Before you commence the dissection of the larynx, make yourself familiar with the cartilages which compose it, and the ligaments which connect them. In other words, examine the frame-work of the larynx as seen in a dry preparation.

Os-hyoides.—This bone, named from its resemblance to the Greek letter Upsilon, is situated between the larynx and the tongue, and serves for the attachment of the muscles of the tongue. It may be felt immediately below, and one inch and a half behind the symphysis of the jaw. Anatomists divide it into a body, two greater and two lesser cornua. The *body* is the thick central portion. Its anterior surface is convex, and marked by the attachment of muscles; its posterior concave

* The larynx being also elevated and drawn forward, a greater space is thus left between it and the vertebræ for the distension of the pharynx.

† This falling of the epiglottis is effected, not by special muscular agency, but by the simultaneous elevation of the larynx and the retraction of the tongue. A perpendicular section through all the parts concerned is necessary to show the working of this beautiful mechanism.

surface corresponds to the epiglottis. The *greater cornua* (right and left) project backwards for about an inch and a half, with a slight inclination upwards, and terminate in blunt ends tipped with cartilage. In young subjects they are connected to the body of the bone by fibro-cartilage; this in process of years becomes ossified. The *lesser cornua* are connected, one on each side to the point of junction between the body and the greater cornua, by means of a little joint lined by synovial membrane, which admits of free motion. They are not larger than a barley-corn, and the stylo-hyoid ligaments are attached to them.

The os-hyoides is connected to the thyroid cartilage by several ligaments, which contain a quantity of elastic tissue. There is:—1. The *anterior thyro-hyoid*, which proceeds from the notch of the thyroid cartilage to the upper and posterior part of the body of the os-hyoides. In front of this ligament there is, in the perfect larynx, a bursa of considerable size, of which the use is to facilitate the play of the thyroid cartilage behind the os-hyoides. 2. The right and left *lateral thyro-hyoid ligaments* extend between the extremities of the greater cornua of the os-hyoides and the ascending cornua of the thyroid cartilage. They often contain a little nodule of cartilage. The space between the hyoid bone and the thyroid cartilage is closed in the recent state by the *thyro-hyoid membrane*. Through this membrane the superior laryngeal nerve and artery enter the larynx.

Cartilages of the Larynx.—The framework of the larynx is composed of five cartilages connected by joints and elastic ligaments, so that they can be moved upon each other by appropriate muscles; the object of this motion being to act upon two elastic ligaments called the “vocal cords,” upon which the voice essentially depends.

Thyroid Cartilage.—This is so called because it shields the beautiful mechanism behind it. It consists of two lateral halves (*alæ*) united at an acute angle which forms the prominence termed “*Pomum Adami*.” This prominence presents a notch at its upper part, as if a portion had been sliced off to allow it to play behind the os-hyoides in deglutition. There is a bursa in front of it. I have seen this bursa as large as a pigeon’s egg. The outer surface of each ala is marked by an

oblique line, which gives attachment to the sterno-thyroid and thyro-hyoid muscles. The inferior border is slightly arched in the middle, and on either side presents a convex prominence, which gives attachment to the crico-thyroid muscle. The superior border is nearly horizontal. The posterior border is nearly vertical. This border terminates above and below in round projections called the upper and the lower cornua. The upper is the longer; the lower articulates with the side of the cricoid cartilage.

Cricoid Cartilage.—This cartilage, named from its resemblance to a ring,* is situated below the thyroid. It is not of equal depth all round. It is narrow in front where it may be felt about one quarter of an inch below the thyroid cartilage: from this part, the upper border gradually rises, so that, posteriorly, the ring is a full inch in vertical depth, and occupies part of the interval left between the alæ of the thyroid cartilage. In the middle of this broad posterior surface is a vertical ridge, on either side of which observe a superficial excavation for the origin of the crico-arytenoidei postici. On its upper part are two oval slightly convex surfaces for the articulation of the arytenoid cartilages. In front, its upper border presents a broad excavation to which the crico-thyroid ligament is attached. The lower border is connected by elastic membrane to the first ring of the trachea.

The thyroid is connected to the cricoid cartilage in front by the crico-thyroid membrane, and laterally by its two inferior horns. Between these two cartilages there is a perfect joint on either side, provided with a synovial membrane, and secured by ligaments. The object of this joint is to permit the approximation of the cartilages. You will understand presently how the movement between them regulates the tension of the vocal cords.

Arytenoid Cartilages.—These cartilages are situated, one on either side, at the back of the cricoid. In the fresh state, before the membranes and muscle have been removed, the space between them resembles the lip of a ewer†; from this resemblance their name is derived. Each is triangular with the apex upwards. Its posterior surface is concave, and gives

* Κρίκος, a ring.

† Ἀρύταινα, a ewer.

attachment to the arytenoid muscle. Its base presents an oval surface, which articulates with the cricoid cartilage. This joint has a very loose capsular and synovial membrane, and permits motion in all directions, like the first joint of the thumb. In front of the base is a tubercle (anterior tubercle), which gives attachment to the vocal cord, and contributes to form part of the boundary of the rima glottidis. At the outer and back part of the base is another tubercle (posterior tubercle), into which certain muscles moving the cartilage are inserted; namely, the crico-arytenoideus posticus, and crico-arytenoideus lateralis. The apex of the cartilage is surmounted by one or two cartilaginous nodules, called "*cornicula laryngis*."

Cordæ Vocales.—These are two elastic cords extending horizontally from the angle of the thyroid cartilage to the base of each of the arytenoid. They diverge as they pass backwards; the space between them is called the "rima glottidis." We shall presently see that, through the muscles which act upon the arytenoid cartilages, these cords can be approximated or removed from each other; in other words, the rima glottidis can be closed or dilated. When sufficiently tightened, and brought parallel by means of certain muscles, the cords are made to vibrate by the stream of the expired air, and thus is produced the voice.

What is the length of the true vocal cords? In the adult male they measure about five-eighths of an inch; in the female about four-eighths. In boys they are shorter; hence their peculiar voice. At puberty, the cords lengthen, and the voice breaks.

Epiglottis.—This is a piece of fibro-cartilage, which projects over the larynx like a valve. Its ordinary position is perpendicular, leaving the glottis free for respiration; but during the elevation of the larynx in deglutition it becomes horizontal, falls over the glottis, and prevents the entrance of food into the larynx. Understand that this falling of the epiglottis is accomplished, not by special muscular agency, but by the simultaneous elevation of the larynx and the retraction of the tongue. Its apex or lower part is attached by the *thyro-epiglottidean* ligament to the angle of the thyroid

cartilage; it is also connected by another ligament (*hyo-epiglottidean*) to the os-hyoides.

The cartilages of the larynx resemble those of the ribs in structure. In the young they are dense and elastic, but they have a tendency to ossify with age. In very old subjects, the thyroid and cricoid cartilages are often completely ossified, and their interior presents an areolar tissue, containing oily matter, analogous to the spongy texture of the bones. The epiglottis is never ossified, on account of its peculiar organization, which resembles that of the ear and the nose.

Let us now examine the larynx in its perfect condition.

Mucous Membrane of the Larynx.—The mucous membrane of the larynx presents a wrinkled appearance, except where it lines the vocal cords, and is connected to the subjacent structures by an abundance of fine cellular tissue, which admits of its being pinched up into large folds. This tissue deserves notice from the rapidity with which it becomes the seat of serous effusion in acute inflammation of the larynx, and thus produces sudden and alarming symptoms of suffocation. From the root of the tongue to the epiglottis the membrane forms three folds (*glosso-epiglottidean*) containing elastic tissue. From the epiglottis it is continued backwards on either side to the apices of the arytenoid cartilages, forming the "*aryteno-epiglottidean*" folds, which bound the entrance into the larynx. In the natural state it is of a pale rose color, and covered by ciliated epithelium.

The mucous membrane of the larynx is remarkable for its acute sensibility. This is requisite to guard the glottis during the passage of the food over it. The glottis is closed during the act of deglutition; but if, during this process, any one attempt to speak or laugh, the glottis opens, and allows the food to go, as it is termed, the wrong way. As soon as the foreign body touches the mucous membrane of the larynx, a spasmodic fit of coughing expels it.


The sub-mucous tissue of the larynx is studded with a profusion of mucous glands. An oblong mass of them lies in the aryteno-epiglottidean fold, and they are particularly numerous about the ventricles. The surface of the epiglottis towards the tongue is abundantly provided with them. Their

ducts pass through the epiglottis, and may be recognized as minute openings on its laryngeal aspect.

Superior Opening of the Larynx.—This is the opening by which the larynx communicates with the pharynx. Its outline is triangular; the front part is bounded by the epiglottis; the sides by the aryteno-epiglottidean folds; the apex presents the peculiar funnel-shape appearance, from which the arytenoid cartilages derive their name.

Inferior Opening of the Larynx, or Rima Glottidis.—Look down into the larynx and observe the horizontal slit in the middle line; this is the “rima glottidis,” or “glottis.” The anterior two-thirds of this slit is bounded by the true vocal cords; the posterior third by the arytenoid cartilages. On each side of the glottis is a little pocket; this is called the ventricle of the larynx, or “*sacculus laryngis*.” Pass the handle of the scalpel into one of these cavities, and observe that it leads into a cul-de-sac, which ascends for a short distance by the side of the thyroid cartilage. The probable use of these cavities is to allow free space for the vibration of the vocal cords. The mucous membrane lining them is so sensitive that if a foreign body accidentally lodges there, the patient has no rest until it is expelled. Lastly, observe what are called the “superior vocal cords.” These are the prominent folds which form the upper boundaries of the ventricles: they are called the “false vocal cords,” because they have little or nothing to do with the production of the voice. They are composed of elastic tissue, like the true vocal cords; but they contain also fatty tissue, which the true ones do not.

Fig. 28.



SHAPE OF THE GLOTTIS
WHEN AT REST.

The glottis admits of being dilated, contracted, and even completely closed by its appropriate muscles. When at rest, its shape is triangular as shown in fig. 28, where the arytenoid cartilages are cut through on a level with the vocal cords. During every inspiration, the glottis is dilated by the crico-arytenoidei

postici; it then becomes spear-shaped (fig. 30). During ex-

piration, it resumes its triangular shape : and this return to a state of rest is effected, not by muscular agency, but by two elastic ligaments shown in fig. 31, which draw the arytenoid cartilages together. Thus then the glottis, like the chest, is dilated by *muscular* tissue ; like the chest also, it is contracted by *elastic* tissue. In speaking or singing, the glottis assumes what is called the vocalizing position—that is, the opening becomes narrower and its edges nearly parallel.

Intrinsic Muscles of the Larynx.—There are nine muscles which act upon the “rima glottidis:” four on each side and one in the middle. The four pairs are—the crico-thyroidei, the crico-arytenoidei postici, the crico-arytenoidei laterales, and the thyro-arytenoidei. The single one is the arytenoideus.

Fig. 29.




DIAGRAM SHOWING THE ACTION OF THE CRICO-THYROID MUSCLE.


M. Crico-thyroideus.—This muscle is situated on the front of the larynx. It *arises* from the side of the cricoid cartilage, ascends obliquely outwards, and is inserted into the inferior border of the thyroid. Its *action* is to tighten the vocal cords. It does this by depressing the thyroid cartilage : for this cartilage cannot be depressed without lengthening these cords, as shown by the dotted line, fig. 29. Its nerve comes from the *superior laryngeal*.

M. Crico-arytenoideus Posticus.—This muscle arises from the posterior part of the crico cartilage, and is *inserted* into

the posterior tubercle of the arytenoid. Its *action* is to dilate the glottis. It does this by drawing the posterior tubercle of the arytenoid cartilage *towards* the mesial line, and therefore, the anterior tubercle (to which the vocal cord is attached) *from* the mesial line (fig. 30). In this movement the arytenoid cartilage rotates as it were upon a pivot, and acts as a lever of the first order; the fulcrum or ideal pivot being intermediate between the power and the weight. This muscle dilates the glottis every time we inspire. Its nerve comes from the inferior laryngeal.

M. Arytenoideus.—This single muscle occupies the inter-

Fig. 30.



GLOTTIS DILATED. MUSCLES DILATING IT REPRESENTED WAVY.

val between the back of the arytenoid cartilages. The fibres pass across from one cartilage to the other. Most of them are transverse, but some cross like the letter X, running from the base of the one to the apex of the other cartilage. *Action.*—By clasping the arytenoid cartilages, they assist in contracting the glottis.


M. Crico-arytenoideus Lateralis.—To expose this muscle, cut away the ala of the thyroid cartilage. It *arises* from the side of the cricoid cartilage, and is *inserted* into the anterior tubercle of the base of the arytenoid. *Action.*—By drawing the arytenoid cartilages inwards, the muscles of opposite sides contract the glottis (fig. 31, p. 149). Its nerve comes from the inferior laryngeal.

M. Thyro-arytenoideus.—This muscle *arises* from the angle

of the thyroid cartilage, runs horizontally backwards, and is *inserted* into the front surface of the base of the arytenoid. Its fibres run parallel with the vocal cord, and some of them are directly inserted into it. Part of the muscle spreads out so as to form a floor for the ventricle of the larynx, and is inserted into the outer border of the arytenoid cartilage. Nerve from the inferior laryngeal.

This muscle relaxes the vocal cord. More than this, it puts the lips of the glottis in the "vocalizing" position; in this the lips of the glottis are parallel, and the chink is reduced to the breadth of a shilling.*

Fig. 31.



GLOTTIS CLOSED. MUSCLES CLOSING IT REPRESENTED WAVY.

The following table shows the action of the several muscles which act upon the glottis:—

Crico-thyroidei . . .	Stretch the Vocal Cords.				
Thyro-arytenoidei . . .	Relax the Vocal Cords, and place them in the vocalizing position.				
Crico-arytenoidei postici . . .	Dilate the glottis.				
Crico-arytenoidei laterales . . .	Draw together the arytenoid cartilages.				} close the glottis.
Arytenoideus . . .	Ditto	Ditto	Ditto		

* *M. aryteno-epiglottideus*.—This consists of a few pale muscular fibres enclosed in the aryteno-epiglottidean fold. Some of them run near the free margin of the fold, others are found lower down. These, being distributed over the ventricle of the larynx, can approximate its walls so as almost to obliterate the cavity. Hence the name "compressor sacculi laryngis" given to this part of the muscle by Mr. Hilton (Guy's Hospital Reports, vol. ii.) who drew attention to it.

The *blood-vessels of the larynx* are derived from the superior and inferior thyroid arteries. The laryngeal branch of the superior thyroid passes through the thyro-hyoid membrane with the corresponding nerve, and divides into branches, which supply the muscles and the mucous membrane. The laryngeal branches of the inferior thyroid ascend behind the cricoid cartilage. A constant artery passes through the crico-thyroid membrane.

The *nerves of the larynx* are the superior and inferior (*recurrent*) laryngeal branches of the pneumogastric.

The *superior laryngeal*, having passed through the thyro-hyoid membrane, divides into branches, distributed to the mucous membrane of the larynx. Its filaments spread out in various directions; some to the anterior and posterior surfaces of the epiglottis, and to the aryteno-epiglottidean folds; others to the interior of the larynx and the vocal cords. A constant filament descends behind the ala of the thyroid cartilage, and communicates with the recurrent. Its external laryngeal branch supplies the crico-thyroid muscle (p. 57).

The *inferior (recurrent) laryngeal nerve* enters the larynx beneath the inferior constrictor, and ascends behind the joint between the thyroid and cricoid cartilages. It supplies all the intrinsic muscles of the larynx, except the crico-thyroid. If the recurrent nerve be divided, or in any way injured, the muscles moving the glottis become paralysed, but its sensibility remains unimpaired. When the nerve is compressed by a tumor—for instance, an aneurism of the arch of the aorta—the voice is changed to a whisper,* or even lost.

Difference between the Male and the Female Larynx.—Until the approach of puberty, there is no great difference in the relative size of the male and female larynx. The male larynx, within two years after this time, becomes nearly doubled in size; the female grows larger, but to a less extent.

The larynx of the adult male is in all proportions about one-third larger than that of the adult female.

The alæ of the thyroid cartilage form a more acute angle in the male; hence the greater projection of the “*pomum Adami*,” and the greater length of the vocal cords, in the male.

* Medical Gazette, Dec. 1843.

The average length of the vocal cords is in the { Male . . . 8 lines.
 { Female . . . 6 "

The average length of the glottis is in the . { Male . . . 12 lines.
 { Female . . . 10 "

The size of the larynx does not necessarily follow the proportions of the general stature; it may be as large in a little person as in a tall one: this corresponds with what we know of the voice.

Crico-thyroid Articulation.—This perfect little joint is provided with a capsule and synovial membrane. There are, besides, two strong ligaments. Both proceed from the cornu of the thyroid cartilage; the one upwards and backwards, the other downwards and forwards to the cricoid. Remember that the only kind of motion permitted is vertical: and that this motion regulates the tension of the vocal cords.

[The cartilages sometimes become ossified in old age. In the museum of the College of Physicians and Surgeons, there is a specimen of ossification of the thyroid, cricoid, and both arytenoids. Foreign bodies are at times arrested in the larynx, and if small, may be lodged in its sinuses. Œdema of the glottis is principally confined to the submucous cellular tissue around the epiglottis, at the margins of the glottis, and the back of the thyroid cartilage. This effusion rarely extends below the true vocal cords, the mucous membrane below this point being attached to the fibrous tissue without the intervention of cellular tissue.]

DISSECTION OF THE TONGUE.

The tongue is a complex muscular organ, subservient to taste, speech, suction, mastication, and deglutition. It is connected by its muscles (genio-hyo-glossus, hyo-glossus, and stylo-glossus), to the symphysis of the jaw, to the os-hyoides, and to the styloid process of the temporal bone. To the soft palate it is connected by the anterior palatine arch (p. 134): and to the epiglottis by folds of mucous membrane; in the middle fold is enclosed a layer of elastic tissue, called the "*glosso-epiglottidean*" ligament. This pulls up the epiglottis


when the tongue is put out of the mouth: hence the rule of never attempting to pass a tube into the œsophagus without pushing back the tongue; otherwise the tube would pass into the larynx.

Direct your attention first to the structure of the upper surface of the tongue—its most sensitive apparatus; next, to the structure of its interior muscular apparatus.

The upper surface of the tongue is composed of structures similar to those of the skin generally; that is to say, it consists of a "cutis," or true skin, with numerous projections, called "*papillæ*," and of a thick layer of epithelium. The cutis is much thinner than that of the skin; it affords insertion to some of the superficial muscular fibres of the tongue, and the blood-vessels form a close network in it before they pass into the *papillæ*.

Papillæ of the Tongue.—The *papillæ* on the tongue are distinguished, according to their size and form, into three kinds, viz., "*papillæ maximæ*, or *circumvallatæ*," "*papillæ fungiformæ*," "*papillæ filiformæ*." (Fig. 32.)

Fig. 32.



1. *Papillæ circumvallatæ*.
2. *Papillæ fungiformæ*.

The *papillæ circumvallatæ* vary in number from seven to twelve. They are arranged at the back of the tongue in two rows, which converge like the branches of the letter V, with the apex backwards, towards the so-called foramen cæcum—a blind pouch in the middle line of the tongue. Each of these *papillæ* is circular, with a central depression, and surrounded by a fossa: the fossa itself is circumscribed by an elevated ring.

The *papillæ fungiformæ*, smaller and more numerous than the *circumvallatæ*, are scattered over the upper surface and sides of the tongue. They vary in shape, some being cylindrical, others having heads like mushrooms; whence their name. Near the apex of the tongue they may be distinguished during life from the other *papillæ* by their redder color; especially if the tongue be

touched with a strong fluid, such as vinegar. They shoot out under the stimulus, and it is probable that the same effect always accompanies the sensation of taste. In scarlatina, and some exanthematous fevers, these papillæ become elongated, and of a bright red color; as the fever subsides, their points acquire a brownish tint, giving rise to what is called "the strawberry tongue."

The *papillæ filiformæ* are the smallest and most numerous. They are so closely aggregated that they give the tongue a velvet-like appearance. Their points are directed backwards, so that the tongue feels smooth, if the finger be passed over it from apex to base; but rough, if in the opposite direction.

All the papillæ are covered with a layer of tessellated epithelium. That which covers the filiform is thicker than the rest, and with a microscope is seen to project from their sides like hair. The various kinds of "fur" on the tongue consist of thickened and soddened epithelium.

Respecting the use of the papillæ, it is probable that they enable the tongue to detect impressions with greater delicacy; and that they are instrumental in detecting different kinds of sensation, whether of taste or touch. From the density and arrangement of their epithelial coat, the filiform papillæ give the surface of the tongue a roughness which is useful in its action upon the food. An apparatus of this kind, proportionately stronger and more developed, makes the tongue of ruminant animals an instrument by which they lay hold of their food. In the feline tribe, the lion and tiger for instance, these papillæ are so sharp and strong that they act like rasps, and enable the animal to lick the periosteum from the bones by a single stroke of his tongue. In some mammalia, they act like combs for cleaning the skin and the hair.

If the papillæ be injected, and examined under the microscope, it is found that they are not simple like those of the skin, but that each is composed of a secondary papillæ: for instance, analyse a "*papillæ circumvallatæ*" and you will see that it consists of an aggregation of smaller papillæ arranged parallel to each other; again, a filiform papilla consists of a central stem, from which minute secondary papillæ shoot off. This elaborate structure escapes observation because it is

buried beneath the epithelium.* Each secondary papilla receives a blood-vessel, which passes nearly to its apex, and returns in a loop-like manner. But the filaments of the gustatory nerve have not, hitherto, been traced beyond the base of the papilla.

Numerous small glands are found in the submucous tissue at the root of the tongue. They are similar in structure and secretion to the tonsillar and palatine glands, so that there is a complete ring of glands round the isthmus faucium. Small round orifices upon their surface indicate the termination of their ducts. Other mucous glands, with ducts from one quarter to half an inch long, are situated in the muscular substance of the tongue.

Glands beneath the Apex of the Tongue.—On the under surface of the apex of the tongue is placed, on either side, a group of glands presumed to be salivary. Considering each group as one gland, observe that it is oblong, with the long diameter from seven to ten lines, parallel with the axis of the tongue. It lies near the mesial line, a little below the ranine artery, on the outer side of the branches of the gustatory nerve, under some of the fibres of the stylo-glossus. Four or five ducts proceed from each group, and terminate by separate orifices on the under surface of the tongue.

Muscular Fibres of the Tongue.—The interior of the tongue is composed of muscular fibre and of a small quantity of fat. The *extrinsic* muscles of the tongue have been described in the dissection of the submaxillary region (p. 26). We have now to examine its *intrinsic* muscles. For this purpose the mucous membrane must be removed from the top of the tongue. It will then be obvious that the great bulk of the organ consists of fibres which proceed in a longitudinal direction, constituting the "*linguales*" muscles. These fibres are not all of equal length; the more superficial are implanted into the skin covering the tongue, those deeply seated run all the way from base to apex; and these are readily exposed by dissecting on the under surface of the tongue immediately on the outer side of the genio-hyo-glossus muscle. By the action of this mass of longitudinal fibres the tongue can be moved

* See Bowman and Todd's "Physiological Anatomy."

so as to reach any part of the mouth. Besides the linguales, there are muscular fibres intermingled variously in the tongue of which it is hard to give a description; one cannot be surprised at this, considering the various and rapid motions of the organ.

If we trace the genio-hyo-glossi of opposite sides into the tongue, we find that some of their fibres ascend directly to the surface; others cross each other in the middle line, intersect the longitudinal fibres, and finally terminate upon the sides of the tongue. Lastly, the fibres of the stylo-glossus may be traced along the side of the tongue to the apex.

In the mesial line near the base of the tongue is found a vertical plane of fibrous tissue, connected behind to the body of the hyoid bone, and lost in front between the muscles. This so-called "*nucleus fibrosus lingue*" is a feeble representative of the lingual bone in some of the lower animals.

[The vertical plane of fibrous tissue is a septum that extends throughout the entire length of the tongue, dividing it into symmetrical halves. This septum is thicker behind than in front. This organ is the seat of congenital deformities, the tip being sometimes bifurcated, sometimes it is wanting. The *frænum lingue*, which passes from the inferior surface to the floor of the mouth, may be too long antero-posteriorly, or too short vertically, giving rise to "tongue tie." The arteries of the opposite sides do not freely anastomose; this has been proved by minute injections, and by removing half of the tongue without hæmorrhage that was troublesome, after tying the lingual artery of the side operated upon.]

The *nerves supplying the tongue* should be followed into its substance. The hypo-glossal nerve supplies all the muscles with motor power. The gustatory or lingual branch of the fifth pair is distributed to the mucous membrane about the apex and sides of the tongue, and endows it with most acute sensibility. Upon this nerve depends the sensation of all ordinary impressions, such as that of hardness, softness, heat, cold, and the like. The glosso-pharyngeal nerve supplies the mucous membrane at the back and sides of the tongue. It is especially a nerve of taste.

DISSECTION OF THE SCALP.

An incision should be made from the nose along the mesial line to the occiput; another at right angles to the first from one side of the head to the other. These incisions must not divide more than the skin, that the subcutaneous vessels and nerves may not be injured.

Strata Composing the Scalp.—The several strata of soft parts covering the skull-cap are: 1, the skin; 2, a thin layer of adipose tissue which contains the cutaneous vessels and nerves, and the bulbs of the hair; and by which the skin is very closely connected to, 3, the broad thin tendon of the occipito-frontalis muscle, (aponeurosis of the scalp); 4, an abundance of loose areolar tissue, which permits the free motion of the scalp upon, 5, the pericranium, or periosteum of the skull-cap.

Immediately beneath the skin, then, we expose the thin stratum of adipose tissue which connects it to the aponeurosis of the scalp. It forms a bed for the bulbs of the hair and for the ramifications of the cutaneous arteries. The toughness of this tissue, in which the arteries ramify, does not readily permit them to retract when divided; hence the hæmorrhage which follows incised wounds of the scalp: hence, also, the difficulty of drawing them out with the forceps. The blood-vessels of the scalp are derived, in front, from the supra-orbital arteries; on the sides, from the temporal; behind, from the occipital and posterior auricular. Trace the leading trunks, and observe that they subdivide into branches, which inosculate freely, and finally form a vascular network among the bulbs of the hair. A few small branches here and there dip down through the aponeurosis of the scalp to the pericranium. The veins accompany the arteries.

Occipito-frontalis Muscle.—This cutaneous muscle is closely connected to the scalp. It consists of two fleshy portions, one on the occiput, the other on the forehead, connected by a broad aponeurosis. The origin of the muscles takes place from the outer two-thirds of the upper curve on the occipital bone, and the adjoining part of the mastoid process. The fibres ascend over the back of the head, and terminate in

the common aponeurosis. The frontal portion, commencing in an arched form from the common aponeurosis near the frontal suture, descends over the forehead, and terminates partly in the skin of the brow, partly in the orbicularis. Some fibres run down the nose under the name of the "pyramidalis nasi." The aponeurosis of the scalp is continued over the temples and side of the head, gradually changing from tendinous into fibro-cellular tissue. This muscle enables us to move the scalp backwards and forwards. But its chief action is as a muscle of expression. It elevates the brows, and occasions those transverse wrinkles in the expression of surprise. Like the other muscles of expression it is supplied by the facial nerve.

Muscles of the Ear.—There are several little muscles to move the cartilage of the ear. In man they are thin and pale; but in animals that possess a more delicate sense of hearing, they are much more developed, for the purpose of quickly directing the cartilage of the ear towards the pulse of the air.

[To dissect the superior muscles, the ear should be drawn down by hooks; their position then will be indicated by a slight prominence between the ear and head. To see the retrahens, the ear should be drawn forwards, when a ridge will denote its situation. The integument alone is all that we should remove, to reach these muscles. Their fibres are so pale at times as to be scarcely appreciated, if not altogether absent.]

M. attrahens aurem and *attollens aurem*.—These little muscles are triangular, and attached by their bases to the aponeurosis of the scalp. The fibres of each converge, and are inserted, those of the former into the front of the helix, those of the latter into the back part of the concha.

M. retrahens aurem.—This proceeds from the base of the mastoid process to the lower part of the concha. All the muscles of the ear are supplied by the auricular branch of the facial nerve.

Nerves of the Scalp.—*The supra-orbital nerve* is a branch of the ophthalmic division of the fifth pair. It emerges from the orbit through the notch of the frontal bone, and subdivides into branches, which are covered at first by the fibres of

the orbicularis and occipito-frontalis ; but they presently become subcutaneous : some of them may be traced over the top of the head as far as the occipital bone.

The superficial temporal nerves ramify in company with the arteries of the same name. Some of them are derived from the inferior maxillary, or third division of the fifth pair ; others from the facial nerve (p. 64).

The posterior auricular nerve is a branch of the facial, and runs with its corresponding artery behind the pinna of the ear (p. 64).

The great occipital nerve is the posterior branch of the second cervical nerve. After passing through the complexus, it appears on the occiput with the occipital artery, and divides into wide-spreading branches. The *lesser occipital nerve*, a branch of the cervical plexus (p. 4), runs near the posterior border of the insertion of the sterno-mastoid.

Points of Surgical Interest.—Raise the aponeurosis of the scalp, and observe the quantity of loose areolar tissue which intervenes between it and the pericranium. This tissue never contains fat. There are many points of surgical interest concerning it : 1. Its looseness accounts for the extensive effusions of blood which one sees after injuries on the head : 2. It admits of large flaps of the scalp being detached from the skull-cap ; but these flaps do not mortify because they carry their blood-vessels with them : 3. In phlegmonous erysipelas of the scalp, it becomes infiltrated with pus and sloughs ; hence the necessity of making incisions : for the scalp will not lose its vitality, and liberate the sloughs, like the skin of other parts under similar conditions, and why ? because its vessels run above the diseased tissue, and therefore its supply of blood is not cut off.

The *absorbent* vessels of the scalp run backwards towards the occiput, where they enter the absorbent glands in that situation. Here, therefore, one finds glandular enlargements when the scalp is diseased.

The brain should now be removed. This organ, with its membranes, will be described in a subsequent part of this work. At present we propose to examine the nerves as they pass out of the base of the skull, and then to dissect the cavernous sinus and the orbit.

Exit of the Cranial Nerves.—The cranial nerves proceed in pairs through the foramina at the base of the skull; they are named, first, second, third, fourth pair, &c. &c., according to their order of succession from before backwards.

The *first pair* are the *olfactory nerves*. These cannot be seen, because the olfactory lobes are removed with the brain, and torn from the delicate filaments which pass through the cribriform plate of the ethmoid bone.

The *second (optic nerve)* passes through the foramen opticum into the orbit.

The *third (motor oculi)* passes through the dura mater close behind the anterior clinoid process.

The *fourth (patheticus)*, a small nerve, passes through the dura mater a little behind the posterior clinoid process.

The *fifth (trigeminal nerve)* passes through an aperture in the dura mater beneath the tentorium cerebelli, just above the point of the petrous portion of the temporal bone.

The *sixth* passes through the dura mater behind the body of the sphenoid bone.

The *seventh*, consisting of the *facial* and *auditory nerves*, passes through the meatus auditorius internus.

The *eighth*, consisting of the *glosso-pharyngeal*, *pneumo-gastric*, and *spinal accessory*, passes through the foramen jugulare. These three divisions do not all pass through the same tube of dura mater. The glosso-pharyngeal has a separate tube anterior to the other two, which have a common one.

The *ninth*, or *hypo-glossal nerve*, passes through the anterior condyloid foramen.

[We prefer Sömmering's division of the cranial nerves into twelve pairs, thus making the facial or portio dura, the seventh; auditory or portio mollis, the eighth; glosso-pharyngeal, the ninth; pneumogastric or par vagum, the tenth; spinal accessory, the eleventh; and the hypo-glossal, the twelfth pair.]

On the inside of the skull, between the dura mater and the bones, observe the ramifications of the arteria meningeae media, with its two veins. This artery, a branch of the internal maxillary (p. 71), enters the skull through the foramen spinosum, and divides into two principal branches; one runs in a groove along the anterior inferior angle of the parietal

bone; the other curves backwards over the temporal bone, and subsequently ramifies on the parietal. It supplies the dura mater and the bones. The position of these arteries renders them liable to be injured in fractures of the skull. As a result of their injury, extravasation of blood may take place between the dura mater and the bones, and cause compression of the brain.

We must now examine the cavernous sinus, and the nerves

Fig. 33.




DIAGRAM OF THE VENOUS SINUSES AT THE BASE OF THE SKULL.

which course through its walls to the orbit; namely, the third, the fourth, the first division of the fifth, and the sixth.

Cavernous Sinus.—One among the many peculiarities of the cerebral circulation is, that the returning blood flows through fibrous canals formed by the dura mater, called "*sinuses*." The general anatomy, the course, and names of these canals will be considered with the anatomy of the brain. But the cavernous sinus, for reasons which will appear pres-

ently, should be examined immediately before the dissection of the orbit.

This sinus, then, (fig. 33), lies by the side of the body of the sphenoid bone. In front it receives the ophthalmic vein which comes through the sphenoidal fissure; behind, the superior and inferior petrosal sinuses run out of it; on the inner side it communicates with the circular sinus which surrounds the pituitary gland (P. in the diagram).

In the outer wall of the cavernous sinus we trace from above downwards the third nerve, the fourth, and the first division of the fifth on their course to the orbit.* On its inner wall we find the sixth nerve, and the internal carotid artery. These objects are said to be contained *in* the cavernous sinus; but clearly understand that none of them are bathed in the blood of the sinus, because they are separated from it by the lining membrane.

Follow the fifth nerve forwards, that you may see how its sensitive root spreads out into a great ganglion, termed after its discoverer, the "Gasserian" ganglion. From this ganglion proceed the primary divisions of the nerve; namely, the *ophthalmic* which passes through the sphenoidal fissure; the *superior maxillary*, through the foramen rotundum; and the *inferior maxillary*, through the foramen ovale. The small motor root of the fifth lies beneath the ganglion, with which it has no communication, and accompanies the inferior maxillary division to supply the muscles of mastication.

Curves of the Carotid Artery.—After the removal of the cavernous sinus, a good view is obtained of the remarkable curves, like the letter S, made by the carotid artery by the side of the sella turcica. The vessel enters the cranium at the apex of the petrous portion of the temporal bone, makes its sigmoid curves, and then passes through the dura mater between the anterior clinoid process and the optic nerve: here it gives off the ophthalmic artery.

Cavernous Plexus.—The superior cervical ganglion of the sympathetic sends up with the carotid artery filaments, which form a plexus round it in its tortuous course through the

* Such is the order in which the nerves are placed in the wall of the sinus. As they enter the orbit, the fourth nerve crosses over the third.

carotid canal, and by the side of the sphenoid. After a careful dissection, we may discover with the naked eye in this plexus very small ganglia, called *carotid* or *cavernous*; but they vary in number, size, and situation. Through these nerves a communication is established between the sympathetic and the nerves which enter the orbit. (Fig. 34.)

DISSECTION OF THE ORBIT.

The roof of the orbit should be removed as far back as the optic foramen. In doing this, be careful not to injure the little pulley on the inner side for the superior oblique.

Periosteum of the Orbit.—The roof being removed, we expose the fibrous membrane which lines the walls of the orbit. It is a continuation of the dura mater through the sphenoidal fissure. Traced forwards, we find that at the margin of the orbit it divides into two layers, one of which is continuous with the periosteum of the forehead, the other

Fig. 34.




DIAGRAM OF THE NERVES OF THE ORBIT.

forms the broad tarsal ligament which fixes the tarsal cartilage.

Fascia of the Orbit.—The fascia of the orbit serves the same purpose that fascia does in other parts. It provides the

lacrimal gland, and each of the muscles, with a loose sheath, thin and delicate at the back of the orbit, but stronger near the eye. In this situation it passes from one rectus muscle to the other, so that their tendinous insertions into the globe are connected by it. From the insertion of the muscle it is reflected backwards over the globe of the eye, and the optic nerve, and separates the eye from the fat at the bottom of the orbit.

Muscles and Nerves of the Orbit.—There are six muscles to move the eye; four of which, running in a straight direction, are called the "*recti*," and are arranged one above, one below, and one on each side of the globe. The remaining two are called from their direction, "*obliqui*," one superior, the other inferior. There is also a muscle to raise the eyelid, termed "*levator palpebræ*." The *nerves* are: the optic, which passes through the optic foramen; the third, the fourth, the first division of the fifth, and the sixth, all of which pass through the sphenoidal fissure. The third supplies all the muscles with motor power, except the superior oblique, which is supplied by the fourth, and the external rectus, which is supplied by the sixth. The ophthalmic division of the fifth divides into a frontal, lacrimal, and nasal branch. The orbit contains, also, a considerable quantity of soft fat which forms a bed for the eye and prevents it from being pulled too far back by its muscles. Upon the quantity of this fat, depends, in some measure, the difference in the prominence of the eyes. Its absorption in disease or old age occasions the sinking of the eyes. The eye is separated from the fat by a fold of the orbital fascia, which, like a "*tunica vaginalis*," enables the globe to roll with the greatest rapidity and precision. Lastly, the orbit contains the lacrimal gland.

After the removal of the periosteum, and the fascia of the orbit, the following objects are seen. In the middle we observe the *frontal* nerve and artery, lying upon the levator palpebræ; on the inner side is the superior oblique muscle with its nerve (the fourth); on the outer side is the lacrimal gland with the *lacrimal* nerve and artery.

Frontal Nerve.—This is one of the three branches of the ophthalmic division of the fifth. It runs forwards upon the upper surface of the levator palpebræ to the supra-orbital

notch, through which it ascends to supply the forehead and scalp. In the orbit it gives off the following branch:—


a. The *supra-trochlear* comes from the frontal, runs above the pulley of the superior oblique, to the inner angle of the orbit, and divides into branches, which supply the skin of the upper eyelid, forehead, and nose. One or two very delicate filaments may be traced through the bone to the mucous membrane of the frontal sinuses.*

Lachrymal Nerve.—This is the smallest of the three divisions of the ophthalmic. It runs along the outer side of the orbit, through the lachrymal gland, and is distributed to the upper eyelid. Its branches within the orbit are—1, filaments to the *lachrymal gland*; 2, a *malar*, which traverses a canal in the malar bone, and supplies the skin of the cheek; 3, one or two nerves which pass down to communicate with the superior maxillary division of the fifth.

Fourth, or Nervus Patheticus.—This nerve enters the orbit above the other nerves which pass through the sphenoidal fissure. It runs directly to the orbital surface of the superior oblique, to which it is exclusively distributed.

Lachrymal Gland.—This gland is situated within the external angular process of the frontal bone. It is about the

size and shape of an almond. Its upper surface is convex, in adaptation to the roof of the orbit; its lower is concave, in adaptation to the eyeball. The anterior part of the gland lies sometimes detached from the rest, close to the back part of the upper eyelid, and is covered by the conjunctiva. The whole gland is kept in place by a capsule† formed by the fascia of the orbit. In structure it resembles the salivary



glands. It consists of an aggregation of small lobes com-

* These filaments have been noticed by Blumenbach, "De Sinibus Frontal."

† This capsule, being a little stronger on the under surface of the gland, is described and figured by Sæmmering as a distinct ligament; "Icones Oculi Humani," tab. vii.

posed of smaller lobules, connected by areolar tissue. The excretory ducts, seven to ten in number, run parallel, and perforate the conjunctiva about a quarter of an inch above the edge of the tarsal cartilage (fig. 35). They are not easily discovered in the human eye, but in that of a horse or bullock they are large enough to admit a small probe. The secretion of the gland keeps the surface of the cornea constantly moist and polished. But if dust, or any foreign substance, irritate the eye, the tears flow in abundance, and wash it off.

All the muscles of the orbit, except the inferior oblique, arise from the margin of the foramen opticum, and pass forwards, like ribands, to their insertions.

Levator Palpebræ.—This muscle *arises* from the roof of the orbit, immediately in front of the optic foramen. It gradually increases in breadth, and terminates in a thin aponeurosis, which is *inserted* into the upper surface of the tarsal cartilage beneath the palpebral ligament. It is constantly in action when the eyes are open, in order to counteract the tendency of the lids to fall. As sleep approaches, the muscle relaxes, the eyes feel heavy, and the lids close.

Obliquus Superior.—This muscle *arises* from the inner side of the foramen opticum. It runs along the inner side of the orbit, and terminates in a round tendon, which passes through a cartilaginous pulley attached to the roof of the orbit. From this pulley the tendon is reflected outwards and backwards to the globe of the eye. It gradually expands, and is inserted into the outer and back part of the sclerotica, between the external and superior recti. The pulley is lined by a synovial membrane which is continued upon the tendon. The *action* of this muscle is to roll the eye on its own axis.

[The superior oblique passes beneath the superior rectus.]

The frontal nerve and levator palpebræ should now be reflected to expose the superior rectus muscle.


Recti Muscles.—These four muscles have a tendinous origin round the foramen opticum, so that collectively they embrace the optic nerve. They diverge from each other, one above, one below, and one on either side of the optic nerve; and are named, accordingly, rectus superior, inferior, externus, and internus. Their broad thin tendons are inserted into the

opposite sides of the sclerotic coat of the eye, about a quarter of an inch from the margin of the cornea (fig. 36).

The *external* rectus not only arises from the circumference of the optic foramen, but has another origin from the inner margin of the sphenoid fissure. Between these origins pass the third nerve, the nasal branch of the fifth, the sixth, and the ophthalmic vein.

The recti muscles enable us to direct the eye towards different points : hence the names given to them by Albinus—attol-

Fig. 36.



lens, depressor, abductor, and abductor oculi. It is obvious that, by the single action of one, or the combined action of two, the eye can be turned towards any direction.

Follow the recti to the eye, in order to see the beautiful shining tendons by which they are inserted. Notice also the "anterior ciliary arteries" which run to the eye along the tendons. The congestion of these little vessels occasions the red zone round the cornea in iritis. We have already mentioned that the tendons are invested by a fascia, which passes from one to the other, forming a loose tunic over the back of the eye. It is this fascia which resists the passage of the hook in the operation for the cure of squinting. Even after the complete division of the tendon, the eye may still be held in its faulty position, if this tissue, instead of possessing its proper softness and pliancy, happen to have become contracted and unyielding. Under such circumstances it is necessary to divide it freely with the scissors.

By removing the conjunctival coat of the eye the tendons of the recti are soon exposed. The breadth and the precise situation of their insertion deserves attention in reference to the operation of strabismus. The breadth of their insertion is about three-eighths of an inch, but the line of this insertion is not at all points equidistant from the cornea. The centre of the insertion is nearer to the cornea by about one line than either end. Taking the internal rectus, which, by the way,

By removing the conjunctival coat of the eye the tendons of the recti are soon exposed. The breadth and the precise situation of their insertion deserves attention in reference to the operation of strabismus. The breadth of their insertion is about three-eighths of an inch, but the line of this insertion is not at all points equidistant from the cornea. The centre of the insertion is nearer to the cornea by about one line than either end. Taking the internal rectus, which, by the way,

we have most frequently occasion to divide in strabismus, we find that the centre of its tendon is, upon an average, three lines only from the cornea, the lower part nearly five lines, and the upper four. It is, therefore, very possible that the lower part may be left undivided in the operation, being more in the background than the rest. The tendon of the internal rectus is nearer to the cornea than either of the others.

The superior rectus must now be reflected: in doing so, observe the branch of the third nerve, which supplies it and the levator palpebræ. After the removal of a quantity of fat, we expose the following parts:—1. The optic nerve. 2. The nasal nerve, the ophthalmic artery and vein, all of which cross over the optic nerve from without inwards. 3. Deeper towards the back of the orbit, between the optic nerve and the external rectus, is situated the “lenticular” ganglion.

Nasal Nerve.—This is one of the three divisions of the ophthalmic branch of the fifth pair (p. 162). It enters the orbit through the sphenoidal fissure between the two origins of the external rectus, and then crosses over the optic nerve towards the inner wall of the orbit. After giving off the *infra-trochlear* branch, the nerve runs out of the orbit through the foramen orbitale anterius, into the cranium, where it lies beneath the dura mater, upon the cribriform plate of the ethmoid bone. However, it soon leaves the cranium through a slit near the “crista galli,” and enters the nose. Here it sends filaments to the mucous membrane of the upper part of the septum, and upper spongy bone; but the main continuation of the nerve runs behind the nasal bone, becomes superficial between the bone and the cartilage, and, under the name of naso-lobular, is distributed to the skin of the end of the nose.

The nasal nerve gives off the following branches in the orbit:—

- a. One or two slender filaments to the *lenticular ganglion*.
- b. One or two *long ciliary* nerves. They run to the back of the globe of the eye, and pass through the sclerotic coat to supply the iris.
- c. *Infra-trochlear* nerve.—This runs below the pulley of the superior oblique, and divides into filaments, which supply the skin of the eyelids and the lachrymal sac.

Optic Nerve.—Having passed through the optic foramen, this nerve proceeds forwards and a little outwards to the globe of the eye, which it enters on the nasal side of its axis. It is invested by a fibrous coat derived from the dura mater. At the optic foramen it is surrounded by the tendinous origins of the recti; in the rest of its course, by loose fat and by the ciliary nerves and arteries.

Ophthalmic Artery.—This artery arises from the internal carotid. It enters the orbit through the optic foramen, outside the optic nerve; occasionally through the sphenoidal fissure. Its course is remarkably tortuous. Situated at first on the outer side of the optic nerve, it soon crosses over it, and runs along the inner side of the orbit to inosculate with the internal angular artery (a branch of the facial). Its branches arise in the following order:—

a. *Lachrymal artery.*—This branch proceeds along the outer side of the orbit to the lachrymal gland. After supplying the gland, it terminates in the upper eyelid.

b. *Supra-orbital artery.*—This branch runs forwards with the frontal nerve along the roof of the orbit, and emerges on the forehead through the supra-orbital notch.

c. *Arteria centralis retinae.*—This small branch enters the optic nerve, and runs in the centre of this nerve to the interior of the eye.

d. *Ciliary arteries.*—These branches proceed tortuously forwards with the optic nerve. They vary from fifteen to twenty in number, and perforate the sclerotic coat at the back of the eye, to supply the choroid coat and the iris. They are sometimes called "*posterior ciliary*," to distinguish them from others named "*anterior ciliary*," which proceed with the tendons of the recti muscles, and enter the front part of the sclerotica. In inflammation of the iris the vascular zone round the cornea arises from enlargement and congestion of the anterior ciliary arteries.

e. *Ethmoidal arteries.*—These branches pass through the foramina on the inner wall of the orbit to supply the mucous membrane of the nose.

f. *Muscular arteries.*—These are uncertain in their origin.

g. *Palpebral arteries.*—These branches proceed from the lachrymal, nasal, and supra-orbital arteries.

h. Nasal artery.—This branch may be considered as the termination of the ophthalmic. It leaves the orbit on the nasal side of the eye, and inosculates with the angular artery (termination of the facial). It supplies the side of the nose and the lachrymal sac.

Ophthalmic Vein.—This commences at the inner angle of the eye, by a communication with the angular and frontal veins. It runs backwards in a more straight course than the artery, receives corresponding branches, and finally passes between the two origins of the external rectus to terminate in the cavernous sinus.

Ophthalmic or Lenticular Ganglion.—This little ganglion (G, p. 162), about the size of a pin's head, is situated at the back of the orbit, between the optic nerve and the external rectus. It receives a branch (*sensitive root*) from the nasal nerve, another (*motor root*) from the lower division of the third nerve, and it also receives (*sympathetic*) filaments from the plexus round the internal carotid artery. The ganglion thus furnished with motor, sensitive, and sympathetic roots, gives off the ciliary nerves. These, from ten to twelve in number, run forward very tortuously with the optic nerve, pass through the back of the sclerotica, and are distributed to the iris. Since the ciliary nerves derive their motor influence from the third nerve, we see that the iris must lose its power of motion when this nerve is paralysed.

Third Nerve, Motor Oculi.—Just before it enters the sphenoidal fissure, the third nerve divides into two branches, both of which pass between the origins of the external rectus. The upper division has been already traced into the superior rectus and levator palpebræ. The lower division supplies a branch to the internal rectus, another to the inferior rectus, and then runs along the floor of the orbit to the inferior oblique muscle.

What is the result of paralysis of the third nerve? Falling of the upper eyelid (ptosis), external squint, dilatation and immobility of the pupil.

Sixth Nerve, Motor Externus.—This nerve enters the orbit between the origins of the external rectus, and terminates in fine filaments, which are exclusively distributed to the ocular surface of this muscle.

Respecting the motor nerves in the orbit, observe that they all enter the ocular surface of the muscles, with the exception of the fourth.

Inferior Oblique Muscle.—This muscle arises by a flat tendon from the superior maxillary bone, near the lower part of the lachrymal groove. It runs outwards and backwards between the rectus inferior and the orbit, then curves upwards between the globe and the external rectus, and is inserted by a broad thin tendon into the outer and back part of the sclerotic coat, close to the tendon of the superior oblique.

Action of the Oblique Muscles of the Eye.—The use of the oblique muscles is to rotate the eye upon its antero-posterior axis, so that, however much the head be moved obliquely to one side or the other, the image of the object may be always kept stationary upon one and the same point of the retina. This was first explained by Hunter. He says:—“When the head is moved towards the right shoulder, the superior oblique muscle of the right side acts, and keeps the right eye fixed on the object, and a similar effect is produced upon the left eye by the action of its inferior oblique muscle. When the head moves in a contrary direction, the other oblique muscles produce the same effect.”*

Orbital Branch of the Superior Maxillary Nerve.—This is at all times of small size, and sometimes absent. It comes from the trunk of the superior maxillary in the speno-maxillary fossa (see diagram), enters the orbit and divides into two branches. Of these, one communicates with the lachrymal branch of the ophthalmic; the other, called the “*temporo-malar*,”† usually divides into two branches, both of which pass through foramina in the malar bone, one to join the temporal branches of the inferior maxillary nerve, the other to supply the skin of the cheek.

Superior Maxillary Nerve and Spheno-palatine Ganglion.—In order to trace this nerve and its branches, we must remove the outer wall of the orbit, so as to expose the speno-maxillary fossa.

* Observations on Certain Parts of the Animal Economy.

† “*Subcutaneus malar*” of Sæmmerring and Meckel.

The superior maxillary nerve is the second division of the fifth cerebral nerve. Proceeding from the Gasserian ganglion (fig. 37), it leaves the skull through the foramen rotundum, passes horizontally forwards across the sphenomaxillary fossa, enters the infra-orbital canal with the corresponding artery, and finally emerges upon the face through the infra-orbital foramen. The branches given off are :—

- a. The *orbital* branch already described.
- b. Branches to the *spheno-palatine ganglion* (Meckel's) situated in the sphenomaxillary fossa.
- c. One or two *palatine* nerves.—These descend through the spheno-palatine ganglion to the hard and soft palate.

Fig. 37.




DIAGRAM OF THE SUPERIOR MAXILLARY NERVE.

1. Spheno-palatine ganglion.

2. Otic ganglion.

d. *Posterior dental* branches, two or three. They descend along the back part of the superior maxillary bone, and divide into smaller branches which pass through holes in the bone in company with minute arteries, and then run up the fangs of the molar teeth to supply the pulp. They also supply the gums, and the lining membrane of the antrum.

e. *Anterior dental* branch.—This arises just before the nerve emerges from the infra-orbital foramen. It descends in a special canal in the anterior wall of the upper jaw, and gives filaments to the fangs of the first molar, canine, and incisor

teeth. It also supplies the gums and the mucous lining of the antrum.

f. The terminal branch, namely the *infra-orbital*, was dissected with the face (p. 65).

At this stage of the dissection make a vertical incision rather on one side of the middle line of the skull, to expose the cavity of the nose. We shall thus be able to dissect the "spheno-palatine ganglion" and its branches.

Spheno-palatine Ganglion.—This little body (fig. 37) called after its discoverer, "*Meckel's ganglion*," is situated in the spheno-maxillary fossa, close to the outer side of the spheno-palatine foramen, immediately below the superior maxillary nerve. Its *sensitive* roots proceed from the superior maxillary; its *motor* from the vidian branch of the facial; and its *sympathetic* from the carotid plexus. Thus supplied, it furnishes branches to the mucous membrane and glands of the hard and soft palate and to the back part of the nose.*

* Meckel's ganglion gives off the following branches:—

a. *Branches to the palate.*—To see these the mucous membrane must be removed from the back part of the nose; we shall then be able to trace the nerves through their bony canals. Their course is indicated by corresponding arteries. There are generally three of these nerves, called by names originally given to them by Meckel, *anterior*, *middle*, and *posterior* palatine. The *anterior*, the large palatine nerve, descends through the palatine canal to the roof of the mouth, and then runs forwards along the hard palate nearly to the gums of the incisor teeth; but one or two branches proceed backwards to supply the soft palate. The *middle* palatine descends either in the same canal with the preceding, or in a smaller one of its own, and terminates exclusively in the mucous membrane and glands of the soft palate. The *posterior* palatine nerve may be traced in a special bony canal down to the soft palate, where it terminates in the mucous membrane and glands. One or two filaments pass into the uvula.

According to Longet (*Anat. et Physiol. du Système Nerveux*: Paris, 1842), the posterior palatine nerve supplies the levator palati and the azygos uvulæ with motor power. In this view of the subject the nerve is considered to be the continuation or terminal branch of the motor root of the ganglion, that, namely, derived from the facial. This opinion is supported by cases in which the uvula is stated to have been drawn on one side in consequence of paralysis of the opposite facial nerve. However, we have not succeeded in tracing the nerve into these muscles.

b. *Spheno-palatine or nasal branches.*—These, three or four in number, pass through the spheno-palatine foramen to the mucous membrane of the nose; but they are very delicate. To see them clearly, the part should have been steeped for some time in dilute nitric acid: afterwards, when well washed, these minute nerves may be easily recognized beneath the mucous membrane

Otic Ganglion.—The otic or auricular ganglion was discovered and described by Arnold, a German anatomist, in 1826.* “It is situated,” he says, “on the inner side of the inferior maxillary division of the fifth pair, immediately below its exit through the foramen ovale. (See diagram, p. 171.) Its inner surface is in contact with the circumflexus palati muscle and the cartilage of the Eustachian tube, and immediately behind it is the middle meningeal artery.” It is always of very small size; and, since we have sometimes failed in finding it, we conclude it is not present in all subjects.†

covering the spongy bones. Most of them ramify upon the outer wall of the nose and the spongy bones. One branch, fig. 37, originally called by Scarpa “*naso-palatine*,” traverses the roof of the nose, distributes filaments to the back part of the septum narium, and then proceeds obliquely forwards along the septum to the foramen incisivum, through which it passes, and finally terminates in the palate behind the incisor teeth.

According to Cloquet, the corresponding nerves of opposite sides unite at the foramen incisivum in a small ganglion which he calls the “*naso-palatine*.” Dissert. sur les Odeurs et les Organes de l’Olfaction : Paris, 1815.

c. *Branches to the pharynx and Eustachian tube.*—In parts prepared for the purpose we may sometimes trace minute filaments to the mucous membrane of the back of the nares, the Eustachian tube, and sphenoidal sinus.

d. *Vidian branch.*—This proceeds from the posterior part of the ganglion, backwards through the vidian canal of the sphenoid bone, traverses the fibro-cartilage at the base of the skull, and here divides into two branches, fig. 38. One joins the sympathetic plexus about the internal carotid artery; the other (sometimes called the *great petrosal*) enters the cranium, runs under the Gasserian ganglion and the dura mater, in a small groove in the petrous portion of the temporal bone, enters the “*hiatus Fallopii*,” and joins the facial nerve. It is probable that the vidian nerve proceeds, not from, but to the ganglion, and that it is the medium through which motor filaments are conveyed to it.

* J. Arnold, Diss. inaug. med. &c. : Heidelbergæ, 1826.

† Respecting its connections with other nerves, Arnold states that the otic ganglion derives filaments (of sensation) from the inferior maxillary, and also from the branch of this nerve, which goes to the internal pterygoid muscle. It also derives a slender filament from the temporo-auricular nerve. Its communication with the sympathetic is established by filaments which proceed from the “*nervi molles*” round the internal maxillary artery. It communicates also with the facial and glosso-pharyngeal nerves by a branch commonly called the lesser *petrosal* nerve. This nerve passes backwards either through the foramen ovale or the foramen spinosum, or through a small hole between the two, and runs beneath the dura mater along a minute groove on the petrous bone (outside the great petrosal nerve). Here it divides into two filaments, one of which joins the facial nerve in the aqueductus Fallopii, the other joins the tympanic branch of the glosso-pharyngeal. These nerves are exceed-

DISSECTION OF THE EIGHTH PAIR OF NERVES AT THE BASE OF THE SKULL.

IN this dissection we propose to examine the the glosso-pharyngeal, pneumogastric, and spinal accessory nerves in the jugular fossa, and the little ganglia and nerves belonging to them in this part of their course. These are very minute points of anatomy, and cannot be followed out with success, unless the nerves have been previously hardened by spirit, and the bones softened by acid. The first thing to be done is to remove the outer wall of the jugular fossa.

Glosso-pharyngeal.—This nerve passes through a separate tube of dura mater in front of that for the other two nerves. Looking at it from the interior of the skull we observe that it is situated immediately in front and rather to the inner side of the jugular fossa.

In its passage through the fossa, the nerve presents two ganglionic enlargements, named respectively the *jugular* and the *petrous*.

The *jugular ganglion* (ganglion superius) has been particularly described by Müller.* It is found upon the nerve immediately after its entrance into the canal of the dura mater, and is so small that its size does not in any direction exceed $\frac{1}{24}$ of an inch. It occupies the outer side of the nerve, and does not implicate all its fibres. According to my observation, this ganglion is frequently absent. It is not introduced in the following diagram.

The *petrous ganglion* (called after its discoverer, the ganglion of Andersch†) is situated upon the glosso-pharyngeal nerve, near the lower part of the jugular fossa. It is oval,

being difficult to trace, not only on account of their minute size, but also because they frequently run in canals in the petrous portion of the temporal bone.

The otic ganglion appears to send a branch to the tensor tympani and to the circumflexus palati muscles.

It may fairly be asked why Arnold called the ganglion, "otic or auricular." He states that by means of a filament from the lesser petrosal it communicates with the auditory nerve. But this must be doubtful, to say the least, since no other anatomist has, so far as we know, traced any connection of the kind.

* Medicin. Zeitung, Berlin, 1833, No. 52.

† Andersch, Fragm. descript. nerv. cardiac, 1791.

Fig. 38.




DIAGRAM OF THE EIGHTH PAIR AT THE BASE OF THE SKULL.

- | | |
|------------------------|--|
| 1. Facial n. | 7. Great petrosal, or vidian n. |
| 2. Glossopharyngeal n. | 8. Branch to Eustachian tube. |
| 3. Pneumogastric n. | 9. Auricular branch of Arnold. |
| 4. Nervus accessorius. | 10. Spheno-palatine. or Meckel's ganglion. |
| 5. Jacobson's n. | 11. Otic ganglion. |
| 6. Lesser petrosal. | 12. Chorda tympani. |

and about one-sixth of an inch long. It is connected by filaments to the pneumogastric and sympathetic nerves, and it gives off the *tympanic* or Jacobson's nerve.* This *tympanic* branch of the glossopharyngeal ascends through a minute canal in the bony ridge which separates the carotid from the jugular fossa, to the inner wall of the tympanum, and terminates in several filaments. One traverses a bony canal to the plexus of sympathetic nerves round the carotid artery; a second goes to the fenestra ovalis; a third to the fenestra rotunda; a fourth is distributed to the mucous membrane of the Eustachian tube; a fifth ascends in front of the fenestra ovalis, and joins the great petrosal nerve in the hiatus Fallopii; a sixth takes nearly a similar course, and under the name of the lesser petrosal nerve, proceeds along the front

* This nerve, though commonly called Jacobson's, was fully described by Andersch.

surface of the petrous bone to the otic ganglion. Thus it would appear that this tympanic branch is distributed to the mucous membrane of the tympanum and the Eustachian tube, and that it communicates with the spheno-palatine ganglion through the great petrosal nerve, and with the otic ganglion through the lesser petrosal.

Pneumogastric Nerve.—This nerve leaves the cranium with the nervus accessorius, through a canal in the dura mater behind that for the glosso-pharyngeal. At its entrance into the canal it is composed of a number of separate filaments; but these soon become embedded in a small ganglion. This ganglion, first described by Arnold,* is about $\frac{1}{12}$ of an inch in diameter. It is connected by filaments to the sympathetic and to the petrous ganglion of the glosso-pharyngeal. But its most singular branch is the one named by Arnold “the *auricular*,” because it is distributed to the pinna of the ear. This branch enters a minute foramen in the jugular fossa, near the styloid process, and then proceeds through the substance of the bone to the aqueductus Fallopii, where it divides into two branches; one joins the facial nerve, the other passes to the outside of the head through a canal between the front of the mastoid process and the meatus auditorius, and is distributed to the cartilage of the ear, and the meatus auditorius.

Immediately below this its first ganglion the pneumogastric nerve is joined by two branches from the nervus accessorius, and consequently becomes after this junction a compound nerve. The second or inferior ganglion of this nerve below the base of the skull has been already examined (p. 81).

Portio Dura through the Temporal Bone.—The portio dura or facial nerve, having arrived at the bottom of the meatus auditorius internus, enters the “*aqueductus Fallopii*.”† This is a tortuous canal excavated through the substance of the temporal bone, and terminating at the stylo-mastoid foramen. When exposed throughout its whole course, we observe that the nerve proceeds from the meatus internus for a short distance outwardly, then makes a sudden bend backwards

* Der Kopftheil. des veget. Nerven Systems. Heidelberg, 1831.

† Fallopius was a distinguished Italian anatomist, and professor at Pavia, 1551

along the inner wall of the tympanum above the fenestra ovalis, and lastly, curving downwards at the back of the tympanum, it leaves the skull at the stylo-mastoid foramen. Its branches in the temporal bone are :—

a. The *great petrosal* nerve (vidian) which runs to the spheno-palatine ganglion (p. 173, fig. 37).

b. The *lesser petrosal* nerve, which runs to the otic ganglion.

c. *Chorda tympani*.—This nerve arises from the facial about one-quarter of an inch before its exit from the stylo-mastoid foramen, ascends for a short distance in a bony canal at the back of the tympanum, and enters that cavity below the pyramid and close to the membrana tympani. It then runs forwards along the tympanum, across the handle of the malleus, to the fissura Glaseri, through which it emerges at the base of the skull, and joining the gustatory nerve, finally proceeds to the submaxillary ganglion.

DISSECTION OF THE NOSE.

PRESUMING that the learner is familiar with the bones composing the skeleton of the nose, we shall now describe—

1. The nasal cartilages ; 2. The general figure and arrangement of the nasal cavities ; 3. The membrane which lines them ; and lastly, the distribution of the olfactory nerves.

Cartilages of the Nose.—Two pieces of fibro-cartilage on either side assist in forming the framework of the external nose ; and one in the centre completes the septum between the nasal fossæ.

The *cartilage of the septum* is placed perpendicularly in the middle line : it may lean a little, however, to one side or the other, and in some rare instances it is perforated, so that the two nasal cavities communicate with each other. Its outline is nearly triangular. The posterior border is received into a groove in the perpendicular plate of the ethmoid and the vomer ; the anterior border is much thicker than the rest of the septum, and is connected, superiorly, with the nasal bones, and on either side with the lateral cartilages.

There are two *lateral cartilages*, an upper and a lower. The

upper is connected, superiorly, to the margin of the nasal and maxillary bones, anteriorly to the cartilage of the septum, and inferiorly to the lower cartilage by means of a tough fibrous membrane. The *lower* is sometimes called the cartilage of the pinna. It is curved upon itself in such a way as to form the boundary of the external opening of the nose. Superiorly, it is connected by fibrous membrane to the upper lateral cartilage; internally it is in contact with its fellow of the opposite side, forming the upper part of the "*columna nasi*;" posteriorly, it is attached by fibrous tissue to the superior maxillary bone: in this tissue, at the base of the ala, are usually found two or three nodules of cartilage, called "*cartilaginee sesamoideae*." We cannot but admire how well these cartilages are adapted to their office. By their elasticity they keep the nostrils continually open, and restore them to their ordinary size whenever they have been expanded by muscular action.

The little muscles moving the nasal cartilages have been described with the dissection of the face (p. 57).

Interior of the Nose.—By making a vertical section through the nasal cavities, a little on one side of the middle line, we expose the partly bony and partly cartilaginous partition (*septum narium*) by which they are divided. We notice that the greatest perpendicular depth of each nasal fossa is about the centre, and that from this point the depth gradually lessens, both towards the anterior and the posterior openings of the nose. Laterally, each fossa is very narrow in consequence of the projection of the spongy bones towards the septum: this narrowness in the transverse direction explains the rapidity with which swelling of the lining membrane from a simple cold obstructs the passage of air.

The boundary of the nasal fossæ is formed by the following bones: Superiorly, are the nasal, the ethmoid, and the body of the sphenoid; inferiorly, are the horizontal plates of the maxillary and palate bones; internally is the smooth and flat septum formed by the ethmoid, the vomer, and the cartilage; externally are the maxillary, the lachrymal, the ethmoid, the palate, and the pterygoid plate of the sphenoid.

Meatus of the Nose.—The outer wall of the nasal cavity is divided by the turbinated bones into three compartments

(meatus) of unequal size; and in these are orifices leading to air-cells (sinuses) in the sphenoid, ethmoid, frontal, and superior maxillary bones. Each of these compartments we should individually examine.

The *superior meatus* is the smallest of the three, and does not extend beyond the posterior half of the wall of the nose. We find in it one wide opening, which leads into the posterior ethmoidal and sphenoidal cells.

The *middle meatus* is larger than the superior. At its anterior part is a long narrow passage (infundibulum), which conducts upwards to the frontal and the anterior ethmoidal cells. About the middle is a small opening which leads into the antrum of the superior maxilla: this opening in the dry bone is large and irregular, but in the recent state it is reduced nearly to the size of a crow-quill by mucous membrane, so that a very little swelling of the membrane is sufficient to close the orifice entirely.

Notice that the orifices of the frontal and the ethmoid cells are so disposed that their secretion will, by its own gravity, pass into the nose. But this is not the case with the sphenoid and maxillary cells; to empty which the head must be inclined on one side. To see all these openings you must raise the respective turbinated bones.

The *inferior meatus* extends nearly along the whole length of the outer wall of the nose. By raising the lower turbinated bone, we observe, towards the front of the meatus, the termination of the nasal duct, through which the tears pass down from the lachrymal sac into the nose. This sac and duct we can now conveniently examine.

Lachrymal Sac and Nasal Duct.—This is the passage through which the tears are conveyed from the lachrymal ducts into the nose (p. 55). The lachrymal sac occupies the groove on the nasal side of the orbit, formed by the lachrymal and superior maxillary bones. The upper end is round and closed; the lower gradually contracts into the nasal duct, and opens into the inferior meatus. When the duct is obstructed, either from thickening of its lining membrane or from disease in the surrounding parts, the sac becomes distended by the tears, and forms a tumor at the inner angle of the eye. The sac is composed of a strong fibrous membrane

which adheres very closely to the bone, and is lined by mucous membrane. Its front surface is covered by the *tendo oculi* and the fascia proceeding from it.

The direction of the nasal duct is downwards and backwards. Its termination is guarded by a valvular fold of mucous membrane; consequently, when air is blown into the nasal passages while the nostrils are closed, the lachrymal sac does not become distended.

Behind the inferior turbinated bone is the opening of the Eustachian tube (p. 139). Into this, as well as into the nasal duct, we ought to practice the introduction of the probe. The chief difficulty is to prevent the probe from slipping into the cul-de-sac between the tube and the back of the pharynx.

Mucous or Schneiderian* Membrane.—This membrane lines the cavities of the nose and the air-cells about it, and adheres very firmly to the periosteum. Observe that at the lower border of the turbinated bones it is disposed in thick and loose folds. Such a fold is sometimes so prominent that it might be mistaken during life for a polypus. The membrane varies in thickness and vascularity in different parts of the nasal cavities. Upon the lower half of the septum and the lower turbinated bones it is much thicker than elsewhere, in consequence of a minute plexus of arteries and veins in the submucous tissue. The veins are especially numerous. Their sudden turgescence produces the temporary obstruction so common in an ordinary cold. In the sinuses, the mucous membrane is thinner, less vascular, and closely applied upon the periosteum.

The great vascularity of the Schneiderian membrane is obviously intended to elevate the temperature of the inspired air, and to pour out a copious secretion which answers the double purpose of partially saturating the air with vapor, and preventing the membrane from becoming too dry. Every one is aware that breathing is performed more comfortably through the nose than through the mouth.

Near the nostrils the mucous membrane is furnished with papillæ and a scaly epithelium, like the skin. In the sinuses

* So named in honor of Schneider, a learned anatomist who first gave an accurate description of this membrane. *De Catarrhis*. Wittenberg, 1660.

and all the lower regions of the nose, the epithelium is columnar, and provided with cilia; but in the upper part of the nose, where the sense of smell resides, we no longer find ciliated epithelium, but a soft pulpy stratum of nucleated cells.*

Blood-vessels of the Interior of the Nose.—The arteries of the nose are, the ethmoidal and nasal branches of the ophthalmic, and the nasal branch of the internal maxillary, which enters the nose through the spheno-palatine foramen. The external nose is supplied by the *arteria lateralis nasi*.

The *veins* of the nose correspond to the arteries. They communicate with the veins within the cranium through foramina in the cribriform plate of the ethmoid bone, also through the ophthalmic vein and the cavernous sinus. These communications explain the relief afforded by hæmorrhage from the nose in cases of cerebral congestion.

Olfactory Nerves.—The olfactory filaments, proceeding from each olfactory ganglion, in number about twenty on either side, pass through the foramina in the cribriform plate of the ethmoid bone. In its passage each filament is invested with a coat derived from the dura mater; so that these filaments, although soft and easily torn within the skull, are, outside it, as firm as any other nerves. To make them more conspicuous, the bones should be steeped in dilute nitric acid. We divide them into an *inner*, a *middle*, and an *outer* set. The *inner* traverse the grooves on the upper part of the septum, and terminate in minute filaments between the periosteum and the mucous membrane. They cannot be traced below the upper half of the septum. The *middle* supply the roof of the nose. The *outer* pass through grooves in the upper and middle turbinated bones, and are lost in the mucous membrane on the convex surfaces of these bones.†

The olfactory are nerves of special sense. The common sensibility of the mucous membrane of the nose is supplied by branches from the fifth pair of nerves; namely, the nasal branch of the ophthalmic (p. 162), and the nasal branch of the spheno-palatine ganglion (p. 172). That the sense of

* See Bowman and Todd, *Physiolog. Anatom.*, cap. xvi.

† See the beautiful plates of Scarpa and Scemmering.

smell is independent of the common sensibility of the nose is proved by experiment and by pathology. For instance, any disease affecting the olfactory nerve, even the inflammation in a common cold, impairs the sense of smell, whereas the common sensation of the part continues equally acute, and becomes even more so, as one may readily ascertain by introducing a foreign body into the nostril.

DISSECTION OF THE MUSCLES OF THE BACK.

THOSE muscles of the back, namely, the trapezius, longissimus dorsi, levator anguli scapulæ, and rhomboidei, which are concerned in the movements of the upper extremity, will be examined in the dissection of the arm. These, therefore, having been removed, we proceed to examine two muscles, named from their appearance "*serrati*," which extend from the spine to the ribs.

Serratus Posticus Superior.—This muscle is situated beneath the rhomboidei. It *arises* from the ligamentum nuchæ,* the spines of the last cervical, and two or three upper dorsal vertebræ, by a sheet-like aponeurosis which makes up nearly half the muscle; the fibres run obliquely downwards, and are *inserted* by fleshy slips into the second, third, fourth, and sometimes the fifth, ribs beyond their angles. Its *action* is to raise these ribs, and therefore to assist in inspiration.

Serratus Posticus Inferior.—This muscle is situated beneath the latissimus dorsi. It *arises* from the strong aponeurosis called "*fascia lumborum*," ascends obliquely, and is *inserted* by fleshy slips into the four lower ribs external to their angles. The tendency of its *action* is to pull down these ribs, and therefore to assist in expiration.

Vertebral Aponeurosis.—This aponeurosis extends from the spines of the vertebræ to the angles of the ribs, and serves to confine the "erector spinæ" in the vertebral groove.

* The ligamentum nuchæ is a rudiment of the great elastic ligament of quadrupeds (termed the *pack-wax*) which supports the weight of the head. It proceeds from the spine of the occiput to the spines of all the cervical vertebræ except the atlas; otherwise it would interfere with the free rotation of the head.

Observe that it proceeds from the upper border of the serratus posticus inferior, and that it is continued beneath the upper serrate muscle, so as to bind down the splenius.

Splenius.—This *arises* from the spines of the four or five upper dorsal and the last cervical vertebra, and from more or less of the ligamentum nuchæ. The fibres ascend and divide into two portions, named, according to their respective insertions, *splenius capitis*, and *splenius colli*.

a. The *splenius capitis* is inserted into the mastoid process, and the superior curved ridge of the occipital bone beneath the sterno-mastoid.

b. The *splenius colli* is inserted by tendinous slips into the transverse processes of the three upper cervical vertebræ.

The *action* of the splenius, taken as a whole, is to draw the head and the upper cervical vertebræ towards its own side: so far, it co-operates with the opposite sterno-mastoid muscle. When the splenii of opposite sides contract, they extend the cervical portion of the spine, and keep the head erect. The permanent contraction of a single splenius may occasion "wry neck." It is necessary to be aware of this, otherwise one might suppose the opposite sterno-mastoid to be affected, considering that the appearance of the distortion is alike in either case.

Erector Spinæ.—The mass of muscle which occupies the vertebral groove on either side of the spine, is, collectively, called "erector spinæ," since it counteracts the tendency of the trunk to fall forwards. We observe that it is thickest and strongest at that part of the spine where it has the greatest weight to support, namely, in the lumbar region; and that its thickness gradually decreases towards the top of the spine.

Fascia Lumborum.—The dense and shining aponeurosis which covers the erector spinæ is called "*fascia lumborum*." Observe that it is attached to the crest of the ilium, to the spines of the lower dorsal, all the lumbar, and sacral vertebræ. Its use is, not only to form a sheath for the erector spinæ, but to give origin to the latissimus dorsi, the serratus posticus inferior, the internal oblique and transverse muscles of the abdomen. If you make a vertical incision through it and reflect it, you will see that on the outer side of the erector spinæ it

is inseparably connected with the tendinous attachments of the internal oblique and transversalis to the transverse processes of the lumbar vertebræ.

The fascia lumborum having been reflected, we observe that the erector spinæ *arises* tendinous from the sacrum and ilium and the spines of the lumbar vertebræ. From this extensive origin the muscular fibres ascend, at first as a single mass. Near the last rib, however, this mass divides into two; an outer, called the "*sacro-lumbalis*;" an inner, the "*longissimus dorsi*." These two portions should be followed up the back; and there is no difficulty in doing so, because the division is indicated by a longitudinal groove, in which we observe the cutaneous branches of the intercostal vessels and nerves.

Sacro-lumbalis.—Tracing the *sacro-lumbalis* upwards, we find that it terminates in a series of tendons which are *inserted* into the angles of the six lower ribs.

Musculus Accessorius.—By turning outwards the *sacro-lumbalis*, we observe that it is continued upwards under the name of "*musculus accessorius ad sacro-lumbalem*." This *arises* by a series of tendons from the angles of the seven or eight lower ribs, and is *inserted* into the angles of the five or six upper ribs.

Cervicalis Ascendens.—This is the cervical continuation of the *musculus accessorius*. It *arises* by tendinous slips from the four or five upper ribs, and is inserted into the transverse processes of the four or five lower cervical vertebræ.

Longissimus Dorsi.—The *longissimus dorsi* (the inner portion of the erector spinæ) terminates in tendons which are *inserted* into the tubercles* at the root of the transverse processes of the lumbar vertebræ, also into the transverse processes of all the dorsal vertebræ, and into the greater number of the ribs (varying from eight to eleven) close to their junction with the transverse processes.

Transversalis Colli.—This is the cervical continuation of the *longissimus dorsi*. It *arises* by tendinous slips from the transverse processes of the second, third, fourth, fifth, and sixth dorsal vertebræ, and is inserted into the transverse processes of four cervical vertebræ.

* Called "anapophyses" by Professor Owen.

Trachelo-Mastoid.—This muscle, situated on the inner side of the preceding, is the continuation of the transversalis colli to the cranium. It arises from the transverse processes of the three or four lower cervical vertebræ, and is inserted by a flat tendon into the back part of the mastoid process beneath the splenius.*

Spinalis Dorsi.—This is a long narrow muscle, situated close to the spines of the dorsal vertebræ, and apparently a part of the longissimus dorsi. It arises from the spines of the two lower dorsal and two upper lumbar vertebræ, and is inserted also by little tendons into the spines of the six or eight upper dorsal vertebræ.

The muscles of the spine hitherto examined are all longi-

* Those who are familiar with the transcendental nomenclature of the vertebræ skeleton, will understand, from the following quotation, the plan upon which the muscles of the back are arranged :—

“ The muscles of the back are either longitudinal or oblique ; that is, they either pass vertically downwards from spinous process to spinous process, from diapophysis to diapophysis, from rib to rib (pleurapophyses) &c., or they extend obliquely from diapophysis to spine, or from diapophysis to pleurapophysis, &c.

“ The erector spinæ is composed of two planes of longitudinal fibres aggregated together, below, to form one mass at their point of origin, from the spines and posterior surface of the sacrum, from the sacro-iliac ligament, and from the posterior third of the iliac crest. It divides into two portions, the sacro-lumbalis and the longissimus dorsi.

“ The former, arising from the iliac crest, or from the pleurapophysis (rib) of the first sacral vertebra, is inserted by short flat tendons into (1.) the apices of the stunted lumbar ribs, close to the tendinous origins of the transversalis abdominis ; (2.) the angles of the eight or nine inferior dorsal ribs ; (3.) it is inserted, through the medium of the musculus accessorius, into the angles of the remaining superior ribs, and into the long and occasionally distinct pleurapophyseal element of the seventh cervical vertebra ; and (4.) through the medium of the cervicalis ascendens, into the pleurapophyseal elements of the third, fourth, fifth, and sixth cervical vertebræ. In other words, the muscular fibres extend from rib to rib, from the sacrum to the third cervical vertebra.

“ The longissimus dorsi, situated nearer the spine than the sacro-lumbalis, is inserted (1.) into the metapophyseal spine of the lumbar diapophysis ; (2.) into the diapophyses of all the dorsal vertebræ, near the origin of the lavatores costarum ; (3.) through the medium of the transversalis colli into the diapophyses of the second, third, fourth, fifth, and sixth cervical vertebræ ; and (4.) through the medium of the trachelo-mastoid into the mastoid process, or the only element of a transverse process possessed by the parietal vertebra. In other words, its fibres extend from diapophysis to diapophysis, from the sacrum, upwards, to the parietal vertebra.”—*Homologies of the Human Skeleton*, by H. Coote, p. 75.

tudinal in their direction. We now come to a series which run obliquely from the transverse to the spinous processes of the vertebræ. And first of the complexus.

Complexus.—This powerful muscle *arises* from the transverse processes of the five or six dorsal and the last cervical vertebræ; also from the articular processes of the fourth, fifth, and sixth cervical vertebræ. It is *inserted* between the two curved lines of the occiput, near the vertical crest. In the centre of the muscle there is generally tendinous tissue.* The muscle is perforated by the posterior branches of the second (the great occipital), third, and fourth cervical nerves. Its *action* is to keep the head erect.

Reflect the complexus to see the “*arteria cervicalis profunda*” (p. 47), and the posterior branches of the cervical nerves.

Transverso-spinalis.—This is the mass of muscle which lies in the vertebral groove after the reflection of the complexus and the erector spinæ. It consists of a series of fibres which extend between the transverse and spinous processes of the dorsal and cervical vertebræ, and is usually divided into the “*semispinalis colli*,” and “*semispinalis dorsi*.”

a. The *semispinalis colli* *arises* from the transverse processes of the five or six upper dorsal vertebræ, and is *inserted* into the spines of the axis and the three or four succeeding vertebræ.

b. The *semispinalis dorsi* *arises* from the transverse processes of the lower dorsal vertebræ, and is *inserted* into the spines of the upper dorsal and the two or three lower cervical vertebræ.

Now reflect part of the *semispinalis dorsi* in order to expose the “*multifidus spinæ*.” This may be considered a part of the preceding muscle, since its fixed points and the direction of its fibres are the same. It consists of a series of little muscles which extend between the spines and transverse processes of the vertebræ from the sacrum to the second cervical vertebra. Those in the lumbar region are the largest. They *arise* by tendinous slips from the transverse processes in the sacral

* The inner border of the complexus is described by some anatomists as a separate muscle, under the name of “*biventer cervicis*,” simply because there is much tendinous tissue in the centre of it.

and dorsal region, and from the articular processes in the lumbar and cervical region. They all ascend obliquely, and are *inserted* into the spines and laminæ of all the vertebræ excepting the atlas. It should be observed that their fibres are not of uniform length; some extend only from vertebra to vertebra, while others extend between one, two, or even three vertebræ.*

The action of these oblique muscles is not only to assist in maintaining the trunk erect, but to bend the spine to one or the other side.

Levatores Costarum.—These little muscles *arise* from the apices of the transverse processes of the dorsal vertebræ, and are *inserted* into the rib below. The direction of their fibres correspond with that of the outer layer of the intercostal muscles. They are muscles of inspiration.

Interspinales.—These muscles extend between the spines of the vertebræ. They only exist in those parts of the vertebral column which are the most movable, namely the cervical and lumbar regions. In the cervical region they are arranged in pairs. In the lumbar region they are single.

Intertransversales.—These muscles extend between the transverse processes in the cervical and lumbar regions. In the neck they are arranged in pairs, like the interspinales, and the corresponding cervical nerve separates one from the other. In the loins they are arranged singly.

We have next to examine the muscles concerned in the movements of the head upon the first and second vertebræ. The complexus must be previously reflected.

Rectus Capitis Posticus Major.—This is a largely developed interspinal muscle. It *arises* from the well marked spine of the second vertebra, and, expanding considerably, is *inserted* below the inferior curved ridge of the occipital bone; in other words, into the spine of the occipital vertebra.

Rectus Capitis Posticus Minor.—This is also an inter-

* Beneath the multifidus spinæ we find, in the dorsal region of the spine only, eleven little flat muscles, called by Theile (Müller's Archives f. Anat. &c., 1839), who first described them, "*rotatores spinæ*." They arise from the upper part of the transverse process, and are inserted into the lower border of the lamina of the vertebra above. These muscles form but a part of the multifidus spinæ.

spinal muscle, but smaller than the preceding. *Arising* from the feebly developed spine of the first vertebra, it expands as it ascends, and is *inserted* into the occipital bone between the inferior curved ridge and the foramen magnum. The *action* of the two preceding muscles is to raise the head.

Obliquus Inferior.—This *arises* from the spine of the second vertebra, and is *inserted* into the transverse process of the first. Its *action* is to rotate the first upon the second vertebra; in other words, to turn the head round.

Obliquus Superior.—This muscle *arises* from the transverse process of the first vertebra, and is *inserted* in the interval between the curved ridges of the occiput. Its *action* is to draw the occiput towards the spine.

Observe that the obliqui and recti muscles of one side form the sides of a triangle, in which we find the branches of the suboccipital nerve, the vertebral artery, and the arch of the atlas.

Rectus Capitis Lateralis.—This small muscle extends between the transverse process of the first vertebra and the *eminentia jugularis* of the occiput; but since this eminence is the transverse process of the occipital vertebra, the muscle must be considered as an intertransverse one.

[The deep muscles of the back are found to present many varieties, both as to their origin and insertion.]

Nerves of the Back.—The posterior branches of the spinal nerves supply the muscles and skin of the back. They pass backwards between the transverse processes of the vertebræ, and divide into external and internal branches. The general plan upon which these nerves are arranged is the same throughout the whole length of the spine; but since there are certain peculiarities deserving of notice in particular situations, we must examine each region separately.

Cervical Region.—The posterior division of the first cervical nerve (the suboccipital) passes between the arch of the atlas and the vertebral artery, and divides into branches which supply the recti and obliqui muscles concerned in the movement of the head upon the first two vertebræ.

The posterior branch (the great occipital) of the second cervical nerve is the largest of the series. It turns upwards beneath the inferior oblique muscle, passes through the com-

plexus, and runs with the occipital artery to the back of the scalp.

The posterior divisions of the six lower cervical nerves divide into *external* and *internal* branches. The *external* are small, and terminate in the splenius, and the continuation of the erector spinæ. The *internal*, by far the larger, proceed towards the spines of the vertebræ; those of the third, fourth, and fifth between the complexus and the semispinalis;* those of the sixth, seventh, and eighth between the semispinalis and the multifidus spinæ: after supplying the muscles they terminate in the skin.

Dorsal Region.—The posterior divisions of the spinal nerves in this region come out between the transverse processes and the tendons attached to them. They soon divide into *external* and *internal* branches. The *external* pass obliquely over the levatores costarum, between the sacro-lumbalis and the longissimus dorsi; and successively increase in size from above downwards. The upper six terminate in the erector spinæ and the levatores costarum; the rest, after supplying these muscles, pass through the latissimus dorsi, and become the cutaneous nerves of the back. The *internal* successively decrease in size from above downwards. They run towards the spine between the semispinalis and the multifidus spinæ. The upper six, after giving branches to the muscles, perforate the trapezius and become cutaneous nerves. The lower ones terminate in the muscles of the vertebral groove.

Lumbar Region.—The general arrangement of the nerves in this region resembles that of the dorsal. Their *external* branches supply the multifidus spinæ; their *internal*, after supplying the erector spinæ, become cutaneous.

Sacral Region—The posterior divisions of the spinal nerves in this region are small. They come out of the spinal canal through the foramina in the back of the sacrum. The upper two or three divide into *external* and *internal* branches. The *internal* terminate in the muscles; the *external* become cutaneous. The last two sacral nerves proceed, without dividing, to the integument.

* The posterior branches of the second, third, and fourth nerves are generally connected, beneath the complexus, by branches in the form of loops. This constitutes the posterior cervical plexus of some anatomists.

The *coccygeal* nerves are exceedingly small, and terminate in the skin.*

Vessels of the Back.—The vessels which supply the back are:—1. Small branches from the occipital; 2. Small branches from the vertebral; 3. The deep cervical; 4. The posterior branches of the intercostal and lumbar arteries.

The *occipital* artery furnishes several small branches to the muscles at the back of the neck: among others, one (called the *princeps cervicis*) descends beneath the complexus, and in some subjects inosculates with the deep cervical artery.

The *deep cervical* artery is the posterior branch of the first intercostal artery. It ascends between the complexus and the semispinalis muscles.

The posterior branches of the intercostal and lumbar arteries accompany the corresponding nerves, and are in all respects similar to them in distribution. They all send a branch into the spinal canal.

The *veins* correspond to the arteries.

Præ-vertebral Muscles.—We have, lastly, to examine three muscles situated in front of the spine; namely, the *longus colli*, the *rectus capitis anticus major*, and the *rectus capitis anticus minor*. In order to have a complete view of the two latter, a special dissection should be made before the head is removed from the first vertebra.

Longus Colli.—This muscle is situated in front of the spine, and extends from the third dorsal to the first cervical vertebra. For convenience of description it is divided into two sets of fibres, of which one extends *longitudinally* from the body of one vertebra to that of another; the other extends *obliquely* between the transverse processes and the bodies of the vertebræ.

The *longitudinal* portion of the muscle arises from the bodies of the two or three upper dorsal and the two lower cervical vertebræ, and is inserted into the bodies of the second, third, and fourth cervical vertebræ.

The *oblique* portion, arising from the transverse processes of the third, fourth, and fifth cervical vertebræ, ascends inwards, and is inserted into the front part or body of the first cervical

* The branching of the posterior divisions of the several spinal nerves has been accurately described by Mr. Ellis, *Med. Gazette*, Feb. 10th, 1843.

vertebra. Other oblique fibres proceed from the bodies of the three upper dorsal vertebræ, and are inserted into the transverse processes of the fifth and sixth cervical vertebræ. The *action* of the muscle, taken as a whole, must be to bend the cervical region of the spine.

Rectus Capitis Anticus Major.—This muscle arises from the transverse processes of the third, fourth, fifth, and sixth cervical vertebræ, and is inserted into the basilar process of the occipital bone, in front of the foramen magnum.

Rectus Capitis Anticus Minor.—This muscle arises from the transverse process of the first cervical vertebra, and is inserted into the basilar process of the occipital bone, nearer to the foramen magnum than the preceding muscle. The *action* of the recti muscles is to bend the head forwards.

LIGAMENTS OF THE SPINE.

The vertebræ are connected by their intervertebral fibro-cartilages, by ligaments in front of and behind their bodies, and by ligaments which extend between their arches and their spines. Their articular processes have capsular ligaments, and synovial membranes.

Anterior Common Ligament.—This is a strong band of longitudinal fibres which extends along the front of the bodies of the vertebræ from the axis to the sacrum. The fibres are not all of equal length. The more superficial extend from a given vertebra to the second or third below it; those a little deeper pass from a given vertebra to the second or third below it; while the deepest of all proceed from vertebra to vertebra. The ligament becomes broader and stronger in proportion to the size of the vertebræ. By making transverse incisions through it in different situations, we observe that its fibres are more firmly adherent to the intervertebral cartilages, and to the borders of the vertebræ, than to the middle of the bones.

Inter-spinous Ligaments.—These bands of ligamentous fibres fill up the intervals between the spines of the dorsal and lumbar vertebræ. They are the most marked in the lumbar region. Those fibres which connect the apices of the spines, being stronger than the rest, are described by some anato-

mists as separate ligaments, under the name of *supra-spinous*. Their use is to limit the flexion of the spine.

Ligaments between the Arches of the Vertebrae.—

These are called, on account of their color, *ligamenta subflava*.—To obtain a good view of them, we ought to remove the arches of the vertebrae with a saw. We then see that they extend from the arch of one vertebra to that of the next, filling up the intervals between them. Their strength increases with the size of the vertebrae. But the chief peculiarity about them is that they are exceedingly elastic. This elasticity answers a double purpose: it not only permits the spine to bend forwards, but materially assists in restoring it to its *curve of rest*. They economise muscular force, like the *ligamentum nuchæ* in animals.

Posterior Common Ligament.—This extends longitudinally in a similar manner to the anterior common ligament, along the posterior surface of the bodies of the vertebrae, from the occiput to the sacrum.

Intervertebral Fibro-Cartilage.—This substance is by far the strongest bond of connection between the vertebrae, and fulfils most important purposes in the mechanism of the spine. Its peculiar structure is adapted to break shocks, and to render the spine flexible and resilient. The best way to see the structure of an intervertebral fibro-cartilage is to make a horizontal section through one of them: say in the lumbar region. We then observe that it is firm and resisting near the circumference, but soft and pulpy towards the centre. The circumferential portion is composed of concentric layers of fibro-cartilage, placed vertically. These layers are attached by their edges to the vertebrae; they gradually decrease in number from the circumference towards the centre, and the interstices between them are filled up by the soft pulpy tissue. The central portion is composed almost entirely of this pulpy tissue; and it bulges when no longer under pressure. Practically it comes to this: the bodies of the vertebrae, in their motions upon each other, revolve upon a ball of fluid tightly girt all round by bands of fibrous tissue. These motions are regulated by the articular processes.

Dissect an intervertebral substance layer after layer in front, and you will find that the circumferential fibres extend ob-

liquely between the vertebræ, crossing each other like the branches of the letter X.

The thickness of the intervertebral cartilages is not the same in front and behind. It is this difference in their thickness, more than that in the bodies of the vertebræ, which produces the several curves of the spine. In the lumbar and cervical regions they are thicker in front; in the dorsal region behind.*

Capsular Ligaments.—Each joint between the articular processes has a capsular ligament and a synovial membrane. The surfaces of the bones are crusted with cartilage.

Motions of the Spine.—Though but little movement is permitted between any two vertebræ (the atlas and axis excepted), yet the collective motion between them all is considerable. The spine can be bent forwards, backwards, or on either side; it also admits of slight rotation. In consequence of the elasticity of the intervertebral cartilages and the ligamenta subflava it returns spontaneously to its natural curve of rest like an elastic bow. Its mobility is greatest in the cervical region, on account of the thickness of the fibro-cartilages, the small size of the vertebræ, the oblique direction of their articulations, and, above all, the horizontal position and the shortness of their spines. In the dorsal region there is very little mobility, on account of the vertical direction of the articular processes, and the manner in which the arches and the spines overlap each other. In the lumbar region the spine

* The structure of the intervertebral cartilages explains the well-known fact, that a man becomes shorter after standing for some hours; and that he regains his usual height after rest. The difference between the morning and evening stature amounts to more than half an inch.

It also explains the fact that a permanent lateral curvature of the spine may be produced (especially in the young) by the habitual practice of leaning to this or that side. Experience proves that the cause of lateral curvature depends more frequently upon some alteration in the structure of the fibro-cartilages than upon the bones. From an examination of the bodies of one hundred and thirty-four individuals with crooked spines, it was concluded that in two-thirds the bones were perfectly healthy; that the most frequent cause of curvature resided in the intervertebral substances; these being, on the concave side of the curve, almost absorbed, and on the convex side preternaturally developed. As might be expected in these cases, the muscles on the convex side become lengthened, and degenerate in structure.—On this subject see *Hildebrandt's Anatomie*, B. ii. s. 155.

again becomes more movable, on account of the thickness of the intervertebral cartilages, and the horizontal direction of the spinous processes.

Ligaments between the Occipital Bone and the first Vertebra.—The occiput is connected to the atlas by an *anterior* (occipito-atlantoid) membrane which passes from the foramen magnum to the front part of the atlas. The thickest part of this is in the middle. A *posterior* (occipito-atlantoid) membrane extends in a similar manner from the posterior border of the foramen magnum to the arch of the atlas. It is perforated by the vertebral artery.

Ligaments between the Occipital Bone and the Axis.—These are the most important; and to see them, we must expose the spinal canal by removing the arches of the upper cervical vertebræ, and the posterior common ligament, which is here very thick and strong. It descends from the basilar process of the occipital bone over the odontoid and transverse ligaments.

Odontoid or Check Ligaments.—The odontoid or check ligaments (fig. 39), are two very strong ligaments which proceed from the sides of the odontoid process to the tubercles on the inner sides of the condyles of the occiput. Their use is to limit the rotation of the head.

Articulation between the Atlas and the Axis.—The odontoid process of the axis forms a pivot upon which the head and atlas rotate. The most important ligament is the *transverse* (fig. 39). It passes behind the odontoid process, and is attached to the tubercles on the inner side of the articular processes of the atlas. Thus it forms with the atlas a ring into which the odontoid process is received. If this transverse ligament be divided we observe that the odontoid process is crusted with cartilage in front and behind, and provided with two synovial membranes.

Articulation of the Ribs.—All the ribs, with the exception of the first and the two last, are articulated with the bodies of two vertebræ, and with the transverse processes.

The head of each rib presents two articular surfaces, corresponding to the bodies of two vertebræ. There are two distinct articulations, each provided with a separate synovial membrane. The ligaments are :—1. An *anterior*, which con-

Fig. 39.




DIAGRAM OF THE ODONTOID AND TRANSVERSE LIGAMENTS.

nects the head of the rib with the vertebræ, and with the intervening fibro-cartilage: this, on account of the divergence of its fibres, is called the "stellate" ligament. 2. An *inter-articular*, which proceeds from the head of the rib to the intervertebral cartilage.

The tubercle of the rib articulates with the transverse process. This articulation has a capsular and synovial membrane, and is secured by the following ligaments:—1. The *posterior costo-transverse* passes from the apex of the transverse process to the tubercle of the rib. 2. The *middle costo-transverse* connects the neck of the rib to the front surface of the transverse process. 3. The *anterior costo-transverse* ascends from the neck of the rib to the lower border of the transverse process above it.

The head of the first rib articulates with a single vertebra.

The eleventh and twelfth ribs articulate each with a single vertebra, and are not connected to the transverse processes.

Connection between the Cartilages of the Ribs and the Sternum.—The cartilages of all the true ribs are received into slight concavities on the side of the sternum. In young subjects we find that the cartilages of the six lower true ribs have distinct articulations provided with synovial membranes. They are secured in front and behind by strong ligamentous


fibres, which proceed from the cartilages and radiate upon the sternum, crossing those of the opposite side.

The *costal cartilages from the sixth to the tenth* are connected by ligamentous fibres.

Articulation of the Lower Jaw.—The condyle of the lower jaw articulates with the glenoid cavity of the temporal bone. The joint is provided with an interarticular fibro-cartilage, and with external and internal lateral ligaments (fig. 40).

The *external lateral ligament* extends from the tubercle of the

Fig. 40.



TRANSVERSE SECTION TO SHOW THE LIGAMENTS AND THE FIBRO-CARTILAGE OF THE JOINT OF THE LOWER JAW. THE DOTTED LINES REPRESENT THE TWO SYNOVIAL MEMBRANES.

zygoma downwards and backwards to the tubercle of the condyle of the jaw.

The *internal lateral ligament* extends from the spinous process of the sphenoid bone to the border of the dental foramen. This "so-called" ligament cannot in any way contribute to the strength of the joint: the articulation of one side performs the office of internal lateral ligament to the other.

The *interarticular fibro-cartilage* is of an oval form, and thicker at the margin than at the centre. It is connected on the outer side to the external lateral ligament, and on the

inner side some of the fibres of the external pterygoid are inserted into it.

There are *two synovial membranes* for the joint. The larger and looser of the two is situated between the glenoid cavity and the fibro-cartilage. The other is interposed between the fibro-cartilage and the condyle of the jaw. They sometimes communicate through a small aperture in the centre of the fibro-cartilage.

The form of the articulation of the lower jaw admits of motion—upwards and downwards, forwards and backwards, and from side to side. A combination of these motions takes place in mastication: during this act the condyles of the jaw describe an oblique rotatory movement upon the glenoid cavity. We see what purposes are served by a piece of fibro-cartilage in this curious joint: it follows the condyle, and adapts a convenient socket for all its movements; more than this, being elastic, it breaks shocks; for shocks here would be almost fatal, considering what a thin plate the glenoid cavity is, and that just above it is the brain.

[In slight depression of the jaw, the condyles remain in the glenoid cavity; if considerable, the condyles glide forwards on the eminentia articularis, carrying their interarticular cartilages along. Should this movement be carried too far, as in a convulsive yawn, the capsule is ruptured, the cartilages being forced forwards, and the condyles pass into the zygomatic fossa.]

THE DISSECTION OF THE ARM.

THE arm being placed at a right angle with the body, make three incisions through the skin; the first, along the sternum; the second, along the clavicle; the third, along the front border of the axilla.

The skin should be carefully separated from the subjacent layer of adipose tissue (called the *superficial fascia*). In doing so, unless the subject be very muscular you will scarcely notice the thin pale fibres of the broad cutaneous muscle of the neck, ("platysma myoides" p. 1) which arises in this tissue.

Cutaneous Nerves.—Numerous nerves run through the subcutaneous tissue to the skin and the mammary gland. They are derived from various sources: some, branches of the superficial cervical plexus, descend over the clavicle: others, branches of the intercostal nerves, come through the intercostal spaces close to the sternum, with a small artery; a third series, also branches of the intercostal nerves, come out on the side of the chest, and run forwards over the outer border of the pectoralis major.

The *supra-clavicular* nerves which descend over the clavicle are subdivided, according to their direction, into *sternal*, *clavicular*, and *acromial* branches (diagram, p. 3). The *sternal* cross the inner end of the clavicle, to supply the skin over the upper part of the sternum. The *clavicular* pass over the middle of the clavicle, and supply the integument over the front of the chest and the mammary gland. The *acromial* branches cross over the outer end of the clavicle, and distribute their filaments to the skin of the shoulder.

Near the sternum you find the *anterior cutaneous branches of the intercostal nerves*. After traversing the pectoralis major, each nerve sends a filament to the skin over the sternum, and a larger one which supplies the skin over the pectoral muscle.

Branches of the internal mammary artery, for the supply of the mammary gland, accompany these nerves. During lactation they increase in size, ramifying tortuously over the surface of the gland. I have seen them nearly as large as the radial at the wrist. They would require a ligature in removal of the breast.

The *lateral cutaneous branches of the intercostal nerves* come out between the digitations of the serratus magnus on the side of the chest. They will be more fully described presently.

Now remove the superficial fascia with the mammary gland from the pectoralis major, by dissecting parallel to the course of its fibres. When the muscle has been fully exposed, observe its shape, the course of its fibres, their origin, and insertion.

Pectoralis Major.—The pectoralis major is the great muscle on the front of the chest. It *arises* from the sternal half of the clavicle, from the front of the sternum, and from the car-

tilages of all the true ribs except the first and last. The fibres converge towards the arm, and terminate in a flat tendon, about two inches in breadth, which is *inserted* into the anterior margin of the bicipital groove of the humerus. Their arrangement, as well as the structure of their tendon, is peculiar. The lower fibres which form the boundary of the axilla are folded beneath the rest, and terminate upon the upper part of the tendon, *i. e.* nearer to the shoulder-joint; whereas the upper fibres descend in front of the lower, and terminate upon the lower part of the tendon. Consequently the upper and lower fibres of the muscle cross each other previously to their insertion.

Now what is the object of this arrangement? It enables all the fibres to act simultaneously when the arm is extended.

The upper part of the tendon sends off an aponeurosis, which straps down the long head of the biceps, and is attached to the great tuberosity of the humerus. The lower part is intimately connected with the fascia of the upper arm, and the tendon of the deltoid.

The chief *action* of the pectoralis major is to draw the humerus towards the chest: as in placing the hand on the opposite shoulder, or in pulling an object towards the body. When the arm is raised and made the fixed point, the muscle assists in raising the trunk, as in climbing.

Between the pectoralis major and the deltoid, the great muscle of the shoulder, there is an interval varying in extent in different subjects, but always more marked towards the clavicle. It contains a small artery—the *thoracica-humeraria*—and the *cephalic vein*, which ascends along the outer side of the arm, and empties itself into the axillary vein. This interval is the proper place to feel for the coracoid process. In doubtful injuries about the shoulder this point of bone is a good landmark in helping us to come to our diagnosis.

The pectoralis major is supplied with nerves by the anterior thoracic branches of the brachial plexus; with blood by the long and short thoracic branches of the axillary artery.

[In some cases the clavicular and sternal portions are separated by areolar tissue. From the sternal portion a fasciculus sometimes descends and unites with the rectus external oblique and even extends to the margin of the latissimus dorsi.

It may also be joined to the brachialis anticus. We have often noticed it blended with the deltoid. If the arms are fixed, this muscle will draw the ribs upwards and outwards, and by thus enlarging the cavity of the chest, assist in the act of respiration.]

Anatomy of the Infra-Clavicular Region.—Reflect from the clavicle the clavicular portion of the pectoralis major. Beneath the portion so reflected part of the pectoralis minor will be exposed. Now between the upper border of this muscle and the clavicle there is an important space, in which you must dissect out and examine the relative position of the following objects:—

a. A strong ligamentous expansion, called the costo-coracoid fascia, which extends from the cartilage of the first rib to the clavicle.

b. The axillary vein, artery, and plexus of nerves.

c. The subclavius muscle enclosed in its fibrous sheath.

d. A short arterial trunk, (the thoracic axis,) which divides into several radiating branches.

e. The termination of the cephalic vein in the axillary.

f. Two or three nerves, (the anterior thoracic,) which descend from the axillary plexus beneath the clavicle, cross in front of the axillary vessels, and subdivide to supply the pectoral muscles.

Costo-Coracoid Fascia.—This fascia extends from the cartilage of the first rib to the coracoid process; between these points it is attached to the clavicle, and forms a complete investment for the subclavius muscle. It presents a crescent-shaped edge which arches over, and protects the axillary vessels and nerves: from this edge is prolonged a fascia which accompanies the vessels into the axilla, enclosing them in a kind of sheath.

Thoracic Axis.—This is the first branch of the axillary artery. It comes off above the pectoralis minor and soon divides into three branches—the superior or short thoracic, the thoracica humeraria, and the thoracica acromialis (p. 210). The *superior* or *short thoracic*, runs between the pectoralis major and minor, supplying ramifications to both. The *thoracica humeraria* descends with the cephalic vein, in the interval between the pectoralis major and deltoid, and ramifies in

both. The *thoracica acromialis* passes over the coracoid process to the under surface of the deltoid, which it supplies, and communicates with the circumflex branches of the axillary artery. All these arteries are accompanied by veins, which most frequently empty themselves into the cephalic, but occasionally into the axillary vein itself.

Cephalic Vein.—The cephalic vein is one of the principal cutaneous veins of the arm. Commencing on the back of the thumb and forefinger, it runs up the radial side of the forearm over the front of the elbow-joint; thence ascending along the outer edge of the biceps muscle, it runs up the interval between the pectoralis major and deltoid to reach the axillary vein.*

Relative Position of the Axillary Vessels and Nerves.—In the space now before us are the great vessels and nerves of the axilla in the first part of their course. They lie at a great depth from the surface. They are surrounded by a sheath of fascia, which descends with them beneath the clavicle, and are situated with regard to each other in the following manner:—The axillary vein lies in front of the artery, and rather to its sternal side. The axillary plexus of nerves is situated above the artery and on a posterior plane. The plexus consists of two or sometimes three large cords, which result from the union of the anterior branches of the last four cervical nerves and the first dorsal (p. 207).

Anterior Thoracic Nerves.—These nerves come from the axillary plexus to supply the pectoral muscles. There are generally two, one for each pectoral muscle: and they enter the muscles with their respective arteries. If you were to trace these nerves to their source, you would find that they proceed from the fifth and sixth cervical nerves. However, the chief point of interest about them is, that they cross over the axillary artery in the first part of its course, where you might be called upon to tie it. Therefore, you would be careful not to include them in the ligature.

Subclavius.—This muscle lies beneath the clavicle en-

* The cephalic vein sometimes runs over the clavicle to join the external jugular; or there may be a communication over the clavicle between these veins.

closed in a strong sheath, which you should open in order to expose it. It *arises* from the cartilage of the first rib by a round tendon, and is *inserted* into a groove on the under surface of the clavicle. Its nerve comes from the brachial plexus. Its *action* is to depress the clavicle, and prevent its too great elevation.

Difficulty of tying the first part of the Axillary Artery.—From this view of the relations of the axillary artery in the first part of its course, some idea may be formed of the difficulty of passing a ligature round it in this situation. In addition to its great depth from the surface, varieties sometimes occur in the position of the nerves and veins, which would render the operation still more embarrassing. For instance, the anterior thoracic nerves may be more numerous than usual, and form by their mutual communications a kind of plexus round the artery. I have often seen a large nerve crossing obliquely over the artery immediately below the clavicle, to form one of the roots of the median nerve.

The cephalic vein may ascend higher than usual, and open into the subclavian; and, as it receives large veins corresponding to the thoracic axis, a concourse of veins would be met with in front of the artery.

Again, it is by no means uncommon to find that a deep-seated vein, (*e. g.* the supra-scapular,) crosses over the artery to join the axillary vein.

DISSECTION OF THE AXILLA.

Sebaceous Glands.—Reflect the skin from the axilla and the side of the chest, taking care not to remove with it the adjacent fascia. In close contact with the skin, near the roots of the hair in the axilla, are numerous sebaceous glands. They are of a reddish-brown color, and rather larger than a pin's head.

Axillary Fascia.—This tough membrane lies immediately beneath the skin of the axilla, and is nothing more than a continuation of the general investment of the muscles. It closes in and forms the floor of the cavity of the axilla. Now

what are its connections? Externally it is strengthened by fibres from the tendons of the pectoralis major and latissimus dorsi, and is continuous with the fascia of the arm; internally, it is prolonged on the side of the chest, over the serratus magnus muscle: in front and behind, it divides so as to enclose between its layers the muscles which form the boundaries of the axilla. Thus the interior layer encloses the two pectoral muscles, and is connected with the coracoid process, the costo-coracoid ligament, and the clavicle; the posterior layer encloses the latissimus dorsi, and passes backwards to the spine.

A subcutaneous artery, sometimes of considerable size, often runs in the substance of the axillary fascia. It generally arises from the brachial, or from the lower part of the axillary, and runs across the floor of the axilla towards the lower edge of the pectoralis major. So far as I know, it has not been hitherto named; but it is worth remembering, because it would occasion much hæmorrhage if wounded in opening an abscess.

The next thing to be done is to remove this axillary fascia and to display the boundaries and contents of the axilla. The dissection of this cavity is difficult. You must proceed cautiously. Bear in mind that the trunk blood-vessels and nerves run through the *upper* part of the axilla; that the long thoracic artery runs along the anterior border, and the subscapular artery along the posterior. Commence working, therefore, in the middle: break down with the handle of the scalpel the loose connective tissue, fat, and absorbent glands, which occupy the cavity. You will soon discover some cutaneous nerves coming out between the ribs. These nerves are the *lateral cutaneous branches of the intercostal nerves*; they perforate the intercostal spaces between the digitations of the serratus magnus, midway between the sternum and the spine, and divide into anterior and posterior branches. The *anterior* turn over the pectoralis major, to supply the skin on the front of the chest and the mammary gland. The *posterior* pass backwards over the latissimus dorsi, and are distributed to the skin covering this muscle and the scapula.

Intercosto-Humeral Nerves.—The perforating branch of the second intercostal nerve requires a special description; it

is larger than the others, and is called the "*intercosto-humeral nerve*," because it supplies the integuments of the arm. It comes through the second intercostal space, traverses the upper part of the axilla, where it receives a branch of the lesser internal cutaneous nerve (nerve of Wrisberg), and terminates in filaments, which are distributed to the skin on the inner side of the arm, as low as the internal condyle. The perforating branch of the third intercostal is also an "*intercosto-humeral*" nerve. It receives a branch from the second, and runs a similar course. The distribution of these nerves may account for the pain down the arm which is sometimes experienced in pleurisy.

Boundaries of the Axilla.—The axilla may be described as a conical space, of which the summit is beneath the clavicle, and the base between the pectoralis major and latissimus dorsi muscles. Now what are its boundaries? On the *inner* side, it is bounded by the ribs, covered by the serratus magnus; on the *outer* side by the humerus, covered by the coracobrachialis and biceps muscles; in *front* by the pectoralis major, *behind* by the latissimus dorsi, teres major, and subscapularis. Its anterior and posterior boundaries converge from the chest, so that the axilla becomes narrower towards the arm. With a full view of the axilla before you, think what serious consequences may follow suppuration here: the matter may burrow under the pectoral muscles; or under the scapula; or it may run up beneath the clavicle and point in the neck.

Axillary Absorbent Glands.—The axillary glands form a continuous chain beneath the clavicle, with the cervical glands. They are from fifteen to twenty in number, of a reddish-brown color and variable size. Most of them lie near some large blood-vessel; others are imbedded in the loose tissue of the axilla; sometimes one or two small ones are observed along the lower border of the pectoralis major. They are supplied with blood by a branch (*thoracica alaris*) of the axillary artery, and by branches from the thoracic and infrascapular arteries.

These glands receive the absorbents from the arm, from the front and side of the chest, and from the outer half of the mammary gland.

You should now reflect the pectoralis major from its inser-

tion, to expose the pectoralis minor and the ramifications of the short and long thoracic arteries. Preserve the arteries as much as possible in connection with the main trunks.

Pectoralis Minor.—This muscle *arises* from the third, fourth, and fifth ribs. The fibres run upwards and outwards, and converge to a strong tendon, which is *inserted* into the anterior surface of the coracoid process. The tendon is connected to that of the coraco-brachialis by a strong fascia which forms a protection for the subjacent axillary vessels and nerves. The *action* of this muscle is to draw the scapula downwards and forwards.

[We may meet a third pectoral muscle, arising from the first and second ribs, and inserted as above; or a fasciculus may arise from the first rib and passing beneath the pectoralis minor, be inserted into the capsule of the shoulder-joint, or become attached to the coraco-brachialis.]

Latissimus Dorsi.—This muscle forms the posterior margin of the axilla. It *arises* from the crest of the ilium, from the spines of all the lumbar, and six lower dorsal vertebræ, and by digitations from the three lower ribs. It is *inserted* by a broad flat tendon (which runs behind the axillary vessels and nerves), into the bottom of the bicipital groove of the humerus.

[This muscle curves around the lower border of the teres major, and is twisted upon itself so that the superior fibres become first posterior, then inferior; the vertical, first anterior, then superior. Its tendon passes in front of that of the teres major. This muscle also crosses the inferior border of the scapular, from which it often receives bands of muscular fibres. The tendons of the teres major and latissimus dorsi are separated by a bursa; a bursa may also intervene between the muscle and angle of the scapula. A slip has been met with passing from the upper border about the middle of the posterior fold of the axilla, across the vessels and nerves to unite with the under surface of the tendon of the pectoralis major, coraco-brachialis, or the fascia over the biceps. This slip passing over the position usually selected for applying the ligature, might mislead the surgeon in the operation of tying the axillary artery. *Action*—To depress the shoulder, bring the arm backwards and inwards; rotate the os-brachia

inwards, so as to turn the palm of the hand backwards. It also assists in raising the body, as in climbing, and may also come into play during respiration.]

Teres Major.—This muscle lies behind the latissimus dorsi, closely connected with it, and assists in forming the posterior boundary of the axilla. It *arises* from the lower angle of the back of the scapula, and is inserted by a broad flat tendon into the posterior margin of the bicipital groove of the humerus. A bursa or sac containing water to prevent friction intervenes between this tendon and that of the latissimus dorsi. The action of this and the preceding muscle is to draw the humerus inwards and backwards.

Subscapularis.—This muscle occupies the internal surface of the scapula. It *arises* from the internal surface of the scapula, and terminates in a strong tendon, which passes under the axillary vessels and nerves, over the inner side of the shoulder-joint, and is *inserted* into the lesser tuberosity of the humerus. Its action is to rotate the humerus inwards.

Serratus Magnus.—This muscle covers the side of the chest like a girth. It *arises* by digitations from the eight upper ribs. Its fibres converge and are *inserted* into the posterior border of the scapula. Its action is to draw the scapula forwards: but of this more hereafter. It is supplied by the following nerve which you see on its outer surface:—

External Respiratory Nerve (of Sir C. Bell.) — This nerve supplies the serratus magnus only. It comes from the fifth and sixth cervical nerves; and runs behind the axillary vessels to the outer surface of the serratus magnus: each digitation receiving a separate filament.*

Having surveyed the muscles which bound the axilla, your next study must be the axillary artery and its branches. But before you begin, notice well the position and direction of the coraco-brachialis muscle; and, if your first essay, reflect the subclavius and pectoralis minor from their insertions.

Course and Relations of the Axillary Artery.—This artery, a continuation of the subclavian, takes the name of “axillary” at the outer border of the first rib. It runs down-

* It may be asked why this nerve is called the “external respiratory?” It was so named by Sir C. Bell, who considered the serratus magnus as *the* external respiratory muscle.

wards and outwards, through the upper part of the axilla, beneath the two pectoral muscles, and along the inner edge of the coraco-brachialis, as far as the lower border of the tendon of the teres major, where it takes the name of *brachial*. In its course downwards, the artery lies successively upon the first intercostal space, the second digitation of the serratus magnus, the loose tissue of the axilla, the tendon of the subscapularis; lastly, upon the tendon of the latissimus dorsi and teres major.

[The axillary artery may give off a much larger branch than usual (one case in ten). This branch then usually forms one of the arteries of the fore-arm, — most frequently the vertical; this vessel may give origin to the subscapula, the two circumflex and the two profundæ, and the brachial plexus may surround this large branch, rather than the main vessel.]

The *axillary vein* in the first part of its course lies in front of the artery, and close to its sternal side: in the lower two-thirds of its course the vein still lies to the sternal side of the artery, but is separated from it by some of the nerves of the brachial plexus.

The axillary plexus, consisting of two or three large nerves is at first situated above the artery, and on a posterior plane;

Fig. 41.




DIAGRAM OF THE BRACHIAL PLEXUS OF NERVES, AND THEIR RELATION TO THE AXILLARY ARTERY.

- | | |
|---------------------------------------|---|
| 1. Ulnar. | 8, 9. Anterior thoracic n. to pectoral muscles. |
| 2. Internal cutaneous. | |
| 3. Lesser cutaneous (n. of Wrisberg.) | 10. N. to serratus magnus. |
| 4. Musculo-spiral. | 11. Median. |
| 5. N. to latissimus dorsi. | 12. Circumflex. |
| 6. N. to teres major. | 13. External cutaneous. |
| 7. N. to subscapularis. | 14. Supra-scapular. |
| | 15. Posterior scapular. |

but the nerves, as they descend, come into closer connection with it, and subdivide in such a way, that, upon the tendon of the subscapularis the artery is surrounded on all sides by nerves.

Axillary or Brachial Plexus of Nerves.—This *plexus* is formed by the anterior branches of the four lower cervical and first dorsal nerves. The plexus is broad at the lower part of the neck, between the scaleni muscles; but it gradually contracts as it descends beneath the clavicle into the axilla. The arrangement of the nerves in the formation of the plexus is generally as follows:—the fifth and sixth cervical unite to form a single cord; the eighth cervical and the first dorsal unite to form another; the seventh cervical runs for some distance alone, and then divides so as to unite with the two other cords. Thus, the plexus consists at the lower border of the first rib of only two large nerves; the one being situated behind the axillary artery, the other above it, and on its acromial side.

Now these two large nerves, by their subdivisions, form a plexus round the axillary artery, where it lies upon the tendon of the subscapularis, and give off the nerves to the arm in the following manner:—

Arrangement of the Nerves round the Axillary Artery.—The *median nerve* arises by two roots, which lie in front of the artery, and converge like the branches of the letter V. On the *outer side* of the artery is the external root of the median, and the external cutaneous nerve; on its *inner side* is the internal root of the median, the ulnar, and the internal cutaneous: *behind* the artery is the circumflex, and the musculo-spiral or radial nerve.*

The axillary plexus gives off some branches above the cla-

* From this arrangement there are occasional deviations. For instance, the

vicle, namely, the supra-scapular, posterior scapular, anterior thoracic, long thoracic, &c.: these were dissected with the neck (p. 49). Below the clavicle, it gives off the following:—

- | | |
|----------------------------|-----------------------------------|
| 1. The external cutaneous. | 5. The lesser internal cutaneous. |
| 2. The median. | 6. The musculo-spiral. |
| 3. The ulnar. | 7. The circumflex. |
| 4. The internal cutaneous. | 8. The subscapular. |


Branches of the Axillary Artery. — The number and origin of these branches often vary, but their general course is in most cases similar; and they usually arise in the order in which they are here described:—

The *thoracic axis* arises above the pectoralis minor, and divides into branches which have been already noticed (p. 200).

The *inferior* or *long thoracic artery*, sometimes called the external mammary, runs along the lower border of the pectoralis minor. It supplies the mammary gland, the serratus magnus and pectoral muscles, and maintains a free anastomosis with the short thoracic, internal mammary, and intercostal arteries.

The *subscapular* is the largest branch of the axillary; it arises opposite the lower border of the subscapularis, and soon divides into an *anterior* and a *posterior* branch.

Fig. 42.



PLAN OF THE BRANCHES OF THE AXILLARY ARTERY.

two roots of the median may embrace the artery higher up in its course, or lower down than usual: or both its roots may lie behind the artery.

A deviation, one which I observed once only in 300 arms, may occur, in which all the nerves are situated above the artery, and on its acromial side, the vessel being entirely free in the whole of its course.

- | | |
|------------------------------|--------------------------|
| 1. Thoracic axis, giving off | 6. Sub-scapular. |
| 2. Short thoracic. | 7. Dorsalis scapulæ. |
| 3. Thoracica acromialis. | 8. Anterior circumflex. |
| 4. Thoracica humeraria. | 9. Posterior circumflex. |
| 5. Long thoracic. | |

a. The *anterior branch* is the continued trunk; it runs along the anterior edge of the subscapularis towards the lower angle of the scapula. Its numerous branches supply the subscapularis, latissimus dorsi, and serratus magnus, and anastomose with the intercostal and thoracic arteries, as well as the posterior scapular (a branch of the subclavian).

b. The *posterior branch* (dorsalis scapulæ), runs to the back of the scapula, through a triangular space bounded by the long head of the triceps, the teres major and subscapularis (5, fig. 43). On the back of the scapula, it divides into branches, which ramify close to the bone, supplying the infraspinatus and teres minor. But its largest branch inosculates freely with the supra-scapular artery (from the subclavian). This is the principal channel through which the collateral circulation is carried on after ligature of the subclavian in the third part of its course. (See diagram, p. 47.)

The *subscapular vein* empties itself into the axillary vein.

Subscapular Nerves.—Three nerves called "*subscapular*" are found on the surface of the subscapularis. They come from the posterior part of the axillary plexus, and supply respectively the latissimus dorsi, teres major, and subscapularis. The *nerve for the latissimus dorsi* (called the *long subscapular*), runs with the anterior branch of the subscapular artery.

The *nerve for the teres major* is either a branch of the preceding, or comes distinct from the plexus. It lies nearer to the humerus than the long scapular.

The proper *nerve of the subscapularis* arises from the plexus higher than the others.

Circumflex Arteries Anterior and Posterior.—There are two circumflex arteries—an *anterior* and a *posterior*, so called from the manner in which they encircle the neck of the humerus. The *posterior circumflex artery* is as large as the subscapular, close to which it is given off; or both may arise by a common trunk from the axillary. It passes backwards

through a quadrilateral space, bounded above by the subscapularis, below by the teres major, externally by the neck of the humerus, and internally by the long head of the triceps (4, fig. 43). It then winds round the back of the neck of the humerus, and is distributed to the under surface of the deltoid.

Besides the deltoid, the posterior circumflex artery supplies the long head of the triceps, the head of the humerus, and the shoulder-joint. It inosculates above with the acromial thoracic artery, below with the ascending branch of the superior profunda (a branch of the brachial), and in front with the anterior circumflex artery. If you cannot find the posterior circumflex artery in its normal situation, look for it (as a branch of the brachial) below the tendon of the teres major.

Fig. 43.




DIAGRAM OF THE ORIGINS OF THE TRICEPS.

- | | |
|--|--|
| 1. Subscapularis. | 5. Triangular space for dorsalis scapulae a. |
| 2. Teres major. | |
| 3. Long head of triceps. | 6. Space for musculo-spiral n., and superior profunda a. |
| 4. Square space for circumflex a. and n. | |

Circumflex Nerve.—The posterior circumflex artery is accompanied by the circumflex nerve which supplies the deltoid. This large nerve comes from the posterior part of the brachial plexus, in common with the musculo-spiral, and, after sending a branch to the teres minor, terminates in the under surface of the deltoid. The nerve supplies, also, the skin covering the deltoid by branches, which turn round the posterior border of the muscle.

The *anterior circumflex artery*, much smaller than the posterior, runs in front of the neck of the humerus, above the tendon of the latissimus dorsi. It passes directly outwards beneath the coraco-brachialis and short head of the biceps, close to the bone, and terminates in the under surface of the deltoid, where it inosculates with the posterior circumflex artery.

The most remarkable branch of the anterior circumflex artery is the one which runs with the long tendon of the biceps up the groove of the humerus. It is called on that account, the *bicipital artery*; it supplies the shoulder-joint and the neck of the humerus.

The *alar thoracic artery* is a small branch, variable in its origin. It may come from the axillary, or the subscapular, or the inferior thoracic. It supplies the axillary glands.

[The subscapular may give off some of the vessels usually derived from the axillary, as the posterior circumflex or large thoracic. It may also arise by two trunks, and the dorsalis scapulæ arise from the axillary artery. The posterior circumflex may arise from the superior profunda, then it ascends behind the tendons of the latissimus dorsi and teres major. This vessel may give off the anterior circumflex, superior profunda, dorsal scapula, &c. We may meet also with two posterior, as well as superior profunda. The latter variety is not so frequent.]

Axillary Vein.—The *axillary vein* is formed by the junction of the *vena comites* of the brachial artery, near the lower border of the subscapularis. It receives the subscapular and the other veins corresponding to the branches of the axillary artery, with the exception of the circumflex, which usually join, either the subscapular, or one of the *venæ comites*. The axillary also receives the cephalic, and sometimes the basilic vein.

Dissection of the Upper Arm.—Let the incision be continued down the inner side of the arm as low as two inches below the elbow. Reflect the skin, and trace the following cutaneous nerves:—

Cutaneous Nerves.—The filaments of the *intercostal-hu-*

meral nerves run down the inner and posterior part of the arm to the olecranon.

The branches of the *internal cutaneous nerve* perforate the fascia about the middle of the inner side of the arm, and subdivide into filaments which supply the anterior and posterior surface of the forearm.

The *lesser internal cutaneous (nerve of Wrisberg,)* perforates the fascia about the lower third of the arm, and supplies the skin over the internal condyle and olecranon.

The *internal cutaneous branch of the musculo-spiral nerve*, sometimes wanting, and always small, comes through the fascia near the middle of the inner side of the arm.

The nerves which perforate the fascia near the middle of the *outer* part of the arm, are the *external cutaneous branches of the musculo-spiral*. They are accompanied by a small branch from the superior profunda artery. They divide into filaments, one of which will be traced down the outer and back part of the forearm to the wrist.

On the outer side of the tendon of the biceps, you find that the *external cutaneous nerve* perforates the fascia, and divides into many branches, which supply the skin of the outer part of the forearm.

Disposition of Veins in front of the Elbow.—The next object of attention should be the disposition of the veins in front of the elbow, where venesection is usually performed. In cleaning these veins, take care not to divide the branches of the internal and external cutaneous nerves which pass both above and below them.

The following is the ordinary arrangement of the superficial veins at the bend of the elbow:—On the outer side is the radial; on the inner side is the ulnar vein; in the centre is the median, which divides into two branches; the external of which, uniting with the radial to form the cephalic vein, is called the *median cephalic*; the internal, uniting with the ulnar to form the basilic, is named the *median basilic*. Near its bifurcation, the median vein communicates by a branch (*medi-ana profunda*) with the deep veins which accompany the arteries of the forearm.


Trace the *cephalic vein* up the arm. It runs along the outer

border of the biceps to the groove between the pectoralis major and the deltoid, where it terminates in the axillary.

The *basilic vein* ascends along the inner side of the arm with the internal cutaneous nerve. Near the upper third of the arm, it perforates the fascia, and empties itself either into the internal vena comes of the brachial artery or into the axillary vein.

Relation of Nerves and Veins at the Elbow.—The principal branches of the nerves pass beneath the veins; but

Fig. 44.



SUPERFICIAL VEINS AND NERVES AT THE BEND OF THE LEFT ELBOW.

many small filaments cross in front which are exposed to injury in venesection.

Relation of Median Basilic Vein to Brachial Artery.—Since the median basilic vein is larger than the median cephalic, and, on account of the strong fascia beneath, more easily compressible, it is usually chosen for venesection: its position, therefore, in reference to the brachial artery, becomes very important. The vein is only separated from the artery by the fascia derived from the tendon of the biceps.

This fascia is in some subjects remarkably thin, or even absent. It sometimes happens that the artery lies above the fascia, in absolute contact with the vein. In choosing, therefore, this vein for venesection, there is a risk of wounding the artery: hence the practical rule, to bleed either from the median cephalic, or from the median basilic above the situation where it crosses the brachial artery.

[The student will often notice that these veins differ from the arrangement above described. Sometimes the median vein will pursue a distinct course up the arm and enter the brachial or axillary.]

Immediately above the internal condyle, in the neighborhood of the basilic vein, we find one or two small *absorbent glands*. Others may lie higher up along the inner side of the arm. I have seen a gland at the bend of the elbow: but never below this joint. These little glands are the first which become tender and enlarged after an injury to the hand.

Muscular Fascia.—The fascia which invests the muscles of the upper arm is a continuation of the fascia from the trunk and the axilla. This membrane varies in density: thus it is thin over the biceps, stronger on the inner side of the arm to protect the brachial vessels and nerves, and strongest over the triceps. What are its connections? At the upper part of the arm it is connected with the coracoid process and the clavicle; it is strengthened at the axilla by an expansion from the tendons of the pectoralis major and latissimus dorsi: posteriorly it is attached to the spine of the scapula. The fascia surrounds the brachial vessels with a sheath, and furnishes partitions which separate the muscles from each other. Of these partitions the most marked are the *external* and *internal intermuscular septa*, which divide the muscles on the anterior from that on the posterior surface of the upper arm. These septa are attached to the projecting ridges of the humerus and to the condyles. The *internal septum*, the most prominent of the two, begins at the insertion of the coracobrachialis, and separates the triceps extensor from the brachialis anticus. The *external septum* commences from the insertion of the deltoid, and separates the brachialis anticus, the supinator longus, and the extensor carpi radialis longior in front, from the triceps behind.

At the lower part of the upper arm the fascia is remarkably strong, especially where it covers the brachialis anticus and the brachial vessels, and is continued over the muscles on the inner side of the forearm. At the back of the elbow, the fascia is attached to the tendon of the triceps, and the olecranon.

Now remove the fascia in order to study the muscles on the front of the arm; namely, the biceps, the coraco-brachialis, and the brachialis anticus.

Biceps.—The biceps, as its name implies, arises by two heads—a long and a short. The *short* or internal *head* arises from the point of the coracoid process of the scapula, by a flat tendon which is common to a slender muscle on its inner side, called the coraco-brachialis. The *long head* of the biceps arises from the upper border of the glenoid cavity of the scapula by a long flat tendon, which traverses the shoulder-joint and passes over the head of the humerus, and down the groove between the two tuberosities. The tendon is retained in the groove by a fibrous bridge derived from the capsule of the joint, and connected with the tendon of the pectoralis major. Divide this bridge, and you will see that the synovial membrane of the joint is reflected round the tendon, and accompanies it for about two inches down the groove, thus forming a sort of synovial fold. The object of this is to facilitate the play of the tendon, and to carry little arteries (from the anterior circumflex) for its supply. The two heads unite about the middle of the arm, and form a single muscle, which terminates in a strong flat tendon of considerable length; this sinks deep into the triangular space at the bend of the elbow, and, after a slight twist upon itself, is *inserted* into the posterior part of the tubercle of the radius. The anterior part of the tubercle, over which the tendon plays, is crusted with cartilage, and a *bursa mucosa* intervenes to prevent friction. The most internal fibres of the muscle are inserted into a *strong, broad aponeurosis*, which is prolonged from the inner border of the tendon to the fascia on the inner side of the forearm. This aponeurosis, called the *semilunar fascia of the biceps*, protects the brachial vessels and the median nerve at the bend of the elbow.

[We meet with several varieties in this muscle. Its two parts

may remain separate as far as the elbow—it may have several heads. In one case we have seen the biceps on one side arise from two distinct tendinous heads, from the upper border of the glenoid cavity, these tendons passing through special sheaths in the capsular ligament.]

The *action* of the biceps is twofold. 1. It is a flexor of the forearm: 2. It is a supinator of the forearm, in consequence of its insertion into the *posterior* part of the tubercle of the radius. Its power of supination is greatest when the arm is bent, because its tendon forms a right angle with the radius. Why does the long tendon pass through the shoulder-joint? It acts like a strap, and confines the head of the humerus in its proper centre of motion. But for this tendon the head of the bone, when the deltoid acts, would be pulled directly upwards and strike against the under surface of the acromion. When the tendon is ruptured or dislocated from its groove, a man can move his arm backwards and forwards, but he cannot raise the smallest weight. The biceps is supplied with blood by an artery (from the brachial), which runs into the middle of it, and then divides into ascending and descending branches. Its nerve comes from the internal cutaneous.

Coraco-brachialis.—This thin muscle is situated at the upper part of the arm, and runs parallel to the inner border of the short head of the biceps. It *arises* by fleshy fibres from the point of the coracoid process, in common with the short head of the biceps, and from a fibrous septum which lies between them. The muscle terminates in a flat tendon, which is *inserted* into the inner side of the middle of the humerus, between the brachialis anticus and the inner head of the triceps. Its action is to draw the humerus forwards and inwards, *e. g.* in bringing the gun up to the shoulder.

Concerning the coraco-brachialis, remember, 1. that the external cutaneous nerve runs through it; 2. that its inner fleshy body is the guide to the axillary artery in the last part of its course; 3. that the brachial artery lies upon its flat tendon of insertion, and can here be effectually compressed by the finger or the tourniquet.

The coraco-brachialis and biceps are covered at their upper part by the deltoid and pectoralis major. The head of the humerus rolls beneath the coraco-brachialis and short origin

of the biceps; and a large *bursa* is interposed between these muscles and the tendon of the subscapularis, which covers the head of the bone.

Brachialis-anticus.—This muscle is situated close to the lower half of the humerus, and is partially concealed by the biceps. Between the two muscles you will find the external cutaneous nerve, which supplies them both.

It *arises* from the humerus by a fleshy digitation on either side the tendon of the deltoid, from the front surface of the bone below this point, and from the intermuscular septa. The muscle, becoming thicker and broader as it descends, covers the front of the capsule of the elbow-joint, and terminates in a tendon, which is *inserted* in a pointed manner into the coronoid process of the ulna. Its action is to bend the forearm.

Now examine the course and relations of the brachial vessels and nerves.

Course and Relations of the Brachial Artery.—The brachial artery, a continuation of the axillary, takes its name at the lower border of the *teres major*. It runs down the inner side of the arm, along the *inner border of the coracobrachialis and biceps*, to the front of the elbow, where it divides near the coronoid process of the ulna into the radial and ulnar arteries.

Thus its direction corresponds with a line drawn from the anterior part of the axilla to the central point between the condyles of the humerus.

In the upper part of its course it lies on the triceps (from which it is separated by the musculo-spiral nerve and superior profunda artery): in the middle it lies on the tendon of the coracobrachialis; in the lower part, on the brachialis anticus.

The artery is accompanied by two veins (*venæ comites*), and the median nerve, all of which are invested in a common sheath of fascia. The median nerve crosses obliquely in front of the artery, lying, near the axilla, on its outer side, near the elbow on its inner side.

The ulnar nerve runs along the inner side of the artery, and is separated from it below by the intermuscular septum. Superficial to the artery, we find the internal cutaneous nerve and the basilic vein.

Observe particularly that the artery is more or less overlapped, in the first part of its course by the coraco-brachialis, lower down by the fleshy belly of the biceps; these muscles in their respective situations are the best guides to the vessel.

About the middle of the humerus, the artery lies for nearly two inches on the tendon of the coraco-brachialis, and is so close to the bone that it can here be effectually compressed; here too it is crossed by the median nerve.


At the bend of the elbow the artery is protected by the semilunar fascia from the tendon of the biceps. It enters a triangular space, bounded by the pronator radii teres internally, and by the supinator radii longus externally. It sinks into this space, with the tendon of the biceps to its outer side, and the median nerve to its inner side; all three rest upon the brachialis anticus muscle. Opposite the coronoid process of the ulna it divides into the radial and ulnar arteries.

[The brachial has been known to descend together with the median nerve, towards the inner condyle as far as the origin of the pronator teres, then to pass through the muscle to regain its usual position at the elbow. This is the course of the vessel in some carnivorous animals. It may divide at almost any point above its usual position; *very rarely* below its usual point of bifurcation; the branch thus given off being the radial in three cases out of four. In one instance only have the radial, ulnar, and interosseous been reported as coming off the radial above the elbow-joint. The arteries thus given off usually run close together. The radial often arising from the inner side of the brachial, and then for a short distance runs parallel with the larger vessel, and crosses over it, opposite to the bend of the elbow, under cover of the fascia; it has been found perforating the fascia and passing under the skin near the elbow. When the ulnar arises high up, it inclines at the lower part of the arm from the brachial towards the inner condyle; it usually lies beneath the fascia and superficially to the flexor muscles. When the interosseous arises from the brachial, it is usually situated behind this vessel. The two unusual arteries either communicate by a branch, or unite. These anomalies are often of surgical importance.]

Two veins, of which the internal is the larger, lie in close contact with the brachial artery, and communicate at frequent intervals by transverse branches.

Branches of the Brachial Artery.—The brachial artery gives off only three branches of importance; all arise from its

Fig. 45.



PLAN OF THE BRANCHES OF THE BRACHIAL ARTERY AND THE ARTERIAL INOSCULATIONS ABOUT THE RIGHT ELBOW-JOINT.

inner side; they are called the profunda superior, the profunda inferior, and the anastomotica magna.

The *profunda superior* arises from the brachial artery, immediately below the tendon of the teres major.* It winds round

* If you cannot find the profunda in its usual place, look for it above the tendon of the latissimus dorsi, where you will probably find that it is given off from a common trunk with the posterior circumflex.

the posterior part of the humerus, between the outer and inner heads of the triceps, with the musculo-spiral nerve (6, p. 211), and a little above the middle of the arm divides into two branches, which run for some distance on either side of the nerve. One of these runs in the substance of the triceps to the olecranon, and anastomoses with the ulnar and interosseous recurrent arteries; the other branch accompanies the musculo-spiral nerve to the outer side of the arm, descends deep in the fissure between the brachialis anticus and supinator radii longus, and terminates in numerous ramifications, some of which pass in front of the external condyle, and others behind it, to inosculate with the radial and interosseous recurrent arteries.

Before its division the superior profunda sends several branches to the triceps, some of which inosculate minutely with the circumflex arteries. These would assist in establishing a collateral circulation if the brachial artery were tied above the origin of the profunda.

The *profunda inferior* arises from the brachial, opposite to the insertion of the coraco-brachialis, or sometimes by a common trunk with the superior profunda. It runs with the ulnar nerve along the inner border of the triceps behind the internal condyle of the humerus, where it inosculates with the posterior ulnar recurrent artery.

The inferior profunda gives branches to the triceps and brachialis anticus, some of which inosculate with the anastomotica magna.

The *medullary artery* of the humerus usually arises from the profunda inferior. It pierces the tendon of the coraco-brachialis, runs obliquely downwards through the bone, and in the medullary canal divides into ascending and descending branches, which anastomose with the other nutrient vessels of the bone derived from the periosteum.

The *anastomotica magna* arises from the inner side of the brachial, about two inches above the elbow, runs tortuously inwards across the brachialis anticus, and divides into branches, some of which pass in front of, others behind the internal condyle, anastomosing with the inferior profunda and the anterior ulnar recurrent arteries.

Numerous unnamed *muscular branches* arise from the outer

side of the brachial artery; one of these, more constant than the rest, supplies the biceps; another runs transversely beneath the coraco-brachialis and biceps, over the insertion of the deltoid, supplying this muscle and the brachial anticus.

Venæ Comites.—The two veins which accompany the brachial artery are continuations of the deep radial and ulnar veins. The internal is usually the larger, since it generally receives the veins corresponding to the principal branches of the artery. In their course they are connected at intervals by transverse branches either in front of, or behind the artery. Near the subscapularis, the *vena comes externa* crosses obliquely over the front of the axillary artery to join the *vena comes interna*, which then takes the name of “axillary.”

Let us now trace the great nerves of the upper arm, which proceed from the brachial plexus near the tendon of the subscapularis; namely, the median, external cutaneous, ulnar, and musculo-spiral or radial.

Median Nerve.—The median nerve arises by two roots, which converge in front of the axillary artery (p. 208). The external root is derived from a trunk in common with the external cutaneous nerve, the internal root from a trunk in common with the ulnar and internal cutaneous nerve. In its course down the arm, the nerve is situated at first on the outer side of the brachial artery, between it and the coraco-brachialis muscle: about the middle of the arm the nerve crosses obliquely over the vessel, or perhaps beneath it, so that at the bend of the elbow it is found on the inner side of the artery, covered by the semilunar fascia from the biceps.*

* I have observed the following *varieties* relating to the median nerve, and its course in regard to the artery:—

a. The roots may be increased in number by one on either side of the artery; or the internal root may be deficient.

b. They may vary in their position with regard to the artery; both may be situated behind the vessel, or one behind, and the other in front of it.

c. The nerve, formed in the usual manner, may be joined lower down by a large branch from the external cutaneous; such a case presents a junction of two large nerves in front of the brachial artery, in the middle of the arm.

d. The nerve in many cases crosses under, instead of over the artery.

e. The nerve sometimes runs parallel and external to the artery; or it may run parallel to, and in front of the artery.

In one hundred arms the relative position of the nerve to the artery in its course down the arm was as follows:—

The eventual distribution of the median nerve is to the two pronators and all the flexors of the forearm (except the flexor carpi ulnaris and half the flexor profundus digitorum), to the muscles of the ball of the thumb, and to both sides of the thumb, fore, and middle fingers, and the radial side of the ring finger.

External Cutaneous Nerve.—This nerve (often called musculo-cutaneous, or perforans Casserii) arises by a common trunk with the external root of the median, on the outer side of the axillary artery. It perforates the coraco-brachialis, and then runs down between the biceps and the brachialis anticus. A little above the elbow-joint, between the tendon of the biceps and the supinator radii longus, the nerve becomes subcutaneous, and, passing under the median cephalic vein, divides into branches, for the supply of the integuments of the forearm.

The external cutaneous nerve, in the upper part of its course, sends branches to the coraco-brachialis and the short head of the biceps, and, as it descends between the biceps and brachialis anticus, it supplies both these muscles. Consequently, if the external cutaneous nerve were divided in the axilla, the result would be inability to bend the forearm.*

Ulnar Nerve.—This nerve arises from the plexus in com-

In 72, the nerve took the ordinary course.

" 20, the nerve crossed obliquely under the artery.

" 5, the nerve ran parallel and superficial to the artery.

" 3, the nerve ran parallel and external to the artery.

These varieties of the median nerve are of practical importance for this reason: whenever, in the operation of tying the brachial artery, we find the nerve out of its normal position, we may expect to find some irregular distribution of the arteries, *e. g.* a high division of the brachial, or even, which I have often seen, a "vas aberrans" coming from the upper part of the brachial, and joining either the radial or ulnar arteries.

* In some instances the external cutaneous nerve descends on the inner side of the coraco-brachialis without perforating the muscle; in these cases it often sends a larger branch than usual to the median nerve.

The trunk of the external cutaneous nerve may come from the median at any point between the axilla and the middle of the arm. In some subjects the nerve is absent: all its branches are then supplied by the median, which is larger than usual. Such anomalies are easily explained by the fact of the two nerves having always a common origin.

mon with the internal cutaneous, and the inner root of the median. It descends along the inner side of the brachial artery, as far as the insertion of the coraco-brachialis. The nerve then diverges from the artery, perforates the internal intermuscular septum, and runs with the inferior profunda artery, behind the internal condyle.

The eventual distribution of the nerve is to the flexor carpi ulnaris, half the flexor profundus digitorum, all the interosseous muscles of the hand, both sides of the little finger, and the ulnar side of the ring finger, also to the muscles of the ball of the little finger.

Previous to the examination of the musculo-spiral nerve we should have some knowledge of the great muscle which occupies the whole of the posterior part of the humerus—viz. the triceps.

Triceps Extensor Cubiti.—This muscle has three distinct origins, named from their position the *external*, *internal*, and *middle* or *long* head (p. 211). The *middle* or *long* head arises by a flat tendon from the inferior border of the scapula, close to the glenoid cavity. The *external* head arises from the humerus, immediately below the great tuberosity. The *internal* head arises from the humerus below the insertion of the teres major. The three heads of the muscle unite near the middle of the arm to form a single fleshy mass, which covers the posterior part of the elbow-joint, and is *inserted* by a thick tendon into the summit and sides of the olecranon.

[The use of this muscle is to extend the forearm and to carry the arm backwards; it also draws up the synovial membrane from between the olecranon process and os-brachia, so as to preserve it from pressure in the extended position of the arm. The long head also protects the inferior part of the capsular ligament, and tends to prevent dislocation at that joint, where it would be most likely to occur. This muscle also comes into play in fracture of the olecranon; it draws the superior portion of bone up from the inferior portion.

Musculo-spiral or Radial Nerve.—This, the largest of the brachial nerves, arises, in common with the circumflex, from the posterior part of the axillary plexus (p. 208). It descends at first behind the axillary artery, and then winds obliquely round the posterior part of the humerus, between the external

and internal heads of the triceps, in company with the superior profunda branch of the brachial artery. About the lower third of the outer side of the arm, the nerve runs deeply imbedded between the brachial anticus and the supinator radii longus. A little above the elbow-joint it divides into its two principal branches,—the *radial*, which accompanies the radial artery along the forearm,—and the posterior interosseous, which perforates the supinator brevis, and supplies all the muscles on the back of the forearm.

What is the distribution of this great nerve? In a word, it supplies *all* the extensors of the forearm, wrist, thumb, and fingers; and all the supinators except one, namely the biceps (supplied by the external cutaneous).

DISSECTION OF THE FRONT OF THE FOREARM.

PROLONG the incision down to the wrist, reflect the skin, and trace the cutaneous nerves and veins (p. 214).

Cutaneous Nerves.—On the *radial* side of the forearm are the superficial radial veins, with filaments of the external cutaneous nerve. On the *ulnar* side are the superficial ulnar veins, with filaments of the internal cutaneous nerve. The external cutaneous branch of the musculo-spiral nerve runs along the outer and back part of the forearm. About the lower third of the radial border of the forearm, the radial nerve becomes superficial, and turns over the radius to supply the back of the hand and fingers. Near the styloid process of the ulna, the superficial branch of the ulnar nerve perforates the fascia to reach the back of the hand.

Cutaneous Veins.—The veins of the hand are placed on its dorsal surface, to be out of the way of pressure. Commencing at the extremity of the fingers, the veins run up *between* the knuckles and unite on the back of the hand in the shape of an arch, with its concavity upwards, from which the veins of the forearm arise.

From the external side of the arch, near the metacarpal

bone of the thumb, the superficial *radial* veins run up the outer and front part of the forearm to the bend of the elbow.

From the internal side of the arch, near the metacarpal bone of the little finger, the superficial *ulnar* veins proceed along the inner and front side of the forearm to the elbow.

Along the middle of the forearm runs the median vein, which is formed by the superficial veins in front of the wrist, and terminates at the elbow by joining the cephalic and the basilic veins.

Fascia of the Forearm.—The fascia which envelopes the muscles of the forearm is much stronger than that of the upper arm, and its strength increases as it approaches the wrist, that the tendons in this situation may be effectually maintained in their proper place.

The fascia is attached to the condyles of the humerus, and is strengthened by fibres from the tendons of the biceps and brachialis anticus.

The aponeurotic expansion from the inner edge of the tendon of the biceps is exceedingly strong. It braces the muscles on the inner side of the arm, and interlaces at right angles with the fibres of the fascia attached to the internal condyle.

Along the forearm the fascia is attached to the ridge on the posterior part of the ulna. At the back of the wrist it forms the posterior annular ligament. On the front of the wrist it is continued into the anterior annular ligament.

The under surface of the fascia gives origin to the muscular fibres in the upper part of the forearm, and furnishes septa, which separate the muscles and form so many distinct sheaths for them.

Remove the fascia from the muscles, and preserve them as much as possible in their proper relative position. But before we examine them in detail, let us direct our attention to the important triangular space at the bend of the elbow.

Triangle at the Bend of the Elbow and Contents.—The pronator teres on the ulnar side, and the supinator longus on the radial, form two sides of a triangular space at the bend of the elbow, of which you must carefully make out the contents; namely:—1. The brachial artery (with its companion veins), dividing into the radial and ulnar; 2. The tendon of the

biceps on its outer side; 3. The median nerve close to its inner side,* 4. The radial recurrent artery; 5. The anterior ulnar recurrent artery; 6. The common interosseous branch of the ulnar artery; 7. The musculo-spiral nerve, deep in the groove between the brachialis anticus and the supinator radii longus, where it divides into the "radial," which accompanies the radial artery, and the posterior interosseous, which perforates the supinator brevis, and supplies the muscles on the back of the forearm.

Muscles of the Forearm.—The muscles of the forearm are arranged in two groups: one group, consisting of pronators and flexors, is attached to the internal condyle; the other, consisting of supinators and extensors, is attached to the external. Examine the inner group first, in the order in which they arise.

Pronator Radii Teres.—This muscle bounds the inner side of the triangular space at the elbow. It *arises* from the anterior surface of the internal condyle, and from the septum between it and the flexor carpi radialis. It has also a small tendinous origin, from the coronoid process of the ulna. From these two origins, which are separated by the median nerve, the muscle proceeds obliquely downwards and is inserted by a flat tendon into a rough surface on the outer and back part of the middle of the radius. In amputating the forearm, it is very desirable to save the insertion of this muscle, that the stump may have a pronator.

Flexor Carpi Radialis.—This muscle *arises* by a thin tendon from the internal condyle, and from the intermuscular septa. The fleshy fibres terminate near the middle of the forearm, in a flat tendon, which runs beneath the anterior annular ligament of the wrist, passes through a groove in the os-trapezium, lined by a synovial membrane, and is *inserted* into the base of the second metacarpal bone. Note that the outer border of its tendon is the guide to the radial artery in the lower half the forearm.

Palmaris Longus.—This slender muscle arises from the tendon of the internal condyle, and from the intermuscular

* The median nerve does not always lie close to the artery; it may be distant as much as half an inch from it. I have seen the nerve lying quite superficial to the artery. I have seen it on the outer side of the vessel.

septa. About the middle of the forearm it terminates in a flat tendon, which descends vertically down the middle of the forearm to the wrist, where it passes over the annular ligament, and is continued into the palmar fascia. This muscle is a tensor of the palmar fascia.*

Flexor Sublimis Digitorum.—This muscle has two distinct origins. The longer origin takes place from the internal condyle, from the coronoid process of the ulna, and from the intermuscular septa: the shorter origin takes place by tendinous and fleshy fibres from an oblique ridge on the front of the radius. This, which is called its *radial origin*, is partly concealed by the pronator teres. The muscle thus formed passes down the middle of the forearm, and divides into four distinct muscular slips: from these, four tendons arise, which pass beneath the annular ligament into the palm, and on to the fingers, where they split to allow the passage of the deep flexor tendons, and are *inserted* into the base of the second phalanges. Its *action*, therefore, is to bend the second joint of the fingers.

[The tendons of the flexor sublimis (perforatus) are enveloped in a large bursa, behind the annular ligament. This bursa also surrounds the deep flexor tendons, and the flexor longus publicis. It ends above and below the annular ligament in a cul-de-sac. It is the seat of that trouble known as *ganglion*, of an hour-glass form, one portion appearing in the palm of the hand, the other in the wrist. It may contain simply synovial fluid, or little bodies like grains of rice.]

Flexor Carpi Ulnaris.—This muscle *arises* from the internal condyle, and from the inner edge of the olecranon: these two origins form an arch, under which the ulnar nerve passes. It also arises from the upper two-thirds of the posterior edge of the ulna, through the medium of the aponeurosis of the forearm. The tendon appears on the radial side of the muscle, about the lower third of the forearm, and receives fleshy fibres on its ulnar side as low as the wrist. It is *inserted* into the os-

* The palmaris longus is sometimes absent. The situation of its muscular portion is subject to variety; sometimes occupying the middle, sometimes the lower third of the forearm. The tendon is in some instances wholly inserted into the anterior annular ligament.

pisiforme, and thence by a strong tendon into the base of the fifth metacarpal bone and the os-unciforme.

The tendon of the flexor carpi ulnaris is the guide to the ulnar artery, which lies close to its radial side, and is *overlapped* by it. As it passes over the annular ligament, the tendon furnishes a fibrous expansion to protect the ulnar artery and nerve.

[The flexor ulnaris not only flexes the hand, but, together with the extensor ulnaris, adducts the hand. Adduction is not as limited as abduction, the ulna being shorter below than the radius.]

Having finished the study of the superficial muscles on the inner side of the forearm, you must notice one of those on the outer side, named supinator radii longus, before you trace the vessels and nerves of the forearm.

Supinator Radii Longus.—This muscle forms the external boundary of the triangular space at the bend of the elbow. It *arises* from the external condyloid ridge of the humerus, commencing a little below the insertion of the deltoid. The muscular fibres terminate about the middle of the forearm in a flat tendon, which is *inserted* into the base of the styloid process of the radius. The inner border of the muscle is *the* guide to the radial artery. It supinates the hand, but acts most powerfully as a flexor of the forearm.

Radial Artery.—The radial artery runs down the radial side of the forearm to the wrist, where it turns over the external lateral ligament of the carpus, beneath the extensor tendons of the thumb, and sinks into the space between the first and second metacarpal bones to form the deep palmar arch. Thus, a line drawn from the middle of the bend of the elbow to the front of the styloid process of the radius indicates its course. In the upper third of the forearm, the artery lies between the pronator teres on the inner, and the supinator longus on the outer side; the fleshy border of the latter overlaps it in muscular subjects. In the lower two-thirds of the forearm the artery is more superficial, and is placed between the tendons of the supinator longus on the outer, and the flexor carpi radialis on the inner side. In its course, it lies successively on the following: first upon the tendon of the biceps; secondly, upon the supinator radii brevis; thirdly, upon the

tendon of the pronator teres ; fourthly, upon the radial origin of the flexor sublimis ; fifthly, upon the flexor longus pollicis ; and, lastly, upon the pronator quadratus. It is accompanied by two veins, which communicate at frequent intervals, and join the venæ comites of the brachial artery at the bend of the elbow.

In the middle third of its course the artery is accompanied by the radial nerve, (a branch of the musculo-spiral) which lies to its outer side. Below this point, the nerve leaves the artery, and passes under the tendon of the supinator longus to the back of the hand.

Thus, in the situation where the pulse is felt, the radial nerve no longer accompanies the artery ; nevertheless, the vessel is not without a nerve, for it is accompanied by a branch of the external cutaneous, which runs superficial to it.

[The radial artery does not vary as much as the ulnar in the forearm. It has been seen lying upon the fibrous expansion from the tendon of the biceps, and the fascia of the forearm, instead of beneath them. In passing around the wrist, it may go over the extensor tendons of the thumb, instead of beneath them. It has been seen to be increased by a branch from the anterior interosseous in front of the wrist, and by the perforating branch of that vessel behind the carpus. An example has been met with where it ceased, after giving off the radial recurrents ; its place near the wrist being supplied by the interosseous.]

The radial artery sends off in the forearm the following branches :—

The *radial recurrent* is given off just below the elbow ; it passes outwards to supply the long and short supinator and the two radial extensors. One of its ramifications runs up with the musculo-spiral nerve, and sometimes forms a delicate inosculation with the superior profunda (p. 220).

The *arteria superficialis volæ* arises from the radial, about half an inch, or more, above the lower end of the radius ; it runs superficially over the anterior annular ligament of the carpus, above or perhaps through the origin of the muscles of the ball of the thumb, into the palm of the hand, where it in-

osculates with the superficial branch of the ulnar, and completes the superficial palmar arch* (p. 238).

The *anterior* and *posterior carpal* arteries are small branches of the radial, which run beneath the tendons, and supply the synovial membrane and bones of the carpus, anastomosing with the interosseous arteries and carpal branches of the ulnar.

Radial Nerve.—The radial nerve, a branch of the musculospiral, is given off above the head of the elbow, deep between the supinator radii longus and brachialis anticus; it descends on the outer side of the radial artery, covered by the supinator radii longus. In the upper third of the forearm the nerve is at some distance from the artery; in the middle third it approaches nearer to it; but in the lower third, the nerve leaves the artery, passes underneath the tendon of the supinator longus, perforates the fascia on the outer side of the forearm, and divides into branches, which supply the back of the thumb, fore, and half the middle finger.

Ulnar Artery.—This artery arises from the brachial at the middle of the elbow, runs obliquely inwards along the ulnar side of the forearm to the wrist, passes over the annular ligament near the pisiforme bone, and entering the palm, forms the superficial palmar arch, by inosculating with the superficialis volæ.

In the upper half of its course the artery describes a gentle curve with the concavity towards the radius, and lies deep beneath the superficial layer of muscles—the pronator teres, flexor carpi radialis, palmaris longus, and flexor sublimis digitorum. In the lower part of its course it descends between the flexor sublimis and flexor carpi ulnaris, of which the tendon partially overlaps it at the wrist. The artery lies for a short distance on the brachialis anticus; in the remainder of its course it lies on the flexor profundus digitorum.

The ulnar nerve is at first separated from the artery by a considerable interval: but about the middle of the forearm it joins the artery, and accompanies it in the rest of its course, lying close to its inner side. Both pass over the anterior an-

* There is great variety in the size and origin of the superficialis volæ; sometimes it is very large, arises higher than usual, and runs to the wrist parallel with the radial; sometimes it is very small, terminating in the muscles of the thumb; or it may be absent.

nular ligament of the carpus, lying close to the pisiform bone,—the nerve being nearer to the bone. A strong expansion from the tendon of the flexor carpi ulnaris protects them in this exposed situation.

Observe particularly the depth of the ulnar artery under the many muscles which cover it in the upper third of its course. In the middle third it is partially overlapped by the flexor carpi ulnaris. In the lower third it lies under the radial border of the flexor carpi ulnaris, which is the *proper guide* to the vessel.

[The ulnar is deficient, as regards size, more frequently than the radial; in such cases the radial compensates for this loss. When the palmar portion is small, it is reinforced by a large median branch. In the upper portion of its course it has been known to receive a large branch from the anterior interosseous.

To tie the radial, in its superior portion, the incision, three inches in length, should be made from the bend of the elbow, obliquely downwards and outwards, avoiding the branches of the median nerve; then, drawing the supinator muscle a little outwards, we will find the vessel. Pass the ligature from the radial to the ulnar side to avoid the nerve.

The ulnar artery is rarely tied in the upper portion of the forearm, on account of its depth beneath the superficial flexors. In tying this vessel the needle must pass from the ulnar to the radial side, so as to avoid the nerve.]

The ulnar artery furnishes the following branches in the forearm:—

The *anterior* and *posterior ulnar recurrent* arise immediately below the elbow-joint,—sometimes by a common trunk, sometimes separately. The *anterior* passes upwards in front of the brachialis anticus, and inosculates with the inferior profunda and anastomotica magna. The *posterior* ascends beneath the flexor sublimis digitorum, to the space between the internal condyle and the olecranon, where it inosculates with the same arteries as the anterior (p. 220).

The *common interosseous* artery arises from the ulnar, about an inch and a half below the division of the brachial; and soon divides into the anterior and posterior interosseous, which we shall examine presently.

A branch (*arteria comes nervi mediani*) almost always accompanies the median nerve. It lies in close contact with the nerve, and sometimes in its very centre: though usually of small size, it may be as large as the ulnar artery itself; and in such a case it passes under the annular ligament with the nerve, to join the palmar arch. This is interesting, because it helps to explain the recurrence of hæmorrhage from a wound in the palm, even after the radial and ulnar arteries have been tied.

The *anterior* and *posterior carpal* branches communicate with corresponding branches from the radial, and supply the synovial membrane and bones of the carpus.

Ulnar Nerve.—This nerve runs behind the internal condyle between the two origins of the flexor carpi ulnaris. In its course down the upper part of the forearm, the nerve is still covered by this muscle, and lies upon the flexor profundus digitorum. About the middle third of the forearm, the nerve joins the ulnar artery, and runs along its inner side over the annular ligament into the palm.

The ulnar nerve supplies only two muscles in the forearm, namely, the flexor carpi ulnaris, and the inner half of the flexor profundus digitorum.

About one inch and a half above the styloid process of the ulna, the nerve gives off a large cutaneous branch to the back of the hand. It crosses under the tendon of the flexor carpi ulnaris, and immediately below the styloid process of the ulna, appears on the back of the hand, where it divides into branches, which supply the back of the little finger, the ring, and half the middle finger.

Median Nerve.—This nerve, at the bend of the elbow, lies on the inner side of the brachial artery. It then passes between the two origins of the pronator teres, and descends along the middle of the forearm, between the flexor sublimis and the flexor profundus digitorum: it enters the palm beneath the anterior annular ligament, and divides into five branches for the supply of the thumb, both sides of the fore and middle fingers, and the outer side of the ring finger.*

* In the upper three-fourths of the forearm the median nerve lies deep between the muscles; but in the lower fourth it is more superficial, and runs very nearly down the centre of the wrist. Its exact position at the wrist is

Immediately below the elbow, the median nerve sends branches to the pronator and all the flexor muscles of the forearm, except the flexor carpi ulnaris and the inner half of the flexor profundus, which are supplied by the ulnar nerve. The *interosseous* branch of the median runs with the anterior interosseous artery, and supplies the flexor profundus digitorum, the flexor longus pollicis, and the pronator quadratus.

Before the median nerve passes beneath the annular ligament, it sends off its *superficial palmar* branch, which passes over the annular ligament and divides into small filaments to supply the skin of the palm and ball of the thumb.

You should now reflect the superficial layer of muscles, to see those more deeply seated. Preserve the principal vessels and nerves.

The deep-seated muscles are the flexor digitorum profundus, and the flexor longus pollicis; beneath both, near the wrist, lies the pronator quadratus. Close to the interosseous membrane run the anterior interosseous artery and nerve.

Flexor Profundus Digitorum.—This is the thickest muscle of the forearm. It *arises* from the upper two-thirds of the anterior surface of the ulna, from the same extent of its internal surface, from the aponeurosis attached to the posterior edge of the ulna, and from the inner two-thirds of the interosseous ligament. About the middle of the forearm it divides into four muscular slips, which terminate in flat tendons. These tendons lie upon the same plane, and pass beneath the annular ligament under those of the superficial flexor, into the palm. On the first phalanx of the fingers, the tendons of the deep flexor perforate those of the superficial, and are *inserted* into the base of the third or ungual phalanx.

[Flexor profundus (perforans). A fasciculus from the tendon of the forefinger often unites with the flexor longus pollicis. This muscle not only bends the last phalanx, but co-operates

between the fleshy border of the outer tendon of the flexor sublimis and the inner border of the tendon of the flexor carpi radialis, and it lies upon one of the tendons of the flexor profundus.

If the tendon of the palmaris longus happen to be broader than usual, it may partially cover the median nerve near the wrist; but most frequently the nerve is immediately beneath the fascia, the tendon lying to its ulnar side.

with the superficial flexor in bending the other phalanges and the wrist.]

Flexor Longus Pollicis.—This muscle is situated on the front surface of the radius. It *arises* from the front surface of the radius between the tubercle and the pronator quadratus, and from the interosseous membrane.* Its tendon proceeds beneath the annular ligament to the last phalanx of the thumb.

[The flexor longus pollicis both flexes and adducts the different joints of the thumb upon the hand, and the hand upon the forearm.]

Pronator Quadratus.—This square muscle *arises* from the lower fourth of the ulna; its fibres pass transversely outwards, and are *inserted* into the lower fourth of the radius. It rotates the radius on the ulna.

[The pronator quadratus has sometimes been absent. In fracture of the lower portion of the radius, the upper fragment is drawn into close relation with the ulna, causing the flexor tendons to be thrust forwards, and giving rise to the projection in the anterior surface of the forearm.]

Interosseus Artery.—Nearly on a level with the insertion of the biceps, the ulnar artery gives off from its outer side the *common interosseous* trunk, which runs backwards for about half an inch, and divides into the *anterior* and *posterior interosseous*.

The *anterior interosseous* artery runs down close to the interosseous membrane, lying between the flexor profundus digitorum and flexor longus pollicis. At the upper edge of the pronator quadratus it divides into two branches; one of which, the smaller of the two, supplies the muscle and the front of the carpal bones, communicating with the anterior carpal arteries from the radial and ulnar; the other, by far the most important, perforates the interosseous membrane, and helps to supply the muscles on the back of the forearm.

The anterior interosseous artery gives off branches to the muscles on either side, and the nutrient arteries which run into the radius and ulna from below upwards, near the centre of the forearm, to supply the medullary membrane.

* Sometimes by a slip from the coronoid process.

Anterior Interosseous Nerve.—This nerve is a branch of the median; it lies close to the artery, supplies the flexor longus pollicis, flexor profundus digitorum, and the pronator quadratus.

DISSECTION OF THE PALM OF THE HAND.

REFLECT the skin from the palm. In doing so, you will observe how closely in the centre of the palm it adheres to the palmar fascia beneath it. On the ball of the little finger and the distal ends of the metacarpal bones, the subcutaneous structure is composed of a dense filamentous tissue, which contains numerous pellets of fat, forming a kind of elastic pad. A similar padding protects the palmar surfaces of the fingers. These cushions on the ends of the fingers defend them in the powerful actions of the hand; they are also useful in subservience to the nerves of touch.

Be careful not to remove a small cutaneous muscle, *palmaris brevis*, situated near the ball of the little finger.

The nerves of the palm are supplied by the palmar branch of the median nerve (p. 234), which passes over the annular ligament; and by others, which run up through minute openings in the palmar fascia with the cutaneous arteries.

Palmar Fascia.—This fascia has a silvery lustre, and, in the centre of the palm, is remarkably dense and strong. It is divided into three portions, a central,—by far the most conspicuous; an external, covering the muscles of the thumb; and an internal, covering the muscles of the little finger. From the deep surface of the fascia two septa dip down, so as to divide the palm into three separate compartments; one for the ball of the thumb, a second for that of the little finger, and a third for the centre of the palm.

The fascia is formed by a prolongation from the anterior annular ligament. It is also strengthened by the expanded tendon of the *palmaris longus*.

The central portion of the fascia is triangular, with the apex at the wrist. About the middle of the palm it splits into four portions, which are connected together by trans-

verse tendinous fibres, extending completely across the palm, and corresponding pretty nearly to the transverse furrow of the skin in this situation.

Examine any one of these four portions of the fascia, and you will find that it splits into two strips which embrace the corresponding flexor tendons, and are intimately connected with the transverse metacarpal ligament. The effect of this is that the flexor tendons of each finger are kept in place in the palm, by a fibrous ring. Between the four divisions of the palmar fascia the digital vessels and nerves emerge, and descend in a line with the clefts between the fingers.

In the hands of mechanics, in whom the palmar fascia is usually very strong, we find that slips of it are lost in the skin at the lower part of the palm, and also for a short distance along the sides of the fingers.

The chief use of the palmar fascia is to protect the vessels and nerves from pressure when anything is grasped in the hand. It also confines the flexor tendons in their proper place.

Between the interdigital folds of the skin, we find aponeurotic fibres to strengthen them, constituting what are called the *transverse ligaments* of the fingers. They form a continuous ligament across the lower part of the palm, in front of the digital vessels and nerves.

Now turn down the palmar fascia towards the fingers, and you will expose the vessels, nerves, and tendons of the palm. The vessels lie above the nerves, and the tendons still deeper. There is an abundance of loose cellular tissue to allow the free play of the tendons. When suppuration takes place in the palm, it is seated in this tissue. Reflect for a moment, what mischief is likely to ensue. The matter cannot get to the surface through the dense palmar fascia, or on the back of the hand; it will therefore run up under the annular ligament, and make its way deep amongst the tendons of the forearm.

Palmaris Brevis.—This small cutaneous muscle is situated on the inner side of the palm. It arises from the inner edge of the central palmar fascia, and terminates in the skin on the inner side of the palm. Its use is to support the pad on the inner edge of the palm: it acts powerfully as we grasp; it

raises the edge of the palm and hollows it, forming the "cup of Diogenes."

Superficial Palmar Arterial Arch.—The ulnar artery having passed over the annular ligament near the pisiform bone, describes a curve across the upper part of the palm beneath the palmar fascia, and, gradually diminishing in size,

Fig. 46.




DIAGRAM OF THE SUPERFICIAL AND DEEP PALMAR ARCHES.

1, 2, 3, 4. Interosseous branches.

inosculates with the superficialis volæ, or some other branch of the radial, to form the superficial palmar arch. The curve of the arch is directed towards the ball of the thumb. How are we to ascertain its exact position in the hand?—most

commonly its greatest convexity descends as low as a horizontal line drawn across the junction of the upper with the middle third of the palm.

In its passage over the annular ligament, the artery is protected in a furrow between the pisiform and unciform bones, and by an expansion from the tendon of the flexor carpi ulnaris to the palmaris longus. The ulnar nerve lies close to its inner side. In the palm, before it crosses the flexor tendons, the artery lies for a short distance upon the muscles of the little finger, and is covered by the palmaris brevis.

Branches of Ulnar Artery in the Palm. — Immediately below the pisiform bone, the ulnar artery gives off the *ulnaris profunda*, which sinks deep into the palm between the abductor and the flexor brevis minimi digiti, to assist in forming the deep palmar arch. It is accompanied by the deep branch of the ulnar nerve.

From the concavity of the arch small branches ascend to the carpus, and inosculate with the other carpal branches of the radial and ulnar arteries.

From the convexity of the arch arise *four digital arteries*, which supply all the fingers excepting the radial side of the index finger. You should observe carefully the course of these arteries, because you may have to open abscesses in the palm. The first descends over the muscles on the inner side of the palm, to the ulnar side of the little finger, along which it runs to the apex. The second, third, and fourth descend nearly vertically between the tendons, *in a line with the clefts* between the fingers; and about half an inch above the clefts each divides into two branches, which proceed along the opposite sides of the fingers nearly to the end of the last phalanges, where they unite to form an arch with the convexity towards the end of the finger; from this arch numerous branches supply the papillæ of the skin.

In the palm of the hand the digital arteries before they divide are joined by a small branch from the corresponding interosseous artery, (a branch of the deep palmar arch).

The digital arteries freely communicate, on the palmar and dorsal aspect of the fingers, by transverse branches, which supply the joints and the sheaths of the tendons. Near the ungual phalanx, a considerable branch passes to the back of

the finger, and forms a network of vessels round the root of the nail.

[The varieties which are met with in the formation of the palmar arches are many. We may have no superficial arch, or it may be diminished and fewer digital branches given off than usual. The branch that supplies the sides of the fore and middle fingers is the one most frequently wanting. The ulnar artery at times, after giving branches to the muscles of the little finger, terminates in the deep arch, and all the four digital arteries are then absent. The deficiencies of the superficial arch are usually compensated for by the deep arch. The palmar interosseous branches, dividing at the clefts of the fingers, and thus forming the digital branches which are wanting from the superficial arch. The superficial arch may be made up by the superficialis volæ, by a large median, or even by a metacarpal branch.]

Ulnar Nerve in the Palm.—The *ulnar nerve* passes over the annular ligament into the palm, on the inner side of the ulnar artery. It lies in a groove between the pisiform and unciform bones, so that it is perfectly secure from pressure in this apparently exposed situation. Immediately below the pisiform bone, the nerve divides into a superficial and a deep palmar branch. The *deep* branch supplies the muscles of the little finger, and accompanies the *ulnaris profunda* artery into the palm, to supply *all* the interosseous muscles. The *superficial* branch divides into two digital nerves, for the supply of both sides of the little finger and the ulnar side of the ring finger.*

Anterior Annular Ligament of the Carpus.—This exceedingly strong and thick ligament confines the flexor tendons of the fingers and thumb, and fastens together the bones

* The more internal of these two nerves sends filaments to the *palmaris brevis*, to the integument on the inner side of the palm, and is joined by a filament from the dorsal cutaneous branch of the ulnar nerve; it then runs along the ulnar side of the palm, internal to its corresponding artery, and is continued along the inner side of the little finger to the extremity. The other nerve descends internal to its corresponding artery, and passes obliquely over the flexor tendons of the little finger towards the cleft between the little and ring fingers, where it subdivides into two branches, which run along the opposite sides of these two fingers to their extremities. It also sends a filament to communicate with the median nerve, behind the superficial palmar arch.

of the carpus. It is attached externally to the scaphoid and trapezium; internally to the pisiform and unciform. Its upper border is continuous with the aponeurosis in front of the wrist; its lower border terminates in the palmar fascia; its anterior surface receives the expanded tendon of the palmaris longus, and gives origin to most of the muscles constituting the ball of the thumb and little finger.

Cut vertically through the ligament, and observe, that, with the carpal bones, it forms a complete elliptical canal, with the broad diameter transversely. This canal is lined by a synovial membrane which is reflected loosely over the tendons. It transmits the superficial and deep flexor tendons of the fingers, the long flexor tendon of the thumb, and the median nerve. The tendon of the flexor carpi radialis does not run with the other tendons, but is contained in a distinct sheath, lined by a separate synovial membrane, formed partly by the annular ligament and partly by the groove in the trapezium.

Median Nerve in the Palm.—In its passage under the annular ligament, the median nerve is enveloped in a fold of synovial membrane. It lies between the tendons of the flexor sublimis and those of the flexor profundus, and is rather nearer to the radial than the ulnar side of the wrist. As soon as it appears in the palm, the nerve lies superficial to all the tendons, gives a branch to the muscles of the ball of the thumb, and then divides into five branches, two for the thumb, the rest for the fore and middle fingers and the radial side of the ring finger.

The two nerves of the thumb proceed, one on each side of the long flexor tendon, to the last phalanx.

The first digital nerve runs along the radial side of the index finger. The second descends towards the cleft between the index and middle fingers, and subdivides into two branches, which supply their opposite sides. The third is joined by a filament from one of the ulnar digital nerves, and then subdivides above the cleft between the middle and ring fingers, to supply their opposite sides.

About an inch and a quarter above the clefts between the fingers, each digital nerve subdivides into two branches, between which the digital artery passes and bifurcates lower

down; therefore a vertical incision down the cleft would divide the artery before the nerve.

In their course along the fingers and thumb, the nerves lie superficial to the arteries, and nearer to the flexor tendons. About the middle of the first phalanx each nerve sends a branch, which runs along the back of the finger nearly to the extremity, communicating with the dorsal branches derived from the radial and ulnar nerves.* Near the ungual phalanx another branch is distributed to the skin around and beneath the root of the nail. Each nerve terminates near the end of the finger in a brush of filaments, with their points directed to the papillæ of the skin.

Flexor Tendons and their Sheaths.—Immediately below the annular ligament the tendons separate from each other; near the metacarpal joints they pass in pairs, through strong fibrous rings (p. 237) formed by the divisions of the palmar fascia. Below the metacarpal joint the two tendons for each finger enter the sheath (theca), which confines them in their course along the phalanges. It is formed by a strong fibrous membrane, which is attached to the ridges on the phalanges, and converts the groove in front of these bones into a complete canal, exactly large enough to contain the tendons. The density of the sheath necessarily varies in particular situations, otherwise there would be an obstacle to the easy flexion of the fingers. To ascertain this, cut open one of the sheaths along its entire length; you will then see that it is much stronger between the joints than over the joints themselves: through these sheaths, inflammation commencing in the integuments of the finger may readily extend to the synovial membrane of the tendon.

In cases of whitlow, when matter forms in the theca, the

* We find upon the cutaneous nerves of the hands and feet little bodies termed, after their discoverer, corpuscles of "Pacini." Some of them will be found, by carefully examining the trunk of a nerve, or one of its smaller branches, in the subcutaneous tissue at the root of a finger. Each corpuscle is attached by a slender fibro-cellular pedicle to the nerve upon which it is situated; through the pedicle a single primitive nerve-fibril passes into the corpuscle. The corpuscle itself is composed of a series of concentric capsules, separated by intervals containing fluid; and the nerve-fibril terminates in a central cavity which exists in the axis of the corpuscle. Their function is unknown.

incision should be made deep enough to lay open this fibro-osseous canal, without which the incision will be of no use. It is obvious that the incision should be made down the *centre* of the finger, to avoid the digital nerves and arteries. If this opening be not timely made, the flexor tendons are likely to slough, and the finger becomes stiff.*

But what protects the joints of the fingers where the flexor tendons play over them? Look into an open sheath, and you will see that in front of the joints the tendons glide over a smooth fibro-cartilaginous structure, called the "*glenoid*" ligament.

To facilitate the play of the tendons, the interior of the sheath, as well as the tendons, are lined by a synovial membrane, of the extent of which it is important to have a correct knowledge. With a probe you may easily ascertain that the synovial membrane is reflected from the sheath upon the tendons, a little above the metacarpal joints of the fingers; that is, nearly in a line with the transverse fold in the skin in the lower third of the palm. Towards the distal end of the finger, the synovial sheath stops short of the last joint, so that it is not injured in amputation of the ungual phalanx.

In dissecting the sheaths of the tendons, I have frequently observed little hernia-like protrusions of the synovial membrane through the sheath, varying in size from a pin's head to a pea, very hard and incompressible: when punctured a reddish substance like apple jelly escapes, and the tumor subsides. I have seen these little tumors in the living hand; they interfere with the proper use of the finger: so far as my experience goes, the best mode of treating them is to evacuate their contents by a subcutaneous puncture, and keep the finger on a splint for a few days afterwards.

* On closer inspection it will be observed that the sheath is composed of bands of fibres, which take different directions, and have received distinct names. The strongest are called the "*ligamenta vaginalia*." They constitute the sheath over the body of the phalanx, and extend transversely from one side of the bone to the other. The "*ligamenta cruciata*" are two slips, which cross obliquely over the tendons. The "*ligamenta annularia*" are situated immediately in front of the joints, and may be considered as thin continuations of the *ligamenta vaginalia*. They consist of delicate fibres, which are attached on either side of the joints to the *glenoid* ligaments, and pass transversely over the tendons.

And now notice how beautifully the tendons are adapted to each other in their course along the finger. The superficial flexor, near the root of the finger, becomes slightly grooved to receive the deep flexor; about the middle of the first phalanx it splits into two portions, through which the deep flexor passes. The two portions reunite below the deep tendon so as completely to embrace it, and then divide a second time into two slips, which interlace with each other and are inserted into the sides of the second phalanx. The *deep* flexor, having passed through the opening of the superficial one, is inserted into the base of the last phalanx.*

In what way are the tendons supplied with blood? Raise and separate the tendons, and you will see that slender folds of synovial membrane (*vincula tendinum*) run up from the phalanges to the tendons; when minutely injected, these folds are full of blood-vessels.

The *tendon of the flexor longus pollicis* lies on the radial side of the other tendons beneath the annular ligament. It passes between the two portions of the flexor brevis pollicis and the two sesamoid bones of the thumb, enters its proper sheath, and is *inserted* into the base of the last phalanx. Its synovial sheath is prolonged from the large bursa of the flexor tendons beneath the annular ligament, and accompanies it down to the last joint of the thumb; consequently the sheath is injured in amputation of the last phalanx.

Bursal Sac of the Carpus.—A large and loose synovial sac, called the bursa of the carpus, facilitates the play of the tendons beneath the anterior annular ligament. It lines the under surface of the ligament and the groove of the carpus, and is reflected in loose folds over the tendons. It is prolonged up the tendons for an inch and a half, or two inches, and forms a “cul-de-sac” above the ligament. Below the ligament the bursa extends into the palm, and sends off prolonga-

* In the museum of the College of Surgeons, a preparation is put up which shows a beautiful piece of animal mechanics about the flexor tendons, namely, that in its passage along the phalanges the deep flexor forms at the first phalanx a kind of little patella for the superficial one; but at the second phalanx the superficial flexor lies deeper than the other, and forms a little patella for it. This is a very pretty mechanical ingenuity to increase the leverage in each case.

tions for each of the flexor tendons, which accompany them down to the middle of the hand. You will easily understand that, when the bursa is inflamed and distended by fluid, there will be a bulging above the annular ligament, and another in the palm, with perceptible fluctuation between them; the unyielding ligament causing a constriction in the centre.*

Lumbricales.—These four slender muscles, one for each finger, are attached to the deep flexor tendons in the palm. All of them *arise* from the radial side of the deep tendon of their corresponding finger: the third and fourth often arise from the adjacent sides of two tendons. Each terminates in a broad thin tendon which passes over the radial side of the first joint of the finger, and is *inserted* into the extensor tendon on the back of the finger. Their *action* is to bend the first joint of the fingers. Being inserted near the centre of motion, they can move the fingers with great rapidity. As they produce the quick motions of the musician's fingers, they are called by anatomists "*fidicinales*."

Muscles of the Ball of the Thumb.—The great strength of the muscles of the ball of the thumb (*unde nomen pollicis*), is one of the distinguishing characters of the human hand. This strength is necessary in order to oppose that of all the fingers. In addition to its strength, the thumb enjoys perfect mobility. It has no less than eight muscles to work it:—namely, an abductor, an opponens, two flexors, three extensors, and an adductor. Take first the muscles of the ball.

Abductor Pollicis.—This is the most superficial of the muscles of the ball. It *arises* from the os scaphoides and annular ligament; and is *inserted* by a flat tendon into the base of the first phalanx of the thumb. Its *action* is to draw the thumb away from the fingers. Reflect it from its insertion to expose the following:—

Flexor Ossis Metacarpi Pollicis or Opponens.—This muscle *arises* from the os trapezium and the annular liga-

* I have met with only one instance in which this bursa communicated with the wrist joint. It communicates always with the synovial sheath of the long flexor of the thumb, in most cases with that of the flexors of the little finger, and but rarely with that of the index, middle, and ring fingers. On this account inflammation of the theca of the thumb or little finger is more liable to be attended with serious consequences than either of the others.

ment, and is inserted into the whole length of the metacarpal bone of the thumb. The *action* of this powerful muscle is to oppose the thumb to all the fingers. Reflect it from its insertion to expose the following :—

Flexor Brevis Pollicis.—This muscle has two origins ; one from the annular ligament and os trapezium, the other from the os trapezoides, os magnum, and the upper end of the third metacarpal bone. It is inserted by two strong tendons into the base of the first phalanx of the thumb. A sesamoid bone is found in each of the tendons. The lower portions of the muscle are separated by the long flexor tendon of the thumb and the arteria magna pollicis. Its *action* is to bend the first phalanx of the thumb.

Adductor Pollicis.—This muscle *arises* from the shaft of the metacarpal bone of the middle finger ; its fibres converge and are *inserted*, along with the inner portion of the flexor brevis pollicis, into the base of the first phalanx of the thumb. Its *action* is to draw the thumb towards the palm, as when we bring the tips of the thumb and little finger into contact.

The adductor pollicis should be considered as one of the palmar interosseous muscles. It is supplied by the deep branch of the ulnar nerve, which we know supplies *all* the other interossei. The other muscles of the ball of the thumb are supplied by the median nerve.

Muscles of the Ball of the Little Finger.—The muscles of the little finger correspond in some measure with those of the thumb. Thus we have an abductor, a flexor brevis, and an opponens minimi digiti.

Abductor Minimi Digiti.—This is the most superficial of the muscles of the little finger ; it *arises* from the pisiform bone, and is *inserted* by a flat tendon into the base of the first phalanx of the little finger. Its *action* is to draw this finger from the rest.

Flexor Brevis Minimi Digiti.—This slender muscle may fairly be considered as a portion of the preceding. It *arises* from the unciform bone and annular ligament, and is *inserted* with the tendon of the abductor into the base of the first phalanx of the little finger. Its *action* is similar to that of the abductor. Between the origins of the abductor and flexor

brevis minimi digiti, the deep branch of the ulnar artery and nerve sinks down to form the deep palmar arch.

[The flexor *brevis minimi* is sometimes absent.]

Opponens Digiti Minimi.—The last two muscles must be reflected from their insertion, to expose the *opponens digiti minimi*. It arises from the unciform bone and the annular ligament, and is inserted along the shaft of the metacarpal bone of the little finger. Its *action* is to draw this bone, the most movable of all the metacarpal bones of the fingers, towards the thumb. Thus it greatly strengthens the grasp of the palm.

Now you must cut through all the flexor tendons, and remove the deep fascia of the palm, to see the deep arch of arteries and its branches.

Branches of the Radial Artery in the Palm.—The radial artery enters the palm at the angle between the first and second metacarpal bones (between the inner head of the flexor *brevis* and the adductor pollicis), and gives off three branches—the *arteria magna pollicis*, the *radialis indicis*, and the *palmaris profunda*, which unites with the ulnar to form the deep arch.

The *arteria magna pollicis* runs in front of the abductor indicis (first dorsal interosseous), close along the metacarpal bone of the thumb: in the interval between the lower portions of the flexor *brevis pollicis*, the artery divides into two branches, which proceed, one on either side of the thumb, and inosculate at the apex of the last phalanx. Their distribution and mode of termination are similar to those of the other digital arteries.

The *arteria radialis indicis* runs along the radial side of the index finger to the extremity, where it forms an arch with the other digital artery, a branch of the ulnar. Near the lower margin of the adductor pollicis, the *radialis indicis* generally receives a branch from the superficial palmar arch.

The *palmaris profunda* is considered as the continuation of the radial artery. It enters the palm between the inner head of the flexor *brevis* and the adductor pollicis, runs across the upper ends of the metacarpal bones, and inosculates with the deep branch of the ulnar artery, thus completing the deep palmar arch. From the curve of the arch small branches as-

pend to supply the bones and joints of the carpus, inosculating with the other carpal arteries. From the convexity of the arch four small branches, called *palmar interosseous* (p. 238, 1, 2, 3, 4), descend to supply the interosseous muscles, and near the clefts of the fingers communicate with the digital arteries. These palmar interosseous branches are sometimes of considerable size, and take the place of one or more of the digital arteries, ordinarily derived from the superficial palmar arch. Other branches called *perforating arteries*, pass between the upper ends of the metacarpal bones to the back of the hand, and communicate with the carpal branches of the radial and ulnar arteries.

Deep Branch of the Ulnar Nerve.—This nerve sinks into the palm with the *ulnaris profunda* artery, between the abductor and flexor brevis minimi digiti. It then runs with the deep palmar arch towards the radial side of the palm, and terminates in the adductor pollicis.

Between the pisiform and unciform bones, the nerve gives a branch to each of the muscles of the little finger. Subsequently, it sends branches to each *interosseous muscle*, and generally to the two internal lumbricales.

The dissection of the remaining muscles of the palm, called, from their position, *interossei*, must be for the present postponed.

DISSECTION OF THE MUSCLES OF THE BACK CONNECTED WITH THE ARM.

MAKE an incision down the spine from the occiput to the sacrum; another from the last dorsal vertebra to the spine of the scapula, and thence along the spine to the acromion. Reflect the skin from the dense subcutaneous tissue.

Cutaneous Nerves of the Back.—These are derived from the posterior branches of the spinal nerves. After supplying the muscles in the vertebral groove, these branches become subcutaneous in nearly a regular series, but not all at an equal distance from the spine. Thus, in the cervical and upper dorsal region, the cutaneous nerves perforate the trapezius

close to the spine: in the lower dorsal and the lumbar region, they perforate the latissimus dorsi in a line nearly corresponding to the angles of the ribs. Each cutaneous nerve then divides into internal and external branches, which supply the integument of the respective regions of the back to which they belong. most of them are accompanied by small arteries. As might be expected, the external branches are the larger, especially in the loins, where some of them descend over the crest of the ilium, and terminate in the skin of the buttock.

Of these cutaneous nerves we need notice only the following:—

The *posterior branch of the second cervical nerve* is called the *great occipital*. It perforates the complexus, and ramifies upon the scalp, with the branches of the occipital artery.

The *cutaneous branch of the third cervical nerve* also supplies the back of the scalp.

The *cutaneous branch of the second dorsal nerve* is the largest of all the dorsal cutaneous nerves. It may be traced outwards towards the spine of the scapula.

The *posterior branch of the second lumbar nerve* perforates the fascia lumborum near the posterior superior spine of the ilium, and runs over the crest of the ilium to supply the skin of the buttock.

Clean the trapezius and latissimus dorsi by dissecting in the course of their fibres.

Trapezius.—Taken alone this muscle is triangular, but with its fellow it presents a trapezoid form. It *arises* from the inner fourth, more or less, of the superior curved line of the occiput; from the ligamentum nuchæ* and from the spines of the seventh cervical, and all the dorsal vertebræ. The fibres converge towards the shoulder. The upper are *inserted* fleshy into the external third of the clavicle; the middle, into the inner border of the acromion and spine of the scapula; the

* The "ligamentum nuchæ" is, in man, only a rudiment of the great elastic ligament which supports the weight of the head in quadrupeds. It extends from the spine of the occiput to the spines of all the cervical vertebræ except the atlas; otherwise it would impede the free rotation of the head. In the giraffe this ligament is six feet long and as thick as a man's forearm. I am told by Professor Quekett, that when divided it shrinks at least two feet.

lower terminate in a thin tendon, which plays over the triangular surface at the back of the scapula, and is inserted into the beginning of the spine. The insertion of the trapezius exactly corresponds to the origin of the deltoid, and the two muscles are connected by a thin aponeurosis over the spine and acromion. If both the trapezius muscles be exposed, you will see that, between the sixth cervical and the third dorsal vertebra, their origin presents an aponeurotic space of an elliptical form.

The fixed point of the muscle being at the spine, all its fibres tend to raise the shoulder. The deltoid cannot raise the humerus beyond an angle of sixty degrees: beyond this the elevation of the arm is principally effected by the action of the trapezius upon the scapula. It is in strong action when a weight is borne upon the shoulders: again it acts powerfully in drawing the scapula backwards, as in preparing to strike a violent blow. It is supplied by the *nervus accessorius* and by the *superficialis colli* artery.

Latissimus Dorsi.—This broad flat muscle occupies the lumbar and lower dorsal region, and thence extends to the arm, where it forms part of the posterior boundary of the axilla. It *arises* from the posterior third of the crest of the ilium, from the “*fascia lumborum*,” from the spinous processes of the sixth lower dorsal vertebræ beneath the trapezius, and lastly, from the three or four lower ribs by digitations, which correspond with those of the external oblique muscle of the abdomen. All the fibres converge towards the axilla, where they form a thick muscle, which is folded beneath the *teres major*, and *inserted* by a broad flat tendon into the bottom of the bicipital groove of the humerus. The tendon is about two inches broad, and lies in front of that of the *teres major*, from which it is separated by a large *bursa*.*

The *latissimus dorsi* draws the humerus inwards and backwards: it also co-operates with the *pectoralis major* in pulling any object towards the body: if the humerus be the fixed point, it raises the body, as in climbing. What is the object of the muscle arising so high up the back? It is that the

* The *latissimus dorsi* sometimes receives a distinct accessory slip from the inferior angle of the scapula.

transverse fibres of the muscle serve the important purpose of strapping down the inferior angle of the scapula. It sometimes happens that the scapula slips above the muscle: this displacement is readily recognized by the unnatural projection of the lower angle of the bone, and the impaired movements of the arm.*

Between the base of the scapula, the trapezius, and the upper border of the latissimus dorsi, a triangular space is observed when the arm is raised, in which the lower fibres of the rhomboideus major and part of the sixth intercostal space are exposed. Immediately above the crest of the ilium, between the free margins of the latissimus dorsi and external oblique, there is, also, an interval in which a little of the internal oblique can be seen.

Fascia Lumborum.—This is a dense shining aponeurosis in the lumbar region, thin and pointed above, broader and stronger below. It consists of tendinous fibres, which are attached to the spines of the six or seven lower dorsal, all the lumbar and sacral vertebræ, and to the posterior part of the crest of the ilium. It serves as the common origin of the latissimus dorsi, the serratus posticus inferior, and the internal oblique and transverse muscles of the abdomen, and it forms the posterior part of the sheath of the erector spinæ. When suppuration takes place in the loins, constituting “lumbar abscess,” in connection with spinal disease, the matter is seated beneath this fascia, and is therefore a long time in coming to the surface.

Reflect the trapezius from its insertion. On its under surface you will see the ramifications of its nutrient artery, the *superficialis colli* (a branch of the posterior scapular). A large nerve, the *spinal accessory*, enters its under surface near the clavicle, and then divides into filaments, some of which ascend, others descend in its substance.

Spinal Accessory Nerve.—This nerve is one of the three

* I have seen an instance of this displacement. There was great projection of the inferior angle of the scapula, especially when the man attempted to raise the arm. He could not raise the arm beyond a right angle, unless firm pressure was made on the lower angle of the scapula, so as to supply the place of the muscular strap. Whether the scapula can be replaced or not, apply a firm bandage round the chest.

divisions of the eighth pair of cerebral nerves. It arises from the lateral part of the cervical portion of the spinal cord by several roots, some of which are as low as the fourth cervical vertebra. Formed by the union of these roots, the nerve enters the skull through the foramen magnum, and leaves it again through the foramen jugulare. It then runs behind the internal jugular vein, traverses obliquely the upper third of the sterno-mastoid, and crosses the posterior triangle of the neck of the trapezius, which it supplies (p. 9).

Beneath the trapezius we have to examine four muscles connected with the scapula, namely, the levator anguli scapulæ, the omo-hyoideus, the rhomboideus major and minor. To clean these muscles, the scapula must be adjusted so as to stretch their fibres.

Levator Anguli Scapulæ.—This muscle is situated at the side of the neck. It *arises* by four tendons from the posterior tubercles of the transverse processes of the four upper cervical vertebrae. The muscular slips to which the tendons give rise, form a single muscle, which descends down the side of the neck, and is *inserted* into the posterior border of the scapula between its spine and superior angle. Its *action* is to raise the posterior angle of the scapula; as, for instance, in shrugging the shoulders. Its nerve comes from the fifth cervical.

Rhomboideus Major and Minor.—These flat muscles extend from the spinous processes of the vertebrae to the base of the scapula. They often appear like a single muscle. The rhomboideus minor, the higher of the two, *arises* by a thin aponeurosis from the spinous processes of the last cervical and the first dorsal vertebrae, and is *inserted* into the base of the scapula opposite its spine. The rhomboideus major *arises* by tendinous fibres from the spinous processes of the four or five upper dorsal vertebrae, and is *inserted* by fleshy fibres into the base of the scapula between its spine and inferior angle. The *action* of these muscles is to draw the scapula upwards and backwards. They are the antagonists of the serratus magnus.

The *nerve of the rhomboid muscles* (posterior scapular) is a branch of the fifth cervical nerve. It passes outwards beneath the lower part of the levator anguli scapulæ, to which

it sends a branch, and is lost in the under surface of the rhomboid muscles.

Omo-hyoideus.—This muscle extends from the scapula to the os hyoides, and consists of two long narrow muscular portions, connected by an intermediate tendon beneath the sterno-mastoid. The posterior portion only can be seen in the present dissection. It *arises* from the scapula, close behind the notch, and from the ligament above the notch. From thence the slender muscle passes forwards across the lower part of the neck, beneath the sterno-mastoid, where it changes its direction and ascends nearly vertically, to be attached to the os hyoides at the junction of the body with the greater cornu (p. 11). Thus the two portions of the muscle form beneath the sterno-mastoid an obtuse angle of which the apex is tendinous, and of which the angular direction is maintained by a layer of fascia, proceeding from the tendon to the first rib and the clavicle. Its *action* is to depress the os hyoides.

Supra-scapular Artery.—This artery (*transversalis humeri*), a branch of the thyroid axis, runs beneath and parallel with the clavicle to the supra-scapular notch, through which it passes to supply the supra-spinatus and infra-spinatus, and then inosculates freely with the dorsalis scapulæ, a branch of the subscapular. It sends off the *supra-acromial* branch, which ramifies upon the acromion, anastomosing with the other acromial arteries derived from branches of the axillary. The supra-scapular vein terminates either in the subclavian or in the external jugular vein.

[The supra-scapular often comes off from the third portion of the subclavian artery.]

The *supra-scapular nerve*, a branch of the fifth and sixth cervical nerve, runs with the corresponding artery, and is distributed to the supra-spinatus and infra-spinatus.

Posterior Scapular Artery.—This artery is a continuation of the "*transversalis colli*," which is usually described as arising from the thyroid axis, but comes just as often from the subclavian in the third part of its course. It runs across the lower part of the neck, above, or between the nerves of the brachial plexus, towards the posterior superior angle of the scapula. Here it takes the name of posterior scapular, and pursues its course beneath the levator anguli scapulæ and the

rhomboid muscles. If you divide the rhomboid muscles near their insertion, you can trace the artery to the inferior angle of the scapula, where it terminates in the rhomboid and serratus magnus.

Numerous muscular branches arise from this artery. One, the *superficialis colli*, is given off near the upper angle of the scapula for the supply of the trapezius.

The *posterior scapular vein* terminates in the subclavian or in the external jugular.

Divide and reflect the latissimus dorsi below the inferior angle of the scapula, and draw the scapula forcibly outwards, in order to have a more perfect view of the extent of the serratus magnus than you had in the axilla. The great abundance of cellular tissue in this situation is necessary for the free play of the scapula on the side of the chest.

Serratus Magnus.—This broad flat muscle intervenes between the scapula and the ribs. It *arises* by fleshy digitations from the eight or nine upper ribs. The four lower digitations correspond with those of the external oblique muscle of the abdomen. All the fibres pass backwards and converge to be *inserted* along the posterior border of the scapula, chiefly near the upper and lower angles.

This is one of the most important of the muscles which regulate the movements of the scapula; it draws the scapula forwards—and thus gives additional reach to the arm; it counteracts all forces which tend to push the scapula backwards; for instance, when a man falls forwards upon his hands, the serratus magnus sustains the shock, and prevents the scapula from being driven back to the spine. Supposing the fixed point to be at the scapula, some anatomists ascribe to it the power of raising the ribs: hence Sir Charles Bell called it the external respiratory muscle, as opposed to the internal respiratory muscle, the diaphragm.

[The serratus magnus may have as many as eleven distinct slips of origin.]

The nerve which supplies the muscle is a branch of the fifth and sixth cervical nerve: it descends along its outer surface (p. 207).

Now divide the serratus magnus near the scapula, and remove the arm by sawing through the middle of the clavicle, and cutting through the axillary vessels and nerves. These

should subsequently be tied to the coracoid process. After the removal of the arm, the precise insertion of the preceding muscles into the scapula should be carefully examined.

DISSECTION OF THE MUSCLES OF THE SHOULDER.

Cutaneous Nerves of the Shoulder. — The *cutaneous nerves of the shoulder* are derived partly from the branches of the cervical plexus which descend over the acromion (page 3), and partly from the circumflex nerve, of which one or two branches turn round the posterior border of the deltoid; others perforate the muscle, each accompanied by a small artery.

The thin layer of fascia upon the surface of the deltoid extends from the strong aponeurosis covering the muscles on the back of the scapula, and is continuous with the fascia of the arm. It dips down between the fibres of the muscle, dividing it into large bundles.

Deltoid. — The great muscle which covers the shoulder-joint is named deltoid, from its resemblance to the Greek Δ reversed. It *arises* from the external third of the clavicle, from the acromion, and from the spine of the scapula down to the triangular surface at its root. This origin, which corresponds to the insertion of the trapezius, is tendinous and fleshy everywhere, except at the commencement of the spine of the scapula, where it is simply tendinous, and connected with the infra-spinous aponeurosis. The muscular fibres descend, the anterior backwards, the posterior forwards, the middle perpendicularly; all converge to a tendon which is *inserted* into a rough surface on the outer side of the humerus, a little above the middle of the shaft. The insertion of the tendon extends one inch and a half along the humerus, and terminates in a V-shaped form, the origin of the brachialis anticus being on either side.

The muscular bundles composing the deltoid have a very peculiar arrangement: a peculiarity arising from its broad origin and its narrow insertion. It consists in the interposition of tendons between the bundles for the attachment of the

muscular fibres. The annexed woodcut shows this beautiful arrangement better than any description. The action of the muscle is not only concentrated upon one point, but its power is also greatly increased, by this arrangement.

Action of the Deltoid.—It raises the arm; but it cannot do so to a greater angle than sixty degrees. The elevation of the arm beyond this angle is effected through the elevation of the shoulder by means of the trapezius and the serratus magnus. Its anterior fibres draw the arm forwards; its posterior, backwards.

This powerful muscle is well supplied with blood, by the anterior and posterior circumflex, the thoracica humeraria, the thoracica acromialis, all from the axillary artery; also by the deltoid branch of the brachial. Its nerve is the circumflex.


The deltoid gives the rotundity to the shoulder. When the head of the humerus is dislocated into the axilla, the fibres of the muscle run vertically to their insertion; hence the flattening of the deltoid, and the greater prominence of the acromion.

It is just below the deltoid that we generally meet with ununited fracture of the humerus; owing to the muscle raising the upper fragment.

Reflect the deltoid from its origin. Observe the ramifications of the circumflex artery and nerve, on its under surface; also the large bursa between it and the head of the humerus. See how the muscle protects the shoulder-joint; how it covers the coraco-acromial ligament, the head, neck, and upper part of the humerus, as well as the tendons inserted into the greater tuberosity.

Bursa under the Deltoid.—A bursa of large size is situated between the deltoid and the head of the humerus. It extends for some distance beneath the acromion and the co-

Fig. 47.



ANALYSIS OF THE DELTOID.

raco-acromial ligament, and covers the great tuberosity of the humerus. I have seen it communicating by a wide opening with the shoulder-joint, but this is a rare exception. Its use is to facilitate the movements of the head of the bone under the acromial arch.

Posterior Circumflex Artery.—This artery is given off from the axillary: it runs behind the surgical neck of the humerus, through a quadrilateral opening, bounded above by the subscapularis, below by the teres major, externally by the neck of the humerus, and internally by the long head of the triceps (p. 211). Its branches terminate on the under surface of the deltoid, anastomosing with the anterior circumflex and acromial thoracic arteries.

From the posterior circumflex, a branch descends in the substance of the long head of the triceps, to inosculate with the superior profunda: this is one of the channels through which the circulation would be carried on, if the axillary were tied in the last part of its course.

Circumflex Nerve.—This nerve, a branch of the axillary plexus, runs with the posterior circumflex artery. It sends a branch to the teres minor, one or two to the integuments of the shoulder, and terminates in the substance of the deltoid. The proximity of this nerve to the head of the humerus explains the occasional occurrence of paralysis of the deltoid, after a dislocation or fracture of the humerus. The nerve is apt to be injured, if not actually lacerated by the pressure of the bone. In the summer of 1840, a man was admitted into the hospital with a severe injury to the shoulder, and died of delirium tremens. On examination the humerus was found broken high up, the capsule of the joint opened and the circumflex nerve torn completely across.*

The muscles on the dorsum of the scapula are covered by a strong aponeurosis, which is firmly attached to the spine and borders of the bone. At the posterior edge of the deltoid it divides into two layers, one of which passes over, the other under, the muscle. Remove the aponeurosis from the muscles, so far as it can be done without injury to the muscular fibres which arise from its under surface.

* See preparation in Museum of St. Bartholomew's Hospital, series 3, number 42.

Infra-spinatus.—This muscle *arises* from the greater part of the infra-scapular fossa. The fibres converge to a tendon, which is at first contained in the substance of the muscle, and then proceeds over the capsule of the shoulder-joint, to be *inserted* into the greater tuberosity of the humerus.

Teres Minor.—This long narrow muscle is situated below the infra-spinatus, along the inferior border of the scapula. It *arises* from the dorsum of the scapula, close to the inferior border. The fibres ascend parallel with those of the infra-spinatus, and terminate in a tendon, which passes over the capsule of the shoulder-joint, and is *inserted* into the greater tuberosity of the humerus. Some of the fibres of the muscle are attached directly into the bone. It is supplied by a branch of the circumflex nerve.

The *action* of the infra-spinatus and teres minor is to rotate the humerus outwards.

Teres Major.—This muscle is closely connected with the latissimus dorsi, and extends from the inferior angle of the scapula to the humerus, contributing to form the posterior boundary of the axilla. It *arises* from the flat surface at the inferior angle of the back of the scapula, and terminates upon a flat tendon, two inches in breadth, which is *inserted* into the inner edge of the bicipital groove of the humerus, behind the tendon of the latissimus dorsi (p. 211). Its *action* is to draw the humerus backwards. By dissecting between the two tendons, you will find that they are separated by a *bursa*.

Supra-spinatus.—This muscle *arises* from the posterior two-thirds of the supra-spinous fossa, and from its aponeurotic covering. It passes underneath the acromion, over the shoulder-joint, and is *inserted* by a strong tendon into the greater tuberosity of the humerus. To obtain a good view of its insertion, the acromion should be sawn off near the neck of the scapula. Its *action* is to assist the deltoid in raising the arm.

Subscapularis.—This muscle occupies the subscapular fossa. It *arises* from the posterior three-fourths of the fossa, from three or four tendinous septa attached to the oblique ridges on its surface. The fibres converge towards the neck of the scapula, where they terminate upon three or four tendons, which are concealed among the muscular fibres, and are

inserted into the lesser tuberosity of the humerus. Its broad insertion is closely connected with the capsule of the shoulder-joint, which it completely protects upon its inner side. Its *action* is to rotate the humerus inwards. The nerve which supplies it is derived from the great cord, which gives off the circumflex and musculo-spiral nerves.

The coracoid process, with the coraco-brachialis and short head of the biceps, form an arch, under which the tendon of the subscapularis plays. There are several *bursæ* about the tendon. There is one of considerable size on the upper surface of the tendon, to facilitate its motion beneath the coracoid process and the coraco-brachialis: this sometimes communicates with the large bursa under the deltoid. Another is situated between the tendon and the capsule of the joint, and almost invariably communicates with it.

Now reflect the muscles from the surface of the scapula, to trace the arteries which ramify upon the bone.

Continuation of Supra-scapular Artery.—This artery, a branch of the thyroid axis, runs under and parallel with the clavicle, and passes above, or, less frequently, through the notch of the scapula, into the supra-spinal fossa: it sends a branch to the supra-spinatus, another to the shoulder-joint, and then descends over the neck of the scapula into the fossa below the spine, where it inosculates directly with the dorsalis scapulæ. Its branches ramify upon the bone, and supply the infra-spinatus and teres minor.

The supra-scapular nerve passes most frequently through the notch of the scapula, accompanies the corresponding artery, and supplies the supra and infra-spinatus.

Dorsalis Scapulæ Artery.—The *dorsal branch of the subscapular artery* (dorsalis scapulæ), passes backwards through a triangular space bounded superiorly by the subscapularis, inferiorly by the teres major, and in front by the long head of the triceps (p. 211). Having reached the dorsum of the scapula, it ascends, close to the bone, and anastomoses with the supra-scapular artery.

The frequent communications about the scapula, between the branches of the subclavian and axillary arteries, would furnish a free current of blood to the arm if the subclavian were tied above the clavicle. (See fig. 7. p. 47.)

Triceps Extensor Cubiti.—This muscle, which was only partially seen in the dissection of the upper arm (p. 211), should now be more fully examined. The long head arises immediately below the glenoid cavity of the scapula, by a strong tendon which is connected with the capsule of the shoulder-joint. The second or external head *arises* from the posterior part of the humerus, below the insertion of the *teres minor*; the third or internal head *arises* from the posterior part of the humerus below the *teres major*. The precise origin of these heads from the humerus may be ascertained by following the superior profunda artery and musculo-spiral nerve, which intervene between them. The three portions of the muscle terminate upon a broad tendon, which covers the back of the elbow-joint, and is *inserted* into the summit and sides of the olecranon; it is also connected with the fascia on the back of the forearm. The reason of this connection is, that the same muscle which extends the forearm may at the same time tighten the fascia which gives origin to the extensors of the wrist and fingers. This is just what we observed in the case of the biceps, and its semilunar expansion in the fascia of the forearm.

Between the tendon and olecranon is placed a *bursa*, commonly of small size, but sometimes so large as to extend upwards behind the capsule of the joint. You must not confound this bursa with that which is always found between the skin and the olecranon, and is so often injured by a fall on the elbow.

By dividing the triceps transversely a little above the elbow, and turning down the lower portion, it will be observed that some of the muscular fibres terminate upon the capsule of the joint. They have been described by some anatomists as a distinct muscle, under the name of "*subanconeus*;" their use is to draw up the capsule, so that it may not be injured during extension of the arm. The *subanconeus* is in this respect analogous to the *subcruræus* muscle of the thigh. Observe the *bursa* under the tendon, and the arterial arch formed upon the back part of the capsule by the superior profunda and the *anastomotica magna* (fig. 45, p. 220).

Trace the continuation of the superior profunda artery (p. 220), and musculo-spiral nerve round the posterior part of

the humerus. They lie close to the bone, and are protected by an aponeurotic arch thrown over them by the external head of the triceps. Observe that the nerve is here placed between the two principal branches of the artery: therefore in amputation of the arm in this situation, you would have to tie two large arteries, one on each side of the musculo-spiral nerve. Trace the branches given off by the nerve to each of the three portions of the triceps, and follow the long branch of the superior profunda artery through the substance of the triceps to the olecranon.

DISSECTION OF THE BACK OF THE FOREARM.

REMOVE the skin from the back of the forearm, hand, and fingers. Observe the subcutaneous *bursa* over the olecranon. It is commonly of considerable size, and, if distended, would appear nearly as large as a walnut. Another *bursa* is sometimes found a little lower down upon the ulna. A subcutaneous *bursa* is generally placed over the internal condyle, another over the external. A *bursa* is also situated over the styloid process of the ulna; this sometimes communicates with the sheath of the extensor carpi ulnaris. Small *bursæ* are sometimes developed in the cellular tissue over each of the knuckles.

The cutaneous veins from the back of the hand and forearm join the venous plexus at the bend of the elbow (see p. 214).

Cutaneous Nerves of the Back of the Forearm and Hand.—They are the external cutaneous branches of the musculo-spiral nerve, branches of the internal cutaneous nerve, and of the external cutaneous. The greater number of these nerves may be traced down to the back of the wrist.

The skin on the back of the hand is united to the subjacent tendons by an abundance of loose connective tissue, in which find large veins, and branches of the radial and ulnar nerve. The dorsal branch of the ulnar nerve passes beneath the tendon of the flexor carpi ulnaris, over the internal lateral liga-

ment of the wrist; and divides upon the back of the hand into filaments, which supply the back of the little finger, the ring finger, and the ulnar side of the middle finger. The radial nerve passes obliquely beneath the tendon of the supinator longus, and subdivides into branches, which supply both sides of the back of the thumb and fore finger, and the radial side of the middle finger.*

The *fascia on the back of the forearm* is composed of fibres interlacing in various directions, and is stronger than that upon the front of the forearm. It is attached to the two condyles of the humerus and to the olecranon, and is strengthened by an expansion from the tendon of the triceps. Along the forearm it is attached to the ridge on the posterior part of the ulna. Its upper third gives origin to the fibres of the muscles beneath it, and separates them by septa, to which their fibres are also attached.

Posterior Annular Ligament.—This should not be considered as a distinct ligament, but rather as a part of the fascia of the forearm, strengthened by oblique aponeurotic fibres on the back of the wrist to confine the extensor tendons. These fibres are firmly attached to the styloid process of the radius, and thence pass obliquely inwards to the inner side of the wrist, where they are connected with the pisiform and uniform bones. Observe that they pass below the styloid process of the ulna, to which they are in no way attached, otherwise the rotation of the radius would be impeded.

From the deep surface of this so-called ligament, processes are attached to the ridges on the back of the radius, so as to form six distinct sheaths for the passage of the extensor tendons. Counting from the radius towards the ulna, the first sheath contains the tendons of the extensor ossis metacarpi and the extensor primi internodii pollicis; the second contains the tendons of the extensor carpi radialis longior and brevior;

* The relative share which the radial and ulnar nerves take in supplying the fingers is not the same in all cases. Under any arrangement the thumb and each finger have two dorsal nerves, one on either side, of which the terminal branches reach the root of the nail. They supply filaments to the skin on the back of the finger, and have frequent communications with the palmar digital nerves. In some instances one or more of the dorsal nerves do not extend beyond the first phalanx; their place is then supplied by a branch from the palmar nerve.

the third contains the tendon of the extensor secundi internodii pollicis; the fourth contains the tendons of the indicator and the extensor communis digitorum; the fifth contains the tendon of the extensor minimi digiti; and the sixth, the tendon of the extensor carpi ulnaris. All the sheaths are lined by synovial membranes, to facilitate the play of the tendons. In a few instances, one or more of them communicate with the wrist-joint.

The *fascia of the metacarpus* consists of a thin fibrous layer, continued from the posterior annular ligament. It separates the extensor tendons from the subcutaneous veins and nerves, and is attached to the radial side of the second metacarpal bone, and the ulnar side of the fifth.

Superficial Muscles on the Back of the Forearm.—The fascia must be removed from the muscles, so far as this can be done without injuring the muscular fibres which arise from its under surface. Preserve the posterior annular ligament. The following muscles are now exposed, and should be examined in the order in which they are placed, proceeding from the radial to the ulnar side of the forearm:—1. The supinator radii longus (already described, p. 229). 2. The extensor carpi radialis longior. 3. The extensor carpi radialis brevior. 4. The extensor communis digitorum. 5. The extensor minimi digiti. 6. The extensor carpi ulnaris. 7. The anconeus.

A little below the middle of the forearm, the extensors of the wrist and fingers diverge from each other, leaving an interval, in which are seen the three extensors of the thumb—the extensor ossis metacarpi pollicis, the extensor primi internodii pollicis, and the extensor secundi internodii pollicis. The two former cross the radial extensors of the wrist, and pass over the lower third of the radius.

Between the second and third extensors of the thumb, we observe a part of the lower end of the radius, which is not covered either by muscle or tendon. This subcutaneous portion of the bone is immediately above the prominent tubercle in the middle of its lower extremity, and, since it can be easily felt through the skin, it presents a convenient place for examination in doubtful cases of fracture.

Extensor Carpi Radialis Longior.—This muscle is partly

covered by the supinator radii longus. It *arises* from the lower third of the ridge leading to the external condyle of the humerus, descends along the outer side of the forearm, and terminates about the middle, in a flat tendon, which passes beneath the extensor ossis metacarpi and primi internodii pollicis, traverses a groove on the outer and back part of the radius, lined by a synovial membrane, and is *inserted* into the carpal end of the metacarpal bone of the index finger. Previously to its insertion, the tendon is crossed by the extensor secundi internodii pollicis.

[The extensor carpi radialis may also be inserted by a separate slip into the third metacarpal bone, and may be blended with the flexor brevis so as to appear one and the same muscle.]

Extensor Carpi Radialis Brevior.—This muscle *arises* from the external condyle. The muscular fibres terminate near the middle of the forearm, upon the under surface of a flat tendon, which descends, covered by that of the extensor carpi radialis longior, beneath the first two extensors of the thumb. The tendon traverses a groove on the back of the radius, on the same plane with that of the long radial extensor, but lined by a separate synovial membrane, and is *inserted* into the carpal end of the metacarpal bone of the middle finger. A *bursa* is generally found between the tendon and the bone.

Extensor Digitorum Communis.—This muscle *arises* from the common tendon attached to the external condyle, from the septa between it and the contiguous muscles, and from its strong fascial covering. About the middle of the forearm the muscle divides into three or four fleshy slips, terminating in as many flat tendons, which pass beneath the posterior annular ligament, through a groove on the back of the radius lined by synovial membrane. On the back of the hand they become broader and flatter, and diverge from each other towards the metacarpal joints of the fingers, where they become thicker and narrower, and give off on each side a fibrous expansion, which covers the sides of the joint. Over the first phalanx of the finger, each tendon again spreads out, receives the expanded tendons of the lumbricales and interossei muscles, and divides at the second phalanx into three portions, of

which the middle is inserted into the upper end of the second phalanx; the two lateral, reuniting over the lower end of the second phalanx, are inserted into the upper end of the third.*

Observe the oblique aponeurotic slips which connect the tendons on the back of the hand. They are subject to great variety. The tendon of the index finger is commonly free; it is situated on the radial side of the proper indicator tendon, and becomes united with it at the metacarpal joint.

The tendon of the middle usually receives a slip from that of the ring finger. The tendon of the ring finger generally sends a slip to the tendons on either side of it, and in some cases entirely furnishes the tendon of the little finger.

It is not only a general extensor of the fingers, but it can extend some of the phalanges independently of the rest: *e. g.* it can extend the first phalanges while the second and third are flexed; or it can extend the second and third phalanges during the flexion of the first.

Extensor Digiti Minimi or Auricularis.—This long slender muscle *arises* from the common tendon from the external condyle, from the septa between it and the contiguous muscles. Its slender tendon runs beneath the annular ligament immediately behind the joint between the radius and ulna, in a special sheath lined by synovial membrane. At the first joint of the little finger, the tendon is joined by that of the common extensor, and both expand upon the first and second phalanges, terminating in the same manner as the extensor tendons of the other fingers.

Extensor Carpi Ulnaris.—This muscle *arises* from the common tendon from the external condyle, from the septum between it and the extensor minimi digiti, and from the aponeurosis of the forearm. The fibres terminate upon a strong, broad tendon, which traverses a distinct groove on the back of the ulna, close to the styloid process, and is inserted into the carpal end of the metacarpal bone of the little finger. Below the styloid process of the ulna, the tendon passes be-

* The extensor tendons are only inserted into the periosteum, whereas the flexor tendons are inserted into the substance of the bone. This accounts for the facility with which the former will tear off the bones in cases of necrosis, while the latter will adhere so tightly as to require cutting before the phalanx can be removed.

neath the posterior annular ligament, over the back of the wrist, and is here contained in a very strong fibrous canal, which is attached to the back of the cuneiform, pisiform, and unciform bones, and is lined by a continuation from the synovial membrane in the groove of the ulna. The *action* of this muscle is to extend the hand, and incline it towards the ulnar side.

[The tendon of the extensor carpi ulnaris may be united to that of the extensor muscle of the little finger.]

In pronation of the forearm, the lower end of the ulna projects between the tendons of the extensor carpi ulnaris and the extensor minimi digiti. A subcutaneous *bursa* is sometimes found above the bone in this situation.

Anconeus.—This small triangular muscle is situated at the outer and back part of the elbow. It is covered by a strong layer of fascia, derived from the tendon of the triceps, and appears like a continuation of that muscle. It *arises* tendinous from the posterior part of the external condyle of the humerus, and is *inserted* into the triangular surface on the upper fourth of the outer part of the ulna. Part of the under surface of the muscle is in contact with the capsule of the elbow-joint. Its *action* is to assist in extending the forearm.

Deep-seated Muscles on the Back of the Forearm.—Detach from the external condyle the extensor carpi radialis brevis, the extensor digitorum communis, the extensor minimi digiti, and extensor carpi ulnaris. Preserve the vessels and nerves which enter under their surface. The deep-seated muscles now exposed are,—1. The supinator radii brevis; 2. Extensor ossis metacarpi pollicis; 3. Extensor primi internodii pollicis; 4. Extensor secundi internodii pollicis; 5. Extensor proprius indicis or indicator.

Supinator Radii Brevis.—This muscle embraces the upper third of the radius. It *arises* from the external lateral ligament of the elbow-joint, from the annular ligament surrounding the head of the radius, from an oblique ridge on the outer surface of the ulna below the insertion of the anconeus, and by fleshy fibres from a triangular excavation below the lesser sigmoid notch of the ulna. The muscular fibres turn over the neck and upper part of the shaft of the radius, and are *inserted* into the upper third of this bone, as far forwards as the ridge

leading from the tubercle to the insertion of the pronator teres.

The muscle is perforated obliquely by the deep branch of the musculo-spiral nerve, and its upper part is in contact with the capsule of the elbow-joint. It is a powerful supinator of the forearm, some of its fibres acting at nearly a right angle to the axis of the radius.

[We occasionally find two muscles corresponding to the supinator brevis.]

Extensor Ossis Metacarpi Pollicis.—This muscle *arises* from the posterior surface of the ulna below the supinator brevis, from the posterior surface of the radius, and from the interosseous ligament. The muscle crosses the radial extensors of the wrist, about three inches above the carpus, and terminates in a tendon, which passes along a groove, lined by synovial membrane, on the outer part of the lower end of the radius, and is *inserted* into the base of the metacarpal bone of the thumb.

Extensor Primi Internodii Pollicis. This muscle *arises* from the posterior surface of the radius, from the interosseous ligament, and from the ulna. It turns over the radial extensors of the wrist, and terminates upon a tendon which passes beneath the annular ligament, through the groove on the outer part of the radius, and is *inserted* into the radial side of the first phalanx of the thumb.

Extensor Secundi Internodii Pollicis.—*Arises* from the posterior surface of the ulna, and from the interosseous ligament. The tendon receives fleshy fibres as low as the wrist, passes beneath the annular ligament in a distinct groove on the back of the radius, proceeds over the metacarpal bone and the first phalanx of the thumb, and is *inserted* into the base of the last phalanx.

The tendons of the three extensors of the thumb may be easily distinguished in one's own hand. The extensor ossis metacarpi, and primi internodii pollicis, cross obliquely over the radial artery, where it lies on the external lateral ligament of the carpus; the extensor secundi internodii pollicis crosses the artery just before it sinks into the palm, between the first and second metacarpal bones, and is a good guide to the

vessel. The *action* of the three extensors of the thumb is implied by their names.

Extensor Indicis, or Indicator.—This muscle *arises* from the posterior surface of the ulna, and from the interosseous ligament. The tendon passes beneath the posterior annular ligament, in the same groove on the back of the radius with the tendons of the extensor digitorum communis. It then proceeds over the back of the hand to the first phalanx of the index finger, where it is united to the ulnar border of the common extensor tendon. By the *action* of this muscle, the index finger can be extended independently of the others.

[The extensor indicis may be a double muscle and send one of its tendons to the middle finger.]

Posterior Interosseous Artery.—This artery comes from the ulnar by a common trunk with the anterior interosseous (p. 220), and supplies the muscles on the back of the forearm. It passes between the radius and ulna above the interosseous membrane, and appears between the supinator radii brevis and the extensor ossis metacarpi pollicis. After supplying branches to all the muscles in this situation, the artery descends, much diminished in size, between the superficial and deep layer of muscles to the wrist, where it inosculates with the carpal branches of the anterior interosseous and those of the radial and ulnar arteries.

But the largest branch of this artery is the *recurrent*. It ascends beneath the anconeus muscle to the space between the external condyle and the olecranon, where it inosculates with the branch of the superior profunda, which descends in the substance of the triceps.

In the lower part of the back of the forearm, a branch of the anterior interosseous artery is seen passing through the interosseous membrane, to reach the back of the wrist (see p. 235).

Posterior Interosseous Nerve.—The nerve which supplies the muscles on the back of the forearm is the *posterior interosseous*, a branch of the musculo-spiral. It passes obliquely through the supinator radii brevis, and descends between the superficial and the deep layer of muscles on the back of the forearm, sending to each a filament, generally in company with a branch of the posterior interosseous artery. It sends

a branch to the extensor carpi radialis brevior, and supplies the supinator brevis in passing through its substance. The supinator radii longus and the extensor carpi radialis longior are supplied by distinct branches from the musculo-spiral.

The continuation of the posterior interosseous nerve descends beneath the tendons of the extensor digitorum communis to the back of the wrist. In this situation the nerve forms a kind of gangliform enlargement, from which filaments are sent to the carpal and metacarpal joints.

Radial Artery on the Back of the Hand.—Trace the continuation of the radial artery over the external lateral ligament of the carpus, beneath the extensor tendons of the thumb, to the upper part of the interval between the first and second metacarpal bones, where it sinks between the origin of the abductor indicis muscle, and, entering the palm, forms the deep palmar arch. In this part of its course it is crossed by filaments of the radial nerve; observe, also, that the tendon of the extensor secundi internodii pollicis passes over it immediately before it sinks into the palm. It supplies the following small branches to the back of the hand:—

a. *Posterior carpal artery.*—This branch passes across the carpal bones, beneath the extensor tendons. It inosculates with the termination of the anterior interosseous artery, and sometimes with a corresponding branch from the ulnar artery. The carpal artery sends off small branches, called the *dorsal interosseous*, which descend along the third and fourth interosseous spaces, beneath the extensor tendons, inosculating near the carpal ends of the metacarpal bones with the perforating branches from the deep palmar arch.

b. The *dorsal interosseous artery* of the second space is generally larger than the others. It descends over the second dorsal interosseous muscle to the cleft between the index and middle fingers, and terminates in small branches, some of which proceed along the back of the fingers, others inosculate with the palmar digital arteries.

c. The *dorsal interosseous artery of the first space*, a branch of variable size, passes over the first interosseous muscle to the radial side of the back of the index finger.

d. The *dorsal arteries of the thumb* are two small branches which arise from the radial opposite the head of the first

metacarpal bone, and run along the back of the thumb, one on either side. They are often absent.

These dorsal interosseous arteries supply the extensor tendons and their sheaths, the interosseous muscles, and the skin on the back of the hand, and the first phalanges of the fingers.

Now remove the tendons from the back and from the palm of the hand: observe the deep palmar fascia which covers the interosseous muscles. It is attached to the ridges of the metacarpal bones, forms a distinct sheath for each interosseous muscle, and is continuous inferiorly with the transverse metacarpal ligament. On the back of the hand the interosseous muscles are covered by a thin fascia, which is attached to the adjacent borders of the metacarpal bones.

Transverse Metacarpal Ligament. — This consists of strong bands of ligamentous fibres, which pass transversely between the digital extremities of the metacarpal bones of the fingers. These bands are intimately united to the fibrocartilaginous ligament of the metacarpal joints, and are of sufficient length to admit of a certain degree of movement between the ends of the metacarpal bones.

Remove the fascia which covers the interosseous muscles, and separate the metacarpal bones by dividing the transverse metacarpal ligament. A *bursa* is frequently developed between their digital extremities.

Interosseous Muscles.—These muscles, so named from their position, extend from the sides of the metacarpal bones to the first phalanges and the extensor tendons of the fingers. In each interosseous space there are two, one of which is an abductor, the other an adductor of a finger. According to this arrangement there are eight in all, inclusive of the adductor pollicis, which is really an interosseous muscle: four of which, situated on the back of the hand, are called dorsal; the remainder, seen only in the palm, are called palmar.

Dorsal Interosseous. — Each *dorsal* interosseous muscle arises from the opposite sides of two metacarpal bones (fig. 48). From this double origin the fibres converge to a tendon, which passes between the metacarpal joints of the fingers, and is inserted into the side of the first phalanx; it is also

connected by a broad expansion with the extensor tendon on the back of the finger.

The *first* dorsal interosseous muscle (abductor indicis) is larger than the others, and occupies the interval between the thumb and fore finger. It arises from the upper half of the

Fig. 48.




DIAGRAM OF THE FOUR DORSAL
INTEROSSEI, DRAWING FROM
THE MIDDLE LINE.

Fig. 49.




DIAGRAM OF THE FOUR PALMAR
INTEROSSEI, DRAWING TOWARDS
THE MIDDLE LINE.

ulnar side of the first metacarpal bone, and from the entire length of the radial side of the second: between the two origins, the radial artery passes into the palm. Its fibres converge on either side to a tendon, which is inserted into the *radial* side of the first phalanx of the index finger and extensor tendon.

The *second* dorsal interosseous muscle occupies the second metacarpal space. It is inserted into the radial side of the first phalanx of the middle finger and its extensor tendon.

The *third* and *fourth*, occupying the corresponding metacarpal spaces, are inserted, the one into the ulnar side of the middle, the other into the ulnar side of the ring finger.

If an imaginary line be drawn longitudinally through the middle finger, as represented by the dotted line in fig. 48 we shall find that all the dorsal interosseous muscles are abduc-

tors from that line; consequently, they separate the fingers from each other.

Palmar Interosseous.—It requires a careful examination to distinguish this set of muscles, because the dorsal muscles protrude with them into the palm. They are smaller than the dorsal, and each arises from the lateral surface of only one metacarpal bone, that, namely, connected with the finger into which the muscle is inserted (fig. 49). They terminate in small tendons, which pass between the metacarpal joints of the fingers, and are inserted, like the dorsal muscles, into the sides of the first phalanges and the extensor tendons on the back of the fingers.

The *first* palmar interosseous muscle arises from the ulnar side of the second metacarpal bone, and is inserted into the ulnar side of the index finger. The *second* and *third* arise, the one from the radial side of the fourth, the other from the radial side of the fifth metacarpal bone, and are inserted into the same sides of the ring and little fingers. The *fourth* interosseous muscle is the adductor pollicis.

The palmar interosseous muscles are all adductors to an imaginary line drawn through the middle finger (fig. 49). They are, therefore, the opponents of the dorsal interosseous muscles, and move the fingers towards each other.

DISSECTION OF THE LIGAMENTS.

Joint between the Clavicle and the Sternum.—The inner end of the clavicle articulates with a shallow excavation on the upper and outer part of the sternum. From the form of the articular surfaces of the bones, it is plain that the security of the joint must depend upon the strength of its ligaments. There are two synovial membranes, and an intervening fibro-cartilage.

The *anterior sterno-clavicular ligament* (fig. 50) consists of a broad band of ligamentous fibres, which pass obliquely downwards and inwards over the front of the joint, from the end of the clavicle to the anterior surface of the sternum.

Fig. 50.




DIAGRAM OF THE STERNO-CLAVICULAR LIGAMENTS.

- | | |
|--|------------------------------------|
| 1. Interclavicular ligament. | 3. Costo-clavicular ligament. |
| 2. Anterior sternoclavicular ligament. | 4. Interarticular fibro-cartilage. |

A similar band, called the *posterior sternoclavicular* ligament, extends over the back of the joint, from the back of the clavicle to the back of the sternum.

The clavicles are connected by the *inter-clavicular ligament*. It extends transversely above the notch of the sternum, and has a broad attachment to the upper border of each clavicle. Between the clavicles it is more or less attached to the sternum, so that it forms a curve with the concavity upwards.

The three ligaments just described are so closely connected, that, collectively, they form a complete fibrous capsule of great strength for the joint.

A ligament, called the *costo-clavicular* or *rhomboid*, connects the clavicle to the cartilage of the first rib. It ascends from the cartilage of the rib to a rough surface beneath the sternal end of the clavicle. Its use is to limit the elevation of the clavicle. There is such constant movement between the clavicle and the cartilage of the first rib that a well-marked *bursa* is commonly found between them.

Interarticular fibro-cartilage.—In order to see this we must cut through the rhomboid, the anterior and posterior ligaments of the joint, and then raise the clavicle. It is nearly circular in form, and thicker at the circumference than the centre, in which there is sometimes a perforation. Inferiorly, it is attached to the cartilage of the first rib, close to the sternum; superiorly, to the upper part of the clavicle and the in-

ter-clavicular ligament. Its circumference is inseparably connected with the fibrous capsule of the joint.

The joint is provided with two synovial membranes : one is placed between the articular surface of the sternum and the inner surface of the fibro-cartilage ; the other, between the articular surface of the clavicle and the outer surface of the fibro-cartilage.

This interarticular fibro-cartilage is a structure highly elastic, without admitting of any stretching. It equalises pressure, prevents shocks, and also acts as a ligament, preventing the clavicle from being driven inwards towards the mesial line.

Observe the relative form of the cartilaginous surfaces of the bones : that of the sternum is slightly concave in the transverse, and convex in the antero-posterior direction ; that of the clavicle is the reverse.

The form of the articular surfaces and the ligaments of a joint being known, it is easy to understand the movements of which it is capable. The clavicle can move upon the sternum in a direction either vertical or horizontal : thus it admits of circumduction. These movements, though very limited at the sternum, become very considerable at the apex of the shoulder.

Connection of the Clavicle with the Scapula.—The outer end of the clavicle articulates with the acromion, and is connected by strong ligaments to the coracoid process of the scapula.

Joint between the acromion and the clavicle.—The clavicle and the acromion articulate with each other by means of two flat oval cartilaginous surfaces, of which the planes slant inwards, and the greater diameters are in the antero-posterior direction.

The *superior ligament*, a broad band of ligamentous fibres, strengthened by the aponeurosis of the trapezius, extends from the upper surface of the acromion to the upper surface of the clavicle.

The *inferior ligament*, of less strength, extends along the under surface of the joint from bone to bone.

An *interarticular fibro-cartilage* is sometimes found in this

joint : but it is incomplete, and seldom extends lower than the upper half. There is only one synovial membrane.

Coraco-clavicular ligament.—The clavicle is connected to the coracoid process of the scapula by two strong ligaments—the *conoid* and *trapezoid*, which, being continuous with each other, might fairly be considered as one. The *trapezoid* ligament is the more anterior. It arises from the back part of the coracoid process, and ascends obliquely outwards to the clavicle, near its outer end. The *conoid* ligament is fixed at its apex to the root of the coracoid process, ascends vertically, and is attached by its base to the clavicle, near the posterior border. Remember that when the clavicle is fractured in the line of the attachment of the coraco-clavicular ligament, there is no displacement of the fractured ends ; these being kept in place by the ligament itself.

Ligaments proper to the scapula.—There are only two : the *coracoid* ligament, which is attached to the margins of the supra-scapular notch, and the *coraco-acromial* or *triangular* ligament, which is attached by its apex to the acromion, and by its base to the outer border of the coracoid process. It is separated from the upper part of the capsule of the shoulder-joint by a large *bursa*.

Shoulder-joint.—The articular surface of the head of the humerus, forming rather more than one-third of a sphere, moves upon the shallow glenoid cavity of the scapula, which is of an oval form, with the broader end downwards, and the long diameter nearly vertical. It is obvious that the security of the joint does not depend upon any mechanical contrivance of the bones, but upon the great strength and number of the tendons which surround it.

To admit the free motion of the head of the humerus upon the glenoid cavity, it is requisite that the *capsular ligament* of the joint be very loose and capacious. Accordingly, we find that the head of the bone, when detached from its muscular connections, may be separated from the glenoid cavity to the extent of an inch, or even more, without laceration of the capsule. This explains the elongation of the arm observed in some cases in which effusion has taken place into the cavity of the joint ; also in some cases of paralysis of the deltoid muscle.

The *capsular ligament* is attached, above, round the circumference of the glenoid cavity; and below, round the surgical neck of the humerus. It is strengthened on its upper and posterior part by the tendons of the supra-spinatus, infra-spinatus, and teres minor muscles; its inner part is strengthened by the broad tendon of the subscapularis, and its lower part by the long head of the triceps. Thus the circumference of the capsule is surrounded by tendons on every side, excepting a small space towards the axilla. If the humerus be raised, it will be found that the head of the bone rests upon this unprotected portion of the capsule; that is, between the tendons of the subscapularis and the long head of the triceps; and through this part of the capsule the head of the bone usually protrudes in dislocations into the axilla.

At the upper and inner side of the joint, a small opening (*foramen ovale*) is observable in the capsular ligament, through which the tendon of the subscapularis passes, and comes in contact with the synovial membrane.

The upper surface of the capsule is, moreover, strengthened by a strong band of ligamentous fibres, called the *coraco-humeral* or *accessory ligament*. It is attached to the root of the coracoid process, expands over the upper surface of the capsule, with which it is inseparably united, and is fixed into the greater tuberosity of the humerus.

Open the capsule to see the tendon of the long head of the biceps. It enters the joint through the groove between the two tuberosities, becomes slightly flattened, and passes over the head of the bone to be attached to the upper border of the glenoid cavity. It is loose and movable within the joint; and it acts like a strap, keeping down the head of the bone. In some cases the tendon having been ruptured by violence, has contracted an adhesion to the bicipital groove.

The tendon of the biceps does not perforate the synovial membrane of the joint. It is enclosed in a kind of tubular sheath, which is reflected over it at its attachment to the glenoid cavity, and accompanies it for two inches down the groove of the humerus. During the earlier part of foetal life it is connected to the capsule by a fold of synovial membrane, which subsequently disappears.

The margin of the glenoid cavity of the scapula is sur-

rounded by a fibro-cartilaginous band of considerable thickness, called the *glenoid ligament*. This not only enlarges the cavity but renders it a little deeper. Superiorly, it is continuous on either side with the tendon of the biceps; inferiorly, with the tendon of the triceps: in the rest of its circumference, it is attached to the edge of the cavity.

Observe that the cartilage covering the head of the humerus is thicker at the centre than at the circumference. The reverse is the case in the glenoid cavity.

The *synovial membrane* lining the under surface of the capsule is reflected around the tendon of the biceps, and passes with it in the form of a cul-de-sac, down the bicipital groove. On the inner side of the joint it always communicates with a bursa beneath the tendon of the subscapularis.

The shoulder-joint has a more extensive range of motion than any other joint in the body; it is, in fact, a kind of universal joint. It is capable of motion forwards and backwards; of adduction, abduction, circumduction, and rotation.

[When the os brachia is dislocated downwards, the lower portion of the capsular ligament is torn, together with the tendon of the subscapularis. So also at times the tendons of the superior and inferior-spinati. When dislocated forwards, the inner surface of the capsule is torn, and also at times the tendon of the subscapularis.]

Elbow-joint.—The elbow-joint is a perfect hinge. The larger sigmoid cavity of the ulna is adapted to the trochlea upon the lower end of the humerus, admitting of simple flexion and extension; while the shallow excavation upon the head of the radius admits not only of flexion and extension, but also of rotation upon the rounded articular eminence (*capitellum*) of the humerus. The joint is secured by two strong lateral ligaments. No ligament is attached to the head of the radius, otherwise its rotatory movement would be impeded. It is simply surrounded by a ligamentous collar, called the annular ligament, within which it freely rolls.


Internal lateral ligament.—This is triangular. Its apex is attached to the internal condyle of the humerus: from this point the fibres radiate, and are inserted into the greater sigmoid cavity of the ulna.

A transverse band of ligamentous fibres extends from the

olecranon to the coronoid process, across a notch observable on the inner side of the sigmoid cavity: through this notch vessels pass into the joint.

External lateral ligament.—This is attached to the external condyle in the humerus; the fibres spread out as they descend, and are interwoven with the annular ligament, surrounding the head of the radius.

Fig. 51.



LIGAMENTS OF THE ELBOW.

The orbicular or annular ligament of the radius (fig. 51), is attached to the anterior and the posterior border of the lesser sigmoid cavity of the ulna. With this cavity, it forms a complete collar, which encircles the head and part of the neck of the radius. The lower part of the collar is narrower than the upper, in order to fit the neck of the radius, and maintain it more accurately in its position.

Some anatomists speak of an *anterior* and a *posterior* ligament for the elbow-joint. But what they describe as ligaments consists merely of a few thin ligamentous fibres spread over the capsule of the joint in front and behind. There is no need of ligaments to limit flexion and extension in this joint: for the coronoid process limits the one, and the olecranon the other.

The preceding ligaments collectively form a continuous capsule for the elbow-joint.

Synovial membrane of the elbow-joint.—Open the joint by a transverse incision through the anterior ligament, and observe the relative adaptation of the cartilaginous surfaces of the bones. The synovial membrane lines the interior of the capsule, and forms a cul-de-sac between the head of the radius and its orbicular ligament. It is widest and most loose under the tendon of the triceps. Where the membrane is reflected from the bones upon the ligaments, there is more or less adipose tissue, particularly in the fossæ on the front and back part of the lower end of the humerus.

Interosseous Ligament or Membrane.—This is an aponeurotic septum, which is stretched between the bones of the forearm, and of which the chief purpose is to afford an increase of surface for the attachment of muscles. The septum is deficient above, in order to permit free rotation of the radius. Its fibres extend obliquely downwards from the radius to the ulna.

The name of *round* or *oblique ligament* is given to a thin band of ligamentous fibres, which extend obliquely between the bones of the forearm in a direction contrary to those of the interosseous membrane. It is attached, superiorly, to the front surface of the ulna, near the outer side of the coronoid process; inferiorly, to the radius immediately below the tubercle. A *bursa* intervenes between it and the insertion of the tendon of the biceps. The use of this ligament is to limit supination of the radius.

[When the radius and ulna are dislocated backwards, the external, internal, and at times the annular ligaments of the radius, are ruptured, though we do not necessarily have all these injuries together. When the ulna alone is dislocated backwards, the annular and accessory are ruptured, which also happens when the radius is dislocated. When dislocation occurs at the wrist, the capsular, and sometimes the lateral ligaments, are torn.]

Radio-carpal or Wrist-joint.—This joint is formed by the lower end of the radius, which articulates with the scaphoid and semilunar bones of the carpus: the lower end of the ulna is excluded from the joint by a triangular fibro-cartilage,

which articulates with a small portion of the cuneiform bone. The joint is secured by an anterior, a posterior, and two lateral ligaments, forming, together, an uninterrupted capsule around it.

The *external lateral ligament* extends from the styloid process of the radius, to the scaphoid bone (p. 279).

The *internal lateral ligament* proceeds from the extremity of the styloid process of the ulna, to the cuneiform bone. Some of its fibres are attached to the pisiform bone.

The *anterior ligament* consists of two or more broad bands of ligamentous fibres, which arise from the lower end of the radius, pass obliquely inwards, and are inserted into the first row of carpal bones.

The *posterior ligament*, weaker than the preceding, arises from the posterior surface of the lower end of the radius, descends obliquely inwards, and is inserted into the posterior surfaces of the first row of the carpal bones.

The *synovial membrane* lines the triangular fibro-cartilage at the end of the ulna, is reflected over the several ligaments of the joint, and thence upon the first row of the carpal bones.

Joint between the Lower Ends of the Radius and Ulna.—The inner surface of the lower end of the radius presents a slight concavity, which rotates upon the convex circumference of the lower end of the ulna: this mechanism is essential to the pronation and supination of the hand. These corresponding surfaces are crusted with a thin layer of cartilage, and are provided with a very loose synovial membrane. The joint is strengthened in front and behind by a thin fibrous capsule, which extends from the anterior and posterior borders of the sigmoid cavity of the radius, to the anterior and posterior surfaces of the styloid process of the ulna. But the principal uniting medium between the bones is a strong fibro-cartilage.

Fibro-cartilage between the radius and ulna.—Saw through the bones of the forearm, and separate them by cutting through the interosseous membrane, and opening the synovial membrane of the joint between their lower ends. So you will get a good view of the fibro-cartilage which connects them (fig. 52). It is triangular in form, and is placed transversely at the lower end of the ulna, filling up the interval caused by the greater length of the radius. Its base is attached to the lower end of

the radius, and its apex to the root of the styloid process of the ulna. It is thin at the base and the centre, thicker at the apex and the sides. Its upper surface is in contact with the ulna, and covered by the synovial membrane of the radio-ulnar joint; its lower surface forming a part of the wrist-joint, corresponds to the cuneiform bone. Its borders are connected with the anterior and posterior ligaments of the wrist. In some instances there is an aperture in the centre.

When, from accident or disease, this fibro-cartilage gets detached from the radius, the consequence is a preternatural projection of the lower end of the ulna.

Fig. 52.




DIAGRAM OF THE WRIST-JOINT.

The *synovial membrane* of this joint is distinct from that of the wrist, except in the case of a perforation through the fibro-cartilage. On account of its great looseness, necessary for the free rotation of the radius, it is often called the "*membrana sacciformis*."

Connection of the Carpal Bones with Each Other.—The bones of the carpus are arranged in two rows, an upper and a lower, adapted to each other, so as to form between

them a joint, connected by anterior, posterior, internal, and external lateral ligaments.

The bones, constituting each row, are united by ligaments placed on their palmar and dorsal surfaces, and by others, placed between the bones, and hence called interosseous. Their contiguous surfaces (those of the pisiform and cuneiform excepted) are covered by the reflections of one synovial membrane.

The *upper row* is united by *transverse* ligaments proceeding from the scaphoid to the semilunar bone, and from the semilunar to the cuneiform, both on their dorsal and palmar surfaces; also by *interosseous* ligaments, proceeding from the semilunar to the bones on either side of it (fig. 52).

The *pisiform bone* is articulated to the palmar surface of the cuneiform bone, to which it is united by a fibrous capsule. Inferiorly it is attached by two strong ligaments, the one to the unciform bone, the other to the carpal end of the fifth metacarpal bone. This little articulation has a distinct synovial membrane.

The *lower row* of carpal bones is connected precisely in the same way as the upper. The dorsal and palmar ligaments pass transversely from one to the other. There are only two interosseous ligaments, one on either side of the os magnum; they are much thicker and stronger than those of the upper row, and unite the bones more firmly together (fig. 52).

Intercarpal Joint.—The upper row of carpal bones is arranged in the form of an arch, so as to receive the corresponding convex surfaces of the os magnum and unciform. Externally to the os magnum, the trapezium and trapezoid bones present a slightly concave surface, which articulates with the scaphoid. In this way a joint capable of flexion and extension only is formed between the upper and lower row. It is secured by anterior, posterior, and two lateral ligaments. The anterior ligament consists of strong ligamentous fibres, which pass obliquely from the bones of the upper to those of the lower row. The posterior ligament consists of oblique and transverse fibres, which connect the dorsal surfaces of the bones of the upper with the lower row.

The lateral ligaments connect, externally, the scaphoid

and trapezium; internally, the cuneiform and unciform bones (fig. 52).

Divide the ligaments, in order to see the manner in which the carpal bones articulate with each other. Their surfaces are crusted with cartilage, and provided with a common *synovial membrane*. This membrane extends, superiorly, between the three bones of the upper row, so as to form two culs-de-sac; inferiorly, it is prolonged into the joint between the carpal and the second and third metacarpal bones.

Joint between Trapezium and the First Metacarpal Bone.—The trapezium presents a cartilaginous surface, convex in the transverse, and concave in the antero-posterior direction, (*i. e.* somewhat saddle-shaped), which articulates with a cartilaginous surface on the metacarpal bone of the thumb, concave and convex in just the opposite directions. This peculiar adaptation of the two surfaces permits the several movements of the thumb, *viz.*, flexion, extension, abduction, and adduction; consequently circumduction. Thus we are enabled to oppose the thumb to all the fingers. The joint is surrounded by a fibrous capsule sufficiently loose to admit free motion, and stronger on the dorsal than on the palmar aspect. The security of the joint is also increased by the muscles which surround it. It has a separate synovial membrane.

Carpo-Metacarpal Joints.—The metacarpal bones of the fingers are connected to the second row of the carpal bones by ligaments upon their *palmar* and their *dorsal* surfaces.

The *dorsal* ligaments are the stronger. The metacarpal bone of the fore finger has two; one from the trapezium, another from the trapezoid bone. That of the middle finger has also two, proceeding from the os magnum, and the os trapezoides. That of the ring finger has also two, proceeding from the os magnum, and the unciform bone. That of the little finger has one only, from the unciform bone.

The *palmar* ligaments are arranged nearly upon a similar plan. The metacarpal bone of the fore finger has one, from the trapezoid bone. That of the middle finger has three, proceeding from the trapezium, the os magnum, and the unciform bone. Those of the ring and little fingers have each one, from the unciform bone.

Besides the preceding ligaments, there is another of considerable strength, called the *interosseous*. It arises from the adjacent sides of the os magnum and the os unciform, descends vertically, and is fixed into the radial side of the metacarpal bone of the ring finger (fig. 52). This ligament isolates the synovial membrane of the two inner metacarpal bones from the common synovial membrane of the carpus.

Separate the metacarpal bones from the carpus, and observe the relative form of their contiguous surfaces. The metacarpal bones of the fore and middle fingers are adapted to the carpus in such an angular manner as to be almost immovable. The metacarpal bone of the ring finger, having a plane articular surface with the unciform bone, admits of more motion. But a still greater degree of motion is permitted between the unciform bone and the metacarpal bone of the little finger; the articular surfaces of each being slightly concave and convex in opposite directions. The greater freedom of motion of the metacarpal bone of the little finger is obviously essential to the expansion and contraction of the palm.

The *carpal extremities of the metacarpal bones of the fingers* are connected with each other by transverse ligaments, both on their dorsal and their palmar surfaces. They are also connected by interosseous ligaments, which extend between the bones, immediately below their contiguous cartilaginous surfaces.

The *digital extremities* of these bones are loosely connected on their palmar aspect by the transverse metacarpal ligament.

Synovial Membranes of the Wrist.—There are six distinct synovial membranes, proper to the lower end of the radius, and the several bones of the carpus: see the diagram (p. 281).

- a. One between the lower end of the radius and the ulna.
- b. One between the radius and the first row of carpal bones.
- c. One between the trapezium and the metacarpal bone of the thumb.
- d. One between the cuneiform and pisiform bones.
- e. One between the first and second rows of carpal bones, (the intercarpal joint). This extends to the metacarpal bones of the fore and middle fingers.

f. One between the unciform bone and the metacarpal bones of the little and ring fingers.

First Joint of the Fingers.—The first phalanx of the finger presents a shallow oval cavity, crusted with cartilage, with the broad diameter in the transverse direction, to articulate with the round cartilaginous head of the metacarpal bone, of which the articular surface is elongated in the antero-posterior direction, and of greater extent on its palmar than its dorsal aspect. This formation of parts permits flexion of the finger to a greater degree than extension; and also a slight lateral movement.

Each joint is provided with two strong *lateral*, and a *palmar* or *glenoid* ligament.

The *lateral* ligaments arise from the tubercles on either side of each metacarpal bone, and inclining slightly forward, are inserted into the sides of the base of the first phalanx of the finger.

The *palmar* or *glenoid* ligament. This name is applied to a thick, compact, fibro-cartilaginous structure, which extends over the palmar surface of the joint. Its lower end is firmly attached to the base of the first phalanx of the finger; its upper end is loosely adherent to the rough surface above the head of the metacarpal bone. On either side it is inseparably connected with the lateral ligaments, so that with them it forms a strong capsule over the front and sides of the joint. Its superficial surface is slightly grooved, to receive the flexor tendons; its deep surface is adapted to cover the head of the metacarpal bone. Two sesamoid bones are found in the glenoid ligament belonging to the joint between the metacarpal bone and the first phalanx of the thumb.

The glenoid ligaments have a surgical importance for the following reason:—In dislocation of the fingers, the facility of reduction mainly depends upon the extent to which the glenoid ligament is injured. If it be much torn there is but little difficulty: if entire the reduction is extremely tedious and sometimes impracticable.


These joints are secured on their dorsal aspect by the passage of the extensor tendon, and the expansion proceeding from it on either side. Their synovial membranes are loose, especially beneath the extensor tendons.

Second and Last Joint of the Fingers.—The corresponding articular surfaces of the phalanges of the finger and thumb are so shaped as to form a hinge-joint, and, therefore, incapable of lateral movement. The ligaments connecting them are similar in every respect to those between the metacarpal bones and the first phalanges. The glenoid ligament of the last joint of the thumb generally contains a sesamoid bone.

THE DISSECTION OF THE ABDOMEN.

Arbitrary Division into Regions.—We divide the abdomen into arbitrary regions, that the situation of the viscera contained in it may be more easily described. For all practical purposes, the ancient division, handed down from the Greek authors, is sufficiently convenient. An imaginary horizontal line is drawn across the abdomen on a level with the cartilage of the ninth rib; another on a level with the anterior spine of the ilium. That part of the cavity above the upper

Fig. 53.



line is called the *epigastric* region; that below the lower one, the *hypogastric*; while the space included between the lines is termed the *mesogastric* or umbilical. These again are subdivided thus:—the sides of the epigastric region, beneath the costal cartilages, are termed respectively the right and left *hypocondria*, and the depression in front just below the ensiform cartilage, is called the *scrobiculis cordis*, or the pit of the stomach. The sides of the umbilical region are named

the *ilia* or flanks, and the back part, the *loins*. The sides of

the hypogastric region are familiarly known as the groins, while the middle of it is called the *pubes*.

[The sides of the umbilical region are also known as the right and left lumbar, while those of the hypogastric are spoken of as the right and left iliac, or inguinal. *The organs contained in these zones* are usually as follows :—**EPIGASTRIC**: Cardiac, and pyloric extremity of the stomach; lesser curvature and a portion of the lower curvature; commencement of the duodenum, and a portion of the liver. *Right hypochondria*: Great lobe of the liver; the gall bladder; the hepatic flexure of the large intestine, and a small portion of the duodenum. *Left hypochondria*: Cul-de-sac of the stomach, sometimes a portion of liver, spleen, left end of the pancreas, and a knuckle of the large intestine. **UMBILICAL**: The small intestines are here attached. *Right iliac*: Ascending colon. *Left iliac*: Descending colon. Behind these organs, we have a region called the lumbar, where are situated the kidneys. **HYPOGASTRIC-PUBIC**: Bladder and rectum: in the female, bladder, uterus and appendages, and rectum. *Right inguinal*: Commencement of large and termination of small intestine. *Left inguinal*: Sigmoid flexure of the colon.]


An incision should be made from the sternum to the pubes, another from the anterior spine of the ilium to the umbilicus, and a third from the umbilicus obliquely outwards, over the cartilages of the ninth and tenth ribs, towards the axilla. The skin should then be dissected from the subjacent adipose and areolar tissue, called the "superficial fascia."

Superficial Fascia.—The subcutaneous tissue of the abdomen presents the same general characters as that of other parts, and varies in thickness according to the condition of the individual. At the lower part of the belly, it admits of separation into two layers; the subcutaneous vessels and nerves ramify between them. Respecting the superficial layer, there is nothing to be observed, except that it contains the fat. The deeper layer is intimately connected with Poupart's ligament or the crural arch; but it is very loosely continued over the spermatic cord and the scrotum, and becomes identified with the superficial fascia of the perineum. These points deserve attention, because they explain how urine, when extravasated into the perineum and scrotum, readily

makes its way over the spermatic cord on to the surface of the abdomen; but from this it cannot travel down the thigh, on account of the connections of the fascia with Poupart's ligament.

Superficial Blood-Vessels and Glands.—Between the layers of the superficial fascia on the groin and the upper part of the thigh, are several absorbent glands and small blood-vessels (fig. 54). The glands are named, according to their situation, inguinal and femoral. The inguinal are often small enough to escape observation. They are of an oval form, with their long axis corresponding to the line of the crural arch (represented by the dark line in fig. 54). They receive the absorbents from the wall of the abdomen, from

Fig. 54.



SUPERFICIAL VESSELS AND GLANDS OF THE GROIN.

- | | |
|--|------------------------------|
| 1. Saphenous opening of the fascia lata. | 5. Superficial pudic a. |
| 2. Saphena vein. | 6. External abdominal ring. |
| 3. Superficial epigastric a. | 7. Fascia lata of the thigh. |
| 4. Superficial circumflexa ili a. | |

the scrotum and penis, and are therefore generally affected in venereal disease.

The *superficial arteries* in the neighborhood arise from the femoral. One, the *epigastric*, ramifies over the lower part of the abdomen; another, the *pudic*, crosses the spermatic cord, and is distributed to the skin of the penis and scrotum; a third, the *circumflexa ilii*, ramifies towards the spine of the ilium. These subcutaneous arteries, the pudic especially, often occasion a free hemorrhage in the operation for strangulated hernia.

The corresponding *veins* join the saphena vein of the thigh. Under ordinary circumstances they do not appear in the living subject; but, when any obstruction occurs in the inferior vena cava, they become enlarged and tortuous, and constitute the chief channels through which the blood would be returned from the lower limbs.*

Cutaneous Nerves.—The skin of the abdomen is supplied with nerves after the same plan as that of the chest; namely, by lateral and anterior branches derived from the intercostal nerves.

a. The *lateral cutaneous nerves* come out between the digitations of the external oblique muscle, in company with small arteries, and divide into anterior and posterior branches. Observe, however, that the lateral branch of the twelfth dorsal nerve is larger than the others, and that it passes over the crest of the ilium to the skin of the buttock. The corresponding branch of the first lumbar nerve has a similar distribution.

b. The *anterior cutaneous nerves* come with little arteries through the sheath of the rectus muscle. They are not only smaller than the lateral nerves, but their number and place of exit are less regular. That which comes through the external abdominal ring, (*ilio-inguinal*) as well as another which comes through the wall of the abdomen just above it, (*ilio-hypogastric*) is derived from the first lumbar nerve. These, however, are but repetitions of the others, and supply the skin of

* A cast, in illustration of this, is preserved in the Museum of St. Bartholomew's Hospital.

the groin and scrotum in the male, and the libium pudendi in the female.*

Muscles of the Abdominal Wall.—Next examine the abdominal muscles. There are three on each side, arranged in strata, named after the direction of their fibres,—the “external oblique,” “internal oblique,” and “transversalis.” They terminate in front in strong aponeuroses, arranged so as to form a sheath for a broad muscle called the “rectus,” which extends perpendicularly on each side the linea alba from the sternum to the pubes (fig. 55, p. 293).

External Oblique.—This muscle is attached to the eight or nine lower ribs, by as many pointed bundles, termed “digitations.” The upper five of these fit in with similar bundles of the serratus magnus, and are obvious during life; the three lower correspond in like manner with the origin of the latissimus dorsi; but they cannot be seen unless the body be turned on the side. The greater part of the muscle descends obliquely forwards, and terminates on the aponeurosis of the abdomen; the remainder proceeds almost perpendicularly from the last ribs, and is attached to the anterior two-thirds of the crest of the ilium.†

The *aponeurosis* of the external oblique increases in strength, breadth, and thickness, as it approaches the lower margin of the belly, this being the situation where the greater pressure of the viscera requires the most effective support. Its tendinous fibres take the same direction as the muscle; they do not, however, stop at the middle line, or linea alba, but cross over those of the opposite side; a decussation which makes the aponeurosis all the stronger.

Poupart's Ligament or Crural Arch.—Along the line of junction of the abdomen and thigh, the aponeurosis extends from the anterior spine of the ilium to the spine of the pubes, and forms an arch over the intermediate bony excavation

* A third small nerve, namely, the genital branch of the genito-crural, comes through the external ring. It is found behind the cord, close to the outer pillar.

† From its position and the direction of its fibres, it is manifest that the external oblique represents, in the abdomen, the external intercostal muscles of the chest.

(p. 296). This, which is termed *the crural arch*,* transmits the great vessels of the thigh, with muscles and nerves, which will be examined hereafter. Near the pubes is an opening in the aponeurosis for the passage of the spermatic cord in the male, and the round ligament of the uterus in the female. It is called the "external abdominal ring." Of this, as well as of the crural arch, you must postpone more particular examination till you come to the parts concerned in inguinal hernia.

The external oblique should be carefully detached from the ribs and the crest of the ilium, and turned forwards as far as this can be done without injuring its aponeurosis or the crural arch. In this way the second muscular stratum will be exposed.

[The external oblique (*descendens*) is spoken of by some as having four insertions,—1, tendinous, into the linea alba; 2, into the spine of the pubes, and reflected for a half inch or more along the linea pectinea, forming Gimbernat's ligament; 3, into the anterior-superior spinous process of the ilium; 4, fleshy, into the anterior two-thirds of the outer lip of the crest of the ilium. This muscle may be defective at various points, thus giving rise to ventral hernia. The student should also notice the *linea semilunaris*, extending from the pubic spine upwards and outwards, about four inches from the linea alba: it is formed from the tendon of the internal oblique, dividing into two portions at the edge of the rectus, for the purpose of enveloping that muscle. The lineal transversæ are those lines seen crossing the rectus, usually three or four in number, being tendinous intersections of that muscle.]

Internal Oblique.—This muscle *arises* from the outer half of the crural arch, and from the anterior two-thirds of the crest of the ilium. From these attachments the fibres ascend obliquely, and are inserted partly into the abdominal aponeurosis, partly into the cartilages of the three or four lower ribs.†

* This was first described by Fallopius, an Italian anatomist, in his "*Observationes Anatomicæ*," published in 1561. It was subsequently described by Poupart in 1705, in the "*Mem. de l'Acad. de Paris*," and is now commonly called in schools "*Poupart's ligament*."

† The internal oblique represents, in the abdomen, the internal intercostal muscles of the chest.

The internal oblique should be detached from the ribs, and from the crest of the ilium; but that portion of it connected to the crural arch must not be disturbed. In removing the internal oblique, we are very apt to cut away the transversalis. To avoid this mistake you should dissect near the crest of the ilium, and endeavor to find an artery which runs between the muscles, and serves as a guide. The artery, called the deep "*circumflexa ilii*," proceeds from the external iliac, and supplies the abdominal muscles. Beneath the internal oblique, too, you will bring into view the continuation of the intercostal nerves and vessels. These it is desirable to preserve.

[The internal oblique (*ascendens*) may, like the external, be deficient, so as to lead to ventral hernia.]

Transversalis.—This muscle *arises* from the outer half of the crural arch, from the anterior two-thirds of the crest of the ilium, from a fascia attached to the transverse processes of the lumbar vertebræ, and, lastly, from the inner surfaces of the six or seven lower costal cartilages, by digitations which correspond with those of the diaphragm. From this manifold origin the fibres proceed horizontally forwards, and terminate in the abdominal aponeurosis.

Rectus.—This muscle lies perpendicularly along the front of the belly, and is enclosed in a sheath formed by the aponeuroses of the lateral muscles of the abdomen. To expose it, therefore, we must slit up its sheath. It *arises* by a flat tendon from the symphysis, and from the upper part of the pubes; and is inserted into the fifth, sixth, and seventh costal cartilages. You notice the tendinous intersections across the muscle called the "*lineæ transversæ*," which are only incomplete repetitions of the ribs in the wall of the abdomen.* Their number varies from three to five, but there are always more above than below the umbilicus. Observe, moreover, that these tendinous intersections adhere closely to the sheath in front, but not behind: consequently, matter formed between the front of the rectus and its sheath would be confined by two intersections; not so on the back of the muscle, for matter might travel down the entire length of it.

How is the sheath of the rectus formed? The front of the sheath comprises the aponeurosis of the external oblique, and

* Some animals—*e. g.* the crocodile—have bony abdominal r.bs.

half the thickness of that of the internal oblique ;* while the

Fig. 55.




DIAGRAM OF A TRANSVERSE SECTION THROUGH THE ABDOMINAL MUSCLES.

back of the sheath comprises the aponeurosis of the transversalis, and the other half of the internal oblique (fig. 55). This however, applies only to the upper three-fourths of the muscle ; the lower fourth has no sheath behind, for all the aponeuroses pass in front of it.

Linea Alba.—

Along the middle line of the belly the several aponeuroses decussate so as form a white tendinous line extending from the sternum to the pubes. This is the "*linea alba* ;" it is but a continuation of the sternum deprived of its earthy matter, in adaptation to the functions of this part of the body. A little below the middle of it is situated what was in the foetus the opening of the passage of the umbilical vessels. After birth, the vessels being no longer required, the opening gradually closes, and becomes plugged by their fibrous remains.

Since the linea alba is the thinnest part of the abdomen, and is free from blood-vessels, it is chosen as a safe line for tapping in dropsy, and for puncturing the bladder in retention of urine.

Pyramidalis.—This little muscle is situated near the pubes, close to the linea alba and has a little sheath of its own. It arises from the upper part of the pubes in front of the rectus, and terminates in the linea alba about midway between the pubes and the umbilicus. It is said to be of use in tightening the linea alba, but it does not appear to have any special purpose in the human subject, for it is sometimes deficient on one or even both sides.

* The line where the internal oblique splits—namely, along the outer border of the rectus—is called the *linea semilunaris*.

By dividing the rectus transversely near the umbilicus, and raising it from its position, we have a complete view of the manner in which the sheath is formed; we observe, too, that this is very indistinct behind the lower fourth of the muscle. Ramifying in the substance of the muscle is a large artery, called the "epigastric," a branch of the external iliac; also the continuation of the internal mammary, which descends from the subclavian.

Nerves of the Abdominal Wall.—These nerves are the continuations of the six lower intercostal nerves, and of the first lumbar. They have the same general course and distribution. We trace them running forwards between the internal oblique and transversalis towards the rectus. They furnish branches to the abdominal muscles, and each gives off its lateral and anterior cutaneous branches, described p. 289.

Action of the Abdominal Muscles.—In consequence of their stratified arrangement, and the different direction of their fibres, the abdominal muscles answer many important purposes:—1. They are the principal muscles concerned in forcible expiration; 2. By compressing the viscera in conjunction with the diaphragm, they are the chief agents in the expulsion of the fæces and urine, also in vomiting, sneezing, laughing, coughing; 3. They act, each in its own way, as movers of the trunk: *e. g.* the right external oblique, co-operating with the left internal oblique, can draw the trunk towards the left side, and *vice versâ*. The recti raise the body from the horizontal position, as any one may ascertain by laying the hand on the abdomen while rising from the ground.

EXAMINATION OF THE PARTS CONCERNED IN INGUINAL HERNIA.

General Idea of the Subject.—Presuming that you have proceeded so far in the examination of the abdominal wall, now turn your attention to the anatomy of the parts concerned in inguinal hernia.

The testicle, originally formed in the loins, subsequently passes through the wall of the abdomen into the scrotum; its blood-vessels and nerves and excretory duct, drawn down

after it, constitute the spermatic cord; and the passage through which it comes through the abdomen is called the "inguinal canal." This canal is not direct, but oblique, that the abdominal wall may the better resist protrusion of the viscera. We have seen that the wall of the abdomen is composed of strata of different structures; and we shall presently find that the spermatic cord, as it passes through each stratum, derives from each a covering similar in structure to the stratum itself. Of these strata there are three: the first—that is, the aponeurosis of the external oblique—we shall call the "aponeurotic stratum;" the second—that is, the internal oblique and transverse muscles—we shall call the "muscular stratum;" the third—namely, the fascia transversalis which lines the under surface of the transverse muscle—we shall call the "fascial stratum." The most intelligible way of investigating the subject is to examine each stratum as it appears on dissection, and then to consider the inguinal canal as a whole. First, then, of the aponeurotic stratum.

Aponeurotic Stratum and External Abdominal Ring.—The lower border of the aponeurosis of the external oblique, extending from the spine of the ilium to the spine of the pubes, constitutes Poupart's ligament, or the crural arch. We have now to examine more in detail the opening in the aponeurosis which gives passage to the spermatic cord; in other words, the "*external abdominal ring*," or the lower aperture of the inguinal canal. This ring, then, is situated immediately above and to the outer side of the spine of the pubes: its size and shape vary a little in different individuals. It is an oval opening, with the long axis directed obliquely downwards and inwards, and will admit the passage of a finger.* These remarks apply to the male; in the female the opening is smaller, and transmits the round ligament of the uterus. The sides of the opening are called its pillars or columns†, and their re-

* To ascertain in the living subject whether the ring be free or otherwise, the best plan is to push the thin skin of the scrotum before the finger; then, by tracking the cord, we can readily pass the finger into the ring behind the fat which covers the pubes.

† These, and many other expressions in the language of anatomy may appear to us at first rather far-fetched or even affected, but we soon conform to them as quite natural and proper, because universally received amongst those whom they most concern.

pective attachments should be thoroughly investigated. The internal pillar is attached to the front of the pubes, not only of its own side, but also of the opposite; so that there is a decussation of aponeurotic fibres in front of the symphysis

Fig. 56.




DIAGRAM OF POUPART'S LIGAMENT, OF THE APONEUROSIS OF THE EXTERNAL OBLIQUE, AND OF THE EXTERNAL ABDOMINAL RING.

just as there is in the linea alba. The external pillar is stronger and much more firmly secured, because there is a greater strain upon it. We find that it has no less than three points of attachment, each of which, being considered as a ligament, has received a special name. One, perhaps the strongest attachment, is to the spine of the pubes (Poupart's ligament). The second extends for nearly an inch along the linea ilio-pectinea (Gimbernat's ligament). The third, and weakest, consists of a few fibres, which pass beneath the internal pillar to the linea alba, and are, in fact, continuous with the aponeurosis of the opposite side: (this is called the triangular ligament). There can be no objection to these ligaments having separate names, provided we understand that they are nothing more than the attachments of the outer pillar of the ring.

For some distance above the opening, observe that the cross fibres of the aponeurosis are strongly marked. These cross fibres, which round off the upper margin of the opening, are so conspicuous that anatomists have called them the *inter-columnar bands*. Their purpose plainly is to prevent the ring from enlarging.

In its passage through the ring, the spermatic cord rests upon the outer pillar, and receives from the margin of the opening a covering, called "*the spermatic fascia*."

Muscular Stratum and Cremaster. — To examine the muscular stratum we must reflect the aponeurosis of the external oblique. This is best done by making a transverse incision through it from the spine of the ilium to the linea alba, and another at right angles to the first, down to the pubes. We can thus turn down a triangular flap of the aponeurosis without injuring either the external ring or the crural arch. This done, the muscular stratum is fairly exposed. Now this stratum consists of the combined fibres of the internal oblique and transverse muscles: these, so far as our present subject is concerned, may be considered as one, having the same origin, direction, and insertion. Their origin is from the outer half of the crural arch; their direction is transversely towards the mesial line, and they terminate upon a thin aponeurosis termed the "*conjoined tendon*," which is inserted into the upper part of the pubes and the linea alba (fig. 57). Observe that this tendon is situated immediately behind the external ring, plainly for the purpose of strengthening the abdominal wall just at a part where, without such provision, the liability to hernia would have been very great.

The spermatic cord passes under this muscular stratum, and derives from it a covering, called the "*cremaster*," or suspensory muscle of the testicle. The fibres of the cremaster are thin and pale, or the reverse, according to the condition of the subject. In bundles, arising from the crural arch, descend most of them in front of the cord, and then arch up again towards the pubes. Thus they form a series of loops of different lengths; some reaching only as low as the external ring, others lower still, while the lowest cover the tunica vaginalis of the testicle. In very muscular subjects you may sometimes succeed in tracing some of the returning fibres up

to the pubes, as shown in the diagram, where the loops are purposely exaggerated: but this is quite the exception.

Fig. 57.




DIAGRAM OF THE LOWER FIBRES OF THE INTERNAL OBLIQUE AND TRANSVERSALIS, WITH THE CREMASTER MUSCLE.

The cremaster is very evident in subjects where there has been a hydrocele or a rupture. Its nerve comes from the first lumbar, which also supplies the internal oblique and transversalis. Its artery is a branch of the deep epigastric.

Fascial Stratum.—This third and last stratum is nothing more than the fascia which lines the transversalis muscle. For many reasons it is most advantageous to examine it from behind: we should therefore turn down a triangular flap of the abdominal wall, just as was done with the aponeurosis of the external oblique.

On the inner surface of the flap, thus reflected, is the peritoneum. If the parts be all kept together and made tense, we notice that the peritoneum is raised into a fold by a thick fibrous cord, which runs up obliquely towards the umbilicus. This cord is the impervious remains of the umbilical artery. If the peritoneal fold be well marked, it appears triangular with the base below, and occasions two pouches, one on either side of it. The depth and distinctness of these pouches will be in proportion to the prominence of the fold itself; but if

it happen that the cord be closely attached to the abdominal wall, then there will be no fold or pouch.* Take care not to confound the remains of the umbilical artery with what is called the "*urachus*." This is another cord, a remnant of embryonic life, which ascends from the bladder directly behind the linea alba to the umbilicus.

Having examined the remains of the umbilical artery and the urachus, now reflect the peritoneum from the abdominal flap. This is done easily enough, for it is loosely connected to the abdominal wall by an abundance of soft cellular tissue, which generally contains a considerable quantity of fat. A tissue of a similar kind connects the peritoneum to the iliac region; and this accounts for the facility with which the peritoneum can be extended so as to envelope even the largest herniæ.

The fascia lining the under surface of the transversalis muscle was first accurately described by Sir A. Cooper, and called by him the "*fascia transversalis*" (fig. 58). Now what are the connections of this important fascia? Speaking generally, one may say that it is attached to the crural arch, thence ascends, and gradually diminishing in thickness, is lost on the under surface of the transversalis muscle. Its inner border is connected to the margin of the rectus and also to the lower margin of the conjoint tendon. Observe that this fascia is nowhere stronger than just behind the external ring, and but for it and the conjoint tendon, there would be a *direct* opening into the cavity of the belly through the ring. You will find that the *outer* half of the fascia is very firmly connected to the crural arch and also to the fascia iliaca: but the *inner* half is loosely connected with the crural arch; and passes down under it over the femoral vessels into the thigh, and forms what is called "the sheath of the femoral vessels." The opening in the fascia through which the spermatic cord passes is called the "internal abdominal ring."†

Internal Abdominal Ring.—This must be accurately ex-

* These peritoneal pouches are considered to favor the occurrence of hernia, and so much importance has been attached to them by some continental surgeons, that they have divided inguinal herniæ into external and internal, according as the protrusion takes place in the outer or the inner fossa.

† Or the inner aperture of the inguinal canal.

amined. First, as to its precise situation. It corresponds with a point midway between the anterior superior spine of the ilium and the symphysis pubis, and about two-thirds of an inch above the crural arch. As to its shape, you had

Fig. 58.




DIAGRAM OF THE FASCIA TRANSVERSALIS SEEN FROM THE FRONT.

better introduce the little finger into it, and judge for yourselves. You will find that it has a defined margin on the pubic side, but none on the iliac, and that on the whole it looks like a little funnel.

In its passage through the internal abdominal ring the spermatic cord receives from the margin of it its first covering, thin and delicate, termed the infundibuliform fascia. (This is not seen in the diagram.) But the chief point of interest concerning the internal ring is, that the epigastric artery ascends close by its inner border.

Having examined the several strata through which the spermatic cord passes, you will do well to replace them in their natural position, and build them up so as to examine the inguinal canal as a whole. You will perceive that its direction is obliquely downwards and inwards. Its length in a well-formed adult male is from one and a half to two inches.

Now, what are its boundaries? It is bounded in front by the aponeurosis of the external oblique; behind, by the fascia transversalis; above, by the lower fleshy fibres of the internal oblique and transversalis; below, by the crural arch.

Course and Relations of the Epigastric Artery.—In a practical point of view this is one of the most important arteries in the body. It arises from the external iliac, just before this vessel passes under the crural arch. It ascends inwards, forms a gentle curve round the inner side of the internal abdominal ring, and consequently on the inner side of the spermatic cord, and then enters the rectus muscle, in which it is gradually lost. In the first part of its course it lies between the peritoneum and the fascia transversalis; afterwards it enters the sheath of the rectus. The artery is accompanied by two veins, of which the larger is constantly found on the inner side of it. They terminate by a single trunk in the iliac vein.

Branches of the Epigastric.—Of the branches of the epigastric artery the most important is the "*pubic*." It runs inwards, just behind the crural arch, towards the pubes, and derives its chief practical interest from the fact that it is liable to be wounded in dividing the stricture in femoral hernia.* But its size varies in different subjects, and is sometimes so small as to escape observation. The second branch is the "*cremasteric*." It supplies the coverings of the cord, and chiefly the cremaster muscle. After giving off other unnamed muscular branches, the main trunk terminates in the rectus by minute inosculations with the internal mammary.

Such is a brief outline of the anatomy of the parts concerned in inguinal hernia. This description applies equally to the female, provided the round ligament be substituted for the spermatic cord. Of course the inguinal canal is proportionably smaller and there is no cremaster.

Nomenclature of the Several Kinds of Inguinal Hernia.—When a piece of intestine escapes along the inguinal canal with the cord, and protrudes through the outer ring, it is called an *oblique* inguinal hernia. If the intestine stops in

* There is a preparation (No. 83, Ser. 17) in the Museum of St. Bartholomew's Hospital quite to the point. The patient had profuse hæmorrhage, which commenced five hours after the operation. He died with peritonitis.

the inguinal canal, it is called an *internal* or incomplete inguinal hernia; such an one is generally of small size, and often difficult of detection. Lastly, if a portion of intestine escape at once through the external ring, then it is called a *direct* inguinal hernia.

Coverings of the Different Herniæ.—An oblique inguinal hernia has all the coverings of the cord. Beneath the skin, therefore, and the subcutaneous tissue, there will be

1. The spermatic fascia, derived from the aponeurosis of the external oblique.
2. The cremaster muscle, derived from the internal oblique and transversalis.
3. The infundibuliform fascia, derived from the fascia transversalis.

The incomplete inguinal hernia will be covered by

1. The aponeurosis of the external oblique.
2. The cremaster.
3. The infundibuliform fascia.

The direct inguinal hernia will be covered by

1. The spermatic fascia.
2. The fascia transversalis.

What becomes, it may be asked, of the fibres of the conjoined tendon of the internal oblique and transversalis? They are either protruded before the peritoneum or permit the hernia to slip between them.

In all cases, or at any rate with very few exceptions, the immediate investment of the intestine is the peritoneum. This constitutes the sac of the hernia. The opening of the sac, communicating with the abdomen, is called its mouth, then comes the neck, and lastly, the body or expanded part of the sac.

Position of the Spermatic Cord in Reference to the Hernia.—The spermatic cord is generally situated behind and on the outer side of the hernial sac. But instances sometimes occur in which there is a different arrangement, and such deviations are easily explained. Since the spermatic vessels and the hernial swelling are more or less connected by cellular tissue, and included, as it were, in a common sheath, we can have no difficulty in understanding that the hernia may separate the component parts of the cord, so that one or other of them

may come to lie on the front of the swelling. A similar displacement is sometimes produced by an old and large hydrocele on the same principle; the increase of the watery tumor affecting the spermatic vessels and the vas deferens in the same way as the growth of rupture. For this reason we ought always, in large herniæ, to be cautious, in cutting down upon the sac, that we do not divide any of the displaced components of the cord. Mr. Hey mentions, as a warning, that he once divided the vas deferens.*

Seat of Stricture.—The stricture may be seated either at the external ring, the internal ring, or at any intermediate part. Sometimes there is a sort of double stricture, one at the external, and the other at the internal ring.

The stricture, however, may be caused by the mouth of the sac itself, independently of the parts outside it; for the peritoneum may become thickened and indurated, and sufficiently unyielding to strangulate the protruded parts. Such changes are liable to be produced by the pressure of a truss.

Direction in which the Stricture should be Divided.—We cannot do better than adhere to the golden rule laid down by Sir A. Cooper, namely, to divide the stricture, in all cases, directly upwards. In this direction, we are least likely to wound the epigastric artery.

Changes Produced by an Old and Large Hernia.—Whoever has the opportunity of dissecting an old hernia of some size, will observe that the obliquity of the inguinal canal is destroyed. The constant dragging of the protruded viscera gradually brings the internal ring nearer to the external, so that at last the one gets quite behind the other, and there is a direct opening into the abdomen. But the position of the epigastric artery with regard to the sac remains unaltered. It is still on the inner side of the neck of the sac.

In herniæ of long standing, all its coverings undergo a change. They become thickened and hypertrophied to such an extent, and so altered from what they once were, that they scarcely look like the same parts.

[In the study of hernia we should bear in mind these several points: the coverings, the rings, the canal and its di-

* Practical Observations, p. 146.

rection, and the epigastric artery (deep). The external ring is situated one-fourth the distance from the symphysis of the pubes to the deep spine of the ilium; the internal ring midway between these two points. The canal is about an inch and a half in length, and passes obliquely upwards and outwards. The epigastric artery passes between the two rings, though nearer the internal ring. In oblique inguinal hernia we will find the following six coverings, passing from without inwards: integument; common superficial fascia; fascia propria of the external ring (spermatic fascia, intercolumnar fascia); cremaster muscle; fascia communis interna (infundibuliform fascia, fascia transversalis); peritoneum (fascia propria or true sac); the epigastric artery being internal to the neck of the sac. In direct hernia we have the same coverings, save the cremaster muscle—the artery then being external to the neck of the sac. The spermatic cord at the neck of the sac is usually outside. The hernia may pass under the cord. When speaking of the inguinal canal, we should have given its boundaries as follows: anteriorly, by the tendon of the external oblique; inferiorly, by Gimbernat's ligament; superiorly, by the conjoined fibres of the internal oblique and transversalis; posteriorly, by the fascia interna and peritoneum.]

Now expose the contents of the abdomen by a crucial incision, and take a general survey of the viscera before they are disturbed from their relative position.

What is Seen on Opening the Abdomen.—In the right hypocondrium the liver is seen projecting more or less below the cartilages of the ribs, and the fundus of the gall-bladder below the edge of the liver, near the end of the ninth costal cartilage. Such is generally the case in the dead body. But remember that in a state of health, the edge of the liver ought not to project much, if at all, below the false ribs: it does so, however, in children, and in women; in children, by reason of the large size of the liver; in women, by reason of the use of stays. In the left hypocondrium is seen more or less of the stomach. Across the umbilical region is a portion of the large intestine called the transverse colon; from this there descends a broad fold of peritoneum, called the great omentum, looking like a curtain of fat over the convolutions

of the small intestines; but the breadth of this fold varies in different instances; sometimes it is contracted and crumpled. The lower part of the belly and part of the pelvis is occupied by the small intestines. The urinary bladder is not apparent, unless distended sufficiently to rise out of the pelvis. In the right iliac fossa is the "caput coli," the commencement of the large intestine; but the ascending part of the large intestine in the right lumbar region, and the descending part of it in the left, are not visible unless distended; they lie contracted at the back of the abdomen.

Such and so much of the viscera are usually seen on opening the abdomen; but a certain latitude must be allowed to this statement. Sometimes more of one organ appears and less of another, according as this or that is distended or hypertrophied. Much depends also upon the amount of pressure which the ribs have undergone during life; so that, all things considered, we seldom see the parts in any two bodies precisely in the same position.

Particular Position of each Viscus.—You should next examine the position of each viscus separately; and, first, that of the liver.

The Liver.—The *liver* occupies the whole of the right hypochondrium, and extends over the epigastric region more or less into the left. Unless the individual be very corpulent, we can ascertain during life the extent to which the liver projects below the costal cartilages, and the general dimensions of the organ may be tolerably well told by percussion. Its under surface overlays part of the stomach, of the duodenum, of the right kidney, and of the transverse colon: its upper surface is convex, and accurately adapted to the arch of the diaphragm. To this muscle the liver is connected by folds of peritoneum, called "ligaments." One of these, nearly longitudinal in direction, and called the "*suspensory*," or, from its shape, the "*falciform*" ligament, is situated a little to the right of the mesial line. The free edge of it in front contains the imperious remains of the umbilical vein called the "*round*" ligament. The suspensory ligament, if traced backwards, leads to another broad fold extending horizontally from the diaphragm to the posterior border of the liver; this constitutes

the "*lateral*" ligament, right or left, according as we trace it on one or other side of the falciform ligament.*

Gall-Bladder.—The gall-bladder is the reservoir for the bile. It is closely confined by the peritoneum in a slight depression on the under surface of the liver. Its lower end or fundus projects beneath the cartilage of the ninth rib. This is important in a practical point of view: it sometimes happens that the gall-bladder, in consequence of some obstruction to its duct, becomes unusually distended, and, under such circumstances, occasions a swelling below the margin of the ribs, which might be mistaken for an hepatic abscess.† The close proximity of the gall-bladder to the duodenum and the transverse colon explains the occasional evacuation of gall-stones by ulceration into the intestinal canal.‡

Stomach.—The great end of the stomach is situated under the shelter of the left hypochondrium, and, when distended, occupies nearly the whole of it; but the narrow or pyloric end extends obliquely across the epigastrium into the right hypochondrium, where it is overlapped by the liver. The position of the great end of the stomach explains the peculiar sonorousness which percussion frequently elicits over the left hypochondrium, and even for some distance up the side of the chest; so that, when the stomach is large and flatulent, it is often very difficult to ascertain how much of the sound belongs to the stomach; how much to the chest. The pressure of the flatulent stomach upon the heart accounts also for the irregular action of this organ observable in cases of indigestion.

Spleen.—The spleen is situated deep in the left hypochondrium, between the stomach and the ninth, tenth, and eleventh ribs. Its outer surface is free and convex, to correspond with the diaphragm and the ribs; its inner surface, where its great vessels enter it, is concave and connected to the great end of the stomach by a peritoneal fold called the "*gastro-*

* The junction of the lateral and the falciform ligaments is described by some authors as the *coronary* ligament.

† See cases in point recorded by Andral, Clin. Méd. tom. iv.; and Graves. Dublin Hospital Report, vol. iv.

‡ See preparations in the Museum, Ser. 16, No. 84.

splenic omentum." Generally, too, the spleen is connected by a small peritoneal fold to the diaphragm.*

Pancreas.—This is the salivary gland of the abdomen. It lies behind the stomach, transversely across the spine about the level of the second lumbar vertebra. Its right end or head is contained within the curve of the duodenum: its left end extends as far as the spleen.

Kidneys.—The kidneys are situated in the lumbar region, nearly opposite the two lower dorsal and the two upper lumbar vertebræ. They lie embedded in fat, partly upon the quadratus lumborum, partly upon the psoas. In contact with the right kidney we have the liver, the second part of the duodenum, and the ascending colon; in contact with the left, are the spleen, the end of the pancreas, and the descending colon.

Renal Capsule.—This body is situated at the top of the kidney. It lies upon the crus of the diaphragm. You will see the right renal capsule by lifting up the liver; the left, by lifting up the spleen, and the great end of the stomach.

General Outline of the Course of the Intestines.—Now trace the course of the alimentary canal from the stomach to the anus, so far as this can be done without injury to its connections. A small part of it only is immovably fixed. Though all the rest of the canal can be moved about with facility, yet it is so connected by folds of peritoneum to the back of the abdomen that it cannot become entangled.

Duodenum.—The first part of the canal termed "*intestinum duodenum*," because it is about twelve inches long, takes a curious course (p. 318). Commencing at the pyloric end of the stomach, the duodenum ascends obliquely as high as the neck of the gall bladder; then, making a sudden bend, it descends in front of the right kidney; lastly, making another bend, it crosses the spine obliquely towards the left side of the second lumbar vertebra: here the canal takes the name of "*jejunum*." The commencement of this part of the intestine is seen by raising the transverse colon. Thus, the duodenum describes a sort of horse-shoe curve. The concavity of this curve is towards the left, and embraces the head or

* Every now and then we find in the gastro-splenic omentum one or more little spleens in addition to the large one.

larger end of the pancreas. For convenience of description, we divide the duodenum into an ascending, a descending, and a transverse portion. The first is completely covered by the peritoneum, and is comparatively loose, that the motions of the stomach may not be restricted; the second and third are only covered by peritoneum in front, and are immovably fixed, for good reasons, to the back of the abdomen.*

Jejunum and Ileum.—Pursuing its course from the left side of the second lumbar vertebra, the intestinal canal forms a number of convolutions, which are loosely connected to the spine by a broad peritoneal fold termed the "*mesentery*." Of these convolutions, the upper two-fifths constitute the "*intestinum jejunum*;" the lower three-fifths the "*intestinum ileum*." This is an arbitrary division; for there is no definite limit between them; but the character of the bowel gradually changes—that is, it becomes less vascular, has fewer folds of the lining membrane, and its coats are therefore less substantial to the feel.

Commencement of Large Intestine.—In the right iliac fossa, the small intestine opens into the left side of the colon; here the large intestine begins: here is the ilio-cæcal valve (fig. 59). Immediately below the junction we observe that the large intestine is expanded into a blind pouch, called the *cæcum* or *caput coli*. Into the back part of this pouch opens a little tube closed at the other end, called the "*appendix vermiformis*." This tube is generally three inches long, about as thick as a tobacco-pipe, and is either coiled up behind the *cæcum*, or connected to it by a peritoneal fold, so as to hang loose in the pelvis.

The commencement of the large intestine is generally confined by the peritoneum to the iliac fossa, in which it lies.†


* There are two reasons why the second and third parts of the duodenum should be fixed and have no mesentery: first, if the duodenum had been loose, it would have fallen down like the other intestines, and have disordered the stomach by dragging it down with it; secondly, had it been loose, it would have been apt to have stretched the bile and pancreatic ducts, and the flow of bile through them would every now and then have been obstructed.

† But this is not invariably so. The bowel is, in some subjects, connected to the fossa, by a fold of peritoneum or a "*meso-cæcum*." I have often seen this fold sufficiently loose to allow the *caput coli* to travel quite over to the left iliac fossa.

Tracing it from this point, we find that it ascends through the right lumbar region in front of the right kidney, and then, crossing the umbilical region towards the left side,* descends in front of the left kidney, down the left lumbar region into the iliac fossa, where it curves like the letter S.† These successive portions of the large intestine are termed respectively the ascending, transverse, descending, and sigmoid parts of its course. Lastly, the bowel enters the pelvis on the left side of the sacrum, and here takes the name of "*rectum*."

Fig. 59.

1. Ileum.
2. Cæcum or caput coli.
3. Appendix vermiformis.



SECTION THROUGH THE JUNCTION OF THE LARGE AND SMALL INTESTINE TO SHOW THE ILEO-CÆCAL VALVE.

This term, so far as concerns the human subject, is misapplied; the canal runs anything but a straight course through the pelvis, for it curves so as to adapt itself to the sacrum.

Looking at the entire course of the colon, observe that it

* This transverse part of the colon, in some instances, makes a coil behind the stomach to the diaphragm; such a state of things, when the bowel happens to be distended, is apt to give rise to symptoms of diseased heart. See some observations in point by Dr. Copland, in *Lond. Med. Gaz.*, 1847, vol. v. p. 660

† The contiguity of the ascending and descending colon to the right and left kidney respectively, explains the occasional bursting of renal abscesses into the intestinal canal.

forms an arch, of which the concavity embraces the convolutions of the small intestines.

Length of the Alimentary Canal.—The small intestines, including the duodenum, vary from sixteen to twenty-eight feet, according to the height of the individual. All this coil of bowel is sustained and kept in place by a simple pedicle of mesentery, which extends from the left side of the second lumbar vertebra, to the right iliac fossa.

In a man five feet high, I found that the small intestines measured twenty-four feet; the large, five feet nine inches. In round numbers we may say that the small and large intestines are from five to six times the length of the body.

Peritoneum.—A certain range of motion being necessary to the abdominal viscera, they are provided with a serous membrane, called the “peritoneum.” This membrane not only facilitates their movements, but maintains each in its proper place. Although its texture is thin, and apparently delicate, yet it is so tough, that if a portion of it were removed and stretched across a hoop, it would support a weight of from forty to fifty pounds without giving way. That it is elastic, too, is proved by the rapidity with which it recovers itself after being distended by dropsy, pregnancy, hernia, &c.

The plan of the peritoneum is like that of all other serous membranes; that is, it forms a closed sac, one part of which lines the containing cavity, the other is reflected over the contained organs. These are respectively termed the parietal and the visceral layers. There is nothing between them,—or, in other words, inside the sac, but just sufficient moisture to lubricate its smooth and polished surface. The viscera, then, are all outside the sac. Some lie altogether behind it, *e. g.* the pancreas and kidneys; others push the visceral layer before them, so as to give rise to membranous folds; and the greater the fold, the greater is the mobility of the viscus which occasions it. The vessels and nerves of the viscus must run up to it through the fold, for they can get to it in no other way.

First let us examine the several folds which connect the viscera either to each other or to the back of the abdomen. This done, we will endeavor to trace the peritoneum as a continuous membrane.

Mesentery.—This is the fold which connects the small intestines to the back of the abdomen. To see it you should raise the omentum and the transverse arch of the colon. You then observe that its attached part or root extends from the left side of the second lumbar vertebra across the spine to the right iliac fossa. The loose part of the mesentery curves, as it were, like a ruffle, and encloses the small intestine from the beginning of the jejunum to the end of the ileum. Between its layers, are the mesenteric vessels, nerves, glands, and lymphatics.

Transverse Meso-Colon.—This broad fold connects the transverse colon to the back of the abdomen. It forms a sort of partition, dividing the abdomen into an upper compartment, containing the stomach, liver, and spleen; and a lower, containing the convolutions of the small intestines. As regards the cæcum, the ascending and descending portions of the colon, they, as a general rule, are bound down by the peritoneum in their respective situations. It covers only two-thirds, or thereabouts, of their anterior surface; the rest is connected by loose cellular tissue to the back of the abdomen.*

Great Omentum.—This broad peritoneal fold proceeds from the lower border of the stomach, like a curtain over the convolutions of the small intestines. From this resemblance its name is probably derived. In corpulent persons it is enormously loaded with fat, and contributes very much to the size of the belly. In thin subjects, on the other hand, it is often quite transparent. Its length, too, varies considerably. In some bodies we find it extending down into the pelvis; in others, contracted and crumpled. Now and then it is found riddled with holes. This last state of parts is attended with some risk, for it may happen that a knuckle of intestine becomes strangulated in one of the holes; and of this, death is the almost certain result.

* It is practicable, therefore, to make an artificial anus by an incision, through the loins, into this portion of the canal. It should be remembered, however, that there are very frequent exceptions to the statement in the text. Indeed, it is by no means uncommon to find both the ascending and the descending colon completely surrounded by peritoneum, and connected in the lumbar regions, respectively, by a right and left lumbar meso-colon. There is also sometimes a meso-cæcum.

Gastro-Hepatic or Lesser Omentum.—This fold connects the stomach to the liver. Its right border is free; and between its layers are the vessels and nerves going to, and the duct coming from, the liver. The duct lies to the right, the hepatic artery to the left, and the vena portæ behind and between them. If the finger be introduced beneath this border, it passes through what is called the "*foramen of Winslow*"* into the lesser cavity of the peritoneum (p. 315).

Foramen of Winslow.—This foramen is situated behind the right edge of the lesser omentum. Very often it is blocked up by adhesions. But if you can get your finger in, you will find that the foramen is bounded above, by the liver; below, by the commencement of the duodenum; in front, by the free border of the lesser omentum; behind, by the vena cava inferior.

Gastro-Splenic Omentum.—This fold proceeds from the great end of the stomach to the spleen, and is continuous below with the great omentum. It contains between its layers the "*vasa brevia*," branches from the splenic artery to the great end of the stomach.

Course of the Peritoneum.—Now let us trace the peritoneum, as a continuous membrane. To say the truth, this is not a very easy matter,—in fact, it has always been considered a sort of anatomical puzzle. Since the peritoneum is a perfect sac, it matters not where we begin, for after all we must come back to the point from which we started. Supposing, then, a longitudinal section were made through the viscera in the middle line of the body, one might trace the peritoneum thus—beginning at the diaphragm, and taking, for brevity's sake, two layers at a time (fig. 60):—

From the diaphragm two layers of peritoneum proceed to the liver, forming its lateral ligaments; they separate to enclose the liver, meet again on its under surface, and pass on, under the name of gastro-hepatic omentum, to the smaller curve of the stomach. Separating, they embrace the stomach, and meeting again at its greater curve, pass down over the small intestines to form the great omentum. At the lower

* Fatal strangulation of a portion of intestine has been known to take place in this foramen.

margin of the great omentum they are reflected upwards (so that the omentum consists of four layers) to the spine near the pancreas, and here the layers diverge from each other. The upper layer ascends in front of the pancreas to the diaphragm; the lower layer proceeds over the arch of the colon and then back to the spine, thus forming the transverse meso-colon.* From thence it is reflected over the small intestines, forming the mesentery. From the root of the mesentery it descends into the pelvis, and invests a part of the rectum (and the uterus in the female). From the rectum it is reflected on to the posterior part of the bladder, and from thence to the wall of the abdomen, along which we trace it up to the diaphragm.

[We have usually found the layers of the omentum to embrace the transverse colon; and passing thence to the spine, forming the transverse meso-colon. Here the two layers separate, the one passing over the pancreas and aorta, to the diaphragm, the other downwards to invest the small intestines.]

Such would be the course of the peritoneum as seen by a longitudinal section. But there are certain lateral reflections which would not be seen except by a transverse section, *e. g.*

* This particular arrangement is demonstrable only in early life. As the child grows, the great omentum becomes adherent to the arch of the colon.

Fig. 60.




DIAGRAM OF THE COURSE OF THE PERITONEUM.

(The arrows show where the bladder can be tapped.)

from the great end of the stomach two layers proceed to the spleen, forming the gastro-splenic omentum : from the transverse meso-colon it is reflected on either side over the ascending and descending colon.

What parts of the Alimentary Canal are only partially covered by Peritoneum?—The descending and transverse portions of the duodenum, the cæcum, the ascending, and descending colon (with exceptional cases), and the lower part of the rectum are covered by the peritoneum only in front. These points are not without practical interest. For instance, since the descending colon is closely connected to the lumbar region by loose cellular tissue, and has not, in the majority of cases, any peritoneal covering behind ; this part of the alimentary canal is available for the purpose of making an artificial anus.* The gut can be reached by a longitudinal incision near the outer border of the quadratus lumborum. Again, the cæcum being very liable to the accumulation of hardened fæces, is apt to engender the formation of an abscess in the cellular tissue which connects it to the right iliac fossa. The matter thus formed may either burst externally, or make its way into the bowel.†

To give another instance. The absence of peritoneum from the posterior part of the cæcum explains how a hernia may take place without a peritoneal sac. That part of the cæcum bare of peritoneum may protrude through the inguinal ring first ; then, if more cæcum should come down, this is likely to be denuded of its peritoneal covering, which is but loosely connected to the gut at the sides. In this way, then, it is possible to have a large cæcal hernia without any peritoneum before it, except just round the neck. The same


* The proposal of opening the intestinal canal, as a last resource, at some point in the abdomen, so as to evacuate its contents and establish an artificial anus, was first made by Littre (*Mém. de l'Acad. des Sciences*, 1720). He proposed to open the sigmoid flexure of the colon in a case of imperforate anus. A successful case of the kind is recorded as having been performed on a boy twenty-four hours after birth by Duset, (*Recueil périod. de la Soc. de Méd. de Paris*, t. iv.) The same child was reported at the age of twelve to be in good health, with an artificial anus in the left iliac fossa. (*Dict. des Sciences Méd.* t. xxiv. See also in vol. xxxv. of *Méd.-Ch. Transactions*.)

† The best account of iliac abscesses is to be found in the "*Leçons Orales*" of Dupuytren.

thing may happen, *mutatis mutandis*, to the sigmoid flexor of the colon when it happens to be fixed in the left iliac fossa like the cæcum.

Lesser Cavity of the Peritoneum.—Returning once more to the peritoneum, anatomists sometimes speak of the lesser cavity of it, as distinguished from the greater, and they say

Fig. 61.



BRANCHES OF THE ABDOMINAL AORTA.

the communication between them takes place through the foramen of Winslow. Now this lesser cavity, or sac of the omentum, is situated behind the stomach. If, in the foetus, air be blown into the foramen of Winslow, we distend the lesser cavity; that is, the air passes behind the stomach and then between the two down and the two up layers, so to speak, of the great omentum. As the child grows, the great omentum becomes adherent to the arch of the colon; and thus the proper sac of the omentum is closed. But the cavity behind the stomach is permanent.

Branches of the Abdominal Aorta.—Our next subject should be the examination of the arteries which supply the viscera. The aorta enters the abdomen between the pillars of

the diaphragm, and then descending in front of the spine, divides opposite the fourth lumbar vertebra into the two common iliac arteries. In this course it gives off its branches in the following order (fig. 61):—

1. The *phrenic*, for the supply of the diaphragm.
2. The *cæliac axis*, a short thick trunk which immediately subdivides into three branches for the supply of the stomach, the liver, and the spleen.
3. The *superior mesenteric*, for the supply of all the small intestines and the upper half of the large.
4. 5. The *supra-renal* and the *renal* arteries.
6. The *spermatic*, for the testicles in the male, and the ovaries in the female.
7. The *inferior mesenteric*, for the supply of the lower half of the large intestine.
8. The *lumbar*, a series of branches, analogous to the intercostals, for the supply of the back part of the abdomen.

Now we shall trace these branches throughout in such order as is most convenient. Let us take the *cæliac axis* first. To dissect this artery and its branches the liver must be well raised, as in fig. 62, and a layer of peritoneum removed from the gastro-hepatic omentum. You will soon find that there is a close network of very tough tissue about all the visceral branches of the aorta. This tissue consists almost entirely of plexuses of nerves, derived from the sympathetic system. Of all these plexuses, the largest surrounds the *cæliac axis* like a ring. It is called the solar plexus,* or, more appropriately, the brain of the abdomen, for it contains a large quantity of ganglionic matter. From this, as from a root, other secondary plexuses branch off, and twine round the arteries; thus forming the phrenic plexus, the coronary, hepatic, splenic, superior mesenteric, renal, &c. It requires a very lean subject, and great anatomical dexterity to trace them.

Cæliac Axis and its Branches.—The *cæliac axis* arises from the front of the aorta, between the pillars of the diaphragm, just above the upper border of the pancreas. It is a very thick trunk which, after a course of about half an inch,

* This plexus is formed by the junction of the two semilunar ganglia described in the Dissection of the Thorax, p. 108.

divides into three branches,—the *coronaria ventriculi*, the *splenic*, and the *hepatic*.

[The *cæliac axis* may be partially covered at its origin by the diaphragm; it may be longer than usual, or, if wanting, its branches will come off separately from the aorta. Both phrenic arteries have been seen arising from this axis—the *hepatic* may arise from the superior mesenteric.]

PLAN OF THE BRANCHES OF THE CÆLIAC AXIS.

Cæliac axis.	{	Coronaria ventriculi, or gastric.	{	pancreaticæ parvæ.	{	
		Splenic.		gastro-epiploica sinistra.		
		Hepatic.		gastro-epiploica dextra.		
				pyloric.		
				cystic		
						pancreatico-duodenales.
						omental.

The *coronaria ventriculi*, the smallest of the three, ascends a little to the left towards the œsophageal end of the stomach, and then curves along the upper border of the stomach towards the pylorus, where it inosculates with the pyloric branch of the *hepatic* artery. It gives branches to the œsophagus as well as to the stomach.

The *hepatic* artery ascends between the layers of the lesser omentum to the transverse fissure of the liver, where it divides into branches for its several lobes. In its course to the liver, it lies to the left of the bile duct, and in front of the portal vein: all three are contained in the right border of the lesser omentum. The *hepatic* gives off:—

a. The *pyloric*, which runs along the upper curve of the stomach and inosculates with the *coronaria ventriculi*.

b. The *gastro-epiploica dextra*, which runs behind the duodenum, then along the great curve of the stomach, and inosculates directly with the *gastro-epiploica sinistra*, a branch of the *splenic*. It gives off:—1. Branches to the pancreas and duodenum (*pancreatico duodenales*). 2. Branches which descend to supply the great omentum.

c. The *cystic*, which supplies the coats of the gall-bladder.

The *splenic*, the largest of the three, proceeds tortuously along the upper border of the pancreas to the spleen, which it

enters by numerous branches. It gives off—1. Several small branches to the pancreas, *pancreaticæ parvæ*. One, rather

Fig. 62.




DIAGRAM OF THE BRANCHES OF THE CÆLIAC AXIS.
(Pancreas in dotted outline behind the stomach.)

- | | |
|--------------------------|-------------------------------|
| 1. Coronaria ventriculi. | 5. Gastro-epiploica dextra. |
| 2. Splenic a. | 6. Gastro-epiploica sinistra. |
| 3. Hepatic a. | 7. Vasa brevia. |
| 4. Pyloric a. | 8. Superior mesenteric a. |

larger than the rest, is called *pancreatica magna*. 2. The *gastro-epiploica sinistra*, which runs along the great curve of the stomach, and inosculates with the *gastro-epiploica dextra*. 3. *Vasa brevia*, which proceed between the layers of the gastro-splenic omentum, to the great end of the stomach.

Thus the stomach is supplied with blood by four channels, which by their inosculations form a main artery along its lesser curve, another along its greater; from these, numerous branches are furnished to both sides of the stomach. The artery of the greater curve also sends down numerous branches to the omentum, which form a beautiful network between its layers. The advantage of this free inosculation of arteries about the stomach must be apparent.

Vena Portæ; its Peculiarities.—The veins which return the blood from the abdominal portion of the alimentary canal, the pancreas, and the spleen, have this peculiarity, that they do not empty themselves into the vena cava, but all unite into one great vein, called the vena portæ, which ramifies throughout the liver, and secretes the bile. The trunk of the vena portæ itself is from three to four inches long; if you trace it downwards you find that it is formed behind the pancreas, by the confluence of the splenic and superior mesenteric veins (fig. 63). Traced upwards you would find that at the transverse fissure of the liver it divides into branches corresponding to the several lobes of the organ. The vena portæ, then, may be compared to the stem of a tree, of which the roots arise in the digestive organs, and the branches spread out in the liver. Eventually all the blood from the liver returns into the vena cava through the venæ cavæ hepaticæ.

The veins which empty themselves into the vena portæ are also peculiar, in that they have no valves. Therefore, if any obstruction arise in the venous circulation through the liver, the roots of the portal vein are apt to become congested; this is a common cause of hæmorrhoids, diarrhœa, hæmorrhage from the bowels, and ascites. Leeches applied to the anus have been long recognized as beneficial in congestion of the liver.




DIAGRAM OF THE VENA PORTÆ.

(The arrow is introduced behind the free border of the lesser omentum.)

Bile Duct.—The hepatic duct is soon joined by the cystic, or the duct from the gall-bladder. The common duct, *ductus communis choledochus*, thus formed, passes along the right edge of the lesser omentum, then behind the first portion of the duodenum, and opens obliquely into the back part of the second portion (p. 318). This common duct is from three to four inches long, and, if distended, would be about the size of a small writing quill.*

The great omentum, with the arch of the colon, must now be turned up over the chest, and the small intestines should be pushed towards the left side. Then, by removing a layer of peritoneum from the mesentery, we expose the mode in which this great artery ramifies so as to supply the small intestines. In making this dissection, the mesenteric glands immediately attract notice. They lie in great numbers between the layers of the mesentery, and vary considerably in size. The fine tubes, called lacteal vessels, which traverse the glands, are too thin and transparent to be seen under ordinary circumstances. But in cases where sudden death has taken place during digestion, they are found distended with chyle, and can be traced into the glands from all parts of the small intestines.† After traversing the glands, they all eventually empty their contents into the receptaculum chyli (p. 105).

Superior Mesenteric Artery and Branches.—This large artery descends beneath the pancreas and over the transverse part of the duodenum (p. 315), and then runs between the layers of the mesentery towards the right iliac fossa, where it terminates in branches for the supply of the cæcum. Thus its course describes a gentle curve from left to right. The left or convex side of the curve gives off a series of branches (from ten to sixteen) to the small intestines; while the concave side furnishes the right colic and the middle colic arteries for the supply of the large intestines (fig. 64).


Now we should trace the branches to the small intestines, in order to see the beautiful series of arches which they form

* That the gall-duct admits of being sometimes distended to a much larger size is proved by the passage of large gall-stones.

† The arrangement of the chyloferous vessels is extremely well displayed in the beautiful plates of Mascagni.

by their mutual inosculations. There are three or four tiers of them, each tier smaller than the preceding. The ultimate branches ramify in circles round the intestine. This circular arrangement of the vessels in the coats of the bowel is practically interesting, because it enables one in almost all cases to distinguish the intestine from the hernial sac.

Fig. 64.



PLAN OF THE MESENTERIC ARTERIES, AND THEIR COMMUNICATION.

The *colic* branches of the superior mesenteric are—the right colic, which proceeds towards the ascending colon, and the middle colic, which ascends between the layers of the meso-colon to the arch. They are arranged after the same plan as those of the small intestines; that is, they inosculate so as to form a series of arches which successively decrease in size, and finally terminate in circles round the bowel.

Remember that the superior mesenteric vein joins the splenic behind the pancreas, and forms the vena portæ (p. 319).

Inferior Mesenteric Artery and Branches.—To trace this artery, the small intestines must be drawn over towards the right side. It is given off about two inches above the bifurcation of the aorta. Descending into the pelvis over the left common iliac artery, it passes between the layers of the

meso-rectum, and, taking the name of superior hemorrhoidal, is finally distributed to the upper part of the rectum. Its branches are—1. the *colica sinistra*, which supplies the descending colon; 2. the *sigmoidea*, which supplies the sigmoid flexure. These inosculate in the form of arches. The *colica sinistra*, too, forms a large arterial arch with the *colica media*, so that there is a chain of arterial communications from one end to the other of the intestinal canal (fig. 64).

The inferior mesenteric vein joins the splenic behind the pancreas.

In order to complete our knowledge of the position of the duodenum and the pancreas, a ligature should now be placed on the upper end of the jejunum, another on the lower end of the sigmoid flexure of the colon, and the intermediate portions of the small and large intestine should be removed. By simply turning up the stomach we expose at once the horse-shoe course of the duodenum round the great end of the pancreas, and the several relations of this portion of the alimentary canal described (p. 304).

Pancreas.—This is the great salivary gland of the abdomen, and is situated immediately behind the stomach (p. 318). It is of a somewhat elongated form. The larger portion, or head, is embraced by the duodenum, and from this the gland extends across the spine to the spleen. Like all other salivary glands, it is most abundantly supplied with blood: it receives numerous branches from the splenic artery, which runs along its upper border; some from the superior mesenteric, which lies beneath it, and others from the gastro-epiploica dextra. Its duct runs through it from end to end, and opens into the back part of the descending portion of the duodenum, close to the opening of the bile duct. Sometimes the two ducts terminate by a common orifice; so that nature intends the fluids to mix. The use of the pancreatic fluid is to dissolve and form an emulsion with the fatty matters of the food.

[The pancreas measures six to eight inches in length, an inch and a half in breadth, and an inch in thickness. Its weight is from two to three ounces. It is found to be larger in the foetus and infant than in the adult. Its color varies with age, from a rosy hue to a pale yellowish color.]

The liver, stomach, duodenum, pancreas, and spleen, should now be collectively removed. For this purpose it is necessary to cut through the ligaments of the liver, the *venæ cavæ hepaticæ*, and the branches of the *cæliac axis*. These viscera, as well as the remainder of the intestinal canal, should be macerated in water, while you examine all that is to be seen at the back of the abdomen; namely, the deep-seated muscles, the aorta and vena cava, the kidneys, the lumbar plexus of nerves, and the sympathetic nerve.

Kidneys and Ureter.—The kidneys are placed in the lumbar region, one on each side of the spine. They lie embedded in more or less fat, on the *quadratus lumborum* and the *psoas* muscles, pretty nearly opposite the two lower dorsal and the two upper lumbar vertebræ. On the top of each is a little body, like a cocked hat, called the renal capsule. We have no difficulty in tracing the ureter, or excretory duct of the kidney. It descends almost vertically on the *psoas* muscle, enters the pelvis over the division of the common iliac artery, and empties itself into the lower part of the bladder, after running obliquely through its coats.

[The kidneys correspond in position to the two last dorsal and the two first lumbar vertebræ; the right being a little lower than the left.]

Remember that in front of the right kidney is the liver, the ascending colon, and the vertical portion of the duodenum; in front of the left, the descending colon, part of the spleen and pancreas. This explains how it is that a renal abscess or a calculus is sometimes evacuated by stool.*

Diaphragm.—This is a partly muscular and partly tendinous arch, so constructed as to form a complete movable partition between the chest and the belly—a floor for the one, and a roof for the other. We cannot see the structure of the

* The proximity of the colon to the ureters gives us a probable explanation of the manner in which pins or other extraneous bodies sometimes find their way into the bladder, and become the nuclei of calculi. In confirmation of this, the following case is related by Velpeau. It occurred at the Hospital of La Pitié. A pin, the head of which was still found in the colon, in which it had produced considerable ulceration, had passed also into the ureter, so that a calculus, of which the pin formed the axis, projected partly within and partly without the canal of the ureter. (Velpeau, *Anat. Chir.* t. ii. p. 175.)

arch until its peritoneal lining is removed. We then observe that there is a broad tendon in the centre, and that muscular fibres converge to it from all sides (fig. 65). What then is the origin of the diaphragm? 1st, from the ensiform cartilage; 2dly, from the inside of the cartilages of the six lower ribs by as many digitations, which correspond with those of the transverse muscle of the abdomen; 3dly, in the interval between the last rib and the spine, they arise from two thin ten-

Fig. 65.




DIAGRAM OF THE DIAPHRAGM, THE OPENINGS IN IT, AND THE PHRENIC ARTERIES.

dinous arches,* thrown, the one over the quadratus lumborum, the other over the psoas muscle; lastly, from the front of the bodies of the lumbar vertebræ by two elongated bundles, called the *crura* of the diaphragm. Between them the aorta enters the abdomen. Observe that both crura have tendinous origins, but that the right crus is a little longer than the left; for it arises from the bodies of the three or four upper lumbar vertebræ and their intervening cartilages, whereas the left does not descend so low by one vertebra.

* These arches are commonly called respectively, the *ligamenta arcuata* "*internum*" and "*externum*." The first extends from the body of the first lumbar vertebra to its transverse process, arching over the psoas; the second extends from the same transverse process to the last rib, and arches over the quadratus lumborum; in point of fact they are nothing more than the upper borders of the sheaths of these muscles, specially thickened for the purpose of giving origin to the fibres of the diaphragm. Where else could the fibres take origin in this situation?

From these various origins, the fibres ascend at first nearly vertically, and then all arch inwards, and are *inserted* into the central tendon.

The *central tendon* is nearly the highest part of the diaphragm. It presents a beautiful glistening surface, owing to the various directions in which its fibres cross each other; and its shape may be rudely compared to a trefoil leaf. The chief point of interest about the tendon is, that, in consequence of its connections with the pericardium (p. 98), it is always maintained pretty nearly on the same level; so that it both supports the heart, and serves as a fixed point for the insertion of the muscular fibres of the diaphragm.

Openings in the Diaphragm.—There are three openings in the diaphragm for the transmission of the aorta, the œsophagus, and the vena cava respectively. The aortic opening is between the crura, close to the spine; it transmits, also, the vena azygos and the thoracic duct, both of which lie rather to the right of the aorta. Trace the crura upwards, and you will observe that the inner fibres of each cross each other in front of the aorta, somewhat like the letter X.* Just above the crossing, and a little towards the left side of it, is the œsophageal opening; this is entirely muscular, whereas the aortic is partly tendinous. The opening for the vena cava is situated in the central tendon, rather to the right of the middle line, and you observe that the vein is intimately connected to its magin, so that it may be kept permanently open. Lastly, there pass through the crus, on each side, the sympathetic, the greater and lesser splanchnic nerves.

Function of the Diaphragm.—The diaphragm is the great muscle concerned in inspiration. We may, therefore, truly say, with Haller, that it is “*musculus post cor nobilissimus*.” During inspiration, the muscular sides of the diaphragm contract and become less arched (as shown by the dotted line in fig. 66); the floor of the chest sinks in consequence, and more room is made for the expansion of the lungs. During expiration the diaphragm relaxes and is forced up to its former position by the pressure of the abdominal muscles on the vis-

* This decussation is not invariable. But the right crus always crosses more or less over the left, so that the crura are never strictly parallel.

cera. This alternate falling and rising of the diaphragm constitutes the mechanical part of breathing. But the diaphragm conduces to the performance of many other functions. Acting in concert with the abdominal muscles, it assists in the expulsion of the fæces and the urine, also in parturition and in vomiting: for in all these operations, we first take in a deep breath, that the diaphragm may be in a state of contraction, and so form a resisting surface against which the viscera may be compressed by the abdominal muscles. Moreover, by its rapid or spasmodic contractions it is one of the chief agents concerned in laughing, sneezing, coughing, hiccough.


The nerves of the diaphragm are the phrenic (p. 91), and the five or six lower intercostal nerves on each side. Its blood-vessels are the two phrenic from the aorta, some branches from the internal mammary (p. 84), and the lower intercostal.

Psoas Magnus.—This long muscle extends from the sides of the lumbar vertebræ into the thigh. In dissecting this muscle take care, 1st, of the sheath in which it is contained; 2dly, of the branches of the lumbar plexus, which emerge from it; and 3dly, of the sympathetic nerve, which runs down the spine close and nearly parallel to its inner border.

Respecting the sheath of the psoas, we have to notice its attachment to the sides of the vertebræ and the brim of the pelvis. It is this which determines the ordinary course of a psoas abscess, namely, beneath the crural arch into the upper part of the thigh; for it is a rare exception when the matter travels into the pelvis.

The psoas *arises* from the bodies and transverse processes

Fig. 66.



DIAPHRAGM FROM ITS UPPER SURFACE.

(The dotted lines show how much it descends when it contracts.)

of the lumbar vertebræ, and their intervening fibro-cartilages; but, observe, only from the projecting borders of the vertebræ, not from the central grooved part; here the fibres arise from a kind of tendinous arch thrown over the lumbar vessels. The muscle descends beneath the crural arch into the thigh, and is inserted by a strong tendon into the back part of the lesser trochanter of the femur.

As it passes under the crural arch, the tendon of the *psoas* lies immediately over the capsule of the hip-joint, and there is a large bursa between them to facilitate the play of the tendon. We ought to be aware that every now and then, even in young subjects, but more frequently in old ones, in consequence of wear and tear, this bursa communicates with the capsule of the hip-joint. The fact is interesting for many reasons: to mention one only, it explains how a *psoas* abscess sometimes makes its way into the hip-joint; a result almost always fatal.

Psoas Parvus.—Once in about eight or ten subjects we meet with a little muscle called the *psoas parvus*. It arises from the last dorsal and the first lumbar vertebra, and the intervening cartilage; thence descending in front of the great *psoas*, it soon ends in a long flat tendon, which spreads out, and is inserted into the brim of the pelvis.

Iliacus Internus and Iliac Fascia.—This muscle occupies the iliac fossa, and is covered by a fascia, which first claims attention. This *iliac fascia*, as it is called, is attached to the crest of the ilium, and indirectly to the brim of the pelvis through its connection with the sheath of the *psoas*. But its most important attachment is to the outer half of the crural arch; here it is directly continuous with the fascia transversalis (p. 300), so that together they present an effectual barrier to the escape of intestine beneath this part of the arch.*

To return to the muscle. It arises from the iliac fossa, from

* The iliac fossæ are very liable to be the seat of suppuration, and the course which the matter takes depends upon its position with regard to the iliac fascia. If the matter be seated in the loose cellular tissue between the peritoneum and the fascia, it usually advances just above the crest of the ilium, or else towards the groin through the inguinal canal; but, if seated beneath the fascia, the chances are that the matter will make its way under the crural arch towards the upper and outer part of the thigh.

the ilio-lumbar ligament,* also from the capsule of the hip-joint. The fibres pass beneath the crural arch, and are inserted into the tendon of the psoas. Thus the two muscles, so far as their action goes, may be considered as one.

The combined *action* of the psoas and iliacus is to assist in raising the body from the recumbent position, and to fix the pelvis steadily on the thigh: this supposes the fixed point to be at the trochanter minor. But supposing the fixed point to be at the spine, then the muscle can raise and rotate the femur outwards. It is this action which is so troublesome to counteract in fractures of the upper third of the femur.

Quadratus Lumborum and its Sheath.—This muscle extends from the crest of the ilium to the last rib, and is contained in a sheath formed for it by the aponeurotic origin of the transversalis (p. 292). The anterior layer of its sheath is attached to the roots of the *transverse processes* of the lumbar vertebræ; and the posterior layer to their summits. The muscle arises from about an inch and a half of the crest of the ilium and from the ilio-lumbar ligament: it ascends nearly perpendicularly, and is inserted into the last rib, and into the front of the transverse processes of the four upper lumbar vertebræ by as many tendinous slips. In addition to these we commonly find that a few fibres take origin from the transverse processes, and run up to the last rib, crossing the front of the other part of the muscle. The principal use of the muscle is to steady the spine; it also steadies the last rib, and enables it to serve as a fixed point for the action of the intercostal muscles and the diaphragm.†

By raising the quadratus, we observe the aponeurotic origin of the transversalis from the summits of the transverse processes: this constitutes the posterior part of its sheath, and separates the muscle from the great erector spinæ.

* This ligament extends from the transverse process of the last lumbar vertebra to the ilium.

† The respective attachments of the quadratus lumborum, the crossing of its fibres, and its mode of action, lead to the inference that it is a large intercostal muscle. It is worth remembering that the outer edge of the quadratus lumborum, in a well-grown adult, is about three inches from the spines of the lumbar vertebræ, and midway between the last rib and the crest of the ilium. It is just outside the edge of this muscle that we can cut down to open the large bowel without wounding the peritoneum.

Before you examine the course of the aorta and its great primary divisions, do not omit to notice that a chain of absorbent glands extends along the brim of the pelvis and the bodies of the lumbar vertebræ, following pretty nearly the course of the great blood-vessels. Generally speaking they are small, and only one here and there attracts observation. They transmit the lymphatics from the lower limbs, the abdominal wall, and the testicle; and all eventually lead to the *receptaculum chyli*, or the beginning of the thoracic duct (p. 105). This is usually found on the right of the aorta, close to the second lumbar vertebra.

Relations of the Abdominal Aorta.—The aorta enters the abdomen between the crura of the diaphragm about the last dorsal vertebra, and descends nearly perpendicularly in front of the spine as low as the fourth lumbar, where it divides, rather to the left of the mesial line, into the two common iliac arteries. The point of division cannot be specified with precision, because it varies in different subjects. But for all practical purposes it is sufficient to know that its division takes place about the level of the highest point of the crest of the ilium. The aorta is crossed in front by the pancreas, the duodenum, and the right renal vein.

Vena Cava Inferior.—The vena cava inferior is formed by the junction of the two common iliac veins. It ascends on the right of the aorta, close to the spine in the greater part of its course. As it approaches the diaphragm, the vena cava passes off a little to the right, to go through its tendinous opening in the diaphragm, and so reach the right side of the heart. Its relations, beginning from below, are :—1. the mesentery; 2. the left renal artery which runs behind it; 3. the duodenum and head of the pancreas; 4. the liver. It receives the lumbar veins, the right spermatic, (the left joins the left renal,) the renal, the capsular, the phrenic, and lastly the hepatic.

[Sometimes the common iliac veins do not unite to form the inferior cava till they reach the level of the renal veins. The hepatic veins may not open into it till after they have passed through the diaphragm. The inferior cava may be on the left side of the aorta, till it receives the left renal vein, and then crosses the artery and resumes its normal position; it

also has been seen on the left side of the aorta altogether. It has also been known to enter the vena azygos and thus did not enter the auricle. In this case the hepatic veins opened into the auricle.]

The branches of the aorta we have still to examine, arise from it in pairs, namely,—the phrenic, capsular, renal, spermatic, and lumbar. See diagram, p. 315.

Phrenic Arteries.—These arteries supply the diaphragm, and arise from the aorta as soon as it comes through the pillars. The right phrenic passes behind the vena cava, the left behind the œsophagus, and both ramify extensively on the corresponding sides of the muscle (p. 324). Their first branches, however, are sent to the renal capsules. Their ultimate ramifications extend to the circumference of the chest, and inosculate with the internal mammary and intercostal arteries. The phrenic *veins* terminate in the inferior vena cava.

The *capsular* arteries proceed to the renal capsules. The capsular veins terminate on the right side in the vena cava, on the left in the renal.

Renal Arteries and Veins.—The renal arteries come from the aorta at right angles, and run transversely to the kidneys. Both are covered by their corresponding veins. The right is necessarily longer than the left, and crosses beneath the vena cava. Each enters its kidney, not as one trunk, but by several branches, corresponding to the original lobes of the organ. The renal veins lie in front of the arteries, and join the vena cava at right angles. The left is longer than the right, and has to cross over the aorta; it receives the spermatic and capsular veins of its own side.

[The renal arteries may be two or more in number, they have been known to come from the common, and also from the internal iliac. One instance is reported where they arose from the aorta by a common trunk. In place of entering the hilus, they may penetrate the kidney at its superior or anterior portion.]

Spermatic Arteries and Veins.—The spermatic arteries arise from the front of the aorta a little below the renal. They descend along the psoas muscle, and then through the inguinal canal to the testicle. In the female, the correspond-

ing arteries proceed between the layers of the broad ligament to the ovaries. Each artery is accompanied by two very tortuous veins, which unite, and then empty themselves, on the right side, into the vena cava; on the left, into the renal vein.

[The spermatic arteries may arise by a common trunk. Two on the same side are often met with, one coming from the renal. Three even have been observed.]

Lumbar Arteries and Branches.—There are five of these arteries on either side: four arise from the aorta, the fifth comes from the *arteria sacra media*. They are strictly repetitions of the intercostal arteries on a small scale, so that “lumbar intercostals” would be an appropriate name for them. They proceed outwards over the bodies of the vertebræ towards the inter-vertebral foramina, and then, like the thoracic intercostals, divide into dorsal and abdominal branches.

The *dorsal* branches pass between the transverse processes to the muscles of the back, and are of a size proportionate to the large developement of these muscles. They also send arteries into the spinal canal.

The *abdominal* branches all run outwards behind the *quadratus lumborum*, except the last, which commonly runs in front. After supplying the *quadratus* and *psoas*, they are lost in the wall of the abdomen.*

The lumbar *veins* empty themselves into the vena cava.

Common Iliac Arteries and Veins.—The two common iliac arteries resulting from the bifurcation of the aorta opposite the fourth lumbar vertebra, diverge from each other at an acute angle, towards the sacro-iliac symphysis, and, after a course of about two inches, divide into the external and internal iliac. They lie close to the vertebræ, and each, at or near its point of division, is crossed by the ureter; the left one is also crossed by the colon. But the most important relation of these arteries, in a practical point of view, is their position with regard to their corresponding veins. In consequence of the vena cava lying on the right of the aorta, you see that the

* Just as the thoracic intercostals, by communicating with the internal mammary, form a vascular ring round the chest, so do the lumbar, by communicating with the epigastric, form a vascular ring, though a less perfect one, round the walls of the abdomen.

right common iliac artery crosses over both the common iliac veins. It is also very closely connected to them. For these reasons it is easier to pass a ligature round the left artery than the right; for though its vein lies on its inner side, it is sufficiently far off to be out of the way of danger.

With the parts now before you, consider what would be the easiest way of performing this operation. Several modes have been recommended. Upon the whole, the best authorities* agree that the artery is most accessible from behind. An incision should be made perpendicularly from the end of the last rib to the ilium; another transversely along the margin of this bone nearly to its spine. We then cut, layer after layer, through the abdominal muscles till the peritoneum is exposed; this is easily raised from the iliac fascia, and with it the ureter is raised too. The application of the ligature is, after all, the most delicate part of the operation. It ought to be placed, as near as possible, midway between the origin and the division of the artery, so that there may be room enough for the formation of a clot on either side.†

[Mott's incision was to commence above the external ring, passing in a curved direction half an inch above Poupart's ligament, then upwards a little above the spine of the ilium. The height at which these arteries arise is subject to variety, as also their place of bifurcation. Their length may vary from an inch and a half to three inches. The minimum length known was less than half an inch, and the maximum four and a half inches. Cruveilhier relates a case where the right was wanting, the internal and external iliacs of that side arising from the aorta. This artery was first tied by Dr. Wm. Gibson, of Philadelphia, in 1812; successfully by Mott in 1827.]

* Consult some observations in point by Sir P. Crampton, in *Med. Chir. Trans.* vol. xvi.

† It is important to be aware that the length of the common iliac artery is apt to vary in different persons. I have seen it from three-fourths of an inch to three and a half inches long. These varieties may arise either from a high division of the aorta, or a low division of the common iliac, or both. It is impossible to ascertain, beforehand, what will be its length in a given instance, for there is no necessary relation between its length and the height of the adult individual. It is often very short in men of tall stature, and *vice versâ*. Anatomists generally describe the left as rather longer than the right; but, from the examination of 100 bodies, I conclude that their average length is the same.

External Iliac Artery.—This artery passes along the brim of the pelvis, on the inner side of the psoas, and then running under the crural arch about midway between the spine of the ilium and the symphysis pubis, takes the name of femoral. The corresponding vein lies close to its inner side, and on a posterior plane. After the removal of the peritoneum, notice that the artery is not bare, but covered by a thin layer of fascia which binds it down to the psoas. There are only two other circumstances of practical interest respecting this artery ; 1st, a slender nerve, the genito-crural, runs close to its outer side ; 2dly, just before it leaves the pelvis it is crossed by the circumflexa ilii vein. The branches given off by this artery are :—

The *epigastric*, already described (p. 301).

The *circumflexa ilii*, which arises just above the crural arch, and runs towards the spine of the ilium in a sheath formed for it by the fascia iliaca.* In the dissection of the abdominal muscles (p. 292), we saw the continuation of it skirting the crest of the ilium between the internal oblique and the transversalis, and sending a branch upwards between these muscles for their supply. The main trunk much reduced in size, inosculates with the ilio-lumbar derived from the internal iliac.

How to Tie the External Iliac.—The easiest way of tying the external iliac is to make a curved incision at the lower part of the belly, beginning a little above the middle of the crural arch, and ending a little beyond the spine of the ilium. The strata of the abdominal muscles, with the fascia transversalis, should then be divided to the same extent ; after which, the peritoneum can be readily raised by the finger from the iliac fossa. It is necessary to make a small incision through the sheath of the vessel, in order to facilitate the passage of the needle. Remember that the vein is closely connected to its inner side,† and that the genito-crural nerve is not far off.

[The external iliac was first tied by Abernethy in 1796.]

* The course of this artery should be borne in mind in opening iliac abscesses.

† This relative position of the vessels must not be always taken for granted. In old subjects, less frequently in adults, it is sometimes found that the external iliac artery runs very tortuously, instead of nearly straight, along the brim of the pelvis. But the vein does not follow the artery in its windings, and

Sympathetic Nerve.—Respecting the general plan upon which the sympathetic nerve is arranged, refer to what has already been said of it in the dissection of the neck (p. 81). Our present business is with the lumbar portion of it.

The abdominal part of the sympathetic nerve descends, on either side, in front of the bodies of the lumbar vertebræ, along the inner border of the psoas. The nerve has a ganglion opposite each lumbar vertebra, so that there are five on each side. Each of these ganglia receives two branches from the corresponding spinal nerve, just as in the chest; and, on the other hand, gives off filaments, of which some twine round the aorta, and accompany the inferior mesenteric and spermatic arteries to the large intestine and the testicle; but the greater number terminate in the hypogastric plexus.

Hypogastric Plexus.—The hypogastric plexus is situated between the common iliac arteries on the last lumbar and first sacral vertebra. It consists of an inextricable interlacement of nerves, partly sympathetic and partly spinal, and is a sort of nerve-centre for the supply of the pelvic viscera, like the solar plexus (round the cæliac axis) is for the abdominal. The minute filaments proceeding from it accompany the visceral branches of the internal iliac artery, and supply the bladder, prostate gland, rectum, and, in the female, the uterus and vagina. Thus we have the vesical, hæmorrhoidal, uterine, and vaginal plexuses. Of these, however, none are seen in an ordinary dissection.

Lumbar Plexus of Nerves.—This plexus is formed by


may possibly lie outside the artery just where we propose to place the ligature.

The mode of performing the operation described in the text is recommended by Sir A. Cooper. Mr. Abernethy, however, who first set the example of tying this artery, in 1796, adopted a somewhat different proceeding. He says: "I first made an incision about three inches in length through the integuments of the abdomen, in the direction of the artery, and thus laid bare the aponeurosis of the external oblique muscle, which I next divided from its connection with Poupart's ligament, in the direction of the external wound, for the extent of about two inches. The margins of the internal oblique and transversalis muscles being thus exposed, I introduced my finger beneath them for the protection of the peritoneum, and then divided them. Next, with my hand, I pushed the peritoneum and its contents upwards and inwards, and took hold of the artery."

the union of the four upper lumbar nerves (fig. 67). It lies over the transverse processes of the corresponding vertebræ, imbedded in the substance of the psoas, so that this muscle must be scraped away before it can be seen. Like the brachial plexus, the nerves composing it successively increase in size from above. Its branches are five in number, and generally arise in the following order :—

The *first lumbar* nerve is, both in course and distribution, analogous to an intercostal nerve, and may therefore be appropriately called the *lumbar intercostal*.* It crosses obliquely over the quadratus lumborum to the crest of the ilium, and

Fig. 67.



PLAN OF THE LUMBAR PLEXUS AND BRANCHES.

then, entering the abdominal wall, proceeds round to the front and terminates in the skin of the hypogastrium. Like the other intercostals, it gives off a lateral cutaneous branch; this is larger than the others, and runs over the crest of the ilium to supply the skin of the buttock.

* It often happens that the first lumbar nerve divides into two branches: of these, the upper is generally termed the ilio-hypogastric; the lower, the ilio-inguinal. But both run nearly a similar course, and have a similar distribution.

The *external cutaneous nerve of the thigh* (c) is generally derived from the second lumbar nerve. It runs through the psoas, then, crossing obliquely the iliacus towards the spine of the ilium, passes beneath the crural arch, and is finally distributed to the skin on the outside of the thigh.*

The *genito-crural nerve* is of small size, and generally comes from the second lumbar. After perforating the psoas, it descends along the outer side of the external iliac artery, and near the crural arch divides into the *genital branch* (g), which runs through the inguinal canal, and is lost in the cremaster muscle,—and the *crural* (e), which proceeds under the crural arch, and is lost in the skin of the upper part of the thigh.

The *anterior crural* (d) is the largest and most important branch of the plexus, and is generally formed by the union of the third and fourth lumbar nerves. It descends in a groove between the psoas and the iliacus, supplies both these muscles, and then, passing under the crural arch, is finally distributed to the extensor muscles and the skin of the thigh.

The *obturator nerve* (f) is next in size to the anterior crural. It proceeds from the third and fourth lumbar nerves, descends by the side of the pelvis to the obturator foramen, and is distributed to the adductor muscles of the thigh.

[We also have an obturator accessory nerve which is of small size, and comes either from the obturator nerve near its origin, or from the third and fourth lumbar. It passes along the inner border of the psoas muscle, and is distributed to the pectineus and hip-joint.]

Postponing the minute anatomy of the abdominal viscera, let us begin the examination of the contents of the pelvis.

DISSECTION OF THE PELVIS.

FIRST refresh your memory of the purposes served by the pelvis, and, what is of more practical importance, of the direction of its axis.

* If the external cutaneous be not seen in its usual situation, look for it as a distinct branch of the anterior crural, nearer the psoas muscle.

The purposes of the pelvis are to protect its own viscera, and to support those of the abdomen; to give attachment to the muscles which steady the trunk; to transmit the weight of the trunk to the lower limbs, and to give origin to the muscles which move them. In adaptation to these functions, the form of the pelvis is that of a perfect arch, with broadly expanded wings at the sides, and projections in appropriate situations to increase the leverage of the muscles. The sacrum, impacted between the ilia, represents the key-stone of the arch, and is capable of supporting not only the trunk, but great burdens besides. The sides or pillars are represented by the ilia; these transmit the weight to the heads of the thigh bones, and are thickest and strongest just in that line, *i. e.* the brim of the pelvis, along which the weight is transmitted. Moreover, to effect the direct transmission of the weight, the plane of the arch is oblique. This obliquity of the pelvis, its hollow expanded sides, its great width, the position and strength of the tuberosities of the ischia, are so many proofs that man is adapted to the erect posture.

The general conformation of the pelvis in the female is modified, so as to be better adapted to utero-gestation and parturition. Its breadth and capacity are greater than in the male. Its depth is less. The alæ of the iliac bones are more expanded. The projection of the sacrum is less perceptible, and consequently the brim is more circular. Above all, the span of the pubic arch is wider. The bones, too, are thinner, and the muscular impressions less strongly marked.

The cavity of the pelvis being curved, the axis, or a central line drawn through it, must be curved in proportion. For all practical purposes, it is sufficient to remember that the axis of the pelvis corresponds with a line drawn from the anus to the umbilicus.*

Contents of the Male Pelvis.—The male pelvis contains the last part of the intestinal canal, named the “rectum,” the

* In a well-formed female the base of the sacrum is about four inches higher than the upper part of the symphysis pubis, and the point of the coccyx is rather more than half an inch higher than the lower part of the symphysis.

The obliquity of the pelvis is greatest in early life. In the foetus, and in young children, its capacity is small; and the viscera, which subsequently belong to it, are situated in the abdomen.

bladder with the prostate gland at the neck, and the vesiculæ seminales. If the bladder be empty, there will also be some of the small intestines in the pelvis ; not so if the bladder be distended.

Course of the Rectum.—The rectum enters the pelvis on the left side of the sacrum, and, after describing a curve corresponding with the concavity of the sacrum, terminates at the anus. Observe, that, in the first part of its course, it is loosely connected to the back of the pelvis by a peritoneal fold, called the "*meso-rectum*;" and that between the layers of this fold the terminal branches (superior hæmorrhoidal) of the inferior mesenteric vessels with nerves and absorbents run to the bowel.

It is worth remembering that the rectum does not take this course in all cases ; sometimes it makes one, or even two lateral curves. In some rare cases it enters the pelvis on the right side instead of the left. Since these variations from the usual arrangement cannot be ascertained during life, they should make us cautious in the introduction of bougies ; for such things have happened as a perforation of the intestine by a bougie.*

Recto-Vesical Pouch.—Whilst the parts are still undisturbed, you should introduce the finger into what is called the *recto-vesical peritoneal pouch* (fig. 68). This is a cul-de-sac formed by the peritoneum in passing from the front of the rectum to the lower back part of the bladder. Now, in the adult male subject, the bottom of this pouch is about one inch distant from the base of the prostate gland :† therefore a part of the under surface of the bladder is not covered by the peritoneum ; and since this part is in immediate contact with the rectum, it is practicable to tap the distended bladder through the front of the bowel without injuring the peritoneum. The operation has, of late years, been revived by surgeons of good repute, and with great success.‡ It is easily

* In old age the rectum has sometimes a zigzag appearance immediately above the anus. These lateral inclinations are probably produced by the enormous distensions to which the bowel has been occasionally subjected.

† The bottom of the pouch is from three to four inches distant from the anus.

‡ See a paper in the *Med. Chir. Trans.* vol. xxxv. by Mr. Cock.

done, and not attended with much risk, provided all the parts be in their regular position. But this is not always the case. It sometimes happens that the peritoneal pouch comes down nearer to the prostate than usual—I have seen it in actual contact with the gland; so that in such a case it would be next to impossible to tap the bladder from the rectum without going through the peritoneum. In children the peritoneum comes down lower than it does in the adult, because the bladder in the child is not a pelvic viscus.

Fig. 68.




DIAGRAM OF THE RELATIVE POSITION OF THE PELVIC VISCERA.

The recto-vesical pouch is permanent. But there is another peritoneal pouch on the front part of the bladder, which is only produced when the bladder is distended. Let us, therefore, blow up the bladder through one of the ureters. It soon fills the pelvis, and then, rising into the abdomen, occasions the pouch in question between it and the abdominal wall. At first, the pouch is shallow, but it gradually deepens as the bladder rises. Now, supposing that the bladder be distended half way up to the umbilicus, which is commonly the case when it has to be tapped, we should find that the bottom of the pouch would be about two inches from the symphysis pubis (fig. 68). Within this distance from the symphysis the bladder may be tapped in the linea alba, without risk of

wounding the peritoneum. Thus, the surgeon has the choice of two situations in which he may tap the bladder,—above the pubes, or from the rectum. Which of the two be the more appropriate must be decided by the circumstances of the case.

DISSECTION OF THE MALE PERINEUM.

EXAMINE first the osseous and ligamentous boundaries of the lower aperture of the pelvis. Looking at a male pelvis (with the ligaments preserved), we observe that this aperture is of a lozenge shape ; that it is bounded in front by the symphysis of the pubes, laterally by the rami of the pubes and ischium ; behind by the coccyx and the great sacro-ischiatic ligaments.

This space, for convenience of description, is divided into two by an imaginary line drawn from one tuber ischii to the other. The anterior forms a nearly equilateral triangle, of which the sides are from three to three and a half inches long ; and since it transmits the urethra, we call it the urethral division of the perineum. The posterior, containing the anus, is called the anal division (fig. 69, 1, 2).*




DIAGRAM OF THE FRAME-
WORK OF THE PERINEUM.

[In twenty-three subjects, Dupuytren found the base of the urethral portion of the perineum of very variable extent ; in some cases not exceeding two inches, and in others, as much as three and a half.

Velpeau has seen the tuberosities of the ischia only one and a half inch apart. The base seldom exceeds, however, two and three-quarters inches.]

The subject is supposed to be placed in the usual position for lithotomy, with a full-sized staff in the bladder, and the

* It is well to bear in mind that the dimensions of the lower outlet of the pelvis are apt to vary in different subjects, and the lithotomist must modify his incisions accordingly.

rectum artificially distended. Observe that a central ridge, named the *raphé*, extends from the anus along the perineum, the scrotum, and the under surface of the penis. In the lateral operation of lithotomy the incision should commence an inch and a quarter above the anus, close to the left side of the *raphé*: it should be carried downwards and outwards for three inches towards the tuber ischii. In the bilateral operation the incision is semilunar, the horns being made on either side between the tuber ischii and the anus, equidistant from these points respectively; while the centre of the incision runs about three-quarters of an inch above the anus.

Anal Glands.—At the anus the skin becomes finer and more delicate, forming a gradual transition towards mucous membrane; and during life it is drawn into wrinkles by the permanent contraction of the cutaneous sphincter. Moreover, the skin at the margin of the anus is richly provided with minute glands,* which secrete an unctuous substance to facilitate the passage of the *fæces*. When this secretion becomes defective or vitiated, the anal cutaneous folds are apt to become excoriated, chapped, or fissured; and then defæcation becomes exceedingly painful.

Subcutaneous Tissue—Fat in Ischio-Rectal Fossæ.—It matters little how we reflect the skin. A convenient way is to make an incision down the *raphé*, around the anus, to the coccyx. Two others may then be made on each side at right angles to the first, the one at the upper, and the other at the lower end of it. Thus the skin of the perineum may be turned back like folding doors. In reflecting the skin we have to notice the characters of the subcutaneous structure.† Now its characters alter in adaptation to the exigencies of each part. On the scrotum the fat constituent of the tissue is entirely and for obvious reasons

* These glands are the analogues of the anal glands in some animals, *e. g.* the dog and the beaver. They are found not only about the anus, but also in the subcutaneous tissue of the perineum, a fact for the demonstration of which we are indebted to Professor Quekett. They are large enough to be seen with the naked eye.

† The probable thickness of this subcutaneous tissue is a point which ought to be determined by the lithotomist in making his first incision. Its great thickness in some cases explains the depth to which the surgeon has to cut in letting out matter from the ischio rectal fossa.

absent, while the fibro-cellular element is most abundant, and during life elastic and contractile. But, as we recede from the scrotum and approach the anus, the fat accumulates more and more, and on either side of the rectum it is found in the shape of large masses, filling up what would otherwise be two deep hollows in this situation, namely the ischio-rectal fossæ. The purpose of this accumulation of fat on each side of the anus, is to permit the easy distention and contraction of the lower end of the bowel during and after the passage of the *fæces*.* Again, over the tuberosity of the ischia we meet with large masses of fat, separated by tough fibrous septa passing from the skin to the bone, so as to make an elastic padding to sit upon. Occasionally, too, there are one or more large *bursæ* interposed between this padding and the bone.

So much respecting the general characters of this subcutaneous tissue of the perineum. Some anatomists describe it as consisting of three, four, or even more layers, but in nature we do not find it so. We may, indeed, divide it into as many layers as we please, according to our skill in dissection; but this only complicates what is, in itself, simple.

Now examine the cutaneous sphincter muscle of the anus, and the superficial vessels and nerves of the perineum.

Cutaneous Sphincter Ani.—The cutaneous sphincter of the anus arises from the point of the coccyx, and from the ano-coccygeal ligament. The muscular fibres surround the anus, and are lost in a pointed manner in the tendinous centre of the perineum (p. 344). It is called the cutaneous sphincter to distinguish it from a deeper and more powerful band of muscular fibres which surround the last inch or more of the rectum, and is situated nearer to the mucous membrane.

Cutaneous Vessels and Nerves.—The cutaneous vessels and nerves of the perineum come from the internal pudic artery and nerve, and chiefly from that branch of it called the *superficialis perinei*. This we shall trace more fully presently.

Besides this, a cutaneous nerve is sent to these parts from the lesser ischiatic, called the *long pudendal nerve* (fig. 70). It

* It is this fat in the ischio-rectal fossæ which renders them so liable to the occurrence of peculiarly fœtid gangrenous abscesses. These should be opened as early as possible, lest they burst into the rectum; and one sees how deep the knife must be introduced in order to reach the seat of the mischief.

comes through the muscular fascia of the thigh a little above the tuber ischii, and ascends, dividing into filaments, which supply the skin of the perineum and scrotum.

As we remove the fat from the ischio-rectal fossa, we observe that a number of vessels and nerves run through it in a transverse direction from the ramus of the ischium towards the anus. These are the *external* or *inferior hæmorrhoidal*, and proceed from the pudic, which may be felt on the inner side of the ischium. They sometimes bleed freely in laying open a fistula.

Superficial or Muscular Fascia of Perineum.—When the subcutaneous structure is fairly cleared off, we come to the fascia which invests the muscles of the perineum, and is, in every sense, analogous to muscular fascia in other parts, for it not only covers the muscles collectively, but provides each with a distinct sheath. At the same time it is much less dense, though more elastic than the strong fascia,—say, for instance, of the limbs: there is, indeed, so much resemblance between it and the deeper strata of the subcutaneous tissue, that you will hardly recognise any distinct demarcation between them. Now what are its connections? 1, it is attached on either side to the anterior lip of the rami of the pubes and ischium: 2, traced forwards it is found to be directly continuous with the *tunica dartos* of the scrotum: 3, traced backwards, we find that, at the base of the urethral triangle, it is reflected beneath the transversus perinei muscle, and joins the “deep perineal fascia,” or “triangular ligament.” This connection explains why, when urine is effused into the perineum, it does not make its way into the ischio-rectal fossa.

[Dr. Gurdon Buck of this city has shown that this fascia forms a sheath for the penis in the way we shall describe, and forming a continuation of the suspensory ligament above. He thus describes the mode of dissection: The penis and scrotum are to be circumscribed by an incision at the distance of three fingers' breadth all around, and crossing the perineum at the anterior margin of the sphincter. The dissection of the skin and subjacent cellular and adipose tissue is to be made towards the penis, on the level of the fascia lata laterally, and of the perineal fascia posteriorly, and carefully continued to the body of the penis, as far as the corona glandis.

This will denude the penis and suspensory ligament of its loose movable investments; make an incision along the dorsum of the penis in the median line, passing through the suspensory ligament and reaching the corona, between the vessels and nerves on either side. The adhesion of the sheath along the dorsum is firm, and requires care to dissect it. The vessels and nerves being raised with it, shows the line of adhesion. The entire corpus cavernosum is enucleated, the muscles of the perineum being raised with the sheath. It is now seen that the suspensory ligament above, and the perineal fascia from below and laterally, form one continuous membrane with the sheath, enclosing the corpus cavernosum in its cavity, and embracing the corpus spongiosum urethra between two layers, one of which passes above, and the other below it. The excavated base of the glands adheres inseparably to the outer surface of the sheath, while by its inner surface it caps the summit of the corpus cavernosum. The adhesions are most firm at the extremity of the corpus cavernosum, along its dorsal surface, and at the insertions of the erector and accelerator muscles.

It is thickest around the corona, along the dorsal surface where it forms the suspensory ligament. The cavity formed by the sheath, and occupied by the corpus cavernosum is limited posteriorly by the triangular ligament. This fascia also sends off septa between the accelerators in the middle, and the erectors on either side. These three distinct sheaths are confounded anteriorly with the common sheath investing the corpus cavernosum. Vol. i., Trans. of the Amer. Med. Ass.]

Remove the fascia in order to see the muscles which cover the bulb of the urethra and the crura of the penis. The bulb of the urethra lies in the middle of the perineum, and is covered by a strong muscle, called the accelerator urinæ. The crura penis are attached, one to each side of the pubic arch, and are covered each by a muscle, called the erector penis. A narrow slip of muscle, called the transversus perinei, extends on either side from the tuber ischii to the *central tendinous point* of the perineum. This point is about equidistant from the anus and the bulb of the urethra, and serves for the fixation of muscular fibres from all quarters of the perineum.

Thus you observe that the muscles of the perineum describe on each side a triangle, of which the sides are formed by the accelerator urinæ and the crus penis respectively, and the base by the transversus perinei. Across this triangle run up from base to apex the superficial perineal vessels and nerves. Let us examine these before the muscles.

Superficial Perineal Vessels and Nerves.—The superficial perineal artery proceeds from the internal pudic as it runs up the inner side of the tuber ischii. Though we cannot see the main trunk, we can easily feel it by pressing the finger against the bone. The artery comes into view a little above the level of the anus, and then passes up the perineal triangle, distributing branches to all the muscles, and is finally lost on the scrotum. The only named branch is a muscular one, called the *transversalis perinei* (fig. 70). This is given off near the base of the triangle, and runs with the transversus perinei muscle towards the central point of the perineum. It is necessarily divided in the first incision in lithotomy, and deserves attention, because it is sometimes of considerable size.

The artery is accompanied by two veins, which are frequently dilated and tortuous, especially in diseased conditions of the scrotum.


The nerves are derived from the internal pudic, follow the course of their corresponding arteries, and give off similar branches. They not only supply the skin of the perineum and scrotum, but each of the perineal muscles.

Accelerator Urinæ.—This muscle embraces the bulb of the urethra. To simplify the understanding of it, let us consider it as arising from a fibrous median raphé beneath the bulb, and from the tendinous centre of the perineum. Starting from this origin, the fibres diverge and are inserted as follows:—The upper ones proceed on either side round the corpus cavernosum penis, like the branches of the letter V, and are fixed on its upper surface: the middle fibres completely embrace the bulb, like a ring, and meet in a tendon on the upper surface of the urethra; while the lower fibres are fixed to the deep perineal fascia (fig. 71). Thus, then, the entire muscle acts as a powerful compressor of the bulb, and expels the last drops of urine from this part of the urethra. By dividing the muscle along the middle line and turning back each

half, we can clearly make out its insertion as above described.

[The name of this muscle is partially a misnomer, for the act of micturition requires a patent condition, and consequently an entire relaxation of the muscles that surround the urethra. It therefore, as we have stated, can only come into play in voiding the last drops of urine that otherwise would remain in the urethra. We prefer the name of ejaculator semenis.]

Fig. 70.



MUSCLES, WITH SUPERFICIAL VESSELS AND NERVES, OF THE PERINEUM.

Erectores Penis.—These muscles are moulded, one upon each crus of the penis. Each muscle arises from the inner surface of the tuber ischii: the fibres ascend, completely covering the crus, and terminate on a strong aponeurosis, which

is gradually lost on the corpus cavernosum penis. The action of these muscles is to compress the root of the penis, and so to contribute in some way to the erection of the organ.

Fig. 71.




DIAGRAM TO SHOW THE ACCELERATOR URINÆ IN PROFILE.

Transversi Perinei.—These muscles are always of insignificant size, and sometimes absent. They arise one on each side from the tuber ischii, and proceed towards the central point of the perineum, where they are blended with the fibres of the accelerator urinæ. We cut through this muscle with its artery in lithotomy.

The next stage of the dissection consists in reflecting and removing the accelerator urinæ from the bulb of the urethra, and the erector penis with the crus penis from the rami of the pubes and ischium. Remove also the transversus perinei. In short, remove everything which obstructs the view of the triangular ligament.

Triangular Ligament of the Urethra.—Understand that the “triangular ligament of the urethra” and the “deep perineal fascia” are synonymous terms. The anatomy of this part has given rise to much useless and perplexing discussion for want of a correct definition of how much fascia is included in this ligament. Some anatomists make two layers of it, anterior and posterior, and say that the membranous part of the urethra is situated between them: others consider the ante-

rior layer only as *the* triangular ligament, and the posterior layer as part of the pelvic fascia, which I think is the proper view to take of it.

The triangular ligament as shown in fig. 72, is a strong and

Fig. 72.




DIAGRAM TO SHOW THE TRIANGULAR LIGAMENT OF THE URETHRA.

resisting membrane stretched across the pubic arch. What are its connections? On each side it is firmly attached to the *posterior* lip of the rami of the pubes and ischium; superiorly, *i. e.* towards the symphysis of the pubes, it is connected with the sub-pubic ligament; inferiorly, it does not present a free border, but is reflected forwards under the transversus perinei and becomes identified with the superficial perineal fascia (p. 343).

The membranous part of the urethra runs through the tri-

angular ligament about one inch below the symphysis pubis. The aperture for it does not present a distinct edge like a round hole, because the ligament is prolonged forwards over the bulb and serves to keep it in position.

Points of Surgical Interest.—The triangular ligament is very important in a surgical point of view, for these among other reasons :—

1. Here the young hand meets with difficulty in introducing a catheter, unless he can hit off the right track through the ligament. The soft and spongy tissue of the bulbous part of the urethra in front of the ligament readily gives way, if violence be used, and a false passage is the result.

2. By elongating the penis, you are much more likely to hit off the proper opening through the ligament.

3. When, in retention of urine, the urethra gives way *anterior* to this ligament, it is this which prevents the urine from traveling into the pelvis. Its connection with the superficial perineal fascia prevents the urine from getting into the ischio-rectal fossa. Neither can the urine make its way into the thighs. The only outlet for it is into the cellular tissue of the scrotum and penis.

4. When suppuration takes place *behind* the ligament, the matter is pent up and should be speedily let out; if not, it will find its way into the cellular tissue of the pelvis and may burst into the urethra or the rectum.

5. We have to cut through the ligament in lithotomy.

[The student should now practice the introduction of the catheter. Taking the instrument in the right hand, lightly held between the thumb and two first fingers, introduce it into the urethra (the penis being raised with the thumb and forefinger of the left hand, so as to efface the curve which the penis forms when it bends down in front of the scrotum), and pass it inwards until the point reaches the bulb; which is about an inch below the arch of the pubes. During this proceeding, the concavity of the catheter is directed towards the symphysis, while the straight portion is held parallel with the abdomen. When the point reaches the bulb, the handle should now be brought to a perpendicular position, so as to form a right angle with the axis of the body. This movement will raise the point of the catheter out of the sinus of the

bulb, after which it may safely be passed onwards through the opening in the triangular ligament into the membranous portion. Gradually depressing the handle now, a little more between the thighs of the patient, it will glide smoothly into the bladder.]

What Lies Behind the Triangular Ligament?—You must now carefully cut away the triangular ligament to see

Fig. 73.




DIAGRAM OF THE PARTS BEHIND THE TRIANGULAR LIGAMENT OF THE URETHRA.

(The anterior fibres of the levator ani are hooked down to show part of the prostate; the rest is tracked by a dotted line.)

what lies behind it. I say *cut* away, because you cannot remove it as a clean layer, owing to its close adhesion to the

parts behind it. Now these parts are shown in fig. 73, namely : 1, the membranous part of the urethra, surrounded by, 2, the compressor urethræ muscle ; 3, Cowper's glands ; 4, the pudic artery and its three terminal branches, *i. e.* the artery of the bulb, the artery of the crus, and the dorsal artery of the penis.

Fig. 74.




DIAGRAM OF THE RELATIONS OF THE COMPRESSOR URETHRÆ SEEN FROM ABOVE.

Compressor Urethræ.—This muscle surrounds and supports the urethra in its passage beneath the pubic arch. It arises from the ramus of the pubes on either side ; from thence its fibres pass, some above, some below the urethra, along the whole length of its membranous part. To obtain the best pe-

rineal view of the muscle, cut through the spongy part of the urethra about three inches above the end of the bulb, and dissect it from the corpus cavernosum penis. Thus, the upper fibres of the constrictor will be exposed; to see the lower, it is only necessary to raise the bulb. The most perfect view, however, of the muscle is obtained by making a transverse section through the rami of the pubes, so as to get at the muscle from above, as shown in fig. 74. We find that it forms a complete muscular covering for the urethra between the prostate and the bulb. It is chiefly through its agency that we are enabled to retain the urine. It is also this muscle which causes spasmodic stricture of the urethra.

[This muscle is sometimes spoken of under the name of Guthrie's or Wilson's muscle, they having described different portions of it.

The direction of the urethra should now be studied. Its course varies in different regions. The *spongy portion* is the most movable, the anterior portion hanging down or taking any position we may please to give it. As it reaches the pubis, we notice it to be more fixed by the suspensory ligament and the crura of the penis. The canal now curves under the symphysis, and pierces the triangular ligament from seven-eighths of an inch to an inch and a quarter below the arch of the pubis. When the body is erect the *membranous portion* ascends with a slight curve, which course is pursued through the prostate. At the termination of the prostatic portion the direction of the urethra is nearly vertically upwards. The lowest part of the canal is that which is in contact with the anterior layer of the deep fascia, when in the erect position.]

Cowper's Glands.—These glands are imbedded, one on either side, immediately behind the bulb, in the substance of the compressor urethræ.* Each consists of a cluster of little glands. Their collective size is about that of a pea, but it varies in different individuals. From each a slender duct runs

* The compressor urethræ was first accurately described and delineated by Santorini (septemdec. tabulæ), and afterwards by Müller in his monograph (Ueber die organ. Nerv. der männlich. Geschlechtsorgane), and by Guthrie in his work on the anatomy and diseases of the neck of the bladder. These authors agree in essentials respecting the muscle, though they differ in minor points.

forwards, and, after a course of about one inch, opens into the under surface of the spongy part of the urethra (p. 351). Their use is to furnish a secretion accessory to the function of generation.

Pudic Artery and its Branches.—The pudic artery is a branch of the internal iliac. It leaves the pelvis through the great ischiatic notch, crosses the spine of the ischium, re-enters the pelvis through the lesser ischiatic notch, and then runs along the inner side of the tuber ischii up towards the pubic arch. About an inch and a half above the tuber ischii, we can feel the trunk of the pudic artery; but we cannot see it, nor draw it out, for it is securely lodged in a fibrous canal. As we trace it upwards, we find that it runs behind the triangular ligament and divides into three principal branches; the artery of the bulb, the artery of the crus, and the dorsal artery of the penis (fig. 73). Its external hæmorrhoidal and superficial perineal branches have been already described (p. 343).

The *artery of the bulb* is of considerable size: it runs through the substance of the compressor urethræ, and before it enters the bulb divides into two or three branches. It also sends a little branch to Cowper's glands. From the direction of this artery it will at once strike the attention that there is great risk of dividing it in lithotomy. If the artery run along its usual level, and the incision be not made too high in the perineum, then indeed it is out of the way of harm. But, supposing the reverse, the artery cannot escape; and its size is such that it might occasion fatal hæmorrhage. If it be asked, how often does the artery run along a dangerous level? I answer, about once in twenty subjects; and there is no possibility of ascertaining this anomaly beforehand.

[The artery of the bulb may be small, wanting, or double. It may arise sooner than usual, cross the perineum farther back than natural, and reach the bulb from behind. It might then be divided in the operation for stone.]

The *artery of the crus penis* is given off after that of the bulb. It ascends for a short distance under cover of the arch, but soon enters the crus.

To see the *dorsal artery* of the penis, we must dissect the penis from its attachment to the symphysis pubis. The artery

should be traced down to the glans. We find that it forms a complete arterial circle round the corona glandis, giving off numerous ramifications to the papillæ on the surface.

[The dorsal artery may arise from an accessory pudic, and has been seen to arise from the deep femoral, passing obliquely upwards and inwards to reach the root of the penis.]

Pudic Nerve.—Respecting the pudic nerve little need be said, except that it corresponds both in its course and branches with the artery. It gives off its external hæmorrhoidal, and its superficial perineal branches,—a small one to the bulb, and another to the crus penis; but the main trunk of the nerve runs with the artery along the dorsum of the penis to the glans, and we cannot fail to be struck with its large size (p. 351). In its passage it supplies the integuments of the penis, and sends one or two branches into the corpus cavernosum. This part of the penis also receives nerves from the sympathetic system.

Ischio-Rectal Fossa.—This is the deep hollow on each side between the anus and the tuber ischii. When all the fat is removed from it, you observe that it is lined on all sides by fascia. Introduce the finger into it to form a correct idea of its extent and boundaries. Externally, it is bounded by the tuber ischii and obturator internus muscle; internally, by the rectum and levator ani; posteriorly, by the gluteus maximus; anteriorly, by the transversus perinei. The fossa is crossed by the external hæmorrhoidal vessels and nerves.

These deep recesses on each side of the rectum explain the great size which abscesses in this situation may attain. The matter can be felt only through the rectum. Nothing can be seen outside. Perhaps nothing more than a little hardness can be felt by the side of the anus. These abscesses should be opened early; else they form a large cavity and possibly may burst into the rectum.


ANATOMY OF THE SIDE VIEW OF THE PELVIC VISCERA.

To make a side view of the pelvic viscera, the left innominate bone should be removed in the following manner:—Detach the peritoneum and the levator ani from the left side of

the pelvis; saw through the pubes about two inches from the symphysis, and cut through the sacro-iliac symphysis; then, by drawing the legs apart, and cutting through the pyriformis, sacro-ischiatic ligaments, and ischiatic nerves, the innominate bone comes away easily. This done, the rectum should be distended with horse-hair, and the bladder blown up through the ureter. A staff should be passed through the urethra into the bladder.

You have already seen how the peritoneum passes from the front of the rectum to the lower part of the bladder (forming the recto-vesical pouch), and thence over the back of the bladder to the wall of the abdomen. You see where the distended

Fig. 75.



VERTICAL SECTION THROUGH THE PERINEUM AND PELVIC VISCERA.

bladder is bare of peritoneum, and that it can be tapped either through the rectum or above the pubes without injury to the serous membrane, as shown by the arrows in fig. 75.

False Ligaments of the Bladder.—The peritoneal connections of the bladder are called its false ligaments; *false* in contradistinction to the *true*, which are formed by the fascia of

the pelvis, and really *do* fasten the neck of the bladder in its proper position. The false ligaments are—1, the *posterior*—two peritoneal folds produced, one on either side the rectovesical pouch, by the fibrous remains of the umbilical arteries; 2, the *anterior*, produced by the passage of the peritoneum from the bladder to the abdominal wall; and, lastly, the *lateral*, produced by its reflexion from the side of the bladder to the side of the pelvis.

Pelvic Fascia.—The pelvic fascia constitutes the true ligaments of the pelvic viscera, supporting and maintaining them in their proper position. To expose it, you need only remove the peritoneum on that side of the pelvis which has not been disturbed. In doing so, notice the abundance of loose cellular tissue which is interposed between the peritoneum and the fascia, evidently for the purpose of allowing the bladder to distend with facility. Whenever urine gains access to this cellular tissue, it is sure to produce the most disastrous consequence; therefore, in all operations on the perineum, it is of the utmost importance not to injure the fascia in question.

Examine, first, to what parts of the pelvis the fascia is attached; secondly, the manner in which it is reflected on the viscera.

Beginning, then, in front, we find that the fascia is attached to the body of the pubes; thence we trace its attachment along the side of the pelvis, just above the obturator foramen, to the greater ischiatic notch; here it becomes gradually thinner, covers the pyriformis muscle, and is lost on the sacrum. Now let us follow it on to the viscera.

From the *pubes* the fascia is reflected over the prostate and the neck of the bladder, so as to form on either side of the symphysis a well-marked band, called the *anterior true ligament* of the bladder. From the *side* of the pelvis it descends as low as the line of origin of the levator ani, and here divides into two layers, inner and outer; the inner is reflected on to the side of the bladder, forming what is called its *lateral true ligament*; the outer lines the external surface of the levator ani.

So far there can be no difficulty. But the most important part of the fascia still remains to be examined. The fact is, that it forms a complete capsule for the prostate gland and

neck of the bladder. Now, how is this capsule constructed? In front, the fascia takes its point of departure from the sides of the pubic arch: from thence it passes backwards, and is gradually lost round the prostate and the neck of the bladder (p. 348). Taking it as a whole, it forms a case, which encloses not only the prostate and neck of the bladder, but also the prostatic plexus of veins, and the beginning of the membranous part of the urethra. Towards the perineum, the case is closed by the triangular ligament. In the operation of lithotomy the greatest care should be taken not to injure the capsule alluded to, for reasons already assigned.

If we were to describe this capsule piecemeal, we should say that it consists of so many ligaments: its upper part would be formed by the anterior ligaments of the bladder, its sides by the lateral ligaments* of the prostate, and its lower part by a horizontal layer of fascia interposed between the prostate and the rectum.†

The use of this capsule is to fasten down the neck of the bladder and the prostate, and to maintain them in their proper position with regard to the pubic arch. But for this the prostate would rise with the distending bladder, elongate the urethra, and thereby materially obstruct the free passage of the urine.‡

General Position of the Pelvic Viscera in the Male.—The pelvic viscera are so surrounded by veins and loose cellular tissue, that he who dissects them for the first time will meet with considerably difficulty in discovering their definite boundaries. The rectum runs at the back of the pelvis. The bladder lies in front of the rectum, immediately behind the

* These ligaments extend from the ramus of the ischium to the side of the prostate, and are sometimes called "ischio-prostatic."


† Denonvilliers is the only anatomist who puts these several portions of the fascia together, and describes them as a whole. He says: "La prostate et la portion membraneuse de l'urètre sont placées au centre, comprises entre des plans fibreux supérieurs, inférieurs, et latéraux, enveloppées des toutes parts, et engainées à la manière des muscles. On conçoit donc comment la portion membraneuse de l'urètre se trouve contenue dans une sorte de caisse irrégulièrement quadrilatère."—*Propositions d'Anat. et de Phys.* Paris, 1837.

‡ In cases of extreme distension of the bladder, the urethra does in some rare instances become elongated to a very considerable extent, so that a very long catheter is required to reach the bladder. Consult M. Deschamps, *Traité de l'Opération de la Taille*, tom. i. p. 221.

symphysis pubis. At the neck of the bladder is the prostate gland through which the urethra passes. In the cellular tissue, between the bladder and the rectum, there is, on each side, a convoluted tube called the "vesicula seminalis;" and on the inner side of each vesicula, is the seminal duct or vas deferens. Before we describe these several parts in detail, it is necessary to say a few words about the large tortuous veins by which they are surrounded.

Plexus of Veins about Prostate and Neck of Bladder.—Beneath the pelvic fascia, about the prostate and the neck of

Fig. 76.



SIDE VIEW OF THE PELVIC VISCERA.

(Taken from a Photograph.)

- | | |
|---|----------------------------------|
| 1. External sphincter. | 7. Vesicula seminalis. |
| 2. Internal sphincter. | 8. Ureter. |
| 3. Levator ani cut through. | 9. Vas deferens. |
| 4. Accelerator urinæ. | 10. Crus penis divided. |
| 5. Membranous part of the urethra
surrounded by compressor muscle. | 11. Triangular ligament. |
| 6. Prostate gland. | 12. Superficial perineal fascia. |
| | 13. Rectum. |

the bladder, are a number of large and tortuous veins, which form, by their mutual communications, the prostatic and the vesical plexuses. They empty themselves into the internal iliac. In early life they are not much developed, but as puberty approaches they gradually increase in size; and any one not familiar with the anatomy of these parts would hardly credit the volume which they sometimes attain in old persons. They communicate freely with the inferior hæmorrhoidal plexus, or veins about the anus, and they receive the blood returning from the penis through the large veins which pass under the pubic arch. This is one of the reasons why the prostatic plexus is so capacious.

If, in lithotomy, the incision be carried beyond the limits of the prostate, the great veins around it must necessarily be divided: these, independently of any artery, are quite sufficient to occasion very serious hæmorrhage.

Rectum Intestinum and its Relations.—The rectum enters the pelvis on the left of the sacrum, and describes a curve corresponding to the axis of the pelvis. Just before its termination, the bowel curves downwards, ^{but} so that the anal aperture is dependent. The rectum is not throughout of equal calibre. Its capacity becomes greater as it descends into the pelvis; and immediately above the sphincter, it presents a considerable dilatation (fig. 75, p. 355). This dilatation is not material in early life, but it increases as age advances. An adequate idea of it cannot be formed unless the bowel be fully distended. Under such circumstances the rectum loses altogether its cylindrical form, and bulges up on either side of the prostate and the base of the bladder. For this reason it is essential that the rectum be always emptied before the operation of lithotomy. *looks back*

The upper part of the rectum is connected to the sacrum by a fold of the peritoneum termed "*meso-rectum*." In this fold, the terminal branch of the inferior mesenteric artery with its vein runs up to supply the bowel. Below the meso-rectum, the gut is connected to the sacrum and coccyx by an abundance of loose cellular tissue, which allows its easy distension. This explains why fistulous passages are so apt to burrow in this direction.

The relations of the front part of the rectum,—that, namely,

1. "You will notice in your morning walks that the faeces are deposited, not on the sidewalk, but up against the wall."

included between the recto-vesical pouch and the anus—are most important. To put this matter in the most practical light, we will suppose the forefinger to be introduced into the anus, and a catheter in the urethra. The first thing felt through the front wall of the bowel is the membranous part of the urethra (fig. 76). It lies just within the sphincter, and is about ten lines in front of the gut. About one and a half or two inches from the anus the finger comes upon the prostate gland; this is in close contact with the gut, and is readily felt on account of its hardness; by moving the finger from side to side we recognize its lateral lobes. Still higher up, the finger goes beyond the prostate, and reaches the “trigone” of the bladder: the facility with which this can be examined depends not only upon the length of the finger and the amount of fat in the perineum, but upon the degree of distension of the bladder; for the more distended the bladder, the better can the prostate be felt. These several relations are practically important. They explain why, with the finger in the rectum, we can ascertain whether the catheter is taking the right direction,—whether the prostate be enlarged or not. We might even raise a stone from the bottom of the bladder so as to bring it in contact with the forceps. The rectum is supplied with blood by the superior, middle, and inferior hæmorrhoidal arteries. The superior come from the inferior mesenteric (p. 321); the middle and inferior from the pudic artery. The corresponding veins join the inferior mesenteric, and consequently the portal system. They are very large and form a very tortuous plexus about the lower part of the rectum. As they have no valves, they are liable to become dilated and congested from various internal causes; hence the frequency of hæmorrhoidal affections.

Large is not so

Bladder.—This viscus, being a receptacle for the urine, must necessarily vary in size, and accordingly the nature of its connections and coats is such as to permit this variation. When contracted the bladder sinks into the pelvis behind the pubic arch, and is completely protected from injury. But, as it gradually distends, it rises out of the pelvis into the abdomen, and, in cases of extreme distention, may reach nearly up to the umbilicus.* Its outline can then be easily felt through

* When the bladder is completely paralysed it becomes like an inorganic

the walls of the abdomen. The form of the distended bladder is oval, and its long axis, if produced, would pass superiorly through the umbilicus, and inferiorly through the end of the coccyx. The axis of a child's bladder is more vertical than in the adult; for in children, the bladder is not a pelvic viscus. It is this which makes lithotomy so much more difficult in their case. An inexperienced operator is apt to forget the axis of the young bladder, and to get out of the groove of the staff, between the bladder and rectum; in short, not to cut into the bladder at all.*

The quantity of urine which the bladder will hold without much inconvenience varies. As a general rule it may be stated at about a pint. A good deal depends upon the habits of the individual; but some persons have, naturally, a very small bladder, and are obliged to empty it more frequently.

In young persons the lowest part of the bladder is the neck, or that part which joins the prostate. But as age advances, the bottom of the bladder gradually deepens so as to form a sort of well or pouch behind the prostate. In old subjects, more particularly if the prostate be enlarged, this pouch is apt to become of considerable depth, and render micturition tedious. It sometimes happens that a stone exists in the bladder, and yet none can be felt; the reason of which may be that the stone has lodged in such a pouch below the level of the neck of the bladder, and therefore escapes the detection of the sound. Under these circumstances it is necessary to place the patient on an inclined plane with the pelvis higher than the shoulders: then the stone falls out of the pouch, and is easily recognisable.


sac, and there seems to be no limit to its distension. Haller found in a drunkard the bladder so dilated that it would hold twenty pints of water. (Elem. Phys. art. Vesica.) Frank saw a bladder so distended as to resemble ascites, and evacuated from it twelve pounds of urine. (Oratio de Signis morborum, &c., &c. Ticini, 1788.)

W. Hunter, in his Anatomy of the Gravid Uterus, has given the representation of a bladder distended nearly as high as the ensiform cartilage.

* In all animals with a bladder, the younger the animal the more elongated is the bladder. This is indicative of its original derivation from a tube, *i. e.* the *urachus*. In the infant, the bladder is of a pyriform shape, as it is, permanently, in the quadruped; but as we assume more and more the perpendicular attitude, the weight of the urine gradually makes the lower part more capacious

Ureter.—This is a tube, about eighteen inches long, which conveys the urine from the kidney to the bladder. In the dissection of the abdomen we saw it descending along the psoas muscle, and crossing the division of the common iliac artery. Trace it down to the bottom of the bladder and you will find that it enters it about one inch and a half behind the prostate, and about two inches from its fellow of the opposite side (fig. 76). The tube perforates the bladder very obliquely, so that the aperture, being valvular, allows the urine to flow into, but not out of the organ. The narrowest part of the ureter is at the vesical orifice; here, therefore, a calculus is more likely to be detained than at any other part of the canal.

Fig. 76.



POSTERIOR VIEW OF THE BLADDER.

Vas Deferens.—This is the tube which conveys the seminal fluid from the testicle into the prostatic part of the urethra. It ascends at the back part of the spermatic cord through the inguinal canal into the abdomen; then, leaving the cord, it curves downwards over the side of the bladder, gradually approaching nearer to the middle line. Before it reaches the prostate it becomes very tortuous, and is joined by the duct of the seminal vesicle. The common duct thus formed (*ductus communis ejaculatorius*) terminates in the lower part of the prostatic portion of the urethra (fig. 68, p. 339). In point of size and hardness the vas deferens has very much the feel of whipcord.*

* The description in the text presumes the bladder to be distended. But

Vesiculæ Seminales.—These are situated, one on either side, between the bladder and the rectum (fig. 68, p. 339). Strictly speaking each is a tube, but so convoluted that it looks more like a little sacculated bladder. When rolled up, the tube is not more than two and a half inches long, but when unravelled, it is more than twice that length, and about the size of a small writing quill. Several blind tubes, or cæcal prolongations, proceed from the main one, after the manner of a stag's horn. The vesiculæ seminales do not run parallel, but they diverge from each other posteriorly like the branches of the letter V, and each lies immediately on the outer side of the vas deferens into which it leads. The function of the vesiculæ seminales is to serve as reservoirs for the semen.

If one of them be opened a brownish colored fluid escapes from it. This is presumed to be in some way accessory to the function of generation.*

[The function of the vesiculæ seminales has not yet been satisfactorily determined; for further information on the physiology and anatomy of these bodies, we would refer to the article on this subject in Todd's Cyclopædia of Anatomy and Physiology.]

Prostate Gland.—The prostate gland is situated at the neck of the bladder, and surrounds the first part of the urethra (p. 339). In the healthy adult it is about the size and shape of a chestnut. Its apex is directed forwards. It is surrounded by a plexus of veins (p. 358), and is maintained in its position by the pelvic fascia (p. 356). Its upper surface is about three-quarters of an inch below the symphysis pubis. The apex of the prostate is about one and a half inches from the anus; the base is about two and a half.

So much for the situation of the prostate. Now for its *relations*. Above it we have the anterior ligaments of the bladder

when the bladder is empty the vas deferens runs down upon the side of the pelvis. In this course it may be seen, through the peritoneum, crossing—1, the external iliac vessels; 2, the remains of the umbilical artery; 3, the obturator artery and nerve; 4, the ureter.

* The vesiculæ seminales are but imperfectly developed till the age of puberty. In a child of three years of age they can hardly be inflated with a blow-pipe.

with the dorsal vein of the penis between them; below, and in contact with it, is the rectum; on each side of it is the levator ani; in front of it we have the membranous part of the urethra (surrounded by its compressor muscle), and the triangular ligament; behind are the neck of the bladder and the vesiculæ seminales with the ejaculatory ducts.

The wide, that is the transverse diameter, is about one inch and a half; the vertical is about half an inch less. But the gland varies in size at different periods of life. In the child it is not developed, or at all events is very small: it gradually grows towards puberty, and increases in size with advancing age.

When you wish to ascertain the size and condition of the prostate during life, the bladder should be at least half full: because the prostate is then pressed down towards the rectum, and more within reach of the finger.

Anatomy of the Urethra in its Passage Under the Pubic Arch.—The urethra, in its passage under the arch of the pubes, is surrounded by muscular fibres; this, therefore, may very properly be called the "*muscular*" part of the urethra, though it commonly goes by the name of the "*membranous*." It comprises that part of the canal intermediate between the prostate and the bulb. It is about one inch in length, but somewhat longer on the upper than the lower surface, in consequence of the encroachment of the bulb. It is about an inch below the symphysis pubis, and nearly the same distance from the rectum; observe, however, that it is not equidistant from the rectum at all points, because of the downward bend which the gut makes towards the anus.*

Levator Ani.—This muscle supports the anus and the lower part of the rectum like a sling. To see it, the pelvic fascia must be reflected from its inner surface. It arises from the ramus of the pubes near the symphysis, from the spine of the ischium, and from a tendinous line extending in a gentle curve between these points of bone; this line being nothing more than the division of the pelvic fascia. From this long origin the fibres descend, and are inserted thus—the an-

* If a clean vertical section were made we should see that the two canals form the side of a triangular space, of which the apex is towards the prostate. This is sometimes called the recto-urethral triangle.


terior passing under the prostate, meet their fellow in the middle line of the perineum in *front* of the anus: the next are lost in the sphincter, while the posterior terminate on the side of the rectum (p. 356).

The function of the levatores ani is to retract the anus and the rectum after it has been protruded in defæcation by the combined action of the abdominal muscles and the diaphragm.

Coccygeus.—This muscle should be regarded as a continuation of the levator ani. It arises from the spine of the ischium, gradually spreads out, and is inserted into the side of the coccyx, which it serves to support.

Internal Iliac Artery and Branches.—From the division of the common iliac, the internal iliac descends into the pelvis, and, after a course of about an inch and a half, divides opposite the great sacro-ischiatic notch into two large branches (fig. 78.) Of these, the posterior gives off the ilio-lumbar, lateral sacral, and glutæal arteries; while the anterior gives

Fig. 78.



PLAN OF THE BRANCHES OF THE INTERNAL ILIAC.

off the obturator, vesical, ischiatic, and pudic: also the uterine and vaginal in the female. Such is their usual order: but these branches, though constant as to their general distribution, vary as to their origin.

[The length of the internal iliac has varied from half an inch to three inches. The lengths of the common and internal iliacs bear an inverse proportion to each other. It may

bifurcate between the upper margin of the sacrum and the upper border of the sacro-sciatic foramen. This artery was first tied by Dr. Stevens, of St. Croix, in 1812, with success, and by Dr. S. P. White of Hudson, New York, in 1827, with a like result. To reach this artery the incision should be in the course of a line extending from the umbilicus to midway between the spine of the pubis and the anterior superior spine of the ilium; it should commence at the outer edge of the rectus, and cease about an inch above Poupart's ligament, so as to avoid the cord.]

a. The *ilio-lumbar* is analogous to the lumbar branches of the aorta. It ascends outwards beneath the psoas, sends branches to this muscle and the quadratus lumborum, and then, running near the crest of the ilium, supplies the iliacus muscle, and finally inosculates with the circumflexa ilii (p. 333).

b. The *lateral sacral* descends perpendicularly in front of the sacral foramina, and near the coccyx inosculates with the middle sacral artery. It gives branches to the pyriformis, the bladder and rectum, and others which enter the sacral foramina for the supply of the cauda equina.

c. The *gluteal* is the largest branch. It passes immediately out of the pelvis through the great ischiatic notch above the pyriformis muscle, and then divides into branches for the supply of the great muscles of the buttock. These we dissect with the thigh.

d. The *obturator* artery runs along the side of the pelvis, below the corresponding nerve, to the upper part of the obturator foramen, through which it passes to be distributed to the muscles of the thigh. In the pelvis it gives off a small branch to the iliacus muscle, and another which ramifies on the back of the pubes.

But the obturator artery does not, in all subjects, take the course above stated. It may arise from the external iliac near the crural arch, or by a short trunk in common with the epigastric.* Under these circumstances, in order to reach the

* In most subjects a small branch of the obturator ascends behind the ramus of the pubes to inosculate with the epigastric. The variety in which the obturator arises in common with the epigastric is but an unusual develop-

obturator foramen, it generally descends on the *outer* side of the femoral ring. Instances, however, every now and then occur, where it makes a sweep round the *inner* side of the ring; so that three-fourths of the ring, or, what comes to the same thing, of the mouth of the femoral hernia, would in such a case be surrounded by a large artery.*

[Cloquet, in his *Descriptive Anatomy*, gives the following table as the result of 250 examinations :

Normal origin of the obturator artery from the inter-	
nal iliac,	160
From the epigastric on both sides,	56
From the epigastric on one side,	28
From the femoral artery,	6]

e. The *ischiatric* artery is smaller than the glutæal. It proceeds over the pyriformis and the sacral plexus, to the lower border of the great ischiatic notch, through which it passes out of the pelvis to the buttock, where it runs with the great ischiatic nerve.

f. The *pudic* artery supplies the perineum and the penis. It passes out of the pelvis through the great ischiatic notch, below the pyriformis, crosses the spine of the ischium, and re-enters the pelvis through the lesser notch. It then ascends on the inner side of the obturator internus muscle towards the pudic arch, where it gives branches to the several parts of the penis. In its passage over the obturator muscle it is enclosed in a strong tube of fascia, and is situated about one inch and a quarter above the tuberosity of the ischium. Here it might for a time be effectually compressed. The branches of the pudic artery were described in the dissection of the perineum (p. 353).

Unfortunately for operative surgery, the pudic artery sometimes of this branch. The branch derives additional interest from the fact, that after ligature of the external iliac it becomes greatly enlarged, and carries blood directly into the epigastric. See a case in *Med. Chir. Trans.* vol. xx. 1836.

* The Museum of St. Bartholomew's Hospital contains two examples of double femoral hernia in the male, with the obturator arising on each side from the epigastric. In three out of the four ruptures the obturator runs on the inner side of the mouth of the sac. (See preparations 55, 69, Series 17.)

times takes a very different course. Instead of passing out of the pelvis, it may run by the side of the prostate gland to its destination; or, which comes to nearly the same thing, one of the large branches of the pudic may take this unusual course, while the pudic itself is regular, but proportionably small. All practical anatomists are familiar with these varieties, and we seldom pass a winter session without meeting with one or two examples of them. It need hardly be said that lithotomy, under such conditions, might be followed by fatal hæmorrhage.

g. The *vesical** and middle *hæmorrhoidal* arteries are variable as to their number and origin. They come from the pudic or the ischiatic, and ramify on the side of the rectum, the bladder, the vesiculæ seminales, and the prostate.

h. The *middle sacral artery* is a very diminutive prolongation of the aorta down to the coccyx. It becomes gradually smaller as it descends, and finally inosculates with the lateral sacral artery. In animals this is the artery of the tail.

Respecting the *veins* in the pelvis they correspond with the arteries. The remarkable plexus of veins about the prostate, neck of the bladder, and rectum, has been described (p. 358).

Nerves of the Pelvis.—We should examine, first, those which proceed from the spinal cord; afterwards those derived from the sympathetic system.


Sacral Plexus.—Five† sacral nerves proceed from the spinal cord through the sacral foramina. The upper four, from their large size, at once attract observation; but the fifth, which perforates the coccygeus muscle, is small, and not easily found. The four upper nerves, together with a branch from the last lumbar (*lumbo-sacral*), form the sacral plexus. The great nerves of the plexus lie on the pyriformis muscle, beneath the branches of the internal iliac artery, and coalesce

* While on the subject of arteries supplying the bladder, it should be mentioned that the upper part of this organ is generally supplied by what is called the *superior vesical*. This is a small artery proceeding from the obliterated umbilical; and from it a still smaller is given off, which accompanies the vas deferens.

† Some anatomists describe a "coccygeal" nerve. It supplies the same parts as the fifth sacral.

to form the *great ischiatic* nerve, the chief branch of which passes out at the back of the pelvis for the supply of the flexor muscles of the inferior extremity. The other branches of the plexus are as follow—

Fig. 79.



PLAN OF THE SACRAL PLEXUS AND BRANCHES.

a. The *superior gluteal* nerve proceeds from the lumbo-sacral, leaves the pelvis with the gluteal artery, and supplies the glutæus medius and minimus, and the tensor fasciæ femoris.

b. The *lesser ischiatic* supplies the glutæus maximus, the skin of the buttock, the perineum, and back of the thigh.

c. The *pudic* nerve runs with the pudic artery, and, like it, supplies the rectum, perineum, and penis. The pudic commonly gives off the branch for the obturator internus, but this is often a distinct branch from the plexus. Under any circumstances the branch in question leaves the pelvis with the artery, and re-enters with it to reach the muscle.

d. The *branches of the pelvic viscera* are very small. They

proceed chiefly from the third and fourth sacral nerves, and form an intricate plexus about the bladder, prostate, and rectum.

e. Lastly, the sacral plexus supplies the levator ani, the coccygeus, the cutaneous sphincter of the anus, the pyriformis, gemelli and quadratus femoris.

Sympathetic Nerve.—From the lumbar region we trace the sympathetic nerve into the pelvis, along the inner side of the sacral foramina. In this part of its course its ganglia vary in number from three to five. The nerves of opposite sides unite in front of the coccyx : here we find the "*ganglion impar*."

The plan upon which the sympathetic nerves are distributed in the pelvis is similar to that in the abdomen. Each ganglion receives one or two filaments from a spinal nerve, and then gives off its branches to the viscera. The visceral branches are exceedingly delicate, and cannot be traced unless the parts have been previously hardened in spirits. They accompany the arteries supplying the respective organs, and are called the vesical, prostatic, and inferior hæmorrhoidal plexuses.

According to the accurate dissections of Muller, the vesical filaments of the sympathetic do not stop at the prostate, but pass on beneath the pubic arch into the corpus cavernosum penis. Thus the erectile tissue of the intromittent organ is brought directly within the influence of the sympathetic system.

STRUCTURE OF THE BLADDER, PROSTATE, URETHRA, AND PENIS.


The parts are presumed to have been collectively taken out from the pelvis, and the partial peritoneal covering of the bladder to have been removed.

Structure of the Bladder.—The bladder is composed of a partial peritoneal coat, a muscular and a mucous ; these are connected by an intermediate layer of cellular tissue, out of which some anatomists make a third coat, and call it the cellular.

The *muscular* is the outside coat. It consists of bundles of muscular fibres crossing each other in various directions.

Their color and development depend upon whether the individual has suffered from irritation of the bladder, or any ob-

Fig. 80.



BLADDER AND URETHRA LAID OPEN BY AN INCISION ALONG ITS UPPER SURFACE.

struction to the flow of urine. The general direction of the fibres is as follows:—An outer layer arises from the circumference of the prostate, and from thence its fibres spread out longitudinally over the bladder. Under this is a layer of circular fibres, more especially near the neck. Towards the sides of the bladder the two sets of fibres have a less definite arrangement, and form a kind of network: these, therefore, are the weakest parts of the bladder, and more liable to the formation of pouches.*

The bladder should be laid open by an incision along its front surface, to examine its interior. In a recently contracted bladder, the mucous membrane is disposed in irregular folds; these, however, disappear when the bladder is distended. Under ordinary circumstances the membrane is perfectly pale, but when inflamed it becomes of a bright red color. If a piece of the membrane, especially when selected from near the neck of the bladder, be placed under the field of a microscope, its surface is seen to be studded with follicles. These follicles secrete the thick ropy mucus in inflammation of the bladder.

The *orifice of the urethra* is situated at the lower and anterior part of the bladder, not at the most dependent part which forms the well behind the orifice, in which urine is so apt to accumulate in old persons. Though it appears small and contracted in the fresh bladder, yet if the little finger be introduced into it, we shall find that it will gradually dilate to a very considerable size. Immediately behind the orifice we observe in some bladders a slight elevation, called the "*uvula*." It is nothing more than a portion of the mucous membrane raised up by an accumulation of the submucous tissue, and is rarely of sufficient size to interfere with the passage of the urine. This is quite a different thing from enlargement of the third or middle lobe of the prostate.

* These pouches arise in the following manner:—A portion of mucous membrane is protruded through one of the muscular interstices so as to form a little sac. This is small at first, but gradually increases in size, because, having no muscular coat, it has no power of emptying itself; generally speaking, several such sacs are met with in the same bladder, and they often contain calculi. Now, if a calculus, originally loose in the bladder, happen to become lodged in a pouch by the side of it, a sudden remission of symptoms may ensue. Occurrences of this kind readily explain the boasted and apparent efficacy of medicine for dissolving a stone.

The *orifices of the ureters* are about an inch and a half behind the urethra, and about two inches apart. See how obliquely these tubes perforate the coats of the bladder, slanting towards each other, and standing out in relief under the mucous membrane.* Moreover, a little ridge proceeds from the orifice of each ureter down to the neck of the bladder, looking like a continuation of the ureter itself. If the mucous membrane be removed from these ridges, we find that they are produced by muscular fibres. Sir Charles Bell,† who first drew attention to them, believed them to be of use in regulating the orifices of the ureters, and named them “the muscles of the ureters.”

The ridges, converging from the ureters, together with an imaginary horizontal line drawn between their orifices, include a triangular area called, by the French anatomists, the “*trigone vesicale*.” The mucous membrane of this area is always firmly adherent to the subjacent tissue, and is therefore perfectly smooth. It is more richly provided with blood-vessels and nerves than the rest of the bladder, and is in consequence endowed with more delicate sensibility. This is the reason why a stone gives more pain when the bladder is empty, and why it is more painful in the erect than in any other position of the body.

The bladder is supplied with blood, by “superior,” “middle,” and “inferior” vesical arteries. The superior come from the umbilical: the middle from the internal iliac: the inferior from the pudic.

Prostate.—Having already examined the form, size, and relations of the prostate, p. 359, you have now only to make out its two lateral symmetrical lobes. There is, also, a third

* This slanting of the ureters serves all the uses of a valve. The urine enters the bladder, drop by drop, but cannot return, because the internal coat is pressed against the other side of the orifice, so as to stop it. When the bladder becomes thickened, in consequence of difficulty in passing the water, it sometimes happens that the ureters lose their valvular direction, so that the urine, when the bladder contracts, is partly forced back up the ureters; the result is, that they become dilated, and so does the pelvis of the kidney.

† Med. Chir. Trans. vol. iii. He says, “These muscles guard the orifices of the ureters by preserving the obliquity of the passage, and pulling down the extremities of the ureters according to the degree of the contraction of the bladder generally.”

or middle lobe.* It unites the lateral lobes, and is situated above the seminal ducts. In the healthy state of parts, this does not appear like a separate lobe; but when abnormally enlarged, it projects towards the cavity of the bladder, and acts like a bar at the mouth of the urethra.

Make a longitudinal incision through the prostate in order to expose the urethra. Notice that the canal does not run exactly in the centre of the gland, but rather nearer to its upper surface; nor is it of the same calibre throughout. It forms a sinus in the interior of the prostate, as though the gland had been purposely hollowed out; and this is what anatomists mean when they speak of the "sinus of the prostate." Along the floor of this sinus is a longitudinal ridge, which is broad and elevated behind, but gradually loses itself forwards in a narrow point. This is called the crest of the urethra, and the most prominent part of it is named the "*caput gallinaginis*," from its supposed resemblance to the head of a woodcock. The seminal ducts open close to each other, one on either side of this prominence (p. 371).

Immediately in front of the *caput gallinaginis*, precisely in the middle line, is a small opening, into which you should pass a probe to ascertain that it leads backwards into a little cul-de-sac or pouch in the substance of the prostate. This pouch is described as the analogue of the uterus, and called the "*utricle*;" but strictly speaking, it is the remains of the primordial sac out of which the parts were formed. It is of a pyriform shape, with the narrowest part at the orifice, and its length is about five or six lines. Practically it deserves attention because in some persons it is large enough to catch the end of a small catheter. Lastly, we have to notice the minute orifices of the proper ducts of the prostate, and these are best seen on the floor of the sinus.† Now the whole substance of

* Attention was first attracted to this middle lobe, in England, by Sir Everard Home, whose account of it is published in the Philos. Trans. for 1806. The preparation prepared by Sir Everard in illustration is preserved in the Museum of the Royal College of Surgeons in London, Physiol. Series, No. 2583 A. But the anatomy and effect of the enlargement of this part of the prostate gland is not a discovery of modern times. It was accurately described by Santorini in 1739, and subsequently by Camper, and is alluded to by Morgagni in the third book of his Epistles.

† In the ducts of the prostate we often find small calculi, of a brown color,

the gland is permeated by the divisions and subdivisions of these ducts. They are not visible to the naked eye, but if one of them were thoroughly traced out with the microscope, it would be seen to terminate in a blind sacculated extremity, upon which the capillaries ramify in rich profusion.*

Modern observations prove that the prostate is more of a muscular than a glandular body. Nearly two-thirds of it is made up of muscular fibre of the non-striped variety. The great mass of this muscular fibre is arranged in a circular manner round the urethra, so as to form a sphincter. The prostate is remarkable for its dilatability. This has been long known to lithotomists, distinguished for their skill and success. If a small incision be made through the anterior part of the gland, the *base being left entire*, the gland may be dilated by the finger to such an extent as to admit of the extraction of even large calculi.

It will naturally suggest itself, that any change in the dimensions of the prostate gland must affect the canal which runs through it and more or less obstruct the flow of urine. For instance, if the entire gland be uniformly enlarged, the length of the prostatic urethra will be increased; if the enlargement preponderate at one part more than another then the canal will deviate more or less from its natural track and assume a more angular or a lateral curve according to the part enlarged. When the middle lobe becomes enlarged, there arises, at the neck of the bladder, a tumor, which will, in proportion to its size, more or less obstruct the passage of the urine. In the efforts made to introduce a catheter into the bladder, it sometimes happens that the end of the instrument is pushed through this hypertrophied lobe.†

Vesiculæ Seminales.—Respecting the internal structure

consisting of phosphate of lime. Cases are sometimes met with, in which these calculi by degrees attain a considerable size, and distend the prostate into a kind of sac, which when examined by the rectum feels not unlike a bag of marbles.

* This was first demonstrated by Mr. Quekett. The same distinguished anatomist has also discovered that the secreting cells of the gland contain calculi of microscopic minuteness. He finds them, almost without exception, in the prostate at every period of life. For further detail concerning them consult the article "Prostate" in Todd's Cyclopædia.

† See the Museum of St. Bartholomew's, Prep. 8 and 21, Series xxix.

of the vesiculæ seminales, we need only observe that their mucous membrane is lined by a scaly epithelium, and that it presents a beautiful honeycomb structure, not unlike that of the gall-bladder; the purpose of this is to increase the extent of the secreting surface.

Cowper's Glands.—The glands of Cowper have been already examined *in situ* in the dissection of the perineum, p. 352. We find them close to the urethra, one on either side, immediately behind the bulb. They consist of an aggregation of smaller glands, of which the collective size is somewhat larger than a pea. Each pours its secretion by a single and very minute duct into the bulbous part of the urethra. The use of these glands appears to be like that of the vesiculæ seminales and the prostate, namely, to pour into the urethra a fluid accessory in some way, not yet understood, to the function of generation. They are found more or less developed in all mammalia, and in some, *e. g.* in the mole, they increase in size periodically with the testicle.

Urethra.—The urethra is the tube which leads from the bladder to the end of the penis, and serves not only as the excretory duct of the bladder, but transmits the secretion of the testicles and the several glands accessory to generation. It is surrounded by different structures in different parts of its course. The first inch, or thereabouts, is surrounded by the prostate gland (p. 374); the second inch, which passes under the pubic arch, is surrounded by the compressor urethræ (p. 364); the remainder of its course along the penis, is surrounded by erectile tissue, termed "corpus spongiosum." Hence anatomists divide it into the prostatic, the muscular or membranous, and the spongy. The length of the whole is about seven or eight inches, but, of course, this will vary according to the condition of the penis.

The general direction of the urethra, when the penis hangs flaccid, is like the letter S reversed; but if the penis be held straight, the canal forms only one curve through the pubic arch, with the concavity upwards. The degree of this curvature varies at different periods of life. In the child the bladder being more in the abdomen than in the pelvis, the curve forms part of a much smaller circle than it does in the adult; but it gradually widens as age increases, and our catheters

are shaped accordingly.* However, the parts, when in a sound state, will yield sufficiently to admit the introduction of a straight instrument into the bladder. Aston Key always used a straight staff in lithotomy.

In its contracted state, the sides of the urethra are in close apposition. When opportunity offers, make transverse sections of it, to see how it looks in the different parts of its course. Through the glans it is flattened vertically, its two sides being in contact; through the prostate, also, it is nearly flat, except at the lower part, where its sides are kept asunder by the verumontanum, which projects upwards, leaving a furrow on each side. But throughout the rest of its course the lining membrane is disposed in longitudinal folds, which project into and accurately close the canal, precisely on the same plan as that by which nature closes the œsophagus, p. 376. These longitudinal folds are plainly seen even when the urethra is slit open; indeed, they do not disappear unless the canal be forcibly stretched contrary to their direction.†

If the urethra be laid open from end to end, you will see that the canal is not of uniform calibre throughout. The external orifice is the narrowest, and the least dilatable part of the urethra; of which the object is, that the urine may be expelled in a jet. Therefore, any instrument which will enter the meatus ought to pass through into the bladder, if there be no stricture. The junction of the membranous with the bulbous part is almost as narrow. The centre of the prostatic and the bulbous part are the largest. In the centre of the glans penis, the canal widens into a little sinus, termed "*fossa navicularis*."

The most *dilatable* part of the urethra is the prostatic. Even the narrowest parts of the canal must admit of considerable dilatation, since calculi of from three to four lines in diameter have been known to pass through it.

* The sharper curve of the urethra in the child was well known to Camper. "In recenter natis, vesica basi sua elatius sita, pedetentum descendit, unde necessario sequitur curvaturum urethræ majorem esse in junioribus quam in adultis."—Demon. Anat. Pathol. lib. ii. p. 13.

† In a well-injected urethra we observe that the ridges of the folds possess very few blood-vessels, while the furrows between them are exceedingly vascular. For the demonstration of this fact the author is indebted to Mr. Quekett.

The seminal ducts open into the prostatic part of the urethra, by the side of the verumontanum. The ducts of Cowper's glands open into the bulbous part. Besides these glands, a number of ducts open into the urethra, proceeding from little glands situated in the submucous tissue. These ducts, called the *lacunæ*, are just large enough to admit a bristle, and they all run in the same direction as the stream of the urine. Most of them are on the lower surface of the urethra, but one, called "*lacuna magna*" is on the upper surface about an inch and a half down the canal.

The mucous membrane of the urethra is laid upon a substratum of areolar tissue. Then comes a very thin layer of elastic fibrous tissue, of which the fibres are arranged both in a longitudinal and a transverse direction—giving the parts that springiness of which we are sensible in introducing the catheter. Outside this is a layer of muscular fibre. It has been demonstrated beyond all dispute that the urethra is surrounded throughout its whole course by muscular fibre of the involuntary kind. Therefore, the whole of the canal having a muscular coat, similar to an intestine, any part of it is liable to a spasmodic contraction.

The urethra is lined by spheroidal epithelium, and near the glands is provided with papillæ: this, therefore, is the most sensitive part.

Lastly, the urethra is provided with a closely-set network of absorbent vessels,—a fact which has been demonstrated by the beautiful quicksilver injections of Panizza.* They run from behind, forwards, and join the absorbents of the glans penis. Eventually, their contents are transmitted down the great trunks on the dorsum penis to the inguinal glands. This satisfactorily explains the pathology of a bubo.

Anatomy of the Penis.—The skin of the penis is remarkably thin and extensible, and is connected to the body of the organ by an abundance of loose cellular tissue, in which fat is never found. At the end of the penis the skin forms an ample fold, called the prepuce, or foreskin, for the protection

* Osservazioni antropo-zootom. &c., Pavia, 1830. This anatomist has also displayed by injections an extremely fine network of absorbents which cover the glans penis. The interstices of this network are smaller than the diameter of the tubes.

of the glans :* and the thin fold which passes from the under surface of the glans to the prepuce is called "*frenum preputii*." If the glans be habitually covered by the prepuce, its surface partakes more of the appearance of mucous membrane than of common skin ; but if the reverse, the cuticular covering of the glans thickens and dries.

In a well-injected glans, we observe that the surface is covered by minute papillæ, just like the end of the finger, which are endowed with keen sensibility by the great dorsal nerves of the penis. Round its margin—" *corona glandis*"—are a number of minute sebaceous glands which secrete an unctuous matter called "*smegma preputii*." In cases of congenital phimosis, this is apt to collect in considerable quantity, and becoming rancid, irritates and excoriates the glans and prepuce, particularly in summer and in hot countries.

The chief bulk of the penis consists of erectile structure, named from the appearance of its interior, "*corpus cavernosum*." In a groove along its under surface runs the urethra, which is itself surrounded by a vascular spongy tissue called the *corpus spongiosum* ; an accumulation of this at the end of the organ forms the glans. These two structures, then,—the corpus cavernosum, and the corpus spongiosum,—together form the penis ; though they appear closely united, yet they are quite distinct from each other, as may be easily ascertained by making a transverse section, as shown, p. 380.

The *corpus cavernosum* constitutes more than two-thirds of the bulk of the penis. It commences posteriorly by two gradually tapering portions, called the *crura*, which are attached, one on each side, along a groove in the descending ramus of the pubes.† The crura converge, come into apposi-


* When the foreskin is, from the time of birth, so tight that the glans can not be uncovered, such a state is called a "congenital phimosis." This condition occasions no inconvenience in childhood, but is apt, after puberty, to become troublesome and painful, so that it is necessary to slit up the prepuce and set the glans at liberty. In persons who have a tight foreskin, it sometimes happens that, when the glans has been uncovered, the prepuce cannot be again drawn over it ; this is called a "paraphymosis." The neck of the glans becomes tightly girt, great distension and inflammation are the consequences, and very serious results may ensue unless the foreskin be reduced.

† While speaking of the attachments of the penis, it should be mentioned that its upper part is connected to the symphysis pubes by an elastic ligament called the "*ligamentum suspensorum penis*."

tion at the root of the penis, and then run on, side by side, to form the body of the organ. Anteriorly, each terminates in a rounded end which is received into a corresponding depression in the glans, which fits on to them like a cap.

If a longitudinal section be made through the corpus cavernosum, we observe that its interior is composed of a delicate reticular structure, surrounded by a thick fibrous coat. This

Fig. 81.



TRANSVERSE SECTION THROUGH THE PENIS.

coat forms a cylinder of adequate strength to protect and support the delicate structure within, and sufficiently elastic to allow the distension of the penis. Moreover, this coat forms a longitudinal partition, dividing the corpus cavernosum into two lateral halves. The partition, however, is only complete near the root of the penis; along the rest of the organ there are a number of gaps in it, so that it has somewhat the appearance of a comb; for this reason it has been called the "*septum pectiniforme*." Through this partition the blood-vessels on one side communicate freely with those on the other: therefore both sides of the penis become erect simultaneously.

And now let us examine the construction of the erectile, spongy-looking tissue inside. The interior of the cylinder is occupied by a number of delicate elastic thread-like septa, which intersect each other in all directions, so as to form a multitude of minute cells. These communicate most freely with each other, as one may readily ascertain by blowing air into the penis. They are not of equal size throughout the penis, for they are much smaller, and their component septa

altogether thicker at the root than they are towards the glans. All the cells communicate freely with the arteries. When the penis is flaccid, they are empty; but when they become distended with blood, the penis becomes erect.

Here two questions arise,—1. how does the blood get from the arteries into the cells? 2. how does it return from the cells?

With regard to the first question. If the arteries of the penis be well injected, we find that the artery of the corpus cavernosum enters the inner side of the crus, and proceeds forwards near the septum, distributing numerous ramifications. These are supported in the middle of the fibrous threads, and after making several spiral turns, as it were for the purpose of being able to accommodate themselves to the varying size of the penis, discharge their blood at once into the cells. Besides the ramifications which feed the cells, there is a capillary system for the nutrition of all the parts concerned.

With regard to the second question. If a colored fluid be injected into the corpus cavernosum, near the glans, it instantly fills all the cells, and returns partly through veins which pass out at the upper surface of the penis and join the great dorsal vein, partly through the deep veins of the penis which leave the inner side of each crus.

Corpus Spongiosum.—The corpus spongiosum is the erectile tissue which surrounds the urethra as it runs along the penis. It commences in the middle of the perineum, in a bulb like form, and at the end of the penis it is continuous with the glans. This is proved by the fact, that if we inject the spongy body, we fill the glans also; not so if we inject the cavernous body. Observe that the urethra does not pass precisely through the middle of the spongy body, but that it runs nearer to its upper surface. The bulb hangs more or less pendulous from the urethra, p. 339. In old persons it hangs lower down than in children, and is consequently more exposed to injury in lithotomy.

The corpus spongiosum has a much thinner external coat than the corpus cavernosum. Its interior is composed of a plexus of minute tortuous veins. This is plainly demonstrated by injecting the dorsal vein of the penis with wax. In this way we not only fill the spongy body, but also the glans and

the large veins which form the plexus round the corona glandis.*

The chief *nerves* of the penis are the pudic. The largest branches run along the dorsum to the outside of the glans : a few only enter the erectile tissue of the organ. This, it has already been mentioned (p. 370), is supplied by filaments of the sympathetic nerve proceeding from the hypogastric plexus.

The *absorbent* vessels proceeding from the glans and the integument of the penis join the inguinal glands. The absorbents of the glans communicate freely all round it : this explains why a venereal sore on one side sometimes affects the inguinal glands on the other.

DISSECTION OF THE FEMALE PERINEUM.

THE *pudenda* in the female consist of folds of the integument, called the labia. Between these is a longitudinal fissure which leads to the orifices of the urinary and genital canals.

Labia Majora.—The pubic region is generally covered by an accumulation of fat, called the *mons Veneris*. From this, two thick folds of skin descend, one on either side, constituting the “labia majora.” Their junction, about one inch above the anus, is called the commissure, or “frenulum labiorum :” it is generally torn in the first labor. The inner layer of the skin of the labium is thinner, softer, and more like mucous membrane than the outer : for this reason, whenever matter forms in the labium, the abscess is almost sure to burst on the inner side. Where the labia are in contact, they are provided with a number of small sebaceous glands, of which the minute ducts are observable on the surface. These glands sometimes inflame and secrete an acrid, purulent matter which creates great irritation and pruritus of the mucous surface of the vulva, often difficult to allay.

Labia Minora.—By separating the external labia, we ex-

* In the Museum of the Royal College of Surgeons there is a beautiful preparation in which the glans penis is injected with quicksilver, clearly showing it to consist of a plexus of veins.—Physiol. Series, No. 2588 A.

pose two smaller and thinner folds of skin, one on either side, termed "*labia minora*," or, by the old anatomists, the *nymphæ*. These folds converge anteriorly, and form a hood for the clitoris, called "*preputium clitoridis*;" posteriorly they are gradually lost on the inside of the *labia majora*. They never contain fat, like the *labia majora*, but are composed of a minute plexus of veins. Between the *nymphæ* and about the clitoris are a number of glands which secrete the same kind of unctuous matter as those about the *corona glandis* in the male.

In young persons, in whom the *nymphæ* are protected from the air and from friction by the external *labia*, they are soft, moist, and of a rose-red color: but when hypertrophied, so as to project beyond the *labia*, they become like the common integument, and acquire a bluish color from the distension of their veins.

Between the *labia minora*, and about a thumb's breadth below the clitoris, is the orifice of the urethra, or "*meatus urinaris*." Immediately below this is the vagina, of which the orifice is partially closed in the virgin state, by a thin fold of skin called the "*hymen*."

Clitoris.—The clitoris resembles in form and structure the penis, but on a very diminutive scale. Like the penis, it is attached to the sides of the pubic arch by two *crura*, fig. 82, each of which is grasped by its own little erector *clitoridis*. The *crura* unite to form the body of the organ, which is tipped by a small glans. The glans is provided with extremely sensitive *papillæ*, and covered by a little *prepuce*. Its dorsal arteries and nerves are exceedingly large in proportion to its size, and have precisely the same course and distribution as in the penis. Its internal structure consists of a plexus of blood-vessels, which freely communicate with those of the *labia minora*, for one cannot be injected without the other.

Urethra.—A smooth channel called the vestibule, three-quarters of an inch in length, leads from the clitoris down to the orifice of the urethra. This orifice is not a perpendicular fissure like that of the penis, but rounded and puckered, and during life it has a peculiar dimple-like feel, which assists us in finding it when we pass a catheter. You should practice the introduction of the catheter in the dead subject, for the

operation is not so easy as might at first be imagined, provided the parts are not exposed. The point of the fore-finger of the left hand should be placed at the entrance of the vagina; and then the catheter, guided by the finger, slips, after a little manœuvring, into the urethra. The canal is about one inch and a half in length, and runs along the upper wall of the vagina, p. 387; indeed the two canals adhere so closely that you can feel the urethra through the vagina like a thick cord. Strictly speaking the urethra is slightly curved with the concavity upwards; but for all practical purposes it may be considered straight. It is necessary, however, to be aware that its direction is not horizontal. In the unimpregnated state of the parts it runs nearly in the direction of the axis of the outlet of the pelvis; so that a probe pushed on in the course of the urethra would strike against the promontory of the sacrum. But after impregnation, when the uterus begins to rise out of the pelvis, the bladder is more or less raised also in consequence of their mutual connection; therefore the urethra, in the latter months of utero-gestation, acquires a much more perpendicular course.


The female urethra is provided with a "compressor" muscle, essentially similar, both in origin and arrangement, to that which surrounds the membranous part of the urethra in the male. It also passes through the triangular ligament, fig. 83; indeed, we cannot convey a better idea of it than by saying that it is just like the male would be if cut short immediately behind the bulb. Though the prostate gland is wanting, yet there are minute glands scattered all round it, especially near the neck of the bladder. At the same time it should be observed, that, in consequence of the wider span of the pubic arch, and the more yielding nature of the surrounding structures, the female urethra is much more dilatable than the male. By means of a sponge tent, it may be safely dilated so as to admit the easy passage of the fore-finger into the bladder. We avail ourselves of this great dilatability in the extraction of calculi.

The mucous coat of the urethra is arranged in longitudinal plicæ. Next to the mucous coat is a layer of elastic and non-stripped muscular fibres intermixed. Outside all is a plexus of veins bearing a strong resemblance to erectile tissue.

Vagina.—The vagina is the canal which leads up to the uterus; at present only the orifice of it can be seen. It is surrounded by a sphincter muscle, which is easily displayed by removing the integument. The muscle is about three-fourths of an inch broad, and connected with the cutaneous sphincter of the anus in such a manner that they together form something like the figure 8.

Bulb of the Vagina.—On each side of the orifice of the vagina, between the mucous membrane and the sphincter, there is a plexus of tortuous veins, termed the bulb of the vagina, from its analogy to the bulb of the urethra in the male. This vaginal bulb extends across the middle line be-

Fig. 82.



BULB OF THE VAGINA.

tween the meatus urinarius and the clitoris, as shown in fig. 82, which I took from an injected preparation in the Musée Orfila, at Paris.

[The vagina is a curved canal; the anterior surface measures about four inches, and the posterior from five to six. It is also flattened from above downwards, so that the opposite surfaces are in contact. Its superior extremity is round, where it is attached to the uterus. The external surface is the narrowest portion of the canal, the middle the largest, and the upper portion intermediate between the two.]

Hymen.—The hymen is a thin fold of skin which, in the virgin extends across the lower part of the entrance of the vagina, about half an inch behind the fourchette. In most instances its form is crescent-shaped, with the concavity upwards. But there are several varieties of hymen; sometimes there are two tegumentary folds, one on either side, so as to make the entrance of the vagina a mere vertical fissure;* or

* Such an one may be seen in the Museum of the College, Phys. Series, No. 2843.

there may be a septum perforated by several openings (*Hymen cribriformis*), or by one only (*Hymen circularis*). Again, there may be no opening at all in it, and then it is called *Hymen imperforatus*. Under this last condition no inconvenience arises till puberty; the menstrual discharge must then necessarily accumulate in the vagina: indeed the uterus itself may become distended, and to such a degree as even to simulate pregnancy.*

The presence of the hymen is not necessarily a proof of virginity, nor does its absence imply the loss of it. Cases are related by writers on midwifery in which a division of the hymen was requisite to facilitate parturition. In Meckel's Museum, at Halle, are preserved the external organs of a female in whom the hymen is perfect even after the birth of a seven months' child.

Cowper's or Duverney's Glands.—At the lower part of the orifice of the vagina is, imbedded in the loose tissue on either side, a gland † about the size of a small pea. Each has a long slender duct, which runs forwards and opens on the inner side of the nympha. In cases of virulent gonorrhœa these glands are apt to become diseased, and give rise to the formation of an abscess in the labium, which is very difficult to heal.

The description of the perineal branches of the pudic vessels and nerves, given in the dissection of the male perineum, applies *mutatis mutandis*, to the female, excepting that they are proportionably small, and that the artery which supplies the bulb of the urethra in the male is distributed to the bulb of the vagina in the female.

DISSECTION OF THE FEMALE PELVIS.

It is presumed that you have some knowledge of the anatomy of the viscera in the male pelvis. Our present concern is, to examine the organs superadded in the female, viz., the

* See Burns' Midwifery.

† See Professor Tiedmann, Von den Duverneyschen Drüsen den Weibs. Heidelberg, 1840.

uterus and its appendages. First, study their position in a general way; secondly, their special anatomy.

General Position of the Uterus and its Appendages.—Looking into the female pelvis, we see the uterus interposed between the bladder in front, and the rectum behind. From each side of the uterus a broad fold of peritoneum extends to the side of the pelvis. These folds are called the *broad ligaments* of the uterus, fig. 84, p. 397. On the posterior surface of the ligament are the ovaries, one on each side. They are completely covered by peritoneum, and suspended to the ligament by a small peritoneal fold. Along the upper part of the ligament we find between its layers a tube about four inches long, called the Fallopian tube. A better name for it is the oviduct, since its office is to convey the ovum into the uterus. For this purpose, one end of it terminates in the uterus, while that nearer to the ovary expands into a wide mouth, furnished all round with prehensile fringes, which, like so many fingers, grasp the ovum as soon as it is ready to escape from the ovary. Lastly there run up to the ovary, between the layers of the broad ligament, the spermatic, or rather the ovarian vessels and nerves which arise from the aorta in the lumbar region, just as they do in the male, because the ovaries are originally formed in the loins.


Look on the anterior surface of the broad ligament; and you will see on either side between its layers what is called the *round ligament* of the uterus. This is a cord which proceeds from the fundus of the uterus, through the inguinal canal, just like the spermatic cord in the male, and terminates in the “mons Veneris.” Besides one or two small blood-vessels, it contains muscular fibres analogous to those of the uterus: these increase very much in pregnancy, so that about the full term, the cord becomes nearly as thick as the end of the little finger.

Side View of the Female Pelvis.—After the removal of the innominate bone, as described at p. 354, the vagina, rectum, and bladder should be moderately distended, and a catheter passed into the urethra. This done, trace the reflections of the peritoneum.

Reflections of the Peritoneum.—From the front of the rectum the peritoneum is reflected on to a small part of the

vagina, thus forming what is called the "recto-vaginal pouch." From the vagina we trace the peritoneum all over the *back*, but only about half way down the *front* of the uterus; thence it is at once reflected over the posterior surface of the bladder, on to the wall of the abdomen.

Fig 83.



VERTICAL SECTION THROUGH THE FEMALE PELVIS.

We can easily understand, that in cases of ascites the water might distend the recto-vaginal pouch, and bulge into the vagina, and that it would be practicable to draw it off through this channel.*

Pelvic Fascia.—To the description of the fascia already given in the dissection of the male pelvis (p. 356), nothing need here be added except that from the side of the pelvis it is reflected over the side of the vagina and the uterus as well as the bladder.

* In the Medical Communications, vol. i., a case is related in which four gallons of fluid were drawn off by tapping through the vagina. The woman immediately afterwards passed urine, which she could not do before. See also a case in Med. and Phys. Journal, vol. ii., p. 412.

It is this fascia which in great measure supports and slings up the uterus in its proper level in the pelvis. Of its efficacy in this respect any one may be convinced by trying to pull down the uterus through the vagina. When, from any cause, the fascia becomes relaxed, there is consequently a greater liability to "prolapsus uteri."

Levator Ani.—For the description of this muscle, see p. 364.

Bladder.—The female bladder is broader transversely, and upon the whole, more capacious than the male. The vesical plexus of veins, as might be expected, is not so large, and there is no prostate gland. But the short urethra has a constrictor muscle, as in the male, and is supported in a similar manner by the pelvic fascia.

Venous Plexus about Vagina.—Though the veins round the neck of the bladder are comparatively small in the female, attention will be immediately attracted by the plexus of large veins which surround the vagina and the rectum. They seem out of all proportion to the size of the arteries. Their congestion in pregnancy sufficiently accounts for the dark color of the vagina and the external organs, and the frequent occurrence of hæmorrhoidal tumors.* It is necessary to remove these veins, with the cellular tissue in which they are imbedded, before a clear view of the parts can be obtained.

Urethra.—Of the urethra we have already spoken at p. 383. But in the side view of the parts we have the opportunity of observing how closely the bladder and urethra are connected to the upper wall of the vagina, and we can have no difficulty in understanding why in cases of protracted delivery, it sometimes happens that the contiguous coats of the bladder and the vagina give way, and that there remains a fistulous communication between them, which continues to be a depending drain for the urine.

Vagina.—At the present stage of the dissection slit open the whole of the vagina along the side, and then you will obtain a clear idea of the manner in which it embraces the lower

* During pregnancy, varicose tumors may form even in the vagina. In the Berlin Med. Zeitung, 1840, No. 11, a case is related of a woman who, at the sixth month, bled to death from the bursting of a large vein in the vagina. Other cases of the kind are related by Siebold.

end of the uterus, and of the extent to which the neck of the uterus projects into it. This proceeding will also make more intelligible what is to be observed respecting the length, axis, and width of the vagina.

The length of the vagina, in an unimpregnated healthy adult, is, on an average, about $4\frac{1}{2}$ inches. It may be more, or it may be less; the difference in each case depending upon the depth of the pelvis, the stature and age of the individual. Owing to the curved direction of the vagina, the anterior wall is about three quarters of an inch shorter than the posterior. The vagina, however, is never so long, that we cannot, during life, feel the neck of the uterus projecting at the top of it; higher up, or lower down, according to circumstances. For instance, it is a little lower in the erect than in the recumbent position: again, in the early months of utero-gestation, the uterus descends a little into the vagina, so that this canal becomes shorter; the reverse holds good when the uterus begins to rise out of the pelvis.

The axis of the vagina is slightly curved, with the concavity upwards: for all practical purposes it may be said to correspond with the axis of the outlet of the pelvis; whereas the axis of the uterus corresponds with that of the brim of the pelvis.

The width of the vagina is not uniform throughout. The narrowest part is at the orifice: it is also a little constricted round the neck of the uterus. The widest part is about the middle: supposing a transverse section to be made through it in this situation, it would present the appearance of a broad horizontal fissure. If, therefore, you would insert the bivalve speculum upon anatomical principles, consequently with the least amount of pain, the blades of the speculum should be vertical when introduced into the orifice of the vagina, and afterwards turned horizontally.

Uterus.—The uterus is the receptacle which receives the ovum, retains it for nine months so as to bring it to maturity, and then expels it by virtue of its muscular walls. Enough has been already said at page 387 of its situation and peritoneal connections. We have now to notice that its axis slants forwards, so that, upon the whole, the axis of the vagina and uterus describes a curve pretty nearly parallel to the axis of

the pelvis The uterus, then, is so placed that it is ready to rise out of the pelvis into the abdomen after the embryo has attained a certain size. The wisdom of this provision becomes manifest in cases where, from some accidental cause, the natural direction of the uterus has been altered, so that, instead of rising freely out of the pelvis, it becomes what is called retroverted; that is, the large end is tilted backwards against the sacrum, while the apex presses forwards against the neck of the bladder. It need hardly be said that such a mal-position more or less obstructs the passage of both urine and fæces.

Respecting the size and shape of the uterus in the unimpregnated state, a better idea is conveyed by a single glance than by any description. Anatomists generally liken it to what a pear would be if it were a little flattened; in other words, it is triangular with the angles rounded. As to its average size, it is about three inches long, two inches broad, and one inch thick, at the upper part; but there is variety in this respect, arising from age, the effect of pregnancies, and other causes.

For convenience of description, we divide the organ into the fundus, the body, and the cervix. The term fundus is applied to that part which lies above the level of the Fallopian tubes (p. 391). The body is the central part, while the cervix is the narrow part which projects into the vagina. The vagina is very closely attached round the neck of the uterus, but you observe that it is attached higher up behind than in front. The mouth of the uterus is at the apex of the neck. It is a transverse fissure, with a slightly prominent lip in front and behind, and, from its fancied resemblance to the mouth of a tench, it was called by the old anatomists the "*os tincae*."

[The average thickness of the walls of the body of the uterus is five lines, of the neck about four. The fundus of the uterus is about a quarter of an inch above the Fallopian tubes. At puberty the organ weighs from six to ten drachms, in the adult one and a half to two ounces, and at full term from one to three pounds.]

Postponing for the present the examination of the interior of the vagina and uterus, let us pass on to the vessels and nerves of these organs.

Uterine and Vaginal Arteries. — In addition to the branches given off in the male (p. 365), each internal iliac artery, in the female, furnishes a branch to the uterus, another to the vagina.

The *uterine* artery proceeds from the anterior division of the internal iliac towards the neck of the uterus, and then ascends tortuously by the side of the uterus, giving off numerous branches to it. The fundus of the uterus is chiefly supplied by branches from the ovarian arteries.

[A branch of the uterine artery called the circular, lies in the loose cellular tissue around the neck of the uterus—another also passes around the neck through its substance. These vessels vary much as regards their size, and if wounded in the operation of dividing the strictured portion of the neck for dysmenorrhœa, will give rise to very troublesome hæmorrhage. We have succeeded in demonstrating, on two occasions, by means of a fine wax injection, the exact position and size of these vessels.]

The *vaginal* artery ramifies along the side of the vagina, and sends branches to the lower part of the bladder and the rectum.

Nerves of the Uterus.—The nerves of the uterus are derived from the sacral nerves, and from the hypogastric plexus (p. 368). They accompany the blood-vessels to the neck of the uterus, and ascend with them along the sides of the organ.

The greater part of the nerves soon leave the vessels, and, subdividing, sink into the substance of the uterus, chiefly about its neck and the lower part of its body. But some very small filaments continue with the vessels, and form around them plexuses, upon which, according to the dissections of Mr. Beck,* minute ganglia are found.

The question whether the nerves of the uterus enlarge during pregnancy like the arteries, has given rise to much discussion, and is still “sub judice.” Great names might be quoted in support of the most opposite opinions. However, in a surgical point of view, the *os uteri* may be said to have no nerves, for it is insensible to the cautery and to the knife.

The *absorbent* vessels of the uterus are very small in its un-

* Philosophical Transactions for 1846.

impregnated state, but greatly increase in size when it is gravid. Those from the fundus and the ovaries proceed with the spermatic vessels to the lumbar glands; and this is the reason why the lumbar glands are sometimes affected in ovarian disease. Those from the body and the lower part of the uterus accompany the uterine artery, and join the glands in the pelvis; some, however, run along with the round ligament to the groin: hence in certain conditions of the uterus, the inguinal glands may be affected.

Structure of the Vagina, Uterus, Ovaries, and Oviducts.—The uterus, vagina, ovaries, and oviducts, should now be collectively removed from the pelvis for the purpose of examining their internal structure. And first of the vagina.

Supposing the vagina to be laid open, we observe that it is lined by a mucous membrane of a pale rose color: and that this membrane is rough and furrowed, especially near the orifice. A more or less prominent ridge runs along its anterior, and another along its posterior wall. From either side of these, which are called "*columnæ rugarum*," proceed a series of transverse ridges with rough jagged margins directed forwards. They are well marked in young females who are still virgins, but repeated parturition and increasing age gradually smooth them down. The use of the vaginal rugæ is to excite the sensibility of the glans in coition. They themselves also possess keen sensibility, for they are richly provided with papillæ.

The mucous membrane has a thick epithelial lining, and in the submucous tissue is an abundant supply of muciparous glands. They increase in number and size towards the uterus, for the probable purpose of facilitating by their secretion the passage of the child. This secretion is thick, creamy, slightly acid, and differs altogether from the secretion of the uterine glands, which is glairy, albuminous, and slightly alkaline. When poured out in excess, and somewhat altered in its character, it constitutes what is called "leucorrhœa."

The chief strength of the vagina depends upon a fibro-cellular coat, which is about one-twelfth of an inch in thickness. If this coat be minutely injected, we find that it is made up almost entirely of the inosculations of blood-vessels; so much so that some anatomists regard it as erectile tissue.

Before the uterus is laid open, examine a little more fully the shape of that portion of the neck which projects into the vagina. First, then, it appears that the back part of the cervix projects into the vagina more than the front; but this merely arises from the vagina being attached higher up behind it. Supposing that the vagina were entirely cut away from the cervix, the anterior lip of the uterus would appear to project a trifle more than the posterior. For this reason, as well as on account of the natural slope forwards of the uterus, the front lip is always felt first in an examination per vaginam.* The length, however, and the general appearance of the vaginal part of the cervix varies according to the age of the individual, and it is also considerably altered by parturition. In the adult virgin it is smooth, plump, round, and projects half an inch: its mouth is a small transverse fissure. But after the stretching which it undergoes in labor, one cannot wonder that it loses its plumpness, that the lips become flaccid and fissured, and the mouth larger than it was before.†

The uterus must now be laid open by a longitudinal incision, to examine its interior. The first thing that attracts attention is the great thickness of the walls. Before you come into the proper cavity in the body of the uterus, you have to slit up a long narrow channel which leads up into it through the neck. This channel is not of the same dimensions throughout: it is dilated in the middle, and gradually narrows towards each end. The upper end, which leads into the body of the uterus, is called "*os internum*," the lower end, which leads into the vagina, "*os externum*." The passage itself is called the cavity or the neck. It remains unchanged in pregnancy for some

* This is the only way to reconcile the discrepancies one meets with in anatomical works, respecting the comparative length of the lips of the uterus. Kraus, Weber, Busch, and others say the anterior is the longer; Mayer, Meckel, Quain, and others, the posterior.

† Instances are recorded in which the neck of the uterus is preternaturally long. It has been known to project, even as much as an inch and a half into the vagina. In such cases it gradually tapers, and terminates in a very narrow mouth. This is said to be one cause of sterility, and it is recommended either to dilate the mouth, or to cut off a portion of the neck. In support of this opinion, it is stated that Dupuytren was once consulted by a lady on account of barrenness; finding the neck of the uterus unusually elongated, he removed a portion of it, and in due time the lady became pregnant. (Hyrtl, Handbuch der top, Anatom.)

time after the cavity in the body has expanded, but gradually disappears with the increasing size of the embryo.

[Dr. Isaac E. Taylor has proved this to be an error by repeated examinations of women at different periods of pregnancy, even within an hour or two before labor. These examinations were conducted by Dr. Taylor and the house staff, at Bellevue Hospital in 1860-61. See his paper on the subject in the Bulletin of the New York Academy of Medicine, vol. i.]

The shape of the cavity in the body of the uterus is triangular, with the apex towards the cervix. In a virgin uterus the cavity is remarkably small, and its sides are convex; but in a uterus which has borne many children, the cavity has lost the convexity of its sides, and has increased in capacity. Each angle at the base is somewhat produced, and leads to the minute opening of the oviduct. This production of the angles is noticed more or less in different females, and is the last indication of the two horns of the uterus in some orders of mammalia.

The interior of the uterus is perfectly smooth at the fundus, but just the reverse at the cervix. Here you remark that there is a central longitudinal ridge, both in front and behind (just as in the vagina), and from these, other closely set ridges curve off laterally, like the branches of a palm-tree. The old anatomists called it "*arbor vitæ*." The roughness produced by these ridges occasions an impression as though we were touching cartilage whenever a metallic sound is introduced into the uterus.

The neck of the uterus is provided with small glands, of which the minute ducts open in the furrows between the ridges referred to. The secretion of these glands is glairy, albuminous, slightly alkaline, and sometimes comes away in such abundance as to be exceedingly troublesome. Soon after conception, the secretion dries up so as to plug the mouth of the uterus, but shortly before and during parturition it is poured out in great quantity, to facilitate the passage of the child. It happens occasionally, that one or more of the ducts of these glands become obstructed, and then dilate into small transparent vesicles, which gradually rise to the surface and burst. These were first described by Na-

both,* and supposed to be true ova : hence their name "*ovula Nabothi*."

The mucous membrane of the uterus is much more delicate than that of the vagina, and is scarcely separable from the subjacent tissue. The greater part of it is lined by a ciliated epithelium, but the epithelium of the cervix is tessellated like that of the vagina. The chief peculiarity, however, about the membrane is this ;—every part of it is covered with minute follicles or tubes arranged at right angles to its surface. These tubes become greatly developed shortly after impregnation, and are presumed to take an active part in the formation of the "*membrana decidua*."

The greater portion of the walls of the uterus consists of muscular fibres of the unstriped or involuntary kind, like that of the bladder and alimentary canal. The texture of the fibres is so close, that it is exceedingly difficult to ascertain their precise arrangement ; indeed, in the unimpregnated uterus it is useless to make the attempt, for they are comparatively atrophied. But in pregnancy, when all the component structures of the organ undergo simultaneous development, the fibres run either longitudinally or in concentric circles. The longitudinal fibres form a thick outside stratum, of which the direction is from the fundus to the neck of the uterus. The concentric circles form the deeper strata, and surround the orifices of the oviducts. Again, there are others which take a more oblique course, so that, upon the whole, their collective disposition is such as to exert equal pressure on all sides when called into operation.

At the same time that they expel the foetus, the muscular fibres perform another very important function ; they close the large venous sinuses developed for its nutrition. Therefore, very little hæmorrhage accompanies the expulsion of the placenta, provided it has been attached to the fundus or the side of the uterus. But every one knows the danger of what is called a "*placenta prævia*." Here the placenta, planted, as it were, over the orifice of the uterus, is attached to a part of the organ which must of necessity expand during labor : thus the reverse of what is common takes place, and every pain

* *De sterilitate mulierum.* Lips. 1770.

increases, instead of checking, the flooding. For the same reason, paralysis of the muscular fibres in immediate connection with the placenta, be it where it may, is likely to be a source of serious hæmorrhage in parturition.

Fallopian Tubes.—The Fallopian tubes or oviducts are situated, one on each side, along the upper border of the broad ligament of the uterus, fig. 84. They are about four or five inches long. One end leads into the uterus; the other terminates in a wide funnel-shaped mouth, surrounded by a

Fig. 84.




DIAGRAM OF THE UTERUS, ITS BROAD LIGAMENTS, THE OVARIES, AND FALLOPIAN TUBES.

- | | |
|--------------------|--|
| 1. Uterus. | 4. Fimbriated extremity of Fallopian tube. |
| 2. Ovary. | |
| 3. Fallopian tube. | 5, 5. Broad ligament. |
| | 6. Vagina. |

fringe, like the opening in a pink flower. This is best seen by floating it in water. One or two of the fringes are generally connected with the outer end of the ovary. If the subject be well injected, you will see that they are richly supplied with blood-vessels from the spermatic artery. If you open the oviduct from the expanded end, and introduce a probe into it, you will find that the tube runs very tortuously at first, then straight into the uterus, gradually contracting in size all the way, so that the uterine orifice scarcely admits a bristle. Its mucous lining is gathered into longitudinal wavy folds especially at the ovarian end: with the microscope we detect that it is provided with a columnar ciliated epithelium.

The free end of the oviduct communicates with the cavity of the peritoneum. This is the only instance in the body where a mucous membrane is directly continuous with a serous one. It explains how the embryo may escape into the peritoneal cavity, though this be an extremely rare occurrence, because such ample provision has been made against it. It also explains what is said to have taken place, namely, the escape of the water in dropsy through the Fallopian tubes.

Ovaries.—The ovaries are the most important parts of the female generative organs, since they contain the ova or the germs. They were called by Galen "*testes muliebres*." We have already seen that they are suspended to the back of the broad ligament of the uterus by a short peritoneal fold, which transmits their proper vessels and nerves: besides this, they are connected to the uterus by a thin cord, called the *ligament* of the ovary. They are of an oblong form, with the long axis transverse, and a little smaller than the testicles. In females who have not often menstruated, their surface is smooth and even: in after life it becomes wrinkled and scarred by the repeated escape of the ova.

The ovary has nearly the same coverings as the testicle. There is first the serous coat, and beneath it the proper fibrous coat, or what is called the "*tunica albuginea*." If a section be made through the ovary, you find that it contains transparent vesicles, bedded in a soft fibrous-looking tissue, remarkably vascular when properly injected, called the "*stroma*" of the ovary.

The transparent vesicles just alluded to are the ovisacs, or "Graafian"* vesicles. They vary in number from eight to twelve, and in size from that of a pin's head to a pea. The smallest are near the centre; but as they advance towards maturity, they gradually approach the surface, increasing at the same time in size. They contain a transparent albuminous fluid. If after puncturing one of the larger vesicles you examine its contents under a microscope, you find suspended in it the true ovum or germ.† It is this minute body which, escaping from the Graafian vesicle on the surface of the ovary,

* So called after De Graaf, a Dutch anatomist, who discovered them in 1672, and believed they were the true ova.

† This was first distinctly pointed out by Von Baer in 1827.

is grasped by the Fallopian tube and conveyed into the uterus. The ruptured vesicle is converted soon afterwards into a yellowish-looking mass called "*corpus luteum*," which persists for a while, and degenerates afterwards into a small fibrous cicatrix.

The ramifications of the ovarian artery through the ovary are remarkable for their convolutions: they run in parallel lines, as in the testicle.

[On holding up the broad ligaments to the light, we sometimes see an appendage to the ovaries, in the shape of a number of small tortuous tubules between the ovary and the Fallopian tube. This is the remains of the Wolffian body of the foetus; it is about an inch broad, with its base to the tube, and apex to the attached portion of the ovary. These tubules are about twenty in number, and each tube is inclosed in a fibrous capsule.]

DISSECTION OF THE ABDOMINAL VISCERA.

HAVING already studied the position and relations of the several abdominal viscera (p. 304), you should now examine their particular configuration and structure.

Liver.—The liver is the largest gland in the body. In an adult male it weighs between four and five pounds. In shape we hardly know what to compare it to. It is thick and round at the back; towards the front it gradually slopes to a thin border. Its surface is entirely covered by peritoneum, except a small part behind, which is connected to the diaphragm by cellular membrane, and just in the hollow for the gall-bladder. On the upper surface of the liver, which is convex, in adaptation to the diaphragm, there is nothing deserving of notice beyond the indication of the division of the organ into a right and a left lobe, the right being the larger.

[The size and weight of this organ, as well as its color, are much affected by disease. The usual measurements and weight are as follows: weight from three to four and a quarter pounds; transverse diameter, ten to twelve inches; antero-posterior, six to seven inches; thickness, at the centre and posterior border, from two to three and a half inches.]

But if the liver be turned on its back, you see that the under surface is irregular, and that there are several fissures in it for the passage of vessels.

Fissures.—First, there is the “*longitudinal fissure*,” dividing the right from the left lobe; it contains the round ligament or the obliterated umbilical vein of the foetus. The continuation of the longitudinal fissure to the posterior border of the liver contains the remains of what was in the foetus, the “*ductus venosus*,” it is therefore called the “*fissure of the ductus venosus*.” Secondly, there is the “*transverse fissure*,” where the great vessels enter the liver. Thirdly, there is the hollow for the gall-bladder. Lastly, in the same line with this, is the channel for the vena cava. Now the relative position of these parts may, perhaps, be best impressed on the memory by comparing them collectively to the letter H. The transverse fissure represents the cross bar of the letter, the longitudinal fissure represents the left bar; the gall-bladder and vena cava together make the right bar.

Lobes.—This comparison to the letter H serves to remind us of certain other parts of the liver, to which particular

Fig. 85.




DIAGRAM OF THE UNDER SURFACE OF THE LIVER.

names have been given. In the upper square of the letter is a tongue-like lobe called “*lobulus Spigelii*,” this is connected to the right lobe by a ridge or tail, called “*lobulus caudatus*.” Again, the area of the lower square is called the “*lobulus*

quadratus." Lastly, the right lobe presents a shallow depression, adapted to the right kidney.

The liver has a thin fibro-cellular coat or capsule, which is best seen on those parts of it not covered by peritoneum. So far it is like other glandular organs. But the capsule does not send down partitions to form a framework for the interior of the organ. At any rate, nothing of the kind is observed with the naked eye when a portion of the capsule is stripped from the surface. Whether there be a cellular framework or not between the lobules, it is certain that it must be exceedingly delicate: this is the reason why the liver is so liable to be lacerated by external violence, or by the action of the abdominal muscles.


Minute Anatomy of the Liver.—The liver is composed of the ramifications of the portal vein and hepatic artery which carry the blood *to* it, and secrete the bile; of the ramifications of the excretory or hepatic duct; of the ramifications of the hepatic veins, which return the blood *from* it, and of nerves and absorbents. Now, it will facilitate the understanding of the subject, if you bear in mind,—1, that the portal vein, hepatic artery, and hepatic duct, ramify together from first to last. They are inclosed in a sheath of loose fibro-cellular tissue, called "*Glisson's capsule*," which enters with them at the transverse fissure: this explains why the portal vein, when cut transversely, does not gape, but partially collapses: 2, that the hepatic veins run from first to last by themselves, and terminate in the vena cava as it passes through the liver. These, having no cellular sheath, always appear, on transverse section, with open mouths.

Now come the questions, What is the distribution of the minute branches of the portal vein? what that of the hepatic vein? These questions are determined by injecting each set of vessels simultaneously with a fluid of a different color,—say a blue fluid for the portal vein, and a yellow for the hepatic. Afterwards, sections in different directions must be made, and examined under the microscope.

In answer to the first question, it may be stated that the minute ramifications of the portal veins map out the substance of the liver into small spaces called *the lobules*. The boundary veins are termed "*inter-lobular*." The spaces can

be seen even with the naked eye. They have a different shape according to the direction in which they are cut: on the surface of the liver, or what comes to the same thing, in a transverse section, they look like mosaic pavement (fig. 86); but in a perpendicular section they somewhat resemble an oak leaf. From the circumference of the lobule, the capillaries of the portal vein penetrate into its area, minutely inosculating with each other, and then freely communicate towards the middle of the lobule with the radicles of the hepatic vein.

Fig. 86.



TRANSVERSE SECTIONS OF THREE LOBULES OF THE LIVER, MAGNIFIED TO SHOW THE PORTAL VENUS PLEXUS.

The second question may be answered thus:—The radicles of the hepatic veins commence within the lobule by free inosculations with those of the portal veins, and converge towards a single vein termed "*intra-lobular*," which runs down the centre of the lobule. The central vein opens at once into an hepatic vein termed "*sub-lobular*," larger or smaller as the case may be, upon which the lobule is sessile. Thus, then, on a perpendicular section, we should see the hepatic veins running into the central vein, like the side veins of an oak leaf do into the midrib (fig. 87); while on a transverse section, they would present a starred appearance.

And now a word or two respecting the ultimate ramifications of the hepatic artery and the hepatic duct.

The *hepatic artery*, entering the liver at the transverse fissure, divides and subdivides with the portal vein, and ulti-

mately ramifies with it between the lobules. A few only of the arterial capillaries enter the lobules. The chief office of the artery appears to be, to supply the coats of the other vessels of the liver,—in other words, the machinery of the organ.

The *hepatic ducts* commence within the lobules by numerous ramifications, which form a close network near the circumference of each lobule. From this network branches proceed on all sides, and accompany the portal veins.


Great doubt still exists as to how the ducts begin. Some say they commence by blind extremities; others by simple channels between the hepatic cells. Be this as it may, their radicles are surrounded by minute cells, which are in fact the essential organs which secrete the bile.

The interior of each lobule, that is, the space left between the several vessels, is filled by the "*hepatic cells*." These are nucleated, and have a mean diameter of about $\frac{1}{1000}$ th of an inch. In some cases they contain fat globules, and when these accumulate in large quantities, they constitute what is called a "fatty liver." The office of the cells is to separate the bile from the blood, in some way which we do not understand: when filled with bile, they burst, and discharge themselves into the bile ducts.

Such, in outline, is the minute anatomy of the liver. Rightly understood, we can have no difficulty in determining, in a section of the organ, whether the portal or the hepatic venous system be congested.*

The present views entertained concerning the functions of the liver may be thus briefly expressed:—1. It purifies the blood by excreting carbon and hydrogen, which, being subsequently re-absorbed, combine with oxygen, and thus help to keep up the heat of the body. 2. It forms sugar, which passes into

Fig. 87.



LONGITUDINAL SECTIONS OF THE LOBULES OF THE LIVER. INTRA-LOBULAR VEINS SEEN JOINING THE SUB-LOBULAR.

* For further information on the subject see the original observations of Kiernan in the *Philosoph. Trans.* for 1833.

the hepatic veins, and being consumed in the process of respiration, helps to maintain animal heat. 3. The bile assists in converting the chyme into chyle, and reducing it into a state fit to be absorbed by the lacteals. 4. The bile acts as a slight natural aperient.

Gall Bladder.—The gall bladder, a reservoir for the bile, is confined by the peritoneum in a slight depression on the under surface of the right lobe of the liver (p. 306). Observe that it is shaped like a pear. Its size varies a little in different subjects; generally speaking, it is about four inches long, and will hold about one and a half ounces of fluid. Its narrow end, or neck, makes a bend downwards, and terminates in a duct, called the "*cystic*," which, after a course of about one inch, joins the hepatic. The common duct, "*ductus communis choledochus*," formed by their union, is about three or four inches long. Remember that it opens into the back of the descending part of the duodenum, after running very obliquely through the coats of the bowel.

Exclusive of its partial peritoneal covering, the gall bladder has only two coats. The outer, consisting of fibro-cellular tissue, contains organic muscular fibres; the inner is the mucous coat.

The *mucous* coat is generally tinged yellow by the bile. Its chief peculiarity is that it is gathered into ridges which give it a honeycomb appearance. It is covered by a columnar epithelium, which secrets an abundance of viscid mucus. Furthermore, observe that, at the bend of the neck of the gall bladder, both its coats project very much into the interior, making the opening considerably narrower than it appears to be outside. Lastly, in the cystic duct, the mucous membrane presents a series of folds, arranged, one after the other, as to form a complete spiral valve. The probable use of this is to prevent the too rapid flow of bile. Besides acting as a reservoir for bile, the gall bladder adds a bitter principle to it, which makes it keep. The bile in the liver is sweet; that in gall bladder is intensely bitter.

[We once met with a liver in an adult male, an emaciated and phthisical subject, where the gall bladder was absent, and without any trace of a fissure for its lodgment. The organ was small, weighing but a pound and three quarters, and con-

sequently, there was no line of demarcation to form a lobus quadratus.]

Spleen.—The spleen is a very vascular spongy organ, varying in size according to the quantity of blood in it, and fluctuating in weight, consistently with health, between five and twelve ounces. Under ordinary circumstances it is of a reddish blue color, owing to the large amount of blood in it. But the proper color of the substance of the spleen is a yellowish grey. This is well seen in cases of cholera, where the organ is bloodless and collapsed; it is also proved by injecting the splenic artery with water, which returns freely by the veins, and washes out all the blood.

Its shape is elliptical. In its natural position it is so placed that the long axis is nearly vertical. The outer side, adapted to the diaphragm and ribs, is convex, while the inner side, adapted to the great end of the stomach, is concave. The blood-vessels enter the concave side by several ramifications, corresponding to the primordial lobes of the organ.

Besides its peritoneal coat, the spleen has a fibrous capsule, which is elastic, to accommodate itself to the varying size of the organ. This capsule not only covers the outside of it, but sends into the interior numerous threads, which cross each other in various directions, and form a network, dividing the spleen into so many chambers. Moreover, it is interesting to observe that at the points where the bands cross they are secured by a small white knot. Besides constituting the general framework of the organ, the capsule provides the vessels with sheaths which support them throughout their ramifications in the interior.

The splenic chambers are filled by what is called the "pulp" of the spleen. This is a soft reddish-brown substance, easily scraped off with a knife. Examined under the microscope, it consists entirely of gland cells of a pale yellow color, and smaller than the corpuscles of the blood. Here and there, however, in the pulp are found other larger cells, of a white color and perfectly spherical form.* Attention was first


* In the human spleen they are about $\frac{1}{80}$ th inch in diameter. But it is useless to look for them unless the subject be exceedingly fresh, for they soon soften and melt in the pulp. It is better, therefore, to examine them in the spleen of a sheep or bullock, in which animals they are about $\frac{1}{40}$ th inch in diameter.

directed to them by Malpighi, and they have therefore been called after him the "Malpighian bodies." They are not free like the smaller cells, but attached, each by a slender pedicle, to one of the threads. The pedicle contains a small artery, which ramifies over the surface of the cell, and then terminates in a brush of capillaries which spread out in the pulp. The interior of the cell is filled with a clear fluid and a multitude of smaller cells, so that it is but a closed sac, containing secreting elements.

How are the blood-vessels distributed? The splenic artery enters the spleen by several branches, which ramify throughout the organ, supported by sheaths derived from the capsule.* These ramifications, however, do not communicate with each other, for, if injection be thrown separately into one branch, it fills part only of the spleen. The smallest ramifications spread out in a brush-like manner through the pulp, and then lead into the veins. The veins themselves form in the pulp a minute network. But this part of the circulation is not yet thoroughly made out.

Thus, it appears that the spleen is essentially a great blood gland: that it consists of chambers filled with secreting gland cells of various size, between which ramify minute arteries and veins. It is presumed that the cells in some way or other elaborate the blood. Some physiologists consider the spleen to be the birth-place of the white corpuscles of the blood. Others have arrived at the

Fig. 88.



SECTION OF THE KIDNEY.

- 1. Ureter.
- 2. Pelvis of the kidney.
- 3, 3, 3. Papillæ.

* The ramifications of the splenic artery may be well seen by washing away the pulp, and floating the flucculent-looking spleen in the water.

conclusion that it serves as the graveyard of the red ones. The precise function of the organ is unknown.

[The measurements of the spleen, in health, are usually, as regards length, four to five inches; width, two to three inches; thickness, one and a half inches.]

Kidney.—With the form and color of the kidney all are familiar. Its weight is about five ounces in the male, rather less in the female. In the majority of cases, the left is heavier than the right.

It is surrounded by a thin capsule, which adheres by minute vessels to its surface, but does not penetrate into its interior. For this reason the capsule can, under ordinary circumstances, be readily stripped off; if not, the presumption is, that its adhesion is the result of disease.

The best way to display the anatomy of the kidney is to make a longitudinal section through it. This shows two distinct structures. The superficial structure, called the "*cortical*," is of a uniform red color, because the blood-vessels are most copiously distributed in it; it is the secreting part of the organ. The deeper structure, called the "*tubular*," consists of the minute tubes which carry off the urine. These tubes, converging from the cortical part, are collected into from ten to sixteen pyramidal bundles (fig. 88): their nipple-like points (termed *papillæ*) consisting of the terminations of a great number of coalesced tubes, project into the pelvis of the kidney; the common receptacle of the secretion from all parts of the organ.*

The pelvis of the kidney is the dilated beginning of the ureter. It is funnel-shaped, and its broad part divides into two principal channels, which again branch out and form from eight to twelve cup-like excavations (called the *calices*). Into each of these calices one, sometimes two, *papillæ* project.


The renal artery runs to the kidney between the vein in front and the ureter behind, and divides into five or six branches, which, again subdividing, make their way between

* Each pyramid represents what was, in the early stage of the kidney's growth, a distinct and independent lobe. In the human subject the lobes gradually coalesce, and no trace of their primordial state remains, except the pyramidal arrangement of the tubes. But in the kidneys of the lower mammalia, of birds and reptiles, the lobes are permanently separate.

the pyramids to the cortical substance. Their minute ramifications terminate in one of two ways, either in plexuses round the tubes, or in the Malpighian bodies.

Malpighian Bodies.—Examine with a microscope a portion of a minutely injected kidney. Observe that the cortical substance is full of red round knots, formed by the coils of minute blood-vessels; these are the *Malpighian** bodies (fig. 89). They are, on an average, about $\frac{1}{130}$ th of an inch in

Fig. 89.



- a. Artery.
- v. Vein.
- c. Capsule.
- d. Urinary tube.

diameter. According to the recent researches of M. Bowman,† each is loosely surrounded by a flask-like capsule, formed by the dilatation of the urinary tube. The continuation of the tube passes off from that part of the capsule opposite to where the artery entered; after making many convolutions, it enters one of the pyramids, and discharges its contents into the pelvis. The vein returning the blood from the Malpighian body leaves the capsule close to where the artery entered; but instead of going at once out of the kidney, as in other organs, it forms a capillary plexus round the convolutions of the tube. The special purpose of this plexus appears to be the

secretion of the solid matter of the urine; while the Malpighian body filters the watery part of the urine into the capsule, and washes the more solid part down the tube.‡

Lastly, the minute urinary tubes, the pelvis of the kidney, and the ureter, are lined by tessellated epithelium.

Supra-Renal Capsule.—This body is situated at the top of the kidney, and in shape resembles a cocked hat. It is surrounded by a thin fibrous covering, which sends down parti-

* So named after Malpighi, a celebrated Italian anatomist who lived during the middle and latter part of the seventeenth century.

† See his paper in the *Philosoph. Trans.* for 1842, Part I.

‡ That the vessel leaving the Malpighian body is a vein, and that a constituent part of the urine is secreted by venous blood, is inferred from two reasons: 1, from the analagous case of the vena portæ, out of which the bile is elaborated in the liver; 2, from the fact that in reptiles the urine is secreted from venous blood.

tions into the interior through furrows observed on the surface.

If a perpendicular section be made through it, we find that it consists of a pretty firm exterior or cortical part, and of an interior soft pulpy substance. The cortical portion is of a yellow color, and forms the principal part of the organ. When examined with the high powers of the microscope, it is found to be composed of independent closed tubes, about $\frac{1}{700}$ th of an inch in diameter, arranged perpendicular to the surface, and covered by the minute ramifications of the arteries. The tubes contain a few nucleated cells, but chiefly minute granules. The central part is of a very dark brown color, and so soft in texture that one might mistake it for a cavity. It appears to consist almost entirely of a plexus of minute veins.*

[The kidneys measure in length from three to five inches; width, two to three; and in thickness a little more than an inch. We may find these organs, united by an isthmus of kidney structure, passing across the spinal column, giving rise to that variety known as the horse-shoe kidney.]

Of late years, the minute structure and functions of the renal capsules have been much investigated, in consequence of the discovery, made by Dr. Addison, of the close relation which exists between disease in these bodies and a brown discoloration of the skin. But their precise function is still unknown.

Stomach and Intestines—The alimentary canal is composed of three coats, differing in structure, and connected together by cellular tissue. First is the *serous* or peritoneal coat, described at p. 313. Secondly, under the serous is a *muscular* coat, upon which the chief strength of the canal depends. It consists of two distinct strata of fibres; the outer stratum is longitudinal, the inner circular. This arrangement not only makes the bowel stronger, but regulates its peristaltic action; for the longitudinal fibres by their contraction, tend to shorten and strengthen the tube, while the circular fibres contract upon and propel its contents to greater advantage. Lastly is the internal or *mucous* lining. This is the most complicated of

* Consult "A Physiological Essay on the Thymus Gland," by Simon, London, 1845.

the three, for it presents different characters in different parts, according to the functions which it has to perform.

Stomach.—The stomach should be distended with air, that we may see its size and shape. Its size will vary in different instances, according to the habits of the individual; and its shape, for want of a better comparison, is likened to the bag of a bagpipe. The great bulge of the organ on the left of the œsophagus, is called the œsophageal or the cardiac end; while the smaller part at which the food passes out is termed the pyloric end. Just before the pylorus, the stomach bulges into a pouch, called "*antrum pylori*." It is analogous to that stomach in the ruminant in which the runnet is found. Again, we speak of the upper margin, or the lesser curve; of the lower margin, or the greater curve; of the anterior and the posterior surface.

Of the *muscular* coat of the stomach, some fibres take a longitudinal, others a circular direction; a few only are oblique. The *longitudinal* fibres are the most superficial; they are continuous with the longitudinal fibres of the œsophagus, and spread out over the stomach. The *circular* come next: they are well marked about the middle of the stomach, but most abundant at the pylorus where they form a powerful sphincter muscle. The *oblique* fibres are nearest the mucous membrane; they are scattered over the sides of the stomach. Examined with the high powers of a microscope, we find that they all belong to the involuntary or non-striped variety. The same observation applies to the muscular coat of the intestines generally.

[At the pylorus we find a valve, sometimes single, sometimes double, formed from the coats of the stomach, chiefly the mucous.]

When the stomach is laid open, observe that the *mucous membrane* is of a pale color, and gathered into longitudinal folds; these disappear when the stomach is full. If a piece of the membrane be cut away, you see beneath it a distinct stratum of fibro-cellular tissue, sometimes called the "*submucous coat*." The object of it is to permit the muscular and mucous coats to move freely on each other, and to serve as a bed, in which the blood-vessels ramify minutely before they enter the mucous membrane. This observation applies not

only to mucous membrane of the stomach, but to that of the alimentary canal generally.

What is the apparatus for the secretion of the gastric juice? If a portion of well-injected mucous membrane, taken from the stomach of any mammalian animal, be examined with one of the low powers of a microscope, you observe that the capillary blood-vessels present a hexagonal arrangement, and that they map the surface into little pits, giving it rather a honey-comb appearance. The pits are, on an average, $\frac{1}{100}$ th part of an inch in diameter. At the bottom of them you see a number of minute pores. These are the orifices of the gastric tubes. In a perpendicular section, we find that the tubes are arranged in parallel bundles at right angles to the surface, and that they terminate in blind sacculated ends set in the submucous tissue. The entire thickness of the mucous membrane is made up of these tubular glands. It is presumed that during digestion, they generate a number of cells containing the gastric juice. As fast as formed, the cells pass into the stomach, discharge their contents, and disappear.

While examining the arrangement of the tubes of the stomach, you should observe how richly they are supplied with blood. The arteries form a stratum of minute inosculation in the submucous tissue, in which the bottoms of the tubes are set: from this stratum the vessels run up between the tubes to the surface of the stomach, where they again inosculate, and form the hexagonal spaces above alluded to.

The mucous membrane of the stomach, and also of the interior of the tubes, is lined by a columnar epithelium. It is exceedingly thin and delicate, and can only be seen in the stomach of an animal recently killed.

Small Intestines.—The small intestines, consisting of the duodenum, jejunum, and ileum, form a tube from sixteen to twenty-eight feet in length, according to the height of the individual. As regards their external character, remark that the duodenum and jejunum are much more vascular than the ileum, and that they may feel much thicker in consequence of the peculiar arrangement of their mucous membrane. The reason of this difference is, that the upper part of the small intestine is more concerned in digestion than the lower. Their peritoneal and muscular coats are the same throughout.

The muscular coat consists of an outer longitudinal and an inner circular layer. The mucous coat requires especial notice.

When the small intestines are cut open from the upper end, the first thing that attracts notice is that the mucous membrane is arranged in close folds or plaits, technically called "*valvulae conniventes*." Now these differ from other folds in the alimentary canal, *e. g.* in the œsophagus and stomach, in that they are not obliterated when the tube is distended. Each fold extends about one half or two-thirds round the intestines; but they are not all of equal size. Observe that they do not commence at the pylorus, but immediately below the openings of the biliary and pancreatic ducts, and that they are the most largely developed in the duodenum and the upper part of the jejunum. Below this part of the tube they gradually decrease in size, and become less closely packed, till they finally disappear near the middle of the ileum. The use of the *valvulae conniventes* is to increase the extent of surface for the absorption of the chyle, and also for secretion.

Next examine what are called the "*villi*" of the small intestine, and afterwards the different kind of glands with which it is provided.

If a portion of the small intestine be carefully washed and placed in water, the surface of the mucous membrane appears like the soft fur or pile upon velvet. This appearance is produced by the villi. Now these are nothing more than extremely vascular projections of the mucous membrane, about a fourth of a line in length, and so close to each other that a square line contains from forty to fifty of them. Their size, however, and their number, bear a direct ratio to that of the *valvulae conniventes*. Their structure is exceedingly simple. In a well injected specimen, you will find that each is furnished with an artery which forms a beautiful network of anastomoses all over it, and then returns its blood by a single vein. Each, too, contains in its interior a lacteal or absorbing vessel, which commences, not, as was formerly supposed, by an open mouth, but by a closed end near the summit of the villus. Lastly, they are covered by a layer of cylindrical epithelium, like the rest of the intestinal mucous membrane. It is generally believed that these cylindrical cells are the agents

in the absorption of the chyle, and that they possess the power of selection.

There are three kinds of *glands** in the small intestines named after their respective discoverers, the glands of Lieberkühn, Brunn, and Peyer. The first are distributed over the whole tract of the mucous membrane; the last two over particular parts.

The *glands of Lieberkühn*† are minute tubes with blind ends, very thickly distributed over both the small and the large intestines. With a microscope,—for they are invisible to the naked eye,—we see their orifices between the villi, like so many minute dots, as if the surface had been pricked with needles. In a vertical section we should observe their depth, and that they are lined by a columnar epithelium.

The *glands of Brunn*‡ are found only in the duodenum. They are just visible to the naked eye, and may be seen to the best advantage by removing the muscular coat. Their structure exactly resembles the pancreas, on a diminutive scale.

The *glands of Peyer*§ abound most in the ileum. Their chief peculiarity is, that they are arranged in groups on that part of the intestine most distant from the attachment of the mesentery. These groups are from one to two inches long, of an oval form. Observe that they increase in size and number as we approach the lower part of the ileum. If a group be examined by dissecting away the muscular coat, you find that the glands are imbedded in the submucous tissue, that they are rather opaque, about three-fourths of a line in diameter, and in shape not unlike a Florence flask. They have no discoverable orifices. When full and ripe they come to the surface, burst, discharge their contents, and die—for what purpose is not precisely known. These are the glands which are so liable to be ulcerated in fever.

But the glands of Peyer are not confined to the ileum.

* A satisfactory examination of the intestinal glands can be made only in specimens quite recent, taking from young persons who have died suddenly, or from a rapidly fatal disease.

† J. N. Lieberkühn, *Diss. de fabric. et actione villorum intestin. tenuium*, 1782.

‡ J. C. Brunn, *Gland. duoden. seu pancreas secundarium*, 1715.

§ Peyer, *De glandulis intestinorum*, 1682. But these glands were described by our countryman, Nehemiah Grew, in 1681.

Glands in all respects like them, except that they are *solitary*, are found scattered in all parts of the small intestine, also in the large.

[We occasionally meet with appendices from the small intestine whose calibre is nearly as great as that of the intestine itself.]

Large Intestine.—The principal external characters of the large intestine are, that it is pouched or sacculated, and that it has attached to it little pendulous portions of fat covered by peritoneum, called "*appendices epiploicæ*." Now the pouches are obviously produced by a shortening of the longitudinal muscular fibres, and by their being collected into three bands, about half an inch wide, nearly equidistant from each other. One of these bands corresponds with the attached part of the circumference of the bowel; another with the front part; a third with what may be called its concavity. If at any given part these bands be divided, the pouches immediately disappear.

The rectum differs from the rest of the large intestine in that its longitudinal muscular fibres are not collected into bands, but distributed equally over its whole circumference. Moreover, both the longitudinal and circular fibres are of considerable strength, like those of the œsophagus, as one might expect from the particular functions which these parts of the alimentary canal have to perform. Lastly, for one inch and a half, or thereabouts, above the anus, the circular fibres are remarkably developed, and constitute the internal "*sphincter ani*."

The mucous membrane of the large intestine differs materially from that of the small. It is altogether more simple. There are no *valvulæ conniventes* or villi; but there are plenty of Lieberkühn's glands; plenty also of the flask-like solitary glands* like those in the ileum. Peyer well describes these glands as studding the mucous membrane, "*tanquam stellæ firmamenti*." As regards the arrangement of the blood-vessels, it deserves to be remarked that a piece of well-injected mucous membrane from the large intestine presents the same


* The solitary glands are more abundant in the cæcum and in the appendix vermiformis than in any other part of the alimentary canal.

hexagonal arrangement on the surface as that of the stomach. That the mucous membrane of the large bowel may be temporarily used as a substitute for the stomach, is proved by the fact of persons having been nourished for six weeks or more, solely by injections.

Ileo-Cæcal Valve.—At the junction of the small with the large intestine the mucous membrane is folded so as to form a valve: but it is not a perfect one, as is proved by pouring water into the large intestine, or by the occasional vomiting of injections. The arrangement of the valve is best examined in a dried preparation. You observe that the opening is a transverse fissure like a button-hole; and that the two flaps are arranged like an upper and a lower eyelid. The flaps of the valve consist of mucous membrane and the circular fibres of the ileum. The longitudinal fibres of the ileum are continued directly on to the cæcum: if these be divided, the ileum can be drawn out and the valve disappears.

Fig. 90.

1. Ileum.
2. Cæcum or caput coli.
3. Appendix vermiformis.



SECTION THROUGH THE JUNCTION OF THE LARGE AND SMALL INTESTINE
TO SHOW THE ILEO-CÆCAL VALVE.

Folds in the Rectum.—In many subjects we observe that transverse or oblique folds of the mucous membrane project into the rectum. We cannot see them to advantage unless the bowel be hardened by alcohol in its natural position.

Three, more prominent than the rest, and half an inch, or thereabouts, in width, were first pointed out by Mr. Houston.* One projects from the upper part of the rectum, opposite the prostate gland; another is situated higher up, on the side of the bowel; while the third is still higher. When thickened or ulcerated, these folds are apt to occasion great pain and obstruction in defæcation.

THE DISSECTION OF THE THIGH.

AN incision should be made along the groin from the spine of the ilium to that of the pubes; another, from the middle of the first, perpendicularly downwards for about six inches.

Superficial Fascia.—Reflecting the skin, we expose the sub-cutaneous adipose and connective tissue, called the “superficial fascia.” Its thickness varies according to the condition of the body. It is described as being divisible into layers. The truth is, that any one who is dexterous with the scalpel can make as many layers as he pleases. Your object must be to have a clear view of the superficial absorbent glands and the cutaneous vessels which are imbedded in this fascia, and ramify in three directions. One ascends over the surface of the abdomen, another towards the pubes, a third towards the ilium; they are named the superficial *epigastric*, the superficial *pubic*, and the superficial *circumflexa ilii* arteries (fig. 91). All three come from the femoral. Each is accompanied by one, sometimes by two veins, which empty themselves, either directly into the femoral vein, or into the great cutaneous vein of the thigh, called the “saphena.”


Femoral Absorbent Glands.—These are easily recognized by their reddish-brown color. Most of them are situated along the outer side of the saphena. They vary in number, and are of an oval form, with the long axis vertical. In a well-injected subject we observe how freely they are supplied with blood-vessels; hence the rapidity with which they enlarge when inflamed. They transmit the absorbents from

* Dublin Hospital Reports, vol. v.

nearly all parts of the lower extremity. They also receive some of the absorbents from the scrotum. This explains why, in some cases of chimney-sweeper's cancer of the scrotum, the femoral, as well as the inguinal glands, become enlarged.

The absorbent vessels which pass in and out of the glands

Fig. 91.



SUPERFICIAL VESSELS AND GLANDS OF THE GROIN. SAPHENOUS OPENING WITH THE CRIBRIFORM FASCIA.

- | | |
|--|------------------------------|
| 1. Saphenous opening of the fascia lata. | 5. Superficial pubic a. |
| 2. Saphena vein. | 6. External abdominal ring. |
| 3. Superficial epigastric a. | 7. Fascia lata of the thigh. |
| 4. Superficial circumflexa ilii a. | |

escape observation, unless specially sought after. In dropsical bodies they become distended, and appear like transparent tubes. They all pass through the femoral ring into the abdomen, and eventually empty themselves into the thoracic duct.

The glands mentioned in the preceding paragraph are all superficial. There are others more deeply seated close to the

great vessels of the thigh : these are much smaller and sometimes cannot be found.

Respecting the *superficial epigastric*, *pudic*,* and *circumflexa ilii* arteries, we have only to observe that their particular ramifications vary in each case, though their general distribution is nearly the same in all. In a surgical point of view they are of importance, since they are exposed to injury in opening abscesses in these parts. The *pudic* is liable to be cut in the operation for femoral hernia ; also in the operation for phimosis, since it runs along the penis to supply the prepuce. Arising, as they all do, directly from so large an artery as the femoral, we cannot wonder if they sometimes bleed profusely ; for it is an admitted fact, that when even a small branch, coming directly from a principal artery, is divided near its origin, it will sometimes pour out as much blood as if an opening were punched out of the trunk as large as the area of the divided branch.†

Saphena Vein.—This is the chief subcutaneous vein of the lower limb. Its roots arising on the inner side of the foot, unite into a single trunk, which ascends *over* the inner ankle, along the inner side of the leg and knee, and then along the inner and front part of the thigh, where you see it passing through an opening in the “fascia lata” to join the femoral vein immediately below the crural arch (fig. 91). In this remarkably long course it receives several tributary veins some of which are often as large as itself ; and, just before its termination, it is joined by the superficial veins of the groin already alluded to. Like all subcutaneous veins, it is provided at intervals with valves to support the column of the blood ; when these, from any cause become inefficient, the veins become enlarged and varicose.

Postponing, for the present, the general muscular fascia of

* The pudic branch referred to in the text is quite superficial. There is another, which runs for some distance beneath the muscular fascia covering the pectineus, and supplies that muscle and the scrotum or the labium pudendi.

† Mr. Liston had occasion to tie the external iliac artery for a supposed injury (by a pistol ball) to the femoral. It was discovered after the death of the patient, that the ball had injured only one of the superficial branches of the femoral, about an inch from its origin. See his paper in the Med. Chir. Trans. vol. xxix. 1846.

the thigh, called "*fascia lata*," let us now examine the opening in it which transmits the saphena vein.

Saphenous Opening in the Fascia Lata.—Immediately below the crural arch is an opening in the muscular fascia of the thigh, which, as it gives passage to the saphena vein, is appropriately called "*saphenous*." Now since this opening is one of the parts concerned in the anatomy of femoral hernia, the mode in which it is formed, its shape, dimensions, borders,—in short, everything connected with it,—is described by anatomical writers with very minute precision. He who dissects these parts for the first time can hardly expect to make them out satisfactorily, still less to make his dissection tally with the description usually given.

Understand that the edge of the saphenous opening does not naturally exist as a defined margin, until the fascia which covers the opening and is blended with the margin has been removed. The term "*cribriform*" has been given to this fascia, because it is riddled with holes for the passage of the superficial vessels and absorbents: of this we will say more anon. At present let us confine our attention to the saphenous opening itself.

Supposing the cribriform fascia to have been removed so as to display the saphenous opening in its simplest form, the opening would appear as represented in fig. 92.

We observe that it is situated just below the crural arch, not far from the pubes; that it is oval with the long axis vertical, and about one and a half or two inches long. In respect to its borders, we find that on the inner or pubic side it is not at all defined, for here the muscular fascia ascends under the femoral vessels, and is continuous with the iliac fascia of the pelvis.* But the outer or iliac border is defined clearly enough. This lies in front of the femoral vessels, and is crescent-shaped, with the concavity towards the pubes. The lower horn of the crescent curves under the saphena vein, and is lost in the fascia on the inner side of the opening. The upper horn† arches over the femoral vein, and is continued

* On the inner side of the femoral vessels the pubic portion of the fascia is attached to the linea ilio-pectinea.

† This upper horn is sometimes called "Hey's ligament," after the surgeon

uninterruptedly into "Gimbernat's ligament," or into that part of the crural arch which is inserted into the linea ilio-pectinea. The upper horn deserves especial attention, because it forms the upper boundary of the aperture through which a femoral hernia takes place; and having a great deal to do with the constriction of the rupture, must be divided for its relief.

Fig. 92.




DIAGRAM OF THE FEMORAL RING.

(The arrow is introduced into the femoral ring.)

This may be easily ascertained by introducing the little finger under the crural arch, on the inner side of the femoral vein,—in other words into the femoral ring (see the arrow in the diagram). See how much the upper horn of the crescent would girt the neck of a hernia, and that its tension is greatly influenced by the position of the limb; for if the thigh be bent and brought over to the other side, the tension of all the parts concerned is materially lessened. This explains the position in which a surgeon places the leg before he attempts to reduce a rupture.*

who first drew attention to it: "Observations in Surgery," by W. Hey, F. R. S. London, 1810.

* We must always bear in mind, that though the crural arch and the adjacent fasciæ have received particular names, they are not, on that account, distinct and separate, but all are intimately connected, and portions merely of one continuous expansion. Thus all the parts are kept in a condition of mutual tension, which depends very much on the position of the thigh.

Cribriform Fascia.—To return to the cribriform fascia. This is simply a thin covering over the saphenous opening. It is prolonged from the outer edge of the opening over the sheath of the femoral vessels, and adheres on the inner side to the fascia lata, over the pectineus muscle. As before observed, it is riddled with holes for the passage of the superficial vessels and absorbents of the groin. Some writers describe this fascia as only a portion of the "superficial fascia;" but I think the more correct view is to consider it as a thin prolongation of the fascia lata itself across the opening. Its chief surgical importance is derived from the fact that it forms one of the coverings of a femoral hernia.

ANATOMY OF THE PARTS CONCERNED IN FEMORAL HERNIA.

THE anatomy of the parts concerned in femoral hernia cannot be thoroughly understood without the assistance of special dissections; indeed, one cannot reasonably expect to make one's self master of the subject, if it be taken in hand merely as something to be investigated by the way in the dissection of the thigh. The following demonstration, therefore, takes for granted that we have the opportunity of seeing the parts, not only on their femoral, but also on their abdominal side.

We propose to treat the different parts of the subject in the following order:—

- a. The formation of the crural arch.
- b. The arrangement of the parts which pass under the arch.
- c. The femoral aperture or ring.
- d. The so-called sheath of the vessels.
- e. The practical application of the subject.

Crural Arch.—The lower border of the aponeurosis of the external oblique muscle extends from the spine of the ilium to the spine of the pubes, and forms, over the bony excavation beneath, what is called the "crural arch." (It is marked by the dark line in fig. 93.) A glance at the relative position of its attachments is sufficient to show that the direction of the arch is oblique; besides which, in consequence of its intimate connection with the muscular fascia of the thigh, the line of

the arch describes a gentle curve with the convexity downwards. The pubic insertion of the arch is a point which we must thoroughly understand. Not only is the arch attached to the spine of the pubes, but also for some distance along the linea ilio-pectinea (fig. 93). This additional attachment, called after its discoverer, *Gimbernats ligament*,* is of material importance, for it is the seat of the stricture in hernia.

Gimbernats Ligament.—The best view of Gimbernats ligament is obtained from within the abdomen. All that is necessary is to remove the peritoneum. You then see that it is triangular, with the apex at the pubes; that the base, directed outwards, is a little curved. Its breadth is from three-quarters of an inch to one inch; but it varies in different subjects, and is always broader in the male than in the female; this is one among many reasons why femoral hernia is less frequent in the male.


Put your finger into the femoral ring and make yourself familiar with the sharp and wiry feel of this ligament: observe, especially, that, as the body lies on the table, the plane of the ligament is perpendicular, and therefore that it *recedes from the surface*.

Arrangement of the Parts which Pass Under the Arch.—The crural arch transmits from the abdomen into the thigh (proceeding in order from the outer side) the following objects, shown in fig. 93: 1. The external cutaneous nerve; 2. The iliacus and psoas muscles, with the anterior crural nerve between them; 3. The femoral artery and vein with the crural branch of the genito-crural nerve. These muscles and vessels completely fill up all that part of the arch which belongs to them. Still there is, on the inner side of the vein, a comparatively vacant space left for the passage of the absorbents; this is called the femoral aperture or ring. The muscles are separated from the vessels by a strong vertical fibrous partition passing from the arch to the bone, which is nothing more than a continuation of the sheath of the psoas. The artery, too, is separated from the vein by a similar, although a much weaker partition, and there is a third close to the inner side of the vein. These three partitions not only keep all the

* Don Antonio de Gimbernats was a Spanish surgeon, who published, in 1793, "New Method of Operating for the Femoral Hernia," Madrid.

parts in their right place, but confine the arch down to the bone, and prevent its being uplifted by any protrusion between it and the muscles and vessels. This, coupled with the

Fig. 93.



POSITION OF PARTS UNDER THE CRURAL ARCH (VERTICAL SECTION).

close attachment of the iliac fascia to the crural arch, explains why a femoral hernia cannot take place in any other situation than on the inner side of the femoral vein.*

Femoral Ring.—The hollow under the crural arch is completely occupied, except for a small space on the inner side of the vein. This weak part is called the “femoral ring;” through this a femoral hernia protrudes. Now in order to form a correct idea of the size of this opening, you should remove the peritoneum, and introduce the end of the little finger into it from the abdomen. This can be done without much force. You find that the finger is girt pretty tightly on all sides: in front is the crural arch; behind is the bone; on the outer side is the femoral vein; on the inner side is the thin wiry edge of Gimbernat’s ligament. Such are the boundaries of the gap which the finger makes. Such are the

* If the partitions from any cause yield, or become slack, then a rupture may descend in front of the vessels, or even (though this is extremely rare) on the outer side of the artery.

boundaries of the femoral ring. Remember, however, that, in the undisturbed condition of the parts, there is no gap; there is only a weak place here, which, when a hernia escapes through it, feels very much like a ring; hence the name "femoral ring."*

The femoral ring is surrounded on all sides by unyielding structures. This accounts for the little benefit afforded by the warm bath in cases of strangulation. Mr. Lawrence says he never saw a strangulated femoral hernia where the warm bath was of any avail.

Sheath of the Femoral Vessels.—The femoral vessels do not descend naked beneath the crural arch, but come down inclosed in a membranous sheath. This sheath appears to be derived immediately from the arch itself, but it is not so: it is a continuation of the fascia transversalis of the abdomen. Now this continuation, uniting with the muscular fascia behind the femoral vessels, forms a funnel, with the wide part uppermost, into which the femoral vessels enter. This is what is meant by the funnel-shaped sheath of the femoral vessels.

[This sheath, which is derived from the common internal fascia, has a septum which divides it into two compartments; the outer portion contains the vessels, while the inner compartment, in the normal state, is empty. It is within this latter portion that the hernia descends, and not, as stated by some authors, within the sheath of the vessels. The prolongation of this fascia descends along the vessels for about an inch and a half.]

To examine this sheath satisfactorily, it is necessary to reflect from its attachment to the crural arch, the upper horn of the saphenous opening, as shown in fig. 94. By this proceeding, you see the fascia transversalis coming down over the femoral vessels, and forming the front part of their sheath. The hind part of the sheath, remember, is formed by the pubic portion of the fascia lata, which runs up behind the vessels to join the fascia iliaca. If asked how the sheath de-

* The femoral ring is naturally occupied by a little fat and cellular membrane, by absorbent vessels, and often by a small absorbent gland. But we have never met with anything deserving the name of a "diaphragm" or membranous septum, such as is described by Cloquet as the "septum crurale."

scends, I should say that, about the lower horn of the saphenous opening, it is gradually lost upon the proper investment of the femoral vessels.

Fig. 94.




DIAGRAM OF THE SHEATH OF THE FEMORAL VESSELS.

Examine for yourselves the texture and the degree of resistance of the sheath by introducing the finger into it from within the abdomen. Try to make an artificial hernia. You will find the sheath offers considerable resistance. If violence be used, the sheath will either give way, like an elastic membrane, or the finger will force a passage through one of the small openings which exist in the sheath for the transmission of the femoral absorbents.

Practically, the sheath is important for many reasons:—1. A femoral hernia descends inside it. 2. It forms, therefore, one of the coverings (*fascia propria*) of the hernia. 3. It contains in its substance bands of fibres running in the same direction as the crural arch, but quite independent of it, as shown in fig. 94; these bands lie over the neck of the sac, and are often the seat of the stricture; it is therefore necessary to divide them before the intestine can be returned.

Practical Application of the Subject.—From what has been said, you ought now to understand—1, at what point a femoral hernia escapes from the abdomen; 2, the course

which it takes, and its relations to the surrounding parts ; 3, the proper mode of attempting the reduction ; 4, the structure and arrangement of its coverings ; and lastly, where the stricture will probably be situated.

The hernia escapes from the abdomen through the femoral ring,—that is, under the weak part of the crural arch, between the femoral vein and Gimbernat's ligament. Here is the mouth of the hernial sac, or that part of it which communicates with the abdomen. It descends for a short distance nearly perpendicularly, and projects as a small tumor in front of the pectineus muscle. Its progress downwards, however, is soon arrested by the very close adhesion of the subcutaneous structures to the lower margin of the saphenous opening. Consequently, if the hernia increase in size, it must rise over the crural arch, where the subcutaneous tissue offers less resistance ; and the bulk of the hernia extends outwards towards the ileum, assuming more or less of an oblong form, with the long axis parallel to the crural arch. Since, then, the body of the hernia forms a very acute angle with the neck, the right mode of attempting its reduction is, to make the hernia retrace its steps,—that is, to draw it first down from the groin, and then to make pressure on it backwards in the direction of the femoral ring.

Now what are its *coverings*? The hernia protrudes before it the peritoneum, technically called the hernial sac.* It next pushes before it the sheath of the femoral vessels, which forms an investment called the "*fascia propria*." In front of this is the cribriform fascia. Lastly, there is the subcutaneous tissue and skin.

[We would enumerate the coverings as follow, from without inwards :—1, integument ; 2, fascia superficialis communis (cribriform fascia) ; 3, fascia lata ; 4, fascia communis interna (fascia transversalis—fascia propria) ; 5, peritoneum. These form the coverings of the neck of the tumor ; the body has but four coverings, it lying *upon*, not under the fascia lata ; for that portion of the fascia that passes over the neck is

* In some cases the fascia propria so much resembles the hernial sac, that it is not easy to distinguish between them. Generally speaking, they are separated by a small quantity of fat.

nothing but the superior border of the saphenous opening, or, as called by some, the femoral ring.]

The *seat of stricture* is at the femoral ring, and the position of the neighboring blood-vessels indicates that the proper direction in which to divide the stricture is, either directly inwards, through Gibernat's ligament, as recommended by Mr. Lawrence, or upwards through Hey's ligament, as recommended by Sir A. Cooper. There is no risk of wounding an artery, supposing the vessels to take their ordinary course. But it occasionally happens (as observed p. 365), that the obturator artery runs round the inner side of the femoral ring; in such a case, the neck of the sac would be encircled by a large blood-vessel.* From the examination of two hundred bodies, I conclude that the chances are about seventy to one against this unfavorable distribution. But the possibility of it has given rise to *this rule* in practice,—not to cut deeply in any one place through the stricture, but rather to notch it in several. By this proceeding we are much less likely to wound the abnormal artery, because it does not run at the base of Gimbernat's ligament, but about a line and a half from the margin of it.

Such is a brief outline of the anatomy of the parts concerned in a femoral hernia. The normal anatomy in each case being nearly the same, one might suppose that all operations for the relief of this kind of hernia would be straightforward and pretty much alike; but this is very far from being the case; indeed, most experienced surgeons will agree that they never take the knife in hand without the expectation of meeting with some peculiarity.

To proceed with the dissection, the incision should be prolonged down the thigh, over the knee, to the tubercle of the tibia. The skin should be reflected, so that the subcutaneous tissue over the whole front of the thigh be fairly exposed; our present object being to examine the cutaneous veins and nerves.

* The Museum of St. Bartholomew's Hospital contains two examples of double femoral hernia in the male, with the obturator arising on each side from the epigastric. In three out of the four herniæ the obturator runs on the inner side of the mouth of the sac. See prep. 55, 69, Series 17.

Cutaneous Veins and Nerves.—The course of the saphena vein, its tributary branches, and the confluence of veins at the saphenous opening, have been described at p. 418. I would merely observe here, that all the larger cutaneous veins of the thigh, and the same obtains of the leg generally, are so situated as to be out of the way of harm. Respecting the cutaneous nerves, a general account is sufficient, since they are not exactly alike in any two subjects. They are usually divided, according to their situation, into *external*, *middle*, and *internal*. All directly, or indirectly, proceed from the lumbar plexus, and may be seen coming through the muscular fascia.

a. The *external cutaneous* nerve is a distinct branch of the lumbar plexus (p. 336). It enters the thigh beneath the crural arch, near the anterior spine of the ilium, and can be traced down the outer side of the thigh to the knee, giving off numerous branches, which subdivide to supply the skin.

b. The *middle cutaneous* nerves, two or more in number, are derived from the anterior crural (which itself is a branch of the lumbar plexus). They perforate the sartorius four or five inches below the crural arch, and descend along the front of the thigh to the knee, distributing branches on either side (p. 431).

c. The *internal cutaneous* nerves, two or more, also branches of the anterior crural, perforate the fascia, one about the middle, another about the lower third of the thigh, and supply the skin on the inner side.*

By careful dissection, you will find that these cutaneous nerves communicate here and there with each other, especially over the patella.†

Fascia Lata, or Muscular Fascia.—Remove the subcutaneous structure to examine the muscular fascia, or “fascia lata” of the thigh. The purpose of this fascia is to clothe the muscles of the thigh collectively, and to form separate sheaths for each; so that it not only packs them together, but maintains each in its proper position. A knowledge of these sheaths is practically important, because they interfere with the pro-

* Besides these, there is the little crural branch of the genito-crural nerve (p. 336), which perforates the fascia about two inches below the crural arch. This is seldom found except by an experienced dissector.

† Some authors speak of a “patellar plexus” of nerves.


gress of deep-seated matter towards the surface, and cause it to burrow in this or that direction according to the part in which it forms.

The fascia is not of equal strength all round the thigh. As one might expect, it is strongest in the most exposed parts; *e. g.* it is comparatively thin and transparent on the inner side, but exceedingly thick and strong all down the outer side: here, indeed, it has the appearance of a dense expanded tendon, strapping down the vastus externus muscle; and it certainly performs the office of a tendon, for we shall presently see that it gives insertion to two powerful muscles,—namely, the tensor fasciæ femoris, and the glutæus maximus (fig. 95).

But what are its attachments? In a general way, it may be said that it is attached to the margin of the bones which constitute the framework of the lower extremity. Beginning from above, we could, if the parts were exposed, trace its attachment along the crest of the ilium, thence along the crural arch to the body of the pubes, and so on down the rami of the pubes and ischium. Proceeding down the thigh, we should see that it penetrates, on either side of the limb, to the linea aspera, forming what are called the “intermuscular septa,” which separate the extensor from the flexor muscles. Below we trace it all round the knee-joint, where it is particularly strong, especially on the outer side, and find that it is attached to the head of the tibia and the fibula. Here let us leave it for the present.

The next stage of the dissection consists in removing the muscular fascia from the front of the thigh, without disturbing the muscles from their relative position. The mass of muscles on the inner side of the thigh consists of the adductors; that in the middle, of the extensors; the long muscle

Fig. 95.



FASCIA ON THE OUTSIDE OF THE THIGH.

1. Tensor fasciæ femoris.
2. Glutæus maximus.
3. Lower fibres of ditto.
4. Fascia lata.

which crosses obliquely over the front is the sartorius. This we will take first, being a sort of landmark to the rest.

Sartorius.—This muscle arises from the anterior superior spine of the ilium and the ridge below. It passes obliquely, like a strap, over the front of the thigh towards the inner side, and then descends almost perpendicularly on the inner side of the knee, where it terminates in a tendon which expands and is inserted into the inner and front part of the tibia just below the tubercle. The tendon appears all the wider on account of its broad connection with the fascia of the leg. A large bursa* is interposed between the tendon and the internal lateral ligament. The chief *action* of the muscle is to fix the pelvis steadily on the thigh. It also crosses one leg over the other, as tailors are in the habit of doing when sitting at work.†

[Meckel reports a case where this muscle was wanting. It may have a middle tendon, and by some is said to be a broader muscle in the negro.]

Scarpa's Triangle.—The sartorius forms with the adductor longus the two sides of a triangle, which has the crural arch for its base. This triangle is called "Scarpa's," in compliment to the anatomist who first tied the femoral artery in it, for popliteal aneurism. You should carefully display the contents of this important surgical space, and study well their relative position. I have attempted to give some idea of them in fig. 96. It contains all the parts which come under the crural arch, namely: the external cutaneous nerve; the psoas and iliacus; the anterior crural nerve and its divisions, especially the long saphenous; the femoral artery with its great branch "the profunda,"‡ which runs down behind it and gives off the "internal" and "external circumflex;" the femoral vein, joined by the profunda vein and internal saphena; the pectineus muscle.


* In persons, females especially, who are in the habit of riding, this bursa sometimes becomes enlarged.

† Hence the name given to it by Spigelius (*De corporis hum. fabric.*), "*Quem ego sartorium musculum vocare soleo, quod sartores eo maximè utuntur, dum crus cruri inter consuendum imponunt.*"

‡ In fig. 96, "the profunda" is drawn more on the outside of the femoral than it is in nature; the internal circumflex cannot be seen because it sinks down between the pectineus and psoas.

The most important practical point about the muscle is, that it overlaps the femoral artery in the middle of the thigh; and the inner border of the muscle is the guide to the vessel in the situation where it is usually tied for a popliteal aneurism.

Fig. 96.



What is the exact spot at which the sartorius reaches the femoral artery? I have examined many bodies in reference to this point, and find that there is no certainty about it. The distance from the crural arch at which the artery and the muscle meet, varies from one and a half to four and a half inches. The best way to find the inner border of the sartorius during life is to make the patient put the muscle into action.

Gracilis.—Passing in the next place to the adductors on the inner side of the thigh, we should examine another long flat muscle, called the gracilis. To dissect it properly, it should be put on the stretch by separating one thigh from the other. It arises by a broad ribbon-like tendon from the pubes close to the symphysis, and from the border of the pubic arch nearly as low as the tuber ischii. The muscle de-

scends perpendicularly on the inner side of the thigh, and terminates on a tendon which spreads out and is inserted into the inner side of the tibia below the tubercle; in fact, just like the sartorius, only below it, so that the one overlaps the other. Both tendons play over the internal lateral ligament of the knee-joint, and there is a bursa common to both. This muscle assists in fixing the pelvis, and, in adducting the thigh, it also helps to bend the knee.

Adductor Muscles.—A strong group of muscles extends along the inner side of the thigh, from the pelvis to the femur. Their chief action is to co-operate in balancing the pelvis steadily on the thigh,—as, for instance, in standing on one leg; or, if the fixed point be reversed, they can draw the thighs together—for example, in riding. Physiologically, they might be considered as one, for they are supplied by the same nerve,—namely, the obturator; but anatomically they are divided into three—the adductor longus, brevis, and magnus. There is also the “pectineus.” Since they lie one above the other, you should reflect them from their origin and examine each in succession, taking care not to divide the branches of the obturator nerve which supply them.

Adductor Longus.—This muscle arises by a tendon from the front of the pubes just below its spine, and, gradually becoming broader, is inserted into the middle third of the linea aspera of the femur. It forms with the sartorius a triangular space, in which are contained the several objects described (p. 430).

Pectineus.—This muscle lies close to, and on the same plane with the adductor longus (p. 430). It arises from the triangular surface of the pubes in front of the linea ilio-pectinea, and is inserted into the ridge leading from the lesser trochanter to the linea aspera.

The adductor longus and pectineus must now be reflected from their origins. Under the pectineus you will see the “obturator nerve,” which supplies all the adductors, except the pectineus, which is supplied by the anterior crural. It leaves the pelvis through the upper part of the obturator foramen, and soon divides into an anterior and posterior branch; the anterior runs in front of the adductor brevis and supplies it, as well as the adductor longus and the gracilis; the posterior

runs behind the adductor brevis, and supplies the obturator externus and the adductor magnus.

Adductor Brevis.—This muscle arises from the front surface of the pubes near the symphysis, widens as it descends, and is inserted into the upper third of the linea aspera. By reflecting it from its origin we expose the following :*—

Adductor Magnus.—This muscle arises from the lower two-thirds of the pubic arch, and from the tuberosity of the ischium. Its fibres spread out, and are inserted, behind the other adductors, into the whole length of the linea aspera; also into the ridge leading from it to the inner condyle. Observe that all the adductor muscles are inserted into the femur by flat tendons which are more or less connected with each other.

About the junction of the upper two-thirds with the lower third of the thigh, the femoral artery passes through an oval opening in the tendon of the adductor magnus.

Psoas Magnus and Iliacus Internus.—As these muscles have been fully described in the dissection of the abdomen (p. 327), it is unnecessary to repeat the description here.

Tensor Fasciæ Femoris.—This muscle is situated at the upper and outer part of the thigh. It arises from the crest of the ilium, close to the anterior spine, descends with a slight inclination backwards, and is inserted between two layers of the strong aponeurosis which is generally described as part of the fascia lata (p. 428). Its chief use is to fix the pelvis steadily on the thigh, and also to rotate the thigh inwards: in this last respect it co-operates with the anterior fibres of the glutæus medius, with which it is almost inseparably connected. Any one may convince himself of this by placing the hand on the hip and rotating the thigh inwards. There are no other muscles to do it, and both are supplied by the same nerve,—namely the superior glutæal.

In order to form an adequate idea of the strength, extent, and connections of the aponeurosis on the outer side of the thigh, we ought to separate it from the vastus externus muscle upon which it lies. There is no difficulty in doing so, for

* Beneath the adductor brevis, and just above the upper border of the adductor magnus, is seen the obturator externus. But we reserve the description of this muscle till the dissection of the external rotators of the thigh (p. 454).

it is united to the muscle by an abundance of loose cellular tissue.* With a little perseverance you can trace the aponeurosis to the linea aspera, the head of the tibia and the fibula. See how completely it protects the outer side of the knee joint.

Extensor Muscles.—The powerful muscles situated between the tensor fasciæ on the outer, and the sartorius on the inner side, are extensors of the leg: all are supplied by branches of the anterior crural nerve. One of them—the “rectus”—arises from the pelvis; the other—the “triceps”—arises from the shaft of the thigh bone by three portions, called the *cruræus*, the *vastus externus*, and *internus*.

Rectus.—This muscle arises from the pelvis by two strong tendons, which soon unite; one from the inferior spine of the ileum, the other from the rough surface of the ileum, just above the acetabulum. To see them, dissect between the sartorius and the tensor fasciæ, and in doing so be careful not to injure the branches of the external circumflex artery. The muscle descends perpendicularly down the front of the thigh, and is inserted into the common extensor tendon. Its structure is very beautiful. A tendon runs down the centre, and the muscular fibres are implanted on either side of it, like the vane on the shaft of a feather.

Triceps Extensor.—This mass of muscle surrounds the greater part of the shaft of the femur; therefore the whole extent of it cannot be seen without completely dissecting the thigh. It consists of an outer, middle, and inner portion, called respectively the *vastus externus*, the *cruræus*, and the *vastus internus*. The *vastus externus* arises by a very strong aponeurosis from the outer side of the base of the great trochanter, and from the outer lip of the linea aspera nearly down to the external condyle. The *cruræus* arises from the upper three-fourths of the front surface of the shaft. The *vastus internus* arises from the upper two-thirds of the inner surface of the shaft, and also from the inner lip of the linea aspera.

Common Extensor Tendon.—The tendon of the rectus,

* When the tissue becomes the seat of suppuration, the matter is apt to extend all down the outside of the thigh, not being able to make its way to the surface by reason of the dense fascia.

gradually expanding, becomes connected on its under surface with the tendon of the cruræus, and on either side with that of the vasti, and is firmly fixed into the upper part and sides of the patella.* From this bone the common extensor tendon, under the name of "*ligamentum patellæ*," descends over the front of the knee-joint, and is inserted into the tubercle of the tibia. But besides this, the lower fibres of the vasti terminate on a sheet-like tendon, which runs wide of the patella on either side, and is directly inserted into the sides of the head of the tibia, so that the knee is completely protected all round. The patella is a great sesamoid bone, interposed to facilitate the play of the tendon over the condyles of the femur; it not only materially protects the joint, but adds to the power of the extensor muscles, by increasing the angle at which the tendon is inserted into the tibia.

Action of the Extensor Muscles.—The extensor muscles of the thigh are among the most powerful in the body. A great power to extend the knee is one of the essential conditions of the erect attitude. Without it, how could we rise from the sitting to the upright position? When erect, how could we walk, run, or spring without it? The rectus, by taking origin from the pelvis, gains a double advantage; for it acts upon two joints simultaneously, bending the thigh while it extends the knee, as when we advance the leg in walking: it also contributes to balance the pelvis on the head of the thigh bone, and thus prevents the body from falling backwards. We cannot have a better proof of the collective power of the extensor muscles than when the patella is broken across by their contraction; an injury which sometimes happens when a man slipping backwards makes a violent effort to recover his balance.

To facilitate the play of the extensor tendon there are two

* A few of the deeper fibres of the cruræus are inserted into the fold of the synovial membrane of the knee-joint which rises above the patella. These are described as a distinct muscle, under the name of the "*sub-cruræus*." Their use is to raise the synovial membrane, so that it may not be injured by the play of the patella. Since the triceps is connected to the lower part of the shaft of the femur only by loose cellular tissue, there is nothing to prevent the distension of the synovial membrane, in cases of inflammation, to the extent of several inches above the patella.

bursæ. One is placed between the ligamentum patella and the tubercle of the tibia, the other between the cruræus and the lower part of the femur. This last is of very considerable size. In early life it is, as a rule, distinct from the synovial membrane of the knee-joint; but after a few years of friction we generally find a wide communication between them.

Bursa over the Patella.—We must here say a few words respecting the structures which cover the patella. The skin over it is exceedingly loose, and in the subcutaneous tissue is a bursa of considerable size. Since this bursa is apt to enlarge and inflame in females who are in the habit of kneeling, it is generally called “the housemaid’s bursa.” Observe that the bursa is not seated exactly on the top of the patella; but that it extends some way down the ligamentum patellæ; indeed, in some cases it is entirely confined to this ligament. This quite corresponds with the position of the tumor which the bursa occasions when enlarged. Generally speaking, in subjects brought for dissection, the wall of the bursa is more or less thickened, and its interior intersected by numerous fibrous cords, which are remnants of the original cellular structure altered by long continued friction. Again, the wall of the bursa does not always form a complete sac; sometimes there is a wide opening in it; this explains the rapidity with which inflammation, in some cases, extends from the bursa into the surrounding cellular tissue.

Below the bursa is a layer of fascia lata, and under this is a beautiful network of arteries. The immediate covering of the bone, or what may be called its periosteum, is a strong expansion derived from the extensor tendon. This is practically interesting for the following reason: in ordinary fractures of the patella from muscular action, the tendinous expansion over it is torn also; the ends of the bone gape widely, and never unite except by ligament. But in fractures from direct mechanical violence, the tendinous expansion, being entire, maintains the fragments in opposition, so that there is commonly a bony union.

Course and Relations of the Femoral Artery.—The femoral artery is a continuation of the external iliac. Passing beneath the crural arch at a point about midway between the spine of the ilium and the symphysis pubis, it descends nearly

perpendicularly along the front and inner side of the thigh. At the junction of the upper two-thirds with the lower third of the thigh, it goes through an opening in the tendon of the adductor magnus, and, entering the ham, takes the name of "popliteal." A line drawn from the point specified of the crural arch to the tubercle of the internal condyle would pretty nearly indicate the course of the artery. Its distance from the surface increases as it descends. In the next place, as to the muscles over which it lies. Immediately under, and for a short distance below the crural arch, it is supported by the inner border of the psoas; lower down it runs in front of the pectineus; still lower, in front of the adductor longus.

In the upper third of the thigh the artery is comparatively superficial, being covered only by the muscular fascia. About the middle third it is more deeply seated, and is covered by the sartorius: here, being situated among powerful muscles, it is protected from pressure by a tendinous canal, which we shall presently examine.

At the crural arch the femoral vein is on the inner side of the artery, but as the vein descends it gradually passes behind the artery. Both lie close together, and are inclosed in a common sheath.

[The femoral vein now and then takes a different course from the artery. Passing up from the popliteal space it may perforate the adductor magnus higher than usual, and uniting with the profunda vein, reach the artery in the groin. It also may be double in a part or the whole length of its course.]

The anterior crural nerve passes beneath the crural arch about $\frac{1}{4}$ th of an inch on the outer side of the artery, and soon divides into branches, some of which are cutaneous, but the greater number supply the extensor muscles. Those branches which more directly concern the artery, and are liable to be injured in the operation of tying it, are as follow:—1. The *long saphenous*. This descends close to the outer side of the artery, and enters the tendinous sheath with it, in the middle third of the thigh (p. 431). Here the nerve crosses over to the inner side of the artery, and leaves it just before it becomes popliteal. The nerve then runs in company with the anastomotica magna artery to the inner side of the knee, where we find it nearly superficial between the gracilis and

the sartorius. Finally, joining the saphena vein, it descends with it to the inner side of the foot. 2. Besides the long saphenous, there are one or two internal cutaneous branches, which cross over the sheath of the artery just where the sartorius begins to overlap it, and therefore in the situation where it is usually tied. They lie so close to the sheath that they would run great risk of being injured unless specially looked after (p. 431). Both supply the skin on the inner side of the thigh.*


Hunter's Canal.—You are already aware that in the middle third of the thigh the femoral artery is contained in a tendinous canal† under the sartorius, called “Hunter's canal.”

This canal at its upper part is rather indistinct; but it gradually becomes stronger towards the opening in the tendon of the triceps. Its boundaries are formed by the tendons of the muscles between which the artery runs. On the inner side are the tendons of the adductor longus and magnus; on the outer side is the tendon of the vastus internus; in front the canal is completed by an aponeurotic expansion thrown obliquely across from the adductors to the vastus internus; as shown in the annexed diagram. In a horizontal section the canal would appear triangular. The adaptation of this shape to the exigencies of the case is manifest when we reflect that the muscles keep the side of the triangle always tight, and thereby prevent any compression of the vessels.

Hunter's canal contains not only the femoral artery and vein, but also the internal saphenous nerve. The vein lies behind the artery, the nerve on its inner side, as shown in fig. 97.

[This artery may be double. Couch has given three ex-

Fig. 97.



SECTION THROUGH HUNTER'S CANAL.

1. Vastus internus.
2. Adductor longus.
3. Aponeurosis thrown across

* One of these nerves generally joins a branch of the obturator near the lower border of the tendon of the adductor magnus.

† Called “Hunter's canal,” because it was in this part of its course that John Hunter first tied the femoral artery for aneurism of the popliteal, in St. George's Hospital, A. D. 1785. The particulars of this interesting case are published in the Trans. for the improvement of Med. and Chir. Knowledge.

amples. Velpeau mentions a fourth; in this case it gave off the branches usually coming from the profunda, and afterwards descended below the knee. Sir C. Bell found the artery dividing into two equal branches, which united to form the popliteal; a similar case has also been reported. It has also bifurcated a little below Poupart's ligament into two vessels, and these became the anterior and posterior tibials. Portal gives a case where, after a high division, the two vessels formed two popliteals.]

After an examination of the course and relations of the femoral artery, it is plain that a ligature can be placed around it, in the upper third of the thigh, with comparative facility; not so easily in the middle third. The artery is tied for an aneurism of the popliteal, just where the sartorius begins to overlap it for three reasons:—1, it is more accessible; 2, the coats of the artery at this distance are less likely to be diseased; 3, the origin of the profunda is sufficiently far off to admit of the formation of a coagulum. An incision, beginning about three inches below the crural arch, should be made about three inches over the line of the artery. The muscular fascia should be divided over a director to the same extent. Then, by gently drawing aside the inner border of the sartorius, we discover the artery inclosed in its sheath with the vein. An opening should be made into the sheath just sufficient to admit the passage of the aneurismal needle, which should be turned round the artery from within outwards. The nerves to be avoided are—the long saphenous, which runs along the outer side of the artery, and the cutaneous nerves which cross obliquely over it.*

Having already traced the superficial branches of the femoral artery in the groin, namely, the superficial epigastric, pudic, and circumflexa ilii, (p. 418,) we pass on now to the profunda.

Profunda Artery and Branches.—The profunda is the chief branch of the femoral, and is the proper nutrient artery of the thigh. It arises from the outer and back part of the femoral, about one and a half or two inches below the crural

* The nerve (sometimes called the short saphenous) which supplies the vastus internus is so far out of the way, that it could not be included in the ligature, except by a very careless operator.

arch, and runs down behind the femoral till it reaches the tendon of the adductor longus ; here the profunda passes behind the adductor, and is finally lost in the hamstring muscles. In most subjects, the profunda, for a short distance after its origin, lies rather on the outer side of the femoral and on a deeper plane, over the psoas muscle : in this situation it might be mistaken for the femoral itself,—indeed, such an error has actually occurred in practice. It soon, however, gets behind the femoral, but is never in actual contact with it, for the two arteries are separated by their corresponding veins.*

[In one case which came under our notice (1865), the profunda was given off six inches below Poupart's ligament ; the circumflex here came off from the femoral.]

The branches of the profunda generally arise in the following order :—1, the internal circumflex ; 2, the external circumflex ; 3, the perforating.

The *internal circumflex* sinks deep into the thigh between the psoas and pectineus. Here you lose it ; but you will find it again in the direction of the back of the thigh, between the adductor magnus and the quadratus femoris. It supplies branches to the muscles in its neighborhood,—namely, to the pectineus, psoas, the adductors, and the quadratus femoris : and inosculates with the obturator and ischiatic arteries, as shown in the diagram, p. 441. It also sends branches to the hip-joint, and sometimes the one which runs through the notch in the acetabulum to the ligamentum teres.


The *external circumflex* runs transversely outwards beneath the sartorius and rectus, and subdivides into three sets of branches, — “ascending,” “transverse,” and “descending.” The ascending run up to the outer side of the ilium, beneath the tensor fasciæ and the glutæus medius, supply these muscles, and inosculate with the terminal branches of the glutæal artery. The transverse are chiefly lost in the vastus externus. The descending, one or more in number, of considerable size, run down between the rectus and the cruræus, supply both

* The point at which the profunda is given off below the crural arch varies very much even in the two limbs of the same body. We have measured it in 19 bodies, or 38 femoral arteries. It varied from $\frac{1}{2}$ to 3 inches. In 22 cases the profunda came off between $1\frac{1}{2}$ and 2 inches ; in 9 this distance was exceeded ; in 7 this distance was less.

these muscles, and may be traced as low as the knee, where they inosculate with the articular arteries.

[The external circumflex may come off from the femoral above the profunda; or we may find two arteries, one from the femoral, and one from the profunda, or both from either of the above vessels. The internal circumflex may also come from the femoral. It has been known to come from the epigastric, circumflex iliac, or external iliac artery.]

Fig. 98.



PLAN OF THE INOSCULATIONS OF THE CIRCUMFLEX ARTERIES

The *perforating* branches of the profunda are so named because they pass through the adductors to supply the hamstring muscles. There are generally three, but they vary both in number and size. The first is the largest. It passes between the pectineus and adductor brevis, and then through the tendon of the adductor magnus. The second passes

through the tendon of the adductor brevis and magnus.* The third through the tendon of the adductor magnus. They not only supply the hamstring muscles—namely, the biceps, semi-tendinosus, and semi-membranosus; but some of their branches turn horizontally round the back of the femur to reach the vastus externus, in the substance of which they communicate by a series of arches.

Arterial Inosculations.—To test your knowledge of the arterial inosculations about the upper part of the thigh, reflect on the answer to the following questions: If the femoral were tied *above* the origin of the profunda, how would the circulation be carried on? The ascending branch of the external circumflex communicates with the glutæal; the internal circumflex communicates with the obturator and ischiatic (see fig. 98). Again—how is the circulation maintained when the femoral is tied *below* the profunda? The descending branch of the external circumflex and the perforating branches of the profunda communicate with the articular branches of the popliteal and the tibial recurrent.†

The continuation of the profunda, much reduced in size, passes through the tendon of the adductor magnus, and is distributed to the short head of the biceps.

The *anastomotica magna* arises from the femoral artery just before it leaves its tendinous canal. It runs in front of the tendon of the adductor magnus, in company with the long saphenous nerve to the inner side of the knee. Here it ramifies extensively over the capsule, and communicates with the other articular arteries.‡

[The anastomotic not unfrequently varies in point of origin as well as in calibre.]

Anterior Crural and Obturator Nerves.—The anterior crural nerve is the largest branch of the lumbar plexus (p.

* This perforating branch usually furnishes the nutritious artery of the femur.

† Read the account of the dissection of an aneurismal limb by Sir A. Cooper, Med. Chir. Trans. vol. ii.

‡ In its course down the thigh the femoral artery gives off several small branches to the sartorius, and one of considerable size for the supply of the vastus internus. We may trace this branch through the substance of the vastus down to the patella, where it joins the beautiful network of vessels on the surface of that bone.

335). It comes from the third and fourth lumbar nerves. It passes under the crural arch in a groove between the iliacus and psoas, about one-quarter of an inch from the outer side of the femoral artery. It soon divides into a number of branches, some of which supply the skin,—others, the extensor muscles of the thigh. The cutaneous branches,—namely, the long saphenous, the middle and the internal cutaneous, have been already described (p. 437). Its muscular branches you should trace into the extensors of the leg; namely the rectus, the two vasti, the cruræus, and subcruræus. But it cannot be said that it supplies them only, for it sends a branch (beneath the femoral vessels), to the pectineus; one also to the sartorius.

The *obturator nerve*, also a branch of the lumbar plexus (p. 335), supplies the adductor muscles. It enters the thigh through the upper part of the obturator foramen with the corresponding artery, and immediately divides into two branches, of which one passes in front of, the other behind the adductor brevis. The front branch* subdivides for the supply of the gracilis, the adductor longus and brevis; the hind one supplies the obturator externus and the adductor magnus. In some bodies you can succeed in tracing a filament of the obturator nerve through the notch of the acetabulum into the hip-joint, and another, which runs near the popliteal artery, into the back part of the knee-joint. I have frequently seen cutaneous branches from the obturator on the inner side of the thigh. This is interesting practically, since it helps to explain the great pain often felt on the inner side of the knee in cases of disease of the hip-joint.

The *obturator artery* after passing through the foramen, divides into two branches, which form a circle round the obturator membrane. These supply the external obturator and adductors of the thigh, and inosculate with the internal circumflex and ischiatic arteries (p. 441). It usually gives off also the little artery to the ligamentum teres of the hip-joint.

* Near the lower border of the adductor longus: a filament from the first branch generally communicates with one of the internal cutaneous branches of the anterior crural nerve. This communication takes place over the femoral artery in Hunter's canal.

DISSECTION OF THE FRONT OF THE LEG.

CONTINUE the incision from the knee down the front of the leg, over the ankle, along the top of the foot to the great toe. Another incision must be made at right angles to the first, on either side of the ankle, that the skin may be reflected from the front and sides of the leg and foot.

Cutaneous Veins and Nerves.—Having already traced the internal saphena vein (p. 418) to the inner side of the knee, you must now follow it down the inner side of the leg, over the inner ankle* to the top of the foot. On the top of the foot notice that the principal veins form an arch, with the convexity forwards, just as on the back of the hand. This arch receives the veins from the toes. From the inner side of the arch the internal saphena originates; from the outer side the external saphena. The latter vein runs behind the external ankle, up the back of the calf of the leg to join the popliteal vein.

The skin on the inner side of the leg is supplied by the long saphenous nerve (p. 437). This nerve becomes subcutaneous on the inner side of the knee, between the gracilis and sartorius. Here it meets the saphena vein, and accompanies it down the leg, distributing its branches on either side, till it is finally lost on the inner side of the foot. The largest branch curves round the inner side of the knee, just below the patella, to supply the skin in this situation.

External Cutaneous Nerve.—This is a branch of the peroneal nerve. It comes through the fascia about the lower third of the outer side of the leg; and descending over the front of the ankle, divides and subdivides over the top of the foot. Trace its subdivisions and you will find that one runs along the inner side of the great toe; the others supply the outside of the second toe, both sides of the third and fourth, and the inside of the little toe.

The outside of the little toe is supplied by the external saphenus nerve, which runs behind the outer ankle with the corresponding vein.

* The French commonly bleed from the internal saphena as it runs over the inner ankle, this being a convenient and safe place for venesection.

The contiguous sides of the great toe, and the toe next to it, are supplied by the termination of the anterior tibial nerve.*

Muscular Fascia and Annular Ligaments.—This is remarkably thick and strong. Besides its general purpose of forming sheaths for the muscles, and straps for the tendons, it gives origin, as in the forearm, to some of the muscular fibres; so that it cannot be removed from the surface of the muscles, near the knee, without leaving them ragged. Observe that the fascia is attached to the head of the tibia and the fibula; that it is identified on the inner side with the expanded tendons of the sartorius, gracilis, and semitendinosus: on the outer side with that of the biceps; consequently, when these muscles act, the fascia is rendered tense. Following it down the leg, you find that it is attached to the edge of the tibia, and that it becomes stronger as it approaches the ankle, in order to form the ligaments which confine the tendons in this situation. Of these ligaments, which are called "*annular*," there are three, as follow:—

a. The *anterior* extends obliquely across the front of the ankle-joint and confines the extensor tendons of the ankle and toes. It consists of two straps which cross each other over the front of the ankle-like braces. One brace goes from the malleolus externus to the scaphoid and internal cuneiform bones; the other runs from the cuboid and os calcis upwards and inwards to the inner border of the tibia: it is the excessive strain of this ligament which occasions the pain in sprains of the ankle. You will see presently that it makes a beautiful pulley for the extensor longus digitorum.

b. The *external* extends from the outer malleolus to the os calcis, and confines the tendons of the peronei muscles, which draw the foot outwards.

c. The *internal* extends from the inner malleolus to the os calcis, and binds down the flexor tendons of the foot and toes.

Remove the fascia, leaving enough of the annular ligaments to retain the tendons in their places.

* Such is the most common distribution of the nerves to the upper surface of the toes. But deviations from this arrangement are frequent.

Muscles on the Front of the Leg.—The muscles on the front of the leg are:—1, the *tibialis anticus*; 2, the *extensor communis digitorum* and *peroneus tertius*; 3, the *extensor proprius pollicis*.

Tibialis Anticus.—The *tibialis anticus* arises from the upper two-thirds of the outer side of the tibia, from the interosseous membrane, and from the fascia which covers it. About the lower third of the leg the fibres terminate on a strong flat tendon, which descends obliquely over the front of the ankle to the inner side of the foot; here it becomes a little broader, and is inserted into the internal cuneiform bone and the metatarsal bone of the great toe. The synovial membrane, which lines the sheath of the tendon beneath the annular ligament, accompanies it to within an inch of its insertion; consequently it is opened when the tendon is divided for club-foot. The action of this muscle is to draw the foot upwards and inwards.* When the foot is the fixed point, it assists in balancing the body at the ankle.

[The *tibialis anticus* flexes the ankle, adducts the foot, and raises the inner edge from the ground.]

Extensor Communis Digitorum.—This muscle lies along the fibular side of the preceding. It arises from the head of the tibia, from the upper two-thirds of the fibula, from the interosseous membrane and the fascia. Its fibres terminate in a penniform manner upon a long tendon which descends over the ankle and divides into four. These diverge from each other, and are inserted into the toes thus:—On the first phalanx, each tendon (except that of the little toe) is joined on its outer side by the corresponding tendon of the *extensor brevis*. The united tendons then expand, and are inserted as on the fingers; that is, the middle part is inserted into the base of the second phalanx; the sides run on to the base of the third.

Immediately below the ankle the anterior annular ligament forms a beautiful pulley through which the tendon of this muscle plays. It is like a sling, of which the two ends are attached to the *os calcis*, while the loop serves to confine the

* It is generally necessary to divide this tendon in distortion of the foot inwards, called "*talipes varus*."

tendon. The play of the tendon is facilitated by a synovial membrane, which is prolonged for a short distance along each of its four divisions. Besides its chief action, this muscle extends the ankle-joint.*

Peroneus Tertius.—This appears like a portion of the preceding. Its fibres arise from the lower part of the shaft of the fibula, and terminate on their tendon like barbs on a quill. The tendon passes through the same pulley with the long extensor of the toes, and, expanding considerably, is inserted into the tarsal end of the metatarsal bone of the little toe.

[The peroneus is sometimes absent; then a tendon from the extensor communis takes its place. It flexes the ankle, abducts the foot, and raises its outer edge.]

The peroneus tertius and the tibialis anticus are important muscles in progression. They raise the toes and foot from the ground. Persons who have lost the use of these muscles are obliged to drag the foot along the ground, or to swing the entire limb outwards in walking.

Extensor Proprius Pollicis.—This muscle lies partly concealed between the tibialis anticus and the extensor longus digitorum. It arises from rather more than the middle third of the fibula, and from the interosseous membrane. The fibres terminate in a penniform manner on the tendon, which runs over the ankle, along the top of the foot to the great toe, where it is inserted into the base of the last phalanx. It has a special pulley beneath the angular ligament, lined by a synovial membrane, which accompanies it as far as the metatarsal bone of the great toe.

[The extensor proprius pollicis flexes the ankle, extends the great toe, and may also abduct the foot and rotate it inwards. The extensor brevis extends the toes, and rotates the anterior part of the foot outwards.]

Extensor Brevis Digitorum.—This muscle is situated on the top of the foot, beneath the long extensor tendons of the toes. It arises from the front part of the os calcis, and from the ligaments uniting this bone to the astragalus. The fibres

* There is often a large bursa between the tendon of the extensor longus digitorum and the outer end of the astragalus. There is much friction here. What is of more importance, this bursa sometimes communicates with the joint of the head of the astragalus; it did so in the last foot I examined.

run obliquely over the foot, and terminate on four tendons, which join the fibular side of the long extensor tendons of the four inner toes.

Now examine the course, relations, and branches of the anterior tibial artery. Since it lies deep between the muscles, it is necessary to separate them from each other: this is easily done by proceeding from the ankle towards the knee.

Course and Relations of the Anterior Tibial Artery.—The anterior tibial artery is one of the two branches into which the popliteal divides at the lower border of the popliteus. It comes forward about $1\frac{1}{4}$ inch below the head of the fibula, and then descends, lying in rather more than the first half of its course upon the interosseous membrane, afterwards along the front of the tibia. It runs beneath the annular ligament over the front of the ankle, and over the instep to the interval between the first and second metatarsal bones, where it sinks into the sole and joins the deep plantar arch. Thus, a line drawn from the head of the fibula to the interval between the first and second metatarsal bones would nearly coincide with its course. In the upper third of the leg it lies deep between the tibialis anticus and the extensor communis digitorum; in the lower two-thirds, between the tibialis anticus and the extensor proprius pollicis. But in *front of the ankle* the artery is crossed by the extensor proprius pollicis, so that on the top of the foot it runs along the outer side of this tendon. Again, just before it sinks into the sole, it is crossed by the short extensor tendon of the great toe.

The artery is accompanied by the anterior tibial nerve (a branch of the peroneal), which runs for some distance on its fibular side, crosses it above the ankle, and lies at the top of the foot on its tibial side. It is accompanied by two veins, one on either side, which communicate at intervals by cross branches.

The best guide to the artery in the upper part of its course is the inner edge of the fibula; on the top of the foot, the outer border of the long extensor tendon of the great toe.

The branches of the anterior tibial are as follow:—

a. The *recurrent* branch ascends close by the outer side of the head of the tibia to the knee-joint, where it inosculates with the other articular arteries derived from the popliteal.

b. Irregular *muscular* branches.

c. The *malleolar* branches, external and internal, ramify over the ankle, supplying the joint, the articular ends of the bones, and the sheaths of the tendons around them.

d. The *tarsal* branch arises near the scaphoid bone, passes beneath the extensor brevis digitorum towards the outside of the foot, supplies the bones and joints of the tarsus, and inosculates with the arteries in the sole.

e. The *metatarsal* branch generally runs towards the outside of the foot, near the bases of the metatarsal bones, and gives off the three outer interosseous arteries. These pass forwards over the corresponding interosseous muscles, supply them, and then subdivide to supply the opposite sides of the upper surface of the toes. They communicate by perforating branches with the plantar arteries.

f. The *dorsalis halucis* is, strictly speaking, the artery of the first interosseous space. It comes from the anterior tibial just before this sinks into the sole, and runs forwards to supply the opposite sides of the first two toes.

[The anterior tibial artery, necessarily depends as to its place of origin upon the division of the popliteal. Where the posterior tibial is absent, it is joined with the peroneal. It has been seen lying upon the popliteus muscle, and between that muscle and the bone, when it has arisen high up from the popliteal. Cases are reported where the vessel has passed outwards along the margin of the fibula, then regained its normal position in front of the ankle joint. Velpeau gives a case where it continued down from the middle of the leg, only covered by fascia and integument. It has also been found to pass round the outer side of the fibula before it reached the anterior portion of the leg. It may be altogether wanting, or much diminished in size, or end at the ankle. The digital branches supplying the first and second toes may come from the internal plantar; and the anterior part of the leg be supplied by perforatory branches from the posterior tibial, in these cases.]

Peronei Muscles.—These muscles are situated on the outer side of the fibula, and are named respectively the peroneus longus and brevis.

Peroneus Longus.—This arises from the outer surface of

the fibula along its upper half. The fibres terminate in a penniform manner upon a tendon, which runs through a groove behind the outer ankle, then along the outer side of the os calcis, and lastly, through a groove on the under surface of the os cuboides deep into the sole. You will see presently that it crosses the sole obliquely, and is inserted into the tarsal end of the metatarsal bone of the great toe. In its course through these several bony grooves the tendon is confined by a fibrous sheath, lined by a synovial membrane. In removing the metatarsal bone of the great toe, you ought if possible to leave the insertion of this tendon.

Peroneus Brevis.—This muscle lies beneath the preceding. It arises from the lower half of the outer surface of the fibula. It terminates on a tendon which runs behind the outer ankle, through the same sheath with the peroneus longus, then proceeds along the outside of the foot, and is inserted into the tarsal end of the metatarsal bone of the little toe.*

The *action* of the peronei is to raise the outer side of the foot.† This movement regulates the bearing of the foot in progression, so as to throw the principal part of the weight on the ball of the great toe. Its action is well exemplified in skating. Again, supposing the fixed point to be the foot, they tend to prevent the body from falling on the opposite side, as when we balance ourselves on one leg.

Peroneal Nerve.—Near the inner side of the tendon of the biceps flexor of the leg, is a large nerve called the peroneal, a branch of the great ischiatic. By reflecting the upper part of the peroneus longus, you will find that this nerve runs round the outer side of the fibula immediately below its head. Here it divides into several branches, as follow:—1. A small articular branch, which accompanies the recurrent artery to the knee-joint; 2. The anterior tibial, which accompanies the corresponding artery; 3. The external cutaneous (p. 444), which comes through the fascia between the peroneus longus and the extensor longus digitorum; 4. Branches which supply

* On the outside of the os calcis there is a slight ridge which separates the tendons of the peronei. Each has a distinct sheath. The short tendon runs above, the long one below the ridge.

† In distortion of the foot outwards, called "talipes valgus," it is generally necessary to divide the tendons of the peronei.

all the muscles in front of the leg, namely, the extensors of the foot and the toes and the peronei.

DISSECTION OF THE GLUTÆAL REGION.

THE incision through the skin should commence at the coccyx, and be continued in a semicircular manner along the crest of the ilium. Another incision should be made from the coccyx downwards and outwards for about six inches below the great trochanter. In reflecting the skin, notice the thick cushion which the subcutaneous adipose tissue forms over the tuberosity of the ischium. I have often seen a large bursa between the cushion and the bone.

Cutaneous Nerves.—These are derived from several sources. Some descend over the crest of the ilium, near the origin of the erector spinæ; they are the posterior divisions of the first and second lumbar nerves. Over the middle of the crest comes the lateral branch of the twelfth dorsal nerve, and often a smaller one from the first lumbar. Other cutaneous nerves come up from below; they are branches of the lesser ischiatic, and proceed from beneath the lower border of the glutæus maximus. Lastly, the skin of the sacrum and coccyx is supplied by the posterior branches of the sacral nerves.

Glutæal Muscles.—Three powerful muscles are situated on the buttock, one above the other, named, according to their relative size, the glutæus maximus, medius, and minimus.

Glutæus Maximus.—This is the largest muscle in the body. Its great size is characteristic of man, in reference to his erect position. Its texture is thick and coarse. It arises from the posterior fifth of the crest of the ilium, from the lower part of the sacrum, the coccyx, and the great sacro-ischiatic ligament. The fibres descend obliquely forwards, and are inserted thus:—the anterior two-thirds terminate on a strong broad tendon which plays over the great trochanter, and joins the aponeurosis on the outside of the thigh (p. 429); the remaining third is inserted into the femur, along the ridge leading from the linea aspera to the great trochanter.

This muscle *extends* the thigh bone upon the pelvis, and is, therefore, one of those most concerned in raising the body from the sitting to the erect position, and in maintaining it erect. It also propels the body—in walking, running or leaping. It is well supplied with blood by the glutæal and ischiatic arteries; with nerves by the lesser ischiatic.

What is seen beneath the Glutæus Maximus.—The glutæus maximus should be reflected from its origin. The best way to do so is to begin at the front border, which overlaps the glutæus medius. This dissection is difficult, and he who undertakes it for the first time is almost sure to damage the subjacent parts. The numerous nutrient vessels which enter its under surface must necessarily be divided. Supposing the object accomplished, what have we exposed beneath? The muscle covering the ilium is the glutæus medius. At the posterior border of this are the several objects which come out of the pelvis through the great ischiatic notch—namely, the pyriformis muscle, above which is the trunk of the glutæal artery with the glutæal nerve, and below which are the greater and lesser ischiatic nerves, the ischiatic artery, and the pudic artery and nerve. Coming through the lesser ischiatic notch, we see the tendon of the obturator externus, and tacked on to it are the gemelli muscles, one above, the other below it. Extending from the tuber ischii to the great trochanter is the quadratus femoris. Lastly, where the tendon of the glutæus maximus plays over the trochanter major, there is a large bursa, simple or multilocular. I have seen it in some subjects sufficiently capacious to hold half a pint of fluid.

Glutæus Medius.—This muscle arises from the surface of the ilium, between the crest and the upper curved line; also from the strong fascia which covers it towards the front. The fibres converge to a tendon which is inserted into the upper and outer surface of the great trochanter: some of the anterior fibres—in immediate connection with the tensor fasciæ—terminate on the aponeurosis of the thigh.

Reflect the glutæus medius to see the third glutæal muscle. The line of separation between them is marked by a large branch of the glutæal artery.

Glutæus Minimus.—This muscle arises from the surface of the ilium below the upper curved line. Its fibres pass over

the capsule of the hip-joint, and converge to a tendon which is inserted into a depression on the front part of the great trochanter. The chief action of this and the preceding muscle is to assist in balancing the pelvis steadily on the thigh, as when we are standing on one leg; with the fixed point at the ilium, it is an abductor of the thigh. The anterior fibres of the glutæus medius co-operate with the tensor fasciæ in rotating the thigh inwards.

Glutæal Vessels and Nerves.—The glutæal artery is the largest branch of the internal iliac (p. 441). Emerging from the pelvis through the great ischiatic notch above the pyriformis muscle, it at once divides into large branches for the supply of the glutæal muscles. Of these, some proceed forwards between the glutæus maximus and medius; others run in curves between the glutæus medius and minimus, towards the anterior part of the ilium. Many of them inosculate with branches of the external circumflex.

The nerve which accompanies the glutæal artery is a branch of the lumbo-sacral nerve (p. 455). It subdivides to supply the glutæus medius and minimus, and the tensor fasciæ; in some subjects it sends a branch to the glutæus maximus; but this muscle is supplied by the lesser ischiatic nerve.

A practical surgeon ought to be able to cut down and tie the glutæal artery as it emerges from the pelvis. The following is the best rule for finding it:—the toes are supposed to be turned inwards. Draw a line from the posterior superior spine of the ilium to the midspace between the tuber ischii and the great trochanter. The artery emerges from the pelvis at the point where the superior third joins the two inferior thirds of this line.*

Now examine the series of muscles which rotate the thigh outwards—namely, the pyriformis, the obturator internus, the gemelli, the quadratus femoris, and the obturator externus.

Pyriformis.—This muscle arises from the front surface of the sacrum between the holes for the sacral nerves, and from the margin of the great sacro-ischiatic notch. The fibres converge to a tendon which is inserted into the upper border of the trochanter major. Its nerve comes from the sacral plexus.

* The operation of tying the glutæal artery was first performed by John Bell. See his "Principles of Surgery," vol. i. p. 421.

[This muscle is often divided by the sciatic nerve. This is frequent when the nerve passes out of the pelvis as two cords; one as usual beneath, the other through the muscle.]

Obturator Internus.—This muscle, of which we can at present see very little more than the tendon, arises within the pelvis from the ischium between the great ischiatic notch and the obturator foramen; also from the obturator membrane. The fibres terminate on four tendons which converge towards the lesser ischiatic notch, pass round it as round a pulley, and then uniting into one, are inserted into the top of the great trochanter. Divide the tendon about three inches from its insertion, to see the four tendons which play over the smooth cartilaginous pulley. There is a large synovial bursa to prevent friction. The nerve of this muscle comes from the sacral plexus; sometimes from the pudic nerve.

Gemelli.—These muscles are accessory to the obturator internus, and are situated, one above, the other below it. The gemellus superior arises from the spine of the ischium: the gemellus inferior from the upper part of the tuberosity. Their fibres are inserted into the tendon of the obturator internus. Both muscles derive their nerves from the sacral plexus.

[Sometimes both gemelli are absent, then again only the superior. The quadratus also is occasionally wanting.]


Quadratus Femoris.—This muscle arises from the ridge on the outer part of the tuber ischii. Its fibres run horizontally outwards, and are inserted into the back of the great trochanter, in a line with the linea aspera. Notice that the lower border of the quadratus femoris runs parallel with the upper edge of the adductor magnus; in fact, it lies on the same plane. Between these muscles is generally seen a terminal branch of the internal circumflex artery. The nerve of this muscle comes from the sacral plexus.

Obturator Externus.—To see this muscle you must reflect the quadratus femoris. It arises from the front surface of the ramus of the pubes and ischium, and from the obturator membrane. The fibres converge to a tendon which runs horizontally outwards over a groove in the ischium, and is inserted into the deepest part of the pit of the great trochanter. Its nerve is a branch of the obturator (p. 443).

Great Ischiatic Nerve.—This large nerve is formed by

the union of all the sacral nerves (fig. 99), and is destined to supply all the flexor muscles of the lower extremity. Emerging from the pelvis through the ischiatic notch, below the pyriformis, it descends over the external rotator muscles of the thigh, along the interval between the tuber ischii and the trochanter major, but rather nearer to the former, so that, in the

Fig. 99.



PLAN OF THE SACRAL PLEXUS AND BRANCHES.

sitting position, the nerve is protected from pressure by this bony prominence. The nerve does not descend quite perpendicularly, but rather obliquely forwards, parallel with the great sacro-ischiatic ligament. It is accompanied by a branch of the ischiatic artery, called the "*comes nervi ischiatici*."* I have known serious secondary hæmorrhage from this branch after an amputation.

* The arteria comes nervi ischiatici runs generally by the side of the nerve, but sometimes in the centre of it. This artery becomes one of the chief chan-

Lesser Ischiatic Nerve.—This comes from the lower part of the sacral plexus. It leaves the pelvis with the greater ischiatic nerve, but on the inner side of it, and in company with the ischiatic artery. The only muscular branches which it gives off are one or more which supply the glutæus maximus. All its other branches are cutaneous. One turns round the lower border of the glutæus maximus, and supplies the skin of the buttock. Another, called the long pudendal, turns inwards towards the perineum, to supply the skin of that region and the scrotum. The continued trunk runs down the back of the thigh beneath the muscular fascia, as low as the upper part of the calf, supplying the skin all the way down.

Ischiatic Artery.—This branch of the internal iliac leaves the pelvis below the pyriformis, and then descends between the tuber ischii and the great trochanter, along the inner side of the great ischiatic nerve. It gives off,—1, Two or more considerable branches to the glutæus maximus; 2, A coccygeal branch, which runs through the great sacro-ischiatic ligament, and then ramifies in the glutæus maximus, and on the back of the coccyx; 3, The comes nervi ischiatici; 4, Branches to the several external rotator muscles; lastly, branches which supply the upper part of the hamstring muscles; and others which inosculate with the internal circumflex and obturator arteries (p. 441).

Pudic Artery and Nerve.—The course of this artery and nerve has been fully described (p. 353). All that we have to observe about them now is, that they pass over the spine of the ischium, and that in a thin subject it is possible to compress the artery against the spine. The rule for finding it is this: rotate the foot inwards, and draw a line from the top of the great trochanter to the base of the coccyx; the junction of the inner with the outer two-thirds gives the situation of the artery.*

nels by which the blood reaches the lower limb after ligature of the femoral. See a preparation in the Museum of St. Bartholomew's Hospital, where the femoral was tied by John Hunter fifty years before the man's death.

* Mr. Travers succeeded in arresting hæmorrhage from a sloughing ulcer of the glans penis by pressing the pudic artery with a cork against the spine of the ischium.

DISSECTION OF THE BACK OF THE THIGH.

THE incision should be continued down the back of the thigh, the ham, and half way down the calf. The skin should be reflected on either side.

Cutaneous Nerves and Veins.—The skin at the back of the thigh is supplied by the lesser ischiatic nerve, which runs down beneath the fascia, as low as the upper third of the calf, distributing branches on either side.

There are no subcutaneous veins of any size at the back of the thigh; for here they would be liable to pressure. But near the ham we find a vein, called the “external saphena.” It ascends up the back of the calf, and joins the popliteal vein.

Muscular Fascia.—Respecting this, there is nothing to be remarked more than that its fibres run chiefly in a transverse direction, that it becomes stronger as it passes over the popliteal space, and that here it is connected with the tendons on either side. Remove it, in order to examine the powerful muscles which bend the leg, called the “hamstrings.”

Hamstring Muscles.—There are three of these, and all arise by strong tendons from the tuber ischii. One, the biceps, is inserted into the head of the fibula; while the other two, namely, the semitendinosus and semimembranosus, are inserted into the tibia. The divergence of these muscles towards their respective insertions occasions the well-known space termed the “ham,” which is occupied by soft fat, and the popliteal vessels and nerves.

Biceps.—This muscle, as implied by its name, has two origins, a long and a short. The long one takes place by means of a strong tendon, from the back part of the tuber ischii; the short one, by fleshy fibres, from the linea aspera of the femur. This, which we can easily see by dissecting on the outside of the thigh, begins at the linea aspera, just below the insertion of the glutæus maximus, and continues nearly down to the external condyle. It joins the longer part of the muscle, and both terminate on a common tendon, which is inserted into the head of the fibula. It also gives off a strong expansion to the fascia of the leg. The tendon covers part of

the external lateral ligament of the knee-joint, and a small bursa intervenes.*

[The short head may be wanting ; again, we may find another long head passing from the tuber ischii down the leg to the tendo Achillis. The relation the peroneal nerve holds to the tendon of the biceps should be noticed, as it might be wounded in tenotomy of this tendon.]

The biceps is not only a flexor of the leg, but rotates the leg, when bent, outwards. It is this muscle which dislocates the knee outwards and backwards in chronic disease of the joint.

Semitendinosus.—This arises, in common with the biceps, from the back part of the tuber ischii. The fibres terminate upon a long tendon, which spreads out below the knee-joint, and is inserted into the inner surface of the tibia below the tendons of the sartorius and gracilis. Like them, it plays over the internal lateral ligament of the knee-joint, and is provided with a bursa to prevent friction.

[The semitendinosus may be divided by one or more tendinous lines.]


The semitendinosus sends off from the lower border of its tendon a very strong fascia to cover the leg, which is attached along the inner edge of the tibia. The middle of the muscle is curiously intersected by an oblique tendinous line.

Semimembranosus.—This muscle arises from the tuber ischii, outside the two preceding, by means of a strong flat tendon which extends nearly half way down the thigh. It forms a bulky muscle which lies on a deeper plane than the others, and is inserted by a thick tendon into the posterior part of the head of the tibia. In connection with this tendon there are three additional points to be noted—1, that its tendon is prolonged under the internal lateral ligament of the knee, and that a bursa intervenes between them ; 2, that it forms the principal constituent of the “ligamentum posticum Winslowii” which covers the back of the knee-joint ; 3, that a fascia proceeds from its lower border, and binds down the popliteus.

* This tendon can be plainly felt as the outer hamstring in one's own person, just above the head of the fibula.

A large *bursa* is almost invariably found between the semi-membranosus and the inner head of the gastrocnemius, where they rub one against the other. It is generally from one and a half to two inches long. The chief point of interest concerning it is, that it occasionally communicates with the synovial membrane of the knee-joint, not directly, but through the medium of another bursa beneath the inner head of the gastrocnemius. From an examination of 150 bodies, I infer that this communication exists about once in five instances; and it need scarcely be said that the proportion is large enough to make us cautious in interfering too roughly with this bursa when it becomes enlarged.*

Action of the Hamstring Muscles.—These muscles produce two different effects, according as their fixed point is at the pelvis or the knee. With the fixed point at the pelvis, it is obvious that they bend the knee; with the fixed point at the knee, they form a very important part of the machinery which keeps the body erect. For instance, if, when standing, you bend the body at the hip, and feel the muscles in question, you find that they are in strong action, to prevent the trunk from falling forwards: they, too, are the chief agents concerned in bringing the body back again to the erect position. In doing this, they act upon a lever of the first order as shown in fig. 100; the acetabulum being the fulcrum F, the trunk w, the weight to be moved, and the power P, at the tuber ischii.



* When the bursa in question becomes enlarged, it occasions a fluctuating swelling of greater or less dimensions on the inner side of the popliteal space. The swelling bulges out, and becomes tense and elastic when the knee is extended, and *vice versa*. As to its shape, it is generally oblong; but this is subject to variety, for we know that the bursæ, when enlarged, are apt to become multilocular, and to burrow between the muscles where there is the least resistance.

To put the action of the muscles of the thigh on the pelvis in the clearest point of view, let us suppose we are standing upon one leg: the bones of the lower extremity represent a pillar which supports the weight of the trunk on a ball-and-socket joint; the weight is nicely balanced on all sides, and prevented from falling by four groups of muscles. In front are the rectus and sartorius; on the inner side, the adductors; on the outer side, the glutæus medius and minimus; behind, the hamstring and glutæus maximus.

The hamstring muscles are supplied with blood by the perforating branches of the profunda, which come through the tendon of the adductor magnus close to the femur. Their nerves are derived from the great ischiatic.

Popliteal Space.—Since the hamstring muscles diverge from each other to reach their respective insertions, they leave between them a space, called the “popliteal.” Now this space is bounded below by the converging origins of the gastrocnemius; its shape, is, therefore, that of a lozenge. It is filled up by a large quantity of fat, which permits the easy flexion of the thigh, and in this fat we find imbedded the popliteal vessels and nerves in the following order: nearest to the surface are the nerves, while the vessels lie close to the bone, the vein being superficial to the artery (fig. 101).

Popliteal Nerve.—If the ischiatic nerve be traced from the buttock, we find that it descends upon the adductor magnus, and that, after being crossed by the long origin of the biceps, it then runs along the outer border of the semimembranosus. Having given branches to the three great flexor muscles, it divides, about the lower third of the thigh (higher or lower in different subjects), into two large nerves—the peroneal and the popliteal.

The *peroneal* runs close by the inner side of the tendon of the biceps* towards the head of the fibula, below which we have already traced it (p. 450), dividing into branches for the peronei and all the extensors of the foot and the toes.


The *popliteal* accompanies the popliteal artery, and is des-

* The nerve is, therefore, very liable to be injured in the operation of dividing the outer hamstring.

tined to supply the flexor muscles on the back of the leg and the sole of the foot.

By clearing out all the fat, we observe that the popliteal vessels enter the ham through an aperture in the adductor magnus, and descend close to the back part of the femur, and afterwards over the back of the knee-joint. At first they are partially overlapped (in muscular subjects), by the semimembranosus; indeed, the outer border of this muscle is a good guide to the artery in the operation of tying it.

Fig. 101.



POPLITEAL SPACE.

Absorbent Glands.—With a little care you may discover two or more absorbent glands, which are situated one on each side of the artery. They deserve attention, because, when enlarged, their close proximity to the artery may give them an apparent pulsation which might be mistaken for an aneurism.

Deferring the course, relations, and branches of the popliteal artery till this vessel is exposed throughout its whole course, pass on now to the great muscles of the calf.

Continue the incision down the centre of the calf to the heel, and reflect the skin.

Cutaneous Veins and Nerves of the Calf.—The chief cutaneous vein and nerve of the calf is called the "*external or posterior saphenous*." The vein commences on the outer side of the foot, runs behind the outer ankle, and then up the calf, receiving numerous tributary veins in its course. It eventually passes through the muscular fascia, and joins the popliteal vein.

The corresponding nerve is formed by two roots,*—one from the peroneal, the other from the popliteal (fig. 101). The roots generally unite about the middle of the calf, and the single nerve coming through the fascia, descends with the saphenous vein, and is finally distributed to the outer side of the foot and the little toe.

Muscles of the Calf.—The great flexor muscle of the foot consists of two portions; the superficial one, called the "*gastrocnemius*," arises from the lower end of the femur; the deep one called the "*soleus*," arises from the tibia and fibula. The force of both is concentrated on one thick tendon, called the "*tendo Achillis*," which is inserted into the heel.

Gastrocnemius.—This muscle arises by two strong tendinous heads, one from the upper and back part of each condyle of the femur (fig. 101). The inner head is larger. The two parts of the muscle descend, distinct from each other, and form the two bellies of the calf, of which the inner is rather the lower. Both terminate rather below the middle of the leg, on the broad commencement of the *tendo Achillis*.

The *gastrocnemius* should be divided transversely near to its insertion, and reflected upwards from the subjacent *soleus*, as high as its origin. By this proceeding you observe that the contiguous surfaces of the muscles are covered by a beautiful glistening tendon, which receives the insertion of

* These two roots are sometimes called, respectively, the *communicans peronei* and *communicans poplitei*.

their fibres, and transmits their collective force to the tendo Achillis.

You observe also the large sural vessels and nerves which supply each head of the muscle. Moreover, to facilitate the play of the inner tendon over the condyle, there is a bursa, which generally communicates with the knee-joint; and in the substance of the outer tendon is commonly found a small piece of fibro-cartilage. Lastly, between the gastrocnemius and soleus is the tendon of the plantaris.

Plantaris.—This little muscle* arises from the rough line just above the outer condyle of the femur (fig. 101). It descends close to the inner side of the outer head of the gastrocnemius, and terminates, a little below the knee, in a long tendon, which we trace down the inner side of the tendo Achillis to the calcaneum.

[The plantaris is sometimes absent. The gastrocnemius and soleus are often affected in middle age with fatty degeneration.]

Soleus.—This muscle arises from the head and upper third of the posterior surface of the fibula, from the oblique ridge on the back of the tibia,† and from about the middle third of the inner border of this bone. The fibres swell out beneath the gastrocnemius, and terminate on a broad tendon, which, gradually contracting, forms a constituent part of the tendo Achillis. The soleus is supplied with blood by several branches from the posterior tibial: also by a large branch from the peroneal. Its nerve comes from the popliteal and enters the top of the muscle. This is an important muscle in a surgical point of view, for two reasons,—1, by reflecting its tibial origin, we reach the posterior tibial artery: 2, by reflecting its fibular origin we reach the peroneal.

The *tendo Achillis* begins about the middle of the leg, and is at first of very considerable breadth, but it gradually con-

* This is the representative of the palmaris longus of the forearm. In man it is lost on the calcaneum, but in monkeys, who have prehensile feet, it is the proper tensor muscle of the plantar fascia. It is remarkably strong in bears and plantigrade mammals.


† The tibial and fibular origins of the soleus constitute what some anatomists describe as the two heads of the muscle. Between them descend the popliteal vessels, protected by a tendinous arch.

tracts and becomes thicker as it descends. The narrowest part of it is about one inch and a half above the heel; here, therefore, it can be most conveniently and safely divided for the relief of club-foot. There is no risk of injuring the deeper-seated parts, because they are separated from the tendon by a quantity of fat. Its precise insertion is into the under and back part of the os calcis. Observe that the tendon previously expands a little, and that between it and the bone there is a bursa of considerable size.

The *action* of the gastrocnemius and soleus is to raise the body on the toes. Since the gastrocnemius passes over two joints, it has the power (like the rectus) of extending the one while it bends the other, and it is, therefore, admirably adapted to the purpose of walking. For instance, by first extending the foot it raises the body, and then, by bending the knee, it transmits the weight from one leg to the other. Supposing the fixed point to be at the heel, the gastrocnemius is also concerned in keeping the body erect, for it keeps the tibia and fibula perpendicular on the foot, and thus counteracts the tendency of the body to fall forwards.

The tendo Achillis in raising the body on tiptoe, acts with

Fig. 102.



great power and under the best leverage; for it acts upon a lever of the *first* order. The fulcrum (which is a movable one) is at the ankle-joint F, fig. 102; the weight, W, is at the toes; the power is at the heel, P. All the conditions are those of a lever of the first order. The power and the weight act in the *same* direction on *opposite* sides of the fulcrum. The pressure

upon the fulcrum is equal to the sum of the pressures applied: *i. e.* $P \times F + W \times F$.

[The student should now refer to the tendons about the the foot, in their connection to the various kinds of deformities coming under the head of club-foot. In *talipes equinus*, the heel is raised by the tendo Achillis. In *talipes varus*, the most frequent deformity, we have the heel raised by the tendo Achillis, the inner portion of the foot elevated by the tibialis

anticus, and the anterior two-thirds twisted in by the tibialis posticus and flexor longus digitorum. The patient then will walk upon the outer border of his foot, or even on the dorsum and external malleoli. In *talipes valgus*, the outer portion of the foot is raised by the peronei, and the patient walks on the internal malleoli. To relieve these deformities we often resort to division of the tendons of the muscles in fault, together with the fascia. Division of the tendo Achillis alone will often suffice.]

Course and Relations of the Popliteal Artery.—After passing through the opening in the tendon of the adductor magnus, the main artery of the lower limb takes the name of “popliteal.” It then descends nearly perpendicularly behind the knee-joint, between the origins of the gastrocnemius, as far as the lower border of the popliteus, where it divides into the anterior and posterior tibial arteries. In its descent it lies, first, upon the lower part of the femur, and here it is slightly overlapped by the semimembranosus; next it lies upon the posterior ligament of the knee-joint, and lastly upon the popliteus. The vein closely accompanies the artery, and is situated superficially with regard to it, and rather to its outer side in the first part of its course. The popliteal nerve runs also in a similar direction with the vein, but is still more superficial and to the outer side (fig. 101). The vessels and the nerve are surrounded by fat, and we generally find one or two absorbent glands in the immediate neighborhood of the artery, just above the joint.

[The popliteal may divide earlier than usual, rarely above the flexure of the joint. It has divided into the anterior tibial and peroneal. The peroneal also has come off from it, together with both tibials. The veins forming the popliteal may not unite as early as usual, and thus two veins may accompany the vessel for a portion, or even the whole length of the artery.]

The *branches* of the popliteal artery are the *articular* and the *sural*.

There are five *articular branches* for the supply of the knee-joint and the articular ends of the bone: the two superior run, one above each condyle, close to the bone; the two inferior run one beneath each lateral ligament of the joint; and

all four proceed towards the front of the capsule. The fifth, called the "*azygos*," enters the joint through the posterior ligament.

Arterial Inosculations.—These articular arteries form, over the front and sides of the joint, a beautiful network of vessels, which communicate superiorly with the descending branch of the external circumflex and the anastomotica magna, and inferiorly with the anterior tibial recurrent. It is mainly through these channels that the collateral circulation is established in the leg after ligature of the femoral artery.

The *sural* arteries proceed one to each head of the gastrocnemius, and are proportionate in size to the muscle. One or two branches are distributed to the soleus. These arteries are accompanied by branches of the popliteal nerve for the supply of the muscles.

You can now examine more fully the insertion of the semimembranosus into the head of the tibia, alluded to p. 458.

Popliteus.—This muscle arises from a depression on the outside of the external condyle by a thick tendon which runs beneath the external lateral ligament of the knee-joint. The muscular fibres gradually spread out, and are inserted into the posterior surface of the tibia above the oblique ridge on this bone. Its action is to rotate the tibia inwards. Notice that the tendon plays over the articulation between the tibia and fibula, that a bursa intervenes to prevent friction, and that this generally communicates by a wide opening with the knee-joint.

Reflect the soleus from its origin, and remove it from the deep-seated muscles, observing at the same time the numerous arteries which enter its under surface. This done, notice the fascia which binds down the deep muscles. It is attached to the margin of the bones on either side, and increases in strength towards the ankle, in order to form an annular ligament to confine the tendons and the vessels and nerves in their passage into the sole of the foot.

Deep Muscles on the Back of the Leg.—There are three:—1, the flexor longus digitorum on the tibial side; 2, the flexor longus pollicis on the fibular; 3, the tibialis posticus upon the interosseous membrane, between and beneath them both.

Flexor Longus Digitorum.—This arises from the poste-

rior surface of the tibia, commencing below the popliteus, and extending to within about four inches of the lower end of the bone. The fibres terminate on a tendon which runs through a groove on the back of the inner ankle, and entering the sole, divides into four tendons for the four outer toes.

Flexor Longus Pollicis.—This is a remarkably powerful muscle. It arises from the lower two-thirds of the posterior surface of the fibula. The fibres terminate on a tendon which runs through a groove on the back of the astragalus, and thence along the sole to the great toe. The action of this muscle is to raise the body on the tip of the great toe. It is essential to the propulsion of the body in walking.

Tibialis Posticus.—This is so concealed between the two preceding muscles that you cannot properly examine it without reflecting them. It arises from the interosseous membrane and from the tibia and fibula. The muscular fibres terminate on a tendon which comes into view a short distance above the inner ankle, and, running through the same groove with the tendon of the flexor longus digitorum, then enters the sole, and is inserted into the scaphoid bone. Its action is to bend and turn the foot inwards.

The precise situation of the tendon of the tibialis posticus is interesting in a surgical point of view, because the tendon has to be divided for the relief of the talipes varus. It lies close to, and parallel with, the inner edge of the tibia, so that this is a very good guide to it. It is necessary to relax the tendon, while the knife is introduced between the tendon and the bone. Its synovial sheath commences about $1\frac{1}{2}$ inch above the end of the internal malleolus, and is consequently opened in the operation.

Course and Relations of the Posterior Tibial Artery.—This artery is one of the branches into which the popliteal divides at the lower border of the popliteus. It descends over the deep muscles at the back of the leg to the interval between the internal malleolus and the os calcis, and, entering the sole, divides into the external and internal plantar arteries. It lies first for a short distance, upon the tibialis posticus, then on the flexor longus digitorum; but behind the ankle it is in contact with the tibia, so that here it can be felt beating during life, and effectually compressed. In the up-

per part of its course, it runs nearly midway between the bones, is covered by the great muscles of the calf: to tie it, therefore, in this situation, is very difficult. But in the lower part of its course, it gradually approaches the inner border of the tibia, from which, generally speaking, it is not more than $\frac{1}{2}$ or $\frac{3}{4}$ inch distant. Here, being comparatively superficial, it may be easily tied. Immediately behind the internal malleolus, we find it between the tendons of the flexor longus digitorum, on the inner side, and the flexor longus pollicis on the outer. It has two venæ comites, which communicate at intervals. Its branches are as follow:—

a. Numerous unnamed branches to the soleus and the deep muscles.

b. The *peroneal* is a branch of very considerable size; often as large as the posterior tibial. Arising about $1\frac{1}{2}$ inch below the division of the popliteal, it descends close to the inner side of the fibula, and then over the articulation between the tibia and fibula to the outer part of the os calcis, where it inosculates with the malleolar and plantar arteries. Observe that all down the leg it is imbedded among the muscles: for it is covered first by the soleus, and afterwards by the flexor longus pollicis. To both these muscles, but to the latter especially, it sends numerous branches, and just above the ankle it gives off a constant one, which runs in front of the tibio-fibular articulation, and inosculates with the other malleolar arteries.

[The posterior tibial depends for its origin upon the division of the popliteal, and, like the anterior tibial, is subject to variations. It may be diminished in size, or absent; this is made up for by a large peroneal in the leg, or by the anterior tibial in the foot. The peroneal artery may arise higher or lower than usual, or may come from the anterior tibial; it is oftener larger than smaller, as to size. It may take the place of the posterior tibial at the lower part of the leg, and supply the foot. The anterior division (peroneal) may be large, and take the place of the anterior tibial in supplying the dorsum of the foot.]

Posterior Tibial Nerve.—This is the continuation of the popliteal. It descends close to its corresponding artery, and, entering the sole of the foot, divides into an external and in-

ternal plantar nerve. Respecting its precise position with regard to the artery, observe, that in the first part of its course the nerve lies superficial to the artery, and rather to its inner side; but behind the ankle the nerve lies on the outer side of the artery, and on the same plane.* Its branches supply the three deep-seated muscles.

DISSECTION OF THE SOLE OF THE FOOT.

MAKE a perpendicular incision down the middle of the sole, and reflect the skin. Notice the peculiar structure of the subcutaneous tissue. It is composed of globular masses of fat, separated by strong fibrous septa, and forms an elastic pad. This is especially marked at the heel, and at the ball of the great and the little toe; these being the points which form the tripod supporting the arch of the foot.

In removing the subcutaneous tissue from the ball of the great and the little toe, we often meet with bursæ, simple or multilocular. They are generally placed between the skin and the sesamoid bones, and have remarkably thick walls. Very frequently we trace an artery and nerve running directly through one of these sacs, and this explains the acute pain produced by their inflammation.

Cutaneous Nerves.—The skin of the heel is supplied by a branch of the posterior tibial nerve: the remainder of the sole by small branches of the plantar nerves which come through the fascia, just as in the palm of the hand.

Plantar Fascia.—This is remarkably dense and strong. It extends from the under and back part of the os calcis to the distal ends of the metatarsal bones: it not only protects the plantar vessels and nerves, but is also one among the many structures which support the arch of the foot. It acts like the string of a bow (p. 484). One sees this exemplified in some cases of distortion, where an unnatural contraction of the plantar fascia makes the arch of the foot too convex, and

* It sometimes happens that the nerve divides it into two plantar branches higher than usual, and then we find that one lies on either side of the artery.

it is necessary to divide the fascia in order to relieve the deformity.

The arrangement of the fascia is just like that in the palm. The central part, where there is the greatest strain, is very strong. The external part which is attached to the proximal end of the fifth metatarsal bone is also very strong. Near the distal ends of the metatarsal bones, the central part divides into five portions; each of these subdivides into two slips, which embrace the corresponding flexor tendons, and are attached to the metatarsal bones and their connecting ligaments. Between the primary divisions of the fascia,—that is, in a line between the toes,—we trace the digital vessels and nerves.

In the interdigital folds of the skin, there are also ligamentous fibres, which run from one side of the foot to the other, and answer the same purpose as those in the hand.

The plantar fascia must be partially removed to examine the muscles. You soon discover that towards the os calcis its removal is not accomplished without some difficulty, because the muscles arise from it.

Superficial Muscles.—After the removal of the fascia three muscles are exposed. All arise from the os calcis and the fascia, and proceed forwards to the toes.* The central one is the flexor brevis digitorum; the two lateral are the abductor pollicis and the abductor minimi digiti.

Abductor Pollicis.—This muscle arises from the inner and back part of the os calcis, and from the internal annular ligament. Its origin arches over the plantar vessels and nerves in their passage into the sole. The fibres run along the inner side of the sole, and terminate on a tendon which is inserted into the inner side of the base of the first phalanx of the great toe, through the medium of the internal sesamoid bone.

Abductor Minimi Digiti.—This muscle has a very strong origin from the under surface of the os calcis; also from its external tubercle. Some of its fibres terminate on a tendon which is inserted into the proximal end of the metatarsal bone of the little toe; but the greater part run on a tendon which is inserted into the outer side of the first phalanx of the little toe.

* They are separated from each other by strong perpendicular partitions, proceeding from the fascia.

Flexor Brevis Digitorum.—This muscle arises from the under surface of the os calcis, between the two preceding. It passes forwards and divides into four tendons, which run superficial to those of the long flexor. Cut open the sheath which contains them; follow them on to the toes, and you will find that each bifurcates over the first phalanx, so as to allow the long tendon to pass; then the two slips, re-uniting, are inserted into each side of the second phalanx. In fact, the same arrangement prevails as in the fingers.

The three superficial muscles should be reflected from their insertions only, so that they can be replaced if necessary. This done, you bring into view the plantar vessels and nerves, the long flexor tendon of the great toe, and that of the other toes, as well as the flexor accessorius.

Tendon of the Flexor Longus Digitorum.—**Musculus Accessorius.**—Tracing this tendon into the sole, you find that an accessory muscle is attached to it. The muscle arises from the inner side of the os calcis, and has often a second tendinous origin from the outer side of the bone. Its fibres run straight forwards, and are inserted into the fibular side of the tendon, so that their action is not only to assist in bending the toes, but to make the common tendon pull in a straight line towards the heel, which, from its oblique direction, it would not do without this accessory muscle. The common tendon then divides into four, one for each of the four outer toes. These run in the same sheath with the short tendons, and after passing through them are inserted into the base of the ungual phalanx. Respecting the manner in which the tendons are confined by fibrous sheaths, and lubricated by a synovial lining, what was said of the fingers (p. 242) applies equally to the toes.

Lumbricales.—These four little muscles are placed between the long flexor tendons. Each, excepting the most internal, arises from the adjacent sides of two tendons, proceeds forwards, and then sinking between the toes, terminates in an aponeurosis which joins the extensor tendon on the dorsum of the toes. Concerning their use refer to p. 245.

Now trace the long flexor tendon of the great toe. From the groove in the astragalus it runs along the groove in the lesser tuberosity of the os calcis, and then straight to the


base of the last phalanx of the great toe. Observe that it crosses the long flexor tendon of the toes, and that the two tendons are connected by an oblique slip; so that we cannot bend the other toes without the great toe.

Plantar Arteries.—The posterior tibial artery, having entered the sole between the origins of the abductor pollicis, divides into the external and internal plantar arteries.

The *internal plantar* artery is very small: it passes forwards between the abductor pollicis and the flexor brevis digitorum to the base of the great toe, where it terminates in small inosculations with the digital arteries. Its chief use is to supply the muscles between which it runs.

The *external plantar* is the principal artery of the sole, and alone forms the plantar arch. It runs obliquely outwards across the sole towards the base of the fifth metatarsal bone; then, sinking deep from the surface, it bends inwards across the bases of the metatarsal bones, and inosculates with the anterior tibial in the first interosseous space. In the first

Fig. 103.



1. Internal plantar.
2. External do.

part of its course, it lies between the flexor brevis digitorum and the flexor accessorius; in the second it lies very deep beneath the flexor tendons and the adductor pollicis, close to the metatarsal bones. Deeply seated as it appears to be, a part of its curve near the fifth metatarsal bone lies immediately beneath the fascia. Here it might be more easily tied than in any other part of its course.

Its chief branches are the four digital arteries, which arise in the arched part of its course. They supply both sides of the fifth, fourth, third, and the outer side of the second toe; the great toe, and the inside of the second, being supplied by the dorsalis hallucis. Concerning the distribution of the digital arteries, refer to the account given of these arteries in the hand; they are in all respects similar (p. 238).

Besides the digital arteries, the arch gives off small branches, which ascend between the metatarsal bones and inosculate with the dorsal interosseous arteries.

Plantar Nerves.—The posterior tibial nerve divides, like the artery, into an external and internal plantar. The *internal* plantar is generally the larger, and supplies nerves to three toes and a half, like the median nerve in the palm. It also supplies the muscles on the inner side of the sole, the flexor brevis digitorum and the two inner lumbricales. The *external* plantar supplies nerves to the fifth toe and the outer side of the fourth, like the ulnar in the palm. Besides these, it furnishes nerves to the abductor and flexor brevis minimi digiti, the two outer lumbricales, the flexor accessorius, and a considerable branch which accompanies the plantar arch for the supply of the adductor pollicis, the transversalis pedis, and all the interossei.

Deep Muscles.—Having traced the principal vessels and nerves, we must divide them with the flexor tendons near the os calcis, and turn them down towards the toes, so as to expose the deep muscles in the sole. These are, the flexor brevis and the adductor pollicis, the flexor brevis minimi digiti, the transversalis pedis, and the interossei.

Flexor Brevis Pollicis.—This muscle arises from the external cuneiform and the cuboid bone. It proceeds along the metatarsal bone of the great toe, and divides into two portions, which run one on each side of the long flexor tendon, and are inserted by tendons into the sides of the first phalanx of the great toe. In each tendon there is a large sesamoid bone. These bones not only increase the strength of the muscle, but both together form a pulley for the free play of the long flexor tendon.

Adductor Pollicis.—This very powerful muscle arises from the cuboid bone and the third and fourth metatarsal bones. Passing obliquely across the foot, it is inserted through the external sesamoid bone into the outer side of the base of the first phalanx of the great toe. This muscle greatly contributes to support the arch of the foot. Like the adductor of the thumb it should be considered as an interosseous muscle.

Flexor Brevis Minimi Digiti.—This little muscle arises

from the base of the fifth metatarsal bone, proceeds forwards along it, and is inserted into the base of the first phalanx of the little toe.

Transversalis Pedis.—This slender muscle runs transversely across the distal ends of the metatarsal bones. It arises by fleshy slips from the four outer toes, and is inserted into the first phalanx of the great toe with the adductor pollicis, of which it ought to be considered a part.

Interossei.—These muscles are arranged nearly like those in the palm; that is to say, they occupy the intervals between the metatarsal bones, and are seven in number, four being on the dorsal aspect of the foot, and three on the plantar. They arise from the sides of the metatarsal bones, and terminate in tendons which are inserted, some into the inner, others into the outer side of the first phalanges of the toes and their extensor tendons. Their use is to draw the toes to or from each other, and they do this or that according to the side of the phalanx on which they act. Now, if we draw an imaginary longitudinal line through the second toe, we find that all the dorsal muscles draw *from* that line, and the plantar *towards* it. This is the key to the action of them all. A more detailed account of these muscles is given in the dissection of the palm, p. 271. Between the tendons of the interossei, that is, between the distal ends of the metatarsal bones, there are little bursæ to facilitate movement. These bursæ are not without interest. They sometimes become enlarged and occasion painful swellings between the roots of the toes. I have seen two instances; both yielded to iodine.

It now only remains to trace the tendons of the peroneus longus and the tibialis posticus. The tendon of the peroneus longus is the deepest in the sole. We find it running through a groove in the cuboid bone obliquely across the sole towards its insertion into the outer side of the base of the metatarsal bone of the great toe. The tendon is confined in a strong fibrous sheath, which is lined throughout by synovial membrane.

The tendon of the tibialis posticus may be traced over the internal lateral ligament of the ankle, and thence under the head of the astragalus to the scaphoid bone into which it is chiefly inserted. One or two slips are sent off to the cuneiform

bones. Observe that the tendon contributes to support the head of the astragalus, and that for this purpose it often contains a sesamoid bone. This is one of the many beautiful provisions for the solidity of the arch of the foot.

DISSECTION OF THE LIGAMENTS.

Ligaments of the Pelvis.—The sacrum is united to the last lumbar vertebra in the same manner as one vertebra is to another. The same observation applies to a union between the sacrum and the coccyx. Refer, therefore, to the description of the ligaments of the spine, p. 191.

The innominate bones are connected to each other in front, constituting the “pubic symphysis;” posteriorly to the sacrum or the spine, constituting the “sacro-iliac symphysis.”


Pubic Symphysis.—This is secured by 1, an anterior ligament, consisting of oblique and transverse fibres; 2, a posterior ligament, less distinct; 3, a sub-pubic ligament, or “ligamentum arcuatum:” it is remarkably strong, and rounds off the point of the pubic arch; 4, an intermediate fibro-cartilage. If you make a perpendicular section through it, you observe that it consists of concentric layers, and that its general structure resembles that between the bodies of the vertebræ. It acts like a buffer, and breaks the force of shocks passing through the pelvic arch.

Sacro-iliac Symphysis.—This is secured by 1, ligamentous fibres in front; 2, ligamentous fibres much more marked, and collected in bundles behind; 3, a very strong ligament called the ilio-lumbar, which extends from the transverse process of the last lumbar vertebra to the crest of the ilium, fig. 105; 4, the greater sacro-ischiatic ligament, one of enormous strength, which extends from the sacrum and coccyx to the tuberosity of the ischium, fig. 104; and 5, the lesser sacro-ischiatic, which extends from the sacrum and coccyx to the spine of the ischium. The two last not only connect the bones, but also, from their great breadth contribute to block up the lower aperture of the pelvis.

Supposing all the preceding ligaments to be divided, we

still find that the ilium adheres most firmly to the sacrum. If they be forcibly separated, it is then seen how they are connected in front by a layer of cartilage, of which the shape is not unlike that of the ear: above and behind this, is the strong *interosseous* ligament, fig. 105, which contributes powerfully to the security of the sacro-iliac joint.

Fig. 104.



Ligaments of the Hip-joint.—This joint is secured by the form of the bones, and by the strength of the powerful muscles which surround it. Being a perfect ball and socket, one might suppose that it would admit of universal motion; but it is not so: for the disposition of its ligaments restricts its range of motion to those directions only which are most consistent with the maintenance of the erect attitude, and the requirements of this part of the skeleton.

Capsular Ligament.—The most important ligament, and that which presents the most interesting peculiarities, is the *capsular*. It is attached above, to the circumference of the acetabulum at a little distance from the margin; below, to the base of the neck of the femur; but observe, to the base of the neck in *front* only: posteriorly the capsule is not attached to the neck, it merely passes round it like a collar. The reason of this is, that there may be no hindrance to the free flexion of the femur.

[Dr. George K. Smith, of Brooklyn, in a paper on the insertion of this capsule, in relation to intra-capsular fracture, has shown that the insertion of this ligament varies so greatly that no two specimens from different subjects can be said to be inserted at the same place. So wide are the differences, that should a transverse fracture occur at a given distance from the posterior inter-trochanteric line, it may be entirely within the ligament in one case, and in another, on the posterior and inferior surfaces of the neck, be half an inch or more external to the capsule. Some locate the posterior insertion of the capsule at the middle of the posterior surface of the neck, others at the posterior inter-trochanteric line; again, some at various points between these two. The most frequent insertion is into the middle of the posterior and inferior surfaces of the neck; in some, a little nearer the head of the bone, in some, more remote, but never as far distant as the shaft of the bone. The capsules of opposite femurs are the same in their insertions.*]

Another important point about the capsule is, that it is not of equal thickness all round. It is made exceedingly thick and strong in front by a broad ligament which extends from the upper margin of the acetabulum down to the anterior inter-trochanteric line. Now this ligament is the thickest in the body: it is called the accessory ligament of the hip. I have shown the direction of its fibres in fig. 104. What is the purpose of it? Plainly, it acts as a strap to prevent the femur being extended beyond a certain limit; in other words it prevents the body from rolling backwards at the hip.


Lay open the capsule, ascertaining the enormous thickness of it in front, and the strong hold it has upon the bones. This proceeding exposes the cotyloid ligament, or "sucker" of the acetabulum, and the ligamentum teres.

Cotyloid Ligament.—The cotyloid ligament is a piece of fibro-cartilage which is attached all round the margin of the acetabulum. Its use is sufficiently apparent. It not only deepens the cavity, but gradually shelves off, so as to embrace the head of the femur like a sucker. It extends over the notch at the lower part of the acetabulum, and in this situation has received the name of the "transverse" ligament.

* American Med. Times, Dec. 14, 1861.

Ligamentum Teres.—The ligamentum teres is exposed by drawing the head of the femur out of the socket. This liga-

Fig. 105.



VERTICAL SECTION THROUGH THE HIP.

ment is not round, as its name would imply, but somewhat flat and triangular. Its base is attached below, to the borders of the acetabular notch; its apex to the fossa in the head of the femur. To prevent pressure on it, and to allow free room for its play, there is a gap at the bottom of the acetabulum. This gap is not crusted with cartilage like the rest of the socket, but is occupied by a soft, yielding fat. The ligamentum teres is surrounded by the synovial membrane. A

little artery runs up with it to the head of the femur. It is a branch of the obturator, and enters the acetabulum through the notch at the lower part.

[The portion of the ligamentum teres which is inserted into the innominate bone, divides into two parts; one can be traced around the lower edge of the cotyloid notch, until we see it lost in the ischium, between the tuberosity and the edge of the acetabulum; the other division passes to the upper end of the notch, where we leave it at the edge of the acetabulum. In dislocation of the head of the femur into the obturator foramen, the two extremities of this ligament are approximated, not necessarily ruptured. In dislocations at this joint the capsular ligament, together with the accessory ligaments, are always more or less torn, even the ligamentum teres, according to Sir A. Cooper, in dislocation into the obturator foramen. The gemelli, pyriformis, obturatorius, and quadratus femoris, have also been completely torn across. When dislo-

cation takes place into the sciatic notch, the head of the bone rests on the piriformis muscle.]

The chief use of the ligamentum teres is to assist in steadying the pelvis on the thigh in the erect position. In this position, the ligament is vertical, and quite tight, fig. 105 : it therefore prevents the pelvis from rolling towards the opposite side, or the thigh from being adducted beyond a certain point. Another purpose served by this ligament is to limit rotation of the thigh, both inwards and outwards. To satisfy yourself of this you must prepare a joint for the purpose. Make a hole through the bottom of the acetabulum large enough to see the working of the ligament ; leaving the capsule entire.

Thus, then, it is evident that the ligaments of the hip are so arranged, that when we "stand at ease," the pelvis is spontaneously thrown into a position in which its range of motion is the most restricted ; for the accessory ligament of the capsule prevents it from rolling backwards, and the ligamentum teres prevents its rolling towards the opposite side. This arrangement economises muscular force in the balancing of the trunk.

The synovial membrane extends down to the base of the neck of the femur in *front*, but only two-thirds down behind. It is laid upon a thick periosteum.

The head of the thigh-bone is kept in the acetabulum by atmospheric pressure ; the amount of this is, of itself, sufficient to keep the limb suspended from the pelvis, supposing all muscles and ligaments to be divided. When fluid is effused into the hip-joint, the influence of the atmospheric pressure is diminished in proportion ; and the corresponding surfaces of the bones being no longer maintained in accurate contact, it sometimes happens that the head of the femur escapes from its cavity, giving rise to what is called a spontaneous dislocation.

Ligaments of the Knee-Joint.—Looking at the skeleton, one would suppose that the knee-joint is very insecure. But nature has amply made up for this apparent insecurity by powerful ligaments, and by surrounding the joint on all sides with a thick capsule formed by the tendons of the muscles which move it.

Capsular Ligament.—Let us first examine the tendons

concerned in the construction of the capsular ligament. In front is the ligamentum patellæ; on either side are the tendons of the vasti; at the back of the joint no less than four tendons contribute to form the capsule,—namely, the tendons of the gastrocnemius, the tendon of the semi-membranosus, and of the popliteus. It deserves to be noticed that the weakest part of the capsule is near the tendon of the popliteus: here, therefore, matter formed in the popliteal space may possibly make its way into the joint, or *vice versâ*.

Exclusive of the capsule, the proper ligaments of the joint are 1, the lateral; and 2, the crucial in the interior.

Internal Lateral.—This is a broad flat band, which extends from the inner condyle to the inner side of the tibia, a

little below its head. A few of the deeper fibres are attached to the inner semilunar cartilage, and serve to keep it in place. In the performance of the several motions of the joint, there is a certain amount of friction between the ligament and the head of the tibia, and consequently a small bursa is interposed.


External Lateral.—This is a strong round band, which extends from the outer condyle to the head of the fibula. In some instances there is a smaller band a little deeper, called the short external lateral ligament.

Posterior.—This, which is generally called “ligamentum posticum Winslowii,” consists of ex-

pansions derived from the tendons at the back of the joint, chiefly, however, from the semimembranosus. It not only closes and protects the joint behind, but prevents its extension beyond the perpendicular.

The joint should be opened above the patella. Observe the great extent of the fold which the synovial membrane forms

Fig. 106.



1. Internal lateral ligament.
2. External ditto.

above this bone.* The use of it is to allow the free play of the bone over the lower part of the femur. The fold extends higher above the inner than the outer condyle: this accounts for the form of the swelling produced by effusion into the joint.

Folds of Synovial Membrane.—Below the patella a slender band of the synovial membrane proceeds backwards to the space between the condyles. It is called the "*ligamentum mucosum*:" the edges of it are called the "*ligamenta alaria*." These are not true ligaments, but merely the remnants of the partition, which, in the early stage of the joint's growth, divided it into two lateral halves.


[The synovial membrane does not pass over the articular surfaces of the joint, as some authors have stated, except perhaps, at the earlier periods of life.]

Immediately outside the synovial membrane there is always more or less fat: this is especially accumulated under the *ligamentum patellæ*. Its use is obviously to fill up vacuities, and to mould itself to the several movements of the joint.

Crucial.—The crucial ligaments, so named because they cross like the letter X, extend from the medial side of each condyle to the head of the tibia. The anterior, *a, b*, passes from the outer condyle forwards to the fossa in front of the spine of the tibia. The posterior, *c, d*, is best seen from behind: it goes from the inner condyle backwards to the fossa behind the spine of the tibia.

Inter-Articular or Semi-Lunar Cartilages.—Between the condyles and the articular surfaces of the tibia there are two incomplete rings of fibro-cartilage, shaped like the letter C. They answer many purposes. They deepen the articular surfaces of the tibia; their mobility and elasticity enable them to adapt themselves to the condyles in the several movements of the joint; they distribute pressure over a greater

Fig. 107.



- a. b.* Anterior crucial ligament.
- c. d.* Posterior ditto.
- e.* Tibio-fibular.

* In performing operations near the knee, the joint should always be bent in order to draw the synovial fold as much as possible out of the way.

surface, upon the principle that a porter places a knot on his shoulders to carry his load with greater ease ; they equalize pressure and prevent shocks. They are thickest at the circumference, and gradually shelve off to a thin margin : thus they fit in nicely between the bones, and adapt a convex surface to a flat one, as shown in fig. 106. Their form is suited to the condyles ; the inner being oval, the outer more circular. The ends of each are firmly attached to the pits in front and behind the spine of the tibia ; but the ends of the internal one are attached further from the spine than the external. Besides this, the cartilages are mutually connected in front by a thin "*transverse*" ligament ; and their circumference is attached round the head of the tibia by fibrous tissue (called the "*coronary*" ligament), yet not so closely as to restrict their range of motion.*

Action of the Ligaments.—The action of the ligaments of the knee-joint is a study of itself. Their respective points of attachment are such, that when the joint is extended, all

Fig. 108.



the ligaments are tight, to prevent extension beyond the perpendicular ; thus muscular force is economized. But when the joint is bent the ligaments are relaxed, enough to admit a slight rotatory movement of the tibia. This movement is more free in an outward than an inward direction ; and it is effected, not by rotation of the tibia on its own axis, but a rotation of the outer head round the inner.

Rotation outwards is produced by the biceps ; rotation inwards by the popliteus.

It is curious that the crucial ligaments, though placed inside the joint, answer the same purposes that the coronoid process and the olecranon do, in the construction of the elbow. They make the tibia *slide* properly forwards and backwards. The anterior especially limits extension ; the posterior, flexion. They not only prevent dislocation in front or behind, but they

* Of the two cartilages the external has the greater freedom of motion, because in rotation of the knee the outer side of the tibia moves more than the inner. Consequently, it is not in any way connected to the lateral ligament ; so far from this, it is separated from it by the tendon of the popliteus, of which the play is facilitated by a bursa communicating freely with the joint. For this reason the external cartilage is more liable to dislocation.

prevent lateral displacement, since they cross each other like braces, as shown in fig. 108.

[When complete dislocation of the patella exists, it rests upon the external condyle, and is thus held by the action of the quadriceps extensor muscle. The synovial membrane, from its extent, is not ruptured. When the tibia is completely dislocated backwards, the attachment of the patella, either above or below, must give way. The crucial and posterior ligaments are also torn. The limb is kept in this position by the flexor of the leg upon the thigh.]

Ligaments Connecting the Tibia and Fibula.—There is a distinct joint between the upper ends of the tibia and fibula, although it admits of but little movement. It is firmly secured by oblique ligaments in front and behind (p. 481). The contiguous surfaces of the bones are crusted with cartilage. In the great majority of instances its synovial membrane is independent; but we now and then meet with cases in which it communicates with the synovial membrane of the knee.

Interosseous Membrane.—The shafts of the bones are connected by the interosseous membrane. The chief purpose of this is to afford additional surface for the attachment of muscles. Its component fibres cross each other like the letter X, and have openings here and there for the passage of blood-vessels.


The lower ends of the tibia and fibula are most firmly connected, for it is essential to the solidity of the ankle-joint that there be no motion between them. They are secured by an oblique ligament in front and behind, and by strong ligamentous fibres which connect their contiguous surfaces, and answer the purpose of rivets. Besides these, a strong fibro-cartilaginous ligament proceeds from the end of the fibula, and is attached along the posterior border of the articular surface of the tibia. The object of it is to deepen the excavation of the tibia, and enable it to adapt itself more accurately to the articular surface of the astragalus.

Ligaments of the Ankle-Joint.—From the form of the bones, it is obvious that the ankle is a hinge joint; consequently, like all others of the kind, its chief security depends

upon the great strength of its lateral ligaments. However, the hinge is not so perfect but that it admits of a slight rotatory motion, of which the centre is on the fibular side, and therefore the reverse of that in the case of the knee.

Internal Lateral.—This ligament, sometimes called, from its shape, “deltoid,” is exceedingly thick and strong, and makes up for the comparative shortness of the malleolus on this side (fig. 109). The great strength of it is proved by the

Fig. 109.



1. Plantar fascia.
2. Calcaneo-scapoid or elastic ligament which supports the head of the astragalus.
3. Internal lateral ligament, called from its shape deltoid.

fact, that in dislocations of the ankle inwards, the summit of the malleolus is more likely to be broken off than the ligament to be torn. It arises from a deep excavation at the apex of the malleolus, radiates from this point, and is inserted into the side of the astragalus, also into the os calcis, and the scaphoid bone. Besides this, some of its fibres are inserted into, and become identified with, the calcaneo-scapoid ligament, so as to brace it up internally (fig. 109).

External Lateral.—This ligament consists of three distinct parts,—an anterior, a posterior, and a middle (fig. 110). All three arise from an excavation near the summit of the external malleolus; the first two are inserted into the front and the back of the astragalus; the middle into the os calcis.

The closure of the joint is completed, in front and behind,

by a thin capsular membrane attached to the bones near their articular surfaces, and sufficiently loose to permit the necessary range of motion.

Besides flexion and extension, the construction of the ankle-joint admits of a very slight lateral movement: but this is only permitted in the extended state of the joint. This is useful to us in the direction of our steps. In adaptation to this movement the internal malleolus is made much shorter than the outer; it is not so tightly confined by its ligaments, and its articular surface is part of a cylinder.

Open the joint to see that the breadth of the articular surfaces of the bones is greater in front than behind. The object of this is to render the tibia less liable to be dislocated forwards over the astragalus. Whenever this happens, the astragalus must of necessity become firmly locked between the malleoli.

Ligaments Connecting the Bones of the Foot.—The astragalus is the key-stone of the arch of the foot, and supports the whole weight of the body. It articulates with the os calcis and the os scaphoides in such a manner as to permit the abduction and adduction of the foot, so useful in the direction of our steps.

The astragalus articulates with the os calcis by two distinct surfaces, respecting which it is necessary to remark, that the anterior is slightly convex, and the posterior slightly concave. This adaptation of itself contributes much to prevent the separation of the bones.

Interosseous Ligament. — But their principal bond of union is a very strong *interosseous* ligament which occupies the interval between them (fig. 110).

In the skeleton you observe that the head of the astragalus

Fig. 110.





DIAGRAM OF EXTERNAL LATERAL LIGAMENT

1. Anterior part.
2. Posterior part.
3. Middle part.
4. Interosseous ligament.

articulates in front with the os scaphoides, but that a part of it is unsupported below. What, then, supports it here in the perfect foot?

Calcaneo-scaphoid.—An enormously strong and elastic ligament which extends from the os calcis to the os scaphoides

Fig. 111.



1. Calcaneo-scaphoid ligament.
2. Calcaneo-cuboid ligament.


(fig. 111). These bones, together with the ligament, form a complete socket for the reception of the head of the astragalus; it is this joint, and not the ankle, which permits the abduction and adduction of the foot. The chief peculiarity about the ligament alluded to is its elasticity. It acts in all respects like a spring, and allows to the key-stone of the arch a certain amount of play which is obviously of great service in preventing concussion of the body. Whenever this ligament loses its elastic property, as is often the case with those who are in the habit of carrying heavy burdens, the arch of the foot gradually yields, and the individual becomes flat-footed.

The os calcis articulates with the os cuboides pretty nearly on a line with the articulation between the astragalus and the os scaphoides.

Calcaneo-cuboid.—The bones are most firmly connected by means of a powerful ligament in the sole, called the *calcaneo-cuboid*, or "*ligamentum longum plantæ*." Some of its fibres extend forwards as far as the third and fourth metatarsal bones.

It would be only tedious repetition to enumerate individu-

Fig. 112.



Interosseous ligaments of the wedge bones.

ally the several ligaments which connect the remaining bones of the tarsus and metatarsus. Let it then suffice to say, that they are firmly braced together by very strong ligaments, both on their dorsal and their plantar surfaces, and by interosseous ligaments which extend be-

tween their contiguous surfaces like rivets. Though there is

very little motion between any two bones, the collective amount is such that the foot is enabled to adapt itself accurately to the ground ; pressure is more equally distributed, and consequently there is a firmer basis for the support of the body. Being composed, moreover, of several pieces, each of which possesses a certain elasticity, the foot gains a general springiness and solidity which could not have resulted from a single bone.

Synovial Membranes of the Tarsus. — Exclusive of the ankle-joint and the phalanges of the toes, the bones of the foot are provided with six distinct synovial membranes,—namely :

a. Between the posterior articular surface of the astragalus and os calcis.

b. Between the head of the astragalus and the bones composing its socket.

c. Between the os calcis and os cuboides.

d. Between the inner cuneiform bone and the metatarsal bone of the great toe.

e. Between the three cuneiform and the adjoining bones (the great toe excepted).

f. Between the os cuboides and the fourth and fifth metatarsal bones.

The phalanges of the toes are connected in all respects like those of the fingers. See, therefore, the description given in the dissection of the hand (p. 285).

THE DISSECTION OF THE BRAIN.

THE most convenient manner of removing the brain is to cut through the scalp, from one ear to the other, so that the anterior part of the scalp may be detached from the skull and pulled down over the face ; the posterior over the back of the head. The skull-cap should then be taken off about half an inch above the supra-orbital ridges. It is better to saw only through the outer table of the skull, and to break through the inner with a chisel. In this way the dura mater and the brain are less likely to be injured.

Three Membranes of the Brain.—Previous to the examination of the brain itself you should study the structure and uses of the three membranes by which it is surrounded,—namely, a tough fibrous membrane termed the “dura mater;” a serous membrane termed the “arachnoid;” and a very vascular membrane termed the “pia mater.”

Dura Mater.—This is the first membrane exposed after the removal of the skull-cap. It is so called from the notion that it gave rise to all the other membranes in the body. Its remarkably tough and fibrous structure adapts it exceedingly well to the four purposes which it serves: 1, it forms the internal periosteum of the skull; 2, it forms, for the support of the lobes of the brain, three partitions, namely, the falx cerebri, the falx cerebelli, the tentorium cerebelli: 3, it forms the sinuses or venous canals which return the blood from the brain: 4, it forms sheaths for the nerves as they leave the skull.

Meningeal Arteries.—These arteries ramify between the dura mater and the skull. Their course may be traced by the grooves which they make in the bones. The most important is the “*arteria meningeae media*,” a branch of the internal maxillary (p. 61). Entering the skull through the foramen spinosum, it winds along a deep groove in the sphenoid, and the anterior inferior angle of the parietal bones, to the top of the skull, giving off branches in all directions. It is accompanied by two veins. The other meningeal arteries are of insignificant size. The *anterior* come from the ethmoidal branches of the ophthalmic; the *posterior* come from the occipital, and enter the skull through the foramen jugulare.

The position of the meningeal arteries renders them liable to injury in fractures of the skull: hence extravasation of blood between the skull and dura mater is one of the common causes of compression of the brain.

Three Partitions Formed by Dura Mater.—Of the partitions formed by the dura mater for the support of the lobes of the brain, two are vertical, and separate the two hemispheres of the cerebrum and those of the cerebellum; the third is nearly horizontal, and supports the posterior cerebral lobes.

Falx Cerebri.—The great vertical partition is named, from

its resemblance to the blade of a sickle, "*falx cerebri*." It divides the cerebrum into two hemispheres—a right and a left. It begins in a point attached to the crista galli, and gradually penetrates deeper as it extends backwards. From its base or broadest part proceeds the horizontal partition, named "*tentorium cerebelli*."


Tentorium Cerebelli.—This forms an arch for the support of the posterior cerebral lobes, so that they may not press upon the cerebellum beneath them. Observe that the tentorium is attached to the transverse ridge of the occipital bone, to the petrous part of the temporal, and to the posterior clinoid processes of the sphenoid.

Falx Cerebelli.—The little partition which separates the lobes of the cerebellum is called "*falx cerebelli*." It is placed vertically in the same line with the falx cerebri, and its point is attached to the edge of the foramen magnum.

Sinuses of the Dura Mater.—It is one of the peculiarities of the cerebral circulation that the blood is returned through canals or sinuses formed by the dura mater. These canals are produced by a splitting of the dura mater into two layers as shown in fig. 113, where 1, represents a vertical section through the superior longitudinal sinus.

They are lined by the same smooth membrane as the rest of the venous system. Since their walls consist of such unyielding structure, and are always on the stretch, it is obvious that they are admirably adapted to resist

the pressure of the brain. There are fifteen of these sinuses; five are pairs, and five are single, as follow :—



The five pairs of sinuses are,—

- The lateral.
- The superior petrosal.
- The inferior petrosal.
- The cavernous.
- The occipital.


The five single sinuses are,—

- The superior longitudinal.
- The inferior longitudinal.
- The circular.
- The transverse.
- The straight.

All of these eventually discharge their blood into the internal jugular veins.

Superior Longitudinal Sinus.—This runs along the upper

edge of the falx cerebri (fig. 114). It begins very small at the crista galli, gradually increases in size in its course backwards, and opposite the tubercle of the occipital bone divides into



1. Superior longitudinal sinus.
2. Inferior longitudinal sinus.
3. Straight sinus.
- 4, 4. Venæ galeni.

the right and left lateral sinuses, the right being in general the larger. Besides numerous veins from the cancellous texture of the skull-cap, the superior longitudinal sinus receives large veins from each hemisphere of the cerebrum. It is interesting to observe that these veins run from behind forwards, contrary to the current of blood

in the sinus, and that they pass through the wall of the sinus very obliquely, like the ureter into the bladder. The probable object of this oblique entrance is to prevent regurgitation of blood from the sinus into the veins of the brain.

Cut open the superior longitudinal sinus: observe that it is triangular, and that its cavity is intersected in many places by slender fibrous cords, termed "*chordæ Willisii*."* Their precise use is not understood.

Glandula Pacchioni.—In the neighborhood of the superior longitudinal sinus we meet with small granulations, sometimes lying singly, sometimes in clusters. They are termed "*glandulæ Pacchioni*,"† and are found in three distinct situations:—1, on the outside of the dura mater, often so large as to occasion depressions in the bones; 2, on the surface of the pia mater; 3, in the interior of the longitudinal sinus, covered by its lining membrane. Their size, number, and appearance differ in different subjects. Nothing is determined concerning their precise nature; but it is presumed that they are morbid products, since they are never observed in very young subjects.

* So called after our countryman Willis, who first described them in his work *De Cerebri Anatome*, 1664.

† After the Italian anatomist, who first described them in 1705. These bodies would appear to originate in the subarachnoid cellular tissue, whence they, in their growth, either perforate the dura mater and hollow out the bones, or make their way into the longitudinal sinus.

Lateral Sinuses.—These are the two great sinuses through which all the blood from the brain is returned into the internal jugular veins. Their course is well marked in the dry skull. The right is commonly the larger. Each proceeds at first horizontally outwards, inclosed between the layers of the tentorium, along a groove in the occipital bone and the infe-

Fig. 115.




DIAGRAM OF THE VENOUS SINUSES AT THE BASE OF THE SKULL.

rior angle of the parietal; it then descends along the mastoid portion of the temporal bone, and again indenting the occipital, turns forward to the foramen lacerum posterius, and terminates in the internal jugular vein.*

Inferior Longitudinal Sinus.—This is of small size. It terminates in the straight sinus at the anterior margin of the tentorium (fig. 114).

* It has in some subjects another outlet, through the foramen mastoideum, or else the posterior condyloid foramen.

Straight Sinus.—This may be considered as the continuation of the preceding. It runs along the line of junction of the falx cerebri with the tentorium cerebelli, and terminates at the divergence of the two lateral sinuses. It receives the two “venæ Galeni” (fig. 114), which return the blood from the lateral ventricles of the brain.

Cavernous Sinus.—This is so called because its interior is intersected by numerous threads. It extends along the side of the body of the sphenoid bone, outside the internal carotid artery. It receives the ophthalmic vein which leaves the orbit through the foramen lacerum orbitale; and it communicates with the circular sinus which surrounds the pituitary gland (p. 159).

Circular Sinus.—This surrounds the pituitary gland (P in the diagram) and communicates on each side with the cavernous.

Petrosal Sinuses.—These lead from the cavernous to the lateral. There are two on each side. The *superior* runs along the upper edge of the petrous bone; the *inferior* along the suture between the petrous and occipital bones.

Transverse Sinus.—This extends from one inferior petrosal to the other, across the basilar process of the occipital bone.

Occipital Sinuses.—These are very small. They run round the margin of the foramen magnum, and open into the divergence of the lateral sinuses.*

Arachnoid Membrane.—This second investment forms the smooth surface of the brain exposed after the removal of the dura mater. It is named “arachnoid” from the delicacy of its tissue, which gives it a resemblance to a spider’s web. It is a serous membrane, and, like all others of the kind, forms a closed sac, one part of which, called the “parietal” layer, lines the under surface of the dura mater; the other, called

* The meeting of the several sinuses opposite the spine of the occipital bone is termed the “*torcular Herophili*,” after the celebrated anatomist who first described it. It is a kind of triangular reservoir, with the base below, and presents six openings—namely, that of the superior longitudinal sinus, those of the two lateral and of the two occipital, and that of the straight sinus. The term “*torcular*” is an incorrect version of the original word *σωλήν* (a canal or gutter), employed by Herophilus.

the "visceral," is reflected over the brain; this reflection takes place along the nerves which leave the base of the skull.

The inner surface of this membrane is perfectly smooth, and lubricated by sufficient moisture to prevent friction, since the brain is alternately rising and falling with a slight pulsating movement, caused in part by the action of the heart, in part by the process of respiration. The parietal layer is of such extreme tenuity that it can only be demonstrated in very early life: indeed, it consists of little more than a layer of epithelium; but the visceral layer is sufficiently obvious. Observe that it is colorless and transparent, and that it is spread uniformly over the surface of the brain, without insinuating itself between the convolutions. On account of its extreme tenuity and its close adhesion to the pia mater, it cannot be readily separated from this membrane; but there are certain parts at the base of the brain termed "sub-arachnoid" spaces, where the arachnoid membrane can be seen distinct from the subjacent tunic.

Pia Mater.—This is the immediate investing membrane of the brain. It is extremely vascular: indeed, its chief use is to form a bed in which the blood-vessels may divide and subdivide before they enter the brain.* It sinks down to the bottom of all the convolutions, and penetrates into the lateral ventricles for the supply of the interior, forming what is called the "velum interpositum" and the "choroid plexus."

Sub-Arachnoid Spaces and Fluid.—The arachnoid membrane is separated in certain situations from the pia mater by a watery fluid contained in the meshes of very delicate areolar tissue. Such interspaces are termed "sub-arachnoid." There is one of these in the longitudinal fissure of the cerebrum, because the arachnoid does not go down to the bottom of it, but passes across below the edge of the falx, at a little distance above the corpus callosum. Again, at the base of the brain there are two of considerable size:—one, called the "middle sub-arachnoid space," is situated between the Pons Varolii

* It is one of the protective provisions of the circulation in the delicate substance of the brain, that the arteries break up into small ramifications on the surface before they penetrate its interior. When carefully injected, it is seen that the blood-vessels of the pia mater form minute angular spaces, from which the vessels pass off at right angles into the brain.

and the commissure of the optic nerves; the other, termed the "posterior," is situated between the cerebellum and the medulla oblongata. In the examination of the spinal cord, too, it will be seen that there is a considerable interval between the arachnoid and the pia mater, also occupied by fluid. The purpose of this fluid is not only to fill up space like fat in other parts, but mechanically to protect the nervous centres from the violent shocks and vibrations to which they would otherwise be liable.

The base of the brain may, in truth, be said to be supported by a bed of water, which insinuates itself into all the inequalities of the surface, and surrounds all the nerves down to the foramina, through which they pass. One sees this water oozing through the ear, in some cases of fracture through the base of the skull, where the fracture runs across the meatus auditorius internus and the petrous portion of the temporal bone. This symptom is the almost certain presage of a fatal result.

Arteries of the Brain.—The brain is supplied with blood by the two internal carotid and the two vertebral arteries.

Internal Carotid.—This artery enters the skull through a canal in the petrous portion of temporal bone, mounts up very tortuously by the side of the body of the sphenoid, and, after giving off the ophthalmic, divides into the anterior and middle cerebral arteries, for the supply of the anterior and middle cerebral lobes.

a. The *anterior cerebral* artery sinks into the longitudinal fissure between the hemispheres, curves round the front part of the corpus callosum, and then runs backwards along the upper surface of it. It distributes branches in all directions. The anterior cerebral arteries of opposite sides run close together, and at the base of the brain are connected by a short branch called the "*anterior communicating*" (fig. 116).

b. The *middle cerebral* artery runs deep in the fissure of Sylvius, distributing large branches to the anterior and middle lobes.

Vertebral Artery.—This artery, after winding backwards along the arch of the atlas, enters the skull through the foramen magnum, and unites with its fellow at the lower border of the pons to form the "*basilar*." This single vessel pro-


ceeds along the middle of the pons, and divides at the upper border of it into the two posterior cerebral arteries for the supply of the posterior cerebral lobes (fig. 116).

In its course along the pons, the basilar gives off on either side—

a. The little *auditory* artery, which enters the meatus auditorius with the auditory nerve.

b. The *superior and inferior cerebellar* arteries, for the supply

Fig. 116.



of the upper and lower surfaces of the cerebellum.

c. The two *posterior cerebral*—the terminal branches (fig. 116)—run backwards, one on the under surface of each posterior cerebral lobe, dividing into numerous branches, which

ultimately inosculate with the other cerebral arteries. But the principal inosculating branch of the posterior cerebral is called the "*posterior communicating*." It proceeds straight forwards to the internal carotid, and thus establishes at the base of the brain that free inosculature of the large arteries called the "*circle of Willis*."

Circle of Willis.—The circle of Willis, as seen in fig. 116, is completed in front, by the anterior communicating artery (between the two anterior cerebral); behind by the posterior communicating (from the posterior cerebral to the carotid). The tortuosity of the large arteries before they enter the brain is intended to mitigate the force of the heart's action; and the circle of Willis provides a free supply of blood from other quarters, in case any accidental circumstance should stop the flow of blood in any of the more direct channels.*

Peculiarities of the Cerebral Circulation.—Besides the remarkable "circle of Willis" just alluded to, we have other peculiarities in the circulation in the brain: namely, the tortuosity of the four great vessels as they enter the skull—their breaking up into minute arteries in the pia mater before they enter the substance of the brain—the formation of the sinuses, p. 489—the great minuteness of the capillaries, and the extreme thinness of their walls; a fact which accounts for the slowness with which a congested brain recovers itself.

General Division of the Brain.—The collective mass of nervous matter, designated under the common term brain, is divided into three parts—the "*cerebrum*," or intellectual brain, which occupies the whole of the upper part of the cranial cavity; the "*cerebellum*," or little brain, which occupies the lower and back part beneath the tentorium; and the "*medulla oblongata*," situated at the base beneath the cerebellum. This

* In many of the long-necked herbivorous quadrupeds a beautiful provision has been made, in the disposition of the internal carotid arteries, for the purpose of equalizing the force of the blood supplied to the brain. The arteries, as they enter the skull, divide into several branches, which again unite, so as to form a remarkable network of arteries, called by Galen, who first described it, the "*rete mirabile*." The object of this evidently is to moderate the rapidity with which the blood would otherwise enter the cranium, in the different positions of the head, and thus preserve the brain from those sudden influxions to which it would under other circumstances be continually exposed.

last part passes out of the skull through the foramen magnum, and is continuous with the spinal cord.

[The weight of the encephalon in the adult male, varies from 46 to 53 ounces. In the female, from 41 to 47 ounces. The encephalon appears to reach its maximum weight at about the fortieth year.]

Cerebrum.—The cerebrum, in man, is so much more developed than the other parts of the brain, that it completely overlaps them. It is of an oval form. It is divided, in the middle line, by a deep "*longitudinal fissure*," into two equal halves, termed the right and left hemispheres.* This fissure is occupied by the falx cerebri (p. 488). The surface of each hemisphere is mapped out by tortuous eminences, termed "the convolutions," which are separated from each other by deep furrows. Many of these furrows are occupied by the larger veins in their course to the sinuses; others are filled by sub-arachnoid fluid. In old age, or in cases of disease where the convolutions are shrunk and atrophied, this fluid sometimes accumulates in considerable quantity. The convolutions themselves are merely folds of the brain for the purpose of increasing the extent of surface upon which may be laid the grey or vesicular nerve substance, now generally admitted to be the source of power. They are not precisely alike on both sides. Their number, disposition, and depth, vary a little in different individuals, and, to a certain extent, may be considered as an index of the degree of intelligence.†


The under surface of each cerebral hemisphere is not only convoluted like the upper, but presents in addition three lobes (fig. 117), an anterior, middle, and posterior, which fit into the

* Examples are now and then met with where the longitudinal fissure is not exactly in the middle line; the consequence of which want of symmetry is, that one hemisphere is larger than the other. Bichat (*Recherches Physiologiques sur la Vie et la Mort*, Paris, 1829) was of opinion that this anomaly exercised a deleterious influence on the intellect. It is a remarkable fact that the examination of his own brain after death, went to prove the error of his own doctrine.

† Those who wish to investigate the cerebral convolutions in their simplest form in the lower classes of mammalia, and to trace them through their successive complications and arrangement into groups as we ascend in the higher classes, should consult M. Leuret, *Anatomie Comparée du Système Nerveux considérée dans ses Rapports avec l'Intelligence*: Paris, 1839; also M. Foville, *Traité de l'Anat. du Système Nerveux, &c.*: Paris, 1844.

base of the skull. The anterior lobe rests upon the roof of the orbit, and is separated from the middle by a cleft called the "*fissure of Sylvius*," which receives the lesser wing of the

Fig. 117.



- 1, 2, 3. Anterior, middle, and posterior lobes of the cerebellum.
4. Cerebellum.
5. Pons Varolii.
6. Medulla oblongata.

sphenoid bone. The middle lobe occupies the middle fossa of the skull formed by the sphenoid and temporal bones. The posterior lobe rests on the arch of the tentorium. Between this and the middle there is no evident boundary; indeed some anatomists, among whom may be mentioned the illustrious Haller, enumerate only two lobes.

By separating the middle from the anterior lobe, we see, deep in the fissure of Sylvius, what is called the "*island of Reil*." It is merely a portion of the under surface of the

hemisphere, mapped out by small and deep convolutions, which dove-tail with those adjoining. It corresponds with the under surface of the corpus striatum, and might, therefore, be called the lobe of that ganglion.

Nomenclature of the Parts at the Base of the Brain.

—We must now notice the several objects seen at the base of the brain in the middle line, proceeding in order from the back part (fig. 116, p. 495). In this description we omit the cerebral nerves. These will be examined hereafter.

Immediately above the medulla oblongata is a square protuberance called the "*Pons Varolii*." This is a transverse commissure which connects the two lobes of the cerebellum, and lies over the basilar groove of the occipital bone. From the anterior border of the pons two round cords of nervous matter, about half an inch thick, diverge from each other,—one towards each hemisphere of the cerebrum. These are the "*crura cerebri*." Winding round the outer side of each crus is a white band—the "*tractus opticus*," or root of the optic nerve. These tracts converge, and uniting in the middle line, constitute the *commissure of the optic nerves*. Between the crura cerebri the surface is perforated by a number of blood-vessels; hence this spot is called the "*locus perforatus*

medius,"—medius, as contra-distinguished from another part of the brain's base presenting similar perforations, situated at the root of the fissure of Sylvius, and called the "*locus perforatus anticus*."* In front of the *locus perforatus medius* are two round white bodies, called the "*corpora mammillaria* or *albicantia*:" in front of these is a simple elevation of the surface, consisting of grey matter, called the "*tuber cinereum*," upon which the commissure of the optic nerves seems, as it were, to ride. From the *tuber cinereum*, and just behind the optic commissure, a conical tube, of a reddish color, termed the "*infundibulum*," descends to the pituitary gland which occupies the *sella turcica* of the sphenoid bone. In front of the optic commissure the *tuber cinereum* is continued forwards under the form of a thin grey layer (*lamina cinerea*) so as to join the anterior end of the corpus callosum. Now these several objects,—namely, the *locus perforatus medius*, *mammillary bodies*, *tuber cinereum*, *infundibulum*, optic commissure, and *lamina cinerea*—collectively constitute the floor of the third ventricle of the brain. Lastly, by gently separating the anterior lobes of the cerebrum, we see the anterior extremity of the corpus callosum, or great transverse commissure which connects the two hemispheres of the cerebrum.

Examination of the Interior.—Let the brain be laid upon its base. We have already noticed that the hemispheres of the cerebrum are divided by a deep longitudinal fissure. By gently separating the hemispheres, we observe at the bottom of this fissure a white band of nervous matter: this is the great transverse commissure of the cerebrum, termed "*corpus callosum*."

White and Grey Matter.—Slice off the hemisphere down to the level of the corpus callosum.† The cut surface presents a mass of white substance surrounded by an undulating layer of grey matter of about one-eighth of an inch thick. Since

* The large number of arteries which enter the brain in these respective situations are destined to supply the two great masses of grey matter in the interior—namely, the optic thalamus and the corpus striatum. The former corresponds with the *locus perforatus medius*, the latter with the *locus perforatus anticus*.

† The section of the brain at the level of the corpus callosum was called by the old anatomists "*centrum ovale majus*:" at a higher level, the section was called "*centrum ovale minus*."

this grey matter forms a sort of bark round the white, it is often called the cortical substance.* Notice the depth of the furrows between the convolutions: they penetrate generally to about one inch; but their depth varies in different individuals: hence it follows that two brains of apparently equal size may be very unequal in point of extent of surface for the grey matter, and therefore in amount of power. To the unassisted eye the cortical layer appears to be made up entirely of grey substance; but with the microscope it may be seen to consist of six layers,—three of grey alternating with three of white.†

Corpus Callosum.—This transverse portion of white substance is the chief connecting medium between the two hemispheres.‡ It is called the great commissure of the cerebrum. Observe that it is about four inches long, and nearer to the front than to the back part of the brain. Its surface is not

* There are two kinds of nervous matter—the grey or vesicular, and the white or fibrous. The grey is exceedingly vascular, and contains the nerve-vesicles, which are generally allowed to be the source of power. The white consists of extremely delicate nerve fibres, is scantily provided with blood-vessels, and is probably the mere conductor of power.


The intense vascularity of the grey matter has been demonstrated by Mr. Quekett and Mr. Smee, who have had the kindness to allow the author to inspect their beautiful preparations. Mr. Smee's injections were made by using a solution of carmine in ammonia mixed with size. The minute arteries enter the surface of the grey matter, and break up in it into a network of the capillaries. These capillaries are peculiar in being quite naked, and not supported, as in other parts, by cellular tissue. Hence it is they are so liable to give way, and allow their contents to escape: hence it is they are fenced round by so many provisions calculated to break the force of the current of blood—namely, the tortuosities of the arteries at the base of the brain, their subdivisions in the pia mater, and the peculiar arrangement of the sinuses.

† These six layers cannot always be demonstrated. Sometimes there are only four, and they are commonly more manifest in the posterior lobes. The external layer is always white. For an account of them, and the best mode of examining them, see Baillarget, *Recherches sur la Structure de la Couche Corticale des Circonvolutions du Cerveau*, insérées dans les *Mémoires de l'Académie de Médecine*, 1840.

‡ In a brain properly hardened by spirit, the fibres may be traced congregating towards the corpus callosum from both hemispheres: hence they were called by Gall the *converging* fibres of the brain. This anatomist applied the above name to all commissural fibres. Those fibres, on the other hand, which ascended from the medulla oblongata into the hemispheres, he named the *diverging*. *Anat. et. Phisiol. du Système Nerv.*: Paris, 1810.

flat, but slightly arched from before backwards. One or more longitudinal ridges, called by the old anatomists the "*raphé*" run along the middle of it (fig. 118). In a fresh brain we can evidently see the transverse fibres, of which the commissure is composed, extending between the hemispheres; these are the "*lineæ transversæ*" of the old anatomists. Notice also the anterior cerebral arteries.

Fig. 118.



UPPER SURFACE OF CORPUS CALLOSUM.

- 1, 1. Lineæ transversæ.
2. Raphé.
3, 3. Anterior cerebral a.

Fig. 119.




DIAGRAM OF LAMINA CINEREA.


- 1, 1. Peduncles of corpus callosum.
2. Lamina cinerea.
3. Commissure of optic nerves.

The anterior part of the corpus callosum turns downwards and backwards, forming a bend called its knee, towards the base of the brain. It becomes gradually thinner, and terminates in two peduncles, which diverge from each other, and are lost, one in each fissure of Sylvius. Between these crura is placed the lamina cinerea (fig. 119). The posterior part of the corpus callosum terminates in a thick round border, under which the pia mater enters into the interior of the ventricles. We cannot obtain a satisfactory view of the arch formed by the corpus callosum, of its terminations in front and behind, and of the relative thickness of its different parts, without

making a perpendicular section through a fresh brain, as shown in diagram 120.*

Lateral Ventricles.—In the next place examine the two cavities called the lateral ventricles, situated, one in each hemisphere of the brain, beneath the corpus callosum. A longitudinal incision should be made on each side, about half an inch from the raphé of the corpus callosum. Be careful

Fig. 120.



VERTICAL SECTION THROUGH THE CORPUS CALLOSUM, AND PARTS BELOW.

not to cut too near the middle line, in order to preserve the delicate partition which descends from the under surface of the corpus callosum, and separates the ventricles from each other. We soon come into these cavities, and, by means of a director, they should be fully laid open. Notice their general form, and then the several objects to be seen in them.

* The corpus callosum is more or less developed in all mammalia, but is absent in birds, reptiles, and fish. It is not absolutely essential to the exercise of the intellect, for it has been found absent in the human subject without any particular intellectual deficiency. See cases recorded by Reil, *Archiv. für die Phys. t. xi.*; and Wenzel, *de plenior Struct. Cereb.* p. 302.

What are they?—The lateral ventricles are two cavities in the general mass of the brain, occasioned by the enormous enlargement and folding backward of the cerebral lobes over the other parts of the central nervous axis. They contain a serous fluid, which, even in a healthy brain, sometimes exists in considerable quantity, but, when preternaturally accumulated, constitutes one form of the disease termed hydrocephalus. The very delicate lining membrane which secrets this fluid is quite distinct from the arachnoid on the outside of the brain, and the epithelium on the surface of it is provided with cilia, of which the vibratile movement may be distinctly seen, especially in the embryo.

With regard to the general shape of the ventricles, they may be compared to two crescents with their backs to each other. Each ventricle extends into the three lobes of which the cerebral hemisphere is composed. That part which extends into the anterior lobe is called the *anterior horn*; it slightly diverges from its fellow on the opposite side. The *posterior horn* may be traced into the posterior lobe; this converges to its fellow.* The *middle horn* runs into the middle lobe, descends towards the base of the brain, making a very curious curve, like a ram's horn,—that is, in a direction backwards, outwards, downwards, forwards, and inwards; the initial letters of which make the memorial word “bodfi.” We can easily follow its windings by cutting through the substance of the middle lobe.


What they contain.—Suppose the roof of the lateral ventricles—in other words the corpus callosum—to be removed, what have we to remark on the floor of their interior? Beginning from the front we notice—1, a grey elevation of a pyramidal form, called the *corpus striatum*; 2, behind this, a white elevation called the *thalamus opticus*. 3. Along the boundary line between the corpus striatum and the thalamus opticus runs a slender, transparent, horn-like substance, called

* The posterior horns are not always equally developed in both hemispheres, and sometimes they are absent in one or in both. They are only found in the brain of man and the quadrumana.

In the carnivora, ruminantia, solipeda, pachydermata, and rodentia, the lateral ventricles are prolonged into the largely developed olfactory lobes. This is the case in the human fœtus only at an early period.

the *tenia semicircularis*. 4. On the back part of the thalamus opticus lies a red fringe called the *choroid plexus*. It is formed by folds of the pia mater continued into the ventricles. 5. Immediately behind, and nearly parallel with the choroid plexus, lies the thin free border of the *fornix*. 6. In the middle or descending horn, the choroid plexus runs down to the base of the brain; and there is also a columnar eminence which

Fig. 121.



TRANSVERSE PERPENDICULAR SECTION THROUGH THE BRAIN.

stands out in relief on the back part of the cavity; to this the fanciful term "*hippocampus major*" has been applied.* 7. In the posterior horn there is a smaller elevation of the surface called the "*hippocampus minor*." These hippocampal elevations are nothing more than deep convolutions which project into the ventricles. By cutting through the white matter on their surface we soon come to the cortical substance of the brain.

Appearance on Perpendicular Section.—If a perpendicular section were made across the middle of the brain, the lateral ventricles would appear as represented in fig. 121. Together with the third or middle ventricle, their shape slightly resembles the letter T. Such a section shows well the radiating fibres of the corpus callosum, the fornix, and the velum interpositum beneath it: also the beginning of the transverse

* The lower end of the hippocampus major is somewhat expanded and indented on the surface, so as to resemble the paw of an animal. Hence the name given to it, "*pes hippocampi*."

fissure at the base of the brain, between the crus cerebri and the middle lobe.

Having learned the names of the parts in the lateral ventricles, now consider them a little more in detail, and also the transparent septum (*septum lucidum*) by which the two lateral ventricles are separated.

Corpus Striatum.—The corpus striatum is so called, because, when cut into, it presents alternate layers of white and grey matter.* It is a much larger mass of grey substance than it appears to be, for only a portion of it projects into the ventricle, cropping out on the surface. What we do see of it looks pear-shaped, with the large end forwards: it tapers gradually to a point as we trace it backwards on the outside of the optic thalamus (p. 510). The under part of the corpus striatum corresponds with the convolution at the base of the brain known as the island of Reil, and also with the locus perforatus anticus,—a spot, we remember, at the root of the fissure of Sylvius, so called on account of the number and size of the blood-vessels which pass in there to supply the mass of grey matter in question.

Tænia Semi-Circularis.—This is a narrow semi-transparent band of white fibres, which skirt the posterior border of the corpus striatum, fig. 123. In front it is connected with the anterior crus of the fornix; behind, it is lost in the middle horn of the lateral ventricle.

Septum Lucidum.—This is a thin and almost transparent partition, which descends vertically in the middle line from the under surface of the corpus callosum, and separates the anterior part of the lateral ventricles from each other. What are its attachments? Above, to the corpus callosum; below, to the reflected part of the corpus callosum and the fornix, p. 502. It is not of equal depth throughout. The broadest part is in front, and corresponds with the knee of the corpus callosum. It grows narrower behind, and altogether disappears where the corpus callosum and the fornix become con-

* The white lines in the corpus striatum are produced by the fibres of the crus cerebri, which traverse this mass of grey matter before they expand to form the hemisphere. The grey matter itself is sometimes called the anterior cerebral ganglion. It is found in all mammalia, in birds, and, to a certain extent, in reptiles. Its precise function is still unknown.

tinuous. Thin as it is, yet the septum consists of two layers which inclose a space called the "fifth ventricle" (fig. 122). Each layer is made up of white substance inside, and of grey outside; the little ventricle between them is closed in the adult, and lined by a delicate serous membrane, which secretes in it a minute quantity of fluid; but in foetal life it communicates with the third ventricle between the pillars of the fornix.*

Now cut transversely through the corpus callosum and the septum lucidum, and turn forwards the anterior half. In this way the ventricle of the septum will appear as in fig. 122. By turning back the posterior half of the corpus callosum we obtain a view of the fornix. This proceeding requires care, or the fornix will be reflected also, since these two arches of nervous matter are here so closely connected.

Fornix.—This, as implied by its name, is a layer of white matter arranged in the form of an arch from before back-

Fig. 122.




DIAGRAM OF THE FORNIX.

(The arrow is in the foramen of Monro.)

wards, beneath the corpus callosum. It is the great longitudinal commissure and lies over the velum (fig. 120, p. 502). Viewed from its upper surface it is triangular, with the base behind, as shown in diagram 122. The broad part is con-

* The development of the septum lucidum commences about the fifth month of foetal life, and proceeds from before backwards *pari passu* with the corpus callosum and fornix.

nected to the corpus callosum. From its anterior narrow part proceed two round white cords, called its anterior crura,—one on either side the mesial line. They do not extend so far forwards as the corpus callosum, but turn downwards to the base of the brain, where they may be traced, one into each of the corpora mammillaria (fig. 120). Immediately behind and below the anterior crura there is a passage through which the choroid plexuses of opposite sides are continuous with each other.

Foramen of Monro.—This aperture is called the “*foramen of Monro*.”* Strictly speaking, it is not a foramen, but only a communication existing between the two lateral and the third ventricles. Look at the posterior part of the fornix, and observe that from each angle of the base is continued a thin white band, which runs along, and is intimately connected with, the concave side of the hippocampus major: to this band the name of “*tænia hippocampi*” is applied; it is sometimes called the posterior crus of the fornix.†

Velum Interpositum and Choroid Plexus.—Reflect the fornix in order to expose the “*velum interpositum*,” which supports it. This is a portion of the pia mater which penetrates into the ventricles through the fissure beneath the posterior border of the corpus callosum, as shown p. 506.‡ The shape of this vascular veil is like that of the fornix itself, and its borders are rolled up so as to form the red fringe called the “*choroid plexus*.” These plexuses consist almost entirely of the tortuous ramifications of minute arteries.§ In front they communicate uninterruptedly with each other through the foramen of Monro; behind they descend with the middle

* Monro, “*Microscopic Enquiries into the Nerves and Brain*.” Edinburgh, 1780.

† The fornix and septum lucidum are absent in fish; they are merely rudimentary in reptiles and birds: but all mammalia have them in greater or less perfection, according to the degree of development of the cerebral hemispheres.

‡ The great fissure beneath the corpus callosum, where the pia mater enters the ventricles, is called the “*transverse fissure*” or the “*fissure of Bichat*.” It extends downwards on each side to the base of the brain, and terminates between the crus cerebri and the middle lobe. Taken as a whole, it has a horse-shoe shape, with the concavity forwards.

§ In preparations where the choroid plexus is well injected, we see that they are covered with beautiful vascular villi. The villi themselves are covered with epithelium.

horns of the lateral ventricles, and become continuous with the pia mater at the base of the brain, between the middle lobe of the cerebrum and the crus cerebri.

Venæ Galeni.—Down the centre of the veil run two large veins called "*venæ Galeni*," which return the blood from the ventricles into the straight sinus.

The velum interpositum, with the choroid plexus, must now be reflected to expose the following parts shown in diagram 123.—1. We have a full view of the thalamus opticus. 2. Between the thalami optici is the "*third ventricle of the brain*," a deep vertical fissure, precisely in the middle line. 3. Behind the fissure is the "*pineal gland*," a vascular body, about the size of a pea. From the gland you trace forwards two slender white cords, called its "*peduncles*,"—one along the inner side of each optic thalamus. 4. Immediately behind the pineal gland are four elevations, two on each side, called the "*corpora quadrigemina*" or "*nates and testes*." 5. These bodies are connected to the cerebellum by two flat bands, one on each side, termed the "*processus a cerebello ad testes*." 6. Between these bands extends a thin layer of grey substance, called the "*valve of Vieussens*," beneath which lies the fourth ventricle. Lastly, there are three commissures, anterior, middle, and posterior, connecting the opposite sides of the brain.

Thalamus Opticus.—This is the square elevation seen on the floor of the lateral ventricle immediately behind the corpus striatum. Though white on the surface, its interior consists of alternate layers of white and grey matter, like the corpus striatum.* The under surface of the thalamus forms the roof of the descending or middle horn of the lateral ventricle. Beneath the posterior part of the thalamus are two small eminences, termed the "*corpus geniculatum internum, and externum*."†

* The elevation called the optic thalamus is occasioned by the interposition of a quantity of grey matter among the fibres of the crus cerebri. Gall termed it the inferior ganglion of the cerebrum, in opposition to the corpus striatum, which he termed the superior ganglion (Anat. de Phys. du Système Nerv. : Paris, 1810). The epithet "optic" applied to the thalamus might lead us to suppose that it presides over vision ; but that it exercises very little influence over sight is rendered probable by comparative anatomy, by experimental physiology, and by pathology.

† The slight eminences, termed "*corpora geniculata*," are produced by small

Third Ventricle.—This is the narrow fissure which exists between the optic thalami, and reaches down to the base of the brain. Its boundaries are rather complicated :—The roof is formed by the velum interpositum ; the floor by certain parts at the base of the brain—namely, the locus perforatus medius, corpora mammillaria, tuber cinereum, infundibulum, commissure of the optic nerves, and lamina cinerea : all of which are best seen in a vertical section, as shown p. 502. In front it is bounded by the anterior crura of the fornix ; behind, it communicates with the fourth ventricle through the “*iter a tertio ad quartum ventriculum* ;” a long canal beneath the corpora quadrigemina.

Commissures.—If you gently separate the optic thalami in a fresh brain, you see that they are connected by a transverse layer of grey matter about half an inch in breadth.

Middle.—This is the middle commissure : it is sometimes called the *soft*, on account of its delicate consistence : in most brains it is generally torn before we have the opportunity of examining it.*

Posterior.—The optic thalami are also connected by a round white cord, called the posterior commissure. It is situated immediately in front of, and rather below the pineal gland.

Anterior.—The corpora striata are connected by a round white cord, called the anterior commissure : it lies immediately in front of the anterior crura of the fornix. If this commissure were traced throughout, we should find that it extends transversely through the corpus striatum of each side ; then arching backwards, its fibres are lost near the surface of the middle cerebral lobe.

Pineal Gland.—This very vascular heart-shaped body is situated immediately in front of the corpora quadrigemina.


accumulations of grey matter beneath the surface. A narrow band of white matter connects the external one with the testes, and a similar band connects the internal one with the testes. These bands are faintly marked in man, but more apparent in the lower animals.

* The soft commissure does not appear to be a very essential constituent part of the brain. It is not found before the ninth month of foetal life ; and in some instances, according to our observation, is never developed. The brothers Wenzel state that it is absent about once in seven subjects (*De pleniori Struct. Cerebri Hom. et Brut. Tubingen, 1812*).

It is wrapped up in the velum. It has two white peduncles or crura, which extend forwards, one on the inner side of each optic thalamus, and terminate by joining the crura of the fornix. In its interior is some gritty matter, consisting of phosphate of lime. Although the pineal gland is found in all mammalia, birds, and reptiles, in the same typical position, its functions are entirely unknown.

Tubercula Quadrigemina. — These are four eminences, situated, two on each side, behind the pineal gland. Though

Fig. 123.



white on the surface, they contain grey matter in the interior: this grey matter is accumulated here for the purpose of giving origin to the optic nerves. A more appropriate term for them would be the "optic lobes," instead of the vulgar name of "nates and testes," handed down from the old anatomists.*

* Eminences homologous to the corpora quadrigemina are found in all vertebrate animals: they are the meso-cephalic lobes; they invariably give origin

Processus a Cerebello ad Testes.—By gently drawing back the overlapping cerebellum we observe two broad white cords, which pass backwards, diverging from each other, from the corpora quadrigemina to the cerebellum (fig. 123). These are the “superior crura of the cerebellum.” They connect the cerebrum and the cerebellum. The space between them is occupied by a thin layer of grey matter which covers the fourth ventricle. This layer is called the valve of Vieussens. The fourth nerve arises from it by several filaments.

Iter a Tertio ad Quartum Ventriculum.—The third ventricle is connected with the fourth by means of a canal large enough to admit a probe, which runs backwards beneath the posterior commissure and the corpora quadrigemina (p. 502). This passage, together with the third and fourth ventricle, are persistent parts of the central canal, which, in early foetal life, extended all down the back of the spinal cord.

Fourth Ventricle.—This cavity is situated between the cerebellum and the posterior part of the medulla oblongata, and pons Varolii. It is only a dilated portion of the primordial canal, alluded to in the last paragraph. To obtain a perfect view of its boundaries you should make a vertical section as in diagram, p. 502. It appears triangular: its boundaries are as follow:—The front or base is formed by the medulla oblongata and pons Varolii; the upper wall by the valve of Vieussens; the posterior wall by the cerebellum itself, and the continuation of the arachnoid membrane on to the back of the spinal cord. The pia mater is prolonged for a short distance into the lower part of the cavity, and forms the “choroid plexus of the fourth ventricle.”*

Calamus Scriptorius.—On the anterior wall of the fourth ventricle,—that is, on the back of the medulla oblongata and

to the optic nerves, and their size bears a direct relation to the power of sight. They are relatively smaller in man than in any other animal. In birds there are only two eminences, and these are very large, especially in those far-seeing birds which fly high, as the eagle, falcon, vulture, &c., who require acute sight to discern their prey at a distance.

* Tiedemann proposes to call the fourth ventricle the first; because, in the foetus, it is formed sooner than any of the others; because it exists in all vertebrated animals, whereas the lateral ventricles are absent in all osseous fishes; and because the ventricle of the septum lucidum is absent in all fishes, in reptiles, and in birds.

pons,—are several objects to be noticed (p. 517). 1. A median furrow, the remains of the primitive axis canal. 2. From the lower part of the furrow two white cords (the *restiform columns* or *inferior crura cerebelli*) pass off from the medulla oblongata, diverging like the branches of the letter V, and enter the lateral lobes of the cerebellum. The divergence of these cords with the median furrow was called by the old anatomists the "*calamus scriptorius*." 3. The floor of the fourth ventricle is covered by grey matter, which is nothing more than the grey substance of the medulla exposed by the divergence of its posterior or restiform columns. On its surface we remark a number of transverse white lines, some of which form part of the origin of the auditory nerves.

[The average weight of the cerebrum is about two and a half pounds. It measures antero-posteriorly about six inches—behind, laterally, which is the largest measurement, five inches, and vertically, it will be found to measure from four to five inches.]

Cerebellum.—This portion of the brain is situated in the occipital fossa, beneath the posterior lobes of the cerebrum, from which it is separated by the tentorium. Its form is elliptical, with the broad diameter transverse. When the arachnoid membrane and the pia mater are removed, you observe that its surface is not arranged in convolutions like those of the cerebrum. It is laid out upon a plan beautifully adapted to produce a much greater superficial extent of grey matter. It consists, in fact, of a multitude of thin plates, disposed in a series of concentric curves, with the concavity forwards. By a little careful dissection it is easy to separate some of the plates from each other, and to see that the intervening fissures increase in depth from the centre towards the circumference of the cerebellum. The fissure at the circumference itself is the deepest of all, and nearly horizontal, so that it seems to divide the cerebellum into an upper and a lower segment.

Upper Surface.—The upper surface of the cerebellum slopes on each side like the roof of a house. The ridge along the middle line is called the "*superior vermiform eminence*." Comparative anatomy proves that this is really the fundamental part of the cerebellum. The sides, called the "*hemi-*

spheres," are merely offsets or wings, superadded for special purposes, and increase in size as we ascend in the vertebrate series, till in man they form by far the largest part of the organ. They are separated posteriorly by a perpendicular fissure, which receives the "falx cerebelli."

Under Surface.—On the under surface of the cerebellum its division into two hemispheres is clearly perceptible. The deep furrow between them is called the "*valley*." The front part of it is occupied by the medulla oblongata. Therefore, to examine the surface of the valley, you must raise the medulla, and separate the hemispheres from each other. Along the middle line of the valley, we observe the "*inferior vermiform eminence*," which is, in point of fact, the under surface of the fundamental part of the cerebellum. Traced forwards, this eminence terminates in a tongue-like body, called the "*uvula*;" * traced backwards, it terminates in a small conical projection, called the "*pyramid*."

Each hemisphere presents on its under surface certain secondary lobes, to which fanciful names have been applied. That portion which immediately overlays the side of the medulla oblongata is called the "tonsil" (*amygdala*): at the anterior part of each hemisphere, near the middle line, is a little lobe called the "*flocculus*," or pneumogastric lobe.

Appearance of the Interior.—To examine the internal structure of the cerebellum, a longitudinal section should be made through the thickest part of one of the hemispheres. There is then seen in the centre a large nucleus of white matter, from which branches radiate into the grey substance in all directions. Each of these branches corresponds to one of the plates of the cerebellum, and from it other smaller branches proceed and again subdivide. This racemose appearance of the white matter in the substance of the grey has been likened to the branches of a tree deprived of its leaves, and is generally known as the "*arbor vitæ*," it is a beautiful

* From either side of the uvula may be traced a thin valve-like fold of white substance, which proceeds in a semicircular direction to the little pneumogastric lobes. These folds are called the semicircular valves, or "*valves of Tarini*," by whom they were first pointed out. (Advers. Anat. prima: Paris, 1750.)

contrivance for bringing an extensive surface of the two kinds of nervous matter into connection with each other.

Corpus Rhomboideum.—In the centre of the white nucleus of each hemisphere is an oval space, circumscribed by a zig-zag line of grey matter. To this the name “corpus dentatum or rhomboideum” has been given. It is displayed both by a vertical and by a horizontal section.

Connections of Cerebellum.—The cerebellum is connected with the cerebro-spinal axis as follows:—With the medulla oblongata, by means of the restiform or sensory tracts of nervous matter: these are called its inferior crura;—with the cerebrum, by means of the processus a cerebello ad testes: these are called its superior crura. The transverse fibres of the pons constitute the middle crura.

Function.—Respecting the function of the cerebellum, the arguments furnished by comparative anatomy render probable that it is a co-ordinator of muscular movements,—*e. g.* in the action of walking, flying, swimming, &c.

Pons Varolii or Tuber Annulare.—This convex eminence (p. 495), is situated at the base of the brain, immediately above the medulla oblongata. It corresponds with the basilar groove of the occipital bone. In its antero-posterior diameter it measures rather more than one inch. Down the middle runs a furrow which lodges the basilar artery.

The Great Commissure of the Cerebellum.—If the pia mater be carefully removed, we see that its fibres proceed transversely from one hemisphere of the cerebellum to the other: hence it is called the commissure of the cerebellar lobes. Throughout the mammalia its size bears a direct ratio to the degree of development of these lobes; therefore it is larger in man than in any other animal.* Understand that only the superficial fibres are transverse; if these be turned off, we see that the anterior fibres of the medulla run under them at right angles into the crura cerebri, like a river under a bridge.

Besides the transverse and longitudinal fibres just described,

* Birds, reptiles, and fishes, have no pons, because there are no lateral lobes to the cerebellum.

the pons contains grey matter which probably gives origin to fresh nerve-fibres; so that it appears reasonable to consider the pons a source, as well as a mere conductor of power.

[The average weight of the cerebellum is six ounces. Transverse diameter from three to four inches; antero-posterior, two to two and a half inches; so also with the vertical measurement.]

Medulla Oblongata.—This term is applied to that part of the cerebro-spinal axis which follows immediately below the pons Varolii, and is continuous with the spinal cord. It is one of the most important divisions of the nervous system. It is not only the highroad for nerves to and from the brain, but it contains the nerve centres which preside over respiration, deglutition, and speech. It is sometimes called the “vital knot.” Injury to the medulla oblongata is immediately fatal. It is about an inch and a quarter long. It lies above the basilar groove of the occipital bone; and descends obliquely backwards to the foramen magnum, where the spinal cord begins. Its posterior surface is received into the fossa between the hemispheres of the cerebellum. Its form is pyramidal with the thickest part above; from thence it gradually tapers into the spinal cord.

[The length of the medulla is from one inch to an inch and a quarter; breadth, from eight to nine lines at the base, its thickness six lines.]

Anterior Aspect.—The anterior surface of the medulla oblongata presents a deep median fissure continuous with that of the cord (fig. 124). Below the pons Varolii this fissure terminates in a cul-de-sac, named the “*foramen cæcum* :” it is occupied by a fold of pia mater. On each side of the fissure there stand out in relief three longitudinal columns. The middle ones are called the “*anterior pyramids*.” External to these are the “*olives*.” Still more external, and towards the posterior part of the medulla are the “*restiform tracts*.”

The *anterior pyramids*, so called from their shape, are narrow below, but gradually increase in breadth as they ascend towards the pons. The fibres of which they are composed are continuous with the anterior columns of the spinal cord, and consist, therefore, of motor fibres. They may be traced through the pons into the lower part of the crura cere-

bri. By gently separating the pyramids about one inch below the pons, you observe that their fibres on each side decussate (fig. 124). This is the explanation of "cross paralysis;" *i. e.*, when one side of the brain is injured the loss of power is manifested on the opposite side of the body. The decussation takes place only between the inner fibres of the pyramid; the outer fibres run straight on without decussating. We shall presently find that these decussating fibres are pro-

Fig. 124.




DIAGRAM OF THE FRONT SURFACE OF THE MEDULLA OBLONGATA.

longations of the deep fibres of the lateral columns of the cord which here come forward to the surface, pushing aside the proper anterior columns.

The *olivary bodies* or *olives* are two oval eminences situated on the outer side of the pyramids. They consist, externally, of white substance; when cut into, their interior presents an undulating line of grey matter, called, from its zigzag shape, "*corpus dentatum*." This grey line forms a circuit, interrupted only on the inner side, so that it nearly isolates the white matter in its centre.

The *restiform tracts* are situated to the outer side of and behind the olives; to see them well you should look at the

back of the medulla. There you observe that they diverge from each other, and pass into the cerebellum, constituting its inferior crura (fig. 125). In consequence of this divergence the grey matter in the interior of the medulla oblongata is exposed; hence it is that the floor of the fourth ventricle is grey. The restiform bodies contain in their interior a considerable portion of grey matter, which is continuous with that in the posterior part of the spinal cord. The fibres of which they are composed are conductors of sensation.*

Fig. 125.




DIAGRAM OF THE FOURTH VENTRICLE AND RESTIFORM TRACTS.

- | | |
|----------------------------|---|
| 1. Thalamus opticus. | 4. Processus a cerebello ad testes. |
| 2. Nates and testes. | 5. Restiform tracts diverging. |
| 3. Origin of fourth nerve. | 6. Origin of seventh or auditory nerve. |

When a longitudinal section is carefully made through the middle of the pons and the medulla oblongata, a number of white fibres are seen passing in a horizontal direction, constituting a kind of septum between the two halves. Some of these septal fibres issuing from the anterior fissure wind round

* Two slender columns are marked off from the back part of the restiform bodies, one on either side the median fissure (fig. 125). These are the *posterior pyramids* of some anatomists. They proceed with the restiform tracts to the cerebellum.

the sides of the medulla, and constitute what are termed the *arciform fibres* of the medulla (p. 516); others, again, issuing from the posterior fissure, and winding round in a similar manner, form some of the transverse fibres seen on the floor of the fourth ventricle.

DISSECTION OF THE SPINAL CORD.

THE proper mode of taking out the spinal cord is to saw through the arches of the vertebræ. When these are removed, the spinal cord, covered by its membranes, is exposed. The first thing we notice is, that the cord does not occupy the whole area of the spinal canal. The dura mater does not adhere to the vertebræ, and is not their internal periosteum as it is in the skull. There intervenes between the bones and this membrane a space filled by a soft, reddish-looking fat, and watery cellular tissue, and by the ramifications of a plexus of veins. A few words about these veins before we proceed.

Spinal System of Veins.—The spine is remarkable for the number of large and tortuous veins which ramify about it, both inside and outside the vertebral canal.* One cannot form an adequate idea of them unless when properly injected. 1. There is a plexus of tortuous veins, both outside and inside the arches of the vertebræ. 2. Two large veins extend all down the spinal canal behind the bodies of the vertebræ: they may with strict analogy be termed the “sinuses” of the spinal canal. They communicate by cross branches like a ladder, and receive the large veins which emerge from the cancellous texture of the bones. 3. There are the proper veins of the spinal cord which lie within the dura mater. All discharge themselves through the intervertebral foramina in the several regions of the spine, as follows:—in the cervical, into the vertebral veins; in the dorsal, into the inter-

* A very accurate description and representation of these veins has been given by Breschet, *Essai sur les Veines du Rachis*, 4to.; *Traité Anatomique sur le Système Veineux*, fol. avec planches.

costal veins; in the lumbar, into the lumbar veins. Note that this complicate system of veins is wholly unprovided with valves; hence it is they are so liable to become congested in diseases of the spine.

Peculiarities of the Membranes of the Cord.—The membranes of the spinal cord, though continuous with those of the brain, differ from them in certain respects, so as to require a separate notice.

Dura Mater.—The “dura mater” of the cord is a tough fibrous membrane like that of the brain, but does not adhere to the bones, because such adhesion would impede the free movement of the vertebræ upon each other. It forms a complete canal, which loosely surrounds the spinal cord, and sends off prolongations over each of the spinal nerves. These prolongations accompany the nerves only as far as the intervertebral foramina, and are then blended with the periosteum.

Cut through the nerves which proceed from the spinal cord on each side, and remove it with the dura mater entire. Then slit up the dura mater along the middle line, in order to examine the arachnoid membrane.

Arachnoid Membrane.—The “arachnoid membrane” of the cord is a continuation from that of the brain. It is not in immediate contact with the pia mater underneath, but is separated from it by a transparent watery fluid contained in the meshes of the subarachnoid tissue: therefore the cord may truly be said to float in a fluid (p. 522).

Cerebro-spinal Fluid.—This cerebro-spinal fluid cannot be demonstrated unless the cord be examined very soon after death, and before the removal of the brain.* The nerves

* The existence and situation of the cerebro-spinal fluid was first discovered by Haller (Element. Phys. Vol. iv. p. 87), and subsequently more minutely investigated by Magendie (Récherches Phys. et Cliniques sur le Liquide Cephalo-rachidien, in 4to. avec atlas: Paris, 1842). This physiologist has shown that if, during life, the arches of the vertebræ are removed in a horse, dog, or other animal, and the dura mater of the cord punctured, there issue jets of a fluid which had previously made the sheath tense. The fluid communicates, through the fourth ventricle, with that in the general ventricular cavity. The collective amount of the fluid varies from one to two ounces or more. It can be made to flow from the brain into the cord, or *vice versâ*. This is proved by experiments on animals, and by that pathological condition of the spine in children termed “spina bifida.” In the latter instance, coughing and crying

proceeding from the cord are loosely surrounded by a sheath of the arachnoid; but this only accompanies them as far as the dura mater, and is then reflected upon that membrane.

Pia Mater.—The pia mater of the cord is the membrane which immediately invests it. It is quite different from that of the brain, since it does not form a bed in which the arteries break up, but serves rather to support and strengthen the cord: consequently it is much less vascular, more *fibrous* in its structure, and more adherent to the substance of the cord. The fibres of which it is composed are rendered very evident by immersion for a time in water. It is prolonged upon the spinal nerves, and forms their investing membrane or “neu-rilemma.”

The pia mater sends off from each side of the cord, along its whole length, a series of ligaments to steady it. They are triangular, their bases being attached to the cord, and their points to the inside of the dura mater (fig. 126).




DIAGRAM OF
THE LIGA-
MENTA DEN-
TATA.

1. Dura mater.
2, 2, 2. Ligamen-
ta dentata.

Ligamenta Dentata.—Thus they look like a series of teeth down the side of the cord: hence the name “*ligamenta dentata*.” There are from eighteen to twenty-two of them on each side, and they lie between the anterior and posterior roots of the spinal nerves.

Spinal Cord.—**Cauda Equina.**—The spinal cord is that part of the cerebro-spinal axis contained in the vertebral canal. It is the continuation of the medulla oblongata, and extends from the foramen magnum down to the upper border of the second lumbar vertebra, where it terminates in a pointed manner, after having given off the great bundle of nerves termed the “*cauda equina*” for the supply of the lower limbs.* The general form of the cord is cylin-

make the tumor swell; showing that fluid is forced into it from the ventricles. Again, if pressure be made on the tumor with one hand, and the fontanelles of the child examined with the other, in proportion as the spinal swelling decreases, so is the brain felt to swell up, accompanied by symptoms resulting from pressure on the nervous axis generally. See some remarks very much to the point, by Dr. Burrows, On Diseases of the Cerebral Circulation, p. 50, 1846.

* Although the nerve substance of the cord itself terminates at the second lumbar vertebra, yet the pia mater is continued as a slender filament, called

drical, slightly flattened in front and behind. It is not of uniform dimensions throughout, but presents a slight enlargement in the lower part of the cervical region; another in the lower part of the dorsal, just where the great nerves of the upper and lower limbs are given off.

Fissures.—The cord is divided into two symmetrical halves by a fissure in front and behind (fig. 127). The *anterior fissure* is the most distinct, and penetrates about one-third of the substance of the cord; it contains a fold of pia mater full of blood-vessels for the supply of the interior. At the bottom of this fissure is a transverse layer of white substance, named the *anterior commissure*, connecting the two anterior halves of the cord. The *posterior fissure* is so much less apparent than the anterior that some anatomists altogether deny its existence; but it can be demonstrated by a careful preparation, and, indeed, penetrates to a greater depth than the anterior, so that it reaches quite down to the grey matter in the centre of the cord.

Besides the anterior and posterior fissures, there run down each half of the cord two very superficial grooves, from which the anterior and posterior roots of the spinal nerves respectively emerge. These constitute the *anterior* and *posterior lateral grooves*. The posterior leads down to the posterior horn of the grey matter in the interior of the cord; the anterior is less distinct than the other, and does not reach down to the anterior horn of grey matter.

Columns.—By these lateral grooves each half of the cord is divided into three longitudinal columns—an anterior, a posterior, and a lateral. The anterior are motor columns, the posterior sensitive, the lateral probably contain both motor and sensitive filaments.

Interior.—A transverse section through the cord (fig. 127) shows that its interior contains grey matter, arranged in the form of two crescents placed one in each half of it, and con-

“flum terminale” down to the base of the coccyx. The explanation of this is, that, at an early period of foetal life, the length of the cord corresponds with that of the vertebral canal; but after the third month, the lumbar and sacral vertebræ grow away, so to speak, from the cord, in accordance with the more active development of the lower limbs. See Tiedemann, *Anatomie und Bildungsgeschichte des Gehirns im Fœtus des Menschen*, &c.; Nürnberg, 1816.

nected across the centre by a portion called the "*grey commissure*."* The posterior horns of the crescents are long and narrow, and extend to the posterior lateral fissure, where they are connected with the posterior roots of the spinal nerves.




DIAGRAM OF A TRANSVERSE SECTION THROUGH THE SPINAL CORD AND ITS MEMBRANES.

The anterior horns are short and thick, and come forwards towards the line of attachment of the anterior roots of the nerves, but do not reach the surface.

Spinal Nerves.—Thirty-one pair of nerves arise from the the spinal cord,—namely, 8 in the cervical region, 12 in the dorsal, 5 in the lumbar, 5 in the sacral, and 1 in the coccygeal.

Two Roots, Sensitive and Motor.—Each nerve comes off by two distinct series of roots,—one from the front, the other from the back of the cord. We are indebted to Sir Charles Bell for the discovery of the fact that the anterior roots consist exclusively of motor filaments, the posterior exclusively of sensitive. All converge and unite in the corresponding intervertebral foramen to form a single nerve, composed of both motor and sensitive filaments.

The posterior or sensitive roots proceed from the posterior lateral groove of the cord, and are thicker and more numerous than the anterior.† But the most remarkable thing about them is, that, previous to their union with the anterior roots, they are collected together and pass through a ganglion. This ganglion is of an oval form, and lies in the intervertebral

* The grey matter in the interior of the cord of man and animals presents somewhat different appearances in its different parts. These have been accurately described and figured by Rolando, *Ricerche Anatomiche sulla Struttura del Midollo Spinale*, con Figure, art. Tratto dal Dizionario Periodico di Medicina, Torino, 1824, 8vo. p. 55.

† The researches of Blandin, *Anat. descript.* t. ii., p. 648, 1838, have led him

foramen, just where the roots of the nerves pass through the dura mater.*

The compound nerve formed by the junction of the two roots divides, outside the intervertebral foramen, into an anterior and a posterior branch, for the diagram of which see p. 110.

Variation in Length of the Roots.—The direction and length of the roots of the nerves vary in the different regions of the spine, because the respective parts of the cord from which they arise are not opposite the foramina through which the nerves leave the spinal canal. In the upper part of the cervical region, the origins of the nerves and their point of exit are nearly on the same level: therefore the roots proceed transversely, and are very short. But as we descend from the neck, the obliquity and length of the roots gradually increase, so that the roots of the lower dorsal nerves are at least two vertebra higher than the foramina through which they emerge. Again, since the cord itself terminates at the second lumbar vertebra, the lumbar and sacral nerves must of necessity pass down from it almost perpendicularly through the lower part of the spinal canal.

Cauda Equina.—To this bundle of nerves the old anatomists have given the name of "*cauda equina*," from its resemblance to a horse's tail.

In brief, then, it appears that the spinal cord consists of two precisely symmetrical parts, separated in front and behind by a deep median fissure; that the two parts are connected at the bottom of the anterior fissure by an anterior or white commissure—at the bottom of the posterior fissure by the posterior or grey commissure; that each part of the cord is divided into three tracts or columns of longitudinal nerve-fibres—an anterior, a posterior, and a lateral—the boundaries between them being the respective lines of origin of the roots

to establish the following relation between the respective size of the anterior and posterior roots of the nerves in the several regions of the spine:—

The posterior roots are to the anterior in the cervical region	::	2	:	1
“ “ “ dorsal “	::	1	:	1
“ “ “ lumbar and sacral	::	1½	:	1

This relation quite accords with the greater delicacy of the sense of touch in the upper extremity.

* The ganglia of the two last sacral nerves lie within the dura mater.

of the spinal nerves ; that the interior of the cord contains grey matter disposed in the form of two crescents placed with their convexities towards each other, and connected by a transverse bar of grey matter, which is the posterior commissure.

Blood-Vessels of the Cord.—The cord is supplied with blood by—1, the *anterior spinal* artery, which commences at the medulla oblongata by a branch from the vertebral of each side, and then runs down the cord, receiving, through the intervertebral foramina, numerous branches in its course from the vertebral, intercostal, and lumbar arteries : 2, the *posterior spinal* arteries, which proceed also from the vertebral, intercostal, and lumbar arteries, and ramify very irregularly on the back of the cord.

It is curious that, on the posterior part of the bodies of the vertebræ the spinal arteries of opposite sides form by their mutual communications a perfect arterial “ladder” down the entire length of the spinal column.* We noticed a similar arrangement of the venous “sinuses” of the spine, p. 518.

Functions of the Columns of the Cord.—The *anterior* columns consist exclusively of motor fibres which originate from the grey matter of the brain or that of the spinal cord, and carry the commands of the will and the power of reflex movement to the muscles.

The *posterior* columns consist exclusively of sensitive fibres, which carry sensations, not, as was formerly believed, direct to the brain, but to the grey matter of the cord, through which *alone*, according to the recent experiments of Brown Séquard,† they are transmitted to the brain. The same experimentalist has also proved another unexpected fact ;—that sensations do not run up on the same side but on the opposite. They cross in the cord ; for instance, if the posterior column on the *right* side were injured, the *left* leg and not the right would be deprived of feeling.

Minute Structure of the Medulla Oblongata and Pons Varolii.—These are among the most complicate parts of the


* A preparation of the arteries in the body of a child, in the Museum of St. Bartholomew's Hospital, shows this point remarkably well.

† See an able article on Brown Séquard's experiments in the British and Foreign Medico-Chirurgical Review by Mr. Thomas Smith.

central nervous system. They contain white and grey matter,* intermixed. The white matter consists in part of a continuation of the longitudinal fibres of the cord, in part of a new system of horizontal fibres. We will endeavor to trace the longitudinal fibres first; then the horizontal.

Anterior Columns of the Cord.—The anterior columns (8), (fig. 128), having reached the lower part of the medulla

Fig. 128.



1. Corpus striatum.
2. Thalamus opticus.
3. Crus cerebri.
4. Locus niger.
5. Pons Varolii.
6. Pyramid.
7. Olive.
8. Anterior columns.
9. Lateral columns.
10. Posterior columns.
11. Corpora quadrigemina.
12. Fillet of Reil.
13. Superior crus of cerebellum.
14. Cerebellum.

DIAGRAM OF THE COURSE OF THE FIBRES OF THE CORD.

oblongata, are not continued straight up through it, but diverge from each other, so as to allow a part of the lateral columns (9), to come forward, and after decussation to form the pyramids (6). In their further progress the fibres of the anterior columns are disposed of thus: a small number of

* The grey matter in the medulla oblongata is collected in three situations:—1. In the olives; 2. In the restiform tracts; 3. On the floor of the fourth ventricle.

them run up and contribute to form the outer portion of their own pyramid; all the rest, after embracing the olive, pass up through the deep strata of the pons, and then divide into two bundles; one of these, called the fillet of Reil (12), mounts over the superior crus of the cerebellum to the corpora quadrigemina, beneath which it meets with the corresponding fillet of the opposite side; the other proceeds with the crus cerebri to the cerebrum.

Lateral Columns of the Cord.—The lateral columns (9), on reaching the medulla oblongata, are disposed of in three ways, as follows:—Some of its fibres come forward between the diverging anterior columns, decussate in the middle line, and then form the pyramid of the opposite side; others ascend with the restiform tract into the cerebellum; a third set ascend along the floor of the fourth ventricle* (concealed by its superficial grey matter), and then along the upper part of the crus cerebri into the cerebrum.

Posterior Columns of the Spinal Cord.—The posterior columns (10), ascend under the name of the restiform tracts at the back of the medulla oblongata, diverge from each other, and are, for the most part, continued into the cerebellum, forming its inferior crura; but some of their fibres run on along the floor of the fourth ventricle (external to the fibres from the lateral columns), then along the upper part of the crura cerebri into the cerebrum.

Horizontal Fibres.—The horizontal fibres in the medulla oblongata and the pons were first accurately described and delineated by Stilling.† Some of them form a raphé, and divide the medulla oblongata and pons into symmetrical halves; others, arising apparently from the raphé, pass outwards in an arched manner through the lateral halves of the medulla; so that, when seen in a transverse section by transmitted light, they describe a series of curves, with the convexity forwards, throughout the entire thickness of the medulla. Some of these transverse fibres appear on the surface over the pyramid and the olive; these have received the name of "*arciform*

* These fibres constitute what are sometimes called the "round cords" of the fourth ventricle.

Ueber die Medulla Oblongata. Erlangen, 1843.

fibres" (p. 516). It is difficult to determine the object of this system of transverse fibres, or what parts they connect. Stilling and Kölliker,* who have deeply studied the subject, are both of opinion that they originate in the restiform tracts, and thence arch forwards,—some on the surface, others through the substance of the medulla, and that they eventually join the fibres of the raphé.

Internal Structure of the Pons Varolii.—The pons consist of transverse and longitudinal white fibres, with a considerable quantity of grey matter in its interior. The superficial layer of fibre is obviously transverse, and connects the two wings of the cerebellum. After removing this first layer, we come upon the longitudinal fibres of the pyramids in their course to the *crura cerebri* (p. 525): these longitudinal fibres however, are intersected by the deep transverse fibres of the pons, which, like the superficial, are continued into the cerebellum. The third and deepest layer of the pons consists entirely of longitudinal fibres, derived partly from the lateral, partly from the restiform tracts of the medulla.

Crura Cerebri.—These are composed of longitudinal fibres, derived from the pyramids, from part of the lateral and restiform columns of the cord, and from the grey matter in the pons Varolii. If one of the *crura* be divided longitudinally, we find in the middle of it a layer of dark-colored nervous matter, called "*locus niger*;" it separates the crus into an upper and a lower stratum of fibres. The lower stratum is tough and coarse, and consists of the continuation of the fibres proceeding from the pyramid and the pons. The upper stratum is much softer and finer in texture, and has received the name of the "*tegmentum*:" it is composed of the fibres proceeding from the lateral and the restiform columns; also from the superior commissure of the cerebellum.

Tracing the fibres of the crus cerebri into the cerebral hemisphere, we find that they diverge from each other, that its lower fibres ascend chiefly through the corpora striata, its upper fibres through the thalami optici. In passing through these great ganglia, the crus receives a very large addition to its fibres: these then branch out widely towards all parts of

* Mikroskopische Anatomie, p. 454.

the hemisphere, in order to reach the cortical substance on the surface.

From what has been said it appears that the crura or roots of the cerebellum and the cerebrum contain part of the motor and part of the sensitive tracts of the spinal cord.

Origin of the Cerebral Nerves.—The cerebral nerves are given off in pairs, named the first, second, third, &c., according to the order in which they appear, beginning from the front. There are nine pairs. Some are nerves of special

Fig. 129.




DIAGRAM OF THE ORIGINS OF THE OLFACTORY AND OPTIC NERVES.

sense,—as the olfactory, the optic, the auditory ; others are nerves of common sensation,—as the large root of the fifth, the glossopharyngeal, and the pneumogastric ; others, again, are nerves of motion,—as the third, the fourth, the small root of the fifth, the sixth, and the facial division of the seventh, the spinal accessory, and the ninth.

First Pair, or Olfactory Nerve.—This nerve arises by three roots,—an outer and an inner, composed of white matter, and a central composed of grey (fig. 129).

The outer root proceeds from the bottom of the fissure of Sylvius, and describes a curve with the concavity outwards.

The inner root arises from the posterior extremity of the internal convolution of the anterior cerebral lobe.

The middle or grey root arises from the posterior extremity of the furrow in which the olfactory nerve is lodged; to see it, therefore, you should turn the nerve backwards.

The olfactory nerve is triangular, that it may fit into a furrow between the convolutions. It proceeds straightforwards under the anterior lobe, and terminates in the olfactory ganglion, which lies on the cribriform plate of the ethmoid bone.

The olfactory ganglion is of an olive-like shape, of a reddish grey color, and very soft consistence. It gives off from its under surface the true olfactory nerves.* For the description of these, see p. 181.

Second Pair, or Optic.—These nerves arise chiefly from the corpora quadrigemina; also from the geniculate bodies and the thalami optici (p. 528). They wind round the crura cerebri to the base of the brain, and, uniting in the middle line, constitute the optic commissure. This commissure rests on the sphenoid bone, in front of the cella turcica. From the commissure each nerve passes through the foramen opticum into the orbit, and terminates in the retina.

At the commissure some of the nerve fibres cross from one side to the other. This decussation affects only the middle fibres of the nerve; the course of the other fibres is shown in the diagram (p. 501). The purpose of this partial crossing is not thoroughly understood. It was ingeniously supposed by Dr. Wollaston† to account for single vision, since the right halves and the left halves of the eyes would derive their nerve-fibres from the same optic nerve.

* Strictly speaking, the olfactory nerve and its ganglion are integral parts (the prosencephalic lobe) of the brain. What in human anatomy is called the origin of the nerve is, in point of fact, the crus of the olfactory lobe, and is in every way homologous to the crus cerebri or cerebelli. In proof of this, look at the enormous size and connections of the crus in animals which have very acute sense of smell. Throughout the vertebrate kingdom there is a strict ratio between the sense of smell and the development of the olfactory lobes. Again, in many animals, these lobes are actually larger than the cerebral, and contain in their interior a cavity which communicates with the lateral ventricles. According to Tiedmann, this cavity exists even in the human foetus at an early period.

† Philosophical Transactions of the Royal Society, 1824.

Third Pair, or Motores Oculorum.—The apparent origin of the third nerve is from the inner side of the crus cerebri, immediately in front of the pons; but its roots penetrate into the crus as deep as the locus niger (p. 525). It passes through the sphenoidal fissure, and supplies all the muscles of the eye, except the superior oblique and the rectus externus.

Fourth Pair, or Pathetici.—The fourth nerve arises from the valve of Vieussens (p. 510). It runs transversely outwards, winds round the crus cerebri, enters the orbit through the sphenoidal fissure, and supplies the superior oblique muscle of the eye.

Fifth Pair, or Trigeminal Nerve.—The fifth nerve arises by two roots of unequal size. The apparent origin of both is from the side of the pons (p. 516); but their real origin is much deeper. The smaller and more anterior of the two consisting of motor fibres only, may be traced into the pyramidal tract in the pons; the posterior and larger root, consisting of purely sensitive fibres, may be traced into the restiform tract of the medulla. The nerve proceeds forwards over the apex of the petrous portion of the temporal bone: here is developed upon the sensitive root the great *Gasserian* ganglion. This root then divides into three branches; the ophthalmic, which passes through the orbital fissure; the superior maxillary, which passes through the foramen rotundum; the inferior maxillary, which passes through the foramen ovale. They all confer common sensibility upon the parts they supply, which comprise nearly the entire head. The small motor root accompanies the inferior maxillary division of the sensitive root, and is distributed to the muscles of mastication.

Sixth Pair, or Abducentes.—The sixth nerve arises from the medulla oblongata, close to the pons (p. 516), leaves the skull through the sphenoidal fissure, and supplies the rectus externus of the eye.

Seventh Pair.—The seventh consists of two distinct nerves—the *portio dura*, or muscular nerve of the face, and the *portio mollis*, or proper auditory nerve. The apparent origin of both is from the lower parts of the pons Varolii (p. 516). The real origin of the *portio dura* is in the lateral columns of the medulla. The real origin of the auditory nerve is from the floor of the fourth ventricle by several filaments. The

seventh pair emerges from the skull through the meatus auditorius internus. For the further description of the portio dura, see p. 64. The auditory nerve is distributed to the internal ear.

Eighth Pair.—This comprises three nerves—the glossopharyngeal, the pneumogastric, and the nervus accessorius (p. 516). The first two arise by several filaments from the restiform tract of the medulla close to the olive. The nervus accessorius arises by a series of roots from the lateral column of the spinal cord, as low down as the fifth cervical nerve. It then ascends into the skull through the foramen magnum, and leaves it, with the other two nerves, through the foramen jugulare. The glossopharyngeal is distributed to the mucous membrane of the pharynx and the back of the tongue (p. 78). The pneumogastric is distributed to the pharynx, the larynx, the heart and lungs, the œsophagus and the stomach. The nervus accessorius supplies the sterno-mastoid and the trapezius. For the further description of these nerves, see pp. 80 and 9.

Ninth Pair, or Hypoglossal.—This nerve arises by several roots from the medulla oblongata, along the groove between the pyramid and the olive. It leaves the skull through the anterior condyloid foramen, and is distributed to the muscles of the tongue and the depressors of the os hyoides and larynx.

THE DISSECTION OF THE EYE.

SINCE the human eye cannot be obtained sufficiently fresh for anatomical purposes, we must have recourse to the eyes of animals—say of the sheep or the pig. The first thing to be done is to remove the conjunctival coat together with the loose connective tissue which unites it to the sclerotica.

Membrana Conjunctiva.—The conjunctiva is the mucous membrane which lines the eyelids. From the back part of the tarsal cartilages it is reflected over the sclerotic coat of the eye by loose folds, so as not to impede the motions of the globe. The palpebral portion of it is very vascular, and pro-

vided with fine papillæ.* The ocular portion has no papillæ, and is nearly colorless, except when inflamed; it then becomes intensely vascular and red, like a piece of scarlet cloth. An abundant supply of nerves has been bestowed upon the membrane for the purpose of giving it that exquisite degree of sensibility necessary to guard the eye (p. 54).

Continued Over the Cornea.—A thin layer of conjunctiva covers the cornea or transparent part of the eye. True, it cannot be separated by dissection in recent eyes; but the corneal conjunctiva possesses the same acute sensibility as the rest of it; changes produced by inflammation of the conjunctiva are often continued over the cornea: we see red vessels injected on its surface, and its texture becomes thickened.†

The human eye is very nearly spherical. It would be quite so, but that the transparent part in front—the cornea—forms a segment of a smaller sphere than the rest. Consequently, the antero-posterior diameter of the ball exceeds the transverse by about one line. The convexity of the cornea, however, varies in different persons, and at different periods of life; this is one cause of the several degrees of near-sight and far-sight.

[*Myopia* and *hypermetropia* are due to an abnormal form of the eyeball. *Presbyopia* depends upon increased hardness of the lens. For example, in *myopia*, the antero-posterior diameter is greater, while the same diameter of the *hypermetropic* eye is shorter than that of the normal eye.]

Coats and Humors of the Eye.—The globe is composed of three coats arranged one within the other. The external coat, called the "*sclerotic*," is fibrous, thick, and strong, so as to form a case to protect the delicate structures within it. The second coat, called the "*choroid*," is composed almost entirely of blood-vessels, and very dark in color. The third coat, called the "*retina*," consists of the expansion of the op-

* These papillæ were first pointed out by Eble, Ueber den Bau und die Krankheiten des Bindehaut des Auges.

† The facts of comparative anatomy confirm this view. In the serpent tribe, which annually shed their skin, the front of the cornea comes off with the rest of the external surface of the body. In the eel the surface of the cornea is often drawn off in the process of skinning. In some species of rodents which burrow under the ground like the mole, the eye is covered with hair like other parts.

tic nerve for the reception of the impression of light. The great bulk of the interior is filled with a transparent humor called the "*vitreous*." Imbedded in the front of this, and just behind the pupil, is the crystalline lens for the purpose of concentrating the rays of light. In front of the lens is placed a movable curtain, called the "*iris*," in order to regulate the quantity of light which shall be admitted through the pupil. The space in which the iris is suspended is filled with a fluid termed the "*aqueous*" humor.

Sclerotic Coat.—The sclerotic is the tough protecting coat of the eye, and consists of glistening fibres interlacing in all directions.* It covers about $\frac{4}{5}$ ths of the globe, the remaining $\frac{1}{5}$ th being completed by the cornea. The thickest part of the sclerotic coat is at the back of the globe (fig. 130): here it is perforated by the optic nerve a little on the nasal side of the axis of vision.† Around the optic nerve it is perforated by the ciliary arteries, veins, and nerves, for the supply of the choroid coat and the iris. Towards the front the sclerotic becomes much thinner, and about a quarter of an inch from the cornea it receives the insertion of the recti muscles; here again it is perforated by the anterior ciliary arteries which creep along the tendons of these muscles.

The inner surface of the sclerotica is coated by a thin layer of areolar tissue impregnated with dark pigment cells. It is sometimes called "*lamina fusca*."

[The tendons of the recti muscles expand, and near the cornea are buried, so to speak, into the sclerotic. These muscles groove this coat before they enter it, and hence the sclerotic is weakest at this situation; therefore the coats of the eye are more liable to give away at this point, either from external violence or internal suppuration.

The sclerotic is very sparingly supplied with capillary blood-vessels. Whether the nerves which run along its inner

* The sclerotic coat of the eye in fishes is of extraordinary thickness and density, for obvious reasons; and in birds this coat is further strengthened by a circle of bony plates, fourteen or fifteen in number, arranged in a series round the margin of the cornea. Similar plates are found in some of the reptiles, and particularly in the fossil ichthyosauri and plesiosauri.

† The optic nerve does not pass through a single large hole in the sclerotic but through a net-work of fibrous tissue. It is also very much constricted just at its entrance.

surface send any filaments into its interior, is not yet fully determined.]

To examine the cornea, it should be removed with the sclerotic coat. This is best done under water, by making a circular cut with scissors about a quarter of an inch from the margin of the cornea. With a little care, it is easy to take off the outer covering of the eye without injuring the dark choroid coat underneath it, or the ciliary ligament, or the iris. In the

Fig. 130.




DIAGRAM OF A VERTICAL SECTION OF THE EYE.

1. Anterior chamber filled with aqueous humor.
2. Posterior chamber.
3. Canal of Petit.

loose watery cellular tissue between the sclerotic and the choroid we observe the ciliary nerves coming forwards towards the iris: their white color makes them very conspicuous on the dark ground.

Cornea.—The cornea is the brilliant and transparent coat which occupies about one-fifth of the front part of the globe, and is, as it were, the window of the eye. However delicate it appears at first sight, yet it is quite as tough and thick as any part of the sclerotica. It is connected to this coat in the firmest possible manner: the margin of the sclerotica is beveled on the inside; that of the cornea on the outside; so that the one overlaps the other (fig. 130).

Structure.—As to the structure of the cornea, it consists of concentric layers of transparent tissue; by squeezing it between the finger and thumb, you can make the layers glide on each other. But these layers are not all composed of the same kind of tissue. The first stratum is conjunctival. The second stratum consists of a remarkably elastic tissue. The third stratum consists of transparent fibrous tissue; upon this the thickness and strength of the cornea mainly depend. The fourth* stratum is again elastic. The fifth and last stratum consists of epithelium. Such is the beautiful structure of the cornea. In its healthy state it contains no blood-vessels; they run back in loops as soon as they reach its circumference.†

[The transparency of the cornea depends upon the parallel condition of its layers and their proper distance from each other being maintained. Should these layers become disarranged in any way,—by pressure, for example,—the transparency is lost. (Squeezing the eye of an ox will illustrate this fact.) The posterior elastic layer is known also under the name of membrane of Demours or Decemet. At the edge of the cornea it breaks up into processes (pillars of the iris), which turn back and blend with the outer border of the iris, choroid, and sclerotic coats. The anterior elastic layer is analogous in structure to the latter, but thinner. The cornea is abundantly supplied with nerves; and the continuity of the sclerotic with the cornea is so perfect, that they cannot be torn apart; maceration, even, will not accomplish this.]

Choroid Coat—After the removal of the sclerotic and cornea, we expose the choroid coat,‡—a white ring, called the "*ciliary ligament*," which bounds its anterior part; also the iris, of which the outer circumference is attached to this ring (fig. 131).


* The great peculiarity of this layer is, that it is perfectly structureless; and when peeled off, it has a remarkable tendency to curl up. Maceration or boiling, or the action of acids, do not render it opaque, like they do the other layers of the cornea. Dr. Jacob calls it the "elastic cornea."—*Med. Chir. Trans.* vol. xii., p. 503.

† For a very elaborate investigation of the structure of the cornea, see Todd and Bowman's *Physiological Anatomy*, Part iii., p. 17.

‡ So called because its outer flocculent surface somewhat resembles the chorion or external investment of the ovum.

The choroid is the soft and flocculent tunic of the eye, remarkable for its dark color and great vascularity. When properly injected, and examined with the microscope, it is found to consist almost wholly of arterial and venous ramifications; the arteries being chiefly arranged on the inner, the

Fig. 131.



SCLEROTIC COAT REMOVED TO SHOW THE CHOROID, CILIARY LIGAMENT, AND NERVES.

veins on the outer surface. The dark color is adventitious, and is owing to a deposit in its texture of coloring matter termed "*pigmentum nigrum*."

Peculiarity of the Choroid Veins.—The veins of the choroid are placed on the outside of the membrane. When successfully injected with mercury, one finds that they are arranged with great regularity in drooping branches (*vasa vorticosa*), like a weeping willow (fig. 131), and that they converge to four nearly equidistant trunks, which, after running backwards for a short distance, perforate the sclerotica not far from the entrance of the optic nerve, and empty themselves into the ophthalmic vein.

The arteries ramify on the inner surface of the choroid: they perforate the sclerotica near the optic nerve, and then divide and subdivide into a very minute network, termed after the Dutch anatomist Ruysch, "*tunica Ruyschiana*."

Posteriorly there is a circular aperture in the choroid for

the passage of the optic nerve. In front the choroid is united to the ciliary ligament: it appears to stop short here; but this is not the case; for, under this ligament, it extends forwards round the circumference of the crystalline lens in a series of plaited folds, called the "ciliary processes."

Pigmentum Nigrum.—This is the coloring matter of the choroid. It is merely adventitious, for, if the choroid be washed in water or spirit, the color is entirely removed, leaving the membrane of a greyish tint. In man this pigment is dark brown, but in most animals it is jet black. Under the microscope, it is found to consist of colored granules contained in minute hexagonal cells. It exists on both surfaces of the choroid, but chiefly on the inner, where it forms a continuous stratum.* The use of the pigment is to absorb the rays of light which pass through the retina, and prevent their being reflected. It serves the same purpose as the black paint with which the inside of optical instruments is darkened. Albinos, in whom the choroid has little or no pigment, are consequently dazzled by daylight, and see better in the dusk.†

Ciliary Ligament.—The ciliary ligament (fig. 131) is a whitish grey ring, about $\frac{1}{8}$ th of an inch broad, situated just beneath the union of the sclerotica and the cornea. It serves as a connecting medium between several structures—namely, the choroid, iris ciliary processes, and sclerotica.‡

* Dalrymple speaks of a very delicate membrane as lining the inner surface of the choroid, for the purpose of keeping the dark pigment in its place. A similar membrane may be detected on the posterior surface of the iris; otherwise the pigment there would be apt to be washed away by the aqueous humor.

† In many of the nocturnal carnivorous quadrupeds, the inner surface of the choroid at the bottom of the eye presents a brilliant color and metallic lustre. It is called the tapetum. By reflecting the rays of light a second time through the retina, it probably causes the animal to see better in the dusk. It is the cause of the well-known glare of the eyes of cats and other animals; and the great breadth of the luminous appearance arises from the dilatation of the pupil.

‡ Some anatomists describe a *ciliary muscle*. They say its point of attachment is at the line of junction of the sclerotic and cornea; from thence it radiates backwards over the outer surface of the ciliary ligament, where its fibres are lost. Its action would be to approximate the ciliary processes and lens towards the cornea.

Sir Philip Crompton has noticed that this muscle is well developed in birds. In them its fibres are of the striped kind, just as the circular fibres of the iris are.

[When the sclerotic has been removed, the ciliary ligament may be seen as a white band, with the muscle of the same name close behind it (*annulus albidus*) outside the front of the choroid. The ligament is about $\frac{1}{40}$ th of an inch thick, and surrounds the iris. Its fibres are stiff, and resemble the properties of elastic tissues. The ciliary muscle is about $\frac{1}{10}$ th of an inch in width of unstriated fibres, which run in a wavy manner, the greater number being in the direction of the meridians, and are readily peeled off; removing these, other fibres are seen to pass in a transverse direction. The action of this muscle, according to H. Müller, is as follows: The circular fibres exert pressure upon the edge of the lens, so the latter becomes thicker,—longitudinal fibres causing tension of the vitreous, so that the posterior surface of the lens is prevented from shifting; and the peripheral pressure is chiefly confined to the anterior surface. Thus the lens is made more or less convex by the action of this muscle.]

Canal of Fontana.—Between the sclerotic, the cornea, and the ciliary ligament is placed a minute circular canal, termed “Fontana’s canal” (fig. 130). It is probably a venous sinus, for it can always be injected from the arteries.

Iris.—The iris is a movable curtain suspended freely in the clear fluid which occupies the space between the cornea and crystalline lens. The iris divides this space into two unequal parts, called the anterior and posterior chambers (fig. 130); these communicate with each other through a circular aperture in the centre, called the pupil.* Its use is to regulate the amount of light which shall be admitted into the eye: for this purpose its inner circumference is capable of dilating and contracting according to circumstances, while its outer circumference is immovably connected to the ciliary ligament.

The color of the iris varies in different individuals, and gives the peculiar tint and brilliancy to the eye. The coloring matter or pigment is contained in minute cells. The posterior surface of the iris, called the *uvea*,† is in all cases covered by a layer of black pigment.

* The size and shape of the pupil vary in different animals. In the bullock, sheep, horse, &c., it is oblong; in carnivorous quadrupeds it is often a mere vertical slit during the day, but dilates into a large circle at night.

† Strictly speaking, the term *uvea* was applied by the old anatomists to the

When the iris is laid under water, and viewed with a low magnifying power, its front surface looks shaggy; a number of fine fibres are seen converging from all sides towards the pupil: many of them unite and form arches. When the pupil is contracted these fibres are stretched, and *vice versâ*. Whether they co-operate in producing the dilatation of the pupil is uncertain.

The contractile power of the iris depends upon muscular fibres of the non-striped kind, arranged some in a radiating, others in a circular manner. The radiating converge towards the pupil; the circular are aggregated round the pupillary margin.*

Its Blood-Vessels.—When minutely injected, the iris appears to be composed almost entirely of blood-vessels;† so much so that some anatomists consider it to be a kind of erectile tissue, and that its power of contracting and expanding depends upon this property alone. Its blood-vessels are derived from two sources—the posterior or long, and the anterior or short ciliary arteries. The posterior perforate the sclerotica round the optic nerve, and then run on upon the choroid to the iris; the anterior proceed from the tendons of the recti, p. 533, and perforate the sclerotica round the margin of the cornea. It is from the enlargement of these latter vessels that the red zone round the cornea is produced in inflammation of the iris.

Its Nerves.—The nerves of the iris, twelve or thereabouts in number, proceed from the lenticular ganglion, and from the nasal branch of the ophthalmic division of the fifth pair, p. 167. They perforate the back of the sclerotica like the arteries, and run along the choroid to the iris.

Membrana Pupillaris.—Until the seventh or eighth month of foetal life, the pupil is closed by a delicate membrane, termed "*membrana pupillaris*." Its vessels are arranged in

choroid and iris collectively, which they very properly considered as one coat, and called the "*χιτων παγοειδής*," because its dark color made it like the berry of the grape.


* The circular fibres of the iris in the bird are of the striped variety, and discernible without difficulty.

† In well-injected preparations one may see that the chief blood-vessels are disposed in two circles on the front surface of the iris, one near the outer, and the other near the inner circumference.

loops, which converge towards the centre of the pupil. It has been lately discovered that this membrane, which has always been regarded as a distinct structure, is identical with the anterior layer of the capsule of the crystalline lens.*

Retina.—To obtain a view of the retina, the choroid coat must be removed while the eye is under water; this is readily done with the forceps and scissors. The optic nerve, having entered the interior of the globe through the sclerotic and the choroid, expands into the delicate nervous tunic called the retina. In passing through the coats of the eye, the nerve becomes suddenly constricted, and reduced to one-third of its diameter; just at this point, too, it projects slightly into the interior of the globe, forming a little prominence to which the term "*papilla conica*" has been applied.† In front the retina terminates in a thin serrated border (*ora serrata*), which fits into corresponding dentations in the posterior margin of the ciliary body.

Fig. 132.



ARTERIES OF THE RETINA.

Canal of Petit (inflated).
Zone of Zinn (exaggerated).

Precisely opposite the pupil there is a bright yellow spot in the retina, about 1-24th of an inch in diameter, fading off gradually at the edges, and having a black spot in the centre. This central spot was believed by its discoverer, the celebrated Soemmering,‡ to be a perforation; but it is now ascertained to depend upon the absence of the yellow color in the centre: so that the dark pigment of the choroid becomes conspicuous. These appearances are lost soon after death, and are replaced by a minute fold, into which the retina gathers itself, reaching from the centre of the spot to the prominence of the optic nerve. The use of this yellow spot is not understood.§

* See a paper by John Quekett in the Transactions of the Microscopic Society of London, vol. iii. p. 9.

† This prominence is remarkable, in that it is insensible to the rays of light.

‡ De foramine centrali, &c., retinae humanæ; in Comment. Soc. Gotting. t. 13.

§ In birds the retina has throughout the yellowish color seen only at one part in the human eye.

Its Elaborate Organization.—Although to the naked eye the retina appears nothing more than a soft semi-transparent membrane, yet, when examined with the microscope, it is found to be most minutely and elaborately organized. It consists of three layers,—an internal vascular layer, an external, termed “*membrana Jacobi*,” a middle or truly nervous layer. In the middle layer, the delicate fibres of the optic nerve spread out and ramify through a stratum of nerve-cells: how they ultimately terminate, whether by loops or in free extremities, is still doubtful. The external layer is formed by the ramifications of the *arteria centralis retinae*, which form a close network of blood-vessels throughout the nervous substance, for its nutrition. After a short maceration in water, the nervous substance can be brushed off with a delicate camel’s-hair pencil, and then, in an injected eye, the web formed by the vessels can be distinctly seen. The larger branches, however, are visible without injection: one of them runs round the free margin of the retina.

Membrana Jacobi.—But the part of the retina most remarkable for the singularity of its organization is the structure which forms its external layer. It is termed the “*membrana Jacobi*.”* When carefully examined with the high powers of a microscope, it is found to be composed of minute cylindrical, transparent rods, arranged like bristles in a brush, at right angles to the surface of the retina. Their outer extremities are imbedded, to a greater or less depth, in the dark pigment of the choroid; so that, when viewed from without, the rods have the appearance of a mosaic pavement.†

Ciliary Body and Processes—These structures are best seen when the globe has been divided by a vertical section into an anterior and a posterior half, the vitreous humor being left undisturbed. We then see a beautiful black disk, about three lines broad, surrounding the lens: it may be regarded as a continuation of the choroid. The posterior boundary of this ring is smooth and flat, and defined by a dentated line.

* After its discoverer, Dr. Jacob, of Dublin, who described it in the *Philosophical Transactions*, 1819.

† See Hannover, *Recherches Microscop. sur le Système Nerveux*, 1844.

The anterior part presents a number of longitudinal plaits or folds, from sixty to seventy in number, alternately long and short, and arranged in a radiated manner round the circumference of the lens. One of these is seen in the diagram p. 534. The entire ring is called the ciliary body, and the folds of it in front, the ciliary processes. The processes are kept in place by being attached to the ciliary ligament; they fit into corresponding depressions in the vitreous humor, and their free ends project for a short distance into the posterior chamber. They consist of convolutions of minute arteries, and their dark color arises from the pigment on their surface. Their use is unknown.

Aqueous Humor.—The aqueous humor consists of a few drops of clear watery fluid, which fills up the space between the cornea and crystalline lens. The iris floats freely in it, and divides the space into two chambers of unequal size—an anterior and a posterior. The posterior is much the smaller of the two: indeed, the iris is so close to the lens that they are separated by a mere film of fluid. This accounts for the frequent adhesions which are apt to take place, during inflammation, between the pupil and the capsule of the crystalline. Some anatomists describe the anterior chamber as lined by a serous membrane, which they call the membrane of the aqueous humor. It is true that there is a delicate layer of epithelium on the posterior surface of the cornea, but nothing like a continuous serous membrane can be demonstrated on the iris or the capsule of the lens. The anterior chamber is remarkable for the rapidity with which it absorbs and secretes, as is proved, in the one case, by the speedy removal of extravasated blood; in the other, by the rapid reappearance of the aqueous humor after the extraction of a cataract.

Vitreous Humor and Hyaloid Membrane.—The vitreous humor is a beautifully transparent, gelatinous-looking substance, which fills up nearly four-fifths of the interior of the globe (p. 534). It consists of a watery fluid contained in the meshes of a cellular structure, called the "*hyaloid membrane*," from its perfect translucency: the cells communicate freely with each other; for, if any part of it be punctured, the humor gradually drains off. The membrane itself is so delicate that it is difficult to obtain it separately; but it may be rendered

slightly opaque by strong spirit or diluted acids. It is of somewhat firmer consistence on the surface, so that it answers the purpose of a capsule for the vitreous humor, and is sufficiently strong to keep it in shape after the stronger tunics of the eye have been removed.*

In the foetus, a branch of the retinal artery runs through the centre of the vitreous humor, and ramifies on the back of the capsule of the lens. It is lodged in a tubular canal in the hyaloid canal; but little or nothing of this is seen in the adult.

Zone of Zinn.—The vitreous humor presents in front a deep depression, in which the crystalline lens is imbedded; around this depression is the "*zone of Zinn*."† This zone is best exposed by peeling off the ciliary body. It then appears like a dark disk and extends from the front margin of the retina nearly to the capsule of the lens: its surface is marked by prominent ridges which correspond with the intervals between the ciliary processes (fig. 130). Various opinions are entertained of the nature of this zone; but it does not appear to be anything more than the impression left by the pigment of the ciliary body upon the hyaloid membrane.

Canal of Petit.—If the transparent membrane between the zone of Zinn and the margin of the lens be carefully punctured, and the point of a small blow-pipe gently introduced, we may succeed in inflating a canal which encircles the lens, and, when inflated, resembles a circle of small glass beads: this is the canal of Petit, or "*canal godronné*" (fig. 130, p. 534). How this canal is formed, whether by the separation of the hyaloid membrane into two layers or not, and what is its use, are questions not easily answered.

Crystalline Lens.—The crystalline lens (fig. 130), is a per-

* The cells of the vitreous humor may sometimes be demonstrated by freezing the eye and then dividing it. The figure and size of the cells are shown by the portions of ice which they contain. Again, by macerating the eye in chromic acid, it is found that the vitreous humor is intersected by 180 delicate partitions, disposed like those in the pulp of an orange—with this difference, however, that the partitions do not quite reach the centre, but leave a cylindrical space in the axis of the humor. Up this space the central artery runs in the foetus.

† Zinn was Professor of Anatomy at Göttingen about the middle of the eighteenth century, and author of "*Descriptio Anat. Oculi Humani*."

fectly transparent solid body, situated immediately behind the pupil, and partly imbedded in vitreous humor. It is convex on both surfaces, but more so behind. Its shape and consistence vary at different periods of life. In early life it is nearly spherical and soft, but it becomes more flattened and firmer with advancing age. In the adult, its transverse diameter is about three-eighths of an inch; its antero-posterior, one-sixth of an inch.

The lens is kept in place by a capsule equally transparent as itself. The capsule is composed of tissue exactly similar to the elastic layer of the cornea. It is at least four times thicker in front than behind, as one might expect, for the sake of more effective support. No vascular connection whatever exists between the lens and its capsule.* The lens protrudes directly the capsule is sufficiently opened. How then is the lens nourished? By means of an extremely delicate layer of nucleated cells on its surface, which absorb nourishment from the capsule. Some anatomists speak of a layer of fluid (*liquor morgagni*) as existing between the lens and its capsule; but no such fluid can be detected during life, and if there be any after death, it is, in all probability, imbibed by the capsule from the aqueous humor.

[The lens measures from side to side, from one-third to half an inch, and from before backwards about one-fourth of an inch. The thickness increases from the circumference to the centre. In infancy the lens is situated farther forward than in the adult, and touches the iris. In the foetus it is nearly spherical; it is also reddish, soft, and not quite transparent. The lens is kept in position by its *suspensory ligament*, a transparent membranous structure, which is placed around the lens at the front of the hyaloid membrane, between the ora serata and the capsule. It unites with the ciliary processes, and hence

* The vessels of the capsule of the lens are derived from the arteria centralis retinae, and in mammalia, can only be injected in the foetal state. In the reptile, however, the posterior layer of the capsule is permanently vascular. According to Quekett, the membrana pupillaris of authors is nothing more than the anterior layer of the capsule. In taking the eye to pieces, it is quite a matter of accident whether the membrane adhere to the iris, or remain in its proper place in front of the lens. See Quekett's paper in the "Transactions of the Microscopic Society of London," vol. iii.

has a fixed point for supporting the lens. We find *in its structure elastic fibres.*]

Minute Structure.—The minute structure of the lens is very remarkable. It is gelatinous in consistence outside, but grows gradually denser towards the centre. After immersion in nitric acid, alcohol, or boiling water, it becomes hard and opaque. One may then see that it is divided into three equal parts, by three lines which radiate from the centre to within one-third of the circumference. Each of these portions is composed of hundreds of concentric layers, arranged one within the other, like the coats of an onion. But this is not all: if we examine any single layer with the microscope, we find that it is made up of fibres about $\frac{1}{5000}$ th of an inch in thickness, and connected together by finely serrated edges. This beautiful dovetailing of the fibres of the lens was first pointed out by Sir David Brewster; and, to see it in perfection, one ought to take the lens of the common codfish

The use of the lens is to bring the rays of light to a focus upon the retina.

DISSECTION OF THE ORGAN OF HEARING.*

THE parts constituting the organ of hearing should be examined in the following order:—1. The outer cartilage; 2. The external meatus, which leads to 3. The tympanum; and lastly, the labyrinth, or internal ear, comprising the vestibule, cochlea, and semicircular canals.

Pinna.—Every one is familiar with the general form of the pinna, or fibro-cartilage of the ear; but anatomists have given names to its different parts. The outer folded border is called the *helix*; the ridge within it, the *ante-helix*. This bifurcates towards the front, and bounds the fossa of the ante-helix. The conical eminence in front of the meatus is termed the *tragus*: from this, hair generally grows. Behind the *tragus*, and separated from it by a deep notch, is the *anti-tragus*. The

* The illustrations of the organ of hearing have been already published in my work on "Human Osteology."

lobule is that part from which the ear-rings are suspended. The deep hollow which collects the sonorous vibrations, and directs them into the meatus, is termed the *concha*. The groundwork of the pinna is a fibro-cartilage, which is attached by an anterior ligament to the zygoma, and by a posterior to the mastoid process.

Muscles of the Pinna.—The muscles which move the cartilage of the ear as a whole have been described (p. 157). Some other little muscles extend from one part of the cartilage to the other; but they are so indistinct, that, unless the body be very muscular, your attempt to find them will be made in vain. The five following are usually described by anatomists,—four on the front of the pinna, and one behind it:—

- a. The *musculus major helix* runs vertically along the front margin of the pinna.
- b. The *m. minor helix* lies over that part of the helix which comes up from the bottom of the concha.
- c. The *m. tragicus* lies vertically over the outer surface of the tragus.
- d. The *m. anti-tragicus* proceeds transversely from the anti-tragus to the helix.
- e. The *m. transversus* is on the back of the pinna: it passes from the back of the concha to the helix.

The arteries of the pinna are derived from the posterior auricular and the temporal. The nerves are furnished by the auriculo-parotidean branch of the cervical plexus, and the temporo-auricular branch of the inferior maxillary.

Meatus Auditorus.—This passage leads down to the membrana tympani, or drum of the ear. It is formed partly by a tubular continuation of the concha, partly by an osseous canal in the temporal bone. It is not a straight tube, but inclines at first upwards and forwards, and then curves a little downwards.* [When we wish to examine the bottom of the ear, therefore, we must draw the external ear upwards and backwards, unless we use a speculum.] Its length is rather

* In order to obtain a correct knowledge of the length and dimensions of the meatus, we ought to make sections through it in different directions, or a cast of it in common plaster.

more than an inch. It is not throughout of the same calibre ; the narrowest part is about the middle : hence the difficulty in extracting foreign bodies which have gained access to the bottom of it. The true skin and the cuticle are continued down it, and, becoming gradually thinner, form a cul-de-sac over the membrana tympani. Only the outer portion is furnished with ceruminous glands, of which the peculiar bitter secretion is for the purpose of keeping the passage moist, and preventing insects from lodging in it.

Tympanum.—The tympanum, or middle ear, is an irregular cavity scooped out of the petrous part of the temporal bone, and lined by mucous membrane. It is filled with air, which is freely admitted through the Eustachian tube ; so that the atmospheric pressure is equipoised on both sides of the membrana tympani. It contains a chain of small bones, of which the use is to communicate the vibrations of the membrana tympani to the internal parts of the ear. For this purpose one end of the chain is attached to the membrana tympani, the other to the fenestra ovalis. At the back part of it is an opening for the passage of air into the mastoid cells. Lastly, a nerve, called the chorda tympani (a branch of the portio dura) runs across it.

[The cavity of the tympanum runs for six or eight lines in the cavity of the petrous bone : entire length about two inches ; diameter a line and a half.]

Membrana Tympani.—The membrana tympani completely closes the bottom of the meatus auditorius. It is nearly circular, and its circumference is set in a bony groove, so that it is stretched somewhat like the parchment of a drum on the outer wall of the tympanum. Its plane is not vertical, but slants from above downwards, forming, with the lower part of the meatus, an angle of 45° : nor is it quite flat, but slightly conical, the apex being directed inwards towards the tympanum, and firmly united to the handle of the little bone called the “malleus.” The structure of the membrane is essentially fibrous ; some of the fibres radiate from the centre, others are circular. Its inner surface is lined by mucous membrane ; its outer surface is covered by an extremely thin layer of the true skin. This sufficiently accounts for the great sensibility of the membrane, and its vascularity when inflamed.

Eustachian Tube.—For a complete account of the Eustachian tube see p. 139. We need only say here that it proceeds from the anterior part of the tympanum downwards and forwards to the pharynx.

Tympanic Bones.—The four little bones in the tympanum are named, after their fancied resemblance to certain implements, the *malleus*, *incus*, *os orbiculare*, and *stapes*. They are articulated to each other by perfect joints, and are so placed that the chain somewhat resembles the letter Z.* Their use is to transmit the vibrations of the membrana tympani to the membrane of the fenestra ovalis, and, through it, to the fluid contained in the internal ear. But they have another use, which would be incompatible with a single bone—namely, to permit the tightening and relaxation of the membrana tympani, and thus adapt it either to resist the impulse of a very loud sound, or to favor a more gentle one.

[The *os orbiculare* is often in the adult a part of the *incus*.]

Muscles of the Tympanum.—These little muscles, by moving the tympanic bones, tighten or relax the membrane of the tympanum. The "*tensor tympani*" runs above and parallel to the Eustachian tube, from the cartilaginous part of which it arises. It passes backwards, and terminates in a round tendon, which enters the fore part of the tympanum through a special bony canal, and is inserted into the short process of the malleus. Its action is to draw the membrana tympani inwards, and thus render it tense. The "*laxator tympani*" arises from the spine of the sphenoid, and is inserted into the long process of the malleus. The "*stapedius*" arises from a tube in the pyramid, and its tendon is inserted into the

* The handle of the malleus is nearly vertical, and attached along its whole length to the upper half of the membrana tympani. The long process (*processus gracilis*) projects at right angles from the body of the bone, runs into the Glasserian fissure, and receives the insertion of the *laxator tympani*. The short process receives the insertion of the *tensor tympani*.

The *incus* or anvil bone, is shaped like a bicuspid molar tooth with unequal fangs. Its broad part articulates with the malleus; its long process articulates with the stapes, or stirrup bone, through the *os orbiculare*; its short process is directed backwards, and its point is fixed in a small hollow at the commencement of the mastoid cells.

The stapes is horizontal, and its base is attached to the membrane covering the fenestra ovalis.

neck of the stapes. Its precise use is not thoroughly understood.

Chorda Tympani.—A branch of the portio dura (“chorda tympani”) crosses the tympanum between the handle of the malleus and the long process of the incus (see p. 177).

On the inner wall of the tympanum—that is, opposite to the membrana tympani—observe in the dry bone the following objects:—Beginning from above there is an opening, called the “*foramen ovale*:” it leads into the vestibule of the internal ear, but is closed in the recent state by a membrane to which is attached the base of the stapes. Below the foramen ovale is a bony prominence called the “*promontory*.”* Still lower is another opening, called the “*foramen rotundum*:” it leads into the tympanic scale of the cochlea, but is closed in the recent state by membrane. Immediately behind the foramen ovale is a small conical eminence, named the “*pyramid*:” there is a minute aperture in the summit of it, from which the tendon of the stapedius emerges. Lastly, at the back part of the tympanum is the opening which leads into the air-cells of the mastoid process.

The tympanum is supplied with blood by a branch of the internal maxillary, which runs in through the fissura Glasseri; 2, by the stylo-mastoid branch of the posterior auricular; 3, by small branches which enter with the Eustachian tube.

Internal Ear.—This, in consequence of its complexity, is very appropriately termed “the labyrinth.” In a general way we may say that it consists of cavities excavated in the most compact part of the temporal bone. These cavities may be divided into three—a middle one, called “the vestibule,” as being a centre in which all communicate with each other; an anterior, named, from its resemblance to a snail’s shell, the cochlea; and a posterior, consisting of three semicircular canals. These cavities are filled with a clear fluid, called the endo-lymph, and contain a membranous expansion (the membranous labyrinth), upon which the filaments of the auditory nerve are expanded.

Vestibule.—The vestibule, or central chamber, communi-

* This promontory is occasioned by the first turn of the cochlea. Upon it ramifies the tympanic plexus of nerves formed by the sympathetic and the glosso-pharyngeal.

cates in front with the cochlea, through the scala cochleæ; behind, with the semicircular canals; on the outside with the tympanum, through the foramen ovale; on the inside, with the bottom of the meatus auditorius internus.*

Semicircular Canals.—The semicircular canals, three in number, are situated above and rather behind the vestibule. Each canal forms the greater part of a circle, and opens at each extremity into the vestibule: therefore, there should be six apertures for them; but, in point of fact, there are only five, since one of the apertures is common to the extremity of two canals. The canals are not precisely of equal diameter throughout; each presents at one end a dilatation termed the "*ampulla*." This dilatation is the most important part of the canal, because it corresponds to a similar dilatation of the membranous sac upon which the auditory nerve expands. Each canal differs in its direction: they are named accordingly *superior*, *posterior*, and *external*. The *superior* s. c. is also the most anterior of the three: its direction is vertical, and runs across the petrous bone: the ampulla is at the outer extremity. The *posterior* s. c. is also vertical, runs parallel to the posterior surface of the petrous bone, and consequently, at right angles to the preceding: the ampulla is at the lower end. The *external* s. c. is horizontal in position, with the convexity of the arch directed backwards: the ampulla is at the outer end.

Cochlea.—The cochlea is the most anterior part of the internal ear: it very closely resembles a common snail's shell, and is placed so that the base of the shell corresponds to the bottom of the meatus auditorius internus, while the apex is directed forwards and outwards. It consists of the spiral convolutions of two parallel and gradually tapering tubes. The partition by which they are separated is termed the "*lamina spiralis*." In the dry bones this partition is only partial; but, in the recent state, it is completed by a membrane. At the very apex of the shell (*helicotrema*) the partition is altogether absent, so that here the tubes communicate with each other. These tubes are called the scales of the cochlea, and are filled

* In some instances there is the opening of a small canal into the vestibule, termed the "*aqueductus vestibuli*." It leads to the posterior surface of the temporal bone, and contains a small vein.

with fluid. The one opens into the vestibule, and is therefore called the vestibular scale; the other leads to the membrane which closes the foramen rotundum of the tympanum, and is termed the tympanic scale. If unwound, they would be about $1\frac{1}{2}$ inch long. Each makes two turns and a half round a central pillar, from left to right in the right ear, and *vice versâ* in the left.

The central pillar of the cochlea is called the "*modiolus*." It is of considerable thickness at the base, but gradually tapers towards the apex. Its interior is traversed by numerous canals, for the purpose of transmitting the filaments of the auditory nerve. One of these canals, larger than the others, runs down the centre of the modiolus nearly to the apex.

The *lamina spiralis*—the partition between the two tubes or scales of the cochlea—is made up, on the inner half, of bone, on the outer half, of membrane. The bony part has a number of minute canals in it, which come off at right angles from the modiolus. They are for the lodgment of the filaments of the auditory nerve in their course to the membranous part, which is the most important element of the cochlea, since it receives the undulations of the fluid in the interior.*

The osseous labyrinth is lined throughout by a delicate fibro-serous membrane, which secretes the fluid called the "peri-lymph."

Membranous Labyrinth.—If the bony labyrinth just now described be properly understood, we can have no difficulty in comprehending the membranous labyrinth in its interior—a structure intended to support the ultimate ramifications of the auditory nerve, and to expose them to the undulations of the fluid in the internal ear.

The membranous labyrinth, then, is nothing more than a sac, situated in the vestibule,† and from this there pass off

* There is an extremely delicate little muscle, termed the "*cochlearis*," for the purpose of tightening or relaxing, according to circumstances, the membranous part of the *lamina spiralis*. It is placed along the outer circumference of the membrane, and, in fact, forms an integrant part of it. Its fibres are of the non-striped kind, like the ciliary muscle of the eye. See Todd and Bowman, *Phys. Anat.* Part iii. p. 79.

† Strictly speaking, the sac in the vestibule is constricted, so as to appear like two sacs of unequal size. The larger of the two is generally called the

three membranous semicircular canals within the bony ones. What was said of the bony canals applies equally to the membranous. They present the same dilatations or ampullæ at one end, and just at this part they nearly fill their bony cases ; but in the rest of their extent the diameter of the membranous canal is not more than one-third that of the bony one.

The membranous labyrinth is protected, inside and out, by fluid. There is the proper fluid in the interior, termed the "endo-lymph," and the thin layer of fluid, the "peri-lymph," between it and the bone.

Distribution of the Auditory Nerve. — The auditory nerve, or *portio mollis* of the seventh pair, passes down the *meatus auditorius internus*, and, at the bottom of it, divides into an anterior and a posterior branch : in other words, a branch for the cochlea, and a branch for the vestibule. These nerves then break up into numerous fasciculi, which pass through the foramina at the bottom of the *meatus* into the osseous labyrinth. Here the filaments are grouped into six bundles, corresponding to the parts which they supply—namely, two for the vestibular sac, one for each of the ampullæ of the semicircular canals, and one for the cochlea.

The cochlear nerve divides into filaments, which run through the canals of the *modiolus*, and then along those of the *lamina spiralis*, in order to terminate upon the membranous part of this lamina. The precise manner in which the filaments terminate is still dubious : according to Breschet,* they communicate and form a series of minute arches.

Respecting the other nerves, little more need be said than that their ultimate ramifications are lost upon the vestibular

common sinus, is nearer to the semicircular canals, and communicates with them. The smaller, called the saccule, is nearer to the cochlea, and communicates with its vestibular scale. Both sacs are filled with the endo-lymph, besides which, each contains a minute quantity of calcareous matter, called by Breschet the otolithes. These masses of cretaceous substance seem to be suspended in the fluid contained in the sacs by the intermedium of a number of nerve-filaments proceeding from the auditory nerve. From the universal presence of these chalky bodies in the labyrinth of all the mammalia, and from their much greater hardness and size in aquatic animals, there is little reason to doubt that they perform some office of great importance in the physiology of hearing.

* *Recherches Anat. et Phys. sur l'Organe de l'Ouïe, &c.* (Mém. de l'Acad. de Med., t. v. fasc. iii. 1836.

sac and upon the ampullæ of the semicircular canals : some of them, however, pass into the sac, and come into contact with the otoconies, or ear-dust, in its interior.

Blood-Vessels of the Labyrinth.—The internal auditory artery—a branch of the basilar—runs with the auditory nerve to the bottom of the meatus, and divides into branches corresponding with the divisions of the nerve. Its ultimate ramifications terminate, in the form of a fine network, on the membranous labyrinth, and on the spiral lamina of the cochlea. The auditory vein pours its blood into the superior petrosal sinus.

DISSECTION OF THE MAMMARY GLAND.

THE form, size, position, and other external characters of the mammary gland, are sufficiently obvious. We would merely observe that the longest diameter of the gland is in a direction upwards and outwards towards the axilla ; that the thickest part is at the centre ; and that the fullness and roundness of the gland depend upon the quantity of fat which is situated about it and between its lobes. Its deep surface is flattened in adaptation to the pectoral muscle, to which it is loosely connected by an abundance of areolar tissue.

It is inclosed by a fascia which not only supports it as a whole, but penetrates into its interior, so as to form a framework for its several lobes : hence it is that, in cases of mammary abscess, the matter is apt to be circumscribed, not diffused.

The *nipple* projects a little below the centre ; it is surrounded by a colored circle termed the “areola :” this circle is of a rose-pink color in virgins, but, in those who have borne children, of a dark brown. It begins to enlarge and grow darker about the second or third month of pregnancy, and these changes continue till parturition. The areola is also abundantly provided with papillæ, and with subcutaneous sebaceous glands, for the purpose of lubricating the surface during the sucking of the child.

[The breasts, at the nipples, are projected *forwards* and *outwards*, and the nipple slightly turned upwards. “This is one of the most beautiful provisions in nature, both for the

mother and child. To the mother because the child rests upon her arm and lap, in the most convenient position for sucking; if the nipple and breast had projected directly forward, the child must have been supported before her by the mother's hands, in a most inconvenient and fatiguing position, instead of reclining upon her side and arm; while the nipple is turned outward for the child to receive it, as the child lies in its mother's arms."*]

Structure.—The gland itself consists of distinct lobes held together by firm connective tissue, and provided with separate lactiferous ducts. Each lobe divides and subdivides into lobules, and the duct branches out accordingly.† Traced to their origin, we find that the ducts commence in clusters of minute cells, and that the blood-vessels ramify upon these cells in rich profusion; altogether, then, a single lobe might be compared to a bunch of grapes, of which the stalk represents the main duct. The main ducts from the several lobes, from fifteen to twenty in number, converge towards the nipple, and, just before they reach it, become dilated into small sacs or reservoirs two or three lines wide; after this they run up to the apex of the nipple, and terminate in separate orifices.

The arteries of the gland are derived from the long thoracic and internal mammary: the nerves come from the cutaneous branches of the intercostal nerves.

DISSECTION OF THE SCROTUM AND TESTIS.

Structure of the Scrotum.—The scrotum is composed of six tunics:—1, The skin; 2, The tunica dartos; 3, A layer of cellular tissue; 4, The spermatic fascia derived from the external abdominal ring; 5, The cremaster, or suspensory muscle; 6, The infundibuliform fascia derived from the internal ring.

I do not mean to say that each of these coverings can be

* Sir Astley Cooper, *Anat. and Diseases of the Breast*, p. 23.

† It is observed, in some cases, that one or more lobules run off to a considerable distance from the main body of the gland, and lie imbedded in the subcutaneous tissue. One should remember this when it is necessary to remove the entire gland.

demonstrated under ordinary circumstances, because they are so blended together; but they can be shown in the case of old and large herniæ when in a state of hypertrophy.

Dartos.—The dartos is a thin layer consisting of muscular fibres of the non-striped kind, like those of the bladder and intestines. Its use is to corrugate the loose and extensible skin of the scrotum, and in a measure to support and brace the testicle.

Layer of Cellular Tissue.—Beneath the dartos is a large quantity of loose cellular tissue, remarkable for the total absence of fat. Together with the dartos, it forms a vertical partition between the testicles, termed "*septum scroti*." It is not a complete partition, since air or fluid will pass from one side to the other. The great abundance and looseness of this

Fig. 133.




DIAGRAM OF THE TESTICLE.

tissue explains the enormous swelling of the scrotum in cases of anasarca, and in cases where the urine is effused into it in consequence of the urethra having given way.

The cremaster muscle and infundibuliform fascia have been described (pp. 297, 300).

Testis.—The testicle is a gland of an oval shape, with flattened sides, suspended obliquely, so that the upper end points forwards and outwards, the lower end in the reverse direction. The left is generally a little the lower of the two, to obviate collision with its fellow. The ordinary weight of the gland is

about six drachms ; but few organs present greater variations in size and weight, even in men of the same age : generally speaking, the left is the larger.

Epididymis.—Along the posterior part of the gland is placed a long narrow body, termed the "*epididymis*:" this is not a part of the testicle, but an appendage to it, formed by the convolutions of its long excretory duct. Its upper larger end is called the "*globus major*," and is connected to the testicle by the efferent ducts ; the lower end, "*globus minor*," is only connected to the testicle by fibrous tissue.

[The left testicle is usually lower than the right for two reasons, says Sir A. Cooper ; the one to permit the disposition of the penis to one side of the median line, the other to allow of the one testicle gliding upon and above the other, in close adduction and crossing of the thighs. The testicle may be arrested in any portion of its course, or may be found in abnormal positions, as in the perineum. The average measurements of the testicle are from twenty-two to twenty-four lines in length, twelve to eighteen in breadth, and from seven to eight lines in thickness.]

Proper Coverings of the Testicle.—The proper coverings of the testicle are—1, a serous membrane, called the "*tunica vaginalis*," to facilitate its movements ; 2, a strong fibrous membrane, called the "*tunica albuginea*," to support and form a case for the glandular structure within ; 3, a delicate stratum of minute blood-vessels which some anatomists have described as a distinct coat under the name of *tunica vasculosa*.

Tunica Vaginalis.—The *tunica vaginalis* is a serous sac, one part of which (*tunica vaginalis propria*) adheres closely to the testicle ; the other (*tunica vaginalis reflexa*) is reflected loosely around it. If the sac be laid open you see that it completely covers the testicle, except behind, where the vessels and ducts are placed (fig. 133) ; and that it also covers part of the outer side of the epididymis. The interior of the sac is smooth and polished, like all other serous membranes, and lubricated by a little fluid. An excess of this fluid gives rise to the disease termed "*hydrocele*."

The *tunica vaginalis* was originally derived from the peritoneum. In some subjects it still communicates with that

cavity by a narrow neck, and is therefore liable to become the sac of a hernia. Such herniæ are termed *congenital*—a bad name, since they do not, as a matter of course, take place at birth, but often in adult age. Sometimes the communication continues through a very contracted canal, open to the passage of fluid only; or the communication may be only partially obliterated, and then one or more isolated serous sacs are left along the cord. Such an one, when distended by fluid, gives rise to hydrocele of the cord.


Tunica Albuginea.—This tunic is a dense, inelastic membrane, composed of fibrous tissue, interlacing in every direction, analogous to the sclerotic coat of the eye. It completely invests the testicle, but not the epididymis. At the posterior part of the gland it penetrates into its substance for a short distance, and forms an incomplete vertical septum, termed, after the anatomist who first described it, "*corpus Highmori*," and subsequently by Sir A. Cooper, the "*mediastinum testis*" (fig. 133). This septum transmits the blood-vessels of the gland, and contains, also, the network of seminal ducts, called the rete testis, shown in diagram 133.

From the mediastinum testis are given off in all directions a number of slender fibrous cords, which traverse the interior of the gland, and are attached to the inside of the tunica albuginea. They serve to maintain the general shape of the testicle, to support the numerous lobules of which its glandular substance is composed, and to convey the blood-vessels into it. These tie-beams (*trabeculi testis*), as well as the mediastinum from which they proceed, are readily seen on making a transverse section through the gland (fig. 134).

Tunica Vasculosa.—Respecting the so-called tunica vasculosa, nothing more need be said than that it consists of a multitude of fine blood-vessels, formed by the ramifications of the spermatic artery, and held together by delicate cellular tissue. It lines the inner surface of the tunica albuginea, and gives off vessels which run with the fibrous cords into the interior of the gland.

Glandular Structure.—When the testicle is cut into, its interior structure looks soft and pulpy, and of a reddish-grey color. It consists of an innumerable multitude of minute convoluted tubes (*tubuli seminiferi*). For convenience of

package they are arranged in lobules, between four and five hundred* in number, of various sizes, and contained in the



TRANSVERSE SECTION
THROUGH THE TESTI-
CLE.

Diagrammatic.

1. Spermatic artery.
2. Vas deferens.
3. Deferential artery.
4. Epididymis.
5. Mediastinum testis.
- 6, 6. Cavity of tunica vaginalis.

(The dots show the reflections
of the tunica vaginalis.)

compartment formed by the fibrous cords proceeding from the mediastinum testis. Only a few of these lobules are shown in diagram 134. Though disposed in lobules, still they communicate with each other, and thus form one vast network of tubes. The secretion from them is carried off by some forty or fifty straight vessels (*vasa recta*), which penetrate the mediastinum testis and there form a plexus of seminal tubes, termed the "*rete testis*." This lies along the back of the gland. From the upper part of the rete the secretion is carried away to the large end of the epididymis by fifteen or twenty tubes, termed, "*vasa efferentia*." These, after forming a vast number of coils, termed, "*coni vasculosi*," which collectively constitute the globus major of the epididymis, ultimately terminate, one after the other, in a single duct, the commencement of the vas deferens.

Commencing, then, in the globus major of the epididymis, the vas deferens descends, making a series of extremely tortuous coils, which alone form the globus minor.† From the lower part of the globus minor the vas deferens ascends, joins the other component parts of the spermatic cord, passes through the inguinal canal, winds round the back part of the bladder, and empties itself into the prostatic part of the urethra. The length of the vas deferens was estimated by Monro at upwards of thirty feet. The same anatomist calculated that the semen, before it arrived at the vas deferens, had to transverse a tube forty-two feet in length.

* This estimate is according to Krause, Müller's Archiv. für Anat. 1837.

† A little blind duct, called *vasculum aberrans*, is sometimes connected either to the epididymis or the vas deferens.

Spermatic Cord.—The spermatic cord is composed of the spermatic vessels, nerves, and absorbents, of the vas deferens, with the little deferential artery (a branch of the superior vesical), of the cremaster muscle and the cremasteric artery. Its coverings have been described with the anatomy of the parts of hernia, p. 295.

The course of the spermatic arteries and veins has been described in the dissection of the abdomen, p. 330. We will only observe that the artery is remarkably tortuous as it descends along the cord, that it enters the back part of the testicle, and breaks up into a multitude of fine ramifications, which spread out on the inner surface of the tunica albuginea. The spermatic *veins* leave the testicle at its back part, and, as they ascend along the cord, become extremely tortuous, and form a plexus termed "*pampiniform*." It is usually stated that these veins are destitute of valves; and this fact is adduced as one of the reasons for the occurrence of "varicocele." But it is certain that the larger veins do contain valves.

The *absorbents* of the testicle terminate in the lumbar glands: hence these glands become affected in malignant disease of the testicles.

The *nerves* of the testicle are derived from the sympathetic. They run down from the abdomen with the spermatic arteries (p. 334). This accounts for the stomach and intestines sympathizing so readily with the testicle, and for the constitutional effects of an injury to it.

Descent of the Testicles.—The testicle is originally developed in the lumbar region, immediately below the kidneys; and it is loosely attached to the back of the abdomen by a fold of peritoneum, termed the "*mesorchium*," along which its vessels and nerves run up to it, as to any other abdominal viscus. From the lower end of the gland there proceeds to the bottom of the scrotum, a contractile cord termed the "*gubernaculum testis*."* By the gradual contraction of this, the organ is brought into the scrotum. It begins to slide

* Mr. Curling considers the gubernaculum testis to be a muscular cord. See his Observations on the Structure of the Gubernaculum, and on the Descent of the Testis in the Fœtus: Medical Gazette, April 10, 1841.

down from the loins about the fifth month, reaches the ring at the seventh, and about the ninth has entered the scrotum. Its original peritoneal coat is retained throughout ; but, as it enters the inguinal canal, the peritoneal lining of the abdomen is pouched out before it, and eventually becomes the tunica vaginalis reflexa. Immediately after the descent of the testis, its serous bag communicates with the abdomen, and in the lower animals continues to do so through life.* But in the human subject the canal of communication soon begins to close. It begins to close at the top first, and the closure is generally complete in a child born at its full time. The final purpose of this is to provide against the occurrence of ruptures, to which man, from his erect attitude, is so much more exposed than animals. At the end of the first month after birth, the canal is entirely obliterated from the abdominal ring to the testis. Sometimes, however, this obliteration fails to take place, or is only partial ; hence may arise congenital hernia, or hydrocele. The possible existence of a communication between the tunica vaginalis and the peritoneal cavity of the abdomen, is the reason why we do not inject hydroceles in infants.

* According to Professor Owen, the African orang-outang (*Simia troglodytes*) is the only exception to this rule. In this animal it is interesting to observe that the lower extremities are more fully developed as organs of support, and there is a ligamentum teres in the hip-joint.

TABLE OF THE CHIEF ARTERIES OF THE HEAD AND NECK.

Arch of the aorta furnishes to the neck.	1. Brachio- cephalic . . .	1. Common carotid . . .	1. External carotid . . .	1. Superior thyroid . . .	{ Hyoid branch laryngeal thyroid.			
				2. lingual	{ Hyoid branch dorsal lingual sublingual ranine.			
				3. facial	{ Inferior palatine br'nch tonsillitic glandular submental inferior labial coronary . { inferior lateral nasal { superior angular.			
				4. occipital	{ Meningeal branch posterior cervical.			
				5. posterior auricular . . .	{ Stylo-mastoid branch auricular mastoid.			
				6. ascending pharyngeal.	{ Pharyngeal branches meningeal.			
				7. temporal	{ Auricular parotid articular transverse facial middle temporal anterior temporal posterior temporal.			
				8. internal maxillary . . .	{ Inferior dental middle meningeal muscular posterior dental infra-orbital spheno-palatine descending palatine vidian pterygo-palatine.			
				2. subcla- vian.	2. internal carotid. . .	1. Arteriæ receptaculi	1. ophthalmic	{ Lachrymal supra-orbital central of the retina ciliary muscular ethmoidal palpebral frontal nasal.
							2. anterior cerebral	
3. anterior communicating								
4. middle cerebral								
5. posterior communicating								
6. choroid.								
2. Left common carotid	2. internal mammary	1. Vertebral	Anterior spinal					
			posterior spinal					
			inferior cerebellar					
			posterior meningeal					
			transverse basilar					
			anterior inferior cerebellar					
			superior cerebellar					
			posterior cerebral.					
			3. Left subcla- vian.	3. thyroid axis . . .	2. internal mammary	Inferior thyroid	Ascending cervical.	
						supra-scapular	{ Supra-spinal infra-spinal.	
transverse cervical. . .	{ Superficial cervical posterior scapular.							
4 superior intercostal.	{ deep cervical.							

TABLE OF THE CHIEF VEINS OF THE HEAD AND NECK.

Brachio-cephalic is formed by the union of	Internal jugular .	1. Lateral sinus .	Superior longitudinal sinus
			inferior longitudinal sinus
			straight sinus
			occipital sinus
			ophthalmic vein
			superior petrosal
			inferior petrosal.
		2. ascending pharyngeal .	Meningeal branches
			pharyngeal.
		3. lingual . . .	Superficial dorsal
			lingual
		Angular	ranine.
			Supra-orbital
		inferior palpebral	frontal
			palpebral
		dorsal and lateral nasal veins	nasal
			Alveolar branches
		anterior internal maxillary	infra-orbital
			descending palatine
		4 facial	naso-palatine
			vidian.
		coronary .	superior
			inferior
		buccal	
		massetine	
		labial	
		submental	
		inferior palatine	
		tonsillitic	
		glandular	
		5. occipital . . .	Mastoid vein
			cervical.
		6. superior thyroid . . .	Thyroid
			laryngeal.
	subclavian.	1. Vertebral . . .	Spinal
			deep cervical
			ascending cervical.
		1. Internal maxillary	Middle meningeal
			inferior dental
		deep temporal	pterygoid
			masseteric.
		2. temporal	Anterior
			posterior
		middle temporal	parotid
			anterior auricular
		transverse facial.	
		3. posterior auricular	Auricular
			stylo-mastoid.
		4. branch to the internal jugular	
		5. supra-scapular . . .	Supra-spinal
			infra-spinal.
		6. transverse cervical	Superficial cervical
			posterior scapular.
		3. anterior jugular.	

TABLE OF THE CRANIAL NERVES.

1. First nerve Filaments to the nose.
2. Second nerve To retina of the eyeball.
3. Third nerve To muscles of the orbit.
4. Fourth nerve To superior oblique muscle.

5. Fifth or tri-facial nerve .

Ophthalmic .	{	Meningeal	{	Lachrymal palpebral.	
		lachrymal			
	{	frontal	{	Supra-orbital supra-trochlear.	
	{	nasal.	{	To lent. ganglion ciliary nerves infra-trochlear nasal.	
ophthalmic or lenticular ganglion . .	{	Connecting branches	{	To nasal nerve to the third nerve to sympathetic.	
		ciliary nerves.			
superior maxillary .	{	Orbital branch.	{	Malar temporal.	
		to Meckel's ganglion			
		posterior dental			
		anterior dental infra-orbital.			
Meckel's ganglion . .	{	Internal branches	{	Nasal naso-palatine.	
		ascending		To the orbit.	
		descending		Anterior palatine posterior external.	
		posterior.		Vidian pharyngeal.	
inferior maxillary .	{	Small or muscular part	{	deep temporal masseteric buccal pterygoid.	
		large or sensory part.	{	Auriculo-temporal	
				{	Articular, and to meatus, parotid, auricular, temporal.
			{	gustatory	To sub-maxillary and sublingual ganglia to hypoglossal to the tongue.
				inferior dental	Mylo-hyoid labial incisor.
otic ganglion .	{	Connecting branches	{	To Jacobson's nerve to the fifth and sympathetic.	
		branches for muscles			
sub-maxillary ganglion . .	{	Connecting branches	{	To the gustatory, chorda tympani, and sympathetic.	
		branches to the glands and the mucous membrane of the mouth.			

TABLE OF CRANIAL NERVES.—Continued.

6. Sixth nerve To external rectus.

7. Seventh nerve	{	Portio dura . . .	{	Connecting branches . . .	{	To join auditory to Meckel's ganglion tympanic and sympathetic nerves the chorda tympani.									
			{	branches for distribution . . .	{	Posterior auricular digastric branches stylo-hyoid branch temporo-facial. . .	{	Temporal malar infra-orbital.							
		{	portio mollis	{	{	{	{	{	{						
										{	nerve to cochlea . . .	{	To the common sac to the sacculæ to the semi-circular canals.		
8. Eighth nerve.	{	Glosso-pharyngeal	{	Connecting branches . . .	{	To vagus to sympathetic. Jacobson's nerve . . .	{	Joins otic ganglion supplies tympanum							
			{	branches for distribution . . .	{	To carotid artery to the pharynx tonsillitic branches muscular lingual.									
			{	pneumo-gastric . . .	{	{	{	{	{						
										{	Connecting branches	{	To glosso-pharyngeal sympathetic and auricular nerves to the hypoglossal		
		{								branches for distribution . . .	{	{	{	{	{
			{	superior laryngeal . . .	{	{	{	{	{						
		{								cardiac nerves	{	{	{	{	{
			{	inferior laryngeal.	{	{	{	{	{						
		{								spinal accessory .	{	{	{	{	{
			{	Connecting branches . . .	{	{	{	{	{						
		{								branches for distribution . . .	{	{	{	{	{
{	Connecting branches .		{	{	{	{	{								
		{						branches for distribution	{	{	{	{	{		
{	To the pneumo-gastric nerve to the sympathetic to loop of atlas to gustatory nerve.		{	{	{	{	{								
		{						Descendens noni thyro-hyoid nerve to the lingual muscles and tongue.	{	{	{	{	{		

TABLE OF THE SPINAL AND SYMPATHETIC NERVES OF THE HEAD AND NECK.

SPINAL NERVES.

The cervical spinal nerves divide into	Anterior branches	The first four form the Cervical Plexus, which gives off . . .	Superficial ascending . . .	{ Small occipital nerve great auricular superficial cervical.
			superficial descending . . .	{ Supra-acromial supra-clavicular supra-sternal.
			deep internal	{ To the pneumo-gastric to the hypoglossal to the sympathetic to rectus major muscle to diaphragm nerves to descendens noni.
	posterior branches . . .	the last four and first dorsal form the Brachial Plexus, which gives off	Branches above the clavicle	{ To join the spinal accessory to the sterno-mastoid to the trapezius to the levator anguli scapulæ.
			branches below	{ The rhomboid nerve to the phrenic nerve supra-scapular nerve subclavian branch posterior thoracic, or respiratory to the scaleni muscles Are dissected with the upper limb.
		Are distributed to the muscles of the back, and give off cutaneous nerves.		

SYMPATHETIC NERVE.

The sympathetic nerve has in the neck	1. Superior cervical ganglion has . . .	Ascending branches, which unite in plexuses	Carotid plexus, which gives	{ Branch to tympanic plexus to the vidian to the sixth and fifth cranial nerves.
			cavernous plexus, which gives branches .	{ To the third cranial nerve to the fourth cranial nerve to the fifth and lenticular ganglion to the carotid artery and branches.
		external branches . . .	{ To join pneumo-gastric and hypoglossal nerves to the spinal nerves.	
		internal branches . . .	{ Pharyngeal branches superficial cardiac nerve.	
		branches to vessels .	Nervi molles.	
	2. Middle cervical ganglion	External branches . . .	To the spinal nerves.	
		internal	{ Middle cardiac nerve, to supply thyroid body and join the external laryngeal.	
		Anterior branches . .	To the subclavian artery.	
	3. Inferior cervical ganglion	External	{ To the spinal nerves forming vertebral plexus.	
		internal	Inferior cardiac nerve.	

TABLE OF THE CHIEF ARTERIES OF THE UPPER LIMB.

The subclavian is continued in the arm by	1. Axillary artery . . .	<ul style="list-style-type: none"> Superior thoracic acromial thoracic . . . { Muscular inferior acromial humeral thoracic. long thoracic alar thoracic sub-scapular { Dorsal artery muscular. } Infra-scapular. anterior circumflex posterior circumflex.
	2. brachial artery . . .	<ul style="list-style-type: none"> To coraco-brachialis superior profunda . . . { Muscular to triceps and anconeus anastomotic. nutritious inferior profunda . . . { Muscular to triceps anastomotic. anastomotic muscular.
	3. radial artery . . .	<ul style="list-style-type: none"> Recurrent muscular superficial volar posterior carpal anterior carpal metacarpal dorsal of the thumb of the index finger princeps pollicis radialis indicis deep arch. { Recurrent perforating interosseous communicating.
	4. ulnar artery . . .	<ul style="list-style-type: none"> Anterior recurrent posterior recurrent interosseous { Anterior { Nutritious muscular. muscular { posterior { Recurrent muscular. dorsal of the hand, or { Dorsal carpal metacarpal or interosseous. anterior carpal superficial arch. { Communicating four digital branches cutaneous muscular.

TABLE OF THE SPINAL NERVES OF THE UPPER LIMB.

BRACHIAL PLEXUS gives off below the clavicle .	Anterior thoracic . . .	{ Superficial deep.
	sub-scapular	{ Superior inferior long.
	circumflex.	{ Articular cutaneous to teres minor to deltoid.
	nerve of Wrisberg	
	internal cutaneous . . .	{ Small cutaneous anterior of forearm posterior of forearm.
	musculo-cutaneous . . .	{ To coraco-brachialis biceps and brachialis anticus cutaneous external of fore- arm articular to carpus.
	median	{ To pronator teres to muscles of forearm, ex- cept flexor ulnaris and part of profundus anterior interosseous cutaneous palmar to muscles of thumb in part five digital branches.
	ulnar	{ Articular to elbow to flexor carpi ulnaris to flexor profundus in part cutaneous branch of fore- arm and palm dorsal cutaneous of the hand superficial palmar division . deep palmar nerve.
	musculo-spiral	{ Internal cutaneous to triceps and anconeus external cutaneous to supinator and extensor radialis longus
		posterior interosseous . . { Muscular articular.
		radial. { Cutaneous of back of thumb and of first two fingers and half the next.

TABLE OF THE ARTERIES OF THE ABDOMEN.

The abdominal aorta gives off	1. Phrenic	Coronary . .	{ Esophageal gastric.
	2. celiac axis . .	hepatic . .	{ Superior pyloric gastro-epiploic . .
			{ Inferior pylor pancreatico- duodenal.
		{ left hepatic branch right hepatic branch.	{ Cystic.
	3. superior mesenteric*	splenic . .	{ Pancreatic vasa brevia splenic left gastro- epiploic.
			{ Pancreatic intestinal ilio-colic right colic middle colic
	4. middle capsular		
	5. renal . .		
	6. spermatic		
	7. inferior mesenteric*	{ Left colic sigmoid superior hæmor- rhoidal.	
	8. lumbar		
	9. middle sacral*	External iliac. . .	{ Epigastric. . .
			{ Pubic branch anastomotic cremasteric.
	10. Common iliac. .	{ Internal iliac. . .	{ circumflex iliac
			{ Ilio-lumbar . .
			{ lateral sacral
			{ gluteal
		{ Parietal branches	{ Coccygeal comes nervi ischiatrici muscular.
			{ sciatic
			{ pudic . . .
			{ Visceral inferior hæmorrhoidal superficial perinæal transverse perinæal artery of the bulb to corpus cavernosum dorsal artery.
		{ visceral branches . .	{ obturator . . . Articular.
			{ Middle hæmor- rhoidal vesical uterine vaginal.

* The branches marked with an asterisk are single.

TABLE OF THE VEINS OF THE ABDOMEN.

INFERIOR VENA CAVA receives	1. Common iliac . . .	Internal iliac . . .	Visceral branches . . .	Hæmorrhoi- dal plexus vesico-pro- static plexus . . . uterine vaginal.	{ Vesical dorsal of the penis deep veins of the penis.
			parietal branches . . .	Obturator pudic . . .	{ Veins of corpus cavernosum of the bulb transverse peri- næal superficial peri- næal inferior hæmor- rhoidal.
		external iliac . . .		sciatic . . .	{ coccygeal comes nervi ischiatrici muscular.
		ilio lumbar lateral sacral middle sacral into the left.	Epigastric circumflex iliac.	gluteal.	
	2. vertebro- lumbar				
	3. right spermatic				
	4. renal . . .	{ Right left	{ capsular spermatic.		
	5. right capsular				
	6. diaphrag- matic	Splenic	Splenic branches vasa brevia pancreatic left gastro- epiploic		
	7. vena portæ by means of the hepatic veins.	superior mesenteric . . .	Intestinal ileo-colic right colic middle colic right gastro- epiploic pancreatic	inferior me- senteric . . .	{ Left colic sigmoid superior hæmor- rhoidal.
		coronary cystic.			

TABLE OF THE SPINAL NERVES IN THE ABDOMEN.

LUMBAR SPINAL NERVES di- vide into .	Posterior branches	Internal . .	Muscular.
		external . .	{ Muscular cutaneous.
	Anterior branches; of these the four first end in the LUMBAR PLEXUS,* which sup- plies	Ilio-hypo- gastric . .	{ Cutaneous of the ilium hypogastric branch.
		ilio-inguinal	{ To integuments of the groin.
		external cu- taneous . .	{ To integuments of the thigh.
		genito-cru- ral	{ Genital branch crural branch.
		anterior cru- ral	{ Branches inside the pelvis . . { To the iliacus muscle. To the femoral artery.
			{ Branches outside the pelvis . . { are noticed in the thigh.
		obturator . .	Accessory . . . { Other offsets are described in the thigh.
SACRAL SPINAL NERVES di- vide into .	Posterior branches unite to- gether and give off. . . .	Muscular and cutaneous filaments.	
		Branches in- side the pelvis . .	{ Visceral to levator ani to obturator in- ternus to the pyriformis
	The anterior branches of the four superior unite with the lumbo-sacral in the SACRAL PLEXUS, and furnish	pudic	{ Inferior hæmor- rhoidal superficial peri- neal (<i>anterior and posterior</i>) muscular to the bulb dorsal of the pe- nis.
		branches outside the pelvis . .	{ inferior hæmor- rhoidal (<i>somet- imes</i>). To the superior gemellus to the inferior ge- mellus and the quadratus articular small sciatic great sciatic.
		{ these are de- scribed in the thigh.	

* The lumbo-sacral gives off the superior gluteal nerve.

TABLE OF THE SYMPATHETIC NERVE OF THE ABDOMEN.

SOLAR PLEXUS* or pre-vertebral centre of the abdomen, furnishes the following plexuses:	Diaphragmatic	Coronary plexus	
	celiac	hepatic	{ Pyloric right gastro-epiploic pancreat.-duodenal cystic.
		splenic	{ Left gastro-epiploic pancreatic.
	superior mesenteric . .	Offsets to small and large intestine.	
	suprarenal		
	renal	Spermatic plexus, filaments to the.	
	aortic	Hypogastric.	
	spermatic		
	inferior mesenteric . .	{ Offsets to the large intestine superior hæmorrhoidal.	
HYPOGASTRIC PLEXUS† ends in the pelvic plexus on each side, which gives the following plexuses:	Inferior hæmorrhoidal		
	vesical	{ Prostatic cavernous deferential to vesiculæ seminales.	
	uterine vaginal.		
GANGLIATED CORD of the sympathetic in the abdomen supplies	External branches . . .	{ To the lumbar and sacral spinal nerves.	
	internal.	{ To aortic plexus to hypogastric plexus to join round middle sacral artery. between the cords on the coccyx, in the <i>ganglion impar</i> .	
* This receives .	{ Great splanchnic nerves part of small splanchnic offset of pneumo-gastric.		This is joined above by . . { The aortic plexus filaments from the lum- bar ganglia.

PNEUMO-GASTRIC NERVE IN THE ABDOMEN.

Pneumo-gastric.	Right	{ Coronary branches to the back of the stomach filaments to join the celiac and splenic plexuses.
	left	{ Coronary branches to the front of the stomach and the hepatic plexus.

TABLE OF THE MUSCLES,

ARRANGED AFTER THE MANNER OF DR. BARCLAY, ACCORDING TO
THEIR ACTIONS.

THE HEAD IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>To either side by</i>
Platysma myoides, Sterno-mastoideus, Rectus anticus major, " " minor,	Part of trapezius, Splenius capitis, Complexus, Trachelo-mastoideus, Rectus posticus major, " " minor, Obliquus capitis superior.	Platysma myoides, Sterno-mastoideus, Part of trapezius, Splenius capitis, " colli, Trachelo-mastoideus, Complexus.
<i>Assisted (when the lower jaw is fixed) by</i>		
Mylo-hyoideus, Genio-hyoideus, Genio-hyo-glossus, Digastrici.		

THE NECK IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Laterally by</i>
Platysma myoides, Sterno-mastoideus, Digastricus, Mylo-hyoideus, Genio-hyoideus, Genio-hyo-glossus, Omo-hyoidei. Sterno-hyoidei, Thyro-hyoidei, Rectus anticus minor, Longus colli.	Part of trapezius, Rhomboides minor, Serratus posticus superior, Splenius capitis, " colli, Complexus, Trachelo-mastoideus, Transversalis colli, Inter-spinales colli, Semi-spinales colli, Rectus posticus major, " " minor, Obliquus capitis superior, " " inferior, Scaleni postici, Levator scapulae.	Various combinations of those muscles which separately move it forwards and back- wards, assisted by the sca- leni, intertransversales, and recti laterales.

THE TRUNK IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Laterally by</i>
Rectus abdominis, Pyramidalis, Obliquus externus abdominis, Obliquus internus, Psoas magnus, " parvus,	Trapezius, Rhomboides major, Latissimus dorsi, Serratus posticus superior, " " inferior, Sacro-lumbalis, Longissimus dorsi, Spinales dorsi, Semi-spinales dorsi, Multifidus spinæ, Inter-transversales dorsi et lumborum	Obliquus externus, " internus, Quadratus lumborum, Longissimus dorsi, Sacro-lumbalis, Serrati postici, Latissimus dorsi.
<i>Assisted (when the arms are carried forwards) by</i>		
Pectoralis major, " minor, Serratus magnus.		

THE SCAPULA IS MOVED

<i>Upwards by</i>	<i>Downwards by</i>	<i>Forwards by</i>	<i>Backwards by</i>
Trapezius, Levator scapulae, Rhomboidi,	Lower part of trape- zius, Latissimus dorsi, Pectoralis minor,	Pectoralis minor, Serratus magnus.	Part of trapezius, Rhomboidi, Latissimus dorsi.

THE HUMERUS IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Inwards by</i>	<i>Rotated inwards by</i>
Part of deltoid, Part of pectoralis major, <i>Assisted in some cir- cumstances by</i> Biceps, Coraco-brachialis.	Part of deltoid, Teres major, " minor, Long head of triceps, Latissimus dorsi.	Part of pectoralis major, Latissimus dorsi.	Subscapularis, <i>Assisted occasionally by</i> Pectoralis major, Latissimus and teres major. <i>Outwards by</i> Supra-spinatus, Infra-spinatus, Teres minor.

THE FOREARM IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Rotated inwards by</i>
Biceps, Brachialis anticus, Pronator teres, <i>Assisted by</i> Flexor carpi radialis, " sublimis, " ulnaris, Supinator longus.	Triceps, Anconeus.	Pronator teres, Flexor carpi radialis, Palmaris longus, Flexor sublimis, Pronator quadratus. <i>Outwards by</i> Biceps, Supinator brevis, Extensor secundi inter- nodii.

THE CARPUS IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Outwards by</i>	<i>Inwards by</i>
Flexor carpi radialis, Palmaris longus, Flexor sublimis, " carpi ulnaris, " profundus, " longus pollicis.	Extensor carpi radi- alis longior, Extensor carpi radi- alis brevior, Extensor secundi in- ternodii, Indicator, Extensor communis digitorum, Extensor proprius pollicis.	Flexor carpi radialis, Extensor carpi radi- alis longior, Extensor carpi radi- alis brevior, Extensor ossis meta- carpi, Extensor primi inter- nodii.	Flexor sublimis, " carpi ulnaris, " profundus, Extensor communis digi- torum, Extensor minimi digiti, Extensor carpi ulnaris.

THE THUMB IS MOVED

<i>Inwards and for- wards, across the palm, by</i>	<i>Outwards and back- wards by</i>	<i>Upwards and for- wards, away from the other fingers, by</i>	<i>Backwards and in- wards, to the other fingers, by</i>
Opponens pollicis, Flexor brevis, " longus.	Extensor ossis meta- carpi pollicis, Extensor primi inter- nodii, Extensor secundi in- ternodii.	Abductor, <i>Assisted by part of the</i> Flexor brevis.	Abductor, Extensor primi inter- nodii, Extensor secundi inter- nodii.

THE FINGERS ARE MOVED

<i>Forwards, or flexed, by</i>	<i>Backwards, or ex- tended, by</i>	<i>Outwards, to radial border, by</i>	<i>Inwards by</i>
Flexor sublimis, " profundus, Lumbricales, Interossei, Flexor brevis digiti minimi, Abductor digiti mi- nimi.	Extensor communis, " minimi digiti. Indicator.	Abductor indicis, " digiti minimi, Interossei.	Abductor digiti minimi, Interossei.

THE THIGH IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Inwards by</i>	<i>Outwards by</i>
Psoas magnus, Iliacus, Tensor vaginae femo- ris, Pectineus, Adductor longus, " brevis.	Gluteus maximus, Part of gluteus me- dius, Pyriformis, Obturator internus, Part of adductor mag- nus, Long head of biceps, Semi-tendinosus, Semi-membranosus.	Psoas magnus, Iliacus, Pectineus, Gracilis, Adductor longus, " brevis, " magnus, Obturator externus, Quadratus femoris.	Tensor vaginae femoris, Gluteus maximus, " medius, " minimus, Pyriformis.

THE THIGH IS ROTATED

<i>Inwards by</i>	<i>Outwards by</i>
Tensor vaginae femo- ris, Part of gluteus me- dius, <i>And, when the leg is extended, by</i> Sartorius, Semi-tendinosus.	Gluteus maximus, Part of gluteus medius, Pyriformis, Gemellus superior, Obturator internus, Gemellus inferior, Quadratus femoris, Obturator externus, Psoas magnus, Iliacus, Adductor longus, " brevis, " magnus, Biceps cruris, slightly.

THE LEG IS MOVED

<i>Backwards, or flexed, by</i>	<i>Extended by</i>
Semi-tendinosus, Biceps, Semi-membranosus, Gracilis, Sartorius, Popliteus.	Rectus, Crureus, Vastus externus, " internus.

THE FOOT IS MOVED

<i>Forwards, or flexed, by</i>	<i>Backwards, or extend- ed, by</i>	<i>Inclined inwards by</i>	<i>Outwards by</i>
Tibialis anticus, Extensor proprius pollicis, Extensor longus digi- torum, Peroneus tertius.	Gastrocnemius, Plantaris, Soleus, Flexor longus digi- torum, Flexor longus pollicis, Tibialis posticus, Peroneus longus, " brevis.	Extensor proprius pollicis, Flexor longus digi- torum, Flexor longus pollicis, Tibialis posticus.	Peroneus longus, " brevis, Extensor longus digi- torum, Peroneus tertius.

THE TOES ARE MOVED

<i>Backwards, or flexed, by</i>	<i>Forwards, or extend- ed, by</i>	<i>Inclined inwards by</i>	<i>Outwards by</i>
Abductor pollicis, Flexor brevis digi- torum, Abductor minimi di- giti, Flexor longus pollicis, " digitorum, " accessorius, Lumbricales, Flexor brevis pollicis, Adductor pollicis, Flexor brevis minimi digiti, Interossei.	Extensor longus digi- torum, Extensor proprius pollicis, Extensor brevis digi- torum.	Abductor pollicis, Interossei.	Abductor pollicis, " digiti minimi, Interossei.

INDEX.

- Abdomen, dissection of, 286.
 division into regions, 286.
 parts exposed on opening, 304.
- Abdominal muscles, 290.
 viscera, position of, 305.
- ABSORBENT GLANDS
 axillary, 204.
 bronchial, 111.
 cervical, deep, 19.
 superficial, 4.
 concatenatæ, 19.
 of the elbow, 215.
 inguinal, 288.
 internal mammary, 86.
 mediastinal, 87.
 intercostal, 111.
 parotid, 62.
 lumbar, 329.
 pelvic, 329.
 pharyngeal, 131.
 popliteal, 461.
 submaxillary, 23.
 thyroid, 17.
- Achillis, tendo, 463.
- Ampullæ of semicircular canals, 550.
- Aneurism of arch of aorta, 94.
- Annulus ovalis, 115.
- Antrum pylori, 410.
- Aorta, arch of, 89.
 abdominal, 315, 329.
 thoracic, 104.
- APONEUROSES
 of abdominal muscles, 290.
 of the scalp, 156.
 pharyngeal, 131.
 of the thigh, 428.
 temporal, 69.
 vertebral, 182.
- Appendices epiploicæ, 414.
- Appendix vermiformis, 309.
 glands in, 414.
- Arachnoid membrane of brain, 492.
- Arachnoid membrane of cord, 519.
 fluid, 493.
- Arbor vitæ in the cerebellum, 513.
 in the uterus, 395.
- Arciform fibres of medulla, 518.
- Arantii corpus, 117.
- Arm, dissection of the, 197.
- Anus, glands of, 341.
 sphincter of, 342, 414.
- ARTERIES
 alar thoracic, 212.
 anastomotica magna of arm, 221.
 thigh, 442.
 angular of the face, 61.
 aorta, abdominal, 315, 329.
 arch of, 89.
 thoracic, 104.
 articular branches of popliteal, 465.
 ascending palatine, 24, 78.
 pharyngeal, 31.
 cervical, 45.
 auditory, 495, 554.
 auricular anterior, 67.
 posterior, 31, 33.
 axillary, 206.
 basilar, 494.
 brachial, 218.
 brachio-cephalic, 91.
 bronchial, 111, 129.
 capsular, 330.
 carotid, common, 13, 92.
 external, 31.
 internal, 78, 494.
 carpal, anterior and posterior, 231,
 233, 269.
 centralis retinæ, 168.
 cerebral anterior, 494.
 middle, 494.
 posterior, 495.
 cerebellar, inferior, 495.
 superior, 495.
 cervicalis profunda, 47, 190.
 ciliary, 168.
 circle of Willis, 496.
 circumflex, anterior and posterior,
 210, 257.
 external and internal, 440.
 circumflexa ilii, 292, 333.
 celiac axis, 316.
 colica, dextra, 332.
 media, 332.
 sigmoidea, 332.
 sinistra, 322.
 comes nervi mediani, 233.

ARTERIES (*continued*)

- comes nervi phrenici, 86.
- ischiatrici, 456.
- common carotid, 13.
- iliac, 331.
- communicating anterior, 494.
- posterior, 496.
- coronaria ventriculi, 317.
- coronary of the lips, 60.
- cremasteric, 301.
- crico-thyroid, 32.
- cystic, 317.
- deep cervical, 47, 190.
- dental, inferior, 72.
- superior, 73.
- digital arteries of hand, 239.
- foot, 472.
- dorsalis hallucis, 449.
- penis, 353.
- scapulæ, 210, 259.
- linguæ, 30.
- epigastric, 301.
- ethmoidal, 168.
- external circumflex, 440.
- iliac, 333.
- facial, 24, 60.
- femoral, 436.
- frenum linguæ, 30.
- gastro-epiploica, dextra, 317.
- sinistra, 318.
- glutæal, 366, 453.
- hæmorrhoidal, superior, 321.
- middle, 365.
- inferior, 343.
- hepatic, 317, 402.
- hyoid branch of lingual, 30.
- superior thyroid, 31.
- iliac, common, 331.
- external, 333.
- internal, 365.
- ilio-lumbar, 366.
- inferior or long thoracic, 209.
- mesenteric, 320.
- profunda, 220.
- infra-orbital, 66, 73.
- innominate, 91.
- intercostal aortic, 109.
- superior, 46.
- intercosto-phrenic, 86.
- internal circumflex, 440.
- carotid, 78, 494.
- iliac, 365.
- mammary, 46, 85.
- maxillary, 71.
- pudic, 348.
- interosseous of fore-arm, 232, 235.
- anterior and posterior, 232, 235.
- dorsal, of carpus, 269.
- palmar, 248.
- ischiatric, 367, 456.
- lachrymal, 168.

ARTERIES (*continued*)

- laryngeal, superior, 31.
- lateralis nasi, 61.
- lateral sacral, 366.
- lingual, 29.
- how to tie, 26.
- long thoracic, 209.
- lumbar, 331.
- magna pollicis, 247.
- malleolar, 449.
- mammary, external, 209.
- internal, 46, 85.
- masseteric, 73.
- maxillary, external, 60.
- internal, 71.
- mediastinal, 86.
- medullary of humerus, 221.
- femur, 442.
- meningeal, anterior, 488.
- middle, 72, 488.
- posterior, 488.
- parva, 72.
- mental, 66.
- mesenteric, inferior, 321.
- superior, 320.
- metatarsal, 449.
- nasal, 73, 169.
- obturator, 366, 443.
- occipital, 31, 33, 190.
- ophthalmic, 168.
- ovarian, 387.
- palatine, ascending, 24.
- descending, 73.
- pterygo, 72.
- palmar arch, superficial, 238.
- deep, 247.
- interosseous, 248.
- palmaris profunda, 247.
- palpebral, 168.
- pancreatico-duodenales, 317.
- pancreaticæ parvæ and magna, 318.
- perforating, of hand, 248.
- profunda, 439.
- peroneal, 468.
- pharyngeal, ascending, 31, 34, 79.
- phrenic, 330.
- plantar, external, 472.
- internal, 472.
- popliteal, 465.
- posterior auricular, 31, 33.
- circumflex, 257.
- interosseous, 268.
- scapular, 45, 46, 253.
- tibial, 467.
- princeps cervicis, 33.
- profunda femoris, 439.
- perforating of, 441.
- inferior of arm, 221.
- profunda superior of arm, 220.
- pterygo-palatine, 74.
- pubic branch of epigastric, 301.
- pudic, internal, 353, 367.

ARTERIES (*continued*)

- pubic, superficial of femoral, 418.
- pulmonary, 111.
- pyloric, 317.
- radial, 229, 247, 269.
 - recurrent, 230.
- radialis indicis, 247.
- ranine, 30.
- recurrent radial, 230.
 - tibial, 448.
 - ulnar, 232.
- renal, 330.
- sacra media, 368.
- septum nasi, artery of, 61.
- short thoracic, 200.
- spermatic, 330.
 - in female, 387.
- splenic, 317.
- sterno-mastoid, 31, 33.
- stylo-mastoid, 33.
- subclavian, 38, 92.
- sublingual, 30.
- sub-mental, 24.
- sub-scapular, 209.
- superficial axillary, 202.
 - circumflexa ilii, 289, 418.
 - epigastric, 289, 418.
 - pubic, 289, 418.
- superficialis colli, 46, 251.
 - perinei, 342.
 - volæ, 230.
- superior hemorrhoidal, 321.
 - mesenteric, 316, 320.
- supra-orbital, 168.
 - scapular, 45, 259, 353.
- sural, 463, 466.
- tarsal, 449.
- temporal, superficial, 67.
 - deep, 73.
- thoracic axis of, 200.
- thoracica acromialis, 201.
 - alaris, 204.
 - humeraria, 200.
 - inferior or long, 209.
 - superior or short, 209.
- thumb, dorsal of, 269.
- thyroid axis, 44, 45.
 - superior, 31.
 - inferior, 45.
 - middle, 17.
- tibial, anterior, 448.
 - posterior, 467.
- tonsillar, 24.
- transversalis colli, 46, 253.
 - faciei, 62.
 - humeri, 253.
 - perinei, 345.
- tympanic, 33, 72.
- ulnar, 231.
 - recurrent, anterior and posterior, 232.

ARTERIES (*continued*)

- ulnaris profunda, 239, 248.
- uterine, 392.
- vaginal, 392.
- vasa brevia to stomach, 318.
- vertebral, 44, 494.
- vesical, 368.
- vidian, 73.
- Arytenoid cartilages, 143.
- Auditory nerve, 530, 551.
- Auricle, right, 114.
 - left, 118.
- Auriculo-ventricular opening, right, 116.
 - left, 119.
- Axis, cæliac, 316.
 - thoracic, 200.
 - thyroid, 44.
- Axilla, dissection of, 202.
- BACK
 - blood vessels of, 190.
 - muscles of, 182.
 - nerves of, 188, 249.
- Bands, intercolumnar, 297.
- Béclard, anecdote of, 20.
- Bell, Sir Charles, his nerves, 49.
- Bichat, fissure of, 507.
- Bile duct, 320.
- Bladder-gall, 404.
- Bladder, urinary, 360.
 - axis of, 361.
 - blood vessels of, 370, 373.
 - female, peculiarities of, 389.
 - ligaments, false, 355.
 - true, 356.
 - structure of, 370.
 - trigone of, 371, 373.
 - uvula of, 372.
- Brachial plexus, 48.
- BRAIN
 - dissection of the, 487.
 - arteries of, 494.
 - fissures of, 497.
 - general division of, 496.
 - membranes of, 488.
 - names of parts at the base, 498.
 - sinuses of, 489, 491.
 - peculiarities of circulation in, 496.
 - white and grey matter of, 499.
- Bransby Cooper, case recorded by, 39.
- Bronchus, right and left, 125.
 - structure, 126.
- Brown Séquard, researches of, 524.
- Brunn's glands, 413.
- Bulb of urethra, 381.
 - artery of, 353.
 - of vagina, 385.
- Burns, Allen, his case, 18.
- BURSÆ
 - of the biceps (arm) 279.
 - under ball of great toe, 469.
 - of the carpus. 244.

BURSÆ (*continued*)

- under deltoid, 256.
- between fingers, 270.
- under glutæus maximus, 452.
- housemaids, 436.
- over the knuckles, 266.
- under latissimus dorsi, 250.
- over patella, 436.
- under ligamentum patellæ, 436.
- over pomum Adami, 142.
- under sartorius, 430.
- under semimembranosus, 459.
- under subscapularis, 259.
- between toes, 474.

Calamus scriptorius, 511.

Camper, his knowledge of the urethra, 377.

CANALS, alimentary, 307.

- of Fontana, 533.
- inguinal, 295.
- of Petit, 543.
- semicircular, 550.

Caput coli, 303.

gallinaginis, 374.

Cardiac plexus of nerves, 111.

Carotid, common, 13.

- external, 31.
- internal, 73.

Caruncula lacrymalis, 56.

Cauda equina, 520, 523.

Cavernous sinus, 160.

Central line of the neck, 19.

Cerebellum, 502.

crura of, 514.

Cerebrum, 497.

- crura of, 498.
- structure of crura, 527.

Cervical fascia, 5.

- ganglia of sympathetic, 82.
- plexus, 34.

superficial branches of, 4.

Chambers of the eye, 542.

Chest, dissection of, 83.

Chordæ vocales, 144.

tendinæ, 116.

Choroid, coat of the eye, 535.

plexus, 504, 507.

Ciliary arteries, 163.

body, 541.

ligament, 535, 537.

muscle, 53.

processes, 541.

Clitoris, 383.

crura of, 383.

Cochlea, 550.

Cæcum, 303.

Cæcal (ileo) valve, 308.

Colon, commencement of, 308.

ascending, 309.

descending, 309.

transverse, 309.

Colon, sigmoid flexure of, 309.

structure of, 414.

Columnæ carnæ, 116.

Commissures of the cerebrum, 500, 509.

of the cerebellum, 514.

Coni vasculosi testis, 558.

Conjunctiva oculi, 531.

Constrictors of the pharynx, 131.

Cord spermatic, 559.

spinal, 518, 520.

Cornea, 534.

Cornicula laryngis, 144.

Corona glandis, plexus of veins round, 379, 381.

Corpora geniculata, 508.

Corpus Arantii, 117.

callosum, 500.

cavernosum, 379.

Highmori, 557.

mammillaria, 499.

luteum, 399.

rhomboideum, 514.

spongiosum, 381.

striatum, 505.

Cowper's glands, 352, 376.

in the female, 386.

Crico-thyroid membrane, 20.

Cricoid cartilage, 143.

Crus penis, 379.

artery of, 335.

clitoridis, 383.

Crura cerebri, 498.

cerebelli, 514.

Crural arch, 290, 421.

Crystalline lens, 543.

Dartos, tunica, 555.

Deglutition, mechanism of, 140.

Digastric triangle, 21.

DUCTS

arteriosus, 122.

communis choledochus, 320, 404.

ejaculatorius, 362.

cystic, 320, 404.

of Cowper's glands, 376.

hepatic, 320, 403.

lachrymal, 164, 179.

nasal, 179.

pancreatic, 322.

parotid, 58, 63.

prostatic, 374.

of Rivinus, 29.

seminal, 374.

of Steno, 63.

sublingual, 29.

submaxillary, 23.

thoracic, 104.

venosus, 122.

of Wharton, 23.

Duodenum, course of, 307.

Dura mater of brain, 488.

uses of, 488.

- Dura mater of brain, arteries of, 488.
sinuses of, 489.
partitions formed
by, 488.
of spinal cord, 519.
- Duverney's glands, 386.
- Ear, dissection of the, 545.
internal, 649.
muscles of the, 157.
- Endocardium, 116.
- Endolymph, 552.
- Epididymis, 556.
- Epiglottis, 144.
ligaments of, 144.
- Eustachian tube, 131, 139, 548.
valve of heart, 115.
- Eye, dissection of, 531.
- Eyelids, structure of the, 54.
- Face, dissection of, 50.
nerves of, 50, 66.
- Fallopian tubes, 397.
- Falx cerebri, 488.
cerebelli, 489.
- FASCIA
axillary, 202.
buccal, 59.
cervical, 5.
costo-coracoid, 200.
cribiform, 419, 421.
of forearm, 226, 262.
iliac, 327.
infundibuliform, 302.
of groin, 288.
lata of thigh, 428.
of the leg, 445.
lumborum, 183, 251.
mammary, 553.
orbital, 162.
parotid, 62.
palmar, 236.
perineal, superficial, 343.
deep, 347.
pharyngeal, 131.
pelvic, 356.
in female, 388.
plantar, 469.
popliteal, 457.
propria of a femoral hernia, 425.
spermatic, 297.
superficial, of abdomen, 286.
thigh, 416.
perineum, 343.
temporal, 69.
transversalis, 299.
of upper arm, 215.
- FIBRO-CARTILAGES
of the ear, 545.
of the eyelids, 54.
intervertebral, 192.
of the lower jaw, 196.
- FIBRO-CARTILAGES (*continued*)
of the knee, 481.
of the nose, 177.
between radius and ulna, 280.
between sternum and clavicle, 272.
tarsal, 54.
of trachea, 126.
- Flexor tendons in the hand, 242.
sheaths of ditto, 242.
synovial lining of, 243.
- Floor of the mouth, how formed, 25.
- Fœtal circulation, 123.
- Fontana's canal, 538.
- FORAMINA
ovale of heart, 115.
tympanum, 545.
rotundum, 545.
of Sœmmering, 540.
Winslow, 312.
Monro, 507.
Thebesius, 115.
- Forearm, back of the, 261.
front of the, 225.
- Fornix, 506.
- Fossa ischio-rectal, 341, 354.
- Frenum linguæ, artery of, 30.
- Frenum preputii, 379.
- Galen, veins of, 508.
- Gall bladder, 306, 404.
- GANGLIA
cervical of sympathetic, 82.
dorsal ditto, 108.
lumbar ditto, 334.
pelvic ditto, 370.
of Andresch, 174.
Arnold's, 81, 173.
cardiac, 111.
Cloquet's, 173.
impar, 370.
jugular, 174.
lenticular, 169.
Gasserian, 530.
Mackel's, 172.
olfactory, 529.
otic, 172.
petrous, 174.
semilunar, 108, 316.
spheno-palatine, 170, 172.
submaxillary, 29.
of Ribes, 82.
- Gastric juice, apparatus for secretion
of, 411.
- Gimbernat's ligament, 296, 422.
- GLANDS
anal, 341.
axillary, 204.
of Brunn, 413.
beneath the apex of tongue, 154.
concatenatæ, 19.
Cowper's, 352, 371, 376.
in female, 386.

GLANDS (*continued*)

- Duverney's, 386.
- epiglottidean, 145.
- femoral, 416.
- inguinal, 288.
- iliac, 328.
- labial, 51.
- lacrimal, 164.
- of Lieberkühn, 413.
- lingual, 154.
- lumbar, 328.
- mammary, 554.
- Meibomian, 56.
- mesenteric, 320.
- molar, 60.
- Naboth's, 395.
- palatine, 139.
- parotid, 62.
- pancreas, 322.
- of Pacchioni, 491.
- pineal, 508, 509.
- pituitary, 499.
- Peyer's, 413.
- popliteal, 461.
- prostate, 373.
- solitary, 414.
- sublingual, 28.
- submaxillary, 23.
- thyroid, 16.
- tracheal, 127.
- Glans penis, 379.
- Glisson's capsule (liver), 401.
- Glottis, 146.
- Gluteal region, dissection of, 451.
- cutaneous nerves of, 451.
- Graafian vesicles, 398.
- Granular lids, 56.
- Gubernaculum testis, 559.

Hamstring muscles, 457.

HEART

- dissection of, 113.
- arteries of, 120.
- right auricle, 114.
- left auricle, 118.
- blood, how supplied with, 120.
- foetal, 122.
- muscular fibres, how arranged, 111, 121.
- nerves of, 111.
- position and form of, 96.
- rings of, 120.
- thickness of its cavities, 121.
- valves, position of, 95.
- vessels, how attached to, 120.
- right ventricle of, 115.
- left ditto, 119.
- weight, 121.

Helicotrema, 550.

HERNIA

- inguinal parts concerned in, 294.
- coverings of, 302.

HERNIA (*continued*)

- inguinal seat of structure, 303.
- changes produced by age in, 303.
- femoral, parts concerned in, 421.
- covering of, 426.
- seat of stricture, 427.
- Herophili, torcular, 492.
- Hey's ligament, 419.
- Hippocampus major, 504.
- minor, 504.
- Horner's muscle, 54.
- Houston, his folds in the rectum, 416.
- Humour, aqueous, 533, 542.
- vitreous, 533, 542.
- Hunter's canal, 438.
- Hyaloid membrane, 542.
- Hymen, 385.

JEJUNUM

- intestinum, 308.
- glands of, 413.
- commencement of, 307.
- peculiarities of, 412.
- Ileo-cæcal valve, 415.
- Ileum intestinum, 308.
- glands of, 413.
- termination of, 309.
- Infundibulum, 499.
- Inguinal canal, 295.
- boundaries of, 301.
- rings, 295, 299.
- Intercostal muscles, 109.
- nerves, 110, 198.
- Intervertebral fibro-cartilages, 192.
- Intestine, small, 307, 411.
- large, 308, 414.
- Intestinal canal, course of, 307.
- length of, 110.
- structure of, 409.
- Intestine, large, characters of, 414.
- course of, 308.
- Iris, 638.
- Ischio-rectal fossa, 341, 354.
- Isthmus faucium, 135.
- glandulæ thyroideæ, 16.
- Iter a tertio ad quartum ventriculum, 509, 511.

JOINTS

- acromio-clavicular, 274.
- ankle, 483.
- carpo-metacarpal, 283.
- crico-thyroid, 151.
- elbow, 277.
- fingers, 285.
- hip, 476.
- inter-carpal, 282.
- knee, 479.
- lower jaw, 196.
- of the metatarsus, 486.
- occipital, 194.
- of pelvis, 475.
- of the ribs, 194.

JOINTS (*continued*)

- shoulder, 275.
- spine, 191.
- between spine and head, 194.
- sterno-clavicular, 272.
- symphysis, pubic, 475.
- sacro-iliac, 475.
- of the tarsus, 485.
- of the thumb, 283.
- tibio-fibular, 483.
- toes, 486.
- wrist, 279, 281.

- Kidneys, position of, 307, 323.
- relations of, 307.
- structure of, 407.

- Labia majora, 382.
- minora, 382.

- Labyrinth, osseous, 549.
- membranous, 551.

- Lachrymal gland, 164.
- duct, 179.

- Lachrymal sac, how to puncture, 52.

- Lacteal vessels, 320.

- Lacunæ of the urethra, 378.

- Lamina cinerea, 499.
- spiral, 550, 551.

- Laryngotomy, observations on, 20, 21.

LARYNX

- dissection of, 141.
- blood-vessels of, 150.
- cartilages of, 142.
- chordæ vocales of, 144.
- muscles of, 147.
- mucous membrane of, 145.
- nerves of, 150.
- sexual differences, 150.
- superior opening of, 146.
- ventricles of, 146.

- Leg, front of the, 444.

- Lens, crystalline, 543.

- capsule of, 544.

- Lenticular ganglion, 169.

LIGAMENTS

- accessory of humerus, 276.
- of hip, 476.
- acromio-clavicular, 274.
- annular, anterior, of wrist, 240.
- posterior ditto, 262.
- annular of ankle, 445.
- of radius, 278.
- arcuata, 324.
- of the atlas and axis, 194.
- of the bladder, 355, 356.
- broad, in the female, 387.
- calcaneo-cuboid, 486.
- scaphoid, 486.
- ciliary, 535.
- conoid, 275, 537.
- coraco-acromial, 274.
- coraco-clavicular, 275.

LIGAMENTS (*continued*)

- coronary of liver, 306.
- knee-joint, 482.
- coraco-humeral, 276.
- costo-clavicular, 275.
- costo-transverse, 195.
- cotyloid of the hip, 477.
- crucial, 481.
- of the fingers, 285.
- falciform of liver, 305.
- Gimbernats', 296, 420.
- glenoid of shoulder-joint, 277.
- fingers, 243.
- dentata, of cord, 520.
- external lateral, of ankle, 484.
- elbow, 277.
- jaw, 196.
- knee, 480.
- wrist, 280.
- Glosso-epiglottidean, 151.
- Hey's, 419.
- hyo-epiglottidean, 145.
- ilio-lumbar, 475.
- interclavicular, 273.
- internal lateral, of elbow, 277.
- ankle, 484.
- jaw, 77, 196.
- knee, 480.
- wrist, 289.
- interosseous, of forearm, 279.
- leg, 483.
- carpus and metacarpus, 282.
- tarsus and metatarsus, 485.
- the pelvis, 476.
- intervertebral fibro-cartilages, 192.
- lateral, of liver, 306.
- nuchæ, 249.
- oblique of radius, 279.
- of occiput, 194.
- odontoid or check, 194.
- orbicular, of radius, 278.
- posticum Winslowii, 458, 480.
- pterygo-maxillary, 58.
- of the ovary, 398.
- Poupart's, 290.
- patellæ, 435.
- of the pubes, 475.
- of the ribs, 194, 195.
- round of liver, 305.
- round of uterus, 387.
- sacro-iliac, 476.
- of spine, 191.
- stellate, 195.
- sterno-clavicular, 272.
- stylo-maxillary, 25.
- subflava, 192.
- suspensory, of liver, 305.
- transverse, of fingers, 237.
- teres, of the hip, 478.
- thyro-epiglottidean, 142.
- thyro-hyoid, 142.
- sacro ischiatic, greater, 475.

LIGAMENTS (*continued*)

- sacro ischiatic, lesser, 475.
- supensorium penis, 379.
- tibio-fibular, 483.
- transverse metacarpal, 236, 270.
- triangular of urethra, 347.
- trapezoid, 275.
- Linea alba, 293.
- semilunaris, 291.
- Lineæ transversæ, 292, 501.
- Liver, position of, 305.
- dissection of, 399.
- Locus niger, 525, 527.
- perforatus anticus, 499.
- medius, 498.
- LUNGS, 124.
- absorbents of, 130.
- lobules of, 127.
- nerves of, 130.
- position and form of, 99.
- root of each, 113, 124.
- structure of, 124.
- Lunula of aortic valves, 118.
- Malpighian bodies in spleen, 406.
- kidney, 408.
- Mamma, dissection of the, 554.
- Mastication, muscles of, 67.
- Meatus auditorius, 546.
- urinarius, in female, 383.
- Mediastina, 86.
- anterior, 87.
- posterior, 84, 103.
- Mediastinum, testis, 557.
- Medulla oblongata, 514.
- minute structure of, 524, 526.
- Meibomian glands, 56.
- Membrana pupillaris, 539.
- Jacobi, 541.
- hyaloidea, 542.
- tympani, 547.
- Membranous labyrinth, 551.
- Membranes of the brain, 488.
- cord, 519.
- MENINGEAL arteries
- anterior, 488.
- middle, 488.
- posterior, 488.
- parva, 71.
- Mesentery, 311.
- Meso-colon, transverse, 310.
- Meso-cæcum, 308.
- Mesorchium, 559.
- Mitral valve, 95, 119.
- Modiolus, 551.
- Mons Veneris, 382.
- Monro, foramen of, 507.
- Morgagni, sinuses of, 133.
- liquor of, 544.
- MUSCLES
- abductor pollicis manus, 245.
- pedis, 470.

MUSCLES (*continued*)

- abductor minimi digiti manus, 246.
- pedis, 470.
- accelerator urinæ, 345.
- accessorius pedis, 471.
- accessorius ad sacrolumbalem, 184.
- adductor pollicis manus, 246.
- pedis, 473.
- adductors of thigh, 432.
- brevis, 433.
- longus, 432.
- magnus, 433.
- anconeus, 266.
- arytenoideus, 148.
- aryteno-epiglottideus, 149.
- of ball of thumb, 245.
- of ball of little finger, 246.
- attollens aurem, 157.
- attrahens aurem, 157.
- azygos uvulæ, 137.
- biceps, flexor cubiti, 216.
- cruris, 457.
- brachialis anticus, 218.
- buccinator, 58.
- cervicalis ascendens, 184.
- ciliary of the eye, 534.
- eyelid, 53.
- circumflexus or tensor palati, 137.
- coccygeus, 365.
- cochlearis, 551.
- complexus, 186.
- compressor nasi, 57.
- urethræ, 351.
- in the female, 384.
- constrictors of pharynx, 131.
- coraco brachialis, 217.
- corrugator supercilii, 54.
- cremaster, 297.
- crico-arytenoideus lateralis, 148.
- posticus, 147.
- thyroid, 147.
- cruræus, 434.
- sub-cruræus, 435.
- DEPRESSOR anguli oris, 51.
- labii inferioris, 51.
- superioris, 58.
- deltoid, 255.
- diaphragm, 323.
- erector penis, 346.
- clitoridis, 383.
- erector spinæ, 183.
- digastricus, 21.
- dilatator narium, 58.
- EXTENSOR brevis digitorum pedis, 447.
- carpi radialis longior, 263.
- brevior, 264.
- carpi ulnaris, 265.
- communis digitorum manus, 264.
- digiti minimi, 265.
- indicis, 268.

MUSCLES (*continued*)

- Extensor communis digitorum pedis, 446.
 ossis metacarpi pollicis, 267.
 primi internodii, 267.
 secundi internodii, 267.
 proprius pollicis pedis, 447.
 external oblique, 290.
 FLEXOR brevis minimi digiti manus, 246.
 minimi digiti pedis, 473.
 pollicis manus, 246.
 digitorum pedis, 473.
 pollicis pedis, 471.
 carpi radialis, 227.
 ulnaris, 228.
 longus digitorum pedis, 466.
 longus pollicis manus, 235.
 pedis, 467.
 ossis metacarpi pollicis, 245.
 profundus digitorum manus, 234.
 sublimis digitorum manus, 228.
 gastrocnemius, 462.
 gemelli, 454.
 genio-hyoid, 26.
 genoio-hyo-glossus, 27.
 glutæus maximus, 451.
 medius, 452.
 minimus, 452.
 gracilis, 431.
 hamstrings, 457.
 iliacus internus, 327.
 hyo-glossus, 26.
 indicator, 268.
 infra-spinatus, 258.
 intercostal, 109.
 internal oblique, 291.
 interosseous of hand, 270.
 foot, 474.
 interspinales, 187.
 intertransversales, 187.
 latissimus dorsi, 205, 250.
 laxator tympani, 548.
 LEVATOR
 anguli oris, 57.
 anguli scapulæ, 252.
 ani, 364.
 menti, 52.
 glandulæ thyroideæ, 17.
 labii superioris et alæ nasi, 57.
 labii superioris, 57.
 palati, 137.
 uvulæ, 137.
 levator palpebræ, 54, 165.
 levatores costarum, 187.
 longissimus dorsi, 184.
 longus colli, 190.
 lumbricales of the hand, 245.
 foot 471.
 masseter, 68.

MUSCLES (*continued*)

- multifidus spinæ, 186.
 mylo-hyoideus, 25.
 obliquus externus, 290.
 internus, 291.
 obliquus inferior capitis, 188.
 superior capitis, 188.
 obliquus inferior oculi, 170.
 superior oculi, 165.
 obturator internus, 454.
 externus, 454.
 occipito-frontalis, 156.
 omo-hyoid, 11, 253.
 opponens digiti minimi, 247.
 opponens pollicis, 245.
 of the pinna of the ear, 546.
 orbicularis, oculi, 52.
 oris, 51.
 palato glossus, 138.
 pharyngeus, 138.
 palmaris brevis, 236, 237.
 longus, 227.
 papillares, 116.
 pectinati auriculæ, 114.
 pectineus, 432.
 pectoralis major, 198.
 minor, 205.
 peroneus brevis, 450.
 longus, 449.
 tertius, 447.
 plantaris, 463.
 platysma myoides, 1.
 popliteus, 466.
 pronator quadratus, 235.
 præ-vertebral, 190.
 pronator radii teres, 227.
 psoas magnus, 326.
 parvus, 327.
 pterygoid, 70, 71.
 pyramidalis nasi, 54.
 abdominis, 293.
 pyriformis, 453.
 quadratus lumborum, 328.
 sheath of, 328.
 femoris, 454.
 rectus abdominis, 292.
 sheath of, how formed, 292.
 capitis anticus major and minor, 191.
 lateralis, 188.
 posticus, major and minor, 187.
 femoris, 434.
 recti muscles of the eye, 165.
 rhomboid, major and minor, 252.
 risorius, 50.
 rotatores spinæ, 187.
 sacro-lumbalis, 184.
 sartorius, 430.
 scalenus anticus, 37.
 medius, 37.

MUSCLES (*continued*)

- scalenus posticus, 37.
- semimembranosus, 458.
- semispinalis colli, 186.
 - dorsi, 186.
- semitendinosus, 458.
- serratus magnus, 206, 254.
 - posticus inferior, 182.
 - superior, 182.
- soleus, 463.
- sphincter ani, external, 342.
 - internal, 414.
- spinalis dorsi, 185.
- splenius capitis, 183.
 - colli, 183.
- stapedius, 545.
- sterno-cleido mastoideus, 7.
 - thyroid, 10.
- stylo-glossus, 77.
 - hyoid, 22.
 - pharyngeus, 77.
- sub-anconeus, 260.
- sub-clavius, 201.
- sub-cruræus, 435.
- sub-scapularis, 206, 258.
- supinator radii longus, 229.
 - brevis, 266.
- supra spinatus, 258.
- temporal, 69.
- tensor fasciæ femoris, 433.
- tensor tarsi, 53.
 - tympani, 548.
 - palati, 137.
- teres major, 206, 258.
 - minor, 258.
- thyro-arytenoideus, 148.
 - hyoid, 12.
- tibialis anticus, 446.
 - posticus, 467.
- trachelo-mastoid, 185.
- transversalis abdominis, 292.
 - colli, 184.
 - pedis, 474.
- transverso-spinalis, 186.
- transversus perinei, 347.
- trapezius, 249.
- triangularis sterni, 85.
- triceps extensor cubiti, 224, 260.
 - femoris, 434.
- vestus internus, 434.
 - externus, 434.
- zygomatichi, 52.

Naboth, glands of, 395.

ovula of, 396.

Nasal duct, 179.

Nates and testes, 508.

NERVES

- accessory, 9, 251.
- anterior crural, 336, 442.
 - dental, 171.
 - thoracic, 49, 201.

NERVES (*continued*)

- auditory, 530, 552.
- auricular of Arnold, 176.
- auriculo-parotidean, 4.
- axillary plexus of, 208.
- brachial plexus of, 48, 208.
- buccal, 76.
- cardiac branches of pneumogas-
tric, 81, 107.
- cerebral, origins of, 528.
- cervical branch of facial, 5.
 - plexus of, 34.
- cervico-facial, 65.
- chorda tympani, 177, 649.
- ciliary, long, 167.
 - short, 169.
- coccygeal, 190.
- circumflex, 211, 257.
- cranial, exit of, 159.
- descendens noni, 15.
- dental anterior, 171.
 - inferior, 76.
 - posterior, 171.
- eighth pair at base of skull, 174, 175, 531.
- external cutaneous, of arm, 223.
 - leg, 444.
 - musculo - spi-
ral, 213.
 - thigh, 336, 428.
- external laryngeal, 33.
 - respiratory, of Bell, 49, 206.
 - saphenous, 462.
- fifth pair, 530.
- fourth pair, 164, 530.
- frontal, 163.
- genito-crural, 336.
- glosso-pharyngeal, 78, 174.
- glutæal superior, 369, 453.
- great ischiatic, 369, 450.
 - occipital, 158, 249.
 - petrosal, 177.
- gustatory, 29, 76.
- hypo-gastric plexus, 334.
- hypo-glossal, 28, 531.
- ilio-hypogastric, 289.
 - inguinal, 289.
- inferior dental, 76.
 - laryngeal, 81, 107.
 - maxillary, 74.
- infra-orbital of the fifth, 66, 172.
 - portio dura, 65.
- trochlear, 167.
- intercostal, 110, 198.
 - costo-humeral, 203.
- internal cutaneous of arm, 213.
 - lesser ditto, ditto, 213.
 - cutaneous of thigh, 428, 437.
- interosseous anterior of forearm, 234, 236.
 - posterior of ditto, 268.

NERVES (*continued*)

- ischiatric, great, 454.
- lesser, 456.
- Jacobson's, 175.
- lachrymal, 164.
- lesser petrosal, 173.
- ischiatric, 369, 457.
- long saphenous, 437.
- pudendal, 342.
- lumbar plexus, 334.
- lumbo-sacral, 368.
- median, 222, 233, 241.
- malar, 65, 164.
- masseteric, 75.
- mental, 66.
- middle cutaneous of thigh, 428.
- molles, 61.
- musculo-spiral or radial, 224.
- mylo-hyoidean, 76.
- maxillary inferior, 74.
- superior, 170.
- nasal, of Meckel's ganglion, 172.
- the ophthalmic, 167.
- naso-lobular, 50.
- naso-palatine, 172.
- neck cutaneous of, 4.
- ninth pair, 531.
- obturator, 336, 443.
- occipitalis minor, 4.
- major, 158, 249.
- olfactory, 181, 528.
- optic, 168, 529.
- ophthalmic branch of the fifth, 162.
- orbital branch of superior maxillary, 170.
- palatine, 171, 172.
- palmar branch of median, 234, 241.
- peroneal, 450, 460.
- petrosal, greater, 177.
- lesser, 177.
- phrenic, 38, 93.
- difference between right and left, 94.
- plantar external, 473.
- plantar internal, 473.
- pneumogastric in the neck, 80.
- chest, 106.
- at base of skull, 176.
- popliteal, 460.
- portio dura, 64, 176, 530.
- mollis, 530, 549.
- posterior auricular, 34, 65.
- dental, 171.
- interosseous, 268.
- saphenous, 462.
- tibial, 468.
- pterygoid, 75.
- pubic, 354, 369, 456.
- pulmonary, 107.
- radial of musculo-spiral, 224.
- below the elbow, 231.
- recurrent, 81, 107.

NERVES (*continued*)

- rhomboid, 49.
- sacral plexus, 368.
- seventh pair, 530.
- sixth pair, 169, 530.
- spermatic, 559.
- spinal, 110, 522.
- splanchnic, 108.
- subscapular, 210.
- superficialis colli, 4.
- perinei, 342.
- superior laryngeal, 32, 81.
- maxillary, 170.
- supra-clavicular, 198.
- orbital, 66, 157.
- scapular, 49, 253.
- trochlear, 164.
- sympathetic system of, 81.
- in the neck, 82.
- chest, 108.
- abdomen, 334.
- in the pelvis, 370.
- temporal, to muscle, 75.
- superficial, 158.
- temporo-auricular, 67, 76.
- temporo-facial, 64.
- malar, 170.
- third pair, 169, 530.
- tibial anterior, 448.
- tympanic, 175.
- ulnar, 223, 233, 240, 248.
- vidian, 173, 177.
- of Wrisberg, 213.
- Nipple, 554.
- Nose
- dissection of, 177.
- blood-vessels of, 181.
- cartilages of, 177.
- meatus of, 178.
- mucous or Schneiderian membrane, 180.
- posterior openings of, 134.
- veins of, 180, 181.
- Nuchæ ligamentum, 249.
- Nymphæ, 383.
- Œsophagus, 105.
- Œsophageal arteries, 106, 111.
- Olives, or olivary bodies, 515, 516.
- Olfactory nerves, 181, 528.
- OMENTUM
- gastro-hepatic, 312.
- gastro-splenic, 312.
- great, 311.
- lesser, 312.
- Optic nerves, 529.
- ORBIT
- dissection of, 162.
- muscles of, 163.
- nerves of, 163.
- periosteum of, 162.
- Os-hyoides, 141.

- Os-tinca, 391.
 Otic-ganglion, 173.
 Otolithes, 551.
 Ovaries, 398.
 Oviducts, 393.

 Pacchioni, glands of, 490.
 Pacini, corpuscles of, 242.
 Palm of the hand, 236.
 PALATE
 hard, 140.
 dissection of, 136.
 arches of, 136.
 glands of, 139.
 muscles of, 137.
 Palpebral arteries, 168.
 Panizza, mercurial injections of, 378.
 Parotid gland, 62.
 relations of, 62.
 Paralysis, cross explanation of, 513.
 Pancreas, 307, 322.
 Patella, structures covering the, 436
 bursa over, 436.
 Pelvic viscera, side view of, 354.
 in the female, 387.
 Pelvis, purposes served by, 336.
 contents of male, 337.
 female, 386.
 Penis, 378.
 corpus cavernosum, 379.
 spongiosum, 381.
 glans, 379.
 Pericardium, 98.
 objects exposed when laid open, 96.
 Peri-lymph of the ear, 552.
 PERINEUM
 male, 340.
 boundaries of, 340.
 cutaneous vessels and nerves of,
 342, 345.
 tendinous centre of, 344.
 triangle of, 341.
 PERITONEUM, 310.
 course of, in male, 312.
 in female, 387.
 lesser cavity of, 315.
 course in the female pelvis, 387.
 Pes anserinus, 63.
 Petit, canal of, 543.
 Peyer's glands, 413.
 Pharyngeal plexus, 78.
 Pharynx, dissection of, 130.
 aponeurosis of, 133.
 muscles of, 131.
 Phrenic nerve, 38.
 Pigmentum nigrum, 536, 537.
 Pinna of the ear, 545.
 Pia mater, 493.
 of cord, 520.
 Platysma myoides, 1.
 Pleura, 101.
 reflections of, 101.

 PLEXUS OF NERVES
 axillary, 208.
 brachial, 208.
 cardiac, 111.
 cavernous or carotid, 161.
 cervical, 34.
 choroid, 507.
 coronary, 112.
 hæmorrhoidal, 334, 368.
 hypogastric, 334.
 lumbar, 334.
 œsophageal, 107.
 pharyngeal, 78.
 prostatic, 370.
 pulmonary, 107.
 sacral, 368.
 solar, 316.
 uterine, 334.
 vaginal, 334.
 vesical, 334, 370.
 Pomum Adami, 142.
 Popliteal space, 460.
 Portio dura nerve, 64, 176.
 Portæ vena, 319.
 Pneumo-gastric nerve, 80.
 Pons Varolii, 498, 514.
 structure of, 524, 526, 527.
 Poupert's ligament, 290.
 Præcordial region, 96.
 Prepuce, 378.
 frenum preputii, 379.
 clitoridis, 383.
 Processus a cerebello ad testes, 508, 511.
 Promontory of the tympanum, 549.
 Prostate gland, 363, 373.
 structure of, 375.
 veins about, 358.
 Pudenda, female, 382.
 Pulsation, real and apparent, of a tu-
 mor, 19.
 Puncta lachrymalia, 55.
 Pylorus, 410.
 Pyramid in the ear, 549.
 Pyramids, anterior, 513, 515.
 posterior, 515.

 Ranula, nature of, 23.
 Receptaculum chyli, 105.
 Recto-vesical pouch, 338.
 Rectum, 338.
 relations of, 359.
 structure of, 414.
 fold of mucous membrane in, 415.
 Reil, island of, 498.
 fillet of, 526.
 Renal capsule, position of, 307.
 structure of, 408.
 Restiform tracts, or columns, 512, 515,
 516.
 Rete testis, 558.
 mirabile of Galen, 496.
 Retina, 540.

- Rima glottidis, 146.
 Ring, external abdominal, 295.
 internal abdominal, 299.
 femoral, 423.
 fibrous, of the heart, 120.
 Risorius, musculus, 50.
 Rule for finding lingual artery, 26.

 Saccule of the labyrinth, 552.
 Santorini's, muscle, 50.
 Saphenous opening of fascia lata, 419.
 SCALP
 dissection of, 156.
 absorbents of, 158.
 blood-vessels of, 156.
 nerves of, 157.
 surgical interest of, 158.
 Scarpa's triangle, 430.
 Sclerotic coat of the eye, 533.
 Semicircular canals, 550.
 Septum lucidum, 505.
 pectiniforme, 380.
 Scrotum, structure of, 554.
 septum of, 555.
 Sinuses of the brain, 489.
 Morgagni (pharyngeal), 133.
 Valsalva, 118.
 Sheath of femoral vessels, 424.
 Shoulder, dissection of, 255.
 Socrates, his description of the eyelids, 57.
 Sœmmering, foramen of, 540.
 Spheno-maxillary region, 67, 69.
 Spheno-palatine ganglion, 172.
 Sphincter ani, internal, 414.
 external, 342.
 Spinal cord, 518, 520.
 nerves, 322.
 Spine, ligaments of, 191.
 Spleen, position of, 306.
 structure of, 405.
 Steno's duct, 63.
 STOMACH, 306.
 blood-vessels of, 318.
 structure of, 410.
 Stylo-maxillary ligament, 25.
 Sublingual salivary gland, 28.
 Submaxillary ganglion, 29.
 gland, 23.
 region, 21.
 Supra-clavicular region, 35.
 Sympathetic nerve, 81.
 in the neck, 82.
 chest, 107.
 abdomen, 334.
 pelvis, 370.
 Symphysis pubis, 475.
 sacro-iliac, 475.

 Tænia semicircularis, 504, 505.
 Hippocampi, 507.
 Tarsal cartilages, 54.

 Tarsal ligaments, broad, 55.
 Temporal region, 67.
 Tendo-Achillis, 462, 463.
 oculi, 52.
 Tentorium cerebelli, 489.
 Testicle, 555.
 coverings of, 556.
 descent of, 559.
 Thalamus opticus, 508.
 Thigh, dissection of, 416.
 back of the, 457.
 Thoracic duct, 105.
 Thyroid, axis of arteries, 44, 45.
 body, 16.
 cartilage, 142.
 Tongue, 151.
 Tonsils, 138.
 Torcular Herophili, 492.
 Trachea, 125.
 Tracheotomy, observations on, 20.
 Transverse meso-colon, 311.
 TRIANGLES of the neck, 8.
 posterior of the neck, 35.
 at the elbow, 226.
 of the perineum, 345.
 of Scarpa, 430.
 Triangular ligament of urethra, 347.
 Tricuspid valve, 116.
 Trigone of bladder, 371, 373.
 Tubes Fallopian, 397.
 Tunica albugenia, 398, 557.
 vaginalis scroti, 556.
 vasculosa scroti, 557.
 Ruyschiana, 536.
 Tuber annulare 514.
 cinereum, 499.
 Tubercula quadrigemina, 510.
 Tympanum of the ear, 547.
 membrane of the, 547.
 muscles of the, 548.
 Tympanic artery, 72.
 bones, 548.

 Umbilical artery, remains of, 298.
 Upper arm, dissection of, 212.
 Uterus, position of, 390.
 appendages of, 387.
 broad ligaments of, 387.
 round ligaments of, 387.
 blood-vessels of, 392.
 nerves of, 392.
 absorbents of, 392.
 structure of, 393.
 Urachus, 299.
 Ureter, 362, 373.
 Urethra, 376.
 in the female, 383, 389.
 Utriculus, 374.
 Uvea, 533.
 Uvula, 136.
 cerebelli, 513.
 vesicæ, 372.

Vagina, 385, 389.

bulb of, 385.

structure of, 393.

Valsalva, sinuses of, 118.

VALVES

of coronary vein, 115.

Eustachian, 115.

ileo-cæcal, 415.

mitral, 95, 119.

pulmonary, 95, 117.

semilunar of aorta, 95, 119.

of Tarini, 513.

of Thebesius, 115.

tricuspid, 95, 112, 116.

of Vieussens, 508.

Valvulae conniventes, 412.

Vas deferens, 362.

Vasa efferentia testis, 558.

recta testis, 558.

vorticosa oculi, 536.

VEINS

anterior jugular, 3.

auditory, 549.

axillary, 207, 212.

azygos major, 104.

minor, 105

basilic, 213.

brachial, 222.

brachio-cephalic, 88.

bronchial, right and left, 129, 130.

capsular, 329.

cava superior, 89.

inferior, 329.

cephalic, 201.

choroid of the eye, 536.

coronary, 120.

common iliac, 331.

dorsalis penis, 381.

external jugular, 2.

saphena, 465.

facial, 61.

femoral, 437.

of Galen, 508.

hæmorrhoidal, 359.

of the hand, 225.

inferior mesenteric, 322.

iliac, common, 331.

external, 331.

innominate, or brachio-cephalic, 88.

intercostal, superior, 47.

internal jugular, 15.

maxillary, 74.

jugular, anterior, 3.

external, 2.

internal, 15.

lingual, 30.

lumbar, 331.

mammary, internal, 46.

maxillary, internal, 74.

median, 214.

basilic, 213.

VEINS (*continued*)

median, cephalic, 213

occipital, 33.

ophthalmic, 169.

pampiniform, plexus of, 555.

pharyngeal, 131

popliteal, 465.

posterior scapular, 46, 254.

saphena, 462.

portal, 319.

phrenic, 330.

profunda femoris, 440.

prostatic plexus of, 358.

radial, superficial, 226.

renal, 330.

saphena, 418, 444.

spermatic, 330, 559.

splenic, 319.

spinal system of, 518.

subclavian, 48.

superior intercostal, 47.

mesenteric, 321.

supra scapular, 46.

thyroid, middle, 17.

superior, 32.

inferior, 46.

ulnar, superficial, 226.

vaginal plexus of, 389.

vena cava, superior, 89.

inferior, 329.

vertebral, 45.

Velum interpositum, 507.

Venesection, anatomy of, 215.

VENTRICLES

lateral of brain, 502.

third ditto, 509.

fourth ditto, 511.

fifth ditto, 506.

of the larynx, 146.

heart, right, 115.

left, 119.

Vermiform, appendix, 308.

eminences of cerebellum, 512, 513.

Vertebral aponeurosis, 182.

Vesiculæ seminales, 363.

structure of, 375.

Vesicles, Graafian, 398.

Vestibule of the ear, 649.

Vieussens, valve of, 508.

Villi of small intestines, 412.

structure of, 412.

Vocal cords, 144.

Wharton's duct, 23.

Whitlow, proper incision in, 243

Willis, circle of, 496.

Winslow, foramen of, 312.

Wrisberg, nerve of, 213.

Zinn, zone of, 542.



