

## **Landmarks, medical and surgical / by Luther Holden.**

### **Contributors**

Holden, Luther, 1815-1905.

### **Publication/Creation**

London : J. & A. Churchill, 1877.

### **Persistent URL**

<https://wellcomecollection.org/works/jtgy8gds>

### **License and attribution**

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>



LANDMARKS  
MEDICAL AND SURGICAL  
—  
LUTHER HOLDEN



# SALEM HOSPITAL LIBRARY




Gift of Dr. Octavius Shreve

Date June, 1932

Book No.







MEDICAL AND SURGICAL  
LANDMARKS



MEDICAL AND SURGICAL

LONDON: PRINTED BY  
SPOTTISWOODE AND CO., NEW-STREET SQUARE  
AND PARLIAMENT STREET

LAMPMAN'S

# LANDMARKS

## MEDICAL AND SURGICAL

BY

LUTHER HOLDEN, F.R.C.S.

MEMBER OF COUNCIL AND OF COURT OF EXAMINERS : SENIOR SURGEON TO  
SAINT BARTHOLOMEW'S AND THE FOUNDLING HOSPITALS



3-C  
H72

LONDON  
J. & A. CHURCHILL, NEW BURLINGTON STREET  
1876





TO  
THE STUDENTS  
PAST AND PRESENT  
OF  
SAINT BARTHOLOMEW'S HOSPITAL


THESE 'LANDMARKS' ARE

Dedicated

BY THEIR SINCERE FRIEND

THE AUTHOR





Digitized by the Internet Archive  
in 2011 with funding from  
Open Knowledge Commons and Harvard Medical School

## P R E F A C E.

---

THESE 'LANDMARKS' have already appeared in Saint Bartholomew's Hospital Reports. They are now republished, with some additions, in the hope that they may be useful to others besides those for whom they were originally intended.

My object has been to collect into a compact form the leading landmarks which help practical surgeons in their daily work. Those relating to the chest and abdomen have been ascertained, with as much precision as natural variations permit, by needles introduced in various directions.

I have to express my acknowledgments to Mr. WALSHAM and to Dr. GODSON for their contributions.





# LANDMARKS

## MEDICAL AND SURGICAL.

---

*‘Chirurgus mente prius et oculis agat quam manu armata.’*

---

I. IN clinical teaching, we often have occasion to point out, on the surface of the living body, what may be called ‘medical and surgical landmarks.’ By ‘landmarks’ we mean surface-marks, such as lines, eminences, depressions, which are guides to, or indications of, deeper-seated parts. We find this practice not only useful, but necessary; because many even advanced students of anatomy are not so ready as they ought to be in their recognition of parts when covered by skin. Men who may be familiar enough with bones, muscles, blood-vessels, or viscera in the dissected subject, are often sadly at fault when they come to put this knowledge into practice.

For instance, ask a student to put his finger on the exact place where he would feel for the head of the radius, the coracoid process of the scapula, the tubercle of the scaphoid bone in the foot; ask him to compress effectually one of the main arteries; to chalk the line of its course; to map on the chest the position of the heart and the several valves at its base; to trace along the walls of the chest the outline of the lungs and pleura; to point out the bony



prominences about the joints, and their relative position in the different motions of the joints; test him about the muscles and tendons which can be seen or felt as they stand out in relief or remain in repose; let him introduce his finger into the several orifices of the body, and say what parts are accessible to the touch: questions such as these, even a good anatomist, unaccustomed to deal with the living subject, might possibly find himself at a loss to answer.

Object in  
view.

2. Our main object, therefore, is to induce in students the habit of looking at the living body with anatomical eyes, and with eyes too at their fingers' ends. The value of this habit cannot be too highly estimated. Is it not of the utmost importance to an operating surgeon that he should have pictures in his mind of the various structures of the body as they lie grouped, connected, and working together? Should he not try at least to see them with the same clearness and accuracy as if they were perfectly transparent?

This habit is within easy reach of any one who has carefully dissected for himself, and acquired the knowledge of what to feel for. Plates will not help him. Let a man examine his own body. Better still that two should work together, each being a 'model' for the other. Best of all, to study with the help of a skeleton.

I would also invite teachers of anatomy to follow the example of Sir C. Bell, who was in the habit of introducing, from time to time, a powerful muscular fellow to his class, 'in order to show them how much of the structure of the body, such as the articulations and the muscles, might be learned without actual dissection.'<sup>1</sup>

At the same time, it is only fair to say that 'landmarks' cannot always be defined with precision. A reasonable latitude must be allowed for natural variations in different persons. Anatomy obvious to the eye and the finger depends in great measure upon the quantity of subcutaneous

<sup>1</sup> Some pertinent remarks on this subject have been made by Mr. C. Heath in a pamphlet 'On Anatomy in relation to Physic.'



fat. In some it stands out beautifully clear ; in others it is masked by corpulency. Selecting, therefore, for study a moderately lean person, let us begin with—

## THE HEAD.

3. The great density of the scalp, more especially at the back of the head, is owing to its intimate connection with the cranial aponeurosis, the scalp vessels and hair bulbs intervening. This density often obscures the diagnosis of tumours on the cranium. A tumour growing upon the head may be either above or below the aponeurosis of the scalp. If below, it will have a firm resisting feel, being bound down by the aponeurosis. Nevertheless its firmness and resistance may not depend simply on its confinement beneath the aponeurosis, but on its having its origin within the skull. Look with suspicion, then, on every tumour on the head that will not readily permit you to move it about, so as to be sure of its connections prior to an attempt at extirpation. Scalp.

The scalp moves freely over the pericranium, to which it is very loosely connected by areolar tissue. When supuration takes place in this tissue free incisions through the dense scalp must be made to let the pus out.

4. The supra-orbital artery can be felt beating just above the supra-orbital notch, and traced for some way up the forehead ; the temporal, about one inch and a quarter behind the external angular process of the frontal bone ; the occipital, near the middle of a line drawn from the occipital protuberance to the mastoid process ; the posterior auricular, behind the apex of the mastoid process. All these arteries can be effectually compressed against the subjacent bone. Arteries of scalp.

5. The skull-cap is rarely symmetrical. This want of symmetry is often obvious. It may occur in men highly gifted, as in the celebrated French anatomist Bichat. As to shape and relative dimensions, no two heads are exactly alike, any more than two faces. It is beside my present Skull-cap.



purpose to go into the question of craniology more than to say that, although the cranium does not follow the brain in all its eminences and depressions so as to be, as it were, a cast of its surface, yet it certainly indicates the dimensions of the great cerebral masses. The frontal and parietal eminences and the occipital region may be taken as a general expression of the development of the corresponding lobes of the brain. To ascertain the relative proportions of these three regions, let a thread be passed from one meatus auditorius to the other, across the frontal, parietal, and occipital eminences respectively.

Frontal  
sinuses.

The 'frontal sinuses' formed by the separation of the two tables of the skull vary much in size in different persons and at different periods of life. This has an important bearing on wounds of the forehead and on trephining in this situation. These 'bumps' do not exist in children, because the tables of the skull do not begin to separate before puberty. From an examination of many skulls in the Hunterian Museum, I find that the absence of the external prominence, even in middle age, does not necessarily imply the absence of the sinus itself, since it may be formed by a retrocession of the inner wall of the skull. In old persons, as a rule, when the sinuses enlarge, it is by the encroachment of the inner table on the brain case. The skull wall here follows the shrinking brain. It is, therefore, important to bear in mind that an adult, and more especially an elderly person, may have a large frontal sinus without any external indication of it.

Neither does a very prominent bump necessarily imply the existence of a sinus at all. The 'bump' may be a mere heaping up of bone, a degradation, as in some Australian skulls.

The same observations apply to the air cells of the mastoid process, which can be felt behind the ear.

Occipital  
protuber-  
ance.

The occipital protuberance, the superior curved line and crest, can be distinctly felt at the back of the head. The protuberance is always the thickest part of the skull-cap, obviously for the purpose of protection.



The posterior inferior angle of the parietal bone, grooved by the lateral sinus, is on a level with the zygoma, and one inch behind the mastoid process.

A line drawn over the head from the root of the nose to the occipital protuberance corresponds with the superior longitudinal sinus. Another line drawn from the occipital protuberance to the front border of the mastoid process corresponds with the lateral sinus. The trunk of the middle meningeal artery runs along the anterior inferior angle of the parietal bone, about one inch and a half behind, and half an inch above, the external angular process of the frontal.

Lines of  
sinuses.

The front borders of the mastoid processes exactly correspond with the middle of the condyles of the occiput, showing how nearly the skull is balanced on the top of the spine.

6. The average thickness of the cap of an adult skull is about  $\frac{1}{8}$  of an inch. The thickest part is at the occipital protuberance, where it is often  $\frac{3}{4}$  of an inch or more, even in an otherwise thin skull. The thinnest part is at the temple, where it may be almost as thin as brown paper. Every one in the habit of making post-mortem examinations knows how much the skull-cap differs in thickness in different persons and in different parts of the same skull. In old persons it is often in some parts not thicker than a shilling, owing to absorption of the *diplœ*. Another point of interest is that the inner plane of the cap does not always correspond with the outer. Hence, in applying the trephine this is not a bad rule—'Think you are operating on the thinnest skull ever seen, and thinner in one half of the circle than the other.'

Thickness  
of skull.

7. The lower level of the anterior and middle lobes of the cerebrum corresponds with a line drawn from the external angular process of the frontal bone to the upper part of the meatus auditorius. Another line drawn from the meatus to the occipital protuberance corresponds with the lower level of the posterior lobe. The lower level of the cerebellum cannot be defined by external examination. It

Levels of  
the brain.



depends upon the extent to which the occipital fossæ bulge into the nape of the neck ; and this bulge varies in different skulls.

### THE FACE.

8. The approaches to the organs of the senses, their ever varying expression, their numerous muscles, and their rich profusion of vessels and nerves, give the face great anatomical importance, which has a most valuable bearing, not only on the practice of surgery, but on the physiognomy of health, and in the diagnosis of disease.

Foramina  
for branches of  
5th nerve.

9. As a surgeon may be called upon to divide either of the three branches of the fifth nerve upon the face, he looks with interest to the precise situations where they leave their bony foramina with their corresponding arteries. The supra-orbital notch or foramen can be felt about the junction of the inner with the middle third of the supra-orbital margin. From this point a perpendicular line drawn with a slight inclination outwards, so as to cross the interval between the two bicuspid teeth in both jaws, passes over the infra-orbital and the mental foramina. The direction of these two lower foramina looks towards the angle of the nose.

Pulley for  
oblique  
muscle.

10. By pressing the thumb beneath the internal angular process of the frontal bone, the cartilaginous pulley of the superior oblique muscle can be distinctly felt. We should be careful not to interfere with this pulley in any operation about the orbit.

Lower  
jaw.

11. The working of the condyle of the jaw vertically and from side to side can be distinctly felt in front of the ear. When the mouth is opened wide, the condyle advances out of the glenoid cavity on to the eminentia articularis, and returns into its socket when the mouth is shut. The muscle which causes this advance is the external pterygoid ; and the object of it is to give the jaw a greater freedom of grinding motion.

The posterior margin of the ramus of the lower jaw corresponds with a line drawn from the condyle to the angle. In opening abscesses in the parotid region, the



knife should not be introduced behind this line for fear of wounding the external carotid artery. Punctures to any depth may be safely made in front of it. They are often necessary where inflammation of the parotid gland ensues after eruptive fevers, and runs on to suppuration. The swelling, tension, and pain are most distressing. Owing to the fibrous framework of the gland, the matter is not in one sac, but diffused. One puncture is not enough. Three or more may be requisite. The blade of the knife should be held horizontally, so as to be less likely to injure the branches of the facial nerve. We are not to be disappointed if no matter flows. The punctures give relief, and matter will exude the next day.

12. A line drawn from the bottom of the lobe of the ear to midway between the nose and the upper lip gives the track of the parotid duct and of the branch of the portio dura which supplies the buccinator. The duct opens into the mouth opposite the second molar tooth of the upper jaw. Parotid duct.

13. Between the root of the zygoma and the ear can be felt the pulsation of the trunk of the temporal artery. The facial artery can be distinctly felt as it passes over the body of the jaw at the anterior edge of the masseter; again near the corner of the mouth close to the mucous membrane: and, lastly, by the side of the ala nasi, up to the inner side of the tendo oculi. By holding the lips between the finger and thumb the coronary arteries are felt under the mucous membrane. The facial vein does not accompany the tortuous artery, but runs a straight course from the inner angle of the eye to the front border of the masseter, just behind the artery. Facial artery.

14. The opening between the eyelids varies in size in different persons; hence more of the eyeball is seen in some than in others, and the eye appears larger. Although human eyes do vary a little in size, yet the actual difference is by no means so great as is generally supposed. The size of the fissure has more to do with it. Contrast the narrow elliptical fissure of the Chinese and Mongolian Eyelids.



ances, and the apparent smallness of their eyes as compared with Europeans. As a rule the external angle of the lid is higher than the internal. When not exaggerated, it gives the face an arch and pleasing expression.

Speaking generally, a prominent eye is a long eye, and *ipso facto* a myopic eye. On the other hand, a flat eye bespeaks a small eye, with deficient refraction.

Evert the lids to see the Meibomian glands; observe their perpendicular arrangement, in the substance of the lid cartilages.

The free borders of the lids are not bevelled, as described by J. L. Petit and most anatomists, 'so as to form with the globe of the closed eye a triangular canal for the flow of the tears.' On the contrary, it is easily seen that the lid margins, when closed, come into accurate contact. Their plane is not exactly horizontal, but slightly inclined upwards.

Every time the eye is shut, the ball moves upwards and inwards, so that the cornea is completely covered by the upper lid. This may be well seen by raising the lid of a sleeping infant; also in cases of low fever, when the lid is not completely closed. The reason of this upturning of the eye is obviously to clear the cornea, and to protect it from the light.

A careful examination of the motion of the lower lid in the act of shutting the eye proves that it is a double motion. The lid is not only slightly raised, but drawn inwards about  $\frac{1}{12}$  of an inch. The object of this second movement is to sweep any particles of dust towards the inner canthus.

Puncta  
lachryma  
lia.

15. The puncta lachrymalia are distinctly visible at the inner angles of the lids. The lower punctum is larger and a little more external than the upper, so that they may not be exactly opposite. The direction, too, of the puncta deserves notice. Their open mouths look a little backwards, ready to imbibe the tears. When their proper bearing is lost, as in facial paralysis or a cicatrix near the lid, the tears overflow the cheek. The length of the



lachrymal canals is from three to four lines. The lower is a little shorter and wider than the upper. As they both make a little angle in their course, soon after their orifices, the lids should be drawn outwards to straighten them when we introduce a probe.

16. To find the lachrymal sac, draw outwards the eye-lids to tighten the tendo oculi, which crosses the sac a little above its middle. A knife introduced just below the tendon close to the edge of the orbit would enter the sac. The angular artery and vein would be on the inner side of the incision. A probe directed in a line with the inner edge of the orbit, *i.e.* downwards, outwards, and backwards, would traverse the nasal duct, and appear in the inferior meatus of the nose.

Lachry-  
mal sac.

The tendo oculi serves many purposes beside giving attachment to the cartilages and muscles of the lids. One purpose is said to be to pump the tears into the lachrymal sac. Place a finger on the tendon, and feel that it tightens every time the lids are closed. The tendon, being intimately connected to the sac, draws, as it tightens, the sac wall outwards and forwards, and in this way it may pump along the lachrymal canals any fluid collected at the angle of the eye.

17. The nasal duct is from six to eight lines long, and narrowest in the middle of its course. Its termination in the inferior meatus lies under the inferior spongy bone, about a quarter of an inch behind the bony edge of the nostril. The appearance of the orifice in the dry bone conveys no idea of its size and shape in life; <sup>1</sup> for it is diminished by a valve-like fold of mucous membrane, so as to become, in most cases, a mere slit, not exceeding a line in diameter.

Nasal  
duct.

The facility with which instruments can be introduced into the nasal opening of the duct depends upon its position as well as its size. This position varies in different instances. Sometimes it opens directly into the roof of the inferior meatus, in which case the hole is large and round, so that tears readily drop into the nose. In other instances

<sup>1</sup> Alluded to by Morgagni, Adv. Anat. xli.



the opening is situated on the outer wall of the meatus, and is then always such a narrow fissure as to be hardly discernible. The practical conclusion then is, that a probe can be easily introduced when the opening is in the roof of the meatus, but not without difficulty and laceration of the mucous membrane when on the outer wall. This difficulty indeed may be increased by the narrowness of the meatus, arising from an unusual curvature of the spongy bone.

Nose.

18. The line where the cartilages of the nose are attached to the nasal and superior maxillary bones can be traced with precision. The close connection of the skin to the cartilages admits of no stretching; hence the acute pain felt in erysipelas and boils on the nose. The external aperture of the nose is always placed a little lower than the floor of the nostril, so that the nose must be pulled up before we can inspect its cavities.

Looking into the nostrils, we find that the left is, in the majority of cases, narrower than the right, owing to the inclination of the septum towards the left. A communication sometimes exists between them, through a hole in the septum, as was the case in the celebrated anatomist Hildebrandt. By dilating the *alæ nasi*, we can get a view of the end of the inferior spongy bone. The middle spongy bone cannot be seen: its attachment to the ethmoid is high up, nearly opposite the *tendo oculi*. The cavities are so much narrowed transversely by the spongy bones, that in the extraction of polypi it is better to dilate the blades of the forceps perpendicularly, and near the septum.

Mouth.

19. What can be seen and felt through the mouth? The upper surface of the tongue, '*speculum primarum viarum*,' is a study in itself. We notice, on its under surface, a median furrow, on each side of which stands out the ranine vein, lying upon the prominent fibres of the *lingualis*. On the floor of the mouth is the '*frenum linguæ*,' in the middle line, with the orifices of the ducts of the submaxillary glands on each side of it. The glands themselves can be felt immediately beneath the mucous membrane.

We can feel the attachment of the '*genio-hyo-glossi*'



behind the symphysis of the jaw. The division of this attachment would enable a surgeon to draw the tongue more freely out of the mouth in any attempt to remove carcinoma extending far back into its root.

There is great difference in the shape of the hard palate; this difference depends upon the depth of the alveolar processes. In some it forms a fine broad arch; in others it is contracted, and rises almost to a point like a Gothic arch, and materially impairs the tone of the voice.

To examine the throat well, the nose should be held so Throat. as to compel breathing through the mouth. Thus the soft palate will be raised, the palatine arches widened, and the tonsils and the back of the pharynx fairly exposed. Pressing the tongue downwards, provided it be done very gently, is also of advantage. Rude treatment the tongue would resist.

Put the finger into the mouth, and feel the anterior border of the coronoid process of the jaw. On the inner side of this process, between it and the tuberosity of the upper jaw, is a recess, where a deeply-seated temporal abscess might burst, or might be opened. On the inner side of the last molar tooth of the upper jaw we can distinctly feel the hamular process of the sphenoid bone; also the lower part of the pterygoid fossa, and the internal pterygoid plate. Behind, and on the outer side of the last molar, can be felt part of the back of the antrum and of the lower part of the external pterygoid plate.

When the mouth is wide open, the pterygo-maxillary ligament forms a prominent fold, readily seen and felt beneath the mucous membrane. A little below the attachment of this ligament to the lower jaw we can easily feel the gustatory nerve, as it runs close to the bone below the last molar tooth. The exact position of the nerve can be ascertained in one's own person by the acute pain on pressure. A division of the nerve, easily effected by a small incision in the right place, gives much temporary relief in cases of advanced carcinoma of the tongue.



To feed a patient in spasmodic closure of the jaw, it is well to know that there exists behind the last molar teeth a space sufficient for the passage of a small tube.

Lift up the upper lip and examine the front wall of the antrum. The proper place in which to tap it is above the second bicuspid tooth, about one inch above the margin of the gum.

Posterior  
nares.

20. A surgeon's finger should be familiar with the feel of the posterior nares, and of all that is within reach behind the soft palate. This is important in relation to the attachment of polypi, to plugging the nostrils, and the proper size of the plug. In the examination of this part of the back of the throat it is necessary to throw the head well back, because, in this position, nearly all the pharynx in front of the basilar process comes down below the level of the hard palate, and can be seen as well as felt. But when the skull is horizontal, *i.e.* at a right angle with the spine, and the hard palate is on a level with the margin of the foramen magnum, the basilar process is concealed from view.

The head, then, being well back, introduce the forefinger behind the soft palate, and turn it up towards the base of the skull. You feel the strong grip of the superior constrictor. Hooking the finger well forwards, you can feel the contour of the posterior nares. Their size depends upon the anterior, but rarely exceeds a small inch in the long diameter, and a small half-inch in the short. The plug for the posterior nares should not be larger than this. Their plane is not perpendicular, but slopes a little forwards. You can feel the septum formed by the vomer, and also the posterior end of the inferior spongy bone in each nostril.

Tonsils.

21. Before taking leave of the throat, look well at the position of the tonsils between the anterior and posterior half arches of the palate. In a healthy state they should not project beyond the level of these arches. In all operations upon the tonsils, we should remember the close proximity of the internal carotid artery to their outer side.



Nothing intervenes but the pharyngeal aponeurosis, and the superior constrictor of the pharynx. Hence the rule in operating on the tonsils always to keep the point of the knife inwards.

In troublesome hemorrhage from the tonsils, after an incision, it is well to know that they are accessible to pressure if necessary by means of a padded stick, or even a finger.

22. A word or two on the lines of the face as indicative of expression. Every one pays unconscious homage to the study of physiognomy, when, scanning the features of a stranger, he draws conclusions concerning his intelligence, disposition, and character. Without discussing how much physiognomy is really worth, there can be no doubt that it is a mistake to place it in the same category as phrenology, since the latter lacks that sound basis of physiology which no one can deny to the former. Features.

The muscles of the features are generally described as arising from the bony fabric of the face, and as inserted into the nose, corners of the mouth, and the lips. But this gives a very inadequate idea of their true insertion. They drop fibres into the skin all along their course, so that there is hardly a point of the face which has not its little fibre to move it. The habitual recurrence of good or evil thoughts, the indulgence in particular modes of life, call into play corresponding sets of muscles which, by producing folds and wrinkles, give a permanent cast to the features, and speak a language which all can understand, and rarely misleads. Schiller puts this well when he says that 'it is an admirable proof of infinite wisdom that what is noble and benevolent beautifies the human countenance; what is base and hateful imprints upon it a revolting expression.'

## THE NECK.

23. We must notice first the direction of the subcutaneous veins. The chief subcutaneous vein is the external jugular. Its course corresponds with a line drawn from the Subcutaneous veins.



angle of the jaw to the middle of the clavicle, where it joins the subclavian. It is made more prominent by putting the sterno-mastoid into action, and pressing in the supra-clavicular fossa. It is exceptionally joined by a branch which runs over the clavicle, and is termed 'jugulo-cephalic.' The anterior jugular generally runs along the front border of the sterno-mastoid.

Parts in  
central  
line.

24. Immediately below and nearly on a level with the lower jaw we feel the body of the os-hyoides, and can trace backwards on each side the whole length of the cornua. They might easily be broken by the grasp of a garotter. Below the body of the os-hyoides is the gap above the thyroid cartilage. This gap corresponds with the anterior thyro-hyoid ligament and the apex of the epiglottis: so that in cases of cut throat in this situation, nearly the whole of the epiglottis lies above the wound.

The projection and depth of the notch in the thyroid cartilage, or 'pomum Adami,' varies in prominence in different persons. Between the notch and the hyoid bone there is a large bursa, to facilitate the play of the cartilage beneath the bone in deglutition. The notch does not appear till puberty, and is, throughout life, much less distinct in the female than the male. The finger can trace the upper borders and cornua of the thyroid cartilage: its lower cornua can be felt by the side of the cricoid.

On each side of the thyroid cartilage we can recognise the lateral lobes of the thyroid gland. On the upper and front part of the gland we can distinctly feel the pulsation of the superior thyroid artery. This pulsation, coupled with the fact that the gland rises and falls with the larynx in deglutition, is the best distinction between a bronchocele and other tumours resembling it.

Below the angle of the thyroid cartilage we feel the interval between it and the cricoid, which is occupied by the crico-thyroid membrane. In laryngotomy we cut through this membrane transversely close to the upper edge of the cricoid cartilage, in order that the incision may be as distant as possible from the attachment of the vocal cords.



25. The projection of the cricoid cartilage is a point of great interest to the surgeon, because it is his chief guide in opening the air passages, and can always be felt even in infants, however young or fat. It corresponds with the fifth cervical vertebra. The commencement of the œsophagus lies behind it: here, therefore, a foreign substance too large to be swallowed would probably lodge, and might be felt externally.

Cricoid  
cartilage.

Again, a transverse line drawn from the cricoid cartilage across the neck would pass over the spot where the omohyoid crosses the common carotid. Just above this spot is the most convenient place for tying the artery.

26. Those who have not directed their attention to the subject are hardly aware what a little distance there is between the cricoid cartilage and the upper part of the sternum. In a person of the average height sitting with the neck in an easy position, the distance is about one inch and a half. When the neck is stretched about three-quarters of an inch more is gained. Thus, we have generally not more than seven or eight rings of the trachea above the sternum. None of these rings can be felt externally. The second, third, and fourth are covered by the isthmus of the thyroid gland. The trachea, it should be remembered, recedes from the surface more and more as it descends, so that, just above the sternum in a short, fat-necked adult, the front of the trachea would be quite one inch and a half from the skin.

27. In the dead subject nothing is more easy than to open the trachea: in the living, no operation may be attended with greater difficulties. In urgent dyspnœa you must expect to find the patient with his head bent forwards, and the chin dropped, to relax as much as possible the parts. On raising his head, a paroxysm of dyspnœa is almost sure to come on, threatening instant suffocation. The elevator and depressor muscles draw the trachea and larynx up and down with a rapidity and a force which may bring the cricoid cartilage within half-an-inch of the sternum. The great thyroid veins which descend in front of the

Trachea.



trachea are sure to be distended. There may be a middle thyroid artery. In children the two lobes of the thymus may rise in front of the trachea as high as the thyroid gland, so that the whole length of the air-tube may be covered by important parts which ought not to be divided. Considering all these possible complications, the least difficult mode of proceeding is to open the trachea just below the cricoid cartilage, and, if more room be requisite, to divide the cricoid itself.

Sterno-  
mastoid  
muscle.

28. The sterno-mastoid muscle is the great surgical landmark of the neck. It stands out in bold relief when the head turns towards the opposite shoulder. Its inner border overlaps the common carotid, which can be easily compressed for a short time against the spine about the level of the cricoid cartilage. The precise course of the artery corresponds with a line drawn from the sterno-clavicular joint to midway between the angle of the jaw and the mastoid process.

Between the sternal origins of the sterno-mastoid is the fossa above the sternum, more or less perceptible in different necks. As it rises and falls, especially in distressed breathing, it was called by the old anatomists 'fonticulus gutturis.' In beautiful necks, as seen in the 'Venus,' it is filled up by fat.

Notice the interval between the sternal and clavicular origins of the sterno-mastoid. A knife introduced a very little way down this interval would wound, slanting outwards the common carotid, slanting inwards the internal jugular vein. These facts are of importance in performing the subcutaneous section of the tendon of this muscle.

Sterno-  
clavicular  
joint.

29. Many important parts lie behind the sterno-clavicular joint. There is the commencement of the vena innominata; behind this comes the division of the arteria innominata on the right side, and the common carotid on the left. Deeper still, the apex of the pleura and lung rises into the neck.

In a child, say of five years old, the arteria innominata often lies in front of the trachea and divides a little higher than the joint: a point to be remembered in tracheotomy.



30. The extent to which the apex of the lung rises into the neck is greater than is generally supposed. Many observations in reference to this point lead to the conclusion that the lung rises behind the sterno-mastoid, on an average, one inch and a half above the clavicle ; in persons with long necks, as much as two inches. The apex of the lung and pleura is covered by the clavicular origin of the sterno-mastoid, the sterno-thyroid, and a part of the scalenus anticus. It is also crossed by the subclavian vessels in the first part of their course. As this cervical portion of lung is peculiarly obnoxious to tubercular disease, it should always be carefully examined. Its condition may be ascertained by percussion on the clavicle.

Apex of lung in the neck.

31. The hollow above the clavicle, between the sterno-mastoid and the trapezius, is very manifest in emaciation and old age. Notice the termination here of the external jugular vein. In some necks only a small depression is visible, particularly when the trapezius has a broad insertion into the clavicle, and comes well forwards, so that its front border gives a graceful contour to the base of the neck.

Supra-clavicular fossa.

32. In this depression, near the border of the sterno-mastoid, we feel the pulsation of the subclavian artery. Here the artery lies upon the first rib, and can be effectually compressed. But the pressure must be made in the right direction, or the artery will be pressed off the rib instead of against it. The plane of the rib is such, that the pressure, to be effectual, must be made in a direction downwards and a little inwards. It is more convenient to stand behind the shoulder and make the pressure with the thumb.

Subclavian artery.

It is worth remembering that the outer border of the sterno-mastoid corresponds pretty nearly with the outer edge of the scalenus anticus, which is the surgical guide to the subclavian artery.

By pressing deeply at the upper part of the supra-clavicular fossa, the transverse process of the seventh cervical vertebra can be distinctly felt.

In long and thin necks, a thin cord is perceptible,



running nearly parallel with and just above the clavicle. It is the posterior belly of the omo-hyoideus. See it rising and falling in breathing, and making tense during inspiration that part of the cervical fascia which lies over the cervical portion of the lung. Thus it may be said to be in all respects a muscle of inspiration, co-operating with the sterno-mastoid and scaleni. In the language of transcendental anatomy, we may say that the central tendon of the omo-hyoid represents a rudimentary cervical rib. Its posterior belly is analogous to a serration of the serratus magnus; its anterior belly to a sterno-hyoid.

### THE CHEST.

33. As a rule, the right half of the chest is larger than the left. Of ninety-two persons of the male sex and good constitutions, seventy-one had the right side the larger; eleven the left; ten had both sides equal. The maximum of difference in favour of the right was  $1\frac{1}{4}$  inch. The measurements were made on a plane with the nipple.

Peculiarities in the female.

34. The chest of the female differs from that of the male in the following points:—Its general capacity is less; the sternum is shorter; the upper opening is larger in proportion to the lower; the upper ribs are more movable, and therefore permit a greater enlargement of the chest at its upper part, in adaptation to the requirements of pregnancy.

35. The top of the sternum is on a level with the second dorsal vertebra; and the available space between the top of the sternum and the spine is hardly more than two inches.

Parts behind first bone of sternum.

36. There is little or no lung behind the first bone of the sternum, the space being occupied by the trachea and large vessels as follows:—

The left vena innominata crosses the sternum just below the upper border. Next come the great primary branches of the arch of the aorta. Deeper still is the trachea dividing into its two bronchi opposite the junction



of the first and second bones of the sternum. Deepest of all is the œsophagus.

About one inch from the upper border of the sternum is the highest part of the arch of the aorta, which lies on the bifurcation of the trachea.

37. The course of the arteria innominata corresponds with a line drawn from the middle of the junction of the first with the second bone of the sternum, to the right sterno-clavicular joint. When the artery rises higher than usual into the neck, its pulsation can be felt in the fossa above the sternum.

38. In fat persons it is often difficult to count the ribs ; hence the following rules may be useful :—

Rules for  
counting  
the ribs.

*a.* Find and mark the transverse projection, slight, but always to be felt, at the junction of the first with the second bone of the sternum. This corresponds with the level of the cartilage of the second rib.

*b.* The nipple in the male is placed, in the great majority of cases, between the fourth and the fifth ribs, about three-quarters of an inch external to their cartilages.

*c.* The lower external border of the pectoralis major corresponds with the direction of the fifth rib.

*d.* A line drawn horizontally from the nipple round the chest cuts the sixth intercostal space midway between the sternum and the spine. This is a useful rule in tapping the chest.

*e.* When the arm is raised, the first visible digitation of the serratus magnus corresponds with the sixth rib. The three lower visible digitations correspond respectively with the seventh, eighth, and ninth ribs.

*f.* The scapula covers the ribs from the second to the seventh, inclusive.

*g.* The eleventh and twelfth ribs can be felt even in corpulent persons, outside the erector spinæ, sloping downwards.

39. Immediately below the clavicle we recognise the triangular interval between the pectoralis major and the deltoid. This space varies in different cases, depending on

Interval  
below  
clavicle.



the distance between the muscles. It is important as a guide to the coracoid process and the axillary artery. In a case of injury to the shoulder, to ascertain whether the coracoid process is broken, carry the arm outwards, to put the deltoid and pectoral muscles on the stretch, and make manifest the space between their opposite borders. Pressing the thumb into the space we can feel the inner side of the coracoid process, the apex being under the fibres of the deltoid; thus it is easy to ascertain whether it be broken. Moreover, this space corresponds with the line of the axillary artery; here its pulsation can be distinctly felt, and here it can be compressed against the second rib.

Internal  
mammary  
artery.

40. The line of the internal mammary artery runs perpendicularly behind the cartilages of the ribs, about half an inch from the sternum. The perforating branch through the second intercostal space is generally the largest.

Outline of  
heart on  
chest wall.

41. To have a general idea of the form and position of the heart, map out its outline on the wall of the chest, as follows :—

*a.* To find the base from which the great vessels arise, draw a transverse line across the sternum about the upper border of the junction of the third rib; continue the line half an inch to the right of the sternum and one inch to the left.

*b.* To find the apex, mark a point about two inches below the left nipple, and one inch to its sternal side. This point will be between the fifth and sixth ribs.


*c.* To find the lower border which lies on the central tendon of the diaphragm, draw a line, slightly curved downwards, from the apex across the bottom of the sternum (not the ensiform cartilage) as far as its right edge.

*d.* To define the right border formed by the right auricle, continue the last line upwards with an outward curve, so as to join the right end of the base.

*e.* To define the left border formed by the left ventricle, draw a line curving to the left, but not including the nipple, from the left end of the base to the apex.



Such an outline (seen in the cut with the angles rounded off) shows that the apex of the heart points downwards and towards the left, the base a little upwards and towards the right; that the greater part of it lies in the left half of



OUTLINE OF THE HEART, ITS VALVES, AND THE LUNGS.

the chest, and that the only part which lies to the right of the sternum is the right auricle. A needle introduced in the third, the fourth, or the fifth right intercostal space close to the sternum would penetrate the lung and the right auricle.

A needle passed through the second intercostal space, close to the right side of the sternum, would, after passing through the lung, enter the pericardium and the most prominent part of the bulge of the aorta.

A needle passed through the first intercostal space,



close to the right side of the sternum, would pass through the lung and enter the superior vena cava above the pericardium.

42. The best definition of that part of the præcordial region which is less resonant on percussion, was given by Dr. Latham years ago in his 'Clinical Lectures.' 'Make a circle of two inches in diameter round a point midway between the nipple and the end of the sternum. This circle will define, sufficiently for all practical purposes, that part of the heart which lies immediately behind the wall of the chest, and is not covered by lung or pleura.'

Apex of  
the heart.

The apex of the heart pulsates between the fifth and sixth ribs, two inches below the nipple, and one inch to its sternal side. The place and extent, however, of the heart's impulse, vary a little with the position of the body. Of this any one may convince himself by leaning forwards, backwards, on this side and on that; feeling, at the same time, the heart. Inspiration and expiration also alter the position of the heart. In a deep inspiration it may descend half an inch, and can be felt beating at the pit of the stomach.

Valves of  
the heart.

43. The aortic valves lie behind the third intercostal space, close to the left side of the sternum.

The pulmonary valves lie in front of the aortic behind the junction of the third rib, on the left side, with the sternum.

The tricuspid valves lie behind the middle of the sternum, about the level of the fourth costal cartilage.

The mitral valves lie behind the third intercostal space, about one inch to the left of the sternum.

Thus these valves are so situated that the mouth of an ordinary sized stethoscope will cover a portion of them all, if placed over the junction of the third intercostal space, on the left side, with the sternum. All are covered by a thin layer of lung; therefore we hear their action better when the breathing is for a moment suspended.

44. Now let us trace on the chest the outline of the



lungs, with as much precision as their expansion and contraction in breathing permit. (See the cut.)

Outline of  
the lungs.

45. The apex of each lung rises into the neck behind the sternal end of the clavicle and sterno-mastoid muscle as much as  $1\frac{1}{2}$  inch, in females rather higher than in males (30). From the sternal ends of the clavicles the lungs converge, so that their thin edges almost meet in the mesial line on a level with the second costal cartilage. Thus there is little or no lung behind the first bone of the sternum. From the level of the second costal cartilage to the level of the fourth, the margins of the lungs run parallel, or nearly so, close behind the middle of the sternum: consequently their thin edges overlap the great vessels at the base of the heart.

Below the level of the fourth costal cartilage the margins of the lungs diverge, but not in an equal degree. The margin of the right corresponds with the direction of the cartilage of the sixth rib: the margin of the left, being notched for the heart, runs behind the cartilage of the fourth. A line drawn perpendicularly from the nipple would find the lung margin about the lowest part of the sixth rib. Laterally, *i.e.* in the axillary line, the lung margin comes down as low as the eighth rib: posteriorly, *i.e.* in the dorsal or scapular line, it descends as low as the tenth.

It should be remembered that, in a deep inspiration, the lung margins descend about  $1\frac{1}{2}$  inch.

In children the lungs are separated in front by the thymus gland. Allowance should be made for this. About the approach of puberty the thymus disappears.

46. The direction of the anterior mediastinum is not straight down the middle of the sternum, but slants a little to the left, owing to the position of the heart. The right pleural sac generally encroaches a little upon the left behind the middle of the sternum. A needle introduced here, in the majority of cases, would go through the right pleura.

Anterior  
mediasti-  
num.

47. The reflection of the pleura from the wall of the



Reflection  
of pleura.

chest on to the diaphragm corresponds with a sloping line drawn from the bottom of the sternum over the cartilages of the ribs down to the lower border of the last rib.

Since the pleura lines the inside of the last rib, a musket ball or other foreign body, loose in the pleural sac, and rolling on the diaphragm, might fall to the lowest part of the sac, which would be between the eleventh and twelfth ribs. The ball might be extracted here. The chest might also be tapped here, but not with a trochar, since a trochar would penetrate both layers of pleura, and go through the diaphragm into the abdomen.

The operation should be done cautiously, by an incision beginning about two inches from the spine, on the outer border of the 'erector spinæ,' on a level between the spines of the eleventh and twelfth dorsal vertebræ. The intercostal artery will not be injured if the opening be made below the middle of the space, which is very wide.<sup>1</sup>

### THE BACK.

Median  
furrow.

48. In a muscular man a furrow, caused by the prominence of the erector spinæ on each side, runs down the middle of the back. The bottom of the furrow corresponds with the interval between the spine of the last lumbar and that of the first sacral vertebra, which is usually small.

Spines of  
vertebræ.

49. A little friction with the fingers down the backbone will cause the spines of the vertebræ to be tipped with red, so that they can be easily counted and any deviation from the straight line detected. Still it is worth remembering that the spine of the third dorsal is on a level with the commencement of the spine of the scapula—the spine of

<sup>1</sup> Special experiments upon this subject were made many years ago by Professor Quekett in the work-rooms of the College of Surgeons.



the seventh dorsal on a level with the inferior angle of the scapula—the spine of the last dorsal on a level with the head of the last rib.

Opposite the spine of the third, in some cases the fourth, dorsal vertebra is the division of the trachea. In front this division is on the level of the junction of the first with the second bone of the sternum.

Division  
of the  
trachea.

The root of the spine of the scapula is marked by a slight dimple in the skin. This is on a level with the third intercostal space. A stethoscope placed on the inner side of this dimple would cover the bronchus, more especially the right, since it is nearer to the chest wall.

50. The place where the kidney is most accessible to pressure is below the last rib, on the outer edge of the erector spinæ.

51. The best incision for opening the descending colon is in a slightly sloping line beginning at the outer edge of the erector spinæ, midway between the crest of the ilium and the last rib, and continued across the flank for three inches or more, according to the amount of subcutaneous fat.

52. In the pit of the neck we can feel the trapezius and the ligamentum nuchæ. By pressing deeply we detect the forked and prominent spine of the second cervical vertebra.

53. The spines of the third, fourth, and fifth cervical vertebræ recede from the surface to permit free extension of the neck, and cannot often be felt. But the spines of the sixth and seventh (v. prominens) stand out well.

54. Notice that most of the spines of the dorsal vertebræ, owing to their obliquity, do not tally with the heads of their corresponding ribs. Thus, the spine of the second dorsal corresponds with the head of the third rib; the spine of the third dorsal with the head of the fourth rib, and so on till we come to the eleventh and twelfth dorsal vertebræ, which do tally with their corresponding ribs. All this, however, is best seen in the skeleton.

55. The spines of the vertebræ may be useful as land-



marks indicative of the levels of important organs ; I have therefore arranged them in a tabular form, thus :—

CERVICAL.	{	7th. Apex of lung : higher in the female than in the male.
		1st.
DORSAL SPINES.	{	2nd.
		3rd. Aorta reaches spine. Apex of lower lobe of lung.
		4th. Aortic arch ends. Angle of bifurcation of trachea. Upper level of heart.
		5th.
		6th.
		7th.
		8th. Lower level of heart. Central tendon of diaphragm.
		9th. Œsophagus and vena cava through diaphragm. Upper edge of spleen.
		10th. Lower edge of lung. Liver comes to surface pos- teriorly. Cardiac orifice of stomach.
		11th. Lower border of spleen. Renal capsule.
		12th. Lowest part of pleura. Aorta through diaphragm. Pylorus.
LUMBAR.	{	1st. Renal arteries. Pelvis of kidney.
		2nd. Termination of spinal cord. Pancreas. Duodenum just below. Receptaculum chyli.
		3rd. Umbilicus. Lower border of kidney.
		4th. Division of aorta.
		5th.

Origins of  
the spinal  
nerves.

56. It is useful to know opposite what vertebræ the spinal nerves in the different regions arise from the spinal cord. They arise as follows :—

The origins of the eight cervical nerves correspond to the interval between the occiput and the sixth cervical spine.

The origins of the first six dorsal nerves correspond to the interval between the sixth cervical and the fourth dorsal spines.

The origins of the six lower dorsal nerves correspond to the interval between the fourth and the eleventh dorsal spines.



The origins of the five lumbar nerves correspond to the interval between the eleventh and twelfth dorsal spines.

The origins of the five sacral nerves correspond to the spines of the last dorsal and the first lumbar vertebræ.

57. The movements of which the spine is capable are threefold: 1, Flexion and extension; 2, lateral inclination; 3, torsion. Flexion and extension are freest between the third and the seventh cervical vertebræ, between the eleventh dorsal and the second lumbar, and between the last lumbar and the sacrum. This is well marked in severe cases of opisthotonos, where the body is supported on the back of the head and heels.<sup>1</sup>

Move-  
ments of  
the spine.

Still better may it be observed when a mountebank bends backwards, and touches the ground with his head.

The lateral movement is freest in the neck and the loins.

The movement of torsion or rotation round its own axis may be proved by the following experiment:—Seated upright, with the back and shoulders well applied against the back of a chair, we can turn the head and neck as far as 70°. Leaning forwards so as to let the dorsal and lumbar vertebræ come into play, we can turn 30° more.

58. There are a few points worthy of observation about the scapula. It covers the ribs from the second to the seventh inclusive. We can feel its superior angle covered by the trapezius. The inferior angle is covered by the latissimus dorsi, which keeps it well applied against the ribs in the strong and athletic; but in weak and consumptive persons the lower angles of the scapulæ project like wings—hence the terms ‘scapulæ alatæ.’

Position  
and mo-  
tions of  
scapula.

A line drawn horizontally from the spine of the sixth dorsal vertebra over the inferior angle of the scapula gives the upper border of the latissimus dorsi. Another line drawn from the root of the spine of the scapula to the spine of the last dorsal vertebra gives the lower border of the trapezius, which stands a little in relief.

59. The sliding movement of the scapula on the chest

<sup>1</sup> See a beautiful illustration of this in Sir C. Bell’s ‘Anatomy of Expression,’ p. 160.



can be best understood only on the living subject. It can move not only upwards and downwards as in shrugging the shoulders—backwards and forwards as in throwing back the shoulders—but it has a rotatory movement round a movable centre. This rotation is seen while the arm is being raised from the horizontal to the vertical position, and is effected by the co-operation of the trapezius with the deltoid. The glenoid cavity is thus made to look upwards, the inferior angle slides forwards, and is well held under the latissimus dorsi.

60. For the medical examination of the back, the patient should sit with the arms hanging between his thighs, to lower the scapulæ as much as possible. In this position the spine of the scapula lies near the septum between the upper and lower lobes of the lung; the apex of the lower lobe being about the level of the third rib.

### THE ABDOMEN.

The student is assumed to be familiar with the arbitrary lines dividing the abdomen into regions.

Abdomi-  
nal lines.

61. The linea alba, or central line of the abdomen, marks the union of the aponeuroses of the abdominal muscles. It runs from the apex of the ensiform cartilage to the symphysis pubis. As this line is the thinnest and least vascular part of the abdominal wall, we make our incision along it in ovariectomy, and in the high operation of lithotomy; in it, we tap the abdomen in ascites, and the distended bladder in retention of urine.

The so-called 'linea semilunaris,' where the aponeuroses of the abdominal muscles divide to form the sheath of the rectus, corresponds with a line, drawn slightly curved, with the concavity towards the linea alba, from the lowest part of the seventh rib to the spine of the pubes. This line would be in an adult about three inches from the umbilicus; but in an abdomen distended by dropsy or other cause, the distance is increased in proportion.



It is important to know the position of the lineæ transversæ, or tendinous intersections across the rectus abdominis. There are rarely any below the umbilicus, and generally three above it. The first is about the level of the umbilicus. The second is about four inches higher—that is, about the level of the lowest part of the tenth rib. These are the principal lines, and they divide the upper part of each rectus into two nearly quadrilateral portions, an upper and a lower: of these, those on the right side are a trifle larger than on the left. We see these muscular squares pretty plainly in some athletic subjects. Much more frequently we see them, too much exaggerated, on canvas and in marble. Artists are apt to overdo them, and make the front of the belly too much like a chess board. It is lucky for them that all the world do not see with anatomical eyes.

To us, as surgeons, a familiarity with the shape and position of these divisions of the rectus is of importance, lest we should, in ignorance, make a mistake in our diagnosis. A spasmodic contraction of one of these divisions, particularly the upper, or a collection of matter within its sheath, has been frequently mistaken for deep-seated abdominal disease.

A transverse line drawn from one anterior superior spine of the ilium to the other is about the level of the promontory of the sacrum. Such a line will always show whether the pelvis is horizontal or not.

62. The umbilicus is not midway between the ensiform cartilage and the pubes, but rather nearer to the pubes. In all cases it is situated above the centre of a man's height. It is a vulgar error to say that when a man lies with legs and arms outstretched, and a circle is drawn round him, the umbilicus lies in the centre of it. This central point is in most persons just above the pubes.

Umbilicus.

In very corpulent persons two deep transverse furrows run across the abdomen. One runs across the navel and completely conceals it. The other is lower down, just above the fat of the pubes. In tapping the bladder above



the pubes in such a case, the trochar should be introduced where this line intersects the *linea alba*.

Although the position of the umbilicus varies a little in different persons, as the abdomen is unusually protuberant or the reverse, still, as a general rule, it is placed about the level of the body of the third lumbar vertebra. Now, since the aorta divides about the lower border of the fourth lumbar, it follows that the best place to apply pressure on this great vessel for the cure of an aneurism of the common iliac is one inch below the umbilicus, and slightly to the left of it. That the aorta can, under favourable circumstances, be compressed under chloroform sufficiently to cure an aneurism below it, is proved by recorded cases, and by none more effectually than by a case related in the second volume of the 'Reports of St. Bartholomew's Hospital.'

It may be asked, why not apply pressure on the aorta above the umbilicus? The answer is, that the aorta above the umbilicus is farther from the surface, and is, moreover, covered by important structures upon which pressure would be dangerous.

Parts be-  
hind *linea*  
*alba*.

63. Let us next consider what viscera lie immediately behind the *linea alba*. For two or three fingers' breadth below the ensiform cartilage there is the left lobe of the liver, which here crosses the middle line. Below the edge of the liver comes the stomach, more or less in contact with the *linea alba*, according to its degree of distension. In extreme distension the stomach pushes everything out of the way, and occupies all the room between the liver and the umbilicus. When empty and contracted, it retreats behind the liver, and lies flat like a piece of intestine in front of the pancreas at the back of the abdomen; thus giving rise to the hollow termed the pit of the stomach. But as the stomach distends, its great curvature comes to the front, making a considerable fulness where there was a pit. Between three and four inches above the umbilicus lies the middle of the transverse colon, occupying space according to its distension. Behind and



below the umbilicus, supposing the bladder contracted, are the small intestines, covered by the great omentum.

64. The peritoneum is in contact with the linea alba all the way down to the pubes, when the bladder is empty. But when the bladder distends, it raises the peritoneum from the middle line above the pubes ; so that with a bladder distended half-way up to the umbilicus, there is a space of nearly two inches above the symphysis where the bladder may be tapped without risk of injury to the peritoneum. For the same reason, we have space sufficient for the successful performance of the high operation for stone. This fact in anatomy must have been well understood by Jean de Dot, the smith at Amsterdam, who, in the seventeenth century, cut himself in the linea alba above the pubes, and took out of his bladder a stone as large as a hen's egg. The stone, the knife, and the portrait of the operator, may be seen to this day in the museum at Leyden.

Peritoneum.

65. The aorta generally divides at a point about one inch and a half below the umbilicus. This point is about the level of the highest part of the crest of the ilium. A line drawn with a slight curve outwards from this point to the crural arch, rather nearer to the pubes than the ilium, gives the direction of the common iliac and external iliac arteries. About the first two inches of this line belong to the common iliac, the remainder to the external. Slight pressure readily detects the pulsation of the external iliac above the crural arch.

Division of aorta.

I have given the length of the common iliac as about two inches. As a rule, this is correct ; but it should be remembered there are frequent deviations. I have seen it any length between three-quarters of an inch and three inches and a half. These varieties may arise either from a high division of the aorta, or a low division of the common iliac, or both. It is impossible to ascertain during life what is its length in a given instance ; for there is no necessary relation between its length and the height of the stature. I have seen it very short in tall men and *vice*



*versâ*. Anatomists generally describe the left as a trifle longer than the right; but from an examination of a hundred bodies, I found their average length to be the same.

66. Mr. Abernethy, who in the year 1796 first set the example of tying the external iliac, made his incision in the line of the artery. But the easiest and safest way to reach the vessel is by an incision, recommended in the first instance by Sir Astley Cooper, and now generally adopted, beginning just on the inner side of the artery, a little above the crural arch, and continued upwards and outwards a little beyond the spine of the ilium. The same incision extended farther in the same direction would reach the common iliac.

Bony prominences.

67. The anterior superior spine of the ilium, the spine of the pubes, and the line of the crural arch are landmarks with which every surgeon should be thoroughly familiar.

Spine of ilium.

68. This spine of the ilium is the spot from which we measure the length of the lower extremity. It is a valuable landmark in determining the nature of injuries to the pelvis and the hip. The thumb easily feels this spine even in fat persons. Its position with regard to the trochanter major should be carefully examined. The best way to do this is to place the thumbs firmly on the opposite spines, and to grasp the trochanters with the fingers. Any abnormal position on one side is thus easily ascertained with the sound side as a guide.

Spine of pubes.

69. The spine of the pubes is the best guide to the abdominal ring. It cannot easily be felt by placing the finger directly over it, since it is generally covered by fat. To feel it distinctly, we should push up the skin of the scrotum and get beneath the subcutaneous fat. If there be any difficulty in finding it, abduct the thigh, and the tense tendon of the adductor longus will lead up to it.

The position of the spine of the pubes is appealed to as a means of diagnosis in doubt between inguinal and femoral hernia. The spine lies outside an inguinal, inside a femoral.



The spine of the pubes is nearly on the same horizontal line as the upper part of the trochanter major. In this line, about one full inch external to the spine, is the femoral ring. Here is the seat of stricture in a femoral hernia.

70. The line of Poupart's ligament, or the crural arch, is in most persons indicated by a slight crescent-like furrow along the skin. It corresponds with a line drawn not straight, but with a gentle curve downwards, from the spine of the ilium to the spine of the pubes. With the help of the preceding landmarks it is easy to find the exact position of the external and internal abdominal rings, and the direction of the inguinal canal.

Crural  
arch.

71. The external abdominal ring is situated immediately above and to the outer side of the spine of the pubes. It is an oval opening with the long axis directed obliquely downwards and inwards. Though its size varies a little in different persons, yet as a rule it will admit the end of the little finger, so that we can tell by examination whether it be free or otherwise. To ascertain this, the best way is to push up the thin skin of the scrotum before the finger: then, by tracking the spermatic cord, the finger readily glides over the crest of the pubes and feels the sharp margins of the ring.

Abdomi-  
nal rings.

The precise position of the internal ring is about midway between the spine of the ilium and the symphysis of the pubes, and about two-thirds of an inch above the crural arch.

72. The position of the external and internal abdominal rings being ascertained, it is plain that the direction of the inguinal canal must be obliquely downwards and inwards, and that its length in a well-formed adult male is from one and a half to two inches, according as we include the openings or not. In very young children the canal is much shorter and less oblique, the inner ring being behind the outer. With the growth of the pelvis in its transverse direction, the anterior spines of the ilia become farther

Inguinal  
canal.



apart, and thus draw the internal ring more and more to the outer side of the external.

Spermatic  
cord.

73. The spermatic cord can be felt as it emerges through the external ring, and its course can be tracked into the scrotum. The vas deferens can be distinctly felt at the back of the cord, and separated from its other component parts.

Epigastric  
artery.

74. The direction of the internal epigastric artery corresponds with a line drawn from the inner border of the internal ring up the middle of the rectus muscle, towards the chest.

In thin persons the absorbent glands which lie along the crural arch can be distinctly felt. They are usually oval, with their long axes parallel to the line of the arch.

Abdomi-  
nal viscera.

75. Now let us see how far we can make out externally the position and size of the abdominal viscera.

To make this examination with anything like success, it is desirable to relax the abdominal muscles. The man should be on his back, the head, shoulders, and thorax being well raised, to relax the recti muscles; and the thighs bent on the abdomen, to relax the several fasciæ attached to the crural arch. To induce complete relaxation, where a very careful examination is desired, chloroform should be given.

In manipulating the abdomen we should not use the tips of the fingers. This is sure to excite the contraction of the muscles. The flat hand should be gently pressed upon it, and with a kind of undulating movement.

76. It is well to bear in mind that the central tendon of the diaphragm is about the level of the lower end of the sternum or the junction of the seventh rib; that the right half of the diaphragm rises to about the level of the fifth rib—that is, about an inch below the nipple; that the left half does not rise quite so high. In tranquil breathing the diaphragm descends about half an inch.

The position of the abdominal viscera varies, to a certain extent, in different persons. In some of them, especially the stomach, their position varies in the same person at different times.



Let us take, first, the largest of the abdominal viscera—the liver.

77. The liver lies under the right hypochondrium, and passes across the middle line over the stomach into the left hypochondrium, generally speaking, as far as the left mammary line. The extent to which it can be felt below the edges of the ribs depends upon whether it is enlarged or not, and also upon the amount of flatus in the stomach and intestines. As a rule, in health its lower thin border projects about half an inch below the costal cartilages, and can be felt moving up and down with the action of the diaphragm; but it requires an educated hand to feel it. An uneducated hand would miss it altogether. That part of it, however, which crosses the middle line below the ensiform cartilage is much more accessible to the feel; here it lies immediately behind the *linea alba*, and in front of the stomach, nearly half-way down to the umbilicus. Here, therefore, is the best place to feel whether the liver be enlarged or pushed down lower than it ought to be. If it be much enlarged and much lower, even the most untutored hand could detect its edge. Liver.

Even if the edge of the liver be felt very much lower than is normal below the ribs, it does not necessarily follow that the liver is enlarged, since it may be pressed down by other causes—for instance, the habit of wearing tight stays.

To what height does the liver ascend? This can only be ascertained by careful percussion of the chest wall. The highest part of its convexity on the right side is about one inch below the nipple, or nearly on a level with the external and inferior angle of the *pectoralis major*. Posteriorly the liver comes to the surface below the base of the right lung, about the level of the tenth dorsal spine.

Roughly speaking, the upper border of the liver corresponds with the level of the tendinous centre of the diaphragm; that is, the level of the lower end of the sternum. Thus a needle thrust into the right side, between the sixth



and seventh ribs, would traverse the lung, and then go through the diaphragm into the liver.

Gall bladder.

78. The gall bladder, or rather the fundus of it, is situated, but cannot be felt, just below the edge of the liver about the ninth costal cartilage, outside the edge of the right rectus muscle.

Stomach.

79. The stomach varies in size more than any organ in the body. When empty and contracted, its plane is vertical, and it lies at the back of the abdomen, overlapped by the left lobe of the liver, and in front of the pancreas. When very full, it turns on its axis and swells up towards the front, coming close behind the wall of the abdomen, occupying most of the left hypochondrium and epigastrium, displacing the other contiguous organs, pushing in every direction, and often interfering with the action of the heart and left lung. Hence the palpitation and distressing heart symptoms in indigestion and flatulence.

The cardiac orifice of the stomach lies to the left of the middle line, just below the level of the junction of the seventh costal cartilage with the sternum.

Pylorus.

80. The pylorus lies under the liver, on the right side, near the end of the cartilage of the eighth rib; but it cannot be felt unless enlarged and hardened by disease, and even then not with certainty.

Spleen.

81. The spleen, if healthy, cannot be felt, so completely is it sheltered by the ribs. It lies on the left side, connected to the great end of the stomach, beneath the ninth, tenth, and eleventh ribs, between the axillary lines—lines, namely, drawn vertically downwards from the anterior and posterior margins of the axilla. Its upper edge is on a level with the spine of the ninth dorsal vertebra, its lower with the spine of the eleventh.

Its position and size, therefore, in health can only be recognised, and but very imperfectly, by a certain dulness on percussion. The greatest amount of dulness would be over the tenth and eleventh ribs; above this the thin edge of the lung would intervene between the spleen and the abdominal wall. If, therefore, the spleen can be distinctly



felt below the ribs, it must be enlarged. In proportion to its enlargement, so can its lower rounded border be detected below the tenth and eleventh ribs, especially when forced downwards by a deep inspiration.<sup>1</sup>

82. The pancreas lies transversely behind the stomach, and crosses the aorta and the spine about the junction of the first and second lumbar vertebræ. The proper place to feel for it, therefore, would be in the linea alba about two or three inches above the umbilicus. Is it perceptible to the touch?—only under very deep pressure, and very favourable circumstances, such as an emaciated and empty abdomen. It is worth remembering that it may be felt under such conditions. Sir W. Jenner says that he has known the pancreas of normal size, in thin persons mistaken for disease—disease of the transverse arch of the colon, or aneurism of the abdominal aorta. Pancreas.

83. The kidney lies at the back of the abdomen, on the quadratus lumborum and psoas muscles, opposite the two lower dorsal and two upper lumbar vertebræ. The right, owing to the size of the liver, is a trifle—say, three-quarters of an inch—lower than the left. The pelvis of the kidney is about the level of the spine of the first lumbar vertebra: the upper border is about the level of the space between the eleventh and twelfth dorsal spines; the lower border comes as low as the third lumbar spine. During a deep inspiration both kidneys are depressed by the diaphragm nearly half an inch. Kidney.

Can we feel a kidney of normal size? The only place where it is accessible to pressure is just below the last rib, on the outer edge of the 'erector spinæ.' I say accessible to pressure, for I have never succeeded in satisfying myself that I have distinctly felt its rounded lower border in the living subject, nor even in the dead, with the advantage of flaccid abdominal walls and the opportunity of making hard pressure with both hands, placed simultaneously, one in front of the abdomen, the other on the back. For these

<sup>1</sup> See some good observations on the position of the enlarged spleen, by Sir W. Jenner, 'Med. Gaz.,' Jan. 16, 1869.



reasons I believe that, although we can easily ascertain its degree of tenderness on pressure, we cannot actually feel it unless it be considerably enlarged.

We must be on our guard not to mistake for the kidney an enlarged liver or spleen or an accumulation of fæces in the lumbar part of the colon.

Large intestine.

84. Let us now trace the large intestine and see where it is accessible to pressure. The 'cæcum,' or 'caput coli,' and the ileo-cæcal valve lie in the right iliac fossa. The ascending colon runs up the right lumbar region over the right kidney. The transverse colon crosses the abdomen three or four inches above the umbilicus. The descending colon lies in the left lumbar region in front of the left kidney. The sigmoid flexure occupies the left iliac fossa.

Throughout the whole of this tortuous course, except at the angles of the transverse colon, the large bowel is accessible to pressure, and we could, under favourable circumstances, detect hardened fæces in it. Within the last few months we have had two instances of this in St. Bartholomew's Hospital. One was a collection of fæces in the transverse colon, which formed a distinct tumour in the abdomen. All the symptoms yielded to large and repeated injections of olive oil. The other case was an accumulation of fæcal matter in the sigmoid flexure, which during life was thought to be malignant disease.

Colotomy.

85. The operation of opening the colon (colotomy) may be done in the right or left loin, below the kidney, in that part of the colon not covered by peritoneum.

The landmarks of the operation are:—(1) The last rib, of which feel the sloping edge; (2) the crest of the ilium; (3) the outer border of the 'erector spinæ.' The incision should be about three inches long, midway between the rib and the ilium. It should begin at the outer border of the 'erector spinæ,' and run not quite transversely, but rather slope in the line of the rib. The edge of the 'quadratus lumborum,' which is the guide to the colon, is about one inch external to the edge of the 'erector spinæ,' or three full inches from the lumbar spines.



The line of the gut is vertical, and runs (for a good two inches) between the lower border of the kidney and the iliac crest.

All the room below the umbilicus is occupied by the small intestines—the coils of the jejunum lying nearer to the umbilicus, those of the ilium lower down—supposing the bladder to be empty. Small intestines.

On the right side, a little below the ninth rib, the colon lies close to the gall bladder, and is, after death, sometimes tinged with bile. Posteriorly, this part of the colon is in contact with the kidney and duodenum.

86. When the bladder distends, it gradually rises out of the pelvis into the abdomen, pushes the small intestines out of the way, and forms a swelling above the pubes, reaching in some instances up to the navel. The outline of this swelling is perceptible to the hand as well as to percussion. More than this, fluctuation can be felt through the distended bladder by tapping on it in front with the finger of one hand, while the finger of the other feels the bottom of the bladder through the rectum. Bladder.

### THE PERINEUM.

The body is supposed to be placed in the usual position for lithotomy.

87. We can readily feel the osseous and ligamentous boundaries of the perineum; namely, the rami of the pubes and ischia, the tuberosities of the ischia, the great sacro-ischiatic ligaments, and the apex of the coccyx. This framework forms a lozenge-shaped space. If we draw an imaginary line across it from the front of one tuber ischii to the other, we divide this space into an anterior and a posterior triangle. The anterior is nearly equilateral, and, in a well-formed pelvis, its sides are from three to three and a half inches long. It is called the urethral triangle. The posterior, containing the greater part of the anus and the ischio-rectal fossa on each side, is called the anal triangle. Bony framework.



Raphé.

88. A slight central ridge of skin, called the raphé, runs from the anus up the perineum, scrotum, and penis. This 'raphé,' or middle line of the perineum, is the 'line of safety' in making incisions to let out matter or effused urine, or to divide a stricture.

Central point of perineum.

89. It is very important to know that a point of the raphé about midway between the scrotum (where it joins the perineum) and the centre of the anus, corresponds with the so-called 'central tendon' where the perineal muscles meet. The bulb of the urethra lies above this point, and never, at any age, comes lower down. The artery of the bulb, too, never runs below this level. Therefore the incision in lithotomy should never commence above it. A knife introduced at this point, and pushed backwards with a very slight inclination upwards, would enter the membranous part of the urethra just in front of the prostate gland; pushed still farther it would enter the neck of the bladder. This point, then, is a very good landmark to the urethra in lithotomy, or, indeed, to any operations on the perineum.

The incision in the lateral operation for lithotomy, beginning below the point indicated, should be carried downwards and outwards between the anus and the ischium, a little nearer to the ischium than the anus.

Triangular ligament.

90. In a thin perineum, we can feel the lower border of the deep perineal fascia or the so-called triangular ligament of the urethra. The urethra passes through it about one inch below the lower part of the symphysis pubis, and about three-quarters of an inch higher than the central tendon of the perineum. It is important to bear in mind these landmarks in introducing a catheter. If the catheter be depressed too soon, its passage will be resisted by the triangular ligament; if too late, it will be likely to make a false passage by running through the bulb.

Anus.

91. One of the most important landmarks which guide a surgeon in his operations about the anus, is a white line which was first pointed out by Mr. Hilton,<sup>1</sup> and can be

<sup>1</sup> 'Lectures on Rest and Pain,' p. 280.



easily recognised in the living body. It shows the junction of the skin and mucous membrane, and is of especial interest because it marks with great precision the linear interval between the external and internal sphincter muscles. It is an exact and truthful landmark, and can be relied upon. From this line the internal sphincter extends upwards, beneath the mucous membrane, for about an inch, becoming gradually more and more attenuated.

The wrinkled appearance of the anus is caused by the contraction of the external sphincter. At the bottom of these cutaneous folds, especially towards the coccyx, we look for 'fissure of the anus.'

92. Many valuable landmarks may be felt by introducing the finger into the rectum, with a catheter at the same time in the urethra. The principal of these landmarks are the following:—

Land-  
marks in  
the rectum.

*a.* You can feel the extent and powerful grasp of the internal sphincter for about one inch up the bowel.

*b.* Through the front wall of the bowel you can most distinctly feel the track of the membranous part of the urethra, exactly in the middle line. This is very important, because you can ascertain with precision whether the catheter has deviated from the proper track.

Urethra.

*c.* About an inch and a half or two inches from the anus, the finger comes upon the prostate gland. The gland lies in close contact with the bowel, and can be detected by its shape and hard feel. The finger, moved from side to side, can examine the size of its lateral lobes.

Prostate  
gland.

*d.* The finger, introduced still farther, can reach beyond the prostate, as far as the apex of the trigone of the bladder. More than this, it can feel the angle between the ductus communes ejaculatorii, which forms the apex of the trigone. This is the precise spot where the distended bladder should be punctured through the rectum. The more distended the bladder, the easier can this spot be felt. Fluctuation is at once detected by a gentle tap on the bladder above the pubes. The trochar must be thrust in the direction of the axis of the distended bladder; that is,



roughly speaking, in a line drawn from the anus through the pelvis to the umbilicus.

*e.* The fold of peritoneum, called the recto-vesical pouch, is about four inches from the anus, therefore it is not within reach of the finger; and we run no risk of wounding it in tapping the bladder if the trochar be introduced near the angle of the trigone.

*f.* The finger can feel one of the ridges or folds of mucous membrane which are situated at the lower part of the rectum. This fold projects from the side, and sometimes from the upper part of the rectum, near the prostate. When thickened or ulcerated, this fold occasions great pain in defæcation; and great relief is afforded by its division.

*g.* Lastly, the finger can examine the condition of the spaces filled with fat on either side of the rectum, called the ischio-rectal fossæ, with a view to ascertain the existence of deep-seated collections of matter, or the internal communications of fistulæ.

In the introduction of catheters it is a good rule to keep the point of the instrument well applied against the upper part of the urethra. By so doing one is much less likely to make a false passage.

Urethra in  
the child.

In children the membranous part of the urethra is, relatively speaking, very long, owing to the smallness of the prostate. It is also more sharply curved, because the bladder in children is more in the abdomen than in the pelvis. It is, moreover, composed of thin and delicate walls. No violence, therefore, should be used in passing a catheter; else the instrument is likely to pass through the coats and make a false passage. Hence the advantage of being able to ascertain through the rectum whether the instrument be in the right track and moving freely in the bladder, which can also be easily felt in children.



### THE THIGH.

93. First mark the anterior superior spine of the ilium, the spine of the pubes, and define the line of the crural arch which extends between them. This line is one of our guides in the diagnosis of inguinal and femoral herniæ. If the bulk of the tumour be above the line, the hernia is probably inguinal ; if below it, femoral. The line is not a straight one drawn from the spine of the ilium to the spine of the pubes, but slightly curved, with convexity downwards, owing to its close connection with the fascia lata of the thigh. In many persons it can be distinctly felt ; in nearly all its precise course is indicated by a slight furrow in the skin.

Crural  
arch.

94. When the thigh is even slightly bent, there appears a second furrow in the skin below that at the crural arch. This second furrow begins at the angle between the scrotum and the thigh, passes outwards, and is gradually lost between the top of the trochanter and the anterior superior spine of the ilium. It runs right across the front of the capsule of the hip joint. For this reason it is a valuable landmark in amputation of the hip. The point of the knife should be introduced externally where the furrow begins, should run precisely along the line of it, and come out where it ends ; so that the capsule of the joint may be opened with the first flap. In suspected disease of the hip pressure made in this line, just below the spine of the ilium, will tell us if the joint be tender. Effusion into the joint obliterates all trace of the furrow, and makes a fulness when contrasted with the opposite groin.

Cutaneous  
furrow at  
the bend  
of the  
thigh.

95. In most persons there is a natural depression over the saphenous opening in the fascia lata, where the saphena vein joins the femoral. The position of this opening is just below the inner third of the crural arch, and about an inch and a half external to the spine of the pubes. This is the

Saphenous  
opening.



place where the swelling of a femoral hernia first appears, and ought to be carefully examined in cases of suspicion.

Femoral  
ring.

96. The position of the femoral ring, through which the hernia escapes from the abdomen, is, on a deeper plane, about half an inch higher than the saphenous opening, and immediately under the crural arch. As the plane of the ring is vertical in the supine position of the body, the way in which we should try to reduce a femoral hernia is by pressure, applied first in a downward direction, afterwards in an upward. The intestine protruded has to pass, so to speak, under the edge of a sharp angle, the angle being the upper horn of the saphenous opening. At the same time we bend the thigh, to relax the fascia as much as possible.

A good way to find the seat of the femoral ring with precision is the following:—Feel for the pulsation of the femoral artery on the pubes; allow half an inch for the femoral vein; then comes the femoral ring.

In performing the operation for the relief of the stricture, the incision through the skin should be about an inch and a half external to the spine of the pubes. Its direction should be vertical, and its middle should be just over the femoral ring.

Absorbent  
glands in  
the groin.

97. The cluster of inguinal and femoral absorbent glands can sometimes be felt in thin persons. The inguinal lie for the most part along the line of the crural arch: they receive the absorbents from the wall of the abdomen, the urethra, the penis, the scrotum, and the anus. The femoral glands lie chiefly over the saphenous opening and along the outer side of the saphena vein: they receive the absorbents of the lower extremity; they receive some also from the scrotum—of which we have practical evidence in cases of chimney-sweepers' cancer.

Trochan-  
ter major.

98. The trochanter major is a most valuable landmark, to which we are continually appealing in injuries and diseases of the lower extremity. There is a natural depression over the hip where it lies very near the surface, and can be plainly felt, especially when the thigh is rotated. Nothing intervenes between the bone and the



skin except the strong fascia of the glutæus maximus and the great bursa underneath it.

The top of the trochanter lies pretty nearly on a level with the spine of the pubes, and is about three-fourths of an inch lower than the top of the head of the femur. A careful examination of the bearing of the great trochanter to the other bony prominences of the pelvis, and a comparison of its relative position with that of the opposite side, are the best guides in the diagnosis of injuries about the hip, and the position of the head of the femur.

99. 'If in the normal state you examine the relations of the great trochanter to the other bony prominences of the pelvis, you will find that the top of the trochanter corresponds to a line drawn from the anterior superior spine of the ilium to the most prominent part of the tuberosity of the ischium. This line also runs through the centre of the acetabulum. The extent of displacement in dislocation or fracture is marked by the projection of the trochanter behind this line.'<sup>1</sup>

Nélaton's  
line.

'Nélaton's line,' as it is termed, is theoretically good. But in stout persons it is not always easy to feel these bony points so as to draw the line with precision. A surgeon must, after all, in many cases trust to measurement by his hands. Thus, let the thumbs be placed firmly on the spines of the ilia, while the fingers grasp the trochanters on each side. Having the sound side as a standard of comparison, the hand will easily detect any displacement on the injured side. Winslow's<sup>2</sup> observation is very much to the point—'Feel whether the injured member answers to the sound.'

The top of the great trochanter is the guide in an operation recently introduced by Mr. Adams, the 'subcutaneous section of the neck of the femur.' The puncture should be made one inch above and one inch in front of the top of the trochanter. The neck of the bone is to be sawn through at right angles to its axis, the saw working in the line of the crural arch.

<sup>1</sup> Nélaton, '*Pathologie chirurgicale*,' t. iv. p. 441. 1848.

<sup>2</sup> Winslow, '*Structure of the Human Body*.' London, 1873.



The spine of the ilium is the point from which we measure the length of the lower limb. By looking at the spines of opposite sides we can detect any slant in the pelvis. By pressure on both spines simultaneously we examine if there be a fracture of the pelvis, or disease at the sacro-iliac joint.

100. 'In reducing a dislocation of the hip by manipulation it is important to bear in mind that, in every position, the head of the femur faces nearly in the direction of the inner aspect of its internal condyle.'<sup>1</sup>

Compres-  
sion of  
femoral  
artery.

101. About a point midway between the spine of the ilium and the symphysis pubis, the femoral artery can be felt beating, and effectually compressed, against the pubes. How should the pressure be applied when the patient lies on the back? In accordance with the slope of the bone—that is, with a slight inclination upwards.

If the Italian tourniquet be used, we should be careful to adjust the counter-pad well under the tuberosity of the ischium. If digital pressure be used, it is easy to command the femoral by slight pressure of the thumb, provided the fingers have a hold on the great trochanter.

Sartorius.

102. The sartorius is the great fleshy landmark of the thigh, as the biceps is of the arm, and the sterno-cleido-mastoideus of the neck. Its direction and borders may easily be traced by asking the patient to raise his leg, a movement which puts the muscle in action. The same action defines the boundaries of the triangle (of Scarpa) formed by the adductor longus and sartorius.

To define the course of the femoral artery, draw a line from midway between the spine of the ilium and the symphysis pubis to the inner side of the knee. This line, observe, applies to the femoral artery, not to the popliteal.

The sartorius begins to cross the artery, as a rule, from three to four inches below the crural arch. The point at which the profunda artery arises is about one and a half or two inches below the arch. Therefore the incision for

<sup>1</sup> Bigelow, 'Mechanism of Dislocation and Fracture of the Hip,' Philadelphia, 1869.



tying the femoral in Scarpa's triangle should commence about a hand's breadth below the crural arch and be continued for three inches in the line of the artery.

To command the femoral in Scarpa's triangle, the pad of the tourniquet should be placed at the apex, and the direction of the pressure should be, not backwards, but outwards, so that the artery may be compressed against the femur.

In the middle third of the thigh the femoral artery lies in Hunter's canal, overlapped by the sartorius. About the commencement of the lower third the artery leaves the canal through the oval opening in the triceps, and enters the popliteal space. The easiest way to reach the artery in Hunter's canal is by an incision along the outer border of the sartorius. Now, this outer border corresponds with a line drawn from the tuberosity of the inner condyle straight up to the middle third of the thigh.

To command the femoral artery in Hunter's canal, the pressure should be directed outwardly, so as to press the vessel against the bone.

103. The bony landmarks of the buttocks which can be distinctly felt are : 1, the posterior superior spines of the ilia ; 2, the spines of the sacral vertebræ ; 3, the two tubercles of the last sacral vertebra ; 4, the apex of the coccyx in the deep groove leading to the anus ; 5, the tuberosities of the ischia on each side of the anus.

Buttocks.

Bony landmarks.

The posterior spines of the ilia are about the level of the second sacral spine, and correspond with the middle of the sacro-iliac symphysis.

The third sacral spine marks the lowest part to which the cerebro-spinal fluid descends in the spinal canal.

The tuberosities of the ischia, in the erect position, are covered by the glutæus maximus. In the sitting position they support the weight of the body, and are only covered by a thick pad of coarse fat. Between this pad and the bones there is a bursa, which I have more than once seen enlarged and inflamed in coachmen.



The prominence of the nates is one of the characteristics of man in connection with his erect attitude. 'Les fesses n'appartiennent qu'à l'espèce humaine.' They are formed of an accumulation of fat over the great muscle of the buttock. From their appearance we may gather some indication of the state of the constitution. They are firm and globose in the vigorous; loose and flaccid in the infirm. Wasting and flattening of one, compared with the other, is an early symptom of disease in the hip.

Fold of  
the but-  
tock.

104. The deep furrow, termed 'the fold of the buttock,' which separates the nates from the back of the thigh, corresponds with the lower border of the glutæus maximus. Its altered direction in disease of the hip is very characteristic. This is the best place to feel the great ischiatic nerve. We find it by pressing deep between the trochanter and the tuber ischii, rather nearer to the latter. When we sit upright, the nerve is not liable to pressure; but it becomes numbed when we sit long sideways.

Glutæal  
artery.

105. To find at what point the glutæal artery comes out of the pelvis, draw a line from the posterior superior spine of the ilium to the trochanter major, rotated inwards. The junction of the upper with the middle third of this line lies over the artery as it emerges from the upper border of the great ischiatic notch.

The point at which the ischiatic artery leaves the pelvis is about half an inch lower than that of the glutæal.

Pudic  
artery.

106. The pudic artery lies over the spine of the ischium. To find it, draw a line from the outer side of the tuber ischii to the posterior superior spine of the ilium. The junction of the lower with the middle third gives the position of the artery. The ischiatic artery lies close to it, but nearer the middle line.

Looking at the course of these arteries it appears that when seated on hard seats the pressure is sustained by the bones; when reclining on soft seats the pressure is sustained more by the soft parts, and reaches the arteries; hence the tendency of modern modes of reposing to drive the blood into the interior of the pelvis and favour the pro-



duction of piles and uterine disorders. A celebrated French accoucheur used to say that the habit of high waists, tight lacing, and easy chairs brought him many thousands a year.

## THE KNEE.

107. The patella ; the tuberosities of the two condyles ; the tubercle of the tibia for the attachment of the ligamentum patellæ ; another (the lateral) tubercle, on the outer side of the head of the tibia ; and the head of the fibula are the chief bony landmarks of the knee. Bony points.

Observe that the head of the fibula lies at the outer and back part of the tibia, and that it is pretty nearly on a level with the tubercle for the attachment of the ligamentum patellæ.

We can also feel the little spur-like projection of bone above the internal condyle which gives attachment to the tendon of the adductor magnus. This spur-like projection corresponds with the level of the epiphysis of the lower end of the femur, and also with the level of the highest part of the trochlea for the patella : facts worth notice in performing excision of the knee.

In reducing a dislocation of the hip, it is important to bear in mind that the inner aspect of the internal condyle in every position of the limb faces nearly in the direction of the head of the femur. (100.)

The external (lateral) tubercle of the tibia is for the attachment of the broad and strong aponeurosis (tendon of the tensor fasciæ), which acts like a brace, so well seen in emaciated persons down the outer side of the thigh, for the lateral support of the pelvis. This tubercle indicates the level to which the condyles of the femur descend, and the lower level of the synovial membrane.

The patella, in extension of the knee, is nearly all of it above the condyles : in flexion, it lies in the intercondyloid fossa, and thus protects the joint in kneeling. Its inner border is thicker and more prominent than the outer, which



slopes down towards its condyle. Hence, probably, the greater disposition of the patella to dislocation externally.

Ligamentum patellæ.

108. The line of the ligamentum patellæ is vertical. Hence any deviation from this line, one way or the other, indicates more or less dislocation of the tibia. There is a pellet of fat under the ligament, which answers a 'packing' purpose—sinking in when the knee is bent; rising when the knee is extended, and bulging on either side of the tendon, almost enough to give the feel of fluctuation.

In a well-formed leg the ligamentum patellæ, the tubercle of the tibia, and the middle of the ankle should be in the same line. A useful point in the adjustment of fractures.

Patellar bursa.

109. The patellar or housemaids' bursa is situated not only over the patella, but over the upper part of the ligament. This is plain enough when the bursa becomes enlarged. There is another subcutaneous bursa over the insertion of the ligament into the tubercle of the tibia. This is quite independent of the deep bursa between the tendon and the bone.

Synovial membrane of knee.

110. The synovial membrane of the knee, when the joint is extended, rises like a cul-de-sac above the upper border of the patella about two inches. It ascends, too, a little higher under the vastus internus than the vastus externus—a fact very manifest when the joint is distended. When the knee is bent this cul-de-sac is drawn down—hence the rule of bending the knee in operations near the lower end of the femur.

The lower level of the synovial membrane of the knee is just above the level of the upper part of the head of the fibula. The tibio-fibular synovial membrane is, as a rule to which there are few exceptions, independent of that of the knee.

Popliteal tendons.

111. The tendons forming the boundaries of the popliteal space can be distinctly felt in flexion of the knee. On the outer side, we have the biceps running down to the head of the fibula. On the inner side, we feel three ten-



dons, disposed as follows:—nearest to the middle of the popliteal space is the semitendinosus, very salient and traceable high up the thigh; next comes the thick round tendon of the semimembranosus; still more internally is the gracilis. The sartorius, which forms a graceful muscular prominence on the inner side of the knee, does not become tendinous until it gets below it.

112. The precise position of the bursa in the popliteal space, which sometimes enlarges to the size of a hen's egg, is between the tendon of the inner head of the gastrocnemius and the tendon of the semimembranosus, just where they rub one against the other. The bursa is from one and a half to two inches long. When enlarged, it makes a swelling on the inner side of the popliteal space, which bulges and becomes tense when the knee is extended, and vice versâ. I examined 150 bodies with a view to ascertain how often this bursa communicates with the synovial membrane of the knee. There was a communication about once in five instances. This should make us cautious in interfering too roughly with the bursa when enlarged.

Popliteal  
bursa.

The hollow of the popliteal space is nearly obliterated when the knee is extended. This is caused by the pressure of the condyles upon the two heads of the gastrocnemius, and the fat in the space.

113. The popliteal artery can be felt beating and can be compressed against the back of the femur, close to which it lies. But pressure, sufficient to stop the blood, should be firm, and should be made against the bone nearer to the inner than to the outer hamstrings. The line of the artery corresponds with the middle of the ham. It lies under cover of the fleshy belly of the semimembranosus, and the outer border of this muscle is the guide to it. An incision down the middle of the ham would fall in with the vessel just above the condyles.

Popliteal  
artery.

114. The peroneal nerve runs parallel with and close to the inner border of the tendon of the biceps. It can be felt

Peroneal  
nerve.



in thin persons. There is a risk of dividing it in tenotomy of the biceps, unless the knife be carefully introduced from within outwards. Below the knee the nerve can be felt close to the fibula just below the head.

### *THE LEG AND ANKLE.*

Bony  
points.

115. The tubercle of the tibia, the sharp front edge called the shin, and the broad, flat surface of the bone covered only by periosteum, can be felt all the way down. The inner edge can be felt too, but not so plainly. The lower third is the narrowest part of the bone and the most frequent seat of fracture.

The head of the fibula is a good landmark on the outer side of the leg, about one inch below the top of the tibia, and nearly on a level with the tubercle. Observe that it is placed well back, and that it forms no part of the knee joint, and takes no share in supporting the weight.

The shaft of the fibula arches backwards, the reverse of the shaft of the tibia. The fact of the bones not being on the same plane should be remembered in flap amputations. The shaft of the fibula is so buried amongst the muscles, that the only part to be distinctly felt is the lower fourth. Here there is a flat surface, subcutaneous, between the peroneus tertius in front, and the two peronei behind. Here is the most frequent seat of fracture.

Malleoli.

116. The shape and relative position of the malleoli should be carefully studied, as the great landmarks of the ankle. The internal malleolus does not descend so low as the outer, and advances more to the front: at the same time, owing to its greater antero-posterior depth, it is on the same plane as the outer behind. The lower border of the inner malleolus is somewhat rounded, and the slight notch in it for the attachment of the lateral ligament can be felt. The outer malleolus descends lower than the inner, for the more effectual locking of the joint on the



outside. Its shape is not unlike the head of a serpent. Viewed in profile, it lies just in the middle of the joint.

In 'Syme's' amputation of the foot at the ankle, the line of the incision should run from the apex of the outer malleolus, under the sole to the centre of the inner.

In a well-formed leg, the inner edge of the patella, the inner ankle, and the inner side of the great toe should be in the same line. Look to these landmarks in adjusting a fracture or dislocation, keeping at the same time an eye upon the conformation of the opposite limb.

There are several strong tendons to be seen and felt about the ankle.

117. Behind is the tendo Achillis. It forms a high relief, with a shallow gutter on each side of it. The narrowest part of the tendon, where it should be divided in tenotomy, is about the level of the inner ankle; below this it expands again to be attached to the lower and back part of the os calcis. Seen in profile, the tendon is not straight, but slightly concave—being drawn in by an aponeurosis which forms a sort of girdle round it. This girdle proceeds from the posterior ligament of the ankle; and, though most of its fibres encircle the tendon, some of them adhere to and draw in its sides. All this disappears when the tendon is laid bare by dissection.

Tendo  
Achillis.

118. Above and behind the malleolus internus we can feel the broad flat tendons of the tibialis posticus and the flexor longus digitorum. The tendon of the tibialis posticus lies nearest to the bone and comes well up in relief in adduction of the foot. It lies close to, and parallel with, the inner edge of the tibia, so that this edge is the best guide to it. When the body is recumbent the plane of the tendon is vertical; therefore in tenotomy the knife should be introduced first perpendicularly between the tendon and the bone, and then turned at right angles to cut the tendon. The tendon has a separate sheath and synovial membrane, which commences about one inch and a half above the apex of the malleolus and is continued to its insertion into the tubercle of the scaphoid bone. The proper place, then, for

Tendons  
behind in-  
ner ankle.



division of the tendon is about two inches above the end of the malleolus.

If the patient were young and fat, so that the inner edge of the tibia could not be distinctly felt, then the best landmark to find the tendon would be a point midway between the front and the back of the leg. An incision in front of this point might injure the internal saphena vein ; behind this point, the posterior tibial artery.

Tendons  
behind  
outer  
ankle.

119. Behind the malleolus externus we feel the two peroneal tendons. They lie close to the edge of the fibula, the short one nearer to the bone. In dividing these tendons, the knife should be introduced perpendicular to the surface, and about two inches above the apex of the ankle, so as to be above the synovial sheaths of the tendons.

Tendons in  
front of  
ankle.

Over the front of the ankle, when the muscles are in action, we can see and feel, beginning on the inner side, the tendons of the tibialis anticus, the extensor longus pollicis, the extensor longus digitorum, and the peroneus tertius. They start up like cords when the foot is raised, and are kept in their proper relative position by strong pulleys formed by the anterior annular ligament. Of these pulleys the strongest is that of the flexor communis digitorum. When the ankle is sprained, the pain and swelling arise from a stretching of these pulleys and effusion into their synovial sheaths. A laceration of one of the pulleys and escape of the tendon are extremely rare.

The place for the division of the tendon of the tibialis anticus, so as not to injure its synovial sheath, is about one inch before its insertion into the cuneiform bone. The knife should be introduced on the outer side, so as to avoid the dorsal artery of the foot.

Now trace the lines of the arteries, and the landmarks near which they divide.

Popliteal  
artery.

120. About one inch and a quarter below the head of the fibula, or say one inch below the tubercle of the tibia, the popliteal artery divides into the anterior and posterior tibial. The peroneal comes off from the posterior tibial about three inches below the head of the fibula.



Consequently we may lay down, as a general rule, that, in amputations one inch below the head of the fibula, only one main artery, the popliteal, is divided. In amputations two inches below the head of the fibula, two main arteries, the anterior and posterior tibial, are divided. In amputations three inches below the head, three main arteries, the two tibials and the peroneal, are divided.

121. The anterior tibial artery comes in front of the interosseous membrane, one inch and a quarter below the head of the fibula, and here lies close to this bone. Its subsequent course is defined by a line drawn from the front of the head of the fibula to the middle of the front of the ankle. This line corresponds pretty nearly with the outer border of the tibialis anticus all the way down. Therefore, if this muscle be put in action, the incision for the ligature of the artery in any part of its course may be defined with the greatest precision. The artery can be felt beating and compressed where it crosses the front of the tibia and ankle.

Anterior  
tibial  
artery.

122. The posterior tibial commences about one inch and a quarter below the head of the fibula. Its subsequent course corresponds with a line drawn from the middle of the upper part of the calf to the hollow behind the inner ankle, where it can be felt beating distinctly about half an inch from the edge of the tibia. A vertical incision down the middle of the calf would reach the artery under cover of the gastro-cnemius and soleus. A vertical incision along the middle third of the leg, about half an inch from the inner edge of the tibia, would enable the operator to reach the artery sideways, by detaching from the bone the tibial origin of the soleus.

Posterior  
tibial  
artery.

123. The internal saphena vein can be always plainly seen over the front of the inner ankle. Its farther course up the inner side of the leg, knee, and thigh to its termination in the femoral is not in all persons manifest.

Saphena  
veins.

The external saphena vein runs behind the outer ankle and up the middle of the calf to empty itself in the popliteal vein.



## THE FOOT.

What are the bony landmarks which guide us in the surgery of the foot?

Points of  
bone.

124. Along the inner side of the foot, beginning from behind, we can feel—1, the tuberosity of the os calcis; 2, the projection of the internal malleolus; 3, the projection of the os calcis, termed ‘sustentaculum tali,’ about one full inch below the malleolus; 4, about one inch in front of the malleolus internus, and a little lower, is the tubercle of the scaphoid bone; the gap between it and the sustentaculum tali being filled by the calcaneo-scaphoid ligament and the tendon of the tibialis posticus, in which there is often a sesamoid bone; 5, the internal cuneiform bone; 6, the projection of the first metatarsal bone; 7, the sesamoid bones of the great toe.

Along the outer side of the foot we can feel—1, the external tuberosity of the os calcis; 2, the external malleolus; 3, the peroneal tubercle of the os calcis, one inch below the malleolus, with the long peroneal tendon below it and the short one above it; 4, the projection of the fifth metatarsal bone.

Lines of  
joints.

125. In fat persons the following rules for finding the joints may be of service as regards the surgery of the foot:—

The level of the ankle joint lies about half an inch above the end of the inner malleolus. This is worth remembering in performing ‘Syme’s’ amputation.

The tubercle of the scaphoid bone is the best guide to the astragalo-scaphoid joint which lies immediately behind it; and the plane of this joint is in the same line as that of the calcaneo-cuboid. Thus a line drawn transversely over the dorsum of the foot, behind the tubercle of the scaphoid, would strike both the joints opened in ‘Chopart’s’ operation.

Place your thumb on the tubercle of the scaphoid, and



measure one inch in front : here you find the joint between the internal cuneiform bone and the metatarsal bone of the great toe.

The line of the calcaneo-cuboid joint lies midway between the external malleolus and the end of the metatarsal bone of the little toe.

The projection of the fifth metatarsal bone is the guide to the joint between it and the cuboid.

Notice that the line of the joints between the metatarsal bones and the first phalanges lies a full inch farther back than the interdigital folds of the skin. This is a point to be remembered in amputating the toes.

126. The line of the dorsal artery of the foot corresponds with one drawn from the middle of the ankle to the interval between the first and second metatarsal bones. The artery can be felt beating over the bones along the outer side of the extensor longus pollicis, which is the best guide to it. Dorsal artery.

127. The synovial sheath of the extensor longus pollicis extends from the front of the ankle, over the instep, as far as the metatarsal bone of the great toe. There is generally a bursa over the instep, above, or it may be, below, the tendon. Bursa.

There is often a large irregular bursa between the tendons of the extensor longus digitorum, and the projecting end of the astragalus, over which the tendons play. There is much friction here. It is well to be aware that this bursa sometimes communicates with the joint of the head of the astragalus.

128. The course of the external plantar artery corresponds with a line drawn from the hollow behind the inner ankle obliquely across the sole nearly to the base of the fifth metatarsal bone : from thence the artery turns transversely across the foot, lying deeply near the bases of the metatarsal bones, till it inosculates with the dorsal artery of the foot in the first interosseous space. Plantar arteries.

The lines of the digital arteries correspond with the lines of the metatarsal bones.



The course of the internal plantar corresponds with a line drawn from the inner side of the os calcis to the middle of the great toe.

Plantar  
fascia.

129. To divide the plantar fascia subcutaneously, the best place is about one inch in front of its attachment to the os calcis. This is the narrowest part of it. The knife should be introduced on the inner side; and the incision will be behind the plantar artery.

The subcutaneous section of the tendon of the abductor pollicis should be made about one inch before its insertion.

### THE ARM.

Clavicle.

130. The line of the clavicle can always be felt, even in the fattest persons. Its direction is not perfectly horizontal, but slightly inclined downwards, when the arm hangs quietly by the side. When the body lies flat on the back, the shoulder not only falls back, but rises a little, the weight of the limb being taken off. Hence the modern practice of treating fractures of the clavicle by the supine position, in the early stage.

On the front surface of the clavicle, not far from its acromial end, there is in many persons of mature age a spine-like projection of bone. So far as I know, it has not been described. A gentleman, himself a surgeon, showed me an instance in his own person. He suspected it was an exostosis.

As a rule the acromio-clavicular joint forms an even plane. But there is sometimes a knob of bone at the acromial end of the clavicle; or it may be only a thickening of the fibro-cartilage, sometimes existing in the joint. In either case this relief might be mistaken for a dislocation, or even for a fracture. A reference to the other shoulder would settle the doubt.

Bony  
points of  
the shoul-  
der.

131. We can distinctly feel the spine of the scapula and the acromion, more especially at the angle where they join behind the shoulder. This angle is the best place from



which to measure in taking the comparative length of the arms.

In some shoulders, though not often, there is a natural symphysis between the spine of the scapula and the acromion. There may indeed be two symphyses and two acromial bones, in which case the acromion has two points of ossification. These natural symphyses might be mistaken for fractures, until we have examined the opposite shoulder, which is sure to present a similar conformation.<sup>1</sup>

Outside the acromion, the arm hanging by the side, we can feel, through the fibres of the deltoid, the upper part of the humerus. It distinctly moves under the hand when the arm is rotated. It is not the head of the bone which is felt, but the tuberosities, the greater externally, the lesser in front. These form the convexity of the shoulder. When the arm is raised, this convexity disappears; there is a slight depression in its place; and the head of the bone can be felt by pressing the fingers high up in the axilla. Tuberosities,

The absence of this prominence formed by the upper part of the humerus under the deltoid, and the presence of a prominence low in the hollow of the axilla, or in front, below the coracoid process, or behind, on the back of the scapula, bespeak dislocation of the head of the bone.

In examining obscure injuries about the shoulder, it is worth remembering that, in the normal relation of the bones, and in every position, the great tuberosity faces in the direction of the external condyle. The head of the bone faces very much in the direction of the internal condyle.

By making deep pressure in front of the shoulder, when the arm is pendent and supine, we can distinguish the bicipital groove. It looks directly forwards, and runs in a line drawn vertically downwards through the middle of the biceps to its tendon at the elbow. We should be aware of this, lest it be mistaken for a fracture.

<sup>1</sup> See Pruge on 'Ossa Acromialia' ('Zeitschrift für rationelle Mediz.'), 3. Reihe, Bd. vii. 1859.



Coraco-  
acromial  
ligament.

132. Under the anterior fibres of the deltoid, we can distinctly feel the position and extent of the coraco-acromial ligament. A knife, passed vertically through the middle of it, goes at once into the shoulder joint and strikes the bicipital groove with the tendon—a point to be remembered in excision.

In persons of an athletic build the triangular form and beautiful structure of the deltoid become conspicuous when the muscle is in action. The depression on the outer side of the arm, indicating its insertion, is the place selected for issues or setons.

The arm being held by an assistant, the anterior and posterior borders of the relaxed deltoid admit of being raised so that the amputating knife can be introduced beneath the muscle for the purpose of making the flap.

Axilla.

133. The anterior border of the axilla, formed by the pectoralis major, follows the line of the fifth rib. In counting the ribs, or in tapping the chest, it is worth remembering that the first visible digitation of the serratus magnus is attached to the sixth rib. The angle of the digitation is directed forwards and corresponds to the upper edge of the rib. The second visible digitation corresponds to the seventh rib; the interval between these digitations, therefore, corresponds to the sixth intercostal space—a convenient place for tapping the chest. (38.)

In the normal state no glands can be felt in the axilla.

Axillary  
artery.

134. When the arm is raised to a right angle, and the head of the humerus thereby depressed, the axillary artery is plainly felt beating, and can be perfectly commanded by pressure made on the inner side of the coraco-brachialis. This muscle stands out in relief along the humeral side of the axilla, and is the best guide to the artery. A line drawn along its inner border—that is, down the middle of the axilla—corresponds with the course of the artery.

The depth and form of the axilla alter in different positions of the arm. In the arm raised and abducted the axilla becomes nearly flat; hence this position is always adopted in operations.



In opening abscesses in the axilla, the incision should be made midway between the borders, and the point of the knife introduced from above downwards.

135. When the arm is extended and supinated, a line drawn from the deepest part of the middle of the axilla down the inner side of the biceps to the middle of the bend of the elbow, corresponds with the course of the brachial artery. The artery can be felt and compressed all the way down ; but nowhere so effectually as midway, where it lies for two inches on the tendon of the coraco-brachialis on the inner flat side of the humerus. The proper direction to apply the pressure is outwards and a little backwards, else the artery is apt to slip off the bone.

Brachial  
artery.

The musculo-spiral nerve and superior profunda artery wind round the back of the humerus about its middle, and come to the front of the external condyloid ridge. Thus, for full three inches above the condyles, there is nothing to interfere with operations on the back of the bone, which is here broad and flat.

136. At the bend of the elbow, the tendon of the biceps can be plainly felt, as well as the pulsation of the brachial artery close to its inner side, before dividing into the radial and ulnar.

Bend of  
elbow.

The bend of the elbow in young children and in persons with fat and round arms presents a semicircular fold of which the curve embraces the lower part of the biceps ; but in muscular persons we see the distinct boundaries of the triangular space, formed by the pronator teres on the inner side, and the supinator longus on the outer. Here can be traced, standing out in strong relief under the thin white skin, the superficial veins, which, in days gone by when bleeding was the fashion, were of such great importance. Their arrangement, although subject to variety, is very much like the branches of the letter M, the middle of the M being at the middle of the elbow. Of these branches the median basilic, which runs over the tendon of the biceps, is the largest and most conspicuous, and is generally selected for venesection ; it crosses the course of the brachial

Cutaneous  
veins.



artery, nothing intervening but the semilunar aponeurosis from the tendon of the biceps.

Land-  
marks of  
elbow.

137. It is of great importance to be familiar with the relative positions of the various bony prominences about the elbow. We can always feel the internal and external condyles. The internal is the more prominent of the two, and a trifle higher.

Olecranon.

We can always feel the olecranon. This is somewhat nearer to the inner than to the outer condyle. Between the olecranon and the internal condyle is a deep impression in which lies the ulnar nerve, commonly called the 'funny bone.'

On the outer side of the olecranon, just below the external condyle, is a pit in the skin, constant even in fat persons. This pit is considered one of the beauties of the elbow in a graceful arm; it is seen in a child as a pretty little dimple. To the surgeon it is most interesting, as in this valley behind the supinator longus and the radial extensors of the wrist he can distinctly feel the head of the radius rolling in pronation and supination of the arm. It is, therefore, one of the most important landmarks of the elbow, since it enables us to say whether the head of the radius be or be not in its right place.

Relations  
of olecra-  
non and  
condyles.

To examine the relative positions of the olecranon and condyles in the different motions of the elbow joint, place the thumb on one condyle, the tip of the middle finger on the other, and the tip of the forefinger on the olecranon. In extension, the highest point of the olecranon is never above the line of the condyles; indeed, it is just in this line. With the elbow at right angles the point of the olecranon is just vertically below the line of the condyles. In extreme flexion the point of the olecranon lies in front of the line of the condyles.

All these relative positions would be altered in a dislocation of the ulna, but not in a fracture of the lower end of the humerus.

Sometimes, though rarely, we meet with a hook-like projection of bone above the internal condyle. It is called



a 'supra-condyloid' process; it can be felt through the skin, with its concavity downwards, and is a rudiment of the bony canal which, in many mammalia, transmits the median nerve and ulnar artery. A third origin of the pronator teres is always attached to it; this origin covers the brachial artery.<sup>1</sup>

The subcutaneous bursa over the olecranon, if dis- Bursæ.  
tended, would be as large as a walnut. A second bursa sometimes exists a little lower down upon the ulna. There is also a small subcutaneous bursa over each of the condyles.

The vertical extent of the elbow joint is limited, above by a line drawn from one condyle to the other; below by a line corresponding to the lowest part of the head of the radius.

138. About one inch below the head of the radius, the Interos-  
seous  
arteries.  
ulnar artery gives off the common interosseous; and this divides, about half an inch lower, into the anterior and posterior interosseous. Thus, in amputating the forearm, say two inches below the head of the radius, four arteries, at least, would require ligature.

By flexion of the elbow to the utmost, the circulation through the brachial artery can be arrested; but the position is painful, and can be tolerated only for a short time.

There is a small absorbent gland just above the inner Absorbent  
gland.  
condyle, in front of the intermuscular septum. It is the first to take alarm in poisoned wounds of the hand.

## THE FOREARM AND WRIST.

139. The edge of the ulna can be felt subcutaneous from Ulna.  
the olecranon to the styloid process; and any fracture could be easily detected. The styloid process of the ulna does not descend so low as the styloid process of the radius, in order

<sup>1</sup> See on this subject a monograph, 'Canalis Supra-Condylloideus Humeri.' By W. Grüber. Petersburg, 1856.



not to impede the free abduction of the hand. Its apex is on a level with the radio-carpal joint. The head of the ulna is plainly felt and seen projecting at the back of the wrist, especially in pronation of the forearm. It then lies between the tendons of the extensor carpi ulnaris and extensor minimi digiti. There is often a subcutaneous bursa over it.

Radius.

140. The upper half of the shaft of the radius is so covered by muscles that we cannot feel it; the lower half is more accessible to the touch, especially just above and below the part where it is crossed by the extensors of the thumb. Its styloid process is readily felt, and made all the more manifest by being covered by the first two extensor tendons of the thumb. It descends lower, and lies more to the front than the corresponding process of the ulna. The relative positions of these styloid processes can be best examined by placing the thumb on one and the forefinger on the other.

Feel for the little bony pulley on the back of the radius which keeps in place the third extensor tendon of the thumb. This and the bone just above it is the place which we examine for suspected fracture (termed Colles's) above the wrist.

Carpus.

141. Below the styloid process of the radius, just on the inner side of the extensors of the thumb, we feel the tubercle of the scaphoid bone.

Between the two is the level of the radio-carpal joint. A little lower we feel the trapezium.

Below the styloid process of the ulna we feel, on the front of the wrist, the pisiform bone; and, on the inner side, the cuneiform.

When the anterior annular ligament is divided for the relief of a diseased carpal bursa, the incision should be made vertically, in the middle of the wrist, in the furrow between the ball of the thumb and that of the little finger.

There are several transverse furrows on the palmar aspect of the wrist. The lowest of these, which is slightly



convex downwards, corresponds with the upper edge of the anterior annular ligament and the intercarpal joint. The line of the radio-carpal joint, as already stated, is on a level with the apex of the styloid process of the ulna.

In forcible flexion of the wrist, the tendon of the flexor carpi radialis and that of the palmaris longus come up in relief. On the outer side of the first-named tendon we feel the pulse, the radial artery here lying close to the radius.

The tendon of the palmaris longus runs down the middle of the wrist, and immediately under it runs the median nerve. In letting out deep-seated matter near the wrist, the incision should be made close to and parallel with the inner edge of the radial flexor tendon, so as to avoid injury to the median nerve.

We can feel the tendon of the flexor carpi ulnaris for some distance above the wrist. It overlies the ulnar artery, and somewhat masks its pulsation.

142. In feeling the pulse, it should be remembered that, Pulse. in some cases, the superficialis volæ arises higher and is larger than usual. In such cases it runs by the side of the radial, and gives additional volume to the pulse. The old writers call it 'pulsus duplex.' When in doubt, therefore, it will be well to feel the pulse in each wrist.

143. The great synovial sheath under the annular ligament common to the flexor tendons of the fingers extends, Great carpal bursa. upwards, about an inch and a half above the edge of the ligament, and, downwards, as low as the middle of the palm. This general synovial sheath communicates with that of the thumb and the little finger; not with that of the index, middle, and ring fingers.

144. On the outer side of the wrist we can distinctly see and feel, when in action, the three extensor tendons of the thumb. Between the second and third there is a deep depression, at the root of the thumb, which the French call the 'tabatière anatomique.' In this depression we can make out—1, the relief of the superficial radial vein; 2, 'Tabatière anatomique.'



the radial artery, in its passage to the back of the hand ;  
3, the upper end of the metacarpal bone of the thumb.

Tendons  
on back of  
wrist.

145. The relative positions of the several extensor tendons of the wrist and fingers, as they play in their grooves over the back of the radius and ulna, can all be distinctly traced when the several muscles are put in action. The length of their synovial sheaths should be remembered. They vary from one inch and a half to two inches and a half. The longest of all are those of the extensors of the thumb. When these sheaths are inflamed and swollen, the motion of the tendons becomes painful and gives rise to a feeling of crepitus, called 'tenalgia crepitans' by some writers. It is said to be met with sometimes in 'pianistes.'

Lines of  
arteries.

146. The course of the radial artery corresponds with a line drawn from the outer border of the tendon of the biceps at the bend of the elbow down the front of the forearm to the front of the styloid process of the radius. In the upper third of its course the artery is overlapped by the supinator longus. To make allowance for this, the incision for the ligature of the artery in this situation should be made, not precisely in the line of its course, but rather nearer the middle of the forearm.

The line of the ulnar artery runs from the middle of the bend of the elbow to the outer side of the pisiform bone. The radial and ulnar arteries can, in most cases, be effectually commanded by pressure well applied at the wrist, in wounds of the palmar arch.

Before we make incisions along the forearm it is always desirable to ascertain whether the ulnar artery, which usually runs under the superficial muscles, may not run abnormally over them ; in which case its pulsations can be felt all down the forearm.



### THE HAND.

147. It is beside the purpose here to examine the question whether the hand can tell more than the arm, the leg, or any other part of the body, about the physical constitution of its owner, and to what use it has been put. Those who are interested in this subject should read a very elaborate treatise by Carus,<sup>1</sup> 'On the Reason and Meaning of the Different Forms of the Hand.' Still less would I indulge curiosity by inquiring whether the professors of chiromancy, relying on the text 'erit signum in manu tua et quasi monumentum ante oculos tuos,' can advance any reasonable pretensions for their assertion that they can read in the furrows of the palm the future destiny of its master.

148. The only furrow in the palm useful as a surgical landmark is that which runs transversely across its lower third, and is well seen when the fingers are slightly bent. This transverse furrow corresponds pretty nearly with the metacarpal joints of the fingers; with the upper limit of the synovial sheaths of the flexor tendons; with the splitting of the palmar fascia into its four slips. The transverse metacarpal ligament lies in the same line with it. Again, a little below this furrow, the digital arteries bifurcate to run along the opposite sides\* of the fingers.

Furrow in palm.

149. By pressing upon the interdigital folds of skin, we can feel the transverse ligament of the fingers, which prevents their too wide separation. The skin of these folds is much thinner on the dorsal than the palmar aspect; hence deep-seated abscesses in the palm very frequently burst on the back of the hand.

Interdigital folds.

150. Concerning the transverse furrows on the palmar surface of the fingers, notice that the first furrow, close to the palm, does not correspond with the first joint. The

Digital furrows.

<sup>1</sup> 'Ueber Grund und Bedeutung der verschiedenen Formen der Hand.' Stuttgart, 1846.



second and third furrows do correspond with their respective joints.

The slight depression observable between the ball of the thumb and that of the little finger corresponds with the middle of the anterior annular ligament.

Palmar  
arterial  
arches.

151. In opening abscesses in the palm, it is important to bear in mind the position of the large arterial arches which lie beneath the palmar fascia. The line of the superficial palmar arch crosses the palm about the junction of the upper with the lower two-thirds—that is, in the line of the thumb separated widely from the fingers. From this, the digital arteries run straight between the lines of the metacarpal bones towards the clefts of the fingers. Incisions, therefore, to let out pus beneath the palmar fascia may safely be made in the lower two-thirds of the palm, provided they run in the direction of the middle line of the fingers. The deep palmar arch lies half an inch nearer the wrist than the superficial.

Digital  
arteries.

152. As the digital arteries run along the sides of the fingers, the incision to open a thecal abscess should be made strictly in the middle line. It should be made not over but between the joints, since the sheath is strongest and thickest over the shafts of the phalanges, and therefore more likely to produce strangulation of the enclosed tendons.

Metacarpal  
joint of  
thumb.

153. The joint of the metacarpal bone of the thumb with the trapezium can be distinctly felt by tracing the dorsal surface of the bone upwards till we come to the prominence which indicates the joint at the bottom of the 'tabatière anatomique.' Supposing, however, there be much swelling, the knife introduced at the angle between the first and second metacarpal bones readily finds the joint if the blade be directed outwards.

Sesamoid  
bones.

154. The sesamoid bones of the thumb can be distinctly felt. Just above them—that is, nearer to the wrist—lies the joint between the metacarpal bone and the first phalanx. We should remember the position of these bones in amputation at this joint. *Mutatis mutandis* the same observations apply to the sesamoid bones of the great toe.



The extensor tendon of the second joint of the thumb crosses the apex of the first interosseous space. Under the tendon, and in the angle between the bones, we feel the radial artery just before it sinks into the palm.

155. The veins on the back of the hand, and their arrangement in the form of arches which receive the digital veins, is sufficiently obvious. The number and arrangement of the arches may vary, but in all hands it is interesting to see that the veins from the fingers run up between the knuckles in order to be out of harm's way.

Subcutaneous veins.

156. Since the dorsal interosseous arteries, like the palmar, run along the interosseous spaces, incisions to let out pus should always be made along the lines of the metacarpal bones.

Interosseous arteries.

The knuckles are formed by the extensor tendons which run over the ends of the metacarpal bones. They are prevented from slipping off, partly by their blending with the capsule of the joint, partly by their connection with the tendons of the lumbricales and interossei.

157. Small subcutaneous bursæ are sometimes developed over the knuckles and the backs of the joints of the fingers. They often become enlarged and unseemly in persons of a rheumatic or gouty tendency.

Digital bursæ.

158. In amputating one of the joints of the fingers, remember that the line of the joint does not correspond with the most convex part of it when bent, but is a trifle nearer to the end of the finger.

Digital joints.

Long and graceful fingers, coupled with thickness and breadth of the sentient pulp at their ends, and too great arching of the nails, have been regarded, ever since the days of Hippocrates, as not unlikely to be indications of a tendency to pulmonary disease.



*PALPATION BY THE RECTUM.*

The following report has been kindly written by Mr. Walsham, one of the Demonstrators of Anatomy, who, having a small hand (somewhat less than seven and a half inches round), has lately had opportunities, in St. Bartholomew's Hospital, of introducing it up the rectum, in the living subject, for the purpose of diagnosis :—

159. 'The hand and greater part of the forearm can be introduced into the rectum and passed through the sigmoid flexure for a considerable distance up the descending colon.

'Once beyond the sphincter, the hand enters a capacious sac, and the following important parts can be felt through its walls :—

'Through the anterior wall the hand first recognises the prostate, which feels like a moderately large chestnut. Immediately behind the prostate, the vesiculæ seminales may be distinguished as two softish masses situated one on either side of the middle line. Internal to them, the whipcord-like feel of the vasa deferentia can be readily traced over the bladder to the sides of the pelvis.

'The bladder is easily recognised, when moderately distended, as a soft fluctuating tumour behind the prostate; when empty it cannot be distinguished from the intestines, which then descend between the rectum and the pubes. The arch of the pubes can well be defined when the bladder is empty.

'Through the posterior wall of the bowel the coccyx and sacrum can be felt, the curve of the sacrum being readily followed by the hand.

'The projecting spine of the ischium on each side of the pelvis is a valuable landmark. From this point the outlines of the greater and lesser sacro-ischiatic foramina can be traced by the fingers; and any new growth, encroaching on the pelvic cavity through these apertures, could be easily detected.



‘If the hand be now pushed farther up the gut, the promontory of the sacrum is reached; the pulsation of the iliac vessels becomes manifest, and the course of the external iliac can be traced along the brim of the pelvis to the crural arch, the loose attachments of the rectum permitting very free movement in this direction. The internal iliac artery can also be followed to the upper part of the great sacro-ischiatic foramen.

‘By semi-rotatory movement, and alternately flexing and extending the fingers, the hand can gradually be insinuated into the sigmoid flexure and passed through this part of the bowel into the descending colon. In the sigmoid flexure the fingers can explore the whole of the lower part of the abdomen, the loose attachment of this portion of the gut permitting the hand to travel freely over the iliac and hypogastric regions.

‘The parts that can here be felt are the bifurcation of the aorta, the division of the common iliac arteries, the iliac fossa, and the crest of the ilium.

‘The hand may now be passed up the descending colon, and can, in some instances, reach the lower half of the left kidney. Probably the existence of a calculus in the kidney might be detected. The cartilage of the last rib can also be touched.

‘In the female, the uterus in the middle line, and the ovaries on either side, can be readily distinguished.

‘In the introduction of the hand into the rectum, in a patient under chloroform, the dilatation of the sphincter ani should be very gradual: first two fingers, then four, and finally the thumb should be passed. It is necessary to use considerable force, and unless care be taken, not only the integumentary edge of the anus, but the sphincter itself, may be lacerated. The introduction is facilitated by the application of the other hand upon the abdomen.

‘When the dilatation has been gradual and the hand not too large, no incontinence of fæces and no very considerable amount of pain has resulted.

‘Dr. G. Simon, in a paper in the “*Archiv für Clinische*



Chirurgie,"<sup>1</sup> states that repeated dilatation of the anus to its maximum does not destroy its contractile power ; that he has often made as many as five examinations of the same rectum ; that in clinical cases he has always permitted one or two of those present to repeat the examination, and that in no instance has permanent incontinence of fæces been the result.

'We have, however, been informed on reliable authority that incontinence of fæces has occasionally followed these repeated examinations.

'What sized hand can be introduced with safety? Dr. Simon states that it ought not to exceed nine inches in circumference.'

Lastly, we think it right to insist upon the important fact that, in some subjects, even a small hand cannot be passed up the rectum beyond the reflection of the peritoneum over the second part of the gut. In such instances the peritoneum offers a resistance like a tight garter, and prevents the farther advance of the hand without great risk of laceration of the parts.

#### EXAMINATION PER VAGINAM.

For this report I am indebted to Dr. Godson, Assistant Physician-Accoucheur, St. Bartholomew's Hospital :—

'The finger introduced through the hymen into the vagina comes upon the carunculæ myrtiformes, which are vascular membranous processes independent of the hymen, variable in number, size, and form. It also feels the transverse ridges known as "rugæ."

'Along the anterior wall of the vagina the finger readily detects the track of the urethra, which feels like a prominent cord and forms an excellent guide to the orifice of the meatus urinarius in passing a catheter. This orifice is in-

<sup>1</sup> Vol. xv. p. 1, 1872.



licated by a slight semicircular prominence, situated about one-third of an inch above the orifice of the vagina. Behind the urethra the finger comes upon the posterior wall of the bladder. But the bladder is not perceptible, as such, to the touch unless distended. With a catheter previously introduced it is much more readily explored.

‘The septum between the vagina and the rectum is so thin that, should the rectum contain faecal matter, its presence becomes at once apparent to the finger.

‘The cervix uteri is felt protruding from the roof of the vagina in a direction downwards and backwards—that is, in a line from the umbilicus to the coccyx. The os uteri is felt, small and round, in the centre of the cervix. The posterior lip feels a little lower than the anterior. The cul-de-sac formed by the vagina in front and behind the cervix should be perfectly elastic to the touch, and not communicate the sensation of a resisting body. Any resistance here bespeaks an abnormal condition.

‘The bony landmarks within reach of a finger, or perhaps two, in a woman who has not borne a child, are the symphysis pubis, the rami of the pubes and ischia. The coccyx and part of the hollow of the sacrum may also be felt, but not without exerting much pressure on the posterior wall of the vagina, which gives considerable pain. If the promontory of the sacrum can be felt, it is a sign that the conjugate diameter of the pelvis is abnormal.

‘The finger in the rectum can detect almost everything which has been mentioned in connection with the vagina. The shape and direction of the cervix uteri are almost as perceptible, and the posterior wall of the uterus can be examined. The peritoneal fold termed recto-vaginal (Douglas’s space) can also be well explored, and anything abnormal detected in this direction—a point of great importance in the diagnosis of diseases and displacements of the uterus.

‘The ovary in its normal state and position cannot be detected by the touch even with the hand firmly pressed on the hypogastrium. If a movable body be felt through



the vaginal roof on one side of the cervix, if this body be exquisitely tender and recede at once from the finger, it is an ovary in a state of prolapse.

‘The fundus of a healthy unimpregnated uterus never rises above the level of the brim of the pelvis, and cannot therefore be felt by pressing the hand on the hypogastrium.

‘The direction of the uterus is subject to changes which cannot be looked upon as abnormal. The fundus may be thrown backwards by a distended bladder, or forwards by a distended rectum. The axis of its cavity is not a straight but a curved line ; and uterine sounds should be shaped to suit it.’













*London, New Burlington Street,  
September, 1876.*

SELECTION

FROM

MESSRS J. & A. CHURCHILL'S

General Catalogue

COMPRISING

ALL RECENT WORKS PUBLISHED BY THEM

ON THE

ART AND SCIENCE

OF

M E D I C I N E



# INDEX

	PAGE		PAGE
Acton on the Reproductive Organs . . . . .	8	De Morgan on the Origin of Cancer . . . . .	18
Adams (W.) on Clubfoot . . . . .	6	De Valcourt on Cannes . . . . .	16
— (R.) on Rheumatic Gout . . . . .	18	Dobell's Lectures on Winter Cough . . . . .	14
Allen on Aural Catarrh . . . . .	6	— First Stage of Consumption . . . . .	14
Allingham on Diseases of Rectum . . . . .	8	Domville's Manual for Hospital Nurses . . . . .	14
Anatomical Remembrancer . . . . .	11	Druitt's Surgeon's Vade-Mecum . . . . .	4
Anderson (McC.) on Eczema . . . . .	19	Dunglison's Dictionary of Medical Science . . . . .	22
— (McC.) on Parasitic Affections . . . . .	19	Elam on Cerebria . . . . .	21
— (A. F.) Photographs of Leprosy . . . . .	20	Ellis's Manual of Diseases of Children . . . . .	12
Arnott on Cancer . . . . .	18	Fayrer's Observations in India . . . . .	4
Aveling's English Midwives . . . . .	14	Fergusson's Practical Surgery . . . . .	4
Balfour's Diseases of the Heart . . . . .	15	Fenwick's Guide to Medical Diagnosis . . . . .	11
Barclay's Medical Diagnosis . . . . .	11	— on the Stomach, &c. . . . .	17
Barker's Puerperal Diseases . . . . .	13	Flint on Phthisis . . . . .	15
Barnes' Obstetric Operations . . . . .	13	Flower's Nerves of the Human Body . . . . .	11
— Diseases of Women . . . . .	13	Foster's Clinical Medicine . . . . .	12
Basham on Renal Diseases . . . . .	8	Fox (T.) Atlas of Skin Diseases . . . . .	19
— on Diseases of the Kidneys . . . . .	8	Fox and Farquhar's Skin Diseases of India . . . . .	19
Beale on Kidney Diseases . . . . .	8	Frey's Histology and Histo-Chemistry of Man . . . . .	9
— on Disease Germs . . . . .	23	Gamgee on Fractures of the Limbs . . . . .	6
Bellamy's Guide to Surgical Anatomy . . . . .	11	Gant on the Science and Practice of Surgery . . . . .	4
Bennet's Winter and Spring on the Mediterranean . . . . .	16	— on Diseases of the Bladder . . . . .	8
— Pulmonary Consumption . . . . .	16	Gaskoin on Psoriasis or Lepra . . . . .	19
Bennett (J. H.) Antagonism of Medicines . . . . .	11	Glenn on the Laws affecting Medical Men . . . . .	20
Bennett (J. R.) on Cancerous and other Intrathoracic Growths . . . . .	18	Gordon on Fractures . . . . .	5
Black on the Urinary Organs . . . . .	8	Habershon on Diseases of the Liver . . . . .	17
— on Bright's Disease . . . . .	8	— on Diseases of the Stomach . . . . .	17
Brodhurst's Orthopædic Surgery . . . . .	6	Hamilton on Syphilitic Osteitis and Periostitis . . . . .	9
Bryant's Practice of Surgery . . . . .	4	Hancock's Surgery of Foot and Ankle . . . . .	6
Buchanan's Circulation of the Blood . . . . .	10	Harley on the Urine . . . . .	8
Bucknill and Tuke's Psychological Medicine . . . . .	21	Harris on Lithotomy . . . . .	7
Buzzard on Syphilitic Nervous Affections . . . . .	9	Hayden on the Heart . . . . .	15
Carpenter's Human Physiology . . . . .	9	Heath's Minor Surgery and Bandaging . . . . .	5
Carter on the Structure of Calculi . . . . .	8	— Diseases and Injuries of the Jaws . . . . .	5
— on Mycetoma . . . . .	19	— Operative Surgery . . . . .	5
Cauty on Diseases of the Skin . . . . .	20	— Practical Anatomy . . . . .	11
Chapman on Neuralgia . . . . .	18	Holden's Landmarks . . . . .	10
Clark's Outlines of Surgery . . . . .	4	Holt on Stricture of the Urethra . . . . .	7
— Surgical Diagnosis . . . . .	6	Holthouse on Hernial and other Tumours . . . . .	7
Clarke's Autobiographical Recollections . . . . .	22	Hood on Gout, Rheumatism, &c. . . . .	18
Clay's Obstetric Surgery . . . . .	12	Hooper's Physician's Vade-Mecum . . . . .	12
Cobbold on Worms . . . . .	20	Horton's Diseases of Tropical Climates . . . . .	16
Coles' Dental Mechanics . . . . .	23	Hutchinson's Clinical Surgery . . . . .	5
Cooper's Surgical Dictionary . . . . .	5	Huth's Marriage of Near Kin . . . . .	10
Coulson on Syphilis . . . . .	9	Jones (C. H.) and Sieveking's Pathological Anatomy . . . . .	10
— on Stone in the Bladder . . . . .	9	— (C. H.) on Functional Nervous Disorders . . . . .	17
Cullingworth's Nurse's Companion . . . . .	14	— (Wharton) Ophthalmic Medicine and Surgery . . . . .	23
Curling's Diseases of the Rectum . . . . .	7		
Dalby on the Ear . . . . .	6		
Dalton's Human Physiology . . . . .	10		
Day on Children's Diseases . . . . .	13		



	PAGE		PAGE
Jordan's Treatment of Surgical In-		Smith (W. R.) on Nursing . . .	14
flammations . . . . .	6	Smith's Dental Anatomy . . .	23
— Surgical Inquiries . . .	6	Squire's Temperature Observations .	18
Kennion's Springs of Harrogate .	16	Steiner's Diseases of Children . .	13
Lee (H.) Practical Pathology . .	9	Stowe's Toxicological Chart . . .	20
— on Syphilis . . . . .	9	Swain's Surgical Emergencies . .	5
Leared on Imperfect Digestion . .	17	Swayne's Obstetric Aphorisms . .	13
Liebreich's Atlas of Ophthalmoscopy	22	Taylor's Principles of Medical Juris-	
Liveing on Megrim, &c. . . . .	17	prudence . . . . .	20
Macdonald's Examination of Water .	21	— Manual of Medical Juris-	
Mackenzie on Growths in the Larynx	15	prudence . . . . .	20
Macnamara on Diseases of the Eye .	22	— Poisons in relation to Medical	
Madden's Health Resorts . . . .	16	Jurisprudence . . . . .	20
Marsden on certain Forms of Cancer	18	Thompson on Stricture of Urethra .	7
Maunder's Operative Surgery . . .	4	— on Practical Lithotomy	
— Surgery of Arteries . . . . .	4	and Lithotrity . . . . .	7
Mayne's Medical Vocabulary . . .	22	— on Diseases of the Urinary	
Meryon's System of Nerves . . .	18	Organs . . . . .	7
Moore's Family Medicine for India .	16	— on Diseases of the Prostate . .	7
Paton on Action and Sounds of Heart	15	— on Calculous Disease . . . .	7
Parkes' Manual of Practical Hygiene	21	Thornton on Tracheotomy . . . .	15
— Issue of a Spirit Ration . . .	17	Thorowgood on Asthma . . . . .	15
Parkin's Epidemiology . . . . .	23	— on Materia Medica . . . . .	12
Pavy on Food and Dietetics . . .	17	Tibbits' Medical Electricity . . .	21
Phillips' Materia Medica and Thera-		Tilt's Uterine Therapeutics . . .	13
peutics . . . . .	12	— Change of Life . . . . .	13
Pirrie's Principles and Practice of		— Health in India . . . . .	16
Surgery . . . . .	4	Tomes' Dental Surgery . . . . .	23
Ramsbotham's Obstetric Medicine		Tufnell's Internal Aneurism . . .	7
and Surgery . . . . .	13	Tuke on the Influence of the Mind	
Reynolds' Uses of Electricity . . .	22	upon the Body . . . . .	21
Richardson's Practical Physic . . .	11	Van Buren on Diseases of the Genito-	
Roberts' Practice of Midwifery . .	13	Urinary Organs . . . . .	9
Ross's Graft Theory of Disease . .	23	Veitch's Handbook for Nurses . .	14
Roy's Burdwan Fever . . . . .	16	Wagstaffe's Human Osteology . .	10
Royle and Harley's Materia Medica .	12	Walker on Egypt . . . . .	16
Rutherford's Practical Histology .	10	Walton's Diseases of the Eye . . .	23
Sabben and Browne's Handbook of		Ward on Affections of the Liver .	17
Law and Lunacy . . . . .	21	Waring's Practical Therapeutics .	12
Sanderson's Physiological Handbook .	10	— Bazaar Medicines of India . .	17
Sansom's Diseases of the Heart . .	15	Waters on Diseases of the Chest . .	14
Savage on the Female Pelvic Organs .	5	Wells (Soelberg) on Diseases of the	
Savory's Domestic Medicine . . .	14	Eye . . . . .	22
Sayre's Orthopædic Surgery . . .	6	— Long, Short, and Weak Sight .	22
Schroeder's Manual of Midwifery . .	13	— (Spencer) on Diseases of the	
Semple on the Heart . . . . .	15	Ovaries . . . . .	14
Sewill's Dental Anatomy . . . . .	23	Wife's Domain . . . . .	14
Shapter's Diseases of the Heart . .	15	Wilks' Pathological Anatomy . . .	10
Shaw's Medical Remembrancer . . .	11	Wilson (E.) Anatomist's Vade-	
Sheppard on Madness . . . . .	21	Mecum . . . . .	11
Sibson's Medical Anatomy . . . . .	11	— on Diseases of the Skin . . .	19
Sieveling's Medical Adviser in Life		— Lectures on Ekzema . . . . .	19
Assurance . . . . .	20	— Lectures on Dermatology . . .	19
Smith (H.) on the Surgery of the		— (G.) Handbook of Hygiene . .	21
Rectum . . . . .	8	Winslow's Obscure Diseases of the	
Smith (E.) on Wasting Diseases of		Brain and Mind . . . . .	20
Children . . . . .	12	Wolff on Zymotic Diseases . . . .	23
— Clinical Studies . . . . .	12		



**THE PRACTICE OF SURGERY:**

a Manual by THOMAS BRYANT, F.R.C.S., Surgeon to Guy's Hospital.  
Second Edition, 2 vols., crown 8vo, with 559 Engravings, 25s. [1876]

**THE PRINCIPLES AND PRACTICE OF SURGERY**

by WILLIAM PIRRIE, F.R.S.E., Professor of Surgery in the University  
of Aberdeen. Third Edition, 8vo, with 490 Engravings, 28s. [1873]

**A SYSTEM OF PRACTICAL SURGERY**

by Sir WILLIAM FERGUSSON, Bart., F.R.C.S., F.R.S., Serjeant-  
Surgeon to the Queen. Fifth Edition, 8vo, with 463 Engravings, 21s.  
[1870]

**OPERATIVE SURGERY**

by C. F. MAUNDER, F.R.C.S., Surgeon to the London Hospital, for-  
merly Demonstrator of Anatomy at Guy's Hospital. Second Edition,  
post 8vo, with 164 Engravings, 6s. [1872]

BY THE SAME AUTHOR.

**SURGERY OF THE ARTERIES:**

Lettsomian Lectures for 1875, on Aneurisms, Wounds, Hæmorrhages,  
&c. Post 8vo, with 18 Engravings, 5s. [1875]

**THE SURGEON'S VADE-MECUM**

by ROBERT DRUITT. Tenth Edition, fcap. 8vo, with numerous En-  
gravings, 12s. 6d. [1870]

**THE SCIENCE AND PRACTICE OF SURGERY:**

a complete System and Textbook by F. J. GANT, F.R.C.S., Surgeon to  
the Royal Free Hospital. 8vo, with 470 Engravings, 24s. [1871]

**OUTLINES OF SURGERY AND SURGICAL PATHOLOGY**

including the Diagnosis and Treatment of Obscure and Urgent  
Cases, and the Surgical Anatomy of some Important Structures and  
Regions, by F. LE GROS CLARK, F.R.S., Consulting Surgeon to St.  
Thomas's Hospital. Second Edition, Revised and Expanded by the  
Author, assisted by W. W. WAGSTAFFE, F.R.C.S., Assistant-Surgeon  
to, and Joint-Lecturer on Anatomy at, St. Thomas's Hospital. 8vo,  
10s. 6d. [1872]

**CLINICAL AND PATHOLOGICAL OBSERVATIONS IN INDIA**

by Sir J. FAYRER, K.S.I., M.D., F.R.C.P. Lond., F.R.S.E., Honorary  
Physician to the Queen. 8vo, with Engravings, 20s. [1873]



**SURGICAL EMERGENCIES**

together with the Emergencies attendant on Parturition and the Treatment of Poisoning: a Manual for the use of General Practitioners, by WILLIAM P. SWAIN, F.R.C.S., Surgeon to the Royal Albert Hospital, Devonport. Second Edition, post 8vo, with 104 Engravings, 6s. 6d. [1876]

**ILLUSTRATIONS OF CLINICAL SURGERY,**

consisting of Plates, Photographs, Woodcuts, Diagrams, &c., illustrating Surgical Diseases, Symptoms and Accidents; also Operations and other methods of Treatment. By JONATHAN HUTCHINSON, F.R.C.S., Senior Surgeon to the London Hospital. In Quarterly Fasciculi. Fasc. I to IV already issued. 6s. 6d. each. [1876]

**FRACTURES OF THE LOWER END OF THE RADIUS,**

Fractures of the Clavicle, and on the Reduction of the Recent Inward Dislocations of the Shoulder Joint. By ALEXANDER GORDON, M.D., Professor of Surgery in Queen's College, Belfast. With Engravings, 8vo, 5s. [1875]

**MINOR SURGERY AND BANDAGING**

a Manual for the Use of House-Surgeons, Dressers, and Junior Practitioners, by CHRISTOPHER HEATH, F.R.C.S., Surgeon to University College Hospital. Fifth Edition, fcap 8vo, with 86 Engravings, 5s. 6d. [1875]

BY THE SAME AUTHOR,

**INJURIES AND DISEASES OF THE JAWS:**

JACKSONIAN PRIZE ESSAY. Second Edition, 8vo, with 164 Engravings, 12s. [1872]

BY THE SAME AUTHOR.

**A COURSE OF OPERATIVE SURGERY:**

with Plates drawn from Nature by M. LÉVEILLÉ, and coloured by hand under his direction. In Five Parts, each to contain 4 Plates. Large 8vo. 7s. 6d. each Part. [1876]

**DICTIONARY OF PRACTICAL SURGERY**

and Encyclopædia of Surgical Science, by SAMUEL COOPER. New Edition, brought down to the present Time by SAMUEL A. LANE, Consulting Surgeon to St. Mary's and to the Lock Hospitals; assisted by various Eminent Surgeons. 2 vols. 8vo, 50s. [1861 and 1872]

**THE FEMALE PELVIC ORGANS**

their Surgery, Surgical Pathology, and Surgical Anatomy, in a Series of Coloured Plates taken from Nature: with Commentaries, Notes, and Cases, by HENRY SAVAGE, M.D. Lond., F.R.C.S., Consulting Officer of the Samaritan Free Hospital. Third Edition, 4to, £1 15s. [1875]



**FRACTURES OF THE LIMBS**

(Treatment of) by J. SAMPSON GAMGEE, Surgeon to the Queen's Hospital, Birmingham. 8vo, with Plates, 10s. 6d. [1871]

**DISEASES AND INJURIES OF THE EAR**

by W. B. DALBY, F.R.C.S., M.B., Aural Surgeon and Lecturer on Aural Surgery at St. George's Hospital. Crown 8vo, with 21 Engravings, 6s. 6d. [1873]

**AURAL CATARRH;**

or, the Commonest Forms of Deafness, and their Cure, by PETER ALLEN, M.D., F.R.C.S.E., late Aural Surgeon to St. Mary's Hospital. Second Edition, crown 8vo, with Engravings, 8s. 6d. [1874]

**PRINCIPLES OF SURGICAL DIAGNOSIS**

especially in Relation to Shock and Visceral Lesions, Lectures delivered at the Royal College of Surgeons by F. LE GROS CLARK, F.R.C.S., Consulting Surgeon to St. Thomas's Hospital. 8vo, 10s. 6d. [1870]

**CLUBFOOT:**

its Causes, Pathology, and Treatment; being the Jacksonian Prize Essay by WM. ADAMS, F.R.C.S., Surgeon to the Great Northern Hospital. Second Edition, 8vo, with 106 Engravings and 6 Lithographic Plates, 15s. [1873]

**ORTHOPÆDIC SURGERY**

Lectures delivered at St. George's Hospital, by BERNARD E. BRODHURST, F.R.C.S., Surgeon to the Royal Orthopædic Hospital. Second Edition, 8vo, with Engravings, 12s. 6d. [1876]

**OPERATIVE SURGERY OF THE FOOT AND ANKLE**

by HENRY HANCOCK, F.R.C.S., Consulting Surgeon to Charing Cross Hospital. 8vo, with Engravings, 15s. [1873]

**ORTHOPÆDIC SURGERY:**

and Diseases of the Joints. Lectures by LEWIS A. SAYRE, M.D., Professor of Orthopædic Surgery, Fractures and Dislocations, and Clinical Surgery, in Bellevue Hospital Medical College, New York. With 274 Wood Engravings, 8vo, 20s. [1876]

**THE TREATMENT OF SURGICAL INFLAMMATIONS**

by a New Method, which greatly shortens their Duration, by FURNEAUX JORDAN, F.R.C.S., Professor of Surgery in Queen's College, Birmingham. 8vo, with Plates, 7s. 6d. [1870]

BY THE SAME AUTHOR,

**SURGICAL INQUIRIES**

With numerous Lithographic Plates. 8vo, 5s.

[1873]



**INTERNAL ANEURISM:**

Successful Treatment of, by Consolidation of the Contents of the Sac.  
By T. JOLIFFE TUFNELL, F.R.C.S.I., President of the Royal College  
of Surgeons in Ireland. With Coloured Plates. Second Edition,  
royal 8vo, 5s. [1875]

**HERNIAL AND OTHER TUMOURS**

of the Groin and its Neighbourhood, with some Practical Remarks on  
the Radical Cure of Ruptures, by C. HOLTHOUSE, F.R.C.S., Surgeon to  
the Westminster Hospital. 8vo, 6s. 6d. [1870]

**DISEASES OF THE RECTUM**

by T. B. CURLING, F.R.S., Consulting Surgeon to the London  
Hospital. Fourth Edition, Revised, 8vo, 7s. 6d. [1876]

**STRICTURE OF THE URETHRA**

and Urinary Fistulæ; their Pathology and Treatment: Jacksonian  
Prize Essay by Sir HENRY THOMPSON, F.R.C.S., Emeritus Professor  
of Surgery to University College. Third Edition, 8vo, with Plates,  
10s. [1869]

BY THE SAME AUTHOR,

**PRACTICAL LITHOTOMY AND LITHOTRITY;**

or, An Inquiry into the best Modes of removing Stone from the  
Bladder. Second Edition, 8vo, with numerous Engravings. 10s. [1871]

ALSO,

**DISEASES OF THE URINARY ORGANS**

(Clinical Lectures). Fourth Edition, 8vo, with 2 Plates and 59  
Engravings, 12s. [1876]

ALSO,

**DISEASES OF THE PROSTATE:**

their Pathology and Treatment. Fourth Edition, 8vo, with numerous  
Plates, 10s. [1873]

ALSO,

**THE PREVENTIVE TREATMENT OF CALCULOUS DISEASE**

and the Use of Solvent Remedies. Second Edition, fcap. 8vo, 2s. 6d.  
[1876]

**STRICTURE OF THE URETHRA**

(the Immediate Treatment of), by BARNARD HOLT, F.R.C.S.,  
Consulting Surgeon to the Westminster Hospital. Third Edition,  
8vo, 6s. [1868]

**LITHOTOMY AND EXTRACTION OF STONE**

from the Bladder, Urethra, and Prostate of the Male, and from the  
Bladder of the Female, by W. POULETT HARRIS, M.D., Surgeon-Major  
H.M. Bengal Medical Service. With Engravings, 8vo, 10s. 6d. [1876]



**THE SURGERY OF THE RECTUM:**

Lettsomian Lectures by HENRY SMITH, F.R.C.S., Surgeon to King's College Hospital. Fourth Edition, fcap. 8vo, 5s. [1876]

**FISTULA, HÆMORRHOIDS, PAINFUL ULCER,**

Stricture, Prolapsus, and other Diseases of the Rectum: their Diagnosis and Treatment, by WM. ALLINGHAM, F.R.C.S., Surgeon to St. Mark's Hospital for Fistula, &c., late Surgeon to the Great Northern Hospital. Second Edition, 8vo, 7s. [1872]

**THE URINE AND ITS DERANGEMENTS**

with the Application of Physiological Chemistry to the Diagnosis and Treatment of Constitutional as well as Local Diseases. Lectures by GEORGE HARLEY, M.D., F.R.S., F.R.C.P., formerly Professor in University College. Post 8vo, 9s. [1872]

**KIDNEY DISEASES, URINARY DEPOSITS**

and Calculous Disorders by LIONEL S. BEALE, M.B., F.R.S., F.R.C.P., Physician to King's College Hospital. Third Edition, 8vo, with 70 Plates, 25s. [1868]

**DISEASES OF THE BLADDER,**

Prostate Gland and Urethra, including a practical view of Urinary Diseases, Deposits and Calculi, by F. J. GANT, F.R.C.S., Surgeon to the Royal Free Hospital. Fourth Edition, crown 8vo, with Engravings, 10s. 6d. [1876]

**RENAL DISEASES:**

a Clinical Guide to their Diagnosis and Treatment by W. R. BASHAM, M.D., F.R.C.P., Senior Physician to the Westminster Hospital. Post 8vo, 7s. [1870]

BY THE SAME AUTHOR,

**THE DIAGNOSIS OF DISEASES OF THE KIDNEYS**

(Aids to). 8vo, with 10 Plates, 5s. [1872]

**MICROSCOPIC STRUCTURE OF URINARY CALCULI**

by H. V. CARTER, M.D., Surgeon-Major, H.M.'s Bombay Army. 8vo, with 4 Plates, 5s. [1873]

**THE REPRODUCTIVE ORGANS**

in Childhood, Youth, Adult Age, and Advanced Life (Functions and Disorders of), considered in their Physiological, Social, and Moral Relations, by WILLIAM ACTON, M.R.C.S. Sixth Edition, 8vo, 12s. [1875]

**URINARY AND REPRODUCTIVE ORGANS**

(Functional Diseases of) by D. CAMPBELL BLACK, M.D., L.R.C.S. Edin. Second Edition. 8vo, 10s. 6d. [1875]

BY THE SAME AUTHOR.

**LECTURES ON BRIGHT'S DISEASE**

delivered at the Royal Infirmary of Glasgow. 8vo, with 20 Engravings, 6s. 6d. [1875]



**GENITO-URINARY ORGANS, INCLUDING SYPHILIS**

A Practical Treatise on their Surgical Diseases, designed as a Manual for Students and Practitioners, by W. H. VAN BUREN, M.D., Professor of the Principles of Surgery in Bellevue Hospital Medical College, New York, and E. L. KEYES, M.D., Professor of Dermatology in Bellevue Hospital Medical College, New York. Royal 8vo, with 140 Engravings, 21s. [1874]

**SYPHILIS**

A Treatise by WALTER J. COULSON, F.R.C.S., Surgeon to the Lock Hospital. 8vo, 10s. [1869]

BY THE SAME AUTHOR,

**STONE IN THE BLADDER:**

Its Prevention, Early Symptoms, and Treatment by Lithotrity. 8vo, 6s. [1868]

**PRACTICAL PATHOLOGY:**

containing Lectures on Suppurative Fever, Diseases of the Veins, Hæmorrhoidal Tumours, Diseases of the Rectum, Syphilis, Gonorrhæal Ophthalmia, &c., by HENRY LEE, F.R.C.S., Surgeon to St. George's Hospital. Third Edition, in 2 vols. 8vo, 10s. each. [1870]

BY THE SAME AUTHOR,

**LECTURES ON SYPHILIS,**

and on some forms of Local Disease, affecting principally the Organs of Generation. With Engravings, 8vo, 10s. [1875]

**SYPHILITIC NERVOUS AFFECTIONS**

(Clinical Aspects of) by THOMAS BUZZARD, M.D., F.R.C.P. Lond., Physician to the National Hospital for Paralysis and Epilepsy. Post 8vo, 5s. [1874]

**SYPHILITIC OSTEITIS AND PERIOSTITIS**

Lectures by JOHN HAMILTON, F.R.C.S.I., Surgeon to the Richmond Hospital and to Swift's Hospital for Lunatics, Dublin. 8vo, with Plates, 6s. 6d. [1874]

**PRINCIPLES OF HUMAN PHYSIOLOGY**

by W. B. CARPENTER, M.D., F.R.S. Eighth Edition by HENRY POWER, M.B., F.R.C.S., Examiner in Natural Science, University of Oxford, and in Natural Science and Medicine, University of Cambridge. 8vo, with 3 Steel Plates and 371 Engravings, 31s. 6d. [1876]

**HISTOLOGY AND HISTO-CHEMISTRY OF MAN**

A Treatise on the Elements of Composition and Structure of the Human Body, by HEINRICH FREY, Professor of Medicine in Zurich. Translated from the Fourth German Edition by ARTHUR E. J. BARKER, Assistant-Surgeon to University College Hospital. And Revised by the Author. 8vo, with 608 Engravings, 21s. [1874]



**HUMAN PHYSIOLOGY:**

A Treatise designed for the Use of Students and Practitioners of Medicine, by JOHN C. DALTON, M.D., Professor of Physiology and Hygiene in the College of Physicians and Surgeons, New York. Sixth Edition, royal 8vo, with 316 Engravings, 20s. [1875]

**HANDBOOK FOR THE PHYSIOLOGICAL LABORATORY**

by E. KLEIN, M.D., F.R.S., Assistant Professor in the Pathological Laboratory of the Brown Institution, London; J. BURDON-SANDERSON, M.D., F.R.S., Professor of Practical Physiology in University College, London; MICHAEL FOSTER, M.D., F.R.S., Prælector of Physiology in Trinity College, Cambridge; and T. LAUDER BRUNTON, M.D., D.Sc., Lecturer on Materia Medica at St. Bartholomew's Hospital; edited by J. BURDON-SANDERSON. 8vo, with 123 Plates, 24s. [1873]

**PRACTICAL HISTOLOGY**

(Outlines of) by WILLIAM RUTHERFORD, M.D., Professor of the Institutes of Medicine in the University of Edinburgh. With Engravings. Crown 8vo, interleaved, 3s. [1875]

**THE MARRIAGE OF NEAR KIN**

Considered with respect to the Laws of Nations, Results of Experience, and the Teachings of Biology, by ALFRED H. HUTH. 8vo, 14s. [1875]

**STUDENTS' GUIDE TO HUMAN OSTEOLOGY**

By WILLIAM WARWICK WAGSTAFFE, F.R.C.S., Assistant-Surgeon and Lecturer on Anatomy, St. Thomas's Hospital. With 23 Plates and 66 Engravings. Fcap. 8vo, 10s. 6d. [1875]

**LANDMARKS, MEDICAL AND SURGICAL.**

By LUTHER HOLDEN, F.R.C.S., Surgeon to St. Bartholomew's Hospital. 8vo, 3s. 6d. [1876]

**THE CIRCULATION OF THE BLOOD**

(Forces which carry on) by ANDREW BUCHANAN, M.D., Professor of Physiology in the University of Glasgow. Second Edition, 8vo, with Engravings, 5s. [1874]

**PATHOLOGICAL ANATOMY**

Lectures by SAMUEL WILKS, M.D., F.R.S., Physician to, and Lecturer on Medicine at, Guy's Hospital; and WALTER MOXON, M.D., F.R.C.P., Physician to, and Lecturer on Materia Medica at, Guy's Hospital. Second Edition, 8vo, with Plates, 18s. [1875]

**PATHOLOGICAL ANATOMY**

A Manual by C. HANDFIELD JONES, M.B., F.R.S., Physician to St. Mary's Hospital, and EDWARD H. SIEVEKING, M.D., F.R.C.P., Physician to St. Mary's Hospital. Edited by J. F. PAYNE, M.D., F.R.C.P., Assistant Physician and late Demonstrator of Morbid Anatomy at St. Thomas's Hospital. Second Edition, crown 8vo, with 195 Engravings, 16s. [1875]



**THE ANATOMIST'S VADE-MECUM:**

a System of Human Anatomy by ERASMUS WILSON, F.R.C.S., F.R.S.  
Ninth Edition, by G. BUCHANAN, M.A., M.D., Professor of Clinical  
Surgery in the University of Glasgow. Crown 8vo, with 371 Engrav-  
ings, 14s. [1873]

**PRACTICAL ANATOMY:**

a Manual of Dissections by CHRISTOPHER HEATH, F.R.C.S., Surgeon  
to University College Hospital. Third Edition, fcap 8vo, with 226  
Engravings, 12s. 6d. [1874]

**MEDICAL ANATOMY**

by FRANCIS SIBSON, M.D., F.R.C.P., F.R.S., Consulting Physician to  
St. Mary's Hospital. Imp. folio, with 21 coloured Plates, cloth, 42s.;  
half-morocco, 50s. [Completed in 1869]

**STUDENT'S GUIDE TO SURGICAL ANATOMY:**

a Text-book for the Pass Examination, by E. BELLAMY, F.R.C.S.,  
Senior Assistant-Surgeon and Lecturer on Anatomy at Charing  
Cross Hospital. Fcap 8vo, with 50 Engravings, 6s. 6d. [1873]

**DIAGRAMS OF THE NERVES OF THE HUMAN BODY**

Exhibiting their Origin, Divisions, and Connexions, with their Distri-  
bution, by WILLIAM HENRY FLOWER, F.R.S., Conservator of the  
Museum of the Royal College of Surgeons. Second Edition, roy. 4to,  
12s. [1872]

**THE STUDENT'S GUIDE TO MEDICAL DIAGNOSIS**

by SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London  
Hospital. Fourth Edition, fcap 8vo, with 106 Engravings, 6s. 6d.  
[1876]

**A MANUAL OF MEDICAL DIAGNOSIS**

by A. W. BARCLAY, M.D., F.R.C.P., Physician to, and Lecturer on  
Medicine at, St. George's Hospital. Third Edition, fcap 8vo, 10s. 6d.  
[1876]

**THE MEDICAL REMEMBRANCER;**

or, Book of Emergencies. By E. SHAW, M.R.C.S. Fifth Edition by  
JONATHAN HUTCHINSON, F.R.C.S., Senior Surgeon to the London  
Hospital. 32mo, 2s. 6d. [1867]

**THE ANATOMICAL REMEMBRANCER;**

or, Complete Pocket Anatomist. Seventh Edition, 32mo, 3s. 6d. [1872]

**ANTAGONISM OF MEDICINES**

(Researches into the) being the Report of the Edinburgh Committee  
of the British Medical Association. By J. HUGHES BENNETT, M.D.  
Post 8vo, 3s. 6d. [1875]

**DISCOURSES ON PRACTICAL PHYSIC**

by B. W. RICHARDSON, M.D., F.R.C.P., F.R.S. 8vo, 5s. [1871]



**HOOPER'S PHYSICIAN'S VADE-MECUM;**

or, Manual of the Principles and Practice of Physic, Ninth Edition  
by W. A. GUY, M.B., F.R.S., and JOHN HARLEY, M.D., F.R.C.P.  
Fcap 8vo, with Engravings, 12s. 6d. [1874]

**CLINICAL MEDICINE**

Lectures and Essays by BALTHAZAR FOSTER, M.D., F.R.C.P. Lond.,  
Professor of Medicine in Queen's College, Birmingham. 8vo, 10s. 6d.  
[1874]

**ROYLE'S MANUAL OF MATERIA MEDICA AND THERAPEUTICS.**

Sixth Edition by JOHN HARLEY, M.D., Assistant Physician to, and  
Joint Lecturer on Physiology at, St. Thomas's Hospital. Crown 8vo,  
with 139 Engravings, 15s. [1876]

**MATERIA MEDICA AND THERAPEUTICS:**

(Vegetable Kingdom), by CHARLES D. F. PHILLIPS, M.D., F.R.C.S.E.  
8vo, 15s. [1874]

**PRACTICAL THERAPEUTICS**

A Manual by E. J. WARING, M.D., F.R.C.P. Lond. Third Edition,  
fcap 8vo, 12s. 6d. [1871]

**THE STUDENT'S GUIDE TO MATERIA MEDICA**

by JOHN C. THOROWGOOD, M.D. Lond., Physician to the City of  
London Hospital for Diseases of the Chest. Fcap 8vo, with Engrav-  
ings, 6s. 6d. [1874]

**THE DISEASES OF CHILDREN**

A Practical Manual, with a Formulary, by EDWARD ELLIS, M.D.,  
Physician to the Victoria Hospital for Children. Second Edition,  
crown 8vo, 7s. [1873]

**THE WASTING DISEASES OF CHILDREN**

by EUSTACE SMITH, M.D. Lond., Physician to the King of the Belgians,  
Physician to the East London Hospital for Children. Second Edition,  
post 8vo, 7s. 6d. [1870]

BY THE SAME AUTHOR,

**CLINICAL STUDIES OF DISEASE IN CHILDREN.**

Diseases of the Lungs; Acute Tuberculosis. Post 8vo, 7s. 6d. [1876]

**OBSTETRIC SURGERY**

A Complete Handbook, giving Short Rules of Practice in every Emer-  
gency, from the Simplest to the most Formidable Operations connected  
with the Science of Obstetricy, by CHARLES CLAY, Ext.L.R.C.P. Lond.,  
L.R.C.S.E., late Senior Surgeon and Lecturer on Midwifery, St.  
Mary's Hospital, Manchester. Fcap 8vo, with 91 Engravings, 6s. 6d.  
[1874]



**THE DISEASES OF CHILDREN**

Essays by WILLIAM HENRY DAY, M.D., Physician to the Samaritan Hospital for Diseases of Women and Children. Fcap 8vo, 5s. [1873]

**COMPENDIUM OF CHILDREN'S DISEASES**

A Handbook for Practitioners and Students, by JOHANN STEINER, M.D., Professor of the Diseases of Children in the University of Prague. Translated from the Second German Edition by LAWSON TAIT, F.R.C.S., Surgeon to the Birmingham Hospital for Women. 8vo, 12s. 6d. [1874]

**THE STUDENT'S GUIDE TO THE PRACTICE OF MIDWIFERY**

by D. LLOYD ROBERTS, M.D., Physician to St. Mary's Hospital, Manchester. Fcap. 8vo, with 95 Engravings, 6s. 6d. [1875]

**PUERPERAL DISEASES**

Clinical Lectures by FORDYCE BARKER, M.D., Obstetric Physician to Bellevue Hospital, New York. 8vo, 15s. [1874]

**OBSTETRIC OPERATIONS,**

including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour; Lectures by ROBERT BARNES, M.D., F.R.C.P., Obstetric Physician and Lecturer on Obstetrics and the Diseases of Women and Children to St. George's Hospital. Third Edition, 8vo, with 124 Engravings, 18s. [1875]

BY THE SAME AUTHOR,

**MEDICAL AND SURGICAL DISEASES OF WOMEN**

(a Clinical History). 8vo, with 169 Engravings, 28s. [1873]

**OBSTETRIC MEDICINE AND SURGERY**

(Principles and Practice of) by F. H. RAMSBOTHAM, M.D., F.R.C.P. Fifth Edition, 8vo, with 120 Plates, 22s. [1867]

**OBSTETRIC APHORISMS**

for the Use of Students commencing Midwifery Practice by J. G. SWAYNE, M.D., Consulting Physician-Accoucheur to the Bristol General Hospital, and Lecturer on Obstetric Medicine at the Bristol Medical School. Sixth Edition, fcap 8vo, with Engravings, 3s. 6d. [1876]

**SCHROEDER'S MANUAL OF MIDWIFERY,**

including the Pathology of Pregnancy and the Puerperal State. Translated by CHARLES H. CARTER, B.A., M.D. 8vo, with Engravings, 12s. 6d. [1873]

**A HANDBOOK OF UTERINE THERAPEUTICS**

and of Diseases of Women by E. J. TILT, M.D., M.R.C.P. Third Edition, post 8vo, 10s. [1866]

BY THE SAME AUTHOR,

**THE CHANGE OF LIFE**

in Health and Disease: a Practical Treatise on the Nervous and other Affections incidental to Women at the Decline of Life. Third Edition, 8vo, 10s. 6d. [1870]



**DISEASES OF THE OVARIES :**

their Diagnosis and Treatment, by T. SPENCER WELLS, F.R.C.S., Surgeon to the Queen's Household and to the Samaritan Hospital. 8vo, with about 150 Engravings, 21s. [1872]

**HANDBOOK FOR NURSES FOR THE SICK**

by ZEPHERINA P. VEITCH. Second Edition, crown 8vo, 3s. 6d. [1876]

**A MANUAL FOR HOSPITAL NURSES**

and others engaged in Attending on the Sick by EDWARD J. DOMVILLE, L.R.C.P., M.R.C.S. Second Edition, crown 8vo, 2s. 6d. [1875]

**THE NURSE'S COMPANION :**

A Manual of General and Monthly Nursing, by CHARLES J. CULLINGWORTH, Surgeon to St. Mary's Hospital, Manchester. Fcap. 8vo, 2s. 6d. [1876]

**LECTURES ON NURSING**

by WILLIAM ROBERT SMITH, L.R.C.S.E, Resident Surgeon, Royal Hants County Hospital, Winchester. With 26 Engravings. Post 8vo, 6s. [1875]

**ENGLISH MIDWIVES :**

their History and Prospects, by J. H. AVELING, M.D., Physician to the Chelsea Hospital for Women, Examiner of Midwives for the Obstetrical Society of London. Crown 8vo, 5s. [1872]

**A COMPENDIUM OF DOMESTIC MEDICINE**

and Companion to the Medicine Chest; intended as a Source of Easy Reference for Clergymen, and for Families residing at a Distance from Professional Assistance, by JOHN SAVORY, M.S.A. Eighth Edition, 12mo, 5s. [1871]

**THE WIFE'S DOMAIN**

The Young Couple—The Mother—The Nurse—The Nursling, by PHILOTHALOS. Second Edition, post 8vo, 3s. 6d. [1874]

**WINTER COUGH**

(Catarrh, Bronchitis, Emphysema, Asthma), Lectures by HORACE DOBELL, M.D., Consulting Physician to the Royal Hospital for Diseases of the Chest. Third Edition, with Coloured Plates, 8vo, 10s. 6d. [1875]

BY THE SAME AUTHOR,

**THE TRUE FIRST STAGE OF CONSUMPTION**

(Lectures). Crown 8vo, 3s. 6d. [1867]

**DISEASES OF THE CHEST :**

Contributions to their Clinical History, Pathology, and Treatment, by A. T. H. WATERS, M.D., F.R.C.P., Physician to the Liverpool Royal Infirmary. Second Edition, 8vo, with Plates, 15s. [1873]



**DISEASES OF THE HEART**

Their Pathology, Diagnosis, Prognosis, and Treatment (a Manual),  
by ROBERT H. SEMPLE, M.D., F.R.C.P., Physician to the Hospital for  
Diseases of the Throat. 8vo, 8s. 6d. [1875]

**DISEASES OF THE HEART AND AORTA**

By THOMAS HAYDEN, F.K.Q.C.P. Irel., Physician to the Mater  
Misericordiae Hospital, Dublin. With 80 Engravings. 8vo, 25s. [1875]

**PHTHISIS**

In a series of Clinical Studies, by AUSTIN FLINT, M.D., Professor of  
the Principles and Practice of Medicine and of Clinical Medicine in  
the Bellevue Hospital Medical College. 8vo, 16s. [1875]

**DISEASES OF THE HEART**

and of the Lungs in Connexion therewith—Notes and Observations  
by THOMAS SHAPTER, M.D., F.R.C.P. Lond., Senior Physician to the  
Devon and Exeter Hospital. 8vo, 7s. 6d. [1874]

**DISEASES OF THE HEART AND AORTA**

Clinical Lectures by GEORGE W. BALFOUR, M.D., F.R.C.P., Physician  
to, and Lecturer on Clinical Medicine in, the Royal Infirmary, Edin-  
burgh. 8vo, with Engravings, 12s. 6d. [1876]

**THE ACTION AND SOUNDS OF THE HEART**

Researches by GEORGE PATON, M.D., author of numerous papers  
published in the British and American Medical Journals. Re-  
issue, with Appendix, 8vo, 3s. 6d. [1874]

**PHYSICAL DIAGNOSIS OF DISEASES OF THE HEART.**

Lectures by ARTHUR E. SANSOM, M.D., Assistant Physician to the  
London Hospital; Physician to the North Eastern Hospital for  
Children. With Engravings, fcap. 8vo, 3s. 6d. [1876]

**NOTES ON ASTHMA;**

its Forms and Treatment, by JOHN C. THOROWGOOD, M.D. Lond.,  
F.R.C.P., Physician to the Hospital for Diseases of the Chest, Victoria  
Park. Second Edition, crown 8vo, 4s. 6d. [1873]

**GROWTHS IN THE LARYNX,**

with Reports and an Analysis of 100 consecutive Cases treated since  
the Invention of the Laryngoscope by MORELL MACKENZIE, M.D.  
Lond., M.R.C.P., Physician to the Hospital for Diseases of the  
Throat. 8vo, with Coloured Plates, 12s. 6d. [1871]

**TRACHEOTOMY,**

especially in Relation to Diseases of the Larynx and Trachea, by  
PUGIN THORNTON, Surgeon to the Hospital for Diseases of the  
Throat. With Photographic Plates and Woodcuts, 8vo, 5s. 6d. [1876]



**WINTER AND SPRING**

on the Shores of the Mediterranean; or, the Genoese Rivas, Italy, Spain, Greece, the Archipelago, Constantinople, Corsica, Sardinia, Sicily, Corfu, Malta, Tunis, Algeria, Smyrna, Asia Minor, with Biarritz and Arcachon, as Winter Climates. By HENRY BENNET, M.D. Fifth Edition, post 8vo, with numerous Plates, Maps, and Engravings, 12s. 6d. [1874]

BY THE SAME AUTHOR,

**TREATMENT OF PULMONARY CONSUMPTION**

by Hygiene, Climate, and Medicine. Second Edition, 8vo, 5s. [1871]

**PRINCIPAL HEALTH RESORTS**

of Europe and Africa, and their Use in the Treatment of Chronic Diseases. A Handbook by THOMAS MORE MADDEN, M.D., M.R.I.A., Vice-President of the Dublin Obstetrical Society. 8vo, 10s. [1876]

**MINERAL SPRINGS OF HARROGATE**

By Dr. KENNION. Revised and enlarged by ADAM BEALEY, M.A., M.D. Cantab., F.R.C.P. Lond. Seventh Thousand. Crown 8vo, 1s. [1875]

**SKETCH OF CANNES AND ITS CLIMATE**

by TH. DE VALCOURT, M.D. Paris, Physician at Cannes. Second Edition, with Photographic View and 6 Meteorological Charts. Crown 8vo, 2s. 6d. [1873]

**EGYPT AS A HEALTH RESORT;**

with Medical and other Hints for Travellers in Syria, by A. DUNBAR WALKER M.D. Fcap 8vo, 3s. 6d. [1873]

**FAMILY MEDICINE FOR INDIA**

A Manual, by WILLIAM J. MOORE, M.D., Surgeon-Major H.M. Indian Medical Service. Published under the Authority of the Government of India. Post 8vo, with 57 Engravings, 8s. 6d. [1874]

**DISEASES OF TROPICAL CLIMATES**

and their Treatment: with Hints for the Preservation of Health in the Tropics, by JAMES A. HORTON, M.D., Surgeon-Major, Army Medical Department. Post 8vo, 12s. 6d. [1874]

**HEALTH IN INDIA FOR BRITISH WOMEN**

and on the Prevention of Disease in Tropical Climates by EDWARD J. TILT, M.D., Consulting Physician-Accoucheur to the Farringdon General Dispensary. Fourth Edition, crown 8vo, 5s. [1875]

**BURDWAN FEVER,**

or the Epidemic Fever of Lower Bengal (Causes, Symptoms, and Treatment), by GOPAUL CHUNDER ROY, M.D., Surgeon Bengal Establishment. New Edition, 8vo, 5s. [1876]



**BAZAAR MEDICINES OF INDIA**

and Common Medical Plants: Remarks on their Uses, with Full Index of Diseases, indicating their Treatment by these and other Agents procurable throughout India, &c., by EDWARD J. WARING, M.D., F.R.C.P. Lond., Retired Surgeon H.M. Indian Army. Third Edition. Fcap 8vo, 5s. [1875]

**SOME AFFECTIONS OF THE LIVER**

and Intestinal Canal; with Remarks on Ague and its Sequelæ, Scurvy, Purpura, &c., by STEPHEN H. WARD, M.D. Lond., F.R.C.P., Physician to the Seamen's Hospital, Greenwich. 8vo, 7s. [1872]

**DISEASES OF THE LIVER:**

Lettsonian Lectures for 1872 by S. O. HABERSHON, M.D., F.R.C.P., Senior Physician to Guy's Hospital. Post 8vo, 3s. 6d. [1872]

BY THE SAME AUTHOR.

**DISEASES OF THE STOMACH: DYSPEPSIA**

Second Edition, crown 8vo, 5s.

**THE STOMACH AND DUODENUM**

Their Morbid States and their Relations to the Diseases of other Organs, by SAMUEL FENWICK, M.D., F.R.C.P., Assistant-Physician to the London Hospital. 8vo, with 10 Plates, 12s. [1868]

**FOOD AND DIETETICS**

Physiologically and Therapeutically Considered. By FREDERICK W. PAVY, M.D., F.R.S., Physician to Guy's Hospital. Second Edition, 8vo, 15s. [1875]

**IMPERFECT DIGESTION:**

its Causes and Treatment by ARTHUR LEARED, M.D., F.R.C.P., Senior Physician to the Great Northern Hospital. Sixth Edition, fcap 8vo, 4s. 6d. [1875]

**THE ISSUE OF A SPIRIT RATION**

during the Ashanti Campaign of 1874; with two Appendices containing Experiments to show the Relative Effects of Rum, Meat Extract and Coffee during Marching, and the Use of Oatmeal Drink during Labour, by EDMUND A. PARKES, M.D., F.R.S., Professor of Hygiene to the Army Medical School, Netley. 8vo, 2s. 6d. [1875]

**MEGRIM, SICK-HEADACHE,**

and some Allied Disorders: a Contribution to the Pathology of Nerve-Storms, by EDWARD LIVEING, M.D. Cantab., Hon. Fellow of King's College, London. 8vo, with Coloured Plate, 15s. [1873]

**FUNCTIONAL NERVOUS DISORDERS**

Studies by C. HANDFIELD JONES, M.B., F.R.C.P., F.R.S., Physician to St. Mary's Hospital. Second Edition, 8vo, 18s. [1870]



**NEURALGIA AND KINDRED DISEASES**

of the Nervous System : their Nature, Causes, and Treatment, with a series of Cases, by JOHN CHAPMAN, M.D., M.R.C.P. 8vo, 14s. [1873]

**THE SYMPATHETIC SYSTEM OF NERVES**

and their Functions as a Physiological Basis for a Rational System of Therapeutics by EDWARD MERYON, M.D., F.R.C.P., Physician to the Hospital for Diseases of the Nervous System. 8vo, 3s. 6d. [1872]

**GOUT, RHEUMATISM**

and the Allied Affections; a Treatise by P. HOOD, M.D. Crown 8vo, 10s. 6d. [1871]

**RHEUMATIC GOUT,**

or Chronic Rheumatic Arthritis of all the Joints; a Treatise by ROBERT ADAMS, M.D., M.R.I.A., Surgeon to H.M. the Queen in Ireland, Regius Professor of Surgery in the University of Dublin. Second Edition, 8vo, with Atlas of Plates, 21s. [1872]

**TEMPERATURE OBSERVATIONS**

containing (1) Temperature Variations in the Diseases of Children, (2) Puerperal Temperatures, (3) Infantile Temperatures in Health and Disease, by WM. SQUIRE, M.R.C.P. Lond. 8vo, 5s. [1871]

**THE ORIGIN OF CANCER**

considered with Reference to the Treatment of the Disease by CAMPBELL DE MORGAN, F.R.S., F.R.C.S., Surgeon to the Middlesex Hospital. Crown 8vo, 3s. 6d. [1872]

**CANCER:**

its varieties, their Histology and Diagnosis, by HENRY ARNOTT, F.R.C.S., Assistant-Surgeon to, and Lecturer on Morbid Anatomy at, St. Thomas's Hospital. 8vo, with 5 Plates and 22 Engravings, 5s. 6d. [1872]

**CANCEROUS AND OTHER INTRA-THORACIC GROWTHS:**

their Natural History and Diagnosis, by J. RISDON BENNETT, M.D., F.R.C.P., Member of the General Medical Council. Post 8vo, with Plates, 8s. [1872]

**CERTAIN FORMS OF CANCER**

with a New and successful Mode of Treating it, to which is prefixed a Practical and Systematic Description of all the varieties of this Disease, by ALEX. MARSDEN, M.D., F.R.C.S.E., Consulting Surgeon to the Royal Free Hospital, and Senior Surgeon to the Cancer Hospital. Second Edition, with Coloured Plates, 8vo, 8s. 6d. [1873]



**ATLAS OF SKIN DISEASES:**

a series of Illustrations, with Descriptive Text and Notes upon Treatment. By **TILBURY FOX**, M.D., F.R.C.P., Physician to the Department for Skin Diseases in University College Hospital. In monthly parts, roy. 4to, each containing Four Coloured Plates. Parts I to V issued since October. 6s. 6d. each. [1875]

**CERTAIN ENDEMIC SKIN AND OTHER DISEASES**

of India and Hot Climates generally, by **TILBURY FOX**, M.D., and **T. FARQUHAR**, M.D., Surgeon Major H.M. Bengal Medical Service (retired). Including Notes on Pellagra, Clou de Biskra, Caneotica, and Aleppo Evil (with 5 Plates), by **H. VANDYKE CARTER**, M.D., Surgeon Major H.M. Indian Medical Service. (Published under the sanction of the Secretary of State for India in Council). 8vo, 10s. 6d. [1876]

**DISEASES OF THE SKIN:**

a System of Cutaneous Medicine by **ERASMUS WILSON**, F.R.C.S., F.R.S. Sixth Edition, 8vo, 18s., with Coloured Plates, 36s. [1867]

BY THE SAME AUTHOR,

**LECTURES ON EKZEMA**

and Ekzematous Affections: with an Introduction on the General Pathology of the Skin, and an Appendix of Essays and Cases. 8vo, 10s. 6d. [1870]

ALSO,

**LECTURES ON DERMATOLOGY**

delivered at the Royal College of Surgeons, 1870, 6s.; 1871-3, 10s. 6d., 1874-5, 10s. 6d.

**ECZEMA**

by **MCCALL ANDERSON**, M.D., Professor of Clinical Medicine in the University of Glasgow. Third Edition, 8vo, with Engravings, 7s. 6d. [1874]

BY THE SAME AUTHOR,

**PARASITIC AFFECTIONS OF THE SKIN**

Second Edition, 8vo, with Engravings, 7s. 6d. [1868]

**PSORIASIS OR LEPRO**

by **GEORGE GASKOIN**, M.R.C.S., Surgeon to the British Hospital for Diseases of the Skin. 8vo, 5s. [1875]

**MYCETOMA ;**

or, the Fungus Disease of India, by **H. VANDYKE CARTER**, M.D., Surgeon-Major H.M. Indian Army. 4to, with 11 Coloured Plates, 42s. [1874]



**FOURTEEN COLOURED PHOTOGRAPHS OF LEPROSY**

as met with in the Straits Settlements, with Explanatory Notes by  
A. F. ANDERSON, M.D., Acting Colonial Surgeon, Singapore. 4to,  
31s. 6d. [1872]

**DISEASES OF THE SKIN**

in Twenty-four Letters on the Principles and Practice of Cutaneous  
Medicine, by HENRY EVANS CAUTY, Surgeon to the Liverpool Dis-  
pensary for Diseases of the Skin, 8vo, 12s. 6d. [1874]

**WORMS:**

a Series of Lectures delivered at the Middlesex Hospital on Practical  
Helminthology by T. SPENCER COBBOLD, M.D., F.R.S. Post 8vo,  
5s. [1872]

**THE MEDICAL ADVISER IN LIFE ASSURANCE**

by EDWARD HENRY SIEVEKING, M.D., F.R.C.P., Physician to St.  
Mary's and the Lock Hospitals; Physician-Extraordinary to the  
Queen; Physician-in-Ordinary to the Prince of Wales, &c. Crown  
8vo, 6s. [1874]

**THE LAWS AFFECTING MEDICAL MEN**

a Manual by ROBERT G. GLENN, LL.B., Barrister-at-Law; with a  
Chapter on Medical Etiquette by Dr. A. CARPENTER. 8vo, 14s.

[1871]

**MEDICAL JURISPRUDENCE**

(Principles and Practice of) by ALFRED S. TAYLOR, M.D., F.R.C.P.,  
F.R.S. Second Edition, 2 vols., 8vo, with 189 Engravings, £1 11s. 6d.

[1873]

BY THE SAME AUTHOR,

**A MANUAL OF MEDICAL JURISPRUDENCE**

Ninth Edition. Crown 8vo, with Engravings. 14s.

[1874]

ALSO,

**POISONS**

in Relation to Medical Jurisprudence and Medicine. Third Edition,  
crown 8vo, with 104 Engravings, 16s. [1875]

**A TOXICOLOGICAL CHART,**

exhibiting at one View the Symptoms, Treatment, and mode of  
Detecting the various Poisons—Mineral, Vegetable, and Animal:  
with Concise Directions for the Treatment of Suspended Animation,  
by WILLIAM STOWE, M.R.C.S.E. Thirteenth Edition, 2s.; on  
roller, 5s. [1872]

**OBSCURE DISEASES OF THE BRAIN AND MIND**

by FORBES WINSLOW, M.D., D.C.L. Oxon. Fourth Edition, post  
8vo, 10s. 6d. [1868]



**PSYCHOLOGICAL MEDICINE:**

a Manual, containing the Lunacy Laws, the Nosology, *Ætiology*, Statistics, Description, Diagnosis, Pathology (including Morbid Histology), and Treatment of Insanity, by J. C. BUCKNILL, M.D., F.R.S., and D. H. TUKE, M.D., F.R.C.P. Third Edition, 8vo, with 10 Plates and 34 Engravings, 25s. [1873]

**MADNESS**

in its Medical, Legal, and Social Aspects, Lectures by EDGAR SHEPPARD, M.D., M.R.C.P., Professor of Psychological Medicine in King's College; one of the Medical Superintendents of the Colney Hatch Lunatic Asylum. 8vo, 6s. 6d. [1873]

**HANDBOOK OF LAW AND LUNACY;**

or, the Medical Practitioner's Complete Guide in all Matters relating to Lunacy Practice, by J. T. SABBEN, M.D., and J. H. BALFOUR BROWNE, Barrister-at-Law. 8vo, 5s. [1872]

**CEREBRIA**

and other Diseases of the Brain by CHARLES ELAM, M.D., F.R.C.P., Assistant-Physician to the National Hospital for Paralysis and Epilepsy. 8vo, 6s. [1872]

**INFLUENCE OF THE MIND UPON THE BODY**

in Health and Disease, Illustrations designed to elucidate the Action of the Imagination, by DANIEL HACK TUKE, M.D., F.R.C.P. 8vo, 14s. [1872]

**A MANUAL OF PRACTICAL HYGIENE**

by E. A. PARKES, M.D., F.R.C.P., F.R.S., Professor of Hygiene in the Army Medical School. Fourth Edition, 8vo, with Plates and Engravings, 16s. [1873]

**A HANDBOOK OF HYGIENE**

for the Use of Sanitary Authorities and Health Officers by GEORGE WILSON, M.D. Edin., Medical Officer of Health for the Warwick Union of Sanitary Authorities. Second Edition, crown 8vo, with Engravings, 8s. 6d. [1873]

**MICROSCOPICAL EXAMINATION OF DRINKING WATER**

A Guide, by JOHN D. MACDONALD, M.D., F.R.S., Assistant Professor of Naval Hygiene, Army Medical School. 8vo, with 24 Plates, 7s. 6d. [1875]

**HANDBOOK OF MEDICAL ELECTRICITY**

by HERBERT TIBBITS, M.D., M.R.C.P.E., Medical Superintendent of the National Hospital for the Paralysed and Epileptic. 8vo, with 64 Engravings, 6s. [1873]



**CLINICAL USES OF ELECTRICITY**

Lectures delivered at University College Hospital by J. RUSSELL REYNOLDS, M.D. Lond., F.R.C.P., F.R.S., Professor of Medicine in University College. Second Edition, post 8vo, 3s. 6d.

[1873]

**AUTOBIOGRAPHICAL RECOLLECTIONS**

of the Medical Profession, being personal reminiscences of many distinguished Medical Men during the last forty years, by J. FERNANDEZ CLARKE, M.R.C.S., for many years on the Editorial Staff of the 'Lancet,' Post 8vo, 10s. 6d.

[1874]

**A DICTIONARY OF MEDICAL SCIENCE**

containing a concise explanation of the various subjects and terms of Anatomy, Physiology, Pathology, Hygiene, Therapeutics, Medical Chemistry, Pharmacology, Pharmacy, Surgery, Obstetrics, Medical Jurisprudence and Dentistry; Notices of Climate and Mineral Waters; formulæ for Officinal, Empirical, and Dietetic Preparations; with the Accentuation and Etymology of the terms and the French and other Synonyms, by ROBLEY DUNGLISON, M.D., LL.D. New Edition, by RICHARD J. DUNGLISON, M.D. Royal 8vo, 28s.

[1874]

**A MEDICAL VOCABULARY;**

being an Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, giving their derivation, meaning, application, and pronunciation, by ROBERT G. MAYNE, M.D., LL.D. Fourth Edition, fcap 8vo, 10s.

[1875]

**ATLAS OF OPHTHALMOSCOPY:**

representing the Normal and Pathological Conditions of the Fundus Oculi as seen with the Ophthalmoscope: composed of 12 Chromolithographic Plates (containing 59 Figures), accompanied by an Explanatory Text by R. LIEBREICH, Ophthalmic Surgeon to St. Thomas's Hospital. Translated into English by H. ROSBOROUGH SWANZY, M.B. Dub. Second Edition, 4to, £1 10s.

[1870]

**DISEASES OF THE EYE**

a Manual by C. MACNAMARA, Surgeon to Westminster Hospital. Third Edition, fcap, 8vo, with Coloured Plates and Engravings, 12s. 6d.

[1876]

**DISEASES OF THE EYE**

A Treatise by J. SOELBERG WELLS, F.R.C.S., Ophthalmic Surgeon to King's College Hospital and Surgeon to the Royal London Ophthalmic Hospital. Third Edition, 8vo, with Coloured Plates and Engravings, 25s.

[1873]

BY THE SAME AUTHOR,

**LONG, SHORT, AND WEAK SIGHT,**

and their Treatment by the Scientific use of Spectacles. Fourth Edition, 8vo, 6s.

[1873]



**DISEASES OF THE EYE**

A Practical Treatise by HAYNES WALTON, F.R.C.S., Surgeon to St. Mary's Hospital and in charge of its Ophthalmological Department. Third Edition, 8vo, with 3 Plates and nearly 300 Engravings, 25s.

[1875]

**OPHTHALMIC MEDICINE AND SURGERY**

a Manual by T. WHARTON JONES, F.R.S., Professor of Ophthalmic Medicine and Surgery in University College. Third Edition, fcap 8vo, with 9 Coloured Plates and 173 Engravings, 12s. 6d.

[1865]

**A SYSTEM OF DENTAL SURGERY**

by JOHN TOMES, F.R.S., and CHARLES S. TOMES, M.A., Lecturer on Dental Anatomy and Physiology, and Assistant Dental Surgeon to the Dental Hospital of London. Second Edition, fcap 8vo, with 268 Engravings, 14s.

[1873]

**A MANUAL OF DENTAL MECHANICS**

with an Account of the Materials and Appliances used in Mechanical Dentistry, by OAKLEY COLES, L.D.S., R.C.S., Surgeon-Dentist to the Hospital for Diseases of the Throat. Crown 8vo, with 140 Engravings, 7s. 6d.

[1873]

**HANDBOOK OF DENTAL ANATOMY**

and Surgery for the use of Students and Practitioners by JOHN SMITH, M.D., F.R.S. Edin., Surgeon-Dentist to the Queen in Scotland. Second Edition, fcap 8vo, 4s. 6d.

[1871]

**STUDENT'S GUIDE TO DENTAL ANATOMY AND SURGERY**

by HENRY SEWILL, M.R.C.S., L.D.S., Dentist to the West London Hospital. With 77 Engravings, fcap. 8vo, 5s. 6d.

[1876]

**EPIDEMIOLOGY;**

or, the Remote Cause of Epidemic Diseases in the Animal and in the Vegetable Creation, by JOHN PARKIN, M.D., F.R.C.S. Part I, 8vo, 5s.

[1873]

**DISEASE GERMS;**

and on the Treatment of the Feverish State, by LIONEL S. BEALE, M.B., F.R.C.P., F.R.S., Physician to King's College Hospital. Second Edition, crown 8vo, with 28 Plates, 12s. 6d.

[1872]

**THE GRAFT THEORY OF DISEASE**

being an Application of Mr. DARWIN's Hypothesis of Pangenesis to the Explanation of the Phenomena of the Zymotic Diseases, by JAMES ROSS, M.D. 8vo, 10s.

[1872]

**ZYMOTIC DISEASES:**

their Correlation and Causation by A. WOLFF, F.R.C.S. Post 8vo, 5s.

[1872]



The following CATALOGUES issued by Messrs CHURCHILL  
will be forwarded post free on application :

1. *Messrs Churchill's General List of nearly 600 works on Medicine, Surgery, Midwifery, Materia Medica, Hygiene, Anatomy, Physiology, Chemistry, &c., &c., with a complete Index to their Titles, for easy reference.*

N.B.—*This List includes Nos. 2 and 3.*

2. *Selection from Messrs Churchill's General List, comprising all recent Works published by them on the Art and Science of Medicine.*

3. *A descriptive List of Messrs Churchill's Works on Chemistry, Pharmacy, Botany, Photography, Zoology, and other branches of Science.*

4. *Messrs Churchill's Red-Letter List, giving the Titles of forthcoming New Works and New Editions.*

[Published every October.]

5. *The Medical Intelligencer, an Annual List of New Works and New Editions published by Messrs J. & A. Churchill, together with Particulars of the Periodicals issued from their House.*

[Sent in January of each year to every Medical Practitioner in the United Kingdom whose name and address can be ascertained. A large number are also sent to the United States of America, Continental Europe, India, and the Colonies.]

---

MESSRS CHURCHILL have a special arrangement with MESSRS LINDSAY & BLAKISTON, OF PHILADELPHIA, in accordance with which that Firm act as their Agents for the United States of America, either keeping in Stock most of Messrs CHURCHILL's Books, or reprinting them on Terms advantageous to Authors. Many of the Works in this Catalogue may therefore be easily obtained in America.



QM  
30  
H72  
1876







