

## **A clinical manual of the diseases of the ear / by Laurence Turnbull.**

### **Contributors**

Turnbull, Laurence, 1821-1900.  
Francis A. Countway Library of Medicine

### **Publication/Creation**

Philadelphia : Lippincott, 1872.

### **Persistent URL**

<https://wellcomecollection.org/works/a4t2ngbm>


### **License and attribution**

This material has been provided by This material has been provided by the Francis A. Countway Library of Medicine, through the Medical Heritage Library. The original may be consulted at the Francis A. Countway Library of Medicine, Harvard Medical School. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>





29. D. 128.  
No. ~~29. D. 128.~~

**BOSTON  
MEDICAL LIBRARY  
ASSOCIATION,  
19 BOYLSTON PLACE,**

---

Received.....


By Gift of.....

*E. H. Clarke M.D.*

















Digitized by the Internet Archive  
in 2012 with funding from  
Open Knowledge Commons and Harvard Medical School



Fig. 1.




Fig. 4.




Fig. 5.




Fig. 3.





Fig. 2.





# DESCRIPTION OF PLATE.

Figure I.—Temporal Bone (Front View).

- |                                        |                                          |
|----------------------------------------|------------------------------------------|
| 1. Squamous portion of temporal bone.  | 23. Posterior auricular artery.          |
| 2. Mastoid portion of temporal bone.   | 24. Branches to mastoid process.         |
| 3. Petrous portion of temporal bone.   | 26. Origin of internal maxillary artery. |
| 4. External auditory canal.            | 32. Portion of auriculo-temporal nerve.  |
| 5. Glasserian fissure.                 | 33. External auriculo-temporal nerve.    |
| 6. Zygomatic process.                  | 50. Portion of auriculo-occipital nerve. |
| 8. Styloid process.                    | 52. Exit of the chorda tympani nerve.    |
| 19. External carotid artery furnishing | 53. Anterior muscle of malleus.          |
| 20. Superficial temporal artery.       | 59. Eustachian tube.                     |
| 22. Stylo-mastoid artery.              |                                          |

Figure II.—Temporal Bone (Interior View).

- |                                        |                                     |
|----------------------------------------|-------------------------------------|
| 2. Portio dura (seventh pair).         | 12. Cavity semicircular canals.     |
| 3. Receptacle for semicircular canals. | 13. Attachment of membrana tympani. |
| 3a. Vidian nerve.                      | 14. Cavity for cochlea.             |
| 4. Genuiculated ganglion.              | 29. Internal meningeal artery.      |
| 5. Branches of auricular artery.       | 31. Nerve of M. Longuet.            |
| 7. Anastomosis of meningeal artery.    | 36. Portion of great petrous nerve. |
| 7a. Anastomosis of auricular artery.   | 48. Seventh pair of nerves.         |
| 11. Fallopian hiatus.                  | 49. External ear (anti-helix).      |

Figure III.—Semicircular Canals and Cochlea.

- |                                   |                                                  |
|-----------------------------------|--------------------------------------------------|
| 1. Superior semicircular canal.   | 11. Auditory nerve divided.                      |
| 2. Inferior semicircular canal.   | 12. Infundibulum.                                |
| 3. Posterior semicircular canal.  | 15. Promontory.                                  |
| 4. Section to show spiral lamina. | 18. Fenestrum rotunda.                           |
| 6. Fenestrum ovale.               | 21. Filaments of cochlear and vestibular nerves. |

Figure IV.—Open Utriculus, etc.

- |                                          |                                           |
|------------------------------------------|-------------------------------------------|
| 1. Open utriculus.                       | 10. Branch of superior vertical canal.    |
| 2. Open sacculus.                        | 11. Branch of horizontal canal.           |
| 3. Superior semicircular canal (memb.).  | 12. Branch of the saccule canal.          |
| 7. Calcareous substance.                 | 13. Branch of posterior vertical canal.   |
| 8. Branch of vestibular nerve furnishing | 14. Termination of nerve filaments to the |
| 9. Branch of utricle.                    | surface of membranous labyrinth           |
|                                          | held suspended in perilymph.              |

Figure V.—Membrana Tympani.

- |                                |                                 |
|--------------------------------|---------------------------------|
| 1. Membrane of the tympanum.   | 9. Base of stapes.              |
| 2. Incus.                      | 10. Malleus.                    |
| 3. Short branch.               | 11. Neck of malleus.            |
| 4. Long branch.                | 12. Short apophysis.            |
| 5. Lenticular bone.            | 13. Handle of malleus.          |
| 6. Stapes.                     | 14. Long process of malleus.    |
| 7. Anterior branch of stapes.  | 15. Tendon of muscle of stapes. |
| 8. Posterior branch of stapes. | 16. Chorda tympani nerve.       |





A  
CLINICAL MANUAL  
OF THE  
DISEASES OF THE EAR.


BY  
LAURENCE TURNBULL, M.D.,

Physician to the Department of Diseases of the Eye and Ear of Howard Hospital of Philadelphia; Permanent  
Member and Corresponding Secretary of the Medical Society of the State of Pennsylvania;  
Permanent Member of the American Medical Association,  
Author of "Lectures on the Electro-Magnetic Telegraph," "The Nature, Causes, and Treatment of  
Nervous Deafness," etc. etc.

With a Colored Lithographic Plate  
AND  
OVER ONE HUNDRED ILLUSTRATIONS ON WOOD.

---

PHILADELPHIA:  
J. B. LIPPINCOTT & CO  
1872.



---

Entered, according to Act of Congress, in the year 1871, by  
J. B. LIPPINCOTT & CO.,  
In the Office of the Librarian of Congress at Washington.

---



## PREFACE.

---

THE small work\* of the author being out of print, and a new edition called for, he has embraced the opportunity to enter more fully into the modern doctrines and practice of otologists. It has been his aim to make the work practically useful, and to lay down clear, broad principles for accurate diagnosis and successful treatment. He has not simply recorded the views and opinions of others, but has added the results of his personal experience in seventeen years private practice, and in the treatment of several thousand cases in the Aural Department of Howard Hospital.

The number of persons who are more or less deaf from age, neglect, or exposure, and who apply too late for successful treatment, is very much on the increase. Their condition has been described, and the best artificial aids for their use have been fully treated of and copiously illustrated.

The subject of rhinoscopy, or the examination of the posterior nares and the pharyngeal orifices of the Eustachian tubes, has received a due share of attention, while an endeavor has been made to represent the appearances of these parts by careful and accurate drawings from nature.

The mode of introducing the Eustachian catheter has

---

\* The Nature, Causes, and Treatment of Nervous Deafness. Philadelphia, 1863.



## PREFACE.

been dwelt upon, and two new forms, with drawings from nature, have been given, the results of numerous experiments on the cadaver, and upon the living subject. By the use of a new form of double-catheter the tube can be more readily washed when it becomes filled with mucus, or when medicated solutions are applied to the middle ear.

The author's object has been to present the subject of diseases of the ear in such a manner that every well-educated physician may approach their treatment with as much confidence as he would the diseases of the heart, lungs, brain, etc. The day has passed for medical men, as a rule, to affirm that they "know but little of the anatomy and physiology of the ear, and nothing about its diseases." A scientific reformation has taken place on the subject of otology, and some of the best men in the profession are devoting their time and talents to its study, in the extended field of the physiological and pathological connection between the organs of sight and hearing, so brilliantly illuminated by the researches of the Marquis of Corti, Profs. Schultz, Helmholtz, and Rüdinger. No small credit is due to the labors of James Jago, M.D., of England, whose original investigations have added much to our knowledge of the various movements which originate in the ear itself, or in adjacent structures, as well as to our knowledge of the true "functions of the tympanum," confirming some of the discoveries of the distinguished Joseph Toynbee, F.R.S., recently deceased, and correcting some of his errors. Advantage has been taken of the important observations of Mr. James Hinton, Aural Surgeon to Guy's Hospital, on nervous deafness; and also of his method of treatment for aural polypi, syphilitic affections of the ear, etc.

For those who desire to study the anatomy and physiology of the ear, more especially the younger members of



the profession, the author has had the beautiful model of Dr. Auzoux lithographed by Messrs. Sinclair, in order to fit the work for a text-book; and for the benefit of those whose active employment in general practice renders it difficult for them to obtain a more complete knowledge of the subject, he has added a résumé of the simplest and most effectual methods of treating aural catarrh and other more frequent and curable affections, which, without proper attention, are liable to terminate in deafness.

It has been the author's endeavor to represent all the more important instruments and apparatus employed in aural surgery, in which he has been kindly assisted by our chief surgical instrument-makers, Messrs. Kolbe and Gemrig, of this city, and several professional friends, who have placed numerous illustrations at his command. In composition and in illustration, he has availed himself of the works of Wilde, Toynebee, Tröltzsch, Moos, Politzer, Weber, Hinton, Triquet, Voltolini, Kramer, Gruber, Knapp, Schwarze, Green, Roosa, Clark, and others, as well as of the American and foreign medical periodicals having any bearing on the subject.

The author gratefully acknowledges his obligations to his friends, Prof. James A. Meigs, Drs. J. Solis Cohen, James Collins, and Charles F. Wittig, for valuable assistance in the preparation of his work for the press.

L. T.

1208 SPRUCE STREET, PHILADELPHIA,

October, 1871.



The first of these is the fact that the  
the second is the fact that the  
the third is the fact that the  
the fourth is the fact that the  
the fifth is the fact that the  
the sixth is the fact that the  
the seventh is the fact that the  
the eighth is the fact that the  
the ninth is the fact that the  
the tenth is the fact that the  
the eleventh is the fact that the  
the twelfth is the fact that the  
the thirteenth is the fact that the  
the fourteenth is the fact that the  
the fifteenth is the fact that the  
the sixteenth is the fact that the  
the seventeenth is the fact that the  
the eighteenth is the fact that the  
the nineteenth is the fact that the  
the twentieth is the fact that the  
the twenty-first is the fact that the  
the twenty-second is the fact that the  
the twenty-third is the fact that the  
the twenty-fourth is the fact that the  
the twenty-fifth is the fact that the  
the twenty-sixth is the fact that the  
the twenty-seventh is the fact that the  
the twenty-eighth is the fact that the  
the twenty-ninth is the fact that the  
the thirtieth is the fact that the  
the thirty-first is the fact that the  
the thirty-second is the fact that the  
the thirty-third is the fact that the  
the thirty-fourth is the fact that the  
the thirty-fifth is the fact that the  
the thirty-sixth is the fact that the  
the thirty-seventh is the fact that the  
the thirty-eighth is the fact that the  
the thirty-ninth is the fact that the  
the fortieth is the fact that the  
the forty-first is the fact that the  
the forty-second is the fact that the  
the forty-third is the fact that the  
the forty-fourth is the fact that the  
the forty-fifth is the fact that the  
the forty-sixth is the fact that the  
the forty-seventh is the fact that the  
the forty-eighth is the fact that the  
the forty-ninth is the fact that the  
the fiftieth is the fact that the  
the fifty-first is the fact that the  
the fifty-second is the fact that the  
the fifty-third is the fact that the  
the fifty-fourth is the fact that the  
the fifty-fifth is the fact that the  
the fifty-sixth is the fact that the  
the fifty-seventh is the fact that the  
the fifty-eighth is the fact that the  
the fifty-ninth is the fact that the  
the sixtieth is the fact that the  
the sixty-first is the fact that the  
the sixty-second is the fact that the  
the sixty-third is the fact that the  
the sixty-fourth is the fact that the  
the sixty-fifth is the fact that the  
the sixty-sixth is the fact that the  
the sixty-seventh is the fact that the  
the sixty-eighth is the fact that the  
the sixty-ninth is the fact that the  
the seventieth is the fact that the  
the seventy-first is the fact that the  
the seventy-second is the fact that the  
the seventy-third is the fact that the  
the seventy-fourth is the fact that the  
the seventy-fifth is the fact that the  
the seventy-sixth is the fact that the  
the seventy-seventh is the fact that the  
the seventy-eighth is the fact that the  
the seventy-ninth is the fact that the  
the eightieth is the fact that the  
the eighty-first is the fact that the  
the eighty-second is the fact that the  
the eighty-third is the fact that the  
the eighty-fourth is the fact that the  
the eighty-fifth is the fact that the  
the eighty-sixth is the fact that the  
the eighty-seventh is the fact that the  
the eighty-eighth is the fact that the  
the eighty-ninth is the fact that the  
the ninetieth is the fact that the  
the ninety-first is the fact that the  
the ninety-second is the fact that the  
the ninety-third is the fact that the  
the ninety-fourth is the fact that the  
the ninety-fifth is the fact that the  
the ninety-sixth is the fact that the  
the ninety-seventh is the fact that the  
the ninety-eighth is the fact that the  
the ninety-ninth is the fact that the  
the hundredth is the fact that the

# CONTENTS.

---

	PAGE
INTRODUCTION, AND CONGENITAL DEFECTS.....	13
CHAPTER I	
Anatomical Description of the Ear and its Functions.....	17
CHAPTER II.	
Physiology of Hearing.....	28
CHAPTER III.	
Acoustics.—Laws of Sound.....	43
Instruments to Assist Hearing.....	49
CHAPTER IV.	
Tables of Classification of Diseases of Ear.....	55
Method of Investigating Diseases of the Ear.....	60
CHAPTER V.	
On the Means of Diagnosis by the Use of Artificial and Daylight for Examining the Membrana Tympani and Auditory Canal—Preven- tion of Deafness—Noises in the Ear, etc.....	65
CHAPTER VI.	
Syringing the Ear for the Removal of Accumulations of Cerumen and Foreign Bodies.....	86
CHAPTER VII.	
Foreign Bodies in the External Auditory Meatus, and their Removal.....	93
Aspergillus in the Auditory Canal.....	104
CHAPTER VIII.	
Polypus of the Ear.....	110
Extraction.....	123
Cauterization .....	126
CHAPTER IX.	
Diseases and Injuries of the Auricle and External Meatus.....	131
Tumors, Abscesses, Hæmatoma, and Scirrhus.....	134
CHAPTER X.	
Pruritus—Acute and Chronic Eczema—Furunculous Abscess—Osseous Tumors—Otitis Externa.....	148



## CHAPTER XI.

	PAGE
Otitis Media Purulentia, Acute, Subacute, Chronic, and Hypertrophic (Suppuration in the Middle Ear).....	167

## CHAPTER XII.

Otitis Media and Interna involving Mastoid Process and Cells—Squam- ous Portion of Temporal Bone—Cerebrum and Cerebellum.....	189
----------------------------------------------------------------------------------------------------------------------------------	-----

## CHAPTER XIII.

Subacute, Complicated, and Chronic Inflammation of the Cavity of the Tympanum—Catarrh of the Middle Ear (Catarrhal Deafness).....	211
Pathology of Chronic Aural Catarrh.....	219
Catarrh of the Tympanum—Paracentesis of Membrana Tympani.....	238

## CHAPTER XIV.

On the Mechanical Appliances used in the Diagnosis and Treatment of Affections of the Eustachian Tube and Middle Ear.....	251
------------------------------------------------------------------------------------------------------------------------------	-----

## CHAPTER XV.

Wounds—Injuries—Perforations of the Membrana Tympani—Acute and Chronic Inflammation.....	278
---------------------------------------------------------------------------------------------	-----

## CHAPTER XVI.

Diseased Throat, Uvula, Tonsils, and Carious Teeth, in their Relations to Disease of the Ear; also Feigned Deafness.....	301
-----------------------------------------------------------------------------------------------------------------------------	-----

## CHAPTER XVII.

Nervous Deafness.....	316
Double Hearing.....	331

## CHAPTER XVIII.

Acute Inflammation of the Facial Nerve within the Fallopian Canal.....	361
Acute and Chronic Inflammation of the Labyrinth.....	364
Treatment of Nervous Deafness.....	368
The Treatment of Nervous Deafness by Electricity.....	373

## CHAPTER XIX.

Deaf Mutism.....	393
Institutions for the Instruction of the Deaf and Dumb in the United States	404
Treatment of Deaf-Mutism.....	408
A Tabular View of the Condition of the Ear in Sixty-five Dissections taken of Deaf Mutes.....	416
On the Choice of Methods for Teaching the Deaf Mute.....	426

## CHAPTER XX.

Résumé of the Simplest and most Effectual Method of Treating the more frequent Affections of the Ear.....	433
BIBLIOGRAPHY.....	453

## INDEX.



# LIST OF ILLUSTRATIONS.

COLORED LITHOGRAPHIC DRAWING FROM MODEL OF AUZOUX.

- FIG. 1. Temporal Bone. Front View.  
 2. Temporal Bone. Interior View.  
 3. Semicircular Canals and Cochlea.  
 4. Open Utriculus, etc.  
 5. Membrana Tympani and Bones of the Ear.  
 6. The Auricle.....*Sappey and Leidy.*  
 7. The Middle Ear.....*Milne Edwards, Carpenter.*  
 8. The Scala Vestibuli and Media Membrana, with Rods of Corti.  
     *Corti, Shultz, Kölliker, Helmholtz, and Gruber.*  
 9. Corti's Fibres in Natural Position.  
 10. Corti's Fibres lying over one another.  
 11. Corti's Organ. From the side.....*Deiters and Gruber.*  
 12. Simple Ear-Trumpet. From Nature.  
 13. Curved Ear-Trumpet.....*Gemrig.*  
 14. Haswell's Ear-Trumpet. From Nature.  
 15. Martineau's Ear-Trumpet. From Nature.  
 16. Conversation Ear-Tube.....*Gemrig's Catalogue.*  
 17. Artificial Auricles.....*Itard.*  
 18. Pocket Ear-Trumpet. From Nature.  
 19. Six Forms of French Ear-Trumpets.....*L'Acoustique, par R. Radau.*  
 20. Toynbee's Illuminating Ear-Speculum.....*Gemrig.*  
 21. Hinton's Modification. From Nature.  
 22. Weber's Reflecting Ear-Speculum.....*Weber.*  
 23. Gruber's Reflecting Ear-Speculum. Modified by Wilde.....*Kolbe.*  
 24. Kramer's Bivalve.....*Kramer.*  
 25. Von Tröltzsch's Reflecting Ear-Mirror.....*Von Tröltzsch, Roosa.*  
 26. Otoscope or Diagnostic Tube.....*Toynbee.*  
 27. Aural Manometer.....*Politzer and Roosa.*  
 28. The Double Otoscope. From Nature.  
 29. Catarrhal Syringe. From Nature.  
 30. Rauchfuss' Laryngeal Powder Insufflator.....*Gemrig.*  
 31. Steam Nebulizer.....*Kolbe's Catalogue.*  
 32. Air-Press.....*Moos.*  
 33. Apparatus for Generating Chloride of Ammonium. From Nature.  
 34. Siegel's Pneumatic Speculum. From Nature.  
 35. Davidson's Syringe, modified. From Nature.  
 36. Double-Ring Syringe. From Nature.  
 37. Ear-Syringe (small). From Nature.  
 38. Ear-Douche.....*Clark's.*  
 39. Ear-Syringe for Patients' Use.....*Wilde, Hewson.*  
 40. Ear-Basin, Wilde's.....*Wilde.*  
 41. Ear-Spout.....*Toynbee.*  
 42. Curette.....*Gross.*  
 43. Author's Ear-Curette. From Nature.  
 44. Author's Angular Forceps. From Nature.  
 45. Angular Forceps.....*Wilde.*  
 46. Epithelial or Raspberry Polypus.....*Toynbee.*  
 47. Fibro-Gelatinous Polypus. From Nature.  
 48. Polypus Snare.....*Wilde.*



FIG. 49. Ring-and-Lever Forceps.....	<i>Toynbee.</i>
50. French Modification of Wilde's Snare.....	
51. English Forceps (Polypus).....	<i>Gemrig.</i>
52. American Forceps (Polypus).....	<i>Kolbe.</i>
53. Ear-Scissors for removal of Hairs, etc.....	<i>Kolbe.</i>
54. Galvano-Cautery.....	<i>Cohen.</i>
55. Thudichum's Nasal Douche.....	<i>Gemrig.</i>
56. Modification of Thudichum's Nasal Douche.....	<i>Gemrig.</i>
57. Tumor of the Lobe of the Ear. From Nature.....	
58. Hagen's Blunt Bistoury, with Spoon.....	<i>Hagen.</i>
58. Osseous Tumor of Left Ear. From Nature.....	
59. Three Osseous Tumors of Right Ear. From Nature.....	
60. Pyogenic Membrane containing Pus. From Nature.....	
61. Necrosis of Temporal Bone. From Nature.....	
62. Destruction of Temporal Bone by Necrosis. Anterior View. From Nature.....	
63. Destruction of Temporal Bone by Necrosis. Posterior View. From Nature.....	
64. Pomeroy's Glass Attachment for the use of Iodine, etc.....	<i>Gemrig.</i>
65. Section of Thickened Mucous Membrane of the Middle Ear.....	<i>Gruber.</i>
66. Instrument for Puncturing the Membrana Tympani.....	<i>Kolbe.</i>
67. Instrument for Incising the Membrana Tympani.....	<i>Kolbe.</i>
68. Eyelets for keeping open Perforation of Membrana Tympani.....	
69. Eyelets for keeping open Perforation of Membrana Tympani.....	
70. Rhinoscopic Mirror.....	<i>Cohen.</i>
71. The Illuminating Apparatus of Lewin, with an Auto-laryngoscopic Mirror.....	<i>Kolbe.</i>
72. Tobold's Illuminating Apparatus, with Stand (Gas).....	<i>Cohen.</i>
73. View of a Section of the Head, with Rhinoscopic Mirrors in Position, altered from Czermak.....	
74. Rhinoscopic Mirrors.....	<i>Wales.</i>
75. Türk's Palate Noose.....	<i>Türk.</i>
76. Wales' Palate Hook.....	<i>Wales.</i>
77. Clamp for Bridge of the Nose.....	<i>Moos.</i>
78. Tongue-Depressor.....	<i>Cohen.</i>
79. View of the Openings of Eustachian Tubes. From Nature.....	
80. Rhinoscopic Image.....	<i>Cohen.</i>
81. Openings of Eustachian Tubes and Rosenmüller's Fossa, seen by Rhinoscopic Mirror.....	<i>Wales.</i>
82. Thudichum's Nasal Speculum. Front View.....	<i>Wales.</i>
83. Thudichum's Nasal Speculum. Profile View.....	<i>Wales.</i>
84. Dr. Cutter's Instrument for Catheterizing the Eustachian Tube through the Mouth.....	<i>Cutter.</i>
85. Eustachian Catheter (full size).....	<i>Kramer.</i>
86. Eustachian Catheter (small size).....	<i>Kolbe.</i>
87. Section of the Head, with Eustachian Catheter introduced into the Eustachian Tube. From Nature.....	
88. Eustachian Tube and Explorer.....	<i>Gemrig.</i>
89. Blowing Air into the Middle Ear and Diagnostic Tube introduced.....	<i>Kramer.</i>
90. Modified Catheters for Eustachian Tube. From Nature.....	
91. Politzer's Air-Bag introduced by the Surgeon.....	<i>Moos.</i>
92. Politzer's Air-Bag, modified by Brunton.....	<i>Brunton.</i>
93. Politzer's Air-Bag, modified by Allen.....	<i>Allen.</i>
94. Double Eustachian Catheter for Cleansing the Middle Ear. From Nature.....	
95. Bishop's Nebulizer.....	<i>Bishop.</i>
96. Section of a Fatty Degenerated Tympanic Membrane.....	<i>Gruber.</i>
97. Platinum Ladle for Melting Nitrate of Silver.....	<i>Wilde.</i>
98. Livingston's Elastic Caustic Probe.....	<i>Wilde.</i>
99. Toynbee's Artificial Membrana Tympani.....	<i>Toynbee.</i>
100. Fahnstock's Tonsil Instrument.....	<i>Charrière.</i>
Fahnstock's Tonsil Instrument. Another View.....	<i>Charrière.</i>
101. Modification of Pulvermacher's Galvanic Chain.....	<i>Dr. Hackley.</i>
102. Kidder's Electro-Magnetic Apparatus.....	<i>Dr. Kidder.</i>
103. Magneto-Electric Machine of Sexton.....	<i>Sexton.</i>
104. Magneto-Electric Machine of Prof. Page.....	<i>Prof. Page.</i>
105. Rheophore of Duchenne.....	<i>Duchenne.</i>
106. Deaf-and-Dumb Alphabet.....	<i>Hutton.</i>
107. Compact Box containing all of the Instruments for Treating Ordinary Cases of Diseases of the Ear.....	<i>Kolbe.</i>



## INTRODUCTION.

---

A RECENT writer in one of our popular magazines\* makes the following statement: "That the knowledge of the ear, its diseases, and their remedies, seem scarcely to have kept pace with the progress made in the other departments of medical science." The skeptical state of certain portions of the medical profession† has had its fruits in a like condition of the mind of the people at large, actually preventing them, in many instances, from receiving aid from those who have devoted both time and talents to the study of an obscure and intricate portion of the human system. A recent instance has occurred in the writer's practice showing its pernicious influences. A son of a wealthy gentleman of this city, aged seventeen, had scarlet fever at the age of *twelve* years, with severe inflammation of the throat, ear, soft palate, and tonsils, producing deafness, with purulent discharge, first from the right ear, and subsequently from both, and, finally, from the nose. He never received any systematic treatment, and even when he became offensive not only to himself, but to his whole family, was *directed to let the disease alone!* Such was the advice of two old and distinguished physicians, also of a celebrated surgeon. And why? No reasons were given, except that it might go to his brain. Fortunately for this youth, a brother studied medicine, and as soon as he began to practice, his first desire was to acquire a knowledge of diseases of the ear, to improve his brother's condition so as to fit him to enter into society. With this view he solicited the writer's aid and counsel. Active and regular treatment of the nose, tonsils, and mucous membrane of his throat was instituted, keeping the ear clean,

---

\* Our Monthly.

† Boylston Prize Essay, 1870.



and opening the closed Eustachian tube. Within three months the discharge from the ear was under control, the nose was well, the tonsils reduced one-half, the voice had lost much of the nasal twang, and the whole aspect of the boy was brighter and better; and instead of keeping to himself, he became able to mix in society, and could hear general conversation. Now, what would he have been if this had gone on for years longer? We are almost sure it would have terminated in pyæmia or phthisis.

The knowledge of the anatomy, physiology, pathology, and treatment of diseases of the human ear is in advance of that of many other organs of the human body, and the works upon the subject at the present time are scientific, clear, and far ahead of those of the last eighteen years, as may be seen by a reference to the bibliography of the subject, and the tables, of success in treatment, appended to this work; for, as well expressed by Prof. Roosa,\* of New York, in reference to the cases presented in these tables which were taken without selection, as they occurred in hospital, clinic, or private practice, "those only are classified as '*cured*' in which the hearing was restored to a normal condition, as tested by the watch, the tuning-fork, and ordinary conversation; and when the membrana tympani (if previously ulcerated) has been healed. Judged by the ordinary use of the term *cured*, this standard is too high. It is higher, for example, than that in ophthalmic practice, where the removal of a cataractous lens—so that the patient gets sight enough to read coarse print—is called a cure of the cataract, although the vision obtained may be only two-thirds of that enjoyed by a person who has never had cataract. In the same manner we speak of *curing* a bone affected with necrosis, by removing the diseased portion, even if considerable deformity be left. If a patient who has for years suffered from an ulcerative process in the cavity of the tympanum recovers to such an extent that the ulcer is healed, and the hearing power improved, he would be said to be cured, under a standard no higher than that usually adopted in medical statistics."

---

\* Roosa, Report of 494 Cases of Disease of the Ear, Am. Jour. Med. Sci., April, 1871.



## CONGENITAL DEFECTS OF THE EAR.

In the body of our work no special reference is made to the subject of congenital defects of the ear. Such cases are rare, and almost always irremediable. In the first class are those in which there is an entire absence of the auricle or in which it exists in a rudimentary state.

The second form consists in the absence not only of the auricle but of the meatus, the position of which is designated only by a slight depression, and yet, in some cases of this kind in which both meatuses were in this condition, the individuals could hear. A similar case is reported in Cooper's *Surgical Dictionary*, seventh edition, page 470, of a child who was wholly destitute of external ears, and in whom no meatus auditorius was visible, still the child could hear, but very imperfectly.

Prof. Allen Thomson published, in the *Edinburgh Journal of Medical Sciences* for April, 1847, a record of some cases of malformation of the external ear, and experiments on the condition of the hearing of the individuals. Three dissections have been made of the class of cases under consideration: one is described by Prof. Jaeger, of Erlangen, one in the museum of the University of Edinburgh, and the third by the late Joseph Toynbee, F.R.S. In the first two instances one ear only was affected. In both (quoting Dr. Thomson) the labyrinth appeared to be quite naturally formed; the cavity of the tympanum and the bony Eustachian tube existed, but were much smaller than usual. The chain of ossicles differed materially from the natural structure, being united (in one of the examples) into one straight and simple piece. The most striking departure from the normal form of the bone consisted in the entire obliteration of the meatus externus, which seemed to be connected with the absence of that portion of the temporal bone which forms the tympanic ring and lower side of the bony canal of the meatus, and the extension backward of the articular or true glenoid portion of the temporal bone to twice its natural breadth. In the late Mr. Toynbee's\* case

---

\* *The Disease of the Ear*, by Jos. Toynbee, F.R.S., with a Supplement, by James Hinton, M.R.C.S. London: H. K. Lewis, 1868, pp. 16, 17.



no meatus or membrana tympani was detected; but in their place a flat surface of bone. Two bones were found, which he considered the analogues of the malleus and stapes. The tensor tympani muscle, and also the Eustachian tube, which opened into the anterior part of the tympanic cavity, were present. The auditory nerve, cochlea, vestibule, and semi-circular canals were healthy.

In these cases there was a certain amount of hearing, which is contrary to the received idea that absence of the meatus externus is always followed with rudimentary formation or absence of the essential portions of the inner ear.

*Treatment.*—A merely cutaneous or cartilaginous closure, with contraction of the osseous wall, may be relieved by opening in the canal with the knife, and subsequent dilatation by bougies of laminaria or compressed sponge; and in this way we may restore the hearing of the patient.

In a recent successfully treated case of this kind of "complete cutaneous occlusion of the left meatus auditorius externus, with hyperostosis of the osseous portion," Dr. W. W. Morland,\* of Boston, after etherizing the patient, made a crucial incision of the membrane, closing the meatus, and removed the four resulting flaps with small curved scissors. He then inserted a piece of compressed sponge, which was kept up until the parts healed, and then had a gold tube made so as to fit the auditory canal with a funnel-shaped opening. The patient's hearing improved, and, as a consequence, the speech also, which had previously been very defective.

---

\* Transactions of the Am. Otol. Society. New York: D. Appleton & Co., 1870. See, also, Hyperostosis of the Cranium and Petrous Bones, by Prof. S. Moos, of Heidelberg, Arch. of Oph. and Otology, vol. ii. No. 7. Also, Symmetrical Formation of Exostosis in both External Auditory Canals of Man, pp. 127, 136, 137.



# DISEASES OF THE EAR.

## CHAPTER I.

### ANATOMICAL DESCRIPTION OF THE EAR AND ITS FUNCTIONS.

THE ear of man is one of the most beautiful and perfect of all his organs. Its most wondrous parts are deeply hidden in the petrous portion of the temporal bone. It is thus protected from external violence, and moreover its mysterious vital actions are thus excluded from the gaze of man.

The ear is divisible into three portions, viz.: the external, middle, and internal.

The external ear or auricle has for its office the collection and augmentation of sounds. It resembles the expanded mouth of a trumpet crushed inwardly, and consists mainly of fibro-cartilage. Its lower pendent portion, called the lobe, consists of connective tissue and fat. To this auricle a thin skin is tightly attached. Connected with it are five small muscles, between the cartilage and the skin, and three larger ones connecting it with the head.


It is divided as seen under Fig. 6.

It is well supplied with blood-vessels from branches of the temporal and external carotid arteries. Its nerves are derived from the great auricular branch of the cervical plexus, the posterior auricular branch of the facial nerve, and the auriculo-temporal branch of the inferior maxillary nerve. Sounds are conducted from the auricle through the external auditory meatus. The latter is a tortuous canal of about an inch in length, directed upward and forward, making a slight turn backward, and finally dips downward and forward to its termination, lined with a skin which is abundantly supplied



with ceruminous glands, which secrete a bitter waxlike substance called cerumen. This, with numerous projecting hairs, serves to exclude foreign matters, which would otherwise interfere with the function of hearing. The meatus is for

Fig. 6.—THE AURICLE.



1, helix; 2, fossa of the helix; 3, antihelix; 4, fossa of the antihelix; 5, concha; 6, tragus; 7, antitragus; 8, entrance of the external auditory meatus; 9, the lobe. The large dots in the concha and fossa of the antihelix are the orifices of sebaceous glands.

about one-third of its length cartilaginous, and in the remaining two-thirds is osseous. These two sections are joined by a membranous layer, giving it the property of extensibility. The anterior wall of the osseous meatus forms the posterior wall of the fossa of the inferior maxilla, and its posterior wall is made up of the mastoid process, so that the canal is separated from the transverse sinus only by two thin plates of bone and the air-cells between them. This canal is closed at its lower extremity by the membrana tympani or "drum of the ear," which separates completely the external from the middle ear, still further occluding the entrance to insects, etc. This is a circular partition,

not flat, but slightly funnel-shaped, its outer depressed surface being directed downward and inward. The greater part of its circumference is inserted into a fine groove in the osseous portion of the canal. The size of the membrane depends upon this osseous ring. The ratio of the length to the breadth in the adult is as four and three-tenths lines to four lines. The angle of inclination to the lower wall of the auditory canal is  $50^{\circ}$ ; that of the plane of the membrane with the upper wall is  $140^{\circ}$ .

This membrane is thin and translucent, and is composed of three layers: an external dermoid, a middle fibrous, and an internal mucous. The external or dermoid layer is really a delicate continuation of the integument lining the auditory canal; the middle or fibrous is composed of several



layers bound together by circular and longitudinal fibres. The internal or mucous layer is a continuation of the lining membrane of the middle ear or tympanum. On examining the membrane from without, we notice a yellowish-white line. This is the handle of the malleus, which is inserted between the laminae of the fibrous layer. According to Politzer, the color of the membrane is a combination, depending on various factors, viz., the kind of light employed, and the shade of the rays reflected from the promontory. Viewed by daylight, it presents a neutral gray mingling with a tint of violet and a light yellowish brown. If examined by artificial light, it acquires an orange hue. It has a peculiar lustre, and has at the anterior inferior portion a bright spot shaped like an isosceles triangle. This was first described by Mr. Wilde, of Dublin. This curious partition is alternately relaxed and made tense by the action of two small muscles, the tensor and laxator tympani.

Helmholtz has shown that this membrane, while it diminishes the amplitude of the waves of vibration, being curved, increases their force.

The cavity of the middle ear or tympanum (see Fig. 7, H) is filled with ever-renewed air, and is deeply situated in the temporal bone. It measures about two lines from the membrana tympani inward, its breadth and height being about half an inch; and Von Trötsch compares it to a hexahedron or cube. The horizontal diameter is therefore the greatest. It is about 13 mm. The height or the vertical diameter anteriorly at the tympanic orifice of the tube is only from 1 to 8 mm.; further back at the malleus it is 15 mm. The distance from the membrana tympani to the labyrinth wall is the least. At the mouth of the tube it is from 3 to  $4\frac{1}{2}$  mm., but if we measure somewhat farther back in the vertical plane of the malleus we get only 2 mm., at the end of the handle of the malleus, which projects very much into the cavity of the tympanum. This is of course the most convex portion of the drum corresponding to the most concave portion externally—the umbo. This chamber is connected directly with the throat by a canal called the Eustachian tube, through which a current of air passes at every act of deglu-



tition. The cavity of the tympanum is lined with a continuation of the mucous membrane of the Eustachian tube. This lining is provided with a tessellated epithelium, while in the Eustachian tube it is ciliated. According to Tröltsch, the membrane lining the tympanum takes the place of a periosteum by transmitting the blood-vessels which supply the bones. This latter fact is important to notice, as any serious affection of this membrane will ultimately react upon the nutrition of the bones forming this cavity, thus resolving every catarrh into a periostitis. If the inflammation become chronic, there is a great tendency to thickening of the mucous membrane and to hypertrophy of the bones. The existence of a mucous cushion in the cavity of the tympanum was first demonstrated by Tröltsch, and is the reason why affections of the middle ear are so numerous in young children. The close contact of the jugular vein to the cavity of the tympanum exposes it to the influence of pus collecting on its floor.

The Eustachian tube is a trumpet-shaped canal, one and a half inches in length, extending from the forepart of the tympanum, obliquely inward, forward, and downward. The upper third, which is connected directly with the tympanum, is an osseous canal, formed in the petrous portion of the temporal bone. The lower two-thirds, communicating with the pharynx, are cartilaginous, and, gradually expanding in their course, terminate in an oval orifice having a prominent border situated at the side of the pharynx, which is seen at Fig. 7, K. It is on a level with the turbinated bone, and therefore a little higher than the floor of the nasal cavity. In children this tube is much shorter and nearly horizontal. In the adult it is narrowest where the cartilaginous portion joins the bone.

The muscles of the tube are, first, the tensor palati, which arise in the fossa, at the root of the internal pterygoid process of the sphenoid bone, and some of its fibres have their origin in the cartilaginous walls of this tube. The contraction of the muscle in the act of swallowing causes the expansion of the portion of the tube opening into the pharynx. The levator palati arises from the under surface of the petrous



portion of the temporal bone, its fibres lying in close contact with the cartilaginous portion of the tube, but not arising from it.

These muscles have their insertion in the soft palate. In raising the soft palate, these muscles act by constricting the pharynx and elevating the velum palati. By this foreign bodies are prevented from finding their way into the upper part of the pharynx and occluding the Eustachian tubes, as in vomiting, eructations; and yet the walls of the tube lie so lightly against each other that in eructations the wall will separate so that air enters the ear.

“The relations and attachments of the muscles of the Eustachian tube have been recently demonstrated very satisfactorily by Rüdinger.\* The tensor-palati muscle is a dilator of the tube; it is inserted along the whole length of the hook of the cartilage, passes forward, inward, and slightly downward, and its fibres spread out along the edge of the soft palate and on the side of the pharynx. In contracting, it draws the hook of the cartilage forward and a little downward, thus enlarging the calibre of the tube. The levator palati takes its origin from the temporal bone just below the osseous tube, and passes along the floor of the tube, some of its fibres arising from the lower end of the cartilage; it is inserted in the uvula, and, in contracting the belly of the muscle which lies along the floor of the tube, becomes thicker; the floor of the tube is raised, and the fibres rising from the cartilage serve to draw the lower end of this away from the opposite wall.

“The palato-pharyngeus rises from the posterior part of the lower end of the cartilage, passes backward, and is inserted on the posterior wall of the pharynx. Its action would be to draw the posterior wall of the tube backward; but, as it is often but slightly developed, it probably only serves to fix the cartilage, so that the other muscles can act more effectively.

“The opening of the tube is thus the result of the action of

---

\* Beiträge zur vergleichenden Anatomie und Histologie der Ohrtrumpete. München, 1870. American Otological Society Transactions, 1870, pp. 9, 10.



these three muscles; the tensor palati, or dilatator tubæ, draws the hook of the cartilage outward, the cartilage becomes less curved and the tube is widened; the levator palati in contracting becomes more horizontal, and draws the lower end of the cartilage inward and upward, thus enlarging the pharyngeal orifice more than 3'''. As soon as these muscles cease acting, the elasticity of the cartilage restores the canal to its former condition.

“The papillæ on the dermoid layer of the membrana tympani, first described by Popper as a normal condition, and mentioned in Dr. Roosa's report of last year (1869), are of interest in connection with the polypoid growths which are sometimes found on the surface of the membrane, and which are undoubtedly hypertrophies of these normal papillary projections.”

We have already seen, in speaking of the membrana tympani (Plate, Fig. 5), the handle of the malleus, one of a chain of small bones connecting the membrana tympani with the oval window or fenestra ovalis. The malleus or hammer-like bone is situated vertically, with its head lodged in the upper part of the tympanum, and its handle, as before described, inserted between the fibrous layers of the membrana tympani. Its head is round and invested with a cartilage for articulation with the second bone.

The malleus has two important processes arising from its neck: the longer one (Plate, Fig. 5, No. 14) is a slender spine projecting nearly at right angles from the neck, and enters the glenoid fissure; the shorter one is a conical eminence at the root of the neck (Plate, Fig. 5, No. 12). The incus or anvil is situated behind the malleus, near the entrance to the mastoid sinuses (Plate, Fig. 5, No. 2). It has also two processes: the short one (Plate, Fig. 5, No. 3), which projects backward, and is connected by a ligamentous band to the posterior part of the tympanum; the long one is curved and tapering (Plate, Fig. 5, No. 4), and descends nearly parallel to the handle of the malleus. Its end, internally, supports an orbicular process (lenticular bone), which articulates with the head of the third bone, the stapes or stirrup. This of all the bones in the body most



resembles the object after which it is named. Its head is flattened, and at the summit possesses a concave articular process for attachment with the incus. All of these bones have movable articulations, surrounded with capsular ligaments and lined with synovial membranes. Between the malleus and incus the joint is hingelike, and between the incus and stapes it is a ball-and-socket joint. These bones are furnished with three small muscles of the striated variety: the stapedius, laxator tympani, and tensor tympani.

The stapedius muscle has its origin in the hollow of the pyramid, and is inserted into the head of the stapes, and regulates the pressure of the latter against the fenestra ovalis. The laxator tympani arises from the spinous process of the sphenoid bone, passes upward and outward, and enters the glenoid fissure, to be inserted into the long process of the malleus. The tensor tympani arises from the contiguous surfaces of the sphenoid and temporal bones, and the end of the cartilage of the Eustachian tube passes through a canal in the temporal bone above the tube and ends in a tendon which enters the tympanum, and is inserted into the neck of the malleus.

The cavity of the tympanum receives its nourishment through the stylo-mastoid branch of the posterior auricular artery (external carotid), which, in its course through the Fallopian canal, also gives branches to the stapedius muscle, coverings of the facial nerve, and mastoid cells. The ascending pharyngeal artery supplies the lining membrane of the tympanum, the Eustachian tube, and the tensor tympani muscle. The middle meningeal artery (internal maxillary), with one or two branches from the internal carotid, are distributed to the tympanic cavity and the Eustachian tube. The nerves of the middle ear are derived from the trigeminal, the facial, the glosso-pharyngeal, the vagus, and great sympathetic. The otic ganglion and the chorda tympani also supply this cavity.

The internal portion of the ear is the most deeply situated of the three parts, and comprises the labyrinth and the internal auditory meatus. The labyrinth is so named from its exceedingly complex character. It is the most important



portion of the organ of hearing, containing as it does the entire distribution of the auditory nerve. It is imbedded in the petrous portion of the temporal bone, and consists of three parts: the vestibule, the semicircular canals, and the cochlea. These may be described as cavities, but it should be remembered that the labyrinth has osseous walls independent of the bony structure inclosing it. The vestibule is an irregularly oval cavity situated between the tympanum and the lower part of the internal auditory meatus, joining the semicircular canals postero-externally and cochlea antero-internally. It communicates with the tympanum through the fenestra ovalis. From the floor of the vestibule springs a slight semicircular crest. This ascends on its inner wall to the roof and terminates in a small pyramidal eminence. This latter presents a group of minute foramina communicating with the internal auditory meatus, and is called the superior cribriform spot. The crest separates two recesses, which, from their shape, are named hemispherical and hemi-elliptical fossæ. A second group of foramina below the centre of the hemispherical fossa communicating with the internal auditory is named the middle cribriform spot. Into the hemi-elliptical fossa open the orifices of the semicircular canals.

The semicircular canals are three in number, and are situated above the inner back part of the tympanum and postero-externally to the vestibule. They are named from their position the superior, posterior, and inferior (Plate, Fig. 3, Nos. 1, 2, 3). The first two are vertical and the last is horizontal. Each canal is rather more than half a circle, communicating with the vestibule by a flasklike dilatation called the ampulla. Two of the undilated extremities conjoin, and with the remaining one likewise open into the vestibule, thus communicating with the latter by five orifices. The ampulla of the posterior semicircular canal has within it a group of minute foramina which communicate with the internal auditory meatus, and is named the inferior cribriform spot. The vestibule and semicircular canals are lined with a delicate membrane of a serous character, which surrounds a secondary structure, the membranous labyrinth (Plate,



Fig. 4), and secretes a liquid called the perilymph. This consists of a pair of communicating pouches occupying the vestibule and semicircular canals and of the same form as the osseous canals containing them. The smaller or spherical saccule (No. 2) occupies the hemispherical fossa; the other or elliptical saccule occupies the hemi-elliptical fossa, and is joined by three membranous semicircular canals (Nos. 11, 12). The membranous labyrinth is composed of an outer fibrinous layer and an inner tessellated epithelium, containing a serous liquid called the endolymph. The vestibular branch of the internal auditory nerve is distributed to the membranous labyrinth, and maintains its floating position in the perilymph. This nerve divides at the bottom of the internal auditory meatus into three branches, one of which, entering the vestibule at the superior cribriform spot, is distributed to the elliptical saccule and the ampullæ of the superior and inferior semicircular canals.

A second, entering at the middle cribriform spot, is distributed to the hemispherical saccule.

The third, entering at the inferior cribriform spot, is distributed to the ampulla of the posterior semicircular canal. Adhering to the inner surface of the spherical and elliptical saccules, where the nervous filaments from the cribriform spots enter, are two white discoidal masses, called otoliths. These appear to be in contact with the nervous filaments of the saccules (Plate, Fig. 4, No. 7). The arteries of the vestibule and semicircular canals are derived from the vestibular branches of the auditory artery which accompany the branches of the vestibular nerve. The *cochlea* (Plate, Fig. 3) is the innermost portion of the labyrinth, and is so named from its resemblance to a snail's shell. Its base is applied to the bottom of the internal auditory meatus, and its apex is directed forward and slightly outward. This osseous tube is an inch and a half in length, wound nearly three times around a central axis, each turn successively rising. The beginning of this tube is connected with the front part of the vestibule, forming the promontory\* of the tympanum. It

---

\* The promontory is a convex eminence on the inner wall, formed by the projection of the cochlea. Back of the upper part of the promontory is the



turns from right to left in the right ear and from left to right in the left, and ends in a closed extremity named the cupola. The axis of the cochlea is a cone whose base forms a spiral tract at the bottom of the internal auditory meatus, and whose apex becomes continuous with the inner wall of the last turn of the canal of the cochlea. It is traversed by numerous fine canals continuous with the foramina of the spiral tract, which transmit the filaments of the cochlear nerve; also cochlear branches of the auditory artery. A central canal extends the length of the axis, transmitting a branch of the latter vessel. An osseous spiral lamina commences below the hemispherical fossa of the vestibule, and winds around the axis of the cochlea and extends about half way across its canal, terminating in a pointed process at its apex. The cochlea is lined by a delicate membrane continuous with and similar to that of the vestibule. This, after investing the free surfaces of the osseous spiral lamina, is extended in a double layer from the free border of the latter across the canal of the cochlea, where it is firmly attached to its outer wall. This extension of membrane, which is named the membranous spiral lamina, together with the osseous spiral lamina, divides the canal of the cochlea into two distinct passages, communicating with each other only at the summit of the cochlea. At the lower extremity of the cochlea, one of these passages terminates at the fenestra rotunda, and is named the tympana scala, while the other opens into the vestibule, and is called the vestibular scala. Both scalæ are filled with a serous liquid, which communicates with the perilymph of the vestibule, but is

---

fenestra ovalis (see Plate, Fig. 3, No. 6) or oval window, which is closed by the base of the stapes. Above the oval window is a ridge formed by the course of the Fallopian canal, which transmits the facial nerve (see Plate, Fig. 2, No. 11). From this ridge there projects forward a hollow conical eminence, called the pyramid.

Behind the lower part of the promontory is a fossa, at the bottom of which is an aperture called the fenestra rotunda or round window (see Plate, Fig. 3, No. 18). This communicates with the cochlea, but is closed by the membrana tympani secundaria, which consists of a fibrous layer invested internally by the lining membrane of the cochlea and externally by that of the tympanum.



prevented from escaping by the secondary tympanic membrane at the round window. The internal auditory nerve gives off the cochlear branch, which is resolved into a multitude of filaments; these, with branches of the corresponding artery (auditory), enter the foramina of the spiral tract at the bottom of the internal auditory meatus. These ascend the canals of the axis of the cochlea and are reflected outwardly to the osseous spiral lamina, within which they form an intricate plexus, becoming associated near its free border with a series of nerve cells. From these emerge the nerve fibres to enter the membranous spiral lamina, where, with peculiar nerve cells, they form a complex arrangement.

The arteries supplying the cochlea are mainly derived from branches of the internal auditory artery. They terminate in a capillary network on the lining membrane and spiral lamina, from which veins originate and join the spiral venous sinus in the axis of the cochlea.




## CHAPTER II.

### PHYSIOLOGY OF HEARING.

By the sense of hearing, sounds produced by bodies in a state of motion are recognized. If these vibrations are regularly and rapidly propagated by the corresponding waves or undulations produced through the surrounding aerial medium, they produce a musical note. Liquids or solids transmit these vibrations more readily. The air is the ordinary medium, and no sound can be propagated in a perfect vacuum. Thus, by ringing a bell in a jar from which the air has been entirely exhausted, no sound is produced. The canal (Fig. 7, D) into which the external ear reflects the

Fig. 7.



sonorous vibrations passes inwardly until it is closed by the membrana tympani or "drum of the ear" (G).

This delicate membrane moves under the influence of



these vibrations, and communicates the motion first to the malleus, the long handle of which is inserted vertically between the laminæ. Its head is articulated with the body of the incus, the long process of the latter being connected with the stapes. The oval extremity of the stapes is attached to the membrane covering the fenestra ovalis or entrance to the labyrinth.

According to Helmholtz, by the transference of the vibrations of the tympanum to the much smaller membrane of the fenestra ovalis, thus throwing the somewhat movable liquid into vibration, mechanical power is gained. The apparently loose connections of the tympanic ossicles would seem to be unfavorable to the transmission of delicate vibrations. Helmholtz shows, however, that the firmness of the bony chain is much greater than has hitherto been supposed. Among other things, he points out that the articular surfaces of the malleus and incus are furnished, on their inferior aspects, with little processes or teeth (*Sperrzähnen*), which are so placed that, when the malleus moves outwardly, the articular surfaces easily separate as far as the somewhat loose capsular ligament will allow. When, however, the malleus moves inwardly, it grasps the incus with great firmness. The consequence of this is that when air is blown into the tympanic cavity, the membrana tympani can move outwardly without dragging the stapes out of the fenestra ovalis.

The functions of the tympanum are to render the vibrations of the membrana tympani quite free, and to isolate the chain of bones in such a manner as to prevent their vibration from being weakened by diffusion through the surrounding parts.

Opinions differ with regard to the function of the Eustachian tube. The one most commonly entertained is that it serves the same purpose as the hole in an ordinary drum, by removing an impediment to the free vibration of the membrane that would be offered by the complete inclosure of the air within. This is at once negatived by the fact, demonstrated by Mr. Toynbee and Dr. Jago,\* that the guttural

---

\* Functions of the Tympanum, by Jas. Jago, M.D., Brit. and For. Med.-Chir. Review, January and April, 1867.



orifice of the tube is usually closed, being only opened in the act of swallowing.

The principal object of the Eustachian tube, which is always found where there is a tympanum, appears to be to maintain an equilibrium between the external air and the air within. Dr. Jago believes that the ingress of aerial undulations from the throat, prevented by the normal condition of closure of the tube, would cause incessant oscillation of the membrana tympani, and that as the moment seized for bringing the tympanum into communication with the fauces must be one at which there can be no respiratory current, the period selected is the instant of swallowing, at which there is a compulsory suspension of respiration. Finally, the same arrangement secures the tympanum from the intrusion of gastric gases, etc. evolved through the fauces. This rule is not absolute, as they enter the tympanum strongly enough to palpably give a momentary protrusion to the membrane. The correlation between the Eustachian tube and the glottis should be thus expressed: the Eustachian tubes are shut whenever the glottis is open, but whatever act closes the latter opens the former.

Dr. Jago has a personal advantage in aural researches, owing to the fact that the guttural orifice of one of his Eustachian tubes can be opened and closed at will. The writer has a friend who has the same power, and who has repeated with him numerous experiments in confirmation of Dr. Jago's views. There are times when one of these tubes opens and remains so without any disposition to close. The writer has suffered from this condition, and the act of blowing the nose is attended with a sensation as though there was a perforation in the membrana tympani. This condition is always attendant on catarrh. When Dr. Jago has suffered from this patency of the tube, he has been able to close it for some moments, by an inspiratory movement with the nose and mouth closed, thus diminishing the aerial pressure in the throat. Again, he has found that a hearty meal would remedy this condition for several hours, if not for the remainder of the day. This, he thinks, is caused by the faucial tissue absorbing moisture directly from the fluids



swallowed. It is now fully fourteen years since the phenomenon of an open Eustachian tube has been understood, and as physician to two charities, and in general practice, he has met with numerous patients who were affected with aural symptoms, yet he has never met with an instance of a permanently patent tube.

The results of numerous experiments have persuaded Dr. Jago that both exhaustive and condensive stretching of the membrana tympani impedes the hearing of all tones, and more so when they are due to larger waves. In this he differs from many observers; and he also thinks that we never enjoy the privilege of hearing through the membrana tympani, or any thread of it, vibrating after the mode of musical strings—that is, in virtue of a given degree of tension. This explanation plainly coincides with the theory which makes the membrana tympani oscillate laterally, carrying with it the ossicular columella. From other experiments evidence was adduced that the membrana tympani will receive sonorous vibrations from a solid rod and (a fact incompatible with the fibre theory) a given tone at any point of its surface. He also concludes with confidence that aerial sonorous vibrations so interpenetrate the membrana tympani that it can perform its functions without lateral vibrations, even though it should not prove that it never performs them in that manner. If therefore the aerial sonorous vibrations make their way into the substance of the membrane, may not the precise constitution of the outer surface be of much acoustic moment?

The vibrations transmitted to the membrane covering the fenestra ovalis act upon the fluid contained within the labyrinth. This consists of the vestibule (Fig. 7, L), the semicircular canals (M), and the cochlea (N). Evidently the vestibule corresponds with the simple cavity constituting the entire organ of hearing in the lower animals. The vestibule is subdivided by a membranous partition into two cavities, in which are found small rounded masses of concretionary particles, collectively named otoconea and singly otoliths. In animals, in which the auditory apparatus is less perfect, they are seen to be quite large, seemingly needful to intensify




sounds. From investigations in comparative anatomy, Goodsir concludes that "the vestibule is that part of the organ by means of which any sound or series or combination of sounds is heard merely as noise." The simplest form of ear, which consists of a vestibule only, probably enables the sensorium merely to become cognizant of sound irrespective of the pitch or harmony of its constituent tones.

The semicircular canals (Fig. 7, M) have their functions variously estimated. It is commonly supposed that they receive the impression by which we distinguish the direction of sound; and it is certainly a powerful argument in support of this view that in almost every instance in which these parts exist at all they hold the same relative positions as in man, their three planes being nearly at right angles to one another. Prof. Samuel Jackson, of the University of Pennsylvania, believes that their function is to suppress sonorous vibrations, or to act as a damper, thus subserving to hearing the same purpose that the pigmentum nigrum does to vision. Flourens concludes, on making sections of the semicircular canals in pigeons, that the nerve supplying them does not minister to the sense of hearing, but to the directions of the movements of the animal; but this is fully explained on the supposition that the normal function of the semicircular canals is to indicate to the animal the *direction* of sounds, and that its movements are partly determined by these; so that a destruction of one or other of them will produce an irregularity of movement, resulting, as it would seem, from a sort of giddiness on the part of the animal, just as when one of the eyes of a bird is covered or destroyed.

M. Dugès surmised that by the cochlea we are especially enabled to estimate the pitch of sounds, and he adduces in support of this idea the fact that the development of the cochlea follows a very similar proportion to the compass of the voice. Dr. Thomas Young regards the cochlea as a micrometer or measurer of sound. Köl liker put forth a similar idea; and Helmholtz has confirmed these views by his researches. Microscopical investigation, with delicate dissections by Corti, Schultze, Köl liker, and others, have elucidated the structure of the cochlea.



"The canal of the cochlea, according to Köl liker, is filled with the fluid of the labyrinth, and contains not only two well-known cavities, the scala tympani and the scala vestibuli (*Sc V* and *Sc T*, Fig. 8), but also a third narrower space, which he proposes to call the scala media (*Sc M*). This is situ-



ated, in a great measure, within the lamina spiralis, and contains the ultimate terminations of the cochlear branches of the auditory nerve. The lamina spiralis consists of an inner bony and an outer membranous portion; the former terminates by a grooved edge, the upper margin of the groove (*zd*) being termed the zona denticulata, in consequence of its presenting a series of well-marked teeth; the lower margin (*hp*), the habenula perforata. From each of these a layer of periosteum is continued to the opposite wall of the cochlea, completing the septum between the scala tympani and scala vestibuli. The upper layer is termed the membrane of Corti (*mC*), the lower one the membrana basilaris (*mb*). Connected with the habenula perforata and lying in the scala media, or space between the two membranes, are certain structures termed the rods of Corti. These (1 and 2) form an angle with one another resembling the gable end of a house, and appear to be connected above by a delicate membrane, the membrana velamentosa. The spaces between the grooves of the bony lamina spiralis and the first set of the rods of Corti, and the space between the outer wall of the cochlea and the second set of the rods of Corti, are filled with large nucleated cells (*cC*), the cells of Claudius. The interval between the two sets of rods and the membrana basilaris (*Sc M*) is probably filled with fluid. The branches of the auditory nerve lie on the under surface of the lamina spiralis, beneath the habenula perforata, through which they pass by many small openings, and (though there is much difficulty in following them at this part) apparently terminate by free and extremely delicate extremities among the cells of Claudius and in the fluid of the scala media.




"The diagram on preceding page will give some idea of the relations of these delicate parts as they appear to be determined by the most recent investigations."\*

To fully illustrate this intricate and yet interesting subject, we have availed ourselves of the description and drawings in the work of Dr. Gruber.†


"If we examine into the minute structure of the membrana basilaris, we find as its foundation a structureless membrane, into which the upper plate of the labium tympanicum is transformed. This, particularly at the outer border, is found covered with a peculiar warty formation which gives the part different transverse diameters. These diameters vary from 0.012 to 0.025 mm. Upon this structureless membrane rests the formation designated as Corti's organ. Following next in order after the labium tympanicum of the sulcus spiralis are the so-called Corti's fibres (teeth of the second row, Corti; little rods, Claudius; arched filaments, Hensen; hearing rods, Henle,—Figs. 9 and 10).

Fig. 9.—CORTI'S FIBRES IN NATURAL POSITION.



a, ascending fibres; b, descending fibres. On their upper ends are seen the little rods, towards their lower ends and towards the concavity of the arch, the nucleus.

Fig. 10.—CORTI'S FIBRES LYING OVER ONE ANOTHER.



The descending fibres are plainly seen, with the little heads or projections placed upon their connecting parts.

"They appear to be the most important formations of the whole Corti's organ. They have their origin in the neighborhood of the foramina of the habenula perforata, and terminate

\* See M. M. Fick, "Anatomie und Physiologie der Binnens-Organen," 1862, p. 122; Carpenter, Principles of Human Physiology, sixth edition, p. 663, London, 1864.

† Lehrbuch der Ohrenheilkunde mit besonderer Rücksicht auf Anatomie und Physiologie, von Dr. Josef Gruber, Vienna, 1870, pp. 130-136.



again in the *membrana basilaris*. They lie side by side along the whole *membrana basilaris*, leaving between them spindle-shaped gaps, in such a manner that when viewed from above their arrangement exhibits a similitude to the keys of a piano. Corti's fibres which, in their chemical composition, show not the slightest resemblance to the *membrana basilaris*, dissolve readily in diluted caustic soda and potassa, and likewise in dilute hydrochloric acid; while alcohol, ether, chromic acid, saturated solutions of salt and sugar only cause them to shrivel (Kölliker); and they are then seen to consist of two parts, which are archlike and united, so that in their route along the whole *ductus cochlearis* they must form a slightly convex and also longitudinal protuberance. According to position, Corti's arches consist of an inner (*a*) and an outer (*b*) Corti's fibre, the former being (according to Claudius) more numerous; so that there are only two outer fibres for three inner ones.

"The inner fibres, which Deiters describes as less flexible than the outer ones, all begin regularly in a line outward from the foramina of the *habenula perforata*. At their commencement they are somewhat broader than they are farther along, and show, according to Kölliker, on one side of their origin a nucleus, which is attached to the fibre by a delicate membrane, which this author believes to be the principal cell constituent.


"The fibre runs, ascending outward and upward (ascending fibre), until it joins the broad end; hence the centre part of Corti's fibre is its narrowest part, and the upper end the broadest. The latter shows a cavity pointing outward and downward, in which fits an arched segment of the descending fibre to be shortly described. These two parts form in Corti's arches the so-called articulations (*coins articulaires internes et externes*, Corti).

"The expanded and inward convex piece of the descending fibre just mentioned lies the highest, yet it does not expose its upper end; while its expanded part above is covered by the outer flat ends of the ascending fibre. In immediate continuation runs still another fibre, so that the upper end of the descending fibre appears turned or lapped over in an arch-



like manner. In their farther course downward, the descending fibres diminish very strikingly, to be finally reinserted with a broader bell-shaped end on the membrana basilaris. This end (according to Kölliker), towards the concavity, and likewise on the starting-point of the ascending fibre, contains a nucleus, which is not always to be found on the single fibres

Fig. 11.—THE ENTIRE MEMBRANA BASILARIS, WITH ALL ITS PARTS LYING EXPOSED FROM ABOVE TO SHOW THE PROPORTION OF NERVES.—CORTI'S ORGAN FROM THE SIDE. (After Deiters.)



*a*, the teeth; *b*, the large cells which fill the sulcus, partly passed forward; *c*, holes or foramina of the habenula perforata; *dq*, inner Corti fibres; *e*, outer Corti fibres; *f*, connecting rods detached; *g*, the first bundles of the transverse nerve fibres; *h*, the second from above respiratory outward: in these are seen the so-called phalanges; *i*, the third; *k*, the fourth; *l*, arising longitudinal nerve fibres, which submerge in the first bundle of the transverse ones; *m*, cylindrical cells on the lower arch of the pars membranacea; *n*, the cells of Claudius on the connecting rods; *o*, the supporting fibrous system of Corti's organ; *p*, a longitudinal nerve fibre on the base of Corti's arch; *q*, cells of Corti; *y*, fusiform cells.



when detached from their insertion. The diameter of the little rods varies from 0.001 to 0.002 mm.\*

"The second formation, which constitutes Corti's organ (see Fig. 11), is the lamina reticularis cochleæ discovered by Kölliker (lamina velamentosa, Deiters). Fig. 8, 1, 2.

"It is named after its discoverer, and is one of the most difficult parts of Corti's organ to delineate, especially if the object be to define its proportions exactly; and concerning no part of the cochlea is there such a variety of opinions as of this. Kölliker describes it, first, as a short, clear plate with fine bordered divisions, corresponding in numbers with the inner Corti's fibres. This plate presents on the border the inner and outer Corti's fibres, and is in close connection with the former, and consists in a union of appendages of the inner fibres, which Kölliker calls plates. After these plates follows a netlike lamella, consisting of longer straight rods (*f*) enlarged at their extremities, whose numbers correspond with those of the outer Corti's fibres, and which are connected with the jointed ends of the same in a manner not satisfactorily solved. These rods, which lie close under the clear plate, show, near the edge of the latter, a slight, sometimes nodular, protuberance. Upon these rods follow smaller ones, situated between the outer ends of the former pieces, in the shape of a biscuit, which Kölliker calls 'inner intermediate parts' (Deiters's phalanges of the first row), (*h*); then, the outer intermediate parts, in shape similar to the former, between the ends of the inner parts (Deiters's phalanges of the second row); and lastly, a row of parts which sometimes appear as four small rectangular portions lying close together, in whose outer ends hair-shaped appendages appear.

---

\* Henle divides the ascending fibres into two species: the first being flattened in a direction perpendicular to the membrana basilaris. The transverse section is elliptical, almost double the length, through the long diameter. The body of the other is cylindrical, only broad toward the ends. He can distinguish these two forms, but believes that the difference is only determined by the position of the objects. The bodies of the descending fibres are always cylindrical.



"The intermediate parts (phalanges) leave between them three rows of foramina, at regular distances, which Kölliker calls inner, middle, and outer holes (rings, Böttcher; circles, Deiters) of the lamina reticularis.

"According to Henle, the lamina reticularis appears partly in the form of a network of fine hyaline threads, with regular, partly four-sided, partly round, meshes, which alternate in three rows, in such a manner that, as a general rule, these little frames, which divide the foramina of the first row from those of the third one, touch upon the middle of the foramina of the second row.

"The outer edge of the third row of meshes lies just over the place of insertion of the outer Corti's fibres, on the membrana basilaris. In other cases (both forms appearing in the same species of animal), the threads of the net are finer, the meshes closer, partly or entirely closed by delicate membrane, with the exception of the three constant rows of round openings.

"Deiters found, sometimes, the round meshes divided by a small archlike cross-piece. As before mentioned, the enlarged rods, which are connected with the joint ends of the descending Corti's fibres, have the form of a ship's rudder, and their somewhat enlarged ends are inserted in depressions of the outer edge of the joint ends of the outer Corti's fibres. These would effect the closing of the innermost meshes of the lamina reticularis.

"While the origin of the lamina reticularis, from the joint ends, is given with certainty, as well as its parallel course with the membrana basilaris, no certain knowledge exists about its insertion. Henle says, under the supposition that it is attached somewhere to the wall of the cochlea, one might assign to it the function of a band, which fastens the little rods in their position. In chemical construction the lamina reticularis resembles Corti's fibres. Besides the structures already mentioned, other cell formations belong to Corti's organ, which converging form and arrangement may partly be designated as epithelium cells; others, however, are not known in their origin, and are named either after their discoverer, or after their peculiar form.



“Here belong the hairy cells (stalky cells of Corti; thorn cells of Leidy; staff cells of Hensen; inner and outer covering cells of Henle). The nomenclature of Henle is derived from their relation to Corti's fibres. He distinguishes the cells here mentioned, according as they are above or under Corti's fibres, as cover and floor cells; if both, according to their position upon the membrana basilaris, *outer* or *inner*. About the hair cells, which constitute the most delicate formation of the whole Corti's organ, in their relation to their neighboring formation, we are instructed in many particulars by Deiters, Kölliker, Hensen, and lately also by Henle. The inner hairy cells are found upon the joint ends of the inner Corti's fibres; they pass backward on the epithelium of the labium tympanum, by which they are partly covered.

“Their form, as Henle remarks, is conical (*m*), whereby the broader end is directed more outward and upward, and carries a bundle of stiff cilia, which may be noticed distinctly upon single cells which evidently lie toward the edge. The inner hair cells, which, as Kölliker remarks, have their seats in three rows alternately on the holes of the membrana basilaris, and likewise have a more conical configuration, run over the outer Corti's fibres downward, and carry on the upper ends just such cilia as the inner hairy cells. The three rows of the outer hairy cells cover one another partly until shaped, and lastly change at the ends to fine thin threads, which are inserted in three rows placed alternately on the membrana basilaris (Deiters).

“On the upper ends adhere the hairy cells which cover the foramina of the lamina reticularis, close to the borders of the foramina of the same, in such a manner that, by their attachment, the upper part of the cell frequently remains closed. These hairy cells, which were known to Corti, although superficially, go by the name of Corti's cells in contradistinction to Corti's fibres.

“Deiters has discovered still other hairy cells, independent of the inner cells, which run in three rows between and outward from those described. Deiters's hairy cells adhere with their inner ends to the outer ends of the so-called phalanges, with their outer ends toward the membrana



basilaris. They are spindle-shaped, and their ends diverge in long, fine threads. The outer end, before its insertion on the membrana basilaris, unites with the outer fine end of one of Corti's cells in such a manner that their insertion is a common one,—the thin end of Deiters's cells uniting the rods. In addition, other cells on the membrana basilaris have more the property of epithelium cells, particularly when treated with water, as they puff up and assume a ball-like form; one sees these especially well under Corti's fibres (Deiters's support tissue; Stützgewebe). Henle's floor cells, on the membrana basilaris, beyond the insertion of the uniting stems, are smaller and larger as such cells are present.

“The outer zone (*zona pectinata*) is that outer portion of the membrana basilaris which unites the inner zone with the wall of the cochlea. It shows on its upper plane, toward the ductus cochlearis, a ribbed, comblike appearance, from whence it derived its name, *membrana pectinata* (Todd and Bowman). These ribs, which, especially on the outer section, do not spread out entirely straight, but are somewhat waving, and which were formerly taken to be an expansion of a fold of the structureless membrane (which, as Henle has lately shown, originate from fibres which have a diameter of 0.0025 mm.), are not always single-radiating, but are wavelike, and bend, especially on the outer ends. To the outer side, toward the wall of the cochlea, slight depressions present themselves on the *zona pectinata*; and here, according to Kölliker, it receives a fibre-mass coming from the lamina spiralis, ossea accessoria, which this author described as *ligamentum spirale*, while Todd and Bowman looked upon this structure as muscular, and named it *musculus cochlearis*.

“According to Henle, the *zona pectinata* is strengthened by several uniting tissue-strings extending on its tympanic surface, running parallel with the longitudinal axis of the lamina spiralis.

“The vestibular wall of the ductus cochlearis, the membrana Reissneri, shows, besides its covering of epithelium, a thin layer of simple connective tissue, and numerous capil-



laries, which stretch from the periosteum of the lamina spiralis close upon the periosteum of the cochlea. From the insertion of Reissner's membrane to the insertion of the membrana basilaris, the periosteum of the cochlea forms the outer wall of the ductus cochlearis. This shows, besides the epithelium, a thin layer of tissue, abundantly supplied with vessels (stria vascularis, Corti), and outward from this a tissue-plate, which consists, according to Kölliker, of cells which have a great resemblance to cartilage cells.

"Besides this outer wall of the ductus cochlearis, we find, on observing further, immediately over the insertion of the membrana basilaris, a protuberance, extending far toward the canal, which, Hensen states, contains a vessel (vas prominens, Hensen), of the presence of which Henle, however, could not convince himself."

The number of the rods of Corti is estimated by Kölliker at three thousand, or about three hundred to each semitone of the octave.

Besides the structures just enumerated, there are within the labyrinth certain fine elastic bristles terminating in sharp points, between the internal nerve fibres. These bristles, discovered by Max Schultze, are eminently calculated to sympathize with those vibrations of the fluid of the labyrinth which correspond to their proper periods. Thrown into vibration, the bristles stir the nerve fibres which lie between their roots. At another point in the labyrinth we have the otoliths already noticed as imbedded among the nervous filaments, which, when they vibrate, exert an intermittent pressure upon the adjacent nerve fibres, thus exciting audition.

Finally, according to Helmholtz, this wonderful organ in the labyrinth, discovered by Marchese Corti, which we have described, is to all appearances a musical instrument, with its cords so stretched as to receive vibrations of different periods and transmit them to the nerve filaments which traverse the organ. "Within the ear of man, without his knowledge or contrivance, this lute of three thousand strings has existed for ages, receiving the music of the outer world and rendering it fit for reception by the brain. Each musical tremor which falls upon this organ selects from its tensioned



fibres the one appropriate to its own pitch, and throws that fibre into unisonant vibration. Thus, no matter how complicated the motion of the external air may be, these microscopic strings can analyze it and reveal its constituents.”\*

“This idea has been confirmed by the recent microscopic researches of M. Victor Hensen† on the hearing of crustaceous fishes. Having placed the decapoda in salt water, charged with strychnia to increase the reflex power of the cerebral nerves, the animal fell into convulsions at the least noise. He has proved afterward that in them hearing takes place through the medium of hearing-bristles, and that each little fibre vibrates in unison with a determined note.”

---

\* Tyndall.

† Helmholtz, *Die Lehre den Tonempfindung*; *L'Acoustique*, par Rodolphe Radau, p. 291. Paris, 1870.



## CHAPTER III.

### ACOUSTICS.—LAWS OF SOUND.

“ACOUSTICS (derived from the Greek verb ἀκούω, “to hear”) treats of the science of sounds, their cause, nature, and phenomena. Sound is the impression (or motion) produced on the sense of hearing by the vibration of sonorous bodies.”

If it be a series of irregular vibrations, it is called a noise; but if it be a succession of rhythmic impulses, a musical tone.

The ear is capable of appreciating the three fundamental qualities of sound, viz., intensity, or amplitude of vibration; pitch, or rapidity of vibration; and timbre, or quality.

Higher tones mingle with the fundamental tone, determining the pitch of a sound, or its value in the musical scale; and it is this intermixture which determines what, for want of a better term, we call the *quality* of the sound.

It is this union of high and low tones that enables us to distinguish the sound of one musical instrument from another.

Most sounds have a series of impulses partaking in a greater or less degree of the character of a musical tone.

Any continuous sound is dependent upon a succession of impulses, and the rapidity with which these succeed one another determines the pitch of the tone. We may ascertain by experiment what number of such impulses are required to give a musical tone of any pitch.

Thus if the instrument called the siren, composed, in its simple form, of a circular plate with a number of apertures at regular intervals, be made to revolve over the top of a pipe through which air is propelled, a succession of short puffs will be allowed to issue from this; and if the revolution be sufficiently rapid, these impulses will unite into a definite tone.\*

These vibrations are transmitted from the external ear by

---

\* Tyndall on Sound.



a special nerve, the auditory, and are recognized by the brain as sound.

Sounds are propagated through the air in waves, which may be transmitted simultaneously in different directions without destroying each other.

This is illustrated in complicated pieces of music played by an orchestra; a practiced ear can readily distinguish the sound of each instrument.

It has been estimated that the number of complete vibrations required to produce an appreciable musical sound, in persons having an acute sense of hearing, may vary from eight for the lowest to thirty-six thousand five hundred for the highest, per second.

Helmholtz has recently fixed the lower limit at sixteen vibrations and the higher at thirty-eight thousand vibrations per second.

The practical range of musical sounds is comprised between forty vibrations and four thousand vibrations per second, which amounts in round numbers to seven octaves.

Weakening of sound, which takes place according to the law of inverse squares, would be in great measure prevented if the sound-wave were so confined as to impede its lateral diffusion.

By transmitting it through a tube with a smooth interior surface, we accomplish this, and the wave thus confined may be transmitted great distances with very little diminution of intensity. This condition is fulfilled and utilized in the speaking-tubes in our dwellings.

"The celebrated French philosopher Biot observed the transmission of sound through the empty water-pipes of Paris, and found that he could hold a conversation in a low voice through an iron tube 3120 feet in length. The lowest possible whisper, indeed, could be heard at this distance, while the firing of a pistol into one end of the tube extinguished a lighted candle at the other.

"The action of sound thus illustrated is exactly the same as that of light and radiant heat. They, like sound, are wave motions. Like sound, they diffuse themselves in space, diminishing in intensity according to the same law. Like



sound also, light and radiant heat, when sent through a tube with a reflecting interior surface, may be conveyed to great distances with comparatively little loss. In fact, every experiment on the reflection of light has its analogue in the reflection of sound.

"The limits of hearing are different in different persons. Dr. Wollaston, who gave the first proof of this, while endeavoring to estimate the pitch of certain sharp sounds, remarked in a friend a total insensibility to the sound of a small organ-pipe, which, in respect to acuteness, was far within the ordinary limits of hearing. The sense of hearing of this person terminated at a note four octaves above the middle E of the piano-forte.

"The squeals of the bat, the sound of a cricket, even the chirp of the common house-sparrow, are unheard by some people who for lower sounds possess a sensitive ear. The ascent of a single note is sometimes sufficient to produce the change from sound to silence.

"The suddenness of the transition," writes Wollaston, "from perfect hearing to total want of perception, occasions a degree of surprise which renders an experiment of this kind with a series of small pipes among several persons rather amusing. It is curious to observe the change of feeling manifested by various individuals of the party, in succession, as the sounds approach and pass the limits of their hearing. Those who enjoy a temporary triumph are often compelled, in their turn, to acknowledge to how short a distance their little superiority extends."\*

Nothing can be more surprising, writes Sir John Herschel in reference to this subject, than to see two persons, neither of them deaf, the one complaining of the penetrating shrillness of a sound, while the other maintains there is no sound at all. Thus, while one person mentioned by Dr. Wollaston could but just hear a note four octaves above the middle E of the piano-forte, others have a distinct perception of sounds full two octaves higher. The chirp of the swallow is about the former limit, the cry of the bat about

---

\* Tyndall on Sound, p. 73.



an octave above it, and that of some insects probably another octave higher.

In "The Glaciers of the Alps," Dr. Tyndall has referred to a case of short auditory range, in crossing the Wengern Alp in company with a friend. The grass at each side of the path swarmed with insects, which to him rent the air with their shrill chirping. His friend heard nothing of this, the insect music lying quite beyond his limit of audition.

Once, while crossing the Stelvio Pass by night in company with a friend, he complained of acute pain in the ears. Dr. Tyndall desired him to swallow his saliva: he did so, and the pain instantly disappeared. By the act of swallowing, the Eustachian tube is opened, and thus equilibrium is established between the external and internal pressure.

Dr. Wollaston was expert in closing the Eustachian tube and leaving the space behind the tympanic membrane occupied by either compressed or rarefied air. He was thus able to cause his deafness to continue for any required time without any effort on his part, always, however, abolishing it by the act of swallowing. A sudden concussion may produce deafness by forcing air either into or out of the drum of the ear. In the summer of 1858, Mr. Tyndall was on the Fee Alp, in Switzerland, where, jumping from a cliff on to what he supposed to be a snow-drift, he came into rude collision with a rock which the snow barely covered. The sound of the wind, the rush of the glacier torrents, and all the other noises which a sunny day awakes upon the mountains, instantly ceased. He could hardly hear the sound of his guide's voice. This deafness continued for half an hour; at the end of which time a suitable act opened the Eustachian tube, and restored, with the quickness of magic, the innumerable murmurs which filled the air around him.

"Light, like sound, is excited by pulses or waves; and lights of different colors, like sounds of different pitch, are excited by different rates of vibration. But in its width of perception the ear exceedingly transcends the eye; for while the former ranges over eleven octaves, but little more than a single octave is possible to the latter."\*

---

\* Tyndall, *op. cit.*, p. 75.



A knowledge of the transmission of sound through gases, liquids, metals, and the various kinds of wood, is of practical importance to the physician and physiologist. The subjoined tables are from the work of Tyndall:

## VELOCITY OF SOUND IN GASES. FROM THE EXPERIMENTS OF DULONG.

	(Temperature 0°C.)	Velocity.
Air . . . . .		1092 feet.
Oxygen . . . . .		1040 "
Hydrogen . . . . .		4164 "
Carbonic acid . . . . .		858 "
Carbonic oxide . . . . .		1107 "
Protoxide of nitrogen . . . . .		859 "
Olefant gas . . . . .		1030 "

## TRANSMISSION OF SOUND THROUGH LIQUIDS. (WERTHEIM.)

Name of Liquid.	Temperature.	Velocity.
River-water (Seine) . . . . .	15° C.	4714 feet.
" " . . . . .	30	5013 "
" " . . . . .	60	5657 "
Sea-water (artificial) . . . . .	20	4768 "
Solution of common salt . . . . .	18	5132 "
Solution of sulphate of soda . . . . .	20	5194 "
Solution of carbonate of soda . . . . .	22	5230 "
Solution of nitrate of soda . . . . .	21	5477 "
Solution of chloride of calcium . . . . .	23	6493 "
Common alcohol . . . . .	20	4218 "
Absolute alcohol . . . . .	23	3804 "
Spirits of turpentine . . . . .	24	3976 "
Sulphuric ether . . . . .	0	3801 "

## VELOCITY OF SOUND THROUGH METALS. (WERTHEIM.)

Name of Metal.	At 20° C.	At 100° C.	At 200° C.
Lead . . . . .	4030	3951	.....
Gold . . . . .	5717	5640	5691
Silver . . . . .	8553	8658	8127
Copper . . . . .	11,666	10,802	9690
Platinum . . . . .	8815	8437	8079
Iron . . . . .	16,822	17,386	15,483
Iron wire (ordinary) . . . . .	16,130	16,728	.....
Cast-steel . . . . .	16,357	16,153	15,709
Steel wire (English) . . . . .	15,470	17,201	16,394
Steel wire . . . . .	16,023	16,443	.....



## VELOCITY OF SOUND IN WOOD. (MM. WERTHEIM AND CHEVANDIER.)

Name of Wood.	Along Fibre.	Across Rings.	Along Rings.
Acacia . . . . .	15,467	4840	4436
Fir . . . . .	15,218	4382	2572
Beech . . . . .	10,965	6028	4643
Oak . . . . .	12,622	5036	4229
Pine . . . . .	10,900	4611	2605
Elm . . . . .	13,516	4665	3324
Sycamore . . . . .	14,639	4916	3728
Ash . . . . .	15,314	4567	4142
Alder . . . . .	15,306	4491	3423
Aspen . . . . .	16,677	5297	2987
Maple . . . . .	13,472	5047	3401
Poplar . . . . .	14,050	4600	3444

The tuning-fork is a familiar instrument with which we may produce an invariable note, its principal use being to regulate the tone of musical instruments.

It is composed of a U-shaped bar of steel, connected at its base with an axis or handle. This, when struck against anything solid or when a bow is drawn across either prong, gives a definite note by its state of intense vibration.

By some modern writers upon diseases of the ear, special importance is attached to this instrument as an aid in diagnosis.

## VIBRATION OF AIR CONTAINED IN TUBES.

In wind instruments with walls of suitable thickness, the column of air contained in the tubes alone enters into vibration. The material of the tube has no influence on the pitch, but affects the quality in a striking and important manner, and the condition of the interior surface, whether rough or polished, or lined with cloth, has no effect upon the intensity of the sound. The shape of the tube has an important influence: if it be of the same size throughout, the sound will be simply condensed and strengthened by the reflections; if it be tapering, the angles will become more and more acute and the note deeper, the sound being increased; if it be conical, the apex receiving the sound and the base giving it exit, as in the speaking-trumpet, the angles will be less and less acute, until the last may be even obtuse.



The exact dimensions of each section of such instruments, therefore, becomes a matter of importance.

Tubes of wood, metal, glass, etc. are not merely conductors of sound, by preventing the rays from diverging and increasing their power by reflection, but the molecules comprising the instrument themselves vibrate.

These vibrations are in harmony with the sound introduced, but varying in their perfection.

It is evident, therefore, that if tubes prevent sounds from being lost in the air by expansion, they must conduct it to an almost indefinite distance; hence the great utility of hearing- and speaking-trumpets.


#### INSTRUMENTS TO ASSIST HEARING.

Ear-trumpets are to the ears what spectacles are to the eyes, but the aid which they render is neither as perfect nor as complete. Yet with all their defects they could not be dispensed with, as they are the only means by which deaf persons have intercourse with their friends, or are enabled to listen to a lecture or sermon; each individual generally has his or her favorite instrument, and would not part with it on any consideration.

Ear-trumpets are bent so as to cause the sound to be deflected from a straight line in passing into the ear; and by making the portion which receives the sound like a funnel, a much larger volume is received for the impression on the impaired auditory nerve.

The simplest form is seen at Fig. 12.

Fig. 12.—SIMPLE EAR-TRUMPET.



Those who are very deaf require a tube from fifteen to eighteen inches in length; those moderately deaf, one not so long.

In treating many deaf persons we must have a variety of




these useful instruments, so as to allow the patient to choose the one most suitable.

Mr. Isaac Williams, of this city,\* manufactures trumpets of every variety to suit the various idiosyncrasies. He has a deaf workman, whom he at times employs to test the various trumpets.

Those sizes that are most used are from three and seven-eighths ( $3\frac{7}{8}$ ) inches in diameter at the larger extremity, the curved tube in all being from twelve to fifteen (12 to 15) inches in length, terminating in a small acorn-like extremity, which can be made of tin, ivory, or silver, to fit the external ear.


The tin japanned trumpet is seen at Fig. 13.

Fig. 13.—EAR-TRUMPET (tin japanned).



The larger trumpet seen at Fig. 14 is one which has been employed by the Rev. Dr. Haswell and his son, the devoted missionaries to Burmah, who are both very deaf.

Fig. 14.—HASWELL'S EAR-TRUMPET (to be had of Messrs. Kolbe and Gemrig, Instrument-makers).



They have employed this form after trying every variety of English and American hearing-trumpets. They prefer it because it allows several persons to converse with them


\* They are to be had also of Kolbe and Gemrig, instrument-makers.



at one time, whilst the large perforated part is turned towards them, and can catch every word distinctly; they have even heard the faint whisper of a dying man. Hence this form, apart from its size, is adapted to clergymen, and many very deaf persons have found it most admirable, enabling them to hear a sermon, lecture, or conversation.

The next one, Fig. 15, is the original trumpet of Harriet

Fig. 15.—MARTINEAU HEARING-HORN (ivory tips, ten inches).



Martineau, which is also a useful instrument, and much employed in England.


Either of the last-mentioned forms can be used in a church or lecture-room. They should be placed upon the floor or a stool, and the sound conveyed from them by being connected with a gutta-percha tube, which can be concealed in the clothing. The trumpet itself can be covered on its outer part with cloth or silk.

#### CONVERSATION-TUBES.

Sometimes the cores of the tubes are made of iron or copper wire, having a funnel-shaped arrangement at one end, and a small tip at the other. These are called "Conversation-Tubes;" the deaf person inserts the tip in his ear, whilst the funnel-shaped extremity is talked into.

The conversation-tube illustrated in Fig. 16 is made of silk or cotton on a wire body covered with gutta-percha, with horn tips.

Fig. 16.—EAR CONVERSATION-TUBE, MADE OF SILK OR COTTON ON A WIRE BODY WITH HORN TIPS.






In speaking into all these instruments, it is of the utmost importance not to elevate the voice, but to speak distinctly.

The auricles invented by the celebrated Itard (Fig. 17) fit into the ears under a cap or bonnet, and can be covered with the hair; they are kept in place by means of a spring passing over the head. They have been found by some to be useful in listening to a general conversation or lecture; but

Fig. 17.



very few can wear them long, on account of the rustling noise and the irritation they create in the external ear.

The following is the experience of one of my patients in regard to the use of auricles or ear-cornets:

“Having used ear-cornets or auricles for six or eight years, I can safely recommend their use.

“These instruments are invaluable in shopping, in the transaction of business, and in social intercourse; as, by wearing them, those persons who are very hard of hearing will only be regarded as moderately so.

“The greatest drawback to their usefulness is that they cannot be worn any length of time, as they press upon and weary the ear, causing at times quite a painful sensation. About three consecutive hours is as long as they can prudently be worn. If a lady be the wearer, she finds it unpleasant at times to have her bonnet tied over them.”


Any form of ear-trumpet should not be indiscriminately worn for use in the early stage of deafness, more especially



those cases arising from debility or congestion of the nervous apparatus. In chronic cases, however, they often prove most serviceable. They are equally advantageous in cases of partial ankylosis of the stapes to the fenestra ovalis. The surgeon should determine, with the assistance of his patient, the proper form to be used, and not trust, as is too often the case, to the instrument-maker alone. He should take the same care in fitting his patient with an ear-trumpet as the ophthalmic surgeon does in fitting his patient with the proper form of eye-glass.

A smaller form, or pocket-trumpet, is seen at Fig. 18, which is capable of being taken apart and so arranged that it may be carried in the pocket.

Fig. 18.—POCKET EAR-TRUMPET.



Many persons employ this kind abroad, as well as a large one at home; it has been found very useful for physicians. That form of ear-trumpet which Von Tröltzsch has found to be the most useful for the greatest number of cases consists of a stranded leather tube, of a foot or more in length, with horn extremities. The end inserted in the ear should be about the size of the meatus, the patient holding it in or on this part. Similar to this leather ear-trumpet is one of pasteboard, which, for the sake of convenience, is made in sections, to be joined together when used. This form may be seen at No. 3, Fig. 19.

The otaphone consists of a silver clamp fitted to the posterior border of the auricle, which has for its object to cause the ear to stand out farther from the head, and thus make easier the reception of sounds.

To distinguish very low sounds, Helmholtz fitted a glass funnel into the ear with softened wax, and was very successful in this way.

Patients living at a distance, who cannot come to the surgeon, by sending a description of the form of deafness, with a wax model of the meatus, can be furnished with an ear-trumpet accurately fitted.

The following illustrations, taken from a recent French




work,\* show, in a comprehensive view, the various forms used in France. The one mostly employed is seen at Fig. 19, No. 1. Another form much employed is that at (2).

Curtis has had constructed trumpets which draw out like a telescope (3).

Itard has indicated a series of other forms. The first is a sort of ellipsoid, furnished with a pavilion and curved tube, which sinks into the ear; the pointed ends are two

Fig. 19.



membranes of goldbeater's skin, which, without augmenting the sound, render it less confused (4).

Another system is represented by (5). To realize it, Itard has taken a spindle-shaped shell, and to it added a little tube to fit the ear; also a pavilion, into which one could introduce, if wished, one or two membranes of goldbeater's skin.

Recently, Mr. Koenig has constructed an acoustic trumpet, which is at the same time a stethoscope (6). It consists of a capsule shut in by a communicating membrane, and is connected with the ear by means of a gum-elastic tube terminating in an ivory end.

When you speak before the membrane, the latter vibrates and pushes the column of air which is in the tube against the tympanum of the ear.

The length of the tubes is four metres, and shorter when the sound seems feeble.

\* *L'Acoustique*, par R. Radau. Paris, 1870.



## CHAPTER IV.

### TABLES OF CLASSIFICATION OF DISEASES OF EAR.

DISEASES of the human ear causing deafness are more frequent in the male than the female; this may be accounted for by the fact that the male is more exposed to the vicissitudes of the atmosphere. According to Sir William Wilde, of two hundred persons affected with deafness from disease of the ear, one hundred and one were males, and ninety nine females. In France, Triquet states that, of the same number of cases, one hundred and twenty-six were males and seventy-four females. In the record kept by the author, of six hundred cases, there were three hundred and twenty-six males and two hundred and seventy-four females.

In Dr. Roosa's table it will be seen that in five hundred cases, including ten deaf mutes, the proportion was only two hundred and ninety-seven males to two hundred and three females. This discrepancy, in this country, arises from the present style of head-dress worn by the ladies in the street, which exposes the ear to every variety of temperature.

Cold, producing catarrh in its various forms and its consequences, is the predisposing cause of deafness in childhood, and throughout all the stages of life.

Diseases of the ear are often the result of constitutional disorders, as, for instance, the gouty, rheumatic, scrofulous, tuberculous, syphilitic, and cancerous. Age is also a frequent cause of deafness; and so good an authority as Von Tröltsch asserts that not more than one out of three persons from twenty to forty years of age possesses good and normal hearing. In our own investigations, we have found that the larger number of persons lose the hearing of one ear and sometimes both, and this we have found at all ages. About one-fifth of the cases of disease of the ear are hereditary or communicated from progenitors, many families having two, three, four, five, and even six members deaf and dumb;



this is owing to too close consanguinity of parents. Then follow, next in order as a cause, the exanthematous disorders, scarlet fever, measles, and smallpox. The other causes are mumps, quinsy, with acute and chronic inflammation of the throat, nasal passages, and middle ear. There is a form of deafness caused by intermittent, typhoid, and typhus fevers, in which the brain is the part usually affected, which, for want of a better name, we call nervous deafness.

A glance at the tables, compiled from various sources and our best authorities on aural surgery, will also serve to give a comparative estimate of the frequency of aural disease and the many forms it assumes, as well as the results of treatment, etc.

## A TABLE

*Of Six Hundred Cases of Diseases of the Ear, from Dr. Turnbull, occurring during 1868, 1869, and 1870, Howard Hospital and Private Practice.*

	Male.	Female.	Whole No.	Cured.	Improved.	Unimproved.	Died.	Not treated.
Injured auricle . . . . .	1	3	4	4	...	...	...	...
Foreign body in external canal . .	2	3	5	5	...	...	...	...
Eczema auriculæ, acute and chronic .	6	14	20	12	6	2	...	...
Inspissated cerumen . . . . .	65	48	113	73	35	5	...	...
Furunculous abscess meat. ext. . .	18	12	30	30	...	...	...	...
Otitis externa . . . . .	12	18	30	16	12	2	...	...
Otitis interna, pur. perf. (chronic) .	2	3	5	...	2	1	2	...
Catarrh, aur. media, acute, perf. . .	68	31	99	37	25	36	1	...
“ “ “ chronic “ . . . . .	63	48	111	12	51	39	...	9
Otitis media, inflam. mastoid process .	18	17	35	10	13	8	4	...
“ “ “ chronic, with caries . . .	6	4	10	...	2	6	2	...
“ “ “ syphilitica . . . . .	9	9	18	2	14	1	1	...
“ “ “ with polypi . . . . .	9	9	18	14	3	1	...	...
Periostitis, with abscess (cerebral). .	2	4	6	...	1	2	3	...
Neuralgia, plex. tymp. . . . .	2	3	5	2	2	1	...	...
Effusion of blood, memb. tympani . .	2	1	3	3	...	...	...	...
“ “ “ middle ear . . . . .	1	1	2	1	...	1	...	...
“ “ “ serum “ “ . . . . .	3	1	4	2	2	...	...	...
Cophosis nervosa, caused by concussion, typhoid, scarlet, and bilious fevers, etc.	5	16	21	...	2	19	...	...
Perforation membrana tympani by blow	3	1	4	3	1	...	...	...
Inflammation membrana tympani, acute	2	1	3	1	2	...	...	...
“ “ “ “ “ chronic . . . . .	18	19	37	11	13	13	...	...
“ “ “ of Eustachian tube . . . .	7	3	10	3	7	...	...	...
Deaf mutes . . . . .	4	3	7	...	1	6	...	...
	326	274	600	241	194	143	13	9



It is only by estimating the causes and morbid conditions developed by unhealthy action that one can form just estimates of the probabilities of success in treatment. To aid in the formation of opinions, with reference to the results of treatment, the effect of location, etc., these figures have been obtained.

A view of the tables will show a great preponderance of chronic affections. Chronic catarrh and chronic inflammation of the membrana tympani are frequent causes of deafness, and to the student of these diseases the question naturally arises, how many of these cases of "dread silence," in this world of sweet harmonies, are the result of neglecting acute attacks that were manageable in their early stage, but, when seen by the aural surgeon, were, alas, too late for remedy?

## A TABLE

*Of Five Hundred Cases of Diseases of the Ear, from Professor Roosa.*

	Whole No.	Cured.	Much imp.	Improved.	Unimproved.	Unknown.	Died.
Furuncular inflammation of external canal	8	7	...	...	...	1	...
Diffuse inflammation of external canal	11	11	...	...	...	...	...
Eczema of external canal	5	4	...	...	...	1	...
Inspissated cerumen	46	42	...	2	1	1	...
Foreign body in external canal	2	2	...	...	...	...	...
Simple acute inflammation of cavity of tympanum. (One side)	21	18	...	1	...	1	1
Simple acute inflammation of cavity of tympanum. (Both sides)	13	11	...	2	...	...	...
Simple chronic inflammation of cavity of the tympanum	245	11	17	56	96	64	1
Acute suppurative inflammation cavity of tympanum. (One side)	8	6	...	...	...	2	...
Acute suppurative inflammation cavity of tympanum. (Both sides)	7	1	5	...	...	1	...
Chronic suppurative inflammation cavity of tympanum. (One side)	45	5	10	19	...	9	2
Chronic suppurative inflammation cavity of tympanum. (Both sides)	66	2	15	27	9	11	2
Diseases of labyrinth	9	...	...	...	...	9	...
Deaf mutes	10	...	...	...	...	11	...
Tinnitus aurium without deafness	3	...	...	...	...	3	...
Hysteria	1	...	...	...	...	1	...
	500	120	47	107	125	95	6



"The time required for treatment is a question that is worthy of consideration. Perseverance is the rule in almost all of the chronic forms of disease. And many that have apparently not been improved, by weeks of careful treatment, will suddenly commence to progress favorably, and eventually be much improved. To have abandoned them would have given a fine opportunity for some potent mystery to have gained the reputation of the cure."

Dr. Roosa gives, in the following table, the duration of five hundred deafness cases before they came under his observation. The figures are valuable, as they are obtained by a distinguished and accurate observer.\*

One year or less . . . . .	177 or about 35 per cent.
Between one and two years . . . . .	35 " " 6.1 "
Three and four years . . . . .	25 " " 5 "
Four and five " . . . . .	19 " " 3 $\frac{1}{2}$ "
Five and ten " . . . . .	107 " " 21 "
Ten and twenty " . . . . .	82 " " 16 "
Twenty and thirty years . . . . .	36 " " 7 "
Thirty and forty " . . . . .	12 " " 2 $\frac{2}{3}$ "
Forty and fifty " . . . . .	4 " " 0.8 "
Fifty " . . . . .	2 " " 0.4 "
Seventy-five " . . . . .	1 " " 0.2 "

The results of treatment may be estimated by a glance at the following table, compiled from various sources:

AUTHOR.	Whole number.	Cured.	Improved and much improved.	Not improved.	Not treated.	Died.
Schwartz . . . . .	322	145	122	31	111	3
Roosa (1870) . . . . .	500	120	{ 47 107	125	95	
Moos (chronic aural catarrh) . . . . .	100	17	64	19	.....	...
Clarke . . . . .	140	62	31	14	33	...
Turnbull . . . . .	600	241	194	143	9	13
Roosa (1871) . . . . .	494	97	{ 65 84	110	Unknown. 137	1

\* New York Med. Journal, Aug. 1869, page 513.



The following table,\* from the Report of Diseases of the Ear, treated in General Hospital in Vienna, by Dr. Josef Gruber, during the year 1868, is of interest, for the sake of comparison. The completeness of this table will make it valuable.

DISEASES.	Remaining at the end of 1867.		Admitted.		Cured.		Improved.		Not cured.		Without treat- ment.		Died		Remaining at the end of 1868.	
	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F
EXTERNAL EAR.																
Freezing of external ear . . . . .	...	...	1	...	1	...	...	...	...	...	...	...	...	...	...	...
Othæmatoma . . . . .	...	...	...	2	...	1	...	1	...	...	...	...	...	...	...	...
Eczema . . . . .	1	1	1	5	1	5	...	...	...	...	...	...	...	...	1	1
Inflammation of mastoid process . . . . .	...	...	1	...	1	...	...	...	...	...	...	...	...	...	...	...
Atresia external auditory canal . . . . .	...	...	2	1	...	1	...	...	...	...	2	...	...	...	...	...
Foreign bodies in internal auditory canal . . . . .	...	...	1	8	1	8	...	...	...	...	...	...	...	...	...	...
Accumulation of cerumen . . . . .	1	3	68	40	64	40	2	1	...	...	...	...	...	...	3	2
Circumscribed inflammation of ex- ternal auditory canal . . . . .	4	2	26	12	23	10	4	...	...	...	...	...	...	...	3	4
Acute inflammation of external ear . . . . .	9	2	21	10	20	9	3	...	...	...	...	...	...	...	7	3
Chronic " " " " . . . . .	2	...	7	4	6	1	...	...	...	...	...	...	...	...	3	3
Acute inflammation of membrana tympani . . . . .	3	1	16	4	13	2	1	...	...	...	...	...	...	...	5	3
Chronic inflammation of membrana tympani . . . . .	2	2	5	8	4	6	...	2	1	...	2	...	...	...	...	2
Thickening of the membrana tym- pani . . . . .	2	2	5	...	...	...	2	1	1	1	2	...	...	...	2	...
Loss of substance of membrana tym- pani . . . . .	...	...	2	...	...	...	...	...	1	...	1	...	...	...	...	...
Rupture of membrana tympani . . . . .	...	...	2	...	1	...	1	...	...	...	...	...	...	...	...	...
Cicatrices of membrana tympani . . . . .	...	...	1	...	...	...	...	...	...	...	1	...	...	...	...	...
Abnormal deformities of the mem- brana tympani . . . . .	...	1	6	2	1	1	3	1	2	...	...	...	...	...	...	1
MIDDLE EAR.																
Acute catarrh . . . . .	9	6	35	21	30	20	3	...	...	...	1	...	...	...	11	6
Chronic " " . . . . .	23	22	90	52	25	8	40	32	5	3	13	5	...	...	30	26
" " with polypus . . . . .	6	2	12	6	9	1	4	3	...	...	...	...	...	...	5	4
Contraction of Eustachian tube . . . . .	10	5	4	2	...	...	4	2	...	...	3	2	...	...	7	3
INTERNAL EAR.																
Acute inflammation of internal ear . . . . .	1	2	5	3	2	...	3	2	...	1	...	...	...	...	1	2
Caries and necrosis of temporal bone . . . . .	1	1	...	2	...	...	...	1	...	1	...	...	1	...	...	1
Affections of labyrinth . . . . .	17	16	37	34	2	1	5	7	6	6	18	19	...	...	23	17
Nervous earache . . . . .	...	...	4	3	4	3	...	...	...	...	...	...	...	...	...	...
" " tinnitus aurium . . . . .	1	1	...	1	...	...	...	...	1	...	...	...	...	...	...	...
Deaf mutism . . . . .	3	2	16	3	...	1	...	...	1	1	15	3	...	...	3	2
	95	71	368	228	208	118	75	53	18	13	57	30	1	...	104	80

\* Monatsschrift für Ohrenheilkunde, No. 9, September, 1869.



*Three Hundred Cases of Diseases of the Ear observed in England, compared with the same number seen in Germany by Kramer.\**

EXTERNAL EAR.				
			ENGLAND.	GERMANY.
Catarrh of the ceruminous glands . . . .			9	46
Inflammation of the cartilage . . . .			4	8
Inflammation of the cellular tissue . . . .			1	1
Inflammation of the periosteum . . . .			0	0
Acute inflammation of the membrana tympani .			1	2
Chronic inflammation of the membrana tympani .			28	55
MIDDLE EAR.				
Catarrhal inflammation without exudation . . . .			0	2
Catarrhal inflammation with free exudation . . . .			7	10
Catarrhal inflammation with free and submucous exudation . . . . .			9	29
Catarrhal inflammation with submucous exudation . . . . .			239	143
Nervous deafness . . . . .			2	4
			<hr/> 300	<hr/> 300

#### METHOD OF INVESTIGATING DISEASES OF THE EAR.

In order to fully comprehend the significance of symptoms and pathological conditions, some method in the order of investigation is necessary.

The following plan† is subjoined as a guide in investigating diseases of the ear. It is based, as far as possible, upon principles that have been established by practical observation and experience.

I. Name, age, and occupation of patient. Residence.

II. State of health, temperament, condition of pulse, etc.

III. If any relations are deaf.

IV. History of the affection; duration; discharge; supposed cause. Former symptoms; if at any time pain in ears or head. Nature of progress, whether rapid or slow; if it has increased by sudden paroxysms, or by imperceptible degrees. Present symptoms; of pain, noises, or discharge; amount of hearing power determined by watch, or as shown by conversation; whether the patient requires to be spoken to distinctly in a room, or within the distance of a yard, or close to the ear; which ear is the worse.

\* Review Kramer and Politzer; Aural Surgery, Dr. Roosa; Boston Medical and Surgical Journal.

† Modified from Toynbee.



*Causes* aggravating deafness: a cold, cold or moist weather, bodily fatigue, or mental excitement; the act of mastication.

*Causes* producing improvement: improved health, dry air, cold weather.

#### V. Result of examination.

Right ear; left ear. The distance at which the watch is heard (the watch brought toward the ear). Ordinary distance in health is from three feet to thirty. Tuning-fork; conduction of sound through the bones. Patient's own voice; mode of speech.

Objective examination; auditory.

*Meatus*: quantity and condition of cerumen; state of the dermis of the osseous wall.

*Membrana tympani*: surface dull or shining, transparent or opaque; state of triangular bright spot; if more or less concave than natural.

Handle of malleus; catheterization and air-bath on hearing.

Eustachian tube: if air is heard, by means of the otoscope, to enter the tympanic cavity naturally during deglutition, the nose and mouth being closed; if the air is heard to enter the tympanic cavity naturally during a forcible attempt at expiration with nose and mouth closed.

Left ear ditto.

State of mucous membrane of the pharynx; condition of tonsils. General condition; hereditary tendency. Rhinoscopy. Examination of posterior nares and Eustachian tube.

#### VI. Previous treatment.

#### VII. Diagnosis.

#### VIII. Treatment.

### DISEASES OF THE EXTERNAL EAR.

I. If ever troubled with erysipelas, and at what age in life?

II. If meatus ever closed, and the ear ever changed its natural form?

III. If ever affected with eczema?

IV. If a thin discharge exuded?

V. Is the external ear sensitive on pressure?

VI. If cysts were ever formed, and the ear disfigured by them?



VII. If there have been any tumors, and their cause? If carcinomatous, and their different stages?

#### EXTERNAL MEATUS.

- I. If the membrana tympani be visible?
- II. The diameter of the external meatus?
- III. If the lamina of the membranes of the meatus are healthy?
- IV. If the entrance of the meatus be protected by hairs?
- V. Shape of the orifice of the meatus?
- VI. If there be the natural projection of the tragus?
- VII. If the meatus can be properly explored?
- VIII. If any foreign body in the meatus? and its character by careful inspection? Symptoms of the patient at the time; whether cough or vomiting occurred, etc.; and whether there is giddiness or congestion in the head?
- IX. Character of membrana tympani? (Its brilliancy.)
- X. If ceruminous glands be normal?
- XI. The color and nature of the substance discharged?
- XII. The cause of its collection?
- XIII. The symptoms when collected,—deafness, pain, crackling, or singing sound heard? if any inflammation by its presence?
- XIV. If it caused any ulceration in the membrana tympani? whether there is inflammation of the dermis? acute or chronic?
- XV. Result of examination: Character of discharge from the ear, and quantity? Causes of said inflammation? and if extending toward the brain? Whether there be simple chronic inflammation, or chronic inflammation of the dermis lining the meatus? The exciting causes? The state of health?

#### POLYPOID GROWTHS.

- I. Their cause? Their position and attachment?
- II. Whether there be a discharge? matter or blood?
- III. Size of polypus by inspection?
- IV. Whether any cerebral irritation?
- V. Character and name of polypus?



## OSSEOUS TUMORS.

- I. Where situated?
- II. Whether caused by rheumatic or gouty diathesis?
- III. How large and how far in the ear they extend?
- IV. Their appearance?
- V. Whether they are painful on pressure or devoid of sensibility?
- VI. Whether there is inflammation?
- VII. If there be a discharge, and if offensive?
- VIII. Whether there is deafness from their position?
- IX. If the ear is distended and a sense of weight is experienced on the affected side of the head?
- X. Whether the patient be of temperate habits?

## MOLLUSCOUS TUMORS.

- I. What part of the ear are the tumors located? Size and extent?
- II. Whether they arise from the dermis? Do they enlarge rapidly?

## SEBACEOUS TUMORS.

- I. Where situated? Size?

## MEMBRANA TYMPANI.

- I. If in a healthy condition? Color, lustre, cone of light.
- II. Character of its three layers? Inclination, curvature.
  - (a) Opacities of epidermoid; (b) dermoid; (c) substantia propria; (d) tendinous or peripheral opacities; (e) mucous membrane; (f) which tissue affected?
- III. If any calcareous degeneration of lamina?
- IV. Whether there is perforation? Size and form? If any artificial tympani would be of service?
- V. If rupture of the membrana tympani? The cause of rupture? Situation, size and form.

## EUSTACHIAN TUBE.

- I. The anatomical character?
- II. Whether obstructed in any way?
  - (a) If at its faucial orifice?
  - (b) If at its tympanic orifice?
  - (c) If in the middle part of the tube?
- III. Causes of obstruction?



IV. If the Eustachian catheter has been used? and if of service?

CAVITY OF TYMPANUM.

- I. Its anatomical character?
- II. If mucous membrane congested?
- III. If inflammation of mucous membrane?
  - (a) Its causes?
  - (b) If chronic inflammation or hypertrophy?
  - (c) If catarrhal inflammation? and
  - (d) Its extent?
- IV. If ulceration of mucous membrane of tympanum?
- V. If rigidity of tympanic mucous membrane?
- VI. If ankylosis of the bones?
- VII. If dislocation of bones?
- VIII. If articulation is perfect?

THE MASTOID CELLS.

- I. Anatomical character?
- II. At what age diseased?
- III. If necrosis of the mastoid processes?

NERVOUS DEAFNESS.

- I. If nervous apparatus diseased?
- II. Causes of the disease?
  - (a) By concussion?
  - (b) By application of cold?
  - (c) Various poisons?
  - (d) Typhus, typhoid, scarlet, or rheumatic fevers, measles, mumps, etc.?
  - (e) Mental excitement?
  - (f) Physical debility?
- III. If any ulceration, caries, or necrosis?

MALIGNANT DISEASES OF THE EAR.

- I. What part of the ear it occurs?
- II. Its causes?
- III. Constitution of patients affected with it?
- IV. Hereditary influences?

DEAF AND DUMB.

- I. Causes, as near as possible?
- II. Condition of ear?



## CHAPTER V.

ON THE MEANS OF DIAGNOSIS BY THE USE OF ARTIFICIAL AND DAYLIGHT FOR EXAMINING THE MEMBRANA TYMPANI AND AUDITORY CANAL — PREVENTION OF DEAFNESS — NOISES IN THE EAR, ETC.


A LARGE number of lamps have been employed as means of illuminating the ear, a few of which we will describe rather as relics of the past than for any good results to be obtained from them; strange as it may seem many persons prefer them to the more simple form which we now employ, as making more impression upon the patient. In 1741, Archibald Cleland, an English army surgeon, published in the "Philosophical Transactions" an account of "instruments proposed to remedy some kinds of deafness proceeding from obstructions in the external and internal auditory passages." The first of these consisted "of a convex glass three inches in diameter, fixed in a handle, into which is lodged a wax candle, which when lighted will dart the collected rays of light into the bottom of the ear." According to Wilde,\* "to this apparatus Bozzini added a concave mirror, but without much effect." Deleau further modified it by placing a lighted taper between two concave mirrors; then followed the lamp of Buchanan, which was improved by Kramer, who substituted an Argand lamp for the feeble wax-light. Later came a number of portable ear lamps, one of which the author employed for a number of years, consisting of a box attached to which was an oil lamp and two metallic reflectors, one of which is set at an angle in the bend of a projecting arm, and throws a stream of light on the membrana tympani through a fixed tubular speculum inserted into the meatus, while the operator looks through a narrow tube containing two small lenses. Another form is that of Clark's, which has been modified by the late Mr. Toynbee by causing the funnel por-

---

Wilde on Diseases of the Ear, pp. 27 and 69.




Fig. 20.—TOYNBEE'S ILLUMINATING SPECULUM.



tion to have a wider flange, and by adding to it a series of tubes which fit into the meatus ; this form is seen at Fig. 20.

Fig. 21.—HINTON'S MODIFICATION OF 'TOYNBEE'S ILLUMINATING SPECULUM SO THAT PHYSICIAN AND STUDENT CAN BOTH SEE AT THE SAME TIME.





Mr. Hinton, the successor of the late Mr. Toynbee, has again modified this speculum by fixing the candle to the funnel with a universal joint, and having a double tube and lens, adding three sizes of tubes for the ear. This instrument was brought for me by Dr. Gerhard during a recent visit to London, and it is claimed for it that when properly adjusted both physician and student, or friend, can see at the same time. It is shown at Fig. 21.

Dr. Weber has arranged a very neat little apparatus on the principle of the ophthalmoscopic mirror by receiving the light from the sun, or a flame on the mirror, which has a hole in the centre through which the operator looks. The light is reflected through a double lens, which, by concentrating the rays, makes the object more distinct, especially if it be

Fig. 22.—WEBER'S REFLECTOR.




minute. The small-sized speculum is supported by a ring attached to the instrument.

Another arrangement is where the mirror is attached to



an elastic band, which holds it in position on the forehead of the operator. The light is placed above the head of the patient, and a small lens can be held between the light and the reflector. The objection to this is that the least motion of the head changes the cone of light, and this interferes with the focus. With artificial light we are able to detect

Fig. 23.



a foreign body, an ulcer, granulation, or in some instances a perforation. But this is not as good as daylight obtained concentrated and reflected by the hand-mirror of Tröltzsch, for with this alone can we detect slight shades of vascu-

larity produced by inflammation or congestion, delicate opacities, minute points of morbid deposit, want of polish or loss of transparency, having fixed in the meatus the tubular conical speculum of Dr. Gruber of Vienna (shown at Fig. 23).

Fig. 24. — KRAMER'S BIVALVE SPECULUM.




These were brought into almost universal use by Sir W. Wilde, who first described the speculum in 1844 as "a small conical tube of silver measuring about an inch and a half in length, five-eighths of an inch in width at the greater aperture, and varying from two to four lines in the clear at the small extremity." The late Mr. Toynbee changed the conical form to the oval, considering this to be the form of the outer meatus. The objection to this form is that in moving it about you give more pain than with the conical. It is well, however, to have a series of sizes and a variety of forms; and in some cases it may be necessary to use the valvular speculum of

Dr. Kramer, which is seen at Fig. 24.

Dr. Politzer has employed hard-rubber specula, being less heavy, and the dark ground of the inner surface favoring a clear definition of the illuminated parts. The conical form will be found most useful to the general practitioner when



Fig. 25.—FULL-SIZE CONCAVE MIRROR OF VON TRÖLTSCHE.





only one set is desired; and should we add to this the concave mirror of Von Tröltzsch (Fig. 25), before alluded to, we can examine the membrana tympani with great satisfaction even on a cloudy or stormy day, and in the evening we can employ a candle, lamp, or gas. The focal distance of this concave glass mirror is about six inches. It is seen at Fig. 25, full size. "The recent advance in our knowledge of aural diseases, which makes it a satisfactory field of study and practice, is due to the adoption of this method."\* Dr. Politzer states: "The most important results in this department (that of the finer pathological changes in the membrana tympani) have been furnished by the thorough investigations of Von Tröltzsch." He compared the observations made on patients by means of a very considerably improved method of examination with corresponding appearances on the cadaver, and showed from the unison between supposed and proven causes of disease how successful were observations of the membrana tympani for the determination of the nature of pathological changes in the auditory apparatus. Now it is well to remember the natural appearance of this membrane in health. "A normal membrana tympani, examined by reflected daylight, is a translucent pearl-gray membrane" (Politzer), with the head and handle of the malleus distinct—not very prominent, however, nor projecting much above the plane of the membrane. On the lower section is a reflection of light of a general triangular shape, its apex resting on the lower extremity of the handle of the malleus. The angle formed by the membrane with the upper wall of the external auditory canal is one of about  $140^{\circ}$ . (Tröltzsch.)† This has been verified by numerous recent examinations, and it agrees with our own results. Of its deviations from this normal standard we have treated in another portion of this work.

---

\* Dr. Roosa, Transactions of New York State Medical Society, 1869, p. 226.


† The Appearance of the Membrana Tympani, Drs. Roosa and Berard, American Journal of Medical Sciences, April, 1867, p. 399.



## ON THE MEANS OF DIAGNOSIS BY AUSCULTATION OF THE EAR.

The devoted and laborious investigations of Laennec were not only applied to the chest, but also extended to the ear. In his treatise he devotes an entire section to the auscultation of this organ; yet his stethoscope had many objections, being applied as it was over such an irregular surface as the external ear. This objection was overcome, however, by means of the flexible stethoscope devised by our lamented friend, the late Dr. Pennock, which, by the addition of two ivory tips, the late Mr. Toynbee made applicable to the diagnosis of diseases of the ear, and it is now called the otoscope (see Fig. 26), or diagnostic tube. It may consist

Fig. 26.—OTOSCOPE.



of a simple elastic tube, or, better, one of coiled metallic wire covered with silk, gutta-percha, or worsted web, having ivory or bone tips, as in Fig. 26. Its use may be exemplified by placing one tip in the subject's ear, when in perfect hearing order, and the other in the physician's ear, when the patient is desired to close the nose and mouth and make a deep inspiration (or inward breathing), when the sound is heard of the *click* of the opening of the Eustachian tube.


If, however, the ear is diseased, and we introduce the diagnostic tube, the patient can, by closing the nose and mouth and swallowing, cause the air to move, but not always to enter the Eustachian tube. If the tube be very moist it creates a vesicular sound; if more normal, the sharpness of the tone is changed; if very dry, we have more of a knocking sound, showing an obstruction; if the sides of the tube be relaxed, we have what may be described as a flap-



ping sound, or the sound may be momentarily obliterated by the alternate collapse and dilatation of the tube; and in some instances no sound whatever is heard, owing to entire obliteration.

The Valsalvian experiment is performed with the little instrument seen in Fig. 27, which is placed in the external

Fig. 27.



meatus. It will sometimes assist us in determining whether the Eustachian tube is open or closed. This instrument, also called the aural manometer, was suggested by Politzer, and consists of a horseshoe shaped glass tube  $1\frac{1}{2}$  mm. in calibre, which when used is partially filled with a solution of carmine in ether. It is firmly retained

in the auditory meatus by means of a nozzle, which must be anointed. By the rising and falling of the liquid the variations in the pressure of the air in the middle ear are indicated. Care, however, must be taken that the motion of the jaw is not mistaken for the movement of the air, as the author has produced this result in his own person whilst the nose and mouth were closed.

If we have a very thin *membrana tympani*, there are the same sounds, with an increased pressure felt upon the *membrana tympani* of the listener; or if it is perforated, there is near us a whistling or loud hissing sound (the *souffle*). If the Eustachian catheter be now introduced, and the physician blows through it, the diagnostic tube still being in position, air may be forced into the middle ear by gradually and cautiously increasing the pressure. The position is indicated in Fig. 86: *a*, patient; *b*, surgeon; *ed*, the rubber diagnostic tube; *e*, the Eustachian catheter.

This pressure may be increased by using the caoutchouc bag of Politzer (Fig. 87) or the air-press, either large or small, which latter is seen in Fig. 32.

If a powerful stream of air pass through the Eustachian tube of normal size, and having a normal degree of moisture,




it creates a sound which "Deleau likens to the falling of rain upon the leaves of a tree, and therefore terms 'bruit de pluie;' while Von Tröltzsch calls it a vesicular murmur, or, if the force of the air be very great, a knocking sound, 'anschlage geräusch;' because we hear the stream of air striking a dry, elastic membrane, the drum, and pressing it somewhat outward."

Dr. E. Leudet\* states that the involuntary objective bruits of the ear are rare. They may be muscular; at times they consist of a kind of crepitation and of crackling, and are identical with those which some persons can produce at will. The involuntary crackling bruit in the ear may coincide with a tic in the face, and seem to be caused by a rhythmical convulsion of the internal muscle of the malleus and chain of ossicles.

#### THE DOUBLE OTOSCOPE.

Having read in some of the German works of a double otoscope, but never having seen one, I have availed myself of a modification of the late Dr. Camman's double flexible stethoscope, which sometimes renders sounds distinct when they are scarcely appreciable with the ordinary instrument. It consists, as in Fig. 28, of two pair of tubes uniting so as to form one direct tube from the patient's to the surgeon's ears. The first pair are made of German silver, gently curved, and supplied at the aural extremities with ivory knobs; these fit securely into the surgeon's ears, and the tubes are firmly held in position by a band of india-rubber stretched across them. As they fit so accurately into the operator's ears, and are held

Fig. 28.



\* Gazette Médicale de Paris, No. 35; Half-Yearly Abstract of Medical Sciences, January, 1870.



there so securely, his hands are free. The other pair fit into the German-silver tubes, thus making them continuous. They are composed of rubber covered with silk, and likewise ending in ivory knobs, which fit accurately, and require to be held in the patient's ears. It is also supplied with an extra single tube, which may be used in one ear of the patient when the two are not wanted.

"With this apparatus the sounds are thus received through both organs of hearing and the faintest sound can be heard. With it we can determine the condition of the labyrinth, as patients are able to hear a tuning-fork placed on the median line of the vertex of the head better in the affected side than in the other.\* Politzer says, in a letter to Von Tröltzsch, that in cases where the patient is not able to state correctly in which ear he hears the tuning-fork most distinctly, he places both ends of a single otoscope (or diagnostic tube) in the auditory canals of the patient, and then the sensation is more decided.

"Lucae used for the same purpose a double otoscope made like the double stethoscope of Scott Allison.

"While the tuning-fork, which has been placed in vibration, is on the vertex of the head, or on the teeth of the patient, we may, by alternately pressing the two arms passing to the patient's ear, determine from which ear the tone is coming, and also distinguish the difference in its magnitude and clearness. Care should be taken that the tuning-fork is vibrating fully, and that the two ends of the three-armed tube are evenly placed in the meatus. The power of perceiving sounds through the bones is much less in advanced age than earlier in life. We can, therefore, draw no direct conclusion from a want of hearing the watch or tuning-fork after a patient is over fifty years of age.

"Lucae has recently described a new method of examining the organ of hearing, for physiological and diagnostic purposes, with the aid of the "interference otoscope." By this name he designates the double otoscope described above, but altered in such a manner that an india-rubber tube is

---

\* Von Tröltzsch, Diseases of the Ear. American edition, p. 261.



inserted in it, which leads to a small metal sound-receiver of the shape of a half paraboloid. The latter is brought in front of a tuning-fork placed on an isolated stand, and this is struck by means of a small wooden hammer. By alternately pressing together one and the other tube of the double otoscope, we may decide how much the sonorous waves are reflected from each ear. This reflection of sounds increases in all changes in the sound-conducting apparatus, which either directly or indirectly cause an increased tension of the membrana tympani."

THE PREVENTION OF DEAFNESS—COMMON HABITS INJURIOUS TO THE EAR AND CAUSING DEAFNESS—NOISES IN THE EAR, ETC.

First, children ought never to be struck (or boxed) on the ears, as injury to the membrana tympani, and even rupture, may follow; or deafness be produced by the sudden jar or shock, dulling the sensibility of the nerve. Teachers should always try a child's hearing before blaming it for inattention. This should be done by placing it a few yards distant and trying whether it can understand what is said to it in a rather low tone. Each ear should thus be tried by a watch while the other is stopped by the finger. The chief cause of deafness is cold. This is to be prevented by avoiding the ordinary causes, and by, first, taking plenty of fresh air; secondly, using cold water to wash the ears, neck, and front of the chest, and then rubbing the parts perfectly dry. Patients should avoid draughts, or sitting in wet clothes, or becoming overheated when in a close room and then passing suddenly into a cooler atmosphere: as from the church, school, concert, theatre, etc. Strong draughts are particularly hurtful to the ear, such as are caused by sitting close to the window of a carriage, car, or building. We should never let rain, sleet, or snow drive into the ear, as they also are very dangerous. Another source of danger is from keeping the interior of the ear too hot, by plugging the meatus with wool or cotton, or covering with hood, fur caps, etc., except in cases of great exposure, as when encountering a driving storm. "We\* have known a case of deafness result from the

---

\* Graphic.



placing of a piece of cotton in each ear before diving; the cotton disappeared and the patient was deaf for five years, and on examination by a physician he found both pieces of cotton firmly pressed against the membrana tympani, and covered with wax. In other instances there were found three separate pieces of cotton pushed down into the passage, where they remained, forming an obstruction to hearing." Pieces of paper are also rolled up and pushed into the ear. Almost all varieties of substances are placed there by children, as stones, grains of indian-corn, wheat, shells, etc. Their removal can always be accomplished by means of syringing with warm water, or the use of the curette. When such accidents happen, no attempt should be made with forceps or probes of any kind, with the exception of a strand of delicate silver wire, which can be employed by the physician, as if he were about to remove a cork from a bottle. It causes much less suffering, however, to the child, to employ the syringe, and also too much haste is thereby avoided. The reflux of the water should be depended on to wash the foreign body out; and this may require syringing for several days. Washing the ear out with soap and water is injurious, keeping the wax moist when it ought to become dry and scaly, increasing its quantity unduly, and collecting the particles of dust always floating in the atmosphere. Introducing the twisted up corner of a towel is particularly hurtful, as it pushes the wax down upon the membrane. Washing should not extend beyond the outer surface, and care should be taken that all the little folds of the outer ear be carefully dried with a soft towel. Another hurtful agent is a piece of sponge attached to a handle (the aurilave) to wash out the ear. This is saturated with water, and becomes an air-tight piston. If there is itching of the ear, a little glycerine introduced on the end of the finger will afford relief, and painting the parts with spirits of wine has been found useful. Earpicks, tweezers, etc. are to be condemned, as are also bodkins, pins, etc., as these may accidentally pierce the delicate membrane. Sportsmen sometimes have this membrane pierced by a twig, on turning suddenly, while getting through a hedge, and farmers often suffer from the



introduction of the beard or straw of wheat or rye. Loud sounds, as the firing of guns and the ringing of large bells, etc., may produce deafness, as often occurs during a cannonade, and in some instances the explosion causes rupture of the membrane.

If an insect should get into the ear, pour in oil or glycerine; these suffocate it, when it should be washed out with a syringe. Boys sometimes injure the ears of their playfellows in snowballing; and bathing, although a very healthful exercise, is often injurious to the hearing, first, by staying too long in the water; second, swimming and floating are dangerous on account of the entrance of cold water into the ear or nose. No cold fluid ought to be allowed to enter the ear, and if water get in, the head should be carefully turned to one side and efforts made at once for its removal. It is, therefore, always safe, in bathing in the sea, to cover the head with some light material, as an oil-silk cap or straw hat tied over the ears. "That the risk to hearing from unwise bathing is not a mere fancy, we have proof in numerous cases of deafness from no other cause in man, and it is also proved by the fact, well known to lovers of dogs, that those animals, if in the habit of jumping or being thrown into water, so that their heads are covered, frequently become deaf. Wet hair will cause deafness, and it should not be left hanging about the ears of children, as for the purpose of curling."\*

We now come to that cause of deafness which results from the desire to keep the ear clean. Nature has her own method of cleansing the ear by means of the ear-wax. In health this wax dries and forms thin, fine scales; these separate from the surface.

It is well, when subjected to violent concussions of air, to take the necessary precautions. Do not stand to the windward of a cannon or large gun, and always keep the mouth open. Furthermore, constant noises, as those made in a factory or foundry, ought, if possible, to be avoided.

It is a bad practice, when in pain, to introduce chloroform,

---

\* Graphic, op. cit.



hartshorn, hot oil, or grease of any kind; pieces of hot fig or onion should never be put in, but warm flannels should be applied with fomentations of hot water or poppy. If the pain does not soon subside, cotton charged with either of the volatile agents may be employed, being always surrounded by a dry piece of cotton that they may not cause much irritation.

NOISES IN THE EAR, OR TINNITUS AURIUM, NERVOUS AND ACOUSTIC.

The ringing sound heard in the ears is not only a symptom of almost all diseases of the ear, but also a very troublesome indication of disease in itself, for after all apparent disorder has passed away, the patient is still distressed and disturbed by a most persistent form of noise, which, in some instances, is a form of mental malady. Tröltsch describes these sounds as a form of irritation of the auditory nerve, known by various names, and produced by quinine, blood diseases (anæmia and chlorosis), temporary and permanent hindrances to the circulation, functional diseases of the nervous system, and abnormal conditions of the ear itself, as we have before stated. More especially in all the conditions that increase the pressure upon the fluid of the labyrinth; but chronic catarrh is the most constant cause of tinnitus, either from mucus in a liquid or dried condition.

It is stated by Politzer, that spasmodic contraction of the tensor tympani is a frequent cause, and Tröltsch\* speaks of a shortening of the tendon of the same muscle as a cause. In diseases of the semicircular canals, as described by Ménière, it is stated to be always accompanied by severe tinnitus. Concussions produce tinnitus; hyperæmic conditions of the naso-pharyngeal space cause this affection, also foreign bodies in the Eustachian tube. The author had under treatment a case caused by swallowing the beard of barley, which passed into the tube, causing otitis media purulenta, and the beard was ultimately discharged through the perforation. Dr. Cohen saw this case with him, and made a rhinoscopic

---

\* Treatise on the Diseases of the Ear, p. 518.



examination, but was not able to remove the offending body in its early stage. The patient recovered under treatment, the membrana tympani healing perfectly and the hearing being normal.

Another case is related by Fleischmann as follows: "A man complained for years of a continuous sound in the ear, and of a very peculiar sensation in the pharynx, as though a hair had gotten into his mouth. On the post-mortem section, a grain of barley was found projecting from the pharyngeal orifice of the tube and reaching into the osseous."\* Cold in the head is sometimes accompanied by temporary tinnitus, which after a time becomes permanent from repeated attacks, and this is always considered an unfavorable symptom. In cases where the venous sinus surrounding the carotid artery becomes ossified the pulsation of it is perceived in the ear, and the same occurs in the narrowing or aneurism of the artery.

Rayer† reports a case of pulsating tinnitus synchronous with the beat of the heart, which was perceived by others than the patient on auscultation, and which was momentarily checked by compression of the mastoid branch of the posterior auricular artery. Many persons are able to produce a crackling sound, which is one of the internal sounds of the ear, and, according to Politzer, results from the sudden drawing away of the membranous portion of the Eustachian tube, which occurs during the contraction of the tensor palati muscle. There is also a similar sound made by certain individuals in the act of swallowing, on the movement of the jaw, and there are also sounds which originate in the ear itself and adjacent structures. An inquiry into their sources and conditions is called by Dr. Jago Entacoustics.‡ One of the points of importance he insists upon

---

\* Von Tröltzsch, op. cit.

† Comptes Rendus des Séances et Mémoires de la Société de Biologie, 1854, p. 163.

‡ Entacoustics, James Jago, M.D., British and For. Med.-Chir. Review, No. 83, July, 1868, p. 225; Functions of the Tympanum, by the same author, British and For. Med.-Chir. Review, April, 1867; On Diseases of the Ear, p. 324.



is, that "a tinnitus aurium frequently finds its way to the labyrinth through the membrana tympani, and that the tympanic wall is liable to the adherence of a tenacious secretion, and it is the removal of this that relieves aural catarrh, and not the simple opening of the Eustachian tube." Another point is, that if the hearing remain good notwithstanding the tinnitus, he believes the nerve to be healthy, and that the membrane and ossicular chain are intact, and that the vessels at fault are situated in the innermost regions. In these cases he believes that the tinnitus rather subsides gradually than suddenly, though a pellicle of mucus may drop from the inner wall of the drum, causing sudden cure. Dr. Jago lays great stress on tinnitus depending upon the numerous arteries, the partial narrowing of the calibre of any one of which may produce a noise. He also makes this remark: If a *souffle* be produced in the temporal artery by a due pressure with the finger in front of the tragus, it becomes much louder if the meatus be stopped; and it may be taken as a safe rule that when deafness depends upon a cause as deeply seated as the drum-head, a *souffle* or vascular tinnitus from vessels lying external to this is not likely to be heard, and where no deafness exists such a *souffle* or tinnitus would be augmented in loudness on stopping the meatus; whilst a sound, issuing from a point more deeply seated than the bottom of the meatus, is not likely in any case to be rendered *appreciably* louder in this way.

At the Howard hospital, in the treatment of all the nervous cases we employ those agents which improve and build up the nervous system. Locally we use the oils of belladonna and hyoseyamus, by dropping in the ears. The truly acoustic cases require the air-douche charged with chloroform, ether, iodine, or carbolic acid. As we have stated, many of these sounds which distress the patient are from a diseased condition of the *naso-pharyngeal mucous membrane*, it being the seat of catarrhal inflammation. This we have been in the habit of treating by the use of a solution of common salt and an abundance of tepid water, by the means suggested first by Professor Weber, and made still more practical by Dr. Thudichum, of London. It consists simply of a portion of



rubber tubing, attached to which is a weight of lead or glass (see Fig. 56), placed at the bottom of a pitcher containing a quantity of the fluid above described; and this is set on a shelf or held higher than the patient's head. The other end of the tube is armed with a nozzle which fits into the nostril of the patient. The patient then bends forward and the fluid passes up one nostril, across, and emerges at the other. There are some important cautions to be observed: first, the fluid must be of the temperature of the body (above  $96^{\circ}$ ); second, the patient must breathe gently with the mouth open; and lastly, must not swallow, else the fluid will pass into the middle ear and cause the following results, well told by a patient in the following letter from

"FREDERICA, KENT CO., DELAWARE.

"MY DEAR SIR,—I find on using the 'nasal douche' as recommended by you, that it affects me somewhat unpleasantly. I find no difficulty in passing the water as directed from one nostril to the other, or back into the throat. On passing the water into the throat the Eustachian tubes apparently are also filled, and give the same sensations I have experienced, when a boy, in swimming, and which we used to call 'bubbles in the ear.' I cannot free my head of the water taken in for some four or five hours after using the douche. I then feel as if I had taken cold. My ears feel sore; pressing the tips of the fingers into the external ear causes a dull pain, apparently about the drum of the ear. This passes off in about twelve hours. . . . I am much more deaf than usual for some hours after using the douche. . . .

"Yours respectfully,

"DR. TURNBULL.

J. R. H."

To this form of medication there are some other objections which have been made by Professors Roosa and Knapp, viz., that otitis media may supervene and perforation of the membrana tympani be caused by excessive sneezing, the result of the use of the douche; but no such results have followed the extensive use of this most valuable means employed by the author and others, in hundreds of cases both of ear disease and of ozæna with or without deafness. To those who are timid




about using it on these accounts, we would recommend the posterior nasal syringe, which is considered by distinguished authority "as being simpler, more thorough, and pleasanter to the patient." It should be made of hard rubber, so that any kind of fluid may be used in it. (It is shown at Fig. 29.) The best astringents to be used are infusions of krameria and

Fig. 29.—POSTERIOR NARES SYRINGE.




sumach, made from the bruised root of the former and the berries of the latter, or by dilution of the fluid extracts.

Fig. 30.—POWDER INSUFFLATOR.



The author has used a warm solution of tannic acid or of alum with good results.

Fig. 31.—STEAM NEBULIZER.




If these means fail in chronic cases, employ the insufflation of powdered cubebs, tannic acid, or cinchona bark. The instrument for this purpose is seen at Fig. 30. The powder



being placed in the opening, the sliding portion is pulled over it, and compression of the elastic portion by the thumb and finger forces out the powder in the form of a fine cloud. This can be alternated with the air-bag and catheter for minute portion of liquids.

If dry powders do not agree with the patient, then resort to the moist spray of steam (Fig. 31) charged with tannic acid, chlorate of potash, or chloride of ammonium (as seen in Fig. 33). By the air-press (see Fig. 32), the vapor of chloride of ammonium can be introduced into the middle ear, and is formed by the direct combination of hydrochloric acid, water, and aqua ammoniæ. In Fig. 33 the form of the apparatus is shown; also the proportion of each article is given on

Fig. 32.—AIR-PRESS.



the bottles. The tube A is to be attached to the catheter, and the tube B is for air to be blown with the mouth, rubber bag, or air-press.

There is another little instrument which has been found useful by some. It was devised by Siegle, and called by him



the "pneumatic aural speculum." It acts by exhausting the meatus of air while the surgeon keeps his eye upon the membrana tympani, so as to see if it moves as a whole or in parts; those parts where it does not move indicating the spots where bands of adhesions occur in the tympanic cavity. We have employed this instrument, but as yet have not had much good to result from it. Mr. Hinton\* states that "in

Fig. 33.—APPARATUS FOR GENERATING CHLORIDE OF AMMONIUM (MURIATE OF AMMONIA).





Fig. 34.—SIEGLE'S PNEUMATIC AURAL SPECULUM.



this respect it has great value. It consists of a box whose external opening is very oblique and covered with glass; to this is attached an india-rubber tube, by which it is exhausted by the mouth or by a syringe. The lower part is of three sizes, and is greased and fitted into the meatus, and is joined to the upper part by a screw, and when in place the membrana tympani can be seen through it and its movements noted." The whole arrangement is seen in Fig. 34.

\* An Outline of the Present Method of Diagnosis in Aural Surgery, by James Hinton, Med. Times and Gazette, Aug. 10th, 1867.



Mr. Hinton has found this instrument useful by virtue of its power over some forms of tinnitus. That this affection very often has its source in an excess of muscular irritability, and other causes leading to an increase of pressure on the labyrinth, he cannot doubt; and the more because, in a large number of cases, slight traction on the membrane by means of this instrument allays it for the moment. Politzer does not speak favorably of the action of the narcotic internal remedies which have been recommended in cases of subjective noises in the ears; but he has, however, observed a very great abatement of the annoyance after the external application of a narcotic, either in drops or as an embrocation, and places no reliance on counter-irritants. Internal remedies have but little influence save in exceptional instances, such as cases of constitutional syphilis, which should be treated by iodide of potassium. An air-bath or pressure of the finger on the mastoid process relieves the tinnitus for a time. If there be catarrh of the nose and adjoining regions, employ Thudichum's douche, first with a solution of common salt, and, after the discharge has become thin and watery, employ a solution of alum (ʒi to warm water Oj), and if the discharge do not cease, use a solution of sulphate of zinc (grs. ij to ʒi of water).

Of the use of electricity in tinnitus we would refer our readers to the article Electricity in Chapter XVIII.



## CHAPTER VI.

### SYRINGING THE EAR FOR THE REMOVAL OF ACCUMULATIONS OF CERUMEN AND FOREIGN BODIES.


SIMPLE as this process may seem, and plain as the indications may appear, yet this operation has been greatly underestimated and much abused. It should be remembered that the patient cannot syringe his own ears with advantage, and the surgeon who relies upon the performance of the operation in this manner need not be surprised to find the results unsatisfactory to himself and annoying to the patient. As the use of the syringe is not unfrequently required, either for the removal of foreign bodies or morbid products and accumulations, we deem it proper to give a few general directions with reference to its systematic employment. First of all, we insist upon having a perfect instrument, nicely adjusted and working smoothly and easily, so that all irregularity of the stream may be prevented. The irregular impulse of a jumping, jerking, knocking stream is unpleasant to the patient. Various forms of syringes for washing out the auditory canal are furnished to the physician.

The modification of Davidson's syringe by the addition of a proper nozzle and shield, as shown in Fig. 35, has been found useful, in the practice of the author, for cleansing the ear and removing foreign bodies, and if the nozzle is bent at right angles can be passed behind the uvula.

Should the elastic syringe not be at hand, the form shown in Fig. 36 may be substituted. This instrument can be made of metal or rubber, as in Fig. 37, and the nozzle should be blunt and smooth. Sharp points and edges may injure the auditory canal, even when care is taken, and they should always be avoided. The nozzle or tube of the syringe should in all cases be narrowed so as to enter



Fig. 35.—DAVIDSON'S SYRINGE.



in readily, yet allow any foreign matter to pass to one side of it.

Fig. 36.—CYLINDRICAL EAR-SYRINGE.




Fig. 37.—RUBBER EAR-SYRINGE.



This syringe should hold at least two fluidounces, or better, four, and the piston and rod should work smoothly and closely. The small nozzle should be so arranged as to be easily removed when the syringe is to be filled. Great advantage is afforded by the rings at the sides of the piston, as they furnish a means of having the instrument completely under control with one hand, while the other is left free to hold the ear of the patient, or for any other manipulation that may be required.


Prof. E. H. Clark, of Boston, has made a modification of Thudichum's (Weber's) nasal douche. Many years of successful practice and close observation of the needs of the aural surgeon give authority to his ideas. The instrument is described as follows: \* "But simpler, better, and less costly

\* Observations on the Nature and Treatment of Polypus of the Ear, p. 68.



than the syringe is an ear-douche which I have modeled after the plan of Thudichum's (Weber's) nasal douche (Fig. 38).

Fig. 38.—CLARK'S EAR-DOUCHE.



It consists of a glass jar (also made of tin), to the side of which, near the bottom, a flexible tube is attached. The jar holds about half a pint. The tube is three or four feet long, with an appropriate nozzle. In using the douche it is only necessary to fill the jar with the liquid to be used and to introduce the nozzle into the orifice of the meatus, and then raise the jar sufficiently high to force a current of the liquid into the ear. By this means a steady and continuous current is secured. It may be made forcible or gentle by elevating or lowering the jar. It cleanses the ear thoroughly and painlessly." This is probably the safest and best form of ear-douche that can be placed in the hands of the patient, and the one that will give the greatest amount of satisfaction.

The late Dr. S. P. Hullihen, of Wheeling, Va., invented an instrument to enable patients to perform the operation without assistance from another person. "It consists of a cylindrical metallic cup about three inches in depth and two and a quarter inches in diameter, with a semicircular piece cut out of its edge to adapt it to the ear, and a small metallic syringe of about half an inch in diameter. This syringe, in the first instruments constructed, was movable, and supported on two horizontal rods running across the cup, by means of which it could be made to slide forward or backward according to the will of the operator, but more recently the inventor has had it fastened permanently to the back part of the cup, thus enabling the patient to use it himself with more facility. The syringe has a large opening at the bottom covered by a valve, and the nozzle is made to come out from



the side, near the bottom, and rising up to the top, is bent forward, so as to project toward the external meatus.

"Dr. H. has further improved the instrument by adding a narrow rim to the outer edge of that portion which is to be pressed against the neck immediately under the ear."\*

In the description of the instruments for cleansing the auditory meatus by means of currents of fluids, the means and precautions for protecting the patient and clothing must not be forgotten. Indeed, too much care cannot be given to neatness in these operations. A large apron of gum-cloth or oil-silk, fitting closely to the neck, is invaluable. A towel should be added to prevent any drops leaking, and thus soiling your patient's clothes.

Wilde's basin (Fig. 40) is of great service in these cases.

Fig. 39.—HULLIHEN'S EAR-SYRINGE.





Fig. 40.—WILDE'S EAR-BASIN.



Its concave part fits accurately the curve beneath the lobe of the ear, and the perforated septum strains the clear water from the dirty.

\* Wilde on Diseases of the Ear, Philadelphia, 1853, p. 85.



If held by the patient in the proper position, closely applied to the cheek, no water can get down by its side. A small cup or bowl may be substituted, and requires to be held up, and closely applied by an assistant.

Toynbee's ear-spout is very serviceable. It consists of a spout-shaped piece held in position with a spring, which passes over the head. Its arrangement is best seen in Fig. 41.

Fig. 41.—TOYNBEE'S EAR-  
SPOUT.



This has been modified by having a piece of wire attached to it, so that it hooks over the auricle and requires no spring to hold it in place (see Box, Fig. 116).

Having completed the arrangements for protection, after deciding that the case is a proper one for the use of currents of fluids, we should select one of the forms of syringes represented in Figs. 35, 36, 37.

Grasping the auricle with the left hand and drawing it slightly upward, outward, and forward, so as to straighten the auditory canal, we inject a steady stream of water into the ear.

In employing the cylindrical or gutta-percha syringe, the thumb should be used to elevate and depress the piston, whilst the fore and middle fingers are inserted through the stationary rings, and thus steady the syringe. In Davidson's syringe, the compression of the bulb will serve to graduate the force of the current.

Warm or tepid water should be always used, as cold fluids cannot be tolerated and ought never to be employed.

It is well to cleanse the ear of pus, etc. by removing as much as possible with small pieces of cotton on the point of a probe, or by a camel's-hair brush, or piece of lint; and should examination reveal a perforation or an ulceration of the tympanum, some other means than the syringe should be resorted to in cleansing the ear.

In syringing the ears of nervous persons or those having tender ears, it is well to fill the concha with warm water.



Mr. Wilde bears strong testimony against indiscriminate syringing. "I am induced," he says, "to make some remarks, in hopes of putting a stop to practices not only useless, but in some cases positively injurious. I allude to the common habit of syringing indiscriminately, and also of probing the ears without proper inspection of the parts. The former is of daily occurrence; thus, a patient laboring under deafness, or perhaps what is worse, violent pain in the ear, is examined, either without the assistance of a speculum, or by means of some of the old divaricating instruments, most probably in a badly-lighted apartment,—at all events, without the membrana tympani being brought into view, a dark cavity being all that the explorer is able to perceive;—it is deemed advisable to try what might come out by squirting hot water into this dark passage for a quarter of an hour or longer; but nothing satisfactory following this operation, the diagnosis that there is no wax in the ear is accordingly made.

"Now there may be a collection of cerumen, which may not be removed by this operation; while, if the cause of pain or deafness is owing to an inflammatory condition of the auditory canal and its membranous extremity, a decided increase of the symptoms is induced by this useless and cruel proceeding: and I have frequently seen inflammation produced by unnecessarily syringing an ear where no wax was present."\*

At the Howard Hospital the author makes it a rule first to examine the ear with a speculum, aided by the sun or reflected artificial light.

If the canal is found to contain any soft living or dead morbid products, the assistant is directed to introduce a few drops of olive oil or glycerine by means of a pipette, and subsequently the ear is carefully washed out with an elastic syringe. If the substance be hard in character, as in cases of inspissated cerumen, a solution of bicarbonate of soda, grains five to ten to the fluidounce of water, is to be preferred. A teaspoonful of this solution is slightly warmed

---

\* Wilde on Diseases of the Ear, American edition, p. 87.



and placed in the ear for one or two nights previous to the operation of syringing.

Simple as the operation of syringing the ear may seem, there are sometimes unpleasant results which could hardly be anticipated. We have seen syncope follow the injection of tepid water into the ear for the purpose of removing a foreign body.

Wilde states "that syringing, or the mere introduction of the speculum, induces at times violent paroxysms of coughing. In others the act of injecting tepid water will produce syncope, although the patients tell us that it is not from the pain they feel. The removal of a granulation, or a small polypus from the meatus, will at times produce sickness of stomach, weakness, and even fainting."



## CHAPTER VII.

### FOREIGN BODIES IN THE EXTERNAL AUDITORY MEATUS, AND THEIR REMOVAL.

THE orifice of the meatus is arranged as though nature had placed a barrier against the ingress of foreign bodies. The obliquity and curved direction of the canal presents another obstacle, while, in the adult, the bristling hairs pointing outward and covered with the bitter secretion of the canal prevent the entrance of insects.

The subject of the removal of foreign bodies from the ear is one of much interest to the profession, as they are constantly devising new methods, while the old and simple plan is always at hand in the use of the syringe and tepid water.

It not unfrequently happens, after the wildest outcry and alarm, on the supposition that something has fallen into the ear, that the most careful examination will reveal nothing, and as soon as the mind is disabused of the false impression, all symptoms of pain and distress disappear.

But if the bodies that get into the ear are not removed, they sometimes fill up the meatus and become lodged there, sooner or later producing unpleasant results.

Bodies with smooth, rounded edges are comparatively harmless, but those with angular surfaces and rough edges may produce deafness. The author has removed a grain of indian-corn from the ear of a boy, which had remained there unnoticed or had been entirely forgotten for four years; and although the hearing was improved, yet the long-continued pressure had permanently thickened the membrana tympani.

A medical friend related to me a rather curious case of a foreign body in the external meatus, which he saw at the



U. S. Army Hospital (South Street, Philadelphia), to which he was attached for a few weeks during his military experience. A young man was in the hospital, doing some duty about the place. He had the external meatus of one of his ears closed by a spent bullet, a convex section of which was visible just outside the commencement of the narrow portion of the meatus. The bullet had lodged there and become firmly impacted. The man declined having it removed, and the surgeon in charge of the case did not insist upon an operation, so that the case was shown to my friend and others as a curious exemplification of some of the vagaries of gunshot missiles.

This man the author examined as late as 1870, and found that he had a slight dark-colored discharge from that ear, but was not entirely deaf, and otherwise enjoyed good health. He was kneeling when he received this spent ball, which sank into the mastoid cells.

It is true that, in the above instance, a foreign substance remained in the ear for a long time without producing severe injury or but slight deafness; yet, in many others that have come under the author's notice, foreign bodies have given rise to inflammation of the meatus resulting in deafness; also to acute and chronic inflammation, with thickening and perforation of the membrana tympani, followed by aural polypi, and even terminating in delirium, convulsions, and death.

The following is a list of some of the bodies which have come under direct notice, introduced either accidentally or intentionally: beads, shells, pins, needles, pieces of paper, wool, cotton, human hairs, bullet, pebbles, buttons, grains of corn, pieces of wood, straw, insects both dead and alive, etc. The means of extraction which have been, and are still, recommended in most works on general surgery and in our journals are not the most simple, but are rather complicated and often prove injurious, so that failure generally results, to the mortification of the physician.

For the comfort of those who have not succeeded in the extraction of a foreign body from the ear, it is stated that the late Sir Benjamin Brodie failed to extract a pea from the



ear in a case which he relates in a report of a clinical lecture published in the London *Lancet*, January 6th, 1844: "There was a case brought into the hospital in which there was some foreign body, I believe a pea, in the external meatus. *I tried all sorts of methods to get it out.* I could not use the forceps, and it nearly filled up the meatus, so that the water injected could not pass behind it, or it was so jammed that the water injected by the syringe would not wash it out. I said let it alone, let it remain there, the pea in all probability will dry and waste of itself, and then it will come out, or when it is decayed it may be washed away with a syringe; but I will make no further efforts to remove it now, for I may drive it into the tympanum and kill the patient."\*

The number of instruments which have been recommended by different writers are very numerous, such as hooks, scoops, curettes, etc. Professor Gross has invented an instrument which he believes will be found useful. It is composed of steel, is about five inches and a quarter in length, very light and delicate, cylindrical, and somewhat rough in the middle, to afford a good hold for the thumb and fingers, spoon-shaped at one extremity, and furnished with a little tooth or prong at the other. This tooth, which projects at a right angle from the shank of the instrument, is exceedingly small, and therefore admits of easy insinuation between the foreign substance and auditory canal. The curette or spoon is also very delicate, and bent considerably more than the ordinary cataract curette. (See Fig. 42.)

Fig. 42.



To facilitate the use of this instrument, chloroform is administered to the extent of entire obliviousness. This is absolutely necessary when the patient is a child or a


---

\* See, also, a similar case in "Defects of Sight," by T. Wharton Jones, edited, with additional remarks on "Defects of Hearing," by the author, p. 177. Philadelphia, 1859.



nervous, excitable adult.\* The objection to this instrument is, that it is made of steel, and, consequently, not being flexible, cannot be altered in shape to suit the circumstances of the case. I have seen and tried almost all the ordinary mechanical contrivances, and have arrived at this conclusion, that the syringe stands first in efficiency for the removal of all kinds of foreign bodies,—all other aids are secondary. This opinion is based upon many years' experience. It is the easiest and safest method, no matter whether the foreign body be large or small, light or heavy, irregular or even, occupying a small space or plugging up the meatus entirely, of recent introduction or of long standing. I have never failed with this instrument, alone, or in conjunction with a soft silver curette (Fig. 43), curved as seen in the cut; the flat end of which, as well as its shaft, can be accommodated to the varying conditions presenting in these cases.

Fig. 43.



The curette of Prof. Gross I find the best instrument for the removal of foreign bodies from the nose; but only secondary in cases of bodies in the ear. In two cases which occurred recently, one a bead and the other a button in the nose, I succeeded in removing them with this instrument.

The use of the syringe is not of recent application, but the credit of calling the attention of the profession to its universal applicability for the removal of foreign bodies from the auditory canal is due principally to Mr. Carpenter, of Castlecomer, Ireland.†

“The first case,” he observes, “brought to me some years back, was one in which the foreign body was a garden-pea, which, as is usual, was pushed in as far as the membrana tympani by the interference of the child’s friends. The instrument I selected was a very small forceps, with blunt

\* American Journal Med. Sciences, Jan. 1842, p. 231.

† Dublin Medical Press, June 10, 1841.



points, with which I could catch the pea, but could not move it, and on endeavoring to do so, caused intolerable pain. It immediately occurred to me to inject cold water with force, in order to, at least, displace the pea. I did so with a two-ounce syringe, and found the pea so far displaced as to lie at the orifice of the meatus, whence I removed it without further trouble or pain to the child. Since that I have extracted many peas in the same way."

About six months after, Mr. Carpenter extracted a pebble, which was so irregular in shape and so large that he was surprised that the child could have borne to introduce it. The meatus was not inflamed, but the pebble was so far in and wedged in its place, that to touch it ever so lightly with any instrument was almost enough to throw the child into convulsions. The first injection failed; but while using a second, the pebble, much to his satisfaction, was forced out upon a napkin.

Dr. James Bolton\* has related a case, interesting both in its history and result, which occurred in a girl aged thirteen years, "in whose meatus a grain of coffee had remained for seven years. He syringed the ear with warm water and Castile soap. On throwing in about the third syringeful, with some force, a black-looking substance shot into the basin, which, on examination, proved, beyond all doubt, to be the half of a coffee grain."

Mr. Edward Hatfield† removed from the ear of a boy by this means a piece of slate-pencil, which had been pushed farther in by the schoolmaster in a fruitless attempt to extract it.

Samuel Cooper‡ relates the case of "a child two years and a half old into one of whose ears a pebble and into the other a French bean had been pushed by another child, and remained there for ten months, causing complete deafness and extreme suffering." By throwing tepid water forcibly into the ear he soon dislodged these foreign bodies, which lay close against the membrane of the tympanum, entirely hidden by the swollen state of the lining membrane.

---

\* Medical Examiner for 1842, p. 613.

† New York Edition London Lancet, July, 1844, p. 97.

‡ Cooper's Surgical Dictionary, art. Ear.



Dr. J. Marion Sims\* gives the following illustrative cases of the ease, safety, and certainty of this method of removing foreign bodies from the auditory canal:

CASE I.—A little negro boy, four or five years old, was sent to Dr. B. R. Jones, Montgomery, Alabama, with a piece of cut-glass in his left ear, and his master had made some fruitless attempts to extract it with a knitting-needle. The meatus was swollen, inflamed, and very tender. I happened to be at the doctor's at the time. We made many ineffectual efforts to get this substance out, using a bent probe, scoop, and forceps. We also used a syringe, but with a view of washing out some coagula of blood, the result of our awkward and clumsy poking with unsuitable instruments. The injury inflicted by our rude attempts at extraction was by no means trifling. After a second attempt on the following day with a pair of long, narrow-bladed, curved forceps with no better success, and just before giving the case over in despair to "bleeding and general antiphlogistic course," I picked up the syringe for the purpose of washing out a coagulum of blood that was hanging from the meatus. I threw in two or three syringefuls of tepid water, but the exuded blood still stuck there. I then requested Dr. Jones to take hold of the external ear and pull it outward, upward, and backward, so as to make the canal straight and allow me to see into the meatus. He did so, and I injected forcibly. My astonishment was equaled only by my delight at seeing the glass "ear-drop," as it proved to be, shoot out with great velocity. It was five lines in length, a little over three in width, and nearly two in thickness, slightly flattened on one side and oval on the other.

Its peculiar shape precludes the possibility of holding it between the blades of the forceps even now. Its size, in connection with the tumefaction of the meatus, shows that it must have been perfectly impacted, and I doubt very much whether it could have been removed at that time by any other instrument.

CASE II.—A little negro girl was sent by Dr. Hugh Henry,

---

\* American Journal of the Medical Sciences, April, 1845, pp. 340, 341.



having a foreign substance in her right ear. It gave her some pain, but as there had been no useless and mischievous probing of the meatus, it was trifling in comparison with that suffered by Case I. I caught hold of the external ear with my left hand, putting it strongly on the stretch, so as to make the meatus a straight channel, while with the other hand I threw in a syringeful of tepid water forcibly against the tympanum. A single injection was sufficient to dislodge the foreign substance so far as to make it easy of access. It was a segment of a berry of the China-tree (*Melia azedarach*).

CASE III. was a negro man Lewis, aged twenty-eight, who had been deaf, with discharge from his ear from the presence of a cherry-stone, which was removed by the same means. It filled completely the bottom of the meatus, and was perfectly impacted and quite immovable, and from his history of the case must have been in the ear for more than *twenty years*.


I will only relate a few cases selected from many occurring in my own hospital and private practice.

CASE I. *A button in the ear*.—June 7th, 1863, a physician brought to my office, in the afternoon, a boy aged six years, who had placed a gilt button in his ear about ten o'clock, A.M. His parents on discovering it took him to this physician's office, and he, after trying several forms of forceps, hook, probe, etc., found himself unable to extract it. He then called in a second physician, who, on examination, would not attempt its extraction for fear of doing more injury to the meatus, which was bleeding, swollen, and very tender. On examining with a speculum and bright light, I could discover the button on the membrana tympani, its eye being toward me in an upward direction, but bloody and flattened, so as to prevent my passing a hook into it. I therefore requested the physician to take hold of the external ear and pull it outward, upward, and backward. While he did so I injected forcibly a continuous stream of tepid water, which changed its position so as to bring it to that portion of the canal where I could seize it with a pair




of forceps, which are shown at Fig. 42, a modification of Wilde's.

Fig. 44.



My only objection to Wilde's angular forceps is that they will not hold securely, being apt to slip; and they twist when much strain is put upon them, which they will not do as modified by the author. Wilde's forceps are seen at Fig. 45.

Fig. 45.



CASE II. *Pebble in the ear.*—May 17th, 1864. Agnes R., aged five, whilst playing with pebbles from a sand-heap, placed one of them in her right ear. It remained there for some time, when she became alarmed, and upon inquiring of one of her companions "if it would kill her," was answered in the affirmative. She confessed to her mother, who became much alarmed, and took her daughter to the family physician, who tried to extract it. Owing to the child's restlessness and screaming, he only inflicted injury on the meatus without being able to extract it. On the 18th she was brought to Howard Hospital, and owing to her excited state it was considered best to place her under the influence of a mixture of ether and chloroform, recommended in such cases in 1859. On examination nothing could be seen but coagulated blood; but after the injection of tepid water, as before described, the pebble was washed out without pain or knowledge of the child. A soothing application was then applied to the ear, and the case discharged.



CASE III. *A grain of corn in each ear for four years.*—March, 1865. Robert B., aged six years, was brought to Howard Hospital by his mother on account of deafness in both ears since the age of two years. The mother could give no information in regard to the cause. It was not the result of cold, measles, or scarlet fever. On examination with a speculum and bright light, some foreign body was discovered in the ear, covering the membrana tympani. The parts were carefully syringed, and in one of the ears was found a mass of softened indian-corn. A second and even a third injection was employed, and a similar softened grain of corn was found in the opposite ear. The hearing was improved after some further treatment, but the membrane in both ears was rendered opaque from the pressure of the indian-corn for so many years.

CASE IV. *A grain of indian-corn in each ear for twenty-four days.*—Ann E. Moore, aged four years, residing in Norristown, Pennsylvania, was sent to me by Dr. W. R., with a grain of indian-corn in each ear, which had remained in for twenty-four days. Efforts had been made by the physician to remove them by means of forceps, etc., but they only pushed them more firmly in, impacting them in the oval or ellipsoidal auditory canal. They were covered with dried blood, and the whole integument was swollen around them.

The child was in a highly irritated, nervous state, from the repeated fruitless attempts. With the assistance of her uncle, A. G. Feather, Esq., I placed the child under the influence of a mixture of ether and chloroform, and then, having the ear held as before mentioned, I injected forcibly several ounces of tepid water, and in a few seconds the water collected behind the body and washed it out, so that I could grasp it with my short angular forceps. The same result followed in the opposite ear, much to the delight of the mother and uncle of the child, who had witnessed the previous abortive efforts.

The child was then allowed to recover from the effects of the anæsthetic; the ear was carefully dried, and a portion of oil of belladonna was dropped in to remove irritation, and the case discharged.



Some bodies in the ear, more especially those that are rounded off and have no sharp edges, will do no harm; but there are many others which we know, from having witnessed the results produced, cause deafness, etc.

The following cases are from Toynbee,\* and were treated by the same means:

CASE I. *Slate-pencil in the ear of a child.*—J. S., aged seven, was brought to me at the St. George's and St. James's Dispensary, on November 28th, 1849, her mother stating that she had pushed a piece of slate-pencil into the right ear. She had complained of pain, and had not slept well. Upon examination by means of a speculum and lamp, a rough piece of slate-pencil was observed lying on the floor of the meatus, one end being apparently in contact with the membrana tympani, while the other looked toward the orifice of the meatus. It was easily removed by the syringe and warm water, and found to be nearly half an inch long and a quarter broad. The patient did not complain of pain, and in a few days the membrana tympani was perfectly healthy.

CASE II. *A percussion-cap in the ear for fifteen years.*—C. E., aged twenty-six, was admitted at St. Mary's Hospital in November, 1852. She said that lately she had felt somewhat deaf in both ears. Upon examination a mass like hardened wax was observed near the orifice of the left meatus. It was removed by the syringe, and proved to be a percussion-cap surrounded by wax. The patient said that she remembered, when about eleven years of age, she put a cap into her ear, but she believed it had been taken out, and never thought of it since. The membrane of the tympanum was concave, the Eustachian tube natural, and the power of hearing nearly gone.

CASE III. From the same authority, showing the injurious effects from the use of instruments in removal of foreign bodies from the external meatus.

"A shell in the meatus; removal by instruments; paralysis of the portia dura nerve. Miss A., aged fourteen, consulted me in May, 1853, on account of deafness of the right ear, and

---

\* Cases of Foreign Bodies removed from the External Meatus, p. 71.



paralysis of the right side of the face. Her father stated that, eight years previously, she had, in play, put a small shell into her ear; that the surgeon, in endeavoring to remove it, forced it deeper into the ear, breaking the shell, and causing intense pain. After its removal there was much discharge from the ear, and in a few days the muscles of the right side of the face lost their power, which they have not regained. Upon examination there was no vestige of the membrana tympani; the mucous membrane of the tympanum was very thick and red, and there was not the slightest power of hearing."

As an agent to assist the syringe in the removal of a shell from the ear, there can be nothing better than a delicate bent wire. By this means shells have been removed with great facility at Howard Hospital, by hooking the bent wire into the opening of the shell; but when they are flat, the reflex stream of water brings them near enough to be seized with a pair of angular forceps.\*

The author is confirmed in this opinion by the most distinguished aural surgeons of the present age, namely, Wilde, Toynbee, Tröltsch, and Hinton.

In cases where the foreign body is irregular or ovoid in shape and fixed firmly, the plan of Jonathan Hutchinson, F.R.C.S., may be adopted.

He has applied the principle of removing a cork from a bottle by a string. The string, in this case, when employed in the ear, is a silver wire of five or six strands, in the form of a loop. This will be less apt to do any injury than the forceps, scoop, etc.

"I have a horror of the use of the scoop [so has the author] so generally employed, because it can only act as a sort of lever, and because I have repeatedly known it to do much damage. We have," Mr. Hutchinson observes, "a specimen in the museum, in which a bean was, by its agency, thrust through the membrana tympani, and produced death by its irritation."

The forceps are little better.

---

\* Transactions American Medical Association, 1865, p. 429.



Another method of removal is proposed by A. G. Brown, by atmospheric pressure (suction), by means of a few inches of French vulcanized rubber attached to an ear syringe so as to produce a vacuum.

The following was treated by Dr. James Collins, of this city:\* F. A., aged five, was presented by his mother, who stated that while at play he had inserted a pebble in his ear, which could be plainly seen at the time, but had been pushed in more deeply by attempts to remove it, made by herself and other members of the family. The child was quite peevish, and complained of pain in the ear. On examination found the auditory canal swollen, and the surface covered with dried blood. A syringeful of tepid water, gently applied, removed this, when there was seen, deep down in the canal, a foreign body, white, hard, and easily recognized as a pebble. Chloroform was administered, and the plan suggested by J. Hutchinson, F.R.C.S., in such cases, adopted for its removal.

After gentle manipulation, the loop of a fine silver wire was passed around the body, and by gentle traction the pebble was removed without difficulty. The ear was filled with warm water, which was allowed to remain several minutes. In two days all pain and swelling had disappeared. The pebble was ovoid, and weighed four grains.

#### ASPERGILLUS IN THE AUDITORY CANAL.

Under the head of foreign bodies in the auditory canal the peculiar vegetable parasite *aspergillus glaucus* may be mentioned. The presence of these parasites produces irritation and inflammation. They accumulate, forming a lardaceous mass.

Dr. Robert Wreden,† of St. Petersburg, in a description of six cases of the affection, thinks they simulate inflammation of the external auditory canal, and the appearance somewhat resembles deposits of cholesterine. The microscope, however, exhibited no cholesterine, but on the contrary

---

\* Med. and Surg. Reporter, Philadelphia, vol. xx. No. 16, p. 308.

† Archiv für Ohrenheilkunde, 1867.



numerous long mushroom tubes with roundish spores. The auditory canals were half filled up with the white lardaceous mass. This was removed by forceps after syringing had failed. The epithelium of the membrana tympani and canal was found to have been removed, the cutis was dark red, and the malleus and light spot on membrane not visible. There was, of course, loss of hearing, and tinnitus aurium, dependent upon the closure of the canal and pressure on the drum.

Mayer reported a case of the same in 1844.

Pacine observed and reported a case in 1851.

Schwartz observed this affection in 1865. He supposes that it is probable that exudation, inflammation, etc. of the canal occurred before the appearance of the mushroom-like growths.

In the year 1870 Dr. Roosa\* reported two cases, and has collected those noticed above. Three cases have since been treated by Dr. J. O. Green, of Boston, and from his experience, a thorough repeated removal, with the use of water, is sufficient to kill the growth.

Dr. Robert Wreden, of St. Petersburg, has subsequently added eight cases, making fourteen in all, and has published a monograph on the subject.†

"There is hardly a doubt, as observed by Roosa, that an affection of the ear caused by the 'aspergillus' has existed for a long time, if not always; but it has not been recognized for the simple reason that diseases of the ear have been, until within a few years, very much neglected, except by isolated practitioners. Since his attention has been turned to the subject, and in this I quite agree with him, he could recall obstinate cases of inflammation of the auditory canal which contained this growth."

The subjective symptoms of the growth of "aspergillus" in the auditory canal are very similar to those from impacted cerumen. There is a sensation of fullness, tinnitus aurium, vertigo, pain, and impairment of hearing. Pain is

---

\* Am. Jour. Med. Sci., Jan. 1870, p. 105.

† Die Myringomyelosis Aspergillina. St. Petersburg, 1868.



not one of the common symptoms of inspissated cerumen, although it does sometimes occur. This may be due, as before stated, to the inflammatory action which always precedes the growth of this fungus. The subjective appearances consist in the adherence of flakes or scales of what may be mistaken for epidermis, on the walls of the canal and the outer surface of the membrana tympani. It requires to be removed by the forceps, and the integument beneath is found reddened and sensitive, and in a few hours the growth reproduces itself. In one variety little black points are seen in the mass, but it requires the aid of the microscope to confirm the diagnosis. The *aspergillus flavescens* and *nigricans*, as well as the *stamphyllium polymorphum*, have been found in the auditory passage. In regard to the etiology, Dr. Wreden says, "All these varieties of *aspergillus* are only varieties of the ordinary mould fungus."

The etiology of this disease is no doubt owing to inflammation and deposit of fungus upon the surface prepared for it by the inflammatory action.

The following is the record of a case from Roosa:

CASE I.—"Was consulted by Mr. S., aged fifty-one, on account of impaired hearing, vertigo, pain in the ears, and tinnitus aurium. Vertigo was the symptom upon which the patient laid the most stress, and of which he was most anxious to be relieved. He said that he was so dizzy whenever he attempted to walk about, as to be unable to attend to his ordinary business. His condition in other respects was excellent. Patient stated that he had heard perfectly well until two months since, when he was attacked with the aural symptoms narrated above, which had been aggravated since their inception. He had been treated by the instillation of oils and so on. He could hear my watch about one inch on the right side and not at all on the other. Both auditory canals were found filled with a tenacious material which could only be removed by the forceps. It was several days before I could remove the firmly-adherent coating to the canal and membrana tympani.

"The morbid product was immediately examined by Dr. Lewis, and found to be a specimen of the *aspergillus*



*flavescens*. Its removal gave the patient great relief, but on the reappearance of the growth, which was in two or three days after its removal, the vertigo and tinnitus returned.

"The membrana tympani was intact, but lustreless and rigid. The Eustachian tube opened sluggishly, and there was all the evidence of aural catarrh, besides the affection of the canal and the outer layer of the membrane of the tympanum.

"The free use of warm water, with an astringent, finally subdued the morbid process in the canal so that the patient was able to make a journey to the South. When he left the doctor's care, October 18th, the auditory canals were entirely free from abnormal secretion, the hearing was improved so that the watch was heard from five to six inches on the right side (thirty inches normal distance), and from one to two on the left. The dizziness was entirely gone, and the tinnitus ceased to be annoying. The catarrh of the inner ear, as shown by rigidity of the membrana tympani, sluggish action of the tubes, and impairment of hearing, still continued."\*

#### MEANS TO PREVENT A REPRODUCTION OF THE PLANT.

Roosa, Green, and Knapp agree that the repeated removal with the use of water (warm) is sufficient to kill the growth. In the monograph of Wreden a long list of agents is given which may be found useful. Some of these are alcohol, bichloride of mercury, acetate of lead, tincture of iodine, and carbolic acid. He gives the preference, however, to the hypochlorate of lime (*calcaria hypochlorosa*), which he recommends to be used of the strength of one or two grains to the ounce of water at each application. Fowler's solution of the arsenite of potassa ranks next to the lime.

The writer prefers in such cases the hyposulphite of soda or the bisulphite, having found them both useful as antiparasitic, both animal and vegetable. The solution is made with water alone, or in combination with carbolic acid and

---

\* American Journal Medical Sciences, op. cit.



glycerine in the proportion of three to five grains to an ounce, employed warm.

In connection with this subject it is well to notice that maggots are sometimes found in the external meatus. The following case, reported by the late Dr. R. P. Thomas,\* will serve as an illustration. "He says that while a lady was standing under a tree, a small fly (probably a common maggot-fly) darted into her right ear, and remained there, in spite of all efforts at extrusion, for fifteen or twenty minutes. Eight hours later a sensation of motion was experienced in the ear; at first faint, but afterward more distinct, and increasing in violence until her powers of endurance were tried to the utmost. Forty-six hours after the ovi deposit was made his attendance was requisite. Her agony from this motion and waving sound in the ear was indescribable. The external meatus proved to be unusually narrow and deep, and therefore rendering an examination with the otoscope unsatisfactory. Having detected a tremulous jelly-like mass on the tympanic membrane, he attempted its removal with an ear-scoop but failed. A resort was then made to injections of strong soapsuds, and in a few minutes a maggot crawled out far enough to be visible, and thus determine its character. Removal by mechanical means without injury to the delicate membrane was impossible, and chemical solutions strong enough to affect their vitality, were deemed hazardous to the same structure. Sweet oil or almond oil will generally float out a bug or other small insect that may stray accidentally into the meatus, but their efficiency in removing parasites that have the power of attaching themselves to an animal membrane being deemed questionable, he resolved finally upon trying a solution of grs. xx of tannin in an ounce of pure glycerine. The reason for its employment was twofold, viz., for exclusion of the air from their breathing pores by the glycerine; and the probability that the violent effect of the tannin upon the delicate surfaces of the parasites would cause such a degree of activity as would bring them in reach of the forceps.

---

\* Am. Jour. Med. Sci., January, 1861.



"The result confirmed the anticipation, and twenty-three maggots, each three-eighths of an inch in length, were removed by a slender pair of forceps in the course of a few minutes. The relief was immediate and complete, in two or three hours the cerebral excitement was allayed, and the following day the patient was entirely free of all uneasiness. Several months have elapsed without manifestations of any injury to the ear or the sense of hearing from the accident."



## CHAPTER VIII.

### POLYPUS OF THE EAR.

ACCORDING to the best authorities,\* both ancient and modern, polypus is understood to be a fleshy tumor, various as to color, form, structure, and consistence of tissue. But there is one common characteristic connecting all varieties of polypi: they all originate and develop themselves upon the mucous margin of the natural orifices,—nose, pharynx, rectum, neck of the uterus, and auditory meatus. It is with this view that authors define polypus of the nose, of the ear, etc. “The membrane of the auditory canal,” says Boyer, “has an analogy to the mucous membranes, and, like them, is subject to polypous excrescences.” “The name of polypus is given to an excrescence developed upon the tegumental membranes.”—Polypi are commonly considered as peculiar to the mucous membrane, yet certain parts of the skin sometimes present excrescences exactly similar to the polypi, as the entrance to the vagina and to the auditory canal.† But the definition which especially applies to the nasal polypi, for which it was created nearly two thousand years ago, is not applicable to polypi of the ear. Triquet,‡ in his work, states that polypi of the ear are nothing else than a sort of fleshy excrescence which originate and develop themselves upon the external and internal membranes of the ear, when affected by chronic and puriform discharges, in the same manner in which we see the inflamed surface of old issues and ulcers sometimes covered with exuding and bleeding flesh, etc. There is nothing similar met with in the etiology of the polypi of the nose; not so, at least, if we are to credit the descriptions given by our best writers. Thus they confound

---

\* Historical Observations of Ancient and Modern Authors, from Triquet.

† Dictionnaire de Médecine, art. Polypi.

‡ Traité pratique des Maladies des Oreilles, p. 309.



two diseases which have really nothing in common but the name, and consequently introduced into practice a deplorable abuse, that of subjecting both diseases to the same treatment. The important differences which, in a few words, Triquet has just pointed out, seem to indicate another plan to be pursued. The ancient authors,—Hippocrates, Celsus, Galen, Ætius, and Paulus Ægineta,—in treating of the polypus, in many passages of their works have intended to be understood as speaking only of polypi of the nose. Thus Hippocrates,\* in a few words, describes three varieties of the polypus which develop themselves in the nasal cavity. Celsus says, “Polypus verò est carnicula quæ narium ossibus inhæret:” “the polypus is a fleshy excrescence which attaches itself to the bone of the nose.” According to Paulus Ægineta, “Polypus is an excrescence, the roots of which extend into all the points of the nasal fossæ.” Such is the tradition which we find reproduced by the surgeons of the Middle Ages,—Guy de Chauliac, Pigray, etc.† But polypi of the ear are not contemplated in their works. We thus arrive at the sixteenth century, represented by A. Paré,‡ who had already perceived that “la fluxion tombée aux oreilles qui fait apostème, engendre ulcère hypersarcorse ou chair supercroissante. Et plus loin, quand cette chair est fort profonde, elle ne se peut guérir.”—“It is true that a concise and very suggestive phrase is thrown in, as it were, by chance, into a chapter only known to contain a few theoretic views of deafness.”

To Fabricius de Hildenis§ we owe the first positive case of polypus of the ear scientifically treated and reported. That case has been imperfectly quoted by all writers subsequent to him, and as it has been the theme of the dogmatical descriptions published since his time, I deem it indispensable to transcribe it accurately.||

---

\* Hippocrate, Œuvres complètes, trad. par E. Littré.

† V. Guy de Chauliac, *Le Maître en Chirurgie*, p. 392; Pigray, *Epitome Chirurgie*, p. 428.

‡ Œuvres complètes, édit. de Malgaigne, t. ii. p. 601.

§ Cent. 3, obs. 1 (anno 1604).

|| *Traité pratique des Maladies de l'Oreille*, t. ii. p. 26.



“CASE I. *Scirrhus fungus protruding from the ear.*—In 1594 a young girl of noble family, Marguerite de Martines, was attacked, at the age of eight years, by the smallpox. The eruption was scarcely perceptible. A year afterward (1595), she felt violent pain, with no less violent shootings in the right ear. A few days later an abscess opened in the meatus, and the discharged matter resembled the offscourings of flesh. Very soon that discharge became yellow, and, after intermitting, it at length established itself permanently, but the pain no longer existed. Five years later (in 1600), the parents, having remarked that the hearing was weak on that side, had the curiosity to examine the diseased ear, and were surprised at finding that it was completely obstructed by a fleshy excrescence (*excrecentiam quamdam carnosam*). The discharge still continued, but as the noble patient showed a great repugnance to surgical operations, her family hesitated to call in the aid of art.

“It was not until the winter of 1604 that the parents consulted Fabricius.

“The fungus excrescence, of which he has left us many designs in his fine work, had the form of the canal; that part which protruded from the ear was irregular, hard, livid, and spreading upon the concha by several muriform tubercles. The operation was appointed for the return of fine weather, and Fabricius employed the ligature on the 23d of May, 1604, to compress the lower part of the tumor. It came away on the third day, but its roots were only destroyed by the aid of numerous cauterizations, and the case was not complete until toward the month of August in that same year, 1604. In other words, the treatment lasted three months.”

Having removed a number of these foreign bodies, the author most fully agrees with the conclusions arrived at by Triquet, who has collected some fifty cases from Fabricius de Hildenis, Leschevin, Scultet, Itard, Kramer, Toynbee, Pelletan, Sédillot, etc. From the Triquet practice, in every instance, there was a discharge before the polypus was known to be in the ear. Secondly, in no instance was there perfect restoration of the function of hearing. Thirdly, if the poly-



pus was situated near the membrana tympani which was almost always more or less destroyed. Fourthly, the most successful treatment was the extraction of the polypus and cauterization. Fifthly, there are three varieties, vascular, gelatinous, and fibrous.

The following is an account of eight cases of polypus of the ear occurring in the author's practice:


CASE I. *Vascular polypus of the left meatus; epithelial of Clarke; extraction; otorrhœa; recovery.*—December 31st, 1857. George M., aged thirty-five years, by occupation a shoemaker, a native of Ireland, was suffering from entire loss of hearing in the left ear, with tinnitus aurium. The patient was of sanguine temperament, with scrofulous enlargement of the glands of the neck. On cleansing the ear by the use of the syringe and warm water, a vascular mass of polypus was discovered, like a strawberry in form, situated about one-half the distance from the membrana tympani. As it was soft, and bled on the slightest touch, I was unable to use the wire ligature, and therefore resorted to the lever-ring forceps of Toynbee. The patient sat on an operating-chair, and the light of the sun was made to shine directly into the ear, which was drawn backward by an assistant so as to dilate and straighten the meatus as much as possible. A large-sized speculum was next introduced, and then, with the right hand, the lever-ring forceps were gradually pushed down to the polypus with the rings in contact. As soon as the rings reached the polypus, they were allowed to separate, and when thus apart were made to surround the growth; when this was effected, the rings were brought into contact by pressing the lever.

The instrument was then withdrawn, bringing with it a part of the polypus.

January 2d. The small portion which remained in the ear resisted the use of astringents and caustics, and was entirely removed by a second operation.

Afterward the hearing distance of left ear was four inches;

Fig. 46. — EPITHELIAL OF CLARKE, AND VASCULAR OF TRIQUET.





membrana tympani opaque. Directed use of iodine externally and internally.

January 9th. Still improving, and otorrhœa less. January 16th, discharged, cured.

CASE II. *Large vascular polypus at the base of the external meatus; removal; perforation of the membrana tympani; otorrhœa; recovery.*—Elwood M. C., aged eighteen, a student, was sent me by his father, October 17th, 1860, a distinguished medical man of a neighboring county, who had himself removed a portion of the polypus. Since then it had so increased as to be a great annoyance to him. He was deaf, and the discharge profuse and offensive.

October 17th. Made an attempt to remove it by using Wilde's snare, but only succeeded in part; the bleeding being very free, was checked by washing out the ear with ice-water. Caustic potash was then applied several times, but failed to remove it.

December 28th. Removed the whole mass by means of the ring forceps of Toynbee; the bleeding was so profuse as to cause fainting, but by means of ice-water and plugging the ear it was stopped. The polypus was situated near the membrana tympani, which was opaque and concave.

He subsequently was appointed a medical officer in the U. S. Navy. He had some deafness in the ear, but no return of the polypus up to date of his death.

CASE III. *Small vascular polypus; otorrhœa; removal.*—M. G., aged sixteen, in February, 1864, became deaf in one ear, with a discharge of blood for months, followed by pus. On examination found a small vascular polypus near the membrana tympani. The removal was attempted with Wilde's snare, which cut through it. I removed as much as possible, and then applied a solution of persulphate of iron, also directed a wash of alum, ʒj, laudanum, ʒj, and water, ʒj. Hearing much improved and polypus did not return.

CASE IV. *Vascular polypus of left meatus; otorrhœa; extraction; recovery.*—Martha G., aged sixteen, a young, delicate girl, consulted me February 18th, 1864. Had been deaf in the left ear, with a free discharge of pus and blood, for seven months. On clearing out the pus, etc., discovered a small vascular



polypus, which was removed by the ring-lever forceps. Directed a tonic, with astringents, which removed all remaining traces of it.

The hearing was improved by pursuing this tonic course for several months.

All the foregoing described polypi were of the vascular character, and racemose in form, attached to the central stem or root. They were covered by tessellated epithelium, and when examined microscopically were found to be made up of soft, fibrous cells, very vascular, and which were easily broken, and found filled with blood and serum.

Some polypi are very small, as in the last case recorded, whilst others are so large as to fill up the entire meatus, and even project from the orifice. They are, as a general rule, attached to the inner half of the tube.

CASE V. *Soft gelatinous polypus on the left meatus for ten years ; extraction ; recovery.*—Mrs. R. L., aged fifty, the wife of a medical bookseller and publisher of this city, consulted me, December 4th, 1864, on account of a persistent discharge, with deafness so great as not to hear the tick of the watch. This she had suffered from for years, and applications had been made of various kinds, by advice of her family physician and others, without any benefit except of a temporary kind.

Her general health is good, but she is of a bilious, nervous temperament. On examining the left ear there was seen, deep in the tube, which was very narrow, a pale-colored polypus, attached to the walls of the meatus, filling up and assuming more of a globular form as it approached the meatus externus.

She did not wish any operation, and I attempted to remove it with caustic potassa, nitrate of silver, etc., but was not successful. After a time she consented. To give as little pain as possible, I removed it with a ring forceps in small portions. By the use of powdered alum, and other mild astringents, the discharge entirely ceased after a treatment of five months. Her deafness in that ear still continued, but she was entirely relieved from the offensive discharge, has no pain in the ear, and has had no return of the polypus.



CASE VI. *Fibro-gelatinous polypus; otorrhœa; removal; recovery.*—November, 1867. Mrs. C. A. R., aged thirty-eight, dressmaker, was sent to Howard Hospital (August, 1863) by her physician. She had suffered from the age of six years from deafness, and more or less discharge, following scarlet fever, but within the last year she has been unable to hear with the right ear, owing to a polypus that had grown so as to protrude outside the ear, the odor of the discharge being so offensive as to interfere with her business; even when sitting at rest, the discharge was so profuse as to pass under her chin and on her neck. On examination, the polypus was found somewhat abraded, and was attached about the middle of the meatus. I therefore resolved to remove it with Wilde's snare, which is the best instrument for the removal of this variety of aural polypus, and which is briefly described on page 124. The patient was seated on a chair with a good sunlight; in this case it was not possible to use the speculum. The cross-bar was pushed forward, and a noose made of the wire at the small extremity, of sufficient size to include the morbid growth, which it was made to surround and press down by means of the stem; the cross-bar was then drawn up smartly to the handle, giving it a single twist, which drew it up from its attachment without cutting it across. Bleeding took place, but by syringing the canal with cold water it was checked. On examination, it was found entirely removed. The patient did not return, but upon making inquiry of her physician, he sent me the following note:

“PHILADELPHIA, November 6, 1867.

“DEAR DOCTOR,—I called to-day on Mrs. C. A. R., and learned from her that, previous to the operation, her hearing was completely obstructed. Since the operation by you her hearing has very much improved, accompanied at times, from taking cold, by a slight discharge, though no appearance of another polypus. With this hasty sketch, believe me to be,

“Yours truly,

“LEWIS S. SOMERS.

“To Dr. TURNBULL, 1208 Spruce Street.”



Under the microscope the interior of the gelatinous polypus is composed of corpuscles and fibrous tissue, proving that the name fibro-gelatinous polypus is the proper one.

No escharotic produces much effect upon fibrous tissue, so that extraction is the best method of removal.

CASE VII. *Large fibro-gelatinous polypus of Toynbee; fibrous of Clarke; otorrhœa; removal; recovery.*—Ann R., aged sixteen, applied at Howard Hospital, August 12th, 1867. She was suffering from pain and discharge, and noise in her ear. On examination, the ear was found very much distended by one of the largest polypi which I have removed. Her statement was, that a similar one had filled up her ear, and had disappeared of itself, but that this one had remained to her distress for six months. It was removed with the snare. The bleeding was very profuse, requiring the use of the persulphate of iron and plugging of the meatus.

The hearing in this case was improved, and the discharge stopped. The form of the polypus is seen at Fig. 47 much reduced in size by being in alcohol for two years. The microscopic appearance is also shown.

Fig. 47.



Mr. Hinton has stated that, as far as his experience goes, polypi in the meatus invariably spring from the dermis, though they also have attachments to the walls of the meatus.

CASE VIII. *Vascular polypus in the left meatus; loss of membrana tympani; otorrhœa; both Eustachian tubes closed; removal; recovery.*—August 15th, 1856. William J. M., aged twenty-eight; otorrhœa in both ears; had scarlet fever at the age of sixteen. Could only hear the tick of the watch when applied over the temporal bone. Both Eustachian tubes closed.

He is bent almost double with rheumatism. Right ear, membrana tympani entirely gone. Left ear filled up with a polypus, covered with yellow pus, which flowed out over the edge of the meatus.

*Treatment.*—Ten grains of iodide of potassium three times



a day, with a wash of zinci sulphatis, ten grains to the ounce, to be dropped into left ear.

After cleansing with cotton, nitrate of silver was applied to the growths by means of Wilde's caustic-holder. Finding, after several trials during August, not much reduction in size, I twisted off portions at three different sittings. I applied powdered alum and then nitrate of silver to remove what was still at the bottom, but found it did not entirely destroy the spongy granulations, after eight months' persevering trial.

February, 1857. Applied a saturated solution of chloride of zinc on a piece of cotton, by means of the speculum, producing contraction of the granulations, so that I could see the membrana tympani, and an orifice from which the whole mass seemed to have sprouted. When I touched the surface of the mass, the patient felt a disposition to cough. His general health is much improved, he is now able to straighten himself, and the rheumatism is gone.

March, 1857. Dilated Eustachian tube by means of catheter and warm air at intervals of two weeks; his hearing was so much improved that he could hear me speak to him at a distance of six or seven feet.

Gradually the membrana tympani closed over, and he was discharged cured.

The case is one of great interest, and shows the good results to be obtained by persevering efforts, assisted by the willing help of the patient and the use of constitutional remedies.

"The truth probably is, that neither sex, age, nor general health of the individuals whose cases we have considered had much to do with the polypi in their ears. They sprang up in consequence of local causes, and grew because they found appropriate soil.

From Dr. Clarke's table, eight of thirteen growths appeared in the left ear, and five in the right ear. It is not fair to infer from this that the left ear is more likely to give birth to polypi than the right one.\*

---

\* Observations on the Nature and Treatment of Polypus in the Ear, by Ewd. H. Clarke, M.D., pp. 32, 33. Boston, 1867.



It would require a very large number of cases to justify any such conclusion. It is probable that they may occur indifferently in either ear. Such, I think, is the common opinion of observers. The inquiry as to which were the earliest symptoms is an important one. We learn from this same table\* that in one instance tinnitus aurium was among the earliest symptoms noticed; in six cases otalgia appeared early; in two or three cases tenderness and swelling were mentioned as among the first indications of difficulty. In all of them otorrhœa was referred to as appearing early; this was the history in all of our cases. Whatever else existed,—pain, tenderness, swelling, and noises,—a discharge from the ear was always one of the first warnings, and sometimes the first and only warning, of impending evil.

“These growths were attached to various parts of the external meatus; more of them to the middle than to any other portion. The auditory canal of the adult is about one and a quarter inches long. Suppose this, for convenience, were divided into three equal parts, called the outer, middle, and inner thirds. Following this subdivision we find that three of these growths were attached to the outer third, three to the inner third, and five to the middle third, and that the remaining two were not recorded, or else not made out. This does not bear out Mr. Wilde’s statement, that in eight cases out of twelve these polypi sprout from the site of the ceruminous glands in the posterior wall of the meatus.

“It is possible, however, that Mr. Wilde may be correct, and a larger number of observations than the above would show such to be the case. I have seen polypi growing from the membrana tympani, and from the cavity of the tympanum, and have extracted them from both places.

“An examination of the intimate structure of these growths shows that they belong to two distinct groups. They were epithelial or fibrous in their character. Those which belong to the fibrous group presented the following characteristics: They were not invested by a distinct membrane, but were covered with a sort of tessellated epithelium. Internally, their

---

\* Clarke’s.



color, when not reddened by infiltrated blood, was a pale pink, or almost of a grayish-white. Their general appearance was homogeneous. Under the microscope some of them exhibited fine parallel fibres. Others, a delicate fibrous structure, in which were numerous minute globules or granules; spindle-shaped or fusiform cells with free nuclei and smaller nucleoli abounded in all of them. Whatever else they contained the predominating elements were those of fibrous tissue in process of development, resembling what Lebert has figured under the head of fibro-plastic growth. Epithelial cells were often found in these specimens, but not in large numbers. The predominating elements were fibro-plastic. The individuals of this group were in reality small fibrous tumors. Their appropriate designation is that of fibrous polypi. They are evidently the same as those which Mr. Toynbee has described under the name of fibro-gelatinous polypi.

“By reference to the table or the cases themselves, we learn that in every instance the meatus was either seriously diseased or bore evidence of previous serious disease, when the patient applied for relief. In a single instance, the meatus was free from redness and tenderness; but in this case the membrana tympani was so largely perforated as to be nearly destroyed, and thus exhibited conclusive evidence of antecedent grave disease. In all the other cases, the meatus was more or less swollen, red, and irritated. In a few instances the tenderness was excessive. We have met with two cases of extensive perforation following the removal of polypus by other surgeons. None of the perforations healed. There remained seven perforations, and in those in which it was not perforated it remained more or less opaque. After extraction the otorrhœa ceased in all but two cases, and in five out of seven of the other cases.

“The next point is the result of treatment upon these growths.

“So far as the polypi were concerned, it was most satisfactory in all of the previous cases.

“In all of them the polypus was so completely removed and eradicated that it did not return. It is true that in some of them years were necessary in order to attain this result;



yet it was fully attained. This accords with my general experience. I do not recollect a single case that fell under my observation in which a similar satisfactory result has not been reached, except when perseverance of the patient has failed, and treatment been renounced, before a radical cure could be effected.

*"The results on the hearing.*—The effect on the hearing was not so satisfactory as it was on the polypi, if we turn to the record. Large fibroid polypi which fill the whole meatus and protrude externally, the membrane being perforated, appear to grow from nearly the whole surface of the tympanic cavity, the entire mucous lining of which seems to partake of the hypertrophy. Growths of this nature are extremely apt to recur, even after apparent cure, but the degree of obstinacy with which they resist treatment is very variable.

"We find that in seven of the previous thirteen cases the hearing distance, as tested by the watch, was the same at the conclusion of treatment as at its commencement. The hearing was not improved by removing the growth. In the remaining six cases the hearing was improved.

"The local treatment consisted of extraction, cauterization, and occasional counter-irritation, besides the use of astringents, syringing, and sometimes of leeches.

"The constitutional treatment consisted chiefly of hygienic measures and tonics, especially iron. Iodide of potassium was used several times. The duration of treatment was recorded in twelve of the thirteen cases. In one of them, where no treatment except extraction of the polypus was requisite, the duration is put down as one day. The longest course of treatment was four years; the average duration was six months.

"By duration of treatment I mean not only the time occupied in the extraction or destruction of the polypus, but also that occupied in leading the tissues into such a condition that there was no danger of its reproduction. It appears from the preceding observations that this cannot be done at once. The extraction of the growth is generally an easy matter; it is not so easy a matter to prevent its return. In a number of these cases the patience, ingenuity, and perse-



verance both of myself and of the patient were severely taxed before any satisfactory result was reached."

From the citations from Dr. Clarke's admirable pamphlet, it will be seen that he makes but two varieties or classes of polypi, viz., fibro-plastic and epithelial polypi.

Fibro-plastic polypi are thus described by the same author: "They have generally a smooth external surface. While in the ear they present a red or light-red appearance, but when removed from it they are a pale pink, and sometimes a grayish color, and firm to the touch, and are easily crushed between the thumb and finger. When touched or pressed in the meatus by a probe, they are slightly sensitive, and rarely bleed profusely. In size they vary from that of a pea to that of a body an inch or an inch and a quarter long and nearly half an inch in diameter. Their shape is as various as their size. Commonly, however, they are globular or pyriform. Growing from the fibrous tissues of the ear, they consequently may be found attached to any part of the outer walls of the external meatus or to the membrana tympani; they rarely spring from the cavity of the tympanum. The outer half of the meatus appears to generate them more readily than its more deeply-seated portion. They are apt to be attached by a pedicle or narrow neck to whatever part they grow from; yet they are occasionally seated with a broad base upon their parent soil. After extraction they collapse slightly. The surface from which they are removed never bleeds freely in consequence of the operation of removal, though some blood, of course, drains from it. Whatever pain accompanies the operation—and it is never severe—results mostly from adjusting the instrument in the meatus, and not from excising or extracting the tumor."

The epithelial polypus, or "raspberry-cellular polypus" of Toynbee, may also be found in any part of the meatus, and appear to be composed of a number of bright scarlet points or buds bound together by filaments, thus giving the appearance of a raspberry growing from the meatus. They are soft and easily crushed, and even when injured slightly bleed profusely. They are irregular in shape and vary greatly in size. They seem to be formed of the epithelial tissues of



the ear, and their frequent location is from the sides or bottom of a sinus, formed by ulceration in the wall of the meatus. "They may spring from the mucous membrane of the cavity of the tympanum, from the mastoid cells, or from the epidermal lamina of the membrana tympani. The formation of these polypi is attended with more or less pain and more or less discharge of blood, and often throwing off an offensive secretion."

"The indications for treatment are twofold: first, to remove or destroy the growth; and secondly, to restore the tissues to a healthy state as soon as possible. The first may be accomplished by retraction or cauterization, or by both of these measures combined, and sometimes by injection into the substance of the tumor. The second indication is answered by counter-irritation, the use of astringents, syringing, leeches, and such constitutional measures as may be required. Each of these points deserves a separate consideration.

"1. *Extraction*.—The ease and thoroughness with which a polypus can be removed from the ear depends as much upon the character of the growth as upon the instrument employed for its retraction. If the growth is soft, bleeding easily when touched, and adherent by a large base, it is not an easy matter to extract the whole at once. It will break up in the grasp of the instrument, and only come away in pieces; if it is not firm, or can be rendered so by the application of astringents to it, and is not attached by a broad base, it can easily and completely be removed by a single operation. As a general rule, fibrous are firmer than epithelial, and are consequently more easily retracted.\*

With very few exceptions, Mr. Hinton believes, it is the best plan to remove polypi by astringents; and I have tried the effect of astringent powders, such as alum and tannin, upon their roots after removal, and have not found them, on the whole, by any means equally satisfactory.

One further remark seems called for on this subject, says Mr. Hinton in his supplement to Mr. Toynbee's work. At page 324 the latter lays it down generally as a rule that

---

\* Clarke, op. cit., p. 451.




polypi should not be interfered with when disease of the brain is threatened, from irritation within or in the neighborhood of the tympanum. This is a rule which needs reconsideration, and, perhaps, even so accurate and thoughtful an observer has been in this particular unduly biased by the unfortunate issue of a single case. When cerebral irritation coexists with polypoid growths of any considerable size occupying the meatus, the removal of such growths should rather be a point engaging the earliest attention of the surgeon, for they are often themselves a chief source of irritation of the nervous centre by hindering the escape of matter. After their removal in such cases, indeed, it is no uncommon thing for the syringe to bring away large semi-solid masses of old discharge, the removal of which is followed by a great mitigation of the symptoms. In truth, the removal of polypi in these cases is only the carrying out of Mr. Toynbee's own principle of securing a free outlet for the discharge, a principle which should be applied without exception."

The instruments for the operation of extraction are various, and must be chosen according to the indications of the case.

In certain cases the diminutive "ligature canula" of Leveret, with silver wire, string, or whipcord, will answer every indication.

The best instrument, and the one which I have always found adapted for the purpose, is Wilde's snare (Fig. 48). It consists of a fine steel stem five inches long, and bent

Fig. 48.



in the centre (so as not to interfere with the direct rays of the sun), with a movable bar sliding on the square portion near the handle, which latter part fits over the thumb. The upper extremity is perforated with holes running parallel with the stem, and loops at the angle serve the same purpose. A fine




wire, fastened to the cross-bar, passes through these loops and holes. It should be of such a length that when the cross-bar is drawn tight to the handle of the ligature, it is fully on the stretch. The best form of wire is sheet-wire of No. 24 grist, with the temper taken out of it by heat.

Mr. Hinton remarks that Wilde's snare may be used of a larger size than that described by him, and is best armed, not with a wire, but with what is known to anglers as "jack-line," because it is more flexible, less brittle: it is easy to apply, gives no pain, and is not apt to fail.

There are several modifications of this instrument, most of them making it straight, but I think that many of them are useless, the original being the most convenient.

For very small vascular growths, the ring-and-lever forceps


Fig. 49.



of Mr. Toynbee is an admirable instrument. (See Fig. 49.)

In some cases the ordinary ring forceps will be found to

Fig. 50.



be suitable; and for others a form of instrument, Fig. 50, was found very useful in removing deep-seated polypi.\*


\* Gazette Hebdomadaire de Médecine et de Chirurgie, Nov. 15, 1867.



At Figs. 51 and 52 are seen two forms of forceps for removing polypi and foreign bodies from the ear.


Fig. 51.

Fig. 52.



The "ear-scissors," Fig. 53, are found useful in many cases, for removal of hairs, detached polypi, etc.

Fig. 53.



Whatever instrument may be chosen for the case, the growth should be secured as near as possible to its base, and then, by a firm grasp and gentle twisting motion, it is to be removed.

#### CAUTERIZATION.

Extraction of polypus is by no means the most important part of the treatment. In most cases it will sprout and grow from its root, unless efficient measures are taken to prevent this. For this purpose caustics in various forms are generally used.

The caustic potassa is considered the best, as it can be used with the greatest safety.

The following plan of Dr. Clarke is recommended: Pack a layer of cotton, previously moistened with acetic acid, closely along the wall of the meatus, and, if possible, a



little beyond the root of the polypus. Melt a piece of the potassa in a test-tube over a spirit- or gas-lamp. Dip the end of a probe—one whose end has been a little roughened is preferable—into the melted caustic. Do this once or twice, until the end of the probe is coated with it. As soon as enough has been collected on the end of the probe, pass it into the meatus and apply it to the growth.

A single application will sometimes be enough to destroy it completely.

The author has found equal parts of caustic soda and lime, moistened with absolute alcohol, a valuable remedy in vascular forms of growths; also tannic acid so moistened and applied.


Nitrate of mercury is much more convenient, but by no means so efficient.

The galvano-cautery, produced by the passage of a galvanic current through a fine platinum wire, was first devised and employed by Middeldorf, of Breslau.

Dr. Mackenzie has made a modification which he recommends for removing large growths without hemorrhage and less pain than the knife. A battery of great intensity is required in connection, and it is necessary to be able to control the current so that the wire shall not be heated until in contact with the part to be burned.

The instrument we would recommend is depicted in Fig.

Fig. 54.



54, and was modeled for Dr. Cohen upon the plan of Mackenzie's laryngeal electrode.

Two copper wires, to be connected with the wires of the battery, pass through an ebony handle to the top of that handle, where they terminate beneath a metallic spring, which,



when pressed down, forms a connection with two other copper wires passing out of the anterior portion of the handle, and which at their terminal extremity are connected by a short piece of thin platinum wire. The copper tubes are isolated by india-rubber jackets; and by twisting these wires the terminal extremity may be turned so that the cauterizing surface shall be present in any desired direction. The polyp to be burned having been reached, the loop of wire is made to encircle it; pressure upon the spring completes the circuit, and in less than two seconds the platinum wire is at a white heat. It requires a battery of great intensity. Dr. Cohen used six cells of the Maynooth battery which answered the purpose admirably.\*

#### INJECTION INTO THE SUBSTANCE OF THE TUMOR.

A solution of the perchloride or persulphate of iron may sometimes be injected into the interior of the tumor with the happiest results, as stated by Dr. Clarke, who has treated one case with success. Two or three drops of either of the above-mentioned substances may be injected into the growth by means of a syringe, the hypodermic being the best to use. The result is sometimes the immediate destruction of the polypus.

#### ASTRINGENTS AND STYPTICS.

The condition of the growth, or the adjacent parts, sometimes renders a certain amount of preparation necessary. In these cases the appearance of the parts must guide us in our selection of measures to be adopted.

If they are not too much inflamed, astringents and styptics will be useful. Astringent solutions of various strengths adapted to the case, poured into the ear, while the patient leans the head down with the affected ear uppermost, will be found more serviceable than the same solutions applied

---

\* Mackenzie on the Use of the Laryngoscope, with additions by Dr. J. Solis Cohen, p. 170.



by means of the syringe. Sulphate of zinc, tannin, alum, creasote, or carbolic acid, and a weak solution of iodine will be found to be some of the most serviceable remedies in these cases." The use of acetate of lead is recommended by Dr. Clarke. In some instances the substances named may be applied in powder to the part, especially in those cases where there is no perforation of the membrana tympani. When powders are used, they should be rendered, so far as practicable, impalpable.

They may be blown into the ear through a quill or through an appropriate instrument made for the purpose.


In the use of astringents it should always be borne in mind that only in cases of chronic inflammation strong solutions or their substitute in powder will be tolerated.

In cases of polypus with otorrhœa, cleanliness is of the greatest importance. Without careful measures to keep the parts thoroughly clean all treatment will be unavailing. Careful syringing with a good instrument will accomplish this object. Frequently the patient is disgusted with an imperfect instrument and a bungling hand working it. Dr. Thudichum's nasal douche (Fig. 55) will answer every purpose if fitted with an appropriate nozzle.

Fig. 55.



Fig. 56.



Another very convenient form of douche, applicable to the same purpose, is seen in Fig. 56, used with an ordinary pitcher on a shelf.

The discharge is best treated, according to Mr. Hinton, by absorbent powders, like "tale," removing the same by

---

\* Clarke, op. cit.



syringing, and sending a current of air by Politzer's method, so as to clear the Eustachian tube; more especially where there is perforation of the membrana tympani. In treating these cases, intercurrent inflammation, attendant upon the presence of the foreign substance in the meatus, must be checked by local and constitutional means. Leeches may sometimes be required, and if the object is to abstract much blood, they should be placed in front of the auricle. In young children it is safer to place them upon the mastoid process. Counter-irritation may be required where there is otalgia of a persistent character, attended with slight swelling and redness. In such cases a blister will be of great service. For this purpose the cantharidated collodion is best to be used, but croton oil and tincture of iodine may also be employed.

“Constitutional measures must be adapted to each case. Polypus is often associated with the strumous diathesis, and it is not easily eradicated, unless a change in the constitutional habit is produced. The weapons of defense of the physician against this protean form of disease are found among the tonics and alteratives and need not be described here.”\*

---

\* Hinton, op. cit.



## CHAPTER IX.

### DISEASES AND INJURIES OF THE AURICLE AND EXTERNAL MEATUS.

WOUNDS and injuries of the auricle are seldom seen except as the result of accident.

Ear-croppings and piercings for punishment have been supplanted by the more modern methods of boxing and pulling.

In Germany, however, the dueling with swords, one of the pleasures of "student-life," gives frequent injuries to the auricle, and these are usually incised wounds.

Wounds of the auricle may be either incised, punctured, lacerated, or contused, and treatment must be adapted to each case. The great object to be kept in view is always to prevent deformity if possible, and until this is observed it is never realized how much the ear contributes to the beauty and adornment of the head.

In cases of incised wounds, the parts should be brought accurately in apposition by means of an interrupted suture, the fine lead wire being preferable. In extensive injuries, where tissues are completely divided, sutures should be applied on both sides, care being taken to pass them through the integument only. Adhesive strips then complete the dressing. To protect the ear from the atmosphere, lint should be nicely placed around the part; this may be retained by a T-bandage, so modified as to suit the case.

In cases where a bandage is not at hand, the auriculo-occipital triangle of Mayor will serve as a useful dressing; a moderate-sized handkerchief being all that is required.

In cases where greater accuracy of adjustment is required, collodion will be found an admirable adjuvant in the dressings, especially if the prejudice against sutures in this part is entertained by the practitioner.



In wounds of the ear, if the parts be almost severed, and the attachment ever so small, we should not abandon the hope of recovering vitality and establishing union. Great attention should be given to the temperature, always endeavoring to keep it near the normal standard by artificial means.

In case the lobe has been destroyed or cut away, the loss may sometimes be supplied by a plastic operation.

The cartilage of the ear may be broken by crumpling up the auricle forcibly and suddenly; in these cases there is more or less inflammatory action, accompanied with pain and subsequent deformity. This curiously-formed organ is covered by a sensitive and delicate skin, and the hot flush and the burning blush are often seen to glow upon the ear, when at the same time a placid face and quiet brow bury all traces of emotion. It is said that even mental impressions affect the ear. Tradition has it, "That when the left ear is red, somebody is speaking well of you; when the right ear, the contrary." The auricle has a wonderful power for reparation after injuries, and one will often be surprised to see how little deformity follows serious injuries of the ear. The operation of piercing the lobes, which is fashionable in common with the savage and enlightened, for reasons alike satisfactory to both, is sometimes followed by grave consequences.

The operation is generally performed by thrusting an awl or needle of proper size through the lobe, which is pressed against a piece of cork held firmly against its back.

Sometimes inflammation of a grave character attacks the auricle in consequence of severe and protracted disease, which causes the patient to repose constantly on one side, and by the continued pressure of even the softest pillow the ear becomes ulcerated, and this sometimes ends in sloughing of the part. This takes place more readily because of the languid state of the circulation and the inability of the capillaries to resist injury, and in phthisis this injury frequently results from the patient being compelled to lie constantly on one side. Other causes of inflammation are the stings of insects, etc. These forms of inflammation are easily managed,



and are to be treated on general principles. One of the most troublesome forms is that induced by cold. In this climate the chilblain of the ear is by no means uncommon; and trifling as it may seem to have the "ears frozen," the subsequent irritation, and even decided injury that may follow, are not unworthy of consideration. Such cases should be treated by restoring the normal circulation by the application of snow or cold water, and the inflammation that follows, by gentle stimulation; one of the best applications is as follows:\*

R.—Nitric Acid, 1 minim;  
Tinct. Opium, 1 drachm;  
Water, 1 ounce. M.  
Sig. Apply externally, once daily.

Phlegmonous inflammation not unfrequently invades the auricle. This runs the usual course, often terminating in abscess, and must be treated on the general principles of surgery. Furuncle of the auricle is sometimes seen, and this must also be treated on general principles.

Gouty inflammation of the auricle, Wilde declares, has been fully established. The same author gives Graves the credit of having pointed out this form of congestion and redness of the auricle in patients of a gouty diathesis. Garrod has called attention to the frequent occurrence of uric acid depositions in the auricle in cases of individuals with an arthritic diathesis. These arthritic deposits were the most frequent of all the external symptoms, and they are therefore of some importance in the diagnosis of gout. Among seventeen patients with gout in whom the deposits of urate of soda were found, Garrod found them on the ear nine times, and at the same time in the vicinity of the joints,—seven times on the auricle alone; and of all the cases where they existed in the joints there was only one in which they could not also be found in one of the ears.

They are most frequently seen on the upper portion of the helix.

---

\* Sissay on Diseases of the Ear, translated by N. R. Smith, M.D. Baltimore, 1829.



They have not been observed as yet on the lower third of the auricle. Their size varies from that of the head of a pin to that of a pea.

TUMORS, ABSCESSSES, HÆMATOMA, AND SCIRRHUS OF THE  
AURICLE.

We have in the auricle tumors of various kinds,—encysted, steatomatous, sebaceous, fibrous, and bloody. The encysted, steatomatous, and sebaceous are generally small, and not frequent, while the fibrous of the lobules of the ear are more common and are sometimes of considerable size. A case of hypertrophy of the lobe is related by Boyer, in which it reached to the neck of the patient. Wilde\* was the first to accurately describe and figure one of these morbid growths, and determined that it was of a “yellowish-white fibrous appearance.” Mr. Paget† also refers to them as one set of fibrous tumors. “These,” he observes, “are trivial things in comparison with the tumors of the jaws, yet they have points of interest, in that they grow after injuries and are very apt to recur after removal. He also reports cases from Mr. Holberton, presented to the museum of St. Bartholomew’s Hospital, in which both lobules were cut off with the tumors, but in or beneath one of the cicatrices a similar tumor formed shortly afterward. This was excised; and in the ten years that have since elapsed there has been no return of the disease. In another case, related by Mr. B. Barrow, two such tumors formed in the same ear after puncture. One of these was cut away, the other was left; a third grew, and the excision of the whole lobule was necessary for the complete extirpation of the disease.” Prof. Alfred Stillé‡ reports an interesting case in a young lady; and describes the tumor when cut “to be smooth and shining, of a pale-pinkish color, and fibrous bands were visible, diverging from its point of attachment to the circumference. The

---

\* Practical Observations on Aural Surgery. June 12, 1853, Dublin; Am. ed., Philadelphia, 1853.


† Lectures on Surgical Pathology, 1847–1852, London; Am. ed., Philadelphia, 1860.

‡ Proceedings of the Pathological Society of Philadelphia, January, 1861.



mass of the tumor was formed by an opaque, elastic substance of an albuminous appearance." Similar instances of tumors of the lobe are recorded by Bruch, Venzetta, Hinton, Gross, Norris, and others. The following is one of a number of cases which occurred in the author's practice: About one year ago (1869) a young mulatto girl, aged eighteen years, bored her ears with an ordinary needle and inserted in them a silk thread. After the ear ceased to swell and the inflammation had passed away, she removed the silk and placed in each ear rings of impure gold. Soon after, the posterior surface of each orifice began to swell and gave her pain. The rings were removed, and not worn again. The swelling gradually increased, but more in the left than in the right ear, and within a few months have grown so rapidly as to become a deformity which she wished removed. The tumors were round, and measured on the left about three-quarters of an inch, on the right half an inch, through their long diameters. On the anterior surface, where the orifice had been, there was a small elevation. Both tumors were covered with the natural skin, and were firm, yet elastic, to the touch, and without pain. The largest of these tumors was removed by carefully dissecting back a portion of the skin covering it, and then cutting into the lobe in the form of a double-convex lens, and the skin was then brought over the wound. A single stitch sufficed. A small space was left to granulate. Subsequently the granulations were touched with nitrate of silver and dressed by a simple ointment and bandage, with isinglass plaster. The parts healed, and the operation produced no deformity. April (1871) the case reported above returned to have the tumor in the opposite ear removed; the one operated on did not return. On microscopical examination the tumor was found to be fibrous, with some fat cells.

Fig. 57.





The figure (57) will show the form of the tumor; although it was of a much larger size, involving the entire lobe. It also shows two rings upon it, as if compressed by a string. This specimen is the largest of a pair, one in each ear of a mulatto girl. Mr. Paget states that "these tumors are penalties attached to the barbarism of ear-rings;" and yet how many thousand wear ear-rings without ever suffering in the smallest degree from such tumors! I am of the opinion that they are owing to some peculiar condition of the system (scrofulous) chiefly, or some galvanic action in the mixed metals employed in the manufacture of the jewelry which is sold for pure gold. I am supported in this opinion by the numbers of such that occurred, not in the true negro but in the mixed blood of the white and black, which are more liable to scrofulous diseases than the true negro. "If," observes Mr. Wilde, "tumors of this description were frequently to follow the irritation of piercing the ears, we should find them much more common, particularly among those nations who wear very large, heavy pendulous ornaments, which is not the case. Dr. Graves has published a case of fatty deposit in the lobes; the patient died subsequently, and Dr. O'Ferrall found upon dissection a fatty degeneration of the liver and fatty deposits in other portions of the body."\* We have also had cases in the pure "white" skin, where the ear-rings were so large as to have torn the lobe in two; also cases where little children had lacerated them; yet they united without deformity, and no tumors followed.

#### HÆMATOMA AURIS.

For many years it has been observed in asylums that the insane are prone to suffer from swellings and indurations of the ear. By some these swellings have been attributed to mechanical injuries, whilst others have urged that this explanation is not applicable in many cases, for the frequent symmetry of the diseased condition makes it important in most cases to determine some other cause.

M. Ferrus, in France, and Bird, in Germany, first called

---

\* Wilde, *op. cit.*, p. 168.



attention to these sanguineous tumors in 1838. Dr. E. R. Hun, of the New York State Lunatic Asylum, reports (*Am. Jour. Insanity*, 1871) twenty-four cases of this affection occurring twenty-three times in man and only once in woman.\*

“Dr. Laycock, from a wider investigation of the symptomatology of the ear, arrives at the following (among other) conclusions: ‘That the states of the circulation, nutrition, and development of the tissues which make up the ear-lobule, and cover the helix, very commonly coincide with similar conditions of the encephalic tissues.’† If this be substantiated, if there be indeed a correlation between the nutrition of the brain and that of the ear, we have an explanation of the occurrence of hæmatoma in insanity. Be this as it may, however, the following example of the symmetrical formation of hæmatoma in a person who had certainly received no injury, is of much interest in reference to her history.

“Mrs. G.,‡ aged twenty-six, the wife of a sailor, married four years, but never had any family. She came to me on March 7th, with a large swelling on the upper part of the right ear. The swelling was at least the size of a walnut, tense, and fluctuating. It has been opened several times during the three or four months which it has been forming, and nothing but blood had ever been obtained. The surgeon on one occasion had put a seton through it. There had been great soreness in it, and the skin and cellular tissue near to the ear were considerably swollen from extension of the irritation. I opened the swelling freely, fully expecting to obtain pus, but none came, only a thin, bloody fluid.

“I kept the opening free from day to day, and the inflammation slowly subsided.

“On March 25th she mentioned that she feared her other ear was about to be affected in a similar way. I found a thickening of the cartilage, with decided fluctuation. There was no redness, but it was very tender on pressure. The

---

\* *The Medical Record*, New York, January 2, 1871, p. 490.

† *Medical Times and Gazette*, March, 1862, p. 289.

‡ Case of hæmatoma auris, under the care of Mr. Hutchinson, *Medical Times and Gazette*, December, 1862, p. 603.



fluctuating part was ill defined; the skin appeared thinner on the inner side of the cartilage. Exactly the same part was affected as in the other ear. I had suspected from the first that the case was similar to the hæmatomata of the ear met with in mania, and this symmetrical occurrence tended to support the idea. On inquiry I learned that a paternal uncle had been insane. Mrs. G. herself was of heavy aspect, with a small forehead and large face. Her ears were large in the upper part and wanting as to the development of the normal curves. Her husband was at sea. She told me that she had of late been very sleepless at nights and somewhat depressed in spirits, with a feeling that something was hanging over her; felt often cold. She had lately lost flesh and become paler, but was still in good general health. Catamenia irregular during last three months. Appetite rather poor."

The above case is not very conclusive, and having seen but very few cases of this disease I wrote to some of the highest authorities, chiefs of our insane institutions, and the following is an abstract of their replies.

Dr. Thomas Kirkbride, the distinguished physician-in-chief of the Pennsylvania Hospital for the Insane at Philadelphia, in a private letter to me on this subject, says: "The swelling has seemed to me the result of an injury to the part, and the best treatment that I have tried is the application of ice to the tumor at the very commencement. This seems to prevent the enlargement which otherwise is pretty sure to occur. I have seen great deformity caused by it. I have never satisfied myself that it occurs without any contusion."

Dr. John Curwin, physician-in-chief of the Pennsylvania State Lunatic Hospital at Harrisburg, thus writes:

"We have no cases of 'hæmatoma auris' at this time in the hospital, and have not had for some time past. My own impression about these affections is that they are the result of a blow or fall upon the ear, and are only developed when the nutrition generally is below par. If Dr. Laycock's theory be true, they would be more frequent than they are. I have not seen a large number of them,—say twenty in eighteen years, not including those I saw at the Pennsylvania Hos-



pital for the Insane at Philadelphia as assistant physician,—and as a rule they are almost invariably confined to males; some say altogether, but I have seen one or two cases in females. I have tried all kinds of means for their removal, but have never found anything equal to the application of tincture of iodine once or twice a day, as the case may demand. This reduces the tumor with less disfiguring than any other means; but the ear rarely resumes the same condition and appearance it had before, a slight degree of thickening always remaining.”

No apology, I trust, is necessary in giving the letter of the chief physician of the “Friends’ Asylum for the Insane at Frankford, Pennsylvania,” as he enters very fully into the subject, and he has had a large number of cases under charge:

“RESPECTED FRIEND,—Thy note of yesterday is at hand. In reply, I may say that I have had a pretty large number of the cases inquired after under observation, and will with pleasure give thee briefly the result of my experience.

“I have never seen a case of the affection in a sane person, nor in a case of curable insanity. I always consider the affection of the *ear* an evidence of an incurable form of insanity, such as I have described in the Journal of Insanity as ‘congestive mania.’ I believe it is always associated with a tendency to congestion of the membranes of the brain or the cerebral substance itself; and I should think, in a general way, that Dr. Laycock’s opinion was correct. The affection seems to be in the perichondrium, which secretes a bloody serum, with which the cellular tissue becomes infiltrated. The best treatment would probably be a lotion of liquor plumbi et tinctura opii, but owing to the difficulty in making any application to the class of cases affected, which are generally unmanageable, I have tried the method which has uniformly proved satisfactory, the swelling disappearing in a few weeks and leaving only a slight pucker under the skin, such as you might suppose a cicatrix in the perichondrium would present. The practice of opening the tumor is altogether unnecessary, and therefore I think un-



scientific. The character and appearance of the lesion would suggest mechanical injury; but as these tumors are never seen in other classes of the insane equally turbulent, or in epileptics, who are especially liable to blows from falling in a fit, there is probably something specific in the cause and character of the affection; and in the cases I have observed I never knew one to be the result of an injury by a blow or fall.

“Thine very truly,

“J. H. WORTHINGTON.

“LAURENCE TURNBULL, M.D.”

Dr. William Kramer, of Berlin, treats of “inflammation of the perichondrium\* as presenting essentially different characters, according to the general state of the patient’s health. In strong constitutions a bright-red swelling; in the debilitated individual, on the other hand, it is bluish-red. The cause of these bloody tumors are unknown, though he must admit that they are especially likely to be produced by violence (blows on the ear), which perhaps explains their frequent occurrence on the left ear. If the inflammation have developed itself in a strong constitution, with synocha, we must recommend rest, cooling diet, mild aperients, and locally cooling and dispersive applications of lead lotion. The necessity of surgical interference with the tumor must be determined by its size and presence or absence of fluctuation.

“Cancerous and nodulated degeneration of the auricle does not often admit of a return to a healthy state. The course of this inflammation is extraordinarily protracted, lasting for months, years, or even through the whole life of the patient; and it may almost always be considered as connected with a more or less strongly-marked scrofulous diathesis. The treatment must be essentially constitutional, and directed to the general condition of the patient. Fat, acids, farinaceous food, and, in general, all

---

\* The Aural Surgery of the Present Day, by Dr. William Kramer. London, 1863.



food difficult of digestion, should be excluded from the diet of these cachectic patients. They require strong, nourishing aliment, both fluid and solid. It is indispensable that the bowels should be kept freely open, but in cachectic individuals this should not be allowed to pass into diarrhœa. Either artificially made baths, or those especially of Kreuznach, Nauheim, and similar places should be taken; and at the same time the Adelaide and Eliza Springs, whey, iodide of potassium, etc. If these means should fail, we may employ, if the patient's strength will permit, Fowler's solution of arsenic, in doses of from two to four drops two or three times a day, a quarter of an hour after meals. The local treatment consists in carefully removing and drying up the diseased secretion by means of a soft sponge dipped in warm water, and subsequently powdering the surface with the semen lycopodii. If the itching and burning be very severe, soft lint dipped in a warm solution of zinc (gr. ii-iv, ad. 3j water) should be placed upon the ear. Thick, hard, purulent scabs are softened, with great relief to the patient, by the application of warm bread or linseed meal poultices. Where all these means fail, relief to the patient can be obtained by the partial or entire removal of the auricle."\*

## CANCER OF THE AURICLE.

True cancer of the auricle must be a very rare disease, for in my hospital and private practice I do not remember seeing more than a single case. This is also the experience of Wilde. Yet I have met with many cases of abscesses and irritable ulcers on the auricle. This diseased condition is to be treated by alteratives and tonics, with the local application of solutions of zinci sulphas, acid. carbolic, or permanganate of potassa. When the elevated edges or nodules require removal, we resort to the sulphate of zinc paste, chloride of zinc, or the compound powder of caustic soda and lime, mixed into a paste with absolute alcohol. In some

---

\* Kramer, op. cit., p. 39.



cases the solution of the protonitrate of mercury may be used.

#### DISEASES OF THE MASTOID AND PRE-AURICULAR REGION.

Wilde,\* "without entering too minutely into the subject of tumors behind the ear," specifies six varieties. "The first is a small gland lying upon the mastoid process, immediately above the insertion of the sterno-cleido-mastoid muscle, and on a level with the auditory canal of the ear, which sometimes enlarges to the size of an almond. It becomes highly irritable and painful to the touch, so as occasionally to resemble a neuroma. It generally occurs in young females." We have seen a number of these tumors, and have found the application of a solution of camphor in an equal weight of chloroform highly beneficial in relieving pain; we also used the solution of iodine in aqua ammonia locally at night. Internally we administered the iodide of iron. If the tumor be large, or if there are several, we have applied, with the happiest effects, the mercurial plaster, with as much camphor as can be incorporated with it and yet not prevent it from adhering. The three varieties next described are not true tumors, but suppurating glands or chronic abscesses, with or without periostitis of the mastoid process. The fifth is a true aneurism of the posterior auricular artery. The sixth tumor described by Wilde as occurring in this region is a malignant fungus, of which he has seen three cases—one in a boy aged ten, the others in adults who had passed the age of fifty. We have only had two such cases to treat. One was in a delicate child, and involved the whole of the mastoid process and side of the neck. The case was brought to the Howard Hospital, but very little could be done for it. The second was a woman aged fifty-six. Her statement was that it began as a tumor behind the ear, and that she had consulted a surgeon, who pronounced it an abscess and cut it open. It discharged some blood, and soon after the operation it began to enlarge, the edges of the ulcer becoming everted. When we examined it the mass of fungus

---

\* The Nature and Treatment of Diseases of the Ear, p. 175.



had penetrated into the meatus through the auricle, and was disposed to bleed on the slightest touch, and there was an ichorous fluid discharging constantly. She was placed upon the use of the following mixture :

R.—Tr. Cinchonæ Comp. fʒj;  
Liq. Potassæ Arsenit. fʒij;  
Syr. Simplex,  
Aquæ, aa fʒij. M.  
Sig. A teaspoonful after each meal.

She also used alternately solutions of sulphate of zinc and carbolic acid and the solution of persulphate of iron as a wash, so as to remove the disagreeable odor and prevent hemorrhage. She continued to attend as an out-door patient for several months, without much increase in the size of the growth.

In the true epithelioma excision or amputation is the treatment in the early stage.

#### MALIGNANT DISEASE OF THE AURICLE.

The following case of malignant disease of the auricle is reported by Dr. Green\* (with remarks), and is worthy of record on account of the success which followed the operation by Dr. Thaxter.

“J. M., aged sixty, night-watchman, entered Dr. Thaxter's wards at the Boston City Hospital on April 21st, 1870. He stated that some eight months before, without having ever had any previous trouble in the ears, an eruption appeared on the right auricle, from his description apparently a moist eczema. From that the present condition has gradually developed itself without pain or symptoms of any kind whatever. No treatment had been adopted from the beginning.

“The lower half of the right auricle was found to be very much enlarged in all directions; the concha and upper part of the lobule were covered by a superficial ulceration, from which an offensive sero-purulent fluid exuded, forming a dark crust. The lobule was occupied by several nodules about the size of hazelnuts, of a purple color and very hard; between

---

\* Transactions of American Otological Society, p. 62.



these were several bleeding fissures. The neighboring ganglia were not affected.

"On examination with the speculum and Tröltsch's mirror, I found that the ulceration did not extend into the meatus, but that the whole inferior wall of the passage was swollen and indurated as far as the edge of the osseous meatus. The deeper parts of the passage were filled by a large mass of loose hairs, which were removed, and the membrana tympani was then seen to be opaque, somewhat swollen and congested along the manubrium. The hearing was considerably diminished on that side, the watch being heard only at the distance of two inches, and there were constant subjective noises.

"The appearance of the disease leaving no doubt as to its character, excision was performed by Dr. Thaxter under ether, the whole auricle being amputated close to the skull. The whole lower half of the cartilaginous meatus was then dissected out down to the edge of the osseous meatus, and found to be indurated. The remaining tissues appeared to be healthy. The bleeding was profuse, but easily checked by tying a few small vessels. Cold-water dressing was applied, and the wound left to granulate.

"The wound healed rapidly, and at the end of a fortnight was not more than one inch in circumference. The process of healing then became slower, and the granulations at the meatus threatening to close that passage, a pledget of lint was kept in it all the time. The granulations were occasionally touched with solid nitrate of silver, and stimulating lotions applied. At the end of four weeks the whole wound was cicatrized, except just at the edge of the meatus, the opening of which was now only about one-eighth of an inch in diameter. The hearing had improved, and the subjective noises were gone; the swelling and congestion of the membrana tympani had disappeared, leaving that membrane dull and somewhat thickened, but otherwise normal.

"The patient now demanded his discharge, promising to return as an out-patient, but no more was seen of him for four months, when, on meeting him, I learned that after leaving the hospital he neglected all dressings, and the ear



soon healed. The meatus I found to be closed by a firm layer of cicatricial tissue, and the hearing power was less than when last examined. There were no signs of any return of the disease, and he expressed himself as perfectly satisfied with the result of the operation.

"In looking up the literature of the subject I have been able to find but little. Neither the text-books on the ear nor on disease of the skin mention the auricle as a likely seat of epithelial disease. In the *Gazette des Hôpitaux*, September 30th, 1869, M. Demarquay describes a case in a robust man, seventy years old, who entered the *Maison Municipale de Santé*. He had lost one leg from disease of ankle, and had had caries of one phalanx; also a hemiplegia on the right side, from which he had nearly recovered. At first a small wart appeared on the middle of the left helix, was scratched off, but immediately returned; it was then cauterized with nitric acid, but gradually extended, and in seven months had formed a brown crust, with a hard, indurated base. Below this crust was an ulcerated, slightly-painful, bleeding surface, which extended in spite of caustics.

"On entrance, the tumor occupied the upper part of the auricle, was 6 cm. long by  $1\frac{1}{2}$  cm. wide, more dependent in the middle than at its extremities; the base was indurated and adherent to the cartilage, its borders hard, defined and red; the surface irregular, and ulcerated. On close questioning, he admitted shooting pains in the meatus, but none in the affected part. The characteristics of the tumor left no doubt of its nature, and it was proposed to excise all the affected parts with the scissors, but the patient refused to submit, and left the hospital.

"M. Demarquay remarks that in such cases all authors are far from partial to the knife. M. Sédillot considers the actual cautery sure of success, and in a patient of his, fifty-five years old, in whom the whole auricle had been destroyed by canceroid in less than three months, and the disease had invaded the auditory canal, the actual cautery applied to the ulceration and repeated many times produced a firm and persistent cicatrix, which he claims could have been obtained in no other way. Velpeau, on the contrary, resorted to excision in



the case of a patient, sixty-seven years old, where the growth had existed only two months. The tumor (size of a filbert) was firm, fleshy, ulcerated on the surface, and movable with the skin on the deeper parts; occasionally there were lancinating pains in it. When small it had been removed, but had recurred. As the neighboring ganglia were not affected, excision was considered practicable.

“Malignant tumors of the auricle offer the peculiarity of rapid development which often renders the ablation of the part indispensable; in this they differ from canceroid of the lip, which develops slowly. Velpeau had already pointed out this fact, and in the case reported by him (*Gazette des Hôpitaux*, 1864), it enabled him to decide on the malignancy of the tumor, and he diagnosed a cancerous tumor containing probably fibro-plastic tissue and the elements of encephaloid cancer. Dr. Campbell states that many of the inhabitants of Nissal have tumors of the auricle, especially those afflicted with goitre. Two of them which he removed occupied the whole auricle, invaded the meatus, and affected the hearing, and their tissue resembled mammary sarcoma.

“Brunner narrates in the *Archiv für Ohrenheilkunde*, vol. v., a case of epithelioma of the ear which he considers began primarily in that organ. For two years the patient, a healthy woman, aged fifty-two, suffered from severe itching in the right ear, and was in the habit of frequently scratching it with a hair-pin. A slight otorrhœa then began, without pain or other symptom. A few weeks after an exposure to cold she had severe pain in the ear, which soon diminished, but continued in slight degree. Two months after this the right facial nerve suddenly became paralyzed. Examination at this time showed the right meatus to be completely filled by a soft, reddish, spongy mass, exactly resembling granulations; the otorrhœa was slight, and more mucous than purulent; the auricle and surrounding tissues of ear normal. A microscopic examination of a small piece of the growth, by Professors Eberth and Billroth, showed an undoubted epithelial carcinoma. As the growth involved the labyrinth wall, no operation was attempted, and the treatment was symptomatic only. The pain increased, ulceration took



place, and the auricle was partially separated. One year after the beginning of the otorrhœa she died. No autopsy was made.

“The case which is reported is interesting, from the disease having extended from the auricle into the cartilaginous meatus and yet having left the deeper parts intact, and from the result of the amputation, which shows that in such cases the complete closure of the meatus must be specially guarded against.”

Gruber\* describes and figures a most formidable disease of the auricle under the term “Angioma,” the edge of the helix being filled with a mass resembling hard cancer, and in the centre is a bloody tumor with an opening resembling an othœtomata. The treatment consisted of the emplastrum diachylon, ʒij, with antimonii et potass. tart. grs. xvij, or the subcutaneous injection of the dilute liq. ferri sesquichlorate; also acupuncture. Where these remedies fail, he recommends the removal of part or the whole of the auricle.

---

\* Lehrbuch der Ohrenheilkunde, p. 409. Wien, 1870.



## CHAPTER X.

### PRURITUS—ACUTE AND CHRONIC ECZEMA—FURUNCULOUS ABSCESS—OSSEOUS TUMORS—OTITIS EXTERNA.

PRURITUS of external auditory canal is frequently observed. Dr. Gruber,\* in inviting attention to this affection, says it is either overlooked or treated of too summarily in treatises on diseases of the ear. Itching of the auditory canal is often met with during the progress, or at the termination, of inflammatory action. The affection indicated resembles in all respects the pruritus cutaneous of Hebra. Intense itching of the meatus constitutes the only objective symptom, the lining membrane of the passage exhibiting no trace whatever of any changed appearance. Gruber notices that in aged persons, suffering from this affection, the passage is often found very dry, and without a vestige of cerumen. That this is no essential feature of the disease is seen in other cases in which the secretion is found in excess. This "pruritus" is often met with in persons of middle age, and especially in those in whom there exists some disturbance of the circulation. Thus, those suffering from hemorrhoids or large varicose veins seem to be liable to it, and even children are not exempt. The itching often comes on periodically, that is, in the evening, the patient having been completely free during the day. Often it is absent for months, and returns again with distressing severity. In Gruber's cases the affection appears to have been strictly local, no other part of the body participating in the irritation. Sometimes, as the result of the constant scratching induced by this affection, an artificial eczema, or diffuse inflammation, is set up.

---

\* British and Foreign Medico-Chirurg. Review, April, 1869; from Allg. Wien. Med. Zeit., Dec. 1868.



The causes of pruritus of the ear are not well known. It is sometimes accompanied by an eruption of papulæ of nearly the same color as the adjoining cuticle. Wilde quotes from the *British and Foreign Medico-Chirurgical Review*, vol. xvii. p. 414. Pruritus of the external meatus auditorius, from hyperæsthesia of the auricular branch of the pneumogastric, is sometimes observed, and is accompanied by cough and vomiting. Sensibility of the terminal portions of the cutaneous nerves may induce distressing itching, which will torment the patient for hours without change of structure being induced by stimulants, changes of temperature, indigestion, etc. Mental excitement or depression seems to have a decided influence upon the disease. Sometimes the affection disappears suddenly, and subsequently returns as unexpectedly with unabated violence; this should always be remembered in our prognosis.

*Diagnosis.*—The character of the itching, which is tormenting, annoying, and periodical, without evidences of inflammation, or other change of structure of the skin, will usually guide to a correct diagnosis. The attacks come on suddenly, and continue for hours, and then as suddenly abate. The violence done during the attack, from the irresistible desire to scratch, increases the irritation.

*Treatment.*—As a palliative treatment, a few drops of glycerine may be used. Mere chronic and habitual sufferers should be taught to pencil the auditory passage with the substance just mentioned, or ointments, such as the citrine or oxide of zinc. And those whose night's rest is liable to be disturbed, should introduce lint imbued with such substances, and leave it in during the night. The ceratum plumbi subacetatis, with equal parts of ungt. aquæ rosarum, will be found of great service in many cases. A more radical means of cure is penciling the meatus with a strong solution of nitrate of silver, gr. x to fʒij of water. This should be done in a good light, so as to avoid the membrana tympani, and must not be continued after signs of reaction. This must be again repeated should the itching reappear. In all cases great attention must be given to the general health, and condition



of the digestive system. Acid products from indigestion seem to produce this affection in those liable to attacks. The same tendency is seen in females dependent upon disturbance of the menstrual function. In these cases special attention must be given to the disturbed organs.

Arsenic, in the form of Fowler's solution, will be found invaluable in cases that have defied all other treatment. Local applications of solution of carbolic acid, acetic acid, bichloride of mercury, or sulpho-carbolate of soda may be used with advantage, ever bearing in mind that judicious consideration of the character and temperament of the patient must be exercised in selection of the remedy. These lessons are learned only by experience, guided by principle, and, although theoretically empirical, form the basis of judicious practice.

#### ACUTE AND CHRONIC ECZEMA.

The acute form of this eruption commences with some disturbance of the general system, and shows itself locally upon the auricle by the discharge of a serous fluid. When the vesicles are rubbed, they peel off, leaving a slightly red surface. The eruption gradually extends over the face; there is itching and heat of skin, and when cloths are applied they become stiff with the secretion, which dries very rapidly. A clinical case may assist the knowledge of this affection.

Mrs. Eliza —, aged thirty-five, applied at the hospital, April, 1870, with acute eczema. She was nursing her infant, and had been attacked four or five weeks before applying for treatment. The eruption commenced behind her ear, then extended over ear and side of face, upper lip, and forehead. Even over her neck, there were numerous clusters of whole and broken vesicles, containing a serous, and some a purulent, matter. There were also scales, which, on removal, left a deep-red surface. There was itching and smarting of the surface, the auricle and meatus red, swollen, and tender. The parts were ordered to be kept moist with mucilage of slippery elm and flaxseed, in each pint of which was dissolved biborate



of soda  $\text{ʒi}$ . A mild anodyne was given to induce sleep, to be followed by a saline laxative in the morning. Believing the eruption to be dependent upon debility, she was advised not to nurse her baby, and was given an infusion of Peruvian bark. This treatment she continued for two weeks with but little benefit, when it was changed for the application of the benzoated oxide of zinc ointment, and five drops of Fowler's solution in the bitter infusion. A month's treatment caused the disappearance of the eruption and a return to her ordinary condition of health.

#### CHRONIC ECZEMA AURIUM.

This is especially a disease of females in middle life, affecting the ear, as a sequela to the cessation of the menstrual flux, and taking the place of the headaches so common during this period. It becomes very annoying from the itching and watery discharge, soiling the collar and neck with scales, and also filling up the auditory canal, often causing great pain.

*Diagnosis.*—In the acute or simple variety we find clusters of minute vesicles without much redness, and in the chronic form we have chaps and fissures pouring out an abundance of ichorous fluid, and at a later period copious desquamations. According to Wilson (and this we can confirm from our own experience), small subcutaneous abscesses form in the integument around the ears, and the neighboring lymphatic glands frequently enlarge; and in one instance under the author's care there occurred sympathetic irritation of the brain.

*Prognosis.*—If the case be seen early, and the constitution is good and strong, the prognosis is favorable; but if the case is of several years' duration, the results are not always as satisfactory as we could desire; still, much can be done to improve and modify, so as to make the patient's condition more comfortable.

*Treatment.*—In the case before referred to, of several years' duration, in which, during the treatment, the brain was very much disturbed, after trying a great variety of remedies, the patient finally was relieved by the use of the following solu-



tion, and ointment of the acetate of lead and mercurials, continued with care for several months :

R.—Bichloride of Mercury, ʒij ;  
Arsenious Acid, ʒss ;  
Hydrochloric Acid, fʒij ;  
Water, fʒviii.

Boil the Arsenious Acid and Hydrochloric Acid in water, and add the Bichloride of Mercury.\*

The dose, from 3 to 10 drops, in water, 3 times a day.

The following case from Wilde is so characteristic of many we see at our public hospitals and clinics, and being related in his inimitable style, we cannot do better than give it at this time :

“M. Q., a female, aged sixty, has been deaf, ‘off and on,’ for several years past, accompanied by noise and wandering pains in her head, with extreme itchiness in the auditory passages. The skin covering the auricle, and the scalp adjacent thereto, is of a fiery red color, speckled with patches of yellow, formed by the exudation which has collected in thin, branny scales all over it. The parts are hot, and in some places sticky, from a thin ichorous matter which exudes from the surface. The auricle has lost its natural shape, its folds and sinuosities being partially obliterated, and it has become hard, thickened, and lumpy. The external auditory aperture, owing to the disease extending into it, has been lessened to a third of its natural size, and is filled with branny scurf. Upon removing the latter impediment, we can obtain but a very partial view of the membrana tympani, which appears to be thickened and opaque. Hearing distance, touching. The state of the parts is nearly the same on both sides.

“Cases of this description, and like this, of long standing, are very hard to manage, because there generally coexists some constitutional taint;—so that while we employ local measures, we must not neglect constitutional means, and strict attention to diet should be enforced.”†

---

\* Pharmacopœia of the Hospital for Skin Diseases. Med. Times and Gazette, April, 1857, p. 336.

† Wilde, op. cit., p. 171.



The ointment before referred to was as follows :

R.—Acetate of Lead,  $\mathfrak{z}_{ss}$  ;  
 Oxide of Zinc,  $\mathfrak{z}_i$  ;  
 Mild Chloride of Mercury,  $\mathfrak{z}_{ss}$  ;  
 Nitrate of Mercury Ointment,  $\mathfrak{z}_i$  ;  
 Lard (washed),  $\mathfrak{z}_{ss}$  ;  
 Pure Palm Oil,  $\mathfrak{z}_{ss}$ . Mix.

With many cases this severe treatment is not absolutely necessary, and the mild mucilaginous applications—the ear being covered by oil-silk—have been attended with success. In another class of cases we resorted, with good results, to the ointment of nitrate of mercury, of double the ordinary strength, which was kindly prepared by Mr. Bullock, of the firm of Bullock & Crenshaw, of this city.

Hebra, quoted by Gruber, recommends the following ointment :

R.—Ol. Oliv. Optim. unc. 15 ;  
 Lithargyri unc. 3 et dr. 6 ;  
 Coq. l. a in ung. moll. ; dein adde  
 Ol. Lavand. drachm 2. M.  
 Ft. Ung.

The following powder\* is recommended by Ausspitz, who advises us to do as little as possible in the acute form. In this we differ from him, for, although we may not obtain a sudden cure, still the treatment is of great relief to the patient :

R.—Flor. Zinci,  $\mathfrak{z}_{ij}$  ;  
 Pulv. Alum.  
 Amyli Pulv.  $\bar{a}\bar{a}$   $\mathfrak{z}_j$ . M.  
 Ft. Pulv.

The powder is to be dusted over the affected parts with a camel's-hair brush.

#### FURUNCULOUS ABSCESS OF THE AUDITORY CANAL.

This is not a frequent affection, but varies with the several seasons, being more prevalent during mild, and consequently damp, winters.

Roosa, in a report of five hundred cases of aural disease, gives only eight of furunculous inflammation of the external auditory canal. In a still more recent report, of four hundred

---

\* Prof. Roosa, Medical Record, p. 79, April 15, 1871.



and ninety-four cases of diseases of the ear, there were fourteen cases of furuncular or circumscribed inflammation of the external auditory canal.\*

Schwartz found thirteen in five hundred and fifty-four cases of disease of the ear.† Pagenstecher,‡ seven in two hundred and twenty cases; and Gruber§ counted thirty-two out of seven hundred and seventy-five, observed in his clinic for diseases of the ear. These statistics from eminent authorities will furnish some aid in forming an estimate of the frequency of this painful affection.

Out of six hundred cases treated by the author, there were thirty-nine of furunculous abscesses. These varied markedly with the influence of our ever-changing temperature of climate, and in position in the course of the auditory canal.

By reference to the anatomy of the external ear, it will be readily noticed that the tissues in which this abscess occurs are tense and unyielding.

It is impossible at all times to distinguish precisely how deeply the inflammatory action has extended. The starting-point is usually one of the hair-follicles in the external portion of the auditory canal, or in one of the sebaceous glands; hence the various positions which they occupy in the canal.

*Etiology.*—The etiology of furunculous abscess is not well established.

Prof. Roosa states that in seven of the eight cases reported, the abscess occurred in persons who were not in good general health.

According to Von Tröltsch,|| furuncles of the auditory canal appear at any time in life, and in the most different constitutions. They often occur as complications of otorrhœa,

---

\* New York Med. Jour., No. 5, Aug. 1869.

† Deutsch Klinik, 1867, II. p. 72.

‡ Praktische Beiträge zur Ohrenheilkunde, von H. Schwartz. Wurtzburg, 1864.

§ Monatsschrift für Ohrenheilkunde, Berlin, No. 9, Sept., p. 128.

|| Von Tröltsch on the Ear. Translated by D. B. St. John Roosa, N. Y. 69, p. 97.



when the ear is very frequently syringed, and also when the affection is left entirely to itself. Hagen\* states that he has observed them in patients from four to sixty-two years of age, and that others have observed them in nursing children.

Rau† “thinks that this is not a frequent affection in children, and appears like aural catarrh, about the time of teething.”

Sex seems to have but little influence, although the weight of authority believes this abscess occurs more frequently among females than males. There seems to be no doubt of some predisposition to this affection among females who suffer from menstrual disturbances, and at the menopause, as well as in phthisical cases.

Wilde‡ states that “these abscesses occur oftener in females than in males, and I am induced to think they appear oftenest in persons in the higher classes than in the lower, and also that they seem to be induced by some peculiar state of the constitution, and very often appear either as a sequela to or a concomitant of furuncles on other parts of the body; particularly about the neck.”

There can be no doubt that irritants, injuries, cold, foreign bodies, collections of wax, constantly picking of the ears with pins, needles, toothpicks, etc. are frequently the exciting causes of these abscesses.

Von Tröltsch, Schwartze, and Von Hagen have observed these abscesses frequently after the use of alum washes for the ear for deeper complications, as catarrh of the membrana tympani, etc. Dr. Roosa is inclined to doubt the peculiar influence attributed to alum solution. The author has observed the tendency of astringents to produce them when used in too strong solution.

*Symptoms.*—The symptoms of this affection vary according to the position it assumes. As a furunculous abscess, having its seat either in a hair-follicle or ceruminous gland, it has in its centre a circumscribed “core,” which must be discharged

---

\* *Praktische Beiträge zur Ohrenheilkunde*, von Dr. R. Hagen, No. III. Leipzig, 1867.

† W. Rau, *Lehrbuch der Ohrenheilkunde*. Berlin, 1856.

‡ Wilde, *Diseases of the Ear*. Phil., 1853, p. 192.



before healing will be accomplished. The pain will be more or less modified by the tissues involved, so that the extent of inflammation will not serve as an index of pain. Cases occur in which the pain is most excruciating ; in others it is scarcely noticeable until the abscess is formed. This difference is fully accounted for by the peculiar formation of the lateral section of the meatus, it being composed partly of cartilaginous walls, and a part consists only of membranous tissue, having an osseous margin, while it is also contiguous to soft parts. Furthermore, on the upper wall, a gusset-shaped piece of integument extends to the osseous wall of the canal, and has just as dense connective tissue as any other part, as well as glands and hairs.

If, now, furuncles should occur in a position where inflamed swollen connective tissue cannot extend itself, and quickly reaches a firm, unyielding basis, viz., the bone, the symptoms depending upon tension of the connective tissue will be much severer. If we reverse the case, such follicular abscesses will be little noticed if situated at the entrance of the auditory canal, or in similar favorable localities. The pain in the beginning is often intense, and, in some instances, increases at night. In milder cases it is, however, scarcely more than a burning, itching, sticking pain, accompanied with fever, loss of appetite, loss of sleep, etc., showing how fully the whole system is affected. Often the acts of chewing or speaking give intense pain. In other cases severe pains radiate from the troublesome centre to adjacent parts, and make the slightest movement of the ear impossible on account of the intense pain. Patients thus affected often persist in describing their pains as neuralgic, and are often encouraged in this by their physicians, to such an extent that it is difficult to persuade them to allow the ear to be examined. So sensitive does the auditory canal become, that they are fearful lest the light from the mirror should cause additional pain. The slightest touch with a probe, on the point of the abscess, is exceedingly painful, and this manœuvre will often aid in diagnosis. They may be seated anywhere around the external meatus, but occur more frequently in the anterior and posterior wall than in the roof



or floor. They vary in size, with a broad base, and their color is but little altered from that of the skin, sometimes being slightly reddened, especially immediately over the swelling. This swelling is not always well defined, and the partial closing of the canal renders it difficult to demonstrate the actual position of the abscess; and when it is situated in the external portion of the canal, the cellular tissue of the external ear becomes at times involved, and the meatus becomes entirely closed.

Without doubt, there is more or less congestion of the membrana tympani in most cases, as is shown by the sense of buzzing and tinnitus.

The hearing in these cases is seldom much impaired, unless, perchance, the abscess be situated near the drum of the ear, or where the swelling materially diminishes the calibre of the external auditory canal.

*Course and Termination.*—Resolution may sometimes take place without discharge of pus, says Von Trötsch. Generally, if left undisturbed, the formation of the abscess progresses tediously and painfully, lasting from two to eight days. The symptoms of pain increase in direct ratio to the duration, until, at last, an abscess is formed. A peculiar exudation at first covers the surface of the abscess, then the tissues give way, and only a few drops of pus with a little blood is discharged. The tormenting pain ceases, and the patient feels a strange sense of relief. Finally, the "core" is discharged, and the opening heals unless another one forms, following rapidly in the steps of its painful predecessor; and frequently the second abscess is forming whilst the first is discharging.

Diagnosis is not difficult if the symptoms already mentioned exist, and yet mistakes occur. In cases of complaint of pain in region of the ear, it is always best to examine the whole canal with direct or reflected light, and explore its whole length by gentle touches of a silver probe. Usually some point of excessive tenderness will be found, thus locating the seat of the abscess.

To illustrate the errors that may arise in cases of this kind, a case from the practice of an acquaintance will serve:



June 20, 1869, Miss B., aged nineteen, applied for advice, and gave the following account: Two days ago, feeling a tingling and itching about the ear, she used a pin to scratch the ear. In doing so she lost her hold of the pin, and she thought it had fallen into the ear; but as it gave her no annoyance it was forgotten. Three or four hours elapsed, and her attention was again arrested by a burning and tingling sensation in her ear, which soon became painful. Remembering the pin, her parents were consulted, and by a light from a lamp it was concluded that the pin could be seen, and various attempts were made to dislodge it by the aid of needles, scoops, etc. The pain became intolerable, and, full of alarm and anxiety, a physician was summoned, who confirmed the diagnosis of patient and parents, stating that the pin had turned, and the head only was visible; also that an operation would be required for its removal. After several hours of suffering, she was induced to consult the physician who gave me the particulars of the case. A circumscribed abscess was found on the floor of the meatus, just at the curve of the canal. The treatment was rapid and effective: puncturing the abscess and application of warm water gave immediate relief.

Prognosis, with reference to each individual abscess, is favorable; bad results seldom follow them, although periosteal inflammation has been occasionally induced by the inflammatory action attendant on their development. They sometimes induce, as a consequence of their successions, thickening of the tissues of the canal and limit its calibre. The saddest feature of these cases is the tendency to recurrence. Sooner or later they come in succession, each one bringing its train of tormenting painful symptoms.

"Von Tröltsch reports a case where the patient, for twelve years, with intervals of ten weeks, and at the highest of two months, suffered from such furuncles, now in one ear and then in the other, and with which there was great febrile disturbance, so that at each attack he was obliged to remain in bed."


Sometimes the inflammation extends to the deeper portions of the connective tissue, also to the tissues covering the mastoid process.



*Treatment.*—The principles of treatment can be summed up in a few words. Incise promptly, favor suppuration and discharge of core, and prevent recurrence by all the means available. Heat and moisture favor suppuration, and may be applied in various forms. The author has found the hypodermic use of morphia and the hydrate of chloral exceedingly valuable as palliatives, especially the latter, which is given in from five- to fifteen-grain doses until the pain is relieved.

So important is early incision that patients who have suffered from the tedious course of the abscess, and subsequently on recurrence of the disease been relieved by early incision, will give up the prejudice against the "horrid knife," and even request its use. So soon as the incision has been made, the ear should be filled with warm water, and the part kept constantly in a warm bath for at least half an hour, or better still, "steam" projected into it by means of the atomizer. If the core can be distinguished, gentle pressure with Daviel's scoop, or Gross's instrument for removing foreign bodies, will facilitate its escape, or Hagen's blunt bistoury, with spoon for evacuating the abscess. (See Fig. 58.)

Fig. 58.



Make your incision at once, without any further ceremony, for often the fear of the knife is greater than the fear of the pain. The knife should be as sharp as a "cataract" knife; for, if surgeons were more careful about keeping the edges of their instruments keen, there would be less pain, and, consequently, less fear. Do not wait until Nature has done



the work, until pus has formed, but cut exactly upon the painful spot; seek only to relieve, and your patient will thank you. When it is over, immediately favor the discharge of blood by the continued injection of warm water. Here Clarke's douche is invaluable, and sufficient injection of warm anodyne fluids should be used to keep the canal clear of pus, blood, cerumen, and epithelium, as they accumulate.

What shall we do to prevent recurrence of these abscesses? To this question there are many answers. Nitrate of silver, preparations of mercury, syringing, oils of fruits, nuts, and animals, and even petroleum, have all had their stanch advocates, and yet recurrent abscesses are as frequent as ever.

We should not, on this account, abandon a case because specifics have failed, but rather seek to modify our treatment to each case.

Should the patient refuse to have the knife used, anodynes must be resorted to. In prophylactic treatment the author has found the best results follow the use of iodide of iron, either in syrup or Blanchard's pills, or pill iodiform et ferri, especially in strumous cases. This must be persevered in for several weeks. The following are a few of the cases treated, and will serve to illustrate the above:

CASE I. *Abscess of the external meatus*.—Miss M. C., aged thirty. September 19th, opened it, and applied heat and warm water. Is a delicate lady; has not had an attack for four years, at which time she had them in both ears; was under my care at that time. Internal treatment, syr. ferri iodidi, in twenty-drop doses, three times a day.

Miss H., aged twenty-two, a cousin of Miss C., had one at the same time, four years ago, and the same treatment was successfully applied.

CASE II.—A few days after, Mrs. G., aged forty-five, called, and was suffering from the same affection. It had opened, after three days' suffering, she having lost the hearing in her left ear by an operation performed by a surgeon of this city in removing a tumor when a child. Introduced the spoon end of a tube and removed about half a teaspoonful of pus;



and directed a warm-water dressing, and the same internal treatment as the previous cases. This abscess was situated about the centre of the canal.

CASE III.—September 29th. Mrs. L., aged forty-five, has been suffering for some days with an abscess in front of the tragus, but pointing inwards. In spite of narcotics, has given her most intense pain. She has a feeling or sense of stuffing, buzzing, and throbbing in the ear, accompanied with fever. The inflammation spread over the whole malar region. She has poulticed it with flaxseed-meal and bread-and-milk, but it will not discharge. Her husband, therefore, brought her for advice. I at once opened it by a free incision with a tenotomy-knife, and a full teaspoonful of laudable pus was discharged. Directed the free use of warm water; also a solution of plumbi acetate et opii to the part. As her health was somewhat broken down, I gave her the syr. ferri iodidi, twenty drops every three hours, with nourishing diet.

The above cases yielded kindly to treatment, and serve as illustrations as regards position of such abscesses in the external auditory canal.

*Osseous tumors* in the auditory canal are not frequent, and these cases will illustrate the subject. Toynbee's treatment consisted in alterative doses of blue mass; the surface washed with a solution of nitrate of silver (5j to 3j) every fourth or fifth day, and afterwards every seventh day; in another case, tincture of iodine was applied to the surface of the tumor, behind the ear, and four grains of iodide of potassium were given three times daily for between two and three months. This latter treatment was found successful in the following case of tumors of the auditory canal and deafness:


F. S., Esq., aged fifty-five, born in England, residence, Walnut Street, Philadelphia, consulted me on March 5th, 1871. His general health was good, but has been deaf in his left ear for a long time, and could not hear his watch, even when pressed close to that ear. He has also become suddenly deaf in the right ear, and can only hear a watch when close to the mastoid process, not distinguishing a loud-ticking watch when pressed on the auricle. He states he had gout several years ago, and, when a boy, an eruption outside and inside of



the ear, with more or less discharge. He had been under the care of a gentleman of some distinction in London, who advised him to have nothing done with the left ear, as no improvement would be likely to ensue. This was ten or twelve years ago, and at present he has in his neck a cicatrix of an old abscess. His daughter was under the writer's care for deafness, and was relieved, two years since; and he has had two other children, both of whom died of tuberculosis of the lungs.

On examination of the right ear, found the lining membrane congested, and near the membrane tympani, a gouty deposit the size of a large pea, projecting half way across the

Fig. 58.




meatus. (See Fig. 58.) The membrana tympani had a sodden and white appearance, and numerous scales of dark-colored epithelium were removed by washing out the ear. The whole space was contracted and dull from an old discharge. The Eustachian tube of that side was not opened by the Valsalvian method, so the Eustachian catheter was introduced and the tube well opened; but no decided improvement of the hearing followed. Applied tincture of iodine to the bony tumor, and ten-grain doses of iodide of potassium internally three times a day.

March 10th. Exhibited the case to Dr. Knight and Prof. Meigs. The latter suggested that the same condition he had no doubt would be found in the interior of the ear, fixing the stapes by osseous connections. After several days' careful washing with a slight astringent and carbolic acid, the writer employed Siegel's aural speculum, and exhausted the air carefully until he felt the motion, but the only improvement was hearing the watch by the auricle. Having so little success with the left ear, he resorted to the other, and found a collection of hardened dark cerumen; this was softened for three nights with a warm alkaline solution. On the



13th the ear was carefully washed, and the dark-brown mass removed, much to the patient's surprise. On examination of the membrana tympani, it was a dull white (no malleus seen, nor bright spot), membrane very concave, and three osseous (see Fig. 59) tumors

Fig. 59.



were found near it. No improvement was made after having air passed into the middle ear, except that for seven days he was able to hear the watch when pressed close, but he observed it was faint. He could also hear the human voice much better, and in the case of a watch, it has been years since he has heard one in that ear.

March 24th. Applied to tumor of left ear a solution of nitrate of silver, seventy grains to the ounce of pure French glycerine, limiting it so as to remove the epithelium. Opened the Eustachian tube by the catheter, desiring him to practice the Valsalvian operation on entering a warm room and not to return for four days.

March 28th. My patient came this day, says simply he is so much better that he is satisfied with his hearing, that it is quite restored. On testing it by my watch, he heard it one foot off, and the voice several feet; his right ear is also much better, and he can hear the watch and the voice several feet.

May, 1871. The improvement is permanent.

#### OTITIS EXTERNA DIFFUSA.

This is an inflammation of the external auditory canal, affecting only the cutis and cutis vera, but it may extend to the outer surface of the membrana tympani, etc.

It is generally an acute affection, but should it become chronic, it may involve the periosteum; if this is the case, we have then a general *otitis*, and pus may be found in the middle ear, and then we class it as *otitis interna*. The



auditory canal, no matter how near the membrana tympani, is not lined by a mucous membrane; we therefore consider both Toynbee and Politzer as not teaching sound doctrine by speaking of a catarrhal inflammation of the auditory canal. The term *catarrh*, according to a proper nomenclature, pertains only to affections of the mucous membrane; hence its use for inflammations of the external auditory canal is not proper, and aural catarrh should only be used for the middle ear.

*Causes.*—In our opinion it occurs chiefly in scrofulous and tuberculous children, and frequently alternates with an affection of the eye; we also fully agree with *Rau* that it always accompanies the dentition of certain children, cutting the hair, and applying cold water to the head, with certain forms of the exanthemata, as measles, chicken-pox; while smallpox and scarlet fever almost always produce otitis media, or interna, with perforation.

*Symptoms.*—In almost every instance it is ushered in with pain in the ear, with or without itching, which the patient desires to relieve with the head of a pin or ear-scoop; or the finger is thrust into the ear, but, owing to the pain, is unable to scratch it with any comfort. This pain increases towards night, when it is so severe as to prevent the patient from sleeping, and the head is pushed into the pillow, in an attempt to avoid the slightest motion, as it increases the deep-seated pain. In very young children the only way of knowing that it is not colic, or some other disease, is by making pressure under the lobe. In other cases but little pain is experienced, and the only indication of the inflammation being there is found in the discharge on the cap or pillow in the morning. This is at first white in color, and very thin, almost like water; after a time it becomes thicker, stringy like mucus, and in some rare cases it is yellow; this latter is only seen when the inflammation has been of a high grade.

If, in the beginning of the attack, we examine the ear by the speculum, we will find the epidermis, with the surface of the membrana tympani, very much injected and swollen. Should this be attempted later, the whole auditory canal will be found filled with white, moist, desquamating cells; these



must be removed with great care, as we frequently find the membrana tympani so soft as to be easily injured. Therefore these scales should be washed out by means of Clarke's modification of Thudichum's douche, a continuous gentle stream, the water being very warm, as rude syringing is also very painful to the patient. In the subsequent examinations it will be found that the canal is so swollen that great care is necessary in introducing the aural speculum each day, so as to see the progress of the disease. The amount of the impairment of hearing depends upon the closure of the canal, and also upon the amount of the membrana tympani involved.

*Prognosis.*—This will depend upon the constitution of the child, and also upon the cause producing the disease; but as a general rule the result is favorable.

*Treatment.*—In the acute form, active purgations, free leeching, with warm spray of solutions of opium, hops, or poppy-head. If a child, a warm poultice of ground flaxseed and elm bark over the whole side of the face, and a dossil of cotton or lint, saturated with an anodyne lotion, in the ear, will give great relief. In the second stage, when there is a free discharge and little or no pain, then we may resort to a solution of the biborate of soda, hyposulphite, or bisulphite, with water and glycerine; with counter-irritation, by an ointment of iodine and iodide of potassium, with lead or mercury, to remove the deposits in the ear. Nothing cold should be applied, but every application should be warm, or about the temperature of the skin.

Long-continued poulticing we consider injurious, but in the very early stage one good poultice quickly produces exudation and free discharge, also a cessation of tension and pain in this form of otitis externa. I fully agree with Von Tröltzsch that poultices should not be applied in deep-seated inflammatory affections of the ear in which there is disturbance of the brain, as they assist in producing profuse and wearying discharges. We now come to the third stage of these affections, and we call these cases subacute, and they are generally found among the poor out-door cases which attend our clinics; these cases require a solution of sulphate of copper or nitrate of silver applied by the surgeon's hand



to the granular surface, and a wash of sulphate of zinc or hydrated chloride of aluminium, two gr. to  $\bar{3}$ i of water, which the patient can use at home, filling up the ear with it, warm, and using a plug of cotton, and allowing it to remain from ten to fifteen minutes, repeated twice a day until the discharge ceases. Great care must be exercised in cold weather for fear of an acute attack supervening on the old one by taking cold, which frequently involves the periosteum, or mastoid cells, or, passing to the cerebellum, causes the death of the patient. In very cold weather the ear should be protected by a cap or pledget of cotton, and I am of the firm opinion that the hair is the natural protector of the ear, and it never should be wet and then allowed to hang in damp folds around it; nor should it be cut very close during cold weather. In warm weather no protection is required for the ears, but the use of an abundance of fresh water to wash the ears, neck, and front of the chest, followed by wiping the parts perfectly dry.



## CHAPTER XI.

### OTITIS MEDIA PURULENTIA, ACUTE, SUBACUTE, CHRONIC, AND HYPERTROPHIC (SUPPURATION IN THE MIDDLE EAR).

THIS affection is usually induced by cold air or water, introduced through the Eustachian tube, or by a perforation in the membrana tympani, and in some instances caused by the feet being exposed to the cold. In several cases we have known it to occur by a wave striking the ear in sea-bathing, or a sudden plunge or dive in very cold water. It is also a sequela in various forms of fever, whooping-cough, measles, and during the irritation of dentition.

*Symptoms.*—Pain in the ear, intermittent in character, commencing at night, and disappearing in the morning; after a time the pain is constant and very much increased in intensity as it reaches the vicinity of the middle ear, involving the membrana tympani and osseous structures. It also extends to the head and neck, attended by fever and a furred tongue. As soon as pus is formed there is more or less of a creep, and in some instances a decided chill. It is always accompanied by deafness, noises, throbbing, with fullness, and soon after is followed by a discharge; first a faint moisture, then a cream-like fluid, and at last an abundant purulent excretion. In young children unable to express their feelings, the only indication is a decided and persistent screaming for hours, which is often confounded with the symptoms of colic, and our diagnosis should be determined by exclusion. Pressure over the antitragus will often confirm the result by the pain which the child suffers; but our only sure guide will be a slight discharge on the pillow in the morning. This discharge has a peculiar odor, resembling decayed cheese. In an account of an interesting case reported by



Prof. H. Knapp,\* the purulent inflammation was caused by the use of *cold water*, by means of Weber's nasal douche, confirmed in cases reported by Roosa and Moos. It is explained thus: the cold water is more apt to cause involuntary swallowing than warm, by opening the Eustachian tubes. Besides the peculiar nature of its origin, the above case of otitis media purulentia was very remarkable for a symptom not much noticed in such cases: viz., *double hearing with both ears*.

Tröltsch and Politzer mention its occurrence; Moos records several cases in his "Klinik der Ohrenkrankheiten" (for notice of which see Nervous Deafness). In the case of Prof. Knapp, the meatus was swollen and red, the membrana tympani injected, soft, and projecting into the meatus, and after a time becoming a greenish-yellow color, with perforation.

*Prognosis.*—Generally favorable unless in case where the blood becomes altered by scarlet, typhoid, or typhus fevers, or in scrofulous or tubercular cases. In the first class, or in healthy individuals (if the case is promptly treated), resolution takes place, and after several weeks the discharge ceases, the membrana tympani heals, and the hearing is restored; but if, on the contrary, the case is neglected, or the constitution is impaired or altered, it passes into chronic otitis media with enlarged perforation. We shall illustrate the treatment by a clinical case.

CASE I.—Mrs. Margaret B., aged forty years, a strong, healthy lady, had been exposed when the thermometer was 13° Fahr., her head being protected with the slight covering of the present fashion. She was out-of-doors most of the day of December 13th, in the evening complained of pain and heaviness in her head, with a decided chill. On the 14th, she had a discharge of serum from the nostrils, with slight relief to the head. On the 15th, the discharge from the nostrils was mucus mixed with blood, which continued also on the 16th, and then suddenly ceased, when the face and gums began to ache. Application of hot water, with warm flaxseed poultices, relieved the pain in the head. On the

---

\* Archives of Ophthalmology and Otology, by Prof. Knapp and Moos. New York, Wm. Wood & Co., 1870, p. 365.



17th, she suffered from great pain in the ears, and which was allayed by steaming the ear with hot water. On the 18th, the ear and throat began to discharge, and the pain returned at night with renewed violence. On the 20th, the left membrana tympani was perforated. Upon that side I could see the point where the perforation was about to take place, and at my visit on the following day I saw the *point* of perforation with the bubble of air under it.

*Treatment.*—A blister in front of the ear, calomel and morphia at night, and a wash of glycerine and water, equal parts, with ten grains of sulphate of zinc to the ounce, which she continued to use until the 25th. During the last week of treatment the wash was changed to two grains of nitrate of silver to the ounce, each, of glycerine and water, with use of the air douche. Feb. 1st, patient able to resume her usual occupations, and after some weeks the discharge gradually reduced, the perforated membrane was healed over, and her hearing almost as good as before her attack.

In this affection we have several symptoms which we have in acute inflammation of the membrana tympani. In the class of patients which present themselves at our clinics with this affection of the ear, there is almost always a feeble condition of the system, accompanied by no fever, and but slight pain; it is therefore properly termed *otitis media purulenta chronica*, or otorrhœa, with perforation of the membrana tympani.

The first appearance which we will always notice is an irritable red meatus, and when the ear has been carefully washed out, so that we can, with more or less effort, see the character of the opening in the membrana tympani, it will be almost invariably found not healthy or ready to heal. The next consideration is, what keeps the membrane open, and why is it not ready to heal, as in the acute form in private practice? It is because there is disease in the middle ear, vascularity, or a granular condition, or some accumulation of pus or mucus not removed, producing an irritation, and preventing permanent cicatrization. The external meatus is usually found filled with pus or mucus, which causes the patient a feeling of fullness. After its



removal, the opening is seen in the membrana tympani, causing a whistling sound when the nose and mouth are closed and we blow outwards or expire. The hearing is always defective, especially if the Eustachian tube is not open. *That* we can determine by taking a mouthful of water, holding it, then closing nose and mouth, and making the effort to swallow; or, a more certain way is by the use of the Eustachian catheter and otoscope, or Politzer's air douche. The patient's voice is almost always affected, and becomes nasal, and there are symptoms of irritation in the nose and throat. When the meatus is carefully examined, it is found inflamed, and covered with softened scales of epithelium, exposing the cutis vera, with here and there masses of dried pus or mucus, with hairs, and, in many instances, small masses of yellowish wax. This chronic affection, owing to the exposure of the mucous membrane of the middle ear, by the perforation in the drum, is very liable to frequent relapses into the subacute stage; we must also remember how very near this inflammation is to the brain when it gets beyond the barrier of the membrana tympani, and act always with caution.

*Prognosis.*—This will depend largely upon the severity of the attack, the general health of the individual, and the duration of the perforation.

If the attack has been one in which there was much mental disturbance, severe or protracted fever, convulsions or blood disease, the prognosis is very unfavorable for decided improvement in hearing, although the discharge may cease and the perforation heal. If the individual is of a scrofulous or tuberculous diathesis, the healing of the perforation rarely occurs, but rather tends to increase, until entire destruction of the membrana tympani takes place, with loss of the small bones. Another termination is in otitis interna, or pyæmia, which finally results in death. If the perforation is large, and has been of several months' duration, it is very difficult to heal even in a healthy person. Another important agent in the prognosis is the character of the discharge: if it is mucus alone, or even muco-purulent, the prognosis is very favorable. If it consists entirely of true pus of a yellowish



tint, the prognosis is not as favorable, and indicates a higher grade of inflammation; while if the pus is sanious, serous, ichorous, or serofulous, the prognosis is very unfavorable. The prognosis, as before stated, in purulent inflammation of the middle ear is always grave in character, although statistics show that the actual number of deaths is not as alarming in proportion to the cases. There exists in every case a liability to sudden extension of the inflammation to the inner ear, causing cerebral complication, which it is impossible for the physician always to prevent. This complication, with that of the mastoid process, will lead to caries and necrosis, especially in youth and middle life. Gruber's advice to the surgeon in all such cases is to inform the family of the dangers which may arise, in order to save his reputation, in case of the sudden outbreak of these symptoms.

We find the majority of general practitioners not well informed of the nature of chronic suppuration of the middle ear, confounding it with chronic suppuration of the external auditory canal, under the old term of "otorrhœa," and rarely ever endeavoring to distinguish the difference by ocular inspection with the aid of proper instruments. Indeed, but few can tell a purulent otitis of the middle ear, with perforation of the membrana tympani, even if shown it, so unaccustomed are they to take the trouble to understand its nature. The statistics of writers on diseases of the ear, and the reports of eye and ear infirmaries and hospitals, show that, instead of external disease of the ear, we have in the majority of cases a most serious disease of the internal ear:

	External Otitis.	Chronic Sup- puration in Middle Ear.	Total.
New York Eye and Ear Infirmary, 1869 . . .	46	324	370
Brooklyn Eye and Ear Hospital, 1869 . . .	17	85	102
Ophthalmic and Aural Institute, 1869 . . .	19	84	103
*Schwartz's Statistics, 166 cases . . .	15	39	54
†Roosa's Statistics, 500 cases . . .	11	111	122
Turnbull's Statistics, 600 cases, Howard Hospital	30	99	129
	<u>138</u>	<u>742</u>	<u>880</u>

\* Vide Archiv für Ohrenheilkunde, B. I.

† Roosa, New York Medical Journal, 1869.



In summing up these figures, it will be seen that there were 742 cases of chronic suppuration of the middle ear to 138 cases of chronic inflammation of the external auditory canal. We might state, in this connection, that we received this day (Oct. 31st, 1870) an illustration of the want of knowledge of the general practitioner under this head. From the neighborhood of this city, a little boy, aged eight years, came to us with chronic "suppuration of the middle ear," affecting both ears. He had been under the care of a most respectable, intelligent, general practitioner, a graduate of one of our best institutions, and yet he never knew that the little fellow had perforation of the membrana tympani, although he had been treating him for seven years and two months, almost the entire life of his patient. According to the statement of the mother, he had never washed the ear or examined the boy with an ear speculum or by reflected light, but gave her a wash, blistered behind the ears, and trusted the cleansing to her.

The first and most important matter in the treatment of acute and chronic suppuration of the middle ear is to wash it out with warm water, to which is added one grain of biborate of soda to each ounce, by means of Clarke's modification of Thudichum's douche, without injury to the delicate perforated membrane.

If there is pain, heat, swelling, and redness, indicating sub-acute inflammation, we must deplete, either locally or generally, or both; but as a rule local depletion is all that is necessary. We prefer leeching, by means of Swedish leeches, applying them as close as possible to the ear; if these cannot be procured, we employ American, in the proportion of six to one Swedish; if neither can be had, we should then resort to very small cut cups or the artificial leech. After the depletion, Canton flannel cloths or patent lint, wrung out of hot water and covered with extra cloth or oil-silk, will materially assist the flow of blood and give relief to the pain in the ear. If these means fail to subdue the pain, resort to morphia sulphatis, either by the mouth or hypodermically, commencing with one-sixteenth of a grain, and gradually increasing to one-eighth or one-quarter or even one-half of a grain, at intervals of three hours. If these means fail, we try the



hydrate of chloral dissolved in lemon, orange, or ginger syrup, beginning with fifteen grains every three hours, largely diluted with water; viz., two ounces of fluid, and increasing the chloral to thirty, forty, or even fifty grains, or combining the morphia and chloral, as they will sometimes act better than either would alone. After the patient has been relieved of the acute pain, and has had some rest, and yet continues to be disturbed with distress in the head, with a hard pulse, we use the foot baths, adding mustard or common salt, and placing the feet of the patient in moderately hot water, adding gradually water of increased temperature as the patient is able to bear, until relief is obtained. If these means fail, we administer an active purgative of calomel, rhubarb, and ginger, or a saline mixture, as the following:

R.—Mass Hydrargyri, gr. x;  
 Extractum Hyoseyami, gr. i;  
 Magnes. Calc. (Husband's), ʒij;  
 Aq. Menth. Pip.  
 Syr. Simplex, aa ʒj. f. Misce.

S. To be taken at one dose, or divided into two.

This relieves the bowels, acts on the liver, and depletes from the intestines with signal relief. In most of the subacute attacks the discharge for the time ceases; and relief is always experienced when it flows freely. Von Tröltzsch recommends filling the ear with lukewarm water, and leaving it in the ear from five to ten minutes, the patient at the same time lying on the other side. We have found also relief from the use of the warm cloths frequently renewed; or in some instances, the elastic bag of Chapman; or in other cases, dry heat by means of a bag of common salt.

After the subacute symptoms have disappeared, we must diminish the discharge and give nature an opportunity to act, but we must not resort to violence; therefore we still recommend the use of the douche or post-nasal syringe to wash away the excess of secretion, for it is bad surgery to remove all the covering from the surface of an ulcer, or even an exudation, as we then bring the air and other irritating substances in contact with it. Clean it by allowing the fluid to flow over it, not to rub it off. One of the greatest evils is



the resort to astringents too soon, and it is better for some time to employ the biborate of soda, or the bisulphite or hyposulphite, a few grains in warm or tepid water, so as to remove whatever would interfere with the healing process; if there is much odor, carbolic acid and water,  $\mathfrak{z}\text{i}$  to  $\mathfrak{z}\text{iv}$ . When the parts become pale or flabby, then it is time, and only then, to give a slight stimulation, as we would to an ulcer; for instance, one or two grains of the acetate of zinc with the one-twelfth or one-sixteenth of a grain of the acetate of morphia, dissolved in the douche three times a day, and allowed to remain in the ear. Opening the Eustachian tube is always necessary by the catheter, and gently passing air through so as to be able to see a drop or two of liquor forced through the cotton, or by resorting to Politzer's method, we have a current of air to assist in cleansing the cavity of pus, mucus, altered lymph, or even blood. This will also assist in bringing the necessary stimulation and wash in contact with the altered mucous membrane of the cavity of the tympanum.

In the case of very young children, Mr. Hinton suggests the use of a piece of small rubber tubing tipped with ivory or bone instead of the air-bag, and this we have used, and the operator is to blow gently the instant the patient swallows. In little children the inflammatory stage is usually a short one, and the most disagreeable feature about the discharge is the odor, which is soon removed by the use of the chlorine-water, prepared by passing chlorine gas into water, or dissolving a portion of the chloride of lime, one drachm to half a pint of water, and filtering it. The writer usually dilutes this in the proportion of a teaspoonful to a cupful of tepid water, gradually increasing it to a tablespoonful; after a time this is replaced by alum, in the proportion of  $\text{gr. ij-v}$  to  $\mathfrak{f}\mathfrak{z}\text{i}$  of water; using morning and evening half a teaspoonful, warm, and allowing it to remain ten minutes, with the ear closed by cotton.

As we frequently find that after a time no improvement takes place, we then have to resort to a stronger stimulant, namely: the nitrate of silver, or sulphate of zinc, or copper, in strength of twenty, thirty, forty, and even sixty grains to the ounce, applied by means of a very delicate piece of sponge, or



cotton, attached to a long, slender wire of aluminium, or silver. The advantages of the sponge over the cotton are that it retains its solution until pressed to the part; at the same time none of the cotton fibres are left in the ear. The delicate sponge is also useful in removing particles of epithelium, pus, mucus or altered cerumen. With the use of the mirror we can also touch vegetations, and limit them to the spot. We have given up the use of the solid stick, as we found we could not limit it, and, in some instances, it destroyed too much by its caustic action. It is true we can control it by the chlorine-water, or a solution of common salt.

Politzer\* states: "Vegetations in the external meatus, on the membrana tympani, or the promontory, which, so long as they remain, keep up the otorrhœa, must be removed, when they can be easily reached, or destroyed, by touching with a concentrated solution of muriate of iron, or, if *hard*, by nitrate of silver in substance." He employs for the latter a simple metallic wire of the length of a sound, bent at right angles at the middle, immersing the point several times in nitrate of silver, melted in a little porcelain cup, till the caustic has hardened upon it in the form of a little ball.

An equally good authority, Wilde,† states: "After a long and fair trial of several means proposed for healing apertures in the membrana tympani, I have come to the conclusion that there is nothing like nitrate of silver. It will not, however, answer to thrust a stick of lunar caustic, fastened to a quill, down the meatus, for the purpose of touching any part of the membrana tympani; to be used with effect, the escharotic should be applied upon the extreme edge of the aperture, or rather within the ring of the opening, every second or third day, so long as the part seems inclined to close; *but the moment we perceive it enlarging*, a day or two after the application, we must desist. As it is not always possible to reduce a pencil of nitrate of silver to the requisite degree of fineness, I have for several years past been in the habit of

---

\* The Membrana Tympani in Health and Disease, Translation, p. 130. New York, 1869.

† The Nature and Treatment of Diseases of the Ear, American ed. Phila., 1853, p. 292.



pointing probes and other instruments used in minor surgery, with nitrate of silver attached to them in the following manner: a portion of the caustic is to be melted to the boiling-point in a small silver or platina ladle, such as is represented at Fig. 96, held over a spirit-lamp. When the caustic has become quite clear, the point of the probe, or instrument, to which it is to be attached, should be heated in the flame of the lamp, and then dipped into the fluid caustic several times, until a sufficient quantity is taken up. The instrument figured at Fig. 97 will be found very convenient for applying nitrate of silver to any surface within the meatus. This porte-caustic is about six inches long, and consists of a silver tube, cut spirally for three-fifths of its length, and having an aperture in the side, or a hollow at its end. When about to be used, its extremity should be coated with caustic, as already directed. The elastic spring prevents any injury to the ear from the starting of the patient, and can also be bent, so as to be applied with ease to any part of the auditory canal. When granulations sprout from a perforation in the membrane,—which is very rare, unless the rupture occurs immediately adjoining its edge,—this method of applying caustic will be found very convenient; but the writer cannot state that it has been very effectual, in his hands.

Before concluding this part of our subject, we must not omit the importance of constitutional treatment in this disease, by means of tonics, alteratives, cod-liver oil, extract of malt, iodine, quinine, and mercury, and also means of counter-irritation by blisters, croton oil, tincture of iodine, and, in some rare cases, the actual cautery; these are all necessary measures of success in removing the discharge. As the tables of our German friends show better success in treatment of this and one or two other forms of diseases of the ear, I have given the views in extenso of Dr. Gruber, one of the most recent writers upon this subject, holding a distinguished position in Germany.

“*Treatment.*—Since not merely the faculty of hearing, but even the life of the patient are in the greatest danger, the utmost circumspection and energy should, as a matter of course, be exercised, and with the best medical experience.



The least neglect may destroy the life of the patient, while a skillful and timely action will be followed by the most happy results. The treatment is mostly symptomatic, and consists in preventing the inflammation from spreading, especially to deeper parts, and in endeavoring to save such parts as have been already attacked. As much as possible the treatment must be antiphlogistic, care being taken to avoid those means which, by impeding the reflux of the blood, increase the hyperæmia of deeper parts. As long as the inflammation is at the stage of hyperæmia, as well as when the patient is suffering extremely, suppuration having occurred, local bleeding may be employed. In the robust, where there is considerable cerebral pressure, venesection will be most serviceable.

“The leeches should be applied around the ear, especially over the mastoid process. It should be remembered that as the depletion is not directly from the inflamed part, and as it has for its object the mitigation of the hyperæmia of the brain and its membranes, to be effective, it should be copious, and the after-bleeding sufficiently maintained.

“As a matter of course, the depletion must be repeated during the course of the disease, as often as the symptoms require and the physical condition will permit. The greatest precaution is to be used with regard to the local application of cold water or ice, as in some patients this may increase the cerebral hyperæmia. This is shown by increased headache and by the symptoms of compression becoming more obvious. It would seem that the hyperæmia of the cranium, in the disease under consideration, frequently remains confined to the region nearest the inflammatory focus, and cold applications to those parts heightens the pain by increasing the hyperæmia, while the effect of cold is beneficial when applied to remote regions of the head, as, for instance, the forehead. The promotion of the reflux of blood from the inflamed parts by the application of warmth, is, perhaps, most serviceable, since it is only thus that, with some patients, warm cataplasms applied to the region of the mastoid process and the corresponding half of the occiput, with or without the simultaneous use of cold to other parts of the head, cause



such extraordinary relief. I have treated patients in whom at the same time ice was applied to the forehead and warm poultices in the vicinity of the diseased ear, in order to somewhat mitigate the raging pain. The revulsion from the head produced by other means will likewise be useful, especially at the beginning of the disease. In this respect the derivative action of the intestinal canal will prove most beneficial, the remedies being chosen according to the strength of the patient and the condition of the intestinal viscera; and it must also be remembered that only a powerful derivation can be of use. In those cases where the latter is not deemed necessary, care should be taken at least to keep the bowels in an open condition. As to internal medicaments, the principle should be adhered to, to avoid all such as would be injurious in concussion of the brain. With regard to purgatives, all those remedies should consequently by all means be avoided which will cause nausea and vomiting. The same principle likewise holds good in the case of patients who labor from idiosyncrasy with regard to some one medicine.

“By derivation by the skin by means of vesication or sinapisms, applied to the region of the mastoid process or to the nape of the neck, the inflammation in the deep-seated parts will not be lessened or even cut short, although it will greatly soothe the pain.

“In cases where these remedies are contraindicated, or are not sufficient to relieve the affection, it is advisable to employ, both externally and internally, those narcotics applicable to other inflammatory processes of the ear, but particularly when the rest of the patient at night is disturbed, and collapse threatened. As acute purulent otitis is a febrile disease, it is superfluous to remark that the strictest bodily and mental rest must be enjoined, and a restricted diet observed. The local treatment must further correspond to the objective symptoms presented in the ear at the time. The local treatment for a high degree of hyperæmia has been given. If the inflammation have passed into suppuration, the local treatment has the further object of giving an outlet to the pus, even by the formation of artificial canals, if it should be im-



possible to evacuate it in the natural way. Unless this is performed timely and adequately, the life of the patient will be exposed to the greatest danger from the proximity of the brain and its membranes in general, and of such important blood-vessels as the internal carotid, jugular vein, and meningeal veins. Should the Eustachian tube be narrowed or closed, or should it not suffice for giving a proper passage to the matter on account of its rapid formation, the artificial perforation of the tympanic membrane—*i.e.* the paracentesis of the membrana tympani, or myringotomy—is indicated. In such cases the phenomena presented by the membrana tympani will indicate the conditions of the organs in the interior ear. The artificial perforation of the tympanum has for its object not only to give an issue to the already formed pus, but also to cause relaxation of the engorged tissues by the division of the blood-vessels of the membrana tympani. Hence it is that, after the operation, the pain will quickly, and in a surprising manner, remit. Myringotomy is, therefore, an operation which causes the patient great relief and comfort, and may, under certain circumstances, be the means of saving life.

“As this operation is performed for two objects, it cannot be a matter of indifference what point for the incision we select. Since exudations will, according to the law of gravitation, always occupy the deepest portion of the cavity of the tympanum, the lower half of the membrane must be the most suitable. It is, however, advisable to give the preference to that segment which presents the greatest amount of swelling. As in the opening of any other abscess, so in that under consideration, the rule should be observed not to make the opening too small, since in that case the evacuation will be incomplete. It is better to make the incision a little too large than too small, since wounds of the membrana tympani, especially on the subsidence of the inflammation of the interior, will heal readily. The practical value of this latter principle is obvious in those cases in which the exudation into the cavity of the tympanum is very considerable, in which case the discharge cannot take place through too small an opening. Should the retraction of the tissue of the membrane follow-



ing a large incision not produce a sufficiently large opening, it may even be necessary to make one of a crucial form.

“The operation of paracentesis is best performed by the myringotome, or with a simple cataract-needle. It may be done under direct illumination in those cases in which the meatus auditorius is wide enough to admit of sufficient light for the illumination of the deeper parts. If, on the contrary, it should be too narrow, the operation should be performed by means of the reflected light, and the reflector with the frontlet mirror should be used, thus allowing the surgeon the free use of both hands for the operation. The head of the patient is, at the same time, to be supported by an assistant, and the largest speculum introduced that the size of the meatus will allow. After the operator has made a proper inspection of the segment of the membrane to be incised, he inserts the myringotome at the most inferior point, and prolonging the incision upward as quickly as possible, in drawing the instrument out he cuts through the membrane to the proper length.

“The matter collected in the cavity of the tympanum will then be evacuated through the aperture. The bleeding from the separated blood-vessels is but very scant, amounting to but a few drops. It can, however, be increased by the careful injection of pure lukewarm water, sometimes practiced for washing out coagula. It is advisable to promote the discharge from the interior ear by the use of the air douche. As long as more pus is being formed at the original focus of inflammation than can be removed in the natural way, the aperture should be kept open by repeated applications of the air douche, and if this fail, by repeated inspection and the insertion of the probe. In case of closure of the opening, which may take place in one night, the repetition of the paracentesis is indicated. It has happened to myself and to other aural surgeons, that the operation had to be repeated twice, and even oftener, in the course of the same morbid process.

“To cause the matter to pass out by Valsalva's method is, in general, not advisable; a stagnation, although transient in the blood-vessels of the brain and its membranes, being pro-



duced by the pressure upon the veins, might exert a prejudicial influence. The patient, suffering under purulent otitis, even when an adult, is, however, generally too weak and languid to perform this movement.

“As long as the organs are engorged with blood, and painful, care should be taken not to drop astringent or other stimulating medicaments into the auditory meatus, or even make injection of such substances into the Eustachian tubes; but astringent gargles, and such as will promote absorption, are applicable in those cases in which an inflammatory swelling of the fauces complicates the disease under consideration. Even the application of a concentrated solution of nitrate of silver has proved beneficial, by diminishing the swelling of the mucous membrane of the Eustachian tube. Injections of astringents into the Eustachian tube ought to be employed with the greatest caution, and cannot be used before the pain is over, the fever ceased, the affection beginning to assume a chronic character, and there are those conditions generally indicating astringent injections. Employed under such circumstances, and with due regard to the rules given under the head of Catarrhal Inflammations, they may prove most beneficial.

“During the course of purulent otitis media, the necessary attention should also be paid to the several parts of the external passage of the ear. In this connection it must be mentioned that the perforation must always be kept open, to allow of the escape of the matter, which, if retained, might decompose, and exert a most injurious influence. If the opening has become contracted from a simple swelling of the soft parts, it should be dilated by the introduction of small pieces of lint, or similar substances. In not a few cases the application of leeches, or even incisions into the soft parts, will be indicated to subdue the swelling. If there should be a complication by inflammation of the external meatus, or periostitis in the region of the mastoid process, the latter is to be treated according to the principles laid down in the respective chapters.

“Should the ominous chills occur, the fever should be treated by larger doses of quinia, five or ten grains, repeated twice



or thrice a day, the body of the patient being properly warmed during the chill, and acids given during apyrexia.

“When otitis depends upon a scrofulous or syphilitic diathesis, absorbent ointments are applicable during its course. In recent exudation their application in the region of the mastoid process is likewise most helpful.

“As in chronic catarrh with perforation of the membrana tympani, so in chronic purulent otitis there will be a most profuse otorrhœa. The effect of the remedies applied to the Eustachian tube must, in such cases, be supported by dropping appropriate medicaments into the external meatus. We make use most frequently of a solution of sulphate of zinc, from two to ten grains to the fluidounce of water, and of borax, alum, or tannin in the same proportion, and occasionally of sulphate of copper, from one to three grains to the fluidounce. Tannin is but seldom used. Where there is excessive swelling, a solution of nitrate of silver, from one to five grains to the fluidounce of distilled water, will not unfrequently be of the greatest service. Schwartze has of late recommended the caustic treatment of profuse otorrhœa, by means of a concentrated solution of nitrate of silver.

“Perforations of the membrana tympani, whether spontaneous or artificial, will generally heal of themselves, provided the inflammation has disappeared, and the loss of substance has not been too great, in the manner described under the head of Myringitis. The cicatrization must be watched in the same way as in the case of simple inflammation of the tympanum, and supported by adequate means.

PLASTIC INFLAMMATION OF THE MUCOUS MEMBRANE OF THE  
MIDDLE EAR (OTITIS MEDIA HYPERTROPHICA).

“This disease differs greatly from the other forms of inflammation of the mucous membrane of the middle ear, as well by the symptoms presented during its course as by its consequences, since its elements of form arising from the inflammatory process bear the germ of a higher organization; whereas those products resulting from catarrhal and purulent inflammation are rapidly given up to destruction. While in the latter the mucous membrane at best simply recovers its



former properties, or, as in the purulent form, is more or less destroyed, and then supplied by cicatricial tissue, in this disease the affected organs are hypertrophied.

“From the dissolution of the newly-formed elements and the increase of the intercellular fluid in catarrhal and purulent inflammation, a more or less consistent mucus or purulent discharge will take place, which, respecting quantity generally, stands in a direct relation to its intensity. In the plastic inflammation, when occurring in its pure form, on the contrary, its fluid product consists only of increased intercellular fluid, with more or less cast off epithelial cells, and is usually but scanty, even in the severest cases. It is true that a more purulent product will occasionally occur upon the free surface, but in such cases the plastic inflammation has supervened on one of the other forms, to which the free product is owing. The disease under consideration occurs most frequently, attacking in its pure form a segment of the middle ear or its whole surface; or as is more often the case, joining from its beginning an already existing catarrhal or purulent inflammation, or the latter forms combining reversely with the plastic inflammation already existing. From the combination of these various forms, as a matter of course, many varieties of disease must arise, differing to a great extent from those of the pure form.

“A large number of those cases hitherto taken for simple chronic catarrh belong to this category. Hypertrophic otitis is owing to the same causes as the catarrhal and purulent forms. It is distinguished from them by the chronic course it generally runs. Its subjective symptoms are so inconsiderable, not only at its beginning, but after it has lasted for years, that the individual affected, and those about him, do not observe any sign of any existing disease of the ear. In other though rarer cases, on the contrary, it will be ushered in with intense and rapidly-developing objective symptoms (generally attacking more important segments), and the patient is tormented by most annoying, morbid, subjective symptoms. We shall know how to designate all these relations better after we have had an insight into the pathological processes that are going on.



"Owing to the injurious influences present, a considerable hyperæmia occurs in a segment or upon the whole of the mucous membrane of the middle ear. This is accompanied by extension, partly also by new formation of blood-vessels and increase of the intercellular fluid. The corpuscles of the connective tissue are increased by subdivision, and perhaps by other processes which cannot yet be stated with certainty. The result is a hypertrophy of the mucous membrane, corresponding to the degree of inflammation present. The epithelium being puffed up to some extent, diminishes somewhat the generally intense redness apparent in the mucous membrane after death. The tissue of the inflamed mucous membrane is generally less moist than in the catarrhal and purulent forms, and the exudation upon the free surface, arising from the protoplasm of the epithelial cells, is quite imperceptible. Sometimes the inflammation causes exuberant epithelial formations, which, undergoing further metamorphosis, constitute the basis of Cholesteatomes.\*

"In the course of this process the newly-formed elements are either equally distributed over the whole inflamed surface or are scattered in groups. They develop themselves at these foci unequally; some will wholly or partly pass into higher organizations. From the division of their processes, which have numerous connections with the equally advanced processes of neighboring cells, a variously arranged, more or less intimately interwoven, soft, connective tissue will proceed.

"This, according to the preceding arrangement of the elements of form, will either be equally distributed over the entire originally inflamed part (resulting in equal hypertrophy of the mucous membrane), or in cases where the new formations occur at circumscribed foci, granular formations on the mucous membrane originate; or in those cases where the further development is going on in a form peculiar to them, they terminate in polypi. It is to be mentioned that, during the further progress of the newly-formed corpuscles, they are frequently not confined to the particular organs in which they originated, but growing, penetrate its epithe-

---

\* Tumors composed essentially of cholesterine and stearine.



lium, if there be any yet existing. If the latter should have been previously destroyed, they sprout out the more readily, tending toward a different surface from that on which they were produced. The same occurred, consecutive to inflammation, in the larger swellings (polypi) previously mentioned. It is particularly marked in the connective filaments and membranes, originated by the process, which so frequently occur here. Especially is this the case in the cavity of the tympanum and the cells of the mastoid process, where they grow from their original seat in the most eccentric directions, connecting with others corresponding in that manner. These effect an exceeding variety of anomalous connections of the organs of the inner ear. On the other hand, we cannot conceal that during the course of the process many of the newly-formed elements undergo retrograde metamorphosis. This consists in the separation into its molecules, passing into fat and being absorbed, or, as will sometimes happen during chronic processes, undergo calcareous degeneration. Even the newly-formed blood-vessels, as well as those simply extended, may waste during the course of the process. They are reduced to simple fibrous filaments, or they may, with their contents, be converted into fat or calcareous material, the entire molecular mass being gradually removed by absorption.

“In this review we shall be enabled, *a priori*, on the one hand, to deduce all the possible symptoms, both subjective and objective; and on the other, to measure the danger to the hearing of the patient that grows out of this form of inflammation, especially at the height of the process. As has been mentioned before, the disease under consideration will be developed unperceived by the patient, unless it presents considerable intensity from its onset, or attacks particularly such segments as, for instance, the tympanum, which are so important for the conduction of sound, but especially those parts which close the round and oval openings into the cavity of the tympanum. Such processes can even cease again without ever having become known to the affected person, the discovery only being made by the post-mortem examination, showing hypertrophic spots on the mucous



membrane, otherwise harmless pseudomembranes and slight obscurations on the tympanum. The case is different when the inflammation from the outset attacks the whole mucous membrane of the middle ear or important segment in considerable intensity. Then there will be no pain, but the patient will complain of difficulty of hearing, and perhaps some subjective noises in the ear. The latter phenomena may be due to causes similar to those of catarrhal and suppurative inflammation. The same is true of the sensation of oppression and fullness in the ear, which is usually combined with more or less pronounced giddiness and depression of mental activity. This disagreeable sensation is the more vexatious to the patient, as it continues longer than in the preceding forms of otitis, in which the moments of relief are more frequent in the course of the disease, the lesion being often mitigated by accidental occurrences. If there be no complication with another disease, and the inflammation under consideration makes the progress above described, the hearing, no less than the other morbid phenomena, will become worse and worse, as soon as the further development of new elements gives rise to such changes as will more and more impede the function of important parts. If the process last long, the severest disturbances of hearing will finally result from the secondary alterations taking place in the labyrinth, owing to the deficient activity of the auditory nerve. Complete deafness, for instance, can proceed from this simple inflammatory process, unless it be counteracted by all means possible; although, on the other hand, it is a fact that this termination is one of the greatest infrequency, the hearing of the affected ear being decidedly interfered with after the close of the morbid process in but a minority of cases. In the majority of cases there remains simply permanent symptoms of a less degree.

“It is evident that these subjective phenomena just described can assume different forms when the hypertrophic inflammation of the ear appears, compared with any of the other forms of otitis. In the complication with suppurative otitis media, for instance, the subjective phenomena that are owing to the latter will become most marked, which must



decidedly alter its clinical appearances. The objective symptoms to be noticed in plastic otitis vary with the attacked part, the intensity of the process or complication with other forms, and, as a matter of course, the previous condition of the affected ear. The Eustachian tube is generally affected sympathetically from its onset. Its orifice, consequently, must thereby become closed or obliterated, and the changes in the tympanum and the ossicles, above described, consecutively take place, according to the quality of the mucous membrane of the cavity of the tympanum, and the various other affections of the tympanum itself. Upon the inspection of the external passage of the ear, different alterations will become observable, which need no further explanation, as they are the result of similar physical lesions to those in the preceding chapters on inflammation of the middle ear. Nor do the results of auscultation upon a single examination offer reliable points. They inform us, at best, about the relations of space in the inner ear. It is not till auscultation has been performed repeatedly and at different periods that we obtain information, in so far as the copious exudation occurring in other forms of inflammation, the symptoms discovered, together with the ocular examination, will vary from those of the hypertrophic inflammation. In this form of inflammation the diagnosis cannot, from its commencement, always be made with that certainty which would be desirable for the physician; only after a longer observation of the patient, or the duration of the disease, can it be obtained with certainty. All the symptoms presented will tend to increase of bulk, provided the inflammation occurs in its usual form, or to hypertrophy accompanied with other symptoms. When plastic otitis occurs in its usual form, and especially when the mucous membrane on the interior wall of the cavity of the tympanum is affected, the prognosis is in general unfavorable. With regard to the future capacity of hearing, sometimes only at its outset can the increasing hypertrophy be checked, or at best, its bad consequences be lessened by a continued careful treatment. It must, however, be remembered that hypertrophic otitis may run its course without causing any striking difficulty of hearing, when



those organs have remained intact, being of greater importance for the conduction of sound. As a matter of course, the prognosis will differ when there are complications with other morbid processes. The treatment should be to check the morbid process powerfully, and, at the same time, palliate the bad consequences of the deafness resulting from the hypertrophy. In the former we must endeavor to arrest the further development of the newly-formed elements of inflammation. If the mucous membrane of the Eustachian tube is principally affected, we should endeavor to keep it open by the application of the air douche, and the insertion of laminaria, or other bougies, as well as to promote absorption by compression, already exercised by the application of the bougie, and so forth. In cases of less violence, perhaps diffused over the mucous membrane of the cavity of the tympanum, injections of a solution of caustic potassa, gr.  $\frac{1}{2}$  to gr. i to f $\frac{3}{4}$ i of distilled water, once a day, or repeated at longer intervals, will be most serviceable, the epithelium being softened by them, and caused to be sooner cast off, after which the air douche must be employed for the evacuation of the free products. Dilute acetic acid, used in a similar mode to the caustic potassa, will be of great service. Acetic or iodic ether, or ammonia vapor, and in syphilitic cases, weak solutions of corrosive sublimate, gr.  $\frac{1}{2}$  to gr. i to f $\frac{3}{4}$ i of water, have likewise given satisfactory results. In cases of greater intensity, the bulk of the mucous membrane being considerably increased from the onset of the morbid process, a more or less concentrated solution of nitrate of silver, grs. i to v to water f $\frac{3}{4}$ i, is indicated. This solution, as well as the preceding medicaments, are to be injected by the Eustachian tube, especially by the spray apparatus of Weber, which is most suitable for this purpose. Catheterization of the Eustachian tube is performed by means of a common tube catheter passed into the cavity of the tympanum, and the spray injected by its bulb."



## CHAPTER XII.

### OTITIS MEDIA AND INTERNA INVOLVING MASTOID PROCESS AND CELLS—SQUAMOUS PORTION OF TEMPORAL BONE—CEREBRUM AND CEREBELLUM.

THE mastoid process in children before the first year is not developed, and merely consists of the horizontal portion, which is so intimately connected with the *cerebrum* that this portion of the brain is primarily affected. When, however, the child becomes the boy or girl, the *cerebellum* is the part usually involved. The most frequent cause of disease of the mastoid cells in childhood is *cold* contracted immediately after birth, followed by a discharge which, if it had been noticed, was preceded by a painful cry of the infant, or, for twenty-four hours, by a languid, drowsy state, accompanied with more or less fever, and, in some instances, brain irritation, which the child never recovers from. As the disease progresses, the tympanum or middle ear becomes filled with mucous or muco-purulent matter, which causes ulceration of the membrana tympani, and it gives way, with relief to the symptoms, and the discharge escapes, and continues for months and even years, to the detriment of the health of the child, or it *may* be cured by proper treatment. In other cases the peculiar form of the mastoid cells preventing the escape of the matter, it is retained, and causes pyæmia, and even death. In older children and in adults the primary causes are scarlet fever, measles, smallpox, scrofulous, and tubercular disease; the secondary cause in almost every case being a *catarrh*, by its direct influence upon the ear with its broken membrane, or through the nose, pharynx, or Eustachian tube, leading to the middle ear, thence by the lateral sinus into the circulation, the cerebellum, or its membranes.

The writer will report a number of cases, most of them



from his private practice, and watched carefully at their homes, with the diagnosis and successful treatment by operative procedure :

CASE I. *Disease of the mastoid process ; no discharge from external meatus ; recovery.*—Martin F., aged seven years, a large and robust boy, but of a tuberculous family (father and four uncles having died of phthisis), was convalescent from scarlet fever, and was discharged well on March 15th, 1858. Was called in haste to see the boy on the 27th. There was great pain and swelling behind the ear, over the mastoid process, but no discharge from the external meatus. He was freely leeches, and purgatives administered, followed by anodynes to relieve the pain. On the 28th, the swelling was on the increase, and extended to the face and eyes, with fever and symptoms of convulsions. It was then proposed to cut down to the bone dividing the periosteum, as the only means of relief, to which the mother consented. An incision was made about an inch long behind the ear, and as nearly as possible parallel with the concha ; as the ear at this time was almost horizontal, a profuse gush of blood followed, mixed with imperfectly-formed pus ; the wound bled for three days. The pain was much relieved, and, by the use of saline mixture with sulphate of morphia, he was able to sleep, which he had not done since the 27th. A poultice was applied on the third night, and by the fourth, pus flowed freely, which was encouraged ; and by the end of the sixth week the wound was disposed to close ; this was prevented by a blister. The discharge continued for four weeks longer, when the wound was allowed to close, the roughness of the bone having disappeared.

Report January, 1871. The young man is now twenty years of age, and has enjoyed good health since, being able to be out in all weather ; is bright and intelligent, and is at work in a cotton-mill ; his hearing in the left ear impaired ; right, very good. Over the mastoid cells there is a depression of a bluish color, from loss of bone. Has never had a severe attack since, excepting now and then a slight discharge from one (deaf) ear, and he is an expert mechanic. Has since lost his mother by phthisis.



In the same neighborhood the writer examined a boy, James R., aged five years, in 1862, who had a similar swelling after an attack of scarlet fever; it opened of itself after long poulticing, and continued to discharge from the back of the ear for twelve months; it then ceased, leaving a deep depression behind the ear with loss of hearing; the boy is imbecile and cannot articulate.

CASE II. *Disease of the mastoid process; perforation of the membrana tympani; recovery.*—Mrs. A., aged thirty-five years, whose case was of the same character as the one just reported, was treated in a similar manner with equally good results.

CASES III., IV. *Abscesses over the mastoid process in mother and child; similarly treated, and both recovered.*—March 25th, 1869, Thomas N., aged five years, applied at Howard Hospital with an abscess over the mastoid process. He is convalescent from an attack of measles. The swelling back of the ear commenced three weeks previously, then subsided, and again began to swell. On examination, there was swelling, redness, and feeble fluctuation. There was no discharge from the ear. On informing the mother what was to be done, she replied, "Well, do what you think is right, as you performed the same operation on my ear when this child was only ten months old. I was at that time three months under your care, and was cured, and remain so." The writer examined the back of her ear and found a deep depression, where there had been a loss of bone. Her hearing in that ear was not perfect, but she was not *very* deaf. The operation was then performed on the boy, when, by the aid of the probe, the bone was found to be denuded of its periosteum. Being of a strumous habit, he was ordered syrup of the iodide of iron, a small poultice of ground flaxseed, and subsequently an ointment of the red oxide of mercury, to dress the part and keep it open.

April 1st. The wound had suppurated freely, but the opening was disposed to close; it was accordingly enlarged, and the previous treatment continued, with good diet and exercise in the open air. By the end of the month he reported himself well, wound healed, and all swelling dis-



appeared. Many other cases of this same class the author has seen while attending his clinic during the last seventeen years.

CASE V. *Perforation of the mastoid process; otorrhœa; removal of necrosed bone; recovery.*—Mary R., aged eight years, a robust-looking child, came under my care early in September, 1861, at the Howard Hospital. It was reported that she had suffered from scarlet fever of a most malignant type, having been in a state of coma for several days. This gradually passed away, when the throat and ears became affected, and a long period elapsed before complete convalescence took place.

*Present condition.*—She is deaf in the right ear, with a constant discharge of offensive pus, etc.; on washing out the parts, the meatus was found to have a white, soft deposit on its surface, with a granular condition near the membrana tympani, with a perforation of three-fourths its size.

A mild wash to be applied of nitrate of silver in solution. The parts kept clean by repeated injections of tepid water. Counter-irritation was to be kept up in front of the ear, and air was to be passed through to keep the Eustachian tube open and force any accumulation of pus out from the middle ear. A guarded prognosis was given as to the ultimate result. Having improved considerably, she ceased her attendance, and the writer saw nothing of her until called in haste to see her on September 21st. The history received from her father, an intelligent man, was as follows: The Sunday previous being a very hot day, the child was sleeping on a sofa, when the father, to cool the house, opened both the front and back doors, which produced a strong draught of cool air upon the sleeping child. After retiring, she was attacked in the middle of the night with intense pain in her ear, causing her to scream and preventing all sleep. The parents applied a blister and used other means, but the relief was of short duration. When the writer was called to her she was suffering intense pain; fever, with alternating chill; pulse, one hundred and thirty; there was swelling over the mastoid region, involving the side of the face and eye.



*Treatment.*—Believing that pus was formed in the mastoid cells and was endeavoring to make its way outward, requiring only an outlet, the writer divided the skin freely, muscle and pericranium, down to the bone. On withdrawing his knife, blood flowed very freely, mixed with imperfectly-formed pus. A poultice was ordered, to be repeated every few hours. A saline mixture of citrate of potassa, containing sulphate of morphia, was directed, to relieve pain.

September 22d, 23d, 24th–30th. Better. Discharging pus from the opening, which being not quite free enough, applied powdered red oxide of mercury to increase the discharge.

November. During this month, visited the case every few days. The opening being disposed to close, introduced a sharp hollow steel probe, so as to perforate the surface of the bone, and applied to it freely the solid nitrate of silver, which increased the discharge.

December. In the early part of this month, the opening was again enlarged, and a piece of bone discharged. On the 19th, a larger piece of carious bone was found by the probe to be movable, and by enlarging the wound, it was removed with some difficulty, the bleeding being very free. A few days after this operation the discharge ceased, and the wound healed. There is a deep depression behind the ear from loss of bone. The membrana tympani has closed somewhat; the child is deaf in that ear, but otherwise well, and continued so for several years. The bone measures six-tenths of an inch in length and three-tenths of an inch in width. It is now in the writer's collection, and has been examined by numerous distinguished surgeons.

In the cases above related, which might have been increased by many others, we have examples of three of the principal forms of ear disease in the mastoid region, which will frequently be met with by any one who devotes much attention to the ear, and occasionally by those who pay but little attention to this department. The writer trusts they will be found useful as guides to diagnosis and treatment.

In the first case, the simple division of the periosteum, with the subsequent application of the blister, was all that was necessary to complete the cure.



In the second case, it required the second enlargement of the opening and breaking down of the bone by the application of nitrate of silver.

The third class of cases was of greater danger to the patient, and it involved a larger number of the cells of the mastoid process. It therefore required a free perforation of the bone, and its removal in a diseased state.

This was the first operation (December, 1861, and published February, 1862) of the kind that had been performed in the United States (so far as the writer is aware), and he has repeated the operation several times since, as, in a case the result of a severe fall, foreign body in the Eustachian tube, etc. It had been performed only eight times in Europe to the above date (and from 1862 to 1871), by Schwartze, Pagenstecher, Hinton (London), Jacoby, Agnew, and Roosa, of New York.\* “Köppe† and Schwartze report a case of epilepsy caused by retained pus in the middle ear, which was relieved by perforation of the mastoid.” In 1871, by Dr. F. Buszard‡ (London). Two successful cases are reported in the London *Lancet* for April 29, 1871, of abscesses in the mastoid process, with perforation of the bone, under the care of Mr. Barwell, of Charing Cross Hospital.

CASE I.—E. F. was only six years of age. History: severe pain in the head, which had the effect of sometimes driving her “half mad,” and at other times of producing stupefaction. Two operations were necessary, the second by a probe, and the T-shaped incision was maintained open by a nail-shaped plug of gentian root until the thirty-sixth day after the first operation.

The second case was of greater interest, on account of the patient having reached the comparative mature age of forty-seven, and as the premonitory symptoms were accompanied by profuse epistaxis.

Now it is the recognized operation in properly-selected

---

\* The Mastoid Process in Inflammatory Affections of the Middle Ear, by D. B. St. John Roosa, M.D., Medical Record, Dec. 1st, 1870.

† Archiv für Ohrenheilkunde, Bd. v. p. 282; Roosa, op. cit.

‡ British Medical Journal, January 28th, 1871, p. 88, and Medical Times, Philadelphia, April 15th, 1871.



cases. It was, perhaps, too frequently and indiscriminately performed at first, and afterwards entirely rejected, and considered illegitimate, although there can be no doubt that cases occur in practice (such as the writer has given) in which the operation ought to be performed in obedience to the fundamental principle that every accumulation of pus in the tissues, especially when in contact with bone, is to be evacuated as soon and as completely as possible, in order to prevent the noxious effects upon neighboring parts and a further spreading of the inflammation.

The tissues adjoining the mastoid process, viz., the dura mater, the internal carotid artery, and internal jugular vein, with the transverse and lateral sinus, are so important, that a morbid process which spreads to them from the middle ear is fraught with danger to life, independently of the circumstance that the sense of hearing may be weakened, or even entirely annihilated. The cells of the mastoid process are in open communication with the cavity of the tympanum; they are covered with the same mucous membrane, and participate in all the affections of that cavity. Pus accumulates the more readily in these cells, as their communication among each other, and with the anterior part of the middle ear, is often very narrow, so that the pus is easily shut in; besides which, a large portion of the mastoid process lies deeper than the membrana tympani and the meatus auditorius. The membrana tympani may, therefore, be perforated, and the secretion of the cavity of the tympanum be allowed to escape freely outward; yet this will not provide an outlet for the pus within the mastoid process, as has been shown in a case before related. If the matter accumulated there does not spontaneously make its way outside by the formation of a fistula behind the ear, which sometimes occurs, the surgeon has no choice but to break through the external layer of bone, in order to gain access to the seat of the pus. The introduction of warm vapors into the Eustachian tube has no decided beneficial effect in severe cases, although, in mild forms of this affection, frequent gargling and repeated blowing of air into the tube, by means of a catheter, or an attachment to Politzer's bag, may serve to prevent the case from



getting worse, and may act favorably altogether. If the bony layer which has to be perforated be very thin and fragile, an ordinary buttoned probe will answer the purpose (or a hollow steel director or probe); but if such be not the case, a hollow chisel ought to be used, and, in every instance, an incision, about an inch long, behind the ear, and parallel with the concha, ought to precede the perforation of the bone, as first recommended by Dr. Wilde, of Dublin.\* This incision should be made, even when "an indistinct sense of fluctuation can be discovered, at least an inch long in the periosteum (and this alone will sometimes be all that is necessary). It is generally required parallel with, and about three-quarters of an inch from, the attachment of the auricle, in order to avoid the posterior aural artery, which, when divided, bleeds profusely. The knife should be drawn from below upward, and, from the swollen state of the parts, the depth to which we are obliged to introduce the instrument is often nearly an inch. The hemorrhage, unless we wish to extract blood, may be arrested by placing a dossil of lint within the incision," saturated with the following styptic of Prof. Paucoast:

R.—Saponis (Castile),  $\mathfrak{z}\text{i}$ ;  
Potassæ Carb.  $\mathfrak{z}\text{ij}$ ;  
Spts. Vin. Rect.  $\mathfrak{f}\mathfrak{z}\text{ij}$ . M.

If further relief be required, the chisel, director, or hollow probe should then be inserted in the middle of the incision, at an equal height with the opening of the ear, and ought to be carried horizontally, and a little forward. In this way the dura mater and transverse sinus are avoided, and the surgeon rapidly arrives in the large cells which are close above and behind the cavity of the tympanum. The instrument must, of course, be used very gently, and with interruptions in some cases for a considerable time, so as to avoid a sudden piercing of the bone.

The first proposal to perform this operation was made by Johannes Riolanus (1649) and Rollfuik (1656), who recommended it for deafness and noises in the ear, caused by ob-

---

\* The Nature and Treatment of Diseases of the Ear, p. 233, Am. ed., Phila., 1853.



struction of the Eustachian tube. Valsalva, in 1704, made injections through a fistulous opening which existed behind the ear; and J. L. Petit was the first to perform perforation of the mastoid process by means of a hollow chisel and hammer. The life of the patient was saved thereby. Dr. Von Tröltsch,\* the distinguished author and Professor of Aural Surgery of the University of Wurzburg, Bavaria, reports eight cases, all of which terminated successfully; so that the operation, if properly and carefully performed on selected cases, is not a dangerous one.

The second case occurred in the practice of Jesser (1776); his patient was a soldier, who had for several years suffered from pain in the ear, with suppuration and deafness. A slight prominence having been formed behind the ear, Jesser made an incision one inch long, after which very little pus came away, and it was only when he had penetrated with the probe into the cells of the mastoid process that much pus was evacuated. The pain in the ear then ceased at once, and never returned, and the hearing was also improved. The mastoid process, on the other side, was also pierced in this patient, with a trocar, for deafness, not for pain, and the hearing improved. The operation was now called "Jesser's operation."

The third case is that of Fielitz. A young girl lost her hearing on one side, in consequence of an acute disease, and suffered for five years, at intervals, from an offensive discharge from the ear, which was always preceded by fever and severe pain in the ear. The mastoid process was pierced with a pointed instrument, and injections were made, after which much pus escaped. The discharge gradually ceased, and the hearing became completely re-established.

The fourth case is that of a patient under the care of Weber (1824). A peasant had suffered for three months from a very painful suppurative otitis, and as an incision behind the ear did not mitigate the pain, a trocar was thrust, three lines deep, and obliquely forward, into the highest prominence of the mastoid process, ten lines behind

---

\* Medical Times and Gazette, Sept. 26th, 1861 (London).



the ear, seventeen lines from the upper, and four lines from the posterior, edge of the mastoid process. Much pus came away, and the pain disappeared immediately; the suppuration ceased four weeks afterwards, and a complete cure was the result.

The fifth case is that of Forget (1849). A young man became affected with otitis in consequence of a cold; copious suppuration in the ear set in, accompanied by paralysis of the facial nerve, but the latter disappeared a year afterwards. Two years after the commencement of the illness, the patient became completely deaf on this side, and the neighborhood of the ear was greatly tumefied. The mastoid process was then pierced by means of a hollow chisel, and much pus escaped. A month afterwards, a sequestered bone, three centimetres high and two centimetres thick, was removed, and from that time the patient began to improve, the suppuration ceased, and the hearing returned.

The sixth and seventh cases occurred in the practice of M. Follin, of Paris (1859). In both cases there was a fistulous opening, which was dilated by that surgeon by means of the trephine; six weeks after, the cure was complete.

The eighth case is one of Dr. Von Tröltsch (1860). A girl, aged ten years, complained, the second day after the eruption of scarlet fever, of violent pain in both ears, which continued to increase until the fifth day, and then began to diminish, in consequence of a discharge from both ears having commenced. The fever took its regular course. The discharge from the ears continued, and after some time the pain became again severe, especially behind the right ear, where a slight swelling was also seen. Three weeks after the commencement of the illness the pain became most intense, and was accompanied with headache and deafness. Pressure on the mastoid process augmented the pain; the skin on the right side was slightly reddened, and the whole neighborhood behind the ear and the upper and lateral region of the neck were diffusely swollen. Both meatuses were filled with thin pus, after the removal of which their walls appeared to be tumefied, and, in the background, a drop of water was pulsating, an evident sign that both membranæ tympani were



perforated. Injections of lukewarm water were then made four times a day, and after the injection, a weak solution of acetate of lead (five drops to the ounce) was dropped in, while mercurial ointment was rubbed in and around, and chiefly behind the ear. During the next few days the discharge was not lessened, and the pain very much increased; the swelling behind the ear became more considerable, was very painful on pressure, and felt œdematous; no fluctuation was perceived.

Six days afterwards, an incision from one and a half to two inches in length was made behind the insertion of the concha, and parallel with it down to the bone. The bleeding was profuse, and pus appeared immediately afterwards in the level of the external opening of the ear. In the evening after the operation, violent pain was felt in the wound, but no longer in the ear; there was no fever. The swelling behind the ear became less on the following day, but again increased on the second day, when the pain also became more violent and the discharge less abundant; on pressure behind the ear, pus escaped from the meatus. Two days afterwards, Dr. Von Tröltsch tried to penetrate the incised wound, which had already become narrowed, by means of a buttoned probe. In this he succeeded only for three lines, and he therefore continued perforating with the probe, and afterwards placed a piece of charpie as deep as possible in the opening. Soon after, much pus, at first thin and afterwards thick, escaped from the opening, while only a trifling quantity was discharged from the meatus. On the following day the patient felt better; the pain had entirely disappeared, and the hearing was improved. Injections were now made daily; the probe could penetrate an inch deep, and the injected water escaped through the meatus. About two months after, all discharge had ceased and the recovery was complete. The girl died eighteen months afterwards, of tuberculosis of the lungs. The post-mortem examination showed the following condition of the organ of hearing:—On the right mastoid process a cicatrix was seen, which corresponded to the incision, and from this proceeded a canal in the mastoid process, with smooth walls, in the form of a cone,



with its base behind, passing from behind forward, four millimetres long, and closed inside. The external meatus was not pathologically changed. The mucous membrane of the tube was pale and covered with a little mucus; the narrowest part of the tube was abnormally dilated; its osseous part was cylindrical and also dilated. The posterior half of the membrana tympani was wanting, except a small strip behind and above. The anterior part of the hole was drawn inward, and grown together with the opposite wall of the cavity of the tympanum. The manubrium mallei was drawn inward, and firmly soldered to the wall of the labyrinth. The cavity of the tympanum, therefore, consisted, properly speaking, of two parts, the anterior of which, lying before the manubrium, was connected with the tube, and the posterior, with the meatus. There was no communication between these two parts, so that water injected through the meatus did not penetrate into the tube. The posterior half of the cavity of the tympanum, which communicated with the meatus, was also completely closed, posteriorly and anteriorly, towards the mastoid process. The stapes was grown together with the upper part of the perforation, but it was movable, and its joint with the crus of the incus was not severed. The malleus and incus were also movable. Between the head of the malleus and the ostium tympanicum of the tube, there was a firm and strong band, horizontally situated, which contained the tendon of the tensor-tympani muscle. This band connected the inner and the outer wall of the cavity of the tympanicum, and communicated with the exudations separating the cavity of the tympanum in two halves. At the left side the meatus did not contain any abnormal secretion. The upper part of the tube was abnormally dilated. The posterior half of the membrana tympani, close to the manubrium of the malleus, showed an indented perforation of the size of a grain of hemp, the posterior edge of which was grown together with the opposite wall of the cavity of the tympanum, and with the capitulum of the stapes.

A series of rather succulent adhesions connected the upper and anterior part of the membrana tympani with the tendon



of the tensor tympani muscle and the opposite wall of the labyrinth. A thin but firm adhesion was seen backward. The mucous membrane of the cavity of the tympanum was thicker, softer, and contained more blood than usual. All the ossicles were freely movable. The cells of the mastoid process were fine, its meshes filled with a reddish-yellow liquid, which only contained a few small nucleated cells and exudation corpuscles. The history of this case gives another striking proof of the fact that very considerable changes may take place in the membrana tympani, and, nevertheless, the hearing of the individual concerned may not be seriously affected.

CASE I. *Of otitis interna of author, followed by death, and a partial post-mortem examination.*—Mary H., aged six years; had scarlet fever when three years of age; was feeble a long time, but her ear was not treated except to keep it clean by washing. About May, 1857, the child began to complain very much, and came to my office. Ordered a small blister, and a solution of one grain of sulphate of zinc to the ounce of water as a wash. Internally, as a tonic, half a grain of sulphate of quinia three times a day. On examination, the right membrana tympani was found to be entirely gone, and there was considerable discharge of unhealthy pus from the ear, indicating caries, or disease of the bones. Having been benefited by the treatment, she did not return to my office until June 6th, when she was suffering intense pain from inflammation of the middle ear, the result of exposure, the discharge being much increased. Being a very willful child, her parents had but little control over her. I directed counter-irritation to be renewed, with the internal use of an opiate to relieve the pain; did not use leeches on account of her feeble condition.

Visited her on the 8th. Pain still very persistent, and head bent to the side of the affected ear. Fearing convulsions, I ordered four leeches to the back of the ear, over the mastoid process, with sedatives internally, opiate fomentations to the ear, with warm foot-baths. In spite of treatment she continued to grow worse, and had considerable discharge of pus from the nose, and was unable to swallow for a day or



two prior to June 26th, when she had a severe attack of convulsions. This was counteracted by the employment of leeches and cold applications, but was soon followed by a state of coma, and she died on the 28th of June. The family being unwilling to allow an examination, the writer was only permitted to remove the temporal bone and ear, which he did by sawing a V-shaped piece, and removing the entire ear. In spite of her anæmic condition, the coverings of the brain were much congested and thickened, with effusion of fluid into the ventricles, and considerable softening of the substance of the cerebellum.

*Eur.*—The membrana tympani was almost gone, yet, strange to state, in the middle ear, although filled up with greenish pus, and the membrane soft and detached, was found the malleus and incus, which he has retained. On cleaning these bones, however, they were found to be ulcerated. The semicircular canals and the cochlea were almost free from disease, which had passed from the middle ear to the lateral sinus and membranes of the brain, causing a low form of inflammation, with softening and effusion, ending in convulsion and death.

What seemed very remarkable in this case was the long period that the small bones remained in the ear, notwithstanding the discharge, showing how slowly the ligaments connecting the bones ulcerate.

CASE II. *Case of chronic otitis interna with acute attack extending to the brain, followed by death; with post-mortem examination, etc.*—Mary E. C., aged fifteen years and eight months, about the year 1866 suffered from an acute attack of otitis externa from cold, involving the membrana tympani. She applied as a patient under the writer's care at Howard Hospital. She was treated by the application of tincture of iodine over the mastoid process, with active purgatives. When the acute symptoms disappeared, the cleansing wash of warm water was changed to a mild astringent one, and she was put on the internal use of the syrup of the iodide of iron. After a few visits she ceased her attendance. She applied at the hospital again on the 14th of December, 1868, with an acute attack of otitis media and interna; the pain was over the



occipital region. The history given by her mother, an intelligent but feeble woman, was as follows: That the girl had recovered from her former attack with some loss of hearing, with now and then some pain and slight discharge from the ear. She was, however, affected at times with a distressing feeling of giddiness, and had to give up part of her occupation of pressing goods, requiring stooping; and at times this feeling was so strong that she was sure, had she not taken hold of some object to support herself, she would have fallen.

At the age of *fourteen* she began to menstruate, and always after the flow came very freely, even more in quantity than her mother at her age; this, however, brought no relief to the symptoms of giddiness, and yet she was bright, cheerful, and her general health seemed otherwise good until her attack.

As the external meatus was full of pus and mucus, she was directed to have warm water to wash the parts, with solution argenti nitratis, applied to diminish the discharge, and leeches at the lower part of the auricle, with foot-baths and anodynes to relieve pain and procure sleep. After the acute symptoms had disappeared, blisters dressed with morphia were freely employed; and, to affect the brain symptoms, large doses of bromide and iodide of potassium. Most of these remedies were rejected by the stomach, as there was constant violent vomiting, against which numerous agents were employed, none of them procuring much relief except small pieces of ice. There was a tendency to constipation, which was relieved chiefly by injections of warm water. The chief point of pain was from the top of the spinal column to the occipital bone. Her pupils were generally contracted, with defective vision; pulse varying from 70 to 90. During the last two weeks the eyes became injected, and unless under the influence of atropia, she was not able to see at all distinctly.

January 26, Prof. James A. Meigs visited the case with me in consultation, and confirmed my diagnosis that an abscess or abscesses had formed on or in the brain. He recommended the use of morphia to relieve pain, and to continue the use of the bromide of potassium and blistering. The ear




still continued to discharge pus and mucus, and only once blood, and a small piece of bone, which was lost before there was an opportunity of seeing it. Her head was kept on a pillow, bent forward on her chest, and if it was moved it gave her great pain; and in one instance, when moved to change an under-garment, it produced such a severe convulsive movement that it was thought she was dead. There was slight internal strabismus, with a tendency in the tongue to be protruded to the left side. She discharged a considerable amount of mucus, etc. by the mouth and nose, but never had a cough or pain in the chest. She swallowed without difficulty; could move her arms and limbs; her tongue was moist until two days before her death, when it became almost black. She died February 15th, 1869, without a struggle, singing a few hours before she passed away. She was deaf in the left ear, with perforation of the membrana tympani, except after a free discharge of pus or blood.

Post-mortem forty-eight hours after death, assisted by Drs. James Collins, J. S. Cohen, and C. S. Turnbull.

The incision was made through skin, fascia, and muscles down to the bone on a line with the ear, and the scalp dissected back to the occiput, and forward, so as to expose the temporal bone, cutting close to the bone and zygomatic process; the cranium was sawn through posterior to the coronal suture, so as to prevent deformity. A second incision was made just above the temporal bone back to the occipital

Fig. 60.




bone, and then the temporal bone sawn down to the base, and by means of the chisel the bone was separated so as to obtain the bony portion of the ear entire to examine it. The brain was then removed, cutting carefully into its membranes, which were slightly congested; veins very turgid; surface of brain congested;

effusion in ventricle, containing three ounces of serum, and vessels much distended, but no abscess in brain or on its sur-



face; slight effusion in sheath of optic nerve. The cerebellum was then removed; it was softened, and the left lobe, situated at B (Fig. 61), when cut into, was found to contain a sac (Fig. 60) of pyogenic membrane containing about two ounces of green pus, which did not separate when allowed to stand. When examined by the microscope there was seen true pus globules. There was thickening of the membranes at this point, and a necrosis of the bone opposite it into the middle ear, as seen in the drawing (Fig. 61).

Fig. 61 —EXAMINATION OF THE BONE REMOVED.



The temporal bone was deficient in size, slightly necrosed, with a small opening at A, through the sinus into the cerebellum on the one side, and into the middle ear on the other. This sinus was filled with dark-colored coagula.

The lining of the whole of the external meatus was soft, elevated, and of a dark color; its surface was denuded of lining membrane beneath its purulent matter. There was but a small rim of membrana tympani, and a part of one of the small bones. Upon filing the bone down to the semicircular canals, it was found that they were also filled with this pus, etc. By making a section with a very delicate saw through the horizontal portion of these cavities, the vestibule and



cochlea were exposed. A portion of the membranous labyrinth was removed and examined, and it was found to contain pus globules. The nerve, as it passed from the external ear to the brain, the portio dura (C), was healthy, but the portio mollis of the seventh pair, the auditory proper, was diseased.

A portion of the brain, under the microscope, was normal in structure, but the cerebellum had lost its beautiful arbor vitæ-like character, its fibres being broken and softened.

This class of ear cases, according to Toynbee, are remarkable and rare, for there is evidence to prove that the disease originated elsewhere; and it is apparent, from the direct communication by blood-vessels between the membranous meatus and the bone forming the lateral sinus, that disease could be easily transferred from the inflamed meatus to the bone beneath. Most of the instances which we have seen, and are also recorded, are cases advancing from the tympanic cavity, or mastoid cells, outward to the meatus, in which process the cerebellum and lateral sinus were implicated.

This affection is termed by Toynbee chronic catarrh of the dermoid meatus and the mucous membrane of the tympanum. We prefer the name at the head of this article, as our dissection proves the disease of the bone, as may be seen by looking at the bone removed.

In a series of dissections of fifteen cases of fatal otorrhœa, Von Tröltsch\* has added some valuable contributions to our knowledge of pathology, and of the course by which these cases become fatal. An abstract of seven of the cases is here given.

In a case of purulent inflammation of one year's duration with polypus, caries of the labyrinth wall of the tympanic cavity had resulted in a fistulous communication through that wall just above the fenestra ovalis, through which the purulent matter had passed into the vestibule and semicircular canals, thence through the internal cribriform plate along the meatus internus to the brain, causing a fatal meningitis of the base of the brain.

In another case a polypus from the mucous membrane of

---

\* Archiv für Ohrenheilkunde, Bd. iv. Heft ii., 1869.



the small cavity, a part of the tympanum above the deeper portions of the external meatus, had projected through the bone into the meatus, so that it could easily have been mistaken for a polypus of the meatus.

A third case offers the peculiarity of multiple unconnected abscesses in the brain, one on the side of the diseased ear, and one on the opposite side; the lateral ventricle on that side being also filled with purulent matter.

In the sixth case a purulent inflammation of right tympanum, *without a perforation of membrana tympani*, led to a perforation of the roof of the tympanic cavity, phlebitis of the vena mastoidea and transverse sinus with metastasis in the lungs. The membrana tympani was found much thickened, and had thus resisted rupture.

In the seventh case a purulent inflammation of middle ear had destroyed the stapes with the membrane of the fenestra ovalis, and through the vestibule caused a pachymeningitis with two abscesses in brain and a phlebitis of the superior petrosal sinus. The phlebitis of this small sinus is explained by the small blood-vessel, which runs inward towards the mastoid process under the superior semicircular canal, entering the bone on the inner surface of the petrous portion, above and between the orifices of the meatus internus and the aqueductus vestibuli. This vessel was first described by Odenius,\* and in two cases he was able to follow it into the superior petrosal sinus, so that we have here the circulation of the middle ear in new direct communication with this sinus.

The following Case III. is of periostitis of the mastoid portion terminating in caries and death, as may be seen in Figs. 62 and 63, taken from a post-mortem specimen in the author's cabinet.

The caries advanced internally, involving the fibrous and osseous portion of the temporal bone, and terminated fatally in abscess of the brain. In such cases, when we cannot see the swelling, and the discharge is from the meatus, of a thick yellow pus, alternating with a thin and watery fluid, and there has been a sudden exposure, followed by intense head-


---

\* Medicinsk. Archiv, Bd. iii. No. 4.



ache, obstinate constipation, pain in the back of the ear, chill with delirium, our suspicions ought to be aroused. The dura

Fig. 62 —ANTERIOR VIEW OF TEMPORAL BONE.



mater is either inflamed or very much irritated; and active measures should be taken to arrest the progress of the mis-

Fig. 63.—POSTERIOR VIEW OF TEMPORAL BONE.



chief by bleeding, active purgation, and revulsives to the feet, in blisters to the mastoid process, etc.

The meatus should not be stopped with cotton, nor should



astrigent injections be resorted to until all the inflammatory action has subsided and passed away. If these active measures are neglected, we find death soon follow; whilst post-mortem examination reveals pus in the mastoid cells, and internal ear, and even portions of the temporal bone destroyed.

The subjoined cases\* exhibit how rapidly grave results may follow from caries of the temporal bone, a result of neglected otorrhœa.

“Dr. E. H. Bennet showed to the Pathological Society of Dublin two specimens illustrative of the fatal results of caries of the temporal bone. Both patients were women, and had, previously to their fatal illnesses, enjoyed good health, with the exception of occasional slight otorrhœa. The first patient, aged forty years, became suddenly excited, violent headache set in, and in two or three days she became comatose, and never rallied. After death, appearances similar to those in fatal cases of cerebro-spinal arachnitis were noticed. An extensive decomposition of lymph had taken place in the pia mater beneath the arachnoid membrane. The lymph lay thickest close to the origin of the seventh nerve of the affected side; and around its insertion into the medulla oblongata, a perceptible induration existed. The lateral sinus was healthy, but the entire petrous portion of the temporal bone was softened and red. The *portio mollis* was adherent in the internal meatus, and the tympanum was destroyed. These latter lesions were evidently of an old date. The inflammatory action in the fatal attack had probably originated in the seat of the primary disease, and spread thence inward along the course of the *portio mollis*. The second patient, aged thirty-five, had been attacked with severe pain in the head. Sudden symptoms of general pyrexia followed, and shortly afterwards complete paralysis of the *portio dura* on the affected side. In this instance, death was clearly due to pyæmia, pleuritis and general pulmonic congestion having set in before death. Here, signs of menin-

---

\* British Medical Journal, December 24th, 1870; Medical News, Philadelphia, February, 1871.



gitis were wanting; for, though the brain was somewhat congested, no purulent deposits had occurred. An encysted abscess was found in the temporal bone, and the petrosal and cavernous sinuses were full of purulent matter. Deposits of pus were discovered in the lungs. A remarkable symptom which had been present during life, was explained by the condition of the cavernous sinuses. This was the extreme distention of both conjunctivæ with serous fluid, the result of passive congestion in the ophthalmic veins. The nerve trunk of the portio dura had apparently been destroyed shortly before death."



## CHAPTER XIII.

### SUBACUTE, COMPLICATED, AND CHRONIC INFLAMMATION OF THE CAVITY OF THE TYMPANUM—CATARRH OF THE MIDDLE EAR (CATARRHAL DEAFNESS).

OF the affections of the ear, none occur more frequently, or require more care in diagnosis and treatment, than inflammation of the middle ear. It assumes three forms,—simple, complicated, and chronic. The simple form is a sequela of acute catarrh in children and adults; also bronchitis, pneumonia, etc. There is evidently swelling of the mucous membrane lining the middle ear, with or without a secretion of mucus or serum, and a discharge of mucus from the nose, with loss of hearing and a feeling of stuffing of the ears, as though a cloth had been placed over them; also an itching sensation, which the patient endeavors to relieve by pulling the ear upward and outward, producing more or less ringing, likened by the patients to the singing of a boiling kettle, puffing of a locomotive, and other similar sounds.

The objective symptoms are almost always a change in the secretion of the meatus auditorius to a yellow or a brownish-yellow. The cerumen assumes a darker tint, becoming almost black; and instead of remaining in thin layers on the sides of the meatus, runs together, forming a compact mass, covering almost, if not entirely, the membrana tympani. The following cases will illustrate the simple, subacute, complicated, and chronic forms. Detailed in this manner, we can best make it plain to our readers.


CASE I. *Simple inflammation of the middle ear; catarrhal deafness following bronchitis.*—Miss L. T., aged eighteen, of New York, while convalescent from a severe attack of acute bronchitis, about the end of the second week gradually became deaf, with roaring in the ears, and had to be spoken to



in a very loud tone of voice. She complained of a sensation of stuffing in the ears and nose, and heaviness of the head. On examination, the meatus was found covered with large scales of epithelium. The cerumen was very dark colored, and a delicate pellicle covered the membrana tympani. The nose and pharynx were swollen and red. On testing her hearing by my watch, she could only hear it with the right ear when in close contact, and with the left at a distance of one inch.

*Treatment.*—A wash of bicarbonate of soda applied warm to the meatus, and a gargle of the same. Internal use of ten grains of chloride of ammonium every two hours in a mixture with sugar and water. The ears were washed out for several days, until all the brown cerumen and epithelium were removed. A solution of sulphate of zinc, one grain to the ounce of water, was afterwards injected to restore tone. This treatment improved the hearing, so that the watch could be heard at ten inches from the right and twelve inches from the left ear. Still, however, the sounds continued, and there was considerable discharge of mucus from the nostrils. Thudichum's douche was then employed, first with a solution of common salt and warm water, and subsequently with a solution of alum, which diminished the discharge after having been used seven or eight days. The hearing, however, was not restored. Politzer's air douche was then tried, first with

Fig. 64.



air, subsequently with ether, and a mixture of ether and chloroform by means of Dr. Pomeroy's glass attachment (see Fig. 64). Five applications of this restored the hearing almost to its normal standard.

A recent letter from her medical attendant states that she continues to hear almost as well as before her attack.

CASE II. *Subacute aural catarrh, the result of an ordinary cold, which was increased by crossing the Atlantic; deafness of one ear for five weeks.*—Rev. B. C. D., aged fifty-five, Honolulu, sent to me by Dr. Goodell. While in England contracted a severe cold, which ran its ordinary course, with the exception of leaving him very deaf, first in both ears, subsequently



the right. On examination, the same conditions were found as in the case before related. In the right ear he could hear my watch at about two inches distance. On closing the left ear, he had to be spoken to in a very loud voice, and it was with difficulty that he could understand. On the left, although the circumstances were the same, he could hear ordinary conversation, and my watch at a distance of thirty-six inches. After the same course of treatment with the right ear, the left ear meanwhile being closed, the watch could be heard at twelve inches, and conversation at twenty feet. As there was considerable roaring in the left, he was supplied with a Politzer's bag and inhaler, with tincture of iodine, to be employed until these symptoms had passed away.

CASE III. *Subacute aural catarrh of several months' duration; deafness unfits the patient for the performance of his duties.*—August 16th, 1870. Rev. M. W. C., aged thirty-six, born in Pennsylvania. While a pupil at the Divinity School, in this city, contracted a severe catarrh, with cough and general impairment of health. Has been under the care of Dr. H. Y. Evans, of this city; his general health had improved by the use of extract of nux vomica and iron; but he is so deaf as to interfere with his studies. Dr. E., therefore, sent him to be relieved.

*Symptoms.*—Left ear has a feeling of fullness, with great discomfort; watch heard only at three inches; in the right, could hear it at three feet; meatus normal; left meatus injected and swollen; membrana tympani congested along the handle of the malleus; light spot not well seen; left eye injected; left Eustachian tube contained mucus; right free.

Employed Politzer's method of inflation, with some relief to the feeling of fullness. Internal treatment, a mixture containing hydrochlorate of ammonia, ten grains, three times a day.

August 18th. Again employed the air douche. The hearing distance for the watch increased to six inches. As there was a nasal tone, with enlarged glands, Thudichum's nasal speculum was applied, and powdered alum was blown up the nostrils; the throat was penciled with glycerine and iodine, also gargled with alum and borax.

August 21st. Treatment has removed the nasal voice,



and by the use of the douche the hearing has improved some fifteen inches; the patient desired to go to the country, and continue his treatment.

February, 1871. He was perfectly cured, and now performs his usual duties as a clergyman of a parish in Pennsylvania.

#### AURAL CATARRH AND ITS COMPLICATIONS.

The first complication with which we are apt to meet is a chronic affection of the Eustachian tube, and one of the most prominent symptoms is the increased resonance or reverberation of the patient's own voice. If the tube of only one side be closed, the tuning-fork will be heard more acutely in that ear. There is also more or less pain as an accompaniment. It is described as a biting or gnawing pain, but is not of long duration. There is also at times pressure and heaviness in the head, and in some rare instances we have noticed vertigo. The objective symptoms are increased, redness of the nostrils, with elongated uvula, and enlarged tonsils, extending to the pharyngeal extremity of the tube, which is seen rounded, swollen, and covered with a gray mucous secretion, when examined by the rhinoscopic mirror.

CASE IV. *Aural catarrh of right ear of twenty five years' duration, of left ear of three months' duration; pharynx, Eustachian tube, and membrana tympani involved.*—Mrs. Y., aged seventy, of Philadelphia. Has been very deaf in the right ear for twenty-five years, in the left ear for three months, so that she could not hear ordinary conversation, unless the voice was so much elevated as to be painful to others. The watch could not be heard on either side; could hear the tuning-fork on both, but best on the right; meatus of both ears very dry; membrana tympani of right ear retracted, opaque; handle of malleus could not be seen, and no light spot; left membrana tympani not so opaque, more normal in curvature; there was a small ulcer in the nostril, and the pharynx was much relaxed and thickened; cough in the morning from accumulated mucus; Eustachian tube of right side evidently closed; left side was open, and gave a very faint, moist sound.

*Diagnosis.*—Catarrh of middle ear involving Eustachian



tubes, membrana tympani, and pharynx, following an attack of pneumonia.

*Treatment.*—Light penciling of the ulcer in the nose with solid nitrate of silver; dressed with ungt. hydrarg. nitr. and cerat. aq. rosar., equal parts. Snuff, for catarrh of pharynx, of powdered cinchona and cubebs. Politzer's douche to inflate the middle ear on the left side, changing it occasionally to the right side, with vapor of iodine, with inhaler, introduced by Dr. Roosa, the object being to diminish the thickened condition of the mucous membrane of the middle ear. A solution of sulphate of zinc was also injected into the Eustachian tube by Bishop's nebulizer; subsequently, by Eustachian catheter with double opening (see Fig. 92). The solution of chloride of ammonium was also used by the same means. Dr. Solis Cohen kindly assisted me.

Internal treatment, fifteen grains of the hydrochloride of ammonium with sugar, three times a day.

Local treatment of pharyngeal mucous membrane by means of gargles, insufflation of powders of tannic acid, etc. This treatment continued through March, with slight improvement. By the 10th of April she could hear her own watch with left ear, and the voices of her children. Throat less injected. On the 20th of April she could hear her watch faintly with the right ear, which she had not done for twenty-five years.

She continued under treatment until July 6th, when she was satisfied with the extent of the improvement. The left ear was entirely restored to its former normal condition, but the improvement in the right did not continue; she lost the power of hearing the watch, but was able to hear general conversation much better than before treatment.

CASE V. *Aural catarrh, with obstruction of the orifice of Eustachian tube, etc.*—E. W. G., aged twenty-eight, of Centre County, Pennsylvania. Has been deaf for five years, but only noticed it during the last three years. Has never had any severe disease except sick headache, and great liability to take cold. The progress of the deafness has been slow, yet getting worse each year. Very much increased during the last twelve months. Cannot hear general conversation.



No hereditary disease. His mother, a sister, and brother are deaf. Has been under treatment occasionally for two years. Visited me for the first time in September, 1869, but had not time to be treated.

*Present condition.*—Has never had pain; general health good, except sick headache from excess of food; bilious habit, yet regular in his bowels; sleeps well; noise in both ears, even in his best ear, after taking a cold from having his hair cut in cold weather; speech thick and nasal. Examination of tonsils found them very much enlarged, with thickening of the uvula, and chronic pharyngeal catarrh; no doubt this condition, acting upon the orifice of the Eustachian tube, displaced it, which was shown by the use of the rhinoscopic mirror. There was also hypertrophy of the glands, with swelling and hyperæmia of the mucous membrane. Right ear, meatus irritable; membrana tympani normal; light spot seen; hearing distance, September 16th, 1870, four inches. Left meatus, no cerumen; appearance, membrana tympani normal; Eustachian tubes open with difficulty, requiring a very strong blast of air through the Eustachian catheter; left, moist; right, dry.

*Treatment.*—Had applied to tonsils various agents to diminish their size; the best result was obtained by painting them with tincture of iodine.

September 17th. Removal of a portion from each tonsil by the knife; then applied the pencil of caustic soda and lime. This was continued every second day until they were reduced to one-half their former size. The Eustachian catheter was introduced every day until the 28th. On that day there was less noise in the ear, and Politzer's apparatus was then employed with success.

On the 29th September, hearing distance of right was seven inches; left, fourteen inches. Hearing distance for general conversation so much improved that his friends congratulated him on the wonderful change.

September 30th. Again employed the air douche, with increased improvement in both ears. He had now to leave for home, owing to the sickness of one of his children, and promised to return if the improvement was not permanent.



This is the third or fourth case in which the removal of enlarged tonsils and uvula has been of most decided advantage both to the voice and hearing. In numerous other cases we have known the tonsils to be entirely removed with so little care, that not the slightest improvement to the hearing of the patient resulted.

CASE VI. *Chronic catarrh of the middle ear, Eustachian tube, and membrana tympani; five years' duration (complete deafness of both ears); no improvement.*—Mrs. Mary S., aged twenty-four, born in Lancaster County, Pennsylvania, of German parentage, has been deaf for five years. General health good, menstruation regular, no children, no deaf relations. According to her statement, five years ago she went to a party, and, after becoming heated, with the same dress on drove to the sea-beach, and there took a bath. This was followed by a severe cold, which left her completely deaf. She applied to numerous irregular (so-called) ear-doctors, who placed various oils, etc. in the external ears without benefit. About one year since she placed herself under the care of Dr. James Collins, a well-qualified surgeon, who has given the subject of aural affections considerable attention. His diagnosis was "chronic aural catarrh," for which he treated her after the most approved method for one year, without any favorable result. He then sent her to the Howard Hospital.

*Objective Symptoms.*—Meatus auditorius of right side dry; membrana tympani resembling parchment; no bright spot; handle of malleus invisible; has pain over the nose, and discharge of an offensive character, indicating ozæna, disease of the mucous membrane lining the nasal bones; can only hear a trip-hammer or the rumbling of a car; a small amount of air entered the Eustachian tube by forcing it with an air-pump or the hand-apparatus. Right ear sensitive to electromagnetic current when filled with salt and water; left, when filled with the same fluid, not sensitive with the conductor in the meatus.

*Diagnosis.*—Aural catarrh (*sclerosis*) of the mucous membrane of the middle ear.

*Prognosis.*—Almost hopeless.

*Treatment.*—Bichloride of mercury and hydrochloride of



ammonium, with iodide of potassium, oil of hyoseyamus percoctum; perforation of the membrana tympani, and injection of fluids into middle ear; application of the induced current and the use of Politzer's and Thudichum's douches, all of which has been carried out without benefit.

The treatment was continued for three months, resulting in but very slight benefit to her hearing, but her catarrh and ozæna were much improved. She was recommended a Haswell's ear-trumpet.

In the abstract of the proceedings of the Otological Congress,\* held at Dresden, September, 1868, furnished by Dr. Roosa, the treatment of non-purulent catarrh of the middle ear was discussed, and Pagenstecher's nomenclature was adopted as the basis:—1. Hypersecretion; 2. Vascular ectasia; 3. Sclerosis.

In the first-named form the use of the catheter with air, vapor of the muriate of ammonia and injections, Politzer's method, internal use of quinine, iodide of potassium and iron, rubbing the skin, treatment of the naso-pharyngeal membrane by gargles and injections, and paracentesis of the membrana tympani, were the remedies spoken of.

In the second form, no irritants should be used. The internal use of acids and the application of the artificial leech were recommended.

In the third form the injection of irritants was advised. One of the members of the Congress had used a solution of atropine (grs. ij to fʒi), poured into the ear, with benefit in some cases, so far as diminishing tinnitus was concerned. This we consider too strong a solution to be employed without a great deal of care. We think even grs. ij to fʒi of water a very strong solution, for fear of a slight perforation or the Rivinian foramen allowing a portion to pass into the throat by the Eustachian tube. Obturation of the canal, as suggested by Politzer, had been practiced; but one delegate (Hinton) had not seen any benefit from this procedure. In the first class of cases almost all recover; in the second, one-half; in the third, one-sixth; and in the fourth, they are obliged to adopt a good hearing-trumpet.

---

\* Transactions American Otological Society, second meeting, 1869.



PATHOLOGY OF CHRONIC AURAL CATARRH (CAVITAS TYMPANI,  
OR MIDDLE EAR).

According to the late Joseph Toynbee,\* the cavity of the tympanum, or middle ear, when diseased, contains the following abnormal contents: pus, mucus, serum, epithelium, lymph, blood, cholesterine, serofulous, oily, and calcareous matter. He describes a case of chronic catarrh, in which he made a post-mortem examination, and found mucus escaping from the tympanum through an orifice in the membrana tympani, not normal, but thick and dark-colored, and distending the cavity (showing one of the most common complications).

No. 279 in his Catalogue. The tympanum nearly full of firm lymph.

No. 280. Right ear.—The tympanum contains a mass of soft, pale substance, which surrounds the stapes, and partly conceals the other ossicles. History.—Male, aged seventy-nine; deaf for four or five years; the deafness came on slowly from cold. The right ear was the worst. He could hear the click of the nail. This man, while living, suffered from a feeling as of stopping up on the right side of the head, and from a noise like the ticking of a clock, which was worse during a cold. The membrana tympani, on examination, was seen to be white, concave, and thick. The act of blowing the nose was accompanied with a feeling of cracking and bursting in the ears.

No. 280a. Right ear.—The meatus contains purulent fluid and thickened epidermis. The membrana tympani had fallen into the promontory, and was of a dark color, and thick. The anterior part of the tympanum is filled with a firm mass, resembling a partially-decolorized clot; the posterior part is filled with a clot of comparatively recent blood, which appears to extend into the mastoid cells. The malleus and incus are freely movable; the base of the stapes is somewhat less movable than natural. History.—Male, aged sev-

---

\* Toynbee's Descriptive Catalogue of Preparations Illustrative of Diseases of the Ear, in the Museum. London: John Churchill, 1857.



enty-nine. At times, when a boy, was hard of hearing, which he attributed to bathing; for fifty years had been decidedly deaf; the deafness varied much, being worse during a cold. He could hear a loud voice, but had a beating noise in his head. Died from cancer of the liver.

Nos. 281, 282. From a girl, aged two and a half years, who died from phthisis. Right ear.—*Membrana tympani* entirely destroyed by ulceration. The tympanum contained some scrofulous matter, although the lining membrane was so swollen as nearly to fill the cavity. Left ear.—*Membrana tympani* thick, white, and soft; the mucous membrane of the tympanum very thick and red. The tympanum was nearly filled with mucus, so dense as to be almost solid. Examined by the microscope, this substance was found to be mucus, in combination with scrofulous matter.

Two other cases, containing the same product, but no history of deafness during life.

No. 286. The tympanum contains calcareous matter, deposited chiefly upon its inferior and posterior walls.

No. 287. The incus dislocated, impacted in the mastoid cells, and surrounded by calcareous matter. No history of deafness.

No. 288. Mucous membrane very thick. No history of deafness.

Nos. 289, 289a. Strong bands of adhesion between the *membrana tympani* and incus. No history of deafness.

No. 305. From a woman, aged sixty-two, who died from gangrene. Deaf in left ear. Left ear.—*Membrana tympani* white, and very concave. Around the point of attachment of the malleus the blood-vessels were enlarged and distended with blood. The tympanum had three parts filled with a thick, tenacious white mass, which was partly the cause of the white appearance of the *membrana tympani*. The mucous membrane of the tympanum was thick and very vascular, and that portion of it which covers the body of the incus was red. In the succeeding twenty-four cases, although the mucous membrane was diseased, there was no history of deafness during life. We shall select only those cases in which a history of deafness is given.



No. 316. A portion of a greatly thickened mucous membrane from the tympanum of a man deaf since early life,—the cause of deafness having been scarlatina.

No. 326. From a woman aged thirty-eight, who died of dropsy. Deaf. Right ear.—*Membrana tympani* dull, more concave; tympanum was nearly filled with a firm gelatinous-looking substance of a red hue, and the mucous membrane of the tympanum was in a highly tumefied condition.

No. 327. Right ear.—*Membrana tympani* white and soft, and in a state of ulceration; mucous membrane of tympanum thick, soft, and ulcerated. The tympanum contained a large quantity of white muco-purulent fluid. The *dura mater* was separated from the mucous membrane of the tympanum by a very thin layer of bone, to which the latter was very slightly attached.

No. 328. Female, aged twenty-nine. Died of jaundice, with diseased liver and kidneys. She was deaf in the right ear, in which she had frequent gatherings, ending in a copious discharge. She also suffered from violent pains in the head, which at times were almost intolerable, but, these were relieved by the discharge.

No. 333. From a woman aged twenty-two. Deaf. Died from disease of the heart. Right ear.—The mucous membrane of tympanum and mastoid cells was much thicker and softer than natural.

No. 338. From a man aged sixty-eight. Deaf in right ear. *Membrana tympani* destroyed; the lower two-thirds of the long process of the malleus removed; the mucous membrane of the tympanum ulcerated; the cavity full of thick soft membrane and offensive discharge.

Fifty cases contained membranous bands of adhesion, and four were deaf prior to death.

No. 341. From a man, aged forty-four, who died of diseased bladder. He had been growing gradually deaf during many years. The Eustachian tube was pervious. Right ear.—The mucous membrane of the tympanum was thick, and bands of adhesion connected the stapes, malleus, and incus to the *membrana tympani*.



No. 342. From a man aged seventy. Died of disease of the heart. Hard of hearing. Right ear.—Membrana tympani of a dull lead color, but much thicker than natural; the tympanum completely full of cellulo-fibrous tissue, which connected the ossicles and the membrana tympani to the promontory.

No. 345, 346. From a man aged seventy. Had been deaf in the left ear since the battle of Trafalgar. For three weeks after the battle, was almost entirely deaf, but afterwards became better. During the four years preceding his death became more deaf; subject to noises of a tinkling character in the left ear during a cold; could hear a watch on pressure over right ear; with the left could hear a click of the finger-nails. Right ear.—Membrana tympani thick and concave. Numerous bands of adhesion connected the ossicles to the tympanic walls and to the membrana tympani. Ossicles not so movable as natural. Left ear.—Membrana tympani destroyed, except a narrow portion posteriorly. The ossicula remain, and are connected with each other and attached to the tympanic walls by broad bands of adhesion. The tympanum contained a secretion of a black color.

Nos. 347, 348. History. —, male, aged sixty-seven. Had been deaf for thirty or forty years after a naval engagement. Was subject to roaring noises in both ears, which sometimes ceased for a day or two. At times there had been a discharge from both ears. Up to about two years before his death, he could hear the click of the finger-nails with each ear, but before he died he became unable to hear the voice. Right ear.—Membrana tympani, lower two-thirds destroyed; the lower margin of the upper third was attached to the thick mucous membrane of the tympanum, which shuts off the mastoid cells from the tympanic cavity. The mastoid cells were full of a dark-colored serous fluid, epithelium, and cholesterine. Mucous membrane of tympanum very thick; labyrinth healthy. Left ear.—In the same state as the right ear, except that the fluid in the mastoid cells was transparent. The stapes was fixed. The mastoid cells were separated from the tympanic cavity by a distinct band between the inner



wall of the tympanum and the upper part of the membrana tympani.

Then follow thirty-three cases in which bands of adhesion were found extending across the tympanum, binding the small bones or the membrana tympani. No deafness recorded.

No. 401 to 410. *The canal for the portio dura nerve imperfect.*

Nos. 410, 410a. *The superior osseous wall hypertrophied.*

From a man, aged eighty-five, who had been deaf for many years. Right ear.—The meatus externus contained a large mass of hardened epidermis, pressing on the membrana tympani, which was opaque. The tympanum contained muco-purulent fluid. The superior wall of the tympanum was greatly hypertrophied, being more than half an inch in thickness. The canal for the carotid artery was greatly contracted. Left ear.—The superior wall of the tympanum was similarly hypertrophied. The canal for the carotid artery was contracted to a less degree.

No. 411. *Superior osseous wall expanded.*

No. 412 to 450. Superior wall partly deficient; the mucous membrane of the tympanum being more or less in contact with the dura mater.

No. 453 to 482. The inferior osseous wall deficient; the mucous membrane of the tympanum being more or less in contact with the outer surface of the jugular vein.

#### CONCLUSIONS IN REGARD TO THE PATHOLOGY OF DISEASES OF THE MIDDLE EAR.

From the cases here given, we may glean the following important facts in regard to the pathology of the middle ear: that it contains abnormal products of inflammation, given in detail above, which have been confirmed by puncturing the membrana tympani during life, and there have been found pus, mucus, and serum; also calcareous deposit upon the membrana tympani, and it is to be inferred that the wall participates in the same deposit, as just shown. There has been seen soft and hard lymph surrounding the malleus, incus, and stapes, and, as a matter of course, preventing their



proper vibration. There are also deposits of blood. These results have been confirmed by Von Tröltsch and others. The mucous membrane is thickened, vascular, and diseased, with bands of adhesion connecting all the bones of the ear, affecting the movements of the same, and the fenestra rotunda and ovalis.

It has also been found that the superior osseous wall of the tympanum has become greatly hypertrophied, and the walls expanded, with the superior osseous wall so deficient that the mucous membrane of the tympanum is brought in contact with the dura mater, and that of the inferior also, so that the mucous membrane is more or less in contact with the outer surface of the jugular vein; both are avenues to disease of the blood and brain.

As the pharynx and Eustachian tubes are apt to become involved, we shall describe some of the changes produced upon these important parts by this disease.

They have been found dilated, the tubes being very large. There are also strictures of the same caused by *hypertrophy of their bony walls*.

No. 487. A male, aged seventy, became deaf four years before his death, after sleeping in a damp bed. He required to be spoken to in a loud voice within two yards. The tympanum was full of a dark slate-colored fluid, which gave a color to the membrana tympani, and the upper wall of the tympanum.

No. 488. Eustachian tube impervious at its entrance into tympanic cavity; membrana tympani very thick.

No. 489. History. —, male, aged sixty-six, had been deaf for about seven or eight years, attributed to cold. The Eustachian tube was so much contracted at the part where the cartilaginous is continuous with the osseous portion, that it only just admitted the passage of an ordinary-sized pin.

No. 491. Eustachian tube and tympanic cavity full of dense fibrine.

No. 494. A man aged fifty-two, who died of dropsy, was deaf, and the stapes was completely ankylosed to the margin of the fenestra ovalis.



There are numerous cases of complete ankylosis of the stapes and other bones; partially ankylosed, one hundred and thirty-four cases in all.

Then fracture of the inferior extremity of the malleus; handle of the malleus carious from a youth, aged ten; effect of scarlatina; case of caries of malleus and incus; also, one of caries of malleus. Then follows caries of incus and stapes.

It will be seen that he makes an important distinction between complete and partial ankylosis; in other cases it was simply more firmly fixed than natural, by bands of adhesion in the tympanum. In those recent cases, when the lymph is soft, proper treatment may overcome this condition.

Von Tröltsch\* observes, under the pathology of aural catarrh, that "the muco-periosteal lining of the osseous middle ear, which for the sake of brevity we call a mucous membrane, has as yet received no complete microscopic examination as to its condition in a state of health." He also states that our knowledge and ideas of the morbid changes in chronic aural catarrh are still in a somewhat crude condition; that they are chiefly confined to what may be observed with the naked eye; and that we do not yet properly understand the finer changes in the tissue of the middle ear.

To illustrate these changes, Prof. Gruber has devoted much attention to this subject, and in his recent work has furnished two illustrations. The first (Fig. 65) is a view of a microscopical section of an inflammatory hypertrophy of the mucous membrane of the tympanic cavity, thickened to five times its normal condition. Chalky masses are seen scattered through the tissue, which, on the addition of muriatic acid, exhibit the usual reaction.

Fig. 65.



The following statistics show the result of treatment of

\* Von Tröltsch on the Diseases of the Ear, p. 282.



chronic catarrh, or chronic inflammation of the middle ear, by Prof. S. Moos, M.D., of Heidelberg:\*

Whole number of cases.	Cured.	Improved.	Unimproved.
100	17	64	19

These we consider very good results, to save from permanent deafness seventeen out of one hundred, or over five per cent.

Dr. H. Schwartz, of Halle,† a highly respected and successful aurist, reports seventy-three cases treated with the following results:

Whole number of cases.	Cured.	Improved.	Unimproved.	Not treated.
73	7	32	8	26

Following these we have the cases of another careful and exact writer, Prof. Roosa,‡ of the University of New York:

Whole number of cases.	Cured.	Much improved.	Improved.	Unimproved.	Unknown.	Died.
245	11	17	56	96	64	1
254½	12	32	55	75	95	

We here give our own|| results. See Table, p. 56.¶

Whole number of cases.	Cured.	Improved.	Unimproved.	Not treated.
111	12	51	39	9

We derive from the table of Prof. Gruber,\*\* of the General Hospital, Vienna (Austria) the following results:

Remaining Dec. 1867.	Admitted.	Cured.	Improved.	Not cured.	Not treated.	Remaining end 1868.
45	142	33	72	8	18	56

We shall also extract from Gruber's valuable work, published in 1870, his observations on prognosis and treatment, as it is our desire to present to our readers the latest views

\* Klinik der Ohrenkrankheiten, von Dr. S. Moos, Docent in der Universität in Heidelberg. Braumüller, Wien, 1866, p. 348.

† H. Schwartz, Halle, Praktische Beiträge zur Ohrenheilkunde. Wurtzburg, 1864, p. 48.

‡ D. B. St. John Roosa, M.D. Transactions of New York Medical State Society, 1868, p. 260.

§ Later published table of four hundred and ninety-four cases of disease of the ear, by Prof. Roosa, April, 1871. Am. Jour. Med. Sciences, p. 392.

|| Turnbull. Clinical Observations, Howard Hospital, etc., 1870.

¶ The author regrets that an error has occurred in his table in the cases of chronic catarrh, which are marked perforated, which is not correct.

\*\* Report of Diseases of the Ear, in General Hospital in Vienna, by Dr. Joseph Gruber, during the year 1868. See table.



upon this frequent and most serious disease of the organ of hearing.

Dr. Gruber\* states that "catarrhal inflammation of the middle ear, in general, admits of a favorable prognosis, provided it has not lasted too long, or has attacked an individual otherwise healthy, and has not produced an injury, difficult to heal or incurable. In the latter case, the condition of the tympanum and auditory ossicles must be particularly considered. A complete recovery can hardly be expected with a total destruction of the membrane of the tympanum; nor can the faculty of hearing be restored when the bones are separated from their connections, or exfoliated. The prognosis is likewise more unfavorable when the individual suffers from constant tinnitus, especially when the latter does not entirely disappear after the application of the air douche,—secondary alterations, which cannot be relieved by art, having taken place in the labyrinth and fenestræ. The prognosis is, however, most favorable in those cases in which, after the application of the douche, the tinnitus has entirely ceased, the faculty of hearing is restored, and the patient is not afflicted with other diseases which might cause a relapse of the aural affection. If the disease has been of long standing, although the patient be otherwise well, a perfect cure cannot be promised, as the douche does not restore the hearing, and the tinnitus will continue.

"The application of the tuning-fork as an auxiliary to prognosis, although recommended by many, is by no means reliable.

"The treatment is not only to relieve the local affection, but also to prevent the possibility of relapse. Concerning the latter particular, the bodily condition of the patient should be carefully considered, and lesions which may present themselves should be appropriately treated. Especially should we endeavor by every available means to counteract such diatheses as the scrofulous, tuberculous, syphilitic, etc., which most frequently keep up obstinate catarrh of the middle ear. In such cases, local applications are from the

---

\* Lehrbuch der Ohrenheilkunde, pp. 469-487.



beginning to be united with suitable constitutional treatment, the latter being even continued for a long time, although the former may have already answered their purpose, since relapses giving rise to an incurable affection are very liable to occur. The local treatment should extend to all the neighboring organs when they are similarly affected, or when their lesions have originated the existing disease. In the latter case, the rhinopharyngeal region should be attentively inspected, and the disorders there exhibited most energetically and skillfully combated; the new formations met with in that region, so frequently causing catarrh of the middle ear, should be extirpated as soon as possible, and the fundamental disease treated appropriately.

“Hypertrophied tonsils are very frequently the fundamental cause of catarrhal inflammations in the pharynx and middle ear, and should not escape the attention of the physician. In such cases, I prefer the extirpation of the whole tonsil, or the excision of a part of it, by the application of caustics or astringents. It is easily performed with Fahnestock's instrument, modified by Linhart, the consecutive treatment being finished within six days, the organ which is removed being possessed of very trifling physiological use; should the patient not submit to the operation, a solution of nitrate of silver,  $\frac{3}{2}$ – $\frac{3}{4}$  to  $\frac{f}{3}$ ss– $\frac{3}{4}$ i of distilled water, or powdered alum, may be applied. For this purpose I frequently employ forceps made of hard caoutchouc. A tuft of lint, or a small sponge, used as a vehicle for the medicament, is brought by means of this instrument to the place that is to be cauterized. On its rings there are two hooks catching each other, by means of which the forceps can be firmly closed. The cauterization is effected once a day, or at longer intervals, to be continued as long as requisite.

“With regard to the employment of gargles in the catarrhal affection of the nasopharyngeal mucous membrane, if they fail to give the desired result, from occlusion of the fauces by muscular action, I advise the patient to bend the head backward as much as possible, and, avoiding the motions usual in gargling, to bring the remedy in contact with the diseased portion alone. To reach the diseased



mucous membrane in the catarrhal affection of both the nostrils and superior part of the pharynx, the injections of remedial fluids through the nostrils recommended by me\* are most serviceable. Solutions of alum, borax, or tannic acid, grs. xx to ʒi, in distilled water, one pound or pint, are most applicable to them. The several injections are to be repeated after the reaction produced by the preceding application has subsided.

“For the treatment of catarrhal affections of the nasopharyngeal mucous membrane, it is not necessary to irritate the other nostril during the injection; but if the fluid is to be injected through the Eustachian tube into the middle ear, it should be introduced by pressing the corresponding ala nasi to the septum nari, in order to bring the injected fluid with more or less force, and to a certain extent to its place of destination. The snuffing up of astringent remedies, which is frequently recommended, is not as effectual as their injection; since, in the former case, they will reach only the anterior third of the nostril. Alum, or some other astringent finely pulverized, together with sugar, one to six, or even in equal parts, is generally used for snuff powders. I have never observed the disadvantage ascribed to injections into the nostrils, namely, that of impairing the sense of smell, to have been of long duration. What has been said about the catarrhal affection of the rhinopharyngeal mucous membrane, as well as the increase of the tonsils, with reference to the catarrh of the middle ear, is applicable, also, to all the other morbid conditions of that region accompanying the diseases of the ear, in so far as they are also to be treated *secundum artem*. To enlarge upon them is out of place in this connection. It must, nevertheless, be mentioned, that especially the healing of pharyngeal ulcers, as well as the appropriate treatment of a polypus, must be instituted from the beginning, as otherwise they

---

\* Incline the head to the affected side; place in the nostril of the same side a glass or elastic syringe (or glass tube), containing about ʒij of the required solution; this is either poured or syringed into the pharynx, where it comes in contact with the orifice of the Eustachian tube. Then inflate the tympanum without swallowing.



will always constitute a new cause of disease. With the treatment of the morbid affection in the pharyngeal region, that of the middle ear is to be connected. The condition of the Eustachian tube is, therefore, to be examined. When it is swollen at its orifice, and no other changes are discernible in its course, the above-mentioned treatment will often suffice to effect a cure. When there is at the same time a catarrhal disease of the pharyngeal mucous membrane, the injections above recommended are to be immediately followed by Valsalva's method. Did the catarrh at the orifice of the Eustachian tube, on the contrary, combine only with a simple pharyngeal affection, the astringent gargles above mentioned must be employed, and the orifice of the tube cauterized. With those who cannot bear cauterization, the astringents may be applied in spray, to which the universal pulverizer, made according to the principle of Richardson's anæsthetic apparatus of the corresponding length, the bend being given to the tube, through which the sprayed fluid is emitted, is applicable. From this time it is advisable frequently to employ the air douche for opening the tube, which will free the patient from the most disagreeable sensation of pressure in the diseased ear, owing to the inward inclination of the membrane of tympanum; after the absorption of the air in the cavity of the tympanum, the hearing is improved, at least momentarily.

"It will occur, not unfrequently, that catarrhal patients do not apply for medical assistance until after the acute inflammation has already disappeared, its product only remaining as a mucous plug in some portion of the tube, generally its orifice, which will be most apt to accompany the rhinopharyngeal catarrh. Such a ball, especially when tough and closing the tube, may be the primary cause of the affection. It should be removed by means of the air douche, or displaced, or made to occupy some section of the middle ear, where it remains harmless, as, for instance, in the cells of the mastoid process, and thus the cure of the disease will suddenly be accomplished. When the patient comes under medical treatment, while the swelling of the mucous membrane and the hypertrophic condition still continue, the air



douche should be applied for the removal of any collection in the middle ear. However, its application is admissible only when absolutely needed, and then with the greatest precaution.

“There will be some cases in which the catheter, under the application of the air douche, is to be introduced even into the tube, and others in which Politzer's method is indicated. With children, and those in whom natural impediments are met with during the introduction of the catheter, only the air douche without the catheter is applicable. I deem the same proceeding applicable to this object where the catarrh in the middle ear is associated with inflammation of the nasal mucous membrane; but the frequent application of the catheter will become most disagreeable to the patient, the mucous membrane of the nostrils being always irritated by its introduction. In such cases, therefore, the air douche without the catheter will be more beneficial. If, on the contrary, only the middle ear is diseased, the pharyngeal or only the nasal mucous membrane being normal, I think Politzer's method, if practiced for a long time, is not beneficial, because the air is pressed by it into the nostrils, and must penetrate in all directions for an escape in the rhinopharyngeal region, as the pharynx is closed below by the motions of deglutition. It will, therefore, spread in the nostril in all directions, passing into all the neighboring spaces. In this way the nasal mucous membrane may be injured; the action of the air forced into the lachrymal duct will injure the latter, and even the conjunctiva. This is, indeed, no theoretical induction. I have too often had patients with whom this method was practiced for a long time, and who complained of various affections of their eyes, especially catarrh of the conjunctiva, and a most annoying dryness, together with a feeling of pressure in the eye. On this account I prefer the air douche, with the application of the catheter, in all those cases in which it can be applied, and in which the rhinopharyngeal mucous membrane is uninjured.

“With regard to the air douche in catarrh of the middle ear, the condition of the tympanum must also be taken into consideration; and it should be remembered that this membrane is often very brittle. Upon the application of the catheter, the



air may be pressed into the cavity of the tympanum with regulated power, which cannot be done in Valsalva's method, the bag being emptied with vehemency in the short moment of deglutition, and the tympanum pressed outward so forcibly that it may rupture. From the indications which we have hitherto stated for the employment of the air douche, it will become apparent that in many cases it will suffice to apply it once, whereas in others it must be employed as often as the secretion collects in the tube and cavity of the tympanum and gives rise to morbid phenomena. In such cases it always has the object of removing the inflammatory product. It may become beneficial, also, in those cases in which there is no collection of morbid matter, but the swelling of the tube may have terminated by closing it, or producing secondary alterations in the cavity of the tympanum. In such cases the air will make its way through the narrowed tube, and, passing into the cavity of the tympanum, press the membrane outward, together with all the ossicles of the ear. In this manner the patient will be freed, for a longer or shorter time, from the annoying subjective symptoms. The pressure which the air exercises during its passage through the tube and into the cavity of the tympanum upon the swollen mucous membrane is, on the other hand, not unequal, in its effect, to the compression of other swollen organs. Consequently it promotes absorption, and to this circumstance no less than to the beneficial influence which the ventilation of the middle ear exerts, I ascribe the complete cure obtained by the skillful application of the air douche, in many cases of acute catarrh complicated with considerable swelling of the mucous membrane.

“The mucous membrane, in the whole middle, is sometimes extraordinarily inflamed from the commencement of the catarrh. This hyperæmia is then not confined to the middle ear, but extends over the labyrinth, and not unfrequently to the deeper organs of the external ear. The subjective noises will, in such cases, be increased, and the objective changes revealed by a high degree of hyperæmia in the tympanum. Local bleeding should then be used, the number of leeches to be applied being in conformity with the objective changes and the strength of the individual. Though I must gener-



ally deprecate the frequent abuse of local and general bleeding in diseases of the ear, yet I cannot help intimating, on the other hand, that by restricting bleeding to those cases in which the vascular depletion tends only to mitigate the pain, sins of omission may be committed which the physician will afterwards deeply regret. When, in catarrhal inflammation, such local bleedings are made under the indications above mentioned, their beneficial effect will be immediately manifest, especially in regard to the noises heard in the ear. If this is not the case, after the depletion has been performed twice or thrice, at intervals, it will be unavailing, although it should be renewed more frequently. If it then fail, it is owing to secondary changes that have taken place in the labyrinth. Cutaneous irritants, particularly the vesicants and ointments so much patronized by the public, in spite of their painfulness, when applied to the skin about the ear are not as beneficial as local bleedings; in many cases they are very inefficient.

“In severe catarrh the bowels will frequently be irregular, and the patient feels much relieved when their action is promoted by laxatives, or stronger saline purgatives, suited to the circumstances. The regular use of mineral water, especially when indicated by constitutional ailments, is likewise attended with benefit, and I have derived very satisfactory results from the use of the springs at Friedrichsthal, Marienbad, Karlsbad, Rakóczy, etc., when there was a great predisposition in the patient, and the local treatment did not suffice. If the catarrhal affection is treated from its beginning according to the rules hitherto laid down, and with the remedies mentioned, the patient being otherwise healthy, and a suitable regimen followed, the disease will yield, at the most, after a few weeks. Should this not be the case, and the disease confined to the mucous membrane, a copious product of inflammation resulting, the injection of astringent fluids into the middle ear may be introduced by means of the catheter. Such injections are, however, injurious, when they are performed at the beginning of the disease, or in cases of hyperæmia of the mucous membrane, since they will increase it, whereas the simple air douche, properly applied, will prove



truly beneficial. Should the catarrh threaten to become chronic, the mode of life of the patient is to be regulated, and whatever has an injurious influence upon the disease is to be forbidden. He should quit his occupation if it influenced the disease.

“Exercise in the open air should be taken; fats, stimulants, and all stimulating food must be interdicted. Sleeplessness should particularly be avoided. I have frequently made the observation, that catarrhs which had lasted for weeks, suddenly disappeared when the patient had left the city to live in the country. This beneficial effect becomes more marked by change of climate. Medicaments applied externally, as a matter of course, have but little influence upon the mucous membrane of the middle ear. As long as the membrane of the tympanum is not perforated, such medication—as, for instance, the dropping of fluids into the external auditory passage—is apt to increase the original morbid process by stimulating the parts with which it comes in contact, and should, therefore, be discarded. It cannot be denied, however, that the application of absorbents around the ear may occasionally be of the greatest, though indirect, service for the relief of secondary ailments. This is especially the case with the preparations of iodine and mercury, being most serviceable in chronic catarrhs depending upon a scrofulous or syphilitic diathesis. The application of medicines upon and over the external ear may promote the cure of the disease, when the membrane of the tympanum is perforated, or the external meatus presents such objective changes as require the application of medicaments. The penetration of such fluids into the cavity of the tympanum, as a matter of course, is subjected to certain conditions, of which we have already treated.

“We must, however, cursorily repeat here, that even with a greater loss of substance in the membrane of the tympanum, the retrogression of the disease cannot be expected from the sole instillation of medicines in the catarrhal inflammation of the whole middle ear. If, in such cases and under such medication, no recovery ensues by the aid of nature, the patient may ineffectually employ the medicines best indi-



cated, even to their own prejudice. The surest method of relieving the inflammatory affection is, in all cases where medicaments are indicated, the injection through the Eustachian tube. By instillation the auricle may be supported, if it is skillfully practiced, and the patient can do it without the assistance of the physician, but not without the knowledge of what has already been said and what will be said afterwards, otherwise the injury done may be greater than the benefit derived. In consecutive inflammations in the external ear-passage and the tympanum, the instillations of remedial fluids may be urgently indicated. In this respect we refer to what we have represented under the treatment of external otitis and myringitis, intimating that the attendant physician must always bear in mind that, in such cases, he has only the consecutive disorders to combat. He must attend principally to the original disease in the middle ear, acting at first more expectantly, unless changes occur which demand a quicker action.

“He will then frequently notice, with the diminution of the inflammatory symptoms in the middle ear, and particularly with the decrease of the fluid which, passing through the opening of the membrana tympani into the external ear, when copious, or of bad quality, constantly irritates the parts with which it comes in contact, that even the irritation thus produced will also gradually disappear, without any direct treatment having been required. On the other hand, the treatment of such conditions as might terminate unfavorably must not be delayed too long. The secondary straightening of the external auditory passage, owing to the swelling of its soft parts, for instance, demands quick dilating in the way before described. Polypous growths, produced from consecutive inflammation in the external ear, must likewise be timely extirpated, because the inflammatory products formed in the depth are prevented from being discharged, thus causing the inflammation to terminate most unfavorably. An important part in the treatment of catarrh in the middle ear was formerly played by the application of medicated vapors. Many authors, as, for instance, Von Tröltsch, still give them credit for a particular beneficial agency. I have, in



former years, made use of them extensively, but in their use I have been convinced that the same result will be obtained in a more simpler mode, from the application of medicines through the Eustachian tube. Since that time I employ medicated vapors but seldom, particularly when the injections of fluids have proven successful, and I wish to do everything to moderate the distress of the patient; but such medication will then generally prove inefficient. This will not, however, imply that the medicated vapors are devoid of all healing power in the disease under discussion. They are, on the contrary, as efficient as the medicated fluids; but I believe the same result may be obtained by the latter means and in a simpler way. The action of vapors is more difficult, on account of their being employed at a higher temperature, which causes the mucous membrane of the tube to swell, whereby it is narrowed, so as soon not to admit the vapors.

“Of the various vapors employed, I must ascribe the greatest efficiency to a solution of common salt,  $\mathfrak{z}\text{ij}$  to  $\mathfrak{z}\text{iv}$  of water. Besides, we have employed the solutions of borax and alum, of the same strength, the vapors of muriate of ammonia, tincture of iodine, the various ethers, acetic and pyroligneous acids, chloroform, pitch- or tar-water, etc. With the exception of those vapors which act as narcotics, contributing to the soothing of the subjective sensations, I should not like to attribute to them any preference, believing that in the future they will be less and less employed. It is, perhaps, principally the high temperature at which they are employed that constitutes their beneficial action, as the simple aqueous vapors would produce the same effect. Of the aeriform bodies, carbonic acid has especially been frequently tried in chronic catarrh, and its action praised. For this purpose it is developed from carbonate of lime and hydrochloric acid. The acid is gradually added to the lime contained in a bottle, the neck of which should be long. Experience, however, has taught that carbonic acid does not render greater service than the use of the medicated fluids. For their application the compressive apparatus before described may be used, or the vapor may be blown directly out of the apparatus into the tube by the



mouth. Such bodies as will readily evaporate, and are applicable without aqueous vapors, may be simply put in a valved bulb, the air impregnated with vapor being driven in through the catheter. When the compressive apparatus is used, or the vapor is blown by the mouth into the tube, the vessel which contains the vapor or aeriform body must be provided with two glass tubes, bent at right angles and fastened in the stopper of the vessel. The end of the tube outside the vessel is to be connected with the compressive apparatus by means of a caoutchouc hose, the end of which is then to be joined to the catheter introduced into the Eustachian tube, in which way the vapors can be driven into the middle ear by means of the compressive apparatus. The vapors may be blown into the ear through the free end of the glass tube, which otherwise is connected with the compressive apparatus. If the fluid to be employed must be warmed, the vessel containing it should be placed in the sand-bath, or over a spirit flame. In the application of warm vapors (those of water are employed at a temperature of  $35^{\circ}$  to  $60^{\circ}$  Rem.), the catheter must be surrounded with a non-conductor—best an elastic tube, or simply some sticking-plaster—to protect the nasal parts from the high temperature.

“We must not fail to notice, in this connection, the pneumatic apparatus to which so much efficiency has been ascribed in the diseases under discussion. I am of the opinion that it can be reduced to the same principle as the above, viz., pressure of air; and our therapeutics has obtained no great result from it, since we have the means to heighten the pressure in a more direct way throughout both the tube and auditory passage as much as we please, without confusing the patient. Acute catarrhs will occasionally subside of themselves, without any assistance; and, in chronic cases, this apparatus will accomplish nothing.

“With reference to the use of various springs, I find that they are of no direct benefit to the diseases of the ear, but are indicated only to cure or relieve the general disorder with which the various inflammatory processes in the mucous membrane are connected, and which of late have been too much neglected by physicians. The use of salt, iodine, or



iron baths must find a frequent indication, according to what has been shown under the head of Etiology. Salt baths are most effectual with individuals who, from the tenderness and great sensibility of their skin, are inclined to relapses; and Pagenstecher declares that the methodical use of warm baths had acted beneficially. He asserts that some patients had, at least, experienced no increase of their deafness from the use of the Weisbaden therms (hot springs)."

CATARRH OF THE TYMPANUM—PARACENTESIS OF MEMBRANA TYMPANI.

The operation of puncturing the membrana tympani has been employed since the time of Sir Astley Cooper, and is still resorted to for relieving a thickened membrana tympani. At one time it was highly lauded, and again much neglected, and was revived by Bonnafont and others, but never founded upon a true scientific and pathological basis.

The following are the conclusions of Schwartz\* and Tröltsch†, two of the most careful and reliable observers, that "among the many reports of favorable results from this operation, none of them can be said to give any sufficient evidence of its real value [of course Von Tröltsch is now speaking of the paracentesis performed in cases of chronic catarrh and not of those where it has been done, as in the opening of an abscess], unless the patients have remained for a long time under observation after the operation. Most of the histories are in this respect imperfect." Schwartz says "that, up to the present time, it is only in very rare cases a permanent success has been seen by trustworthy observers." Mr. James Hinton‡ reports of seven cases for the most part very favorably. Five in 1870, three of which were very successful, would throw some doubts upon the conclusion arrived at by Schwartz, who only operated upon eight cases of chronic catarrh; he also states in his pamphlet,

---

\* Die Paracentese des Trommelfells, von Dr. H. Schwartz. Halle, 1868.

† Treatise on the Diseases of the Ear. Translated by Dr. Roosa, New York, 1869.

‡ Guy's Hospital Reports, 1869-70.



that the operation is useful in a limited number of chronic cases. Mr. Hinton, as far as the author can judge, is perfectly honest and willing to relate his failures, as on p. 227: "Since most of the cases I have referred to have been successful ones, I think it is better briefly to relate one in which I was entirely unsuccessful. Such cases, no doubt, will continue to occur, but I believe that a good diagnosis, such a one as we may very reasonably hope to attain, will reduce them to an exceedingly small percentage." We shall give an abstract of three successful cases:

"CASE I. *Deafness, with recurrent attacks of inflammation in each tympanum, following scarlatina six years previously; evacuation of small dense masses of mucus by incision.*—May 14th, 1868. M. G., aged twelve, a healthy boy; at age of six had scarlatina, followed by discharge from right ear only, lasting about fourteen days. Ever since he had been decidedly deaf in the right ear, and with the exception of two years, had been more or less so in the left, and subject also to frequent attacks of pain, during which any sound, as music, etc., became painful. Watch\* was heard on contact on right side, and one inch on the left; the tuning-fork was heard well upon the vertex, louder in the better ear. Each membrane appeared very concave; there existed also a small white *dimple* anteriorly, apparently a healed perforation. The throat was pale. He could not inflate the tympanum. Air was blown through the Eustachian tube when he swallowed, without much difficulty, entering the tympanum with a sharp sound, and improving the hearing on the right side to three inches, and on the left to twenty-four, the membranes appearing afterwards convex posteriorly. Tincture of chloride of iron, sol. iodine, salt, and water to be drawn through the nose.

"Jan. 24th, 1869. The right ear has remained very deaf, and lately been subject to pain in the left ear for three or four days, attended with discharge. In a week, by treatment, the redness and pain were gone; watch heard each side at

---

\* The hearing distance of Mr. Hinton's watch is not stated. The late Mr. Toynbee's was three feet. Some must employ a watch with a much louder tick; hence the variety in the results.



fourteen inches. I had formed, however, the decided opinion that the continued liability to pain and deafness depended upon the presence of retained secretion in the tympanum, and that its removal afforded the only prospect of permanent benefit. I made the usual incision in the right membrane, viz., between the position of the malleus and incus, and extending from within half a line of the upper to within a line and a half of the lower border of the membrane. I then, after passing air through the Eustachian tube, injected through it, by means of the catheter attached to an elastic bag, a solution of bicarbonate of soda (grs. xx to fʒi). The effect produced by this was that there protruded from the incision a white shining mass of mucus. For the next four days the injections of air and solution of soda through the Eustachian tube were repeated, and with the effect of washing out through the perforation small dense masses of white mucus; on the fifth day the incision was healed. The same treatment was then adopted for the left ear, and with very similar results. The improvement of hearing on each side was decided, but not complete; accordingly the incision was repeated, and with precisely similar results,—the gradual forcing out of small dense masses of mucus; twice more on each side. No irritation whatever followed, and the hearing of both ears became, so far as I could judge, absolutely perfect, the lowest whisper being heard at the distance of six yards. The right ear, which had been continuously deaf for six years, appears to have recovered as completely as the left."

Mr. Hinton believes that relaxation of the membrana tympani, accompanied with thinning of its substance, is a very frequent effect of the protracted presence of mucus within the tympanum, and that when mucus is present within the tympanum, the membrane is more depressed, at least in parts, than protruding outward. He has also known of alternate healing and giving way of the membrane in which mucus had accumulated. On this ground he has adopted the practice of treating all perforations of the membrane, acute as well as chronic (unless particular conditions forbid it), by washing out the tympanum with a warm solution of bicarbonate of soda and chlorate of potash, syringed freely



through the perforation. This is done by means of the Eustachian catheter, but in the opposite direction, by applying the syringe to the meatus, the nozzle being well covered with elastic tubing, so as to fit pretty closely without causing pain; the stream may be sent with considerable force, provided the Eustachian tube be open, and will escape by the nostril, the head, of course, being bent forward. This cleansing should be repeated every second or third day until all appearance of an accumulation has ceased, and then a solution of sulphate of zinc (grs. ij-x add fʒi of water) may be substituted and used in the same way.

CASE II.—This is an instance in which, after perforation, masses of mucus were syringed out with good results. Among the complications which Mr. Hinton notices, is a frequent obstruction of the Eustachian tube. This is for the most part dependent upon the swelling of the mucous lining, and an inspissated secretion; this is sufficiently treated by inflation during the act of swallowing, and by the passage of alkaline and astringent fluids employed to wash out the cavity of tympanum. This is by no means always successful in relieving the case; sometimes the constriction is of the most obstinate character, and requires for its removal vigorous and persevering treatment,—by the introduction of bougies of laminaria or very small elastic tubes passed through the Eustachian catheter and along as far as possible; air or fluid being then injected through them so as to exert a direct pressure upon the constricted part.

He then records Case III., in which mucus accumulation took place in each tympanum. On the right side deafness existed since childhood, with obstruction of the Eustachian tube. Three times the incision was repeated; also the treatment by bougies, with perfect recovery. The only unpleasant result after the introduction of a laminaria bougie was, that the air escaped into the cellular tissue, causing some swelling of the soft palate and uvula.

These were pricked by the point of a knife, and pressure made, and in two days all swelling had subsided. The treatment, however, was omitted for three weeks, and then resumed by the injection of a solution of carbonate of potash.



He began in August, 1868, and it was not until February 2d, 1869, that a small stream of air entered the tympanum; but

Fig. 66. — INSTRUMENT FOR PUNCTURING MEMBRANA TYMPANI.



on the following day the tube closed again, and this continued for some weeks. On the 9th of March the mem-

brane was incised for the last time, and a yellow viscid matter was washed out; but it was not until April 16th that the tendency of the tube to close was overcome. On that day he syringed into the tympanum a solution of the nitrate of silver (grs. v ad. f3j), and from that time inflation was easy. The watch was heard at forty inches, and a low whisper at six yards.

The author has resorted to the operation in chronic cases of catarrh of the tympanum in several instances, but, so far, his success has not been what he could have wished. When it is desirable to make a simple puncture, the instrument Fig. 66 was employed; but if a portion of the membrana

Fig. 67. — INSTRUMENT FOR INCISING THE MEMBRANA TYMPANI.



tympani was to be removed, or a large incision made, he used the sickle-shaped instrument, Fig. 67.

The operation in some cases is attended with considerable pain; to allay this, and the nervous feeling attending the dread of the operation, it is well to administer the nitrous oxide gas, or a mixture of sulphuric ether and chloroform, so as to produce a gentle sleep.

#### ACCUMULATION OF SERUM IN THE TYMPANIC CAVITY.

In the valuable monograph of Politzer on the membrana tympani, in "Health and Disease," the translators\* have added a supplement from him, of the diagnosis and treatment of the accumulation of a fluid resembling serum, in the tympanic cavity, but not involving the transparency of the membrana tympani; that it admits light enough into the cavity so that this fluid can be seen, being of a dark-gray color on the membrana tympani. The boundary between the parts is marked by a fine black line, having the appearance of a

\* Translated by A. Matthewson, M.D., and H. G. Newton, M.D., New York. William Wood & Co., New York, 1869. Pp. 161-169.



black hair lying upon the membrane. By causing the patient to lie down, the black line changes its position, showing it is fluid. When air was forced through the Eustachian tube by Politzer's method, and the membrana tympani inspected, the following changes occurred: the lateral portions of the membrane bulged out, and instead of the distinct limitation in the colors before described, the whole field was covered with rings (or air-bubbles in the fluid),\* of various sizes, having dark contours. The other symptoms were deafness, heaviness, and pressure in the head, with loud ringing.

The treatment in the majority of Dr. Politzer's cases consisted in the employment of his method for inflating the middle ear, with the head bent over to the opposite side, and somewhat forward. In the cases in which the membrane was much shrunken, and presented a dark greenish-yellow color, Dr. Politzer, after using the air douche with only temporary success, performed paracentesis, and immediately afterwards forced air through the ear by his method, thus driving the matter out into the external meatus.

The late Mr. Toynbee discovered, in a post-mortem examination, the first indication of serum with or without mucus or lymph. Voltolini† followed Toynbee, more especially in his inaugural essay (*Examination of the Ear in the Cadaver: Breslau, 1862*), by describing the peculiar characteristics of the mucous membrane of the tympanum as those of a serous membrane. Politzer, again treating of this subject (1869), after numerous observations, mentions a new characteristic, that, partly from the deflection of the rays of light reflected from the promontory, and partly from the yellowish color of the fluid in the tympanum, the gray color of the membrane is often tinted a light bottle-green.

Some of the patients had the sensation of the movement of a fluid in the middle ear when the head was inclined forward, backward, or sideways. This symptom was noticed

---

\* Von Tröltzsch was the first to recognize, by inspection, the presence of mucus bubbles in the tympanic cavity, in 1862.

† Serous Accumulations in the Tympanum: *Archives of Ophthalmology and Otology*, by Professors Knapp & Moos, vol. i. No. 2, p. 557. New York, W. Wood & Co., 1870.



in a case under the writer's care, and was relieved by Politzer's method.

During an interview between Dr. Moos and Politzer, in 1869, the latter mentioned a still more characteristic symptom,—the decided improvement in hearing following the use of the air douche; which, however, generally sank to the original minimum in a very short time, especially when the air douche was not regularly continued. Dr. Moos\* has had the opportunity of observing seven cases of serous exudation into the tympanum. Five of these were on both sides, and four were relapsing. The following is an abstract of these interesting cases:

CASE I.—The result of a wound. Paracentesis on both sides; relapse; final recovery. Duration of treatment, 3d of May to the 22d of August.

CASE II.—Serous accumulation in the left tympanic cavity, affording no indication on the tympanic membrane; paracentesis but once performed; permanent recovery. Duration of treatment, May 10th to September 5th.

CASE III.—Serous accumulation in the tympanic cavity on both sides; paracentesis on both sides; repeated on the right; otorrhœa of short duration; formation of furuncles in both external ears; recovery. Duration of treatment, 20th of March to the 28th of April. Hearing distance not given at the termination of treatment. Watch employed—normal hearing distance thirty feet. This patient only heard it at nine inches. This case is somewhat remarkable as being from Philadelphia.

CASE IV.—Serous accumulation in the right tympanic cavity; small quantity of serum; the demarkation indicated by two divergent descending lines; paracentesis; subsequent otitis media and formation of furuncles in external meatus. Duration of treatment, 1st of June to June 12th. Hearing distance of watch prior to treatment, one foot; after, two feet.

CASE V.—Serous accumulation in the tympanic cavity on both sides; paracentesis on both sides; hearing restored; continuance of subjective noises on the left side. Hearing distance at the commencement of treatment,—right side, six inches

---

\* Archives of Ophthalmology and Otology, op. cit., pp. 557-602.



(in place of six feet); left side, with the watch, one foot (in place of thirty feet); voice, at a distance of six paces. Duration of treatment, May 25th to the 11th of June. The patient had the normal degree of hearing in the right, and four inches in the left.

CASE VI.—Serous accumulation in the tympanic cavity on both sides; absence of the characteristic appearance of the membrana tympani; paracentesis and relapse on both sides, followed by the first appearance of characteristic symptoms on the left side; repetition of the paracentesis on both sides; repeated myringitis, resulting from operation on the right side. Duration of treatment, from 8th of June to July 14th. Hearing distance prior to operation,—right ear, five inches; left ear, seven inches. After July 14th, hearing distance six inches. Perforation remains.

CASE VII.—Rapid accumulation of serum in both tympanic cavities; paracentesis on both sides; sudden restoration of hearing. Duration of treatment not given. Hearing distance prior to treatment, one and a half inches; after air douche, at eighteen inches. The hearing distance reached four feet.

*Remarks.*—With the exception of a boy nine years of age, all the subjects were adult males.

“The conjectural duration of the disease extended from four days to three weeks.

“The immediate cause could be determined in one case only,—a draught of air upon the head when perspiring.

“In the remaining cases the affection must be considered a secondary one, an extension of the catarrhal inflammation from the nares and pharynx to the Eustachian tube.

“Continuous subjective noises were present in four cases; interrupted in two cases,—in one so severe that the continued singing had a disturbing effect upon the mental action of the patient.

“In the majority of the cases these noises were removed by the introductory treatment. In one case they disappeared entirely only after a long course of treatment following paracentesis; in another case they persisted, and made an unfavorable symptom for the prognosis. Where they had once existed, and been dispelled by treatment, they returned



on the occurrence of a relapse, and remained, as a rule, during its whole course. Vertigo was present in two cases; in one of these it continued after paracentesis had been performed, and was probably owing to gastric trouble.

"During the time immediately following the paracentesis, we should not omit to make the patient remain in bed, or keep his room, at least until the perforation has closed, and in unfavorable weather, under all circumstances. Early treatment of the pharynx and nasal passages is necessary.

"On the whole, the results are so favorable, and the cure so often permanent; that the treatment of serous accumulations in the tympanum (middle cavity of the ear) may be numbered among the most satisfactory in aural surgery."

#### NEW METHOD OF PREVENTING THE CLOSURE OF AN ARTIFICIAL PERFORATED MEMBRANA TYMPANI.

The following case is worthy of notice, from the success of a new method of treatment employed by Politzer to prevent the artificial perforation of the membrana tympani closing:\*

"On the 22d of October, 1868, Anna P., aged forty-eight, a native of Vienna, first presented herself at Politzer's clinique. Two years before, according to her account, she was suddenly seized with severe pain in the left ear; the pain lasted for eight days, and was followed by a copious secretion. This also ceased in a few days, but deafness, and the passage of air through the meatus on blowing the nose, continued for some weeks. An application to the meatus, ordered by a physician, did much to diminish the secretion, and decidedly improved the hearing; this improvement lasted but about two months, and at the end of that time she began to experience troublesome sensations in the ear, such as loud ringing and rushing sounds.

"A marked degree of deafness was observed at the date of admission; this had lasted for a year.

"The trouble in the right ear dated from three months before admission; it commenced in the same manner as in the left ear; the pain continued, notwithstanding the appear-

---

\* Clinique in Aural Surgery, observed and reported by C. J. Blake, M. D., Boston Medical and Surgical Journal.



ance of a purulent secretion on the first day, for a period of three weeks. The secretion and the sensation of a hissing sound in the ear were still present upon admission.

"On examination, the left membrana tympani appeared of a dull-gray color, the short process of the malleus and the folds running outward from it were very sharply marked, and the manubrium mallei drawn inward and less clearly defined than usual; below the manubrium was a dark and depressed spot of an oval form, about 1''' long by  $\frac{1}{2}$ ''' wide, which was recognized as the cicatrix of a previously existing perforation. The ticking of a watch could be heard only when the watch was brought in contact with the ear, or when it was pressed upon the side of the head; and the voice at no greater distance than 3'. After the injection of air into the Eustachian tube by Politzer's method, the cicatrix was seen to have been forced outward and the distance at which the voice was heard was increased by one foot, but it was still necessary to bring the watch in contact with the ear. In the right membrana tympani, situated below and anteriorly to the manubrium mallei, was a circular perforation of the size of a pin's head, through which a drop of thick pus and a few air-bubbles were forced by the employment of Valsalva's method. After removal of the pus, the watch was heard at a distance of 2'', and when pressed upon the head in front of the ear; the voice at a distance of five feet. The tuning-fork, placed over the sagittal suture, was most clearly heard in the left ear, because the obstructions to the passage of vibrations from without retained those which were communicated to the tympanic cavity through the medium of the bones of the head. The results of the examination showed the existence of purulent catarrh of the tympanic cavity of the right ear, with perforation of the membrana tympani. The same process had existed in the left ear, but the purulent secretion from the mucous membrane of the tympanic cavity had ceased more than a year previously, and the perforation in the membrana tympani had been closed by a cicatricial tissue. The fact that a decided improvement in hearing followed the injection of air, favored the supposition that the deafness resulted neither from closure of the Eustachian tube nor from



a collection of mucus in the tympanic cavity, but from (old) adhesions between the ossicles and the tympanic walls.

"The previous history showed that as long as the patient was able to force air outward through the meatus, that is to say, as long as the perforation existed, the loss of hearing on this side was but slight, whereas the closure of the opening was followed by decided deafness. These facts justified the conclusion that either the malleus or incus was attached to the wall of the tympanic cavity by firm bands of connective tissue. So long as the perforation in the membrana tympani existed, the vibrations could pass through it, and, striking upon the base of the stapes, be communicated to the labyrinth. After the closure of the opening, the vibrations impinging upon the membrana tympani were no longer communicated to the stapes, because of the resistance of the malleus and incus. The indication for treatment in this case was the removal of the thin cicatrix, in order that vibrations might be admitted to the tympanic cavity through the perforation.

"The excision of a portion of the membrana tympani in certain cases of disease of the ear was first recommended by Riolan, in the seventeenth century; but from the doubtful results of the operation, very little regard was paid to it by the more celebrated surgeons of the time. The first and most important scientific communications concerning the artificial perforation of the membrana tympani date from the beginning of the present century, when Sir Astley Cooper and Himly, simultaneously and independently of each other, reintroduced, and, encouraged by the favorable results at first obtained, strongly recommended it. They became convinced, after long trial, of the difficulty of making the artificial opening a permanent one, and finally abandoned the attempt; but in the mean time their favorable reports having reached the continent, the operation was performed in both France and Germany on hundreds of cases, in almost every one of which, however, the first gain proved to be but temporary, for sooner or later the opening became closed by cicatricial tissue.\* Pieces of catgut cord, splinters of whalebone, etc.

---

\* Schwartz. Studien und Beobachtungen über die Künstl. Die Paracentese des Trommel-fells. Archiv für Ohrenheilkunde. 2 Bd.



were introduced, and the edges of the perforation touched with caustics, but without favorable result. Bonnafont,\* in one case, introduced a silver canula, and allowed it to remain forty-five days, but the closure of the perforation followed its removal. Many instruments have been devised for the purpose of removing a portion of the membrana tympani. Among the latest are those of Wilde, Toynbee, and Gruber; but none of them has succeeded in gaining a decidedly favorable result. The operation proposed by Menière, of perforating the membrana tympani by means of caustics, seems to have been equally unsuccessful; and concerning sphyrotomy,† we have as yet no definite reports of success. The indications for the operation in the present case induced Dr. Politzer to make the attempt of establishing a permanent artificial perforation by means of a new method; and on October 26th, he operated as follows: a large-sized aural speculum having been introduced, and the membrana tympani illuminated by means of a reflector, an incision was made with a paracentesis knife through the cicatrix parallel to its long diameter, and a 'laminaria-stäbchen'‡ introduced, and allowed to remain for two or three hours, by which time it had sufficiently expanded. Its removal showed a circular opening in the membrana tympani, through which a hard rubber eyelet, having several rims, was pushed into the tympanic cavity to the depth of  $\frac{1}{2}$ ". The pain resulting from the operation was very slight. Through the opening of the eyelet, the mucous membrane covering the promontory, of a light reddish-yellow color, could be plainly seen; and the patient voluntarily declared not only that she heard better, but that the sensation of noises in the ear had very much diminished. It having been ascertained that the eyelet was firmly fixed, the

---

\* *Traité théorique et pratique des Maladies de l'Oreille.* Paris, 1860.

† Sphyrotomy, the operation of excising the handle of the malleus; myringo-dectomy being the excision of a portion of the membrana tympani. Both these means are employed as remedies for impairment of hearing and *tinnitus aurium*; the first was proposed by Dr. Warden of St. Petersburg, and the latter by Prof. Joseph Gruber of Vienna.

‡ *Laminaria*, a species of alga, possessing the expansive properties of pressed sponge, and employed for the same purpose.



edge of the perforation having slipped into one of the grooves between the rim, the hearing power was now carefully tested, and the watch, which it had been necessary before the operation to bring in contact with the ear, was now heard at a distance of 9'', and the voice at a distance of forty feet, instead of three feet.

"On the 10th of February, Dr. Blake had opportunity of examining the case above described. The eyelet, which is about the diameter of a common steel knitting-needle, projects two lines or more from the membrana tympani, and rests upon the posterior inferior wall of the meatus; it is closely encircled, and firmly held in place by the edges of the perforation. The patient does not notice its presence in the least, and is entirely relieved from the previous unpleasant sensations in the ear. The ticking of a watch is heard at the distance of 14'', and words spoken in the ordinary tone of voice, throughout the length of Oppolzer's ward, a distance of about ninety feet.

"The right ear having been treated with inflation, Politzer's method, and the application of pulv. aluminis, the purulent exudation ceased, and the perforation became closed by cicatricial tissue about four weeks since. On examination, the cicatrix, 2''' in diameter, is readily seen, and the hearing power is normal. In this ear, the disease having been treated in season, the ossicles retain their normal relations to each other, and to the walls of the tympanic cavity.

"The eyelets (made by Leiter of Vienna) are of hard rubber, and from 1''' to 1½''' in length; the longitudinal opening is oval in form, and about ½''' in its longest diameter. On

the outer surface are either one (Fig. 68) or two (Fig. 69) grooves. In the outer end is a

small transverse opening for the passage of a silk thread, by which the eyelet can be withdrawn in case of its slipping from the pincette during the operation.

"The pincette used in introducing the eyelet terminates in two sharp points, which being inserted in the longitudinal opening, and then expanded, give a much firmer hold than if the eyelet were grasped from the outside."



## CHAPTER XIV.

### ON THE MECHANICAL APPLIANCES USED IN THE DIAGNOSIS AND TREATMENT OF AFFECTIONS OF THE EUSTACHIAN TUBE AND MIDDLE EAR.

THE intimate connection between the throat and the middle ear, the mucous structures of which are continuous through the Eustachian tube, and the knowledge that many aural affections are extensions of pharyngeal disease, have directed a great deal of attention to the means of gaining access from the pharynx to the middle ear, and given rise to the introduction of numerous ingeniously devised instruments for facilitating the purpose.

Some of these we have already noticed, and we will now describe the most important and practically useful of these appliances, and the manner of their employment.

And first we will describe the rhinoscopic mirror, a small piece of looking-glass or polished metal firmly attached to a convenient handle, at such an angle that when carried behind the soft palate it will receive the image of the tissues in front of it above, and to either side, according as it is inclined more to the right or to the left, forward or backward. (See Fig. 70.) The employment of the rhinal or pharyngeal mirror enables us to inspect the condition of the pharyngeal portion of the Eustachian tube and the parts adjacent. Such an examination should never be neglected in cases of disease of the middle ear, and should always precede, if possible, the use of the Eustachian catheter, as it enables us to recognize the position of the entrance of the tube, so that we may then reach it at once, instead of searching for it, as we too frequently do.

The examination of the upper region of the pharynx is by no means an easy one at all times, and cases will be met where it seems impossible to make a satisfactory exploration.



The value of the method is so great, however, that it must become incorporated into the regular manipulation of the

Fig. 70.





Fig. 71. — THE ILLUMINATING APPARATUS OF LEWIN, WITH AN AUTO-LARYNGOSCOPIC MIRROR.



aurist, or he will have the mortification of being unable to accomplish results equal to those attained by his competitors.

*Rhinoscopy* requires, in the first place, a good light. We will suppose the day to be a fine one and the sun to be shining into the examining-chamber. The patient is to be placed, sitting or standing, as may be most convenient, so that the light shall fall directly into the open mouth, and illuminate the fauces. The operator takes a position in front of the patient, but in such manner that his hand shall not cast a shadow into the throat, and proceeds with the examination in the manner to be described presently. If the light is unfavorable for a direct illumina-




tion in this way, it may be reflected into the mouth from a small looking-glass, so placed as to receive the direct rays from the sun and deflect them into the mouth. This reflector may be placed on the window-sill, and throw the light upon the opposite wall in such manner that when the patient is seated with his back to the wall the rays fall at once into his mouth.

Another plan of using reflected light is to condense some of the diffused light of the apartment upon a concave mirror, and thence reflect it into the throat. This is the manner most frequently employed; and the reflector may be attached by a universal joint to a band encircling the forehead of the operator, or it may be attached to a stand or placed upon a table.

So rarely do we have the opportunity of making these examinations by sunlight that we prefer the use of artificial light, and we may use gas or coal oil, as in Fig. 71, the latter of which furnishes the brighter, whiter, and more constant flame. Some practitioners have arranged an oxyhydrogen light for this purpose. Gas will be found sufficiently available for all ordinary and almost all extraordinary cases.

Direct illumination by artificial light is rarely employed in these examinations. The process consists in placing a condensing lens before the light, the rays of which are thrown directly into the mouth of the patient. The light is between the operator and the patient; and the instruments are carried

Fig. 72.—TOBOLD'S ILLUMINATING APPARATUS.





to one side or other of the light. The eyes of the operator are shielded from the direct rays of the light by a screen passed around the chimney of the lamp.

Reflected artificial light, however, is the method which will be found most convenient. Here the light from a lamp is received upon a concave reflector, as in the examination by diffused sunlight. The mere light of the lamp received upon the reflector is not sufficient to answer every indication; and although there are many cases in which this will be all that is required, it will be better to employ light which has been condensed before it is received upon the reflector.

Many instruments have been devised for this purpose, but the best and most convenient one will be found to be that of Tobold, depicted in Fig. 72, attached to a convenient stand, permitting of instantaneous movement in any direction, and with the reflecting apparatus above the source of illumination, as arranged for office work, by Dr. Cohen of this city.


According to the plan advised and practiced by Semeleder, the mirror is connected with the frame of a pair of spectacles in such a manner as to support it steadily before the eye.

For the purpose of conducting a rhinoscopic examination we need much the same apparatus as for laryngoscopic examination. The light is to be thrown rather lower in the fauces, so that it will shine brightly on the spot where the mirror is to be placed. The desired object is to place a mirror behind the uvula and soft palate, with its reflecting surface upward and forward, so as to illuminate the posterior entrance of the nasal passages and the parts in the immediate neighborhood, as may be seen in Fig. 73, A; by this arrangement, a ray of light, C, striking upon the mirror is thrown upon the vault of the pharynx at D. By still further depressing the handle and raising the mirror to the position in B, the ray of light, E, impinging upon the mirror, is reflected into the posterior nares at F, which point becomes visible. Now, by sliding or inclining laterally the mirror, the Eustachian orifices are shown at k, with sinus of Rosenmüller and the lateral pharyngeal walls. The primary requisite to successful examination is, that there should exist space



enough between the velum and the posterior wall of the pharynx to admit the mirror. Sometimes the hard palate extends so far back that there appears to be but a rudi-

Fig. 73.




mentary soft palate. Under such circumstances it would be impossible to make an examination of this kind.

It is essential that the soft palate should hang free from the posterior wall of the pharynx. Now, when the mouth is kept open for purposes of examination, there is an involuntary disposition to breathe through it, instead of through the nostrils as is ordinarily the case. The moment air is inhaled through the mouth, the palate becomes closely applied to the posterior wall of the pharynx, and thus the communication between the nares and the mouth is completely cut off. It is, therefore, necessary that means be adopted to secure the preservation of sufficient space between the palate and the pharynx; and this is to be done by the attempt to breathe through the nostrils while the mouth is kept open. When



the respiratory current is received through the nostrils, the palate of necessity becomes relaxed and removed from the pharynx, and the desired space is gained. This action of

Fig. 74. — RHINO-SCOPIC MIRRORS.



the palate is readily studied by observing it in a mirror. It is often difficult for the patient to succeed in breathing quietly through the nostrils with the mouth open; but a little practice, especially if the attention be called to the effect as watched in the looking-glass, will almost always accomplish the result. Should this plan not succeed, he may be told to emit what are called "nasal sounds," such as the French "en," and as the expired air passes through the nares the object will be attained. Should this fail, we resort to forcible separation by means of a broad blunt hook passed under the parts and then drawn forward by the examiner. This plan is often only partially successful on account of the spasmodic action of the palatine muscles which follows the procedure, and which is to be overcome only by repeated contact of the instrument until its presence and pressure is tolerated, or until the irritability of the muscles is exhausted. The same amount of time and patience devoted to the acquisition of the required mode of breathing will insure the success of the latter and more desirable method of securing space enough between the pharynx and the palate for the introduction of a mirror.

The mirrors to be used may be the same as employed for examination of the larynx, or like those at Fig. 74. The majority of authors insist that the mirror must be attached to its stem at a right angle or nearly so. There is no necessity for this, nor is such a mirror as convenient as one of the ordinary laryngeal mirrors. An-




other point insisted upon is that the mirror shall be about half an inch or so in diameter. There is no reason for employing so small a mirror; for greater convenience, we ought to have a series of mirrors of different sizes, as seen in Fig. 74, unless for exceptional cases. An ordinary laryngeal mirror of the diameter of seven-eighths of an inch or an inch will usually answer; but we occasionally encounter cases where a mirror of an inch and a quarter in diameter can be employed with the greatest freedom. It is only where but little space exists between the palate and the pharynx that small mirrors are needed; and in such cases, of course, mirrors of three-quarters, five-eighths, or half an inch are required; but such cases are the exception, and not the rule. That the distance to the posterior wall of the pharynx is greater than would at first sight be supposed, may be readily proved by taking a pencil or any similar object and passing it into the mouth until its extremity reaches the pharynx.

A mirror soldered to the shank at a right angle is of most use when the head is decidedly inclined forward, as has been recommended by some observers, who contend that the palate thus drops somewhat forward of its own accord. This is not often the case. If the head is kept erect, as it should be, and a mirror at a right angle to the shank is used, the hand of the examiner is much more apt to be in his own way than if the mirror is used at an obtuse angle; and if a vertical position of the reflecting surface of the mirror is desired, it is readily produced by lowering the handle, which at the same time lowers the hand, so that it is still less likely to intercept the light. On the other hand, when the object is to gain a view of the upper portion of the posterior nares, as it is called, or the vault of the pharynx, in a mirror at a right angle to its shank, the handle will have to be raised so that the mirror may not thus intercept vision. The ordinary laryngeal mirror is the best for rhinoscopic purposes. It is only a perspective image of the parts in front and above it that can be procured at best, and is easier of manipulation and more suited to examination of all parts than the mirror usually recommended.



It has been thought desirable, as a rule, to resort to mechanical means for raising the uvula and pulling it with the palate forward, in order to facilitate examination and increase the nasopharyngeal space. For this purpose a flat hook about three-tenths of an inch broad, and made of silver or other metal, is used.

Fig. 75.—TÜRCK'S PALATE NOOSE.



This hook, fenestrated or not, according to fancy, and having a turn-up edge of one to two-tenths of an inch, is passed beneath the uvula, which is received upon its broad surface and hooked under the velum, which is then gently drawn upward and forward.

Sometimes the palate is seized between the broad lips of forceps, fashioned like the palate hook.

I find but few patients who can bear these instruments without some little training, and I have discarded their use except in cases where the uvula is very long and thus obstructs the image, or in certain cases of pathological alterations, where I wish to examine everything with extraordinary minuteness.

Fig. 76.—WALES'S PALATE HOOK.



Türk proposed to remove these difficulties by adopting the use of the little contrivance shown in cut (Fig. 75), consisting of a small silver tube, about an inch long, perforated near one extremity by a small hole, and of sufficient diameter to permit the passage of a doubled thread, the "bight" of which is drawn through the small hole at one end of the tube, while the free ends depend from the other. In employing this noose, the loop of the thread is hitched over the uvula, the tube is then shoved down upon it, and the free extremities of the thread are drawn forward from the mouth until sufficient traction is made upon the velum, when they are to be attached to a band passing around the head.

Neither of these plans has succeeded in accomplishing

Neither of these plans has succeeded in accomplishing



the object in view, and indeed, from the irritability of the points about the throat, it is difficult to devise any instrument that will possess, in its application, advantages more than counterbalancing its inconvenience. Dr. Wales has found, in the little hook shown in Fig. 76, all the advantages procurable from such an instrument. We have not employed this palate hook, so that we cannot give an opinion in regard to it. It is made of light German-silver wire in the form shown. The points of the hook are intended to catch on the posterior surface of the velum on either side of the uvula, the latter fleshy process being sustained in the small circular bend of wire at the base projecting horizontally, and doubled, as it were, out of the way upon the anterior surface of the velum. The hook allows of the most thorough examination of the posterior surface of the soft palate, as the fine wire covers but little of the surface. The handle of the hook is shaped into an oval, with which the elongated uvula may be removed to one side or the other, as circumstances require, and held against the palate. The wire permits the hook to be bent into any desirable shape; for instance, if the surgeon wishes to use the hand holding the hook for some other purpose, he has only to expand the two wires, forming the handle into an oval, and bend this over the nose, when the palate will be securely held forward. This plan, of course, will be restricted to complex and difficult cases of rhinoscopic manipulation.\*

If an elongated uvula prevents examination, the best procedure for the patient's ultimate benefit is to retrench it. If any form of palate hook is employed, it must first be warmed, on the same principle upon which we warm a catheter before introducing it into the urethra.

Now, if the observer employ a palate hook with one hand and a mirror with another, how shall he introduce the catheter, should such procedure be necessary? To assist manipulation, therefore, with the palate hook in position, methods have been devised by Kramer, Wagner, and others, by which the end of the palate hook shall rest upon a con-

---


\* Medical and Surgical Reporter, p. 125, July 2, 1870.



trivance suspended from a band encircling the head, or a clamp spanning the bridge of the nose, as in Fig. 77. These inelegant contrivances are said to be tolerably well borne. I have never had occasion to resort to them.

The space between the velum and pharynx may be gradu-


Fig. 77.



ally enlarged to some extent by successive drawings forward of the velum, by means of a broad blunt hook, and repeated at intervals for several days or weeks.

An appliance all but absolutely necessary for successful examination, however, is a tongue depressor. In examinations, I very rarely resort to an instrument of this kind, letting the patient hold the tip of the tongue with a napkin as far out as possible; but in examinations of the posterior nasal passages, it facilitates the manœuvre, by securing space between the lower border of the uvula and the tongue for the introduction of the mirror and its subsequent manipulation. A tongue depressor for

Fig. 78.



this purpose should be long enough to extend to the base of the tongue, and should not be fenestrated,—one is represented in Fig. 78. It is made of one piece of hard rubber, and the handle portion is bent under, so that the hand holding it and the hand of the

instrument itself need not be in the way. It should be applied so as to depress the base of the tongue and bring it forward, and then it may be confided to the patient himself, so as to leave the operator's hands free for the mirror and palate hook, or mirror and instrument, as the case may be. *Rhinoscopes*—instruments combining mirror and tongue depressor—have been invented by Voltolini, Wagner, Mackenzie, Semiock, and others; but I have had no experience with them, having rarely encountered a case where the reflector and mirror used for laryngeal examinations, with the addition of the tongue depressor, has not enabled me to fulfill every indication of diagnosis or treatment.




Those portions of structure to examine which we resort to rhinoscopy, are the posterior surface of the uvula and velum; the posterior aspect of the nasal passages, to wit, the septum, turbinated bones, and meatuses; the pharyngeal openings of the Eustachian tubes; the vault or roof of the pharynx; and the upper portion of the posterior and lateral walls of the pharynx. These structures cannot all be examined at the same moment. I have occasionally encountered a case in which I could employ a mirror an inch and a quarter in diameter, and thus see the image of both posterior nares and both Eustachian openings at the same time. Most frequently, however, we can examine but one side at a time, though a little more may be seen in the mirror. The extreme lower portion of the septum and nares being covered by the velum, is not seen; but the upper three-fourths of the space can almost always be examined. The parts are seen more or less in perspective, this effect increasing with the horizontalization of the reflecting surface of the mirror.

If a good view is obtained, we procure such an image as is presented in Fig. 79.

It is very essential to become familiarized with the appearances represented in the rhinoscopic mirror, because this portion of the body is rarely dissected by the student, and the idea of relation of parts, as seen in the skull deprived of soft tissues, is not realized in the examination under consideration. I have known the reflection of a back tooth to be taken for the nares of that side, with the turbinated bones covered with mucus; and the ashy color was not, at first sight, unlike the appearance of the posterior nasal region in a recently boiled skull.

If we examine the engraving (Fig. 80), we find the prominent object to be a bright columnar ridge in the centre, gradually expanding above. This is the septum narium. It is of a pale-yellow or yellowish-pink color at its narrow portion,

Fig. 79.




1, Eustachian tube, enlarged; 2, the middle turbinated bones.



but as it expands, it takes the red color of the pharyngeal mucous membrane above. If we follow the outline of the expanding portion of this septum, we define on each side the corresponding posterior nasal opening, the lower portion

Fig. 80.—RHINOSCOPIC IMAGE.



1, septum; 2, free space of nasal passage; 3, superior meatus; 4, middle meatus; 5, superior turbinate bone; 6, middle do.; 7, inferior do.; 8, position of the opening into the Eustachian tube; 9, fossa of Rosenmüller; 10, lateral wall of pharynx; 11, superior wall of pharynx; 12, posterior surface of velum.

of which image is cut off by a horizontally-curved projecting red ridge, which, with as much as is reflected below, is the image of the posterior portion of the velum. Following this velum round on either side, we observe it rising over the outer portion of each nasal opening, and then bending sharply over a well-marked protuberance, formed by the internal pterygoid plate of the sphenoid bone, beneath which is the entrance to the pharyngeal orifice of the Eustachian tube. Following this protuberance, we observe it marking a canal with a fossa at its termination as it runs backward, upward, and outward, which is the fossa of Rosenmüller, and lies between this lateral prominence and the posterior wall of the pharynx. In this fossa the Eustachian catheter often becomes engaged during its introduction. To return to the central portion of the image, the parts in shadow each side of the septum represent the free cavity of the nares on each side respectively. Following this shadow from below upward on either side, we see it terminate in a larger shadow, which represents the superior meatus; the bright portion above this is the upper turbinate bone,



the lower portion of which we see turning down to become lost in a central portion, which is the middle turbinated bone partly covered by another prominent object, which is the image of the inferior turbinated bone, above which, and between it and the middle turbinated bone to the outside, we observe the shadow representing what is seen of the middle meatus. Occasionally we can discern the position of the inferior meatus just below the lower turbinated bone.

Ordinarily we cannot see deeply into the cavity of the nose; sometimes, especially when the mirror is in a vertical plane, and much of it hidden behind the velum, we can see by the side of the middle turbinated bones for a greater or less distance. The color of the mucous membrane of the cavity of the nose is a pale grayish-red or yellow; the space between the outline of the nares and the position of the Eustachian tube is yellowish-red; the opening of the Eustachian tube is red, and the other parts have the usual red color of the mucous membrane of the pharynx. The precise tint of each structure varies with many circumstances, as the position of patient, amount of shadow, and character of the disease.

Mucus is easily distinguished, and when present usually hangs about the turbinated bones, the septum, and the Eustachian opening.

In gaining a view of the posterior nares, the first thing seen is the image of the posterior surface of the uvula and palatine arches, which have much the general appearance of the outline of the image afforded by the septum and outline of the nasal openings, and if at the same time the image of a molar tooth is reflected, the deception, to one unfamiliar with the subject, is very great. If, when such an image is obtained, the handle of the mirror is gradually depressed, or the mirror slipped up farther behind the velum, we see the image of the velum gradually extending upward and then backward, not unlike a shelf, on top of which and a little in the rear we begin to recognize the true image of the septum and nares appearing in the mirror.


By turning our mirror a little to one side or the other, we



bring the pharyngeal orifice of the Eustachian tube more completely into view. This represents, when seen under favorable circumstances, a projecting nipple, in the centre of which is a distinct depression, the Eustachian orifice. In Fig. 79 is a rhinoscopic view of my own Eustachian tube; and Fig. 81 is that of Dr. Wales, who has kindly permitted me to use it.

We occasionally may succeed in passing the Eustachian

Fig. 81.



A, roof of the pharynx; B, superior turbinated bone; C, middle turbinated bone; D, inferior turbinated bone; E, Eustachian tube; F, septum narium.

catheter with one hand while holding the rhinoscopic mirror in position with the other, and in this way may direct the point of the catheter into the orifice under sight; but cases that can tolerate this double manipulation are rare, and experts who are able to perform it still rarer. Still, the fact that it can be occasionally done is indisputable, and it is therefore well to attempt the manipulation in cases that appear favorable.

“Before closing what we have to say on the instruments required in rhinoscopy, it will be proper to call attention to the fact that there is an anterior as well as a posterior rhinoscopy, which in certain cases may be had recourse to with great advantage. Dr. Markusovszki invented a speculum for this purpose, resembling, in its mode of construction, an ear speculum with two compartments; but its size is much greater, and the divisions are larger and flatter. With this instrument, and a concave mirror to illuminate the nasal apertures, Czermak has been able to inspect the inferior turbinate bone and its free border, which passes to the extent



of seven-twentieths of an inch above the floor of the nasal fossæ. In one case wherein the nasal cavity was very capacious, he saw up to the posterior part of the pharyngo-nasal cavity, and upon a cadaver he employed a small mirror, its reflecting surface being turned above and in front, and arranged in the space situated under the inferior turbinate bone, in such a manner as allowed a clear view of the nasal orifice of the lachrymal canal, which was marked by a hog's bristle, that had been previously introduced.

"This sort of exploration has hitherto been neglected, as the specula employed have been ill constructed to obtain the desired results. Voltolini employs short polished tubes, ear specula, introduced into the nostrils, and a concave reflector. He has thus succeeded in getting a glimpse of the posterior wall of the pharynx, and the swellings at the mouth of the Eustachian tubes.

"I have employed several sorts of specula, but none seem to me to deserve the same commendation as the one devised by Dr. Thudichum, illustrated in the cut, Figs. 81 and 82.

THUDICHUM'S SPECULUM—FRONT AND PROFILE VIEWS.

Fig. 82.





Fig. 83.



He describes it as consisting of two valves held together by a spring, and when compressed, form a canal of oval bore, about three-fourths of an inch in length. The valve which expands the ala is from one-quarter of an inch to three-fifths longer than the valve which is placed against the septum. This latter inner valve should never be long enough to reach



the sensitive part of the mucous membrane of the septum. It follows from this that each nostril requires its own speculum. As the nostrils of different persons vary greatly in size, different sizes of specula must be at hand in practice. The speculum is inserted into the nostril while closely compressed; the moment the spring is liberated, the nostril is fully expanded, and if the operator now draws the spring (which serves also as a handle) upward, the cavity of the nose can be fully inspected. The speculum, therefore, is designed exclusively to expand the atrium or membro-cartilaginous part of the nostril; if it were allowed to enter beyond this, and to reach the isthmus formed by the unyielding cartilaginous septum and the nasal margin of the ascending branch of the upper maxillary bone, it would cause pain, and narrow, by its own thickness, this small passage. The new specula cause no pain, widen the area to the utmost extent, and admit of the introduction through them, into any part of the nose, of instruments for slight, delicate, or severe operations.”\*

The Eustachian catheter is an instrument of frequent employment, intended to enable us to pass a current of air, of medicated vapor, of steam, of spray, of water, of powder into the Eustachian tube, or through it into the cavity of the tympanum. These instruments are passed either through the mouth or through the nostril, the latter being the method most proper and most used.

The instrument for catheterizing the tube through the mouth is that of Dr. Cutter, of Boston (see Fig. 84), which is intended to be used with the rhinoscopic mirror. The terminal portion can be turned to any angle, so as to meet any desired radius, which may facilitate its introduction. We have made several attempts to employ this instrument, but as they were unsuccessful, we soon reverted to our usual method of catheterizing through the nose.

The catheter generally employed is that of Kramer (see Fig. 85).

---

\* Med. and Surgical Reporter, op. cit.



Fig. 86 is a small Eustachian catheter for use in young children.

Fig. 84.




Fig. 85.






Fig. 86.





The rules laid down by authors for the manner of introducing the Eustachian catheter are very unsatisfactory. The position of the pharyngeal orifice of the tube differs, as can be demonstrated by a rhinoscopic examination, and therefore no one rule is of universal application. Experts in aural surgery, who make themselves familiar by hard study with the anatomical arrangements of the upper portion of the pharynx, are very often able to find the entrance of the tube by a sort of knack which nothing but experience can give. The general practitioner, who has perhaps never seen this portion of the body exposed, for it is rarely dissected in the study of general anatomy, and whose ideas of the relations of the parts are gained from the skull and from engravings, will seldom succeed in the manipulation; hence the discredit

Fig. 87.—SECTION OF THE HEAD, WITH EUSTACHIAN CATHETER INTRODUCED.



A, showing the turn which the catheter should make in entering the pharyngeal orifice of the Eustachian tube; B, showing the position of Rosenmüller's fossa, and the necessity for withdrawing the catheter before entering the Eustachian tube.

which has fallen upon the instrument. The fault is not in the catheter, but too often in the operator.


The most popular rule is that of Kramer, to touch the posterior wall of the pharynx, then turn the back of the in-



strument toward the external auditory orifice and pull it toward you, it will slip over the posterior portion of the cartilage of the tube, and if it is only allowed to ride on the anterior face of this posterior lip of the cartilage, it can then be pushed on into the orifice of the tube, as shown in Fig 87. We prefer to turn the beak of the instrument as soon as we enter the pharynx, to graze the lateral wall with it as we proceed backward, mounting the anterior lip of the tube. In this way we avoid catching the instrument in the fossa of Rosenmüller. If the catheter remains in position, it is probably in the tube. This can often be determined positively by making a rhinoscopic examination, which will show us if it is in the orifice; if not, it will show us the relation the beak of the catheter bears to the orifice, and we can readily make the correction. Where the tube is impervious, so that a current of air cannot be passed into it, this would be almost our only method of verification,—an important point if we wish to force a stylet through the tube.

As an instrument of diagnosis the Eustachian catheter is used to determine the permeability of the Eustachian tube, as in Fig. 88. This is done by blowing into the catheter, producing a certain sensation easily recognized by the patient. A flexible tube is sometimes used in this connection, and in addition a second tube, reaching from the ear of the patient to the ear of the operator, is employed, so that the latter may hear the effect of the current, as in Fig. 89. The whole instrument is known as Toynbee's otoscope. The accompanying illustrations exhibit these tubes and their modes of employment.

Fig. 88.—EXPLORER, WITH EUSTACHIAN CATHETER ATTACHED.




#### IMPERVIOUS EUSTACHIAN TUBE.

“Of the means more especially used to open an impervious tube, the nitrate of silver of the strength of forty or eighty grains to the ounce, applied to the mouth of the tube by



means of the Eustachian catheter, is considered by Dr. Pomeroy,\* of New York, one of the best. He recommends



1, the patient; 2, Eustachian catheter; 3, the surgeon; 4, the india-rubber diagnostic tube or Toynbee's otoscope.

its use in the following manner, and we have tried the method and approve of it:

"The end of the catheter containing the ring should be supplied with a thimble-shaped piece of rubber, similar to that used on atrophine-dropper; the beak of the instrument is immersed in the solution, and one or several drops may be drawn up, the end freed from excess of solution by wiping on blotting-paper, then passed up to the Eustachian tube, or perhaps a little way within, and the contents discharged. Or, after loading the instrument, a bit of cotton may be wound upon the end, and when introduced the thimble-shaped bulb may be slightly pressed, so as to moisten the cotton when a swab is extemporized, and being exactly in position, the remedy is readily applied."

---

\* The Treatment of Naso-Pharyngeal Catarrh in its Relation to Diseases of the Ear, by O. D. Pomeroy, M.D., Medical Record, March 15th, 1871, p. 30.




## THE PHARYNGEAL OPENINGS OF THE EUSTACHIAN TUBE.

Having had the opportunity of examining and studying some interesting cases of cleft palate, we have come to the following conclusions: that there is a peculiarity of articulation or nasal voice. This is accompanied by more or less deafness. There is also disease of the mucous membrane, which is congested, harder and dryer than natural, owing to constant exposure to the air.

The muscles have not as much power to control the pharyngeal openings of the tube, so that in the act of swallowing they remain so long open that in two cases particles of food passed along the dilated tube; and was stated by one of the patients to have been removed from the external meatus. In both these cases perforation of the membrana tympani had taken place.

In the case of Mary N., aged fifteen years, of Rome City, Indiana,—a patient of Dr. Cohen,—by simply drawing back the cheek we could distinctly see through the cleft both mouths of the Eustachian tubes, and watch their action. These openings were triangular or obliquely oval, looking downward, outward, and forward. They showed that, to enter these openings by the catheter, required it to be curved for three-fourths of an inch in depth, in a downward direction, for the terminal two inches, with the tip curved one-eighth of an inch to the right or left, to enter with facility into the mouth of either tube.\* (See Fig. 90.) It also required a slight rotary movement to the right for the left opening, and vice versa to enter; but once entered, it retained its position firmly, in spite of violent shaking of the head. The ordinary catheter of Kramer† is of six and a half

Fig. 90.



\* These catheters were made by G. P. Pilling, gold and silver surgical instrument-maker of this city.

† Kramer on Diseases of the Ear. Translated for New Sydenham Society, London, 1863, p. 27.



inches in length, with a beak bent laterally for three-quarters of an inch at an angle of  $144^{\circ}$ ; this could not be introduced in this case without painful pressure upward of the soft tissues of the nose; and yet all the parts were perfectly normal in appearance. Although this was an independent observation to study the feasibility of an improvement in the curve of the catheter, it was found, on consulting some old works, that *Saissy* pictured a somewhat similarly-shaped catheter in his celebrated "Essay on the Diseases of the Ear," published originally as an article in the "Dictionnaire des Sciences Médicales," in 1819, and translated with additions by the distinguished surgeon, Prof. Nathan R. Smith, of Baltimore,

Fig. 91.





Fig. 92.



from the posthumous edition published by "Montain" in 1827.



Both the catheter of Saissy and that employed in the case referred to, differ in their curvature from the catheter of Toynbee.


In one other case examined by Dr. Cohen one year after a successful operation for closure of both hard and soft palate, he found a perforation of the membrana tympani. The deafness was moderate, but marked; and on questioning the patient as to the entanglement of food in the Eustachian tube, it appeared that it had also occurred in this case previous to the operation; but that since the operation there had been no difficulty of this kind.

These observations do not confirm those of Dieffenbach, that the impairment of hearing was always completely relieved after closure of the fissure of the palate by sutures. They show, however, the importance of an early operation, before the tissues have received such profound alteration that they are unable to return to their normal condition.

Politzer has devised an apparatus, which we have before described and frequently employed, for examining and inflating the middle ear through the Eustachian tube by the nose, which, though less exact than the catheter, often answers the purpose. His method consists in placing a tube attached to an elastic bag with a valve at its base into one nostril, compressing the other upon it, and forcing a current of air as the patient swallows saliva, or a mouthful of water held for that purpose. This is of easy application, and suitable to practitioners who cannot manage the catheter.

Fig. 91 exhibits the instrument, and its manner of application; Fig. 92 shows Dr. Brunton's\* modification, which

Fig. 93.




\* Braithwaite's Retrospect.



the patient can employ without the aid of the physician, the compression of the bag being made by the foot the instant the patient swallows, which is well illustrated in the cut.

Fig. 94.



A still more recent change in the form of this apparatus has been made by Dr. Peter Allen,\* which we have received from John Weiss & Son, of London; but we can express no opinion of its utility until further trials. It is described as follows: "I have still further improved Politzer's most invaluable appliance, by substituting a *nasal pad*, which is pressed *against* the opening into the nostrils, for the tube which he inserted into one of them. Mounted on a strong piece of covered copper wire are two air pads, which can be so approximated or separated as to stop up conveniently the nasal orifices. The metal serves as a handle for the surgeon or patient. Through each pad runs a hole, and these holes communicate with two short bits of india-rubber tubing, joining into a single tube. Into this tube the pipe of the inflating-bag is inserted, and the apparatus being thus complete," is shown at Fig. 93.

For the introduction of minute quantities of liquids, or the spray of the same into the middle ear, we have several instruments. It is most generally accomplished through the Eustachian tube. A very convenient instrument will be found in the spray catheter,† in which Richardson's apparatus is placed within a catheter, which is movable in any direction. The difficulty in the use of this instrument is that the end gets clogged by mucus, and thus arrests the operation, requir-

\* Lectures on Aural Catarrh, by Peter Allen, M.D., Aural Surgeon to, and Lecturer on Aural Surgery at, St Mary's Hospital. London, 1871, pp. 78, 79.

† Made by Messrs. Codman & Shurtleff, of Boston.




ing the removal of the catheter; to avoid this, rapid pumping will sometimes dislodge the mucus.

A still better form of catheter is shown at Fig. 94, which can be freed from obstructing mucus without removal from the Eustachian tube. The tube is double, so that one stream is passing in while another is passing out. No. 1 is the exit tube; No. 2, double orifice for entrance and exit of fluids; No. 3, tube for air-bag, or syringe for introducing liquids.

We can with this double catheter wash out the pharyngeal extremity of the Eustachian orifice, or introduce a few drops of liquid and spray it by means of the air-bag and drive it in this state into the middle ear.

Another form for the same purpose is called Bishop's\* nebulizer, with which he states lotions of any strength, up to the point of saturation, may be used; but the quantity re-

Fig. 95.—BISHOP'S NEBULIZER FOR EUSTACHIAN TUBE.



quired to come into contact with the whole lining membrane is so small that there is not the least fear of mechanical injury, and by slightly turning the point of the instrument the pharynx, the nasal passages, and the parts connected therewith, may all be subjected to the action of the remedy. The apparatus, as seen at Fig. 95, consists of a small graduated glass syringe, similar to that used for subcutaneous injection. The solution is gently forced drop by drop into a

---

\* On a New Method of Applying Remedial Agents to the Cavity of the Tympanum, by Edward Bishop, M.D., pp. 8, 9.



cylinder, and at the point where it leaves the nozzle of the syringe it is caught by a current of air sent by a pump worked by the hand. This drives the fluid forward in a pulverized state. The apparatus is then attached to an ordinary Eustachian catheter, previously introduced,—the end of the catheter being covered with a fine gauze wire.

Before closing what we have written upon the subject of the Eustachian catheter, we take pleasure in noticing briefly the form devised by Dr. Henry D. Noyes, of New York,\* “especially when the catheter is introduced by the opposite nostril:”

“I may with modesty remark that I have not found the forms figured by Kramer for the ordinary instrument the most suitable; the error, in my judgment, being that the beak is too long, and the sweep of the curve too great.

“The catheter of  $1\frac{1}{2}$ ”, or a little more than 3mm. in size, I have come to prefer. For a larger size I would have the beak a little larger, the curve the same. Curved in such a manner, the catheter finds entrance readily to the mouth of the tube, and when fairly engaged can be pushed up into it by the thumb and forefinger.

“The change needful is simply to give the beak a double curve. Take the elongated beak generally recommended, and, at three-eighths of an inch from the point, bend it in a plane transverse to the plane of the primary curve. This bend will always be upward when the catheter lies flat upon the floor of the nostril,—but the same instrument will not serve for both sides, because the secondary curve is in opposite directions when used for the right or left sides.

“When introducing the catheter, it is needful to keep the point close to the septum as well as to the floor of the nostrils. Arrived at the posterior edge of the septum, the beak should wind closely around it,—curving obliquely across and turning upward, so as to point toward the Eustachian orifice.

“It is better always to use the left hand for the left Eustachian tube, and the right hand for the right.

---

\* Transactions of the American Otological Society, 1870, pp. 81-84.



"I find it better not to push the point back to the pharynx and then draw forward, but rather to aim at once for the ostium when the mark on the straight portion of the instrument shows that the beak is near its destination. I always inspect the nostril with speculum and mirror before putting in the catheter, so as to learn the peculiarities of the passage; and it is advisable to dip the end of the catheter in glycerine to lubricate it. I always give the patient the usual directions about holding the mouth a little open, to aid relaxation of the soft palate."



## CHAPTER XV.

### WOUNDS—INJURIES—PERFORATIONS OF THE MEMBRANA TYMPANI—ACUTE AND CHRONIC INFLAMMATION.

WOUNDS of this membrane are rare, owing to its situation, the depth of the meatus, and its tortuous canal.

It is injured in fractures, by blows on the side of the head, falls, concussions, or sharp vibrations of air.

Sir Astley Cooper relates a case of laceration of the membrane by a blow on the side of the head, extending from above downward across the entire membrane.

I have seen a number of cases in which there were both lacerations and cicatrices of the membrana tympani. As a general rule, this form of wound heals with great facility, without treatment. The same result from disease would take a much longer time to heal. There is a form of deafness, the result of loud, sharp reports, to be found in both officers and men belonging to our artillery corps; but in raw recruits it assumes the form of hemorrhage from the meatus from laceration of the membrane.

### RUPTURE OF MEMBRANA TYMPANI BY STRANGULATION.

Dr. Alfred Taylor, in his Medical Jurisprudence (Philadelphia edition, 1856, p. 575), mentions a case of *suicidal* strangulation in which constriction had been produced by a ribbon, and the violence applied was sufficient to produce bleeding from one ear. On dissection, there was found to have resulted a rupture of the membrane of the drum of the ear.

Von Tröltsch mentions two instances of perforation by a blade of straw; and I have met with a similar injury in a boy. Another case which came under my observation was



where the membrana tympani was ruptured by a blow from a pillow, thrown with great force against the ear.

*Treatment.*—Absolute rest, depletion of a local character, if there be pain, one-quarter to one-half grain of morphia injected hypodermically, with a solution of sulphate of zinc, gr. i to x in one pint of water, applied warm, by means of Clarke's douche. Cotton or wool should be placed in the meatus, to exclude the air, and careful attention required until union is complete, for if neglected it is apt to be tedious. The membrane, if properly treated, will heal without difficulty.

The manubrium, or the handle of the malleus, is inclosed for about three-fourths of its length between the laminæ of the membrana tympani.

This delicate bone was first found to have been broken and united in the case of one of the lower animals. Soon afterward a similar fracture was discovered by Mènière\* in a gardener, who had accidentally thrust a twig of a pear-tree into his ear. A very extensive laceration of the membrana tympani took place, and the little bones of the ear could be plainly seen, and their movements distinguished. It is stated that this injury healed of itself, without any special treatment.

Prof. Von Trötsch also reports a case of united fracture, caused by the thrust of a pen-handle. The translator of Von Trötsch's work furnished a case from Dr. Weir, of New York, of ununited fracture of the handle of the malleus.

"The man received a fall, which caused unconsciousness and bleeding from the ear. He had pain in his ear for sixteen hours, running along the forehead to the ear; was laid up for a month, and the pain passed away. The tinnitus aurium is very great. On examining the right ear, the handle of the malleus seems to have been fractured just below the short process, where an irregularity is seen, and the lower fragment moves with great freedom upon the

---

\* Gazette Médicale de Paris, 1856, No. 50; Von Trötsch, Treatise on Diseases of the Ear, translated by John Roosa, M.D., Am. ed., pp. 151-153.



upper. The irregularity disappears after inflating the ear by the Valsalvian method. On the posterior part of the drum a patch of increased whiteness is seen, and possibly it indicates the site of a rupture of the membrana tympani."

INFLAMMATION OF THE MEMBRANA TYMPANI (MYRINGITIS),  
ACUTE AND CHRONIC.

Having already treated of this important membrane when considering acute affections of the meatus auditorius, otitis, acute and chronic, externa and interna, we shall not devote as much space to the acute as to the chronic affections of the membrane; as the latter subject has assumed great importance from the investigations by Toynbee into the change produced by this form of inflammation, in which he has been followed by Wilde, Von Tröltsch, and still more recently by Politzer, who has surpassed them all in the care, accuracy, and completeness of his investigations of the subject in reference to a true diagnosis.

Acute affections of the ear are very amenable to therapeutic agents; but in chronic affections of all organs it is most difficult to arrive at a correct diagnosis and successful treatment,—the ear being no exception to this rule. As is well observed by Politzer, "An accurate examination of the membrana tympani is indispensable for a complete and exhaustive diagnosis;" and we can fully agree with him, when he says that, in many cases which he has examined, there are changes in the membrane to which no impairment of function whatever corresponds. On the other hand, experience teaches us that, in some cases, with a marked degree of deafness, the membrana tympani does not present the least departure from the normal condition; so that the condition of the membrane should only be taken in conjunction with all other symptoms, but is never to be depended upon alone as a safe guide in diagnosis.

In acute inflammation of this membrane, there is always great vascularity and pain; usually the attack of pain is in the night, and is of the most excruciating character, producing at times delirium. It is increased by chewing or swallowing, or by the least pressure behind the ear, which is



the only means of diagnosing it in little children. There is a feeling of fullness and bursting within the ear. If not treated promptly, the pain extends to the throat and mastoid region, or may pass into the brain. Almost every patient has some form of his own for expressing the noise or tinnitus aurium felt in the ear. In every instance there is deafness, either partial or complete, of the side affected, as in most of the cases it is one ear that has been exposed. The chief cause of this disorder is cold, but other agencies will produce it, as foreign bodies, attacks of various diseases, the exanthemata, bronchitis, typhoid and typhus fevers, and phthisis. In almost every instance there is more or less swelling of the lining membrane of the external meatus, and therefore we cannot, in the very early stage, see the membrana tympani. From numerous examinations, where they can be borne by the patient, or when under the influence of chloroform or ether, it is seen that the membrana tympani loses its polish and its semi-transparency, and assumes a pink, and subsequently a rose tint, with numerous vessels crossing it, forming a network, commencing at the insertion of the malleus, with here and there patches of ecchymosis. If it progresses, exudation of lymph and muco-purulent secretion takes place with perforations, either by abscess or slough. As soon as the perforation occurs, there is temporary relief. If resolution takes place, the vascularity decreases, and the membrane ultimately recovers its natural appearance, but is apt to assume a muddy or opaque character, or is covered with numerous spots.

*Treatment.*—Our sheet-anchor is local depletion, with purgatives, and relief to the pain by morphia, or hydrate of chloral, or the hypodermic injection of sulphate of morphia; the use of moist heat by the spray or douche; subsequently blisters, with iodine and tonics, to promote absorption of the deposits. The adhesions are to be broken up by the use of warm water by the catheter or Politzer's apparatus. After all inflammation has disappeared, the air douche must be employed, and the opacity dusted with calomel, while we administer internally the iodide of potassium, in five-to ten-grain doses, three times a day, largely diluted with water,



until local inflammation is again set up, when we must let the parts rest for a time.

#### CHRONIC INFLAMMATION OF THE MEMBRANA TYMPANI.

In this form of inflammation there is one important element wanting, and that is pain, and there are but few exceptions to this general rule. It is characterized by a general thickening and opacity of the membrane, progressing for months or even years, giving no warning except a gradual and slowly increasing deafness, which is sometimes noticed by the friends of the patient before he is himself aware of it. Sometimes we have a collapse or falling-in of the membrana tympani, with anomalies in the transparency and color. These anomalies are in part owing to the arrangement of the several layers as before noticed,—an external dermoid and an internal mucous: the dermoid is an extremely delicate continuation of the lining of the external meatus, which, in its transition to the membrane, consists of an epidermic layer, a Malpighian mucous layer, and the cutis. The middle fibrous layer consists of two separate laminae, an external radiate, and an internal circular.

The first form of chronic disease of the membrane, according to Politzer, is softening of the epidermic layer, the appearance of which becomes dull and lustreless, and is generally the result of serous exudation. It occurs in the beginning of acute catarrh of the cavity of the tympanum, in the rare forms of idiopathic myringitis, and in the outset of otitis externa.

Then follows congestion of the dermoid layer, which is generally seen on the handle of the malleus, indicating chronic inflammation of the tympanic cavity. There is also a circular wreath of vessels near its periphery, from which little converging branches run toward the centre of the membrane; this is especially the case in acute catarrh of the tympanic cavity, or in relapses of chronic catarrh.

*Case of chronic inflammation of the tympanic cavity, with collapse of membrana tympani.*—June 4th, 1869. E. M. D., aged fifty-eight, residence Chelton Hills, Pa., applied on account of deafness of right ear, which has been of eighteen months'



duration,—the result of cold. Has been under treatment for one year without benefit.

*Examination.*—Right ear: cannot hear the ordinary watch close to the meatus when the other ear is closed, or on the temporal bone, but can hear the tuning-fork. Left ear: can hear the watch\* fifteen inches; can only hear one sound or one individual voice at a time; but if more than one, the sound becomes confused, and he has a sensation of closing or shutting in the ear. His hearing, when young, was so acute that he could distinguish, even in an adjoining apartment, the faintest sound, so that his friends had to be very careful what they said when he was near. Meatus dry, but of normal color; membrana tympani very concave; reflection of light feeble; handle of the malleus not well seen from congestion of its vessels; sounds from Eustachian tube of left side normal; of right side faint and moist.

*Rhinoscopic examination.*—Mucous membrane swollen; orifice of Eustachian tube covered with gray mucus; follicles of the pharynx red and enlarged.

*Treatment.*—Chlorate of potash five grains three times daily; iodine and ether, by means of Politzer's air douche, through the Eustachian tube.

June 18th, 1869. Treatment continued; no change noticed.

Sept. 21st. Slight improvement, could hear the watch close to right ear.

October 6th. Directed the use of Politzer's air douche, so as to cause the air to reduce the concave character of the membrana tympani. Prescribed an alterative mixture, as follows:

R.—Iodide of Potassium, grs. v;  
 Sol. of Arsenite of Potassa, gtt. v;  
 Muriate of Ammonia, grs. x;  
 Extract of Licorice, grs. x;  
 Water, ℥ij. M.

Sig. Three times a day.

He continued to take this mixture at intervals for a month, having the current from Kidder's battery applied to the ear

---

\* Ordinary hearing distance of this watch is from two and a half to three feet.



every third day in the following manner: the conductor leading to the positive pole or anode was held in the patient's left hand, whilst the opposite conductor or cathode, ending in a funnel-shaped conductor, was placed in the meatus of the right ear, which was filled with warm salt-and-water. The current was by this means brought to bear upon the membrana tympani, and made to pass directly through the head, instead of making a short cut through the skin. Subsequently the battery was applied in the same manner every sixth day.

Oct. 13th. Has rheumatism from exposure,—not the first attack; directed bath of bicarbonate of potassa, and continuance of mixture, with astringents to the throat.

Oct. 26th. Can hear the watch about three inches from the ear.

Nov. 17th, 1869. Is in high spirits; hearing distance for watch ten inches; membrana tympani less concave; now able to hear any one of his family at table five or six feet off. Same treatment continued up to January, 1870, when he was discharged much improved.

February, 1871. Hearing continues good, his brother-in-law informing the author that he considered his hearing all right.

Next to congestion of membrana tympani, Politzer places ecchymoses that occur in catarrh of the middle ear.

He considers their recognition, by inspection of the membrane, to be very easy. They appear as irregular hemorrhagic spots, in varying number, sometimes sharply defined, sometimes faint, and especially frequent behind the manubrium.\*

In still higher grades of acute catarrh, especially when accompanied by purulent secretion, the membrane, before perforation, appears so reddened that individual ramifications of vessels can no longer be seen. The same good authority states that the general opacities of the membrana tympani depending on a lack of transparency in the *substantia propria*, arise either by immediate transmission from the

---

\* Politzer, translation, op. cit.




other layers of the membrane, or are secondary to affections of the deeper-lying structures; and they also occur independently in old age. "On examination in the living subject, the membrane thus changed appears flattened and uneven, in consequence of several unequally-clouded portions. The color is dirty bluish-white like lustreless porcelain, or the yellow predominates, and the membrane may be compared to a lamina of leather."

OPACITIES FROM CHANGES IN THE MUCOUS LAYER OF THE MEMBRANA TYMPANI.

In acute catarrh of the cavity, without perforation, the membrana tympani has a bluish-red color, on account of congestion of its mucous layer. In chronic cases, and these, according to Dr. Politzer, are the most frequent, the membrane is either bluish-white, with a tendinous opacity, or dirty-gray, and usually dull, having the appearance of ground glass, or glass covered with vapor upon one side.

The following illustration is from an interesting microscopical section of a diseased membrana tympani, in which

Fig. 96.



Section of a fatty degenerated tympanic membrane of a patient suffering from a chronic catarrh of the cavity of the tympanum. *e*, epidermis; *d*, dermoid layer; *r*, radiating layer; *c*, circular layer; *s*, layer of mucous membrane.

we have simple fatty degeneration, from Prof. Gruber,\* according to whose investigations it is a condition frequently connected with chronic aural catarrh.

\* *Lehrbuch der Ohrenheilkunde*, p. 461.



In the dermis, membrana propria, and mucous membrane are seen partially crystalline and partially scaly masses of fat, by which the membrane becomes very tender, and tears readily.

Atheromatous or calcareous deposits, according to Wilde,\* are seated in the anterior vibrating portion. This deposit is of a yellowish color, and has a sharp, well-defined edge, totally different from that of a lymphic exudation, which generally shades off into the surrounding membrane. If scratched with a cataract-needle, it will be found to be gritty, its composition being carbonate of lime. Wilde believes that he was the first to notice this peculiar condition, which he stated in his essay upon Otorrhœa, in 1843. Toynbee has given three illustrations of this form of deposit, and states that it is found in the fibrous lamina, and that in some instances the whole of the membrana tympani is converted into calcareous matter. This change may occur at all periods of life, sometimes taking place when the remaining portion of the membrane is healthy. In other cases he considers it symptomatic of calcareous deposit within the tympanic cavity. In others, it follows, and appears to be produced by, chronic inflammation of the dermoid layer with or without catarrh. Politzer says, the knowledge of their occurrence appears to be of earlier date than is generally supposed. In an old work of Cassebohm† (1734), he finds a description of a preparation of the membrana tympani of an old woman, upon which there were calcareous deposits in front of the manubrium and behind it. Von Tröltsch gives a more accurate account. He says, such calcareous deposits are, for the most part, distinctly separated from the surrounding tissue, and with their yellowish-gray or clear, whitish appearance, are not to be mistaken. They occur even in childhood, and are not at all infrequent.

With the exception of a few cases in which the hearing was tolerably good, he has hitherto found them only associated with a high degree of deafness; so that the existence

---

\* Wilde, Practical Observations on Aural Surgery, p. 276.

† Tractatus quatuor anatomici de Auri humani, 1734.



of similar earthy processes on the membrane of the fenestra rotunda, or about the foot of the stapes, may be inferred. Moos, of Heidelberg, was the first to study carefully the gradual formation of these calcareous deposits; the case selected was an old woman seventy years of age, in the course of a chronic catarrh of the middle ear. Since that statement, Dr. Politzer has had a similar opportunity of following up the development of these deposits.

The following case is from Politzer's clinic, and is worthy of notice, from the fact of the patient's retaining an almost normal degree of hearing, notwithstanding the existence of a serious lesion:

"Woman, aged twenty, a patient with tuberculosis, in Oppolzer's ward, Vienna General Hospital. The aural examination was made in this case, not from any symptoms of disease of the ear, but in accordance with the general custom in this clinique. With both ears, the watch was heard at a distance of one foot, and the voice, in a whisper; the length of the ward about ninety feet. Examination of the membrana tympani showed in each a perforation about  $2\frac{1}{2}$ " in length, and situated behind the manubrium mallei; the remainder of the membrana, with the exception of a small rim at the periphery, was infiltrated with calcareous deposit. When a child, the patient had severe pain in both ears, followed by a discharge, but no deafness.

"Death occurred a few days after this examination, and an autopsy confirmed the observations above described. In the fresh state, the calcareous mass was so firm as to be immovable under a strong pressure, and the deposit had not only impregnated all the layers of the membrani tympani, but extended above the surface of the mucous membrane into the tympanic cavity, and the malleus was so firmly imbedded that no vibration could have been transmitted by it to the other ossicles. The patient was able to hear, and had heard since childhood, inasmuch as the perforation permitted vibrations to pass into the tympanic cavity, and fall directly on the stapes, which was freely movable. The labyrinth was perfectly normal. Had the opening been closed by cicatricial tissue, it would have been necessary to make an artificial perforation,



in order to permit the passage of vibrations and to give to the patient the degree of hearing which, as the case stood, she possessed up to the time of her death. The importance of retaining or reproducing a perforation of the membrana tympani in some, and by no means infrequent diseases of the ear, is illustrated by this case. It also shows that opacity in the membrana tympani alone is not always a cause of deafness.”\*

Schwartz has confirmed the previous observations. They are not always an attendant upon, or an absolute cause of, deafness, for Politzer has found them after death in persons whose hearing had been normal. He regards them as the product of otorrhœas that have passed away. Besides the circumscribed opacities, we find tendinous, gray, oblong, and uniformly faint or striped areas, which are separated by either normal or slightly-clouded portions; a condition such as is often met with in chronic catarrhs, and after otorrhœas, but also frequently enough in persons of normal hearing, especially in old age.

“As regards diagnostic value, the result of clinical investigations may be summed up in the following points:

“1. Diseases of the membrana tympani are, for the most part, combinations of the signs of disease of the external and middle, less frequently of the inner ear.

“2. Opacities occur frequently in aural disease, according to the unanimous testimony of authors, and, in many cases, afford important data for a diagnosis.

“3. The fact, however, that analogous opacities occur also in persons with normal hearing, diminishes their diagnostic value not a little. Nevertheless, they deserve, in given cases, full consideration; since, taken in connection with other signs, with the nature and course of the case, and the degree of functional impairment, they often essentially facilitate the diagnosis.”†

---

\* C. G. Blake. M.D., Boston Med. and Surg. Journal.

† Politzer, op. cit., p. 69.



## PERFORATION OF THE MEMBRANA TYMPANI.

The most frequent cause of this solution of continuity of the membrana tympani is purulent catarrh of the middle ear, and the disease which produces the largest number of cases of purulent catarrh is scarlet fever; next to this is cold and its consequences; then follow phthisis, measles, pneumonia, typhus and typhoid fevers, pertussis, and, in a small number of cases, eczema, puncture, etc. In almost all there is a softening of the tissue; and the pus, or mixed pus and mucus, in the cavity of the tympanum or middle ear, exercises a strong pressure on this softened membrane, until laceration takes place.

In phthisis, the cough will often cause a sudden pressure of air in the cavity, which may terminate in a rupture of both membranes. I recently watched the ruptured membrane in a case of phthisis, and there was a gradual increase of the size of the perforation every few days, until there was but little left of it. In this disease pain seldom, if ever, attends the destruction of the membrane. In the post-mortem examination of such cases, Politzer has found the mucous membrane of the middle ear pale, seldom hyperæmic, and covered with a thin layer of creamy muco-purulent secretion; and in the membrana tympani, usually at its lower segment, an irregular perforation, from a line to a line and a half in diameter, the border of the opening being softened and collapsed in folds.

The occurrence of the perforation of the membrana tympani is usually indicated by pain in the ear, accompanied with more or less deafness. Examination of the ear with a speculum in a bright sunlight, or with the concave mirror, shows the perforation either in the centre or through the lower half of the membrane. In acute cases the author has seen the first faint dark line or spot which indicates perforation. This was shown to be such by the patient blowing air through it, on closing the nose and mouth, and making forced expiration. The cases most difficult to heal are those following scarlatina; while those produced by acute catarrh, or puncture, heal more readily, if the perforation is not of large size. When more than one-half of the membrane is destroyed, it rarely heals.



Recently I have had under treatment two cases which healed in three months; but the majority require many months, and even years, to accomplish the closure. In another place the exanthematous affections of the middle ear have been fully treated of, and the great importance pointed out of giving early attention to them.

The treatment of simple subacute perforations will depend somewhat upon the cause; but generally they require antiphlogistic measures: leeches or cups, with warm fomentations and brisk purgation. Opium or its salts, with or without antimony, at bedtime, will hasten the cure. The after-treatment will consist of blisters behind the ear, with a mild astringent wash of sulphate or acetate of zinc, with the addition of tinctura opii; this treatment applies to all cases of acute perforations.

*Chronic perforation.*—In cases of perforation of long standing, we resort to solutions of nitrate of silver, 20 to 30 grs. to fʒi of water, or solutions of iodine or carbolic acid, applied with great care, by means of a brush or piece of cotton, not oftener than every third day. At the same time we should attend carefully to the constitutional treatment by alteratives and tonics; and keep the pharyngeal orifice of the Eustachian tube open and free from mucus, by means of Politzer's air-bag or its modifications, so that there may be a free connection with the throat. Thudichum's nasal douche will be found useful, if the nose and throat are affected, as is frequently the case.

Syringing, as a general rule, should be forbidden, and, when necessary, the modification of Thudichum's apparatus by Dr. Clarke should be preferred for the purpose. The danger from syringing is the liability of increasing the size of the perforation. The pus and other matters that accumulate in the ear should be removed at least daily by the use of a little cotton or soft sponge moistened in tepid water; and, to remove fetor, a few drops of Burnett's solution of chloride of zinc, diluted (one part to twelve of water), or a weak solution of carbolate of zinc should be used. Mr. Hinton recommends drying-powders, such as magnesia, chalk, etc. Solutions of the acetate of lead are very objectionable, as



there is danger of deposits of lead being formed, even when the solution is rendered acid; although the danger is thought to be lessened by this measure. The application of solid nitrate of silver to the edges of the perforation is attended with risk of increasing the size rather than leading to its diminution. The local application should simply stimulate the edges of the perforation to a sufficient degree to increase the activity of the circulation, and favor the closure by granulation. This is done with great facility by melting the salt in the platinum ladle (Fig. 97), and applying it by Livingstone's elastic caustic probe (Fig. 98).

Should suppuration or a free discharge from the middle ear exist, this must be gotten rid of before the opening can be closed. The proportion of cases in which closure takes place is one in five; such, at least, accords with the experience of the author.

Two cases of perforation, successfully treated, are added, by way of illustration:

CASE I.—Mrs. Anna A., American lady, aged thirty-five, married. Whilst attending the opera had a seat near an open door. In the course of the following night she was attacked with a pain in the ear, which was described as radiating into the head, causing intense suffering, in spite of the free leeching, purgation, and anodynes, both local and general, which had been used by her physician. She was not relieved until a discharge took place from her ear, which continued for some time, when she became alarmed at her deafness,

Fig. 97.




Fig. 98.



and consulted the author. Upon examination, I discovered a small linear perforation, situated about the centre of the membrana tympani. The walls of the meatus were red, and covered with a discharge, which I removed. The surface of the membrana tympani was injected with blood. The edge



of the perforation was touched with a solution of nitrate of silver, gr. xxx to fʒi of water, every third day. A solution of sulphate of zinc, gr. i to fʒi of water with fʒss glycerine, was dropped into the ear by a competent nurse, and removed by careful syringing. A blister was applied behind the ear, and Politzer's bag employed to keep the middle ear free, by inducing a discharge from the pharyngeal orifice. Gradually the diffuse redness of the membrana tympani disappeared, and, under the influence of the blister and the wash, the aperture became closed in a month, without much diminution of the hearing power on that side.

I saw the patient one year afterward; she has had but one return of the catarrhal affection, which was promptly relieved by an astringent wash.

CASE II. was much more tedious, requiring six months to heal the perforation, but this was owing to the debility of the general system caused by the disease. M. E. T., German, aged forty-nine; had contracted a remittent fever whilst visiting the Sulphur Springs, Va. His fever was very high, brain affected, etc. He convalesced, and was able to ride about, when his ear began to discharge. This continued for some time before he sent for me. When seen, the meatus was red and swollen, and no examination could be made for several days. When the swelling subsided, I was able to see, by the aid of a speculum and Von Tröltsch's reflector, a perforation at the bottom of the membrane, through which, with some difficulty, air could be forced by expiration. The treatment was the same as the preceding case. Fomentations and leeching, however, were required to reduce the swelling of the meatus. It was six months before the aperture healed up; and although he hears ordinary conversation, yet when the watch is applied to that ear, it can be heard only at one-half the distance that it can be heard by the sound ear. A visit to Germany during the summer of 1868 has improved his general health and, with it, his hearing. I directed him during the extreme cold weather to obtain from W. Oakford a pair of his ear-protectors, and wear them in windy cold weather. They are made of velvet, lined with oil-silk, and by means of an elastic cord, going over the head,



kept in place. They allow the air to circulate in the ear, without confining any discharge.

In 1859, the author published an account of "The Use of the Artificial Membrana Tympani, by the late Mr. Toynbee," and reported some cases of his own; but at that time the use of this little apparatus was not well understood, because the designer of it was himself in error in regard to its proper application and theory of action. Mr. Yearsley certainly deserves most of the credit for the practical application of his piece of wet cotton, and also for his theory of its mode of action, which he published to the world in 1848; and subsequently, on June 6th, 1857,\* he states: "The more experience I have of this important invention the more I am satisfied that my theory of its *modus operandi* is correct, and Mr. Toynbee is decidedly wrong, and that so surely as he succeeds in shutting up the opening in the membrane, as surely does he fail in improving the hearing, whatever be the material employed, whether it be the original wetted cotton, or the substitution of vulcanized rubber." Again, in Dec. 1857, he stated the true theory of its action, which was subsequently accepted by Mr. Toynbee (being led, he tells us, to do so by witnessing a demonstration by Mr. Julius Erhard), that hearing might be improved by the pressure of cotton-wool on an entire membrane, and coming now to the conclusion that it was by restoring contact occasioned by some disconnection, or loss in the ossicular chain. Thus, as the Supplement points out,† he approached the opinion long held by Mr. Yearsley, that the cotton-wool supported the ossicula. As the matter now stands, Mr. Yearsley is the inventor of this application, and thus states the case in his own words: He maintains that to be successful the orifice in the *membrana tympani* must neither be filled in, nor completely covered by, the cotton, and that an opening must be left along the walls of the *meatus*, down to the site of the *membrana tympani*.

The following is a condensed statement of answers to questions addressed to his patients: I apply it only to the end

---

\* London Med. Journ. and Gaz., June, 1857.

† The Diseases of the Ear, by the late J. Toynbee, F.R.S., with a Supplement by James Hinton, M.R.C.S., Aural Surgeon to Guy's Hospital. London, 1868.



of the passage, and on one side. (1) Were I to cover the passage entirely, I should be more deaf than I am. (2) If I do not introduce the cotton down to the (opening), and place it on one side, I find it more than useless. (3) I find it to have the desired effect by placing it on one side only, so as to preserve an opening down to the tympanum. (4) I still apply the cotton remedy, as at first applied by you, at the side and the end of the passage. (5) I do not cover the external part of the drum, but I place it sideways toward the back, leaving a small opening.

One of the chief objects of the cotton is, according to Mr. Yearsley, to rupture the air bubble (indicated to the patient by a distinct *click*); and secondly, to prevent the re-formation of bubbles of air in the discharge in and around the perforated membrane. Simple as it would appear, it is an operation requiring the most delicate tact to manipulate the nice adjustment of the cotton. A very convenient way of applying it is to pass a linen or silk thread through the centre of a bunch of cotton-wool, and tie it so as to hold the wet cotton by the thread. The free ends of the thread can then be passed through a straw quill, or silver tube, two inches long, and by this means the cotton or wool can be carried down through the meatus to the edge of the perforation, but not to cover it entirely. When it is accurately adjusted, the tube is withdrawn, and the cotton left in position. The cotton can also be adjusted by means of a pair of forceps, in little cylinders on the handle of the malleus in case of nervous deafness without perforation, as recommended by Dr. Erhard,\* who claims to have been the first to have found the method of curing deafness by pressure on the membrana tympani on his own ear, and to have published it in 1849.

This is not a new idea, for in both modern and ancient literature there are cases related where deaf patients have accidentally found that by introducing a foreign body into the meatus the hearing was temporarily improved.

Mènière† relates an interesting case: An old judge had

\* *Rationelle Otiatrik*, pp. 274, 275. Erlangen, 1859.

† *Traité des Maladies d'Oreille*, par Kramer, traduit par Mènière. Paris, 1848, p. 526.



been accustomed, for at least sixteen years, by pressure of a blunt gold needle against the membrana tympani, to improve his hearing for an hour or two. Mènière examined the ear during this state of things, and found the membrana tympani uninjured, and that the pressure was upon the handle of the malleus. He states that he had seen several similar cases, and considered them cases of nervous deafness.

It will be remembered that in long-continued discharges from the ear of a purulent character, there is often a separation of the little bones. As most of these cases commence in the middle ear, the loosening of the capsule of the joint of the long process of the incus takes place, separating it from the stapes; but if the membrana tympani, with the incus, be pressed against the stapes, the connection will be restored. Should there be no handle of the malleus or incus left, there is not much prospect of any improvement by pressure; still, if the stapes is alone, with a portion of the membrane left, a partial success will follow. In two cases, out of many in which the author employed the artificial membrana tympani of Mr. Toynbee, the improvement in hearing must have followed from the contraction of the artificial tympani, as in every instance it was his endeavor to fit the membrane carefully and accurately over the opening. It is almost impossible, from the shrinking of the material, to preserve a shut sack. At Fig. 99 may be seen the form of the artificial membrane of Toynbee.

CASE I. *Chronic purulent catarrh, or chronic otitis interna, with perforation of the membrana tympani; application of an artificial membrane with success.*—James Riddle, aged twenty-four, a native of Ireland, by occupation a silk weaver, applied January 21st, 1858, at "Western Clinical Infirmary,"\* on account of a troublesome discharge and deafness in his right ear, the result of scarlet fever, of which he had a severe attack at the age of thirteen. He had become so deaf in the right ear as not to hear the tick of a watch when placed in immediate contact.

On examination with Wilde's speculum and a good light,

---

\* Former name of Howard Hospital.



the meatus was found filled with mucus, pus, etc.; upon removal of which, by careful syringing with warm water, a large opening was found in the membrana tympani, near its centre, and the membrane lining the meatus thickened and contracted. Astringent and stimulating injections were now employed to change and alter the secretions, and as there were some enlarged cervical glands, he was ordered five grains of iodide of potassium three times daily, in a bitter infusion, with counter-irritation behind the ear by cantharidal collodion. In the other ear there was thickening of the membrana tympani, and, from this cause, deafness.

January 26th. Discharge has moderated, but hearing not improved. Found, on examination, that air did not pass through the Eustachian tube. Introduced the Eustachian catheter at two different sittings.

Jan. 29th. Same treatment; improving.

February 9th. Discharge still less. Introduced the moistened cotton; but, after remaining in for a few days, it increased the discharge. Although removed and cleansed each night, he complained that the itching was so intolerable that he desired its removal, which was done, and injections resumed.

Feb. 12th. Introduced an artificial membrana tympani, made by Mr. Kolbe, of this city, which gave him no pain, and his hearing was slightly improved. (See Fig. 99.)



Feb. 15th. The discharge is again on the increase; has blackened the silver wire, and caused the artificial membrane to wrinkle and change its color, so that it was removed, and the ear allowed to rest.

Feb. 18th. Again introduced the membrane. It is more comfortable to-day; can hear best in the open air, and reintroduced the membrane himself, finding out the right spot; for, if he pushes it too far, it is of no use to him. He has been testing his powers of hearing by a clock.

Feb. 24th. Returned to-day, and states that he cannot hear so well; when, upon examination, found the Eustachian tube blocked up with mucus accumulations, which were again



removed with a series of injections of warm air; and on filling up the ear with a weak solution of sulphate of copper, and by making an effort to swallow, with the nose and mouth closed, air passed up in bubbles through the liquid.

This effort to swallow with nose and mouth closed he was directed to make, when his artificial membrane was out, so as to keep the tube pervious.

*Conclusion.*—This patient continued to visit the Infirmary twice a week until 17th of March, when he left for the country, having but slight discharge, and being able to hear conversation with ease and comfort, and even the ticking of an ordinary clock across a room.

CASE II. *Scrofulous inflammation of the membrana tympani; perforation; and successful use of artificial membrane.*—Wm. C. N., sailor, aged twenty, native of Philadelphia, came under my care in the beginning of October, 1860, sent to me by Professor J. A. Meigs. I found his hearing very defective in right ear; the membrana tympani had almost all disappeared, save a small ledge. In this ear I advised him to use an artificial membrana tympani, of Mr. Toynbee's, which I obtained of Mr. Weiss, of London. With the left ear he was unable to hear his watch, unless in contact with his temporal bone. I introduced the Eustachian catheter with great difficulty and with much care; the air passed, apparently for the first time, into the middle ear; it was heard by the use of the otoscope by Drs. Burpee, Tryon, and Buchanan. I continued to use it, but he complained of much pain over the brow and eye; while holding his mouth and nostrils closed, passes air into the cavity himself, and finds his hearing improved by it, being able to hear his watch in contact with his ear.

Treatment continued for one month; no more improvement. Artificial tympani having given way, a second one (made by Mr. Kolbe under my direction) was introduced, which he states he cannot hear without.

The internal treatment consisted of Fowler's solution of the arsenite of potassa, with cod-liver oil.

The following is his statement after being under treatment about two months:



“DR. L. TURNBULL.

“DEAR SIR,—I have no remembrance of ever hearing with my left ear. If the ear was perfect in the performance of its function, it must have been while I was very young. My right ear was perfect until October, 1859. Whilst I could hear with it, it was very acute; so much so that I was marked as being quick at hearing.

“In March, 1858, I contracted a severe cold, which by neglect settled on my left lung. About five weeks after I contracted the cold, swellings began to appear on my neck, first on the right side then on the left. These proved to be scrofulous swellings. In June, 1859, one of them opened, and remained open till last April. In October last, whilst my neck was in the above condition, I experienced a sudden loss of hearing, and my right ear commenced discharging matter, which it has kept up ever since,—although the discharge at present is very small. I was advised to syringe it daily with lukewarm water and Castile soap or milk. About last March I commenced to use cotton and glycerine; I continued to use these until June; I then procured some wool, and every two or three days would put a small piece dipped in glycerine close to, or rather on, the drum-head. This improved my hearing very much; I continued to use it for seven or eight weeks, and latterly the ear became painful; I went to Dr. J. A. Meigs, who examined my ears with a speculum. He said the right ear was very much inflamed; that there was a perforation in the membrane; that the upper part hung down like a curtain over it. The left ear had the membrane very much thickened; also the lining membrane of the tube. He told me to introduce with a quill a few drops of a solution of nitrate of silver (two grains to one ounce) in my right ear daily, so as to allay the inflammation; also to use with a brush citrine ointment in my left ear, for the purpose of thinning the membrana tympani. I did so for a week, and went to him again. My left ear became very painful under the action of the citrine ointment. In washing it out every morning, filmy matter would come from it. Upon examining the right ear again, there was a great decrease of inflammation, but the opening was much larger.



The left ear had become much inflamed. Dr. Meigs then told me to stop using the ointment in the left ear and use the caustic solution until the inflammation was brought down, when I was to begin again to use the ointment. For the right ear I was to increase the strength of the solution. I asked Dr. Meigs the cause of the defects. He stated that a small ulcer had formed on the membrana tympani, and perforated it. Upon looking at it first, he thought he could close the opening by touching it with caustic; but when he examined it again, the upper part of the membrane had resumed its place, and the opening below was much larger. At first it was about as large as o, the second time O. The defect in the left ear was owing to a thickening of the membrana tympani caused by a severe cold.

"At present I use nothing in my right ear, except the artificial membrane which you gave me. It improves my hearing considerably. My left ear has improved somewhat under your treatment. I can now hear the ticking of a watch, which I could not do before.

"Yours respectfully,

"W. M. C. N.

"PHILADELPHIA, Oct. 1860."

"*Chronic purulent inflammation of the tympanum* is often an accompaniment of perforation of the membrana tympani, and the following are Lucae's\* methods of treating chronic purulent inflammation of the tympanum. In the majority of these cases he has found that they were accompanied by granulations, not only in the middle ear, but along the Eustachian tube and in the pharynx, and this condition of tube and pharynx keeps up the inflammation in the ear. He directs treatment more especially to the Eustachian tube, and has seen the best results from the insufflation of dry powders through the catheter into the tube. For this purpose he recommends muriate of ammonia, sulphate of zinc, alum, acetate of lead, and sulphate of copper. The addition of a small amount of camphor to any of these he thinks is of

---

\* Monatschrift f. Ohrenheilk., No. 4, 1870; Berliner Klinisch. Wochenschrift, No. 6, 1870; Transactions of the Am. Otol. Society, 1870.



advantage.. The powdered sulphate of copper is the most valuable in granular pharyngitis. The advantages of the powder over strong solutions are, that we get the greatest action in the tube where it is needed, without the risk of inflammation of the tympanum from the unexpected entrance of considerable quantities of fluid. Where only a very mild action is required, Lucae prefers the injection of liquids by Gruber's method. For the granulations in the tympanum, sulphate of copper in substance or solution (from two to five grains to an ounce) is highly recommended.

"One cause of the frequent failure to cure in these cases is the insufficient or imperfect application of the astringent instillations. To remedy this, Lucae uses what he calls a prolonged ear-bath, by means of which he keeps the solutions applied to the desired parts for any length of time, without interfering with the occupation of the patient. A medium-sized glass tube is bent at a right angle, with a long arm about one inch in length, and a short arm about one-half inch in length. The long arm is covered with rubber tubing, and inserted tightly into the meatus; into the shorter arm, which stands upright, the solution is instilled drop by drop, till the tube is nearly full. The patient is now able to move about, and the solution can be thus retained as long as desired. Previous to these applications the ear is to be thoroughly cleansed by the double current (catheter, see Fig. 94) already described. At first, the solution is retained for half an hour twice a day, but the length of time can be increased to several hours if desired."

"Nasiloff\* has described, as a new form of *inflammation of the membrana tympani (myringitis villosa)*, a shaggy condition of the dermoid layer of that membrane, which seems to have been an inflammation and hypertrophy of these papillæ. Kessel also accepts this name, and describes† a condition of the membrana tympani similar to that given by Nasiloff."

---

\* Centralblatt f. d. Med. Wissensch., 1867, No. 11.

† Archiv für Ohrenheilkunde, vol. v. p. 250, and Transactions of the Am. Otol. Society, 1870.



## CHAPTER XVI.

DISEASED THROAT, UVULA, TONSILS, AND CARIOUS TEETH, IN THEIR RELATIONS TO DISEASE OF THE EAR; ALSO FEIGNED DEAFNESS.

HAVING in a previous chapter treated of the anatomy and pathology of the posterior portion of the naso-pharyngeal space and of the necessity of a rhinoscopic examination, with the catheterization of the Eustachian tube, we shall in this chapter direct our attention to the anterior portion of the nasal passage and mucous membrane lining the pharynx, arches of the soft palate, uvula, and tonsils. The importance of this subject cannot be overestimated, for it is a well-recognized fact, demonstrated by a large practice, that in a considerable number of cases the catarrhal deafness is either directly or indirectly the result of acute or chronic inflammation of this portion of the mucous membrane which lines these parts, and is extended to the orifice of the Eustachian tube, middle ear, and mastoid cells. In such cases of deafness we must endeavor, both by topical and constitutional means, to restore the throat to a healthy condition; for on the slightest increase of disease in this part in certain individuals their deafness is augmented. An enlarged pendulous uvula filled with serum, causing cough, must be reduced by cutting off a portion, followed by the application of astringents and stimulants, to restore it to its normal condition.

The throat in the acute catarrhal inflammation is red, tumid with injection of the superficial blood-vessels, with fever and pain, and more or less secretion, which becomes thickened; in the chronic form there is either a great accumulation of mucus or extreme dryness. The acute condition is to be treated by depletion, direct or from the skin and intestines, mucilaginous and anodyne drinks, astringent gargles, etc.; the chronic form, by those agents that will



restore tone and vigor to the capillaries and build up the broken-down constitution.

*Enlargement of the tonsils.*—This depends upon a semi-solid organizable material, effused into the tissue connecting the cellules or lacunæ of which the gland is composed. In the adult it becomes organized and much more firm; in children there is in most instances a reabsorption; but in some cases it remains permanent.

The symptoms of enlarged tonsils are more or less impediment to the breathing, compelling the individual to sleep with the mouth open; snoring in sleep; a thickness or peculiar twang of the voice; some difficulty in swallowing, and often *deafness*. The tonsils may either project toward the middle of the throat and almost touch, or they may grow upward and downward, so that in looking into the open mouth but a small portion of the enlargement is seen unless the finger be passed into the fauces or a hook inserted and the gland drawn up.

If the enlargement is in an upward direction, it may obstruct respiration through the nose and also the passage of air through the Eustachian tubes.

Dyspnœa and difficulty of swallowing depend upon congestion or inflammation of the tonsils, to which persons having such enlargement are very subject, and which inflammatory attacks extend to the cervical glands and mucous membrane of the fauces. Deafness in almost all cases is due, according to the best authorities, to chronic thickening of the mucous membrane of the fauces near the enlarged tonsils. Tröltsch\* says this is particularly shown by examinations on the dead body. In recent subjects we often find the mucous membrane of the middle ear, at the same time with that of the pharynx, in a state of congestive swelling, hyperæmia, and hyper-secretion. Johannes Müller says the mucous membranes have a great tendency to communicate their affections along their course.† “We see,” observes Tröltsch, “that affections of the mucous membrane are often extended

---

\* Tröltsch on the Ear, translated by Prof. Roosa. New York, 1869.

† Handbuch der Physiologie, 1844.



'per continuitatem.' Catarrhal inflammation of the conjunctiva and lachrymal sac occurs from coryza, and the inflammation of the buccal cavity, in typhus fever, extends through Wharton's duct to the little glandular canal of the parotid."

It is well known that constitutional diseases—typhus fever, tuberculosis, and the acute exanthemata—very often extend themselves from the pharynx to the mucous membrane of the ear. If the tonsils, therefore, are abnormally large, he (Tröltsch) advises their removal; for if they are not themselves the seat of frequent inflammation and abscesses, they retain the chronic congestion of the pharynx by their presence, since they act as foreign bodies and prevent the normal action of the muscles of the throat. They also push the broad portion of the soft palate upward, and thus, but not in a direct manner, as is generally believed, press the anterior lip of the Eustachian tube against the posterior.

In advising the removal of the tonsils, we should be governed by their condition. If they are not much diseased, or if their enlargement cause no inconvenience to the patient, they should not be removed. If, however, there is impediment to breathing, a nasal voice, frequent attacks of sore throat, and, above all, deafness, then their removal should be effected, no matter what is the age of the sufferer. The operation is simple, and not often dangerous. It is stated that Dr. James Yearsley, of London, has operated in more than *three thousand cases*, without a single accident or unfavorable result. M. Guersant\* says he has operated on more than *one thousand children*, and can hardly reckon more than three cases out of this number in which formidable bleeding occurred. On the other hand, he has operated on about twelve or fifteen adults, and must confess that four or five at least of them claimed his attention, and it was not possible, without the use of the actual cautery or the sesquichloride of iron, to arrest the bleeding.

After relating two instances in which the hemorrhage was

---

\* The Surgical Treatment of Diseases of Infancy and Childhood. By. T. Holmes, M.A., Cantab., p. 555.




not only copious but absolutely alarming, Prof. Gross says: \* "Since the publication of the first edition of this work, I have met with two other cases of severe bleeding after this operation,—one in a youth of eighteen and the other in a man of thirty. The hemorrhage was quite copious, but finally yielded to the application of the tincture of the sesquichloride of iron, aided by a full anodyne."

Erichsen refers to a case in which the hemorrhage was effectually arrested, after the failure of all other means, by a gargle of spirits of turpentine suspended in mucilage. We have had cases in which there was profuse hemorrhage, but by the use of ice, tannic acid, or dry persulphate of iron the bleeding was checked.

There are three methods of removing the tonsils, viz., with the forceps and knife, with the tonsillotome or guillotine, and with some powerful caustic paste. The first mode is applicable to all cases, the second to children, and the third is to be used where the patient has a dread of the knife.

The forceps should be long and hooked, and the knife should be a probe-pointed bistoury, which should be guarded to within about an inch and a half of its point. The enlarged tonsil is seized with the forceps and drawn as far as possible into the mouth. It is then divided by passing the knife from below upward, only cutting off the upper portion drawn into the mouth, being careful not to wound the tongue or soft palate.

Fig. 100.



The principal tonsil instruments are Mathieu's, Physick's, and Fahnstock's. This latter instrument, which has been modified by Charrière, of Paris, is the one we employ. It

\* Gross's System of Surgery, vol. ii. p. 375.



consists (as in Fig. 100) of two blades, sliding on one another, and terminating in a ring. The posterior edge of the upper ring is sharp. On the upper blade a groove is chiseled, along which travels an arm, which carries a double-pronged fork; and this arm is so constructed that, as it passes over the ring, the fork is raised a certain distance away from ring.

The frame is provided with a handle on either side, for the first two fingers of the surgeon's hand, and one behind for his thumb. The frame is so connected with the fork and blades that, by pressing the thumb forward, the operator first projects the fork, and then the upper ring, as in Fig. 100. The instrument being put into the mouth, usually depresses the tongue sufficiently. It is then turned, with the lower ring toward the tonsil and pressed upon it, so as to draw the tonsil into the ring; then, by pressure of the thumb, the prongs of the fork are struck into the enlarged tonsil, and as the fork passes forward, it draws the gland farther into the ring; then, by continuing the pressure of the thumb, the upper ring slides forward and cuts off the part within the ring, as in lower part of figure. As observed by T. Holmes,\* the whole affair is over in a moment, and the main recommendation of the guillotine is that it only requires one hand, so that any opposition not too resolute can be overcome, the mouth kept open, or the tongue depressed with the other; while, in operating with both hands, the surgeon often finds himself foiled just at the moment of commencing his incision, by the child shutting its mouth in terror, or in beginning to struggle. Besides, when once the fork has been struck into the tonsil, the operation is over; while, in operating with the knife, the child sometimes tears himself loose after the section has been begun, with a scrap of tonsil hanging down in his throat. It is well to have several sizes of rings of the instrument, as often one will not fit into the mouth of the child or youth. In all operations for excision of the tonsils in which chloroform (or ether, or a mixture of the two) is employed, and in many cases where the child is merely a little unsteady and frightened, the gag, devised by

---

\* Surgical Treatment of the Diseases of Infancy and Childhood.



Mr. T. Smith, for operations on the soft palate, is recommended.

It is made of stout wire, bent to the shape and soldered on to a tongue-plate. The horizontal portion fits inside the lower teeth, and the plate keeps the tongue out of the operator's way. The upper branches fit inside the upper teeth, lying close under the superior maxilla, and, when open, maintain the jaws apart. They are opened and closed by a screw. The whole is kept in position by a strap passing around the child's head; the branches connecting the upper and lower portions lie in the corners of the mouth and keep it widely distended. A somewhat similar instrument was devised by Dr. Mussy, of Cincinnati, and by Mr. Hohlmann, of the firm of Tiemann & Co., of New York, a figure of which may be seen at p. 448, Transactions Am. Med. Association, vol. xx. 1869.

With respect to hemorrhage, Dr. Holmes writes: "It has not been my lot to witness any formidable hemorrhage in either of these operations." He also regards the after-treatment quite unimportant. Prof. Gross considers it prudent to confine the patient to a moderately warm apartment; at all events, he should avoid cold air, and, if necessary, on account of the resulting inflammation, he should take an active cathartic. For want of attention to this, he states that "several lives have been lost that might otherwise have been saved."

The third plan for removing enlarged tonsils is that proposed by W. J. Smith\* and Dr. Morell Mackenzie,† of London. Smith's method consists in the repeated application of the potassa fusa to their surface by means of a small metal dish, mounted on a handle. The dish being heated over the flame of a spirit-lamp, is filled with potassa fusa by rubbing a heated stick of that substance into it, and is then pressed on the surface of the tonsil. Thus a portion is destroyed; and when the slough has separated and the symptoms produced by the caustic have subsided, it is to be re-

---

\* On the Treatment of Enlarged Tonsils at any Period of Life without the Operation of Excision. London, 1865.

† Medical Mirror, 1864.



applied. The interval will probably be about a week, and the number of applications required, even in cases of no inordinate extent, seems, according to Holmes, to be not less than five or six. He also states that the application causes considerable pain in the parts, in the ears and throat, with swelling of the glands, etc.; but no alarming symptoms have ever occurred, beyond a temporary spasm of the glottis, twice repeated in a child, in whom, however, the treatment was successfully carried out. He also claims for it a superior efficacy in cases in which the enlarged tonsils are flat, deeply buried behind the arches of the palate, and growing upward toward the opening of the Eustachian tube.

Dr. Mackenzie gave the name of "London paste" to his agent, which consists of a mixture of equal parts of *caustic soda and lime*. Forty cases are cited (1864) in which it had been used by him with complete success. We have tested this paste to a limited extent, so far, with success. It is moistened with absolute alcohol on a piece of glass or porcelain, and applied by means of a glass rod, being careful that the paste be not too thin. The pain is not generally severe; the paste acts at once, adhering to the tonsils and causing a deepening of the color, and in a day or two an eschar is formed.

In one instance there was pain following the application, which continued for two days, and the child, a girl ten years of age, would not again submit to its use; yet so large a quantity had been employed that the tonsils were reduced nearly one-half. If by any means too much is employed, the parts should be washed, or the throat gargled with vinegar and water, followed by olive oil, which removes the unpleasant taste. If the tonsils are very much indurated, a good plan is to remove a small portion by the knife, and then apply the escharotic paste. In several instances we have succeeded by this plan in removing altered or nasal voice, irritable throat, and deafness, especially in young lads.

#### DEAFNESS FROM CARIOUS TEETH.

Deafness is often dependent upon the want of teeth, and sometimes an *artificial set* will restore the hearing.

In other instances pain and even deafness will be caused



by decayed teeth. Mr. Cattlin\* describes a case in which the loss of hearing was evidently reflex paralysis, caused by a carious tooth:

"In the month of September, 1843, a lady in the second stage of consumption consulted me concerning a diseased right lower molar. She had for about three months suffered acute pains in the tooth, ear, and side of the neck. When I saw her she had become deaf for four days. The inflamed tooth was extracted, and her hearing returned within an hour after the operation."

In another instance a diseased molar tooth caused ulceration of the auditory canal. I shall quote Mr. Hilton's† account of it:

"Some years ago Dr. Addison had a very offensive discharge from the auditory canal of one of his ears, which annoyed him very much, and below the external ear was a small gland, enlarged. He had tried various remedies for this discharge, and had gone, I believe, to some surgeons who attended especially to the ear; but, as far as I could learn, no good resulted from any of their applications. Upon examining the ear from which the offensive discharge proceeded, I found a slight ulceration upon the floor of the auditory canal. On arguing the question out between us, we came to the conclusion that the ulceration probably depended upon a diseased molar tooth in the lower jaw on the same side. We had that tooth extracted, and in a very short time the ulcer healed; the discharge and morbid secretion disappeared from the auditory canal, and as soon as that ulceration was cured the enlarged gland subsided."

#### OTALGIA (EARACHE, OR NEURALGIC PAIN IN THE EAR).

This will often occur in children without any apparent inflammation or warning. Tincture of opium applied in the ear with an equal quantity of olive oil will generally relieve it. In the adult suffering from intense pain in the ear we

---

\* Transactions Odontological Society, vol. iii. p. 308, London, 1863; Guy's Hospital Reports, Third Series, vol. xiii. p. 94, London, 1868.

† On the Influence of Mechanical and Physiological Rest, etc., by John Hilton, F.R.S. London, 1863, pp. 194-199.



have found useful equal parts of glycerine and chloroform, or the oil of belladonna, or hyoscyamus, alone or combined with morphia. M. Emile Duval\* says "that he has in person found relief in severe earache, after other means had been tried in vain, from the use of a mixture of equal parts of chloroform and laudanum; a little being introduced on a piece of cotton. The first effect produced is a sensation of cold; then there is numbness, followed by scarcely perceptible pain and refreshing sleep."

In cases of very severe neuralgic pain radiating from a tooth to the ear, the following pill will often be found useful:

Sulphate of Quinia, gr. i ;

Extract of Belladonna, gr.  $\frac{1}{2}$  ;

Sulphate of Morphia, gr.  $\frac{1}{2}$  .

Make into one pill.

S. Give one every three or four hours.

If this do not relieve the pain, apply a lotion of acetate of lead, chloroformi and pulv. camphoræ, partis equalis. Should the pain still continue, administer hypodermically one-third of a grain of the solution of the acetate or sulphate of morphia.

#### FEIGNED DEAFNESS.

As deafness may occur without any change in the appearance of the external ear, it is a favorite disease to simulate. The modes of detecting it are obvious. A watch should be set on the suspected person day and night, and things calculated to excite interest or apprehension should be uttered in his presence and the effect carefully noted. It has been recommended to place the fingers on the pulse while any bad news or threat of punishment is being uttered. The patient should be called sharply or unexpectedly by name, or in a whisper, or he should be roused from sleep and spoken to, or a piece of money should be let fall close to him. All these and similar tests have failed, and sometimes even the loudest noises have not produced any apparent effect. In one case related by Dunlop, a pistol was fired off close to the ear without effect; but upon putting the man to sleep with opium the imposition was detected on the repetition of the firing.

---

\* Gaz. Méd. de Lyon, i., July, 1861.



(Anæsthesia produced by ether or chloroform has been advantageously used.)\*

*Dumbness.*—This is sometimes assumed with great perseverance. As a general rule, it may be stated that if a man not deaf can move his tongue, he is not dumb. Some add to feigned dumbness a pretended mutilation of the tongue, which they effect by rolling it back into the throat, and scratching it to make it bleed. A relation of Dr. Guy's detected this fraud in the case of a man who, by a written paper, describing his captivity among the Algerines and his mutilation by them, had excited great commiseration. Pretending to give entire credence to the man's tale, and taking a half-crown out of his pocket, he requested the man to show him his tongue once more, upon which he thrust his finger suddenly against the root and the tongue rolled out. My own observations lead me to believe, however, that this test is not always to be relied upon.

*Deaf-dumbness.*—The combination of deafness and dumbness, though never occurring in a person previously in possession of the faculties of hearing and speech, is sometimes obstinately pretended. In one or two remarkable instances, men who have pretended that they were born deaf and dumb have resisted every test, and have been detected only by those having personal experience with the really deaf and dumb.†

Dr. Casper (*Handbuch der Gerichtlichen Medicin*, tome i.) gives the two following instances of feigned deafness: A female, convicted of theft and imprisoned, complained that the dampness of her cell had rendered her deaf. Even when addressed in the loudest tones she appeared not to hear. Suspecting her veracity, the doctor pretended to believe her, and to prescribe for the disorder. On one occasion, while visiting her, he exclaimed aloud, "Good heavens, there is vermin here!" And suddenly turning towards the prisoner, said, in a very low voice, "Madam, you have a louse on your right sleeve." The woman instantly turned, and examined her right arm, with a motion of evident disgust.

---

\* Loss of Speech and Hearing successfully treated by Inhalation of Ether, by J. H. Hutchinson, M.D., *Amer. Jour. Med. Sciences*, vol. xlvii. p. 412.

† Guy's *Forensic Medicine*. London, 1869, p. 153.



The next instance is that of an old woman who was tried for an assault upon a female with a sickle. The prisoner pretended to be sick, feeble, and completely deaf; and no amount of voice on the part of the court sufficed to make her understand anything. Casper was thereupon called in, who, from the bearing of the woman, at once suspected her of feigning. Approaching her ear, he shouted, "You are accused of having grievously wounded Mrs. Lemke!" "It's a lie," replied the prisoner. "But," rejoined the doctor, "Mrs. Lemke would not say so were it not true, for"—suddenly dropping his voice into a whisper—"you know Mrs. Lemke is not a liar." "She is a liar!" exclaimed the prisoner aloud, pushed by the spirit of revenge to gratify her hatred even at the expense of exposing her own deception.

A plan adopted by some of our army surgeons, during the late war, to detect malingerers feigning deafness, was to drop a large book or other body suddenly behind the suspected person. If deaf, his attention would be attracted by the jar, and he would turn around to ascertain the cause of the concussion, but if a malingerer, he would not move a muscle. In the early history of the New York Deaf and Dumb Institution, it is stated that a man presented himself as a deaf mute, and as having been educated in France. Some doubt was entertained as to the truth of his statements, and it was decided to test his pretensions to deafness by dropping on the floor a metal basin containing heavy things to make a noise. The experiment was made, and he stood the test without flinching, as was supposed. He was treated kindly and dismissed with presents. Shortly afterwards he was heard from, as being in prison for some offense against the law, and it was found that he could speak and hear as well as anybody. In fact, a true deaf mute would have been startled by the concussion. In general, people pay but little attention to jars. A few years since, in England, an opposite inference was drawn from a similar experiment, and an impostor detected.\*

---

\* Annual Report of Pennsylvania Institution for the Deaf and Dumb, 1868, p. 19.



FEIGNED MUTISM DETECTED BY A STRONG AND PROLONGED  
SHOCK OF ELECTRICITY.\*

"On the 16th day of July, 1859, a boy about fourteen years of age was taken up in the street for begging, and committed to the House of Refuge by one of the aldermen of the city. At the time of his arrest he intimated to the officer that he was unable to speak. He could, however, evidently hear perfectly well. He wrote his name in the alderman's office as John Thompson, born in the State of New York, and gave his age as thirteen years. At the Refuge he also gave them to understand that he could not speak, and that he had lost the ability to articulate sounds in his early infancy. He was found to be able to read with facility anything that was written for him, and he wrote a quite distinct and well-formed hand himself. My colleague, Dr. Haskell, who was on duty at the time of his admission, examined him carefully, and desired that he should be closely watched, so as to detect if possible whether his mutism was feigned or not. From that date, however, up to the close of February, a period of nearly eight months, during which he was constantly at work or at play with the other boys, no one ever heard him utter a single distinct word save once, shortly after his admission, when he was locked in his room for some misdemeanor. He then called to a boy passing the door to know if the bell (for dinner) had rung. When accused immediately afterwards with having uttered these words, he admitted that he had, but indicated his inability to repeat the same or any other words, even under the threats of punishment. His whole conduct, subsequently, was so very consistent with his statements that those of the managers and officers with whom I conversed in reference to his case, expressed themselves convinced of the boy being either unable, or under a delusion as to his ability to talk. Indeed, one of the managers, who took a special interest in the boy, wrote me in February a letter in regard to him, requesting

---

\* Dr. A. Hewson, Transactions College of Physicians. Philadelphia, June, 1861.



me to examine him carefully as to the possibility of any surgical operation being performed to restore his voice; and if such was not possible, that I should recommend his being sent to the Deaf and Dumb Asylum. On my first interview with the boy I found him remarkably bright, with an expression of face quite characteristic of deaf mutes. He had a restlessness of eye and remarkable quickness of manner. He communicated with me very freely with the single-hand alphabet, which he stated he had learned whilst in the institution from one of the teachers. To my inquiry as to where he had learned to read and write so well, he gave me to understand that his mother had shown him the letters and taught him their meaning; that she had been very devoted and assiduous in her efforts to teach him. The teacher from whom he stated he had learned the mute alphabet, assured me afterwards that he had never shown him the letters but twice. His organs of speech, on careful examination, were found by me to be in a perfect state. His palate, tongue, fauces, and larynx were, as far as I could ascertain, perfectly natural, and to my frequent solicitations for him to utter a single word or letter, I got only a negative shake of the head. The boy evidently would not attempt to articulate a sound, and the fact that he had spoken once in the institution satisfied me that there was merely an unwillingness, and not an inability, to speak. I also made frequent attempts to get him to utter some sounds. At first he would not, then he gave utterance to a squeaking guttural sound. Then, by dint of perseverance, I got him to pronounce after me some labial sounds, such as *b*, some of the hissing sounds, such as *c* and *s*; but I never could get him to attempt the distinct lingual and guttural sounds by which I hoped he would betray himself. My conviction of his feigning was not in any way, however, diminished by these interviews, although my researches in works on feigned diseases and medical jurisprudence were not rewarded by the discovery of a single case approaching to it; but, on the contrary, in the appendix of Mr. Wilde's work on the Ear (see p. 438, American edition), I there saw some cases of real mutism so analogous to this one that I might well have had some misgivings as to the



correctness of my opinion, especially with the strong testimony of the experienced officers of the institution in opposition to me. The fact, however, that the boy had been heard to speak, and admitted that he had, was an insurmountable one, and two expedients presented themselves to me by which I thought I might be able to determine the truth. These were a strong shock of electricity, and the intoxication of anæsthetic agents. I determined to try the electricity first, and accordingly on Saturday, the 20th of March, with the assistance of my friend, Dr. Charles Lee, I administered a strong shock of the induced current from a large magneto-electrical apparatus, by means of moistened conductors applied over the larynx. This application, I assured him, would bring his voice back, and I told him I would continue it until he made some effort to speak—until he uttered some articulate sound. He held out for some time, at least twenty minutes, although the application was evidently very painful. Finally, however, he cried in quite a suppressed tone, 'Enough!' I then ceased; and then feeling satisfied that the boy had been severely punished, I did not resume the application, although he would not utter another word. On the following Monday I intended to repeat the battery, and, if it was not more successful, to put him under the influence of ether. When he saw me, however, on that day approaching in the hall, with the battery and bottle of ether in hand, and in company with my colleague, Dr. Haskell, he came forward and said, in a slow, measured cadence, and as though he articulated with difficulty, 'How do you do, sir?' and I was informed that he had been gradually recovering his voice since I had used the battery on Saturday. He reiterated, in answer to my inquiries, the facts in reference to his history. Some of those circumstances I was confident were not correct, and I told him so, but he assured me he was telling the truth. He continued to improve, and in the course of a few days it was discovered that he could read poetry in a very creditable manner; and when questioned as to where he had learned to read with such good measure, he quickly answered that he had noticed the boys reading poetry at the public school



near Brooklyn. He now became quite a lion, and when I took him before the managers to induce them to resort to some coercion to get at the truth in his case, I found the most experienced of them under the conviction that I had either brought the voice to a child where it had never been developed, or that the child had been suffering under hallucination that he could not speak. These impressions prevailed for six weeks, when one day the boy, of his own free will, gave the superintendent the following statement in his own handwriting:

“‘PHILADELPHIA HOUSE OF REFUGE,  
“‘May 4th, 1860.

“‘Having feigned to be deaf and dumb for about nine months, and made several erroneous statements while in this institution in regard to my history, I now truthfully declare that my object in doing so was to avoid the possibility of detection as a runaway from my parents, who reside in Alleghany City, opposite Pittsburg, on the Alleghany River, in Pennsylvania. I left Alleghany City on the 1st of July, 1859. I took with me a box partly full of lozenges, so as to make my parents believe I would sell them. When I got to the river I commenced to call the boys' attention by daring feats; at last I got a thick plank, and went out into the middle of the Alleghany River. I floated down the river on it, and the boys on the shore thought I was drowned. I suppose I must now come back to Mr. S. S. East,—he was a newsdealer in Alleghany City. He always trusted me, and I now owe him two dollars and a half, which I would like to pay very much. He resides near the depot. I then traveled partly through Ohio, and from Ohio to New York City. Stayed there two or three days, and then came to Philadelphia on Monday, July 16th, 1859. I am not induced to make the above statement through fear of punishment, but that I may be relieved of a great uneasiness which I feel on account practicing such a deception.

“‘CHARLES L. C., alias JOHN THOMPSON.’”



## CHAPTER XVII.

### NERVOUS DEAFNESS.

I HAVE endeavored to bring together the important facts in regard to one of the most obscure forms of diseases of the human ear, namely, "nervous deafness."

*Definition.*—By "nervous deafness" is generally understood "a diseased condition or impaired function of labyrinth and auditory nerve;" but, in my judgment, it should also include affections of the nerves which supply the external ear, membrana tympani, and middle ear in their reflex influence. The term *nervous*, in a truly scientific sense, is not correct, but it is the best term we have at present for indicating the probable origin of this form of deafness.

*Etiology.*—This form of disease is often the result of hereditary taint, being communicated from parent to child.

In one family, consisting of two sisters and one brother, every member was deaf, in another two, and in a third there was one deaf in the family, for three generations.

Sedgwick\* has observed that a surgical instrument-maker, aged twenty-nine years, became deaf, like his father, at the age of eighteen years; and Dr. Camp informed him of a case occurring in his practice of nervous deafness affecting a father and son.

One of the most common occasional causes is cold, more especially applied to the head uncovered, or by sudden force to the ears; heat from the sun, when the head is exposed, or with but a slight covering, if the system is exhausted, producing *coup de soleil* and deafness, from congestion of the brain. Then we have the reflex influence of

---

\* Sedgwick on the Influence of Sex in Hereditary Disease, B. and F. Medico-Chirurg. Review.



convulsions in adults or children, or from apoplexy or paralysis, producing pressure, congestion, inflammation, or atrophy of the seventh pair of nerves or its branches. Then follow direct injuries from falls or blows upon the ear, of a severe character, producing concussion of the nerves. Another prolific cause is the poison of typhus, typhoid, and scarlet fevers; rheumatism, gout, measles, or mumps, intermittent fever, hysteria, epilepsy, and syphilis.

We have also this form of deafness symptomatic or associated with disorders of the digestive organs. It is likewise the result of old age.

Toynbee classifies the nervous disorders of the ear according to their cause:

"The first class (to which belong those cases where the special nervous apparatus of the organ is alone affected) may be subdivided into diseases arising from—(1) Concussion. (2) The application of cold. (3) Various poisons: as that of typhus, scarlet, or rheumatic fevers, of measles and mumps, of gout, of an accumulation of bile in the blood, and of quinine in large doses.

"And the second (where the brain, conjointly with the ear, seems to be injured), into diseases arising from—(1) Excess of mental excitement. (2) Physical debility."

With regard to the last, Mr. Hinton remarks:

"It is difficult to me to accept *debility*, nervous or otherwise, as in itself a cause of nervous deafness. I do not find that the cases of deafness, which appear to me nervous in their character, occur at all specially in the weak. Nor does my observation show me that among specially weak people defect of hearing is particularly common.

"I have, however, been much struck by the frequency with which the symptoms of a passive nerve lesion—a condition suggesting the idea of a functional paresis—coincides with demonstrable tympanic disorder, past or present.

"There are, however, exceptions to this rule, and not very unfrequently we meet with a gradually increasing inability to hear in persons, especially of the female sex, in whom no traces of physical disorder or mental strain can be discovered.



"It is remarkable how small a mental shock suffices in certain persons to give, as it were, the finishing blow to the power of hearing. One patient, whose hearing was previously weak, found herself almost totally deaf after a quarrel between two relations who both appealed to her; another, on visiting a sister dangerously ill, was told on entering the house that she was dead. She heard the information perfectly, but on reaching an up-stairs room was deaf. In both these cases the affection has continued more or less; but both of them also, when seen by me some years afterwards, showed slight signs of former tympanic disease."\*

*Case of nervous deafness from mental shock.*—The wife of Mr. S., a missionary of the Bible Society of Philadelphia, has been deaf for many years. She is of exceedingly nervous temperament. She says that while a young lady she was suddenly called home from a visit among her relatives on account of the illness of her father. Upon being taken to her room she asked how he was, and hearing no reply, thought he was very ill; and calling the doctor into an adjoining room, asked how her father was. Receiving, as she thought, no reply, she feared the worst, and implored him not to keep her in suspense, but tell her how her father really was; when the doctor laid his hand on her head, and said, "Why, my child, you must be deaf!" And so she was; and has remained hard of hearing ever since, perhaps some twenty years or more.

The following is her own account, sent in September, 1863, with the examination:

"I was what is called a delicate child, but not known to have any nervous affection. I was obliged to earn my own support before I was twelve years old; I therefore led a very sedentary life. I think it was in my sixteenth summer I received some distressing news, which instantly caused almost total deafness. I used remedies to brace my nervous system, and in a few weeks could hear, if spoken to in a louder tone than for ordinary conversation. The following summer I spent several weeks in the country, and a short

---

\* Mr. Hinton's Supplement to Toynbee, p. 459.



time at the seashore. My bodily health improved very much, and I gained ten pounds in flesh. My hearing returned; but in a few weeks after my return home it again became bad, and, though I have tried various remedies since, I have never received any permanent benefit. I will be forty-four years of age next December. Ever since I have been hard of hearing I have had noises of some kind in my head, either of a buzzing, ringing, or roaring sound, and can hear best when there is a great noise, as when an organ is playing very loud; I can then hear, though spoken to in a very low tone; and I have often noticed, when riding in an omnibus over the stones, that I could hear conversation better than others whose hearing was good. My not hearing persons when they first speak to me has always had an unfavorable effect, as the very effort to hear seems to make me more deaf. I have been married sixteen years.

“AMANDA M. S.”

*Examination.*—External meatus natural; membrana tympani good color, slightly milky tint; middle ear, both Eustachian tubes open, but a good deal of mucus accumulation. No treatment was of any permanent benefit.

*Frequency of the disease.*—Kramer, in his early work, states that out of 2000 patients 1074 were classed as affections of the auditory nerve; now, with the advance in pathological anatomy, especially in the study of exudations, he has reduced the number to 4 in 1000.\* In Wilde's table of 2385 cases, only 244 were true cases of “nervous deafness.” In the still later table of Triquet, of 163, 46 were truly nervous. Toynbee publishes 200 cases of deafness, and only about 35 or 36 of nervous are given; still, he considers that the “nervous apparatus, which receives the sonorous undulations from the tympanum and conveys them to the brain,—one of the most delicate structures in the human body,—is liable to many functional and organic derangements.”

Of some fourteen hundred cases of ear disease, which I

---

\* See The Aural Surgery of the Present Day, by Dr. W. Kramer. Translated for the New Sydenham Society.



have treated in the Howard Hospital and in private practice, I find that the proportion of cases in which there was no lesion of the external or middle ear is about twenty in one thousand.

*Influence of sex.*—In Kramer's table we find 581 are males and 447 females. In Wilde's the sexes are nearly equal; while Triquet had 18 females and 28 males.

These statistics do not sustain the generally received opinion that females are more affected with nervous deafness than males.

*Physiology.*—The auditory nerve or portio mollis of the seventh pair has its origin from the medulla oblongata by two roots. The labyrinth receives nerves from no other source than the portio mollis, unless we suppose the portio intermedia to consist of filaments from the facial, which accompany the ramifications of that nerve into that part of the ear.

Mr. Bowman considers that the portio mollis is the nerve of hearing, and that it is abundantly proved by the following arguments:—1. The distribution of the nerve to the internal ear, to which no other nerve of any importance is distributed. 2. Its softness of texture and cribriform character, which distinguish it from ordinary nerves of sensation or motion. 3. Diseased states of it, or of parts immediately near its origin, affect the sense of hearing, whilst a paralytic state of the portio dura or of the fifth does not affect the sense. I do not consider this argument of Mr. Bowman's entirely conclusive, as the accessory nervous apparatus plays a most important and necessary part in this function. We find its analogy in the eye. If the supraorbital branch is cut, vision will often be lost, while the optic is not injured; so also, injury, pressure, inflammation of the branches of the tympanic nerve of Jacobson, from the glosso-pharyngeal, and from the otic ganglion, produce loss of hearing of "nervous character," and only affecting the auditory nerve secondarily, or by reflex action.

The tympanum receives branches from the facial and glosso-pharyngeal, and probably from the sympathetic.

The facial, in its passage through the aqueduct of Fallopius, gives off the chorda tympani, which, according to Mr. Bow-



man, seems to have no physiological connection with the tympanum or its contents. Yet it has a connection, if we admit, with Arnold, that there is an anastomosis between the acoustic nerve and the nerve of Wrisberg, into which the chorda tympani continues. The chorda tympani passes along the membrana tympani, and if a moderately strong induction current is sent into the ear, the patient will sometimes experience a peculiar sound and a painful sensation of sticking in the ear, and a drawing-together sensation in the anterior half of the tongue. Duchenne, Erdmann, Baierlacher, and other observers have proved that this sensation in the tongue is produced by the galvanic irritation of the chorda tympani, which, passing on the inner side of the membrana tympani, joins, immediately after its exit by the Glaserian fissure, the lingual branch of the fifth. These two nerves do not lie together in one, but are connected by filaments throughout their whole course.

The stapedius muscle also receives a branch from the facial nerve. The anastomosis of Jacobson results from the subdivision of the tympanic branch of the glosso-pharyngeal nerve, which enters the cavity of the tympanum below, and, passing over the promontory, gives off branches to the membranes of the fenestræ and Eustachian tube, and to the otic ganglion.

A branch is described by Arnold as proceeding from the otic ganglion to the tensor tympani muscle.

The external ear is supplied by the facial nerve as regards its muscular apparatus, and by the fifth pair as regards its sentient surfaces.

*Pathology.*—There exist at present very many indications by which the morbid conditions which have been found on dissection of the nervous structures of the ear may, with some degree of certainty, be connected with symptoms observed during life. Mr. Toynbee found the auditory nerve diseased fifteen times; ten times atrophied, twice with blood effused around it, once suppurating, and once having a tumor attached to it, and once absent. Disease of the labyrinth he observed seventy-three times; most frequently atrophied, thickening, deficient endolymph, excess or deficiency of



otoconie, deposits of pigment membranous bands, exostosis, the endolymph opaque or red, or the blood-vessels enlarged.\*

"Voltolini found most frequently thickening of the membranous structures, cretaceous deposits, deficiency and excess of otoconie, deposit of pigment, amyloid degeneration of the auditory nerve, once a fibroid tumor in the cupola of the cochlea, and once a sarcoma of the nerve.†

"In Mr. Hinton's dissections he has found more or less decided morbid conditions of the nervous structures by no means very uncommon. Most frequent has been an enlargement and congestion of the vessels of the labyrinth, often of a very extreme degree. This he has observed forty-one times, and chiefly in connection with inflammatory affections of the tympanum. Excluding cases of deaf-dumbness, he has found the membranous labyrinth obviously atrophied sixteen times, soft and swollen ten times, and seeming fatty twice; the endolymph opaque or red seventeen times, seeming too abundant three times, and too scanty twice; the otoconies in apparent excess fives times, and in apparent deficiency eight times; twice the labyrinth has contained pus, and once cholesterine; once the sacculus had undergone bony degeneration; once the lamina spiralis was thickened; once the cochlea contained a fibroid mass, three times an excess of pigment, and twice extravasated blood. The auditory nerve he has seldom found decidedly diseased. Twice the internal meatus was narrowed by bony deposit, and three times the fibres seemed atrophied or softened.

"Of all the appearances observed, the frequency of vascular congestion, and the extreme degree it reaches, without the existence of any symptoms which have commanded attention, seem to him the most important."‡

The late Joseph Toynbee, F.R.S., has made the greatest number of sections of the ear, and states that the occurrence is very rare of caries of the labyrinth, for this reason: that ulceration of the mucous membrane of the tympanum is far

---

\* Medico-Chirurgical Transactions, vol. xxxviii.; Toynbee's Catalogue, p. 119.

† Von Tröltzsch, *Ohrenheilkunde*, third edition, p. 385.

‡ Hinton on Nervous Deafness, *Guy's Hosp. Rep.*, 1868.



from a common disease, while, under the influence of chronic inflammation and of the secretion collected in the tympanum, the membranes both of the fenestra rotunda and ovalis become thickened and turgid. Mr. Toynbee has met with one case in which there was ulceration of the membranous labyrinth; he reports four other cases. In two of them the labyrinth became carious or necrosed, and the dead bone discharged, without the production of any symptoms of cerebral disturbance. The first occurred under the care of Mr. Hinton, in a man aged fifty-five, who suffered from a discharge from the right ear for some years, and at last the cochlea was discharged entire. The second case occurred to Mr. Shaw, in a boy aged seven years, who had suffered from otorrhœa in both ears, following a severe attack of scarlet fever, which he had two years and a half before. He had paralysis of the muscles of the left side of the face; for a year he had been completely deaf in both ears. The greater part of the petrosal portion of the temporal bone, including the meatus auditorius internus and labyrinth, was removed by the forceps, being separated by necrosis. He entered the Middlesex Hospital, July 31st, 1855, and was discharged by the end of September. The ulcer in the tube cicatrized; there was paralysis of the face; deafness, and a trifling discharge from both ears. He left the hospital in good health. In the other cases death resulted, the brain becoming seriously involved.

The following most interesting case, which also recovered, is quoted from Mr. Wilde. He says: "I am indebted to Sir Philip Crampton for an examination of one of the most extraordinary pathological dissections of diseased bone perhaps in existence, consisting of the entire internal ear, cochlea, vestibulum, and semicircular canals, with a small portion of the inner wall of the tympanum, which he drew forth from the meatus of a young lady, who, after the most urgent symptoms of inflammation of the brain, with paralysis of the face and leg, and total deafness of one side, recovered from the head symptoms and the paralysis of the extremities, after a copious discharge from the ear."\*

---

\* The Nature and Treatment of Diseases of the Ear, p. 358.



Let us, with Von Tröltsch, examine the analogous relations in other organs; for instance, that of vision, and, at the same time, study the sources of nutrition, and the mode of development of the inner ear.

“The inner ear is entirely independent of the other portions of the organ of hearing, both as regards its source of nutrition and its development. Its artery, the internal auditory, does not come from the external carotid, as do the vessels which supply the external and middle ear, but from the brain. It comes either directly from the basilar or from the anterior cerebral branch of the latter. There does not appear to be any direct communication between the vessels of the middle and external ear, so that secondary disturbances of nutrition of the labyrinth can only result from alterations in the brain, or from morbid conditions in the interior of cranium, but not from affections of the outer ear, except when both are involved. The process of development also shows the independent character which pertains to the labyrinth from the very beginning. Whilst the tympanic cavity with the Eustachian tube are developed from the bronchial apparatus, or, as Arnold and Von Baer assert, from a process of the pharyngeal mucous membrane, the labyrinth is developed from the ear vesicle of Emmert, or projecting process of the cerebral vesicle. The labyrinth is formed much sooner than the petrous portion of the temporal bone, and its ossification proceeds independently. Even afterwards, the nutrition of the part of the petrous portion of the temporal bone which contains the labyrinth is seen to be independent of that of other parts; and I have mentioned instances of isolated necrosis of this portion. Let us now examine the analogous relations of the eye, and we will find that there, as is well known, the diseases of the retina and optic nerve are far less frequent than affections of the outer coats and refracting media, and yet here the causes for a production of disturbances in the nutrition of the nervous apparatus exist in a far greater degree than they do in the ear. The retina and the entrance of the optic nerve lie in an elastic globe, which is exposed not only to internal but to external influences; the retina is



not only connected with the brain, but with the choroid and the vitreous body, whilst the nervous apparatus of the ear is only connected, as regards nutrition, with the brain, and being enveloped in a bed of the hardest bone, and considerably removed from the surface of the body, it is unaffected by any external influence except the severest wounds and concussions.

“It is, therefore, highly probable that primary affections of the labyrinth, or idiopathic ‘nervous deafness,’ are incomparably less frequent than affections of the outer and middle ear, and that, when secondary, they are most frequently caused by affections of the brain, or by changes in those parts which also belong to the tympanum, and which, as we have seen, give rise to very numerous pathological phenomena, viz., the round and oval windows, which have both a covering of the periosteum of the labyrinth. Abscesses or caries may find their way from without, as we have already said. In no case, however, should the diagnosis ‘nervous deafness’ be made, unless no change, to which we can attribute the symptoms, can be discovered in those parts which are explorable by our senses. This demands a perfect knowledge of the normal condition of the parts, and a faculty to discern the slightest change,—*e.g.* in the membrana tympani. Many even experienced aurists are deficient in these requisites, and very considerable abnormalities in the just-mentioned organ have been overlooked. With regard, then, to the frequency of the diagnosis ‘nervous deafness,’ we remark, that in all the branches of medicine, the less one’s anatomical knowledge and exact research, the oftener one considers diseases as ‘nervous;’ but the greater one’s progress in these two requisites, the more means one has of explaining the symptoms, the less often does one make such a diagnosis, which, as a general rule, is nothing better than a makeshift. The severity of the deafness does not necessarily indicate an affection of the labyrinth; affections of the middle ear may cause the most severe deafness, as observation teaches us.”\*

---

\* Von Tröeltsch, Anatomie des Ohres, pp. 93, 94, 95.



The nervous apparatus of the ear suffers, as is well known, in many ways from affections occurring during infancy, especially from the morbid conditions associated with convulsions. Considering the severity of the cerebral disturbance often resulting from diseases even of the tympanum alone, especially in children, it seems to be deserving of consideration, whether, in many of the cases in which cerebral disease is supposed to result in deafness, it would not rather be found that disease of the internal ear had induced the symptoms of cerebral disease. Apart from Flourens' experiments—confirmed and extended by the more recent ones of Czermak on doves—on the effects produced by injury to the vestibule and canals, there is evidence of the serious results to the nervous centres which acute disease of these parts involves.

As bearing upon this subject, Mr. Hinton avails himself of Dr. Wilks' kindness to report two cases which were under his care:

"The child of a medical man, aged two, was taken ill on March 9th, with feverish symptoms, etc., and a powder with calomel was given it. On the three following days continued very ill, constantly crying out, as if in pain, and with much febrile disturbance.

"On March 14th Dr. Wilks saw him; found him very ill, very restless, constantly moving about, as if in great pain; skin hot, lips parched, pulse 140. There were no symptoms indicating any cerebral mischief; but the bowels were relaxed, the abdomen full and soft. It was thought, therefore, that his complaint might be dysentery in an early stage, and two grains of Dover's powder were ordered.

"March 19th. Appeared better; had slept, and was quieter. After this, however, the former symptoms reappeared; extreme restlessness, and screaming, as if in pain. There was evidently some source for this, but it was difficult to discover. For a week he thus continued; throwing his head about, as if in pain, but with no other cerebral symptoms; and the child appeared quite sensible. He then passed a lumbricus, and it was hoped that another purge might relieve him, but it had no such effect.



"March 22d. Appeared worse; never quiet for a moment; lying in his nurse's lap, throwing himself about, his arms, legs, and head, as if endeavoring to gain repose from some internal agony. He then began to make curious movements with his mouth, constantly thrusting his tongue out, and licking his lips. For nearly another fortnight the child thus continued; with constant restlessness, throwing his head about, as if in great pain, and placing his hand over his head and face. The father then lanced the gums, but with no effect. About a month after the commencement of his illness, and when the child was wasted to a mere shadow, a discharge occurred, first from one ear and then from the other. For a day or two there appeared no relief; but after this the child began to get better; ceased to cry, took its food, and grew stout. He is at the present time in good health, and his hearing said to be good.

"The case was a difficult one, and no decided opinion could be given during the child's illness; but after the occurrence of the discharge from the ear, it was thought most probable that an acute otitis had been the disease.

"Jas. B., aged twelve, admitted into Guy's Hospital, under Dr. Wilks, January 12th, 1861, with severe cerebral symptoms. He was apprentice to a bootmaker at Peckham. On inquiry being made in reference to injuries, it was said whilst in the workhouse he had often been struck by the master, and he also lately had falls whilst sliding, but it was not known that he had hurt his head. A fortnight before admission he was seized with rigors, vomiting, etc., with febrile symptoms. These continued until two days before he came in, when he became deaf. He was then excessively ill, and in a drowsy state, rolling his head about as if in pain. Was perfectly deaf, the loudest noise not perceived; he was, however, apparently quite sensible. Tongue thickly coated; bowels costive; occasional vomiting. Leeches had been applied to the temples; was ordered calomel, one-half grain every four hours. He continued much the same until the 15th, when purging came on, and the mercury was stopped. After this he slightly improved, and on the 19th he was much better. The febrile symptoms were disappearing, but he



was as deaf as ever. He rapidly improved, and on the 31st he had left his bed. On examining the ears, the membrana tympani appeared quite perfect. The boy soon after left. He called upon me about eight months afterwards; he was in good health, but he appeared entirely deaf."

The following case shows somewhat of the subsequent history of a case probably very similar to the last:

"Laura E., aged thirty, healthy, but very nervous. Lost her hearing entirely at about a year old, at which time severe cerebral symptoms set in; the eyes being turned up, and the body stiff for several days; no distinct convulsions; both jugulars are said to have been opened, and mercury was rubbed in. Diarrhœa followed, which seemed to relieve the head, but the hearing was gone. An eminent physician said that if it were recovered it would be at the time of cutting the second molars, and at that time, about the age of seven, there came on a slight perception of sound. She was then again treated with mercury, which is supposed to have induced a nervous weakness, without result on the hearing. She appears now totally deaf in the right ear; the left very gradually advanced till about the age of twenty, and with it and the aid of the eyes she can understand short sentences, but cannot repeat numbers shouted into her ear; distinguishes sounds by the quality of the vibration,—*e.g.* she perceives thunder in the head, guns in the feet. She felt the explosion at Hounslow, some years back, at a distance of fifteen miles, when no one else in the neighborhood was aware of it. Some tinnitus; does not hear better in a noise. It is said that she heard (comparatively) well on the mountains in Switzerland. Throat healthy.

"Meatus narrow; each membrana tympani bright; inflates left with a slight moist sound, not right. The right Eustachian tube being closed, it was in two or three sittings gently made pervious. No change ensued."

"This is an example of very many cases, and, in respect to some at least of them, is it not natural to doubt whether an affection of the brain, or its membranes, of so severe a character, could have passed away entirely, leaving no effects except an almost total abolition of the function of the audi-



tory nerve? What should induce so exact a limitation of an inflammation affecting parts in absolute continuity? Would it not be more reasonable to suppose an inflammation, possibly beginning in the tympanum, and, by extension inward, involving the vestibule and cochlea?—a result which might well have ensued in the first case mentioned by Dr. Wilks, if the membrana tympani on each side had not given way. It is not uncommon to find inflammatory affections of the tympanum inducing abscess in the cerebral substance without disease of the intervening parts,—the bone and dura mater being in no visible way affected. Is it at all improbable that a parallel extension of morbid action might take place in respect to the nearer and more closely related labyrinth? I am the more persuaded that the causation of the symptoms, in some of these cases, is that here suggested, because I have, in the dissection of the ears of deaf mutes, met with appearances in the vestibule that seemed indicative of former inflammatory action, especially a roughness and contraction of the first turn of the cochlea at its junction with the vestibule.

“It is not only in this extreme form, however, that deafness results from apparent ‘cerebral’ affections in childhood.”

“Among the multitude of children who apply for assistance, in consequence of disease of the tympanum, or Eustachian tube, there occurs every now and then a case which, on a hasty examination, would pass as an ordinary one, but would be sure in time to arrest the surgeon’s attention by its refusal to take the ordinary course. Remedies do not tell. Perhaps the Eustachian tubes are obstructed, but opening them has little effect; if the membranes are perforated, the condition of the exposed mucous lining of the tympanum may be improved, but the hearing does not advance. Then, perhaps, it begins to be noticed that the child has a peculiar, inexpressive look, unlike both the keen inquisitiveness of the congenital deaf mute and the self-absorbed restlessness of the imbecile; and, on testing more minutely, it is found that the tuning-fork is very badly heard, and that, so far as can be elicited from children, the other symptoms of a nerve affection are present. In these



cases it is often found, on inquiry, that there have been, probably some years before, more or less defined convulsive or 'cerebral' attacks. For the most part these attacks seem to occur during the first two or three years of life. Sometimes, however, they are deferred to a somewhat later period."\*

In view of numerous clinical observations, there is no doubt that nervous deafness results from "cerebro-spinal meningitis."

*Symptoms.*—In forty-six cases of nervous deafness reported by Triquet,† the symptoms may be summed up :

1. *Anatomical symptoms.*—The pavilions and external meatus were found healthy forty-five times. In one case there was a slight serous exudation, indicating some catarrhal complication. In twelve cases the membrana tympani had lost their usual transparency. They were cloudy; and in one case the membrana tympani had a parchment-like aspect. In twenty-two cases cerumen was found in the meatus in more or less noticeable quantity, of good consistency, and the color and appearance of honey. The Eustachian tubes were found healthy and free in forty-six cases.

2. *Physiological symptoms.*—The initial deafness has always been considerable. In twenty-seven cases the patients could not hear the watch, when in contact, at the commencement of the treatment; and in the remaining nineteen cases eleven patients heard the watch only when in contact; and in eight others the watch was heard at a distance of from four to ten centimetres, on an average.

Of over forty-six patients, nineteen had deep-toned intra-aural sounds; four had shrill-toned sounds or whistlings; one patient had strange sounds without any indication; and twelve had no intra-aural sounds.

Erethismus of the ear is noted eleven times; twice it was very marked, and nine times it was completely wanting. There is no mention made of it in the other thirty-five cases. Headache has been noted twenty-four times. Paracousie, cophosis or double hearing, is noted only twice. The senile circle existed in all the cases, in a degree more or less marked.

---

\* Hinton, Guy's Hosp. Rep., Third Series, vol. xiii.

† Traité Pratique des Maladies de l'Oreille, p. 449.



## DISSONANCE OF THE ORGANS OF CORTI.\*

"The dissonance of the organs of Corti may have reference to a single or to a whole succession of tones; it may be transient or permanent.

"It has hitherto been observed in diseases of tympanic origin, in severe concussion of the labyrinth; and it may result from a blow upon the ear, or intensity of sound, and also by influences acting upon the whole nervous system. From these general observations we will, in proceeding, mention the special cases or facts in historical order.

## DOUBLE HEARING (PARACUSIS DUPLICATA).

"Under the head of derangements of the organs of Corti, we find, as longest known, that of double hearing, which affects the hearing of both ears as well as that of one only.†

Sauvages‡ relates a case of double hearing, which developed itself in consequence of a catarrh, and with it disappeared. Itard treated a woman who was troubled with intermittent deafness; whenever the hearing returned again, the patient heard all sounds double.

"Von Gumpert observed in himself a paracusis duplicata monauricularis. Here, the symptom appeared to present itself, after inflammation of the ear, the origin of which could not be anatomically located. The variation in tone fluctuated between a *third*, a *fourth*, and an *octave*. Von G. could distinguish both tones; the deeper tone appearing close to him, the higher at a distance of two or three yards from the diseased ear. Von Gumpert also observed in himself a double hearing of words. These phenomena, which, during the whole continuation of the disorder, were at no time accompanied by ringing, whizzing, or rushing in the ear, continued

---

\* Klinik der Ohrenkrankheiten, von Dr. S. Moos. Wien, 1866, pp. 319, 320.

† This is termed by Prof. Knapp "diplacusis binauricularis," in analogy with "diplopia binocularis." See Hemholtz's Theory, p. 40.

‡ Bressler, Die Krankheiten des Kopfes und der Sinnesorgane. Berlin, 1840, Bd. ii. s. 375.



for eight days without interruption, and seemingly disappeared after dropping in *oleum hyoscyamus coctum*.\*

"For an accurate personal examination of these symptoms, made four weeks after a severe purulent inflammation of the middle ears, we are indebted to Von Wittich.†

"The tone of a tuning-fork struck the disordered ear at exactly a half-tone higher than the sound ear, and so with all the moderately high tones; when the third octave was whistled with the mouth, or struck upon the piano, it was heard doubled, with a difference of half a tone. These experiments continued the same, whether the outer auditory passages of the diseased ears were filled with wadding, or water, or otherwise rendered tense through pressure of air on the drum of the tympanum. A tuning-fork placed on the teeth, produced first the natural tone, and then by degrees the next half-tone was heard ringing; if placed upon the forehead, the tone became a half-tone higher, increasing as it neared the disordered ear.

"With two tuning-forks, one of which gave a half-tone higher than the other, only one tone was heard with both ears when the higher was sounded near the healthy ear, and the lower near the diseased ear.

"If the organs of Corti, through their really peculiar arrangement of fibres, bring about the sensation of a tone of fixed number of vibrations and continuance, it is accountable how another tone becomes apparent through exudation in the cavity of the tympanum, and through a thereby altered pressure of the fluid of the labyrinth on the nerve filament. So that for the tone *a*, the filament for *b* sounds, whilst when *b* was struck, the filament *a* was vibrated.

"These explanations of Wittich's may be adopted for all those disorders of the organs of Corti which belong to the diseases of the middle ear.

"So, also, in the following case: A celebrated tenor consulted me, because, fourteen days after a violent cold, he was troubled with a symptom which completely interrupted him

---

\* Made by boiling the fresh leaves of the *hyoscyamus* in olive oil.

† A case of double hearing observed in himself. Königsberg Medical Chronicle, vol. iii. pp. 40-45.



in the exercise of his calling. The patient said that he heard the third of every note he sang at the same time. Examination showed as the cause a simple catarrh of the middle ear, with considerable deafness of both sides. After nine applications of the catheter, together with injections of ammoniæ murias, — ten grains to half a fluidounce, — the patient was completely cured."

In other cases the theory of Wittich's was not so applicable; for example:

"V. S., an officer, sixty years old, very musically inclined, for four years visited the sea-shore, on account of emphysema of the lung. Since that time he suffered under a gradually increasing deafness. This had developed itself entirely without pain, beginning with a rushing sound on both sides, which had lately ceased. Of late years the patient had suffered much from catarrhal complaints of throat and chest, and especially a tolerably severe asthma.

"The examination showed a considerable amount of epithelial scales in both external ears, extending pretty well into the tympanic membrane, and spread from the centre to the circumference, as also a strongly marked circular, muddy margin. Injection of the vessels of the handle of the malleus; dullness of the light spot. The tubes of both sides are slightly pervious; catarrh of pharynx. Understands conversation two and a half paces to the right, one foot to the left. The watch at six feet is not heard at all (also when applied directly to the bones). After three weeks' treatment, the patient was able to hear low muffled sounds for more than thirty-six feet from him,—end of August, 1863, dismissed.

"On the 14th of May, 1864, the patient appeared again, and reported as follows: 'About a month after, at an evening musical entertainment at my house, I had a pretty violent attack of asthma. I went into another room, in order to relieve the attack as quickly as possible by means of the inhalation of chloroform. Directly after the narcosis I was much deafer; it came on with various strange noises, upon both sides, and I heard all tones above the first *a* struck doubled; at present, however, it is the second *e* struck



doubled, and all octaves from that on.' Found the patient much worse as to hearing. Could now distinguish only very loud conversation at the length of one room and a half to the right, and the length of one room to the left. A watch, distinguishable at twenty-five to thirty feet, was heard two inches to the right, and only in contact on the left side. His perception of sounds now gradually became worse, and on the 21st of April, 1865, he could hear only loud conversation at one to two paces from him, on both sides, and the watch, on contact, on both sides.

"All musical tones were heard false and confused; and music, for which the patient had before entertained such liking, is now an abomination to him.

"The dissonance of the organs of Corti may not be considered in this case as a secondary disorder, brought about or dependent upon a chronic process of inflammation in the tympanum.

"We must suppose rather that as the former had already existed ten years, in consequence of the narcotic effects of the chloroform, sudden changes in the labyrinth arose, which caused the double hearing; the nature of which changes it is difficult to determine."

*Diagnosis of nervous deafness.*—To arrive at a correct diagnosis, we must first examine with care the auricle, external meatus, membrana tympani, handle of the malleus, so as to discover any change in color or form from that found in a perfectly normal ear. Then follow in order the throat, uvula, and tonsils, and the Eustachian tube, passing air through it with the aid of the catheter and explorer, with the otoscope to determine the internal condition of the tube. Rhinoscopy should also be practiced to examine the condition of the orifice of the Eustachian tube, and in some cases may be available to determine whether the Eustachian catheter has actually been introduced. Then we take into consideration the diseases of the labyrinth, namely, acute and chronic inflammation, with inflammation of the facial nerve within the Fallopian canal and obscure affections of the brain. In this manner we reduce the actual number of apparent affections of the inner ear to which deafness may be attributed.



“Erhard in Germany, and Bonnafont in France, give the preference over this objective diagnosis of nervous deafness to a functional subjective. Thus, according to Erhard, ‘the common physiologico-pathological symptom of all nervous deafness is a lessened conduction (of sound) by the bones of the head, varying in degree, so that a cylinder watch is not heard, and a case watch only seldom, and then only feebly.’ The uncertainty is further increased by the declaration, that the ‘sound-conducting power of the bones of the head is lessened with age, and also by a thick diploë, a thick scalp and soft parts;’ that finally, ‘those persons only can be determined to be affected with nervous deafness whose conducting power of the bones of the head is deficient, and who are under forty years of age.’

“Bonnafont is still better satisfied with his means of diagnosis, for he believes, and takes credit to himself for having made a new and great diagnostic discovery, ‘that the sensibility of the auditory nerve must be perfectly normal if the watch is heard on all parts of the cranium;’ and ‘when the watch is heard only at the mastoid and zygomatic process, the hardness of hearing is still curable; if the watch is heard nowhere on the skull, but the tuning-fork (at the third octave) is heard five centimetres from the ear, or at least at any point of the skull to which it may be applied, he considers the curability of the hardness of hearing as very doubtful; finally, if the tuning-fork is heard only a short distance from the ear, or when applied to the skull, the incurability of the functional disturbance of the ear cannot be doubted.’ Here again arises the difficulty that the patients themselves must determine whether they hear or do not hear the watch or tuning-fork.”\*

Mr. Hinton, surgeon aurist of Guy’s Hospital,† has published a selection of twenty-three cases, from which we have availed ourself of some affections classed as “nervous deafness,” and gives the following signs by which any given case of deafness may be inferred to depend upon a disease of the nervous system :

---

\* Von Trötsch, *op. cit.*

† Guy’s Hospital Reports, Third Series, vol. xiii. 1868, p. 152.



“1. *The history.*—This is in many cases of itself decisive, especially when it connects the defect of hearing with other nervous derangements. The history, however, may be very misleading if too much relied upon; some of the most marked forms of nervous deafness, especially those of sudden origin, being liable to be closely simulated by simple mechanical obstructions. Among the most frequent causes of nerve-deafness, besides blows on the head and the effect of extremely loud or long-continued noises, are fever, sunstroke, mumps, diphtheria, parturition, prolonged residence in India. Convulsions in childhood stand as the origin of many cases, but it is a question whether, in a large proportion of these, the convulsive seizures have not been an effect of the ear disease.

“2. Certain peculiarities of the hearing; such as the hearing better in a noise (as when riding), or hearing worse on excitement, depression or fatigue, or on any special attempt to listen. A much better hearing for some sounds than for others, as for example, for the watch as compared with the voice, is a suspicious circumstance; so also is a great diminution. It might have been supposed (inasmuch as we hear our own voice partly through the cranial bones, and its resonance is decidedly increased by closure of the meatus) that some value might be attached to the patient's own manner of speaking, whether loud or low; since this seems to depend mainly on the degree in which his voice is heard. But though as a rule, at least in the more recent and less extreme cases, this supposition seems to hold good, there are too many exceptions to allow it any practical value. Many patients with recent tympanic deafness complain that their own voice sounds loud to them; and the loud and harsh sounds uttered by deaf mutes is an extreme instance of the tendency to raise the voice in those whose nervous faculty of hearing begins to fail.

“3. A certain value is attached to the degree of deafness, which if it be very excessive cannot depend on absence of conduction; but it is difficult to draw the line.

“It seems certain that disease of the conducting apparatus may reduce the hearing to a lower pitch than its congenital absence necessarily involves.



"4. Certain tests can be applied to the hearing by means of a tuning-fork placed on the head, which will sometimes give almost or quite decisive information. Little value can be attached to the mere fact that a tuning-fork of any particular pitch is heard badly, or even not at all, through the cranial bones, since not only do various diseases of the tympanum produce this effect, but I have known at least one person, with healthy ears and perfect hearing for the voice, entirely unable to hear a tuning-fork upon the head. But since this peculiarity is rare, an inability to hear vibrations thus imparted to the cranial bones may be fairly held as a suspicious circumstance, and at least the converse may be held valid that, if the tuning-fork be well heard, the deafness is not likely to be nervous. But the chief use of the tuning-fork in diagnosis rests on the simple fact that its vibrations, when imparted to the bones, *pass outward* freely through the tympanum and external meatus, and are therefore reinforced by reflection when either the tympanum or the meatus is rendered impervious to sound. If the meatus of a healthy ear be closed by the finger, the sound of a tuning-fork placed on the head becomes louder, and the same is the case with sounds produced by our own vocal organs, when the mouth is closed, the escaping vibrations being thrown back upon the labyrinth.

"If the meatus be obstructed by wax, the ear being otherwise healthy, the effect is the same; or, again, if the passage of vibrations through the tympanum be hindered by closure of the Eustachian tube. It may, indeed, be laid down as fairly ascertained that whatever diminishes the permeability of the tympanum or meatus for sonorous vibrations, causes the bone-conducted sounds to be, *cæteris paribus*, better heard. If, therefore, a patient comes to us deaf on one side only, we have at hand an immediate means of determining, with an extreme degree of probability, the seat of the disease. We place a tuning-fork upon his head, and inquire on which side it is best heard, of course being careful not to accept a hasty answer. Generally it will be found loudest on the deafer side, the conducting media being most often in fault. If it is heard best, or only, by the better



ear, the nervous system is in some way concerned in the deafness. It often happens, however, that no decisive information can be obtained in this way, the tuning-fork not being heard unequally on the two sides, although the condition of the ears may be very different. This may be accounted for in many ways, such as inaptness of the patient to discern, or a balancing of morbid conditions producing opposite effects in the ear itself.

"Of course, when both ears are pretty equally involved, this method is not applicable.

"But in all cases the effect of closing the meatus by the finger, while the tuning-fork is placed upon the head, gives important information. If the conducting media are healthy, the sound is rendered louder by the closure; if they are impermeable to vibrations, no effect is produced."\*

*Prognosis.*—The prognosis is generally unfavorable; still, there are cases which are improved by a rational mode of treatment; but it must be borne in mind that nervous deafness is a symptom which may arise from many different lesions, and that, consequently, the same means of treatment, blindly or empirically followed, cannot be applicable in all cases.

ILLUSTRATIVE CASES OF NERVOUS DEAFNESS ARISING FROM  
CONCUSSION.

"B. H., aged forty-four, from a boy thinks his hearing was not sharp, though it was good enough for all purposes, and he never had any disease of the ears. At the age of twenty-two, being then in perfect health and in possession of all his faculties, he was about to call on the parents of the lady who afterwards became his wife; but on the morning of the proposed visit he woke up almost entirely deaf; 'they could not make him hear anything,' he said. He had neither pain nor noise in the ears, but there was a stuffed feeling in them, for which they were shortly afterwards syringed, with no effect. He had gone to bed hearing well. In about a week he began to improve, and was much better in a month; but about two years after he was troubled again

---

\* Hinton, op. cit.



by defective hearing, chiefly on the right side, which, within the last two or three years, has become worse. There has been all the while an occasional singing in the ears, mostly in the right.

"He was a perfectly healthy man, of fair complexion; his hair had turned gray within the last two years. His health had always been good; four or five years ago he had been depressed about business, and suffered much from his throat, but was not then more deaf; now finds himself made much worse by any excitement, especially of an annoying kind; no members of his family are deaf; there is some tendency to gout, but he never had an attack of it. He is moderate in his habits, and a great walker, though lately he has become rather stout. Is more deaf after dinner, or after wine. Does not hear better in a noise. Is a great sportsman.

"My watch was heard on the right side only on contact with the meatus; on the left at two inches. The tuning-fork placed on the head was heard very imperfectly, and less on the right side than the left. The right membrana tympani appeared perfectly healthy; the left, the better ear, was slightly white. He inflated each tympanum with ease, diminishing for the time the hearing of the watch.

"Thus far the symptoms appeared to me entirely unaccountable. Unquestionably we all of us in our turn have felt, or may hope to feel, happily nervous under circumstances like my patient's. But growing stone deaf under them, unless it be to an unfavorable reply, is a phenomenon which pathology refuses to accept. But on carrying my inquiries further, I ascertained that at the age of eighteen (four years, therefore, before the access of the sudden deafness) he had fallen a distance of twenty feet down the stairs of his college, and cut the vertex severely. He was ill for some time after, and the right pupil was dilated at the time, but neither his eyesight nor his hearing was affected. The right pupil is still somewhat larger than the left, but reacts well to light.

"It appeared to me that this circumstance probably contained the key of the case, and that the deafness was due



primarily, and in chief degree, to the apparently innocuous concussion, and only in a minor way to the nervous excitement which seemed to be its cause. The supposition is, that the fall—a cause which so frequently jars the auditory nervous apparatus into an almost complete paralysis—had in this case jarred it into a condition of extreme liability to this condition. Other cases that I have met with seem to justify this influence. A butcher's boy applied in May, 1864, suffering from a high degree of deafness, of a month's duration, with no apparent disease of the organ. He had been thrown out of a cart a week before the hearing became impaired. Another case may be worth reporting more in detail.

“Mary B., a healthy woman, aged thirty-six, single, applied at Guy's Hospital, in April, 1864, on account of deafness, which was almost complete. Four years ago she had fallen down-stairs and struck the back of her head. She was insensible for several days, and on recovering consciousness, found herself quite deaf on the left side, and very considerably so on the right. After three or four months the hearing began to return in the right ear, and soon became moderately good. Loud sounds, however, were painful and confusing to her, and she would become more deaf during a cold, or if excited or worried. The fall also destroyed the sense of smell.

“This condition continued upward of three years, until eight weeks ago, when, after returning from a walk during hot weather, but unattended with fatigue, she became again suddenly deaf in the right ear. Nothing had occurred to agitate her beyond a pleasing invitation. She had worked and stooped more than usual a fortnight before, and the day before, after riding in a cab, had felt somewhat sick. At the time of becoming deaf she had pain in the head. Leeches and blisters had been applied, and medicine given, which produced ‘twitchings;’ she had been galvanized also without result. She complained of tinnitus, like bells or a piano, in both ears. This had previously existed on the left side only. She was unable to repeat words shouted into her ear. The tympana, so far as could be ascertained, were perfectly healthy. Eighteen months afterwards she had



become entirely deaf. Tinnitus less; health good. Each membrane was natural in appearance, except a little whiteness at the upper and posterior parts. Other cases also I have met with in which deafness caused by a blow on the head has ceased for a time and suddenly returned.”\*

Rev. J. M. H., a missionary, aged fifty-nine. First had symptoms of dullness of hearing in August, 1859. In trying to mark out a window in a chapel which he was building in Birmah, put a chair on a table, and in stepping back, forgot that the chair was not on the floor, and fell, making a clean cut on the occiput. The wound healed very quickly, having been dressed by a native with adhesive strips. The sensation in the head was like a bowl of jelly. Having frequent headaches afterwards, the pain all centred in that spot. Over a year after the fall, the deafness was noticed, and gradually becoming complete, with paralysis agitans.

In 1861 and 1862, he could not hear the voices of his family. He used instruments in 1863: first a gutta-percha tube, from England, which did him no good. Hears his own voice perfectly. The trumpet he now uses is the form illustrated in article under Fig. 14, Hearing-Trumpets, and is one of the best forms for such cases. No treatment was of any avail. All parts of the ear were apparently healthy.

There is a variety of nervous deafness arising from compression of the auditory nerve in the internal ear. There is severe and constant headache, vertigo; noises of various kinds within the ear; impaired sight, with or without weakness of the mental faculties, especially of the memory, etc.

Another form is from *cold and its consequences*.—R. W., aged sixty-two, foreman in a large factory, was very much exposed during the winter (1859), contracting repeated attacks of ordinary catarrh, which left him deaf, with no pain. He could hear my watch, indistinctly, close to the left ear; close to the right, rather better; but ordinary conversation in a room he could not hear. He has had a constant humming sound, like that of a saw- or planing-mill, in his ear. No member of his family is deaf; his general health is good.

---

\* Hinton, Guy's Hosp. Rep., xiii. p. 152.



*Examination.*—The external meatus normal; membrana tympani natural form; the handle of the malleus is readily seen; the Eustachian tubes free. On shutting the nose and mouth, and making the effort to swallow, the air is found by means of the otoscope to strike upon the membrana tympani in the middle ear.

In this case a great variety of treatment had been tried, but without benefit.

The following case is that of an intelligent and well-educated physician, in active practice, who consulted me in June, 1862. The only change from the normal condition of his ear was a slight opacity of the membrana tympani. He had consulted almost every distinguished physician and surgeon in this country, but without receiving any benefit. "My history," he writes, "briefly stated, runs thus: when twelve or fourteen years of age, I had typhoid fever, was very low; lay six months; after which I have never been robust. From eighteen to twenty-six I was troubled greatly with dyspepsia for several years; much annoyed at times with general headache; took cold easily, which almost always affected my head. In the summer of 1851 I suffered much with dyspeptic constipation, with headache and a sense of fullness, with great thirst, etc. In October of the same year I had a severe attack of congestion of the brain. In February or March of 1852, soon after a full dinner, while sitting in my room, an unpleasant sensation came into my right ear. The next day I had some ice-cold water poured into my ear, which produced severe pain for a moment, but since then I have never had but very slight pain in the ear. In the summer of 1859 I experienced a similar feeling, as above described, in my left ear; to either I have applied nothing except the water spoken of. I have done nothing by any other medical treatment, except to use for about two months a mixture composed of—

R.—Hydrarg. Bichloridi, gr. ss;  
Potassii Iodidi, ℥ij;  
Syr. Sarsap. Comp.,  
Aquæ, āā f̄ij. M.

Sig. A dessertspoonful one-half hour after breakfast and supper.

With the occasional use of a mild laxative to keep the bowels open.



"In 1860 and 1861 I threw aside tobacco and coffee; both seemed to increase my difficulty of hearing, and also increased the pain in the head."

NERVOUS DEAFNESS PRODUCED BY THE ACTION OF MORBID  
POISONS.

*Syphilis.*—The following cases will illustrate this class. The first was seen in consultation with Dr. Condie, and the other is taken from Hutchinson.\*

John A. McC., aged thirty-two, a bookbinder, of good general health, but almost totally deaf in both ears; born in the United States; mother living and well; no other member of the family deaf. His deafness commenced in the left ear, and, according to the statement of his physician, he had secondary syphilis about this time, was treated by specific remedies for it, and recovered from all the symptoms but the deafness, which increased, eventually extending to the other ear. Has ringing in the ear, as if he had just heard the loud report of a cannon. No pain in head or ear.

*Examination.*—Cannot hear my watch in contact with his ear, or over his mastoid process; but if I place it upon a board, with his face down upon it, he can hear it; he also can hear at times the clock strike in the room, also the sound of cars or wagons in the street. External meatus of right ear normal; left ear narrowed. Some one, he states, advised the use of water of ammonia dropped into this ear, after he had become deaf, which caused a discharge and narrowing of the meatus, slight opacity in both membrana tympani, and greater dryness than natural. The Eustachian tubes pervious. There is not disease sufficient to account for the state of extreme deafness. The throat red and inflamed; tonsils very much reduced in size. It is now some two years, and although Dr. C. has tried both specific and other treatment, no improvement has resulted.

---

\* A Clinical Memoir on Certain Diseases of the Eye and Ear consequent on Inherited Syphilis, by Jonathan Hutchinson, F.R.C.S. London: John Churchill, 1863, pp. 182, 183.



"Susan S., aged thirteen, aspect of syphilis fairly marked, and teeth characteristic, converging toward each other, very short, with a vertical notch or cleft in their free edges, not being so wide there as at their necks. Numerous white deposits of long standing in both corneæ. Both pupils notched, and irregular when dilated with atropine. Nine brothers and sisters are living, none have died. The patient is the third in family, the two elder ones being girls; one of the elder ones has had inflamed eyes.

"She began to lose her hearing in February, 1861. There was no discharge, and not any material earache. She was very much troubled with noises and singing in her ears. The left was rather the worse, but both were affected. She can now just hear a watch pressed over her right ear, but cannot hear it on the left. The ear is dry internally, but there is no other peculiarity visible. The tonsils are not enlarged.

"*General comments.*—It will be seen that all the cases reported by Dr. Jonathan Hutchinson (twenty-one in number), in which the ears were inspected, support the belief that the deafness of syphilitic children is due either to disease of the nerve itself, or to some changes in non-accessible parts of the auditory apparatus. Its symmetry in all the cases would point to a central cause. In none were there found adequate changes in the membrana tympani, although in none was that membrane quite normal. In all the Eustachian tubes were pervious: my belief, therefore, is that the deafness in these cases is due either to disease of the nerves or of their distribution in the labyrinth. These cases constitute the analogues of syphilitic retinitis and of white atrophy of the optic nerves."\*

*Gout.*—The poison of gout may also give rise to nervous deafness and other peculiar symptoms in the head.

---

\* In confirmation of Dr. Hutchinson's observations, Prof. Jos. Gruber has found, in a post-mortem examination of a soldier who died from typhus fever while laboring under syphilitic deafness, there was considerable hyperæmia of the mucous membrane of the drum, and the membranous labyrinth appeared much thickened. The liquid within the labyrinth was sanguinolent and abundant. (*Lehrbuch der Ohrenheilkunde.* Wien, 1870, p. 617.)



*"Presumed immobility of the stapes from gout.*—A. M., aged thirty, a healthy man, has been gradually becoming deaf for ten or twelve years. For about two years the deafness has caused him inconvenience; never any pain in the ears, nor tinnitus; does not hear better in a noise; is worse after a hearty meal, or during a cold, if severe. Smokes three or four pipes daily; has traveled much by rail. No relatives deaf; there is much gout in his father's family, but he never had any symptom of it; sight good; throat slightly relaxed.

"Watch, each ear, seven inches; heard also on the forehead; tuning-fork well heard, sounding a little louder on the right side; no effect from closing either meatus. Right meatus dry; membrana tympani bright, of natural curve; somewhat milky posteriorly. Left meatus contained several loose flakes of epidermis; membrana tympani concave, irregular, and its surface rough. Could inflate the right tympanum, not the left. By means of the catheter the left Eustachian tube was opened, so that he inflated both tympana, which had at first no effect upon the hearing; subsequently it appeared somewhat to diminish it. When air was thus forced into the tympanum the malleus was not seen to move, but on each side the membrane yielded very freely at the upper and posterior part.

*"Presumed affection of the auditory nerve from gout.*—Miss N., aged fifty-two, of languid and worn appearance, had been affected for three or four years with a noise as of rushing water in the right ear, and for about six months with a similar sound in the left ear, but much more intense. It is constant, and occasions her great distress; at times a sudden whistle, like that of a steam-engine, will come in each ear, especially the left, and last a few seconds. During the same period her hearing also has become impaired. No cause has been observed, except that she has long suffered much from severe pains in the head, which begin generally from the right eye, affecting only half of the head at a time, but involving both sides before they leave her. These return regularly every four or five weeks. The deafness and tinnitus are somewhat worse during the attacks. Two years ago she had smallpox, which left her weak, but did not affect the ears. There is gout



on both sides, and she has had slight attacks of it in the knuckles. No relatives deaf.

"Watch, right, four inches; left, contact; tuning-fork not well heard; louder on the *right* (the better) side; closing either meatus increases the sound; membrana tympani bright and natural; Eustachian tubes pervious; throat healthy; inflation of the tympanum produced a slight temporary improvement of hearing for the watch, and exhaustion of air from the meatus gave a little relief for the moment to the tinnitus.

"Probably in both of these cases, in spite of their difference, the gouty diathesis was a chief element.\*

"*Distressing sensations produced in the ears by gout.*—D. T., Esq., aged fifty-four, consulted me in June, 1857. He said that for the last four or five years he had been subject to attacks of gout, which had at times caused him great inconvenience, and the disease had recently made so much progress as to make him fearful that his brain was weakened by its influence. He added that he was never really clear-headed, excepting just after an attack of gout, when he supposed his blood was temporarily freed from the poison. He had rapidly aged in the course of the previous two years. An extremely distressing symptom had lately presented itself in the form of a peculiar sensation of vacancy in the ears, accompanied sometimes by a low humming sound. There was no deafness, and the patient applied to me only on account of the sensations in the ear. On examination, small deposits of gouty matter were observed in the substance of the right upper eyelid, the surface of the meatus externus was of a bright red color, the circumference of the membrana tympani was red, and there was congestion of the long process of the malleus, while the surface of the tympanic cavity was very bright, air passed freely, and with the natural sound, into it. The hearing power was perfect. By what, therefore, could the distressing symptoms be caused? Were they the result of congestion of the nerve?—a condition which, it seemed to me probable, might render

---

\* Hinton, Guy's Hosp. Rep.



the nerve so exquisitely sensitive that the ordinary sounds ever floating in the air might become a source of excitement to the ear. Being aware, also, from previous experience in similar cases, that pressure on the external meatus, so as to shut out or diminish the sound in the meatus, would remove the symptom complained of, I closed with my fingers each external meatus, and the unpleasant symptoms at once disappeared. On subsequently exerting a gentle pressure on the ears, by the introduction of cotton steeped in water into each, the patient was enabled to leave in comparative comfort. For the purpose of preventing the recurrence of the symptoms, it was, of course, requisite to diminish the congestion, for which purpose two leeches were applied below each ear, small doses of colchicum administered, and strict attention to diet enjoined. The quantity of wine was decreased from four to two glasses daily, and in lieu of beef and mutton, of which he had been in the habit of partaking very abundantly, he was ordered to live principally on poultry, game, and fish, with abundance of farinaceous food and vegetables. The result of this treatment was the removal of the distressing symptoms in the ears, and the gradual disappearance of the attacks of gout.\*

"D. W., aged fifty, was sent to Mr. Hinton, in November, 1865, by Mr. Paget, under whose care he had long been, 'with various symptoms of *disorder* of his nervous system, with chronic catarrh of the bladder.' He had always been delicate, but had never had syphilis, nor gone to any excess. First noticed himself a little deaf about fifteen years ago, shortly after a box on the ear from one of his children. A singing tinnitus, now constant, has existed about the same time. Left ear always the worst. Some treatment was early tried in vain, and the deafness continued about the same until three weeks ago, when, without any cause, he became very much more deaf in the course of two or three days. He had been troubled with giddiness for a few weeks previously, which has latterly been worse. His manner was that of extreme nervousness, and his study of his own symp-

---

\* See case of F. S., p. 161.



toms amounted almost to hypochondria. Family healthy; has had no gout; but Mr. Paget having thought he observed indications of a gouty tendency, he has found, on inquiry, that his mother's father suffered from it. Four or five years ago, while hunting, he had an attack of double vision, being otherwise well. Mr. Bowman pronounced the eyes right, but that the nerves did not act in unison. This passed off in two or three weeks, but returned about a year ago, and has continued since at intervals, when the eyes are directed to the side; has *muscæ* occasionally; sees well by each eye with a glass. The right pupil is somewhat larger than the left. No albuminuria.

"He required a raised voice on the right side; on the left could just repeat a monosyllable shouted into the ear. A loud watch was not heard, nor an ordinary tuning-fork placed on the head, except when the right meatus was closed. Right *membrana tympani* thick, and somewhat concave; redness along the malleus; left, thick, rigid appearance, decidedly too concave, and drawn inward. The right Eustachian tube was pervious, and inflating the tympanum somewhat dulled the hearing. The left tube was closed.

"Although, from the history of the case, the nervous constitution of the patient, and the fact that the tuning-fork placed on the vertex was heard on the right side only, it was considered very doubtful whether any improvement to hearing would result from restoring the perviousness of the left Eustachian tube, it was held right to attempt it, not only on account of the deafness, but also with a view of removing a possible cause of nervous irritation in the pressure on the labyrinth, which exclusion of air from the tympanum seems to involve. The obstruction proved to be considerable, but by the aid of bougies the tube was rendered pervious in the course of a month, so that he could freely inflate the tympanum, the *membrana tympani* yielding with a jerk before the air, and resuming a natural appearance. A good deal of mucus was withdrawn from the region of the tube, and warm iodine vapor injected a few times. The hearing seemed at times improved, but ultimately remained about



as at first, and the proceeding was considered a failure. In about a week, however, the hearing began to improve, and in the course of two months it became, without further treatment, in both ears as good as it had been before the last access of deafness.

"In this case, also, there existed both tympanic and nerve disease; and it is an example of a very numerous, if not the most numerous, class of nervous affections of the ear. As a rule, these coincide, so far as my observation extends, with disorder more or less considerable, of the conducting apparatus; and in this fact, scarcely less than in the difficulty of the exploration of the organ, it seems probable to me, is to be found a reason for the great differences of opinion that have existed as to the comparative frequency of affections of the labyrinth and auditory nerve; and in part, also, the doubt which envelops their causes and their nature. It is not always easy, in any case in which the history does not at once vouch for the seat of the disease, to convince one's self that it lies beyond the tympanum, certain affections of which, such as ankylosis of the stapes, may give no visible indication whatever of their presence, but it has been doubly hard to detect a nervous defect through, and in spite of, a disordered conducting apparatus. Even the advance of diagnostic knowledge, revealing morbid conditions of the tympanum before undetected, has tended thus, in some degree, and for a time, to mislead."\*

"*Noises and deafness after rheumatic fever.*—Miss B., aged thirty-six, consulted me in 1851. She stated that ten years previously she had an attack of rheumatic fever, which was followed by dullness of hearing in the right ear, and accompanied by a constant whizzing sound and a pulsation, which extended over the head. The left ear has lately become dull. The watch was heard only when pressed upon the right ear, or at a distance of six inches from the left. Neither ear presented any appearance of disease, except a slight dullness of the surface of the left membrana tympani. The treatment consisted in applying mustard-plasters and stimu-

---

\* Hinton, Guy's Hosp. Reports, Third Series, vol. xiii. p. 160.



lating liniments to the nape of the neck and the ethereal solution of cantharides behind the ears. The result of two months' perseverance was such an improvement of the hearing power that conversation could be more distinctly perceived, and the watch could be heard at half an inch from the right ear.

*"Total deafness following an attack of rheumatic fever.—*Mr. M. G., aged seventeen, a year and a quarter before consulting me had a bad attack of rheumatic fever, which was followed by noises in both ears and gradually increasing deafness, so that in a month after the fever he could not hear any sounds. Since the attack, has now and then heard loud sounds for a very short time; but when he saw me he was so deaf that he could not hear anything, even when the poker and tongs were knocked against each other. The ears had been syringed, blistered, and galvanized without any effect. There was no appearance of disease in either ear. The case was at once regarded by me as incurable.\*

*"Partial deafness following an attack of typhus fever.—*Miss A. M., aged sixteen, saw me on March 1st, 1851. Eleven years previously she had an attack of typhus fever, and during the illness became so deaf as not to be able to hear the human voice. After the symptoms of fever had disappeared, the power of hearing slowly returned, until she was able to hear when loudly spoken to close to the head. There was no appearance of disease in either ear."

*Deafness from the use of sulphate of quinia.—*It is not uncommon for hardness of hearing, or even deafness, to supervene upon the administration of sulphate of quinia. This deafness is sometimes incurable, as the following case shows:

"In 1852, a colleague of mine, chief assistant at the Val de Grâce, presented to me his father, who, for five years, had been affected by complete deafness in both ears; my colleague hoping that I could effect a cure by electrifying the inner ear. The following are the particulars which were given to me in this case: M. X., captain of infantry, had, while in Africa, an intermittent fever, which was followed

---

\* Toynbee, Diseases of the Ear. London, 1868, pp. 359-362.



by a considerable hypertrophy of the spleen, which resisted treatment by quinine in the ordinary dose. The dose, consequently, had to be largely increased, and the treatment long continued. After that treatment, obstinate headache supervened, accompanied by intra-aural noises and hardness of hearing. At the end of some five or six months, the headache disappeared; but the intra-aural sounds continued, and the deafness gradually increased. The patient was subjected to various kinds of treatment, all of which failed, although they were directed by men well instructed in their professional specialty. On examining M. X., I ascertained, as more experienced men had previously, that the Eustachian tube was free on both sides, as also was the external auditory conduit. Nevertheless, the deafness was complete. M. X. could not hear even explosive noises; he only perceived the shock that they produced. I immediately resorted to my Faradisation of the ear. M. X., at each intermittence, perceived neither noise nor sensation in the depth of the ear, or in the tongue, although I had gradually raised the intensity of the current to a much higher than the ordinary degree. After ten trials, which produced no improvement, I abandoned that treatment, which, indeed, after the first electric exploration of the ear, offered me but little chance of success.

“The fact of the incurableness of that case of deafness by the sulphate of quinia, adds importance to the therapeutic fact which I am about to relate:

“In bed No. 13, Saint Martha's ward, lay a young girl, who, for a long time, had suffered under a tertian intermittent fever; the spleen being greatly enlarged, and the disease of long standing. M. Briquet gave the patient a gramme of sulphate of quinia, every twenty-four hours, during nine successive days. From the first day she had a commencement of intra-aural noises, which increased from day to day, and became complicated by a deafness so decided that the ticking of a watch, placed in contact with the ear, could not be heard on either side, nor could even the loudest tones of voice be heard. The patient remained in that state for a fortnight, when M. Briquet, seeing no other means of



cure, requested me to subject his patient to Faradisation of the ear. The operation was performed in my usual manner. On the excited side, the patient heard a sound, and felt numbness in the tongue, followed by prickings. Scarcely was the operation at an end when, on the excited side, she heard the tickings of the watch and the conversation of the persons around her. On the opposite side, the deafness still persisted. On the following day only that side was in its turn subjected to the electric excitation, which provoked the same sensations as the first operation had provoked in the other ear, but not so freely and completely. On the third day I made another electrical operation, which sufficed to restore the hearing to its normal state. From that moment, also, the intra-aural sounds ceased. The patient remained a fortnight after her cure, so that its permanence could be well ascertained.”\*

“*Mumps.*—The peculiar poison which causes the disease generally known by the name of mumps, is very often the source of complete deafness, which, however, usually occurs in one ear only. In these cases the nervous apparatus is evidently affected, as the deafness comes on suddenly, is usually complete, and, as a general rule, no appearance of disease can be detected in the meatus, membrana tympani, or tympanic cavity. When the nerve is not wholly paralyzed, and some, although it may be a very slight degree of, hearing remains, the only plan of treatment which can be recommended is the use of gentle counter-irritation over and around the ears, at the same time that the ear is exercised by means of the elastic speaking-tube.

“*Bile.*—The circulation of bile mixed with the blood is sometimes a cause of nervous deafness.”†

“*Repressed eruption.*—A patient had for several months been troubled with an intra-aural noise, analogous to that of the wheels of a mill. At the commencement of his infirmity, he attributed to the influence exerted on his ears by his business, those sensations which troubled him both by day and by night, not ceasing even when he slept. And he

---

\* Duchenne.

† Toynbee.



at length became so jaded and worn out by these abnormal sounds that he gave up his employment,—that of a miller. But the change produced no relief, and he came to us for advice. On inspecting the outer and exploring the inner ear, we perceived the integrity of both, with the exception of some cicatrices which we noticed upon the pavilion of the left ear. Our patient told us that these cicatrices proceeded from some cutaneous affection with which he had been troubled during several years. Some astringent topical applications had freed him from that affection, and very soon afterwards the aural disturbances commenced. It was beyond all doubt that we had to do with a transmigrating lesion. The ease with which we succeeded in restoring the organ of hearing to its physiological condition, confirmed us in that belief. In fact, we applied a large blister to the nape of the neck, and the raw surface took the character of the primitive cutaneous affection in proportion as the auricular lesion dispersed. Here the acoustic lesion had supervened slowly, and without real pain, for we cannot give that name to the annoyance caused by the subjective sensations. Moreover, the hearing was perfect; it is even to be observed that it was sharper, the patient perceiving the least sounds, even rustling.”\*

*Nervous deafness from mental excitement.*—J. M., aged sixty, a native of England, by occupation a manufacturer. He stated that he was totally deaf in the right ear for twenty-five years, and in the left for twelve years, and that the cause of the deafness was a sudden loss of property; but, upon close inquiry, he also stated that he had a brother and sister as deaf as himself. On testing his power of hearing, by a variety of hearing-trumpets, I found it was not increased by them; neither could he hear the watch when applied over his mastoid process, or between his teeth. Everything had to be written. If the floor was struck, he felt the vibration. He hears a constant noise, as if a wheel were going round, all the time. General health good.

*Examination.*—External meatus normal; membrana tym-

\* *Traité pratique Maladies de l'Oreille*, par Dr. E. H. Triquet. Paris, 1857, p. 436.



pani good color, no defect; Eustachian tube free; throat, pharynx, and uvula slightly thickened and inflamed; tonsils not enlarged.

Has tried all kinds of treatment without benefit. He was sent to me by Dr. J. J. Woodward, U.S.A.

*Hysteria*.—In hysterical deafness you have the usual symptoms of hysteria affecting the senses, but especially the hearing, with intra-aural sounds, as buzzing, ringing of bells, etc., while parts of the ear appear healthy.

In the front rank of the nervous paralysis which Duchenne has seen cured surely and rapidly by the process of Faradisation, he places the deafness which is connected with the hysteric condition. He relates the following case:

*"Hysteric deafness of several months' standing rapidly cured by Faradisation of the motor muscles of the small bones and of the chorda tympani.*—A young girl, habitually quite regular, and otherwise generally healthy, occupying the bed No. 12 in Saint Martin's ward of l'Hôpital de la Pitié, was cured there, at the beginning of April, 1855, of a typhus fever; after which she was not regular again, though the general health appeared very good. During her convalescence at the hospital she had a very warm dispute, just after which she was seized with sobbing and choking, which, however, did not proceed so far as a complete hysterical fit. On the following day, at the very same hour in which on the previous day she had had the dispute which had so much affected her nervous system, the patient fell into a sort of cataleptic state, in which she continued for a whole hour and a half. That species of attack recurred several times during the remainder of her stay in the hospital,—a term of about six weeks. On leaving, she returned to domestic service; but her menstruation did not recur, and by degrees she perceived a constantly diminishing sensibility on her right side, at first in the arm, and then in the leg. At the same time the motion of the limbs was less, as was the strength, so that at length she was obliged to quit her employment as a domestic servant, and return to the hospital. At the time of her doing so, she could not keep in her left hand any object that was placed there. Under the care of M. Briquet, it was ascertained,



by touching, pinching, and electrifying, that her whole left side was affected by cutaneous and muscular anæsthesia (trunk, limb, and face being all affected), and a muscular weakening so great as to make it difficult for her to walk, and to embarrass her considerably in the use of her arms. If any object was placed in her hand when she was not looking at it, she immediately let it fall. The organs of the senses were equally affected in the left side; the sight was considerably weakened, and smell, taste, and *hearing were abolished*. The mucous membranes of the nose, the ear, and the conjunctiva were completely anæsthetized, as was the skin of the pavilion and of the external auditory conduit. The patient was still further affected by continual and very various intra-aural sounds (buzzing, ringing of bells, and whistlings more or less strong and shrill), which completely deprived her of sleep.

“M. Briquet was so good as to permit me to try in this case the therapeutic action of the various modes of electric excitation, which he had seen me successfully apply to a considerable number of patients in his ward who were laboring under similar functional disturbances. In three visits, daily, the electro-cutaneous excitation restored the sensibility of the trunk and the arm (the lower limb was reserved by M. Briquet that he might make comparative trial of the oil of seeds of *Croton Tiglium*). I must add that after the electro-cutaneous excitations, the menstruation, which had been suspended from the commencement of the typhus fever, was restored to its normal regularity. It sufficed for me to pass the metallic rheophores along the nasal and lingual mucous membrane, to restore, with the general sensibility, the senses of taste and of smell. *I also restored the sensibility of the pavilion and of the external auditory conduit; but without modifying the functional disturbance of the hearing.* Then the patient being so placed as to give her auditory canal a vertical direction, I filled it with plain water, and, having plunged a metallic thread into that liquid, and taking care that the metal should touch neither the walls of the auditory conduit nor the membrane of the tympanum, and having placed the other wet rheophore upon the mastoid process, I passed



intermittent currents from my inductive apparatus, with an interval of half a second. The current, extremely weak at first, was gradually augmented in strength, until a disagreeable, though not actually painful, sensation was experienced in the depth of the auditory canal. At each intermission, the patient immediately heard a slight dry, crackling sound on the excited side; but *she experienced no sensation* in the tongue. Immediately after the operation, she could distinguish the ticking of my watch at a very considerable distance, which previously she could not even when the watch was placed in contact with her ear; and she could hear the voice as well on the left as on the right.

"On the following morning I learned that the cure had been but temporary (it lasted an hour), but I ascertained that the hearing was weakened, and that the intra-aural sounds were neither so strong nor so annoying. I renewed, in the same manner as in the previous day, the operation upon the ear; it produced the same physiological effects, and the hearing immediately became normal. From the time at which the patient entered the hospital, her nervous crises had become more frequent; recurring every week, and sometimes lasting for one or two hours each attack. The cutaneous sensibility and the muscular strength, which had been for some time preserved, were lost again, and for several weeks the patient had her left arm contracted. However, she experienced no disturbance as to her hearing, and a few months afterwards left the hospital perfectly cured of her deafness.

"This species of deafness is not always complete; it is generally complicated by intra-aural sounds more or less strong and annoying. Generally the disturbance of the hearing exists only on the left side, as do the other disorders of the sensibility and mobility; but it sometimes also occurs that the deafness is on both sides, though greater on the left.

"It must not be inferred from the foregoing case that hysteric deafness is always a slight affection. I have met with cases which for years have obstinately resisted every kind of treatment. However, although I am not prepared to give the precise statistics of all the cases of hysteric deaf-



ness which I have subjected to electro-therapeutic experiment, I may say, approximately, that hysterical deafness is curable by that treatment in eight cases out of ten. Numerous cases of that form of deafness may be happily modified by peripheric excitation. In fact, I have seen such disappear before the simple Faradisation of the skin of the pavilion or of the external auditory conduit, and even of regions more or less distant from the ear. In those cases the same result would be obtained by any other kind of excitation; as, for instance, exciting injections into the external auditory conduit or of the Eustachian tube, or blisters applied in the vicinity of the ear. The reason is, that in hysterical disorders, as in certain general nervous disorders of the same nature, excitation of a very limited space of the cutaneous surface causes the disorder to disappear as if by magic.”\*

“*Bodily debility*.—It is difficult to draw a correct comparison between the number of cases dependent upon excess of mental excitement and those arising from overtasking the body; but from the data before me, perhaps those of the class now to be considered are the most common. They occur in both sexes, but are more frequently met with in the female; and present great variety of form as well as cause. Sometimes they are temporary, and produced by a long walk, heated rooms, late hours, etc., when noises, with diminished hearing power, come on, but disappear after rest. Other cases, and even some of those which come on suddenly, may, however, remain more or less permanent. Thus, cases have been met with, in my experience, in which patients have become totally deaf after the administration of too violent a purgative, or after an attack of diarrhoea or cholera, and after the nervous exhaustion attendant upon childbirth; in some instances of the latter, the deafness has begun with the birth of the first child, and increased with each successive birth, until at last the nervous power was wholly lost. Perhaps the most common cause of nervous deafness from physical debility is the want of proper care in the management of young persons, and particularly girls, when they are grow-

---

\* Duchenne.



ing fast. In hospital practice, young nurse-maids, who carry heavy children, and whose night's rest is often disturbed, and youths just entering laborious situations, are found to suffer. Any cause, in fact, which reduces the nervous energy of the body to a state too low for the due regulation of the functions of the various organs of the frame, may be followed by a manifest depression of the nervous power of the ears, which shows itself not merely in diminished power of hearing, but often by singing and other sensations in the ear, and sometimes by severe pain, like *tic douloureux*. In cases where debility of the nervous system of the ear is the result of a debilitated state of the body, the pulse, as a general rule, is weak, and there are symptoms of previous or present indigestion. Generally no unhealthy condition of the organ itself is apparent; though in recent cases the cerumen may be softer and more abundant than usual, and in old standing cases may even be absent."

*"Deafness produced by want of sufficient sleep.*—In the early part of 1855, a young gentleman, aged fourteen, was brought to me by his father, on account of gradually increasing deafness. He appeared to be in tolerable health, and was at school in the neighborhood of London. No cause could be assigned for the affection, which had so far advanced as to cause him great discomfort from his inability to hear what his masters said to him. On examination, it became apparent that the deafness depended upon debility of the nervous system, for there was no history of any other disease, nor was there any appearance indicative of disease. The patient had, however, occasional noises in the ears following overexertion, and he certainly was deafer when he was tired. On inquiry, I could discover no special cause for the deafness, as he followed the same rules and regulations which were pursued by all the boys in the school. I prescribed internally quinine, and a stimulating liniment externally, giving directions that he should not be overworked. In a month's time the boy was seen again, but remained in much the same state, so I requested to be allowed to see the lady with whom he boarded, in order to ascertain further particulars as to his mode of living. On the most minute



questioning, no sufficient cause could be detected, except that, being very desirous to prepare his lessons well, he sat up so late that when the time arrived for getting up, he was so sleepy as to be roused with difficulty. It was at once rendered probable that the debility of the nervous apparatus of the ear might be dependent upon the want of sleep; and I therefore requested his friends to see that his duties were so relaxed that he could take as much sleep as he required, and gave directions that he should go to bed at eight o'clock, and sleep until he awoke of his own accord. The result was that for several successive nights he slept for fourteen hours, and by degrees the number was reduced to ten, which was his usual allowance for three weeks, at the end of which time he returned to me, when, to the gratification of all, it was found that his hearing was nearly restored, and he was no longer styled the 'deaf boy' at school. This patient has been seen by me twice or thrice since, at considerable intervals, in consequence of the deafness returning; but each time it was evident that the nervous system had been too much exhausted, and the administration of quinine, with less work, and an increase in the amount of sleep, soon restored the hearing.

*"Total deafness produced by the nervous shock consequent upon successive accouchements.—Mrs. B., aged forty, pale, and of a nervous temperament, consulted me in 1850, on account of complete deafness in both ears. She stated that she had married in India ten years previously, and at the time of her marriage she could hear perfectly well. On the occasion of her first confinement, previous to which her hearing was still perfect, she suffered a good deal from exhaustion, and this was followed by a great degree of deafness, so that she could scarcely hear what was said to her, even when the voice was much raised. Upon getting up, and growing stronger, the deafness was so much relieved that she merely required to be spoken to a little louder than usual."*\*

The following case, occurring in my own practice, confirms the foregoing :

---

\* Toynbee, pp. 367, 369.



Mrs. Anna H., aged twenty-six, general health good, pulse 72, no deafness in her family. Has been deaf for four years, slowly progressing after the birth of every child, and never has had any pain or discharge, but is constantly annoyed by rushing and roaring noises and sounds as of machinery. Is only able to understand her children by the motions of their lips and by means of signs.

The supposed cause was a severe cold, which was followed by slight deafness for two years, with pain in the top of the head, temples, and eyes. Three years ago, she had an attack of partial paralysis of the hand, side, and face, when about six months pregnant. This was stated to have been treated by bathing with salt and water, etc.

On examination, the meatus was found natural and the cerumen of ordinary amount, the membrana tympani transparent, less convex than natural; the handle of the malleus not seen.

By the otoscope, air was observed to pass through the Eustachian tubes.

In the left ear the membrana tympani was more opaque and the meatus narrow.

Her previous treatment had consisted in merely syringing the ears and the introduction of glycerine.

“*Nervous deafness produced by overexertion.*—E. Clarke, aged thirty-one, a tall, muscular carter, was admitted under my care at St. Mary's Hospital, on January 27th, 1859. He stated that fourteen years before, when out of health, deafness came on in the left ear, accompanied by noises, sometimes like a kettle singing, at others like the ringing of bells. On recovering his health, he found himself perfectly deaf in the left ear, which has remained so ever since. Eleven weeks ago, when a good deal exhausted by hard work, he took a bad cold, during which singing came on in the right ear, with loud noises, like the ringing of bells, and were accompanied with so serious an amount of deafness that he required to be spoken to in a loud voice, within a yard of his head. The patient's pulse was weak, and he had a worn aspect, as from exhaustion of the nervous system. On examination, no appearance of disease was visible in



either ear, and the Eustachian tubes were pervious. Two grains of quinine were ordered to be taken twice daily, and a stimulating liniment to be rubbed over the surface of the ears, at the back of the neck, and down the spine. In the course of a week the noises decreased greatly, and in a fortnight they had wholly disappeared, while the hearing power gradually increased; so that when he left the hospital at the end of six weeks, to use his own words, 'he at times hears quite nicely.' '\*

#### ACUTE INFLAMMATION OF THE FACIAL NERVE WITHIN THE FALLOPIAN CANAL.†

"There occurs suddenly a cutting, drawing pain, without previous ear affection, in the cheek and the ear of one side, increasing by pressure on the stylo-mastoid foramen, and accompanied with paralysis of the muscles of the corresponding half of the face. Considerable febrile disturbance attends this affection from the very first day, and soon afterwards deafness and sounds in the ear are ushered in. If the disease is not checked, pain in the head, delirium or coma, and chills, and febrile paroxysms, set in, and finally death. In more favorable cases, an abscess forms under the mastoid process, the opening of which and the exit of the pus exerts a beneficial influence, the pains in the ear and cheek, the paralysis of the face and the fever, the hardness of hearing and the sounds in the ear disappear, and the recovery of the patient takes place.

"If the disease terminates in death, we find, upon dissection, the facial nerve, throughout its course, from the stylo-mastoid foramen outward, spongy, soft, and swollen, the auditory nerve softened to liquefaction; pus in the labyrinth, and in the root of the seventh and eighth nerves of hearing; a large quantity of serum between the membranes of the brain, the cerebrum softened; the membrana tympani and the meatus, on the contrary, in their natural condition. This last circumstance is the most valuable point in the diagnosis be-

---

\* Toynbee, p. 371.

† Translated from Kramer's Aural Surgery of the Present Day.



tween the affection under consideration and acute and chronic inflammation of the labyrinth, attended, as the latter usually also are, with paralysis of the muscles of the side of the face corresponding to the affected ear; whilst this paralysis, in its turn, affords us the best means of distinguishing inflammation of the facial nerve from nervous otalgia.

"An 'exaltation of the hearing' has been described as an especial characteristic symptom of inflammation of the facial nerve, and the consequent paralysis of the muscles of the face, and this phenomenon has been ascribed to a paralysis of the tensor tympani muscle, which is supplied by the facial with a nervous branch (Landouzy, Longet). If we consider, however, that in the case adduced as proof of this, the inflammatory process was only of short duration and moderate development, the ensuing paralysis of the facial nerve, on the contrary, of very long continuance, and in no wise corresponding to the only sometimes accompanying 'exaltation of hearing,' and if also we consider that the same 'exaltation of hearing' takes place in very many other diseases of the ear, we must believe this theory to be entirely unfounded.

"Acute inflammation of the facial nerve is a very rare affection, and is only caused by the violent effect of cold; for instance, washing the face, whilst covered with perspiration, in cold water, cold currents of air directed against the head, when greatly heated, etc.

"The prognosis is doubtful, and through neglect of the patient or physician, the inflammation may extend to the brain and its membranes, and life be endangered.

"The treatment must be strongly antiphlogistic; the necessity of blood-letting must be determined by the general symptoms, and whether the brain is involved. Otherwise, it is sufficient to apply a large number of leeches, which may even be repeated, to the neighborhood of the stylo-mastoid foramen, to rub gray mercurial ointment into the surrounding parts, and, on the occurrence of painful swelling behind the angle of the lower jaw, to apply hot emollient poultices of linseed-meal, and to open with the bistoury, as soon as possible, the abscess, on the slightest appearance of fluctuation.



"If the inflammation communicates itself to the brain and its membranes, with the well-known symptoms, leeches and powerful purgatives, and ice-cold applications to the occiput, are necessary, to prevent, if yet possible, exudations into the cavity of the skull.

"If the inflammation is subdued, without, however, the paralysis of the muscles of the face being prevented, the inunction of pustulating ointment or iodine ointment is proper, in order to promote the absorption of the exudations in the Fallopian canal, and thus to remove the paralysis of the muscles of the face."

#### ACUTE INFLAMMATION OF THE LABYRINTH.

"The pains are in most cases deeply seated in the ear, and attended with violent fever, loss of consciousness, and convulsions, in other cases only moderately severe, and extending over the temporal region, and the corresponding half of the head, and attended with but slight fever, incapability of holding the head erect, drowsiness, vomiting; a sero-purulent discharge takes place from the ear within the first twenty-four hours from the commencement of the attack, but no alleviation is produced, and death may take place in a few days, or in a few weeks, with the well-known symptoms of inflammation of the brain.

"In either case there is found, on dissection, injury to or destruction of the membrana tympani, the small bones of the ear, and the bony labyrinth, sero-purulent effusions into the middle and inner ear, hardening of the cortical substance of the cerebrum and cerebellum of the affected side, and similar results, which show, beyond all doubt, the acute inflammatory character of the affection.

"It may be caused by mechanical injury of the membrana tympani extending to the labyrinth, by sharp-pointed instruments; for instance, by the incautious insertion of a pin for the purpose of cleaning out the ear, or by awkward and forcible attempts to extract, in spite of the struggles of the patient, foreign bodies by means of hooks, forceps, levers, and similar instruments.



"The diagnosis is confirmed by the knowledge of the cause of the pain, and by the discovery by sunlight of laceration or injury of the membrana tympani, and of the filling of the tympanum with blood or bloody serum.

"The prognosis is very unfavorable in cases attended with serious brain symptoms, such as profound coma, etc., and even in those in which the inflammatory symptoms are strongly marked, there is great danger to life.

"The treatment must in all cases be energetically antiphlogistic; besides a cool and elevated position of the head, and the maintenance of quiet in the neighborhood of the patient, cooling drinks, antiphlogistic and purgative medicines, copious and frequently repeated bleedings are necessary around the affected ear, and at the nape of the neck, ice-cold applications to the occiput. If the pain obstinately persists, the ear must be surrounded with warm poultices of bruised meal, day and night, whilst the ice-cold applications are continued to the occiput.

"If by these means the inflammation in the ear and membranes of the brain, the violent pains, etc., are subdued, the life of the patient is saved, but the hearing of the affected side is generally lost.

#### CHRONIC INFLAMMATION OF THE LABYRINTH.

"After an apparently harmless flow from one or both ears, varying greatly in quantity and quality, and attended with a greater or less degree of deafness, has lasted for several years, the patient suddenly complains of dull pains in one ear, extending with greater or less rapidity to the temporal region, the top of the head, and the nape of the neck. Sometimes, but less frequently, however, the pains in the ear are very acute, and extend to the adjoining cheek, the tongue, the soft parts of the neck, and the upper arm, in such a radiating manner that they have been frequently confounded with '*tic douloureux*.' In these cases, the meatus is filled near its entrance, or even in its deeper portion, with extremely sensitive fleshy vegetations, by which the examination with ear specula or probes is rendered very difficult, or even impossi-



ble. In all cases, the pains in the head are rendered unbearable by walking or riding on stone pavements, by taps on the head with the finger, etc., the spirits become depressed, followed by drowsiness, with inability to hold the head erect, and paralysis of the muscles of the face, and even of the upper arm of the corresponding side. Sooner or later are added vague febrile disorders, returning with or without heats and morning sweats, from which is developed typhus fever, with coma, delirium, convulsions, vomiting, etc., and the patient is delivered by death from his sufferings. Long before this stage, the discharge from the affected ear assumes an ichorous and offensive character, and the hardness of hearing is changed to complete deafness.

“If the patient is in a condition in which ocular inspection of the affected ear at a window by strong sunlight is possible, the above-mentioned fleshy growths are discovered, in a few cases, in the meatus, but far oftener the membrana tympani is either partly or entirely destroyed, the small bones of the ear consumed, the tympanum filled with dirty pus, and its lining membrane of a dark-red color, and covered with fungous granulations, or partially destroyed; with the probe, the carious surface of the bone may be distinctly felt. If death ensues, there is, in addition to the caries of the labyrinth, a similar condition of the Fallopian canal, whilst that portion of the dura mater which covers the petrous portion of the temporal bone is found injected, loosened, thickened, and covered with pus, the cerebrum and cerebellum (most frequently the latter) changed in various ways by the inflammatory process, congested, softened, and containing abscesses, which communicate by carious openings in the petrous portion of the temporal bone with the labyrinth and tympanum. These lesions are of so important a character, and the primary ear affection (chronic inflammation of the perforated membrana tympani) thrown so much in the background by long continuance and neglect, that during the existence of this life-endangering complication with secondary brain symptoms, it has been so little thought of that, in explaining the appearances presented



on dissection, a primary character has been assigned to them, instead of to the ear affection (Itard, *Otorrhœa cerebialis primaria*).

"In a very few cases, the periostitis of the petrous portion of the temporal bone does not extend to the dura mater and brain, but terminates in the exit of the pieces of carious bone. The most remarkable instance of the kind known was that of a boy seven years of age, from whose left ear, which had been discharging pus for several years, a carious piece of bone, 1" long and  $\frac{1}{2}$ " thick, was removed, on which the meatus auditorius, 3-5'" deep, the diploë of the petrous portion of the temporal bone, and the two windows were distinctly recognized. After the removal of a second piece of bone, the sore in the external meatus healed within four weeks without any unfavorable symptoms, so that the boy, although, to be sure, completely deaf in that ear, and with paralysis of one side of the face, was in a condition to be discharged from the hospital.

"The chief point in the diagnosis between chronic and acute inflammation of the labyrinth is the slow development of the first from a long-existing inflammation of the membrana tympani, which is probably always perforated, and of the adjoining lining membrane of the tympanum.

"It is seldom that we can discover the causes of the extension of chronic inflammation of the membrana tympani to the mucous membrane of the tympanum, labyrinth, etc., except in the case of taking cold; the abatement of the discharge from the ear, which sometimes takes place at the same time with the occurrence of the life-endangering symptoms, cannot be considered as the cause of the latter, but as the simultaneous effect of some increased irritation of the lining membrane of the tympanum.

"The prognosis is in a high degree unfavorable, especially in cases where polypus and fungous growths are found in the meatus or on the mucous membrane of the tympanum, above all, however, in cases in which typical chills have taken place several times in twenty-four hours, for this is the indication of a fatal formation of pus in the cavity of the skull. Superficial caries in the tympanum is not dangerous



to life; even paralysis of the muscles of the side of the face corresponding to the affected ear is not always incurable. If the danger to life is averted, the hearing is nevertheless destroyed in the affected ear.

"The treatment must be prompt and energetic locally, antiphlogistic; an elevated position of the head, in a cool room, on a horse-hair pillow; copious evacuations by means of calomel, etc.; leeches in as great numbers as the strength of the patient will bear, applied in the neighborhood of the affected ear, and to the occiput; ice-cold applications to the latter; finally, a powerful pustulating ointment between the occiput and the nape of the neck, applied until a plentiful eruption is produced, which should be kept discharging according to circumstances."\*

---

\* Kramer, pp. 15-120.



## CHAPTER XVIII.

### TREATMENT OF NERVOUS DEAFNESS.

WE know of no special remedies for nervous deafness. Its treatment can only be founded upon a correct diagnosis, and a knowledge of general principles. Yet we would invite the attention of our readers to the modes of treatment we have employed, with such modifications as are indicated by variations due to the age, constitution, habit, occupation, etc. of the patient.

"Inasmuch as marked improvement sometimes occurs in this variety of deafness, we should never neglect in the beginning to exhibit the so-called resorbents (among which remedies good nourishment certainly ranks high). Injections of iodide of potassium (grs. ij to xx, distilled water fʒi) through the catheter may be of value in recent cases. In old cases, such as usually come to the aural surgeon, I have never seen any decided benefit either from local or general treatment. Only once have I seen any great benefit to the hearing from local treatment (the use of Politzer's method and the catheter). In this case, however, before any treatment was begun, some of the hearing power had returned after absolute deafness had existed for six months."\*

The treatment of the first class of cases, those which are hereditary, or the result of mental emotion, consists in the use of alteratives and tonics. If these fail, and there is exalted sensibility, then we resort to the use of sedatives and anodynes applied to the Eustachian tube or to the external meatus, such as acetate of morphia, gr. 1-12 to 1-8, dissolved in ʒi of water, with a few drops of acetic acid, to which is

---

\* Von Tröltsch.



added a small portion of eau de cologne. This may be applied on a small pledget of cotton, or by means of a nebulizer, or one of the apparatus before described. Sulphuric ether or chloroform may be used in the same manner.

The second form of nervous deafness results from cold and its consequences. The treatment is commenced by leeching or dry cupping below the ear, by the application of tincture of iodine, with the internal use of small doses of mild purgatives, followed by tonics, as sulphate of quinia, iodide of iron, with cod-liver oil. Change of air to the seashore in extreme hot weather, and later in the season to the mountains, may be of service. For those whose means will not permit them to travel, resort may be had to the salt-bath, friction to the skin, and gymnastic exercises, and care, during the prevalence of easterly winds or extreme cold, to cover the ears. In chronic cases, warm air should be blown by Politzer's air-bag through the nose, with or without the vapor of iodine. Or by means of the nebulizer a solution of sulphate of zinc, grs. iij-v, to water f $\bar{z}$ i, alternating with a solution of muriate of ammonia, grs. xx to water f $\bar{z}$ i, may be employed daily for three or four weeks, or till benefit is derived.

In the third variety of nervous deafness, arising from compression or white softening of the auditory nerve in the internal ear, there is severe and constant headache, vertigo, noises of various kinds within the ear, double hearing, impaired sight, with or without weakness of the mental faculties.

In the treatment of this class of cases, the general condition of the patient must be always borne in mind. If the patient is plethoric, venesection is indicated by cups to the nape of the neck, or leeches at the base of the mastoid process, to be followed by the internal administration of acetate of potassa and sulphate of magnesia, in small and repeated doses, to diminish the excess of serum in the blood. Should these means fail, we may employ alterative doses of some form of mercurial alone, or in combination with iodine or iodide or bromide of potassium. A favorite remedy is the protiodide of mercury, gr. i, in combination with extract of



belladonna, gr. 1-12, or opium, gr.  $\frac{1}{4}$ . If the case should become chronic, we may employ with advantage strychnine from 1-30 to 1-20 of a grain twice a day. We apply the galvanic or electro-magnetic current, a full account of which will be given under the proper head.

The fourth variety comprises a very large class of cases, the result of the poisoning of the blood by typhus, typhoid, scarlatina, cerebro-spinal meningitis, or inflammation of the labyrinth.

In the treatment of these cases our chief dependence is on changing or modifying the blood by tonics and alteratives, with counter-irritation and nervous stimulation. One of the most efficient agents known is strychnia or its preparations. Commencing the treatment with the administration of the sixtieth of a grain of the sulphate of strychnia, increasing the dose until the patient experiences its physiological effects; occasionally intermitting it for one or two weeks, and thus continuing its use for months. In case this treatment should fail to benefit the patient, small doses of Fowler's solution of the arsenite of potassa, gtt. iij thrice daily, in water, will prove useful. This should be given for one week, and then omitted for an equal period, and thus continued for some time, with regular intermissions.

In other cases, in which there is a scrofulous diathesis associated with these affections, chloride of barium will be found a valuable remedy. The solution, in pure water, should be administered in doses of five drops, twice daily, until nausea, purging, or vertigo are produced, always remembering that the remedy is a powerful one, and should be discontinued on the appearance of unpleasant symptoms. We may try the acutal cautery, at a white heat, behind the ear and on the nape of the neck, repeating this at intervals.

If all these means are ineffectual, we resort to the galvanic current of Grove's, Daniell's, or Bunsen's battery (or Siemens-Halske's modified elements, after Brenner's arrangement), one pole applied within the meatus by means of an insulated electrode, in a weak solution of common salt, the other pole being placed in the hand of the patient or over the mastoid process.



The fifth class of cases are seen in connection with hysteria, or as a result of large doses of quinine. In the first variety we depend upon purgatives, antispasmodics, and emmenagogues. Failing in these, we resort to the electro-magnetic current.

In regard to the treatment of those cases which are dependent upon large doses of quinine, it is important that all the eliminating organs should be in active operation, with the persevering use of galvanism or electro-magnetism.

The sixth class of cases are the result of the poison of syphilis. These cases must be treated with specific remedies, but with great care, combining them with tonics, especially the tartrate of iron and potassa.

In young persons who are deaf, the supposed cause being inherited syphilis, and who have the peculiar condition of the teeth described by Mr. Hutchinson\* as pathognomonic of this disorder, viz., the central upper incisors being usually short and narrow, widely separated from each other, with a broad vertical notch in their edges, and having their corners rounded off; the treatment should consist principally in attention to the patient's general health. With regard to drugs, we believe that mercury judiciously used is to be our chief dependence. In children and young persons, the best mode of employing it is by inunction, the mercurial ointment being rubbed into the thighs and armpits twice a week, for twenty minutes at a time. For those who are more advanced in life, the following mixture may be prescribed:

R.—Bichloride of Mercury, gr. i;  
Iodide of Potassium,  $\mathfrak{z}\text{i}-\mathfrak{z}\text{ij}$ ;  
Water,  $\mathfrak{f}\mathfrak{z}\text{iv}$ ;  
Oil of Bitter Almonds, gtt. ij;  
Syrup,  $\mathfrak{f}\mathfrak{z}\text{iv}$ ;  
Brandy,  $\mathfrak{f}\mathfrak{z}\text{ij}$ .

Mix the Bichloride with the Oil of Almonds and Brandy, and then add the other ingredients.

Sig. A tablespoonful to be taken twice a day, after meals.

We should be careful to have the patient well covered by

---

\* A Clinical Memoir on Certain Diseases of the Eye and Ear Consequent on Inherited Syphilis. London, 1863, p. 204.



warm clothing, if the weather be cold, and he should also have a full and varied diet, avoiding green vegetables or acid fruits.

The foregoing prescription may also be used for young persons, diminishing the dose in proportion to the age. After its use for a month or six weeks, the iodide of iron is to be substituted for the bichloride of mercury.

Hinton\* states that "among the poorer classes of the community (London), the number of persons whose hearing has been destroyed by hereditary syphilis is unhappily very large. At Guy's Hospital it has furnished more than one-twentieth of the aural patients. Evidently it is the disease, or one of the diseases, which Sir William Wilde† described as affecting the ears in early life, subsequently to or alternating with an inflammatory affection of the eye. The patients present the now familiar aspect of hereditary syphilis, and have, in every case I have met with, suffered from impaired vision before the deafness has arisen. This deafness makes its appearance generally between the tenth and sixteenth years; about but not precisely coinciding with the period of puberty. The great majority of cases that I have seen have been in females. In one case the deafness was at first greatly aggravated at the catamenial periods, but it became afterwards almost complete."

"Besides their other symptoms, but too well marked, especially among the poor, patients suffering from this disease may, as a rule, at least when they are young, be at once distinguished by the amount of deafness which they exhibit. I know of no other affection, except fever, which in a person under twenty brings on a deafness so rapid and so nearly complete. In the course of a few weeks a girl, previously hearing well, will, without pain or known cause, become unable to distinguish words. Perhaps her eyes, which have been long inflamed, have about the same time been cured. On examination, it is found that a tuning-fork placed on the head is heard for a very short time, or not at all. The

---

\* Diseases of the Ear, by Joseph Toynbee, with a Supplement, by James Hinton, Aural Surgeon to Guy's Hospital. London, 1868, p. 461.

† Aural Surgery, p. 273, Otitis in Connection with Ophthalmia.



meatus is free from wax, the membrana tympani looks somewhat white and rough; it may be flat or too concave, but generally has a dried-up look, as if its juices were deficient. The throat is by no means always unhealthy; the Eustachian tube may or not be pervious. In the latter case, the opening of the tube, by the catheter or other means, affords some relief, which, however, is apt not to be permanent."

#### NERVOUS DEAFNESS CAUSED BY WANT OF FOOD.

In a case of nervous deafness, the only cause for which was the habit of going for many hours without food, Toynbee\* found good results from the use of creasote and morphia. His treatment consisted in making gentle counter-irritation over each ear, the back of the neck, and down the spine, and in administering creasote and morphia, in doses of two or three minims of the former to one-twelfth of a grain of the latter, twice a day. This treatment was continued for four months, while at the same time every means for restoring the general health were resorted to, and at the end of the treatment the hearing power was so greatly improved that the patient could both hear and take part in general conversation.

#### THE TREATMENT OF NERVOUS DEAFNESS BY ELECTRICITY.

The first form of electricity employed in medicine was frictional, obtained chiefly from glass, which, owing to its being so easily dissipated, its too rapid and incontinuous action, confined with great difficulty to conductors, renders its employment rare as a therapeutic agent.

The second form was galvanism or electricity developed by chemical action. Thus, by means of a galvanic current passed through the eye, we produce the effect of light, through the auditory nerve that of sound, and the nerves of smell and taste may be similarly acted on. Dr. Wilson Philip has asserted that he could excite the secretion of the

---

\* Diseases of the Ear, by Joseph Toynbee, p. 372.



gastric juice by sending a current along the divided pneumo-gastrics.

The first apparatus for generating chemical electricity was the pile of Professor Volta, of Pavia, a description of which is published in the *Philosophical Transactions*, London, 1800; although the discovery of galvanism is due to Galvani, Professor of Anatomy at Bologna, who found that by forming a chain of conducting substances between the outside of the muscles of the leg and the crural nerve of a frog, convulsions might be produced. Galvani previously entertained the false idea that the contractions of the muscles of animals were dependent on electricity.

The invention of the pile by Volta was the result of profound thought on the development of electricity at the surface of contact of different metals. The galvanic pile of Volta consisted of an equal number of silver coins and pieces of zinc of the same form, with circular disks of card, soaked in salt-water; of these he formed a pile or column by placing them alternately. If the uppermost disk of metal, either copper or silver, be touched with the finger, previously moistened, while a finger of either hand is applied to the lowest disk, a distinct shock is felt, which is increased with the number of plates.

Instead of the moist conductor, we now use liquids of various kinds, and electricians have devised numerous forms of batteries, but all are based on the important principle discovered by Volta. Still, to Galvani is due the discovery of animal electricity, for he demonstrated that the contact of the nerves of the frog would produce a current of electricity.


In 1827, Nobili confirmed the results of Galvani by means of a delicate galvanometer; but Von Humboldt, Matteucci, and Du Bois-Reymond developed the important facts which are the foundation of electro-physiology, and which led to the application of the galvanic current to the treatment of disease.

The important discovery of induced currents by Faraday, gave Duchenne the means of prosecuting his valuable investigations, and also fired Remak to repeat Duchenne's experi-



ments, and to improve his method by exciting the motor nerves, and not the points of entrance of the nerves into the muscles.

Fig. 101.




This simple apparatus is still employed, only substituting copper for silver and cloth for card, as in Pulvermacher's chain, or in Fig. 101, modified by Dr. Hackley, or in the forms altered by Prof. Hammond and Dr. Palmer. All these batteries are inconstant and not steady in action. They are chiefly applicable to the skin. Prof. Daniell was the first to invent a galvanic battery capable of constant and steady action. He amalgamated the plate of zinc with mercury, preventing local electric action; and used porous partitions, thus freely transmitting the electric current, but preventing the passage of the sulphate of zinc to the copper plate.

Another form of battery was proposed and put in operation by Prof. Grove, of London, which is an improvement even upon Daniell's, in respect to the amount of force generated in a small space. Prof. Smee has also invented a battery, bearing his name, which is very nearly constant. It yields electricity in great quantity, and is always ready for use. This form has been adopted by one of our most intelligent physicians and mechanics, Dr. Jerome Kidder, of New York City, in his electro-magnetic apparatus, also in the form of a compound Smee's battery, which is convenient and portable, and yields all the electricity that is required for operations upon organs as sensitive as the eye and ear.



He has also applied the principles discovered by the lamented Faraday by furnishing his beautiful machine with

Fig. 102.



coils adapted to vary the quantity and tension to any desired degree,—advantages for medical purposes.

By means of two systems of helices or coils of wire, equal alternating currents are produced on their circuits, being closed and opened by a double-armed vibrating arrangement. There are six currents, any two of which can be combined as a negative and positive. The currents from the poles A and D (the bundles of wires in the new machine being covered with a copper cylinder) can be used with a small



wet sponge or with the apparatus of Duchenne (see Fig. 105), placed in the ear with warm water and salt. We have also employed the magneto-electric machine (see Fig. 103, also

Fig. 103.





Fig. 104 of Sexton, modified by Neff), which is especially powerful and effective, and may be substituted for the

Fig. 104.




apparatus which I have described. This form of apparatus presents the advantages of being always ready for use and requiring no initial voltaic current to set it in motion. In it there is but little chemical influence on the organs of taste



and smell, while there is a perceptible influence on the sense of touch, and muscles of the body are caused to contract. The magneto-electric machine at Fig. 103 will not readily decompose the fluids of the body, being useless in removing tumors of any kind, yet in Page's machine (Fig. 104), in which there is a double magnet and intense coils of wire, the current becomes sufficiently powerful for electrolysis or imparting magnetism to iron.

In applying the interrupted or constant current to the ear, we employ a button conductor outside the ear, a sponge, or the apparatus of M. Duchenne.

Fig. 105.—RHEOPHORE OF DUCHENNE.



“That the patient might be spared from pain, M. Duchenne got M. Charrière to construct an auricular rheophore (Fig. 105, A), in which the conducting metallic thread is insulated by an ivory envelope, and cannot penetrate to the membrane of the tympanum. Having waited until the noise had ceased that is produced by the impression of the liquid upon the membrane of the tympanum, place the auricular rheophore in connection with one of the conductors of an inductive apparatus (galvanic or magneto-electric), and close the circuit by placing upon the mastoid process the other wetted rheophore, which itself communicates with the second conductor of that apparatus.

“The inductive apparatus which serves for this experiment is appropriate to the delicacy of the organ upon which it acts; that is to say, the minimum of its power must be scarcely appreciable when the metallic rheophores are applied upon the extremity of the tongue, and it must be in our power to graduate it with precision and upon a scale of great extent.



"The rheophore being placed in the auditory conduit, which was previously half filled with water, and the apparatus being at its minimum, as intimated above, we perceived at the very moment of the intermittence a small, dry sound, like the crackling of parchment, which could clearly be traced to the bottom of the external auditory conduit. The intermittences having occurred with extreme rapidity, the sounds approached to the likeness of those made by the beating of a fly's wings between a pane of glass and a curtain. The intensity of the little sounds which have just been described increased in direct proportion to the increase of the current. Sensation was perfectly limited to the point at which the sound appeared to originate. After a certain time of excitation, and at a certain degree of tension of the current, a kind of tickling in the right side of the tongue, at the junction of its posterior third with its middle third, was plainly felt. Still gradually raising the force of the current, felt the tickling sensation as gradually progressing toward the point of the tongue, then experienced a numbness and a disagreeable pricking, which did not go so far as actual pain."\*

Brenner† has published the following observations in regard to the effect of electricity on the organ of hearing. "If the cathode is placed in the auditory passage, filled with water, and the anode is connected with any other part of the body, there arises, when the circuit is closed, a strong sensation of sound, which continues during the flow of the current, but gradually dies away when the circuit is opened.

"If the anode be placed in the ear, no sound is heard either at the moment of making the connection or during the continuance of the current, yet when the circuit is broken a slight sound is perceived. These reactions Schwartze and Lucae have not been fully able to establish; on the contrary,

---

\* M. Duchenne (de Boulogne), "De l'Électrisation localisée et de son Application à la Physiologie à la Pathologie et à la Thérapeutique." Paris, 1855.

† Zur Electrophysiologie und Electropathologie des N. Acusticus, Petersburger Med. Zeitung, 1866.



the latter has observed that when the cathode (zinc or negative pole) is placed in the ear, and the anode (positive or copper pole) on the neck or hand, a painful drawing sensation is perceived in the ear at the closing and during the flow of the current, which immediately ceases with the opening of the circuit. When the anode (copper pole) is in the ear, a less painful drawing is produced, which also vanishes when the current is interrupted. When the induction current is used, there is, moreover, observed a sensation as of a roaring and rushing wind, which is produced by the presence of water in the auditory passage. There is also a tickling, pricking feeling, which by means of a very intense current may be made unendurable. At the same time there is perceived an unpleasant taste on the corresponding side of the tongue, probably in consequence of the irritation of the chorda tympani, which descends from the cavity of the drum of the ear against the gustatory nerve, and in common with it reaches the submaxillary salivary gland."

Altheus has remarked that there is an increase in the flow of saliva.

"Remak\* found that the galvanic current possesses far more therapeutic properties than the induced currents, and that very often where the former may be beneficial, and effect a cure, the latter will be useless or even injurious. One of the chief advantages of the galvanic current is that, in a painless way, and without shock, we can introduce into the system a large amount of electricity, which is often indispensable, considering the enormous resistance of the body to the electric current. Remak published numerous articles on the galvanic current, but his principal work is '*Galvanotherapie der Nerven und Muskelkrankheiten.*' Berlin, 1858."

I have employed the interrupted current in my own person and in numerous patients. There was never much pain, and the suffering was not more than a feeling of warmth in the ear, with more or less noise, like that of an insect buzz-

---

\* Galvano-Therapeutics: the Physiological and Therapeutical Action of the Galvanic Current upon the Acoustic, Optic, Sympathetic, and Pneumogastic Nerves, by William Neftel, M.D. New York: Appleton & Co., 1871.



ing. If the uncovered button is applied to the skin of the tragus, it is always more or less painful, and if kept long will produce a blister.

Dr. Neftel states that he has made numerous experiments on the healthy and diseased auditory apparatus, and can fully confirm all the statements of Brenner,\* as has also been done by Erb† and by Hagen.‡

“According to Brenner, the pain is quite characteristic for each electrode. When the positive electrode is in the ear, the pain is accompanied by a sensation of pressure, entering from without toward the tympanum. When, on the contrary, the negative electrode is applied, a sensation of something apparently pressing from within outward is felt, so that the individual on whom the experiment is made can distinguish with great precision the direction of the current.

“Vertigo very often attends the application of the electrodes to the fossa auriculo-maxillaris, and to the head; but this never occurs, no matter how strong the current may be, when the electrodes are so applied that a supposed line connecting them is parallel to the plane passing through the antero-posterior diameter of the head. As soon, however, as the line connecting the electrodes forms an angle with this plane, vertigo will invariably take place, and will be most intense when a right angle is formed.

“In both methods the current traverses a more or less oblique plane of the nerve,—*i.e.* is not axial, because the *sympathetic* is also galvanized when the current deviates from the antero-posterior plane. On account of the sensitiveness of the membrana tympani, Brenner considers it better not to introduce the ear-electrode into the meatus, but to apply a button-shaped electrode to the tragus, pressing it inward, I suppose on a sponge, so as to occlude the meatus. The indifferent one must not be applied to the temple or mastoid process, as was formerly done, but rather to a distant part, as the neck or hand.

---

\* R. Brenner, Untersuchungen und Beobachtungen, auf dem Gebiete der Electrotherapie, Bd. i. Abt. i. Leipzig, 1868 und 1869.

† Knapp and Moos's Archives of Ophthalmology and Otology.

‡ Hagen, Electro-Otiatresche. Leipzig, 1866.



"Pflüger\* has discovered that in the electrotonic state the irritability of the nerve undergoes a striking change; it is wonderfully exalted in the region of the cathode (catelectrotonus), and lowered in the region of the anode (anelectrotonus). The reason why sensation is exalted is that alkalies, which are deposited at the negative pole, excite pain, while acids, which are deposited at the positive pole, soothe irritation. Hence the positive pole (anelectrotonus) soothes, the negative (catelectrotonus) excites." The fact is given by the Germans. Its explanation is due to MM. Legros and Onimus.†

"If we apply the cathode of a weak current to the ear, and the anode to the neck or the hand, closing the circuit in the metallic part in which a commutator is intercalated, a pricking, burning sensation is felt at first, followed by the other symptoms accompanying galvanization of the ear. On increasing the intensity of the current, a sound is perceived, usually a distinct metallic ringing (alkali at the cathode excites again). This acoustic perception is most distinct at the closing of the circuit, lasting but a few seconds of the current duration, and not reappearing at the breaking of the circuit. On reversing the current direction by means of the commutator, so that the ear-electrode becomes the anode (acid at anode soothes; no sensation), and the indifferent one, the cathode, no acoustic sensation is perceived, either at the closing of the current or during its continuance. The sensation appears again, however, at the breaking‡ of the circuit, when it is like that perceived at the cathodic closing, though feebler and of less duration. This reaction of the acoustic may be expressed by the following formula (Brenner):

*KaCS*: The closing (*C*) of a galvanic current produces an acoustic sensation (sound, *S*) when the auditory organ is under the influence of the cathode (*Ka*).

*KaDS*>: This acoustic sensation continues with dimin-

\* Pflüger, Untersuchungen über die Physiologie des Electrotonus, Berlin, 1859; Neftel's Galvano-Therapeutics, p. 32.

† Observations on the Effects of Electric Currents, translated by J. Solis Cohen. Philadelphia, 1870.

‡ Disturbing the soothing effects of the acid at positive pole (anode).



ishing ( $>$ ) intensity, and soon completely disappears during the continuance (duration,  $D$ ) of the current.\*

*KaO*—: At the opening ( $O$ ) of the circuit no acoustic sensation is perceived.

*AnC*—: At the closing of the circuit, when the auditory organ is under the influence of the anode ( $An$ ), no acoustic sensation is felt.

*AnD*—: In the anodic duration no acoustic sensation.

*AnOS*: At the anodic opening, however, the same sound is heard as at the cathodic closing, but of less intensity ( $S$ ), and less prolonged.

Brenner has also proved that the reaction of the acoustic nerve is produced by the galvanic current directly affecting the nerve itself, and not through a reflex action from the trifacial, as supposed by Benedikt and others. To use besides the battery only a commutator and current selector for changing the number of elements, while the current flows uninterruptedly, is considered by Neftel a comparatively rough proceeding, or at least a very imperfect mode of treating the acoustic apparatus. It is more or less painful, and is not sufficient for diagnostic and therapeutic purposes. To make the galvanization of the auditory apparatus more exact, and very much milder, it is necessary to intercalate the rheostat† as an accessory circuit.‡

In Brenner's formula the number of elements is marked by Roman figures, and the numbers of resistances in the rheostat by Arabic.

The lower current intensities usually call for the murmurs,

---

\* Disturbance of course greatest at first.

† *Brenner's Stopper Rheostat*.—It consists of three metallic disks on a stand B, C, D. On the circle B, they are numbered by units or quantities from 0 to 10; on the circle C, by tens from 0 to 100; on the circle D, by hundreds from 0 to 1000. The resistance to the passage of the current is made by wires of different lengths, and it can be increased or diminished without breaking the circuit. A wire runs across the disks. To use the rheostat, connect one wire with the poles of the battery and the other wire with one of the electrodes. The radii from these disks are called stoppers, and when they are all at 0 there is no resistance; if at 10 it will make it equivalent to one more, or eleven elements (or pairs of cells of the battery).

‡ One of these rheostats, made in Berlin, we have in operation in our office, with a battery of twenty cups of zinc and carbon.



the higher ones for musical sounds. This harmonizes with the results of Helmholtz's researches, according to which the organs for the perception of murmurs are situated within the vestibule, and those for the perception of musical sounds in the cochlea.

The following interesting case is reported by Dr. M. Meyer,\* and shows the good effects of applying electricity for deafness in conjunction with other remedies.

"Carl M., aged nine years, a lively and smart boy, of a rather scrofulous diathesis, showed suddenly, in May, 1862, signs of difficult hearing, which alarmed his relatives, the more so as already several members of the family suffered from a similar difficulty. Dr. Ehrhard, being of the opinion that the trouble was a rheumatic affection, the examination of the ear showing no disease, ordered corrosive sublimate and baths of potash. After eight days the little patient could hear even a low conversation, musical sounds, rappings, etc.; however, the hearing was very slight. Iodide of potassium was then substituted, and the potash-baths continued. Now this peculiar symptom appeared, viz., that the baths had no action upon the skin, although in the first week they produced a copious diaphoresis. Yet the perception of spoken words improved while other sounds were absolutely unheard.

"The vibrating tuning-fork held against the skin produced no sensation; its application to the thorax caused the surprising discovery that there was an anæsthesia not only of the face, but also of the whole body, especially of the upper portion. As the continued administration of the above-mentioned remedies brought about no change, on the 14th of July, upon the advice of Privy Councillor Romberg, my aid was sought with regard to the application of electricity. One application of the electric brush to the face, forearms, and hands sufficed to remove the anæsthesia in its whole extent, and with it also to relieve the difficulty of hearing. The increased temperature of the skin following this operation, and the free perspiration now ensuing each time after the use of

---

\* Electricity in its Relations to Practical Medicine, by Dr. Moritz Meyer, translated by Wm. A. Hammond, M.D., New York.



a potash-bath, caused in a short time a complete restoration of the sense of hearing, so that the patient could perceive both words and sounds as well as ever."\*

The conclusions arrived at by Dr. Meyer in the use of electricity for many years are:

"1. The *interrupted* current is applicable in those cases in which we wish—

"(a) To excite the muscles, the nerves of sense, the sensory or the motory nerves.

"(b) To produce contraction of the blood or lymphatic vessels.

"(c) Or to affect certain organs supplied from the sympathetic.

"2. The use of the *constant* current is indicated in those cases in which we wish—

"(a) To excite the nerves of sense or the skin.

"(b) To destroy the outer skin or mucous membrane. .

"(c) To produce an increase of warmth.

"(d) To induce a chemical process and also blood coagulation.

"In galvanizing a nerve it is advisable often to change the direction of the current, because the conductivity of a current flowing long in the same direction decreases, while with a change in the direction it increases."

Dr. Neftel† reports eight cases treated by the galvanic current, an abstract of which we give as follows:

CASE I. *Hyperæsthesia‡ of the auditory nerve, with tinnitus aurium; catarrh of the middle ear, and hardness of hearing.*—There is no mention of improvement of hearing, but the most distressing of all symptoms, the noise, disappeared, and did not return.

CASE II.—Same condition of the acoustic nerve as in Case I., with noises and headache. Deafness in left ear from infancy, with anatomical lesions in the middle ear. Improvement of hearing and rapid cure of the noise in the head, and

\* Meyer, op. cit., pp. 342, 343.

† Op. cit., p. 67.

‡ It reacts to a milder current longer and more powerful.



headache. His concluding words are: "I continue to treat the patient once a week; his hearing is normal; he has no headache, no noises in the head; and the noises in the ear are very slight, and disappear entirely after each galvanization." (Do they not, however, return?—for the writer has treated cases which he thought were cured, but the noises returned.)

CASE III.—Same condition of the acoustic nerve as in the preceding cases; hardness of hearing of one ear and deafness of the other (no details given of hearing distance), a case of otitis media purulenta. Politzer's method of inflation was used. The conclusions are as follows: "I repeated the galvanic treatment daily during one week, the patient feeling remarkably well, the heaviness of the head, the tinnitus aurium, and discharges from the ear disappeared; even the ear-wax, the secretion of which had entirely ceased for many years, returned, and the hearing became normal."

CASE IV. Also a case of *hyperæsthesia*,\* *with loss of the olfactory sense; tinnitus aurium, and hardness of hearing; return of the olfactory sense, and improvement of hearing.*—The following is the conclusion: "The noises have returned, but in so slight a degree that they do not interfere with his irritable disposition. After repeated treatment his hearing has much improved [hearing distance not stated], and his olfactory sense is perfect." This is a remarkable case, as the results of Drs. Beard and Rockwell's treatment have been either negative or partial, and in no case complete.†

CASE V. *Tinnitus aurium, with two different noises; torpor of left acoustic nerve; hyperæsthesia of the right ear; left ear completely deaf; improvement of hearing after treatment, with disappearance of dizziness and one kind of noise.*—Neftel says: "The hearing has improved wonderfully, so that the lady is able to converse, and can hear such murmurs as, for instance, the

---

\* The excitability of the acoustic nerve may be either morbidly increased (Brenner's hyperæsthesia) or it may be greatly diminished (torpor of the acoustic nerve).

† Anosmia, or Loss of Sense of Smell. A Practical Treatise on the Medical and Surgical Uses of Electricity, p. 584. W. Wood & Co., New York, 1871.



scraping of a pen, etc., which formerly she could not at all perceive."

CASE VI. *Facial paralysis of right side, with hardness of hearing.*—The treatment consisted in galvanization with proper precautions. In the course of a month the patient recovered from both the paralysis and hardness of hearing.

CASE VII. *Torpor of the auditory nerve, lasting from childhood, in a man aged sixty.*—Improvement of hearing after treatment.

CASE VIII. Also a case of *torpor of the acoustic nerve; total deafness in right ear for two years, and failure of hearing in the left.*—Immediately after the first treatment loud conversation could be heard with the right ear from a considerable distance. The hearing continued to improve.

Dr. Neftel concludes as follows: As can be seen from the foregoing cases, to which I could add a considerable number of others, the treatment of affections of the acoustic nervous apparatus is accomplished by the same means by which they have been diagnosticated.

Every unprejudiced observer will agree with the closing sentence of Brenner: "Henceforth a diseased ear cannot be considered fully examined, and the means for its cure exhausted, so long as for these purposes the galvanic current has not been used in a scientific manner."\*

Dr. S. Moos† publishes the following remarkable case of recovery from nervous deafness, which we have condensed:

Miss S. D., aged nineteen, took a severe cold on February 9th, 1869, resulting in an attack of acute articular rheumatism, which run its usual course during four weeks. In the fifth week nervous (hysterical) symptoms appeared, which lasted until the seventh week, when they began to decline. At the end of the seventh week, the patient complained of a terrible pain behind the left ear. The organ of hearing was now very sensitive, every noise caused pain, and the acuteness of hearing was very great. In the eighth week reconvalescence seemed to set in. The sensitiveness of the organ of hearing

---

\* Neftel, op. cit., p. 81.

† Archives of Ophthalmology and Otology, vol. i., No. 2, p. 464.



decreased, accompanied by hardness of hearing, which increased to such a degree that at the end of the ninth week she was perfectly deaf, and it was necessary to communicate with her in writing. In the eleventh week, the patient became affected with hystero-epileptic attacks, with pains in the abdomen and back, loss of consciousness, tetanus, and at last clonic cramps. On the 3d of May, she complained for the first time of a severe pain in the brain, on the left side, accompanied by excessive hyperæsthesia of the scalp, which was decidedly unilateral. There was hyperæsthesia of the face, also unilateral, but not as painful as the scalp. On the 5th of May, he used a constant current of 5 Meidinger's elements to the sympathetic nerve of the neck for three minutes. On the 22d of May, the patient heard her own voice with her left ear, and the next day heard it with the right. From the 18th to the 24th of June, conductibility of the bones was noticed for the first time.

On the 12th July, the hearing distance for the watch was, on the right, ten feet; on the left, nine feet. July 27th, she was sent to spend six weeks in the Black Forest. There she recovered completely.

Dr. Moos, in conclusion, says: "May this case toll the funeral knell for all those opponents of a therapeutic agent which, when employed according to proper indications, may yet prove a rational and grateful means not only for certain muscular and nervous affections, but also for some ear diseases, though their number may be limited!"

Each individual operator prefers a special battery. We are in the habit of using zinc and carbon elements, so as to introduce into the circuit any number. The conducting wires being well insulated, we have a current-selector, so that we can increase or diminish its intensity by adding or removing any number of elements. To produce the current density, by introducing resistance, we employ Brenner's rheostat, which we have already described, by means of which from ten to twenty-one thousand units of resistance can be introduced into the accessory circuit. There is also a commutator, by means of which the direction of the current can be reversed; also a galvanometer, indicating the



direction, its approximate intensity, its constancy, and fluctuation.

Brenner\* details eleven cases of diseases of the ear treated by the galvanic current.

"In one case of thickening of the drum, the current caused absorption.

"In one case of hyperæsthesia, with tinnitus aurium and anatomical changes in the middle ear, the tinnitus was rapidly cured.

"In one case of hyperæsthesia, after the use of quinine, there was recovery.

"In one case of hyperæsthesia, with tinnitus aurium and catarrh of the middle ear, the tinnitus was cured.

"In one case of obstinate subjective symptoms of various kinds, there was improvement under great difficulties of application.

"In one case of noises in the head and ears of ten years' standing, with important anatomical changes in the ear, there was improvement.

"Of deafness, two cases were improved, one was much improved, and one was cured. The case which recovered was one of facial paralysis, with anomalous reaction of the auditory nerves.

"In all the cases there were anatomical changes. In some cases the treatment was quite persistent."

*Case of nervous deafness tested by Brenner's formula; no improvement in the noises or the deafness.*—Miss E. G., aged thirty-five years; has been healthy, but of an extremely nervous and sensitive temperament. Is in business, but owing to her deafness it is almost broken up. The deafness has continued to increase in spite of prolonged treatment under a number of physicians; also has constant noises in both ears. (No cause except catarrh.)

---

\* Op. cit., Band i. 2 Abth. p. 233, et seq. Brenner also mentions the fact that he failed in seventeen cases of tinnitus, loc. cit., p. 235, *Medical and Surgical Electricity*, Beard and Rockwell; Wm. Wood & Co., New York. Also R. Brenner's *Researches and Observations in Electro-Therapeutics. On the Effects of the Electro-Current on the Auditory Apparatus in Healthy and Diseased Conditions. An attempt at founding a system of Rational Electro-Otiatrics*. Leipzig, 1868.



Left ear, July 31st, 1871. Acuteness of hearing for speech very much diminished; requires to be spoken to in a very loud voice.

Ordinary watch cannot be heard on left side; meatus, membrana tympani, and orifices of Eustachian tubes normal; voice not natural; loud-ticking watch was heard on right, two inches; tuning-fork best on the left.

External arrangement of conductor on external ear, without water in the meatus, and without rheostat; no impression with 5—10—20—40 cells. With warm salt and water in the meatus with rheophore and the rheostat, the following was the unsatisfactory results:\*

July 24th, 5 P.M.

20 cells, Ka in ear.	20 cells, 50 resist, bell and hammer,
" 20 resist.	both.
" 30 " K cl, bells and hammer, but do not remain.	K cl, hammer.
K o.	" 60 " no reaction.
A o.	" 30 " "
A cl, hammer.	" 100 " K cl, hammer.
A Dur, "	K D, "
" 40 " K o, "	" 200 " "
	15 " 150 " "
	20 " 30 " K cl, sounds.
	K D, confusion.

July 27th.

20 cells, 0 no reaction.	An D, no sound.
" 10 resist.	K cl, "
" 50 " K cl, slight sound of hammer.	20 cells, 70 resist K cl, slight sound.
K D, no sound.	d, no sound.
An d, "	Ano, "
" 60 " K cl, slight sound.	cl, twice.
An D, " "	d, no sound.
An O, " "	" 100 " K cl, slight sound.
A cl, no sound.	A cl, "

\* In the table, a, means anode; k, kathode; o, opening; cl, closing; d, duration.



July 31st, 1871, 5 P.M.

	K, in left ear.	20 cells, 20 resist	K cl.
	A, in hand.		K d.
20 cells, 10 resist	K cl.		K o.
	K D.		An cl, slight sound of hammer.
	K O.		An d.
	An cl.		An O.
	An d.	" 30 "	K cl.
	An O.		An cl.
	K d.		An d.
	K o.		An O.
	An cl.	20 cells, 100 resist	An cl, sound, bell.
	An d.	" 210 "	K cl, hammer.
	An o.	" 220 "	K cl, no results.
20 cells, 40 resist	K cl.		K d, "
	K d.		K o, "
	K o.		

*Case of nervous deafness, the result of gout, tested by Brenner's formula, with no noise in ears.*—A gentleman of leisure, of good family, aged forty, predisposed to hereditary gout, and of nervous temperament, began to become deaf without any apparent cause. In about three years there was complete deafness on one side, and insensibility to any sounds, except those spoken directly in front of the ear, on the other side. An ordinary watch not heard ticking in contact with the ear. An ordinary tuning-fork not heard by bony transmission, though a very large one could be heard on the best side; and by trying a number of them varying in pitch, some of the higher ones could be heard more readily than others. The worst side was apparently insensible to sound of any kind, until the vibration of large tuning-forks was intensified by means of Helmholtz's attuned metallic resonators placed in the ear; which induced sensation of sound in each instance, but not sufficiently distinct to distinguish their musical character. No tinnitus aurium in either ear at any time.

General health tolerably good, but some nervous debility and mental depression on account of the infirmity. External meatus healthy in color, and curvature normal, but exhibiting a gouty concretion below handle of malleus; tympanic membrane, Eustachian tubes, and their pharyngeal orifices normal in every respect.



The best ear responds promptly to Brenner's formula for the normal acoustic reaction to the electric current. No sonorous reaction whatever, normal or abnormal, on the part of the worst ear, although as many as thirty cells were intercalated in the circuit, producing intense pain, not followed, however, by any after-effects, good or ill.

The large trumpet (Fig. 14) permits the patient to hear ordinary conversation, with some effort, as the sounds are confused by the resonance of the instrument.

Treatment with strychnia, internally, one-sixteenth of a grain, and locally, by the electric current, negative in a strictly professional sense, though the patient and his family claim some slight improvement from it.



## CHAPTER XIX.

### DEAF MUTISM.

THE term "*Deaf and Dumb*," or more properly "Deaf Mute," is applied to a person born deaf, and therefore unable to speak. Of deaf mutes there are two great classes:

1st. Those who are congenitally deaf, or have never spoken.

2d. Those who become deaf before the fourth or fifth year, but who have spoken. These are the "semi-mutes," and constitute one-half of the whole number of deaf and dumb. Many so-called deaf mutes are as intelligent and bright as persons possessed of all their senses. The deaf mutes of our day have many advantages over those of the last century, as they are now taught conventional signs in the schools devoted to that object, the alphabetical signs, the articulation of sounds, reading and writing, and reading on the lips. The boys are taught various trades, whilst the girls are instructed in plain sewing, dress-making, and domestic employments. History informs us that so-called civilized and refined people, such as the Egyptians, Greeks, Hebrews, and Romans, looked upon the deaf and dumb as allied to the brutes. We are told that in some portions of the East even at the present day they are obliged to perform the hardest and most degrading duties. "It was reserved for Christianity," says Wilde,\* "to conceive the noble idea, and for modern genius and perseverance to perform the exalted work, of developing the faculties of the unfortunate deaf mutes, of giving to them the knowledge of good and evil, of rendering them independent, and of awakening within them feelings and aspirations which, but for the beneficial enlightenment of education, would forever have remained unknown to them."

---

\* On the Physical, Moral, and Social Condition of the Deaf and Dumb, by W. R. Wilde, M.D. London, 1854.



They were the special objects of Christ's miraculous love, and we find many striking examples of it in the New Testament, but none more so than in St. Mark :\* " And they bring unto him one that was deaf, and had an impediment in his speech ; and they beseech him to put his hand upon him. And he took him aside from the multitude, and put his fingers into his ears, and he spit, and touched his tongue ; and looking up to heaven, he sighed, and saith unto him, Ephphatha, that is, Be opened. And straightway his ears were opened, and the string of his tongue was loosed, and he spake plain."

The first practical attempt to teach the deaf mute was made by Petro de Ponce, a Benedictine monk, at Sagahun, about the middle of the sixteenth century. He taught the two mute sons of a Castilian nobleman. But it is to the Abbé de l'Epée, of France, that the deaf mute is most indebted, as he devoted his whole life and fortune to devising his system of instruction, which he brought to great perfection in the "Royal Parisian Institution." He was born November 12th, 1712, and died 1789.

"It is stated that a casual circumstance drew his attention to the education of the deaf and dumb. Calling on a lady one day on business, he was told that she was not at home. Being invited to walk in, he entered the house and found two young females occupied in needle-work. Making the usual salutations, he was surprised at receiving no reply. While wondering at what appeared to him a want of politeness, their mother came in and solved the mystery, by informing him her daughters were deaf mutes. His heart was filled with compassion for them when he thought of their isolated condition. He gave himself up to devise some plan by which their condition could be ameliorated. An ecclesiastic named Vanin had commenced instructing them. Death removed him, and their condition appeared hopeless when De l'Epée appeared. He invited them to come daily to his abode, and from this time he dedicated his life to the cause of the deaf and dumb. He reflected that savages of

---

\* St. Mark, vii. 32-35.



different languages were able, by means of signs, to communicate to a certain extent with each other.\*

"It struck him forcibly that signs were the natural language of the deaf and dumb. He adopted them as his plan for instruction, enlarged and improved them, and reduced them to a system. But his precious life was too short to complete it. Fortunately, he had instructed the Abbé Sicard, and when on the 23d of December, 1789, he was called, in the seventy-seventh year of his age, from time to eternity, there survived a man worthy to succeed him.† This venerable Christian had aroused public attention to the wants of the deaf and dumb, had vanquished the prejudices that existed against them, and had called forth a generous sympathy in their behalf. They became the objects of public benevolence and private beneficence. Numerous schools were founded for them. Pious, learned, and capable teachers arose to instruct them.

"The most eminent of these was Roch Ambroise Cucurron Sicard. He was born at Fousseret, near Toulouse, on the 20th of September, 1742. He was educated in that city, and on the completion of his studies entered into holy orders.

"The good and learned Archbishop of Bordeaux sent him to

---

\* "Major Long, in his Expedition to the Rocky Mountains (Philadelphia, 1823, vol. i. p. 578, etc.), gives upwards of one hundred Indian signs. Among which are 'Man, finger held vertically.' 'Sun, the forefinger and thumb are brought together at the tip so as to form a circle, and held up toward the sun's track. To indicate any particular time of the day, the hand with the sign of the sun is stretched out toward the east horizon, and then gradually elevated, to show the ascent of that luminary, until the hand arrives in the proper direction, to indicate the part of the heavens in which the sun will be at the given time.' 'Fire, the act of striking fire with the flint and steel is represented, after which ascent of the smoke is indicated by closing the fingers and thumb of the right hand, holding them in a vertical position with the hand as low as convenient, the hand is then gradually elevated, and the fingers and thumb a little expanded, to show the ascent and expansion of the volume of smoke.'

"Indians, on several occasions, visited the Pennsylvania Institution for the Deaf and Dumb, and were able to hold a conversation with the pupils by signs."—*Memorial of A. B. Hutton, by James J. Barclay, Esq., Secretary Pennsylvania Institution.*

† He deprived himself of the comfort of fuel for a whole winter, in order that he might have larger means to aid the deaf and dumb.



Paris, and had him placed under the instruction of De l'Epée. He soon entered enthusiastically and energetically into the views of his master, the value of which he was not slow to perceive and appreciate. Endowed with a vivid and fertile imagination, he had remarkable skill to clothe abstract ideas with sensible forms; he had a peculiar talent for pantomime, which is the appropriate language of deaf mutes. Gifted with a pliant and active spirit, he eagerly sought and discovered new and varied means to express and explain his ideas or his precepts. It seemed that to teach the mute was his natural vocation, so admirably was he qualified for it. Adopting the fundamental principles of De l'Epée, he seized, above all, this primitive idea,—that the instruction of deaf mutes was a translation, that mimic signs was their mother tongue, that the conventional language of society was to them a foreign one, which, by the aid of translation, might be taught them. He reproduced this idea under new forms; he reflected upon it incessantly. The language of signs became to him a natural language; so much did he love it, and so skillful was he in its use, that he not only used it, but he enlarged it, enriched it, moulded and remoulded it."

"Sicard, now admirably trained to teach, was recalled from Paris by the Archbishop of Bordeaux, and placed at the head of the school established by him.

"Sicard's fame as an instructor soon spread, and his success rendered him celebrated, and gained him clerical advancement. He was made Vicar-General of Condom, and a canon of Bordeaux."\*

In 1760, Samuel Heinicke,† of Germany, developed the "Artificial Method," which had for its principal aim the development of the power of speech, which exists in all save a few deaf mutes, and the training of the eye of the mute to interpret the duty of the ear. This is now called the "German Method." In England, the first person who conceived a plan for instructing the deaf mute was Dr. John Wallis, Professor of Mathematics in the University of Oxford. In

---

\* Memorial, op. cit.

† Heinicke, a Saxon by birth, born in 1723, and died in 1790.



1662 he exhibited to the Royal Society his first pupil. Thomas Braidwood opened a school on the system of Wallis, in Edinburgh, in 1760. From this sprang all the institutions of a like nature in Great Britain. In 1783 Braidwood removed his academy to Hackney, near London. He died in 1806, but his institution was continued by his widow and grandchildren until 1816. The work which, according to Wilde, gives the fullest account of his mode of instruction is "*Vox Oculis Subjecta*," a dissertation on the most curious and important "art of imparting speech and the knowledge of language to the naturally deaf and (consequently) dumb," with a particular account of the academy of Messrs. Braidwood, of Edinburgh, and a proposal to perpetuate and extend the benefits thereof, as published in London, in 1783, by an American gentleman whose son Braidwood had educated.

Articulation was one of the chief means employed at this academy, and it would appear that some success was attained in speaking by the pupils educated there. To assist in imparting a knowledge of vocalization, we are told that the master employed a small instrument composed of "a small, round piece of silver, a few inches long, the size of a tobacco-pipe, flattened at one end, with a ball as large as a marble at the other," for placing the tongue of the pupil in the right position. Soon after special instruction was provided by the government and private benevolence.

In 1807, the deaf-and-dumb daughter of Dr. Cogswell, of Hartford, Conn., attracted the attention and enlisted the sympathies of the late Rev. Thomas H. Gallaudet, "a man who united the true principles of charity with distinguished scholarship and unwearied assiduity and patience."\* He subsequently proceeded to Europe, in order to acquire the art of imparting instruction to deaf mutes; and having remained for some time a pupil under the Abbé Sicard, he returned to the United States, accompanied by M. Clerc, who was then (in connection with Massieu, the favorite pupil of Sicard) one of the mute assistants of the Parisian Institution.

---

\* Op. cit., Wilde.



"When Mr. Scixas retired (November 3d, 1821) from the Pennsylvania Institution for the Deaf and Dumb, M. Clerc, with the consent of the Directors of the American Asylum, took temporary charge of it.

"He reorganized it, introduced the method of Abbé Sicard, and gave instruction and advice to the assistant teachers until May, 1822, when he was recalled to Hartford, and was succeeded by Lewis Weld, the pupil of Gallaudet and Clerc.

According to Mr. Barclay, M. Clerc was remarkable for the grace with which he made signs. They were intelligible to those who were not familiar with the language of signs.

"In 1818 he was introduced to a young Chinese, who was alike unacquainted with the French and English. M. Clerc knew nothing of the Chinese. Yet, by means of signs, they were enabled to communicate with each other; and M. Clerc learned from his Chinese acquaintance "many facts respecting the place of his nativity, his parents, and their family; his former pursuits in his own country, his residence in the United States, and his notions concerning God and a future state." By the aid of appropriate signs, M. Clerc ascertained the meaning of about twenty Chinese words.\*

---

\* During the period when Mr. Dunn's Collection, the "Chinese Museum," was in Philadelphia, a Chinese whom Mr. Dunn had brought to this city, and who attended at the Museum, often visited the Pennsylvania Institution for the Deaf and Dumb, and was enabled to hold conversation with the pupils by means of signs.

About two years ago, some Japanese students from Rutgers College visited the Institution, and one of the older boys conversed with them without difficulty, in signs, to the great interest and pleasure of the visitors.

While Mr. Stansbury, formerly of the New York Institution, was in St. Petersburg, he visited the Institution for the Deaf and Dumb in that city. He knew nothing of the Russian language. He told a short story, in signs, to the pupils, who immediately wrote it on their slates.

In 1819, while Mr. Gallaudet was at Cornwall, Conn., he had an interview with a number of young persons from the South Sea Islands, and from different tribes of North American Indians. He held an interesting conversation with them in the language of signs. A number of questions were asked and correctly answered, and the meaning of a number of Owhyheean words ascertained. It appeared that not a few of the signs used in the instruction of the deaf and dumb were the same as those used by the South Sea Islanders, to supply the deficiency of, or give emphasis to, their own comparatively defective language.



“Advancing years rendered the labors of teaching too onerous for M. Clerc, and after forty-one years of arduous duty, he retired from the chair which he had so long and so ably filled. His valued services were not forgotten by the directors of the asylum.”\*

At the time a silver pitcher and salver were presented to Mr. Gallaudet, a like memorial of affection was also presented to M. Clerc.

Mr. Lewis Weld discharged his various and important duties in a manner highly satisfactory to the board and advantageous to the pupils, until October, 1830, when, on the resignation of Mr. Gallaudet, he was invited to take charge of the American Asylum, the scene of his early labors.

When Mr. Weld retired from the Pennsylvania Institution, in 1830, Mr. Hutton† was appointed his successor.

\* Memorial, op. cit.

† ABRAHAM B. HUTTON, Esq.—The object of this notice, although not a medical man, was so associated with medical matters in his capacity of principal of the Pennsylvania Institution for the Instruction of the Deaf and Dumb, that a short notice of him is considered proper.

Mr. A. B. Hutton was born in Albany, New York, September 10th, 1798, and received his literary instruction at Union College, where he graduated in 1817. Subsequently he entered as a student in the Theological Seminary at Princeton, New Jersey, in 1819. Owing to an affection of his throat he was obliged to relinquish the study of divinity. In March, 1822, he entered the Pennsylvania Institution for the Deaf and Dumb as an assistant teacher. In this capacity he gave such entire satisfaction that, when Mr. Lewis Weld, the principal, was called to take charge of the American Asylum for the Deaf and Dumb, at Hartford, Mr. Hutton was appointed, on the 4th of September, 1830, as his successor.

Mr. Hutton possessed all the requisites for his new position, having a cultivated mind and refined taste, with gentleness and firmness combined, also great aptness in teaching. His attainments in natural philosophy, mechanics, and chemistry were very considerable, and he took great pleasure in lecturing on these subjects, both to the advantage of the pupils and the entertainment and instruction of the numerous visitors at their weekly exhibitions. But everything was subservient to the great object of his life,—the successful instruction of the deaf mute. To this subject he devoted all his energies, and was alive to every improvement. The success with which his labors were crowned, and the thorough appreciation of his efforts by the managers of the institution, his friends, and the community at large, are the best evidences of his ability and fidelity.

For several years he suffered from an affection of his kidneys, and yet he was always at his post. In the spring of 1869 the disease increased, but



“He made himself master of the science of instructing the deaf and dumb. After a full and careful examination of the two systems of instruction, he became convinced of the superiority of that of signs with its auxiliaries, dactylology, writing, etc., over artificial articulation and lip-reading, and found no reason to suggest any alteration of the Sicard system introduced into the Institution by Clerc. That system he pursued to the close of his life, fully satisfied that it was the best system. His large experience and ripe judgment entitled his opinion to great consideration. For upwards of forty years he discharged the arduous duties of principal with great success and fidelity.

“Laurent Clerc died a few months ago (1870), at the age of eighty-three, much respected by a large number of pupils, who are scattered all over this country.”\*

The first institution for the instruction of deaf mutes in the United States was opened at Hartford in 1817. Since then the number has increased to thirty, and the interest in the good work has also increased in the same proportion. It is known that there is now (1870) one deaf mute in every two thousand of the whole population of the United States. According to the census of 1870, the population of Pennsylvania is three million five hundred and eleven thousand five hundred and forty-three, which, by the above ratio, will give seventeen hundred and fifty-five as the whole number of deaf mutes of all ages in the State. The number of annual births in the United States being in the proportion of about one to twenty of the population, we may safely assume eighty-eight as the number born or made deaf yearly in Pennsylvania.†

The semi-centennial anniversary of the Pennsylvania Institution took place February 28th, 1871, and we had the

---

he did not leave the city until July, 1870, when he made a visit to his sister, and soon after died on the spot where he always wished to die, ministered to by the affectionate hands of those he loved.

The directors of the Pennsylvania Institution, feeling great confidence in the experience, ability, zeal, and high moral character of Mr. Joshua Foster, have elected him principal, to fill the place of the lamented Mr. Hutton.

\* Mr. Barclay's Memorial.

† Report of the Pennsylvania Deaf and Dumb Institution, 1870.



privilege of enjoying the gathering of the pupils from all parts of the country, and the pleasure which they took in this interesting occasion, with their success in getting up a variety of amusements. We were informed that the number of deaf mutes who have enjoyed the privileges of the institution since its commencement, fifty years ago (including the present inmates), is nearly fourteen hundred.

In their last Annual Report,\* the directors stated their intention of introducing more systematic instruction in articulation. For this purpose, they appointed Miss Rebecca H. Cropper; and to prepare her fully for the situation, they were, through the considerate kindness of the principal and directors of the New York Institution, enabled to place her under the tuition of Mr. B. Englesman, whose reputation as an instructor in this art deservedly stands very high. After remaining under his instruction for two months, she returned to this city, and at the commencement of the present term began a course of teaching the semi-mutes. "Sufficient time has not elapsed to enable the directors to speak decidedly on the subject, but nothing has occurred to change the opinion they have already expressed. In their last report they gave not only their own, but the views of Dr. Kitto, Dr. Peet, Rev. Dr. Day, Mr. Weld, the Rev. Mr. Turner, Mr. E. M. Gallaudet and Miss Montgomery. They now add an extract from a letter, dated Doncaster, August 20th, 1870, to the secretary, from the able and experienced head of the Yorkshire Institution for the Deaf and Dumb, Charles Baker, Esq.:

"You are aware that the Braidwood family made this [articulation] their vehicle for the instruction of their pupils, so that my earliest predilections were in its favor; but, so early as 1815, it was stated in one of the reports of the Institution at Birmingham, that 'the children are taught to read and write, and in some instances to speak.' We, therefore, conclude that, in those early days, the teaching of articulation was the exception. When I became a resident of that institution, in 1818, there were only three cases of pupils who could articulate, and they were with difficulty understood.

---

\* Pennsylvania Institution for the Deaf and Dumb. Philadelphia, 1870.



The general instruction was carried on by signs and spelling on the fingers, with writing and other auxiliaries. There is no institution in these islands where speech is made even a principal mode of conveying instruction, and probably not one in which it is not partially cultivated. In the London Asylum, signs, pictures, the manual alphabet, and the usual means adopted elsewhere, are all employed; at the same time it puts forth articulation as its chief basis.

“At Donaldson’s Hospital, Edinburgh, professedly an institution for teaching by means of speech, all the other means are resorted to. None of the other institutions in this country profess to make speech their basis, but I have reason to think that in all of them semi-mutes and the semi-deaf have the power of speech carefully cultivated; and this is our case. I have always two or three pupils who read vocally, and whose reading is well understood by those accustomed to them, and also by strangers, after they have listened for a time, and overcome their monotony and faults of pronunciation. Nothing that I have ever seen or read on the subject would induce me to give up the substantial advantages which our pupils derive from the modes of instruction we pursue for the illusory benefits which it is supposed articulation, with reading on the lips, would confer on them. We know that a solid foundation is laid, and that constant and uniform improvement will follow under moderately favorable circumstances in after-life, and I would not renounce such solid advantages for the showy and adventitious acquirements, so liable to fall into disuse, which speech offers. In the course of my long experience, I have met with but two instances of deaf mutes having acquired speech who have persistently used it for common purposes, while I know of hundreds who have preferred the manual alphabet and writing as the *readiest* way of making themselves understood.

“Doubtless in a large institution where articulation is made a regular branch of study, a few may be found who will make the acquisition, but these will be very rare instances, and the time required must sadly interfere with the ordinary curriculum of the institution, for which the pupils thus engaged must inevitably suffer. I would suggest, in



order to complete the education of such a class, that, provided their improvement warrants it, a seventh year should be added to their course. Such an exception your directors could authorize."

The following remarks have been furnished by Mr. Ather-ton Blight, a member of the Board:

"At my visit to the Institution for the Deaf and Dumb, in London, I made particular inquiries of the principal with regard to the subject of articulation. He told me that in cases of semi-mutes, where the children had their hearing at birth, but lost it a few years afterwards, they endeavored to do all in their power by special instruction to nurse and improve the germ of capacity for speaking which each child might possess. He, in common with the large majority of teachers in Great Britain, considered it a hopeless task to attempt to teach articulation to children who were born deaf.

"The sign language, he assured me, was almost universally taught in Great Britain.

"Upon attempting to converse with one of the women teachers who was introduced to me as the best example they had of a mute who could articulate, it was with great difficulty that I could understand a single word.

"I also visited the Dutch Jewish school, for the special purpose of teaching *articulation*; together with another gentleman, I examined several of the children very carefully, and we both came to the conclusion, from the examples of one or two years' standing, that, except in the case of a bright, intelligent semi-mute, the task was utterly fruitless and hopeless."

The following is a list of thirty-three of the thirty-four institutions in the United States, with their locations, chief executive officers, and number of pupils, from the Tenth Annual Report of the Columbia Institution, June 30th, 1867, with such additions and alterations as to bring the record up to 1871:



*Institutions for the Instruction of the Deaf and Dumb in the United States.*

NAME.	LOCATION.	Date of opening.	CHIEF EXECUTIVE OFFICER.	Number of teachers.	Number of pupils during year last reported.
1 American Asylum.....	Hartford, Conn.....	1817	Edward C. Stone, M.A., principal.....	11	200
2 New York Institution.....	New York City.....	1818	Isaac L. Peet, M.A., principal.....	19	479
3 Pennsylvania Institution.....	Philadelphia, Penn.....	1821	Joshua Foster, principal.....	9	230
4 Kentucky Institution.....	Danville, Ky.....	1823	J. A. Jacobs, M.A., principal.....	4	80
5 Ohio Institution.....	Columbus, Ohio.....	1829	Gilbert O. Fay, M.A., superintendent.....	8	180
6 Virginia Institution*.....	Staunton, Va.....	1839	John C. Covell, M.A., principal.....	3	47
7 Indiana Institution.....	Indianapolis, Ind.....	1844	Thomas McIntire, M.A., superintendent.....	8	182
8 Tennessee School.....	Knoxville, Tenn.....	1845	Joseph H. Ijams, B.A., principal.....	3	61
9 North Carolina Institution*.....	Raleigh, N. C.....	1845	Willie J. Palmer, M.A., principal.....	3	41
10 Illinois Institution.....	Jacksonville, Ill.....	1846	Philip G. Gillet, M.A., principal.....	10	282
11 Georgia Institution.....	Cave Spring, Ga.....	1846	W. O. Conner, principal.....	2	35
12 South Carolina Institution*.....	Cedar Spring, S. C.....	1849	J. M. Hughston.....	2	22
13 Missouri Asylum.....	Fulton, Mo.....	1851	W. D. Kerr, M.A., superintendent.....	5	57
14 Louisiana Institution*.....	Baton Rouge, La.....	1852	J. A. McWhorter, M.A., superintendent.....	3	54
15 Wisconsin Institute.....	Delavan, Wis.....	1852	George S. Weed, M.A., principal.....	5	91
16 Michigan Asylum*.....	Flint, Mich.....	1854	Egbert L. Bangs, M.A., principal.....	6	124
17 Iowa Institution.....	Iowa City, Iowa.....	1855	Benjamin Talbot, M.A., principal.....	4	96
18 Mississippi Institution.....	Jackson, Miss.....	1855	(Buildings destroyed by fire in 1864)†	...	...
19 Texas Institution.....	Austin, Texas.....	1856	J. A. Van Nostrand, M.A., principal.....	2	27
20 Columbia Institution.....	Washington, D. C.....	1857	Edward M. Gallaudet, M.A., president.....	9	118
21 Alabama Institution.....	Talladega, Ala.....	1858	Joseph H. Johnson, principal.....	1	29
22 California Institution*.....	San Francisco, Cal.....	1860	Warring Wilkinson, M.A., principal.....	1	37
23 St. Bridget's Institution (Catholic).....	St. Louis, Mo.....	1860	Sister Stanislaus, principal.....	1	25
24 Kansas Institution.....	Olathe, Kansas.....	1862	Lewis H. Jenkins, principal.....	2	20
25 St. Mary's Asylum (Catholic).....	Buffalo, N. Y.....	1862	Sister Mary Ann, superintendent.....	4	54
26 Minnesota Institution.....	Fairbault, Minn.....	1863	Jonathan L. Noyes, M.A., superintendent.....	1	27
27 Institution for the Improved Instruction of Deaf Mutes.....	New York.....	1867	F. A. Rising, principal.....	3	26
28 Arkansas Institute.....	Little Rock, Ark.....	1867	Elmore P. Caruthers, M.A., principal.....	3	36
29 Maryland Institution.....	Frederick City, Md.....	1868	Charles W. Ely, M.A., principal.....	5	88
30 Nebraska Institute.....	Omaha, Neb.....	1869	William M. French, principal.....	2	21
31 West Virginia Institute.....	Romney, West Va.....	1870	Horace H. Hollister, M.A., principal.....	2	22
32 National Deaf-mute College†.....	Washington, D. C.....	1864	Edward M. Gallaudet, M.A., president.....	123	2005
33 Clarke Institution for Deaf Mutes¶.....	Northampton, Mass.....	1867	Miss Harriet B. Rogers, principal.....	5½	133
				5	49

\* These have departments for blind.

† The college is a distinct organization within the Columbia Institution.

‡ Resident professors.

§ First Institution in the United States which employed the articulate method of instruction.



The college referred to in the list of institutions is a distinct organization of a national character within the Columbia Institution, with a primary department, and a full college faculty, under the presidency of Edward M. Gallaudet, M.A., a son of the late distinguished Rev. Thomas H. Gallaudet. It was organized in 1864, and bids fair to further high intellectual development in deaf mutes, so as to fit them to become teachers in other institutions. Another son of this worthy gentleman is rector of St. Ann's Church, New York, a church in which the mute language is employed, and at which they can receive all the rights and privileges of our holy religion.

"The three graduates of Columbia College of last year (1869) were all quickly provided with suitable employment. Two have been appointed instructors in other institutions for the deaf and dumb; but, in addition, one of them has received a commission to prosecute researches with the microscope, in which he is a great adept, while the third has entered the patent-office, in the service of the government, and become an assistant examiner.

"It is expected that as authors, editors, scientific investigators, as well as in other walks of intellectual life, this class of men, hitherto almost wholly excluded, and ranked as pariahs, may, some of them, soon rise to the position of leaders."

It may be well to introduce at this place a notice of another effort for the deaf and dumb at Pittsburg, Pennsylvania.

"Into a mission Sunday-school connected with Dr. Brown's church there strayed sometimes a little colored deaf mute; though often dismissed, he as often returned, and this was looked on as a providential indication that some pains should be taken to have him instructed. An educated deaf mute was found, who undertook to become the boy's teacher. Then it was determined that, as there was a competent teacher, a class of deaf mutes should be collected, and, the number both of scholars and teachers increasing, a separate room was obtained for their accommodation, and ultimately a day-school was organized as a part of the common school



system of the city.”\* It opened in September, 1869, and has now (1871) thirty-two pupils taught by the sign language.

Besides the two great classes which we have enumerated in the beginning of this chapter, there is still another small division into “dumb but not deaf,” or dumb only. The majority of these “dumb only” prove, on examination, to be idiotic, paralytic, or both.

Localities seem to exert some influence in the matter of deaf mutism, as more congenital cases are from the country; but those who become dumb after birth are most common in cities and towns, thus showing the deleterious influences of cities.

Climate has a certain influence, more especially in low damp valleys, like those of Switzerland, where we find the largest number of deaf mutes compared with population. Belgium, on the contrary, contains the smallest proportion of deaf mutes, and this is accounted for by its flatness, and the great fertility of its soil. In regard to the sexes in our American institutions, the proportion is—males, 124; females, 102. In the Irish, in the investigations of Dr. Wilde, he gives the proportion of 100 males to 74.61 females in the congenital; while in the acquired, or semi-mutes, the proportion is 100 males to 91.46 females. Certain writers have asserted that the organs of speech in deaf mutes are either malformed, or from want of exercise are undeveloped; but this has been proven by recent observers not to be so. Among the chief predisposing causes of congenital deaf mutism is the close relationship or consanguinity of parents. This matter has been set at rest by the results of the Irish census, for in 154 cases, where the parents were related in the degree of first, second, or third cousins, the results of these intermarriages were 100 cases; 86 congenital and 14 acquired deaf mutes, and 4 of these were dumb only, and 4 were dumb and idiotic.

The following interesting facts are furnished by Mr. Joshua Foster,† and were obtained by an examination of the records

---

\* America and the Americans, by W. G. Blaikie, D.D., Sunday Magazine.

† Principal (Report for 1870, p. 34).



of the Pennsylvania Institution for the Deaf and Dumb at Philadelphia. The number of families in which more than one of the children is deaf and dumb appears to be one hundred and forty-seven:

Families in which there are two children deaf and dumb	. .	98
“ “ three “ “	. .	36
“ “ four “ “	. .	9
“ “ five “ “	. .	3
“ “ six “ “	. .	1
Total . . .	.	147

In five of these families both parents are mutes. In one the mother is a mute, and has had two husbands, both hearing. She had a daughter by each husband, both of whom are “deaf and dumb.”

The number of deaf mutes who have enjoyed the privileges of this institution since its commencement, fifty years ago, is (including the present inmates) nearly fourteen hundred.

“From laryngoscopic examination of twenty deaf mute children, from eight to fourteen years of age, Prinz\* arrives at the following conclusions: 1. Mobility of the muscles which open and close the larynx existed, and the power of vibration of the cords was perfect. 2. From deficient control on the part of the ear a perfect adaptation of the cords did not always follow the attempt to pronounce a vowel, but after instruction this improved. 3. In most of the cases there was a slight catarrh of the cords, referred, by Prinz, to a strain from using unnecessary force.”

Many deaf mutes have married and reared families of children; a number of the men have married hearing women, and several of the girls have been married to hearing men; but most of them have preferred partners who were, like themselves, mutes. The children of these parents in most cases hear and speak; where there are exceptions to this rule, it is believed that both parents have almost invariably been congenital mutes.

It has also been asserted that deaf mutism is an infirmity

\* Archiv f. Heilkunde, iv. p. 413. Transactions of the American Otological Society, p. 13, 1870. New York, D. Appleton & Co.



of the poor, the result of their unhealthy dwellings, bad and insufficient food, impure air, want of clothing, and other causes which elicit scrofulous manifestations. But if this were the case we would find more mutes in the cities than in the rural districts.

Almost all deaf mutes have a certain amount of hearing; and Itard makes the following division:

1st. Those that can hear the human voice as sounds, but are unable to distinguish words, amounting to about one-tenth of the whole.

2d. Those who can distinguish loud noises, such as clapping the hands, the ringing of bells, thunder, cannon firing, etc., who amount to five-tenths.

3d. Those who are completely deaf, numbering about four-tenths of the whole.

They are all, however, particularly sensitive to vibration or jar. Thus, in the Imperial Deaf and Dumb Institution at Paris, the movements of the pupils are regulated by the beat of the drum. Also in the New York Institution, containing the largest number of mutes of any establishment in the world, there has been made a successful experiment of forming an alphabet of drum-taps, by which intelligence may be conveyed in the dark, or at considerable distance, to any number who had learned the alphabet.

Cases of mutism also occur in connection with blindness. The earliest record is that in St. Matthew, xii. 22: "There was brought unto him one possessed with a devil, blind and dumb, and he healed him, insomuch as the blind and dumb both spake and saw." Several cases have been brought to the notice of the medical world, as, per example, those of James Mitchell, Julia Brace, and the well-known Laura Bridgman.

#### TREATMENT OF DEAF MUTISM.

The treatment of the deaf mute is divided into—1st. Prophylactic. 2d. Medical. 3d. At home. 4th. In school.

The first important matter is to place the mute or semi-mute in the healthiest part of the country, free from dampness or any malarious emanations, where there is access to



good and abundant food, warm clothing, healthful exercise, with cheerful company, and regulated gymnastic exercise.

2d. *Medical treatment.*—We are of the opinion that, as a general rule, congenital deaf mutism is not curable. If the child, however, has shown any signs of hearing, or has ever spoken, Wilde, and several good authorities, advise that every possible pains should be taken to keep up the articulation, whether heard by the child or not. The late Mr. Toynbee's treatment consisted in the use of trumpets, whereby the nervous apparatus may be gradually excited so as to become sensitive to ordinary sonorous undulations and external stimulants.

In cases where, by disease subsequent to birth, either the membrana tympani or the mucous membrane lining the tympanum has been thickened, counter-irritation over the mastoid process will aid the use of trumpets; and in those where the membrana tympani has been partially or wholly destroyed by ulceration, and where there is a constant discharge of mucus, pus, etc. from the surface of the tympanic mucous membrane, it is desirable to syringe the ears occasionally with a weak astringent, so as to prevent the membrane becoming ulcerated and the bone which it covers carious. The artificial drum, made of gutta-percha\* (or a piece of wet cotton), may also be resorted to.

Triquet, not depending upon his own judgment in treatment, gives the competent conclusions of M. Valade-Gabel: "The deaf mute who presents the most favorable conditions for treatment is he whose accidental deafness has supervened at the age at which the child begins to hear and speak, and who still retains some faint evidence of hearing and speech. If the organic lesion, the first cause of the infirmity, be seated beyond the nerve centres; if the child be intelligent, and have no brother or sister in the same state as himself; if he be the child of healthy parents, who have no connection by consanguinity, and if he have never previously been under treatment, the chances of cure are numerous; and if all these conditions are met with in the same

---

\* Toynbee on the Diseases of the Ear, by Hinton, p. 408. London, 1868.



subject, the chances almost reach to a certainty. On the contrary, they decrease in value in proportion as one or more of those conditions are wanting, and when all are wanting we should entertain scarcely any hope.”\*

Triquet agrees with this opinion of M. Valade-Gabel in every point, and we have already declared our own in the most positive terms. One word more. If the child has shown any signs of hearing and of speech, if deafness has come on in the first months of life under the influence of catarrhal or eruptive fever, whooping-cough, coryza, or angina, there can be no doubt that medical treatment should be resorted to at the earliest moment possible. But if the child has lost hearing while in the uterus, or after convulsions, brain fever, etc., if he has never given signs of hearing or of speech, however weak those signs might be, the chances of the success of treatment will be extremely uncertain, and we should not conceal that fact from the parents.

Triquet cites many instances of the so-called cures of deaf mutes which are to be found in the records of medical literature, but on account of certain defects of detail and their not being well authenticated it is not necessary for us to copy them. We will only refer to one case, which, coming from a gentleman of the highest authority in the profession and thoroughly qualified in this special department, is worth more than all the so-called “cured cases.” Dr. Von Tröltsch, in the fourth edition of his work on the Ear, writes: “As a matter of course, medical treatment must be introduced as soon as possible with the systematic instruction. I could relate to you from my practice several cases in which deaf mutism was obviously prevented, or was checked, or caused to retrograde, when in a condition of development. For instance, there is under my treatment at present a child four and a half years old, who, from the first months of his existence, has suffered from a profuse discharge from both ears, and is conscious only of loud sounds. Until within a few months, when I saw him

---

\* Introduction à l'Étude médicale et philosophique de la Sourdemuteté. Paris, 1859.



for the first time, he was able to produce only inarticulate barking and other sounds, which were unintelligible even to the mother, so that he was already properly considered a deaf-mute child. Under local treatment the profuse discharge from the ears soon decreased, and with the decrease of the discharge the child manifestly commenced to notice noises which were made around him, and especially the words of bystanders, and also made attempts to imitate what was said. These attempts were encouraged, and the child was employed as much as possible in speaking words and sentences. In this manner I succeeded not only in decreasing the degree of impairment of hearing, but after a few months the child possessed a tolerably distinct and at any rate quite intelligible language. Without these local applications, and the correct guiding care of those about him, the child would certainly soon have *been counted among the deaf and dumb.*" The same affection of the ear which only makes an adult hard of hearing may deprive the child at the same time of language, and cause him to remain during his whole future life in a lower state of social and mental development.

Dr. Wm. Kramer, of Berlin, in his *Aural Surgery of the Present Day*, published by the New Sydenham Society, London, 1863, p. 136, gives the following case :

"Among the diseases of the middle ear, when the membrana tympani has been whole, I have found 'exclusively interstitial exudation,' a frequent concomitant, the removal of which, in a completely (acquired) deaf mute, produced such recovery of the hearing power in the right ear that the little patient heard and repeated all letters, syllables, words, and short sentences, which were spoken with the mouth closely applied to his ear; but he was not on that account fitted to receive education with children in the perfect possession of their senses, nor to be removed from the category of deaf mutes."

This is the class of cases for which public schools should be provided to teach them articulation.

*A semi-mute improved by treatment.*—Elizabeth Ann P., aged ten years, suffered from abscesses in her ears from



the age of fifteen months, and deafness as a consequence. At the age of three years she was brought to the ear clinic of Howard Hospital. I made a careful examination, and directed a tonic course, also that she should be taught the sign language, and gave the hope that at the age of ten or twelve years she might be able to hear to a degree.

On March, 20th, 1869, she was again brought to me, when she was able to hear my watch, also her mother's voice when she placed her mouth over her ear, also heard sound of piano, etc.

*Examination.*—External meatus filled with dark secretion. Membrana tympani of both ears opaque; was not able to see the handle of the malleus. Eustachian tube of right ear filled with mucus, left ear very dry and narrowed. Throat irritable, tonsils had been removed; directed cleaning of the ear with syringe and alkaline solution, use of Politzer's air-douche, with iodine vapor and astringents to throat, the preparations of iron and cod-liver oil; also gave directions to cultivate her voice and articulation, watching her mother's mouth and the other members of her family, and to report further progress.

Several other cases have occurred in the author's clinic of young children with otitis externa and media which have not only recovered from the disease, but, by judicious after-treatment, have had their voice return and been saved from the fate of the deaf and dumb.

#### ON THE CONDITION OF THE EAR IN THE DEAF AND DUMB, AS REVEALED BY DISSECTION.

It is highly desirable that careful post-mortem examinations should be made of the ears of deaf-and-dumb patients, whose cases have been carefully noted during life, in order that the organ may be compared with the amount of hearing possessed by the patient.

Pathological investigations into the condition of the ear in the deaf and dumb have been of late years conducted so as to enable us to draw some general deductions as to the state of the ear in deaf mutes. The late Joseph Toyn-



bee, F.R.S.,\* gave the details of five dissections performed by himself, and followed them with a tabular view of the morbid appearances found in all the other dissections of similar cases that had been met with by him.†

"*Dissection 1.*—A woman, aged forty, deaf, and consequently dumb, from birth. For the last ten years of her life she was insane, and was confined in a lunatic asylum. The petrous bones were sent to Mr. Toynbee, immediately after the death of the patient, by the late Mr. Crosse, of Norwich.

"Right ear.—The meatus externus, membrana tympani, and tympanic cavity were in a healthy state. Labyrinth.—The anterior limb of the superior semicircular canal communicated with the vestibule, but was rather larger than natural, and its shape somewhat irregular. Examined as it passes inward, it was observed to terminate in a cul-de-sac, after having attained only half its usual length. The portion of bone in the situation usually occupied by the inner part of the canal was of an ivory whiteness, and could be distinctly distinguished from the surrounding bone. The greater part of the posterior semicircular canal was absent, its anterior and posterior extremities presenting two cul-de-sacs, half a line in length. The external semicircular canal, the vestibule, and cochlea were in a normal state.

"Left ear.—The superior semicircular canal is in the same incomplete state as that of the right ear.

"*Dissection 2.*—J. C., aged fifty, died from fever. He was born deaf, and had been educated as a deaf mute.

"Right ear.—Meatus externus in a natural state, excepting that one part of the osseous parietes was quite rough. Membrana tympani thicker than natural, and perforated, a small red polypus was attached to its dermoid layer. A membranous band connected the stapes, incus, and tensor tympani muscle. The osseous semicircular canals were large, but they did not contain any membranous tubes. The cochlea

---

\* The Diseases of the Ear, by J. Toynbee, edited by Hinton. London, 1859.

† This table has been added to by Prof. S. Moos, M.D., of Heidelberg, who has collected twenty-nine additional cases.



was in a normal state, as were also the tympanic muscles and nerves.

"Left ear.—The osseous semicircular canals did not contain any membranous tubes. One of these osseous canals was so contracted as to admit of the passage through it of a fine wire only.

"*Dissection 3.*—R. B., aged sixteen, a girl, bright and intelligent. Her parents stated that she was born deaf; but the child, on the contrary, said that she had heard sounds. Her teacher thought the child's statement the more likely to be correct, as she pronounced many words according to their proper sounds. None of her relatives were deaf mutes. Upon making a careful dissection of each ear, no deviation from the normal condition could be detected in the meatus externus, membrani tympani, or tympanic cavity. The stapes adhered to the fenestra ovalis with its usual degree of firmness. The membranous labyrinth in the vestibule was healthy, as was also that in the semicircular canals, with this exception, that in the middle of the superior semicircular canal in the right ear was a quantity of otoconie (ear-crystals), which completely obstructed the tube.

"From the occurrence of an accident to the left ear, it was not possible to ascertain its condition.

"*Dissection 4.*—A boy, aged fifteen, who had been wholly deaf from his birth. No deviation from the normal state in either ear could be detected.

"*Dissection 5.*—For the opportunity of making this dissection, Mr. Toynbee was indebted to Dr. Ormerod, of Brighton. The ears were taken from a young woman deaf and dumb, with discharge from each ear. She died with tubercular inflammation of the brain.

"Right ear.—The membrana tympani was absent; the mucous membrane of the tympanum red, and of extreme thickness, so as to fill the whole of the tympanic cavity, and to conceal the stapes; the tympanum was distended with dark-colored blood. The petrous bone was so intensely hard that it could with difficulty be cut by the bone-forceps. The cochlea, to the naked eye, had a natural appearance, and when submitted to microscopic examination, no deviation from the



normal structure could be detected, excepting in that portion of the lamina spiralis which is near to the vestibule. This, instead of being composed of a delicate osseous lamina, and a fine membrane, the two making a delicate septum between the scala tympani and scala vestibuli, presented a mass of solid bone, filling up the scala tympani, and concealing from view the membrana fenestra rotunda, the inner surface of which it wholly covered. The outer surface of this membrane could, however, be distinctly seen from the tympanic cavity, and it appeared natural. The semicircular canals contained more otoconie than was natural.

“Left ear.—The membrana tympani was absent. The mucous membrane of the tympanum thick and red, as in the right ear. The semicircular canals also contained more otoconie than natural, especially the posterior one, which, at its junction with the posterior limb of the superior canal, was wholly distended with crystals to the extent of half a line. The lamina spiralis of the cochlea was of a deep red color, and blood was effused in both scala. The part of the lamina spiralis near to the vestibule was in the same state as the same portion in the right ear.”

In the subjoined tables the condition of the ears in deaf mutes, as revealed by various reported dissections, is shown. (See Tables.)



*A Tabular View of the Condition of the Ear in Sixty-five Dissections taken of Deaf Mutes.\**

MEATUS EXTERNUS.	MEMBRANA TYMPANI.	TYMPANUM.	LABYRINTH.	NERVE.	NAME OF OBSERVER.
Absent.	.....	.....	.....	Soft as mucus.	Itard.
do.	.....	.....	.....	.....	Fabricius.
do.	.....	.....	.....	.....	do.
.....	.....	.....	.....	.....	Itard.
.....	Destroyed.	Containing vegetations from the mucous membrane; ossicles absent.	.....	.....	do.
.....	do.	do.	.....	.....	do.
.....	Partially destroyed.	Containing scrofulous matter.	.....	.....	Cock.
.....	do.	.....	.....	.....	do.
.....	do.	.....	.....	.....	do.
.....	.....	Containing calcareous concretions.	.....	.....	Itard.
.....	.....	Full of gelatinous matter.	Full of gelatinous matter.	.....	do.
.....	.....	Containing a yellow fluid.	.....	Harder than natural.	Rosenthal.
.....	.....	Anchylous of stapes to fenestra ovalis.	.....	.....	Valsalva.
.....	.....	All the ossicles wanting.	.....	.....	Reimarus.
.....	.....	.....	Vestibule full of caseous matter.	.....	Haighton.
.....	.....	.....	Cochlea consisting of only one turn and a half.	.....	Mundini.
.....	.....	.....	Vestibule, cochlea, and semicircular canals absent.	.....	Meckel.
.....	.....	.....	Semicircular canals wanting.	.....	Murer.
.....	Partially destroyed.	All the ossicles wanting.	Two of the semicircular canals imperfect.	.....	Cock.
.....	.....	.....	do.	.....	do.
.....	.....	.....	.....	Atrophied.	Sylvius.
.....	.....	.....	.....	Indurated.	Ackermann.
.....	.....	.....	.....	Wanting.	Morgagni.
.....	.....	.....	Fenestra rotunda closed by bone in each ear.	.....	Cock.
.....	.....	.....	One semicircular canal incomplete in one ear only; the other ear healthy.	.....	Thurnam.
.....	.....	.....	Aquæductus vestibuli very large.	.....	Dalrymple.
.....	.....	.....	.....	Very hard.	Rosenthal.
.....	.....	.....	Semicircular canals absent in one ear.	.....	Murer.
.....	.....	.....	Filled with caseous matter.	Half its usual size.	Haighton.
.....	.....	.....	.....	Atrophied.	Hoffmann.
.....	.....	.....	.....	Compressed by a tumor.	Duverney.
.....	.....	.....	A portion of one of the membranous semicircular canals filled with otoconie.	.....	Toynbee.
.....	.....	.....	The superior and posterior semicircular canals incomplete in right ear; the superior incomplete in the left ear.	.....	do.
.....	.....	.....	Membranous semicircular canals absent.	.....	do.
.....	.....	.....	.....	.....	do.
.....	Destroyed.	Mucous membrane thick.	Lamina spiralis near the vestibule filled the scala tympani.	Otoconie obstructing the canals.	do.

\* Where no entry is made, it is to be assumed that that part of the organ was in a healthy state.



To the tabular arrangement, made by Toynbee, we will add, from Moos,\* the following, omitting the observations made upon monsters, such as hemi- and anencephali:

EXTERNAL EAR.	MEMBRANE OF THE TYMPANUM.	CAVITY OF THE TYMPANUM.	LABYRINTH.	NERVE.	NAME OF OBSERVER.
.....	.....	The walls rough and uneven, fenestra rotunda and promontory wanting.	.....	.....	Schallgruber.
.....	.....	Fenestra rotunda closed by bony material.	.....	.....	Ribes.
.....	.....	Fenestra ovalis too narrow.	.....	.....	Römer, Otto.
.....	.....	Stirrup wanting.	.....	.....	Deleau, Jr.
.....	.....	The cavity of the tympanum filled with a mucous mass; no trace of any ossicles.	Vestibulum, cochlea, semicircular canals, and both fenestra wanting.	.....	Montain, Saissy.
.....	.....	.....	The semicircular canals of both ears terminating blindly, not communicating with the vestibule.	.....	Bochdalek.
.....	.....	.....	The interior half of the superior semicircular canal wanting; the posterior canal wanting completely.	.....	Schallgruber.
.....	.....	.....	The semicircular canals preternaturally filled with bony substance.	.....	Hg.
.....	.....	.....	Semicircular canals thickened, hard as ivory, and developed at the expense of these cavities.	.....	Bochdalek.
.....	.....	Stirrup wanting.	Fenestra rotunda wanting.	.....	Clarus.
.....	Thickened.	Malleus and incus of both sides atrophied; atrophy of the promontory on the right side, a bony process; from the plate of the stapes to both sides; anchylosis of the stapes; the eminentia pyramidalis wanting. On the left side the fenestra ovalis closed completely, by the atrophy of the stirrup, the arms of of which were no thicker than a hair.	The terminations of the new vestibuli and cochlea dissolved into a brownish pulp. On the left side the horizontal semicircular canal was wanting.	Neurilemma on both sides thickened; the soft oleaginous substance wanting.	Hyrtl.
.....	On the right side, smaller than normally, but three times the thickness.	On the right side, the base of the stapes no larger than the os Sylvii; fenestra ovalis contracted.	On the right side, only one turn and a half to the cochlea, the others running together into a common cupola, into which the atrophied modiolus projected; the spiral lamina entirely wanting; the semicircular canals small and narrow; the osseous substance around the labyrinth very hard and compact.	.....	do.
.....	The same on the left side.	On the left side, the stirrup wanting, the fenestra ovalis closed by an osseous plate attached to its whole circumference.	On the left side, there was the same appearance as on the right.	.....	
.....	.....	On both sides, the external wall wanting, the entire cavity wasted; the ossicles wanting, intimated only by a bony stem.	The semicircular canals and vestibule wasted; the fenestra rotunda closed by coalition; cochlea making only one turn and a half, otherwise normal.	Thinner than normal.	do.

\* "Klinik der Ohrenkrankheiten," pp. 341-348. Wien, 1866.



EXTERNAL EAR.	MEMBRANE OF THE TYMPANUM.	CAVITY OF THE TYMPANUM.	LABYRINTH.	NERVE.	NAME OF OBSERVER.
.....	.....	Anchylous of the malleus and stapes into one bone; the stapes attached to the wall of the cavity of the tympanum; complete immobility of the ossicles; fenestra rotunda wanting.	.....	.....	Gellé.
.....	.....	.....	.....	Thickening of the ependyma in the fourth ventricle; striæ auditoreæ wasted, probably in consequence of meningitis interna.	Meyer.
.....	.....	Grown together with the promontory; the plate of the stirrup immovable; chorda wanting on right side.	.....	.....	Triquet.
.....	.....	The whole auditory apparatus without any anomaly.	.....	.....	do.
.....	.....	Only one branch to the stirrup on the left side.	Fenestra rotunda of minimum size on the left side; arrested development of the semicircular canals on both sides, and of the cochlea on one side.	.....	do.
.....	Very horizontal.	Somewhat narrowed, the promontory not distinctly developed; fenestra rotunda wanting; the branches of the stapes very small, and its plate hardened and very narrow.	Horizontal semicircular canal wanting; vestibule narrowed; modiolus and spiral lamina wanting.	Internal auditory meatus narrowed, the nerve itself wanting.	Nuhn.
.....	.....	Stapes and os Sylvii wanting; the papillary process entirely wanting.	The whole internal ear completely wanting on both sides.	Nerve completely wanting on both sides; no trace of any opening for its entrance.	Michel.
Meatus externus narrower than normal.	.....	On the right side, stirrup immovable, fenestra rotunda wanting. On the left side, stapes adherent, but less firmly than on the right. Fenestra rotunda rudimentary in form and extremely narrow.	The scala tympani of the well-formed cochlea opens into the vestibule, instead of into the tympanum, through the fenestra rotunda.	.....	Dardel, Schweitzer.
.....	.....	.....	Degeneration by chalky deposition.	.....	Voltolini.
.....	.....	.....	Obscuration, thickening, and calcination of the membrane lining the labyrinth.	.....	Virchow.
.....	.....	On the right side, foramen rotundum overlaid with a pseudo-membrane; stapes in the same condition, and immovable. On the left side, the same appearance, but more marked, showing greater vascular injection.	On the left side, no trace of any otoliths; the small sacks, of an obscure, yellow appearance, with a great number of capillaries. On the right side, many otoliths in the sack, and in the left horizontal canal.	The ependyma of the fourth ventricle of the brain much thickened; striæ not discernible; auditory nerves partly atrophied.	Voltolini.
.....	.....	.....	The membranous canals atrophied, containing a great many otoliths; these were wanting in the thickened sacks.	.....	do.



EXTERNAL EAR.	MEMBRANE OF THE TYMPANUM.	CAVITY OF THE TYMPANUM.	LABYRINTH.	NERVE.	NAME OF OBSERVER.
.....	.....	Horizontal position of both the malleus and incus; anchylosis of the ossicles.	.....	Atrophy of the auditory nerve on both sides.	Voltolini.
.....	.....	.....	.....	Atrophy of the auditory nerves.	do.
.....	Completely destroyed.	All the ossicles present; stapes immovable; osseous tuberculous.	On both sides, superior and posterior canals partly closed by bony material; aditus ad cochleam narrowed; the membranous labyrinth thickened by pseudo-membrane, and unrecognizable; cochlea thickened, and of a gray color, from the presence of pigment.	Atrophy of the auditory nerve on both sides.	do.
.....	.....	.....	.....	do.	do.



## DIRECTIONS FOR TEACHING DEAF MUTES AT HOME.

The following directions, drawn up by one of our most valued teachers, now deceased,\* are of the utmost importance, and should be begun with the child as early as possible.

It is very important to the deaf mute that his parents and friends should cultivate the language of signs, and encourage him in the use of them as early as possible. Let them observe the child, and imitate the signs he makes. When he is pleased with anything, invent a sign for the thing, and repeat that sign many times afterwards. Distinguish different persons by signs, suggested by a scar, mole, beard, or any little peculiarity which the person may possess. Imitate the actions of riding, sewing, eating, mowing, cutting, throwing, sowing, etc. For "good," kiss the hand. For "bad," bring the hand to the lips, turn the palm down and throw it from you. For "glad," pat the heart rapidly, with a cheerful expression of countenance. For "sorry," rub the clinched hand on the heart, with a sad expression of countenance. For "black," draw the end of the forefinger along the eyebrow. For "red," touch the lips with the forefinger. For "love," cross the hands and press them on the heart. For "hate," push both hands, the palms out, from the heart, as if repelling something from the left side. For "lie," move the forefinger across the mouth horizontally. For "true," place the forefinger perpendicularly across the lips and thrust it forward.

These are very few examples, merely as suggestive hints. The child can be taught to spell on the fingers at a very early age by the use of the deaf-and-dumb alphabet. (See p. 421.)

Any person can take an object, as a hat, pick out the letters h-a-t from the deaf-and-dumb alphabet, and learn to place the fingers in the true position for each letter, no matter how slowly it is done. Let the child imitate until he can make the letters of the word in order, without assistance; at the same time show the object. Do this very often, until

---

\* The late A. B. Hutton, M.A., Principal of the Pennsylvania Institution for the Instruction of the Deaf and Dumb.



# DEAF AND DUMB ALPHABET.

a



b



c



d



e



f



g



h



i



j



k




l



m



n



o



p



q



r



s



t



u



v




w




x



y



z



&





the child has learned to spell the word when the hat is presented to it, or to go and bring the hat when the word is spelled to it. Then take another object, as "pin," go through the same process, until it is thoroughly learned by frequent repetition every day. So with *arm, cat, dog, chair*, etc. The following ten short words, the names of common objects, contain every letter of the alphabet, viz., adze, fan, map, cow, box, jar, sky, hat, quill, glove. After the power of spelling the names of many common objects has been acquired, the names of familiar persons may be taught. In a similar way, the child may be taught to write the names of things on a slate quite early. Let him imitate the form of the letters for one word, as "hat," and repeat it many times, until he can write it as readily as he can spell it on the fingers. Take another word and go over the same process. Point to each letter, and require to mark the sign for the letter on the fingers. By frequent repetition, the ability to write the names of many things and to form all the letters of the alphabet will be acquired. It is best to make the child form the letters as round as possibly, and not to take off the pencil until the word is completed. The habit thus early begun will save much time, and enable the writer to accomplish more in a given time, and with more ease, than can possibly be done on any other principle. If it is desired to go further, write the name of the child, as "*John* sees a chair," "*John* sees a table." Let him copy the sentence, explaining by signs the word *sees*, and pointing to the chair or table, and also to the child. Then let him write "*John* sees," and let him select another object to fill up the blank; and finally, let him cover his slate with sentences thus formed. Help him to objects out of the house as well as in. Encourage him to write as many such sentences as he can. All this may be the work of years, but the advantage to the child cannot be estimated. A little attention thus bestowed every day, will accomplish all this and probably much more.

#### INSTRUCTION OF THE DEAF MUTES IN SCHOOL.

It is by signs that the process of teaching the deaf mutes is chiefly conducted. When we look at the Chinese charac-



ters on a tea-box, we can see no meaning in them, and might so look forever without becoming any wiser. So, also, with the mute. Our written or printed words are as inexplicable to him as the Chinese characters are to us, and inspection alone could never afford any clue to their meaning. An interpreter, or a book, would speedily convey to us the meaning of the characters, through the medium of our language, with which we have been familiar from early infancy. But the deaf mute has no language. To enable him, therefore, to learn the meaning of our words, he must acquire a language through which he can get that meaning. Every mute of tolerable capacity makes use of motions to indicate assent or denial, approbation or repugnance, as well as some common objects and familiar actions. On these motions, limited and imperfect as they are, we graft by degrees a system of signs, which enables us finally to communicate considerable knowledge on many subjects, and to develop and call into exercise the faculties of the mind. These signs convey thought, and have no resemblance to words, but they enable us to define words, explain their relations to other words, give their arrangement in sentences, and the different meanings which are attached to them. This language of signs can only be acquired from the living teacher. Incomprehensible as it may seem to a speaking person, unacquainted with the subject, that thought, however abstruse or refined, may be conveyed by various motions of the arms, it is nevertheless true, and a system of these motions is the grand means of instructing the deaf and dumb. This being premised, a class of ten or twenty mutes is furnished with large slates, on which to write with chalk, crayon, or pencil. The instructor presents an object, or a picture of one, or makes a sign for it. He then teaches them to write the name, presenting each letter by the manual alphabet. When they can all write it, it is erased, and rewritten a number of times, till it is impressed upon the memory. Some information may be communicated respecting the object. Questions may be asked to induce the pupil to think. In this way a number of nouns are taught, so that when a concise sign is made for one of them, it will be readily written.



In the same way words expressive of the qualities and properties of bodies may be taught. When such words are presented with appropriate nouns, the pupils write them in connection. They are then required to give examples of similar combinations from their own resources. This is the first attempt at composition. Another step will be to make signs for actions, and teaching their names. Then the use of these words, in combination with the words already familiar, as "a boy sees a horse," "a boy sees a strong horse."

Again, some of the words expressive of the relations of objects may be taught; as, "a lady sits on a chair," "a bird flies into a cage."

Other words and other ideas are presented to them. They endeavor to express the ideas in writing, using the words and forms of arrangement which had been taught. These sentences are corrected, and the pupils are required to give examples of their own. These original efforts are also corrected. The connections of language, the abstract terms, the phrases and the idioms are successively taught. Series of sentences, anecdotes, narratives, etc. are written off and explained by signs. These are copied by the pupils and studied as evening lessons, and in school are written from memory, or recited by signs.

There are other evening exercises, such as writing a number of original sentences, or single words, composition on particular subjects, letters, etc. From time to time the elementary principles of arithmetic and geography are taught. Indeed, our illustrations of words and principles are drawn from the sciences and the whole range of human knowledge, so that in the course of their education a great amount of knowledge is communicated to them. The subjects of arithmetic, geography, grammar, history, etc. cannot be taught systematically until the latter part of their course, when they are supposed to have acquired a considerable command of written language.

Moral and religious subjects have also a large share of attention. Much useful information is communicated by lectures addressed in the language of signs to all when assembled together.



It will readily be inferred from these statements that much will depend upon the capacity of the pupil, his attention, and his diligence. There can be no set course or limited periods for certain studies, which, when completed, make an educated person.

The longer the mute is under instruction, the greater will be his command of language. It will also be perceived that much depends upon the knowledge, ingenuity, and tact of the teachers in the use of signs. The language of signs is the all-important instrument by which the educator is to reach the mind of the mute pupil in his early and his later efforts. By this alone can he lead the pupil to reflect on his own mental operations, feelings, motives, emotions, and passions, and thus learn the thoughts, feelings, etc. of others, and to understand and use the language employed to express ideas on these subjects. When this point is reached, the pupil may relinquish entirely and forever, if he please, the use of signs. A new instrument has been given him by which he may explore the world of books, and communicate with his fellow-men to an unlimited extent.

From this point, self-education may be carried on and continued to the end of life through written language. It should be remarked, however, that a large number of mutes do not reach this point from want of capacity; yet the acquisitions, even of such, are probably as valuable in proportion as those made by the more gifted. The proboscis of the fly is doubtless as important to the little insect as the trunk of the elephant is to that sagacious and majestic animal.

The deaf mute, when the study of language is begun in our institutions, has passed the favorable age for getting words and phrases.\* They delay the commencement of the work till the age of ten or twelve, that they may work upon the higher faculties, but only in accordance with nature. This makes the task more difficult, but the results are more satisfactory.

---

\* Deaf and dumb children are not received where the sign language is taught, until the age of ten years; while in the schools where the articulate method is employed, the pupil is only required to be five years old—that is, before the articulating muscles have become rigid.



There is a marked difference in the docility of mutes, arising partly from their natural temperament and disposition, but greatly from their training, or rather from their want of training. Parents should treat their mute children with as much impartiality and restraint as their other children.

#### ON THE CHOICE OF METHODS OF TEACHING THE DEAF MUTE.

It is perfectly natural for the entirely deaf to use pantomime in expressing their wants, and the teaching of such persons in classes was a great boon conferred upon them; yet, after having acquired all that was taught by the natural method, they were still unfitted to mingle in general society or to make their way in the world. By the artificial method introduced by Heinicke, the bright mute can learn to talk, and can receive intelligence, by watching the motions of the lips of others. In very few instances only are the vocal organs defective. In a large number of cases examined by Dr. Morell Mackenzie, of London, he has only met with four instances in which they were not fitted for speech. This has been confirmed by Prof. Louis Elsberg, of New York, and by Dr. J. Solis Cohen, of Philadelphia, gentlemen well qualified to give an opinion, having examined such cases.

The whole difficulty is in the ear, as it conveys "no sound which the child can imitate, and thus produce coherent speech;" but, as I have stated before, there are certain visible signs which the lips and countenance portray whenever we utter a word; thus the deaf mute may be said to see the words. They could not thus be seen unless they were formed according to certain regular well-defined positions and motions of the visible organs of speech. These certain of the mutes can be taught to imitate, and, consequently, to speak. This is no fanciful theory, but a statement of facts which came under my own observation during a recent visit, to witness the articulate method of instruction, at No. 330 East Fourteenth Street, New York,\* now (1871) located on Broad-

---

\* Institution for the Improved Instruction of Deaf Mutes.



way between Forty-fourth and Forty-fifth Streets. I was accompanied by Prof. Elsberg, and we were at once ushered into the school-room. The pupils, as a class, had to me a very intelligent look. The first pupil brought forward was a little girl of nine years of age. The principal caught her eye, and spoke to her in a whisper, without any special contortions of the mouth, and without hesitation the child replied, by writing upon the blackboard, "This gentleman is from Philadelphia, and he is a doctor." He then called upon a boy, who was a congenital mute, who also repeated his words with more effort. I was then requested to test their proficiency, by asking them a question in arithmetic; which I did; and, although I wore a moustache covering my lips, two of them at once comprehended my question, and answered it promptly. Other questions were answered by them, and they could read the lips of the principal (Prof. F. A. Rising) the whole distance of the school-room, some twenty-five or thirty feet. There were also children who were day-scholars; and one very intelligent little fellow delighted us and his parent very much by being able to say "father and mother." He also called out "Home," when it was time to go. There was great diversity in the sounds which they uttered, some being more or less disagreeable; but none of them as harsh as an animal, to which it has been compared. The consonants were the most difficult to pronounce; but the principal showed great ingenuity in his efforts to make them understand them by the vibration on the head and on the neck. He gives the following explanation of the *method*:

"*Explanation of the articulate method.*—At the risk, on the one hand, of being unnecessarily explicit, and on the other, of not clearly and concisely illustrating the mechanical formation of letters, the following attempt to explain some of the sounds of the alphabet is made. It is hoped that the parents of little ones, who are deaf, will thus acquire some hints which will enable them to teach their children to utter, not alone the sounds here described, but all the sounds of the different letters and words easy of pronunciation.

"The foundation of the system rests upon the fact that articulation is simply a *mechanical process*, the result of certain



well-defined positions and movements of the vocal organs, together with a more or less forcible expulsion of air from the lungs. Of the former, his sense of vision renders the mute cognizant; of the latter, his sense of feeling.

"All children are imitative,—the mute child is particularly so,—and it is upon the possession of this faculty, *imitation*, that our success in teaching him depends. We proceed, then, at the outset, by making simple gestures with the arms, the easier gymnastics in short; then we walk, look in various directions, sit, stand, etc., all which operations he readily performs.

"Having thus engaged his attention, place him directly in front of you, press your upper teeth in a marked manner upon the lower lip, hold a bit of paper in your hand, and blow; he sees the paper fly away, is amused, imitates the process, and has given his first letter, *f*. Again, swell your lips out with air, open them quickly, the paper flutters away; this is *p*. Should he be too forcible, or not sufficiently so, in forming this letter, then imitate a person smoking a pipe; the *puff*, *puff*, conveys the idea of *p* exactly.

"*T*. Place the tongue between the teeth, force the air out so that it shall come in one volume,—best felt on the back of the pupil's hand,—and *t* is produced.

"*H* is simply the expiration of breath with slightly increased force, the mouth opened naturally.

"*M*. With closed lips the force of the letter *m* is felt by the pupil, who must apply his finger to the side of his nose; the vibration indicates the rapid expulsion of air from the lungs.

"*N*. In the same manner the mute learns the nasal *n*, but is shown the mouth opened slightly and the tongue pressing against the back of the upper teeth.

"These six consonants are the ones perhaps most easily learned.

"*B* is formed like *p*, but is accompanied with sound, the throat is both seen and felt to dilate, and the expulsion of breath is also less forcible, the puff is gentler.

"*D* resembles *t*, although it is better in the beginning to teach the pupil to place the point of the tongue back of the



upper teeth; the vocalization is made apparent as in the case of *b*, also the less decided emission of breath.

"*V* is simply a vocalized *f*, and as in the case of *z*, and the sub-vocal *th*, the vibration is most sensibly perceived by pressing the palm of the pupil's hand upon the crown of your head.

"*O*. The vowel *o* is attended with no difficulty. Place the child's hand firmly on your chest that he may feel the vibration, while he sees your mouth assume the shape of the letter produced.

"*E*. Place the pupil's fist under the angle of the jaw, the peculiar and forcible vibration there felt he readily produces.

"*A*. Press the palm of the pupil's hand against the chest, and show him your tongue firmly set against the lower teeth, and somewhat arched; with this position the letter *a*, as in fate, is produced. (The German method is to teach *a* as in far, first, which is accomplished by holding the tongue down with the finger, or better, by a paper divider; this is, however, entirely unnecessary in teaching the English language.)

"*I*. To form *i*, open the mouth very wide, and, as you produce sound, close it quickly.

"*U* is made like *o*, except that the lips are nearly closed.

"As soon as the pupil has learned the sounds of the vowels, require him to join a consonant with each in succession, making easy syllables, as *fa*, *fe*, *fi*, *fo*, *fu*, and reversing the letters, *af*, *ef*, etc. Next teach him simple words, the names of objects, which he can see, or pictures of them which can be shown him, as *map*, *mat*, *cat*, *top*, etc. The next step is to embody these in the form of a sentence, as 'What is that?' 'That is a top.' 'What is that?' 'That is a cat.' 'What are those?' 'Those are maps.' The point cannot be too strongly insisted upon that the child should write as well as speak everything he learns. This constant practice strengthens the memory, and insures steady and permanent progress."

I am satisfied that this system must improve and develop their lungs and chest. This system is not a recent experiment, except in a few institutions, but has been in active



operation in Germany from 1760 to the present time, and I trust ere long will be introduced throughout this country. This method has lately been introduced, in part, into the Pennsylvania Deaf and Dumb Institute at Philadelphia, and has been in operation at Northampton, Mass., and in New York, since 1867. Last winter (1870) this latter was placed upon the same footing as the old New York Asylum by the State Legislature, which is highly creditable to its intelligence, and to its care for the indigent mute, who can now be taught what was only to be obtained by the rich at great expense, time, and trouble.

Let me, in this connection, give a few more extracts from the report of the principal of the Columbia Institution, who recently made a visit to Europe to examine into the whole subject; and who is not in favor of the artificial method, yet he endeavors to give a fair statement of what he saw of this mode of instruction:

“The institution at St. Hippolyte du Fort, France. Here the manual alphabet is unknown by the pupils; articulation is taught to all. The director is M. Martin, whose wife is also a teacher. They admitted that schools conducted on the natural method had done a good work, and were even willing to admit that as a means of affording instruction this method is superior to the artificial. They held to the view, however, that the power of communicating freely in speech with their fellow-men, which was acquired by mutes under the latter system, was so great a boon as to justify a lower standard in the intellectual training of the deaf and dumb. Prof. Gallaudet states that he conversed orally with the pupils, and with the best articulators had little difficulty in making himself understood. The utterances of some were unintelligible, but those of others very distinct; and he remembers that one in particular, who was said to have been born deaf, spoke well, and read from his lips with ease.

“The Jewish institution at Vienna, Austria, is sustained entirely by private contributions within their own circle, and open only to their children. This school is conducted on the most rigid principles of the artificial method. Articulation is undertaken with all the pupils, and on his visit to the insti-



tution it was demonstrated to his satisfaction that oral conversation on familiar subjects is maintained with ease between the teachers and a large proportion of the pupils.

“The following extract from a sketch of the institution, prepared by M. Deutsch, the director, will show what is attempted in the course of instruction :

““In this institution the manual alphabet or finger language, artificial mimicry,\* and conversation by pantomime have been entirely excluded. The natural gesture is only used as a starting-point, as the first medium of understanding, which in the instruction only serves to represent real ideas obtained by personal observation through actual perception, but not to produce ideas or notions by means of signs. The method prevailing in this institution tends to make oral language and written language independent of the pantomime,—that is to say, the direct channel and medium of thought. The written language is assumed as the basis of instruction. It begins with a single word which conveys a subject, action, or quality, and leads the pupil to a connected idea. The *copia verborum* obtained in this manner is used through a regular course of exercises to form a simple sentence, and continued up to the explanation of the abstract idea. The understanding of the abstract idea is obtained in a natural way, by plain and combined sentences as well as through short descriptions and narratives. This is very different from the true elementary method made use of with other intelligent children. The instruction of religion commences only then, when the pupil has advanced so far as to understand the plain expressions of thought. Thus his religious education undergoes no especial difficulty, as the reading of the Holy Scriptures is already begun. Upon oral language a decided value is placed, and therein important results are obtained.

““The pupil not born deaf enjoys the particular advantage of acquiring the verbal language with greater facility. Read-

---

\* This is understood to refer to the exercises of the school-room. Among the pupils, during their hours of relaxation, the sign language and manual alphabet are freely used in conversation.



ing the lips has obtained such a degree of perfection that the director is able to deliver his Sabbath sermons with exhortation orally, in which the pupils of the higher class can take part.'

"The examination I was allowed to make of the attainments of the pupils in this institution was altogether unrestrained. I was requested to indicate what pupils should be examined, and, in two cases, I purposely selected those whose personal appearance led me to suppose they were rather below than above the average of intelligence. At my suggestion, the director dictated to a pupil a short account of the coronation of the Emperor of Austria as King of Hungary, which had occurred at Pesth a few days before. M. Deutsch held his hands behind his back and spoke in a whisper, without any special contortions of the mouth and without repetitions, the boy writing with great rapidity, sometimes finishing a word while his eye was already on the mouth of the director to catch what was to follow. My companion in the visit was a German professor under whose tuition I was studying in Vienna. He assured me that what was written on the black-board was an exact transcript of what M. Deutsch had spoken, absolutely without mistake. I think I do not exaggerate when I say that this exercise was performed as rapidly as would have been possible had the boy been in the possession of his hearing, and could not have been more speedily accomplished had the communication between the teacher and pupil been by means of the sign language as used in our best American institutions.

"I happened to find in this institution a boy of ten years, from Baltimore, with whom I had been previously acquainted. He is one of three mutes in the same family, born totally deaf, and with his sister has been in the school of M. Deutsch about three years. This little child comprehended with considerable readiness simple words which I addressed to him in German, repeating them after me in a clear but unmusical voice, and wrote with quickness and precision the following, which was dictated to him at my suggestion by the director in German: 'This gentleman is from America, and when he returns to America he will see your dear parents and tell them



that he met you, and that you were well and improving in your studies.' With regard to this boy, I should venture the opinion that he read from the lips (so far as his range of study had extended) with as great readiness as any pupil of his age and standing in the institution. The President of the Northampton school visited, in 1870, M. Deutsch, who for twenty-five years has had this school under his charge, and found him with all the enthusiasm of his youth, and aided by a corps of excellent teachers, with a class of about one hundred pupils.

#### THE INSTITUTION AT ROTTERDAM, HOLLAND.

"An address on the subject of deaf-mute instruction, delivered before the 'ninth congress (scientific) of the Netherlands,' in Ghent, last August, by Mr. Hirsch, the director of the Rotterdam school, so clearly defines his position as a radical supporter of the artificial method that I\* will quote a few paragraphs from it before proceeding to describe my visit to him and his establishment:

"The first and principal fact that has been made patent to society is the possibility of developing intellectually, morally, and religiously the deaf and dumb. As to the means by the aid of which instruction can and ought to be imparted to them, opinions are very diverse, often very contradictory. Those diversities and contradictions of opinion have given rise to differences in methods of instruction, and to dissensions between the schools of France and of Germany.

"The object to be attained is to render possible the admission of the deaf mute into society by teaching him to see—that is, to understand—the movements of the lips and to speak in his turn.

"To attain this end the act of seeing or comprehending and of speaking must be the exclusive principle of instruction, and neither the palpable alphabet nor the language of signs can have any connection with it.

"It is true that the language of natural signs is the first means employed by the teacher to enter into relations with

---

\* Report of Edward M. Gallaudet, M.A.



the pupil, but he does not make use of this method for any length of time, and it is abandoned as soon as it can be superseded by speech.

“The daily observations which I have made for more than thirty years that I have devoted to the deaf and dumb, have convinced me that *the art of seeing speech in the movements of the mouth is the most important* of all the branches of instruction, and that therefore it should be most sedulously cultivated.

“Next to the art of seeing or understanding, the act of speaking is the principal object of the instruction of the deaf and dumb. By this system ninety-nine out of every hundred deaf mutes may be taught, and their progress will depend entirely on the talent and patience of the teacher; this truth, too long and often too coldly doubted, is now penetrating everywhere.

“This school was one of the few where I was unfortunate in calling at the season of vacation. I was not, therefore, able to satisfy myself by personal examination as to what extent the attainments of his pupils *en masse* would confirm the remarkable claims he makes in the above paragraphs.

“I had, however, an opportunity of examining an individual case in a manner quite novel, and which put the oral and visual abilities of the pupil to what I conceive to be a very severe test.

“Just as I was leaving Mr. Hirsch, after having held a long conversation with him, in which he urged with much earnestness, and even eloquence, the advantages of his system, a young man about twenty-five years of age entered, who was introduced to me as Mr. Edward Polano, the son of a physician, and who with his sister constituted the first class taught by Mr. Hirsch in Rotterdam. I was told that these persons were born totally deaf, and that they have never at any time gained the slightest power of hearing.

“Mr. Hirsch in introducing Polano to me used the German language, and in telling him who I was used the Dutch.

“As I shook hands with the young man I said, looking him full in the face, ‘Sprechen Sie Deutsch?’ His answer was promptly, ‘Ja wohl.’ Immediately I added, ‘Parlez-vous Français?’ and his answer was as immediate, ‘Un peu.’ Without a moment’s pause I added, ‘Sprechen Sie Englisch?’



He hesitated a few seconds and then said distinctly, 'Very little,' adding, with a smile, 'This is a pleasant day; I am glad to see you,' and saying in German that was the extent of his knowledge of English.

"Mr. Hirsch then retired to the other side of the room, a distance of some twenty feet, and speaking in a whisper, told young Polano in Dutch that my father was the first teacher of deaf mutes in America, that my mother was deaf and dumb, and that none of my brothers or sisters were deaf. Polano understood him perfectly and required no repetition.

"As I was under the necessity of parting from Mr. Hirsch at this time in order to take a train for Cologne, there was no further opportunity there for me to test Polano's powers of articulation and lip-reading. But I asked him if he would not walk with me to my hotel, and he replied, 'Mit vergnügen.'

"I will give in English the greater part of what passed between us after starting on our walk, premising the remark that *all* our conversation was in *oral German*, without the use of a single sign.

"As we left the house of Mr. Hirsch, Polano said, 'What hotel are you staying at?' I replied, 'The Hôtel des Pays Bas.' 'Oh, I know it,' said he. 'Do you know my name?' he asked. 'Yes,' said I, 'it is Polano.' 'That is right,' said he, and we exchanged cards. 'Do you not believe that I was born deaf?' he inquired. 'Oh, yes,' said I, and added immediately, 'Do you talk with your sister by signs or with the voice?' 'With the voice,' replied he; 'I prefer it.' 'Isn't it very warm to-day?' said he. 'Very warm,' was my answer.

"Presently I remarked, 'I think we are not going right, for my hotel.' Oh, yes,' said he, 'we are right; did not you say you were stopping at the Hôtel des Pays Bas?' 'Yes,' I answered, 'that is the name of my hotel.' 'Then we are quite right,' said he, adding, 'I live in Rotterdam, you remember, and know the city well.'

"We walked on farther, when, being quite sure we were astray, I repeated that I feared we were wrong, adding that we were following quite a different course from that I took in going from my hotel, and asking if there were two hotels



of the name Pays Bas in Rotterdam. He said he thought not; and so we kept on.

"Growing quite certain we were wrong, I stopped and insisted we were not right, and said I feared I should be too late for the Cologne train if we did not reach my hotel soon.

"He seemed much troubled, and asked me if I would prefer to take a carriage. I said I would; and so we hailed a cab-driver, and Polano asked him if there were two Hôtels des Pays Bas in Rotterdam. The cabman replied that there were; and mentioned that one was Adler's. I then remembered that was the name of the proprietor of my hotel, and so we jumped into the cab and told the driver to go to Adler's Hôtel des Pays Bas.

"Polano said as we rattled over the stones, in a voice that I perfectly understood, 'I hope my mistake will not make you too late for your train; I did not know there were two hotels of the same name here.'

"On reaching my hotel I paid my bill and got my luggage very hurriedly, and then we hastened on in the carriage to the railway station. On the way I took out my watch, and Polano said, 'Is that an American watch?' On my replying in the affirmative he seemed much interested, and wanted to look at it.

"Just before we reached the railroad station, I asked him how much I ought to pay the driver, and he said he thought one florin was quite enough. He asked me when I should come to Rotterdam again, and I said I hoped in a few years. I asked him when I should see him in America. This question I had to repeat a second time, when he replied with a shrug, that it cost too much money; that, perhaps, by-and-by, when he was rich, he would go. I told him he must come to see me in Washington, if he came to America. He replied he certainly would.

"As we reached the railroad station, he said he hoped I would excuse him for making me so much trouble about getting to my hotel.

"As I handed a porter some money for taking my luggage, he remarked, 'You paid him too much.' He accompanied me to the railroad carriage, and bade me good-by, and in a moment the train moved.



"All this I have described was done in the greatest hurry. From the time I left Mr. Hirsch, Polano and I were either walking at a rapid pace through crowded streets, or riding over the pavements in a carriage, and yet what conversation we had was carried on with perfect ease, and without any resort whatever to the language of signs.

"The circumstances of my interview with Polano were of such a nature as to induce me to accord cheerfully the merit of notable and praiseworthy success to Mr. Hirsch in this case; asking you, however, to bear in mind that the young man and his sister were private pupils of Mr. Hirsch during a period of eleven years, and were, therefore, in the enjoyment of advantages secured at a cost far beyond what can reasonably be demanded at the hands of public legislators or almoners of private benevolence in behalf of the great mass of deaf mutes, coming as they do from families of the poor."\*

The Jews' Home and school for the deaf, in London, was founded by the Baroness Rothschild in 1867, and is the only school in Great Britain where articulation is taught as it should be. It was originally intended for Jews only, but others are now admitted. W. Van Praag, the sole teacher, was educated by Dr. Hirsch, of Rotterdam.

In November, 1869, a free public school for deaf mutes was opened in Boston, taught by articulation, and classes in articulation and lip-reading have been organized in the State institutions of New York, Ohio, Illinois, Wisconsin, and at the American Asylum in Hartford.

What is wanted is the combination of the two systems. The sign language is admitted by every one of intelligence to be a most valuable aid in the instruction of the congenital deaf mute; yet in the present advanced stage of civilization, where education of the highest order is given free as the air we breathe, why should the deaf mute be debarred from his full share of this good gift, if he be only capable of receiving it? I trust that physicians and others will aid in the consummation of this noble work, and recommend to all the institutions of this country that they be provided with proper instructors in the artificial method.

---

\* Report, op. cit.




## CHAPTER XX.

### RÉSUMÉ OF THE SIMPLEST AND MOST EFFECTUAL METHOD OF TREATING THE MORE FREQUENT AFFECTIONS OF THE EAR.

For the benefit of those whose active employment in general practice renders it impossible for them to obtain a more complete knowledge of the subject, the author has added a résumé of the simplest and most effectual methods of treating those more frequent and curable affections of the ear, which, without proper attention, are liable to cause deafness. To those who would give more attention to the subject, he would recommend the perusal of this and other works. He has also had prepared a compact box containing the most necessary instruments and apparatus, which may be obtained at a very moderate cost of Messrs. Kolbe or Gemrig, of Philadelphia. A view of this is given at Fig. 106.

Fig. 106.



The most important instruments are the mirror of Von Tröltsch, and the conical tubular specula of Dr. Gruber, of Vienna. The latter are three or four in number, of different



sizes, are made of German-silver, and, as they must be always clean and bright, are kept in a bone-case small enough to be carried in the vest-pocket. Those who are able to devote more time to the subject we would refer to our chapter on the various forms of lamps, etc. for illuminating the ear and membrana tympani. The concave glass mirror is to be either held in the hand or attached by the forehead-band. It has a central aperture for the eye, and the focal distance is six inches. In examining the ear, one of the conical specula is placed in the auditory canal by first drawing the auricle or external ear upward, and gently rotating the speculum until it is secure without being held. The patient is to be placed opposite a window, so that the light will be reflected from the mirror into the ear. The ear thus examined is in the shadow, and the mirror reflects the rays of light, illuminating the whole auditory canal. By changing the position of the speculum, any part of the canal can be examined. If the conical speculum is employed alone, then the light from a clear sky, or a white cloud, or the sun itself, is allowed to penetrate the ear, by placing the patient directly in front of the window, and inclining his head to the opposite side, the operator being careful to keep his own head out of the way. The great advantage of the concave mirror is, that an examination can be made on a cloudy day, or at night, by having a candle, lamp, or gas-burner held to one side and a little behind. This same mirror can be employed to throw light into any narrow canal, or to view the larynx, or pharynx, or the fundus of the eye.

With this mirror we can inspect the color, form, transparency, and curvature of the membrana tympani. If the membrane contains any foreign bodies upon it, if it is darker or lighter than natural, *these* are all matters of importance in obtaining a knowledge of its diseases. With the tuning-fork placed upon the vertex, we can test the condition of the auditory nerve. If it is heard best in the deaf ear, we can pronounce that there is no nervous deafness.

We then determine by the Valsalvian method if the Eustachian tubes are open, and by introducing the otoscope into the patient's meatus, and desiring him to make a strong effort



at expiration (mouth and nose closed), we can judge that if free it will give a crackling sound, with a certain "thug" (or impulse), as it is called, which is neither dry nor moist, and which can be heard by the surgeon.

As an inspection of the membrana tympani and the Valsalvian operation do not give us very accurate results, the Eustachian catheter can then be resorted to, or Politzer's bag, which will be found in the box. As it requires considerable time to become expert in the use of the catheter, the active practitioner had better resort to Politzer's method for inflating the middle ear, or tympanum, and also in applying remedies in the numerous cases of deafness from cold, or aural catarrh, with which he will often meet. The operation is simply inserting the nozzle (attached to a short piece of tubing, which is connected with an elastic bag) into the patient's nostrils and closing them both, swallowing with the mouth shut (previously having taken some water),—the surgeon compressing the bag at intervals as the patient swallows to assist him in his efforts. When employed with little children, a simple piece of tubing is all that is required, the surgeon blowing through it.

*The injured auricle, or external ear,* is treated on general surgical principles, avoiding the cartilages, passing ligatures only through the integuments, and supporting the stitches by means of a pad of cork, covered with lint, or some other agent, with collodion or adhesive plaster. In a recent case, where there was almost entire separation of the auricle by a blow from a shovel, the vitality of the part was found so great that a union took place by simply applying, as a first dressing, pledgets of lint, which adhered and formed a perfect covering with the blood.

*In gangrene* we would recommend the yeast-poultice, which is both antiseptic and a stimulant, with the addition of charcoal and a small portion of the oleum terebinthinæ. Deformity of the ear in childhood is often caused by malposition of the external ear, as, for instance, from wearing a cap or hat in the wrong place, and in some cases it may be hereditary.

*Foreign bodies* in the external canal should almost always



be washed out, near to the orifice of the meatus, by the syringe and warm water, then removed by a soft silver curette, or angular forceps.

We should never take the statement of a patient for granted, that there is a foreign body in the ear. Always before washing it out, or introducing an instrument, there should be an examination with the speculum and a bright light.

If the foreign body is a living one, a little warm glycerine should be poured in, or olive oil, so as to kill the insect, and afterwards the ear must be washed out with the syringe and warm water.

*Eczema auriculæ* (acute and chronic).—This troublesome affection is treated, in its acute form, by soothing applications of bland fluids, the ear being covered with oil-silk, and internal tonics, with alteratives of iron, arsenic, or mercury, being given. In the chronic form, the ointments of the protonitrate of mercury, or the benzoated oxide of zinc, must be used, with cod-liver oil in the case of children.

*Inspissated cerumen*.—That there is cerumen in the auditory canal is to be determined by careful inspection, and it is sometimes overlooked by being thin and dark, and adhering to the membrana tympani. Soft cerumen should be broken up and removed by the curette; but if hard, must be softened by a warm solution of bicarbonate of soda in rose-water (1 gr. to fʒi), and then removed by the syringe or curette. This will not always relieve the hearing, as attention must be given to the condition of the lining membrane of the auditory canal, membrana tympani, and middle ear; and applications are often necessary of astringents and alteratives, with the use of the air-douche of Politzer, or the Eustachian catheter. By a reference to the table, p. 56, it will be seen that the whole number treated was 113, and of these 73 were cured—that is, their hearing was entirely restored; 35 were improved, and in 5 no improvement followed the removal of the apparent cause.

Accumulation of cerumen accompanies a large number of permanent defects in hearing, followed by and being the cause of deafness; producing pressure on the membrana tympani, the delicate bones of the ear, affecting the laby-



rinth, semicircular canal, and cochlea, which latter, as you know, contains the nerve of hearing.

*Furunculous abscess in the external meatus.*—These abscesses are exquisitely painful, slow in development, and form a core which is discharged with difficulty. There is also a plugging up of the canal by a swelling of its tissues. The most successful treatment has been the application of moist warmth to the parts, and a free incision into the abscess as soon as its presenting-point could be determined. As the general system in these cases is almost always below the healthy standard, it is proper to recommend the use of a very nourishing diet, with a moderate amount of wine or malt extract, and tonics of iron and quinine in a simple infusion of gentian.

*Otitis externa.*—This is sometimes known as “otorrhœa externa,” and was supposed to be the most frequent form of affections of the ear; but we now know that it is rather rare, and it will be seen there were only 30 cases in 600. It extends from the orifice of the auditory canal to the external surface of the membrana tympani, and is apt to follow violence to the ear, foreign bodies, abscesses, etc. The most successful treatment was proper counter-irritation, by means of small blisters, croton oil, or tincture of iodine, keeping the parts cleansed daily by the use of a mild astringent wash of sulphate of zinc or alum, and now and then the application of a solution of nitrate of silver—x to xl to fʒi of glycerine—applied to the abraded surface by means of a delicate piece of sponge on a flexible holder, while we must keep the Eustachian orifices open and clean, and the middle ear inflated by Politzer’s air-bag, before described, as these parts are more or less affected by continuity of surface.

*Aspergillus.*—This is a vegetable fungus which is found growing in the external auditory canal, and on the surface of the membrana tympani, and resembles “otitis externa,” and requires a careful removal by the forceps, and the parts kept clean by a weak solution of the hyposulphate of soda, or chlorate of lime, gr. x to fʒi of water. The true presence of this growth can only be detected by placing a small portion under the microscope.



*Acute aural catarrh of the middle ear, with perforation.*—The treatment of this affection consists in keeping the parts cleansed by Clarke's or Thudichum's douche, with astringents, tonics, and counter-irritants, with occasional depletion by leeches or small cups. After the acute stage has passed, we must resort to Politzer's method of passing air into the middle ear, so as to restore the hearing.

This is an affection usually amenable to treatment, and to which the term otorrhœa is given. Of ninety-nine cases, thirty-seven were cured—that is, the discharge was checked, but the hearing was not restored in all of these to the normal standard. We, however, save the patient's life by checking the discharge, and relieving the inflammation.

*Chronic otitis media, or chronic suppuration in the middle ear.*—These cases are very numerous, and are also termed otorrhœa, and many of these have existed for years as the results of catarrh, teething, whooping-cough, measles, and scarlet fever, and most of them have large perforations in the membrana tympani, with granulations and ulcerations in the middle ear. In nearly every instance this disease has broken down the constitution of the individual. There is also a discharge from the nose, eruptions on the head and face, with ulceration, and a thickening of the tonsils, uvula, soft palate, and the pharyngeal orifices of the Eustachian tubes. There are numerous remedies for this form of disease of the ear, showing that the treatment, with any one of them, has not been successful. The treatment consists in restoring the system to its normal condition, and cleansing all the parts involved with warm fluids.

*Chronic catarrh of the middle ear, with or without perforation of the membrana tympani.*—If the membrane be perforated we can relieve the noises and discharge more effectually than if it be thickened and the cavity filled with mucus. There should be applied agents in a solution, or vapor of iodine, etc., to give tone to the parts, removing any excessive discharge by a post-nasal syringe every day, and Thudichum's douche. Also employing tonics, alteratives, with good diet to build up the system. In the cavity of the middle ear, closed by the thickened membrana tympani (accompanied by



various kinds of moist or dry sounds, with deafness), artificial perforation of the membrane will be found, in some cases, successful, driving out from the middle ear by the Eustachian catheter, or air-douche, the accumulation of mucus, pus, etc., and subsequently employing slightly stimulating and astringent washes of zinc, alum (if the sounds are moist), and now and then a solution of caustic potash, or bicarbonate of soda, etc. (if the sounds are dry), or a few grains to fʒi of water in the form of spray. If there is sclerosis or dryness of the middle ear, we resort to steam, which was recommended, in 1856, by Dr. W. Rau, and again revived within the last few years. It is passed into the middle ear by means of the Eustachian catheter, which is introduced through the nose into the mouth of the tube, and then held in position by an assistant, or catheter-holder. A second tube from a glass flask with a nozzle is placed within the catheter, and by means of Politzer's bag the steam is sent in jets, not a continuous stream, for fear of heating the hard-rubber catheter and burning the patient's soft palate. The steam can be generated in the flask by a spirit-lamp or heated sand, or by a small gas-stove, or a generator, made of tin or copper, with a valve and two elastic tubes, one for the catheter and the other for the air-bag, or press. With this and various other means which we cannot enter into in a résumé, we may be able to cure our cases; but if not taken in their early stage, the number of cures will be comparatively small, say 12 out of 111; but, as is shown in the table, 51, or nearly one-half, were improved.

*Otitis media*, with inflammation of the mastoid cells, is generally the result of pus being retained in the middle ear (from inflammation acute or subacute, following exposure to cold, while the patient is suffering from otorrhœa), the pus seeking an outlet by the internal track which it inflames, softens, and ultimately ulcerates these soft cells; and unless the inflammation is relieved, or a new outlet made by means of an incision over the mastoid process, the results are always very serious. This in some cases is all the treatment that is necessary; but in more severe cases, the bone is to be broken down and the outlet made larger, by means of a gouge or chisel. In some instances this has been neglected, and, as a



consequence, extensive caries ensued which no treatment of ours could cure; still, in some few instances, by removal of a part or a whole of the dead bone, the patient's life is saved. Another complication, which in some instances is hereditary, is *syphilitic otitis*, which requires a protracted and specific treatment, which, as may be seen in our table, is far from being satisfactory, only 2 being cured of 10 cases. In the complication, *otitis media with polypi*, the results are more satisfactory, 14 cases being cured out of 18.

The treatment was, removal of the polypus by Wilde's snare, polypus forceps, and caustic potash, packing the meatus with cotton saturated with oil, or diluted acetic acid, which, with other proper measures, was continued until the mucous membrane was in a healthy state.

*Periostitis, with cerebral abscess*, is almost invariably fatal; yet in one case we had all the symptoms in the case, and notwithstanding it recovered.

*Neuralgia of the plexus tympani* is a very protracted disease. It is treated by sedatives of morphia, belladonna, chloroform, and tonics of nux vomica, with quinine, to give tone and vigor to the nervous system.

*Effusion of blood upon the membrana tympani* is generally the result of an exudating process, with the development of a glovelike cast of thickened epithelium, and in other cases is the result of a fall or blow on the side of the head. In three cases there was no perforation; in the fourth, there *was* perforation with a discharge of serum for some weeks, with attachment of the membrana tympani to the promontory, and entire loss of hearing on one side. In the three cases the blood was removed, and the membrane restored to a state of health.

*Effusion of serum into the tympanic cavity, or middle ear*: instead of the usual mucous secretions, there are changes in the layers of the coats of the membrana tympani, causing opacity, or, in some instances, the membrane is so clear that one-half of it appears of a higher shade than the lower part, which is of a dark gray. The boundary between the two is marked with a dark-blue line; this line, according to Politzer, has the appearance of a black hair lying upon the membrane.



The treatment was the air-douche, used every second day, with restoration of the mucous membrane of the nares and pharynx to a healthy condition.

*Perforation of the membrana tympani*, by blows or foreign bodies, usually healed readily if treated with care and without irritating applications. If of long standing, with edges covered with granulations, they are difficult to cure. For these, use a solution of nitrate of silver xx to xl grains to the ounce of water, applied by the surgeon on a pledget of cotton or lint, to the edges of the perforation, the parts being cleansed with tepid water, and air introduced by the Eustachian catheter to rid the middle of any accumulation.

*Acute inflammation of the membrana tympani*.—This affection is accompanied with violent pains in the ear, occurring suddenly, with more or less fever and suppuration. Meatus dry, membrana tympani much inflamed, sensitive, dull, flat, and thickened. Prognosis favorable, if treated promptly. *Perforation* one of the results, but this heals with care and perseverance. Our treatment is perfect rest, free leeching, cupping, or bleeding from the arm, with injections of warm water, glycerine, or opium; and toward the termination, mild astringents and counter-irritants, by tincture of iodine, of small blisters, with the use of air-douche.

*In chronic inflammation of the membrana tympani* there is no fever, and so little pain that the patient is not aware of his disease except by loss of hearing. There is no secretion of cerumen, the membrana tympani is more or less red, or very opaque, dense, with polypi or granulations upon it. Prognosis not favorable. Local and general treatment. The mucous membrane of the nares, pharynx, and Eustachian tube being in almost every case more or less softened and secreting abundantly. A portion of tannic acid, or pulverized alum, is blown in through a rubber tube of the size of a large catheter, which is introduced into the pharyngeal space, in the neighborhood of the Eustachian tube, through the lower nasal opening. Instillation of solution of sulphate of zinc, or powders of protochloride of mercury, are blown upon the membrane, while the patient is given iodide of potassium or bichloride of mercury; the Eustachian catheter or Polit-



zer's air-douche being employed every second or third day. *Inflammation of the Eustachian tube* is to be treated by the means already spoken of under the head of acute aural catarrh; and the chronic form under that of chronic catarrh of the middle ear, with or without perforation. There are numerous kinds of apparatus described at p. 251 et seq., by means of which this portion of the ear is reached and applications made.

*Purulent otitis interna, with perforation.*—In the acute form this occurs in children, and presents all the symptoms of meningitis, but is distinguished from the ordinary form of this disease by the labyrinth being affected, and the meninges. It is apt to terminate fatally, and is almost always followed by complete deafness. The treatment should be depletion by bleeding or leeching, with sedatives, followed by counter-irritation by blisters and tincture of iodine, the internal use of the bromides of sodium or potassium, with the tincture of the sesquichloride of iron. It will be noticed that of 5 cases 2 died, 2 chronic cases were improved, and 1 being unimproved was sent to a deaf-and-dumb institution.

#### A GENERAL SUMMARY OF INFLAMMATORY AFFECTIONS OF THE INNER EAR.\*

“When deafness occurs almost suddenly, under apoplectiform symptoms, in a previously healthy person, and the apoplectiform symptoms disappear rapidly and completely, but the deafness remains, then the seat of the disease is the cavity of the labyrinth, and its pathology probably hemorrhagic or serous effusion.

“The deafness resulting from *injuries* on the head is caused by fissures or fractures of the petrous pyramid, with consequent hemorrhage and inflammation in the membranous labyrinth, or, perhaps, by isolated rupture and hemorrhage in the soft parts of the labyrinth, by ‘*contrecoup*.’

“The chain of symptoms known as *Ménière's disease*, viz.,

---

\* A Clinical Analysis of the Inflammatory Affections of the Inner Ear, by Prof. H. Knapp, Archives of Ophthalmology and Otology, vol. ii. No. 1, pp. 199–283. W. Wood & Co., N. Y. In passing through the press we give his General Summary and Literature on this subject to 1871.



deafness and tinnitus aurium, occurring under symptoms of apoplectiform cerebral congestion, may be a *primary* or a *secondary* affection of the labyrinth.

“The *former* is to be assumed when, in a previously healthy person, the symptoms on the part of the labyrinth—tinnitus, contraction of the range of audition, and hardness of hearing—are prominent and persistent, while the general symptoms—headache, paleness of the face, fainting, obscuration of the visual field, vertigo, nausea, vomiting, and disturbances of equilibrium—cannot be referred to actual disease of the brain or the constitution, and are transitory. The pathology of Ménière’s disease is most probably hemorrhagic and serous inflammation within the whole labyrinthine cavity. *Increase of intra-auricular pressure* may be considered a primary consequence of it, upon which other symptoms—vertigo, nausea, vomiting, disturbance of equilibrium, etc.—depend. Glaucoma of the eye is, perhaps, a counterpart to Ménière’s disease.

“Ménière erred in confining the disease to the *semicircular canals*, the accompanying deafness being evidence that the cochlea is involved. That the disease is neither an affection of the trunks of the auditory nerves, nor of their centre in the brain, is demonstrated by the deafness for high and low tones—a contraction of the range of audition—which I have witnessed. This symptom can be explained only by changes in those parts of the cochlea which are subservient to the perception of high and low tones. It is analogous to the contraction of the visual field in glaucoma.

“Ménière’s disease may be *secondary* to affections of the *tympanum*, the *brain*, and the *general system*. I have reported typical cases in proof of this assertion.

“In a chronic *aural catarrh* the whole chain of the symptoms of Ménière’s disease may occur as an acute complication,—caused by exposure,—disappear, and leave the chronic catarrh in its previous condition, but with increased deafness and tinnitus. In such cases the inflammatory symptoms of the labyrinth may be well marked without noticeable aggravation of the otitis media.

“The deafness, mostly total, occurring in the course of



acute *meningitis*, and sporadic or epidemic *cerebro-spinal meningitis*, is due to *purulent otitis interna*, a complication—perhaps a metastasis—of the purulent inflammation of the meninges, in the same way as purulent choroiditis is not an infrequent complication of the same diseases. Voltolini's opinion, that the whole affection in such cases is an inflammation of the labyrinth alone,—*otitis labyrinthica*, mistaken for meningitis,—lacks both foundation on pathological anatomy and clinical analogy.

“Among the *general diseases* which are liable to produce deafness, *constitutional syphilis* furnished a valuable example in a case, showing a pure attack of Ménière's disease, in a patient who soon after had an analogous affection of his eyes. This being exudative irido-choroiditis, chiefly of a serous character, it may be inferred that the labyrinthine affection also was a serous exudative inflammation. *Severe fevers*—for instance, *typhus* or *typhoid*; *acute exanthemata*: variola, scarlatina, rubeola; *eclampsia infantum*; *erysipelas*; *pustular* and *eczematous eruptions on the head*; *mumps*, *angina Ludwigi*, and *sunstroke* are all apt to produce deafness without material changes in the middle ear. The most probable anatomical lesion in all these cases seems to be serous or hemorrhagic inflammation in the labyrinth.

“The lesion producing deafness after childbirth may be of the same nature or metastatic, as in severe puerperal fevers; anæmia, in certain cases.

“We may, therefore, *subdivide otitis interna into three kinds*:

“1. *Hemorrhagic*: sudden and traumatic deafness.

“2. *Serous exudative*, or *simple*: idiopathic Ménière's disease, and the deafness from syphilis, severe fevers, acute exanthemata, erysipelas, and other constitutional diseases.

“3. *Purulent*: a *primary* disease of traumatic origin, perhaps also spontaneous and *sui generis* (Voltolini), and *secondary* in simple and cerebro-spinal meningitis, puerperal fever, and other pyæmic diseases.

“The *diagnosis* of inflammation of the labyrinth is mainly supported by the symptoms of permanent irritation and paralysis of the auditory nerve:—tinnitus aurium and deafness,—accompanied by transitory cerebral symptoms:—headache,



paleness of the face, vertigo, fainting, nausea, vomiting, disturbance of equilibrium,—without marked changes in the middle or outer ear. Secondary inflammation of the labyrinth is to be assumed when, in the general diseases enumerated above, tinnitus aurium and deafness, without symptoms of otitis media, are well marked.

“The *prognosis* of idiopathic otitis interna varies with the nature and cause of the inflammation. In traumatic cases the hearing may be partially or almost totally restored. The secondary affections of the labyrinth in typhoid fever, syphilis, and other constitutional diseases may terminate favorably, or, on the other hand, leave all possible degrees of hardness of hearing, even total deafness being not infrequent. Idiopathic serous exudative otitis interna, if we may call Ménière’s disease by that name, is mostly fatal to the hearing. If the first attack be not severe and do not destroy the hearing more or less completely, other attacks will follow at indefinite intervals, and gradually result in deafness. The course of this disease bears great analogy to glaucoma. Purulent otitis interna is almost always followed by complete deafness. Only in exceptional cases of cerebro-spinal meningitis hearing has been restored.

“The *treatment* of the labyrinthine affections is still more in its infancy than is its pathology. Antiphlogosis and derivation, both locally and constitutionally, with due regard to the nature of the general symptoms, are indicated in all recent cases. Leeches behind the ears, aperient remedies, among which calomel may be preferred; sudorifics, digitalis, etc. and absorbents; hydrargyrum, iodide of potassium, etc.; afterwards vesicants over the mastoid region should be recommended. Electricity, according to different methods, has nearly always failed; Brenner’s method, however, claiming success in exceptional cases, may be tried. I have employed it in nearly all the reported cases, but without a shade of improvement. If the increase of intra-auricular tension were better known as a fundamental and detrimental symptom in certain inflammations of the inner ear, we should endeavor to get access to the cavity of the labyrinth with proper instruments, perhaps best by one of the fenestræ, in order to evacuate a part of the perilymph.



(The papers marked with \* have been read by the writer\* in the original, the others in extracts.)

"\*1. Paul Mérière. *Gaz. Méd. de Paris*, 1861, pages 29, 55, 88, 239, 379, 597.

"\*2. Von Tröltsch, *Treatise*. Lecture 28th.

"\*3. A. Politzer, Lesion of the Labyrinth. *Arch. f. Ohrenh.* II. pp. 88-99. 1865.

"\*4. Dr. Voltolini, Injury of Head; Complete Deafness; Death; Autopsy. *Monatsschr. f. Ohrenh.*, 1869, pp. 109 and 110.

"5. Delstanche, Lettres sur l'Otologie. *Journ. de Brux.*, 1862, pp. 227, 441, and 543. In \*extract in Schmidt's *Jahrb.*, Bd. 118, pp. 331 and 332.

"6. S. Moos, Sudden Deafness. *Wien Med. Wochenschr.*, Nos. 41, 42, and 43. 1863.

"\*7. S. Moos, *Clinic of Aural Diseases*, pp. 313 and 314.

"\*8. J. Orne Greene, Obstinate Vertigo, Nausea, and Vomiting dependent on Disease of the Ear. *Bost. Med. and Surg. Journ.*, Jan. 21, 1869, vol. 79, p. 388, etc.

"9. Flourens, *Recherches Expérimentales sur les Propriétés et les Fonctions du Système Nerveux*. Second ed. Paris, 1842, p. 438.

"10. Harless. *Wagner's Handwörterbuch der Physiologie*, IV. p. 422.

"11. Czermak. *Comptes Rendus*, 1860, LI. p. 821; and, *Jenaische Zeitschrift*, III. 1867, p. 101.

"12. Brown-Séquard, *Course of Lectures on the Physiology and Pathology of the Central Nervous System*. Philadelphia, 1860, p. 194.

"13. Vulpian, *Leçon sur la Physiologie Générale et Comparée du Système Nerveux*. Paris, 1860, p. 600.

"14. Schiff, *Lehrbuch der Physiologie des Menschen*. Lahr, 1858-59, p. 399.

"\*15. Fr. Goltz, Ueber die physiologische Bedeutung der Bogengänge des Ohrlabyrinths. *Pflüger's Arch.*, 1870, III. p. 172, etc.

"\*16. A. Trousseau, *Lectures on Clinical Medicine*. From



the French by J. R. Cormack. 1870. Vol. III. p. 548, etc. (Vertigo ab aure læsa.)

"17. Hillairet, Lésion de l'Oreille Interne: Action Réflexe sur le Cervelet et les Pédoncules. *Comptes Rendus, et Mémoire de la Société de Biologie*. Third Series. Vol. III. p. 148. For the year 1861. Paris, 1862.

"\*18. A. Lucae on the Production of Sound by the Cranial Bones. *Arch. f. Ohr.*, V. pp. 82 and 117.

"\*19. Voltolini, The Acute Inflammation of the Membranous Labyrinth, commonly mistaken for Meningitis. *Monatsschr f. Ohr.* Berl., Oct., 1867, pp. 9-14.

"\*20. Idem, l. c. Juni, 1868, p. 91, etc.

"\*21. Idem, l. c. July, 1870, p. 91, etc., and August, 1870, p. 103.

"\*22. Reichel, Otitis intima sive labyrinthica. The Acute Inflammation of the Membranous Labyrinth, an Idiopathic Disease hitherto commonly mistaken for Meningitis. *Berl. Klin. Wochensch.*, Nos. 24 and 25, 1870.

"\*23. Moos, *Klinik der Ohrenkrankheiten*. Wien, 1866, pp. 322-327.

"\*24. Itard, *Traité des Maladies de l'Oreille*. Paris, 1842, p. 226.

"25. Huhllings Jackson, Epileptic or Epileptiform Seizures occurring with Discharge from the Ear. *British Med. Journ.*, June 26, 1869.

"\*26. Koeppe and Schwartz, Two Cases of Reflex Epilepsy in Diseases of the Ear. *Arch. f. Ohrenheilkunde*, VI. p. 282, etc.

"\*27. Jos. Gruber, *Text-book of Otology*. Wien, 1870, p. 617."



# BIBLIOGRAPHY

OF THE

## MORE IMPORTANT ARTICLES ON THE

# EAR AND ITS DISEASES,

PUBLISHED

FROM 1683 TO 1871.\*

---

DU VERNEY: *Traité de l'Organe de l'Ouïe, contenant la Structure, les Usages et les Maladies de toutes les Parties de l'Oreille.* Paris, 1683. (Avec XVI planches.) Translated and edited, London, 1737.

TSCHUDIUS, J. CH.: *Otoiatria s. Aurium Medicina.* Basle, 1715, 4to.

CLELAND, A., and WHATHAN, J.: *Essays in Phil. Trans.* London, 1741 and 1755.

LESCHÉVIN: *Mémoire sur la Théorie des Maladies de l'Oreille et sur les Moyens que la Chirurgie peut employer pour leur Curation. Mém. sur les Sujets proposés pour les Prix de l'Académie royale de Chirurgie.* Paris, 1778. Tom. iv. P. 1. Nouvelle édit., Paris, 1819. Tom. iv. P. 1.

DESMONCEAUX, M.: *Traité des Maladies des Yeux et des Oreilles.* Paris, 1786. Tom. ii. Seconde édition, Paris, 1806.

KRITTER, J. FR., und LENTIN, L. FR. B.: *Ueber das schwere Gehör und die Heilung der Gehörfehler. Mit Anmerkungen und Zusätzen aus alle den bisher bekannt gewordenen Beobachtungen herausgegeben von Christian Friedrich Niceus.* Leipzig, 1794.

WILDBERG, C. J. L.: *Versuch einer anatomisch-physiologisch-pathologischen Abhandlung über die Gehörwerkzeuge des Menschen. Mit Kupfern.* Jena, 1795.

TRAMPEL: *Wie erhält man sein Gehör gut, und was fängt man damit an, wenn es fehlerhaft geworden ist?* Pyrmont, 1800. Zweite Aufl., mit Anmerk. und einer Vorrede von K. Th. Menke. Hannover, 1822.

SAUNDERS, JOHN CUNNINGHAM: *The Anatomy of the Human Ear, illustrated by a Series of Engravings of the Natural Size; with a Treatise on the Diseases of that Organ, the Causes of Deafness, and their Proper Treatment.* London, 1806, fol. Edit. 2, London, 1817. Edit. 3, 1821, Philadelphia edition, by Wm. Price, M.D., one of the surgeons of the Pennsylvania Hospital.

CURTIS, JOHN HARRISON: *A Treatise on the Physiology and Diseases of the Ear, etc.* London, 1817. Ed. 6, 1836.—*Abhandlung über den gesunden*

---

\* Some references not found in this list have been quoted in full in the body of the work.



und kranken Zustand des Ohres, nebst einer kurzen Uebersicht vom Baue und den Verrichtungen dieses Organs. A. d. Engl. übers. und mit praktischen Anmerkungen begleitet von Heinrich Robbi. Mit einem Kupfer. Leipzig, 1819.

CURTIS, JOHN HARRISON: Cases illustrative of the Treatment of Diseases of the Ear, both Local and Constitutional. London, 1818.—Interessante Krankenfälle, ein Beitrag zur besseren Behandlung örtlicher oder constitutioneller Gehörleiden u. s. w. A. d. Engl. übers. und mit Anmerkungen versehen von Heinrich Robbi. Leipzig, 1823.

DELEAU: Tableau des Maladies de l'Oreille, qui engendrent la Surdit . Commercay, 1820, fol.

ITARD, J. M. G.: Trait  des Maladies de l'Oreille et de l'Audition. Deux tom. Paris, 1821. 2e  dit., Paris, 1842.—Die Krankheiten des Ohres und des Geh rs. A. d. Franz s. Mit 3 Tafeln Abbildungen. Weimar, 1822. Chirurgische Handbibliothek, Bd. 4.

CASPER, J. L.: Die Krankheiten der Ohren und des Geh rs nach ausl ndischen Beobachtern bearbeitet. Rust's Magazin f r die gesammte Heilkunde, Bd. 10. S. 445–470, Bd. 11. S. 203–229, Bd. 12. S. 120–145.

BECK, KARL JOSEPH: Die Krankheiten des Geh rorganes. Ein Handbuch zum Gebrauche seiner Vorlesungen. Heidelberg und Leipzig, 1827.

ITARD, J. M. G.: De la Surdit . Monograph, pp. 506. Paris, 1828.

TOD, DAVID: The Anatomy and Physiology of the Organ of Hearing; with Remarks on Congenital Deafness, the Diseases of the Ear, some Imperfections of the Organ of Speech, and the Proper Treatment of these Several Affections. London, 1832.

KRAMER, W.: Erfahrungen  ber die Erkenntniss und Heilung der langwierigen Schwerh rigkeit. Berlin, 1833.

KRAMER, W.: Die Erkenntniss und Heilung der Ohrenkrankheiten. Zweite sehr verb. u. verm. Aufl. seiner "langwierigen Schwerh rigkeit." Mit Abbildungen in Kupferstich. Berlin, 1836.—Zweite g nzlich umgearbeitete und sehr vermehrte Auflage. Mit 2 Kupfertafeln. Berlin, 1849; London, trans. and ed., 1837.

VON VERING, JOSEPH RITTER: Aphorismen  ber Ohrenkrankheiten. Wien, 1834.

LINCKE, C. G.: Sammlung auserlesener Abhandlungen und Beobachtungen aus dem Gebiete der Ohrenheilkunde. Erste bis f nfte Sammlung. Leipzig, 1836–1841.

LINCKE, C. G.: Handbuch der theoretischen und praktischen Ohrenheilkunde. 3 Bde. Leipz., 1837–1845.

WILLIAMS, JOSEPH: Treatise on the Ear, including its Anatomy, Physiology and Pathology. London, 1840.

BRESSLER: Die Krankheiten des Seh- und Geh rorgans. Berlin, 1840. A. u. d. T.: Die Krankheiten des Kopfs und des Sinnesorgane. Bd. 2.

FABRICI, P.: Ueber die am Ohr vorkommenden Operationen. Uebersetzt von Lincke. Leipz., 1841.

POLANSKY: Grundriss zu einer Lehre von den Ohrenkrankheiten. Wien, 1842.

PILCHER, G.: A Treatise on the Structure, Economy, and Diseases of the Ear. London, 1838, and 2d ed. 1842. Philadelphia edition, 1843.



JONES, T. WHARTON, M.D.: Ear and Hearing. *Cyclopædia of Practical Surgery*. Ninth Part. London, 1841.

RUETE, C. G. T.: *Klinische Beiträge zur Pathologie und Physiologie der Augen und Ohren*. Nach der numerischen Methode bearbeitet. Erstes Jahreshft. Braunschweig, 1843.

DUFTON, WILLIAM: *The Nature and Treatment of Deafness and Diseases of the Ear*. London, 1844.

VON GAAL, GUSTAV: *Die Krankheiten des Ohres and deren Behandlung*. Wien, 1844. Philadelphia edition, 1848.

FRANK, MARTELL: *Praktische Anleitung zur Erkenntniss und Behandlung der Ohrenkrankheiten; ein Handbuch der praktischen Ohrenheilkunde, mit 156 in den Text eingedruckten Holzschnitten*. Erlangen, 1845.

KRAMER, W.: *Beiträge zur Ohrenheilkunde*. Mit 19 statistischen Tabellen. Berlin, 1845.

SCHMALZ, E.: *Erfahrungen über die Krankheiten des Gehöres und ihre Heilung*. Mit 4 Tafeln in gr. Folio. Leipzig, 1846.

SCHMALZ, E.: *Beiträge zur Gehör- und Sprach-Heilkunde*. 3 Hefte. Leipz., 1846-1848.

HUBERT-VALLEROUX, M. E.: *Essai théorique et pratique sur les Maladies de l'Oreille*. Paris, 1846.

KRAMER, W.: *Ueber den Werth ohrenärztlicher Erfahrungen, mit besonderer Bezugnahme auf Schmalz's "Erfahrungen" und "Beiträge."* Berlin, 1847.

VON WALTHER, PH. FR.: *Krankheiten der Organe des Hörsinnes*. *System der Chirurgie*. Bd. 2. Freiburg, 1847. S. 217-330.

HARVEY, W., and BUCHANAN, THOMAS: *A New and Improved Synoptical Table of the Diseases of the Human Ear, with their Symptoms, Causes, and Treatment*. London, 1848.

KRAMER, W.: *Die Ohrenheilkunde in den Jahren 1849 und 1850*. Ein wissenschaftliches Zeitbild. Berlin, 1851.

BRYAN, J.: *Treatise on the Anatomy, Physiology, and Diseases of the Ear*. Philadelphia, 1852.

WILDE, WILLIAM R.: *Practical Observations on Aural Surgery and the Nature and Treatment of Diseases of the Ear*. London: John Churchill. 1 vol. 8vo. Republished 1853, Philadelphia, edited by A. Hewson, M.D.

This work did more for the advancement of a true knowledge of the pathology and treatment of diseases of the ear than any of its predecessors.

WITH, O. H.: *Ohrentönen und nervöse Schwerhörigkeit*. [Ear-Sounds\* and Nervous Deafness.] Altona: Wendemann. 8vo.

LOEWE, WILH.: *Erkenntniss und Behandlung der Taubheit*. [The Pathology and Treatment of Deafness.] Pasewald: Braune. 8vo.

COE: *Ohrenleiden, mit Betheiligung des N. vagus*. [Affections of the Ear, with Participation of the Vagus Nerve.] *Assoc. Journ.*, 104.

GUCKELBERGER: *Beiträge zur pathologischen Anatomie der Entzündung des Gehörorgans*. [Contributions to the Pathological Anatomy of the Ear.] *Zeitschr. f. Chir. u. Geburtsh.*, vii. 3.

PILCHER: *The Membrana Tympani in its Physiological Relations*. *Med. Times and Gaz.*, Feb. 1854.

---

\* Part of this list was published in the *Trans. Am. Med. Assoc.*, 1865, by the author.



HOUSSELLE: Mittheilung über Toynbees künstliches Trommelfell. [A Communication concerning Toynbee's Artificial Memb. Tympani.] Deutsche Klin., 43.

ERHARD: Der künstliche Tensor tympani, oder Taubheit heilbar durch Druck. [The Artificial Tensor Tympani, or Deafness curable by Pressure.] Deutsche Klin., 52. 1854.

GULL, DR.: Cases of Phlebitis, Pneumonia, and Pleurisy after Chronic Diseases of the Ear. Med.-Chir. Transact., xxxviii. 157. 1855.

SANTESSON: Fall von Caries der Pars petrosa am linken Ohr u. s. w. [Case of Caries of the Petrosal Bone of the Left Ear, etc.] Hygiea, Bd. 14.

ERHARD, J.: Reform der Ohrenheilkunde. [Reform of Ear Therapeutics.] Berlin: Förstner. 8vo. 1855.

ERHARD, J.: Das Gehör und die Schwerhörigkeit. [Hearing and Hardness of Hearing.] Berlin: Förstner. 8vo. 1855.

RINNE: Beiträge zur Physiologie des menschlichen Ohres. [Contributions to the Physiology of the Human Ear.] Prag. Vjhrschr., xlv. 71; xlvi. 45. 1855.

RAU, W.: Lehrbuch der Ohrenheilkunde für Aerzte und Studirende. [Manual of Aural Surgery for Physicians and Students.] Berlin: Peters. 8vo. 1856.

HARVEY, WM., F.R.C.S.: The Ear in Health and Disease. London: H. Renshaw. 2d ed., 8vo. 1856.

V. TSCHARNER: Ueber die Trommelfellentzündung. [On Inflammation of the Membrana Tympani.] Schw. Kant. Zeitschr., 2. Heft.

HINTON'S Cases illustrating the Pathology of the Ear. Med.-Chir. Trans., xxxix. 101. 1856.

MARTINO: Abnorm grosse Ohrmuscheln und Verkleinerung derselben durch blutige Operation (Otomiosie). [Abnormally Large Auricle, and its Reduction by a Bloody Operation.]

BRUNETTI: Offenes Sendschreiben an Hrn. Prof. Schuh, über die Methode des Prof. Vanzetti, fremde Körper aus dem Ohr zu entfernen. [A Letter to Prof. Schuh, concerning Prof. Vanzetti's Method of removing Foreign Bodies from the Ear.] Wien. Med. Wochenschr., vi. 16. 1856.

BONNAFONT: Neues einfaches Instrument zur Ausziehung fremder Körper aus dem äuss. Gehörgange. [A New Simple Instrument for the Extraction of Foreign Bodies from the Meatus.] L'Union, 123.

MÉNIÈRE: Fraktur des Hammergriffs. [Fracture of the Malleus.] Gaz. de Paris, 50.

TRIQUET: Chronische Otitis mit Zerstörung der Trommelfelle auf beiden Seiten; bedeutende Otorrhöe und Granulationen in beiden Trommelhöhlen. Anwendung zweier künstlicher Trommelfelle. [Chronic Otitis, with Destruction of the Membrana Tympani on both Sides; considerable Otorrhœa and Granulations in both Tympanic Cavities. Application of two artificial Membrana Tympani.] Gaz. des Hôp., 105. 1856.

MÉNIÈRE: Abhandlung über in den verschiedenen Theilen des Gehörorgans beobachtete Knochensequester. [Essay on Sequesters of Bone observed in the Various Parts of the Organ of Hearing.] Gaz. de Paris. 1856.

MARTIN: Fall von Tumor erectilis am Ohre. [Case of Erectile Tumor in the Ear.] Gaz. des Hôp., 102.

MÉNIÈRE: Ueber die Auskultation zur Diagnose von Ohrkrankheiten.



[On Auscultation in the Diagnosis of Ear-Diseases.] Bull. de l'Acad., t. xxii. 16, 731.

LAUWERS: Bemerkungen über die Anwendung der Auscultation zur Diagnose von Ohrkrankheiten. [Observations on the Application of Auscultation to the Diagnosis of Ear-Diseases.] Gaz. Hebd., iv. 20. 1856.

CERUTTI, GIUSEPPE: Del Cateterismo della Tuba Eust., etc. [On Catheterism of the Eustachian Tube, etc.] Torino: Biancardi. 4to.

MELCHIOR: Entzündung des äussern und innern Ohres in der Kindheit u. s. w. [Inflammation of the External and Internal Ear in Childhood, etc.] Journ. f. Kinderkr., Hefte 7-12.

GLÜCK, J.: Krankheiten der Luftwege, welche das Gehör beeinträchtigen. [Diseases of the Respiratory Passages which injure the Hearing.] Oesterr. Zeitschr. f. prakt. Heilk., iii. 31, 39, 46. 1856.

VALERIO, PASQUALE: Nuovo Metodo di Cura per Uotirrea. [New Method of Cure for Otorrhœa.] Napoli: G. Gioja. 4to.

KRAMER, W.: Die Gegensätze zwischen Seh- und Hörorgan. [The Contrasts between the Organ of Sight and that of Hearing.] Deutsche Klin., 41.

SCHMALZ: Erklärung über den Wolff'schen Apparat zur Einleitung von wässerigen Dämpfen in die Ohrtrompete, und über die Heilung eines Taubstummen. [Description of the Apparatus of Wolff for the Application of Watery Vapor to the Eustachian Tube for the Cure of Deafness.] Med. Centr. Ztg., xxvi. 5. 1856.

BELLMONT: Fall von Neuralgia Chordæ Tympani, Verwechslung mit Katarrh des Mittelohres. [Case of Neuralgia of the Chorda Tympani Nerve, alternating with Catarrh of the Middle Ear.] Schw. Monatschr., Dec. 1856.

GENDRIN: Neue Untersuchungsart des Gehörorgans, anwendbar zur Diagnose verschiedener Erkrankungen dieses Organs. [New Mode of Exploring the Ear, applicable to the Diagnosis of various Diseases of this Organ.] Gaz. des Hôp., 105. 1856.

SCHNEPF: Geschichtliche Bemerkungen über die Anwendung der Auscultation zur Diagnose der Krankheiten des Ohres. [Historical Observations on the Application of Auscultation to the Diagnosis of the Diseases of the Ear.] Gaz. de Paris, 43.

RENU, WOLF: Versuche über die Unterscheidung differenter Schallstärken. [Researches concerning the Distinguishing of Sounds of different Strengths.] Arch. f. phys. Heilk., ii. 185. 1856.

ERHARD, J.: Ueber Schwerhörigkeit, heilbar durch Druck. [On Hardness of Hearing curable by Pressure.] Berlin: Förstner. 8vo.

KRAMER, W.: Ueber Erhard's "Schwerhörigkeit, heilbar durch Druck." [On Erhard's Hardness of Hearing curable by Pressure.] Deutsche Klin., 49.

KRAMER, W.: Die Ohrenheilkunde in den Jahren 1851-1855. [Aural Surgery in the Years 1851 to 1855.]

NOTTINGHAM, JOHN: Diseases of the Ear. London: John Churchill. 8vo. 1857.

TOYNBEE's Catalogue of Preparations illustrative of Diseases of the Ear. 1857.

V. TRÖLTSCHE: Beiträge zur Anatomie des menschlichen Trommelfells. [Contributions to the Anatomy of the Human Membrana Tympani.] Zeitschr. f. wiss. Zoologie, ix. 91. 1857.



WOLFF, PH. H.: Entgegnung auf die Erklärung des Hrn. Schmalz über meinen Apparat zur Einleitung von Dämpfen in die Ohrtrompete u. s. w. [A reply to the foregoing.] Das. 10.

V. TRÖLTSCHE: Die Untersuchung des Gehörorgans in der Leiche. [The Examination of the Ear in the Dead Body.] Virchow's Arch., xiii. 6, 513. 1858.

KRAMER, W.: Die akute Entzündung des Trommelfells. [Acute Inflammation of the Membrana Tympani.] Deutsche Klin., 22, 23. 1858.

DELEAU: Paralyse des N. facialis, willkürlich hervorgebracht in einem Falle von Erkrankung des mittlern Ohres. [Paralysis of the Facial Nerve voluntarily produced in a Case of Disease of the Middle Ear.] Bull. de l'Acad., xxiii. 193, Janv. 1858.

CLARKE, EDW.: Causes, Effects, and Treatment of Perforation of Membrana Tympani. Amer. Journ. Med. Sci., Jan. 1858. Comments of Von Tröltsch in the Med.-Chir. Monthly Journ., 1859, vol. i.

KRAMER, W.: Der diagnostische Schlauch und die Krankheiten des mittlern Ohres. [The Diagnostic Tube and the Diseases of the Middle Ear.] Deutsche Klin., 52, fig. 1858.

KRAMER, W.: Taubstummheit geheilt durch Herrn San.-Rath Bamberger. [Deaf-muteness cured by Dr. Bamberger.] Deutsche Klin., 2.

KRAMER, W.: Toynbee und Erhard. Eine kritische Abhandlung. [Toynbee and Erhard. A Critical Essay.] Deutsche Klin., 11.

ERHARD: Berichtigung zu "Toynbee und Erhard." [A Correction of "Toynbee and Erhard."] Deutsche Klin., 21.

MEYER, H.: Zur Anatomie der Taubstummheit. [On the Anatomy of Deaf-muteness.] Virchow's Arch., xiv. 5, 6, p. 551. 1858.

V. TRÖLTSCHE: Anatomische Beiträge zur Ohrenheilkunde. Sektion von 16 Schwerhörigen. [The Dissection of the Ears of 16 Persons who were hard of hearing.] Virchow's Arch., xvii. 1, 2, p. 1. 1859.

CZERMAK: Inspektion des Cavum pharyngo-nasale. [Inspection of the Naso-pharyngeal Space.] Wiener Med. Wochenschr., 32. 1859.

VOLTOLINI: Ein Erleuchtungsapparat zur Untersuchung kranker Ohren, auch anderer Höhlen des menschlichen Körpers. [An Illuminating Apparatus for the Examination of Diseased Ears, and also of other Cavities of the Human Body.] Virchow's Arch., xvii. 1, 2, p. 193. 1859.

TURNBULL, L., M.D.: Otorrhœa as a Sequel of Scarlatina: Its Treatment; with illustrative Cases. Medical and Surgical Reporter, Phila., April, 1858. Comments of D. F. Condie, M.D., in the North American Med.-Chir. Review, Jan. 2, 1859.

VOLTOLINI: Zur Funktion des Steigbügels und dessen Ankylose im ovalen Fenster. [On the Function of the Stirrup and its Anchylosis in the Oval Window.] Deutsche Klin., 34, 35, 36. 1859.

BONAFOS and LAZURNU: Otorrhœa; Death; Induration of the Inferior Peduncle of the Cerebellum. Ann. par Jamain et Waker, p. 219. 1860.

TOYNBEE: Acute Caries of the Tympanic Cavity. Med.-Chir. Trans., p. 217. 1860.

BILLROTH: A New Instrument for the Extraction of Foreign Bodies from the Ear. Deutsche Klinik, No. 32. 1860.

V. TRÖLTSCHE, A. FR.: Die Anatomie des Ohres in ihrer Anwendung auf



die Praxis und die Krankheiten des Gehörorganes. Beiträge zur wissenschaftlichen Begründung der Ohrenheilkunde. [The Anatomy of the Ear in its Practical Application to the Diseases of the Organ of Hearing. Contributions to a Scientific Foundation of Aural Medicine and Surgery.] Würzburg. 1861.

This little work is one of the most valuable contributions to aural surgery which have appeared for some time.

"Deutsche Klinik," Nos. 28 and 30, by Hermann Schwartz. 1861. The best brochure on diseases of the ear occurring in typhus fever.

TURNBULL, L., M.D.: Report of Three Cases of Otitis Interna successfully treated. Med. and Surg. Reporter, Philadelphia, Feb. 15 and 22, 1862.

HYRTL: A Case of Fracture of the Neck of the Malleus in a Prairie Dog. Wiener Med. Wochenschr., No. 11. 1862.

AGNEW, C. R.: Case of Necrosis of Cochlea and Labyrinth. Amer. Med. Times, vol. vi., No. 16, p. 183. Tröltsch, Diseases of the Ear, Amer. edition. 1863.

Berlin Medical Society Experiments, Virchow, Chairman. Schmidt's Year Book. New Sydenham Society, 1863. Tröltsch, Diseases of the Ear, Amer. edition, p. 157. 1863.

YEARSLEY, J.: Deafness practically illustrated; being an Exposition of the Nature, Causes, and Treatment of Diseases of the Ear. Sixth edition. London: John Churchill & Sons. 1863.

BRENNER: On Galvanism in Nervous Noises of the Ear. Petersburg Med. Zeit., No. 8. 1864.

TRIQUET: Statistics of Diseases of the Ear. Gaz. des Hôp., No. 137. These relate to 895 cases treated by Triquet at his clinic during four years, 1860-64.

VOLTOLINI: Post-mortem Appearances in the Ear of the Deaf and Dumb. Virchow's Archiv, vol. xxxi., No. 2. 1864. This is the fourth series of autopsies, consisting of 19 cases. Voltolini dwells much upon the importance of making a more exact examination of the labyrinth.

GRUBER: Cases of Necrosis of the Petrous Bone. Wien. Allg. Zeit., Nos. 41, 43, 45. 1864. An extraordinary and unique case is given, in which the cochlea, with a portion of the wall of the tympanum, was discharged from the meatus of each ear, the boy recovering.

KRAMER: Criticism on Politzer's Mode of removing Obstruction of the Eustachian Tube. Deutsche Klinik, No. 34.

SIEGLE: Pneumatic Ear-Speculum. Ibid., No. 37. A new instrument for determining with exactitude the amount of mobility remaining in the membrana tympani. 1864.

GROSS, S. D.: On Extraction of Foreign Bodies from the Ear and Nose. Amer. Journ. of the Med. Sci., Oct. 1864. His instrument consists of a delicate, light cylindrical bar of steel, five and a half inches long, somewhat roughened at its middle to afford a good hold, spoon-shaped at one extremity, and having at the other a very small tooth or prong, projecting at a right angle from the shank.

HILTON'S "Two Cases of Chronic Impairment of Hearing improved after Scarlet Fever and Erysipelatous Sore Throat," and an account of the "Dissection of the Ear in a Case of Deafness." The dissection of the ear is in a case in which death occurred suddenly during an attack of rheumatic fever,



after the patient had been deaf for sixteen years. *Guy's Hospital Reports*, Third Series, vol. x. London, 1864.

PAGENSTECHER: On Trephining the Mastoid Process and Petrous Bone. *Arch. f. klin. Chir.*, iv. 523. 1864.

ROOSA, D. B. ST. JOHN: On Extraction of Foreign Bodies from the Ear. *Am. Journ. Med. Sci.* 1864.

ERHARD, J.: *Vade Mecum for Clinical Otiatry*. Lissa: Günther. 1864.

SCHWARTZE, H.: *Practical Contributions to Aural Surgery*. Würzb.: Stahel.

WEBER, E.: *Brief Communications on Aural Surgery, etc.* Berlin.

VIRCHOW, R.: On Malformations of the Ear. *Arch. f. path. Anat.*, vol. xxx. p. 221.

OPPENHEIMER, Z.: On the Diagnosis of Ear-Diseases. *Loc. cit.*, p. 240.

OCKEL, GLAMA, and WREDEN: Report of the Cases at the Maximilian-heilanstalt for the Years 1858-61. *Petersb. Med. Zeitschr.*, v. 129, 257; vi. 257, 321.

BÖKE, J.: Cases. *Wien. Med. Halle*, 1864, pp. 379, 391.

ERHARD: On Examination of the Ear with reference to Soldiers and Medico-legal Questions. *Wien. Med. Wochenschr.*, 1864, pp. 730, 761.

WEBER, J. E.: Cases. *Berl. Klin. Wochenschr.*, 1864, p. 164.

DUPLAY, S.: An Examination of recent Publications on the Anatomy, Physiology, and Pathology of the Ear. *Arch. Gén.*, 1864, ii. 327, 576.

LUCAE, A.: *Anatomico-Physiological Contributions*. *Arch. f. path. Anat.*, vol. xxix. p. 33.

BRENNER: On the Use of Galvanism. *Loc. cit.*, vol. xxviii. p. 197.

HINTON: Improvement in hearing Voice, but not in hearing Watch. *Med. Times and Gaz.*, 1863, ii. 644.—Case. *Guy's Hospital Reports*, x. 340. On Insufflation of Alum in Chronic Catarrh of Meatus, etc. *Med. Times and Gaz.*, 1864, i. 11.

RUSCHENBERGER, W. S. W., M.D., U.S.N.: Description of a Syringe for Washing the Auditory Canal. *Amer. Journ. of the Med. Sci.*, vol. xlviii. p. 86. 1864.

FOLLIN: Case of Trepanning the Mastoid Process. *Gaz. des Hôp.*, 1864, p. 4.

TRIQUET: On Bleeding from the Ear. *Loc. cit.*, p. 26.

GRUBER: On the Diseases of the Muscles of the Ear. *Wien. Med. Halle*, 1864, p. 36.

BÖKE, J.: On Cancer of the Ear. *Loc. cit.*, 1863, pp. 428, 438.

POLITZER's Method of Inflating the Tympanum. *Med. Times and Gaz.*, 1864, i. 10.

KRAMER, W.: On the Introduction of Fluids, etc. into the Tympanic Cavity. *Deutsche Klin.*, 1864, p. 113. On Politzer's Method of Inflating the Tympanum. *Loc. cit.*, p. 325.

HERMANN, A.: On a New Mode of Catheterizing the Eustachian Tube. *Wien. Med. Halle*, 1864, p. 339.

ANDERSON, T. MCCALL: A Short Sketch of Obstructions in Eustachian Tube, etc. *Glasgow Med. Journ.*, xii. 1.

POLITZER, A.: On the Artificial Membrane. *Wien. Med. Halle*, 1864, p. 5, etc.



SCHREIBER: On Perforations of the Membrana Tympani. Loc. cit., pp. 323, 342.

GRUBER, J.: On Myringotomy. Allg. Wien. Med. Zeit., 1864, p. 305, etc.

DARDEL: Anomalous Condition of the Internal Ear in Congenital Deaf-mutism. Schweiz. Zeitschr. f. Heilk., iii. 155. 1864.

A Year-Book of Medicine, Surgery, and their Allied Sciences, 1863 and 1864, for New Sydenham Society, London, 1864 and 1865.

GUERSANT, M.: Removal of Foreign Bodies from the Ear in Children. Bull. de Thérap., Dec. 30, 1864.

TRIQUET: Perforations of the Membrana Tympani. Gaz. des Hôp., No. 29.

AUSPITZ: Eczema of the External Ear. Archiv der Ohrenheilkunde, No. 2.

TRIQUET, M.: Chilblains of the Ear. Gaz. des Hôp., No. 13.

SCHWARTZE: Report on Aural Medicine during the Last Decennary. Archiv für Ohrenheilkunde, No. 3.

WELCKER: Osseous Growth and Closure of the External Meatus. Ibid.

KRAMER, DR. W.: Ohrenheilkunde und Ohrenärzte in England und Deutschland. Ein Nachtrag zur Ohrenheilkunde der Gegenwart. Berlin, 1865. Pp. 96. [Aural Surgery and Aural Surgeons in England and Germany. A Supplement to Aural Surgery of the Present Day.]

Archiv für Ohrenheilkunde, herausgeg. von Dr. v. Tröltsch (Würzburg), Dr. Politzer (Wien), und Dr. H. Schwartze (Halle), 1864, '5, '6, '7. The communications upon the ear in this volume are numerous and valuable.

HELMHOLTZ, PROF. H.: Die Lehre v. den Tonempfindungen als physiologische Grundlage f. die Theorie der Musik. Mit in den Text eingedr. Holzst. 2. Ausg. gr. 8. (XI u. 606 S.) Braunschweig, 1865: Vieweg & Sohn. New (3d) edition, 1870.

MACH, DR. ERNST: Bemerkungen über die Accommodation d. Ohres. (Abdr. aus den Sitzungsber. d. k. Akad. d. Wiss.) Lex.-8vo. (4 S.) Wien, 1865: Gerold's Sohn.

MACH, DR. ERNST: Untersuchungen über den Zeitsinn d. Ohres. Mit 3 (eingedr.) Holzschn. (Abdr. aus d. Sitzungsber. d. k. Akad. d. Wiss.) Lex.-8vo. (18 S.) Ebd., 1865.

POLITZER, DR. ADAM: Die Beleuchtungsbilder d. Trommelfells im gesunden u. kranken Zustande. Klinische Beiträge zur Kenntniss u. Behandlg. der Ohren-Krankheiten. Mit 24 chromolith. Trommelfell-Bildern (auf 2 Taf.) u. 13 in den Text gedr. Holzschn. gr. 8. (VII u. 143 S.) Wien, 1865: Braumüller. Translated and edited by Drs. Mathewson and Newton, New York, 1869.

RÜDINGER, PROSECT. DR.: Ein Beitrag zur Anatomie u. Histologie der Tuba Eustachii. Mit 1 (lith.) Abbildg. (in 4.) (Abdr. aus dem ärztlichen Intelligenz-Blatte.) gr. 8. (16 S.) Ebd., 1865.

BOCKÉ, JULIUS: Ueber d. Primäre Entzündung der Trommelfells. Wien. Med. Presse, vii. 19, 20. [The Primary Inflammation of the Membrana Tympani.]

DUMONT, O.: Obhämatom am beiden Muscheln. Berlin. Klin. Wochenschr., iii. 28. [Double Hæmatoma.]

FOX, CORNELIUS: Relations of Ear and Larynx. Lancet, i. 17, April, 1866.



HAGEN, R.: I. Praktische Beiträge zur Ohrenheilkunde. [Practical Treatises on the Diseases of the Ear.] Leipzig, 1866.

HAGEN, R.: II. Der seröse Ausfluss aus dem äussern Ohr nach Kopfverletzungen. Leipzig. [Serous Exudation of External Ear after Injury to Head.]

HILL: Der gegenwärtige Zustand d. Taubstummen-Bildungs-Wesens in Deutschland. Weimar. [The Present Condition of the Deaf and Dumb Institutions of Germany.]

WENDT, H.: Ueber d. Affectionen d. Mittel-Ohrs im kindl. Alter. Zeitschr. f. Med., Chir. und Geburtsh., N. F., v. 2, p. 100. [Affections of the Middle Ear in Childhood.]

WOLF, PHIL. HEIN.: Die nervöse Schwerhörigkeit u. ihre Behandlung durch ein neu Method. 2. Auf. mit 3 lithog. Taf. Berlin. [Treatment of Nervous Deafness by a New Method.]

ZAUFAL, EMANUEL: Zur patholog. Anatomie des Gehörorgans. Arch. f. Ohrenheilkunde, ii. 3, p. 174. [Pathology and Anatomy of the Organs of Hearing.]

BÜCHTING, ADOLPH, Buchhandler: Bibliotheca Otiatrica oder Verzeichniss aller auf dem Gebiete der Ohrenheilkunde in den letzten 20 Jahren, 1847-1866. Nordhausen, 1867.

FRIEDRICH, DR.: Die Krankheiten des Gehörorganes. Eine allgemeinfassl. Anleitung, Ohrenzwang, Ohrensausen, Schwerhörigkeit u. Taubheit zu verhüten u. zu heilen. 8vo. (VIII u. 110. S.) Quedlinburg, 1866.

REINECKER, D.: Ueber Othämatom. Würzburg. Med. Ztschr., vii. 1, p. 13. [On Hæmatoma.]

ROOSA, B. ST. JOHN: Use of Artificial Membrana Tympani. Amer. Jour. Med. Sci., cl. p. 106.

SCHWARTZE, HERM.: Ueber subjective Gehörempfindungen. Berlin. Klin. Wochenschr., iii. 12-13.

TOYNBEE, JOS.: Treatment of Perforation of Membrana Tympani. Lancet, i., March 12, 1866.

BERHOL, JOHN: Pathology of Sounds in the Ear. Lancet, ii., July 3-4, 1866.

BABELEI: Ueber Verhütung der Taubstummheit. Journ. de Brux., 3e sér., ii. p. 83, Févr. [Protection from Deafness.]

BERTIN, E.: Ueber Anwendung der comprimirten Luft bei Behandlung der Taubheit. Gaz. de Par., xlix. 781. [On the Use of Compressed Air in the Treatment of Deafness.]

BONIG, F.: Klinisch-otiatrische Mittheilungen. Hannover. Zeitschr. für prakt. Heilk., iv. 1, p. 17. [Abstract of Ear Clinic.]

BREARD, GEORGE, M.D., of New York: The Membrana Tympani and Pharynx in Deaf-Mutism. Am. Jour. Med. Sci., April, 1866.

CHIMANI, R.: Ueber die Wirkungsweise d. Politzer'schen Verfahrens bei Behandlung d. eiterigen Mittelohrkatarrhs mit Perforation d. Trommelfells. Wien. Med. Wochenschr., xvi. 35, 36. [On the Mode and Efficiency of Politzer's Method in the Treatment of Purulent Catarrh of the Middle Ear by Perforation of the Membrana Tympani.]

COESFELD, H.: Zur Entfernung der Ohrenschmalzpropfe. Berl. Klin. Wochenschr., iv. 21. [On the Removal of Impacted Cerumen.]



ERHARD: Brauchbare Hörrohre. Berl. Klin. Wochenschr., iv. 31. [Useful Ear-Trumpets.]

FARNECK: Fall ein. Fremderkörper in äusserm Gehörgange. Arch. für Ohrenheilk., ii. 4, p. 300. [Case of a Foreign Body in the External Auditory Canal.]

FANT: Nothwendige ohrenärztliche Geräthschaften. Bayer. Aerztl. Intel., Bd. 28. [Necessary Apparatus for Otology.]

GRUBER, JOS.: Beiträge zur Anatomie d. Schläfenbeins und ihrer Anwendung auf praktische Ohrenheilkunde. Wien. Med. Wochenschr., xvii. 53, 54, 73. [Contributions to the Anatomy of the Temporal Bones, etc.]

GRUBER, JOS.: Ueber ein neues Verfahren zur Eintragung medicamentöser Flüssigkeiten in einem mittleren Ohrtheil ohne Zuhülfenahme eines Katheters. Oesterr. Zeitschr. für prakt. Heilk., xiii. 42. 1866. [On a New Method of introducing Medicated Fluids into the Middle Ear without the Use of a Catheter.]

HAGEN, A.: Ueber das Politzer'schen Verfahren und dessen Verwerthung für die Einführung von Wasserdampfen in der Trommelhöhle. Wien. Med. Presse, vii. 47, 48. [On Politzer's Method for the Introduction of the Vapor of Water into the Tympanic Cavity, and its Value.]

HAGEN, R.: Die Pflege d. Ohres im gesunden und kranken Zustande. Leipzig. 1866. [The Care of the Ear in Health and Disease.]

HAGEN, R.: Praktische Beiträge zur Ohrenheilkunde. III. Die circumscribed Entzündung d. äussern Gehörganges. Mit Holzschn. Leipzig. [Practical Observations on Diseases of the Ear. Circumscribed Inflammation of the External Auditory Passage.]

HERMANN, ADOLF: Caries d. Trommelhöhle u. d. hintern Wendung d. Canalis Caroticus: Arrosion der in letzterem verlaufenden Carotis cerebralis. Tod durch Verblutung. Wien. Med. Wochenschr., xvii. 30, 31, 32. 1866. [Caries of the Tympanic Cavity, involving the Posterior Curvature of the Carotid Canal. Erosion of the Carotis cerebralis. Death from Hemorrhage.]

HILL: Grundzüge eines Lehrplans für Taubstumm-Anstalten u. der Unterricht in desselben betr. Einrichtungen. Weimar. [Principles of a Plan of Instruction in Institutions for the Deaf and Dumb.]

LUCAE, AUG.: Ueber eine neue Methode zur Untersuchung des Gehörorgans zu physiologischen u. diagnostischen Zwecken mit Hülfe des Interferenz-Otoskops. Arch. für Ohrenheilkunde, iii. 2, No. 3, p. 186. [A New Method of investigating the Organs of Hearing for Physiological and Diagnostic Purposes, etc.]

MOOS: Ueber das subjective Hören wirklicher musikalischer Töne. Virchow's Archiv, xxxix. 2, p. 280. [On the Subjective Hearing of Musical Tones.]

PETERS: Fälle von Katarrh a. Mittelohrs, aus Dr. F. Weber's Ohrenklinik zu Berlin, mit besonderer Rücksicht auf Gruber's Einspritzungsverfahren. Wien. Med. Wochenschr., xvii. 69, 70. 1866. [Cases of Catarrh of the Middle Ear, from Dr. Weber's Ear-Clinic in Berlin, with Special Reference to Gruber's Mode of injecting the Middle Ear.]

POLITZER, A.: Diagnose und Therapie der Ansammlung seröser Flüssigkeit in der Trommelhöhle. Wien. Med. Wochenschr., xvii. 16. [Diagnosis and Therapeutics of the Collection of Serous Fluids in the Middle Ear.]



POLITZER, A.: Ueber luftdichte Obturation des äusseren Gehörganges als Heilmittel bei chron. Mittelohrkatarrhen. *Wien. Med. Wochenschr.*, xvii. 78-85. 1866. [On Hermetic Obturation of the External Auditory Passage as a Remedy for Chronic Catarrh of the Middle Ear.]

SCHWARTZE, H.: Statist. Ber. die in d. Poliklinik zu Halle-am-S. im Wintersem. 1865-66 und Sommer 1866 untersuchten und behandelten Ohrenkranken. *Archiv für Ohrenheilk.*, iii. 1, p. 22. [Statistical Report on the Ear-Patients examined and treated in the Polyclinic at Halle-on-the-Saale in the Winter of 1865-66 and the Summer of 1866.]

SCHWARTZE, H.: Ueber Indikationen und therapeutischen Werth der künstl. Perforation d. Trommelfells. *Arch. für Ohrenheilk.*, ii. 4, p. 239. [The Indications and Therapeutic Value of Artificial Perforation of the Membrana Tympani.]

SCHWARTZE, H.: Fall einseitiger Taubheit: schnelle Heilung durch Heurteloup's Blutegel. Pp. 298. 1866. [A Case of One-sided Deafness: Rapid Cure by means of Heurteloup's Leech.]

SCHMELCK, CHR.: Ueber chron. Ohrenkatarrh und dessen Behandlung, *Norsk Mag.*, xxi. 2, p. 65. [On Chronic Aural Catarrh and its Treatment.]

TOLMATSCHEW, N.: Fall von Othaematom. *Deutsche Klin.*, xvi. [Case of Hæmatoma.]

WATSON, W. SPENCER: Examination of the Membrana Tympani. *Med. Times and Gazette*, Aug. 31, p. 239. 1867.

WENDT, H.: Beiträge zur Verallgemeinerung d. Ohrenheilkunds. *Zeitschr. für Med., Chir. u. Geburtsh.*, N. F., vi. 2, p. 87. [Treatises on the Generalization of Treatment of Diseases of the Ear.]

WERNER: Chronischer Ohrenfluss; Tod unter meningitischen Erscheinung; ausgedehnte Thrombosenbildung in Hirnblutleitern und in der Diploë des Schläfenbeins; klein cariöse Stelle in der Tiefe des Felsenbeins. *Zeitschr. für Wundärzte und Geburtsh.*, xx. 3, p. 204. [Chronic Otorrhœa; Death, with Meningitic Symptoms; Thrombosis of the Diploic Structure; Caries.]

BONNAFONT: Fall von vollständiger Taubheit bedingt durch Verstopfung des äussern Gehörganges durch eine nahe am Trommelfell sitzende Exostose: Heilung durch Trepanation. *Gaz. des Hôp.*, 64. [Case of Complete Deafness induced by Occlusion of External Auditory Canal by an Exostosis near the Membrana Tympani. Recovery of Hearing by Trepanning, etc.] 1867.

BONNAFONT: Ueber Operation der Oberpolypen mittels Ecrasement. *Bull. de l'Acad.*, xxx. p. 57. [Operation for Ear-Polypus by the Ecraseur.]

BROWN, A. GARDINER: Removing of Foreign Bodies from External Auditory Canal. *Med. Times and Gaz.*, Aug. 8, 1868.

COUSIN: A. Politzer's Verfahren zur Injection von Luft und medikamentären Dämpfe in Tuba Eustachii und Trommelhöhle. *Bull. de Thér.*, lxxiv. p. 161. 1868. [Poltizer's Results from Injection of Air and Medicated Vapors in Eustachian Tube and Middle Ear.]

DELSTANCHE: Ueber Otorrhœe; deren Folgen und Behandlung. *Jour. de Brux.*, xlvi. p. 314. [Otorrhœa, its Consequences and Treatment.]

GROSSMANN: Purulente Basilar meningitis in Folge recidivirender Otitis interna, mit Gangrän der Ohrmuschel umgebenden Weichtheile. *Mon.-Schr. f. Ohrenheilkunde*, ii. 68. [Purulent Basilar Meningitis in Connection with Otitis Interna, with Gangrene of External Ear and Adjacent Soft Parts.]



GRUBER: Rückblicke auf die wissenschaftliche Leistungen in der Ohrenheilkunde. Mon.-Schr. f. Ohrenheilk. 1868. [Retrospect of the Scientific Results in Otolology.]

GURIN: Insekten im äussern Gehörgang: (Soc. de Chir.) Gaz. des Hôp. [Insects in the External Auditory Canal.]

GIAMPIETRO, EDOARDO: Ueber die Anwendung des Otoskops zur Diagnosis von Ohrenkrankheiten. Il Morgagni, x. 3-6. [The Use of the Otoloscope in Diagnosis of Diseases of the Ear.]

GRUBER, JOSEF: Ueber Autophonie und Tympanophonie: Beiträge zur Auscultation des Gehörganges. Monat-Schr. f. Ohrenheilk., ii. 8.

GRUBER, J.: Untersuchung über die Anatomie des äussern Gehörganges, mit besondere Rücksicht auf die prakt. Ohrenheilkunde. Monat-Schr. f. Ohrenhkl., i. 5-6. [Investigation of the Anatomy of the External Ear, with Special Reference to Practical Otolology.]

GUYE, A. A.: Ueber d. Beziehungen zwischen dem Pharynx und der Trommelhöhle. Nederl. Tijdschr., 2. Afl. [On the Connection between the Pharynx and the Tympanic Cavity.]

HELMHOLTZ: Mechanik der Gehörknöchelchen u. d. Trommelfells. Archiv für Physiologie, i. 1; und Monat-Schr. für Ohrenheilk., ii. 10. 1868. [Mechanism of the Small Bones of the Ear and of the Membrana Tympani.]

HENKE, W.: Der Mechanismus der Gehörknöchelchen. Henle und Pf.'s Zeitschr. für rationale Med., 3. B., xxxi. 1868. [Mechanism of the Small Bones of the Ear.]

KLOTZ, H. G.: Ueber Ohrenpolypen. Inaug.-Diss. Leipzig.

KRISHABER, MONTZ: Stummheit ohne Taubheit. Gaz. Hebdom., 2e sér., v. 39. [Dumbness without Deafness.]

KUHNEL, FRIEDRICH MAX: Die Paracentese d. Trommelfells. Inaug.-Diss. Leipzig. 1868. [Perforation of the Membrana Tympani.]

LEFORT: Fremder Körper im Ohre bei einem 4jähr. Kind. Gaz. des Hôp., 57, p. 227. 1868. [Foreign Body in the Ear of a Child Four Years old.]

LAREYS, R.: Sicher u. billige Heilung d. Schwerhörigkeit, ein Rathgeber für Alle welche an Gehörkrankheiten leiden u. s. w. Leipzig. 1868. [Safe and Cheap Cure of Deafness: A Counselor for all who suffer from Hardness of Hearing.]

MOSETRY: Zur Casuistik a. fremd. Körper in Ohren. Wien. Med. Presse, ix. 32. [Foreign Bodies in the Ear, etc.]

MITTENDROP, H. W.: Zur Histologie u. Entwicklung d. Schnecke. Monat-Schr. für Ohrenheilk., ii. 11. 1868. [On the Histology and Development of the Cochlea.]

MARCHAL, CH.: Purulente Otorrhöe mit Otitis Strumosa. L'Union, 46.

OEKEL: Insekten in äussern Gehörgänge. Petersb. Med. Zeitschr., xii. 6, 363. [Insects in the External Auditory Canal.]

OFFENGER, H.: Ueber einige auffallende Missbildungen des äussern und innern Ohres. Henle u. Pf.'s Zeitschr. f. rationale Med., 3. B., xxxi. 1868. [On certain Remarkable Malformations of the External and Internal Ear.]

POLITZER, ADAM: Ueber willkürliche Contractionen des Musc. Tensor Tympani. Archiv für Ohrenheilk., iv. 1868. [On Voluntary Contractions of the Tensor Tympani Muscle.]

PRAT: Ueber Behndl. d. Otorrhöe durch Irrigation mit warmen Was-



sers. Bull. de l'Acad., xxxiii. 1858. [Treatment of Otorrhœa by Irrigation with Warm Water.]

REHSAMEN, AUG.: Zum Mechanismus der Tuba Eustachii. Monat-Schr. für Ohrenheilk., ii. 3. [On the Mechanism of the Eustachian Tube.]

RÜDINGER: Ueber anomales Offensein der Tuba Eustachii bei alten Leute. Mon.-Schr. für Ohrenheilk., ii. 9. [On Anomalous Patulency of the Eustachian Tube in Old Persons.]

SCHWARTZE, HERMANN: Die Paracentese d. Trommelfells. Archiv für Ohrenheilk., iii. 4, P. 218. [Paracentesis of the Membrana Tympani.]

SCHMIDKAM, JOS.: Experimentelle Studien zur Physiologie des Gehörorgans. Inaug.-Diss. Kiel. [Experimental Study of the Physiology of the Organs of Hearing.]

SCHURIG, EM.: Günstige Erfolge der künstlichen Ruptur einer Trommelfellnarben mittels des Politzer'schen Verfahrens. Wien. Med. Presse, ix. 20. [Favorable Result from Artificial Rupture of Cicatrix of the Membrana Tympani, etc.]

SMITH, THOMAS: Investigations on Deafness, etc. St. Barthol. Hospital Reports, Rep. iii., p. 145.

TOYNBEE, JOS.: Cerebral Symptoms in Certain Affections of the Ear. St. George's Hosp. Rep., i., 1868.

WREDEN, ROBERT: Otitis Media Neonatorum. Mon.-Schr. für Ohrenheilk., ii.

WEBER, F. E.: Ueber d. Gehörstörungen b. ganz oder theilweis zerstörtem Trommelfell u. defektem Gehörknöchelchen. [On Defects of Hearing from Total or Partial Destruction of the Membrana Tympani and the Loss of the Bones of the Middle Ear.]

WEBER, F. E.: Otitis Parasitica. Mon.-Schr. für Ohrenheilk., ii. 1868.

WEBER, F. E.: Zur Therapie d. Mittelohr in Specie der Paukenhohlen-Affektionen. Zur Physiologie der Tuba Eustachii. Monat.-Schr. f. Ohrenheilk., 5 and 7. 1868.

ALLEN, PETER: Concerning certain Functions of Middle Ear and its Analogues, etc. Lancet, i. 10, March and May, 1868.

BRUNTON, JOHN: Improved Apparatus for Use of Politzer's Method of Air Douche of Eustachian Tube. Glasgow Med. Journal, Nov. 1868.

BERTUCH, WM.: Ueber Rigidität u. Synostose der Steigbügel. Vorhofverbindung. Inaug.-Diss. Halle, 1869. [Rigidity and Synostosis of the Stapes, and Vestibular Connection.]

BARETY, ALEXANDER, and JOSEPH RENAUT: Anatomical Changes in Otitis Interna of New-Born Children. Arch. Phys., ii. 3, p. 374. 1869.

BRUNNER: Fälle von Othaematom bei Gesunden entstanden, in einem Falle nach traumat. Einwirkung, im andern durch Einwirkung von Kalte. Epithelial Carcinoma des Ohres. [Cases of Hæmatoma in Healthy Persons, in One Case after Traumatic Influence, in another from Cold and Epithelioma of the Ear.]

BUHEL, TH.: Ein Fall von doppelseitiger Caries des Felsenbein mit günstigem Ausgange. Inaug.-Diss. Würzburg, 1869. [Case of Caries of Temporal Bones of both Sides, with Favorable Termination.]

CHIMANI, R.: Ueber die Häufigkeit der Krankheiten d. Gehörganges, u. ihren Einfl. auf d. Militärtauglichkeit. Militär. Arztl., iii. 6-9. 1869. [Fre-



quency of Diseases of the Ear, and their Influence upon Fitness for Military Service.]

COUSIN, A.: Ueber einige auf d. äussere Ohr anwendbar Mittel bei Behandlung verschiedener Krankheiten des Gehörapparats. *Bull. de Thér.*, lxxvii., Oct. 1869. [Concerning certain Remedial Agents applicable in the Treatment of the External Ear, and in the Treatment of other Affections of the Hearing Organs.]

DEMARQUAY: Ueber Cancroid der Ohrmuschel. *Gaz. des Hôp.*, 114.

EULENBERG, ALBERT: Ueber ein noch nicht beobachtete Modification der galvan. Reaktion des N. acousticus. *Arch. für klin. Med.*, v. 5-6, p. 547. [Concerning a Modification of Galvanic Action upon the Auditory Nerve not previously observed.]

ERHARDT: Ueber d. Luft in d. Trommelhöhle. *Berl. Klin. Wochenschr.*, v. 51, p. 531. [Air in the Tympanic Cavity]

ERB, W.: Die galvan. Reaktion des nervösen Gehörapparats in gesund u. kranken Zustände. *Arch. für Augen- u. Ohrenheilk.*, i. 1, 156.

GOTTSTINE: Beitrag zur Ohrenheilkunde. *Arch. für Ohrenheilkunde*, iv. 2, p. 65. 1869. [Contribution to Otology.]

GRUBER, JOS.: Ueber d. Anatomie d. äussern Gehörganges, mit besondere Rücksicht auf d. prakt. Ohrenheilkunde. *Monat-Schr. für Ohrenheilk.*, ii. 12. [Anatomy of the External Auditory Canal, with Especial Regard to the Practice of Otology.]

GRAY: Acute Brain-Symptoms in a Case of Chronic Disease of Internal Ear. Acute Meningitis, Abscess, etc. *Med. Times and Gaz.*, February 27, 1869, p. 223.

GELLÉ: Doppelseit Taubheit ohne nachweisbare Veränderungen. Neuralgie des 7. Nervenpaares und Gehörnerven. *L'Union*, 30. [Deafness without any Demonstrable Cause. Neuralgia of Seventh Pair of Nerves, and Great Anæmia.]

HAGEN, R.: Praktische Beiträge zur Ohrenheilkunde. Die Carbolsäure u. ihre Anwendung in d. Ohrenheilkunde. Leipzig, 1869. [Practical Contributions to Otology. Carbolic Acid and its Uses in Diseases of the Ear.]

HAGEN: Praktische Beiträge zur Ohrenheilkunde. 6. Heft: Casuistisch Belege f. d. Brenner'sche Methode der galvanischen Acousticusreizung. Mit Holzschnitt. Leipzig, 1869.

HINTON, JAMES: Case of Perforation of Cells of Mastoid Process; Formation of Abscess under Sterno-Cleido-Mastoid. *Med.-Chir. Trans.*, li. p. 231.

GREEN, J. ORNE, M.D.: Progress of Otology for 1869-70. *Transactions of the Am. Otol. Society*. D. Appleton & Co., New York, 1870.

MORLAND, WM. W., M.D., of Boston: Deafness in Connection with Pregnancy and the Puerperal Condition.

HINTON, JAMES: Two Cases of Inflammation of the Tympanum, with Development of Polypus; one ending fatally.

MOOS, S.: Alterations of Taste and Sensibility in the Tongue by the Application of an Artificial Tympanum in Case of Large Perforation in both Membrana Tympani. *Archives of Ophthalmology and Otology*. By Prof. H. Knapp and Prof. Moos. Vol. i., No. 1. New York, Wm. Wood & Co., 1869.

MOOS, S.: Investigation on the Relation between the Handle of the Malleus and the Membrana Tympani (with chromo-lithographs). *Arch. Oph. and Otol.*, loc. cit.



- Moos, S.: Two Fatal Cases of Ear-Disease. *Arch. Oph. and Otol.*, loc. cit.
- Moos, S.: Two Cases of Ear-Disease in Court. Doubtful Responsibility. *Arch. Oph. and Otol.*, loc. cit.
- Moos, S.: On the Medico-Legal Significance of Atrophy of the Tympanum produced by Hardened Cerumen. *Arch. Oph. and Otol.*, loc. cit.
- Moos, S.: A Simple Expedient for the Diagnosis of One-Sided Simulated Deafness. *Arch. Oph. and Otol.*, loc. cit.
- Moos, S.: Peculiar Disturbances of Hearing after Cerebro-Spinal Meningitis; Considerable Improvement by the Galvanic Current. *Arch. Oph. and Otol.*, loc. cit.
- POLITZER, ADAM: Ueber bewegliche Exsudate in der Trommelhöhle. *Wiener Med. Presse.* 1869
- ZAUFAL, EMANUEL: Ueber das Vorkommen seröser Flüssigkeit in der Paukenhöhle (Otitis media serosa). *Archiv für Ohrenheilkunde*, Band v. 1869.
- Moos, S.: Serous Accumulations in the Tympanum. *Arch. Oph. and Otol.*, vol. i. No. 2, 1870.
- WEIR, ROBERT F., M.D.: Ingrowing Hairs from the Tragus. *Trans. Am. Otol. Soc.* New York, D. Appleton & Co., 1870.
- ROBERTSON, C. A., M.D., Albany: A Case of Malignant Disease of the Ear. *Trans. Am. Otol. Soc.*, loc. cit.
- AGNEW, C. R., M.D., of New York: Sinus through Mastoid from Old Suppurative Otitis Media; and Trepanning. *Trans. Am. Otol. Soc.*, loc. cit.
- NOYES, HENRY D., M.D., of New York. Otitis Media following the Use of a Bougie in Eustachian Tube; Introduction of an Eyelet into Membrana Tympani; Results. *Trans. Am. Otol. Soc.*, loc. cit.
- GREEN, J. ORNE, M.D., of Boston: A Case of Epithelioma of the Auricle and Meatus, loc. cit.
- MORLAND, WM. W., M.D., of Boston: Foreign Bodies in the Ear; Importance of a Correct Diagnosis; Methods of Removal; Cases. *Trans. Am. Otol. Soc.*, loc. cit.
- GREEN, JOHN, M.D., of St. Louis, Mo.: A Case of Aspergillus in the External Auditory Meatus. *Trans. Am. Otol. Soc.*, 1870, loc. cit.
- KNAPP, H., M.D., of New York: Congenital One-Sided Bony Closure of the External Auditory Canal, with Rudimentary Development of the Auricle. *Trans. Am. Otol. Soc.*, loc. cit.
- ROOSA, D. B. ST. JOHN, M.D., of New York: Hemiplegia occurring in Coincidence with Chronic Suppuration of the Middle Ear. *Trans. Am. Otol. Soc.*, loc. cit.
- PARDEE, CHARLES INSLEE, M.D., of New York: Observations on the Use of Steam in the Treatment of Diseases of the Middle Ear. *Trans. Am. Otol. Soc.*, loc. cit.
- GREEN, JOHN, M.D., of St. Louis, Mo.: Observations upon the Physiology of the Eustachian Tube made during a Short Exposure to an Atmospheric Pressure of Sixty Pounds to the Square Inch. *Trans. Am. Otol. Soc.*, loc. cit.
- BLAKE, C. J., of Boston: A New Form of Wire Snare for the Removal of Aural Polypi, modified from that of Wilde. *Arch. of Oph. and Otol.*, vol. i. No. 2. New York, 1870.
- BUCK, ALBERT H., of New York: On the Mechanism of the Ossicles of the Ear. *Arch. Oph. and Otol.*, vol. i. No. 2. New York, 1870.
- Moos, S.: Historical and Critical Remarks concerning the Deafness fol-



lowing Meningitis Cerebro-Spinalis. Arch. Oph. and Otol., vol. i. No. 2. New York, 1870.

MOOS, S.: Sudden Hemorrhage into Right Tympanum accompanying Angina Diphtheritica. Arch. Oph. and Otol., vol. i. No. 2. New York, 1870.

MOOS, S.: A Case of Idiopathic Diphtheria of the External Meatus. Arch. Oph. and Otol., vol. i. No. 2. New York, 1870.

KAISER, H., of Dieburg: The Mechanism of the Organ of Hearing. Arch. Oph. and Otol., vol. i. No. 2. New York, 1870.

MOOS, S.: A Preliminary Notice on the Anatomy and Physiology of the Eustachian Tube. Arch. Oph. and Otol., vol. i. No. 2. New York, 1870.

BORBERG, AUGUST: Ueber Wölbungsanomalien des Trommelfelles. Inaug.-Diss. Halle. Druck von W. Plötz. 8vo. 34 S. 1870.

DEUTSCH, HEINRICH: Verwandtschafts-Ehen u. Taubstummheit. Mon.-Schr. für Ohrenheilk., iv. 1. 1870.

FLEURY: Einführung einer Nadel in d. Ohr; Meningitis; Tod nach 5 Tagen. (Mitgetheilt von Raphael Percepid.) Gaz. des Hôp., 15. 1870.

FOX, CORNELIUS B.; CLARKE, J. LOCKHART; BUSH, W.: On Ear-Cough and its Causes. Brit. Med. Journ., Dec. 18, 1869, Jan. 15, 1870.

GIAMPIETRO, EDOARDO: Ueber chron. Katarrh d. äussern Ohres. Gazz. Lomb., 52.

GOLTZ, FR.: Ueber die physiolog. Bedeutung der Bogengänge des Ohr-labyrinthes. Arch. für Physiol., iii. 2 u. 3, p. 172. 1870.

GRUBER, JOS.: Lehrbuch der Ohrenheilkunde, mit besond. Rücksicht auf Anatomie u. Physiologie. Mit 88 (eingedr.) Holzschn. u. 2 chromolith. Taf. Wien, 1870. Gerold's Sohn. gr. 8. XVI u. 647 S. 1870.

GRUBER, JOS.: Ueber Syphilis des Gehörorganes. Wien. Med. Presse, xi. 1, 3, 6. 1870.

GRUBER, JOS.: Zur Kenntniss über die Verbindung des Hammers mit d. Trommelfell. Mon.-Schr. für Ohrenheilk., iv. 1. 1870.

HECKSCHER: Fremder Körper in der Tuba Eustachii. Mon.-Schr. für Ohrenheilk., iv. 1. 1870.

HESEL, JULIUS ADOLPH: Ueber Ohrpolypen. (Inaug.-Diss.) Halle. Druck von W. Plötz. 8vo. 38 S mit 4 Tabellen.

JAGO, JAMES: On the Functions of the Tympanum. Brit. Rev., xlv. (89) p. 229. Jan 1870.

KESSEL, J.: Beitrag zum Baue der Paukenhöhlenschleimhaut des Hundes u. der Katze. (Centralblatt f. d. med. Wiss., 6, 1870.) Mon.-Schr. f. Ohrenheilk., iv. 2. 1870.

LENT: Statistik der Taubstummen d. Reg.-Bez. Köln. Köln, 1870. Du Mont-Schauberg. gr. 8. IV u. 40 S.

LUCAE, AUGUST: Zur Behandlung der Otitis purulenta chronica. Berl. Klin. Wochenschr., vii. 6. 1870.

MAYER, LUDWIG: Ueber die fremden Körper im Ohre. Mon.-Schr. für Ohrenheilk., iv. 1, 2. 1870.

POLITZER, ADAM: Ueber ein Höhlensystem zwischen Trommelfell und Hammerhals. Wien. Med. Wochenschr., xx. 15, 16, 18. 1870.

ROSSI, EMILIO DE: L'Otoscopie binoculaire. Mon.-Schr. für Ohrenheilk., iii. 12. 1870.

RÜDINGER, N.: Beiträge zur vergleichenden Anatomie u. Histologie der



Ohrtrompete. München, 1870. Lentner. gr. 4. VIII u. 59 S. mit 11 chromo-lith. Taf., nach mikro-photogr. Aufnahmen.

SCHMIDDEKAM u. HENSEN: Experimentelle Studien zur Physiologie des Gehörorgans. Mon.-Schr. für Ohrenheilk., iii. 11, 12. 1870.

SONRIER: Atresie d. äussern Gehörganges. Gaz. des Hôp., 17. 1870.

STOKES, DR.: Chronic Otorrhœa; Disease of the Middle Ear; Thrombosis of the Sinus petrosi laterales and Sinus cavernosi of Right Side. Brit. Med. Journ., February 12, p. 155. 1870.

TILLMANN, ROB. HERM.: Ueber Facialislähmung bei Ohrenkrankheiten. (Inaug.-Diss.) Halle, 1870. Druck von W. Plötz. 8vo. 39 S.

WEBER, FR. EUG.: Sektionsbefunde bei Schwerhörigen. Mon.-Schr. für Ohrenheilk., iii. 12, 1869; iv. 1, 1870.

WEBER, FR. EUG.: Ueber d. chron. Entzündungs-Zustände des äussern Gehörganges u. die anat. Verhältnisse desselben. Mon.-Schr. f. Ohrenheilk., iii. 11.

WREDEN, R.: Fälle von Fremdkörpern im Ohr. Mon.-Schr. für Ohrenheilk., iii. 12.

WREDEN, R.: Zur Casuistik der angeborenen Missbildungen des Ohres. Mon.-Schr. für Ohrenheilk., iv. 2. 1870.

ZAUFAL, EMANUEL: Sektion der Gehörorgane zweier durch Koblendampf erstickten. Wien. Med. Wochenschr., xix. 104.

BOUISSON: Ueber Amputation der Ohrmuschel. Gaz. de Par., 24. 1870.

BURGER, EM.: Ueber das Einbringen von Flüssigkeit u. Dämpfen durch die Tuba in die Paukenhöhle u. über die Wirkungsweise der Luftdouche. Arch. für Ohrenheilk., v. 4, p. 272. 1870. Oesterr. Zeitschr. für prakt. Heilk., xvi. 22, p. 374.

COLLES, W.: Trepanning of the Mastoid Process. Dubl. Journ., 1. (99) p. 32, Aug. 1870.

DERBLICH (Lemberg): Beobachtungen auf dem Gebiete der Ohrenkrankheiten. I. Die Krankheiten d. Trommelhöhle. Wien. Med. Wochenschr., xx. 31.

JONES, SAMUEL J., M.D.: Report on Otology. Chicago Med. Examiner, Oct. 1870.

HOLMES, E. L., M.D.: A Statistical Report of Diseases of the Ear. Chicago Med. Journ., Sept. 1870.

GREENE, J. ORNE, JR., M.D.: Restoration of the Membrana Tympani after its almost complete Destruction from Chronic Inflammation. Boston Med. and Surg. Journal, Nov. 24, 1870.

ERHARD: Zur Genesis der Gehörkrankheiten. Deutsche Klin., 34, 1870.

EYSELL, A.: Beiträge zur Anatomie des Steigbügels u. seiner Verbindungen. Arch. für Ohrenheilk., v. 4, p. 237. 1870. [Investigations into the Mode of Attachment of the Base of the Stirrup with the Oval Window.]

GRUBER, JOS.: Ueber Ohrenschnalzanhäufung im äussern Gehörgange. Mon.-Schr. für Ohrenheilk., iv. 6. 1870.

GRUBER, JOS.: Anatomische Bemerkungen über das Trommelfell. Mon.-Schr. für Ohrenheilk., iv. 7, 8.

HEDINGER JUN.: Zur Elektrotherapie. Würtemb. Corr.-Bl., xl. 12.

KESSEL, J.: (a) Zur Myringitis villosa. (b) Ueber Form- u. Lageverhältnisse eigenthümlicher, an d. Schleimhaut des menschl. Mittelohres vorkommender Organe. Arch. für Ohrenheilk., v. 4, pp. 250, 254.



MÜLLER, ALWIN: Der Mittelohrkatarrh u. dessen Behandlungsweisen. (Inaug.-Diss.) Leipzig. Druck von C. G. Naumann. 8vo. 15 S.

NICOL, PATRICK: Hæmatoma. Brit. and F. M. C. Rev., xlv. (91) p. 191. July, 1870.

POLITZER, ADAM: Klonischer Krampf der Muskeln der Tuba Eustachii. Wien. Med. Presse, xi. 28. 1870. [A Case of Clonic Spasm of the Muscles of the Eustachian Tube.]

POEITZER, AD.: Pflanzliche Parasiten im Ohre. (Wien. ärztl. Verein.) Wien. Med. Wochenschr., xx. 28, p. 598; Oesterr. Zeitschr. f. prakt. Heilk., xvi. 21, p. 260; Wien. Med. Presse, xi. 22, p. 416. 1870.

POLITZER, AD.: Therapie der beweglichen Exsudate in der Trommelhöhle. Wien. Med. Wochenschr., xx. 35, 37, 39.

POLITZER, AD.: Physiological Experiments on the Functions of the Membrana Tympani and Small Bones of the Ear. (Communicated by Dr. Edwin Millingen.) Med. Times and Gazette, Aug. 6, 1870.

REICHEL, G.: Otitis acuta intima sive labyrinthica, die akute Entzündung d. häutigen Labyrinthes, eine selbstständige Erkrankung, bisher gewöhnlich für Meningitis gehalten. Berl. Klin. Wochenschr., vii. 24, 25. 1870.

RÜDINGER, N.: Ueber die Nerven u. das Epithel in den Ampullen u. Säckchen des häutigen Labyrinthes u. über eigenthüml. Zellenreihen in den häutigen Bogengängen bei *Salmo hucho*. Mon.-Schr. für Ohrenheilk., iv. 5, p. 65. 1870.

RÜDINGER, N.: Verbindung d. Fussplatte des Steigbügels mit dem ovalen Fenster beim Menschen. Mon.-Schr. für Ohrenheilk., iv. 7. 1870.

SCHMIEDT: Ueber die Ménière'schen Labyrinthaffektion. Berl. Klin. Wochenschr., vii. 28. 1870.

SCHURIG: Ueber Paracentese des Trommelfells. Jahresber. d. Ges. f. Natur- u. Heilk. in Dresden, Juni, 1869, bis Mai, 1870, p. 34.

SCHWARTZE, HERMANN: (a) Zur Pathologie der Synostose des Steigbügels. (b) Beiträge zur Pathologie u. pathol. Anatomie des Ohres. Mon.-Schr. für Ohrenheilk., v. 4, pp. 257, 292. 1870.

VOLTOLINI, R.: Nekrotische Ausstossung des vollständigen Labyrinths der rechten Seite. Mon.-Schr. für Ohrenheilk., iv. 6. 1870.

VOLTOLINI, R.: Zur akuten Entzündung des häutigen Labyrinthes. (Otitis intima s. labyrinthica.) Mon.-Schr. für Ohrenheilk., iv. 7, 8. 1870.

WEBER, FR. EUGEN: Kopf- u. Ohrhalter mit Ohrspiegel. Mon.-Schr. für Ohrenheilk., iv. 5, p. 70. 1870.

WEBER, FR. EUGEN: (a) Ueber die durch das Eindringen von Flüssigkeiten in die Paukenhöhle hervorgerufenen entzündl. Erscheinungen, deren Prophylaxis u. Behandlung. (b) Zur Anwendung des Spiritus vini rectificatiss. in der Ohrenheilk. Mon.-Schr. für Ohrenheilk., iv. 6. 1870.

WREDEN, R.: (a) Mehrjähriges unbemerktes Verweilen von zwei Birnenkernen im Ohr. (b) Zweimonatl. Verweilen eines Knochenkügelchens im Ohr. (c) Achtzehnmonatl. Verweilen einer Muschel im Ohr. Mon.-Schr. für Ohrenheilk., iv. 6. 1870.

DALBY, W. B.: Cases of Perforation of Membrana Tympani, i. 17, April; Lancet, ii. 8, Aug. 1870.

DELSTANCHE, CH.: Ueber Anwendung des continuirl. Wasserstromes in der Ohrenheilkunde. Journ. de Brux., li. p. 229, Sept. 1870.



GRUBER, JOSEF: Anatomische Bemerkungen über das Trommelfell. Mon.-Schr. für Ohrenheilk., iv. 9, 10. 1870.

GRUBER, JOSEF: Zur Lehre von den Pilzen im Gehörorgane. Mon.-Schr. f. Ohrenheilk., iv. 9. 1870.

KAISER, H.: Die Mechanik der Gehörfunktion Arch. f. Augen- u. Ohrenheilk., i. 2, p. 100.

POLITZER, ADAM: Therapie der beweglichen Exsudate in der Trommelhöhle. Wien. Med. Wochenschr., xx. 41, 42.

RÜDINGER: Beiträge zur Histologie des Gehörorganes. Mit 5 Taf. in Holzschn. München. Lentner. gr. 8. III u. 30 S. 1871.

RÜDINGER: Der Meniscus in dem Hammer-Ambosgelenk. Mon.-Schr. f. Ohrenheilk., iv. 9. 1871. [Mode of Attachment of the Base of the Stapes.]

VOLTOLINI, R.: Beschaffenheit des Gehörorgans eines Hemicephalus. Mon.-Schr. für Ohrenheilk., iv. 9.

WEBER, FR. EUGEN: Tenotomie des Tensor tympani. Mon.-Schr. für Ohrenheilk., iv. 10. 1871.

WENDT, HERMANN: Beiträge zur pathologischen Anatomie des Ohres. Arch. d. Heilk., xi. 6, p. 562. [Cases of Cerebral Abscess, Thrombosis of the Sinus, and Meningitis in Ear-Patients.] 1871.

BEZOLD, FRIEDR.: Partielle Nekrose d. knöchernen Gehörganges u. d. Paukenhöhlenwand. Mon.-Schr. für Ohrenheilk., iv. 4. 1871.

BONNAFONT: Apparat zur Injektion von Flüssigkeiten in d. Mittelohr u. zur Adspiration solch. aus demselben. L'Union, 51.

American Annals of the Deaf and Dumb. Edited by Edward A. Fay. Vol. xvi. No. 1, January, 1871, Washington, D.C.

HECKSCHER: Beiträge zur prakt. Ohrenheilkunde. Mon.-Schr. f. Ohrenheilk., iv. 3. 1871.

HUMPERDINCK, G.: Die Sprachlaute, physiol. u. sprachwissenschaftlich betrachtet. Siegburg. Berlin. Calvary u. Co. 4to. 23 S. m. 1 Taf. 1871.

BRUNNER, G.: Beiträge zur Anatomie und Histologie des mittleren Ohres. Wilhelm Engelmann, publisher. Leipzig, 1871.

LOEWENBERG, B.: Anatomie et Physiologie de l'Oreille. La Lame spirale du Limaçon et l'Organe de Corti, d'après des Recherches originales. Paris. Impr. Martinet. 8vo. 56 pp. avec 4 pl. Journ. de l'Anat. et de la Physiol. 1871.

LUCAE, AUGUST: Einfaches Stativ für d. Ohrenspiegel. Berliner Klin. Wochenschr., vii. 14. 1871.

MAYER, LUDWIG: Ueber die fremden Körper im Ohre. Mon.-Schr. für Ohrenheilk., iv. 3-5. 1871.

GUDDEN u. LUDWIG MEYER: Ueber den mikroskopischen Befund in traumatisch gesprengten Ohrknorpel. Virchow's Archives, 1870, 1871, Nos. 51 and 52.

OLLIVIER, AUGUSTE: Ueber Gehirnbrunnens nach Otorrhöe. Gaz. de Par., 17. 1871.

PÉTREQUIN, J. E.: Vues nouvelles sur la Composition chimique du Cérumen, et son Rôle dans certaines Maladies de l'Oreille. Lyon. Impr. Regard. 8vo. 98 pp. 1871.

POLITZER, ADAM: Beiträge zu den Hörprüfungen mittels der Stimmgabel. Wien. Med. Presse, xi. 12, 13. 1871.

POLITZER, ADAM: Ueber d. Funktion d. Trommelfells u. d. Gehörknöch-



elchen. (K. k. Gesellsch. d. Aerzte zu Wien.) Oesterr. Zeitschr. für prakt. Heilk., xvi. 18, p. 307; Wien. Med. Presse, xi. 17, p. 321; Wien. Med. Wochenschr., xx. 24, p. 463. 1871.

MILLINGEN, EDWIN: Treatment of Otitis Purulenta, from A. Politzer's Clinic. Med. Times and Gaz., March 26, 1871.

PRAT: Physiologie de l'Audition. Paris. Adr. Delahaye. 8vo. 16 pp.

VOLTOLINI, R.: Ob die Tuba Eustachii des Menschen beständig offen oder geschlossen ist. Mon.-Schr. f. Ohrenheilk., iv. 3. 1871.

WASHINGTON, N. C.: Radical Cure of Otitis Media Purulenta. St. Louis Med. and Surg. Journ., N. S., vii. 2, p. 116, March, 1871.

WEBER, FRIEDR. EUGEN: Zur Casuistik d. tödtlich endenden chron. Otorrhöen. Deutsche Klin., 17, 1871.

WEBER, FRIEDR. EUGEN: (a) Ueber die durch das Eindringen von Flüssigkeiten in die Paukenhöhle hervorgebrachten entzündl. Erscheinungen, deren Prophylaxe u. Behandlung. (b) Zur Therapie der Mittelohrleiden. Mon.-Schr. für Ohrenheilk., iv. 4. 1871.

WENDT, HERM.: (a) Ueber schlauchförmige Drüsen der Schleimhaut der Paukenhöhle. (b) Beiträge zur pathol. Anatomie des Ohres. Arch. d. Heilk., xi. 3, p. 252 fig. 1871.

ALLEN, PETER, DR.: Inflating the Cavity of the Tympanum. Dr. Allen exhibited to the Medical Society of London "An Improvement on Politzer's most Invaluable Appliance." Dr. Tilbury Fox stated that he had been perfectly relieved by Dr. Allen's method of treating catarrhal deafness. Med. Times and Gazette; Med. Cosmos, May 15, 1871.

BUCK, ALBERT H., New York: Case of Angioma Cavernosum. Archives of Ophthalmology and Otology. By Prof. H. Knapp and Prof. S. Moos. Vol. ii. No. 1. New York, Wm. Wood & Co., 1871.

BRUNNER, GUSTAV, of Zürich: On Alteration in the Hearing of One's own Voice. Arch. Oph. and Otol., 1871.

Moos, S.: Hyperostosis of the Cranium and Petrous Bones. Translated by Albert H. Buck. Arch. Oph. and Otol., 1871, op. cit.

POLITZER, ADAM: Cicatrices of the Membrana Tympani. Reported by Mr. E. Millingen. Med. Times and Gaz., March 4, 1871.

Moos, S.: Rare Anomalies, both Congenital and Acquired, occurring in the Ear of a Deaf-Mute. Arch. Oph. and Otol., op. cit.

ELSBERG, L., of New York: On the Employment of the Nasal Douche. Arch. Oph. and Otol., op. cit.

Moos, S.: The Diagnosis of Complete Paralysis of the Auditory Nerve. Arch. Oph. and Otol., op. cit.

BRUNNER, GUSTAV: On Vertigo occurring in Affections of the Ear. Arch. Oph. and Otol., op. cit.

Moos, S.: Four Cases of Gunshot Wound of the Ear. Arch. Oph. and Otol., op. cit.

Moos, S.: Case of Contraction of the Left Tensor Tympani synchronous with the Act of Mastication. Arch. Oph. and Otol., op. cit.

FITCH, W. H., M.D.: Paracentesis Tympani. Chicago Med. Examiner, Feb. 1871.

WEBER, F. EUGEN: Ueber Sekretionen und Ansammlungen von Flüssigkeiten in der Paukenhöhle; deren Wesen, Diagnose und Therapie. Mon.-Schr. f. Ohrenheilk., v. 1. 1871.



ZAUFAL: Ueber die Operation der Paracentese des Trommelfells. Wien. Med. Wochenschr., xxi. 8. 1871.

MILLINGEN, EDWIN: On the Accumulation of Serum in the Tympanic Cavity. Diagnosis and Treatment. Med. Times and Gaz., March 25, 1871.

BUCK, ALBERT H., M.D.: Cases illustrating some of the more Unusual Forms of Ear-Disease. The Medical Record, No. 136, New York, Oct. 16, 1871, p. 362.

FORSTER, COOPER: Naso-Pharyngeal Polypi. The Danger of their Removal without the Laryngoscopic Mirror. Lancet, May 20, 1871; The Phila. Med. Times, Aug. 18, 1871.

WEBER, F. E.: Treatment of Otorrhœa by Spirits of Wine in Cases in which there is neither Caries, Perforation of Membrana Tympani, nor Polypous Growth. The ear must be well syringed out; then, the patient lying down, as much of the pure spirit as the ear will contain is poured in and allowed to remain five minutes. After the spirit flows out, the meatus must be dried and plugged so as to prevent the access of air. At first it will be applied three times a day, then twice, and is to be continued some time after the otorrhœa has ceased. Better results are obtained by adding equal parts of water and proof-spirits. Berl. Klin. Wochenschr., January 9; Med. Times and Gaz., and Phila. Medical Times, Sept. 1, 1871.

SPENCER, H. N.: Treatment of Sclerosis of the Middle Ear by Use of Steam. He employed Dr. Pardee's method in five cases of sclerosis (plastic inflammation) of the middle ear, having an average duration of ten years, which were markedly relieved after a course of treatment varying from twenty days to three months. St. Louis Med. and Surg. Journ., July; Phila. Med. Times, July 15, 1871.

BENEDIKT, M.: Elektr. Untersuchung u. Behandlung Gehörkranker. Mon.-Schr. für Ohrenheilk., v. 1. 1871. (Vgl. Wien. Med. Presse, 37-48. 1870.)

BOETTCHER, ARTH.: Ueber Entwicklung und Bau des Gehörlabyrinths nach Untersuchungen an Säugethieren. Mit 12 Kupfertaf. Leipzig, 1871. Engelmann. gr. 4. IV. u. 203 S.

FITCH, W. H., M.D.: Paracentesis Tympani. Chicago Med. Examiner, Feb. 1871. (A case of serous accumulation in the tympanum.)

BRUNNER, GUST.: Beiträge zur Anatomie u. Histologie d. mittlern Ohres. Mit 4 Kupfertaf. Leipzig. Engelmann. gr. 8. III u. 39 S.

COLLINS, JAS., M.D.: Cases of Aural Disease. Reported by John W. Millick. Phila. Med. and Surg. Reporter, xxiv. 2, p. 35, Jan. 1871.

ERHARD: Zur Genesis d. Gehörkrankheiten. Mon.-Schr. f. Ohrenheilk., iv. 12.

POLITZER, ADAM: Ueber Trommelfellnarben. Wien. Med. Wochenschr., xxi. 1, 2. 1871.

SCHAPRINGER, A.: Ueber die Contraktion des Trommelfellspanners. (Sitz.-Ber. d. k. Akad. d. Wiss.) Wien, Gerold's Sohn. Lex-8. Mon.-Schr. Ohrenheilk., iv. 12.

VOLTOLINI, R.: (a) Die Krankheiten des Trommelfells. (b) Ueber d. Perforation d. Trommelfells. Mon.-Schr. f. Ohrenheilk., iv. 12, 1870; v. 1, 1871.

BURGER, EMANUEL: Ueber Pathologie u. Therapie der subjektiven Ohrengeräusche. Wien. Med. Wochenschr., xxi. 13; Wien. Med. Presse, xii. 23.



FICK: Ueber die Helmholtz'sche Theorie der Gehörempfindungen durch die Membrana basilaris. Verh. d. phys.-med. Ges. zu Würzburg, N. F., ii. 1 u. 2, p. 9.

GRUBER, JOSEF: Bericht über die im Wiener Krankenhause im Jahre 1869 behandelten Ohrenkranken. Mon.-Schr. für Ohrenheilk., v. 2, 3.

GRUBER, JOSEF: Ueber Anomalien in der Spannung des Trommelfells. Mon.-Schr. für Ohrenheilk., v. 3, 4, 5. 1869.

LUCAE, AUGUST: Die Schallleitung durch die Kopfknochen u. ihre Bedeutung für d. Diagnostik d. Ohrenkrankheiten. Würzburg, 1870. Stachel. gr. 8. 48 S.

POLITZER, A.: Ueber endotische Geräusche. (Aerztl. Verein.) Wien. Med. Wochenschr., xxi. 10. 1871.

POLITZER, ADAM: Zur Physiologie des Schallleitungs-Apparats. Wien. Med. Wochenschr., May 27, 1871.

REICHERT, C. B.: Zur feinern Anatomie d. Gehörschnecke beim Menschen u. den Säugethieren. (Zweite Abtheilung.) Arch. f. Anat., Physiol. u. wissenschaftl. Med., i. p. 117.

SCHMALZ, ED.: Ueber Erhaltung d. Gehöres, oder: Das wichtigste über den Bau und die Verrichtung des Gehör-Organes, über die Krankheiten d. Ohres u. Gehöres, über die Verhütung derselben, über das dabei zu beobachtende Verhalten, sowie über den Ersatz u. die Erleichterung des Gehöres; nebst den bei schriftlichem Verkehre zu beantwortenden Fragen. Mit 3 Kupfertafeln. 12. Aufl. Leipzig. Arnold. gr. 8. VIII u. 80 S. 1871.

WOLF, OSKAR: Sprache und Ohr. Vieweg und Sohn, Braunschweig, 1871. [Speech and the Ear.]

DOUGHTY, WM. H.: The Nasal Douche in the Treatment of Naso-Pharyngeal Catarrh, with Report of a Case of Rupture of the Drum of the Right Ear. Richmond and Louisville Med. Journal, Feb. 1871.

PRAT, M. LE DOCT.: L'Action anatomique, physiologique, thérapeutique des Irrigations d'Eau tiède sur la Membrane du Tympan. Bulletin de l'Académie Impériale de Médecine. Tome xxxv. No. 11, page 500. 1871.

BURNETT, DR. CHAS. H.: Three Cases of Fibrous Tumors of the Auricle. Report of Proceedings of Path. Society of Philadelphia. Amer. Journal of the Medical Sciences, Jan. 1871.

HINTON, JAMES: On the Treatment of Perforations of the Membrana Tympani. Guy's Hosp. Reports, Series III. vol. xvi. 1870-71.

MOXHAY, MR., of Reading: Remarkable Deformities of the Ears and Face in a Young Girl. Two perfect ears and three rudimentary auricles. British Med. Journal, Nov. 12, 1870.

BÖKE, JULIUS: Ueber totale Taubheit bei Kindern, in Folge von Erkrankung des Centralnervensystems. Wiener Med. Presse, xii. 6-8, 1871.

SMIRNOFF, G.: Om Katarr i mellersta örat. Akad. afhandl. Helsingfors, 1870. 8vo. 52 pp. Nord. med. Ark., iii. 1, Nr. 5, p. 27. 1871.

VOLTOLINI, R.: Die Krankheiten des Trommelfelles. Entzündung desselben. Mon.-Schr. f. Ohrenheilk., v. 2.

VOLTOLINI, R.: Die polypösen Wucherungen der Schleimhaut im Cavum pharyngo-nasale als Ursache der Schwerhörigkeit. Mon.-Schr. f. Ohrenheilk., v. 5.

WEBER, F. E.: (a) Bedeutende und dauernde Hörverbesserung durch Herstellung einer persistent gebliebenen Trommelfellöffnung—neue Methode—



rechterseits; Tenotomie des Tensor tympani mit günstigem Erfolg linkerseits. (b) Ueber die Perforation des Trommelfelles. Mon.-Schr. für Ohrenheilk., v. 2. [(a) A Case of Very Obstinate and Annoying Tinnitus Aurium cured by Division of the Tendon of the Tensor Tympani Muscle. (b) Perforation of the Membrana Tympani.]

WEBER, F. E.: Oleum Terebinthinæ bei akuten Mittelohraffektionen. Mon.-Schr. für Ohrenheilk., v. 3. Berlin, 1871.

WEBER, F. E.: Operation einer Atresie des linken Meatus audit. extern. durch Galvanokaustik. Mon.-Schr. für Ohrenheilk., v. 4.

WEBER, F. E.: Ueber die Natur der wichtigsten u. häufigsten Gehörkrankheiten. (Ges. d. Heilk. in Berlin.) Deutsche Klin., 19, p. 173.

ZAUFAL: Anwendung des sulphocarbolsauren Zinkoxydes bei Otorrhœe. (Verein d. deutschen Aerzte in Böhmen.) Wien. Med. Wochenschr., xxi. 18.

BUCK, A. H., M.D., of New York: Report on the Progress of Otology, 1870, 1871. Trans. Am. Otol. Society, 1870-71. Alfred Mudge & Son, 34 School Street, Boston, Mass.

BUCK, A. H., M.D., of New York: Case of Horny Growth from the Auricle. Op. cit.

LAIGHT, CHARLES, M.D., of New York: Necrosis of the Mastoid Cells in Immediate Neighborhood of External Auditory Meatus. Op. cit.

WEIR, R. F., M.D., of New York: A Singular Case of Nervous Deafness treated unsuccessfully by Galvanism. Op. cit.

WEIR, R. F., M.D., of New York: Tumor of the Cavity of the Tympanum with Otitis Media Serosa. Op. cit.

SEELY, W. W., M.D., of Cincinnati: Some Cases of Polypus and their Treatment. Op. cit.

BLAKE, C. J., of Boston: Parasitic Growths in the External Meatus. Op. cit.

MORLAND, W. W., M.D., of Boston: Case of Abscess of Cerebellum. Op. cit.

SPENCER, H. N., M.D., of St. Louis: A Case of Reproduction of the Membrana Tympani. Op. cit.

PARDEE, CHARLES J., M.D., of New York: On the Conclusions to be drawn from the Secretions in Chronic Purulent Inflammation. Op. cit.

POMEROY, O. D., M.D., of New York: Results of the Aural Examination of Sixty Lunatics. Op. cit.

POMEROY, O. D., M.D., of New York: Investigations on Inflation. Op. cit.

BERTHOLET, DR. R. M., of Philadelphia: Fibrous Tumor of the Auricle. Op. cit.

GREEN, J. ORNE, M.D., of Boston: Five Cases of Fatal Otorrhœa, with Remarks on the Course by which such Cases lead to Death. Op. cit.

ALLEN, PETER: Lectures on Aural Catarrh, or the Commonest Forms of Deafness and their Cure. London, 1871.

HUN, E. R., M.D.: Hæmatoma Auris. Albany. Am. Journ. of Insanity, July, 1870.

HARLAN, GEORGE C., M.D.: Catarrh of Tympanum from Use of Thudichum's Nasal Douche. Philadelphia Medical Times, Aug. 1, 1871.

KEEN, W. W., M.D.: Necrosis and Removal of the entire Petrous Portion of the Temporal Bone. Taste, smell, and hearing were destroyed, and none of these special senses returned or improved. Philadelphia Medical Times, June 15, 1871.

Schmidt's Jahrbucher. Leipzig, 1869, 1870, 1871.



# INDEX.

- Abscess, in cerebellum, 202.  
 cerebrum, 207.  
 external auditory canal, 160, 161.  
 temporal bone, 193-210.
- Acoustics, 43-48.
- Acuteness of hearing, 44, 45.
- Addison, Dr., of London, case of ulceration of auditory canal, 308.
- Adhesions in cavity of tympanum, 219, 220.
- Air-douche, Brunton's modification, 272, 273.  
 Politzer's modification, 273, 274.
- Air-press, 83.
- Allen's (London) modification of air-douche, 274.
- Alum, cause of furunculous abscess in catarrhal affections of the nasopharyngeal, 155.  
 in chronic purulent inflammation of tympanum, 299.  
 in diseases of mucous membrane, 229.  
 in treatment of polypus, 129.
- Ammonia, vapor of muriate of, in middle ear, 83, 84.  
 use, in affections of the Eustachian tube, 188.
- Anatomical description of the ear, 17-27.
- Anchylolysis of the stapes, 221, 222.
- Angioma of the auricle (Gruber), 147.
- Arnold, of Paris, on development of the inner ear, 324.
- Arteries supplying the cochlea, 27.
- Artery, stylo-mastoid, 23.
- Artillerists, rupture of membrana tympani, 278.
- Aspergillus flavescens, case of, 106, 107.  
 glaucus, in auditory canal, 104.  
 how to prevent its reproduction by treatment, 107.  
 Knapp, Prof. H., on its removal, 107.
- Astringents, in otitis media purulenta, 173, 174.  
 and styptics in treatment of polypus, 128.
- Astringents, in otitis externa, 166  
 objections to strong solutions of, 129.  
 use too early in cases of otorrhœa, 174.
- Auricle, anatomy of, 17, 18.  
 calcareous deposits in, 133.  
 case of chronic eczema of auricle (Wilde), 152.  
 contusions of, 132.  
 eczema, acute and chronic of, 150-153.  
 gout in, 133.  
 malformations, 15.  
 treatment of, 16.  
 physiognomic importance of, 19.  
 tumors of, 134.  
 wounds of, 440.
- Auricles for the deaf, 52.
- Auditory canal, anatomy of, 18.  
 exostoses of, 161.  
 glands of, 18.  
 length, 17.  
 lining membrane of, 18.  
 mode of examination of, 65-70.  
 narrowing of, 152.  
 physiology of, 18.
- Aural disease, importance of study of, 56.  
 surgery, 145, 146.
- Aural catarrh, acute, 211.  
 chronic, 214.  
 complications, 214.  
 pathology, 219-223.  
 prognosis and treatment (Gruber), 227.  
 subacute, 213.  
 treatment of acute, 227-238.  
 chronic, 24.
- Auscultation of ear, 71.
- Auto-laryngoscopic mirror and illuminating apparatus of Lewin, 252.
- Baierlacher (Paris), opinions of on galvanism, 321.
- Barrow, Mr. B. (London), case of tumor of lobule, 134.
- Baths, hot, and iron, 238.
- Bathing, effect upon the ears, 77.



- Battery of Grove, Daniell, Smee, 375.  
136.
- Biot (Paris) on transmission of sound, 44.
- Bird (Germany), hæmatoma auris, Bishop, Dr. (London), nebulizer, 275.
- Blisters in aural disease, 169.
- Bonafont on diagnosis of nervous deafness, 335.  
paracentesis of membrana tympani, 249.
- Bougies, Eustachian, 188.  
of laminaria, 188.
- Bowman, Wm. (London), opinions on nerves of the ear, 40.
- Boxing ears, results of, 75.
- Boyer, Baron (Paris), on polypus, 110.
- Brenner's eleven cases of disease of the ear treated by galvanic current, 389.  
observations in regard to the effect of electricity upon the organs of hearing, 379, 381, 383.
- Bristles of Max Schultze, 41.
- Bruch, (Germany), tumors of lobe, 135.
- Brunton, Dr. (London), modification of Politzer's air-douche, 273.
- Bullet in the ear, 94.
- Camp, Dr., case of hereditary deafness, 316.
- Cancer of the auricle, 140.  
cases of, 141.
- Carbolic acid in otorrhœa, 174.
- Caries of petrous portion of temporal bone, 207, 208.  
temporal bone, fatal results from, 205-209.
- Carious teeth as a cause of deafness, with cases, 307, 308.
- Cassebohm, Dr. (London), case of calcareous deposit, 286.
- Cases of abscess in cerebellum, 204.  
cerebrum, 207.  
aural catarrh, 211-218.  
bony tumor of auditory canal, 161.  
chronic catarrh, 341.  
concussion causing nervous deafness, 338.  
gout (nervous deafness), 345.  
hemorrhage from external ear, 114-196.  
inspissated cerumen, 91.  
nervous deafness (concussion), 338.  
nervous deafness caused by sulphate of quinia, 350.
- Cases of post-aural inflammation, 142.  
removal of sequestrum from mastoid process, 193.  
rheumatic fever, 349.  
syphilitic deafness, 343.  
tinnitus aurium, 78.  
use of artificial membrana tympani, 293-299.
- Casper, Dr. (Germany), case of feigned deafness, 310.
- Catarrh, acute aural, 211.  
catheter and air-douche in, 231.  
chronic, 217.  
definition of, 211,  
middle ear with perforation, 443.  
mineral springs in, 237.  
naso-pharyngeal, 229.  
pathology of, 219-223.  
prognosis of, 227.  
subacute, 213.  
tubal, 197-229.  
treatment of, 228-240.  
use of astringent fluids and vapors in, 233-235.  
with or without perforation of membrana tympani, 443.
- Catheter, Eustachian, author's, 271-274.  
and explorer, 269.  
Dr. Cutter's, 266.  
Dr. Kramer's, 266.  
Dr. Noyes's, 276.  
form and use, 266.
- Cauterization, 126.
- Cavity of tympanum, deficiency in floor of, 19.  
complicated inflammation of, 211.  
chronic inflammation of, 211.  
mucous membrane of, 20.  
physiology of, 19.  
relations of, 21.  
section of, 28.  
size, anatomy of, 19.  
subacute inflammation of, 211.
- Cerumen, appearance and form of, 91,  
caution in the use of end of towel, 76.  
diminution of, 9.  
inspissated, 91-93.  
treatment of, 441.  
removal of accumulations of, 86, 92.  
solution to soften, 91.
- Cerebrum, abscess of, 207.  
disease in, caused by aural affections, 206, 207.
- Chauliac, Guy de (Paris), on polypus, 111.
- Children, care of their ears, 75.  
injurious habits of, 76.



- Chills in purulent otitis media, 181.  
their treatment in otitis media, 181.
- Cholesterine in middle ear, 184.
- Chorda tympani nerve, 321.
- Chronic catarrh, table of cases, 226.
- Clamp for holding Eustachian catheter, 260.
- Clark (Prof. E. H., Boston), aural douche, 88.  
cases of deafness, 58.  
perforation of membrana tympani, 290.  
modification of Thudichum's (Weber's) nasal douche, 87.  
polypi, 118-129.
- Cleansing the ear, 76.
- Cleland, A., method of examining the ear, 65.
- Climate, influence of, upon deaf mutes, 406.
- Cleft palate, 271-273.  
importance of operating early for closure, 273.
- Cochlea, anatomy of, 25.  
canal of, 33.  
function of, 32.  
microscopical investigations of, 33-42.
- Cohen, Dr. J. S. (Phila.), galvanic cautery, 127.
- Cold, effects of upon the ear, 75-77.  
bathing, 77.  
deafness from, 75-77.
- College for deaf mutes, 401.
- Columbia Institution for Deaf Mutes, 405.
- Compressed sponge in dilating meatus, XVI.
- Compressed air-pump, 83.
- Counter-irritation, 178.
- Concha (see Auricle).
- Concussions, violent, of air, precautions, 77.
- Conversation-tubes, 51.
- Corti, Marchese, organ of, 33, 35, 40.  
rods, fibres, etc. of, 32.
- Coughing produced by syringing the ears, 92.
- Curette of Prof. Gross, 95.  
silver, 96.
- Curwin, Dr. John (Harrisburg), case of hæmatoma auris, 138.
- Outter, D. E. (Boston), Eustachian catheter, 266.
- Czermak (Berlin), rhinoscopy, 264.
- Day-school for deaf mutes, Pittsburg, 405.
- Deaf mutes, Abbé de l'Epée's efforts to teach them by signs, 394  
artificial method of Heinicke, 396, 397.  
definition and history of, 393.  
Dr. Cogswell's daughter, Hartford, Conn., 397.  
influence of climate, etc., 406.  
Pennsylvania Deaf and Dumb Institution, 400, 401.  
Rotterdam Institution for, 433.  
Sicard and his success, 395.  
treatment at home, 410.  
at school, 408.  
medical, 409.  
prophylactic, 410.  
examination by laryngoscope of, 408, 409.  
first practical attempt to teach, 394.  
list of institutions in the United States, 404.  
number of, in Pennsylvania and United States, 395-400.\*
- Deafness caused by adhesions in cavity of tympanum, 221.  
bathing in cold water, 75-78.  
boxing the ears, 76.  
caused by carious teeth, 307, 308.  
causes of, 55, 56.  
congenital, 303.  
definition of, 153.  
disease of glands, 142.  
eczema of auditory canal, 150-153.  
foreign bodies in the ears, 93.  
from aural catarrh; acute, sub-acute, and chronic, 201-214, 217.  
from impacted cerumen, 91.  
from rupture of the membrana tympani, 278, 279.  
from tumors in the auditory canal, 161.  
furunculous abscess a cause of, 101.  
injuries of the head, 447.  
narrowing of auditory canal, 152.  
nervous, 316.  
treatment by mechanical means, 49-54.
- Deglutition, effect on pharyngeal orifice of Eustachian tube, 30.
- Deiter's cells of the minute structure of the cochlea, 40.

\* Having omitted in the body of the work to mention the number of deaf mutes in the United States, we here extract it from the advanced sheets of the Census. Total number, 16,258. Of these, 8916 are males, and 5289 are females.



- Dentition, causing deafness and otorrhœa, 164.
- Diagnosis, of the various forms of disease of the ear, 65.  
by sunlight and artificial light, 66-70.  
importance of, 172.  
means used in, 66-70.
- Diagnostic tube, 72.
- Dieffenbach (Berlin), opinions on deafness in cleft palate, 273.
- Diseases of the ear, the result of constitutional disorders, 55.  
cold, etc., 55.  
in the male and female, 55.  
method of investigating, 60, 61.  
table of, 56-59.  
treatment of malignant, 143  
the mastoid and pre-auricular region, 142.
- Dissections, showing conditions of ear in deaf and dumb, 412-419.
- Double hearing, 331.
- Double tube Eustachian spray catheter, 275.
- Douche, air, Politzer's description and mode of use, 273.  
Clark's aural, 81.  
modifications, 274.  
risks in using nasal, 129.  
Thudichum's nasal, 88.  
used with catheter, 273.
- Draughts of cold air on the ear, 75.
- Duchenne (Paris) on nervous structure of the ear, 321.
- Duchenne's (Paris) rheophore, 378.
- Ductus cochlearis, 40, 41.
- Dugès (Paris) on the cochlea, 32.
- Dulong (Paris), experiments of velocity of sounds in gases, 47.
- Dunlop, Dr., case of feigned deafness, 309.
- Duval (M. Emile, Paris) on earache, 309.
- Ear, ache or neuralgic pain in the, 308, 309.  
anatomy and physiology of, 17-43.  
diagnosis of diseases of, 64.  
drops, 309.  
east winds, effect upon the, 77.  
external mode of examining, 61.  
general care of, 75.  
itching of, 76-116.  
rings, effect upon, 136.  
scissors, 126.  
spout of Toynbee, 90.  
table of diseases of, 56.  
trumpets, 49-54.
- Eczema auriculæ, 441.
- Eczema auriculæ, chronic, 151  
auditory canal, 150.  
of auricle, 150.  
treatment of, 151-153.
- Effusion of serum into tympanic cavity, 445.  
blood upon membrana tympani, 445.
- Electricity, electro-magnetic, 373-380.  
galvanic, 380.  
in nervous deafness, 373.  
in tinnitus aurium, 386-389.  
magneto-electric, 377.
- Entacoustics, Dr. Jago on, 79, 80.
- Epithelial cancer of the auricle, 145-147.
- Epithelial polypus of Clarke, 113.  
vascular, Triquet, 113.
- Erdmann (Paris), opinions of galvanic electricity, 321.
- Erhard on cotton as an artificial drum, 294.  
diagnosis of nervous deafness, 335.
- Eustachian tube, acute inflammation of, 217.  
anatomy of, 20.  
calibre, 21.  
catheter, 266, 267.  
catheterism, 269.  
catheter introduced, 268.  
catheter, use of, 269.  
chronic inflammation of, 214.  
closure of, 224-337.  
dilatation of, 224.  
explorer of, with catheter attached, 269.  
impervious, 224, 269, 270.  
mechanical appliances for, 251.  
method of examining, 71-73  
mode of introducing catheter into, 268, 269.  
mucous membrane of, 20.  
muscles of, 21.  
papillæ of, 22.  
pathology of, 224.  
pharyngeal openings of, 271.  
physiology of, 29.  
section of, 28.  
statistics of disease of, 56.  
structure of, 20, 21.
- Examination of the ear described, 68.
- Exanthematous otitis, 303.
- Exostosis in auditory canal, xvi. 161-163.
- External auditory canal, inflammation of, 165.
- Fabricius Hildenis, case of polypus, 111.



- Facial nerve, acute inflammation of, 361.  
 Facial paralysis, 102.  
 Fallopian canal, 26.  
 Faradisation of the ear, 376.  
 Fatal results from caries of temporal bone, 209.  
 Feigned deafness, 309.  
 Feigned dumbness and deaf-dumbness, 310-315.  
 Fenestra ovalis, 26.  
 Fenestra rotunda, 26.  
 Ferrus (France), hæmatoma auris, 136.  
 Fever, deafness attending and consequent upon, dumbness following, 412.  
 Fick, M. M., fibro-plastic polypi, 127. extraction of, 123. investigations of Corti's organ, 34.  
 Fluids in the external auditory canal, 77.  
 Fluids injected into middle ear, 274.  
 Foramen of Rivinius, 218.  
 Forceps, angular, "Wilde," 160. for removing polypi and foreign bodies, 126.  
 Foreign bodies, in external canal, 440. in Eustachian tube, 93, 94. list of, 94.  
 Fossa of Rosenmüller, 262.  
 Fracture of malleus, 279. annular cartilage, 131.  
 French hearing-trumpets, 54.  
 Fright, a cause of deafness and deaf-dumbness, 318.  
 Frost-bites of external ear, 133.  
 Function of Eustachian tube, 29.  
 Fungous growths of ear, 104-107. malignant, of ear, 142, 143.  
 Furunculous abscesses of the auditory canal, 153. with cases and treatment, 161. in external meatus, 160.  
 Gallaudet, Edward M., account of deaf and dumb institutions in Europe, 433.  
 Galvanic cautery, 127.  
 Galvanic electricity, Remak, in diseases of the ear, 380.  
 Gangrene of ear, 440.  
 Gargling in ear diseases, 228, 301.  
 Garrod, Dr. (London), gouty inflammation of auricle, 133.  
 Gases from stomach, introduction into tympanum, 30.  
 Glands, post-aural, diseases of, 142.  
 Glosso-pharyngeal nerve, 23.  
 Glycerine in discharge from ear, 116-119. and tannic acid, 108.  
 Goodsir, T. (Edinburgh), on the vestibule, 32.  
 Gouty inflammation of external ear, 133.  
 Granular membrana tympani, 175.  
 Growths, morbid, in meatus, 118. in tympanum, 119. treatment of, 175. tumors in auditory canal, 161.  
 Gruber, Prof., of Vienna, 176. on chronic catarrh of middle ear, 227. otitis media purulenta, 176. pruritus, 148. its prognosis and treatment, 148-153. treatment of chronic catarrh of middle ear, 228. tables of diseases of the ear, 59.  
 Guersant, M., on removal of tonsils, 303.  
 Guy, Dr. (London), case of feigned deafness, 310.  
 Hæmatoma auris, 136, 137.  
 Hearing, certain peculiarities of, 336. double, 331.  
 Helmholtz, Prof., physiology of hearing, 29.  
 Hemorrhage from ears, 116, 196. from tonsils, 304.  
 Henle on the fibres of Corti and lamina reticularis, 38.  
 Hensen's microscopic researches, 42.  
 Hereditary tendency to deafness from aural disease, 55. to deaf-dumbness, 55.  
 Herschel, Sir John, on influence of sound on different persons, 45.  
 Hewson, Dr. A. (Phila.), case of feigned mutism, 312.  
 Hildenis, Fabricius de (Paris), on polypus, 111.  
 Hinton, Prof., of London, modification of Toynbee's speculum, 67. nervous deafness, cases of, 338. on syphilitic deafness, 372. treatment of polypi, 123. use of Siegel's pneumatic speculum, 54, 84, 85.  
 Hippocrates, Celsus, Galen, Ætius, Paulus Ægineta, opinions upon polypus, 111.  
 Holberton, Mr. (London), tumors of lobule, 134.  
 Holmes, Dr. (London), on cauterization of enlarged tonsils, 306.



- Hun, Dr. E. R. (New York), hæmatoma auris, 137.  
 Hysterical deafness, 354.
- Idiotic deaf and dumb, 406.  
 Illuminating apparatus, Lewin and Tobold's, 252, 253.  
 Imposture of deaf and dumb, 310.  
   detection of, 311.  
 Incisions, post-aural mode of performing, 196.  
 Incus, description of, 22.  
 Induced electric current to recover voice, 314.  
 Infantile aural catarrh, 167.  
 Inflammation of auricles in the insane, 136-140.  
   auditory canal, 163.  
   Eustachian tube, 214-231.  
   of brain in acute otitis, 327-330.  
   internal ear, 447-450.  
   mastoid cells, 189, 191-200.  
   membrana tympani, 280-283.  
 Injection of astringents into Eustachian tube, 181.  
   nose, 233.  
   substance of polypi, 128.  
   tympanum, 233.  
 Injuries of external ear, 131, 132.  
   labyrinth, 363.  
   membrana tympani, 278.  
   tympanum, 440.  
 Insects in the ear, mode of removal, 77.  
 Inspissated cerumen, 441.  
 Institutions in United States for deaf and dumb, 400, 401, 404, 405.  
 Instruction of deaf mutes at home, 422.  
   in school, 420.  
 Instruments for incising membrana tympani, 242.  
   to assist hearing, 49.  
 Insufflator for powders, 82.  
 Intermarriage as a cause of deaf mutism, 407.  
 Internal ear, anatomy of, 23.  
   diseases of, 201-447.  
   treatment of, 450.  
 Itard (Germany), auricles of, 523.  
 Itching of the meatus, 76.
- Jackson, Prof. Samuel, function of semicircular canals, 32.  
 Jacobson's nerve, anastomosis, 320, 321.  
 Jago, Dr., functions of the tympanum and Eustachian tube, 29-31.  
 Jugular vein, contiguity to cavity of tympanum, 224.  
   ulceration of, 224.
- Kidder's electro-magnetic apparatus, 375, 376.  
 Kirkbride, Dr. Thomas (Phila.), cases of hæmatoma auris, 138.  
 Knapp, Prof. H., a general summary of inflammatory affections of the inner ear, 447-451.  
   case of otitis media purulenta, 168.  
   double hearing, 331.  
   opinions concerning the use of nasal douche, 81.  
   purulent inflammation of middle ear, 168.  
 Koenig's acoustic trumpet and stethoscope, 54.  
 Kölliker, Prof. (Würzburg), on the physiology, etc. of cochlea, 32, 33, 34, 41.  
 Kramer's bivalve aural speculum, 68.
- Labyrinth, acute inflammation of, 363.  
   classification of its diseases, with treatment, 325, 450.  
   chronic, 364.  
   osseous and membranous, 25.  
 Laennec on aural auscultation, 71.  
 Laminaria, bougies of, 249.  
 Lamps, aural, 66-68.  
 Larva in the ear, 108.  
 Laryngoscopic examination of deaf mutes, 407.  
 Larynx, irritation of, from touching polypi, 118.  
 Laycock, Dr. (Edinburgh), case of hæmatoma auris, 137.  
 Leeches, mode of applying, 172.  
 Leschevin (Paris), fungus in meatus auditorius, 112.  
 Lewis, Dr. (Boston), aspergillus, 106.  
 Light, artificial, 66.  
 Light spot, 19.  
 Lip reading by the deaf and dumb, 427-437.  
 Lobe, fibrous tumor of, 134-136.  
 London paste for removal of tonsils, 307.  
 Lucae on conduction of sound by the bones of the head, 74.  
   on a new method of examining the ear, 74, 75.
- Mackenzie, Dr. (London), modification of galvano-cautery, 127.  
   vocal organs of the deaf and dumb, 426.  
 Maggots in the ear, and their removal, 108, 109.



- Magneto-electric machine of Neff, Sexton, and Page, 377.
- Malformations of the auricle, 440.
- Malgaigne, Prof. (Paris), on polypus, 111.
- Malignant disease of the auricle, 143, 144.
- Malleus, description of, 22.  
fracture of, 279.
- Manometer, aural, 72.
- Marchese Corti, his discovery, 41.
- Markusovszki, Dr. (Poland), speculum, 264.
- Martineau, Harriet, ear-trumpet, 51.
- Mastoid process and cells, disease of, 190-200.
- Mastoid process or cells in child and adult, 189.  
in chronic aural catarrh, 189, 190.  
operations upon diseased, 193-199.
- Mayor, Dr. (Paris), handkerchief dressing, 131.
- Meatus externus, anatomy of, 17, 18.  
classification of its diseases, 62.  
malformations of, 15, 16.  
diseases of, 131. (See Auditory canal.)
- Membrana tympani, anatomy and physiology, 18-31.  
artificial and substitute for, 293-299.  
cases of perforation, 291.  
collapse of, 282.  
diseased, 286-288.  
ecchymosis of, 284.  
illumination, mode of, 66-69.  
inflammation, acute and chronic, 280.  
injuries of, 278.  
loss of, 293.  
microscopical section, 285.  
movements, 31.  
papillæ and inflammation of, 22, 300.  
paracentesis, 238-242.  
pathology of acute aural catarrh, 285, 286.  
perforation, 287-289, 446.  
treatment of, 291-299, 446.  
Poltzer on color of, 19, 284.  
Rupture of, 278.
- Ménière on fracture of handle of malleus, 279.
- Meyer, Dr., electricity applied in deafness, 384.
- Middeldorf (Breslau), galvano-cautery, 127.
- Middle ear, anatomy and physiology of, 19.  
mechanical applications for, 251.  
table of diseases of, 59.
- Mineral springs in chronic catarrh, 237.
- Mirror, concave, for examining the ear, 70.
- Moos, Prof. S. (Heidelberg), air-press, 82.  
cases of chronic aural catarrh, 58.  
double hearing, 331.  
nervous deafness, 387.  
table from, 58, 226.
- Mucous cushion in middle ear, 20.
- Müller, Johannes, opinions of, 302.
- Muscles, levator palati, 21.  
stapedius, 22.  
tensor tympani, 21.
- Muscles of Eustachian tube, 21.
- Mussy, Dr. (Cincinnati), tonsil instrument, 306.
- Myringitis, 289.
- Myringo-dectomy, 249.
- Nares, syringe for, 82.
- Naso-pharyngeal mucous membrane, 229, 263.  
catarrh of, 229.
- Nasal clamp, 260.
- Nasal douche, objection to use, 81.  
modifications of, 83.  
objections to its use, 81.  
Thudichum's, 129.
- Nasiloff, inflammation of membrana tympani, 300.
- Nebulizer, nasal, 84.  
Bishop's, 275.  
steam, 82.
- Neftel, Dr., cases treated by the galvanic current, 385-387.
- Nerves of middle ear, facial, 23.
- Nervous deafness, treatment, 316, 368.  
electricity in, 373.
- Neuralgia of plexus tympani, 445.
- Neuralgic pain in ear from catarrh, 308.
- Nitrate of silver in furuncles, 160.  
perforation of membrana tympani, 118.
- Noises in the ear, or tinnitus aurium, 78.  
treatment by electricity, 385-390.
- Norris, Dr. G. W. (Phila.), tumors of lobe, 135.
- Noyes, Dr. Henry D., modification of Eustachian catheter, 276.
- Opacities in the membrana tympani, 284-288.
- Osseous tumors in auditory canal, cases, treatment, etc., 161-163.
- Ossicula auditus, 22.  
mode of action, 29.
- Otalgia, 308.



- Otaphone, 53. (See *Hæmatoma auris*.)  
 Othatomata, 136.  
 Otitis, hypertrophic, Gruber, 182-188.  
   externa, 163, 442.  
   diffuse, 163.  
   interna, case, with post-mortem, 201.  
   with inflammation of mastoid cells, 444.  
   prognosis, 206.  
   treatment, 450.  
   media purulenta, 167-172.  
   and interna, 189.  
   chronica, 173.  
   subdivision of, 449.  
   syphilitic, 445.  
 Otorrhœa, description of, 167-188.  
   description of by Gruber, 176-182.  
   discharge, 130.  
   prognosis, 170.  
   results if neglected, 209.  
   report of fatal cases from Von Trötsch, 206, 207.  
   treatment of chills and fever following, 181.  
   with polypus, 129.  
 Otoscope, 71.  
   double, 73.  
   of Toynbee, 270.
- Paget, Mr. (London), Lectures on Surgical Pathology, 134.  
 Pain in the ear, 77, 78, 308, 309.  
 Palate noose, Türcks, 258.  
   hook, 258.  
     directions how to use, 258-280.  
   Moos' cases of, 244, 245.  
 Paracentesis of the membrana tympani, 179.  
   Hinton's cases of, 241.  
 Paré, Ambrose (Paris), on polypus, 111.  
 Pelletan (Paris) on fungus in meatus, 112.  
 Perforation of the membrana tympani, 289.  
   prognosis and treatment, 290, 446.  
 Periostitis of external auditory canal, 207, 208.  
   mastoid process, 193.  
   with cerebral abscess, 445.  
   by rhinoscopy, 252.  
 Pharyngeal orifice of Eustachian tube, 261.  
 Pharyngitis, acute, chronic, and granular, 301.  
 Pharynx, cauterization of, 301.  
   examination of, 263-266.
- Phlebitis from aural disease, 207.  
 Physiology of hearing, 28-42.  
 Pincette for introducing the eyelets into membrana tympani, 250.  
 Politzer's air-douche, and its modifications, 273.  
   hard rubber eyelet, 249, 250.  
   method of inflating ear, 273.  
   on chronic catarrh, 212.  
   on size and color of membrana tympani, 71.  
   on spasmodic contraction of tensor palati, 80.  
 Polypi, aural, 110, 111.  
   fibro-gelatinous, 116.  
   large pulsating, 117.  
   nasal, 110.  
   removal of, 112.  
   soft gelatinous, 115.  
 Posterior nasal syringe, 83, 84.  
 Post-mortem examination of the ear, 205-208.  
   inner ear, 413-416.  
 Poulticing the diseased ear, 193.  
 Prevention of deafness, 75.  
 Probing Eustachian tube, 188.  
   the ear, 76.  
 Promontory, description of, 25, 26  
 Pruritus of external auditory canal, 148-150.  
 Pyæmia, cause of death in otorrhœa, 209.
- Quinine, nervous deafness from use of, 350.  
   use in chills accompanying otitis, 181.
- Radau, R. (Paris), *L'Acoustique*, 54.  
 Reflected light (artificial), 253, 254.  
 Reflex irritation from foreign bodies in the ear, 80.  
 Résumé of the simplest and most effectual method of treating the more frequent diseases of the ear, 438, 439.  
 Rheophore of Duchenne, 378.  
 Rheostat, Brenner's, 383.  
 Rhinoscopic image, 262.  
   mirrors, 255-257.  
     how to use them, 254, 255.  
 Rhinoscopy, 250-266.  
 Richardson's spray apparatus, 274.  
 Ring-and-lever forceps (of Toynbee), 125.  
 Rods of Corti, number and use of, 41.  
 Roosa, Prof. St. John, table of five hundred cases of diseases of ear, 57.



- Rosenmüller's fossa, 262-268.  
 Rudinger on the muscles of the Eustachian tube, 21.
- Scarlet fever as a cause of deafness, 117.
- Schultze, Max, bristles of, 41.
- Schwartz, table of diseases of the ear, 58.  
     fungous growths in the ear, 105.  
     on paracentesis of membrana tympani, 238.
- Scrofulous diseases of the ear, 130, 297.
- Scultet (Paris), fungous growth in meatus, 112.
- Sebaceous glands in auricle, 18.  
     meatus, 18.
- Sedgwick, Dr., influence of sex on hereditary deafness, 316.
- Sédillot (Paris), fungous growths in meatus, 112.
- Semeleder, mode of using the rhinoscopic mirror, 255.
- Semicircular canals, description of, 24.  
     Flourens's experiments, 32.
- Semioc, Dr. (New York), instrument, 260.
- Sequestrum, removal from internal ear, 192.
- Serum in tympanic cavity, 242, 243.
- Sissay, Dr. (Lyons), on Diseases of the Ear, 133.
- Smith, Prof. N. R. (Baltimore), trans. of Sissay on Ear Diseases, 133.
- Somers, Dr. S. L. (Phila.), fibro-gelatinous polypus, 116.
- Sound, laws of, 43.
- Sounds in the middle ear, 78-81.  
     velocity of, in gases, liquids, and solids, 47, 48.
- Specula, aural, 68.  
     aural, in tinnitus, 84, 85.
- Speculum, Thudichum's, 265.
- Sphyrotomy, mode of operation, 249.
- Spray catheter of Codman and Shurtleff, 274.  
     double, 274, 275.
- Stillé, Prof. Alfred (Phila.), tumor of auricle, Proc. Pathological Society of Phila., 134.
- Sunlight in examining the ear and membrana tympani, 68.
- Suppuration of the middle ear, 167.
- Synechia in cavity of tympanum, 220.
- Syphilis, a cause of nervous deafness, 343, 371, 449.
- Syringes, various forms of, 87.
- Syringing the ear for the removal of cerumen and foreign bodies, 86-92.  
     the ear producing cough, 92.
- Tables of disease of the ear by Dr. Gruber, 56.  
     Dr. Roosa, 57.  
     Dr. Turnbull, 59.  
     of the institutions for the instruction of the deaf and dumb in the United States, 404.  
     of the transmission of sound through solids, liquids, and gases, 47, 48.
- Tabular view of conditions of the ear in deaf and dumb, 416-418.
- Taylor, Dr. Alfred (London), on rupture of membrana tympani by strangulation, 278.
- Teaching the deaf mute, on the choice of methods, 426.
- Tensor tympani (see muscles of the ear), 21.  
     palati, 21.
- Thomas, Dr. R. P. (Phila.), case of maggot in meatus, 108.
- Tinnitus aurium, 79-87.
- Tobold's illuminating apparatus, 253.
- Todd, R. B. (London), on the physiology of the ear, 40.
- Tones, power of hearing various, 46.
- Tongue depressor, 260.
- Tonsils, enlarged, as a cause of deafness, 302.  
     method of removal, 303.  
     by caustics, 306, 307.  
     Prof. Gross on hemorrhage from, 304.
- Toynbee on function of Eustachian tube, 29.
- Toynbee's artificial membrana tympani, 295.  
     ear-spout, 90.  
     illuminating speculum, 67.
- Triquet (Paris), cases of diseases of the ear, 55.  
     nervous deafness, 330.  
     polypus of the ear, 110.
- Tuberculosis with otorrhœa, 287.
- Tumors of the lobe, 134.
- Tuning-fork in diagnosis of nervous deafness, 48, 337.  
     modification of Lucae's double, 74.
- Türck, Dr. (Vienna), palate noose, 258.
- Turnbull, Dr., table of six hundred cases of disease of the ear, 56.



- Tympanum, chronic purulent inflammation with perforation, 299.
- Tyndall, Prof. (England), on sound, 41.
- Typhoid fever cause of deafness, 342.
- Umbo (see *Membrana tympani*), 19.
- Uvula, elongation and oblique position of, 301.
- Valsalvian mode of opening Eustachian tube, 72.  
experiment, 72.
- Vapors, injection of, into middle ear, 235.
- Vapors of chloride of ammonium, 83, 84.
- Vascular polypus of left meatus, 113.  
at base of external meatus, 114.
- Venous sinus, inflammation of, 210.
- Venzetta (Italy) on tumors of the lobe, 135.
- Vertigo in aural disease, 203, 448.
- Vibration of air in tubes, 48.
- Virchow's cases of dissections of the ears of deaf mutes, 415.
- Voltolini on inflammation of labyrinth, 449.
- Voltolini's cases of dissections of the ears of deaf mutes, 415, 416.
- Vomiting in disease of the labyrinth, 448.
- Von Baer (Paris) on development of Eustachian tube, 324.
- Von Tröltsch, Prof. (Würzburg), on the form of tympanum, 19.  
analogous relations of eye and ear in their mode of development, 324, 325.  
calcareous deposits in *membrana tympani*, 286.  
diagnosis of nervous deafness, 335.  
fatal otorrhœa, 206.  
furuncular inflammation, 157, 158.  
medical treatment of a deaf mute, 411.  
mucous bubbles in the tympanic cavity, 243.  
paracentesis of *membrana tympani* in catarrh of tympanum, 238.
- Von Tröltsch, Prof. (Würzburg), on pathology of aural catarrh, 225.  
perforation of the mastoid process, 197, 198.  
treatment of nervous deafness, 368.
- Wagner, Dr. (Germany), nose clamp, 260.
- Wales, Dr. P., U.S.N. (Phila.), palate hook, 259.
- Warm water, its use in aural disease, 81-90.
- Washing the ear, how it should be done, 71.  
in syringing the ear, 92, 93.
- Water douche to Eustachian tube, 81.
- Weber, aural reflector, 67.
- Wertheim (Germany), experiments on transmission of sound through liquids, metals, and wood, 47.
- Wetting the hair a cause of deafness, 77.
- Wilde, Sir William, ear basin, 91.  
collapse of *membrana tympani*, 19.  
modification of Gruber's speculum, 69.  
polypus snare, 124.  
on indiscriminate syringing, 92.  
on light spot on drum, 19.  
thickening of *membrana tympani*, 19.
- Wilks, Dr. (London), acute otitis, with cerebral disturbance, 326, 327.
- Wollaston, Dr. (England), on acuteness of hearing, limits of hearing, closing the Eustachian tube, 45, 46.
- Wounds and injuries of the *membrana tympani*, 278, 279.  
of the auricle, 131.
- Wreden, Dr., on exsection of the handle of the malleus, 249.  
fungous growths in ear, 105.
- Yearsley on artificial *membrana tympani*, 293, 294.  
removal of tonsils, 313.
- Zona pectinata, 40.



# IMPORTANT MEDICAL WORKS

RECENTLY PUBLISHED BY

J. B. LIPPINCOTT & CO.,

713 and 717 MARKET STREET, Philadelphia.

## *A New and Revised Edition of the United States*

Dispensatory. The DISPENSATORY OF THE UNITED STATES OF AMERICA. By GEORGE B. WOOD, M.D., and FRANKLIN BACHE, M.D. Thirteenth Edition. *Carefully Revised and Corrected, with Numerous Important Additions.* One Volume, Royal 8vo. Superfine paper. Sheep. \$10.00.

## *Hand-Book of Operative Surgery.* By John H.

PACKARD, M.D., Sec. of the Coll. of Physicians of Phila., etc. With Fifty-four Steel Plates, and Numerous Illustrations on Wood. 8vo. Extra cloth. \$5.00.

"Its distinguishing feature is abundant illustration, and in this respect it is unequaled by any hand-book with which we are acquainted. . . . To students and practitioners who have not the fre-

quent opportunity of witnessing practical surgery, the volume before us will prove of unquestionable value."—*N. Y. Med. Record.*

## *Medical Diagnosis with Special Reference to Prac-*

tical Medicine. A Guide to the Knowledge and Discrimination of Diseases. By J. M. DA COSTA, M.D. Illustrated with Engravings on Wood. *Third Edition, Revised and Enlarged.* 8vo. Fine cloth, \$6.00; sheep, \$7 00.

"To digest and memorize the matter of all its pages will well repay the labor it costs. It may be welcomed as a text-

book which will do credit to our national medical literature."—*Amer. Jour. of Med. Sciences.*

## *A Compend of Materia Medica and Therapeutics.*

For the Use of Students. By JOHN C. RILEY, A.M., M.D., Prof. of Mat. Med. and Therap. in the Nat. Med. Coll., etc. 8vo. Extra cloth. \$3.00.

"Prof. Riley has done his work very well. . . . The descriptions of the various articles of the Materia Medica

are very concise and remarkably accurate."—*N. Y. Med. Journal.*



PUBLICATIONS OF J. B. LIPPINCOTT & CO.

*A Handbook of Medical Microscopy.* By Joseph

G. RICHARDSON, M.D., Microscopist to the Pennsylvania Hospital, Philadelphia. 12mo. Fine cloth. \$2.25.

This work is designed to supplement by minuteness of detail and accuracy of illustration the practical instruction so generally necessary for an advantageous use of the instrument as an aid to the diagnosis of disease. It should be in the hands of every medical man who either possesses a microscope or contemplates purchasing one, since, in the latter case, it will often save many times its own cost by advice

in regard to the selection of apparatus; while the advanced student and practitioner of medicine will find in it brief but precise directions for the various kinds of research, according to the most recent discoveries, as well as for the avoidance of those errors of interpretation which are sometimes so mortifying to the inexperienced observer.

*Rand's Medical Chemistry. A New Edition of*

Elements of Medical Chemistry. By B. HOWARD RAND, M.D., Professor of Chemistry in the Jefferson Medical College. Carefully Revised, with Additions. Illustrated. 12mo. Extra cloth, \$2.00; sheep, \$2.50.

"We commend it to the student as the best *vade-mecum* we have yet seen. We shall also adopt it as a text-book in our

College of Pharmacy."—*St. Louis Med. Reporter.*

"Prof. Rand has long held, as he has deserved, a high position as a teacher of Chemistry. The simplicity, clearness, and absence of all superfluity, which characterize his lecture-room demonstrations, are seen in this book from the beginning to the end. We would especially remark the due apportionment of space

to its different subjects. . . . Medical students may feel themselves greatly indebted to Prof. Rand for furnishing them with a text-book so portable, convenient, and concise, and yet so lucid and full, as to the really essential elements of chemistry."—*Medical and Surgical Reporter, Philadelphia.*

*Opium and the Opium-Appetite. With Notices of*

Alcoholic Beverages, Cannabis Indica, Tobacco, Coca, Tea, and Coffee, in their Hygienic Aspects and Pathologic Relations. By ALONZO CALKINS, M.D. 12mo. Extra cloth. \$1.75.

This volume forms an elaborate treatise respecting the character and use of stimulants. Its subject is one that is daily

assuming greater importance, and in which all classes of thoughtful readers are more or less interested.

*Microscopical Manipulation, being the Subject-*

matter of a Course of Lectures delivered before the Quekett Microscopical Club, *January—April, 1869.* By W. T. SUFFOLK, F.R.M.S. Illustrated with Forty-nine Engravings and Seven Lithographs. Crown 8vo. Fine cloth. \$2.00.

"Mr. Suffolk gives an explanation of the construction of the microscope, and clear, practical directions for preparing slides and other apparatus, mounting and

preserving objects, and conducting a great variety of experiments and researches. The book is well illustrated."—*N. York Tribune.*







