Essays on the morbid anatomy of the human eye / by James Wardrop ...; illustrated by plates.

Contributors

Wardrop, James, 1782-1869. Francis A. Countway Library of Medicine

Publication/Creation

Edinburgh: Printed by George Ramsay and Co., for Archibald Constable and Co., Edinburgh; London: John Murray, Fleet-Street, 1808-1818.

Persistent URL

https://wellcomecollection.org/works/rwev6afa


License and attribution

This material has been provided by This material has been provided by the Francis A. Countway Library of Medicine, through the Medical Heritage Library. The original may be consulted at the Francis A. Countway Library of Medicine, Harvard Medical School. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



28.9.33


Add.
De Waeloworth

Digitized by the Internet Archive in 2011 with funding from Open Knowledge Commons and Harvard Medical School









ESSAYS

ON

uold

THE MORBID ANATOMY

OF THE

HUMAN EYE.

BY JAMES WARDROP,

OF THE ROYAL MEDICAL AND CHIRURGICAL SOCIETIES, AND ONE OF THE SURGEONS OF THE PUBLIC DISPENSARY OF EDINBURGH.

ILLUSTRATED BY PLATES.

EDINBURGH:


Printed by George Ramsay and Company,

FOR ARCHIBALD CONSTABLE AND COMPANY, EDINBURGH;

AND JOHN MURRAY, FLEET STREET,

LONDON.

1808.



10777

adding real County was a

RORE

ANDREW WARDROP, M. D.

F. R. S.

FELLOW OF THE ROYAL COLLEGE OF SURGEONS, AND FORMERLY
ONE OF THE SURGEONS OF THE ROYAL INFIRMARY
OF EDINBURGH, &c. &c.

THIS ESSAY IS INSCRIBED,

AS A TESTIMONY OF THE AUTHOR'S GRATITUDE,

FOR THE PROFESSIONAL ADVANTAGES DERIVED,

AND THE BENEFITS EXPERIENCED,

FROM THE KINDNESS

OF

AN AFFECTIONATE RELATIVE,

AND

SINCERE FRIEND.

ANDREW WARDROE M D

F. R. S.

THE REAL PROPERTY OF STREET, SECOND STREET, SECOND

THIS ESSAY IS INSCRIBED.

AND A PRESTURE OF THE AUTHOR'S CHAPTERING

FOR THE PROPERTY AND ADVANTABLE SERVICE.

AND THE BINEFIES EXPERS SACES.

THOM THE MINDSHE

20

STREET STATES TO STATE OF THE S

CHEN.

CONTRACT MARRIED

CONTENTS.

	Post interest boots to release the transfer of	age
Prelimi	ary Observations.	
CHAP	I. General Remarks on the Cornea,	.1
	II. Inflammation of the Cornea,	. 5
1	1. Of the Inflammation of the Conjunc-	
	tiva covering the Cornea,	.7
	2. Of the Inflammation of the proper	
	substance of the Cornea,	14
	3. Of the Inflammation of the Membrane	
	which lines the internal surface of	
	the Cornea,	20
	III. Of the Pterygium,	22
	IV. Of Fleshy Excrescences of the Cornea,	30
	V. Of Pustules of the Cornea,	37
	VI. Of the Abscess of the Cornea and Anterior	
	Chamber,	16
	VII. Of Ulcers of the Cornea,	54
	III. Of Wounds of the Cornea,	64
	IX. Of Foreign Bodies adhering to the Cornea,	67

CONTENTS.

Page
X. Of Ossification of the Cornea,72
XI. Of the Speck of the Cornea,
1. Of the Varieties of the Speck of the
Cornea,
2. Of the Formation of Specks of the
Cornea, 87
3. Appearances of Speck on Dissection, 92
4. Causes of Speck of the Cornea, 94
XII. Of the Staphyloma,
XIII. Of Alterations in the Form of the Cornea, 114
XIV. Of the Effusion of Blood between the La-
mellæ of the Cornea, and into the Ante-
rior Chamber,
Explanation of the Plates,
Plate I
Plate II
Plate III
Plate IV148
Plate V
Plate VI
Plate VII

PREFACE.

The object of the following essays is to describe the various morbid alterations in the structure of the human eye, and to illustrate, by engravings, those which are most remarkable.

In the accurate and detailed view which Dr Baillie has given of the morbid anatomy of some of the most important parts of the body, he has not described the diseases of this organ; and, as no attempt has yet been

made in this country to treat of the pathology of the human eye, little apology seems necessary for the present undertaking. Several excellent practical treatises and detached essays, indeed, have at different times been published; but during the last thirty years, the diseases of the eye do not appear to have excited the same attention here as on the continent.

RICHTER, of Göttingen, has, perhaps, given the best description of these diseases, and laid down the most judicious practical rules that have yet been suggested. Beer, of Vienna, has contributed many useful hints; Voigtel and Sybel have collected a great store of facts connected with the morbid anatomy of the

eye; and Conradi, Scarpa, and others, have likewise enlarged our knowledge of the treatment of the diseases of this organ. None of these authors, however, have delineated the morbid changes of structure which they have described: A few drawings only are to be found in the works of Beer and Scarpa, and in some periodical publications.

The importance of a work, the object of which is to supply these defects, is sufficiently obvious.

The opportunities which the eye affords to the pathologist, from the variety in its structure and situation, of discriminating all its morbid changes, and of observing their progress, renders it peculiarly interesting; and as there is no organ, the

loss of which can be productive of so many disadvantages, and so various and bitter calamities, without entirely destroying the existence of the individual, its diseases claim the most patient investigation, and deserve the minute attention of medical men.

Although the following pages are devoted exclusively to the investigation of one organ, the author is fully sensible that the task which he has undertaken is attended with considerable difficulty; but if they shall, in any degree, contribute to the attainment of the end proposed, or if they even excite the attention of medical men to so interesting a subject, the author will deem his labours well rewarded. It is only by continu-

ed attention, and patient investigation of the changes produced by disease in all the organs which compose the human body, that we can expect to enlarge the knowledge of the morbid anatomy of each, whilst, at the same time, it is also by a detailed account of the diseases of each separate organ that we can arrive at any general conclusions. But the field of medical science is very extensive, and to explore it with success, requires the co-operative efforts of many individuals; for it is only in proportion as facts are accumulated, and the various morbid appearances investigated, that the phenomena of disease can be understood, the morbid actions explained, the science of medicine freed from erroneous theories and hypotheses, and its practice liberated from the rash and unskilful hand of empiricism.

The late progress which has been made in pathological science, is a sufficient stimulus to exertion, and affords every reason to expect ultimate success. For, to use the words of Dr Reid, "We may, by caution and humility, avoid error and confusion. The labyrinth may be too intricate, and the thread too fine to be traced through all its windings; but if we stop where we can trace it no further, and secure the ground we have gained, there is no harm done; a quicker eye may, in time, trace it further."

In the following essays, the author has not only stated what he

has had an opportunity of observing himself, but he has endeavoured to collect information from the works of the respectable authors whose names he has mentioned, and from every other source to which he could find access. He ought also to acknowledge himself indebted to the liberality of many of his medical friends, who have afforded him opportunities of examining diseases which were either uncommon, or in any way worthy of notice. His field of observation has likewise been much enlarged, from the extensive opportunities which he has enjoyed in consequence of being professionally connected with Dr WARDROP, who, at an early period, directed his attention to this subject, and from

whom he has derived great assistance during the prosecution of it.

For the drawings which accompany this work, the author is, in a particular manner, indebted to Mr Syme, an ingenious artist of this place. He has combined the art of the painter with the skill of the anatomist; and as he has taken the trouble to retouch all the impressions of the plates, there is a truth and accuracy preserved in the colouring, which is seldom to be met with in works of this kind.

In some of the drawings, several diseases are represented; for every opportunity was embraced of delineating an eye affected with more than one complaint, in order to avoid multiplying the number of

engravings. Those who are not much accustomed to examine the morbid appearances of the eye, may begin with consulting the plates, as they will thence be enabled to form a general idea of the disease represented, and the subsequent account will thus become more clear and intelligible.

If the public shall approve of this essay, it is the author's intention to prosecute his plan, by considering the remaining diseases of the eye and its appendages, and the treatment which such diseases require.

Edinburgh, }
April, 1808.

engravings. Those who are and manch accusioned to examine the morbid appearances of the eye nor begin with consulting the plates as they will thence be enabled to toral a general idea of the disease cape will thus become more clear and to telligible.

If the public shall approve of this essay, it is the author's internion of the enables of this essay, it is the author's internion.

ing the remaining diseases of the eye and its suppendages, and the tentinent which such diseases

PRELIMINARY OBSERVATIONS.

All animals are composed of a certain number of organs, which, under the influence of the vital principle, produce those wonderful phenomena that distinguish living organized bodies. An acquaintance with the relative position, magnitude, and direction, of these organs, is one of the objects of our anatomical researches: On it the surgeon builds all his theories, and it guides his hand in every operation, But, for the advancement of physiological and pathological science, some-

thing more than this species of anatomy is necessary. It can only be considered as the first step towards a knowledge of the functions, and it does not even serve this purpose except under particular circumstances. When the anatomist divided the body into regions and districts, and shaped his inquiries to suit his unnatural divisions, every organ appeared insulated and detached, the most minute parts might have been discovered and described, but their mutual connections and sympathies were unknown. Anatomy and physiology were then disjoined; the former was imperfect, and the latter could scarcely be said to exist.

A more minute and philosophical examination of the structure and properties of the different organs, led the way to a knowledge of some of their functions, and pointed out the principles which should regulate the investigations of every rational physiologist. Haller was among the first to avail himself of the advantages of this plan. It conducted him to all his important discoveries; and

it has determined the progress of every scientific inquirer since his time. To it we are indebted for almost every improvement that has been made in this branch of science; and it is the only method by which we can hope still further to augment our knowledge. In pursuing this track, the labours of modern anatomists have been well rewarded. They have freed physiology and pathology from the nonsensical conjectures by which they were so long debased, so that they now begin to assume their rank among the sciences, and in some cases to afford a safe guide to the medical practitioner.

No one in our day has exerted himself more successfully in this field than the late celebrated Bichât. His Anatomie Generale is one of the most remarkable productions that has ever been produced in medical science. It has unfolded a path of investigation which was scarcely ever trodden before, and laid the foundation of a new anatomy and a new physiology. I cannot pretend here to do justice to the merits

of this work, nor to give a correct view of the facts and reasonings by which his doctrines are supported. They are as numerous and various as are the parts and functions of the living body. But, as I propose, in examining the pathological anatomy of the eye, to adopt some of the principles which he has established, the following observations are necessary, in order to explain the purport and tendency of the classification which I mean to adopt.

Most of the organs of our body are made up of a variety of elementary parts, or textures, each of which, in whatever situation it is found, affords uniformly the same physical properties. These are the elementary parts, which, by the diversity of their combinations, produce all the modifications of structure and functions which the different organs of animals exhibit. The study of these elementary parts, independent of the organs which they concur to form, is the object of general anatomy.

This method of considering organized bodies, is not an unnatural abstraction, nor a speculative refinement. It arises from the essential nature of their constitution, and it accords with every phenomenon with which we are acquainted. We may trace it in the observations of many of the older anatomists; and it may be considered as the basis of some of the most ingenious physiological theories of the late celebrated Mr John Hunter. Although, therefore, at first sight, it may have the appearance of being arbitrary and artificial, it is nevertheless, I am persuaded, founded on the most approved principles of philosophical investigation. A knowledge of the qualities of the different parts of which our organs are composed, must afford the surest means of acquiring information concerning the functions of these organs, and of becoming acquainted with the changes which they undergo in disease. On these principles Bichât has founded his anatomical system. To numberless experiments upon living animals, he added all the

information which could be acquired by dissection. He employed chemical re-agents to supply the deficiencies of the scalpel, and examined with minuteness all the varieties of morbid structure. By these means he endeavoured to fix the characters of the elementary textures, and then proceeded to investigate their combinations, as they are naturally presented to us in the different organs.

Of these textures, he has enumerated twenty-one, all of which he has shewn to be differently organized; and hence he proves the dissimilarity of their properties, both in health and in disease. This is the ground-work of the whole fabric, and to it we must ultimately recur in every attempt, to account either for the natural or morbid appearances which are to be met with among organized beings.

I mean not at present to enter more minutely upon the consideration of the elementary textures, my object being merely to shew, in a general manner, the effect of this anatomical arrangement, upon our patho-

logical theories. In our notions of all local affections, its influence is obvious; but in those diseases, where there is no evident change of structure, and where all the parts of the body seem to be disordered simultaneously, there is little room for the inquiries of the pathologist. It is, accordingly, in the former class of affections that the utility and advantages of general anatomy are most apparent.

By this view of the subject we have learned that diseases at their commencement are generally confined to one texture, the others of which the organ is composed remaining sound. This important truth is made manifest in many affections of the Eyes; but there is no part of the body, from which illustration of the same doctrine may not be deduced. At different times we see inflammation attacking the conjunctiva, or the various textures which form the cornea; at others, it is seated in the iris, in the capsule of the crystalline lens, the sclerotic coat, &c. The same is true of the different membranes

of the brain; of the mucous, serous, and muscular textures, which compose the stomach, and intestinal canal; of the cellular texture of the lungs; of the mucous texture of the bronchi; the serous one of the pleura, &c. &c.

But diseases are not only confined to one individual texture of any organ, as in the cases just mentioned; the symptoms and morbid changes are likewise uniformly the same in textures of a similar structure, in whatever part of the body these textures may happen to be found. Thus the serous membranes, which invest the lungs, the brain, the heart, the abdominal viscera, have one common character, when affected with any specific disease; so also have the mucous membranes, whether we trace them in the mouth, the nose, the vagina, urethra, or covering the eye-ball; and the same is observable of every individual texture which enters into the composition of our bodies. Dr Carmichael Smyth*, in this country, and

^{*} Vide Medical Communications and Inquiries, Vol. II.

afterwards Pinel *, in France, did much in pointing out the variety in the phenomena of inflammation, in some of the different textures of the body. It was an attempt highly worthy of the authors; but they did not lay down any system, or draw any general conclusions, nor did they attempt to trace the same analogy in other diseases. Thus, although the morbid changes of some of the textures have been ascertained with tolerable accuracy, we are still ignorant of many of the others. This is a field which has hitherto been little explored. It is of boundless extent, and presents inexhaustible subjects of investigation to the genius and industry of future inquirers.

Besides the symptoms and morbid changes which are common to all textures whose structure is similar in the natural state, there are others which are determined from the particular functions of the organ in which the diseased texture exists: For example,

^{*} Vide Nosographie Philosophique.

when any of the *serous* membranes are inflamed, the nature of the pain, the degree of fever, and the duration of the symptoms, are the same in whichsoever one it may have taken place. But to these symptoms are added, cough, difficulty of breathing, &c. when it happens to be connected with the organs of respiration, as in the case of *pleuritis*; costiveness, stranguary, delirium, loss of vision, when the intestines, the bladder, the brain, or the eye, are involved in the disease.

This view of the subject naturally suggests a correspondent division of the symptoms. The first class are general, and characterise a whole genus of textures; the second are in a manner accessory, and depend upon the relative situation, or the particular functions of the organ, into the composition of which the affected texture enters.

The foregoing remarks will tend to explain the object of general anatomy, and the important purposes to which it may be applied, both for illustrating pathology and therapeutics. But here we must set bounds

to this theory;—the history and progress of diseases shews that we ought not to confine our observations within such narrow limits. The principles which I have stated, indeed, account admirably well for the propagation of some affections, and for some of the sympathies which subsist between different parts of the body; but there are other disorders which advance in a different manner. In some diseases which are termed chronic, for example, the whole structure of an organ becomes gradually altered, although the primary affection was confined to one of its component textures. This is often to be observed in cancer, scrofula, lues venerea, &c. When cancer attacks the mamma, it is, at its commencement, generally confined to a small portion of that organ, but if allowed to proceed, it ultimately involves the whole glandular, cellular, and cutaneous textures, in one common mass of disease.

The author, from whom I have adopted some of the foregoing remarks, has, with wonderful ingenuity, illustrated and established the theory which I have in part described. In his hands morbid anatomy has assumed a new aspect; and he has pointed out a method of classifying the numerous facts which that science embraces, with a degree of accuracy and precision that was never known before. In prosecuting this science, we ought to examine the symptoms and changes of structure which are to be found in every individual texture, in whatever organ or region it may exist. After ascertaining the alterations proper to every system, we shall be better prepared to investigate the maladies as they take place in different organs of the body, or in different regions.

These general observations will be sufficient to give an outline of the principles of a pathological system, founded on the basis of anatomical knowledge; and when we apply them to the investigation of the morbid anatomy of the Eye, they will be found to afford a happy illustration of the system which I have ventured to adopt. For this beautiful organ is not only composed of a great va-

riety of textures, but the transparency and ready examination of many of its parts in the living body, admit of a great minuteness and accuracy of observation; and the various morbid changes can be seen going on much more distinctly than in any other part of the body.

The parts which form the eye-ball, and which are immediately connected and subservient to the performance of its functions as an organ of vision, as they present a great variety in their structure, they are necessarily liable to a proportionate variety in their morbid changes. The external covering of the eye-ball, eye-lids, and lacrymal passages, or conjunctiva, being a mucous membrane, we will find that it is subject to all the diseases of the mucous membranes in other parts of the body. The diseases of the cellular membrane, which lies underneath the conjunctiva, are analogous to those of the cellular membrane in other organs. The sclerotic coat, the iris, the choroid coat, the crystalline lens, the optic nerve, the retina, and

the different parts which compose the cornea, are also liable to morbid changes similar to those of the textures in other organs to which they are analogous, the various phenomena being more or less modified from the peculiarity of the functions of the organ.

In the descriptions of the diseases of the eye, which are given in the following essays, I shall divide them into two great classes. The first shall contain an account of the morbid changes of each separate structure, which enters into the composition of that organ; and, the second, an account of those diseases which have a specific character, and have symptoms peculiar to themselves, in whatever structure they appear, or which, when they attack the eye, affect a greater or less number of its different parts at the same time.

Under the first class will be comprehended, the diseases of the Cornea, Iris, Aqueous, Vitreous, and Crystalline humours, Optic Nerve and Retina, Choroid coat, Sclerotic

coat, Conjunctiva and Cellular membrane, Tarsi and Palpebræ, Lacrymal gland and Caruncle.

Under the second class will be included, Rheumatism, Cancer, Scrofula, Lues Venerea, Exanthematous Ophthalmia, &c.

As to the order in which these subjects are to be treated, I shall follow nearly that in which the different parts have been just now mentioned, as it appears to me to be the most simple arrangement, to begin with the parts which first present themselves when we examine the organ in its natural state.

1272 AMERICA COLONIA LA COLONIA L court of the training and the training of the court of th A SERVICE SERVICE DE LES PROPERTORS DE LA COMPANION DE LA COMP THE RESIDENCE OF THE PARTY OF T

CHAP. I.

GENERAL REMARKS ON THE CORNEA.

The cornea, in its natural state, possesses properties different from any of those classes of simple textures or membranes, into which anatomists have divided the component parts of organized bodies.

Some have conceived, that, in its structure, it much resembles the nails; but a very superficial examination will be sufficient to shew, that there is little similarity between these two textures. The nails are not, like the cornea, separated by a particular fluid into distinct layers. They are not endowed with an equal degree of sensibility, nor are they liable to all those diseases which occur in the cornea.

Others have attempted to demonstrate, that the cornea is merely a continuation of the sclerotic coat; but the functions and diseases of these two membranes differ so much from each other, that they clearly indicate a difference of structure and qualities. The sclerotic coat exhibits all the properties which fibrous membranes possess. It has a shining, opake, white, colour. It is composed of numerous filaments, running in every direction, which are closely interwoven with one another, and it cannot be divided into layers. The cornea is organized in a manner totally different. It is formed of an indeterminate number of concentric laminæ, which are easily separable from one another, either by the knife, or by maceration. They are, also occasionally separated, in particular diseases, by the effusion of blood or pus. From these circumstances, the cornea may be considered as a membrane sui generis.

But, though the cornea possesses properties peculiar to itself, yet it will be found, that the structure, functions, and morbid changes, of some of the parts of this membrane, have a striking analogy to those of other textures in the animal economy.

Anatomists have shewn, that the external surface of the cornea is a continuation of the tunica conjunctiva; this covering seems referable to that class of membranes which have been denominated *mucous*. It can be separated from the subjacent parts by dissection; it is of a softer texture, and tends more rapidly to putrefaction than the substance of the cornea. It is also nourished by the same vessels which supply the conjunctiva, covering the sclerotic coat, as may be seen when it becomes inflamed.

The internal surface of the cornea is lined by a membrane that forms part of the cavity which contains the aqueous humour, and perhaps it assists in its secretion *. According to the arrangement which I have adopted, it may, with propriety, be considered as a membrane of the *serous* class.

^{*} Tunica humoris aquei.

The substance of the cornea, which is situated between these two membranes, is composed of concentric cellular lamellæ, in the cells of which a peculiar fluid is deposited by the exhalents. This analogy in the two surfaces of the cornea to mucous and serous membranes, which is observed in the sound eye, is also remarked in the morbid changes of this organ. Some of the diseases to which the cornea is liable, are the same with those which occur in membranes of the serous and mucous textures; and they are accompanied, not only with those symptoms which characterize the diseases of these textures in all other parts of the body, but by a series of phenomena peculiar to the cornea.

In the description of the diseases of the cornea, I shall begin with inflammation, as it is the morbid affection which occurs most frequently, and as many of the other diseases to which the cornea is liable, are either preceded or accompanied by one or other of the various modifications of that state.

CHAP. II.

OF THE INFLAMMATION OF THE CORNEA.

The cornea in the sound eye is perfectly transparent, and none of its vessels admit the red particles of the blood; but, when inflammation takes place, vessels carrying red blood may be distinctly seen ramifying through it, and every change which takes place in the number, in the mode of distribution, or in the colour of the contents of these vessels, may be accurately examined. Besides an increase in the vascularity, inflammation of the cornea is attended with more or less dimness of vision, pain in the eye, in-

tolerance of light, and those symptoms which usually take place when any other part of the organ of vision is inflamed.

The inflammation, however, does not always affect all the textures of which the cornea is composed; and though, in many instances, it may be difficult to discriminate the precise extent of the disease, yet, in others, it may be clearly shown, that the inflammation is entirely confined to one of the textures.

As this distinction involves very important consequences, both in a pathological and practical point of view, we shall consider inflammation of the cornea under three separate heads:

First, As affecting the conjunctiva, which covers the cornea externally.

Secondly, As being confined to the proper substance of the cornea; and,

Thirdly, As affecting the membrane which lines the internal surface of the cornea.

1. Of the Inflammation of the Conjunctiva covering the Cornea.

It is by no means unusual to observe, in those who have had long or repeated attacks of inflammation of the conjunctiva covering the sclerotic coat, that a considerable degree of obscurity and vascularity extends from it over the cornea. This took place in many of our troops who suffered from ophthalmia in Egypt, and who have, since their return home, been subject to frequent inflammatory attacks in their eyes. Many cases have also occurred where the same effect was produced by the ophthalmia which has been so frequent among the troops in this country. The obscurity of the cornea, in the cases to which I allude, appeared to be confined to the external surface. Red vessels became ramified upon its surface, and they were much more numerous in proportion to the degree of opacity, than they ever are in the

speck of the cornea. I have observed, in a variety of cases of ophthalmia in this country, where the conjunctiva covering the sclerotic coat was attacked with inflammation, that red vessels branched out from a particular portion of the affected part, and extended a little way over the edge of the cornea; but these vessels generally disappeared when the more violent inflammatory symptoms were removed. In all these cases, however, the inflammation commenced and extended from the conjunctiva, covering the white of the eye, to that part of it which covers the cornea. But in other instances, the inflammation first appears in the conjunctiva, covering the cornea, and is almost entirely confined to it alone. When this takes place, it very much resembles a speck on a superficial examination, but the commencement, progress, and final termination of these two diseases are very different. Inflammation of the conjunctiva of the cornea begins at the circumference, and gradually extends from this over the surface of the cornea, whereas a

speck commonly commences at the centre of the cornea, or at some distance from its circumference, and extends in every direction from that point. In the inflammation of the conjunctiva covering the cornea, the opaque part becomes elevated above the natural surface of the cornea, and is highly vascular; whereas, in speck, the form of the cornea remains unaltered; and, although one or more vascular trunks advance to the speck, yet the distribution of their branches becomes in general imperceptible.

Inflammation of the conjunctiva of the cornea may also be mistaken for a pustule of that membrane; for, in the latter disease, the conjunctiva adjacent to the pustule is sometimes distinctly vascular, and elevated above its natural surface. But if the progress of the two affections be attentively observed, they will be found to be different; for a pustule of the cornea, like a speck, begins at one point, extends from it as a centre, and most frequently terminates in an ulcer.

The most remarkable symptom of inflammation of the external covering of the cornea, is the appearance of an elevated and whitish coloured spot or streak, in some point, near the union of the cornea with the sclerotic coat, accompanied by a greater or less degree of inflammation of the adjacent conjunctiva. The blood vessels in the inflamed part are of a bright scarlet colour, run in a straight direction, and are more numerous towards the diseased spot, some of them being ramified through it. This opaque spot gradually extends across the cornea, acquiring, as it proceeds, an oblong form, with a rounded obtuse point. It continues elevated above the natural surface, is distinctly defined, and remains nearly of the same breadth till it reaches the centre of the cornea. If it passes beyond this, it becomes broader, and the breadth increases as it extends, but I have never, in any instance, observed it pass. as far as the opposite edge of the cornea. The degree of vascularity also increases in proportion as this spot becomes larger; and if a

magnifying glass be used, the whole of the diseased part will be seen crowded with red vessels*.

When the inflammatory symptoms abate, and the progress of the complaint is checked, the vessels distributed on the diseased portion of conjunctiva acquire a purplish colour, gradually diminish both in size and in number, and run in a tortuous direction; the elevation of the membrane subsides, and if the inflammation has neither been very violent, nor of long duration, its transparency is completely restored. But this is not always the way in which it terminates, as a degree of obscurity sometimes remains after all the inflammatory symptoms and appearances of vascularity have abated. In other instances there is both a slight degree of obscurity, and some red vessels. In one case, where the accompanying inflammation was very violent, the diseased portion of conjunctiva separated completely from the subjacent cornea, came

^{*} Compare fig. 2. Plate I. and fig. 1. Plate II.

off in the form of an opaque slough, leaving a slight degree of obscurity in that part of the cornea over which it was situated.

After the eye has been once affected with this disease, it is very liable to subsequent attacks; and although the cornea regains its transparency, and the red vessels disappear, yet every fresh inflammation is more obstinate than the former, a larger portion of the conjunctiva becomes inflamed, and even after the inflammatory symptoms disappear, the conjunctiva remains interwoven with varicose vessels, flaccid and opaque, having the appearance of a new formed membrane *.

Most of the cases of this disease which I have had an opportunity of observing, were in children. In one instance, however, it occurred in a lady nineteen years old. I have seen another very remarkable example of this disease in a strong healthy man, about thirty-five years of age, under the care of Mr Johnston, surgeon at Kirk-

^{*} Vide Pterygium.

ealdy. In this instance it was accompanied with very violent inflammatory symptoms at its commencement; and although the whole of the corneal substance had become cloudy, from the commencement of the disease, yet the patient himself, by examining daily his eye at a glass, gave a very accurate account of the progress of the opacity of the conjunctiva covering the cornea, and described it as beginning at the edge of the cornea, and extending gradually to a little way past its centre. When I saw him, all the inflammatory symptoms had disappeared, the substance of the cornea had regained its transparency, and the affected portion of the corneal conjunctiva remained thickened, of a brownish colour, and interwoven with blood vessels, the trunks of which were formed into a cluster, at the union of the cornea with the sclerotic coat.

participated to the severity of the other

2. Of the Inflammation of the Proper Substance of the Cornea.

If the whole of the proper substance of the cornea be inflamed, the red vessels enter at every part of its circumference. They are always most numerous at the margin, and few of them advance to the centre. I have seen them so numerous, as to form a red band or ring round the outer edge, leaving the central part transparent. If the inflammation be confined to one spot, the red vessels appear in clusters on the adjacent portions, both of the cornea and sclerotic coat.

From the commencement of the inflammation, the cornea loses its transparency and lustre, and becomes so dim and clouded, as greatly to interrupt vision. There is either an equal degree of obscurity over the whole cornea, or one part is more opaque than another; but the degree of it is, in general, proportioned to the severity of the other symp-

violent, I have seen small vesicles, filled with red blood, formed between the layers of the cornea; and in some cases, blood effused into the aqueous humour, tinging it of a red colour. In the first, or active stage of the inflammation, the blood-vessels on the sclerotic coat are of a bright scarlet colour. They run in a very straight direction, and pass over the edge of the cornea. Each trunk can be readily distinguished, and the branches into which it is divided come off at very acute angles, and not until the trunk reaches the circumference of the cornea.

Inflammation of the cornea is also accompanied by more or less general fever, pain in the eye, extending to the head, intolerance of light, increased secretion of tears, and impaired vision.

Although inflammation may originate in the cornea, it is always accompanied with a preternatural degree of redness of the sclerotic coat; and it is often impossible to determine whether the cornea or sclerotic coat is primarily affected. In cases where the cornea is wounded, it cannot be doubted, that the inflammation originates in the wounded part, however far it may afterwards extend; but, in many instances, both the cornea and sclerotic coat seem to have an equal share in the disease. It is probable that this ready communication of the diseased action between these two coats, depends chiefly on their vascular connection; for the vessels of the substance of the cornea are all derived from the sclerotic coat, and, as has been before mentioned, those of the conjunctiva covering the cornea, are continuations of the vessels of the conjunctiva covering the sclerotic coat.

The symptoms of inflammation of the cornea generally suffer, whether by the application of remedies, or from the progress of the disease, a very remarkable change after a certain period. The red vessels of the sclerotic coat and cornea, increase in size and in number, change in colour, and are distri-

buted in a different manner. Instead of running in straight lines, and sending off their branches at acute angles, they become tortuous, anastomose, and form net-works. The bright colour changes into a darker and more purple hue, and the eye appears as if glazed, losing all its lustre and expression. The pain, too, and general fever, which were severe, are now alleviated, or cease altogether, and the free admission of light is not attended with much uneasiness. If the inflammation has been occasioned by an wound, such as that which is made in the extraction of the cataract, this change takes place generally several days after the operation. The change approaches more slowly in those cases where the inflammation arises from other causes, and rarely happens till after a period of some weeks, or even months, from its commencement. In many cases, indeed, no such distinction can be made, as, throughout the whole progress of the disease, there is a constant change taking place in the abatement or increase of all the symptoms.

I have seen a few cases where there were a number of varicose vessels in the cornea, in consequence of previous inflammation; and the patients, in describing the state of their vision, said they saw small lines, or streaks, passing across the eye, which they sometimes observed to vary in their size, and to have a kind of irregular vermicular motion, such as the blood-vessels in the cornea may be supposed to have.

If an attempt be made to divide the enlarged vessels as they pass over the cornea, (a practice recommended in the treatment of the disease), they are found to lie deep, and they cannot be so easily raised from the cornea or sclerotic coat with the forceps, or elevated by a pointed instrument, as may be done in the natural state of these parts. In order to complete the division of one of these vessels, it becomes necessary to remove a considerable portion of the substance of the cornea, or sclerotic coat. Soon after such an operation has been performed, lymph is effused on the surface of the wound; and it

often happens, that vessels shoot through this lymph, forming a medium of communication between the divided extremities; so that, after the effects of the operation seem to have abated, the vessel remains in one continued trunk, and appears as if no division had been made. In other cases, the minute ramifications of the divided vessel which remain, anastomose on the cornea, so that red blood continues to be circulated through the trunk.

Inflammation of the cornea occurs at every period of life, and, like other inflammatory affections, is most frequent among the young and plethoric. It most frequently arises from wounds and other external causes; but the cornea is also often inflamed in cases of ophthalmia arising from lues venerea, small-pox, scrophula, and various eruptive diseases; the inflammation being, in all those cases, modified according to the specific causes from which it originated.

Inflammation of the cornea, when proper means are employed for its removal, most frequently terminates by the disappearance of all its symptoms. Sometimes, however, during the progress of the violent inflammation, purulent matter is formed between the lamellæ of the cornea, which, by ulceration, either discharges itself internally into the anterior chamber, or externally, leaving an ulcer of the cornea. More frequently, after all the inflammatory symptoms have disappeared, a part of the cornea does not regain its natural transparency, so that a *speck* remains, which continues to be nourished by a number of red vessels.

3. Of the Inflammation of the Membrane which lines the Internal Surface of the Cornea.

Although, from analogy, it is highly probable that the membrane which lines the internal surface of the cornea may become inflamed, and the inflammation neither affect the proper substance of the cornea, nor its external covering, yet I have never been

able to observe an instance with sufficient accuracy where this took place. In cases of ophthalmia from lues venerea, there is often a peculiar muddiness of the cornea, apparently deep seated, followed by the effusion of lymph between the cornea and iris; and it is by no means improbable that, in these cases, the lymph has been effused, and the opacity of the cornea produced by the inflammation of its internal layer. I have also frequently observed cases where the iris and internal surface of the cornea had formed adhesions, in which the substance of the cornea did not appear to have been inflamed. The inflammation, or at least some of the morbid changes of the internal layer, may also be concerned in the cases where the quantity of the aqueous humour is either increased or diminished; and it must often participate in the inflammation which originates in the proper substance, or in the external covering of the cornea.

CHAP. III.

OF THE PTERYGIUM.

The word *Pterygium** is employed to denote all those morbid changes, in which that portion of the conjunctiva covering any part of the cornea or sclerotic coat becomes thickened, vascular, and opaque; and some authors have attempted to introduce a number of terms to characterise the numerous varieties of the disease, considering each as a distinct species.

^{*} Das Augenfell of the Germans,-L'Ongle of the French.

[†] See Traité sur les Maladies de l'Oeil, par Antoine Maitre-Jan,—Also Anfangsgründe der Wundarzneykunst, von August. Gottlieb Richter. Dritter Band.

It has been already mentioned, that inflammation may be confined to the conjunctiva covering the cornea. If this inflammation continues for a long time, or if there are repeated attacks of it, the affected portion of conjunctiva assumes the appearances of a new formed membrane, even after all the inflammatory symptoms have abated. This new formed membrane constitutes what has commonly been denominated the membranous pterygium, or pannus*.

It also sometimes happens, that the portion of the conjunctiva which covers the sclerotic coat, becomes preternaturally thick, and the cellular membrane which connects the thickened part of it with the sclerotic coat, is so much relaxed, that it may be very easily moved backwards and forwards; and when the eye-ball is placed in particular positions, it forms itself into folds, and becomes as if wrinkled. This thickening and relaxation of the conjunctiva extends, in some cases,

^{*} Pterygium tenue, Ungula, l'Onglet, see Plate III. fig. 1.

round the whole circumference of the white of the eye*, in others, it is confined to a small part of it.

In the first case, the disease has the appearance of a dull white-coloured fold, all round the edge of the cornea, and the eye loses its shining appearance and lustre, and becomes of a yellowish colour. When the disease increases, the fold gradually extends over the cornea, approaching towards the centre.

If the disease be confined to a particular part of the conjunctiva, it is observed, at its commencement, like a small globule of fat; or condensed cellular substance, situated near the junction of the cornea and sclerotic coat; and this spot, extending imperceptibly along the surface of the conjunctiva, at length passes over the cornea. After it has extended a little way, the conjunctiva on the adjoining part of the sclerotic coat becomes puckered, and appears

^{*} See Richter's Anfangsgründe.

⁺ Das Fettfell of the Germans,-Pterygium pingue.

as if it were drawn forcibly over the cornea. The portion of it which lies on the sclerotica is commonly loose, and can be easily elevated; but that which is on the cornea adheres more firmly. This species of pterygium has generally a triangular form *, one of the angles of the triangle either advancing towards the cornea, or covering it, and the base lying on the sclerotic coat. Sometimes the thickening of the conjunctiva is first perceived on the cornea, the conjunctiva covering the sclerotica remaining quite sound +. A pterygium is always considerably elevated above the surface of the adjacent cornea, but the degree of its thickness varies from that of a thin membranous film to a thick fleshy mass ‡. In some cases it has been found thick and coriaceous, and in others as hard as parchment, and even cartilaginous §. The cellular substance under

^{*} See Plate III. fig. 2.

⁺ Vide Richter's Anfangsgründe.

[‡] See Plate III. fig. 1, 2, and 3.

[§] Vide Richter's Anfangsgründe:

the conjunctiva sometimes participates in this disease; in other instances it does not seem to be affected.

In those pterygia which have a membranous-like appearance, the red vessels are generally few in number, and run in straight lines from the sclerotic coat towards the centre of the cornea. In those which are thicker, and which have a fleshy appearance, there is a general red tinge given, from the vessels being very numerous.

Pterygia arise commonly at the great or nasal angle of the eye-ball; they are formed also at the small or temporal angle, and they sometimes occur in both places, in the same eye *. I have seen one case in which there were two pterygia on each eye. They are formed very rarely on the under part of the eye-ball.

This disease seldom extends farther than the centre of the cornea, when it begins in

^{*} Vide Traitè Pratique des Maladies des Yeux, par A. Scarpa, fig. 3. Plate IV.

a single point. But when two pterygia arise from two opposite points of the same eye, they sometimes spread over the cornea till they nearly meet, and then form a complete obstruction to vision; the imperfection of vision in this disease being in proportion to the thickness of the pterygium, and to its approximation to the pupil.

When pterygium arises in the nasal angle of the eye, it seems almost always to attach itself to the semilunar membrane, and in many cases it also adheres to, and involves the caruncula lacrymalis.

Pterygia occur most frequently in people advanced in life. They are, however, also met with in children: And I have seen one instance in which the disease was observed immediately after birth.

Many have supposed that the pterygium was a particular kind of expansion or growth from the caruncula lacrymalis, or from the semilunar membrane; but from the variety in its appearances, and from observations on the progress of the disease, it would seem that

its connection with these parts is accidental; and that pterygia arise from a variety of causes, some of which we are, perhaps, not able satisfactorily to explain. It has been already mentioned, that the thin membranous pterygium is the consequence of repeated attacks of inflammation of the conjunctiva covering the cornea. The manner in which the common triangular-shaped pterygium is formed is much more singular, and appears to have no analogy to any morbid change in other organs which have a similar structure. The constancy in the regularity of its triangular form ought to be referred, says Scarpa, the celebrated Professor of Pavia, to the adhesion of the lamina of the conjunctiva covering the cornea becoming stronger, in proportion as it advances from the circumference to the centre of the cornea; for, in consequence of such structure, and different degree of cohesion which exists in the sound eye, it should necessarily follow, that, in the first place, the progress of the pterygium ought to be, in every case

of the disease, much slower upon the cornea than upon the white of the eye; secondly, that, from the greater resistance which the pterygium always meets with in proportion as it extends towards the centre of the cornea, it ought, from mechanical necessity, to assume a triangular form, the base of the triangle corresponding to the white of the eye, and the apex to the centre of the cornea. Its progress is, in almost every case, very slow; it arises without any evident cause, and gradually increases, without pain or inconvenience, until it acquires a considerable bulk, and encroaches on the sphere of the pupil. In many instances, it remains for years without undergoing any perceptible alteration.

CHAP. IV.

OF FLESHY EXCRESCENCES OF THE CORNEA.

Besides pterygia there are excrescences of a different description, which are called Caruncles*, or fleshy excrescences of the cornea. Of these there are two very distinct kinds. One of them appears at birth, or soon after it, and resembles the nævi materni so frequent on the skin of various parts of the body. The second species has a greater analogy to the fungi which grow from mucous surfaces, and they arise commonly from previous ulceration.

^{*} Carunculæ Corniæ, -- Excroissances des Chairs, of the French, -- Fleishgewächses of the Germans.

I have had an opportunity of examining two very remarkable examples of tumours of the cornea, which appeared at birth. The first was that of a girl of eight or ten years of age, on whose left eye there was a conical shaped mass, the base of which grew from about two thirds of the cornea, and a small portion of the adjoining sclerotic coat. It was firm and immoveable, had a rough granulated appearance externally, and, from its brownish colour, did not appear to be very vascular. It was very small when it was first observed, and it increased in size in proportion with the other parts of the body.

The second case I saw along with Dr Monro, junior. The patient was upwards of fifty years old, and the tumour had been observed from birth. It was about the bulk of a horse bean, and only a small portion of it adhered, and seemed to grow from the cornea; the other part was situated on the white of the eye, next the temporal angle of the orbit. Its surface had not the particular appearance which was in the girl's eye; it

was smooth like a pterygium, and seemed to be covered by the conjunctiva, having the natural colour of that membrane. But the singularity in this case was, that a considerable number of very long and strong hairs, upwards of twelve in number, grew from the middle part of it, passed through between the eye-lids, and hung over the cheek. The patient remarked that these hairs did not appear until he advanced to his sixteenth year, at which time also his beard grew *.

Dr Barron of St Andrews met with a similar case when at Lisbon, with the following account of which he has favoured me:

"The disease took place in a boy of fifteen

"years of age. It was a flat tumour, about

"one-third of an inch in diameter, with a

"perfectly circular base. More than one half

"of it was situated on the cornea, and the

"rest on the conjunctiva, adjoining to the

"temporal angle of the orbit. Its surface

"was smooth and shining, and from its

^{*} See Plate IV. fig. 1.

"centre grew two hairs, similar to those in
"the tarsus of the upper eye-lid. In colour
"it resembled the white part of the conjunc"tiva. It was, however, rather more of a
"pink hue. The disease was of five years
"duration, and it was, at times, accompanied
"with excessive pains above the orbit, and
"in the temple of the side affected."

Mr Crampton of Dublin, in his Essay on the Entropeon, p. 7. also mentions, that he once saw "a tuft of very strong hairs proceeding from the sclerotica;" and De Gazelles saw a case where there was a single hair growing from the cornea*.

I have in my possession, a preparation of a disease of this kind in an ox's eye, where a thick tuft of black hair grows out from and covers about one-third of the cornea, and some hairs are also observed growing from the semilunar membrane. A similar excrescence had formed in the other eye of this animal.

Such tumours greatly resemble those spots covered with hairs, which are so frequent in different parts of the surface of the body, particularly the face. I remember to have seen the description of a very curious case of a child, where a tumour, covered with hair, appeared in the pharynx.

A fungous tumour, arising from the surface of an ulcer on the cornea, is a rare disease. It, however, frequently happens, that an ulcer destroys the whole thickness of the cornea, allowing a portion of the iris to protrude, and the protruded portion has, in many cases, been the origin of large excrescences. The following case is a remarkable example: "The largest excrescence I ever saw," says Maitre-Jan*, " arose from an ulcer which occupied partly the opaque, and partly the transparent cornea. It was so large as to advance beyond the eye-lids,

^{*} Traité des Maladies des Yeux.

like a mushroom, and cover the whole eveball." After mentioning the means employed to remove it, he adds, "that when the excrescence was consumed to the level of the cornea, I then observed, that its base only occupied one half of the small angle; that the cornea was ulcerated and broken; and that the roots of the excrescence passed from thence, and had their attachment to the uvea." Voigtel * quotes the case of a boy from Mohrenheim, in whom, after a violent inflammation of the eye, a small white point appeared on the inferior part of the cornea, which gradually grew into a hard cartilaginous tumour, of the bulk of a pea. Its base covered one half of the cornea, and its surface was interwoven with large blood vessels. Beer + describes a fatty and fleshy swelling of the cornea

^{*} See Handbuch der pathologischen Anatomie, von F. G. Voigtel. Halle, 1804.

⁺ Practische Beobachtungen über den grauen Staar und die Krankhieten der Hornhaut, von Joseph Beer. Vienna, 1791.

which was as large as a cherry-stone; and Plaichner * relates a case, where a spongy tumour, the size of a hen's egg, grew from the cornea, after the removal of a fleshy swelling.

^{*} Dissertatio de Fungo Oculi, 1780.

CHAP. V.

OF PUSTULES OF THE CORNEA *.

Pustules are small tumours, which are formed both on the cornea and sclerotic coat, but they occur most frequently near the junction of these membranes.

A pustule of the cornea commonly first appears like a dusky, yellow, or reddish spot, a little elevated above the surface of the cor-

^{*} They are called Pustulæ Corneæ—Eiterblattern by the Germans, when they contain matter. Phlyctenæ Corniæ,—Phlyctides,—Wasserblattern, when they contain water.

nea, and, in a short time, it becomes a distinct conical tumour. The adjacent part of the cornea is always more or less dim, and a considerable degree of inflammation * accompanies it, which is either confined to the white of the eye contiguous to the pustule, or is spread over the whole eye-ball. Whilst the pustule is forming, the inflammation is generally confined to that part of the white of the eye which is in its immediate vici-The vessels, instead of being of a nity. bright scarlet colour, as in the inflamed cornea, are of a pale livid hue. They appear superficial, and can be readily elevated by a pointed instrument. Each trunk can be distinguished, for they are never so numerous as to appear confused, or like one red mass, an appearance so common when a portion of the conjunctiva is inflamed +.

They sometimes run in various directions, anastomose freely with one another, forming net-works on the white of the eye.

^{*} Ophthalmia Pustulosa, or Catarrhalis.

⁺ See Plate I. fig. 3.

If the inflammation and pustule remain for some time, the pustule generally advances to suppuration. When suppuration takes place, the apex of the pustule ulcerates, and frequently a chalky white spot appears at the centre of the ulceration, and the opacity of the cornea at the same time daily increases around it. In other cases the opaque matter separates, and leaves behind it a deep ulcerous excavation *.

Sometimes the suppuration proceeds more like a common pimple or *phlegmon* of the skin; a small quantity of a thick matter collects within the pustule, and when it is discharged, a conical tumour remains, which has a depression at the apex. When the pustule contains a watery fluid, the fluid is most frequently absorbed in a gradual manner, but at other times the pustule breaks, and an ulcer is formed.

If, in either of these cases, the contents of the pustule are artificially discharged, all

^{*} See Ulcer of the Cornea.

the accompanying inflammatory symptoms are much increased *.

Most frequently there is only one pustule, and only one eye affected, but in some cases there are several, both on the cornea and sclerotic coat of each eye.

The disease, at its commencement, is almost invariably accompanied with the sensation of a mote in the eye, and the whole conjunctiva covering the sclerotic coat has often a yellowish and shining glassy colour before the redness appears. There is often also a degree of redness and swelling, chiefly of the upper eye-lid, and the tarsi are found adhering together in the morning from the exudation of a yellow matter among the ciliæ. There is frequently an unusual dryness felt in the eye; but if it is exposed to a bright light, or if any attempt is made to use it, the secretion of tears is increased.

This species of inflammation is always accompanied with a much greater degree of

^{*} Vide Richter's Anfangsgründe.

general fever, in proportion to the severity of the other symptoms, than any other ophthalmia. The pain is rarely acute till the pustule ulcerates, but, if that takes place, it is commonly very severe.

As the pustule disappears, and the subsequent ulcer heals, the inflammatory symptoms generally abate. It is not, however, unfrequent to find, that although this has taken place, the inflammation returns on any slight irritation; and I know several instances of people who are very subject to repeated attacks of this disease. In some of the cases where it returns frequently, the pustule seldom ulcerates, but disappears gradually, after having remained a few days. In one case, after a second attack of the inflammation, a quantity of a yellow lymph was effused near the margin of the cornea; and I have seen two instances where a small quantity of purulent matter formed in the anterior chamber. It was readily distinguished by its yellow colour at the inferior margin of the cornea, and it diminished daily as the inflammation abated.

Pustules of the cornea are met with in people of all ages, but they are more common in the young than in those advanced in life. This disease appears to be more frequent in particular places, and in particular seasons of the year. In Edinburgh, where it is not a very frequent complaint, I have generally observed it in the winter months.

During the winter of 1804, I saw a great number of patients with this disease in the clinical school of Dr Beer, at Vienna, for diseases of the eyes; and in all of them it was accompanied with such a degree of fever, as to require attention almost solely to the general treatment. I have, in several cases, been able to trace its origin to the sudden change from a very warm to a cold atmosphere. At one of the theatres of Vienna, where a very cold stream of air passed immediately behind the orchestra, it was observed that those people who were placed

near it were very often affected with this complaint.

It appears to me highly probable that the pustules of the cornea, and white of the eye, very much resemble the small pustules, or aphthæ, as they are called, which are so frequently observed in the cavity of the mouth, on the tongue, lips, and on the internal surface of the alimentary canal. Although these aphthæ may be, perhaps, considered as a morbid change of the mucous membranes, yet the phenomena attending the disease are very different from those which are observed in the common inflammation of a mucous surface. From the swelling which takes place, and from the tendency of this swelling to suppuration, there appears to me to be a striking analogy between this inflammation and inflammation in the cellular membrane in other parts of the body, or the common phlegmon. It is, therefore, by no means improbable that pustules are formed in the cellular texture, which enters into the composition, or which lies immediately

underneath the conjunctiva which covers the eye-ball.

As some remarks of Professor Himly of Brunswick illustrate very strikingly this analogy, I shall finish this description by quoting his words: "At a time," says he, "when aphthæ of the throat were very frequent at this place, I also found many small vesicles, beginning with an inflammation of the sclerotic coat, and also sometimes, but more rarely, of the cornea. Once I saw a whole family affected with this disease, one after another. It was a true catarrhal affection, and in some cases these vesicles disappeared by diaphoretic medicines, in some by blisters, camphor, and antimony, without any local application, except mucilaginous ones. I think that it is just the same disease as aphthæ of the intestinal canal, of the corona of the glans penis, and other fine continuations of the external skin. Those on the cornea become worse if they are opened; and if they open themselves, and form ulcers, they generally dry up by means of borax and white vitriol, but

if they are neglected, they cause sometimes considerable ulcers, which are very obstinate and hurtful to the cornea *."

* Vide Bemerkungen über einige Augenkrankheiten, von Professor Himly zu Braunschweig, p. 402, of the 1st volume of Loder's Journal.

CHAP. VI.

OF THE ABSCESS OF THE CORNEA AND AN-TERIOR CHAMBER.

Purulent matter may be collected either between the lamellæ of the cornea, when the disease is termed unguis or onyx, or in the anterior chamber, along with the aqueous humour, when it gets the name of hypopion or empyesis oculi *.

When the matter is collected between the lamellæ of the cornea, it first appears like a small spot; and instead of resembling a speck in colour, it is of the yellow hue of common pus. As the quantity of the matter increases, this spot becomes broader, and it

^{*} Die Eyterung des Auges, or Der Hornhaut Apostem of the Germans.

does not alter its situation from the position of the head. If it is situated among the external layers of the cornea, or immediately below the corneal conjunctiva, a tumour is formed anteriorly, which, if touched with the point of a probe, the contained fluid can be felt fluctuating within it, or if the eye is looked at side-ways, there is readily perceived an alteration in the form of the cornea.

When the matter collects between the interior lamellæ, it does not produce any evident alteration in the external form of the cornea, but if it is touched with the point of a probe, a fluctuation can be more or less distinctly perceived, and the spot alters its form, and becomes somewhat broader.

Such collections of matter appear on every part of the cornea. Sometimes they alter their situation by degrees, and sink downwards; and sometimes they change both their situation and form. They very seldom cover more than one-fourth or one-third of the cornea; but, in one instance, I saw the

matter so extensive, as to be spread over nearly the whole corneal surface.

If the quantity of matter is small, it is often completely absorbed during the abatement of the inflammatory symptoms, and it generally leaves no vestige behind it; In other cases, the cornea is eroded externally, producing an ulcer and subsequent opacity *. In some few instances, the internal lamellæ of the cornea give way, and the matter escapes into the anterior chamber. If an artificial opening is made, in order to discharge the matter, it often does not flow out readily; and it is sometimes so tenacious, and contained in a cavity apparently so irregular, that it neither escapes spontaneously, nor can it be evacuated by art.

When the matter is collected in the anterior chamber, it generally appears like a small yellow globule between the iris and cornea; and as its specific gravity is greater than that of the aqueous humour, it generally

^{*} Vide Ulcer of the Cornea.

occupies the inferior part of the cavity *. In some cases I have observed the matter appearing like small opaque flocculi diffused through the aqueous humour, and these, by gradually uniting together, formed a drop of purulent matter, which sunk to the inferior part of the cornea. As the quantity of matter increases, the spot becomes larger, and often assumes a semilunar form at the edge of the cornea; sometimes it collects in such a large quantity as to pass through the pupil, and fall behind the iris. When this happens, the cornea not only becomes opaque, but also loses all its natural firmness of texture; and in some very far advanced cases of the disease, when attempting to discharge the matter, I have found it quite soft, and easily torn to pieces,

Unless when in great quantity, the matter is generally absorbed in proportion as the inflammatory symptoms are alleviated; but if it remains a long time, it sometimes ulcerates the cornea, or becomes inspissated into a tough, light-coloured mass, which remains after all the inflammatory symptoms have disappeared.

Schiegel mentions a case of abscess of the anterior chamber, in which the pores of the cornea were so open that the matter oozed out in the form of threads. The following are his words, "By the application of the decoction * in a case of hypopion, the matter discharged itself in an uncommon manner. Whilst using it, the fine pores of the cornea were opened, and the matter oozed out in the form of fine threads. On the second day the distended cornea was considerably flatter, the oozing out of the matter continued without interruption, and in four days nearly two drams of matter had passed through the pores †."

These abscesses are commonly the effect of violent ophthalmia, occasioned by blows,

^{*} Decoct. Fol. Malvæ cum Opio.

⁺ Vide Magazin für die Wundarzneiwissenschaft herausgegeben, von J. Arneman, 2d Band. 2d Stück.

or injuries of the eye-ball. Richter, however, remarks, that purulent matter sometimes collects in venereal and scrophulous patients, without any preceding inflammatory symptoms. Dr Rutherford mentioned to me the case of a woman who was under his care in the Royal Infirmary of Edinburgh, who had a very considerable collection of matter in the anterior chamber, accompanied with very little or no inflammation. The matter altered its form and place, according to the position of the head, and, during the day, the agitation of the body, produced from walking, mixed the matter with the aqueous humour, and rendered the whole anterior chamber turbid. Janin relates a very curious case, where there was not only the absence of the inflammatory symptoms, but where the disease recurred periodically. " About the beginning of the month of March 1757," says he, "Peter Valis consulted me about a periodical blindness with which he had been affected for twelve months, during the first fifteen days of every month; and, after

that time had elapsed, his eyes were restored to their natural state. I examined the organ, in order to ascertain the cause of that singular kind of blindness, and I observed that the anterior chamber of both eyes was filled with a yellow-coloured matter, so thick as neither to allow the colour of the iris, nor the state of the pupil to be seen through it. The most remarkable circumstance in this case was, that the conjunctiva was very little inflamed, and the eye not painful *."

Richter saw a man who was blind every morning, and it was always remarked that, while the paroxysm lasted, the aqueous humour was quite turbid.

It has been a subject of dispute, to account for the source of the matter which is formed in hypopion. It appears most probable, that as there is no ulcerated surface from which it can be derived, it is the produce of some secreting organ. There are

^{*} Vide Memoires sur l'Oeil, par Janin, p. 412.

numerous examples of the natural secretion of surfaces being altered by diseases. It is very remarkable in the mucous membranes, and in those of the serous class, to which the membrane which contains and secrets the aqueous humour belongs. When they are inflamed, it is a very common morbid appearance to observe their surfaces covered with a matter, varying from the consistence of a thick coagulated lymph to that of a thin yellow puriform fluid.

on any specific differences in the nature

CHAP. VII.

OF ULCERS OF THE CORNEA*.

ULCERS in the cornea have been divided by some authors into a number of species, from differences in their size, in their duration, in the degree of the severity of the accompanying symptoms, and from the various causes from which they have been supposed to originate. But as these divisions are not founded on any specific differences in the nature of the disease, and as, instead of elucidating the subject, and assisting us in our inquiries, they lead to erroneous conclusions, and ren-

^{*} Helcoma .- Das Geschwüre der Hornhaut of the Germans.

der more complex a subject in itself simple, I shall omit mentioning them, referring those who wish for information on the subject to the works of Wallis*, Maitre-Jan, Mauchart †, and Rowley ‡.

The most frequent variety of ulcer of the cornea is that which remains after the cornea has suppurated and burst, either in consequence of a pustule or of an abscess. When a pustule suppurates, the central part of it generally first gives way, and, as the disease continues, the ulceration extends in all directions from that point. Ulcers of this kind are generally circular, and the edges rounded and smooth, having sometimes the appearance of a small artificial dimple §; in other instances, they have an irregular shape, and

^{*} A Treatise on the Diseases of the Eye, by George Wallis, M. D. 1785.

⁺ Burcard, David Mauchart, de ulceribus corneæ. Tubingæ, 1742.

[‡] A Treatise on the Diseases of the Eye and Eyelids, by William Rowley, M. D. 1790.

[§] See Plate IV. fig. 3.

their edges are more jagged and acute. Their size is very various; in some cases they do not appear larger than a depression made by the point of a pin; whilst in others they cover a much larger portion of the cornea. It seldom, however, happens that they spread over a large surface, for those of the most malignant kind are more apt to increase in depth than in breadth. They are generally superficial, and do not extend deeper than the external lamellæ; at other times, they penetrate the whole thickness of the cornea, and allow the aqueous humour to escape. Most frequently the part of the cornea contiguous to the ulcer becomes more or less obscure; and in some cases red vessels may be also traced in it. It is not unusual, however, to observe a small ulcer without any perceptible obscurity of the cornea.

The surface of the ulcer sometimes retains the natural transparency of the cornea; in some cases the whole, or a part of the surface of the ulcer, is covered with an opaque white matter, much resembling wet chalk *; and in others, where the ulceration spreads with great rapidity, the cornea loses not only its transparency, but its tenacity and firmness, and becomes like a piece of wet pasteboard, separating in the form of sloughs.

When the aqueous humour is discharged, from the whole thickness of the cornea having been destroyed, it often happens that the iris falls forwards, and a fold of it is insinuated through the opening in the cornea made by the ulcer. If the ulcer heals speedily, and the aqueous humour is regenerated, the iris sometimes resumes its natural situation; but more frequently an adhesion takes place between it and the cornea, so that ever afterwards the pupil remains drawn from its natural situation, and of an irregular form.— Sometimes the ulcer is long in healing, so that the aqueous humour continues constantly to ooze through it, and a fistula of the cornea is formed. In cases where a very great

^{*} See Plate II. fig. 2.

portion of the cornea is destroyed, not only the aqueous humour, but also the vitreous humour and crystalline lens make their escape, completely destroying the whole eye-ball. A woman, in whom, by a violent attack of inflammation, the cornea was destroyed and came off in large sloughs, a transparent tumour was formed by a prolapsus of the vitreous humour through the opening of the cornea; by which means she was enabled to see with considerable distinctness for several days, until the vitreous humour began to be absorbed, and the eye-ball to collapse. A case very similar to this was, some time ago, under Mr Thomson's care in the Royal Infirmary of Edinburgh. A man had a cancerous sore on the under eye-lid, which, in spreading, inflamed the eye-ball, and ulcerated nearly the whole cornea. Through this ulcer a portion of vitreous humour was pushed, forming a large transparent tumour, which enabled him, for several days, to distinguish minute objects with tolerable distinctness.

Ulcers often take place from the formation of abscesses, or collections of purulent matter betwixt the layers of the cornea. When the contained matter is evacuated by that part of the cornea being destroyed which formed its external covering, the ulcer which is produced has, like those already described, sometimes a colourless surface; at other times it is covered with an opake white matter, and it is subject to the same variations in form, size, depth, &c.

Ulcers sometimes, but very rarely, take place after wounds of the cornea.

They are seldom accompanied with much swelling of the cornea, except in children, in whom I have sometimes observed the cornea all round the ulcer considerably tumefied *.

They are generally accompanied with acute pain, which is much aggravated by exposure to light, or even by the most careful motion of the eye-lids. I recollect having seen only one instance, of an ulcer of

^{*} See Plate IV. fig. 3.

considerable size and depth, attended with little pain or uneasiness.

Ulcers, such as have been described, have a striking analogy to those which form on mucous surfaces, or on those parts of the body where the skin is very thin and inflected inwards *, as on the mouth, lips, internal surface of the nose, tip of the tongue, &c. The pustules or aphthæ † which precede them, have the same tendency to ulcerate rather than to form matter; they discharge an acrid serum instead of pus; they spread rapidly, and are also attended with acute pain.

Ulcers are also formed in consequence of the action of corrosive substances destroying the vitality of the cornea, and producing an ulcer where the dead portion separates from the living. Lime getting into the eyes is the most frequent accident which produces this effect; but the application of the nitrate of silver, muriate of antimony, &c. produces

^{*} Vide Scarpa Traité sur les Maladies des Yeux.

⁺ See Pustules of the Cornea.

one nearly similar. When lime falls within the eye-lids, those parts of the surface of the cornea to which it has been applied, become covered with an opaque white scale, accompanied by inflammation over all the external parts of the eye-ball and lids*; but the degree of thickness of the slough and violence of the inflammatory symptoms vary, according to the quantity of the lime, and the length of time it has been applied.

The circumstances attending this accident are strikingly illustrated in the case from which the drawing of fig. 3. Plate II. was taken. Nearly the whole external lamellæ of the cornea was destroyed from the application of the lime; and, from the small share of sensibility which the cornea possesses in its healthy state, the process of separation of the dead parts went on very slowly, and lasted several months.

After the violent inflammatory symptoms were subdued, the chalky matter began to

^{*} See Plate II. fig. 3.

separate at the union of the cornea with the sclerotic coat, and numerous small red vessels were seen at the place where the separation was going on. The process of separation proceeded from the circumference to the centre of the cornea—small flakes of the white matter could be daily observed to be coming away; and, after the lapse of several months, the whole disappeared, and the cornea regained nearly its natural transparency.

The healing process of ulcers shows that the cornea, like most other textures of the body, is capable of cicatrization. In most cases, this process seems to advance little farther after the principal part of the cavity of the ulcer is filled up, and a plain surface formed; so that, even after all the symptoms of disease have abated, a small dimple or inequality of the cornea remains. If the ulcer has been small or superficial, the depression is almost imperceptible; but when it has been of considerable size, the inequality of surface is distinct, and if it is situated opposite to the

pupil, it is very apt to render vision obscure. The portion of cicatrized cornea, it ought to be remarked, is not only unequal, but is by no means in all cases transparent, for it often happens, that an ulcer leaves a *speck* of the most opaque and incurable kind.

OF WOUNDS OF THE CORNEA.

ten without leaving the least trace of any oi-

by adhesion, a slight opacity only remains,

ly from the lips of the wound not being ad-

insted with mulicient acquisers at the times

tion has absted, and a few months elapsed,

the cicarrix appears like a bair sticking on

the cornea; and even this can only be disco-

se dimpuration to mences, the edges

OF THE COURSE.

The portion of cicatrized comes, it ought

by no means in all cases transparent, for it

CHAP. VIII.

OF WOUNDS OF THE CORNEA.

Wounds of the cornea unite readily, and often without leaving the least trace of any cicatrix behind them. When the wound heals by adhesion, a slight opacity only remains, and commonly there is nothing to be observed but a little elevation or inequality of the edges, which, in many cases, arises chiefly from the lips of the wound not being adjusted with sufficient accuracy at the time the wound was made. When all inflammation has abated, and a few months elapsed, the cicatrix appears like a hair sticking on the cornea; and even this can only be dis-

tinguished in certain lights by an attentive eye.

When a wound of the cornea, in place of healing by adhesion, goes through the more tedious process of suppuration, a considerable time elapses before the parts resume their natural form; and there always remains an obscurity, extending from the divided edges over more or less of the adjacent cornea.-When the suppuration commences, the edges of the wound swell, and are separated from one another, to a considerable distance, by a yellow, tough matter, resembling lymph, which sometimes hangs down from the wound in the form of flakes. As the healing process. goes on, the quantity of this matter diminishes, and the edges gradually approach each other, until a firm and complete cicatrix is formed. It is generally several weeks before this process is completed, supposing every thing to go on uninterrupted; but it often happens that, when wounds are accidentally inflicted, or even in the incision of the cornea, made in the operation of extracting the crystalline lens, a

portion of the iris falls forwards, and either adheres to the edges of the wound, or passes completely through it. This last accident is always attended with disagreeable effects; for, besides the permanent defect in the form of the pupil, and the distressing symptoms of pain and inflammation, which the strangulation of a portion of the iris never fails to produce, the lips of the wound are seldom allowed to close altogether, so that the aqueous humour drills out, until a very tedious process of cicatrisation is completed.

In most cases of wounds of the cornea, the opacity does not extend far beyond the edges of the wound, except in those cases where the inflammation has been long protracted. It seems to be confined to the cut surfaces; and in the incision made for the extraction of the cataract, where the knife runs between the layers of the cornea for a considerable extent, and does not make an incision perpendicular to the plane of the spherical surface, the remaining opacity is of a very considerable breadth, pointing out the precise form of the incision.

CHAP. IX.

OF FOREIGN BODIES ADHERING TO THE CORNEA.

When a foreign body once adheres to the cornea, all attempts in rubbing the eye-lids, or winking and shutting them forcibly, tend more to imbed it firmer, rather than to remove it. From the external layer of the cornea being soft and yielding, the foreign body, if it be small, soon forms to itself a seat, and the constant flow of tears and disposition to shut the eye-lids, produced by its irritation, soon bring on a violent inflammation, which never abates, until the body is either removed by art, or comes away by a tedious process of suppuration.

When the foreign body is removed by art, it leaves behind it a depression, which is often discoloured; the colouring matter, however, is soon absorbed, and the depression is generally filled up, and all surrounding opacity removed in a few weeks.

In a very few hours after a foreign body adheres to the cornea, the adjacent portion of cornea becomes opaque, and the opacity extends according to the violence of the inflammatory symptoms, which the irritation of the new substance creates. I have observed this opacity form very rapidly, and to a great extent, in the eyes of animals, from a similar cause; but in animals, opacities are more rapidly formed, and are more speedily removed.

Sometimes a foreign body remains imbedded in the cornea for a long time, and is the source of constant inflammation and pain, till suppuration takes place around it, and allows it to drop out.

It sometimes happens that, after a body is imbedded in the cornea, a layer of a new substance is formed over it, so that it does not excite any inflammation, but remains through life in a kind of sac. I have observed this process begin and be completed in a case where a small portion of the iris had been pushed through an ulcer of the cornea. The cornea near the prolapsed iris became obscure, and the opaque matter was daily effused from the circumference towards the centre of the opening, so as finally to cover the prolapsed iris so completely, that it appeared afterwards like a common speck of the cornea.

It is by no means an uncommon thing to hear of extraneous substances, as musket-balls, &c. remaining during life in different parts of the body, by forming to themselves, in a similar manner, a kind of sac. In one case, I found a piece of whin-stone inclosed in a sac of cellular membrane, lying close to the sclerotic coat, which had remained for ten years prior to the person's death, without his experiencing the least uneasiness from it, or even suspecting its presence.

Manniske of Frankenhausen mentions a curious case in Loder's Journal *, where a body, which stuck in the conjunctiva covering the white of the eye, gradually advanced to the central part of the cornea. I shall quote the case in his own words:

"A priest requested my assistance concerning a speck on the eye, in the year 1792. He had on the cornea of the right eye a dark speck, which impaired much his vision, of which he gave me the following account: Two years before, he found suddenly a little pain in the eye. By examination he remarked, on the white of the eye, below the upper-lid, a black spot; it did not hurt his sight, and the pain soon went away, so he took no further notice of the accident. Some time having elapsed, he was aware that this spot had changed its situation, and had appeared at the union of the cornea with the sclerotic coat. Now the case appeared to

^{*} Vide Journal für die Chirurgie, &c. Von Just. C. Loder, 2d. Band. 1st. Stuck. 1799.

him doubtful. He asked the opinions of many medical men, and made use of many external and internal medicines, without any effect. The speck continued its progress very slowly, but uninterruptedly; it came forwards on the cornea, approached towards the pupil, and at last covered a portion of it. The patient was in this situation when I saw him. There was a prominent spot above the cornea, which felt hard, and was the size of a small lens, but longer than it was broad. Many small red vessels appeared like streaks around it. The patient had no pain. The undescribable hardness of the spot, along with its situation, made me think that it was a foreign body fastened in the eye. I made an incision on the spot from without inward, and saw, with the assistance of a microscope, a black body lying in the incision. I removed it with the point of the knife, from the small hole it had formed for itself in the cornea, and found it to be a hard wing-case of a beetle."

CHAP. X.

OF OSSIFICATION OF THE CORNEA.

The deposition of bony matter is a morbid change, which has been met with in almost every organ of the body. I have seen only one instance of it in the cornea; and in that case, the whole eye was changed in its form, and the cornea had become opaque. On macerating the cornea, a piece of bone, weighing two grains, oval shaped, hard, and with a smooth surface, was found between its lamellæ. A piece of bone was also found between the choroid coat and retina of the same eye. When dissecting a cornea for an ana-

tomical purpose, of which no history could be obtained, I found several gritty particles and inequalities on its internal surface.

Walter had in his museum a piece of cornea, taken from a man sixty years of age, in which a bony mass was enveloped, of a long shape, three lines in length, two lines broad, and weighing two grains *.

A very curious case, of a piece of bony matter, which was formed in the substance of the cornea, or immediately behind it, was communicated to me by Mr Anderson, Surgeon at Inverary. "The patient was a woman of about 31 years of age. Upon carefully examining her right eye, I observed a substance of a whitish appearance, in the under part of the globe of the eye, arising from the inside of the sclerotic coat, and extending upwards, below the cornea, over a great part of the iris, to very near the pupil. It had seemingly communicated much irritation to the eye, and induced a degree

^{*} Anatom. Museum, B. I. S. 139. No. 275.

of inflammation, severe pain, almost a constant flow of tears, inability to bear the light, with a considerable diminution of sight. The ball of the eye performed its natural rotatory motion, but was less in size than the left eye. This complaint had been occasioned fifteen years ago, by a fall at the root of a tree, by which one of the roots struck against her right eye, but did not cut any part of it.

From this period, the said substance had begun to grow, and gradually increased in size; but the pain in the eye, and other symptoms, were sufferable until about nine months ago, when the complaint became more violent. I advised her to submit to an operation, for the purpose of extracting the said substance that appeared to injure her eye, to which she readily agreed. I made an incision into the cornea, in the manner recommended for the extraction of the cataract, then raised the flap of the cornea with a flat crooked probe, and, with the same instrument, turned out a small piece of bone. The upper part of the bone

was as thin as a piece of paper; at the under part it was thicker, porous and brittle, of an irregular semilunar form, and about the size of half an ordinary silver sixpence. The upper part was quite detached, but the under part slightly adhered to some part of the globe out of sight; but it was easily extracted without requiring the use of the knife to separate its adhesions. From the unsteadiness of the patient, she would not permit me to examine from what part the ossification originated. I am consequently at a loss, whether to suppose it took its growth from any of the coats of the eye, or if any osseous matter might, in consequence of the accident, collect within the said coats, and, in the course of fifteen years, form a complete bone."

The cornea varies very much in its firmness of texture, at different periods of life, and in different individuals, at the same age. In the fœtus, its cellular lamellæ adhere to one another loosely, so that it is thick and spongy, whilst in old people it is sometimes extremely hard and coriaceous: And Angely * remarks, that he has sometimes met with it in old people as hard as a piece of wet cartilage.

* Commentatio Medica de Oculo, originisque lacrymalibus.

Auctore, J. L. Angely.—Erlangæ, 1803.

CHAP XI.

OF THE SPECK OF THE CORNEA *.

1. Of the Varieties of the Speck of the Cornea.

Although the cornea is naturally quite transparent, yet it often happens, that the means which are employed to preserve it in that state, are not sufficient to prevent it from being rendered obscure by disease. The obscurities or specks to which it is liable, are observed to be of very different shades, varying from the slightest perceptible cloudiness or mist, to a dense white, or

^{*} Macula Corneæ, - Die Verdunklung der Hornhaut, or Die Flecke of the Germans.

pearl-coloured opacity. These great varieties in the degree of the obscurity, as well as those which occur in the shape, the mode of formation, the relative position and extent of corneal specks, have induced some authors to consider them as distinct species, and to distinguish them by particular names. As the meaning, however, of these names is very ambiguous, there being scarcely two authors who use the same word to denote. the same variety of the disease; and also, as it will appear that there is such a close resemblance between some, and such a similarity of character in all the forms in which specks appear, these arrangements are not only useless, but inconsistent. Instead, therefore, of introducing a variety of names, or attempting to arrange in a systematic manner, the varieties of this morbid alteration of structure, we shall consider it in whatever form it may appear, or in whatever degree, under the general name of obscurity of the cornea, or corneal speck.

The first, and most simple variety or form of corneal speck is, when a particular part of the cornea loses its natural transparency, and appears clouded, objects appearing to the patient as if seen through a mist or smoke. Such obscurities are either undefined, or distinctly circumscribed, and have either an equal degree of opacity throughout, or one part is more opaque than the rest. They are most commonly of a circular or rounded form, but, in some cases, their shape is very irregular. Their size varies from the smallest spot to such an extent as to occupy the whole cornea *.

In a second form of the corneal speck the opacity is of a darker shade; it gives the cornea a bluish, or, in some parts, a white milky appearance. It is seldom equally opaque through its whole extent, being generally more so at the centre, and becoming gradually of a lighter shade towards the mar-

^{*} See Plate V. fig 1.

gin. In some instances, the shade is very unequal in the different parts of the speck *.

In a third form of the corneal speck, the cornea becomes of the opaque glistening white colour of common pearl; and the opacity generally extends through the whole of the lamellæ of the cornea, so that if even several of those layers which are external be removed, the remaining ones continue to form a complete obstruction to the entrance of light. Specks of this description sometimes produce a slight thickening of the cornea, and are accompanied with adhesions between the cornea and iris. They are almost always distinctly circumscribed, and, if of any considerable size, they are nourished by one or more red vessels †.

In the first form of speck, the iris can be seen through the diseased portion of the cornea; but, in the second and third forms of the disease, the degree of opacity is such,

^{*} See Plate V fig. 2.

⁺ See Plate V. fig 3, and Plate VI. fig. 1 and 3.

that nothing can be accurately distinguished behind it. If there is an active inflammation accompanying the speck, the red vessels will be seen in a cluster on that part of the sclerotic coat nearest to it; and some of the branches may often be traced passing over the edge of the cornea, and terminating in the substance of the speck. As the accompanying inflammation abates, the number of the red vessels on the cornea commonly diminishes; but sometimes one or more trunks remain, and are distributed on the speck. In some cases there are large specks, with numerous bloodvessels supplying them during the continuance of active inflammation, and although the opacity remains extensive after the inflammation abates, yet no red vessels continue to nourish it. The number of bloodvessels is in no case in proportion to the extent or degree of the opacity during any stage of the accompanying inflammation; for we frequently observe a net-work of blood-vessels on a cornea which has very little obscurity, and at other times there is a large opaque spot with only one, or even without a single red vessel supplying it.

Specks appear on every part of the cornea, but, as far as I have been able to observe, they occur most frequently towards its centre.

They vary in number: Commonly there is only one; but it frequently happens that there are two, three, or more distinct spots on one cornea, all of which differ in their size, shape, and in the degree of opacity.

Specks impede the function of vision in proportion to the degree of their obscurity, and according to their situation. Even a speck of the lightest shade, which is hardly perceptible to a common observer, if it is placed directly opposite the pupil, materially injures the sight; whereas those of the most opaque kind, if placed beyond its circumference, diminish the sphere, but not the accuracy of vision. In those cases where the speck is of a moderate size, and placed towards the centre of the cornea, the patient sees better in a

dull than in a clear light; for in a clear light the pupil contracts so much that it is covered by the speck, and the rays of light are prevented from entering it; but in a dull light it dilates and becomes larger than the speck, so that the rays of light enter by its edge. Specks impede vision more when they are situated on the under than on the upper half of the cornea.

Those who have a speck on the cornea are apt to have a squint. In one case, this was so remarkable, that I found it impossible for the eye-ball to be turned, so as to render more than one half of the cornea visible.

In some cases, where the speck is situated towards one side of the cornea, the pupil seems to have a tendency to dilate towards that portion which remains transparent. I observed this in an adult who had completely lost the sight of one eye, and had a large speck formed on the temporal and central part of the cornea of the other eye soon after birth. The pupil was very considerably dilated towards the nasal part,

which remained transparent; so that, by this effort of nature in drawing the pupillar opening from the opaque to the transparent part of the cornea, the patient was enabled to guide himself through the streets, and in twilight he could see large objects around him with considerable accuracy. I observed the same thing also take place in a young woman's eye, under precisely similar circumstances.

Specks seldom produce any very sensible alteration in the external form of the cornea. In some cases where the inflammation has been very violent, or where the disease has been of long duration, the cornea becomes thickened, so that its internal surface comes in contact with the iris, and adheres to it, suffering at the same time a slight increase in its convexity; and in many cases a cornea, which for some time was only opaque, becomes at last staphylomatous *.

Besides those varieties of specks of the cornea which have been described, there are

^{*} Vide Staphyloma, and Plate VI. fig. 3.

others that occur much less frequently, the mode of formation and appearances of which are somewhat different.

I have remarked a few instances where the cornea acquired a very peculiar mottled appearance. In one case this was the consequence of an inflammation of the eye, which came on during a mercurial course, and, in another, it was the effect of an inflammation brought on by lightning. In both these cases, some parts of the cornea retained their transparency, while the rest of it was covered by a number of small white rounded spots, varying in their degree of obscurity.

In some cases the opacity, instead of being formed towards the central part of the cornea, or at some distance from its circumference, begins at the place of the junction of the cornea and sclerotic coat, and gradually extends towards the centre of the cornea, forming an opaque ring around its circumference. Most authors have described this as an appearance only to be remarked in the eyes of old people,

and have given it the name of arcus senilis. But I have observed it at all periods of life, and it may be seen in several of the drawings which were taken from young subjects *. It generally appears like a bluish ring at the edge of the cornea, running round the whole circumference. It is almost universally to be found in the eyes of old people, and it increases in breadth as the person advances in life; but it is never attended with any impediment or inconvenience to vision.

In a few instances I have observed a cloudiness gradually extending from several points, or from the whole circumference of the cornea towards its centre, which, in most cases, went off along with the inflammatory symptoms by which it was accompanied; but in others it was more permanent.

^{*} See Plate I. fig. 1, Plate III. fig. 3, Plate V. fig. 3.

2.—Of the Formation of Specks of the Cornea.

Specks most commonly are either preceded or accompanied by inflammation of the cornea. In children they occur very frequently during the progress of the more severe cases of the purulent ophthalmia, also in small-pox and measles, in that peculiar inflammation of the eyes which is accompanied with eruptions of the head, and in almost all those inflammations in which the cornea participates. Lues venerea and scrofula, each of which produce specific inflammations of the different parts of the eye-ball, also occasion various degrees of obscurity of the cornea. Likewise wounds, accidentally inflicted on the cornea, or in the operation of extraction of the cataract, if they do not unite without suppuration, generally leave a very opaque mark; and ulcers, if they have been deep, or of long duration, are followed by a white pearl-coloured speck.

When the inflammation accompanying a speck abates, the speck most commonly diminishes, and the opacity which remains becomes more distinctly defined, and less opaque, or, if it is small, it entirely disappears.

In some instances the cornea acquires a very remarkable degree of obscurity, when the inflammatory symptoms are apparently mild, and where there is very little perceptible redness. I have observed several instances of this kind, and in all of them the obscurity came on by very slow degrees, and was attended by no pain. In the 2d vol. of the London Medical Communications, there is a description and history of two children by Mr Farr of Deptford, where the corneæ acquired a very remarkable degree of obscurity, without much apparent inflammation. A very accurate account of several cases of this kind is also given by Mr Ware of London, in his excellent Treatise on Ophthalmia.

The length of time necessary for the formation of specks varies much in different instances. In some cases they are formed very slowly, and do not acquire any great degree of opacity, even when the inflammatory symptoms are extremely violent. Most frequently they require several weeks or even months before they become either large or very opaque; so that when they are of very considerable size, they have generally been formed after repeated inflammatory attacks, each attack adding to their size and degree of opacity. Sometimes they are formed very rapidly, as in cases where the cornea is wounded, or where a foreign body has adhered to it.

Specks are formed at every period of life, but they occur much more frequently in young people; probably because in them the cornea is much softer and more spongy, and, also, as they are more subject to various inflammatory complaints of the eye. The specks which are formed rapidly are generally most speedily removed, whilst those whose

progress is slower disappear in a very gradual manner. They are removed much more quickly in children than in old people, and in them also a much greater degree of obscurity can be made altogether to disappear. When applications are employed with the view of removing specks, the obscurity generally diminishes much more rapidly for a short time at first than at any future period. Dr Vetch, in his ingenious treatise on the Egyptian Ophthalmia *, gives an account of a singular instance, which shews the rapidity with which some kinds of specks disappear under particular circumstances. It occurred in a man during his convalescence from ophthalmia. "Some pectoral symptoms, to which he had been long subject, suddenly assumed the appearance of pulmonary consumption, which proceeded in a rapid manner towards its last stage: five days previous to his death, he was seized with a violent

^{*} See an Account of an Ophthalmia, &c. by John Vetch, M. D. 1807.

aggravation of the hectic fever and the other symptoms; and his death was hourly expected. At this time, to the surprise of all his attendants, the opacities, by which the vision in both eyes had been long obstructed, disappeared with an amazing rapidity; and, a short time before his death, his vision became nearly as distinct as ever." -A case, though of a different kind from that now quoted, deserves to be mentioned in this place, as it strikingly illustrates the same remarkable power in the absorbent system. The patient had a very perceptible obscurity of the crystalline lenses of both eyes, which had considerably impaired his sight. This complaint had continued for several months, when he was seized with a pain of the breast, attended with fever and spitting of blood. These symptoms continued for several weeks, and, during that period, the obscurity of the crystalline lenses altogether disappeared, and his vision was restored. I saw him many months afterwards, when his eyes continued well.

3.—Appearances of Speck on Dissection.

WHEN the cornea is examined after death, no change of structure can be observed in those cases where there had been a mere cloudiness, or general opacity during life; for even before death, more especially if it is slow and lingering, the fluid which, in the natural state, is deposited between the lamellæ of the cornea, exudes, forming an obscure layer over its anterior surface, and the aqueous humour oozes out, giving the cornea an unequal puckered appearance *. Indeed, it is from this change in the eye that approaching death is often foreseen; for whenever the cornea begins to collapse, and becomes turbid, the eye loses all its lustre and intelligence, and gives that awful expression to the whole countenance, which has been called facies hypocratica.

^{*} Vide Anatomie Descriptive, par Xav. Bichât.

When the cornea has been much more opaque, no other change is to be perceived after death, than a diminution of the transparency, either of the external lamellæ, or of the whole substance of the cornea. I have had many opportunities, in the living body, of taking off layers of very opaque specks, and I have never been able to observe any other change, except that, in some of those which have been of long duration, the cornea had acquired a degree of hardness much greater than that of a sound cornea.

In most cases, too, a speck bleeds when a piece of it is removed in the living body; and I have observed this happen, even when no red vessels passing into it could be detected by the naked eye. An incision made in the healthy cornea, gives little or no uneasiness, but, when the portion of a speck is removed, it often excites acute pain.

4.—Causes of Specks of the Cornea.

As the deposition or effusion of the albuminous part of the blood is a common effect of inflammation in many organs of the body; and as this change produces a diminution of the transparency of the serous membranes, it is probable, from the analogy in the natural structure, that it is a similar change which takes place in the cornea, during the formation of specks. This effect of inflammation is very remarkable in the pleura, after an attack of pleurisy; in the peritoneum after peritonitis; and in the membranes of the brain after phrenitis; for on dissection it is invariably found, that there is not only an effusion of albumen on the surface of the inflamed membrane, but the membrane has become thicker, changed its colour, and lost its transparency.

It is perhaps difficult to determine, whether this new matter, added to the diseased membrane, be effused among its layers, but it appears to me the most probable opinion. It has been already remarked *, that the cornea is composed of a number of concentric cellular lamellæ, and that there is a fluid deposited in the cells by exhalation. It is, therefore, in the cellular structure in which this fluid is contained, that the albumen may be deposited, the cells being like as many serous surfaces, and therefore subject to similar morbid changes.

An opacity may also be produced by the cornea losing its vitality. This change may arise either from the inflammation of the cornea being so violent as to terminate in the death of the part, or from the action of caustic substances, such as lime, lunar caustic, &c †.

It is also probable, that the cornea may become obscure, in consequence of an alteration in the quantity of the contents of the eye-ball, producing a change in the arrangement of its component particles. In the

^{*} See General Remarks on the Cornea.

⁺ See Ulcer of the Cornea.

dead body this change is very remarkable, for if pressure be applied to the eye-ball, or if the ophthalmic veins be injected with quicksilver, or pure water, so that the quantity of the contents of the eye-ball is increased, the cornea is found to lose its natural transparency, and to acquire a milky colour *. As it was probable, from this curious phenomenon in the dead eye, that, in the living body, some opacities of the cornea might arise from an increase in the quantity of the contents of the eye-ball, it occurred to me, that, in cases of this kind, the opacity might be removed, by making an incision through the cornea, and discharging the aqueous humour. That a certain degree of opacity of the cornea in the human eye, is sometimes produced by a mere derangement of its component particles, is proved from the immediate effects which followed the discharge of the aqueous humour, in cases where the opacity had a ge-

^{*} See The Muscular Motions of the Human Body, by John Barclay, M. D. 1808.

neral clouded appearance over the whole cornea, besides some defined spots at particular places. In some cases of this kind, the instant I discharged the aqueous humour, by a small opening made in the cornea, I observed that all the general obscurity disappeared, and nothing remained but the more opaque spots, which became more distinctly circumscribed *.

In the lower animals, I have had several opportunities of observing the cornea affected with a general obscurity and cloudiness, which was not like common speck, but more resembled that obscurity so easily produced artificially in the dead eye, by injecting the veins, or by the application of pressure. A disease of the cornea very similar, is common among sheep which have made long journies, or have been much fatigued; and although the opacity of the cornea in them is

G

^{*} See Observations on the Effects of Evacuating the Aqueous Humour in Inflammation of the Eyes, in the Edinburgh Medical and Surgical Journal for January 1807.

accompanied with little apparent inflammation, or fulness of vessels, its transparency is quickly restored by the common practice of the shepherds, which is opening a vein at the inferior part of the orbit, and allowing the blood to flow over the eye *. The same kind of obscurity of the cornea is by no means unfrequent among dogs and horses.

^{*} See Dr Duncan's Essay on the Diseases of Sheep, in the Transactions of the Highland Society of Scotland,

CHAP. XII.

OF THE STAPHYLOMA.

When the cornea, besides losing its transparency, swells to such a degree that its internal surface comes in contact with and adheres to the iris, and when it forms a prominent tumour externally, the disease has generally been called Staphyloma *.

This term, however, has had very extensive applications, having been employed by some authors to denote not only various morbid changes of the cornea, but also a va-

^{*} Das Staphylom, - Der Forfall der Hornhaut of Richter.

riety of tumours involving other parts of the organ of vision. I shall, follow Richter, and limit its signification to those changes which produce an alteration in the structure and form of the cornea. When the structure of the whole cornea is changed, it has been called the *staphyloma totale*, and when the disease is confined to a particular part, *staphyloma partiale*.

In the staphyloma, where the whole cornea participates in the disease, it generally assumes a more or less conical form, loses entirely its natural transparency, and vision is completely destroyed. The opacity is often most remarkable towards the apex of the tumour, and is generally of a pearl-white colour, diffused through the whole corneal substance *. In some instances I have observed the opacity confined to one-half of the cornea, and this was generally the inferior one †.

^{*} See Plate VI. fig. 2

⁺ Plate VI. fig. 3.

The internal surface of the cornea adheres to the iris in almost every case of staphyloma; for as the internal lamellæ, as well as those which are external, increase in thickness, they come in contact with the iris, and the accompanying inflammation produces a permanent adhesion between them; so that, in this disease, the anterior chamber is often found almost entirely obliterated. This happens most frequently in children, as in them the cornea is much thicker than in adults, and is very nearly in contact with the iris even in the sound eye.

In staphyloma, the pupil is hid according to the situation and degree of the opacity of the cornea; but in most cases it is altogether obliterated, and even in those where a transparent portion of the cornea is opposite to it, the vision is much impaired; for as the eye has lost its form as an optical instrument, the change in its refractive power must render objects seen through it very indistinct.

If the tumour is recent, and has not acquired a very large size, the diseased por-

tion of the cornea, when cut into, will be found to be softer and more spongy than the transparent part; and generally, if the incision penetrates into the anterior chamber, only a very small quantity of aqueous humour flows out. This is more particularly the case in children. For when the tumour is large, and when it forms in an adult, the cornea often becomes extremely hard, and much thinner than natural, and the quantity of aqueous humour, instead of being diminished, is greater than in the sound eye. Scarpa mentions his having found the cornea, in some cases of staphyloma, as hard as parchment, or even converted into bone: Richter* mentions a case where it resembled a cartilaginous excrescence, and Beer + describes and gives a delineation of one which was so thick and tough, that, in cutting it away, he could scarcely penetrate it with the

^{*} Chirurgische Bibliothek, Vol. 8. S. 76.

⁺ Practische Beobachtungen über den grauen Staar und die krankheiten der Hornhaut. Wien 1791.

knife. There is generally one or more red vessels ramified over the sound part of the cornea (vasa nutrientia), which are continuations of vessels on the sclerotic coat *.

After an wound has been made through the cornea, penetrating into the anterior chamber, and a small portion of cornea removed, so as to prevent it from healing up immediately, a violent inflammation generally ensues. A suppuration afterwards takes place, and the sore gradually heals. Sometimes, even after this operation, the tumour continues to increase in bulk, but, in other cases, it produces a permanent diminution and collapse of the whole organ. In a case where a small portion of the apex of a large staphyloma was removed, and scarcely any humour could be observed to escape during the operation, and when, consequently, it produced no immediate change upon its bulk, the contents of the eye soon after began to be absorbed,

^{*} See Plate VI. fig. 2, and Plate VII. fig. 3.

and in a few months were so much diminished, that the eye-ball could be completely covered by the eye-lids.

When vision is altogether destroyed, and the disease has extended far, the humours collect in great quantities, and, in some cases, form a tumour of an enormous size *. In such cases the eye does not retain its conical form; for when the cornea has become so much distended, and when the humours have insinuated themselves between its lamellæ, one part yields more readily than another; the tumour becomes more irregular in its form, hangs over the under eye-lid, and, in many cases, has the appearance of a large swelling composed of several smaller ones. A change also takes place in the colour of the tumour. Instead of the opaque white, or pearly coloured opacity, it becomes of a dark blue colour, and sometimes one or more of the smaller swellings are semi-transparent. A fluid can be felt fluctuating within them,

^{*} Haller's Dissertationes Chirurg. Tom. 1. p. 25.

and they appear as if ready to burst. The blood-vessels increase both in number and size along with the bulk of the tumour, and they will sometimes be seen ramifying and forming net-works in a most beautiful manner over its whole surface *. If the bulk of the swelling prevents the eye-lids from closing, the exposure to bodies floating in the air, and the contact of the ciliæ excite always more or less inflammation over the eye-ball and lids; and at the same time it often happens, that the tears are of an acrid and irritating quality, inflaming and excoriating the outer surface of the palpebræ and cheek. In this situation the patient is often relieved by a part of the tumour giving way, and allowing the contents of the eye-ball to escape. Even when this has taken place, and after the inflammation which it has occasioned has abated, the disease sometimes returns as before, and a large staphylomatous tumour is again formed. More fre-

^{*} See Plate VII. fig. 3.

quently, however, the organ remains collapsed into a whitish mass, in which no vestige remains of the natural structure.

The sudden relief which is produced by the discharge of the contents of an eye affected with staphyloma, may be accounted for on the same principles as the practice of evacuating the aqueous humour in violent inflammation of the eyes, or the rupture of a common abscess. This effect is so strikingly illustrated, and so accurately described in the following case of a medical gentleman who had been seized with violent ophthalmia which brought on staphyloma, and deprived him of his sight, that I shall give the account of his case in his own words. "When my eyes were examined, I was told that there was a considerable elevation of the cornea of both of them, but particularly of the left; indeed this was quite evident to my own sense of touch. In July 1800 I returned to England; and, on my voyage from London to Edinburgh, I was much surprised one morning when I awaked,

to find that the aqueous humour was completely discharged from the left eye; the eye being quite soft. The opening by which it had escaped soon closed, and the humour again collected; but I conceived that the cornea was not quite so prominent as formerly. From this circumstance I concluded, that if the humour were again evacuated, the eye might assume its natural form, or at least approach nearer to it. In March 1801, I attempted to make a small perforation in the cornea with a large needle, and succeeded easily in evacuating the aqueous humour. This, I think, was repeated at least three times, and the eye gradually acquired its present shape. I wished to perform the same upon the right eye, but was baffled, on account of its greater sensibility. If I can place any dependence on my own feelings, I think the left eye was never perfectly free of inflammation until the above mentioned evacuations were effected; though they were made without having the most distant idea that such would be the result of them.

About three years ago, when rubbing the right eye with my finger, it suddenly became soft, part of the aqueous humour having been discharged. Since that time, two or three drops have flowed out in the space of every week. But if, as it sometimes happens, the discharge does not take place in the space mentioned, I am in general seized with a pain immediately above the right eye-brow, accompanied with the sensation of tension, and uneasiness of the eye itself: These symptoms, however, vanish when the accidental evacuation takes place.

When the humour was first discharged, a small degree of pressure upon the eye, when soft, gave considerable uneasiness; and the pain seemed to be situated in the bottom of the eye. The pain, however, is not now excited by the same degree of pressure."

After a staphyloma has burst, and a new tumour is formed, it commonly resembles the original one in every respect; and in some cases, this alternate discharge and regeneration has taken place two or three times before the eye finally collapsed. Mr Thomson informs me that he has remarked several cases, where a fistulous opening remained long after the tumour burst, through which a humour, resembling the aqueous, continued to be discharged.

Staphyloma most frequently occurs in children after the small-pox, measles, purulent ophthalmia, or violent inflammation of the eyes from other causes. The cornea being in them more spongy in its texture, and much thicker than in adults, is probably the cause why, as in specks, they are more subject to this disease *. Scarpa mentions, that he never saw a staphyloma of such a size as to project beyond the eye-lids, which did not begin during childhood. I have, however, seen several instances, particularly where the disease arose in consequence of an wound, in which the tumour extended beyond the eye-lids, and did not commence till the pa-

^{*} Angely, Commentatio Medica de Oculo, &c. Erlangæ, 1803.

no doubt rare. Staphylomas which arise after wounds in the eye with sharp pointed instruments, present nearly the same appearances as those which have been already described. In such cases, the swelling is generally more irregular in its shape, the coats of it are thinner, the disease advances more rapidly, and is attended with much more pain and inflammation.

Staphyloma generally attacks only one eye; in some cases it affects both, and it is not unfrequent to see a person with two staphylomas, in consequence of small-pox, &c. I have seen one case, where the sympathy between the two eyes was very remarkable. The person received a blow with a pike on one eye, which produced staphyloma, and more than a year afterwards, the other eye became inflamed, and the cornea of it also gradually became staphylomatous.

Staphylomas are variable in their progress; sometimes they grow suddenly to a certain size, and afterwards remain stationary; some-

times they grow progressively larger till they burst; and often they increase in bulk as if by starts. From small-pox particularly, and also in consequence of the purulent ophthalmia, staphylomas grow rapidly, and the coats of the tumour appear extremely thin; but, after some time, they seem to acquire additional thickness and strength, the tumour remaining of the same size.—Richter* remarks, that the swelling and thickening of the cornea sometimes go away with the accompanying inflammation.

Staphylomas sometimes also occur, where a small portion of the cornea is affected, in which case they are called *Staphyloma partiale*. Sometimes there is only one swelling of this kind, and sometimes several of them arise in the same cornea, and form an irregular shaped mass, which has been compared to a cluster of grapes; hence the name *staphyloma*. Such swellings vary from the size of a pin head to that of a small

^{*} Richter's Anfangsgründe.

pea. In some cases they are transparent, communicate with the anterior chamber, and contain a quantity of the aqueous humour, or even a portion of the iris. More commonly they are horny or warty excrescences, and if cut off, they generally grow again *.

Tumours of this kind are often observed where there have been wounds or ulcers of the cornea; but they are also met with where there has been no such previous cause.

There is also a formidable disease, a few examples of which I have seen, and traced the progress, which, perhaps, may be with more propriety considered as a variety of staphyloma than any other disease. It occurred in adults, whose eyes had been exposed to much fatigue, and suffered from deep-seated, long continued inflammation; and I have seen one or two cases of this disease, where, from its commencement, and during its whole progress, there was neither pain nor any inflammatory symptom. Whilst the sphericity of

^{*} Richter's Anfangsgründe.

the cornea increased, the aqueous humour became turbid, and substances were seen floating in it, resembling flakes of the black pigment. The crystalline lens also became opaque, and appeared to separate into pieces, and moulder down. The sclerotic coat became preternaturally distended, and instead of retaining its pearly white colour, it assumed a dark blue, or almost black shade. At last, a large prominent tumour was formed, more spherical than the common staphyloma, which terminated by bursting, and in a total loss of the organ.

CHAP. XIII.

OF ALTERATIONS IN THE FORM OF THE CORNEA.

The form of the cornea varies in different people, and in the same individual at the different periods of life. In many it is so convex, that objects are seen very indistinctly, unless when held close to the eye, or viewed with the assistance of concave glasses. This preternatural convexity of the cornea is sometimes the failure of original organization, at other times it is produced from some morbid change. The cornea is always more convex at the earlier periods of life, and, as far as I have been able to ob-

serve, it is very liable to acquire its greatest degree of convexity about the age of puberty, people being most apt to become *short-sighted* at that period.

When people advance in life, the cornea gradually loses its convexity, in consequence, perhaps, of the quantity of humours within the eye-ball being diminished. In one case, of a girl eight years of age, the corneæ of both eyes were observed to be remarkably flat, and her vision very imperfect from her infancy. In people much enfeebled by considerable evacuations, by numerous bleedings, or by disease, the quantity of the aqueous humour diminishes, the convexity of the cornea is lessened, and their sight is enfeebled, so that they are not able to see objects but at a distance *.

The cornea is also observed to collapse at the approach of death, particularly if it is slow and lingering †.

^{*} Vide Anatomie Medicale, par Portal.

⁺ Vide Anatomie Descriptive, par Xav. Bichât.

From the ingenious experiments of Mr Everard Home, and the late Mr Ramsden, recorded in the Philosophical Transactions, it appears that the sphericity of the cornea is altered according to the distance at which objects are viewed.

The power which the eye has of changing its form, is also very remarkable in those people who have had the operation of the extraction of the cataract performed; for, when the eye has recovered from the inflammation and irritability occasioned by that operation, so as to allow the person to look at objects, it requires, most commonly, the assistance of a lens of two and a half inches focus to read a common printed book; but, if after a few weeks, a lens of a lesser power be used, the eye, by gradually accommodating itself to the change, will see very distinctly with a glass of a much smaller power than what was first necessary. Thus a person who has had this operation performed, generally finds his sight improve for many months after it; and the change appears to arise chiefly from the cornea gradually acquiring a greater degree of convexity, from the action of the muscles of the eye-ball.

Besides these slighter changes in the form of the cornea, it sometimes increases to such a size, or collapses so much, whilst, at the same time, it retains its transparency, that the functions of the eye are much interrupted, or even entirely destroyed. In the first case, the disease has been called by some authors the *staphyloma pellucidum*; in the second *Rhytidosis*.

Levéillé †, the French translator of Scarpa's work on the diseases of the eye, has described a case where the cornea of both eyes became of a conical form. Two examples of a similar disease have fallen within my own observation; but only one eye was affected in each of them. In both cases, the conical figure of the cornea was very remarkable, and the apex of the cone was in the centre of the cornea. When the eye was viewed

⁺ Vide Traité sur les Maladies des Yeux, traduit de l'Italien de Scarpa, par J. B. F. Levéillé, Tom. II. p. 179.

laterally, the apex resembled a piece of solid crystal, and when looked at directly opposite, it had a transparent sparkling appearance, which prevented the pupil and iris from being distincly seen.

One of these cases occurred in a lady upwards of thirty years of age, and the changes produced in her vision were very remarkable. At one, or one and a half inch distance, she could distinguish small objects distinctly, when held towards the temporal angle of the eye, although it required considerable exertion; but the sphere of vision was very limited.

On looking through a small hole in a card, she could distinguish objects held very close to the eye, and could even read a book.

At any distance greater than two inches vision was very indistinct, and at a few feet she could neither judge of the distance nor the form of an object.

When she looked at a luminous body at a distance from her eye, such as a candle, it was multiplied five or six times, and all the

images were more or less indistinct. She could never find any glass sufficiently concave to assist her vision. She did not remark this complaint in her eye, until she was about sixteen years of age, and she does not think that it has undergone any change since that time.

On mentioning to my ingenious friend Dr Brewster this case, which appeared to me so remarkable, he had the politeness to examine the eye, and to favour me with the following letter, giving a most satisfactory and philosophical explanation of all the phenomena:

fig. 1.) as a regular surface, resembling very much that of a hyperboloid; for the only indistinctness occasioned by a cornea of this form, would arise from the concentration of the rays before they fell upon the retina.

When I had the pleasure of examining the eye itself, the difficulty of explanation was in no respect diminished. In every aspect in which the cornea could be viewed, its section appeared to be a regular curve, increasing in curvature towards the vertex; a form which could produce no derangement in the refraction of the incident rays. As the disease was evidently seated in the cornea, which projected to an unnatural distance, it did not seem probable that there was any defect in the structure of the crystalline lens. I was therefore led to believe, that the broken and indistinct images which appeared to encircle luminous objects, arose from some eminences in the cornea, which could not be detected by a lateral view of the eye, but which might be rendered visible by the changes which they induced upon the image

of a luminous object that was made to traverse the surface of the cornea. I therefore held a candle at the distance of fifteen inches from the cornea, and keeping my eye in the direction of the reflected rays, I observed the variations in the size and form of the image of the candle. The reflected image regularly decreased when it passed over the most convex parts of the cornea; but when it came to the part nearest the nose, it alternately expanded and contracted, and suffered such derangements, as to indicate the presence of a number of spherical eminences and depressions, which sufficiently accounted for the broken and multiplied images of luminous objects."

Mr Phipps of London, who was consulted in this case, remarked to me, that he has had the opportunity of watching the progress of several cases where the cornea had become conical, and that he never found it in persons younger than fourteen or sixteen. "It is a change," he observed, "which sometimes takes place at the age of puberty;

and, when the cone is once complete, the disease seldom makes any further progress, or suffers any other change, than that the apex sometimes becomes opaque."

Beer * mentions, "That there is a kind of staphyloma worthy of remark, which I have seen in more than one case of hydrophthalmia." He adds, "The cornea, in such cases, is inconceivably distended, but it does not lose its transparency; when it is punctured, it is found to be very thin; it has happened also that sometimes it has burst. The patients, notwithstanding the transparency of the cornea, saw little or none at all.

"I had the last year an opportunity to observe a remarkable case of this kind in a woman, who, after an inflammation of the brain, was seized with a violent inflammation of the left eye, which was properly treated, and got well. Soon after, she was attacked with a very violent pain in the left half of the head, and afterwards with an uncommon weakness of sight. The eye was a

^{*} Pratische Beobachtungen über den Grauen Staar, &c.

little red afterwards, and then swelled; the pupil was very much dilated, and contracted but slowly. During this attack, the iris gradually changed its colour, and became at last quite red. The stinging pain soon became heavy and beating, and the patient lost the sight of this eye altogether, whilst the cornea continued to expand. I was at last forced to make an incision into the cornea, to prevent its bursting, but the aqueous humour was soon renewed.

"I was forced to repeat the operation; and the constant application of a cold vinous infusion of bark after it, prevented the eye from again filling. The organ afterwards remained shrunk; the iris retained its reddish colour; the pupil remained much dilated and immoveable, and she never recovered her sight *."

Richter + says he never saw this disease; Burgman + saw a very remarkable case, where

^{*} See Plate VII. fig. 2.

⁺ Richter's Anfangsgründe.

[#] Haller Disputationes Chirung, Tom. II.

the corneæ of both eyes of a person who was hanged, were so prodigiously extended, that they reached down to the mouth like two horns.

The corrugation or collapse of the cornea, arises either from a diminution in the quantity of the contents of the eye-ball, or from some disease of the cornea, or sclerotic coat. Violent inflammation of the eye, wounds and ulcers of the cornea, penetrating into the anterior chamber, are frequently followed by this disease.

After violent deep-seated ophthalmia, it is not unfrequent to observe the eye-ball altered in its form, and become smaller; and the change appears to arise chiefly from a diminution of the anterior chamber. Sometimes the change is only such that the cornea becomes nearly a plane surface, and comes into contact with the iris. In other cases it becomes puckered, and forms a furrow, sometimes at one part, at other times completely across the cornea.

The adjoining part of the sclerotic coat takes sometimes the same form, and is corrugated in a similar manner. It seldom occurs that there is such a total change in the appearance of the cornea, that the original division between it and the sclerotic coat cannot be distinguished. I have seen two cases where no such line of division could be observed. Reil* could see no remains of a cornea in a blind eye, which had grown smaller for eighteen years. "The eye-ball appeared to be formed of four parts; two deep furrows divided it, which probably arose from the action of the four straight muscles."

It also sometimes happens, that the cornea is lessened to half its natural size, or is altogether wanting, from an original malconformation †.

It very frequently occurs, that the eyeball does not recover its natural form after

^{*} Archiv für die Physiologie, von Joh. Christ. Reil.

⁺ Vide Handbuck der Pathologischen Anatomie, von Voigtel.

the cornea or sclerotic coat has been wounded; for although the wound heals soon after the accident, yet there is often so much inflammation excited, and such extensive injury done to the other parts of the eye-ball, that the humours are neither again collected in sufficient quantity, nor does the retina recover its tone.

Ulcers are still more apt to produce a permanent change in the form of the cornea than wounds. When they destroy its whole thickness, and allow the aqueous humour to escape, they heal so slowly that the relative situation of parts is altered: the iris comes in contact with the cornea, so that the aqueous humour oozes out until the ulcer is healed, the eye ever afterwards remaining of an unnatural shape.

Vision, in such cases, is always more or less impaired, and generally altogether destroyed.

CHAP. XIV.

OF THE EFFUSION OF BLOOD BETWEEN THE LAMELLÆ OF THE CORNEA, AND INTO THE ANTERIOR CHAMBER*.

The effusion of blood between the lamellæ of the cornea, or into the anterior chamber, is generally the consequence either of violent inflammation or of wounds.

When treating of inflammation of the cornea, I took notice that it sometimes happened that blood was effused between its lamellæ. When this takes place, a dark red-coloured circumscribed spot appears on some part of the cornea, which never alters its situation, but remains stationary until it is absorbed.

^{*} Hypoæma-Das Blutaug of the Germans.

When the internal parts of the eye are much inflamed, the aqueous humour often loses its natural transparency, and becomes tinged of a red colour from the mixture of blood. The red shade turns deeper as the quantity of effused blood increases, and in very violent ophthalmia, the blood is sometimes effused in such a quantity as to render the aqueous humour so opaque, that the iris and pupil cannot be distinguished. An wound of the iris, whether made accidentally or during the operation of the extraction or depression of the cataract, is very often followed by an effusion of blood into the anterior chamber; and it also happens that blood is effused, not only when an wound has been received, but even some days afterwards, probably in consequence of the subsequent inflammatino*.

I have seen several instances of an effusion of blood from an improper use of the

^{*} This disease has been called Cataracta secundaria Cru-

couching needle, and I believe it is not an unfrequent accident. The blood, however, is, in such cases, readily absorbed, and is followed by no bad consequence.

It sometimes happens, that blood is effused into the anterior chamber, from a violent blow on the eye. A very remarkable case of this kind was under the care of Mr Campbell, surgeon of the Inverness-shire militia: In consequence of a blow with the fist, a drop of blood collected in the anterior chamber, at the junction of the cornea and iris, but in a few days it was altogether absorbed.

There are also diseases of the iris, and of the internal parts of the eye, which are attended with hemorrhage, and which tinge the aqueous humour with blood. Voigtel * mentions the following case, where the quantity of blood was so great, as to distend the eye-ball, and burst it. "A man, fifty years of age, had by degrees lost the sight


^{*} Des Pathologischen Anatomie

of the left eye; -a year afterwards, a round white speck formed on the cornea, and in three months it changed to a blue colour; for other nine months it remained in the same state, and then became inflamed, but the inflammation soon went off. One day, after this, the patient felt his eye so much. distended, that it appeared to him as large as a hen's egg. This sudden swelling was accompanied with acute pain; and when slightly pressed, the pain became extremely violent, and darted through the whole head; at the same moment, blood flowed from the eye-lid, when the pain began to lessen and went off altogether in half an hour. The bleeding lasted two hours, and there were from five to six ounces of blood discharged the eye-ball was afterwards completely destroyed."

EXPLANATION

OF THE

Plates.






Inflammation of the Conjunctiva:



Inflammation of the Corner and Solerotic coats



Pustular Inflammation:



Thyme owner ad; not pino;

EXPLANATION

OF THE

PLATES.

PLATE I.

The three figures of this Plate are grouped together, in order to contrast more strongly the variety in the appearances of the peculiar inflammation which is represented in each.

Fig. 1.—Represents the conjunctiva covering the sclerotic coat, and lining the eye-lids, inflamed to a considerable degree, in the left eye of a young boy, eight days after the first attack. The inflammation increased for se-

veral days afterwards, so that this drawing may be said to represent the disease in its first or active stage. The upper eye-lid is elevated, in order to show a larger surface of the eye-ball. The colour, mode of distribution, and general appearance of the bloodvessels, were accurately copied from nature. They were of a deep scarlet hue; they did not seem to run to any particular point, but anastomosed freely with each other, and gave off branches in all directions. They appeared lying loose, and quite superficial, and could be readily elevated with forceps, or on the point of a needle. The conjunctiva was slightly swelled. The cornea remained quite transparent, and none of the red vessels passed over it. There was a light blue circle at the circumference of the cornea, resembling the arcus senilis of old age.

The palpebræ were considerably swelled, and had a dark purplish hue externally. Their internal surface had a villous appearance, and was very turgid with red vessels, which were so numerous as not to be distin-

guished in separate trunks. The *ciliæ* were glued together with a puriform fluid, and several globules of pus were seen on the under eye-lid, floating among the tears, and collected towards the angles of the eye.

Fig. 2.—This drawing represents the inflammation of the cornea and sclerotic coat, and the commencement of a corneal speck. The general expression given to this eye differs much from that of Fig. 1. and this arises chiefly from the appearance of the bloodvessels. The inflammation is confined, or is at least much more remarkable on one half of the eye-ball. There are numerous blood-vessels on the inferior part of the sclerotic coat; which pass over the transparent cornea and form on it a red cluster. They are of a deep scarlet colour, tinged with brown; they run in nearly straight lines on the sclerotic coat, and do not give off any branches till they approach near the cornea; each trunk is seen distinct, and they appear deeper, or more in the substance of the sclerotic coat, than the vessels in Fig. 1. There is a small circumscribed speck near the centre of the cornea, of a circular form,—a bloodvessel is seen running towards it; the speck is rather more opaque at the centre than at the circumference. The whole anterior chamber had that muddy, turbid appearance, which disappears instantaneously when the aqueous humour is evacuated: The eye-lids were turgid with vessels, but they were not much swelled, nor of the livid colour, nor covered with a puriform fluid, as those in Fig. 1.

This drawing was taken from a young woman who had had symptoms of inflammation in the eye for about six weeks, which came on suddenly, and without any known cause; —it was accompanied with violent pain in the temples, and in the eye-ball.

Fig. 3.—This drawing represents a pustule of the cornea, which has advanced to ulceration, and the peculiar inflammation which accompanies this disease. The blood-vessels are of a pale livid colour, and are more nu-

merous on the white of the eye adjoining the pustule, than at any other part. They also appear superficial when compared with those in Fig. 2.; and they anastomose freely with each other, each trunk being easily distinguished. The small white spot on the centre of the pustule, is the chalky looking matter so frequently covering ulcers of the cornea. There is also a diffused opacity round the cornea, immediately contiguous to the spot. The eye-lids are not affected, and there is no appearance of purulent matter among the ciliæ.—The drawing was taken from a young girl, six days after the commencement of the disease, who had been subject to frequent attacks of the disease, all of which were speedily removed by the application of the vinous tincture of opium.

PLATE II.

Fig. 1.—Represents the inflammation of a portion of the conjunctiva covering the cornea, as described in page 7th. The diseased portion appears of an opaque white colour, a little elevated above the natural surface of the cornea, and extending from the sclerotic coat to beyond the centre of the cornea. It has an oblong form, except near its extremity, where it becomes broader, and two large vascular trunks pass to the cornea, and are ramified into a number of minute branches when they reach the extremity of the opaque

Toflamation of the Conjunction of the Comes




Mer of the Come



Death of the Corner from Line



P. Syme adad from



spot. The whole cornea is a little clouded. The eye-lids are swelled and turgid with blood. A comparison of this Figure with Fig. 2. Plate I. will illustrate the difference in the appearances of the inflammation of the conjunctiva of the cornea with the common corneal speck.

This drawing was taken from a young lady about eighteen years of age, who had suffered from inflammation in her eye, attended with pain in the head and temple, intolerance of light, increased secretion of tears, &c. which had continued during three months. She formerly had four similar attacks, each lasting between three and four months. In this case the vessels were divided going to the corneal conjunctiva, by introducing a very sharp pointed curved needle behind them, then elevating them, and cutting away the elevated portion with a pair of scissars. This operation was followed by an increase of the inflammatory symptoms for a few days, but they gradually abated; the diseased portion of conjunctiva came away in the

form of a slough; and, after the lapse of a considerable time, the transparency of the cornea was nearly completely restored.

Fig. 2.—Represents a cornea which has two ulcers, having an opaque white appearance, resembling a piece of wet chalk, and, on a superficial examination, liable to be mistaken for a common speck. The edges of these ulcers are rounded and smooth, and elevated above the level of the surface of the ulcer. The rest of the cornea is quite transparent. On the nasal side of the white of the eye, there were a considerable number of blood-vessels, terminating in an ecchymosis. The eye-lids are a little swelled and turgid. This drawing was taken from a gentleman upwards of fifty years of age, who I saw along with Dr Monro, junior. The ulcers arose in consequence of a blow on the eye, which produced a violent degree of inflammation that lasted four months.

Fig. 3.—Shows the opacity of the cornea produced from the action of lime, as described in page 61.

The patient was a soldier who had suffered from ophthalmia during the Egyptian campaign, and who had since been subject to weak eyes. A quantity of lime accidentally got into one of them, which gave excruciating pain in the eye and head, produced a great degree of swelling and redness of the whole conjunctiva, and rendered about two-thirds of the cornea so obscure as entirely to destroy his sight. After the violent inflammatory symptoms were abated by bleeding at the temples, scarifying the eye-lids, &c. the process of separation of the dead portion of conjunctiva. commenced, and the drawing was taken when it had advanced a little way. The slough at the junction of the cornea and sclerotic coat began to separate, and vessels branched out, so that the division between these two coats, which at one period

could not be distinguished, now became visible. The separation went on from the circumference towards the centre, leaving the portion of the cornea underneath very vascular. When the white slough was touched with a pointed instrument, it appeared as if loose and moveable, and was very hard and brittle.

The whole cornea, except a small line at its upper part, is seen covered with the opaque matter, and there is also some remains of it on parts of the sclerotic coat. The conjunctiva on the sclerotic coat was a little tumefied, and instead of lying loose and moveable, it was hard and firm when cut with a scarificator. In some places it is considerably inflamed, and through the white portions small ramifications of red vessels may be observed.

Various applications were made use of during the treatment of this patient, but none had such a remarkable effect in abating the inflammatory symptoms and pain, and promoting the separation of the slough, as a solution of the nitrate of quicksilver. I had an opportunity of observing the progress of the case for five months, and, at the end of that period, the whole of the slough had separated, except some small ragged portions towards the centre of the cornea. There was a degree of obscurity apparently deep seated in the cornea, but his sight was so much recovered, that, with this eye, he could distinguish large objects.

PLATE III.


The three figures contained in this plate are intended to explain the most remarkable appearances of *pterygium*, varying in thickness from the thin membranous pellicle to the fleshy, cartilaginous excrescence.

Fig. 1.—Represents a thin membrane, covering about one-half of the cornea and sclerofic coat, which had nearly destroyed vision. It is interwoven with blood vessels, which appear lying superficial, and of a very considerable size; both the vessels and the membrane are of a dull crimson colour. The inferior tarsus of this eye is also diseased,


Membranen Lierygian



Common, trungular, Pterygium



· Theoling Perygium



P. Syme and wall prove

the edge of it is of an irregular form, and the natural figure of the cartilage is destroyed; almost all the ciliæ have dropped out, and in several places the edge of the tarsus is of a bright crimson tint. The gentleman, from whose eye this drawing was taken, had been, during a course of many years, repeatedly attacked with acute inflammation in the eye. He was under the care of Drs Monro and Wardrop; and it was proposed, that the portion of the conjunctiva, which was filled with vessels, should be removed close to the circumference of the cornea. In order to do this, Dr W. introduced a curved needle underneath them, elevated them, and then snipped off the elevated portion with scissars. The portion of the membrane, which remained on the cornea after this was done, appeared very loose, and it was readily dissected off with a scalpel. A few vessels, which were then seen in the substance of the cornea, increased in size for some days after the operation, but they gradually diminished, and in a short time his sight began to improve; and

after several months, though his other eye was completely lost, it was so much recovered, that he was able with it to read and write. Two years after the operation, his eye continued well.

Fig. 2.—Is a drawing of the most usual appearance of the pterygium, where the triangular form is well marked. Its base adheres to the semilunar membrane, and its apex extends a little way over the cornea. The vessels, in this case, instead of being of a dull crimson colour, as in Fig. 1. were of a pale scarlet. They are in considerable numbers, all small, and running in nearly straight lines towards the apex of the pterygium. The new formed body appears of considerable thickness, and lies loose, except at its base and apex. I had an opportunity of observing the progress of this case for upwards of eight years. When I first saw it, it had the appearance of a small globule of fat near the junction of the cornea and sclerotic coat, and it gradually became larger, so that its base adhered to the semilunar fold, and

its apex passed over the edge of the transparent cornea.

Fig. 3.—This drawing was taken from the eye of a young gentleman, who had the common triangular shaped pterygium from early life. Its growth having become rapid, a surgeon attempted to remove it by repeated scarifications; but these, instead of causing it to diminish, made it grow more rapidly. The mass was so large, as to separate the two tarsi, and involved the semilunar membrane and lacrymal caruncle. The surface of the pterygium was very irregular and rough, and some of the most prominent parts were white and hard; the rest of the tumour was of a bright red colour, bordering on vermilion; round the anterior extremity there was a ring of opaque cornea, more like the arcus senilis than common speck.

PLATE IV.

Fig. 1.—Was taken from the case described in page 31, where a fleshy excrescence, adhering to the conjunctiva of the cornea and sclerotic coat, has a tuft of hair growing from it.

Fig. 2.—Represents a collection of puriform fluid in the anterior chamber, in a man about forty years of age. There is also a considerable degree of cloudiness of the cornea, and its surface in some parts was eroded. The conjunctiva, covering the sclerotic coat, and lining the palpebræ, has become


Tuner of the Cogundica with hair guning from it




Paulent matter in the anterior chamber with opacify of the Cornea



Dep Weer with thicking of the Comes



Flympin'




very red, from violent inflammation, and some of the red vessels may be seen passing over the cornea.


When mining coals, a piece of the rock struck this patient's eye, and the drawing was made twenty-one days after the accident. It gave him great pain, and the pain daily increased, and became particularly severe in the forehead and temples, so as entirely to deprive him of sleep. The vision of this eye was destroyed, and he could only distinguish very large objects with the other eye, in consequence of an opaque speck of the cornea, which had formed early in life. I made an incision through the cornea, similar to that for the extraction of the cataract, and several drops of a puriform matter rushed out along with the aqueous humour. The effect of this discharge was almost instantaneous; for though the operation caused considerable smarting, and the fixing of the eye-ball produced a good deal of uneasiness, yet as soon as the contents of the anterior chamber were discharged, the

deep seated pain in the eye-ball, and the oppressive pain in the head, vanished. The pain never again returned, and the cornea daily regained so far its transparency, by scarification of the eye-lid, and the application of stimulating ointments, as to enable him again to be employed as a miner.


Fig. 3.—Represents an ulcer of the cornea, the appearances of which are very different from that of Fig. 3. Plate II. Instead of the superficial ulceration, and the surface being covered with an opaque white matter, there is a deep hollow cavity, of a conical form, the apex inclining towards the pupil. There is a degree of muddiness of the cornea, and it is very considerably swelled and thickened. This ulcer was the consequence of a tedious inflammation in a child's eye. By the continued use of the vinous tincture of opium for some weeks, the cavity of the ulcer was filled up, and the transparency of the cornea much restored.




Obscurity of the Comea accomp? with Enhopion.



Circumsorbed Speck with a blood versal passing into it .
and the under Eye lid inverted.



Part contoured Speck with adhesion of the Tris



P. Sym al Nat. pense!

PLATE V.

Fig. 1.—Shows the cornea in the first form of corneal speck, where it has become nebulous, or has a general cloudiness diffused over it. The degree of opacity is such, that the iris and pupil can be seen through it very indistinctly, and some parts are rather more opaque than others. The obscurity of the cornea was, in this instance, occasioned by that disease of the tarsi, called entropeon, in which the tarsi are turned in upon the eyeball. All the ciliæ have dropped out; the cartilages have lost their natural form, and their

edges become ragged. In this case, in order to restore the two tarsi into their natural situation, the ingenious operation of Mr Crampton of Dublin was employed, viz. dividing them at their junction, at the temporal angle of the eye, and after thus liberating the two eye-lids, keeping them in their natural situation, in the manner Mr Crampton points out *.

Fig. 2.—This drawing is intended to represent a corneal speck, of the second form. The speck is distinctly circumscribed, but not of an equal degree of opacity throughout. It is so situated, as to hide nearly the whole pupil. Towards its circumference, the iris may be observed shining through it. One small vascular trunk creeps along the sclerotic coat, passes over the edge of the cornea, and is lost on the speck. The patient had been seized twenty-two months before with a violent inflammation in his

^{*} Vide an Essay on the Entropeon, by Philip Crampton, M. D. London, 1805.


eye, after which the speck was formed, and, at the same time, the inferior eye-lid was inverted.

The tarsus is completely out of view, and is turned in upon the eye-ball; but the integuments of the eye-lids were so loose, that it could be easily put into its natural situation, and it remained there till, by a convulsive twitch of the eye-lids, it was again thrown inwards. The integuments of the upper eyelid are loose and puckered, from the constant winking, and corrugation of the eye-brow, which always accompanies this disorder. This and the former figure show two very different varieties of the entropeon, and explain also the very different modes of treatment necessary to be employed in each.

The tarsus was restored to its natural situation by removing a portion of the skin of the eye-lid, and keeping the divided edges of the wound in close contact, by three ligatures and adhesive plasters. When this was done, the inflammation speedily abated, and the speck diminished in opacity; thin layers

of it were also removed with a scalpel, so as to destroy the red vessel which passed into it; and by this treatment the speck became so small in a few weeks, that the man, from being quite blind, having lost completely the sight of the other eye, was enabled to walk about the streets, and distinguish objects with considerable accuracy.


Fig. 3.—Shows the appearance of a very thick pearl-coloured speck, of the third form, where, in consequence of an adhesion of the iris to the cornea, the pupil is drawn from the centre of the eye, and contracted to a very small point, so as nearly to destroy vision. The boy, from whom the drawing was taken, had lost also the sight of the other eye, from a violent inflammation after measles. It is one of those few cases where an artificial pupil may be attempted to be made, but, in this instance, the boy was too young to submit to the operation.



Undefined Speek with ad herior of the Tris.



Complete Stapley lone of the Comea



Partial Saphylonia of the Cornea



P. Lynn and, not pince.

PLATE VI.

Fig. 1.—Represents a corneal speck of the most opake, and, indeed, of the most incurable kind. It occupies the centre of the cornea, and is of a very irregular shape. It is of a pearl-white colour, and is nourished by several red-vessels. The cornea has lost its smooth spherical surface, from the cicatrix of an ulcer on the central part of the opacity. The speck is so large and opaque as to allow only a very small portion of the iris to be distinguished through it at its circumference. The cornea and iris also adhere to one another. The woman, from

whom the drawing was taken, had suffered repeatedly from violent attacks of inflammation in the eyes for three years.

Fig. 2.—Represents a complete staphyloma of the cornea in a man twenty-five years of age. It was produced by a violent inflammation, which followed a wound of the eye with a sharp-pointed instrument, two years before. It is of such a bulk as not to be entirely covered by the eye-lids. There still remains a line of division between the cornea and sclerotic coat. The cornea is formed into a tumour, nearly globular, and of an opaque white, and, in some parts, of a bluish, pearl colour. Over several parts of it are seen ramifications of red vessels. The sclerotic coat has lost its natural whiteness and lustre, and has a greenish hue. The palpebræ are inflamed, and many of the ciliæ have dropped out.

Fig. 3.—Is an example of what has been called the partial staphyloma; as only a portion

of the cornea is affected. The distinction is a useless one, both in a pathological and practical point of view. In this case, where the disease came on in consequence of the other eye having become staphylomatous from an wound, a portion of the cornea became obscure, and more prominent than natural; and although a part of the cornea immediately opposite the pupil remained transparent, yet its increased sphericity destroyed vision. The speck was of a pearl-colour, and was nourished by a blood-vessel: An adhesion had taken place between the cornea and the iris.

PLATE VII.


Fig. 1.—Is an outline taken from the eye of Miss——, as described in page 119, when viewed laterally, showing a cornea having assumed a conical form, whilst, at the same time, it remained transparent.

Fig. 2.—This is an outline of the case described by Beer, as quoted in page 122, in which the cornea is very much distended from hydrophthalmia.


Fig. 3.—This figure is a magnified view of a staphyloma, showing the mode of distribu-




Cornea distanded from Hydrophthalmia



Map of the Bloodrefiels of a Staphyloma





tion of its blood-vessels. The appearances of the two large trunks which advance to the extremity of the tumour were very remarkable, from the difference in their colour at different points. About one half of the trunk was of a pale livid hue; the contiguous portion became suddenly of a deep crimson, and the minute ramifications were of a reddish brown tint.

It is difficult to give a satisfactory explanation of this appearance, but to the physiologist the fact appears interesting. The difference in the quantity and in the chemical qualities of the blood in the trunks and extremities of the vessels, and the difference in the thickness of the coats of the vessels in the different parts, may all have a certain share in producing this appearance.

George Ramsay & Co. Printers.

tion of its blood-reseds. The repairmence of the two two large troubs which ever over 10 tips extinmity of the thousand water very premitted to the things of things of the things of the things of the things of the things of th

The addition of this appreciation of the collars of the collars of this appreciation of this appreciation of this appreciation of the collars of the collars

AND STREET, ST

ESSAYS

ON

THE MORBID ANATOMY

OF THE

HUMAN EYE.

By JAMES WARDROP, F. R. S. ED.

ILLUSTRATED BY PLATES.

VOL. II.

LONDON:

PRINTED FOR ARCHIBALD CONSTABLE AND COMPANY, EDINBURGH.

1818.

ESSLYS

THE MOUBID ANATOMY

HUMAN DEF

BETAMES WARDEGE P. H. S. E.

PARTE THE REAL CONTROL OF THE PARTY AND THE

II POW

MANUAL OF SECOND OF STREET

RIET

To JOHN WISHART, ESQUIRE.

MY DEAR SIR,

Allow me the pleasure of prefixing your name to a work, the subject of which has occupied so much of your attention.

It is peculiarly gratifying to me to have an opportunity of paying this public tribute of respect and esteem to your professional acquirements and your private virtues; and, at the same time, of expressing the pride I feel in reflecting, that a similarity of pursuits should have tended to mature the intercourse of our very early years into a friendship which has been uniform, and which, I fondly trust, will continue uninterrupted during the remainder of our lives.

I ever am,

MY DEAR SIR.

Your sincere Friend,

JAMES WARDROP.

Charles Street, St James's Square.

To JOHN WISHART, Esquere.

Mi Dean Small was well and the

Allow me the pleasure of prefixing your mone to drawk, the subject, of which has decupled so much of your attention.

Opportunity of paying this puedic tribute of respect
oned esteem to your projessional acquirements, and
your private virtues; used, at the some time, of exyour private virtues; used, at the some time, of extivity of private I feel in reflective, used a same
tiping of private time and know tender to materials
tiping of private truly circle and and a familiar
tiple time, will continue an acquire, and and a familiar
time, will continue an acquire, and and a familiar
time of our time.

problished a Freezont the Union age of the Relation of the Rel

Serial and American Street Street, and the last

St. Janes Street, Squared

ADVERTISEMENT.

The Publication of this Volume has been delayed from several causes which the Author could not have foreseen, and a detail of which can little interest the Public.

Several works have appeared since part of the present Volume was printed, so that the Author has unavoidably been prevented making use of the information they contain. There are some valuable additions in a new edition of Scarpa's Work; and Dr Beer of Vienna has published a Treatise on the Diseases of the Eye, consisting of two large octavo volumes, with some excellent engravings, which is certainly the most complete Work on this subject which has appeared in any country.

Charles Street, St James's Square, August 1818.

ADVERTISEMENT.

The Paldienion of this Volume has been delegablicon serviced causes which the Author could not have foreseen, and a detail of which can little interest the Public.

Several works have appeared since part of the present Science was printed, so that the Author has analysidably been prevented makeing used the ingused the ingustalists of Scarps's Work and Or Beer of Viennahas published a Treatise of the Disesses of the With some excellent engravings, which is certainly the most complete Work on this subject, which has appeared in any country.

Charles Street, St Jener's Square, Juguet 1818.

CONTENTS.

Page
CHAP. XV. General Remarks on the Aqueous Humour
and its Capsule,1
XVI. Of Inflammation of the Capsule of the Aque-
ous Humour, and its Consequences, 6
1. Of the Symptoms of Inflammation,ib.
2. Of the Effusion of Albumen,
3. Of Adhesions,15
XVII. Of Ossification of the Capsule of the Aque-
ous Humour,
XVIII. Of the Diseases of the Aqueous Humour,19
XIX. General Remarks on the Iris,23
XX. Of the Inflammation of the Iris,31
XXI. Of Alterations in the Form of the Pupil,37
XXII. Of the Permanent Pupillary Membrane,43
XXIII. Of the Undulatory Motion of the Iris,46
XXIV. Of Tumours of the Iris,49
XXV. Of the Prolapsus of the Iris,51
XXVI. Of Wounds and Lacerations of the Iris,53
XXVII. General Remarks on the Choroid Coat,58
XXVIII. Of the Inflammation of the Choroid Coat,62
XXIX. Of the Dropsy of the Choroid Coat,64
XXX. Of Ossification of the Choroid Coat

CONTENTS.

Page
CHAP. XLV. Of Amaurosis,165
1. Amaurosis from Diseases of the Brain, 167
2. Amaurosis from Poisons and Narco-
tics,176
3. Amaurosis from Wounds of the Fron-
tal Nerve, 179
4. Of Amaurosis from Disorders of the
Primæ Viæ,182
5. Symptoms of Amaurosis,185
XLVI. Of Night-Blindness,
XLVII. Of Coloured Vision,196
XLVIII. Of Squinting,202
1. Squint from Chylopoetic Derangement, 207
2. Squint from Worms,208
3. Squint from Organic Affections of the
Head,209
4. Squint from Teething,210
5. Squint from Imitation and Habit,211
6. Squint from unusual Position of the
Head,212
7. Squint from Disparity in the two Eyes, 213
8. Squint from Opacities, 214
XLIX. Of Double Vision, 216
L. Of Ocular Spectra,221
LI. Of the Disparity in the Vision of the two
Eyes,227
LII. Of the Involuntary Motion of the Eye-Ball, 232
LIII. General Remarks on the Sclerotic Coat, 236
LIV. Of Inflammation of the Sclerotic Coat,242
LV. Of Staphyloma of the Sclerotic Coat, 245

		Page
Explana	tion of the Plates	249
A Tormer's	Plate VIII	251
	Plate IX	254
	Plate X	257
Li Marido	Plate XI	259
	Plate XII	263
resiste gr	Plate XIII	267
	Plate XIV	270
	Plate XV	273

the auterior portion of the iris, passes through

ion of the tries and is united with the cap-

tior lamina: A complete bag is thus form-

conned bide CHAP. XV. is forming ali mi

GENERAL REMARKS ON THE AQUEOUS HUMOUR AND ITS CAPSULE.

pericardium, peritonaemas or vaginal coat,

When treating of the structure of the Cornea, it was observed, that its internal surface is lined by a polished, transparent, and firm membrane, forming part of that cavity which contains the aqueous humour.* This membrane, which has been called the capsule of the aqueous humour, lines not only the cornea, but the whole of the anterior chamber. After leaving the interior surface of the cornea, it extends over

branes cand ofce

the anterior portion of the iris, passes through the pupil, is reflected over the posterior portion of the iris, and is united with the capsule of the crystalline lens, forming its anterior lamina. A complete bag is thus formed, which in all respects is analogous, both in its natural structure and morbid changes, to *serous* cavities in other parts of the body.

The capsule of the aqueous humour, like the other serous membranes, as the pleura, pericardium, peritonæum, or vaginal coat, forms a circumscribed cavity, which is constantly kept moist by a fluid; and it is extremely probable, that the aqueous humour which it contains is exhaled from the surface of that membrane, in the same manner as the fluids are supplied in other serous cavities. Like serous membranes, too, the capsule of the aqueous humour is subject to particular morbid changes of structure. When inflamed, it acquires an additional thickness; albumen is effused on its surface; adhesions form between opposite portions of the membrane; and ulceration seldom if ever takes

place. It has also been found ossified; and the qualities of the fluid which it contains are subject to alterations from the natural state.

The aqueous humour itself performs several important functions. Like the fluids of serous cavities, it lubricates and allows the free motions of the contained parts, readily admitting of the contraction and dilatation of the iris, and yielding to any changes which may take place in the eye, to enable it to view objects at different distances.

The aqueous humour, besides, forms a refracting medium for the rays of light which enter into the eye; and perhaps it has also, along with the crystalline lens, and vitreous humour, an acromatic power.

The aqueous humour in the sound eye does not exceed five drops; but the quantity seems to vary in different eyes. In the fœtus, it is of a rosy colour, and during the first month after birth it remains reddish and turbid. Both its colour and consistence vary at different periods of life. In youth it is very thin, clear, and transparent; but it becomes

thicker, and less transparent, as we advance in years. In old age it is frequently whitish. * It is somewhat heavier than water, and possesses a small degree of viscidity. It is void of smell, unless it be kept for some time, and then it smells like the white of eggs become putrid. It is almost insipid, having merely a slight saltishness, which is only perceptible in old animals † According to the analysis of Chenevix it consists of water, holding small quantities of albumen, gelatine, muriate of soda, and phosphate of lime in solution.

The aqueous humour possesses a remarkable power of dissolving many substances accidentally deposited in it, and thus becomes a means of removing extraneous matter by which vision might be impaired. Instances have been observed, where the point of a knife, or other instrument which has been accidentally broken while performing operations on the eye, has been rapidly oxi-

^{*} See Porterfield on the Eye.

dated, and afterwards absorbed, when deposited in the anterior chamber. The solvent power of the aqueous humour is also strikingly illustrated in those cases, where portions of the lens or its capsule being detached, and left floating in that fluid, are gradually absorbed.

From the view which has now been given of the natural structure and functions of the capsule of the aqueous humour, its pathology becomes extremely simple, as its different morbid changes can be illustrated by comparing them with the changes of structure in other serous membranes; the symptoms in the diseases of this capsule being subject only to those modifications which arise from the peculiar functions of the eye as the organ of vision

Men considering inflammation of the corners of the corners of the chartens of the external or

CHAP. XVI.

OF INFLAMMATION OF THE CAPSULE OF THE AQUEOUS HUMOUR, AND ITS CONSE-QUENCES.

1.—Of the Symptoms of Inflammation.

of the natural structure

Inflammation and its consequences are the most remarkable morbid changes to be observed in the capsule of the aqueous humour. The former, as far as I know, has never been described by any author; and the latter, when noticed, have always been mentioned as distinct diseases.

When considering inflammation of the cornea,* it was observed, that the inflammation sometimes affected the external or

mucous covering of that organ; sometimes its proper substance; and in other cases it was confined to its internal membrane. Since these observations were published, I have been able, distinctly, to discriminate the inflammation of the capsule of the aqueous humour from affections of the adjacent structures; the inflammation, in such cases, either extending over the whole surface, or having been limited to a particular portion of that capsule.

When the capsule of the aqueous humour becomes inflamed, the disease is characterized, both by the changes which take place in the anterior chamber, and by the appearances of the inflamed vessels. The delicate membrane becomes opaque, and more or less of an albuminous deposition takes place on its interior surface, producing a muddiness or turbidity in the anterior chamber, and giving an appearance as if the eye-ball were unusually full and prominent.

Besides the diffused muddiness, there are

often one or more spots which distinctly denote an opacity of the substance of the cornea. These do not resemble any of the common forms of speck, but have a mottled appearance; and around the more opaque white central points of these specks, there is a kind of disk, very like what is to be perceived in some agates, and what are commonly called the eyes of pebbles—an appearance, no doubt, produced from the opaque capsule of the aqueous humour lying behind the opaque portion of cornea.

The effusion of albumen takes place, not only on that portion of the capsule of the aqueous humour which forms the interior surface of the cornea, but also in many instances on the surface of the iris, and in the pupillar opening; so that the pupil, instead of retaining its lustre and its circular form, becomes more or less dim, its form irregular, and its motions limited.

The appearance of the blood-vessels is different from that in the other species of ophthalmia. There is not that general redness of the white of the eye and internal palpebral membrane, accompanied with the puriform discharge, which characterizes the inflamed conjunctiva; nor is there that opacity of the cornea, and increase in the number of the blood-vessels, at a particular part, which take place, either in inflammation of the substance of the cornea, or of its mucous covering. A red circle of minute vessels appears on that part of the sclerotic coat, corresponding with the attachment of the iris, leaving the sclerotica between the cornea and these vessels nearly of its natural pale colour.

Besides the red circle of inflammation which has been described, and which is probably produced by an increase in the size and number of those vessels which form the connection between the iris and sclerotic coat, some vessels of the conjunctiva are frequently enlarged. These appear as insulated trunks, and can readily be made to glide from place to place, by moving the flaccid conjunctiva, whilst those of the sclerotica,

from being distributed throughout the substance of that tunic, appear deeply seated, and cannot be elevated with a pointed instrument, as may be done with those of the conjunctiva, or subjacent cellular membrane.*

The vessels are of a bright red colour during the active stage of the inflammation, and gradually assume a more crimson hue as the inflammatory symptoms subside. A difference, too, may sometimes be discriminated in the colour of the vessels of the affected sclerotica and conjunctiva.

The eye-lids participate little in this disease. There is sometimes an increased flow of tears, but the patient suffers very little pain on exposure to light, so that the eye-lids are kept open. The vision is more or less dim; and what ought particularly to be noticed, is a sensation of distention and fulness in the eye-ball, accompanied with a dull aching pain, generally in the forehead,

^{*} See Plate VIII. fig. 1.

sometimes also in the back part of the head:
—symptoms, it may here be remarked,
which are instantly and permanently relieved by evacuating the aqueous humour.

The constitutional symptoms accompanying the inflammation of the capsule of the aqueous humour, vary much in their degree of severity. Sometimes the pulse is very frequent and hard, the skin hot and dry, the tongue loaded, and the functions of the alimentary canal disordered. In other cases the disease, almost from its commencement, assumes a chronic form, and, after continuing a certain period, participates in any peculiarity of the patient's constitution.

An excellent illustration of the disease which has now been described in the human eye, may be observed in the horse, the eyes of that animal being subject to a peculiar inflammation, which seems to be entirely confined to the capsule of the aqueous humour. It attacks the organ very rapidly, and I have sometimes seen two-thirds of the anterior chamber filled with albumen

in less than twenty-four hours from its commencement. Horses are very subject to this kind of inflammation; and it affects every description of them, and at every age, though it is most frequent amongst those that are young and high bred, and those in high condition. Most commonly each eye is affected alternately, but sometimes both at the same time. An animal thus affected, is very subject to a relapse; and the returns of the disease are often far distant. The albumen is generally completely and rapidly absorbed, at least during the first attack; but ultimately the pupil remains covered with a portion of it, the edge of the iris adheres to this matter, and vision is thus destroyed. bedinsely need won and abiday

2.—Of the Effusion of Albumen.

THE effusion of albumen invariably accompanies the inflammation of all serous surfaces, and more or less of this substance

may remain after the inflammatory symptoms have subsided. It has already been mentioned* as being highly probable, that specks of the cornea are produced from an albuminous effusion into its cellular texture; and, when deposited in the anterior chamber, it is to be considered as one of the characteristic symptoms of the capsule of the aqueous humour having been previously inflamed.

During the continuance of the inflammatory symptoms, there is generally so much muddiness diffused over the whole anterior chamber, that no distinct portions of albumen can be distinguished, unless they be of a large size; but when this turbid state goes off, flakes of albumen will sometimes be perceived loosely attached to the surface of the iris, and, in other instances, the whole surface of the anterior chamber is covered with a thin layer of it. Albumen, when effused into the anterior chamber, assumes a varie-

^{*} See Vol. J. p. 94.

ty of appearances besides that of flakes. In some cases it floats in the aqueous humour, appearing like a thick smoke or cloud;* in others it is deposited in streaks, having a reticulated appearance;† and in others it resembles a purulent fluid.‡

The colour of the albumen is a straw yellow; and this, combined with the natural colour of the iris, produces a remarkable change in the appearance of the eye. If the iris be blue, the addition of the yellow albumen produces a green hue; § and when the natural colour of the iris is brown, the admixture of lymph gives it a lighter tint,—changes which are remarkably striking, when the diseased is compared with the sound eye.

If the portions of albumen which have been deposited during an attack of inflammation, be not afterwards absorbed, they become organized; and, in many cases, ves-

^{*} See Plate IX. fig. 1.

⁺ See Plate IX. fig. 3.

[†] See Plate IX. fig. 2.

[§] See Plate IX. fig. 2.

sels conveying red blood can be distinguished ramifying through them. This may be particularly observed, when a quantity of albumen is effused into the aperture of the pupil, as, in such cases, one or more red vessels can be seen coming off from the edge of the iris, to be distributed in its substance. *

3.—Of Adhesions.

When the capsule of the aqueous humour has been throughout inflamed, the albumen effused during the inflammation becomes the medium of permanent adhesions between different portions of the inflamed membrane. Thus, the edge of the pupil will adhere to the capsule of the crystalline lens; opposite points of the pupil will adhere to one another; and portions of albumen will sometimes form a connection between the iris and adjacent cornea. In this manner are pro-

^{*} See Plate VIII. fig. 3.

duced an infinite variety in the forms of the pupil; and as these changes take place during an attack of inflammation, the aperture of the pupil is not only irregular, but generally very much contracted.

But, besides the deviation from the natural shape, and the alteration commonly produced in the colour of the iris, from an admixture of yellow albumen, the aperture of the pupil seldom, if ever, remains transparent. In some cases, thin delicate webs may be distinguished passing across it; and these may be perceived varying in all the different degrees of obscurity, from that now described to a solid albuminous cake, in which bloodvessels may be traced ramifying through its substance.

A mere alteration in the form of the pupil has little effect in diminishing the powers of vision; but a very slight degree of obscurity, produced by the deposition of albumen, materially injures the sight.

CHAP. XVII.

OF OSSIFICATION OF THE CAPSULE OF THE AQUEOUS HUMOUR.

From the morbid changes which are known to take place in the pleura, in the peritonæum, in the vaginal coat of the testicles, in the pia mater, and in all other serous membranes, it might be expected that the capsule of the aqueous humour would occasionally be found ossified. One case has been already noticed,* where a thin shell of bone was formed within the anterior chamber; and I have no doubt but, in that instance, part of the cap-

^{*} See Vol. I. page 73.

sule which lines the cornea was ossified. A similar case has since come within my observation, thin laminæ of bone being several times discharged from the anterior chamber through ulcers formed in the cornea.

I have also had an opportunity of examining a case, under Mr Wishart's care, where that portion of the capsule of the aqueous humour which is reflected over the iris, was almost entirely converted into a bony shell.

It will afterwards be noticed, that, in several instances, the capsule of the lens has been found converted into bone, and in one case, the hyaloid membrane or capsule of the vitreous humour was found ossified. These facts make it the more probable, that, in the other cases, the osseous laminæ were formed in the capsule of the aqueous humour, all these membranes belonging to the serous class.

doubt but, in that instance, part of the cap-

CHAP. XVIII.

OF THE DISEASES OF THE AQUEOUS HUMOUR.

The aqueous humour is subject to few morbid changes, and as these are to be considered rather as symptoms of other affections than as distinct diseases, a few general observations only will be here necessary.

In some diseases of the eye, the quantity of the aqueous humour is increased, in others it is diminished. The former takes place in staphyloma, and in the disease which has been called *hydrophthalmia*; but I have never had an opportunity of seeing an instance

where there was a preternatural collection of aqueous humour, without its being accompanied with a disease in the coats of the eye. *

The quantity of the aqueous humour is diminished in old people, and in many of those diseases where the form of the anterior chamber is altered or destroyed.

It has already been remarked, that an albuminous fluid is sometimes mixed with the aqueous humour. Blood is also sometimes observed diffused through it; † and it is said that, in pregnant women, milk has been seen in the anterior chamber, but such a case has never come within my observation.

Woolhouse relates that he saw bladders of air, and it has even been asserted, that particles of quicksilver have been remarked in the aqueous humour. ‡

^{*} See "Staphyloma," Chap. XI. and "Alterations in the Form of the Cornea," Chap. XII. Vol. I.

⁺ Das Blutaug.

[†] Sybel in Reil's Archiv. für die Physiologie.

Prochaska mentions a case, where the aqueous humour was so acrid, that it tarnished the extracting knife.*

It may not be out of place to mention here, that there is a disease which frequently affects the eyes of horses in India, occasioned by a worm which, in size and colour, from the description I have received, may be compared to the common ascaris. It seems to be generated in the anterior chamber, and can be distinguished swimming about in the aqueous humour with great vigour. It produces a good deal of irritation and inflammation, the effects of which ultimately destroy the organ. The natives of India cure the disease by making an incision through the cornea, and extracting the worm through it. Though I have never had an opportunity of examining an eye affected with this singular complaint, yet the circumstantial accounts received from several accurate observers, leave no doubt in my mind of its actual

^{*} Voigtel's Handbuch, II. Band. p. 110.

occurrence; and the fact accords with what is known to take place regarding the formation of worms in many parts of the human body, and still more frequently in the inferior animals.

by affects the eyes of horses in India occa-

to be repented in the anterior chamber, and

devent turowed entropy to baryes not ad

district examining an eye affected, with this

antha at the horas and of Milan at 1768

CHAP. XIX.

GENERAL REMARKS ON THE IRIS.

The functions of the Iris are of considerable importance to vision. Besides regulating the quantity of light falling upon the retina by changes in the size of the aperture of the pupil, the contraction and dilatation of that opening enables the eye to perceive objects more distinctly at different distances. *

The iris, as has already been mentioned, is covered on both sides by the capsule of the aqueous humour, † and between these

^{*} Vide Wood's Optics.

two serous laminæ there are numerous bloodvessels, which form a net-work, and freely anastomose round the pupil. The iris is also plentifully supplied with nerves, the connections of which, as shall afterwards be mentioned, explain many of the phenomena in diseases of the eye.

Motions of the Iris.—Though no muscular fibres can be demonstrated in the iris of the human eye, yet, from the motions it performs, their existence can scarcely be doubted. This opinion is corroborated from some animals, and particularly some species of the parrot, possessing a voluntary power of contracting and dilating the pupil; and from muscular fibres having been observed in the iris of some of the larger animals.

Perhaps the motions of the iris have some analogy to those motions in plants which are produced by the influence of the sun's rays. Its motions in an eye where there is a complete cataract, or where the aperture of the pupil is drawn aside from a part of the cor-

nea* which has become opaque till it gets opposite to a transparent portion, makes this opinion probable. The iris, too, in many cases, retains its power of contraction and dilatation, even where the retina is completely insensible, so that it is likely that its motions do not altogether depend upon impressions made on the retina. In other cases of amaurosis, the motions of the iris are completely destroyed.

In some cases of amaurosis affecting one eye, the iris of that eye contracts and dilates, whilst corresponding changes are going on in the iris of the sound eye, from the influence of different degrees of light; but if the sound eye be excluded from light, the iris of the diseased eye remains immoveable. The motion, therefore, of which the blind eye is susceptible, must entirely depend on that sympathy which is known to subsist between the two eyes, and between corresponding structures in each of these organs.

^{*} See Vol. I. page 83.

Colour of the Iris.—The iris is of various colours, and is differently variegated in almost every individual. Some irides are of a light, and some of a dark blue shade, others are light brown, hazle brown, or of a very dark shade, variously streaked. In the Albino, the iris is of a rosy-red, and when the iris is red, the choroid coat is always of the same colour. The red colour is frequent in the lower animals, particularly in those who seek their prey during the night.

In an Albino from Surinam, the iris was observed of a fiery red colour, and variegated with blue and white streaks. The pupil also appeared very red.*

I have seen some people have a dark-brown angular spot on the iris of one or both eyes. A gentleman, who had a very distinct mark of this kind, said that there was a similar one on the irides of several branches of his family. †

^{*} See Mémoires de l'Academie de Sciences. Paris, 1734.

⁺ This variation in the colour of the iris produces what is called the "ring-eye" in the horse.

Sometimes the colour of the iris is different in the eyes of the same individual. Borelli mentions having known two people, each of whom had a blue and a black iris. Sybel saw a man, the iris of whose right eye was of a dark-brown, and that of the left a dark blue colour. This variety in the colour of the iris is sometimes accompanied with a corresponding variety in the colour of the ciliæ and eye-brows of the two eyes.

The difference in the original colour of the iris seems chiefly to depend on the colour and quantity of the pigmentum. Those whose irides are dark have it in greatest quantity, and in the Albino this colouring matter is entirely wanting, the red shade arising from the blood-vessels shining through the delicate and transparent membrane.

Of the Pupil.—In man, the natural figure of the pupil is circular, and is placed nearly in the centre of the iris, whereas in some animals it is of an oblong form, being so situated as to be most useful in enabling the

individual to extend his sphere of vision in those directions most necessary to his peculiar habits of life. Thus, in the graminivorous animals it is vertical, and in beasts of prey, as cats, it is a perpendicular chink.

The pupil also varies in its size in different people; and it may sometimes be observed different in the two eyes of the same individual, this probably arising from a difference in the convexity of the two eyes.

The pupil is generally larger in children than in adults. The pupil is also large in those who have black eyes, and are of a dark complexion; and, on the contrary, in those who are very fair and have light blue eyes, it is commonly a good deal smaller. The reason of this seems to be, that, when the eye-lashes are black, the eyes are better shaded from the light, and little light will be reflected from their inner surface upon the eye; therefore the pupil, which always becomes dilated when the light is faint, will keep wider than in those who, being of a fair complexion, have their eye-lashes white;

for white eye-lashes, by reflecting much light into the eye, must make the pupil contract.*

When the anterior chamber is small, the pupil is small also; and when it is of a full size the pupil is always larger. Hence the variety in the apparent bulk and brilliancy of the eye-ball.

In some individuals the pupil scarcely contracts at all in a bright light, whilst, in others, it is excited by the slightest glimmer. In people who live much in dark places, the pupils are more dilated than in those who are exposed to vivid lights, or employed in looking at minute objects. Men who live in countries covered with snow have the pupils constantly contracted, and see clearly; whilst those who inhabit clouded and moist climates have the pupils much dilated.

The effects of the extracts of belladonna and hyoscyamus, in producing a temporary dilatation of the pupil, are very curious;

^{*} See Porterfield on the Eye.

and it is remarkable, that this effect takes place not only when these substances have been applied to the conjunctive membrane, but to distant parts. Raius* mentions a case where a dilatation of the pupil took place each time the leaves of the belladonna were applied to a cancerous ulcer underneath the eye; and when an artificial dilatation of the pupil is wanted, it is easily effected by the application of the extract of one or other of these plants to the eye-brow.

Diseases.—Besides the diseases of the serous surface, the other structures which enter into the composition of the iris are subject to morbid changes; these, however, are not numerous. A variety of changes also take place in the motions and form of the pupil; but these are chiefly to be considered merely as symptoms of other diseases.

^{*} Historia Plantarum, p. 680.

CHAP. XX.

OF THE INFLAMMATION OF THE IRIS.*

Inflammation affects either the serous surface of the iris, or its proper substance, or it may affect both at the same time. †

From what has been already said of inflammation of the capsule of the aqueous humour, ‡ it is unnecessary to detail those symptoms and appearances which are produced by inflammation of that portion of the membrane which covers the iris. The thickening, the discoloration, and the quan-

^{*} Ophthalmia Uveæ, Iritis.

⁺ In this respect the iris may be compared to the lungs, inflammation of that organ being either confined to the pleura or to the parenchyma of the lungs, or affecting both these structures.

[#] See Chap. xvi.

tity of albumen effused on the surface of the iris, all take place, and vary according to the violence and duration of the inflammatory symptoms.

In those cases where inflammation seems to be confined to the iris, the proper substance of that organ will be found chiefly affected. But, although the origin of the complaint be confined to this part of the structure of the iris, yet when the symptoms are severe, its serous surfaces will also, in many cases, be observed to participate.

It has been noticed, when describing the symptoms of inflammation of the capsule of the aqueous humour, that the patient suffers little pain on exposure to light, the more remarkable symptoms being a pain in some part of the head, with a sense of fulness in the eye-ball; whereas, when the proper substance of the iris is inflamed, extreme pain on exposure to light is one of the most striking characters of the inflammation. It may here be remarked, that the apparent redness of the eye-ball forms no certain criterion of the degree of pain which a patient

suffers in any species of ophthalmia; for those parts of the organ, which, in their natural state, are most influenced by light, become peculiarly sensible to it when inflamed, whilst the exposure to light seems to produce but a slight impression upon the other parts, whose functions are not so immediately connected with vision, and which, from being most external, assume the reddest colour. Thus it happens that the iris, whose province, from the peculiar sensibility with which it is endowed, is to regulate the quantity of light falling upon the retina, suffers from exposure to it in a more remarkable manner than any other part of the eye-ball, when affected with inflammation. Besides acute pain in the eye, inflammation of the iris is usually accompanied with a deepseated pain at the bottom of the orbit, which occasionally extends through different parts of the head, and is chiefly seated in the brow. Perhaps this degree of pain may, in a great measure, be produced by the inflammation extending to the choroid

coat; the vascular connections between that tunic and the iris being very numerous.

When the iris is inflamed, the pupil will be found very much contracted, even in a dull light; its edge becomes puckered, loses its acuteness, and appears turned back into the posterior chamber.

From the quantity of pigment which enters into its structure, the iris does not acquire that red colour which most other parts of the body do when they become inflamed, the colour produced by the inflammation being such as would arise from a mixture of red blood with the natural colour of the iris, whatever that may happen to be. In some cases, however, the red colour appears distinct, the blood-vessels being so numerous as to shine through the iris. Janin relates a case of violent inflammation of the iris, where it became of a blood-red hue, resembling a piece of raw flesh.* Beer saw it blood-red in a case of nephritis. Conradi

^{*} See Traité des Maladies de l'Oeil.

also once remarked the iris become of a blood-red colour, after a wound of the eye.*

When the serous membrane of the iris participates in the inflammation, besides the symptoms which have now been mentioned, more or less of an albuminous deposition takes place on its surface, producing discoloration of the iris, irregularity and opacity of the pupil, and a turbidity in the anterior chamber; appearances which have been particularly described, when treating of inflammation of the capsule of the aqueous humour. † The appearances of the blood-vessels in these two species of inflammation are very similar, the peculiar red circle which they form around the eye-ball, at a small distance from the junction of the cornea and sclerotic coat, and the deep situation of these vessels, being very striking in both cases,-a similarity which, no doubt, arises from the blood-vessels which supply the two structures, being both branches of the same vascular trunks.

^{*} Arneman's Magazin, B. I. S. 1. + See Chap. xvi.

Inflammation of the iris is excited by a variety of causes. It is particularly apt to be occasioned by exposure to bright and dazzling lights. It sometimes accompanies phrenitis; and, as far as I have been able to observe, the iris is more liable than any other part of the eye to be affected with venereal inflammation.*

When inflammation is excited in the iris of one eye, it sometimes happens that the iris of the other eye is in like manner affected. In one instance this was remarkable; the iris of one eye being inflamed in consequence of a punctured wound, soon after which the iris of the other eye became similarly affected.

This sympathy, which subsists not only between the two eyes, but between the similar parts of each of these organs, is illustrated by many other diseases, as well as those of the iris.

^{*} See Venereal Ophthalmia.

CHAP. XXI.

OF ALTERATIONS IN THE FORM OF THE PUPIL.

ui soldier ommit esw aloida, ogsde aslover

the mis that the ander-part of the iris wall

to different degrees of light. Heil "saw a man,

NEITHER the size of the pupil, nor the extent of the motions of the iris, are always equal in both eyes of the same person; and, in some instances, I have observed this difference to be very considerable, and usually accompanied with a disparity of the vision in the two eyes.

Malconformation.—It sometimes happens that there is an original malconformation in the figure of the human pupil. In a child of a few months old the pupils of both eyes were

observed to be of an irregular oval form, and they readily altered their size when exposed to different degrees of light. Reil * saw a man, in whose right eye the pupil extended downwards to a small sharp point. In the left eye the pupil was round, but the upper edge of it was so situated in the centre of the iris, that the under-part of the iris was scarcely to be observed. Kühn + saw a young woman born with a pupil of a rectangular shape, which was immoveable in the middle of the eye, not being lessened by a bright, nor enlarged by a dull light. Himly has observed cases where the pupillar opening was not circular, but indented to the depth of a line. In one eye he observed seven such indentations, so that the margin of the pupil formed seven small semicircles. The vision of this eye was sound, the motions of the iris natural, and there had been no injury of the eye. This peculiarity in the form of the iris he conceives to

^{*} See Archiv. für die Physiologie.

⁺ See Naturhistorische Bemerkungen.

arise from an irregularity in the disappearance of the pupillary membrane.

Malconformations in the pupil are sometimes hereditary. Block* knew a family in whom the father, with his children, nephews, and nieces, had long-shaped pupils. One of the daughters and one of the nephews had in one eye the pupil oval, and in the other it was circular. Conradi + mentions, that he knew a father, daughter, and granddaughter, in whom the under margin of the pupil was, as it were, cut out; the pupil consequently was not round, but oval, and run oblong to a point at the lower part. Hagström t saw a whole family, in which each member had such a large oval-shaped pupil, that the under edge of the iris had quite vanished. Acrell § saw a similar deformity, which was hereditary.

Sometimes there is more than one open-

^{*} See Medic. Bemerkungen.

⁺ See ibidem.

[†] Handbuch d. Path. Anat. §, 517.

[§] See Medic. Bemerkungen.

ing in the iris, from original malconformation. In the cabinet of Meckel there is a calf's eye preserved, in the iris of which there are two regular formed pupils close to one another.

Sometimes a double pupil is met with in one eye whose other parts are single; and sometimes both lens and pupil have been found double.*

Change from Disease.—The form of the pupil undergoes various changes from diseases of the iris, and also of other parts of the eye.

In some people the pupil is not quite circular, from a portion of the iris having lost its power of contraction and dilatation; at least I have observed several cases, in which one part of the edge of the pupil remained immoveable, whilst the others dilated and contracted, no adhesions or any alteration in the structure of the parts being perceptible. In some cases this want of motion seemed to be only at one point, and

^{*} See Voigtel's Handbuch.

in others it extended to a large portion of the circumference of the pupil. The same loss of power may sometimes be observed in a particular portion of the iris, where it has sustained some injury from the extracting knife.

When the iris has suffered from inflammation, the pupil is frequently found not exactly in the center of the iris, and it seldom regains its circular form and natural extent of motion, the edge of the iris forming adhesions sometimes to the capsule of the crystalline lens or to the cornea.

In a number of instances, by the shape and mobility of the pupil, an opinion may be formed of the sensibility of the retina. It sometimes happens that the pupil is dilated to such a degree, that the iris appears merely as a streak round the circumference of the cornea; but this is always the symptom of some other disease. A great dilatation of the pupil may generally be observed where there is a wound of the ciliary ligament; and I have always observed the dilata-

tion equally great in those cases where vision has been destroyed by a wound of the frontal nerve; an effect which may probably be accounted for from the connection which the frontal branch of the fifth pair of nerves has with the nerves of the iris.

activing the center of the link, and it seldom

recions cometimes to the carryle of the

raion of the pupil may generally be observed

CHAP. XXII.

month. It is very vascular, receiving its

partition between the two chambers, is

supply of blood from the iris.

When it is not absorbed before

OF THE PERMANENT PUPILLARY MEM-BRANE.*

All the natural openings have been found obstructed from original malconformations of structure. The meatus of the external ear, the nostrils, the vagina, the rectum, and the urethra, have been found obliterated; and the same thing has been met with, though very rarely, in the pupil.

The membrane which shuts up the pupillary opening in the fœtus, and forms a

Sinizesis Congenita.

partition between the two chambers, is seldom visible at birth, but may be seen in a still-born child of the sixth or seventh month. It is very vascular, receiving its supply of blood from the iris.

When it is not absorbed before birth, it appears in the form of an opaque web, which is easily distinguished from cataract by its being vascular, from the size of the pupil being unaltered, and from its being on the same plane with the iris.

In one case the pupillary membrane remained perfectly apparent until the sixth week, at the end of which period it spontaneously disappeared. Wrisberg * saw a boy four years of age, in whom this membrane was perfectly entire. Cheselden operated on a boy ten years old, who had this membrane imperforated. Borthwick † mentions an obliteration of the pupil in which a complete tunic of a hard nature separated the two chambers of the eye, and adhered around

^{*} See Haller, Nov. Comment. Gott. Tome II.

⁺ See Med. and Phys. Com. Edin. Vol. I.

the margin of the iris. M. De Wenzel also mentions a case of an adult, where the pupillary membrane of one eye remained unabsorbed.*

* See Manuel de l'Oculiste, p. 416.

resident apparently uninjured and the built

ed to have a very singular andulators pice

deep dies state, of the organ, vision dock

longer retains the power of contracting and

tion, being agirated to and the fixed a price

cloth exposed to a fluctuating wind,

CHAP. XXIII.

OF THE UNDULATORY MOTION OF THE IRIS.

In some cases where the operation for cataract has been performed, and where the iris remains apparently uninjured and the pupil of its natural form, the iris has been observed to have a very singular undulatory motion, being agitated to and fro like a piece of cloth exposed to a fluctuating wind.

From this state of the organ, vision does not seem to be injured, but the pupil no longer retains the power of contracting and dilating so freely as usual. This appearance has, in most cases, been observed soon after the operation, and has continued afterwards during life.

In people who have been born blind of amaurosis, or with cataracts, the same singular appearance may often be observed; and I have also remarked it in cases of blindness produced from injuries.

In some cases where the iris is subject to this undulatory motion, the whole eye-ball has also a peculiar goggling and involuntary action. In some cases of cataract, too, the opaque lens partakes of the tremulous motion. *

It is difficult to give a satisfactory explanation of this curious diseased motion of the iris, and it may not, in all cases, depend on the same cause. It is not improbable, that, in some instances, it may arise from an inert or paralytic state of the muscular fibres of the iris, the consequence of injury, so that the membrane hangs loose and unsupported, partaking of all those undulations of the aqueous humour

^{*} The " Cataracte Tremblante" of the French.

which take place when the form of the eyeball changes to look at objects at different distances. In cases of cataract where the lens is dislocated, and lies loose in the anterior chamber, the motions which the iris then receives, makes it probable that its undulatory motions may also be produced by the fluctuation of the aqueous humour.

has also a peculiar goggling and involuntary action. In some cases of cataract, too, the opaque lens partakes of the tremulous motion.

nation of this curious diseased motion of the

the same cause. It is not improbable, that, in

paralytic state of the muscular fibres of the iris, the consequence of injury, so that the mem-

brane hangs loose and unsupported particking

of all those undulations of the aqueous homour.

" The " Catacacte Traphien (c), of the French.

CHAP. XXIV.

OF TUMOURS OF THE IRIS

SMALL tumours, which have been called polypi, are sometimes, though rarely, found on the iris.

In the cabinet of Meckel, there is a preparation where the pupil is shut up by a polypous excrescence.

Beer mentions cases where "fleshy tumours" were found growing from the inner margin of the iris. *

I have seen only one example of a tumour growing from the iris, and, in this instance,

^{*} Lehre der Augenkrankheiten.

though of long standing, it had acquired but a small size. It seemed, however, extremely vascular, for, frequently without any external cause, it bled profusely, and would, in a short time, fill the anterior chamber with blood. Tumours of this description may be sometimes observed growing from the iris of the horse.

CHAP. XXV.

OF THE PROLAPSUS OF THE IRIS. *

A portion of the iris is occasionally prolapsed through wounds of the cornea.

But the most frequent cause of a prolapsus of the iris, is when an ulcer erodes the cornea, and penetrates into the anterior chamber. The aqueous humour then escapes, and, by the contents of the posterior chamber being pressed forwards, a portion of the iris is thrust through the ulcer of the cornea, forming a small projecting tumour.

When a portion of the cornea is destroy-

^{*} Der Vorfall der Regenbogenhaut-Hernia Iridis. Myocephalon-Tête de Mouche.

ed from ulceration, or gives way, from the over-distention which takes place during a violent attack of ophthalmia, the prolapse of the iris being, in this case, accompanied with a discharge of the aqueous humour, the inflammatory symptoms become greatly alleviated. But this accident is often succeeded by great pain; for the iris is not only kept preternaturally stretched by being dragged from its natural situation, but becomes strangulated, and violent inflammation ensues.

If the inflammatory symptoms be not alleviated by a proper treatment, and if the tumour be left untouched, a complete adhesion is formed between it and the contiguous cornea. The size of the pupil is more or less diminished, its form and situation altered, and, after a tedious process, the prolapsed portion of the iris is at last covered by a layer of albumen, which defends it from the action of external agents.

CHAP. XXVI.

the edges of the wound seldem adhere. and

OF WOUNDS AND LACERATIONS OF THE IRIS.

THE iris is occasionally wounded both from accidents, and during operations.

Simple wounds of the iris are not attended with pain, neither do they excite that degree of inflammation which has generally been supposed to follow wounds of that part of the eye. But when they are accompanied with more extensive injury, and when any portion of the iris is displaced, the consequences are often alarming.

Wounds of the iris are always followed by more or less inflammation, accompanied by an effusion of lymph; but, as that membrane is exposed to constant motions, from the dilatations and contractions of the pupil, the edges of the wound seldom adhere.

The iris is frequently torn from blows and injuries of the eye, and the laceration generally takes place at the union of the iris with the ciliary ligament. This sometimes happens in such a manner, that a second pupil is formed, whilst, in other instances, the disunion is much more extensive; and I have seen nearly the whole iris dragged to the centre of the eye, being almost entirely detached from the ciliary ligament. In one case, where a thorn penetrated the eye, after the subsequent inflammation was removed, the whole of the iris was found to have disappeared, except a very narrow stripe, which extended across the eye-ball. * The vision of this eye remained extremely indistinct, unless assisted with a convex glass, or by looking through a small hole made in a card; by either of which means the patient could read a small printed book. Richter saw in a man, who had fallen from his horse on his head, a complete want of the natural pupil; but there was an opening on the upper part of the iris, which was occasioned by a separation of that tunic from the ciliary ligament. Through this opening the patient could see objects clearly at ten and twelve paces distant; but when he approached near, he saw only the under half. * Beer saw in a young woman, who fell without injury from the second floor, the pupils in both eyes become immediately immoveable, and in one of them the iris separated from the ciliary ligament in such a manner as to form two pupils, through which the patient saw only the half of all objects. †

The easy manner by which the iris can be detached from its union with the ciliary ligament, makes this accident by no means

^{*} See Chirurg. Bib.

⁺ See Lehre der Augenkrankheiten.

unusual, and led Scarpa to think of detaching the iris from the ciliary ligament, in order to form an artificial pupil in those cases where such an operation was required.

After the iris has been disunited from the ciliary ligament, a permanent opening is formed between the two chambers, and this new pupil does not interfere with the functions of the old one, nor does it in any manner disturb vision, unless when the injury is very extensive.

In those cases which I have had an opportunity of examining, the form of the new opening was unchangeable, neither contracting nor dilating by alterations of light. M. De Wenzel, * however, mentions a case where the contrary was observed. "I have seen an instance," says he, "of a laceration of the iris which presented a very singular appearance. The laceration had taken place at the inferior and nasal part of the iris, and formed a second pupil, which was nearly

^{*} See Manuel de L'Oculiste.

oval. This pupil had existed more than twenty years, and had never affected the sight, and it had been formed without any pain or disease, and without any accident that could be recollected. A beginning opacity could easily be perceived in the crystalline lens of both eyes; but I was much surprised to remark in the right eye, where the disunion of the iris had taken place, a motion of contraction and dilatation in the new formed pupil. These motions were very apparent, and they were also observed by several other persons who were present."

CHAP. XXVII.

pain or disease, and without any accident

GENERAL REMARKS ON THE CHOROID COAT.

THE structure of the choroid coat in some respects resembles that of the iris, which has led to the opinion that they were continuations of one another.

Besides having on its interior surface a layer of black pigment, the choroid coat is composed of a thin delicate membrane, divisible into two laminæ. Its exterior surface is smooth, having no adhesion to the sclerotic coat, but from a few vessels, and, like other serous cavities, it is kept constantly moist by an exhaled fluid. The interior surface has no connection with the retina, a slimy fluid

being interposed between the black pigment and the nervous expansion. The choroid coat may therefore be considered as a double serous membrane, forming two distinct serous cavities; for it is not probable that the two cavities communicate in the sound eye, as air blown through a wound of the sclerotic coat passes readily every where between the sclerotic and choroid coats, but never enters into the anterior chamber.

The choroid coat is extremely vascular, and it is plentifully supplied with nerves, the principal branches of which pass as far as the ciliary ligament, and are distributed on the iris and ciliary processes.

There is no reason to suppose that the choroid coat has, like the iris, any muscular fibres, nor are the functions of these two membranes at all similar.

From the number and particular disposition of the vessels of the choroid coat, it probably performs some important office. It is generally supposed to be the organ for the secretion of the black pigment; and, in some cases of disease, where there was an inordinate secretion of the black pigment, I have found the choroid coat thickened.*

The choroid coat, however, is probably concerned also in some other function, for it is to be found in the Albino, and in those animals in whose eyes the black pigment is entirely awanting.

The diseased changes which have been met with in the choroid coat are not only extremely few, but they have been rarely met with. Like other serous membranes, it is subject to inflammation; a preternatural quantity of fluid has been found in both its cavities; and it is subject to ossification.

Klinkosch has found the choroid coat entirely awanting, from an original malconformation. When the eye-ball is affected with fungus hæmatodes, the choroid coat appears to be intimately connected with the

^{*} See Fungus Hæmatodes of the Eye.

diseased changes which take place in that formidable complaint; being generally found very much thickened and redder than natural, and in some parts completely blended with the diseased mass.*

In eyes that have long been disorganized, the choroid coat has been found extremely tender, and of a pale colour, with a very small quantity of black pigment. In one case, I found the natural polished exterior surface of the choroid coat destroyed at one spot, having acquired a granulated appearance.

Portal † says that he has found hydatids between the choroid coat and the retina.

^{*} See Fungus Hæmatodes of the Eye.

[†] See Anatomie Médicale par Portal.

CHAP. XXVIII.

OF THE INFLAMMATION OF THE CHOROID COAT.

The choroid coat is occasionally inflamed, the inflammation, in some instances, originating in that tunic, and in others, extending from some contiguous inflamed part.

Inflammation of the choroid coat is attended with great intolerance of light, and pain at the bottom of the orbit, darting through the head. There is little external redness, and the inflamed vessels which appear on the white of the eye are deep-seated, being those belonging to the sclerotic coat. The pupil is contracted, and the anterior chamber appears large, from the iris having a

preternatural inclination backwards. disease occurs most frequently in strong plethoric people, and is always attended with a considerable degree of symptomatic fever.

In diseased eyes, I have several times found, on dissection, the choroid coat turgid with red vessels. Larrey * found the interior surface of the choroid coat of a brownish colour, subsequent to an attack of inflammation, and this was probably produced from the effusion of albumen. In one case, the inflammation of the choroid coat was so violent, that suppuration took place, a puriform fluid being discharged through an opening in the sclerotic coat.

There are some cases of painful and defective vision, which are usually considered as diseases of the retina, but which are probably connected with affections of the vessels of the choroid coat.

^{*} See Mémoires de Chirurgie Militaire et Campagnes par Larrey, Tom. I. p. 220.

CHAP. XXIX.

thoric people, and is always attended with

OF THE DROPSY OF THE CHOROID COAT.

It has already been remarked, that the choroid coat may be considered as a double serous membrane, forming two distinct cavities, each of which is constantly kept moist by exhalation, in the same manner as similar cavities in other parts of the body. The exhaled fluid being sometimes collected in these cavities in a preternatural quantity, the disease may properly be denominated the dropsy of the choroid coat.

The fluid may be collected either in the cavity between the choroid coat and the retina, or between the choroid and sclerotic coats. From this it is probable that the func-

tions of the two laminæ of the choroid coat are independent of each other, and that the one may be diseased whilst the other remains unchanged; an observation which may be made in other organs where there are two different surfaces, as the intestines, lungs, and bladder.

I have had an opportunity of dissecting several eyes where a serous fluid had collected between the choroid coat and the retina. In these cases, the retina and the vitreous humour were displaced and compressed by the morbid collection of water; the vitreous humour being more or less absorbed, whilst the retina was shrivelled up, and formed a white bundle, extending through the centre of the eye-ball, from the entrance of the optic nerve to the posterior part of the crystalline capsule. The choroid coat was unchanged.*

Mr Ware mentions a case of this disease, where, on dissection, he found "a considerable quantity of a yellow-coloured fluid, as

E

^{*} See Plate XV.

thin as water, accumulated between the choroid coat and the retina; the retina itself being collapsed, and resembling a cone of a white colour, the apex of which was at the entrance of the optic nerve, and the basis surrounded the crystalline humour, the vitreous humour being absorbed." * Verle and Zinn describe similar cases. †

It will be difficult to distinguish this disease in the living body, by any symptoms or appearances of the eye, as these much resemble some other affections of the posterior chamber. ‡ When the accumulation of water has taken place slowly, the loss of vision has been gradual, and the accompanying pain and redness have not been severe. In other cases, the water is collected quickly, and is accompanied with great pain, particularly in the head; the pupil becomes much dilated, and when the disease

[·] See Surgical Observations on the Eye, Vol. I. p. 510.

⁺ See Zinn de Oculo, p. 25. Scarpa also mentions a case of this disease, but he describes it as a dropsy of the vitreous humour.

I Ware's Cases.

has far advanced, there is an appearance of an opaque body behind it, from the retina being compressed, which, in one instance, was mistaken for cataract, and an attempt made to couch it.

In the disease now described, the water has always been found collected between the choroid coat and retina; but, in a few instances, a watery fluid has been observed between the choroid and sclerotic coats; and this has led some authors to suppose that there was a communication between the cavity formed by these two tunics and the anterior chamber.*

^{*} See Zinn de Oculo Humano.

CHAP. XXX.

OF OSSIFICATION OF THE CHOROID COAT.

I have met with a few instances where a thin cup of bone was found between the sclerotic coat and the retina.* The ossifications in all these cases were exactly similar. At the bottom of the cup, there was a small round perforation, through which the retina passed to expand on the interior surface of the osseous shell. The retina was in immediate contact with the interior surface of the bone, but between the sclerotic coat and the ossification there was a very thin, tender, and pale-coloured membranous expansion, the only vestige of the choroid coat.

Similar cases are mentioned by Haller, Morgagni, Walter, Pellier, Morand, Scarpa, and Günz.

That the ossifications, in the cases which have been mentioned, were a diseased change in the choroid coat, is extremely probable, both from their situation, and from their resemblance to the choroid coat in form. Besides, ossification is a morbid change, to which all serous membranes are subject. It has been already noticed, that this change has been met with in the capsule of the aqueous humour;—and bony matter has also been found deposited in the capsules of the crystalline lens and vitreous humour.

CHAP. XXXI.

PROCESSES AND THEIR DISEASES.

The ciliary processes, like the choroid coat and the iris, are covered with a black pigment, and float in a serous cavity, into which they may be considered as forming prolongations. But neither the structure nor the functions of the ciliary processes are fully understood. It is well known, that they are extremely vascular, and that they are very plentifully supplied with nerves; but whether they are also muscular, or whether they are employed in regulating the position of the crystalline lens, to enable the eye to look at objects at different distances, physiologists

have not yet been able to determine. From their structure, however, it is extremely probable that their functions are important.

Diseases.—The diseases of the ciliary processes are little known. They probably participate in general inflammation of the internal parts of the eye; and they are involved in many of those diseases to which the adjacent structures are subject.

Wounds of the ciliary processes are said by Beer to be always followed by blindness, and an extraordinary dilatation of the pupil. It is a fact well established, that a wound of the frontal branch of the fifth pair of nerves is generally followed by complete blindness, with a great dilatation of the pupil; and from whatever cause the blindness may in such cases be produced, the connection between the ciliary nerves, and the frontal branch of the fifth pair, might lead us to expect that a wound of the ciliary nerves themselves would have a like influence on the retina. I have observed several cases of wounds which penetrated the interior portion of the sclerotic coat, followed by a complete amaurosis, accompanied with a pupil very much dilated. In such cases, it is probable that the ciliary processes were injured; affording an additional proof of the accuracy of Beer's observations on the danger of such wounds in operations of the eye.

by Beer to be always followed by blind-

and from whatever cause the blindness may

CHAP. XXXII.

GENERAL REMARKS ON THE CRYSTALLINE LENS AND ITS CAPSULE.

The crystalline lens is enveloped in its capsule in the same manner as the aqueous humour is contained in a serous cavity. The crystalline lens cannot, however, be altogether compared to the fluids which the other serous membranes exhale, though it is probable that the watery fluid, lying between the crystalline lens and its capsule,* moistens these surfaces in the same manner as the pericardium or surface of the pleura exhale their respective fluids. But the capsule of the crystalline lens cannot be considered as a simple serous cavity, for the pupillar por-

[·] Liquor Morgagni.

tion forms part of the cavity of the aqueous humour, and the neural portion is in a similar manner united with, and derives a covering from, the capsule of the vitreous humour.*

It is extremely probable, that the crystalline lens itself is organized, and that it is nourished in the same manner as other parts of the body. Albinus conceived that he had detected vessels running into it; and the changes which disease produces in its structure confirm this opinion. The lens, too, like the aqueous humour, forms part of an optical instrument; and its capsule serves the important purposes of its secretion, of its nourishment, and of retaining it in its proper position.

Like the capsule of the aqueous humour, the crystalline capsule is liable to inflammation, and to that thickening, loss of transparency, and those other changes which inflammation produces in all membranes of the

^{*} These terms, neural and pupillar, are used by Dr Barclay as synonymous with anterior and posterior.

serous class. It is also, like them, sometimes converted into bone; and the fluid which it secretes, if the lens can be so denominated, is subject to changes, some of which are very probably the effects of disease in the capsule.

The capsule of the lens may be considered as consisting of two portions, a pupillar portion, the exterior lamina of which forms part of the cavity of the aqueous humour; and a neural portion, which, in a similar manner, forms part of the capsule of the vitreous humour.

The pupillar and neural portions of the capsule will be found to undergo changes in their structure separately; the exterior lamella of the pupillar portion being frequently inflamed along with the other portions of the capsule of the aqueous humour, and the neural portion being chiefly affected in other diseases.

In children newly born, the crystalline lens is extremely soft throughout, and it grows firmer as they advance in years, the central portion having always the greatest degree of consistence. In old age, this firmness is sometimes equal in every part of the lens.

In the same proportion as the lens increases in density, it diminishes in size.

Zinn has remarked, that the figure of the lens varies at the different periods of life, being nearly spherical in the fœtal eye, and becoming gradually more flat till about the age of thirty, after which its form does not appear to alter. *

Sometimes the crystalline Iens has an unusual degree of convexity, from whence arises one kind of short-sightedness. In other instances, the lens has been found unusually flat, producing long-sightedness. † The crystalline lens is sometimes entirely awanting. It has also been found double; in which cases the pupil was also double. Its form, too, sometimes varies, having been observed of a triangular shape. ‡ Heister

[·] Zinn de Oculo Humano.

⁺ See Porterfield on the Eye.

[†] Voigtel's Handbuch.

found it divided into different portions; and Morgagni saw a lens, a portion of which was awanting.

As people advance in life, the lens generally loses its perfect transparency, acquiring a yellowish hue; and this sometimes increases to such a degree, that it acquires, in old age, the colour of amber.

But, besides this change of colour, which may be considered as that natural decay to which the lens, like all other parts of the body, is subject, it often becomes completely opaque, forming a cataract.

The lens is also subject to changes in its consistence, as well as in its colour, sometimes degenerating into a watery fluid, and sometimes being converted into bone. Its volume, too, is augmented or diminished, being sometimes found small and shrivelled, and in other instances large, and of an irregular form.

Disease also produces changes in the lens similar to the effects of maceration, exhibiting the radiated appearance of its laminæ, or dividing its surface into isoscular triangles.

The crystalline fluid, or liquor Morgagni, is sometimes preternaturally abundant; and its consistence and other qualities are also subject to changes from disease. Wenzel says, that he has found this humour resembling the meconium of the primæ viæ of infants.* More of this fluid has always been found in the eyes of animals long dead, than in those newly killed. Portal, on examining a number of eyes of animals immediately after death, could find none of this fluid, which led him to believe "that, like the fluids of other serous cavities, it exists nowhere in its natural state, but as a kind of vapour which condenses and augments from different causes." †

The crystalline lens is also subject to external injuries from sharp-pointed instruments occasionally puncturing its capsule, and blows on the eye dislodging it from its natural situation.

^{*} Manuel de l'Oculiste, Tom. I. p. 148.

⁺ Portal, Anatomie Médicale, Tom. IV. p. 440.

CHAP. XXXIII.

Another ciass of cataracts have been

minated secondary; but, as these are the

OF CATARACT OF THE CRYSTALLINE LENS.

The word cataract has been used as a general name for all those diseases of the eye where an opacity is formed behind the pupil; it is, however, employed here only to signify those diseases of the crystalline lens and capsule, in which these organs lose that perfect transparency which they naturally possess. Following the nosological system which I have hitherto adopted, of considering the diseases of each structure separately, the first class of cataracts are treated of with the diseases of the lens; and the second with those of the capsule.

Another class of cataracts have been denominated secondary; but, as these are the effects of operations on the eye, they will be, with more propriety, described in another place. *

When the crystalline lens becomes opaque, the opacity begins either at the central part of the lens, extending towards its circumference, or a general obscurity pervades the whole of the lens from the commencement of the disease. † It has been said, that sometimes the liquor Morgagni alone becomes opaque; but, in most cases, every part of the lens is diseased. These differences have led to the distinction of two species of cataract, the former having been called interstitial, and the other mixed cataract. ‡

The Colour of crystalline cataracts, as they appear in the eye, is very various.

^{*} See Volume III. Of the Treatment of Cataract.

[†] Glaucosis of Hippocrates—Gutta opaca of the Arabians—Suffusis—Der graue Staar, or Krystall Staar, of the Germans—Cataracta vera Crystallina.

¹ Cataracta Interstitialis et Cataracta Mixta.

The most usual colour of them is a bluish white or grey; * some are of an opaque chalky white; + sometimes they are clouded in different parts, having the appearance of a flake of snow; sometimes they have a greenish hue, and others are of a dull yellow, or amber colour; sometimes the opaque crystalline has a striated or radiated appearance, resembling that division of the sound lens into different segments, which is produced by chemical agents. ‡ These differences, which have been observed in the colour of cataracts, have also given rise to several of those trivial distinctions which have been considered sufficient to constitute distinct species of the disease. §

Cataracts are seldom of the same colour in the eye-ball, as after they are removed from it. Those lenses which appear white or grey in the eye, are generally dark yellow, or amber-coloured when extracted; and those

^{*} See Plate XII. Fig. 3.

⁺ See Plate XI. Fig. 1. and 2.

[‡] See Plate XI. Fig. 3.

[§] Cataracta virgata fibrosa, or striata. Cataracte Barrée.

VOL. II.

which have a yellow hue when in the eye, often appear of a dusky white after they are removed.

The consistence of the lens also varies very much in different cases of cataract. Sometimes it retains its natural texture; sometimes it acquires a caseous, * and in other cases a gelatinous consistence, and sometimes almost the whole lens is converted into a milky fluid.† In other instances, it becomes firmer and harder than natural, ‡ having been in some cases found converted into a chalky matter, and in others into bone. §

Like the natural lens, the central portion of the opaque one is usually firmer and harder than the part which is more external.

Sometimes the consistence of the crystalline cataract is not the same in both the eyes of the same person.

It is of much practical importance to be able to distinguish, in the living eye, these

^{*} Cataracta caseosa.

⁺ Milchstaar - Cataracta lactea or purulenta.

[‡] Cataracta crystallina solida.

[§] See Ossification of the Lens.

differences in the consistence of the diseased lens; for, according as the cataract is hard or soft, so ought the treatment to be varied. An accurate knowledge of these differences can only be acquired by experience, and habits of attentive discrimination.

It has generally been remarked, that the fluid or milky cataract is the most frequent in children, and that the solid or concrete cataract is most common in the adult. Soft cataracts, however, are by no means unusually met with in those advanced in life; and I have, in a few instances, known the lenses of young people converted into a chalky matter, and also into bone.

There are scarcely any diagnostic marks of a soft and hard cataract which can be altogether depended on. The soft cataract will, however, be in general distinguished, from the large size of the opaque body, its near approach towards the plane of the iris, or edge of the pupil, its white colour, and from having points, streaks, or inequalities, which vary their appearances at different times. The blindness, too, by which soft cataracts are accompanied, is always considerable; for, when the whole lens is opaque, the patient can derive no benefit from the great dilation of the pupil, which is produced by shading the eyes, or by the influence of belladonna, and can seldom see more than differences in the intensity of light.

When the cataract is hard, the opaque body is neither so large nor so close to the edge of the pupil as when it is soft, so that a sufficient number of rays of light can enter, and the patient is still capable of distinguishing some objects from the side of the eye. The cataract has the same shade of colour throughout the whole lens, and its natural smooth surface may be remarked; the motions of the pupil are extremely lively, and it seldom remains much enlarged.* The opacity behind the pupil is, at the commencement of the disease, observed in the middle, and then extends, very slowly, towards the circum-

^{*} Beer's Lehre der Augenkrankheiten.

ference of the lens. The colour of the hard cataract is usually grey, passing more or less to a bluish hue.

In cases of simple opacity of the lens, the pupil continues to contract and dilate, according as the quantity of light to which it is exposed is increased or diminished; and the retina continues to retain its sensibility, though its functions become necessarily limited. When the opacity is confined to the central part of the lens, the vision is little impaired, and in some people, who have naturally a large pupil, although a very considerable portion of the lens becomes opaque, yet the sight of the eye continues good. If the opacity of the lens is more general, objects appear to the patient, from the commencement of the disease, as if seen through a mist or smoke; and, in proportion as the opacity increases, vision becomes more obscure, till even the largest objects cannot be distinguished, and the patient can merely perceive a difference in the quantity of light, and in bright colours. An eye, affected with cataract, never becomes perfectly blind, for the lens does not acquire, even in the most advanced state of the disease, that degree of obscurity which prevents all rays of light from passing through it.

In most cases, the patient with cataract can distinguish objects better in a dull, than in a bright light. This arises from the pupil being more dilated in a moderate light, and thus allowing rays to pass to the retina through the edge of the lens, which still remains transparent. For the same reason, a person with cataract sees most distinctly when he shades his eyes, and exposes the object to a bright light; and it is on the same principle that the belladonna, by producing an unnatural dilatation of the pupil, gives a temporary relief, even in those cases of cataract where nearly the whole lens has become opaque.

When the exterior edge of the lens is less obscure than its centre, the patient then sees those objects more distinctly which are placed by his side, than those held opposite to him. If, however, the obscurity has not affected the middle of the lens, but some part of its edge, any circular body then appears to have its edge imperfect.

Near objects can be more distinctly seen than those which are more remote; and thus a person affected with cataract appears shortsighted.

The progress of cataract is very different in different instances of the disease. Sometimes it proceeds so slowly, as not to destroy vision for many years; at other times a complete obscurity of the lens takes place very rapidly. Several instances have come within my own observation, where a patient has, in a few hours, become quite blind with this disease. Richter and Eschenbach relate cases where people labouring under gout, which suddenly retroceded, were entirely deprived of their sight in one night, from the formation of cataracts.

The loss of vision from cataract is seldom uniformly progressive; for, sometimes after remaining a while stationary, it becomes suddenly more imperfect; and, by a succession of rapid alterations, is ultimately destroyed.

Cataract of the lens commonly affects, first, only one eye, and after the disease has advanced to a greater or less degree, the other eye is attacked. In other cases, both eyes are simultaneously affected. I have generally observed, that when cataract takes place in young people, and affects only one eye, the other eye continues, during life, free from disease. This is also the case in those cataracts which arise from an injury, particularly when no affection of the sound eye takes place soon afterwards. Richter mentions the case of a man who received a wound of the eye, which was followed by a cataract, and soon afterwards a cataract formed in the other eye. A similar sympathy has been observed with regard to the iris, * and it may be remarked in all similar structures of these organs.

When there is a cataract in each eye, the disease is not always of the same

^{*} See Chap. XX. page 36.

kind in both; sometimes the lens is affected in one eye, and in the other the capsule is opaque. Sometimes both the capsule and lens are opaque in one eye, whilst in the other there is only a crystalline cataract; and sometimes one eye is affected with cataract, whilst the vision of the other is destroyed by amaurosis.

Both sexes are equally subject to cataract, and no age is exempt from the disease; but it is more common in infants, and those advanced in years, than at the middle period of life.

In most cases, the crystalline cataract appears to be merely a *local* disease, though, in many instances, an opacity of the lens accompanies or succeeds other diseases of the eye. Cataract is frequently observed in gouty and rheumatic constitutions; and in such people, there is often reason to suspect that the opacity of the lens is more or less connected with the general constitutional affection.

In some cases, cataract appears to be

hereditary.* Many examples of hereditary cataract are recorded by different authors, and a considerable number have fallen under my own observation. I have several times known brothers and sisters, and, in one instance, twins, affected with cataract. I have also known a father, son, and grandfather, affected with it. Richter extracted a cataract from a man, whose father and grandfather were both blind from that complaint. + Maitre-Jean and Janin have both met with similar cases. Richter also saw three children born of the same parents, all of whom had cataracts at three years of age; and Janin saw a whole family of six persons blind from this

In some cases, where the cataract is hereditary, the opacity is confined to the lens; in others, both the lens and capsule are opaque. The hereditary disposition is observed both in the infant and in the adult,

^{*} Cataracta hereditaria.

^{*} Observations on the Cataract.

and to prevail equally in both sexes. The same kind of cataract generally prevails in the same family, where different branches are afflicted with the disease.

In some instances, cataract is congenital. It is by no means unusual to see infants born with cataracts; and these consist either of opacities of the crystalline lens, or of its capsule, * or both of these are at the same time affected.

In the congenite crystalline cataract, the lens is found converted either into a milky fluid or soft substance, into which the whole lens is changed; or into a firmer structure than natural, which alteration is generally confined to the central part of the lens, leaving the edges quite transparent. † In the latter case, the opaque portion is generally of a chalk-white colour, there is usually a distinct line of separation between the opaque and transparent portions of the lens, and the shape of the opacity is irre-

^{*} See Capsular Congenite Cataract.

⁺ See Plate XI. Fig. 1. and 2.

gular. In those cases where I have had an opportunity of examining lenses thus diseased, the opaque part was found to consist of a substance exactly resembling moistened chalk in colour and consistency. In other instances, where the disease affects the central part of the lens alone, the opaque portion is of a pale grey colour.

The congenite cataract commonly affects both eyes; but sometimes the disease affects the crystalline of one eye, whilst, in the other, the capsule of the lens is opaque; and sometimes both the capsule and crystalline are opaque in one eye, and in the other only one of these parts is affected.

The disease is never observed to make any progress after birth, neither does it ever suffer any amendment, unless in some cases, where, from an accidental blow or injury, the capsule of the lens is lacerated, and the lens undergoes a gradual process of absorption, leaving behind it the opaque capsule.*

^{*} See Scarpa, Traité de Maladies de l'Oeil.

In some cases, the congenital cataract is absorbed with great rapidity, even in a few minutes after it is mixed with the aqueous humour.

Vision varies in this, as in cataract of the lens, the patient being sometimes only able to distinguish different degrees of light, whilst others see with considerable distinctness.

In some cases of congenite cataract, the eye-ball has a peculiar involuntary rolling or goggling motion, and sometimes the cataract itself has, at the same time, a tremulous motion. *

Cataract of the lens is not unfrequently combined with other diseases. In most cases, where amaurosis accompanies this disease, there is a preternatural dilatation and immobility of the pupil, and the opaque lens appears behind it of a large size, and generally of an equal shade of opacity. † In some cases, where these two diseases are combined, the pupil remains of its natural size, and con-

^{*} Cataracte Branlante of the French.

⁺ See Plate XII. Fig. 2.

tinues to be influenced by light; but the presence of the disease in the nerve is always pointed out by the loss of vision being antecedent to the opacity of the lens, by the blindness being to a greater degree than would take place in simple cataract, and by sparks of fire, and spots appearing before the eyes, pains of the brow and head, with other symptoms of diseased retina.

Cataract of the lens is often accompanied with opacities, both of the cornea and of the crystalline capsule; and there are many derangements of the eye, where, along with other diseased changes, the lens becomes opaque.

The changes which take place in the structure of the crystalline lens, and the various appearances which it assumes in cataract, have not been well explained. The dimness and amber hue which have been mentioned * as common in those advanced in life, probably arise from a want of that balance in the secreting and absorbent systems, which

^{*} See Chap. XXXII. page 77.

is necessary to preserve the different parts of the body perfect and entire; and an unusual degree of that decay may, in some cases, produce a complete cataract. In those cases of cataract, which are the immediate result of a wound or injury of the lens, it is not improbable that the obscurity is produced in the same manner as a piece of glass or transparent crystal becomes opaque when it is struck or bruised; or a cataract of this description may be imitated by penetrating the lens with a needle, after it has been removed from the eye. Whatever, too, is capable of causing a coagulation of the albumen or gelatine, of which the sound lens is composed, will destroy its transparency, a change which will probably be produced whenever, from any cause, its organization is destroyed, or when it comes in contact with the aqueous humour se sew bas sliniga ni omis omos rot

for dissection by the late Mr Allan Burns of Glasgow; a man no less remarkable for his eagerness and liberality in diffusing know-

See Ossification of the Capanie of the Lens.

CHAP. XXXIV.

OF OSSIFICATION OF THE CRYSTALLINE LENS.

It is not unusual to hear of cases where the crystalline lens has been supposed to have been converted into bone; but in all those which I have had an opportunity of examining, the lens was found inclosed in an osseous shell, which, in all probability, was the capsule of the lens ossified, excepting in one instance.*

In this case, the eye had been preserved for some time in spirits, and was sent to me for dissection by the late Mr Allan Burns of Glasgow, a man no less remarkable for his eagerness and liberality in diffusing know-

^{*} See Ossification of the Capsule of the Lens.

ledge, than for his industrious professional zeal. A greyish spot was observed on ic.noving the sclerotic from the choroid coat. On dissecting back the choroid coat, the pupillar portion of the posterior chamber was filled with a white pulpy mass; and, on dividing the crystalline lens, its central portion was found converted into a hard bone.* The external laminæ of the lens were soft, but those nearer the centre became more consolidated, the central portion itself being of a deep brown colour, perfectly osseous, and exhibiting a laminated structure.

It has already been noticed, it that the lens has been found converted into a substance resembling moistened chalk in colour and consistence.

^{*} See Plate XI. Fig. 5.

⁺ See page 83, Chap. XXXIII. and Plate XI. Fig. 1. and 2.

CHAP. XXXV.

OF WOUNDS AND DISPLACEMENTS OF THE LENS.

THE lens sometimes becomes obscure, both from blows on the eye-ball, and from sharp-pointed instruments penetrating it.

When a blow produces an obscurity of the lens, it is probable that the change will be found to arise from the capsule having burst, and the lens being exposed to the action of the aqueous humour.

A diminution of the transparency of the lens may also be produced by the mechanical changes from the puncture of a hard body on a substance possessing the tenacity and transparency of the lens.

When the lens is wounded by an instrument penetrating the anterior chamber, generally more or less of the opaque lens escapes, by degrees, through the wound of the capsule, forming an opaque cloud, which passes through the pupil, and extends into the anterior chamber. * In some accidents of this kind, I have seen the whole opaque crystalline pass, by degrees, through the wound of the capsule, into the anterior chamber, and be gradually and completely absorbed. This process has been imitated with great success, in order to produce the absorption of some kinds of cataracts.

In some cases, cataracts arising from an injury of the lens, remain unaltered during life; but, I have generally observed, that in those instances the wound has been made in such a manner, as not to form a communication between the anterior chamber and cavity of the capsule of the lens.

It sometimes happens, that, in conse-

^{*} See Plate IX. Fig. 1.

quence of a blow on the eye, the crystalline lens is thrown out of its capsule, and passes into the anterior chamber, in which case, it always becomes rapidly opaque.*

In most cases where the lens is luxated from disease of the internal parts of the eyeball, it comes into the anterior chamber inclosed in its capsule. †

^{*} See Maladies de l'Oeil, par Maitre-Jean, p. 271.

⁺ See Cataract of the Capsule of the Lens.

CHAP. XXXVI.

OF INFLAMMATION OF THE CAPSULE OF THE LENS.

It has already been noticed, that the capsule of the lens is composed of two portions, which, in some respects, differ in their structure; the neural portion being much thinner, softer, and weaker than the pupillar.* From what I have been able to observe, it appears to me extremely probable, that, not only each portion, but that each of the laminæ of these two portions may be separately diseased; in the same manner as has been mentioned, when treating of the

^{*} See Chap. XXXII. page 74.

diseases of the choroid coat, a membrane having two distinct serous surfaces, it is observed that each texture is subject to diseased changes, unconnected and distinct from the others.*

Inflammation affects either the exterior lamina of the capsule of the lens, or the neural portion of the capsule, or it affects the whole interior lamina of the capsule at the same time.

It has already been observed, that, when inflammation has extended over the capsule of the aqueous humour, that portion of the capsule which forms the exterior lamina of the pupillar portion of the capsule of the crystalline lens also participates in the disease. † The affection of this portion of the capsule is distinguished by the effusion of albumen on its surface obliterating the pupil, and rendering it irregular, contracted, and unchangeable; and by adhesions taking place between the edges of the iris and the

^{*} See Chap. XXIX.

⁺ See Chap. XVI.

newly effused albumen. These changes may sometimes be observed when the iris and adjacent parts are little affected. At other times, the albumen is also effused over the whole surface of the iris, and into the anterior chamber. *

The adhesions formed between the iris and capsule of the lens are, in some instances, very partial, being easily destroyed by the application of belladonna to the eye; but, in others, they are firmer, and extend round the whole edge of the pupil, a partition being thus formed between the two parts of the anterior chamber. Hence the iris is pressed forwards towards the cornea, from the aqueous humour not having a free communication through the opening of the pupil.

When the capsule of the lens is extracted in this state, it will be found sometimes to have acquired a most extraordinary degree of thickness; but the thickness and opacity

^{*} See Chap. XVI. page 8.

are alone confined to the pupillar portion, the neural portion of the capsule retaining its natural pellucidity.* It is this state of the eye that has been described by Mr Wathen as a species of spurious cataract, and called by him the "Adventitious" cataract.†

Inflammation of the neural portion of the capsule, and of the proper capsule of the lens, can only be known from its effects in destroying the transparency and delicate structure of these parts.

When the capsule of the lens is wounded, the whole interior lamina usually becomes opaque; but the exterior lamina of the pupillar portion seldom becomes affected; and this is known from there being no appearances of albuminous deposition in the pupil or iris;—no adhesions of the iris to the capsule;—and the colour of the opaque capsule, instead of having the buffy hue produced by albumen, appears like an opaque white and thin floating membrane.

^{*} See Plate XI. Fig. 4.

⁺ See Wathen on Cataract, page 143.

Inflammation and opacity of the neural portion of the capsule are only to be distinguished, as far as I have been able to observe, in "secondary" cataract.*

* See Plate XIII. Fig. 5.

the capsule of the lens, it may readily be

CHAP. XXXVII.

OF CATARACT OF THE CAPSULE OF THE LENS. *

From the general observations which have been made on the structure and functions of the capsule of the lens, it may readily be perceived how it should frequently be the seat of opacities. Indeed there is, perhaps, no disease of that membrane which is not either accompanied or followed by a diminution in its transparency.

These opacities, of which there are several varieties, have all been, by most authors, denominated *capsular* cataracts.

Sometimes the whole capsule of the lens becomes opaque; and, in other in-

^{*} Cataracta capsularis vel membranacea. Der Kapselstaar.

stances, the opacity is confined to a portion of it. This is either the pupillar or neural portion; and the former of these, which is much thicker than the other in its natural state, sometimes exhibits a still greater inequality when diseased. That the two portions of this capsule should present different diseased appearances, may have been suspected from the difference in their connections, the pupillar portion being covered with a portion of the reflected capsule of the aqueous humour, and participating in its diseases, whilst the neural portion is, in a similar manner, connected with the capsule of the vitreous humour.*

The appearances of capsular cataracts are very various. Some of them are of a pearl white colour, and glistening; —others are of a dull milky whiteness; —others are mottled with small white spots upon a more opaque ground, —or they have a reticulated appear-

^{*} See Plate XI. Fig. 1.

⁺ See Plate XIII.

[†] Plate XIII. Fig. 2.

ance. Sometimes they have large white spots, and sometimes they appear like a delicate white web or flake of snow.* Their texture, too, is very various, being sometimes soft, and very pulpy; sometimes brittle, and easily ruptured; and sometimes very tough and elastic. In two or three instances, this membrane has been found in adults of considerable thickness, very much resembling cornea when long immersed in water. +

I have observed several examples of capsular cataract where the opacity was confined to the central portion of the capsule, an opaque white point appearing in the centre of the pupil, with several shades or degrees of opacity around it, forming circles, so that the diseased capsule resembled the section of a calculous concretion composed

^{*} See Practical Observations on the Formation of an Artificial Pupil in several deranged States of the Eye. To which are annexed, Remarks on the Extraction of Soft Cataracts, and those of the Membraneous kind, through a Puncture in the Cornea. By Benjamin Gibson, Surgeon to the Manchester Infirmary, &c. &c. London, 1811.

⁺ Ibid.

of concentric laminæ.* When a cataract of this kind is extracted, the pupillar portion of the capsule is found very much thickened, and the lens is sometimes quite transparent.

Though the neural portion of the capsule of the lens cannot be separated by dissection from the vitreous humour, the connection between these two capsules is sometimes completely destroyed by disease; and the lens inclosed in its capsule can be easily separated, i or is altogether detached, and allowed to float in the cavity of the aqueous humour, passing and repassing through the pupil, according to the position of the head. ‡ In almost all the cases of this kind which I have had an opportunity of observing, the capsule of the lens had become detached only after a considerable change had taken place in the structure of the parts contained in the posterior chamber; the vitreous hu-

^{*} See Plate XII. Fig. 4.

⁺ Cataracta cystica, —elastica. Balgstaar.

[†] Cataracta luxata.

mour being changed in its colour and consistence, and the functions of the retina destroyed.

In most cases, when the lens is found in the anterior chamber, it is extremely probable that it remains inclosed in its capsule, otherwise it would be absorbed by the action of the aqueous humour.

In some cases of cataract, where the lens, as well as the capsule, is opaque, and where the iris has an undulatory motion, the cataract is also moveable, the motion appearing to be communicated to it from adhesions between the capsule and iris. * In some cases this variety of cataract is accompanied with a change in the structure of the vitreous humour and its capsule, the humour losing its natural transparency and consistence, and the connection between the capsule and the ciliary processes being more or less destroyed. This disorganization of the eye is generally the consequence of violent inflam-

^{*} Cataracta tremulans.—Cataracte branlante—Zitterstaar.

mation, and is always attended with total blindness.

Most frequently the capsular cataract is congenital. Though it is generally accompanied by an opacity of the crystalline lens, yet in many cases the capsule is alone found in the eye, the lens having been absorbed. * This change in the congenital cataract is not unfrequent; for, in some cases, the capsule, without any evident cause, gives way, and allows the lens, which is usually in children degenerated into a milky fluid, to pass through the pupil, and mix with the aqueous humour, where it is absorbed. In other cases, this change is the effect of a blow on the eye. Mr Gibson gives an account of a child born blind, who received a blow on the eye, which was immediately followed "by a muddy white appearance, so as entirely to obscure the iris. On examining the other eye, I found a cataract, which, from its appearances, and being congenital, I concluded was of a milky kind. On inspecting the

^{*} Cataracta membranacea primitiva of Scarpa.

eye two or three days afterwards, a rent could be seen in the capsule, where the fluid had been discharged."* In several instances of congenital cataract, where, from the appearances of the eye, there was reason to suspect the lens was absorbed, the parents of the child have been able to recollect a period when a particular change took place in the cataract; appearances which were produced by the escape of part of the soft lens into the anterior chamber. In some cases, the whole lens is absorbed in this manner, except a very small portion.

When the lens undergoes this natural process of absorption, the pupillar portion of the capsule comes in contact with the neural portion, so that there appears only one membrane, more or less opaque, at some distance from the plane of the iris.

In one instance, I observed congenite cataracts in a child, the appearances of which were very remarkable. A substance of a regular *pyramidal* form, and of an opaque

^{*} Mr Gibson's work formerly quoted.

white colour, occupied about two thirds of the pupil of each eye. The base of the pyramid rested on the capsule of the crystalline lens, and its apex passed through the pupillar opening, advancing forwards till it apparently came in contact with the transparent cornea. The rest of the pupil was quite transparent, the iris had its natural motions, and the child appeared to see so well, that no operation, in order to remove the opaque bodies, was thought advisable. *

A case, in many respects similar to that now mentioned, is narrated by M. Wenzel. "The capsules of both eyes of a girl born blind were very opaque, and each formed a pyramid, which projected through the pupil, and extended as far as the cornea, but did not touch it." †

^{*} See Plate XII. fig. 6.

⁺ See Manuel de l'Oculiste, par M. Wenzel. Paris, 1808.

CHAP. XXXVIII.

OF OSSIFICATION OF THE CAPSULE OF THE LENS.

It has already been observed, that ossification is a diseased change to which all the serous membranes are subject. It is met with in the pia mater, in the pleura, in the peritoneum, and in the vaginal coat. I have found the capsule of the aqueous humour converted into bone; and I have also met with several examples of complete ossification of the capsule of the crystalline lens.

When the capsule of the crystalline lens becomes ossified, the ossification generally takes place in its pupillar portion, and in some cases the whole capsule is converted ally retains the natural form of the capsule, and contains the lens, the transparency of which is generally destroyed.* In some cases, where the capsule is ossified, the lens itself has been previously absorbed or diminished in bulk, so that the ossification is of a more irregular form; the capsule having been shrivelled previous to its conversion into bone. The ossified capsule is sometimes extremely thin and brittle; † in other cases, the ossification is thicker, and its texture firmer.‡

Ossification of the capsule of the lens is usually met with in people advanced in life. It is a change of structure which always takes place very slowly, and is usually accompanied with diseased changes in other parts of the eye. I have met with it in eyes where there was ossification of the

^{*} See Chap. XXXII.

⁺ See Plate XI. fig. 6.

[#] See Plate XIV. fig. 1. and 2.

choroid coat;—in eyes where the hyaloid membrane was converted into bone;—in some cases of fungus hæmatodes, and in staphyloma.*

Examples of ossification, where the diseased change seems to have taken place in the capsule, and not in the lens itself, as has been supposed, are mentioned by different authors. Morgagni met with a case where " there was nothing in the eye which could have been certainly known to be the crystalline or vitreous humour, but only a few drops of turbid and brown water. Not far, however, from the natural situation of the lens, I found a hard little body, in the size and form of its circumference, not much differing from the crystalline, but a little larger and convex on its anterior surface; and on its posterior it was concave, so that it represented a little shield." +

^{*} Ossification of the capsule of the lens may be frequently found in the diseased eyes of horses.

⁺ Epist. xiii. Art. 10.

Morand, * Janin, † and others, mention cases, in all of which the ossification appeared to have taken place in the capsule, and not in the lens itself.

Mr Gibson relates a case where the capsule of the lens was partially ossified. "It felt gritty to the couching needle, and produced a noise as if the instrument passed over a piece of dry parchment." ‡

aqueous his nature and erraculture lens, from

^{*} Mémoires de l'Acad. R. Anno 1730.

⁺ Traité de Maladies de l'Oeil.

[‡] See Practical Observations, &c. p. 121, by Benjamin Gibson, Manchester, 1812.

CHAP. XXXIX.

GENERAL REMARKS ON THE VITREOUS HUMOUR AND ITS CAPSULE.

THE capsule of the vitreous humour, or hyaloid membrane, is a serous cavity, and may justly be compared to the capsules of the aqueous humour and crystalline lens, from its structure, functions, and diseases.

The hyaloid membrane, in the adult, has the most perfect pellucidity, and is so extremely thin, that, in the natural state, it is scarcely visible. In the fœtus it is tinged of a rosy colour.

The anterior portion of the hyaloid membrane has been described as composed of two laminæ; * the one uniting with the pupillar portion of the capsule of the lens, and the other forming the neural portion of that capsule. After these two laminæ separate, they again unite, a cavity being thus formed, called the canal of Petit, which has no communication with any of the contiguous cavities, and the functions of which are not understood; nor has it ever been observed to have any share in diseases.

The interior surface of the hyaloid membrane does not altogether resemble the capsules of the other humours, for, instead of consisting of a simple cavity, numerous prolongations pass from it, which form cells communicating with one another, and in which is contained a fluid perfectly transparent. This cellular structure is probably intended for giving the vitreous humour a consistence, elasticity, and form; to keep the lens in its proper situation, and to permit of a certain degree of motion. It may also assist

^{*} See Chap. XXXII.

in preserving the shape of the eyeball, when its cavity is penetrated.

Like the aqueous humour, it is somewhat thicker, heavier, and more viscid than water, and, though it may appear to have more consistence, this deception arises from the globules which are contained in small cells communicating with each other, so that, when one is torn, the contents of those contiguous are slowly discharged.

The chemical qualities of the vitreous humour are nearly the same as those of the aqueous humour, and its refracting power is also similar.

The vitreous humour is penetrated by an artery arising from the central artery of the retina, and passes through its middle in a serpentine direction, giving off branches which communicate with one another; whilst the principal trunk terminates in the capsule of the lens.

Though the vitreous humour resembles, in many respects, the aqueous, it does not seem to possess the same solvent powers. This is strikingly illustrated by comparing the difference in the rapidity with which the crystalline and capsular cataracts are absorbed when in contact with these humours.

It is extremely probable that the fluid contained within the cells of the hyaloid membrane, is exhaled from the arteries in the same manner as the aqueous humour and fluid of the crystalline lens are exhalations from the vessels of their respective capsules.

The capsule of the vitreous humour is subject to morbid changes similar to those of other serous membranes, and the fluid itself is sometimes altered in its quantity, colour, and other qualities.

The vitreous humour does not lose its pellucidity like the crystalline lens, but retains, even in advanced age, its perfect transparency.

When a part of the vitreous humour is evacuated, it is replaced by an aqueous fluid, which seems to answer all the purposes of vision. This regeneration takes place even when the quantity discharged has been very

considerable; and, in experiments on animals, the whole has been taken out, and replaced by a watery fluid, the sight being perfectly restored.

CHAP. XL.

OF THE DISEASES OF THE VITREOUS HUMOUR
AND ITS CAPSULE.

The diseases of the vitreous humour and its capsule, like those of the aqueous humour, are few, and occur very seldom; and they generally accompany other diseases.

The vitreous humour is sometimes perceived to be much thinner in its consistence than natural. * When this takes place, the hyaloid membrane, with its various prolongations, are absorbed, so that the bulb of the eye resembles one bag filled with water; the

^{*} Synchysis, Dissolutio humoris vitrei, Atraphie de l'Oeil.

sclerotic coat yielding to the slightest touch, and the cornea losing its convexity. This sometimes happens in eyes which have suffered from severe internal inflammation, and it is always accompanied by other diseased changes.

A portion of the vitreous humour generally degenerates into a watery fluid after the use of the couching needle.

Morgagni dissected an eye "which was smaller than natural, and the pupil appeared white. Having cut into the back part of the sclerotica, a quantity of limpid water flowed out, into which water a great part of the vitreous humour seemed to have degenerated, the remaining part appearing natural."

The vitreous humour also becomes, in some instances, thicker than natural. Morgagni mentions having dissected the eyes of a man whose vision had been very imperfect for some time previous to his death, "in one of which the vitreous humour ad-

^{*} Epist. lviii. Art. 16.

hered to the finger, and formed itself into long threads, following the fingers as they were drawn asunder. This was not the case with the vitreous humour of the other eye, but in both eyes the lens was diseased."*

The vitreous humour is sometimes more or less diminished in quantity. An absorption of it sometimes takes place after the eye has suffered from violent attacks of inflammation, more particularly when the inflammation has affected the choroid coat and iris. In the dropsy of the choroid coat, † the vitreous humour is absorbed in proportion as the morbid accumulation of water takes place between the choroid coat and retina; and, in those cases where the retina has been so compressed as to form a cone, no remains of vitreous humour could be observed. ‡

In some instances there is an increase in the quantity of the vitreous humour. This

^{*} Epist. xiii. Art. 15.

⁺ See Dropsy of the Choroid Coat, Chap. XXIX.

[‡] See Maitre-Jan, p. 241.

happens not unfrequently in staphyloma; for, in this disease, the increase in the bulk of the eyeball will generally be found to arise more from an increase in the quantity of the vitreous, than of the aqueous humour. An increase in the quantity of the vitreous or aqueous humour has generally been treated of as a distinct disease, and denominated *Hydrophthalmia*; but I have never seen a dropsy of the eye without an accompanying disease of the sclerotic coat or cornea; and it seems to me that it is an example of staphyloma which Scarpa has described, in order to illustrate the pathology of hydrophthalmia. *

A change is sometimes to be observed in the colour of the vitreous humour. Morgagni † saw it converted into a few drops of a turbid brown water; and Scarpa found in its place a few drops of a glutinous, bloody water.

In the living eye, the colour of the poste-

^{*} Treatise on the Diseases of the Eye:

⁺ Letter xiii. 9.

rior chamber sometimes appears changed; but it is difficult to discriminate, in many instances, whether this change of colour arises from a change in the colour of the vitreous humour, or from some coloured body shining behind it. In the commencement of fungus hæmatodes, the buff-coloured substance formed at the bottom of the eye produces an appearance as if the colour of the vitreous humour was changed; and I have remarked a similar appearance from the effusion of albumen in the posterior chamber.

In some cases the vitreous humour acquires a dull greenish colour, accompanied with insensibility of the retina, a species of amaurosis which has generally been called Glaucoma.*

The capsule of the vitreous humour, like that of the aqueous humour and crystalline lens, sometimes becomes opaque. † Maitre-Jan saw it in the eye of a horse, like a beautiful net-work. I have observed the

^{*} See Amaurosis.

⁺ Cataracta Hyaloidea.

same appearance in a few instances, where the lens had been imprudently extracted, and where violent inflammation followed the operation.**

The capsule of the vitreous humour is sometimes found ossified. In one case, besides the capsule of the lens being ossified, I found several large, but thin, scales of bony matter, dispersed in an irregular manner throughout the vitreous humour, which, in all probability, were ossifications of the hyaloid membrane.

On examining an eye, the natural form of which was destroyed, the cornea having become quite opaque, and the anterior chamber almost entirely obliterated, I found, on dissecting back the sclerotic coat, the posterior chamber occupied by a hard irregular-shaped mass, with the choroid coat and iris adhering to it. On removing the choroid coat, no remains of the retina could be detected. The bony mass was composed of

^{*} Cataracta secundaria Arachnoidea.

⁺ See Plate XIV. fig. 2.

two distinct portions, slightly connected with one another.* One portion was in the situation of the crystalline lens, exactly resembled it in form, and, being composed of a thin hollow shell of bone, appeared to be the capsule of the lens ossified. † The other portion occupied the site of the vitreous humour; and, from the inequality of its surface, being composed of numerous osseous laminæ irregularly disposed, I conceived it to be portions of the hyaloid membrane ossified. A similar case is related by Morgagni, and another by Scarpa. ‡

tion of the external air; the cliactory derves,

^{*} See Plate XIV. fig. 1.

⁺ See Ossification of the Capsule of the Lens, Chap. XXXVIII.

[‡] Treatise on the Diseases of the Eye.

CHAP. XLI.

GENERAL REMARKS ON THE OPTIC NERVES
AND RETINA.

The structure of the different nerves is various, being, in this respect, unlike the arteries and veins, which throughout the system are the same, whatever be their size, their course, or functions. Thus, the nerves of the organs of sense have a structure calculated to receive impressions of external objects, according to the particular office which each organ is intended to perform. The auditory nerve is distributed in a fluid, which is agitated by every undulation of the external air; the olfactory nerves, and nerves of touch, are spread underneath

a delicate skin; the gustatory nerves terminate in numerous papillæ; whilst the retina is expanded around a transparent and refracting cavity, to receive the impressions of the rays of light.

The optic nerves are remarkable for being the only nerves, besides the olfactory, which arise from the cerebrum; and which, except the auditory and the olfactory nerves, pass to the organ they are intended to supply, without forming any communication with other nerves.

The optic nerves are larger, and the medullary matter is in much greater proportion in them than in any of the other nerves.

The optic nerves are also softer than any of the nerves within the cranium. Between their origin and place of union, they are as pulpy as the medullary substance from which they emanate, but their consistence is a little increased from their union to the optic foramina.*

^{*} Bichât, Anat. Des. Tom. III. p. 155.

Until the optic nerves unite, they derive no covering from the pia mater; but at the place of their union, canals are formed by this membrane, which are filled with the pulp, and which are prolonged to the retina. When they pass through the optic foramina, they receive an additional tunic; the dura mater dividing at this place into two laminæ, one of these uniting with the periosteum which lines the orbit, whilst the other forms a very dense and strong neurilema, which envelopes the nerve as far as the sclerotic coat. The pia mater, which forms the canals in which the pulp is contained, passes within the globe, forming the vascular lamina of the retina. Hence the sympathy observed in the diseases of these parts, from the connection of the coverings of the optic nerves. How often does pain in the head and in the different sinuses accompany affections of the eyes, and morbid feelings in the eyes, diseases of the membranes of the brain?

The optic nerves, besides being remarkable

for having their origin in the cerebrum,—
for containing a greater quantity of medullary matter than other nerves,—for having no
anastomoses,—from being enveloped in one
general covering, instead of being composed
of bundles of filaments,—are also singular in their structure, having an artery*
passing through their axis, in place of being supplied from the surrounding vascular
trunks. †

The optic nerves, at the place of their union in the sella turcica, were long believed to cross each other; that going to the left eye, being supposed to come from the right thalamus; and that from the left thalamus, to go to the right eye. Vesalius, Aquapendente, and others, had observed that the optic nerves sometimes remained separate from one another during their whole course; and the man in whom Vesalius saw this formation had strong vision. The same thing was

^{*} The arteria centralis retinæ.

⁺ Bichât, Anat. Gen. Tom. III. p. 154.

observed in animals;* and it was also found, that, in dissecting diseased eyes, the disease always extended along the side corresponding with the affected eye.

But, though the optic nerves do not seem to cross each other, there is little doubt that there is an intermixture of some of their medullary fibres; for it is much more probable that the remarkable sympathy between the two eyes arises from this intermixture of the medullary fibres at the union of the nerves in the sella turcica, than from any intermixture of other nerves in the brain. ‡

The retina has two surfaces, the structure of which are very different, the one which is contiguous to the choroid coat being vascular, and the other, which is medullary, being in contact with the capsule of the vitreous humour.

The first of these laminæ, of which the retina may be said to be composed, is a very

^{*} See Briggs's Ophthalmographia. Lugd. Batavor. 1686.

⁺ See Diseases of the Optic Nerve, Chap. XLIV.

[‡] Anatomie Medicale, par A. Portal.

delicate membrane, of a fine cellular structure; and it adheres slightly to the choroid coat. Injections shew that it is very vascular; and it is supplied with nerves from the ophthalmic ganglion, in like manner as the neurilema of the optic nerve is supplied with branches from the same ganglion.

The other lamina of the retina, which seems to be entirely composed of medullary matter, is so intermingled with the vascular membrane, that the two cannot be distinctly separated.

Besides the vascular and medullary lamina of the retina, there is a peculiar part of its structure called the Macula Lutea, discovered by Sömmerring, which has hitherto been only observed in the human eye, and the functions of which have not been even conjectured.* The macula lutea, which is observed near the centre of the retina, appears in the form of a spot of a yellow colour, contiguous to which is also a fold and a

^{*} See Icones Oculi Humani.

small hole. From the observations of Michaelis, it is extremely probable that this part of the retina has an important share in some diseases. In eyes which had become opaque, the spot could not be detected;—in a staphylomatous eye it was scarcely visible;—in an amaurotic eye this spot was black; *—and Beer has observed it changed in glaucoma.

In most diseases of the nerves, there is merely an alteration in their sensibility. Seldom can any change be detected in their structure. As the natural structure of the optic nerves and retina cannot be compared to that of any other organ, pathological researches can receive little advancement from that analogy which I have so frequently employed in investigating the morbid changes of other parts of the eye. The morbid alterations in the structure of the nervous system have hitherto been less successfully investigated than those of any of the

Mémoires de la Societé Medicale d'Emulation, anno 1798.

other textures which compose the human body. These researches are attended with peculiar difficulties, more especially in the retina; not only because the parts are extremely minute, and cannot be seen like other diseases in the living eye, but as very small deviations from the natural structure, which materially derange its functions, might escape an attentive observer. Important, therefore, as the diseases of the nerve of vision undoubtedly are, much remains to be done by future observers in this part of pathology; and it is only by the collection of accurate histories and dissections that we can expect to be able to refer the various combinations of symptoms which are observed in affections of this nerve to their respective morbid changes.

other textures which compose the inman body. These researches are attended with

peculiare difficulties, more especially in the

retina; not only because the parts are ex-

tremely minutes and cannot be seen ung

CHAP. XLII.

OF THE SYMPATHIES OF THE EYES.

The eyes, like the ears, the mammæ, the testes, and all organs which are in pairs, have a sympathy with each other, both in health and disease; and this is, perhaps, more remarkable between the eyes than any of the other organs, from the optic nerves being intimately interwoven. Diseases which were originally confined to one eye are transmitted to the second eye; and even when an eye sustains a slight injury, the other frequently becomes weak and irritable.

This sympathy between the eyes has not

escaped common observation; and there is a disease, frequent in the eye of the horse, having the appearance of a specific inflammation, which usually first affects one eye and then the other, almost always sooner or later destroying vision. It is known among some Farriers, that, if the eye first affected with this disease suppurate and sink in the orbit, the disease does not attack the other eye, or subsides if it had commenced in it. Thus they have adopted a practice of destroying altogether the diseased eye, in order to save the other; which is rudely done by putting lime between the eyelids, or thrusting a nail into the cavity of the eye-ball, so as to excite violent inflammation and suppuration. I have frequently succeeded in saving one eye of the horse by adopting this practice; but I destroyed the eye by simply making an incision in the cornea, and discharging through it the lens and vitreous humour. In some diseases of the human eye, where the disease makes a similar progress, first affecting one eye and then the other with complete blindness, the practice so successful in animals might, by judicious discrimination, be beneficially adopted.

But, besides this general consent between the two eyes, there is a striking sympathy in the corresponding textures of each organ. This is well exemplified in diseases. the lens be opaque in one eye, it is probable that, sooner or later, the lens of the other will become similarly affected. St Ives mentions the case of a man who received an injury of his eye, which produced a cataract, and some time after the lens of the sound eye became obscure; and what is curious, when the cataract of the wounded eye was extracted, the cataract in the other disappeared. The same thing is observed in the other textures. A young man received a wound in the eye, which produced violent inflammation in the iris; and in three weeks the iris of the other eye became inflamed. When treating of staphyloma, * a case is mentioned where a large staphyloma came

^{*} See Volume I. Chap. XII.

on after a punctured wound of the eyeball, and some time afterwards the other eye also became staphylomatous.

Thus it usually happens, that not only both eyes are apt to suffer when one is injured or diseased, but that, when both are diseased, either at the same time or at different periods, they are generally affected with similar diseases.

Besides this remarkable sympathy between the two eyes, depending, no doubt, on the intermixture of some of the fibres of the optic nerves at the place of their union, the eyes have a great sympathy with other parts, from the nervous connection which the retina has by the medium of the ophthalmic ganglion.

This connection explains many of the phenomena that take place in diseases of the eyes, which, in a pathological point of view, are extremely interesting, and a knowledge of which may lead to important improvements in the treatment of diseases.

The distribution of the first branch of the

how sudden exposure to a bright light, or looking at the sun, produces sneezing; how strong odours increase the flow of tears; how, during certain headachs affecting the scalp, the eyes become red and painful, the tears flow, light cannot be tolerated, and the eyelids have involuntary twitches; how contusions about the internal and upper part of the orbit so frequently inflame and irritate the eye; and how wounds of the frontal, infra-orbital, and other branches of nerves, which form anastomoses with the ophthalmic ganglion, are sometimes followed by amaurosis.

The communication between the sixth pair with the great sympathetic, explains the connection between the eyes and the abdominal viscera, and this connection makes it probable that these nerves send branches to the globe of the eye, an opinion some have denied. Petit found, that, when the par vagum was divided in animals, the eye of that side lost its lustre, and the pupil enlarged. Hence wounds and compression of the great sym-

pathetic nerve, whether in the neck, chest, or abdomen, occasion convulsions of the eyes, and even blindness; and children who have worms, have often dilated pupils, and other amaurotic symptoms. Besides, how often are diseases of the eyes sympathetic of affections of the chylopoetic viscera?

The nerves of the eye have also a great influence on those of the stomach. If an instrument be turned rapidly before the eyes, it sometimes produces vomiting, and injuries of the eye often have a similar effect.*

On introducing the couching-needle into a woman's eye, at the moment it pierced the coats she shrieked loudly, and was in a short time afterwards attacked with violent vomiting.

Inflammation rarely occurs in parts most

and the retina, being seldom inflam

incident to vision.

^{*} Portal, Anatomie Medicale.

CHAP. XLIII.

OF INFLAMMATION OF THE RETINA, AND ITS CONSEQUENCES.

1.—Of Inflammation of the Retina.

It has already been observed, that the retina, besides being composed of medullary pulp, has an exterior cellular lamina interwoven with numerous blood-vessels. Changes in the structure and functions of these vessels produce a variety of the morbid phenomena incident to vision.

Inflammation rarely occurs in parts most supplied with nerves; the brain, the tongue, and the retina, being seldom inflamed.*

When the retina is affected with inflammation, the disease is marked by painful vision; intolerance of light; sparks of fire, or drops of a red colour falling before the eyes; little external redness; pain darting through the head; with more or less constitutional derangement.

The disease is generally brought on by excessive use, or by much exposure of the eyes to bright and dazzling lights; or it accompanies or participates in inflammatory affections of the brain and its meninges.

In some instances, inflammation appears to be confined to the retina; whilst in others, the choroid coat or the iris seem to be also affected.

Inflammation of the retina usually attacks both eyes.

Besides this active inflammation, to which the retina is subject, its exquisite sensibility and the delicacy of its functions, render the slightest deviations from the healthy state perceptible. Hence it is, that some people cannot expose their eyes to the common light of day, without inconvenience; that sparks of fire appear before the eyes of some people; that strange images, which have been compared to flies, tadpoles, and such like, float before the eyes of others; and that vision becomes impaired in a variety of different ways. In all these affections of the retina, it is extremely probable that each arises from some particular change in the state of the vessels of the vascular lamina; but what the changes are which produce the different morbid symptoms, has not been at all satisfactorily ascertained, and must require the nicest pathological research.

In most cases of ophthalmia, in whatever texture of the eye inflammation commences, the retina generally participates; the degree being always indicated by the increase of the sensibility of the eye to light.

2.—Of the Morbid Sensibility of the Retina.*

In some people the eyes acquire a morbid sensibility to light, where there is no reason to suppose that the retina is inflamed.

This is very remarkable in those people who have long been confined in dark places, the eyes not being able to endure the ordinary light of day; whilst, at the same time, they can distinguish objects in an almost inconceivable obscurity.

Buffon relates a remarkable instance of an officer who was thrown into a dark dungeon, where the light never entered, and with no opening but a hole at the top, which was always kept closed, except when opened by the keeper to put down provisions. After remaining in it some weeks, the unfortunate man began to think he saw

^{*} Nyctalopia—Visus nocturnus—Cocitas diurna—Amblyopia meridiana—Vespertina acies—Dysopia luminis.—Photophobia.

some little glimmering of light. This internal dawn seemed to increase from time to time, so that he not only began to discover the parts of his bed and other large objects, but at length could perceive the mice that frequented his cell. After some months confinement, he was set free; but such was the effect of the darkness upon his eyes, that he could not, for some days, venture to leave his dungeon, and was obliged to accustom himself, by degrees, to endure the light of day.*

Instances have also been narrated, and with every probability of truth, of prisoners who had been for many years confined in dungeons, having been found altogether blind when they were exposed to the light of day. This calamity was said to have happened to some of those who were released from the Bastile, when that awful prison was destroyed.

This morbid sensibility to light is, in a

^{*} Goldsmith's Animated Nature, Vol. II. page 161.

lesser degree, exemplified by the improper use of shades and coverings to the eyes. It often happens that people wear a shade to relieve diseases of the eye, which are not attended with increased sensibility to light; but this finally takes place, and can always be relieved by gradually exposing the eye to the ordinary light of day.

In like manner, as vision becomes impaired by the exclusion of the eyes from light, so any inordinate exposure produces an increased sensibility in these organs. inhabitants of those Northern regions, which are constantly covered with snow, keep their eyelids nearly closed, and do not see distinctly during the day. To prevent their eyes from being injured, and to enable them to see more distinctly, these people wear an instrument consisting of a thin and light piece of wood, which covers both eyes, and in which there is a long, but narrow, horizontal slit or chink opposite to each eye. By this ingenious contrivance, the eyes are guarded from all lateral and dazzling light,

whilst the chink is sufficiently wide to allow of a pretty extensive range of vision on the surface of the earth.

Travellers on the hot and sandy plains of Africa find their vision equally injured, as those who live in countries covered with snow; and they are in the habit of wearing a piece of black crape before their eyes, to diminish the quantity of light, as well as to prevent the particles of sand falling into the eyes.

The inhabitants of some of the Eastern countries, where the sun shines with dazzling splendour, have a practice which enables them to see more clearly, whilst, at the same time, it beautifies the countenance, by giving brilliancy to the eyeball. A black pigment, composed of finely levigated oxide of antimony, mixed with oil, is spread over the edge of the eyelids and roots of the ciliæ, and renewed as occasion requires.

Various other causes render the eyes susceptible of distinguishing minute objects in obscure light. Boerhaave mentions the case of a man who could read during the night when he drank too freely.* Richter saw a man who had an inflammation in the eye, in consequence of a blow, who could read in a dark night. † A woman, who had suppressed menses, was blind during the day. The pupils became so contracted in a clear light, that they almost disappeared. As soon as the menstrual discharge returned, the disease subsided. ‡

It has been told of Tiberius Cæsar, that, if he awoke in the night-time, he could see all objects as clearly as if they had been illuminated with a bright light; but that, in a short time, the objects grew fainter and fainter, till at last they vanished. The same thing is also told of Alexander. Asclepiodorus is said to have read books in the night-time, without the assistance of any kind of light; and Hieronimus Cardanus assures us, that, when he was young, he

^{*} De Morbis Oculorum.

⁺ Wundartzneikunst, 3. Band. p. 478.

[†] Pellier.

needed not the assistance of a candle to read in the dark.* Similar cases are mentioned by Willis, Briggs, and other writers; the affection being denominated *Nyctalopia*, or night-vision, and treated of as a distinct disease.

* Porterfield on the Eye.

tabe awake in the night time, he

became your test in the resemble

The Empe dung is also told of Alexander

desideer us, that, when he was ,

CHAP. XLIV.

OF THE DISEASED CHANGES OF THE RETINA AND OPTIC NERVES.

1.—Of the Diseased Changes of the Retina.

The retina has seldom been found changed in its structure. The morbid changes are observed to take place either in the vascular membrane of the retina, or in its medullary lamina, or both these may be at the same time affected.

When inflammation has taken place, I have seen the retina assume a buff colour, which was probably produced by the effusion of albumen.

Morgagni found the retina quite indurated and white, adhering firmly to the capsule of the vitreous humour.*

The retina has been thickened, and changed in its structure, in eyes that have become disorganized. †

Beer has observed, in some cases of amaurosis, the vessels of the vascular membrane of the retina in a varicose state. This is a change which very probably takes place in those cases of amaurosis where there are symptoms of congestion in the head, and where the disease is relieved by depletion.

When the vitreous humour was evacuated from a lady's eye affected with amaurosis, a profuse hæmorrhage came on soon after the operation, which probably arose from a varicose state of the vessels of the retina, as well as of the choroid coat. In this instance, the vitreous humour had degenerated into an aqueous fluid; and varicose vessels were

^{*} Ep. xiii. Art. 9.

[†] Morgagni, Ep. lii. Art. 30.

observed on the sclerotic coat towards its posterior part.

It is not improbable, that, in many cases of impaired vision, where figures of various forms appear before the eyes, these symptoms arise from a change in the vessels of the retina.

The medullary portion of the retina has been found entirely awanting in persons who have long been affected with amaurosis.

It has already been observed, that Michaelis found the *macula lutea* in an amaurotic eye converted into a black spot.

Beer has made several dissections of that species of amaurosis which has been called *Glaucoma*, in all of which he found, that the obscurity in the vitreous humour commenced at the *macula lutea*, extending from it throughout the whole humour.

Walter found the retina altered in its appearance, and become tough.

Guerin found in a man who had been

blind ten years, "the retina thin, and scarcely observable, but very tough." *

In fungus hamatodes, the structure of the retina and optic nerve is changed in a remarkable manner, the whole cavity of the eyeball becoming filled with a substance resembling medullary matter, and the optic nerve changed in its form, colour, and structure. †

2.—Of the Diseased Changes of the Optic Nerves.

The optic nerves have frequently been found diseased, changes having been observed both in the neurilema and in the medullary portion of these nerves.

Cheselden and Kaltschmidt found the optic nerves very small in children who died of hydrocephalus. ‡

^{*} Malad. des Yeux.

⁺ See Fungus Hæmatodes of the Eye.

[‡] Voigtel's Handbuch.

The optic nerve is sometimes very much elongated, from tumours pressing the eyeball out of its natural situation; and, in several instances where this took place in a considerable degree, the functions of the retina continued unimpaired.

Morgagni found the "optic nerve of a perfectly sound eye discoloured, and flattened like a piece of tape." *

In a man who had never been blind, Morgagni found the optic nerve had lost its natural colour, and had become thin, from the orbit to its union. †

Paw found in the optic nerve a large hydatid, which had produced amaurosis. ‡

In Mr Heaviside's Museum, there is a preparation of the optic nerve of an amaurotic eye, where a *tumour* of considerable bulk has grown from the neurilema. §

Wandeler found, in a young man who had

+ Ep. xiii. Aut. 6.

6 Voigtel's Haudbuch.

^{*} Epist. lxvii. Art. 21.

[†] Epist. lxiii. Art. 8.

[†] Obser. Anat. Rarior. Ob. ii.

[§] See Plate XV. fig. 1.

amaurosis, "a hard swelling in the optic nerve."

Calculous concretions have sometimes been found in the optic nerve. Walter found in the left optic nerve of a maniac, just before it passes through the optic foramen, a calculous concretion, of a rounded and flattened shape, and of two lines in diameter. *

Morgagni found, on opening the head of a woman who had been blind, and who complained of excruciating pain in her head, a stone the bulk of a pea, "in the very substance of the optic nerve."

Morgagni found a large scrofulous tumour at the origin of the optic nerves. +

Ferro found the optic nerves surrounded and covered with albumen, in a case of amaurosis, which he considered as the effect of gout. ‡

Matter resembling chalk has been observed surrounding the optic nerve. §

^{*} Museum Anatom.

⁺ Ep. xiii. Art. 6.

[‡] Voigtel's Handbuch.

[§] Voigtel's Handbuch.

Bonnetus relates the case of a gentleman who had been blind from his childhood, and after whose death it was found, that "both optic nerves were not obstructed and contracted, but contorted." *

In a man who had been blind in the right eye, the optic nerve was found of a brownish colour, and thin for about one finger's breadth from the eye. It contained no pulp, but a fluid of a viscid consistence, and of a muddy grey colour. When this fluid was squeezed out, the neurilema remained in the form of a tube, the sides of which were thicker than natural. Beyond this portion, the pulp was firmer, but discoloured, and the nerve thinner as far as the union of the two nerves. †

Cesalpinus found, in dissecting the body of a man who had the sight of one eye very feeble, "one of the optic nerves attenuated, and the other of its natural form. The vision was weak in that eye which had the dis-

^{*} Sepulchretum Anatom.

⁺ Voigtel's Handbuch. Erster Band.

eased nerve, the patient having received a wound of the head on that part." *

A boy nine years of age fell on his head, which was followed by pain in the head, continued fever, and blindness. "The optic nerves were found, on dissection, very much wasted, and so fragile, that they dissolved by touching them with the hand." †

The optic nerves undergo great changes in fungus hæmatodes, and these changes are usually to be observed in the medullary portion of the nerve. ‡

Malacarne saw an instance where, besides the whole eye, the optic nerves, with the thalami and striated bodies, were awanting. In a child born with one eye, there was only one optic nerve. There were two at the origin, but in a short way they contracted, and formed into one trunk.

^{*} Sepulchretum Anatom.

⁺ Voigtel's Handbuch.

[†] See Fungus Hæmatodes of the Eye.

[§] Voigtel's Handbuch.

A case of Amaurosis is related by Morgagni, where "a remarkable bladder, full of the most limpid watery matter, occupied the optic nerves at the place of their crossing."*

3.—Of Changes in the Structure of the Optic Nerve, when the Eye has been destroyed.

Changes from the healthy structure take place in organs when their functions are not performed. If the motion of a joint be destroyed, the muscles diminish in size, and their fibres become more delicate and of a paler colour; and, if the circulation through an artery be interrupted, the canal of the vessel becomes obliterated. Thus, also, when the retina can no longer receive the impressions of external objects, in consequence of the eye-ball being destroyed, the Optic nerve loses its natural appearance. The bulk of the

^{*} Ep. xiii. Art. vi.

nerve is generally diminished. Instead of being round and firm, it becomes thin and flaccid. It loses, too, its natural opaque white colour, becoming of a pale yellow, and appearing more or less transparent.

These changes have usually been observed to extend only as far upwards as the union of the two nerves.

Mr Allan Burns dissected the head of a woman who had been blind from infancy, the cornea of both eyes being quite opaque. The Optic nerve on one side was small and pellucid from the eye-ball to where it joins its fellow. On the other side the nerve was quite healthy from the eye to the place of union, but beyond this it was small and disorganized. The preparation is now in my possession.

Bichât examined two cases where one of the eyes had sunk; the Optic nerves of the diseased eyes were sensibly shrunk as far as the place of union, but beyond this both nerves were of the same size.*

^{*} Anat. Descrip. Vol. III. p. 153.

Morgagni found the Optic nerve of an eye that had been disorganized, more slender than that of the sound eye, of a more compact structure when cut into, and of a brownish colour. "This change of structure and of colour extended to the place where the two nerves unite, but at this point, and above it, nothing but soundness could be seen on both sides." *

Vesalius dissected a woman where one eye was destroyed, and the other remained perfect. The Optic nerve of the diseased eye, from its origin in the brain to its insertion in the eye, appeared much shrunk on being compared with the nerve of the sound eye. †

Morgagni found, on dissecting the Optic nerve of an eye which was shrunk, and which had probably been long blind, the nerve "extenuated to a very great degree, and cineritious. For about a finger's breadth from the eye, it contained no medullary

^{*} Epist. xiii. Art. 29.

[†] Bonnetus, Sepulch. Anatom.

matter, but only a cineritious, turbid, glutinous humour. This being squeezed out by a slight compression, the cavity was left empty, so that the coat of the nerve seemed to be more like the coats of some canal. The coat was thicker than usual near to that place where the two nerves unite; here all difference disappeared, and both seemed perfectly sound at their union, and up to their origin.*

When an organ of sense is totally destroyed, the ideas which were received by that organ seem to perish along with it, as well as the power of perception.

willed and problem been

^{*} Epist. xiii. Art. 8.

⁺ Darwin's Zoonomia, Vol. I.

CHAP. XLV.

OF AMAUROSIS. *

The term Amaurosis has had a very extensive application in nosology, having been employed to denote all those diseases of the eye, in which the functions of the retina have become imperfect or destroyed; the eye appearing in other respects sound. Amaurosis, therefore, in its usual acceptation, signifies a Symptom of disease, as well as a distinct affection.

It has already been noticed, that Amaurosis accompanies almost all those changes of

^{*} Gutta Serena, Black Cataract, Palsy of the Retina.—Suffusio nigra.—Der Schwarze Staar.

structure which have been met with in the retina and optic nerves;* and amaurotic symptoms have also been observed accompanying some of the diseases of the choroid coat, † and internal parts of the eye-ball. The variety of the morbid changes of these parts is sufficient to account for the number of the symptoms of amaurosis which authors have enumerated; and so seldom has this disease become the subject of pathological research, that scarcely any attempt has ever been made to refer any particular assemblage, or combination of symptoms, to the concomitant diseased change.

Besides those cases of Amaurosis, where the paralytic state of the retina accompanies a change of structure in some of the parts within the orbit, amaurotic symptoms also arise from affections of the Brain and nerves, and from Sympathy with other diseased organs, more especially the alimentary canal.

^{*} See Chap. XLIV.

⁺ See Chap. XXIX. and Chap. XXX.

1.—Amaurosis from Diseases of the Brain.

Amaurosis accompanies many affections of the Brain.

In all those diseases where the quantity of blood in the encephalon is increased, or the actions of the arteries changed, the functions of the retina are more or less affected, and, in some cases, a complete Amaurosis is produced.

From the course of the ophthalmic arteries,
—the situation of the circulus arteriosus,—and
the peculiar distribution of these arteries in
the different tribes of animals, there can be
no doubt that an equal and uniform current
of blood in the internal parts of the eye is
essential to vision. It is, therefore, easy to
conceive, how any disturbance in that circulation should interfere with the functions of
the retina.*

^{*} Ware's Works, Vol. I. p. 428.

Thus, those who are of a plethoric habit, when they hang down the head, or by any means increase the circulation of the blood, frequently excite the appearance of Spots before their eyes, and sometimes complete blindness.

It was observed by Sauvages,* that the pulsations of the optic artery might be perceived by looking attentively on a white wall, well illuminated. A kind of net-work, darker than the other parts of the wall, appears, and vanishes alternately with every pulsation. This change of colour of the wall he ascribes to the compression of the retina, by the diastole of the artery. The various colours produced in the eye by the pressure of the finger, or by a stroke on it, as mentioned by Sir Isaac Newton, seem likewise to originate from unequal pressure on various parts of the retina.

Richter mentions the case of a plethoric person, who, when he held his breath, and

^{*} Nosol. Meth. Class VIII. Ord. I.

looked at a white wall, perceived a kind of net-work, which alternately appeared and disappeared with the diastole and systole of the arteries; this phenomenon being probably occasioned by the plethoric state of the vessels of the retina. * In a staphylomatous eye, on the cornea of which was distributed two large red vessels, the patient could distinctly perceive a vermicular motion.

Mr George Young saw a cadet at Woolwich, who, from being obliged to wear a very tight neckcloth, and collar of the coat tightly clasped, had his sight much impaired; the sphere of vision was diminished, the pupils were dilated, and he had the appearance of flies constantly floating before his eyes.

Amaurosis is frequently brought on by keeping the eyes with great attention, and for a long time, fixed on minute objects, more particularly if the objects are not diversified. A frequent change in the objects which we look at has a material effect in refreshing the eye.

^{*} Richter's Anfangsgründe.

Tailors and milliners, who are employed at very minute needle-work, and who, when at night, have the objects brightly illuminated, frequently become Amaurotic.

Vision is particularly injured, when the objects are looked at with one eye. I have seen several cases of officers of the navy, who lost the sight of the eye with which they had been in the habit of looking through the telescope.

I have known several instances of soldiers and sailors who had become blind after having undergone great fatigue.

Richter relates a case of a man who became suddenly blind while carrying a heavy burden up stairs. Another man, who worked very hard for three successive days, became blind at the end of the third day.*

The effects of intoxicating liquors on the brain and retina are well known. Boër-haave relates the case of a man, who, whenever he was intoxicated, laboured un-

^{*} Anfangsgründe, 3. Band.

der Amaurosis. The disease came on by degrees, increasing with the quantity of wine; and, after the intoxication went off, his vision returned.*

Amaurosis has also been known to arise from other causes which increase the quantity of blood in the encephalon.

The cessation of any accustomed hæmorrhage, such as from the nose or rectum, has been observed to produce Amaurosis.

Pechilini saw a young woman who had Amaurosis during a suppression of the menstrual discharge, which was removed the moment menstruation returned. †

Rolfini mentions examples of women, who, as often as they became pregnant, were blind to the time of their delivery. ‡

Beer saw a woman who was amaurotic during three successive Pregnancies. The disease commenced immediately after each impregnation, and between the third and

^{*} De Morbis Oculorum.

⁺ Ob. Med.

[‡] Disputatio de Gutta Serena.

fourth months she became quite blind. After the two first confinements vision was restored; but after the third, it never returned.*

Portal saw a woman who became a-maurotic after the delivery of her first child; after her second, she became deaf; and, after the third, almost dumb. She had not lost any blood during either of these pregnancies, but was bled during the next, and she had no new affection, the former, at the same time, diminishing. †

Amaurosis has been known to come on after the healing up of ulcers, or drying up of eruptions.

Beer says he has frequently seen Amaurosis produced among the Polish Jews, from cutting off the Plica Polonica.

Amaurosis is sometimes produced from a sudden cessation of the secretion of milk.

Amaurosis has been known to be occasioned by a sudden fright, and other violent mental emotions.

^{*} Lehre der Augenkrankheiten. II. Band .- Wien, 1817.

⁺ Anatomie Medicale.

Amaurotic symptoms frequently come on before or after a fit of Apoplexy. Bonnetus, in a young man who died paralytic, and who, besides losing the use of one side of the body, lost the sight of the eye of the same side, found "the anterior cavity of the cerebrum filled with blood, and a quantity of serum." *

Amaurosis often remains after Fevers, accompanied by affections of the brain. In a gentleman who had a severe attack of fever at the Isle of France, accompanied with a very marked determination of blood to the head, not only a complete and permanent Amaurosis remained, but for many months he was also deprived of the senses of Taste and Smell.

Instances have been observed, where a complete amaurosis has come on from a stroke of Lightning. I have only seen one case of this kind, and the sight was restored by the repeated application of small blisters over the frontal nerve.

Sepulchretum Anatomicum.

I have known two instances where Amaurosis came on after a coup-de-soleil.

Injuries of the Brain are sometimes followed by amaurosis. Richter saw a man who received a smart blow on the ear, and immediately lost his sight.

Richter mentions a person who lost his sight during a violent fit of Vomiting.

Hildanus saw an Amaurosis arise from Sneezing.

Amaurosis is a common symptom of internal Hydrocephalus, both eyes being generally affected; the accumulation of water in the ventricles probably compressing both thalami.

Tumours of the brain frequently produce Amaurosis. Ford found the optic thalamus, in a case of amaurosis, enlarged to the size of a hen's egg. In a young woman, who had lost the sight first of the left, and then of the right eye, subsequent to a violent fever and severe pain in the head, Plater found a tumour in which the optic nerves were involved at that part where they emerge from the brain.

Dr Kühl of Leipsic met with a case of Amaurosis, in a young man of healthy appearance, who died after being blind a few months. On dissection, a tumour, as large as a hen's egg, was found where the optic nerves decussate, part of which was firm, and part filled with a fluid. Bonnetus relates a case where an abscess of the mamillary process, which proved fatal, was accompanied with amaurosis. Bonnetus also found "a steatomatous tumour, the bulk of a fist, between the cerebrum and the cerebellum, producing amaurosis." *

Exostosis has been sometimes met with growing from that part of the sphenoid bone, forming the sella turcica, and on which the conjoined optic nerves lie. In the Josephine Academy, there is a preparation of these parts, taken from a boy who had been affected with Amaurosis; and where a spicula of bone was found growing up from the spot just mentioned,

Sepulchretum Anatomicum.

transfixing the conjunction of the optic nerves.

2.—Amaurosis from Poisons and Narcotics.

Many of those substances which act directly on the nervous system, have an influence upon the Retina, and produce amaurotic symptoms.

Bosman narrates a case, where Amaurosis came on from the saliva of a serpent getting on the face.

A young healthy man, when lying awake early in the morning, perceived a large spider on the corner of his bed. When the animal was immediately over his head, his wife seized it, and it emitted a drop of liquid which fell directly on his eye. He rubbed the eye, and immediately found, that he could see nothing, at the same time there did not appear any external change. *

^{*} Boerhaave de Morbis Oculorum.

In one case, an amaurosis proceeded from the bite of a mad dog.*

Scarpa has observed, that when the infusion of digitalis, stramonium, tobacco, and several other medicines of the same class, are taken into the stomach, they sometimes produce an amaurotic blindness.

A case is mentioned by Conradi, where Amaurosis was produced by the application of the water of the lauro-cerasus every day. A quantity of it was put into the eye, to remove a speck on the cornea; in a short time, the pupil became dilated, and the iris paralytic, with a complete loss of vision. The eye recovered by the frequent application of blisters.

An Amaurosis was in one case produced by the injection of the juice of the belladonna; and was cured by the use of volatile remedies. †

^{*} London Med. Obs. Vol. III.

⁺ Observations on the effect of the Belladonna, &c. &c. by Mr Wishart, in the Edinburgh Medical and Surgical Journal, Vol. III.

Beer himself had a complete Amaurosis, which lasted only a few hours, in consequence of a mistake in the quantity of laudanum, given as a glyster.

In the Mallacca Islands, there is an amaurosis well known to strangers, which is attributed to eating hot barley. In this state, the barley is supposed to throw out a narcotic vapour.

Beer has known Amaurosis produced from the poison of lead, both used as a cosmetic, and taken internally.

Beer mentions that he had seen a woman, who was attacked with violent vomiting, as often as she drank chocolate, and
for several hours afterwards remained stoneblind. He considered this occurrence as
the consequence of the exertion of vomiting; but having had occasion to see her
affected with vomiting, from a different
cause, without the blindness being produced,
he prohibited the chocolate, and she had no
return of the blindness.

3.—Amaurosis from Wounds of the Frontal Nerve.

Wounds of the eye-brow, injuring the Frontal nerve, have frequently been known to produce Amaurosis; the amaurotic symptoms coming on, in some cases, instantaneously, and sometimes long after the wound has been healed.

Morgagni mentions having seen a case of Amaurosis produced by a wound above the eye-brow, which, he observes, explains a passage of Hippocrates. "The sight," says the Father of Medicine, "is obscured in wounds which are inflicted on the eye-brow or a little higher." *

This curious fact I have seen illustrated in several instances, and is an example of the sympathy which exists between parts whose nerves have a direct communication, the ophthalmic branch of the fifth pair send-

^{*} Letter xiii. Art. 5.

ing off the frontal nerve, and also a twig to join the third pair to form the lenticular ganglion.

It is only when the Frontal nerve is wounded or injured, and not divided, that Amaurosis takes place; for, as shall afterwards be observed, amaurosis following a wound of this nerve, may sometimes be cured, by making a complete division of the trunk nearest its origin.

Portal saw a child, who received a slight puncture on the forehead, with the point of a knife, which was followed by a considerable convulsion of the upper eye-lid. This ceased when a small incision was made at the place of the puncture.*

A gentleman received an oblique cut in the forehead, which, from its direction and depth, must have injured the Frontal nerve. The wound was not accompanied by any severe symptoms, and soon healed. But afterwards, the vision of this eye began to fail,

^{*} Anatomie Medicale, Tom. IV. p. 167.

and in a few months was completely destroyed; the pupil was much dilated, the iris was not influenced by variations of light, and had slight tremulous motions.

A sailor got a blow on the edge of the orbit, from a ram-rod during an engagement, at the place where the Frontal nerve passes on the brow. The vision of that eye was instantly destroyed, and when I saw him several years after the accident, the eye remained amaurotic, with a dilated and immoveable pupil. A cataract had formed in the other eye.

An officer, at the siege of Badajos, received a deep wound on the eye-brow by a piece of a shell, which, from its direction, must have injured the Frontal nerve. Great inflammation and pain succeeded the wound, the vision of the eye became gradually imperfect, and, after a few months, was entirely lost. The pupil was very much dilated and immoveable, and the crystalline lens opaque.

Wounds of the Infraorbitary nerve, and portio dura, are sometimes followed by amau-

rotic symptoms. Beer mentions an instance of the former, and I saw an officer in whom a ball wounded some branches of the latter nerve, which was followed by Amaurosis.

This connection between the branches of the fifth pair of nerves and retina affords an useful channel for applying remedies in diseases of the retina. *

4.—Of Amaurosis from Disorders of the Primæ Viæ.

The sympathy which exists between the Chylopoetic viscera, and various parts of the nervous system, particularly the brain, and organs of sense, has long been well known to physiologists. The headachs produced by improper food, and the vomiting brought on by injuries of the brain, are sufficient

^{*} Since the foregoing part of this volume was printed, I have been informed that M. J. Ribes has discovered, that there are filaments which go directly from the atlantal extremity of the great sympathetic nerve, to join the ophthalmic ganglion.

proofs of the reciprocal sympathy of these organs.

The senses of hearing, taste, and smell, are also affected in many of the diseases of the digestive organs; and very frequently these affections influence the functions of the Retina. These diseases ought strictly to be considered as sympathetic, or symptomatic, forming a class very different from those where the amaurotic symptoms arise from a change of structure in any of the parts which compose the organ of vision.

Amaurotic symptoms frequently accompany worms in the intestines. A child who had worms, besides amaurotic symptoms, had nearly lost altogether the sense of hearing, but recovered both senses as soon as the disease of the bowels was removed. Vandermonde relates the case of a girl, who, from worms, and disordered primæ viæ, lost her Vision and Speech. *

Beer remarks, that he has never seen

^{*} Journal de Medicine, Tom. X.

Amaurosis, with complete blindness, from disorders of the primæ viæ, except from worms.

Amaurotic symptoms frequently affect those who are subject to Biliary complaints.

A lady who was subject to violent headachs, and vomiting of bile, sometimes became perfectly blind for several hours.

Richter mentions a man who lost his sight a few hours after being in a violent passion, and recovered it again the next day, by taking an emetic.

A woman became blind whenever she was troubled with what are termed Acidities in the stomach.

A dropsical woman became blind on the water being discharged from her abdomen.*

Those who are afflicted with what are called Bilious headachs, have sometimes amaurotic symptoms. In the common megrim, when the pain affects the frontal branch of the fifth pair, the retina is very liable to be influenced.

^{*} Angfangsgrunde.

This sympathy is strikingly illustrated in wounds of the frontal nerve, and it is probable, that the same nervous connection between the frontal branch of the fifth pair, and the alimentary canal, and between that canal and the retina, depends on the distribution and connections of the fifth pair of nerves. * A gentleman who had long been subject to headachs, affecting chiefly the frontal nerve of one side, and whose digestive organs were always imperfectly performing their functions, became amaurotic in the corresponding eye, the pains in the frontal nerve at the same time ceasing.

Bartholin † relates the case of a monk who lost his sight as often as he shaved his beard, and recovered it again when the beard began to grow. With permission of his superiors, he gave up shaving!

5.—Symptoms of Amaurosis.

I have already remarked how varied and

^{*} See Sympathies of the Eyes, Chap. XLIV.

⁺ Epist. Med. Cent. 3. Epist. 67, n. m. 27.

multiplied are the symptoms of Amaurosis, and how deficient our knowledge is when any attempt is made to refer particular assemblages of symptoms to the morbid changes which accompany them.

It may in general be said, that in amaurotic eyes are observed all varieties of change in the size, form, position, and motions of the Pupil.

That opening is sometimes so much enlarged, that the iris can scarcely be seen, and atother times it is unnaturally contracted.

Sometimes the iris is perfectly immoveable; at other times it retains its natural sensibility. In some instances, that sensibility is increased; the motions of the iris pointing out how far the ciliary nerves are affected, a circumstance of importance to be known. *

The pupil is sometimes regular, sometimes irregular, in its form.

Sometimes the pupil is removed from its natural position, upwards, downwards, inwards, or outwards, but most frequently in a

^{*} See Wenzel on Cataract.

diagonal direction, between upwards and inwards.*

The colour behind the pupil is sometimes black, clouded, greenish, a muddy amber, or of a horny colour.

Sometimes Amaurosis comes on suddenly, and at other times very slowly.

Amaurosis is sometimes periodical, but most frequently permanent. Beer has had several patients who, in the course of from three to four years, have had complete Amaurosis from four to six times.

Vision is sometimes altogether destroyed in Amaurosis, the patient not being able to distinguish between sunshine and complete darkness; and in some instances it is only imperfect.

The imperfections of vision consist sometimes in a diminution of the power to distinguish small objects, these appearing to be concealed by a mist or smoke; † or a diminution in the sphere of vision, ‡ so that the

^{*} Beer. Lehre der Augenkrankheiten.

⁺ Visus Nebulosus.

[†] Hæmyopia-Visus interruptus vel dimidiatus.

person only sees a part of any object. Sometimes objects become distorted;* and a variety of images and colours appear before the eyes of those afflicted with Amaurosis.

A person with Amaurosis sometimes becomes short-sighted, and at other times longsighted.

Amaurotic patients, when they wish to look at a small object, frequently require to have it brightly illuminated. ‡

Amaurosis is, in some instances, accompanied with double vision.

Sometimes the disease is accompanied with, and sometimes it is without, pain.

In some cases of Amaurosis there is an involuntary rolling motion of the eye-ball, accompanied with a peculiar gaze, as if the eye was in search of light.

" But thou

Revisitest not these eyes, that roll in vain To find thy piercing ray, and find no dawn, So thick a drop serene hath quench'd their orbs!"

There is one symptom which has been considered as always accompanying an

^{*} Visus Defiguratus.

⁺ See Ocular Spectra, Chap. L.

[‡] Lighthunger.

Amaurosis,—a peculiar cast, or a confirmed Squint of the diseased eye. Richter has observed, that this is the only symptom which can be depended on when the veracity of the patient cannot be trusted.

In some cases of Amaurosis, where the blood-vessels of the retina and choroid coat seem to be in a varicose state, a corresponding change may be observed in the vessels of the white of the eye.

It is not unusual for a Cataract to be formed in an amaurotic eye; and sometimes the vitreous humour is also disorganized.*

Amaurosis affects people at all periods of life.

Sometimes Amaurosis is Congenital, and, when this is the case, it is usually accompanied both with cataract and the involuntary motion of the eye-ball.

Amaurosis affects eyes of all colours, but is generally supposed to be most prevalent in black.

Amaurosis may be hereditary. I have

^{*} Glaucoma.

known several instances of this kind, and similar cases have been recorded. *

Beer knew a family, in the third generation, of which all the females became amaurotic at the cessation of the catamenia, except those branches who had children. The male branches shewed a disposition to the disease, but none became blind.

Sometimes only one eye is affected with Amaurosis, and sometimes both become blind at the same time. I think it may be remarked, as a general observation, that, when only one eye becomes at first amaurotic from a sympathetic affection, there is little danger of the other eye becoming blind; but that, when Amaurosis is produced by any organic change in one eye, the other is in danger of becoming sympathetically affected. This observation coincides with what has already been said of the sympathy which exists between the corresponding textures of each organ. †

^{*} Edinburgh Medical and Surgical Journal, Vol. XII.

⁺ See page 140. Chap. XLII.

CHAP. XLVI.

OF NIGHT-BLINDNESS.*

Those afflicted with Night-blindness, or Hæmeralopia, have their sight impaired from
the time when the sun sets till it rises the following morning, vision being perfect during
the day.

I have never observed this disease in Britain, but it occurs frequently on the continent of Europe, and in the more southern latitudes.

In some countries it is endemic, and in

^{*} Hæmeralopia.—Disopia tenebrarum.—Ambliopia crepuscularis.—Visus diurnus.—Cæcitas nocturna.—Aveuglement de nuit.—Nachtblindheit.

others it appears as an epidemic at certain seasons of the year.*

In this disease no changes can be observed in the appearance of the eyes, except some slight alteration in the motions of the iris. In some instances the pupil is a little dilated, and loses its natural mobility during the day, whilst, in others, the functions of the iris are perfectly natural, except during the night, when the pupil is always more or less enlarged. †

This disease always attacks both eyes simultaneously.

The progress of Hæmeralopia is sometimes slow, at other times sudden. In the beginning of this affection the person may be able to distinguish objects a short time after sunset, or he may see a little by clear moonlight; but, as the disease advances, vision becomes more imperfect, and he is finally unable to discover even large

^{*} Recueil periodique de la Société de Médecine, Tom. II. Journal de Médecine pour l'année 1756.

[†] Boyer, Maladies Chirurgicales, Tom. V. Paris, 1816.

objects, being only able to distinguish light from darkness.

In some instances the person becomes blind at twilight, and on the following morning at sun-rise he perfectly recovers his sight.

It is a circumstance very remarkable, that those affected with this disease see distinctly in an obscure day, whilst the sun is above the horizon; but, whenever the sun sets, however bright may be its reflected light, such patients see objects very imperfectly. Thus many of them, in a cloudy moment, know the precise time of the sun's setting, though that period cannot be distinguished by other people.*

This disease has generally been observed to continue from three weeks to eight months, when no remedies have been used.

Sometimes Hæmeralopia gets well spontaneously; in other instances, vision becomes more or less impaired; and it has been

^{*} Boyer, Maladies Chirurgicales, Tom. V.

known to terminate in permanent blindness.

Those who have had one attack of this complaint are very apt to have a relapse every year, and at the same season. Boyer relates the case of a man, forty-three years of age, who had been attacked with it every spring from the age of twenty-three.

People of all ages, and eyes of all colours and forms, are equally subject to this disease.

Hæmeralopia is accompanied with pain and weight in the head, which almost always increase towards the evening.

Symptoms of plethora frequently accompany this disease, the patients complaining of giddiness, particularly when the head is bent downwards, or they are occasionally affected with a dimness of sight during the day.*

Night-blindness seems, in most instances, to be connected with derangement of the prime viæ.

^{*} Essay on Hæmeralopia by Mr Bamfield. Medico-chirurgical Transactions, Vol. V.

It is said to affect those labouring under Scurvy, and not to yield until the scorbutic affection is removed.

Night-blindness has been supposed to arise from a mere inability of the retina, to be excited by the slighter stimulus of the evening light, the difference between the splendour of the sun's light, and that of the moon or candles, being immense. * In like manner, as it is observed, that some animals, as owls, cats, &c. see only in a very moderate light; and there are other tribes, as fowls, which become very blind at twilight.

^{*} The strength of moonlight is computed to be 300 000th part of that of daylight.

CHAP. XLVII.

OF COLOURED VISION. *

There is a remarkable peculiarity in vision, which may rather be considered as a congenital imperfection, than a disease. It has been denominated Coloured Vision, the person affected with it having no perception, or at least a very faint one, of some of the principal colours.

When a number of objects of different colours are placed together before a person whose vision is imperfect in this respect, he can distinguish differences between most of

^{*} Suffusio colorans.—Visus coloratus.—Crupsia.—Farben-schen.

the colours, without knowing exactly in what that difference consists; and, in general, if the colours are presented to him separately, he is unable to give them names, or to say to what class they belong.

He has a distinct perception of colours purely yellow or blue, such as gamboge or Prussian blue. These colours he never mistakes, either when seen separately, or when lying amongst other colours; and all other colours seem to him only modifications of these. The colours which appear in the solar spectrum, for example, are to him only two, viz. blue and yellow, whereas, to a perfect eye, there appears in that image to be seven.

He experiences great difficulty in distinguishing the different kinds of green, most of those modifications of colour called green appearing to him either yellow or blue, in proportion as one or other of these colours predominates. Grass appears to be yellow, generally not so light as gamboge, but near the colour of an orange; and, if a piece of

red cloth, and another piece of the colour called olive, be presented to him, though he can perceive some difference between them when they are both together, yet, when seen separately, he continually mistakes one for the other. The appearances of green to his eye depend in some measure upon the quantity of light which falls on objects of that colour, some kinds of green cloth, when the sun shines strongly upon them, appearing yellow; but, in a shaded place, and even in common day-light, they appear like an impure blue.

The different kinds of red create in him equal embarrassment; for vermillion, and all the varieties of red which incline to that colour, appear to him to be yellow, whilst carmine, and all the reds which incline to that colour, appear to him to be blue. An officer's red coat, for example, appears to him to be yellow, and his sash blue. Red ink, when fluid, appears yellow; and, when washed on paper and dried, it appears blue.

These are the appearances of colour to his

eye when seen by day-light; when they are viewed by candle-light, the only difference is, that carmine, and all objects of that class of red, appear yellow, like the vermillion reds; the appearances of vermillion reds, and of all other colours, are the same as during the day.

These defects exist in some people to a much greater degree than in others; and those in whom vision is thus defective, are frequently led to make very absurd mistakes with respect to the appearances of objects. A man, who was employed in a woollendraper's shop, made some ridiculous errors in selling the cloths, from having this defect in his vision.

This imperfection of vision is sometimes Hereditary. Several branches of a noble family in this country have been remarkable for having it. A similar example is given in the Philosophical Transactions,* and another in the Medico-chirurgical Transactions.†

^{*} Vol. LXIII. Part II.

⁺ Vol. VII. p. 477.

It is not improbable that this defect of vision arises from a greater sensibility of the retina to the impressions of the blue and yellowmaking rays, than to those of any of the others. This may depend upon the refractive power of the humours, by which the rays of these two colours are more accurately united on the retina than rays of any other colour, and, consequently, the images formed there of objects reflecting these colours are more distinct than those formed of objects reflecting the other colours. When the colour of a body is compounded of several colours, the superior correctness of the image formed by the blue and yellow rays reflected from it, may cause the sensations which those colours excite to preponderate over the sensations caused by the other colours, and thus may cause in the mind of the observer the perception of that compound colour to be different from the perception of the same colour to another person, whose eye forms images of external objects differently.

The imperfection of vision which has here

been described, though it does not occur frequently, yet is by no means unusual, some striking examples having fallen within my own observation. From these, with the able assistance of Mr Narrien, of the Royal Military College at Sandhurst, the foregoing account of this curious peculiarity of sight has been collected.

When this state of vision is better understood, it is not improbable that it may lead to an explanation of some of the phenomena of light and colours, and explain, more satisfactorily than has hitherto been done, some of the functions of the retina.

that not only the papil, out the whole cor-

nea was hid. The other eye being quite blind.

corner of the squinting eye outwards, which

she had acquired the newer of doing by

CHAPTER XLVIII.

OF SQUINTING.

A PERSON is said to Squint when the axes of the two eyes are not directed to the same point.

The degree of Squint varies in different instances; sometimes the eye having merely a slight cast, whilst in others the eye-ball is very much turned inwards. In one instance, an eye squinted to such a degree, that not only the pupil, but the whole cornea was hid. The other eye being quite blind, the patient could only see by turning the cornea of the squinting eye outwards, which she had acquired the power of doing by

^{*} Strabismus.—Strabositas.—Louche.—Luscitas.—Visus obliquus.—Das Schielen.—Les yeux de Travers.—Distortio oculorum.

pressing with the point of her forefinger in the internal angle till the cornea was exposed. Whilst with her finger she retained the eye-ball in this position, she could see distinctly, whereas, without this contrivance, she was perfectly blind. The squint was attributed to the effects of severe headachs, to which the woman had been subject for many years.

In all cases of squinting, the affected eye is either turned inwards towards the nose, or in the opposite direction, outwards; it has never been observed turned upwards or downwards. When it is considered that the elevation or depression of the globe of the eye are produced by similar muscles in the two eyes, whilst the lateral movements depend on the actions of the adductor of the one, and the abductor muscle of the other eye, we can conceive the frequency of the lateral squint, and why a squint upwards or downwards never exists, or is at least extremely rare.

It has been said by some,* that Squint al-

^{*} Mémoires de l'Academie, 1745.

ways arises from a disparity in the vision of the two eyes, or from a change in the structure of one eye. This, however, is not the case, for sometimes both organs squint where there are specks on the cornea, or from an alteration in the position of the pupils. Neither does a squint always take place where there is a disparity in the vision of the two eyes; for, in many instances where the sight of one eye is diminished, no squint takes place; and, as is observed in another place, * few people see equally well with both eyes.

In some cases of Squint, I have not been able to detect any decided difference in the vision of the two eyes.

The derangement in vision, produced by Squint, is seldom considerable. Some people squint without being conscious of it; others have their vision rendered indistinct, and some have double vision; this being particularly the case where the squint is recent, and not to any great degree. For, as the patient

^{*} See Chap. LI. on the Disparity of the two Eyes.

accustoms himself to look at objects only with the sound eye, the sensation transmitted by the other eye is feeble, and becomes gradually effaced; and this fault may be also corrected by the sense of touch,—a proof that single vision with two eyes is not produced from the images of an object falling on corresponding points in the two retinæ, but the result of habit. Where the Squint is very considerable, the two eyes cannot see the same object, and, consequently, double vision cannot take place.

Buffon observed some people who squinted, make use of the two eyes alternately, according as the objects were placed more or less distant; with the feeble eye they looked at near objects; and with the stronger, objects at a distance.

In most cases the Squint may be removed from the one eye to the other, by shutting up, for a certain length of time, the sound eye, and making use only of the eye which squints. When this has been done, and both eyes opened, the eye which originally

squinted will now be the one whose axis is directed to the object, whilst that which was sound now squints. This was strikingly illustrated in a young lady, who long had a complete squint in one eye from a corneal speck, and whose other eye was attacked with pustulous ophthalmia. The violence of the symptoms made her keep the inflamed eye covered with a shade, whilst the other, with which she saw but imperfectly, served the common purposes of life. In a few days, to the astonishment of all who knew her. what formerly was the sound eye, now became the squinting one; but, as the inflammation subsided, and she began to make use of that eye, the squint gradually left it, and returned to the one originally affected.

A Squint may arise from various causes. It is sometimes produced from Chylopoetic derangement,—from Worms,—from organic affections of the Encephalon,—from Dentition,—from Imitation or Habit,—from long position of the head in one posture,—from the transparency of some of the parts of the

eye-ball being destroyed,—or from a disparity in the vision of the two eyes.

1.—Squint from Chylopoetic Derangement.

Besides changes in the sensibility of the retina, which accompany some diseases of the organs of digestion, I have observed well marked cases of squint, which arose from similar causes.

In cases of this description, the Squint has usually come on gradually, varying in degree at different times; and in some instances, the state of the eye forms an accurate measure of the degree of derangement in the alimentary canal.

This cause of Squint is particularly frequent among young people, though it is met with at different periods of life. A lady thirty years of age, who had frequently during several years the primæ viæ much deranged, had at last a severe attack of pain in her head, extending to the right eye;

the bowels having been at this time unusually confined. The affected eye then squinted, and she had double vision. In proportion as the functions of the bowels were restored, the squint diminished, and before the patient ultimately recovered, every stage of amendment, or occasional change, was distinctly marked by a corresponding diminution or increase in the squint.

2.—Squint from Worms.

It is by no means unusual to see children squint who have Worms. In such cases the squint usually comes on gradually, and whilst it continues, the degree varies on different days, and even several times in the same day. Besides the squint, all the symptoms of worms are present in such cases.

The Squint generally disappears when the worms are removed, whilst in other cases it does not leave the patient.

3.—Squint from Organic Affections of the Head.

In many organic affections of the brain and its meninges, the disease at some period or other is accompanied by a Squint. This is very remarkable in Hydrocephalus, and in diseases which affect parts contiguous to the thalami and tracti optici. Abscesses, and the various kinds of tumours which are formed about these parts, are generally accompanied with a squint; and so are also many affections of the adjacent bone. Indeed, Squint is one of the symptoms which often points out the seat of disease to be in the encephalon.

A blow on the head has in many instances produced a Squint; the squint, in such cases, probably arising from the effects of the blow on the optic thalamus. In some cases, the squint is one of the immediate effects of the injury; but in others, it does not take place

till some time after the accident. In Squints arising from injuries of the head, the distortion sometimes goes off along with the other effects of the accident; but in some cases it has been permanent.

A Squint has also been known to arise from a wound of the Frontal nerve. The nervous anastomosis, which has already been noticed, between the frontal branch of the fifth pair, and other nerves distributed within the orbit,* affords a satisfactory explanation of the peculiar effects of such an accident.

A Squint often accompanies amaurosis.

4.—Squint from Teething.

During the process of Teething, there are a variety of nervous affections occasionally produced, and one of these is a squint.

The squint in such cases is sometimes

^{*} See Chap. XLII.

slight, and goes away when the teething process is completed. In other instances the squint is to a greater extent, and is permanent.

5.—Squint from Imitation and Habit.

Some children have been known to acquire a habit of squinting; and there are many instances of squinting from imitation.

A child two years of age acquired a complete Squint, from living for a few weeks in a house where there was a young lady who squinted. In another instance, a child got a squint whilst teething, and her sister, by frequently imitating her, at last acquired the same habit. A child who suffered from a violent fit of hooping-cough during the night, was observed to squint the following morning. In a short time one of her sisters acquired a squint. I have also known instances of children squinting, whose mothers or nurses squinted.

6.—Squint from unusual Position of the Head.

It sometimes happens, that when a person is long confined in one posture, and the head in such a position, that only one of the eyes can see the window, or can be chiefly employed in looking at objects, the other eye acquires a Squint. But this squint is usually temporary, disappearing by a change in the position of the head, or when both eyes are used to look at objects.

A young lady was confined for several weeks to bed, with a disease of the hip-joint, in a room lighted by a window on one side. She was in the habit of turning her eyes constantly to the light, and thus contracted a squint; which was easily cured, by simply changing her position in the room.

Another young person, from remaining ten days on one side, acquired a squint, which lasted seven years; it was afterwards removed by looking a great deal in the opposite direction.

7.—Squint from Disparity in the two Eyes.

In those who have a great disparity in the vision of the two eyes, it sometimes, though rarely happens, that the person squints. In most instances, where the vision of the two eyes is unequal, there is only a peculiar cast of the eye, as it is usually termed.

There is nothing more remarkable in the history of squinting, and it is the same with regard to double vision, than that, as far as the pathologist can discover, the same disease in the eye does not always produce a squint. Many have a considerable disparity in the vision of the two eyes who have no squint, whether this disparity arises from a speck on the cornea,—from any particular species of cataract,—from differences in the sphericity of the cornea,—or from some

difference in the sensibility of the two re-

I have known a few instances of people, advanced in life, suddenly acquiring a squint, generally accompanied with double vision, when no cause for the disease could be detected.

8.—Squint from Opacities.

When there is a partial obstruction to the passage of the rays of light through the pupil, the disease sometimes produces a squint. This generally happens either from corneal Speck, or from Cataract.

In cases of corneal speck, accompanied by a squint, the speck is so situated on the nasal portion of the cornea, that unless the eye-ball is rolled very much inwards, the rays of light from an external object cannot pass through any part of the pupil to reach the retina. The eye-ball, therefore, being kept constantly inwards, acquires a Squint. In one case where the nasal portion of each cornea had become obscure, the person had a most inveterate Squint in both eyes.

It is chiefly those who have cataracts from birth who Squint. In that disease, it often happens, that only a portion of the pupil is obscure; and when this is the nasal part, the eye, in order to look at objects, is always rolled inwards.

CHAP. XLIX.

OF DOUBLE VISION. *

It has been demonstrated by writers on Optics, that when an object is situated at the concourse of the optic axes of the two eyes, it is seen by both eyes in one and the same direction, and hence single vision is obtained: But if, from disease, the axis of one eye be so distorted that it cannot be directed to an object at the same time with the axis of the other eye, there is produced a double image of that object.

Dyplopia.—Suffusio multiplicans.—Visus duplicatus.—
 Bèvue.—Doppelsichtichkeit.

Thus has arisen the opinion of there being corresponding points in the two retinæ. The phenomena of diseases of the eye, however, refute this doctrine, and afford striking illustrations of the powers of habit in teaching us to see single with two eyes. It is by no means unusual for a person to acquire a squint suddenly, * which is at first accompanied with double vision, and to find by degrees objects appear single. A person had one of his eyes distorted by a blow, and for some time every object appeared to him double; but by degrees the most familiar ones became single, and in time all objects appeared so, though the distortion continued. In like manner, it sometimes happens, that the pupil of one eye is deformed, and its position altered, so that the person sees double; but after a while vision, even under these circumstances, becomes single.

Double vision arises either from some or-

^{*} See Chap. XLVIII. page 201.

ganic change in the eye, or from some derangement in the functions of the retina. *

It has been already noticed, that those who squint often have double vision, when the squint is recent; it has also been observed, that when a change takes place in the position of the pupil of one eye, or when there is more than one aperture in the iris, vision becomes multiplied. The same thing takes place from alterations in the form of the cornea and lens.

In the first of these cases of double vision, the patient sees double only when he looks with two eyes, whilst in the others he sees all objects double with the diseased eye.

A person affected with double vision seldom sees the two objects equally distinct. Commonly the most distinct image is the real one; consequently, the person who has this defect in vision does not often commit mistakes in discriminating the real from the false

^{*} May not the sensation of two bodies, which is produced by the presence of one between two fingers when crossed, be explained on the same principle?

image. This, however, is not always the case.

Double vision sometimes takes place when a person has strained his eyes by looking at minute objects in candle-light; but this disappears when he shuts his eyes for a moment.

Some people only see objects double which are at a distance, and sometimes objects appear double only when looked at in certain directions.

Double vision sometimes affects those who make frequent use only of one eye. I have known several instances of officers of the navy who became affected with double vision from fatigue caused by looking through telescopes.

A Catholic clergyman experienced a kind of double vision whilst he held down his eyes to read, the letters appearing to cross each other. This confusion ceased when he shut one eye, or when he elevated the book to the same height, or higher up than his eyes. He found, however, some difficulty if the book

was placed at the same time to the left. Morgagni conceived that this arose from a weakness in the abductor muscles and depressor muscle of the right eye.*

A young man was frightened and dazzled by lightning when looking at the clouds. An hour afterwards he saw double. On examining his eyes he was found to squint.

A gentleman eat a small piece of pastry between meals, and had immediately uneasy feelings about the stomach. On walking in the open air his sight became dim, and he had double vision, accompanied with severe headaches, all which went off the following day.

^{*} Letter xiii. a. 20.

⁺ Leinch, de visu duplicato.

CHAP. L.

OF OCULAR SPECTRA. *

A PERSON affected with this disease sees one or more bodies floating before his eyes; which are, apparently, situated between the eye and the surrounding objects.

These Spectra assume various forms; sometimes resembling black spots, flies, or insects, pieces of net-work, or streaks. Sometimes they are of a red colour, and resemble flowers with numerous petals, or falling stars.

The Spectra are either fixed or moveable,

^{*} Muscæ Volitantes.—Images before the eyes. Hypochyma.—Suffusio.—La Berlue.

⁺ Visus Lucidus .- Photopsia.

and their appearance is either temporary or permanent.

Sometimes the disease is so slight, and of so short a duration, as to be little remarked by the patient; whereas, in other instances, it continues many years, and is a great hinderance and inconvenience to vision.

The Spectra that are fixed are so called, because they always keep the same position, in respect to the axis of vision. If a person with these spots keeps his eyes unmoved, by looking steadily at the corner of a cloud, at the same time that he observes the spectra, he will be convinced that they have no motion but what is given to them by the eye in pursuit of them.

Those Spectra which are moveable float about even when the eye is stationary, and are sometimes above, below, or at the side of the axis of vision; but they are never in it. These are sometimes seen when the eye-lids are closed.

Spectra appear either in one or in both eyes.

Short-sighted people are least liable to this affection.

Spectra are most troublesome when the sky is clear and serene, or when the person looks at a resplendent object. They are also most distinct when the person looks directly up towards the sky, or at distant objects; and in proportion as the object looked at is near, the less distinct the spectra are, and when at the nearest point, they entirely disappear.

These ocular Spectra arise from different causes, and depend either on a morbid condition of the retina, or an opacity of some of the parts of the eye which are naturally transparent. In the latter case, the opacity must be in the posterior part of the vitreous humour, because experiments, and the principles of optics, prove that no opacity of the aqueous, crystalline, or anterior part of the vitreous humour, can throw a partial shadow on the retina.

It has already been observed, * that, under

^{*} See Chap. XLV.

certain circumstances, the pulsations of the arteries of the retina may be seen; and that various colours produced in the eye by pressure of the finger, or by a stroke on it, seem to arise from unequal pressure on different parts of the retina. It is, therefore, extremely probable, that one and by far the most frequent cause of Spectra is a change in the state of the vessels of the retina. In some cases, this seems to depend on an inflamed state of the vascular membrane of the retina; * in some, on a state of plethora; and in others, a state of what is usually called debility. Hence, in all cases of inflammation, affecting the internal structure of the eye-ball, various spectra appear, as sparks of fire or drops of blood. Spectra almost constantly appear before the eyes of those who have plethoric symptoms, and they generally precede a fit of apoplexy. Black spots are often perceived on the bed-clothes by patients with fevers, or inflammation of the

^{*} See Chap. XLIII.

Spectra are also observable when the eyes are weakened by fatigue, or exposed to bright and dazzling lights; and it frequently happens, that persons of delicate constitutions are affected with them, more particularly those whose digestive organs are deranged.

The influence of diseases of the primæ viæ on the functions of the retina has already been noticed, * and a variety of curious appearances, affecting vision from particular kinds of food, or from whatever disturbs the digestive process, are frequently observed.

There are ocular spectra, as Dr Darwin observes, which are derived from another source. After looking at any luminous object, as the sun, for a short time, so as not to fatigue the eye very much, part of the retina becomes less sensible to smaller quantities of light. Hence, when the eye is turned on less luminous parts of the sky, a dark

spot is seen resembling the shape of the sun. It must, therefore, be from habit and want of attention, that we do not see such spots in all objects every hour of the day.*

*Zoonomia.

ready licentuations, and a variety of con-

ons appearances, diffecting vision from parel

There are obther spectra as Dr. Darwin

observes; which one derived from anothers

faligues there you were part of the rest

CHAP, LI.

OF THE DISPARITY IN THE VISION OF THE TWO EYES.

It seldom happens that any of those organs of the body which are in pairs are both equally perfect. The two kidneys and the two testes are generally different in size; and there is often an unequal secretion of milk from the two mammæ. There is frequently a disparity between the two nostrils; and few people hear with equal acuteness with both ears.

Besides this inequality in those organs of

which there are pairs in the system, the two sides of the body differ from one another. In all animals, as well as in man, a distinct difference may be observed in the two sides of the countenance; the food is usually masticated only by the teeth of one side of the mouth; and there is generally a remarkable difference in the strength and vigour of the limbs of the right compared with the left side of the body.

These observations are exemplified in the Eyes, for the vision of the two eyes will seldom be found equally perfect, and in most people the right is stronger than the left eye.

Throughout the animal kingdom, those animals who have two or more eyes employ them to extend their sphere of vision, but man makes use of both his eyes chiefly as one organ, little advantage being probably derived from his being endowed with a pair, except that of diminishing the risk of injuries and disease.

Few are aware of the disparity that often

exists between the two eyes until, from some accident, they are led to make the comparison; and it is by no means unusual to meet with instances of people who, in making this experiment, have discovered that one eye was entirely blind.

It will generally be found, that not only the right is more perfect than the left eye, but when a person is apparently looking at an object with both eyes, generally only one of the eyes, and that the right one, is actually directed to the object. To demonstrate this, let a spot be covered with the point of the finger when looking at it with both eyes. If the left eye be closed, the point of the finger will continue to appear to cover the spot, and to preserve the same relative situation; but if the right eye be closed and the left opened, then the relative situation of the point of the finger and spot appear altered, the spot being uncovered; proving that, in directing the finger to cover the spot, the right eye had alone been employed.

It might be expected, that a difference in the vision of the two eyes would produce a confusion in the image. This, however, is not the case, as appears both by the foregoing experiment, and from the effects of disease. If the lens be extracted from one eye, the vision of the other eye being unimpaired, the imperfect sight of the one eye does not usually interfere with the perfect vision of the other; and, as has already been mentioned, when treating of double vision, the inconvenience arising from that imperfect sight is soon cured by habit, even when it has arisen from some organic change.

The disparity of the two eyes depends on different causes. In some instances it is produced from a difference in the convexity of the cornea, some people having both a long and a short-sighted eye. In other instances, the difference appears to arise from some imperfection in the retina.

I once saw a child in whom there was a

cover the spot, the right eye had alone been

distinct difference in the size of the two eyes; the parents remarked this at its birth.

Congenital differences in the two eyes are sometimes hereditary,

munion. This is often accompanied with an

list-eve odt do goiton mailion vantaulosof

CHAP. LII.

OF THE INVOLUNTARY MOTION OF THE EYE-BALL.

In some people the eye-ball has a constant tremulous motion, which is usually congenital; but, in some instances, it comes on at a later period of life.

I have already mentioned,* that the iris is liable to a kind of undulatory or tremulous motion. This is often accompanied with an involuntary rolling motion of the eye-ball, but each of these affections in some cases exists separately.

The involuntary motion of the eye-ball often accompanies amaurotic affections and congenital cataract; and, as far as I have been able to observe, this state of the eye-ball always indicates, to a greater or less degree, some disease of the retina.

The involuntary rolling of the eye-ball varies in the rapidity of the motions in different instances. Sometimes it is a mere trembling, whilst in others the movements are both rapid and extensive. Buffon saw it to such a degree, that it was impossible to know the point to which the person wished to direct his eyes.*

I have seen a boy, about ten years of age, who has this involuntary motion of both eyes, accompanied with no other disease; when an infant, his eyes were discerned to move with such rapidity that the cornea could never be distinguished, nor the state of the anterior chamber ascertained. Since his birth, the rapidity as well as the extent of the motions

^{*} Boyer, Oeuvres Chirurgicales.

have gradually diminished, and the disease now exists only in a slight degree.

The involuntary motion of the eye-ball is generally much increased by mental agitation.

This disease always affects both eyes.

In the Albino the eyes have a trembling motion.

Two sisters, who had hair of a remarkably white colour, and whose eyes seemed tender to ordinary light, had their eyes constantly agitated.

Those who have this affection have usually some imperfection in their vision, but this varies according to the nature of the accompanying disease. In general, they require to hold the object very close to the eye, and knit the eye-lids.

A child, on whom I had operated for congenital cataracts, and whose eyes had this involuntary rolling motion, when he looked at an object minutely, brought it near the nose, and pressed the eye-ball with the point of the finger; by which contrivance alone he could distinguish small bodies. This ap-

peared to me to improve his vision more by keeping the eye steady, than altering the length of the axis of the eye-ball. A lady, however, who had not this motion in her eye, could see a small object distinctly, but, in order to read, she was obliged to squeeze, in a similar manner, the eye-ball with the point of her finger.

The Selectic coat is a membrane of the fibrate class, resembling very much in structure, and being considered by some analogists as a continuation of the neutricina of

It is of considerable thickness, though not equally so throughout, being stronger posteriorly, and becoming gradually minner an-

the optic nerve.

The chief use of the Sclerotic coat seems to be, to give that form which it is necessary for the eye to possess as an optical ment; and to serve this purpose the more

CHAP. LIII.

GENERAL REMARKS ON THE SCLEROTIC COAT.

THE Sclerotic coat is a membrane of the fibrous class, resembling very much in structure, and being considered by some anatomists as a continuation of the neurilema of the optic nerve.

It is of considerable thickness, though not equally so throughout, being stronger posteriorly, and becoming gradually thinner anteriorly.

The chief use of the Sclerotic coat seems to be, to give that form which it is necessary for the eye to possess as an optical instrument; and, to serve this purpose the more completely, a considerable portion of it is found to be in some animals osseous.

The sclerotic coat is firm and tough, being composed only of one lamina, the fibres of which are firm, white, silvery, and tendinous; irregularly disposed, both in longitudinal and transverse directions.

The sclerotic coat is pierced by numerous holes for the passage of arteries, veins, and nerves; the posterior ciliary arteries passing through the openings which are formed near the optic nerve, and the anterior ciliary vessels, through those a little behind the cornea. The veins which form the vasa vorticosa of the choroid coat and the ciliary nerves, penetrate the sclerotica obliquely about its middle.

The blood-vessels which are distributed in the sclerotic coat itself, are few in number, and they have but few capillary ramifications; a peculiarity of structure belonging to fibrous textures.*

In examining some of the diseases of the

^{*} Bichât, Anatomic Descriptive, Tom. II. p. 430.

eye, it is of great importance to be able to discriminate the changes which take place in these different sets of vessels on the sclerotic coat, as the appearances of inflammation or turgescence in these are different, when either the iris, the choroid coat, the sclerotic coat itself, or the conjunctiva covering it, is affected.

The external surface of the sclerotic coat is covered by a delicate cellular tissue, which lies between it and the conjunctiva. Its interior surface is smooth and polished, having no connection with the choroid coat but by some vessels and nerves; and it is kept constantly moist by an exhaled fluid.

The Sclerotic coat has little or no sensibility in its natural state, and no nerves have been traced in it.

As it does not seem to be concerned in the performance of any of the peculiar functions of the eye, merely serving as a cup to contain the parts destined for the sense of sight, the sclerotic coat is not liable to many diseases.

The sclerotic coat is occasionally affected

by Inflammation, and it is subject to Staphyloma.

Sometimes the sclerotic coat becomes flaccid and yielding when pressed by the finger, so that if the humours be discharged, instead of preserving a globular shape, it collapses. I have never observed this change without a complete disorganization of the internal parts of the eye.

Sometimes the Sclerotic coat becomes unusually tense, so that the eye-ball feels harder than natural.

The Sclerotic coat is capable of great extension. This is shown in staphyloma and in fungus hematodes.

I have known the Sclerotic coat ruptured from a blow. A woman got a sharp blow with a fist on the eye, and there was a distinct fissure formed in the sclerotic coat, at a little distance from the cornea, and parallel to it. The conjunctiva covering this cleft had not given way, so that the transparent parts within the posterior chamber shone through the fissure. A man struck his eye

against a stick, and ruptured the superior part of the sclerotic coat to the extent of half an inch, in nearly a semicircular direction. A portion of vitreous humour within its capsule protruded through the wound.*

When the sclerotic coat collapses, it usually becomes very much thickened, and whiter than natural.

Morgagni observed, in an eye which was shrunk, the sclerotic coat unusually hard and thick.

When the vitreous humour is absorbed, and the sclerotic coat collapses, it usually becomes puckered at four points, forming the ball into as many segments. This is probably produced from the action of the recti muscles.

The sclerotic coat, like other fibrous membranes, has been found Ossified. Blazius found "a bony scale inherent in the sclerotic coat." †

Mr George Young saw a child who was born with no vestige of an eye-ball, ex-

^{*} See Edinburgh Medical and Surgical Journal, 1816.

[†] Morgagni Epist. lii. a. 30.

cept a body like a small pea, which had some resemblance to a shrunk sclerotic coat.

seat of inflammation, but if frequently parti-

tom that the red vessels are scated in the

or tris invergentianied. An the first case, the

VOL. II.

CHAP. LIV.

OF INFLAMMATION OF THE SCLEROTIC

The sclerotic coat is seldom the primary seat of inflammation, but it frequently participates in inflammation of the adjacent textures. Though what is commonly called the white of the eye becomes inflamed, it is seldom that the red vessels are seated in the sclerotic coat. The redness usually arises either from the conjunctiva, the choroid coat, or iris being inflamed. In the first case, the vessels are distributed over the whole surface of the eye, being moveable with the folds

of the conjunctiva: In the second, the posterior ciliary arteries and veins are enlarged; and in the third, the anterior ciliary vessels are chiefly affected.

It has already been observed, that the Sclerotic coat is more or less inflamed when the substance of the cornea is affected with inflammation, * and as Rheumatism usually affects fibrous membranes in other parts of the body, it is extremely probable that in the Rheumatic Ophthalmia the disease is chiefly seated in the sclerotic coat; for when Rheumatism attacks the eye, I have frequently observed, that the blood-vessels on the white of the eye appeared neither to be those of the conjunctiva inflamed, nor the ciliary vessels, but a general redness pervading the whole sclerotic coat. In corroboration of this opinion, it may be remarked, that when the Rheumatic Ophthalmia subsides, the sclerotic coat has generally lost its natural pearl whiteness, and acquired a dingy yellow

^{*} See Chap. II. Vol. I.

hue. The pain, too, in Rheumatic Ophthalmia, is usually seated in the head, and parts of the face surrounding the orbit, probably arising from the similarity, or, as some have supposed, from the continuation of structure between the sclerotic coat, dura mater, and periosteum.

As in inflammation of other fibrous membranes, there is no subsequent thickening, opacity, or adhesions, where the sclerotic coat has been inflamed; but it sometimes happens that puriform matter collected in the posterior chamber ulcerates, and forms an opening through the sclerotic coat.

selectic cost has generally lost its natural

pearl waiteness, and acquired a dings yellow

CHAP. LV.

In a few instances where the eve

OF STAPHYLOMA OF THE SCLEROTIC COAT.

STAPHYLOMA of the sclerotic coat is a rare disease. Richter says he has seen only a single instance of it, and Scarpa never observed one; several cases, however, have fallen within my observation.*

A young man, who had at a distant period suffered from severe inflammation in his eye, had the anterior chamber much diminished, the pupil was irregular, and the vision destroyed. On the sclerotic coat of this eye were several small tumours at a little distance from its union with the cor-

^{*} See Plates of Staphyloma, Vol. I.

nea. These tumours were of a dark blue colour; felt soft to the touch; and seemed to contain a fluid within a delicate membrane.

In a few instances where the eye had suffered from internal ophthalmia, I have observed the whole eye-ball enlarge, and the sclerotic coat become distended, acquiring the blue appearance of staphyloma at that part where it unites with the cornea.

Staphylomatous tumours sometimes form on the sclerotic coat in consequence of injuries. A small tumour of this kind is represented in Plate X. fig. 3.

There is another disease of the sclerotic coat described by Scarpa, under the name of Staphyloma, of which I have never seen an example. In two eyes affected with this disease which Scarpa dissected, the eyeball had assumed an oval shape, and was larger than the other. On the posterior hemisphere, and on the temporal side of the entrance of the optic nerve, the sclerotic coat was elevated in the form of an oblong tumour

the size of a small nut. The humours were changed; the retina was deficient within the staphyloma; and the choroid and sclerotic coats forming the tumour were so thin as to admit the light.

the size of a small mut. "The humours were changed; the retina was deficient within the staphyloma; and the rehorold and sclerotict coats forming the tumour were so thin as to admit the lights? The same secretary was a secretary and a sec

There is not be Secrete, and in the Secrete and Secret

EXPLANATION


OF THE

Plates.


EXPLANATION

STATE OF THE PARTY.

. Eninka



Inflammation of the Capsule of the aqueous humor.



Albumen effused on the Tris.



Albumen effused in the Papil & in which a red vefsel is distributed.



EXPLANATION

OF THE

PLATES.

PLATE VIII.

This Plate is intended to illustrate the appearances and effects of inflammation of the Capsule of the Aqueous Humour.

Fig. 1.—Represents the peculiar appearances which the blood-vessels assume in that affection, as well as the kind of muddiness and opacity of the anterior chamber. The drawing was made from the eye of a boy thirteen years of age, whilst the inflammation was in its active stage.

The vascular trunks divide into different branches, which do not reach the cornea, but penetrate the sclerotic coat at some distance from its margin, and form a red ring parallel to its circumference. These are probably the anterior ciliary vessels enlarged. The inferior hemisphere of the cornea is clouded and obscure, and the pupillar opening filled with albumen.

The appearances in the drawing may be compared with the other inflammations which have been delineated in Plate I. Volume I.

Fig. 2.—Shows a portion of concreted albumen adhering to the surface of the iris, whilst, at the same time, the whole iris has acquired a greenish colour, from a deposition of albumen throughout its substance, as well as from a thickening of the capsule of the aqueous humour.

In some cases, this albuminous effusion disappears along with the inflammatory symptoms; but in others, particularly where these have not been speedily subdued, the albumen remains permanent, as in the example here represented.

Fig. 3.—The albuminous effusion is confined to the pupillar opening, which, in this case, is completely filled with a portion of a yellowish coloured substance, to which a red vessel may be seen coming off from the edge of the iris, to be distributed through it. The form of the pupil is also altered from adhesions having taken place between the albumen and edge of the iris.

These appearances were the consequence of tedious inflammation, and are frequently observed in eyes that have suffered from gout.

Fig. I .- This drawing was ande from the


eye of a hoy who, six weeks before, had a hog's bristle thrust into the anterior chamber. An opaque white matter, resembling a fake of snow, extends from the the least which is all so opaque, through the pupil and concerned by in contact with the cornea. Some inflammation which followed the injury soon above antion which followed the injury soon above

PLATE IX.


THE three figures of this plate are intended to illustrate different effects of Injuries of the anterior chamber of the eye.

Fig. 1.—This drawing was made from the eye of a boy who, six weeks before, had a hog's bristle thrust into the anterior chamber. An opaque white matter, resembling a flake of snow, extends from the lens, which is also opaque, through the pupil, and comes nearly in contact with the cornea. Some inflammation which followed the injury soon abat-


Albumen offused from a wound of the Capsule of the lons.



Albumen effused in the anterior Chamber after the operation of Couching .



Albumen offused in the anterior Chamber, after the operation of Extraction.




ed, and all this opaque matter was ultimately absorbed. It is probable, that, in this case, the bristle had penetrated not only the cornea, but also the capsule of the lens, thus allowing the thinner parts of the lens to come through the wound into the anterior chamber. This appearance is very different from that produced by inflammation of the capsule of the aqueous humour, and precisely resembles the effects of the operation of Keratonixis, where the capsule of the lens is wounded by the point of the needle. I have several times had an opportunity of observing similar effects of injuries to that represented in this drawing. In all of them the opaque matter has been ultimately absorbed.

In Fig. 2.—The whole anterior chamber is filled with a yellow matter, whilst the conjunctiva covering the eye-ball is so much inflamed, as to produce what has been called Chemosis. This eye had been couched a few days before the drawing was taken, and though vision was not restored, yet the ef-


fused albuminous fluid was finally absorbed.

Fig. 3.—This represents the appearances of a net-work of lymph, as referred to in p. 14, after the extraction of a cataract. Towards the inferior edge of the cornea, a white semicircular cicatrix is perceptible; the pupillar opening is of a full size, and throughout the whole anterior chamber may be perceived filaments of coagulated albumen interwoven in various directions, so as to form a sort of net-work. This case is narrated by Beer.


Flakes of coagulated albumen are often found floating in the anterior chamber, or loosely attached to the iris, of the eyes of horses and other animals.



Tres prolapsed through two Ulcers of the Cornea



Laceration of the Tris



Laceration with Absorption of the Jus



PLATE X.

Fig. 1.—Is intended to illustrate the appearances of Prolapsus of the Iris, in consequence of ulceration of the cornea.

In this case, two Ulcers, such as have been delineated in Plate II. fig. 2., had penetrated the cornea, given an outlet to the aqueous humour, and portions of the Iris passed through the opening. The whole cornea has become clouded and vascular, and round each portion of iris there is an opaque white circle, produced from a substance resembling wet chalk;—an appearance rather unusual.

VOL. II.

The whole white of the eye is considerably inflamed.


The two next Figures are intended to illustrate the effects of Injuries of the Iris.

Fig. 2.—Represents a small portion of iris torn from its attachment with the ciliary ligament. In this instance, the laceration was the consequence of a blow with a whip, which did not produce any other injury, and was followed by no defect in vision.

Fig. 3.—In this case, the whole Iris, except a small portion, is destroyed, the person having had his eye wounded by a thorn.

A small Staphylomatous tumour has arisen at that part of the sclerotic coat where the thorn entered the eye.

This man could see so as to be enabled to read, by looking through a small hole made in a card.






PLATE XI.

No. 1. and 2.—Are drawings of two Congenital Cataracts, in both of which the central part of the lens is converted into an opaque white or chalky-looking substance.

In No. 1.—This central opacity has a triangular form, and is imbedded in the transparent lens.

In No. 2.—The central opacity has a more oval form. Exterior to it there is another opacity, resembling what is met with in the more common kind of cataract, and the ex-

ternal margin of the lens remains perfectly transparent.

No. 3—Shows the radiated appearance which the crystalline lens sometimes assumes in Cataract. The rays come from the centre of the lens, but are lost towards its circumference. The central part is of an amber colour, and of a firm consistence, whereas, towards the circumference, the amber hue fades into an opaque leaden grey, the lens becoming much softer, and nearly pulpy.

The drawing was made from a lens recently extracted from a middle-aged patient; its appearance in the eye being of a pale grey colour.

No. 4.—Shows a Capsular Cataract after being removed from the eye. In this case, the anterior portion of the capsule has become perfectly opaque, and acquired a great degree of thickness, being as thick as a com-


mon wafer. The posterior portion, which is of its natural thinness and transparency, is raised with the point of a pair of forceps. This cataract was removed from a young man who had suffered severely from ophthalmia. When the inflammatory symptoms had subsided, a dark yellow opaque body appeared behind the pupil, resembling Plate VIII. fig. 3. the pupil being immoveable from its adhesions to the opacity. This opaque substance was the thickened capsule, on extracting which, the vitreous humour was found converted into a watery fluid, and the retina had lost its sensibility.

The two remaining figures show the appearances of Ossification of the Lens and Capsule.

No. 5.—Is a perpendicular section of a lens, showing its appearance when ossified, as described in Chap. XXXIV. The Ossification is seen commencing in the centre of the lens, and extending towards its circum-

ference in the form of concentric bony laminæ. The central portion was a dark brown coloured and hard bone; the exterior laminæ were of a paler colour, and more friable.

No. 6.—Represents an Ossification of the Capsule of the lens. In this case, nearly the whole capsule, particularly its anterior portion, was converted into a shell of bone. Where the shell is not complete, the dried lens is seen lying within it. The Ossification was in this case of the thickness of a piece of paper, and had all the external characters of bone. The case is particularly described in Chap. XXXVIII.






PLATE XII.

THESE figures show several different kinds of Cataract.

No. 1.—Shows the appearances of the more usual form of the Capsular Cataract in the adult, where the anterior portion of the capsule has become much thickened, and of a pearl-white colour.

No. 2.—In this eye may be perceived the usual appearances of *Glaucoma*, or of cataract combined with a dilated and im-

moveable pupil. The lens has a bluish-grey colour, but not of an equal degree of opacity, and the shade varied so much when the eye was inspected in different directions, that those who looked at it compared it to a cat's eye.

The pupil was permanently dilated, and at one part a portion of the iris is eroded. The lens was extracted, and found to be soft. The lady from whom the drawing was taken had an incipient cataract and Amaurosis of the other eye.

No. 3.—Exhibits one of the more usual appearances of the Crystalline Cataract. The lens is of an opaque bluish-grey colour, and of an equal shade throughout. The iris is beautifully streaked. The other eye was sound. The drawing was made from a girl three years of age, and the cataract had been perceived several months.

No. 4.—This figure represents the peculiar form of cataract which may be denominated

Laminated, (see p. 109.) There is, in the pupil, a very opaque white central spot, and around it a less opaque ring, the limits of both being accurately defined.

No. 5.—This figure represents a Cataract formed in consequence of an Injury of the lens. A needle had penetrated the anterior chamber of a young lady's eye, torn the edge of the iris, and punctured the lens. The pupil is seen of an irregular oblong form, and is separated into two portions by a slender filament of iris, detached from the edge of the pupil; behind one portion lies the opaque lens, whereas the other remains perfectly transparent.


In cases of this description, the wound which renders the lens opaque, seems to destroy its organization.

The drawing was made three months after the accident.

No. 6.—This represents a lateral view of a conical-shaped opaque body, whose base

rests on the capsule of the crystalline lens, and whose apex reaches the cornea. The pupil was quite moveable, vision was impaired, and both eyes had been affected with the disease from birth. The drawing was taken from a boy four years of age.

This peculiar form of cataract, which I have denominated *Pyramidal*, is described in Chap. XXXVIII. p. 112.






PLATE XIII.

In this Plate are illustrated some of the diseases of the Crystalline Capsule.

No. 1.—This is an example of the anterior portion of the Crystalline Capsule become very much thickened and opaque from inflammation. The pupil has lost its circular form, and is immoveable, whilst the opaque body behind it is of a pale grey colour.

No. 2.—Congenital Cataract in which both

the Lens and its Capsule are opaque. There is a green opacity throughout the lens, and in the capsule there are numerous spots. The pupil is dilated by belladona, so that the whole opacity is exposed to view. This figure is copied from the work of the late Mr Saunders, as exhibiting very distinctly the appearances of one form of diseased capsule.

No. 3.—In this figure are contrasted the appearances of an opacity of a portion of the Capsule of the lens, and of the Cornea. The opaque portion of capsule hangs from the upper edge of the pupil, whilst the opacity of the cornea is opposite to the inferior portion of the iris. In this case, the operation of extraction was performed in the usual manner, and no bad effects arose from the incision being carried through the diseased portion of cornea. Whilst making efforts to remove the capsule, the lens escaped. The capsule thus let loose floated upwards behind

the iris, and the pupil afterwards remained transparent, and vision was restored.

No. 4.—In this instance, the whole edge of the Iris adhered to the Capsule of the lens, and the iris had acquired a convexity, being pressed forwards in its middle part. The capsule has become quite opaque, and a small red vessel passes into it from the edge of the iris. These appearances are very frequent after attacks of Gouty inflammation of the eye.

No. 5.—This drawing shows the appearance of the Pupil and Capsule of the lens, after extraction. The irregularity in its form did not proceed from any injury done to the Iris during the operation, but from a portion of the capsule of the lens, which was pushed through it, adhering to the wound of the cornea, and become opaque.

No. 6. — Circumscribed opacity of the Crystalline Capsule, in consequence of a punctured Wound.


PLATE XIV.

Fig. 1.—Represents an Ossification found within an eye-ball, which appeared to be completely disorganized. The patient died of phthisis, but no history could be obtained of the diseased eye. The general form and shape of the eye were completely changed; the cornea was opaque, its margin not distinctly marked, and the anterior chamber almost entirely obliterated. The sclerotic coat was easily separated from the parts within, and upon removing it, the cavity was found occupied by a hard irregular-shaped mass










of bone. The Ossification was covered by the choroid coat and iris, these having lost their natural appearance, and no vestige of retina could be detected.

The Ossification consisted of two different portions. The upper one was smooth, and a thin hollow shell, which, from its rounded form and position, must have been the Capsule of the Lens ossified. The inferior portion was a very irregular-shaped mass, and appeared to be the Capsule of the Vitreous humour converted into bone.

Fig. 2.—Ossification of the capsule of the lens, and hyaloid membrane, described in Chap. XL. the drawing having been made from a preparation where the parts had been dried, and kept in turpentine, in the collection of Mr Blizzard, to whom I am indebted for this representation. The ossified capsule of the lens is seen in the centre of the preparation, and is readily distinguished from the other pieces of bone by its smooth surface, and more opaque white colour.

Fig. 3.—A thin cap of Bone which was found between the sclerotic coat and retina. A few thin and easily torn cellular laminæ were observed on the external surface of this ossification, and seemed to be the only remains of the choroid coat. At the apex is seen a small rounded perforation, through which passed the retina, to be expanded on the interior surface of the ossification. This eye was Amaurotic, but the particular history of the case was not known. The gentleman was also blind of the other eye, but in it there were no appearances of diseased structure.

In these three figures, the Ossifications in the two first resemble one another, that of the third being quite different.






PLATE XV.

Fig. 1.—Is the drawing of a Tumour on the Optic nerve from Mr Heaviside's collection, for which I am indebted to Mr Howship. The tumour appears to have formed in the Neurilema of the nerve, and no further history of the patient was known, than that he was Amaurotic of this eye.

Fig. 2.—Represents a vertical section of the eye-ball, described in Chap. XXIX. p. 67, where a watery fluid had collected between the choroid coat and retina, in

such quantity, as to compress the retina into a chord, and produce a complete absorption of the vitreous humour.

This drawing represents the anterior portion of the eye-ball, so that the compressed retina is distinctly seen terminating at, and closely surrounding the neural portion of the crystalline capsule.

a, a, a, The outline of the sclerotic coat.

b, b, b, The choroid coat.

c, The Retina covering the neural portion of the crystalline capsule.

d, Section of the Retina compressed into a chord.

The eye in this instance appeared, on a superficial examination, to be affected with cataract, a white substance being seen behind the pupil. An attempt was made to couch what was conceived to be the opaque lens,—a fruitless operation, which gave great pain.

THE END.

Printed by George Ramsay and Company. . 1818.

