### **Human physiology (Volume 2).**

#### **Contributors**

Dunglison, Robley, 1798-1869. National Library of Medicine (U.S.)

### **Publication/Creation**

Philadelphia: Lea and Blanchard, 1844.

#### **Persistent URL**

https://wellcomecollection.org/works/ex27ug6f


#### License and attribution

This material has been provided by This material has been provided by the National Library of Medicine (U.S.), through the Medical Heritage Library. The original may be consulted at the National Library of Medicine (U.S.) where the originals may be consulted.


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.











3375F

# Dunylison's Practice of Medicine,

OR A TREATISE

# ON SPECIAL PATHOLOGY AND THERAPEUTICS.

### BY ROBLEY DUNGLISON, M.D.

PROFESSOR OF THE INSTITUTES OF MEDICINE, ETC., IN THE JEFFERSON MEDICAL COLLEGE, PHILADELPHIA, LECTURER ON CLINICAL MEDICINE, AND ATTENDING PHYSICIAN AT THE PHILADELPHIA HOSPITAL, ETC.

#### CONTAINING

The Diseases of the Alimentary Canal — the Diseases of the Circulatory Apparatus —
Diseases of the Glandular Organs — Diseases of the Organs of the Senses — Diseases
of the Respiratory Organs — Diseases of the Glandiform Ganglions — Diseases of
the Nervous System — Diseases of the Organs of Reproduction — Diseases involving
various organs, &c., &c.

### In Two Volumes, Octavo.

"We have great pleasure in welcoming this important addition to American Medical Literature, indicating, as it does, untiring research, experienced judgment, and enlarged and accurate observation. So far as our knowledge extends, this work affords the most comprehensive view of the present state of Medical Science, which has yet been offered to the profession. The author has succeeded in being 'essentially eclectic.' He has wisely avoided the introduction of controversy, although he has not failed to evince, in his treatment of some topics, that he lacks not the skill by which the mazy subtleties of false reasonings are so successfully detected. The not insignificant results of the professor's own practical experience, are given with an unobtrusive propriety, that strikingly contrasts with the bearing of some modern authors, whose aim appears to be the presentation, not of the science, but of themselves, their practice, and their views to the public. The style is that of an accurate and practised writer, and the volumes exhibit throughout evident marks of careful preparation. The publishers have, of course, discharged their duties to the work, d brought it out in a beautiful style. We predict that it will meet as ready and extensive a e as it deserves, and the parties most interested in the fate of the work, must be very unreasonble if that does not prove amply satisfactory." — New York Lancet, Feb. 1842.

"It" (the work) "makes a part of the great scheme of labor-saving and power-giving means and contrivances of the day. It brings together and presents to its readers, under a succinct and condensed form, a vast amount of medical matter, which no physician could collect for himself, without a most tedious and toilsome examination of detached works, and which, from a want of leisure and books, the great body of physicians in the United States could never collect. By its copious references to authorities, it serves to those who are solicitous of further information on given topics, as an index to libraries to which they may have access. It presents to the physicians of the United States a very valuable and praiseworthy example of literary research, ambition and perseverance."

— The Western Journal of Medicine and Surgery, June, 1842.

"We hail the appearance of this work, which has just been issued, with no ordinary degree of pleasure. Comprised in two large and closely printed volumes, it exhibits a more full, accurate and comprehensive digest of the existing state of medicine than any other treatise with which we are acquainted in the English language. It discusses many topics—some of them of great practical importance, which are entirely omitted in the writings of Eberle, Dewees, Hosack, Graves, Stokes, McIntosh, and Gregory; and it cannot fail, therefore, to be of great value, not only to the student, but to the practitioner, as it affords him ready access to information of which he stands in daily need in the exercise of his profession. It has been the desire of the author, well-known as one of the most abundant writers of the age, to render his work strictly practical; and to this end he has been induced, whenever opportunity offered, to incorporate the results of his own experience with that of his scientific brethren in America and Europe. To the former, ample justice seems to have been done throughout. We believe this constitutes the seventh work which Professor Dunglison has published within the last ten years; and, when we reflect upon the large amount of labor and reflection which must have been necessary in their preparation, it is amazing how he could have accomplished so much in so short a time."—Louisville Journal.

"The science of Medicine and the art of Medicine are both very well posted up, not up to the beginning of the century or to the close of the late war — or to the first year when the Professor began to lecture (as would be the case were some learned professors to print their lectures) but up to the present time, up to 1842. The student will here find a clear distinct digest of medical science — the practitioner will here see at a glance all that has been done, on any important point in practical medicine upon which he may need information. We wanted something of the kind very much, and this supplies the want. The text books which teachers have been obliged to place in the hands of their students, "faute de mieux," are all extremely defective. Those of Eberle and Dewees are evidently not the productions of medical scholars; that of Mackintosh, though full of practical sagacity, is exceedingly defective, and besides abounds in whims and notions; that of Gregory, though perhaps upon the whole the best, though full enough upon the art of medicine is lamentably deficient in the science of disease. The work of Professor Dunglison is superior to either, and will, we hope, soon supersede them. Most assuredly it shall with those over whose medical studies we may have control." — New York Medical Gazette, March 2, 1842.

### MEDICAL LEXICON, BROUGHT UP TO 1842.

A NEW DICTIONARY OF MEDICAL SCIENCE; Containing a concise account of the various Subjects and Terms, with the French and other Synonymes, and Formulæ for various Officinal and Empirical Preparations, &c. Third edition, brought up to 1842. By ROBLEY DUN-GLISON, M.D., Professor in the Jefferson Medical College, &c. In One Volume, royal 8vo.

"The present undertaking was suggested by the frequent complaints, made by the author's pupils, that they were unable to meet with information on numerous topics of Professional Inquiry,

— especially of recent introduction, — in the medical dictionaries accessible to them.

"It may, indeed, be correctly affirmed, that we have no dictionary of medical subjects and terms which can be looked upon as adapted to the state of the science. In proof of this the author need but to remark, that he has found occasion to add several thousand Medical Terms, which are not to be met with in the only medical lexicon at this time in circulation in this country.

"The present edition will be found to contain many hundred Terms more than the first, and to have experienced numerous Additions and Modifications.

"The author's object has not been to make the work a mere lexicon or dictionary of terms, but

"The author's object has not been to make the work a mere lexicon or dictionary of terms, but

to afford, under each, a condensed view of its various medical relations, and thus to render the work an epitome of the existing condition of Medical Science."

This New Edition includes, in the body of the work, the Index or Vocabulary of Synonymes that was in the former Editions printed at the end of the Volume, and embraces many Corrections, with the addition of many New Words.

### A NEW WORK --- DUNGLISON'S

### THERAPEUTICS AND MATERIA MEDICA.

GENERAL THERAPEUTICS AND MATERIA MEDICA, ADAPTED FOR A MEDICAL TEXT-BOOK, BY ROBLEY DUNGLISON, M.D., Professor of Institutes of Medicine, &c., in 2 vols. 8vo. - Just ready.

A second edition of the work on General Therapeutics, being called for by the publishers, the author has deemed it advisable to incorporate with it an account of the different articles of the Materia Medica. To this he has been led by the circumstance, that the departments of General Therapeutics and Materia Medica are always associated in the Medical Schools. The author's great object has been to prepare a work which may aid the Medical Student in acquiring the main results of modern observation and reflection; and, at the same time, to be to the Medical Practitioner a trustworthy book of reference.

Throughout, he has adopted the Nomenclature of the last edition of the Pharmacopæia of the United States, a work which ought to be in the hands of every practitioner as a guide in the pre-paration of medicines; and he has endeavored to arrange the articles in each division, as nearly

as he could, in the order of their efficacy as Therapeutical agents.

### MEDICAL REMEDIES.

NEW REMEDIES. THE METHOD OF PREPARING AND ADMINISTERING THEM; THEIR EFFECTS UPON THE HEALTH AND DISEASED ECONOMY, &c. &c., by PROFESSOR ROBLEY DUNGLISON. Fourth edition, brought up to 1843. In one volume octavo.

This work contains articles that have been recently introduced into the Materia Medica; or old articles, that have received new applications; some of these are in the general works on Materia Medica, but their properties are only briefly referred to. In this work, the experience of individuals is extensively given, with reference to the original papers. Under Iodine, for example, all the information — pharmaceutical and therapeutical — up to the time of the publication of the work, is afforded, with the prescriptions that have been proposed by various observers; and interpretated with it the result of recent experience and is therefore each successive edition has incorporated with it the result of recent experience, and is therefore " new.

# HUMAN

# PHYSIOLOGY.

WITH

UPWARDS OF THREE HUNDRED ILLUSTRATIONS.

BY

# ROBLEY DUNGLISON, M.D.,

PROFESSOR OF THE INSTITUTES OF MEDICINE, ETC., IN JEFFERSON MEDICAL COLLEGE,
PHILADELPHIA; ATTENDING PHYSICIAN AND LECTURER ON CLINICAL
MEDICINE AT THE PHILADELPHIA HOSPITAL; SECRETARY TO THE AMERICAN
PHILOSOPHICAL SOCIETY, ETC.,

" Vastissimi studii primas quasi lineas circumscripsi." - HALLER.

FIFTH EDITION,

GREATLY MODIFIED AND IMPROVED.

IN TWO VOLUMES.

VOL. II.



PHILADELPHIA:
LEA AND BLANCHARD.
1844.

QT D916h 1844 v. 2

[Entered, according to the Act of Congress, in the year 1841, by Robley Dunglison, in the Clerk's Office of the District Court for the Eastern District of Pennsylvania.]

Printed by Barrington & Haswell.

# CONTENTS OF VOL. II.

### BOOK II.

118 - 12 Alle Alle Alle Alle Alle Alle Alle All	AGE
CHAP. III. Respiration	13
1. Anatomy of the Respiratory Organs	13
2. Of Atmospheric Air	22
3. Physiology of Respiration	26
a. Mechanical Phenomena of Respiration	26
1. Inspiration	27
2. Expiration	32
3. Respiratory Phenomena concerned in certain Func-	0,0
tions	35
4. Respiratory Phenomena connected with Expression	38
b. Chemical Phenomena of Respiration	42
c. Cutaneous Respiration, &c	60
d. Effects of Section of the Cerebral Nerves on Respira-	-
tion	61
e. The Respiration of Animals	64
CHAP. IV. Circulation	66
1. Anatomy of the Circulatory Organs	67
a. Heart	67
b. Arteries	74
c. Intermediate or Capillary System	77
d. Veins	83
2. Blood	88
3. Physiology of the Circulation 1	112
a. Circulation in the Heart 1	14
b. Circulation in the Arteries 1	30
c. Circulation through the Capillaries 1	36
d. Circulation in the Veins 1	42
e. Forces that Propel the Blood 1	44
f. Accelerating and Retarding Forces 1	53
g. The Pulse 1	60
h. Uses of the Circulation 1	65
i. Transfusion and Infusion 1	66
4. Circulatory apparatus in animals 1	68
CHAP. V. Nutrition 1	71
VI. Calorification 1	86
VII. Secretion 2	21
1. Anatomy of the Secretory Apparatus 2	21
2. Physiology of Secretion 2	224
1*	

			PAGE
04		-	233
Of the Exhalations		-	234
1. Internal Exhalations		_	234
a. The Serous Exhalation			235
b. Serous Exhalation of the Cellular Membrane			235
c. Adipous Exhalation of the Cellular Membrane			241
d. Exhalation of the Marrow	•		242
e. Synovial Exhalation -	24		240
f. Exhalation of the Colouring Matter of the Ski	n and	u	049
other parts	-	-	243
g. Areolar Exhalation		•	244
2. External Exhalation	-	. 85	244
a. Cutaneous Exhalation or Transpiration -		-	244
b. Pulmonary Transpiration	-	7.	254
c. Exhalation of the Mucous Membranes -			258
Follicular Secretions	-		258
a. Mucous Follicular Secretion		-	258
b. Follicular Secretion of the Skin -	-	6	259
Glandular Secretions		-	260
a. Secretion of the Tears	-		260
b. Secretion of the Saliva		-	261
c. Secretion of the Pancreatic Juice -		-	- 263
d. Secretion of the Bile			265
e. Secretion of Urine	12.	1	281
	10	200	297
f. Connexion between the Stomach and the Kidney	3		300
Glandiform Ganglions		100	
a. The Spleen		7:	300

### BOOK III.

#### REPRODUCTIVE FUNCTIONS.

CHAP.	I.	Generation			- 307
		1. Generative Apparatus		-	315
		a. Genital Organs of the Male	1. 1-		- 315
		1. Sperm	2		325
		b. Genital Organs of the Female			- 331
		1. Menstruation			342
		c. Sexual Ambiguity			- 351
		2. Physiology of Generation			355
		a. Copulation	0.		- 356
		b. Fecundation	- 3		360
		c. Theories of Generation			- 381
		1. Epigenesis			381
		2. Evolution			- 384
		a. Conception	-		398
		b. Superfectation			- 408
		c. Pregnancy		355	410
		d. Signs of Pregnancy			- 419
		e. Duration of Pregnancy			425
		P Ti	No contract		- 428
		g. Lactation		100	
CHAP.	II.	Fætal Existence. — Embryology			40.4
		1. Special Anatomy and Histology of the Fœtus	de la		- 443 443
					443

### CONTENTS.

								TD	AGE
	a. Dependencies of the Fœtus	-				-		- 19	443
	b. Development of the Ovum		-		-		-		458
	c. Peculiarities of the Fœtus	4 11		-		-		-	475
	2. Physiology of the Fœtus		-		-		-		482
		-		_		- 11		-	482
	a. Animal Functions -					30			484
	b. Functions of Nutrition		-		-		-		
	c. Functions of Reproduction	110011		-				-	509
	DOOK IV								
	BOOK IV								
CTAR	I. Ages		-		-		-		510
CHAP.		320						-	510
	1. Infancy						-		510
	a. First period of Infancy	- n			-		-		
	b. Second period of Infancy or firs	t Denu	tion			*		2	515
	c. Third period of Infancy	-	-		-		-		520
	2. Childhood	-				-		-	520
	3. Adolescence	-	-		-0		-		524
	4 Visility or Manhood							-	527
	4. Virility or Manhood	100	40.0				14		529
	5. Old Age	-	-		-		-		
CHAP.	II. Sleep	-				*		-	534
	1. Dreams	-	-		-		-		539
	2. Waking Dreams			-		-		-	547
	3. Revery		-			- 2	-		554
Com	III. Correlation of Functions -								555
CHAP.	1. Mechanical Correlations		12		123		120		555
	1. Mechanical Correlations	100	8.		18			25	556
	2. Functional Correlations -			-				m.	561
	3. Sympathy	-	-		-				
-	a. Sympathy of Continuity	- 5		-		-		*	562
	b. Sympathy of Contiguity	-			-		-		562
	c. Remote Sympathies -	-		-		*		-	564
	d. Imagination	-			-		-		564
	e. Superstitions connected with	Symna	thy					-	566
	C. Ameta by which Sympothy	is accon	anlie	hod					568
	f. Agents by which Sympathy	alind	upire	mou		_			570
CHAP.	IV. Individual Differences amongst Man	nemu				-		The same	
	1. Temperaments -		-		-		-		570
	a. Sanguine Temperament -	-						-	571
	b. Bilious or Choleric Tempera	ment	-				•		572
	c. Melancholic or Atrabilious T	Cempera	amer	nt		-		-	573
	d. Phlegmatic, Lymphatic or P	ituitous	Te	mpe	ram	ent			573
	a. I megmane, hymphanic of a			-		-		-	573
	e. Nervous Temperament			200	-		1220		575
	2. Idiosyncrasy	-	120		70		100		
	3. Of Natural and Acquired Differ	ences		*		-		-	577
	1. Natural Differences	-	-		-		200		577
	a. Peculiarities of the Female	3 -				-		-	577
	2. Acquired Differences		-		-		-		581
	1. Habit			-		-		-	581
		_	-		-		-		585
	2. Association -	1000	7		7	-		20	586
	3. Imitation -	-	-	•					587
	4. Varieties of Mankind		-		-				
	1. Division of the Races			-		-		-	590
	a. Caucasian Race	-					-		591
	b. Ethiopian Race -	-		-				-	593
	c. Mongolian Race		-		-		-		594
	d. American Race -			-				-	595
	O Origin of the Different I	Pages	200	22.5	_	1	11/20		597
	2. Origin of the Different l	races							-

#### CONTENTS.

																PAGE
CHAP.	V.	Of	Life		-		-		-		-		-		-	605
	1000		Instinct		-	-		-		-		-		-		611
			Vital Pro	operties	-		-		-		-		-		-	619
		3.	Life of th	e Bloo	d	11/2		-		-		-		-		624
CHAP.	VI.			-	-		-		-		-		-		-	630
			Death fr	om Old	Age	-						-		-		630
			Acciden				-		-		-		-			633
			a. Death			the	He	art		-				-		633
			b. Death	beginn	ing in	the	Bra	in	-				-			634
			c. Death	beginn	ing in	the	Lu	ngs		12		-		-		634
INDEX		-	-	-	-		-		-							645

### LIST OF ILLUSTRATIONS IN VOL. II.

FIG.							PAGE
145.	The thorax		-		-	-	14
146.	Anterior view of the thorax, after Wil	son			-	-	14
147.	Anatomy of the heart and lungs, after	Wilson	-	-		-	16
148.	Reflections of the Pleura -	-	-		-	-	20
149.	Do. Do	100	- 1	-		-	21
150.	Section of the thorax and abdomen						27
151.	Thoracic and abdominal viscera		-		-		36
152.	Do. Do. of the	ostrich	-	-	-	-	65
153.	Heart of the dugong	15-07	-		+	-	66
154.	The right and left hearts separated	-	-	-	-	-	67
155.	Pulmonic heart	-	-	-	-	-	68
156.	Section of the pulmonic heart -	-	-	-	-		68
157.	Semicircular valves closed -		-	-	-	-	69
158.	View of the heart in sitû -	-	-0		-	-	72
159.	Circulation in the web of the frog's fo	ot, after	Wagner	- "	-	-	79
160.	Portion of the web of the frog's foot, a	fter Wa	gner	-	-		79
161.	Capillaries of the web of the frog's foo	ot, after 1	Wagner	-	-		81
162.	Bloodvessels of the lung of a live new	t, after V	Wagner			-	82
163.	Ramifications of the splenic artery in	the splee	n -	to all	-	1-	84
164.	Portal system, after Wilson -	-		-	-		88
165.	Corpuscles of human blood from a veir	n, beaten	so as to	separate	the fib	rin,	
	after Wagner	-	-	-	-	-	93
166.	Blood corpuscles of the edible frog, Ra	ana escul	lenta, aft	er Wag	ner		93
167.	Hydraulic apparatus, after Venturi	-	-	-	-		121
168.	Hæmadynamometer of Poisseuille	-	- 1/1	-101	-	-	130
169.	Section of a forcing pump -	-	2	-	-		133
170.	A small venous branch, from the web	of a frog	's foot, a	fter Wa	gner	-	140
171.	Large vein of frog's foot, after Wagne	r	-	-	-		141
172.	Diagram illustrating the circulation	-	-		-	-	142
173.	Diagram of vena contracta, from Vent	turi	-		-		149
174.	Do. adjutage, Do.		-		-	-	150
175.	Circulation in the frog	100	129		-	-	169
	Circulation in fishes		-		-		169
177.	Interior of the leech, after Sir E. Hom	e -			-	-	170

			PAGE
FIG.			176
	Globular structure of cellular tissue		177
179.			177
180.			178
181.		-	180
	Primary organic cell, after Todd and Bowman		185
	Tattooed head of a New Zealand chief -		222
184.	Secreting arteries, and nerves of the intestines -	man -	235
	Portion of areolar tissue inflated and dried, after Todd and Bow		236
	Fat vesicles, after Todd and Bowman Bloodvessels of fat vesicles, after Todd and Bowman -		236
	Fat vesicles from an emaciated subject, after Todd and Bowman	-	238
	Diapnogenous apparatus from the palm of the hand, after Wagn		245
	Sebaceous and ceruminous glands, after Wagner -		260
191	A small portion of the parotid of a new-born infant, filled with	mercury,	
101.	after Wagner		261
192.	Biliary and pancreatic ducts		264
	Abdominal and pelvic viscera	-	266
194.	Lobules of the liver, after Wagner	-	268
195.	Connection of lobules of liver with hepatic vein, after Wagner		268
	Transverse section of the lobules of the liver, after Kiernan	-	269
197.	Horizontal section of three superficial lobules, showing the two	principal	
	systems of bloodvessels, after Kiernan	-	269
198.	Horizontal section of two superficial lobules, showing interlobular	plexuses	
	of biliary ducts, after Kiernan		269
	First stage of hepatic venous congestion, after Kiernan	-	270
	Second Do, Do,	-	270
	Portal venous congestion, after Kiernan	-	270
	Under surface of liver, after Wilson		271
	Section of a kidney. — Supra-renal capsule	0 000	282
	Portion of kidneys of new-born infants, after Wagner -		283
	Small portion of kidney magnified 60 diameters, after Wagner	- 1	284
	A side view of viscera of male pelvis in sitû, after Wilson	-	286
	Entrance of the ureter into the bladder	-	292
	Section of the spleen		302
	Transverse section of testis, after Wilson	-	317
	Male organs, after Sir C. Bell	-	317
	Human testis injected with mercury, after Lauth		318
	Plan of the structure of the testis and epididymis, after Lauth		318
	Posterior view of male bladder, after Wilson	-	321
	Section of the vesiculæ seminales, &c.		321
	Section of bladder, prostate, and penis, after Wilson	-	322
	Section of the penis	- / /	324
	Spermatozoa, after Wagner -	-11	329
	Lateral view of pelvic viscera in the female	- 1 1	333
	Anterior view of the female organs		334
	Female organs of generation, after Wilson		335
	Section of the uterus		336
422.	Nerves of the uterus, after Dr. R. Lee		338

FIG.			PAGE
223.	Fallopian tube	all the	339
224.	Section of ovary		339
225.	Ripe ovum of rabbit, taken from the Graafian vesicle, after Wag	ner -	341
	Longitudinal division of the ovary		342
	Ovary of a female dying during menstruation -		347
	Tubal pregnancy		364
229.	Corpus luteum in the third month, after Montgomery -	7-0	377
230.	Corpus luteum at the end of the ninth month, after Montgomery		377
231.	Corpora lutea, after Sir E. Home	110 710	379
232.	Do. Do	-	380
233.	Villi of lining membrane of uterus, after Von Baer -	-	410
234.	Section of the uterus about eight days after impregnation, after	Wagner	412
235.	Do. when the ovum is entering its cavity,	Do.	412
236.		00	413
237.	Extra-uterine pregnancy	-	415
238.	Arteries of the impregnated uterus	- 10 10 -	417
	Cervix uteri at three months -		422
240.	Do. six months	-	422
241.	Do. eight months		422
242.	Do. nine months	-	423
243.	Natural labour	-	429
	Rotation of the head		430
	Breech presentation		431
	Arm presentation		432
	Milk ducts in human mamma, after Sir A. Cooper -		434
	Commencement of milk ducts, after Sir A. Cooper -		435
249.	Transverse section of the uterus and placenta, after Dr. J. Reid	. ;	449
250.	Connection between the maternal and fœtal vessels, after Dr. J.	Reid -	450
	Uterine surface of the placenta		451
	Fætal Do		451
	Knotted umbilical cord		454
254.	Diagram of the fœtus and membrane about the sixth week, after	Carus -	455
255.	Ovarium of the laying hen, after Sir E. Home -		459
256.	New laid egg with its molecule, after Sir E. Home -		460
257.	Section of a hen's egg within the ovary, after T. W. Jones		460
258.	Egg, thirty-six hours after incubation, after Sir E. Home		461
259.	Egg, opened three days after incubation, after Sir E. Home		463
	Egg five Do. Do		464
	Egg ten Do. Do		464
	Embryo of the egg, after Sir E. Home		465
263.	Embryo eighteen days old, after Sir E. Home -		465
264.	Ovum, fourteen days old, after Velpeau -		466
265.	Ovum and embryo, fifteen days old, after Maygrier -		467
266.	1 - 11 - Con Managing		467
	Fœtus at forty-five days		467
	Fœtus at two months		468
269	Corpora Wolffiana, with kidney and testes, from embryo of bird	s -	468
270.	Fœtus at three months, in its membranes -		469
-			

#### LIST OF ILLUSTRATIONS.

FIG.								PAGE
	Full period of utero-gestation			-		-		473
	Fœtus at full term -		-	-		-		475
	Section of the thymus gland at	the o	ighth mor	nth afte	er Sir A	. Coope	r -	476
	Circulatory organs of the fœtus				-		-	478
	Descent of the testicle -	s, alter	W HSUH		-			481
276.	Do							481
						-		486
	Decidual cotyledons, after Mon		ery			-		518
	Front view of the temporary to							518
	The separate temporary teeth				!	ear and	ina	0.0
280.	Permanent rudiment given off	from	the tempo	orary in	an inc	sor and	III a	501
	molaris, after T. Bell			-	-			521
281.	Temporary tooth and permane	nt rud	iment, af	ter T. E	Bell	-	-	521
282.	Temporary teeth and permaner	nt rud	iments, at	fter T.	Bell	-		522
283.	Do.			Do.	-		-	522
284.	Upper and lower teeth of the l	left sid	le of the j	aws, af	ter T. I	Bell		523
285.	Upper and lower teeth			-	-	-	-	524
286.	Skull of the aged, after Sir C.	Bell		-	-			530
287.	Physiognomy of the aged, afte	r Sir (	C. Bell	-	-	-		530
288.	Curves indicating the developm	nent o	f the heig	hts and	weigh	t of man	and	
	woman at different ages, a			-	-	-		532
289.	Caucasian variety -		-	-	-			591
290.	Negro variety -			-		-	-	593
	Mongolian variety -				-	-		594
	American variety -	-		-	-			595
	Curve indicating the viability of	or exis	tibility at	differen	nt ages	after On	etelet	637
	o		troiting ore	*********	as many	miner of c	A PATARA	001

# HUMAN PHYSIOLOGY.

# BOOK II.

### CHAPTER III.

RESPIRATION.

The consideration of the function of absorption has shown us how the different products of nutritive absorption reach the venous blood. By simple admixture with this fluid they do not become converted into a substance, capable of supplying the losses, sustained by the frame from the different excretions. Nothing is better established than the fact, that no being, and no part of any being, can continue its functions unless supplied with blood, which has become arterial, by exposure to air. It is in the lungs, that the absorbed matters undergo their final conversion into that fluid, by a function, which has been termed hæmatosis, and which is the great object of that we have now to investigate - Respiration. This conversion is occasioned by the venous blood of the pulmonary vessels coming in contact with the air in the air-cells of the lungs, during which contact, the blood gives to the air some of its constituents, and, in return, the air parts with its elements to the blood.

To comprehend this mysterious process, we must be acquainted with the pulmonary apparatus, as well as with the properties of atmospheric air, and the mode in which the contact between it and the blood is effected.


#### 1. ANATOMY OF THE RESPIRATORY ORGANS.

The thorax or chest contains the lungs, which are the great agents of respiration. It is of a conical shape, the apex of the cone being formed by the neck, and the base by a muscle, which has already


been referred to, more than once, - the diaphragm.

The osseous framework, Figs. 145 and 146, is formed, posteriorly, of twelve dorsal vertebræ: anteriorly, of the sternum, originally composed of eight or nine pieces; and laterally, of twelve ribs on each side, passing from the vertebræ to, or towards, the sternum. Of these, the seven uppermost extend the whole distance from the spine to the breast-bone, and are called the true or sternal ribs; sometimes, the vertebro-sternal. They become larger as

VOL. II. - 2



The Thorax. c. c. c. c. The ribs.



Anterior view of the Thorax. 1. Superior piece of sternum. 2. Middle piece. 3. Inferior piece, or ensiform cartilage. 4. First dorsal vertebra. 5. Last dorsal vertebra. 6. First rib. 7. Its head. 8. Its neck, resting against transverse process of first dorsal vertebra. 9. Its tuberosity. 10. Seventhor last true rib. 11. Costal cartilages of true ribs. 12. Two last false ribs—floating ribs. 13. The groove along lower border of rib for lodgment of intercepts and perve—(Wilson) costal vessels and nerve .- (Wilson.)

they descend, and are situate more obliquely in regard to the spine. The other five, called false or asternal, do not proceed as far as the sternum; but the cartilages of three of them join that of the seventh true rib, whilst the two lowest have no union with those above them, and are therefore called floating ribs. These false ribs become shorter and shorter as they descend; so that the seventh true rib may be regarded as the common base of two cones, formed by the true and false ribs respectively.

The different bones, constituting the thorax, are so articulated as to admit of motion, and thus to allow of dilatation and contraction of the cavity.

The motion of the vertebræ a. Sternum or breast-bone. b. b. The spine. on each other has been described under another head. It is not materially concerned in the respiratory movements. The articulation of the ribs with the spine and sternum demands attention. They are articulated with the spine in two places, at the capitulum or head, and at the tubercle. In the former of these, the extremity of the ribs, encrusted with cartilage, is received into a depression, similarly encrusted, at the side of the spine. One half of this depression is in the body of the upper vertebra; the other half in the one beneath it; and, consequently, partly in the intervertebral fibro-cartilage between the two. The joint is rendered secure by various ligaments; but it can move readily up and down on the spine. In the first, eleventh, and twelfth ribs, the articulations are with single vertebræ respectively. In the second articulation, the tubercle of the rib, also encrusted with cartilage, is received into a cavity in the transverse process of each corresponding vertebra; and the joint is rendered strong by three distinct ligaments. In the eleventh and twelfth ribs, this articulation is wanting. The articulation of the ribs with the sternum is effected by an intermediate cartilage, which becomes gradually longer, from the first to the tenth ribs, as seen in Figs. 145 and 146. The end of the cartilage is received into a cavity at the side of the sternum; and the junction is strengthened by an anterior and posterior ligament. This articulation does not admit of much motion; but the existence of a synovial membrane shows, that it is destined for some.

The cavity of the thorax is completed by muscles. In the intervals between the ribs are two planes of muscles, whose fibres pass in inverse directions, and cross each other. These are the intercostal muscles. The diaphragm forms the septum between the thorax and abdomen Above, the cavity is open; and through


the opening numerous vessels and nerves enter.

The muscles, concerned in the respiratory function, are numer-The most important of these is the diaphragm. It is attached, by its circumference, around the base of the chest; but its centre rises into the thorax; and, during its state of relaxation, forms an arch, the middle of which is opposite the inferior extremity of the sternum. It is tendinous in its centre, and is attached by two fasciculi, called pillars, to the spine, - to the bodies of the two first lumbar vertebræ. It has three apertures; one before for the passage of the vena cava inferior; and two behind, between the pillars, for the passage of the œsophagus and aorta. The other great muscles of respiration are the serratus posticus inferior, the serratus posticus superior, the levatores costarum, the intercostal muscles, the infra-costales and the triangularis sterni or sterno-costalis; but, in an excited condition of respiration, all the muscles, that raise and depress the ribs, directly or indirectly, participate - as the scaleni, sterno-mastoidei, pectoralis, (major and minor,) serratus major anticus, abdominal muscles, &c.

In the structure of the *lungs*, as Magendie<sup>a</sup> has remarked, nature has resolved a mechanical problem of extreme difficulty. The problem was, — to establish an immense surface of contact between the blood and the air, in the small space occupied by the lungs. The admirable arrangement adopted consists in this, — that each of the minute vessels, in which the pulmonary artery terminates, and the pulmonary veins originate, is surrounded on every side by the air. The lungs are two organs of considerable size, situate in the lateral parts of the chest, and subdivided into lobes and lobules, the shape and number of which cannot be readily determined. They are termed right and left, respectively, according to the side of the cavity of the chest which they occupy. The former consists of three lobes; the latter of two. Each of

these exactly fills the corresponding cavity of the pleura; and they

Fig. 147.



Anatomy of the Heart and Lungs.

1. Right ventricle; vessels to right of figure are middle coronary artery and veins; and those to its left, the anterior coronary artery and veins. 2. Left ventricles. 3. Right auricle. 4. Left auricle. 5. Pulmonary artery. 6, 6. Right pulmonary artery. 7. Left pulmonary artery. 8. Remains of ductus arteriosus. 9. Arch of aorta. 10. Superior vena cava. 11. Right vena innominata, and arteria innominata. 12. Right subclavian artery and vein. 13. Right common carotid artery and vein. 14. Left vena innominata. 15. Left carotid artery and vein. 16. Left subclavian artery and vein. 17. Trachea. 18. Right bronchus. 19. Left bronchus. 20, 20. Pulmonary veins; 18, 6, 20, form the root of right lung; and 7, 19, 20, root of left. 21. Superior lobe of right lung. 22. Its middle lobe. 23. Its in ferior lobe. -(Wilson.)

are separated from each other by a duplicature of the pleura-(the serous membrane that lines the chest, and is reflected over the lungs); - and by the heart. The colour of the lungs is generally of a marble blue; and the exterior is furrowed by figures of an hexagonal shape. The appearance is not, however, the same at all ages, and under all circumstances. In infancy, they are of a pale red; in youth, of a darker colour; and in old age, of a livid blue.

The elements that compose the lungs are; — the ramifications of the trachea; those of the pulmonary artery and of the pulmonary

veins, besides the organic elements, that appertain to every living structure, — arteries, veins, lymphatics, nerves and cellular tissue. The ramifications of the windpipe form the cavity of the organ of respiration. The trachea is continuous with the larynx, from which it receives the external air conveyed to it by the mouth and nose. It passes down to the thorax, at the anterior part of the neck, and bifurcates opposite the second dorsal vertebra, forming two large canals, called bronchi or bronchia. One of these goes to each lung; and, after numerous subdivisions, becomes imperceptible: hence, the multitudinous speculations that have been indulged regarding the mode in which the bronchial ramifications terminate. Malpighia believed that they form vesicles, at the inner surface of which the pulmonary artery ramifies. Reisseisen describes the

\* Epist. de Pulmon, i. p. 133.

b Ueber den Bau der Lungen, u. s. w., Berlin, 1822; also, in Latin, Berl. 1822.
See, likewise, Horner, Special Anatomy and Histology, 6th edit. ii. 161, Philad. 1843.

vesicles as of a cylindrical, and somewhat rounded figure; and he states, that they do not communicate with each other. Helvetius, on the other hand, affirmed, that they end in cells, formed by the different constituent elements of the lungs, — the cells having no determinate shape, or regular connexion with each other; whilst Magendieb asserts, that the minute bronchial division, which arrives at a lobe, does not enter it, but terminates suddenly as soon as it has reached the parenchyma; and, he remarks, that as the bronchus does not penetrate the spongy tissue of the lung, it is not probable that the surface of the cells, with which the air comes in contact, is lined by a prolongation of the mucous coat, which forms the inner membrane of the air-passages. Certain it is, that the most attentive examination has failed to detect its presence.

The ramifications of the pulmonary artery are another constituent element of the lung. This vessel arises from the right ventricle of the heart, and, at a short distance from that organ, divides into two branches; one passing to each lung. Each branch accompanies the corresponding bronchus in all its divisions; and, at length, becomes capillary and imperceptible. Its termination, also, has given rise to conjecture. Malpighi conceived it to end at the mucous surface of the bronchi, in an extremely delicate network, which he called rete mirabile. This was also the opinion of Reisseisen. According to others, the pulmonary artery, in its ultimate ramifications, is continuous with two kinds of vessels, - the capillary extremities of the pulmonary veins, and the exhalants engaged in the secretion of the pulmonary transpiration. Bichate admits, at the extremities of the pulmonary artery, and between that artery and the veins of the same name, vessels of a more delicate character, which he conceives to be the agents of hæmatosis. and which he calls the capillary system of the lungs. All that we know is, that the air gets a ready access to the blood in the pulmonary artery; but, with regard to the precise arrangement of the means of such access, we are ignorant. The same may be said of the third constituent of the lungs - the pulmonary veins. Their radicles manifestly communicate freely with those of the pulmonary artery; but they equally escape detection. When we observe them, they are found uniting, to constitute larger and larger veins, until they ultimately end in four large trunks, which open into the left auricle of the heart.

In addition to these organic constituents, the lung, like other organs, receives arteries, veins, lymphatics, and nerves. It is not nourished by the blood of the pulmonary artery, which is not adapted for that purpose, seeing that it is venous. The bronchial arteries are its nutritive vessels. They arise from the aorta, and are distributed to the bronchi.

Around the bronchi, and near where they dip into the tissue of the lung, lymphatic glands exist, the colour of which is almost

<sup>\*</sup> Mémoires de l'Académ. pour 1718, p. 18. Anatomie Descriptive, vol. iv., Paris, 1801.

black, and with which the few lymphatic vessels, that arise from the superficial and deep-seated parts of the lung, communicate. Haller has traced the efferent vessels of these glands into the thoracic duct.

The nerves, distributed to the lungs, proceed chiefly from the eighth pair or pneumogastric. A few filaments of the great sympathetic are also sent to them. The eighth pair - after having given off the superior laryngeal nerves, and some twigs to the heart - interlaces with numerous branches of the great sympathetic, and forms an extensive nervous network, called the anterior pulmonary plexus. After this, the nerve gives off the recurrents, and interlaces a second time with branches of the great sympathetic, forming another network, called the posterior pulmonary plexus. It then proceeds to the stomach, where it terminates. (See Fig. 108, vol. i., p. 418.) From these two plexuses the nerves proceed, that are distributed to the lungs. These accompany the bronchi, and are spread chiefly on the mucous membrane of the air-tubes. lung likewise receives some nerves directly from the three cervical ganglions of the great sympathetic, and from the first thoracic ganglion. In addition to these, a distinct system of nerves — the respiratory system, described in the first volume of this work is supposed by Sir Charles Bell to be distributed to the multitude of muscles, which are associated in the respiratory function, in a voluntary or involuntary manner. This system includes one of the nerves just referred to - the eighth pair - and the phrenic nerves, which are distributed to the diaphragm. The various nerves composing it are intimately connected, so that, in forced or hurried respiration, in coughing, sneezing, &c., they are always associated in action.b

Lastly, the lungs are constituted, also, of cellular tissue, which has been termed interlobular tissue; but it does not differ from

cellular tissue in other parts of the body.

Such are the constituent elements of the pulmonary tissue; but, with regard to the mode, in which they are combined to form the intimate texture of the lung, we are uninstructed. We find, that the lobes are divided into lobules, and these, again, seem to be subdivided almost indefinitely, forming an extremely delicate spongy tissue, the areolæ of which can only be seen by the aid of the microscope. It is generally thought, that the areolæ communicate with each other, and that they are enveloped by the cellular tissue which separates the lobules. Magendiec inflated a portion of lung, and dried and cut it in slices, in order that he might examine the deep-seated cells. These appeared to him to be irregular, and to be formed by the final ramifications of the pulmonary artery, and the primary ramifications of the pulmonary veins; the cells of one lobule communicating with each other, but not with those of an-

\* Elem. Physiologiæ, viii. 2.

b Müller's Handbuch, u. s. w., Baly's translation, p. 348, Lond. 1838.
c Précis, &c., ii. 309.

other lobule. Professor Horner, of the University of Pennsylvania, has attempted to exhibit that this communication between the cells is lateral. After filling the pulmonary arteries and the pulmonary veins with minute injection, the ramifications of the bronchi, with the air-cells, were distended to their natural size by an injection of melted tallow. The latter, being permitted to cool, the lung was cut into slices and dried. The slices were subsequently immersed in spirit of turpentine, and digested, at a moderate heat, for several days. By this process, all the tallow was removed, and the parts, on being dried, appeared to exhibit the air-cells empty, and, seemingly, of their natural size and shape. Preparations, thus made, appear to show the air-cells to be generally about the twelfth of a line in diameter, and of a spherical shape, the cells of each lobule communicating freely, like the cells of fine sponge, by lateral apertures. The lobules, however, only communicate by branches of the bronchi, and not by contiguous cells. This would seem to negative the presumption of some anatomists and physiologists, as Blumenbach, Cuvier, &c., - that each air-cell is insulated, communicating only with the minute bronchus, that opens into it; whilst it confirms the views of Haller, Monro (Secundus), Boyer, Sprengel, Magendie, and others; but it is impossible to decide positively, where all is so minute. Many anatomists, and perhaps with accuracy, by the term air-cell, mean simply the ultimate termination of a bronchus.b

The surface afforded by the air-cells is immense. Hales supposed them to be polyhedral, and about one-hundredth part of an inch in diameter. The surface of the bronchi he estimated at 1035 square inches; and that of the air-cells at 20,000. Keill estimated the number of cells to be 1,744,186,015; and the surface 21,906 square inches; and Lieberkühn has valued it at the enormous amount of 1500 square feet! All that we can derive from these mathematical conjectures is, that the extent of surface is surprising, when we consider the small size of the lungs themselves.

Recently, Professor Horner has published an account of various experiments, which exhibit the ready communication between the pulmonary air-vessels and the pulmonary veins. By fixing a pipe into the human trachea, and permitting a column of water to pass gently, he found that the air-cells became distended with water; and that the left side of the heart filled, and the aorta discharged water freely from its cut branches.

American Journal of the Medical Sciences, for Feb. 1832, p. 538, and Op. cit.

b See, on the Minute Anatomy of the Lungs, Wagner, Elements of Physiol., by R. Willis, Lond. 1842, and Bourgery, Gazette Méd. Juill. 16, 1842, and Brit. and For. Med. Rev. Oct. 1842, p. 546; and W. Addison, Proceedings of Royal Society, March 17, 1842, and Brit. and For. Med. Rev. Oct. 1842, p. 574.


Statical Essays, vi. p. 241. d Tentam. Med. Phys. p. 80.

<sup>·</sup> Blumenbach, in Elliotson's Physiology, p. 197, Lond. 1835.

Amer. Journ. of the Medical Sciences, April, 1843, p. 332; and Special Anatomy and Histology, 6th edit. ii. 163.

This experiment he repeated on human lungs on different occasions, and with like results. Very little water flowed from the pulmonary artery. In the sheep and the calf, however, when the experiment was practised upon them after they had been pretty thoroughly evacuated of blood, the water passed freely through both the pulmonary veins and the pulmonary arteries. Dr. Horner is disposed to infer, that his experiments exhibit a communication of the pulmonary air-vesicles by a direct route with the pulmonary bloodvessels, especially the veins; but this may well be questioned. It is possible, that such a communication may really have been made by the force of the column of water: and if not so, the passage of the fluid from the air-cells to the bloodvessels might have been effected through the pores, as in ordinary imbibition, which, we have elsewhere seen, is readily accomplished in the lungs, but not more readily perhaps than in the case of serous and other tissues under favourable circumstances. Hemorrhage by transudation occurs we know most rapidly at times through the coats of vessels; and a thinner fluid would of course transude more easily. It can scarcely be doubted, from Dr. Horner's experiments, that a certain arrangement exists between the air-vesicles and the pulmonary veins in man, which admits of a more ready imbibition and transudation; but what that arrangement is, admits of question.

Each lung is covered by the pleura, — a serous membrane analogous to the peritoneum, — and, in birds, a prolongation of the latter. This membrane is reflected from the adjacent surface of the lung to the pericardium which covers the heart, and is then spread over the interior paries of the half of the thorax to which it belongs; lining the ribs and intercostal muscles, and covering the convex or upper surface of the diaphragm. There are, consequently, two pleuræ, each of which is confined to its own half of




the thorax, lining its cavity and covering the lung. Behind the sternum, however, they are contiguous to each other, and form the partition, called mediastinum, which extends between the sternum and spine. In Fig. 148, the dotted lines exhibit the boundaries of the two cavities of the pleura, and the middle space between is the mediastinum. Within this septum, the heart, enveloped by the pericardium, is situate, and separates the pleuræ considerably from each other. Anatomists generally subdivide the mediastinum into two regions; one pass-

ing from the front of the pericardium to the sternum, called the anterior mediastinum; the other, from the posterior surface of the pericardium to the dorsal vertebræ, — the posterior mediasti-

num; and, by some, the part, which is within the circuit of the

first ribs, is termed superior mediastinum. The second of these contains the most important organs,—the lower end of the trachea, æsophagus, aorta, vena azygos, thoracic duct, and pneumogastric nerves. The portion of the pleura, covering each lung, is called pleura pulmonalis; that, which lines the thorax, pleura costalis. The mode, in which the two are connected to form one whole, is shown by the dotted line in Fig. 149, representing a vertical section of the chest. It is obvious, that, as in the case of the abdomen, the viscera are not in the



Reflections of the Pleura.

cavity of the pleura, but external to it; and that there is no communication between the serous sac of one side and that of the other.

The use of the pleura is to attach the lungs, by their roots, to their respective cavities, and to facilitate their movements. To aid this effect, the membrane is always lubricated by a fluid, exhaled from its surface. The other surface is attached to the lung in such a manner, that air cannot get between it and the parietes of the thorax.

Dr. Stokes, of Dublin, has described a proper fibrous tunic of the lungs. In a healthy state, this capsule, although possessing great strength, is transparent, a circumstance in which it differs from the fibrous capsules of the pericardium, and which, Dr. Stokes thinks, has probably led to its having been overlooked. It invests the whole of both lungs; covers a portion of the great vessels; and the pericardium seems to be but its continuation, — endowed, in that particular situation, with a greater degree of strength, for purposes that are obvious. It covers the diaphragm where it is more opaque; in connexion with the pleura, it lines the ribs; and, turning, forms the mediastina, which are thus shown to consist of four layers, — two serous and two fibrous. It seems, that Dr. Hart, of Dublin, has, for years, demonstrated this tunic to his class.

It was, at one time, the prevalent belief, that air always exists in the cavity of the chest. Galen supported the opinion by the fact, that, having applied a bladder, filled with air, to a wound, which had penetrated the chest, the air was drawn out of the bladder at the time of inspiration. This was also maintained by Hamberger, Hales, and numerous others. The case, alluded to by Galen, is insufficient to establish the position, inasmuch as we have no evidence, that the wound did not also implicate the pulmonary tissue. Since the time of Haller, who opposed the prevalent doctrine by observation and reasoning, the fact of the ab-

On Diseases of the Chest, Part i. p. 460, Dublin, 1837; or Dunglison's American
 Medical Library edition, p. 301, Philad. 1837.
 Statical Essays, ii. 81.

sence of air in the cavity of the pleura is generally considered to be entirely established. It is obvious, that its presence there would materially interfere with the dilatation of the lungs, and thus be productive of much inconvenience; besides, anatomy instructs us, that the lungs lie in pretty close contact with the pleura When the intercostal muscles are dissected off, and the pleura costalis is exposed, the surface of the lungs is seen in contact with that transparent membrane; and, when the pleura is punctured, the air rushes in, and the lungs retire, in proportion as the air is admitted. This occurs in cases of injuries inflicted upon the chest of the living animal. Moreover, if a dead or living body be placed under water, and the pleura be punctured, so as not to implicate the lungs, it has been found by the experiments of Brunn, Sprögel, Caldani, Sir John Floyer, Haller, and others, that not a bubble of air escapes, - which would necessarily be the case, if air were contained in the cavity of the pleura.b

### 2. ATMOSPHERIC AIR.

The globe is surrounded every where, to the height of fifteen or sixteen leagues, by a rare and transparent fluid called air; the

total mass of which constitutes the atmosphere.

Atmospheric air, although invisible, can be proved to possess the ordinary properties of matter; and, amongst these, weight. It also partakes of the character of a fluid, adapting itself to the form of the vessel in which it is contained, and pressing equally in all directions.

As air is possessed of weight, it results, that every body on the earth's surface must be subjected to its pressure; and as it is elastic or capable of yielding to pressure, the part of the atmosphere near the earth's surface must be denser than that above it. As a body, therefore, ascends, the pressure will be diminished; and this accounts for the different feelings experienced by those who ascend lofty mountains, or voyage in balloons, into the higher strata of the atmosphere. Dr. Edwards ascribes part, at least, of the effect produced upon the breathing, at great elevations, to the increased evaporation which takes place from the skin and lungs; and in many aërial voyages great inconvenience has certainly been sustained from this cause.

The pressure of the atmosphere at the level of the sea is the result of the whole weight of the atmosphere, and is capable of sustaining a column of water thirty-four feet high, or one of mercury of the height of thirty inches, — as in the common barometer. This is equal to about fifteen pounds avoirdupois on every square inch of surface; so that the body of a man of ordinary stature, the

\* Element. Physiol. viii. 2

b Bostock's Physiology, 3d edit. p. 305, Lond. 1836; and Adelon, Physiologie de l'Homme, edit. cit. iii. 144.

See the author's Elements of Hygiène, p. 39, Philad. 1835; and art. Atmosphere,
 by the author, in Amer. Cyclop. of Pract. Med. iii. 529, Philad. 1836.
 De l'Influence des Agens Physiques, &c. p. 493, Paris, 1824.

surface of which Haller estimates to be fifteen square feet, sustains a pressure of 32,400 pounds. Yet, as the elasticity of the air within the body exactly balances or counteracts the pressure from without, he is not sensible of it.

The experiments of Davy, Dalton, Gay Lussac, Humboldt, Despretz, and others, have shown, that pure atmospheric air is composed essentially of two gases, oxygen and azote, which exist in it in the proportion of 21 of the former to 79 of the latter: Dr. T. Thomson, whose analysis is one of the most recent and satisfactory, says 20 of oxygen to 80 of azote or nitrogen; and these proportions have been found to prevail in the air whencesoever taken; - whether from the summit of Mont Blanc, the top of Chimborazo, the sandy plains of Egypt, or from an altitude of 23,000 feet in the air.ª Chemical analysis has not been able to detect the presence of any emanation from the soil of the most insalubrious regions, or from the bodies of those labouring under the most contagious diseases, malignant and material as such emanations unquestionably must This uniformity in the proportion of the oxygen to the nitrogen in the atmosphere has led to the conclusion, that as there are many processes, which consume the oxygen, there must be some natural agency, by which a quantity of oxygen is produced equal to that consumed. The only source, however, by which oxygen is known to be supplied, is the process of vegetation. A healthy plant absorbs carbonic acid during the day; appropriates the carbon to its own necessities, and gives off the oxygen with which it was combined. During the night an opposite effect is produced. The oxygen is then taken from the air, and carbonic acid given off; but the experiments of Davy and Priestley show, that plants, during the twenty-four hours, yield more oxygen than they consume. It is impossible to look to this as the great cause of equilibrium between the oxygen and azote. Its influence can extend to a small distance only; and yet the uniformity has been found to prevail, as we have seen, in the most elevated regions, and in countries whose arid sands never admit of vegetation.

In addition to the oxygen and azote, — the principal constituents of atmospheric air, — another gas exists in very small proportion, but is always present. This is carbonic acid. It was found by De Saussure on Mont Blanc, and by Humboldt in air brought down, by Garnerin, the aëronaut, from the height of several thousand feet. The proportion is estimated by Dalton not to exceed the  $\frac{1}{1000}$ th or  $\frac{1}{1400}$ th of its bulk. In one of the wards of La Pitié, in Paris, which had been kept shut during the night, M. Felix Leblanc<sup>b</sup> found a larger portion of carbonic acid, nearly  $\frac{3}{1000}$ ths; and in one of the dormitories of La Salpetrière, the air yielded  $\frac{8}{1000}$ ths; the largest proportion found by him in the air of hospitals. In the lecture room of the Sorbonne, which is capable of

<sup>\*</sup> Art. Atmosphere, (Physical and Chemical History,) by Dr. R. M. Patterson, in Amer. Cyclopedia of Practical Medicine and Surgery, vol. ii. p. 526, Philad. 1836.
\* Gazette Med. de Paris, 11 Juin, 1842.

containing 1000 cubic inches of air, after a lecture an hour and a half long, and at which 900 persons were present, the oxygen was found to have lost 1 in every 100, although two doors were open; whilst the carbonic acid was increased, in rather a greater ratio. In a ward in an institution for children, although the door was half open, and there was an open space in the roof, the air was found to contain 3 ths of carbonic acid, and there was a proportional

diminution of oxygen.

These, then, may be looked upon as the constituents of atmospheric air. There are certain substances, however, which are adventitiously present in variable proportions; and which, with the constitution of the atmosphere as to density and temperature, are the causes of general or local salubrity, or the contrary. Water is one of these. The quantity, according to De Saussure, in a cubic foot of air, charged with moisture, at 65° Fahr., is 11 grains. amount in the atmosphere is very variable, owing to the continual change of temperature to which the air is subject; and even when the temperature is the same, the quantity of vapour is found to vary, as the air is very rarely in a state of saturation. The varying condition as to moisture is indicated by the hygrometer. From a comparison of numerous observations, Gay Lussac affirms, that the mean hygrometric state of the atmosphere is such, that the air holds just one-half the moisture necessary for its saturation. In his celebrated aërial voyage, he found the air to contain but oneeighth of the moisture necessary for saturation. This is the greatest degree of dryness ever noticed.

It has been presumed, that the hygrometric condition of the atmospheric air has more agency in the production of disease than either the barometric or thermometric. It is not easy to say which exerts the greatest influence: probably all are concerned, and when we have a union of particular barometric, thermometric, hygrometric, electric, and other conditions, we have certain epidemics existing, which do not prevail under any other combination. When the air is dry, we feel a degree of elasticity and buoyancy; whilst, if it be saturated with moisture, - especially during the heat of summer, - languor and lassitude, and indisposition to mental or cor-

poreal exertion are excited.

In addition to aqueous vapour, numerous emanations from animal and vegetable substances must be generally present, especially in the lower strata of the atmosphere; by which the salubrity of the air may be more or less affected. All living bodies, when crowded together, deteriorate the air so much as to render it unfit for the maintenance of the healthy function. If animals be kept crowded together in ill-ventilated apartments, they speedily sicken. The horse becomes attacked with glanders; fowls with pep, and sheep with a disease peculiar to them if they be too closely folded. is probably a principal cause of the insalubrity of cities compared with the country. In them, the air must necessarily be deteriorated by the impracticability of due ventilation, and this, with the want of due exercise, is a fruitful cause of cachexia — and of tuberculous cachexia; hence, also, it is, that in work-houses and manufactories, diseases dependent on this condition of constitution are prevalent. One of the greatest evidences we possess of the positive insalubrity of towns is the case of the young. In London, the proportion of those that die annually under five years of age to the whole number of deaths is as much as thirty-eight per cent., and under two years, twenty-eight per cent.; in Paris, under two years of age, twenty-five per cent.; and in Philadelphia and Baltimore, rather less than a third. These estimates may be considered approximations; the proportions varying somewhat, according to the precise year in which they have been taken. Manifest, however, as is the existence of some deleterious principle, in these cases, it has always escaped the researches of the chemist.

Lastly. Air is indispensable to organic existence. No being, animal or vegetable, - can continue to live without a due supply of it; nor can any other gas be substituted for it. This is proved by the fact, that all organized bodies cease to exist, if placed in vacuo. They require, likewise, renovation of the air, otherwise they die; and if the residual air be examined, it is found to be diminished in quantity, and to have received a gas, which is totally unfit for life, — carbonic acid. The experiments of Hales prove this as regards vegetables; whilst Spallanzani and Vauquelin have confirmed it in the case of the lower animals. The necessity for the presence of air, and its due renewal, - as regards man and the upper classes of animals, - is sufficiently obvious. Not less necessary is a due supply of air to aquatic animals. They can be readily drowned, when the air in the water is consumed, if prevented from coming to the surface: if the fluid be put under the receiver of an air-pump, and the air be withdrawn, or if the vessel be placed so that the air cannot be renewed, the same changes are found to have been produced in the air; and hence the necessity for making holes through the ice, where small fish-ponds are frozen over, if we be desirous of preserving the fish alive. The necessity for the renewal of air is not, however, alike imperative in all animals. Whilst the mammalia, birds, fishes, &c., speedily expire, when placed under the receiver of an air-pump, if the receiver be exhausted; the frog is but slightly incommoded. It swells up almost to bursting, but retains its position, and when the air is admitted, seems to have sustained no injury. This exception, afforded by the amphibious animal to the ordinary effects of destructive agents, we have already had occasion to refer to more than once; and it is strikingly exemplified in the fact, now indisputable, that the toad has been found alive in the substance of trees and rocks, where no access of air appeared practicable.

The influence of air on mankind is most interesting and important in its hygienic relations, and has accordingly been a topic of

VOL. 11. - 3

study since the days of Hippocrates. In other works, it has been investigated, at considerable length, by the author.a

### 3. PHYSIOLOGY OF RESPIRATION.

a. Mechanical Phenomena of Respiration. - Within certain limits, the function of respiration is under the influence of volition. The muscles, belonging to it, have consequently been termed mixed, as we can at pleasure increase or diminish their action, but cannot arrest it altogether, or for any great length of time. If, by a forced inspiration, we take air into the chest in large quantity, we find it impossible to keep the chest in this condition beyond a certain period. Expiration irresistibly succeeds, and the chest resumes its pristine situation. The same occurs if we expel the air as much as possible from the lungs. The expiratory effort cannot be prolonged indefinitely, and the chest expands in spite of the effort of the will. The most expert divers do not appear capable of suspending the respiratory movements longer than 95 or 100 seconds. Dr. Lefèvre found the average period of the Turkish divers to be 76 seconds for each man. These facts have given rise to two curious and deeply interesting topics of inquiry; - the cause of the first inspiration in the new-born infant? and of the regular alternation of inspiration and expiration during the remainder of existence? The first of these questions will fall under consideration when we investigate the physiology of infancy; the latter will claim some attention at present. Haller attempted to account for the phenomenon by the passage of the blood through the lungs being impeded during expiration, - a reflux of blood into the veins, and a degree of pressure upon the brain being thus induced. Hence, a painful sense of suffocation arises, in consequence of which the muscles of inspiration are called into action by the will, for the purpose of enlarging the chest, and, in this way, removing the impediment. The same uneasy feelings, however, ensue from inspiration, if too long protracted: the muscles cease to act, and, by their relaxation, the opposite state of the chest is induced. Whytte conceived, that the passage of the blood through the pulmonary vessels is impeded by expiration, and that a sense of anxiety is thus produced. The unpleasant sensation acts as a stimulus upon the nerves of the lungs and the parts connected with them, which arouses the energy of the sentient principle; and this, by acting in a reflex manner, causes contraction of the diaphragm, enlarges the chest, and removes the painful feeling. The muscles then cease to act, in consequence of the stimulus no

Elements of Hygiène, pp. 33 to 305, Philad. 1835; and American Cyclopedia of Practical Medicine and Surgery, art. Atmosphere, p. 527, Philad. 1836.


b Loudon's Magazine of Nat. Hist. p. 617, Dec. 1836; and Dunglison's Amer. Med. Intelligencer, p. 30, April 15, 1837. Elementa Physiologiæ, viii. 4, 17.

d An Essay on the Vital and other Involuntary Motions of Animals, sect. viii., Edinb.

longer existing. These, and all other methods of accounting for the phenomena, are, however, too pathological. From the first moment of respiration the process appears to be accomplished without the slightest difficulty, and to be as much a part of the instinctive extra-uterine actions of the frame, as circulation, digestion, or absorption. It is obviously an internal sensation, after respiration has been once established; and, like all internal sensation.

sations, is inexplicable in our existing state of knowledge. The part which developes the impression is probably the lung, through its ganglionic nerves; the pneumogastric nerves convey the impression to the brain or spinal marrow, which calls into action the muscles of inspiration. We say, that the action of impression arises in the lungs, and this, from some internal cause, connected with the office to be filled in the economy: but in so saying we sufficiently exhibit our total want of acquaintance with its nature.

The movements of inspiration and expiration, which, together, constitute the function of respiration, are entirely accomplished by the dilatation and contraction of the thorax. The air enters the chest when the latter is expanded; and it is driven out when the chest is restored to its ordinary dimensions; — the thorax thus seeming to act like an ordinary pair of bellows with the valve stopped:



Section of the Thorax and Abdomen.

a. Thorax. b. Abdomen. c. Relaxed diaphragm.

when the sides are separated, the air enters at the nozzle, and it is forced out when they are brought together.

#### 1. INSPIRATION.

The augmentation of the capacity of the thorax, which constitutes inspiration, may be effected to a greater or less extent, according to the number of muscles which are thrown into action. The chest may, for example, be dilated by the diaphragm alone. This muscle, as we have seen, in its ordinary relaxed condition is convex towards the chest, as in Figures 150 and 151. When, however, it contracts, it becomes more horizontal; and assumes the position indicated by the dotted line d, Fig. 150, in this manner augmenting the cavity of the chest in a vertical direction. The sides or lateral portions of the diaphragm, which are fleshy and correspond to the

See, also, Dr. M. Hall's Lectures on the Nervous System, p. 55, Lond. 1836; or Amer. Edit, p. 36, Philad. 1836.

lungs, descend more, in this movement, than the central, tendinous portion, which is moreover kept immovable by its attachment to the sternum, and its union with the pericardium. In the gentlest of all breathing, the diaphragm appears to be the sole agent of inspiration; and in cases of inflammation of the pleura costalis, or of fractured rib, our endeavours are directed to the prevention of any elevation of the ribs by which the diseased part can be put upon the stretch. Generally, however, as the diaphragm descends, the viscera of the abdomen are compressed; the abdominal muscles assume the position of the double dotted line f, and the ribs and the breast bone are raised so that the latter is protruded as far as the dotted line e. When the diaphragm acts, and, in addition, the ribs and sternum are raised, the cavity of the chest is still farther augmented.

The mechanism, by which the ribs are raised, has been productive of more controversy than the subject merits. Hallera asserted, that the first rib is immovable, or at least admits of but trifling motion when compared with the others; and he denies that the thorax makes any movement, as a whole, of either elevation or depression; affirming that the ribs are raised successively towards the top of the cavity; and this to a greater extent as they are more distant from the first. Magendie, on the other hand, denies that they are elevated in this manner; and endeavours to show that they are all raised at the same time; that the first rib, instead of being the least movable, is the most so; and that the disadvantage, which the lower ribs possess in the movement, by their admitting of less motion in their posterior articulations, is compensated by the greater length of these ribs. This compensation he considers to have its advantages; for as the true ribs, with their cartilages and the sternum, usually move together, and the motion of one of these parts almost always induces that of the rest, it would follow, that if the lower ribs were more movable, they could not execute a more extensive movement than they do; whilst the solidity of the thorax would be diminished. By the elevation, then, of the ribs, and the depression of the diaphragm, the chest is augmented, and a deeper inspiration effected than when the diaphragm acts singly. In this elevation of the ribs, we see the advantage of their obliquity as regards the spine. Had they been horizontal, or inclined obliquely upwards, any elevation would necessarily have contracted the thoracic cavity, and favoured expiration instead of inspiration.

The muscles chiefly concerned in inspiration are the intercostals, and those muscles which arise, either directly or indirectly, from the spine, head, or upper extremities, and which can, in any manner, elevate the thorax. Amongst these, are the scaleni antici and postici, the levatores costarum, the muscles of the neck, which

are attached to the sternum, &c.

As no air exists in the cavity of the pleura, it necessarily happens, that, when the capacity of the chest is augmented, the residuary air, contained in the air-cells of the lungs after expiraton, is rarefied;

<sup>&</sup>lt;sup>a</sup> Elementa Physiologiæ, viii.

and, in consequence, the denser air without enters the larynx by the mouth and nose, until the air within the lungs has attained the density, which the residuary air had, prior to inspiration, - not that of the external air, as has been affirmed.a At the time of inspiration, the glottis opens by the relaxation of the arytenoidei muscles, as Legallois proved by experiments, performed at the Ecole de Médecine of Paris. On exposing the glottis of a living animal, the aperture is found to dilate very distinctly at each inspiration, and to contract at each expiration. If the eighth pair of nerves be divided low down in the neck, and the dilator muscles of the glottis, which receive their nerves from the recurrents - branches of the eighth pair — be thus paralysed, the aperture is no longer enlarged during inspiration, whilst the constrictors — the arytenoidei muscles which receive their nerves from the superior laryngeal, - given off above the point of section - preserve their action, and close the glottis more or less completely.

When the air is inspired through the mouth, the velum is raised, so as to allow the air to pass freely to the glottis; and, in forced inspiration, it is so horizontal, as to completely expose the pharynx to view. The physician takes advantage of this, in examining morbid affections of those parts, and can often succeed much better in this way than by pressing down the tongue. On the other hand, when inspiration is effected entirely through the nose, the velum palati is depressed, until it becomes vertical, and no obstacles exist to the free entrance of the air into the larynx. In such case, where difficulty of breathing exists, the small muscles of the alæ nasi are frequently thrown into violent action, alternately dilating and contracting the apertures of the nostrils; and hence this

is a common symptom in pulmonary affections.

Mayow conceived, that the air enters the lungs in inspiration as it would a bladder put into a pair of bellows, and communicating with the external air by the pipe of the instrument. The lungs, however, are not probably so passive as this view would indicate. In cases of hernia of the lungs, the extruded portion has been observed to dilate and contract in inspiration and expiration. Reisseisen believed this to be owing to muscular fibres, which Meckel and himself conceived to perform the whole circuit of the bronchial ramifications. These are not, however, generally admitted by anatomists, and the phenomenon is usually ascribed to the bronchi having in their composition the highly elastic tissue, which is an important constituent of the arteries. Laennecd affirms, that he has endeavoured, without success, to verify the observations of Reisseisen; but that the manifest existence of

Animal Physiology, Library of Useful Knowledge, p. 100, Lond. 1829.

b Œuvres, p. 177, Paris, 1824. ° Tractatus Quinque, p. 271, Oxon. 1674. d On the Diseases of the Chest, &c., 4th edit., Lond. 1834: reprinted in this country, Philad. 1835. See, also, Dr. Watson, Lectures in Lond. Med. Gaz. Feb. 11, 1842; and the author's Practice of Medicine, 2d edit. vol. ii. art Asthma, Philad. 1844.

circular fibres in branches of a moderate size, and the phenomena, presented by many kinds of asthma, induce him to consider the temporary constriction and occlusion of the minute bronchial ramifications as a thing well established. In the trachea, an obvious muscular structure exists in the posterior third, where the cartilages are wanting. There it consists of a thin muscular plane, the fibres of which pass transversely between the interrupted extremities of the cartilaginous rings of the trachea and bronchi, to which a layer of longitudinal fibres may at times be seen superadded. The use of the transverse muscular tissue, as suggested by Dr. Physick, and, since him, by Cruveilhier and Sir Charles Bell, is to diminish the calibre of the air-tubes in expectoration; so that the air having to pass through the contracted portion with greater velocity, its momentum may remove the secretions that are adherent to the mucous membrane. The explanation is in-

genious and probably just.

Magendie asserts, that the lung has a constant tendency to return upon itself, and to occupy a smaller space than that which it fills; and, that it consequently exerts a degree of traction on every part of the parietes of the thorax. This traction has but little effect upon the ribs, which cannot yield: but upon the diaphragm it is considerable. It is, indeed, in his opinion, the cause, why that muscle is always tense, and drawn so as to be vaulted upwards; and when the muscle is depressed during contraction, it is compelled to draw down the lungs towards the base of the chest, so that they are stretched, and, by virtue of their elasticity, have a more powerful tendency to return upon themselves, and to draw the diaphragm upwards. If a puncture be made into the chest in one of the intercostal spaces, the air will enter the chest through the aperture, and the lung will shrink. By this experiment, the atmospheric pressure is equalized on both surfaces of the lung, and the organ assumes a bulk determined by its elasticity and weight. Owing to this resiliency of the lungs, and to their consequent tendency to recede from the pleura costalis, there is less pressure upon all the parts against which the lungs are applied; and, accordingly, the heart is not exposed to the same degree of pressure as the parts external to the chest; and the degree of pressure is still farther reduced, when the chest is fully dilated, the lungs farther expanded, and their elastic resiliency increased.

Many physiologists have pointed out three degrees of inspira-

b Dr. Goddard, in Wilson's Anatomist's Vade-mecum, Amer. edit., p. 484, note. Philad. 1843.

Ge Horner's Lessons in Pract. Anat., p. 179, Philad. 1836; and Special Anat. and Histology, 4th edit., Philad. 1843.

Georgia Philos. Transact. for 1832, p. 301.

J. Carson, on the Elasticity, of the Lungs, in Philos. Transactions for 1820; and Inquiry into the Causes of Respiration, &c. 2d edit., Lond. 1833.

a See, on this subject, Trousseau and Belloc on Laryngeal Phthisis, translated by Dr. Warder, of Cincinnati, for Dunglison's Amer. Med. Library, p. 81, Philad. 1839; and Carpenter, Human Physiology, § 527, Lond. 1842.

tion, but it is manifest that there may be innumerable shades between them:—1. Ordinary gentle inspiration, which is owing simply to the action of the diaphragm; or, in addition, to a slight elevation of the chest. 2. Deep inspiration, where, with the depression or contraction of the diaphragm, there is evident elevation of the thorax, and, lastly, forced inspiration, when the air is strongly drawn in, by the rapid dilatation, produced by the action of all the respiratory muscles that elevate the chest directly or in-

directly.

Trials have been instituted for determining the quantity of air taken into the lungs at an inspiration; and considerable diversity, as might be expected, exists in the evaluations of different experimenters. We have just remarked, that, in the same individual, the inspiration may be gentle, deep, or forced; and, in each case, the quantity of air inspired will necessarily differ. There is, likewise, considerable diversity in individuals, as regards the size of the chest; so that an approximation can alone be attained. The following table sufficiently exhibits the discordance of authors on this point. Many, however, of the estimates, which seem so extremely discrepant, may probably be referred to imperfection in the mode of conducting the experiment, as well as to the causes above mentioned:

	Cubic inches at each Inspiration.	transmitted at and	Cubic inches at each Inspiration.
Reil,	42 to 100	Fontana,	35
Menzies, Sauvages,	The second second	Richerand, Dalton,	30 to 40 30
Hales, Haller,	maintent di	Herholdt, Jurine and Coathupe,	20 to 29 20
Ellis,	40	Allen and Pepys,	161
Sprengel,	desting output	J. Borelli,	15 to 40
Sömmering,	Data mineral	Goodwin,	14
Thomson,	Transfer and	Sir H. Davy,	13 to 17
Bostock,	And services onto	Abernethy and Mojon,	12
Jurin,	35 to 38	Keutsch,	6 to 12b

In passing through the mouth, nasal fossæ, pharynx, larynx, trachea, and bronchi, the inspired air acquires pretty nearly the temperature of the body; and, if the air be cool, the same quantity by weight occupies a much larger space in the lungs, owing to its rarefaction in those organs. In its passage, too, it becomes mixed with the halitus, which is constantly exhaled from the mucous membrane of the air-passages; and, in this condition, it enters the air-cells, and becomes mixed, by diffusion, with the residuary air in the lungs after expiration.

a Dr. Marshall Hall has devised a pneumatometer for this purpose. See art. Irri-

tability, in Cyclop. of Anat. and Physiol. July, 1840.

b Bostock's Essay on Respiration, Liverpool, 1804; and Elementary, System of Physiology, 3d edit. p. 314, Lond. 1836; and C. T. Coathupe, in Philos. Magaz. June, 1839.

It is obvious, that if we knew the exact capacity of the lungs in an individual when in health, we might be able to determine the extent of solidification in pulmonary affections by the diminution in their capacity. Owing, however, to our want of this requisite preliminary knowledge, the test is not of much avail.

### 2. EXPIRATION.

An interval, scarcely appreciable, elapses after the accomplishment of inspiration, before the reverse movement of expiration succeeds; and the air is expelled from the chest. The great cause of this expulsion is the restoration of the chest to its former dimensions; and the elasticity of the yellow tissue composing the bronchial ramifications, which have been put upon the stretch by the air rushing into them during inspiration. The restoration of the chest to its dimensions may be effected simply by the cessation of the contraction of the muscles, that have raised it, and the elasticity of the cartilages, which connect the bony portions of the ribs with the sternum or breast-bone. In active expiration, however, the ribs are depressed by the action of appropriate muscles, and the chest is thus still farther contracted. The chief expiratory muscles are the triangularis sterni, the broad muscles of the abdomen, rectus abdominis, sacro-lumbalis, longissimus dorsi, serratus posticus inferior, &c. Hallera conceived that the ribs, in expiration, are successively depressed towards the last rib; which is first fixed by the abdominal muscles and quadratus lumborum. The intercostal muscles then act and draw the ribs successively downwards. Magendieb contests the explanation of Haller; and the truth would seem to be, that the muscles, just mentioned, participate with the intercostals in every expiratory movement. By this action, the capacity of the chest is diminished; the lungs are correspondently pressed upon, and the air issues by the glottis. It has been already remarked, that, during expiration, the arytenoidei muscles contract, and the glottis appears to close. Still, space sufficient is left to permit the exit of the air.

It has been asked - is the air expired precisely that which has been taken in by the previous inspiration? It is impossible to empty the lungs wholly by the most forced expiration. A portion still remains; and hence it has been assumed, that the use of inspiration is to constantly renew the air remaining in the air-cells. On this subject we are not well informed; but it is probable that the lighter and more rarefied air mixes, by diffusion, with the newlyarrived and denser medium. Many experiments have been made to determine the change of bulk which air experiences by being respired. According to Sir Humphry Davy,e it is diminished, by a single inspiration, from \frac{1}{70}th to \frac{1}{100}th part of its bulk. Cuvier makes it about 1 th; Allen and Pepys a little more than a half per cent. Berthollet from 0.69 to 3.70 per cent.; and Bostock

c Researches, Chemical and Philosophical, p. 431, Lond. 1800.

<sup>&</sup>lt;sup>a</sup> Element. Physiol. viii. 4. b Précis, &c. ii. 324.

th,—as the average diminution. Assuming this last estimate to be correct; and forty cubic inches to be the quantity of air drawn into the lungs at each inspiration, it will follow, that half a cubic inch disappears each time we respire. This, in a day, would amount to 14,400 cubic inches, or to rather more than eight cubic feet. The experiments of MM. Dulong and Despretz make the diminution considerable. The latter gentleman placed six small rabbits in forty-nine quarts of air for two hours, at the expiration of which time the air had diminished one quart. A portion of the

inspired air must consequently have been absorbed.

Attempts have been made to estimate the quantity of air remaining in the lungs after respiration; but the sources of discrepancy are here as numerous as in the cases of inspiration or expiration. Goodwyna estimated it at 109 cubic inches; Menziesb at 179; Jurine at 220; Fontanad at 40; and Cuvier, after a forced inspiration, at from 100 to 60. Davye concluded, that his lungs, after a forced expiration, still retained 41 cubic inches of air; after a natural expiration they contained 118 cubic inches; after a natural inspiration, 135; after a forced inspiration, 254; by a full forced expiration after a forced inspiration, he threw out 190 cubic inches; after a natural inspiration, 78.5; after a natural expiration, 67.5. It is impossible, from such variable data, to deduce any thing like a satisfactory conclusion; but if we assume with Bostock, (and Dr. Thomsonf is disposed to adopt the estimate,) 170 cubic inches as the quantity, that may be forcibly expelled, and that 120 cubic inches will be still left in the lungs, we shall have 290 cubic inches as the measure of the lungs in their natural or quiescent state; to this quantity, 40 cubic inches are added by each ordinary inspiration, giving 330 cubic inches as the measure of the lungs in their distended state. Hence it would seem, that about one-eighth of the whole contents of the lungs is changed by each respiration; and that rather more than two-thirds can be expelled by a forcible expiration. Supposing, that each act of respiration occupies three seconds, or that we respire twenty times in a minute, a quantity of air, rather more than 23 times the whole contents of the lungs, will be expelled in a minute, or about four thousand times their bulk in twenty-four hours. The quantity of air, respired during this period, will be 1,152,000 cubic inches, about 6662 cubic feet. Such is Bostock's estimate. It is the residuary air, that gives to the lungs the property of floating on the surface of water, after they have once received the breath of life, and no pressure, that can be employed, will force out the air, so as to make them sink. Hence, the chief proofs, whether a child has been born alive or dead, are deduced from the lungs. These proofs constitute the docimasia

<sup>&</sup>lt;sup>a</sup> Op. citat. p. 36. <sup>b</sup> Op. citat. p. 31.

<sup>Philosoph. Trans. vol. xxx. p. 758.
Op. citat. p. 411.
Philos. Trans. for 1799, p. 355.
System of Chemistry, vol. iv.</sup> 

<sup>&</sup>amp; Mr. Jennings, Transactions of the Provincial Medical and Surgical Association, vol. 1.

pulmonum, Lungenprobe, or Athemprobe, ("Lung-

proof or Respiration-proof,") of the Germans.

Expiration, like inspiration, has been divided into three grades; ordinary, free, and forced; but it must necessarily admit of multitudinous shades of difference. In ordinary passive respiration, expiration is effected solely by the relaxation of the diaphragm. In free active respiration, the muscles that raise the ribs are likewise relaxed, and there is a slight action of the direct expiratory muscles. In forced expiration, all the respiratory muscles are thrown into action. In this manner, the air makes its way along the airpassages through the mouth or nostrils or both; carrying with it a fresh portion of the halitus from the mucous membrane. This it deposits, when the atmosphere is colder than the temperature acquired by the respired air, and if the atmosphere be sufficiently cold, as in winter, the vapour becomes condensed as it passes out, and renders expiration visible.

If the whole time occupied by a respiratory act — that is, from the beginning of one inspiration to the beginning of the next — be represented by 10, the time occupied by the inspiratory movement has been estimated approximately at 5; that of the expiratory at 4; and of the pause between the expiratory and succeeding inspi-

ratory movement at 1.a

The number of respirations in a given time differs considerably in different individuals. Dr. Hales reckons them at twenty. A man, on whom Menzies made experiments, breathed only fourteen times in a minute. Sir Humphry Davy made between twentysix and twenty-seven in a minute. Dr. Thomson about nineteen, and Magendie fifteen. Our own average, is sixteen. The average, deduced from the few observers, that have recorded their statement, - or twenty per minute, - has generally been taken; but we are satisfied it is above the truth; eighteen would be nearer the general average; and it has accordingly been admitted by many. Eighteen in a minute give twenty-five thousand nine hundred and twenty in the twenty-four hours. The number is influenced, however, by various circumstances. The child and the female, and perhaps also the aged, breathe more rapidly than the adult male. MM. Hourmann and Dechambreb examined two hundred and fifty-five women, between the ages of sixty and ninety-six, the average number of whose respirations was 21.79 per minute. We find as much variety in man as we do in the horse: whilst some men are short, others are long-winded; and this last condition may be improved by appropriate training; to which the pedestrian and the prize-fighter, equally with the horse, are submitted for some time before they are called upon to exhibit their powers. In sleep, the respiration is generally deeper,

<sup>\*</sup> Walshe, Physical Diagnosis and Diseases of the Lungs, Amer. Edit. p. 11, Philad.

b Archiv. Génér. de Médecine, Nov. 1835; and British and Foreign Med. Rev., April, 1836, p. 555.

less frequent, and appears to be effected greatly by the intercostals and diaphragm. Motion has also a sensible effect in hurrying the respiration, as well as distension of the stomach by food, certain mental emotions, &c.: its condition during disease becomes, also, a subject of interesting study to the physician, and one that has been much facilitated by the acoustic method, introduced by Laennec. To his instrument — the stethoscope — allusion has already been made. By it or by the ear applied to the chest, we are able to hear distinctly the respiratory murmur and its modifications; and thus to judge of the nature of the pulmonary affection when existent. But this is a topic that appertains more especially to pathology.

### 3. RESPIRATORY PHENOMENA CONCERNED IN CERTAIN FUNCTIONS.

There are certain respiratory movements concerned in effecting other functions, which require consideration. Some of these have already been topics of discussion. Adelonb has classed them into: First. Those employed in the sense of smell, either for the purpose of conveying the odorous molecules into the nasal fossæ; or to repel them and prevent their ingress. Secondly. The inspiratory action employed in the digestive function, as in sucking. Thirdly. Those connected with muscular motion when forcibly exerted: and particularly in straining or the employment of violent effort. Fourthly. Those concerned in the various excretions, either voluntary, - as in defecation and spitting; or involuntary, - as in coughing, sneezing, vomiting, accouchement, &c.; and lastly, such as constitute phenomena of expression — as sighing, yawning, laughing, crying, sobbing, &c. Some of these, that have already engaged attention, do not demand comment; others are topics of considerable interest, and require investigation.

1. Straining. — The state of respiration is much affected during the more active voluntary movements. Muscular exertion, of whatever kind, when considerable, is preceded by a long and deep inspiration; the glottis is then closed; the diaphragm and respiratory muscles of the chest are contracted, as well as the abdominal muscles which press upon the contents of the abdomen in all directions. At the same time that the proper respiratory muscles are exerted, those of the face participate, owing to their association through the medium of particular nerves. By this series of actions, the chest is rendered capacious; and the force, that can be developed is augmented, in consequence of the trunk being rendered immovable as regards its individual parts, — thus serving as a fixed point for the muscles that arise from it, so that they are

enabled to employ their full effect.c


The physiological state of muscular action, as connected with the mechanical function of respiration, is happily described by

Adelon, Physiologie de l'Homme, iii. 185.
 Ibid. p. 190; and art. Effort, in Dict. de Med. vii. 319, Paris, 1823.

Shakspeare, when he makes the 5th Harry encourage his soldiers

at the siege of Harfleur.a

In the effort required for effecting the various excretions, a similar action of the respiratory muscles takes place. The organs, from which these excretions have to be removed, exist either in the thorax or abdomen; and in all cases, they have to be compressed by the parietes of those cavities. See Fig. 151. A full inspiration is first made; the expiratory muscles, with those that close the glottis, are then forcibly and simultaneously contracted, and by this means the thoracic and abdominal viscera are compressed. Some difference, however, exists, according as the viscus



Thoracic and Abdominal Viscera.

A. Right lung. B. Left lung. C. Right ventricle of the heart. D. Right auricle of the heart. E. Vena cava superior. F. F. Subclavian veins. G. G. Internal jugular veins. H. Ascending aorta. I. Pulmonary artery. K. Diaphragm. L. L. Right and left lobes of the liver. M. Ligamentum rotundum. N. Fundus of gall-bladder. O. Stomach. P. Spleen. Q. Q. Situation of the kidneys, behind the intestines. R. R. Small intestines.

to be emptied, is seated in the abdomen or thorax. In the evacuation of the fæces, the lungs are first filled with air; and, whilst the muscles of the larvnx contract to close the glottis, those of the abdomen contract also: and as the lung, in consequence of the included air, resists the ascent of the diaphragm, the compression bears upon the large intestine. The same happens in the excretion of the urine, and in accouchement.

2. Coughing and Sneezing. - When the organ, that has to be cleared, is in the thorax, - as in coughing to remove mucus from the air-passages - the same action of the muscles of the abdomen is invoked; but the glottis is open to allow the lently from the lungs,

and, in its passage, sweeps off the irritating matter and conveys it

a "Stiffen the sinews, summon up the blood; Now set the teeth, and stretch the nostrils wide; Hold hard the breath and bend up every spirit To its full height." King Henry V. iii. 1. out of the body. To aid this, the muscular fibres, at the posterior part of the trachea and larger bronchial tubes, contract, so as to diminish the calibre of these canals; and, in this way, expectoration is facilitated. The action differs, however, according as the expired air is sent through the nose or mouth; in the former case, constituting sneezing; in the latter, coughing. The former is more violent than the latter, and is involuntary; whilst the latter is not necessarily so. In both cases the movement is excited by some external irritant, applied directly to the mucous membrane of the windpipe or nose; or by some modified action in the very tissue of the part, which acts as an irritating cause. In both cases the air is driven forcibly forward, and both are accompanied by sounds that cannot be mistaken. In these actions, we have striking exemplifications of the extensive association of muscles, through the medium of nerves, to which we have so often alluded. The pathologist, too, has repeated opportunities for observing the extensive sympathy between distant parts of the frame, as indicated by the actions of sneezing and coughing, especially of the former. If a person be exposed for a short period to the partial and irregular application of cold, so that the capillary action of a part of the body is modified, as where we get the feet wet, or sit in a draught of air, a few minutes will frequently be sufficient to exhibit sympathetic irritation in the Schneiderian membrane of the nose, and sneezing, Nor is it necessary, that the capillary action of a distant part shall be modified by the application of cold. We have had the most positive evidence, that if the capillary circulation be irregularly excited, even by the application of heat, whilst the rest of the body is receiving none of its influence, inflammation of the mucous membrane of the nasal fossæ and fauces follows with no less certainty.

3. Blowing the Nose. — The substance that has to be excreted by this operation is composed of the nasal mucus, the tears sent down the ductus ad nasum, and the particles deposited on the membrane by the air, in its passage through the nasal fossæ. Commonly, these secretions are only present in quantity sufficient to keep the membrane moist, the remainder being evaporated or absorbed. Frequently, however, they exist in such quantity as to fall by their own gravity into the pharynx, where they are sent down into the stomach by deglutition, are thrown out at the mouth, or make their exit at the anterior nares. To prevent this last effect more especially, we have recourse to blowing the nose. This is accomplished by taking in air, and driving it out suddenly and forcibly, closing the mouth at the same time, so that the air may issue by the nasal fossæ and clear them; the nose being compressed so as to make the velocity of the air greater, as well as

to express all the mucus that may be forced forwards.

4. Spitting differs somewhat according to the part in which the mucus or matter to be ejected is seated. At times, it is exclusively in the mouth; at others in the back part of the nose, pharynx or

VOL. II. - 4

larynx. When the mucus or saliva of the mouth has to be excreted, the muscular parietes of the cavity, as well as the tongue, contract so as to eject it from the mouth; the lips being at times approximated, so as to render the passage narrow, and impel the sputa more strongly forward. The air of expiration may be, at the same time, driven forcibly through the mouth, so as to send the matter to a considerable distance. The practised spitter sometimes astonishes us with the accuracy and power of propulsion of which he is capable. When the matter to be evacuated is in the nose, pharynx, or larynx, it requires to be brought, first of all, into the mouth. If in the posterior nares, the mouth is closed, and the air is drawn in forcibly through the nose, the pharynx being at the same time constricted so as to prevent the substances from passing down into the œsophagus. The pharynx now contracts, from below to above, in an inverse movement from that required in deglutition, and the farther excretion from the mouth is effected in the manner just described.

Where the matters are situate in the air-passages, the action may consist in coughing; or, if higher up, simply in hawking. A forcible expiration, unaccompanied by cough, is, indeed, in many cases, sufficient to detach the superfluous mucous secretion from even the bronchial tubes. In hawking, the expired air is forcibly sent forwards, and the parts about the fauces are suddenly contracted so as to diminish the capacity of the tube and propel the matters onwards. The noise is produced by their discordant vibrations. Both these modes bear the general name of expectoration.

When these secretions are swallowed, they are subjected to the digestive process; a part is taken up, and the remainder rejected; so that they belong to the division of recremento-excrementitial

fluids of some physiologists.

### 4. RESPIRATORY PHENOMENA CONNECTED WITH EXPRESSION.

It remains to speak of the expiratory phenomena that strictly form part of the function of expression, and depict the moral feeling

of the individual who gives utterance to them.

1. Sighing consists of a deep inspiration, by which a large quantity of air is received slowly and gradually into the lungs, to compensate for the deficiency in the due aëration of the blood which precedes it. The most common cause of sighing is mental uneasiness; it also occurs at the approach of sleep, or immediately after waking. In all these cases, the respiratory efforts are executed more imperfectly than under ordinary circumstances: the blood consequently does not circulate through the lungs in due quantity, but accumulates more or less in these organs, and in the right side of the heart; and it is to restore the due balance, that the deep inspiration is now and then established.

2. Yawning, oscitancy, oscitation or gaping, is likewise a full, deep, and protracted inspiration, accompanied by a wide separation of the jaws, and followed by a prolonged and sometimes sono-

rous expiration. Yawning is excited by many of the same causes as sighing. It is not, however, the expression of any depressing passion, but is occasioned by any circumstance that impedes the necessary aëration of the blood; whether this be retardation of the action of the respiratory muscles, or the air being less rich in oxygen. Hence we yawn at the approach of sleep, and immediately after waking. The inspiratory muscles, fatigued from any cause, experience some difficulty in dilating the chest; the lungs are consequently not properly traversed by the blood from the right side of the heart: oxygenation is, therefore, not duly effected, and an uneasy sensation is induced, which is put an end to by the action of yawning, which allows the admission of a considerable quantity of air. We yawn at the approach of sleep, because the agents of respiration, becoming gradually more debilitated, require to be now and then excited to fresh activity, and the blood needs the necessary aëration. Yawning on waking seems to be partly for the purpose of stimulating the respiratory muscles to greater activity, the respiration being always slower and deeper during sleep. It is, of course, impossible to explain, why the respiratory nerves should be those that are chiefly concerned, under the guidance of the brain, in these respiratory movements of an expressive character. The fact, however, is certain; and it is remarkably proved by the circumstance, that yawning can be excited by even looking at another affected in this manner; nay, by simply looking at a sketch, and by even thinking of the action. The same also applies to sighing and laughing, and especially to the latter.

3. Pandiculation or stretching is a frequent concomitant of yawning, and appears to be established instinctively to arouse the extensor muscles to a balance of power, when the action of the flexors has been predominant. In sleep, the flexor muscles exercise that preponderance which, in the waking state, is exerted by the extensors. This, in time, is productive of some uneasiness; and, hence, at times during sleep, but still more at the moment of waking, the extensor muscles are roused to action, to restore the equipoise; or, perhaps, as the muscles of the upper extremities, and those concerned directly or indirectly in respiration, are chiefly concerned in the action, it is exerted for the purpose of arousing

the respiratory muscles to increased activity.

By Dr. Good, a yawning and stretching have been regarded as morbid affections and amongst the signs of debility and lassitude:

—"Every one," he remarks, "who resigns himself ingloriously to a life of lassitude and indolence will be sure to catch these motions as a part of that general idleness which he covets. And, in this manner, a natural and useful action is converted into a morbid habit; and there are loungers to be found in the world, who, though in the prime of life, spend their days as well as their nights in a perpetual routine of these convulsive movements, over which they have no power; who cannot rise from the sofa without stretching

<sup>\*</sup> Study of Medicine, class 4, ord. 3, gen. 2, sp. 6.

their limbs, nor open their mouths to answer a plain question without gaping in one's face. The disease is here idiopathic and chronic; it may perhaps be cured by a permanent exertion of the will, and ridicule or hard labour will generally be found the best reme-

dies for calling the will into action."

4. Laughing is a convulsive action of the muscles of respiration and voice, accompanied by a facial expression, which has been explained elsewhere. It consists of a succession of short, sonorous expirations. The air is first inspired so as to fill the lungs. To this succeed short, interrupted expirations, with simultaneous contractions of the muscles of the glottis, so that this aperture is slightly contracted, and the lips assume the tension, necessary for the production of sound. The interrupted character of the expirations is caused by convulsive contractions of the diaphragm, which constitute the greater part of the action. In very violent laughter, the respiratory muscles are thrown into such forcible contractions, that the hands are applied to the sides to support them. The convulsive action of the thorax likewise interferes with the circulation through the lungs; the blood, consequently, stagnates in the upper part of the body; the face becomes flushed; the sweat trickles down the forehead, and the eyes are suffused with tears; but this is apparently owing to mechanical causes; not to the lachrymal gland being excited to unusual action, as in weeping. At times, however, we find the latter cause in operation, also.

5. Weeping. The action of weeping is very similar to that of laughing; although the causes are so dissimilar. It consists in an inspiration, followed by a succession of short, sonorous expirations. The facial expression, so diametrically opposite to that of laughter,

has been depicted in another place.

Laughter and weeping appear to be characteristic of humanity. Animals shed tears, but this does not seem to be accompanied with the mental emotion that characterizes crying in the sense in which we employ the term. It has, indeed, been affirmed by Steller, that the phoca ursina or ursine seal; by Pallas, that the camel; and by Von Humboldt, that a small American monkey, shed tears when labouring under distressing emotions. The last scientific traveller states, that "the countenance of the titi of the Orinoco, — the simia sciurea of Linnæus, — is that of a child; the same expression of innocence; the same smile; the same rapidity in the transition from joy to sorrow. The Indians affirm, that it weeps like man, when it experiences chagrin; and the remark is accurate. The large eyes of the ape are suffused with tears, when it experiences fear or any acute suffering."

Shakspeare's description of the weeping of the stag, - "That from the hunter's aim had ta'en a hurt,"

a Nov. Comm. Academ. Scient. Petropol. ii. 353.

b Sammlungen Historisch. Nachricht. über die Mongolischen Völkerschaften, Th. i. 177.

Recueil d'Observations de Zoologie, &c. i. 333.

is doubtless familiar to most of our readers.<sup>a</sup> We have less evidence in favour of the laughter of animals. Le Cat,<sup>b</sup> indeed, asserts, that he saw the chimpansé both laugh and weep. The ourang-outang, carried to Great Britain from Batavia, by Dr. Clarke Abel, never laughed; but he was seen occasionally to

weep.c

6. Sobbing still more resembles laughing, except that, like weeping, it is usually indicative of the depressing passions; and generally accompanies weeping. It consists of a convulsive action of the diaphragm; which is alternately raised and depressed, but to a greater extent than in laughing, and with less rapidity. It is susceptible of various degrees, and has the same physical effects upon the circulation as weeping. Dr. Wardrop<sup>d</sup> considers laughter, crying, weeping, sobbing, sighing, &c., as efforts made with a view to effect certain alterations in the quantity of blood in the lungs and heart, when the circulation has been disturbed by mental emotions.

- 7. Panting or anhelation consists in a succession of alternate, quick, and short inspirations and expirations. Their physiology, however, does not differ from that of ordinary respiration. The object is, to produce a frequent renewal of air in the lungs, in cases where the circulation is unusually rapid; or where, owing to disease of the thoracic viscera, a more than ordinary supply of fresh air is demanded. We can, hence, understand, why dyspnæa should be one of the concomitants of most of the severe diseases of the chest; and why it should occur whenever the air we breathe does not contain a sufficient quantity of oxygen. The panting, produced by running, is owing to the necessity for keeping the chest as immovable as possible, that the whole effort may be exerted on the muscles of locomotion; and thus suspending, for a time, the respiration, or admitting only of its imperfect accomplishment. This induces an accumulation of blood in the lungs and right side
  - \* "The wretched animal heaved forth such groans,
    That their discharge did stretch his leathern coat
    Almost to bursting; and the big round tears
    Coursed one another down his innocent nose\*
    In piteous chase; and thus the hairy fool,
    Much marked of the melancholy Jaques,
    Stood on th' extremest verge of the swift brook,
    Augmenting it with tears."

As You Like It, ii. 1.

b Traité de l'Existence du Fluide des Nerves, p. 35.

d On the Nature and Treatment of Diseases of the Heart, part i. p. 62, Lond. 1837.

Lawrence, Lectures on Physiology, Zoology, and the Natural History of Man, p. 236, Lond. 1814.

<sup>\* &</sup>quot;The alleged big round tears," which course one another down the innocent nose of the deer, the hare and other animals when hotly pursued, are in fact only sebaceous matter, which, under these circumstances, flows in profusion from a collection of follicles in the hollow of the cheek." — Fletcher's Rudiments of Physiology, part. ii. b. p. 50, Edinb. 1836.

of the heart; and panting is the consequence of the augmented action necessary for transmitting it through the vessels.

# b. Chemical Phenomena of Respiration.

Having studied the mode in which air is received into, and expelled from, the lungs, we have now to inquire into the changes produced on the venous blood - containing the products of the various absorptions - in the lungs, as well as on the air itself. These changes are effected by the function of sanguification, hæmatosis, respiration, in the restricted sense in which it is employed by some, arterialization of the blood, decarbonization, aëration, atmospherization, &c. With the ancients this process was but little understood. It was generally believed to act as the means of cooling the body; and, in modern times, Helvetius revived the notion, attributing to it the office of refrigerating the blood, - heated by its passage through the long and narrow channels of the circulation, - by the cool air constantly received into the lungs. The reasons, that led to this opinion, were: that the air, which enters the lungs in a cool state, issues warm; and that the pulmonary veins, which convey the blood from the lungs, are of less dimension than the pulmonary artery, which conveys it to them. From this it was concluded, that the blood, during its progress through the lungs, must lose somewhat of its volume or be condensed by refrigeration. The warmth of the expired air can, however, be readily accounted for; and it is not true that the pulmonary veins are smaller than the pulmonary artery. The reverse is, indeed, the fact; and it is equally obvious, that the doctrine of Helvetius does not explain how we can exist in a temperature superior to our own: this ought, in his hypothesis, to be impracticable.a

Another theory, which prevailed for some time, was:—that during inspiration the vessels of the lungs are unfolded, as it were, and that thus the passage of the blood from the right side of the heart to the left, through the lungs, is facilitated. Its progress was, indeed, conceived to be impossible during expiration, in consequence of the considerable flexures of the pulmonary vessels. The discovery of the circulation of the blood gave rise to this theory; and Hallerb attaches considerable importance to it, when taken in connexion with the changes effected upon the blood in the vessels. It is inaccurate, however, to suppose, that the circulation of the blood through the lungs is mechanically interrupted, when respiration is arrested. The experiments of Williamsc and Kayd would seem to show, that the interruption is owing to the non-conversion of venous into arterial blood, and to the non-adaptation of the radicles of the pulmonary veins for any thing

Adelon, Physiologie de l'Homme, edit. cit. iii. 201. Element. Physiol. viii. Edinb. Med. and Surg. Journ. vol. lxxvii, 1823.

d Edinb. Med. and Surg. Journ. vol. xxix.; and Physiology and Pathology, &c., of Asphyxia, Lond. 1834.

but arterial blood, owing to which causes stagnation of blood supervenes in the pulmonary radicles. Numerous other objections might be made to this view. In the first place, it supposes, that the lungs are emptied at each expiration, and, again, if a simple deploying or unfolding of the vessels were all that is required, any gas ought to be sufficient for respiration, — which is not the fact.

In these different theories, the principal object of respiration is overlooked — the conversion of the venous blood and its various absorptions, conveyed to the lungs by the pulmonary artery, into arterial blood. This is effected by the contact of the inspired air with the venous blood; in which they both lose certain elements, and gain others. Most physiologists have considered that the whole function of hæmatosis is effected in the lungs. Chaussier, however, has presumed, that the air, in passing through the cavities of the nose and mouth, and the different bronchial ramifications, experiences some kind of elaboration, by being agitated with the bronchial mucus; similar to what he conceives to be effected by the mucus on the aliment in its passage from the mouth to the stomach; but his view is conjectural in both one case and the other.

Legallois, again, thought, that hæmatosis commences at the part, where the chyle and lymph are mixed with the venous blood, or in the subclavian veins. This admixture, he conceives, occurs more or less immediately; is aided in the heart, and the conversion is completed in the lungs. To this belief he was led by the circumstance, that when the blood quits the lungs it is manifestly arterial, and he thought, that what the products of absorption lose or gain in the lungs is too inconsiderable to account for the important and extensive change; and that therefore it must have commenced previously. Facts, however, are not exactly in accordance with the view of Legallois. They seem to show, that the blood of the pulmonary artery is analogous to that of the subclavian veins; and hence it is probable, that there is no other action exerted upon the fluid in this part of the venous system, than a more intimate admixture of the venous blood with the chyle and lymph in their passage through the heart.

The changes, wrought on the air by respiration, are considerable. It is immediately deprived of a portion of both of its main constituents — oxygen and azote; and it always contains, when expired, a quantity of carbonic acid greater than it had when received into the lungs, along with an aqueous and albuminous ex-

halation to a considerable amount.

b Adelon, Physiologie, de l'Homme, iii. 205.

Annales de Chimie, iv. 115.

<sup>\*</sup> See the article "Asphyxia," by the author, in the "American Cyclopedia of Practical Medicine and Surgery," part. x., Philad. 1836; and Practice of Medicine, 2d edit. vol. i., Philad. 1844; also, Dr. W. P. Alison, in the Fourth Report of the British Association, and in Edinb. Med. and Surg. Journ for Jan. 1836; and Dr. J. Reid, on the order of succession in which the vital actions are arrested in Asphyxia, Edinb. Med. and Surg. Journ., April, 1841, p. 437.

Oxygen is consumed in the respiration of all animals, from the largest quadruped to the most insignificant insect; and, if we examine the expired air, the deficiency is manifest. Many attempts have been made to estimate the precise quantity consumed during respiration; but the results vary essentially from each other; partly owing to the fact, that the amount consumed by the same animal in different circumstances is not identical. Menziesa was probably the first that attempted to ascertain the quantity consumed by a man in a day. According to him, 36 cubic inches are expended in a minute; and, consequently, 51840 in the twenty-four hours, equal to 17496 grains. Lavoisier makes it 46048 cubic inches, or 15541 grains. This was the result of his earlier experiments; and, in his last, which he was executing at the time when he fell a victim to the tyranny of Robespierre, he made it 15592.5 grains; corresponding largely with the results of his earlier observations. The experiments of Sir Humphry Davye coincide greatly with those of Lavoisier. He found the quantity consumed in a minute to be 31.6 cubic inches; making 45504 cubic inches, or 15337 grains in twenty-four hours. The results obtained by Messrs. Allen and Pepysd make it much less. They consider the average consumption to be, in the twenty-four hours, under ordinary circumstances, 39534 cubic inches, equal to 13343 grains. Now, if we regard the experiments of Lavoisier and Davy, between which there is the greatest coincidence, to be an approximation to the truth, it will follow, that in a day, a man consumes rather more than 25 cubic feet of oxygen; and as the oxygen amounts to only about one-fifth of the respired air, he must render 125 cubic feet of air unfit for supporting combustion and respiration. The experiments of Crawford, Jurine, Lavoisier and Séguin, Prout, Fyfe, and Edwards, have proved, that the quantity of oxygen consumed varies according to the condition of the functions and of the system generally. Séguinf found, that muscular exertion increases it nearly fourfold. Prout, who gave much attention to the subject, was induced to conclude, from his experiments, that moderate exercise increases the consumption; but if the exercise be continued so as to induce fatigue, a diminished consumption takes place. The exhilarating passions also appeared to increase the quantity; whilst the depressing passions and sleep, the use of alcohol and tea, diminished it. He discovered, also, that the quantity of oxygen consumed is not uniformly the same during the twenty-four hours. Its maximum occurred between 10 A.M. and 2 P.M., or generally between 11 A.M. and 1 P.M.; its minimum commenced about 82 P.M., and it continued nearly uniform till about 32 A.M. Dr. Fyfeh found, that the quantity was likewise dimi-

Dissertation on Respiration, p. 21, Edinb. 1796.

Mémoir. de l'Académ. des Sciences, 1789, 1790.
 Researches, &c. p. 431.
 Philos. Transact. for 1808.

De l'Influence des Agens Physiques sur la Vie, p. 410, Paris, 1824; or Hodgkin and Fisher's translation.
 β Mém. de l'Académ. des Sciences, 1789 and 1790.
 β Annals of Philos. ii. 330, iv. 331, and xiii. 269.

h Annals of Philos. iv. 334, and Bostock's Physiol. i. 350.

nished by a course of nitric acid, by a vegetable diet, and by affecting the system with mercury. Temperature, also, has an influence, Crawforda found, that a Guinea-pig, confined in air at the temperature of 55°, consumed double the quantity which it did in air at 104°. He also observed, in such cases, that the venous blood, when the body was exposed to a high temperature, had not its usual dark colour; but, by its florid hue, indicated that the full change had not taken place in its constitution, in the course of circulation. We may thus understand the great lassitude and yawning, induced by the hot weather of summer; and the languor and listlessness which are so characteristic of those who have long resided in torrid climes. Dr. Prout conceives, that the presence or absence of the sun alone regulates the variation in the consumption of oxygen which he has described; but the deduction of Dr. Fleming appears to us more legitimate, — that it keeps pace with the degree of muscular action, and is dependent upon it. Consequently, a state of increased consumption is always followed by an equally great decrease, in the same manner as activity is

followed by fatigue.

The disagreement of experimenters, as respects the removal of nitrogen or azote from the air, during respiration, is still greater than in the case of oxygen. Priestley, Davy, Humboldt, Henderson, Cuvier, Pfaff, and Thomson, found a less quantity exhaled than was inspired. Spallanzani, Lavoisier and Séguin, Vauquelin, Allen and Pepys, Ellis, and Dalton, inferred that neither absorption nor exhalation takes place, - the quantity of that gas, in their opinion, undergoing no change during its passage through the aircells of the lungs; whilst Jurine, Nysten, Berthollet, and Dulong and Despretz, on the contrary, found an increase in the bulk of the azote. In this uncertainty, most physiologists have been of opinion that the azote is entirely passive in the function. The facts, ascertained by Dr. W. F. Edwards,d of Paris, shed considerable light on the causes of this discrepancy amongst observers. He has satisfactorily shown that, during the respiration of the same animal, the quantity of azote may, at one time, be augmented, at another, diminished, and, at a third, wholly unchanged. These phenomenahe has traced to the influence of the seasons, and he suspects that other causes have a share in their production. In nearly all the lower animals that were the subjects of experiment, an augmentation of azote was observable during summer. Sometimes, indeed, it was so slight that it might be disregarded; but, in numerous other instances, it was so great as to place the fact beyond the possibility of doubt; and, on some occasions, it almost equalled the whole bulk of the animal. Such were the results of his observations until the close of October, when he noticed a sensible diminution in the nitrogen of the inspired air, and the same continued throughout the whole of winter and the beginning of spring. Dr.

Do. cit. p. 387.
Bostock, op. citat. 3d edit. p. 356.
Philosophy of Zo
Op. citat. p. 462.

b Philosophy of Zoology, Edinburgh, 1822.

Edwards considers it probable, that, in all cases, both exhalation and absorption of azote are going on; that they are frequently accurately balanced, so as to exhibit neither excess nor deficiency of azote in the expired air, whilst, in other cases, depending, as it would appear, chiefly upon temperature, either the absorption or the exhalation is in excess, producing a corresponding effect upon

the composition of the air of expiration.

But not only has the respired air lost its oxygenous portion, it has gained, as we have remarked, an accession of carbonic acid, and, likewise, a quantity of serous vapour. If we breathe through a tube, one end of which is inserted into a vessel of lime-water, the fluid soon becomes milky, owing to the formation of carbonate of lime, which is insoluble in water. Carbonic acid must consequently have been given off from the lungs. In the case of this gas, again, the quantity, formed in the day, has been attempted to be computed. Jurine conceived, that the amount, in air once respired in natural respiration, is in the enormous proportion of 1/6th or 1/2th. Menzies, that it is 1 th; and, from his estimate of the total quantity of air respired in the twenty-four hours, he deduced the amount of carbonic acid formed to be 51840 cubic inches, equal to 24105.6 grains. Lavoisier and Séguin, in their first experiments, valued it at 17720.89 grains; but in the very next year they reduced their estimate more than one-half; - to 8450.20 grains; and, in Lavoisier's last experiment, it was farther reduced to 7550.4 grains. Sir Humphry Davy's estimate nearly corresponds with that of the first experiment of Lavoisier and Séguin, - 17811.36 grains; and Messrs. Allen and Pepys accord pretty nearly with him. These gentlemen found, that atmospheric air, when inspired, issued on the succeeding expiration, charged with from 8 to 6 per cent. of carbonic acid gas; but this estimate exceeds considerably that of Dr. Apjohn, of Dublin, who, in his experiments, found the expired air to contain only 3.6 per cent of this gas. The experiments and observations of Crawford, Prout, Edwards, and others, to which we have referred - as regards the consumption of oxygen, under various circumstances - apply equally to the quantity of carbonic acid formed, which always bears a pretty close proportion to the oxygen consumed. These experiments also account, in some degree, for the discrepancy in the statements of different individuals on this subject.

One fatal objection, however, to all these estimates is, that more carbon must be given off than is received in the daily food. The experiments of Mr. Coathupe, which were carefully conducted, make the amount of carbonic acid generated in the 24 hours, about 17.856 cubic inches, that is, 2.616 grains or 5½ ounces of solid carbon. Liebig found the proportion of carbon expired by himself to

a Mémoir. de l'Academ. des Sciences, p. 609, Paris, 1790.

Edinb. Med. and Surg. Journal, Jan. 1831; Dublin Hospital Reports, vol. v.; and Select Medico-Chirurgical Transactions, p. 230, Philad. 1831. See, also, Mr. Coathupe, Philos. Magaz. June, 1839.

Philos. Magazine, June, 1839.

be 8½ ounces daily; by a soldier, 13½ ounces; by prisoners in close confinement, 7 ounces; and by a boy who took considerable exercise, 9 ounces." Recently, further experiments have been made on this subject by competent observers. Professor Scharling, of Copenhagen, found, that, at the age 35, he exhaled 7.7 ounces avoirdupois of carbon in the twenty-four hours - seven of which were passed in sleep. A soldier, 28 years of age, exhaled 8.15 ounces; a lad, of 16, 7.9 ounces; a young woman, aged 19, 5.83 ounces; a boy, 9½ years old, 3.069 ounces; and a girl, 10 years old, 4.42 ounces. In the two last, the time spent in sleep was 9 hours. From all his experiments, Professor Scharling deduces, that males exhale more carbonic acid than females; and children comparatively more than adults. MM. Andral and Gavarret have been engaged in an interesting set of experiments on this subject. Their first object was to ascertain the modifying influence of age, sex, and constitution on the quantity of carbonic acid exhaled from the lungs. To determine this, their observations were made under circumstances as uniform as possible. Each experiment, too, was repeated several times on the same subject. The apparatus employed was so devised as to enable the respirations to be freely performed: no portion of the expired air was again inspired; and the greatest care was taken to analyze the expired air with accuracy. The general results obtained by these observers were as follows: -1. The quantity of carbonic acid exhaled by the lungs in a given time varies according to the age, sex, and constitution. 2. In both male and female, the quantity undergoes modification, according to the ages of the individuals experimented upon, quite independently of their weights. 3. In all periods of life, there is a difference between the male and female in the amount of carbonic acid exhaled from the lungs in a given time: cæteris paribus, man exhales a much larger quantity than woman. Between the ages of 16 and 40, the former exhales nearly twice as much as the latter. 4. In man, the quantity exhaled goes on regularly increasing from 8 to 30 years of age; and a remarkable augmentation takes place at puberty. After 30, it begins to decrease; and the decrease continues becoming more and more marked as the individual approaches nearer and nearer to extreme old age; so that, at this last period, it returns to the standard at which it was about the age of 10. 5. In woman, the exhalation augments up to the period of puberty, according to the same law as in man: the increase then suddenly ceases, and the quantity continues at this low standard, with little variation so long as the catamenia regularly appear; but as soon as they cease, the exhalation of carbonic acid from the lungs undergoes a considerable augmentation, after which it decreases as in man, according as age advances. 6. During gestation, the amount of carbonic acid exhaled is raised tempora-

a Graham's Elements of Chemistry, Amer. Edit. p. 686, Philad. 1843.

b Annales des Sciences Naturelles, Février, 1843; and Brit, and For. Med. Rev. for July, 1843, p. 285.

rily to the standard which it attains after the cessation of the catamenia. 7. In both sexes, and at all ages, the quantity of carbonic acid exhaled by the lungs is greater in proportion to the strength of the constitution, and the development of the muscular system.

The following table exhibits the quantity of solid carbon calculated to be exhaled in one hour at different ages: the gramme is

equal to about 151 grains.

MALE.					FEMALE.					British on Auditor of
8 years.			5 g	rammes.	8 years.		177	5 g	rammes.	The same standard con
15 -		-	-	8.7	12-38	-	-	-	6.4	tinues in women during the
16 -	-	(4)	-	10.8	38-50	-	-		8.4	whole of the menstrual pe
18-20		-	-	11.4	50-60			-	7.3	riod: but if the catamenia
20-30		-	-	12.2	60-80	-		-	6.8	be temporarily suppressed
30-40	-	-	-	12.2	82	-	-	-	6.0	or pregnancy occur, it rise
40-60	-	-	-	10.1						to the standard it attain
60-80			-	9.2						after their entire cessation
102	-	-	-	5.9						namely, 8.4 grammes.

These numbers express the averages, the maximum amount being often considerably greater. In a young man of athletic system, and sound constitution, the quantity of carbonic acid exhaled in an hour was 14·1 grammes; a man of 60, equally vigorous for his age, exhaled 13·6 grammes; and one of 63, 12·4 grammes. An old man, of 92, who preserved a remarkable degree of energy, and who had possessed unusual vigour in his youth, was found to exhale 8·8 grammes per hour; whilst the same amount appeared to be the ordinary standard in a man of 45; who, unlike the last, had a feeble system, although he was in equally good health. How far these variations were connected with differences in the capacity of the chest, and with the number of the respiratory movements, MM. Andral and Gavarret propose to investigate subsequently.

It has been a question amongst physiologists, whether the quantity of carbonic acid gas given out its equal in bulk to the oxygen taken in. In Priestley's experiments, the latter had the preponderance. Menzies and Crawford found them to be equal. Lavoisier and Séguin supposed the oxygen, consumed in the twentyfour hours, to be 15661.66 grains; whilst the oxygen, required for the formation of the carbonic acid given out, was no more than 12924 grains; and Sir Humphry Davy, in the same time, found the oxygen consumed to be 15337 grains, whilst the carbonic acid produced was 17811.36 grains; which would contain 12824.18 grains of oxygen. The experiments of Allen and Pepys seem, however, to show that the oxygen which disappears is replaced by an equal volume of carbonic acid; and hence it was supposed that the whole of it must have been employed in the formation of this acid. They, consequently, accord with Menzies and Crawford; and the view is embraced by Dalton, Prout, Ellis, Henry, and other distinguished individuals. On the other hand, the view of

Experiments, &c. on different kinds of Air, vol. iii.
 Bostock's Physiology, 3d edit. p. 352, Lond. 1836.

those, who consider that the quantity of carbonic acid produced is less than that of the oxygen which has disappeared, is embraced by Thomson, and by Dulong and Despretz. In the carnivorous animal, they found the difference as much as one-third; in the herbivorous, on the average, only one-tenth. The more recent experiments of Dr. Edwards have shown, that here, also, the discordance has not depended so much upon the different methods and skill of the operators, as upon a variation in the results arising from other causes; and he concludes, that the proportion of oxygen consumed, to that employed in the production of carbonic acid, varies from more than one-third of the volume of carbonic acid to almost nothing; that the variation depends upon the particular animal species, subjected to experiment; upon its age, or upon some peculiarity of constitution, and that it differs considerably in the same individual at different times.

It would appear, then, that the whole of the oxygen, which respiration abstracts from the air, is not accounted for, in all cases, by the quantity of carbonic acid formed; and that, consequently, a portion of it disappears altogether. It has been supposed by some, that a part of the watery vapour, given off during expiration, is occasioned by the union of a portion of the oxygen of the air with hydrogen from the blood in the lungs; but this view is entirely conjectural. This subject, with the quantity of vapour combined with the expired air, will be the subject of inquiry under the head of Secretion.

The air likewise loses, during inspiration, certain foreign matters that may be diffused in it. In this way, medicines have been attempted to be conveyed into the system. If air, charged with odorous particles,— as with those of turpentine,— be breathed for a short time, their presence in the urine will be detected; and it is probably in this manner, that miasmata produce their effects on the frame. All these substances pass immediately through the coats of the pulmonary veins by imbibition, and, in this way, speedily affect the system.

These changes, produced in the air during respiration, are easily shown, by placing an animal under a bell-glass until it dies. On examining the air, it will be found to have lost a portion of its oxygen and azote and to contain carbonic acid and aqueous vapour.

Let us now inquire whether the changes produced in the respired air be connected with those effected on the blood in the lungs. In its progress through the lungs, this fluid has been changed from venous into arterial. It has become of a florid red colour; of a stronger odour; of a higher temperature by nearly two degrees, according to some, but others have perceived no

Magendie, Précis de Physiologie, ii. 343; and Dr. J. Davy, in Philos. Transact. for 1814.

VOL. II. - 5

difference, whilst others, again, have found it of lower temperature; of less specific gravity, in the ratio of 1053 to 1050 on the average, according to Dr. John Davy, b and it coagulates more speedily, according to most observers, but Thackrahe observed the contrary.d That this conversion is owing to the contact of air in the lungs we have many proofs. Lowere was one of the first, who clearly pointed out, that the change of colour occurs in the capillaries of the lungs. Prior to his time, the most confused notions had prevailed on the subject, and the most visionary hypotheses had been indulged. On opening the thorax of a living animal, he observed the precise point of the circulation at which the change of colour takes place, and he showed, that it is not in the heart, since the blood continues to be purple, when it leaves the right ventricle. He then kept the lungs artificially distended, first with a regular supply of fresh air, and afterwards with the same portion of air without renewing it. In the former case, the blood experienced the usual change of colour. In the latter, it was returned to the left side of the heart unchanged.

Experiments, more or less resembling those of Lower, have been performed by Goodwyn, Cigna, Bichat, Wilson Philip, and

numerous others, with similar results.

The direct experiments of Priestleyh more clearly showed, that the change, effected on the blood, was to be ascribed to the air. He found, that the clot of venous blood, when confined in a small quantity of air, assumed a scarlet colour, and that the air experienced the same change as from respiration. He afterwards examined the effect produced on the blood by the gaseous elements of the atmosphere separately, as well as by the other gaseous fluids that had been discovered. The clot was reddened more rapidly by oxygen than by the air of the atmosphere, whilst it was reduced to a dark purple by nitrogen, hydrogen, and carbonic acid.

Since Priestley's time, the effect of different gases on the colour of venous blood has been investigated by numerous individuals. The following is the result of their observations, as given by Thénard. It must be remarked, however, that all the experiments have been made on blood, when out of the body; and it by no means follows, that precisely the same changes would be accomplished if the fluid were circulating in the vessels.

\* Wagner's Elements of Physiology, by R. Willis, § 180, Lond. 1842.

b Physiological and Anatomical Researches, vol. ii.

· Inquiry into the Nature and Properties of the Blood, p. 42, Lond. 1819.

- d See, on this subject, J. Müller's Handbuch, u. s. w., Baly's translation, p. 323; and Burdach's Physiologie als Erfahrungswissenschaft, Band. iv. s. 381, Leipz. 1832.

  Tractatus de Corde, &c. c. iii. Amstelod. 1761.
  - The Connexion of Life with Respiration, &c., Lond. 1788.
    Recherches Physiol. sur la Vie et la Mort, Paris, 1800.
  - Experiments, &c. on different kinds of Air, &c. Lond. 1781.

Traité de Chimie, &c. 5e édit. Paris, 1827.

Gas.	Colour.	Remarks.
Oxygen	Rose red.	The blood employed
Atmospheric air	Do.	had been beaten, and,
Ammonia	Cherry Red.	consequently, deprived of
Gaseous oxide of carbon	Slightly violet red.	its fibrin.
Deutoxide of azote	Do.	The state of the s
Carburetted hydrogen -	Do.	DIESTAND TOWN TO SEE
Azote	Brown red.	CIT IN CO. OF CO. LANS.
Carbonic acid	Do.ª	South or when the state of the state of
Hydrogen	Do.	The state of the s
Protoxide of azote	Do.	Charles Allinois Coll Land P
	CDeep violet, passing	THE BUTTE THE PARTY OF THE PART
Arsenuretted hydrogen -	gradually to a green-	CHAIR THE AS DE SENSE
Sulphuretted hydrogen -	(ish brown.	Delegation of many from
Chlorohydric acid gas -	Maroon brown.	
Sulphurous acid gas -	Black Brown.	These three gases coa-
A STATE OF THE PARTY OF THE PAR	(Blackish Brown,	gulated the blood at the
Chlorine	2 passing by degrees	same time.
Walter State of the last of th	(to a yellowish white.)	THE RESERVE THE PERSON NAMED IN

It is sufficiently manifest, then, from the disappearance of a part of the oxygen from the inspired air, and from the effects of that gas on venous blood out of the body, that it forms an essential part in the function of sanguification. But we have seen, that the expired air contains an unusual proportion of carbonic acid. Hence carbon, either in its simple state or united with oxygen, must have been given off from the blood in the vessels of the lungs.

To account for these changes on chemical principles has been a great object with chemical physiologists at all times. At an early period, the conversion of venous into arterial blood was supposed to be a kind of combustion; and, according to the notion of combustion then prevalent, it was presumed to consist in the disengagement of phlogiston; in other words, the abstraction or addition of a portion of phlogiston, made the blood, it was conceived, arterial or venous; and the removal of phlogiston was looked upon as the principal use of respiration. This view was modified by Lavoisier, who proposed one of the chemical views to be now mentioned.

Two chief chemical theories have been framed to explain the mode in which the carbon is given off. The first is that of Black, Priestley, Lavoisier, and Crawford; — that the oxygen of the inspired air attracts carbon from the venous blood, and that the carbonic acid is generated by their union. The second, which has been supported by La Grange, Hassenfratz, Edwards, Müller,

- Müller says he agitated blood with hydrogen, but could perceive no change of colour. Handbuch, u. s. w., Baly's translation, p. 322, Lond. 1838.
  - b Lectures on the Elements of Chemistry, by Robison, ii. 87, Edinb. 1803.
  - c Philosoph. Transact. for 1776, p. 147.
  - d Mém. de l'Acad. des Sciences, pour 1777, p. 185.
  - o On Animal Heat, 2d edit. Lond. 1788.
  - f Annales de Chimie, ix. 269.
- De l'Influence des Agens Physiques, &c. p. 411, Paris, 1823; and Hodgkin and Fisher's translation.

  Physiology, by Baly, p. 537.

Bischoff, Magnus, and others, - that the carbonic acid is generated in the course of the circulation, and is given off from the venous blood in the lungs, whilst oxygen gas is absorbed. former of these views is still maintained by many physiologists. It is conceived, that the oxygen, derived from the air, unites with certain parts of the venous blood, - the carbon and the hydrogen, - owing to which union carbonic acid and water are found in the expired air; the venous blood, thus depurated of its carbon and hydrogen becomes arterialized; and, in consequence of these various combinations, heat enough is disengaged to keep the body always at the due temperature. According to this theory, as we have seen in the views of Priestley, Lavoisier, &c., respiration is assimilated to combustion. The resemblance, indeed, between the two processes is striking. The presence of air is absolutely necessary for respiration; in every variety of respiration the air is robbed of a portion of its oxygen; and hence a fresh supply is continually needed; and respiration is always arrested before the whole of the oxygen of the air is exhausted, and this partly on account of the residuary azote, and the carbonic acid gas given off during expiration. Lastly, it can be continued much longer when an animal is confined in pure oxygen gas than in atmospheric air. All these circumstances likewise prevail in combustion. Every kind of combustion requires the presence of air. A part of the oxygen is consumed; and, unless the air is renewed, combustion is impossible. It is arrested, too, before the whole of the oxygen is consumed, owing to the residuary azote. and the carbonic acid formed; and it can be longer maintained in pure oxygen than in atmospheric air. Moreover, when the air has been respired, it becomes unfit for combustion, - and conversely. Again, the oxygen of the air, in which combustion is taking place, combines with the carbon and hydrogen of the burning body; hence the formation of carbonic acid and water; and as, in this combination, the oxygen passes from the state of a very rare gas, or one containing a considerable quantity of caloric between its molecules, to the condition of a much denser gas, or even of a liquid, the whole of the caloric, which the oxygen contained in its former state, can no longer be held in the latter, and it is accordingly disengaged; hence the heat, which is given off. In like manner, in respiration, the oxygen of the inspired air, it is conceived, combines with the carbon and hydrogen of the venous blood, giving rise to the formation of carbonic acid and water; and, as in these combinations, the oxygen passes from the state of a very rare to that of a denser gas, or of a liquid, there is a considerable disengagement of caloric, which becomes the source of the high temperature maintained by the human body. M. Thénarda admits a modification of this view, - sanguification being owing, he conceives, to the combustion of the carbonaceous parts of the venous blood, and probably of its colouring matter, by the oxygen of the air.

<sup>\*</sup> Traité de Chimie, edit, estat.

This chemical theory, which originated chiefly with Lavoisier, and La Place and Séguin, was adopted by many physiologists with but little modification.

Mr. Ellis imagined, that the carbon is separated from the venous blood by a secretory process; and that, then, coming into direct contact with oxygen, it is converted into carbonic acid. The circumstance that led him to this opinion was his disbelief in the possibility of oxygen being able to act upon the blood through the animal membrane or coat of the vessel in which it is confined. It is obvious, however, that to reach the blood circulating in the lungs, the oxygen must, in all cases, pass through the coats of the pulmonary vessels. These coats, indeed, offer little or no obstacle, and, consequently, there is no necessity for the vital or secretory action suggested by Mr. Ellis. Priestley and Hassenfratz exposed venous blood to atmospheric air and oxygen in a bladder. In all cases, the parts of the blood, in contact with the gases, became of a florid colour. The experiments of Faust, Mitchell, and others (vol. i., p. 44), are, in this aspect, pregnant with interest. They prove the great facility with which the tissues are penetrated by the gases, and confirm the facts developed by the experiments of Priestley, Hassenfratz and others.

The second theory, — that the carbonic acid is generated in the course of the circulation, — was proposed by La Grange, in consequence of the objection he saw to the former hypothesis — that the lung ought to be consumed by the perpetual disengagement of caloric taking place within it; or, if not so, that its temperature ought, at least, to be superior to that of other parts. He accordingly suggested, that, in the lungs, the oxygen is simply absorbed, passes into the venous blood, circulates with it, and unites, in its course, with the carbon and hydrogen, so as to form carbonic acid and water, which circulate with the blood and are finally exhaled

from the lungs.

The ingenious and apparently accurate experiments of Dr. Edwardsa prove convincingly, not only that oxygen is absorbed by the pulmonary vessels, but that carbonic acid is exhaled from them. When he confined a small animal in a large quantity of air, and continued the experiment sufficiently long, he found, that the rate of absorption was greater at the commencement than towards the termination of the experiment; whilst at the former period, there was an excess of oxygen present, and at the latter an excess of carbonic acid. This proved to him that the diminution was dependent upon the absorption of oxygen, not of carbonic acid. His experiments, in proof of the exhalation of carbonic acid, ready formed, by the lungs, are decisive. Spallanzani had asserted, that when certain of the lower animals are confined in gases, containing no oxygen, the production of carbonic acid is uninterrupted. Upon the strength of this assertion, Edwards confined frogs in pure hydrogen, for a length of time. The result indicated, that carbonic

Op. citat. p. 437, and Messrs. Allen and Pepys, in Philos. Transactions for 1829.

acid was produced, and, in such quantity as to show, that it could not have been derived from the residuary air in the lungs; as it was, in some cases, equal to the bulk of the animal. The same results, although to a less degree, were obtained with fishes and snails, - the animals on which Spallanzani's observations were made. The experiments of Edwards were extended to the mammalia. Kittens, two or three days old, were immersed in hydrogen: they remained in this situation for nearly twenty minutes, without dying, and on examining the air of the vessel after death, it was found, that they had given off a quantity of carbonic acid greater than could possibly have been contained in their lungs at the commencement of the experiment. The conclusion, deduced by Dr. Edwards, from his various experiments, is, "that the carbonic acid expired is an exhalation proceeding wholly or in part from the carbonic acid contained in the mass of blood." Several experiments were subsequently made by M. Collard de Martigny, who substituted azote for hydrogen; and, in all cases, carbonic acid gas was given out in considerable quantity.

These and other experiments would seem, then, to show, that, in the lungs, carbonic acid is exhaled, and that oxygen and azote are absorbed. They would also seem to prove the existence of carbonic acid in venous blood, respecting which so much dissi-

dence has existed amongst chemists.

Allusion has already been made to the fact, that gelatin is not met with in the blood, and to the idea of Dr. Prout, b that its formation from albumen must be a reducing process. This process, he considers to be one great source of the carbonic acid, which exists in venous blood. Gelatin contains three or four per cent. less carbon than albumen; it enters into the structure of every part of the animal frame, and especially of the skin; the skin, indeed, contains little else than gelatin. Dr. Prout considers it, therefore, most probable, that a large part of the carbonic acid of venous blood is formed in the skin, and analogous textures. "Indeed," he adds, " we know that the skin of many animals gives off carbonic acid, and absorbs oxygen; - in other words, performs all the offices of the lungs; - a function of the skin perfectly intelligible, on the supposition, that near the surface of the body, the albuminous portions of the blood are always converted into gelatin." Gmelin, Tiedemann, and Mitscherlich,e and Stromeyer, affirm, however, on the strength of experiments, that the blood does not contain free carbonic acid gas, but that it holds a certain quantity in a state of combination, which is set free in the lungs, and commingles with the expired air. The views of Gmelin and Tiedemann, and Mits-

d Schweigger's Journal für Chimie, u. s. w., lxiv. 105,

<sup>&</sup>lt;sup>2</sup> Journal de Physiologie, x. 111. For an account of various experiments relative to the exhalation of carbonic acid, during respiration in gases which contain no oxygen, see Müller's Handbuch, u. s. w., Baly's translation, p. 338, Lond. 1838.

b Bridgewater Treatise, Amer. Edit. p. 280, Philad. 1834.
c Tiedemann und Treviranus, Zeitschrift für Physiol. B. v. H. i.

cherlich on this subject are as follows. It may be laid down as a truth, that the greater part, if not all, of the properties of secreted fluids are not dependent upon any act of the secreting organs, but are derived from the blood, which again must either owe them to the food, or to changes effected on it within the body. These changes are probably accomplished in part, during the process of digestion, but are doubtless mainly effected on the lungs by the contact of the blood with the air. Now, most of the animal fluids, when exposed to the air, generate, by the absorption of oxygen, acetic or lactic acid, and this is aided by an elevated temperature like that of the lungs. In their theory of respiration, the azote of the inspired air is but sparingly absorbed; by far the greater proportion remaining in the air-cells. The oxygen, on the other hand, penetrates the membranes freely, mingles with the blood, combines partly with the carbon and hydrogen of that fluid, and generates carbonic acid and water, which are thrown off with the expired air, whilst the remainder combines with the organic particles of the blood, forming new compounds, of which the acetic and lactic acids are some; these unite with the carbonated alkaline salts of the blood, and set free the carbonic acid, so that it can be thrown off by the lungs. The acetate of soda, thus formed during the passage of the blood through the lungs, is deprived of its acetic acid by the several secretions, especially by those of the skin and kidneys, and the soda again combines with the carbonic acid. which, during the circulation of the blood through the body, is formed by the decomposition of its organic elements. Carbonate of soda is thus regenerated and conveyed to the lungs, to be again decomposed by the fresh formation of acids in those organs.

Almost the same view is entertained by MM. Dumas and Boussingault, and it is esteemed by Professor Grahama to have a high degree of probability; and another view, in many respects similar, is held by Professor Arnold.b As it is more than probable, he remarks, that the carbonic acid occurs in the venous blood, united with some substance from which it is separated, with greater or less rapidity, by the contact of the atmospheric air; and as, further, the carbonate of the protoxide of iron greedily withdraws oxygen from the atmosphere, at the same time parting with its carbonic acid and becoming changed into a peroxide, it may reasonably be supposed, he thinks, that the carbonic acid of the venous blood is united with the iron of the red colouring matter, and that it is set free during the act of respiration, by the reciprocal action of the blood and air. The protoxide, at the same time, by absorption of oxygen, becomes a peroxide, which, during the circulation of the blood through the capillaries, again parts with its oxygen. Carbon is at the same time eliminated from the blood,

Elements of Chemistry, Amer. Edit. by Dr. Bridges, p. 687, Philad. 1843.
 Lehrbuch der Physiologie des Menschen, Zurich, 1836-7; and Brit. and For.
 Med. Rev. Oct. 1839, p. 481. See, on this subject, Dr. J. Davy, Researches, Physiological and Anatomical, in Dunglison's Amer. Med. Lib. Edit. p. 80, Philad. 1840,

and unites with the liberated oxygen to form carbonic acid, which is thrown out by the lungs, whilst oxygen is again absorbed. This is the view embraced by Liebig, who has affirmed, that the amount of iron present in the blood, if in the state of protoxide, is sufficient to furnish the means of transporting twice as much carbonic acid as can possibly be formed by the oxygen absorbed in

the lungs.

Chaussier and Adelon, bagain, regard the whole process of hæmatosis as essentially organic and vital. They think, that an action of selection and elaboration is exerted both as regards the reception of the oxygen and the elimination of the carbonic acid. But their arguments on this point are unsatisfactory, and are negatived by the facility with which oxygen can be imbibed, and with which carbonic acid transudes through animal membranes. In their view, the whole process is effected in the lungs, as soon as the air comes in contact with the vessels containing the venous blood. The imbibition of oxygen they look upon as a case of ordinary absorption; the transudation of carbonic acid as one of exhalation; both of which they conceive to be, in all cases, vital actions, and not to be

likened to any physical or chemical operation.

Admitting, then, that the oxygen and a portion of nitrogen absolutely enter the pulmonary vessels, of which we appear to have direct proof, are they, it has been asked, separated from the air in the air-cells, and then absorbed; or does the air enter undecomposed into the vessels, and then furnish the proportion of each of its constituents necessary for the wants of the system, the excess being rejected? Could it be shown that such a decomposition is actually effected at the point of contact between the pulmonary vessels and the air in the lungs, it would seem, at first, to prove the notion of Ellis, and of Chaussier and Adelon, that an action of selection, or of vitality is exerted; but the knowledge we have attained of late, of the transmission of gases through animal membranes, would suggest another explanation. The rate of transmission of carbonic acid is greater than that of oxygen; and that of oxygen greater than that of azote (see vol. i., p. 46). We can hence understand, that more oxygen than azote may pass through the coats of the pulmonary bloodvessels, and can comprehend the facility with which the carbonic acid, formed in the course of the circulation, permeates the same vessels, and mixes with the air in the lungs. Sir Humphry Davy is of opinion, that the whole of the air is absorbed, and that the surplus quantity of each of the constituents is subsequently discharged. In favour of this view, he remarks that air has the power of acting upon the blood through a stratum of serum, and he thinks that the undecomposed air must be absorbed before it can arrive at the blood in the vessels. It is

b Physiologie de l'Homme, edit. cit. iii. 254.

<sup>&</sup>lt;sup>a</sup> Animal Chemistry, Webster's Amer. Edit., p. 261, Cambridge, 1843.

An Enquiry into the Changes induced on Atmospheric Air, &c., Edinb. 1807; and Further Enquiries, Edinb. 1816.

obvious, however, from the different penetrating powers of the gases — oxygen and azote — which compose it, that the proportion of those constituents cannot be the same in the interior as at the exterior of the pulmonary vessels. Muller, however, accords with Davy, and supposes that the air, on entering the lungs, is decomposed in consequence of the affinity of the oxygen for the red particles of the blood; carbonic acid being formed, which is exhaled in the gaseous form, along with the greater part of the azote.

It has been remarked, that when oxygen is applied to venous blood, the latter assumes a florid colour. On what part of the blood, then, does the oxygen act? The general belief is, upon the red globules. The facts, hereafter stated in the description of venous blood, have shown, that these globules appear to consist of a colourless nucleus, surrounded by a coloured envelope; that both of these are devoid of colour, whilst they exist as chyle and lymph; but that, in the lungs, the contact of air changes the envelope to a florid red. Some, indeed, have believed, that both the envelope and its colour are added in the lungs. The coloration of the blood. consequently, seems to be effected in the lungs; but whether this change be of any importance in hæmatosis is doubtful. Several tissues of the body are not supplied with red blood: in many animals, the red colour does not exist; and, in all, it can perhaps only be esteemed an evidence, that the other important changes have been accomplished in the lungs. Recently, the opinion has been revived, that the oxygen of the air acts upon the iron, which Engelhart and Roseb have detected in the colouring matter, but how we know not. It is asserted, that if the iron be separated, the rest of the colouring matter, which is of a venous red colour, loses the property of becoming scarlet by the contact of oxygen.

A different view of arterialization has been advanced by Dr. Stevens. According to him, the colouring matter of the blood is naturally very dark; is rendered still darker by acids, and acquires a florid hue from the addition of chloride of sodium, and from the neutral salts of the alkalies generally. The colour of arterial blood is ascribed by him to hematosin reddened by the salts contained in the serum; the characters of venous blood to the presumed presence of carbonic acid, which, like other acids, darkens hematosin; and the conversion of venous into arterial blood to the influence of the saline matter in the serum being restored by the separation of carbonic acid. If we take a firm clot of venous

<sup>&</sup>lt;sup>a</sup> Handbuch, u. s. w., Baly's translation, p. 334, Lond. 1838; see, also, Magnus, in Annales de Chimie et de Physique, Nov. 1837.

b Edinb. Med. and Surg. Journal, for Jan. 1827.

Observations on the Healthy and Diseased Properties of the Blood, Lond. 1832; and Proceedings of the Royal Society, for 1834-5, p. 334. See, also, Dr. Robert E. Rogers, in Amer. Journ. of the Med. Sciences, p. 282, Aug. 1836; and Mr. Ancell, Lectures on the Physiology and Pathology of the Blood, Feb. 1, 1840, p. 686.

blood, cut off a thin slice, and soak it for an hour or two in repeatedly renewed portions of distilled water; in proportion as the serum is washed away, the colour of the clot deepens, and, when scarcely any serum remains, the colour, by reflected light, is quite black. In this state, it may be exposed to the atmosphere, or a current of air may be blown upon it without any change of tint whatever; whence it would follow, that when a clot of venous blood, moistened with serum, is made florid by the air, the presence of serum is essential to the phenomenon. The serum is believed, by Dr. Stevens, to contribute to this change by means of its saline matter; for when a dark clot of blood, which oxygen fails to redden, is immersed in a pure solution of salt, it quickly acquires the crimson tint of arterial blood, and loses it again when the salt is abstracted by soaking in distilled water. The facts, detailed by Stevens, are confirmed by Mr. Prater, and by Dr. Turner, of the London University. The latter gentleman, assisted by Professor Quain, of the same institution, performed the following satisfactory experiment. He collected some perfectly florid blood from the femoral artery of a dog; and on the following day, when a firm coagulum had formed, several thin slices were cut from the clot with a sharp penknife, and the serum was removed from them by distilled water, which had just before been briskly boiled, and allowed to cool in a well-corked bottle. The water was gently poured on these slices, so that while the serum was dissolved, as little as possible of the colouring matter should be lost. After the water had been poured off, and renewed four or five times, occupying in all about an hour, the moist slices were placed in a saucer, at the side of the original clot, and both portions were shown to several medical friends, all of whom unhesitatingly pronounced the unwashed clot to have the perfect appearance of arterial blood, and the washed slices to be as perfectly venous. On restoring one of the slices to the serum, it shortly recovered its florid colour; and another slice, placed in a solution of bicarbonate of soda, instantly acquired a similar tint; - yet, as we have seen, the carbonate of soda is considered by Messrs. Gmelin, Tiedemann, and Mitscherlich, to exist in venous or black blood!

In brightening, in this way, a dark clot by a solution of salt or a bicarbonate, Dr. Turner found the colour to be often still more florid than that of arterial blood; but the colours were exactly alike when the salt was duly diluted. Dr. Turner remarks, that he is at a loss to draw any other inference from this experiment, than that the florid colour of arterial blood is not due to oxygen, but, as Dr. Stevens affirms, to the saline matter of the serum. The arterial blood, which was used, had been duly oxygenized within the body of the animal, and should not in that state have lost its

<sup>\*</sup> Experim, Inquiries in Chemical Physiology, part i. on the Blood, Lond. 1832.

b Elements of Chemistry, 5th edit. by F. Bache, p. 609, Philad. 1835.

tint by the mere removal of its serum; and he adds, the change from venous to arterial blood appears, contrary to the received doctrine, to consist of two parts essentially distinct: one is a chemical change, essential to life, accompanied by the absorption of oxygen, and the evolution of carbonic acid; the other depends on the saline matter of the blood, which gives a florid tint to the colouring matter after it has been modified by the action of ovxgen. "Such," says Dr. Turner, "appears to be a fair inference from the facts above stated; but being drawn from very limited observations, it is offered with diffidence, and requires to be confirmed or modified by future researches." But we are perhaps scarcely justified in inferring from the experiments of Stevens, Turner, and others, more than the fact, that a florid tint is communicated to blood by sea-salt, and by the neutral salts of the alkalies in general, and indeed by admixture with sugars; whilst acids render it still darker. The precise changes that occur during the arterialization of the blood in the lungs are still unknown; and if we rely on the recent experiments of Gmelin, Tiedemann, and Mitscherlich, venous blood cannot owe its colour to free carbonic acid, because none is to be met with in it; whilst the presence of the carbonates of alkalies ought to communicate the florid hue

Since Dr. Stevens first published his opinions, the subject has been farther investigated by Dr. William Gregory, and by Mr. Irvine. They introduced portions of clot, freed by washing from serum, into vessels containing pure hydrogen, nitrogen, and carbonic acid, placed over mercury. As soon as the strong saline solution came in contact with them, the colour of the clot, in all the true gases, changed from black to bright red, and the same change was found to take place in the Torricellian vacuum. On repeating these experiments with the serum of the blood, and a solution of salt in water of equal strength with the serum, no change took place until atmospheric air, or oxygen gas, was admitted. It therefore appears — as properly inferred by the late Mr. Egerton A. Jennings, from whom we have an interesting "Report on the Chemistry of the Blood as Illustrative of Pathology,"a — that though saline matter may be necessary to effect the change of colour from that of venous to that of arterial blood, still, with so dilute a saline solution, as that which exists in serum, the presence of oxygen is likewise necessary. Dr. Davyb dissents. however, from those conclusions, and is disposed to infer, from all the facts with which he is acquainted, that the colour of the blood, whether venous or arterial, that is, dark or florid, is independent of the saline matter in the serum, considered in relation to agency; and that, according to the commonly received view, oxygen is the

<sup>\*</sup> Transactions of the Provincial Medical and Surgical Association, vol. iii., Worcester and London, 1835.

b Researches, Physiological and Anatomical, Dunglison's Amer. Med. Lib. Edit. p. 96, Philad. 1840.

cause of the bright hue of the arterial fluid, and its consumption and conversion into carbonic acid, the cause of the dark hue of the venous — the saline matter being negative in regard to colour, and its chief use being, in his opinion, "to preserve the red globules from injury, prevent the solution of their colouring matter, retain their forms unchanged, and to bear them in their course

through the circulation."

The slight diminution, if it exist, in the specific gravity of arterial blood is considered, but we know not on what grounds, to depend on the transpiration, which takes place into the air-cells, and which was formerly thought to be owing to the combustion of oxygen and hydrogen. This will engage us in another place, as well as the changes produced in its capacity for heat, on which several ingenious speculations have been founded, to account for ANIMAL TEMPERATURE. The other changes are at present inexplicable, and can only be understood hereafter by minute chemical analysis, and by an accurate comparison of the two kinds of blood, — venous and arterial.

It is manifest, from the preceding detail, that our knowledge regarding the precise changes effected upon the air and the blood by respiration is by no means definite. We may, however, consider the following points established. In the first place: - the air loses a part of its oxygen, and of its azote; but this loss varies according to numerous circumstances. 2dly. It is found to have acquired carbonic acid, the quantity of which is also variable. 3dly. The bulk of the air is diminished; but the extent of this likewise differs. 4thly. The blood when it attains the left side of the heart, has a more florid colour. 5thly. This change appears to be caused by the contact of oxygen. 6thly. The blood in the lungs gets rid of a quantity of carbonic acid. 7thly. The oxygen taken in is more than necessary for the carbonic acid formed. Sthly. The constituents of the air pass directly through the coats of the pulmonary vessels, and certain portions of each are discharged or retained, according to circumstances. Lastly. A quantity of aqueous vapour, containing albumen, is discharged from the lungs.a

# c. Cutaneous Respiration, &c.

A question, again, has arisen, whether any absorption and exhalation of air, and conversion of blood from venous to arterial, take place in any other part of the body than the lungs. The reasons, urged in favour of the affirmative of this question, are;—that, in the lower classes of animals, the skin is manifestly the organ for the reception of air; that the mucous membrane of the lungs evidently absorbs air, and is simply a prolongation of the skin, resembling it in texture; and lastly, that when a limited quantity of air has been placed in contact with the skin of a living animal, it has been absorbed, and found to have a Bostock's Physiology, 3d edit. p. 361 and 377, Lond. 1836.

experienced the same changes as are effected in the lungs. Mr. Cruikshanka and Mr. Abernethy analyzed air, in which the hand or foot had been confined for a time, and detected in it a considerable quantity of carbonic acid. Jurine, having placed his arm in a cylinder hermetically closed, found, after it had remained there two hours, that oxygen had disappeared, and that 0.08 of carbonic acid had been formed. These results were confirmed by Gattoni.e On the other hand, Drs. Priestley, Klapp, and Gordon, could never perceive the least change in the air under such circumstances. Perhaps in these, as in all cases where the respectability of testimony is equal, the positive should be adopted rather than the negative. It is probable, however, that absorption is effected with difficulty; and that the cuticle, as we have elsewhere shown, is placed on the outer surface to obviate the bad effects which would be induced by heterogeneous gaseous, miasmatic, or other absorption. We have seen that some of the deleterious gases, as sulphuretted hydrogen, are most powerfully penetrant, and, if they could enter the surface of the body with readiness, unfortunate results might supervene. In those parts where the cuticle is extremely delicate, as in the lips, some conversion of the venous blood into arterial may be effected, and this may be a great cause of their florid colour. According to this view, the arterialization of the blood occurs in the lungs chiefly, owing to their formation being so admirably adapted to the purpose, and it is not effected in other parts, because their arrangement is unfavourable for such result.g

### d. Effects of the Section of the Cerebral Nerves on Respiration.

It remains for us to inquire into the effect produced on the lungs by the cerebral nerves distributed to them, — or rather, into what is the effect of depriving the respiratory organs of their nervous influence from the brain. The only cerebral or encephalic nerves, distributed to them, are the pneumogastric or eighth pair of Willis, which, we have seen, are sent, as their name imports, to both the lungs and the stomach. The section of these nerves early suggested itself to physiologists, but it is only in recent times that the phenomena resulting from it have been clearly comprehended. The operation appears to have been performed as long ago as the time of Rufus of Ephesus, and was afterwards repeated by Chirac,

<sup>\*</sup> Experiments on the Insensible Perspiration, &c., Lond. 1795. See, also, Edwards, Sur l'Influence des Agens Physiques, p. 12; or Hodgkin and Fisher's translation.

b Surgical and Physiological Essays, part ii. p. 115, Lond. 1793.

C Dict. des Sciences Médicales, art. Peau.

Experiments and Observations on different kinds of Air, ii. 193, and v. 100, Lond.
 1774.
 Inaugural Essay on Cuticular Absorption, p. 24, Philad. 1805.

f Ellis's Inquiry into the Changes of Atmospheric Air, &c., p. 355, Edinb. 1837. See, also, Madden's Experimental Inquiry on Cutaneous Absorption, p. 130, Lond. 1838.

Müller's Handbuch, u. s. w., Baly's translation, p. 334, Lond. 1838.

VOL. II. - 6

Bohn, Duverney, Vieussens, Schräder, Valsalva, Morg agni, Haller, and numerous other distinguished physiologists.<sup>a</sup> It is chiefly, however, in very recent times, and especially by the labours of Dupuytren, Dumas, De Blainville, Provençal, Legallois, Magendie, Breschet, Hastings, Broughton, Brodie, Wilson Philip, and Dr. John Reid, that the precise effects upon the respiratory and

digestive functions have been appreciated.

When the nerves are divided in a living animal, on both sides at once, the animal dies more or less promptly; at times, immediately after their division, but sometimes it lives for a few days; Magendie says never beyond three or four. The effects produced upon the voice, by their division above the origin of the recurrents, have been referred to under another head (vol. i., p. 421). Such division, however, does not simply implicate the larynx, but necessarily affects the lungs, as well as the stomach. As regards the larynx, precisely the same results, according to Magendie, are produced by dividing the trunk of the pneumogastric above the origin of the recurrents, as by the division of the recurrents themselves: the muscles, whose function it is to dilate the glottis, are paralyzed; and, consequently, during inspiration, no dilatation takes place; whilst the constrictors, which receive their nerves from the superior laryngeal, preserve all their action, and close the glottis, at times so completely, that the animal dies immediately from suffocation. But if the division of these nerves should not induce instant death in this manner, a series of symptoms follows, considerably alike in all cases, which go on until the death of the animal. These phenomena are the following: - respiration is, at first, difficult; the inspiratory movements are more extensive and rapid, and the animal's attention appears to be particularly directed to them; the locomotive movements are less frequent, and evidently fatigue; frequently, the animal remains entirely at rest; the formation of arterial blood is not prevented at first, but soon, on the second day for instance, the difficulty of breathing augments, and the inspiratory efforts become gradually greater. The arterial blood has now no longer the vermilion hue which is proper to it. It is darker than it ought to be. Its temperature falls. Respiration requires the exertion of all the respiratory powers. At length, the arterial blood is almost like the venous, and the arteries contain but little of it; the body gradually becomes cold, and the animal dies. On opening the chest, the air-cells, the bronchi, and frequently even the trachea, are found filled by a frothy fluid, which is sometimes bloody; the substance of the lung is tumid; the divisions and even the trunk of the pulmonary artery are greatly distended with dark, almost black, blood; and extensive effusions of serum and even of blood are found in the parenchyma of the lungs. Experiments have, likewise, shown that, in proportion as these symptoms appeared, the animals consumed less and

a Haller. Element. Physiologiæ.

b Précis, &c. 2de édit. ii. 355.

less oxygen, and gave off a progressively diminishing amount of carbonic acid.a

From the phenomena that occur after the section of these nerves on both sides, it would seem to follow, that the first effect is exerted upon the tissue of the lungs, which, being deprived of the nervous influence they receive from the brain, are no longer capable of exerting their ordinary elasticity or muscularity, which soever it may be. Respiration, consequently, becomes difficult; the blood no longer circulates freely through the capillary vessels of the lungs; the consequence of this is, that transudation of its serous portions, and occasionally effusion of blood, owing to rupture of small vessels, takes place, filling the air-cells more or less; until, ultimately, all communication is prevented between the inspired air and the bloodvessels of the lungs, and the conversion of the venous into arterial blood is completely precluded. Death is then the inevitable and immediate consequence. The division of the nerve on one side affects merely the lung of the corresponding side: life can be continued by the action of one lung only. It is, indeed, a matter of astonishment how long some individuals have lived when the lungs have been almost wholly obstructed. Every morbid anatomist has had repeated opportunities for observing, that, in cases of pulmonary consumption, for a length of time prior to dissolution, the process of respiration must have been wholly carried on by a very small portion of lung.

From his experiments on this subject, Sir Astley Cooper infers, that the pneumogastric nerve is most important; — 1st, in assisting in the support of the function of the lungs, by contributing to the changing of the venous into arterial blood; 2dly, in being necessary to the act of swallowing; and 3dly, in being very essential to the digestive process; whilst Dr. John Reid is of opinion, that the pulmonary branches seem to be the nerves, chiefly concerned in transmitting to the medulla oblongata the impressions which excite respiratory movements, and are thus principally afferent nerves; but it is possible, he adds, that they contain motor filaments also.<sup>b</sup>

The experiments of Dr. Wilson Philip<sup>e</sup> and others moreover show, — what has been more than once inculcated, — the great similarity between the nervous and galvanic fluids. When the state of dyspnæa was induced by the division of the pneumogastric nerves, the galvanic current was passed from one divided extremity to the other, and, in numerous cases, the dyspnæa entirely ceased. The results of these experiments induced him to try the effect of galvanism in cases of asthma. By transmitting

<sup>\*</sup> See Sir Astley Cooper, in Guy's Hospital Reports, part i. 468, Lond. 1836; also Dr. J. Reid, Edinb. Med. and Surg. Journ. p. 163, for Jan. 1838.

Edinb. Med. and Surg. Journ. April, 1839.
 Experimental Inquiry into the Laws of the Vital Functions, &c. 2d edit. p. 223,
 Lond. 1818; also, Journal of Science and Arts, viii. 72; and art. Galvanism, in Ure's Dict. of Chemistry, 2d edit. Lond. 1823.

its influence from the nape of the neck to the pit of the stomach, he gave decided relief in every one of twenty-two cases, four of which occurred in private practice, and eighteen in the Worcester

Infirmary.

Sir A. Coopera instituted similar experiments on the phrenic nerves. As soon as these were tied, the most determined asthma was produced; breathing went on by means of the intercostal muscles; the chest was elevated to the utmost by them; and in expiration the chest was as remarkably drawn in. The animals did not live an hour; but they did not die suddenly, as they do from pressure on the carotid and vertebral arteries. The lungs appeared healthy, but the chest contained more than its natural exhalation. He also tied the great sympathetic; but little effect was produced: the animal's heart appeared to beat more quickly and feebly than usual. The animal was kept seven days, and one nerve was ulcerated through, and the other nearly so at the situation of the ligatures. No particular alteration of any organ was observed, on examination. Lastly, Sir Astley tied all three nerves on each side, the pneumogastric, phrenic, and grand sympathetic: the animal lived little more than a quarter of an hour, and died of dyspnæa.

From these experiments, he infers, that the sudden death, which he found in his experiments to take place from pressure on the sides of the neck, cannot be attributed to any injury of the nerves, but to an impediment to the due supply of blood to the great

centre of nervous influence.

The centre of the respiratory movements is now considered to be the upper part of the medulla oblongata. Into it the pneumogastric nerves, which appear to be the chief excitors of respiration, may be traced, and from it the different motor or efferent nerves, which Sir Charles Bell has grouped together in his respiratory system, proceed either directly or indirectly. Of these, the most important is the phrenic.<sup>b</sup>

# e. Respiration of Animals.

In concluding the subject of respiration, we may briefly advert to the different modes in which the process is effected in the classes of animals, and especially in birds, the respiratory organs of which constitute one of the most singular structures of the animal economy. The lungs themselves, — as in the marginal figure of the lungs, &c., of the ostrich, (Fig. 152,) — are comparatively small, and are adherent to the chest, — where they seem to be placed in the intervals of the ribs. They are covered by the pleura only on their under surface, so that they are, in fact, on the outside of the cavity of the chest. A great part of the thorax, as well as of the abdomen, is occupied by membranous air-cells, into which the lungs open by considerable apertures. Besides these 2 Op. cit. p. 475.

Dr. Carpenter, Human Physiology, p. 138, Lond. 1842.

cells, a considerable portion of the skeleton forms receptacles for

air, in many birds; and if we break a long bone of a bird of flight, and blow into it, the body of the bird being immersed in water, bubbles of air will escape from the bill. The object, of course, of all this, is to render the body light, and thus to facilitate its motions. Hence the largest and most numerous bony cells are found in such birds as have the highest and most rapid flight, as the eagle. The barrels of the quills are likewise hollow, and can be filled with air, or emptied at pleasure. In addition to the uses just mentioned, these receptacles of air diminish the necessity of breathing so frequently, in the rapid and long-continued motions of several birds, and in the great vocal exertions of singing birds.

In fishes, in the place of lungs we find branchiæ or gills, which are placed behind the head on each side, and have a

Thoracic and Abdominal Viscera of the Ostrick.

a. Heart, lodged in one great air-cell. b. The stomach. c. The intestines, surrounded by large air-cells. d. The trachea dividing into bronchi. c. c. The lungs. 1, 2, 3.f. f. Other great air-cells, communicating with other cells and with the lungs. g, g. The openings by which such communication is made.

moveable gill-cover. By the throat, which is connected with these organs the water is conveyed to the gills, and distributed through them: by this means, the air, contained in the water, which, according to Biot, Von Humboldt, and Provençal, Configliachi, and Thomson, is richer in oxygen than that of the atmosphere, having from 29 to 32 parts in the 100, instead of 20 or 21, comes in contact with the blood circulating through the gills. The water is afterwards discharged through the branchial openings, — aperturæ branchiales, — and consequently, they do not expire along the same channel as they inspire.

Lastly, in the insect tribe, — in the white-blooded animal, — we find the function of respiration effected altogether by the surface of the body; at least, so far as regards the reception of air, which passes into the body through apertures termed stigmata,

\* Mémoir, de la Société d'Arcueil, i. 252, and ii. 400; Annals of Philosophy, v. 40; and Richerand's Elèmens de Physiologie, 18ème édit. par M. Bérard, ainé, p. 141, Bruxelles, 1837.

<sup>b</sup> Dr. Thomson found that 100 cubic inches of the water of the river Clyde contained 3.113 inches of air; and that the air contained 29 per cent. of oxygen. Edinb. New Philosoph. Journal, xxi. 370, Edinb. 1836.

the external termination of trachew or air-tubes, whose office it is

to convey the air to different parts of the system.

In all these cases, we find precisely the same changes effected upon the inspired air, and especially, that oxygen and azote have disappeared, and that carbonic acid is contained in nearly equal bulk with the azote in the residuary air.<sup>a</sup>

### CHAPTER IV.

#### CIRCULATION.

THE next function to be considered is that by which the products of the various absorptions, converted into arterial blood in the lungs, are distributed to every part of the body, — a function of the most important character to the physiologist, and the pathologist, and without a knowledge of which it is impossible for the latter

to comprehend the doctrine of disease.


Assuming the heart to be the great central organ of the function, the circulatory fluid must set out from it, be distributed through the lungs, undergo aëration there, be sent to the opposite side of the heart, whence it is distributed to every part of the system by efferent vessels, and be thence returned by the veins or afferent vessels to the right side, from which it set out, — thus performing a complete circuit.

The lower classes of animals differ essentially, as we shall find hereafter, in their organs of circulation: whilst in some, the apparatus appears to be confounded with the digestive; in others, the blood is propelled without any great central organ; and in others, again, the heart is but a single organ. In man, and in the upper classes of animals, the heart is double; — consisting of two sides, or really of two hearts, separated from each other by a septum.

Fig. 153. In the dugong, the two ventricles are almost entirely detached from

each other.b

As all the blood of the body has to be emptied into this central organ, and to be subsequently sent from it, and as its flow is continuous, two cavities are required in each heart,—the one to receive the blood, the other to propel it. This last contracts and dilates alternately. The cavity or chamber of each heart, which receives the blood, is called auricle, and the vessels that transport it thither are the veins; the cavity by which the blood is pro-



Heart of the Dugong.

D. Right auricle. E. Right ventricle.
L. Left auricle. L. Left ventricle. F.
Pulmonary artery. A. Aorta.

Roget's Animal and Vegetable Physiology, ii. 221, Amer. Edit.; and Tiedemann, Traité Complet de Physiologie de l'Homme, par Jourdan, p. 302, Paris, 1831.


b Roget, op. cit., ii. 200, Philad. 1836.

jected forwards is called ventricle, and the vessels, along which the blood is sent, are the arteries. One of these hearts is entirely appropriated to circulation of venous blood, and hence has been called the venous heart, — also the right or anterior heart, from its situation, — and the pulmonary from the pulmonary artery arising from it. The other is for the circulation of arterial blood, and is hence called the arterial heart, also the left or posterior, from its situation, and the aortic heart, because the aorta arises from it. In Fig. 154, the two hearts are separated from each other, and shown to be distinct organs in the adult, although in the subject they seem to form but one. Between the two, after birth, there is

not the slightest communication; and consequently, all the blood, with the exception of that sent out by the coronary artery, and returned by the coronary vein, which has to attain the left side of the heart, must make the circuit through the lungs.

The whole of the vessels communicating with the right heart, contain venous blood; those of the left side, arterial blood.

If we consider the heart to be the centre, two circulations are accomplished before the blood, setting out from one side of the heart, performs the whole cir-



The Right and Left Hearts, separated.

a, a. Venæ cavæ ascendens, and descendens. b. Right auricle. c, c. Right ventricle. d. Pulmonary artery. c. Pulmonary veins. f. Left auricle. g. Left ventricle. h, h. Aorta. The arrows indicate the course of the blood.

cuit. One of these consists in the transmission of the blood from the right side of the heart, through the lungs, to the left; the other in its transmission from the left side, along the arteries, and, by means of the veins, back to the right side. The former of these is called the lesser or pulmonic, the latter the greater or systemic circulation. The organs, by which these are effected, will require a more detailed examination.

#### 1. ANATOMY OF THE CIRCULATORY ORGANS.


The circulatory apparatus is composed of the organs, by which the blood is put in motion, and along which it passes during its circuit.

### a. Heart.

To simplify the consideration of the subject, we shall consider

the heart double; and that each system of circulation is composed of a heart; of arteries, through which the blood is sent from the heart; and of veins, by which the blood is returned to it. At the minute termination of each of these is the capillary system. We shall first describe the central organ, as forming two distinct hearts; and afterwards the two united.

The pulmonic, right, or anterior heart, called also the heart of black blood, is composed of an auricle and a ventricle. The auricle,




Pulmonic Heart.

A. Right auricle with its venæ cavæ. B. Right ventricle. C. Pulmonary artery.

so termed from some resemblance to an ear, is situate at the base of the organ, and receives the whole of the blood returning from various parts of the body by three veins; the two venæ cavæ, and the coro-The vena cava descendens terminates in the auricle in the direction of the aperture by which the auricle communicates with the ventricle. The vena cava ascendens. the termination of which is directed more backwards, has the remains of a valve which is much larger in the fætus, called the valve of Eustachius. The third vein is the cardiac or coronary; it returns the blood from the heart which has been carried thither by the coronary artery. In the septum between the right and

left auricle, there is a superficial depression, about the size of the point of the finger, which is the vestige of the foramen ovale,—an important part of the circulatory apparatus of the fœtus. The opening, through which the auricle projects its blood into the ven-



Section of the Pulmonic Heart.

A. Right auricle. B. Right ventricle. C. Pulmonary artery.

tricle, is situate downwards and forwards, as seen in Fig. 156. The inner surface of the proper auricle, or that which more particularly resembles the ear of a quadruped, - the remainder being sometimes called the sinus venosus or sinus venarum cavarum, - is distinguished by having a number of fleshy pillars in it, which, from their supposed resemblance to the teeth of a comb, are called musculi pectinati. They are mere varieties, however, of the columnæ carneæ of the ventricles.

The right ventricle or pulmonary ventricle is situate in the anterior part of the heart; the base and apex corresponding to those of the heart. Its cavity is generally greater than that of the left side, and its parietes not so thick, owing to their merely having to force the blood through the lungs. It communicates with the auricle by the auriculo-ventricular opening - the ostium venosum; and the only other opening into it is that which communicates with the interior of the pulmonary artery. The opening, between the auricle and ventricle, is furnished with a tripartite valve, called tricuspid or triglochin; and the pulmonary artery has three others, called sigmoid or semilunar. From the whole edge of the tricuspid valve, next the apex of the heart, small, round, tendinous cords, called chordæ tendineæ, are sent off, which are fixed, as represented in Fig. 156, to the extremities of a few strong columnæ carnew. These tendinous cords are of such a length as to allow the valve to be laid against the sides of the ventricle, in the dilated state of that organ, and to admit of its being pushed back by the blood, until a nearly complete septum is formed, during the contraction of the ventricle. The semilunar or sigmoid valves are three in number, situate around the artery. When these fall together, there must necessarily be a space left between them. To obviate the inconvenience, that would result from the existence of such a free space, a small granular body is attached to the middle of the margin of each valve; and, these coming together, as at

A, Fig. 157, when the valves are shut down, complete the diaphragm, and prevent any blood from passing back to the heart. These small bodies are termed, from their reputed discoverer, corpuscula Arantii, and also corpuscula Morgagnii; or, from their resemblance to the seed of the sesamum, corpuscula sesamoidea. The valves, when shut, are concave towards the lungs, and convex towards the ventricle. Immediately above them the artery bulges out, forming three sacculi





Semilunar Valves closed.

or sinuses, called sinuses of Valsalva. These are often said to be partly formed by the pressure of the blood upon the sides of the vessels. The structure is doubtless ordained, and is admirably adapted for a specific purpose, namely, to allow the free edges of the valves to be readily caught by the refluent blood, and thus to facilitate their closure. Within the right ventricle, and especially towards the apex of the heart, many strong eminences are seen, which are called columnæ carneæ (Fig. 156). These run in different directions, but the strongest of them longitudinally with respect to the ventricle. They are of various sizes, and form a beautifully reticulated texture. Their chief use probably is, to strengthen the ventricle and prevent it from being over-distended; in addition to which they may tend to mix the different products of absorption.

The corporeal, left, aortic or systemic heart, - called also the heart of red blood, - has likewise an auricle and a ventricle. The left auricle is considerably thicker and stronger but smaller than the right; and is like wise divided into sinus venosus and proper auricle, which form a common cavity. The columns in the latter are like those of the right auricle, but less distinct. From the under part of the auricle, a circular passage, termed ostium arteriosum or auricular orifice leads to the posterior part of the base of the cavity of the left ventricle. The left auricle receives the blood from the pulmonary veins. The left or aortic ventricle is situate at the posterior and left part of the heart. Its sides are three times thicker and stronger than those of the right ventricle, to permit the much greater force which it has to exert; for, whilst the right ventricle merely sends its blood to the lungs, the left ventricle transmits it to every part of the body. It is narrower and rounder, but considerably longer, than the right ventricle, and forms the apex of the heart. The internal surface of this ventricle has the same general appearance as the other, but differs from it in having its columnæ carneæ larger, more numerous, firmer, and stronger. In the aperture of communication with the corresponding auricle, there is here, as in the opposite side of the heart, a ring or zone, from which a valve, essentially like the tricuspid, goes off. It is stronger, however, and divided into two principal portions only: the chordæ tendineæ are also stronger and more numerous. This valve has been termed mitral, from some supposed resemblance to a bishop's mitre. At the fore and right side of the mitral valve, and behind the commencement of the pulmonary artery, a round opening exists, which is the mouth of the aorta. Here are three semilunar valves, with their corpuscula Arantii, exactly like those of the pulmonary artery, but a little stronger; and, on the outer side of the semilunar valves, are the sinuses of Valsalva, a little more prominent than those of the pulmonary artery.

The structure of the two hearts is the same. A serous membrane covers both, which is an extension of the inner membrane

of the pericardium.

The substance of the heart is essentially muscular. The fibres run in different directions, longitudinally and transversely, but most of them obliquely. Many pass over the point, from one heart to the other, and all are so involved as to render it difficult to unravel them. The cavities are lined by a thin membrane, the endocardium, which differs somewhat in the two hearts; — being in one a prolongation of the inner coat of the aorta, and in the other of the venæ cavæ. On this account, the inner coat of the left heart is but slightly extensible, more easily ruptured, and considerably disposed to ossify; that of the right heart, on the other hand, is very extensible, not readily ruptured, and but little liable to ossify. M. Deschamps has described a membrane which is situate between the endocardium and the cellular tissue that lines

<sup>&</sup>lt;sup>a</sup> Gazette Médicale de Paris, No. 10, and Encyclographie des Sciences Médicales, Avril, 1840, p. 281.

the muscular structure at its inner surface, and belongs essentially to the elastic fibrous tissue. The tissue of the heart is supplied with blood by the cardiac or coronary arteries—the first division of the aorta; and their blood is conveyed back to the right auricle by the coronary veins. The nerves, which follow the ramifications of the coronary arteries, proceed chiefly from a plexus, formed

by the pneumogastric nerves, and great sympathetic.

In both hearts, the auricles are much thinner and more capacious than the ventricles; but they are themselves much alike in structure and size. The observation, that the right ventricle is larger than the left, is as old as Hippocrates, and has been attempted to be accounted for in various ways. Some have ascribed it to original conformation; others to the blood being cooled in its passage through the lung, and therefore occupying a smaller space when it reaches the left side of the heart. Hallera and Meckelb assert that it is dependent upon the kind of death; that if the right ventricle be usually more capacious, it is owing to the lung being one of the organs that yields first, thus occasioning accumulation of blood in the right cavities of the heart; and they state that they succeeded, in their experiments, in rendering either one or the other of the ventricles more capacious, according as the cause of death arrested first the circulation in the lung or in the aorta; but the experiments of Legallois, and Seiler, especially of the former, - with mercury poured into the cavities, on dogs, cats, Guineapigs, rabbits, in the adult, the child, and the still-born fœtus, have shown, that, except in the fœtus, the right ventricle is more capacious, whether death have been produced by suffocation, in which the blood is accumulated in the right side of the heart, or by hemorrhage; and Legalloisf thinks that the difference is owing to the left ventricle being more muscular, and, therefore, returning more upon itself.5 The capacity of each of the ventricles in the full sized heart, may be estimated at about two fluid ounces.

The two hearts, united together by a median septum, form, then, one organ, which is situate in the middle of the chest, (see Fig. 147,) between the lungs, and consequently in the most fixed part of the thorax. Figure 158 is modified from one carefully made from nature by Dr. Pennock.<sup>h</sup> It represents the normal po-

sition of the heart and great vessels.

According to Carus, the weight of the heart compared with that of the body is as 1 to 160. M. J. Weber found the proportion, in one case, as 1 to 150; Dr. Clendinningk that of the male 1 to 160;

\* Element. Physiol., iv. 3. 3.

Handbuch der Menschlichen Anatomie, Halle, 1817, s. 46; or the translation from the French version, by Dr. Doane, Philad. 1832.
 Clendinning, Report to the British Association, Lond. Med. Gaz. Nov. 13, 1840.

d Dict. des Sciences Médicales, v. 440.

e Art. Herz, in Anat. Physiol. Real Wörterb. iv. 32, Leipz. 1821.


CEuvres, Paris, 1824. & Burdach, Physiologie, u. s. w., iv. 214.

h Medical Examiner, April 4, 1840.

Introduction to Comp. Anat. translated by R. T. Gore, Lond. 1827.

Hildebrandt's Handbuch der Anatomie, von E. H. Weber, Braunschweig, 1831. Band. iii. s. 125. k Journal of the Statistical Society of London, July, 1838.

that of the female 1 to 150; and Laënnec considered the organ to be of a healthy size when equal to the fist of the individual. Cruveilhier estimates the mean weight at six or seven ounces. Bouillauda weighed the hearts of thirteen subjects, in whom, from the general habit, the previous state of health, and the mode of death, there was every reason to believe they were in the natural state. The mean was eight ounces and three drachms. From all his data he is led to fix the mean weight of the heart, in the adult, from the 25th to the 60th year, at from 8 to 9 ounces.



S. Outline of Sternum. C. C. Clavicles. 1, 2, 3, 4, 5, 6, &c. The ribs. 1', 2', 3', 4', 5', 6', &c. Cartilages of the ribs. 4''. Right and left nipples. a. Right ventricle. b. Left ventricle. c. Septum between the ventricles. d. Right auricle. e. Left auricle. f. The aorta. f'. Needle passing through aortic valves. g. Pulmonary artery. g'. Needle passing through valves of pulmonary artery. h. Vena cava descendens. i. Line of direction of mitral valve; the dotted portion posterior to the right ventricle. i'. Needle passed into the mitral valve at its extreme left. k. Line of tricuspid valve. o. Trachea.

Dr. Clendinning carefully examined nearly four hundred hearts of persons of both sexes, and of all ages above puberty. The result was about nine ounces avoirdupois, much less than that obtained recently by Dr. John Reid, who found the average weight of the male heart — of 89 weighed — to be 11 oz. and 1 dr.; and of the female heart — of 53 weighed — to be 9 oz. and ½ dr.

<sup>a</sup> Traité Clinique des Maladies du Cœur, &c. Paris, 1835,

See, also, Dr. Gross, Elements of Pathological Anatomy, ii. 124, Boston, 1839.
 Lond. and Edinb. Monthly Journ. of Med. Science, April, 1843, p. 322.

The dimensions of the heart, according to Lobstein and Bouillaud, are as follows:—Weight of heart, 9 to 10 ounces; length from base to apex, 5 inches 6 lines; breadth at the base, 3 inches; thickness of walls of left ventricle, 7 lines; do. at a finger's breadth above the apex, 4 lines; thickness of walls of right ventricle, 2½ lines; do. at apex, ½ a line; thickness of right auricle, 1 line; do. of left auricle, ½ a line. M. Bizot<sup>a</sup> has given the following measurements, taking the average of males from 16 to 89 years.

					Base.	Middle.	Apex.
Left ventricle,	1	-	-	11129	44 lines	51	33
Right ventricle.		-	100	-	115	13	11

In the female, the average thickness is something less.

Recently, Dr. Rankingb has published the results of measurements, evidently made with accuracy, of upwards of 100 hearts, care being taken to exclude all those that exhibited any trace of organic change. The following are the mean admeasurements. Of 15 male hearts, the mean circumference was 927 ths inches; of 17 female hearts, 816 ths inches. The mean length of the male heart was 416 ths inches; of the female, 436 ths. The mean thickness of the left ventricle in the male was 27ths of an inch; in the female, 23ths; of the right ventricle, in the male, sths; in the female, feths. The septum ventriculorum has, in the male, a mean thickness of  $\frac{2}{3}$ ths of an inch; in the female,  $\frac{1}{3}$ ths. The aortic orifice, in the male, had a mean circumference of 234ths inches; the right auriculo-ventricular orifice, 435 ths inches; the left auriculo-ventricular orifice, 345 ths inches. The corresponding parts of the female were relatively less. Dr. Ranking infers, that the heart of the male is larger than that of the female, - that the length of the healthy heart to its circumference is rather less than 1 to 2 — that the thickness of the parietes of the right ventricle to the left is as 1 to 3 nearly: — that the pulmonary artery is slightly wider than the aorta; and, lastly, that the right auriculo-ventricular opening is considerably larger than the left.

It need scarcely be said, that the weight and dimensions of the organ must vary according to the age, sex, &c. of the individual. M. Bizote found, that the influence of stature on its size was slight; and not such as might have been expected à priori; as in individuals of the male sex above sixty inches, and in females above fifty-five inches in height, the mean dimensions of the organ, especially its breadth, were less than in persons of lower stature. He found the width of the shoulders to furnish a better proportionate standard of its measures, — the distance between the acro-

<sup>&</sup>lt;sup>a</sup> For an account of the results of M. Bizot's researches, to ascertain the dimensions of the heart and arteries, see Memoires de la Société Médicale d'Observation, Paris, 1837; and Hope on the Diseases of the Heart, Amer. Edit., by Dr. Pennock, p. 234, Philad. 1842.

b London Medical Gazette, No. xxiv. 1842.

Mémoires de la Société Médicale d'Observation de Paris, tom. 1ère, Paris, 1836.
 VOL. II. — 7

mial point of the clavicles, and the length and breadth of the heart

increasing in tolerably regular ratio.

The heart is surrounded by its proper capsule, called the pericardium - a fibro-serous membrane, which is composed of two layers. The outermost of these is fibrous, semitransparent, and inelastic; strongly resembling the dura mater in its texture. Its thickness is greater at the sides than below, where it rests upon the diaphragm; or than above, where it passes along the great vessels which communicate with the heart. The inner layer is of a serous character, and lines the outer, giving the polish to its cardiac surface; it is then reflected over the heart, and adheres to it by cellular substance. Like other serous membranes, it secretes a fluid, which is termed liquor pericardii, to lubricate the surface of the heart. This fluid is always found in greater or less quantity after death; and a question has arisen regarding the amount that must be considered morbid. This must obviously vary according to circumstances. It seldom, however, in the healthy condition, is above a tea-spoonful. When its quantity is augmented, along with inflammation of the membrane, the disease hydropericarditis exists. The great use of the pericardium is probably to keep the heart constantly moist by the exhalation effected from it; and, also, to restrain the movements of the heart, which, under the influence of the emotions, sometimes leaps inordinately. If the pericardium be divided in a living animal, the heart is found to bound, as it were, from its ordinary position; and hence the expression — "leaping of the heart," during emotion is physiologically accurate.

## b. Arteries.

The arteries are solid, elastic tubes, which arise, by a single trunk, from the ventricle of each heart, and gradually divide and subdivide, until they are lost in the capillary system. The large artery, which arises from the left ventricle, and conducts the blood to every part of the body, - even to the lungs, so far as regards their nutrition, - is, as we have seen, the aorta; and that, which arises from the right ventricle and conveys the venous blood to the lungs, for aëration, is the pulmonary artery. Neither the one nor the other is a continuation of the proper tissue of the ventricles; the inner membrane is alone continuous, the muscular structure of the heart being united to the fibrous coat of the arteries, by means of an intermediate fibrous tissue. The aorta, as soon as it quits the left ventricle, passes beneath the pulmonary artery, is entirely concealed by it, and ascends to form a curvature with the convexity upwards, the summit of which rises to within threequarters of an inch or an inch of the superior edge of the sternum. This great curvature is called the cross or arch of the aorta. The vessel then passes downwards, from the top of the thorax to nearly as far as the sacrum, where it divides into two trunks, one of which proceeds to each lower extremity. In the whole of this course, it lies close to the spine, and gives off the various branches that convey arterial blood to the different parts of the body. Of the immense multitude of these ramifications, an idea may be formed, when we reflect, that the finest pointed needle cannot be run into any part of the surface of the body, without blood, — probably both arterial and venous, — flowing. The larger arteries are situate deeply, and are thus remote from external injury. They communicate freely with each other, and their anastomoses are more frequent as the arteries become smaller and farther from the heart. At their final terminations, they communi-

cate with the veins and the lymphatics.

It has been a common, but erroneous belief, that the branches of the aorta, when taken collectively, are of much greater capacity than the parent trunk, and that this inequality goes on augmenting; so that the ultimate divisions of an artery are of much greater capacity than the trunk of the vessel. Hence, the arterial system has been considered to represent, in the aggregate, a cone, whose apex is at the heart, and the base in the organs; but as all the minute arterial ramifications are not visible, it is obviously impracticable to discover the ratio between their united capacity and that of the aorta at its origin; yet the problem has been attempted. Keil, by experiments made upon an injected subject, considered it to be as 44507 to 1. J. C. A. Helvetius and Sylva as 500 to 1. Senac estimated, not their capacities but their diameters, and he conceived the ratio of these to be as 118,490 to 90,000; and George Martine affirmed, that the calibre of a parent arterial trunk is equal to the cube root of the united diameters of the branches.a It will be shown, however, hereafter, from the observations of M. Poiseuille and of Mr. Ferneley, that the notion of the much greater capacity of the branches than of the parent trunk is a fallacy. This subject will be referred to hereafter.

The pulmonary artery strongly resembles the aorta. Its distribution has been already described as a part of the respiratory

organs (page 16).

The arteries are composed of different coats in superposition, respecting the number of which anatomists have not been entirely of accord. Some have admitted six, others five, others four, but at the present day, three only are perhaps generally received; — first, an external or cellular, called also nervous, and cartilaginous by Vesalius, and tendinous by Heister, which is formed of condensed cellular substance, and has considerable strength and elasticity, so that if a ligature be applied tightly round the vessel, the middle and internal coats may be completely cut through, whilst the outer coat may remain entire. Scarpa is not disposed to admit this as one of the coats. He considers that it is only an exterior

\* Haller, Elementa Physiologiæ, tom. iv.

b See, on the Histology of the Arteries, Henle, and Mr. Paget, Brit. and For. Med. Rev. July, 1842, p. 284; and Mandl, Manuel d'Anatomie générale, p. 187, Paris, 1843.

envelope, to retain the vessel in situ. The next coat is the middle, muscular or proper coat, the character of which has been the subject of much discussion. It is composed of yellow, circular fibres, which do not appear individually to pass entirely round the vessel. This coat was, at one time, almost universally believed to be muscular. Such was the opinion of Hunter; a and hence the muscularity of the arteries was regarded as an agent in the circulation. Careful examination does not, however, exhibit the characters of ordinary muscular tissue. The latter is soft, extensible, contractile, and of a red colour: the arterial tunic is firm. solid, elastic, easily ruptured, and of a yellow colour, resembling not a little the muscular coat of the intestines. Henleb advances the opinion, that its structure is intermediate between cellular and muscular tissue; its microscopic elements being broad and very flat, slightly granulated fibres or bands, which lie in rings around the internal membrane, and are about 0.003 lines in diameter. These with a system or network of dark streaks constitute the middle coat of the artery. Nysten, Magendie, and Müllere applied the galvanic stimulus to it, but without effect; and it is known, that this is the most sensible test of irritability. The middle coat appears to be a tissue of a peculiar character, the base of which is formed by the tissu jaune, or yellow tissue of the later comparative anatomists; itself composed essentially of gelatin.f It is proper to remark, that the heart also seems equally unsusceptible of the galvanic stimulus; or at least is not affected by it like the voluntary muscles. In the cases of two executed criminals, which the author had an opportunity of observing, although all degrees of galvanism were applied, half an hour after the drop fell, no motion whatever was perceptible; yet the voluntary muscles contracted, and continued to do so for an hour and a half after execution. The same fact is recorded in the galvanic experiments of Dr. Ure, detailed in another part of this work, (vol. i., p. 368,) and is attested by Bichat, Treviranus and others. Humboldt, Pfaff, J. F. Meckel, Wedemeyer, and J. Müller, however, affirm the contrary. The last observer states, that with a single pair of plates he excited contractions not only in a frog's heart, which had ceased to beat, but also in that of a dog, under similar circumstances. Into the subject of the cause of the heart's action, we shall, however, inquire presently. Müllerh suggests, that in the capability to contract under the influence of cold

a On the Blood, p. 124, Lond. 1794.

e Handbuch der Physiologie, Baly's translation, p. 205, Lond. 1838.

s Handbuch, u. s. w. translation, p. 205.

b Casper's Wochenschrift, May 23, 1840, and Brit. and For. Med. Rev. Oct. 1840, p. 551.

Recherches de Physiologie, &c. p. 325, Paris, 1811.

d Précis, 2d edit. ii. 387, Paris, 1825.

f Eulenberg, De Telâ Elasticâ Diss. Berol. 1836, cited by Mandl, Manuel d'Anatomie générale, p. 181, Paris, 1843.

h Archiv. für 1836, and Lond. Med. Gaz. May, 1837.

. .. .. .. . .

as exhibited in the experiments of Schwann, referred to hereafter, the contractile tissue of the arteries resembles that of the dartos, and that which is found in many parts of the skin, as about the nipple and follicles, although the physical characters of the latter are so different from elastic tissue. The third or inner coat is smooth and polished, and is a continuation of the membrane which lines the ventricles. It has an epithelial lining, resembles the serous membranes, and is lubricated by a kind of serous exhalation.<sup>a</sup>

The arteries receive the constituents that belong to every living part, - arteries, veins, lymphatics, and nerves. These arteries proceed not from the vessels themselves, which they nourish, but from adjacent trunks, as we have remarked of the vasa vasorum, to which class they really belong. The nerves proceed from the great sympathetic, form plexuses around the vessels, and accompany them through all their ramifications. By some anatomists, the arteries of the head, neck, thorax and abdomen, are conceived to be supplied from the great sympathetic, whilst those of the extremities are supplied from the nerves of the spinal marrow. It is probable, however, that more accurate discrimination might trace the dispersion of the twigs of the great nervous system of involuntary motion on all these vessels. The organization of the arteries renders them very tough and extremely elastic, both of which qualities are necessary to enable them to withstand the impulse of the blood sent from the heart, and to react upon the fluid so as to influence its course. It is, likewise, by virtue of this structure, that the parietes retain their form in the dead body, - one of the points that distinguish them from the veins.

The vitality of the arteries is inconsiderable. Hence their diseases are by no means numerous or frequent; an important fact, seeing that their functions are eminent, and their activity incessant.

## c. Intermediate, Peripheral or Capillary System.

The capillary or intermediate vessels are the vessels of extreme minuteness, by some considered to be formed by the termination of the arteries and the commencement of the veins; by others to be a distinct set of vessels. This system forms a plexus, which is distributed over every part of the body, and constitutes, in the aggregate, what is meant by the capillary system. It admits of two great divisions, one of which is situate at the termination of the branches given off from the aorta, and is called the general capillary system; the other at the termination of the branches of the pulmonary artery, — the pulmonic capillary system. Although the capillary system of man does not admit of detection by

a For some speculations as to the agency of this secretion in the production of the buffy state of the blood, &c., see M. Romain Gérardin, in Journal des Connaisances Médico-Chirurgicales, Mars, 1836.

the unaided sight, its existence is evidenced by the microscope; by injections, which can develope it artificially in almost every organ; by the application of excitants, and by inflammation. The parietes of the vessels frequently cannot be distinguished from the substance of the organs; - the colour of the blood, or the matter of the injection alone indicating their course. In some parts, as in the white textures, these vessels do not seem to admit the red particles of the blood, whilst, in others, the red particles always circulate. This diversity has given rise to the distinction of the capillaries into red and white; but there are probably no white capillaries. It is difficult, indeed, to conceive of the red particles being arrested at the mouths of the white vessels - if such existed, without their preventing the entrance of fluid into them. The true cause of the whiteness appears to be the small quantity of blood, which they receive, and it is only when the network is very close, and the quantity of blood passing through them great, that a perceptible colour is produced by the red particles.b There are certain textures, again, which receive neither the one nor the other, - the corneous and epidermous, for example, which are probably nourished by transudation of nutritive matter from the vessels through the cells of the tissue.

The ancients were of opinion, that the arteries and veins are separated by an intermediate substance, consisting of some fluid effused from the blood, and which they called, in consequence, parenchyma.d The notion is, indeed, still entertained, and is considered to be supported by microscopical observations. In the examination of delicate and transparent tissues, currents of moving globules are seen with many spaces of apparently solid substances, resembling small islets, surrounded by an agitated fluid. If it be irritated, by thrusting a fine needle into it, the motion of the globules becomes more rapid, new currents arise where none were previously perceptible, and the whole becomes a mass of moving particles, the general direction of which tends towards the points of irritation. But, although a part of the apparatus of intermediate circulation may be arranged, as we shall see presently, in this manner, there are reasons for the belief, that a more direct communication between the arteries and the veins exists also. substance of an injection passes from one set of vessels into the other without any evidence of intermediate extravasation. The blood has been seen, too, passing in living animals, directly from the arteries into the veins. Leeuenhoeke and Malpighi, on ex-

e Mr. Toynbee, Proceedings of the Royal Society, No. 48.

d Galen., Administrat. Anatom. vi. 2.

<sup>&</sup>lt;sup>2</sup> See Berres, in Magendie on the Blood, Select Medical Library Edit., p. 170, Philad.

b Carpenter, Human Physiology, § 479, Lond. 1842. See, also, Mandl, Manuel d'Anatomie générale, p. 194, Paris, 1843.

<sup>\*</sup> Select Works, containing his Microscopical Discoveries, by Samuel Hooke, p. 90, Lond. 1778.

f Epist. de Pulmonibus, 1661, and Haller, Element. Physiol.

the unaided sight, its existence is evidenced by the microscope; by injections, which can develope it artificially in almost every organ; by the application of excitants, and by inflammation. The parietes of the vessels frequently cannot be distinguished from the substance of the organs; - the colour of the blood, or the matter of the injection alone indicating their course. In some parts, as in the white textures, these vessels do not seem to admit the red particles of the blood, whilst, in others, the red particles always circulate. This diversity has given rise to the distinction of the capillaries into red and white; but there are probably no white capillaries. It is difficult, indeed, to conceive of the red particles being arrested at the mouths of the white vessels - if such existed, without their preventing the entrance of fluid into them. The true cause of the whiteness appears to be the small quantity of blood, which they receive, and it is only when the network is very close, and the quantity of blood passing through them great, that a perceptible colour is produced by the red particles.b There are certain textures, again, which receive neither the one nor the other, - the corneous and epidermous, for example, which are probably nourished by transudation of nutritive matter from the vesselse

through the cells of the tissue.

The ancients were of opinion, that the arteries and veins are separated by an intermediate substance, consisting of some fluid effused from the blood, and which they called, in consequence, parenchyma.d The notion is, indeed, still entertained, and is considered to be supported by microscopical observations. In the examination of delicate and transparent tissues, currents of moving globules are seen with many spaces of apparently solid substances, resembling small islets, surrounded by an agitated fluid. If it be irritated, by thrusting a fine needle into it, the motion of the globules becomes more rapid, new currents arise where none were previously perceptible, and the whole becomes a mass of moving particles, the general direction of which tends towards the points of irritation. But, although a part of the apparatus of intermediate circulation may be arranged, as we shall see presently, in this manner, there are reasons for the belief, that a more direct communication between the arteries and the veins exists also. The substance of an injection passes from one set of vessels into the other without any evidence of intermediate extravasation. The blood has been seen, too, passing in living animals, directly from the arteries into the veins. Leeuenhoeke and Malpighi, on ex-

d Galen., Administrat. Anatom. vi. 2.

<sup>\*</sup> See Berres, in Magendie on the Blood, Select Medical Library Edit., p. 170, Philad.

b Carpenter, Human Physiology, § 479, Lond. 1842. See, also, Mandl, Manuel d'Anatomie générale, p. 194, Paris, 1843.

Mr. Toynbee, Proceedings of the Royal Society, No. 48.

Select Works, containing his Microscopical Discoveries, by Samuel Hooke, p. 90,
 Lond. 1778.
 Epist. de Pulmonibus, 1661, and Haller, Element. Physiol.

here figured, according to Wagner, a are furnished with distinct

parietes.

The capillary vessels have been esteemed by some to belong chiefly to the arteries, the venous radicles not arising almost imperceptibly from the capillary system, as the arteries terminate in it, but having a marked size at the part where they quit this system, which strikingly contrasts with the excessive tenuity of the capillary arterial vessels; whilst between the capillary system and the arteries, there is no distinct line of demarcation. The opinion of Bichatb was, that this system is entirely independent of both arteries and veins; and Autenriethe imagined, that the minute arteries unite to form trunks, which again divide before communicating with the veins, so as to represent a system analogous to that of the vena portæ. The experiments of Dr. Marshall Halld on the batrachia, which were performed with signal care, led him to the following conclusions, which agree with those of Bichat, so far as regards the independent existence of a capillary system. The minute vessels, he says, may be considered as arterial, so long as they continue to divide and subdivide into smaller and smaller branches. The minute veins are the vessels that gradually enlarge The true capillary from the successive addition of small roots. vessels are distinct from these. They do not become smaller by subdivision, or larger by conjunction, but they are characterized by continual and successive union and division, or anastomoses, whilst they retain a nearly uniform diameter. The last branches of the arterial system, and the first root of the venous, Dr. Hall remarks, may be denominated minute, but the term "capillary" must be reserved for, and appropriated to, vessels of a distinct character and order, and of an intermediate station, carrying red globules, and perfectly visible by means of the microscope.

The capillary arteries are distinct in structure — as we shall see they are in office — from the larger arteries. All the coats of these minute vessels diminish in thickness and strength, as the tubes lessen in size, but more especially the middle coat, which, according to Wedemeyer, may still be distinguished by its colour in the transverse section of any vessels whose calibre is no less than the tenth of a line; but entirely disappears in vessels too small to receive the wave of blood in a manifest jet. But while the coats diminish, the nervous filaments, distributed to them, increase; the smaller and thinner the capillary, the greater the proportionate quantity of its nervous matter. The coats of the capillaries, becoming successively thinner and thinner, at length disappear altogether, and the vessels — many of them at least — terminate in membraneless canals or interstitial passages, formed in the substance of the tissues. The blood is contained — according to Wester the proportion of the substance of the tissues. The blood is contained — according to Wester the proportion of the substance of the tissues.


\* Elements of Physiology, by R. Willis, Lond. 1842.

b Anatomie Générale, tom. i.
d A Critical and Experimental Essay on the Circulation, &c., Lond. 1830; Amer. Edit. Philad. 1835.

demeyer, Gruithuisen, Döllinger, Carus, and others, in the different tissues, in channels, which it forms in them; even under the microscope, the stream is seen to work out for itself, easily and rapidly, a new passage in the tissues, which it penetrates, and it has been esteemed certain, that in the figura venosa of the egg, the blood is not surrounded by vascular parietes.

Most histologists of the day are disposed, however, to believe that the capillaries are provided with distinct coats. Such, as has been seen, appeared to Wagner to be the case in the frog's foot, when magnified 45 diameters; and it has even been announced, that these are composed of a fibrous structure, analogous to the mus-

cular.



Capillaries of the Web of the Frog's Foot.

a. Deep venous trunk, composed of three principal branches, b, b, b; and covered with a rete of smaller vessels.—(Wagner.)

Fig. 161, also from Wagner, exhibits the vascular rete and circulation of the web of the hind foot of a frog — Rana temporaria, magnified 110 times, in which the parietes are very distinct. In Fig. 162, from Wagner, which represents a portion of a live newt, magnified 150 diameters, the capillaries are exceedingly delicate, and their walls by no means so distinct. The arterial and venous


a Tiedemann, loc. cit.

Burdach, op. cit. iv. 191. See, also, Dr. Marshall Hall and others, Lond. Lancet,

Nov. 5, 1842, p. 197.

See J. Müller's Handbuch, u. s. w., Baly's translation, p. 216; and Geddings, art. Arteries, in American Cyclopedia of Practical Medicine, ii. 305, Philad. 1836.

trunks and the capillaries that form the medium of communication between them are well seen, as well as the islets of the substance of the lung, on which a granular or cellular texture is indistinctly



Bloodvessels of the Lung of a Live Newt.

b a. Pulmonary vein receiving blood from another vein, c; itself made up of two branches. d. Pulmonary artery anastomosing with the pulmonary veins by means of capillary vessels .-

perceptible.a Dr. Carpenterb is of opinion, that the mode of origin of the capillaries, also refutes the supposition, that they are mere passages channelled out of the tissues through which they convey the blood. He thinks there can be no doubt, that they are produced, in any newly forming tissue, not by the retirement of the cells, one from the other, so as to leave passages between them, but by the formation of communications among certain cells, whose cavities become connected with each other, so as to constitute a plexus of tubes, of which the original cell-walls become the parietes. But farther observations are needed on this matter. The whole affair of the arrangement of cells in the building up of tissues has appeared to the author to have been adopted too hastily; and opinions will doubtless experience vast modifications hereafter.

b Human Physiology, § 477, Lond. 1842.

<sup>\*</sup> See, on this subject, Mr. Paget, Brit. and For. Med. Rev., July, 1842, p. 286.

Of these fine capillaries, — the diameter of which, in parts finely injected, varies from the  $\frac{1}{1000}$ th to the  $\frac{1}{4000}$ th, and the  $\frac{1}{3000}$ th of an inch, and even more - some, according to Wedemeyer, communicate with veins. In the others, there are no visible openings or pores in the sides or ends, by which the blood can be extravasated preparatory to its being imbibed by the veins. There is nowhere apparent a sudden passage of the arterial into the venous stream: no abrupt boundary between the division of the two systems. The arterial streamlet winds through long routes before it assumes the nature, and takes the direction, of a venous streamlet. The ultimate capillary rarely passes from a large arterial into a large venous branch. Many speculations have, however, been indulged, regarding the mode in which the vascular extremities of the capillary system are arranged. Bichat regarded it as a vast reservoir, whence originate, besides veins, vessels of a particular order, whose office it is to pour out, by their free extremity, the materials of nutrition, - vessels, which had been previously imagined by Boerhaave, and are commonly known under the appellation of exhalants. Mascagnid supposed that the final arterial terminations are pierced, towards their point of junction with the veins, by lateral pores, through which the secreted matters transude; - but these points will farther engage attention under the heads of Nutrition, and Secretion.

## d. Veins.

The origin of the veins, like that of all capillary vessels, is imperceptible. By some they are regarded as continuous with the capillary arteries; Malpighie and Leeuenhoekf state this as the result of their microscopic observations on living animals; and it has been inferred, from the facility with which an injection passes from the arteries into the veins. According to others, cells exist between the arterial and the venous capillaries, in which the former deposit their fluid and whence the latter obtain it. Others, again, substitute a spongy tissue for the cells.

A question has also been asked, — whether the veins terminate by open mouths; or whether there may not be more delicate vessels, communicating with their radicles, similar to the exhalants, which are presumed to exist at the extremities of the arteries, and which are regarded as the agents of exhalation. All this is, however, conjectural. It has already been observed, that the mesenteric veins have been considered by some to terminate by open mouths in the villi of the intestines; and the same arrangement has been conceived to prevail with regard to other veins. Ribes

a Müller, op. cit. p. 211.

b Krause, in Müller's Archiv. Heft. 1, 1837; and Brit. and For. Med. Rev., July, 1838, p. 218; also, Henle, Allgemeine Anatomie, S. 176, Leipz. 1841.


See, on this subject, Wilbrand, Physiologie des Menschen, s. Rap. xxviii. and xxix. Leipz. 1840.

<sup>&</sup>lt;sup>d</sup> Vasor. Lymph. Corpor. Human. Histor., Sen. 1817; and Prodromo della Grande Anatomie, Firenz. 1819.

<sup>\*</sup> Opera., Lond. 1687.

Opera., Lugd. Bat. 1722.

concludes, from the results of injecting the veins, that some of the venous capillaries are immediately continuous with the minute arteries, whilst others open into the cells of the laminated tissue, and into the substance of the different organs.



Ramifications of the Splenic Artery in the Spleen.

When the veins become visible, they appear as an infinite number of tubes, extremely small, and communicating very freely with each other; so as to form a very fine network. These vessels gradually become larger and less numerous, but still preserve their reticular arrangement; until, ultimately, all the veins of the body empty themselves into the heart, by three trunks—the vena cava inferior, the vena cava superior, and the coronary vein. The first of these receives the veins from the lower part of the body, and extends from the fourth lumbar vertebra to the right auricle; the second receives all the veins of the upper part of the body; and into it the subclavian opens, into which the chyle and lymph

are discharged. It extends from the cartilage of the first rib to the right auricle. The coronary vein belongs to the heart exclusively. Between the superior and inferior cava a communication

is formed by means of the vena azygos.

Certain organs appear almost wholly composed of venous radicles. The spleen is one of these. Fig. 163 represents the ramifications of the splenic artery, in the substance of that organ; and if we consider, that the splenic vein has corresponding ramifications, the viscus would seem to be almost wholly formed of bloodvessels. The same may be said of the corpus cavernosum of the penis and clitoris, the nipple, urethra, glans penis, &c. If an injection be thrown into one of the veins that issue from these different tissues, they are wholly filled by the injection; which rarely occurs, if the injection be forced into the artery. Magendie<sup>a</sup> affirms, that the communication of the cavernous tissue of the penis with the veins occurs through apertures two or three millimètres — in. 0·117 — in diameter.

In their course towards the heart, particularly in the extremities, the veins are divided into two planes; - one subcutaneous or superficial; the other deep-seated, and accompanying the deep-seated arteries. Numerous anastomoses occur between these, especially when the veins become small, or are more distant from the heart. We find, that their disposition differs according to the organ. In the brain, they constitute, in great part, the pia mater; and enter the ventricles, where they contribute to the formation of the plexus choroides and tela choroidea. Leaving the organ we find them situate between the laminæ of the dura mater; when they take the name of sinuses. In the spermatic cord, they are extremely tortuous, anastomose repeatedly, and form the corpus pampiniforme; around the vagina, they constitute the corpus reliforme; in the uterus, the uterine sinuses, &c. The veins have three coats in superposition, according to most anatomists: but many modern anatomists are disposed to assign them six.b The outer coat is cellular, dense, and very difficult to rupture. The middle coat has been termed the proper membrane of the veins. The generality of anatomists describe it as composed of longitudinal fibres, which are more distinct in the vena cava inferior than in the vena cava superior; in the superficial veins than in the deep-seated; and in the branches than in the trunks. Magendie states, that he has never been able to observe the fibres of the middle coat; but that he has always seen a multitude of filaments interlacing in all directions; and assuming the appearance of longitudinal fibres, when the vein is folded or wrinkled longitudinally, which is frequently the case in the large veins. It exhibits no signs of muscularity; even when the galvanic stimulus is applied; yet Magendie suspects its chemical nature to be fibrinous. It was remarked, in

a Précis, &c. ii. 238.

<sup>&</sup>lt;sup>5</sup> See, on the researches of recent histologists, Mr. Paget, Brit. and For. Med. Rev. July, 1842, p. 285.

VOL. 11. - 8

an early part of this work, (vol. i., p. 38,) that the bases of the cellular and muscular tissue are, respectively, gelatin, and fibrin; and that the various resisting solids may all be brought to one or other of these tissues. The middle coat of the veins doubtless belongs to the former, and is a variety of the tissu jaune of the French anatomists. Magendie merely states its fibrinous nature to be a suspicion; and, like numerous suspicions, it may be devoid of foundation. Yet we have reason to believe, that it is contractile. Broussaisa affirms, that this action is one of the principal causes of the return of the blood to the heart. He conceives, that the alternate movements of contraction and relaxation are altogether similar to those of the heart; but that they are so light as not to have been rendered perceptible by any process in the majority of the veins, although very visible in the vena cava of frogs. where it joins the right auricle. In some experiments by Sarlandière on the circulation, he observed these movements to be independent of those of the heart. After the heart was removed, the contraction and relaxation of the vein continued for many minutes in the cut extremity, and even after the blood had ceased

The inner coat is extremely thin and smooth at its inner surface, and has an epithelial lining. It is very extensible, and yet presents considerable resistance; bearing a very tight ligature without being ruptured. In many of the veins, parabolic folds of the inner coat exist, like those in the lymphatics, and inservient to a similar purpose: the free edge of these valves is directed towards the centre of the circulation; showing that their office is to permit the blood to flow in that direction, and to prevent its retrogression. They do not seem, however, in many cases, well adapted for the purpose; inasmuch as their size is insufficient to obliterate the cavity of the vein. By most anatomists, this arrangement is considered to depend upon primary organization; but Bichat conceives it to be wholly owing to the state of contraction, or dilatation of the veins at the moment of death. Magendie, however, affirms, that he has never seen the distension of the veins exert any influence on the size of the valves; but that their shape is somewhat modified by the state of contraction or dilatation; and this he thinks probably misled Bichat.c Moreover, they are covered by the epithelial coat and consist of tissue like that of fibrous membrane, which, as Hunterd observed, shows, that they are not duplicatures of the lining membrane, Their number varies in different veins. As a general rule, they are more numerous, where the blood proceeds against its gravity, or where the veins are very extensible, and receive but a feeble support from the circumambient parts, as in the extremities. They are entirely wanting in the veins of the deep-seated viscera; in

b See, on this subject, the remarks on the circulation in the veins.

<sup>\*</sup> Op. citat., Amer. translation, p. 391.

e Précis, &c. ii. 241.
Henle; and Mr. Paget, Brit. and For. Med. Rev. July, 1842, p. 284.

those of the brain and spinal marrow, and of the lungs; in the vena portæ, and in the veins of the kidneys, bladder and uterus. They exist, however, in the spermatic veins; and, sometimes, in the internal mammary, and in the branches of the vena azygos. On the cardiac side of these valves, cavities or sinuses exist, which appear externally in the form of varices. These dilatations enable the refluent blood to catch the free edges of the valves, and thus to depress them, so as to close the cavity of the vessel; serving, in this respect, precisely the same functions as the sinuses of the pulmonary artery and aorta in regard to the semilunar valves.

The valves exist in veins of less than a line in diameter."

The three coats united form a solid vessel, - according to Bichat devoid of elasticity, but in the opinion of Magendie, belastic in an The elasticity is certainly much less than that eminent degree. of the arteries. The veins are nourished by vasa vasorum, or by small arteries, which have their accompanying veins. Every vessel, indeed, in the body, if we may judge from analogy, appears to draw its nutriment, not from the blood circulating in it, but from small arterial vessels, hence termed vasa vasorum. This applies not only to the veins, but to the arteries. The heart, for example, is not nourished by the fluid constantly passing through it; but by vessels, which arise from the aorta, and are distributed over its surface, and in its intimate texture. The coronary arteries and their corresponding veins are, consequently, the vasa vasorum of the heart. In like manner, the aorta and all its branches, as well as the veins, receive their vasa vasorum. There must, however, be a term to this; and if our powers of observation were sufficient we ought to be able to discover a vessel, which must derive its support or nourishment exclusively from its own stores:

The nerves that have been detected on the veins, are branches

of the great sympathetic.


The capacity of the venous system is generally esteemed to be double that of the arterial. It is obvious, however, that we can only arrive at an approximation, and that not a very close one. The size and number of the veins is generally so much greater than that of the corresponding arteries, that when the vessels of a membranous part are injected, the veins are observed to form a plexus, and, in a great measure, to conceal the arteries: in the intestines, the number is more nearly equal. The difficulty of arriving at any exact conclusion, regarding the relative capacities of the two systems, is forcibly indicated by the fact, that, whilst Borelli conceived the preponderance in favour of the veins to be as four to one, Sauvages estimated it at nine to four, Haller at sixteen to nine, and Keill at twenty-five to nine.

There is one portion of the venous system, to which allusion has already been made, which is peculiar. We mean the abdominal venous or portal system. All the veins, that return from the digestive organs, situate in the abdomen, unite into a large trunk,

<sup>\*</sup> Henle, op. cit.

<sup>c</sup> Elementa Physiologiæ, Lausan. 1757-1766.

b Précis, &c., ii. 243.



Portal System.

1. Inferior mesenteric vein: it is traced by means of dotted lines behind the pancreas (2) to terminate in splenic vein (3)-4. Spleen. 5. Gastric veins, opening into splenic vein. 6. Superior mesenteric vein. 7. Descending portion of the duodenum. 8. Its transverse portion which is crossed by superior mesenteric vein and by a part of trunk of superior mesenteric artery. 9. Portal vein. 10. Hepatic artery. 11. Ductus communis choledochus. 12. Divisions of duct and vessels at transverse fissure of liver. 13. Cystic duct leading to gall-bladder. — (Wilson.)

called the vena portæ. This, instead of passing into a larger vein - into the vena cava, for example - proceeds to the liver, and ramifies, like an artery in its substance. From the liver, other veins, called suprahepatic, arise, which empty themselves into the vena cava; and which correspond to the branches of the hepatic artery as well as to those of the vena portæ. The portal system is concerned only with the veins of the digestive organs situate in the abdomen; as, the spleen, pancreas, stomach, intestines and omenta. The veins of all the other abdominal organs, - of kidney, supracapsules, &c., are not connected with it. The first part of the vena portæ is call-

ed, by some authors, vena portæ abdominalis vel ventralis, to distinguish it from the hepatic portion, which is of great size, and has been called the sinus of the vena portæ.

## 2. BLOOD.

It is not easy to ascertain the total quantity of blood, circulating in both arteries and veins. Many attempts have been instituted for this purpose, but the statements are most diversified, partly owing to the erroneous direction followed by the experimenters, but, still more, to the variation that must be perpetually occurring in the amount of fluid, according to age, sex, temperament, activity of secretion, &c. Harvey and the earlier experimenters formed their estimates, by opening the veins and arteries freely on a living animal, collecting the blood that flowed, and comparing this with the weight of the body. The plan is, however, objectionable, as the whole of the blood can never be obtained in this manner, and the proportion discharged varies in different animals and circum-

BLOOD. 89

stances. By this method, Moulins found the proportion in a sheep to be  $\frac{1}{23}$ d; King, in a lamb,  $\frac{1}{20}$ th; in a duck,  $\frac{1}{30}$ th; and in a rabbit,  $\frac{1}{30}$ th. From these and other observations, Harvey concluded, that the weight of the blood of an animal is to that of the whole animal as 1 to 20. Drelincourt, however, found the proportion in a

hog to be nearly 10th; and Moor, 12th.a

An animal, according to Sir Astley Cooper, b generally expires, as soon as blood, equal to about 1th of the weight of the body, is abstracted. Thus, if it weigh sixteen ounces, the loss of an ounce of blood will be sufficient to destroy it; ten pounds will destroy a man weighing one hundred and sixty pounds; and, on examining the body, blood will still be found - in the small vessels especially - even although every facility may have been afforded for draining them. Experiments have, however, shown, that no fixed proportion of the circulating fluid can be indicated as necessary for the maintenance of life. In the experiments of Rosa, asphyxia occurred in young calves when from three to six pounds, or from \frac{1}{32}d to 10 th of their weight, had been abstracted, but in older ones not until they had lost from twelve to sixteen pounds, or from 11th to 4th of their weight. In a lamb, asphyxia supervened on a loss of twenty-eight ounces, or 1 th of its weight, and in a wether, on a loss of sixty-one ounces, or 23d of its weight. Blundelle found that some dogs died after losing nine ounces, or their weight; and others withstood the abstraction of a pound, or 10th of their weight; and Piorry affirms, that dogs can bear the loss of 1 th of their weight, but if a few ounces more be drawn, they succumb.

From all the experiments and observations, Burdach<sup>d</sup> concludes, that, on the average, death occurs when \$\frac{3}{4}\$ths or \$\frac{3}{6}\$ths, of the mass of blood is lost, although he has observed it in many cases, as in hæmoptysis, to supervene on the loss of \$\frac{1}{4}\$th, and even of \$\frac{1}{8}\$th.

The following table exhibits the computations of different physiologists, regarding the weight of the circulating fluid — arterial and venous.

lbs.				lbs.
Harvey,	Wagner,	-		20 to 25
Lister,	Quesnai,			- 27
Moulins,	F. Hoffmann,			- 28
Abildguard, ) Blumenbach, 7	Haller, -	1-	-	28 to 30
Lobb, 10	Young, -	120	-	- 40
Lower,	Hamberger,	4	2	- 80
Sprengel, 10 to 15 Miller and Burdach, - 20	Keill, -	-	4	- 100

Although the absolute estimate of Hoffmann is perhaps below

\* Haller, Elementa Physiologiæ, iv. 2, seq.

b Principles and Practice of Surgery, p. 33, Lee's edition, Lond. 1836.

Researches, Physiological and Pathological, p. 66 and 94, Lond. 1825.
 Die Physiologie als Erfahrungswissenschaft, iv. 101 and 334, Leipzig, 1832.

e Haller, op. citat.; and Herbst, Comment. Historico-critica, &c., de Sanguinis Quantitate, Gotting. 1822.

the truth, his proportion is probably nearly accurate. He conceives, that the weight of the blood is to that of the whole body as 1 to 5. Accordingly, an individual, weighing one hundred and fifty pounds, will have about thirty pounds of blood; one of two hundred pounds, forty; and so on. Of this, one-third is supposed to be contained in the arteries, and two-thirds in the veins. The estimate of Haller is, perhaps, near the truth; the arterial blood being, he conceives, to the venous, as 4 to 9. If we assume, therefore, that the whole quantity of the blood is thirty pounds in a man weighing one hundred and fifty pounds, — which is perhaps allowing too much, — nine pounds, at least, may be contained in the

arteries, and the remainder in the veins.a

An ingenious plan, proposed by M. Valentin for estimating the quantity of blood in the body, affords an approximation to the truth, and confirmatory of the estimate made from other data. Having weighed an animal, and determined the proportion of solid matter in a portion of its blood, he injects into its vessels a given quantity of distilled water, which soon becomes mixed with the blood. He then takes away a fresh portion of blood, and ascertains the proportion of solid matter in it. The relation between the amount of solid matter in the blood first taken, and that in the blood diluted with the given quantity of water, enables him to calculate the quantity of blood in the body of the animal. In this manner, M. Valentin found the ratio of blood to the weight of the body to be in the male dog as 1 to 4.36 in the male sex; and 1 to 4.93 for the female, and adapting these proportions to M. Quetelet's table of the weight of the human subject at different ages, he inferred, that the mean quantity of blood in the male adult, at the

a Good's Study of Medicine, Physiological Proem to class Hæmatica; Haller, Op. citat.; Rudolphi, Grundriss der Physiol. i. 40; Brandt, in art. Blut, in Encyclopäd. Wörterb. der Medic. Wissenschaft. v. 598, Berlin, 1830; and Mr. Ancell, Lectures on the Physiol. and Pathology of the Blood, in Lond. Lancet, May 16, 1840, p. 257.

b The following question and solution are given, in order to show, how the quantity

of blood may be estimated in the manner proposed by Valentin.

A portion of blood, (=1190 grains,) drawn from a dog, yielded 24.54 per cent. of solid matter. After injecting 10905 grains of water into the bloodvessels, a portion of blood drawn yielded 21.86, (or, by another trial, 21.89,) per cent. of solid matter. What was the amount of blood in the body at the commencement of the experiment?

Let x be the amount of blood after the first experiment. Then, since it contained

24.54 per cent. of solid residue, the amount of solid matter in it was 2454 x.

After injecting the water the whole amount of the diluted blood was x + 10905; and, (by the experiment,) the solid matter which it contained was = .2186 (x + 10905). But the solid matter was of the same amount in both cases. Therefore we have,

or, 
$$\begin{array}{c} \cdot 2454 \ x = \cdot 2186 \ (x+10905) \\ \cdot 2454 - \cdot 2186) \ x = \cdot 2186 + 10905 \\ \text{or,} \qquad x = \frac{2373 \cdot 8330}{\cdot 0268} = 88576 \ \text{grs.} \end{array}$$

Add for the blood first drawn, - - - 1190

And we get - - - - 89766 grs.
the weight of blood in the body, at the commencement of the experiment.
The ratio 21.89 per cent. gives - - 91269 grs.
And the mean of the two is - - 90517 "

time when the weight of the body may be presumed to be greatest, namely, at 30 years, should be about 34½ pounds; and that of the female at 50, when the weight is generally greatest, at about 26 pounds.<sup>a</sup>

The blood strongly resembles the chyle in its properties; — the great difference consisting in the colour; and the venous blood, and the chyle, and the lymph become equally converted into the

same fluid - arterial blood - in the lungs.

Venous blood, which chiefly concerns us at present, is contained in all the veins, in the right side of the heart, and in the pulmonary artery; - organs which constitute the apparatus of venous circulation. As drawn from the arm, its appearance is familiar to every one. At first, it seems to be entirely homogeneous; but, after resting for some time, it separates into different portions. The colour of venous blood is much darker than that of arterial: - so dark, indeed, as to have had the epithet black blood applied to it. Its smell is faint and peculiar; by some compared to a fragrant garlic odour, but it is sui generis; its taste is slightly saline, and also peculiar. It is viscid to the touch; coagulable, and its temperature has been estimated at 96° Fahrenheit; simply, we believe, on the authority of the inventor of that thermometric scale, who marked 96° as blood heat. This is too low by at least three or four degrees. Rudolphi, and the German writers in general, estimate it at 29° of Réaumur, or "from 98° to 100° of Fahrenheit;" whilst, by the French writers in general, its mean temperature is stated at 31° of Réaumur, or 102° of Fahrenheit; Magendie, who is usually very accurate, fixes the temperature of venous blood at 31° of Réaumur, or 102° of Fahrenheit; and that of arterial blood at 32° of Réaumur, or 104° of Fahrenheit. 100° may perhaps be taken as the average. This was the natural temperature of the stomach in the case related by Dr. Beaumont, which has been so often referred to in these pages. In many animals, the temperature is considerably higher. sheep it is 102° or 103°; but it is most elevated in birds. duck it is 107°. On this subject, however, further information will be given under the head of Calorification.

The specific gravity of the blood is differently estimated by different writers. Hence it is probable, that it varies in different individuals, and in the same individual at different periods. Compared with water its mean specific gravity has been estimated, by some, to be as 1.0527, by others, as 1.0800, to 1.0000. It is stated, however, to have been found as high as 1.126; and, in disease, as low as 1.022. It has, moreover, been conceived, that the effect of disease is, invariably, to make it lighter; and that the more healthy the individual, the greater is the specific gravity of the

<sup>\*</sup> Müller's Physiology, by Baly, Book ii. Sect. I.

b Grundriss der Physiologie, i. 143, Berlin, 1821. Precis, &c. ii. 229,

<sup>4</sup> Experiments, &c. on the Gastric Juice, &c. p. 274, Plattsburg, 1833.

blood; but our information on this point is vague. That it is not always the same in health is proved by the discrepancy of observers. Boyle estimated it at 1.041; Martine, at 1.045; Jurin, at 1.054; Muschenbroek, at 1.056; Denis, at 1.059; Senac, at 1.082; and Berzelius at from 1.052 to 1.126; J. Müller, at from 1.0527 to 1.057; and Mandl, at from 1.050 to 1.059. The average may perhaps be stated at 1.050. A part of the discrepancy may be owing to the specific gravity not having been always taken at the same temperature. Dr. B. Babington found experimentally, that four degrees of temperature corresponded with a difference of 001 of specific weight: consequently, if one author states the specific gravity of blood at about its circulating temperature — say 98° of Fahr. — while another states it at 60° Fahr. — the usual standard — the former will make it .0095 lighter than the latter.

When blood is examined with a microscope of high magnifying powers, it appears to be composed of numerous, minute, red particles or corpuscles, commonly called red globules, blood corpuscles and blood disks, suspended in the serum. These red particles have a different shape and dimension, according to the nature of the animal. In the mammalia, they are circular; and, in birds and cold-blooded animals, elliptical. In all animals, they are affirmed, by some observers, to be flattened, and marked in the centre with a luminous point, of a shape analogous to the general shape of the globule. Prof. Giacomonic of Padua, has, however, affirmed, that the red globules, swimming in serum, - which have been described by so many writers on the circulating fluid, - exist only in their imaginations. As in every case which rests on microscopic observation, the greatest discrepancy prevails here, not only as regards the shape but the size of the globules. These were first noticed by Malpighi,d and were afterwards more minutely examined by Leeuenhoek,e who at first described them, correctly enough, in general terms; but, subsequently, became hypothetical, and advanced the phantasy, that the red particles are composed of a series of globular bodies, descending in regular gradations; each of the red particles being supposed to be composed of six particles of serum; a particle of serum of six particles of lymph, &c. Totally devoid of foundation, as the whole notion was, it was implicitly believed for a considerable period, even until the time when Haller wrote. Hewsonf described the globules as consisting of a solid centre, surrounded by a vesicle, filled with a fluid; and to be "as flat as a guinea." Hunter, on the other hand, did not regard them as solid bodies, but as liquids, possessing a central attraction, which determines

b Dr. Babington, in Cyclop. Anat. and Physiol. parts iv. and v. c Encyclogr. des Sciences Medicales, Avril, 1840, p. 529.

4 Opera, Lond. 1687.

Experimental Inquiries, part iii. p. 16, Lond. 1777.

Burdach, Die Physiologie als Erfahrungswissenschaft, iv. 15, Leipz. 1832.

e Martine, in Edinb. Med. Essays, ii. 74; and Phil. Transact. for 1694.

On the Blood, by Palmer, reprinted in Select Med. Lib. p. 63, Philad. 1840.

BLOOD. 93

Fig. 165.

beaten so as to separate the Fibrin, mag-

c. A blood corpuscle undergoing change as

nified 900 times.

their shape. Della Torres supposed them to be a kind of disk

or ring, pierced in the centre; whilst Monro conceived them to be circular, flattened bodies, like coins, with a dark spot in the centre, which he thought was not owing to a perforation, as Della Torre had imagined, but to a depression. Cavallo, b again, conceived, that all these appearances are deceptive, depending upon the peculiar modification of the rays of light, as Corpuscles of Human Blood from a vein, affected by the form of the particle; and he concluded, that they are face. b. On edge a three quarter view. simple spheres. Amici found them B. Congeries of blood corpuscles in columns. of two kinds, both with angular by exposure to air. b. A lymph corpuscle margins; but, in the one, the cen(Wagner.) tre was depressed on both sides;

whilst, in the other, it was elevated. The observations of Dr. Young, of Sir Everard Home and Mr. Bauer, and of MM. Prévost and Dumas, e accord chiefly with those of Hewson. All these gentle-

men consider the red particles to be composed of a central globule, which is transparent and whitish, and of a red envelope, which is less transparent. Still more recently, however, Dr. Hodgkin and Mr. Lister, have denied that they are spherical, and that they consist of a central nucleus inclosed in a vesicle. They affirm, on the authority of a microscope, which, on comparison, was found equal to a

Fig. 166.

Blood Corpuscles of the edible frog, Rana esculenta. a, a a, b. Blood corpuscles, b. Seen edgewise. c. Lymph corpuscle. d. Altered by dilute acetic acid. — (Wagner.)

celebrated one, taken a few years ago to Great Britain by Professor Amici,5 that the particles of human blood appear to consist of circular, flattened, transparent cakes, their thickness being about 1th part of their diameter. These, when seen singly, appear to be nearly or quite colourless. Their edges are rounded, and being the thickest part, occasion a depression in the middle, which

- a Philos, Trans. for 1765, p. 252.
- <sup>b</sup> An Essay on the Medicinal Properties of Factitious Air, &c. p. 237, Lond. 1798.
- c Introduct. to Med. Literature, p. 545.
- d Philosoph. Transact. for 1811-18; and Lectures on Comp. Anat. iii. 4, Lond. 1823.
- Annales de Chimie, &c. xxiii. 50, 90; and Journal of Science and Arts, xvi. 115. Philosoph. Magazine and Annals of Philosophy, ii. 130, Lond. 1827; also, Hodgkin, Catalogue of the Preparations in the Anatomical Museum of Guy's Hospital, introd. to sect. xi. P. i. Lond. 1829.
  - Edinb. Medical and Surgical Journal, xvi. 120.

exists on both surfaces. The view of these gentlemen, conse-

quently, appears to resemble that of Monro.

Amidst this discordance, it is difficult to know which view to adopt.a The belief in their consisting of circular, flattened, transparent bodies, with a depression in the centre, and that they consist of an external envelope and of a central nucleus, the former of which is red and gives colour to the blood, appears to have the greatest weight of authority in its favour. The nucleus is devoid of colour, and appears to be independent of the envelope; as, when the latter is destroyed, the central portion preserves its original shape. The nucleus is much smaller than the envelope, being, according to Dr. Young, only about one-third the length, and one-half the breadth of the entire particle. According to Sir Everard Home, the globules, when enveloped in the colouring matter, are 1700th part of an inch in diameter, requiring 2,890,000 to a square inch; but when deprived of their colouring matter, they appear to be 2000 th part of an inch in diameter, requiring 4,000,000 of globules to a square inch. According to these measurements, the globules, when deprived of their colouring matter, are not quite one-fifth smaller. The views of MM. Prévost and Dumas, who have investigated the subject with extreme care and signal ingenuity, are deserving of great attention. They conceive the blood to consist essentially of serum, in which a quantity of red particles is suspended; that each of these particles consists of an external red vesicle, which incloses, in its centre, a colourless globule; that, during the progress of coagulation, the vesicle bursts, and permits the central globule to escape; that, on losing their envelope, the central globules are attracted together; that they are disposed to arrange themselves in lines and fibres; that these fibres form a network, in the meshes of which they mechanically entangle a quantity of both the serum and the colouring matter; that these latter substances may be removed by draining, and by ablution in water; that, when this is done, there remains only pure fibrin; and that, consequently, fibrin consists of an aggregation of the central globules of the red particles, while the general mass, that constitutes the crassamentum or clot, is composed of the entire particle. So far this seems satisfactory; but, we have seen, Dr. Hodgkin does not recognise the existence of external vesicle, or of central globule; and he affirms, contrary to the notion of Sir Everard Home and others, that the particles are disposed to coalesce in their entire state. This is best seen, when the blood is viewed between two slips of glass. Under such circumstances, the following appearances, according to Dr. Hodgkin, are perceptible. When human blood, or that of any other animal having circular particles, is examined in this manner, considerable agitation is, at first, seen to take place among

See, on this subject, Paine, Medical and Physiological Commentaries, vol. i, Appendix on the Microscope, p. 899, New York, 1840; Mr. Paget. British and For. Med. Rev. July, 1842, p. 260; and Mr. T. Wharton Jones, Brit. and For. Med. Rev. Oct. 1842, p. 585.

b Lectures on Comparative Anatomy, iii. 4, and v. 100, Lond. 1828.

BLOOD.

the particles; but, as this subsides, they apply themselves to each other by their broad surfaces, and form piles or rouleaux, which are sometimes of considerable length. These rouleaux often again combine amongst themselves. - the end of one being attached to the side of another, - producing at times, very curious ramifica-

tions. (See Fig. 165, B.)

The belief in the particles being flattened disks is now generally received; and the form of the disk is found to be altered by various substances. Its external envelope admits readily the endosmose of fluids, so that, if placed in water, they may assume a truly globular shape. In examining the blood, consequently, it is advisable to dilute it with a fluid of as nearly as possible the same character as the serum. In the particles of the blood of the frog as represented in Fig. 166 — a nucleus is observed projecting somewhat from the central portion: this is rendered extremely distinct by the action of acetic acid, which dissolves the rest of the particle, and renders the nucleus more opaque. It then appears to consist of a granular substance.a .

Microscopical discordances are no less evidenced by the estimates, which have been made of the size of the red globules; yet all are adduced on the faith of positive admeasurements. Leaving out of view the older, and, consequently, it might be presumed, less accurate observations, the following table will show their diameter in human blood, on the authority of some of the most emi-

nent microscopic observers of more recent times.

Sir E. Home and Bauer, with colouring \ \frac{1}{1700} th part of an inch,
Eller, 1930
Sir E. Home and Bauer, without colour-
ing matter, 5
Jurin, 2000
Müller, 2300 to 3500
Mandl, 2625 to 1 3150
Hodgkin, Lister, and Rudolphi, - 3000
Sprengel, 3000 to 1 3500
Cavallo,
Blumenbach and Senac, 3330
Tabor, 3600
Milne Edwards,
Wagner,
Kater, 1 6000
D
Haller Wollaston and Weber
5000
Young,

\* Carpenter, Human Physiology, § 570, Lond. 1842; see, also, Dr. G. O. Rees, and Mr. S. Lane, in Guy's Hospital Reports, No. xiii. Oct. 1841.

<sup>b</sup> Burdach, Die Physiclogie als Erfahrungswissenschaft, iv. 21, Leipz. 1832. See, also, Ancell, Lectures on the Physiology and Pathology of the Blood, in Lond. Lancet, Dec. 7, 1839, p. 380.

The blood of different animals is found to differ greatly in the relative quantity of the red globules it contains, the number seeming to bear a pretty exact ratio with the temperature of the animal. The higher the natural temperature, the greater the proportion of particles; and arterial always containing a much greater

proportion than venous blood.

The blood corpuscles of the greater part of the mammalia have the same shape as those of man; but their size varies greatly in different families. It would appear, from the researches of Mandl, a that of the mammalia the elephant has the largest globules, (100 th of a millimètre,) and the ruminantia the smallest; that the family of camels is the only one, whose corpuscles are not round like those of the other mammalia, but elliptical like those of birds,

reptiles, and fishes.b

The chemical constitution of the blood corpuscles is not definitely settled. Two proximate principles have been discovered in them—
hematosin and globulin. The former, as mentioned hereafter, has been supposed to be the colouring matter. The latter, which differs from the globulin of Laennec,—an impure hematosin mingled with some albumen,— is the main constituent of the globules, and is the same as the blood-casein of Simon. It has not been separated, but is presumed to differ but little in its properties from protein. They are supposed to be produced originally in the germinal membrane of the embryo; but throughout the remainder of existence, they are formed in the blood from the chyle.

In addition to the red globules, colourless or white corpuscles are observed in the blood. These were observed by Müller in the blood of frogs; and by Mandle in that of the mammalia. They are small, colourless corpuscles, finely granulated; insoluble in water, and strongly refracting light. According to Mandl, they may be separated into two species, - some round, and containing two or three granules, which become more evident when they are treated with acetic acid: these are the true lymph globules, described already (vol. i., p. 625); the others, generally also round; sometimes oblong; and at others irregular; the edges slightly notched; and the surface finely granulated. They appear to be composed of a multitude of small molecules, from  $\frac{1}{1000}$ th to  $\frac{1}{1200}$ th of a millimètre in diameter; some are also found single. These globules are seen forming under the microscope, when the blood, placed between two glasses, is attentively examined. They are, in Mandl's opinion, produced by the coagulation of the fibrin, and hence they are called by him fibrinous globules. Recently, however, he has abandoned this name, "be-

Manuel d'Anatomie générale, p. 248, Paris, 1843.

c Carpenter, Human Physiology, § 575, Lond. 1842.

4 Wagner, Elements of Physiology, by Willis, § 96, Lond. 1842.

b Op. citat., and Annales des Sciences Naturelles, 1824 and 1825. See, on the blood corpuscles of different animals, Burdach, Op. cit. p. 21; and Gulliver, Appendix to Gerber's Elements of the General and Minute Anatomy of Man and the Mammalia, Lond. 1842.

<sup>·</sup> Gazette Médicale, 1837; and Manuel d'Anatomie générale, p. 252, Paris, 1843.

BLOOD.

cause it rests on a chemical character, that requires confirmation; and because it is not drawn from anatomical characters, which ought chiefly to fix the attention of the microscopist." He now terms them white granulated corpuscles.a Dr. Barry and Mr. Addison, however, think, that the colourless corpuscles, - which have generally been regarded as lymph corpuscles - are formed from the central portion of the blood corpuscles: they consider them to hold an intermediate position between the true red corpuscles, and the greatly modified forms of corpuscles, which, in their view, are the bases of the tissues, as well as of pus and other globules. The

subject requires further investigation.b

When blood is drawn from a vessel, and left to itself, it exhales, so long as it is warm, a fetid vapour consisting of water and animal matter, of a nature not known. This vapour is what has been called the halitus of the blood, - by Plenck, the gas animale sanguinis, which he conceives to be composed of carbon and hydrogen, and to be inservient to many supposititious uses in the economy. The odour exhaled by the blood would appear to preserve the same general characters under all circumstances. After a time, the blood coagulates, giving off, at the same time, it has been said, a quantity of carbonic acid gas. This disengagement is not evident, when the blood is suffered to remain exposed to the air, except, perhaps, by the apertures or canals formed by its passage through the clot; but it can be collected by placing the blood under the receiver of an air-pump, and exhausting the air. On this fact, however, observers do not all accord. The experiments of Vogel, Brande, Sir E. Home, and Sir C. Scudamore, are in favour of such evolution; and the last gentleman conceives it even to be an essential part of the process; but other distinguished experimenters have not been able to detect it. Neither Dr. John Davy,h nor Dr. Duncan, Jr., nor Dr. Christison, could procure it during the coagulation of the blood. Dr. Turner suggests that the appearance of the carbonic acid, in the experiments of Vogel, Brande, and Scudamore, might easily have been occasioned by casual exposure to the atmosphere, previous to the blood being placed under the receiver; but we have no reason for believing, that this source of fallacy was not guarded against as much by one set of experimenters as by the other. Our knowledge, on this point, is confined then to the fact,

Manuel d'Anatomie générale, p. 554, Paris, 1843.

Ancell, Lectures on the Blood, Lancet, Jan. 18, 1840, p. 608.

F Philosophical Transactions for 1820, p. 6; and an Essay on the Blood, p. 107, Lond. 1824.

Elements of Chemistry, 5th edit. by Dr. Bache, p. 607, Philad. 1835.

b See, on this point, Carpenter, Human Physiology, Amer. Edit., p. 419, Philad. 1843; and Andral, Hématologie Pathologique, p. 33, Paris, 1843.

<sup>4</sup> Annales de Chimie, t. xciii. Philosophical Transactions for 1818, p. 181. f Lectures, &c. iii. 8.

h Phil. Trans. for 1823, p. 506; and Edinb. Med. and Surg. Journ. xxix. 253. Since that time, however, Dr. Davy has succeeded in extricating it both from venous and arterial blood. See his Researches, Physiological and Anatomical, Dunglison's Amer, Med. Lib. Edit. p. 82, Philad. 1840.

that, by some, carbonic acid gas has been found exhaled during the process of coagulation; - by others, not. More recent experiments, by Stromeyer, and by Gmelin, Tiedemann, and Mitscherlich, would seem to decide, that the blood does not give off any free carbonic acid, but that it holds a certain quantity in a state of combination; and that this combination is intimate is shown by the fact, mentioned by Müller, that blood, artificially impregnated with carbonic acid, yields no appreciable quantity of the gas, when subjected to the air-pump. M. Magnus, however, found, in his experiments, that not only venous, but arterial blood, contains carbonic acid, oxygen, and azote, and that, as regards carbonic acid, arterial blood contains more than venous; and he accounts for the failure of those, who have attempted to elicit carbonic acid from venous blood by the air-pump, to the air in the receiver not having been sufficiently rarefied. Prof. C. H Schultz. of Berlin — who believes that the vesicles of the blood, in a perfect state, are composed of a membranous covering, whose interior is filled with an aëriform fluid in the midst of which is found the nucleuse - succeeded in so evident a manner by the following simple method in extracting air from the blood, "that it is impossible to doubt there exists a great quantity of air in the vesicles." He completely filled a bottle with warm blood, flowing immediately from the vein of a horse, and hermetically sealed the bottle so that the cork was plunged into the blood, thus absolutely preventing the contact of air. The blood, in cooling, diminished in volume, and thus produced a perfect vacuum in the upper part of the bottle; and in proportion as this took place, bubbles of air arose from the blood and filled the vacuum. Chemical analysis of this air demonstrated that it was carbonic, acid. In arterial blood, he found oxygen mixed with more or less carbonic acid.f The experiments of Dr. Stevens, and of Dr. Robert E. Rogers, also exhibit, that carbonic acid is contained in the blood. The latter observer found, when a portion of venous blood was placed in a bag of some membrane, and the bag was immersed in an atmosphere of some gas - as oxygen, hydrogen, or nitrogen - that carbonic acid was pretty freely evolved.

Whilst the blood is circulating in the vessels, it consists of liquor sanguinis and red corpuscles; but during coagulation, it separates into two distinct portions; a yellowish liquid, called the serum; and a red solid, known by the name of the clot, cruor, crassamentum, coagulum, placenta, insula or hepar sanguinis. The proportion of the serum to the crassamentum varies greatly in

a Schweigger's Journal für Chemie, u. s. w. lxiv. 105.

b Tiedemann und Treviranus, Zeitschrift für Physiologie, B. v. H. i.; Geddings's North Amer. Archives for July and August, 1835; and British and Foreign Med. Review, No. 9, p. 590, April, 1836.

d Annales de Chimie et de Physique, Nov. 1837. See, also, Carpenter, Human Physiology, § 541, Lond. 1842.

<sup>Lond. Lancet, Aug. 10, 1839, p. 713.
Philos. Transact. for 1834-5, p. 334. See, also, Dr. M. Edwards, in article "Blood," Cyclopædia of Anat. and Phys. Part. iv. p. 404, Lond. 1836.</sup> 

Amer. Journ. of the Med. Sciences, Aug. 1836, p. 283.

BLOOD. 99

different animals, and in the same animal at different times, according to the state of the system. The latter is more abundant in healthy, vigorous animals, than in those that have been impoverished by depletion, low living, or disease. Sir Charles Scudamore found, by taking the mean of twelve experiments, that the crassamentum amounted to 53.307 per cent. in healthy blood.

The difference between living and coagulated blood may be ex-

pressed in a tabular form as follows: -

The serum is viscous, transparent, of a slightly yellowish hue, and alkaline, owing to the presence of a little free soda. Its smell and taste resemble those of the blood. Its average specific gravity has been estimated at about 1.027. But on this point, also, observers differ. Dr. John Davyd found it to vary from 1.020 to 1.031. Martine, Muschenbroek, Jurin, and Haller, from 1.022 to 1.037: Berzelius and Wagner, from 1.027 to 1.029; Christison, from 1.029 to 1.031; Lauer, from 1.009 to 1.011; whilst Thackrahh found the extremes to be 1.004 and 1.080. At 158° of Fahrenheit. it coagulates; forming at the same time, numerous cells, containing a fluid, which oozes out from the coagulum of the serum, and is called the serosity. It contains, according to Bostock, about 1th of its weight of animal matter, together with a little chloride of sodium. Of this animal matter, a portion is albumen, which may be readily coagulated by means of galvanism; but a small quantity of some other principle is present, which differs from albumen and gelatin, and to which Marceti gave the name muco-extractive matter, and Bostock, uncoagulable matter of the blood — as a term expressive of its most characteristic property. Serum preserves its property of coagulating, even when largely diluted with According to Brande, it is almost pure liquid albumen,

- \* Roget's Outlines of Physiology, American Edition, by the author, § 385, Philad. 1839.
- b B. Babington, art. Blood, in Cyclop. of Anat. and Physiol., Lond. 1836; T. W. Jones, Brit. and For. Med. Rev., Oct. 1842, p. 588; and Mandl, Manuel d'Anatomie générale, p. 255, Paris, 1843.

c Bostock's Physiology, &c. edit. cit. p. 287; Marcet, in Medico-Chirurg. Trans.

iii. 363; and Mandl, Manuel d'Anatom. générale, p. 230, Paris, 1843.

d Researches, Physiological and Anatomical, Dunglison's Amer. Med. Lib. Edit. p. 11. Philad. 1840.

Elements of Physiology, by R. Willis, § 103, Lond. 1842.

On Granular Degeneration of the Kidneys, p. 61, London, 1839, or Dunglison's American Med. Library Edit., Philad. 1839.

5 Hecker's Annalen, xviii. 393; and Burdach, op. cit. iv. 29.

h Inquiry into the Nature and Properties of the Blood, &c. Lond. 1819.

i Medico-Chirurg. Transact. ii. 364.

Philosoph. Transact. for 1809, p. 373.

united with soda, which keeps it fluid. Consequently, he affirms, any reagent, which takes away the soda, will produce coagulation; and by the agency of caloric, the soda may transform a part of the albumen into mucus. The action of the galvanic pile coagulates the serum, and forms globules in it analogous to those of the blood.

From the analysis of serum, by Berzelius,a it appears to consist in 1000 parts; - of water, 903; albumen, 80; substances soluble in alcohol, - as lactate of soda and extractive matter, chlorides of sodium and potassium, 10; substances soluble in water, -as soda and animal matter, and phosphate of soda, 4; loss, 3. Marcet assigns it the following composition: - water, 900 parts; albumen, 86.8; chlorides of potassium and sodium, 6.6; muco-extractive matter, 4; carbonate of soda, 1.65; sulphate of potassa, 0.35, and earthy phosphates, 0.60; - a result, which closely corresponds with that of Berzelius, who states that the extractive matter of Marcet is lactate of soda, united with animal matter. One of the most recent analyses is by M. Lecanu.b According to him, 1000 parts contain, - water, 906 parts; albumen, 78; animal matter, soluble in water and alcohol, 1.69; albumen combined with soda, 2.10; crystallizable fatty matter, 1.20; oily matter, 1; chlorides of sodium and potassium, 6; subcarbonate and phosphate of soda, and sulphate of potassa, 2.10; phosphate of lime, magnesia and iron, with subcarbonate of lime and magnesia, 0.91; loss, 1.º Occasionally, the serum presents a whitish hue, which has given rise to the opinion that it contains chyle; but it would seem that this is fatty matter, and that it is always present. In the serum of the blood of spirit drinkers, Dr. Traill found a considerable portion of this substance, which has been considered to favour the notion, that the human body may, by intemperance, become preternaturally combustible; and has been used to account for some of the strange cases of spontaneous combustion, or rather of preternatural combustibility, which are on record. Dr. Christison has likewise met with fat mechanically diffused through the serum, like oil in an emulsion. On one occasion, he procured five per cent. of fat from milky serum, and one per cent. from serum which had the aspect of whey.d

The crassamentum or clot is a solid mass, of a reddish-brown colour, which, when gently washed for some time under a small stream of water, separates into two portions, — colouring matter and fibrin. As soon as the blood is drawn from a vessel, the colouring matter of the red globules leaves the central nucleus free; these then unite, as we have seen, and form a network, containing some of the colouring matter, and many whole globules. By washing the clot in cold water, the free colouring matter and the globules can be removed, and the fibrin will alone remain. When freed from the colouring matter, the fibrin is solid, whitish, insipid, inodorous, heavier than water, and without action on vege-

THE THE 1337.

Medico-Chirurg. Transactions, iii. 231.

Journal de Pharmacie, xvii.; and Annales de Chimie, &c., xlviii. 308.
 See, also, Boudet, in Journal de Pharmacie, Juin, 1833; and Annales de Chimie,
 4 Edinb. Med. and Surg. Journal, xvii. 235, and xxxiii. 274.

BLOOD. 101

table colours; elastic when moist, and becoming brittle by desiccation. It yields, on distillation, much carbonate of ammonia, and a bulky coal, the ashes of which contain a considerable quantity of phosphate of lime, a little phosphate of magnesia, carbonate of lime, and carbonate of soda. One hundred parts of fibrin, according to Berzelius, consist of carbon, 53.360; oxygen, 19.685; hydrogen, 7.021; azote, 19.934. Fibrin has been designated by various names. It is the gluten, coagulable lymph, and fibre of the blood of different writers. Its specific gravity is said to be greater than that of the serum; but the difference has not been accurately estimated, and cannot be great. The red particles are manifestly, however, heavier than either, as we find them subsiding during coagulation to the lower surface of the clot, when the blood has flowed freely from the orifice in the vein. Fibrin appears to be the most important constituent of the blood. It exists in animals, in which the red particles are absent, and is the basis of the muscular tissue.

The colouring matter of the blood, called, by some, cruorin, hematin, hematosin, zoo-hematin, hemachroin, globulin (of Lecanu,) and rubrin, has been the subject of anxious investigation with the analytical chemist. We have already remarked, that it resides in distinct particles or globules; and, in the opinion of the best observers, in the envelope of those globules. The globules themselves are insoluble in serum, but their colouring principle is dissolved by pure water, acids, alkalies, and alcohol. Raspaila asserts, that the globules or nuclei are entirely soluble in pure water, but MM. Donné and Boudet, who repeated his experiments, declare that they are wholly insoluble, and Müller is of the same opinion. Great uncertainty has always existed regarding the cause of the colour of the globules. 'As soon as the blood was found to contain iron, the peroxide of which has a red hue, the colour of the red globules was ascribed to the presence of that metal. Fourcroy and Vauqueline held this opinion, conceiving the iron to be in the state of subphosphate; and they affirmed, that if this salt be dissolved in serum by means of an alkali, the colour of the solution is exactly like that of the blood. Berzelius, however, showed, that the subphosphate of iron cannot be dissolved in serum by means of an alkali, except in very minute quantity; and that this salt, even when rendered soluble by phosphoric acid, communicates a tint quite different from that of the red globules. He found that the ashes of the colouring matter always yield oxide of iron in the proportion of 1/200 th of the original mass; whence it was inferred, that iron is somehow or other concerned in the production of the colour; but the experiments of Berzelius did not indicate the state in which that metal exists in the blood. He could not detect its presence by any of the liquid tests.

Chimie Organique, p. 368, Paris, 1833.

Handbuch der Physiologie, Baly's translation, p. 105, Lond. 1838.

System. Chym. ix. 207. 4 Med. Chir. Trans., iii. 213.

The views of Berzelius, and the experiments on which they were founded, were not supported by the researches of Mr. Brande." He endeavoured to show, that the colour of the blood does not depend upon iron; for he found the indications of the presence of that metal as considerable in the parts of the blood that are devoid of colour, as in the globules themselves; and in each it was present in such small quantity, that no effect, as a colouring agent, could be expected from it. He supposed that the tint of the red globules is produced by a peculiar, animal colouring principle, capable of combining with metallic oxides. He succeeded in obtaining a compound of the colouring matter of the blood with the oxide of tin; but its best precipitants are the nitrate of mercury and corrosive sublimate. Woollen cloths, impregnated with either of these compounds, and dipped in an aqueous solution of the colouring matter, acquires a permanent red dye, unchangeable by washing with soap. The conclusions of Brande have been supported by Vauquelin, but the fact, connected with the presence of iron, seems to have been decided by many observers. Engelharte demonstrated. that the fibrin and albumen of the blood, when carefully separated from colouring particles, do not contain a trace of iron; whilst he procured iron from the red globules by incineration. He also succeeded in proving the presence of iron in the colouring matter by the liquid tests; for on transmitting a current of chlorine gas through a solution of red globules the colour entirely disappeared, white flocks were thrown down, and a transparent solution remained, in which the peroxide of iron was discovered by the usual reagents. The results, obtained by Engelhart, as regards the quantity of iron, correspond with those of Berzelius. These facts have since been confirmed by Rose,d of Berlin; and Würzer, of Marburg, by pursuing Engelhart's method, by liquid tests, has detected the existence of the protoxide of manganese, likewise. The proportion of iron does not appear to be more than one-half per cent. ; yet, as it is contained only in the colouring matter, there is some reason for believing, that it may be concerned in the coloration of the blood, although probably in the form of oxide. The sulpho-cyanic acid has been detected in the saliva; and this acid, when united with the peroxide of iron, forms a colour exactly like that of venous blood; so that it has been presumed it may be connected with the coloration of the blood, but this is not probable; for Dr. Stevens found, that venous blood is darkened by the sulpho-cyanic acid. M. Lecanus has subjected the hematosin or colouring matter

Philosophical Transactions for 1812, p. 90.
Annales de Chimie et de Physique, tom. i. p. 9.

c Edinb. Med. and Surg. Journal, Jan. 1827; and Turner's Chemistry, 5th Amer. Edit. p. 605, Philad. 1835.

d Poggendorf's Annalen, vii. 81; and Annales de Chimie, &c. xxxiv. 268.

c Schweigger's Journal, lviii. 481.

r See, on this subject, Hermstädt, in Schweigger's Journ. 1832; J. Müller, op. cit. p. 125, and Brandt, art. Blut, in Encyclopäd. Wörterb. der Medicinisch. Wissenschaft. v. 606, Berlin, 1830.

8 Annales de Chimie et de Physique, xlv. 5.

BLOOD. 103

to analysis, and found it to be composed of: — loss, representing the weight of the animal matter, 97.742; subcarbonate of soda, alkaline chlorides, subcarbonates of lime and magnesia, and phosphates of lime and magnesia, 1.724; peroxide of iron, 0.534. The result of his researches induces him to conclude, that the colouring matter is a compound of albumen with some colouring substance yet unknown. This substance yielded on analysis: — loss, 98.26; peroxide of iron, 1.74; and M. Lecanu suggests, that it may result from the combination of some animal matter with certain ferruginous compounds, analogous to the cyanides.

After all, therefore, our ignorance on this subject is still great; and all that we seem to know is, that the peroxide of iron is contained in the colouring matter of the blood. Scherer has found, that the iron may be wholly dissolved by the agency of acids, and that the animal matter, boiled afterwards in alcohol, colours the

spirit deeply red.

Of late, the origin of the blood corpuscles has been an interesting topic with the histologist, but it is far from being determined.a Some of the views will be referred to under Nutrition. The redness of the fluid is one of its most obvious characteristics; and we are induced to esteem the change effected in the lungs, as regards colour, of eminent importance. It is, however, no farther so, than as it indicates the accomplishment of the conversion of venous into arterial blood. There is nothing essential, connected with the mere coloration. In the insect, the blood is transparent; in the caterpillar, of a greenish hue; and in the internal vessels of the frog, vellowish. In man, it differs according to numerous circumstances; and the hue of the skin, which is partly dependent upon these differences, thus becomes an index of the state of individual health or disease. In the morbus caruleus, cyanopathy or blue disease, the whole surface is coloured blue, especially in those parts where the skin is delicate, as on the lips, - owing to a communication existing between the right and left sides of the heart, so that the blood can pass from one to the other, without proceeding through the lungs. The appearance, too, of the jaundiced is familiar to all.

The formation of the clot, and its separation from the serum, are manifestly dependent upon the fibrin; which, by assuming the solid state, gives rise to the coagulation of the blood;—a phenomenon, which has occasioned much fruitless speculation and experiment; yet, if the views of Raspail<sup>b</sup> were proved to be correct, it would be sufficiently simple. The alkaline character of the blood, and the production of coagulation by a dilute acid leave no doubt, in his mind, that an alkali is the menstruum of the albumen of the blood. The alkaline matter, he thinks, is soda,

b Chimie Organique, p. 373.

<sup>&</sup>lt;sup>a</sup> For various views on this subject, see Carpenter, Human Physiology, § 575-578, Lond. 1842; M. Barry, Philosoph. Transact. 1842; and Lond. Edinb. and Dub. Philos. Mag. May, 1843; M. Donné, Comptes rendus, 7 Mars, 1841; and T. W. Jones, Proceedings of the Royal Society, No. 56, Lond. 1843.

but more especially ammonia, of which, he says, authors take no account; but whose different salts are evident under the microscope. Now, "the carbonic acid of the atmospheric air, and the carbonic acid, which forms in the blood by its avidity for oxygen, saturate the menstruum of the albumen, which is precipitated as a clot. The evaporation of the ammonia, and, above all, the evaporation of the water of the blood, which issues smoking from the vein, likewise set free an additional quantity of dissolved albumen, and the mass coagulates the more quickly as the blood is

less aqueous."

The process of coagulation is influenced by exposure to the air. Hewson affirmed, that it is promoted by such exposure, but Hunter was of an opposite opinion. If the atmospheric air be excluded, - by filling a bottle completely with recently drawn blood, and closing the orifice with a good stopper, - coagulation is retarded. Yet Sir C. Scudamore mentions the singular fact, that if blood be confined within the exhausted receiver of an airpump, the coagulation is accelerated; and MM. Gmelin, Tiedemann, and Mitscherlicha found that, under such circumstances, both venous and arterial blood coagulated as perfectly as under ordinary circumstances. The presence of air is certainly not essential to the process. Experiments have also been made on the effect produced by different gases on the process of coagulation; but the results have not been such as to afford much information. It is asserted, for example, by some, that it is promoted by carbonic acid, and certain other of the irrespirable gases, and retarded by oxygen: by others, the reverse is affirmed; whilst Davyb and M. Schröder van der Kolke informs us, that they could not perceive any difference in the period of the coagulation of venous blood, when it was exposed to azote, nitrous gas, oxygen, nitrous oxide, carbonic acid, hydrocarbon, or atmospheric air.

The time, necessary for coagulation, is affected by temperature. It is promoted by warmth; retarded, but not prevented, by cold. Hewson froze blood, newly drawn from a vein, and afterwards thawed it: it first became fluid, and then coagulated as usual. Hunter made a similar experiment with the like result. It is obviously, therefore, not from simple refrigeration that the blood coagulates. Sir C. Scudamore found, that the blood, which begins to coagulate in four minutes and a half, in a temperature of 53° Fahr., undergoes the same change in two minutes and a half at 98°; and that, which coagulates in four minutes at 98° Fahr., becomes solid in one minute at 120°. On the contrary, the blood,

<sup>\*</sup> Tiedemann und Treviranus, Zeitschrift für Physiol. B. v. Heft i. and in Op. citat. See, also, Mandl, Archiv. génerale de Méd. ix. 278, Paris, 1840; and Manuel d'Anatomie générale, p. 227, Paris, 1843.

b Researches, &c. chiefly concerning nitrous oxide, p. 380, Lond. 1800; and Dr. John Davy, Researches, Physiological and Anatomical, Dunglison's Amer. Med. Libr. Edit. p. 48, Philad. 1840.

<sup>&</sup>lt;sup>c</sup> Dissert. sistens Sang. Coag. Histor. Gröning. p. 81, 1820; and Burdach, op. citat. iv. 37.

which coagulates firmly in five minutes at 60° Fahr., will remain quite fluid for twenty minutes at the temperature of 40° Fahr., and requires upwards of an hour for complete coagulation. The observations of Gendrina were similar. As a general rule, it would seem, from the experiments of Hewson, b Schröder van der Kolk, and Thackrah, that coagulation takes place most readily at the temperature of the body. During the coagulation, a quantity of caloric is disengaged. Fourcroye relates an experiment, in which the thermometer rose no less than 11° during the process; but as certain experiments of Hunter appeared to show, that no elevation of temperature occurred, the observation of Fourcroy was disregarded. It was, however, confirmed by some experiments of Dr. Gordon, of Edinburgh, in which the evolution of caloric during coagulation was rendered more manifest by moving the thermometer during the formation of the clot, first into the coagulated, and afterwards into the fluid part of the blood: he found, that by this means he could detect a difference of 6°; which continued to be manifested for twenty minutes after the process had commenced. In repeating the experiment on blood, taken from a person labouring under inflammatory fever, the thermometer was found to rise 12°. Sir C. Scudamore affirms,h that the rate at which blood cools is distinctly slower than it would be were no caloric evolved; and that he observed the thermometer to rise one degree at the commencement of coagulation. On the other hand, Dr. John Davy, Mr. Thackrah, and Schröder van der Kolk, accord with Mr. Hunter in the belief, that the increase of temperature, from this cause, is very slight or null, whilst Raspail asserts that the temperature falls.k Again we have to deplore the discordance amongst observers; and it will perhaps have struck the reader more than once, that such discordance applies as much to topics of direct observation as to those of a theo-The discrepance, regarding anatomical and retical character. physical facts, is even more glaring than that which prevails amongst physiologists in accounting for the corporeal phenomena; a circumstance, which tends to confirm the notion promulgated by one of the most distinguished teachers of his day, (Dr. James Gregory,) that "there are more false facts in medicine, (and the remark might be extended to the collateral or accessory sciences.) than false theories."

There are certain substances, again, which, when added to the

\* Hist. Anatom. des Inflammations, ii. 426, Paris, 1826.

4 Inquiry into the Nature, &c. of the Blood, p. 38, Lond. 1819.

h An Essay on the Blood, p. 68, Lond. 1824.

Müller's Physiology, Baly's translation, p. 98, Lond. 1838.

k Chimie Organique, p. 361.

b Experiment. Inquiries, i. 19, Lond. 1774.

Annales de Chimie, xii. 147.
 A Treatise on the Blood, &c. p. 27, Lond. 1794.
 Annals of Philosophy, vol. iv. 139.

<sup>&</sup>lt;sup>1</sup> Researches, Physiological and Anatomical, Dunglison's Amer. Med. Lib. Edit. p. 6, Philad. 1840.

blood, prevent or retard its coagulation. Hewson found, that the sulphate of soda, the chloride of sodium, and the nitrate of potassa were amongst the most powerful salts in this respect. The muriate of ammonia and a solution of potassa have the same effect. the contrary, coagulation is promoted by alum, and by the sulphates of zinc and copper.a How these salts act on the fibrin, so as to prevent its particles from coming together, it is not easy to explain. But these are not the only inscrutable circumstances that affect the coagulation of the blood. Many causes of sudden death have been considered to have this result : - lightning and electricity; a blow upon the stomach; injury of the brain; the bites of venomous animals; certain narcotico-acrid vegetable poisons; also, excessive exercise and violent mental emotions, when they suddenly destroy, &c. Many of these affirmations doubtless rest on insufficient proof. Sir C. Scudamore, for example, asserts that lightning has not this effect. Blood, through which electric discharges were transmitted, coagulated as quickly as that which was not electrified; and, in animals, killed by the discharge of a powerful galvanic battery,

the blood in the veins was always found in a solid state.

M. Mandl has summed up the results of modern experiments on this subject as follows. First. The alkalies—potassa, soda, and ammonia - completely prevent coagulation: lime retards it. Secondly. The soluble alkaline salts — combinations of soda, potassa, ammonia, magnesia, baryta and lime, with carbonic, acetic, nitric, phosphorie, tartaric, citric, boracic, sulphuric and cyanohydric acid, also the chlorides, in very small quantity, favour coagulation. On the other hand, these substances in concentrated solution retard it, and even prevent it entirely. The most active salts are the carbonates; the least so, the combinations of chlorine, and sulphates. 0.007 of carbonate of soda retards coagulation for several hours, whilst the sulphates do not act in the proportion of 14 per 1000. The action of a salt is more marked in proportion as it reddens more the blood; whilst the combinations of chrome, chlorine and iodine do not redden it, and do not prevent its coagulation. When water is added to blood thus liquefied by a salt it coagulates again - the fibrin being precipitated. Thirdly. Metallic salts decompose the blood; some causing coagulation; others preventing it. Fourthly. Very dilute vegetable acids favour coagulation; when a little more concentrated, they prevent it; and when highly concentrated, they decompose it like the mineral acids. Fifthly. The action of the vegetable substances has not been sufficiently studied: some affirm, for instance, that narcotics prevent coagulation; others that they favour it. The same doubt exists in regard to the action of poisons; it is generally believed, however, that they - as well as lightning, a violent discharge of electricity, the instantaneous destruction of the nervous system, &c .- prevent coagulation. Sixthly. Very dilute solutions of gum Arabic, sugar, albumen, milk, &c., appear to act

a Magendie, Lectures on the Blood, in London Lancet, reprinted in Bell's Select Medical Library, Philad. 1839.

only in a mechanical manner by preventing the approximation of

the coagulated particles.

We shall find, hereafter, that the presumed action of some of these agents has been considered evidence that the blood may be killed; and, consequently, that it is possessed of life. All the phenomena, indeed, of coagulation, inexplicable in the present state of our knowledge, have been invoked to prove this position. The preservation of the fluid state, whilst circulating in the vessels although agitation, when it is out of the body, does not prevent its coagulation - has been regarded of itself, sufficient evidence in favour of the doctrine. Dr. Bostock, indeed, asserts, that perhaps the most obvious and consistent view of the subject is, that fibrin has a natural disposition to assume the solid form, when no circumstance prevents it from exercising this inherent tendency. it is gradually added to the blood, particle by particle, whilst that fluid is in a state of agitation in the vessels, it has no opportunity, he conceives, of concreting; but when it is suffered to lie at rest, either within or without the vessels, it is then liable to exercise its natural tendency. It is not our intention, at present, to enter into the subject of the vitality of the blood. The general question will be considered in a subsequent part of this work. We may merely observe, that, by the generality of physiologists, the blood is presumed, either to be endowed with a principle of vitality, or to receive from the organs, with which it comes in contact, a vital impression or influence, which, together with the constant motion, counteracts its tendency to coagulation. Even Magendie, - who is unusually and properly chary in having recourse to this method of explaining the notum per ignotius, - affirms, that instead of referring the coagulation of the blood to any physical influence, it should be considered as essentially a vital process; or, in other words, as affording a demonstrative proof, that the blood is endowed with life.

Within a few years, Vauquelin has discovered in the blood a considerable quantity of fatty matter, of a soft consistence, which he, at first, considered to be fat; but Chevreul, after careful investigation, declared it to be identical with the matter of the brain and nerves, and to form the singular compound of an azoted fat. Cholesterin has been detected in it by Gmelin, and by Boudet. Prévost and Dumas, Ségalas, and others have likewise demonstrated the existence of urea in the blood of animals, from which the kidneys had been removed. Chemical analysis is, indeed, adding daily to our stock of information on this matter; and is exhibiting to us, that many of the substances, which compose the tissues, exist in the very state in the blood in which we meet with them in the tissues. This is signally shown in the analysis of the

Physiology, 3d edit. p. 271, Lond. 1836.

b J. Müller's Handbuch, u. s. w. Baly's translation, p. 97, Lond. 1838.

c Précis, &c., ii. 234. d Bostock's Physiology, p. 294. c Chimie, iv. 1163.

Journ. de Pharmacie, Paris, 1833, and Annales de Chimie, lii. 337.

blood by M. Lecanu, who found it, in one analysis, to be composed—in 1000 parts—of water, 785.590; albumen, 69.415; fibrin, 3.565; colouring matter, 119.626; crystallizable fatty matter, 4.300; oily matter, 2.270; extractive matter, soluble in alcohol and water, 1.920; albumen combined with soda, 2.010; chlorides of sodium and potassium, alkaline phosphate, sulphate, and subcarbonates, 7.304; subcarbonate of lime and magnesia, phosphates of lime, magnesia, and iron, peroxide of iron, 1.414; loss, 2.586. On this analysis, Dr. Prout has remarked, that gelatin is never found in the blood, or any product of glandular secretion, and he adds, that a given weight of gelatin contains at least three or four per cent. less carbon than an equal weight of albumen. Hence, the production of gelatin from albumen, he conceives, must be a reducing process. We have seen, under the head of Respiration, what application he makes of these considerations.

Dutrochet believed that he had formed muscular fibres from albumen by the agency of galvanism; and he supposed that the red particles of the blood formed each a pair of plates, the nucleus being negative, the envelope positive: but Müller has shown, that all the appearances, which he attributed to different electric properties of the blood are explicable by the precipitation of the albumen and fibrin, in consequence of the decomposition of the salts of the serum, and of the oxidation of the copper wire used in the experiments, — both the decomposition of the salts and the oxidation of the copper being the usual effects of galvanic action. With the galvanometer he was unable to discover any electric current in the blood: he perceived no variation in the needle of the multiplicator, even when he inserted one wire into an artery of a living

animal, and the other into a vein.

Lastly,—some interesting experiments and considerations on the blood have been published by Dr. Benjamin G. Babington. The principal experiment was the following. He drew blood, in a full stream, from the vein of a person labouring under acute rhematism, into a glass vessel filled to the brim. On close inspection, a colourless fluid was immediately perceived around the edge of the surface, and after a rest of four or five minutes, a bluish appearance was observed forming an upper

<sup>a</sup> Annales de Chimie et de Physique, xlviii. 308, and Journal de Pharmacie, Sept. 1831.

b For an analysis of the Blood by Boudet, see Journal de Pharmacie, Juin, 1833.
c Bridgewater Treatise, Amer. Edit. p. 280, Philad. 1834; see, also, J. Müller, op.

citat. 133; Thomson, Animal Chemistry, p. 349, Edinb. 1843.

d See, also, Bellingeri Experimenta in Electricitatem Sanguinis, Urinæ et Bilis Animalium, Aug. Taurin. 1826 — cited by J. Müller, loc. cit.; and Burdach, Die Physiologie als Erfahrungswissenschaft, iv. 15 und 103, Berlin, 1832.

e Handbuch, u. s. w., Baly's translation, p. 133.

<sup>t</sup> Med. Chirurg. Transact. vol. xvi. part 2, Lond. 1831; art. Blood (Morbid Conditions of the) in Cyclop. Anat. and Physiol., Lond. 1836 — reprinted in Dunglison's American Medical Library and Intelligencer, for April, 1837. See, also, on this subject, the views of Dr. John Davy, in his Researches, Physiological and Anatomical, Dunglison's Amer. Medical Library edit., p. 13, Philad. 1840.

BLOOD, 109

layer on the blood, which was owing to the subsidence of the red particles to a certain distance below the surface, and the consequent existence of a clear liquor between the plane of the red particles and the eye. A spoon, previously moistened with water, was now immersed into the upper layer of liquid, by a gentle depression of one border. The liquid was thus collected quite free from red particles, and was found to be an opalescent, and somewhat viscid solution, perfectly homogeneous in appearance. By repeating the immersion, the fluid was collected in quantity, and transferred to another vessel. That, which Dr. Babington employed, was a bottle, holding about 180 grains, of globular form, with a narrow neck and perforated glass stopper. The solution, with which the globular bottle was filled, though quite homogeneous at the time it was thus collected, was found, after a time, to separate into two parts, viz., into a clot of fibrin, which had the precise form of the bottle into which it was received, and a clear serum, possessing all the usual characters of the fluid. From this experiment, Dr. Babington infers, that buffed blood, to which we shall have to refer under another head, consists of only two constituents, the red particles, and a liquid to which he gives the name, liquor sanguinis - the plasma of Schultza - so called by him, because he esteems it to be the true nutritive and plastic portion of the blood, from which all the organs of the body are formed and nourished.

It has long been observed, that the blood of inflammation is longer in coagulating than the blood of health, and that the last portion of blood drawn from an animal, coagulates the quickest. The immediate cause of this buffy coat is thus explained by Dr. Babington. The blood, consisting of liquor sanguinis and insoluble red particles, preserves its fluidity long enough to permit the red particles, which are of greater specific gravity, to subside through the liquor sanguinis. At length, the liquor sanguinis separates, by a general coagulation and contraction, into two parts, and this phenomenon takes place uniformly throughout the liquor. That part of it, through which the red particles had time to fall, furnishes a pure fibrin or buffed crust, whilst the portion, into which the red particles had descended, furnishes the coloured clot. This, in extreme cases, may be very loose at the bottom, from the great number of red particles collected there, each of which has supplanted its bulk of fibrin, and consequently diminished its firmness in that part. There is, however, with this limitation, no more fibrin in one part of the blood than another.

It is a well known fact, that the shape of the vessel, into which the blood is received, influences the depth of the buff. The space, left by the gravitation of the red particles, bears a proportion to the whole perpendicular depth of the blood, so that in a shallow vessel scarcely any buff may appear, whilst the same blood in a

See an analysis of Professor Schultz's Views, in Lond. Lancet, Aug. 1839, p. 712.
 Dr. J. Davy, op. citat. p. 45.

vol. II. - 10

deep vessel would have furnished a crust of considerable thickness; but Dr. Babington asserts, that even the quantity of the crassamentum is dependent, within certain limits, on the form of the vessel. If this be shallow, the crassamentum will be abundant; if approaching the cube or sphere in form, it will be scanty. The difference is owing to the greater or less distance of the coagulating particles of fibrin from a common centre, which causes a more or less powerful adhesion and contraction of these particles. This is a matter of practical moment, inasmuch as blood is conceived to be thick or thin, rich or poor, in reference to the quantity of crassamentum; and pathological views are entertained in consequence of conditions, which after all depend not perhaps on the blood

itself, but on the vessel into which it is received.

To remove an objection, that might be urged against a general conclusion deduced from the experiment cited, - that it was made upon blood in a diseased state, Dr. Babington received some healthy blood into a tall glass vessel, half filled with oil, which enabled the red particles to subside more quickly than would otherwise have been the case. This blood was found to have a layer of liquor sanguinis, which formed a buffy coat, whilst a portion of the same blood, received into a similar vessel, in which there was no oil, had no buff. Hence, it would appear, that healthy blood is similarly constituted as blood disposed to form a buffy coat, the only difference being, that the former coagulates more quickly than the latter. Dr. J. Davy, however, has observed, that inflammatory blood, in some instances, does not coagulate more slowly than healthy blood, and as from the experiments of J. Müller it would appear that the presence of fibrin in the blood favours the subsidence of the red particles, Müller was led to infer, that the formation of the buffy coat may arise from the blood containing a larger quantity of fibrin, which the blood of inflammation is known to do. So that the principal causes, he thinks, of the subsidence of the red particles and the formation of the buffy coat in inflammatory blood appear to be - the slow coagulation of the blood and the increased quantity of fibrin.

The most correct view is, perhaps, that of Andral, that the essential condition of the buffy coat is an increase in the quantity of fibrin in proportion to the red particles. Hence, if there be an absolute increase of fibrin, the red particles remaining the same, as in inflammation; or, if there be a diminution in the proportion of the red particles, the fibrin remaining the same, the buffy coat may

result.

\* Philosophical Transactions, for 1822.

cine, Book 3, 2d edit., Philad. 1844.

b Op. citat. p. 117; and Magendie, Leçons sur le Sang, &c., or translation in Lond. Lancet, and in Bell's Med. Libr. edition, p. 77, Philad. 1839.

c Dr. J. Davy, Researches, &c. p. 28; see, connected with the buffy coat and the life of the blood, the remarks in the latter part of this volume, under the head of Life.
d Hématologie Pathologique, p. 75, Paris, 73. See, also, Carpenter, Human Physiology, § 589, Lond. 1842, or Amer. Edit. p. 430; and the author's Practice of Medi-

BLOOD. 111

It need scarcely be said, that venous blood must differ some what in its character in the different veins. In its passage through the capillary or intermediate circulation, the arterial blood is deprived of several of its elements, but this deprivation is different in different parts of the body. The blood, for example, which returns from the salivary glands, must vary from that which returns from the kidneys. In the blood of the abdominal venous system, the greatest variation is observed. Professor Schultza has, of late, inquired into the chemical and physiological differences between the blood of the vena portæ and that of the arteries and other veins. He found, that it is not reddened by the neutral salts, or by exposure to the atmosphere, or to oxygen; that it does not generally coagulate; that it contains 5.23 per cent. less fibrin; proportionably more cruor, and less albumen; and has twice as much fat in its solid parts as that of the arteries and the other veins; the proportions being as follows:

Blood of the vena portæ - - 1.66 per cent.
— of the arteries - - 0.92
— of the other veins - - 0.83

The character and quantity of the different constituents of the blood, as well as its coagulation, vary greatly in disease; and the investigation is one of the most important in the domain of pathology. It is one which has attracted the attention of modern pathologists, and especially of MM. Andral and Gavarret, who have endeavoured to detect the changes that occur in disease in the amount of the organic elements of the fluid. These the author has referred to in their appropriate places in another work.c The usual proportions of each element, in 1000 parts of healthy blood, are as follows: Fibrin, 3; red particles, 127; solid matter of serum, 80; water, 790. The proportion of fibrin may perhaps vary, within the limits of health, from 2½ to 3½ parts in a thousand. The amount of red particles appears to be subject to greater variation, within the limits of health, than that of the fibrin. The maximum is about 140, but this is connected with a plethoric condition; the minimum is about 110. Strength of constitution contributes most to raise the globules towards the maximum; whilst debility, congenital or acquired, diminishes them towards the minimum proportion. The solid matter of the serum likewise varies, but

Rust, Magazin für die gesammt. Heilkund. Bde. 44, H. i. See, also, Ancell's Lectures on the Physiology, &c. of the Blood, Feb. 1, 1840, p. 682; and Prof. Schultz,

in Lond. Lancet, Aug. 10, 1839, p. 717.

c Practice of Medicine, 2d edit. Philad. 1844.

b See, on this subject, Babington, op. citat.; G. O. Rees's Analysis of the Blood and Urine, in Health and Disease, Lond. 1836; Mr. E. A. Jennings's Report on the Chemistry of the Blood, as illustrating its Pathology, in Transact. of the Provincial Medical and Surgical Association, vol. iii., Lond. 1835; Giacomini, De la Nature, de la Vie, et des Maladies du Sang, — Monograph. in Encyclographie des Sciences Médicales, Avril, 1840; L. Mandl, Manuel d'Anatomie générale, 271, Paris, 1843; MM. Andral and Gavarret, Archiv. Général. Série 3, tom. viii. p. 501; and Andral, Hématologie Pathologique, Paris, 1843.

there is a certain point of diminution below which they do not pass in health.a

## 3. PHYSIOLOGY OF THE CIRCULATION.

The blood, contained in the circulatory apparatus, is in constant motion. The venous blood, brought from every part of the body, is emptied into the right auricle; from the right auricle it passes into the corresponding ventricle; the latter projects it into the pulmonary artery, by which it is conveyed to the lungs, passing through the capillary system into the pulmonary veins. These convey it to the left auricle; from the left auricle it enters the corresponding ventricle; and the left ventricle sends it into the aorta, along which it passes to the different organs and tissues of the body, through the general intermediate or capillary system, which communicates with the veins; these last vessels return the blood to the part whence it set out. This entire circuit includes both the lesser and

the greater circulation.

It was not until the commencement of the seventeenth century, that any precise ideas were entertained regarding the general circulation. In antiquity, the most erroneous notions prevailed; the arteries being generally looked upon as tubes for the conveyance of some aërial fluid to, and from, the heart, whilst the veins conducted the blood, but whither or for what precise purpose was not understood. The names, given to the principal arterial vessel the aorta - and to the arteries, sufficiently show the functions originally ascribed to them, both being derived from the Greek, and, "air," and raper, "to keep;" and this is farther confirmed by the fact, that the trachea or windpipe was originally termed an artery, - the appropria toaxes of the Greeks, - the aspera arteria of the Latin writers. In the time of Galen, however, the arteries were known to contain blood; and he seems to have had some notions of a circulation. He remarks, that the chyle, the product of digestion, is collected by the meseraic veins and carried to the liver, where it is converted into blood; the supra-hepatic veins then convey it to the pulmonary heart; thence it proceeds in part to the lungs, and the remainder to the rest of the body, passing through the median septum of the auricles and ventricles. This limited knowledge of the circulation continued through the whole of the middle ages; the functions of the veins being universally misapprehended; and the general notion being, that they also convey blood from the heart to the organs; from the centre to the circumference. It was not until after the middle of the sixteenth century, that the lesser circulation, or that through the lungs, was comprehended, by the labours of Michael Servetusb - who fell a

a Andral, Hématologie Pathologique, p. 29, Paris, 1843.

b See the unnoticed theories of Servetus, by Geo. Sigmond, M.D., Lond. 1828; Hecker, Lehre vom Kreislauf von Harvey, Berlin, 1831; Haller, Element. Physiol. iv. 4, 17; Sprengel's Hist. de Médecine, par Jourdan, tom. iv.; Dr. J. R. Coxe. Inquiry into the Claims of Dr. W. Harvey to the Discovery of the Circulation, &c., Philad. 1834; and Gerdy, in art. Circulation, Diction. de Médecine, 2de édit. viii. 68, Paris, 1834.

victim to the persecution and intolerance of Calvin, - and of Andrew Caesalpinus, and Realdus Columbus. It has, indeed, been imagined, that they possessed some notion of the greater circulation. However this may have been, all nations unite in awarding to Harvey the merit, if not of entire originality, of at least of having first clearly described it. The honour of the discovery is, therefore, his; and by it his name has been rendered immortal, for its importance to the physiology and pathology of the animal fabric is overwhelming. How vague and inaccurate must have been the notions of the earlier pathologists regarding the doctrine of acute diseases, in which the circulation is always largely affected, diseases, which, according to the estimate of some writers, constitute two-thirds of the morbid states to which mankind are liable. It was in the year 1619, that Harvey attained a full knowledge of the circulation; but his discovery was not promulgated until the year 1628; in a tract, to which the merit of clearness, perspicuity and demonstration has been awarded by all.a Yet so strong is the force of prejudice, and so difficult is it to discard preconceived notions, that it was remarked, according to Hume, b that no physician in Europe, who had reached forty years of age, ever, to the end of his existence, adopted Harvey's doctrine of the circulation; and Harvey's practice in London diminished extremely for a time from the reproach drawn upon him by that great and signal discoverv.c

Of the truth of the course of the blood, as established by Harvey, we have numerous and incontestable evidences, which it may now be almost a work of supererogation to adduce. Of these the following are some of the most striking. First. If we open the chest of a living animal, we find the heart alternately dilating and contracting so as manifestly to receive and expel the blood in reciprocal succession. Secondly. The valves of the heart, and of the great arteries, which arise from the ventricles, are so arranged as to allow the blood to flow in one direction, and not in another; and the same may be said of those veins, which are directed towards the heart. The tricuspid valve permits the blood to flow only from the right auricle into the corresponding ventricle; the sigmoid valves admit it to enter the pulmonary artery, but not to return; and, as there is, in the adult, no immediate communication between the right and left sides of the heart, the blood must pass along the pulmonary artery and by the pulmonary veins to the left auricle. The mitral valve, again, is so situate, that the blood can only pass in one direction from auricle to ventricle; and, at the mouth of the aorta, the same valvular arrangement exists, as at the month of the pulmonary artery, permitting the blood to proceed along the artery, but preventing its reflux. Thirdly. If an artery and a

<sup>\*</sup> Exercitat. Anatom. de Motû Cordis et Sanguinis, Francof, 1628, Glasguæ, 1751.

b History of England, chap. lxii.
c See, also, Purkinje, in art. Circulatio Sanguinis, in Encyclop. Wörterb. der Medicin.
Wissenschaft. vii. 695, Berlin, 1831.

vein be wounded, the blood will be observed to flow from the part of the vessel nearest the heart in the case of the artery; from the other extremity in that of a vein. The ordinary operation of bloodletting at the flexure of the arm affords us an elucidation of this. The bandage is applied above the elbow, for the purpose of compressing the superficial veins, but not so tightly as to compress, also, the deep-seated artery. The blood passes along the artery to the extremity of the fingers, and returns by the veins, but its progress back to the heart by the subcutaneous veins being prevented by the ligature, they become turgid; and, if a puncture be made, the blood flows freely. If, however, the ligature be applied so forcibly as to compress the main artery, the blood no longer flows to the extremity of the fingers; there is none, consequently, to be returned by the veins; they do not rise properly; and if a puncture be made no blood flows. This is not an unfrequent cause of the failure of an inexperienced phlebotomist. If the bandage, under such circumstances, be slackened, the blood will resume its course along the artery, and a copious stream will issue from the orifice, which did not previously transmit a drop. This operation, then, exhibits the fact of the flow of blood along the arteries from the heart, and of its return by the veins. From what has been said, too, it will be obvious, that if a ligature be applied to both vessels, the artery will become turgid above the ligature, the vein below it. Fourthly. The microscopical experiments of Leeuenhoek, Malpighi, Spallanzani, and others, have exhibited to the eye the passage of the blood in successive waves by the arteries towards the veins, and its return by the latter. Lastly. The fact is farther demonstrated by the effects of transfusion of blood, and of the injection of substances into the vessels; both of which operations will be alluded to in another place.a

In tracing the physiological action of the different parts of the circulatory apparatus, we shall follow the order observed in the anatomical sketch; and describe, in succession, the circulation in the heart, in the arteries, in the capillary vessels, and in the veins; on all of which points there has been much interesting diversity of opinion, and much room for ingenious speculation, and for farther

improvement.

## a. Circulation in the Heart.

It has been already observed, that when the heart of a living animal is exposed, it is remarked to undergo alternate contraction and dilatation. The mode, in which the circulation through the heart is accomplished, is generally considered to be as follows:—the blood is received into the two auricles at the same time, and is transmitted into the two great arteries synchronously.

In order that the heart shall receive blood, it is necessary that the auricle should be dilated. This movement is partly, perhaps, effected by virtue of the elasticity which it possesses in its structure.

Bostock's Physiology, 3d edit. p. 213, Lond. 1836.

Let us suppose it to be once filled; the stimulus of the blood excites it to contraction, and the blood is thus sent into the corresponding ventricle. As soon, however, as it has emptied itself, the stimulus is withdrawn: and, by virtue of its elasticity, it returns to the state in which it was prior to contraction. An approach to a vacuum is thus formed in the cavity, and the blood is solicited towards it from the veins, until it is again filled and its contraction is renewed. When the right auricle contracts there are four channels by which the blood might be presumed to pass from it, - the two terminations of the venæ cavæ, the coronary vein, and the auriculo-ventricular communication. The constant flow of blood from every part of the body prevents it from readily returning by the venæ cavæ, whilst the small quantity, which, under other circumstances, might have entered the coronary vein, is prevented by its valve. To the flow of the blood through the aperture into the ventricle, which is in a state of dilatation, there is no obstacle, and accordingly it takes this course, raising the tricuspid valves.

It may be remarked, that physiologists are not entirely of accord regarding the reflux of blood into the venæ cavæ. Some think that this always occurs to a slight extent; others, that it is never present in the physiological or healthy state. Its existence is unequivocal, where an obstacle occurs to the due discharge of the blood into the ventricle. For example, if any impediment exists to the flow of blood along the pulmonary artery, either owing to mechanical obstruction or to diminished force of the ventricle, the reflux will be manifested by a kind of pulsation in the

veins, which Haller has called the venous pulse.

The blood, having attained the right ventricle, by the effort exerted by the contraction of the auricle, and by the aspiration exerted by the dilatation of the cavity through the agency of its elastic structure, the ventricle contracts. Into it there are but two apertures, - the auriculo-ventricular, and the mouth of the pulmonary artery. By the former, much of the blood cannot escape, owing to the tricuspid valve, which acts like the sail of a ship, the blood distending it, as the wind does a sail, and the chordæ tendineæ retaining it in position, so that the greater part of the blood is precluded from reflowing into the auricle. This auriculoventricular valve is not, however, as perfect as that of the left heart. The observations of Mr. Kinga show, that whilst the structure of the mitral valve is adapted to close accurately all communication between the left auricle and ventricle during the contraction of the latter, that of the tricuspid valve is designedly calculated to permit, when closed, the flow of a certain quantity of blood into the auricle. The comparatively imperfect valvular function of the tricuspid was shown by various experiments on recent hearts, in which it was found, that fluids, injected through the aorta into the left ventricle, were perfectly retained in that

<sup>\*</sup> Guy's Hospital Reports, No. iv. for April, 1837; see, also, P. Blakiston, Lond. Med. Gaz., Aug. 1841.

cavity, by the closing of the mitral valve, but that when the right ventricle was similarly injected through the pulmonary artery, the tricuspid valves generally allowed the escape of the fluid in streams, more or less copious, in consequence of the incomplete apposition of their margins. This peculiarity of structure in the tricuspid, Mr. King regards as an express provision against the mischiefs that might result from an excessive afflux of blood to the lungs, - the tricuspid thus acting as a safety valve, and being more especially advantageous in incipient diseased enlargements of the right ventricle. The only other way the blood can escape from the right ventricle is by the pulmonary artery, the sigmoid valves of which it raises. These had been closed like flood-gates, during the dilatation of the ventricle; but they are readily pushed outwards, by the column transmitted from the ventricle. Such is the circulation through one heart, - the pulmonic. The same explanation is applied to the other, - the systemic; and hence it is, that the structure, as well as the functions of the heart, is so much better comprehended, by conceiving it to

be constituted of two essentially similar organs.

The above description is that which is usually given of the circulation through the heart. There is great reason, however, for the belief, that too much importance has been assigned to the distinct contraction of the auricles. If we examine their anatomical arrangement we discover, that there are no valves at the mouths of the great veins which open into them, and that, although, in the proper auricular or dog's ear portion, muscular fibres, and columns, - somewhat analogous to those of the columnæ carneæ of the ventricles, and probably destined for similar uses - exist, the parietes of the main portions of the auricles - or those that constitute the venous sinuses - are but little adapted for anything like energetic contraction. In experiments on living animals observation shows, that the rhythmic acts of dilatation and contraction are more signally exhibited by the ventricle, whilst in some monsters the auricles are wanting, and in birds they are very M. d'Espine, too, considers the auricles, in receiving or transmitting blood, to have only a vermicular motion, not one of contraction; and in an interesting case of monstrosity, described by Dr. T. Robinson, of Petersburg, Virginia, no distinct systole and diastole of the auricles could be detected. Besides, if we admit both an active power of dilatation and of contraction in the ventricles, any similar action of the auricles would seem to be superfluous. In the state of active dilatation of the ventricles, the blood is drawn into their cavities; and as soon as they enter into contraction, the auriculo-ventricular valves prevent the farther entrance into them of the blood arriving in the auricles by the large veins, and give occasion to the distension of the auricles; in this way, the dilatation of the auricles, synchronous with the contraction of the ventricles, is accounted for. As soon as the ven-

American Journal of the Medical Sciences, No. xxii. for February, 1833.

tricle has emptied itself of its blood, it dilates actively; the blood then passes suddenly from the auricle into its cavity through the

auriculo-ventricular opening.

From careful experiments instituted by Drs. Pennock and Moore, a they drew the following conclusion: — the ventricles contract and the auricles dilate at the same time, occupying about one-half the whole time required for contraction, diastole, and repose. Immediately at the termination of the systole of the ventricle, its diastole succeeds, occupying about one-fourth of the whole time, synchronously with which the auricle diminishes, by emptying a portion of its blood into the ventricle, but without muscular contraction. The remaining fourth is devoted to the repose of the ventricles, near the termination of which the auricle contracts actively, with a short, quick motion, thus distending the ventricles with an additional quantity of blood; this motion is propagated immediately to the ventricles, and their systole takes place, thus rendering their contractions almost continuous.

From the termination of their diastole to the commencement of their systole, the ventricles are in a state of perfect repose, their cavities remaining full, but not distended; whilst those of the

auricles are partially so, during the whole time.

It appears probable, that the great use of the auricles —in which we include the sinuses — is to act as true "sinuses" or gulfs for the reception of the blood proceeding from every part of the body; — and that little effect is produced on the circulation by their varying condition.<sup>b</sup>

The state of the heart in which the ventricles are dilated is termed its Diastole; that, in which they are contracted, its Systole.

Since the valuable improvement, introduced by Laënnec in the discrimination of diseases of the chest by audible evidences, it has been discovered, that the heart is not in a state of incessant activity, but that it has, like other muscles, its intervals of repose. If we apply the ear or the stethoscope, to the præcordial region, we hear, first, a dull, lengthened sound, which according to Laennec is synchronous with the arterial pulse, and is produced by the contraction of the ventricles. This is instantly succeeded by a sharp, quick sound like that of the valve of a bellows or the lapping of a dog. To convey a notion of these sounds, Dr. C. J. B. Williams employs the word lubb-dup. The first word of the compound expressing the protracted first sound — and the latter the short second sound. The latter sound corresponds to the interval between two pulsations, and is owing to the contraction of the auricles. The space of time, that elapses between this and the sound of the contraction of the ventricles, is the period of repose. The relative

Medical Examiner, Nov. 2, 1839, and Dunglison's American Medical Intelligencer, Dec. 16, 1839, p. 277.

See, on this subject, Elliotson's Human Physiology, p. 174, Lond. 1840.
 A Treatise on the Diseases of the Chest, translated by Dr. Forbes, 4th edit. Lond. 1834.

duration of these periods is as follows: - one half, or somewhat less, for the contraction of the ventricles; a quarter, or somewhat more, for the contraction of the auricles; and the remaining quarter for the period of total cessation from labour. So that in the twentyfour hours the ventricles work twelve hours and rest twelve; and the auricles work six and rest eighteen. Such is the view of Laënnec; but it is manifestly erroneous. Ocular observation on living animals, as Dr. Alisona has remarked, shows that the emptying of the auricle precedes that of the ventricle, and that the interval of rest is between the contraction of the ventricle, and the next contraction or emptying of the auricle: between the contraction of the auricle, and that of the ventricle, there is no appreciable interval. Pucheltb thinks it most probable that the first sound is caused by the impulse of the blood against the walls of the ventricle during the contraction of the auricles, and the second by the impulse of the blood against the commencement of the arteries during the contraction of the ventricles. M. d'Espine thinks that the first sound is produced by the contraction of the ventricle, and that the second is owing to their dilatation. Our knowledge, indeed, of the. cause of the sounds rendered by the heart, is sufficiently imprecise this is farther proved by the circumstance, that Magendie ascribed the first sound to the shock or impulsion of the apex of the heart during its diastole, and the second to the impulsion of the base of the heart during its systole; but the results of more recent experimentsd lead him to infer, that the first sound is owing to the contraction of the ventricles, and to the impulse of the apex of the heart against the ribs, and the second sound to a similar impulse produced by their dilatation. Rouanete ascribes the first or dull sound to the shock or impulse of the tricuspid and mitral valves against the auriculo-ventricular orifices, and the second or clear sound to the succussion of the blood in the distended agree and pulmonary artery backwards against the semilunar valves, during the dilatation of the ventricles; and a similar opinion is entertained. by Dr. Hopef and by Messrs. Mayog and Bouillaud.h Mr. Carlisle and Dr. Williams referred the first sound, with Laënnec, to the systole of the ventricles, and the second to the obstacle presented by the semilunar valves to the return of the blood from the

. Outlines of Physiology, Lond. 1831.

b System der Medicin. th. i. Auflage 2te, s. 149, Heidelb. 1835.

c Revue Médicale, Oct. 1831. d Annales des Sciences Naturelles, 1834.

· Ibid. No. xcvii.

s Outlines of Human Pathology, p. 465, Lond. 1836.

h Journal Hebdomad. No. ix., 1834.

Report of the Third Meeting of the British Association for the Advancement of Science; and Amer. Journal of Med. Sciences, p. 477, for Feb. 1835.

A Rational Exposition of the Physical Signs of Diseases of the Lungs and Pleura,

Amer. Edit. Philad. 1830.

f For an elaborate account of the Experiments on the Action and Sounds of the Heart up to the period of the publication of the work, see Dr. Hope, Treatise on Diseases of the Heart, 3d edit. p. 9, Lond. 1839; or Amer. Edit. by Dr. Pennock, Philad.

arteries into the heart, - and Messrs. Corrigan, Pigéaux, b Stokese and Mackintoshd thought the first sound to be owing to the systole of the venous sinuses, and the second to the systole of the ventricles - an opinion, which Burdache thinks is best founded, but

which, as we have seen, is manifestly erroneous.f

In a case of ectopia cordis, recently described by Cruveilhier, s by applying the finger to the origin of the pulmonary artery, a distinct vibratory thrill was perceived, which corresponded with the ventricular diastole; but no such thrill could be felt by the finger when it was applied to any part of the base of the ventricles. He inferred, therefore, that the first sound cannot be dependent upon the action of the auriculo-ventricular valves. The greatest intensity of the first sound was, indeed, in the same situation as the greatest intensity of the second, that is, at the origin of the large arteries. Dr. Carpenterh thinks the results of these observations of M. Cruveilhier, clearly establish, that the principal cause of the first sound exists at the entrances to the arterial trunks; and it does not seem to him, that any other reason can be assigned for it than the prolonged rush of blood through their orifices, and the throwing back of the semilunar valves, which, in suddenly flapping down again, produce the second sound. M. Cruveilhier states it, in his opinion, to be a uniform occurrence, that disease of the semilunar valves modifies both sounds; - a fact, which the author has long noticed. Without expressing an opinion as to the validity of M. Cruveilhier's conclusion regarding the two sounds of the heart, Dr. Forbes evidently regards it with favour, under the view long maintained by him, that although certainly characteristically different, the two sounds have so great a similarity, and are so allied in time and place, that he could not readily bring his mind to believe, that they do not both depend upon one and the same cause, slightly modified, or, at least, on the different play of the same parts.

Drs. Pennock and Moore, who agree in the main with Dr. Hope, found the first sound, the impulse, and the systole of the ventricles to be synchronous; and the second sound to be synchronous with the diastole of the ventricles. The first sound, they suggest, may be a combination of that caused by the contraction of the auricles, the flapping of the auriculo-ventricular valves, the rush of blood from the ventricles, and the sound of muscular contraction. In four of their experiments, when the heart was removed from the

a Dublin Medical Trans., vol. i., New Series.

e Edinb. Med. and Surg. Journal, vol. xxxiv.

b Bulletin des Sciences Médicales, par Férussac, xxv. 272.

d Principles of Pathology, &c., 2d Amer. edit., ii. 6, Philad. 1837. Die Physiologie als Erfahrungswissenschaft, iv. 219, Leipz. 1832. f Müller's Handbuch, u. s. w., Baly's translation, p. 176, Lond. 1838.

<sup>5</sup> Gazette Méd. de Paris, 7 Aout, 1841, p. 535; and Brit. and For. Med. Rev., Oct. h Human Physiology, § 486, Lond. 1842.

Translation of Laënnec, 4th edit.; and Brit. and For. Med. Rev., loc. cit.

body, the ventricles cut open and emptied of their contents, and the auriculo-ventricular valves elevated, a sound resembling the first was still heard, which they attribute chiefly to muscular contraction. The second sound they refer exclusively to the closure of the semilunar valves by the refluent blood from the aorta and pulmonary artery. "This," they remark, "is proved by the greater intensity of this sound over the aorta than elsewhere, the blood having a strong tendency to return through the valvular opening; by the greater feebleness of the sound over the pulmonary artery, which is short, and soon distributes its blood through the lungs, thus producing but slight impulse upon the valves in the attempt to regurgitate; by the disappearance of the sound when the heart becomes congested and contracts feebly; and, finally, on account of its entire extinction when the valve of the aorta was elevated." The main results of the experiments of Drs. Pennock and Moore accord closely with what the author has entertained and taught on this subject, but the views of M. Cruveilhier are well worthy of attention. The whole matter is still open for further investigation. Whilst these sheets, indeed, are passing through the press, a case of thoracic ectopia is published by M. Monod, bin which it is stated, that the maximum of intensity of the first sound did not occur at the base of the ventricles, but at the middle of their fleshy walls; and M. Monod thinks, that it was caused by the shock of the walls of the ventricles against the internal fleshy columns at the moment of contraction. As to the second sound he is of opinion, that it was owing to the return of the wave of blood against the semilunar valves.

It has been a question with physiologists, whether the cavities of the heart completely empty themselves at each contraction. Senac, and Thomas Bartholine, from their experiments, were long ago led to answer the question negatively. On the other hand, Haller, entertained an opposite opinion, — suggested, he remarks, by his experiments; but, perhaps, notwithstanding all his candour, connected, in some manner, with his doctrine of irritability, which could not easily admit the presence of an irritant in a cavity which had ceased to contract. It has been remarked by Magendie, that if we notice the heart of a living animal, whilst it is in a state of action, it is obvious, that the extent of the contractions cannot have

a See Elliotson, Human Physiology, part i. p. 175, Lond. 1840; Hope, op. citat.; Gerhard, on the Diagnosis of Diseases of the Chest, Philad. 1836; Bouillaud, Traité Clinique des Maladies du Cœur, Paris, 1835; Raciborski, Manual of Auscultation, by Fitzherbert, p. 102, Lond. 1835; Piorry, Traité de Diagnostie, § 273, 2de édit., Bruxelles, 1838; C. J. B. Williams, Lectures on the Chest, in Lond. Lancet, reprinted in Bell's Select Medical Library, Philad. 1839; Drs. Williams, Todd and Clendinning, in Lond. Med. Gaz. Dec. 10, 1836; and W. H. Walshe, Physical Diagnosis of Diseases of the Lungs, Amer. Edit. Philad. 1843.

<sup>&</sup>lt;sup>b</sup> Bullet. del. Académ. Royale de Méd. 7, Février, 1843; and Edinb. Med. and Surg. Journ., July, 1843.


c Traité de la Structure du Cœur, &c. 2d édit. Paris, 1774.

d Dissertat. de Corde, Hafn. 1648. Clement. Physiol. iv.

the effect of completely emptying the ventricles; but it must, at the same time, be admitted, that such experiments are inconclusive, inasmuch as they exhibit to us the action of the organ under powerfully deranging influences, and such as could be readily conceived to modify materially the extent of the contractions. They certainly are insufficient to prove, that, whilst an animal is in a physiological condition, the auricles and ventricles are not emptied of their contents by their contraction. The quantity of blood consequently propelled at each ventricular contraction, must fall short of two fluid ounces.

The objection that has been urged against the opposite view, that there would always be stagnant blood in the cavities of the heart, is not valid. The experiments of Venturi, have shown, that even in

an ordinary hydraulic apparatus, the motion of a stream, passing through a vessel of water, is communicated to the fluid, which is at rest in the vessel, so that an incessant change is produced. -- Let us suppose a stream of water



to enter the vessel D E F B, Fig 167, which is full of fluid, by the pipe A C, and that opposite to this pipe is the tube S M B R. The stream will pass up this tube higher than the vessel, and discharge itself at B V. At the same time, the fluid in the vessel will be observed to be in motion, and, in a few seconds, the level in the vessel will fall from D B to H M.

During the systole of the heart, the organ is suddenly carried forward; and although it appears to be rendered shorter, its point strikes the left side of the chest opposite the interval between the fifth and seventh true ribs; producing what is called the "beating of the heart." The cause of this phenomenon was, at one period, a topic of warm controversy. Borelli, Winslow, and others, affirmed, that it was owing to the organ being elongated during contraction; but to this it was replied by Bassuel, that if such elongation took place, the tricuspid and mitral valves, kept down by the columnæ carneæ, could not possibly close the openings between the corresponding auricles and ventricles. Experiments by

Sur la Communication Latérale du Mouvement dans les Fluides, Paris, 1798;
 and Sir C Bell, in Animal Mechanics, p. 35, Library of Useful Knowledge, Lond. 1829.
 De Motû Animalium, Lugd. Bat. 1710.
 Magendie, Précis, &c. ii, 395.

Drs. Pennock and Moore, a exhibited to them, that the expulsion of the blood from the ventricles was effected by an approximation of the sides of the heart, and not by a contraction of the apex towards the base; and that, during the systole, the heart performs a spiral movement and becomes elongated. Senacb ascribed the beating of the heart to three causes, and his views have been adopted by most physiologists: - 1, to the dilatation of the auricles, which occurs during the contraction of the ventricles; 2, to the dilatation of the aorta and pulmonary artery by the introduction of the blood, sent into them by the ventricles; and 3, to the straightening of the arch of the aorta, owing to the blood being forced against it by the contraction of the left ventricle. Dr. William Hunter, considered the last cause quite sufficient to explain the phenomenon, and many physiologists have assented to his view. Sir David Barryd instituted some experiments upon this subject. He opened the thorax of a living animal, and by passing his hands into the cavity, endeavoured to ascertain the actual condition of the heart and great vessels, as to distension and relative position. He performed seven experiments of this kind, from which he concluded, that the vena cava is considerably increased in size during inspiration, which he ascribes, as will be better understood hereafter, to the partial vacuum then formed in the chest. He supposes, that the force exerted by the venous blood on entering the heart, in consequence of the expansion of the chest and the great vessels behind the heart, pushes the organ forwards, and thus causes it to strike against the ribs. Drs. Pennock and Moore, however, in their experiments, found the impulse to be synchronous with and caused by the contraction of the ventricles, and that, when felt externally, it arose from the striking of the apex of the heart against the thorax. This is probably the true explanation; yet Müller thinks that great uncertainty rests as to whether the impulse be produced during the contraction or dilatation of the ventricles.

The systole of the heart is admitted by all to be active. Some physiologists are disposed to think the diastole passive, — that is, the effect of relaxation of the fibres or of the cessation of contraction. Pechlin, Perrault, Hamberger, d'Espine, Alison, and numerous others, have supported an opposite view; — affirming that direct experiment on living animals shows, that positive effort is exerted at the time of the dilatation of the cavities; — a view strikingly confirmed by the case of monstrosity related by Dr. Robinson.<sup>g</sup> His opinion is, that the force of the diastole was in

<sup>2</sup> Med. Examiner, Nov. 2, 1839.

John Hunter, Treatise on the Blood, &c.

Handbuch, u. s. w., Baly's translation, p. 175, Lond. 1838.

b Traité de la Structure du Cœur, &c., Paris, 1749.

Exper. Researches on the influence of atmospheric pressure upon the circulation, Lond. 1826.

Op. citat.

Amer. Journal of the Medical Sciences, No. xxii. Feb. 1833. See, also, a case of partial Ectopia Cordis, by Dr. John O'Bryen, in Lond. Lancet, for July 7, 1838, p. 520; Cruveilhier, op. cit.; and Monod, Bullet de l'Acad. Royale de Méd., 7 Fév. 1843; and Edinb. Med. and Surg. Journ., July, 1843.

that case, equal to, if not greater than that of the systole. In the case observed by Cruveilhier, the diastole had the rapidity and energy of a very active movement, overcoming pressure exerted upon the heart, so that the hand closed upon it when contracted was opened with violence. It has been suggested, that if the course of all the fibres, composing the muscular parietes of the organ, were better known, this apparent anomaly might perhaps be as easily explained as in the ordinary case of antagonist muscles. It is probable, however, that the active force, exerted in the dilatation of these cavities, is that of elasticity; and that when the contraction of the muscular fibres has ceased, this is aroused to action, and promptly restores the organ to its previously dilated condition. According to this view, the natural state would be that of dilatation. We shall see, hereafter, that this elasticity is probably one of the agents of the circulation of the blood along the vessels.

The cause of the heart's action has been a deeply interesting question to the physiologist, and, in the obscurity of the subject, has given rise to many and warm controversies. From the first moment of fætal existence, at which the heart becomes perceptible, till the cessation of vitality, it continues to move. By many of the ancients this was supposed to be owing to an inherent pulsific virtue, which enabled it to contract and dilate alternately,—a mode of expression, which, in the infancy of physical science, was frequently employed to cover ignorance, and which has been properly and severely castigated by Molière. It was in ridicule of the same failing that Swift represents the action of a smokejack as depending on a meat-roasting power.

Descartes<sup>d</sup> imagined that an explosion took place in the ventricles as sudden as that of gunpowder. With equal nescience, the phenomenon was ascribed, by Van Helmont<sup>e</sup> to his imaginary archæus; and by Stahl,<sup>f</sup> and the rest of the animists, to the anima, soul or intelligent principle, which is supposed to preside over all the mental and corporeal phenomena. Stahl was, however, one of the first that attempted any rational explanation of the heart's action. Its muscular tissue; the similarity of its contractions to those of ordinary muscles, with the exception of their not being

a Haller, Elémenta Physiologiæ, ii. 6.

b "Mihi a doctore

Domandatur causam et rationem quare
Opium facit dormire.
A quoi respondeo;
Quia est in eo
Virtus dormitiva,
Cujus est natura
Sensus assoupire."

Le Malade Imaginaire, Intermède iii.

· Fletcher's Rudiments of Physiology, P. ii. a, p. 52, Edinb. 1836.

d Tract. de Homine, p. 167, Amst. 1677. Ortus Medicin, &c. Amstel. 1648.
Theoria vera Medica, Hal. 1737; Sprengel's Hist. de Médecine, par Jourdan, v. 195, Paris, 1815.

voluntary; the fact of its action being modified by the passions, &c., led him to liken its movements to those of muscles. He admitted, that, generally, we possess neither perception of, nor power over, its motions; but he affirmed, that habit alone had rendered them involuntary; in the same manner as certain muscular twitchings or tics, which are at first voluntary, may become irresistible by habit. A strong confirmation of this opinion was drawn from the celebrated case of the honourable Colonel Townshend, (called by Adelona and other French writers, Captain Towson,) who was able, (not all his life, as Adelon asserts, but a short time before his death,) to suspend the movements of his heart at pleasure. This case is of so singular a character, in a physiological as well as pathological point of view, that we shall give it in the words of Dr. George Cheyne, one of the physicians who attended him, and whose character for veracity is beyond suspicion. "Colonel Townshend, a gentleman of excellent natural parts, and of great honour and integrity, had, for many years, been afflicted with constant vomitings, which had made his life painful and miserable. During the whole time of his illness he had observed the strictest regimen, living on the softest vegetables and lightest animal food; drinking asses' milk daily, even in the camp; and for common drink, Bristol water, which, the summer before his death, he had drunk on the spot. But his illness increasing, and his strength decaying, he came from Bristol to Bath in a litter, in autumn, and lay at the Bell Inn. Dr. Baynard, who is since dead, and I were called to him, and attended twice a day for about the space of a week: but, his vomitings continuing still incessant, and obstinate against all remedies, we despaired of his recovery. While he was in this condition, he sent for us early one morning; we waited on him with Mr. Skrine, his apothecary (since dead also); we found his senses clear, and his mind calm; his nurse and several servants were about him. He had made his will and settled his affairs. He told us he had sent for us to give him some account of an odd sensation he had for some time observed and felt in himself, which was that, composing himself, he could die or expire when he pleased, and yet by an effort, or somehow, he could come to life again; which it seems he had sometimes tried before he had sent for us. We heard this with surprise; but as it was not to be accounted for from tried common principles, we could hardly believe the fact as he related it. much less give any account of it; unless he should please to make the experiment before us, which we were unwilling he should do, lest, in his weak condition, he might carry it too far. He continued to talk very distinctly and sensibly above a quarter of an hour, about this (to him) surprising sensation, and insisted so much on our seeing the trial made, that we were at last forced to comply. We all three felt his pulse first; it was distinct, though small and

Physiologie de l'Homme, edit. cit. iii. 302.
Treatise on Nervous Diseases, p. 307.

thready; and his heart had its usual beating. He composed himself on his back, and lay in a still posture, some time. While I held his right hand, Dr. B. laid his hand on his heart, and Mr. S. held a clean looking-glass to his mouth. I found his pulse sink gradually, till at last I could not feel any, by the most exact and nice touch. Dr. Baynard could not feel the least motion in his heart, nor Mr. Skrine the least soil of breath on the bright mirror he held to his mouth. Then each of us, by turn, examined his arm, heart and breath, but could not by the nicest scrutiny discover the least symptom of life in him. We reasoned a long time about this odd appearance as well as we could; and all of us judging it inexplicable and unaccountable; and finding he still continued in that condition, we began to conclude indeed that he had carried the experiment too far, and at last were satisfied that he was actually dead, and were just ready to leave him. This continued about half an hour, by nine o'clock in the morning, in autumn. As we were going away, we observed some motion about the body, and upon examination found his pulse and the motion of his heart gradually returning; he began to breathe gently, and speak softly; we were all astonished, to the last degree, at this unexpected change, and after some further conversation with him, and among ourselves, went away fully satisfied as to all the particulars of this fact, but confounded and puzzled, and not able to form any rational scheme, that might account for it. He afterwards called for his attorney, added a codicil to his will, settled legacies on his servants, received the sacrament, and calmly and composedly expired about five or six o'clock that evening."

It is manifest that this case — unaccountable as it is, in many respects — can add no weight to the views of the Stahlians. It is as strange, as it is inexplicable. The opinion, with them, that the heart's action is a muscular function, was accurate. The error lay in placing it amongst the voluntary functions. It belongs to the involuntary class, equally with many of the muscles concerned in deglutition, and with those of the stomach and intestines; and how well is it for us, as Sir Charles Bell has remarked, that the actions of this and other organs, directly instrumental to the organic functions, are placed out of our control! "A doubt — a moment's pause of irresolution — a forgetfulness of a single action at its appointed time — would otherwise have terminated our existence."

In an oriental journal, Mr. H. M. Twedel<sup>b</sup> has published a case, even more extraordinary than that of Col. Townshend, — of a Hindoo, thirty years of age, who "is said, by long practice, to have acquired the art of holding his breath, by shutting the mouth, and stopping the interior opening of the nostrils with the tongue." This man submitted to be buried for a month, and was dug out alive at the expiration of that period. "He was taken out in a perfectly

<sup>\*</sup> See Fletcher's Rudiments of Physiology, part ii. b, p. 71, Edinb. 1836.

b India Journal of Medical and Physical Sciences; and Amer. Journ. of the Medical Sciences, p. 250, Nov. 1837.

senseless state — his eyes closed, his hands cramped and powerless — his stomach shrunk very much, and his teeth jammed so fast together, that they were forced to open his mouth with an iron instrument to pour a little water down his throat. He gradually recovered his senses, and the use of his limbs, and was restored to

perfect health!

The doctrine of Hallera on the heart's action rested upon the vis insita or irritability to which he referred all muscular contractions, whether voluntary or involuntary. This property, as stated in another place, he conceived to be possessed by muscles as muscles, independently of all nervous influence. The heart, being a muscle, enjoyed it of necessity; and the irritant, which developed it incessantly, was the blood. In evidence of this, he observes, that its contractions are always more forcible and rapid, when the blood is more abundant; and that they occur successively in the cavities of the heart as the blood reaches them. So completely did Haller assign the heart's action to this irritability, that he denied the nerves any influence over it; resting his belief on the admitted facts, - that the heart will continue to beat after decapitation; after the division of the spinal marrow in the neck; and of the nerves distributed to the organ; and, even, after it has been entirely removed from the body. How far the opinions of this great man are correct, respecting the power of contraction residing in the heart, as he conceived it to do in other muscles, we shall inquire presently. The heart, however, is, doubtless, indirectly, under the nervous influence. We see it affected in the various emotions; sometimes augmenting its action violently, at others retarding it. These circumstances have led some individuals to adopt a kind of intermediate opinion, and to regard the nervous influence as one of the conditions necessary for all muscular contraction, just as the due circulation of blood is one of those conditions; and to admit, at the same time, the separate existence of a vis insita. Sommering,b and Behrendsc have, indeed, asserted that the cardiac nerves are not distributed to the tissue of the heart, but merely to the ramifications of the coronary arteries; and hence, that these nerves are not concerned in the functions of the organ, but only in its nutrition; but this is denied by Scarpa, and by the generality of anatomists.e

Although the emotions manifestly affect the heart, direct experiments exhibit but little influence over it on the part of the nerves. This, indeed, we have seen, is one of the grounds for the doctrine of Haller. Willisf divided the eighth pair of nerves; yet the action of the heart persisted for days. Similar results followed the sec-

<sup>\*</sup> Op. citat.

Op. citat.

Dissert. quâ Demonstrat. Cor. Nervis Carere, Mogunt. 1792; and in Ludwigii Script. Neurol. Min. i. 1.

Tabulæ Neurologicæ, &c., Ticin. 1794.

Seiler, in art. Herz, in Anat. Phys. Real. Wörterb. iv. 33, Leipz. 1821; and Mül-

ler's Handbuch, Baly's translation, p. 190, Lond. 1838.

Cerebri Anat. cap. xxiv. in Oper., Genev. 1776.

tion of the great sympathetic. Magendie<sup>a</sup> states, that he removed, on several occasions, the cervical ganglions, and the first thoracic; but was unable to determine any thing satisfactory from the operation in consequence of the immediate death of the animal from such extensive injury as was inevitable. He observed, however, no direct influence on the heart.

We have numerous examples of the comparative independence of the organ, as regards the encephalon. Decapitated reptiles have lived for months; and anencephalous infants or those born with part of the brain only have vegetated during the whole period of pregnancy, and for some days after birth. Legalloisb kept several decapitated mammiferous animals alive; and maintained the heart in action, (having taken the precaution to tie the vessels of the neck for the purpose of preventing hemorrhage,) by employing artificial respiration, so as to keep up the conversion of venous into arterial blood, and thus to insure to the heart a supply of its appropriate fluid. We find, too, that in fracture of the skull in apoplexy, and in congenerous affections, the functions of the heart are the last to be arrested. The result of his own experiments led Legallois to infer, that the power of the heart is altogether derived from the spinal marrow; and he conceived, that through the cardiac nerves it is influenced by this portion of the cerebro-spinal axis, and is liable to be affected by the passions, because the spinal marrow is itself influenced by the brain. Dr. Wilson Philipe has, however, shown, that the facts do not warrant the conclusions; and he has exhibited, by direct experiment, that the brain has as much influence as the spinal marrow over the motions of the heart, when the circumstances of the experiment are precisely the same. The removal of the spinal marrow, like that of the brain, if the experiment be performed cautiously and slowly, does not sensibly affect the motion of the heart, - the animal having been previously deprived of sensibility. In these experiments, the circulation ceased quite as soon without, as with, the destruction of the spinal marrow. Loss of blood appeared to be the chief cause of its cessation; and pain would have contributed to the same effect, if the animal had been operated on, without having been previously rendered insensible.

Mr. Clift, the ingenious conservator of the Museum of the Royal College of Surgeons of London, made a series of experiments to ascertain the influence of the spinal marrow on the action of the heart in fishes, and he found, that, whether the heart be exposed or not, its action continues long after the brain and spinal marrow are destroyed, and still longer when the brain is removed without injury to its substance. Similar results were obtained by Treviranus on the frog, and by Saviole on the chick in ovo. Zinn and Ent too

<sup>&</sup>lt;sup>a</sup> Précis, &c. ii. 401. See, also, Sir A. Cooper, in Guy's Hospital Reports, i. 470, Lond. 1836; Müller, op. citat, p. 198; and Burdach, op. citat. iv. 457.

<sup>Sur le Principe de la Vie, p. 138.
An Experimental Inquiry into the Laws of the Vital Functions, &c. p. 62, Lond.
1817.
Philosoph. Transact. for 1815.</sup> 

found, that after the destruction of the cerebellum, to which Willis

ascribed the heart's action, it continued to beat.a

All these facts plainly exhibit, that, although the heart is indirectly influenced by the brain or spinal marrow, it is not directly acted upon by either one or the other, and that its action can be maintained for some time, after the destruction of one or both, provided artificial respiration be kept up; but this last agent is unnecessary: the heart will continue to beat, even after it has been removed from the body. In the case of the rattlesnake, Dr. Harlanb observed the heart, torn from the body, continue its contractions for ten or twelve hours; and in the monstrous fœtus, observed by Dr. T. Robinson, eits motion continued for some time after the auricles and ventricles had been laid open; the organ roughly handled, and thrown into a basin of cold water. We are compelled, then, if we do not admit the whole of the Hallerian doctrine of irritability, to presume, that there is something inherent in the structure of the heart, which enables it to contract and dilate, when appropriately stimulated; and it is not even necessary, that this should be by the fluid, to which it is habituated. It is certain, that the organ, when separated from the body, may be stimulated to contraction, by being immersed in warm water, or pricked with a sharp-pointed instrument. In some experiments by Sir B. Brodie, d he emptied the heart of its blood, and found that it still contracted and relaxed Similar experiments were instituted by M. Mayo, and with like results, from which he concludes that the alternations of contraction and relaxation in the heart depend upon something in its structure. The conclusion is, indeed, irrefutable, if we add to these evidences the results of some experiments by Prof. J. K. Mitchell, of Philadelphia. In 1823, being engaged in dissecting a sturgeon - Acipenser brevirostrum? - its heart was taken out and laid on the ground, and, after a time, having ceased to beat, was inflated with the breath, for the purpose of drying it. Hung up in this state, it began again to move, and continued for ten hours to pulsate regularly, though more and more slowly; and when last observed in motion, the auricles had become so dry as to rustle when they contracted and dilated. He subsequently repeated the experiment with the heart of a Testudo serpentaria or snapper, and found it to beat well under the influence of oxygen, hydrogen, carbonic acid, and nitrogen, thrown into it in succession. Water also stimulated it, - perhaps more strongly, - but made its substance look pale and hydropic, and, in one minute, destroyed action beyond recovery.

<sup>\*</sup> Burdach, Physiologie als Erfahrungswissenschaft, &c. iv. 454, Leipz. 1832. See, also, Müller, op. citat. p. 191.

Medical and Physical Researches, p. 103, Philad. 1835.
 Amer. Journ. of the Med. Sciences, No. xxii. Feb. 1833.

d Cooke's Treatise on Nervous Diseases, Introd. p. 61, Lond. 1820-23, Amer. Edit., Boston, 1824.

Outlines of Human Physiology, 4th edit. p. 46, Lond. 1837.
 American Journal of the Medical Sciences, vii. 58, Philad. 1830.

It has been supposed, that when the heart is empty of blood, the contact of air with its cavities is the stimulus by which its irritability is excited, but Dr. John Reid, found, when he placed a frog's heart in a state of activity under the receiver of an air-pump, that its action still continued after the receiver had been exhausted.

The heart is the generator of one of the forces that move the blood. This force has been the subject of much calculation, but the results have been so discordant as to throw discredit upon all mathematical investigations on living organs; a circumstance which renders it unnecessary to state the different plans that have been pursued in these estimations. They are all given in the elaborate work of Haller, to which the reader, who may be desirous of examining them, is referred. Borellid conceived the force exerted by the left ventricle to be equivalent to 180,000 pounds; Senace to 40 pounds; Halesf to 51 pounds 5 ounces; Juring to 15 pounds 4 ounces; whilst Keillh conceived it not to exceed from 5 to 8 ounces! The mode adopted by Hales has always been regarded the most satisfactory. By inserting a glass tube into the carotid of various animals, he noticed how high the blood rose in the tube. This he found to be, in the dog, 6 feet 8 inches; in the ram, 6 feet 51 inches; in the horse, 9 feet 8 inches; and he estimated, that, in man, it would rise as high as 7½ feet. Now, a tube, whose area is one inch square and two feet long, holds nearly a pound of water. We may therefore reckon the weight, pressing on each square inch of the ventricle, to be, on a rough estimate, three pounds and three-quarters, or four pounds; and if we consider, with Michelotti, the surface of the left ventricle to be fifteen square inches, it will exert a force, during its contraction, capable of raising sixty pounds.i Its extent is more frequently, however, estimated at 10 square inches, and the force developed would therefore be forty pounds; but this is, of course, a rude approximation. In such a deranging experiment, the force of the heart cannot fail to be modified; and it is so much affected by age, sex, temperament, idiosyncrasy, &c. that the attainment of accurate knowledge on the subject is impracticable. The indefinite character of our information on this matter is sufficiently shown by the investigations of Poiseuille, which led him to suppose, that the force with which the heart propels the blood in the human aorta is about 4 pounds, 3 ounces, and 43 grains. By means of an instrument, which, from its use,

\* Carpenter, Human Physiology, § 480, Lond. 1842.

b Cyclop. of Anat. and Physiol. ii. 611.

- c Elementa Physiologiæ, tom. ii. Lausann. 1757-1766,
  - d De Motû Animalium, part ii. Ludg. Bat. 1710. c Traité de la Structure du Cœur, Paris, 1749.

f Statical Essays, &c., 4th edit. vol. ii.

- Philosophical Transactions, for 1718 and 1719.

  Tentamina Medico-Physica, &c. Lond. 1718.
- Martini, Lezioni di Fisiologia, tomo sesto, p. 420, Torino, 1828; and Arnott's Elements of Physics, Amer. Edit. vol. i. 2d edit. Philad, 1835.

Magendie's Journal de Physiologie, x. 241.

he terms hæmadynamometer, the same physiologist has endeavoured to show, that the blood is urged forward with as great a momentum in a small artery, far from the heart, as in any important branch near it. In other words, that there is a uniform amount of pressure exerted by the blood upon the coats of the arteries in every part of the body;—those in the immediate vicinity of the heart being distended by an equal force with those the most remote from it. M. Poiseuille made the experiment on the carotid, and on the muscular branch of the thigh of the horse, and notwithstanding the very great dissimilarity in the diameter, and distance from the heart, of the two tubes, the displacement of the mercury was exactly the same in both. This infer-

Fig. 168.

Hæmadynamometer.

ence, if correct, - and the experiments have been repeated by Magendieb with corresponding effects, - are important in a therapeutical point of view, as they would lead to the belief, that if it be desirable to lessen the quantity of the circulating fluid, it is of little consequence what vessel is opened. The instrument, employed of M. Poiseuille, consists of a bent glass tube, of the form represented in the marginal figure, filled with mercury in the lower bent part a, d, e. The horizontal part b, provided with a brass head, is fitted into the artery, and a small quantity of a solution of carbonate of soda is interposed between the mercury and the blood, which is allowed to enter the tube with the view of preventing the coagulation of the blood. When the blood is allowed to press upon the fluid in the horizontal limb, the rise of the mercury towards e, measured from the level to which it has fallen towards d, gives the pressure under which the blood moves.

## b. Circulation in the Arteries.

The blood, propelled from the heart by the series of actions we have described, enters the two great bloodvessels; — the pulmonary artery from the right ventricle, and the aorta from the left; the former of which sends it to the lungs, the latter to every part of the system; and, in both vessels, it is prevented from returning into the corresponding ventricles by the depression of the

Magendie, Journal de Physiologie, ix. 46; and art. Circulation, Cyclop. of Anat.

b Leçons sur le Sang, &c., or translation in Lond. Lancet, Sept. 1838 to March 1839; in Bell's Select Medical Library, p. 57, Philad. 1839.

semilunar valves. We have now to inquire into the circumstances, which act upon it in the arteries, or whether it be the contraction of the ventricle, which is alone concerned in its pro-

gression.

Harveya and the whole of the mechanical physiologists regarded the arteries as entirely passive in the circulation, and as acting like so many lifeless tubes; the heart being, in their view, the sole agent in the circulation. We have, however, numerous reasons for believing that the arteries are concerned, to a certain degree, in the progression of the blood. If we open a large artery, in a living animal, the blood flows in distinct pulses; but this effect gradually diminishes as the artery recedes from the heart, and ultimately ceases in the smallest arterial ramifications; -seeming to show, that the force, exerted by the heart, is not the only one concerned in propelling the blood through these vessels. It is manifest, too, that if the action of the heart were alone concerned, the blood ought to flow out of the aperture, when the artery is opened, at intervals coinciding with the contractions of the heart; and that during the diastole of the artery, no blood ought to issue. This, however, is not the case, notwithstanding the authority of Bichat, and some others is in its favour. The flow is uninterrupted, but in jets or pulses, coinciding with the contractions of the ventricles. Again, if two ligatures be put around an arterial trunk, at some distance from each other, and a puncture be made between the ligatures, the blood flows with a jet, - indicating that compression is exerted upon it; and if the diameter of the artery be measured with a pair of compasses, before and after the puncture, it will be found manifestly smaller in the latter case; an experiment which shows the fallacy of a remark of Bichat, that the force with which the arteries return upon themselves is insufficient to expel the blood they contain. An experiment of Magendiec exhibits this yet more clearly. He exposed the crural artery and vein in a dog, and passed a ligature behind the vessels, tying it strongly at the posterior part of the thigh, so that the blood could only pass to the limb by the artery, and return by the vein. He then measured, with a pair of compasses, the diameter of the artery; and, on pressing the vessel between his fingers, to intercept the course of blood in it, the artery was observed to diminish perceptibly in size below the part compressed, and to empty itself of the blood it contained. On readmitting the blood, by removing the fingers, the artery became gradually distended at each contraction of the heart, and resumed its previous dimensions.

These facts prove, that the arteries contract; but the kind of contraction has given occasion to much discussion. It has been imagined, by some physiologists, that their proper coat is muscular, and that they exert a similar action on the blood to that of the

b Magendie, Précis, &c. ii. 388.

<sup>\*</sup> Exercitatio Anat. De Motû Cordis et Sanguinis, &c., Rotterd. 1648.

<sup>&</sup>lt;sup>e</sup> Journal de Physiologie, i. 111; and Précis, &c. ii. 386.

heart; dilating to receive it from that organ, and contracting to propel it onwards; - their systole being synchronous with the systole of the auricles and the diastole of the ventricles, and their diastole with that of the auricles and the systole of the ventricles. The principal reasons, urged in favour of this view, are; - the fact of the circulation being effected solely by the arteries in acardiac fœtuses, and in animals which have no heart; - the assertion of MM. Lamure and Lafosse, that they noticed, in an experiment with the carotid artery, similar to that described above, that the vessel continued to beat between the ligatures; - the affirmations of Verschuir, Bikker, Giulio, and Rossi, Thomson, Parry, Hastings,e Wedemeyer, and numerous others,f that when they irritated arteries with the point of a scalpel or subjected them to the electrical and galvanic influences, they exhibited manifest contractility; and lastly, the fact, that the pulse is not perfectly synchronous in different parts of the body, which ought to be the case, were the arteries not possessed of any distinct action.

The chief objection to the views, founded on the muscularity of the middle coat, is the want of evidence of the fact. In the anatomical proem to the function of the circulation, it was stated, that this coat does not seem to consist of the fibrous or muscular tissue; and that the experiments of Magendie, Nysten, and others, had not been able to exhibit any contraction, on the application of the ordinary excitants of muscular irritability. The chemical analyses of Berzeliusg and Youngh also show, that the transverse fibres differ

essentially from those of proper muscles.

Again, if an artery be exposed in a living animal, we observe none of that contraction and dilatation which is perceptible in the heart; although a manifest pulsation is communicated to the finger placed over it. The phenomena of the pulse will engage attention speedily. We may merely remark, at present, that the pulsations are manifestly more dependent upon the action of the heart than upon that of the arteries. In syncope, they entirely cease; and whilst they continue beneath an aneurismal tumour, because the continuity of the vessel is not destroyed, they completely cease beneath a ligature, so applied around an artery as to cut off the flow of blood. Bichat attached an inert tube to the carotid artery of a living animal, so that the blood could flow through it: the same kind of pulsation was observed in it as in the artery. To this he adapted a bag of gummed taffeta, so as to simulate an aneurismal tumour: the pulsations were evidenced in the bag. If, again, arterial blood be passed into a vein, the

De Arteriar, et Venar, vi Irritabili, &c., Groning, 1766.

b Elemens de Médec. Operat., Turin, 1806.


- Lectures on Inflammation, p. 83, Edinh. 1813; also, 2d Amer. Edit., Philad. 1831.

  d On the Arterial Pulse, p. 52, Bath, 1816.
- · On Inflammation of the Mucous Membrane of the Lungs, p. 20, Lond. 1820.
- <sup>f</sup> See Henle, Allgemein. Anatom. u. s. w. S. 513, Leipz. 1841. <sup>g</sup> View of the Progress of Animal Chemistry, p. 25, Lond. 1813. <sup>h</sup> An Introduction to Medical Literature, p. 501, Lond. 1813.

latter vessel, which has ordinarily no pulsation, now begins to beat; whilst, if blood from a vein be directed into an artery, the latter ceases to beat.<sup>a</sup>

Another class of physiologists have reduced the whole of the arterial action to simple elasticity; a property, which the yellow tissue that composes the proper membrane of the artery, seems to possess in an unusual degree. Such is the opinion of Magendie.b "Admitting it to be certain," he remarks, "that contraction and dilatation occur in the arteries, I am far from thinking, with some authors of the last century, that they dilate of themselves, and contract in the manner of muscular fibres. On the contrary, I am certain, that they are passive in both cases, that is, that their dilatation, and contraction are the simple effect of the elasticity of their parietes, put in action by the blood, which the heart sends incessantly into their cavity," -and he farther remarks, that there is no difference, in this respect, between the large and the small arteries. As regards the larger arteries, it is probable, that this elasticity is the principal but not the only action exerted; and that it is the cause, why the blood flows in a continuous, though pulsatory, stream, when an opening is made into them; thus acting like the reservoir of air in certain pumps. In the pump A B, represented

in the marginal figure, were there no air-vessel C, the water would flow through the pipe E at each stroke of the piston, but the stream would be interrupted. By means of the air-vessel, this is remedied. The water, at each stroke, is sent into the vessel; the air contained in the air-vessel is thus compressed, and its elasticity thereby augmented; so that it keeps up a constant pressure on the surface of the water, and forces it out of the vessel, through the pipe D, in a nearly uniform stream. Now, in the heart, the contraction of the ventricle acts like the depression of the piston; the blood is propelled into the artery in an interrupted manner, but the elasticity of the bloodvessel presses upon the blood,



in the same manner as the air in the air-vessel presses upon the water within it; and thus the blood flows along the vessel in an uninterrupted, although pulsatory, stream.

There are many difficulties, however, in the way of admitting the whole of the action of the arteries in the circulation to be de-

<sup>2</sup> Adelon, Physiol. de l'Homme, edit. cit. iii. 380; and art. Circulation, in Dict. de Médecine, lère édit. v. 321, Paris, 1822.

<sup>5</sup> Précis, &c. edit. cit. ii. 387.

VOL. II. - 12

pendent upon simple elasticity. The heart of a salamander was opened by Spallanzani, a yet the blood continued to flow through the vessels for twelve minutes after the operation. The heart of a tadpole was cut out, yet the circulation was maintained for some time in several of the vascular ramifications of the tail. The heart of the chick in ovo was destroyed immediately after contraction; the arterial blood took a retrograde direction, and the momentum of the venous blood was redoubled. The circulation continued in this manner for eighteen minutes. Dr. Wilson Philipb states, that he distinctly saw the circulation in the smaller vessels, for some time after the heart had been removed from the body, and a similar observation was made by Dr. Hastings.c The latter gentleman states, that in the large arterial trunks, and even in the veins, he has noticed, in the clearest manner, their contraction on the application of varions stimulants, both chemical and mechanical. It is, moreover, well known, that if a small living artery be cut across, it will soon contract, so as to arrest hemorrhage; and that, whilst an animal is bleeding to death, the arteries will accommodate themselves to the decreasing quantity of blood in the vessels, and contract beyond the degree to which their elasticity could be presumed to carry them; and that after death they will again relax. Dr. Parry found, that the artery of a living animal, if exposed to the air, will sometimes contract in a few minutes to a great extent; in such case, only a single fibre of the artery may be affected, narrowing the channel in the same way as if a thread were tied round it.

The experiments, which have been instituted for the purpose of discovering the dependence of the arterial action on the nervous system, have likewise afforded evidences of their capability of assuming a contractile action, and have led to a better comprehension of those cases of what have been called local determinations of blood. Dr. Philip found, that the motion of the blood in the capillaries is influenced by stimulants applied to the central parts of the nervous system, which must be owing to the capillaries possessing a power of contractility, capable of being aroused to action by the nervous influence. The experiments of Sir Everard Homed are, however, more applicable, as they were directed to the larger arteries, respecting which the greatest doubts have been entertained. The carotid artery of a dog was laid bare; the par vagum and great sympathetic, which, in that animal, form one bundle, were separated from it by a flattened probe, for one-tenth of an inch in length; the head and neck of the dog were then placed in an easy position, and the pulsations of the carotid artery were attended to by all present for two minutes, in order that the eye might be accustomed to their force in a natural state. The nerve, passing

d Lectures on Comparative Anatomy, iii. 57, Lond. 1823.

Experiments on the Circulation, &c., translated by R. Hall, Lond. 1801.

b An Experimental Inquiry into the Laws of the Vital Functions, Loud. 1817; and

Lond. Med. Gazette, for March 25th, 1837, p. 952. Cop. citat. p. 51.

over the probe, was then slightly touched with caustic potassa. In a minute and a half, the pulsations of the exposed artery became more distinct. In two minutes, the beats were stronger; in four minutes, their violence was lessened; and in five minutes, the action was restored to its natural state. The experiment was repeated, with analogous results, upon a rabbit. In this animal, the par vagum was separated from the intercostal nerve; and it was found, that, when the former nerve alone was irritated, no increase took place in the force of the action of the artery. "The carotid artery," says Sir Everard, "was chosen as the only artery in the body of sufficient size, that can be readily exposed, to which the nervous branches, supplying it, can be traced from their trunk. This experiment was repeated three different times, so as to leave no doubts respecting the result." These experiments demonstrate, that, under the nervous influence, an increase or diminution may take place in the contraction of an artery; and they aid us in the explanation of those cases, in which the circulation has been accomplished, where the heart has been altogether wanting or completely defective in structure.

Sir Everard instituted some farther experiments, with the view of determining whether heat or cold have the greatest agency in stimulating the nerves to produce this effect upon the artery. The wrist of one arm was surrounded by bladders filled with ice; and after it had remained in that state for five minutes, the pulse of the two wrists was felt at the same time. The beats in that which had been cooled were found to be manifestly stronger. A similar experiment was now made with water, heated to 120° or 130° of Fahrenheit. The pulse was found to be softer and feebler in the heated arm. When one wrist was cooled and the other heated, the stroke of the pulse, in the cooled arm, had much greater force than that of the heated one. These experiments were repeated upon the wrist of several young men and young women of different

ages, with uniform results.

Lastly, we have remarked, and shall have occasion to refer to the matter again, that certain animals, which have no heart, have circulating vessels in which contraction and dilatation are perceptible. This is the case with the class vermes of Cuvier, and can be seen very distinctly in the lumbricus marinus or lug, the leech, &c. The fact has been invoked both by the believers in the muscular contractility of arteries, and by those who conceive the contractility to be peculiar; but our acquaintance with the intimate structure of the coats of the vessels, in those animals, is too minute for us to assert more than that they are manifestly contractile. In an interesting case of acardiac fætus examined by Dr. Houston, of Dublin, it seemed impossible that the heart of a twin fætus could have occasioned the movement of blood in the acardiac one; and

a See a case of this kind in the human fætus, by Dr. J. S. B. Jackson, of Boston, in the American Journal of the Medical Sciences, for February, 1838, p. 362; and another by Dr. Houston, in the Dublin Journal of Med. Sciences, No. xxix. See, also, Prof. Graves, Lond. Med. Gaz, June 30, 1838, p. 562.

hence that there must have been some power in the vessels of the

latter to effect the circulation through it.

From these and other considerations, the majority of physiologists have admitted a contractile action, not simply in the capillary vessels, but in the larger arterial trunks; and, at the present day, the most general and satisfactory opinion appears to be, that, in addition to the highly elastic property possessed by the middle coat, it is capable of being thrown into contraction; that, in the larger vessels, this contraction is but little exerted, the action of the artery being mainly produced by its elasticity; but that, in the smaller arterial ramifications, the contractility is more apparent; and, in the capillary vessels, is scarcely equivocal. To this action of contractility, necessarily connected with the life of the vessel, and differing from both muscular contractility and simple elasticity, Dr. Parrya gave the name tonicity.

## c. Circulation through the Capillaries.

The agency of the capillary vessels in the circulation has been a subject of contention. It was the opinion of Harvey, and it is embraced by J. Müller, b that the action of the heart is alone sufficient to send the blood through the whole circuit; but we have seen, that, even when aided by the elasticity and contractility of the arterial trunks, the pulsations of the heart become imperceptible in the smaller arteries; and, hence, that there is some show of reason for the belief, that, in the capillary vessels, the force may be entirely spent. Were we, indeed, to admit that the force of the heart were sufficient to send the blood through a single capillary circulation, it would be difficult to admit that it could send it through two - as in the portal circulation. Bichat regarded the capillaries as organs of propulsion, and alone concerned in returning the blood to the heart through the veins. Dr. Marshall Hall, on the other hand, denies that we have any proof of irritability in the true capillaries; and Magendied conceives the contraction of the heart to be the principal cause of the passage of the blood through these vessels. In support of this view he adduces the following experiment. Having passed a ligature round the thigh of a dog, so as not to compress the crural artery or vein, he tied the vein near the groin, and made a small opening into the vessel. The blood immediately issued with a considerable jet. He then pressed the artery between the fingers, so as to prevent the arterial blood from passing to the limb. The jet of venous blood did not, however, stop. It continued for some moments, but went on diminishing, and the flow was arrested, although the vein was filled throughout its whole extent. When the artery was examined during these events, it was observed to contract gradually, and at length became completely empty when the course of the blood in

b Handbuch, u. s. w., Baly's translation, p. 220, Lond. 1838.

<sup>2</sup> On the Arterial Pulse, p. 52, Bath, 1816.

c A Critical and Experimental Essay on the Circulation, &c. p. 78, Lond. 1831. Reprinted in this country, Philad. 1835.

the vein ceased. At this stage of the experiment, the compression was removed from the artery; the blood immediately passed into the artery, and, as soon as it had reached the final divisions, it began to flow again through the opening in the vein, and the jet was gradually restored. On compressing the artery again, until it was emptied, and afterwards allowing the arterial blood to pass slowly along the vessel, the discharge from the vein occurred, but without any jet; which was resumed, however, as soon as the

artery was entirely free.

This experiment is not so convincing to us as it appears to be to M. Magendie. The chief fact, which it exhibits, is the elastic, and probably contractile, power of the arteries. It might have been expected, à priori, under any hypothesis, that the quantity of blood discharged from the vein would hold a ratio with that sent by the artery; and, consequently, the experiment appears to us to bear but little on the question regarding the separate contractile action of the capillaries. It is difficult, indeed, to believe, that such an action does not exist. In addition to the circumstance, already mentioned, of the absence of pulsation in the smaller arteries, almost every writer on the theory of inflammation considers the fact of a distinct action of the capillaries established, and leaves to the physiologist the by no means easy task of proving it. Dr. Wilson Philip placed the web of a frog's foot in the microscope, and distinctly saw the capillaries contract upon the application of those stimulants, which produce contraction of the muscular fibre. The result of Dr. Thomson'sb experiments, in investigating the subject of inflammation, were the same, as well as those of Dr. Hastings.c The facts, which we have already referred to, regarding the continuance of the circulation in the minute vessels after the heart has been removed, are confirmatory of the same point; as well as the observation of Dr. Philip, that the blood in the capillaries is influenced by stimulants applied to the central parts of the nervous system. The experiments of Thomson, Philip, and Hastings, were repeated by Wedemeyer, with great care. The circulation in the mesentery of the frog, and in the web of its foot, being observed through the microscope, it was evident, that no change occurred in the diameter of the small arteries, or in that of the capillaries, so long as the circulation was allowed to go on in its natural state; but as soon as excitants were applied to them, an alteration of their calibre was perceptible. Alcohol arrested the flow of blood without inducing much apparent contraction of the vessels. Chloride of sodium, in the course of three or four minutes, caused the vessels to contract one-fifth of their calibre, which contraction was followed by dilatation of the vessels, and a gradual retardation and stoppage of the blood. In a space of time varying from ten

A Treatise on Febrile Diseases, 3d edit. ii. 17, Lond. 1813; and Medico-Chirur. Transact. vol. xii. p. 401.

Lectures on Inflammation, p. 83, Edinb. 1813.
 Untersuch. über die Krieslauf, u. s. w., Hannover, 1828; and Edinb. Med. and Surg. Journ. vol. xxxii.

to thirty seconds, and sometimes immediately after the application of the galvanic circle, the vessels contracted, some one-fourth, others one-half, and others three-fourths of their calibre. The contraction sometimes continued for a considerable time, occasionally several hours; in other instances, it ceased in ten minutes, and the vessels resumed their natural diameter. A second application of galvanism to the same capillaries seldom caused any material contraction.a Schwannb likewise found, that when cold water was poured on the vessels of a frog, which had been previously in a warm atmosphere, the capillaries immediately contracted, but after a time regained their diameter. Farther, Mr. Hunter found that on exposing arteries to the air, they contracted so much as to occasion obliteration of their tubes; and it is well known, that when arteries - as the temporal - are divided, hemorrhage may be arrested by the spontaneous contraction of the divided extremity, a contraction, which, as remarked by Dr. Carpenter, is much greater than could be accounted for by simple elasticity of tissue, and is more marked in small than in large vessels.c

All these facts prove the existence of a vital power in the capillaries, capable of modifying, to considerable extent, the flow of

blood through them.d

Again, of this independent action of the capillary vessels we have, every day, proofs in local inflammation; in which there is increased redness of a part, without the general circulation exhibiting the slightest evidences of augmented action or excitement. In the natural state, the vessels of the tunica conjunctiva covering the white of the eye, receive little blood; but if any cause of irritation exists, as a grain of sand entering between the evelids, we find blood rapidly sent into them, giving the appearance, which has been termed, not inappropriately, "bloodshot."e In the experiments of Kaltenbrunner, which were fully confirmed on repetition, the blood was at first observed, in inflammation, streaming to the irritated part, in consequence of which the capillary vessels became distended; afterwards irregularity of circulation occurred in the gorged capillary system; and subsequently complete arrestation of the circulation, and disorganization. These phenomena are of themselves sufficient to prove the existence of the separate action of the capillaries, and when taken in conjunction with other facts, are overwhelming. The blush of modesty, and the paleness of guilt, the hectic glow, and the trans-

c Human Physiology, § 502, Lond. 1842.

· Thomson's Lectures on Inflammation, Edinb. 1813.

<sup>\*</sup> Richerand's Elémens de Physiologie, 13éme édit., par Bérard ainé, Edit. Belge, p. 125, Bruxelles, 1837.

b Müller's Archives, 1836, and Lond. Med. Gazette, May, 1837; and Müller's Handbuch, u. s. w., Baly's translation, p. 206.

<sup>&</sup>lt;sup>4</sup> Purkinje, in Encyclop. Wörterb. der Medicin. Wissenschaft. vii. 660, Berlin, 1831; Prof. Graves, Lond. Med. Gazette, June 30, 1838, p. 561; Gerdy, Dict. de Méd. viii. p. 60; Dubois d'Amiens, L'Experience, No. clxxi. 1840; and Dr. Calvert Holland, Edinb. Med. and Surg. Journ., July, 1842.

<sup>&</sup>lt;sup>1</sup> Experimenta circa Statum Sanguinis et Vasorum in Inflammatione, p. 23, Monach. 1826.

lucency of congelation, are all circumstances, that go to establish

the same point.

The contractile power of the capillaries is doubtless modified by the condition of the ganglionic nerves distributed to them, which, as we have seen, are observed to increase as the size of the vessels and the thickness of their coats diminish. Their influence is, indeed, strikingly evinced in actions, which are altogether nervous, as in the flushed countenance occasioned by sudden mental emotions. By some, however, the whole capillary circulation has been ascribed to the motive faculty inherent in the globules of the blood; whilst others, again, have asserted, that the "electrogalvanic power," - or in other words - the nervous power, generated in the nervous system, and acting on the blood globules through the parietes of the capillary system of vessels is the immediate agent directing the movements or circulation in the capillaries: all this, however, enters into the inscrutable question of what is the cause of life in the fluids or tissues, — a question to be agitated, but not solved, in a subsequent part of this volume.

But, not only has a vital power of contraction been conceded to the capillaries; it has been imagined, that they possess what the Germans call a Lebensturgor (turgor vitalis) or vital property of expansibility or turgescence. Such appears to be the opinion of Hebenstreitb and of Prus; and it has been embraced, in this country, by Professor Smith of Yale College, by his son, Professor N. R. Smith of Baltimore, in his excellent work on the "Arteries," and by Professor Hodge of Philadelphia. The idea has been esteemed to be confirmed by the fact of excitants having been seen under the microscope, by Hastings, Wedemeyer, and others, to occasion not only contraction but dilatation of the capillaries. The phenomena, observed in the erectile tissues, have likewise been considered to favour the hypothesis; but, in answer to these arguments, it may be replied, that the irregular excitation, produced in the parts by the application of powerful stimulants, might readily give occasion to an appearance of expansibility under the microscope, without our being justified in inferring, that these vessels possess an innate vital property of expansibility; and, in many of the cases, in which ammonia and galvanism were applied by Thomson, Hastings, Wedemeyer and others, the action of contraction ought rather to be esteemed physical or chemical, than vital.f The results of the application of such excitants, as diluted alcohol, dilute solutions of ammonia and of chloride of sodium, can alone be adduced as evidences of vital action

e De l'Irritation, &c., Paris, 1825.

d Surgical Anatomy of the Arteries, 2d edit., Baltimore, 1835.

f Burdach's Physiologie als Erfahrungswissenschaft, B. iv. Leipz 1832.


Alison's Outlines of Physiology, Supplement to 2d edit., Edinb. 1836; and A Short Inquiry into the Capillary Circulation, &c. by James Black, M.D., Lond. 1825.

b Dissert. de Turgore Vitali, Lips. 1795; Hildebrandt's Physiologie, Auflag. 5, § 84; and Tiedemann's Physiologie, trad. par Jourdan, p. 625, Paris, 1831.

e North Amer. Med. and Surg. Journal, June, 1828. See, also, Prof. Graves, in Lond. Med. Gazette, for June 30, and July 7, 1838.

on the part of these vessels. The dilatation of the capillary system and of the smaller arteries, which has been remarked on the contact of these agents, is not, as Oesterreicher, has remarked, the primary effect: it is the consequence of the afflux of blood to the irritated part, as is also demonstrated in the experiments of Kaltenbrunner on inflammation, to which allusion has been made. Lastly, attentive observation of the phenomena presented by the erectile tissues must lead to the conclusion that the turgescence of the vessels is not the first link in the chain of phenomena; excitation is first induced in the nerves of the part - generally through the influence of the brain, and from thence, perhaps, through the sympathetic nerve, - and the afflux of fluid supervenes on this.b The vital expansibility of the capillaries cannot then, we think, be regarded as proved, or probable.

The circulation through the capillaries has long been an interesting topic of microscopic research. According to Wagner, a magnifying power of from two to three hundred diameters is required to make out the particular details. According to him, the blood in mass, or in the larger channels, is seen to flow more rapidly than in the smaller. Here the blood corpuscles advance with great rapidity, especially in the arteries, and with a whirling motion, and form a closely crowded stream in the middle of the vessel, without ever touching its parietes. With a little attention, a narrower and clearer, but always very distinct, space is seen to



seen. - (Wagner.)

remain between the great middle current of blood corpuscles and the walls of the vessel, in which a few white corpuscles, or what Wagner considers to be lymph corpuscles (?) (see vol. i., p. 625) are moved onwards, but at a much slower rate. These white corpuscles swim in smaller numbers in the transparent liquor sanguinis, and glide slowly, and in general smoothly, though they sometimes advance by fits and starts more rapidly, but with intervening pauses; and, as a general rule, at least ten or twelve times more slowly than the corpuscles of the A Small Venous Branch from the Web of a Frog's central stream. The clear space, Foot, magnified 350 diameters. b, b. Cells of pavement epithelium, contain. filled with liquor sanguinis and ing nuclei. In the space between the current of white corpuscles, is obvious in all oval blood-corpuscles, and the walls of the vessel, the round transparent lymph globules (?) are the larger capillaries, whether arterial or venous, but it ceases to


\* Versuch einer Darstellung der Lehre vom Kreislaufe des Blutes, Nürnberg, 1826. <sup>b</sup> Rullier, art. Expansibilité in Dict. de Mêd. viii. 423, Paris, 1823; and Purkinje, in art. Circulatio Sanguinis, in Encyclopad. Wörterb. der Medicin. Wissenschaft. b. vii. s. 667, Berl. 1831.

Elements of Physiology, translated by R. Willis, § 122, Lond. 1842.

be apparent on the smaller intermediate vessels which admit but one or two rows of blood corpuscles (Fig. 160). In these vessels, the two sets of globules proceed pari passû; but, according to Wagner, it is easy to see, that the blood corpuscles glide more readily on-

wards - the white corpuscles seeming often to be detained at the bendings of vessels, and at the angles, where anastomosing branches are given off; where they remain adherent for an instant, and then suddenly proceed onwards. These phenomena are observed in every part of the peripheral systemic circulation; but an exception appears to exist in the pulmonic circulation; the capillaries there being filled with both kinds of globules to their very walls.

It is in this — the intermediate — part of the sanguiferous system, that most important functions take place. In the smallest ar-



Large Vein of Frog's Foot, magnified 600 diameters.
b, c. Blood corpuscles. a, a. Lymph corpuscles (?)
principally conspicuous on the clear space near the
parietes of the vessel.—(Wagner.)

tery, we find arterial blood; and in the smallest vein, communicating with it, blood always possessing the venous properties. Between those points, a change must have occurred, precisely the reverse of that which happens in the lungs. It is in this very part, too, that nutrition, secretion, and calorification are effected. In the explanation of these functions, we shall find it impossible not to suppose a distinct and elective agency in the tissues concerned; and as it is by such agency, that the varying activity of the different functions is regulated, we are constrained to believe, that the capillary vessels may be able to exert a controlling influence over the quantity and velocity of the blood circulating in them.


In disease, the agency of this system of vessels is an object of attentive study with the pathologist. To its influence in inflammation, we have already alluded; but it is no less exemplified in the more general diseases of the frame, — as in the cold, hot, and sweating stages of an intermittent. Local, irregular capillary action is, indeed, one of the most common causes of acute affections, and these generally occur in some organ, at a distance from the seat of the deranging influence. It is a common and just observation, that getting the feet wet, and sitting in a draught of air, are more certain causes of catarrh than sudden atmospheric vicissitudes, which apply to the whole body; and so

extensive is the sympathy between the various portions of this system of vessels, that the most diversified effects will be produced in different individuals exposed to the same common cause; one may have inflammatory sore throat; another, ordinary catarrh; another, inflammation of the bowels; — according to the precise predisposition, existing in the individual at the time, to have one structure morbidly affected rather than another; — but these are interesting topics, which belong more strictly to the pathologist.

By the united action, then, of the heart, the arteries, and the capillary or intermediate system of vessels, the blood attains the veins. We have now to consider the circulation in those vessels.

### d. Circulation in the Veins.

It has been already observed, that Harvey considered the force of the heart to be of itself sufficient to return the blood, sent from the left ventricle, to the heart; whilst Bichat conceived the whole



propulsory effort to be lost in the capillaries, and the transmission of the blood along the veins to be entirely effected by the agency of the capillary system. It is singular, that an individual of such distinguished powers of discrimination should have been led into an error of such magnitude. It is a well-known principle in hydrostatics, that although water, when uncon-

fined, can never rise above its level at any point, and can never move upwards; yet, by being confined in pipes or close channels of any kind, it will rise to the height from which it came. Hence the water or blood in the vessel A, Fig. 172, which may be considered to represent the right auricle, would stand at the same height as that in the vessel B, which we may look upon as the left ventricle, were they inanimate tubes. We need be at no loss, therefore, in understanding how the blood might attain the right auricle, when the body is erect, by this hydrostatic principle alone; but we have seen, that the force exerted by the heart, by the arteties, and by the capillary system is superadded to this, so that the blood would rise much higher than the right auricle, and consequently exert a manifest effort to enter it. It may be remarked, also, that the left ventricle is not the true height of the source, but the top of the arch of the aorta, which is more elevated, by several inches, than the right auricle. A similar view is embraced by Dr. Billing, but Dr. Carpenter, - in commenting on the author's

b Human Physiology, § 516, Lond. 1842.

First Principles of Medicine, Amer. Edit., p. 36, Philad. 1842.

observations on this subject—suggests, that the influence of this hydrostatic force would scarcely be felt through the plexus of capillary vessels; "for the interposition of a system of tubes even of much larger calibre would be, by the friction created between the fluid and their walls, an effectual obstacle to the rapid ascent of a current, which had so slight an impetus as that derived from its previous fall." The author did not mean, however, to say more than that the blood "might attain" the right auricle by the hydrostatic force alone; he did not wish to convey the idea, that the circulation could be carried on without the aid of an additional force; but that a slight effort only on the part of the heart and arteries might be needed to enable the blood to perform its entire circuit.

Are we then to regard the veins as simple elastic tubes? This is the prevalent belief. Their elasticity is, however, much less than that of the arteries. Some physiologists have conceived them to possess also contractile properties. Such is the opinion of Broussais, who founds it, in part, upon certain experiments by Sarlandière, already referred to, in which contraction and relaxation of the venæ cavæ of a frog were seen, for many minutes after the heart was removed from the body. These pulsations of the venæ cavæ and also of the pulmonary veins in their natural state have been seen by numerous observers - by Steno, Lower, Wepfer, Borrachius, Whytt, Haller, Lancisi, Müller, Marshall Hall, Flourens, J. J. Allison, and others. The experiments of Dr. Allison, in reference to the venæ cavæ and pulmonary veins, appeared to him, to prove, - that they pulsate near the heart in the four classes of the vertebrata, — that in dying animals they pulsate long after the auricle and ventricle have ceased to beat, - that they will also beat even in quadrupeds, for hours after they have been separated from the heart and from the body; - and that they can be stimulated to contract either when in or out of the body, by mechanical and galvanic means, especially by the latter, after all motion has ceased for some time.

It seems to be very doubtful, whether the veins generally possess any sensible contraction like that of the venæ cavæ and pulmonary veins near the heart, for although irritated by galvanic and mechanical stimuli by Haller, Nysten, Müller, J. J. Allison, and others, no motion whatever could be detected in them. Gerber affirms, that the fibres of the middle coat of the veins bear a stronger resemblance to those of muscular tissue than do those of the corresponding part of the arteries, which resemble more the ordinary elastic fibres; but Dr. Carpenterd thinks it not improbable, that his observations were made on portions of the veins near the heart, which partake of its contractility.

<sup>\*</sup> Traité de Physiol. &c., Drs. Bell and La Roche's translat. p. 391, Philad. 1832. See also, Müller's Handbuch, u. s. w., Baly's translation, p. 171, Lond. 1838.

b See the experiments of the last named gentleman proving the existence of a venous pulse independent of the Heart and Nervous System, in Amer. Journal of the Medical Sciences, Feb. 1839, p. 306.

c J. Müller, Elements of Physiology, p. 170, Lond. 1838, and J. J. Allison, op. cit. p. 320. d Human Physiology, § 514, Note. Lond. 1842.

In the experiments of Dr. Marshall Hall, on the circulation in the web of the frog's foot, he was almost invariably able to detect, with a good microscope, a degree of pulsatory acceleration of the blood in the arteries at each contraction of the heart; and he is disposed to conclude, from his observations, that the natural circulation is rapid, and entirely pulsatory in the minute arteries, and slow and equable in the capillary and venous systems. But whenever the circulation was in the slightest degree impeded, the pulsatory movement became very manifest at each systole of the heart, and it was seen in all the three systems of vessels - arterial, capillary, and venous. He observed, that in the arteries there was generally an alternate, more or less rapid flow of the globules at each systole and diastole of the ventricle; and that in the capillaries and veins the blood was often completely arrested during the diastole, and again propelled by a pulsatory movement during the systole; all which he esteems conclusive proof, that the power and influence of the heart extend through the arteries to the capillaries, and through these to the veins, even in the extreme parts of the

That the veins are possessed of elasticity is proved by the operation of bloodletting, in which a part of the jet, on puncturing the vein, is owing to the over-distended vessel returning upon itself; but that this property exists to a trifling extent only is shown by the varicose state of the vessels, which is so frequently

seen in the lower extremities.

## e. Forces that propel the Blood.

From the inquiry into the agency of the different circulatory organs in propelling the blood, it is manifest, that the action of the heart, the elasticity of the arteries, and a certain degree of contractile action, in the smaller vessels more especially, a distinct action of the capillary vessels, and a slight elastic and perhaps contractile action on the part of the veins, must be esteemed the chief agents. Of these, the action of the heart and capillaries, and the contraction of the arteries and veins, can alone be regarded as sources of motion, the elasticity of the vessels being simple directors, not generators of force. But there is another agency, which is probably more efficient than has been generally conceived. This is the suction power of the heart, or derivation, as it has been termed, to which attention has been chiefly directed by Haller, Wilson, Carson, Zugenbühler, Schubarth, Platner, Blumenbach, and others; but which is not assented to by Oesterreicher, Müller, and

\* Essay on the Circulation, ch. i. Lond. 1831, and Philad. 1835.

b Elem. Physiol. ii. lib. vi.

c Enquiry into the Moving Powers employed in the Circulation of the Blood, Lond. 1784.

4 Inquiry into the Causes of the Motion of the Blood, 2d edit. Lond. 1833.

 Tiedemann's Traité de Physiol. par Jourdan, p. 347; and Burdach's Physiologie als Erfahrungswissenschaft, iv. 270, Leipz. 1832.

Lehre vom Kreislauf des Blutes, Nürnberg, 1826, Handbuch, u. s. w., Baly's translation, p. 173,

some others.a It is presumed, that the muscular fibres of the heart are mixed up with a large quantity of cellular tissue; and that, whilst the contraction of the cavities is effected by the action of the muscular fibres, dilatation is produced by the relaxation of the contracted fibres, and the elasticity of the cellular tissue; so that when the heart has contracted, and sent its blood onwards, its elasticity instantly restores it to its dilated condition; a vacuum is formed, and the blood rushes in to fill it. This action has been compared by Dr. Bostock, and by Dr. Southwood Smith, Prof. Turner, and others, to that of an elastic gum bottle, which, when filled with water, and compressed by the hand, allows the fluid to be driven from its mouth with a velocity proportionate to the compressing force. But the instant the pressure is removed, elasticity begins to operate, and if the mouth of the bottle be now immersed in water, a considerable quantity of that fluid will be drawn up into the bottle, in consequence of the vacuum formed within it. The existence of this force is confirmed by Döllinger, - who, when examining the embryo of birds, saw the blood advance along the veins, whilst the venous trunks poured it into the auricles at the moment when they dilated to receive it; as well as by Dr. T. Robinson, who was forcibly struck with the activity with which the diastole was effected, in the case of monstrosity more than once referred to. Dr. Carpenters thinks it very doubtful "how far the auricles have such a power of active dilatation as would be required for this purpose;" but the question need not regard the auricles. It is but necessary to suppose, that an action of power of dilatation exists in the ventricles; and this is now generally admitted. He farther remarks, that it has been shown experimentally by Dr. Arnott and others, that no suction power exerted at the farther end of a long tube, whose walls are as deficient in firmness, as those of the veins are, can occasion any acceleration in a current of fluid transmitted through it; for the effect of the suction is destroyed at no great distance from the point at which it is applied by the flapping together of the sides of the vessel; but in answer to this it may be observed, that it remains to be shown, that such flapping of the sides would necessarily occur in the veins, which are living vessels, and constantly receiving blood from the capillaries under the action of vital forces.

Another accessory force, which has been invoked, is the suction power of the chest, or the *inspiration of venous blood*, as it has been termed. This is conceived to be effected by the same mechanism as that which draws air into the chest. The chest is di-

\* Carpenter, Human Physiology, § 575, Lond. 1842.

b Physiology, 3d edit. p. 251, Lond. 1836.

Animal Physiology, (Library of Useful Knowledge,) p. 83, Lond. 1829.

d Edinb. Medico-Chirurg, Transact iii. 225.

Denkschriften der Königl. Akademie der Wissenschaft. zu München, vii. 217;
 and Burdach, op. citat, p. 272.

American Journal of the Medical Sciences, No. xxii.

F Human Physiology, § 515, Lond. 1842.

VOL. II. - 13

lated during inspiration; an approach to a vacuum occurs in the thorax; and the blood, as well as the air, is forcibly drawn towards that cavity. On the other hand, during expiration, all the thoracic viscera are compressed; the venous blood is repelled from the chest, and the arterial blood reaches its destination with greater celerity, owing to the action of the expiratory muscles being added to that of the left ventricle. Haller,a Lamure, and Lorryc had observed, that the blood, in the external jugular vein, moves under manifestly different influences, during inspiration and expiration. Generally, when the chest is dilated in inspiration, the vein empties itself briskly, becomes flat, and its sides are, occasionally, accurately applied against each other; - but, during expiration, the vein rises and becomes filled with blood; - effects, which are more evident, when the respiratory movements are more extensive. The explanation of this phenomenon, by Haller and Lorry, is the one given above.

To discover whether the same thing happens to the venæ cavæ, Magendie introduced a gum elastic catheter into the jugular vein. so as to penetrate the vena cava and even the right auricle; - the blood was observed to flow from the extremity of the tube at the time of expiration only. During inspiration, air was rapidly drawn into the heart, giving rise to the symptoms to be mentioned hereafter, which attend the reception of air into that organ. Similar results were obtained, when the tube was introduced into the crural vein in the direction of the abdomen. So far as regards the larger venous trunks, therefore, the influence of respiration on the circu-

lation is sufficiently evidenced.d

It can be easily shown, by opening an artery of the limbs, that expiration manifestly accelerates the motion of arterial blood; especially in forced expiration, and during violent exertion. In animals, subjected to experiment, it is impracticable to excite either the forced expiration or the violent effort at pleasure; but we can, as a substitute, compress the sides of the chest with the hands, according to the plan recommended by Lamure, when the blood will be found to flow more or less copiously in proportion to the pressure exerted. It occurred to Magendie, that this effect of respiration on the course of the blood in the arteries might influence the flow along the veins. To prove this, he passed a ligature around one of the jugular veins of a dog. The vessel emptied itself beneath the ligature, and became turgid above it. He then made a slight puncture, with a lancet, in the distended portion; and in this way obtained a jet of blood, which was not sensibly modified by the ordinary respiratory movements, but became of triple or quadruple the size, when the animal struggled. As it might be objected to this experiment, that the effect of respiration was not transmitted by the arteries to the open vein, but rather by the veins that had remained free, which might have conveyed the blood, repelled

\* Elementa Physiologiæ, tom. ii.

4 See, also, Poiseuille, in Magendie's Journal de Physiologie, viii. 272.

Mém. de l'Acad. des Sciences, pour 1749.
Magendie, Précis, &c. ii. 416.

from the vena cava, towards the tied vein, by means of anastomoses, the experiment was varied. The dog has not, like man, large internal jugular veins, which receive the blood from the interior of the head. The circulation from the head and neck is, in it, almost wholly confined to the external jugular veins, which are extremely large; the internal jugulars being little more than vestiges. By tying both of these veins at once, Magendie made sure of obviating, in great part, the reflux in question; but, instead of this double ligature diminishing the phenomenon under consideration, the jet became more closely connected with the respiratory movement; for it was manifestly modified even by ordinary respiration, which was not the case when a single ligature was employed.

From these and other experiments, Magendie properly concludes, that the turgescence of the veins must not be ascribed, with Haller, Lamure, and Lorry, simply to the reflux of the blood of the venæ cavæ into the branches opening directly or indirectly into them; but that it is partly owing to the blood being sent in larger quan-

tity into the veins from the arteries.a

In the same manner are explained — the rising and sinking of the brain, which, as was observed in an early part of this work, (vol. i., p. 79,) are synchronous with expiration and inspiration. During expiration, the thoracic and abdominal viscera are compressed; the blood is driven more into the branches of the ascending aorta, and it is, at the same time, prevented from returning by the veins: owing to the combination of these causes, the brain is raised during expiration. In inspiration, all this pressure is removed; the blood is free to pass equally by the descending, as by the ascending, aorta; the return by the veins is ready, and the brain therefore sinks. We can thus, also, explain why the face is red and swollen during crying, running, straining, and the violent emotions; and why pain is augmented in local inflammations of an extremity, - as in cases of whitlow, and when respiration is hurried or impeded by running, crying, &c. The blood accumulates in the part, owing to the compound effect of increased flow by the arteries, and impeded return by the veins. The same explanation applies to the production of hemorrhage by any violent exertion; and Bourdon' affirms, that he has always seen hemorrhage from the nose largely augmented during expiration; diminished at the time of inspiration, and arrested by prolonged inspiration; — a therapeutical fact of some interest.

It is manifest, then, that the circulation is modified by the movements of inspiration and expiration, — the former facilitating the

a Précis, &c. ii. 421.

b This motion of the brain must not be confounded with that which is synchronous with the contraction of the left ventricle; and which is owing to the pulsation of the arteries at the base of the brain.

c Recherches sur le Mecanisme de la Respiration et sur la Circulation du Sang, Paris, 1820.

d Dr. Clendinning's Report to the Brit. Association, 1839-40, Lond. Med. Gazette, Nov. 13, 1840, p. 270.

flow of blood to the heart by the veins, and the latter encouraging the flow by the arteries; and we shall see hereafter, that there is great reason for the belief, that the dilatation of the chest, - which constitutes the first inspiration of the new-born child, - is a great cause of the establishment of the new circulation; the same dilatation, which causes the entrance of air into the air-cells, soliciting the flow of blood, or the "inspiration of venous blood," as Magendiea has termed it. In a paper read before the Royal Society of London, in June, 1835, Dr. Wardrop, b after remarking, that he considers inspiration as an auxiliary to the venous, and expiration to the arterial circulation, attempts, on this principle, to explain the influence exerted on the circulation, and on the action of the heart, by various modes of respiration, whether voluntary or involuntary, in different circumstances. Laughter, crying, weeping, sobbing, and sighing, he regards as efforts made with a view to effect certain alterations in the quantity of blood in the lungs and heart, when the circulation has been disturbed by mental emotions. The influence of ordinary respiration can, however, be but trifling; yet it has been brought forward by Sir David Barrye as the efficient cause of venous circulation. His reasons for this belief are, - the facts just mentioned, regarding the influence of inspiration on the flow of blood towards the heart, and certain ingeniously modified experiments, tending to the elucidation of the same result. He introduced one end of a spirally convoluted tube into the jugular vein of an animal, and plunged the other into a vessel filled with a coloured fluid. During inspiration, the fluid passed from the vessel into the vein; during expiration, it remained stationary in the tube, or was repelled into the vessel. Dr. Bostockd remarks, that he was present at some experiments, which were performed by Sir David, at the Veterinary College in London, and it appeared sufficiently obvious, that when one end of a glass tube was inserted either into the large veins, into the cavity of the thorax, or into the pericardium, - the other end being plunged into a vessel of coloured water, - the water was seen to rise up the tube during inspiration, and to descend during expiration. The conclusion of Sir David from these experiments is most comprehensive; - that "the circulation in the great veins depends upon atmospheric pressure in all animals possessing the power of contracting and dilating a cavity around that point, to which the centripetal current of their circulation is directed ;" and he conceives, that as, during inspiration, a vacuum is formed around the heart, the equilibrium of pressure is destroyed, and the atmosphere acts upon the superficial veins, propelling their contents onwards to supply the vacuum.

Independently of other objections, there are a few, which ap-

a Précis, &c. ii. 416.

Experimental Researches on the Influence of Atmospheric Pressure upon the Circulation of the Blood, &c., Lond. 1826.

on the Nature and Treatment of the Diseases of the Heart; with some new views of the Physiology of the Circulation, Lond. 1837.

d Physiology, 3d edit. p. 330, Note, Lond. 1836.

pear to us convincing against this sole agency of ordinary respiration in effecting venous circulation. According to Sir David's hypothesis, blood ought to arrive at the heart at the time of inspiration only; and as there are, in the average, seventy-two contractions of the heart for every eighteen inspirations; or four contractions, or -- what is the same thing - four dilatations of the auricle for each respiration; one of these only ought to be concerned in the propulsion of blood, whilst the rest should be bloodless; yet we feel no difference in the strength of the four pulsations. It is clear, too, if we adopt Sir David's reasoning, that, of the four pulsations, two, and consequently two dilatations of the auricles; must occur during expiration, at which time the capacity of the chest is actually diminished: moreover, holding the breath ought to suspend the circulation; and the respiratory influence cannot be invoked to explain the circulation in the fœtus or in aquatic animals. At the most, therefore, respiration can only be regarded as a feeble auxiliary in the circulation. In favour of Dr. Barry's opinion of the efficiency of atmospheric pressure in causing the return of blood by the veins, he adduces the fact, - already referred to, under the head of Absorption,— that the application of an exhausted vessel over a poisoned wound prevents the absorption of the poison; but this, as we have seen, appears to be a physical effect, which would apply equally to any view of the subject.


In all these cases, the elastic resilience of the lungs, by contributing to diminish the atmospheric pressure from the outer surface of the auricles, may, likewise, as suggested by Dr. Carson, have some agency in soliciting the blood into these cavities, but the

agency cannot be great.

There is another circumstance of a purely physical nature, which may exert some influence upon the flow of the blood along the veins; viz., the expanded termination of the venæ cavæ in the right auricle. To explain this, it is necessary to premise a detail


of a few hydraulic facts. If an aperture A, Fig. 173, exist in a cistern X, the water will not issue at the aperture by a stream of uniform size; but, at a short distance from the reservoir, it will be contracted as at B, constituting what has been termed the vena contracta. Now. it has been found, that if a tube, technically called an adjutage, be attached to this aperture, so as to accurately fit the stream, as at A B, Fig. 174, as much fluid will flow from the reservoir as if the aperture alone existed.

Again, if the pipe B C be attached



<sup>&</sup>lt;sup>2</sup> Philosoph. Transact. for 1820, and An Inquiry into the Causes of Respiration, &c. 3d edit. Liverpool, 1833.

to the adjutage A B, the expanded extremity at A will occasion



the flow of water, from the reservoir, to be greater than it would be, if no such expanded extremity existed, in the ratio, according to Venturi, of 12.1 to 10;

and if to the tube B C, a truncated conical tube C D be attached, the length of which is nearly nine times the diameter of C; and the diameter of C to that of D be as 1 to 8; the flow of water will be augmented in the proportion of 24 to 12.1; so that, by the two adjutages A B and C D, the expenditure through the pipe B C is increased in the ratio of 24 to 10. This fact, - the result of direct experiment, and so important to those who contract to supply water by means of pipes, — was known to the Romans. Private persons, according to Frontinus, were in the habit of purchasing the right of delivering water in their houses from the public reservoirs, but the law prohibited them from making the conducting pipe larger than the opening allowed them in the reservoir, within the distance of fifty feet. The Roman legislature must, therefore, have been aware of the fact, that an adjutage with an expanded orifice, would increase the flow of water; but they were ignorant that the same effect would be induced beyond the fifty feet.

Let us apply this law to the circulation. In the first place, at the origin of the pulmonary artery and aorta, there is a manifest narrowness, formed by the ring at the base of the semilunar valves (see Fig. 156); and this might be conceived unfavourable to the flow of the blood along those vessels during the systole of the ventricles; but from the law, which has been laid down, the narrowness would occupy the natural situation of the vena contracta, and, therefore, little or no effect would be induced. The discharge would be the same as if no such narrowness existed. We have seen, again, that the vena cava becomes of larger calibre as it approaches the right auricle, and finally terminates in that cavity by an expanded aperture. This may have a similar effect with the expanded tube C D, Fig. 174, which doubles the ex-

penditure.b

In making these conjectures, — some of which have been adduced by Sir Charles Bell, — it is proper to observe, that, in the

\* Lat. Oudendorp, Lugd. Bat. 1731.

b Venturi, Sur la Communication Latérale du Mouvement dans les Fluides, Paris, 1798; and Pouillet, Elémens de Physiologie, i. 205, Paris, 1832.

opinion of some natural philosophers, the effect of the adjutage is entirely due to atmospheric pressure, and that no such acceleration occurs, provided the experiment be repeated in vacuo. Sir Charles Bella conceives, that "the weight of the descending column in the reservoir being the force, and this operating as a vis à tergo, it is like the water propelled from the jet d'eau, and the gradual expansion of the tube permits the stream from behind to force itself between the filaments, and disperses them, without producing that pressure on the sides of the tube, which must take place, where it is of uniform calibre." It is on this latter view only, that these singular hydrostatic facts can be applied to the doctrine of the circulation.

In addition to the movements, impressed on the blood by the parietes of the cavities in which it moves, it has been considered by many physiologists, - as by Harvey, Glisson, Bohn, Albinus, Rosa, Tiedemann, G. R. Treviranus, Bogerson, Alison, and others, to possess a power of automatic or self-motion. Broussaisf asserts, that he has seen experiments, - originally performed by P. A. Fabre, which showed, that the blood, in the capillary system, frequently moves in an opposite direction to that given it by the heart, - repeated by M. Sarlandière, on the mesentery of the frog. In these, the blood was seen to rush for some moments towards the point irritated, and, when a congestion had taken place there, they remarked, that the globules took a different direction, and traversed vessels which conveyed them in an opposite course, and, a few seconds afterwards, these were again observed to return with equal rapidity to the point from which they had been repelled. Tiedemanng has collected the testimonies of various individuals on this point. Haller, b Spallanzani, Wilson Philip, G. R. Treviranus, k and others, have remarked, by the aid of the microscope, that the blood continued to move in the vessels of different animals, but chiefly of frogs, for some time after the great vessels had been tied, or the heart itself removed; - a fact which Tiedemann, also, often witnessed. C. F. Wolff, Rolando, Döllinger, and Pander, Prévost and Dumas, Von Baer, and others, saw globules of blood in

Animal Mechanics, p. 40, in Library of Useful Knowledge, Lond. 1829.

b Tiedemann, Traité Complet de Physiologie de l'Homme, traduit par Jourdan, i. A Treatise on Inflammation, &c., Lond. 1832. 348, Paris, 1831. d Edinburgh Med. and Surg. Journal for Jan. 1836. Also, Prof. R. J. Graves, in

Lond. Med. Gazette, June 30, 1838, p. 561.

o Dr. S. Smith's Philosophy of Health, p. 398, Lond. 1835; and Messrs. Emmerson

and Reader, in Edinb. Med. and Surg. Journal, April, 1836.

<sup>f</sup> Traité de Physiologie, &c., translated by Drs. Bell and La Roche, 3d edit. p. 374, b Oper. Minor. i. 115, Sect. 8. g Op. citat. Exper. on the Circulation, &c., in Eng. by R. Hall, Lond. 1801.

Philos. Transact. 1815; and Medico-Chirurg. Trans. vol. xii.

k Vermischte Schriften, i. 102.

Theoria Generationis, Hal. 1759.

m Dizionario Periodico di Medicina, Torino, 1822-1823.

Dissert. sist. Hist. Metamorphoseos quam Ovum Incubatum prioribus quinque Diebus subit, Wirceb. 1817.

Annales des Sciences Naturelles, tom. xii. p. 415, Dec. 1827.

P Ueber Entwickelungsgeschichte der Thiere, u. s. w. Th. i. Königsberg, 1828. See,

motion in the incubated egg, before the formation of either vessels or heart; and Hunter, Gruithuisen, and Kaltenbrunner observed - in the midst of the cellular tissue of inflamed parts, in tissues undergoing regeneration, and during the cicatrization of wounds,bloody points placed successively in contact with each other, forming small currents, which represented new vessels, and united to those already existing. The fact, indeed, that the embryo forms its own vessels, and that blood in motion can be detected before vessels are in esse, is a sufficient proof, -- were there no other, - that the globules of the blood possess the faculty of motion, a either in themselves, or by virtue of an attraction exerted upon them by the solid parietes in which they move. The idea of spontaneous motion in a fluid, independently of attraction or repulsion from the sides of another object, Müllerb thinks is inconceivable; but as Tiedemanne has remarked, if we admit this faculty in animals provided with a heart, the progression of the blood must be mainly owing to that viscus; for, after the heart ceases to act, the circulation is soon arrested. The blood, too, only remains fluid, and possesses the faculty of motion, whilst it is in connexion with the living body. When taken from the vessel in which it circulates, it soon coagulates, and loses its motive power. This motion has, by some, - and, according to Brandt, not without grounds, - been presumed to be owing to electro-chemical agency.

Burdache has properly observed, that the old but perfectly correct saying, "ubi stimulus, ibi affluxus," means nothing more than that where the vital activity of an organ is augmented, more blood will be drawn to it; whence it naturally follows, that the progression of blood in the capillaries must be, in some measure, dependent on the activity of the vital manifestations in the tissue. It has been already shown, that if the capillary action be excited by stimulants, a greater flow of blood takes place into that system of vessels; and, as the functions of nutrition and secretion are accomplished by that system, it is obvious, that any increase in the activity of these functions must attract a larger afflux of fluids, and, in this manner, modify the circulation independently of the heart and larger vessels. But this, again, can have but a subor-

dinate influence on the general circulation.

Lastly, Rapsails resolves the whole of the circulation, as he does also, Dr. Allen Thomson, on the Formation of New Bloodvessels, Edinb. 1832; and art. Circulation, in Cyclopædia of Anat. and Physiology, p. 7, Lond. 1836; Alison's Outlines of Physiology, 2d edit., Supplement, Lond. 1836; and Purkinje, in art. Circulatio Sanguinis, Encycl. Wörterb. u. s. w. B. vii. s. 676, Berl. 1831.

Mr. Ancell Lectures on the Physiology and Pathology of the Blood, &c., in Lond.

Lancet, Oct. 26, 1839, p. 147.

b Handbuch, u. s. w., Baly's translation, p. 224, Lond. 1838. See, also, Magendie on the Blood, Amer. Edit. in Bell's Select Medical Library, p. 150, Philad. 1839.

c Op. cit. p. 349.

- d Art. Blut, in Encyclopad. Wörterb. der Medicinisch. Wissenschaft. v. 596, Berlin, 1830.
  - · Die Physiologie als Erfahrungswissenschaft, &c., Band. iv., Leipz. 1832.

C. D. Shultz, Der Lebensprocess im Blute, Berl. 1822.

6 Chimie Organique, p. 364, Paris, 1833.

every other function, into a double action of aspiration and expiration by the tissues concerned. As the blood is the bearer of life to every part of the organism, and of nourishment and reparation to the organs, - to prevent its destination being annulled, a part of the fluid, he says, must be absorbed by the surfaces, which it bathes: these surfaces must attract nutritive juices from the blood, and they must return to the blood the refuse of their elaboration,in other words, they must aspire and expire. Now, this double function cannot take place without the fluid being set in motion, and this motion must be the more constant and uniform as the double function is inherent in every molecule of the surface of the vessels. In this way he accounts for the mercury, placed in a tube communicating with an artery, being kept at the same height near to, or at a distance from, the heart; because, he says, it is not the action of the heart which supports it, but the action of the parietes of the vessels. Every surface, which aspires, provided it is flexible, must be, in its turn, he conceives, attracted by the substance aspired, and, consequently, by the act of aspiration alone, the motions of systole and diastole of the heart and arteries may be explained. When their inner parietes aspire — or assimilate the fluid — the heart will contract; when, on the contrary, they expire, owing to the mutual repulsion between the heart and the fluid, the former will dilate; and, as the movement of the heart is energetic on account of its size, its movements will add to the velocity of the circulation in the arteries, which will, therefore, besides their proper actions of aspiration and expiration, present movements isochronous with the pulsations of the heart. "Add to this accessory cause of arterial pulsations, the movements impressed by the aërial aspiration, which takes place in the lungs, and the circulation of the blood will no longer present insurmountable problems."

All this, we need scarcely say, is ingenious, but nothing more.

# f. Accelerating and Retarding Forces.

The above are the chief accelerating causes of the circulation. There are some others, which at times accelerate, and at others retard; and others, again, that must always be regarded as impeding influences. All these are of a physical character, and apply as well to inert hydraulic machines, as to the pipes of the

human body.

1. Friction always acts as a retarding force. That, which occurs between a solid and the surface on which it moves, can be subjected to calculation, but not so with a fluid; inasmuch as all its particles do not move equally: whilst one part is moving rapidly, another may be stationary, moving slowly, or even in a contrary direction, as is seen in rivers, where the middle of the scream always flows with much greater velocity than the sides. The same thing happens to water flowing through pipes; the water, which is in contact with the sides of the pipe, moves more slowly

than that at the centre. This retarding force is much diminished by the polished state of the inner surface of the blood vessels, as is proved by the circumstance, that if we introduce an inert tube into an artery, the blood will not flow through it for any length of time. Poiseuille infers, from his investigations, that a layer of serum lines the interior of the capillary vessels, which may have some effect in retarding the blood globules in their progress through the intermediate system. Yet the viscosity of the blood, within certain limits, would seem to enable it to pass through the capillary system. Magendie, indeed, pronounces it to be an indispensable

condition for its free circulation through the capillaries.b

2. Gravity may either be an active or retarding force, and is always exerting itself, in both ways, on different sets of vessels. If, for example, the flow of blood to the lower extremity, by the arteries, be aided in the erect attitude by the force of gravity, its return by the veins is retarded by the same cause. Every observer must have noticed, that the pulse of an individual in health beats slower when he is in the recumbent, than in the erect, attitude. This is owing to there being no necessity for the heart to make use of unusual exertions for the purpose of forcing the blood, against gravity, towards the upper part of the body. In therapeutics, the physician finds great advantage from bearing this influence in mind; and, hence, in diseases of the head, — as in inflammation of the brain, in apoplectic tendency, ophthalmia, &c., - he directs the patient's head to be kept raised; whilst, in uterine affections, the horizontal posture, or one in which the lower part of the body is raised even higher than the head, is inculcated; and in ulcers or inflammatory diseases of the lower extremities, the leg is recommended to be kept elevated. Every one, who has had the misfortune to suffer from whitlow, must have experienced the essential difference, as regards the degree of pain, produced by position. If the finger be held down, gravity aids the flow of blood by the arteries, and retards its return by the veins: the consequence is turgescence and painful distension; but if it be held higher than the centre of the circulation, the flow by the arteries is impeded, whilst its return by the veins is accelerated, and hence the marked relief afforded.c

3. Curvatures. — Besides friction, the existence of curvatures has considerable effect on the velocity and quantity of the fluid passing through pipes. A jet will not rise as high from the pipe or adjutage of a reservoir, if there be an angular turn in it, as if the bend were a gradual curve or sweep. The expense of force, produced by such curvatures in arteries, is seen at each contraction of the ventricle, — the tendency in the artery to become straight producing an evident movement, which has been called the locomotion of the artery, and has been looked upon, by some, as the principal cause of the pulse. This motion is, of course,

a Biblioth. Universelle, Novemb. 1835.

Mr. Moseley, in Lond. Medical Gazette, April 15, 1837.

b Magendie, Lectures on the Blood, edit. cit. p. 102, Philad. 1839,

more perceptible the nearer it is to the heart and the greater the vessel; hence it is more obvious at the arch of the aorta; and we can now understand why this arch should be so gradual. We have a striking example of the force, used in this effort at straightening the artery, in the case of the popliteal artery, when the legs are crossed, and a curvature is thus produced. The force is sufficient to raise a weight of upwards of fifty pounds at each contraction of the ventricle, notwithstanding it acts at the extremity of so long a lever. This fact is sufficient to exhibit the inaccuracy of the notion of Bichat and Bricheteau, that the curvatures in the arteries can have no effect in retarding the flow of blood. Such could only be the case, Bichat thinks, if the vessels were empty at each systole.

4. Anastomoses. — The anastomoses of vessels have, doubtless, also some influence on the course of the blood; but it is impossible to appreciate it. The superficial veins are especially liable to have the circulation impeded by compression in the different postures of the body; but, by means of the numerous anastomoses that exist, if the blood cannot pass by one channel, it is diverted into others. Although, however, a forcible compression may arrest or retard the flow by these vessels, a slight degree of support prevents dilatation of the vein by the force of the blood passing into it, and thus favours its motion. The constant pressure of the skin is hence serviceable to the circulation through the subcutaneous veins; and if, by any means, the pressure is diminished, especially in those parts in which the blood has to make its way contrary to gravity — as in the lower extremities — varices or dilatations of the vessels arise, which are remedied by the mecha-

nical compression of an appropriate bandage.

Attempts have been made to calculate the velocity with which the blood proceeds in its course; and how long it would take for a globule of blood, setting out from the left side of the heart, to attain the right side. It is clear, that the data are, in the first place, totally insufficient for any approximation. We know not the exact quantity of blood, contained in the vessels; the quantity sent into the artery at each contraction of the ventricle; the relative velocity of the arterial, venous, and capillary circulations; and if we knew them at any one moment, they are liable to incessant fluctuations, which would preclude any accurate average from being deduced. Were these circumstances insufficient to exhibit the inanity of such researches, the varying estimates of different observers would fully establish it. These assign the time occupied in the circulation from two minutes to fifteen or twenty hours! Moreover, the distance the globules have to traverse must be very various. In the heart, the passage from one side to the other by the coronary vessels is very short, whilst if the blood have to proceed to a remote part of the body, the distance must be considerable.

<sup>2</sup> Clinique Médicale, p. 145, Paris, 1835; and the author's translation in his American Medical Library, Philad. 1837.

Were we to regard the vascular system as forming a single tube - by knowing the weight of the blood and the quantity which the left ventricle is capable of sending forward at each contraction, we could calculate with facility the period that must elapse before the whole mass is distributed. Thus, if we estimate the quantity propelled forward at each contraction of the ventricle to be two ounces; and the whole mass of blood to be 30 pounds, it will require, on an average, about 240 beats of the heart to send it onwards; which can be accomplished in little more than 3 minutes;2 vet, notwithstanding the absence of the requisite data, a recent writer has gone so far as to affirm the average velocity of the blood in the aorta, to be about eight inches per second; whilst "the velocity in the extreme capillaries is found to be often less than one inch per minute." A similar estimate was made by Dr. Young;b Hales, too, estimated the velocity of the blood leaving the heart at 149.2 feet per minute, and the quantity of blood passing through the organ every hour at twenty times the weight of the blood in the body; but the judicious physiologist knows well, that in all operations, which are partly of a vital character, the results of every kind of calculation must be given with caution and humility. In the larger animals, as the whale, the quantity of the fluid circulating in the aorta must be prodigious. Dr. Hunter, in his account of the dissection of a whale, states that the aorta was a foot in diameter, and that ten or fifteen gallons of blood were probably thrown out of the heart at each stroke; so that this vessel is in the whale actually larger than the main pipe of the old water-works at London Bridge; and the water, rushing through the pipe, it has been conceived, had less impetus and velocity than that gushing from the heart of this leviathan.d

These estimates, as to the velocity of the circulatory current, are probably far beneath the truth, inasmuch as experiments have shown, that substances introduced into the venous circulation may be detected in the remotest parts of the arterial circulation in animals larger even than man in less than thirty seconds. Ten seconds after having injected a solution of nitrate of baryta into the jugular vein of a horse, Mr. Blake drew blood from the carotid of the opposite side; after allowing this to flow for five seconds, he received the blood that flowed during the next five seconds into another vessel; and that which flowed after the twentieth second, by which time the action of the heart had stopped, was received into a third vessel. No trace of baryta could be detected in the blood that flowed between the tenth and fifteenth second; but it was discovered in that which flowed between the fifteenth and twen-

e Edinb. Med. and Surg. Journal, Oct. 1841.

For various estimates, see Burdach, Die Physiologie als Erfahrungswissenschaft, iv. 253, Leipzig, 1832; Dr. Allen Thomson, Cyclop. of Anatomy and Physiol. art. Circulation; and Müller's Handbuch, u. s. w., Baly's translation, p. 186.

Med. Literature, Lond. 1813.
 Paley's Natural Theology; and Animal Physiology, p. 75, Library of Useful Knowledge, Lond. 1829.

tieth. From the results of these and other experiments, Dr. Carpenter thinks it difficult to resist the conclusion, that some other force than its contractions must have a share in producing the

movement of the blood through the vessels.a

The velocity of the circulating fluid in the minute vessels is generally thought to be less than in the larger; band their united calibres to be much greater than that of the trunk with which they communicate.c Were this the case, the diminution of velocity would be in accordance with a law of hydrodynamics; - that when a liquid flows through a full pipe, the quantity which traverses the different sections of the pipe, in a given time, must be every where the same; so that where the pipe is wider the velocity diminishes; and, on the contrary, where it is narrower the velocity increases. This would not seem, however, to be consistent with the experiments of Poiseuille, already referred to, which appear to show, that the pressure exerted on the blood in different parts of the body - as measured by the column of mercury, which the blood in different arteries will sustain - is almost exactly the same.

The cause of error in the common belief, that the capacity of the arterial tubes increases in proportion to their distance from the heart, has been explained by Mr. Ferneley.d It is true, he observes, that the sum of the diameters of the branches is considerably greater than that of the trunk. Thus a trunk, 7 lines across, may divide into two branches of 5 lines each, or a trunk of 17 into three branches of 10, 10, and 91; but when their areas are compared, which is the only mode of arriving at their calibres, the correspondence is as close, he thinks, as can be reasonably expected, when the nature of the measurement is taken into account. In the first case, the area of the trunk is represented by the square of 7 — that is 49; whilst the area of each branch will be 25, and the sum of the two will be 50. In the second instance, the area of the trunk will be 17 squared, or 289; whilst that of the branches is the sum of 100, 100, and 90½, making 290½. This will be more strikingly seen from the following table: -

		Trunk.	Branches.			
	Diameter.	Square of Diameter.	Diameter	Sum of Squares of Diameter.		
I.	9	81	7.5 + 5	81-25		
II.	7.2	51-64	6+4	52		
III.	3.5	12.25	3+2	13		
IV.	7.0	49	5 + 5	50		
V.	17	289	10+10+9.5	290.25		
VI.	10	100	7+7+2	102		
VII.	4.5	20.25	3.5+3	21.25		
VIII.	8	64	4+7	65		

a Human Physiology, § 491, Lond. 1842.

b Dr. J. R. Graves, in Lond. Med. Gazette for June 30, 1838, p. 559.

<sup>6</sup> Horner, Special Anatomy and Histology, ii. 172, Philad. 1843.

d Lond. Med. Gaz. Dec. 7, 1839.

e Lond. Med. Gazette, Dec. 7, 1839; and Brit. and For. Med. Rev., April, 1840, Carpenter, Human Physiology, § 476, Lond. 1843.

VOL. II. - 14

It will be observed, that the sum of the squares of the diameters of the branches is in every case slightly more, than the square of the diameter of the trunk. This was found to be somewhat greater in subsequent experiments made by Mr. Paget.<sup>a</sup> The following table gives the ratio of the area of each arterial trunk to the joint area of its branches, as observed by him.

							Trunk.		Branches.
Arch of the aorts	19.0	112/1	112	11219	102/1	100	1	1	1.055
Innominata -	43	4-11	11-11	1-11	184	10-1	1	:	1.147
Common carotid	-				0000		1	100	1.013
External carotid		-		-	-		1	:	1.19
Subclavian -			113	NAME OF	12	-	1	:	1.055
Abdominal aorta	to the l	ast lui	mbar a	arterie	s -	17 -51	1	:	1.183
					Die i	13	1	:	-893
Common iliac		victory.	100	10 - 12	-	-	1	100	-982
External iliac				-		100	1	1:	1.15

It would appear, that where the aorta divides into the common iliacs, or at the division next lower down, the stream is always contracted; an effect, which, Mr. Paget suggests, must be to accelerate the circulation not only in the iliacs themselves, but in the arteries given off from the trunk above them, — as the mesenteric and the renal.

From what has been said, regarding the curvatures and angles of vessels, it will be understood, that the blood must proceed to different organs with different velocities. The renal artery is extremely short, straight, and large, and must consequently transmit the blood very differently to the kidney, from what the tortuous carotid does to the brain; or the spermatic artery to the testicle. A different impulse must, consequently, be given to their corresponding organs by these different vessels. A great portion, however, of the impulse of the heart must fail to reach the kidney, short as the renal artery is, owing to its passing off from the aorta at a right angle; and, hence, the impulse of the blood on the kidney may not be as great

as might be imagined at first sight.

The tortuosity of the carotid arteries is such as to greatly destroy the impetus of the blood; so that but trifling hemorrhage takes place when the brain is sliced away on a living animal, although it is presumed, that one eighth of the whole quantity of blood is sent to the encephalon. Dr. Rush supposed, that the use of the thyroid gland is to break the afflux of blood to the brain; for which its situation between the heart and the head appeared to him to adapt it; and he adduced, as farther arguments, - first, the number of arteries which it receives, although effecting no secretion; secondly, the effect on the brain, which he conceived to be caused by diseases, and by extirpation, of the thyroid; the operation having actually occasioned, in his opinion, in one case, inflammation of the brain, rapidly terminating fatally; and, thirdly, the fact that goître is often accompanied by idiotism. The opinion, however, is so entirely conjectural, and some of the facts, on which it rests, so questionable, that it does not demand serious examination.

This leads us to remark, that the thyroid gland, as well as other organs, with whose precise functions we are totally unacquainted as the thymus, spleen, and supra-renal capsules, - have been conceived to serve as diverticula or temporary reservoirs to the blood, when, owing to particular circumstances, that fluid cannot circulate properly in other parts. Lieutaud having observed, that the spleen is always larger when the stomach is empty than when full, considered that the blood, when digestion is not going on, reflows into the spleen, and that thus this organ becomes a diverticulum to the stomach. The opinion has been indulged by many, with more or less modification. Dr. Rush's view was yet more comprehensive. He regarded the spleen as a diverticulum, not simply to the stomach, but to the whole system, when the circulation is violently excited, as in passion, or in violent muscular efforts, at which times there is danger of sanguineous congestion in different organs; and in support of his view, he invoked the spongy nature of the spleen; the frequency of its distension; the large quantity of blood distributed to it; its vicinity to the centre of the circulation; and the sensation referred to it, in running, laughing, &c. Broussais has still farther extended the notion of diverticula. He affirms, that they always exist in the vicinity of organs, whose functions are manifestly intermittent. In the fœtus, the blood does not circulate through the lungs as when respiration has been established: diverticula, he, therefore, considers to be necessary: these are the thymus and thyroid glands. The kidneys do not act in utero: hence the use of the supra-renal capsules as diverticula. At birth, these organs are either wholly obliterated, if the organs to which they previously acted as diverticula have continuous functions; or they are only partly obliterated, if the functions be intermittent. Thus, the spleen continues as a diverticulum to the stomach, because its functions are intermittent through life; and the thymus disappears, when respiration is established: the liver and the portal system he regards as a reservoir, inservient to the reception of the blood in cases of impediment to the circulation in different parts of the body.

These notions are entirely hypothetical. We shall see, hereafter, that our ignorance of the offices of the spleen, thymus, &c., is extreme; and we have already shown, that much more probable uses can be assigned to the portal system. The insufficiency of Broussais's doctrine of diverticula is strikingly evidenced by the fact, that whilst the thymus gland disappears gradually in the progress of age, the thyroid remains, as well as the supra-renal

capsules.b

The erectile tissues offer a variety in the circulation, which requires some comment. Examples of these occur in the corpora cavernosa of the penis and clitoris; and in the nipple.

Commentaries des Propositions de Pathologie, &c., Paris, 1829; or Drs. Hays and Griffith's translation, p. 214, Philad. 1832.
 Adelon, Physiologie de l'Homme, tom. iii. 328, 2de édit. Paris, 1829.

They appear, according to Gerber, to consist of a plexus or rete of varicose veins enclosed in a fibrous envelope, with relatively minute interspaces, which are occupied and traversed in all directions by arteries, nerves, contractile fibres, and by elastic, fibrous and cellular tissue. Of the particular arrangement of vessels in the corpora cavernosa, mention will be made hereafter: the mode of termination of the arteries in the erectile tissues has not been sufficiently studied, nor are views uniform in regard to their mode of action; some being of opinion, that they afford examples of vital expansibility; but, as before remarked (page 138), the excitation is first induced in the nerves of the part, generally through the influence of the brain, and the turgescence of vessels is a consequence.

The arrangement of the portal system of the liver is also pecu-

liar, and has been given already (p. 87).

## g. The Pulse.

We have had occasion, more than once, to refer to the subject of the pulse, or to the beat felt by the finger when applied over any of the larger arteries. Opinions have varied essentially regarding its cause. Whilst most physiologists have believed it to be owing to distension of the arteries, caused by each contraction of the left ventricle; some have admitted a systole and diastole of the vessel itself; others, as Bichat and Weitbrecht, have thought that it is owing to the locomotion of the artery; others, that the impulse of the heart's contraction is transmitted through the fluid blood, as through a solid body; and others, as Dr. Young and Dr. Parry, that it is owing to the sudden rush forward of the blood in the artery without distension.

Bichat was one of the first, who was disposed to doubt, whether the dilatation of the artery, which was almost universally admitted, really existed; or if it did, whether it were sufficient to explain the phenomenon; and, since his time, numerous experiments have been made by Dr. Parry, the result of which satisfied him, that not the smallest dilatation can be detected in the larger arteries, when they are laid bare during life; nor does he believe, that there is such a degree of locomotion of the vessel as can account for the effect produced upon the finger. He ascribes the pulse to "impulse of distension from the systole of the left ventricle, given by the blood, as it passes through any part of an artery contracted within its natural diameter." Dr. Bostock appears to coincide

Elements of General Anatomy, by Gulliver, p. 298, Lond. 1842.

Mandl, Manuel d'Anatomie générale, p. 197, Paris, 1843.
 Comment. Acad. Imper. Scient. Petropol. ad Ann. 1734 and 1735, Petrop. 1740.

d Croonian Lectures, in Philos. Transact. for 1809, Part i.

e An Experimental Inquiry into the Nature, Causes, and Varieties of the Arterial Pulse, by Caleb Hillier Parry, Lond. 1816; also, Additional Experiments on the arteries of warm-blooded animals, &c., by Ch. Henry Parry, M.D., &c., Lond. 1819; Dr. Prichard, in Transact. of the Provincial Med. and Surg. Association, iv. 16, Lond. 1836.

(Physiology, 3d edit, p. 246, Lond. 1836.

PULSE. 161

with Dr. Parry, if we understand him rightly, or at all. "According to this (Dr. Parry's) doctrine," he remarks, " we must regard the artery as an elastic and distensible tube, which is at all times filled although with the contained fluid not in an equally condensed state, and that the effect produced upon the finger depends upon the amount of this condensation, or upon the pressure which it exercises upon the vessel, as determined by the degree, in which it is capable of being compressed. Where there is no resistance to the flow of the blood along the arteries, there is no variation, it is conceived, in their diameter, and it is only the pressure of the finger or some other substance against the side of an artery that produces its pulse.4

Most of the theories of the pulse take the contractility of the artery too little into account. In pathology, where we have an opportunity of observing the pulse in various phases, we meet with sensations communicated to the fingers, which it is difficult to explain upon any theory, except that of the compound action of the heart and arteries. The impulse is obviously that of the heart, and although the fact of distension escaped the observation of Bichat, Parry, Weitbrecht, Lamure, Döllinger, Rudolphi, Jäger,c and others, we ought not to conclude, that it does not occur. It is, indeed, difficult for us to believe, that such an impulse can be communicated to a fluid filling an elastic vessel without pulsatory distension supervening. In opposition, too, to the negative observations of Bichat and Parry, we have the positive averment of Dr. Hastings, and of Poiseuille, Oesterreicher, Segalas, and Wedemeyer, that the alternate contraction and dilatation of the larger arteries was clearly seen.e

The pulsations of the different arteries are pretty nearly synchronous with that of the left ventricle. Those of the vessels near the heart may be regarded as almost wholly so, but an appreciable interval exists in the pulsations of the more remote vessels.f

We have remarked, that the arterial system is manifestly more or less affected by the nerves distributed to it; that it may be stimulated by irritants, applied to the great nervous centres, or to the nerves passing to it; and this is, doubtless, the cause of many of the modifications of arterial tension, that we notice in disease. No inflammation can affect any part of the system, for any length of time, without both heart and arteries participating, and affording us unequivocal signs of such inflammation. This, however, is a subject that belongs more especially to pathology.

\* Good's Study of Medicine, Physiological Proem to the class Hæmatica.

b Grundriss der Physiologie, &c. Berlin, 1821.

· Tractatus Ahatomico-physiologicus de Arteriarum Pulsû. Vircerb. 1830.

4 Repertoire générale d'Anatomie, &c. par Breschet, 1829, tom. vi. and vii. and Magendie's Journal de Physiol. viii. and ix.

 Burdach's Physiologie als Erfahrungswissenschaft, tom. iv. Leipz. 1832. For a mode of estimating the arterial distension, see Poiseuille, in Magendie's Journal de Physiologie, ix. 44, and Jules Herison's description of an instrument — the Sphygmameter - which makes the action of the arteries apparent to the eye.

Dr. Allen Thomson, art. Circulation, Cyclopædia of Anatomy and Physiology, P. vii. p. 664, Lond. 1836; and Müller's Handbuch, u. s. w., Baly's translation, p. 200.

The ordinary number of pulsations, per minute, in the healthy adult male, is from seventy to seventy-five; but this varies greatly according to temperament, habit of life, position, - whether lying, sitting, or standing, &c. Dr. Guy, from numerous observations, found the pulse, in healthy males, of the mean age of 27 years, in a state of rest, 79 when standing; 70, sitting, and 67, lying; the difference between standing and sitting being 9 beats; between sitting and lying, 3 beats; and between standing and lying, 12 beats. When all exceptions to the general rule were excluded, the numbers were: standing, 81; sitting, 71; and lying, 66; the difference between standing and sitting being 10 beats; between sitting and lying, 5 beats; and between standing and lying, 15 beats. The effect produced upon the pulse by change of posture. Dr. Guy ascribes to muscular contraction, whether employed to change the position of the body, or to maintain it in the same position. In children, the difference between the pulse in the sitting and lying position is often very marked. In a boy, six years of age, now before the author, it amounts to fifteen beats; and Dr. Evansone states, that he has often found the pulse, which at night (during sleep) was 80, full and steady, up to 100 or even 120 during the day, small and hurried, - and this in children six or seven years of age and in perfect health.

In some individuals in health, the number of beats is singularly few. The pulse of a person, known to the author, was on the average thirty-six per minute; and Lizzarid affirms, that he knew a person in whom it was not more than ten. It is not improbable, however, that in these cases, obscure beats may have taken place intermediately, and yet not have been detected. In a case of pericarditis, in which the author felt great interest, the pulse exhibited a decided intermission every few beats, yet the heart beat its due number of times; the intermission of the pulse at the wrist consisting in the loss of one of the beats of the heart. It was not improbable but that in this case the contractility of the aorta was unusually developed by the inflammatory condition of the heart; and that the flow of blood from the ventricle was thus occasionally diminished or entirely impeded. The quickest pulse, which Dr. Elliotsone ever felt, was 208, counted easily, he says, at the heart,

though not at the wrist.

The pulse of the adult female is usually from ten to fourteen beats in a minute quicker than that of the male. In infancy, it is generally irregular, intermitting, and always rapid, and it gradu-

e Practical Treatise on the Management and Diseases of Children, by Messrs. Evan-

son and Maunsell: Amer. Edit. by Dr. Condie, p. 19, Philad., 1843.

Dr. M. Donnel, in Dublin Journal of Med. and Chemical Science, Sept. 1835.
 Guy's Hospital Reports, No. vi. April, 1838, p. 92. See, also, Dr. John M. B. Harden, Amer. Journ. of the Med. Sciences, April, 1843, p. 340.

<sup>&</sup>lt;sup>4</sup> Raccolta D'Opusculi Scientifici, p. 265; and Good's Study of Medicine, Physiological Proem to class iii. Hæmatica. See Cases of Slowness of the Pulse, by Mr. Mayo, Lond. Med. Gaz. May 5, 1838, p. 232.

Human Physiology, p. 215, Lond. 1840.
 Guy, in Guy's Hospital Reports, iii. 112.

ally becomes slower in the progrees of age. It is, of course, impossible to arrive at any accurate estimate of its comparative frequency at different periods of life, but the average of the following numbers, on the authority of Heberden, a Sommering, and Müller, c may, on the whole, be regarded as approximations.

Ages.					Number of beats per minute, according to			
In the embryo,	112	01	112	CON THE	Heberden.	Sömmering.	Müller.	
At birth,					130 to 140	Do.	Do.	
One month, -					120	distantanting	diam'r.	
One year,					120 to 108	120	115 to 130	
Two years, -				-	108 to 90	110	100 to 115	
Three years, -					90 to 80	90	90 to 100	
Seven years, -					72	मार्क्षका व्याप्त हैं।	85 to 90	
Twelve years,					70	CONT. See MOST	ord of the	
Puberty,	-		-	-	inno to the second	80	80 to 85	
Adult,						70	70 to 75	
Old age,				-	A STATE OF THE PARTY.	60	50 to 65	

Researches by MM. Hourmann and Dechambre, do not, however, accord with these estimates in respect to the smaller number of pulsations in the aged. MM. Leuret and Mitivié had suspected an error in this matter from an examination of 71 of the aged inmates of the Bicêtre and La Salpétrière. MM. Hourmann and Dechambre examined 255 women between the ages of 60 and 96,

and found the average number of the pulse to be 82.29.

M. Trousseau, from repeated observations, infers that but little stress ought to be laid on the pulse in the diagnosis of disease in infants. He found, that during the first two weeks, it may vary from 78 to 150; during the second fortnight, from 120 to 164; one to two months, from 96 to 132; two to six months, 100 to 162; six to twelve months, 100 to 160; and from twelve to twenty-one months, 96 to 140. From the observations of MM. Billard, Valleix and others, it would seem that the pulse of the fœtus at the moment it is expelled from the uterus often falls to 83 in the minute, and that, in some minutes afterwards, it rises to 160. In the course of the first day, it falls again to 127, and continues to diminish during the ten first days, the average being then from 87 to 90. These are, however, only the averages: the variations

2 Med. Transact. ii. 21. Also, Blumenbach's Elements of Physiology, by Elliotson

p. 89, 4th edit., Lond. 1828.

Handbuch der Physiologie, Baly's translation, p. 171, Lond, 1838.

Archiv. Général. de Med. pour 1835.

b See, also, Quetelet sur l'Homme, &c., ii. 80, Paris, 1835; Dr. Knox, in Edinb. Med. and Surg. Journ., April, 1837; Dr. Gorham on the Pulse of Infants, in Lond. Med. Gazette, Nov. 25, 1837; Valleix, Clinique des Maladies des Enfans Nouveaunés, Paris, 1838; and Dr. J. M. B. Harden, op. cit.

d Dr. Gorham assigns 130 as the mean number of the pulse from five months to two years old; and 107.63 from two to four years of age, whence the number continues almost the same up to the tenth year.

f Journ. des Connaiss. Méd. Chir. Juillet & Août, 1841; and Amer. Journ. Med. Sciences, Oct. 1841, p. 458, and Jan. 1842, p. 199.

are very great. The sex appeared to have some influence. In infants, from eight days to six months old, the average number of pulsations for boys was 131; for girls, 134; from six to twentyone months, the average for boys was 113; for girls, 126. The state of sleeping or waking had a greater influence. In infants from fifteen days to six months old, the average of the pulse was 140 during waking, 121 during sleep. When the child cried or struggled, he has known it rise from 112 to 160 and 180. On the whole, M. Trousseau concludes, that the pulse of children at the breast varies from 100 to 150. After the first two months, it is a little more frequent in females than in males; and it is about

20 higher in the waking than in the sleeping state.

The pulse - strange to say - may be wholly absent, without the health seeming to be interfered with. A case of this kind is referred to by Prof. S. Jackson, as having occurred in the mother of a physician of Philadelphia. The pulse disappeared during an attack of acute rheumatism, and could never again be observed during her life. Yet she was active in body and mind, and possessed unusual health. In no part of the body could a pulse be detected. Dr. Jackson attended her during a part of her last illness - inflammation of the intestines; no pulse existed. She died whilst he was absent from the city, and no examination was made to elucidate the cause of this remarkable phenomenon.

Between the number of pulsations and respirations there would not appear to be any fixed relation. In many individuals the ratio in health is 4 to 1, but in disease it varies greatly. Dr. Elliotson alludes to a case of nervous disease in a female, at the time in no

danger, whose respiration was 106, and pulse 104.

Dr. Knox has made some observations on the pulsations of the heart and on its diurnal revolution and excitability, d from which he infers: 1. The velocity of the heart's action is in a direct ratio with the age of the individual, - being quickest in young persons, slowest in the aged. There may be exceptions to this, but they do not affect the general law. 2. There are no data to determine the question of an average pulse for all ages. 3. There is a morning acceleration and an evening retardation in the number of the pulsations of the heart, independently of any stimulation by food, &c. 4. The excitability of the heart undergoes a daily revolution; that is, food and exercise affect the heart's action most in the morning and during the forenoon, least in the afternoon, and least of all in the evening. Hence it might be inferred that the pernicious use of

b Quetelet, op. citat. p. 87.

<sup>\*</sup> The Principles of Medicine, founded on the Structure and Functions of the Animal Organism, p. 492, Philad. 1832. See, also, a case of complete disappearance of the beating of the heart, in Gazette Médicale, Nov. 21, 1836; and analogous cases in Parry on the Pulse, Bath, 1816; in Medico-Chirurg. Review, xix. 285, and April, 1836.

e Human Physiology, p. 215, Lond. 1835. See, also, Dr. Ch. Hooker, of New Haven, Conn., in Boston Medical and Surgical Journal, for May 16, 23, &c., 1838. d Edinburgh Medical and Surgical Journal, April, 1837. See, also, Dr. Stratton, in Edinb, Med. and Surg Journ, for Jan. 1843.

USES. 165

spirituous liquors must be greatly aggravated in those who drink before dinner. 5. Sleep does not farther affect the heart's action than by a cessation of all voluntary motion, and by a recumbent position. 6. In weak persons, muscular action excites the action of the heart more powerfully than in strong and healthy individuals; but this does not apply to other stimulants, to wine, for example, or to spirituous liquors. 7. The effects of the position of the body, in increasing or diminishing the number of pulsations, is solely attributable to the muscular exertion required to maintain the body in the sitting or erect posture; the debility may be measured by altering the position of the person from a recumbent to a sitting or the erect position. 8. The most powerful stimulant to the heart's action is muscular exertion. The febrile pulse never equals this.<sup>a</sup>

## h. Uses of the Circulation.

The chief uses of the circulation are, — to transmit to the lungs the products of absorption, in order that they may be converted into arterial blood; and to convey to the different organs this arterial blood, which is not only necessary for their vitality, but is the fluid by which the different processes of nutrition, calorification and secretion are effected. These functions will engage us next. We may remark, in conclusion, that the agency of the blood, as the cause of health or insalubrity, has had greater importance assigned to it than it merits; and that although the blood may be the medium, by which the source of disease is conveyed to other organs, we cannot look to it as the seat of those taints that are commonly referred to it. "Upon the whole," says Dr. Good, "we cannot but regard the blood as, in many respects, the most important fluid of the animal machine; from it all the solids are derived and nourished, and all the other fluids are secreted; and it is hence the basis or common pabulum of every part. And as it is the source of general health, so is it also of general disease. In inflammation, it takes a considerable share, and evinces a peculiar appearance. miasms of fevers and exanthems are harmless to every part of the system, and only become mischievous when they reach the blood; and emetic tartar, when introduced into the jugular vein, will vomit in one or two minutes, although it might require, perhaps, half an hour if thrown into the stomach, and in fact it does not vomit till it has reached the circulation. And the same is true of opium, jalap, and most of the poisons, animal, mineral and vegetable. If imperfectly elaborated, or with a disproportion of some of its constituent principles to the rest, the whole system partakes of the evil, and a dysthesis or morbid habit is the certain consequence; whence tabes, atrophy, scurvy, and various species of gangrene. And if it becomes once impregnated with a peculiar taint, it is wonderful to remark the tenacity with which it retains

a Dr. Guy, op. citat.; and in Edinb. Med. and Surg. Journal, p. 90, Jan. 1841.

b Loc. citat.

it, though often in a state of dormancy and inactivity for years, or even entire generations. For as every germ and fibre of every other part is formed and regenerated from the blood, there is no other part of the system that we can so well look to as the seat of such taints, or the predisposing cause of the disorders I am now alluding to; often corporeal, as gout, struma, phthisis: sometimes

mental, as madness; and occasionally both, as cretinism." This picture is largely overdrawn. Setting aside the pathological allusions, which are erroneous in assigning to the blood what properly belongs to the system of nutrition, how can we suppose a taint to continue for years, or even entire generations, in a fluid which is perpetually undergoing renovation, and, at any distinct interval, cannot be presumed to have one of its quondam particles remaining? Were all hereditary diseases derived from the mother, we could better comprehend this doctrine of taints; inasmuch as, during the whole of fætal existence, she transmits the pabulum for the support of her offspring: the child is, however, equally liable to receive the taint from the father, who supplies no pabulum, but merely a secretion from the blood at a fecundating copulation, and from that moment cannot exert any influence upon the character of his progeny. The impulse to this or that organization or conformation must be given from the moment of union of the particles, furnished by each parent at a fecundating intercourse; and it is probable, that no subsequent influence is exerted even by the mother. She affords the pabulum, but the embryo accomplishes its own construction, as independently of the parents as the chick in ovo. (See, on the morbid influences of the blood, the author's Practice of Medicine, 2d edit., Philad. 1844.)

# i. Transfusion and Infusion.

The operation of Transfusion,—as well as of Infusion of medicinal agents,—was adduced in an early part of this chapter, to prove the course of the circulation to be by the arteries into the veins. Both these operations were suggested by the discovery of Harvey. The former, more especially, was looked upon as a means of curing all diseases, and of renovating the aged ad libitum. The cause of every disease and decay was presumed to reside in the blood, and consequently, all that was conceived to be necessary was to remove the faulty fluid, and to substitute pure blood obtained from a healthy animal in its place.

As a therapeutical agent, the history of this operation does not belong to physiology. The detail of the fluctuation of opinions regarding it, and its total disuse, are given at some length in the Histories of Medicine, to which we must refer the reader.<sup>a</sup> There are some interesting physiological facts, however, that cannot be passed over. MM. Prévost and Dumas found that the vivifying

a Sprengel, Hist. de Méd. par Jourdan, iv. 120, Paris, 1815; Dr. J. P. Kay, art. Transfusion, Cyclopædia of Practical Medicine, P. xxi. 246, Lond. 1834; and E. Gräfe, art Infusion und Transfusion, Encyclop. Wörterb. der Medicin. Wissenschaft. xviii. S. 434, Berlin, 1838.

power of the blood does not reside so much in the serum as in the red particles. An animal bled to syncope was not revived by the injection of water or of pure serum at a proper temperature; but if blood of one of the same species was used, the animal seemed to acquire fresh life, at every stroke of the piston, and was at

length restored.

Transfusion has been revived by Dr. Blundell, of London, and by MM. Prévost and Dumas; the first of whom has employed it with safety, and he thinks with happy effects, in extensive uterine hemorrhage. All these gentlemen remark, that it can only be adopted with perfect safety, in animals of like kinds, or in those, the globules of whose blood are of similar configuration. MM. Prévost and Dumas, Dieffenbach and Bischoff, all agree as to the deadly influence of the blood of the mammalia when injected into the veins of birds. This influence, according to Müller, is in some way connected with the fibrin of the blood. Experiments have certainly shown, that blood deprived of fibrin acts most injuriously

when injected into the vessels."

The introduction of the practice of infusing medicinal agents into the blood was coëval with that of transfusion. Both, indeed, are affirmed to have been commenced in 1657, at the suggestion of Sir Christopher Wren. It is a singular fact, that in cases of infusion, medicinal substances are found to exert their specific actions upon certain parts of the body, precisely in the same manner as if they had been received into the stomach. Tartar emetic, for example, vomits, and castor oil purges, not only as certainly, but with much greater speed; for whilst the former, as before remarked, requires to be in the stomach for fifteen or twenty minutes, before vomiting is excited, it produces its effect in one or two minutes, when thrown into the veins. Dr. E. Hale, Junr. of Boston. has published an interesting pamphlet on this subject. In it he traces the history of the operation, and details several interesting experiments upon animals; and one upon himself, which consisted in the introduction of a quantity of castor oil into the veins.

In this experiment, he did not experience much inconvenience immediately after the injection; but very speedily he felt an oily taste in the mouth, which continued for a length of time, and the

medicine acted powerfully as a cathartic.

Considerable difficulty was experienced in the introduction of

\* Medico-Chirurgical Transactions, ix. 56; and x. 296; and appendix to Ashwell's Practical Treatise on Parturition, London, 1828.

b Biobliotheque Universelle, xvii. 215.

Die Transfuson dies Blutes Berlin, 1828.

Müller's Archiv. 1835; cited in Baly's translation of J. Müller's Handbuch, u. s. w. Magendie on the Blood, in Lond. Lancet, and Bell's Select Medical Library Edit. p. 154, Philad. 1839. See, also, on the different effects of transfusion of arterial and venous blood on animals, Bischoff, in Müller's Archiv. Heft iv. 1838, and Brit. and For. Med. Rev. April, 1839, p. 548.

5 Boylston Medical Prize Dissertations, for the years 1819 and 1821, p. 100, Bos-

ton, 1821.

the oil, to which circumstance Magendie ascribes Dr. Hale's safety; for it is found, by experiments on animals, that viscid fluids, such as oil, are unable to pass through the pulmonary capillaries, in consequence of which the circulation is arrested, and death follows. Such also appears to have been the result of

the experiments of Dr. Hale with powdered substances.

The injection of medicines into the veins has been largely practised at the Veterinary School of Copenhagen, and with complete success, — the action of the medicine being incomparably more speedy, and the dose required much less. It is rarely employed by the physician, except in his experiments on animals; but it is obvious that it might be had recourse to, with happy effects, where narcotic and other poisons have been taken, and where the mechanical means for their removal are not at hand.<sup>b</sup>

#### 3. CIRCULATORY APPARATUS IN ANIMALS.

In concluding this subject, a brief allusion to the circulatory apparatus of other parts of the animal kingdom may be interesting and instructive.

In the mammalia in general, the inner structure of the heart is the same as in man, but its situation differs materially; and, in some of them, as in the stag and pig, two small flat bones, called bones of the heart, exist, where the aorta arises from the left ventricle. In the amphibious mammalia and the cetacea, it has been supposed that the foramen ovale, situate in the septum between the auricles, is open as in the human fætus, to allow those animals to pass a considerable time under water without breathing; but the observations of Blumenbach, Cuvier, and others seem to show, that it is almost always closed. Sir Everard Home found it open in the sea otter, in two instances; but these are regarded by naturalists as exceptions to the general rule.

In several of the web-footed mammalia and cetacea, as in the common otter, the sea otter, and the dolphin, particular vessels are found to be always greatly enlarged and tortuous; — a structure which has been chiefly noticed in the vena cava inferior, and which is supposed to serve the purpose of a diverticulum, whilst the animal is under water, or to receive a part of the returning

blood, and to retain it until respiration can be resumed.

In birds, the structure of the heart universally possesses a singular peculiarity. Instead of the right ventricle having a membranous valve, as in the left, and as in all the mammalia, it is provided with a strong, tense, and nearly triangular muscle, which aids in the propulsion of the blood from the right side of the heart into the lungs. This is presumed to be necessary, in consequence of their lungs not admitting of expansion like those of the mammalia, and of their being connected with numerous air-cells.

\* Précis, &c. ii. 430.


b See on the Infusion of different Medicinal Agents, Mr. Blake, Edinb. Med. and Surg. Journal, April, 1839.

The heart of reptiles or amphibia in general consists either of only one ventricle, or of two ventricles; which freely communicate, so as essentially to constitute but one. The number of auricles always corresponds with that of the ventricles. That the cavities - auricular or ventricular - are, however, single, although apparently double, is confirmed by the fact, that, in all, there is only a single artery proceeding from the heart, which serves both for the pulmonic and systemic circulations. After this vessel has left the heart, it divides into two branches, by one of which a part only of the blood is conveyed to the lungs, whilst the other proceds to different parts of the body. These two portions are united in the heart, and after being mixed together are sent again through the great artery.


In these animals, therefore, aëration is obviously less necessary than in the higher classes; and we can thus understand many of their peculiarities;

how the circulation may continue, when the animal is so situate as to be incapable, for a time, of respiration; and the great resistance to ordinary deranging influences, by which they are characterized.

The marginal figure (Fig. 175) represents the circulatory apparatus of the frog; in which E is the ventricle and D the auricle. From the former arises the aorta F, which soon divides into two trunks. These, after sending branches to the head and neck, turn downwards, (O and P,) and unite in the single trunk A. This vessel sends arteries to the body and limbs, which ultimately terminate in veins, and unite to form the vena cava C. From each of the trunks into which the aorta bifurcates at its origin, arise the arteries F. These are distributed to the lungs, and communicate with the pulmonary veins, which return




In the tadpole state, the circulation is branchial, as in fishes. The heart then sends the whole of its blood to the branchiæ or gills, and it is returned by veins following the course of the dotted lines M and N, (Fig. 175,) which unite to form the descending VOL. II. - 15



Circulation in the Frog.


Fig. 176.



Circulation in Fishes.

aorta. As the lungs undergo their development, small arterial

branches arise from the aorta and are distributed to those organs, and in proportion as these arteries enlarge, the original branchial arteries diminish, until ultimately they are obliterated, and the blood flows wholly through the enlarged lateral trunks, O and P, which, by



Insects appear to be devoid of bloodvessels. Cuvier examined all the organs in them, which, a. Respiratory cells. in red-blooded animals, are most vascular,
Two large arteries. without discovering the least appearance of a b. b. Two large arteries. without discovering the least appearance of the c. c. Mucous glands. d. d. d. Glands connected with the bloodvessel, although extremely minute ramitesticles. c. c. Testicles. fications of the trachea were obvious in every f. Penis. g. Uterus.—(Sir part. Insects, however, both in their perfect E. Home.)

and larva state, have a membranous tube running along the back, in which alternate dilatations and contractions are perceptible; and which has been considered as their heart; but it is closed at both ends, and no vessels can be perceived to originate from it. the innumerable ramifications of the trachea convey the air, and thus, as Cuvier has remarked, "le sang ne pouvant aller chercher l'air, c'est l'air qui va chercher le sang;" ("the blood not being able to go in search of the air - the air seeks the blood.") Carus has, however, discovered a continuous circulation through arteries and veins in a few of the perfect insects, and especially in some larvæ.

Lastly, in many genera of the class vermes, particularly amongst the molluscous and testaceous animals, there is a manifest heart, which is sometimes of a singular structure. Some of the bivalves are affirmed to have as many as four auricles; whilst many animals, as the leech and Lumbricus marinus, have no heart; but circulating vessels exist, in which contraction and dilatation are perceptible.

The marginal figure, (Fig. 177,) of the interior of a leech, given by Sir Everard Home, will exhibit the mode of circulation and respiration in that animal. There is no heart, but a large vessel

Fig. 177.



Interior of the Leech.

exists on each side of the animal. The water is received, through openings in the belly, into the cells or respiratory organs, and passes out through the same.<sup>2</sup>

#### CHAPTER V.

#### NUTRITION.

THE investigation of the phenomena of the circulation has exhibited the mode in which arterial blood is distributed over the body in minute vessels, not appreciable by the naked eye, and often not even with the microscope, and so numerous, that it is impossible for the finest-pointed instrument to be forced through the skin without penetrating one, and perhaps several. We have seen, likewise, that, in the capillary system of vessels, this arterial blood is changed into venous; and it was observed, that in the same system, parts are deposited or separated from the blood, and certain phenomena accomplished, into the nature of which we have now to inquire, beginning with those of nutrition, which comprise the incessant changes that are taking place in the body, both of absorption and deposition, and which effect the decomposition and renovation of each organ. Nutrition is well defined by Adelonb as the action, by which every part of the body, on the one hand, appropriates or assimilates to itself a portion of the blood distributed to it; and, on the other, yields to the absorbing vessels a portion of the materials that previously composed it. The precise character of the apparatus, by which this important function is accomplished, we have no means of knowing. All admit, however, that the old matter must be taken up by absorbents, and the new be deposited by arteries, or by vessels, continuous with them. As the precise arrangement of these minute vessels is not perceptible by the eye, even when aided by powerful instruments, their arrangement has given rise to much controversy. Whilst some have imagined lateral pores in the capillary system of vessels, for the transudation of nutritive deposits; others have presumed, that inconceivably small vessels are given off from the capillary system, which constitute a distinct order, and whose function is to exhale the nutritive substance. Hence, they have been termed exhalants or nutritive exhalants; but the physiological student must bear in mind, that whenever the term is used by writers, they do not always pledge themselves to the existence of any distinct set of vessels, but merely mean the capillary vessel, whatever may be its nature, which is the agent of nutrition, and conveys the blood to the different tissues.

b Physiologie de l'Homme, tom. iii. p. 359, 2de édit., Paris, 1829.

<sup>\*</sup> See, on all this subject, Roget's Animal and Vegetable Physiology, Amer. Edit. . ii., 137, Philad. 1836; art. Circulation, by Dr. Allen Thomson, in Cyclopædia of Anat. and Physiol., p. 641, Lond. 1836; and Carpenter, Principles of General and Comparative Physiology, 2d edit., Lond. 1841.

In investigating the physiology of nutrition, two topics necessarily divide our attention; 1st. The action of decomposition, by which the organ yields to the absorbing vessels a portion of its constituents; and 2dly. The action of composition, by which the organ assimilates a part of the arterial blood that enters it, and supplies the loss, which it has sustained by the previous action of decomposition. The former of these actions obviously belongs to the function of absorption; but its physiology was deferred, in consequence of its close application to the function we are now considering. It comprises what is meant by interstitial, organic or decomposing absorption, and does not require many comments, after the long investigation of the general phenomena of absorption into which we entered. The conclusion, at which we then arrived, was, - that the chyliferous and lymphatic vessels form only chyle and lymph respectively, refusing the admission of all other substances; that the veins admit every liquid which possesses the necessary tenuity; and that, whilst all the absorptions, - which require the substance acted upon to be decomposed and transformed, - are effected by the chyliferous and lymphatic vessels, those that demand no alteration are accomplished through the coats of the veins by imbibition. It is easy, then, to deduce the agents to which we refer the absorption of decomposition. As it is exerted on solids, and as these cannot pass through the coats of the vessel in their solid condition, it follows, that other agents than the veins must accomplish the process; and, again, as we never find in the lymphatic vessels any thing but lymph, and as we have every reason to believe that an action of selection is exerted at their extremities, similar to that of the chyliferous vessels on the heterogeneous substances exposed to them, we naturally look to the lymphatics as the main, if not the sole, organs, concerned in the absorption of solids.

It has been maintained, by some physiologists, that the different tissues are endowed with a vital attractive and elective force, which they exert upon the blood; — that each tissue attracts only those constituents of which it is itself composed; and thus, that the whole function of nutrition is an affair of elective affinity; but this, obviously, cannot be the force that presides over the original formation of the tissues in the embryo. An attraction cannot be exerted by parts not yet in existence. To account for this, it has been imagined, that a peculiar force is destined to preside over formation and nutrition, and to this force various names have been assigned. By most of the ancients it was termed facultas formatrix, nutrix, auctrix; by Van Helmont, Blas alterativum; and by Bacon, motus assimilationis. It was the fucultas vegetativa of Harvey; the anima vegetativa of Stahl; the puissance du moule intérieur of Buffon; the vis essentialis of C. F. Wolff;

De Generatione, Hal. 1759.

De Generatione Animalium, Lond. 1651, p. 170.

d Theoria Medica Vera. Hal. 1708. C Histoire Naturelle, tom. ii.

and the Bildungstrieb or nisus formativus of Blumenbach and most of the German writers. This force is meant, when writers speak of the plastic force, force of nutrition, force of formation, and force of vegetation. Whatever difference there may be in the terms selected, all appear to regard it as charged with maintaining, for a certain length of time, living bodies and all their parts, in the possession of their due composition, organization, and vital properties, and of putting them in a condition, during a certain period of their existence, to produce beings of the same kind as themselves. It is obvious, however, that none of these terms elucidate the intricate phenomena of nutrition, and that none express more than — that living bodies possess a vital force under the action of which formation and nutrition are accomplished.

Under the idea, that all the vessels of the intermediate system are possessed of coats, it is not easy to comprehend how either nutrition or secretion can be accomplished. Lateral pores, as we shall see under the head of Secretion, have been imagined, but this supposed arrangement, provided it existed, which has not been, and cannot easily be, demonstrated, b would not materially elucidate the subject; but if we adopt the opinion, before referred to, that many of the vessels of the capillary system consist of membraneless or coatless vessels, we can comprehend, that by the elective and attractive forces possessed by the tissues and exerted by them on the blood, materials may be obtained from that fluid as it passes through the intermediate system, which may be inservient to the nutrition of the various tissues that are bathed by it, - the mode in which the blood is distributed through the tissues resembling that in which the water of a river is distributed through a marsh, conveying to the vegetable bodies that flourish on its surface, the materials for their nutrition. To adopt the language of an intelligent and philosophical writer, "In every part of the body, in the brain, the heart, the lung, the muscle, the membrane, the bone, each tissue attracts only those constituents of which it is itself composed. Thus the common current, rich in all the proximate constituents of the tissues, flows out to each. As the current approaches the tissue, the particles appropriate to the tissue feel its attractive force, obey it, quit the stream, mingle with the substance of the tissue, become identified with it, and are changed into its own true and proper nature. Meantime, the particles which are not appropriate to that particular tissue, not being attracted by it, do not quit the current, but, passing on, are borne by other capillaries to other tissues, to which they are appropriate, and by which they are apprehended and assimilated. When it has given to the tissues the constituents with which it abounded, and received from them particles no longer useful, and which would become noxious, the blood flows into the veins, to be re-

The Philosophy of Health, by Dr. Southwood Smith, vol. i., p. 405, London, 1835.

a Comment. Societ. Götting. tom. viii.; Elliotson's Blumenbach's Physiology, 4th edit., Lond. 1828, p. 490; and Tiedemann's Traité de Physiologie, par Jourdan, p. 405, Paris, 1831.

Mandl, Manuel d'Anatomie générale, p. 188, Paris, 1843.

turned by the pulmonic heart to the lung, where, parting with the useless and noxious matter it has accumulated, and replenished with new proximate principles, it returns to the systemic heart, by which it is again sent back to the tissues." Particles of blood are seen to quit the current and mingle with the tissues; particles are seen to quit the tissues, and mingle with the current; but all that we can see, as Dr. Smith has remarked, with the best aid we can get, does but bring us to the confines of the grand operations that go on, of which we are altogether ignorant. It would not seem, however, to be necessary for the nutrition of certain parts, that they should receive capillary vessels. There are tissues, commonly termed extra-vascular, in the substance of which, neither injection nor the microscope has exhibited the existence of bloodvessels, which would seem to derive their nourishment from blood flowing in the vessels of adjacent tissues by imbibition. To these belong the crystalline, the epidermis and epithelium, hair,

nails, enamel of the teeth, &c., &c.a

We have said, that the main, if not the sole, agents of the absorption of solids, are the lymphatics. Almost all admit, that they receive the product of absorption; but all do not admit, that the action of taking up solid parts is accomplished immediately by the absorbents. They who think, that a kind of spongy tissue or "parenchyma" is situate at the radicles of the absorbent vessels, believe that this sponge possesses a vital action of absorption, when bodies, possessing the requisite constitution and consistence, are put in contact with it; but they maintain that the solid parts of the body are broken down by the same agents—the extreme arteries - which secreted them, and that, when reduced to the proper fluid condition, they are imbibed by the parenchyma, and conveyed into the lymphatics. If the existence of this sponge were demonstrated, the above explanation would be the only one, perhaps, that could be admitted; for the sponge could scarcely be conceived to do more than imbibe; it could not break down the solid textures, and reduce them to lymph - the only fluid, which, as we have seen, is ever met with in the lymphatic vessels. But its existence is altogether supposititious. Besides, the arrangement has not been invoked in favour of the chyliferous vessels, which are so analogous in their organization and functions to the lymphatics. It has not been contended, that the arteries of the intestinal canal form the chyle from the alimentary matters in the small intestine, and that the office of the chyliferous vessels is restricted to the reception of this chyle, imbibed and brought in contact with their radicles by this ideal sponge or parenchyma.

We have before shown, that there is every reason for the belief, that a vital action of selection and elaboration exists at the very radicles of the chyliferous vessels; and the same may be inferred of the radicles of the lymphatics. The great difficulty is in believing how either exhaling artery or absorbing lymphatic can reduce

Mr. J. Toynbee, Philos. Transact. 1841; and Medico-Chirurg. Rev., April, 1842, p. 426; also, Carpenter, Human Physiology, Amer. Edit., p. 461, Philad. 1843.

the solid matter - of bone, for example - to the necessary constitution and consistence to enter the lymphatics; but we can conceive, that the latter as readily as the former, by virtue of its vital properties - for the operation must be admitted by all to be vital - and by means of its contained fluid, may soften the solid so as to admit of its being received into the vessel. We leave, then, wholly unexplained, the mysterious operation by which these absorbents are enabled to reduce to their elements, bone, muscle, tendon, &c., and to recompose them into the form of lymph. Dr. Bostocka fancifully suggests, that the first step in this series of operations is the death of the part, by which expression he means, that it is no longer under the influence of arterial action. "It therefore ceases to receive the supply of matter which is essential to the support of all vital parts, and the process of decomposition necessarily commences." The whole of his remarks on this subject are eminently gratuitous, and appear to be suggested by his extreme unwillingness to ascribe the process to any thing but physical causes. If there be, however, any one phenomenon of the animal economy, which is more manifestly referable to vital action than another, it is the function of nutrition, both as regards the absorption of parts already deposited, and the exhalation of new; and it is wise to confess our utter ignorance of the mode in which it is accomplished. We know that the blood contains most of the principles that are necessary for the nutrition of organs, and that it must contain the elements of all. Fibrin, albumen, fat, osmazome, salts, &c., exist in it, and these are deposited, as the blood traverses the tissues; but why one should be selected by one set of vessels, as by the exhalants of bone, and another by another set, and in what manner the elements of those, not already formed in the blood, are brought together, is totally unknown to us. Blood has been designated as "liquid flesh," - chair coulante, but something more than simple transudation through vessels is necessary to form it into flesh, and to give it the compound organization of fibrin, gelatin, osmazome, &c. - in the form of muscular fibre and cellular membrane — which we observe in the muscle.

Nothing, perhaps, more clearly exhibits our want of knowledge on the subject than the following vague attempt at solving the mystery by one of the most distinguished physiologists of the age. "Some immediate principles, that enter into the composition of the organs or of the fluids, are not found in the blood, — such as gelatin, uric acid, &c. They are consequently formed at the expense of other principles, in the parenchyma of the organs, and by a chemical action, the nature of which is unknown to us, but which is not the less real, and must necessarily have the effect of

developing heat and electricity."

Histologists have not been content with endeavouring to reduce the different organized textures to primary fibres and filaments, but, by the aid of the microscope, they have attempted to discover the particular arrangement and mode of formation of the

Physiol., edit. cit. p. 625.

constituent corpuscles. The discovery of that valuable instrument gave the impulse, and very soon the scientific world was presented with the results obtained by numerous observers. These observations have been, from time to time, continued until the present day. It is, however, to be regretted, that, until recently, our information, derived from this source, was not as accurate as was desirable. From different quarters the most discordant statements were presented, exhibiting clearly, either that the narrators employed instruments of very diffrent powers, or that they were blinded, or had the vision depraved, by preconceived theories or hypotheses. One of the very first effects of the discovery of the microscope was the detection of a globular structure of the primitive tissue of the body, by Leeuenhoek, an announcement that gave rise to much controversy, which has continued, indeed, till the present time, and has engaged the attention particularly of Prochaska, b Fontana, Sir Everard Home, Mr. Bauer, the brothers Wenzel,d Dr. Milne Edwards, MM. Prévost and Dumas,e Dutrochet, Hodgkin, Raspail, and others. The observations and experiments of Dr. Edwards especially occasioned at the time much interesting speculation and inquiry. They may perhaps be taken as the foundation on which the believers in the globular structure rest their opinions. His views were first published in 1823, in a communication, entitled " Mémoire sur la Structure élémentaire des principaux Tissues Organiques des Animaux;" and in a second article in the Annales des Sciences Naturelles, for December, 1826, entitled " Recherchés microscopiques sur la Structure intime des Tissues Organiques des Animaux." He examined all the principal textures of the body, the cellular tissue, the membranes, tendons, muscular fibre, nervous tissue, the skin, the coats of the bloodvessels, &c. When the cellular tissue was viewed through a powerful lens, it seemed to consist of cylinders; but, by using still higher magnifying powers, these cylinders were found to be formed of rows of globules all of the same size, that is,

Fig. 178.



about the 7300 th or 1000 th of an inch in diameter (Fig. 178); separated from each other, and lying in various directions; crossing and interlacing; some of the rows straight, others bent, and some twisted, forming irregular layers, united by a kind of network. The membranes, which consist of cellular tissue, were found to present exactly the same kind of arrangement, The muscular fibre, when examined in the like manner, was found to be formed of globules also 1000 th part of an inch in

Opera Omnia, &c., Lugdun. Batav. 1722.
 De Structurâ Nervorum, Vind. 1779.
 Sur les Poisons, ii 18.
 De Structurâ Cerebri, Tubing. 1812.


· Bibliothèque Universelle des Sciences et Arts, t. xvii.

t In Drs. Hodgkin and Fisher's translation of W. Edwards, Sur les Agens Physiques, Lond. 1832; and Hodgkin's Lectures on the Morbid Anatomy of the Serous and Mucous Membranes, p. 26, Lond. 1836; or Dunglison's Amer. Med. Lib. Edit. Phil. 1838.

diameter. Here, however, the rows of globules are always parallel. The fibres never intersect each other like those of cellular tissue, and this is the only discernible difference, — the form and size of the globules being alike. The size of the globules, and the linear arrangement they assume, seemed to be the same in all animals that possess a muscular structure. (Fig. 179.)

The nervous structure has, by almost all observers, been esteemed globular — and one of the recent observers has satisfied

himself that this is certainly the most uniform ap-The examination of Dr. Edwards pearance. vielded similar results. It seemed to be composed of lines of globules of the same size with those that form the cellular membrane and the muscles; but holding an intermediate place as to the regularity of their arrangement, and having a fatty matter interposed between the rows. regard to the size of the globules, Dr. Edwards, however, differed materially from an accurate and experienced microscopic observer, Mr. Bauer, b who asserted that the cerebral globules are of various sizes. (Fig. 180.) From the results of his own diversified observations, Dr. Edwards concluded, that "spherical corpuscles, of the diameter of



of those parts, and the functions for which they are destined."

The harmony and simplicity, which would thus seem to reign through the structures of the animal body, attracted great attention to the labours of Dr. Edwards. The vegetable kingdom was subjected to equal scrutiny; and, what seemed still more astounding, it was affirmed, that the microscope proved it also to be constituted of globules exactly like those of the animal, and of the same magnitude, and of an






Fig. 179.





a Dr. Luigi Calori, in Bulletino dello Scienze Medich. di Bologna, Sett. 1836, p. 152.

b Philosoph, Transact. for 1818; and Sir E. Home's Lectures on Comparative Anatomy, vol. iii. lect. 3, Lond. 1823.

animal differs, as we have seen, from the vegetable, in composition; and this difference must exist not only in the whole, but in its parts; so that, even were it demonstrated, that the globules of the beings of the two kingdoms are alike in size, it would by no means follow, that they should be identical in intimate composition.

The discordance which we have deplored, is strikingly applicable to the case before us. The appearance of the memoir of Dr. Edwards excited the attention of Dutrochet, and in the following year his "Researches" on the same subject were published, in which he asserts, that the globules, which compose the different structures of the invertebrated animals, are considerably larger than those of the vertebrated; that the former appear to consist of cells, containing other globules still smaller; and hence he infers, that the globules of vertebrated animals are like wise cellular, and contain series of still smaller globules. Dr. Edwards, in his experiments, found, that the globules of the nervous tissue, whether examined in the brain, in the spinal cord, in the ganglia, or in the nerves, have the same shape and diameter, and that no difference can be distinguished in them, from whatever animal the tissue is taken. Dutrochet, on the other hand, considers, with Sir Everard Home and the brothers Wenzel, that the globules of the brain are cellules of extreme minuteness, containing a medullary or nervous substance, which is capable of becoming concrete by the action of heat and of acids. This structure, he remarks, is strikingly evi-

Fig. 181.

denced in certain molluscous animals; and he instances the small pulpy nucleus, forming the cerebral hemisphere of the *limax rufus*, and the *helix pomatia*, composed of globular, agglomerated cellules, on the parietes of which a considerable number of globular or ovoid corpuscles are perceptible. (Fig. 181.)

M. Dutrochet, again, did not find the structure of the nerves to correspond with that of the brain. He asserts, that the elementary fibres, which enter into their composition, do not consist simply of rows of globules, according to the opinion of Edwards and others, but that they are cylinders of a diaphanous substance, the surface of which is studded with globular corpuscles, and that, as these cover the whole surface of the cylinder, we are led to believe that they are situate internally. After detailing this difference of structure between the brain and the nerves, the former consisting chiefly of nervous corpuscles, the latter chiefly of cylinders or fibres, Dutrochet announces the hypothesis, which exhibits too many indications of having been formed prior to his microscopic investigations, - that these cerebral corpuscles are destined for the production of the nervous power, and that the nervous fibres are tubes, filled with a peculiar fluid, by the agency of which nervimotion is effected. For further developments of the analysis of Dutrochet, the reader is referred to the work itself, which exhibits all the author's ingenuity and enthu-

siasm, but can scarcely be considered historical.

The beautiful superstructure of Dr. Edwards, and the ingenuity of Dutrochet, were, however, most fatally assailed by subsequent experiments, with a microscope of unusual power, by Dr. Hodgkin. The globular structure of the animal tissues, so often developed, and apparently so clearly and satisfactorily established by Dr. M. Edwards, is, we are told by Dr. Hodgkin, a mere deception; and the most minute parts of the cellular membrane, muscles, and nerves had again to be referred to the striated or fibrous arrangement. A part of the discrepancy between Messrs. Edwards and Dutrochet may be explained by the fact of the former using an instrument of greater magnifying power than the latter, who employed the simple microscope only. It has been observed, that when Dr. Edwards used an ordinary lens, the arrangement of a tissue appeared cylindrical, which, with the compound microscope, was distinctly globular. The discordance between Messrs. Edwards and Hodgkin was reconcilable with more difficulty. On the whole subject, indeed, minds were kept in a state of doubt, and the rational physiologist waited for ulterior developments. Messrs. Prévost and Dumas, and Dr. Edwards, farther affirmed, that all the proximate principles - albumen, fibrin, gelatin, &c. - assume a globular form, whenever they pass from the fluid to the solid state, whatever may be the cause producing such conversion. M. Raspailb ranged himself among those who considered, that the ultimate structure of all organic textures is vesicular, and that the organic molecule, in its simplest form, is an imperforate vesicle, endowed with the faculty of inspiring gaseous and liquid substances, and of expiring again such of their decomposed elements, as it cannot assimilate; - properties, which he conceived it to possess under the influence of vitality. His views contain, perhaps, the germ of those that follow, and that now occupy the minds of observers.

The microscopical researches of Schwann and Schleidene led them to affirm, that the new-forming tissues of vegetables originate from a liquid gum or vegetable mucus, and those of animals probably from the liquor sanguinis, which consists essentially of fibrin, after transudation from the capillary vessels. This substance itself, in a state fully prepared for the formation of the tissue, is termed by them intercellular substance and Cytoblastema. In the

Op. citat. p. 466. See, also, Grainger's Elements of General Anatomy, London, 1829.

b Op. citat. § 126.
c Mikroskopische Untersuchungen über die Uebereinstimmung in der Struktur und dem Wachstum der Thiere und Pflanzen, von Dr. Th. Schwann und Schleiden, in Müller's Archiv. p. 137, 1838; and an interesting notice of the same, in Brit. and For. Med. Review, p. 495, April, 1840. Sec, also, Müller, Henle, Carpenter, Wagner, and Mandl.

Fig. 182.



Primary organic cell, showing the cell, membrane, the nucleus and nucleolus.—(Todd and Bowman.)

first instance, it exhibits minute granular points, which grow and become more regular and defined from the agglomeration of the minuter granules around the larger, constituting nuclei or cytoblasts, having, when fully formed, and in fact formed before them, one or more well-defined bodies within them, called nucleoli. From the cytoblasts, cells—primordial cells—are formed. A transparent vesicle grows over each, and becomes filled with fluid; this gradually extends and becomes so large that the cytoblast appears like a small body within its walls. The form of the cells is at first

irregular, then more regular, and they are alternately flattened by pressure against each other, assuming different forms in different tissues. Such is their description of the vegetable cells from which all the tissues of plants take their origin. In like manner, the tissues of animals are formed from a fluid, in which first nucleoli, then nuclei or cytoblasts, and then cells are developed. The globules of lymph, pus, and mucus, according to them, are cells with their walls distinct and isolated from each other; horny tissues are cells with distinct walls, but united into coherent tissues; bone, cartilage, &c., are formed of cells whose walls have coalesced; cellular tissues, tendon, &c., are cells which have split into fibres, and muscles, nerves and capillary vessels are cells of which both the walls and cavities have coalesced.

According to the fashionable doctrine of the day, these cells possess an independent life, and a limited duration, which has no immediate connection with that of the organism, and the continual decomposition, which is taking place in the living body, is connected with the death of the cells of which the several parts are constructed; and for the reintroduction of which into the circulating fluid, the lymphatic system appears to be specially destined. By virtue of this vital power, they not only attract but change the substance brought into contact with them, or in other words, have a power of self-nutrition. That this is probably independent of the nerves is shown by an experiment of Dr. Sharpey, in which the reproduction of a portion of the tail of a salamander took place, although it was cut off, after the organ had been completely paralysed by dissecting out at its root a portion of the spinal cord, together with the arches of the vertebræ.

The doctrine of the development of all the organic tissues from cells is embraced at the present day by almost all histological inquirers; yet there are some who doubt it; and others, who by no means regard it as applicable to all the tissues. Thus, Mandl, one

a Carpenter, Human Physiology, § 645, Amer. Edit., Philad. 1843.

b Ancell, Lectures on the Physiology, &c. of the Blood, in Lond. Lancet, p. 157, April 25, 1840.

C Manuel d'Anatomie générale, p. 549, Paris, 1843.

of the latest histologists, objects to the term cytoblastema as applicable to the matrix or organizing material of the tissues, because it necessarily involves the supposition that it gives origin to cells. According to him, the elements, which are developed in the blastema - as he prefers to call it - do not generally deserve the name of cells, inasmuch as they may either liquify as in the glands; consolidate as in the amorphous membranes; or become transformed directly into fibres, as in the cellular tissue. Mr. Gulliver, a too, has inferred from his observations, that the mere extension of the parietes of the cells is not essential to the formation of all tissues, since fine fibres or fibrils are found in fibrin that has coagulated even out of the body. He has, also, given several figures to exhibit the analogy of structure between false membranes and fibrin coagulated after death, or after the removal of the blood from the body. Schwann, on the other hand, lays down the rule, which he considers of universal application, that all the organic tissues, however different they may be, have one common principle of development as their basis, viz., the formation of cells; - that is to say, nature never unites molecules immediately into a fibre, a tube, &c.; but she always, in the first instance, forms a round cell, or changes, when it is requisite, the cells into the various primary tissues as they present themselves in the adult state; but "how," says Mr. Gulliver, "is the origin of the fibrils, which I have depicted in so many varieties of fibrin, to be reconciled with this doctrine? and what is the proof that these fibrils may not be the primordial fibres of animal textures? I could never see any satisfactory evidence, that the fibrils of fibrin are changed cells; and indeed, in many cases, the fibrils are formed so quickly after coagulation, that their production, according to the views of the eminent physiologist just quoted [Schwann], would hardly seem possible. Nor have I been able to see, that these fibrils arise from the interior of the blood-disks, like certain fibres delineated in the last interesting researches of Dr. Barry."

Mr. T. Wharton Jones, also, in a paper read to the Royal Society, on the 8th of December, 1842, has considered the notion entertained by Dr. Barry, that a fibre exists in the interior of the blood corpuscles, and that these fibres, after their escape from thence, constitute the fibres which are formed by the consolidation of the fibrin of the liquor sanguinis, to be wholly erroneous. He

regards the appearance as altogether illusive.

Dr. Carpenter, in remarking on Mr. Gulliver's figures, all of which, as he properly observes, clearly show, that a small portion of coagulated fibrin contains a far larger number of fibres than we

Proceedings of the Royal Society, No. 56.
 Philosophical Transactions for 1842.

Appendix to Gerber's Anatomy, Atlas, p. 60, and Figs. 244-6, Lond. 1842.
 Lond. and Edinb. Philosoph. Magazine, Oct. 1842; see, also, Andral, Hématologie Pathologique, p. 77, Paris, 1843.

See his interesting report on the Origin and Functions of Cells, in Brit. and For.
 Med. Rev. for Jan. 1843, p. 277.

VOL. II. - 16

can imagine to be contained in the number of blood-disks that would fill the same space, states, that he has lately discovered a very interesting example of a membrane composed almost entirely of matted fibres, which so strongly resembles the delineations of fibrous coagula given by Mr. Gulliver, that he cannot but believe in the identity of the process by which they are produced. This is the membrane enclosing the white of the egg, and forming the animal basis of the shell. If the shell be treated with dilute acid, a tough membrane remains, exactly resembling that which lines it; and if the hen have not been supplied with lime, there is no difference between the two membranes even without the action of acid on the outer one. Each of these membranes consists of numerous laminæ of most beautifully matted fibres intermixed with round bodies exactly resembling exudation cells. It is in the interstices of these fibres that the calcareous particles are deposited, which give density to the shell. These membranes, according to Dr. Carpenter, are formed around the albumen, which is deposited upon the surface of the ovary during its passage along the oviduct, from the interior of which the fibrinous exudation must take place.

It is clear, then, that this doctrine of the origin of all the tissues from cells, cannot be considered established. Nor can ideas be esteemed more fixed in regard to the character of the matrix or blastema. Mandla affirms, that we know not whether it is the albumen or fibrin of the blood. Others, and perhaps the majority of the present day, ascribe it to fibrin, between which, as we have elsewhere seen, and albumen, there is, according to Mulder, Liebig, and others, an almost identity of chemical composition. Fibrin, however, is considered to possess much higher vital properties, and the change of albumen into fibrin has been esteemed the first important step in the process of assimilation.b In the alimentary mass in the small intestines, albumen is found, but not fibrin: in the chyliferous vessels, however, the latter is met with, and its proportion increases as the chyle and lymph proceed onwards in the vessels; whilst that of the albumen diminishes. Such, however, is not rigorously the fact, for on referring to the table slightly modified from that of Gerber, which has been given elsewhere (vol. i., p. 609), it will be seen, that in the afferent lacteals between the intestines and mesenteric glands, the albumen has been found in minimum quantity; in the efferent or central lacteals, from the mesenteric glands to the thoracic duct, in maximum quantity; and in the thoracic duct in medium quantity; whilst the fibrin goes on progressively increasing in the progress of the chyle and lymph onwards. On the other hand, the fat was found to diminish progressively; so that there appears to be more probability that the fibrin is formed from the fat, directly or indirectly, than from the albumen. It would seem not improbable, that some organized substance like pepsin, or diastase in plants, is secreted from the parietes of the chyliferous vessels, b Carpenter, Brit. and For. Med. Rev., Jan. 1843, p. 269. . Op. cit. p. 548.

which occasions a change in the elements of those constituents of the chyle, and is the earliest step of animalization. That there is an essential difference, however, between fibrin and albumen, notwithstanding their affirmed similarity in chemical composition, is shown by the fact, that effused fibrin has a tendency to spontaneous coagulation, whilst albumen requires the agency of heat; and that, as we have seen, there is an appearance of distinct organization in coagulated fibrin. This difference in properties would necessarily induce the belief, that the two substances differ more perhaps in chemical composition than the results of the analyses of Mulder, Liebig, and others, would seem to indicate; and such appears to be proved by the recent analysis of MM. Dumas and Cahours, which have been conducted on a very extensive scale, and show, that the proportion of carbon is seven per cent. less in fibrin than in albumen; whilst that of azote is from eight to nine per cent. more. A correct idea, these gentlemen think, may be formed of the elementary composition of fibrin by considering it a compound of casein, albumen, and ammonia.a

Such is the state of uncertainty in which we are compelled to rest in regard to this important function. None of the views can be esteemed established. They are in a state of transition; and all, perhaps, that we are justified in deducing is, that the vital property, which exists in organizable matters — in the fibrinous portion of the blood, and in the blastema that is furnished by the parents at a fecundating union — gives occasion to the formation of cells, in some cases, of fibres in others; and that the tissues are farther developed through the agency of this cell-life or fibre-life, so as to constitute all the tex-

tures of which the body is composed.b

It is the action of nutrition, that occasions the constant fluctuations in the weight and size of the body, from the earliest embryo condition till advanced life. The cause of the development or growth of organs and of the body generally, as well as of the limit, accurately assigned to such development, according to the animal or vegetable species, is dependent upon vital laws that are unfathomable. Nor are we able to detect the precise mode in which the growth of parts is effected. It cannot be simple extension, for the obvious reason that the body and its various compartments augment in weight as well as in dimension. The rapidity with which certain growths are effected is astonishing. The Bovista giganteum has been known to increase, in a single night, from a mere point to the size of a large gourd, estimated to contain 48,000,000,000 cellules; and supposing twelve hours to have been necessary for its growth, the cells in it must have been produced at the rate of 4,000,000,000 an hour, or more than 66,000,000 a minute, - the greater part of the elements necessary

a Med. Examiner, Oct. 14, 1843, p. 232.

b See, on all this subject, Carpenter, Brit. and For. Med. Rev., Jan. 1843, p. 259.

for this astonishing formation being obtained from the air. But these rapid growths possess little vitality, and their decay is almost as rapid as their productions. Analogous growths — but not to the like extent — occur in the human body, and the same

remark applies to them.

In the large trees of our forests we find a fresh layer or ring added each year to the stem, until the full period of development; and it has been supposed that the growth of the animal body may be effected in a similar manner, both as regards its soft and harder materials, - that is, by layers deposited externally. That the long bones lengthen at their extremities is proved by an experiment of Mr. Hunter. Having exposed the tibia of a pig, he bored a hole into each extremity of the shaft, and inserted a shot. The distance between the shots was then accurately taken. Some months afterwards, the same bone was examined, and the shots were found at precisely their original distance from each other; but the extremities of the bone had extended much beyond their first distance from them. The flat bones also increase by a deposition at their margins, and the long bones by a similar deposition at their periphery, - additional circumstances strongly exhibiting the analogy between the successive development of animals and vegetables. Exercise or rest, freedom from, or the existence of, pressure produce augmentation of the size of organs, or the contrary; and there are certain medicines, as iodine, which occasion the emaciation of particular organs only -- as of the female mammæ. The effects of disease is likewise, in this respect, familiar and striking.

The ancients had noticed the changes effected upon the body by the function we are considering, and attempted to estimate the period at which a thorough conversion might be accomplished, so that not one of its quondam constituents should be present. By some, this was supposed to be seven years; by others, three. It is hardly necessary to say, that in such a calculation we have nothing but conjecture to guide us. The nutrition of the body and of its parts varies, indeed, according to numerous circumstances. It is not the same during the period of growth as subsequently, when the absorption and deposition are balanced, so far at least as concerns the augmentation of the body in one direction. Particular organs have, likewise, their period of development, at which time the nutrition of such parts must necessarily be more active, - the organs of generation, for example, at the period of puberty; the enlargement of the mammæ in the female; the appearance of the beard and the amplification of the larynx in the male, &c. All

these changes occur after a determinate plan.

See the author's General Therapeutics, p. 419, Philad. 1836; and his General

Therapeutics, and Mat. Med. vol. ii. Philad. 1843.

M. Truman, Food and its influence on Health and Disease, &c., p. 229, Lond. 1842.

Carpenter, Animal Physiology, pp. 118, 130, Lond. 1843; and Human Physiology, Amer. Edit., by Dr. Clymer, p. 411, Philad. 1843.

The activity of nutrition appears to be increased by exercise, at least in the muscular organs; hence the well-marked muscles of the arm in the prize-fighter, of the legs in the dancer, &c. The muscles of the male are, in general, much more clearly defined; but the difference between those of the hard-working female and of the inactive male may not be very apparent.

The most active organs in their nutrition are the glands, muscles, and skin, which alter their character — as to size, colour and consistence — with great rapidity; whilst the tendons, fibrous membranes, bones, &c., are much less so, and are altered more slowly



Tattooed Head of a New Zealand Chief.

by the effect of disease. A practice, which prevails amongst certain professions and people, would seem at first sight to show that the nutrition of the skin cannot be energetic. Sailors are frequently in the habit of forcing gunpowder through the cuticle with a pointed instrument, and of figuring the initials of their names upon the arm in this manner: the particles of the gunpowder are thus driven into the cutis vera and remain for life. The operation of tattooing, or of puncturing and staining the skin, prevails in many parts of the globe and especially in Polynesia, where it is looked upon as greatly ornamental. The art is said to be carried to its greatest perfection in the Washington or New Marquesas Islands; where the wealthy are often covered with various designs from head to foot; subjecting themselves to a most painful operation for this strange kind of personal decoration. The operation consists in puncturing the skin with some rude instrument, according to figures previously traced upon it, and then rubbing into the punctures a thick dye, frequently composed of the ashes of the plant that furnishes the colouring matter. The marks, thus made, are indelible. Magendie asks: - "How can we reconcile this phenomenon with the renovation, which, according to authors," (and he might have added, according to himself,) "happens to the skin?" It does not seem to us to be in any manner connected with the nutrition of the skin. The colouring matter is an extraneous substance, which takes no part in the changes constantly going on in the tissue in which it is imbedded; and the circumstance seems to afford a

b Précis, &c. edit. cit. ii. 483.

Lawrence's Lectures on Physiology, &c., p. 411, Lond. 1819.

negative argument in favour of venous absorption. Had the substance possessed the necessary tenuity it would have entered the veins like other colouring matters; but the particles are too gross for this, and hence they remain free from all absorbent influence.

## CHAPTER VI.

## CALORIFICATION.

THE function, which we have now to consider, is one of the most important to organic existence, and one of the most curious in its causes and results. It has, consequently, been an object of interesting examination with the physiologist, both in animals and plants, and as it has been presumed, by a large class of speculatists, to be greatly owing to respiration, it has been a favourite topic with the chemist also. Most of the hypotheses, devised for its explanation, have, indeed, been of a chemical character; and hence it will be advisable to premise a few observations regarding the physical relations of caloric or the matter of heat, - an imponderable body, according to common belief, which is generally distributed throughout nature. It is this that constitutes the temperature of bodies, - by which is meant, the sensation of heat or cold. which we experience, when bodies are touched by us; or the height at which the mercury is raised or depressed by them, in the instrument called the thermometer; - the elevation of the mercury being caused by the caloric entering between its particles, and thus adding to its bulk; and the depression being produced by the abstraction of caloric.

Caloric exists in bodies in two states ; - in the free, uncombined or sensible, and in the latent or combined. In the latter case, it is intimately united with the other elementary constituents of bodies, and is neither indicated by the feelings nor by the thermometer. It has, consequently, no agency in the temperature of bodies; but, by its proportion to the force of cohesion, it determines their condition; - whether they shall be solid, liquid or gaseous. In the former case, caloric is simply interposed between the molecules, and is incessantly disengaged, or abstracted from surrounding bodies; and, by impressing the surface of the body or by acting upon the thermometer, it indicates to us their temperature. Equal weights of the same body, at the same temperature, contain the same quantities of caloric; but equal weights of different bodies, at the same temperature, have by no means the same quantities. The quantity, which one body contains, compared with that in another is called its specific caloric, or specific heat; and the power or property, which enables bodies to retain different quantities of caloric, is called capacity for caloric. If a pound of water, heated to 156°, be mixed with a pound of quicksilver at 40°, the resulting temperature is 152°,—instead of 98°, the exact mean. The water, consequently, must have lost four degrees of temperature, and the quicksilver gained 112°; from which we deduce, that the quantity of caloric, capable of raising one pound of mercury from 40° to 152° is the same as that required to raise one pound of water from 152° to 156°; or, in other words, that the same quantity of heat, which raises the temperature of a pound of water four degrees, raises the same weight of mercury one hundred and twelve degrees. Accordingly, it is said, that the capacity of water for heat is to that of mercury, as 28 to 1; and that the specific heat is

twenty-eight times greater.

All bodies are capable of giving and taking free caloric, and consequently, all have a temperature. If the quantity given off be great, the temperature of the body is elevated. If it take heat from the thermometer, it is cooler than the instrument. In inorganic bodies, the disengagement of caloric is induced by various causes: such as electricity, friction, percussion, compression, the change of condition from a fluid to a solid state; and by chemical changes, giving rise to new compounds, so that the caloric, which was previously latent, becomes free. If, for example, two substances, each containing a certain amount of specific heat, unite, so as to form a compound, whose specific heat is less, a portion of caloric must be set free, and this will be indicated by a rise in the temperature. It is this principle, which is chiefly concerned in some of the theories of calorification.

The subject of the equilibrium and conduction of caloric has already been treated of, under the sense of touch (vol. i., p. 103); where several other topics were discussed, that bear more or less upon the present inquiry. It was there stated, that inorganic bodies speedily attain the same temperature, either by radiation or conduction; so that the different objects in an apartment will exhibit the same degree of heat by the thermometer; but the temperature of animals, being the result of a vital operation, they retain the degree of heat peculiar to them, with but little modification from external temperature. There is a difference, however, in this respect, sufficient to cause the partition of animals into two great divisionsthe warm-blooded and the cold-blooded; the former comprising those whose temperature is high, and but little influenced by that of external objects; - the latter those whose temperature is greatly modified by external influences. The range of the temperature of the warm-blooded — amongst which are all the higher animals — is limited; but of the cold-blooded extensive. The following table exhibits the peculiar temperature of various animals in round numbers; - that of man being 98° or 100°, when taken under the tongue. The temperature in the axilla is something less. In the

<sup>\*</sup> Turner's Elements of Chemistry, 5th Amer. Edit., by Prof. F. Bache, from 5th London, p. 5, Philad. 1835.

latter situation, Dr. Edwardsa found it to vary, in twenty adults, from 96° to 99° Fahrenheit, the mean being 97°.5.

Animals.		Observers.	Temperature.
	U 154	Capt. Lyon.b	107
Arctic fox,		Do. 7	
Arctic wolf,		Pallas.e 5	105
Squirrel,	E 11 10 20	Do. 7	mich algeria
Hare,	S	Scoresby.d 5	104
Whale,	-	Pallas.	103
Arctomys citillus, zizil, - in summer,		Pallas.	80 to 84
Do. when torpid, -	5 5	Prévost and Duma	
Goat,	NO.	Do. ?	
Bat, in summer,		Do. }	102
Musk,		Do.	101 or 102
Marmota babac, - Bobac,		Do.	101
House mouse,		Do.	101 or 102
Arctomys marmota, marmot - in summer,		Do.	43
Do. when torpid, -		Delaroche.	100 to 104
Rabbit,		Capt. Lyon.	100
Polar Bear,		Martine.	
Dog,	10 11	Do.	
Cat,		Do.	100 to 103
Swine,		Do. (	100 10 100
Sheep,	-	Do. )	
Ox,			100 to 102
Guinea-pig,	100 H 34	Delaroche.	99
Arctomys glis,		Pallas.	98
Shrew,	-	Do.	96
Young wolf,		Do.	30
Fringilla arctica, Arctic finch, -		Braun.s	111
Rubecola, redbreast,	-	Pallas.	110 111
Fringilla linaria, lesser red poll, -	-	Do.	110 or 111
Falco palumbarius, goshawk,		Do. 3	110
Caprimulgus Europæus, European goat-st	ucker,	Do. 5	100 4- 110
Emberiza nivalis, snow-bunting, -	- 0 - 7	Do.	109 to 110
Falco lanarius, lanner,	the state of	Do.	
Fringilla carduelis, goldfinch, -		Do.	100
Corvus corax, raven,		Despretz.b >	109
Turdus, thrush, (of Ceylon,)	-	J. Davy.	
Tetrao perdrix, partridge,	- 1 -	Pallas.	
Anas clypeata, shoveler,	-	Do.	
Tringa pugnax, ruffe,		Do.	
Scolopax limosa, lesser godwit, -		Do.	
Tetrao tetrix, grouse,		Do.	108
Fringilla brumalis, winterfinch, -	2 0 150	Do.	STREET, LOT
Loxia pyrrhula,		Do.	
Falco nisus, sparrowhawk,		Do.	
Vultur barbatus,		Do.	
THE RESERVE OF THE RE	200 -	The second second second second	Description of the later of the

De l'Influence des Agens Physiques, &c., Paris, 1824; or Hodgkin and Fisher's translation, Lond. 1832.

b Parry's Second Voyage to the Arctic Regions.

4 An Account of the Arctic Regions, Edinb. 1820.

· Bibliothèque Univers. xvil. 294.

Annales de Chimie, xxvi. 337, Amst. 1824.
Edinb. Philos. Journal, Jan. 1826.

Nov. Species Quadruped. de Glirium Ordine, Erlang. 1774.

Med and Philos. Essays, Lond. 1740; and De Similibus Animalibus et Animal. Calore, &c., Lond. 1740. 8 Nov. Comment. Acad. Petropol. xiii. 419.

Anser pulchricollis, -	+ dro	emo	e nd	1-box	Pallas.	1	
Colymbus auritus, dusky grebe	0,	-	1-00	3-571	Do	1000	107
Tringa vanellus, lapwing, (wo		1.)	-	-	Do.	7	107
Tetrao lagopus, ptarmigan,		-	-	1	Do.	)	
Fringilla domestica, house-spar	row.		1- 1	14	Do.		107 to 111
Strix passerina, little owl,	-	-	-	-	Do.	7	
Hamatopus estralagus, sea-pie,					Do.		
Anas penelope, wigeon, -				-	Do.	(	106
Anas strepera, gadwall, -	-				Do.	1	
Pelecanus carbo,	-			1271	Do.		o tent with
Falco ossifragus, sea-eagle,				h-orn	Do.	3	
Fulica atra, coot,				-	Do.	(	105
Anas acuta, pintail-duck,					Do.	(	THE PROPERTY.
Falco milvus, kite, (wounded,)				Litte	Do.	3	12 10 10 10
Merops apiaster, bee-eater,	-	-	1000	1110	Do.	8	104
Goose,		-	1	1	Martine.	3	
Hen,					Do.	1	an an annual season
Dove,					Do.	>	103 to 107
Duck,	0-10	0 11		Barrie	Do.	1	
Ardea stellaris,				400	Pallas.	3	
Falco albicollis,					Do.	1	103
Picus major,					Do.	(	100
Cossus ligniperda,					Schultze.	,	89 to 91
Shark,	3111	13.11		E TO	I. Davy.		83
Torpedo Marmorata, -	31 4	SE CONTRACTOR	251 1	920	Rudolphi.a		74
a or pour ereur moraru,					readorbur.		12

According to the above table it will be observed, that the inhabitants of the Arctic regions — whether belonging to the class of mammalia or birds — are among those whose temperature is highest. That of the Arctic fox is, indeed, probably higher than the amount given in the table, being taken after death, when the temperature of the air was as low as — 14° of Fahrenheit, and when loss of

heat may be supposed to have taken place rapidly.

The temperature of the smaller insects it is, of course, impracticable to indicate; but we can arrive at an approximation in those that congregate in masses, as the bee and the ant; for it is impossible to suppose, with Maraldi, that the augmented temperature is dependent upon the motion and friction of the wings and bodies of the busy multitudes. Juch found, that when the temperature of the atmosphere was — 18° of Fahrenheit, that of a hive of bees was 44°: in an ant-hill, the thermometer stood at 68° or 70°, when the temperature of the air was 55°; and at 75°, when that of the air was 66°; and Hausmann° and Renggerd saw the thermometer rise, when put into narrow glasses in which they had placed scarabæi and other insects. Berthold detected the elevation of heat only when several insects were collected together, not in one isolated from the rest. This, Mr. Newport affirms, must have arisen

· De Animal, Exsanguium Respiratione, p. 65.

· Tiedemann, op. citat. p. 511.

a Grundriss der Physiol., &c., band i. 166; Tiedemann's Traité de Physiologie, par Jourdan, p. 498, Paris, 1831; and P. H. Bérard, art. Chaleur Animale, in Dict. de Méd. 2de édit. vii. 177, Paris, 1834.

<sup>&</sup>lt;sup>4</sup> Physiologische Untersuchung. über die Insecten, p. 40, Tübing, 1817.

r Philosoph. Transact. for 1837, part ii. p. 259. See Dr. W. B. Carpenter, Principles of General and Comparative Physiology, p. 372, Lond. 1839; and some experi-

from his having ascertained the temperature only whilst the insect was in a state of rest: for Mr. Newport found, that although during such a state, the temperature of the insect was very nearly or exactly that of the surrounding medium, yet when the insect was excited or disturbed, or in a state of great activity from any cause, the thermometer rose in some instances, even to 20° Fahr. above the temperature of the atmosphere, — for instance, to 91°, when

the heat of the air was 71°.

The power of preserving their temperature within certain limits, is not, however, possessed exclusively by animals. The heat of a tree, examined by Mr. Hunter, was found to be always several degrees higher than that of the atmosphere, when the temperature of the air was below 56° of Fahr.; but it was always several degrees below it when the weather was warmer. Some plants develope a considerable degree of heat, during the period of blooming. This was first noticed by De Lamarck in the Arum italicum. In the Arum cordifolium, of the Isle of Bourbon, Hubert found, when the temperature of the air was 80°, that of the spathe or sheath was as high as 134°; and Bory De St. Vincent observed a similar elevation, although to a less degree, in the Arum esculentum, esculent arum or Indian kale.

The animal body is so far influenced by external heat as to rise or fall with it; but the range, as we have already remarked, is limited in the warm-blooded animal, -- more extensive in the cold-blooded. Dr. Currie found the temperature of a man plunged into sea-water at 44° sink, in the course of a minute and a half after immersion, from 98° to 87°; and, in other experiments, it descended as low as 85°, and even to 83°.e It was always found, however, that, in a few minutes, the heat approached its previous elevation; and in no instance, could it be depressed lower than 83°, or 15° below the temperature at the commencement of the operation. Similar experiments have been performed on other warm-blooded animals. Hunter found the temperature of a common mouse to be 99°, when that of the atmosphere was 60°: when the same animal was exposed, for an hour, to an atmosphere of 15°, its heat had sunk to 83°; f but the depression could be carried no farther. He found, also, that a dormouse, - whose heat in an atmosphere at 64°, was 81½°, - when put into air, at 20°, had its temperature raised, in the course of half an hour to 93°; an hour after, the air being at 30°, it was still 93°; another hour after. the air being at 19°, the heat of the pelvis was as low as 83°, an experiment, which strongly proves the great counteracting influence exerted, when animals are exposed to an unusually low ments proving the same point in Dr. M. Paine's Medical and Physiological Commentaries, ii. 76, New York, 1840. .

Philos. Transact. 1775 and 1778.
 Voyage dans les Quatre Principales Iles des Mers d'Afrique, ii. 66.

d Sir J. E. Smith, Introduction to the Study of Botany, 7th edit., by Sir W. J. Hooker, p. 47, Lond. 1833; Tiedemann, p. 492; and Carpenter, op. citat. p. 368.
Philos. Transact. for 1792, p. 199.

temperature. In this experiment, the dormouse had maintained its temperature about 70° higher than that of the surrounding medium, and for the space of two hours and a half. In the hibernating torpid quadruped the reduction of temperature, during their torpidity, is considerable. Jennera found the temperature of a hedgehog, in the cavity of the abdomen, towards the pelvis, to be 95°, and that of the diaphragm 97° of Fahrenheit, in summer, when the thermometer in the shade stood at 78°; whilst in the winter, the temperature of the air being 44°, and the animal torpid, the heat in the pelvis was 45°, and of the diaphragm 48½°. When the temperature of the atmosphere was at 26°, the heat of the animal, in the cavity of the abdomen, where an incision was made, was reduced as low as 30°; but - what singularly exhibits the power, possessed by the system, of regulating its temperature, - when the same animal was exposed to a cold atmosphere of 26° for two days, its heat, in the rectum, was raised to 93°, or 67° above that of the atmosphere. At this time, however, it was lively and active, and the bed, on which it lay, felt warm. In the cold-blooded animal, we have equal evidence of the generation of heat. Hunter found, that the heat of a viper, placed in a vessel at 10°, was reduced in ten minutes, to 37°; in the next ten minutes, the temperature of the vessel being 13°, it fell to 35°; and in the next ten minutes, the vessel being at 20°, to 31°. In frogs, he was able to lower the temperature to 31°; but beyond this point it was not possible to lessen the heat, without destroying the animal.

In the Arctic regions, the animal temperature appears to be steadily maintained, not withstanding the intense cold that prevails; and we have already seen, that the animals of those hyperborean latitudes possess a more elevated temperature than those of more genial climes. In the enterprising voyages, undertaken by the British government for the discovery of a northwest passage, the crews of the ships were frequently exposed to the temperature of -40° or -50° of Fahrenheit's scale; and the same thing happened during the disastrous campaign of Russia in 1812, in which so many of the French army perished from cold. The lowest temperature noticed by Captain Parry was -55° of Fahrenheit. Captain Franklin, on the northern part of this continent, observed the thermometer on one occasion - Feb. 7, 1827, - as low as -58° of Fahrenheit; and M. Von Wrangele states that, in January, on the north coast of Siberia, the cold reaches -59° of Fahrenheit; but the greatest observed natural cold was marked by Cap-

<sup>\*</sup> Hunter on the Animal Economy, p. 99, 2d edit., Lond. 1792, or Owen's edit. in Bell's Select Med. Lib. p. 165, Philad. 1840. See, also, Müller's Handbuch, u. s. w., Baly's translation, p. 76, Lond. 1838.

Journal of a Voyage for the Discovery of a Northwest Passage, Amer. Edit. p. 130,

d Narrative of a Second Expedition to the Shores of the Polar Sea, &c., Amer. Edit. p. 245, Philad. 1835.

e Reise des Kaiserlich. Russischen Flotten, Lieutenants Ferdinand Von Wrangel, längs der Nordküste von Siberien, u. s. w., Berlin, 1839, translated in Harper's Family Library.

tain Back<sup>a</sup> in his expedition to the Arctic regions: on the 17th of January, 1834, the thermometer stood at —70° of Fahrenheit, or 102° below the freezing point!

During the second voyage of Captain Parry, the following temperatures of animals, immediately after death, were taken princi-

pally by Captain Lyon.

1821.  Nov. 15. An Arctic fox 1063° 14°  Dec. 3. Do 1013 5  Do 3								Temperature of the			
Nov. 15. An Arctic fox $\frac{106\frac{3}{4}^{\circ}}{100}$ - $\frac{14^{\circ}}{100}$ Dec. 3. Do $\frac{101\frac{1}{2}}{100}$ - $\frac{5}{100}$ - $\frac{5}{100}$	1821.							Animal.	1111	Atmosphere.	
Dec. 3. Do. $\frac{101\frac{1}{2}}{100} = \frac{5}{100}$		An Arctic fo	X	191	1 -1	30,533	14.5	10630	200	- 14°	
The state of the s	Dec. 3.	Do.	201	No.	45	465744	1000		-		
		Do.	- '	-	-	-	-		-		
111 201	11.	Do.	211	-	-	1955011	-	1014	-	- 21	
15. Do 99 <sup>3</sup> 15	15.	Do.	- 11	-	R.J.	1	11-11		-		
17. Do 98 10	17.	Do.	- 11	100	131	I Phy			-		
19. Do 99\frac{3}{4} 14		Do.	11	Litera	1000	10300		994		- 14	
1822.										Hadayl W	
Jan. 3. Do $104\frac{1}{2}$ - $-23$			-	-				and the second s	-		
9. A white hare 101 21				O FILL	YALL	missin.	11:20)		-		
10. An Arctic fox 100 15				07:111	1572	1151	1873		10		
17. Do 106 32			-	1121	to Sur	1000	BIEN		10		
24. Do 103 27	24.		-		-		-		15.00		
Do 103 27			-		100	*			-		
Do. 102 - 25	Separation of the last		-	-	-	111			15		
27. Do 101 32		( N - 200 10 2	-11	11:11	19714	11/09	1		150		
Feb. 2. A wolf 105 27	Feb. 2.	A wolf	10	tinto	With 1	115	IUMI	105	-	- 27	

These animals must, therefore, have to maintain a temperature at least 100° higher than that of the atmosphere, throughout the whole of winter; and it would appear as if the counteracting energy becomes proportionately greater as the temperature is more depressed. It is, however, a part of their nature to be constantly eliciting this unusual quantity of caloric, and therefore they do not suffer. Where animals, not so accustomed, are placed in an unusually cold medium, the efforts of the system rapidly exhaust the nervous energy; and when this becomes so far depressed as to interfere materially with the function of calorification, which we shall find is to a certain extent under the nervous influence, the temperature sinks, and the individual dies lethargic - or, as if struck with apoplexy. The ship Endeavour, being on the coast of Terra del Fuego, on the 21st of December, 1769, Messrs. Banks, Solander, and others were desirous of making a botanical excursion upon the hills on the coast, which did not appear to be far distant. The party, consisting of eleven persons, were overtaken by night on the hills, during extreme cold. Dr. Solander, who had crossed the mountains which divide Sweden from Norway, knowing the almost irresistible desire for sleep produced by exposure to great cold, more especially when united with fatigue, enjoined his companions to keep moving, whatever pains it might cost them, and whatever might be the relief promised by an indulgence in rest. "Whoever sits down," said he, "will sleep, and whoever sleeps will wake no more." Thus admonished, they set forward, but

<sup>\*</sup> Narrative of the Arctic Land Expedition to the mouth of the Great Fish river, &c., in the years 1833, 1834, and 1835, Lond. 1836.

b Op. citat. p. 157.

whilst still upon the bare rock, and before they had got among the bushes, the cold suddenly became so severe as to produce the effects that had been dreaded. Dr. Solander himself was the first who found the desire irresistible, and insisted on being suffered to lie down. Mr. Banks, (afterwards Sir Joseph,) entreated and remonstrated in vain. The doctor lay down upon the ground, although it was covered with snow; and it was with the greatest difficulty that his friend could keep him from sleeping. Richmond, one of the black servants, began to linger and to suffer from the cold, in the same manner as Dr. Solander. Mr. Banks, therefore, sent five of the company forward to get a fire ready at the first convenient place they came to; and himself, with four others, remained with the doctor and Richmond, whom, partly by persuasion and partly by force, they carried forward; but when they had got through the birch and swamp, they both declared they could go no farther. Mr. Banks had again recourse to entreaty and expostulation, but without effect. When Richmond was told, that if he did not go on, he would, in a short time, be frozen to death, he answered, that he desired nothing but to lie down and die. Dr. Solander was not so obstinate, but was willing to go on, if they would first allow him to take some sleep, although he had before observed, that to sleep was to perish. Mr. Banks and the rest of the party found it impossible to carry them, and they were consequently suffered to sit down, being partly supported by the bushes, and, in a few minutes they fell into a profound sleep. Soon after, some of the people, who had been sent forward, returned with the welcome intelligence, that a fire had been kindled about a quarter of a mile farther on the way. Mr. Banks then endeavoured to rouse Dr. Solander, and happily succeeded, but, although he had not slept five minutes, he had almost lost the use of his limbs, and the soft parts were so shrunk, that his shoes fell from his feet. He consented to go forward with such assistance as could be given him, but no attempts to relieve Richmond were successful. He, with another black left with him, died. Several others began to lose their sensibility, having been exposed to the cold near an hour and a half, but the fire recovered them.a

The preceding history is interesting in another point of view besides the one for which it was more especially cited. Both the individuals, who perished, were blacks, and it has been a common observation, that they bear exposure to great heat with more impunity, and suffer more from intense cold, than the white variety of the species. As regards inorganic bodies, it has been satisfactorily shown, that the phenomena of the radiation of caloric are connected with the nature of the radiating surface; and that those surfaces, which radiate most, possess, in the highest degree, the absorbing power; in other words, bodies that have their temperatures most readily raised by radiant heat are those that are most

<sup>\*</sup> See, on the effects of Cold, Sir H. Halford, in Lond. Med. Gazette, for March 11, 1837, p. 902.

VOL. II. - 17

easily cooled by their own radiation. In the experiments of Professor Lesliea it was found, that a clean metallic surface produced an effect upon the thermometer equal to 12; but when covered with a thin coat of glue its radiating power was so far increased as to produce an effect equal to 80; and, on covering it with lampblack, it became equal to 100. We can thus understand why, in the negro, there should be a greater expense of caloric than in the white, owing to the greater radiation; not because as much caloric may not have been elicited as in the white. In the same manner we can understand that, owing to the greater absorbing power of his skin, he may suffer less from excessive heat than the white; and this is perhaps the great use of the dark rete mucosum. To ascertain, whether such be the fact, the following experiments were instituted by Sir Everard Home. He exposed the back of his hand to the sun at twelve o'clock, with a thermometer attached to it, another thermometer being placed upon a table with the same exposure. The temperature, indicated by that on his hand, was 90°; by the other, 102°. In forty-five minutes, blisters arose, and coagulable lymph was thrown out. The pain was very severe. In a second experiment, he exposed his face, eyelids, and the back of his hand to water heated to 120°; in a few minutes they became painful; and, when the heat was farther increased, he was unable to bear it; but no blisters were produced. In a third experiment, he exposed the backs of both hands, with a thermometer upon each, to the sun's rays. The one hand was uncovered; the other had a covering of black cloth, under which the ball of the thermometer was placed. After ten minutes, the degree of heat of each thermometer was marked, and the appearance of the skin examined. This was repeated at three different times. The first time, the thermometer under the cloth stood at 91°, the other thermometer at 85°; the second time, they indicated respectively 94° and 91°; and the third time, 106° and 98°. In every one of these trials, the skin, that was uncovered, was scorched, whilst the other had not suffered in the slightest degree. From all his experiments, Sir Everard concludes, that the power of the sun's rays to scorch the skin of animals is destroyed, when applied to a black surface; although the absolute heat, in consequence of the absorption of the rays, is greater.

When cold is applied to particular parts of the body, the heat of those parts sinks lower than the minimum of depressed temperature. Although Hunter was unable to heat the urethra one degree above the maximum of elevated temperature of the body, he succeeded in cooling it 29° lower than the minimum of depressed temperature, or to 58°. He cooled down the ears of rabbits until they froze; and when thawed, they recovered their natural heat and circulation. The same experiment was performed on the comb and wattles of a cock. Resuscitation was, however, in no instance

On Heat, Edinb. 1804; and Dr. Stark, in Philosoph. Transact. part ii. for 1833.
Lect. on Comp. Anat. iii. 217, Lond. 1823.

practicable where the whole body had been frozen.a The same observer found, that the power of generating heat, when exposed to a cooling influence, was possessed even by the egg. An egg, which had been frozen and thawed, was put into a cold mixture along with one newly laid. The latter was seven minutes and a half longer in freezing than the other. In another experiment, a fresh-laid egg, and one which had been frozen and thawed, were put into a cold mixture at 15°; the thawed one soon rose to 32°, and began to swell and congeal; the fresh one sank to 291°, and in twenty-five minutes after the dead one, it rose to 32°, and began to swell and freeze. All these facts prove, that when the living body is exposed to a lower temperature than usual, a counteracting power of calorification exists; but that, in the human species, such exposure to cold is incapable of depressing the temperature of the system lower than about 15° beneath the natural standard. In fish, the vital principle can survive the action even of frost. Captain Franklin found, that those which they caught in Winter Lake, froze as they were taken out of the net; but if, in this completely frozen state, they were thawed before the fire, they recovered their animation. This was especially the case with a carp, which recovered so far as to leap about with some vigour after it had been frozen for thirty-six hours.b

On the other hand, when the living body is exposed to a temperature greatly above the natural standard, an action of refrigeration is exerted; so that the animal heat cannot rise beyond a certain number of degrees; - to a much smaller extent in fact than it is capable of being depressed by the opposite influence. Boerhaave maintained the strange opinion, that no warm-blooded animal could exist in a temperature higher than that of its own body. In some parts of Virginia, there are days in every summer, in which the thermometer reaches 98° of Fahrenheit; and in other parts of this country, it is occasionally much higher. The meteorological registers show it to be at times as high as 108° at Council Bluffs, in Missouri; at 104° in New York; and at 100° in Michigan; d whilst in most of the states, on some days of summer, it reaches 96° or 98°. At Sierra Leone, Messrs. Watt and Winterbottom, e saw it frequently at 100°, and even as high as 102° and 103°, at some distance from the coast. Adamson saw it at Senegal as high as 1081°. Brydone affirms, that when the sirocco blows in Sicily the heat rises to 112°. Dr. Chalmers observed a heat

<sup>&</sup>lt;sup>a</sup> Sir E. Home's Lect. &c. iii. 438.

<sup>&</sup>lt;sup>b</sup> See, also, Dr. W. B. Carpenter, art. Life, in Cyclop. of Anat. and Physiol. Sept. 1840.

c "Observatio docet nullum animal quod pulmones habent, posse in aere vivere, cujus eadem est temperies cum suo sanguine." Element. Chemiæ, i. 275, Lug. Bat. 1732; and Haller, Element. Physiologiæ, tom. ii.

<sup>&</sup>lt;sup>4</sup> Meteorological Register, for the years 1822, 1823, 1824, and 1825, from observations made by the surgeons at the military posts of the United States. See, also, a similar register for the years 1826, 1827, 1828, 1829, and 1830, Philad. 1840.

<sup>·</sup> Account of the Native Africans, vol. i., p. 32 and 33.

Lawrence's Lectures on Physiology, &c., p. 306, London, 1819.

of 115° in South Carolina; Humboldtb of 110° to 115° in the Llanos or Plains near the Orinoco; and Captain Tuckey asserts, that on the Red Sea he never observed the thermometer at midnight under 94°; at sunrise under 104°; or at mid-day under 112°. In British India it is asserted to have been seen as high as 130°.

So long ago as 1758, Governer Ellisd of Georgia had noticed how little the heat of the body is influenced by the external atmosphere. "I have frequently," he remarks, "walked an hundred vards under an umbrella with a thermometer suspended from it by a thread, to the height of my nostrils, when the mercury has rose to 105°, which is prodigious. At the same time I have confined this instrument close to the hottest part of my body, and have been astonished to observe, that it has subsided several degrees. Indeed I never could raise the mercury above 97° with the heat of my body." Two years after the date of this communication, the power of resisting a much higher atmospheric temperature, was discovered by accident. MM. Duhamel and Tillet, in some experiments for destroying an insect, that infested the grain of the neighbourhood, - having occasion to use a large public oven, on the same day in which bread had been baked in it, were desirous of ascertaining its temperature. This they endeavoured to accomplish by introducing a thermometer into the oven at the end of a shovel. On being withdrawn, the thermometer indicated a degree of heat considerably above that of boiling water; but M. Tillet, feeling satisfied that the thermometer had fallen several degrees in approaching the mouth of the oven, and seeming to be at a loss how to rectify the error, a girl, - one of the servants of the baker, and an attendant on the oven, - offered to enter and mark with a pencil the height at which the thermometer stood within the oven. The girl smiled at M. Tillet's hesitation at her proposition, entered the oven, and noted the temperature to be 260° of Fahrenheit. M. Tillet, anxious for her safety, called upon her to come out; but she assured him she felt no inconvenience from her situation, and remained ten minutes longer, when the thermometer had risen to 280° and upwards. She then came out of the oven, with her face considerably flushed, but her respiration by no means quick or laborious.

These facts excited considerable interest, but no farther experiments appear to have been instituted, until, in the year 1774, Dr. Geo. Fordyce, and Sir Charles Blagden made their celebrated trials with heated air. The rooms, in which these were made,

b Tableau Physique des Regions Equatoriales.

Philosophical Transactions for 1775, p. 111.

Account of the Weather and Diseases of South Carolina, London, 1776.

e Prof. Jameson, in British India, Amer. edit. iii. 170, New York, 1832. For an account of the highest temperature, observed in different climates, see the author's "Elements of Hygiène," p. 96; art. Atmosphere, by the author, in the Cyclopædia of Practical Medicine, Philadelphia, 1836; Annuaire du Bureau des Longitudes, 1825; and Elémens de Physique, par Pouillet, iv. 637, 2de édit., Paris, 1832.

<sup>Philosophical Transactions, 1758, p. 755.
Mémoir. de l'Académie des Sciences de Paris, 1762, p. 186.</sup> 

were heated by flues in the floor. Having taken off his coat, waistcoat, and shirt, and being provided with wooden shoes, tied on with list, Dr. Fordyce went into one of the rooms, as soon as the thermometer indicated a degree of heat above that of boiling water. The first impression of this heated air upon his body was exceedingly disagreeable; but in a few minutes all uneasiness was removed by copious perspiration. At the end of twelve minutes he left the room very much fatigued, but not otherwise disordered. The thermometer had risen to 220°. In other experiments, it was found, that a heat even of 260° could be borne with tolerable ease. At this temperature, every piece of metal was intolerably hot; small quantities of water, in metallic vessels, quickly boiled; and streams of moisture poured down over the whole surface of his body. That this was merely the vapour of the room, condensed by the cooler skin, was proved by the fact, that when a Florence flask, filled with water of the same temperature as the body, was placed in the room, the vapour condensed in like manner upon its surface, and ran down in streams. Whenever the thermometer was breathed upon, the mercury sank several degrees. Every expiration - especially if made with any degree of violence - communicated a pleasant impression of coolness to the nostrils, scorched immediately before by the hot air rushing against them when they inspired. In the same manner, their comparatively cool breath cooled their fingers, whenever it reached them. "To prove," says Sir Charles Blagden, "that there was no fallacy in the degree of heat shown by the thermometer, but that the air which we breathed, was capable of producing all the well-known effects of such an heat on inanimate matter, we put some eggs and beef-steak upon a tin frame, placed near the standard thermometer, and farther distant from the cockle than from the wall of the room. In about twenty minutes, the eggs were taken out roasted quite hard; and in forty-seven minutes, the steak was not only dressed, but almost Another beef-steak was rather overdone in thirty-three minutes. In the evening, when the heat was still greater, we laid a third beef-steak in the same place; and as it had now been observed, that the effect of the heated air was much increased by putting it in motion, we blew upon the steak with a pair of bellows, which produced a visible change on its surface, and seemed to hasten the dressing; the greatest part of it was found pretty well done in thirteen minutes." In all these experiments, and similar ones were made in the following year, by Dobson,a of Liverpool, the heat of the body, in air of a high temperature, speedily reached 100°; but exposure to 212°, and more, did not carry it higher. These results are not, however, exactly in accordance with those of MM. Berger and Delaroche, b deduced

a Philosophical Transactions for 1775, p. 463.

b Expér. sur les Effects qu'une forte Chaleur produit sur l'Economie, Paris, 1805; and Journal de Physique, lxiii. 207, lxxi. 289, and lxxvii. 1.

from experiments performed in 1806. Having exposed themselves, for some time, to a stove, - the temperature of which was 39° of Réaumur, or 120° of Fahrenheit, - their temperature was raised 3° of Réaumur, or 63° of Fahrenheit; and M. Delaroche found, that his rose to 4° of Réaumur, or 9° of Fahrenheit, when he had remained sixteen minutes in a stove heated to 176° of Fahrenheit. According to Sir David Brewster,a - the distinguished sculptor, Chantry, exposed himself to a temperature yet higher. The furnace which he employs for drying his moulds, is about 14 feet long, 12 feet high, and 12 feet broad. When raised to its highest temperature, with the doors closed, the thermometer stands at 350°, and the iron floor is red-hot. The workmen often enter it at a temperature of 340°, walking over the iron floor with wooden clogs, which are, of course, charred on the surface. On one occasion, Mr. Chantry, accompanied by five or six of his friends, entered the furnace, and after remaining two minutes, they brought out a thermometer, which stood at 320°. Some of the party experienced sharp pains in the tips of their ears, and in the septum of the nose, whilst others felt a pain in the eyes. In some experiments of Chabert, who exhibited his powers as a "Fire King," in this country as well as in Europe, he is said to have entered an oven with impunity, the heat of which was from 400° to 600° of Fahrenheit. Experiments have shown, that the same power of resisting excessive heat is possessed by other animals. Drs. Fordyce and Blagden shut up a dog in a room, the temperature of which was between 220° and 236°, for half an hour; at the end of this time a thermometer was applied between the thigh and flank of the animal; and in about a minute the mercury sank to 110°; but the real heat of the body was certainly less than this, as the ball of the thermometer could not be kept a sufficient time in proper contact; and the hair, which felt sensibly hotter than the bare skin, could not be prevented from touching the instrument. The temperature of this animal, in the natural state, is 101°.

We find in the case of aquatic animals, astonishing cases of adaptation to the medium in which they live. Although man is capable of breathing air, heated to above the boiling point of water with impunity, we have seen, that he cannot bear the contact of water much below that temperature. Yet we find certain of the lower animals—as fishes—living in water at a temperature, which would be entirely sufficient to boil them if dead. In the thermal springs of Bahia, in Brazil, many small fishes are seen swimming in a rivulet, which raises the thermometer to 88°, when the temperature of the air is only 77½°. Sonnerat, again, found fishes existing in a hot spring at the Manillas, at 158° Fah.; and MM. Humboldt and Bonpland, in travelling through the province of Quito in South America, perceived them thrown up alive, and apparently in health, from the bottom of a volcano, in the course of its

<sup>\*</sup> Letters on Natural Magic, p. 281, Amer. Edit., New York, 1832.

explosions, along with water and heated vapour, which raised the thermometer to 210°, or only two degrees short of the boiling point. Dr. Reeve found living larvæ in a spring, whose temperature was 208°; Lord Bute saw confervæ and beetles in the boiling springs of Albano, which died when plunged into cold water; and Dr. Elliotson knew a gentleman, who boiled some honeycomb, two years old, and, after extracting all the sweet matter, threw the refuse into a stable, which was soon filled with bees.

When the heating influence is applied to a part of the body only, as to the urethra, the temperature of the part, it has been affirmed, is not increased beyond the degree to which the whole body may be raised.

From all these facts, then, we may conclude, that when the body is exposed to a temperature, greatly above the ordinary standard of the animal, a frigorific influence is exerted; but this is effected at a great expense of the vital energy; and hence is followed by considerable exhaustion, if the effort be prolonged. In the cold-blooded animal, the power of resisting heat is not great; so that it expires in water not hotter than the human blood occasionally is. Dr. Edwards found that a frog, which can live eight hours in water at 32°, is destroyed in a few seconds in water at 105°: this appears to be the highest temperature that cold-blooded animals can bear.

Observation has shown, that although the average temperature of an animal is such as we have stated in the table, particular circumstances may give occasion to some fluctuation. A slight difference exists, according to sex, temperament, idiosyncrasy, &c. MM. Edwards and Gentil found the temperature of a young female half a degree less than that of two boys of the same age. Edwardse tried the temperature of twenty sexagenarians, thirty-seven septuagenarians, fifteen octogenarians, and five centenarians, at the large establishment of Bicêtre, and he observed a slight difference in each class. John Davyd found, that the temperature of a lamb was a degree higher than that of its mother; in five new-born children, the heat was about half a degree higher than that of the mother, and it rose half a degree higher in the first twelve hours after birth. Dr. Holland, too, found that the mean temperature of forty infants exceeded that of the same number of adults by 13°: twelve of the children possessed a temperature of 100° to 1034°. Edwards, on the other hand, found, that, in the warm-blooded animal, the faculty. of producing heat is less, the nearer to birth; and that in many cases, as soon as the young dropped from the mother, the temperature fell to within a degree or two to that of the circumambient air;

Animal Physiology, Library of Useful Knowledge, p. 3.

b Physiology, p. 247, Lond. 1840; also, Hodgkin and Fisher's Appendix to their translation of Edwards, Sur l'Influence des Agens Physiques, &c., p. 467; and Carpenter, Principles of General and Comparative Physiology, p. 154, Lond. 1839.

De l'Influence des Agens, &c., p. 436, Paris, 1826.

<sup>4</sup> Philosoph. Transact. p. 602, for 1814.

e An Inquiry into the Laws of Life, &c., Edinb. 1829; and Despretz, in Edinb. Journal of Science, &c., iv. 185.

and he moreover affirms, that the faculty of producing heat is at its minimum at birth, and that it increases successively to the adult age. His trials on children, at the large Hôpital des Enfans of Paris, and on the aged at Bicêtre, showed that the temperature of infants, one or two days old, was from 93° to 95° of Fahrenheit; of the sexagenarian from 95° to 97°; of the octogenarian 94° or 95°; and that, as a general rule, it varied according to the age. In his experiments connected with this subject, he discovered a striking analogy between warm-blooded animals in general. Some of these are born with the eyes closed; others with them open: the former, until the eyes are opened, he found to resemble the cold-blooded animal; the latter - or those born with the eyes open - the warmblooded. Thus, he remarks, the state of the eyes, although having no immediate connexion with the production of heat, may yet coincide with an internal structure influencing that function, and it certainly furnishes signs, which indicate a remarkable change in this respect; for, at the period of the opening of their eyes, all young mammalia have nearly the same temperature as adults. Now, in accordance with analogy, a new-born infant, at the full period, having its eyes open, should have the power of maintaining a pretty uniform temperature during the warm seasons; but, if birth should take place at the fifth or sixth month, the case is altered; the pupil is generally covered with the membrana pupillaris, which places the young being in a condition similar to that of closure of the eyelids an animals. Analogy, then, would induce us to conclude, that, in such an infant, the power of producing heat should be inconsiderable, and observation confirms the conclusion; although we obviously have not the same facilities, as in the case of animals, of exposing the infant to a depressed temperature. The temperature of a seven months' child, though well swathed, and near a good fire, was, within two or three hours after birth, no more than 89.60 Fahrenheit. Before the period at which this infant was born, the membrana pupillaris disappears; and it is probable, as Dr. Edwards has suggested, if it had been born prior to the disappearance of the membrane, its power of producing heat might have been so feeble, that it would scarcely have differed from that of mammalia born with their eyes closed.a

The state of the system, as to health or disease, also influences the evolution of heat. Dr. Francis Home, of Edinburgh, took the heat of various patients at different periods of their indispositions. He found that of two persons, labouring under the cold stage of an intermittent, to be 104°; whilst, during the sweat and afterwards, it fell to 101°, and to 99°. The highest degree, which he noticed in fever, was 107°. We have often witnessed the thermometer at 106° in scarlatina and in typhus, but it probably rarely

<sup>\*</sup> Op. cit., and Analytical Review of Hodgkin and Fisher's translation, by the author of this work, in Amer. Journ. of Medical Sciences, p. 150, for May, 1834.

b Medical Facts and Experim., Lond. 1759.

<sup>&</sup>lt;sup>c</sup> Currie's Medical Reports, Liverpool, 1798, and Philad. 1808.

exceeds this, although it is stated to have been seen as high as 1120," and this is the point designated as "fever heat," on Fahrenheit's scale. M. Edwards alludes to a case of tetanus, in a child, the particulars of which were communicated to him by M. Prévost of Geneva, in which the temperature rose to 110.75° Fahrenheit.b Huntere found the interior of a hydrocele, on the day of operation, to raise the mercury to 92°; on the following day, when inflammation had commenced, it rose to 99°. The fluid, obtained from the abdomen of an individual, tapped for the seventh time for dropsy of the lower belly, indicated a temperature of 101°. Twelve days thereafter, when the operation was repeated for the eighth time, the temperature was 104°. Dr. Granvilled has asserted that the temperature of the uterine system sometimes rises as high as 120° — the elevation seeming to bear some ratio to the degree of action in the organ. We have frequently been struck with the seemingly elevated temperature of the vagina under these circumstances, but cannot help suspecting some inaccuracy in the observations of Dr. Granville, the temperature, he indicates, being so much higher than has ever been noticed in any condition of the system. Under this feeling, several experiments were made, at the author's request, by Dr. Barnes, one of the resident physicians of the Philadelphia Hospital, which exhibit only a slight difference between the temperature of the vagina and that of the uterus during parturition. In two cases, the temperature of the labia was 100°, and in a third 105°; whilst that of the uterus was 100°, 102°, and 106° respectively. Dr. James Currie had himself bled; and during the operation, the mercury of a thermometer, which he held in his hand, sank, at first slowly and afterwards rapidly, nearly 10°; and when he fainted, the assistant found, that it had sunk 8° farther.

MM. Edwards and Gentil assert, that they have likewise observed diurnal variations in the temperature of individuals, and these produced, apparently, by the particular succession in the exercise of the different organs; as where intellectual meditation was followed by digestion. These variations, they affirm, frequently amounted to two or three degrees, between morning and evening.

M. Chevalliers has investigated the temperature of the urine on issuing from the bladder. This he did not find to vary from external temperature, but it was effected by rest, fatigue, change of regimen, remaining in bed, &c. The lowest temperature, which was observed on rising in the morning, was about 92°; the highest,

<sup>&</sup>lt;sup>a</sup> G. T. Morgan's First Principles of Surgery, p. 80, Lond. 1837.

b Edwards, op. citat. p. 490. c On the Blood, &c. p. 296, Lond. 1794. d Philos. Transact. p. 262, for 1825; and Sir E. Home, in Lect. on Comp. Anat. v. 201, Lond. 1828.

<sup>·</sup> Dunglison's American Medical Intelligencer, Feb. 15, 1839, p. 346.

See Purkinje, in art. Calor Animalis, in Encyclop. Wörterbuch der Medicin. Wissenschaft. Band. vi. s. 530, Berlin, 1831.

s Essai sur la Dissolution de la Gravelle, &c. p. 120, Paris, 1837.

after dinner, and when fatigued, 99°. In the case of another person, the temperature of the urine was never lower than 101°, and

occasionally upwards of 102°, when he was fatigued.

Such are the prominent facts connected with the subject of animal heat. It is obvious, that it is altogether disengaged by an action of the system, which enables it to counteract, within certain limits, the extremes of atmospheric heat and cold. The animal body, like all other substances, is subjected to the laws regarding the equilibrium, the conduction, and the radiation of caloric; but, by virtue of the important function we are now considering, its own temperature is neither elevated nor depressed by those influences to any great amount. Into the seat and nature of this mysterious process, and the various ingenious theories that have

been indulged, we shall now inquire.

Physiologists have been by no means agreed, regarding the organs or apparatus of calorification. Some, indeed, have affirmed that there is not, strictly speaking, any such apparatus; and that animal heat is a result of all the other vital operations. Amongst those, too, who admit the existence of such an apparatus, a difference of sentiment prevails; some thinking, that it is local, or effected in a particular part of the body; others, that it is general, or disseminated through the whole economy. Under the name caloricité, Chaussier admits a primary vital property, by virtue of which living beings disengage the caloric on which their proper temperature is dependent, in the same manner as they accomplish their other vital operations, by other vital properties; and in support of this doctrine, he adduces the circumstance, that each living body has its own proper temperature; which is coexistent only with the living state; is common to every living part; ceases at death; and augments by every cause, that excites the vital activity. It has been properly objected, however, to this view, that the same arguments would apply equally to many other vital operations, - and that it would be obviously improper to admit, for each of these functions, a special vital principle. The notion has not experienced favour from the physiologist, and is, we believe, confined to the individual from whom it emanated.\*

Amongst those who admit that calorification is a local action, some have believed, that the caloric is disengaged in a particular organ, whence it is distributed to every part of the body; whilst others conceive, that every part disengages its own caloric and has

its special temperature.

So striking a phenomenon as animal temperature could not fail to attract early attention; and accordingly, we find amongst the ancients various speculations on the subject. The most prevalent was, — that its seat is in the heart; that the heat is communicated to the blood in that viscus, and is afterwards sent to every part of the system; and that the great use of respiration is to cool the

<sup>&</sup>lt;sup>a</sup> Art. Caloricité, by Coutanceau, in Dict. de Médecine, tom. iv. Paris, 1822; and Adelon's Physiologie de l'Homme, tom. iii. 407, 2de édit., Paris, 1829.

heart; but this hypothesis is liable to all the objections, which apply to the notion of any organ of the body acting as a furnace. - that such organ ought to be calcined; and it has the additional objection, applicable to all speculations, regarding the ebullition and effervescence of the blood as a cause of heat, that it is purely conjectural, without the slightest fact or argument in its favour. It was not, indeed, until the chemical doctrines prevailed, that any thing like argument was adduced in support of the local disengagement of heat: the opinions of physiologists then settled almost universally upon the lungs; and this, chiefly, in consequence of the observation, that animals, which do not breathe, have a temperature but little superior to the medium in which they live; whilst man and animals that breathe have a temperature considerably higher than the medium heat of the climate in which they exist, and one which is but little affected by changes in the thermal condition of that medium; and, moreover, that birds. which breathe, in proportion, a greater quantity of air than man, have a still higher temperature than he. Mayow, whose theory of animal heat was, in other respects, sufficiently unmeaning, affirmed, that the effect of respiration is not to cool the blood, as had been previously maintained, but to generate heat, which it did by an operation analogous to combustion. It was not, however, until the promulgation of Dr. Black's doctrine of latent heat, that any plausible explanation of the phenomenon appeared. According to that distinguished philosopher, a part of the latent heat of the inspired air becomes sensible; consequently, the temperature of the lungs, and of the blood passing through them, must be elevated; and, as the blood is distributed to the whole system, it communicates its heat to the parts as it proceeds on its course. But this view was liable to an obvious objection, which was, indeed, fatal to it, and so Black himself appears to have thought, from his silence on the subject. If the whole of the caloric were disengaged in the lungs, as in a furnace, and were distributed through the bloodvessels, as heated air is transmitted along conducting pipes, the temperature of the lungs ought to be much greater than that of the parts more distant from the heart; so great indeed, as to consume that important organ in a short space of time.

The doctrine, maintained by Lavoisiere and Séguin, was;—
that the oxygen of the inspired air combines with the carbon and
hydrogen of the venous blood, and produces combustion. The
caloric given off is then taken up by the bloodvessels, and is distributed over the body. The arguments, which they adduced in
favour of this view, were:— the great resemblance between
respiration and combustion, so that if the latter gives off heat, the
former ought to do so likewise;— the fact that arterial blood is

<sup>\*</sup> Tract. quinque, Oxon. 1674.

b Bostock's Physiology, 3d edit. p. 440, Lond. 1836.

Mém. de l'Acad. des Sciences pour 1777, 1780, and 1790.

somewhat warmer than venous; - and certain experiments of Lavoisier and La Place, which consisted in placing animals in the calorimeter, and comparing the quantity of ice which they melted, and, consequently, the quantity of heat, which they gave off, with the quantity of carbonic acid produced; and finding, that the quantity of caloric, which would result from the carbonic acid formed, was exactly that disengaged by those animals. Independently, however, of other objections, this hypothesis is liable to those already urged against that of Black, which it closely resembles. The objection, that the lungs ought to be much hotter than they really are - both absolutely and relatively-was attempted to be obviated by Dr. Crawfordb in a most ingenious and apparently logical manner. The oxygen of the inspired air, according to him, combines with the carbon given out by the blood, so as to form carbonic acid. But the specific heat of this is less than that of oxygen: and accordingly, a quantity of latent caloric is set free; and this caloric is not only sufficient to support the temperature of the body, but also to carry off the water - which was supposed to be formed by the union of the hydrogen and the oxygen - in the state of vapour, and to raise the temperature of the inspired air considerably. So far the theory of Crawford was liable to the same objections as those of Black, and Lavoisier and Séguin. He affirmed. however, that the same process by which the oxygen of the inspired air is converted into carbonic acid, converts likewise the venous into arterial blood; and as he assumed from his experiments, that the capacity for caloric of arterial blood is greater than that of venous, in the proportion of 1.0300 to 0.8928; he conceived, that the caloric, set free in the formation of the carbonic acid, in place of raising the temperature of the arterial blood, is employed in saturating its increased capacity, and in maintaining its temperature at the same degree with the venous. According to this view, therefore, the heat is not absolutely set free in the lungs, although arterial blood contains a greater quantity of caloric than venous; but when, in the capillaries, the arterial becomes converted into venous blood, or into blood of a less capacity for caloric, the heat is disengaged, and occasions the temperature of the body.

If the facts, which served as a foundation for this beautiful theory of animal heat, were not false, the deductions would be irresistible; and, accordingly, it was at one time almost universally received, especially by those who consider that all vital operations can be assimilated to chemical processes; and it is still favoured by many. "The animal heat," observes a recent writer, "has been accounted for in different ways by several ingenious physiologists; from the aggregate of their opinions and experiments, I deduce, that heat is extricated all over the frame, in the capillaries,

Dr. Billing, First Principles of Medicine, 2d edit., p. 19, Lond. 1837.

Memoir, de l'Acad. des Sciences pour 1780.

b Exp. &c. on Animal Heat, 2d edit. Lond. 1788; and Fleming's Philosophy of . Zoology, i. 387, Edinb. 1822.

by the action of the nerves, during the change of the blood, from scarlet arterial to purple venous; and also whilst it is changing in the lungs from purple to scarlet. There is a perpetual deposition by the capillary system of new matter, and decomposition of the old all over the frame, influenced by the nerves; in this decomposition there is a continual disengagement of carbon, which mixes with the blood returning to the heart, at the time it changes from scarlet to purple; this decomposition, being effected by the electric agency of the nerves, produces constant extrication of caloric; again, in the lungs that carbon is thrown off and united with oxygen, during which caloric is again set free, so that we have in the lungs a charcoal fire constantly burning, and in the other parts a wood fire, the one producing carbonic acid gas, the other carbon, the food supplying through the circulation the vegetable (or what answers the same end, animal) fuel, from which the charcoal is prepared which is burned in the lung." Numerous objections have, however, been made against the view of Crawford. In the first place, it was objected, that our knowledge is limited to the fact, that oxygen is taken into the pulmonary vessels, and carbonic acid given off, but that we had no means of knowing whether the one goes immediately to the formation of the other. Dr. Crawford had inferred from his experiments, that the specific heat of oxygen is 4.7490; of carbonic acid, 1.0454; of azote, 0.7936; and of atmospheric air, 1.7900; but the more recent experiments of Delaroche and Bérard make that of oxygen, 0.2361; of carbonic acid, 0.2210; of azote, 0.2754; and of atmospheric air, 0.2669; a difference of such trifling amount, that it has been conceived the quantity of caloric, given out by oxygen during its conversion into carbonic acid, would be insufficient to heat the residual air, which is expelled in breathing, to its ordinary elevation. Secondly. The elevation of temperature of one or two degrees, which appears to take place in the conversion of venous into arterial blood, although generally believed, is not assented to by all. The experiments instituted on this point have been few and imprecise; and the most recent by MM. Becquerel and Breschet, made by introducing delicate thermometers into the auricles of the heart of dogs, invariably gave the temperature of arterial blood only a few fractions of a degree higher than that of venous blood. Thirdly. M. Dulong, - on repeating the experiments of Lavoisier and La Place. for comparing the quantities of caloric, given off by animals, in the calorimeter, with that which would result from the carbonic acid, formed during the same time in their respiration - did not attain a like result. The quantity of caloric disengaged by the animal was always superior to that which would result from the carbonic acid formed. Fourthly. The estimate of Crawford, regarding the specific heat of venous and arterial blood, has been contested. He made that of the former, we have seen, 0.8928; of the latter,

<sup>&</sup>lt;sup>a</sup> See, also, Elliotson, Human Physiology, p. 238, Lond. 1840.

b Comptes Rendus, Oct. 1841. Magendie's Journal de Physiologie, iii. 45.

1.0300. The result of the experiments of Dr. John Davy<sup>a</sup> give 0.903 to the former, and 0.913 to the latter; and in another case, the result of which has been adopted by Magendie, the specific heat of the venous was greater than that of the arterial blood, in the proportion of .852 to .839. Granting, however, that the case is as stated by Crawford, it is insufficient to explain the phenomena. It has, indeed, been attempted to show, that if the whole of the caloric, set free in the manner mentioned, were immediately absorbed, it would be insufficient for the constitution of the arterial blood; and that, instead of the lung running the risk of being calcined, it would

be threatened with congelation.

Of late, the theory of combustion has received the powerful support of Liebig, and many able elucidations and expansions from that able chemist. According to him, the carbon and hydrogen of the food, in being converted, through the agency of oxygen, into carbonic acid and water, must give out as much heat as if these gases were burned in the open air. The temperature of the human body is essentially the same in the torrid as in the frigid zone; but as the body may be regarded in the light of a heated vessel, which cools with the greater rapidity the colder the surrounding medium, the fuel, necessary to maintain its heat, must vary, he maintains, in different climates. How unequal must be the loss of heat at Palermo, where the external temperature is nearly equal to that of the body, and in the polar regions, where the external temperature is from 70° to 90° lower. In the animal body, the food is the fuel, and with a proper supply of oxygen we obtain the heat during its oxidation or combustion. In winter, when we take exercise in a cold atmosphere, and when, consequently, the amount of inspired oxygen increases, the necessity for food containing carbon and hydrogen increases in the same ratio, and, by gratifying the appetite thus excited, we obtain the most efficient protection against piercing cold. A starving man is soon frozen to death; and every one, according to Liebig, knows, that the animals of prey in the arctic regions far exceed, in voracity, those of the torrid zone. Our clothing is merely an equivalent for a certain amount of food. Were we to go naked, like certain savage tribes, or if, in hunting or fishing, we were exposed to the same degree of cold as the Samoyedes, we should be able with ease to consume sixteen pounds of flesh, and perhaps a dozen tallow candles into the bargain, as warmly clad travellers have related with astonishment of these people. We should then, also, be able to take the same quantity of brandy or train-oil without bad effects, because the carbon and hydrogen of these substances would only suffice to keep up the equilibrium between the external temperature and that of our bodies. The whole process of respiration is clearly exhibited when we view the condition of man or animals under abstinence from all food. Oxygen is abstracted from the air, and carbonic acid and water expired, because the number of respira-\* Philos. Transactions for 1814.

tions remains unaltered. With the continuance of the abstinence, the carbon and hydrogen of the body are seen diminishing. The first effect of abstinence is the disappearance of the fat, which can be detected neither in the scanty fæces nor urine; its carbon and hydrogen are thrown off by the skin and lungs, in the form of a compound with oxygen. These constituents, then, have served for the purposes of respiration. Every day, 321 ounces of oxygen are inspired; and these must remove their equivalents of carbon to form carbonic acid. When this combination ceases to go on, respiration terminates: death has taken place. The time required for starving an animal to death depends on its fatness, on the state of its activity, on the temperature of the air, and on the presence or absence of water. That the quantity of heat evolved by the combination of 13.9 ounces of carbon is amply sufficient to account for the temperature of the human body may be estimated by figures; an ounce of carbon, according to the experiments of Despretz, burned, would evolve 14067 degrees of heat; and 13.9 oz. would therefore give out 195531.3 degrees of heat. This would suffice to boil 67.9 pounds of water at 32°, or to convert 11.4 pounds of water at 98.3° into vapour. If we consider, then, the quantity of water vaporized through the skin to be, in twentyfour hours, 48 ounces or 3 pounds, there will remain, after deducting the necessary amount of heat, 144137.7 degrees of heat, which are dissipated by radiation, by heating the expired air, and by the excrementitious matters.a

These views of Liebig have necessarily attracted the attention of the chemical physiologist, and have met with unqualified support from some, whilst they have been as much condemned by others, who appear to have a horror at the introduction of chemical explanations to account for vital phenomena. Yet it cannot be contested, that the function of calorification is an act of vital chemistry, and, therefore, although the views of Liebig may fail to convince, they certainly have taken the proper direction, and, all must admit, have been plausibly and ably supported. It has been objected, that if even the theory of Liebig be allowed to be applicable to mammalia, birds and reptiles, it is by no means so to animals that respire by means of branchiæ or gills, all of which consume but little oxygen, comparatively speaking, yet many of them devour very great quantities of food. Even the largest and most voracious of the reptiles, as the alligators, crocodiles, &c., which consume enormous quantities of food, under a burning climate too, breathe feebly with their vesicular lungs, and consume but little oxygen. Fishes, too, whose blood is but imperfectly oxygenized by their branchial apparatus, are perhaps amongst the most voracious of animals; yet, according to the above theory, they ought to eat little, because they consume little oxygen. These and other facts were early urged by M. Virey,c

Animal Chemistry, Amer. Edit. by Webster, p. 33, Cambridge, 1842.

b See, on Liebig's views, Mr. Ancell, London Lancet, 1842-3.

<sup>·</sup> Journal de Pharmacie, Mai, 1842.

as objections to the views of the Professor of Giessen. It may be replied, however, that in such cases, a large portion of the carbon must pass off in the excrements. There is no country in the world, according to Madame Calderon de la Barco, where so much animal animal food is consumed as in Mexico, and there is no country in which so little is required. To this and to want of exercise she ascribes the early fading of beauty in the higher classes, the decay of teeth, and the over-corpulency so common amongst them; and in regard to the last she is, doubtless, correct.

To the statement of Liebig respecting the greater voraciousness of the animals of prey of the arctic regions, it has been urged, b that a Bengal tiger or cape hyæna requires, in proportion to its size, quite as much aliment as any of the arctic carnivora, and that the vultures of Hindostan and Persia exceed, perhaps, all other animals in gluttony. The voraciousness of the shark, too, even within the tropics, is proverbial. "Those who ride over the Pampas in South America," says Dr. Graves, "at the rate of one hundred miles a day, exposed to a burning sun, subsist entirely on boiled beef and water, without a particle of vegetable food of any kind, and yet they attain to an extraordinary condition, and capability of enduring violent and long-continued exertion. Liebig's theory must be very ductile, if it can explain how it happens, that an exclusively animal diet agrees with man quite as well at the equator as within the arctic circle." Numerous facts, indeed, can be brought forward of an opposite tendency to those of Liebig, and which render it impracticable for us, in the present state of our knowledge, to embrace all his positions. Under Respiration, the theory, supported by him, that the blood corpuscles are the carriers of oxygen from the lungs to the tissues; and the conveyers of carbonic acid back from the tissues to the lungs, was mentioned. It would seem probable, therefore, that if the amount of blood corpuscles should become diminished from any cause, the function of calorification ought to be impaired to the like extent. To discover what effect was produced on the temperature of the living body by diminution in the quantity of blood corpuscles, Andral recently performed some experiments, which showed that the temperature remained normal, even in cases in which the corpuscles had experienced the greatest diminution in number. In the axilla, the temperature was 98° or 99° of Fahrenheit in persons, the proportion of whose blood corpuscles was not higher than 50, 40, 30, and even 21 parts in the 1000; the healthy ratio being 127. Indeed, notwithstanding the great depression in anæmic patients, the heat rose as usual, when they were attacked with fever, to which they are as subject as other individuals.c

But the theories of combustion have been most seriously assailed

Life in Mexico, vol. i., p. 152, Boston, 1842.

Dr. R. J. Graves, A System of Clinical Medicine, p. 57, Dublin, 1843.
 Andral, Hématologie Pathologique, p. 60, Paris, 1843.

by other experiments, tending to show, that the function of calorification is derived from the great nervous centres. When an animal is decapitated, or when the spinal marrow, or the brain, or both, are destroyed, the action of the heart may still be kept up, provided the lungs be artificially inflated. In such case, it is found, that the usual change in the blood, from venous to arterial, is produced; and that oxygen is absorbed and carbonic acid exhaled as usual. Sir Benjamin Brodie, in performing this experiment, directed his attention to the point, - whether animal heat, under such circumstances, be evolved, and the temperature maintained, as where the brain and spinal marrow are entire - and he found, that although the blood appeared to undergo its ordinary changes, the generation of animal heat seemed to be suspended; and consequently, if the inspired air happened to be colder than the body, the effect of respiration was to cool the animal; so that an animal, on which artificial respiration was kept up, became sooner cold than one killed at the same time and left undisturbed. The inference, deduced from these experiments, was, that instead of circulation and respiration maintaining the heat, they dissipate it; and that as the heat is diminished by the destruction of the nervous centres, its disengagement must be ascribed to the action of those centres, and particularly to that of the encephalon.

M. Chossatb endeavoured to discover the precise part of the nervous system engaged in calorification; but the results of his experiments were not such as to induce him to refer it exclusively, with Sir B. Brodie, to the encephalon. He divided the brain, anterior to the pons varolii, in a living animal, so that the eighth pair of nerves were uninjured. Respiration, consequently, continued, and inflation of the lungs was unnecessary. Notwithstanding this serious mutilation, the circulation went on; and Chossat observed distinctly, that arterial blood circulated in the arteries. Yet the temperature of the animal gradually sank, from 104° Fahr., - its elevation at the commencement of the experiment, — to 76°, in twelve hours, when the animal died. It seemed manifest to M. Chossat, that, from the time the brain was divided, heat was no longer given off, and the body gradually cooled, as it would have done after death. He, moreover, noticed, that the time, at which the refrigeration occurred most rapidly, was that in which the circulation was most active, - at the commencement of the experiment. In other experiments, M. Chossat paralysed the action of the brain by a violent concussion, and by injecting a strong decoction of opium into the jugular vein, - keeping up respiration at the same time artificially. The results were the same. From these experiments, he drew the conclusion, that the brain has a direct influence over the production of heat. His next experiments were

a Philos. Trans. for 1811 and 1812.

Sur la Chaleur Animale, Paris, 1820; Adelon, op. cit. iii. 416; Coutanceau, in art. Chaleur Animale, Dict. de Méd. iv. 17, Paris, 1822; and P. H. Bérard, art. Chaleur Animale, Dict. de Méd. 2de édit. vii. 206, Paris, 1834.

directed to the discovery of the medium through which the brain acts, - the eighth pair of nerves, or spinal marrow. He divided the eighth pair of nerves in a dog, and kept up artificial respiration. The temperature sank gradually, and, at the expiration of sixty hours, when the animal died, it was reduced to 65° of Fahrenheit. Yet death did not occur from asphyxia or suspension of the phenomena of respiration, as the lungs crepitated, exhibited no signs of infiltration, and were partly filled with arterial blood. The animal appeared to M. Chossat to expire from cold. As, however, the mean depression of heat was less than in the preceding experiments, he inferred that a slight degree of heat is still disengaged after the section of the eighth pair, whilst, after injury done to the brain directly, heat is no longer given off. Again, he divided the spinal marrow beneath the occiput, and although artificial respiration was maintained, as in the experiments of Brodie, the temperature gradually fell, and the animal died ten hours afterwards, its heat being 79°; and as death occurred in this case so much more speedily than in the last, he inferred, that the influence of the brain over the production of heat is transmitted rather by the spinal marrow than by the eighth pair of nerves. In his farther experiments, Chossat found, that when the spinal marrow was divided between each of the twelve dorsal vertebræ, the depression of temperature occurred less and less rapidly, the lower the intervertebral section; and at the lowest it was imperceptible; he, therefore, concluded, that the spinal marrow does not act directly in the function, but indirectly through the trisplanchuic nerve. To satisfy himself on this point, he opened a living animal on the left side, beneath the twelfth rib, and removed the suprarenal capsule of that side, dividing the trisplanchnic where it joins the semilunar plexus. The animal gradually lost its heat, and died ten hours afterwards in the same state, as regarded temperature, as when the spinal marrow was divided beneath the occiput.

Desiring to obtain more satisfactory results, - the last experiment applying to only one of the trisplanchnic nerves, - he tied the aorta, which supplies both with the materials on which they operate, beneath the place where it passes through the arch of the diaphragm, at the same time preventing asphyxia by inflating the lungs. The animal lost its heat much more rapidly, and died in five hours. In all these cases, according to Chossat, death occurred from cold; the function, by which the caloric, constantly abstracted from the system by the surrounding medium, is generated, having been rendered impracticable. To obtain a medium of comparison, he killed several animals by protracted immersion in cold water, and found, that the lowest temperature, to which the warm-blooded could be reduced, and life persist was 79° of Fahrenheit. M. Chossat also alludes to cases of natural death by congelation, which he conceives to destroy in the manner before suggested, by impairing the nervous energy, as indicated by progresLastly: — on killing animals suddenly, and attending to the progress of refrigeration after death, he found it to be identical with that which follows direct injury of the brain, or division of the

spinal marrow beneath the occiput.

A view somewhat analogous to this of M. Chossat, has been embraced by Sir Everard Home. He conceives, that the phenomenon is restricted to the ganglionic part of the nervous system, and he rests the opinion chiefly upon the position, that there are certain animals, which have a brain, or some part equivalent to one, but whose temperature is not higher than that of the surrounding medium; whilst, on the other hand, all the animals that

evolve heat are provided with ganglia.

The doctrines of Brodie, Chossat, and Home have been considered by the generality of the chemists, - by Brande, Thomson, c and Paris, d - to be completely subversive of the chemical doctrines, which refer the production of animal heat to the respiratory function; and their position, — that it is a nervous function, — has seemed to be confirmed by the facts attendant upon injury done to the nerves of parts, and by what is witnessed in paralytic limbs, the heat of which is generally and markedly inferior to that of the sound parts. But there are many difficulties in the way of admitting, that the nervous system is the special organ for the production of animal temperature. Dr. Wilson Philip, from a repetition of the experiments of Sir Benjamin Brodie, was led to conclude, that the cause, why the temperature of the animal body diminished more rapidly, when artificial inflation was practised, than when the animal was left undisturbed, was - too large a quantity of air having been sent into the lungs; and he found, that when a less quantity was used, the cooling process was sensibly retarded by the inflation. The experiments of Legallois, Hastings,g and Williams,h although differing from each other in certain particulars, corroborate the conclusion of Dr. Philip, and, what is singular, they would appear to show, that the temperature occasionally rises during the experiment; facts, which would rather confirm the view that respiration is greatly concerned in the evolution of heat.

Many of the facts, detailed by Chossat, are curious, and exhibit the indirect agency of the nervous system, but his conclusion, that the trisplanchnic is the great organ for its development, is liable to the objections already urged regarding the theory, which looks upon the heart, or the lungs as furnaces for the disengagement of

b Manual of Chemistry, vol. iii. c System of Chemistry, vol. iv.

<sup>d</sup> Medical Chemistry, p. 327, Lond. 1825.

Annales de Chimie, iv. 5, Paris, 1817.

Philos. Trans. p. 257, for 1825; Journal of Science and Arts, xx. 307; and Lect. on Comparative Anat. v. 121 and 194, Lond. 1828.

An Experimental Inquiry into the Laws of the Vital Functions, 3d edit. p. 180.

Wilson Philip, op. cit.; and Journal of Science, &c. xiv. 96.

\* Edinb. Medico-Chirurgical Transact, ii. 192.

caloric, that they ought to be consumed in a short space of time by the operation. All the facts, however, exhibit, that, in the upper classes of animals, the three great acts of innervation, respiration and circulation are indirectly concerned in the function; not that any one is the special apparatus. M. Edwards has attempted to show, that it is more connected with the second of these than with either of the others. Thus, animals, whose temperature is highest, bear privation of air the least; whilst cold-blooded animals suffer comparatively little from it; and young animals are less affected by it than the adult. Now, the greater the temperature of the animal, and the nearer to the adult age, the greater is the consumption of oxygen. He farther observed, that whilst the seasons modify calorification, they affect also respiration; and that if, in summer, less heat be elicited, and in winter more, respiration con-

sumes less oxygen in the former season than in the latter.

The experiments of Legallois, as well as those instituted by Edwards, led the latter to infer, that there is a certain ratio between heat and respiration, in both cold-blooded and warm-blooded animals, and in hibernating animals, both in the periods of torpidity and of full activity. When the eighth pair of nerves is divided in the young of the mammalia, a considerable diminution is produced in the opening of the glottis; so that, in puppies, recently born, or one or two days old, so little air enters the lungs, that when the experiment is made in ordinary circumstances, the animal perishes as quickly as if it were entirely deprived of air. It lives about half an hour. But, if the same operation be performed upon puppies of the same age, benumbed with cold, they will live a whole day. In the first case, M. Edwards thinks, and plausibly, the small quantity of air is insufficient to counteract the effect of the heat; whilst, in the other, it is sufficient to prolong life considerably, and he deduces the following practical inferences applicable to the adult age, and particularly to man. A person, he remarks, is asphyxied by an excessive quantity of carbonic acid in the air he breathes; the pulse is no longer perceptible; the respiratory movements cannot be discerned, but his temperature is still elevated. How should we proceed to recall life. Although the action of the respiratory organs is no longer visible, all communication with the air is not cut off. The air is in contact with the skin, upon which it exerts a vivifying influence; it is also in contact with the lungs, in which it is renewed by the agitation constantly taking place in the atmosphere, and by the heat of the body, which rarefies it. The heart continues to beat, and maintains a certain degree of circulation, although not perceptible by the pulse. The temperature of the body is too high to allow the feeble respiration to produce upon the system all the effect of which it is capable. The temperature must then be reduced: the patient must be withdrawn from the deleterious atmosphere; be stripped of his clothes, in order that the air may have a more extended action upon his skin; be exposed to

the cold, although it be winter, and cold water be thrown upon his face until the respiratory movements reappear. This is precisely the treatment adopted in practice to revive an individual from a state of asphyxia. If, instead of cold, continued warmth were to be applied, it would be one of the most effectual means of extinguishing life. This consequence, like the former, is confirmed by experience. In sudden faintings, where the pulse is weak or imperceptible, the action of the respiratory organs diminished, and sensation and voluntary motion suspended, persons, the most ignorant of medicine, are aware, that means of refrigeration must be employed, - such as exposure to air, ventilation, and sprinkling with cold water. In violent attacks of asthma, also, when the extent of respiration is so limited that the patient experiences a sense of suffocation, he courts the cold air even in the severest weather; opens the windows; breathes a frosty air, and finds himself relieved.

As a general rule, an elevated temperature accelerates the respiratory movements, but the degree of temperature, requisite to produce this effect, is not the same in all. The object of this accelerated respiration is, that more air may come in contact with the lungs, in

a given time, so as to reanimate what the heat depresses.

It is proper to remark, however, that we meet with many exceptions to the rule endeavoured to be laid down by M. Edwards. as regards the constant ratio between heat and respiration. Experiments on the lower animals, and pathological cases in man, have shown, that lesions of the upper part of the spinal marrow give occasion, at times, to an extraordinary development of heat. In the case of a man at St. George's Hospital, London, labouring under a lesion of the cervical vertebræ, Sir B. Brodie observed the temperature to rise to 111°, at a time when the respirations were not more than five or six in a minute. a Drs. Graves and Stokesb give the case of a patient who laboured under a very extensive development of tubercles, had tubercular abscesses in the upper portions of both lungs, and general bronchitis. In this case, at a period when the skin was hotter than usual, and the pulse 126, the respirations were only 14 in the minute; besides, as Dr. Alisone has remarked, the temperature of the body is not raised by voluntarily increasing or quickening the acts of respiration; but by voluntary exertions of other muscles, which accelerate the circulation, and thus necessitate an increased frequency of respiration: a fact, which would seem to show that calorification is dependent not simply on the application of oxygen to the blood, but on the changes that take place during circulation, and to the maintenance of which the oxygenation of the blood is one essential condition.

c Outlines of Physiol., Lond. 1831.

Lond. Med. Gazette, for June, 1836, and G. T. Morgan's First Principles of Surgery, p. 85, Lond. 1837.

Dublin Hospital Reports, vol. v. See, also, Dr. Graves, Clinical Lectures, Dunglison's Amer. Med. Lib. Edit. p. 126, Philad. 1838; and Dr. John Davy, Researches, Physiological and Anatomical, Amer. Med. Lib. Edit. p. 89, Philad. 1840.

Moreover, in the fœtus in utero, there is, of course, no respiration; yet its temperature equals, and indeed is said to even exceed, that of the mother; and we know that its circulation is more rapid, and its nutrition more active.a That innervation is indirectly concerned in the phenomenon is proved by the various facts which have been referred to; and Legallois, although he does not accord with Sir B. Brodie, conceives, that the temperature is greatly under the influence of the nervous system, and that whatever weakens the nervous power, proportionally diminishes the capability of producing heat. Dr. Philip, too, concluded from his experiments, that the nervous influence is so intimately connected with the power of evolving heat, that it must be looked upon as a necessary medium between the different steps of the operation. He found, that if the galvanic influence be applied to fresh drawn arterial blood, an evolution of heat, amounting to three or four degrees, takes place, whilst the blood assumes the venous hue and becomes partly coagulated. He regards the process of calorification as a secretion; and explains it upon his general principle of the identity of the nervous and galvanic influences, and of the necessity for the exercise of such influence in the function of secretion.b

Mr. H. Earle found the temperature of paralysed limbs to be slightly lower than that of sound limbs, and the same effect upon calorification is observed to supervene on traumatic injuries of the nerves. In a case of hemiplegia, of five months' duration, under the author's care at the Blockley Hospital, the thermometer in the right — the sound — axilla of the man stood at 962°; in the axilla of the paralysed side at 96°. The difference in temperature of the hands was more signal. The right hand carried the mercury to 87°, whilst the left raised it no further than 79½°. In another case - that of a female - of two weeks' duration, accompanied with signs of cerebral turgescence, the temperature in the axilla of the sound side, was 100°; in that of the paralysed 98.25°; of the hand of the sound side, 94°; of the other 90°.d It is a general fact, that the temperature of the paralysed side in hemiplegia is less than that of the sound side; yet the irregularity of nervous action is so great, and the power of resistance to excitant or depressing agents so much diminished, that the author has not unfrequently found the temperature to be more elevated.e

Lastly, that the circulation is necessary to calorification, we have evidence in the circumstance that if the vessels, proceeding to a part be tied, animal heat is no longer disengaged from it.

<sup>&</sup>lt;sup>a</sup> See, also, on the connexion of respiration with calorification, P. H. Bérard, art. Chaleur Animale, in Dict. de Méd. 2de édit. vii. 175, Paris, 1834; Dr. Southwood Smith's Philosophy of Health, vol. ii. Lond. 1838; and Mr. Newport on the Temperature of Insects, and its Connexion with the Functions of Respiration and Circulation in this class of invertebrated animals, Philos. Transact. part ii. 4to. p. 77, Lond. 1837.

b Ley, in Appendix to Essay on Laryngismus Stridulus, &c. p. 374, Lond. 1836.

<sup>·</sup> Medico-Chirurgical Transactions, vii. 173, Lond. 1819.

<sup>&</sup>lt;sup>4</sup> See Nasse, Untersuchungen zur Physiologie und Pathologie, Bonn. 1835-6.

<sup>e</sup> See the author in his American Med. Intelligencer, Oct. 15, 1838, p. 225.

It is manifest, then, that in animals, and especially in the warm-blooded, the three great vital operations are necessary for the disengagement of the due temperature, but we have no sufficient evidence of the direct agency of any one, and we see heat elicited in the vegetable, in which these functions are at all events rudimental; and the existence of one of them—innervation—perhaps more than doubtful.<sup>a</sup>

The view of those, who consider, that the disengagement of caloric occurs in the intermediate system or system of nutrition of the whole body, appears to us the most consistent with observed phenomena. These views have varied according to the physical circumstances, that have been looked upon as producing heat. By some, it has been regarded as the product of effervescence of the blood and humours; by others, as owing to the disengagement of an igneous matter, or spirit from the blood; by others, to an agitation of the sulphureous parts of the blood; whilst Boerhaaveb and Douglase ascribe it to the friction of the blood against the parietes of the vessels, and of the globules against each other. In favour of the last hypothesis, it was urged, that animal heat is in a direct ratio with the velocity of the circulation, the circumference of the vessels, and the extent of their surface; and that thus we are able to explain, why the heat of parts decreases in a direct ratio with their distance from the heart; and the greater heat of the arterial blood, in the lungs, was accounted for, by the supposition, that the pulmonary circulation is far more rapid. Most of these notions are entirely hypothetical. The data are generally incorrect, and the deductions characteristic of the faulty physics of the period in which they were indulged. The correct view, it appears to us, is, that caloric is disengaged in every part, by a special chemico-vital action, which, in animals, is greatly under the nervous influence. In this manner, calorification becomes a function executed in the whole system of nutrition, and therefore appropriately considered in this place. It has been remarked by Tiedemann,d that the intussusception of alimentary matters, and their assimilation by digestion and respiration, the circulation of the humours, nutrition and secretion, the renewal of materials accompanying the exercise of life, and the constant changes of composition in the solid and liquid parts of the organism, - all of which are under the nervous influence, - participate in the evolution of heat, and we deceive ourselves when we look for the cause in one of those acts only. In some experiments by Dr. Robert E. Rogers,e he found that when recently drawn venous

Dr. W. B. Carpenter, Principles of General and Comparative Physiology, p. 379, Lond. 1839.

<sup>&</sup>lt;sup>b</sup> Gerard Van Swieten, Comment in Boerhaav. Aphorism., &c. § 382, 675, Ludg. Bat. 1742-1772. On Animal Heat, p. 47, Lond. 1747.

d Traité de Physiologie, &c. trad. par Jourdan, p. 514, Paris, 1831; Ley, op. cit. Appendix, p. 303; and Collard de Martigny, in Journal Complémentaire des Sciences Médicales, xliii. 268, Paris, 1832.

<sup>·</sup> Amer. Journ. of the Med. Sciences, p. 297, for Aug. 1836.

blood contained in a freshly removed pig's bladder was immersed in oxygen gas, there was a remarkable elevation of temperature. Dr. Davya performed some experiments which led to the same results. In one of these, he took a very thin vial, of the capacity of eight liquid ounces, and carefully enveloped it in badly conducting substances - namely, several folds of flannel, of fine oiled paper, and of oiled cloth. Thus prepared, and a perforated cork being provided, holding a delicate thermometer, two cubic inches of mercury were introduced, and immediately after it was filled with venous blood kept liquid by agitation. The vial was then corked, and shaken. The thermometer included was stationary at 45°. After five minutes, during which it remained stationary, it was withdrawn; the vial closed by another cork, was transferred inverted to a mercurial bath, and 12 cubic inch of oxygen was introduced. The common cork was returned, and the vial was well agitated for about a minute: the thermometer was now introduced; it rose immediately to 46°, and by continuing the agitation, it rose further to 46.5°, and very nearly to 47°. This experiment was made on the blood of the sheep. These and other experiments of a similar character, in Dr. Davy's opinion, appear to favour the idea, that animal heat is owing, first, to the fixation or condensation of oxygen in the blood in the lungs, in its conversion from venous to arterial; and secondly, to the combinations into which it enters in the circulation in connexion with the different secretions and changes essential to animal life.

It is by the theory of the general evolution of caloric in the capillary system, or in the system of nutrition, that we are capable of accounting for the increased heat that occurs in certain local affections, in which the temperature greatly exceeds that of the same parts in health. By some, it has been doubted, whether, in cases of local inflammation, any such augmentation of temperature exists, but the error seems to have arisen from the temperature of the part, in health, having generally been ranked at blood heat; whereas, we shall find, that it differs essentially in different parts. Dr. Thomson found, that a small inflamed spot, in his right groin, gave out, in the course of four days, a quantity of heat, sufficient to have heated seven wine-pints of water from 40° to 212°; yet the temperature was not sensibly less than that of the rest of the body at the end of the experiment, when the

inflammation had ceased.b

Of the mode in which heat is evolved in the capillaries, it is impossible for us to arrive at any satisfactory information. The result alone indicates, that the process has been accomplished. In the present state of our knowledge, we are compelled to refer it to some vital action, of the nature of which we are ignorant; but which seems to be possessed by all organized bodies, — vegetable as well as animal. By supposing, that calorification is effected

Proceedings of the Royal Society for 1837-8, No. 34, and Researches, Physiological and Anatomical, Dunglison's American Med. Lib. Edit. p. 89, Philad. 1840.
 Annals of Philosophy, ii. 27.

in every part of the body, we can understand why different portions should have different temperatures; as the activity of the function may vary, in this respect, according to the organ. Chopart and Dessault found the heat of the rectum to be 100°; of the axilla and groin, when covered with clothes, 96°; and of the chest, 92°. Dr. Davyb found the temperature of a naked man, just risen from bed, to be 90° in the middle of the sole of the foot; 93° between the inner ankle and tendo-achillis; 91.5° in the middle of the shin; 93° in the calf; 95° in the ham; 91° in the middle of the thigh; 96.5° in the fold of the groin; 95° at three lines beneath the umbilicus; 94° on the sixth rib of the left side; 93° on the same rib of the right side; and 98° in the axilla. MM. Edwards and Gentil found the temperature of a strong adult male, to be, in the rectum and mouth, 102°; in the hands, 100°; in the axilla and groins, 98°; on the cheeks, 97°; in the prepuce and the feet, 96°; and on the chest and abdomen, 95°. All these experiments, it is obvious, concern only temperature of parts, which can be readily modified by the circumambient medium. To judge of the comparative temperature of the internal organs, Dr. Davy killed a calf, and noted the temperature of different parts, both external and internal. The blood of the jugular vein raised the thermometer to 105.5°; and that of the carotid artery to 107°. The heat of the rectum was 105.5°; of the metatarsus, 97°; of the tarsus, 90°; of the knee, 102°; of the head of the femur, 103°; of the groin, 104°; of the under part of the liver, 106°; of the substance of that organ, 106°; of the lung, 106.5°; of the left ventricle, 107°; of the right, 106°; and of the substance of the brain, 104°.

In the case of fistulous opening into the stomach, observed by Dr. Beaumont, the thermometer indicated a difference of three-fourths of a degree between the splenic and pyloric orifices of the stomach; the temperature of the latter being more elevated. It is not easy to account for these differences without supposing that each part has the power of disengaging its own heat, and that the communication of caloric is not sufficiently ready to prevent the

difference from being perceptible.

It was stated early in this section, that man possesses the power of resisting cold as well as heat within certain limits, and of preserving his temperature greatly unmodified. Let us inquire into the direct and indirect agents of these counteracting influences. As the mean temperature of the warmest regions does not exceed 85° of Fahrenheit, it is obvious that he must be constantly disengaging caloric to the surrounding medium: — still, his temperature remains the same. This is effected by the mysterious agency which we have been considering, materially aided, however, by several circumstances, both intrinsic and extrinsic. The external envelope of the body is a bad conductor of caloric, and therefore

b Philosoph. Transact. for 1814.

<sup>3</sup> J. Hunter, Observations on the Animal Œconomy, Lond. 1786.

Exp. and Observations on the Gastric Juice, p. 274, Plattsburg, 1833.

it protects the internal organs, to a certain extent, from the sudden influence of excessive heat or cold. But the cutaneous system of man is a much less efficient protection than that of animals. In the warm-blooded animals, in general, the bodies are covered with hair or feathers. The whale is destitute of hair; but, besides the protection which is afforded by the extraordinary thickness of its skin, and the stratum of fat—a bad conductor of caloric - with which the skin is lined, as the animal constantly resides in the water, it is not subjected to the same vicissitudes of temperature as land animals. The seals, bears, and walruses, which seek their food in the same seas, sleep on land. They have a coating of hair to protect them. In the cases of some of the birds of the genus Anas, of northern regions, we meet with a singular anomaly, the whole of the circumference of the anus being devoid of feathers, but to make amends for this deficiency, the animal has the power of secreting an oleaginous substance, with which the surface is kept constantly smeared. It may be remarked, that we do not find the quantity of feathers on the bodies of birds to be proportionate to the cold of the climates in which they reside, as is pretty universally the case regarding the quantity of hair on the mammalia.

Man is compelled to have recourse to clothing, for the purpose of preventing the sudden abstraction or reception of heat. This he does by covering himself with substances which are bad conductors of caloric, and retain an atmosphere next to the surface, which is warmed by the caloric of the body. He is compelled, also, in the colder seasons, to have recourse to artificial temperature, and it will be obvious, from what has already been said, that the greater the degree of activity of any organ or set of organs, the greater will be the heat developed; and in this way muscular exertion

and digestion influence its production.

By an attention to all these points, and by his acquaintance with the physical laws relative to the development and propagation of caloric, man is enabled to live amongst the arctic snows, and to exist in climates, where the temperature is frequently for a length of time upwards of 150° lower than that of his own body. The contrivances adopted in the polar voyages, under the direction of Captain Parry and others, are monuments of ingenuity directed to obviate one of the greatest obstacles to prolonged existence in inhospitable regions, for which man is naturally incapacitated, and for which he attains the capability solely by the exercise of that superior intellect with which he has been vested by the Author of his being. In periods of intense cold, the extreme parts of the body do not possess the necessary degree of vitality to resist congelation, unless they are carefully protected. In the disastrous expedition of Napoleon to Russia, the loss of the nose and ears was a common casualty; and, in arctic voyages, frost-bites occur in spite of every care." When the temperature of the whole body

<sup>&</sup>lt;sup>a</sup> Larrey, Mémoires de Chirurgie Militaire et Campagnes, tom. iv. p. 91, 106, and 123, Paris, 1817.

sinks to about 78° or 79°, death takes place, preceded by the symptoms of nervous depression, which have been previously

depicted.

The counteracting influence, which is exerted, when the body is exposed to a temperature greatly above the ordinary standard of the animal, is as difficult of appreciation as that by which calorification is effected. The probability is, that, in such case, the disengagement of animal heat is suspended; and that the body receives heat from without, by direct, but not by rapid, communication, owing to its being an imperfect conductor of caloric. Through the agency of this extraneous heat, the temperature rises a limited number of degrees; but its elevation is generally considered to be checked by the evaporation, constantly taking place through the cutaneous and pulmonary transpirations. For this last idea we are indebted to Franklin, and its correctness and truth have been maintained by most observers. MM. Berger and Delaroche put into an oven, heated to from 120° to 140°, a frog, one of those porous vessels, called alcarazas - which permit the transudation of the fluid, within them, through their sides - filled with water at the animal heat, and two sponges, imbibed with the same water. The temperature of the frog at the expiration of two hours, was 99°; and the other bodies continued at the same. Having substituted a rabbit for the frog, the result was identical. On the other hand, having placed animals in a warm atmosphere, so saturated with humidity that no evaporation could occur, they received the caloric by communication, and their temperature rose; whilst inert, evaporable bodies, put into a dry stove, became but slightly warmed; - much less so, indeed, than the warm-blooded animals in the moist stove. Hence they concluded, that evaporation is a great refrigerative agent when the body is exposed to excessive heat; and that such evaporation was considerable, was shown by the loss in weight, which animals sustain by the experiment.

Recently, however, it has been contested, that the cutaneous evaporation has any effect in tempering the heat of the body; whilst it is admitted that the elimination from the system of a certain quantity of aqueous matter is all important; — whatever arrests it being the source of morbid phenomena. MM. Becquerel and Breschet<sup>e</sup> found, when the hair of rabbits had been shaved off, and the skin covered with an impermeable coating of strong glue, suet, and resin, that the animals died very soon afterwards, and, they thought, by a process of asphyxia in consequence of the transpiration from the skin being prevented. In these experiments, they found to their surprise, that the temperature of the animals, instead of rising, fell considerably. Thus, the temperature of the first rabbit, before it was shaved, and covered with the impermeable

Works, iii. 294, Philad. 1809; or Spark's edit. vi. 213, Boston, 1838.

c Comptes Rendus, Oct. 1841.

b Delaroche, in Journal de Physique, lxxi. 289, Paris, 1810; Coutanceau, Dict. de Méd. v. 23; and Magendie's Précis, ii. 509.

coating, was 38° centigrade; but immediately after the coating was dry, the temperature of the muscles of the thigh and breast had fallen to 24.5 centigrade. In another rabbit, the coating on which was put on with more care, - as soon as the coating was dried, the temperature was found to have fallen so much that it was only three degrees above that of the surrounding atmosphere, which was on that day 17° centigrade. An hour after this, the animal died. These experiments clearly exhibit the importance of the functions executed by the skin. Dr. Carpentera thinks they place in a very striking point of view the importance of the cutaneous surface as a respiratory organ, and enable us to understand how, when the aërating power of the lungs is nearly destroyed by disease, the heat of the body is kept up to its natural standard by the action of the skin. "A valuable therapeutic indication, also," he adds, "is derivable from the knowledge which we thus gain of the importance of the cutaneous respiration; for it leads us to perceive the desirableness of keeping the skin moist in those febrile diseases in which there is great heat and dryness of the surface, since aëration cannot properly take place through a dry mem-

Dr. Edwards, in his experiments on the influence of physical agents on life, found, that warm-blooded animals have less power of producing heat, after they have been exposed for some time to an elevated temperature, as in summer, - whilst the opposite effect occurs in winter. He instituted a series of experiments. which consisted in exposing birds to the influence of a freezing mixture, first in February, and afterwards in July and August, and observing in what degree they were cooled by remaining in this situation for equal lengths of time; the result of which was, that the same kind of animal was cooled six or eight times as much in the summer as in the winter months. This principle he presumes to be of great importance in maintaining the regularity of the temperature at the different seasons; even more so than evaporation, the effect of which, in this respect, he thinks, has been greatly exaggerated. From several experiments on yellow hammers made at different periods in the course of the year, it would result, that the averages of their temperature ranged progressively upwards from the depth of winter to the height of summer, within the limits of five or six degrees of Fahrenheit, and the contrary was observed in the fall of the year. Hence, Dr. Edwards infers, and with every probability, that the temperature of man experiences a similar fluctuation.b

When exposed to high atmospheric temperature, the ingenuity of man has to be as much exerted as in the opposite circumstances. The clothing must be duly regulated according to physical princi-

b De l'Influence des Agens Physiques, p. 489; and Hodgkin and Fisher's translation, Lond. 1832.

<sup>\*</sup> Human Physiology, § 726, Lond. 1842. See, also, Dr. R. Willis, Proceedings of the Royal Society of London, No. 56.

ples,<sup>a</sup> and perfect quietude be observed, so that undue activity of any of the organs that materially influence the disengagement of animal heat may be prevented. It is only within limits, that this refrigerating action is sufficient. At a certain degree, the transpiration is inadequate, the temperature of the animal rises, and death supervenes.

## CHAPTER VII.

#### SECRETION.

We have yet to describe an important and multiple function, which also takes place in the intermediate system — in the very tissue of our organs — and which separates from the blood the various humours of the body. This is the function of secretion, — a term literally signifying separation, and which has been applied both to the operation and the product. Thus, the liver is said to separate the bile from the blood by an action of secretion, and the bile is said to be a secretion.

The organs that execute the various secretory operations differ greatly from each other. They have, however, been grouped by anatomists into three classes, each of which will require a general notice.

#### 1. ANATOMY OF THE SECRETORY APPARATUS.

The secretory organs have been divided into the exhalant, the

follicular, and the glandular.


The remarks made respecting the exhalant vessels, under the head of nutrition, will render it unnecessary to allude, in this place, to any of the apocryphal descriptions of them, especially as their very existence is supposititious. Many, indeed, imagine them to be nothing more than the minute radicles of ordinary arteries.

The follicle or crypt has the form of an ampulla or vesicle, and is situate in the substance of the skin and mucous membranes; secreting a fluid for the purpose of lubricating those parts. In the exhalant vessel, the secreted fluid passes immediately from the bloodvessel, without being received into any excretory duct; and, in the follicle, there is essentially no duct specially destined for the excretion of the humour. The follicle is membranous and vascular, having an internal cavity into which the secretion is poured; and the product is excreted upon the surface beneath which it is situate, either by a central aperture, or by a very short duct — if duct it can be called — generally termed a lacuna.

The gland is of a more complex structure than the last. It consists of an artery which conveys blood to it; of an intermediate

<sup>\*</sup> See the chapter on Clothing in the author's "Elements of Hygiène," p. 388, Philad. 1835.

body, — the gland, properly so called, — and of an excretory duct to carry off the secreted fluid, and to pour it on the surface of the skin or mucous membrane. The bloodvessel, that conveys to the gland the material from which the secretion has to be effected,



Secreting Arteries, and Nerves of the Intestines.

a a. A portion of the intestine. b b. Part of the aorta. c cc. Nerves following the branches of the aorta to supply the intestine.

enters the organ, at times, by various branches; at others, by a single trunk, and ramifies in the tissue of the gland; communicating at its extremities with the origins of the veins and of the excretory ducts. These ducts arise by fine radicles at the part where the arterial ramifications terminate; and they unite to form larger and less numerous canals, until they terminate in one large duct, as in the pancreas; or in several, as in the lachrymal gland; the duct generally leaving the gland at the part where the bloodvessel enters. Of this we have a good exemplification in the kidney.

The pavement and the cylinder epithelium, as well as all the intermediate forms, are met with in the different glands. These are not necessarily a continuation of that of the cutaneous system; on the contrary, the epithelium of the latter is often seen changing

its form at its entrance into the gland."

Besides the vessels above mentioned, veins exist, which communicate with the vessels that convey blood to the gland, both for the formation of the humour and the nutrition of the organ, and which return the residuary blood to the heart. Lymphatic vessels are likewise there; and nerves, — which proceed from the gan-

\* Mandl, Manuel d'Anatomie générale, p. 450, Paris, 1843.

glionic system, — form a network around the secreting arteries, as in Fig. 184, accompany them into the interior of the organ, and

terminate, like them, invisibly.

Bordeu<sup>a</sup> was of opinion, that the glands, judging from the parotid, are largely supplied with nerves. The nerves, however, do not all belong to it, some merely crossing it in their course to other parts. Bichat, from the small number sent to the liver, was in-

duced to draw opposite conclusions to those of Bordeu.

These may be looked upon as the great components of the glandular structure. They are bound together by cellular membrane, and have generally an outer envelope. The intimate texture of these organs has been a topic of much speculation. It is generally considered, that the final ramifications of the arterial vessels, with the radicles of the veins and excretory ducts, and the final ramifications of the lymphatic vessels and nerves, form so many small lobules, composed of minute, granular masses. Such, indeed, is the appearance the texture presents, when examined by the naked eye. Each lobule is conceived to contain a final ramification of the vessel or vessels that convey blood to the organ, a nerve, a vein, a lymphatic, and an excretory duct, - with cellular tissue binding them together. When the organ has an external membrane, it usually forms a sheath to the various vessels. The lobated structure is not equally apparent in all the glands. It is well seen in the pancreas, and in the salivary and lachrymal. The precise mode in which the bloodvessel, from the blood of which the secretion is effected, communicates with the excretory duct, does not admit of detection. Some have supposed, that between the termination of the bloodvessel and the commencement of the duct. a secretory vessel, or a spongy tissue specially charged with the function, exists, which conveys the secreted humour into the excretory duct. Of this, however, we have no evidence. Professor Müllere maintains, that the glandular structure consists essentially of a duct with a blind extremity, on whose parietes plexuses of bloodvessels ramify, from which the secretions are immediately produced, -- a view which is confirmed by the pathological appearances, in a case of disease of the portal system, that fell under the author's observation, and is referred to under the Secretion of Bile. The opinion of Malpighid was similar. He affirmed that such glands as the liver are composed of very minute bodies, called acini, from their resemblance to the stones of grapes; that these acini are hollow internally, and are covered externally by a network of bloodvessels; and that these minute bloodvessels pour into the cavities of the acini the secreted fluid, from which it is

<sup>\*</sup> Sur les Glandes in Œuvres Complètes, par M. Richerand, Paris, 1818.

b Anat. Général. tom. ii.

e De Glandular, Secernent, Structurà Penitiori, &c., Lips. 1830; or the English Edit. by Mr. Solly, Lond. 1839.

d Opera Omnia, &c., p. 300, Lugd. Batav. 1687.

subsequently taken up by the excretory ducts. Ruysch, however, held, that the acini of Malpighi are merely convoluted vessels, and that they are continuous with the excretory ducts. In Malpighi's view, the secretory organ is a mere collection of follicles; in Ruysch's, simply an exhalant membrane variously convoluted. "The chief, if not the only difference," says a popular writer, "between the secreting structure of glands and that of simple surfaces, appears to consist in the different number and the different arrangement of their capillary vessels. The actual secreting organ is in both cases the same, - capillary bloodvessel; and it is uncertain whether either its peculiar arrangement, or greater extent in glandular texture, be productive of any other effect than that of furnishing the largest quantity of bloodvessels within the smallest space. Thus convoluted and packed up, secreting organ may be procured to any amount that may be required, without the inconvenience of bulk and weight."c

It is manifest then, that the simplest form of the secretory apparatus is this simple capillary vessel; and that the follicles and

glands are structures of a more complex organization.

### 2. PHYSIOLOGY OF SECRETION.

The uncertainty which rests upon the intimate structure of secreting organs, and upon the mode in which the different bloodvessels communicate with the commencement of the excretory duct, envelopes the function, executed by those parts, in obscurity. We see the pancreatic artery pass to the pancreas, ramify in its tissue, become capillary, and escape detection; and we see other vessels becoming larger and larger, and emptying themselves into vessels of greater magnitude, until, ultimately, all the secreted humour is contained in one large duct, which passes onwards and discharges its fluid into the small intestine. Yet if we follow the pancreatic artery as far back as the eve can carry us, even when aided by glasses of considerable magnifying power, or if we trace back the pancreatic duct as far as practicable, we find, in the former vessel, always arterial blood, and in the latter, always pancreatic juice. It must, consequently, be between the part at which the artery ceases to be visible, and at which the pancreatic duct becomes so, that secretion is effected; and we cut the knot by asserting, that it occurs in the very tissue, parenchyma, or in the capillary system of the secretory organ. Conjecture, in the absence of positive knowledge, has been busy, at all times, in attempting to explain the mysterious agency by which we find such various humours separated from the same fluid; and, according as chemical, or mechanical, or exclusively vital doctrines

<sup>\*</sup> Epist. Anatom. qua Respondet Viro Clarissimo Hermann. Boerhaav. p. 45, Lugd. Batav. 1722.

<sup>&</sup>lt;sup>b</sup> Dr. Southwood Smith, in Animal Physiology, p. 115; Library of Useful Knowledge, Lond. 1829.

e See Béclard, in art. Glande, Dict. de Méd. x. 256, Paris, 1824; and Carpenter, Human Physiology, § 649, London, 1842.

have prevailed in physiology, the function has been referred to one or other of those agencies. The general belief, amongst the physiologists of the sixteenth and seventeenth centuries, was, that each gland possesses a peculiar kind of fermentation, which assimilates to its own nature the blood passing through it. The notion of fermentation was, indeed, applied to most of the vital phenomena. It is now totally abandoned owing to its being purely imaginary, and inconsistent with all our ideas of the vital operations. When this notion passed away, and the fashion of accounting for physiological phenomena on mechanical principles usurped its place, the opinion prevailed, that the secretions are effected through the glands as through filters. To admit of this mechanical result, it was maintained, that all the secreted fluids exist ready formed in the blood, and that, when they respectively arrive at the different secretory organs, they pass through, and are received by the excretory ducts. Descartes and Leibnitz were warm supporters of this mechanical doctrine, although their views differed materially with regard to the precise nature of the operation. Descartes supposed, that the particles of the various humours are of different shapes, and that the pores of the glands have respectively a corresponding figure; so that each glaud permits those particles only to pass through it which have the shape of its pores. Leibnitz, on the other hand, likened the glands to filters, which had their pores saturated with their own peculiar substance, so that they admitted this substance to pass through them, and excluded all others, as paper, saturated with oil, will prevent the filtration of water. The mechanical doctrine of secretion was taught by Malpighi and by Boerhaave, and it continued to prevail even till the time of Haller. All the secretions were conceived to be ready formed in the blood, and the glands were looked upon as the sieves or strainers to convey off the appropriate fluids or humours. In this view of the subject, all secretion was a transudation through the coats of the vessels, - particles of various sizes passing through pores adapted to them.d

The mechanical doctrine of transudation, in this shape, is founded upon supposititious data; and the whole facts and arguments are so manifestly defective, that no refutation is necessary. It is now, indeed, wholly abandoned. MM. Magendie and Fodéra have, however, revived the mechanical doctrine of late years, but under an essentally different form; and one applicable especially to the exhalations. The former gentleman, believing that many of the exhalations exist ready formed in the blood, thinks, that the character of the exhaled fluid is dependent upon the physical arrangement of the small vessels, and his views repose upon the following experiments. If, in the dead body, we inject warm

<sup>\*</sup> Tractatus de Homine, p. 18, Amstel. 1677. b Haller, Element. Physiol. vii. 3. Prælectiones Academicæ, &c. edit. A. Haller, § 253, Gotting. 1740-1743.

d Mascagni, Nova per Poros Inorganicos Secretionem Theoria. Rom. 1793, tom. ii. Précis, &c. edit. cit. ii. 444.

water into an artery passing to a serous membrane, as soon as the current is established from the artery to the vein, a multitude of minute drops may be observed oozing through the membrane, which speedily evaporate. If, again, a solution of gelatin, coloured with vermilion, be injected into all the vessels, it will often happen, that the gelatin is deposited around the cerebral convolutions, and in the anfractuosities, without the colouring matter escaping from the vessels, whilst the latter is spread over the external and internal surface of the choroid. If, again, linseed oil, also coloured with vermilion, form the matter of the injection, the oil, devoid of colouring matter, is deposited in the articulations, that are furnished with large synovial capsules, no transudation takes place at the surface of the brain, or in the interior of the eye. Magendie asks, if these be not instances of true secretion taking place post-mortem, and evidently dependent upon the physical arrangement of the small vessels; and whether it be not extremely probable, that the same arrangement must, in part at least, preside over exhalation during life? Fodéra, to whose experiments on the imbibition of tissues we had occasion to allude under the head of absorption, embraces the views of Magendie. If the vessels of a dead body, he remarks, be injected, the substance of the injection is seen oozing through the vessels; and if an artery and a vein be exposed in a living animal, a similar oozing through the parietes is observable. This is more manifest if the trunk, whence the artery originates, be tied, - the fluid being occasionally bloody. If the jugular veins be tied, not only does ædema occur in the parts above the ligatures, but there is an increase of the salivary secretion. It is not necessary to adduce the various experiments of Fodéra, relating to this topic, or those of Harlan, Lawrence and Coates, or of Dutrochet, Faust, Mitchell. and others. They are of precisely the same character as those that we have previously described regarding the imbibition of tissues; and transudation is only imbibition or soaking from within to without: Magendie and Fodéra, indeed, conclude, that one primary physical cause of exhalation is the same as that of absorption, - namely, imbibition.

Another physical cause, adduced by Magendie, is the pressure experienced by the blood in the circulatory system, which, he conceives, contributes powerfully to cause the more aqueous part to pass through the coats of the vessels. If water be forcibly injected, by means of a syringe, into an artery, all the surfaces, to which the vessel is distributed, as well as the larger branches and the trunk itself, exhibit the injected fluid oozing in greater abundance according to the force exerted in the injection. He farther remarks, that if water be injected into the veins of an animal, in sufficient quantity to double or treble the natural amount of blood, a considerable distension of the circulatory organs is produced; and,

<sup>&</sup>lt;sup>2</sup> Magendie's Journal de Physiologie, iii. 35; and Recherches, &c. sur l'Absorption et l'Exhalation, Paris, 1824.

consequently, the pressure, experienced by the circulating fluid, is largely augmented. If any serous membrane be now examined, — as the peritoneum, — a serous fluid is observed issuing rapidly from its surface, which accumulates in the cavity, and produces a true dropsy under the eyes of the experimenter, and, occasionally, the colouring part of the blood transudes at the surface of certain organs, as the liver, spleen, &c. Hamberger, again, broached the untenable physical hypothesis, that each secreted humour is deposited in its proper secretory organ, by virtue of its

specific gravity.a

It is obvious, that all these speculations proceed upon the belief, that the exhalations exist ready formed in the blood; and that, consequently, the act of secretion, so far as concerns them, is one of separation or of secerning, - not of fresh formation. That this is the case with the more aqueous secretions is probable, and not impossible with regard to the rest. Organic chemistry is subject to more difficulties in the way of analysis than inorganic; and it can be readily understood, that, in a fluid so heterogeneous as the blood, the discovery of any distinct humour may be impracticable. Of course, the elements of every fluid, as well as solid, must be contained in it; and we have already seen, that not merely the inorganic elements, but the organic, or compounds of organization, have been detected by the labours of Chevreul and others. There are, indeed, some singular facts connected with this subject. MM. Prévost and Dumas, having removed the kidneys in cats and dogs, and afterwards analyzed the blood, found urea in it - the characteristic element of urine. This principle was contained in greater quantity, the longer the period that had elapsed after the operation; whilst it could not be detected in the blood, where the kidneys existed. The experiment was soon afterwards repeated by Vauquelin and Ségalase with the same results. The latter introduced urea into the veins of an animal, whose kidneys were untouched; he was unable to detect the principle in the blood; but the urinary secretion was largely augmented after the injection. Whence he concludes that urea is an excellent diuretic. More recently, MM. Gmelin and Tiedemann, in association with M. Mitscherlich, have arrived, experimentally, at the same conclusions as MM. Prévost and Dumas. The existence of urea in the fluid ejected from the stomach of the animal was rendered probable, but there were no traces of it in the fæces or the bile. The animal died the day after the extirpation of the second kidney. They were totally unable to detect either urea, or sugar of milk in the healthy blood of the cow. These circumstances would favour the idea,

<sup>\*</sup> See Adelon, Physiologie de l'Homme, 2de édit. iii. 455, Paris, 1829; and Richerand, Elemens de Pyysiologie, 13ème édit. par M. Bérard ainé, p. 176, Bruxelles, 1837.

b Annales de Chimie, tom. xxii. and xxxiii. 90.

Magendie, Précis, &c., ii. 478.
 Tiedemann und Treviranus, Zeitschrift für Physiol. B. v. Heft i.; and Brit. and Foreign Med. Review, p. 592, for April, 1836.

that certain of the secretions may be formed in the blood, and may simply require the intervention of a secreting organ to separate them; but the mode in which such separation is effected is entirely inexplicable under the doctrine of simple mechanical filtration or transudation. It is unlike any physical process, which can be imagined. The doctrines of filtration and transudation can apply only to those exhalations, in which the humour has undergone no apparent change; and it is obviously impossible to specify these, in the imperfect state of our means of analysis. In the ordinary aqueous secretions, simple transudation may embrace the whole process; and, therefore, it is unnecessary to have recourse to any other explanation; especially after the experiments instituted by Magendie, supported by pathological observations in which there has been partial ædema of the legs, accompanied by more or less complete obliteration of the veins of the infiltrated part, - the vessels being obstructed by fibrinous coagula, or compressed by circumjacent tumours. It is obvious, that ascites or dropsy of the lower belly may be frequently occasioned by obstruction of the portal circulation in the liver, and that in this way, we may account for the frequency with which we find a union of hydropic and hepatic affections in the same individual. The same pathological doctrine, founded on direct observation, has been extended to phlegmatia dolens or swelled leg; an affection occurring in the puerperal state, and which has often been found connected with obstruction in the great veins that convey the blood back from the lower extremity. It may not, consequently, be wide of the truth — if not wholly accurate — to consider the secretions, with Dr. Billing, b to be "vital transudations from the capillaries into the excretory ducts of the glands, by pores invisible to our senses, even when aided by the most perfect optical instruments."

The generality of physiologists have regarded the more complex secretions—the follicular and the glandular—as the results of chemical action; and under the view, that these secretions do not exist ready formed in the blood, and that the elements alone are contained in that fluid, it is impossible not to admit that chemical agency must be exerted. In support of the chemical hypothesis, which has appeared under various forms,—some, as Keill, presuming that the secretions are formed in the blood, before they arrive at the place appointed for secretion; others, that the change is effected in the glands themselves,—the fact of the formation of a number of substances from a very few elements, provided these be united in different proportions, has been adduced. For example: take the elementary bodies, oxygen and azote. These, in one proportion, form atmospheric air; in another, nitrous oxide; in another, nitric oxide; in a fourth, hyponitrous acid; in a fifth,

<sup>Dr. W. Philip, in Lond. Med. Gazette for March 25th, 1837, p. 952.
First Principles of Medicine, Amer. Edit. p. 55, Philad. 1842.</sup> 

<sup>·</sup> Tentamina Medico-Physica, iv.; and Haller, Element. Physiolog. &c. vii. 3.

nitrous acid; in a sixth, nitric acid, &c., substances which differ as much as the various secretions differ from each other and from the blood. Many of the compounds of organization likewise exhibit by their elementary composition, that but a slight change is necessary, in order that they may be converted into each other. Dr. Prouta has exhibited this close alliance between three substances - urea, lithic acid, and sugar - and has shown how they may be converted into each other, by the addition or subtraction of single elements of their constituents. Urea is composed of two atoms of hydrogen, and one of carbon, oxygen, and azote respectively; by removing one of the atoms of hydrogen and the atom of nitrogen, it is converted into sugar; by adding to it an additional atom of carbon, into lithic acid. Dr. Bostock,b - who is disposed to push the application of chemistry to the explanation of the functions as far as possible, - to aid us in conceiving how a variety of substances may be produced from a single compound, by the intervention of physical causes alone, supposes the case of a quantity of the materials adapted for the vinous fermentation being allowed to flow from a reservoir, through tubes of various diameters, and with various degrees of velocity. "If we were to draw off portions of this fluid in different parts of its course or from tubes, which differed in their capacity, we should, in the first instance, obtain a portion of unfermented syrup; in the next, we should have a fluid in a state of incipient fermentation; in a third, the complete vinous liquor; while, in a fourth, we might have acetous acid." Any explanation, however, founded upon this loose analogy, is manifestly too physical: this Bostock admits, for he subsequently remarks, that "if we adopt the chemical theory of secretion, we must conceive of it as originating in the vital action of the vessels, which enables them to transmit the blood, or certain parts of it, to the various organs or structures of the body, where it is subjected to the action of those reagents, which are necessary to the production of these changes." The admission of such vital agency, in some shape, seems to be indispensable.

Attempts have been made to establish secretion as a nervous action; and numerous arguments and experiments have been brought forward in support of the position. That many of the secretions are affected by the condition of the mind is known to all. The act of crying, in evidence of joy or sorrow; the augmented action of the salivary glands at the sight of pleasant food; the increased secretion of the kidney during fear or anxiety, and the experimental confirmation, by Mr. Hunter, of the truth of the common assertion — that the she-ass gives milk no longer than the impression of the foal is on her mind; the skin of her foal, thrown over the back of another, and frequently brought near her, being sufficient to renew the secretion, — sufficiently indicate, that the

Medico-Chirurg. Transact. viii. 540.
 Physiol. 3d edit. p. 519, Lond. 1836.

organs of secretion can be influenced through the nervous system in the same manner as the functions of nutrition and calorification.a

The discovery of galvanism naturally suggested it as an important agent in the process, - or rather suggested, that the nervous fluid strongly resembles it. This conjecture seems to have been first hazarded by Berzelius, and by Sir Everard Home; b and. about the same time, an experiment was made by Dr. Wollaston, which he conceived to throw light upon the process. He took a glass tube, two inches high, and three-quarters of an inch in diameter; and closed it at one extremity with a piece of bladder. He then poured into the tube a little water, containing 1/240th of its weight of muriate of soda, moistened the bladder on the outside, and placed it upon a piece of silver. On curving a zinc wire so that one of its extremities touched the piece of metal, and the other dipped into the liquid to the depth of an inch, the outer surface of the bladder immediately indicated the presence of pure soda; so that, under this feeble electric influence, the muriate of soda was decomposed, and the soda, separated from the acid, passed through the bladder. M. Fodérad performed a similar experiment, and found, that whilst ordinary transudation frequently required an hour before it was evidenced, it was instantaneously exhibited under the galvanic influence. On putting a solution of prussiate of potassa into the bladder of a rabbit, forming a communication with the solution by means of a copper wire; and placing on the outside, a cloth soaked in a solution of sulphate of iron, to which an iron wire was attached; he found, by bringing these wires into communication with the galvanic pile, that the bladder or the cloth was suddenly coloured blue, according as the galvanic current set from without to within, or from within to without; - that is, according as the iron wire was made to communicate with the positive pole, and the copper wire with the negative, or conversely. But it is not necessary that there should be any communication with the galvanic pile. If an animal membrane, as a bladder containing iron filings, be immersed in a solution of sulphate of copper, the sulphuric acid will penetrate the membrane to reach the iron, with which it forms a sulphate, and the metallic copper will be deposited on the lower surface of the membrane; the animal membrane in such case, offering no obstacle to the action of the ordinary chemical affinities.

e Philosoph. Mag. xxxiii. 438.

d Magendie's Journal de Physiologie, iii. 35; and Researches, &c., sur l'Absorption

et l'Exhalation, Paris, 1824.

<sup>\*</sup> For several examples of the same kind, see Fletcher's Rudiments of Physiology, part ii. b. p. 10, Edinb. 1836; Burdach, Physiologie, u. s. w. § 522; Dr. A. Combe, on Infancy, Lond. and Philad. 1841; and Carpenter, Human Physiology, § 427, 428, Lond. 1842.

b Lectures on Comp. Anat. iii. 16, Lond 1836; and v. 154, Lond. 1828.

<sup>&</sup>lt;sup>e</sup> Dr. Robert E. Rogers, in the American Journal of the Medical Sciences, p. 291, for August, 1836. See, also, the observations of Prof. Mitchell and Dr. Draper, elsewhere referred to.

With some of the chemical physiologists, there has been a disposition to resolve secretion into a mere play of electric affinities. Thus, M. Donnéa affirms, that from the whole cutaneous surface is secreted an acid humour, whilst the digestive tube, except in the stomach, secretes an alkaline mucus; and hence, he infers, that the external acid, and the internal alkaline membranes of the human body represent the two poles of a pile, the electrical effects of which are appreciable by the galvanometer. On placing one of the conductors of the instrument in contact with the mucous membrane of the mouth, and the other in contact with the skin, the magnetic needle, he affirms, deviated fifteen, twenty, and even thirty degrees, according to its sensibility; and its direction indicated, that the mucous or alkaline membrane took negative, and the cutaneous membrane, positive electricity. He further asserts, that, between the acid stomach and the alkaline liver, extremely powerful electrical currents are formed. These experiments do not, however, aid us materially in our solution of the phenomena of secretion. They exhibit merely electric phenomena dependent upon difference of chemical composition. This is, indeed, corroborated by the experiments of M. Donné himself on the secretions of vegetables. He observed electrical phenomena of the same kind in them, but, he says, electric currents in vegetables are not produced by the acid or alkaline conditions of the parts as in animals, the juice of fruits being always more or less acid. Experiments of M. Biot, however, show, that the juices, which arrive by the pedicle, are modified in some part of the fruit, and M. Donné thinks it is perhaps to this difference in the chemical composition of the juices of the two extremities, that the electrical phenomena are to be attributed. The effects of the section of the pneumogastric nerves on the functions of digestion and respiration have been given elsewhere, at some length. It was there stated, that when digestion was suspended by their division, Dr. Wilson Philip was led to ascribe the suspension to the secretion of the gastric juice having been arrested; an opinion, which Sir B. Brodie had been induced to form previously, from the results of experiments, which showed, that the secretion of urine is suspended by the removal or destruction of the brain; and that when an animal is destroyed by arsenic, after the division of the pneumogastric nerves, all the usual symptoms are produced, except the peculiar secretion from the stomach. Sir B. Brodie did not draw the conclusion, that the nervous influence is absolutely necessary to secretion, but that it is a step in the process, and the experiments of Magendiec on the effect of the division of the nerve of the fifth pair on the nutritive secretion of the cornea, confirm the position. We have, indeed, numerous evidences, that the nervous system cannot be indispensable to secretion. In all animals, this power must exist, yet there are some in which no nervous system is apparent. Dr. Bostockd has given references, in a

4 Physiology, edit. cit. p. 525, Lond. 1836.

<sup>\*</sup> Annales de Chimie, &c. lvii. 400; and Journal Hebdomad., Fév. 1834.

Lond. Med. Gazette for March 18, and March 25, 1837. Précis, &c. ii. 489.

note, to cases of monstrous or deformed fætuses, born with many of their organs fully developed, yet where there was no nervous system. It may be said, however, that, in all these cases, an organic nervous system must have existed; but setting aside the cases of animals, we have the most indisputable testimony of the existence of secretion in the vegetable, in which there is no nervous system, or, at the most, a rudimental one; yet the function is accomplished as perfectly, and perhaps in as multiple a manner, as in man. It is manifest, therefore, that this is one of the vital actions occurring in the very tissue of organs, of which we have no more knowledge than we have of the capillary actions in general. All that we know is, that in particular organs various humours are secreted from the blood, some of which can be detected in that fluid, others not, but we are ignorant of the precise agency,

by which this mysterious process is effected.

Recently, the fashionable doctrine of cells has been applied to secretion. We have elsewhere stated the mode in which cells are stated by Mr. Goodsira to effect absorption (vol. i., p. 605). Secretion, according to him, is accomplished in a similar manner. During the progress of the development and growth of cells formed by the secreting organs, they appropriate certain substances with which they may be in relation; and when their term of existence is over - which is longer or shorter, in different cases - they discharge their contents into the excretory channels. An acinus, according to Mr. Goodsir, is at first a single nucleated cell. From the nucleus of this cell others are produced. From these, again, others arise in the same manner. The parent cell, however, does not dissolve away, but remains as a covering to the whole mass, and is appended to the extremity of the duct. Its cavity, therefore, as a consequence of its mode of development, has no communication with the duct. The original parent cell now begins to dissolve away, or to burst into the duct at a period when its contents have attained their full maturity, - this period varying in different glands, according to a law or laws impressed upon each of them. Mr. Goodsir considers growth and secretion to be identical - the same process under different circumstances. His views are worthy of the attention of the histological inquirer.

In cases of vicarious secretion, we have the singular phenomenon of organs assuming an action for which they were not destined. If the secretion from the kidney, for example, be arrested, urine is occasionally found in the ventricles of the brain, and, at other times, a urinous fluid has been discharged by vomiting or by cutaneous transpiration: the capillaries of these parts must, consequently, have assumed the functions of the kidney, and to this they must have been excited by the presence of urea, or of the elements of the urinary secretion in the blood - a fact, which exhibits the

a Transactions of the Royal Society of Edinburgh, 1842. See, also, Brit. and For. Med. Rev. Oct. 1842, p. 566; Carpenter, ibid. Jan. 1843, p. 279; and Mandl, Manuel d'Anatomie générale, p. 507, Paris, 1843. b Haller, Elementa Physiologiæ, lib. vii. S. i. § 9.

important influence, that the condition of the blood must exert on the secretions, and, indeed, on nutrition in general. It is thus that many of our remedial agents, alkalies, the preparations of iodine, &c. — produce their effects. They first enter the mass of blood, and, by circulating in the capillary system, induce a modification of its functions. There are other cases, again, in which the condition of the blood being natural, the vessels of nutrition may take on morbid action. Of this we have examples in the ossification of organs, which, in the healthy condition, have no osseous constituent; in the deposition of fat in cases of diseased ovaria; and in the altered secretions produced by any source of irritation in a

secreting organ.

In describing the physiology of the different secretions, one of three arrangements has usually been adopted; either according to the nature of the secreting organ, the function of the secreted fluid, or its chemical character. The first of these has been followed by Bichat and by Magendie, who have adopted the division into exhaled, follicular and glandular secretions. It is the arrangement followed by Lepelletier, except that he substitutes the term perspiratory for exhaled. According to the second, embraced by Boyer, b Sabatier, c and Adelon, they are divided into recrementitial secretions, or such as are taken up by internal absorption and re-enter the circulation, and into excrementitial, or such as are evacuated from the body, and constitute the excretions. Some physiologists add a third division - the recremento-excrementitial, - in which a part of the humour is absorbed and the remainder ejected. Lastly, the division according to chemical character, has been followed, with more or less modification, by Plenck, Richerand, Blumenbach, Youngh and Bostock: the last of whom, one of the most recent writers, has eight classes : - the aqueous, albuminous, mucous, gelatinous, fibrinous, oleaginous, resinous, and saline. To all of these classifications cogent objections might be made. The one we shall follow is the anatomical, not because it is the most perfect, but because it is the course that has been usually adopted throughout this work.

#### EXHALATIONS.

All the exhalations take place in the areolæ and internal cavities of the body, or from the skin and mucous membranes:—hence their division into internal and external. The former are recrementitial, the latter recremento-excrementitial. To the class of internal exhalations belong: 1. The serous exhalation.

· Précis de Physiologie, 2de édit. ii. 343, Paris, 1825.

b Anatomie, 2de édit. i. 8, Paris, 1803. CTraité Complet d'Anatomie, Paris, 1791.

d Physiologie de l'Homme, edit. cit. iii 438.

The Chemico-Physiological Doctrine of the Fluids, &c. translated by Dr. Hooper, Lond. 1797.

Elémens de Physiologie, 13ème édit. chap. vi. Bruxelles, 1837.

g Physiology, by Elliotson, 4th edit. Lond. 1828.

Medical Literature, p. 104, Lond. 1813.
Physiology, 3d edit. p. 48, Lond. 1836.

2. The serous exhalation of the cellular membrane. 3. The adipous exhalation of the cellular membrane. 4. The exhalation of the marrow. 5. The synovial exhalation. 6. The exhalation of the colouring matter of the skin, and of other parts; and 7. The areolar exhalation. To the class of external exhalations belong: 1. That of the skin or cutaneous transpiration. 2. The exhalation of the mucous membranes.

#### 1. INTERNAL EXHALATIONS.

### a. Serous Exhalation.

This is the fluid secreted by the serous membranes that line the various cavities of the body; - as the pleura, pericardium, peritoneum, arachnoid coat of the brain, and tunica vaginalis testis. Rudolphia asserts, that serous membranes are incapable of inflammation, are not vascular, and do not secrete; but that the secretions of shut sacs take place from the subjacent parts, and transude the serous membranes, which, in his view, are, consequently, a kind of cuticle. In a physiological consideration this is not of moment; and anatomically it only concerns the layer that covers the surface, whether it resemble the cuticle or not. From these membranes a fluid is exhaled, which is of an albuminous character, considerably resembling the serum of the blood, except in containing less albumen. M. Donnéb says it is always alkaline in the healthy state. This is owing to the presence of carbonate or albuminate of soda. It contains 7 or 8 per cent. of albumen and salts. In health, this fluid never accumulates in the cavities; the absorbents taking it up in proportion as it is deposited; but if, from any cause, the exhalants should pour out a larger quantity than usual, whilst the absorbents are not proportionably excited, accumulation may take place; or the same effect may ensue if the exhalants pour out no more than their usual quantity, whilst the absorbents do not possess their due activity. Under either circumstance, we have an accumulation or dropsy. The exhaled fluid probably transudes through the parietes of the arteries, and re-enters the circulation by imbibition through the coats of the veins. If we kill an animal and open it immediately afterwards, this exhalation appears in the form of a halitus or vapour, and the fluid is seen lubricating the free surface of the membrane. This, indeed, appears to be its principal office; by which it favours the motion of the organs upon each other.

The serous exhalations probably differ somewhat in each cavity, or according to the precise structure of the membrane. The difference between the chemical character of the fluid of the dropsy of different cavities would lead to this belief. As a general rule, according to Dr. Bostock, the fluid from the cavity of the abdomen contains the greatest proportion of albumen, and that from

the brain the least; but many exceptions occur to this.d

a Grundriss der Physiologie, 113. See, also, Hodgkin's Lectures on the Morbid Anatomy of the Serous and Mucous Membranes, P. i p. 27, Lond. 1836.


Journal Hebdomad. Février, 1834.
 Op. citat. p. 485.
 Burdach's Physiologie als Erfahrungswissenschaft, v. 184, Leipz. 1835.

# b. Serous Exhalation of the Cellular Membrane.

The nature of the primary cellular, areolar or fibro-cellular

membrane has been given in an early part of this work (vol. i., p. 38). As we observe it is not properly cellular, but is composed of a network of fibres, and of lamellæ formed by the adhesion of fibres laid side by side; and these interwoven so as to leave numerous interstices and cavities amongst them, having a tolerably free communication with each other.<sup>a</sup>

These interstices, wherever existing, are kept moist by a serous fluid, analogous to that exhaled from serous membranes, and which appears to have the same uses,—that of facilitating the motion of the lamellæ, or fibres on each other, and consequently of the organs, between which the cellular tissue is



Portion of areolar tissue inflated and dried, showing the general character of its larger meshes; magnified twenty diameters. — (Todd and Bowman.)

placed. When this secretion collects, from the causes mentioned in the last section, the disease, called ædema or anasarca, is induced.

## c. Adipous Exhalation of the Cellular Membrane.

Considerable diversity of opinion has prevailed regarding the precise organ of the secretion of fat. Haller supposed, that the substance exists ready formed in the blood, and that it simply transudes through the pores of the arteries; and Chevreul and others have given some countenance to this opinion, by the circumstance of their having met with fatty matter in that fluid. Anatomists have, likewise, been divided upon the subject of the precise tissue into which the fat is deposited; some believing it to be the ordinary cellular tissue, into which it is dropped by the agency of appropriate vessels; others, as Malpighib and William Hunter, believing in the existence of a peculiar adipous tissue, consisting, according to Béclard, of small bursæ or membranous vesicles, which inclose the fat, and are found in the areolæ of the cellular tissue. These vesicles are said to vary greatly in size: generally, they are round and globular; and, in certain subjects, receive vessels, that are very apparent. They form so many small sacs without apertures, in the interior of which are filaments

\* For the histology of the cellular membrane, see Todd and Bowman, Physiological Anatomy and Physiology of Man, London, 1842; Carpenter, Human Physiology, Amer. Edit. by Dr. Clymer, § 638, Philad. 1843; Gerber, General Anatomy, by Gulliver, p. 213, Lond. 1842; and Mandl. Manuel d'Anatomie, p. 111, Paris, 1843.

b De Omento, Pinguedine, et Adiposis Ductibus, in Oper. Lond. 1687.


Medical Observations and Inquiries, vol ii. Lond. 1757.

4 Art. Adipeux, in Dictionnaire de Médecine, tom. i.; and Elements of General Anatomy, translated by Togno, p. 128, Philad. 1830.

arranged like septa. In fatty subjects, these adipous vesicles are very perceptible, being attached to the cellular tissue and neighbouring parts by a vascular pedicle.

The fat, according to modern histologists, originates from fat-

Fig. 186.



Fat vesicles, assuming the polyhedral form from pressure against one another. The capillary vessels are not represented. — From the omentum; magnified about 300 diameters.—(Todd and Bowman.)

cells, which are usually of a spherical or spheroidal shape, but sometimes when closely pressed together without the intervention of any intercellular substance, they become polyhedral.a The adipous tissue is formed by a membrane of extreme tenuity which forms the vesicle that includes the fat. The membrane is homogeneous and transparent, about the 2 1000 th of an inch thick, and is moistened by a watery fluid, for which it has a greater attraction than the fat it contains. Each vesicle is from the 300th to the  $\frac{1}{800}$ th of an inch in diameter. When the fat vesicles exist in any number, their arrangement is generally lobular, with an investment of cellular tissue, which favours motion, and the distribution of

the bloodvessels. These enter the interlobular clefts, ramify through

Fig. 187.



Bloodvessels of Fat Vesicles.

A. Minute flattened fat-lobule, in which the vessels only are represented. a. The terminal artery. v. The primitive vein. b. The fat vesicles of one border of the lobule, separately represented. Magnified 100 diameters.—c. Plan of the arrangement of the capillaries on the exterior of the vesicles: more highly magnified.—(Todd and Bowman.)

their interior as a solid capillary network, occupy the angles formed by contiguous sides of the vesicles, and anastomose with one another at the points where these angles meet.

M. Raspail<sup>b</sup> affirms, that there is the most striking analogy between the nature of the adipous granules and that of the amylaceous grains. As in the case of fecula, each granule is pous composed of at least one integument, and an inclosed substance,

\* Mr. Paget, Brit. and For. Med. Review, July, 1842, p. 262.

b Chimie Organique, p. 183, Paris, 1833.

both of which are as slightly azoted as fecula; and both fecula and fat are equally inservient to the nutrition of the organs of development: whenever there is excess of life and activity, the fat is seen to disappear, and whenever there is rest, it accumulates in its reservoirs. If a portion of fat be examined, it is found to consist of an outer vesicle with strong membranous parietes, containing small adipous masses readily separable from each other, each invested with a similar, but slighter, vesicular membrane; and these, again, contain others still more minute, until ultimately we come to the vesicles that invest the adipous granules themselves. Each of these masses adheres, at some point of its surface, to the inner surface of the vesicle that incloses it by a hilum in the same manner as the grain of fecula. All the vesicles, but especially the outermost and strongest, have a reddish vascular network on their surface, the vessels of which augment in size as they approach the part where the vesicle is adherent, and there they open into one of the vessels of the larger vesicle that incloses them.

The arrangement of this tissue, as well as the quantity of fat, varies in different parts of the body. It is always found in the orbit, on the sole of the foot, and at the pulps of the fingers and toes. The subcutaneous cellular tissue, and that covering the heart, kidneys, &c., also generally contain it: but it is never met with in the eyelids, scrotum, or within the cranium.<sup>a</sup>

Fat is exhaled by the secretory vessels in a fluid state; but after it is deposited, it becomes more or less solid. According to the researches of Chevreul<sup>b</sup> and Braconnot, human fat is almost always of a yellow colour; inodorous, and composed of two portions; - the one fluid, and the other concrete, which are themselves composed, but in different proportions, of two new immediate principles, to which the former chemist gave the names elain or olein, and stearin, respectively. Subsequently, the organic elements of fat were considered to be stearin, margarin, and olein; the two former, which are solid when separate, being dissolved in the latter, at the ordinary temperature of the body. Recent chemistry has, however, shown, that the fat contained in the cells of the adipous tissue is composed of a base termed glycerin, with stearic and margaric acids, - stearin being esteemed a bi-stearite of glycerin; and olein or elain, an oleate of glycerin.º These proximate principles are sometimes seen spontaneously separated within the human fat vesicle. The stearin collects in the form of a small star on the inner surface of the membrane, as in

<sup>\*</sup> Geddings, art. Adipose Tissue, in Amer. Cyclop. of Pract. Medicine, part ii. p. 215. Philad. 1833; and Craigie, art. Adipose Tissue, in Cycl. Anat. and Physiol. part i. p. 56, Lond. 1835.

b Recherches Chimiques sur les Corps Gras, &c., Paris, 1823; W. T. Brande, art.

Fat, Cyclopæd. Anat. and Physiol. August, 1837, p. 231.

See page 32, vol. i.: and Thomson, Chemistry of Animal Bodies, p. 135, Edi

c See page 32, vol. i.; and Thomson, Chemistry of Animal Bodies, p. 135, Edinb., 1843.

Fig. 188.

Fat vesicles from an emaciated subject.

a. a. The cell-membrane. b. b. b. The solid portion collected as a star-like mass, with the elain in connection with it, but not filling the cell.—(Todd The cell-membrane. and Bowman.)

the marginal figure at b, b, b, the elain occupying the remainder of the vesicle, except where there is an unusually small quantity of fat, when a little aqueous fluid is seen interposed between the elain and the cell-membrane.

> It is probable, that chemical analysis would exhibit the fat to vary in different parts of the body, as its sensible properties are manifestly different. Sir Everard Home, a on loose analogies and inconclusive arguments, has advanced the opinion, that it is more than probable, that fat is formed in the lower portion of the intestines, and from thence is carried, through the medium of the circulating blood,

to be deposited in almost every part of the body. "When there is a great demand for it, as in youth, for carrying on growth, it is laid immediately under the skin, or in the neighbourhood of the abdomen. When not likely to be wanted, as in old age, it is deposited in the interstices of muscular fibres, to make up in bulk for the wasting of these organs." M. de Blainvilleb is of opinion, that fat is derived from venous blood, and that it is exhaled through the coats of the vessels. This opinion he founds on the mode in which the fat is distributed in the omenta along the course of the vein; and he affirms, that he has seen it flow out of the jugular vein in a dead elephant. But this last fact, as Lepelletier has judiciously remarked, proves nothing more than that the fat taken up by the absorbents, from the vesicles, in which it had been deposited by the exhalants - had been conveyed into the venous blood with other absorbed matters. It in no wise shows, that the venous blood is the pabulum of the secretion, or that the veins accomplish it.

The uses of the fat are both general and local. The great general use is, by some physiologists, conceived to be, - to serve as a provision in cases of wasting indisposition; when the digestive function is incapacitated for performing its due office, and emaciation is the consequence. In favour of this view, the rapidity with which fat disappears after slight abstinence has been urged, as well as the facts, connected with the torpidity of animals, which are always found to diminish in weight during this state. Professor Mangili, of Pavia, procured two marmots from the Alps, on the 1st of December. The larger weighed 25 Milanese ounces; the smaller only 22 th; on the 3d of January, the larger had lost 3ths of an ounce, and the smaller 17ths. On the 5th of February, the larger weighed only 22%; the smaller 21. Dr. Monro kept a hedgehog from the month of November to the month of March

<sup>\*</sup> Lect, on Comp. Anat. i, 468, Lond. 1814, and vol. vi., Lond. 1828; and Philos. Transact. 1821, p. 34. b De l'Organisation des Animaux, &c., Paris, 1825. Physiologie Médicale et Philosophique, ii. 496, Paris, 1832.

following, which lost, in the meanwhile, a considerable portion of its weight. On the 25th of December, it weighed 13 ounces and 3 drachms; on the 6th of February, 11 ounces and 7 drachms; and on the 8th of March, 11 ounces and 3 drachms. The loss was 13 grains daily.4 The local uses of fat are chiefly of a physical character. On the sole of the foot it diminishes the effects of pressure, and its use is the same on the nates: in the orbit, it forms a kind of cushion, on which the eyeball moves with facility; and when in certain limits it gives that rotundity to the frame, which we are accustomed to regard as symmetry. Dr. Fletcher, indeed, considers its principal use to be, to fill up interstices, and thus to give a pleasing contour to the body. In another place, it was observed, that fatty substances are bad conductors of caloric; and hence that it may tend to preserve the temperature of the body in cold seasons; a view which is favoured by the fact, that many of the arctic animals are largely supplied with fat beneath the common integuments; and it has been affirmed, that fat people generally suffer less than lean from the cold of winter.

It is obviously impracticable to estimate, accurately, the total quantity of fat in the body. It has been supposed that, in an adult male of moderate size, it forms to the whole weight; but it is doubtful whether we ought to regard this as even an approximation; the data being so inadequate. In some cases of polysarcia or obesity, the bulk of the body has been enormous. The case of a girl is detailed, who weighed 256 pounds, when only four years old. A man of the name of Bright, at Malden, England, weighed 728 pounds; and the celebrated Daniel Lambert, of Leicester, England, weighed 739 pounds a little before his death, which occurred in the fortieth year of his age.d The circumference of his body was three yards and four inches; of his leg one yard and one inch. His coffin was six feet four inches long; four feet four inches wide; and two feet four inches deep. The public journals of this countrye have recorded the death of a Mr. Cornelius, who weighed 720 pounds. Dr. Elliotsonf says he saw a female child, but a year old, who weighed sixty pounds. She had begun to grow fat at the end of the third month. In these cases, the specific gravity of the body may be much less than that of water.s It is said, that some time ago there was a fat lighterman, on the river Thames, " who had fallen overboard repeatedly, without any farther inconvenience than that of a good ducking; since though

\* Fleming's Philosophy of Zoology, ii. 59, Edinb. 1822.

b Rudiments of Physiology, part iii., by Dr. Lewins, p. 71, Edinb. 1837.

Philos. Transact. No. 185.

d Good's Study of Medicine, Class vi. Ord. 1. Gen. 1. Sp. 1.

e Philadelphia Public Ledger, Oct. 4, 1841.

Human Physiology, Lond. 1841, P. i. 301; and Fletcher's Rudiments of Physio-

logy, Edinburgh, 1835, P. i. p. 123.

s For various cases of great obesity, see Choulant, in art. Fettbereitung, in Pierer, op. citat. B. 3. s. 53; A. L. Richter, in Encyclopädisches Wörterbuch der Medicinischen Wissenschaften, Band. i. s. 733, art. Adiposis, Berlin, 1828; and Fletcher's Rudiments of Physiology.

he knew nothing whatever of the art of swimming, he always continued to flounder about like a firkin of butter, till he was picked

up."a

In some of the varieties of the human family we meet with singular adipous deposits. In the Bosjesman female vast masses of fat accumulate on the buttocks, which give them the most extravagant appearance. The projection of the posterior part of the body, in one subject, according to Barrow, b measured five inches and a half from a line touching the spine. "This protuberance," he remarks, "consisted of fat, and when the woman walked, had the most ridiculous appearance imaginable, every step being accompanied with a quivering and tremulous motion, as if two masses of jelly were attached behind." The "Hottentot Venus," who had several projections, measured more than nineteen inches around the haunches; and the projection of the hips exceeded 63 inches. Dr. Somerville found on dissection, that the size of the buttocks arose from a vast mass of fat, interposed between the integuments and muscles, which equalled four fingers' breadth in thickness. It is singular, that, according to the statement of this female, which is corroborated by the testimony of Mr. Barrow, the deposition does not take place till the first pregnancy. Pallasd has described a variety of sheep - the ovis steatopyga or "fat-buttocked," - which is reared in immense flocks by the pastoral tribes of Asia. In it, a large mass of fat covers the nates and occupies the place of the tail. The protuberance is smooth beneath, and resembles a double hemisphere, when viewed behind; the os coccygis or rump-bone being perceptible to the touch in the notch between the two. They consist merely of fat; and, when very large, shake in walking like the buttocks of the female Bosjesman. Mr. Lawrencee remarks, that there are herds of sheep in Persia, Syria, Palestine and some parts of Africa, in which the tail is not wanting as in the ovis steatopyga, but retains its usual length, and becomes loaded with fat.

In the view of Liebig, the abnormous condition, which causes the deposition of fat in the animal body, depends on a disproportion between the quantity of carbon in the food, and that of the oxygen absorbed by the skin and lungs. In the normal condition, the quantity of carbon given out is exactly equal to that which is taken in the food, and the body experiences no increase of weight from the accumulation of substances containing much carbon and no azote; but if the supply of highly carbonized food be increased, then the normal state can only be preserved, by exercise and labour, through which the waste of the body is increased, and the supply of oxygen accumulated in the same proportion. The production of fat, Liebig maintains, is always a consequence of a

<sup>&</sup>lt;sup>2</sup> Fletcher, op. citat. p. 71. <sup>c</sup> Medico-Chirurgical Transactions, vii. 157. <sup>d</sup> Spicilegia Zoologica, fasc. xi. p. 63.

Lectures on Physiology, Zoology, &c. p. 427, Lond. 1819.

Animal Chemistry, Webster's edit. p. 85, Cambridge, Mass., 1842.

deficient supply of oxygen, for oxygen is absolutely indispensable for the dissipation of the excess of carbon in the food. "This excess of carbon, deposited in the form of fat, is never seen in the Bedouin or in the Arab of the desert, who exhibits with pride to the traveller his lean, muscular, sinewy limbs, altogether free from fat; but in prisons and jails it appears as a puffiness in the inmates, fed, as they are, on a poor and scanty diet: it appears in the sedentary females of oriental countries; and finally it is produced under the well known conditions of the fattening of domestic animals." In accordance, too, with his views of animal heat, already referred to, Liebig considers that in the formation of fat there is a new source of heat. The oxygen set free in the action is given out in combination with carbon and hydrogen; and whether this carbon and hydrogen proceed from the substance that yields the oxygen, or from other compounds, still there must have been generated by this formation of carbonic acid or water, as much heat as if an equal weight of carbon or hydrogen had been burned in air or in oxygen gas. Whether the view of Liebig be admitted or not it is certain that the circumstances, which favour obesity, are absence of activity and of excitement of all kinds; hence, for the purpose of fattening animals in rural economy, they are kept in entire darkness, to deprive them of the stimulus of light, and to favour sleep and muscular inactivity. Castration - by abolishing one kind of excitability - and the time of life at which the generative functions cease to be exerted, especially in the female, are favourable to the same result.

# d. Exhalation of the Marrow.

A fluid, essentially resembling fat, is found in the cavity of long bones, in the spongy tissue of short bones, and in the areolæ of bones of every kind. This is the marrow. The secretory organ is the very delicate membrane which is perceptible in the interior of the long bones, lining the medullary cavity, and sending prolongations into the compact substance, and others internally, which form septa and spaces for the reception of the marrow. The cells, thus formed, are distinct from each other. From the observations of Howship, it would seem probable, that the oil of bones is deposited in longitudinal canals, that pass through the solid substance of the bone, and through which its vessels are transmitted. This oil of bones is the marrow of the compact structure, the latter term being generally restricted to this secretion when contained in the cavities of long bones; that which exists in the spongy substance being termed, by some writers, the medullary juice. The medullary membrane, called also the internal periosteum, consists chiefly of bloodvessels ramifying on an extremely delicate cellular tissue, in which nerves may likewise be traced.

Berzelius examined marrow obtained from the thigh-bone of an ox, and found it to consist of the following constituents: — pure

<sup>\*</sup> Medico-Chirurg. Trans. vii. 393.

adipous matter, 96; skins and bloodvessels, 1; albumen, gelatin,

extractive, peculiar matter, and water, 3.ª

The marrow is one of the corporeal components, of whose use we can scarcely offer a plausible conjecture. It has been supposed to render the bones less brittle; but this is not correct, as those of the fœtus, which contain little or no marrow, are less brittle than those of the adult; whilst the bones of old persons, in which the medullary cavity is extremely large, are more brittle than those of the adult. It is possible that it may be placed in the cavities of the bones, — which would otherwise be so many vacant spaces, — to serve the general purposes of the fat, when it is required by the system. The other hypotheses, that have been entertained on the subject, are not deserving of notice.

### e. Synovial Exhalation.

Within the articular capsules, and the bursæ mucosæ, - which have been described under the head of muscular motion, - a fluid is secreted, which is spread over the articular surfaces of the bones, and facilitates their movements. Haversb considered this fluid to be secreted by synovial glands, - for such he conceived the reddish cellular masses to be, that are found in certain articulations. Hallere strangely regarded the synovia as the marrow, which had transuded through the spongy extremities of the bones; but, since the time of Bichat, every anatomist and physiologist has ascribed' it to the exhalant action of the synovial membrane, which strongly resembles the serous membranes in form, structure, and functions, and whose folds constitute the projections, which Havers mistook for glands.d This membrane exists in all the movable articulations, and in the channels and sheaths in which the tendons play. The generality of anatomists regard the articular capsules as shut sacs; the membrane being reflected over the incrusting cartilages. Magendie, however, affirms, that he has several times satisfied himself, that the membranes do not pass beyond the circumference of the cartilages. From the inner surface of these membranes the synovia is exhaled, in the same manner as in other serous cavities.

Marguerone analyzed the synovia, obtained from the posterior extremity of the ox, and found it to consist of fibrous matter, 11.86; albumen, 4.52; chloride of sodium, 1.75; soda, 0.71; phosphate of lime, 0.70; and water, 80.46. M. Donné says it is always alkaline in health; but in certain diseases it sometimes becomes acid.

<sup>2</sup> Thomson, Chemistry of Animal Bodies, p. 256, Edinburgh, 1843.

b De Ossibus, serm iv. c. 1; and Osteologia Nova, Lond. 1691.

Element. Physiol. iv. 11.

Béclard, Elements of General Anat. by Togno, p. 140.

Annales de Chimie, xiv. 123; also, Thomson, Chemistry of Animal Bodies, p. 505,
 Edinburgh, 1843.

Journal Hebdomad. Février, 1834.

f. Exhalation of the Colouring matter of the Skin and other parts.

The nature of the exhalation, which constitutes the colouring matter of the rete mucosum, has already engaged our attention, when treating of the skin, under the Sense of Touch. It is presumed to be exhaled by the vessels of the skin, and to be deposited beneath the cuticle, so as to communicate the colours that characterize the different races. Such are regarded as the secretory organs by most anatomists and physiologists; but Gaultier, whose researches into the intimate constitution of the skin have gained him much celebrity, is of opinion, that it is furnished by the bulbs of the hair; and he assigns, as reasons for this belief, that the negro, in whom it is abundant, has short hair; that the female, whose hair is more beautiful and abundant than that of the male, has the fairest skin; and that when he applied blisters to the skin of the negro, he saw the colouring matter oozing from the bulbs of the hair, and deposited at the surface of the rete mucosum. Breschet, again, describes a particular chromatogenous apparatus, for producing the colouring matter, which is composed of a glandular or secreting parenchyma, situate a little below the papillæ, and having excretory ducts, which pour the colouring principle on the surface of the true skin. This mingles with the soft and diffluent mucous matter, the admixture producing the "pretended reticular body of Malpighi," and the epidermis or cuticle. He describes particular organs, or a "blennogenous apparatus," for the secretion of this mucous matter. They consist of a glandular parenchyma, or organ of secretion, in the substance of the true skin; and of excretory ducts, which issue from the latter and deposit the mucous matter between the papillæ.

The composition of this pigment cannot be determined with precision, owing to its quantity being too small to admit of examination. Chlorine deprives it of its black hue, and renders it yellow. A negro, by keeping his foot for some time in water impregnated with this gas, deprived it of its colour, and rendered it nearly white; but, in a few days, the black colour returned with its former intensity. This experiment was made with similar results on the fingers of a negro. Blumenbache thought, that the mucous pigment was formed chiefly of carbon; and the notion has received

favour with many.d

The colour appears to be owing to pigment cells, of which the pigmentum nigrum of the eye is wholly composed. They are considered to exhibit, usually, the original form of the cell with little alteration. On the choroid coat of the eye they form a kind of pavement, and have somewhat of a polyhedral shape. In the human skin, they are scattered through the ordinary epidermic

<sup>\*</sup> Recherches sur l'Organisation de la Peau de l'Homme, &c., Paris, 1809, and 1811.

Nouvelles Recherches sur la Structure de la Peau, Paris, 1835.

c Instit. Physiol. § 274; and Elliotson's translation, 4th edit. Lond. 1828.

d Carpenter, op. cit. § 616.

cells, and the colour of the skin is determined by that of their contents.<sup>a</sup> Pigment granules are amongst the most minute structures of the body, being not more than  $\frac{1}{2\sqrt{6}}$  of the part of an inch in their largest diameter, and about one-fourth as much in thickness.<sup>b</sup>

The uses of the pigment of the skin — as well as of that which lines the choroid coat of the eye, the posterior surface of the iris, and of the ciliary processes — are detailed in another place.

### g. Areolar Exhalation.

Under this term, Adelond has included different recrementitial secretions effected within the organs of sense, or in parenchymatous structures, — as the aqueous, crystalline, and vitreous humours of the eye, and the liquor of Cutugno, all of which have already engaged attention; the exhalation of a kind of albuminous, reddish, or whitish lymph into the interior of the lymphatic ganglions, and into the organs, called, by Chaussier, glandiform ganglions, and by Béclard, sanguineous ganglions; — namely, the thymus, thyroid, supra-renal capsules, and spleen. We know but little, however, of the fluids formed in these various parts. They have never been analyzed, and their uses are inappreciable.

By some physiologists, a fluid is supposed to be exhaled from the inner coat of the arterial, venous and lymphatic vessels. We are unaware, however, of the nature of this fluid. Its chief use is presumed to be to lubricate the interior of the vessel, and to prevent adhesion between it and the fluid circulating within it.

#### 2. EXTERNAL EXHALATIONS.

# a. Cutaneous Exhalation or Transpiration.

A transparent fluid is constantly exhaled from the skin, which is generally invisible in consequence of its being converted into vapour as soon as it reaches the surface; but, at other times, owing to augmentation of the secretion, or to the air being loaded with humidity, it is apparent on the surface of the body. When invisible, it is called the insensible transpiration or perspiration; when perceptible, sweat. In the state of health, according to Thénard, this fluid reddens litmus paper; yet the taste is rather saline—resembling that of common salt—than acid. Wagner, indeed, affirms that it generally shows alkaline reaction; and at other times does not affect vegetable blues; but the sweat of many parts of the body,—the armpits for example,—is said always to react like an alkali. Allusion has already been made

a Carpenter, Human Physiology, § 616, Lond. 1842.

b Henle; and Paget, in Brit. and For. Med. Rev., July, 1842, p. 266.

See, on the subject of the Pigments, Burdach's Physiologie als Erfahrungswissenschaft, v. 176, Leipz. 1835.

d Physiologie de l'Homme, 2de édit. tom. iii. 483, Paris, 1829.

· Traité de Chimie, tom. iii.

Elements of Physiology, by R. Willis, § 204, Lond. 1842.

to the views of M. Donné, a who considers, that the external acid,

and the internal alkaline membranes of the human body represent the two poles of a pile, the electrical effects of which are appreciable by the galvanometer. The smell of the perspiration is peculiar, and becomes almost insupportable when concentrated, and especially when subjected to distillation. The fluid is composed, according to Thénard, of much water, a small quantity of acetic acid, chloride of sodium, and perhaps of potassium, a very little earthy phosphate, a trace of oxide of iron, and an inappreciable quantity of animal matter. Berzeliusb regards it as water, holding in solution the chlorides of potassium and sodium, lactic acid, lactate of soda, and a little animal matter; and Anselmino, c as consisting of a solution of osmazome, chlorides of sodium and calcium, acetic acid and an alkaline acetate, salivary matter, sulphates of soda and potassa, and calcareous salts, with mucus, albumen, sebaceous humour, and gelatin in variable proportions. Raspaild regards the sweat as an acid product of the disorganization of the skin. After evaporation upon a clean glass plate, fragments of epidermic cells are generally observed in it, and crystals are left behind, which are those of its contained salts.

In a memoir presented to the Académie Royale des Sciences, of Diapnogenous Apparatus from the Palm of the Hand. Paris, MM. Breschet and Roussel de Paris, MM. Breschet and Roussel de Vauzème have shown, that there exists in the skin an apparatus for the secretion of the sweat, consisting of a secretion of the sweat, consisting of a which the gland is imbedded.—(Wagner.) glandular parenchyma, which se-

Fig. 189.

cretes the liquid, and of ducts, which pour it on the surface of the body. These ducts are said to be arranged spirally, and to open very obliquely under the scales of the epidermis. To this appara-

a Journal Hebdomad., Février, 1834.

b Medico-Chir, Trans. iii. 256, and Bostock, ibid. xiv. 424.

Lepelletier, Physiologie Médicale et Philosophique, ii. 452, Paris, 1832. Also, Thomson, Chemistry of Animal Bodies, p. 523, Edinburgh, 1843.

d Chimie Organique, p. 505, Paris, 1832.

tus they apply the epithet "diapnogenous:" the ducts they call "sudoriferous or hidrophorous." The marginal figure, (Fig. 189,)

exhibits the apparatus magnified.

Numerous experiments have been instituted for the purpose of discovering the quantity of transpiration that takes place in a given time. Of these, the earliest were by Sanctorius, - for which he is more celebrated than for any other of his labours, - after whom the transpiration was called Perspirabile Sanctorianum. For thirty years, this indefatigable experimentalist weighed daily, with the greatest care, his solid and liquid ingesta and egesta, and his own body, with the view of deducing the loss sustained by the cutaneous and pulmonary exhalations. He found, that every twenty-four hours, his body returned sensibly to the same weight, and that he lost the whole of the ingesta; - five-eighths by transpiration, and three-eighths by the ordinary excretions. For eight pounds of ingesta, there were only three pounds of sensible egesta, which consisted of forty-four ounces of urine, and four of fæces. It is lamentable to reflect, that so much time was occupied in the attainment of such insignificant results. The self-devotion of Sanctorius gave occasion, however, to the institution of numerous experiments of the same kind; as well as to discover the variations in the exhalation, according to age, climate, &c. The results of these have been collected by Haller, but they afford little instruction; especially as they were directed to the transpiration in general, without affording us any data to calculate the proportion exhaled from the lungs compared with that constantly taking place by the cutaneous surface. Rye,d who dwelt in Cork, lat. 51° 54', found, in the three winter months - December, January, and February — that the quantity of urine was 3937 ounces; of the perspiration, 4797: in the spring months - March, April, and May — the urine amounted to 3558; the perspiration to 5405; in the summer months - June, July, and August - the urine amounted to 3352; the perspiration to 5719; and in the three autumnal months - September, October, and November the quantity of urine was 3369; that of the perspiration 4471. The daily average estimate, in ounces, was as follows: -

			MAN TO MAN			Urine.	Perspiration.	
Winter,	-		-	ob l-con	mile ly	427		53
Spring,		Garana	MATE IN	July Soul	aurism d	40	1941	60
Summer,	100	Virginia - California		1002.00	200	37	100	63
Autumn,	-			N. P. Sand		37	-	50

thus making the average daily excretion of urine, throughout the year to be a little more than 39 ounces; and of the transpiration, 56 ounces. Keill, on the other hand, makes the average daily perspiration, 31 ounces; and that of the urine 38; the weight of

Elem. Physiol. xii. 2, 10.

<sup>&</sup>lt;sup>2</sup> Breschet, Nouvelles Recherches sur la Structure de la Peau, Paris, 1835; also, Plumbe, on Diseases of the Skin, p. 12, Lond. 1837.

b Ars Sanctorii de Staticà Medicina, cum Comment. Martini Lister, Logd. Bat. 1711.

<sup>&</sup>lt;sup>4</sup> Rogers, on Epidem. Diseases, Appendix, Dubl. 1734. <sup>e</sup> Tentamina Medico-Phys. — Appendix, Lond. 1718.

the fæces being 5 ounces, and that of the solid and liquid ingesta, 75 ounces. His experiments were made at Northampton, England, lat. 52° 11'. Bryan Robinsona found, as the result of his observations in Ireland, that the ratio of the perspiration to the urine was, in summer, as 5 to 3; in winter, as 2 to 3; whilst in April, May, October, November, and December, they were nearly equal. In youth, the ratio of the perspiration to the urine was as 1340 to 1000; in the aged, as 967 to 1000. Hartmann, when the solid and liquid ingesta amounted to 80 ounces, found the urine discharged 28 ounces; the fæces 6 or 7 ounces; and the perspirable matter, 45 or 46 ounces. De Gorter, b in Holland, when the ingesta were 91 ounces, found the perspiration to amount to 49 ounces; the urine to 36; and the fæces to 8. Dodart asserts, that in France, the ratio of the perspiration to the fæces, is as 7 to 1; and to the whole egesta as 15 to 12 or 10. The average perspiration in the twenty-four hours, he estimates at 33 ounces and 2 drachms; and Sauvages, in the south of France, found that when the ingesta were 60 ounces in the day, the transpiration amounted to 33 ounces; the urine to 22; and the fæces to 5. But most of these estimates were obtained in the cooler climates, - the "regiones boreales," - as Hallerd has, not very happily, termed them.

According to Lining, whose experiments were made in South Carolina, lat. 32° 47′, the perspiration exceeded the urine in the warm months; but in the cold, the latter had the preponderance. The following table gives the average daily proportion of the urine and perspiration, for each month of the year, in ounces, —

as quoted by Haller.

sundinoris.				Urine.		Perspiration.		
December,	and a contract of	III PI	nils-	70.81	Drive.	27111	42.55	
January,	discount.	1-11	S COUNTY	72.43	10810	di-to	39.97	
February,				77.86		10600	37.45	
March,	ACCEPTED IN	-	THE PERSON NAMED IN	70.59		1000	43.23	
April,	Septemp	III SELL	- utaru	59.17	DB DE	1814791	47.72	
May,	DI HOUR	AND FIRM	de-pa	56-15	12 (253)	HELDE	58-11	
June,	monte del	Passile	CHARLES	52.90	m-le	mix11	71.39	
July,			Spate.	43.77	me no	7 510	86.41	
August,	- nan			55-41	-		70.91	
September,	no sum	40000	naime	40 60	2000	WHI.	77.09	
October,	21-11		100	47.67		-	40.78	
November,	miner U		16 300	63.16	110	Las I	40.97	

After the period at which Haller wrote, no experiments of any moment were adopted for appreciating the transpiration. Whenever trials were instituted, the exhalation from both the skin and the lungs was included in the result, and no satisfactory means were adopted for separating them, until Lavoisier and Séguinf made their celebrated experiments. Séguin inclosed himself in a bag of gummed taffeta, which was tied above the head, and had an aperture, the edges of which were fixed around the mouth by a mix-

a Dissertation on the Food and Discharges of Human Bodies, Dublin, 1748.

ь De Perspiratione Insensibili, Lugd. Bat. 1736.

Mémoir. de l'Acad. des Sciences, ii. 276.

d Op. citat.

<sup>·</sup> Philos. Transact. for 1743 and 1745.

<sup>1</sup> Mémoir. de l'Académ. des Sciences de Paris, Paris, 1777 and 1790.

ture of turpentine and pitch. By means of this arrangement, the pulmonary transpiration alone escaped into the air. To estimate its quantity, it was merely necessary for M. Séguin to weigh himself in the sac, by a very delicate balance, at the commencement and termination of the experiment. By repeating the experiment out of the sac, he determined the total quantity of the transpired fluid; so that, by deducting from this the quantity of fluid exhaled from the lungs, he obtained the amount of the cutaneous transpiration. He, moreover, kept an account of the food, which he took; of the solid and liquid egesta; and as far as he was able, of

every circumstance that could influence the transpiration.

The results — as applicable to Paris — at which Lavoisier and Seguin arrived, by a series of well-devised and well-conducted experiments, were the following: - First. Whatever may be the quantity of food taken, or the variations in the state of the atmosphere, the same individual, after having increased in weight by the whole quantity of nourishment taken, returns daily, after the lapse of twenty-four hours, to nearly the same weight as the day before; - provided he be in good health; his digestion perfect; that he is not fattening, or growing; and avoids all kind of excess. Secondly. If, when all other circumstances are identical, the quantity of food varies; or if—the quantity of food being the same the effects of transpiration differ; the quantity of the excrements augments or diminishes, so that every day at the same hour, we return nearly to the same weight; proving, that when digestion goes on well, the causes, which concur in the loss or excretion of the food taken in, afford each other mutual assistance; - in the state of health one charging itself with what the other is unable to accomplish. Thirdly. Defective digestion is one of the most direct causes of the diminution of transpiration. Fourthly. When digestion goes on well, and the other causes are equal, the quantity of food has but little effect on the transpiration. Séguin affirms, that he has very frequently taken at dinner two pounds and a half of solid and liquid food; and at other times, four pounds; yet the results, in the two cases, differed but little from each other; provided only, that the quantity of fluid did not vary materially in the two cases. Fifthly. Immediately after dinner, the transpiration is at its minimum. Sixthly. When all other circumstances are equal, the loss of weight, induced by insensible transpiration, is at its maximum during digestion. The increase of transpiration during digestion compared with the loss sustained when fasting, is, on an average, 23 grains per minute. Seventhly. When circumstances are most favourable, the greatest loss of weight, caused by insensible transpiration, was, according to their observations, 32 grains per minute; consequently 3 ounces, 2 drachms and 48 grains poids de marc, per hour; and 5 pounds in twenty-four hours; under the calculation, that the loss is alike at all hours of the day, which is not, however, the fact. Eighthly. When all the accessory circumstances are least favourable, provided only that digestion

is properly accomplished, the smallest loss of weight is 11 grains per minute; consequently, 1 ounce, 1 drachm and 12 grains per hour; and 1 pound, 11 ounces and 4 drachms in the twenty-four hours. Ninthly. Immediately after eating, the loss of weight caused by the insensible perspiration, is 101 grains per minute, during the time at which all the extraneous causes are most unfavourable to transpiration; and 1910 grains per minute, when these causes are most favourable and the internal causes are alike. "These differences," says M. Séguin, "in the transpiration after a meal, according as the causes influencing it are more or less favourable, are not in the same ratio with the differences, observed at any other time when the other circumstances are equal; but we know not how to account for the phenomenon." Tenthly. The cutaneous transpiration is immediately dependent both on the solvent virtue of the circumambient air, and on the power possessed by the exhalants of conveying the perspirable fluid as far as the surface of the skin. Eleventhly. From the average of all the experiments, it seems, that the loss of weight caused by the insensible transpiration is 18 grains per minute; and that, of these 18 grains, 11, on the average, belong to the cutaneous transpiration, and 7 to the pulmonary. Twelfthly. The pulmonary transpiration, compared with the volume of the lungs, is much more considerable than the cutaneous, compared with the surface of the skin. Thirteenthly. When every other circumstance is equal, the pulmonary transpiration is nearly the same before and immediately after a meal; and if, on an average, the pulmonary transpiration be 171 grains per minute before dinner, it is 1770 grains after dinner. Lastly. Every other intrinsic circumstance being equal, the weight of the solid excrements is least during winter.

Although these results are probably fairly deduced from the experiments; and the experiments themselves were almost as well conceived as the subject admits of, we cannot regard the estimates as more than approximations. Independently of the fact, that the envelope of taffeta must necessarily have retarded the exhalation by shutting off the air, and caused more to pass off by pulmonary transpiration, the perspiration must incessantly vary according to circumstances within and without the system: some individuals, too, perspire more readily than others: and this is dependent, as we have seen, upon climate and season, — and likewise upon the quantity of fluid received into the digestive organs. From all these and other causes, Bichat is led to observe, that the endeavour to determine the quantity of the cutaneous transpiration is as vain. as to endeavour to specify what quantity of water is evaporated every hour, by a fire, the intensity of which is varying every instant. To attempt, however, the solution of the problem, experiments were like wise undertaken by Cruikshank, and by Abernethy. Their plan consisted in confining the hand, for an hour, in an air-

\* Experiments on the Insensible Perspiration, p. 5, Lond. 1795.

tight glass jar, and collecting the transpired moisture. Mr. Abernethy, having weighed the fluid collected in the glass, multiplied its quantity by 38½, the number of times, he conceived the surface of the hand and wrist to be contained in the whole cutaneous surface. This gave 21 pounds, as the quantity exhaled from the skin in the twenty-four hours, upon the supposition, that the whole surface perspires to an equal extent. These experiments have been repeated by Dr. William Wood, a of Newport, England, with some modifications. He pasted around the mouth of a jar one extremity of a bladder, the ends of which were cut away, and the hand being passed through the bladder into the jar, the other extremity was bound to the wrist with a ligature, not so tight, however, as to interfere, in any degree, with the circulation. The exact weight of the jar and bladder had previously been ascertained. During the experiment, cold water was applied to the outer surface of the jar, to cause the deposition of the fluid accumulated within. The result of his experiments was as follows:

Exp.	Time of day.		Temperature in apartment.			Pulse per minute.			Fluid collected in an hour.		
1.	Noon.			66°	7 -17	1141	84	1	14	32 grs.	
2	Do.			66	19	- 1	78		1	32	
3	Do.		1012	66		1.00	78	1		26	
4.	Do.	1.0	HAI	61	-	- 9	84			32	
5.	9 P. M.			62		-	80	1	-	26	
6.	Do.	13	76	62		19-5	75	19.		23	
	Mean.			63.8			79.8			28.5	

The next thing was to estimate the proportion, which the surface of the hand and wrist bears to the whole surface of the body. Abernethy reckoned it as 1 to 381, whilst Cruikshank computed it as 1 to 60! Dr. Wood does not adopt the estimate of either. He thinks, however, that the estimate of the former as regards the surface of the hand and wrist, which he makes seventy square inches, is near the truth, having found it to correspond both with his own measurements, and the reports of glovers. Mr. Abernethy's estimate of the superficial area of the whole body - 2700 square inches, or above eighteen square feet, he properly regards as too high. Perhaps the most general opinion is, that it amounts to sixteen square feet, or 2304 square inches; but Haller did not think it exceeded thirteen square feet or 2160 square inches. Dr. Wood adopts the former of these, and is disposed to think, that the proportion of the surface of the hand and fingers, taken to the extremity of the bone of the arm, does not fall short of 1 to 2, which if we adopt the ratio of the quantity, that he found transpired per hour, gives, for the whole body, about forty-five ounces, or nearly four pounds troy in the twenty-four hours. This is considerably above the result of the experiments of either Séguin, or Abernethy; yet, on reviewing the experiments, Dr. Wood is not disposed to think it far from the truth.

Upwards of fifty years ago, Dr. Dalton, of Manchester, under-

An Essay on the Structure and Functions of the Skin, &c., Edinb. 1832.

took a series of experiments similar to those of Sanctorius, Keill, Hartmann and Dodart. The first series of experiments he made upon himself, in the month of March, for fourteen days in succes-The aggregate of the articles of food consumed in this time was as follows, - bread, 163 ounces avoirdupois; oaten cake, 79 ounces; oatmeal, 12 ounces; butcher's meat, 542 ounces; potatoes, 130 ounces; pastry, 55 ounces; cheese, 32 ounces; - Total of solid food, 5252 ounces; averaging 38 ounces daily: - of milk, 435½ ounces; beer, 230 ounces; tea, 76 ounces; — Total, 741½, averaging 53 ounces of fluid daily. The daily consumption was, consequently, 91 ounces; or nearly six pounds. During the same period, the total quantity of urine passed was 680 ounces; and of fæces, 68 ounces — the daily average being, — of urine, 48½ ounces; of fæces, 5 ounces; making 531. If we subtract these egesta from the ingesta, there will remain 37½ ounces, which must have been exhaled by the cutaneous and pulmonary transpirations, on the supposition that the weight of the body remained stationary. To test the influence of difference of seasons, Dr. Dalton resumed his investigations, in the month of June of the same year. results were as might have been anticipated, - a less consumption of solids and a greater of fluids; a diminution in the evacuations and an increase in the insensible perspiration. The average of solids consumed per day, was 34 ounces; of fluids, 56 ounces; - total, 90 ounces; the daily average of the evacuations - urine, 42 ounces; fæces, 4½, - leaving a balance of nearly 44 ounces, for the daily loss by perspiration, or one-sixth more than during the cooler season. He next varied the process, with the view of obtaining the quantity of perspiration, and the circumstances attendant upon it more directly. He procured a weighing beam, that would turn with one ounce. Dividing the day into periods of four hours in the forenoon, four or five hours in the afternoon, and nine hours in the night - or from ten o'clock at night to seven in the morning, he endeavoured to find the perspiration corresponding to these periods respectively. He weighed himself directly after breakfast, and again before dinner, observing neither to take nor part with any thing in the interval, except what was lost by perspiration. The difference in weight indicated such loss. same course was followed in the afternoon and in the night. This train of experiments was continued for three weeks in November. The mean hourly losses by transpiration were; -in the morning, 1.8 ounce avoirdupois; - afternoon, 1.67 ounce; night, 1.5 ounce. During twelve days of this period, he kept an account of urine corresponding in time with perspiration. The ratio was as 46 to 33.

From the whole of his investigations on this subject, Dr. Dalton concludes; — that of six pounds of aliment taken in the day, there appears to be nearly one pound of carbon and azote together, the remaining five pounds are chiefly water, which seems necessary as a vehicle to introduce the other two elements into the circula-

Manchester Memoirs, vol. v.

tion, and also to supply the lungs and membranes with moisture; that very nearly the whole quantity of food enters the circulation, for the fæces constitute only 1/18th part, and of these a part — bile — must have been secreted; that one great portion is thrown off by the kidneys, — namely, about half of the whole weight taken, but probably more or less according to climate, season, &c.; that another great portion is thrown off by means of insensible perspiration, which may be subdivided into two portions, one of which passes off by the skin — amounting to one-sixth part, and the other five-sixths are discharged from the lungs in the form of car-

bonic acid, and of water or aqueous vapour.

Since the time of Lavoisier and Séguin, Dr. Edwards has made some experiments, for the purpose of illustrating the effect produced upon cutaneous transpiration by various circumstances to which the body is subjected. His first trials were made on coldblooded animals, in which the cutaneous transpiration can be readily separated from the pulmonary, owing to the length of time, that they are capable of living without respiring. All that is necessary is to weigh the animal before and after the experiment, and to make allowance for the ingesta and egesta. In this way he discovered, that the body loses successively less and less in equal portions of time; that the transpiration proceeds more rapidly in dry than in moist air; in the extreme states nearly in the proportion of 10 to 1; that temperature has, also, considerable influence, - the transpiration at 68° of Fahrenheit, being twice as much; and, at 104°, seven times as much as at 32°. He likewise found, that frogs transpire, whilst they are in water, as is shown by the diminution which they experience while immersed in that fluid, and by the appearance of the water itself, which becomes perceptibly impregnated by the matter excreted by the skin. In warm-blooded animals, he found, as in the coldblooded, the transpiration become less and less in proportion to the quantity of fluid evaporated from the body; and he observed the same difference between the effects of moist and dry air, and between a high and a low temperature. The effects of these agents were essentially the same on man as on animals. He found, that the transpiration was more copious during the early than the latter part of the day; that it was greater after taking food; and, on the whole, appeared to be increased during sleep.b

Whenever the fluid, which constitutes the insensible transpiration, does not evaporate, owing to the causes referred to at the commencement of this article, it appears on the surface in the form of sensible perspiration or sweat. It has been supposed by some physiologists, that the insensible and sensible perspirations are two distinct functions. Such appears to be the opinion of Haller, and

b For various estimates relative to the quantity of the cutaneous transpiration, see

Burdach's Physiologie als Erfahrungswissenschaft, v. 196, Leipz. 1835.

Sur l'Influence des Agens Physiques, Paris, 1822; or Hodgkin and Fisher's translation, Lond. 1832; see, also, an analytical review of the same, by the author of this work, in the Amer. Journ. of the Med. Sciences, for May, 1834.

of Edwards, who regards the former as a physical evaporation,—the latter as a vital transudation or secretion, but no sufficient reason seems to exist, why we should not regard them as different degrees of the same function. It is, indeed, affirmed, that the sweat is generally less charged with carbonic acid than the vapour of transpiration; and that it is richer in salts, which are deposited on the skin, and are sometimes seen in the form of white flocculi;

but our knowledge on this matter is vague.

Particular parts of the body perspire more freely, and sweat more readily than others. The forehead, armpits, groins, hands, feet, &c., exhibit evidences of this most frequently; some of these, perhaps, owing to the fluid, when exhaled, not evaporating readily, - the contact of air being impeded. It is presumed, likewise, that the sweat has not every where the same composition. Its odour certainly varies in different parts of the body. In the armpits and feet it is more acid: in the violent sweats, accompanying acute rheumatism, this acidity always attracts attention; and in the groins, its odour is strong and rank. It differs, too, greatly in individuals, and especially in the races. In the red-haired, it is said to be unusually strong; and in the negro, during the heat of summer, it is alliaceous and overwhelming. By cleanliness, the red-haired can obviate the unpleasant effect, in a great measure, by preventing undue accumulation in the axilla, groins, &c.; but no ablution can remove the odour of the negro, although cleanliness can detract from its intensity. Each race appears to have its characteristic scent; and, according to Humboldt, the Peruvian Indian, whole smell is highly developed by education, can distinguish the European, the American Indian, and the negro, in the middle of the night, by this sense alone. Some physiologists have doubted, whether this odorous matter of the skin belongs properly to the perspiration, and have presumed it to be the product of specific organs. This is, however, conjectural; and the experiments of Thénard, as well as the facts we have just mentioned, would rather seem to show, that the matter of sweat itself has, within it, the peculiar odour. The fact of the dog tracing its master to an immense distance, and discovering him in a crowd, has induced a belief, that the scent may be distinct from the matter of sweat; but the supposition is not necessary, if we admit the matter of perspiration to be itself odorous.

Besides the causes before referred to, the quantity of perspiration is greatly augmented by running or by violent exertion of any kind; especially if the temperature of the air be elevated. Warm fluids favour it greatly, and hence their use, alone or combined with sudorifics, where this class of medicines is indicated.

a Edwards, Sur l'Influence des Agens Physiques, &c.; or Drs. Hodgkin and Fisher's translation, Lond. 1832; also, the author's Elements of Hygiène, p. 63, Philad. 1835; his General Therapeutics, p. 278, Philad. 1836; or his General Therapeutics and Materia Medica, vol. i., Philad. 1843.

VOL. II. - 22

Magendie<sup>a</sup> conceives, that being readily absorbed, they are also readily exhaled. This may be true; but the perspiration breaks out too rapidly to admit of this explanation. When ice-cold drinks are taken in hot weather, the cutaneous transpiration is instantaneously excited. The effect, consequently, must be produced by the refrigerant influence of the cold medium on the lining membrane of the stomach, — this influence, being propagated, by sympathy, to every part of the capillary system. The same explanation is applicable to warm drinks, whilst the hot exert a sympathetic effect on the skin by virtue of their stimulant properties exerted on the mucous membrane.

With regard to the uses of the insensible transpiration, it has been supposed to preserve the surface supple, and thus to favour the exercise of touch; and also, by undergoing evaporation, to aid in the refrigeration of the body. It is probable, however, that these are quite secondary uses under ordinary circumstances, and that the great office, performed by it, is to remove a certain quantity of fluid from the blood: hence it has been properly termed the cutaneous depuration. In this respect, it bears a striking analogy to the urine, which is the only other depuratory secretion, with the exception of the pulmonary transpiration, which we shall find essentially resembles the cutaneous. Being depuratory, it has been conceived, that any interruption to the transpiration must be attended with the most serious consequences; accordingly most diseases have, from time to time, been ascribed to this cause. There is, however, so great a compensation existing between the urinary and cutaneous depurations, that if one be augmented, the other is decreased, - and conversely. Besides, it is well known, that disease is more apt to be induced by partial and irregular application of cold than by frigorific influences of a more general character. The Russian vapour-bath exemplifies this; the bather frequently passing with impunity from a temperature of 130° into cold water. The morbific effect - in these cases of fancied check given to perspiration - is derangement of the capillary vessels engaged in the important functions of nutrition, calorification, and secretion, and the extension of this derangement to every part of the system.c

As the sensible transpiration or sweat is probably only the insensible perspiration in increased quantity, with the addition ofsalts, and other matters that are not evaporable, its uses demand

no special notice.

# b. Pulmonary Transpiration.

The pulmonary transpiration, to which we have so often alluded, bears a striking analogy to the cutaneous. Sir B. Brodie and Magendie, from the examination of cases of fistulous opening into

\* Précis de Physiologie, 2de édit. ii. 455.

See Dr. Robert Willis, Proceedings of the Royal Society, No. 56, Lond. 1843.
 This subject has been expatiated upon by the author, in the works referred to, p. 253, Note.

the trachea, deny that it comes from the lungs, believing it to be formed by the moist mucous lining of the nose, throat, &c.; but this view has been disproved by Paoli and Regnoli, in the case of a young female, whose trachea had been opened, and in whom, at the temperature of 39° Fahr., watery vapour was distinctly ex-

pired through the canula.

Mojona strangely supposes the vapour of the breath to be a watery fluid secreted by the thyroid gland, and suspended in the respired air, from its volatility, caused by the presence of caloric. At one time, it was universally believed to be owing to the combustion of the air, with the hydrogen and carbon given off from the lungs; but we have elsewhere shown, that no such combustion occurs; and, besides, the exhalation takes place, when gases, containing no oxygen, have been respired by animals. It is now almost universally admitted to be exhaled into the air-cells of the lungs from the pulmonary artery chiefly, but partly from the bronchial arteries, distributed to the mucous membrane of the air-passages. Much of the vapour, Dr. Prout conceives, - and Dr. Carpentere thinks, with much probability, — is derived from the chyle in its passage through the lungs; and thus, he considers, the weak and delicate albumen of the chyle is converted into the strong and perfect albumen of the blood.

Several interesting experiments have been made on this exhalation, by Magendie, Milne Edwards, Breschet, and others. If water be injected into the pulmonary artery, it passes into the aircells in myriads of almost imperceptible drops, and mixes with the air contained in them. Magendie<sup>4</sup> found, that its quantity might be augmented at pleasure on living animals, by injecting distilled water, at a temperature approaching that of the body, into the venous system. He injected into the veins of a small dog a considerable amount of water. The animal was at first in a state of real plethora, the vessels being so much distended that it could scarcely move; but, in a few minutes, the respiration became manifestly hurried, and a large quantity of fluid was discharged from the mouth, the source of which appeared evidently to be

the pulmonary transpiration considerably augmented.

Not only, however, is the aqueous portion of the blood exhaled in this manner; but experiment shows, that many substances, introduced into the veins by absorption, or by direct injection, issue by the lungs. Weak alcohol, a solution of camphor, ether and other odorous substances, when thrown into the cavity of the peritoneum or elsewhere, were found, by Magendie, to be speedily absorbed by the veins, and conveyed to the lungs, where they transuded into the bronchial cells, and were recognised by the smell in the expired air. Phosphorus, when injected, exhibited

<sup>·</sup> Leggi Fisiologiche, &c., translated by Skene, p. 76, Lond. 1827.

Bostock's Physiology, 3d edit. p. 359.
 Human Physiology, § 549, Lond. 1842.

this transmission in a singular and evident manner. Magendie, on the suggestion of M. Armand de Montgarny, "a young physician," he remarks, "of much merit," now no more, injected into the crural vein of a dog, half an ounce of oil, in which phosphorus had been dissolved; scarcely had he finished the injection, before the animal sent through the nostrils clouds of a thick, white vapour, which was phosphoric acid. When the experiment was made in the dark, these clouds were luminous. M. Tiedemann injected a drachm of the expressed juice of garlic into a vein in the thigh of a middle-sized dog: in the space of three seconds the breath smelt strongly of garlic. When spirit of wine was injected, the exhaled vapour was recognised when the injection

was scarcely over.

MM. Breschet and Milne Edwardse made several experiments, for the purpose of discovering, why the pulmonary transpiration expels so promptly the different gases and liquid substances received into the blood. Considering properly, that exhalation differs only from absorption in taking place in an inverse direction, these gentlemen conjectured, that it ought to be accelerated by every force, that would attract the fluids from within to without; and such a force they conceive inspiration to be, which, in their view, solicits the fluids of the economy to the lungs, in the same mechanical manner as it occasions the entrance of air into the aircells. In support of this view, they adduce the following experiments. To the trachea of a dog, a pipe, communicating with a bellows, was adapted, and the thorax was largely opened. Natural respiration was immediately suspended; but artificial respiration was kept up by means of the bellows. The surface of the air-cells was, in this way, constantly subjected to the same pressure; there being no longer diminished pressure during inspiration, as when the thorax is sound, and the animal breathing naturally. Six grains of camphorated spirit were now injected into the peritoneum of the animal; and, at the same time, a similar quantity was injected into another dog, whose respiration was natural. In the course of from three to six minutes, the odorous substance was detected in the pulmonary transpiration of the latter; but in the other it was never manifested. In the first animal, they now exposed a part of the muscles of the abdomen, and applied a cupping-glass to it; when the smell of the camphor speedily appeared at the cupped surface. Their conclusion was, that the pulmonary surface, having ceased to be subjected to the suction force of the chest, during inspiration, the exhalation was arrested, whilst that of the skin was developed as soon as an action of aspiration was exerted upon it by the cupping-

Précis, &c. ii. 348.

c Recherches Expérimentales sur l'Exhalation Pulmonaire, Paris, 1826.

b Tiedemann and Treviranus Zeitschrift für Physiologie, vol. v., part ii.; and British and Foreign Medical Review, i. 241, Lond. 1836.

Into the crural veins of two dogs;—one of which breathed naturally, and the other was circumstanced as in the last experiment,—they injected the essential oil of turpentine. In the first of these, the substance was soon apparent in the pulmonary transpiration; and, on opening the body, it was discovered, that the turpentine had impregnated the lung and the pleura much more strongly than the other tissues. In the other animal, on the contrary, the odour of the turpentine was scarcely apparent in the vapour of the lungs; and, on dissection, it was not found in greater quantity in the lungs than in the other tissues;—in the pleura than in the peritoneum.

From the results of these experiments, MM. Breschet and Edwards conclude, that each inspiratory movement constitutes a kind of suction, which attracts the blood to the lungs; and which causes the ejection, through the pulmonary surface, of the liquid and gaseous substances that are mingled with that fluid, more than through the other exhalant surfaces of the body. In their experiments, these gentlemen did not find that the exhalation was effected with equal readiness in every part of the surface, when the cupping-glass was applied in the manner that has been mentioned. The skin of the thigh, for example, did not indicate the odour of camphorated alcohol, as that of the region of the stomach.

The chemical composition of the pulmonary transpiration is probably nearly identical with that of the sweat; appearing to consist of water, holding in solution, perhaps, some saline and albuminous matter; but our information, on this matter, derived from the chemist, is not precise. Collard de Martigny's experiments make it consist, in 1000 parts, — of water 907, carbonic acid 90; animal matter — the nature of which he was unable to determine — 3 parts. Chaussier found, that by keeping a portion of it in a close vessel, exposed to an elevated temperature, a very evident putrid odour was exhaled on opening the vessel. This could only have arisen from the existence of animal matter in it.

The pulmonary transpiration being liable to all the modifications which affect the cutaneous, it is not surprising, that we should meet with so much discordance in the estimates of different individuals, regarding its quantity in a given time. Hales valued it at 20 ounces in the twenty-four hours; Sanctorius, Menzies, and Dr. William Wood, at 6 ounces; Abernethy at 9 ounces; Lavoisier and Séguing at 17½ ounces, poids de marc; Thomson at 19 ounces, Dalton at from 1 pound 8¾ ounces, to 20½ ounces

- \* Magendie's Journal de Physiologie, x. 111.
- b Statical Essays, ii. 322, Lond. 1767. Comedicina Statica, Aphor. v.
- d Dissertation on Respiration, p. 54, Edinb. 1796.
- . Essay on the Structure, &c., of the Skin, Edinb. 1832.
- f Surgical and Physiol. Essays, p. 141, Lond. 1793.
- 6 Mem. de la Société Royale de Médecine, pour 1782-3; Annal. de Chimie, v. 264; and Mém. de l'Acad. des Sciences, pour 1789.
  - System of Chemistry, vol. iv. Manchester Memoirs, 2d series, ii. 29.

avoirdupois, and Carpenter, at from 16 to 20 ounces. The uses it serves, in the animal economy, are identical with those of the cutaneous depuration. It is essentially depuratory. Experiments, some of which have been detailed, have sufficiently shown, that volatile substances introduced, in any way, into the circulatory systemare rapidly exhaled into the bronchial tubes—if not adapted for the formation of arterial blood. Independently, therefore, of the lungs being the great organ of respiration, they play a most important part in the economy, by throwing off those substances, that might be injurious, if retained.

#### c. Exhalation of the Mucous Membranes.

The mucous membranes, like the skin, which they so strongly resemble in their structure, functions, and diseases, exhale a similar transpiratory fluid; which has not, however, been subjected to chemical examination. It is, indeed, almost impracticable to separate it from the follicular secretions, poured out from the same membrane; and from the extraneous substances, almost always in contact with it. It is probably, however, similar to the fluid of the cutaneous and pulmonary depurations, both in character and use.

#### FOLLICULAR SECRETIONS.

The follicular secretions must, of necessity, be effected from the skin or the mucous membranes; as the follicles or crypts are met with there only. They may, therefore, be divided into two classes; — 1st, the mucous follicular secretion; and 2d, the cutaneous follicular secretion.

# a. Mucous Follicular Secretion.

The whole extent of the great mucous membranes, — lining the alimentary canal, the air-passages, and the urinary and genital organs, — is the seat of a secretion, the product of which has received, in the abstract, the name of mucus; although it differs somewhat according to the situation and character of the particular follicles, whence it proceeds. Still, essentially, the structure, functions, and product are the same. According to M. Donné, its character is alkaline in health; in disease, often acid. In the history of the different functions, in which some of the mucous membranes are concerned, the uses of this secretion have been detailed; and in those that will hereafter have to engage attention, in which other mucous membranes are concerned, their uses will fall more conveniently under notice then. But few points will, therefore, require explanation at present.

The mucus, secreted by the nasal follicles, seems alone to have been subjected to chemical analysis. Fourcroy and Vauquelind

<sup>\*</sup> Ibid. vol. v.

b Human Physiology, § 549, Lond. 1842.
c Journal Hebdomad. Février, 1834.
d Journal de Physique, xxxix. 359.

found it composed of precisely the same ingredients as the tears. According to the analysis of Berzelius, a its contents are as follows: -- water, 933.7; mucus, 53.3; chlorides of potassium and sodium, 5.6; lactate of soda, with animal matter, 3.0; soda, 0.9; albumen and animal matter, soluble in water, but insoluble in alcohol, with a trace of phosphate of soda, 3.5. According to Raspail, mucus is the mere product of the healthy and daily disorganization, or wear and tear of the mucous membranes. Every mucous membrane, he affirms, exfoliates in organized layers, and is thrown off, more or less, in this form; whilst the serous membranes either do not exfoliate, or their exfoliation (excoriation) is resolved into the liquid form, to be again absorbed by the organs; but this - like many other of M. Raspail's speculations — is a generalization which does not appear to be warranted by the facts; the slightest examination, indeed, exhibits, that the general physical character of the mucus is very different from that of the membranes which form it: still, mucus, when examined by a microscope of high magnifying powers — and this the author had an opportunity of doing through the hydroxygen microscope - does present, here and there, appearances of shreds similar to those described by Raspail. These have been esteemed by recent histologists to be detached epithelium cells, with granulated globular particles, that are regarded as characteristic of the secretion from the surface of mucous membranes.

Although mucus is classed as a follicular secretion, it would seem to be formed in mucous membranes in which no follicles can be detected, as in those lining the frontal and other sinuses of the cranium. Mandl, who first stated the belief in the identity of structure of the globules of mucus and pus, and of the red corpuscles of the blood, describes mucus as composed of a viscid liquid in which are swimming, besides a greater or less number of lamellæ of epithelium, special elements, which he calls globules of mucus. These are of two kinds, — the one consisting of mammillated corpuscles, 0.005 to 0.006 of a millimètre in diameter; the other, from 0.01 to 0.02 of a millimètre in diameter, — the latter being true cells, composed of an envelope and a nucleus.

The great use of mucus, wherever met with, is to lubricate the

surface on which it is poured.e

### b. Follicular Secretion of the Skin.


This is the sebaceous and micaceous humour, observed in the skin of the cranium, and in that of the pavilion of the ear. It is

Med. Chirurg. Transactions, tom. iii.; also, Thomson, Chemistry of Animal Bodies, p. 507, Edinb. 1843.

<sup>Chimie Organique, p. 246, and p. 504, Paris, 1832.
Carpenter, Human Physiology, § 640, Lond. 1842.
Manuel d'Anatomie générale, p. 478, Paris, 1843.</sup> 

Burdach's Physiologie als Erfahrungswissenschaft, v. 235, Leipz. 1835.

also the humour, which occasionally gives the appearance of small



Sebaceous and Ceruminous Glands.

A. Section of skin magnified three diameters. b, b. Hairs. c, c. Superficial sebaceous glands. a, a. Larger and deeper seated glands by which the cerumen appears to be secreted. B. A ceruminous gland more largely magnified, formed of the convoluted tube a, forming the excretory duct b. c. A small vessel, and its branches. c. A hair from the meatus auditorius, perforating the epidermis at e, and at d, contained within its double follicle, e, e. a, a. Sebaceous follicles of the hair with their excretory ducts. — (Wagner.)

worms beneath the skin of the face. when it is forced through external aperture of the follicle; and which, when exposed to the air, causes the black spots sometimes observable the face. The cerumen is, likewise, a follicular secretion, as well as the whitish. odorous fatty matter which forms under the prepuce of the male, and in external parts

of the female, where cleanliness is disregarded. The humour of Meibomius is also follicular, as well as that of the caruncula la-

chrymalis.

The use of this secretion is, — to favour the functions of the part over which it is distributed. That which is secreted from the skin, is spread over the epidermis, hair, &c., giving suppleness and elasticity to the parts, rendering the surface smooth and polished, and thus obviating the evils of abrasion that might otherwise arise. It is also conceived that its unctuous nature may render the parts less permeable to humidity.<sup>a</sup>

#### GLANDULAR SECRETIONS.

The glandular secretions are seven in number; those of the tears, saliva, pancreatic juice, bile, urine, sperm, and milk.

# a. Secretion of the Tears.

The lachyrmal apparatus, being a part of that accessory to vision, was described under another head (vol. i., p. 196). As we meet with the tears, they are not simply the secretion of the lachrymal Burdach, op. cit. vi. 245.

gland, but of the conjunctiva, and occasionally of the caruncula lachrymalis and follicles of Meibomius. They have a saline taste; mix freely with water; and, owing to the presence of free soda, communicate a green tint to the blue infusion of violets. Their chief salts are the chloride of sodium, and phosphate of soda. According to Fourcroy and Vauquelin, the animal matter of the tears is mucus; but it is presumed, by some, to be albumen or an analogous principle — dacryolin.

When the tears are examined with the microscope, globules of

mucus, and debris of the epidermis are seen in them.b


This secretion is more influenced by the emotions than any other; and hence it is concerned in the expressions of lively joy or sorrow, especially of the latter.

#### b. Secretion of the Saliva.

The salivary apparatus has like wise engaged attention elsewhere. It consists of a parotid gland on each side, situate in front of the ear, and behind the neck and ramus of the jaw; a submaxillary, beneath the body of the bone; and a sublingual, situate immediately beneath the tongue; - the parotids and submaxillary glands having each but one excretory duct; - the sublingual

several. The structure of the salivary glands in man greatly resembles that of the mammary glands. The marginal figure (Fig. 191) exhibits their structure in man. All these glands pour their respective fluids into the mouth, where it collects, and becomes mixed with the exhalation of the mucous membrane of the mouth, and the secretion from its follicles. It is this mixed fluid that has been generally analyzed by the chemist. When collected, without the action of sucking, it is of a specific gravity varying from 1004 to 1009, translucent, slightly opaque, very frothy, and ultimately deposits a nebulous sediment. Even with of a new-born infant, filled with the purest saliva there is always found mercury and magnified five diameters. - (Weber.) mixed a few epithelial cells, derived from





the mucous lining of the mouth, or from the excretory ducts of the secreting glands. It usually contains free alkali; in rare cases, during meals, Professor Schultz,d of Berlin, found it acid; and, during fasting, it is occasionally neutral. Mitscherlich,e indeed, affirms that it is acid whilst fasting, but becomes alkaline during

Journal de Physique, xxxix. 256. See, also, Thomson, Chemistry of Animal Bodies, p. 511, Edinb. 1843.

b Mandl, Manuel d'Anatomie générale, p. 489, Paris, 1843.

Dr. Julius Vogel, in Wagner's Elements of Physiology, § 144, Lond. 1842.

4 Hecker, Wissenschaftliche Annalen, B. ii. H. 1, § 32, 1835.

 Rullier and Raige-Delorme, in art. Digestion, Dict. de Médecine, 2de édit. x. 300, Paris, 1835.

eating, - the alkaline character disappearing at times with the first mouthful of food. The average amount of the secretion in the twenty-four hours does not probably exceed four ounces.a According to Berzelius, its constituents are - water, 992.2; peculiar animal matter, 2.9; mucus, 1.4; chlorides of potassium and sodium, 1.7; lactate of soda, and animal matter, 0.9; soda, 0.2. Drs. Bostocke and Thomas Thomsond think, that the "mucus" of Berzelius resembles coagulated albumen in its properties. In the tartar of the teeth, which seems to be a sediment from the saliva, Berzelius found 79 parts of earthy phosphate; 12.5 of undecomposed mucus; 1 part of a matter peculiar to the saliva, and 7.8 of an animal matter soluble in chlorohydric acid. This animal matter, according to the microscopic experiments of Raspail, is composed of the deciduous fragments from the mucous membrane of the cavity of the mouth; and he considers, that the saliva is nothing more than an albuminous solution, mixed with different salts, which are capable of more or less modifying its solubility in water, and of shreds or layers of tissue. MM. Leuret and Lassaignef analyzed pure saliva, obtained from an individual labouring under salivary fistula, and found it to contain, - water, mucus, traces of albumen, soda, chloride of potassium, chloride of sodium, carbonate and phosphate of lime; - and Messrs. Tiedemann and Gmeling affirm, - and their analysis agrees pretty closely with that of Van Settenh - that the saliva contains only one or two-hundredths of solid matter, which are composed of a peculiar substance, called salivary matter or ptyalin, osmazome, mucus, perhaps albumen, a little fat containing phosphorus, and the insoluble salts - phosphate and carbonate of lime. Besides these, they detected the following soluble salts; - acetate, carbonate, phosphate, sulphate, muriate, and sulpho-cyanate of potassa.i Treviranusi thinks the saliva contains a peculiar acid, to which he gives the name Blausäure, properly combined with an alkali; but its chemical properties resemble the sulpho-cyanic acid so greatly, that according to Kastnerk they may be taken for each other. Messrs. Tiedemann and Gmelin, and M. Donné, found the saliva invariably alkaline, when the functions of the stomach were well performed. The latter gentleman considered acidity of the stomach a diagnostic symptom of gastritis; and Dr. Robt. Thomson<sup>m</sup> found

a Mitscherlich, Rust's Magazine, Bd. xxxvi. S. 491.

b Medico-Chirurgical Transactions, iii. 242; also, Thomson, Chemistry of Animal Bodies, p. 389, Edinb. 1843.

c. Physiol. ed. cit. p. 487. d System of Chemistry, vol. iv.

º Nouveau Système de Chimie Organique, p. 454. f Recherches, &c., sur la Digestion, p. 33, Paris, 1826.

5 Recherches, &c., sur la Digestion, par Jourdan, Paris, 1827.

h De Salivâ ejusque Vi et Utilitate, Groning. 1837; and Brit. and For. Med. Rev., Jan. 1839, p. 236.

Magendie's Précis Elémentaire, ii. 63; and Burdach's Physiologie, u. s. w., v. 251, Leipz. 1835. i Biologie, Band. iv. § 330.

Ficinus, in art. Speichel, in Pierer's Anat. Physiol. Real Wörterbuch, vii. 634, 1 Archives Générales, Mai & Juin, 1835.

m Records of General Science, Dec. 1836.

the acid reaction in all cases of inflammation of the mucous and serous membranes. With the view of testing these experiments, Mr. Laycocka instituted numerous experiments, and affords the tabular results of no less than 567 observations. His deductions do not accord with those of M. Donné. They are as follows:—

1. The saliva may be acid without any apparent disease of the stomach, and when the individual is in good health. 2. It is alkaline during different degrees of gastric derangement, as indicated by the tongue. 3. It may be alkaline, acid and neutral, when the gastric phenomena are the same; and, consequently, acidity of the saliva is not a diagnostic mark of gastric derangement; and, lastly—in general it is alkaline in the morning, and acid in the evening.

When saliva is examined by the microscope, it presents, besides a considerable number of lamellæ of epithelium, globules in variable quantity, which, according to Mandl, proceed partly from the muciparous glands of the mouth, and partly from the salivary glands. These cannot, however, be distinguished from each other.

As the salivary secretion forms a part in the processes preparatory to the stomachal digestion, its uses have been detailed in the first volume of this work.

A soft, whitish or yellowish matter, of greater or less thickness, is constantly deposited on the teeth, which, unless attention is paid, accumulates, and sometimes adheres to them with great force, constituting hard and dry concretions, known under the name of tartar or tartar of the teeth. Different views have existed in regard to its origin. Some have supposed it to be a secretion, others a deposition from the saliva, which is the most probable opinion; and others, that it is an exhalation from the capillary vessels, to which the mucous membrane of diseased gums is liable. Recently, however, it has been affirmed by M. Mandl, to be a collection of calcareous skeletons of infusoria, agglutinated by means of dried mucus.

### c. Secretion of the Pancreatic Juice.

The pancreas or sweetbread, (Fig. 193, G,) secretes a juice or humour, called succus pancreaticus or pancreatic juice. Its texture resembles that of the salivary glands: and hence it has been called by some the abdominal salivary gland. It is situate transversely in the abdomen, behind the stomach, towards the concavity of the duodenum; is about six inches in length, and between three and four ounces in weight. From the results of six examinations, Dr. Gross<sup>d</sup> gives the following as the mean weight and dimensions. Weight, 2½ ounces; length, 7 inches; breadth at the body and


Lond. Med. Gazette, Oct. 7, 1837. See, also, for a detailed account of the saliva, Dr. S. Wright, Lond. Lancet, vols. 1 and 2, for 1841-2.

<sup>Manuel d'Anatomie générale, p. 488, Paris, 1843.
Gazette des Hopitaux, 8 Août, 1843, p. 363.</sup> 

<sup>4</sup> Elements of Pathological Anatomy, ii, 357, Boston, 1839.

splenic extremity, 16½ lines; breadth at the neck, 12 lines; at the head, 2 inches and 3 lines; thickness at the body, neck, and splenic extremity, 4 lines; thickness at the head, 8 lines. Bécourt found the average length of thirty-two to be 8 inches; and the weight between 3 and 4 ounces." It is of a reddish-white colour, and firm consistence. Its excretory ducts terminate in one, - called the duct of Wirsung, - which opens into the duodenum, at times separately from the ductus communis choledochus, but close to it; at other times, being confounded with, or opening into it.b

The amount of fluid, secreted by the pancreas, does not seem



Biliary and Pancreatic Ducts.

a. Hepatic duct, formed by a branch from right, and one from the left lobe of liver. b. Fundus of gall-bladder. c, d. Body and neck of gall-bladder. e. Cystic duct. f. Ductus communis choledochus. g, g. Trunk and branches of pancreatic duct. h. Termination of ductus communis choledochus and ductus pancreaticus.

to be considerable. Magendie, in his experiments, was struck with the small quantity discharged. Frequently, scarcely a drop issued in half an hour; and, occasionally, a much longer time elapsed. Nor did he find that the flow, according to the common opinion and to probability, was more rapid whilst digestion was going on. It will be readily understood, therefore, that it cannot be an easy task to collect it. De Graaf, a Dutch anatomist, affirms, that he succeeded, by introducing into the intestinal end of the excretory duct, a small quill, terminat-

ing in a phial fixed under the belly of the animal. Magendie,d however, states, that he tried this plan several times, but without success; and he believes it to be impracticable. The plan he adopts is to expose the intestinal orifice of the duct; to wipe, with a fine cloth, the surrounding mucous membrane; and as soon as a drop of the fluid oozes, to suck it up by means of a pipette or small glass tube. In this way, he collected a few drops, but never sufficient to undertake a satisfactory analysis. Messrs. Tiedemann and Gmeline make an incision into the abdomen; draw out the duodenum, and a part of the pancreas; and, opening the excretory duct, insert a tube into it; and a similar plan was adopted successfully on a horse by MM. Leuret and Lassaigne. The difficulty, experienced in collecting a due quantity, is a probable cause of some of the discrepancy amongst observers, regard-

Précis Elémentaire, i. 462; J. P. Mondière, op. cit. · Tract. de Pancreat. Lugd. Bat. 1761; Haller, Elem. Physiol. lib. xxii.

Recherches sur le Pancreas, ses Fonctions et ses Alterations Organiques, Thèse, Strasbourg, 1830, cited by Mondière, Archives Générales de Médecine, Mai, 1836.

Précis, &c., ii. 462. Researches, &c., i. 41. Researches, &c., p. 49.

ing its sensible and chemical properties. Some of the older physiologists affirm it to be acidulous and saline; others assert that it is alkaline." The majority of those of the present day compare it with the saliva, and affirm it to be inodorous, insipid, viscid, limpid, and of a bluish-white colour. The latest experimenters by no means accord with each other. According to Magendie, it is of a slightly yellowish hue, saline taste, devoid of smell, occasionally alkaline, and partly coagulable by heat. MM. Leuret and Lassaigne found that of the horse, - of which they obtained three ounces, - to be alkaline, and composed of 991 parts of water in 1000; of an animal matter, soluble in alcohol; another, soluble in water; traces of albumen and mucus; free soda; chloride of sodium; chloride of potassium, and phosphate of lime. In their view, consequently, the pancreatic juice strongly resembles saliva. MM. Tiedemann and Gmelin succeeded in obtaining upwards of two drachms of the juice in four hours; and, in 100 parts, they found from five to eight of solid. These solid parts consisted of osmazome; a matter which became red by chlorine; another analogous to casein, and probably associated with salivary matter; much albumen; a little free acid, probably the acetic; the acetate, phosphate, and sulphate of soda, with a little potassa; chloride of potassium, and carbonate and phosphate of lime: - so that, according to these gentlemen, the pancreatic juice differs from the saliva in containing: - a little free acid, whilst the saliva is alkaline; much albumen, and matter resembling casein; but little mucus and salivary matter, and no sulpho-cyanate of potassa."

The precise use of the pancreatic juice in digestion — as we have previously seen — is not determined. Brunner<sup>d</sup> removed almost the whole pancreas from dogs, and tied and cut away portions of the duct; yet they lived apparently as well as ever. It is not presumable, therefore, that the secretion can be possessed of very im-

portant uses.

# d. Secretion of the Bile.

The biliary secretion is, also, a digestive fluid, of which we have spoken in the appropriate place. The mode, however, in which the process is effected, has not yet been investigated. The apparatus consists of the liver, which accomplishes the formation of the fluid; the hepatic duct, — the excretory channel, by which the bile is discharged; the gall-bladder, in which a portion of the bile is retained for a time; the cystic duct, — the excretory channel of the gall-bladder; and the ductus communis choledochus, or

b Fourcroy and Thomson in their Systems of Chemistry.

Burdach's Physiologie, u. s. w., v. 257, Leipz. 1835; and Thomson, Chemistry

of Animal Bodies, p. 403, Edinb. 1843.


<sup>&</sup>lt;sup>3</sup> Haller, Elem. Physiol. vi. 10; and Seiler, in art. Pancreas, Pierer's Anat. Physiol. Real Wörterb., Band. vi. 100, Altenb. 1825.

<sup>&</sup>lt;sup>4</sup> Experimenta nova circa Pancreas, Amstel. 1683; and J. T. Mondière, op. cit., in a Mémoire, crowned by the Société Médicale d'Emulation, of Paris, and entitled "Recherches pour servir a l'Histoire Pathologique du Pancreas."

vol. II. - 23

choledoch duct, formed by the union of the hepatic and cystic ducts, and which conveys the bile immediately into the duodenum.

The liver, (A, A, Fig. 129, and A, A, Fig. 193,) is the largest gland in the body; situate in the abdomen beneath the diaphragm, above the stomach, the arch of the colon, and the duodenum; filling the whole of the right hypochondrium, and more or less of the epigastrium, and fixed in its situation by duplicatures of the peritoneum, called ligaments of the liver. The weight of the human liver is generally, in the adult, about three or four pounds. Some make the average about five pounds; a but this is a large estimate. Of 60 male livers weighed, Dr. John Reidb found the average weight to be 52 oz. 122 dr.: and of 25 female, 45 oz. 32 dr.



Abdominal and Pelvic Viscera.

In disease, however, it sometimes weighs twenty or twenty-five pounds; and, at other times, not as many ounces. Its shape is irregular, and it is divided into three chief lobes, the right, the left, and the lobulus Spigelii. Its upper convex surface touches every where the arch of the diaphragm. The lower concave surface corresponds to the stomach, colon, and right kidney. On the latter surface, two fissures are observable; - the one passing from before to behind, and lodging the umbilical vein in the fætus -- called the horizontal sulcus or fissure, great fissure or fossa umbilicalis; the other, cutting the last at right angles, and running from right to left, by which the different nerves and vessels proceed to and from the liver, and called the principal fissure, or sulcus transversus.

The liver itself is composed of the following anatomical elements: A, A. Concave surface of liver turned up. 1. The hepatic artery, a branch of wards, and to the right side. B. Lobulus Spi. 1. The hepatic artery, a branch of gelii. Between B and C, the porta of the liver the cœliac, which ramifies minutely D. Ligamentum rotundum. E, F. Gall-bladder. G. The pancreas. H. The spleen. I, through the substance of the organ. The ribs. K, K. The kidneys. L, L. Renal The minuter branches of this artery waite arteries. Q, Q. Common iliac arteries are arranged somewhat like the R. Vena cava. S. The spermatic veins. U. Common iliac veins. V. End of colon hairs in a painter's brush, and have X. Commencement of the rectum. Y, y hence been called the penicilli of Urinary bladder. the liver. Mr. Kiernane believes,

W. J. E. Wilson, art. Liver, Cyclop. of Anat. and Phys. Sept. 1840. b Lond, and Edinb. Monthly Journ. of Med. Sciences, April, 1843, p. 323.

Philosophical Transactions for 1833, p. 711. See, also, Elliotson's Human Physiology, p. 93, Lond. 1840.


that the blood, which enters the liver by the hepatic artery, fulfils three functions: - it nourishes the organ; supplies the excretory ducts with mucus; and, having fulfilled these objects, it becomes venous; enters the branches of the portal veins, and not the radicles of the hepatic, as usually supposed, and contributes to the secretion of bile. 2. The vena porta, which we have elsewhere seen to be the common trunk of all the veins of the digestive organs and of the spleen. It divides like an artery, its branches accompanying those of the hepatic artery. Where the vein lies in the transverse fissure, it is of great size, and has hence been called sinus venæ portæ. The possession of two vascular systems, containing blood, is peculiar, perhaps, to the liver, and has been the cause of some difference of opinion, with regard to the precise material arterial or venous - from which the bile is derived. According to Mr. Kiernan, the portal vein fulfils two functions: it carries the blood from the hepatic artery, and the mixed blood to the coats of the excretory ducts. This vessel has been called the vena arteriosa, because it ramifies like an artery, and conveys blood for secreting: but, as Mr. Kiernan has observed, it is an arterial vein in another sense, as it is a vein to the hepatic artery, and an artery to the hepatic vein. 3. The excretory ducts or biliary ducts. These are presumed to arise from acini, communicating, according to some, with the extremities of the vena portæ; according to others, with radicles of the hepatic artery; whilst others have considered, that the radicles of the hepatic ducts have blind extremities, and that the capillary bloodvessels which secrete the bile, ramify on them. This last arrangement of the biliary apparatus in the liver was well shown in an interesting pathological case, which fell under the care of Professor Hall, in the Baltimore Infirmary, and which was examined after death in the author's presence. The particulars of the case have been detailed, with some interesting remarks by Professor Geddings.a In this case, in consequence of cancerous matter obstructing the ductus communis choledochus, the whole excretory apparatus of the liver was enormously distended; the common duct was dilated to the size of the middle finger: at the point where the two branches that form the hepatic duct emerge from the gland, they were large enough to receive the tip of the middle finger; and as they were proportionately dilated to their radicles, in the intimate tissue of the liver, their termination in a blind extremity was clearly exhibited. These blind extremities were closely clustered together, and the ducts, proceeding from them, were seen to converge, and to terminate in the main trunk for the corresponding lobe. At their commencement, the excretory ducts are termed pori biliarii. These ultimately form two or three large trunks, which issue from the liver by the transverse fissure, and end in the hepatic duct. 4. Lymphatic vessels. 5. Nerves, in small number, compared with the size of the liver, some proceeding from the North American Archives of Medical and Surgical Science, for June, 1835, p. 157.

eighth pair; but the majority from the solar plexus, and following the course and divisions of the hepatic artery. 6. The supra-hepatic veins or venæ cavæ hepaticæ, which arise in the liver by imperceptible radicles, communicating, according to common belief, with the final ramifications of both the hepatic artery and vena portæ; according to Mr. Kiernan occupying the centre of the lobules - and hence termed intralobular veins. They return the superfluous blood, carried to the liver by these vessels, by means of two or three trunks, and six or seven branches, which open into the vena cava inferior. These veins generally pass, in a convergent manner, towards the posterior margin of the liver, and cross the divisions of the vena portæ at right angles. 7. The remains of the umbilical vein, which, in the fœtus, enters at the horizontal fissure. This vein, after respiration is established, becomes converted into a ligamentous substance, called, from its shape, ligamentum rotundum or round ligament.

The parenchyma, or substance formed by these anatomical elements, it is difficult to describe; and although the term livercoloured is used in common parlance, it is not easy to say what

are the ideas attached to it.

The views of Mr. Kiernan, in regard to the intimate structure of the liver, which have been embraced by so many anatomists,



may be understood by the accompanying illustrations, taken from his communications on the subject. The acini, to which allusion has been made, are termed by him lobules. Fig. 194, A, exhibits some of the cells of which the lobules are composed, seen under a magnifying power of 200 diameters.

B represents a longitudinal section of a lobule with ramifications of the hepatic vein: and Fig. 195, the connection of the lo-

Fig. 195.




bules with the same vein; - the centre of each being occupied by a venous twig - or interlobular vein. Fig. 196 represents the lobules as seen on the surface of the liver when divided transversely. In this, B, exhibits the interlobular spaces; E, interlobular fissures; D, intralobular veins occupying the centres of the lobules; and E, smaller veins termi-

Connection of Lobules of Liver with Hepatic nating on the central veins. Fig. 197, exhibits a similar section of

three lobules, showing the arrangement of the two principal systems of bloodvessels; a, a, being intralobular veins; and b, b, in-

Fig. 196.

terlobular plexus formed by branches of the vena porta. Lastly, Fig. 197 represents a horizontal section of two superficial lobules, showing the interlobular plexus of biliary ducts: a, a, being the intralobular veins; b, b, trunks of biliary ducts, proceeding from the plexus, which traverses the lobules; c, interlobular tissue; and d, the parenchyma of the lobules.



It is from this arrangement of the Transverse Section of the Lobules of the Liver. - (Kiernan.) bloodvessels and biliary ducts, that

Mr. Kiernan infers the bile must be secreted from the portal Fig. 197.


vessels, which alone surround the lobules; the intralobular ramifications of the hepatic veins conveying back the blood to the heart, which has been inservient to the secretion. The views of Mr. bd Kiernan have been generally adopted by anatomists. Wagner, however, whilst he regards the beautiful figures and descriptions of Mr. Kiernan as the best he has seen, asserts, that "they very certainly also include many mistakes;" whilst Krause "combats the views of Kiernan, Horizontal Section of Three Superficial Lobules. holding them to be hypothetical."a



showing the two principal Systems of Bloodvessels .- (Kiernan.)


Mr. Kiernan has proceeded to deduce interesting pathologi-

cal inferences from the anatomical arrangement which he conceives to exist; thus, he considers that the lobules may be congested by accumulation of blood in the hepatic or in the portal venous system; and that this may be detected by a minute inspection of the lobules. The precise causes of this are referred to in another work.b The marginal




Horizontal Section of Two Superficial Lobules, showing interlobular Plexus of Biliary Ducts .- (Kiernan.)

Wagner, Elements of Physiology, by R. Willis, § 195, Lond. 1842. b See the author's Practice of Medicine, vol. ii. 2d edit., Philad. 1844. illustrations will be sufficient here. Fig. 199 represents the lo-




First Stage of Hepatic Venous Congestion .- (Kiernan.)

tal venous congestion. This is Fig. 200.



Second Stage of Hepatic Venous Congestion .- (Kiernan.)

peritoneum, which is very thin, transparent, easily lacerable, and



Portal Venous Congestion .-- (Kiernan.)

bules in the first stage of what he terms hepatic venous congestion, or congestion of the terminations of the hepatic vein: b, representing the interlobular spaces and fissures. In Fig. 200, the lobules are in the second stage. B and C, representing the interlobular spaces; D, congested intralobular or hepatic veins; E, congested patches extending to the circumference of the lobules; and F, the uncongested portions. Lastly, in Fig. 201, the lobules are in a state of por-This is not a common occurrence. It has been seen by Mr. Kiernan in children only. The view of Mr. Kiernan has also been held to explain the diversity of the statement of anatomists as to the relative position of the red and yellow substances, which have been considered to compose the liver: the red is the congested portion of the lobules, whilst the yellow is the non-congested portion in which the biliary plexus appears more or less

The liver has two coats; —

distinctly.a

vascular, and is the seat of the secretion, operated by serous membranes in general, It does not cover the posterior part, or the excavation for the gall-bladder, the vena cava, or the fissures in the concave surface of the liver. The inner coat is the proper membrane of the liver. It is thin, but not easily torn, and it covers not only every part of the surface of the liver, but also the large vessels that are proper to the organ. The

a Carpenter, Human Physiology, § 658, Lond. 1842.

condensed cellular substance, — which unites the sinus of the vena portæ and its two great branches, the hepatic artery, the common biliary duct, lymphatic glands, lymphatic vessels, and nerves in the transverse fossa or fissure of the liver, — was described by Glisson as a capsule; and hence has been called the *capsule* of Glisson. It connects, indeed, the various anatomical elements of the liver

together.

The gall-bladder, (Figs. 129, 192, 193, and 202,) is a small membranous pouch, of a pyriform shape, situate at the inferior and concave surface of the liver, to which it is attached, and above the colon and duodenum. A quantity of bile is usually found in it. It is not met with in all animals. It is wanting in the elephant, horse, stag, camel, rhinoceros, dilatation of the gall-

Fig. 202.

Under Surface of Liver.

stag, camel, rhinoceros, and goat; in certain of the cetacea; in some birds, as the ostrich, pigeon, and parrot; and is occasionally wanting in man. No traces of it are met with in the invertebrata. It may be looked upon as a column. — (Wilson.)

1. Right lobe. 2. Left lobe. 3. Lobus quadratus. 4. Lobus Spigelii. 5. Lobus caudatus. 6. Longitudinal fissure, in which is seen the rounded cord; the remains of the umbilical vein. 7. Pons hepatis. 8. Fissure for the ductus venosus; the obliterated cord of the ductus is seen passing backwards to be attached to the coats of inferior vena cava, 9. 10. Gall-bladder lodged in its fossa. 11. Transverse fissure, containing from before backwards, the hepatic duct, hepatic artery, and portal vein. 12. Vena cava. 13. A depression corresponding with curve of colon. 14. A double depression produced by right kidney and its supra-renal capsule. 15 Rough surface on posterior border of liver left uncovered by peritoneum; the cut edge of peritoneum surrounding this surface forms part of coronary ligament. 16. Notch on posterior border, corresponding with vertebral column. — (Wilson.)

ducts, and adapted for the reception and retention of bile. Its largest part, or fundus, Figs. 192 and 193, is turned forwards; and, when filled, frequently projects beyond the anterior margin of the liver. Its narrowest portion, cervix or neck is turned backwards, and terminate in the cystic duct. Externally, it is partly covered by the peritoneum, which attaches it to the liver, and to which it is, moreover, adherent by cellular tissue and vessels. Internally, it is rugous; the folds being reticulated, and appearing somewhat like

the cells of a honey-comb.

Anatomists have differed with regard to the number of coats proper to the gall-bladder. Some have described two only;—the peritoneal and mucous; others have added an intermediate cellular coat; whilst others have reckoned four;—a peritoneal coat,—a thin stratum of muscular fibres, passing in different directions, and of a pale colour,—a cellular coat, in which a number of bloodvessels is situate, and an internal mucous coat. The

existence of the muscular coat has been denied by perhaps the generality of anatomists; but there is reason for believing in its existence. Amussat saw muscular fibres distinctly in a gall-bladder dilated by calculi; and Dr. Monro, the present Professor of Anatomy in the University of Edinburgh, asserts, that he has seen it contract, in a living animal, for half an hour, under mechanical irritation, and assume the shape of an hour-glass. The mucous coat forms the rugæ to which we have already alluded. In the neck, and in the beginning of the cystic duct, there are from three to seven — sometimes twelve — semilunar duplicatures, which retard the flow of any fluids inwards or outwards. These are sometimes arranged spirally, so as to form a

kind of valve, according to Amussat.º

On the inner surface of the gall-bladder, especially near its neck, numerous follicles exist; the secretion from which is said to fill the gall-bladder, when that of the bile has been interrupted by disease, as in yellow fever, scirrhus of the liver, &c. The hepatic duct, Fig. 192, a, is the common trunk of all the excretory vessels of the liver; and makes its exit from that organ by the transverse fissure. It is an inch and a half in length, and about the diameter of an ordinary writing quill. It is joined, at a very acute angle, by the duct from the gall-bladder — the cystic duct, Fig. 192, e, to form the ductus communis choledochus. The cystic duct is about the same length as the hepatic. The ductus communis choledochus is about three, or three and a half, inches long. It descends behind the right extremity of the pancreas, through its substance; passes for an inch obliquely between the coats of the duodenum, diminishing in diameter; and ultimately terminates by a yet more contracted orifice, on the inner surface of the intestine, at the distance of three or four inches from the stomach. The structure of all these ducts is the same. The external coat is thick, dense, strong, and generally supposed to be of a cellular character; the inner is a mucous membrane, like that which lines the gallbladder.d

The secretion of bile is probably effected like the other glandular secretions; but modified, of course, by the peculiar structure of the liver. We have seen that the organ differs from every other secretory apparatus, in having two kinds of blood distributed to it: arterial blood by the hepatic artery; and venous blood by the vena portæ. A question has consequently arisen — from which of these is the bile formed? Anatomical inspection throws no light on the subject; and, accordingly, argument is all that can be adduced on one side or the other. The most common and the oldest opinion is, that the bile is separated from the blood of the

\* See, also, M. Barth, L'Expérience, Nov. 19, 1840.

· Magendie's Précis, &c., ii. 464.

Elements of the Anatomy of the Human Body, Edinb. 1825.

<sup>&</sup>lt;sup>4</sup> For a full account of the Normal Anatomy of the Liver, see W. J. E. Wilson, Cyclop. Anat. and Physiol., Sept. 1840; and his Anatomist's Vade Mecum, Amer. Edit. by Dr. Goddard, p. 520, Philad. 1843.

vena portæ; and the chief reasons adduced in favour of this belief, are the following: First. The blood of the portal system is better adapted than arterial blood for the formation of bile, on account of its having, like all venous blood, more carbon and hydrogen, which are necessary for the production of a humour as fat and oily as the bile; and, as the experiments of Schultz<sup>a</sup> and others have proved, that the portal blood contains more fat than that of the other veins and arteries, it has been imagined, by some, that the blood, in crossing the omentum, becomes loaded with fat. Secondly. The vena portæ ramifies in the liver, after the manner of an artery, and evidently communicates with the secretory vessels of the bile. Thirdly. It is larger than the hepatic artery; and more in proportion to the size of the liver; the hepatic artery seeming to be merely for the nutrition of the liver, as the bronchial artery is for

that of the lung.

In answer to these positions, it has been argued; that there seems to be no more reason why the bile should be formed from venous blood than the other fatty and oleaginous humours, - the marrow and fat for example, - which are derived from arterial blood. It is asked, too, whether, in point of fact, the blood of the vena portæ is more rich in carbon and hydrogen? and whether there be a closer chemical relation between the bile and the blood of the vena portæ, than between the fat and arterial blood?b The notion of the absorption of fat from the omentum, it is properly urged, is totally gratuitous. Secondly. The vena portæ does not exist in the invertebrated animals, and yet, in a number of them, there is an hepatic apparatus, and a secretion of bile. Thirdly. Admitting that the vena portæ is distributed to the liver after the manner of an artery; is it clear, it has been asked, that it is inservient to the biliary secretion? Fourthly. If the vena portæ be more in proportion to the size of the liver than the hepatic artery, the latter appears to bear a better ratio to the quantity of bile secreted: and, Lastly. It is probable, as has been shown in another place, that the liver has other functions connected with the portal system, in the admixture of heterogeneous liquids absorbed from the intestinal canal.c Similar views to those here expressed are entertained by Professor Giacomini, of Padua. He considers, 1. That the system of the vena portæ belongs less to the circulatory than to the absorbent system. 2. That the fluid which it contains is not real blood, but one that has not yet undergone the process of organic assimilation; and, 3. That the liver is an excretory organ, which deprives the fluid of the vena portæ of the principles that pass into the bile and are improper for assimilation.

d Encyclograph. des Sciences Médicales, Avril, 1840.

<sup>&</sup>lt;sup>2</sup> Rust's Magazine, B. xliv.; or Gazette Médicale, Aug. 15, 1835. See, also, Lafargue, Bullet. Médical. du Midi, Février, 1839.

<sup>&</sup>lt;sup>6</sup> R. Willis, Lond. and Edinb. Monthly Journ. of Med. Science, Sept. 1841, p. 628.
<sup>6</sup> Adelon, art. Foie, (Physiol.) in Dict. de Méd. ix. 193, Paris, 1824; and Physiol. de l'Homme, tom. iii. 505, Paris, 1829.

In the absence of accurate knowledge, derived from direct experiment, physiologists have usually embraced one or the other of these exclusive views. The generality, as we have remarked, assign the function to the vena portæ. Bichat, on the other hand, ascribes it to the hepatic artery. Broussaisa thinks it probable, that the blood of the vena portæ is not foreign to the formation of the bile, since it is confounded with that of the hepatic artery in the parenchyma of the liver; "but to say with the older writers, that the bile cannot be formed but by venous blood, is, in our opinion," he remarks, "to advance too bold a position, since the hepatic arteries send branches to each of the glandular acini, that compose the liver." Magendie likewise concludes, that nothing militates against the idea of both kinds of blood serving in the secretion; and that it is supported by anatomy; as injections prove, that all the vessels of the liver, - arterial, venous, lymphatic, and excretory, - communicate with each other. Mr. Kiernan, as we have seen, considers that the blood of the hepatic artery, after having nourished the liver, is inservient to the secretion, but not until it has become venous, and entered the portal veins. He, - with all those that coincide with him in the anatomical arrangement of these parts - denies that there is any communication between the ducts and the bloodvessels; and he asserts that if injections pass between them, it is owing to the rupture of the coats of the vessels. Experiments on pigeons, by M. Simon, of Metz, showed, that when the hepatic artery was tied, the secretion of bile continued, but that if the veins of the porta and the hepatic veins were tied, no trace of bile was subsequently found in the liver. It would thence appear, that in these animals the secretion of bile takes place from venous blood; but inferences from the ligature of those vessels have been very discordant. In two cases, in which Mr. Phillips tied the hepatic artery, the secretion of bile was uninterrupted: yet the same thing was observed in three other cases, in which the ligature was applied to the trunk of the vena portæ.

The view, that ascribes the bile to the hepatic artery, has always appeared to the author the most probable. It has all analogy in its favour. We have no disputed origin as regards the other secretions, excepting recently in the case of the urinary secretion. They all proceed from arterial blood; and function sufficient, we think, can be assigned to the portal system, without conceiving it to be concerned in the formation of bile. (See vol. i., p. 617.) We have, moreover, pathological cases, which would seem to show that bile can be formed from the blood of the hepatic artery. Mr. Abernethyd met with an instance, in which the trunk of the vena

<sup>&</sup>lt;sup>a</sup> Traité de Physiologie, &c., Drs. Bell and La Roche's translation, 3d edit. p. 456, Philad. 1832. See, also, Conwell's Treatise on the Liver, p. 54, London, 1835.

<sup>&</sup>lt;sup>b</sup> Carpenter, Human Physiology, § 662, Lond. 1842.

c Edinb. Med. and Surg. Journal, xc. 229; and Mayo's Outlines of Human Physio logy, 3d edit. p. 130, Lond. 1833.

d Philosoph. Transact. vol. lxxxiii.

portæ terminated in the vena cava; yet bile was found in the biliary ducts. A similar case is given by Mr. Lawrence; and the present Professor Monrob details a case communicated to him by the late Mr. Wilson, of the Windmill Street School, in which there was reason to suppose, that the greater part of the bile had been derived from the hepatic artery. The patient, a female, thirteen years old, died from the effects of an injury of the head. On dissection, Mr. Wilson found a large swelling at the root of the mesentery, consisting of several absorbent glands in a scrofulous state. Upon cutting into the mass, he accidentally observed a large vein passing directly from it into the vena cava inferior, which, on dissection, proved to be the vena portæ; and on tracing the vessels entering into it, one proved to be the inferior mesenteric vein: and another, which came directly to meet it, from behind the stomach, proved to be a branch of the splenic vein, but somewhat larger, which ran upwards by the side of the vena cava inferior, and entered that vein immediately before it passes behind the liver. Mr. Wilson then traced the branches of the trunk of the vessel corresponding to the vena portæ sufficiently far in the mesentery and mesocolon, to be convinced, that it was the only vessel that returned the blood from the small intestines, and from the cæcum and colon of the large intestines. He could trace no vein passing into the liver at the cavity of the porta; but a small vein descended from the little epiploon, and soon joined one of the larger branches of the splenic vein. The hepatic artery came off in a distinct trunk from the aorta, and ran directly to the liver. It was much larger than usual. The greater size of the hepatic artery. in this case, would favour the idea, that the arterial blood had to execute some office, that ordinarily belongs to the vena portæ. Was this the formation of bile? The case seems, too, to show, that bile can be formed from the blood of the hepatic artery. In Professor Hall's patient, (p. 267,) the vena portæ and its bifurcation were completely filled with encephaloid matter, so that no blood could pass through it to the liver; the secretion of the bile could not, consequently, be effected through its agency. It has been presumed, however, that in such cases, portal blood might still enter the liver through the extensive anastomoses, which Professor Retzius, of Stockholm, found to exist between the abdominal veins. That gentleman observed, when he tied the vena portæ near the liver, and threw a coloured injection into the portion below the ligature, that branches were filled, some of which, proceeding from the duodenum, terminated in the vena cava; whilst others, arising from the colon, terminated in the left emulgent vein. In subsequent investigations, he observed an extensive plexus of minute veins ramifying in the cellular tissue on the outer surface of the

Medico-Chirurgical Transact. iv. 174.
 Elements of Anatomy, Edinb. 1825.

e Ars Berättelse af Setterblad, 1835, s. 9; cited in Zeitschrift für die Gesammte Heilkunde, Feb. 1837, s. 251; also, Müller's Handbuch, u. s. w., Baly's translation, p. 185, Lond. 1837.

peritoneum, part of which was connected with the vena portæ, whilst the other part terminated in the system of the vena cava. In a successful injection, these veins were seen anastomosing very freely, in the posterior part of the abdomen, with the colic veins, as well as those of the kidneys, pelvis, and even with the vena cava. The arrangement, pointed out by Retzius, accounts for the mode in which the blood of the abdominal venous system reaches the cava, when the vena portæ is obliterated from any cause; and it shows the possibility of portal blood reaching the liver so as to be inservient to the biliary secretion, but does not, we think, exhibit its probability.

It would seem, then, that the portal system is not absolutely necessary to the formation of bile, inasmuch as it has occasionally been found absent; yet a recent writer considers it "a most puerile question" to ask whether the secretion can be effected from venous blood! "Had not," he adds, "secretion been destined to take place from the blood of the vena portarum, nature would not have been at the pains to distribute it through the liver; the peculiar arrangement is already an answer to the question; the end

of it is, as I have said, to economise arterial blood." Yet, as before remarked, a sufficient function can be assigned to the portal

system without supposing that it has any agency in the secretion of bile.

When bile is once secreted in the tissue of the liver, it is received into the minute excretory radicles, whence it proceeds along the ducts until it arrives, from all quarters, at the hepatic duct. difference of sentiment exists regarding the flow of the bile from the liver and gall-bladder into the duodenum. According to some, it is constantly passing along the choledoch duct; but the quantity is not the same during digestion as at other times. In the intervals, a part only of the secreted bile attains the duodenum; the remainder ascends along the cystic duct, and is deposited in the gall-bladder. During digestion, however, not only the whole of the secretion arrives at the duodenum, but all that which had been collected in the interval is evacuated into the intestine. In support of this view it is affirmed, that bile is always met with in the duodenum; that the gall-bladder always contains more bile when abstinence is prolonged, whilst it is empty immediately after digestion.

The great difficulties have been, to explain how the bile gets into the gall-bladder, and how it is expelled from that reservoir. In many birds, reptiles, and fishes, the hepatic duct and the cystic duct open separately into the duodenum; whilst ducts, called hepato-cystic, pass directly from the liver to the gall-bladder. In man, however, the only visible route, by which it can reach that reservoir, is by the cystic duct, the direction of which is retrograde; and, consequently, the bile has to ascend against gravity. The spiral

<sup>&</sup>lt;sup>2</sup> Dr. R. Willis, Lond. and Edinb. Monthly Journal of Med. Science, Sept. 1841, p. 628. See, also, Dr. Bostock, Ibid., May, 1842, p. 403.

valve of Amussat has been presumed to act like the screw of Archimedes, and to facilitate the entrance of the refluent bile, but this appears to be imaginary. It is, indeed, impossible to see any analogy between the corporeal and the hydraulic instrument. The arrangement of the termination of the choledoch duct in the duodenum has probably a more positive influence. The embouchure is the narrowest part of the duct, the ratio of its calibre to that of the hepatic duct having been estimated at not more than one to six, and to the calibre of its own duct as one to fifteen. This might render it impracticable for the bile to flow into the duodenum as promptly as it arrives at the embouchure; and, in this way collecting in the duct, it might reflow into the gall-bladder. Amussat, indeed, affirms, that this can be demonstrated on the dead body. By injecting water or mercury into the upper part of the hepatic duct, the injected liquid was found to issue both by the aperture into the duodenum, and by the upper aperture of the cystic duct'

into the gall-bladder.

With regard to the mode in which the gall-bladder empties itself during digestion, it is probably by a contractile action. We have seen, that it has not usually been admitted to possess a muscular coat, but that it is manifestly contractile. The chyme, as it passes into the duodenum, excites the orifice of the choledoch duct; this excitement is propagated along the ducts to the gall-bladder, which contracts; but according to Amussat, it does not evacuate its contents suddenly, for the different planes of the spiral valve are applied against each other, and only permit the flow to take place slowly. This he found was the case, in the subject, when water was injected into the gall-bladder, and pressed out through the cystic duct.a Other physiologists have presumed, that although the bile is secreted in a continuous manner, it only flows into the duodenum at the time of chylification; at other times, the choledoch duct is contracted, so that the bile is compelled to reflow through the cystic duct into the gall-bladder; and it is only when the gallbladder is filled, that it passes freely into the duodenum. Independently, however, of other objections to this view, vivisections have shown, that if the orifice of the choledoch duct be exposed, whatever may be the circumstances in which the animal is placed, the bile is seen issuing guttatim at the surface of the intestine. That the flow of bile from the gall-bladder, however, is dependent upon the presence of aliment in the intestines, is shown by the fact, that the reservoir is almost always found turgid in those who have died from starvation; the secretion formed at the ordinary slow rate having gradually accumulated for want of demand. This fact, it has been properly remarked, is important in juridical inquiries.b

The biliary secretion, which proceeds immediately from the liver, —hence called hepatic bile, differs from that obtained from the gall-

Adelon, Physiol. de l'Homme, iii. 494; and art. Foie, op. citat. Carpenter, Human Physiology, § 664, Lond. 1842.

bladder, which is termed cystic bile. The latter possesses greater bitterness, is thicker, of a deeper colour, and is that which has been usually analyzed. It is of a yellowish-green colour, viscid, and slightly bitter. Its chemical properties have been frequently examined; yet much is still needed, before we can consider the analysis satisfactory. It has been examined by Boerhaave, Verheyen, Baglivi, Hartmann, Macbride, Ramsay, Gaubius, Cadet, Fourcroy, Maclurg, Thénard, Berzelius, Chevreul, Leuret and Lassaigne, Frommherz and Gugert, Schultz, Vogel, John, Treviranus, Tiedemann and Gmelin, &c., &c. Thenard'sa analysis of 1100 parts of human bile is as follows: - water, 1000; albumen, 42; resinous matter, 41; yellow matter, 2 to 10; free soda, 5 or 6; phosphate, chloride of calcium, and sulphate of soda, phosphate of lime, and oxide of iron, 4 or 5. According to Chevallier, it contains also a quantity of picromel, now called bilin. Berzeliusb calls in question the correctness of Thénard's analysis, and gives the following: water, 908.4; picromel, 80; albumen, 3.0; soda, 4.1; phosphate of lime, 0.1; common salt, 3.4; phosphate of soda, with some lime, 1.0. The results of Dr. Davy'se analysis of healthy bile were as follows: - water, 86.0; resin of bile, 12.5; albumen, 1.5. Lastly, the experiments of Gmelin, for which he is highly complimented by Berzelius, d although Berzelius considers, that some of the products may have been formed by the reaction of elements upon each other - yielded the following results: - an odorous material, like musk; cholesterin; oleic acid; margaric acid; cholic acid; resin of bile; taurin (gallen-asparagin); picromel; colouring matter, now called biliverdin; osmazome; a substance, which, when heated, had the odour of urine; a substance resembling birdlime, gleadin; albumen (?); mucus of the gall-bladder; casein, or a similar substance; ptyalin, or a similar matter; bicarbonate of soda; carbonate of ammonia; acetate of soda; oleate, margarate, cholate, and phosphate of potassa and soda; chloride of sodium and phosphate of lime. Cadete considered bile as a soap with a base of soda, mixed with sugar of milk, - a view, which Raspail, f Demarcay, and others think harmonizes with observed facts. Every other substance met with in the bile, Raspail looks upon as accessary. Lastly, it has been more recently analysed by Mura-

Mém. de la Société d'Arcueil, i. 38, Paris, 1807.

b Med. Chirurgical Transactions, iii. 241.
c Monro's Elements of Anatomy, i. 579.

d Henle, art. Galle, in Encyclop. Wörterb. u. s. w., p. 126, and his Allgemein. Anat. u. s. w., or French translation by Jourdan, p. 80, Paris, 1843; art. Bile, by Mr. Brande, in Cyclop. of Anat. and Physiol. part iv. 374, Lond. 1835; Orfila, art. Bile, in Dict. de Médecine, i. 355, Paris, 1821; and Burdach's Physiologie als Erfahrungswissenschaft, v. 260, Leipz. 1835.

e Experiences sur la Bile des Hommes, &c. in Mém. de l'Académ. de Paris, 1767.

t Chimie Organique, p. 451, Edinb. 1833.

s Annal. der Pharmac. xxvii., cited by Liebig, Animal Chemistry, Webster's edit., p. 305, Cambridge, Mass. See, also, Prout, on Stomach and Renal Diseases, p. 510, 4th edit., Lond. 1843; and Thomson, Chemistry of Animal Bodies, p. 412, Edinb. 1843.

tori, who assigns it the following constituents; — water, 832; peculiar fatty matter, 5; colouring matter — biliverdin — 11; cholesterin combined with soda, 4; picromel of Thénard, 94.86; extract of flesh (Estratto di carne), 2.69; mucus, 37; soda, 5.14; phosphate of soda, 3.45; phosphate of lime, 3; and chloride of sodium, 1.86. The proportion of solid matter in the bile is usually from 9 to 12 per cent., nearly the whole of which consists of cholesterin and bilin. The cholesterin is almost altogether composed of carbon and hydrogen. Bilin contains nitrogen.

The specific gravity of bile, at 6° centigrade, according to Thénard, is 1.026. Schultz found that of an ox, after feeding, at 15°

to be 1.026; of a fasting animal, 1.030.

Hepatic and cystic bile do not appear to differ materially from each other, except in the greater concentration of the different elements in the latter. Leuret and Lassaigneb found them to be alike in the dog. Orfila, however, affirms, that human hepatic bile does

not contain picromel.

When bile is examined with the microscope, it is seen to contain a few, and but a few, globules of mucus, proceeding, according to Mandl, from the muciparous glands of the gall-bladder; lamellæ of cylinder-epithelium swimming in an amorphous liquid, and small yellowish globules. At times, some crystals of cholesterin are also observed in it.

The great uses of the bile have been detailed under the head of digestion. It has been conceived to be a necessary depurative excretion; separating from the blood matters that would be injurious if retained. This last idea is probable; but our knowledge of the precise changes, produced in the mass of blood by it, are extremely limited. The view has been ingeniously contended for by MM. Tiedemann and Gmelin, who regard the function of the liver to be supplementary to that of the lungs - in other words, to remove hydro-carbon from the system. The arguments, adduced in favour of their position, are highly specious, and ingenious. The resin of the bile, they say, abounds most in herbivorous animals, whose food contains a great disproportion of carbon and hydrogen. The pulmonary and biliary apparatuses are in different tribes of animals, and even in different animals of the same species, in a state of antagonism to each other. The size of the liver, and the quantity of bile are not in proportion to the quantity of food and frequency of eating, but inversely proportionate to the size and perfection of the lungs. Thus, in warm-blooded animals, which have large lungs, and live always in the air, the liver, compared with the body, is proportionally less than in those that live partly

Bulletino Mediche di Bologna, p. 160, Agosto et Settembre, 1836.

b Recherches, &c. sur la Digestion, Paris, 1825.

e Elém. de Chimie, Paris, 1817.

d Mandl, Manuel d'Anatomie générale, p. 501, Paris, 1843.

Die Verdaung nach Versuch. &c. traduit par Jourdan, Paris, 1827. See, also, Lafargue, Bullet. Medical du Midi, Février, 1839.

in water. The liver is proportionally still larger in reptiles, which have lungs with large cells incapable of rapidly decarbonizing the blood, — and in fishes, which decarbonize the blood but slowly by the gills, and, above all, in molluscous animals, which effect the same change very slowly, either by gills, or by small imperfectly developed lungs. Again, - the quantity of venous blood, sent through the liver, increases as the pulmonary system becomes less perfect. In the mammalia, and in birds, the vena portæ is formed by the veins of the stomach, intestines, spleen, and pancreas; in the tortoise, it receives also the veins of the hind legs, pelvis, tail, and the vena azygos: in serpents, it receives the right renal, and all the intercostal veins; in fishes the renal veins, and those of the tail and genital organs. Moreover, during the hibernation of certain of the mammalia, when respiration is suspended, and no food taken, the secretion of bile goes on. Another argument is deduced from the physiology of the fætus, in which the liver is proportionally larger than in the adult, and in which the bile is secreted copiously, as appears from the great increase of the meconium during the latter months of utero-gestation. Their last argument is drawn from pathological facts. In pneumonia and phthisis, the secretion of bile, according to their observations, is increased; in diseases of the heart the liver is enlarged; and in the morbus cæruleus, the liver retains its fætal proportion. In hot climates, too, where, in consequence of the greater rarefaction of the air, respiration is less perfectly effected than in colder, a vicarious decarbonization of the blood is established by an increased flow of bile. That the separation of the bile from the blood, however, is not an indispensable function, is shown by Dr. Blundell, who gives the cases of two children that lived for four months, apparently well fed and healthy, and on opening their bodies, it was found that the biliary ducts terminated in a cul-de-sac, and that consequently not a drop of bile had been discharged into the intestines. It is proper to remark, however, that azote is given off likewise in the bile, - being one of the constituents of bilin. Admitting, then, that the bile is essentially a depurative secretion, it by no means follows, that this depuration must be effected upon the blood of the hepatic artery. It is more probably exerted upon that of the portal system. The veins of the stomach and small intestines necessarily absorb much heterogeneous matter; this may be separated by the liver, along with other products that might be injurious if they passed into the blood.

The recent views of Liebigb on this function, as well as on that of the urinary secretion, are ingenious, and, if not true, are at least plausible. Venous blood, before reaching the heart, passes through the liver; arterial blood through the kidney; and both these

b Animal Chemistry, Webster's edit., p. 57, Cambridge, Mass. 1843.

<sup>\*</sup> Stokes, Theory and Practice of Medicine, Dunglison's American Medical Library Edition, p. 104, Philad. 1837.

organs separate from the blood the substances that are incapable of serving for the nutrition of the tissues. The compounds which contain the nitrogen or azote of the transformed tissues are collected in the urinary bladder, and, not being inservient to any further use, are expelled from the body. Those, again, which contain the carbon of the transformed tissues, are collected in the gall-bladder, in the form of a compound of soda, the bile, which is miscible with water in every proportion, and which, passing into the duodenum, mixes with the chyme. All those parts of the bile, which, during the digestive process, do not lose their solubility, return, during that process, into the circulation in a state of extreme division. The soda of the bile, and the highly carbonized portions, which are not precipitated by a weak acid, retain the capability of being taken up by the absorbents of the small and large intestines, - a capability which has been directly proved by the administration of enemata containing bile - the whole of the bile having disappeared along with the fluid injected into the rec-Liebig affirms, that the constituents of the bile cannot be recognized in the fæces of carnivorous animals; whence he infers that the whole of the bile has been reabsorbed; and, he believes, in order that its hydro-carbon may pass off by the lungs. This can scarcely, however, apply to man; and Liebig admits, that in the herbivora a certain portion of the elements of the bile can be discovered in the fæces. Certainly, a marked difference is observable in the fæces of a patient whose biliary ducts are obstructed. As to the precise change effected in the bile in order to fit it for being reabsorbed, Liebig leaves us wholly in the dark. His observations on this matter afford room for interesting reflection; but they can only at present be regarded in the light of suggestions.

If the excretion of the bile be prevented from any cause, we know that derangement of health is induced; but it is probable that its agency in the production of disease is much overrated; and that, as Broussais has suggested, the source of many of the affections termed bilious is in the mucous membrane lining the stomach and intestines; which, owing to the heterogeneous matters constantly brought into contact with it, must be peculiarly liable to be morbidly affected. When irritation exists there, we can easily understand how the secretion from the liver may be consecutively modified; the excitement spreading directly along

the biliary ducts to the secretory organ.

# e. Secretion of Urine.

This is the most extensive secretion accomplished by any of the glandular structures of the body, and is essentially depurative; its suppression gives rise to formidable evils. The apparatus consists of the kidneys, which secrete the fluid; the ureters, which convey the urine to the bladder; the bladder itself, which serves as a re-

servoir for the urine; and the urethra, which conveys the urine

externally. These will require a distinct consideration.

The kidneys are two glands situate in the abdomen; one on each side of the spine, (Fig. 193, K, K,) in the posterior part of the lumbar region. They are without the cavity of the peritoneum, which covers them at the anterior part only, and are situate in the midst of a considerable mass of adipous cellular tissue. The right kidney is nearly an inch lower down than the left, owing to the thick posterior margin of the right lobe of the liver pressing it downwards. Occasionally, there is but one kidney; at other times, three have been met with. They have the form of the haricot or kidney-bean, which has, indeed, been called after them; and are situate vertically - the fissure being turned inwards. If we compare them with the liver, their size is by no means in proportion with the extensive secretion effected by them. Their united weight does not amount to more than six or eight ounces. Of 65 male kidneys, weighed by Dr. John Reid, the average weight was found to be 5 oz. 7 dr. for the right kidney; 5 oz. 111 dr. for the left. Of 28 female kidneys, the right weighed 4 oz. 13 dr.; the left, 5 oz. 2 dr. The left kidney generally weighs more than the right at all ages. They are hard, solid bodies, of a brown colour. The sanguiferous vessels, which convey and return the blood to them, as well as the excretory duct, communicate with the kidney at the fissure.

The anatomical constituents of these organs are; — 1. The renal artery, which arises from the abdominal aorta at a right angle,

Fig. 203.

Section of a Kidney. Supra-renal Capsule.

1. Supra-renal capsule. 2. Cortical or vascular portion. 3, 3. Tubular portion. 4, 4. Calices receiving the apices of cones. 5, 5, 5. Infundibula. 6. Pelvis. 7. Ureter.

and, after a short course, enters the kidney, ramifying in its substance. 2. The excretory ducts, which arise from every part of the tissue, in which the ramifications of the renal artery terminate, and end in the pelvis of the kidney. (Fig. 203.) 3. The renal veins, which receive the superfluous blood, after the urine has been separated from it, and terminate in the renal or emulgent vein, which issues at the fissure and opens into the abdominal vena cava. 4. Of lymphatic vessels, arranged in two planes - a superficial and a deep-seated, which terminate in the lumbar glands. 5. Of nerves, which proceed from the semilunar ganglion, solar plexus, &c., and which surround the renal artery as with a network, following


it in all its ramifications. 6. Of cellular membrane, which, as in

Lond. and Edinb. Monthly Journ. of Med. Sciences, April, 1843, p. 323.

every other organ, binds the parts together. These anatomical elements, by their union, constitute the organ as we find it.


When the kidney is divided longitudinally, it is seen to consist of two substances, which differ in their situation, colour, consistence, and texture. The one of these, and the more external, is called the cortical or glandular substance. It forms the whole circumference of the kidney; is about two lines in thickness; of less consistence than the other; of a pale red colour; and receives almost entirely the ramifications of the renal artery. The other and innermost is the tubular, medullary, uriniferous, conoidal, or radiated substance. It is more dense than the other; less red; and seems to be formed of numerous minute tubes, which unite in conical bundles of unequal size, and the base of which is turned towards the cortical portion; the apices forming the papillæ or mammillary processes, and facing the pelvis of the kidney. The papillæ vary in number, from five to eighteen; are of a florid colour; and upon their points or apices are the terminations of the uriniferous tubes large enough to be distinguished by the naked eye. Around the root of each papilla a membranous tube arises called calix or infundibulum: this receives the urine from the papilla, and conveys it into the pelvis of the kidney, which may be regarded as the commencement of the ureter.

The cortical part of the kidney is the most vascular; and the plexus formed by the tubuli uriniferi appears to come there in closest relation with the plexus formed by the renal capilla- a. ries. In the marginal illustrations of the two portions as they appear in the new-born infant. A exhibits them of the natural size: a a, are the corpora Malpighiana or Malpighian bodies appearing as points in the cortical substance; b, is one of the papillæ. These Malpighian bodies are scattered through the plexus formed by the bloodvessels and uriniferous tubes. Each one of these, when examined by a high magnifying power, is found to consist of a convoluted mass of minute bloodvessels. In these - it was at one time supposed - the uriniferous tubes originate; but the examinations of Müller and Huschke have seemed to show, that they are Portion of Kidneys of New-born Infants. only capable of injection from the arteries or veins. They are found in the kidneys of most, if not all, A small portion of A of the vertebrata. In the cortical substance, ac-magnified a a. Corpora Maipighiana b Tubuli cording to Wagner, the tubuli can be traced, al- uriniferi. -(Wagner.) though with difficulty, winding among the vascular plexuses, or skeins, mostly looped towards the margin of the organ, and running



. J. Müller, Baly's edit. or Bell's Amer. abridgment, p. 435, Philad. 1843; Carpenter, Human Physiology, § 667, Lond. 1842.

Elements of Physiology, by R. Willis, § 193, Lond. 1842.



Small Portion of Kidney magnified 60 Diameters.

a. Coccal extremity of a tubulus; bb, loops of tubuli; cc, bifurcated of the gland, contubuli; def, tubuli converging towards the papille; ggs, corpora ducting the blood-Malpighiana; h, arterial trunk.

organ, and constituting small chambers in the cortical portion in,

Lond. and Edinb. Monthly Journ. of Med. Science, May, 1842.

into one another, or having blind or cœcal extremities; more rarely enlarged and club-shaped as in Fig. 12, and occasionally cleft (Fig. 205). entire cortical substance, according to Wagner, consists of convolutions of the uriniferous tubes; which present a nearly uniform diameter, and, on an average, may be from about the 60th to the 50th of a line. Mr. Goodsir, a however, without denying the existence of occasional blind extremities of the tubuli uriniferi, the result probably, he thinks, of arrested development, states that he has never seen the ducts terminate in this way. He describes a fibrocellular or framework, which, pervading every part of the gland, and particularly its cortical portion, performs the same office in the kidney, which the capsule of Glisson does in the liver, forming a basis of support to the delicate structure vessels through the each of which a single ultimate coil or loop of the uriniferous ducts is lodged. Mr. Goodsir believes, that the urine is formed at first within the epithelium cells of the ducts, and that these burst, dissolve, and throw out their contents, and are succeeded by others, which perform the same functions. The urine of man has not been detected by Mr. Goodsir within the cells, which line the ducts, but he has submitted to the Royal Society of Edinburgh a memoir, already referred to (p. 232), in which he has endeavoured to show, that the urine, bile and milk, as well as the other more important secretions in the lower animals, are formed within the nucleated cells of the ducts themselves; and he is of opinion, that the urine of man is poured at first into the cavities of the nucleated cells of the human kidney. More recently, Mr. Bowmana has affirmed, that the kidney is furnished with a true portal system, and that the urine, like the bile, is secreted — in part at least from blood, which is traversing at the time a second set of capillaries. According to him, each of the exceedingly tortuous and convoluted urinary conduits terminates, at its final extremity, by a contracted neck, which leads into a little chamber or cyst, in which is contained the true glandule of Malpighi, which consists of a tuft or coil of capillary bloodvessels, totally naked, that originates in one of the ultimate branches of the renal artery, and terminates in an efferent vessel. Several of these latter form, by their anastomosing ramifications, the plexus that surrounds each urinary conduit and tubule; the urinary conduits being lined by thick epithelium, and their necks furnished with vibratile cilia. All the blood of the renal artery, according to him, - with the exception of a small quantity distributed to the capsule, surrounding fat, and the coats of the larger vessels - enters the capillary tufts of the corpora Malpighiana; thence it passes into the capillary plexus surrounding the uriniferous tubes, and finally leaves the organ through the branches of the renal vein. Thus, there are in the kidney two perfectly distinct systems of capillary vessels; the first, that inserted into the dilated extremities of the uriniferous tubes, and in immediate connection with the arteries - the Malpighian bodies: - the second, that enveloping the convolutions of the tubes, and communicating directly with the veins. The efferent vessels of the Malpighian bodies, that carry the blood between these two systems, are termed by Mr. Bowman the portal system of the kidney. The views of Mr. Bowman have been embraced by some histologists; but they stand in need of further observation.

In the quadruped, each kidney is made up of numerous lobes, which are more or less intimately united according to the species. In birds, the kidneys consists of a double row of distinct, but connected, glandular bodies, placed on both sides the lumbar vertebræ.


The ureter is a membranous duct, which extends from the

Proceedings of the Royal Society, No. lii. Feb. 3, 1842; and Philos. Transactions, Pt. 1, p. 57, Lond. 1842.

b Aldridge, Dublin Journal of Med. Science, cited in Amer. Journ. of the Medical Sciences, Oct. 1843, p. 445.

kidney to the bladder. It is about the size of a goosequill; de-

Fig. 206.



A side view of Viscera of Male Pelvis in sitû.

1. Divided surface of os pubis. 2 Divided surface of sacrum. 3. Body of bladder. 4. Its fundus; from apex is seen passing upwards the urachus. 5. Base of bladder. 6. Ureter. 7. Neck of bladder. 8, 8. Pelvic fascia; fibres immediately above 7 are given off from pelvic fascia and represent anterior ligaments of bladder. 9. Prostate gland. 10. Membranous portion of urethra, between two layers of deep perineal fascia. 11. Deep perineal fascia formed of two layers. 12. One of Cowper's glands between two layers of deep perineal fascia, and beneath membranous portion of urethra. 13. Bulb of corpus spongiosum. 14. Body of corpus spongiosum. 15. Right crus penis. 16. Upper part of first portion of rectum. 17. Recto-vesical fold of peritoneum. 18. Second portion of rectum. 19. Right vesicular seminalis. 20. Vas deferens. 21. Rectum covered with descending layer of pelvic fascia, just as it is making its bend backwards to constitute third portion. 22. A part of levator ani muscle investing lower part of rectum. 23. External sphincter ani. 24. Interval between deep and superficial perineal fascia; they are seen to be continuous beneath the figure. —(Wilson.)

scends through the lumbar region; dips into the pelvis by crossing in front of the primitive iliac vessels and the internaliliac; crosses the vas deferens at the back of the bladder, and, penetrating that viscus obliquely, terminates by an orifice, ten twelve lines behind that of the neck of the bladder. At first, it penetrates two of the coats only of that viscus; running for the space of an inch between the mucous and muscular coats, and then entering the cavity. The ureters have two

coats. The outermost is a dense fibrous membrane; the innermost a thin mucous layer, continuous at its lower extremity with the inner coat of the bladder; and at the upper end supposed, by some, to be reflected, over the papillæ, and even to pass for some distance into the tubuli uriniferi.

The bladder is a musculo-membranous sac, situate in the pelvis; anterior to the rectum, and behind the pubes. Its superior end is called the upper fundus; and the lower end, the inferior fundus or bas-fond; the body being situate between the two. The part where it joins the urethra is the neck. The shape and situation of the organ are influenced by age and by sex. In very young infants, it is cylindroid, and rises up almost wholly into the abdomen. In the adult female, who has borne many children, it is nearly spherical; has its greatest diameter transverse, and is more capacious than in the male. Like the other hollow viscera, the bladder consists of several coats. 1. The peritoneal coat, which covers only the fundus and back part. Towards the lower por-

tion the organ is invested by cellular membrane, which takes the place of the peritoneal coat of the fundus. This tissue is very loose, and permits the distension and contraction of the bladder. 2. The muscular coat is very strong; so much so, that it has been classed amongst the distinct muscles, under the name detrusor urinæ. The fibres are pale, and pass in various directions. Towards the lower part of the bladder, they are particularly strong; arranged in fasciculi, and form a kind of network of muscles inclosing the bladder. In cases of stricture of the urethra, where much effort is necessary to expel the urine, these fasciculi acquire considerable thickness and strength. 3. The mucous or villous coat is the lining membrane, which is continuous with that of the ureters and urethra, and is generally rugous, in consequence of its being more extensive than the muscular coat without. It is furnished with numerous follicles, which secrete a fluid to lubricate Towards the neck of the organ, it is thin and white, though reddish in the rest of its extent. A fourth coat, called the cellular, has been reckoned by most anatomists, but it is nothing more than cellular tissue uniting the mucous and muscular coats. The part of the internal surface of the bladder, situate immediately behind and below its neck, and occupying the space between it and the orifices of the ureters, is called the vesicle triangle, trigonus Lieutaudi or trigone vésical. The anterior angle of the triangle looks into the orifice of the urethra, and is generally so prominent, that it has obtained the name uvula vesicæ. It is merely a projection of the mucous membrane, dependent upon the subjacent third lobe of the prostate gland, which, in old people, is frequently enlarged, and occasions difficulty in passing the catheter. The neck of the bladder penetrates the prostate gland, but, at its commencement, it is surrounded by loose cellular tissue, containing a very large and abundant plexus of veins. The internal layer of muscular fibres is here transverse; and they cross and intermix with each other, in different directions, forming a close, compact tissue, which has the effect of a particular apparatus for retaining the urine, and has been called the sphincter. Anatomists have not usually esteemed this structure to be distinct from the muscular coat at large; but Sir Charles Bella asserts, that if we begin the dissection by taking off the inner membrane of the bladder from around the orifice of the urethra, a set of fibres will be discovered, on the lower half of the orifice, which, being carefully dissected, will be found to run in a semicircular form around the urethra. These fibres make a band of about half an inch in breadth, particularly strong on the lower part of the opening; and having ascended a little above the orifice, on each side, they dispose of a portion of their fibres in the substance of the bladder.

<sup>\*</sup> Anatomy and Physiol. 5th Amer. Edit., by Dr. Godman, ii. 375, New York, 1829. See, also, on this subject, Dr. Horner, Special Anatomy and Histology, vol. ii., 6th edit., Philad. 1843; and Dr. Goddard, in Wilson's Anatomist's Vade Mecum, Amer. Edit., p. 531, Philad. 1843.

A smaller and somewhat weaker set of fibres will be seen to complete their course, surrounding the orifice on the upper part. The arteries of the bladder proceed from various sources, but chiefly from the umbilical and common pudic. The veins return the blood into the internal iliacs. They form a plexus of considerable size upon each side of the bladder, particularly about its neck. The lymphatics accompany the principal veins of the bladder, and, at the under part and sides, pass into the iliac glands. The nerves

are from the great sympathetic and sacral.

The urethra is the excretory duct of the bladder. It extends, in the male, from the neck of the bladder to the extremity of the gland; and is from seven to ten inches in length. In the female it is much shorter. The male urethra has several curvatures in the state of flaccidity of the penis; but is straight, or nearly so, if the penis be drawn forwards and upwards, and if the rectum be empty. The first portion of this canal, which traverses the prostate gland, is called the prostatic portion. Into it open, - on each side of a caruncle, called the verumontanum, caput gallinaginis or crista urethralis, - the two ejaculatory ducts, those of the prostate, and a little lower, the orifice of Cowper's glands. Between the prostate and the bulb is the membranous part of the urethra, which is eight or ten lines long. The remainder of the canal is called the corpus spongiosum or spongy portion, because surrounded by an erectile spongy tissue. It is situate beneath the corpora cavernosa, and passes forward to terminate in the glans; the structure of which will be considered under Generation. At the commencement of this portion of the urethra is the bulb of the urethra, Fig. 206; the structure of which resembles that of the corpora cavernosa of the penis - to be described hereafter. The dimensions of the canal are various. At the neck of the bladder, it is considerable; behind the caput gallinaginis it contracts, and immediately enlarges in the forepart of the prostate. The membranous portion is narrower; and, in the bulb, the channel enlarges. In the body of the penis, it diminishes successively, till it nearly attains the glans, when it is so much increased in size as to have acquired the name fossa navicularis. At the apex of the glans it terminates by a short vertical slit. Mr. Shawa has described a set of vessels, immediately on the outside of the internal membrane of the urethra, which, when empty, are very similar, in appearance, to muscular fibres. These vessels, he remarks, form an internal spongy body, which passes down to the membranous part of the urethra, and forms even a small bulb there. Dr. Horner, however, says, that this appeared to him to be rather the cellular membrane connecting the canal of the urethra with the corpus spongiosum. The whole of the urethra is lined by a very vascular and sensible mucous membrane, which is continued from

Manual of Anatomy, ii. 118, Lond. 1822.

b Lessons in Practical Anatomy, 3d edit. p. 272, Philad. 1836; and Amer. Journ. of the Medical Sciences, p. 538, for Feb. 1832.

the inner coat of the bladder. It has, apparently, a certain degree of contractility, and therefore, by some anatomists, is conceived to possess muscular fibres. Sir Everard Home, from the results of his microscopical observations, is disposed to be of this opinion. This is, however, so contrary to analogy, that it is probable the fibres may be seated in the tissue surrounding it. The membrane contains numerous follicles, and several lacunæ, one or two of which, near the extremity of the penis, are so large as occasionally to obstruct the catheter, and to convey the impression that a stricture exists.

The prostate and the glands of Cowper, being more concerned

in generation, will be described hereafter.

There are certain muscles of the perineum, that are engaged in the expulsion of the urine from the urethra; and some of them in defecation, and in the evacuation of the sperm likewise; as the acceleratores urinæ or bulbo-urethrales, which propel the urine or semen forward; the transversus perinei or ischio-perinealis, which dilates the bulb for the reception of the urine or semen; the sphincter ani, which draws down the bulb, and thus aids in the ejection of the urine or sperm; and the levator ani, which surrounds the extremity of the rectum, the neck of the bladder, the membranous portion of the urethra, the prostate gland, and a part of the vesiculæ seminales, and assists in the evacuation of the bladder, vesiculæ seminales, and prostate. A part of the levator, which arises from the pubis and assists in inclosing the prostate gland, is called by Sömmering compressor prostatæ. Between the membranous part of the urethra, and that portion of the levator ani which arises from the inner side of the symphysis pubis, a reddish, cellular, and very vascular substance exists, which closely surrounds the canal, has been described by Mr. Wilson under the name compressor urethræ, and is termed, by some of the French anatomists, muscle de Wilson. By many, however, it is considered to be a part of the levator ani. Amussat asserts, that the membranous part of the urethra is formed, externally, of muscular fibres, which are susceptible of energetic contraction, and Magendieconfirms his assertion.

With regard to the urinary organs of the female:— the kidneys and ureters have the same situation and structure as those of the male. The bladder, also, holds the same place behind the pubis, but rises higher when distended. It is proportionally larger than that of the male, and is broader from side to side, thus allowing the greater retention to which females are often necessitated. The urethra is much shorter, being only about an inch and a half, or two inches long, and it is straighter than in the male, having only a slight curve downwards between its extre-

<sup>\*</sup> For Dr Horner's views on the origin of the acceleratores urinæ, see his Special Anatomy and Histology 6th edit., vol. ii., Philad. 1843.

b Lectures on the Structure and Physiology of the Urinary and Genital Organs, Lond. 1821. Precis, &c., ii., 472.

VOL. II. - 25

mities, and passing almost horizontally under the symphysis of the pubis. It has no prostate gland, but is furnished, as in the male, with follicles and lacunæ, which provide a mucus to lubricate it.

In birds in general, and in many reptiles and fishes, the urine, prior to expulsion, is mixed with the excrement in the cloaca. Nothing analogous to the urinary organs has been detected in the lowest classes of animals, although in the dung of the caterpillars

of certain insects, traces of urea have been met with.

The urine is separated from the blood in the kidneys; and it has until recently been the universal belief, that it is secreted from arterial blood; Mr. Bowman, however, in accordance with his views on the minute anatomy of the kidney, already given, (p. 285,) has attempted to show, that it is separated from venous blood. His main conclusions are as follows: - First. The epithelium, lining the tubes, is the proper organ that secretes the characteristic products of the urine from the blood; and it does this by first assimilating them into its own substance, and afterwards giving them upon its free surface. Secondly. These proper urinous products require for their solution a large quantity of water. Thirdly. This water is furnished by the Malpighian tufts of capillaries, placed at the extremity of the uriniferous tubes; and Fourthly. A farther use of the Malpighian bodies seems to be that of sharing in regulating the amount of water in the body. He thus makes a striking analogy between the liver and kidneys both in structure and function; and expresses his belief, first, that diuretic medicines act specially on the Malpighian bodies, and that many substances, especially salts, which, when taken into the system, have a tendency to pass off by the kidneys with rapidity, in reality escape through the Malpighian bodies; secondly, that certain morbid products occasionally found in the urine, such as sugar, albumen, and the red particles of the blood, also, in all probability, pass off through this bare system of capillaries. According to Raspail, the urine is a kind of caput mortuum, ejected into the urinary bladder by those glands. They separate carbon and nitrogen from the kidneys, in the form of cyanogen, which unites with oxygen to produce cyanic acid; and this combines with ammonia — itself a compound of nitrogen and hydrogen — to form urea, which is the characteristic element of the urinary secretion; and they separate and excrete superfluous fluid from the body. The proofs of this separation are easy and satisfactory; but with regard to the mode in which the operation is effected, we are in the same darkness that hangs over the glandular secretions in general. The transformation must, however, occur in the cortical part of the organ; for the tubular portion seems to consist only of a collection of excretory ducts; and, if we cut into it, urine oozes out.

The urinary secretion takes place continuously. If a catheter

A Chimie Organique, p. 505, Paris 1833.

b Carpenter, Human Physiology, § 668, Lond, 1842.

be left in the bladder, the urine drops constantly; and in cases of exstrophia of the bladder - a faulty conformation, in which the organ opens above the pubes, so that a red mucous surface, formed by the inner coat of the bladder, is seen in the hypogastric region, in which two prominences are visible, corresponding to the openings of the ureters - the urine is seen to be constantly passing out at these openings.a After the secretion has been effected in the cortical substance, it flows through the tubular portion, and issues guttatim through the apices of the papillæ into the pelvis of the kidney, whence it proceeds along the ureter to the bladder. When the uriniferous cones are slightly compressed, the urine issues in greater quantity, but, instead of being limpid, as when it flows naturally, it is thick and troubled. Hence a conclusion has been drawn, that it is really filtered through the hollow fibres of the medullary or tubular portion. If this were the case, what must become of the separated thick portion? Ought not the tubes to become clogged up with it? And is it not more probable, that compression, in this case, forces out with the urine some of the blood that is connected with the nutrition of the organ? The fresh secretion constantly taking place in the kidney causes the urine to flow along the tubuli uriniferi to the pelvis of the organ, whence it proceeds along the ureter, if we are in the erect attitude, by virtue of its gravity; the fresh fluid, too, continually secreted from the kidney, pushes on that which is before it; and, moreover, there is not improbably some degree of contractile action exerted by the ureters themselves; although, as in the case of the excretory ducts in general, such a power has been denied them. These are the chief causes of the progression of the urine into the bladder, which is aided by the pressure of the abdominal contents and muscles, and, it is supposed, by the pulsation of the renal and iliac arteries; but the agency of these must be trivial.

The orifices of the ureters form the posterior angles of the trigone vésical, and are contracted somewhat below the size of the ducts themselves. They are said, by Sir Charles Bell, to be furnished with a small fasciculus of muscular fibres, which runs backwards from the orifice of the urethra, immediately behind the lateral margins of the triangle, and, when it contracts, stretches the orifice of the ureter so as to permit the urine to enter the bladder with facility. As the urine enters, it gradually distends the organ until the quantity has attained a certain amount. It cannot reflow by the ureters, on account of the smallness of their orifices and their obliquity; and as the bladder becomes filled,—owing to the duct passing for some distance between the muscular and mucous coats,—the sides are pressed against each other, so that the cavity is obliterated. (Fig. 207.) As, however, the ureters have a ten-

<sup>\*</sup> See note of a case of this kind, by the author, in Amer. Med. Intelligencer, i. 137; and another, by Dr. Pancoast, ibid. p. 147, Philad. 1838.

b Anatomy and Physiology, 5th American Edit., by Godman, ii. 381, New York, 1827.

dency to lose this obliquity of insertion, in proportion as the bladder is emptied, the two bands of muscular fibres which run from the back of the prostate gland to the orifices of the ureters, not only assist in emptying the bladder, but, at the same time, pull

Fig. 207.

C. Vesical orifice of the ureter.

down the orifices of the ureters, and thus tend to preserve the obliquity.2 Moreover, when we are in the erect attitude, the urine would have to enter the ureters against gravity. These obstacles are so effective, that if an injection be thrown forcibly and copiously through the urethra into the bladder, it does not enter the ureters. On the other hand, equally powerful impediments exist to its being discharged through the urethra. The inferior fundus of the bladder is situate lower than the neck; and the sphincter presents a degree of resistance, which requires the bladder to contract forcibly on its contents, aided by the abdomi-Entrance of the Ureter into the Bladder. A. Cavity of the bladder. B. Ureter. nal muscles, to overcome it. Magendieb considers the contraction of the

levatores ani to be the most efficient cause of the retention of the urine; the fibres which pass around the urethra pressing its

sides against each other and thus closing it.

The urine accumulates in the bladder until the desire arises to expel it: the number of times that a person in health and in the middle period of life, discharges it in the twenty-four hours, varies; whilst some evacuate the bladder but twice, others may be compelled to do so as many as twelve or fourteen times. Nine times, according to Dr. Thomas Thomson, is a common number. The quantity, too, discharged at a time varies. The greatest quantity observed by Dr. Thomson was 25½ cubic inches, or somewhat less than a pint, the most common quantity being from seven to nine cubic inches. During the stay of the urine in the bladder, it is believed to be deprived of some of its more aqueous portions by absorption, and to become of greater specific gravity, and more coloured; it is here that those depositions are apt to take place which constitute calculi; although we meet with them in the kidneys and ureters also. As in every excretion, a sensation first arises, in consequence of which the muscles required for the ejection of the secreted matter are called into action. This sensation arises whenever the urine has accumulated to the necessary extent, or when it possesses irritating qualities, owing to extraneous substances being contained in, or deposited from it; or if the bladder be unusually irritable from any morbid

a Sir C. Bell, op. cit., and in Medico-Chirurgical Transactions, vol. iii.

b Précis, &c. edit. cit. ii. 473. British Annals of Medicine, p. 6, Jan. 1837.

cause, the sensation may be repeatedly - nay, almost incessantly - experienced. The remarks, that have been made on the sensations accompanying the other excretions, are equally applicable here. The impression takes place in the bladder; such impression is conveyed to the brain, which accomplishes the sensation; and, consecutively, the muscles, concerned in the excretion, are called into action by volition. Physiologists have differed regarding the power of volition over the bladder. Some have affirmed that it is as much under cerebral control as the muscles of locomotion; and they have urged, in support of this view, that the bladder receives spinal nerves, which are voluntary; that it is paralysed in affections of the spinal marrow, like the muscles of the limbs: and that a sensation, which seems destined to arouse the will, is always the precursor of its action. Others again have denied, that the muscular fibres of the bladder are contractile under the will; and they adduce the case of other reservoirs, - the stomach and the rectum, for example, - whose influence in excretion we have seen to be involuntary; as well as the fact that we no more feel the contraction of the bladder than we do that of the stomach or intestines; and they affirm, that the action of the bladder itself has been confounded with that of the accessory muscles, which are manifestly under the influence of the will, and are important agents in the expulsion of the fluid from the bladder. The views, last expressed, appear to be most accurate, and the catenation of phenomena seems to be as follows: the sensation to expel the urine arises; the abdominal muscles are thrown into contraction by volition; the viscera are thus pressed down upon the pelvis; the muscular coat of the bladder is, at the same time, stimulated to contraction; the levatores ani and the sphincter fibres are relaxed, so that the resistance of the neck of the organ is diminished, and the urine is forced out through the whole extent of the urethra, being aided in its course, especially towards the termination, by the contractile action of the urethra itself, as well as by the levatores ani and acceleratores urinæ muscles. These expel the last drops by giving a slight succussion to the organ, and directing it upwards and forwards; an effect which is aided by shaking the organ to free it from the drops that may exist in the part of the canal near its extremity. The gradually diminishing jet, which we notice, as the bladder is becoming empty, indicates the contraction of the muscular coat of the organ; whilst the kind of intermittent jet, coincident with voluntary muscular exertion, indicates the contraction of the urethral muscles. When we feel the inclination to evacuate the bladder, and do not wish to obey it, the same muscles, - the levatores ani, the acceleratores urinæ, and the fibres around the membranous portion of the urethra and the neck of the bladder, - are thrown into contraction, and resist that of the bladder.

Such is the ordinary mechanism of the excretion of urine. The contraction of the bladder is, however, of itself sufficient to expel

its contents. Magendie<sup>a</sup> affirms, that he has frequently seen dogs pass urine when the abdomen was opened, and the bladder removed from the influence of the abdominal muscles; and he farther states, that if, in a male dog, the bladder, with the prostate and a small portion of the membranous part of the urethra, be removed from the body, the bladder will contract after a few moments, and project the urine, with an evident jet, until it is en-

tirely expelled.

Urine — voided in the morning, urina sanguinis, by a person who has eaten heartily, and taken no more fluid than sufficient to allay thirst — is a transparent, limpid fluid, of an amber colour, saline taste, and a peculiar odour. Its specific gravity is estimated by Chossat at from 1.001 to 1.038; by Cruikshank and Wagner, b from 1.005 to 1.033; by Prout, from 1.015 to 1.025; by Gregory. from 1.005 to 1.033; by Christison, on the average, 1.024 or 1.025; by F. D'Arcet, from 1.001 to 1.060 [?]e by Rayer, on the average, 1.018; and by Dr. Bostock and Martin Solon, 1.020; by Elliotson, from 1.015 to 1.025; and Dr. Thomson found it in an individual, from 50 to 60 years of age and in perfect health, to be, on the average, 1.013; the lowest specific gravity, during ten days, being 1 004; and the highest, 1.026. Dr. Thomson has published some tables showing the quantity of urine passed at different times during ten days by the individual in question, and the specific gravity of each portion. These tables do not accord with the opinion generally entertained, that the heaviest urine is voided on rising in the morning. No generalization can, indeed, be made on this subject. The temperature of the urine when recently voided, varied in one case from 92° to 95°. It is slightly acid, for it reddens vegetable blues. Although at first quite transparent, it deposits an insoluble matter on standing; so that urine, passed at bed-time, is found to have a light cloud floating in it by the following morning. This substance consists, in part, of mucus from the urinary passages; and, in part, of the super-urate of ammonia, which is much more soluble in warm than in cold water.

The following table, drawn up by M. Becquerel as far as 1032, and completed from the observations of Dr. Golding Bird, k exhibits, at a single inspection, the amount of solids and water present in 1000 grains of urine of any particular density; and from the quantity of urine passed in twenty-four hours it is easy to calculate how much solid matter the patient is parting with in that period.

- <sup>a</sup> Précis, ii. 474. b Elements of Physiology, by R. Willis, § 200, Lond. 1842.
- Burdach, Die Physiologie als Erfahrungswissenschaft, v. 271, Leipz. 1835.
   On Granular Degeneration of the Kidneys, p. 34, Edinb. 1839; or Dunglison's American Med. Library edit., Philad. 1839.

c L'Expérience, No. lv., Aug. 1838.

Traité des Maladies des Reins, &c., tom. i., Paris, 1839.

5 De l'Albuminurie, ou Hydropisie Causée par Maladie des Reins, Paris, 1838.

Human Physiology, p. 293, Lond. 1835.
British Annals of Medicine, p. 5, Jan. 1837.
Lond. Med. Gazette, Feb. 10, 1843, p. 678.

Op. citat. p. 6.

Density.	Water in 1000 grains.	Solids in 1000 grains.	Density.	Water in 1000 grains.	Solids in 1030 grains.
1001	998-35	1.65	1022	963-7	36-3
1002	996-7	3.3	1029	960-4	39.6
1004	993-4	6.6	1026	957-1	42.9
1006	990-1	9.9	1028	953.8	46.2
1008	986-8	13.2	1030	950.5	49.5
1010	983-5	16.5	1032	947-2	52.8
1012	980-2	19-8	1034	943-9	56.1
1014	976-9	23.1	1036	940-6	59.4
1016	973-6	26.4	1038	937-3	62.7
1018	970-3	29.7	1040	934.	66.
1020	967-	33.	1042	930-7	69.3
	940 miles 58	sanduar L	1044	927-4	72.6
	Andrewson I	and annual	1046	924-1	75-9

The urine is extremely prone to decomposition. When kept for a few days, it acquires a strong smell, which, being sui generis, has been called urinous; and as the decomposition proceeds, the odour becomes extremely disagreeable. The urine, as soon as these changes commence, ceases to have an acid reaction, and the earthy phosphates are deposited. In a short time, a free alkali makes its appearance; and a large quantity of the carbonate of ammonia is generated. These phenomena are owing to the decomposition of urea, which is almost wholly resolved into carbonate of ammonia.

Dr. Henrya affirms, that the following substances have been satisfactorily proved to exist in healthy urine, - water, free phosphoric acid, phosphate of lime, phosphate of magnesia, fluoric acid, uric acid, benzoic acid, lactic acid, urea, gelatin, albumen, lactate of ammonia, sulphate of potassa, sulphate of soda, fluate of lime, chloride of sodium, phosphate of soda, phosphate of ammonia, sulphur and silex. One of the most elaborate analyses has been given by Berzelius. He states it to consist - in 1000 parts - of water, 933.00; urea, 30.10; sulphate of potassa, 3.71; sulphate of soda, 3.16; phosphate of soda, 2.94; chloride of sodium, 4.45; phosphate of ammonia, 1.65; muriate of ammonia, 1.50; free lactic acid, lactate of ammonia, animal matter, soluble in alcohol, and urea not separable from the preceding, 17.14; earthy phosphates, with a trace of fluate of lime, 1.00; lithic acid, 1.00; mucus of the bladder, 0.32; silex, 0.03. Dr. Proute found 100 parts to consist of lithic acid, 90.16; potassa, 3.45; ammonia, 1.70; sulphate of potassa, with a trace of chloride of sodium, .95; phosphate of lime, carbonate of lime, and magnesia, 80; and animal matter, consisting of mucus and a little colouring matter, 2.94.d M. Ras-

a Elements of Experimental Chemistry, vol. ii.

c Annals of Philos. v. 415. See, also, an analysis by Prof. Thomas Thomson, in

Records of General Science, ii. 3.

b Med. Chirurgical Transact. vol. iii.; and Annals of Philos. ii. 423; and The Kidneys and Urine, by J. J. Berzelius, translated from the German, by M. H. Boyé, and F. Leaming, M.D., p. 97, Philad. 1843.

<sup>4</sup> For the appearances presented by the urine under the microscope, see Quevenne, and Vigla, l'Expérience, Dec. 1837, Janvier, 1838, & Mars, 1838; Donné, ibid. Janvier,

paila thinks it "possible" that uric acid is merely a mixture of organic matter (albumen) with an acid cyanide of mercury; so that the results of analysis may differ according as the analyzed substances may have been more or less separated from the organic matter. The physical and chemical characters of uric acid, he thinks, accord very well with this hypothesis.

The yellowish-red incrustation, deposited on the sides of chamber utensils, is the uric or lithic acid. This is the basis of one of the

varieties of calculi.

The quantity of urine, passed in the twenty-four hours, is very variable. Boissier states it at 22 ounces; Hartmann at 28; Dr. Robt. Willisb at from 30 to 40; Prout at 32; Robinson at 35; Von Gorter at 36; Keill at 38: Rye at 39; Bostock at 40; Sanctorius at 44; Stark at 46; Dalton at 48½; Haller at 49; Christison at from 35 to 50; Becquerel at about 46; Dr. Thomas Thomson at 53; and Lining at from 56 to 59 ounces.c On the average, it may be estimated perhaps at two pounds and a half; hence the cause of the great size of the renal artery, which, according to the estimate of Haller, conveys to the kidney a sixth or eighth part of the whole blood. Its quantity and character vary according to age, and, to a certain extent, according to sex. We have already seen, under the head of cutaneous exhalation, how it varies, according to climate and season; and it is influenced by the serous, pulmonary, and cellular exhalations likewise: one of the almost invariable concomitants of dropsy is diminution of the renal secretion. Its character, too, is modified by the nature of the substances received into the blood. Rhubarb, turpentine, and asparagus materially alter its physical properties; whilst certain articles stimulate the kidney to augmented secretion, or are "diuretics."

The urine does not appear to be intended for any local function. Its use seems to be restricted to the removal of the elements of the substances, of which it is composed, from the blood; hence, it is solely depuratory and decomposing. How this decomposition it accomplished we know not. We have already referred to the experiments, performed by MM. Prévost and Dumas, Ségalas, Gmelin, Tiedemann and Mitscherlich, in which urea was found in the blood of animals whose kidneys had been extirpated: an inquiry has consequently arisen - how it exists there? Prior to these experiments, it was universally believed, that its formation is one of the mysterious functions executed in the intimate tissue of the kidney.d It is proper to add, however, that neither Pré-1838; and for some new researches on Human Urine, see Lecanu, Journal de Pharmacie, Nov. and Dec. 1839; Dr. G. Bird, op. cit.; Mandl, Manuel d'Anatomie, générale, p. 504, Paris, 1843; and Prout, on the Nature and Treatment of Stomach and Renal Diseases, 4th edit., p. 517, Lond. 1843. 2 Op. citat., p. 507.

d Annales de Chimie, xliii. 61, and Raspail's Chimie Organique, p. 506.

b Urinary Diseases and their Treatment, Bell's Library Edit., p. 14, Philad. 1839.
c Burdach, op. citat. v. 271. See, also, Séméiotique des Urines, ou Traité des Alterations de l'Urine dans les Maladies, &c., Paris, 1841; and Thomson, Chemistry of Animal Bodies, p. 459, Edinb. 1843.

vost and Dumas, Tiedemann and Gmelin nor M. Lecanua could detect the smallest trace of this substance in the blood of animals placed under ordinary circumstances. It is, according to Wöhler, a cyanate of ammonia, and contains a very large proportion of azote, so that, it has been imagined, the kidney may possibly be the outlet for an excess of azote, or for preventing its accumulation in the system — in the same manner as the lungs and liver have been regarded as outlets for superfluous carbon. The quantity of azote, discharged in the form of urea, is so great, even in those animals whose food does not essentially contain this element, that it has been conceived a necessary ingredient in the nutrition of parts, and especially in the formation of fibrin, which, we have seen, is a chief constituent of the blood, and of every muscular organ. The remarks, made on the absorption of azote during respiration, indicate how it is received into the system; and it has been presumed, that the superfluous portion is thrown off in the form of urea. The experiments of MM. Prévost and Dumas, and of the other physiologists, would certainly favour the conclusion, that urea may exist ready formed in the blood, and that the great function of the kidney may be to separate it along with the other constituents of the urine.

Adelone ascribes the source of the urea to the products of interstitial decomposition. He conceives, that, in this shape, they are received into the blood, and that the office of the kidneys is to separate them. All this is necessarily conjectural, and it must be admitted, that our knowledge of the subject is by no means ample, and that we must wait for farther developments. Certain it is, that the removal of the constituents of the urinary secretion from the blood is all-important. Experiments on animals have shown, that if it be suppressed by any cause for about three days, death usually supervenes, and the dangers to man are equally imminent. Yet there are some strange cases of protracted suppression on record. Haller mentions a case in which no urine had been secreted for twenty-two weeks, and Dr. Richardsond one of a lad of seventeen, who had never made any and yet felt no inconvenience.

# a. Connexion between the Stomach and the Kidneys.

In consequence of the rapidity with which fluids, received into the stomach, are sometimes voided by the urinary organs, it has been imagined, either that vessels exist, which communicate directly between the stomach and bladder, or that the fluid passes through the intermediate cellular tissue, or by means of the anastomoses of

a Etudes Chimiques sur le Sang Humain, Paris, 1837.

b Berzelius, The Kidneys and Urine, translated by M. H. Boyé and F. Leaming, M.D., p. 73, Philad. 1843; Graham's Elements of Chemistry, Amer. Edit. by Dr. Bridges, p. 671, Philad. 1843; and Thomson, Chemistry of Animal Bodies, p. 77, Edinb. 1843.

c Physiologie de l'Homme, tom. iii., Paris, 1829.

d Philos. Transact. for 1713; and Elliotson's Blumenbach's Physiology, 4th edit., Lond. 1828.

the lymphatics. In support of the opinion, that a more direct passage exists, the assertion of Chirac. - that he saw the urinary bladder become filled with urine, when the ureters were tied, and that he excited urinous vomiting, by tying the renal arteries, - is adduced. It has been farther affirmed, that the oil, composing a glyster, has been found in the bladder. Darwin," having administered to a friend a few grains of nitrate of potassa, collected his urine at the expiration of half an hour, and had him bled. The salt was detected in the urine, but not in the blood. Brande made similar experiments with the prussiate of potassa, from which he inferred that the circulation is not the only medium of communication between the stomach and the urinary organs, without, however, indicating the nature of the supposed medium; and this view is embraced by Sir Everard Home, Wollaston, Marcet, and others. Lippi, of Florence, thinks he has found an anatomical explanation of the fact. According to him, the chyliferous vessels have not only numerous inosculations with the mesenteric veins, either before their entrance into the mesenteric glands, or whilst they traverse the glands, but, when they attain the last of those glands, some of them proceed to open directly into the renal veins, and into the pelves of the kidneys. At this place, according to him. the chyliferous vessels divide into two sets; the one, ascending, and conveying the chyle into the thoracic duct; the other, descending, and carrying the drinks into the renal veins and pelves of the kidneys. He affirms, that the distinction between these two sets is so marked, that an injection, sent into the former, goes exclusively into the thoracic duct, whilst if it be thrown into the latter it passes exclusively to the kidneys. These direct vessels Lippi calls vasa chylopoietica urinifera. If the assertions of Lippi were anatomical facts, it would obviously be difficult to doubt some of the deductions: other anatomists have not, however, been so fortunate as he; and, consequently, it may be well to make a few comments. Some of these chylopoietica urinifera, he affirms, open into the renal veins. This arrangement, it is obvious, cannot be invoked to account for the shorter route, - the royal road to the kidney: the renal vessel is conveying the blood back from the kidney, and every thing that reaches it from the intestines. must necessarily pass into the vena cava, and ultimately attain the kidney through the renal artery. The vessels, therefore, that end in the renal veins, must be put entirely out of the question, so far as regards the topic of dispute; and our attention be concentrated upon those that terminate in the pelvis of the kidney. Were this termination proved, we should be compelled, as we have remarked, to bow to authority; but not having been so, it may be

Illustrazioni Fisiologiche e Patologice del Sistema Linfatico-Chilifero, &c. Firenz. 1825.

a Zoonomia, xxix. 3.

b Philosophical Transactions, xcviii. 51, and ci. 163, for 1808 and 1811; and Lectures on Comparative Anatomy, i. 221, Lond. 1814; and iii. 138, Lond. 1823.

stated as seemingly improbable, that the ducts in question should take the circuitous course to the pelvis of the kidney, instead of

proceeding directly to the bladder.

We know then, anatomically, nothing of any canal existing between the stomach and the bladder; and we have not the slightest evidence, - positive or relative, - in favour of the opinion, that there is any transmission of fluid through the intermediate cellular tissue. We have, indeed, absolute testimony against it. MM. Tiedemann and Gmelin, having examined the lymphatics and cellular tissue of the abdomen, in cases where they had administered indigo and essence of turpentine to animals, discovered no traces whatever of them, whilst they could be detected in the kidney. The facts, again, referred to by Chirac, are doubtful. If the renal arteries be tied, the secretion cannot be effected by the kidney; yet, as we have seen, in the case of extirpated kidneys, urea may exist in the blood, and, consequently, urinous vomitings be possible. If the ureters be tied, the secretion being practicable, death will occur if the suppression be protracted; and, in such case, the secreted fluid may pass in the vessels, and readily give a urinous character to the perspiration, vomited matters, &c., &c. The experiments of Darwin, Brande, Wollaston, and others only demonstrate, that these gentlemen were unable to detect in the blood that which they found in the urine. Against the negative results attained by these gentlemen, we may adduce the positive testimony of Fodéra, an experimentalist of weight, especially on those matters. He introduced into the bladder of a rabbit a plugged catheter, and tied the penis upon the instrument to prevent the urine from flowing along its sides. He then injected into the stomach a solution of the ferrocyanate of potassa. This being done, he frequently removed the plug of the catheter, and received the drops of urine on filtering paper: as soon as indications of the presence of the salt appeared in the urine by the appropriate tests, - which usually required from five to ten minutes after its reception into the stomach, - the animal was killed; and, on examining the blood, the salt was found in the serum taken from the thoracic portion of the vena cava inferior, in the right and left cavities of the heart, in the aorta, the thoracic duct, the mesenteric glands, the kidneys, the joints, and in the mucous membrane of the bron-Magendie, too, states, as the result of his experiments, -First. That whenever prussiate of potassa is injected into the veins, or is exposed to absorption in the intestinal canal, or in a serous cavity, it speedily passes into the bladder, where it can be readily recognised in the urine. Secondly. That if the quantity of prussiate injected be considerable, it can be detected in the blood by reagents; but if the quantity be small, it is impossible to dis-Thirdly. That the same thing cover it by the ordinary means.

b Précis, &c. ii 477.

a Recherches Expérimentales sur l'Absorption et l'Exhalation, Paris, 1824.

happens if the prussiate of potassa be mixed with the blood out of the body. Fourthly. That the salt can be detected, in every

proportion, in the urine.

We may conclude, therefore, with Dr. Hale, who has written an interesting paper on this subject, that the existence of any more direct route from the stomach to the bladder than the circulatory system and the kidneys is disproved; and the absorption of fluids must be considered to be effected through the vessels described under the Absorption of Drinks.

Such are the glandular secretions, which we shall consider in this place. There are still two important fluids, the *sperm* and the *milk*, whose uses will have to be detailed under the next class of functions.

#### GLANDIFORM GANGLIONS.

There are several organs, — as the spleen, thyroid, thymus, and supra-renal capsules, — which are termed glands by many anatomists; but which Chaussier has termed glandiform ganglions. Of the uses of these we know little or nothing. Yet it is necessary, that the nature of the organs, and their fancied functions should meet with notice. The offices of the thyroid, thymus, and supra-renal capsules, — being chiefly confined to fætal existence, — will not require consideration here.

## a. The Spleen.

The spleen is a viscus of considerable size, situate in the left hypochondriac region, (Fig. 193, H,) beneath the diaphragm, above the left kidney, and to the left of the stomach. Its medium length is about four and a half inches; its thickness two and a half inches; and its weight about eight ounces. It is of a soft texture, somewhat spongy to the feel, and easily torn. In a very recent subject, it is of a greyish-blue colour; which, in a few hours, changes to a purple, so that it resembles a mass of clotted blood. At its inner surface, or that which faces the stomach and kidney, a fissure exists, by which the vessels, nerves, &c., enter or issue from the organ.

The anatomical elements of the spleen are:—1. The splenic artery, which arises from the cœliac, and after having given off branches to the pancreas and the left gastro-epiploic artery, divides into several branches, which enter the spleen at the fissure, and ramify in the tissue of the organ, so that it seems to be exclusively formed by them. Whilst the branches of the artery are still in the duplicature of the gastro-splenic omentum, and before they ramify in the spleen, they furnish the vasa brevia to the stomach. The precise mode of termination of the arteries

Boylston Prize Dissertations for the years 1819 and 1821, Boston, 1821.
 Dr. Gross, Elements of Pathological Anatomy, ii. 344, Boston, 1839.

in the spleen is unknown. The communication of the arteries with the veins does not, however, appear to be as free as in other parts of the body, or the anastomoses between the minute arteries as numerous. If, according to Assolant, one of the branches of the splenic artery be tied, the portion of the spleen to which it is distributed dies; and if air be injected into one of these branches, it does not pass into the other; so that the spleen would appear to be a congeries of several distinct lobes; and in certain animals the lobes are so separated as to constitute several spleens. A similar appearance is occasionally seen in the human subject. 2. The splenic vein arises by numerous radicles in the tissue of the spleen: these become gradually larger, and less numerous, and leave the fissure of the spleen by three or four trunks, which ultimately, with veins from the stomach and pancreas, unite to form one, that opens into the vena portæ. It is without valves, and its parietes are thin. These are the chief constituents. 3. Lymphatic vessels, which are large and numerous. 4. Nerves, proceeding from the cœliac plexus: they creep along the coats of the splenic artery, - upon which they form an intricate plexus, - into the substance of the spleen. 5. Cellular tissue, which serves as a bond of union between these various parts; but is in extremely small quantity. 6. A proper membrane, which envelopes the organ externally; adheres closely to it, and furnishes fibrous sheaths to the ramifications of the artery and vein; keeping the ramifications separated from the tissue of the organ, and sending prolongations into the parenchyma, which gives it more of a reticulated than spongy aspect. 7. Of blood, according to many anatomists; but blood differing from that of both the splenic artery and vein; containing, according to Vauquelin, less colouring matter and fibrin, and more albumen and gelatin, than any other kind of blood. This, by stagnating in the organ, is conceived to form an integrant part of it. Malpighib believed it to be contained in cells; but others have supposed it to be situate in a capillary system intermediate to the splenic artery and vein.c

Assolant and Meckel<sup>d</sup> believe, that the blood is in a peculiar state of combination and of intimate union with the other organic elements of the viscus, and with a large quantity of albumen; and that this combination of the blood forms the dark brown pulpy substance, contained in the cells formed by the proper coat, and which can be easily demonstrated by tearing or cutting the spleen, and scraping it with the handle of a knife. These cells and the character of the tissue of the spleen are exhibited in the marginal


<sup>\*</sup> Recherches sur la Rate, Paris, 1801.

b Op. Omnia, part ii. Lond. 1687; and Op. Posthum. p. 42, Lond. 1697.

c Seiler, in art. Milz, in Pierer's Anat. Physiol. Wörterbuch, B. v. s. 322, Altenburg, 1832.

d Handbuch, &c., traduit par Jourdan, iii. 476, Paris, 1825.

VOL. II. - 26



Section of the Spleen.

figure. (Fig. 208.) In addition to the pulp, there is an abundance of rounded corpuscles, varying in size from an almost imperceptible magnitude to a line or more in diameter. By Malpighi, these were conceived to be granular corpuscles, and, by Ruysch, a simply convoluted vessels. Andralb affirms, that by repeated washings, the spleen is shown to consist of an infinite number of cells, which communicate on the one hand together, and, on the other, directly with the splenic veins. The latter, when the inner surface of the large subdivisions of the splenic veins are examined, appear to have a great number of perforations, through which a probe passes directly into the cells of the organ. The farther the subdivisions of the vein examined are from the trunk, the larger are these perfora-

tions, and still farther on, the coats of the vein are not a continued surface, but are split into filaments, which do not differ from those forming the cells, and are continuous with them.c Recently, M. Bourgery has maintained that the fibrous envelope of the spleen sends off a multitude of lamellæ, which penetrate into its interior, forming irregular spaces of unequal dimensions. These short spaces he calls splenic vesicles. In the septa, a number of lymphatic glands exists. The capillaries of the arteries communicate directly with those of the veins; but, according to M. Bourgery, there are, in addition, veins with patulous orifices. The interior of the vesicles is filled with a soft substance of a deep red colour, in which the small white corpuscles, discovered by Malpighi, are suspended. M. Mandld suggests, that the white corpuscles may be analogous to the intestinal villi, in which the lymphatics originate by a cæcal extremity. Besides the proper membrane, the spleen also receives a peritoneal coat; and, between the stomach and the organ, the peritoneum forms the gastro-splenic

2 Meckel, op. citat.

Précis, d'Anatomie Pathologique, tom. ii. part i. p. 416, Paris, 1832.

d Manuel d'Anatomie générale, p. 518, Paris, 1843.

<sup>•</sup> See, also, Sir E. Home's Lectures on Comparative Anatomy, iii. 148; Dr. Warner on the Distribution of the Splenic Vein, in the spleen of the ox and sheep, in American Journal of the Med. Sciences, Feb. 1836, p. 541; and Weber's Hildebrandt's Handbuch der Anatomie, u. s. w., Band iv.s. 328, Braunschweig, 1822: and on the Minute Structure of the Spleen, Bourgery, Gaz. Méd., Juin 11, 1842; and Brit. and For. Med. Rev. Oct. 1842, p. 541.

SPLEEN. 303

epiploon or gastro-splenic ligament, in the duplicature of which are situate the vasa brevia.

Lastly: the spleen is capable of distension and contraction; and is possessed of little sensibility in the healthy state. It has no

excretory duct.

The hypotheses, which have been indulged on the nature of the spleen, are beyond measure numerous and visionary; and, after all, we are in the greatest obscurity as to its real uses. Many of these hypotheses are too idle to merit notice; such are those, that consider it to be the seat of the soul, - the organ of dreaming, - of melancholy and of laughter, - of sleep and the venereal appetite,—the organ that secretes the mucilaginous fluids of the joints, that serves as a warm fomentation to the stomach, and so on. It was long regarded as a secretory apparatus, for the formation of the atrabilis, — of a fluid intended to nourish the nerves, - of the gastric juice, - of a humour intended to temper the alkaline character of the chyle or bile, &c. The absence of an excretory duct would be a sufficient answer to all these speculations, if the non-existence of the supposititious humours were insufficient to exhibit their absurdity. MM. Tiedemann and Gmelina consider its functions to be identical with those of the mesenteric glands. They regard it as a ganglion of the absorbent system, which prepares a fluid to be mixed with the chyle and effect its animalization. In favour of the view, that it is a part of the lymphatic system, they remark, that it exists only in those animals that have a distinct absorbent system; that its bulk is in a ratio with the development of the absorbent system; that the lymphatics predominate in the structure of the organ; that its texture is like that of the lymphatic ganglions; and lastly, that, on dissecting a turtle, they distinctly saw all the lymphatics of the abdomen passing first to the spleen, then leaving that organ of larger size, and proceeding to the thoracic duct. In support of their second position, that it furnishes some material towards the animalization of the chyle, they adduce; - the large size of the splenic artery, which manifestly, they conceive, carries more blood to the spleen than is needed for its nutrition; and they affirm, that, in their experiments, they have frequently found, whilst digestion and chylosis were going on, the lymphatic vessels of the spleen gorged with a reddish fluid, which was carried by them into the thoracic duct, where the chyle always has the most rosy hue; and they add, that a substance injected into the splenic artery, passes readily into the lymphatics of the spleen. Lastly, after extirpating the spleen in animals, the chyle appeared to them to be more transparent; no longer depositing coagula; and the lymphatic ganglions of the abdomen seemed to have augmented in size. Views, similar to these, have been maintained by Sir Everard Home.b

b Philosoph. Transactions for 1808 and 1811; and Lect. on Comp. Anatomy, loc. cit.

Versuche über die Wege auf welchen Substanzen aus dem Magen und Darmkanal ins Blut gelangen, p. 86, Heidelb. 1820.

Chaussier, as we have seen, classes the spleen amongst the glandiform ganglions, and affirms, that a fluid is exhaled into its interior, of a serous or sanguineous character, which, when ab-

sorbed, assists in lymphosis.

Many, again, have believed, that the spleen is a sanguineous, not a lymphatic ganglion, but they have differed regarding the blood on which it exerts its action; some maintaining, that it prepares the blood for the secretion of the gastric juice; others, for that of the bile. The former of these views is at once repelled by the fact, that the vessels which pass from the splenic artery to the stomach, leave that vessel before it enters the spleen. The latter has been urged, of late, by M. Voisin.<sup>a</sup> He thinks, the principal use of the spleen is to furnish to the liver, blood containing those materials that enter into the composition of the bile; but this view, also, rests on very uncertain data and deductions.

Since the period of Haller, the blood of the splenic vein has been presumed to differ essentially from that of other veins, which has led to the belief, that some elaboration is effected in the spleen so as to fit the blood for the secretion of the bile. It has been described as more aqueous, more albuminous, more unctuous, and blacker than other venous blood; to be less coagulable, less rich in fibrin, and the fibrin it does contain to be less animalized. Yet these affirmations have been denied; and even were they admitted, we have no positive knowledge, that such changes better adapt

it for the formation of bile by the liver.

The ideas that have existed, regarding its acting as a diverticulum for the blood, have been mentioned under the head of Circulation. By some, it has been supposed to act as such in the intervals of digestion; or in other words, to be a diverticulum to the stomach: by others, its agency in this way is believed to apply to the whole circulatory system, so that when the flow of blood is impeded or arrested in other parts, it may be received into the spleen. Such a view was entertained by Dr. Rush. It has been embraced by many others.

It is hard to say which of these speculations is the most ingenious. None can satisfy the judicious physiologist, especially when he considers the comparative impunity consequent on extirpation of the organ. This was an operation performed at an early period. Pliny affirms that it was practised on runners to render them more swift. From animals the spleen has been repeatedly removed, and although many of these have died in con-

c Coxe's Medical Museum, Philad. 1807. See, also, on this subject, art. Milz. in Pierer, op. cit. s. 328; and Mr. Hake, Proceedings of the Royal Society, No. 39,

Inne 20, 1839.

Nouvel Aperçu sur la Physiologie du Foie, et les Usages de la Bile, Paris, 1833.
 Dr. W. Stukely, Of the Spleen, its Description and History, Uses and Diseases,
 &c., Lond. 1722; and Dr. G. Dermott, Medical Times, April 3, 1841.

<sup>&</sup>lt;sup>4</sup> Dr. O'Beirne, Dublin Journal of Medical Science, Nov. 1842, p. 219; Carpenter, Human Physiology, § 709, Lond. 1842; and Dr. Hargreave, Dublin Medical Press, Aug. 10, 1842, p. 81.

SPLEEN. 305

sequence of the operation, several have recovered. Adelonb refers to the case of a man who was wounded by a knife under the last false rib of the left side. Surgical attendance was not had until twelve hours afterwards; and as the spleen had issued at the wound, and was much altered, it was considered necessary to extirpate it. The vessels were tied; the man got well in less than two months, and has ever since enjoyed good health. Sir Charles Belle asserts, that an old pupil had given him an account of his having cut off the spleen in a native of South America. spleen had escaped through a wound, and had become gangrenous. He could observe no effect to result from the extirpation. T. Chapman, Esq. of Purneah, in India, has related a case of excision of a portion of the spleen by Dr. Macdonald of that station. A native, about thirty years of age, was gored in the abdomen by a buffalo, and through the wound, which was about three inches in length, a portion of the spleen protruded. Six days afterwards, the man sought advice from Dr. Macdonald, who removed the spleen with the knife, and the patient rapidly recovered.

Dr. O'Brien, in an inaugural dissertation, published at Edinburgh, in 1818, refers to a case, which fell under his own management. The man was a native of Mexico: the spleen lay out, owing to a wound of the abdomen, for two days before the surgeon was applied to. The bleeding was profuse: the vessels and other connexions were secured by ligature, and the spleen separated completely on the twentieth day of the wound. On the forty-fifth day, the man was discharged from the hospital, cured; and he remarked to some one about this time, that "he felt as well as

ever he did in his life."e

Dulaurens, Kerckring, Baillie, and others, refer, also, to cases, in which the spleen has been found wanting in man, without any

apparent impediment to the functions.

The experiments, which have been made on animals, by removing the spleen, have led to discordant results. Malpighi says, that the operation was followed by increased secretion of urine; Dumas, that the animals had afterwards a voracious appetite; Mead, and Mayer, that digestion was impaired, that the evacuations were more liquid, and the bile more watery; Tiedemann and Gmelin, that the chyle appeared more transparent and devoid of clot; Professor Coleman, that the dogs,—the subjects of the experiment,—were fat and indolent. A dog, whose spleen was removed by Mr. Mayo, became, on recovering from the wound,

<sup>2</sup> J. H. Schulze, de Splene Canibus Exciso, Hal. 1735: Morgagni, Animad. Anat. iii. Animad. xxv. Lugd. Bat. 1741.

b Physiol. de l'Homme, 2de édit. tom. iii. Paris, 1829.

Anat. and Physiol., 5th Amer. Edit., by Dr. Godman, ii. 363, New York, 1827.
 India Journal of Medicine, vol. viii. p. 1; and London Medical Gazette, for May, 20th, 1837, p. 285.
 Blundell's Researches, Physiol. and Pathol. Lond. 1825.
 Markid Anatomy 2d edit p. 260. Lond. 1727; Medical's Handbuch der Pathologie.

f Morbid Anatomy, 2d edit. p. 260, Lond. 1797; Meckel's Handbuch der Pathologischen Anatomie, B. i. s. 608, Leipz. 1812.

s Outlines of Human Physiology, 4th edit. p. 107, Lond. 1838; and Outlines of Human Pathology, p. 128, Lond. 1836.

26\*

fatter than before: in a year's time it had returned to its former condition, and no difference was observed in its appearance or habits from those of other dogs. Similar results followed the experiments of Dr. Blundell, Mr. Dobson, and Mr. Eagle; and the last gentleman states, that an offer had been made him of a "smart sum of money" by a dealer in Leadenhall Market, if he would tell

him his method of fattening animals.

Dupuytren extirpated the spleen of forty dogs on the same day, without tying any vessel, but merely stitching up the wound of the abdomen, - vet no hemorrhage supervened. In the first eight days, half the dogs operated on, died of inflammation of the abdominal viscera induced by the operation, as was proved by dissection. The other twenty got well without any accident, at the end of three weeks at the farthest. At first they manifested a voracious appetite, but it soon resumed its natural standard. They fed on the same aliment, the same drinks, took the same quantity of food, and digestion seemed to be accomplished in the same time. The fæces had the same consistence, the same appearance, and the chyle appeared to have the same character. Nor did the other functions offer any modification. Dupuytren opened several of these dogs some time afterwards, and found no apparent change in the abdominal circulation, - in that of the stomach, epiploon, or liver. The last organ, which appeared to some of the experimenters to be enlarged, did not seem to him to be at all so. The bile alone appeared a little thicker, and deposited a slight sediment.b

These circumstances render it extremely difficult to arrive at any theory regarding the offices of this anomalous organ. It is manifestly not essential to life, and therefore not probably inservient to the purposes assigned it by Tiedemann and Gmelin. Its office must evidently be of a supplementary or vicarious nature; and this would accord best, perhaps, with the notion of its serving as a diverticulum; the blood speedily passing, after the organ has been extirpated, into other channels; c a view, which, as elsewhere remarked,d is somewhat confirmed by the splenic enlargements consequent on repeated attacks of intermittent, - the blood, which has receded from the surface, accumulating perhaps in this organ. It must be admitted, however, that our knowledge of the function is of a singularly negative and unsatisfactory character, and this is strikingly exemplified by the suggestion of Paleye - who was certainly not predisposed to arrive at such a conclusion - that the spleen "may be merely a stuffing, a soft cushion to fill up a vacuum or hollow, which, unless occupied, would leave the pack-

age loose and unsteady."

a Lond. Lancet, Oct. 8, 1842, p. 58, and Dec. 10, 1842, p. 406.

Natural Theology, c. 11.

Bostock, Physiology, 3d edit. p. 580, Lond. 1836. See, also, Hodgkin, in Appendix to translation of Edwards, De l'Influence des Agens Physiques, &c. Lond. 1833. Practice of Medicine, vol. i., 2d edit., Philad. 1844.

## BOOK III.

### REPRODUCTIVE FUNCTIONS.

The functions, which we have been hitherto considering, relate exclusively to the individual. We have now to investigate those that refer to the preservation of the species, and without which living beings would soon cease to exist. Although these functions are really multiple, it has been the custom with physiologists to refer them to one head — generation — of which they are made to form the subordinate divisions.

## CHAPTER I.

#### GENERATION.

The function of generation, much as it varies amongst organized bodies, is possessed by them exclusively. When a mineral gives rise to another of a similar character, it is at the expense of its own existence; but the animal and the vegetable produce being

after being, without any curtailment of theirs.

The writers of antiquity considered, that all organized bodies are produced in one of two ways. Amongst the upper classes of both animals and vegetables, they believed the work of reproduction to be effected by a process, which is termed univocal or regular generation, (generatio homogenea, propagatio;) but in the very lowest classes, as the mushroom, the worm, the frog, &c., they conceived that the putrefaction of different bodies, aided by the influence of the sun, might generate life. This has been termed equivocal or spontaneous generation, (generatio heterogenea, æquivoca, primitiva, primigena, originaria, spontanea;) and is supposed to have been devised by the Egyptians to account for the swarms of frogs and flies, which appeared on the banks of the Nile after its periodical inundations.a Amongst the ancients, the latter hypothesis was almost universally credited. Pliny unhesitatingly expresses his belief, that the rat and the frog are produced in this manner; and at his time it was generally thought, that the bee, for example, was derived, at times from a parent, but at others from putrid beef.b

The passage of Virgil<sup>c</sup> — in which he describes how the shepherd Aristæus succeeded in producing swarms of bees from the entrails of a steer, exposed for nine days to putrefaction — is probably fa-

miliar to most readers, and exhibits the same belief.

\* Fleming's Philosophy of Zoology, i. 24, Edinb. 1822; Burdach's Physiologie als Erfahrungswissenschaft, 2te Auflage, i. 8, Leipz. 1835; and Purkinje, in art. Erzeugung, Encyclop. Wörterbuch, xi. 512, Berlin, 1834.

b " Apes nascuntur partim ex apibus, partim ex bubulo corpore putrefacto." - Varro.

De Re Rustica, iii. 16. See, also, Plinii Hist. Natural.

c Georgic. lib. iv. l. 295.

The hypothesis of equivocal generation having been conceived, in consequence of the impracticability of tracing ocularly the function in the minute tribes of animals, it naturally maintained its ground uninterruptedly, as regarded those animals, until better means of observation were invented. The difficulty, too, of admitting regular generation as applicable to all animals, was augmented by the fact, not at first known to naturalists, that many of the lower tribes conceal their eggs, in order that their nascent larvæ may find suitable food; but the existence of evident sexual organs in many of these small species induced physiologists, at an early period, to believe, that they also might be reproduced by sexual intercourse: direct proofs were not, however, obtained until the discovery of the microscope; after which the investigations of Redi, Vallisnieri, Swammerdam, Hooke, Réaumur, Bonnet and others, clearly demonstrated that many of the smallest insects have eggs and sexes, and that they reproduce like other animals. In the case of plants it has been supposed that the growth of the fungi amongst dung, and of the various parasitical plants that appear on putrid flesh, fruit, &c. furnishes facts in support of the equivocal theory; but the microscope exhibits the seeds of many of these plants, and experiments show that they are prolific. The characters, by which the different species and varieties are distinguished, although astonishingly minute, are fixed, exhibiting no fluctuation, such as might be anticipated did these plants arise by spontaneous generation, or by the fortuitous concourse of atoms.

The animalcules, that make their appearance in water in which vegetable or animal substances have been infused or are contained, would seem, at first sight, to favour the ancient doctrine. In these cases, however, the species, again, have determinate characters; presenting always the same proportion of parts; and appearing to transmit their vitality to their descendants in a manner not unlike animals and vegetables higher in the scale. The explanation, offered by the supporters of the univocal theory for those obscure cases in which direct observation fails us, is that their seeds and eggs are so extremely minute, that they can be borne about by the winds, or by birds; be readily deposited, and, when they find a soil or nidus, favourable to their growth, can undergo develop-Thus, the soil, in which alone the monilia glauca flourishes, is putrid fruit; whilst the small infusory animal the vibrio aceti or vinegar eel - requires, for its growth, vinegar that has been for some time exposed to the air. "That the atmosphere," says Dr. Good, is freighted with myriads of insect eggs, that elude our senses; and that such eggs, when they meet with a proper bed, are hatched in a few hours into a perfect form, is clear to any one who has attended to the rapid and wonderful effects of what, in common language, is called a blight, upon plantations and gardens. I have seen, as probably many, who may read this work, have also, a hop-ground completely overrun and desolated b Study of Medicine, Cl. i. Ord. 1. Gen. x. Sp. 3. \* Fleming, op. citat. p. 24.

by the aphis humuli or hopgreen-louse, within twelve hours after a honey-dew (which is a peculiar haze or mist, loaded with a poisonous miasm) has slowly swept through the plantation, and stimulated the leaves of the hop to the morbid secretion of a saccharine and viscid juice, which, while it destroys the young shoots by exhaustion, renders them a favourite resort for this insect, and a cherishing nidus for the myriads of little dots that are its eggs. The latter are hatched within eight-and-forty hours after their deposit, and succeeded by hosts of other eggs of the same kind; or, if the blight takes place in an early part of the autumn, by hosts of the young insects produced viviparously; for in different seasons of the year, the aphis breeds both ways. Now it is highly probable, that there are minute eggs or ovula of innumerable kinds of animalcules floating in myriads of myriads through the atmosphere, so diminutive as to bear no larger proportion to the eggs of the aphis than these bear to those of the wren, or the hedge-sparrow; protected, at the same time, from destruction by the filmy integument that surrounds them, till they can meet with a proper nest for their reception, and a proper stimulating power to quicken them into life; and which, with respect to many of them, are only found obvious to the senses in different descriptions of animal fluids. "The same fact occurs in the mineral kingdom: stagnant water, though purified by distillation and confined in a marble basin, will, in a short time, become loaded on its surface or about its sides with various species of confervas; while the interior will be peopled with microscopic animalcules. So, while damp cellars are covered with boletuses, agarics and other funguses, the driest brick walls are often lined with the lichens and mosses. We see nothing of the animal and vegetable eggs or seeds by which all this is effected; but we know, that they exist in the atmosphere, and that this is the medium of their circulation."

This view of the extraneous origin of the seeds of the confervas, &c., is corroborated by an experiment of Senebier. He filled a bottle with distilled water, and corked it accurately; not an atom of green matter was produced, although it was exposed to the light of the sun for four years; nor did the green matter, considered as the first stage of spontaneous organization, exhibit itself in a glass of common water, covered with a stratum of oil." It is proper, however, to remark, that the observation of others invalidates the results of this experiment. Burdach, b assisted by Hensche, and along with Professor von Baer, poured water on marble in a glass vessel, the remainder of the vessel being filled with atmospheric air, oxygen or hydrogen, and placed it in the light of the sun, or in warm sand. No green matter was perceptible, but there was a slimy substance with white threads, part of which had a ramified appearance, and part that of coral. On the other hand, pieces of granite, newly broken from the midst of a block,

a See, also, Wrisberg, Observ. de Animal, Infusor.

b Die Physiologie als Erfahrungswissenschaft, 2te Auflage, i. 23, Leipz. 1835.

produced — with fresh distilled water, and oxygen or hydrogen, in the sun — green matter, with threads of the confervas; but in the warmth of digestion flocculi only. He next took some mould, which he dug up, and which was inodorous, and apparently free from all foreign matter; boiled it in a considerable quantity of water, and reduced the decoction to the consistence of a thick, partly pulverulent extract. This gave, with common water and atmospheric air — in bottles with ground stoppers, tied over with bladder — in the sun, numerous infusory animalcules and green matter; but with distilled water and oxygen or hydrogen, green

matter only appeared at the bottom of the bottles.

The subject of intestinal worms has been eagerly embraced by the supporters of the doctrne of equivocal generation, who are of opinion, that the germs need not be received from without; whilst the followers of the univocal doctrine maintain, that they must always be admitted into the system. The first opinion includes amongst its supporters the names of Needham, Buffon, Patrin, Treviranus, b Rudolphi, Bremser, d Himly, and other distinguished helminthologists. The latter comprises those who believe in the Harveian maxim, - omne vivum ex ovo.º To support the latter opinion, it has been attempted to show, that the worms, found in the human intestines, are precisely the same as others that have been found out of the body; but the evidence, in favour of this position, is by no means strong or satisfactory. Linnæus affirms, that the twnia vulgaris, — of a smaller size, however, — has been met with in muddy springs; and the ascarides vermiculares in marshes, and in the putrescent roots of plants. Gadd affirms, that he met with the tænia articulata plana osculis lateralibus geminis in a chalybeate rivulet; Unzer, tænia in a well; and Tissot says, that he found a tænia, exactly like the human, in a river; whilst Linnæus, Leeuenhoek, Schäffer and others affirm, that they have found the distoma hepaticum in water; but O. F. Müller, who took extraordinary pains in the comparative examination of the entozoa, which infest the human body, with those that are met with in springs, — states, that he has frequently detected the planariæ, but never saw one like the distoma hepaticum.

On the other hand, the supporters of the equivocal theory have laboured, with a good deal of success, to show, that a difference is always discoverable between the worms found without, and those found within, the body; but were it demonstrated to a mathematical certainty, that such difference exists, it would not be an invincible argument against the correctness of the univocal theory; as difference of locality, food, &c., might induce important changes in their corporeal development, and give occasion to the diversity.

An account of some New Microscopical Discoveries, 8vo., Lond. 1745.

b Biologie, tom. ii. 264.

· Entozoorum sive Vermium Intestinalium Historia Naturalis, tom. i. p. 370.

d Ueber Lebende Würmer im Lebenden Menschen, Wien, 1819.

e See the author's Commentaries on the Diseases of the Stomach and Bowels of Children, p. 20, Lond. 1824.

Rudolphi, op. citat.

which is occasionally perceptible amongst these parasites. Yet, if we admit that the germs of the entozoa are always received from without, their occurrence in different stages of development, in the fœtus in utero, is a circumstance difficult of explanation. Small, indeed, must be the germ, which, when received into the digestive organs of the mother, can pass into her circulation, be transmitted into the vessels of the fœtus, be deposited in some viscus, and there undergo its full development; yet such ceses have occurred, if the theory be correct. Certain it is, - however the fact may be accounted for, - that worms have been found in the fætus, by individuals whose testimony cannot be doubted. Eschholz saw them in the egg of the hen. Fromann found the distoma hepaticum in the liver of the fætal lamb; Kerckring, ascarides lumbricoides in the stomach of a fœtus six and a half months old; Brendel, tæniæ in the human fætus in utero; Heim, tæniæ in the newborn infant; Blumenbach, tæniæ in the intestine of the new-born puppy; and Göze, Bloch, and Rudolphi, the same parasite in

sucking lambs.b

Perhaps the conclusion of Cuviere is the soundest and most consistent with analogy, - that these parasites "propagate by germs so minute as to be capable of transmission through the narrowest passages; so that the germs may exist in the infant at birth." We have seen, that not simply the germs, but the animals themselves have been found at this early period of existence. The scientific world was, at one time, astounded by the assertion of Mr. Crosse, that he had succeeded in forming infusory animalcules from solutions of granite, silex, &c. by the aid of galvanism; but the only mode of explaining the phenomenon is, that organized matters, perhaps ova, were existent in the solutions, which became developed under the galvanic influence. The whole matter is involved in insuperable difficulties; yet the univocal theory is, in all respects perhaps, most admissible as regards the whole of the living creation: d still there are many distinguished naturalists who conceive it probable, that spontaneous generation occurs in the lowest divisions of the living scale. Amongst these may be mentioned De Lamarck, Raspail, Burdach, Treviranus, Wrisberg, Schweigger, Gruithuisen, Von Baer, and Adelon seems to accord with them. It has been properly remarked by a recent writer on this subject, who is himself inclined to admit the existence of spontaneous generation among some species of cryptogamic plants, infusorial animalcules, and entozoa, that "it must be held in recollection,

\* Spicilegium Anatom. Obs. lxxix. p. 154, Amstel. 1670.

Müller's Physiology, Baly's translation, Part i. p. 14, Lond. 1838.

Part xiii. p. 432, Feb. 1838.

b Stannius, art. Entozoa, in Encyclop. Wörterb. der Medic. Wissensch. xi. 276, Berlin, 1834; also, Grätzer, Die Krankheiten des Fætus, p. 107, Breslau, 1837.

c Règne Animal, p. 27. See, also, Ehrenberg, in Nov. Act. Nat. Cur. tom. x.; and

Fletcher's Rudiments of Physiology, Part ii. a, p. 13, Edinb. 1836; and Hodgkin's
 Lect. on the Morbid Anat. of the Serous and Mucous Membranes, p. 218, Lond. 1836.
 Dr. Allen Thomson, in art. Circulation, Cyclopædia of Anatomy and Physiology,

that many of these productions, after their first origin, propagate their species as parents, - that the so-called spontaneous kind of generation is to be looked upon as no more than an exception to the general law of reproduction, - and that, therefore, extreme caution is necessary in admitting any organized body to be the product of spontaneous generation upon the mere negative evidence of the absence of its seeds or ova." The views of De Lamarck .regarding the formation of living bodies, are strange in the ex treme; and exhibit to us, what we so frequently witness, that, in order to get rid of a subject, which is difficult of comprehension. the philosopher will frequently adopt suppositions, that require a much greater stretch of the imagination to invent, and present stronger obstacles to belief than those for which they have been substituted. M. De Lamarck maintains, that the first organized beings were formed throughout by a true spontaneous generation; their existence being owing to an excitative cause of life, probably furnished by the circumambient medium, and consisting of light and the electric fluid. When this cause meets with a substance of a gelatinous consistence, dense enough to retain fluids, it organizes the substance into cellular tissue, and a living being results. This process, according to De Lamarck, is occurring daily at the extremity of the vegetable and animal kingdoms. The being, thus formed, manifested originally, according to him, three faculties of life; - nutrition, growth, and reproduction, - but only in the most simple manuer. The organization soon, however, became more complicated, for it is, he remarks, a property of the vital movement to tend always to a greater degree of development of organization; to create particular organs, and to divide and multiply the different centres of activity; and, as reproduction has constantly preserved all that had been acquired, numerous and diversified species have been, in this manner, formed, possessing more and more extensive faculties. So that, according to this system, nature was directly concerned only in the first draughts of life; participating indirectly in the existence of living bodies of a more complex character; and these last proceeded from the former, after the lapse of an enormous time, and an infinity of changes in the incessantly increasing complication of organization; - reproduction continuing to preserve all the acquired modifications, and improvements.

The simplest kind of generation does not require sexual organs. The animal, at a certain period of existence, separates into several fragments, which form so many new individuals. This is called fissiparous generation or generation by spontaneous division. We have examples of it in the infusory animal cules, — as the vinegar eel or vibrio aceti. A somewhat more elevated kind of reproduction is the gemmiparous, — common in the vegetable kingdom, — which consists in the formation of buds, sporules, or germs on some part of the body. These, at a particular period.

\* Philosophie Zoologique, vol. i., Paris, 1830.

drop off, and form as many new individuals, and according as the germs are developed at the surface of the body, or internally, the gemmiparous generation is said to be *external*, or *internal*. In these two varieties, the whole function is executed by a single individual.

Higher up in the scale, we find special organs for the accomplishment of generation — male and female. In those animals, however, that possess special reproductive organs, some have both sexes in the same individual, or are hermaphrodite or androgynous, as is the case with almost all plants, and with some of the lower tribes of animals. In these, again, we notice a difference. Some are capable of reproduction without the concourse of a second individual; others, although possessing both attributes, require the concourse of another; the male parts of the one uniting with the female parts of the other. Both, in this way, become impregnated. The helix hortensis or garden snail affords us an instance of this kind of reproduction. They meet in pairs, according to Shaw, and, stationing themselves an inch or two apart, launch several small darts, not quite half an inch long, at each other. These are of a horny substance, and sharply pointed at one end. The animals, during the breeding season, are provided with a little reservoir for them, situate within the neck, and opening on the right side. On the discharge of the first dart, the wounded snail immediately retaliates on its aggressor, by throwing a similar dart; the other renews the battle, and, in turn, is wounded. When the darts are expended, the war of love is completed, and its consummation succeeds.

In the superior animals, each sexual characteristic is possessed by a separate individual, — the species being composed of two individuals, male and female, and the concourse of these individuals, or of matters proceeding from them, being absolutely necessary for reproduction. But here, again, two great differences are met with in the process. Sometimes the fecundating fluid of the male sex is not applied to the ovum of the female, until after its ejection by the latter, as in fishes. In other cases, the ovum cannot be fecundated after its ejection, and the fluid of the male sex is applied to it whilst still within the female, as in birds and the mammalia. In such case, the male individual is furnished with an organ for penetrating the parts of the female, and, in this kind of generation, there must be copulation. Again, where there is copulation, the following varieties may exist. First. The ovum, when once fecundated, may be immediately laid by the female, and may be hatched out of the body, constituting oviparous generation. Secondly. Although the process of laying may commence immediately, the fecundated ovum may pass so slowly through the excretory passages, that it may be hatched there, and the new individual may issue from the womb of the parent possessing the

<sup>&</sup>lt;sup>2</sup> Zoology, or Systematic Natural History, Lond. 1800; Dr. Good, in Physiol. Proem to class v. Genetica; and Purkinje, op. citat. xii. 521.

VOL. II. - 27

proper formation. This constitutes ovo-viviparous generation, of which we have examples in the viper and salamander. Thirdly. The fecundated ovum may be detached from the ovary soon after copulation, but, in place of being ejected, it may be deposited in a reservoir, termed a womb or uterus; be fixed there, attract fluids from the organ adapted for its development, and thus, increasing at the expense of the mother, be hatched, as it were, in this reservoir, so that the new individual may be born under its appropriate form. In such case, moreover, the new being, after birth, may be for a time supported on a secretion of the mother — the milk. These circumstances constitute viviparous generation: in which there are copulation, fecundation, gestation or pregnancy, and lactation or suckling. Lastly. There are animals, which, like the kangaroo, opossum, and wombat, are provided with abdominal pouches, into which the young, born at a very early stage of development, are received, and nourished with milk secreted from glands, contained within these pouches. Such animals are termed marsupial or marsupiate.a

There is considerable difference in animals as regards the nurturing care afforded by the parents to the young. Amongst the oviparous animals, many are satisfied with instinctively depositing their ova in situations and under circumstances favourable to their hatching, and then abandoning them, so that they can never know their progeny. This is the case with insects. Others, again, as birds, subject their ova to incubation, and, after they have been hatched, administer nourishment to their young during the early period of existence. In the viviparous animal, these cares are still more extensive; the mother drawing from her own bosom the

nutriment needed by the infant, or suckling it.

There are yet other varieties in the generation of animals. In some, it can be performed but once during the life of the individual; in others, we know it can be effected repeatedly. Sometimes one copulation fecundates but a single individual; at others, several generations are fecundated. A familiar example of this fecundity occurs in the common fowl, in which a single access will be sufficient to fecundate the eggs for the season. In the insect tribe, this is still more strikingly exemplified. In the aphis puceron or green plant louse, through all its divisions, and in the monoculus pulex, according to naturalists, a single impregnation suffices for at least six or seven generations. There is, in this case, another strange deviation from the ordinary laws of propagation, viz., that, in the warm summer months, the young are produced viviparously, and, in the cooler autumnal months, oviparously. A single impregnation of the queen bee will serve to fecundate all the eggs she may lay for two years at least - Huberb believes for the whole of her life, but he has had numerous proofs of the former

· Virey, Philosophie d'Histoire Naturelle, p. 475, Paris, 1835.

b Nouvelles Observations sur les Abeilles, Paris, 1814 ; or English translation ; also, Good, op. citat.

She begins to lay her eggs forty-six hours after impregnation, and will commonly lay about three thousand in two months, or at the rate of fifty eggs daily.

Lastly, the young are sometimes born with the shape which they have always to maintain; at others, under forms, which are, subsequently, modified materially, as in the papilio or butterfly

genus.

Reproduction in the human species requires the concourse of both sexes; these sexes being separate, and each possessed by a distinct individual — male and female. All the acts comprising it may be referred to five great heads. 1. Copulation, the object of which is to apply the fecundating principle, furnished by the male, to the germ of the female. 2. Conception or fecundation, the prolific result of copulation. 3. Gestation or pregnancy, comprising the sojourn of the fecundated ovum in the uterus, and the development it undergoes there. 4. Delivery or accouchement, which consists in the detachment of the ovum; its excretion and the birth of the new individual: and lastly, lactation, or the nourishing of the infant on the maternal milk.<sup>a</sup>

### 1. GENERATIVE APPARATUS.

The part, taken by the two sexes in the process of generation, is not equally extensive. Man has merely to furnish the fluid necessary for effecting fecundation, and to convey it within the female. He, consequently, participates only in copulation and fecundation; whilst, in addition, the acts of gestation and lactation are accomplished by the female. Her generative apparatus is therefore more complicated, and consists of a greater number of organs.

# a. Genital Organs of the Male.

The generative apparatus of the male comprises two orders of parts: — those which secrete and preserve the fecundating fluid, and those which accomplish copulation. The first consists of two similar glands — the testes — which secrete the sperm or fecundating fluid from the blood. 2. The excretory ducts of those glands — the vasa deferentia. 3. The vesiculæ seminales, which communicate with the vasa deferentia and urethra; and 4. Two canals, called ejaculatory, which convey the sperm from the vesiculæ seminales into the canal of the urethra, whence it is afterwards projected externally. The second consists of the penis, an organ essentially composed of erectile tissues, and capable of acquiring considerable rigidity. These parts will require a more detailed notice.

Testes. — The testicles are two glands situate in a bag sus
See, on the various kinds of generation, Burdach, op. citat. p. 40; Roget's Animal and Vegetable Physiology, Amer. Edit., p. 409, Philad. 1834; Prof. T. Rymer Jones, art. Organs of Generation, in Cyclop. of Anat. and Physiol. part xiii. p. 406, Lond. Feb. 1838; and Dr. Allen Thomson, art. Generation, ibid. p. 433; Carpenter, Principles of General and Comparative Physiology, 2d edit. Lond. 1841; and Müller, Elements of Physiology, by W. Baly, p. 1430, Lond. 1842.

pended beneath the pubes, called the scrotum; the right being a little higher than the left. They are of an ovoid shape, compressed laterally, their size being usually that of a pigeon's egg, and their weight about seven and a half, or eight drachms : - Sir A. Cooper says about an ounce, and Mr. Curlinga six drachms. The left testis is generally larger than the right. Like other glands, they receive arterial blood by an appropriate vessel, which communicates with the excretory duct. The spermatic artery conveys the blood, from which the secretion has to be operated, to the testicle. It arises from the abdominal aorta at a very acute angle, is small, extremely tortuous and passes down to the abdominal ring, through which it proceeds to the testicle. When it reaches this organ, it divides into two sets of branches, some of which are distributed to the epididymis, and others enter the testicle at its upper margin, and assist in constituting its tissue. The excretory ducts form, in the testicle, what are called the seminiferous vessels or tubuli seminiferi. These terminate in a white cord or nucleus-situate at the upper and inner part of the organ, where the excretory duct commences - which is called the corpus Highmorianum or sinus of the seminiferous vessels. Besides these anatomical elements of the testes, there are also - 1. Veins, termed spermatic, which return the superfluous blood back to the heart. These arise in the very tissue of the organ, and form the spermatic plexus, the divisions of which collect in several branches, that pass through the abdominal ring, and unite into a single trunk, which subsequently divides again into another plexus, termed corpus pampiniforme. This has been described as peculiar to the human species, and as a diverticulum for the blood of the testicle, whose functions are intermittent. These veins ultimately terminate on the right side in the vena cava, and on the left in the renal vein. 2. Lymphatic vessels, in considerable number, the trunks of which, after having passed through the abdominal ring, open into the lumbar glands. 3. Nerves, partly furnished by the renal and mesenteric plexuses and by the great sympathetic, partly by the lumbar nerves, and which are so minute as not to be traceable as far as the tissue of the testicle. An outer membrane or envelope to the whole organ, called tunica albuginea or peritestis. This is of an opaque white colour, of an evidently fibrous and close texture; it envelopes and gives shape to the organ, and sends into the interior of the testicle numerous filiform, flattened prolongations, which constitute incomplete septa or partitions. These form triangular spaces, filled with seminiferous vessels, which pass, with considerable regularity, towards the superior margin and the corpus Highmorianum.

These elements, united, constitute the testicle, the substance of which is soft, of a yellowish-gray colour, and divided by prolongations of the tunica albuginea, and the tunica vasculosa of Sir A. Cooper, into a number of lobes and lobules. It seems to be formed

a Practical Treatise on Diseases of the Testis, &c., p. 9, Lond. 1843.


of an immensity of very delicate, tortuous filaments, interlaced and convoluted in all directions, loosely united, between which are ramifications of the spermatic arteries and veins.

According to Monro, secundus, the seminiferous tubes of the testicle do not exceed the 100 th part of an inch in diameter, and when filled with mercury, the 120th part of an inch. He calculated, that the testis consists of 62,500 tubes, supposing each to be one inch long; and that, if the tubes were united, they would be 5208 feet and four inches long. Lauth estimates their length at 1750 feet, and Krause at 1015. The tubuli seminiferi finally terminate in straight tubes, lis; the most external layer is called vasa recta, which unite near the centre of the testis, in a complicated arrangement, bearing the name rete testis or rete tinum testis, giving off nument, bearing the name rete testis or rete tinum testis, giving off nument, bearing the name rete testis or rete ducts proceed upwards and backwards to penetrate the corpus Highmorianum and the tunica albuginea. These ducts are called vasa efferentia. Each of them is afterwards convoluted upon itself, so as to form a conical body, called conus vasculosus, having its base backwards; and at its base, the tube of each cone enters the tube of which the epididymis is formed. The epididymis

is the prismatic arch, B, C, Fig. 210, which rests vertically on the back of the testicle and adheres to it by the reflection of the tunica vaginalis, so as to appear a distinct part from the body of the testis. It is enlarged at both ends; the upper enlargement being formed by the coni vasculosi, and B called the globus major; the lower called the globus minor. epididymis is

Fig. 209.

Transverse Section of Testis. tunica vaginalis reflexa; and that in contact with the organ, tunica vaginalis propria. 2. Tunica albuginea. 3. Mediasdiated direction to internal surface of tunica albuginea, The cut extremities of vessels below the number belong to the rete testis; and those above to arteries and veins of the organ. 4. Tunica vasculosa, or pia mater testis. 5. One of the lobules, consisting of con-volutions of tubuli seminiferi, and terminating by a single duct - the vas rectum. Corresponding lobules are seen between other fibrous cords of mediastinum. 6. Section of epididymis. - (Wilson.)




Male Organs.

formed by a single Left Hand Fig. The testicle covered by its membranes, and seeming like one body. Right Hand Fig. The testicle freed from convoluted tube, the its outer coat. A. Body of the testicle. B. Commencement of the epididymis, or globus major. C. The small head or globus mifourth of a line in dia-nor. D. The vas deferens. — (Sir C. Bell.)


a Elements of the Anat. of the Human Body, by Monro, tertius, ii. 179, Edinb. 1825.

meter. When the tube attains the lower end of the globus minor, it becomes less convoluted, enlarges, turns upwards, and obtains the name of vas deferens.



Human Testis injected with Mercury.

a. a. Lobules formed of seminiferous tubes. b. Rete testis. c. Vasa efferentia. d. Plexuses of the efferent vessels passing into the head of the epididymis c. f. Body of the epididymis. g. Its appendix; its tail or cauda. i. Vas deferens. — (Lauth.)



Plan of the Structure of the Testis and Epididymis. a, a. Tubuli seminiferi. a,\* a\*. Their anastomoses. The other references as in the last figure. - (Lauth.)

Into the angle, made by the epididymis where it terminates in the vas deferens, is poured the secretion of the appendix or vasculum aberrans, which closely resembles a single lobule in structure. Its use is unknown.

The marginal figures exhibit the arrangement of the testis and its ducts.

The testes of most animals, that procreate but once a year, are comparatively small during the months when they are not excited. In man, the organ before birth, or rather during the greatest part of gestation, is an abdominal viscus; but, about the seventh month of fætal existence, it gradually descends through the abdominal ring into the scrotum, which it reaches in the eighth month, by a mechanism to be described hereafter. In some cases it never descends, but remains in the cavity of the abdomen, giving rise to considerable mental distress in many instances, and exciting the idea, that there may be a total absence of the organs, or that if they exist, they cannot effect the work of reproduction. The uneasiness is needless, provided there be other evidences of virility, the descent appearing to be by

no means essential. It has been sufficiently demonstrated that individuals, so circumstanced, are capable of procreation. In many animals, the testicles are always internal; whilst, in some, they appear only in the scrotum during the season of amorous excitement. Fodéré has indeed asserted, that the crypsorchides or testicondi, or those whose testes have not descended, are occasionally remarked for the possession of unusual prolific powers and sexual vigour.a

Dr. Marshall states, that in the examination of 10,800 recruits he found 5 in whom the right, and 6 in whom the left testicle was not apparent. He met with but one instance in which both testi-

cles had not appeared.b

It appears, that there is a set of barbarians at the back of the Cape of Good Hope, who are generally possessed of but one testicle, or are monorchides; and Linnæus, under the belief, that this is a natural defect, has made them a distinct variety of the human species. Mr. Barrow has noticed the same singularity; but Dr. Goode thinks it doubtful, whether, like the want of beard amongst the American savages, the destitution may not be owing to a barbarous custom of extirpation in early life. The deviation is not, however, more singular than the unusual formation of the nates and of the genital organs of the female in certain people of those regions, to which we shall have to refer. The possession of a single testicle appears to be sufficient for procreation. Occasionally three testicles exist.d

At times, the testicles are extremely small, but capable of executing all their functions. Mr. Wilsone was consulted by a gentleman, on the point of marriage, respecting the propriety of his entering into that state, whose penis and testicles very little exceeded in size those of a youth of eight years of age. He was twenty-six years old, but had never experienced sexual desire until he became acquainted with the lady whom he proposed to make his wife; after which he had repeated erections, with nocturnal emissions. He married; became the father of a family; and when he was twenty-eight years old, the organs had increased to the usual size

of those of the adult.

In certain cases, the testes are drawn up against the abdominal ring so as to encourage the idea, that there are no testes in the scrotum, and Professor Grossf has given the cases of two boys, one fourteen, the other eleven years of age, who were said to have been castrated, and a medical practitioner deposed to the absence of the testes; which, however, were found to be situate in the groin,

b Hints to Young Medical Officers in the Army, p. 83. Physiological Proem to class Genetica, Study of Medicine, vol. iv.

d See a case of this kind by Dr. F. Macann, in Edinb. Med. and Surg. Journ., Jan. 1843; and others, referred to in Curling, Practical Treatise on Diseases of the Testis, &c., p. 50, Lond. 1843.

· Lectures on the Structure and Physiology of the Male Urinary and Genital Organs, &c. Lond., 1821.

a " Ces organes paraissant tirer du bain chaud ou ils se trouvent plongés plus d'aptitude à la secrétion que lorsqu'ils sont descendus au dehors dans leurs enveloppes ordinaires !" - Traité de Médecine Légale, i. 370, Paris, 1813.

t Western Journal of Medicine and Surgery, May, 1841, p. 355.

a little below the external ring, whence, by a little traction, they

could be easily forced down into the scrotum.

The testicle is connected with the abdominal ring by means of the spermatic cord, a fasciculus of about half an inch in diameter, which can be readily felt through the skin of the scrotum. It is formed, essentially of the vessels and nerves that pass to or from the testicles; — the spermatic artery, spermatic veins, lymphatics and nerves of the organ, and the vas deferens, or excretory duct. These are bound together by means of cellular tissue; and, externally, a membranous sheath of a fibrous character envelopes the cord, and keeps it distinct from the surrounding parts, and especially from the scrotum. When the cord has passed through the abdominal ring, its various elements are no longer held toge-

ther, but each passes to its particular destination.

The scrotum or purse is a continuation of the skin of the inner side of the thighs, the perineum, and the penis. It is symmetrical, the two halves being separated by a median line or raphe. The skin is of a darker colour here than elsewhere; is rugous, studded with follicles, and sparingly furnished with hair. This may be considered its outermost coat. Beneath this is the dartos, - a reddish, cellular membrane, which forms a distinct sac for each testicle; and a septum — the septum scroti — between them. Much discussion has taken place regarding the nature of the dartos; some supposing it to be muscular, others cellular. Breschet and Lobstein affirm, that it does not exist in the scrotum before the descent of the testes, and they consider it to be formed by the expansion of the gubernaculum testis. Meckel, however, suggests, that it constitutes the transition between the cellular and muscular tissues, and that there exists between it and other muscles the same relation that there is between the muscles of the superior and inferior animals. It consists of long fibres considerably matted together, and passing in every direction, but which are easily separable by distension with air or water, and by slight maceration.a The generality of anatomists conceive it to be of a cellular character, yet it is manifestly contractile, corrugates the scrotum, and probably consists of a muscular tissue also. Professor Horner, bindeed, affirms, that he dissected a subject in January, 1830, in which the fibres were evidently muscular, although interwoven. Beneath the dartos a third coat exists, which is muscular: - it is called the cremaster or tunica erythroïdes. It arises from the lesser oblique muscle of the abdomen, passes through the abdominal ring, aids in the formation of the spermatic cord, and terminates insensibly on the inner surface of the scrotum. It draws the testicle upwards. The cellular substance, that connects the dartos and cremaster with the tunica vaginalis, has been considered by some as an additional coat, and termed tunica vagi-


b Special Anat. and Histology. 6th edit. Philad. 1843; and Lessons on Practical Anatomy, 3d edit. p. 273, Philad. 1836.

<sup>\*</sup> Meckel's Handbuch, u. s. w., Jourdan's French translation, iii. 622, Paris, 1825, and Weber's Hildebrandt's Handbuch der Anatomie, iv. 382, Braunschweig, 1832.

nalis communis. The tunica vaginalis or tunica elytroïdes is a true serous membrane, enveloping the testicle and lining the scrotum; having, consequently, a scrotal and a testicular portion. We shall see, hereafter, that it is a dependence of the peritoneum, passing before the testicle in its descent, and afterwards becoming separated from any direct communication with the abdomen.

The vas deferens or excretory duct of the testicle commences at the globus minor of the epididymis, (C, Fig. 210,) which is itself, we have seen, formed of a convoluted tube. This, when unfolded, according to Monro, measures as much as thirty-two feet. As soon as the vas deferens quits the testicle, it joins the spermatic cord, passes upwards to the abdominal ring, separates from the bloodvessels on entering the abdomen, and descends downwards and inwards to the posterior and inferior part of the bladder, passing between the bas-fond of the latter and the ureter. It then converges towards its fellow along the under extremity of the bladder, at the inner margin of the vesicula seminalis of the same side, and ultimately opens into the urethra near the neck of the bladder. (Fig. 206 and Fig. 213.) At the base of the prostate, it receives a canal from the vesicula, and continues its course to the urethra under the name of ejaculatory duct. The vas deferens has two coats, the outermost of which is very

The inner coat is thin, and belongs to the class of mucous membranes. The vesiculæ seminales, 19, Fig. 206, and Fig. 214, are considered to be two convoluted tubes, - one on each side, - which are two inches or two inches and a half long, and six or seven lines broad at the fundus, and are situate at the lower fundus of the bladder, between it and the rectum and behind the prostate gland. At the anterior extremities they approach each other very closely, being separated only by the vasa deferentia. When inflated and dried, they present the appearance of cells; but



Posterior View of Male Bladder.

1. Body of bladder.

2. Fundus.

3. Inferior fundus or base.


4. Urachus.

5. 5. Ureters.

6. 6. Vasa deferentia.


7. 7. Vesiculæ seminales. Triangular area, corresponding with trigonum vesicæ through which bladder would be pierced, in puncturing bladder through rectum. The dotted line forming base of this triangular area, marks extent of recto-vesical fold of peritoneum. — (Wilson.)

firm and almost cartilaginous; but its structure is not manifest. The inner coat is thin, and be-



Section of the Vesicula Seminales, &c.
V. Section of vas deferens. S. Section of vesicula seminalis. E. Section of ejaculatory duet.

are generally conceived to be tubes, which, being convoluted, are brought within the compass of the vesiculæ. When dissected and stretched out, they are four or five inches long by about one-fourth of an inch in diameter. Amussat, however, denies this arrangement of the vesiculæ: he affirms, that he has discovered them to



Section of Bladder, Prostate, and Penis.

1. Urachus attached to upper part of fundus of bladder. 2. Recto-vesical fold of peritoneum, at its point of reflection from base of bladder upon anterior surface of rectum. 3. Opening of right ureter. 4. A slight ridge, formed by muscle of ureter, and extending from termination of ureter to commencement of urethra. This ridge forms lateral boundary of trigonum vesicæ. 5. Commencement of urethra; the elevation of mucous membrane immediately behind figure is uvula vesicæ. Constriction of bladder at this point is neck of bladder. 6. Prostatic portion of urethra.

7. Prostate gland; difference of thickness of gland, above and below urethra, is shown. 8. Isthmus, or third lobe of prostate; immediately beneath which ejaculatory duct is seen passing. 9. Right vesicula seminalis; vas deferens is seen to be cut short off, close to its junction with ejaculatory duct. 10. Membranous portion of urethra. 11. Cowper's gland of right side, with its duct. 12. Bulbous portion of urethra: throughout whole length of urethra of corpus spongiosum, numerous lacunæ are seen. 13. Fossa navicularis. 14. Corpus cavernosum, cut somewhat obliquely to right side, near its lower part. Character of venous-cellular texture is shown. 15. Right crus penis. 16. Near upper part of corpus cavernosum, section has fallen a little to left of middle line; a portion of septum pectiniforme is consequently seen. This figure also indicates thickness of fibrous investment of corpus cavernosum, and its abrupt termination at base of (17) glans penis. 18. Lower segment of glans. 19. Meatus urinarius. 20. Corpus spongiosum. 21. Bulb of corpus sponglosum.—

(Wilson.) 1. Urachus attached to upper part of fundus of bladder. 2. Recto-vesical

be formed of a minute canal of considerable length, variously convoluted, the folds of which are united to each other by cellular filaments, like those of the spermatic vessels. At the anterior part, termed neck, a short canal passes off, which

of contracting in the act of ejaculation; and an internal lining, of a white, delicate character, a little like that which lines the interior of the gall-bladder, and supposed to be mucous. Although the vesiculæ are manifestly contractile, no muscular fibres have been detected in them. They are found filled, in the dead body, with an opaque, thick, yellowish fluid, very different, in appearance, from the sperm ejaculated during life.

The prostate gland, (Fig. 206 and Fig. 215,) is an organ of very dense tissue, embracing the neck of the bladder, and penetrated by the urethra, which traverses it much nearer its upper than its lower surface. The base is directed backwards, the point forwards, and its inferior surface rests upon the rectum, so that, by passing the finger into the rectum, enlargements of the organ may be detected. The prostate was once universally esteemed glandular, and it is still so termed. It is, now, generally and correctly regarded as an agglomeration of several small follicles, filled by a viscid whitish fluid. These follicles have numerous minute excretory ducts, which open on each side of the caput gallinaginis.

The glands of Cowper are two small, oblong bodies; of the size of a pea; of a reddish colour, and of a somewhat firm tissue. They are situate anterior to the prostate, parallel to each other, and at the sides of the urethra. Each has an excretory duct, which creeps obliquely in the spongy tissue of the bulk, and opens before the

verumontanum.

The male organ or penis consists of the corpus cavernosum and corpus spongiosum; parts essentially formed of an erectile tissue, and surrounded by a very firm elastic covering, which prevents over-distension, and gives form to the organ. The corpora cavernosa constitute the great body of the penis. They are two tubes which are united and separated by an imperfect partition. Within them a kind of cellular tissue exists, into which blood is poured, so as to cause erection. The posterior extremities of these cavernous tubes are called crura penis. These separate in the perineum, each taking hold of the ramus of the pubis; and, at the other ex tremity, the cavernous bodies terminate in rounded points under the glans penis. The anatomical elements of the internal tissue of the corpora cavernosa are, — the ramifications of the cavernous artery, which proceeds from the internal pudic; those of a vein bearing the same name; and probably, nerves, although they have not been traced so far. All these elements are supported by filamentous prolongations from the outer dense envelope. A difference of opinion prevails amongst anatomists with regard to the precise arrangement of these prolongations. Some consider them to form cells, or a kind of spongy structure, on the plates of which the ramifications of the cavernous artery and vein and of the nerves terminate, and into which the blood is extravasated. Others conceive, that the internal arrangement consists of a plexus of minute arteries and veins, supported by the plates of the outer membrane, interlacing like the capillary vessels, but with this addition, that in place of the minute veins becoming capillary in the plexus, they are of greater size, forming very extensible dilatations and networks, and anastomosing freely with each other. If the cavernous artery be injected, the matter first fills the ramifications of the artery, then the venous plexuses of the cavernous bodies, and it ultimately returns by the cavernous vein, having produced erection. The same effect is caused still more readily by injecting the cavernous vein. J. Müller, who has investigated the structure of the male organ, has discovered two sets of arteries in the organ differing from each other in size, mode of termination, and uses: the first he calls Rami nutritii, which are distributed upon the parietes of the veins and throughout

the spongy substance, differing in no respect from the nutritive arteries of other parts. The second set he calls arteriæ helicinæ. They differ from the nutritive vessels in form, size, and distribution. They are short, and are given off from the larger branches as well as from the finest twigs of the artery; most of them come off at a right angle, and project into the cavity of the spongy substance, either terminating abruptly or swelling out into a clublike process without again subdividing. Almost all these arteries have this character, that they are bent like a horn, so that the end describes half a circle or somewhat more. These arteries have a great resemblance to the tendrils of the vine, whence their name - arterix helicinx. A minute examination of them, either with the lens or with the microscope, shows, that, although they at all times project into the venous cavities of the corpora cavernosa, they are not entirely naked, but are covered with a delicate membrane, which under the microscope appears granular.a The views of Müller are embraced by Krause and Hyrtl, but the researches of Valentine are not in accordance with them. The result of numerous examinations has convinced him, that the helicine arteries are not peculiar vessels, but merely minute arteries that have been divided or torn, and that the real distribution of the vessels of the corpora cavernosa follows in every respect the most simple laws. The investigations of Müller have led him to infer, that, both in man and the horse, the nerves of the corpora caver-

Fig. 216.

Section of the Penis. A. External membrane or sheath of the penis. B. Corpus cavernosum, D. Corpus spongiosum urethræ.

nosa are made up of branches proceeding from the organic as well as the animal system, whilst the nerves of animal life alone provide the nerves

of sensation of the penis.d

Attached to the corpora cavernosa, and running in the groove beneath them, is a spongy body of similar structure, - the corpus spongiosum urethræ, - through which the urethra passes. It commences, posteriorly, at the bulb of the urethra, - already described under the Secretion of Urine, - and terminates anteriorly in the glans, which is, in no wise, a dependency of the corpora cavernosa, but is separated from them by a portion of their outer membranes; so that erection may take place in the one, and not

a For a farther description of these vessels, see J. Müller, art. Erectiles Gewebe, Encycl. Wörterbuch der Medic. Wissench. xi. 452, Berlin, 1834; Handbuch der Physiologie, u. s. w., Baly's translation, Lond. 1838; and Abhandlungen der Königlich. Akademie der Wissenschaft. zu Berlin, s. 93, Berlin, 1837; also, Dr. Hart, in art. Erectile Tissue, in Cyclop. Anat. and Physiol. part x., p. 146, June, 1837.

b Mandl, Manuel d'Anatomie générale, p. 197, Paris, 1843.

· Müller's Archiv. für Anatomie, u. s. w., and Lond. Med. Gazette, June 23, 1838, p. 543; and Henle, Allgemein. Anatom. s. 485, Leipz. 1841.

d Lond. Med. Gazette, April 23d, 1836, and Abhandlung, u. s. w., s. 117.

SPERM. 325

simultaneously in the other; and injections into the corpora cavernosa of the one do not pass into those of the other. The glans appears to be the final expansion of the erectile tissue which surrounds the urethra. The posterior circular margin of the glans is called the corona glandis, and behind this is a depression termed the cervix, collum or neck. Several follicles exist here, called the glandulæ odoriferæ Tysoni, which secrete an unctuous humour called the smegma præputii, which often accumulates largely, where cleanliness is not attended to.

The penis is covered by the skin, which forms, towards the glans, the prepuce or foreskin. The cellular tissue, which unites it to the organ is lax, and never contains fat. The inner lamina of the prepuce being inserted circularly into the penis, some distance back from the point, the glans can generally be denuded, when the prepuce is drawn back. The under and middle part of the prepuce is attached to the extremity of the glans by a duplicature, called the frænum præputii, which extends to the orifice of the urethra. The skin is continued over the glans, but it is greatly modified in its structure, being smooth and velvety, highly delicate, sensible, and vascular.

Lastly. In addition to the acceleratores urinx, the transversus perinei, the sphincter ani, and the levator ani muscles, which we have described as equally concerned in the excretion of urine and semen, the erector penis or ischio-cavernosus muscle is largely connected with the function of generation. The genital organs of man are, in reality, merely an apparatus, for a glandular secretion of which the testicle is the gland; the vesiculæ seminales are supposed to be the reservoirs; and the vas deferens and urethra the excretory ducts; — the arrangement which we observe in the penis being for the purpose of conveying the secreted fluid into the parts

of the female.

#### 1. SPERM.

The sperm, sperma or semen is secreted by the testicles from the blood of the spermatic artery, by a mechanism, which is no more understood than that of secretion in general. When formed, it is received into the tubuli seminiferi, and passes along them to the epididymis, the vas deferens, and the vesiculæ seminales, where it is generally conceived to be deposited, until it is projected into the urethra, under the venereal excitement. That this is its course is sufficiently evidenced by the arrangement of the excretory ducts, and by the function which the sperm has to fulfil. De Graaf, however, adduces an additional proof. On tying the vas deferens of a dog, the testicle became swollen under excitement, and ultimately the vas deferens gave way between the testicle and the ligature. The causes of the progression of the sperm through the ducts are, — the continuity of the secretion by the testicle, and a

De Virorum Organ. Gener. Inserv., in Med. Oper. Omn. Amstel. 1705.

contraction of the excretory ducts themselves. These are the effi-

cient agents.

It has been a question with physiologists, whether the secretion of the sperm be constantly taking place, or whether, as the function of generation is accomplished at uncertain intervals, the secretion may not likewise be intermittent. It is impossible to arrive at any positive conclusion on this point. It would seem, however, unnecessary for the secretion to be effected at all times; and it is more probable, that when the vesiculæ seminales are emptied of their contents, during coition, a stimulus is given to the testes by the excitement, and they are soon replenished. This, however, is more and more difficult in proportion to the number of repetitions of the venereal act, as the secretion takes place at best but slowly. By some, the spermatic and pampiniform plexuses have been regarded as diverticula to the testes during this intermission of action. The sperm passes slowly along the excretory ducts of the testicle, owing partly to the slowness of the secretion, and partly to the arrangement of the ducts, which, as we have seen, are remarkably convoluted, long, and minute. The use of the vesiculæ seminales has been disputed. The majority of physiologists regard them to be reservoirs for the sperm, and to serve the same purpose as the gall-bladder in the case of the bile. Others, however, have supposed, that they secrete a fluid of a peculiar nature, the use of which may probably be to dilute the sperm; and others, again, infer, that they are both seminal reservoirs, and secreting organs, furnishing mucus, or perhaps some other fluid for admixture with the semen. Dr. John Davy found spermatozoa in the fluid of the vesiculæ, but except in two instances no animalcules could be seen in the fluid expressed from the divided substance of the testes. He invariably found, however, extremely minute, dense spherules, which he conjectures to be the ova of the spermatozoa. These are manifestly not essential to the function of generation, as they do not exist in all animals. The dog and cat kind, the bear, opossum, sea-otter, seal, &c., possess them not; and there are several in which there is no direct communication between the duct and the vas deferens, which open separately into the urethra. This circumstance, however, with the fact, that they generally contain, after death, a fluid of different appearance and properties from those of the sperm, - with the glandular structure, which their coats seem to possess, in many instances, - is opposed to the view, that they are simple reservoirs for the semen, and favours that which ascribes to them a peculiar secretion. Where this communication between the duct of the vesicles and the vas deferens exists, a reflux of the semen may take place, and an admixture between the sperm and the fluid secreted by them. It is not improbable, however, as Adelone suggests, that all the excretory ducts

Dr. John Davy, in Edinb. Med. and Surg. Journ. for July, 1838, p. 12; and Researches, Physiological and Anatomical, Dunglison's Amer. Med. Libr. Edit. p. 363, Philad. 1840.
 Ibid., p. 373.

<sup>·</sup> Physiologie de l'Homme, 2de édit. iv. 15, Paris, 1829.

SPERM. 327

of the testicle may act as a reservoir; and in the case of animals, in which the vesiculæ are wanting, they must possess this office exclusively. If we are to adopt the description of Amussat as an anatomical fact, the vesiculæ themselves are constituted of a convoluted tube, having an arrangement somewhat resembling that

which prevails in the excretory ducts of the testes.a

That the excretory ducts of the testes may serve as reservoirs is proved by the fact, that impregnation is practicable after thorough castration. This has been doubted both as regards animals and man, but there is no question of the fact as regards the former.b The author's respectable friend, Dr. Pue, of Baltimore, related to him unquestionable instances of the kind. In one case, a boar was observed on one side of a hedge, striving to get at some sows in heat on the other side. The boar was castrated, and no inconvenience being apprehended, he was turned loose into the field with the sows. In five minutes after the operation, he had intercourse with one of the sows, and subsequently with others. The first sow brought forth a litter, but none of the others were impregnated. In another case, after a horse had been castrated, it was recollected, that the male organ had not been washed - which, it seems, is looked upon as advisable. To save inconvenience, it was suggested, that the same effect might be produced by putting him to a mare, then in the stable, and in heat. This was done, and, in due time, the mare brought forth a foal, unequivocally the result of this sexual union. Mr. Walton Hamilton, - a great breeder of horses, in Saratoga county, New York, - informed the author's friend, Mr. Nicholas P. Trist, United States consul at the Havana, that he, also, had known several instances of impregnation after castration.c

It is to be presumed, that the power of procreation can exist for a short time only after the operation; yet a secretion may take place from the lining membrane of the ducts, and vesiculæ, and from the prostate and other follicles, but this secretion cannot supply the place of the sperm. Sir A. Cooper gives the case of a man, who stated to him, that for nearly the first twelve months after complete castration, he had emissions in  $coit\hat{u}$ , or the sensation of emissions. Afterwards, he had erections and intercourse at distant intervals, but without the sensation of emission.<sup>d</sup>

It has been asked, how does it happen, that the sperm, in its progress along the vas deferens, does not pass directly on into the urethra by the ejaculatory duct, instead of reflowing into the spermatic vesicles? This, it has been imagined, is owing to the existence of an arrangement at the opening of the ejaculatory duct into the urethra, similar to that which prevails at the termination of the

d Observations on the Structure and Diseases of the Testis, Lond. 1830; and J. Müller, in art. Erection, Encyclopad. Wörterb. der Medic. Wissensch. xi. 460, Berl. 1834.

<sup>\*</sup> Magendie's Précis, &c. ii. 348. b Varro, De Re Rusticâ, lib. ii. cap. 5. c See some remarks, by the author, in his American Medical Intelligencer, p. 146, July 15, 1837; and by Dr. Warrington, ibid. p. 244, Oct. 1.

choledoch duct in the duodenum. It is affirmed, by some, that the prostate exerts a pressure on the ductus ejaculatorius, and that the opening of the duct into the urethra is smaller than any other part of it; by others, that the ejaculatory ducts are embraced, along with the neck of the bladder, by the levator ani, and consequently, that the sperm finds a readier access into the ducts of the vesiculæ seminales.

The sperm — lac maris, male's milk, propagatory or genital liquor, vitale virus, vital or quickening venom - is of a white colour, and of a faint smell, which, owing to its peculiar character, has been termed spermatic. This smell would seem to be derived from the secretions of the vesiculæ seminales, prostate, and mucous follicles of the urethra, as pure semen taken from the epididymis or vas deferens does not possess it." It is of a viscid consistence, of a saline, irritating taste, and appears composed of two parts, the one more liquid and transparent, and the other more grumous. In a short time after emission, these two parts unite and the whole becomes more fluid. When examined chemically, the sperm appears to be of an alkaline, and albuminous character. Vauquelinb analyzed it and found it to be composed, - in 1000 parts, - of water, 900; animal mucilage, 60; soda, 10; calcareous phosphate, John's analysise accords with this. Berzelius affirms, that it contains the same salts as the blood along with a peculiar animal matter - spermatine. After citing these analyses, Raspaild ob-. serves, that if any thing be capable of humiliating the pride of the chemist, it is assuredly the identity he is condemned to discover amongst substances, which, notwithstanding, fulfil such different

No analysis has been made of the sperm as secreted by the testicle. The fluid examined has been the compound of the pure sperm and the secretions of the prostate gland and of those of Cowper. The thicker, whitish portion is considered to be the secretion of the testicles;—the more liquid and transparent, the

fluids of the accessory glands or follicles.

Some authors have imagined, that a sort of halitus or aura is given off from the sperm, which they have called the aura seminis, and have considered to be sufficient for fecundation. The fallacy of this view will be exhibited hereafter. Others have discovered, by the microscope, numerous minute bodies in the sperm, seminal animalcules, — spermatozoa or zoospermes, — which they have conceived to be important agents in generation.

By careful examination, according to Wagner, fother minute,

2 Curling, Practical Treatise on Diseases of the Testis, &c., p. 37, Lond. 1843.

b Annales de Chimie, ix. 64.

c Chemische Tabellen des Thierreichs, s. 169, Nürnberg, 1814; cited in Burdach's Physiologie, u. s. w., i. 111.

d Chimie Organique, p. 386, Paris, 1833.

Adelon, in art. Generation, of Dict. de Méd. tom. x.; and Physiologie de l'Homme, iv. 17. See, also, Burdach's Physiologie, i. 112, for various opinions on this subject; and M. Donné, in Gazette Médicale de Paris, Juin 3, 1837.

f Elements of Physiology, translated from the German, by Robert Willis, M.D., part i. p. 4, Lond. 1841. See, also, Mandl, Manuel d'Anatomie générale, p. 494, Paris,

1843.

329 SPERM.

round, granulated bodies may almost always be detected; which are, in all cases, much less numerous than the spermatozoa. These bodies he distinguishes by the names seminal granules - granula seminis. Both elements of the sperm are suspended in a small quantity of perfectly homogeneous fluid, transparent and clear as water. "Pure semen, therefore, in its most perfect state, consists principally of seminal animalcules and seminal granules, both of which are enveloped in a small quantity of fluid." This fluid

Wagner calls liquor seminis; and he suggests, in connexion with the discoveries of Schwann and Schleiden, referred to at page 179 of this volume, whether, in the development of the spermatozoa, the liquor seminis may not be regarded as a matrix, (Zellenkeimstoff, cystoblastema, Schwann,) in which the granular nuclei are developed as cytoblasts, which again put forth their covering or cyst as a cellular wall: the finely granular contents would then have to be considered as the cell-fluid. The cytoblasts disappear as soon as the spermatozoa are evolved in their contents, and the cells burst and cast out the animalcules, as the cells of the algæ scatter abroad their sporules.a These animalcules, however, have been denied to be peculiar to this fluid, and have been regarded as infusory animalcules, similar to those met with Spermatoroa, magnified from in all animal infusions; by others, they have 900 to 1000 diameters. been esteemed organic molecules of the sperm. 900 to 1000 diameters. from the was deferens of a man shortly after death. in all animal infusions; by others, they have M. Virey, b-a physiologist, strangely fantastic in a. Flat surface. b. Profile his speculations, — conceives, that as the pollen view, margin presenting. of vegetables is a collection of small capsules, surface presenting. d. Sper-containing within them the true fecundating not unlike a proboscis from principle, which is of extreme subtilty, the pre-its body. c. Seminal gratended spermatic animalcules are tubes contain-nules. ing the true sperm, and the motion we observe in them is owing to the rupture of the tubes; whilst Raspaile is led to think, that they are mere shreds, (lambeaux,) of the tissues of the generative organs, ejaculated with the sperm, which describe involuntary movements by virtue of the property they possess of aspiring and expiring. In confirmation of this, he states, that if we open an ovary of the mussel, we may observe, alongside the large ovules, myriads of moving shreds, whose form and size are infinitely varied,

Fig. 217.

and which possess nothing resembling regular organization. They bear evident marks of laceration. Now, these shreds, he conceives, may affect greater regularity in certain classes of animals

<sup>2</sup> Wägner, op. cit. p. 27. b Art. Generation, in Dict. des Sciences Médicales; and Philosophie d'Histoire c Op. citat. p. 389. Naturelle, Paris, 1835.

of a more elevated order; but, he concludes, that however this may be, the spermatic animalcules, which have hitherto been classed amongst those incertæ sedis, may be provisionally placed in the genus cercaria — that is, amongst infusory, agastric animals having a kind of tail — which Raspail considers the simplest of animated beings, and to live only by "aspiration and expiration." Wagner also remarks, that the expression cercaria seminis, applied to the spermatozoon, can only be a collective title, and that the manifold forms of spermatozoa, which he has found to occur in the seminal fluid of a great number of animals, must be viewed in the light of so many different species. Ehrenberg refers them to the haustellate entozoa."

The author has examined the sperm with microscopes of high magnifying power, but without being able to satisfy himself, that the minute bodies, contained in it, are animalcular. In a powerful hydroxygen microscope, not the slightest appearance of animalcules presented itself, but this may have been owing to the great intensity of light. Sir Everard Home and Mr. Bauer were equally unsuccessful, and they were led to conclude, that the appearance of living animalcules in the semen is not real, but the effect of a microscopic deception. Wagnere considers, that they are essential elements of the seminal fluid, and bear a specific relation to the generative act, and that they are thus far comparable to the blood-globules; which present themselves in the same manner as essential typically organized constituents of the blood amid the liquor sanguinis, just as the spermatozoa present themselves amid the liquor seminis. The question of their animality he considers to be undetermined as their internal organization had not been detected. In the appendix, however, to Dr. Wagner's work, Dr. Willisd remarks, that in the examination of the spermatozoa of the bear, Dr. Valentine had settled the question of the organization and consequently the true animal nature of the seminal animalcule; but the question is not regarded as settled by this statement. Dr. Carpenter, indeed, considers, that there is but little reason to consider them independent animalcules. He esteems them to be bodies having an inherent power of motion, not exceeding in their activity ciliated epithelium cells, and even blood corpuscles; and he thinks there is no evidence, that their function is any higher than that of the pollen-tube of plants, which conveys into the ovulum the germ of the first cells of the embryo. The presence of the spermatozoa, whole or broken into fragments, may aid in detecting nocturnal emissions, and be of assistance in cases of alleged rape.g

<sup>a</sup> For a full account of these animalcules, see Wagner, op. citat. p. 6.

5 J. Müller, Elements of Physiology, translated by Baly, p. 1477, Lond. 1842.

b Lect. on Comp. Anat. v. 337, Lond. 1828. Cop. citat. p. 34. d Ibid. p. 228. Nov. Act. Acad. C. L. Natur. Curios. vol. xi. 1839.

Human Physiology, § 733, Lond. 1842; or Principles of General and Comparative Physiology, § 607, Lond. 1841; and Todd and Bowman, Physiological Anatomy, &c., of Man, p. 66, Lond. 1843.

It would seem that the spermatozoa are not found solely in the sperm; for Mr. Lloyd, of Saint Bartholomew's Hospital, London, stated recently to the Medico-Chirurgical Society, that in two cases of common hydrocele, in which he examined microscopically the fluid withdrawn by tapping, he had found numerous spermatozoa. He counted forty of these in one drop of the fluid. Some of them were observed to retain their power of motion for three hours after the fluid had been withdrawn. In the fluid of many other cases of hydrocele, Mr. Lloyd was unable to detect these animalcules.

The agency of the sperm in fecundation will be considered hereafter. It may be observed, however, that in all examinations of it, whether by the microscope or otherwise, we must bear in mind, the caution to which we have adverted more than once, as applicable to the examination of animal fluids in general, — that we ought not to conclude positively, from the results of our observations of the fluids when out of the body, that they possess precisely the same characteristics when in it; and this remark is especially applicable to the sperm, which varies manifestly in its sensible pro-

perties a short time after it has been excreted.

The sperm being the great vivifying agent, — the medium by which life is communicated from generation to generation, — it has been looked upon as one of the most important if not the most important of animal fluids; and hence it is regarded, by some physiologists, as formed of the most animalized materials, or of those that constitute the most elevated part of the new being — the nervous system. The quantity of sperm secreted cannot be estimated. It varies according to the individual, and to his extent of voluptuous excitement, as well as to the degree of previous indulgence in venereal pleasures. Where the demand is frequent, the supply is larger; although when the act is repeatedly performed, the absolute quantity at each copulation may be less.<sup>b</sup>

# b. Genital Organs of the Female.

The genital organs of the male effect fewer functions than those of the female. They are inservient to copulation and fecundation only. Those of the female, — in addition to parts, which fulfil these offices, — comprise others for gestation and lactation.

The soft and prominent covering to the symphysis pubis — which is formed by the common integuments, elevated by fat, and, at the age of puberty, covered by hair, formerly termed tressoria — is called the mons veneris. The absence of this hair has, by the vulgar, been esteemed a matter of reproach; and it was formerly the custom, when a female had been detected a third time in incontinent practices, in the vicinity of the Superior Courts of Westmin-

<sup>\*</sup> Provincial Medical Journal, cited in Medical Examiner, July 22, 1843, p. 168.

b Theophrastus, Pliny, and Athenæus assert, that with the help of a certain herb, an Indian prince was able to copulate seventy times in twenty-four hours! — Theophr. I. c. v., Plin. I. xxvi. c. 9, and Athenæus, I. i. c. 12. See, also, art, Cas rares, in Dict. des Sciences Médicales.

ster, to punish the offence by cutting off the tressoria in open court. Occasionally its growth is excessive. Below this, are the labia pudendi or labia majora, which are two large, soft lips, formed by a duplicature of the common integument, with adipose matter interposed. The inner surface is smooth, and studded with sebaceous follicles. The labia commence at the symphysis pubis, descend to the perinæum, which is the portion of integument, about an inch and a half in length, between the posterior commissure of the labia and the anus. This commissure is called the franum labiorum, frænulum perinæi or fourchette. The opening between the labia is the vulva or fossa magna. At the upper junction of the labia, and within them, a small organ exists, called the clitoris or superlabia, which greatly resembles the penis. It is formed of corpora cavernosa, and is terminated anteriorly by the glans, which is covered by a prepuce, consisting of a prolongation of the mucous membrane of the vagina. Unlike the penis, however, it has no corpus spongiosum, or urethra attached to it; but it is capable of being made erect by a mechanism similar to that which applies to the penis; and it has two erector muscles, the erectores clitoridis, - similar to the erectores penis. Anciently, if a female was detected a fourth time in incontinence in the vicinity of the Superior Courts of Westminster, the clitoris was amputated in open court. Extending from the prepuce of the clitoris, and within the labia majora, are the labia minora or nymphæ, the organization of which is similar to that of the labia majora. They gradually enlarge as they pass downwards, and disappear when they reach the orifice of the vagina.

A singular variety is observed in the organization of those parts amongst the Bosjesmen or Bushmen, the tribe to whose peculiarities of organization we have already had occasion to refer. Discordance has, however, prevailed regarding the precise nature of this peculiarity, some describing it as existing in the labia, others in the nymphæ, and others again, in a peculiar organization; some deeming it natural, others artificial. Dr. Somerville, who had numerous opportunities for observation and dissection, asserts, that the mons veneris is less prominent than in the European, and is either destitute of hair, or thinly covered by a small quantity of a soft, woolly nature; that the labia are very small, so that they seems at times to be almost wanting; that the loose, pendulous, and rugous growth, which hangs from the pudendum, is a double fold; and that it is proved to be the nymphæ, by the situation of the clitoris at the commissure of the folds, as well as by all other circumstances; and that they sometimes reach five inches below the margin of the labia; Le Vaillantd says nine inches. Cuviere

<sup>&</sup>lt;sup>a</sup> Chitty's Practical Treatise on Medical Jurisprudence, part i. p. 390, Amer. Edit., Philad. 1836.

<sup>b</sup> Chitty, op. cit. part i. p. 391, Amer. Edit., Philad. 1836.

Medico-Chirurgical Transactions, vii. 157.
 Voyage dans l'Intérieur d'Afrique, p. 371.

e Mémoir. du Museum, iii. 266; and Broc, Essai sur les Races Humaines, p. 87, Paris, 1836.

examined the Hottentot Venus, and found her to agree well with the account of Dr. Somerville. The labia were very small; and a single prominence descended between them from the upper part. It divided into two lateral portions, which passed along the sides of the vagina to the inferior angle of the labia. The whole length was about four inches. When she was examined naked by the French Savans, this formation was not observed. She kept the tablier, ventrale cutaneum, or, as it is termed by the Germans, Schürze ('apron,') carefully concealed, either between her thighs, or yet more deeply; and it was not known, until after her death, that she possessed it. Both Mr. Barrowa and Dr. Somerville deny that the peculiarity is artificially excited.

In warm climates, the nymphæ are often greatly and inconveniently elongated, and amongst the Egyptians and other African

tribes, it has been the custom to extirpate them, or to diminish their size. This is what is meant by circumcision in the female.

The vagina is a canal, which extends between the vulva and the uterus, the neck of which it embraces. It is sometimes called the vulvo-uterine canal, and is from four to six inches long, and an inch and a half, or two inches, in diameter. It is situate in the pelvis, be-




Lateral view of Pelvic Viscera in the Female.

tween the bladder before, and the rectum behind; it is slightly curved, with the concavity forwards, and is narrower at the middle than at the extremities. Its inner surface has numerous — chiefly transverse — rugæ, which become less in the progress of age, after repeated acts of copulation, and especially after accouchement. It is composed of an internal mucous membrane, supplied with numerous mucous follicles, of a dense cellular membrane, and, between these, a layer of erectile tissue, which is thicker near the vulva; but is, by some, said to extend even as far as the uterus. It is termed the corpus spongiosum vaginæ. It is chiefly situate around the anterior extremity of the vagina, below the clitoris, and

<sup>a</sup> Travels, pp. 279, 280; see, also, Lawrence's Lectures on Physiology, Zoology, &c. Lond. 1819; and Dr. D. D. Davis, in Principles, &c., of Obstetric Medicine, i. 54, Lond. 1836.


at the base of the nymphæ; and the veins of which it is constituted are called plexus retiformis. The upper portion of the vagina, to a small extent, is covered by the peritoneum. The sphincter or constrictor vaginæ muscle surrounds the orifice of the vagina, and covers the plexus retiformis. It is about an inch and a quarter wide, and ordinarily about six inches in length; arises from the body of the clitoris, and passes backwards and downwards, to be inserted into the dense, white substance, in the centre of the perineum, which is common to the transversi perinei muscles, and the anterior point of the sphincter ani.



Anterior view of the Female Organs.

Near [the external aperture of the vagina is the hymen or virginal, or vaginal valve, which is a more or less extensive, membranous duplicature, of variable shape, and formed by the mucous membrane of the vulva, where it enters the vagina, so that it closes the canal, more or less completely. It is generally very thin, and easily lacerable: but is sometimes extremely firm, so as to prevent penetration. It is usually of a semilunar shape; sometimes oval from right to left, or almost circular, with an aperture in the middle, whilst, occasionally, it is entirely imperforate, and of course prevents the issue of the menstrual flux. It is easily destroyed by mechanical violence of any kind, as by strongly rubbing the sexual organs of infants by coarse cloths, and by ulcerations of the part; hence its absence is not an absolute proof of the loss of virginity, as it was of old regarded by the Hebrews. Nor is its presence a positive evidence of continence. Individuals

have conceived, in whom the aperture of the hymen has been so small as to prevent penetration. Its general semilunar or crescentic shape has been considered to explain the origin of the symbol of the crescent assigned to Diana — the goddess of chastity. Around the part of the vagina, where the hymen was situate, small, reddish, flattened, or rounded tubercles — the carunculæ myrtiformes seu hymenales — afterwards exist, which are of various sizes, and are formed, according to the general opinion, by the remains of the hymen. Béclard and J. Cloquet consider them to be folds of the mucous membrane. Their number varies from two to five, or six.



Female Organs of Generation.

1. Upper part of vagina. 2. Os uteri, projecting into vagina; posterior lip is seen to be longer and larger than anterior. 3. Cervix uteri. 4. Body of uterus. 5. Its fundus. 6. Broad ligament of left side, having enclosed between its layers (7) Fallopian tube, and (8) round ligament. On right side broad ligament is removed, so as to bring more clearly into view structures which it contains. 9. Fallopian tube. 10. Its fimbriated extremity. 11. One of its fimbriæ attached to ovary. 12. Ovary attached by its ligament to upper angle of uterus. 13. Round ligament.— (Wilson.)

At either side of the entrance of the vagina, beneath the integument covering its inferior part, as well as the superficial perineal fascia, and the constrictor vaginæ muscle, are situate the glands of Duverney. The space they occupy lies between the lower end of the vagina, the ascending ramus of the ischium, the crus clitoridis, and the erector clitoridis muscle. The excretory duct is at the anterior edge of the superior part of the gland, and runs beneath the constrictor vaginæ, horizontally forwards and inwards, to the inner face of the nympha, opening in front of the carunculæ myrtiformes, in the midst of a number of small mucous follicles. These glands secrete a thick, tenacious, grayish-white fluid, which is emitted in considerable quantity at the termination of sexual intercourse, and—it has been suggested—through the spasmodic contraction of the constrictor vaginæ muscle, under which they lie.


a Chitty, op. citat. p. 389, and Beck's Medical Jurisprudence, 6th edit., Philad. 1838. For an elaborate description of the Hymen and Caruncles, see Devilliers, Revue Médicale, Mai, 1840; and Encyclograph. des Sciences Médicales, Juin, 1840, p. 65. See, also, Virey, Gazette Médicale, No. xxiii., Paris, 1840.

Dictionnaire de Médecine, &c. art. Caroncule, Paris, 1821.
 Mr. Taylor, Dublin Journal of Med. Sciences, vol. xiii. 1838.

The uterus is a hollow organ, for the reception of the fœtus, and its retention during gestation. It is situate in the pelvis, between the bladder - which is before, and the rectum behind, and below the convolutions of the small intestines. Fig. 218 gives a lateral view of their relative situation, and Fig. 219, of their position, when regarded from before. It is of a conoidal shape, flattened on the anterior and posterior surfaces; rounded at the base, which is above, and truncated at its apex, which is beneath. It is of small size; its length being only about two and a half inches; its breadth one and a half inch at the base, and ten lines at the neck; its thickness about an inch. It is divided into the fundus, body, and cervix or neck. The fundus is the upper part of the organ, which is above the insertion of the Fallopian tubes. The body is the part between the insertion of the tubes and the neck; and the neck is the lowest and narrowest portion, which projects and opens into the vagina.

At each of the two superior angles are - the opening of the





Fallopian tube, the attachments of the ligament of the ovary, and that of the round ligament. The inferior angle is formed by the neck, which projects into the vagina to the distance of four or five lines, and terminates by a cleft, situate crosswise, called os tincæ, os uteri or vaginal orifice of the uterus. The aperture is bounded by two lips, which are smooth and rounded in those that have not had children; jagged and rugous in those who are mothers, - the anterior lip being somewhat thicker than the posterior. It is from three to five lines long, and is generally more or less open, especially in those who have had children.a The internal cavity of the uterus is very small in proportion to the bulk of the organ, owing to the thickness of the parietes, which

almost touch internally. It is divided into the cavity of the body, and that of the neck. (Fig. 221.) The former is triangular. The tubes open at its upper angles. The second cavity is more long than broad; is broader at the middle than at either end, and at the upper part, where it communicates with the cavity of the

See, on the Neck of the Uterus in the Young Female, Dr. Marc D'Espine, in Archiv. Général. de Médecine, Avril, 1836; and Dunglison's American Medical Intelligencer, i. 103, Philad. 1838.

body of the uterus, an opening exists, called the *internal orifice* of the uterus: the *external orifice* being the os uteri. The inner surface has several transverse rugæ, which are not very prominent. It is covered by very fine villi, and the orifices of several mucous follicles are visible.

The precise organization of the uterus has been a topic of interesting inquiry amongst anatomists. It is usually considered to be formed of two parts, a mucous membrane internally, and the proper tissue of the uterus, which constitutes the principal part of the substance. The mucous membrane has been esteemed a prolongation of that which lines the vagina. It is very thin; of a red hue in the cavity of the body of the organ; white in that of the neck. Chaussier, Ribes, and Madame Boivin, however, deny its existence. Chaussier asserts, that having macerated the uterus and a part of the vagina in water, in vinegar, and in alkaline solutions; and having subjected them to continued ebullition, he always observed the mucous membrane of the vagina stop at the edge of the os uteri; and Madame Boivin, - a well-known French authoress on obstetrics, who has attended carefully to the anatomy of those organs during pregnancy, - says, that the mucous membrane of the vagina terminates by small expansible folds, and by a kind of prepuce, under the anterior lip of the os uteri. In their view, the inner surface of the uterus is formed of the same tissue as the rest of it.a The proper tissue of the organ is dense, compact, not easily cut, and somewhat resembles cartilage in colour, resistance, and elasticity. It is a whitish, homogeneous substance, penetrated by numerous minute vessels. In the unimpregnated state, the fibres, which seem to enter into the composition of the tissue, appear ligamentous and pass in every direction, but so as to permit the uterus to be more readily lacerated from the circumference to the centre than in any other direction. The precise character of the tissue is a matter of contention amongst anatomists. To judge from the changes it experiences during gestation, and by its energetic contraction in delivery, it would seem to be decidedly muscular, or at least capable of assuming that character; but, on this point, we shall have occasion to dwell hereafter.

The uterus has, besides the usual organic constituents, — arteries, veins, lymphatics, and nerves. The arteries proceed from two sources; — from the spermatic, which are chiefly distributed to the fundus of the organ, and towards the part where the Fallopian tubes terminate; and from the hypogastric, which are sent especially to the body and neck. Their principal branches are readily seen under the peritoneum, which covers the organ; they are

<sup>\*</sup> Velpeau, Traité Elémentaire de l'Art. des Accouchemens, i. 77, Paris, 1829, 2d Amer. Edit. by Prof. Meigs, Philad. 1838; and Adelon, Physiologie de l'Homme, 2de édit. iv. 26.

VOL. II. - 29

very tortuous; frequently anastomose, and their ramifications are lost in the tissue of the viscus, and on its inner surface. The veins empty themselves partly into the spermatic, and partly into the



Nerves of the Uterus.

hypogastric. They are even more tortuous than the arteries; and, during pregnancy, they dilate and form what have been termed the uterine sinuses. The nerves are derived partly from the great sympathetic, and partly from the sacral pairs.


The uterus is sometimes absent.<sup>a</sup>

The appendages of the uterus are:—1. The ligamenta lata or broad ligaments, which are formed by the peritoneum. This membrane is reflected over the anterior and posterior surfaces and over the fundus of the uterus; and the lateral duplicatures of it form a broad expansion, and envelope the Fallopian tubes and

ovaria. These expansions are the broad ligaments. (See Fig. 220, and Fig. 219.) 2. The anterior and posterior ligaments, which are four in number and are formed by the peritoneum. Two of these pass from the uterus to the bladder, - the anterior; and two between the rectum and uterus, - the posterior. 3. The ligamenta rotunda or round ligaments, (Fig. 220,) which are about the size of a goosequill, arise from the superior angles of the fundus uteri, and, proceeding obliquely downwards and outwards, pass out through the abdominal rings to be lost in the cellular tissue of the groins. They are whitish, somewhat dense cords, formed by a collection of tortuous veins and lymphatics, of nerves, and of longitudinal fibres, which were, at one time, believed to be muscular, but are now generally considered to consist of condensed cellular tissue. 4. The Fallopian or uterine tubes; two conical, tortuous canals, four or five inches in length; situate in the same broad ligaments that contain the ovaries and extending from the superior angles of

<sup>\*</sup> For a number of such cases, see Dr. Chew, Amer. Journ. Med. Sciences, May, 1840, p. 39.

the uterus as far as the lateral parts of the brim of the pelvis. (Figs. 219, 220, and 223.) The uterine extremity of the tube (Figs. 221 and 223,) is extremely small, and opens into the uterus by an aperture so minute, as scarcely to admit a hog's bristle. The other extremity is called the pa-



Fallopian Tube.

vilion. It is trumpet-shaped, fringed, and commonly inclined towards the ovary, to which it is attached by one of its longest fimbria. This fringed portion is called corpus fimbriatum or morsus diaboli. The Fallopian tubes, consequently, open at one end into the cavity of the uterus, and at the other through the peritoneum into the cavity of the abdomen. They are covered externally by the broad ligament, or peritoneum; are lined internally by a mucous membrane, which is soft, villous, and has many longitudinal folds; and between these coats is a thick, dense, whitish membrane, which is possessed of contractility; although muscular fibres cannot be detected in it. Santorini asserts, that in robust females the middle membrane of the tubes has two muscular layers; an external, the fibres of which are longitudinal, and an internal, whose fibres are circular.

Recently, a memoir was read by M. Raciborskib to the Académie Royale des Sciences of Paris, in which he states it to be a general rule, that the extremities of the Fallopian tube in domestic animals are so placed during the act of fecundation as to envelope the entire ovary, either directly by means of the open trumpetshaped extremity, or indirectly by the aid of the fimbriated extremity. In women, however, the fimbriated extremity of the tube embraces but a small portion of the ovary; and M. Raciborski thinks, that this anatomical peculiarity is the cause of extra-uterine conception being so much more common in women than in do-

mestic animals. In the latter, indeed, it is very rare.

The ovaries, (Fig. 224,) are two ovoid bodies, of a pale red colour, rugous, and nearly of the size of the testes of the male. They are situate in the cavity of the pelvis, and are contained in the posterior fold of the broad ligaments of the uterus. At one time they were conceived to be glandular, and were called the female testes; but as soon as the notion prevailed

Fig. 224.

Section of Ovary.

<sup>\*</sup> Weber's Hildebrandt's Handbuch der Anatomie, Band. iv. s. 422, Braunschweig, 1832.

b Gazette Médicale de Paris, 25 Juin, 1842.

that they contained ova, the term ovary or egg-vessel was given to them. The external extremity of the ovary has attached to it one of the principal fimbriæ of the Fallopian tube. The inner extremity has a small fibro-vascular cord inserted into it: this passes to the uterus to which it is attached behind the insertion of the Fallopian tube, and a little lower. It is called the ligament of. the ovary, and is in the posterior ala of the broad ligament. It is solid, and has no canal. The surface of the ovary has many round prominences, and the peritoneum — forming the indusium - envelopes the whole of it, except at the part where the ovary adheres to the broad ligament. The precise nature of its parenchyma or stroma is not determined. When torn or divided longitudinally, as in Fig. 226, it appears to be constituted of a cellulo-vascular tissue. In this, there are from fifteen to twenty spherical vesicles — ovula Graafiana, folliculi Graafiania or ovicapsules. Roedererb asserts that he found in the ovary of one woman thirty, in that of another about fifty. These are filled with an albuminous fluid, which is colourless or yellowish, and may be readily seen by dividing the vesicles carefully with the point of fine scissors. The fluid from the ovary of a mare was examined by Lassaigne, and found to contain albumen, with chlorides of sodium and potassium.c

In the lower animals, the ovarium consists of a loose tissue, containing many cells, in which the ova are formed, and from which they escape by the rupture of the cell-walls; in the higher animals, as in the human female, the tissue of the ovarium is more compact, and the ova, except when they are approaching maturity, can only be distinguished by the aid of a high magni-

fying power.d

The microscopic analysis of the ovum has engaged the deep attention of modern histologists, who have greatly extended our knowledge in regard to it, although we have still much to learn. In man and the mammalia, the ova, even when advanced, are exceedingly minute, which is owing to the small quantity of vitellus that enters into their composition. The ripest ovum in the ovary of the human subject, and of the mammalia, generally does not measure more than from the fifteenth to the twentieth part of a line in diameter; it rarely happens, that they are as much as 1 th of a line. Hence, they are distinguished with difficulty by the naked eye. Under the microscope, the Graafian vesicle is found to consist of an external and internal membrane. The former - the tunic of the ovisue of Dr. Barry - is extremely vascular; the latter, the ovisac of the same observer - the membrana propria of some - is smooth and velvety, and derives its vessels from the former. The cavity, inclosed by these membranes, is far from

Dupuy's Journal de Médec. Veterin. for July, 1826, p. 336.

d Carpenter, Human Physiology, § 739, Lond. 1842.

<sup>Weber's Hildebrandt's Handbuch der Anatomie, B. iv. 458, Braunschweig, 1832.
Stannius, art. Eierstock, in Encyclop. Wörterb. x. 188, Berlin, 1836.</sup> 

being filled by the ovum; it contains, besides, a whitish or yellowish albuminous mass, which consists chiefly of granules, from the two hundredth to the three hundredth part of a line in diameter, connected together by a tenacious fluid, and forming the tunica granulosa of Dr. Barry.a Its density is unequal, and, towards some part of the periphery of the vesicle, these granules are accumulated in a disk-like form, making a slight prominence, in which there is a depression. The disk is termed by Von Baer discus proligerus; and it has been called discus vitellinus, and stratum proligerum. The prominence is called cumulus. Dr. Barry likewise observed certain granular cords, resembling both in appearance and function the chalazæ of the egg, which he has called retinacula, but which Bischoff does not admit. A small cup-like cavity in the cumulus receives the ovum. The ovum is surrounded

by a thick white ring, which has been called zona pellucida, and which has been considered to be a membrane, but, according to Mr. T. W. Jones, is now pretty generally acknowledged to be "the optical expression of the circumferential doubling of a thick transparent membrane, which incloses the volk." Within this, is a granular layer the vitellus, the larger granules of which are superficial and compact; whilst, internally, it is a clear albuminous fluid almost devoid of granules. Imbedded in the vitellus, but nearer its circumference than Ripe Ovum of Rabbit, taken from the Granfian its centre, is the germinal vesicle or vesicle of Purkinje, which appears like a clear ring, of very small size, and measures, in man spot.— (Wagner.) and the mammalia, not more than

Fig. 225.

Vesicle.

a, a. Proligenous disks, the edges of which

th part of a line in diameter. Upon a particular part of the germinal vesicle is observed the macula germinativa or germinal spot, which presents itself as a rounded granular formation attached to the inner wall of the germinal vesicle. All these parts are represented in the marginal figure. Wagner thinks the germinal vesicle may be viewed as a cell — a primary cell — of which the germinal spot forms the nucleus, and that it would perhaps be well to style the germinal spot the germinal nucleus.

The experiments of Carusc have shown, that the vesicles exist

a Philos. Transact. for 1838.

c Gazette Médicale de Paris, Aug. 12, 1837.

<sup>&</sup>lt;sup>b</sup> Human Physiology, translated by R. Willis, p. 43, Lond. 1841. See, also, the works of the various ovologists referred to hereafter, and especially an elaborate report on the ovum of man and the mammalia before and after fecundation, by Mr. T. W. Jones, in Brit. and For. Med. Rev., Oct. 1843, p. 513.

even in the fœtus. In these vesicles, we shall see hereafter, the ovum is contained. The arteries and veins of the ovaries belong to

Fig. 226.



Longitudinal Division of the

the spermatics. The arteries pass between the two layers of the broad ligament to the ovarium, assuming there a beautiful convoluted arrangement, very similar to the convoluted arteries of the testis. These vessels traverse the ovary nearly in parallel lines, as in the marginal figure, forming numerous minute twigs, which have an irregular knotty appearance, from their tortuous condition, and appear to be chiefly distributed to the Graafian vesicles.<sup>a</sup> The nerves of the ovaries, which

are extremely delicate, are from the renal plexuses; and their lym-

phatics communicate with those of the kidneys.

Such is the anatomy of the chief organs concerned in the function of generation. Those of lactation we shall describe hereafter.

#### 1. MENSTRUATION.

Before proceeding to the physiology of generation, there is one function, peculiar to the female, which will require consideration. This consists in a periodical discharge of blood from the vulva, occurring from three to six days in every month, during the whole time that the female is capable of conceiving, or from the period of puberty to what has been termed the critical age. This discharge is called the catamenia, menses, flowers, &c., and the process menstruction. It seems to be possessed by the human species alone. MM. Geoffroy St. Hilaire, and F. Cuvier, however, assert, that they have discovered indications of it in the females of certain animals; but it is usually denied that this is any thing more than the exudation of a bloody mucus. In some females, menstruation is established suddenly, and without any premonitory symptoms; but, in the greater number, it is preceded and accompanied by some inconvenience. The female complains of signs of plethora, or general excitement, - indicated by redness and heat of skin, heaviness in the head, oppression, quick pulse, and pains in the back or abdomen; whilst the discharge commences drop by drop, but continuously. During the first twenty-four hours, the flow is not as great as afterwards, and is more of a serous character; but, on the following day, it becomes more abundant and sanguineous. and gradually subsides, leaving, in many females, a whitish, mucous discharge, technically termed leucorrhea, and, in popular language, the whites.

E. Rigby, System of Midwifery, Amer. Edit. p. 23, Philad. 1841.
 Blumenbach, De Gener. Human. Variet. ed. 3, p. 51, Gotting. 1795.

e Dr. Allen Thomson, art. Generation, in Cyclop. of Anatomy and Physiology, part xiii. p. 441, Feb. 1838.

The quantity of fluid lost, during each menstruation, varies greatly, according to the individual and to the climate. Its average is supposed to be from six to eight ounces in temperate climes. By some, it has been estimated as high as twenty, but this is an exaggeration. The fluid proceeds from the interior of the uterus, and not from the vagina. At one time, it was believed, that in the intervals between the flow of the menses, the blood gradually . accumulates in some parts of the uterus, and when these parts attain a certain degree of fulness, they give way and the blood flows. This office was ascribed to the cells, - which were conceived to exist in the substance of the uterus between the uterine arteries and veins, - and, by some, to the veins themselves, which, owing to their great size, were presumed to be reservoirs, and hence were called uterine sinuses. The objection to these views is, that we have no evidence of the existence of any such accumulation; and that when the interior of the uterus of one who has died during menstruation is examined, there are no signs of any such rupture as that described: the enlarged vessels exist only during pregnancy or during the expanded state of the uterus; the veins, in the unimpregnated uterus, being extremely small, and totally inadequate for such a purpose.

The menstrual fluid is a true exhalation, effected from the inner surface of the uterus. This is evident from the change in the lining membrane of the organ during the period of its flow. It is rendered softer and more villous, and exhibits bloody spots, with numerous pores from which the fluid may be expressed. An injection, sent into the arteries of the uterus, also readily transudes through the lining membrane. The appearance of the menstrual fluid in the cavity of the uterus, during the period of its flow; its suppression in various morbid conditions of the organ; and the direct evidence, furnished to Ruysch, Blundell, Sir C. Clarke, and others, in cases of prolapsus or of inversio uteri, where the fluid has been seen distilling from the uterus, likewise show that it is a uterine ex-

halation.

It has been a question, whether the fluid proceeds from the arteries or veins; and this has arisen from the circumstance of its being regarded as mere blood; but it is in appearance not much like blood; and it may be distinguished from it by the smell, which is sui generis, and also by not being coagulable. "It is," says Mr. Hunter, "neither similar to blood taken from a vein of the same person, nor to that which is extravasated by accident in any other part of the body, but is a species of blood, changed, separated, or thrown off from the common mass by an action of the vessels of the uterus, similar to that of secretion, by which action the blood loses the principle of coagulation and, I suppose, life." The principle of coagulation does not exist, owing, accord-

a Principles and Practice of Obstetricy, Amer. Edit. p. 49, Washington, 1834.
 b On Diseases of Females, attended with Discharges, Amer. Edit. Philad. 1824; see, also, Dr. D. Davis, in Principles, &c., of Obstetric Medicine, i. 256, Lond. 1836.

ing to Lavagna, Toulmouche, and J. Müller, to absence of fibrin, Retziuse asserts, that he has detected in it free phosphoric and lactic acids, by the presence of which, he conceives, the fibrin is kept in a state of solution and prevented from coagulating. The fluid has the properties, according to Brande, of a very concentrated solution of the colouring matter of the blood in a dilute serum.d Dr. - Burowe examined twelve ounces of menstrual blood, which had been retained in the uterus by an imperforate hymen. The fluid was of a dirty reddish brown colour, of the consistence of syrup, very adhesive, and entirely devoid of odour. It abounded in albumen, and was very little susceptible of putrefaction. When examined with the microscope, almost all the blood-globules were found to have lost their regular form, and to resemble the granules, observed in pus which has been for a long time exposed to the air, or retained within the cavity of an abscess. These blood-globules were suspended in a transparent fluid. On stirring the blood for a considerable time, no perceptible change was produced to the eye, but under the microscope numerous delicate transparent lamellæ were seen floating in the serum, which Dr. Burow regarded as portions of fibrin, a substance sparingly present — as has been remarked - in menstrual blood. The red colour of the menstrual fluid was found by Remakf to be owing to the presence of blood-globules, and the intensity of the colour to their number.

M. Bouchardats has recently undertaken a new analysis of the menstrual fluid, obtained from a female who permitted a speculum to remain in the vagina for ten hours in order that an ounce might be procured. Without this precaution, the fluid becomes mixed with vaginal mucus, and urine, as the presence of ammoniaco-magnesian phosphate demonstrates. The following were the results of the analysis: - Water, 90.8; fixed matters, 6.92. The fixed matters were composed of -fibrin, albumen, and colouring matter, 75.27; extractive matter, 0.42; fatty matter, 2.21; salts, 5.31; mucus, 16.79. The female was a patient of M. Brière de Boismont, who considers, that the large proportion of water was due to the delicacy of her frame, and to her subsisting on vegetable diet (?). Another specimen of menstrual fluid, examined by M. Donné, presented the following appearances under the microscope. 1. Abundance of the ordinary corpuscles of the blood. 2. Vaginal mucus, formed of epidermic scales from the mucous membrane. 3. Mucous globules, furnished by the neck

8 Brière de Boismont, De la Menstruation, &c., Paris, 1842; and Provincial Med. Journal, July 30, 1842.

a Brugnatelli, Giornale di Fisica, &c. p. 397, 1817; Desorméaux, in Dict. de Médec. xvi. 181.

Handbuch der Physiologie, Baly's translation, p. 256, Lond. 1837, and p. 1481,

c Ars Berättelse af Setterblad, 1835, Seite 19 - quoted in Zeitschrift für die Gesammte Medicin. Marz, 1837, S. 390.

d Philos. Transact. cii. 113; and Blundell, op. cit. p. 46.

Müller's Archiv. No. vii. 1840; and Brit. and For. Med. Rev., July, 1840, p. 287. Medicinische Zeitung, Dec. 25, 1839. See, also, Mr. Ancell, Lectures on the Physiology and Pathology of the Blood, April 25, 1840, p. 149.

of the uterus (?). So far, therefore, as these examinations go, they would show, that there is a resemblance between the catamenial discharge and arterial blood. The fact of injections, sent into the arteries, transuding through the inner lining of the uterus is in favour of the exhalation taking place from the arteries, and the analogy of all the other exhalations is confirmatory of the position. Still there are many eminent physiologists and obstetricians who

regard the discharge as a monthly hemorrhage.

The efficient cause of menstruation has afforded ample scope for speculation and hypothesis.\* As its recurrence corresponds to a revolution of the moon around the earth, lunar influence has been invoked; but, before this solution can be admitted, it must be shown, that the effect of lunar attraction is different in the various relative positions of the moon and earth. There is no day in the month, in which numerous females do not commence their menstrual flux, and, whilst the discharge is beginning with some, it is at its acme or decline with others. The hypothesis of lunar influence must therefore be rejected. In the time of Van Helmont, it was believed, that a ferment exists in the uterus, which gives occasion to a periodical intestine motion in the vessels, and a recurrence of the discharge; but independently of the want of evidence of the existence of such a ferment, the difficulty remains of accounting for its regular renovation every month. Local and general plethora have been assigned as causes, and many of the circumstances, that modify the flow, favour the opinion. The fact of, what has been called, vicarious menstruation has been urged in support of this view. In these cases, instead of the menstrual flux taking place from the uterus, hemorrhages occur from various other parts of the body, as the breast, lungs, ears, eyes, nose, &c. It does not seem, however, that in any of these cases, the term menstruation is appropriate; inasmuch as the fluid is not menstrual, but consists of blood periodically extravasated. Still, they would appear to indicate, that there is a necessity for the monthly evacuation or purgation, Reinigung, as the French and Germans term it; and that if this be obstructed, a vicarious hemorrhage may be established; yet the loss of several times the quantity of blood from the arm, previous to, or in the very act of, menstruation does not always prevent, or interrupt the flow of the catamenia; and in those maladies, which are caused by their obstruction, greater relief is afforded by the flow of a few drops from the uterus itself, than by ten times the quantity from any other part. Some of the believers in local plethora of the uterus have maintained, that the arteries of the pelvis are more relaxed in the female than in the male, and the veins more unvielding; hence, that the first of these vessels convey more blood than the second return. It has been also affirmed, that whilst the arteries of the head predominate in man, by reason of his being more dis-

Dr. D. Davis, op. cit. i. 259.

b Opera, edit. 4, p. 440, Lugd. Bat. 1667; and Haller, Elementa Physiologiæ, vii. 1.

posed for intellectual meditation, the pelvic and uterine arteries predominate in the female, owing to her destination being more especially for reproduction. Setting aside all these gratuitous assumptions, it is obvious, that a state, if not of plethora, at least of irritation, must occur in the uterus every month, which gives occasion to the menstrual secretion; but, as Adelona has properly remarked, it is not possible to say, why this irritation is renewed monthly, any more than to explain, why the predominance of one organ succeeds that of another in the succession of the ages. The function is as natural, as instinctive to the female, as the development of the whole sexual system at the period of puberty. That it is connected most materially with the capability of reproduction is shown by the fact, that it does not make its appearance until puberty, the period at which the young female is capable of conceiving, and that it disappears at the critical time of life, when conception is impracticable. It is arrested, too, as a general rule, during pregnancy and lactation; and in amenorrhœa or obstruction of the menses, fecundation is not readily effected. In that variety, indeed, of menstruation, which is accomplished with much pain at every period, and is accompanied by the secretion of a membranous substance having the shape of the uterine cavity, conception may be esteemed impracticable. Professor Hamilton, of the University of Edinburgh, was in the habit of adducing this, in his lectures, as one of two circumstances - the other being the want of a uterus - that are invincible obstacles to fecundation. Yet, in the case of dysmenorrhæa, of the kind mentioned, if the female can be made to pass one monthly period without suffering, or without the morbid secretion from the uterine cavity, she may become pregnant, and the whole of the evil be removed: for, the effect of pregnancy being to arrest the catamenia, the morbid habit is usually got rid of during gestation and lactation, and may not subsequently recur.

Gall<sup>b</sup> strangely supposed, that some general, but extraneous cause of menstruation exists,—not the influence of the moon; and he affirms, that, in all countries, females generally menstruate about the same time; that there are, consequently, periods of the month in which none are in that condition; and he affirms, that all females may, in this respect, be divided into two classes:—the one comprising those who menstruate in the first eight days of the month, and the other, those who are "unwell"—as it is termed by them, in some countries—in the last fortnight. He does not, however, attempt to divine what this cause is. We are satisfied, that his positions are erroneous. Attention to the matter has led us to the knowledge, already expressed, that there is no period of the moon, at which the catamenial discharge is not taking place in some; and we have not the slightest reason for supposing, that, on the average, more females are menstruating


<sup>Physiologie de l'Homme, 2de édit. iv. 48, Paris, 1829.
Sur les Fonctions du Cerveau, iv. p. 355.</sup> 

at any one part of the month than at another. It would seem, however, that there are circumstances in the economy, which, as in the case of fevers, give occasion to something like periodicity at intervals of seven days; — for example, Mr. Roberton, of Manchester, England, asserts, that of 100 women, the catamenia returned every fourth week in 68; every third week in 28; every second week in 1; and at irregular intervals in 10; these varieties usually existing as family and constitutional peculiarities.

It is scarcely necessary to notice the visionary speculations of those who have regarded menstruation as a mechanical consequence of the erect attitude; or the opinion of Roussel, that it originally did not exist, but that being produced artificially by too succulent and nutritious a regimen, it was afterwards propagated from generation to generation; or, finally, that of Aubert, who maintained, that if the first amorous inclinations were satisfied, the resulting pregnancy would totally prevent the establishment of menstruation. The function, it need scarcely be repeated, is instinctive, and forms an essential part of the female constitution.

Of late, M. Gendrin<sup>d</sup> has revived a view entertained by Mr. Cruikshank, Dr. Power, and others, that menstruation is dependent upon changes occurring periodically in the ovary. Many cases have been observed by Cruikshank, Robt. Lee, Gendrin and others, in which, on the dissection of females who have died during menstruation, evidences have been afforded of the rupture of an

ovarian vesicle. and of a small irregular rupture or cicatrix in the coats of the ovarium, as represented in the marginal figure, which communicated with the remains of a Graafvesicle :h whence it has inferred, that during the whole of that period of



Ovary of a Female dying during Menstruation.

- See art. Generation, by Dr. Allen Thomson, in Cycl. Anat. and Physiol.. part xiii.
   p. 440, Feb. 1838.
   Edinb. Med. and Surg. Journal, xxxviii. 237.
  - Système Physique, &c., de la Femme, p. 13, Paris, 1809.
     Traite Philosophique de Médecine Pratique, Paris, 1838-9.
  - See Brit. and For. Med. Review, Oct. 1840, p. 592.
     An Essay on the Periodical Discharge, &c., Lond. 1832.
- s M. Negrier, cited in Dunglison's Amer. Med. Intelligencer, July 15, 1840, p. 121. See, also, Dr. Robt. Lee, art. Ovaria, Cyclopædia of Pract. Medicine; Dr. Laycock, on the Nervous Diseases of Women, p. 42, Lond. 1840; and Dr. R. Willis, in note to Wagner's Elements of Physiology, p. 69, Lond. 1841.
- b Churchill, Theory and Practice of Midwifery, Amer. Edit. by Prof. R. M. Huston, p. 71, Philad. 1843.

when the capability of conception continues, there is a constantly successive development of vesicles and their contained ovules in the ovary, and that, at each epoch of menstruation, a vesicle having reached the surface of the ovary becomes the foyer of a peculiar organic action, in which all the organs of generation participate; and that the result of this action is the rupture of the vesicle, and the loss of the infecund ovum, either by expulsion from the uterus or by destruction in the ovary. Still more recently, M. Raciborski, a in a letter to the Academié Royale de Medecine of Paris, has maintained, as the result of his researches on menstruation-First. That there exists the most intimate connection between the Graafian vesicles and menstruation. When these vesicles arrive at their full development, menstruation commences, and when they are destroyed, it ceases. Secondly. At each menstrual period, a follicle projects like a nipple on the surface of the ovary, where it afterwards bursts, without requiring for that purpose any venereal excitement. Thirdly. The rupture of the follicles in general appears to take place at the period when the menstrual discharge is stopping; and Fourthly. The ovaries do not act alternately, as has been affirmed: in this respect not seeming to be under any fixed law. It must be admitted, then, that there certainly would seem to be a close and intimate relation between the number of cicatrices on the surface of the ovaries, and the number of times a female has menstruated.b

The age, at which menstruation commences, varies in individuals and in different climates. It has been esteemed a general law, that the warmer the climate, the earlier the discharge takes place, and the sooner it ceases; but there is reason for doubting the correctness of this prevalent belief.c With us, the most common period of its commencement is from thirteen to seventeen years. Mahomet is said to have consummated his marriage with one of his wives, "when she was full eight years old."d Of 450 cases, observed at the Manchester Lying-in Hospital, in England, emenstruation commenced in the eleventh year in 10; in the twelfth in 19; in the thirteenth in 53; in the fourteenth in 85; in the fifteenth in 97; in the sixteenth in 76; in the seventeenth in 57; in the eighteenth in 26; in the nineteenth in 23; and in the twentieth in 4. Menstruction commonly ceases in the temperate zone at from forty to fifty years. These estimates are, however, liable to many exceptions. In rare cases, the catamenia have appeared at a very early age, even in childhood; and again, the menses, with powers of fecundity, have continued, in particular instances, beyond the ages that have been specified: some of these protracted cases have had regular catamenia; in others, the discharge, after a long suppression, has returned. Of 77 individuals, they ceased in 1 at the age

Bullet. de l'Académ. Royal. de Méd. Jan. 1843; and Edinb. Med. and Surg. Journ., April, 1843.

See, on this subject, Mr. Girdwood, London Lancet, March 4, 1843, p. 825.

<sup>Dr. A. Thomson, op. citat. p. 442.
Prideaux's Life of Mahomet, p. 30, Lond. 1718.</sup> 

e Roberton, op. citat.

of 35; in 4 at 40; in 1 at 42; in 1 at 43; in 3 at 44; in 4 at 45; in 3 at 47; in 10 at 48; in 7 at 49; in 26 at 50; in 2 at 51; in 7 at 52; in 2 at 53; in 2 at 54; in 1 at 57; in 2 at 60; and in 1 at 70. Of the 10,000 pregnant females, registered at the Manchester Hospital, 436 were upwards of 40 years of age; 397 from 40 to 45; 13 in their 47th year; 8 in their 48th; 6 in their 49th; 9 in their 50th; 1 in her 52d; 1 in her 53d; and 1 in her 54th. Mr. Roberton asserts, that as far as he could ascertain, — and especially in the three cases above 50 years, — the catamenia continued up to the period of conception.<sup>a</sup>

The following table has recently been published by M. Brière de Boismont. It is founded on the results of 2352 cases. It will be seen from it, that by far the greatest number of women begin

to menstruate during the 14th or 15th year.

Age.	Paris, 1200 cases by Meniers.	Paris, 85 cases by Marc D'Es- pine.	Lyons, 432 cases by Pétrequin,	Marseilles, 68 cases by Marc D'Es- pine.	Manchester, 450 cases by Roberton.	Göttingen, 137 cases by Osiander.
5	1	0	0	0	0	0
7	1	0	0	0	0	0
8	2	0	0	0	0	0
9	10	1	0	0	0	0
10	29	0	5	0	0	0
11	93	3	14	6	10	0
12	105	14	26	10	19	3
13	132	6	47	13	53	8
14	194	18	50	9	85	21
15	190	14	70	16	97	32
16	141	7	79	8	76	24
17	127	6	58	4	57	11
18	90	5	38	2	26	18
19	35	8	21	0	23	10
20	30	3	9	0	4	8
21	8	. 0	5	0	0	1
22	8	0	1	0	0	0
23	4	0	0	0	0	1
24	0	0	3	0	0	00

Mr. Roberton<sup>d</sup> has attempted, and successfully, to show, that the age of puberty is as early in the cold, as in the tropical regions of the earth; and, that were marriages to take place in England, at as juvenile an age as they do in Hindusthan, instances of very early fecundity would be as common in England as they are in that country. He is of opinion, that early marriage and early intercourse between the sexes, where found prevailing generally, "are to be attributed, not to any peculiar precocity, but to moral and political degradation, exhibited in ill laws and customs, the enslavement more or less of the women, ignorance of letters, and

<sup>&</sup>lt;sup>2</sup> See Montgomery, on the Signs and Symptoms of Pregnancy, p. 160, Lond. 1837.

b De la Menstruation, &c., Paris, 1842.

c For a similar table, formed from 5062 cases, see Dr. W. A. Guy, Principles of Forensic Medicine, part i. p. 79, Lond. 1843.

d Edinb. Med. and Surg. Journal, Oct. 1832, and July, 1842; and Lond. Med. Gaz., July 21, 1843.

VOL. II. - 30

impure or debasing systems of religion." He has also shown, from statistical evidence, that menstruation does not occur more

early in the negress than in the white female.

In the statement sent to Parliament by Bartholomew Mosse, when endeavouring to procure a grant for the Dublin Lying-in Hospital, he mentions, that 84 of the women delivered under his care were between the ages of 41 and 54; 4 of these were in their 51st year, and 1 in her 54th. A relation of Haller had two sons after her fiftieth year; and children are said to have been born even after the mother had attained the age of sixty. Holdefreund relates the case of a female, in whom menstruation continued till the age of seventy-one; Bourgeois till the age of eighty; and Hagendorn till ninety; however, it is probable that these were not cases of true menstruation, but perhaps of irregularly periodical dis-

charges of true blood from the uterus or vagina.c

As a general rule, the appearance of the menses denotes the capability of being impregnated, and their cessation the loss of such capability. Yet, females have become mothers without ever having menstruated.d Fodérée attended a woman, who had menstruated but once - in her 17th year, - although 35 years of age, healthy, and the mother of five children. Morgagni instances a mother and daughter, both of whom were mothers before they menstruated. Sir E. Homef mentions the case of a young woman, who was married before she was seventeen, and having never menstruated, became pregnant; four months after her delivery, she became pregnant again; and four months after the second delivery, she was a third time pregnant, but miscarried: after this she menstruated for the first time, and continued to do so for several periods, when she again became pregnant; and Mr. Harrison, at a meeting of the Westminster Medical Society remarked. that he knew an instance in which the mother of a large family had never menstruated; - yet Dr. Deweesh and Dr. Campbelli assert, that there is not a properly attested instance on record, of a female conceiving, previous to the establishment of the catamenia: the latter gentleman admits, however, that when an individual has once been impregnated, she may conceive again, several times in succession, without any recurrence of the catamenia between these different conceptions, - because he has known a case of this kind, but not of the other!

During the existence of menstruation, the system of the female

Philosoph, Transact. cvii. 258; and Lect. on Comp. Anat. iii. 298.

5 Lond. Lancet, Jan. 19, 1839, p. 619.

<sup>\*</sup> See a case of this kind in Transylvania Journal, ix. 185, Lexington, 1836.

b See a case by Dr. Strassberger, of recurrence of menstruation at 80, after it had ceased at 42 years of age, in Medicin. Zeitung, s. 248, Nov. 30, 1836.

c Dr. D. Davis, Principles, &c. of Obstetric Medicine, i. 239, London, 1836; and C. W. Mehliss, Ueber Virilescenz und Rejuvenescenz Thierischer Körper, s. 75, Leipz. 1838. d Churchill, op. cit. e Médecine Légale, i. 393, Paris, 1813.

Compend. System of Midwifery, 8th edit. Philad. 1836.
 Introduction to the Study of Midwifery, Edinb. 1833.

den and irregular checks of transpiration should be avoided, as well as every kind of mental and corporeal agitation, otherwise the process may be impeded, or hysterical and other troublesome affections be excited. The sacred volume exhibits the feeling entertained towards the female, whilst performing this natural function. Not only was she regarded "unclean" in antiquity; she was looked upon, as Dr. Elliotson has remarked, as mysteriously deleterious. In the time of Pliny, a female, during menstruation, was considered to blight corn, destroy grafts and hives of bees, dry up fields of corn, cause iron and copper to rust and smell, drive dogs mad, &c., &c.; and it is firmly believed by many, in England, that meat will not take salt if the process be conducted by a female so circumstanced.

The temperature of the vagina does not appear to be affected

by menstruation or pregnancy.c

## c. Sexual Ambiguity.

The sexual characteristics, in the human species, are widely separate; and the two perfect sexes are never, perhaps, united in the same individual. Yet such an unnatural union has been supposed to exist; from the fabulous son of 'Equals and Aspedican, — Mercury and Venus, — to his less dignified representative of modern times:—

"Nec fæmina dici, Nec puer ut possent, neutrumque et utrumque videntur." — Ovid.4

We have already remarked, that in the lower animals, and in plants, such hermaphrodism is common; but, in the upper classes and especially in man, a formation, which gives to an individual the attributes of both sexes has never been witnessed. Monstrous formations are occasionally met with; but if careful examination be made, it can usually be determined to what sex they rather belong. Cases, however, occur, in which it is extremely difficult to decide, although we may readily pronounce that the being is totally incapable of the function of reproduction. The generality of cases are produced by unusual development of the clitoris in the female, or by a cleft scrotum in the male. Only two instances of this kind have fallen under the observation of the author, both of which were females, as they usually are. One of these has been described by the late Professor Béclard, of Paris, whose details we borrow.

\* Elliotson's Blumenbach, p. 465, Lond. 1828. b Histor. Natural. xxvii.

· Fricke, Zeitschrift für die Gesammte Medicin, Nov. 1838.

<sup>4</sup> "Both bodies in a single body mix,

A single body with a double sex." — Addison.

Art. Hermaphrodite, in Dict. de Médecine, tom. xi.

o Dr. Willis, in Wagner's Elements of Physiology, part i. p. 55, Philad. 1841. See a case of supposed hermaphrodism, by Robt. Perry, with the dissection, by Sir Astley Cooper, in Guy's Hospital Reports, for April and October, 1840.

Marie-Madeleine Lefort, aged sixteen years, seemed to belong to the male sex, if attention were paid merely to the proportions of the trunk, limbs, shoulders, and pelvis; to the conformation and dimensions of the pelvis; to the size of the larynx; the tone of the voice, the development of the hair; and to the form of the urethra, which extended beyond the symphysis pubis. An attentive examination, however, of the genital organs showed, that she was of the female sex. The mons veneris was round and covered with hair. Below the symphysis pubis was a clitoris, resembling the penis in shape, twenty-seven millimètres, or about an inch long, in the state of flaccidity; and susceptible of slight elongation during erection; having an imperforate glans, hollowed beneath by a duct or channel, at the inferior part of which were five small holes, situate regularly on the median line. Beneath and behind the clitoris, a vulva existed, with two narrow, short and thin labia, furnished with hair, devoid of any thing like testicles, and extending to within ten lines of the anus. Between the labia was a very superficial cleft, pressure upon which communicated a vague sensation of a void space in front of the anus. At the root of the clitoris was a round aperture, through which a catheter could not be passed into the bladder. It could be readily directed, howver, towards the anus, in a direction parallel to the perineum. When the catheter was passed a little backwards and upwards to the depth of eight or ten centimètres, it was arrested by a sensible obstacle, but no urine flowed through it. It seemed to be in the vagina. At the part where the vagina stopped, a substance could be distinguished through the parietes of the rectum, which appeared to be the body of the uterus. Nowhere could testicles be discovered. She had menstruated from the age of eight years: the blood issuing in a half coagulated state through the aperture at the root of the clitoris. She experienced, too, manifest inclination for commerce with the male, and a slight operation only would probably have been necessary to divide the apron, closing the vulva from the clitoris to the posterior commissure of the labia. The urethra extended in this case for some distance beneath the clitoris, as in the penis. From all the circumstances, M. Béclard concluded that the person, subjected to the examination of the Société de Médecine of Paris, was a female; and that she possessed several of the essential organs of the female; - the uterus and vagina whilst she had only the secondary characters of the male; - as the proportions of the trunk and limbs; that of the shoulders and pelvis; the conformation and dimensions of the pelvis; the size of the larynx; the tone of the voice; the development of the hair; the urethra extending beyond the symphysis pubis, &c.

In the year 1818, an individual was exhibited in London, who had a singular union of the apparent characteristics of the two sexes. The countenance resembled that of the male, and she had a beard, but it was scanty. The shape, however, of the body and limbs was that of the female. The students of the Anatomical

Theatre of Great Blenheim Street, London, of whom the author was one, offered her a certain sum, provided she would permit the sexual organs to be inspected by the veteran head of the school — Mr. Brookes: to this she consented. She was, accordingly, exposed before the class; and her most striking peculiarities exhibited. The clitoris was large, but not perforate. Mr. Brookes, desirous of trying the experimentum crucis, passed a catheter into the vagina, and attempted to introduce another into the urethra; but fearing discovery, and finding that the mystery of her condition was on the point of being unveiled, she started up and defeated the experiment. No doubt existed in the mind of Mr. Brookes, that there were two distinct canals, — one forming the vagina; the other the urethra, — and that she was consequently female.

One of the most complete cases of admixture of the sexes was laid by Rudolphi before the Academy of Sciences of Berlin, at the sitting of Oct. 22, 1825." It was met with in the body of a child, which died, it was said, seven days after birth, but the development of parts led to the supposition, that it was three months old. The penis was divided inferiorly; the right side of the scrotum contained a testicle; the left side was small and empty. There was a uterus, which communicated at its superior and left portion with a Fallopian tube, behind which was an ovary destitute of its ligament. On the right side, there was neither Fallopian tube, nor ovary, nor ligament, but a true testicle, from the epididymis of which arose a vas deferens. Below the uterus was a hard, flattened ovoid body, which, when divided, exhibited a cavity with thick parietes. The uterus terminated above in the parietes of this body, but without penetrating its cavity. At its inferior part was a true vagina, which terminated in a cul-de-sac. The urethra opened into the bladder, which was perfect; and the anus, rectum and other organs were naturally formed. Rudolphi considered the ovoid body, situate beneath the uterus, to be the prostate, and vesiculæ seminales in a rudimental state.

A curious case of doubtful sex was seen by the author's friend—Dr. Pue, of Baltimore. It occurred in a negro, who had the appearance of a woman, and passed for such. On examination, at the request of her owner, Dr. Pue found no signs of female organs of generation. There was a penis of some development—the size of a boy's, 12 or 14 years old. The glans was well formed, but the urethra did not terminate at the extremity of the organ. It opened at the frænum. There was some appearance of scrotum, but no testes were perceptible. Menstruation took place regularly, but with great pain, the fluid passing through the urethra. During this period, the breasts—which were natural, and the size of a girl's of 14—became tender and swoller, and she had at this time only a desire for copulation. The pubes was well covered with hair. Dr. Pue's impression was, that her inclinations were for the

<sup>\*</sup> Amer. Journ. of the Med. Sciences, p. 499, Feb. 1832.

male. A very interesting case has likewise been described by Professor Mayera of Bonn. The mixed attributes of male and female were well marked. On the one hand, there was a testicle slightly atrophied, a penis and a prostate gland; on the other, a vagina and uterus, with its Fallopian tubes, and on the left side a

body analogous to an ovary.

Cases like the above, — and we have such on the authority of Baillie, Verdier, Giraud, Ackermann, Handy, Sir E. Home, Pinel, Maret, Sue, Bouillaud, and others, — have led to the belief, that hermaphrodism is possible. This is the opinion of Tiedemann, Meckel, and many other physiologists, but it does not rest on any observed examples. The varieties of these sexual vagaries are extremely numerous; and form occasionally the subject of medico-

legal inquiry.

Instances of animals being brought forth, whose organs of generation are preternaturally formed, sometimes occur, and they also have been commonly called hermaphrodites; but such examples have been rarely investigated. Monstrous productions, having a mixture of the male and female organs, seem to occur most frequently in neat cattle, and have been called free-martins. When a cow brings forth twin calves, one a male and the other apparently a female, the former always grows up to be a perfect bull, but the latter appears destitute of all sexual functions, and never propagates. This is the free-martin. It was the opinion of Mr. Hunter, that the free-martin never exhibits sexual propensities; but this has been controverted by Mr. Allnatt. A clergyman of great respectability informed that gentleman, that he had bred a free-martin upon his estate, which had not only shown the natural desire for the male, but had actually admitted him - of course, ineffectually. Mr. Allnatt farther remarks, that he had seen, the day before, working in harness, a true, apparently female, adult free-martin, which occasionally manifested its male propensities in an intelligible manner. When the cows amongst which this freemartin is kept, exhibit their inclination for the male, the creature unlike the spayed heifer or common ox - is peculiarly on the alert, and has been observed to leap the cows like the entire male. A gentleman of Mr. Allnatt's neighbourhood had a true free-martin, which had received the bull several times, but had never propagated. After death the animal was examined. Scarcely a vestige of uterus could be discovered. From Mr. Hunter's observationsd it would seem, that in all the instances of free-martins, which he examined, no one had the complete organs of the male and

a Gazette Médicale de Paris, Sept. 24, 1836; and Philadelphia Med. Examiner, April 10, 1841, p. 232.

b Dr. Simpson, art. Hermaphroditism, Cyclop. of Anat. and Physiol. vol. ii.; Marc, art. Hermaphrodisme, Dict. de Medecine, 2de édit. xv. 241, Paris, 1837; Beck, Med. Jurisprud. 6th edit. vol. i. Philad. 1838; Mr. Pilcher, Lancet, March 17, 1838, p. 915; and W. A. Guy, Principles of Forensic Medicine, part i. p. 36, Lond. 1843.

c London Medical Gazette for July 2d, 1836.

<sup>4</sup> Animal (Economy, edit. by Mr. Owen, Lond. 1838.

female, but partly the one and partly the other; and, in all, the

ovaria and testicles were too imperfect to perform their functions. In noticing this phenomenon, Sir Everard Homea remarks, that it may account for twins being most commonly of the same sex; "and when they are of different sexes," he adds, "it leads to inquire whether the female, when grown up, has not less of the true female character than other women, and is incapable of having children." "It is curious," says Sir Everard, "and in some measure to the purpose, that, in some countries, nurses and midwives have a prejudice, that such twins seldom breed." The remark of Sir Everard is signally unfortunate, and ought not to have been hastily hazarded, seeing that a slight examination would have exhibited, that there is no analogy between the free-martin and the females in question; and, more especially, as the suggestion accords with a popular prejudice, highly injurious to the prospects and painful to the feelings of all who are thus circumstanced. In the London Medical Repository, Mr. Cribb, of Cambridge, England, has properly observed, that the external characters and anatomical conformation of the free-martin are totally unlike those of the human female. In external appearance, the free-martin differs considerably from the perfectly formed cow; - the head and neck, in particular, bearing a striking resemblance to those of the bull. Mr. Cribb has, however, brought forward the most decisive evidence in favour of the fallacy of the popular prejudice, by the history of seven cases, which are of themselves sufficient to put the matter for ever at rest. Of these seven cases, which are all that he had ever known, of women, born under the circumstances in question, having been married, - six had children. The case of an orang-outang, described by Dr. Harlan, affords

The case of an orang-outang, described by Dr. Harlan, affords the nearest approach to a complete union of the sexes in the same individual, which has been recorded. It possessed ovaries, Fallopian tubes, uterus and vagina, and also testes, epididymis, vasa

deferentia, and a highly erectile penis.

### 1. PHYSIOLOGY OF GENERATION.

In man and the superior animals, in which each sex is possessed by a distinct individual, it is necessary, that there should be a union of the sexes, and that the fecundating fluid of the male should be conveyed within the appropriate organs of the female, in order that, from the concourse of the matters furnished by both sexes, a new individual may result. To this union we are incited by an imperious instinct, established within us for the preservation of the species, as the senses of hunger and thirst are placed within us for the preservation of the individual. This has been termed the desire or instinct of reproduction; and for wise purposes, its gratification is attended with the most pleasurable feelings, which man

Medical and Physical Researches, p. 22, Philad. 1835.

Philosoph. Transactions for 1799; and Lectures on Comparative Anatomy, iii.
 Sept. 1823, and ibid. 1827.

or animals can experience. Prior to the period of puberty, or whilst the individual is incapable of procreation, this desire does not exist; but it suddenly makes its appearance at puberty, persists vehemently during youth and the adult age, and disappears in advanced life, when procreation becomes again impracticable. It is strikingly exhibited in those animals, in which generation can only be effected at particular periods of the year, or whilst they are in heat; — as in the deer, during the rutting season.

The views, which have been entertained, regarding the seat of this instinct - whether in the encephalon or genital organs - were considered under the head of the mental and moral manifestations. It was there stated, that Cabanis and Broussais make the internal impressions to proceed from the genital organs, but to form a part of the psychology of the individual; and that Gall assigns an encephalic organ - the cerebellum - for its production, and ranks the instinct of reproduction amongst the primary faculties of the mind. In farther proof of the idea, which refers it to the encephalon, it may be remarked, that the instinct has been observed in those, who, owing to original malformation, have wanted the principal part of the genital organs, whilst it has continued in the case of eunuchs not castrated till after the age of puberty. In opposition to this view, it has been urged, that simple titillation of the organs will excite the desire. This, however, may be entirely dependent upon association, in which the brain is largely concerned. In many cases, the desire is produced through the agency of vision; when the brain must necessarily be first excited, and, through its influence the generative apparatus. The cause of the desire has, by some, been ascribed to the presence of sperm, in the requisite quantity, in the vesiculæ seminales; but in answer to this, it is urged, that eunuchs, under the circumstances above mention, and females, in whom there is no spermatic secretion, have the desire. The fact is, we have no more precise knowledge of the nature of this instinct, than we have of any of the internal sensations or moral faculties. We know, however, that it exhibits itself in various degrees of intensity, and occasionally assumes an opposite character — constituting anaphrodisia.

a. Copulation. — In the union of the sexes, the part performed by the male is the introduction of the penis, — the organ for the conveyance of the sperm to the uterus, — and the excretion of that fluid, during its introduction. In the flaccid state of the organ this penetration is impracticable; it is first of all necessary, that under the excitement of the venereal desire, the organ should attain a necessary state of rigidity, which is termed erection. In this state the organ becomes enlarged, and raised towards the abdomen; its arteries beat forcibly; the veins become tumid; the skin more coloured, and the heat augmented. It becomes also of a triangular shape, and these changes are indicated by an indescribable feeling of pleasure. Erection is not dependent upon volition. At times, it manifests itself against the will; at others

it refuses to obey it; yet it requires, apparently, the constant excitement of the encephalic organ concerned in its production; the slightest distraction of the mind causing its cessation. The modest and retiring spouse is, at times, unable to consummate the marriage for nights, perhaps weeks; yet, he is only temporarily impotent; for the inclination and the consequent erection supervene sooner or later. Pills of crumb of bread, and a recommendation to the individual not to approach his wife for a fortnight, whatever may be his desire, have, in almost all cases, removed the impotence! The state of erection is not long maintained, except under unusual excitement; the organ soon returning to its ordinary flaccidity. Its cause is a congestion of blood in the erectile tissue of the corpora cavernosa, urethra, and glans. Swammerdamm and De Graaf cut off the penis of a dog during erection, and found the tissue gorged with blood, and that the organ returned to its flaccid condition, as the blood flowed from it. The same fact, according to Adelon, has been observed in the human subject, where erection has continued till after death. Mr. Callaway, of Guy's Hospital, London, has described the case of an individual, who, in a state of inebriation, had communication three times with his wife the same night, without the consequent collapse succeeding, although emission ensued each time. This condition persisted for sixteen days, not withstanding the use of appropriate means: at this time, an opening was made with a lancet into the left crus of the penis, below the scrotum, and a large quantity of dark, grumous blood, with numerous small coagula escaped. By pressing the penis, the corpora cavernosa were immediately emptied, and each side became flaccid; the communication by the pecten or septum penis permitting the discharge of the contents of both corpora by the incision into the left crus. After recovery, the person remained quite impotent, the organ being incapable of erection, probably owing, as Mr. Callaway suggests, to the deposition of coagulable lymph in the cells of the corpora cavernosa preventing the admission of blood, and the consequent distension of the organ. Artificial erection can, likewise, be induced in the dead body by injections, so that but little doubt need exist, that the enlargement and rigidity of the penis, during erection, are caused by the larger quantity of blood sent into it. The great difficulty has been, to account for this increased flow. The older writers ascribed it to the compression of the internal pudic vein against the symphysis pubis, owing to the organ being raised towards the abdomen by the ischio-cavernosi muscles; and as the cavernous vein empties its blood into the internal pudic, stagnation of blood in the corpora cavernosa ought necessarily to result from such compression, and consequent distension of the organ; whilst the cavernous arteries, being firmer, could not yield to the compression, and would, therefore, continue to convey the blood to the penis. It is obvious, how-

Physiologie de l'Homme, edit. citat. iv. 57.
 London Medical Repository, for April, 1824.

ever, that here, - as in every case, where the erectile tissue is concerned, - the congestion must be of an active kind: the beating of the arteries and the coloration of the organ indicate this; and, besides, compression of the pudic vein cannot precede erection; it must, if it occur at all, be regarded rather as a consequence of erection than as its cause. The case of the female nipple affords us an instance of erectility, where no compression can be invoked, and where the distension must be caused by augmented flow of blood by the arteries. If the nipple be handled, particularly whilst the female is under voluptuous excitement, it will be found to enlarge, and to become rigid, or to be in a true state of erection. The correct opinion is, that irritation of this erectile tissue is the first link in the chain of phenomena constituting erection. The feeling of pleasure is certainly experienced there, prior to, and during erection; and this irritation, like every other, solicits an increased flow of blood into the erectile tissue, which, by organization, is capable of considerable distension. The erectile tissues of the corpora cavernosa, and of the corpus spongiosum urethræ, and glans, are all concerned in the process; but in what precise manner, physiologists are not entirely agreed. Some have supposed, that the blood is effused into the cells, and is consequently out of the vessels. Another view, supported by some of the most eminent anatomists and physiologists, is, that the blood simply accumulates in the venous plexuses of the corpora cavernosa. seems to have been the opinion of Tiedemann, Stieglitz, and J. Müller, a and of Cuvier, Chaussier, and Béclard, from their injections; and the rapidity, with which erection disappears, favours the notion. The discovery of the helicine arteries of the penis by Professor Müller, described at page 324 of this volume, has led to the inference that the peculiar arrangement may be concerned in the function in question, but in what manner the circulation in the male organ or its erection is modified by them is not appreciated: b Valentin, indeed, - as we have seen, denies their existence.

It has been asked again, whether this accumulation of blood be, as we have remarked, an increased afflux by the arteries, or a diminished action of the veins; or these two states combined.º The last opinion is probably the most correct. The arteries first respond to the appeal; the organ is, at the same time, raised by the appropriate muscles; its tissue becomes distended, the plexus of veins turgid, and the return of blood impeded. In this way the organ acquires the rigidity, necessary for penetrating the parts of the female. The friction, which then occurs, keeps up the

c Chaussier and Adelon, in art. Erection, Dict. des Sciences Médicales, xi. 461,

Berlin, 1834; also, Adelon's Physiologie de l'Homme, iv. 57.

a Art. Erection, in Encyclop. Wörterb. der. Medicin. Wissenschaft. xii. 460, Berlin,

b Wagner, Elements of Physiology, translated by R. Willis, p. 75, London, 1841. See, on the Causes of Erection, Krause in Müller's Archiv. H. i. 1837; and London Med. Gazette, for Aug. 1837.

voluptuous excitement, and the state of erection. This excitement is extended to the whole generative system; the secretion of the . testicles is augmented; the sperm arrives in greater quantity in the vesiculæ seminales; the testicles are drawn up towards the abdominal rings by the contraction of the dartos and cremaster, so that the vas deferens is rendered shorter, and, in the opinion of some, the sperm filling the excretory ducts of the testicle, is in this manner forced mechanically forwards towards the vesicles. When these have attained a certain degree of distension, they contract suddenly and powerfully, and the sperm is projected through the ejaculatory ducts into the urethra. At this period the pleasurable sensation is at its height. When the sperm reaches the urethra, the canal is thrown into the highest excitement; and the ischiocavernosus and bulbo-cavernosus muscles, with the transversus perinei, and levator ani, are thrown into violent contraction; the two first holding the penis straight, and assisting the others in projecting the sperm along the urethra. By the agency of these muscles and of the proper muscular structure in the urethra, the fluid is expelled, not continuously, but in jets, as it seems to be sent into the urethra by the alternate contractions of the vesiculæ These muscular contractions are of a reflex character, being independent of the will, and incapable of being controlled by any exertion of it. They are induced, as in deglutition, by a special excitant, — the food in one case; the sperm in the other.

The quantity of sperm discharged varies materially according to the circumstances previously mentioned: its average is estimated at about two drachms. Along with the true sperm, the fluids of the prostate and of the glands of Cowper are discharged; so as to constitute the semen as we meet with it. When the emission is accomplished, the penis gradually returns to its ordinary state of flaccidity; and it is usually impracticable, by any effort, to repeat the act without the intervention of a certain interval of repose, to enable the due quantity of sperm to collect in the spermatic vessels, and vesicles. In some persons, however, the excitability is so great, and the secretion of sperm so ready, that little or no interval is required between the first and second attempt.

This comprises the whole of the agency of the male in the func-

tion of gestation.

In man the emission of sperm is soon effected; but in certain animals it is a long process. In the dog, which has no vesiculæ seminales, the penis swells so much, during copulation, that it cannot be withdrawn until the emission of sperm removes the erection.

In the female, during copulation, the clitoris is in the same state of erection as the penis; as well as the spongy tissue, lining more especially the entrance of the vagina; and it is in these parts, particularly in the clitoris, that pleasure is experienced during sexual desire, and during copulation. This feeling persists the

whole time of coition, and ultimately attains its acme, as in the case of the male, but without any spermatic ejaculation. It is not owing to the contact of the male sperm, - for it frequently occurs before or after emission by the male, but is dependent upon some inappreciable modification in the female organs, - in the ovaries or Fallopian tubes, it is supposed by some physiologists. In most - if not in all - cases, an increased discharge suddenly takes place, during the orgasm, from the mucous follicles of the vagina and vulva, but chiefly from the glands of Duverney, the admixture of which with the male sperm is supposed to have some connection with impregnation, and to be perhaps the vehicle for the fecundating principle of the sperm. After the kind of convulsive excitement into which the female is thrown, a sensation of languor and debility is experienced, as in the male, but not to the same extent, - and, in consequence of no spermatic emission taking place in her, she is capable of a renewal of intercourse more speedily than the male, and can better support its frequent

repetition.

b. Fecundation. - An admixture having, in this manner, been effected between the materials furnished by the male and those of the female; after a fecundating copulation conception or fecundation results, and the rudiments of the new being are instantaneously constituted. The well-known fact, that, after the removal of the testicles, the individual is incapable of procreation although the rest of the genital organs may remain entire, is of itself sufficient to show, that the fecundating fluid is the secretion of those organs, and that this fluid is indispensable. Physiologists have not, however, been satisfied with this fact. Spallanzania examined frogs with great attention, whilst in the act of copulation, both in and out of water; and he observed, that, at the moment when the female deposited her eggs, the male darted a transparent liquor through a tumid point which issued from its anus. This liquor moistened the eggs, and fecundated them. To be certain that it was the fecundating agent, he dressed the male in waxed taffeta breeches; when he found, that fecundation was prevented, and that sperm enough was contained in the breeches to be collected. This he took up by means of a camel's hair pencil, and all the eggs which he touched with it were fecundated. Three grains of this sperm were sufficient to render a pound of water fecundating; and a drop of a solution which could not contain more than the 2,994,687,500th part of a grain was enough for the purpose. To diminish the objection, that the frog is too remote in organization from man to admit of any analogical deduction, Spallanzani took a spaniel bitch, which had engendered several times; shut her up some time before the period of heat, and waited until she exhibited evidences of being in that condition, which did not happen until after a fortnight's seclusion. He then injected into the vagina and uterus, by means of a common syringe warmed to 100° of Fahrena Sur la Génération, par Senebier, Genév. 1783.

heit, nineteen grains of sperm obtained from a dog. Two days afterwards she ceased to be in heat, and, at the ordinary period, she brought forth three young ones, which not only resembled her but the dog from which the sperm had been obtained. This experiment has been repeated by Rossi, of Pisa, and by Buffolini, of Cesena, with similar results.<sup>a</sup> The success of an analogous experiment on the human species rests on the authority of John Hunter. He recommended an individual, affected with hypospadias, to inject his sperm by means of a warm syringe. His wife afterwards

became pregnant.b

In some experiments on generation, Prévost and Dumas fecundated artificially the ova of the frog. Having expressed the fluid from several testicles, and diluted it with water, they placed the ova in it. These were observed to become tumid and developed; whilst other ova, placed in common water, merely swelled up, and in a few days became putrid. They observed, moreover, that the mucus, with which the ova are covered in the oviduct, - the part corresponding to the Fallopian tube in the mammalia, - assists in the absorption of the sperm, and in conducting it to the surface of the ovum; and that in order to succeed in these artificial fecundations, the sperm must be diluted: if too much concentrated its action is less. They satisfied themselves likewise, that the chief part of the sperm penetrates as far as the ova, as animalcules could be detected moving in the mucus covering their surface, and these animalcules they conceive to be the active part of the sperm. It is not, however, universally admitted, that the positive contact of sperm with the ovum is indispensable to fecundation. Some physiologists maintain, that the sperm proceeds no further than the upper part of the vagina; whence it is absorbed by the vessels of that canal, and conveyed through the circulation to the ovary. This is, however, the most improbable of all the views that have been indulged on this topic; for if such were the fact, impregnation ought to be effected as easily by injecting sperm into the bloodvessels, - the female being, at the time, in a state of voluptuous excitement. It has been directly overthrown, too, by the experiments of Dr. Blundelle on the rabbit, who found, that when the communication between the vagina and the uterus was cut off, impregnation could not be accomplished, although the animal admitted the male as many as fifty times, generally at intervals of two or three days or more. Yet, it is evident - Dr. Blundell remarks - that much of the male fluid must have been deposited in the vagina, and absorbed by veins or lymphatics.d Others have presumed, that when the sperm is thrown into the vagina, a hali-

<sup>&</sup>lt;sup>a</sup> Adelon, Physiologie de l'Homme, iv. 66, Paris, 1829; and Wagner, Elements of Physiology, by Dr. R. Willis, p. 68, Lond. 1841.

b Sir E. Home, Lect. on Comp. Anat. iii. 315; and Burdach's Physiologie, u. s. w. 506.

e Principles and Practice of Obstetricy, Amer. Edit., Washington, 1834.
 d See an experiment with a similar object, by Dr. Harlan, Medical and Physical Researches, p. 627, Philad. 1835.

VOL. II. - 31

tus or aura - the aura seminis - escapes from it, makes its way to the ovary, and impregnates an ovum. Others, again, think, that the sperm is projected into the uterus, and that in this cavity it undergoes admixture with the germ furnished by the female; whilst a last class, with more probability in their favour, maintain that the sperm is thrown into the uterus, whence it passes through the Fallopian tube to the ovary, the fimbriated extremity of the tube, at the time, embracing the latter organ. The late Dr. Dewees, suggested, that after the sperm is deposited on the labia pudendi or in the vagina, it may be taken up by a set of vessels which, he admitted, have never been seen in the human female whose duty it is to convey the sperm to the ovary. This conjecture he conceives to have been in part confirmed by the discovery of ducts, leading from the ovary to the vagina, in the cow and sow, by Dr. Gartner, of Copenhagen. The objections that may be urged against his hypothesis, Dr. Dewees remarks, "he must leave to others." We have no doubt, that his intimate acquaintance with the subject could have suggested many that are pertinent and cogent. It will be obvious, that if we admit the existence of the ducts, described by Gartner, it by no means follows, that they are certainly inservient to the function in question. Independently, too, of the objection that they have not been met with in the human female, it may be urged that if we grant their existence, there would seem to be no reason, why closure of the os uteri, after impregnation, or interruption of the vulvo-uterine canal, by division of the vagina - as in the experiments of Dr. Blundell on rabbits - or division of the Fallopian tubes, should prevent subsequent conception, in the first case during the existence of pregnancy; in the two last, for life. These vessels ought, in both cases, to continue to convey sperm to the ovary, and extra-uterine pregnancies or superfætation ought to be constantly occurring.

MM. Prévost and Dumas<sup>b</sup> are amongst the most recent writers, who maintain that fecundation takes place in the uterus, and they assign the following reasons for their belief. First. That in their experiments, they always found sperm in the cornua of the uterus, and they conceive it natural, that fecundation should be operated only where sperm is. Secondly. That in those animals, whose ova are not fecundated until after they have been laid, fecundation must necessarily be accomplished out of the ovary; and Thirdly, that in their experiments on artificial fecundation, they have never been able to fecundate ova taken from the ovary. In reply to the first of these positions it has been properly remarked by Adelone that the evidence of MM. Prévost and Dumas, with regard to the presence of sperm elsewhere than in the uterus, is only of a negative character; and that, on the other hand, we have the positive

b Annales des Sciences Naturelles, iii. 113.

A Compendious System of Midwifery, 7th edit. Philad. 1835.

<sup>·</sup> Physiologie de l'Homme, 2de édit. iv. 68, Paris, 1829.

testimony of physiologists in favour of its existence in the Fallopian tubes and ovary. Haller asserts, that he found it there; and MM. Prévost and Dumas afford us evidence against the position they have assumed respecting the seat of fecundation. They affirm, that on the first day after copulation, the sperm was discoverable in the cornua of the uterus, and that it was not until after the lapse of twenty-four hours, that it had attained the summits of the cornua. Once they detected it in the Fallopian tubes; - a circumstance which is inexplicable under the view, that fecundation is accomplished in the uterus. Leeuenhoek and Hartsöker also found it in some cases in the Fallopian tube; and still more recently, Bischoff, Wagner, and Dr. M. Barry have discovered spermatozoa in the fluid collected from the surface of the ovary, and within the capsular prolongations of the Fallopian tubes that enclose the ovaria; and recently, Dr. Barry has found, in two cases, spermatozoa within the ovum of the rabbit taken from the Fallopian tube. They were within the thick transparent membrane - zona pellucida - brought with the ovum from the ovary.

In reply to the second argument it may be remarked, that analogies drawn from the inferior animals are frequently very loose and unsatisfactory, and ought consequently to be received with caution. This is peculiarly one of these cases; for fecundation, in the case adduced, is always accomplished out of the body, and analogy might with equal propriety be invoked to prove, that in the human female fecundation must also be effected externally. In answer to the third negative position of MM. Prévost and Dumas, the positive experiments of Spallanzani may be adduced, who succeeded in producing fecundation in ova, that had been previously

separated from the ovary.

The evidence, that conception takes place in the ovary, appears to us convincing. Ovarian pregnancy offers irresistible proof of it. Of this, Mr. Stanley of Bartholomew's Hospital has given an instructive example; and a still more extraordinary instance is related by Dr. Granville. Other varieties of extra-uterine pregnancy are confirmative of the same position. At times, the fœtus is found in the cavity of the abdomen,—the ovum seeming to have escaped from the Fallopian tube when its fimbrated extremity grasped the ovary to receive the ovum and convey it to the cavity of the uterus. At other times, the fœtus is developed in the Fallopian tube,—as in the marginal figure, (Fig. 228)—some impediment having existed to the passage of the ovum from the ovarium to the uterus. This impediment can, indeed, be excited artificially so as to give rise to tubal pregnancy. Nuck applied a

<sup>\*</sup> Elements of Physiology, translated by Dr. R. Willis, p. 66, Lond. 1841.

b Philosophical Transactions for 1839, p. 315.
c Proceedings of the Royal Society, Dec. 8, 1842; and Lond. Med. Gazette, April 7, 1843, p. 73.

d Medical Transactions, vol. vi.

e Philosophical Transactions, for 1820.



Tubal Pregnancy.

ligature around one of the cornua of the uterus of a bitch, three days after copulation; and he found, afterwards, two fœtuses arrested in the Fallopian tube between the ligature and the ovary. Von Baera detected an ovule in its passage along the Fallopian tube in a bitch, and

Raspail asserts, that he once met with an ovule, still attached to

the ovary, which contained an embryo.b

It is obvious, then, from these facts, either that fecundation occurs in the ovarium, or else that the ovum, when fecundated in the uterus, travels along the Fallopian tube to the ovarium, and from thence back again to the uterus, which is not probable.c Besides, that the ovaries are indispensable agents in the function of generation is shown by the well-known fact, that their removal, by the operation of spaying, not only precludes reproduction but takes away all sexual desire. A case is detailed of a natural defect of this kind in an adult woman, who had never exhibited the slightest desire for commerce with the male, and had never menstruated. On dissection, the ovaria were found deficient;

and the uterus was not larger than an infant's.d

But, to prevent impregnation, it is not even necessary that the ovaries should be removed. It is sufficient to deprive them of all immediate communication with the uterus, by simply dividing the Fallopian tubes. On this subject, Haightone instituted numerous experiments, the result of which was, that after this operation a fætus was in no instance produced. The operation is much more simple than the ordinary method of spaying by the removal of the ovaries, and it has been successfully practised, at the recommendation of the author, on the farm of his friend Thomas Jefferson Randolph, Esq. of Virginia. It does not seem that the simple division of the Fallopian tubes takes away the sexual desire, as Haighton supposed. Dr. Blundell has proposed this division of the tubes, and even the removal of a small portion of them, so as to render them completely impervious, when the pelvis is so contracted as not to admit of the birth of a living child in the seventh month; and he goes so far as to affirm, that the operation is much less dangerous than delivery by perforating the head, when the pelvis

\* De Ovi Mammalium et Hominis Genesi, Lips. 1827,

b Chimie Organique, p. 262, Paris, 1833.

· Granville's Graphic Illustrations of Abortion, p. iii. Lond. 1834.

Philosoph. Transactions for 1805. e Philosoph. Transact. for 1797. is greatly contracted. We have already remarked, that sperm has been found in the cavity of the uterus, and even in the Fallopian Fabricius ab Acquapendente maintained, that it could not be detected there; and Harvey contended, that in the case of the cow, whose vagina is very long, as well as in numerous other animals, the sperm cannot possibly reach the uterus, and that there is no reason for supposing that it ever does. In addition, however, to the facts already cited, we may remark, that Mr. John Huntera killed a bitch in the act of copulation, and found the semen in the cavity of the uterus, conveyed thither, in his opinion, per saltum, Ruyschb discovered it in the uterus of a woman taken in adultery by her husband and killed by him; and Hallere in the uterus of a sheep killed forty-five minutes after copulation. An interesting case, in relation to this point, has been published by Dr. H. Bond,d of Philadelphia. A young female, after having passed a part of the night with a male friend, destroyed herself early in the morning, by taking laudanum. On cutting open the uterus, it was found to be thickly coated with a substance having the appearance, and the strong peculiar odour, of the sperm. One of the Fallopian tubes was laid open, and found to contain, apparently, the same matter, but it was not ascertained whether it possessed the seminal odour. More recently, Bischoff,e in his investigations, found few or no spermatozoa in the vagina of bitches and rabbits, after coitus, but the uterus was quite full.

Blumenbach supposes, that, during the venereal orgasm, the uterus sucks in the sperm. It is impossible to explain the mode in which this is accomplished, but the fact of the entrance of the fluid into the uterus, and even as far as the ovarium seems unquestionable. This Dr. Blundellg admits, but he is disposed to think, that, in general, the rudiments from the mother, and the fecundating fluid meet in the uterus; as, in his experiments on rabbits, he found - from the formation of corpora lutea, the development of the uterus and the accumulation of water in the uterine cavity that the rudiments may come down into the uterus, without a previous contact of the semen. His experiments, however, appear to us to prove nothing more, than that infecund ova may be discharged from the ovarium, and that if they are prevented from passing externally, owing to closure of the vagina or cervix uteri, the uterine phenomena, alluded to, may occur. They do not invalidate the arguments already adduced to show, that the ovum must be fecundated in the ovarium. It has been suggested by Dr.

Philosoph. Transact. for 1817.

b Thesaur. Anat. iv.; and Adversaria Anat. Med. Chirurg. Dec. 1.

c Element. Physiol. viii. 22.

<sup>4</sup> American Journal of the Medical Sciences, No. xxvi. for February, 1834, p. 403. · Entwickelungsgeschichte des Kaninchen-Eies, u. s. w. Leipz. 1842; cited by Mr. T. W. Jones, in Brit. and For. Med. Rev., Oct. 1843, p. 513.

Elements of Physiology, by Elliotson, 4th edit. p. 467, Lond. 1828.

Principles, &c., of Obstetricy, Amer. Edit. p. 56, Washington, 1834.

Bostock, a that the ciliary motions, which have been observed by Purkinje, Valentin,b and others in the mucous membranes of the air-passages, and which have been likewise detected in the generative organs, and whose office appears to be to propel substances along them, may have something to do with the propulsion of the sperm towards the ovary; and J. Müllere affirms, that "the mode, in which the semen is conducted so far through the female generative organs, is no longer a problem requiring solution; for the discovery of the ciliary motion affords a solution of it." Dr. Sharpey,d however, remarks, that the direction of these motions, in those organs, is from within outwards, so that he conceives it to be difficult to assign any other office to them than that of conveying outwards the secretion of the membrane, unless we suppose that they also bring down the ovum into the uterus. Prof. Wagnere considers, that the sperm reaches the ovary, partly by the ciliary motions, which begin in the cervix uteri; partly by the contractions of the tubes, and partly by the motility of the spermatozoa. Future observations may shed farther light on this subject.

Granting, then, that conception occurs in the ovarium, and that sperm is projected into the uterus, with or without the actions referred to; in what manner does the sperm exert its fecundating agency on the ovarium? It is manifestly impossible, that the force of projection from the male can propel it, not only as far as the cornua of the uterus, but also through the narrow media of communication between the uterus and ovary by the Fallopian tubes. This difficulty suggested the idea of the aura seminis or aura seminalis, which, it was supposed, might readily pass into the uterus, and through the tubes to the ovary. Haighton, indeed, embraced an opinion more obscure than this, believing, that the semen penetrates no farther than the uterus, and acts upon the ovaria by sympathy; and this view has been adopted by some

distinguished individuals.

In opposition to the notion of the aura seminis, we have some striking facts and experiments. In all those animals, in which fecundation is accomplished out of the body, direct contact of the sperm appears necessary. Spallanzani, and MM. Prévost and Dumas found, in their experiments on artificial fecundation, that they were always unsuccessful when they simply subjected the ova to the emanation from the sperm. Spallanzani took two watch-glasses, capable of being fitted to each other, the concave

Physiology, 3d edit. p. 654, Lond. 1836.

d Art. Cilia, Cyclop. of Anat. and Physiology, part vii. p. 633, July, 1836. See, also, Müller, Elements of Physiology, by Baly, p. 857, Lond. 1838; Todd and Bowman, Physiological Anat. and Physiol. of Man, p. 62, Lond. 1843; and vol. i. p. 472

Müller's Archiv. B. i.; and translation in Dublin Journal of Med. and Chem.
 Science, May, 1835; and in Edinb. New Philos. Journal, for July, 1835.
 Elements of Physiology, translated by Baly, p. 1491, Lond. 1842.

e Elements of Physiology, translated by R. Willis, part i. p. 72, Lond. 1841.

surface of the one being opposed to that of the other. Into the lower he put ten or twelve grains of sperm, and into the upper about twenty ova. In the course of a few hours, the sperm had evaporated, so that the ova were moistened by it; yet they were not fecundated, but fecundation was readily accomplished by touching them with the sperm that remained in the lower glass. A similar experiment was performed by MM. Prévost and Dumas. They prepared about an ounce and a half of a fecundating fluid from the expressed humour of twelve testicles, and as many vesiculæ seminales. With two and a half drachms of this fluid they fecundated more than two hundred ova. The remainder of this fluid was put into a small retort, to which an adopter was attached. In this, forty ova were placed, ten of which occupied the hollowest part, whilst the rest were placed near the beak of the retort. The apparatus was put under the receiver of an air-pump, and air sufficient was withdrawn to diminish the pressure of the atmosphere one-half. The rays of the sun were now directed upon the body of the retort, until the temperature within rose to about 90°; after the lapse of four hours, the experiment was stopped, when the following were the results. The eggs, at the bottom of the adopter, were bathed in a transparent fluid, the product of distillation. They had become tumid as in pure water, but had undergone no development. The eggs, near the beak of the retort, were similarly circumstanced, but all were readily fecundated by the thick sperm, which remained at the bottom of the retort. No aura, no emanation from the sperm consequently appeared to be capable of impregnating the ova. Absolute contact was indispensable.a This is probably the case with the human female, and if so, the sperm must proceed from the uterus along the Fallopian tube to the ovarium. The common opinion is, that during the intense excitement at the time of copulation, the tube is raised, and its digitated extremity applied to the ovarium. The sperm then proceeds along it, - in what manner impelled we know not, - and attains the ovary. According to Blundell and others, during the time of intercourse, the whole of the tube is in a state of spontaneous movement. Cruikshank pithed a female rabbit, when in heat, and examined the uterine system very minutely. The external and internal parts of generation were found black with blood; the Fallopian tubes were twisted like writhing worms, and exhibited a very vivid peristaltic motion; and the fimbriæ embraced the ovaria, like fingers laying hold of an object, so closely and so firmly as to require some force, and even slight laceration to disengage them. Haller states, that by injecting the vessels of the tube in the dead body, it has assumed this kind of action. De Graaf, too, affirms, that he has found the fimbriated extremity adhering to

<sup>. \*</sup> Rudolphi, art. Aura Seminalis, in Encyclopäd. Wörterbuch der Medicinisch. Wissenschaft, B. iv. s. 452, Berlin, 1830; and Hüter, art. Empfangniss, ibid. x. 628, Berlin, 1834.

b Dr. Allen Thomson, art. Generation, in Cyclop. of Anat. and Physiol. part xiii. p. 462, Feb. 1838.

the ovary, twenty-seven hours after copulation; and Magendie, that he has seen the extremity of the tube applied to a vesicle.

As excitement would appear to be necessary to cause the digitated extremity of the Fallopian tube to embrace the ovary, it would seem probable, that a female could not be impregnated without some consciousness of the sexual union. This may be imperfect, as during sleep, or when in a state of stupor — either from spirituous liquors or narcotics — but still some feeling must probably be engendered in order that fecundation may take place.

As the aura seminis appears to be insufficient for impregnation, it is obviously a matter of moment, that the sperm should be projected as high up into the vagina as possible. It has been often observed, that where the orifice of the urethra does not open at the extremity of the glans, but beneath the penis, or at some distance from the point, the individual has been rendered less capable of procreation. In a case, that fell under the care of the author, the urethra was opened opposite the corona glandis by a sloughing syphilitic sore, and the aperture continued, in spite of every effort to the contrary. The individual was married, and the father of three or four children; but after this occurrence he had no increase of his family. Many medico-legal writers have considered, that when the urethra terminates at some other than its natural situation, impotence is the necessary result, - and that although copulation may be effected, impregnation is impracticable. Zacchias,a however, gives a positive case to the contrary. Belloc, too, asserts, that he knew of a person, in whom the orifice of the urethra terminated at the root of the frænum, who had four children that resembled the father, two having the same malformation; and Dr. Francis refers to the case of an inhabitant of New York, who, under similar circumstances, had two children. Many such cases are, indeed, on record. We cannot, therefore, regard it as an absolute cause of impotence, but the inference is just, that if the semen be not projected far up into the vagina, and in the direction of the os uteri, impregnation is not likely to be accomplished; a fact, which it might be of moment to bear in mind, where the rapid succession of children is an evil of magnitude. Yet, notwithstanding this weight of evidence, it has been affirmed by Professor Heim,d that impregnation may take place by the simple contact of the sperm with the lower part of the abdomen. The answer to this view, by D'Outrepont, appears to us, however, overwhelming. Heim relies on statements made by the parties that no

d Wochenschrift für die Gesammte Heilkunde; and Gazette Médicale, Sept. 26,

<sup>Quæstiones Medico-Legales, Lugd. 1674.
Cours de Médecine Légale, p. 50, Paris, 1819.</sup> 

c Beck's Elements of Medical Jurisprudence, 6th edit. p. 71, Philad. 1838; and Traill's Outlines of a Course of Lectures on Med. Jurisprudence, p. 26, Edinb. 1840, or American Edition, by the author, Philad. 1841; and W. A. Guy, Principles of Forensic Medicine, part i. p. 44, Lond. 1843.

Neue Zeitschrift für Geburtskunde, von Busch, d'Outrepont und Ritgen, B. iv.
 H. ii., Berlin, 1836; or Dunglison's Amer. Med. Intelligencer, p. 275, Nov. 1, 1837.

penetration existed; but D'Outrepont properly observes, that whenever this has been alleged in a case of pregnancy, he has found, when the parties were strictly questioned, that one or both of them admitted, that the male organ *might* have been in the vagina, or at least within the vulva.

The part, then, to which the sperm is applied is the ovary, and probably, according to Bischoff, the liquor sanguinis is the substance that passes through the parietes of the follicle by imbibition, in order to reach the ovum. Mr. Jones states, that although he is not prepared to deny this, yet when he takes into consideration all the evidence on the subject, he is of opinion, that there is no proof, that fecundation takes place until the ovum has escaped from the Graafian follicle, and comes into direct contact with

the sperm.a

Let us now inquire into the changes experienced by this body after a fecundating copulation. Fabricius ab Acquapendente,b having killed hens a short time after they had been trodden, examined their ovaries, and observed, - amongst the small yellow, round granula, arranged racemiferously, which constitute those organs, - one having a small spot, in which vessels became developed. This increased in size, and was afterwards detached, and received by the oviduct; becoming covered, in its passage through that tortuous canal and the cloaca, by particular layers, especially by the calcareous envelope; and being ultimately extruded in the form of an egg. Harvey, in his experiments on the doe, made similar observations. He affirms, positively, that the ovary furnishes an ovum, and that the only difference, which exists amongst animals in this respect, is, that, in some, the ovum is hatched after having been laid, whilst, in others, it is deposited in a reservoir a womb - where it undergoes successive changes. De Graafd instituted several experiments on rabbits, for the purpose of detecting the series of changes in the organs from conception till delivery. Half an hour after copulation, no alteration was perceptible, except that the cornua of the uterus appeared a little redder than usual. In six hours, the coverings of the ovarian vesicles or vesicles of De Graaf seemed reddish. At the expiration of a day from conception, three vesicles in one of the ovaries, and five in the other, appeared changed, having become opaque and reddish. After twenty-seven, forty, and fifty hours, the cornua of the uterus and the tubes were very red, and one of the tubes had laid hold of the ovary; a vesicle was in the tube, and two in the right cornu of the uterus. These vesicles were as large as mustard seed. They were formed of two membranes, and were filled by a limpid fluid. On the fourth day, the ovary contained only a species of envelope, called, by De Graaf, a follicle; this appeared to be the capsule, which had contained the ovum. The ovum itself was

<sup>\*</sup> T. W. Jones, Report on the Ovum of Man and the Mammifera, in Brit. and For. Med. Rev., Oct. 1843, p. 525.

\* De Generatione, p. 228.

\* Tom. i. 310.

in the cavity of the uterus, had augmented in size, and its two envelopes were very distinct. Here it remained loose until the seventh day, when it formed an adhesion to the uterus. On the ninth day, De Graaf observed a small opaque point, a kind of cloud, in the transparent fluid that filled the ovum. On the tenth day, this point had the shape of a small worm. On the eleventh, the embryo was clearly perceptible; and from this period, it underwent its full development, until the thirty-first day, when delivery took place. Malpighia and Vallisnierib also observed, in their experiments, that after a fecundating copulation, a body was developed at the surface of the ovary, which subsequently burst, and suffered a smaller body to escape. This was laid hold of by the tube, and conveyed by it to the uterus. It is not, however, universally admitted, that this body is the impregnated ovum; some affirming, that it is a sperm similar to that of the male; and others, that it is an amorphous substance, which, after successive developments, becomes the new individual.c Haller exposed the females of sheep and of other animals to the males on the same day; and killed them at different periods after copulation, for the purpose of detecting the whole series of changes by which the vesicle is detached from the ovary and conveyed to the uterus. Half an hour after copulation, one of the vesicles of the ovary appeared to be prominent; to have on its convexity a red, bloody spot, and to be about to break; in an hour or more, the vesicle gave way, and its interior seemed bleeding and inflamed. What remained of the vesicle in the ovary, and appeared to be its envelope, gradually became inspissated, and converted into a body of a yellowish colour, to which Haller gave the name corpus luteum. The cleft, by which the vesicle escaped, was observable for some time, but, about the eighth day, it disappeared. On the twelfth day, the corpus luteum became pale and began to diminish in size. it continued to do until the end of gestation; and ultimately became a small, hard, yellowish or blackish substance, which could always be distinguished in the ovarium, by the cicatrix left by it. Its size was greater, the nearer the examination was made to the period of conception. In a bitch, for example, on the tenth day, it was half the size of the ovary; yet it proceeded, in that case. from one vesicle to the other. In multiparous animals, as many corpora lutea existed as fœtuses.

The experiments of Haller<sup>d</sup> have been frequently repeated and with similar results. Magendie,<sup>e</sup> whose trials were made on bitches, observed, that the largest vesicles of the ovary were greatly augmented in size, thirty hours after copulation; and that the tissue of the ovary, surrounding them, had acquired greater consistence, had changed colour, and become of a yellowish-gray. This part

De Formatione Pulli in Ovo, Lond. 1673; and De Ovo Incubato, Lond. 1686.

b Istoria della Generazione dell'Uomo, Discorsi Academ. iv. Venez. 1722-1726.

c Adelon, Physiologie de l'Homme, iv. 74.

d Element. Physiologiæ, xxix. Précis de Physiologie, edit. citat. ii. 534.

was the corpus luteum. It, as well as the vesicles, increased for the next three or four days; and seemed to contain, in its areolæ, a white, opaque fluid, similar to milk. The vesicles now successively ruptured the external coat of the ovary, and passed to the surface of the organ, still adhering to it, however, by one side. Their size was sometimes that of a common hazlenut, but no germ was perceptible in them. The surface was smooth, and the interior filled with fluid. Whilst they were passing to the uterus, the corpus luteum remained in the ovary, and underwent the changes referred to by Haller. Insimilar experiments, instituted by MM. Prévost and Dumas," no change was perceptible in the ovary during the first day after fecundation; but, on the second day, several vesicles enlarged, and continued to do so for the next four or five days, so that from being two or three millimètres in diameter, they attained a diameter of eight. From the sixth to the eighth day, the vesicles burst, and allowed an ovule to emerge, which often escaped observation, owing to its not being more than half a millimètre in diameter, but was clearly seen by MM. Prévost and Dumas by the aid of the microscope. This part they term ovule, in contradistinction to that developed in the ovary, which they call vesicle. The latter has the appearance, at its surface, of a bloody cleft, into which a probe may be passed; and in this way it can be shown, that the vesicle has an interior cavity, which is the void space left by the ovule after its escape from the ovarium into the Fallopian tube. On the eighth day, in the bitch, the ovule passes into the uterus. All the ovules do not, however, enter that cavity at the same time; -an interval of three or four days sometimes occurring between them. When they attain the uterus, they are at first free and floating; and if examined with a microscope magnifying twelve diameters, they seem to consist of a small vesicle, filled with an albuminous, transparent fluid. If examined in water, their upper surface has a mammiform appearance, with a white spot on the side. This is the cicatricula. These ovules speedily augment in size, and, on the twelfth day, fœtuses can be recognised in them.

Similar experiments, with like results, have been made by Von Baer, b Seiler, c and others; and much minute attention has been paid to the subject by recent histologists, — by Wagner, Barry, Bischoff and T. W. Jones more especially. The subject, however, is altogether in a transition state; and it is difficult to say positively what is the condition of the ovum from the period post coitum until it leaves the ovary. After the intercourse of the sexes, an increased flow of blood takes place to the ovaries; the vascular membrane of the Graafian vesicle enlarges; the granules, mingled

a Annales des Sciences Naturelles, iii. 135.

De Ovi Mammalium, &c. Epist. Lips. 1827.
 Das Ei und Die Gebärmutter des Menschen, Dresd. 1832; Burdach's Physiologie
 als Erfahrungswissenschaft, Th. ii. Leipz. 1828; and Purkinje, art. Ei, in Encyclop.
 Wörterb. der Medicin. Wissensch. x. 107, Berlin, 1834.

with its contents, become greatly developed, and changed; and thickening and general increase of its walls, particularly of the base and sides supervene; the ovum and other contents of the follicle are by this pressed forwards, or against that aspect of the follicle, which is in contact with the peritoneum, and which now becomes thinner and thinner, and finally bursts, so that the ovum escapes, and a cavity is left in the ovary, which is soon obliterated by the growth on all sides of the inner membrane of the follicle; a reddish, fleshy-looking mass sprouts from the walls towards the shrinking cavity, and the rent by which the ovum had escaped is finally closed. The follicle, thus altered, constitutes the *corpus luteum*—to be described presently. Such is the description of Wagner.<sup>a</sup>

From the above facts, then, we may conclude, that the sperm excites the vesicles in the ovaries to development; that the ova, within the germinal part, burst their covering, are laid hold of by the Fallopian tube, and conveyed to the uterus, where they re-

main during the period of gestation.

In the passage of the ova along the Fallopian tube it has generally been believed that they experience but little change. They carry off, according to Wagner, b a small portion of the granular stratum of the vesicle with them, which appears hanging to them at first as an irregular, ragged discoidal appendage; but this is soon detached. The ovum gains a little in size; but there is still no trace of any special spot where the embryo arises. It would seem, however, from the researches of British embryologists, that the outer membrane of the ovum, which it acquired in the ovary, becomes swoln with moisture in the tube, and assumes the appearance of a thick, gelatinous-looking membrane - the proper chorion; and by and by, the zona pellucida having disappeared, the newly-formed chorion comes to be the only investment of the yolk.º All the observations on the ovum in the Fallopian tubes have, however, been made on animals. Of the human ovum whilst in these tubes, nothing is known.d

The exact time, required by the ovum or ova to make their way into the uterus, has not been accurately determined. Cruikshanke found, that in rabbits forty-eight hours were necessary. Haightonf divided one of the Fallopian tubes in a rabbit; and, having exposed the animal to the male, he observed that gestation occurred only on the sound side. On making this section after copulation, he found, that if it were executed within the two first days, the descent of the ovules was prevented; but if it were delayed for sixty hours, the ovules had passed through the tube and were in the cavity of the uterus. A case, too, is quoted by writers on

Elements of Physiology, translated by R. Willis, p. 136, Lond. 1842.

R. Willis, Ibid. p. 138.

Ibid. p. 142.

d T. W. Jones, Brit. and For. Med. Rev., Oct. 1843, p. 542.

e Philosoph. Transactions, for 1797. f Ibid. lxxxvii. 304.

this subject, on the authority of a surgeon named Bussières, who observed an ovoid sac, about the size of a hazlenut and containing an embryo, half in the Fallopian tube and half adherent to the ovary. The minuteness of the calibre of the Fallopian tube is not as great a stumbling-block in the way of understanding how this passage is effected, as might appear at first sight. The duct is, doubtless, extremely small in the ordinary state; but it admits of considerable dilatation. Magendie<sup>b</sup> asserts, that he once found it half an inch in diameter. Moreover, the size of the ovum, as

we have just seen, is only  $\frac{1}{200}$ th part of an inch.

The period that elapses between a fecundating copulation and the passage of the ovum from the ovarium to the uterus, is different in different animals. In the sheep it occurs, according to Hallere and Kuhlemannd on the seventeenth day. In rabbits, it is uncertain, but occurs generally on the third or fourth day after copulation; in bitches on the fifth, according to some, but not till after the lapse of ten or twelve days, according to others; f and in the human female, perhaps about the same time; yet Mr. Burns infers from analogical evidence, that we should be more justified in the belief that the ovum, in the human female, does not enter the uterus until at least three weeks after conception.8 Maygrier refers to a case of abortion twelve days after copulation; the abortment consisting of a vesicle, shaggy on its surface and filled by a transparent fluid. One of the most instructive cases that we possess on this subject is given by Sir Everard Home, and although, as Dr. Granville has remarked, it has lately been the fashion to doubt the accuracy of the case, or to esteem it morbid, there is reason to believe it correct, from the circumstance of Mr. Bauer's microscopic examination of the ovulum, and description of its structure corresponding with the more recent discoveries of Professor Boer. A servant-maid, twentyone years of age, had been courted by an officer, who had promised her marriage, in order that he might more easily accomplish his wishes. She was but little in the habit of leaving home, and had not done so for several days, when she requested a fellowservant to remain in the house, as she was desirous of calling upon a friend, and should be detained some time. This was on the 7th of January, 1817. After an absence of several hours, she returned

\* Adelon, Physiologie de l'Homme, edit. cit. iv. 77. b Précis, &c. edit. cit. p. 536.

c Element. Physiol. viii. 59.

d Observ. quædam circa Negotium Generationis in Ovibus fact. Gotting. 1753.
 e Recherches sur la Génération des Mammifères par Coste, suivies de Recherches sur la Formation des Embryons, par Delpech et Coste, Paris, 1834.

Wagner, Elements of Physiology, by Willis, p. 137, Lond. 1841.

The Principles of Midwifery, &c. 3d edit. p. 132, Lond. 1814.

h Philosophical Transactions for 1807, p. 252; and Lectures on Comparative Anatomy, iii. 288, Lond. 1823.

Graphic Illustrations of Abortion, &c., p. vii. Lond. 1834.

Weber's Hildebrandt's Handbuch der Anatomie, iv. 465, Braunschweig, 1832; Dr. Allen Thomson, in art. Generation, Cyclop. of Anat. and Physiol. P. xiii. p. 454, Feb. 1838; and Dr. John Davy, Researches, Physiological and Anatomical, Dunglison's Amer. Med. Libr. Edit., p. 379, Philad. 1840.

VOL. II. - 32

with a pair of new corsets, and other articles of dress which she had purchased. In the evening she got one of the maid-servants to assist her in trying on the corsets. In the act of lacing them, she complained of considerable general indisposition, which disappeared on taking a little brandy. Next day, she was much indisposed. This was attributed to the catamenia not having made their appearance, although the period had arrived. On the following day, there was a wildness in her manner, and she appeared to suffer great mental distress. Fever supervened, which confined her to her bed. On the 13th, she had an epileptic fit, followed by delirium, which continued till the 15th, when she expired in the forenoon. On making inquiries of her fellow-servants, many circumstances were mentioned which rendered it highly probable, that on the morning of the 7th, when she was immediately on the point of menstruating, her lover had succeeded in gratifying his desires; and that she had become pregnant on that day, so that, when she died, she was in the seventh or eighth day of impregnation. Dissection showed the uterus to be much larger than in the virgin state, and considerably more vascular. On accurately observing the right ovarium, in company with Mr. Clift, Sir Everard noticed, upon the most prominent part of its outer surface, a small ragged orifice. This induced him to make a longitudinal incision in a line close to this orifice, when a canal was found leading to a cavity filled with coagulated blood, and surrounded by a narrow yellow margin, in the structure of which the lines had a zigzag appearance. The cavity of the uterus was then opened, by making an incision through the coats from each angle; and from the point where these incisions met, a third incision was continued down through the os uteri to the vagina. The os uteri was found completely blocked up by a plug of mucus, so that nothing could have escaped by the vagina; the orifices leading to the Fallopian tubes were both open, and the inner surface of the cavity of the uterus was composed of a beautiful efflorescence of coagulable lymph resembling the most delicate moss. By attentive examination, Sir Everard discovered a small, spherical, transparent body concealed in this efflorescence, which was the impregnated ovum. This was submitted to the microscopic investigations of Mr. Bauer, who made various drawings of it, and detected two projecting points, which were considered to mark out, even at this early period, and before the ovum was attached to the uterus, the seat of the brain and spinal marrow. This case shows, that an ovum had left the ovarium, and that it was in the interior of the uterus, prior to the seventh or eighth day after impregnation. Weber and Von Baer have each recorded a case in which there was an opportunity for examining the embryo, probably eight days after a fecundating copulation; but no ovum was detected either in the uterus or tubes.

On comparing the degree of advancement of the fœtus in the ovum, described by different observers, with that of the fœtus in

the dog, cat, and sheep, at known periods, Dr. Allen Thomsona hazards the opinion, that the human ovum does not arrive in the uterus before the eleventh or twelfth day after conception. Valentin, indeed, thinks the twelfth or fourteenth day. From this discrepancy, however, amongst observers, it is manifest that our knowledge on the matter is by no means fixed or definite.

But, it has been asked, is it a mere matter of chance, which of the ovarian vesicles shall be fecundated; or are there not some that are riper than the rest, and that receive, by preference, the vivifying influence of the sperm? MM. Prévost and Dumas have shown, that such is the case with oviparous animals. They found, in their experiments, that not only were the vesicles of the ovaries of frogs of different sizes, but that the largest were always first laid, whilst the smallest were not to be deposited until subsequent years. In all the animals, whose eggs were fecundated externally, they seemed evidently prepared or maturated. We have, too, the most indubitable evidence that birds - although unquestionable virgins - may lay infecund eggs. Analogy would lead us to believe, that something similar may happen to the viviparous animal, and direct observation has confirmed the position. In all cases in which an ovum has escaped, a cavity of course is left in the ovarium, which is filled up, in the manner already mentioned, by a growth from the inner membrane of the Graafian follicle, which, thus altered, constitutes the corpus luteum.

Not longer ago than the year 1808, the existence of the corpora lutea in the ovaria was held to be full proof of impregnation. In that year, Charles Angus, Esq., of Liverpool, England, was tried at the Lancaster Assizes, for the murder of Miss Burns, a resident of his house.d The symptoms previous to her decease, and the appearances observed on dissection, were such as to warrant the suspicion that she had been poisoned. The uterine organs were also found to be in such a state as to induce a belief, that she had been delivered a short time before her death, of a fætus, which had nearly arrived at maturity. It was not, however, until after the trial, that the ovaria were examined, in the presence of a number of physicians, and a corpus luteum was distinctly perceived in one of them. The uterus was taken to London and shown to several of the most eminent practitioners there, all of whom appear to have considered that the presence of a corpus luteum proved the fact of pregnancy beyond a doubt.d Such, indeed, is the positive averment of Haller, an opinion which was embraced by Haighton, who maintained that they furnish "incontestable proof" of previous impregnation. It was this belief, - coupled with the

Art. Generation, in Cyclop. of Anat. and Physiol. P. xiii. p. 454, Feb. 1838.

<sup>&</sup>lt;sup>b</sup> Hüter, art. Empfangniss, in Encyclopäd. Wörterb. der Medicin. Wissensch. x. 629, Berlin, 1834.

c F. Renaud, Lond. and Edinb. Monthly Journal of Medical Science, June, 1842.

d Edinb. Med. and Surg. Journal, v. 220.

e Beck's Medical Jurisprudence, 6th edit. i. 247, New York, 1838.

f Element. Physiolog. xxix. 1. 8 Philosoph. Transact. lxxxvii. 159.

fact, that division of the Fallopian tubes, in his experiments, prevented impregnation, whilst corpora lutea were found, notwithstanding, in the ovary, — which led him to the strange conclusion, that the semen penetrates no farther than the uterus, and acts upon

the ovaria by sympathy.

Sir Everard Homea affirms, that corpora lutea exist independently of impregnation. Upon examining," said he, "the ovaria of several women, who had died virgins, and in whom the hymen was too perfect to admit of the possibility of impregnation, there were not only distinct corpora lutea, but also small cavities around the edge of the ovarium, evidently left by ova, that had passed out at some former period;" and he affirms, that whenever a female quadruped is in heat, one or more ova pass from the ovaria to the uterus, whether she receives the male or not. view of the subject appears to have been first propounded by Blumenbach, b who remarks that the state of the ovaria of females, who have died under strong sexual passion, has been found similar to that of rabbits during heat; and he affirms, that in the body of a young woman, eighteen years of age, who had been brought up in a convent, and had every appearance of being a virgin, Vallisnieri found five or six vesicles pushing forward in one ovarium, and the corresponding Fallopian tube redder and larger than usual, as he had frequently observed in animals during heat. Bonnet, he adds, gives the history of a young lady, who died vehemently in love with a man of low station, and whose ovaria were turgid with vesicles of great size. It has been already remarked, under Menstruation, that the periodical recurrence of that function has been supposed to consist in the production and development of vesicles in the ovary, that is, of a matured ovum which is periodically brought forward either to be expelled with the menstrual flux, or to be destroyed in the ovary; a view which was entertained by Cruikshank, and has been recently urged strongly by Gendrin.

Buffon, again, maintained, that instead of the corpus luteum of Haller being the remains of the ovule, it is its rudiment; and that the corpus exists prior to fecundation, — as he also found it in the virgin. Lastly, Dr. Blundelle states, that he has in his possession a preparation, consisting of the ovaries of a young girl, who died of cholera under seventeen years of age, with the hymen, which nearly closed the entrance of the vagina, unbroken. In these ovaries, the corpora lutea are no fewer than four; two of them being a little obscure, but easily perceptible by an experienced eye. The remaining two are very distinct, and differ from the corpus luteum of genuine impregnation merely by their more diminutive size and the less extensive vascularity of the con-

c Researches, Physiol. and Pathological, p. 49, Lond. 1825.

<sup>&</sup>lt;sup>2</sup> Philosoph. Transact. for 1817 and 1819; and Lectures on Compar. Anat. iii, 304. <sup>3</sup> Comment. Soc. Roy. Scient. Gotting. ix. 128; and Elliotson's edit. of Blumenbach's Physiology, 4th edit., p. 468, Lond. 1828.

tiguous parts of the ovary. "In every other respect," says Dr. Blundell, " in colour and form, and the cavity which they contain, their appearance is perfectly natural, indeed, so much so, that I occasionally circulate them in the class-room, as accurate specimens of the luteum upon the small scale." Mr. Stanleya confirms the fact of the corpora lutea of virgins being of a smaller size than those that are the consequences of impregnation; and Dr. Montgomeryb says, that he has seen many of these virgin corpora lutea. "as they are unhappily called," and has preserved several specimens of them, but not in any instance, did they present what he would regard as even an approach to the assemblage of characters belonging to the true corpus luteum, -the result of impregnation; from which, according to him, they differ in the following particulars: - 1. There is no prominence or enlargement of the ovary over them. 2. The external cicatrix is almost always wanting. 3. There are often several of them found in both ovaries, especially in subjects who have died of tubercular diseases,

such as phthisis, in which case they appear to be merely depositions of tubercle, and are frequently without any discoverable connection with the Graafian vesicles. 4. They present no trace whatever of vessels in their substance, of which they are in fact entirely destitute, and of course cannot be injected. 5. Their (Montgomery.)

Fig. 229.

texture is sometimes so infirm, that it seems to be merely the remains of a coagulum, and at

others appears fibro-cellular, like that of the internal structure of the ovary; but never presents the soft, rich, lobulated, and regularly glandular appearance, which Hunter meant to express, when he described them as "tender and friable like glandular flesh." 6. In form they are often triangular, or square, or of some figure bounded by straight lines; and 7. They never present, either the central cavity, or the radiated or stelliform white line, which

results from its closure.c Figures 229 and 230 represent the corpus luteum in the third, and at the end of the ninth month, respectively.



Corpus Luteum at the end of the ninth month. - (Montgomery.)

\* Transactions of the Royal College of Physicians of London, vol. vi.

b An Exposition of the Signs and Symptoms of Pregnancy, &c., p. 245, Lond. 1837, or Dunglison's Amer. Med. Lib. Edit., Philad. 1839; and art. Signs of Pregnarcy and Delivery, in Cyclop. of Pract. Medicine, Lond. 1833.

· See Dr. E. Rigby, System of Midwifery, American Edit., p. 27, Philad. 1841;

and Dr. R. Paterson, Edinb. Med. and Surg. Journ. Jan. 1840.

Recently, however, Dr. William Davidson, of Edinburgh, has published three dissections of females - not one of whom was pregnant - and in each, corpora lutea were found. They all possessed the characters assigned them by Dr. Montgomery, -a central cavity, or fibrous coagulum; an oval shape, and a radiated white cicatrix in the centre, just about the central body; - the body being at the same time immediately under the peritoneal coat. This last point is dwelt upon by Dr. Robert Lee, who maintains, that false corpora lutea are never observed in immediate connection with the peritoneum, a small portion of stroma intervening. One of the females had been in a weakly condition for some years, and had no children; another was unmarried, and had menstruated three days previous to her death; of the third, there was no history, but all the organs were healthy, and the Fallopian tube and uterus were in every respect natural. Dr. Davidson expresses his confident opinion, that in none of the cases had there been impregnation prior to the appearance of these bodies; and he refers to Professors Alison, Allen Thomson, John Reid, and Mr. Goodsir, in proof of the accuracy of his statement, and of their perfect resemblance to true corpora lutea. Dr. Davidson states, as the result of his investigations, that he is led to believe, that impregnation cannot take place without the appearance of a true corpus luteum. but that a true corpus luteum may appear independently of impreg-

It is not yet decided at what period the central cavity disappears or closes up to form the stellated line. Dr. Montgomery thinks he has invariably found it existing up to the end of the fourth month. He has one specimen in which it was closed in the fifth month, and another in which it was open in the sixth, but later than this he has never found it.

The structure of the corpus luteum is of a peculiar kind, and is not distinctly seen in small animals, or in those that have numerous litters; but in the cow, which commonly has only one calf at a birth, it is so large, according to Sir Everard Home, that, when magnified, the structure can be made out. It is a mass of thin convolutions, bearing a greater resemblance to those of the brain than to any other organ. Its shape is irregularly oval, with a central cavity, and in some animals, its substance is of a bright orange-colour, when first exposed. The corpora lutea are found to make their appearance immediately after puberty, and they continue to succeed each other, as the ova are expelled, till the period arrives when impregnation can no longer be accomplished. Sir Everard's theory, regarding these bodies, is, that they are glands, formed purposely for the production of ova, — and a similar view is entertained by Seiler, that they exist previous to, and are uncon-

b Lec. on Comp. Anat. iii. 303.

Lond. and Edinb. Monthly Journal of Med. Science, Dec. 1841.

<sup>1</sup> as Ei und die Gebärmutter des Menschen, u. s. w., Dresd. 1832; and Stannius,

nected with, sexual intercourse, and, when they have fulfilled their office of forming ova, they are removed by absorption, whether the ova be fecundated or not.

Figures, 231, a and b, afford an external and internal view of a human ovary, that did not contain the ovum, from which a child had been developed. It was taken immediately after the child was born. The corpus luteum is nearly of the full size. a and b, Fig. 232, afford an external and internal view of the ovarium, in which the impregnated ovum had been formed. The latter figure exhibits how much the corpus luteum had been broken down. In it we see a new corpus luteum forming.

From all these facts, then, we are perhaps justified in concluding with Sir Everard Home, and Messrs. Blundell, Saumarez, Cuvier, and others,e that something resembling a corpus luteum may be produced independently even of sexual intercourse, by the mere excitement of high carnal desire, during which it is probable, that the digitated extremity of the Fallopian tube embraces the ovary, a vesicle bursts its covering, and a yellow body remains. The ovule is conveyed along the tube into the uterus, but, being infecund, it undergoes no farFig. 231.





Corpora Lutea. - (Sir E. Home.)

ther development there; so that unimpregnated ova may, under such circumstances, be discharged, as we observe in the oviparous animal.

When pregnancy is over, the corpus luteum gradually diminishes in size and disappears. Dr. Montgomery remarks, that the exact period of its total disappearance he is unable to state, but

art. Eierstock, in Encycl. Wörterb. der Medicin. Wissensch. x. 193, Berl. 1834. See also, Weber's Hildebrandt's Handbuch der Anatomie, iv. 464, Braunschweig, 1832.

\* Op. cit. iii. 304.

<sup>b</sup> A New System of Physiology, i. 337; see Granville's Graphic Illustrations of Abortion, part vi. Lond. 1834; and Dr. Allen Thomson, in art. Generation, Cyclop. Anat. and Physiol. part. xiii. p. 450, for Feb. 1838.

c For a history of the opinions entertained at various times regarding the corpus luteum, see Dr. Paterson, Edinb. Med. and Surg. Journal, April, 1841, p. 402.

Fig. 232.





Corpora Lutea .- (Sir E. Home.)

that he has found it distinctly visible so late as the end of five months after delivery at the full time, but not beyond this period. It would appear, therefore, that in a few months after the termination of pregnancy, all traces of the corpus luteum are lost, and that, consequently, it will be impossible to decide how frequently impregnation has taken place, merely by examining the ovaries.<sup>a</sup>

c. Theories of Generation. - We have now endeavoured to demonstrate the part performed by the two sexes in fecundation. We have seen that the material furnished by the male is the sperm; that afforded by the female an ovum. The most difficult topic of inquiry yet remains, - how the new individual results from their commixture? Of the nature of this mysterious process we are indeed profoundly ignorant; and if we could make any comparison between the extent of our ignorance on the different vital phenomena, we should be disposed to decide, that the function of generation is, perhaps, the least intelligible. The new being must be stamped instantaneously as by the die. From the very moment of the admixture of the materials, at a fecundating copulation, the embryo must have within it the powers necessary for its own

formation, and under impulses communicated by each parent—as regards likeness, hereditary predisposition, &c. From this moment the father has no communication with it; yet we know, that it will resemble him in its features and in its predispositions to certain morbid states,—whilst the mother probably exerts but a slight and indirect control over it afterwards, her office being chiefly to furnish the homunculus with a nidus, in which it may work its own formation, and with the necessary pabulum. We have seen, that even so early as the seventh and eighth day after fecundation, two projecting points—it has been asserted—are

Rigby, op. cit. p. 27, Philad. 1840. See, also, T. W. Jones, in Brit. and For. Med. Rev., Oct. 1843, p. 528.

observed in the ovum, which indicate the future situations of the brain and spinal marrow. Our want of acquaintance with the precise character of this impenetrable mystery will not, however, excuse us from passing over some of the ingenious hypotheses, that have been entertained on the subject. These have varied according to the views that have prevailed respecting the nature of the sperm; and to the opinions indulged regarding the matter furnished by the ovary. Drelincourt, who died in 1697, collected as many as two hundred and sixty hypotheses of generation; but they may all, perhaps, be classed under two, — the system of epi-

genesis and that of evolution.

1. Epigenesis. — According to this system, which is the most ancient of all, the new being is conceived to be built up of materials furnished by both sexes, the particles composing those materials having previously possessed the arrangement necessary for constituting it, or having suddenly received such arrangement. Still, it is requisite that these particles should have some controlling agent to regulate their affinity, different from any of the ordinary forces of matter; and hence a force has been imagined to exist, which has been termed cosmic, plastic, essential, nisus formativus -the Bildungstrie b of the Germans-force of formation, &c. Hippocratesb maintains, that each of the two sexes possesses two kinds of seed, formed by the superfluous nutriment, and by fluids constituted of materials proceeding from all parts of the body, and especially from the most essential, — the nervous. Of these two seeds, the stronger begets males, the weaker females. In the act of generation, these seeds become mixed in the uterus, and by the influence of the heat of that organ, they form the new individual - by a kind of animal crystallization - male or female, according to the predominance of the stronger or the weaker seed. Aristotlee thought, that it is not by seed that the female participates in generation, but by the menstrual blood. This blood he conceived to be the basis of the new individual, and the principles furnished by the male to communicate to it the vital movement, and to fashion it. Empedocles, Epicurus, and various other ancient physiologists, contended, that the male and female respectively contribute a seminal fluid, which equally co-operate in the generation and development of the fætus, and that it belongs to the male or female sex, or resembles more closely the father or the mother, according as the orgasm of the one or the other predominates, or is accompanied by a more copious discharge.d Lactantius, in quoting the views of Aristotle on generation, fancifully affirms, that the right side of the uterus is the proper chamber of

De Generatione Animalium, &c. i. 19.

<sup>\*</sup> Novem Libelli de Utero, Conceptione, Fœtû, &c., Lugd. Bat. 1632.

b περι χονες; in Oper. Omnia. edit. A. Foësio. Genev. 1657-1662.

d "Semper enim partus duplici de semine constat;

Atque utrique simile est magis id quodcumque creatur."

the male fœtus, and the left of the female, - a belief, which appears to be still prevalent amongst the vulgar, in many parts of Great Britain. But he adds, if the male or stronger semen should, by mistake, enter the left side of the uterus, a male child may still be conceived; yet as it occupies the female department, its voice, face, &c., will be effeminate. On the contrary, if the weaker or female seed should flow into the right side of the uterus, and a female fœtus be engendered, it will exhibit evidences of a masculine character. The idea of Aristotle, with regard to the menstrual blood, has met with few partisans, and is undeserving of notice. That of Hippocrates, notwithstanding the objections, which we now know to apply to it, - that the female furnishes no sperm, and that the ovaria are probably in no respect analogous to the testes of the male, - has had numerous supporters amongst the moderns, being modified to suit the scientific ideas of the time, and of the individual. Descartes, for example, considered the new being to arise from a kind of fermentation of the seed furnished by both sexes. Pascal, that the sperm of the male is acid, and that of the female alkaline; and that they combine to form the embryo. Maupertuisa maintained, that, in each seed, parts exist. adapted for the formation of every organ of the body, and that, at the time of the union of the seed in a fecundating copulation, each of the parts is properly attracted and aggregated by a kind of crystallization.b

The celebrated hypothesis of the eloquent but too enthusiastic Buffone is but a modification of the Hippocratic doctrine of epigenesis. According to him, there exists in nature two kinds of matter, - the living and the dead; the former perpetually changing during life, and consisting of an infinite number of small, incorruptible particles or primordial monads, which he called organic molecules. These molecules, by combining in greater or less quantity with dead matter, form all organized bodies; and without undergoing destruction, are incessantly passing from vegetables to animals, in the nutrition of the latter, and are returned from the animal to the vegetable by the death and putrefaction of the former. These organic molecules, during the period of growth, are appropriated to the development of the individual; but, as soon as he has acquired his full size, the superfluous molecules are sent into depot in the genital organs, each molecule being invested with the shape of the part sending it. In this way he conceived the seed of both sexes to be formed of molecules obtained from every part of the system. In the commixture of the seeds, during a fecundating copulation, the same force that assimilates the organic molecules to the parts of the body for their nourishment and increase, causes them, in this hypothesis, to congregate for the formation of the new individual; and, according as the molecules of

a Venus Physique, Paris, 1751.

Adelon, Physiologie de l'Homme, iv. 85, Paris, 1829.
 Histoire Naturelle, tom. xvii. &c., Paris, 1799.

The ingenuity of this doctrine was most captivating; and it appeared so well adapted for the explanation of many of the phenomena of generation, that it had numerous and respectable votaries. It accounted for the circumstance of procreation being impracticable, until the system had undergone its great development at puberty. It explained why excessive indulgence in venery occasions emaciation and exhaustion; and why, on the other hand, the castrated animal is disposed to obesity, — the depot having been removed by the mutilation. The resemblance of the child to one parent rather than to the other was supposed to be owing to the one furnishing a greater proportion of organic molecules than the other; and as more males than females are born, the circumstance was ascribed to the male being usually stronger, and therefore fur-

nishing a stronger seed, or more of it.

Prior to this hypothesis, Leeuenhoeka had discovered what he considered to be spermatic animalcules in the semen; but Buffon contested their animalcular nature, and regarded them as his vital particles or organic molecules; whilst he looked upon the ovarian vesicle as the capsule that contained the sperm of the female. The opinions of Buffon were slightly modified by Blumenbach, b and by Darwin.c The former, like Buffon, divided matter into two kinds, possessing properties essentially different from each other; the inorganic and the organized; the latter possessing a peculiar creative or formative effort, which he called Bildungstrieb or nisus formativus, - a principle in many respects resembling gravitation, and endowing every organ, as soon as it acquires structure, with a vita propria. This force he conceived to preside over the arrangement of the materials, furnished by the two sexes in generation. Darwin preferred to the term organic molecules that of vital germs, which he says are of two kinds, according as they are secreted or provided by male or female organs, whether animal or vegetable. In the subdivision, however, of the germs the term molecule is retained; but it is limited to those of the female; the vital germs or particles, secreted by the female organs of a bud or flower, or the female particles of an animal, being denominated by him molecules with formative propensities; whilst those secreted from the male organs are termed fibrils with formative appetencies. To the fibrils he assigns a higher degree of organization than to the molecules. Both, however, he asserts, have a propensity or appetency to form or create, and "they reciprocally stimulate and embrace each other and instantly coalesce; and may thus popularly be compared to the double affinities of chemistry."

Subtile as these hypotheses are, they are open to forcible objections of which a few only will suffice. The notion of this occult

<sup>\*</sup> Arcana Naturæ, Lugd. Bat. 1685.

b Ueber den Bildungstrieb, Gotting. 1791; Comment. Societat. Gotting. tom. viii.; and Efliotson's Blumenbach's Physiology, 4th edit. p. 490, Lond. 1828.

c Zoonomia, Lond. 1796.

force is identical with that, which, we shall see hereafter, has prevailed as regards life in general, and it leaves the subject in the same obscurity as ever. What do the terms plastic, cosmic, or vegetative force, or Bildungstrieb express, which is not equally conveyed by vital force, - that mysterious property, on which so many unfathomable processes of the animal body are dependent, and of the nature or essence of which we know absolutely nothing? The objection, urged against the doctrine of Hippocrates, - that we have no evidence of the existence of female sperm, - applies equally to the hypotheses that have been founded upon it; and even were we to grant, that the ovarium is a receptacle for female sperm, the idea, that such sperm is constituted of organic molecules, derived from every part of the body, would still be entirely gratuitous. We have no facts to demonstrate the affirmative; whilst there are many circumstances, that favour the negative. The individual, for example, who has lost some part of his person nose, eye, or ear, or has had a limb amputated, still begets perfect children; yet whence can the molecules, in such cases, have been obtained? It is true, that if the mutilation affect but one parent, the organic molecules of the lost part may still exist in the seed of the other; but we ought, at least, to expect the part to be less perfectly formed in the embryo, which it is not. Where two docked horses are made to engender, the result ought, à fortiori, to be imperfect, as the organic molecules of the tail could not be furnished by either parent, yet we find the colt, in such cases, perfect in this appendage. An elucidative case is also afforded by the fætus. If we admit the possibility of organic molecules constituting those parts that exist in the parents, how can we account for the formation of such as are peculiar to fœtal existence. Whence are the organic molecules of the navel-string, or of the umbilical vein, or of the ductus venosus, or the ductus arteriosus, or the umbilical arteries obtained? These and other objections have led to the abandonment of the theory of Buffon, which remains merely as a monument of the author's ingenuity and elevation of fancy.

2. Evolution. — According to this theory, the new individual pre-exists in some shape in one of the sexes, but requires to be vivified by the other, in the act of generation; after which it commences the series of developments or evolutions, which lead to the formation of an independent being. The great differences of sentiment, that have prevailed under this view, have been owing to the part, which each sex has been conceived to play in the function. Some have considered the germ to exist in the ovary, and to require the vivifying influence of the male sperm to cause its evolution. Others have conceived the male sperm to contain the rudiments of the new being, and the female to afford it merely a nidus, and pabulum during its development. The former class of physiologists have been called ovarists; — the latter sperma-

<sup>&</sup>lt;sup>2</sup> C. Windischmann, art. Evolutions Theorie, Encyclopäd. Wörterb. der Medicinisch. Wissensch. xi. 615, Berlin, 1834.

tists, seminists, and animalculists. The ovarists maintain, that the part furnished by the female is an ovum from the ovary; and this ovum they conceive to be formed of an embryo and of particular organs for the nutrition and first development of the embryo; and adapted for becoming, after a series of changes or evolutions, a being similar to the one whence it has emanated. The hypothesis was suggested by the fact, that in many animals but a single individual is necessary for reproduction; and it is easier, perhaps, to conceive this individual to be female than male; as well as by what is noticed in many oviparous animals. In these, the part, furnished by the female, is manifestly an ovum or egg; and in many, such egg is laid before the union of the sexes, and is fecundated, as we have seen, externally. By analogy, the inference was drawn, that this may happen to the viviparous animal also. The notion is said, but erroneously, to have been first of all advanced by Joseph de Aromatariis.4 It was developed by Harvey, b who strenuously maintained the doctrine omne vivum ex ovo. The anatomical examinations of Sylvius, Vesalius, Fallopius, De Graaf, Malpighi, Vallisnierie and others, - by showing, that what had been previously regarded as female testes, and had been so called, were organs containing minute vesicles or ova, and hence termed, by Steno, ovaria, - were strong confirmations of this view, and startling objections to the ancient theory of epigenesis; and the problem appeared to be demonstrated, when it was discovered, that the vesicle or ovum leaves the ovarium, and passes through the Fallopian tube to the

The chief arguments that have been adduced in favour of this doctrine are: - First. The difficulty of conceiving the formation, ab origine, of an organized body, as no one part can exist without the simultaneous existence of others. Secondly. The existence of the germ prior to fecundation in many living beings. In plants, for example, the grain exists in a rudimental state in the flower, before the pollen, which has to fecundate it, has attained maturity. In birds, too, the egg must pre-exist, as we find that those, which have never had intercourse with the male, can yet lay. This is more strikingly manifest in many fishes, and in the batracia or frog kind; where the egg is not fecundated until after ex-Spallanzani, moreover, asserts, that he could distinguish the presence of the tadpole in the unfecundated ova of the frog; and Haller, that of the chick in the infecund egg; at least he has seen them containing the yolk, which, in his view, is but a dependence of the intestine of the fœtus, and if the yolk exist, the chick exists also. Thirdly. The fact, before referred to, that in certain animals, a single copulation is capable of fecun-

a Epist. de Generatione Plantarum ex Seminibus, Venet. 1625.

b Exercitationes de Generatione Animalium, Lond. 1651.

De Organis Mulierum, &c. Lugd. Bat. 1672.
 Append. ad Opera Omnia, Lugd. Bat. 1687.

<sup>·</sup> Istoria della Generazione dell'Uomo, Discorsi Acad. i.-iv., Venez. 1722-1726.

vol. II. - 33

dating several successive generations. In these cases, it is argued, the germ of the different generations must have existed in the first. Fourthly. The fact of natural and accidental encasings, inclusions, or emboîtements; as in the bulb of the hyacinth, in which the rudiments of the flower are distinguishable; in the buds of trees, in which the branches, leaves, and flowers, have been detected in miniature, and greatly convoluted; in the jaws of certain animals, in which the germs of different series of teeth can be detected; in the volvox, a transparent animal, which exhibits several young ones encased in each other; in the common egg, which occasionally has another within it; and in the instances on record, in which human fœtuses have been found in the bodies of youths, of which there is a striking example in the Museum of the Royal College of Surgeons, of London; and a similar case in a boy of fourteen years of age, has been related by Dupuytren. A most singular case of the kind occurred to M. Velpeau. A tumour was removed from the scrotum of a young man aged 27, which was found to contain almost all the elements of a human body. Its exterior was evidently tegumentary, and the greater part of its substance was a mixture of lamellæ and fibres like cellular, adipose, muscular and fibrous tissues. In the interior, there were two cysts filled with a substance like albumen or the vitreous humour: another cyst, as large as a partridge's egg. contained a greenish semi-fluid matter like meconium; and a fourth contained a dirty yellow grumous mass surrounded by hairs: the mass consisted of sebaceous matter and scales of epidermis; the hairs had no bulbs. A tuft of hair, which protruded externally from a kind of ulcer at the posterior part, - and which, with the fact of the tumour being congenital, induced M. Velpeau to consider it to be fætal, - proceeded from the cyst that contained the meconium-like substance, and gave the opening into it somewhat the appearance of an anus. In the midst of all these, numerous perfectly organized portions of a skeleton were found, consisting of bones resembling more or less the clavicle, scapula, humerus, sphenoid bone, sacrum, portions of vertebræ, and others whose names could not be determined. A peculiarity of this case of monstrosity by inclusion was, that the second fœtus did not act as a foreign body in the other, but had a separate and independent existence and power of growth within itself. The tumour had its own colour and consistence, and a sensibility entirely independent of the person to whom it was attached. The man himself pierced it several times with a knife without feeling the least pain; and vet, all the wounds that were made in it bled, inflamed and cicatrized like those of any other part of the body. Perhaps, the explanation of these extraordinary cases by Dr. Blundellb is as philosophical as any that could be devised. A seed or egg, he re-

2 Gazette Médicale, Fév. 15, 1840.

<sup>&</sup>lt;sup>b</sup> Principles and Practice of Obstetricy, edited by Dr. Castle, Lond. 1834; American edition, Washington.

marks, though fecundated, may lie for years without being evolved. A serpent may become inclosed under the eggshell of the goose; the shell probably forming over it as the animal lies in the oviduct of the bird. These facts Dr. Blundell applies to the phenomenon in question. When the boy was begotten, a twin was begotten at the same time, - but, while the former underwent his development in the usual manner, the impregnated ovum of his companion lay dormant, and, unresistingly, became closed up within the fraternal structure, as the viper in the eggshell. For a few years, these living rudiments generally lie quiet within the body, and ultimately become developed so as to occasion the death of both. "The boy," he remarks - speaking of one of the cases - "became pregnant with his twin brother, his abdomen formed the receptacle, where, as in the nest of a bird, the formation was accomplished." Cases of this kind of arrest of development occasionally occur, where two or more ova are fecundated at the same time, or in succession. To this we shall refer under Superfætation." Fifthly. The fact of the various metamorphoses that take place in certain animals. Of these we have the most familiar instances in the batracia, and in insects. The forms which they have successively to assume are evidently encased. In the chrysalis, the outlines of the form of the future butterfly are apparent; and in the larva we observe those of the chrysalis. The frog is also apparent under the skin of the tadpole. Sixthly. The fact of artificial fecundation, which has been regarded, by the ovarists, as one of the strongest proofs of their theory; the quantity of sperm employed, as in the experi ments of Spallanzani, already detailed, being too small, in their opinion, to assist in the formation of the new individual, except as a vivifying material. Lastly. They invoke the circumstance of partial reproduction, of which all living bodies afford more or less manifest examples; - as the reproduction of the hair and nails in man; of the teeth in the rodentia; of the tail in the lizard; of the claw in the lobster; of the head in the snail, &c., &c. All these phenomena are, according to them, owing to each part possessing, within itself, germs destined for its reproduction, and requiring only favourable circumstances for their development. The partisans of the doctrine of epigenesis, however, consider these last facts as opposed to the views of the ovarists; and they maintain, that, in such cases, there is throughout a fresh formation.

The chief objections, that have been urged against the hypothesis of the ovarists, are: — First. The resemblance of the child to the father — a subject which we shall refer to presently. The ovarists cannot of course deny that such resemblance exists; and they ascribe it to the modifying influence exerted by the male sperm, but without being able to explain the nature of such influence. They affirm, however, that the likeness to the mother is more frequent and

a See a case of "Abdominal Enadelphia or Monstrosity by Inclusion," by M. Roux, du Var, with reflections by M. Geoffroy Saint-Hilaire, in Gazette Médicale de Paris, No. 35, Août 27, 1836; and Researches on Monstrosity by Inclusion in Animals, suggested by a case of the kind in a Human Fætus, by M. Charvet, in Archives Générales de Médecine, Nov. 1838.

evident. Certain cases of resemblance are, indeed, weighty stumbling blocks to ovism, or to the doctrine of a pre-existent germ in the female. It is a well-known fact, that six-fingered men will beget six-fingered children. How can we explain this upon the principle of the pre-existence of the germ in the female, and of the part played by the male sperm being simply that of a vivificative agent; and must we suppose, in the case of monstrosities, that such germs have been originally monstrous? Secondly. The production of hybrids is one of the strongest counter-arguments. They are produced by the union of the male and female of different species. Of these, the mule is the most familiar instance - the product of the ass and the mare. This strikingly participates in the qualities of both parents, and, consequently, the pre-existing germ in the female must have been more than vivified by the sexual intercourse. Its structure must have been altogether changed, and all the germs of its future offspring annihilated as the mule is seldom fertile. If a white woman marry a negro, the child is a mulatto, and if the successive generations of this woman be continually united to negroes, the progeny will ultimately become entirely black; or, at least the white admixture will escape recognition. As a general rule, the offspring of different races has an intermediate tint between that of the parents: and the proportions of white and black blood, in different admixtures, have even been subjected to calculation, in countries where negroes are common. The following table represents these proportions, according to the principles sanctioned by custom.

Parents.	Offspring.	Degree of Mixture.
Negro and white,	mulatto,	white, ½ black.
White and mulatto,	terceron,	4 - 4 -
Negro and mulatto,	Sgriffo or zambo, or black terceron,	1 - 1 -
White and terceron,	quarteron,	2 - 1 -
Negro and terceron,	black quarteron,	1 - 7 -
White and quarteron,	quinteron,	5 - 16 -
Negro and quarteron,	black quinteron,	18 - 15 -

The two last are considered to be respectively white and black; and of these the former were white by law, and consequently free, in the British West India Islands.<sup>a</sup> All these cases exhibit the influence exerted by the father upon the character of the offspring, and are great difficulties in the way of supposing that the male sperm is simply a vivifier of the germ pre-existing in the female.<sup>b</sup> Thirdly. The doctrine of the ovarists does not account for the greater degree of fertility of cultivated plants and of domesticated animals. Fourthly. The changes, induced by the succession of ages on the animal and vegetable species inhabiting the surface of

<sup>\*</sup> Lectures on Physiology, Zoology, and the Natural History of Man, p. 299, Lond. 1819.

b Rudolphi, Grundriss der Physiologie, u. s. w. B. i. s. 54, Berl. 1823; Burdach, Physiologie als Erfahrungswissenschaft, u. s. w., B. i. s. 516; and Berndt, art. Bastardthiere in Enc. Wörterb. u. s. w., B. v. s. 53, Berl. 1830.

the globe, have been adduced against this hypothesis. In examining the geological character of the various strata that compose the earth, it has been observed by geologists that many of these contain imbedded the fossil remains of animals and vegetables Now, those rocks on which others rest are the oldest, and the successive strata above these are more and more modern, and it has been found that the organic fossil remains in the different strata differ more and more from the present inhabitants of the surface of the globe in proportion to the depth we descend; and that the remains of those beings, that have always been the companions of man, are found only in the most recent of the alluvial

deposits, - in the upper crust of the earth.

In the older rocks the impressions are chiefly of the less perfect plants - as the ferns and reeds; and of the lower animals - the remains of shells and corals; whilst fish are uncommon. In the more recent strata, the remains of reptiles, birds and quadrupeds are apparent, but all of them differ essentially from the existing kinds, and in none of the formations of more ancient date has the fossil human skeleton been met with. The pretended human hones, conveyed by Spallanzani from the Island of Cerigo the ancient Cythera — are not those of the human species any more than the bones of the Homo diluvii testis of Scheuchzer; and the skeleton of the savage Galibi, conveyed from Guadaloupe and deposited in the British Museum, is imbedded in a calcareous earth of modern formation. Hence it has been concluded, that man is of a date posterior to animals in all countries where fossil bones have been discovered. These singular facts, furnished by modern geological inquiry, have been attempted to be explained by the supposition, that the present races of animals are the descendants of those whose remains are met with in the rocks, and that their difference of character may have arisen from some change in the physical constitution of the atmosphere, or of the surface of the earth, producing a corresponding change in the forms of organized beings. It has been properly remarked, however, by Dr. Fleming, that the effect of circumstances on the appearance of living beings is circumscribed within certain limits, so that no transmutation of species was ever ascertained to have taken place, whilst the fossil species differ as much from the recent kinds, as the last do from each other; and he adds, that it remains for the abettors of the opinion to connect the extinct with the living races, by ascertaining the intermediate links or transitions. will probably ever be impracticable. The difference, indeed, between the extinct and the living races is in several cases so extreme, that many naturalists have preferred believing in the occasional formation of new organized beings. Linnæus was bold enough to affirm, that, in his time, more species of vegetables were in existence than in antiquity, and hence, that new vegetable species

33\*

Principles of Geology, by Charles Lyell, Esq., F. R. S. i. 241, 4th edit. Lond. 1835.
 Philosophy of Zoology, i. 26, Edinb. 1822.

must necessarily have been ushered into being; and Wildenow embraced the views of Linnæus. De Larmack, a one of the most distinguished naturalists of the day, openly professes his belief, that both animals and vegetables are incessantly changing under the influence of climate, food, domestication, the crossing of breeds, &c., and he remarks, that if the species now in existence appear to us fixed in their characters, it is because the circumstances that modify those species demand an enormous time for action, and would consequently require numerous generations to establish the fact. The manifest effect of climate, food, &c., on vegetables and animals, he thinks, precludes the possibility of denying those changes on theoretical considerations; and what we call lost species are, in his view, only the actual species before they experienced modification. It is proper, however, to observe, that the representations on the wall of one of the sepulchres in the valley of Beban el Molook, at Thebes, which are regarded by Champollion as having been executed upwards of two thousand years before the Christian era, enable the features of the Jew and of the negro, amongst others, to be recognised as easily as the representations of their descendants of the present day; so that, for the space of at least three thousand eight hundred years, no modification of the kind referred to by Lamarck seems to have occurred in the human

Another explanation has been offered for these geological facts, and for the rotation, which we observe in the vegetable occupants of particular soils in successive years. It has been supposed, that as the seeds of plants and the ova of certain animals are so excessively minute as to penetrate wherever water or air can enter; and as they are capable of retaining the vital principle for an indefinite length of time, of which we have many proofs, and of undergoing evolution whenever circumstances are favourable, the crust of the earth may be regarded as a receptacle of germs, each of which is ready to expand into vegetable or animal forms, on the occurrence of conditions necessary for their development. This is the hypothesis of panspermia or dissemination of germs, according to which the germs of the ferns and reeds were first expanded, and afterwards those of the staminiferous or more perfect vegetables; and, in the animal kingdom, first the zoophyte, and gradually the being more elevated in the scale; the organized bodies of the first period flourishing, so long as the circumstances, favourable to their development, continued, and then making way for the evolution of their successors, - the changes effected in the soil by the growth and decay of the former probably favouring the evolution of the latter; which, again, retained possession of the soil so long as circumstances were propitious.b The changes

<sup>&</sup>lt;sup>a</sup> Philosophie Zoologique, edit. cit. tom. i.; and Lyell's Principles of Geology, ii. 407.

<sup>&</sup>lt;sup>b</sup> Fleming, op. citat. i. 28; and Purkinje, art. Erzeugung, in Encycl. Wörterb. der Medicin. Wissensch. xi. 537, Berlin, 1834.

that take place in forest vegetation are favourable to this doctrine. If, in Virginia, the forest trees be removed so as to make way for other growth, and the ground be prepared for the first cultivation, the Phytolacca decandra or poke, which was not previously perceptible on the land, usurps the whole surface. When Mr. Madison went with Gen. Lafayette to the Indian treaty, they discovered, that wherever trees had been blown down by a hurricane, in the spring, the white clover had sprung up in abundance, although the spot was many miles distant from any cleared land: and it has often been remarked, that where, during a drought in the spring, the woods have taken fire and the surface of the ground has been torrefied, the water-weed has made its appearance in immense quantities, and occupied the burnt surface. The late Judge Peters, having occasion to cut ditches on his land, in the western part of Pennsylvania, was surprised to find every subterraneous tree that was met with, different from those at the time occupying the surface; and President Madison informed the author. that in the space of sixty or seventy years, he had noticed the following spontaneous rotation of vegetables: - 1. Mayweed: 2. Blue centaury; 3. Bottle-brush-grass; 4. Broomstraw; 5. White clover; 6. Wild carrot; and the last is now giving way to the blue grass.a

The doctrine of panspermia is, however, totally inapplicable to the viviparous animal, in which the ovum is hatched within the body, and which, consequently, continues to live after the birth of its progeny; and the facts furnished by geology seem clearly to show, that the development of the animal kingdom has been successive, not simultaneous; but, under what circumstances the different animals were successively ushered into being, we know not.

Lustly, as regards the ovarists themselves; - they differ in essential points: whilst some are favourable to the doctrine of the dissemination of germs, believing, as we have seen, that ova or germs are disseminated over all space, and that they only undergo development under favourable circumstances, as when they meet with bodies capable of retaining them, and causing their growth, or which resemble themselves; others assert, that the germs are inclosed in each other, and that they are successively aroused from their torpor, and called into life, by the influence of the seminal fluid; so that not only did the ovary of the first female contain the ova of all the children she had, but one only of these ova contained the whole of the human race. This was the celebrated system of emboîtement des germes or encasing of germs, of which Bonnet was the propounder, and Spallanzani the promulgator. Yet how monstrous for us to believe, that the first female had, within her, the germs of all mankind, born, and to be born; or to conceive, that a grain of Indian corn contains within it all the seed,

b Considerat. sur les Corps. Organises vol. i. and ii., Amst. 1762.

<sup>&</sup>lt;sup>a</sup> Prichard, Researches on the Physical History of Man, i. 39; and Carpenter, Principles of General and Comparative Physiology, 2d edit. Lond. 1841.

that may hereafter result from its culture. In this strange hypothesis — as Professor Elliotson<sup>a</sup> observes — there must have been an uncommon store of germs prepared at the beginning, for the ovaria of a single sturgeon have contained 1,467,500 ova.<sup>b</sup> Many of the ovarists, again, and they alone have any thing like probability in their favour, believe, that the female forms her own ova, as the male forms his own sperm, by a secretory action; and; so far as the female is concerned in the generative process, we shall find that this is the only philosophical view; but it is imperfect in not admitting of more than a vivifying action in the

materials furnished by the male.

About the middle of the seventeenth century, Hamme or Van Hammen, Leeuenhoeke and Hartsöker, discovered a prodigious number of small moving bodies in the sperm of animals, which they regarded as animalcules.e This gave rise to a new system of generation, directly the reverse of that of Harvey, - that of generation ab animalculo maris. As, in the Harveian doctrine, the germ was conceived to be furnished by the mother and the vivifying influence to be alone exerted by the male, so, in this doctrine, the entire formation was regarded as the work of the father, the mother affording nothing more than a nidus, and appropriate pabulum for the homunculus or rudimental fœtus.f The spermatic doctrine was soon embraced by Boerhaave, Keill, Cheyne, Wolff, Lieutaud, and others.8 The pre-existing germ was accordingly now referred exclusively to the male; and, by some, the doctrine of emboîtement or encasing was extended to it. In support of this hypothesis, the spermatists urged, - that the animalcules they discovered were peculiar to the semen, and that they exist in the sperm of all animals capable of generation; that they differ in different species, but are always identical in the sperm of the same animal, and in that of individuals of the same species; that they are not perceptible in the sperm of any animal until the age at which generation is practicable, whilst they are wanting in infancy and decrepitude; that their number is so considerable, that a drop of the sperm of a cock, scarcely equal in size to a grain of sand, contains 50,000; and lastly, that their minute size is no obstacle to the supposition, that generation is accomplished by them ; - the disproportion between the trees of our forest and the seed producing them being nearly if not entirely as great as that between the animalcule and the being it has to develope. Leeuenhoek estimated the dimensions of those of the frog at about the 1-10,000th

Petit, Mémoir. de l'Académ. des Sciences, 1733.
 Oper. iii. 285, and iv. 169, Lugd. Bat. 1722.

a Blumenbach's Physiology, by Elliotson, 4th edit. p. 494, Lond. 1828.

d Journal des Sçavans, pour 1678; and Essai de Dioptrique, p. 227, Paris, 1694. Haller, Element. Physiol. vol. vii. 27; Rudolphi, art. Animalcula Seminalia, in Encyclopäd. Wörterb. der Medecin. Wissenschaft, ii. 597, Berlin, 1828; and Sprengel, Histoire de Médecine, par Jourdan, iv. 309, Paris, 1815.

Mohrenheim, Nova Conceptionis atque Generationis Theoria. Regiom. 1794. Bostock's Physiology, 3d edit. p. 642, Lond. 1836.

part of a human hair, and that the milt of a cod may contain 15,000,000,000,000,000 of them. The difficulty with the spermatists or animalculists was to determine the mode in which the homunculus attains the ovary, and effects the work of reproduction. Whilst some asserted it to be only requisite, that the sperm should attain the uterus, whither it attracted the ovum from the ovarium; others imagined, that the animalcule travelled along the Fallopian tube to the ovary; entered one of the ovarian vesicles; shut itself up there for some time, and then returned into the cavity of the uterus, to undergo its first development, through the medium of the nutritive substance contained in the vesicle: and a celebrated pupil of Leeuenhoek even affirmed, that he not only saw these animalcules under the shape of the tadpole, as they were generally described, but that he could trace one of them, bursting through the envelope that retained it, and exhibiting two arms, two legs, a human head and a heart !a

Although this doctrine was extremely captivating, and, for a time, kept the minds of many eminent philosophers in a state of delusive enthusiasm; insomuch that Dr. Thomas Morgan, in a work, published in 1731, thus expresses himself regarding it : - " That all generation is from an animalculum pre-existing in semine maris, is so evident in fact, and so well confirmed by experience and observation, that I know of no learned men, who in the least doubt of it;" it was, subsequently, strongly objected to by many; and the great fact on which it rested - the very existence of the spermatic animalcules - was, and is, strenuously contested. Linnæuse discredited the observations of Leeuenhoek. Verheyen denied the existence of the animalcules, and undertook to demonstrate, that the motion, supposed to be traced in them, was a mere microscopic delusion : - whilst Needhamd and Buffon regarded them as organic molecules.e Of late years, MM. Prévost and Dumas have directed their attention to the subject; and their investigations, as on every other topic of physiological inquiry, are worthy of the deepest regard. The results of their examinations led them to confirm the existence of these animalcules, and likewise to consider them as the direct agents of fecundation. By means of the microscope they detected them in all the animals, whose sperm they examined, and these were numerous. Whether the fluid were observed after its excretion by a living animal, or after its death, in the vas deferens or in the testicle, the animalcules were detected in it with equal facility. They consider these bodies to be characteristic of the sperm, as they found them only in that secretion; being wanting in every other humour of the body, even in those

<sup>a</sup> Adelon, Physiologie de l'Homme, edit. cit. iv. 94.

b Mechanical Practice of Physic, Lond. 1735.

d New Microscopical Discoveries, Lond. 1745.

e Haller, Element. Physiologiæ, xxvii.

e Bostock's Physiology, 3d edit. p. 643, London, 1836.

<sup>&#</sup>x27; Mêm. de la Société Physique de Genève, i. 180, and Annales des Sciences Naturelles, tom. i. and ji.

that are excreted with the sperm, as the fluids of the prostate, and of the glands of Cowper; and although similar in shape, and size, and in the character of their locomotion in the individuals of the same species, they are of various shapes and dimensions in different species. In passing through the series of genital organs these animalcules experience no change, being as perfect in the testicle as at the time of their excretion; and they controvert the remark of Leeuenhoek, that they are met with apparently of different ages. The animalcules were manifestly endowed with spontaneous motion, which gradually ceased, - in the sperm obtained during life by ejaculation, in the course of two or three hours; in that taken from the vessels after death, in fifteen or twenty minutes, and in eighteen or twenty hours, when left in its own vessels after death. In farther proof of the position, that these animalcules are the fecundating agents, MM. Prévost and Dumas assert, that they are only met with whilst reproduction is practicable: - that, in the human species, they are not found in infancy or decrepitude; and, in the majority of birds, are only apparent in the sperm, at the periods fixed for their copulation; facts which, in their opinion, show, that they are not mere infusory animalcules. MM. Prévost and Dumas moreover affirm, that they appear to be connected with the physiological condition of the animal furnishing them; their motions being rapid or languishing, according as the animal was young or old, or in a state of health or disease. They state, also, that in their experiments on the ova of the mammiferous animal, they observed animalcules filling the cornua of the uterus, and remaining there alive and moving, until the ovule descended into that organ, when they gradually disappeared; and they argue in favour of the influence of these animalcules, - that the positive contact of the sperm is necessary for fecundation, and that the aura seminis is totally insufficient; that the sperm, in twenty-four hours, loses its fecundating property, and it requires about this length of time for the animalcules to gradually cease their movements and perish; and, lastly, that having destroyed the animalcules in the sperm, the fluid lost its fecundating property. One of these experiments consisted in killing all the animalcules in a spermatised fluid,whose fecundating power had been previously tested, - by repeated discharges of a Leyden phial: another consisted in placing a spermatized fluid on a quintuple filter, and repeating this until all the animalcules were retained on the filter; when it was found, that the fluid, which passed through, had no fecundating power, whilst the portion retained by the filter had the full faculty; a result that had been obtained by Spallanzani, who found, besides, that he was capable of effecting fecundation with water in which the papers, used as filters, had been washed. M. Donnéa has investigated the mode in which the zoospermes are affected in the blood, the milk, the vaginal and uterine mucus in the healthy state, in the purulent matter of chancres, and of blennorrhæa, in the 2 Gazette Médicale de Paris, No. xxii., 3 Juin, 1837.

saliva, urine, &c. He observed these animalcules continue to live, and to move in some of those fluids, whilst in others they died instantaneously. For instance, the blood, the milk and pus did not affect them: in the mucus of the vagina and uterus they generally lived well; but in the saliva and urine they died almost instantaneously. M. Donné, too, affirms, that there are cases in which the mucus of the vagina and uterus acquires properties that are deleterious to the zoospermes, and he is of opinion, that this is a cause of sterility. This deleterious property, according to M. Donné, occasionally resides in the vaginal mucus; but at others, in a still higher degree in the mucus of the uterus; and he endeavoured to discover, whether the mucus of the two membranes presented any peculiar characters or signs of disease, and he affirms. that he particularly noticed the excessive acidity of the one, and the marked alkaline character of the other. The mucus secreted by the vagina as far as the orifice of the os uteri differed from that which flowed from within the cervix uteri, independently of its physical characters, by a different reaction, M. Donné found the vaginal mucus always acid — the uterine always alkaline, and he thinks, that the deleterious influence exerted by them on the zoospermes depended on excess of acidity in the one, and excess of alkali in the other. All this, however, it need scarcely be said, requires substantiation. Professor Wagner, a who has entered at great length into the consideration of the spermatozoa, accords with the general conclusions of M. Donné: some of his experiments, however, instituted for the most part on the spermatozoa of the lower animals, led him to different conclusions. He found, for example, that they almost always lived in saliva; also in urine when it was kept warm and was not too concentrated. He has repeatedly detected them in the urine of persons, whom he suspected of masturbation. Dr. John Davyb affirms, that on examining the fluid from the urethra after stool in a healthy man, he had always detected spermatozoa in it; and Dr. Robt. Willise asserts, that under the same circumstances, and even after the mere evacuation of the bladder, he has several times discovered spermatozoa in the fluid of the urethra; but the subjects of his observation were never strong or healthy men; they mostly laboured under anomalous nervous symptoms, which, he thinks, were in all likelihood connected with an irritable or disordered state of the vesiculæ seminales and prostate part of the urethra.

MM. Prévost and Dumas, and Rolando, conjecture, that the spermatic animalcule forms the nervous system of the new being, and that the ovule furnishes only the cellular framework in which the organs are formed; but this is mere hypothesis. The essays of these ingenious experimenters would seem to prove the existence of peculiar animalcules in the sperm, and their apparent agency

<sup>\*</sup> Elements of Physiology, translated by Dr. Willis, p. 20, Lond. 1841.

b Edinburgh Medical and Surgical Journal, vol. i.

Wagner, op. citat. p. 21, (note.)

in the generative process; yet, as we have before seen, (page 330 of this volume,) all this has been questioned, and Raspaila is disposed to regard the fancied animalcules as mere shreds of the tissues of the generative organs ejaculated with the sperm.

It is scarcely necessary to remark, that all the objections, which were urged against the system of the ovarists, as regards the proof in favour of an active participation of both sexes in the work of reproduction, are equally applicable to the views of those animal-culists, who refer generation exclusively to the spermatic ani-

malcule.b

Such are the chief theories that have been propounded on the subject of generation. It has been already observed, that the particular modifiations are almost innumerable. They may all, however, be classed with more or less consanguinity under some of the doctrines enumerated. Facts and arguments are strongly against any view that refers the whole process of formation to either sex. There must be a union of materials furnished by both, otherwise it is impossible to explain the similarity in conformation to both parents, which is often so manifest. Accordingly, this modified view of epigenesis is now adopted by most physiologists: - that at a fecundating copulation, the secretion of the male is united to a material, furnished by the ovarium of the female; that from the union of these elements the embryo results, impressed, from the very instant of such union, with life, and with an impulse to a greater or less resemblance to this or that parent, as the case may be; and that the material, furnished by the female, is as much a secretion resulting from the peculiar organization of the ovarium. as the sperm is from that of the testicle, - life being susceptible, in this manner, of communication from father to child, without a necessity for invoking the incomprehensible and revolting doctrine of the pre-existence of germs. This admixture of the materials furnished by both sexes accounts for the likeness that the child may bear to either parent, whatever may be the difficulty in understanding the precise mode in which they act in the formation of the fœtus. It has been attempted, however, by some, to maintain, that the influence of the maternal imagination during a fecundating copulation may be sufficient to impress the germ within her with the necessary impulse; and the plea has been occasionally urged in courts of justice. Of this we have an example in a well-known case, tried in New York, between 30 and 40 years ago. A mulatto woman was delivered of a female bastard child, which became chargeable to the authorities of the city. When interrogated, she stated that a black man of the name of Whistelo was the father, who was accordingly apprehended, for the purpose of being assessed with the expenses. Several physicians, who were summoned before the magistrates, gave it as their opinion that it was not his child, but the offspring of a white man. Dr. S.

2 Chimie Organique, p. 389, Paris, 1833.

b Burdach's Physiologie als Erfahrungswissenschaft, 2te Auflage, i. 112, Leipz. 1835.

Mitchill, however, who, according to Dr. Beck, seemed to be a believer in the influence of the imagination over the fætus, thought it probable that the negro was the father. Owing to this difference of sentiment, the case was carried before the mayor, recorder, and several aldermen. It appeared in evidence, that the colour of the child was somewhat dark, but lighter than the generality of mulattos, and that its hair was straight, and had none of the peculiarities of the negro race. The court very properly decided in favour of Whistelo, and of course against the testimony of Dr. Mitchill, who, moreover, maintained, that as alteration of complexion has occasionally been noticed in the human subject, -as of negroes turning partly white, -and in animals, so this might be a parallel instance.a The opinion does not seem entitled to much greater estimation than that of the poor Irishwoman, in a London police report, who ascribed the fact of her having brought forth a thick-lipped, woolly-headed urchin to her having eaten some black potatoes during her pregnancy! It is obvious, that the effect of the maternal imagination can only be invoked — by those who believe in its agency on the future appearance of the fætus — in the case of those animals in which copulation is a part of the process. Where the eggs are first extruded and then fecundated, all such influence must be out of the question; and even in the viviparous animal, we have seen, experiments on artificial impregnation have shown, that not only has the bitch been fecundated by sperm injected into the vagina, but that the resulting young have manifestly resembled the dog, whence the sperm had been obtained. But the strongest case in favour of the influence of the maternal imagination is given by Sir Everard Home. An English mare was covered by a quaga, — a species of wild ass from Africa, which is marked somewhat like the zebra. This happened in the year 1815, in the park of Earl Morton, in Scotland. The mare was only covered once; went eleven months, four days, and nineteen hours, and the produce was a hybrid, marked like the father. The hybrid remained with the dam for four months, when it was weaned and removed from her sight. She probably saw it again in the early part of 1816, but never afterwards. In February, 1817, she was covered by an Arabian horse, and had her first foal - a filly. In May, 1818, she was covered again by the same horse, and had a second. In June, 1819, she was covered again, but this year missed; but in May, 1821, she was covered a fourth time, and had a third; all being marked like the quaga. Similar facts have been alluded to by other writers. Hallerd remarks, that the female organs of the mare seem to be corrupted by the unequal copulation with

<sup>&</sup>lt;sup>a</sup> Beck's Medical Jurisprudence, 6th edit. i. 500, Philad. 1838. See, also, for some ridiculous stories of this kind, Demangeon, Du Pouvoir de l'Imagination sur de Physique et le Moral de l'Homme, p. 201, Paris, 1834.

b See page 360 of this volume.

c Philosoph. Transact. for 1821, p. 21; and Lectures on Comparative Anatomy, iii. 307.

d Element. Physiol. viii. 104.

the ass, as the young foal of a horse from a mare, which previously had a mule by an ass, has something asinine in the form of its mouth and lips; and Bechera says, that when a mare has had a mule by an ass, and afterwards a foal by a horse, there are evidently marks, in the foal, of the mother having retained some ideas of her former paramour, - the ass; whence such horses are commended on account of their tolerance and other similar qualities. It has been even affirmed, that the human female, when twice married, bears children occasionally to the second husband, which resemble the first both in bodily structure and mental powers.b The mode in which the influence is exerted, in this and similar cases, is unfathomable; and the fact itself, although indisputable, is astounding. Sir Everard Homee thinks, that it is one of the strongest proofs of the effect of the mind of the mother upon her young that has ever been recorded. Although we are totally incapable of suggesting any satisfactory solution, it appears to us more probable, that the impression must have been made in these cases on the genital system, and probably upon the ovarian vesicle,

rather than upon the mind of the animal.d

d. Conception. - Conception usually occurs without the slightest consciousness on the part of the female; and hence the difficulty of reckoning the precise period of gestation. Certain signs, as shivering, pains about the umbilicus, &c., are said to have occasionally denoted its occurrence; but these are rare exceptions, and the indications afforded by one are often extremely different from those presented by another. In those animals, in which generation is only accomplished during a period of generative excitement, the period of conception can be determined with accuracy; for, in by far the majority of such cases, a single copulation will fecundate, — the existence of the state of heat indicating, that the generative organs are ripe for conception. In the human female, where the sexual intercourse can take place at all periods of the year, conception is by no means as likely to follow a single intercourse: for, although she may be always susceptible of fecundation, her genital organs are perhaps at no one time so powerfully excited as in the animal during the season of love. It is not for the physiologist to inquire into the morbid causes of sterility in either male or female; nor is it desirable to relate all the visionary notions which have prevailed regarding the circumstances that favour conception. It would certainly seem more likely to supervene when the venereal orgasm occurs simultaneously in both parties; and

b See art. Generation, by Dr. Allen Thomson, Cyclop. Anat. and Physiol. P. xiii. p. 468, for Feb. 1838. Lectures, &c. iii. 308.

a Physic. Subterran. Lips. 1703.

Sprengel's Hist. de la Médecine, Jourdan's translation, i. 231, Paris, 1815; Dr. A. Thomson, op. cit. p. 427; Adelon, Physiologie de l'Homme, 2de édit. p. 81, Paris, 1829; Meigs, Philadelphia Practice of Midwifery, p. 83, Philad. 1838; and Seiler, art. Erzeugung, in Pierer, Anat. Phys. Wörterb. ii. 802, Leipz. und Altenb. 1818; and Breschet, in l'Art de Procréer les Sexes à Volonté, par Millot, Paris, 1829.

when the sperm is thrown well forwards towards the mouth of the uterus. We have already shown, that preternatural openings of the urethra, which interfere with this projection of the sperm in

the proper direction, render fecundation less probable.

It has been generally affirmed by writers, that conception is apt to take place more readily immediately after menstruation; either, it has been imagined, because the uterus continues slightly open, so as to admit the sperm more easily into its cavity, or because the whole apparatus is in a state of some excitement. This opinion is problematical; and, accordingly, a female is in the habit of reckoning from a fortnight after her last menstrual period; for as she might have fallen with child immediately after one menstruation, or not until immediately preceding the following menstruation, a difference of three weeks might occur; and she, therefore, takes the middle point between those periods; that is, ten days or a fortnight after her last menstruation, or, what is the same thing, ten days or a fortnight before the first obstructed menstruation. Sir Everard Home, a however, differs on this topic from the generality of physiologists, — affirming that, in the human species, the fulness of the vessels of the womb, prior to menstruation, corresponds with the state of heat in the female quadruped, and shows that, at that period, the ova are most commonly fit for impregnation. "The females in India," he observes, "where from the warmth of the climate, all the internal economy respecting the propagation of the species goes on more kindly than in changeable climates, reckon ten months as the period of utero-gestation. In the Apocrypha, the wisdom of Solomon, chap. vii. v. 2, - 'And in my mother's womb was fashioned to be flesh in the time of ten months.' This circumstance seems to prove, that immediately before menstruation, when all the appendages of the womb are loaded with blood, the ova and ovaria are more frequently ready for impregnation in the climates most congenial for propagation; and therefore the mode of reckoning is from the previous menstruation, which is ten months before the birth." Dr. Carpenter thinks it is "quite certain" that there is a greater aptitude for conception immediately before and after menstruation, than there is at any intermediate period. There is no subject, however, which is less certain; Raciborski states, that of fifteen women, who specified accurately the period of their last menstruation, as well as that of their last sexual commerce, conception took place in five from two to four days before the period at which the catamenia were due. In seven, it dated from a union two or three days after menstruation; in two, it took place during the period: and in one only as long as ten days after. Nor is the author prepared to agree with Dr. Carpenter, that there is good reason to believe, that in women the sexual feeling becomes stronger at the menstrual period.c

\* Lectures on Comparative Anatomy, iii. 297.

These were of course lunar months; although Sir Everard clearly does not think so. Carpenter, Human Physiology, Amer. Edit. by Dr. Clymer, § 742, Philad. 1843.

It has been attempted to ascertain what age and season are most prolific. From a register kept by Dr. Bland, of London, it would appear, that more women between the ages of twenty-six and thirty years, bear children than at any other period. Of two thousand one hundred and two women delivered, eighty-five were from fifteen to twenty years of age; five hundred and seventy-eight from twenty-one to twenty-five; six hundred and ninety-nine from twenty-six to thirty; four hundred and seven from thirty-one to thirty-five; two hundred and ninety-one from thirty-six to forty; thirty-six from forty-one to forty-five; and six from forty-six to forty-nine. At Marseilles, according to Raymond, women conceive most readily in autumn, and chiefly in October; next in summer; and lastly in winter and spring, - the month of March having Morand says, that July, May, June, and fewest conceptions. August, are the most frequent months for conception; and November, March, April, and October, successively, the least frequent. At the Havana, according to tables by the author's friend, Don Ramon de la Sagra, the monthly number of births, amongst the white population, during a period of five years, - from 1825 to 1829 inclusive - was in the following order: - October, September, November, December, August, July, June, April, May, January, March, and February. February, January, March and April are, therefore, the most frequent months for conception at the Havana - June, July, May and September the least so. Mr. Burnsb asserts, that the register for ten years of an extensive parish in Glasgow renders it probable that August and September are most favourable for conception. M. Villermé, from an estimate founded on eight years' observations in France, comprising 7,651,437 births, makes the ratio of conceptions as follows; - May, June, April, July, February, March and December, January, August, November, September and October: - and lastly, Dr. Gouverneur Emerson, who has employed himself most profitably on the Medical Statistics of Philadelphia, has furnished a table of the number of births, during each month, for the ten years ending in 1830; according to which, the numbers are in the following order: December, September, January, March, October, August, November, February, July, May, April and June, - the greatest number of conceptions occurring, consequently, in April, January, and May, - the least in October, August and September.

The human female is uniparous, — one ovum only, as a general rule, being fecundated: numerous other animals are multiparous, or bring forth many at a birth. The law, however, on this subject is not fixed. Occasionally, the human female will bring forth twins, triplets, or quadruplets, whilst the multiparous animal is not always delivered of the same number. It is impossible to account for those differences. The ovarists refer them to the female; the

<sup>Historio Economico-Politica y Estadistica de la Isla de Cuba. Habana, 1833.
Principles of Midwifery, 3d edit. p. 126, Lond. 1814.</sup> 

<sup>.</sup> Amer. Journ. of the Med. Sciences, for Nov. 1831.

animalculists to the male; and facts have been found to support both views. Certain females, who have been frequently married, have been multiparous with each husband; and analogous facts have occurred to males under similar circumstances. Ménage cites the case of a man, whose wife brought him twenty-one children in seven deliveries; and the same individual having impregnated his servant-maid, she brought forth triplets likewise. In 1755, it is asserted, a peasant was presented to the Empress of Russia, who was seventy years of age, and had been twice married. His first wife had fifty-seven children at twenty-one births. In four deliveries she had four children at each; in seven, three; and in six, two. This appears to be the ne plus ultra of such cases! In the Hospice de la Maternité, of Paris, it has been observed, that twins occur once in about eighty cases. In the Westminster Hospital, the same ratio has been found to prevail. In the British Lying-in-Hospital, the proportion was not greater than 1 in 91; whilst in the Dublin Lying-in Hospital, the cases were nearly twice as frequent, or about 1 in 57. Dr. Collinsa properly remarks on the singular circumstance, that in Ireland, the proportional number of women giving birth to twins is nearly a third greater than in any other country of which he had been able to obtain authentic records. He states the proportion in France to be one in every 95 births; in Germany, one in 80; in England, one in 92; in Scotland, one in 95; and in Ireland, one in 62. Of 129,172 women delivered in the Lying-in Hospital, Dublin, 2062 gave birth to twins; 29 produced three at a birth, which is in the proportion of one in 4450; and one only gave birth to four. In this country, the average, according to Dr. Dewees, is about 1 in 75. Triplet cases were found to occur in the Hospice de la Maternité, of Paris, about once in 9000 times; and in the Dublin Hospital once in 5050 times; the balance being largely in favour of the prolific powers of the Irish. Dr. Dewees affirms, that in more than 9000 cases, he has not met with an instance of triplets. Of 36,000 cases in the Hospice de la Maternité not one brought forth four children; yet there are cases on record where five have been born at a birth. Beyond this number the tales of authors ought perhaps to be esteemed fabulous.

In referring to the following table, it will be found to prevail, as a general rule, amongst quadrupeds, that the largest and most formidable bring forth the fewest young, and that the lower tribes are unusually fruitful; the number produced compensating, in some measure, for their natural feebleness, which renders them constantly liable to destruction. On the other hand, were the larger species to be as prolific as the smaller, the latter would

a A Practical Treatise on Midwifery, Lond. 1835: republished in Bell's Select Library, Philad. 1838.

b Dr. Garthshore, in Philos. Transact. for 1787, Kleinert's Repertorium, cited in Amer. Journ. of the Med. Sciences, Feb. 1838, p. 459; also, Bulletino delle Scienze Mediche, Agosto e Settembre, 1838, cited in Brit. and For. Med. Rev., Oct. 1839, p. 564.

soon be blotted from existence. What would have been the condition of animated nature, if the gigantic mastodon, once the inhabitant of our plains, could have engendered as frequently and as numerously as the rabbit. For wise purposes, it has also been ordained, that the more formidable animals seldom begin the work of reproduction until they have nearly attained their full size; whilst those that bring forth many commence much earlier. Lastly, there is some correspondence, likewise, between the duration of gestation and the size of the animal.

Animals.	Duration of ges- tation.			Duration of ges- tation.	Number of young.	
Ape	about 9 months,	1	Lioness -		4 or 5	
Bat \		2	Tigress -		4 or 5	
Rat	5 or 6 weeks,	5 or 6	Cat	8 weeks,	4 or 5	
Mouse	told to the state of	6 to 10	Seal		2	
Hare	30 days,	4 or 5		11 months)	111233	
Rabbit	Do.	Do.	Mare	and some	1	
Guinea-pig	3 weeks,	5 to 12	TOTAL CO. L. C.	days,		
Squirrel -	6 weeks,	4 or 5	Ewe	5 months,	1 or 2	
Mole		4 or 5	Goat	44 months,	1, 2, or 3	
Bear		2 or 3	Cow	9 months,	1 or 2	
Otter	9 weeks,	4 or 5	Reindeer -	8 months,	2	
Bitch	9 weeks,	4 to 10	Hind	Do.	1 or 2	
Ferret	6 weeks,	6 or 7			6 to 12	
Wolf	10 weeks,	5 to 9	Sow	4 months,	and more	
Opossom -	to weeks,	4 or 5	Camel	12 months,	1	
Kangaroo -		1	Walrus -	9 months,	1	
Jackall -	112 7 137 181	6 to 8	Elephant -	2 years,	1	
Fox	10 weeks,	4 or 5	Whale -	9 or 10 months,	1 or 2a	

Conception being entirely removed from all influence of volition, it is obviously impracticable, by any effort of the will, either to modify the sex of the fœtus, or its general physical and moral characters. Yet idle and absurd schemes have been devised for both one and the other. The older philosophers, as Hippocrates and Aristotle, believed that the right testicle and ovary furnished rudiments of males; and the same organs, on the left side, those of females: some of the old writers, de Re Rustica, assert that this was the result of their experiments with the ram. These statements gave rise to a pretended "art of procreating the sexes at pleasure," which has even been seriously revived in our own time. Mr. John Hunter published an experiment in the Philosophical Transactions, which was instituted for the purpose of determining whether the number of young be equally divided between the ovaria. He took two sows from the same litter, deprived one of an ovarium, and counted the number of pigs each produced during its life. The sow with two ovaria had one hundred and sixty-two: the spayed sow only seventy-six. Hence he inferred, that each ovarium had nearly the same proportion. In this experiment, he makes no mention of the interesting fact regarding the

<sup>2</sup> See, on this subject, Dr. Laycock, on the Nervous Diseases of Women, p. 48, Lond. 1840.

proportion of the males in the two cases, and whether they were not all of the same sex in the sow that had been spayed. Had his attention been drawn to this point, the results would have been sufficient to arrest the strange hypothesis brought forward by Millot, a who boldly affirmed that males are produced by the right ovarium, and females by the left; asserting, that he could so manage the position of the woman during copulation, that she should certainly have a boy or girl, as might have been determined upon: and he published the names of mothers, who had followed his advice, and had succeeded in their wishes! A case, related by Dr. Granville, of London, to the Royal Society, b has completely exhibited the absurdity of this doctrine. A woman, forty years of age, died at the Hospice de la Maternité of Paris, six or seven days after delivery, - of what had been supposed to be a disease of the heart. The body was opened in the presence of Dr. Granville, and the disease was found to be aneurism of the aorta. On examining the uterus, it was discovered to be at least four times the size of what it is during the unimpregnated state. It had acquired its full development on the right side only, where it had the usual pyriform convexity; whilst the leftformed a straight line scarcely half an inch distant from the centre, although it was more two inches from the same point to the outline of the right side. The Fallopian tube and the ovarium, with the other parts on the right side, had the natural appearance; but they were not to be found on the left. Yet this woman had been the mother of eleven children of both sexes; and a few days before her death had been delivered of twins; - one male and one female. M. Jadelot, too, has given the dissection of a female, who had been delivered of several children - boys and girls; yet she had no ovary or Fallopian tube on the right side. Lepelletierd asserts that he saw a similar case in the Hospital of Mans, in 1825, and the Recueils of the Société de Médecine, of Paris, contains the history of an extra-uterine gestation, in which a male fœtus was contained in the left ovary.

Independently of these decisive cases, it has been found, that when one of the ovaries has been entirely disabled by disease, the other has conceived of both sexes. In rabbits, an ovary has been removed; yet both male and fætuses have subsequently been engendered; and if the gravid uterus of one of those animals be examined, male and female fætuses will be found in the same cornu of the uterus, all of which, owing to peculiar construction of the uterus,—the cornu forming the main part of the organ,—must manifestly have proceeded from the corresponding ovary. We are totally unaware, therefore, of the circumstances that give rise to the sex of the new being, although satisfied that it is in no respect influenced by the desires of the parents. We shall see,

a Millot, l'Art de Procréer les Sexes a Volonté ; nouvelle édit., par M. Breschet, Paris, 1829. b Philos. Transact. for 1808, p. 308.

Sir E. Home, Lect. on Comp. Anat. iii. 300.
 Physiologie Médicale et Philosophique, iv. 333, Paris, 1833.

hereafter, that some distinguished physiologists believe, that the sex is not settled at the moment of conception, and that it is determined at a later period, after the embryo has undergone a

certain development.

It is an ancient opinion, which seems to be in some measure confirmed by what we notice in certain animals, that the character of the offspring is largely dependent upon the moral and physical qualities of the parent; and a Dr. Robert, of Paris, in a dissertation under the pompous title of Megalanthropogenesis, has fancifully maintained, that the race of men of genius may be perpetuated by uniting them to women possessed of the same faculties. Similar views are maintained by Claude Quillet.a It is, also, an old opinion, that the procreative energy of the parents has much to do with the mental and corporeal activity of the offspring. Hence it is, that bastards have been presumed to excel in this respect. Such is the view of Burton, b and the same idea is put, by Shakspeare, into the mouth of Edmund.º This, we have no doubt, is erroneous. Much depends upon the condition of the parents as regards their organization and strength of constitution. The remark — " fortes creantur fortibus et bonis" — is true within certain limits; but we have no proof, that the ardour of the procreative effort can have any such influence; and the ratio of instances of bastards, who have been signalized for the possession of unusual vigour - mental or corporeal - to the whole number of illegitimates, is not greater than in the case of those born in wedlock.d It would appear, too, that the number of male children is greater in cases of legitimate than of illegitimate birth. Mr. Babbage has compared the ratio in different countries, from which he has deduced the following table: -

	Legitimate Births.		Number of Births	Illegitima	te Births.	Number of Births
	Females.	Males.	observed.	Females.	Males.	observed.
France,	10,000	10,657	9,656,135	10,000	10,484	673,047
Naples,	10,000	10,452	1,059,055	10,000	10,267	51,309
Prussia,	10,000	10,609	3,672,251	10,000	10,278	212,804
Westphalia,	10,000	10,471	151,169	10,000	10,039	19,950
Montpellier,	10,000	10,707	25,064	10,000	10,081	2,735
Mean,	10,000	10,575		10,000	10,250	The Landing

From the statistics of the Gebäranstall, the Imperial Lying-in

<sup>a</sup> Quilleti Callipædia, sive de Pulchræ Prolis Habendæ Ratione, &c. Lond. 1708. <sup>b</sup> Anatomy of Melancholy, vol. ii.

With base? with baseness? bastardy? base? base? Who in the lusty stealth of nature take More composition and fierce quality Than doth, within a dull, stale, tired bed Go to the creating a whole tribe of fops Got 'tween sleep and wake!" — King Lear, i. 2.

Elliotson's Blumenbach's Physiology, 4th edit. p. 496, Lond. 1828.

e Brewster's Journal of Science, New Series, No. 1; and Quetelet, Sur l'Homme, i. 47, Paris, 1835.

Hospital of Vienna, which is the great receptacle of illegitimate conceptions, it would seem that the number of female children born actually exceeds there that of the males. Of 21,212 children born there in the seven years prior to 1838, the sexes were in the propor-

tion of 10,584 males to 10,628 females.a To elucidate the effect of the condition of the parent on the future progeny, M. Girou de Buzareinguesb gave a violent blow to a bitch, whilst lined, in consequence of which she was paraplegic for some days. She brought forth eight pups, all of which, except one, had the hind legs wanting, malformed, or weak. It has been attempted to show, also, that the corporeal vigour of the parents has much to do even with the future sex. M. Girou instituted a series of experiments on different animals, but especially on sheep, to discover, whether a greater number of male or female lambs may not be produced at the will of the agriculturist. The plan, adopted to insure this result, was to employ very young rams in that division of the flock whence it was desired to obtain females; and strong and vigorous rams, of four or five years of age, in that from which males were to be procured. The result would seem to show, that the younger rams begat females in greater proportion, and the older, males. M. Girou asserts, that females commonly predominate amongst animals, that live in a state of "polygamy," and it is asserted, that the same fact has been observed, in Turkey, and Persia, in our own species; but statistical facts are wanting on this subject. From the researches of Hofackere and Sadler,d it would seem, that, as a general rule, when the mother is older than the father, fewer boys are born than girls, and the same is observed where they are of equal age, but the greater the excess of age on the part of the father, the greater will be the ratio of boys born.e The fact de-

Father younger than mother, Father and mother of equal age, Father older by 1 to 6 years, 6 to 9	103·4 124·7	Father younger than mother, Father and mother of equal age, Father older by 1 to 6 years, 6 to 11	86.5 94.8 103.7 126.7
9 to 18	143.7	11 to 16	147.7
18 and more,	200.0	16 and more,	163.2

duced from the observations of these gentlemen, has been characterised by a recent writer as "one of the most remarkable contributions that have yet been made by statistics in physiology. The following table gives the average results obtained by them; the numbers indicating the proportion of male births to 100 females.

a Austria: its Literary, Scientific and Medical Institutions, by W. R. Wilde, p. 22, Note, Dublin, 1843.

b Mémoire sur les Rapports des Sexes, &c. Paris, 1836; and a farther Memoir, in Revue Médicale, 1837; and Encyclograph. des Sciences Médicales, Janv. 1838. See, also, a notice of the first Memoir by Dr. G. Emerson, in Amer. Journ. of the Medical Sciences, p. 171, May, 1837. Annales d'Hygiène, p. 537, July, 1829.

d The Law of Population, ii. 343, Lond. 1830; and Quetelet sur l'Homme, i. 53,

See, on this subject, Mr. A. Walker, Intermarriage, Amer. Edit., p. 219, New York, 1839.
Carpenter, Human Physiology, § 771, Lond. 1842.

It appears, that the general proportion of males born to the females is every where pretty nearly the same. The calculations of Hufeland give the numbers in Germany as 21 to 20; those of Girou, in France, make the proportions as 21 to 19.69; and in Paris as 21 to 20.27; and the census of Great Britain, taken in 1821, estimates them as 21 to 20.066. In the Dublin Lying-in Hospital, during ten years, the ratio was as 21 to 19.33; and in the Eastern District of the Royal Maternity Charity of London, during the year 1830, it was as 21 to 19.64. In Philadelphia, according to the tables of Dr. Emerson, the proportion from 1821 to 1830, was as 21 to 19.43. In the whole of Europe the proportion is estimated as 106 to 100.b Although, however, a greater number of males may be born, they seem more exposed to natural or accidental death; for amongst adults the balance is much less in their favour. and, indeed, the number of adult females rather exceeds that of the males. Dr. Emersone states, that of the children born in Philadelphia, during the ten years included between 1821 and 1830. amounting, according to the returns made to the Board of Health. to 64,642, there were 2,496 more males than females. But, notwithstanding the males at birth exceeded the females about 71 per cent., the census of 1830 shows, that by the fifth year, the male excess is reduced to about 5 per cent., and at ten years to only 1 per cent.; and that reduction still going on, the females between the ages of 10 and 15, exceed the males about 8 per cent.; and between 15 and 20, 7.3 per cent.; facts, which clearly authorize the deduction of Quetelet,d that during the early stages of life there are agencies operating to reduce the proportion of the male sex. Dr. Emerson's investigations exhibit clearly, that the greater liability of males to accidents did not furnish a sufficient reason for their greater mortality; - the deaths, reported in the Philadelphia bills, under the head of casualties, constituting but a small proportion of the whole mortality; and this - when burns and scalds are included - being more considerable in the case of the female. The gross male mortality under the twentieth year, for the three years above mentioned, exceeded the female in the ratio of 7.94 per cent. The diseases, which seemed to be particularly obnoxious to the male sex, were, according to the Philadelphia bills, the following arranged in the order of their decreasing mortality : - Inflammation of the brain, inflammation of the bowels, bronchitis, croup, inflammation of the lungs, fevers of all kinds (except scarlet), convulsions, general dropsy, dropsy of the head, and small-pox. To these sources of mortality may be added those under the head of

a Amer. Journ. of the Med. Sciences, for Nov. 1835.

c American Journal of the Medical Sciences, for Nov. 1835, p. 56.

d Sur l'Homme, i. 156, Paris, 1835.

b Quetelet, Sur l'Homme, i. 43, Paris, 1835. See, also, Transactions of the Statistical Society of London, vol. i. part i. Lond. 1837; and a notice of the volume by Dr. Emerson, in American Journal of the Medical Sciences, p. 444, for Feb. 1838; Dr. Allen Thomson, art. Generation, Cyclop. Anat. and Physiol. part xiii. p. 478, Feb. 1838; and Burdach's Physiologie als Erfahrungswissenschaft, i. 587, 2te Auflage, Leipz. 1835.

casualties, and others vaguely designated debility, decay, &c. The few cases in which the deaths of females predominated were consumptions, dropsy of the chest, scarlet fever, burns and scalds,

and hooping-cough.a

It would appear that about one infant in twenty is still-born. The cause of this is a difficult inquiry; as well as that of the greater ratio - double - in cities than in the country; in some cities than in others; amongst male infants rather than females; in the winter than in the summer; amongst the illegitimate rather than the legitimate. It is an interesting topic of investigation for the medical statistician.b The following table, embracing the statistics of the Imperial Lying-in Hospital of Vienna, which, as before remarked, is the great receptacle for illegitimate conceptions, is interesting on account of the number of cases observed. It is given by Mr. Wilde, who states, that it was collected and arranged with much care, from the unpublished records of the hospital for the eight years ending Dec. 31, 1840, and exhibits the results of 25,906 deliveries, and 26,149 births.

## NUMBER OF DELIVERIES, 25,906.

Children,	Single b Twins, 2 Triplets, Total	48 tin	nes, 496, or 1 in 15, or 1 in	
Sex in 23,413 births,	SBoys, 11 Girls, 11	,717 Z ,696 S	Proportion of Males to 100 Females,	100-17
Sex of still-born children in 2201 births,	Soys, Girls, Total,	48 45 - 93	Proportion of Males to 100 Females,	106-66
Total still-born in 23,413 Died before the ninth day	pirths,	100	939, or 1 in 1482, or 1 in	
Sexes in 95 of these,	S Boys, Girls,	49 7	Proportion of Males to 100 Females,	106-52
Abortions and premature in 25, 705,	deliveries	3	674, or 1 in	38.13

It will be observed, that the ratio of still-born is smaller than the average; yet it is much higher than in the whole of the Austrian dominions, in which the proportion, according to Mr. Wilde, is 1 in 30.62. It is difficult, however, to believe, that his statistics can be accurate, when we observe the ratio in Linz stated to be 1 in 155.35 only !d

\* See, on this subject, W. Farr, in Third Report of the Registrar-general of Births,

Deaths and Marriages in England, p. 72, Lond. 1841.

b See an article on this subject by the author, in American Med. Intelligencer, for Sept. 1, 1837, p. 203, and ibid. Oct. 1, 1836, p. 252; Dr. Avery, in Transact. of the Med. Society of the State of New York, iii., part ii. p. 179, Albany, 1837; Quetelet, Sur l'Homme, p. 122, Paris, 1835; Brit. and For. Med. Review, July, 1837, p. 234; Dr. Emerson, in American Journ. of the Medical Sciences, Feb. 1838, p. 444; and Prof. Rau, of Bern, Ueber die Unnatürliche Sterblichkeit der Kinder in ihrem ersten Lebensjahre, Bern, 1836, cited in Brit. and For. Med. Rev. April, 1839, p. 593; and Wilde's Austria.

c Austria, &c., &c., p. 223, Dublin, 1843.

e. Superfætation .- It has been an oft agitated question, whether, after an ovule has been impregnated and passed down into the cavity of the uterus, another ovule may not be fecundated; so that the products of two conceptions may undergo their respective developments in the uterus, and be delivered at an interval corresponding to that between the conceptions. Many physiologists have believed this to be possible, and have given it the name of superfætation or superfecundation. The case, cited from Sir Everard Home, of the young female, who died on the seventh or eighth day after conception, exhibits that the mouth of the womb is at an early period completely obstructed by a plug of mucus; and that the inner surface of the uterus is lined by an efflorescence of coagulable lymph, the nature of which will be described under the next head. When such a change has been effected, it would seem to be impossible for the male sperm to reach the ovary; and accordingly, the general belief is, that superfectation is only practicable prior to these changes, and where there is a second vesicle ripe for impregnation. Of this kind of superfætation it is probable, that twin and triplet cases are often, if not always, examples; one ovule being impregnated at one copulation, and another at the next. It seems also to be common in animals. The dog-breeders have often remarked, that a bitch, after having been lined, will readily admit a dog of a very different kind to copulate with her; and where this has occurred, two different descriptions of puppies have been brought forth, - some resembling each of the fathers. Sir Everard Homeb states, that a setter bitch was lined in the morning by a pointer. The bitch went out with the game-keeper, who had with him a Russian setter of his own, which also lined her in the course of the afternoon. She had a litter of six puppies; two only of which were preserved. One of these bore an exact resemblance to the pointer, the other to the Russian setter, - the male influence being predominant in

Of this kind of superfætation or double conception we have several instances on record, of which the following are amongst the most striking, the male parents of the respective fœtuses having differed in colour. The first is the well-known case, cited by Buffon,c of a female at Charleston, South Carolina, who was delivered in 1714 of twins, within a very short time of each other. One of these was black, the other white. This circumstance led to an inquiry, when the woman confessed, that on a particular day, immediately after her husband had left his bed, a negro entered her room, and compelled her to gratify his wishes, under threats of murdering her. Several cases of women in the West India Islands having had, at one birth, a black and a white child, are recorded; and Dr. Moseleyd gives the following case, which is very analo-

d A Treatise on Tropical Diseases, p. 111.

a Art. Zwillinge, in Pierer's Anat. Physiol. Real. Wörterb. Band viii., Altenb. 1829. b Lect. on Comp. Anat. iii. 302. c Hist. Nat. de l'Homme, Puberté.

gous to that described by Buffon. A negro woman brought forth two children at a birth, both of a size, one of which was a negro, the other a mulatto. On being interrogated, she said, that a white man, belonging to the estate, came to her hut one morning before she was up, and that she received his embraces soon after her black husband had quitted her. Sir Everard Homea likewise asserts, that a particular friend of his "knows a black woman, who has two children now alive, that are twins and were suckled together; one quite black, the other a mulatto. The woman herself does not hesitate in stating the circumstances: one morning just after her husband had left her, a soldier, for whom she had a partiality came into her hut, and was connected with her, about three or four hours after leaving the embraces of her husband." One of the author's pupils, Dr. N. J. Huston, then of Harrisonburg, Virginia, also communicated to him the case of a female who was delivered, in March, 1827, of a negro child and a mulatto on the same night. Where negro slavery exists, such cases are sufficiently numerous.<sup>b</sup> So far, therefore, as regards the possibility of a second vesicle being fecundated, prior to the closure of the os uteri by the tenacious mucus and the flocculent membranous secretion from the interior of the uterus, or by the decidua, no doubt, we think, can be entertained; but, except in cases of double uterus, it would seem to be impracticable afterwards; although cases have been adduced to show its possibility. Still, these may perhaps be explained under the supposition, that the uterine changes, above referred to, may not be as rapidly accomplished in some cases as in others; and, again, the period of gestation is not so rigidly fixed, but that children, begotten at the same time, or within twenty-four hours, may still be born at a distance of some weeks from each other. A case happened to the author in which nearly three weeks elapsed between the birth of twins, in whose cases the ova were probably fecundated either at the same copulation or within a few hours of each other.

It may happen, too, that although two ova may be fecundated, both embryos may not undergo equal development. One, indeed, may be arrested at an early stage, although still retaining the vital principle. In such a case, the other will generally be found larger than common. A case of this kind occurred in the practice of Professor Hall, of the University of Maryland. On the 4th of October, 1835, a lady was delivered of a female fætus, 2 inches and 10 lines in length. This occurred about half-past eight in the morning; and, at two o'clock on the following morning, she was delivered of a second child, which weighed 9½ pounds. The fætus, whose development was arrested, was seen by the author.

<sup>2</sup> Op. citat.

b See, for an enumeration of cases, Beck's Medical Jurisprudence, 6th edit. i. 222; and Dr. Allen Thomson, in Cyclop. Anat. and Physiol., part xiii. p. 469, for Feb. 1838. Wagner, Elements of Physiology, by Willis, p. 79, Lond. 1841.

VOL. II. — 35

When first extruded, it gave no evidences of decay, and in colour and general characters resembled the fœtus of an ordinary abortion."

f. Pregnancy. - When the fecundated ovum has been laid hold of by the fimbriated extremity of the Fallopian tube, and through this channel, - perhaps by the contraction of the tubes and the ciliary motions of its lining membraneb - has reached the cavity of the uterus, it forms a union with this viscus, to obtain the nutritive fluids, that may be required for its development, and to remain there during the whole period of pregnancy or utero-gestation; — a condition which will now require some consideration.

Immediately after a fecundating copulation, and whilst the chief changes are transpiring in the ovary, certain modifications occur in the uterus. According to some, it dilates for the reception of the ovum. Bertrandi found this to be the case in extra-uterine pregnancy, and in females whom he opened at periods so near to conception, that the ovum was still floating in the uterus. Its substance appeared at the same time redder, softer, less compact, and more vascular than usual. In the case to which we have more than once alluded from Sir Everard Home, the lining of the uterus was covered by a beautiful flocculent appearance, about the seventh or eighth day after impregnation. The soft flocculent membrane, which forms in this way, is the membrana caduca seu decidua, decidua externa, first described by Hunter; the epicho-

Fig. 233.

rion of Chaussier; the tunica exterior ovi, t. caduca, t. crassa; membrana cribrosa; membrana ovi materna, membrana mucosa; decidua cellularis et spongiosa, of others.

In a case, observed by Von Baer at a very early The dark shade, period, when the decidua was still in a pulpy state, and between the villi of the lining membrane of the uterus, which the villi, is the deci-dua. The uterine in the unimpregnated state are very short, were found vessels are seen ex-tending into the de. to be remarkably elongated; and between the villi, cidua and forming and passing over them, was a substance not organized, but merely effused, and evidently the decidua at an extremely early age.d Others have supposed, that the decidua is composed of the inner portion of the mucous membrane itself, which undergoes a considerable change in its character.

The arrangement of this membrane has given rise to some discussion.f The opinions of most of the anatomists of the present

b Wagner op. citat. p. 137. c Lect. on Comp. Anat. iii. 209, Lond. 1823. 4 Von Baer, op. citat.; and Wagner's Physiology, by Willis, p. 184, Lond. 1841. · Weber and Sharpey, cited in Müller, Elements of Physiology, by Baly, p. 1574, Lond. 1842.

For similar cases, of which many are on record, see Dr. Samuel Jackson, formerly of Northumberland, Pa., now of Philadelphia, in American Journal of the Medical Sciences, May 1838, p. 237, and May, 1839, p. 256; also, Dr. J. G. Porter, ibid. Aug. 1840, p. 307; and Mr. Streeter, Lond. Lancet, Oct. 30, 1841.

Weber's Hildebrandt's Handbuch der Anatomie, iv. 486 & 515, 1832; Purkinje, art. Ei, Encyclop. Wörterb. der Medicin. Wissensch. x. 107, Berlin, 1834; W. Hunter's Anatomical Description of the Human Gravid Uterus and its Contents, Lond. 1794; and Carus, Zur Lehre von der Schwangerschaft, u. s. w., Abth. ii. s. 5.

day are in favour of one of two views. It is maintained by some, that one of the first effects of conception is to cause the secretion of a considerable quantity of a sero-albuminous substance from the inner surface of the uterus; so that the organ becomes filled with it. At first, when the ovum arrives in the uterus, it falls into the midst of this secretion, gradually absorbing a part by its outer surface for its nutrition. The remainder is organized into a double membrane, one corresponding to the uterus, the other adhering to the ovum. This sero-albuminous substance has been assimilated, both to the white, with which the eggs of birds become invested in passing through the oviduct, and to the viscid substance, that envelopes the membranous ova of certain reptiles. It is conceived by some to plug up both the orifices of the Fallopian tubes, and that of the uterus; and according to Krummachera and Dutrochet,b it has been seen extending into the tubes; whilst the remains of that, which plugged up the os uteri, has been recognised in the shape of a nipple on the top of the aborted ovum. To this substance, Breschete has given the name Hydropérione. By others, it is held, that the decidua is slightly organized even prior to the arrival of the ovum, lining the whole of the cavity and being devoid of apertures; so that when the ovum passes along the tube and attains the cornu of the uterus, it pushes the decidua before it; the part so pushed forwards constituting the tunica decidua reflexa or ovuline, and enveloping the whole of the ovum except at the part where the decidua leaves the uterus to be reflected over it. This is the seat of the future placenta. Such is the opinion of Velpeau, Wagner, and others. An objection to it, however, is the difficulty of so small a body pushing the decidua before it; and a still stronger is the assertion of Professor Sharpey, that the structure of the decidua and of the decidua reflexa is different. Hence, it has been thought more probable, that the latter is almost entirely a new production, the growth of which is simultaneous with the enlargement of the ovum, and that the decidua vera has no more share in its formation, than as supplying, through its vessels, the necessary materials.g At the point of reflection of the decidua reflexa, there is a thick stratum of a substance precisely similar to the decidua reflexa, which attaches the ovum to the side of the uterus, and which blends intimately on the outer side of the reflex

<sup>&</sup>lt;sup>2</sup> Diss. Sistens Observationes quasd. Anatom. circa Velamenta Ovi humani. Duisb. 1790.

b Mém. de la Société Médicale d'Emulation, viii. P. i. 1817.

Etudes Anatomiques, Physiologiques, et Pathologiques, de l'Œuf dans l'Espèce Humaine, &c., Paris, 1832.

d Traité Elémentaire de l'Art des Accouchemens, i. 231, Paris, 1829, or Prof. Meigs's translation, 2d edit. 246, Philad. 1838; also, Velpeau, Embryologie ou Ovologie Humaine, Paris, 1833; or a copious Analysis of the same, by the author, in Amer. Journ. Med. Sciences, Aug. 1834, p. 389.

o Op. citat. p. 188.

s Carpenter, Human Physiology, § 748, Lond. 1842.


fold with the decidua vera. This thick stratum is termed the decidua serotina, from its appearing to have been formed at a

Fig. 234.

pregnation.

a. Cervix. b, b. Orifices of Fallopian tubes,

cyst is in contact with the whole surface of the uterine cavity, and



its Cavity.

flexa. - (Wagner.)

later period. It is represented in Fig. The view of Mr. Burnsa differs from this in supposing that the decidua consists of two layers, the innermost of which has no aperture, so that the ovum on attaining the cornu of the uterus pushes it forwards, and forms the decidua protrusa or decidua reflexa. Impregnation, Velpeau says, occasions a specific excitation in the uterus, promptly followed by an exhalation of coagulable matter. This Section of the Uterus about eight days after Im Concretes, and is soon transformed into a kind of cyst or ampulla, filled c. Decidua vera. d. Cavity of uterus. - (Wag- with a transparent or slightly rosecoloured fluid. This species of

sometimes extends into the commencement of the tubes, and most frequently into the upper part of the cervix uteri, in the form of solid, concrete cords; but is never, he says, perforated naturally, as Hunter, Bojanus, Lee and others have maintained. The decidua uteri, according to Velpeau, retains a pretty considerable thickness, especially around the placenta, until the end of gestation: the decidua reflexa, on the contrary, becomes insensibly thinner and thinner, so that at the full period it is at times of extreme tenuity. Towards the third or fourth month—a little sooner or later - they touch and Section of the Uterus when the Ovum is entering press upon each other, and re-

main in a more or less perfect Ovum, f, surrounded by its chorion g. a. main in a more or less perfect Cervix. b, b. Fallopian tubes. c. Decidua State of contiguity, until the exvera. d. Cavity of the uterus. c. Decidua re-pulsion of the secundines but pulsion of the secundines; but, Velpeau asserts, they are never

confounded; and such appears to be the view of Bischoff.b The decidua - the true as well as the reflected - is esteemed by Velpeau a simple concretion - a layer without regular texture - the

b Wagner, op. citat. p. 190 (note).

Principles of Midwifery, 3d edit. p. 147, Lond. 1814.

product of an excretion from the lining membrane of the uterus; on this account, he terms it, "anhistous membrane" (from ar, privative, and b ioros, "a web") or " membrane texture." There has, ? indeed, been a striking dissatisfaction with the name "decidua." Besides the appellatives already given, Dutrochet has proposed to call it épione, Breschet, périone, Seiler, membrana uteri interna evoluta and Burdach, nidamentum.a

A difficulty exists in understanding how the decidua is formed continuously over the orifice of the Fallopian tubes, and over the upper surface of the cervix uteri. A new production must evidently take place there. By some, however, it is not presumed to exist in the latter situation; but a plug of gelatinous matter is found there, as in



some, however, it is not Section of the Uterus with the Ovum somewhat advanced.

presumed to exist in the fallopian tubes. c, c. Decidua vera, prolonged at c 2, into latter situation; but a fallopian tube. d. Cavity of uterus, almost completely occupied by ovum (compare with Fig. 235). c, c. Angles at which decidua vera is reflected. f. Decidua serotina. g. Allantois. ter is found there, as in h. Umbilical vesicle. i. Amnion. k. Chorion, lined with outer fold of serous tunic. —(Wagner.)

The use of the decidua is, in Velpeau's opinion, to retain the fecundated ovum to a given point of the uterine cavity; and if his views of its arrangement were correct, the suggestions would be good. In favour of this arrangement, a good deal might be said. If there were apertures in the decidua corresponding to the Fallopian tubes, it would seem, that the ovum ought more frequently to fall into the serous-albuminous matter about the cervix uteri, and attachment of the placenta over the os uteri ought, perhaps, to occur more frequently than it is known to do. Under M. Velpeau's doctrine, the attachment of the placenta ought to be near the

<sup>\*</sup> Burdach, Die Physiologie als Erfahrungswissenschaft, B. ii.; Bojanus Isis, 1821, H. iii.; and Dr. Robt. Lee's Remarks on the Pathology and Treatment of some of the most important Diseases of Women, London, 1833.

cornu of the uterus, which is, in fact, the case. Of 34 females, who died in a state of pregnancy at the Hôpital de Perfectionnement, an examination of the parts exhibited, that, in twenty, the centre of the placenta corresponded to the orifice of the Fallopian tube; in three, it was anterior to it; in two, posterior; in three, beneath; and in six, near the fundus of the uterus. It is not so easy to subscribe to his assertions regarding the "anorganic" nature of the decidua. Many excellent observers have affirmed, not only that this membrane exists between the placenta and the uterus, which M. Velpeau's view, of course, renders impossible, but that numerous vessels pass between it, the uterus, and the placenta. We know, too, that the safest and most effectual mode of inducing premature labour is to detach the decidua from the cervix uteri, or, in other words, to break up the vessels that form the medium of communication between it and the lining membrane of the uterus. It may be said, indeed, that the mere separation of the "anorganic pellicle" — as M. Velpeau designates it — is a source of irritation, and may excite the uterus to the expulsion of its contents, and this is possible; but he affirms, that no tissue attaches the decidua to the uterus; and that it adheres to the inner surface of the organ merely in the manner of an excreted membraniform shell (plaque). The views of Lepelletiera and Raspail coincide with those of Velpeau as to the decidua being an excretion; but those of Raspail are modified by his peculiar opinions. He maintains, that the surfaces of an organ — whether external or internal - having once fulfilled their appropriate functions, become detached and give place to the layer beneath them; and we have before remarked, that he considers the secretions of the mucous and serous membranes to be constituted of the detritus of those membranes. Now, that which happens to the intestinal canal and the bladder must likewise happen, he affirms, to the uterus, and as, at the period of gestation, it surpasses in development, elaboration, and vitality, every other living organ, it ought necessarily to cast off its layers, in proportion as they have executed the work of elaboration. These deciduous layers constitute the decidua, on which, he says, traces of a former adhesion to the parietes of the uterus, and of the three apertures into the organ, may be met

But the very existence of a decidua reflexa has been denied. It is so by Jörg, Samuel, and by Dr. Granville, who affirms, that it is now scarcely admitted by one in ten of the anatomists of the European continent.e He refers to a specimen of an impregnated uterus in the Museum of the Royal College of Surgeons of London, which exhibits distinctly a round ovum, suspended naturally within the decidua, as a globe may be supposed to hang from some point

Physiologie Médicale et Philosophique, iv. 339, Paris, 1833.

b Chimie Organique, p. 270, Paris, 1833.

Das Gebärorgan des Menschen, u. s. w. Leipz. 1808.

De Ovorum Mammal. Velament. Wirceb. 1816; and art. Ei, in Encycl. Wörterb. der Med. Wissensch. x. 107, Berlin, 1834.

· See, also, Dewees, Compendious System of Midwifery.

of the interior of an oblong sac; and to two specimens, in the collection of Sir Charles Clarke, exhibiting an ovulum, which has already penetrated about an inch into the cavity of the uterine decidua; but neither in these, nor in the specimen of the Royal College, is any part of the uterine decidua pushed forward. The ovum appears to have its natural covering; and, in the College specimen, there is a large space between them and the deciduous lining of the uterus. Dr. Granville regards the decidua reflexa to be the external membrane of the ovum, to which Professor Boër, of Königsberg, gave the name "cortical membrane," and which Dr. Granville terms cortex ovi. It has received various names. By Albinus, it was termed involucrum membranaceum; by Hoboken, membrana retiformis chorii; by Roederer, membrana filamentosa; by Blumenbach, membrana adventitia; and by Osiander, membrana crassa. To this membrane—and to the decidua uteri, as connected with the placenta - we shall have to refer hereafter.

Such is the uncertain state of our information on this interesting topic of intra-uterine anatomy.




Fig. 237.

Extra-Uterine Pregnancy.

a. The uterus, its cavities laid open. b. Its parietes thickened, as in natural pregnancy. c. A portion of decidua separated from its inner surface. d. Bristles to show the direction of the Fallopian tubes. c. Right Fallopian tube distended into a sac which has burst, containing the extrauterine ovum. f. The fœtus. g. The chorion. h. The ovaries; in the right one is a well marked corpus luteum. i. The round ligament.

The decidua manifestly does not belong to the ovum; for it not only exists prior to the descent of the ovum, into the uterus, but is

a Graphic Illustrations of Abortion, &c. p. v. Lond. 1834.

b Burdach, Physiologie als Erfahrungswissenschaft, ii. 75, who refers to the various views on the subject of the decidua reflexa. See, also, Velpeau, in op. cit.; and Purkinje, art. Ei, in Encyclop. Wörterb. der Medicin. Wissensch. x. 107, Berlin, 1834.

even formed, according to Breschet, in all cases of extra-uterine pregnancy. (See Fig. 237.) Chaussier saw it in several cases of tubal gestation. It existed in a case of abdominal pregnancy, cited by Lallemant, and, according to Adelon, Evrat affirms, that one is secreted after every time of sexual intercourse, — which is apocryphal. Recently, Dr. Robert Lee has shown, that the decidua is not formed within the uterus in all cases of extra-uterine gestation. In ten cases detailed by him, and in one other cited from Chaussier, the decidua was seen distinctly surrounding the ovum

in the Fallopian tube.

When the ovum attains the interior of the uterus, which it does in the first five or six days after conception, it forms, in a short space of time, a connexion with the uterus by means of the placenta, in the mode to be mentioned hereafter. During the development of the embryo, it is requisite that the uterus should be correspondently enlarged, in order to afford room for it, as well as to supply it with its proper nutriment. These changes in the uterine system will engage us exclusively at present. In the first two months, the augmentation in size is not great, and chiefly occurs in the pelvis; but, in the fourth, the increase is more rapid. The uterus is too large to be contained in the pelvis, and consequently rises into the hypogastrium. During the next four months, it increases, in every direction, occupying a larger and larger space in the cavity of the abdomen, and crowding the viscera into the flanks and the iliac regions. At the termination of the eighth month, it almost fills the hypogastric and umbilical regions; and its fundus approaches the epigastric region. After this, the fundus is depressed and approaches the umbilions, leaving a flatness above, which has given rise to the old French proverb :- En ventre plat enfant y a. During the first five months of utero-gestation, the womb experiences but little change, maintaining a conoidal shape. After this, however, the neck diminishes in length, and is ultimately almost entirely effaced. The organ has now a decidedly ovoid shape, and its bulk is, according to Haller and Levret, eleven and a half times greater than in the unimpregnated state. Its length, at the full period, has been estimated at about a foot; its transverse diameter at nine inches; its circumference on a level with the Fallopian tubes, at twenty-six inches; and, at the uterine portion of the cervix uteri, thirteen inches. Its weight, which, prior to impregnation, was from fourteen to eighteen drachms, is, at this time, from a pound and a half to two pounds.

Whilst the uterus is undergoing expansion, the size and situation of the parts attached to it also experience modification. The broad ligaments are unfolded; the ovaries and Fallopian


\* Repert. Général d'Anatomie, p. 165, pour 1828.

c Lond. Med. Gazette, June 5, 1840.

b Physiologie de l'Homme, 2de édit. iv. 110, Paris, 1829.

tubes are raised a little, but are subsequently applied against the sides of the uterus. The vagina is elongated. The round ligaments yield to the elevation of the organ as far as their length will permit; but, ultimately, they draw the uterus forward, so that the great vessels of the abdomen are not injuriously compressed. The parietes of the abdomen are so much distended that the cuticle yields; hence, an appearance of cicatrices always exists on the abdomen of one who has borne children; and occasionally, the fasciculi of the abdominal muscles separate so as to give rise to ventral hernia.

The changes, produced in the uterus, are not limited to simple dilatation of its tissue. Its condition has experienced various alterations, dependent upon the new mode of nutrition it has assumed. The whole organ has undergone, not only extension, but inspissation of its parietes. In its unimpregnated condition, it is about four lines thick; in the third month of utero-gestation, five. Its arteries enlarge as well as its veins, which latter form large dilatations at the inner surface. These have been called



Arteries of the Impregnated Uterus.

uterine sinuses. Its nerves are greatly increased in size, as well as its lymphatics; and its proper tissues from being hard, whitish, and incontractile, has become red, soft, spongy, and capable of energetic contraction. A difference of sentiment has existed with regard to the nature of the new tissue of the uterus; some comparing it to the middle coat of arteries; other describing it as partly cellular and partly muscular; but an immense majority esteeming it to be muscular. The respectable name of Blumenbacha is in the minority. The facts in favour of its muscularity appear to us overwhelming. It is clearly muscular in the mam-

\* Instit. Physiol. § 547. See, also, Dr. Ramsbotham, Edinb. Med. and Surg. Journal, xxxix. 456.

miferous animal. Thus, in the rabbit, the muscularity of the uterus, according to Blundell,a is far more conspicuous than that of the intestines; the fibres can be seen coarse and large, and their motion can be observed, if they be examined immediately after the rabbit is killed. The same acute physiologist remarks, that, when developed by pregnancy, the muscularity of the organ is so clear, that if you take a portion of it, and show it to any anatomist, asking him what its nature is, he will unhesitatingly reply - it is muscular. This experiment, he says, he once made himself. He took a portion of the impregnated uterus, showed it to Mr. Green and Mr. Key - "excellent judges on this point," -and, without mentioning the womb, he asked them to tell him what was the structure, when they immediately declared it to be muscular. A similar experiment had previously been made upon Mr. Else, who had made up his mind as to the non-muscularity of the uterus. A small portion was taken to him for his opinion of the precise nature of the tissue submitted to him. It was from the uterus at an advanced period of utero-gestation. After carefully examining it, he gave for answer, that in his opinion it was muscular; but as it was detached from its natural locality, he could not say to what part of the body it belonged. When told, however, that it was a piece of the uterus, he examined it again, and then said that it could not be muscle, for there were no muscular fibres in the uterus.b The arrangement of the fibres is not clearly understood. Generally, perhaps, they are described as running externally, in a longitudinal direction, from the fundus to the neck: beneath this plane is another with circular fibres; but within this the fibres are interlaced in inextricable confusion. Some anatomists, however, enumerate as many as seven superposed planes. The fibres are of much lighter colour than those of ordinary muscles, are more like those of the bladder and intestines, and are collected in very flat and loose fasciculi.c The development of this structure would not seem to be limited to the pregnant condition. It appears to occur whenever the uterus in increased in size, as has been remarked by Dr. Horner, and by Lobstein. The muscular layers are thickest at the fundus uteri. At the cervix uteri, they are extremely small and indistinct.

After the ovum has attained the interior of the uterus, and entered the flocculent decidua, it becomes connected, in process of time, with the uterus, by means of a body to be described hereafter, called the *placenta*, which is attached to the uterus, and communicates with the fœtus by a vascular cord that enters its umbilicus. The seat of the attachment of the placenta — we have seen —

Désormeaux, art Grossesse, in Dict. de Méd. x. 380, Paris, 1824.

d Lessons in Practical Anatomy, p. 304, Philad. 1836.

Principles and Practice of Obstetricy, Amer. Edit. p. 67, Washington, 1834.
 Dr. D. Davis's Principles and Practice of Obstetric Medicine, ii. 850, Lond. 1836.

e Fragment d'Anatomie Physiologique sur l'Organisation de la Matrice dans l'Espèce Humaine, Paris, 1803.

is not always the same. Frequently, it is near one of the cornua of the uterus; but occasionally it is implanted over the os uteri. The diversity of position has given occasion to difference of opinion regarding the causes that influence it. By some, it has been presumed, that, in whatever part of the uterus the ovum lodges, when it quits the Fallopian tube, there an adhesion is formed. By others, it has been said, that as the ovum pushes the decidua at the mouth of the Fallopian tube before it into the uterus, the attachment of the placenta must be near the orifice of the tube. Such would, indeed, appear to be the fact in the majority of cases, but we see so many irregularities in this respect, as to preclude us

from assigning any very satisfactory reason for it.

g. Signs of Pregnancy. - Along with the changes that supervene in the generative apparatus during pregnancy, the whole system commonly sympathises more or less in the altered condition. Some females, however, pass through the whole course of gestation with but very slight or no disturbance of the ordinary functions; whilst, with others, it is a period of perpetual suffering. One of the earliest and most common signs is suppression of the catamenial discharge; but, of itself, this cannot be relied on, as it may result from disease. Soon after impregnation, the digestive and cerebral functions exhibit more or less modification. The female is affected with nausea and vomiting, especially in the morning after rising; the appetite is most fastidious; substances, which previously excited loathing being at times desired or longed for with the greatest avidity; whilst on the contrary, cherished articles of diet can no longer be regarded without disgust. The sleep is apt to be disturbed; the temper to be unusually irritable, even in those possessed of signal equanimity on other occasions. The mammæ enlarge, and become knotty, and sometimes lancinating pains are felt in them; and a secretion of a whitish serum can often be pressed from the nipple. The areola around the nipple becomes of a darker colour in the first pregnancy than it is in the virgin state; and it is darker during each successive pregnancy, than when the female is not pregnant. There is, also, a puffy turgescence, not alone of the nipple, but of the whole of the surrounding disk, with a development of the small follicles around the nipple.a These appearances constitute one of the best single proofs of the existence of pregnancy; but it is obvious, that for accurate discrimination, it is necessary to be aware of the hue in each particular case in the unfecundated state; and, moreover, instances are on record of a well-marked areola occurring in persons who are not pregnant, as well as of an entire absence of areola in those who are. Dr. Guy remarks, that Dr. J. Reid showed him a case of enlarged mammæ, with distinct areolæ and mucous follicles, in a female who had never been, and was not at that time,

<sup>\*</sup> Montgomery's Exposition of the Signs and Symptoms of Pregnancy, &c., p. 62, Lond. 1837, or Dunglison's Amer. Med. Libr. Edit. Philad. 1839; and Hamilton's Practical Observations on Midwifery, Amer. Med. Libr. Edit. p. 44, Philad. 1837.

pregnant.a The author has had numerous opportunities for appreciating the insufficiency of these evidences when taken singly.

It has been affirmed by Dr. Kluge, of Berlin, and by M. Jacquemin, of Paris, that a bluish tint of the vagina, extending from the os externum to the os uteri, is a sure test of pregnancy. According to Kluge, this discoloration commences in the fourth week of utero-gestation, increases until the time of delivery, and ceases with the lochia. M. Jacquemin, on examining the genitals of prostitutes, in compliance with the police regulations of Paris, observed the same peculiarity of colour in the same situation in those that were pregnant: he describes it as a violet colour, or like lees of wine, and so distinct as never to deceive him, being sufficient of itself, and independently of the other signs of pregnancy, to determine the existence of that state. Parent-Duchateletb affirms, that he was present when M. Jacquemin's accuracy in this matter was successfully put to the test: in the investigation, he examined no less than 4500 prostitutes. Dr. Montgomery, however, - from limited observation it is true, - found, that whilst in some cases the bluish colour was very obvious, in others it was so slight as to be scarcely, if at all, perceptible. There is nothing more probable, than that the capillary circulation of the mucous coat of the vagina may be modiffied along with that of the interior of the uterus during pregnancy, so as to give occasion to a change of colour, like that mentioned by those eminent observers; but it may be doubted whether the test can often be available, especially in private practice.

Attention has been paid to the condition of the urine during uterogestation,d but although a difference has appeared to exist between it and the urine of an unimpregnated female, it has not generally been esteemed distinctive.e M. Donné, indeed, affirms, that during pregnancy it contains less uric acid and phosphate of lime than in the natural state, a difference explicable on considering the elements that are necessary for the formation of the organs of the fœtus. The crystallization of the salts of the urine is thus so remarkably modified, that by simple inspection, without examining the females, he has in more than thirty cases recognized the state of pregnancy at different periods. Of late years, the urine of the pregnant female has been found to contain a peculiar substance, which, when it is allowed to stand, separates and forms a pellicle on the surface. To observe this, the urine must be allowed to stand from two to six days, when minute opaque bodies are observed to rise from the bottom to the surface of the fluid, where they gradually agglomerate and form a continuous layer over the sur-

b De la Prostitution dans la Ville de Paris, i. 217, 218, Paris, 1837.

6 Op. citat. p. vi. Lond. 1837.

d Galen and Savanorola, cited by Fodéré, Méd. Légal. i. 435, Paris, 1837.

<sup>\*</sup> See Dr. J. Reid, Lond. Lancet, Dec. 22, 1838; and Dr. W. A. Guy, Principles of Forensic Medicine, p. 72, Lond. 1843.

Montgomery, op. cit. p. 157, E. Rigby, System of Midwifery, p. 57, Lond. 1841
Gazette Médicale de Paris, 29 Mai, 1841.

face, which is so consistent, that it may be almost lifted off the urine by raising it by one of its edges. To this pellicle the name kiestein, or more properly kyestein, has been given. It is whitish, opalescent, slightly granular, and may be compared to the fatty substance, which swims on the surface of soups, after they have been allowed to cool. When examined by the microscope, it has the aspect of a gelatinous mass without determinate form: at times, cubical crystals are observed in it, but this appearance is only seen when it has stood for a long time, and may be esteemed foreign to The kiestein remains on the surface for several days; the urine then becomes turbid, and small opaque masses become detached from the kiestein and fall to the bottom of the fluid; the pellicle then soon becomes destroyed. Various experiments have been made on this matter by Nauche, a Eguisier, Dr. Golding Bird, c Mr. Letheby, d Dr. Stark, and the author's friend Dr. E. K. Kane, of Philadelphia, now physician to the Chinese Embassy; and, at the author's request, by Drs. McPheeters and Perry, resident physicians at the Philadelphia Hospital. They show, that when taken in conjunction with other phenomena, the appearance of the kiestein is certainly a great aid in the diagnosis of pregnancy. Mr. Letheby found unquestionable evidence of kiestein in 48 out of 50 cases between the 2d and 9th month of utero-gestation, and was unable to account for its absence in the two exceptions. The result of Dr. Kane's observations, which the author had an opportunity of examining from time to time, and for the accuracy of which he can vouch, was deduced by Dr. Kane as follows. First. Kiestein is not peculiar to pregnancy, but may occur whenever the lacteal elements are secreted without a free discharge from the mammæ. Secondly. Although it is sometimes obscurely developed and occasionally simulated by other pellicles, it is generally distinguishable from all others. Thirdly. Where pregnancy is possible, the exhibition of a clearly defined kiestenic pellicle is one of the least equivocal proofs of that condition; and Fourthly. When this pellicle is not found in the more advanced stages of supposed pregnancy, the probabilities, if the female be otherwise healthy, are as 20 to 1 (81 to 4) that the prognosis is incorrect. The author has distinctly noticed in some of the cases the cheesy odour of kiestenic urine described by Dr. Bird.

Along with the above signs, the uterus gradually enlarges; and, about the end of the fourth calendar month or the eighteenth week, sooner or later, quickening, as it is usually but erroneously termed, takes place, or the motion of the child is first felt. Prior to this, — from the moment, indeed, of a fecundating copulation, — the

La Lancette Française, and Lond. Lancet, No. clxvii , p. 675.

d Lond. Med Gazette, Dec. 24, 1841.

Edinb. Med. and Surg. Journal, Jan. 1842.

b La Lancette Française, Février 21, 1839. See, also, L'Experience, Juillet 25, 1839. Guy's Hospital Reports, April, 1840.

f American Journal of the Med. Sciences, p. 37, Philad. 1842.

<sup>5</sup> Amer. Med. Intelligencer, March 15, 1841, p. 369.

VOL. II. - 36

female is quick with child, but it is not until this period that the


Fig. 239.



fœtus has undergone the development necessary for its movements to be perceptible. This occurrence establishes the fact of gestation, whatever doubts may have previously existed. Where there is much corpulence, or where the fluid, surrounding the fœtus is in such quantity as to throw obscurity around the case, it may be necessary, for the purpose of verifying the existence of pregnancy, to institute an examination per vaginam. This can rarely afford much evidence, prior to the percentage of the case of the percentage of the case of the case

Cervix uteri at three months. riod of quickening; but, after this, the ex-

amination, by what the French term the mouvement de ballotte-



Cervix uteri at six months.

ment, may indicate the presence or the contrary of a fœtus in the womb. This mode of examination consists in passing the forefinger of one hand into the vagina, — the female being in the erect attitude, — and in giving the fœtus a sudden succussion by means of the other hand placed upon the abdomen. In this way, a sensation is communicated to the finger in vaginâ, which is often of an un-

equivocal character. During the latter months, the cervix uteri exhibits the changes depicted in

the marginal figures.



Cervix uteri at eight months.

Of late years<sup>a</sup> the application of the stethoscope has been used as a means of discrimination in doubtful cases. By applying this instrument to the abdomen of a pregnant female, after the fifth month, the pulsations of the heart of the fœtus are audible. This instrument

may also exhibit when the pregnancy is multiple, by indicating the pulsations of two or more distinct hearts, according as the conception is double, triple, &c. It would appear, however, that auscultation affords but two main signs of pregnancy, — the pulsations of the fœtal heart, and a murmur, which, according to some, should, correctly speaking, be designated the "uterine souffle or murmur." This murmur has been supposed to take place in the uterine artery, which serves for the nutrition of the placenta, and

<sup>&</sup>lt;sup>a</sup> Dr. E. Kennedy's Observations on Obstetric Auscultation, &c., Dublin, 1833; De Kergaradec, Mémoire sur l'Auscultation Appliquée à l'Etude de la Grossesse, &c. Paris, 1822; Dr. J. C. Ferguson, in Hays's Select Medico-Chirurgical Transactions, p. 172, Philad. 1831; Montgomery, op. cit. p. 120, Lond. 1837; and Nägele, Die geburtshülfliche Auscultation, Mainz, 1838; noticed in Brit. and For. Med. Rev. Oct. 1839, p. 368.

even to indicate the situation of the placenta: a others have con-

sidered, that it is is not connected with the placenta, but depends upon the increased vascularity and peculiar arrangement of the uterine vessels during the gravid state; but it has been questioned whether it be ever produced except by pressure on the vessels of the mother. It is, indeed, positively stated,



Cervix uteri at nine months.

that the sound has been heard in cases of uterine and other tumours where there was no pregnancy. In one case of fibrous tumour of the uterus, the author heard it distinctly.

In addition to the placental or uterine murmur and the sounds of the fætal heart, a third sound is occasionally to be heard, and one which is considered to be seated in the umbilical cord. This has been termed the "funic bellows' sound." It is of the bellows species, is synchronous with the first sound of the fætal heart, and appears to depend upon diminution of the caliber of the umbilical arteries, either through pressure, or stretching of the funis, or both combined. It is affirmed, too, by Prof. Nägele, that the movements of the fætus may be distinguished by the stethoscope at a very early period of pregnancy.

The pulsations of the fætal heart vary, according to a number of observers, from 120 to 180 in the minute. Nägele, however, affirms, that he has occasionally found them, but momentarily, to sink as low as 50 or 60. Professor Hamilton refers to various cases, in which Drs. Sidey and Moir attended particularly to the action of the fætal heart previous to breathing, in all of which it was 60 or less in a minute, before the establishment of respiration. Professor Hamilton affirms, that almost half a century has elapsed since he remarked, that in infants who did not breathe upon birth, but in whom the pulsation of the cord continued, the action of the heart did not exceed sixty pulsations in the minute till breathing took place, when it became so frequent that it could not be numbered. This led him to take every opportunity — when he had

Dr. Doherty, Dublin Journal of Med. Science, March, 1840.
 E. Rigby, System of Midwifery, Amer. Edit., p. 90, Philad. 1841.

Raciborski, Manual of Auscultation, translated by Fitzherbert, p. 145, Lond. 1835; E. Rigby, System of Midwifery, p. 54, Lond. 1841; Dr. Hope, Treatise on Diseases of the Heart, Amer. Edit. by Dr. Pennock, Philad. 1842; and Prof. Huston, in Churchill, System of Midwifery, Amer. Edit., p. 136 (note).

d Dietrich, Medicinische Zeitung, No. xxxvii. 1839; cited in Brit. and For. Med.

Review, July, 1840, p. 274; and E. Kennedy, op. cit. p. 121.

Dublin Journal of Medical Science, Jan. 1838.
Practical Observations on Various Subjects relating to Midwifery, Amer. Med. Library edit. P. i. pp. 51, 97, and P. ii. p. 123, Philadelphia, 1837-8.

occasion to introduce his hand into the uterus to extract the infant - to endeavour to ascertain the action of the fætal heart before birth, and he in no instance discovered it to be more frequent than in the still-born infant whose cord beats. Yet neither Dr. Hamilton, nor any one of the gentlemen referred to, denies that pulsations are heard by the stethoscope, varying from 120 to considerably upwards in the minute, and which have been referred to the fætal heart. The truth would seem to be, that in the cases examined by Professor Hamilton, owing to the influence of the parturient efforts on the function of innervation, and through it on the circulation of the fœtus, the pulsations of the fœtal heart were unusually depressed; but in every case, he would, doubtless, have found them isochronous with those of the umbilical cord, had he made the experiment. It is obvious, indeed, that they must be so, seeing that the umbilical arteries are but a part of the circulatory apparatus of the fœtus. In a case observed, at the author's request, by Dr. Vedder, resident physician at the Philadelphia Hospital, it was noticed, that whilst the uterus was quiescent, the pulsations of the fætal heart numbered 140 per minute; but that immediately succeeding a pain they were only 96, and then gradually rose to 140. After delivery, the cord and fætal heart beat respectively 134 in the minute.a Von Hoefft,b of St. Petersburg, has seen the influence of uterine contraction on the circulation of the fœtus exhibited in the most marked manner. During slight pains, the pulsations of the fætal heart continue, but during more violent contractions, especially after the discharge of the waters, they are wholly interrupted, so that the fœtus may be presumed to be in a state of temporary asphyxia in the last periods of labour, and great danger may threaten it if the pains continue for a long time without interruption. The observations of Professor Hamilton ought, therefore, to have no weight with the observer. They were imperfect, inasmuch as the pulsations of the fætal heart were not attended to, whilst he numbered the beats of the cord; and, consequently, they in no respect conflict with the observations of other obstetrical physiologists, which show, that the sounds usually heard during pregnancy, and referred to the fætal heart, are actually owing to the pulsations of that organ.

A case of triplets has been published in which the pulsations of

three distinct fætal hearts were clearly detected.

Lastly, many uneasy feelings, attendant upon gestation, are owing to the increased size of the uterus. These occur more particularly during the latter half of pregnancy. The parietes of the abdomen may not yield with the requisite facility, so that pain may be experienced, especially at the part where the soft parietes join the false ribs. The pressure of the uterus upon the vessels and nerves of the lower extremities occasions enlargement of the

a Dunglison's Amer. Med. Intelligencer, Jan. 15, 1839, p. 311.

b Ibid. April 15, 1839, p. 31; also, Néue Zeitschrift für Geburtskunde, B. vi. s. 1. c Dr. Bell Fletcher, Lond. Lancet, Sep. 11, 1841.

veins of the legs; transudation of the serous part of the blood into the cellular tissue, so as to cause considerable swelling of the feet and ankles; numbness or pricking of the lower limbs, and the most violent cramps, especially when the female is in the recumbent posture, so that she may be compelled to rise suddenly from bed several times in the course of the night. The same pressure exerted on the bladder and rectum, especially during the latter months, brings on a perpetual desire to evacuate the con-

tents of these reservoirs."

h. Duration of Pregnancy.—The duration of human pregnancy has given rise to much discussion amongst medico-legal and obstetrical writers; and opinions still fluctuate largely. In the years 1825-6, a case occurred before the House of Lords, which exhibits this discordance in a striking point of view. It was the Gardner Peerage cause, in which the principal accoucheurs of the British metropolis were examined, - including Sir Charles M. Clarke, Drs. Blegborough, D. Davis, A. B. Granville, Conquest, Merriman, Hopkins, Blundell, and Power. Of seventeen medical gentlemen, who gave evidence, five maintained the opinion, that the period of human utero-gestation is united to about nine calendar months, - from thirty-nine to forty weeks, or from two hundred and seventy to two hundred and eighty days, - and of course considered it to be an impossibility, that the claimant could have been the product of a three hundred and eleven days gestation. On the other side, of twelve medical gentlemen, all of whom appeared to agree that nine calendar months is the usual term of utero-gestation, most of them maintained the possibility, that pregnancy might be protracted to nine and a half, ten, or even eleven calendar months, and were, of course, in favour of the claimant in the cause.b

The difficulty, which arises in fixing upon the precise term, is owing to the impracticability, in ordinary cases, of detecting the time of conception. The sensations of the female are most fallacious guides; and accordingly, as has been previously remarked, she is usually in the habit of reckoning from ten days after the disappearance of the catamenia; but it is manifest, that impregnation might have taken place on the very day after their cessation, or not until a day prior to the subsequent period; so that in this way, an error of at least ten days might occur in the estimate; and again, it does not always happen, that the menstruation, immediately succeeding, is arrested. The period of quickening, which generally happens about the eighteenth week of uterogestation, does not afford us more positive evidence, seeing that it is liable to vary; being experienced by some females much ear-

a See, on the Signs of Pregnancy, Dr. T. R. Beck, Amer. Journ. of the Medical

Sciences, Jan. 1843, p. 112.

b The Medical Evidence, relative to the Duration of Human Pregnancy, as given before a Committee of the House of Lords, in 1825 and 1826, with notes, by Robert Lyall, M.D. Lond. 1826. See, also, Beck's Med. Jurisprudence, 6th edit. i. 477, Philad. 1838.

lier, and, by others, somewhat later. We are, however, justified in stating, that the ordinary duration of human pregnancy is ten lunar months or forty weeks; but we have no less hesitation in affirming, that it may be protracted, in particular cases, much beyond this. We find in animals, where the date of impregnation. can be rigidly fixed, whilst the usual term can be determined without difficulty, that numerous cases are met with in which the period is protracted, and there is no reason to doubt, that the same thing happens occasionally to the human female. Earl Spencer has communicated the results of his observations on cows for a number of years to the English Agricultural Society.b Of 764 cases, 314 calved before the 284th day, and 310 after the 285th; so that the probable period of gestation, he thinks, ought to be considered 284 or 285 days, and not 270, as stated in the book upon Cattle, published under the superintendence of the Society for the Diffusion of Useful Knowledge. The following table exhibits the results of Lord Spencer's observations:

No. of days of gestation.		Cows calved.	No. of days of gestation.				Cows calved.			
220		1 (11)	-	1	276	BRAN	11140	min	15	M
226	1	1		1	277	O ( Sel		1 192	14	
233	DI.	-	-	1	278	114	120	1121	18	
234	112	111-11		1	279	111-51	-	-	32	
235		-		1	280			-	35	
239	-	-	-	1	281	-		-	39	
242	-	1951		1	282	91111	1	1	47	
245	1	120	1	2	283	1	19:11	18	54	
246	11.3	NA.	1000	2	284	-	-	-	66	
248			-	1	285		-	-	74	
250	-	-	-	1	286	11128		-	. 60	
252			-	2	287	121	1930	1934	52	
253	-		-	1	288	-		-	42	
254		100-11	1	1	289	-		-	45	
255	-	-	-	2	290	-			23	
257	-	-	-	2	291	-		1	31	
258	-		-	3	292	112	100	50	16	
259	1	11 280	1 343	1	293				10	
262	11-11	10-01		1	294	-	-	1	8	
263	1.		-	2	295	-	150		7	
266	-			1	296	1924	1200	12	6	
268		-	-	2	297	124	THE T	HEER	2	
269	-	1	1	2	299	1	-	116	1	
270	100	10000	-	5	304	-			1	
271			0.00	6	305	-	5.18		1	
272			-	3	306			000	2	
273		1000	-	3	307	THE WAY	1	1	3	
274	1		-	5	313	1000	188	-	1	
275	110	10-0	-	5	-				1	

The observations of Earl Spencer<sup>c</sup> have suggested an interesting question, in regard to man, on which we have no knowledge, but which is worthy of investigation. He has noticed, that cows in calf

b Journal of the English Agricultural Society, part ii. 1839.

c Dr. Hall, Lond. Med. Gazette, May 6, 1842.

<sup>&</sup>lt;sup>2</sup> Tessier, in Mémoir. de l'Académie Royale des Sciences, An. 1817, tom. ii. p. 1; Hamilton, op. cit. p. 56; and Montgomery, op. cit. p. 267.

to a particular bull belonging to him, carried their calves about four days longer than cows in calf to any other bull; — the average

period of gestation being in them 2904 days.

In a case detailed by Dr. Dewees, an opportunity occurred for dating with precision the time of fecundation. The case is likewise interesting in another respect, as demonstrating, that fecundation does not necessarily arrest the succeeding catamenial discharge. The husband of a lady, who was obliged to absent himself many months, in consequence of the embarrassment of his affairs, returned one night clandestinely; his visit being known only to his wife, her mother, and Dr. Dewees himself. The lady was, at the time, within a week of her menstrual period; and, as the catamenia appeared as usual, she was induced to hope, that she had escaped impregnation. Her catamenia did not, however, make their appearance at the next period; the ordinary signs of pregnancy supervened; and in nine months and thirteen days, or in two hundred and ninety-three days from the visit of the husband, she was delivered.b In his evidence before the House of Peers, in the case before alluded to, Dr. Granville stated his opinion, that the usual term of utero-gestation is as we have given it; but he, at the same time, detailed the case of his own lady, in whom it had been largely protracted. Mrs. Granville passed her menstrual period on the 7th of April, and on the 15th of August following she quickened; - that is, four months and six or seven days afterwards. In the early part of the first week in January, her confinement was expected, and a medical friend desired to hold himself in readiness to attend. Labour pains came on at this time, but soon passed away; and Mrs. G. went on till the 7th of February, when labour took place, and the delivery was speedy. The child was larger and stronger than usual, and was considered by Dr. Granville, - as well as by Dr. A. T. Thomson, Professor of Materia Medica in the University of London, - to be a ten months' child. Now, if, in this case, we calculate, that conception occurred only the day before the interruption of menstruation, three hundred and six days must have elapsed between impregnation and birth; and if we take the middle period between the last menstruation and the interruption, the interval must have been three hundred and sixteen, or three hundred and eighteen days.

The limit, to which the protraction of pregnancy may possibly extend, cannot be assigned. It is not probable, however, that it ever varies largely from the ordinary period. The University of Heidelberg allowed the legitimacy of a child, born at the expiration of thirteen months from the date of the last connubial intercourse; and a case was decided by the Supreme Court of Friesland, by which a child was admitted to the succession, although it was

A Compendious System of Midwifery, 7th Edit. Philad. 1835.

b See a case of protracted gestation communicated by Dr. James R. Manley to Dr. T. R. Beck, in Amer. Journ. of the Medical Sciences, Jan. 1831, p. 59; and the case of Innes v. Innes, cited by Dr. Beck, in Amer. Journal of the Medical Sciences, July, 1841, p. 235. See, also, Dr. W. A. Guy, Principles of Forensic Medicine, part i. p. 167, Lond. 1843.

not born till three hundred and thirty-three days from the husband's death; or only a few days short of twelve lunar months. These are instances of the ne plus ultra of judicial philanthropy, and, perhaps we might say, credulity. Still, although extremely improbable, we cannot say that they are impossible. This much, however, is clear, that real excess over two hundred and eighty days is by no means frequent; and we think, in accordance with the civil code now in force in France, that the legitimacy of an infant born three hundred days after the dissolution of marriage may be contested; although we are by no means disposed to affirm, that if the character of the woman be irreproachable, the decision should be on the side of illegitimacy. Professor Hamilton, indeed, says he is "quite certain," that the term allowed by the French code is too limited, and, he is inclined to regard ten calendar months, which he believes to be the established usage of the Consistorial Court of Scotland, as a good general rule, liable to exceptions, upon satisfactory evidence that menstruation had been obstructed for a certain period.a

Much uncertainty exists as to the earliest period at which a child is viable or capable of carrying on an independent existence. The period is generally fixed at near the end of the seventh month, but children have lived for some time, which have been born earlier. Much evidence was brought forward in a recent case in Scotland, to show that it is possible for a child to live some months, which has been born at the conclusion of 24 weeks of utero-gestation. In that case, the Presbytery decided in favour of the legitimacy of an infant born alive within 25 weeks after marriage. The difficulty, in such cases, of fixing the exact date of conception must necessarily render all computation in regard to the precise

age of the child uncertain.b

i. Parturition.—At the end of seven months of utero-gestation, and even a month earlier, the fœtus is capable of an independent existence; provided, from any cause, delivery should be hastened. This is not, however, the full period, and although labour may occur at the end of seven months, the usual course is for the fœtus to be carried until the end of about nine calendar months. If it be extruded prior to the period at which it is able to maintain an independent existence, the process is termed abortion or miscarriage; if between this time and the full period, it is called premature labour. From certain records, abortion or premature labour has been estimated to occur, on the average, once in 78½ cases. In the Gebäranstalt of Vienna, the inmates of which are chiefly unmarried, the ratio appears to be more than double, or 1 in 38·13.

With respect to the causes, that give rise to the extrusion, we are in utter darkness. It is in truth as inexplicable as any of the

a Op. cit., Amer. Med. Library edit. p. 59.

b See Beck's Medical Jurisprudence, 6th edit., Philad. 1838; Carpenter, Human Physiology, § 755, Lond. 1842; and Dr. W. A. Guy, Principles of Forensic Medicine, Part i. p. 177, Lond. 1843.

Churchill, Theory and Practice of Midwifery, Amer. Edit. by Prof. Huston, p. 167, Philad. 1843.

d Wilde, Austria, &c. p. 222, Dublin, 1843.

other instinctive operations of the living machine. Yet although this is generally admitted, the discussion of the subject occupies a considerable space in the works of some obstetrical writers. Our knowledge appears to be limited to the fact, that when the fœtus has undergone a certain degree of development, and the uterus a corresponding distension, its contractility is called into action, and the uterine contents are beautifully and systematically expelled. Nor can we always fix upon the degree of distension, that shall give occasion to the exertion of this contractile power. Sometimes, it will supervene after a few months of utero-gestation so as to produce abortion; at other times, it happens when the fœtus is just viable; and at others, again, and in the generality of cases, it is not elicited until the full period. In cases of twins, the uterus will admit of still greater distension before its contractility is aroused.

In regard to the action of the muscles specially concerned in the process, it would seem, that the diaphragm and abdominal muscles are excited to action through an excitor influence conveyed from the uterus to the spinal cord; whilst the contractions of the uterus take place independently of all connection with the nervous centres, — like the peristole of the intestines, and the systole of the ventricles of the heart. The fœtus has been observed to be expelled after the cessation of the respiratory movements of the mother. This, as has been suggested, has probably occurred in consequence of the uterine fibres retaining their power of contraction longer than those of the muscles supplied by the cerebro-spinal nerves.

A day or two preceding labour, a discharge is occasionally observed from the vagina of a mucous fluid more or less streaked with blood. This is termed the show, because it indicates the commencement some dilatation of the neck, or mouth of the womb,-the forerunner of labour or tra-The external vail. organs, at the same time, become tumid and flabby. The orifice of the uterus, if an examination be made, is perceived to be enlarging; and its edges are thinner. Along with this, slight

\* Dewees, Comp. of Midwifery.




Natural Labour.

b Carpenter, Human Physiol. p. 153, Lond. 1842.

grinding pains are experienced in the loins and abdomen. After an uncertain period, pains of a very different character come on, which commence in the loins, and appear to bear down towards the os uteri. These are not constant, but recur, at first after long intervals, and subsequently after shorter,—the body of the uterus manifestly contracting with great force, so as to press the ovum down against the mouth of the womb, and to dilate it. In this way, the membranes of the ovum protrude through the os uteri with their contained fluid, the pouch being occasionally termed the bag of waters. Sooner or later, the membranes give way, the waters are discharged, and the uterus contracts so as to embrace the body of the child, which was previously impracticable, except through the medium of the liquor amnii.

At the commencement of labour, the child's head has not entered the pelvis, the occiput being generally towards the left acetabulum; but, when the uterine contractions become more violent, and are accompanied by powerful efforts on the part of the abdominal muscles, the head enters the pelvis, the mouth of the womb becomes largely dilated, and the female is in a state of agitation and excitement, owing to the violence of the efforts, and the irresistible desire she has of assisting them as far as lies in her power. When the head has entered the pelvis, in the position described, in which the long diameter corresponds to the long diameter of the pelvis, it describes, laterally, an arc of a circle, the face passing into the hollow of the sacrum, and the occiput behind the arch of the pubis. By the continuance of the pains, the head presents at the vulva.



Rotation of the Head.

The pains now become urgent and forcing. The os coccygis is pushed backwards, and the perineum is distended. - at times so considerably as to threatten, and even to effect laceration; the anus is also forced open and protruded: the nymphæ and carunculæ of the vagina are effaced; the labia separated, and the head clears the vulva, from the occiput to the chin, experiencing a vertical rotation as depicted in Fig. 244. When the head is extruded, the shoulders and

rest of the body readily follow on account of their smaller dimensions. The child, however, still remains attached to the mother by the navel-string, which has to be tied, and divided at a few fingers' breadth from the umbilicus.

After the birth of the child, the female has generally a short interval of repose; but after a time, slight bearing down pains are experienced, owing to the contraction of the uterus for the separation of the placenta, and of the membranes of the ovum, called

the secundines or afterbirth.

The process of parturition is accomplished in a longer or shorter time in different individuals, and in the same individual in different labours, according to the particular conditions of the female and fœtus. The parts, however, when once dilated, yield much easier afterwards to similar efforts, so that the first labour is generally

the most protracted.

After the separation of the secundines, the female is commonly left in a state of debility and fatigue; but this gradually disappears. The uterus also contracts; its vessels become tortuous, small, and their orifices are plugged up. For a short time, blood continues to be discharged from them; but as they become obliterated by the return of the uterus to its usual size, the discharge loses its sanguineous character, and is replaced by one of a paler colour, called the *lochia*, which gradually disappears, and altogether ceases in the course of two or three weeks after delivery.

For a day or two after delivery, coagula of blood form in the interior of the uterus, especially in the second and subsequent labours,

which excite the organ to contraction for their expulsion. These contractions are accompanied with pain, and are called after-pains; and as their object is the removal of that, which interferes with the return of the uterus to its proper dimensions, it is obvious that they ought not to be officiously interfered with.

Whilst the uterus is contracting its dimensions, the other parts gradually resume the condition they were in prior to delivery; so that, in the course of three or four weeks, it is impracticable to pronounce positively,



Breech Presentation.

whether delivery has recently taken place or not.

Labour, as thus accomplished, is more deserving of the term in the human female than in animals; and this is partly owing to the large size of the fætal head, and partly to the circumstance, that in the animal the axis of the pelvis is the same as that of the body, whilst in the human female, the axis of the brim, as represented by the dotted straight lines in Fig. 244, forms a considerable angle with that of the outlet. In rare cases, the child is extruded without labour-pains. The author was called in the night, to a female, who declared to him, that she was awoke by a slight abdominal uneasiness, when she found both the child and secundines expelled; and other cases of a like kind are on record. These facts should be borne in mind in charges of infanticide.

The position of the child — with the face behind and the occiput before — constitutes the usual presentation in natural labour. Of twelve thousand six hundred and thirty-three children, born at the Hospice de la Maternité of Paris, twelve thousand one hundred and twenty, according to M. Jules Cloquet, were of this presentation; sixty-three had the face turned forward; one hundred and ninety-eight were breech presentations; (see Fig. 245;) in one hundred and forty-seven cases





the feet presented; and in three, the knees. All these, however, are cases in which labour can be effected without assistance: the knee and feet presentations being identical as regards the process of delivery with that of the breech. But, whenever any other part of the fœtus presents, the position is unfavourable, and requires that the hand should be introduced into the uterus, with the view of bringing down the feet, and

converting the case into a foot presentation. The marginal figure of a presentation of the right superior extremity, sufficiently shows, that labour could not be accomplished without the efforts of art.

The following table, drawn up from data furnished by Velpeau, will show the comparative number of presentations, according to the experience of the individuals mentioned.

T. Lewis, Lond. and Edinb. Monthly Journ. of Med. Science, July, 1842; Mr. Rawson, Lond. Lancet, Nov. 27, 1841.

TABLE
EXHIBITING THE RATIO OF PRESENTATIONS IN 1000 CASES.

	According to							
	Merri- man.	Bland.	Mde. Boivin.	Mde. Lacha- pelle.	Nägele.	Lovati.	Hospital of the Faculté.	Boer.
Regular, or of the vertex,	924	944	969	933	933	911	980	14
I. Occipito anterior,	908		944	910		895	1111111111	
a. Occipito-cotyloid (left)		1973	760	717	- Later	537	1000	
Do. (right)	10000		179	209	1177		month	
b. Occipito-pubian,		15583	0.29	MY BY	THE REAL PROPERTY.	Holman !	6083	
II. Occipito posterior,	COUNTY.	TO STATE OF	9.4	9	PORTE !			
a. Fronto-cotyloid (left)	10m	15.33	5.3	7.3		119	100	
b. Do. (right)	112 031	120 11	4.4	2.9	OUTSIN		1877	
Face presentation,	2.2	2.6	3.6	4.6	100000	4350.00	1000	8.8
Mento-iliac (right)	1333	1377	1000	2.6	La com	H. France	Den 23	
Of the pelvis,	36	28	29	36	47	1200	19	29
Of the foot,	12.7	9.4	The sale	14	1000	11/2 11	1 11.	10.3
Of the knees,		1	0.19	0.40		100	90/	
Of the breech,	23	13	18	22				19
Of the trunk,	1	1	4.6	5.3	4.8			
Requiring Forceps,	6.6	4.7	4.6	3.4	36	The Street	1384	5.7
Turning,	16	4.7	1211-7	7.8	7.2	1 000	0)778	5.9
Cephalotomy,	3.3	5.2	4.77	0.53	2.4	13	A Same	1.5

It is found, that the period of the twenty-four hours has some influence upon the process of parturition; about five children being

born during the night for four during the day.b

The parturient and child-bed condition is not devoid of danger to the female; yet the mortality is less than is generally, perhaps, imagined. In some of the great lying-in institutions it has been, however, enormous, and in the Gebüranstalt, of Vienna, it is still estimated at 1 in 30.87! The number of deaths, during labour and subsequently, connected therewith, has been stated to be in Berlin as 1 in 152; In Königsberg, as 1 in 168; and in Wirtemberg, as 1 in 175; a proportion much less than during the last century. Dr. Collinse states, that of 16,414 women, delivered in the Dublin Lying-in Hospital, 164 died, or in the proportion of 1 in 100; and if, he observes, from this number we deduct the deaths from puerperal fever, which may be considered accidental, the proportion becomes greatly diminished, viz. to 1 in 156 deliveries; and again,

b Quetelet, Sur l'Homme, i. 102, Brux. 1835; Dr. Buek, Nachricht von dem Gesundheits-zustande der Stadt Hamburg, von N. H. Julius, s. 157, Hamburg, 1829; and Dunglison's Amer. Med. Intelligencer, Sept. 1, 1837, p. 213.

c See an article on the subject, by the author, in his Amer. Med. Intelligencer,

Sept. 1, 1837, p. 264.

d Casper, Beitrage zur Medicinisch. Statistik, u. s. w. Berl. 1825; and Elements of Medical Statistics, by Dr. B. Hawkins, Lond. 1829. See, also, Quetelet, i. 130.

Velpeau, Traité Elémentaire de l'Art des Accouchemens, Paris, 1829; or Meigs's translation, 2d edit. Philad. 1838. See, also, on the same subject, Dr. Collins, Practical Treatise on Midwifery, Lond. 1835; Churchill, Theory and Practice of Midwifery, Amer. Edit. by Prof. Huston, Philad. 1843; and Wilde, Austria, &c., &c., p. 223, Dublin, 1843.


Op. citat. p. 366.
 VOL. II. — 37

if we subtract the deaths from causes not the results of childbirth, the mortality, from effects arising in consequence of parturition, is vastly reduced, viz. to 1 in 244. In the year 1839, childbirth was fatal to 2915 women throughout England and Wales. Of 1,000,000 females living, 368 died from this cause in 1838, and 372 in 1839. About 5 births in 1000, it was estimated, were fatal to the mother. In 1840, the ratio was greater, about 1 in 187.

The further details of this subject belong more appropriately to

obstetrics.

j. Lactation. — When the child has been separated from the mother, and continues to live by the exercise of its own vital powers, it has still to be dependent upon her for the nutriment adapted to its tender condition. Whilst in utero this nutriment consisted of fluids placed in contact with it, but, after birth, a secretion serves this purpose, which has to be received into the stomach



Milk Ducts in Human Mamma.

The ducts are filled with wax. — (Sir Astley Cooper.)

and undergo the digestive process. This secretion is the milk. It is prepared by the mammæ or breasts, the number, size, and situa-

b W. Farr, in Fourth Annual Report, &c., &c., p. 219, Lond. 1842.

<sup>&</sup>lt;sup>a</sup> W. Farr, in Third Annual Report of the Registrar-general of Births, Deaths and Marriages in England, p. 74, London, 1841.

tion of which are characteristic of the human species. Instances are, however, on record of three or more distinct mammæ in the same individual.a Two such cases are described by Dr. G. C. M. Roberts, of Baltimore. At times, two nipples are met with on one breast. Three cases of the kind are given by Tiedemann, and one recently, by Dr. Chowne. In some instances, the supernumerary breasts have been on other parts of the body.d

Each breast contains a mammary gland, surrounded by the fat of the breast, and resting on the pectoralis major muscle. It is formed of several lobes, united by a some what dense cellular tissue,

and consisting of smaller lobules, which seem, again, composed of round granulations, of a rosy-white colour, and of about the size of a poppy seed. These granula or acini, according to Reil,e cannot be distinguished in the mammæ of the virgin. The glandular granula give origin to the excretory ducts, called tubuli lactiferi or galactophori, which are tortuous, extensible, and transparent. These enlarge and unite as exhibited in a mercurial injection. — (Sir Astley Cooper.) with each other, so that those of each lobe





remain distinct from, and have no communication with, the ducts of any other lobe. All these finally terminate in sinuses, or reservoirs, near the base of the nipple, which are fifteen or eighteen in number, and open on the nipple, without having communication with each other.

The size and shape of the breast are chiefly caused by the cellular tissue in which the mammary gland is situate: this is covered by a thin layer of skin, which is extremely soft and delicate, and devoid of folds. In the middle of the breast is the tubercle, called the nipple, mammella, or teat, - a prominence consisting of an erectile spongy tissue, differing in colour from the rest of the breast.

The nipples do not project directly forwards, but forwards and outwards, for wise purposes, which have been thus depicted by Sir Astley Cooper: - " The natural obliquity of the mammella or nipple forwards and outwards, with a slight turn of the nipple upwards, is one of the most beautiful provisions in nature both for the mother and her child. To the mother, because the child rests upon her arm and lap, in the most convenient position for sucking;

<sup>&</sup>lt;sup>a</sup> Dr. Robt. Lee, London Medical Gazette, Jan. 20, 1838; Medico-Chirurg. Transact. vol. xxi., Lond. 1838; and Mr. Thursfield, Lond. Med. Gaz. March 3, 1828, p. 898.

<sup>&</sup>lt;sup>b</sup> Baltimore Medical and Surgical Journal, ii. 497, Baltimore, 1834.

c Lond. Lancet, July 2, 1842, p. 465.

<sup>4</sup> Art. Cas Rares, in Diction. des Sciences Medicales; Journal de Physiologie, par Magendie, Janv. 1827; Hedenus, art. Brust (weibliche,) in Encyclop. Wörterbuch der Medicin. Wissenschaft. vi. 352, Berlin, I832; art. Brustwarze, ibid. p. 406; Davis's Principles and Practice of Obstetric Medicine, ii. 777, Lond. 1836; Petrequin, in Gazette Médicale de Paris, No. xiii. Avril 1, 1837; and Chowne, op. cit.

<sup>·</sup> Schlemm, art. Brüste, in Encyclop. Wörterb. der Medicin. Wissenschaft. vi. 332,

On the Anatomy of the Breast, p. 12, London, 1840.

for if the nipple and breast had projected directly forwards, the child must have been supported before her in the mother's hands in a most inconvenient and fatiguing position, instead of its reclining upon her side and arm. But it is wisely provided by nature, that when the child reposes upon its mother's arm, it has its mouth directly applied to the nipple, which is turned outwards to receive it, whilst the lower part of the breast forms a cushion upon which the cheek of the infant tranquilly reposes."

The erection of the nipple, which is so manifest during the process of suckling, and can be readily produced by handling it, has been supposed to be owing to an arrangement similar to that of the corpora cavernosa penis, or to a venous circle surrounding the nipple; but Sir Astley Cooper attributes it simply to an afflux of

blood into the capillaries of the part.

Around the nipple is the arcola, which is of a rosy hue in youth, but becomes darker in the progress of life, and the capillary system of which is so delicate as to blush, like the countenance, under similar emotions. The changes, produced on the arcola by gestation, have been already described. The skin, at the base of the nipple, and on its surface, is rough, owing to the presence of a number of sebaceous follicles, called by Sir Astley Cooper the "tubercles of the arcola," which secrete a fluid for the lubrication of the part, and for defending it from the action of the secretions of the mouth of the infant during lactation. Numerous arteries, veins, nerves and lymphatics, — the anatomical constituents of organic textures in general, — also enter into the composition of the mammæ and nipples.<sup>b</sup>

The mammary gland of the male is analogous to that of the

female, but much smaller.

The secretion of milk is liable to longer intermissions than any other function of the kind. In the unmarried and chaste female, although the blood, whence milk is formed, may be constantly passing to the nipple, no secretion takes place from it. It is only during gestation and some time afterwards, as a general rule, that the necessary excitation exists to produce it. Yet although largely allied to the generative function, — the mammæ undergoing their chief development at puberty and becoming shrivelled in old age, — the secretion may arise independently of impregnation; for it has been witnessed in the unquestionable virgin, in the superannuated female, and even in the male sex. The fact as regards the unimpregnated female is mentioned by Hippocrates. Baudelocquestates, that a young girl at Alençon, eight years old, suckled her brother for the space of a month. Dr. Gordon Smithd refers to a manuscript in the collection of Sir Hans Sloane, which gives an

Art. d'Accouchement, i. 188, Paris, 1822. 6 Forensic Medicine, p. 484.

<sup>&</sup>lt;sup>2</sup> Prof. Sebastian, Tijdschrift voor Natuurlijke Geschiedenis en Physiologie, door J. Van der Hoeven en W. H. de Vriese, 2de Deel, bl. i., Amsterdam, 1835.

See, on the intimate structure of the Mammary Glands, Sir Astley Cooper, op. cit.; Mr. Solly, art. Mammary Glands, Cyclop. of Anatomy and Physiology, part xxi. April, 1841; and Dr. Carpenter, Human Physiology, § 681, Lond. 1842.

account of a woman, at the age of sixty-eight, who had not borne a child for more than twenty years, and who nursed her grandchildren, one after another.a Professor Hall, of the University of Maryland, related to the author the case of a widow, aged fifty, whom he saw giving suck to one of her grandchildren, although she had not had a child of her own for twenty years previously. The secretion of milk was solicited by putting the child to her breast during the night, whilst weaning it. Dr. Francis, of New York, describes the case of a lady, who, fourteen years previously, was delivered of a healthy child after a natural labour. "Since that period," he remarks, "her breasts have regularly secreted milk in great abundance, so that, to use her own language, she could at all times easily perform the office of a nurse :" and Dr. Kennedy, of Ashby-de la Zouch, has described the case of a woman, who menstruated during lactation, suckled children uninterruptedly through the full course of forty-seven years, and, in her eightyfirst year, had a moderate, but regular supply of milk, which was rich, and sweet, and did not differ from that yielded by young and healthy mothers. In a note, with which the author has been favoured by Dr. Samuel Jackson - formerly of Northumberland county, Pa., now of Philadelphia - a case is described, of a lady, certainly above sixty-five years of age, who nursed one of her daughter's twins. She had not borne a child for many years, and was suddenly endowed with a full flow of milk. A lady of Northumberland observed to Dr. Jackson, "that she could not but admire the beautiful fulness and contour of her bosom." Dr. Richard Clarke,d of Union Town, South Alabama, gives the case of a lady, who had never borne a child, and who was requested to take charge of an infant, during the illness of its mother. In the course of the night, the infant became restless and fretful, and the lady - to quiet it - put her nipple into its mouth. This was done from time to time, until the milk began to flow. An interesting fact, connected with this case was, that some time afterwards she conceived, and at the expiration of the usual term was delivered of a fine child. Dr. Clarke refers to other cases, which would appear to show, in another form, the intimate and mysterious sympathy that exists between the mammæ and the uterus. But these, and cases of a similar nature, of which there are many on record,e do not possess the same singularity as those of the function being executed by the male. Yet we have the most unquestionable authority in favour of the occurrence of such instances. A Bishop of Corkf relates the case of a man who suckled his child after the

See a similar case, by Mr. Temple, in North of England Med. and Surg. Journal,
 i. 230.
 Medico-Chirurgical Review for July, 1832.

c A similar case is given by Audubert, in Journal de la Société de Médecine, Pratique de Montpellier; and Encyclographie des Sciences Médicales, Février, 1841, p. 299.

d Dunglison's American Medical Intelligencer, April 16, 1838, p. 19.

Elliotson's Blumenbach, 4th edit. p. 509, Lond. 1828.
 Philos. Trans. xli. 813.

death of his wife. Humboldt adduces one of a man, thirty-two years of age, who nursed his child for five months on the secretion from his breasts; Captain Franklina gives a similar instance; and Professor Hall, of the University of Maryland, exhibited to his obstetrical class, in the year 1827, a coloured man, fifty-five years of age, who had large, soft, well-formed mammæ, rather more conical than those of the female, and projecting fully seven inches from the chest; with perfect and large nipples. The glandular structure seemed to the touch to be exactly like that of the female. This man, according to Professor Hall, had officiated as wet-nurse, for several years, in the family of his mistress, and he represented, that the secretion of milk was induced by applying the children, entrusted to his care, to the breasts, during the night. When the milk was no longer required, great difficulty was experienced in arresting the secretion. His genital organs were fully developed.b

It appears, therefore, that the secretion of milk may be caused, independently of a uterus, by soliciting the action of the mammary glands, but that this is a mere exception to the general rule, according to which the secretion is as intermittent as gestation

itself.

We have noticed, as one of the signs of pregnancy, that the breasts become enlarged and turgid, denoting the aptitude for the formation of the fluid; and it not unfrequently happens, that, towards the middle and latter periods of pregnancy, milk will distil from the nipples. This fluid, however, as well as that which flows from the breasts during the first two or three days after delivery, differs somewhat from milk, containing more serum and butter, and less caseum, and it is conceived to be more laxative, so as to aid the expulsion of the meconium. This first milk is called colostrum, protogala, &c., and, in the cow, constitutes the biestings or beastings. Generally, about the third day after confinement, the mammæ become tumid, hard, and even painful, and the secretion from this time is established, the pain and distension soon disappearing.

It is hardly necessary to discuss the views of Richerand, who considers the milk to be derived from the lymph; of others, who derive it from the chyle; of Raspail, who is disposed to think, that the mammary glands are in connexion, by media of a communication yet unknown, with the mucous surface of the stomach, and that they subtract, from the alimentary mass, the salts and

a Narrative of a Journey to the Polar Sea, p. 157.

Nouveaux Elémens de Physiologie, 7ème édit., Paris, 1817.

b For similar cases, see C. W. Mehliss, Ueber Virilescenz und Rejuvenescenz thierischer Körper, s. 41 & 71, Leipz. 1838; Belloc, Cours de Médec. Légale, p. 52, Paris, 1819; Foderé, Traité de Médecine Légale, i. 440; Coxe's Medical Museum, i. 267; Beck's Medical Jurisprudence, 6th edit. i. 188, Philad. 1838; and Montgomery, on the Signs and Symptoms of Pregnancy, p. 70, London, 1837, or Dunglison's Amer. Medical Library Edition, Philadelphia, 1838.

organizing materials which enter into the composition of the milk; or of Girard of Lyons, who gratuitously asserts, that there is in the abdomen an apparatus of vessels, — intermediate between the uterus and mammæ, — which continue inactive, except during gestation, and for some time after delivery, but, in those conditions, are excited to activity. All these notions are entirely hypothetical, and there is no reason for believing, that this secretion differs from others, as regards the kind of blood from which it is separated. The separation takes place in the tissue of the gland, and the product is received by the lactiferous ducts, along which it is propelled by the fresh secretion continuously arriving, and by the contractile action of the ducts themselves, the milk remaining in the ducts and sinuses, until the mammæ are, at times, considerably distended and painful.

The excretion of the milk takes place only at intervals. When the lactiferous ducts are sufficiently filled, a degree of distension and uneasiness is felt, which calls for the removal of the contained fluid. At times, the flow occurs spontaneously; but, commonly, only when solicited either by sucking or drawing the breast, the secretion under such circumstances being very rapid, and the contraction of the galactophorous ducts such as to project the milk through the orifices in a thready stream.

Milk is a highly azoted fluid, composed of water, caseum, sugar of milk, certain salts, — as chloride of sodium, phosphate, and acetate of potassa, with a vestige of lactate of iron and earthy phosphate, — and a little lactic acid. According to Berzelius, cow's milk consists of cream, and milk properly so called, — the cream consisting of butter, 4.5; cheese, 3.5; whey, 92.0; — and the whey, of milk and salt, 4.4; — the milk containing water, 928.75; — cheese, with a trace of butter, 28.01; sugar of milk, 35.00; chloride of potassium, 1.70; phosphate of potassa, 0.25; lactic acid, acetate of potassa, and lactate of iron, 6.00; and phosphate of lime, 0.30.°

Raspaild defines milk to be an aqueous fluid, holding albumen and oil in solution by means of an alkali, or alkaline salt, which he suggests may be the acetate of ammonia, — and, in suspension, an immense number of albuminous and oleaginous globules. The following table exhibits the discrepant results of the investigations of Brisson, Boyssou, Stipriaan Luiscius and Bondt, Schübler, and John, in 1000 parts of the milk of different animals — as given by Burdach.

Adelon's Physiologie de l'Homme, 2de édit. iv. 141, Paris, 1839.

b Medico-Chirurgical Transactions, vol. iii.

c See, on the Microscopic Examination, &c. of the Milk, Donné, Du Lai, et en particulier de celui des Nourrices, &c., Paris, 1837, or notice thereof in Brit. and For. Med. Rev., July, 1838, p. 181. See, also, Donné, Comptes rendus, Sept. 18, 1839, or Dunglison's Amer. Med. Intelligencer, Jan. 1, 1840, p. 306; and Nasse, Müller's Archiv. 1840, Heft iii., or Brit. and For. Med. Rev., Jan. 1841, p. 228.

d Chimie Organique, p. 345, Paris, 1833.

Physiologie als Erfahrungswissenschaft, B. ii. and v. 2te Auflage, s. 259, Leipz.
 1833. See, also, Carpenter, Human Physiology, § 684, Lond. 1842.

Observers.	Specific gravity.	Butter.	Cheese.	Sugar of milk.	Water.	Extract.
≝ (Brisson,	10409	38-24	51.26	20.73	886-19	3.45
Boyssou,	10350	58.12	153.75	41.87	746.25	11.07 11120
Brisson, Boyssou, Luiscius, John,	10350	54.68	31.25	39.06	875.00	ile io
Brisson,	10324	a language	16.81.561	AND N	000.00	3.45
Brisson, Boyssou, Luiscius, Schübler,		24.88	39.40	31.33	900.92	3.49
Luiscius,	10280	26.87	89.37	30.62	853-12	34199111
Schübler,		24.00	50.47	77.00	848-53	1221500
John,	Serial and Serial	23.43	93.75	39-06	843.75	Sand
Brisson, Boyssou, Luiscius, John,	10341	72822		00.00	000.05	3.45
Boyssou,	-	29.95	52.99	20.73	892-85	5.40
Luiscius,	10360	45.62	91.25	43.75	819·37 849·39	TO STATE OF
g ( John,	The faile	11.71	105.45	23.43	849.99	Om had
Brisson, Boyssou, Luiscius, John,	10364		-			10.00
Boyssou,		0.57	18.43	32.25	938-36	10.36
2 Luiscius,	10450	0.00	16.25	87.50	896-25	William He
g ( John,	San your	0.00	64.84	35-15	900.00	PRONO
≝ ( Brisson,	10355	Party Lab	LYN HA	THE REAL PROPERTY.	1000	0.01
Boyssou,	A SPHES	0.92	19-58	39.97	932-60	6.91
Brisson, Boyssou, Luiscius, John,	10230	0.00	33.12	45.00	921.87	halata
~ ( John,	B 54 34	0.00	11.71	46.87	941.40	(IS)(III)
Brisson,	10203	San Ball	1	The second of	-	0.00
Boyssou,		32.25	11.52	46.08	903.92	6.91
Luiscius,	10250	30.00	26.87	73-12	870.00	1 mg
John,	the secretary	23.43	15.62	39.06	921.87	7

From this table, an approximation may be made, as to the main differences between the milk of those animals, but it is not easy to explain the signal discrepancy amongst observers as to the quantity of the different materials in the milk of the same animal. Much, of course, may be dependent upon the state of the milk at the time of the experiment, but this can scarcely account for the whole discrepancy.

From a great number of experiments, MM. Deyeux and Parmentier<sup>b</sup> classed six kinds of milk, which they examined, according to the following table, as regards the relative quantity of the ma-

terials they contained.

Caseum. Butter.		Sugar of Milk.	Serum.	
Goat. Sheep. Cow.	Sheep. Cow. Goat.	Woman. Ass. Mare.	Ass. Woman.	
Ass.	Woman.	Cow.	Mare.	
Woman. Mare.	Ass. Mare.	Goat. Sheep.	Goat. Sheep.	

<sup>\*</sup> See, also, MM. O'Henry and A. Chevalier, Journal de Pharmacie, Juin et Juillet, 1839; M. Lecanu, ibid., Avril, 1839; MM. D'Arcet et Petit. Revue Médicale, Fév. ou Mars, 1839; and Sir Astley Cooper and Dr. G. Bird in Sir Astley Cooper on the Anatomy of the Breast, Lond. 1840.

b Précis d'Expér., &c. sur les différentes espèces de Lait, Strasbourg, an vii. 1790.

Human milk, therefore, contains more sugar of milk and less cheesy matter than that of the cow; hence it is sweeter, more liquid, less coagulable, and incapable of being made into cheese. It has also albuminous, oleaginous, and saccharine ingredients combined, so as to adapt it admirably as an aliment for the young; and of all the secreted fluids it appears to be most nearly allied to blood in its composition.<sup>a</sup>

M. Romanet<sup>b</sup> has lately affirmed, that the globules in cow's milk are wholly formed of butter, which exists as a pulp, enveloped in a white, translucent, elastic and resistent pellicle; and that the cyst is broken in churning, by which the butter escapes, and the pellicles floating about separately constitute the white particles

that give consistence to buttermilk.

When human milk is first drawn, it is of a bluer colour than that of the cow. It resembles rather whey, or cow's milk much diluted with water. If allowed to rest, it separates, like the milk of other animals, into cream and milk,—the quantity of the cream being one-fifth to one-third of that of the milk. The milky portion, however, appears semitransparent like whey, instead of being white and opaque like that of the cow. During the first days of its remaining at rest, abundance of cream and a little curd are separated, and lactic acid is developed. The specific gravity of human milk was found by Dr. Rees to be 1.0358, and the solid contents 12 per cent. The specific gravity of the cream was 1.021.°

Casein — the azoted constituent of milk — is distinguished from fibrin and albumen by its great solubility, and by not coagulating when heated. This is regarded by Liebigd as "the chief constituent of the mother's blood. To convert casein into blood no foreign substance is required, and in the conversion of the mother's blood into casein no elements of the constituents of the blood have

been separated."

The quantity and character of the milk differ according to the quantity and character of the food, —a circumstance, which was one of the greatest causes of the belief, that the lymphatics or chyliferous vessels convey to the mammæ the materials for the secretion. The milk is, however, situate in this respect like the urine, which varies in quantity and quality, according to the amount and kind of solid or liquid food taken. The milk is more abundant, thicker, and less acid, if the female lives on animal food, but possesses the opposite qualities when vegetable diet is used. It is apt, also, to be impregnated with heterogeneous matters, taken up from the digestive canal. The milk and the butter of cows indicate unequivocally the character of their pasturage, especially if they have fed on the turnip, wild onion, &c. Medicine, given to

d Animal Chemistry, Amer. Edit. p. 51, Cambridge, 1842.

Dr. G. O. Rees, art. Milk, Cyclop. of Anat. and Physiol.

b Comptus rendus, Avril, 1842. Cooper, op. cit.

the mother, may in this way act upon the infant. Serious almost fatal - narcotism was induced in the infant of a professional friend of the author, by a dose of morphia administered to his wife.

It would seem, that occasionally the secretion of the two glands differs. A case has been related in which the milk of the right breast had a distinctly salt taste, whilst that of the opposite breast

was of the ordinary sweetness.b

The quantity of milk secreted is not always in proportion to the bulk of the mammæ: a female whose bosom is of middle size often secretes more than another in whom it is much more developed: - the greater size being usually owing to the larger quantity of adipous tissue surrounding the mammary gland, and this tissue is in nowise concerned in the function. The ordinary quantity of milk that can be squeezed from either breast at any one time, and which must consequently have been contained in its tubes and

reservoirs, is about two ounces.c

The secretion of milk usually continues until the period when the organs of mastication of the infant have acquired the necessary development for the digestion of solid food: it generally ceases during the second year. For a great part, or the whole of this time, the menstrual flux is suspended; and if both the secretions, - mammary and menstrual, - go on together, the former is usually impoverished, and in small quantity. This, at least, is the general opinion. Some, however, think, that no general rule can be established on the subject; and that the condition of the child's health ought to be the only guide. M. Gendrin would on no account permit a woman to continue nursing after the catamenia had returned. The subject has recently been closely investigated by M. Raciborski,d who laid the results before the Academie Royale de Médecine on the 31st of May, 1843. The inferences are, - that, contrary to generally received opinions, the milk of nurses who menstruate during suckling does not differ sensibly, in physical, chemical or microscopic characters, from that of nurses whose catamenia are suspended; - that the only difference, which can be detected between these two kinds of milk, is, that in most cases the milk of menstruating nurses contains less cream during the menstrual periods than in the intervals between those periods; hence arises the bluish appearance presented occasionally by such milk; and that a nurse should never be rejected merely because she menstruates.

Whilst lactation continues, the female is less likely to conceive;

Sciences, Oct. 1843, p. 455.

a See, on this subject, and also for an analysis of the Milk, Simon, Journal de Pharmacie, Juin, 1839, and Sir Astley Cooper on the Anatomy of the Breast, Lond. 1840. b Hartmann, in Mittheilungen aus dem Archiv. der Gesellschaft practischer Aerzte zu Riga, Ie sammlung. Leipz. Riga & Mitau, 1839, cited in Brit. and For. Med. Rev. Oct. 1841, p. 430. c Carpenter, Human Physiology, § 692, Lond. 1842. d Dublin Medical Press, Aug. 2d, 1843; cited in the American Journal of the Med.

and hence the importance, — were there not even more weighty reasons, — of the mother's suckling her own child, in order to prevent the too rapid succession of children. From observations, made at the Manchester Lying-in Hospital, on one hundred and sixty married women, Mr. Roberton<sup>a</sup> concludes, that in seven out of eight women, who suckle for as long a period as the working classes in England are in the habit of doing — about fifteen and a half months on the average — there will be an interval of fifteen months between parturition and the commencement of the subsequent pregnancy; and that, in a majority of instances, when suckling is prolonged to even nineteen or twenty months, pregnancy does not take place till after weaning. When menstruation recurs during suckling, it is an evidence that the womb has, again, the organic activity, which befits it for impregnation.

## CHAPTER II.

## FŒTAL EXISTENCE. - EMBRYOLOGY.

The subject of feetal existence forms so completely a part of the function just considered, that its investigation naturally succeeds that of the part performed by the parents in its production, and more especially as the development of the ovum is synchronous with the uterine changes that have been pointed out.

## 1. SPECIAL ANATOMY AND HISTOLOGY OF THE FŒTUS.

a. Dependencies of the Fætus.—These are the parts of the ovum, that form its parietes, attach it to the uterus, connect it with the fætus, and are inservient to the nutrition and development of the new being.

They are generally conceived to consist, — First, of two mem branes, according to common belief, which constitute the parietes of the ovule, and which are concentric; the outermost, called the chorion, — the innermost, filled with a fluid, in which the fætus is placed, and called the amnion or amnios. By Boer and Granville, a third and outer membrane has been admitted, — the cortical membrane or cortex ovi. Secondly, of a spongy, vascular body, situate without the chorion, covering about one-quarter of the ovule, and connecting it with the uterus, — the placenta. Thirdly, of a cord of vessels, — extending from the placenta to the fætus, the body of which is penetrated at the umbilicus, by the vessels, called the umbilical cord or navel string; and lastly, of three vesicles — the umbilical, allantoid, and erythroid, which are considered to be concerned in fætal nutrition.

<sup>&</sup>lt;sup>a</sup> Edinb. Med. and Surg. Journal, Jan. 1837. See, on the same subject, Mr. Laycock, Dublin Med. Press, Oct. 26, 1842, or Amer. Journ. Med. Sciences, Jan. 1843, p. 184.

<sup>b</sup> Graphic Illustrations of Abortion, Lond. 1834.

1. The chorion - which has received various names - is the outermost of the membranes of the ovum. About the twelfth day after conception, according to Velpeau, it is thick, opaque, resisting, and flocculent at both surfaces. These flocculi, in the part of the ovum that corresponds to the tunica decidua reflexa, aid its adhesion to that membrane; but, in the part where the ovum corresponds to the uterus, they become developed to constitute the placenta. At its inner surface, the chorion corresponds to the amnion. These two membranes are, however, separated during the earliest period of fœtal existence, by a gelatinous or albuminous fluid; but at the expiration of three months, the liquid disappears and they are afterwards in contact. By many anatomists, the chorion is conceived to consist originally of two laminæ; and by Burdachd these have been distinguished by different names; the outer lamina being called by him exochorion; the inner endochorion. Velpeau denies this, and asserts, that he has never been able to separate them, even by the aid of previous maceration.

As the placenta is formed on the uterine side of the chorion, the membrane is reflected over the fætal surface of that organ, and is continued over the umbilical cord, as far as the umbilicus of the fætus, where it is confounded with the skin, of which it consequently appears to be a dependence. As pregnancy advances, the chorion becomes thinner, and less tenacious and dense, so that at the full period, it is merely a thin, transparent, colourless membrane, much more delicate than the amnion. Haller, Blumenbach and Velpeau affirm it to be devoid of vessels; but, according to Wrisberg, it receives some from the umbilical trunks of the fœtus, and, according to others, from the decidua. Dutrochet conceives it to be an extension of the fætal bladder. Its vascularity, according to Dr. Granville, is proved by its diseases, which are chiefly of an inflammatory character, ending in thickening of its texture; and he affirms, that there is a preparation in the collection of Sir Charles Clarke, which shows the vessels of the chorion as evidently as if they were injected.

2. The amnion lines the chorion concentrically. It is filled with a serous fluid, and contains the fætus. In the first days of fætal existence, it is thin, transparent, easily lacerable, and somewhat resembling the retina. At first, it adheres to the chorion only by a point, which corresponds to the abdomen of the fætus; the other portions of the membranes being separated by the fluid already mentioned, called the false liquor amnii. Afterwards, the membranes coalesce, and adhere by very delicate cellular fila-

e Purkinje, art. Ei, Encyclopäd. Wörterbuch der Medicinisch. Wissenschaft. x. 149, Berlin, 1834.

Haller, Element. Physiol. viii. 188; Burdach's Physiologie als Erfahrungswissenschaft, ii. 57; and Velpeau, Embryologie, Paris, 1833.
 Embryologie ou Ovologie Humaine, Paris, 1833.

e See, also, Weber's Hildebrandt's Handbuch der Anatomie, iv. 492, Braunschweig, 1832; and Purkinje, art. Ei, in Encyclop. Wörterb. der Medicin. Wissensch. x. 144, Berlin, 1834.

ments; but the adhesion is feeble, except at the placenta and umbilical cord. In the course of gestation, this membrane becomes thicker and tougher; and, at the full period, it is more tenacious than the chorion, elastic, semitransparent and of a whitish colour. Like the chorion, it covers the fætal surface of the placenta, envelopes the umbilical cord, passes to the umbilicus of the fætus, and

commingles there with the skin.a

It has been a question, whether the amnion be supplied with bloodvessels. Velpeau denies it: Haller and others have maintained the affirmative. Haller asserts, that he saw a branch of the umbilical artery creeping upon it. The fact of the existence of a fluid within it, which is presumed to be secreted by it, would also greatly favour the affirmative. But, admitting that it is supplied with bloodvessels, a difference has existed with regard to the source whence they proceed; and anatomical investigation has not succeeded in dispelling it. Monro affirms, that on injecting warm water into the umbilical arteries of the fætus, the water oozed from the surface of the amnion. Wrisberg asserts, that he noticed the injection to stop between the chorion and amnion; and Chaussier obtained the same results as Monro, by injecting the vessels of the mother.

The amnion contains a serous fluid, the quantity of which is in an inverse ratio to the size of the new being; so that its weight may be several drachms, when that of the fœtus is only a few grains. At first, the liquor amnii, - for so it is called, - is transparent; but, at the full period, it has a milky appearance, owing to flocculi of an albuminous substance held in suspension by it. It has a saline taste, a spermatic smell, and is viscid and glutinous to the touch. Vauquelin and Bunivab found it to contain, water, 98.8; albumen, chloride of sodium, soda, phosphate of lime, and lime, 1.2. That of the cow, according to these gentlemen, contains amniotic acid; but Prout, Dulong, and Labillardiére and Lassaigne were not able to detect it. Dr. Rees analysed several specimens. He found its specific gravity to be about 1.007 or 1.008, and its mean composition in two cases at 7½ months to be as follows: water, 986.8; albumen (traces of fatty matter), 2.8; salts soluble in water, 3.7; albumen from albuminate of soda, 1.6; salts soluble in alcohol, 3.4; lactic acid, urea, 1.7. Total, 1000, The salts consisted of chloride of sodium, and carbonate of soda, with traces of alkaline sulphate and phosphate.c Dr. Vogtd analyzed it at two different periods of pregnancy, at 31 months and 6 months, and found the constituents to vary as follows:

Velpeau, Embryologie, &c. Paris, 1833.

b Annales de Chimie, tom. xxxiii.; and Memoir. de la Société Médicale d'Emulation, iii. 229.

c Ancell, Lectures on the Physiology and Pathology of the Blood, April 25, 1840, p. 154.

d Müller's Archiv.; and Brit. and For. Med. Review, July, 1838, p. 248.

VOL. II. - 38

Water	31 Months. 978.45	0 00	Sin can	6 Months. 990-29
Alcoholic extract, consisting of uncertain	3.69	10:31	de	0.34
Chloride of sodium	5.95		1000	2.40
Albumen (as residuum)	10.79	-		6.77
Sulphate and phosphate of lime, and loss -	0.14	-	No. 1	0.30
	1000-		It de	1000-
Specific gravity	1.0182			1.0092

No inferences can, however, be drawn from these cases as to the proportion of solid matters at different periods of utero-gestation, inasmuch as the subject of the first case died of an inflammatory disease; the other in a state of cachexia. Prouta found some sugar of milk in the liquor amnii of the human female; Berzelius detected fluoric acid in it; Scheele, free oxygen; b and Lassaigne, in one experiment, a gas resembling atmospheric air; in others, a gas composed of carbonic acid and azote. J. Müller, however, was never able to detect oxygen in it. The chemical history of this substance is, consequently, sufficiently uncertain, nor is its origin placed upon surer grounds; - some physiologists ascribing it to the mother, others to the fœtus; - opinions fluctuating, according to the presumed source of the vessels, that supply the amnion with arterial blood. It has even been supposed to be the transpiration of the fœtus, or its urine. One reply to these views is, that we find it in greater relative proportion when the fœtus is small. Meckel thinks, that it chiefly proceeds from the mother, but that, about the termination of pregnancy, it is furnished in part by the fætus. The functions, however, to which, as we shall see, it is probably inservient, would almost constrain us to consider it a secretion from the maternal vessels; and what perhaps favours this notion is the fact, that if a female be made to take rhubarb for some time prior to parturition, the liquor amnii will be found tinged with it.e It is interesting, also, to recollect, that, in the experiments of Dr. Blundell, - which consisted in obliterating the vulvo-uterine canal in rabbits, and, when they had recovered from the effects of the injury, putting them to the male, - although impregnation did not take place, the wombs, - as in extra-uterine pregnancy, - were evolved, and the waters collected in the uterus. The fluid, consequently, must, in these cases, have been secreted from the interior of the uterus. May not the liquor amnii be secreted, in this manner, throughout the whole of gestation, and pass through the membranes of the ovum by simple imbibition? and may not the fluid secretions of the fœtus, which are discharged into the liquor amnii, pass through the membranes, and enter the system of the mother, in the same way?

Annals of Philosophy, v. 417.

b Dissert. de Liquoris Amnii Arteriæ Asperæ Fætuum Humanorum Naturâ et Usu, &c., Copenh. 1799.

c Archiv. Général. de Méd., ii. 308.
d Handbuch der Physiologie, i. 305, Berlin, 1833.

<sup>·</sup> Granville, Graphic Illustrations of Abortion, p. xxi., Lond. 1834.

The quantity of the liquor amnii varies in different individuals, and in the same individual at different pregnancies, from four ounces to as many pints. Occasionally, it exists to such an amount as to throw obscurity even over the very fact of pregnancy. An instance of this kind, strongly elucidating the necessity of the most careful attention on the part of the practitioner in such cases, occurred in the practice of a respectable London practitioner, — a friend of the author. The abdomen of a lady had been for some time enlarging by what was supposed to be abdominal dropsy: fluctuation was evident, yet the case appeared to be equivocal. A distinguished accoucheur, and a surgeon of the highest eminence, were called in consultation, and after examination the latter declared, that "it was an Augean stable, which nothing but the trocar could clear out." As the lady, however, was even then complaining of intermittent pain, it was deemed advisable to make an examination per vaginam. The os uteri was found dilated and dilating, and in a few hours after this formidable decision, she was delivered of a healthy child, the gush of liquor amnii being enormous. After its discharge, the lady was reduced to the natural size, and the dropsy, of course, disappeared.

3. The cortical membrane or cortex ovi is, according to Boer and Granville, the one, which is usually regarded as a uterine production, and denominated the decidua reflexa. It surrounds the ovule when it descends into the uterus, and envelopes the shaggy chorion. This membrane is destined to be absorbed during the first months of utero-gestation, so as to expose the next membrane to the contact of the decidua, with which a connexion takes place in the part where the placenta is to be formed. In that part, Boer and Granville consider, that the cortex ovi is never altogether obliterated, but only made thinner; and in process of time it is converted into a mere pellicle or envelope, which not only serves to divide the filiform vessels of the chorion into groups or cotyledons, in order to form the placenta, but also covers those cotyledons. This Dr. Granville calls the membrana propria.

4. Placenta. — This is a soft, spongy, vascular body, formed at the surface of the chorion, adherent to the uterus, and connected with the fætus by the umbilical cord. The placenta is not in existence during the first days of the embryo state; but its formation commences, perhaps, with the arrival of the embryo in the uterus. In the opinion of some, the flocculi, which are at first spread uniformly over the whole external surface of the chorion, gradually congregate from all parts of the surface into one, uniting with vessels proceeding from the uterus, and traversing the decidua, to form the placenta; the decidua disappearing from the uterine surface of the placenta about the middle of pregnancy, so that the latter comes into immediate contact with the uterus. In the opinion of others, the placenta is formed by the separation of the layers of the chorion, and by the development of the different vessels, that

A Graphic Illustrations of Abortion, part iv. Lond. 1835.

creep between them. Velpeau<sup>a</sup> maintains, that the placenta forms only at the part of the ovule, which is not covered by the true decidua, and which is immediately in contact with the uterus; and that it results from the development of the granulations that cover this part of the chorion; these granulations or villi, according to Velpeau, being gangliform organs containing the rudiments of the placental vessels. Others, again, regard it as formed by the growth of the vessels of the uterus into the decidua serotina.

The mode in which the placenta is attached to the uterus has always been an interesting question with physiologists; and it has been revived, of late, by Messrs. Lee, b Radford, and others. A common opinion has been, that the large venous canals of the nterus are uninterruptedly continuous with those of the placenta. Wharton and Reuss,d and a number of others, conceive that, at an early period of pregnancy, the part of the uterus, in contact with the ovum, becomes fungous or spongy, and that the fungosities, which constitute the uterine placenta, commingle and unite with those of the chorion so intimately, that laceration necessarily occurs when the placenta is extruded; and Dubois goes so far as to consider milk fever as a true traumatic disease, produced by such rupture! The opinion of Messrs. Lee, Radford, Velpeau and others is, that the maternal vessels do not terminate in the placenta; but that apertures - portions scooped out, as it were, - exist in their parietes, which are closed up, according to the two first gentlemen. by the true decidua; - according to Velpeau, by a membranule or anorganic pellicle, which he conceives to be thrown out on the fungous surface of the placenta, or by some valvular arrangement, the nature of which has not been discovered; but these apertures have no connexion, in his opinion, with any vascular orifice, either in the membrane or the placenta. The mode, therefore, in which these authors consider the placenta to be attached to the uterus is, so far as it goes, somewhat unfavourable to the idea generally entertained, that the maternal vessels pour their fluid into the maternal side of the placenta, whence it is taken up by the radicles of the umbilical vein. Whatever blood is exhaled must necessarily pass through the decidua, according to Lee and Radford; or through the pellicle, according to Velpeau. More recently, Dr. Leee has somewhat modified his views, and now believes, that the circulation in the human ovum, in the third month of gestation, is carried on in the following manner: - The maternal blood is conveyed by the arteries of the uterine decidua into the interstices of the placenta and villi of the chorion. The blood, which has circulated in the placenta, is returned into the veins of the uterus by the oblique

2 Embryologie ou Ovologie Humaine, p. 63.

on the Structure of the Human Placenta, Manchester, 1837.

c London Medical Gazette, Dec. 1838; or Amer. Journ. of the Med. Sciences, May, 1839, p. 189.

Philosoph. Transactions for 1832; and Remarks on the Pathology and Treatment of some of the most important Diseases of Women, Lond. 1833.

<sup>6</sup> Novæ quædam Observationes Circa Structuram Vasor. in Placent. Human. et peculiarem hujus cum Utero Nexum, Tubing. 1784.

openings in the decidua covering the placenta. The blood, which has circulated between the villosities of the chorion, passes through the openings in the decidua reflexa into the cavity between the two deciduous membranes, whence it is taken up by the numerous apertures and canals that exist, according to him, in the uterine decidua, and so passes into the veins of the uterus. Biancini' maintains, that a number of flexuous vessels connect the uterus directly with the placenta, which are developed immediately after the period of conception. These utero-placental vessels, he says, are not prolongations of the uterine vessels, but a new production. Recently, Dr. John Reidb has carefully examined into this point of anatomy. On separating the adhering surfaces of the uterus and placenta, cautiously under water, he satisfied himself, but not without considerable difficulty, of the existence of the utero-placental vessels described by the Hunters. After a portion of the placenta had been detached in this manner, Dr. Reid's attention was attracted towards a number of rounded bands passing between the uterine surface of the placenta and the inner surface of the uterus, several of which could be drawn out in the form of tufts from the mouths of the uterine sinuses. On slitting up some of the uterine sinuses with the scissors, these tufts could be seen ramifying in their interior. These were ascertained to be prolongations of the fætal placental vessels, and to protrude into the open mouths of certain of the uterine sinuses, and in those placed next the inner surface of the uterus only. These tufts were surrounded externally by a

soft tube, similar to the soft wall of the utero-placental vessels, which passed between the margin of the open mouths of the uterine sinuses and the edges of the orifices in the decidua through which the tufts protruded into the sinuses. On examining the tufts, as they lay in the sinuses, it was evident, that though they were

Fig. 249.

examining the tufts, as they lay in the sinuses, it was evident that they that they they were

so far loose, and could be floated about, yet they were bound down firmly at various points by reflections of the inner coat of the venous system of the mother upon their outer surface. Dr. Reid farther satisfied himself that the interior of the placenta is composed of numerous trunks and branches, each including an artery, and an accompanying vein, every one of which, he believes, is closely ensheathed in prolongations of the inner coat of the vascular system of the mother, or at least in a membrane continuous with it. According to this view of the structure of the placenta, the inner coat of the vascular system of the mother is prolonged over each individual tuft, so that when the blood of the

b Edinb. Med. and Surg. Journal, Jan. 1841, p. 4.

<sup>&</sup>lt;sup>a</sup> Sul Commercio Sanguigno tra la Madre e il Feto, Pisa, 1833.

mother flows into the placenta through the curling arteries of the uterus, it passes into a large sac formed by the inner coat of the vascular system of the mother, which is intersected in many thousands of different directions, by the placental tufts projecting into it like fringes, and pushing its thin wall before them in the form

Fig. 250.

Connection between the Maternal and Fatal Vessels.

a. Curling artery. b. Uterine vein. c. Placenta, d. Placental tufts with inner coat of vascular system of the mother enveloping them. — (J. Reid.)

of sheaths, which closely envelope both the trunk and each individual branch composing these tufts. From this sac, the maternal blood is returned by the utero-placental veins without having been extravasated, or without having left the maternal system of vessels. Into this sac in the placenta, containing the blood of the mother, the tufts of the placenta hang like the bronchial vessels of certain aquatic animals, to which they have a marked analogy. This sac is protected and strengthened on the fætal surface of the placenta by the chorion, on the uterine surface by the decidua vera, and

on the edges or margin by the decidua reflexa. In this view, the fætal and maternal portions are every where intimately intermixed with tufts of minute placental vessels, their blunt extremities being found lying immediately under the chorion covering its fætal surface, as well as towards its uterine surface. The discovery of the prolongations of the fetal placental vessels into some of the uterine sinuses, Dr. Reid thinks, is principally valuable, as it presents us with a kind of miniature representation of the whole structure of the placenta; and the reason why the placental tufts are not perceptible on the uterine surface of the placenta expelled in an accouchement is, that they are so strongly bound down by the reflection of the inner coat of the uterine sinuses that they are torn across. Professors Alison, Allen Thomson, and J. Y. Simpson inspected the preparations of Dr. Reid, and expressed themselves satisfied that the placental tufts were prolonged into the uterine sinuses, and that the inner coat of the veins was prolonged upon them. Dr. Sharpey, too, confirms the views of Dr. Reid from his own observation of impregnated uteri; and Dr. Churchilla states, that in a recent visit to Edinburgh, Dr. Reid showed him one of the portions of uterus and placenta on which his investigations were made, and there was no difficulty in demonstrating the tufts dipping into the uterine sinuses. "No doubt," he adds, "farther observations are necessary for the perfect elucidation of the subject; but I certainly think, that as far as our knowledge extends, it is in favour of the opinion adopted by Dr. Reid and the later physiologists."

A somewhat similar view to that of Dr. Reid is entertained by Prof. E. H. Weber.

<sup>&</sup>lt;sup>a</sup> The Theory and Practice of Midwifery, Amer. Edit. by Prof. Huston, p. 110, Philad. 1843.

b Hildebrandt's Handbuch der Anatomie des Menschen, iv. 496, Braunschweig, 1832; Wagner, op. cit. p. 201, and Dr. Reid, op. cit. p. 11. See, also, Mr. Dalrymple, Medico-Chirurg. Transactions, vol. xxv., Lond. 1842.


In whatever manner originally produced, the placenta is distinguishable in the second month, at the termination of which it covers two-thirds, or at the least, one-half of the ovum; after this, it is observed to go on successively increasing. Prior to the full, term, however, it is said to be less heavy, more dense, and less vascular, owing - it has been conceived - to several of the vessels that formed it having become obliterated and converted into hard, fibrous filaments; a change, which has been regarded as a sign of maturity in the fætus, and a prelude to its birth. At the full period, its extent has been estimated at about one-fourth

of that of the ovum; its diameter from six to nine inches; its circumference twenty-four inches; its thickness from an inch to an inch and a half at the centre, but less than this at the circumference; and its weight, with the umbilical cord and membranes, from twelve to twenty ounces. All this is subject, however, to much variation. It is of a circular shape, and the cord is usually inserted into its centre. It may be attached to any part

two surfaces, that which corresponds to the uterus is divided into irregularly rounded lobes or cotyledons, and it is covered by a soft and delicate cellulo-vascular membrane, which by many is considered to be the decidua vera.b Wrisberg,c Lobstein,d and Desormeaux,e however, who consider, that the decidua disappears from behind the placenta about the fourth or fifth month, regard it as a new membrane; and Bo-



of the uterus, but is usually found towards the fundus.



a Burdach's Physiologie als Erfahrungswissensch. ii. 403.

b See Dr. John Reid, Edinb. Med. and Surg. Journal, Jan. 1841, p. S.

· Observ. Anat. Obstetric. de Structura Ovi et Secundinar. Human. &c. Gotting. d Essai sur la Nutrition du Fœtus, Strasbourg, 1802.

· Art. Œuf Humain, in Dict. de Médecine.

janus, believing it to be produced at a later period than the decidua vera, gives it the name of membrana decidua serotina. (See Fig. 236, page 413 of this volume.) Breschet, again, maintains that two laminæ—decidua vera and decidua reflexa— are found intervening between the uterus and placenta, whilst Velpeau maintains that the true decidua never exists there!

The fætal or umbilical surface is smooth, polished, covered by the chorion and amnion, and exhibits the distribution of the umbilical vessels, and the mode in which the cord is attached to the

organ.

The following are the anatomical constituents of the placenta, as usually described by anatomists. First. Bloodvessels, from two sources, the mother and the fœtus. The former proceed from the uterus, and consist of arteries and veins, of small size but considerable number. The vessels, which proceed from the fœtus, are those that constitute the umbilical cord; - viz. the umbilical vein, and the umbilical arteries. These vessels, after having penetrated the fætal surface of the placenta, divide in the substance of the organ, so that each lobe has an arterial and a venous branch, which ramify in it, but do not anastomose with the vessels of other lobes. Secondly. Expansions of the chorion, which are described by some as dividing into cellular sheaths, and accompanying the vessels to their final ramifications; - an arrangement which is, however, contested by others. Thirdly. White filaments, which are numerous in proportion to the advancement of pregnancy, and which seem to be obliterated vessels. Fourthly. A kind of intermediate cellular tissue, serving to unite the vessels together, and which has been regarded, by some anatomists, as an extension of the decidua accompanying those vessels. Lastly. A quantity of blood poured into this intermediate cellular tissue. which may be removed by washing. In addition to these constituents, a glandular structure has been presumed to exist in it; as well as lymphatic vessels.c Fohmann,d affirms, that the umbilical cord, in addition to the bloodvessels, consists solely of a plexus of absorbents, which may be readily injected with mercury. This has been done also by Dr. Montgomery, of Dublin. These lymphatics of the cord communicate with a network of lymphatics, seated between the placenta and the amnion, the termination of which Fohmann could not detect, but he thinks they pass to the uterine surface of the placenta. These vessels proceed to the umbilicus of the child, and chiefly unite with the subcutaneous lymphatics of the abdominal parietes; follow the superficial veins; pass under the crural arches; ramify on the iliac glands; and terminate in the thoracic duct. Lobstein and Meckel say that they have never been able to detect lymphatics in the cord.

a Isis, von Oken, für 1821.

Mémoir. de l'Académ. Royal. de Médec. tom. ii. Paris, 1833.
 Granville, op. citat. p. xix.

d Sur les Vaisseaux Absorbans du Placenta, &c. Liège, 1832; and Amer. Journ. May, 1835, p. 174.

Chaussier and Ribes, and Mr. Cæsar Hawkins describe nerves in the placenta proceeding from the great sympathetic of the fætus.

The uterine and the fætal portions of the placenta are generally described as quite distinct from each other, during the two first months of fætal life; but afterwards they constitute one mass. Still, the uterine vessels remain distinct from the fætal; the uterine arteries and veins communicating freely with each other, as well as the fætal arteries and veins; but no direct communication existing between the maternal and fætal vessels. Until of late, almost every obstetrical anatomist adopted the division of the placenta into two parts, constituting - as it were - two distinct placentæ, - the one maternal, the other fætal. Messrs. Lee, Radford, and others have, however, contested this point, and have affirmed, with Velpeau, that the human placenta is entirely feetal. The very fact, indeed, of the existence of a membrane, or — as M. Velpeau calls it - a "membranule," between the placenta and the uterus, would destroy the idea of any direct adhesion between the placenta and uterus, and make the placenta wholly fætal. Yet the point is still contested, — by those especially, who consider that the maternal vessels ramify on one surface of the placenta, and the fætal on the other.c

It is generally supposed, that the placenta is most frequently attached to the right side of the uterus, but Nägeled found the opposite to be the fact in his examinations. In six hundred cases, which he carefully auscultated, the placenta was found in two hundred and thirty-eight cases on the left side, and in one hundred and forty-one on the right.

5. Umbilical cord. — From the fætal surface of the placenta a cord of vessels passes, which enters the umbilicus of the fætus, and has hence received the name umbilical cord, as well as that of navel string. It forms the medium of communication between the fætus and the placenta. During the first month — Pockelse says the first three weeks — of fætal existence, the cord is not perceptible; the embryo appearing to be in contact, by the anterior part of its body, with the membranes of the ovum. Such, at least, is the description of most anatomists; but Velpeauf says it is erroneous. The youngest embryo he dissected had a cord. At a fortnight and three weeks old, the length is three or four lines; and he thinks his examinations lead him to infer, that at every period of fætal development, the length of the cord is nearly equal to that of the body, if it do not exceed it a little.

von Oken, 1825.

a Journal Universel des Sciences Médicales, i. 233.

b Sir E. Home, Lect. on Comp. Anat. v. 185, Lond. 1828; Lond. Lancet, June, 1833; and Messrs. Mayo and Stanley, Report on the Preparations of Impregnated Uteri in the Hunterian Museum, Lancet, June 22, 1833.


See Weber's Hildebrandt's Handbuch der Anatomie, iv. 495, Braunschweig, 1832.
 Die Geburtshülfliche Auscultation, Mainz, 1838; cited in Brit. and For. Med.

Rev., Oct. 1839, p. 371.

• Neue Beiträge zur Entwickelungsgeschichte des Menschlichen Embryo, in Isis,

<sup>1</sup> Embryologie, ou Ovologie Humaine, Paris, 1833.

In an embryo, a month old, Béclarda observed vessels creeping, for a certain space, between the membranes of the ovum, from the abdomen of the fœtus to a part of the chorion, where the rudiments of the future placenta were visible. During the fifth week, the cord is straight, short, and very large, owing to its containing a



Knotted Umbilical Cord.

portion of the intestinal canal. It presents, also, three or four dilatations. separated by as many contracted portions or necks: but these gradually disappear; the cord lengthens, and becomes smaller, and occasionally it is twisted, knotted. and tuberculated in a strangely inexplicable manner.

(Fig. 253.) After the fifth week it contains - besides the duct of the umbilical vesicle - the omphalo-mesenteric vessels, and a portion of the urachus, or of the allantoid, and of the intestines. At about two months, the digestive canal enters the abdomen: the urachus, the vitelline canal - to be mentioned presently - and the vessels become obliterated, so that, at three months, as at the full period, the umbilical cord is composed of three vessels,the umbilical vein, and two arteries of the same name, - of a peculiar jelly-like substance, and it is surrounded, as we have seen, by the amnion and chorion. The vessels will be more particularly described hereafter. They are united by a cellular tissue, containing the jelly of the cord, or of Wharton, a thick albuminous secretion, which bears some resemblance to jelly, and the quantity of which is very variable. In the fœtus, the cellular tissue is continuous with the sub-peritoneal cellular tissue; and in the placenta, it is considered to accompany the ramifications of the vessels. The length of the cord varies, at the full period, from eight inches to fifty-four. The most common length is eighteen inches.b

It has been already remarked, that Chaussier, Ribes, and Hawkins have traced branches of the great sympathetic of the fœtus as far as the placenta; and the same has been done by Durr, Rieck,d

and others.

6. Umbilical vesicle. - This vesicle, called also vesicula alba and intestinal vesicle, appears to have been first carefully observed

a Embryologie, ou Essai Anatomique sur le Fétus Humain, Paris, 1821.

b Dr. Churchill, in Dublin Journal of Med. Science, March, 1837. Dissert. Sistens Funicul. Umbilic. Nervis Carere, Tubing. 1815. d Utrum Funiculus Umbilicalis Nervis polleat aut careat. Tubing. 1816.

by Albinus.a Dr. Granville, however, ascribes its discovery to Bojanus, whilst others have assigned it to Diemerbroeck.d It was unknown to the ancients; and, amongst the moderns, is not universally admitted to be a physiological condition. Osiander and Döllinger class it amongst imaginary organs; and Velpeau remarks, that out of about two hundred vesicles, which he had examined, in fætuses under three months of development, he had met with only thirty in which the umbilical vesicle was in a state, that could be called natural. Under such circumstances, it is not easy to understand how he could distinguish the physiological from the pathological condition. If the existence of the vesicle be a part of the physiological or natural process, the majority of vesicles ought to be healthy or natural; yet he pronounces the thirty in the two hundred to be alone properly formed; and, of consequence, one hundred and seventy to be morbid or unnatural. This vesicle is described as a small pyriform, round or spheroidal sac; which, about the fifteenth or twentieth day after fecundation, is of the size of a common pea. It probably acquires its greatest dimensions in the course of the third or fourth week.






Diagram of the Fatus and Membranes about the sixth week.

a. Chorion b. The larger absorbent extremities, the site of the placenta. c. Allantois. d. Amnion. c. Urachus. é. Bladder. f. Vesicula umbilicalis. g. Communicating canal between the vesicula umbilicalis and intestine. h. Vena umbilicalis. ii. Arteria umbilicales. l. Vena omphalomeseraica. k. Arteria omphalo-meseraica. n. Heart. c. Rudiment of superior extremity. p. Rudiment of lower extremity.—(Carus.)

After a month, Velpeau always found it smaller. About the fifth sixth or seventh week it is about the size of a coriander seed.

Annotat. Académic. lib. i. p. 74.

b Graphic Illustrations of Abortion, p. xii. Lond. 1834.

· Meckel's Archiv. iv. s. 34; and Journ. Complément du Dict. des Sciences Médi-

cales, ii. 1818, p. 84.

d Opera, p. 304, Ultraject, 1672; see, also, Mackenzie, in the Edinburgh Medical and Surgical Journal for January, 1836, p. 46. For a list of those who have described it, see Purkinje, in art. Ei, of Encyclopad. Medicin. Wörterb. x. 156, Berlin, 1834; and Weber's Hildebrandt's Handbuch der Anatom. Band. iv.

After this, it becomes shrivelled and disappears insensibly. It seems to be situate between the chorion and amnion, and is commonly adherent either to the outer surface of the amnion, or to the inner surface of the chorion, but, at times, is situate loosely

between them. It is seen in Fig. 236 and Fig. 254.

The characters of the vitelline pedicle, as Velpeau terms it, which attaches the vesicle to the embryo, vary according to the stage of gestation. At the end of the first month, it is not less than two, nor more than six lines long, and about a quarter of a line broad. Where it joins the vesicle, it experiences an infundibuliform expansion. Its continuity with the intestinal canal appears to be undoubted. Up to twenty or thirty days of embryonic life, the pedicle is hollow, and, in two subjects, M. Velpeau was able to press the contained fluid from the vesicle into the abdomen, without lacerating any part. Generally, the canal does not exist longer than the expiration of the fifth week, and the obliteration appears to proceed from the umbilicus towards the vesicle. The parietes of the vitelline pouch — as M. Velpeau also calls it, from its analogy to the vitelline or yolk-bag of the chick — are strong and resisting; somewhat thick, and difficult to tear.

As the umbilical vesicle of brutes has been admitted to be continuous with the intestinal canal, anatomists have assigned it and its pedicle three coats. Such is the view of Dutrochet. Velpeau has not been able to detect these in the human fœtus. He admits,

however, a serous surface, and a mucous surface:

The vesicle is evidently supplied with arteries and veins, which are generally termed omphalo-mesenteric or omphalo-mesaraic, but, by Velpeau, vitello-mesenteric, or, simply, vitelline. The common belief is, that they communicate with the superior mesenteric artery and vein; but Velpeau says he has remarked, that they inosculate with one of the branches of the second or third order of those great vessels (canaux), — with those, in particular, that are distributed to the cæcum. These vessels he considers to be the vessels of nutrition of the umbilical vesicle.

The fluid, contained in the vesicle, which Velpeau terms the vitelline fluid, has been compared, from analogy, to the vitellus or yolk of egg. In a favourable case for examination, Velpeau found it of a pale yellow colour; opaque; of the consistence of a thickish emulsion; different in every respect from serosity, to which Albinus, Boerhaave, Wrisberg and Lobstein compared it, and from every other fluid in the organism; and he regards it as a nutritive substance—a sort of oil—in a great measure resembling that which

constitutes the vitelline fluid of the chick in ovo.f

7. Allantoid vesicle or allantois. - This vesicle - called also

\* Purkinje, art. Ei, in op. cit. x. 157.

c Haller. Elementa Physiol. viii. 208.

d Descript. Anat. Embryonis, Gotting. 1764. Cop. cit. p. 42. See, on this and other topics of Embryology, Von Siebold's Journal für Gebürtshülfe, xiv. Heft 3, Leipz. 1835; and Brit. and For. Med. Review, i. 241, Lond. 1836.

b Sec, also, R. Wagner, Elements of Physiology, translated by Robert Willis, M.D. p. 194, Lond. 1841.

membrana furciminalis and membrana intestinalis - has been alternately admitted and denied to be a part of the appendages of the human fœtus, from the earliest periods until the present day. It has been seen by Emmert, Meckel, Pockels, Velpean, Von Baer, Burdach and others a It is situate between the chorion and amnion, and communicates, in animals, with the urinary bladder by a duct called urachus. It has been observed in the dog, sheep, cow, in the saurian and ophidian reptiles, birds, &c. M. Velpeaub was never able to detect any communication with the bladder in the human subject, and he is compelled to have recourse to analogy to infer, that any such channel has in reality existed. From all his facts - which are not numerous or forcible - he "thinks himself authorized to say," that from the fifth week after conception till the end of pregnancy, the chorion and amnion are separated by a transparent colourless, or slightly greenish-yellow layer. This layer, instead of being a simple serosity, is lamellated, after the manner of the vitreous humour of the eye. It diminishes in thickness, in proportion to the development of the other membranes. The quantity of fluid, which its meshes inclose, is, on the contrary, in an inverse ratio with the progress of gestation. Becoming gradually thinner, it is ultimately formed into a homogeneous and pulpy layer, by being transformed into a simple gelatinous or mucous layer (enduit), which wholly disappears, in many cases, before the period of accouchement. Between the reticulated body, as Velpeau terms it, and the allantoid of oviparous animals, he thinks, there is the greatest analogy. Yet the fluid of the allantoid is very different from the urine, which is supposed, by some, to exist in the allantoid of animals. This fluid, we shall find, has been considered inservient to the nutrition of the new being, but, after all, it must be admitted, that our ideas regarding the vesicle, in man, are far from being determinate.

Pockels, of Brunswick, as existing in the human subject. It had been before observed in the mammalia. According to Pockels, it is pyriform; and much longer than, though of the same breadth as, the umbilical vessel. Within it, the intestines are formed. Velpeau, however, asserts, that he has never been able to meet with it; and he is disposed to think, that none of the embryos, depicted by Pockels, and by Seiler, were in the natural state. Such, too, is the opinion of Weber, and it is not noticed by the later embryologists.

According to most obstetrical physiologists, when pregnancy is

Purkinje, art. Ei, in Encyc. Wört. der Medicin Wissensch. xi. 151, Berlin, 1834.

b Embryologie ou Ovologie Humaine, Paris, 1833. See, also, Burdach, Die Physiologie als Erfahrungswissenschaft, ii. 530; Weber's Hildebrandt's Handbuch der Ana-

tomie, iv. 509, Braunschweig, 1832; and Meckel's Handbuch, u. s. w., Jourdan and Breschet's French translation, iii. 768, Paris, 1825.

c Isis, von Oken, p. 1342, 1825.

Das Ei und Die Gebärmutter des Menschen, u. s. w., p. 24, Dresd. 1832.
 Weber's Hildebrandt's Handbuch der Anatomie, iv. 518, Braunsch. 1832.

vol. II. - 39

multiple, the ova in the uterus are generally distinct, but contiguous to each other. By others, it has been affirmed, that two or more children may be contained in the same ovum, but this appears to require confirmation. The placenta of each child, in such multiple cases, may be distinct; or the different placentæ may be united into one, having intimate vascular communications with each other. At other times, in twin cases, but one placenta exists. This gives origin to two cords, and at others to one only, which afterwards bifurcates, and proceeds to both fœtuses. Maygrier, however, affirms unconditionally, that there is always a placenta for each fœtus; but that it is not uncommon, in double pregnancies, to find the two placentæ united at their margins; the circulation of each fœtus being distinct, although the vessels may anastomose. This was the fact in a case of quadruple pregnancy, communicated by M. Capuron to the Académie Royale de Méde-

cine, in Jan. 10th, 1837.

b. Development of the Ovum. — Prior to the consideration of the development of the human ovum, it may be well to refer to the changes that the egg undergoes during incubation; in which we have an opportunity of observing the transmutations at all periods of fætal formation, independently of any connexion with either parent. The subject has engaged the attention of physiologists of all ages; but it is chiefly to those of more modern times - as Hunter, Cuvier, Dutrochet, Pander, Rolando, Sir Everard Home, Prévost and Dumas, Von Baer, Kuhlemann,d Döllinger, D'Alton, Oken, Purkinje, Rathke, C. F. Wolff, Breschet, Burdach, Reichert, Carus, Krause, Seiler, Bojanus, Meckel, E. H. Weber, Bernhardt and Valentin, Coste, Owen, Sharpey, Velpeau, Flourens, Allen Thomson, T. W. Jones, Bischoff, Schwann and Schleiden, J. Müller, Rudolph Wagner and Martin Barry, the two last of whom received, about the same time, medals for their researches, the former from the Institute of France, and the latter from the Royal Society of London - that we are indebted for more precise information on the subject; although, unfortunately, they are by no means of accordance on many points. The investigations of Sir Everard Home, aided by those of the excellent microscopic observer, Mr. Bauer, are accompanied by engravings, some of which we shall borrow in elucidation of the following brief description.

Theegg of a bird, - of a hen for example, - consists of two descriptions of parts: - those which are but little concerned in the development of the new being, and which remain after the chick is hatched, - as the shell and the membrane lining it, - and such as

a Nouvelles Demonstrations d'Accouchemens, Paris, 1822-26.

<sup>b</sup> Journal de Physique, p. 88, for 1819.

e Isis, 1829, p. 407.

<sup>·</sup> Beiträge zur Entwickelungsgeschichte des Hühnchens im Ei, Wurzb. 1817. d Observ. quædam circa Negotium Generationis in ovibus fact. Gotting. 1753.

Symbolæ ad Ovi Avium Historiam ante Incubationem, p. 16, Wratislav. 1825; and art. Ei, in Encyclopad. Wörterb. der Medicin. Wissenschaft. Band. x. 107, Berl. 5 Sir E. Home's Lectures on Comp. Anat. iii. 4275

undergo changes along with those of the chick and co-operate in its formation, - as the white, the yolk, and the cicatricula or molecule. The shell is porous, to allow of the absorption of air through it; and of the evaporation of a part of the albumen or white. In the ovarium it is albuminous, but in the cloaca becomes calcareous. The membrane, membrana testæ, that lines the shell, is of a white colour, and consists of two layers, which separate from each other at the greater end of the egg, and leave a space filled with air, owing to the evaporation of the white and the absorption of air. This space is larger the older the egg, and is called the folliculus aëris or air-chamber. The albumen or white does not exist whilst the egg is attached to the ovary. It is deposited between the yolk and the shell, as the egg passes through the oviduct. Of the white there are two distinct kinds; - the outermost, thin and fluid, which evaporates in part, and is less abundant in the old than in the fresh laid egg, and another, situate within the last, which is much denser, and only touches the shell at the smaller extremity of the egg by a prolongation of its substance, which has

been called the ligament of the white. The yolk or yolk-ball, vitellus, seems to be at first sight a semifluid mass without organization; but on closer examination, it is found to consist of a volk-bag, two epidermic membranes, which envelope it as well as the cicatricula or mole cule. Two prolonga tions of these membranes, knotty, and terminating in a flocculent extremity in the albumen, called chalazæ or poles, are attached to the two ends of the egg and thus suspend it. It is also surrounded by a proper membrane; and lastly, under the epidermic coats of the volk, and upon its proper coat lies the



Ovarium of the laying Hen; natural size. The Ova at different stages of increment.—(Sir E. Home.)

cicatricula, macula, tread of the cock, or gelatinous molecule from which the future embryo is to be formed. It is found before the volk leaves the ovarium.

The external membrane of the yolk, when it quits the yolk-bag, is very thin and delicate; its surface is studded over with red dots, which disappear in its passage along the oviduct. When this membrane is removed, there is a natural aperture in the thick, spongy covering under it, through which is seen the cicatricula or

Fig. 256.



molecule, surrounded by an areola, halo, or circulus. On examination, this areola proves to be nothing more than that part of the surface of the yolk, which is circumscribed by the

margin of the aperture.

The molecule or cicatricula itself, (Fig. 256,) has a granulated appearance; and, according to Sir Everard Home, a is made up, in the centre, of globules an inch in part of an inch in diameter, surrounded by circles of a mixed substance; about two-thirds consisting of the same small globules, and one-third of larger oval globules, about To at h part of an inch in diameter; the last resembling in shape the oval red globules of the New laid Egg with its Molecule, &c .- (Sir blood in the bird. Besides the glo-

appears in drops, when the parts are immersed in water. Oval globules and oil are also met with in the yolk itself, but in small proportion and devoid of colour. If the ovum, according to

Fig. 257.

Section of a hen's egg within the Ovary.

a. The granular membrane form ing the periphery of the yolk. b. Vesicle of Purkinje imbedded in the cumulus. c. Vitellary membrane. d. Inner and outer layers of the capsule of the ovum. c. Indusium of the ovary. — (T. W. Jenes.)

Valentin, be lacerated and its contents minutely examined, the cicatricula is found like a grayish white disk, which in its whole periphery is dense, granular and opaque, but in the centre presents a clear non-granular, and perfectly diaphanous point. Purkinje found, that when he removed the dark granular mass, by suction with a small tube, there remained a perfectly transparent vesicle, filled with a pellucid lymph, which had a decidedly spherical form, but, being extremely delicate, was very easily lacerated, and its fluid escaped. As he found this, which later naturalists have named - after its discoverer - the ' Purkinjean vesicle,' in the ova of the ovary, but could not see it in ova, which had already entered the

\* Op. cit. iii. 426.

b Handbuch der Entwickelungsgeschichte des Menschen, u. s. w. Berlin, 1835 Edinburgh Med. and Surg. Journ. April, 1836, p. 393; and Bernhardt, Symbolse ad ovi Mammalium Historiam ante Prægnationem. Wratisl. 1834. See, also, Dr. A

oviduct, he gave it the name "germinal vesicle." The granular membrane, - its thickened portion, the so called cicatricula, - and the germinal vesicle, constitute those parts of the ovum, which pass immediately into the original foundation of the embryo, the

blastoderma or "germinal membrane."

When the egg leaves the ovarium, (Fig. 255,) the ovarial yolk-bag gives way at the median line, and the yolk drops into the commencement of the oviduct. The volk-bags are exceedingly vascular, the outer membrane of the yolk being connected to them by vessels and fasciculi of fibres, but being readily separable from them. During the first hours of incubation no change is perceptible in the egg, but, about the seventh, the molecule is evidently enlarged, and a membrane, containing a fluid substance, is observable. This membrane is the Amnion, Colliquamentum, Sacculus Colliquamenti, Nidus pulli, Areola pellucida Wolffii, transparent pellucid area. At this time, a white line is perceptible in the molecule, which is the rudimental fœtus; and, even at this early period, according to Sir Everard Home, the brain and spinal marrow can be detected. The areola has extended itself; and the surface, beyond the line which formed its boundary, has acquired the consistence of a membrane, and has also a distinct line by which it is circumscribed. This Sir Everard calls the outer areola. In the space between these two areolæ are distinct dots of an oily matter.

In twelve hours the rudiments of the brain are more distinct, as well as those of the spinal marrow. A dark line or primitive trace or streak may now be discovered in the cicatricula towards the centre of the transparent area, lying in the transverse axis of the egg, and swollen at the extremity, which lies to the left when the smaller end of the egg is turned from us. The larger extremity indicates the place where the head is afterwards formed, and occupies nearly the centre of the transparent area. As incubation proceeds, the whole cicatricula expands: towards the twelfth or fourteenth hour the germinal membrane divides into two layers Egg thirty-six hours after Incubation. — of granules, the uppermost being (Sir E. Home.)



entitled the serous or animal layer; the lower the mucous or vege-

Thomson, art. Generation, in Cyclopæd. of Anatomy and Physiology, Feb. 1838, part xiii. p. 452; Burdach, Physiologie als Erfahrungswissenschaft, 2te Auflage, i. 87, Leipz. 1835; Mr. Thomas Wharton Jones, on the First Changes in the Ova of the Mammifera, in Philos. Transact. part ii. for 1837, p. 339; and Dr. Martin Barry, Philosoph. Transact. 1838, pp. 301, 341; 1839, p. 307, and 1839-40.

\* Lectures on Comparative Anatomy, iii. 426.

tative layer, - the former being more extended than the latter. In a few hours later, the separation of the germinal membrane becomes more distinct, and between the serous and the mucous layers there appears a new layer, called the vascular. In one or other of the three primitive layers - serous, mucous, or vascular - is contained the germ of all the tissues and organs of the body, and it is conceived, that their histogeny admits of a distinct grouping. The serous layer, for example, is presumed to give origin to the organs of animal life - the brain, spinal marrow, the organs of the senses, the skin, muscles, tendons, ligaments, cartilages, and bones; from the mucous layer originate the organs of vegetative life - the intestinal canal, lungs, liver, spleen, pancreas, and other glands; and from the vascular layer the heart and vascular system are presumed to arise. It is not decided to which layer the reproductive system should be assigned. According to Dr. Barry,a however, there does not occur in the mammiferous ovum any such phenomenon as the splitting of a germinal membrane into the "so called serous, vascular, and mucous laminæ. Nor is there any structure entitled to be denominated a germinal membrane, for it is not a previously existing membrane, which originates the germ, but it is the previously existing germ which, by means of a hollow process, originates a structure having the appearance of a membrane."

In thirty-six hours, the head is turned to the left side. The cerebrum and cerebellum appear to be distinct bodies. The iris is perceptible through the pupil. The intervertebral nerves are nearly formed; those nearest the head being the most distinct. A portion of the heart is seen. At this period, under the inner areola, apparently at the termination of the spinal marrow, a vesicle begins to protrude, which is seen earlier in some eggs than in others. The white of egg is found to be successively absorbed by the yolk, so that the latter is rendered more fluid and its mass augmented. The first appearance of red blood is discerned on the surface of the volkbag towards the end of the second day. A series of points is observed, which form grooves; and these closing constitute vessels, the trunks of which become connected with the chick. The vascular surface itself is called figura venosa, area or area vasculosa; and the vessel, by which its margin is defined, vena terminalis, and circulus venosus. The trunk of all the veins joins the vena portæ, whilst the arteries, that ramify on the yolk-bag, arise from the mesenteric artery of the chick, and have hence been called omphalomesenteric.

In two days and a half, the spinal marrow has its posterior part inclosed; the auricles and ventricles of the heart are perceptible, and the auricles are filled with red blood. An arterial trunk from the left ventricle gives off two large vessels, — one to the right side of the embryo, the other to the left, — sending branches over the whole of the areolar membrane, which is bounded on each side by a large trunk carrying red blood; but the branches of the two trunks

<sup>\*</sup> Op. cit.; and Wagner's Elements of Physiology, p. 153, (note,) Lond. 1841.

do not unite, there being a small space on one side, which renders the circle incomplete. This Sir Everard Home calls the areolar circulation.

In three days, the outer areola has extended itself over one-third of the circumference of the yolk, carrying the marginal arteries along with it to the outer edge, but diminished in size. The brain is much enlarged; the cerebellum being yet the larger of the two. The spinal marrow and its nerves are most distinctly formed; and the eve appears to want only the pigmentum nigrum. The right

ventricle of the heart contains red blood; the arteries can be traced to the head; the rudiments of the wings and legs are formed, and the vesicle is farther enlarged, but its vessels do not carry red blood. It has forced its way through the external covering of the yolk, and opened a communication through this slit, by which a part of the albumen is admitted to mix itself with the yolk, and gives it a more oval form. At this period, the embryo is generally found to have changed its position and to be wholly turned on the left side.

In four days, the vesicle is more optic nerve and pigmentum nienlarged, and more vascular, its



grum of the eye are visible. The outer areola extends half over the yolk, with which a larger portion of the white is now mixed.

In five days, the vesicle has acquired a great size and become exceedingly vascular; the yolk, too, has become thinner, in con-

sequence of its admixture with more of the albumen.

In six days, the vascular membrane of the areola has extended farther over the yolk. The vesicle, at this time, has suddenly expanded itself in the form of a double night-cap over the yolk, and its coverings are beginning to inclose the embryo, the outermost layer being termed the chorion, the innermost the middle membrane. The amnion contains a fluid in which the embryo is suspended by the vessels of the vesicular membrane. The brain has become enlarged so as to equal in size the body of the embryo. Its vessels are distinctly seen. The two eyes equal the whole brain in size. The parietes of the thorax and abdomen have begun to form; and the wings and legs are nearly completed, as well as the bill. At this period muscular action has been noticed.

In seven days, the vesicle - having extended over the embryo, - has begun to inclose the areolar covering of the yolk, and a pulsation is distinctly seen in the trunk that supplies the vesicular

bag with blood. The pulsations were, in one case, seventy-nine in a minute, whilst the embryo was kept in a temperature of 105°; . but when the temperature was diminished, they ceased, and when again raised to the same point, they were reproduced. The muscles of the limbs now move with vigour.


In eight days, the anastomosing branches of the vesicular circu-

lation have a strong pulsation.

In nine days, the vesicle has nearly inclosed the yolk.



Egg five days after Incubation .- (Sir E. Home.)



Egg ten days after Incubation .- (Sir E. Home.)

In ten days, no portion of the volk is observable on the outside of the vesicle. The embryo being taken out of the amnion, - now become full of water, - the thorax is found to be completely formed, and the roots of the feathers very distinct.

The contents of the egg, during the formation of the embryo, become much diminished in quantity, and the void space is gradually occupied by a gas, which was examined by Mr. Hatchett, and found to be atmospheric air deposited at the great end of the egg between the layers of the membrane lining the shell. Even prior to incubation, there is always a small portion of air in this place which is supposed to be employed in aërating the blood, from the time of its first acquiring a red colour, till

superseded in that office by the external air acting through the egg-shell upon the blood in the vessels of the vesicular membrane,

with which it is lined.

Between the period of fourteen and eighteen days, the yolk becomes completely inclosed by the areolar membrane; and, at the expiration of the latter period, the greater part of the yolk is drawn into the body, as in Fig. 263. At twenty days, the chick is completely formed, the yolk is entirely drawn in, and only portions of the membrane belonging to the vesicle are seen externally. The yolkbag has a narrow tube, ductus vitellarius, ductus vitelli intestinalis seu apophysis, half an inch long, connecting it with the intestine, eight inches above the openings of the cæca into the gut.

The whole of these changes, which, in the viviparous animal, are effected within the womb of the mother, take place in the incubated chick by virtue of its own powers; and without any assistance except that of the atmospheric air and of a certain degree of warmth. In the course of incubation, the volk becomes constantly thinner and paler, by the admixture of the white; and, at the same time, innumerable fringe-like vessels, with flocculent extremities, of a singular structure, form on bag, and hang into the yolk. portions of the vesicular and areolar membranes and The office of these is presumed two diameters. — (Sir E. Home.)

to be, to absorb the yolk and to convey it into the veins of the yolkbag, where it is assimilated to the blood and applied to the nutrition of the new being. Blumenbach states, that in numerous and varied microscopical examinations of the yolk-bag, in the latter weeks of incubation, he thinks he has observed the actual passage of the volk from the vellow flocculent vessels of the inner surface of the bag into the bloodvessels which go to the chick. He has, at all events, seen manifest yellow streaks in the red blood contained in those veins. When the chick has escaped from the shell, the yolk, we have seen, is not exhausted, but is received into the abdomen, and as it communicates with the intestinal tube, it is, for some time, a source of supply to the young animal, until its strength is equal to the digestion of its appropriate food. The highly vascular chorion is manifestly an organ

of aëration, like the placenta of the mammalia. a



Embryo of the Egg. the inner surface of the yolk- showing the opening in the abdomen, from which




Embryo eighteen days old. Half the natural size. -- (Sir E. Home.)

For a minute and recent account of the development of the chick in ovo, illustrated by numerous engravings, see R. Wagner, Elements of Physiology, translated by R Willis, p. 80, Lond. 1841; also, Burdach, Die Physiologie als Erfahrungswissenschaft,. 2te Band, 2te Auflage, Leipz. 1837; Towne, Guy's Hospital Reports, Oct. 1839, p. 385; and T. W. Jones, Brit. and For. Med. Rev., Oct. 1843, p. 513.

The development of the mammalia greatly resembles that of the bird; and it is in the former, that the histogeny of the impregnated ovum, at the earliest periods of intra-uterine existence, has been studied. There can be no doubt, as Wagner has remarked,a that researches on the mammalia, carefully conducted and used with discretion, are much better calculated to throw light on the primary formation of the human embryo than any amount of necessarily unconnected observations on human ova cast off at an early period, and in the great majority of cases obviously diseased. In treating the whole of the subject, Wagner considers the development of the chick to be the safest guide, and he consequently refers to it continually in his interesting description of the development of the mammiferous ovum. In the higher oviparous animals, the germinal membrane absorbs the nutritious matter from the yolk, and prepares it for the use of the embryo itself, by converting it into blood; but after the yolk has been exhausted, the yolk-bag is taken into the body, and is gradually removed by absorption. In the mammalia, the quantity of yolk, laid up for the nutrition of the embryo, is comparatively inconsiderable, and it is only required until the ovum has contracted a union with the uterus; after which the yolk-bag is separated from the embryo, and thrown off as useless. Still, as Dr. Carpenter has remarked, "the early processes are the same in mammiferous as they are in oviparous animals; and the development of man, of a bird, of a reptile, or of a fish, takes place, up to a certain point, upon the same general plan."

The human ovum certainly does not reach the uterus until towards the termination of a week after conception. On the seventh or eighth day, it has the appearance referred to in the case so often cited from Sir Everard Home; the future situations of the brain and the spinal marrow being recognisable with the aid of a powerful microscope. On the thirteenth or fourteenth day, according to

Fig. 264.



Maygrier, the ovum is perceptible in the uterus, and of about the size of a pea, - according to Pockels, of a Spanish nut,e containing a turbid fluid, in the midst of which an opaque point is suspended, the punctum saliens. Fig. 264 represents an ovum, which is figured

by Velpeau, and could not have been more than fourteen days old, unless the midwife, who gave it to him, and who was herself the subject of the miscarriage, deceived him, and she appeared to have had no reason for so doing. Good descriptions and representations of the human ovum within the first month from conception are, however, as Wagnerd has remarked, very scarce, and many of the accounts appertain to diseased ova, or to monstrous or distorted embryos thrown off by abortion. Its weight has

d Elements of Physiology, translated by R. Willis, p. 161, Lond. 1841.

b Human Physiology, § 758, Lond. 1842. a Op. cit. c Isis, von Oken, December, 1825; and Granville's Graphic Illustrations of Abortion, part vii., Lond. 1834.


been valued at about a grain; - length one-twelfth of an inch.

On the twenty-first day, the embryo appears under the shape of a large ant, according to Aristotle; of a grain of lettuce; of a grain of barley, according to Burton; of the malleus of the ear, according to Baudelocque; - or one-tenth of an inch long, according to Pockels. At this period the different parts of the embryo have a little more consistence; and those that have afterwards to form bone assume the carti- ovum and Embryo, fifteen days laginous condition. On the thirtieth day, some feeble signs of the principal organs





and of the situation of the upper limbs are visible; - length four or five lines.



Ovum and Embryo, twenty-one days old .- (Maygrier.)

About the forty-fifth day the shape of the child is determinate; and it now, in the language of some anatomists, Fig. 267.

ceases to be the embryo, and becomes the fætus. According to others it does not become entitled to the latter name until after the beginning of the fourth month."

The limbs resemble tubercles, or the shoots of vegetables; the body lengthens, but preserves its oval shape, the head bearing a considerable proportion to the rest of the body. The base of the Fætus at forty-five trunk is pointed and elongated. Blackish points, or lines, indicate the presence of the eyes, mouth, and nose; and

<sup>\*</sup> Valentin, in art. Fœtus, Encyclopad. Wörterb. der Medicin. Wiesensch. xii. 355, Berlin, 1835.

Fig. 268.




Fœtus at two months.

similar, parallel points correspond to the situa-

tion of the vertebræ. Length ten lines.
In the second month, most of the parts of the fætus exist. The black points, which represent. ed the eyes, enlarge in every dimension; the eyelids are sketched, and are extremely transparent; the nose begins to stand out; the mouth increases, and becomes open; the brain is soft and pulpy; the heart is largely developed, and opaque lines set out from it; which are the first traces of large vessels. Prior to this period - very early in-

deed - substances or bodies are perceptible, which were first de-



Corpora Wolffiano, with Kidney and Testes, from Embryo of Birds.

scribed, as existing in the fowl, by Wolff,a and in the mammalia by Oken, b and which have been called by the Germans, after their discoverers, Wolffische oder Okensche Körper, ("bodies of Wolff or Oken.") According to J. Müller, they disappear in man very early, so that but slight remains of them are perceptible after the ninth or tenth week of pregnancy. They cover the region of the kidneys and renal capsules, which are formed afterwards, and they are presumed to be the organs of urinary secretion during the first periods of fætal existence. The fingers and toes are now distinct.

In the third month, the eyelids are more a. Kidney. b, b. Ureters. c. developed and firmly closed. A small hole Corpus Wolffianum. d. Its excretory duct. c, c. Testicles. At is perceptible in the pavilion of the ear. the summit are seen the supra. The alæ nasi are distinguishable. The lips are very distinct, and approximate, so that

the mouth is closed. The genital organs of both sexes undergo an extraordinary increase during this month. The penis is very long; the scrotum empty, frequently containing a little water. The vulva is very apparent, and the clitoris prominent. The brain, although still pulpy, is considerably developed, as well as the spinal marrow. The heart beats forcibly. The lungs are insignificant; the liver very large, but soft and pulpy, and appears to secrete scarcely any bile. The upper and lower limbs are developed. Weight two and a half ounces; length three and a half inches.

During the fourth month, all the parts acquire great advancement and character, except perhaps the head and the liver, which increase less in proportion than the other parts. The brain and spinal marrow acquire greater consistence: the muscular system,

<sup>\*</sup> Theoria Generationis, Hal. 1759.

b Oken und Kieser, Beiträge zur Vergleichend. Zoologie, Anatomie und Physiologie, H. i. p. 74, Bamberg und Wurzburg, 1806; and Rathke, in Weber's Hildebrandt's Handbuch der Anatom. iv. 440. Valentin, op. citat. p. 380.

which began to be observable in the preceding month, is now distinct; and slight, almost imperceptible, movements begin to manifest themselves. The *length* of the fætus is, at the end of one hundred and twenty days, five or six inches; the *weight* four or five ounces.

During the fifth month, the development of every part goes on; but a distinction is manifest amongst them. The muscular system is well-marked, and the movements of the fœtus unequivocal. The head is still very large, compared with the rest of the body, and is covered with small, silvery hairs. The eyelids are glued together. Length, seven to nine inches; weight six or eight ounces. If the fœtus be born at the end of five



Fætus at three months, in its membranes.

months, it may live for a few minutes.

In the sixth month, the derma begins to be distinguished from the epidermis. The skin is delicate, smooth, and of a purple colour; especially on the face, lips, ears, palms of the hands and soles of the feet. It seems plaited, owing to the absence of fat in the subcutaneous cellular tissue. The scrotum is small, and of a vivid red hue. The vulva is prominent, and its lips are separated by the projection of the clitoris. The nails appear, and towards the termination of the month, are somewhat solid. Should the feetus be born now, it is sufficiently developed to breathe and cry, but it dies in a few hours. Length, at six months, ten or twelve inches. Weight under two pounds.

During the seventh month all the parts are better proportioned. The head is directed towards the orifice of the uterus, and can be felt by the finger introduced into the vagina, but it is still very movable. The eyelids begin to separate, and the membrane, which previously closed the pupil — the membrana pupillaris — to disappear. The fat is more abundant, so that the form is more rotund. The skin is redder, and its sebaceous follicles are formed, which secrete a white, cheesy substance — the vernix caseosa —

which covers it; and the testicles are in progress to the scrotum. The length at seven months is fourteen inches; the weight under

three pounds.

In the eighth month, the fætus increases proportionably more in breadth than in length. All its parts are firmer and more formed. The nails exist; and the child is now certainly viable or capable of supporting an independent existence. The testicles descend into the scrotum; the bones of the skull, ribs, and limbs are more or less completely ossified. The length is sixteen inches; the

weight four pounds and upwards.

At the full period of nine months, the organs have acquired the development necessary for the continued existence of the infant. Length eighteen or twenty inches; weight six or seven pounds. According to Dr. Granville, length 22 inches; weight from five to eight pounds. Dr. Deweesa says the result of his experience, in this country, makes the average weight above seven pounds. He has met with two ascertained cases of fifteen pounds, and several, which he believed to be of equal weight. Dr. Moore, of New York, had several cases, where the weight was twelve pounds; and a case occurred in that city in 1821, of a fætus, born dead, which weighed sixteen and a half pounds. Dr. Traillb once weighed a child at the moment of birth, which weighed 14 pounds; Mr. Park, of Liverpool, found another to weigh 15 pounds; and a case has been recorded by Mr. J. D. Owens, in which a still-born child measured 24 inches in length, and weighed seventeen pounds twelve ounces.c Recently, Dr. Storer,d of Boston, has published the following results of observations. Of 30 children, 14 females weighed 112 pounds, or averaged 8 pounds each; and 16 males weighed 1454 pounds, or averaged 81 pounds each. The largest child seen by Dr. Storer was a male, and weighed 13 pounds: the next in weight was 121 pounds. One weighed 11, one 101, and two 10 pounds each.

Where there are twins in utero, the weight of each is usually less than in a uniparous case, but their united weight is greater. M. Dugès, of Paris, found that in 444 twins the average weight was four pounds, and the extreme weights three and eight pounds.

The whole of this description amounts to no more than an approximation to the truth. The facts will be found to vary greatly in individual cases, and, therefore, according to individual experience; and this accounts for the great discordance in the statements of different observers. This discordance is strongly exemplified in the following table containing the estimates of the length

2 Compendious System of Midwifery, 8th edit. Philad. 1836.

New England Quarterly Journal of Medicine and Surgery, July, 1842.

<sup>5</sup> Outlines of a Course of Medical Jurisprudence, 2d édit. p. 16, Edinb. 1840, or Amer. Edit. with notes by the author of this work, p. 27, Philad. 1841.

• Lond. Lancet, Dec. 22, 1838.

Burdach's Physiologie als Erfahrungswissenschaft, Band ii.; Valentin, art. Fætus, in Encycl. Wörterb. der Medicin. Wissensch. xii. 372, Berlin, 1835; and Trattato Générale di Ostetricia, di Dr. Asdrubali, 2de edizione, i. 152, Roma, 1812.

and weight of the fœtus at different periods of intra-uterine existence, as deduced by Dr. Becka from various observers, and as given by Maygrierb on his own authority, and by Dr. Granvillec as the averages of minute and accurate observations made by Autenrieth, Sömmering, Bichat, Pockels, Carus, &c., confirmed by his own observations made on several early ova, and many fœtuses examined in the course of seventeen years' obstetrical practice. It is proper to remark, that the Paris pound, poids de marc, of sixteen ounces, contains 9216 Paris grains; whilst the avoirdupois contains only 8532.5 Paris grains; and that the Paris inch is 1.065977 English inch.

1	Beck.	Maygrier.	Granville.	Beck.	Maygrier.	Granville.
SI 532119	noth sim wo	Length.	College (	Weight.		
At 30 days, 2months, 3 do. 4 do. 5 do. 6 do. 7 do. 8 do.	3 to 5 lines 2 inches 3½ do. 5 to 6 do. 7 to 9 9 to 12 12 to 14 16	10 to 12 lines 4 inches 6 do. 8 do. 10 do. 12 do. 14 do. 16 do.	1 inch 3 inches 9 inches 12 do. 17 do.	2 ounces 5 drachms 20 grains 11 ounce 4 or 5 do. 7 or 8 do 16 ounces 1 to 2 pounds 2 to 3 do. 3 do. 2 to 4 pound 3 to 4 do. 4 to 5 do.		

The difficulty must necessarily be great in making any accurate estimate during the early periods of fætal existence; and the changes in the after months are liable to considerable fluctuation. Chaussier affirms, that after the fifth month, the fœtus increases an inch every fifteen days, and Maygrier adopts his estimate. The former gentleman has published a table of the dimensions of the fœtus at the full period, deduced from an examination of more than fifteen thousand cases. From these we are aided in forming a judgment of the probable age of the fœtus in the latter months of utero-gestation; - a point of interest with the medico-legal inquirer. At the full period, the middle of the body corresponds exactly with the umbilicus, or a very little below it; at eight months, it is three-quarters of an inch, or an inch higher. At seven months it approaches still nearer the sternum; and at six months it falls exactly at the lower extremity of that bone; hence, if we depend upon these admeasurements, should the middle of the body of the fætus be found to fall at the lower extremity of the sternum, we may be justified in concluding that the fœtus is under the seventh month, and consequently not viable or rearable.

The following are the results of some observations made by Mr. Taylor, Lecturer on Medical Jurisprudence at Guy's Hospital, and Dr. Geoghegan, Professor of Medical Jurisprudence in the Royal College of Surgeons in Ireland.

Medical Jurisprudence, 6th edit. i. 276, Philad. 1838. See, also, Dr. W. A. Guy, Principles of Forensic Medicine, Part i. p. 100, Philad. 1843.

Principles of Forensic Medicine, Part i. p. 100, Philad. 1843.

b Nouvelles Demonstrations d'Accouchemens, Paris, 1822-26.

e Graphic Illustrations of Abortion, &c., p. xi., Lond. 1834. See Seiler, art. Embryo, in Anat. Phys. Real Wörterb. ii. s. 530, Leipz. und Altenb. 1818.

Case. Whole length.			1.	Attachment of the Umbilical Cord.		
1		184	-	A quarter of an inch below	w the centre.	
2	-	20		Half an inch	Do.	
3	-	175	-	Half an inch nearly	Do.	
4	-	165	III DI	Half an inch	Do.	
5	1	19	11.	Do.	Do.	
6	-	17	-	A little below the centre.		
7	-	18	-	Exactly at the centre.		
8	-	17	-	Do. Do.		
9	114	203	-	A little below the centre.		
10	-	191		Do. Do.		
11	27	183		Exactly at the centre.a		

A striking circumstance connected with the development of the fœtus, is the progressive diminution in proportion between the part of the body above the umbilicus and that below it. At a very early period of fœtal life, (see Figs. 267, and 268,) the cord is attached near the base of the trunk; but the parts beneath become gradually developed, until its insertion ultimately falls about the middle of the body.<sup>b</sup>

The following table of the length and weight (French), and central point of the fætus at different ages is given by M. Lepelletier. It still farther exhibits the discordance to which we have alluded

above.

Month.	Length.	Weight.	at the junction of the head and trunk.		
1	5 or 6 lines	9 to 15 grains			
2	18 to 20 lines	6 to 8 drachms	at the upper part of the sternum.		
3	2 to 3 inches	2 or 3 ounces	at the upper extremity of the xiphoid cartilage.		
4	5 to 6 inches	10 to 16 ounces	at the middle of the xiphoid cartilage.		
5	7 to 9 inches	1 to 2 pounds	at the lower extremity of the xiphoid cartilage.		
6	9 to 12 inches	2 to 3 pounds	several lines below the xiphoid car- tilage.		
7	12 to 15 inches	3 to 4 pounds	equidistant between the cartilage		
8	15 to 18 inches	4 to 6 pounds	an inch above the umbilicus.		
9	16 to 20 inches	6 to 8 pounds	at the umbilicus.		
Photos	Extremes,	Extremes,	Cheron His court of marriagality		
CHILDREN .	12 to 15 inches (Millot.)	2 to 16 pounds (Voistel.)	No. of Lands of the State of th		

The position of the fœtus in utero, and the cause of such position at various periods of utero-gestation, have been topics of some interest. In the early weeks, it seems to be merely suspended by the cord; and it has been conceived, that because the head is heavier, it is the lowest part. It is difficult, however, to admit this as the cause of the position assumed in such an immense majority of cases, or to fancy, that the nice adaptation of the fœtal

<sup>T. R. Beck, Amer. Journ. of the Med. Sciences, Oct. 1842, p. 489.
See Carus, Lehrbuch der Gynäkologie, Th. ii. s. 29, Leipz. 1828.
Physiologie Médicale et Philosophique, iv. 451, Paris, 1833.</sup> 

position to the parts through which the child has to pass is simply dependent upon such a mechanical cause. Gravity can afford us no explanation why the face in 12,120 cases out of 12,533, has been found turned to the right sacro-iliac synchondrosis, and the occiput to the left acetabulum; and in the 63 of these cases in which the face was turned forwards, and in the 198 breech presentations, are we to presume, that the whole effect was owing to mere difference of weight in those parts that were lowest?

The common position of the fætus, at the full period, is exhibited in the next illustration. The body is bent forward, the chin resting on the chest, the occiput towards the brim of the pelvis, the arms approximated in front, and one or both lying upon the face; the thighs bent upon the abdomen, the knees separated, the legs crossed, and drawn up, and the feet bent upon the anterior surface of the legs; so that the oval, which it thus forms, has been estimated at about ten inches in the long diameter; the head, at the full period, resting on the neck, and even on the mouth, of the womb, and the breech corresponding to the fundus of the organ.

From the first moment of a fecundating copulation, the minute matters, furnished by both sexes, when commingled, commence the work of forming the embryo. For a short time, they find in

the ovum the necessary nutriment, and subsequently obtain it from the uterus. The mode in which this action of formation is accomplished is as mysterious as the essence of generation itself. When the impregnated ovum is first seen, it seems to be an amphorous, gelatinous mass, in which no distinct organs are perceptible. In a short time, however, the brain and spinal marrow, and bloodvessels, make their appearance; but which of these is first developed is undecided.

Sir Everard Home,a

Fig. 271.

Full period of Utero-Gestation.

\* Lect. on Comp. Anat. iii. 292, and 429.

- from his observations of the chick in ovo, as well as from the microscopic appearances, presented by the ovum in the case of the female who died on the seventh or eighth day after impregnation, in which a rudimental brain and spinal marrow were perceptible, - decides, that the parts, first formed, bear a resemblance to brain, and that the heart and arteries are produced in consequence of the brain having been established. Malpighi, Brera, Pander, Tiedemann, Prévost and Dumas, Velpeau, Rolando, and Schröder van der Kolk,a also assign the priority to the nervous system. Meckel, however, admits no primitive organizing element, but believes, that the first rudiments of the fætus contain the basis of every part. On the other hand, the researches of Serres, on the mode of development of the nervous system, have induced him to be in favour of the earlier appearance of the bloodvessels,b and this view appears to be supported by the fact, that if an artery of the brain be wanting, or double, the nervous part to which it is usually distributed is also wanting, or double. The acephalous fœtus has no carotid or vertebral arteries; whilst the bicephalous or tricephalous have those vessels double or treble: with these views Dr. Granvillec accords, and he lays it down as a law, that the nerves invariably appear after the arteries which they are intended to accompany. A like discordance exists in the views regarding the precedence in formation of the bloodvessels. blood is clearly formed before the heart. It appears at distinct points from it, and acquires a motion independently of it. The veins are formed first; the heart next, and lastly the arteries. This is the view of the generality of physiologists, but a distinguished Italian physiologist — Rolando — assigns the precedency to the arteries. Farther experiments are demanded on these interesting, but intricate points of organogenia.e Messrs. Geoffroy Saint-Hilaire, Meckel, Serres, Tiedemann, and others are of opinion, that the development of the embryo takes place from the sides towards the median line - from the circumference towards the centre; but Velpeau considers, that the median line is first formed.k The spinal marrow is the first portion of the nervous system which appears; and this system, he thinks, precedes every other.

Histogeny, or the development of the different tissues, is a topic of deep interest to the student of general anatomy, and has en-

a Observationes Anatom. Pathol. et Pract. Argum. 8vo. Amstel. 1826; cited in Edinb. Med. and Surg. Journal, for April 1, 1836.

b See, also, Medici, in Bulletino delle Scienze Mediche, Dicembre, 1836, p. 389; and Richerand, Nouveaux Eléments de Physiologie, 13ème édit. par M. Bérard ainé, § 206, Bruxelles, 1837.

<sup>4</sup> See Von Baer, Ueber Entwickelungsgeschichte der Thiere, u. s. w., Königsberg, 1837; also, Prof. R. J. Graves, in Lond. Med. Gaz. for June 30th, 1838, p. 591.

\* Adelon, Physiologie de l'Homme, 2de édit. iv. 353, Paris, 1829.

f Philosophie Anatomique, Paris, 1818-22.

# Handbuch der Anatomie, Band, i.

b Recherches d'Anatomie Transcendante, &c. Paris, 1832.

† Anatomie du Cerveau, traduit par Jourdan, Paris, 1823.


1 Embryologie ou Ovologie Humaine, Paris, 1833.

k See, also, H. Mayo's Outlines of Physiology, p. 385, Lond. 1833; and Granville, Graphic Illustrations of Abortion, p. xii. Lond. 1834.

gaged the attention of some of the most excellent anatomists of modern times. Amongst these, Rolando, Tiedemann, Ackermann, Serres, Velpeau, Walther, Béclard, Rosenmüller, Geoffroy Saint-Hilaire, Meckel, Burdach, Rudolph Wagner, Valentin, Bischoff, Schwann and Schleiden, are especially conspicuous. The nature and limits of this work will preclude us, however, from entering into this investigation any farther than to point out some of the most striking peculiarities of fætal life. The fullest treatise on the subject is that of Burdach, and one of the most recent that of Rudolph Wagner.

c. Peculiarities of the Fætus. - The head of the fætus is large

in proportion to the rest of the body, and the bones of the skull are united by membrane; the sagittal suture extends down to the nose, so as to divide the frontal bone into two portions; and where this suture unites with the coronal, a quadrangular space is left, filled up by membrane, which is called the anterior fontanelle or bregma. Where the posterior extremity of the sagittal suture joins the lambdoidal, a triangular space of a similar kind is left, called the posterior fontanelle or posterior bregma. It is important for the obstetrical practitioner to bear in mind the shape of these spaces, as they indicate to him whether the anterior or posterior part of the head is the presenting part. The pupil of the eye, in a very young fœtus, is entirely closed by a membrane, called membrana pupillaris, which arises from the inner margin of the iris, and continues there till the seventh month, when it gradually vanishes by absorption. It is a vascular substance, and, like the iris, to which it is attached, separates the two chambers of the eye from each other. Wachendorffd first described it in



Fatus at full term.

a, a. Divided integuments. c, c. Divided ribs and intercostal muscles. c, c. Lobes of thymus gland g, g, h, h. Lungs. i. Right auricle of the heart. k. Right ventricle. n, o. Right and left lobes of the liver. p. Stomach. q, q. Small intestines. r. The colon. s. Bladder of urine inflated. t. The urachus. u, u. The umbilical arteries. v. The umbilical vein. w. The umbilicus. z. The collapsed umbilical cord.

Die Physiologie als Erfahrungswissenschaft, 2ter Band. 2te Auflage, Leipz. 1837.

· Elements of Physiology, translated by Willis, Lond. 1841.


Weber's Hildebrandt's Handbuch der Anatomie, iv. 467, Braunschweig, 1832; Purkinje, art. Ei, Encyclop. Wörterb. der Medicinisch. Wissensch. xi. 107, Berlin, 1834-5; Valentin, art. Fætus, ibid. xii. 255, Berl. 1835. An abstract of the views of Serres, is given in Amer. Journal of the Med. Sciences, Feb. 1837, p. 473. See, also, Fletcher's Rudiments of Physiology, part i. Edinb. 1836.

<sup>4</sup> Commerc. Litterar. Noric. 1740; and Valentin, art. Fætus, Encycl. Wörterb. n. s. w., xii. 376.

1738; and both he and Wrisberg detected vessels in it. Its vascularity was denied by Bichat, but it has been satisfactorily demonstrated by J. Cloquet.<sup>a</sup> The membrane is manifestly connected with the process of formation of the delicate organ to which it is attached; and according to Blumenbach,<sup>b</sup> it keeps the iris

expanded, during the rapid increase of the eyeball.

In the upper part of the thorax of the fætus, a large gland, or rather glandiform ganglion exists, called the thymus, and by Joseph Frank, corpus incomprehensibile. It is situate in the superior mediastinum, and lies over the top of the pericardium and the arch of the aorta. It has two long cornua above, and two broad lobes below. Its appearance is glandular, and colour very variable. In the progress of age it diminishes, so that in the adult it is wasted, and in old age can scarcely be discovered amongst the cellular tissue. Krause<sup>c</sup> states, that he has found it in almost all individuals between twenty and thirty years of age, and very often larger than in young children. It is surrounded by a thin, cellular capsule, which sends prolongations into its interior, and divides it into lobules of unequal size, in which several vesicles are distin-



Section of the Thymus Gland at the Eighth Month.

1. Cervical portions of the gland; independence of two lateral glands is well marked. 2. Secretory cells seen upon cut surface of section; these are observed in all parts of section. 3, 3. Pores or openings of secretory cells and pouches; they are seen covering whole internal surface of great central cavity or reservoir. Continuity of reservoir in lower or thoracic portion of gland with cervical portion, is seen in the figure.— (Sir Astley Cooper.)

guishable, filled with a milky Sir Astley Cooper fluid. states, that the lobules may be drawn out and separated from each other in the manner of a string of beads, when their enveloping capsule has been removed. The ordinary weight of the thymus has been estimated at half an ounce, but this is probably above the average. Dr. Roberts, d of New York, found the average weight in the full grown fœtus to be 229 grains. The thymic arteries proceed from the inferior thyroid, internal mammary, bronchial, mediastinal, &c. The nerves proceed from the pneumogastric, diaphragmatic, and inferior cervical ganglia. It has no excretory duct; and is one of the

Mémoir sur la Membrane Pupillaire, Paris, 1817.

b Institut. Physiolog. § 262. See, also, Elliotson's Translation, and Weber's Hildebrandt's Handbuch der Anatomie, iv. 84.

c Müller's Archiv. Heft 1, 1837.

d Amer. Journ. of the Med. Sciences, Aug. 1837, Nov. 1838, and Oct. 1841; New York Journ. of Med. and Surg., Jan. 1840; and New York Med. Gaz., July 21, 1841. See, also, Dr. C. Lee, in Amer. Journ. of the Med. Sciences, Jan. 1842, p. 138.

most obscure, in its physiology, of any organ of the body, although, like the lymphatic glands, probably connected with the function of nutrition.

The thyroid gland, which has been described in another place, and whose functions are equally obscure, is also largely developed


in the fœtus; as well as the supra-renal capsules.

The lungs, not having received air in respiration, are collapsed and dense, containing no more blood than is necessary for their nutrition. They are of a dark colour, like liver, and do not fill the cavity of the chest. Their specific gravity is greater than that of water, and consequently they sink in that fluid. On cutting into them, no air is emitted, and no hemorrhage follows. The absolute weight, however, of the lungs is less; no more blood, as we have seen, being sent to them than what is necessary for their nutriment; whilst, after respiration is established, the whole of the blood passes through them: the vessels are consequently filled with blood, enlarged, and the organs themselves increased in absolute weight. Ploucquet asserts, from experiments, that the weight of the lungs of a full-grown fœtus, which never respired, is to that of the whole body, as 1 to 70; whilst, in those, in which respiration has been established, it is as 1 to 35; the absolute weight being thus doubled. These numbers cannot, however, be considered to afford a satisfactory average, - the exceptions being numerous; but they show that, as might be expected, the absolute weight is less, prior to the establishment of respiration. Careful and extended observations have satisfactorily shown, that although an increase in the weight of the lungs is generally found after respiration has been completely established, it is by no means the case when the inspirations have been feeble, as they often are for some hours and days after birth; and, on the other hand, it is not unusual to meet, in infants that have not breathed, with lungs as heavy as in the average of those that have respired. b The subject is one of great interest, connected with infanticide, and has been treated in a competent manner, by Dr. J. B. Beck, in Beck's Elements of Medical Jurisprudence, - decidedly, in our opinion, one of the best medico-legal works in existence.c

It is, however, in the circulatory system of the fætus, that we meet with the most striking peculiarities. The heart is proportionably larger and more conical than in the adult. The valve of Eustachius—at the left side of the mouth of the inferior vena cava, where this vessel joins the sinus venosus,—is larger than at an after period, and is supposed to direct the principal part of the blood of that cava directly through the opening which exists between the right and left auricle. This opening, which is called the foramen ovale or foramen of Botal, is in the septum between the

Gulliver, in Gerber's General Anatomy, p. 100, Lond. 1842.
 Dr. Guy, Edinb. Med. and Surg. Journ., vol. lvi. and lvii.

See, also, J. B. Beck, Researches in Medicine and Medical Jurisprudence, New York, 1835.



Circulatory Organs of the Fatus.

1. Umbilical cord, consisting of umbilical vein and two umbilical arteries; proceeding from placenta (2). 3. Umbilical vein dividing into three branches; two (4, 4) to be distributed to liver; branches; two (4, 4) to be distributed to liver; and one (5), ductus venosus, which enters inferior vena cava (6). 7. Portal vein returning blood from intestines, and uniting with right hepatic branch. 8, Right auricle; course of blood is denoted by arrow, proceeding from 8 to 9, left auricle. 10. Left ventricle; blood following arrow to arch of aorta(11), to be distributed through branches given off by arch to head and upper extremities. Arrows 12 and 13, represent return of blood from head and upper extremities through jugular and subclavian veins, to superior vena cava (14), to right auricle (8), and in course of arrow through right ventricle (15), to pulmonary artery (16). 17. Ductus arteriosus, which appears to be a proper continuation of pulmonary artery, to be a proper continuation of pulmonary artery, the offsets at each side are right and left pulmo-nary artery cut off; these are of extremely small size as compared with ductus arteriosus. Ductus arteriosus joins descending aorta (18, 18), which divides into common iliacs, and these into internal iliacs, which become umbilical arteries (19), and return blood along umbilical cord to placenta; while other divisions, external iliacs (20), are continued into lower extremities. Arrows at termination of these vessels mark return of venous blood by veins to inferior cava .-(Wilson.)

auricles, and is nearly equal in size to the mouth of the inferior cava. It is situate obliquely, and has a membrane, forming a distinct valve, and somewhat of a crescentic shape, which allows part of the blood of the right auricle to pass through the opening into the left auricle, but

prevents its return.

The pulmonary artery, instead of bifurcating as in the adult, divides into three branches; - the right and left going to the lungs of the corresponding side, whilst the middle branch, - to which the name ductus arteriosus has been given, opens directly into the aorta; so that a great part of the blood of the pulmonary artery passes directly into that vessel. From the internal iliac arteries, two considerable vessels arise, called the umbilical arteries. These mount by the sides of the bladder, as in Fig. 272, on the outside of the peritoneum and perforate the umbilious in their progress to the umbilical cord and placenta. The umbilical vein, which is also a constituent of the cord, and conveys the blood from the placenta to the fætus, arises from the substance of the placenta by a multitude of radicles, which unite together to form it. Its size is considerable. It enters the umbilicus, (Fig. 272,) passes towards the inferior surface of the liver, and joins the left branch of the vena porta hepatica. Here a vessel arises called the ductus venosus, which opens into the vena cava inferior, or joins the left vena hepatica where that vein enters the cava. A part only of the blood of the umbilical vein goes directly to the vena cava; the remainder is distributed to the right

and left lobes of the liver, especially to the latter.

The digestive apparatus exhibits few peculiarities. The bowels, at the full period, always contain a quantity of greenish, or deep black, viscid fæces, to which the term meconium has been applied, owing to their resemblance to the inspissated juice of the poppy (unxay, "a poppy"). It appears to be a compound of the secretions from the intestinal canal and bile, and frequently contains down or fine hairs mixed with it. The liver is very large; so much so as to occupy both hypochondriac regions; and the right and left lobes are more nearly of a size than in the adult. Prior to birth it would seem to be the only decarbonizing organ; the lungs being inactive, but as soon as respiration is established, less blood is sent to the liver; and the diminution takes place with great rapidity, and is usually evidenced a short time after birth by the comparative paleness of its substance. Hence it has been supposed that the weight of the liver might aid in the determination of the question, whether a child had breathed or not. It has been shown, however, that although the liver, as a general rule, weighs less after respiration has been established, it by no means exists when the inspiration has been feeble for hours or days after birth; and, on the other hand, it is not uncommon to meet, in infants that have not breathed, with livers as light as in the average of those that have.a

The urinary bladder is of an elongated shape, and extends almost to the umbilicus. The muscular coat is somewhat thicker and more irritable than in the adult, and it continues to possess more power during youth. The common trick of the schoolboy — of sending the jet over his head — is generally impracticable in more advanced life. From the fundus of the bladder, a ligament of a conical shape, called the urachus, (Fig. 272,) ascends between the umbilical arteries to the umbilicus; becoming confounded in this place with the abdominal aponeuroses, according to Bichat, and forming a kind of suspensory ligament to the bladder. It is sometimes found hollow in the human fætus, but such a formation Bichat considers to be preternatural. In the fætal quadruped, it is a large canal, which transmits urine to the allantois, of which, as well as of other fætal peculiarities, we have previously treated.

Lastly, the genital organs require some notice. The successive development of this part of the system has given rise to some singular views regarding the cause of the sex of the fœtus. During the first few weeks, the organs are not perceptible; but, about the termination of the fifth week, a small, cleft eminence appears, which is the rudiment of the scrotum or the vulva, according to the sex. In the sixth week, an aperture is perceptible, common to the anus and genital organs, in front of which is a projecting tubercle. In the seventh and eighth weeks, this tubercle seems to

<sup>\*</sup> Guy, op. cit.; and Carpenter, Human Physiology, § 661, Lond. 1842.

be tipped by a glans, and grooved beneath by a channel which extends to the anus. In the eleventh and twelfth weeks, the perineum is formed and separates the anus from the genital organs. In the fourteenth week, the sex is distinct; but there still remains, for some time, a groove beneath the clitoris or penis, which becomes closed in the former, and is made into a canal in the latter. The striking similarity between the male and female organs has led Tiedemanna to conclude, that the female sex is the male, arrested at an inferior point of organization. In his view, every embryo is originally female; the cleft, described above, being the vulva, - the tubercle, the clitoris; to constitute the male sex, the cleft is united so as to form a raphe, the labia majora are joined to form the scrotum, the nymphæ to form the urethra, and the clitoris is transformed into a penis. In support of this opinion, Tiedemann asserts, that the lowest species of animals are almost all females; and that all the young acephali and aborted fœtuses, which have been examined, are of that sex. Others, again, assert, that the sexes are originally neuter, and that the future sex is determined by accidental circumstances, during the first week of fætal life; whilst Velpeau is disposed to believe, that they are all male; the infrapubic prolongation existing in every embryo, although there may be neither labia majora nor scrotum. But admitting that the embryo belongs to either the one or the other sex, or is neutral, we must still remain at a loss regarding the influences that occasion the subsequent mutations; and it seems impossible not to admit, that although an apparent sexual identity may exist amongst different embryo's, there must be an impulse seated somewhere, which gives occasion to the sex being ultimately male or female, as it causes the young being to resemble one or other parent in its outward form, or internal configuration; and if our means of observation were adequate to the purpose, a distribution of arteries or nerves might probably be detected, which could satisfactorily account, ab initio, for the resulting sex. In the absence of such positive data, Geoffroy St. Hilaire has suggested, that the difference of sex may be owing to the distribution of the two branches of the spermatic artery. If they continue in approximation, proceeding together, - the one to the testicle, the other to the epididymis, - the individual is male; if they separate, — the one going to the ovary, the other to the cornu of the uterus, - the individual is female. degree of predominance of the cerebro-spinal system, he thinks, determines the approximation or separation of the two arterial branches. This predominance being greater in the male, the spermatic arteries are more feeble and consequently in greater proximity; and conversely in the female.b


a Anatomie der Kopflosen Missgeburten, p. 54, Landshut, 1813.

b Adelon, Physiologie de l'Homme, 2de, édit. iv. 375; J. Müller, Bildungsgeschichte der Genitalien, Berlin, 1830; E. H. Weber's Hildebrandt's Anatomie, iv. 446; Valentin, art. Fætus, Encycl. Wörterb. der Medicin. Wissensch. xii. 381, Berlin, 1835; Coste, Annales d'Anatomie et de Physiologie, tom. ii. No. ii., Mars, 1838; and London Medical Gazette, May 12, 1838, p. 304.

Leaving these phantasies of the generalizing anatomists on a subject regarding which we must, probably, ever remain in the dark, let us inquire into the phenomena of the descent of the testes in the fætus. In the early months of fætal life, the testicle is an abdominal viscus, being seated below the kidney. About the middle of the third month of utero-gestation, it is about two lines long, and is situate behind the peritoneum, which is reflected over its ventral surface. At this time, a sheath of peritoneum may be observed, passing from the abdominal ring to the lower part of the testicle, and containing a ligament, called gubernaculum testis, which is considered to be formed of elastic cellular tissue, proceeding from the upper part of the scrotum, and from the part of the general aponeurosis of the thigh near the ring, and of some muscular fibres coming from the internal oblique and transversalis muscles.

Fig. 275.

Fig. 276.



Descent of the Testicle.

A. Testicle. B. Peritoneal covering or tunica vaginalis testis. C. Peritoneum of the loins.

D. Peritoneum descending before the testicle.

F. Peritoneum lining the abdomen.

Descent of the Testicle.

A. Testicle in the scrotum. B. Prolongation of the peritoneum. C. Peritoneum lining the abdomen. D. Peritoneum forming the tunica vaginalis. E. Cavity of the peritoneum. F. Kidney.

The head of the fœtus in utero being the lowest part, the testis has necessarily to ascend into the scrotum, and consequently some force must be exerted upon it. This is supposed to be effected by the contraction of the gubernaculum testis, yet it does not contain any fibrous structure, until after the descent of the testis has commenced. About the seventh month, the testes are in progress towards the scrotum, and have attained the inner ring. Fig. 275 exhibits one leaving the abdomen and entering the scrotum, into which it generally passes in the ninth month. In this descent, the organ abandons successively one portion of the peritoneum to pass behind another immediately below, until the lowest part of the pouch, formed by the peritoneum, around the testicle, as in Fig. 276,

becomes the tunica vaginalis testis; whilst the portion of peritoneum, that descended before the testicle, becomes, when the testicle

has fully descended, the second coat or tunica vaginalis.

As soon as the testicle has reached the lower part of the scrotum, the neck of the pouch approaches a closure, and this is commonly effected at birth. Sometimes, however, it remains open for a time, the intestines pass down, and congenital hernia is thus induced.<sup>a</sup>

The testes have not always descended into the scrotum at birth, even at the full period; of 97 new-born infants, Wrisberg found both testes in the scrotum in 67: one or both in the canal in 17; one testis in the abdomen in 8; and both testes were in the cavity of the abdomen in 3. Sometimes, as has been remarked elsewhere, one or both testes remain through life in the abdomen, a condition which is natural in the ram.

## 2. PHYSIOLOGY OF THE FŒTUS.

In investigating this interesting point of human physiology, we shall inquire into the functions, in the order we have adopted respecting the functions of the adult. Over many of the topics that will have to engage attention, the deepest obscurity rests; whilst the hypotheses, indulged regarding them, have been of the most fanci-

ful and mystical character.

a. Animal Functions.—The external senses in general are manifestly not in exercise during fætal life; of this there can be no doubt, as regards the sense of sight; and the same thing probably applies to the taste, smell, and hearing. With regard to tact, however, we have the best reason for believing, that it exists, particularly towards the latter periods of utero-gestation. The cold hand, applied over the abdomen of the mother, will instantly elicit the motions of the child. The brain and nervous system of the fætus must, therefore, have undergone the development necessary for the reception of the impression made through the medium of the mother; to convey such impression to the percipient organ, and to accomplish perception.

The existence of most of the internal sensations or wants would of course be supererogatory in the fætal state, where the functions, to which they minister after birth, are themselves wanting. It is probable, that there is no digestion except of the mucous secretions of the tube; no excretion of fæces or urine, and certainly there is no pulmonary respiration. It is not unlikely, however, that internal impressions, originating in the very tissue of the organs, may be communicated to, and appreciated by, the brain.

b Comment, Soc. Reg. Scient, Gotting. 1778; and Curling, op. cit. p. 66.

<sup>&</sup>lt;sup>2</sup> See Oesterreicher, De Gubernaculo s. d. Hunteriano, 1828; and Neue Darstellung der Lehre der Ortsveränderung der Hoden, 1830; also, Rathke, Abhandlungen zur Bildungs und Entwickelungsgeschichte des Menschen, u. s. w., Th. i., 1832 — cited by Valentin, art. Fætus, in Encyclop. Wörterbuch der Medicin. Wissenschaft. xii. 381, Berlin, 1835; and J. B. Curling, on the Structure of the Gubernaculum and on the Descent of the Testis, in London Lancet, April 10, 1841, p. 70; and in Practical Treatise on the Diseases of the Testes, &c., p. 32, Lond. 1843.

We have strong reason for believing, that pain may be experienced by the fœtus; for if it be destroyed by any sudden influence in the latter periods of pregnancy, death will generally be preceded by irregular movements which are manifest to the mother, and frequently lead her to anticipate the result. Adelon asks, whether it may not be affected, under such circumstances, with convulsions, similar to those that animals experience when they die suddenly, especially from hemorrhage? It is impossible to reply to this ques-

tion, but that the child suffers appears evident.

The most elevated of the functions of relation — the mental and moral faculties - would seem to be needless to the fœtus, and consequently little, if at all, exercised. Bichat and Adelon,a considering that its existence is purely vegetative, are of opinion that they are not exerted at all. Cabanis, however, suggests, that imperfect essays may, at this early period, be made by virtue of the same instinct that impels animals to exercise their organs prior to the period at which they are able to derive service from them; as in the case of the bird, which will shake its wings before they are covered with feathers, and when yet incapable of bearing them. It is difficult to deny to the fœtus all intellectual and moral manifestations. They must doubtless be obscurely rudimental; still, we may conceive that some may exist, if we admit that the brain is in a state for the perception of impressions, that tact is practicable, and instinct in full activity. We find, moreover, that the power of motion, voluntary as well an involuntary, exists certainly after the fifth month, and probably much earlier, could it be appreciated. During the latter months of utero-gestation, the motion of the fœtus appears to be almost incessant, and can be distinctly felt by placing the hand upon the abdomen. At times, indeed, it is manifest to the sight. The cause of these movements is by no means clear. It is probable, that they are instituted for the purpose of inducing a change in positions, which may have become irksome, and for assuming others; for we have already remarked, that the fœtus readily appreciates any sudden succussion given to it through the mother, - hence that it possesses tact, and, as we can readily understand, may experience fatigue from the maintenance of an inconvenient posture. This impression is conveyed to the brain, which sends out volition to the appropriate muscles, and the position is changed. All this proves, that the cerebral functions are exerted, but for a few definite objects only.

The function of expression is of course almost, if not entirely, null in the fœtus. There are cases upon record, where children are said to have cried in utero, so as to be heard distinctly, not only by the mother, but by those around her. Indeed, the objection, that an infant may respire before it is born, and yet not come into the world alive, — in which case there will be buoyancy and dilatation of the lungs, has been brought forward against the doci-

<sup>Physiologie de l'Homme, 2de édit. iv. 420, Paris, 1829.
Rapport du Physique et du Moral de l'Homme, Paris, 1802.</sup> 

masia pulmonum or lung-proof of infanticide. We would not be understood as believing these cases to be mere fabrications, or the phenomenon impossible, - except, indeed, whilst the membranes are in a state of integrity. When they have given way, and the child's mouth presents towards the os uteri, breathing and the vagitus may be practicable and may have occurred; but very positive and unexceptionable testimony is required to establish such an astounding event.a

b. Functions of Nutrition. - These functions are not as numerous in the fœtus as they are in the adult. Their object is, however, the same; - the formation of the various parts of the organized machine, and their constant decomposition and renovation.

One of the least tenable hypotheses, that have been entertained regarding the embryo at its first formation, is - that, for the first month — and why the period is thus limited is not apparent the vitality of the fœtus is not independent, but is a part, -an offset, as it were, - of that of the mother; that it has no separate powers of existence; no faculty of self-evolution; and that its organs are nourished by the plastic materials, which it incessantly derives from the maternal blood.b It appears manifest, that from the very moment of the union of the materials furnished by both sexes at a fecundating copulation, the elements of the new being must exist; and it must possess, within itself, the faculty of selfevolution; otherwise, how can we understand the phenomena that take place in the ovarium after fecundation? It is admitted that this last organ furnishes the unfecundated ovum; and that the sperm must come in contact with this ovum; after which, fecundation is accomplished, and immediately the ovum undergoes a farther development; escapes, in due time, from the viscus in which it was formed; is laid hold of by the Fallopian tube; passes through that canal, and is deposited in the interior of the uterus, with which it ultimately contracts adhesions. But all this requires time. The ovum does not probably reach the uterus, in the human female, until about the end of a week; and some time must still elapse before such adhesions are effected; and, consequently, before any thing like maternal blood, whence the plastic materials are derived, according to the view in question, could be sent to it. During the whole of this time, the embryo doubtless derives its nourishment from the albuminous matters with which it is surrounded in the ovum itself, in the same manner as the young of the oviparous animal obtains the nutriment necessary for its full development, during incubation, from the matters surrounding it; in which case the supply of fresh plastic materials, derived from the maternal blood, is obviously impossible.º But, in due time, after it has attained the interior of the uterus, it is compelled to absorb appropriate nutriment from the

a See a case of this kind by Dr. Collins, in his Practical Treatise on Midwifery, &c. Lond. 1835; also, Beck's Medical Jurisprudence, 6th edit. Philadelphia, 1838.

b Geddings, art. Acephalous, Amer. Cycl. of Pract. Medicine, P. ii. p. 142, Phila-

delphia, 1833.

Granville's Graphic Illustrations of Abortion, p. vii., London, 1834.

mother; the minute quantity existing in the ovum, at this early period, being totally insufficient for the development which the fætus is destined to attain. In this last respect, the human ovum differs from the eggs of oviparous birds, which are hatched out of the body, and contain sufficient nutriment for full fætal evolution. In treating of this subject, Dr. Granville has the following remarks - "What stronger proof need be required of the existence of an adherent life principle in the ovulum, which is, at one time at least, indeed, I suspect throughout the period of gestation, independent of any connexion with the parent mother. Yet none of the earlier writers who adopted the ovarian theory of generation have ever asked themselves this question --- what supports, the vitality of a fecundated ovulum after it has left the ovarium, and previously to its becoming connected with the womb? fact, the subject had never been mooted, before the more modern physiologists took it up and satisfactorily explained it:" - and he adds in a note, "the whole of the English physiologists, writers on midwifery and lecturers, whether ancient or modern, are entirely silent on this important stage of embryonic life." It is a topic, however, discussed at length in the first edition of this work,

and much expanded in the subsequent edition.a

Since the time of Hippocrates, Aristotle, and Galen, different anatomists and physiologists have asserted, that the umbilical vein is the only channel through which nutriment reaches the fœtus, or, in other words, that the whole of the nourishment which the fœtus receives is from the placenta; but the facts, to which allusion has already been made, are sufficient to overturn this hypothesis. It is impossible, that the placenta can have any agency until it is in esse. Such an explanation of the process of feetal nutrition could only hold good after the first periods, and then, as we shall see, it is sufficiently doubtful. Accordingly, some of the most distinguished of modern physiologists, who have devoted their attention to embryology, have completely abandoned the idea of placental agency during the first months; and they, who have invoked it at all, have usually done so, as regards the after periods only. On all this subject, however, we have the greatest diversity of views. Lobstein, for example, affirms, that the venous radicles of the rudimental placenta obtain nutritive fluids from the mother during the first days only, until the period when the arteries are formed; but that, after this, all circulation between the uterus and placenta ceases, and the fluid of the umbilical vesicle, the liquor amnii, and the jelly of the cord, are the materials of nutrition. Meckel<sup>c</sup> thinks the placenta is never the source of nutritive materials. He regards it as an organ for the vivification of the blood of the fœtus, analogous to the organ of respiration in the adult; and nutrition is, in his opinion, accomplished by the matter of the

Human Physiology, 1st edit. ii. 365, Philad. 1832, and 2d edit. ii. 413, Philad.
 1836.
 Essai sur la Nutrition du Fœtus, Strasbourg, 1802.

<sup>.</sup> Handbuch, u. s. w., Jourdan and Breschet's French translation, iii. 784.

umbilical vesicle in the beginning, by the liquor amnii until midterm, and by the jelly of the cord until the end. According to Béclard, nutrition is effected during the first weeks, by the fluid of the umbilical vesicle; afterwards by the liquor amnii, and the jelly of the cord; and as soon as the ovum becomes villous, and developes the placenta, by that organ. Dr. Montgomery has described several "decidual cotyledons," as he terms them, which are best seen about the second or third month, and are not to be found at the advanced period of gestation. These are small cuplike elevations on the external surface of the decidua vera, having the appearance of little bags, the bottoms of which are attached to or embedded in its substance; they then expand a little, and again grow smaller towards their outer or uterine end, which, in by far the greater number of them, is an open mouth when separated

Fig. 277.



Decidual Cotyledons. - (Montgomery.)

from the uterus; how it may be when adherent he does not profess to say, nor does he offer any decided opinion as to their precise nature or uses; but from having on more than one occasion observed within their cavity a milky or chylous fluid, he is disposed to consider them reservoirs for nutrient fluids separated from the maternal blood, to be thence absorbed for the support and development of the ovum. "This view," says Dr. Montgomery, "seems strengthened when we consider, that at the early periods of gestation, the

ovum derives all its support by imbibition, through the connexion existing between the decidua and the villous processes covering the outer surface of the chorion." Adelone is of opinion, that two sources of nutrition ought alone to be admitted,—the umbilical vesicle, which is the sole agent for nearly two months, and the placenta for the remainder of the period. M. Velpeaud equally thinks, that the nutriment of the ovum is derived from different sources at different periods of intra-uterine existence. The embryo, he says, is at first but a vegetable, imbibing the surrounding humours. The villi of its circumference, which are true cellular spongioles, obtain nutritive principles in the Fallopian tube and the uterus, to keep up the development of the vesicles of the embryo; after which the new being is nourished like the chick in ovo, or rather like the plantule, which is, at first, altogether deve-

Signs and Symptoms of Pregnancy, p. 133, Lond. 1837.
 Physiologie de l'Homme, 2de édit., iv. 397, Paris, 1829.

Embryologie, ou Essai Anatomique sur le Fétus Humain, Paris, 1821.

<sup>&</sup>lt;sup>4</sup> Embryologie ou Ovologie, &c., Paris. 1833; and Traité Elémentaire de l'Art des Accouchemens, Professor Meigs's translation, 2d edit. p. 213, Philad. 1838. See, also, Broussais, Traité de Physiologie, &c., Drs. Bell and La Roche's translation, 3d Amer. edit. p. 535, Philad. 1832.

loped at the expense of principles inclosed in its cotyledons. It gradually exhausts the vitelline substance contained in the umbilical vesicle. The emulsive substance of the reticulated sac of the allantoid pouch is also gradually absorbed. The end of the second month arrives; the vessels of the cord are formed, and the placenta is developed; by its contact with the uterus, this organ obtains reparatory materials, elaborates them, and forms from them a fluid more or less analogous to blood, and this fluid is absorbed

by the radicles of the umbilical vein.

We find, consequently, some of the most distinguished physiologists of the age denying - as it would seem that every one ought to deny - that the nutrition of the fœtus takes place solely by means of blood sent by the mother to the fætus. If we search into the evidence afforded us by transcendental anatomy, we find that amidst the various singular monstrosities met with, there would appear to be but one thing absolutely necessary for fætal development, - an absorbing surface, surrounded by a nutritive substance capable of being absorbed. Head, heart, - every thing, in short, except organic nervous system, vessels, and cellular tissue, - may be wanting, and yet the fœtus may grow so as to attain its ordinary dimensions. We have the most incontestable evidence, that neither the placenta nor umbilical cord is indispensably necessary for fætal development. Adelon disposes of this in the most summary manner, by affirming, that "there is no authentic instance of a fætus, devoid of umbilical cord and placenta, attaining full uterine growth." The case is not, however, got rid of so easily. The marsupial animals breed their young without either placenta or cord. The embryos are inclosed in one or more membranes, which are not attached to the coats of the uterus, and are supplied with nourishment from a gelatinous matter by which they are surrounded. Thomas Bartholin, moreover, during his travels in Italy, saw an individual, forty years old, who was born without anus, penis, or umbilicus; and M. Velpeau cites cases from Ruysch, Samson, Chatton, Rommeil, Denys, Fatio, V. Geuns, Sue, Penchienati, Franzio, Desgranges, Kluyskens, Pinel, Mason, Osiander, Dietrich, Von Froriep, and Voisin; but as these cases militate against his views of embryotrophy, he attempts to diminish their force by affirming, that the observations, which he had made, satisfy him, that all the fœtuses thus born had died in utero, in consequence of the destruction of the cord, or the closure of the umbilicus; or else, that the umbilicus existed, but was hidden or lost in the extroversion of the bladder, almost always remarked in those that lived. Now, passing by the singular deduction of M. Velpeau, that his observations have satisfied him of the incorrectness of observations made by men, many of whom have long since passed away, - long before he existed, - as well as the facts relating to the marsupial animal, and that the fætus, in extra-uterine pregnancies, has frequently no placenta, - with the case cited by Dr. Good, from Hoffmann, of a fætus born in good health and vigour, with the funis sphacelated and divided into two parts; and one

by Stalpart Van der Wiel,a of a living child, exhibited without any umbilicus as a public curiosity, - a case, observed by Dr. Goodb himself, appears to us to be impregnable. The case in question occurred in 1791. The labour was natural; the child. scarcely less than the ordinary size, was born alive; cried feebly once or twice after birth, and died in about ten minutes. The organization, both internal and external, was imperfect in many parts. There was no sexual character whatever; neither penis nor pudendum; nor any interior organ of generation. There was no anus, no rectum, no funis, no umbilicus. The minutest investigation could not discover the least trace of any. With the use of a little force, a small, shrivelled placenta, - or rather the rudiment of a placenta, - followed soon after the birth of the child. without a funis or umbilical vessels of any kind, or any other appendage by which it appeared to have been attached to the child. In a quarter of an hour afterwards, a second living child was protruded into the vagina and delivered with ease, being a perfect boy, attached to its placenta by a proper funis. The body of the first child was dissected in the presence of Dr. Drake of Hadleigh, and of Mr. Anderson of Sunbury, to both of whom Dr. Good appeals for the correctness of his statement. In the stomach, a liquor was found resembling the liquor amnii. How could nutrition have been effected, then, in this case? Certainly not by blood sent from the mother to the child, for no apparatus for its conveyance was discoverable; and are we not driven to the necessity of supposing, that the food must have been obtained from the fluid within the ovum? This case, - with the arguments already adduced, - seems to constrain us to admit, that the liquor amnii may have more agency in the nutrition of the new being than is generally granted. Professor Monro, amongst other reasons, -- all of which are of a negative character, - for his disbelief in this function of the liquor amnii, asserts, that if the office of the placenta be not that of affording food to the embryo, it becomes those, who maintain the contrary doctrine, to determine what other office can be allotted to it, and that till this is done, it is more consistent with reason to doubt the few and unsatisfactory cases, at the time brought forward, than to possess ourselves with facts directly contradictory of each other. The case given by Dr. Good, since Professor Monro's remarks were published, is so unanswerable, and so unquestionable, that it affords a positive fact, of full or nearly full fætal development, independently of placenta and umbilical cord; and the fact must remain, although our ignorance of the functions of the placenta be "dark as Erebus."

The following case, with which the author was obligingly favoured by his friend, Dr. Wright, of Baltimore, has an instruc-

\* Observat. Rar. Med. Chirur. cent. ii. p. 1, Obs. 32.

b Case of Preternatural Fætation, with Observations, read before the Medical Society of London, Oct. 20, 1794; and Study of Medicine, in Physiological Proem to Class v. Genetica.

c Edinburgh Medical Essays, ii. 162.

tive bearing upon the subject. The condition of the placenta was such as to lead that intelligent observer to conclude, that any circulation between the mother and the fœtus, through the placenta, was impracticable.

" Baltimore, September 26th, 1835.

" DEAR SIR, - In compliance with your request, I offer you the following plain and short statement of a case, which occurred in my practice, at the date indicated.

"On the 6th of December, 1833, I was requested to visit Mrs. T-, of this city, a young woman of large form, good constitution, and generally excellent health. She had been married about fifteen months, and I was now called to attend her first labour. She had felt occasional labour pains through the day, and was delivered of a fine, vigorous, female infant, in about four hours from the time of my call. The labour was, in all respects, natural, and as easy as is common - or consistent - with a first parturition. After the birth of the child, an hour, perhaps, was passed in waiting for secondary pains to effect the expulsion or favour the removal of the placenta, but no movement of this kind having then occurred, a gentle examination was made to ascertain whether that body might be easily and properly taken away. The vagina contained nothing more than the funis, - the outlet of the uterus was open, soft and extensible. The cord was gently followed into the uterine cavity, and the cake found near its fundus, retaining a close connexion with the uterus. The placental mass was large and firm, presenting to the touch a peculiar feeling - as of a dense sponge, full of coarse, granular or gravelly particles. Deeming it now proper to relieve the patient fully, a cautious effort was made to detach the placenta from the uterus, in order to its manual extraction. In pursuing this design, it was found, that the adhering surface of the former consisted of a uniform calcareous lamina or plate, rough to the finger, and exciting such a sensation or feeling, as would be caused by a sheet of coarse sand-paper. When the mass was detached, and brought away, the laminar surface, just referred to, was found to be a calcareous plate, uniformly covering the whole of the attached portion of the cake, - the entire surface of the utero-placental connexion. The calcareous matter, thus distributed, was thin, and readily friable, but, as before remarked, it appeared to constitute a uniform and superficial covering. The correspondent uterine surface the part from which the placenta had been separated - felt rough, but comparatively soft, imparting nothing distinctly of the calcareous or gritty feel. Out of the body, the placenta felt heavy, and emineutly rough throughout. When compressed or rubbed together, the large amount of nodular or granular matter, dispersed through its substance, was not only manifest to the touch, but a very audible crepitation or grating sound could be thus elicited from any, and every part of the mass.

"In this uncommon instance of placental degeneracy, both the mother and child were perfectly healthy and well. The latter, indeed, was remarkable for its fine size, perfect nutrition and vigour. From the condition of the cake, and the character of its adhesion to the uterus, I apprehended a more than ordinary liability to secondary affection, in the form of puerperal fever, - and whether influenced or not by the circumstances detailed, secondary fever did ensue on the third day from delivery, attended by the usual signs of puerperal hysteritis, which affection, however, was happily subdued

by general and topical bleeding, calomel, &c.

"With sincere regard, yours, "T. H. WRIGHT.

" PROFESSOR DUNGLISON.

" P.S. - The child, referred to, is living, and, from its birth to the present, has considerably exceeded the common bulk of children at the same age. The mother is now far advanced in her second pregnancy."a

Since this letter was written, the same lady has been delivered of another healthy child; and although the maternal surface of the placenta was of the same calcareous character, the deposition was not to the same extent as in the first pregnancy. A similar

a See an analogous case by Madame Buisson Dauthez, in Gaz. Méd. de Paris, Juillet, 1842.

case has been described by Mr. Gilbert, of Beaminster, England.a

Amongst those physiologists, who admit the liquor amnii to be a fluid destined for fætal nutrition, a difference prevails, regarding the mode in which it is received into the system. Osiander, b Brugmans, Van den Bosch, Fohmann, Carusd and others think, that it is absorbed through the skin. In the fætal state, the cuticle is extremely thin; and, until within a month or two of the full period, can be scarcely said to exist. There is consequently not that impediment to cutaneous absorption, which, we have seen, exists in the adult. The strong argument, however, which they offer in favour of such absorption is the fact, that the fœtus has been developed, although devoid of both mouth and umbilical cord; and Professor Monro, in opposing the function ascribed to the liquir amnii, refers to cases of monstrous formation, in which no mouth existed, nor any kind of passage leading to the stomach. Others, as Hallere and Darwin, are of opinion, that the fluid enters the mouth and is sent on into the stomach and intestines; and in support of this view they affirm, that the liquor amnii has been found in these viscera; - that it has been shown to exist in the stomach and pharynx. Heister, on opening a gravid cow, which had perished from cold, found the liquor amnii frozen, and a continuous mass of ice extending to the stomach of the fœtus.

The physiologists, who believe that the liquor amnii is received into the stomach, differ as to what happens to it in that organ. Some suppose, that it is simply absorbed without undergoing digestion; others, that it must first be subjected to that process. According to the former opinion, it is simply necessary, that the fluid should come into contact with the mucous membrane of the alimentary passages; and they affirm, that if digestion occur at all, it can only be during the latter months. Others, however, conceive that the waters are swallowed or sucked in, and that they undergo true digestion. In evidence of this they adduce the fact of meconium existing at an early period in the intestinal canal. which they look upon as evidence that the digestive function is in action; and in farther proof of this, they affirm, that on opening the abdomen of a new-born infant the chyliferous vessels were found filled with chyle, which could not, they say, have been formed from any other substance than the liquor amnii; and lastly, that fine silky down has been found in the meconium, similar to that which exists on the skin of the fœtus, and which is conceived to have entered the mouth along with the liquor amnii. These rea-

Lancet, for Dec. 16, 1837, p. 418.

b Handbuch der Entbindungskunde, tom. i. p. 237.

5 Adelon, Physiologie de l'Homme, iv., 389, Paris, 1829.

De Natura et Utilitate Liquoris Amnii, Ultraject. 1792.
 Lehrbuch der Gynäkologie, Th. ii. s. 27, Leipz. 1828.
 Elementa Physiologiæ, viii. 205.

<sup>&</sup>lt;sup>f</sup> Zoonomia, vol. i., London, 1796. See Meckel's Handbuch, u. s. w., Jourdan and Breschet's translation, iii. 784, Paris, 1825.

sons have their weight, but they cannot explain the development in the cases above alluded to, in which there was no mouth; and of course they cannot apply to the acephalous fœtus. Moreover, it has been properly remarked, that the presence of meconium in the intestinal canal — admitting that it is the product of digestion, which is denied by many — merely proves that digestion has taken place, and the same may be said of the chyle in the chyliferous vessels: neither one nor the other is a positive evidence of the digestion of the liquor amnii. Both might have proceeded from the stomachal secretions. It has also been affirmed, that meconium exists in the intestines of the acephalous fœtus; and in those in which the mouth is imperforate. Lastly, with regard to the down discovered in the meconium, it has been suggested as possible that it may be formed by the mucous membrane of the intestine, which so strongly resembles the skin in structure and functions.

Others have supposed that the liquor amnii is received through the respiratory passages, from the circumstance, that, in certain cases, the fluid has been found in the trachea and bronchia; some presuming, that it readily and spontaneously enters at the nostrils and passes to the trachea and bronchia; others, that it is forced in by the pressure of the uterus; and others, again, that it is introduced by the respiratory movements of the fœtus. Views have differed in this case, also, regarding the action exerted upon it after introduction; some presuming that it is absorbed immediately; others, that it is inservient to a kind of respiration; and that, during feetal existence, we are aquatic animals, - consuming the oxygen or atmospheric air, which Scheele, Lassaigne, and others have stated to exist in the fluid. It is scarcely necessary to oppose seriously these gratuitous speculations. The whole arrangement of the vascular system of the fœtus, so different from that which is subsequently established, and the great diversity in the lungs, prior and subsequent to respiration, would be sufficient to refute the idea, had it even been shown, that the liquor amnii always contains one or other of these gases, which is by no means the fact. The case of the acephalous fœtus is also an obstacle to this view as strong as to that of the digestion of the liquor amnii.

As if to confirm the remark of Cicero — "nihil tam absurdam, quod non dictum sit ab aliquo philosophorum," — it has been advanced by two individuals of no mean pretensions in science, that the liquor amnii may be absorbed by the genital organs or by the mammæ. Lobsteine supports the former view; Okend the latter. He asserts, that the fluid is laid hold of by the mammæ, is elaborated by them, and conveyed from thence into the thymus

<sup>\*</sup> De Liquoris Amnii Utilitate, Copenhag. 1795.

<sup>&</sup>lt;sup>b</sup> Archiv. Général. de Med. ii. 308.

c Essai sur la Nutrition du Fœtus, p. 102, Strasbourg, 1802.

d Zeugung, p. 162, Bamberg, 1805.

gland, the thoracic duct, and the vascular system of the fætus! Hence, in this opinion, the necessity of both sexes possessing nip-

ples before birth.a

Of these various opinions, the one that assigns the introduction of the fluid to the agency of the cutaneous absorbents appears to carry with it the greatest probability. It must be admitted, however, that the whole subject is environed with obscurity, and requires fresh, repeated, and accurate experiments and observa-

tions to enlighten us.

But it may now be asked, with Monro, what are the nutritive functions performed by the placenta? We have before alluded to the different views, entertained regarding the connexion between the placenta and the uterus. Formerly, it was universally maintained, that vessels pass between the mother and the maternal side of the placenta, and that others pass between the fœtus and the fætal side, but that the two sides are so distinct, as to justify their being regarded as two placents, - the one maternal, the other fætal, - simply united to each other. At one time, again, it was supposed, that a direct communication exists between the maternal and fætal vessels, but this notion has long been exploded. We have decisive evidence, that the connexion is of the most indirect nature. Wrisberg made several experiments, which showed that the fluid of the fætal circulation is not drained when the mother dies from hemorrhage. It has been affirmed, too, that if the uterine arteries be injected, the matter of the injection passes into the uterine veins, after having been effused into the lobes of the placenta. If, on the other hand, the injection be thrown into the umbilical arteries, the matter passes into the umbilical vein, and is effused into the fætal side of the placenta, but does not pass into the uterine vessels. When, however, an odorous substance, like camphor, is injected into the maternal veins of an animal, the fœtal blood ultimately assumes a camphorated odour, and when animals have been fed on madder during gestation, the colouring matter has been found in the fœtus; b also when the human female has taken rhubarb, evidence of it has been found in the liquor amnii, in the serum of the blood of the umbilical vessels, and in the first urine of the infant.c Magendied injected this substance into the veins of a gravid bitch, and extracted a fœtus from the uterus at the expiration of three or four minutes: the blood did not exhibit the slightest odour of camphor; whilst that of a second fætus, extracted at the end of a quarter of an hour, had a decidedly camphorated smell. This was the case, also, with the

<sup>&</sup>lt;sup>a</sup> Hedenus, in art. Brust (weibliche,) Encyclopäd. Wörterb der Medicinish. Wissensch. vi. 351, Berlin, 1831. See, especially, on the whole subject of fætal nutrition, Adelon, Physiologie de l'Homme, 2de édit. iv. 383, Paris, 1829; and Meckel's Handbuch, u. s. w., Jourdan's French translation, iii. 781; or Doane's English translation from the French, Philad. 1832.

b Dr. Musey, in Amer. Journ. of the Med. Sciences, for November, 1829.
c Granville's Graphic Illustrations of Abortion, &c. p. xx. Lond. 1834.
d Précis Elémentaire, ii. 552.

other fætuses. Such communication may, however, have been owing to the same kind of transudation and imbibition, of which we have spoken under the head of absorption, and may consequently be regarded as entirely adventitious; and the fact of the length of time, required for the detection of the odorous substance favours this idea; for if any direct communication existed between the mother and the fœtus, the transmission ought certainly to have been effected more speedily. The transmission of substances from the fætal to the maternal placenta is yet more difficult. Magendie was never able to affect the mother by poisons injected into the umbilical arteries and directed towards the placenta; and he remarks, in confirmation of the results of the experiments of Wrisberg, that if the mother dies of hemorrhage, the vessels of the fœtus remain filled with blood. They who consider, that there is no maternal and fætal portion of the placenta, or, rather, that it is all fætal, of course believe, where the matter of injection, thrown into the uterine vessels, has passed into the cells of the placenta, that it has been owing to the rupture of parts, by the force with which the

injection has been propelled.

Another fact, that proves the indirect nature of the connexion which exists between the parent and child, is the total want of correspondence between the circulation of the fœtus and that of the mother. By applying the stethoscope to the abdomen of a pregnant female, the beating of the fætal heart is observed to be twice as frequent as that of the mother. (See p. 422 of this volume.) Again, examples have occurred in which the fœtus has been extruded with the placenta and membranes entire. In a case of this kind, which occurred to Wrisberg, the circulation continued for nine minutes; in one described by Osiander, for fifteen minutes; in some, by Professor Chapman, from ten to twenty minutes; and in one, by Professor Channing, of Boston, and Dr. Selby, of Tennessee, where a bath of tepid water was used to re-suscitate the fœtus, for an hour. Marsond and Flajani relate cases in which life continued for the same time: Dr. Nehr, of Rehau, in Bavaria, has given a case in which the circulation of the child was unequivocal for seven hours after the sudden and decided death of the mother; and other cases are referred to by D'Outrepont in his comments on this one.f In other cases of a similar kind, where the child could scarcely breathe and was in danger of perishing, the life of the placenta was maintained by keeping it in water of a temperature nearly equal to that of the body, and the child has been saved. All these facts prove demonstratively, that the fœtus carries on a circulation independently of

4 Lond. Med. Gazette, August, 1833.

Granville, op. cit. part x. Lond. 1834.

Begin and General Anatomy, 5th edit. ii. 277, Philad. 1839.

Neue Zeitschrift für Geburtskunde, von Busch, D'Outrepont und Ritgen, Band. iv. Heft 1, s. 58, Berlin, 1836.

<sup>&</sup>lt;sup>1</sup> Ibid. s. 60. See, also, Diction. des Sciences Médicales, tom. xvii. p. 442; and Osiander's Handbuch der Entbindungskunde, B. iii. Abth. 2, s. 157.

that of the mother; and whatever passes between the fætal and maternal vessels is probably exhaled from the one and absorbed by the other, as the case may be. The fluid sent to the fœtus is supposed by some, - indeed by most physiologists - to be the maternal blood, modified or unmodified. Schreger, a however, and others maintain, that the communication of any nutritious fluid from the mother to the fœtus, and conversely, takes place by means of lymphatics, and not by bloodvessels; and that the maternal vessels exhale into the spongy tissue of the placenta the serous part of the blood, which is taken up by the lymphatics of the fætal portion,

and conveyed into the thoracic duct.

It has been before remarked, that Lobsteinb and Meckele suppose, that the gelatinous substance of the cord is one of the materials of fætal nutrition. This opinion they found on the circumstance of the albuminous nature of the substance, and the great size which it gives to the cord at the early periods of fœtal life, as well as on the great development of the absorbent vessels of the fœtus, that proceed from the umbilicus to the anterior mediastinum; whilst others invoke, also, the fluids of the umbilical and allantoid vesicles. All these speculations regarding the various sources of nutritive matter are sufficient evidence of the uncertainty that prevails on this interesting topic. It is manifest, however, that we cannot regard as nutritive matters those substances which are secreted by the fœtus itself. It is impossible, that any development can occur without the reception of materials from without. We have seen, that when the ovum passes from the ovarium to the uterus, it contains, within it, a molecule, and fluids which are probably destined for the nutrition of the new being, and which afford the necessary pabulum for the increase, that occurs between the period of impregnation and that at which an adhesion is formed between the ovum and the inner surface of the uterus. The mother, having furnished the nutritive material in the ovum, she must continue to provide it in the uterus; and as soon as the vascular communication is formed between the exterior of the ovum and the interior of the uterus, nutritive elements are doubtless received by the embryo; - for otherwise it would perish from inanition. then can be the nature of these elements? Do they consist of maternal blood, laid hold of by the fœtus at this early period, when no circulatory system is apparent; or are the bloodvessels distributed to the membranes of the ovum, to enable them to continue the secretion of that nutritive matter which they took with them from the ovarium, and which must necessarily have had a maternal origin? The latter certainly is the more probable supposition, and it is, as we have seen, an argument in favour of the amnion being supplied with blood from the uterus, rather than from the fœtus; for, if we admit it to be in any manner inservient to nutrition, its production must be extraneous to the body which it has

a De Functione Placentæ Uterinæ, Erlang. 1795. b Essai sur la Nutrition du Fœtus, Strasbourg, 1812.

<sup>·</sup> Handbuch, u. s. w., Jourdan and Breschet's translation, iii. 785, Paris, 1825.

to nourish. These observations apply equally to the jelly of the umbilical cord, which is probably secreted by the membranous envelopes, and may consequently be regarded as a nutritive material derived from the parent. Both, it is true, might be secreted by the fœtus from fluids furnished by the mother, and be placed in

depot, as the fat is in after existence.

Transcendental anatomy, then, instructs us, that the placenta and umbilical cord are not indispensable to fœtal nutrition; and compels us to infer with Meckela - one of the most eminent of modern anatomists and physiologists - that the human placenta may have no direct agency in embryotrophy. We are, therefore, necessarily driven to the conclusion, before laid down, - that, in order for a fætus to be developed in utero, it is but necessary, that there shall be an absorbing surface, surrounded by a nutritive substance, which will admit of being absorbed. Now, the cutaneous envelope of the fœtus - monstrous or natural - is such a surface, and the liquor amnii such a fluid; and the matter of the umbilical vesicle, with the jelly of the cord, when these parts exist, and possibly some material derived through the placenta — after it exists — may lend their aid; but the participation of these last agents is - to say the least — doubtful. The function of the placenta probably is, to admit of the feetal blood being shown to that circulating in the maternal vessels, so that some change may be effected in the former, which may better adapt it for serving as the pabulum, whence the secretions, from which the fætal organs have to be elaborated, may be formed. According to Dr. Reid, the blood of the mother, contained in the placental sac already described (p. 450 of this volume), and the blood of the fœtus contained in the umbilical vessels, can readily act and react on each other, through the spongy and cellular walls of the placental vessels and the thin sac ensheathing them, in the same manner as the blood in the branchial vessels of aquatic animals is acted upon by the water in which they float. If we admit this, however, it is obvious, that the nutritive fluid, when received into the system, will have to be made into blood by the action of the fætus, in a manner, bearing some analogy to what occurs in the adult, or in the simplest of living beings, in which the nutritive fluid is absorbed at the surface of the body. Of the mode in which such conversion is effected we are in the same darkness that envelopes all the mysterious processes which are esteemed organic and vital; but that the fœtus is capable of effecting it we have irrefragable proof in the oviparous animal, where there can be no communication, after the egg is laid, between the embryo and the parent. Yet we find it forming its own blood from the yolk surrounding it, and undergoing its full and regular development from causes seated in itself alone.

Of those physiologists, who consider that the mother sends her blood to the placenta, to be taken up by the fætal vessels, all do not conceive that it is in a state adapted for the nutrition of the

Loc. citat. b Edinb. Med. and Surg. Journ. Jan. 1841, p. 8.

new being: some are of opinion that the placenta, or the liver, or both, modify it, but in a manner which they do not attempt to explain. In favour of such an action being exerted by the placenta, they state that it is clearly the organ which absorbs the fluid, and that every organ of absorption is necessarily one of elaboration; - a principle which we have elsewhere proved to be unfounded; and, moreover, that the blood conveyed to the fætus by the umbilical vein, differs essentially in colour from that conveyed to it by the umbilical arteries, - a fact, which, we shall see, can be accounted for more satisfactorily. In support of the view, that a second change is effected in the liver, they affirm, that a great part of the fætal blood ramifies in the substance of that organ before it reaches the heart; a part only going by the ductus venosus; and that the great size of the liver, during fætal life, when its function of secreting bile can be but sparingly exerted, is in favour of this notion.2 The opinion, that some change is effected upon the blood in the liver, is certainly much more philosophical and probable than the belief of Haller, that the object of its passage through that organ is to deaden the force with which the mother projects the fluid into the fætal vessels. We have seen, that it is extremely doubtful, whether she transmit any; and that if she

does, the communication is very indirect. M. Geoffroy Saint-Hilaire appears to think, that the blood of the mother, which he conceives to be sent through the placenta to the fœtus, is unfitted for fœtal life, before it has undergone certain modifications. The blood, according to him, which leaves the placenta, proceeds in part to the liver, and the remainder to the heart. In the liver it forms the material of the biliary secretion, or at least of a fluid, which, when discharged into the intestines, irritates them, and provokes a copious secretion from the mucous or lining membrane. This mucus, according to M. Saint-Hilaire, is always met with in the stomach and intestines of the fætus; and the presence of meconium, and of other excrementitious matters in the intestines, shows, that digestion must have taken place. This digestion he considers to be effected upon the mucus, secreted in the manner just mentioned; and, in support of its being inservient to sanguification, he, affirms, that its quantity is too great for the simple purpose of lubricating the parts; that mucus is the first stage of all organic compounds; that it predominates in all young beings; is the foundation of every organ; more capable of assimilation than any other substance, &c. But independently of the whole of this view being entirely hypothetical, it cannot be esteemed probable, that the fœtus is nourished by one of its own secretions. All secretions must be formed from blood. Blood must, therefore, pre-exist in the fœtal vessels, and the process, indicated by Saint-Hilaire, be unnecessary.

b Philosophie Anatomique, Paris, 1818-22.

<sup>&</sup>lt;sup>a</sup> Velpeau, Traité de l'Art des Accouchemens; or Meigs's translation, 2d edit. p. 224, Philad. 1838.

Allusion has already been made to the opinions of Schreger on the nutrition of the fœtus. These were developed in a letter, written by him, in 1799, to Sömmering.a He considers, that all communication of nutritious matter between the mother and fœtus occurs through the lymphatics which he has described as existing in considerable numbers in the placenta and umbilical cord. The red blood, flowing in the maternal vessels, is too highly charged with carbon, and with other heterogeneous substances, he thinks, to serve for the nutrition of the fætus. Its serous part, which is purer and more oxygenized, is therefore alone exhaled. The uterine arteries pour this serum into the spongy texture of the placenta, whence it is taken up by the lymphatics of the fætal portion. These convey it along the umbilical cord to the thoracic duct, whence it passes into the left subclavian, vena cava superior, right auricle and ventricle, ductus arteriosus, and aorta; and, by the umbilical arteries, is returned to the placenta. In this course, it is mixed with the blood, and becomes itself converted into that fluid. When it attains the placenta, the blood is not poured into the cells of that organ, to be transported to the mother, but it passes into the umbilical vein, the radicles of which are continuous with the final ramifications of the umbilical arteries. Lateral pores, however, exist in the latter, which suffer fluids to escape, that cannot be elaborated by the fœtus, or that require again to be submitted to the maternal organs, before they are fitted for its support. These fluids, according to Schreger, are not absorbed by the veins of the uterus, but by the lymphatics of that viscus, which are so apparent in the pregnant state, and have been injected by Cruikshank, Meckel, &c. In his view, therefore, the conversion of the serous fluid into blood is chiefly effected in the lymphatic system, and it has been a favourite hypothesis with many physiologists, that those organs, regarding whose functions we are so profoundly ignorant, and whose development is so much greater during intra-uterine than extra-uterine existence, - as the thymus, and thyroid glands, and the supra-renal capsules, - are, in some way, connected with the lymphosis or hæmatosis of the fœtus. We have already referred to the conjectures, that these organs are diverticula for the blood of those parts, the functions of which are not exerted until an after period of existence. Broussaisb makes the thyroid a diverticulum to the larynx; the thymus a diverticulum to the lungs, and the supra-renal capsules to the kidneys. Notwithstanding these ingenious speculations, however, our darkness, with regard to the true functions of these singular organs, is not the less impenetrable.

To conclude. The most plausible opinion we can form on this intricate subject is, that the mother secretes the substances, which

<sup>\*</sup> Epistol. ad S. Th. Sömmering, De Functione Placentæ Uterinæ, Erlang. 1799. See, also, Richerand, op. cit., 13ème édit. § ccvi.

b Commentaires des Propositions de Pathologie, &c. Paris, 1829, or Drs. Hays and Griffith's translation, p. 214, Philad. 1832.

are placed in contact with the fætus, in a condition best adapted for its nutrition; that in this state they are received into the system, by absorption, as the chyle or the lymph is received in the adult, — undergoing modifications, in their passage through the fætal placenta, as well as in every part of the system where the elements of the blood must escape for the formation of the various tissues.

With regard to the precise nutritive functions executed in the fætal state, and first as concerns digestion, it is obvious, that this cannot take place to any extent, otherwise excrementitious matter would have to be thrown out, which, by entering the liquor amnii, would be fatal to its important functions, and probably to the very existence of the fætus. Yet, that some digestion is effected, is manifest from the presence of meconium in the intestines, which is probably the excrementitious matter, arising from the digestion of the mucous secretions of the alimentary canal.

2. Respiration, as accomplished by lungs, does not exist; and we have already seen, that the idea of the fœtus possessing the kind of respiration of the aquatic animal is inadmissible. An analogous function to the respiration of the adult, however, occurs, as respects the changes effected upon the blood. It is probable, that the blood is sent to the placenta to be aërated there, as it is in the lungs in extra-uterine existence. Such was the opinion of Sir Everard Home, of Girtanner, Stein, and we may say, such is the opinion of many of the most enlightened physiologists of the present day. The chief arguments adduced in support of this opinion are, - the absolute necessity for aëration to every living being, animal or vegetable; the no less necessity for a free circulation of blood - along the umbilical cord to and from the placenta to the life of the fœtus; and the analogy of birds, in which the umbilical vessels are inservient to respiration by receiving the external air through the pores of the shell, so that if the shell be greased, respiration is prevented, and the chick dies.b

The sensible evidences of these changes being effected by the placenta are not like those, which we possess regarding the aëration of the blood in its passage through the lungs of the adult, where the venous differs so essentially from the arterial blood. It is indeed asserted, in works of anatomy, that "the effete blood of the umbilical arteries becomes regenerated in the placenta, assumes a brighter hue, and is returned to the fœtus by the umbilical vein," but this is not in accordance with experiment and observation. Bichate made numerous dissections of young

<sup>\*</sup> Meckel's Handbuch, u. s. w. Jourdan and Breschet's Fr. translation, iii. 793, Paris, 1825; Meigs's translation of Velpeau, edit. cit. p. 225; and Granville's Graphic Illustrations of Abortion, p. xviii. Lond. 1834.

b Varnishes of organic animal matters, as albumen, gelatin, &c., have no effect in preventing the transmission of air. See Mr. Towne, Guy's Hospital Reports, Oct. 1839, p. 385.

c Anatomie Générale, ii. 344, Paris, 1818; and Researches Physiologiques sur la Vie et la Mort, p. 190, Paris, 1806.

pigs whilst yet in utero, and he uniformly found the blood of the arteries and veins presenting the same appearance and resembling the venous blood of the mother. Not the slightest difference was observed between the blood of the aorta and that of the vena cava, or between that of the carotid artery and of the jugular vein. He made the same observations in three experiments of a similar kind on fætuses of the dog. He also frequently examined human fœtuses that had died in utero, and always found the same uniformity between the arterial and venous blood: hence he concludes, that there is no difference between the arterial and venous blood of the fœtus, at least in appearance. Similar experiments by Autenrietha furnished like results. Dr. Granville, too, affirms, that he has never been able to detect the least difference between the arterio-umbilical and venous-umbilical blood in the many cases he has examined, and it is important to bear this fact in mind. inasmuch as it may be received as one of the evidences that a foetus has not respired. The apparent identity, however, between the blood passing to the placenta by the umbilical arteries and that returning by the cord cannot be real. The slightest reflection will show, that they must be different, and such is the opinion, from observation, of Bostock, Jeffrey, and others. It is from the blood, carried by the umbilical vein and distributed over the body, that all the organs of the fœtus have to derive the materials of their nutrition and development; and being deprived of these materials, the fluid must necessarily be different in the umbilical arteries from what it was in the umbilical vein. The researches of more modern chemistry have not been directed to the fætal blood, but Fourcrove analyzed it, and found it to differ materially from the blood of the child that had respired. He asserts, that its colouring matter is darker, and seems to be more abundant; that it is destitute of fibrin and phosphoric salts, and is incapable of becoming florid by exposure to the influence of atmospheric air. It has been found, too, that the globules of the fætal blood do not resemble those of the blood of the mother. The fact, however, of the similarity in appearance between the arterial and venous blood of the fœtus, is no evidence that respiration is not one of the fœtal functions, inasmuch as the same thing is observed in fishes. Under the head of circulation it was remarked, that the coloration of the blood is perhaps of no farther importance than as indicating that the vital change of aëration has taken place in the lungs. In this case, we have the vital change effected without any such coloration. Yet

a Dissertatio Sistens Experimenta circa Calorem Fœtûs, et Sanguinem Ipsius Instituta, Tubing. 1799; also, Velpeau, by Meigs, p. 224.

b Graphic Illustrations, &c., p. 20. See, also, J. Müller's Handbuch, u. s. w., Baly's translation, p. 318, Lond. 1837; Richerand, Elémens de Physiologie, 13ème édit. par Bérard ainé, § ceviii.; Cuvier, Leçons d'Anatomie Comparée, iv. 298, Paris, 1799; and Adelon, Physiologie de l'Homme, 2de édit. iv. 405.

<sup>&</sup>lt;sup>6</sup> Dr. Chapman, Philadelphia Journal of the Med. and Phys. Sciences, i. 10; and the Physiology of the Fœtus, Liver and Spleen, by George C. Holland, M.D., p. 154, Lond. 1831.

<sup>6</sup> Annales de Chimie, vii. 165.

how, it may be asked, is the modification in the blood produced where no placenta and no umbilical cord exist? And can we suppose that in such cases the aëration is effected by the liquor amnii containing an unusual quantity of oxygen, as has been presumed by some physiologists? We have before remarked, (p. 446 of this volume,) that J. Müller, was unable to detect oxygen in the liquor amnii, and he found that when fish were placed in it, they died as soon as in oil.<sup>a</sup> These are embarrassing questions—more easily propounded than answered. By some, it has been presumed, that the liver, even in the adult, performs a function supplementary to that of the lungs, and the great size of the organ, in the fœtus, has been conceived to favour the idea, that it may separate carbon and other matters freely from the system, and in this way, be depuratory; but the grounds for this presumption are not, we think,

impregnable.

3. It is in the fætal circulation that we observe the most striking peculiarities of intra-uterine existence. Of its condition at the very earliest periods we know little that is not conjectural. We will, therefore, consider it as it is effected during the last months of uterogestation. From the sketch already given of the circulatory organs of the fœtus, it will be recollected, - 1st. That the two auricles of the heart communicate by an aperture in the septum, called the foramen ovale, which has a valve opening towards the left ventricle. 2dly. That near the orifice of the vena cava inferior is the valve of Eustachius, so situate as to direct the blood of the cava into the foramen ovale. 3dly. That the pulmonary artery has a vessel passing from it into the aorta, -- the ductus arteriosus. 4thly. That two arteries, called umbilical, proceed from the internal iliacs to the umbilicus and placenta; and lastly, That the umbilical vein from the placenta pours part of its blood into the vena porta; and part passes by the ductus venosus, — a fætal vessel, — into the inferior cava.

The course of the circulation, then, is as follows: - the blood of the umbilical vein, - the radicles of which communicate with those of the umbilical arteries in the placenta, - proceeds along this vein to the umbilicus, and thence to the liver. A part of this traverses the ductus venosus, enters the vena cava inferior, and becomes mixed with the blood from the lower parts of the fœtus; the remainder passes into the vena portæ, is distributed through the liver, and, by means of the hepatic veins, is likewise poured into the vena cava. In this manner it attains the right auricle. Owing to the arrangement of the valve of Eustachins, the blood passes immediately through the foramen ovale into the left auricle, - without being mixed with the fluid proceeding from the upper parts of the body, - into the right auricle through the vena cava superior. The left auricle is consequently as much developed as the right, which it would scarcely be, did it receive only the blood from the lungs. Were it not as large, it is obvious, that it would be insufficient to carry on the circulation when the

whole of the blood passes through the lungs, and is poured into it

after respiration is established.

The above are the opinions of Wolff and Sabatiera regarding the use of the Eustachian valve. According to this view, if the valve did not exist, the aërated blood, conveyed to the heart by the ductus venosus, instead of being directed into the left auricle through the foramen ovale, would pass into the right auricle, and thence, - in part, at least, - into the right ventricle; from which it would be transmitted, through the pulmonary artery and ductus arteriosus, into the descending aorta; so that no part of the body, above the opening of the duct into the aorta, could receive the aërated blood, whilst much of that, which passes along the aorta, would be returned to the placenta by the umbilical arteries. But as the blood is directed into the left auricle by the Eustachian valve, it passes from thence into the left ventricle, and is forced by it into the aorta, which distributes it to every part of the system, and thus conveys the regenerated fluid to every organ. Dr. Wistarb also suggested, that, without this arrangement of the Eustachian valve, the coronary arteries, distributed to the heart, would be unfit for supporting the life of that organ, inasmuch as they would be deprived of a regular supply of revivified blood. From the left auricle, the blood passes into the left ventricle, and from the left ventricle into the ascending aorta, and to the upper parts of the body, from which it is brought back, by the vena cava superior, into the right auricle; thence it is transmitted into the right ventricle, and, by the contraction of the ventricle, into the pulmonary artery. By this vessel it is sent, - the greater part through the ductus arteriosus into the descending aorta, and a small part to the lungs. From the' lungs, it is returned into the left auricle by the pulmonary veins. Through the descending aorta, the blood, conveyed in part by the ductus arteriosus, and in part by the contraction of the left ventricle, is distributed, partly to the lower extremities, from which it is returned by corresponding veins into the vena cava inferior, and partly by the umbilical arteries to the placenta.

This view of the circulation supposes, — what is disputed, — that the blood of the vena cava superior and of the vena cava inferior does not undergo admixture in the right auricle; whence it would follow, that some parts of the body receive a purer blood than others, — the upper parts, as the head and neck, receiving that which flows immediately from the placenta, whilst the lower parts do not obtain it until it has circulated through the upper. Under any view it is manifest, that not the whole of the blood is distributed to the organ of aëration, as in the adult, but a part

only, as in the batrachia.

Bichat and Magendiec contest the explanation of Wolff and Sa-

Traité Complet d'Anatomie, ii. 224, and iii. 387; and Mémoir de l'Académ. des
 Sciences, pour 1744.
 System of Anatomy, 3d edit. edited by Dr. Horner, ii. 76, Philad. 1823.

Précis Elémentaire de Physiologie, 2de édit. ii. 550, Paris, 1825.

batier, regarding the use of the valve of Eustachius and the non-admixture of the blood of the two cavæ in the right auricle. In their opinion, the two bloods do commingle; but - owing to the existence of the foramen ovale, and the arrangement of the valve of Eustachius - the left auricle is filled simultaneously with the right: and, consequently, the same kind of blood must be distributed to both the upper and lower portions of the body. The uses of the foramen ovale and ductus arteriosus are explained as follows. As the left auricle receives but little blood from the lungs, it could furnish but a small quantity to the left ventricle, did it not receive blood through the foramen ovale; and again, as the lung is exerting no function, during the state of fœtal life, the blood is sent along the pulmonary artery and ductus arteriosus into the aorta. so that the contraction of both ventricles is employed in propelling the blood along the aorta to the lower parts of the body and to the placenta. Without this union of forces, it is conceived, that the blood could not be urged forward as far as the placenta.

Experiments, by Dr. John Reid, a favour the views of Wolff and Sabatier. He took a fætus of about seven months, and threw simultaneously a red-coloured injection up the vena cava inferior. and a yellow-coloured one down the vena cava superior. On tracing the red injection upwards, it was found to have passed through the foramen ovale and to have filled the left side of the heart, without any intermixture with the yellow, except very slightly at the posterior part of the right auricle. Not a drop of the yellow appeared to have accompanied the red into the left side of the heart. From the left side of the heart it ascended the aorta. and filled all the large vessels going to the head and upper extremities. The injection, in all these vessels, had not the slightest tinge of vellow. On tracing the yellow injection downwards, he found it filling the right auricle and the right ventricle, whence it proceeded along the pulmonary artery, and filled the ductus arteriosus, and branches going to the lungs. On entering the aorta. it passed down that vessel, filling it completely without any admixture of red, so that all the branches of the thoracic and abdominal aorta were filled with the yellow. From this and other experiments of a similar kind, Dr. Reid infers, that the blood, returning from the placenta, passes principally to the head and upper extremities; and that the lower part of the body is principally supplied by blood returning by the vena cava superior; or, in other words, by blood which has already gone the circuit of the body.

Recent observations by Dr. T. Williams have convinced him, notwithstanding the opposing experiments of Dr. Reid, that the Eustachian valve is mechanically inefficient as a means for preserving the individuality of the two currents from the vena cava as they traverse the right auricle; and he affirms, that at the period of its diastole, when the auricle has attained a moderate limit of

Edinb. Med. and Surg. Journal. xliii. 308.
 Lond. Med. Gazette, March 31, 1843.

distension, it may be readily demonstrated, that the two streams must freely intermingle. Hence it is not true — he infers — that the difference of quality is so great, as is generally taught by anatomists, between the blood that is distributed to the two portions

of the body in the fœtus.

After all, the great difference between the adult and fætal circulation is,—that in the former, a part of the blood only proceeds to the organ of sanguification; that the aërated blood is poured into the right auricle instead of the left; that, instead of proceeding through the lungs, a part of the blood gets at once to the left side of the heart, whence it is sent to the head and upper extremities, and the remainder goes directly from the pulmonary artery into the aorta; and that a part of the aortic blood proceeds to the lower extremities, and the remainder goes to the placenta, from which it is returned into the inferior cava.

4. With regard to the nutrition, (properly so called,) of the fœtus, it is doubtless effected in the same manner as in the adult; and our ignorance of the precise nature of the mysterious process is equally great. During the whole of fœtal existence, it is energetically exerted, and especially during the earlier periods. Sömmering has asserted, that the growth of the fœtus fluctuates; that in the first month it is greatest; in the second, less; in the third, greater; less, again, in the fourth; and then greater until the sixth, when it diminishes until birth.

There is, by the way, a singular circumstance connected with the nutrition of the fœtus, which cannot be passed over without a slight notice, although, in its details, it belongs more properly to pathological anatomy. Owing to inappreciable causes, the different parts of the fœtus, or some particular part, may be preternaturally developed, or be defective, so as to give rise to anomalies of conformation, or what have been termed monstrosities. Three kinds of monsters may be considered to exist. The first comprises such as are born with an excess of parts, as with two heads on one trunk, two trunks to one head; with four arms and four legs; twins with a band uniting them, as in the case of the Siamese twins, &c. The second includes those in which parts are defective, as acephali, anencephali, &c.; and the third, those in which there is perversion of the formative process, so as to produce various modifications in the direction and situation of organs, - as where the heart is on the right side, the liver on the left, &c.; where, in other words, there is transposition of the viscera. In these cases there is - to use the language of the German pathologists - superabundant, defective, or perverted action of the force of formation - the Bildungstrieb - to which we have more than once alluded.

The hypotheses, that have been advanced to account for these formations, as well as for those in which the parts are irregularly developed, may be reduced to three; the others, that have been entertained, having no probability in their favour. First. They

have been attributed to the influence of the imagination of the mother over the fœtus in utero. Secondly. To accidental changes experienced by the fætus at some period of uterine existence; and Thirdly. To some original defect or fusion of the germs. The first of these causes has been a subject of keen controversy amongst physiologists, at all periods. We have seen, that the mother transmits to the fœtus the materials for its nutrition; and that, to a certain extent, the nutrition is influenced by the character of the materials transmitted; so that if these be not of good quality or in due quantity, the fœtus will be imperfectly nourished, and may even perish. Any violent mental emotion may thus destroy the child. by modifying the quantity or quality of the nutritive matter sent to it. Small-pox, measles, and other contagious diseases can also be unquestionably communicated to the fœtus in utero; so that the life of the fætus is indirectly but largely dependent upon the condition of the mother. But the maternal influence has been conceived to extend much beyond this; and it has been affirmed, that the excited imagination of the mother may occasion an alteration in the form of particular parts of the fœtus, so as to give rise to nævi, and to all kinds of mother's marks, as they have been termed. These may consist of spots resembling raspberries, grapes, &c.; or there may be deficient formation of particular parts, and some of the cases that have been adduced in favour of their having been induced by impressions, made upon the mother during pregnancy, are sufficiently striking. There are numerous difficulties, however, in the way of accepting the cause assigned. If a child be born with nævi of any kind, the recollection of the mother is racked to discover whether some event did not befall her during gestation to which the appearance may be referred, and it is not often difficult to discover some plausible means of explanation. Cases have occurred in which the mother, when a few months advanced in pregnancy, has been shocked by the sight of a person who had lost a hand, and the child has been born with the same defect. A young female, a few months gone with child, visited a brother in one of the hospitals of London, who was wounded in the side. His condition affected her extremely. Her child was born with a deep pit in the same part that was wounded in the brother. These are samples of the thousands of cases that have been recorded, or that have occurred to different individuals." Similar instances have even been related of the inferior animals. In the extracts from the minute book of the Linnean Society of London, an account is given, by Mr. George Milne, F.L.S., of the effect of the imagination of a female cat on her young. One afternoon, whilst Mr. Milne and his family were at tea, a young female cat, which had arrived at the middle of gestation, was lying on the hearth. A servant, by accident, trod very heavily on her tail; she screamed violently, and from the noise emitted, it was evident, that

<sup>&</sup>lt;sup>2</sup> J. B. Demangeon, Du Pouvoir de l'Imagination sur le Physique et le Moral de l'Homme, Paris, 1834.

a considerable degree of terror was mingled with the feeling from the injury. From so common a circumstance no extraordinary result was expected; but, at the full time, she dropped five kittens, one of which was perfect, but the other four had the tail remark-

ably distorted; and all distorted in the same manner.a

Are we to consider these and similar cases of malformation or monstrosities to be dependent upon the influence of the maternal imagination upon the fœtus in utero? Or are we to regard them as coincidences, the cause being inappreciable, but such as we find to give occasion to vicious organization, where no coincidence with excited imagination can be discovered? Under the head of generation we have combated the notion, that the mother's fancy can have any effect—as to sex or likeness—during a fecundating copulation. Let us see, then, what we have to admit in a case where a female is, we will suppose, four months advanced in pregnancy, when she is shocked at the appearance of one who has lost his arm, and the child is born with a like defect. It has been seen, that the communication between the mother and the fætus is of the most indirect character; that the circulation of the fœtus is totally distinct from that of the mother; and that she can only influence the fœtus through the nutritive material she furnishes — whatever be its character - and, consequently, that such influence must be exerted upon the whole of the fœtus, and not upon any particular part. Yet, in the supposititious case we have taken, the arm must have been already formed, and the influence of the mother's fancy must have been exclusively exerted upon its absorbents, so as to cause them to take up again that which had been already deposited! The case we have assumed is not environed with more difficulty than any of the kind. It is a fair specimen of the whole. Yet how impracticable for us to believe, that the effect can be in any way connected with the assigned cause, and how much more easy to presume, that the coincidence, in such cases, has been accidental. Cases of hare-lip are perpetually occurring, yet we never have the maternal imagination invoked; because, it is by no means easy to discover any similitude between the affection and common extraneous objects. Moreover, in animals of all kinds - even in the most inferior, as well as in plants - monstrous formations are incessantly happening where maternal imagination is out of the question. As a cause of monstrosities, therefore, the influence of the maternal imagination has been generally regarded as an inadmissible hypothesis. By many, it has been esteemed ridiculous; yet it manifestly receives favour with Sir Everard Home; b and Professor Elliotson, and Burdach appear inclined to favour it; but on the whole we are justified in adopting the opinion of Dr. Blundell,

<sup>\*</sup> Fleming's Philosophy of Zoology, vol. i, Edinb. 1822.

Philos. Transactions for 1825, and Lect. on Compar. Anat. v. 190, Lond. 1828.
 See his translation of Blumenbach's Physiology, 4th edit. p. 497, Lond. 1828.

<sup>&</sup>lt;sup>d</sup> Die Physiologie als Erfahrungswissenschaft, B. ii.; see, also, Pierquin, in Magendie's Journal de Physiologie, x. 364.

VOL. II. - 43

which has been embraced by Drs. Allen Thompsona and Wagner, that it is contrary to experience, reason and anatomy, to believe, that the strong attention of the mother's mind to a determinate object or event can cause a determinate or a specific impression upon the body of her child, without any force or violence from without; and that it is equally improbable, that, when the imagination is operating, the application of the mother's hand to any part of her own body, will cause a disfiguration or specific impression on a corresponding part of the body of the child! The third hypothesis, with regard to defective germs, we have already canvassed under Generation, and attempted to prove it insufficient. The second, consequently, alone remains, and is almost universally adopted. Independently of all disturbing influences from the mother, the fœtus is known to be frequently attacked with spontaneous diseases, such as dropsy, ulceration, gangrene, cutaneous eruptions, &c. Some of these affections occasionally destroy it before birth. At other times, it is born with them; and hence they are termed connate, or congenital.c

The following table, drawn up by Mr. Wilde, from the records of the Gebäranstalt, of Vienna, exhibits the malformations observed

there in 23,413 births.

Clubfoot -	La la	12000	A log by	THE WAY	16 ca	ses, o	ronce	in 1463-31
Hare lip -	HOUSE	(14)	11200	Man M	20	122601	102911	1170-65
simple	- 1700		111-211	TORS:	9	13437	TOUR OF	2601-44
with cleft pala	te	-			11		10-11	2128-45
Spina bifida -			-	-	5	-	-	4682-6
Hydrocephalus	TIGHT	-	monn	-	6	-	-	3902-16
Six fingers -	- 1.00	A 5) III	TOLON!	TORRES	3	12 7	111	7804-33
Inperforate anus			16-303	-	2	11-01	1201	11706-5
Hemicephalus					1	1	-	23413
Acephalus -	-	-	-		1	-		23413
Umbilical hernia	TO THE		THE PARTY	19:31	1	10000	HEAL	23413
Without eyes -	a with	1.50	1 123/17	10(2011)	2	Digit!	0000	11706.5
Wanting superior part	of vert	tex	law in	1111-1111	1	10000	1943176	23415
Lenticular cataract	-				1	-	-	23413
Wanting one upper ex	tremity	7		-	2	-	-	11706-5
With plurality of finge	rs and	toes	Halion 1	The last	5	-	THE REAL PROPERTY.	4682-6
Hydrocephalus with sp	ina bif	ida, an	d closed	anus	1	1000	Digital Co	23413
Clubfoot and closed an	us	-	DELI-SU	TIL-)I	1	1900h	( - DIS	23413

In a population, consequently, chiefly illegitimate, 88 deviations from the natural type occurred in 23,413 births, or about 1 in every 266 cases.

According to the last census of the United States, (1840,) the proportion of deaf and dumb, amongst the whites, was 1 in 2123; amongst the coloured, 1 in 2933.

Where a part has been wanting, the nerve, or bloodvessel, or

d Austria, &c., p. 224, Dublin, 1843.

a Art. Generation, Cyclop. of Anat. and Physiol. P. xiii. p. 477, February, 1838.
b Elements of Physiology, translated by Dr. Willis, p. 227, note, Lond. 1841.

c J. Grätzer, Die Krankheiten des Fætus, Berlin, 1837, and Dr. W. C. Roberts, Amer. Journ. of the Med Sciences, Aug. 1840, and Oct. 1841.

Prof. G. Tucker, Progress of the United States in Population and Wealth, in Fifty Years, p. 77, NewYork, 1843.

both, proceeding to it, have likewise been found wanting; so that the defect of the organ has been thus explained; without our being able, however, to understand the cause of the deficiency of such nerve or bloodvessel.

In some of the cases of montrosities, a fusion of two germs seems to have occurred. Two vesicles have been fecundated and subsequently commingled, so that children have been produced with two heads and one trunk, or with two trunks and one head, &c., &c. This view appears certainly more probable than that the whole class of monsters by excess - including those commonly so called, and those usually considered as double monsters - owe their origins to different degrees of one common fault, and, consequently, the explanation of their origin ought to be the same for all: "that no kind of fusion can account for the production of supernumerary individual organs, the rest of the body being single; but that it is not impossible, that excess of power in one ovum, which all admit can alone explain the lower degrees of duplicity, may, in proportionally lower degree, produce the more completely double monsters, or even two such separate individuals as are sometimes found with a single amnion." The case of the supernumerary individual organs appears to the author, however, to be easily separable, and to admit of a distinct explanation from that of duplicity, or, as he regards it, fusion of ova. At times, where two ova appear to have been fecundated, the development of the one has been arrested, whilst the other has gone on. Two instances of this kind we have alluded to under Superfætation, and under the theory of generation by evolution.

This interesting department of pathological anatomy has become, of late years, of moment as elucidating the formation of the animal body, which appears to be effected—even in the production of these anomalies—according to a unity of organic composition, and to laws of development but little appreciated until the present century. The labours of Geoffroy and Isidore Saint-Hilaire, Serres, Sömmering, Meckel, Tiedemann, Béclard, Breschet and others, have, indeed, as Cuvier remarked respecting some of them—occasioned the accumulation of an infinity of facts and views, which, even if we do not admit all that their authors contend for,

cannot fail to be of solid advantage to science.b

5. The animal temperature of the fotus cannot be rigidly determined. The common belief is, that it is some degrees lower

\* Brit. and Foreign Medical Review, Oct. 1841, p. 388.

See, especially, Geoffroy St. Hilaire, Philosophie Anatomique, Paris, 1818, and 1822; Isidore St. Hilaire, Histoire Générale et Particulière des Anomalies de l'Organisation, Paris, 1832-36; Serres, Recherches d'Anatomie Transcendante, Paris, 1832; and Précis d'Anatomie Transcendante, appliquée à la Physiologie, &c. Paris, 1842. See, also, an epitome of the views of modern Teratologists, in Brit, and For. Med. Rev. July, 1839, p. 1; Mr. North, Lond. Lancet, March 7, 1840, or Dunglison's Amer. Med. Intelligencer, Aug. 1 and 15, 1840; and an analysis of Professor Vrolik's work on Double Monsters, Amsterd. 1840, in Brit. and For. Med. Rev., Oct. 1841, p. 374.

than that of the mother; and it is affirmed, that the temperature of the dead fœtus is higher than that of the living. The fœtus must, therefore, possess the means of refrigeration. Edwards found, in his experiments, that the temperature of young animals is inferior to that of the adult; which is in accordance with the general belief regarding that of the fœtus in utero. In some cases, as in those of the kitten, puppy and rabbit, if the young be removed from the mother and exposed to a temperature of between 50° and 70°, their temperature will sink, - as happens to the cold-blooded animal, - to nearly the same degree. The faculty of producing heat he found to be at its minimum at birth; but it progressively increased, until in about fifteen days the animal acquired the power in the same degree with the adult. This was not the case, however, with all the mammalia. It seemed to be confined to those animals that are born blind; in which a state of imperfection probably exists in other functions. It was the same with birds as with the mammalia: birds, hatched in a defective state, as regards their organs generally, have the power of producing heat defective; whilst others, born in a more perfect condition, have the organs of calorification more capable of exercising their due functions. In a case in which the mother, who had borne five children, was confident that her period of gestation was less than 19 weeks; but probably from the length and weight of the fœtus it was 25 weeks, the power of calorification of the infant was so low, that artificial heat was constantly needed to sustain it: under the influence of the heat of the fire, however, it evidently became weaker, whilst the genial warmth of a person in bed rendered it lively and comparatively strong.2

Opinions with regard to the temperature of the human infant vary. Haller asserts that it has less power of producing heat than the adult, and such is the opinion of Despretz, Edwards, and the generality of physiologists. The latter gentleman estimated it at 94.25° of Fahrenheit. On the other hand, Dr. John Davy affirms, that the temperature of young animals generally, and that of a new-born child, which he particularly examined, was higher than in the adult. It is impossible to account for this discordance; but the general results of experiments will be found to agree with those of Edwards. The fœtus certainly possesses the power of forming or separating its own caloric; otherwise its temperature should correspond with that of the mother, which we have elsewhere seen

is not the fact.

6. That the secretions are actively exercised in the fœtus is proved by the circumstance, that all the surfaces are lubricated nearly as they are subsequently. The follicular secretion is abundant, and at times envelopes the body with a layer of sebaceous matter—the vernix caseosa—of considerable thickness. Vau-

4 Philos. Transact. p. 602, for 1814.

<sup>\*</sup> Edinb. Medical and Surgical Journal, vol. xi.

De l'Influence des Agens Physiques, Paris, 1824.

quelin and Buniva have asserted, that this is a deposit from the albumen of the liquor amnii; but, in reply to this, it may be urged, that we do not find it except on the body of the fætus. It is not on the placenta or umbilical cord, and is most abundant on those parts of the fætus, where the follicles are most numerous. The fat also exists in quantity after the fifth month. The greatest question has been with regard to the presence, in the fætal state, of some of the secretions which are of an excrementitious character. For example, — by some, the urinary secretion is supposed to be in activity from the earliest period of intra-uterine existence, and its product to be discharged into the liquor amnii. Such is the opinion of Meckel. The circumstances that favour it, are—the fact of the existence of the kidneys at a very early period; and that at the full time the bladder contains urine, which is evacuated soon after birth. On analysis, this is found to be less charged with

urea and phosphoric salts than in after life. Of the meconium we have already spoken. It is manifestly an excretory substance, produced, probably, by the digestion of the fluids of the alimentary canal, mixed with bile. Some, indeed, are of opinion, that it is altogether a secretion from the liver, and intended to purify the blood sent from the mother, so as to adapt it for the circulation of the fœtus. Into the value of the theory on which this notion rests, we have inquired at some length. The notion itself scarcely requires farther notice. We may remark, however, that it has been conceived by Dr. Lee, that the fætal liver secretes an abundant albuminous and nutritive matter, which fills the hepatic ducts, the duodenum and small intestines; whilst in the stomach we find only an acid fluid, and in the large intestines meconium. The meconium, evacuated after birth, has been found composed of water, about two-thirds; of a substance of a vegetable nature, but sui generis, about one-third; and mucus, a few hundredth parts.º It appears, consequently, to be less azoted than the

Lastly, the cutaneous respiration is supposed to be a fætal excretion, and to be poured into the liquor amnii; but although this is probable, we have no positive evidence on the subject.

c. Functions of Reproduction. — These require no consideration. They are inactive during the fætal state, except that the testicles and the mammæ appear respectively to secrete a fluid which is neither sperm nor milk, and is found in the vesiculæ seminales in the one case, and in the lactiferous ducts in the other. It would appear also that the ovarian vesicles are already existent at this early period.<sup>d</sup>

Annales de Chimie, tom. xxxiii.; and Mémoir. de la Société Medicale d'Emula-

<sup>5</sup> Handbuch, u. s. w.; or French translation by Jourdan and Breschet, iii. 780; or the English translation by Dr. S. A. Doane. Philad. 1832; and Dr. Robert Lee, in Medico-Chirurgical Transactions, vol. xix., Lond. 1835.

Bouillon-Lagrange, in Annales de Chimie, lxxxvi. and lxxxvii.

d Carus, Gazette Médicale de Paris, Aug. 12, 1837.

excrement of the adult.

510

# BOOK IV.

AGES.

# CHAPTER I.

AGES.

UNDER this head we have to include the modifications that occur in the functions during the life of man, from birth until dissolution. The different ages may be separated as follows: - infancy, comprising the period from birth till the second dentition; — childhood, that between the second dentition and puberty; - adolescence, that between puberty and manhood; - virility, that between youth and old age; - and old age.

#### 1. INFANCY.

The age of infancy extends from birth to the second dentition. or until about the seventh or eighth year. By Hallé, and after him by Renauldin, Rullier, Adelon, and others, this has been again subdivided into three periods, which are somewhat distinct from each other, and may therefore be adopted with advantage. The one comprises the period between birth and the first dentition, - generally about seven months; a second embraces the whole period of the first dentition, or up to about two years, and the third includes the whole interval, that separates the first from the second dentition.

a. First period of Infancy. — As soon as the child is ushered into the world, it assumes an independent existence, and a series of changes occurs in its functions, of the most sudden and surprising character. Respiration becomes established, after the manner in which it is to be effected during the remainder of existence; and the whole of the peculiarities of fætal circulation cease,the organs of these peculiarities being modified in the manner to be described presently. The first act after the child is extruded is to breathe, and at the same time to cry. What are the agencies, then, by which this first inspiration is effected, and this disagreeable impression made upon the new being at the moment when it makes its appearance amongst us? This has been an interesting topic of inquiry amongst physiologists. A few of the hypotheses, that have been indulged, will be sufficient to exhibit the direction which the investigation has taken.

Whytt,d - whose views were long popular, and still have supporters, - conceived, that, before birth, the blood of the fœtus is properly prepared by the mother; and that when, after birth, it

· Physiologie de l'Homme, 2de édit. iv. 425, Paris, 1829.

Art. Age, Dictionnaire des Sciences Médicales. b Art. Age, Dict. de Médecine, i. 381, Paris, 1821.

<sup>4</sup> An Essay on the Vital and other Involuntary Motions of Animals, Sect. ix. 109; Edinb. 1751.

no longer receives the necessary supply, an uneasy sensation is experienced in the chest, which may be looked upon as the appetite for breathing, in the same manner as hunger and thirst are appetites for meat and drink. To satisfy this appetite, the brain excites the expansion of the chest, to prevent the fatal effects that would ensue if the lungs were not immediately aroused to action. This appetite is supposed to commence at birth, owing to the circulation being quickened by the struggles of the fœtus at that period, and to an additional quantity of blood passing through the lungs, which excites them to action, and seems to be the immediate cause of the appetite.a Hallerb ascribes the first inspiration to the habit which the fœtus has acquired, whilst in the uterus, of taking into the mouth a portion of the liquor amnii; and he supposes, that it still continues to open its mouth, after leaving the mother, in search of its accustomed food. The air, consequently, rushes into the lungs, and expands them; the blood is distributed through them, and a regular supply of fresh air is needed to prevent the blood from stagnating in its passage from the right to the left side of the heart. Dr. Wilson Philip, regards the muscles of inspiration as entirely under the control of the will; and he thinks, that they are thrown into action by the uneasy sensation experienced by the new being, when separated from the mother, and having no longer the necessary changes produced upon the blood by her organs. Adelon thinks it probable, that the series of developments, occurring during gestation, predisposes to the establishment of respiration. According to him, the lungs gradually increase in size during the latter months; the branches of the pulmonary vessels become enlarged, and the ductus arteriosus less; so that the lungs are prepared for the new function they have to execute. In addition to these alterations, he conceives, that the process of accouchement predisposes to the change; that, by the contractions of the uterus, the circulation of the blood must be necessarily modified in the placenta, and, consequently, in the fœtus; -- for he is a believer in the doctrine, that the fœtus receives blood from the mother by the placenta. Owing to this disturbance in the circulation, more blood is sent to the lungs; and, when the child is born, it is subjected to new and probably painful impressions. "For instance," he remarks, "the external air, by its coldness and weight, must cause a disagreeable impression on the skin of the infant, as well as on the origin of all the mucous membranes; and, perhaps, the organs of the senses being, at the same time, suddenly subjected to the contact of their proper irritants, receive painful impressions from them. These different impressions being transmitted to the brain, they are reflected to the different dependencies of the nervous system, and, consequently, into the nerves of the inspiratory muscles: these muscles, thus

b Elem. Physiol. viii. 5. 2.

<sup>&</sup>lt;sup>a</sup> See Dr. Marshall Hall, in Lect. on the Nervous System, p. 55, Lond. 1836.

e Quarterly Journal of Science, &c., vol. xiv. 100.

512 AGES

excited, enter into contraction, in the same manner as the heart is stimulated to renew its contractions, during syncope, when we

inspire a stimulating vapour."

None of these views satisfactorily explain the true physiology of the first inspiration; nor is it probable that any can be devised which has not its difficulties. That which has been embraced by Dr. Bostocka appears to us to be liable to fewer objections than any we have seen; and to explain the process, as far perhaps as is practicable, on mechanical principles. The first respiratory act, according to him, seems to be purely mechanical, and to result from the change of position which the child undergoes at birth. From the mode in which it rests in utero, everything is done that position could accomplish, to diminish the dimensions of the chest; and any change in this position must have the effect of liberating the lungs from a portion of the pressure which they sustain. The head cannot be raised from the breast, nor the knees removed from the abdomen, without straightening the spine; and the spine cannot be reduced to a straight line without elevating the ribs, and permitting the abdominal viscera to fall; but the ribs cannot rise nor the diaphragm descend, without enlarging the chest; and, as the chest enlarges, the lungs, which are the most elastic organs of the body, expand their air-cells, hitherto collapsed by external pressure, and the external air rushes in. The same cause is considered to account for the new circulatory movement. The blood, which, in the fœtus, had passed through the foramen ovale and the ductus arteriosus without visiting the lungs, is solicited from its course by the expansion of the chest, which draws the blood through the pulmonary artery as forcibly as it does air through the windpipe. The blood, thus exposed to the air in the lungs, becomes arterialized, and, from this moment, the distinction between arterial and venous blood is established. The circulation, through the vessels peculiar to the fætal condition, now ceases, even without any ligature being placed upon the umbilical cord.

The sudden and important changes supervening in this manner guide us to the decision of an interesting medico-legal inquiry,—viz. whether, in a case of alleged infanticide, the child has respired or not;—in other words, whether it has been born alive or dead? After respiration has been established, the lungs, from being dark-coloured and dense, become of a florid red hue; are light and spongy, and float on water: on cutting into them, the escape of the air from the air-cells occasions a crepitus, and a bloody fluid exudes; there is an approach to closure of the foramen ovale; the ductus arteriosus is empty, as well as the ductus venosus; and the absolute weight of the lungs may be doubled. (See page 477

of this volume.)

The different conditions of the organs of circulation, the cord and the skin, at different periods, beginning at the first and ending

<sup>&</sup>lt;sup>a</sup> Elementary System of Physiology, 3d edit. p. 323, Lond. 1836; Dr. Southwood Smith, in Animal Physiology, p. 101, Library of Useful Knowledge.

with the thirty-fifth day, have been summed up as follows by M. Devergie.a They may enable us to judge approximately of the age of the young infant, in questions of a medico-legal character. On the first day. Cord beginning to wither. Foramen ovale, ductus arteriosus, ductus venosus, and umbilical vessels open. Second day. Withering of the cord complete. Foramen ovale closed in two out of eleven cases; partially closed in one out of seven. Ductus arteriosus beginning to close. Umbilical arteries obliterated to a greater or less extent. Umbilical veins and ductus venosus still open. Third day. Desiccation of the cord. Foramen ovale sometimes closed. Ductus arteriosus obliterated in one in eleven cases. Umbilical arteries very often obliterated. Umbilical vein and ductus venosus still open. Fourth day. Cord beginning to fall off. Foramen ovale closed in about one-third of the cases. Ductus arteriosus still open in the majority of cases. Umbilical arteries closed, but sometimes open near the iliacs. Umbilical vein and ductus venosus much contracted. Fifth day. Separation of the cord with rare exceptions. Foramen ovale closed in more than half the cases. Ductus arteriosus closed in about half the cases. Umbilical vessels closed; vein occasionally open. Separation of the cuticle advanced. Eighth day. Entire separation of the cord, with commencing cicatrization. Foramen ovale closed in threefourths of the cases. Ductus arteriosus completely obliterated in half the cases. Umbilical vessels closed. Ninth to eleventh day. Cicatrisation of the umbilicus often complete; sometimes, however, there is an oozing of mucus from the cord for many days, so that the cicatrix is retarded. Separation of the cuticle on the trunk, chest, and abdomen, and at the articulations. Twentieth to twentysixth day. Separation of the greater part of the cuticle. Thirtieth to thirty-fifth day. Separation of the entire cuticle, excepting that of the hands and feet, which is often delayed until the fortieth day.b

Respiration having been once thoroughly established, the individual enters upon the period of infancy, which has now to engage

our attention.

The animal functions, during this period, undergo considerable development. The sense of tact is little evinced, but it exists, as the child appears sensible to external cold. At first, the touch is not exerted under the influence of volition, but, towards the termination of the period, it begins to be active. The taste is almost always null at first. Adelone thinks, that it is probably exerted on the first day as regards the fluids, which the infant sucks and drinks. We have daily evidence, however, that at an early period of existence, the most nauseous substances, provided they are not irritating, will be swallowed indiscriminately, and without the slightest repugnance; but, before the termination of the period

\* Médecine Légale, art. Infanticide.

Physiologie de l'Homme, edit. cit. iv. 433.

b Guy, Principles of Forensic Medicine, Part. i. p. 149, Lond. 1843.

we are considering, the taste becomes inconveniently acute, so that the exhibition of nauseous substances, as of medicine, is a matter of more difficulty. The smell is probably more backward than any of the other senses; the development of its organ being more tardy, the nose being small, and the nasal sinuses not in existence. In the first few weeks, sight and hearing are imperfectly exerted, but, subsequently, they are in full activity. The internal sensations, being instinctive, exist; all those at least that are connected with the animal and nutritive functions. Hunger and thirst appear during the first day of existence; the desire of passing the urine and fæces is doubtless present, notwithstanding they appear to be discharged involuntarily; and the morbid sensation of pain is often experienced, especially in the intestinal canal, owing to flatus, acidity, &c. During the first part of the period, the child exhibits no mental and moral manifestations; but, in the course of a few weeks, it begins to notice surrounding objects, especially such as are brilliant, and to distinguish between the faces to which it has been accustomed and those of strangers; awarding the smile of recognition or of satisfaction to the former, the look of gravity and doubt to the latter. Locomotion is, at this time, utterly impracticable, as well as the erect attitude. The muscular system of the child is not yet sufficiently developed; the spinous processes of the vertebræ are not formed, and it has not learned to keep the centre of gravity - or rather the vertical line - within the base of sustentation. The function of expression is at the early part of the period confined to the vagitus or squalling, which indicates the existence of uneasiness of some kind; but, before the termination of the period, the infant unites smiles and even laughter to the opposite expressions, and attempts to utter sounds, which cannot yet be considered as any attempt at conventional language. Sleep is largely indulged. Soon after birth, it is almost constant, except when the child is taking nutriment. Gradually, the waking intervals are lengthened; but, throughout, much sleep is needed, owing to the frail condition of the nervous system, which is soon exhausted by exertion however feeble, and requires intermission of action.

After birth, the child has to subsist upon a different aliment from that with which it was supplied whilst in the maternal womb. Its digestion, therefore, undergoes modification. The nutriment is now the milk of the parent, or some analogous liquid, which is sucked in, in the manner described under the head of Digestion. For this kind of prehension, the mouth of the infant is well adapted. The tongue is very large, compared with the size of the body, and the want of teeth enables the lips to be extended forward, and to embrace the nipple more accurately and conveniently. The action of sucking is doubtless as instinctive as the appetite for nutriment, and equally incapable of explanation. The appetite appears to be almost incessant, partly owing to the rapidity of growth demanding continual supplies of nutriment, and partly, perhaps,

owing to a feeling of pleasure experienced in the act, which is generally the preclude to a recurrence of sleep, broken in upon, apparently, for the mere purpose of supplying the wants of the system, or the artificial desire produced by frequent indulgence. Often, we have the strongest reason for believing, that the great frequency of the calls of the appetite is occasioned by the habit, with many mothers, of putting the child constantly to the breast: whilst in those children that have been trained, from the earliest period of existence, to receive the nutriment at fixed hours only, the desire may not recur until the lapse of the accustomed interval. Digestion is, at this age, speedily accomplished; the evacuations being frequent, - two or three or more in the course of the day. -of a yellow colour, something like custard, or curdy, and having by no means the offensive smell, which they subsequently possess. During the first days after birth, they are dark and adhesive, and consist of the meconium, already described. Young mothers are apt to be alarmed at this appearance which is entirely healthy, and always exists. The respiration of the infant is more frequent than in the adult, nearly in the proportion of two to one, and it is chiefly accomplished by the muscles that raise the ribs, on account of the great size of some of the abdominal viscera, which do not permit the diaphragm to be readily depressed. The stethoscope exhibits the respiration to be much more sonorous; so characteristic, indeed, is it, in this respect, that it has been called "puerile," by way of distinction. It appears to indicate a greater degree of activity in the respiratory function. The circulation is more rapid; the pulsations at birth being nearly twice as numerous as in the adult. Nutrition is very active in the development of the different organs. Calorification becomes gradually more energetic from the time of birth. The recrementitial secretions, as well as the excrementitial, are as regularly formed as in the adult; but the products vary somewhat. The urine, for instance, is less charged with urea, and contains benzoic acid; the perspiration is acidulous, &c., &c. Adelona asserts, that these excretions are frequently insufficient for the necessary depuration, and that nature, therefore, establishes others that are irregular and morbid, in the shape of cutaneous efflorescences, &c. These can scarcely be regarded as depurations; unless we consider all cutaneous eruptions. that are connected with gastric or digestive irritation, to be thus induced, which is more than problematical; especially as most of them are neither pustular nor vesicular, and therefore, not accompanied by any sensible exudation.

b. Second period of Infancy or first Dentition. — This period embraces the whole time of dentition, and is considered to include the age between seven months and two years. In it, the external senses are in great activity, and continually furnishing to the intellect the means from without for its development. The internal sensations are likewise active. From these united causes, as well

Physiologie de l'Homme, 2de édit. iv. 436, Paris, 1829.

as from the improved cerebral organization, the intellect is more strengthened during this period than perhaps during any other. The senses are continually conveying information; perception is, therefore, most active, - as well as memory; whilst imagination and judgment are feeble and circumscribed. The faculty of imitation is strong, so that, by hearing the spoken language, and appreciating its utility, the child endeavours to produce similar sounds with its own larynx, and gradually succeeds, - the greater part of its first language consisting of imitations of sounds emitted by objects, which sounds are applied to designate the object itself, in the manner we have seen elsewhere. The affective faculties are likewise unfolded during this period, but generally those of the selfish cast are predominant, and require the most careful attention for their rectification. Even at this early time of life, the effect of a well-adapted education is striking, and spares the child from numerous inconveniences, to which unlicensed indulgence in its natural passions would inevitably expose it. The general feeling is, that the infant is not yet possessed of the necessary intelligence to pursue the course that is indicated; but it is surprising how soon it may be made to understand the wishes of its instructor, and with what facility it may be moulded, at this tender age, in almost any manner that may be desired. During this period, the child is capable of standing erect and of walking. Previous to this, these actions were impracticable, for the reasons already stated, as well as owing to the weight of the thoracic and abdominal viscera, - to the spine having but one curvature, the convexity of which is backwards, - to the smallness of the pelvis, and its inclination forwards, so that it scarcely supports the weight of the abdominal viscera, and to the smallness of the lower limbs and to the feebleness of their muscles, which are insufficient to prevent the trunk from falling forward. These imperfections are, however, gradually obviated, and the child commences to support itself on all-fours; a position assumed much more easily than the biped attitude, owing to the centre of gravity being situate low, and the base of sustentation being large. In this attitude, he moves about for some time, or his locomotion is effected by pushing a chair before him, or by being steadied by his nurse. Gradually, he passes from place to place on his feet, by laying hold of surrounding objects, and, in proportion as the bones and muscles become developed, and the obstacles to progression are removed, he succeeds in walking alone; but it is some time before he is capable of running or leaping. Perhaps, the average period, at which the infant begins to walk, is about twelve months; but we see great difference in this respect. When once the infant is fairly on his legs, the whole of his waking hours is spent in incessant activity and amusement. His functions of expression are commensurate with his intellectual development, which we have seen is great in this period. Sleep, which is now more interrupted, is still imperiously and frequently demanded, the nervous system being devoid of that strength, which it subsequently

possesses, and therefore requiring repose.

One of the most important changes going on at this age concerns the function of digestion. This is the process of dentition, which usually commences about the seventh month, and continues till the end of the second year at least. Prior to the appearance of the teeth, mastication is of course impracticable; and the food, best adapted for the delicate powers of the infant, is that afforded by the maternal breast, or a substitute which resembles it as closely as possible. The appearance, however, of teeth would seem to indicate, that the infant is about to be adapted for more solid aliment. As early as the second month of utero-gestation, if the jaws be carefully examined, the germs of the teeth are perceptible in their substance, under the form of membranous follicles of an oval shape, attached by their deep-seated extremity to a vascular and nervous pedicle, and by their superficial extremity to the gum. The cavity of these follicles is at first filled with a colourless limpid fluid; but a kind of vascular and nervous papilla or pulp soon forms in it, which commences at the deep-seated portion of the follicle, proceeds towards the other extremity, and ultimately fills it, —the fluid diminishing in proportion to the increase of the pulp. About the termination of the third month, ossification begins, and a little sooner in the lower than in the upper jaw. This consists, at first, in a deposition of ivory matter on the surface of the pulp and at its top, which goes on increasing in width until it covers the whole of the dental pulp with a shell of bone. It augments, also, in thickness at the expense of the dental pulp, which becomes gradually less and less. When the bony shell has extended as far as the neck of the tooth, the external membrane or sac of the tooth — for the follicle consists of two membranes - attaches itself closely, but not by adhesion, to the part. The inner membrane becomes much more vascular, and the enamel is secreted by it. A thickish fluid is observed to be poured out from the inner surface, which is soon consolidated into a dark, chalky substance, and afterwards becomes white and hard.


At birth, the coronæ of the incisors are formed; those of the canine are not completed; and the molares have only their tubercles. The root or fang is formed last of all. As ossification proceeds, the corona of the tooth presses upon the gum, a portion of the follicle being interposed, which is gradually absorbed as

well as the gum; and the tooth issues.

The age at which the teeth make their appearance varies. Occasionally, children have been born with them, whilst in other cases they have not pierced the gum until after the period we are considering. Generally, the middle incisors of the lower jaw appear about the seventh month, and subsequently those of the upper jaw; next the inferior and superior lateral incisors in succession; then the first lower molares, and the first upper; next the inferior


and superior canine teeth, successively; and, lastly, the second





1. Front view of the temporary Teeth.

Fig. 279.



2. The separate temporary Teeth of each Jaw.

a. Central incisor. b. Lateral incisor. c. Canine. d. First molaris. ed of late years c. Second molaris.

molares of each jaw. The approximate times of their appearance are thus estimated by Mr. Thomas Bell.a

From five to eight months, the four central incisors.

From seven to ten, the four lateral incisors.

From twelve to sixteen, the four anterior molares.

From fourteen to twenty, the four canine.

From eighteen to thirtysix, the four posterior molares.b

The subject of the intimate anatomy development of the teeth has been investigat-

by many observers, whose contributions are well worthy of perusal. Amongst the most important are those of Fränkel, Raschkow, Retzius, Arnold, Goodsir, Owen, and Nasmyth; and a recent writer has

The Anatomy, Physiology, and Diseases of the Teeth, Lond. 1829. See, also, Dr. Ashburner, Lectures on Dentition, Med. Gaz. 1833-34.

b See M. Trousseau, Gaz. Méd. de Paris, 4 Fév. 1842; and Amer. Journ. of the Med. Sciences, July, 1842, p. 165.

c Edinb. Med. and Surg. Journ. li. See, also, for an account of his views, Car-

penter, Human Physiology, § 632, Lond. 1842.

4 J. Müller, Elements of Physiology, by Dr. Baly, p. 391, and part iv. chap. 2, appendix; and Nasmyth, Researches upon the Development and Structure of the Teeth, London, 1839 and 1841; Mr. Robert Nasmyth, in Lond. and Edinb. Monthly Journal of Med. Science, Jan. 1843, p. 40; and R. Owen, Odontography, part i. Lond. 1840. See, also, Brit. and For. Med. Rev., July, 1839, p. 158, and Oct. 1841, p. 491.

remarked, that in no organs have the results of recent microscopic researches been so unexpected or so brilliant.a These researches have shown the teeth to be composed of three main constituents. 1. The crusta petrosa, which differs in its minute structure in no respect from common osseous tissue. It forms the outermost layer of the teeth, visibly surrounds the whole fang, and extends, according to Mr. Nasmyth, in a very thin layer over the enamel of the crown. 2. The enamel, which invests only the crown of the teeth, and is composed of solid prisms or fibres about 1 th of an inch in thickness set side by side upright on the ivory; and 3. The dentine or ivory, which forms the chief mass and body of the teeth, which is composed of a fibrous basis, traversed by very fine, branching, cylindrical tubuli, which run in an undulating course from the pulp cavity on the interior of which they open towards the adjacent part of the exterior of the tooth. The basis of the intertubular substance, according to Henle, is composed of bundles of flat, pale, granular fibres, the course of which is parallel to that of the tubules.

Dentition is necessarily a physiological process, but it is apt to be a cause of numerous diseases. The whole period of its continuance is one of great nervous susceptibility, - so that the surgeon never operates during it, unless compelled, - and we can understand, that the pressure, exerted by the tooth on the gum, and the consequent inflammation and irritation, may lay the foundation for numerous diseases. More are doubtless ascribed to the process than it is entitled to, but still they are sufficiently numerous; and all require in their management the free division of the distended gum, so as to set the presenting part of the tooth at liberty. Whilst the teeth are appearing, the muscular structure of the body generally is acquiring strength, and the salivary organs are described by anatomists as becoming much more developed. The food of the child is now diversified, and it begins to participate in the ordinary diet of the table. The excrementitious matters are consequently altered in their character, particularly the alvine, which become firmer, and acquire the ordinary fæcal smell; the urea is still, however, in the generality of cases, in less proportion than in the adult. The other functions require no particular mention.

The number of deaths, during this period, is great. The bills of mortality of London, as has been elsewhere remarked, show, that the deaths, under two years of age, are nearly thirty per cent. of the whole number. In Philadelphia, during a period of twenty years ending with 1826, the proportion was rather less than a third. The cholera of infants is the great scourge of our cities during the summer months, whilst in country situations it is comparatively rare; and it is always found to prevail most in crowded alleys, and in the filthiest and impurest habitations. There is

Mr. Paget, Brit, and For. Med. Rev., July, 1842, p. 270.

something in the confined and deteriorated atmosphere of a town, which seems to act in a manner directly unfavourable to human life, and to the life of the young especially. This is not confined to man. It is applicable also to the animal. Experiments were instituted by Jenner, and since him by Dr. Baron, which show that a privation of free air and of their natural nourishment has a tendency to produce disorganization and death. Dr. Baron placed a family of young rabbits in a confined situation, and fed them with coarse green food, such as cabbage and grass. They were perfectly healthy when put up. In about a month, one of them died, - the primary step of disorganization being evinced by a number of transparent vesicles on the external surface of the liver. In another, which died nine days after, the disease had advanced to the formation of tubercles in the liver. The liver of a third, which died four days later, had nearly lost its true structure, so completely was it pervaded by tubercles. Two days afterwards, a fourth died; a number of hydatids was attached to the lower surface of the liver. At this time, Dr. Baron removed three young rabbits, from the place where their companions had died, to another situation, dry and clean, and to their proper accustomed food. The lives of these were obviously saved by the change. He obtained similar results from experiments of the same nature performed on other animals.b

c. Third period of Infancy. — This requires no distinct consideration; — the growth of the child and activity of the functions going on as in the preceding period, but gradually acquiring more and more energy. Within this period, a third molar tooth appears, which is not, however, temporary, but belongs to the per-

manent set.

During the whole of infancy, the dermoid texture — both skin and mucous membranes — is extremely liable to be morbidly affected; hence, the frequency of eruptive diseases, and of diarrhæa, aphthæ, croup, bronchitis, &c., many of which are of very fatal tendency. Owing, also, to the susceptibility of the nervous system, convulsions, hydrocephalus, and other head affections are by no means infrequent.

#### 2. CHILDHOOD.

Childhood may be considered to extend from the seventh to the fifteenth year, or to the period of puberty; and it is particularly marked by the *shedding* of the first set of teeth, and the appearance of the second. It is manifest, that in the growth of the jaws with the rest of the body, the teeth, which, for a time, may have been sufficient in magnitude and number, must soon cease to be

b See the author's Elements of Hygiène, p. 138, Philad. 1835; and art. Longevity, in American Quarterly Review, viii. 380, Philad. 1830.

Delineations of the Origin and Progress of various changes of Structure which occur in Man, and some of the inferior Animals, Lond. 1828.

so; hence, the necessity of a fresh set, which may remain permanently. The process for the formation of the permanent teeth is similar to that of the milk or temporary teeth; yet it presents some remarkable points of difference; and it affords us another surprising instance of the wonderful adaptation of means to definite objects, of which we have so many in the human body.


This process is well described by Mr. Bell, a whose opportunities for observation have been unusually numerous, and whose zeal and ability in his profession, as well as in the prosecution of natu-

ral science, are well known.

The rudiments of the permanent teeth are not original, and independent, like those of the temporary. They are derived from the latter, and continue, for a considerable time, attached to, and intimately connected with them. At an early period in the forma-

tion of the temporary teeth, the investing sac gives off a small process or bud, containing a portion of the essential rudiments, namely, the pulp covered by its proper membrane. This constitutes the rudiment of the permanent tooth. It commences in a small thickening on one side of the parent sac, which gradually becomes more and more temporary in an incisor.

b. Permanent rudiment given off from the circumscribed, and at length as- temporary in a molaris. - (T. Bell.) sumes a distinct form, though still



a. Permanent rudiment given off from the

connected with it by a pedicle. For a time, the new rudiment is contained within the same alveolus as its generator, which is exca-


vated by the absorbents for its reception. It is not, according to Mr. Bell, in consequence of the pressure of the new rudiment upon the bone, that this absorption is occasioned, but by a true process of anticipation; for he states, that he has seen, in the human subject - and still more evidently in the foal - the commencement of the excavation before the new sac was formed, and, consequently, before any pressure could have taken place on the parietes of the socket. The absorption does not, indeed, begin in the smooth surface of the socket, but in the cancelli of bone immedi-Temporary tooth ately behind it. By degrees, a small recess is thus and permanent formed in the paries of the alveolus, in which the new Bell.) rudiment is lodged, and this excavation continues to



increase with the increasing size of the rudiment; whilst, at the same time, the maxillary bone becomes enlarged, and the temporary tooth, advancing in its formation, rises in the socket. The new cell is thus gradually separated from the other, both by being itself more and more deeply excavated in the substance of the bone,

<sup>\*</sup> The Anatomy, Physiology, and Diseases of the Teeth, Lond. 1829.

and also by the formation of a bony partition between them, as seen in the marginal figure, 281, which exhibits the connexion between the temporary tooth and the permanent rudiment, as it exists after the former has passed through the gum. As the temporary tooth grows and rises in the jaw, the connecting cord or pedicle elongates, and although the sac, from which it is derived, is gradually absorbed, it still remains attached to the neck of the temporary tooth. The situation of each permanent rudiment, when its cor-




Temporary Teeth and Permanent Rudiments. -

responding temporary tooth has made its appearance through the gum, is deeper in the jaw and a little behind the latter, as represented in the marginal illustrations, (Figs. 282 and 283,) of the upper and lower jaw after the whole of the temporary teeth have passed through the gum. From these, it will be understood, how the upper part of the sac of the permanent rudiment, being.

by means of the cord connected with the gum, gradually assumes the same relation to the gum as was originally sustained by the temporary rudiment.

The ossification of the permanent teeth commences from




Temporary Teeth and Permanent Rudiments. - (T. Bell)

the third to the sixth month after birth, — for the incisors and first molaris; about the ninth month, for the canine teeth; about three years, for the second molaris; at three years and a half, for the fourth; and, at ten years, for the fifth; but all this is liable to much variation.

The permanent teeth, during their formation, are crowded together in the jaw; but, as soon as they have advanced to a certain point, and can no longer be contained within their own alveoli, absorption of the anterior parietes of those cavities takes place, and the teeth are allowed to come in some measure forwards. In consequence of such absorption, it frequently happens, that not only the socket of the corresponding temporary tooth, but that of the tooth on each side is opened to the permanent one. Absorption now occurs in the root of the temporary tooth, — generally at the part nearest its successor, and this gradually proceeds as the latter advances, until the root is completely removed,

when the crown falls off, leaving room for the permanent tooth to

supply its place. does not seem, that this absorption of the root is produced by pressure on the part of the permanent tooth, as it often happens, according to Mr. Bell, that the root of the temporary tooth is wholly absorbed, and the crown falls out spontaneously, long before the succeeding tooth has approached the vacant space. As a general rule, however, the actions must be regarded consentaneous; and Mr. Bell thinks, that this absorption resembles that, already referred to, for the formation of a new cell to receive the perma-



Upper and lower Teeth of the left side of the Jaws. - (T. Bell.)


nent pulp, and that it may be termed, like it, a "process of anticipation." In both instances, the existence, though not the pressure, or even the contact, of the new body is necessary to excite the action of the absorbent vessels; and we, accordingly, find, that in those cases, by no means unfrequent, in which the temporary teeth retain their situation in the mouth, with considerable firmness, until adult age, the corresponding permanent ones have not been formed.

The following are the periods at which the permanent teeth generally make their appearance. They are extremely irregular, however, in this respect: the estimate must, consequently, be regarded as a general approximation only.

	* *								
Anterior or first great	mola	res.	-	-		-			6½ years.
Middle incisors,				-	1	-	-		6
Lateral incisors,			-		-	- 1	1 -11		8
Anterior bicuspids, or	first	esse	r mola	ares.					9
Posterior bicuspids, or	r seco	nd I	esser	mola	res.				10
Canine teeth, -	_			-		-		-	11 or 12
Second great molares,		-	-	-		-	1000		12 or 13
Third great molares of		es 81	abient	iæ.	-				17 to 20

When these have all appeared, the set is complete, consisting of thirty-two teeth, sixteen in each jaw, — the number of temporary teeth having been only twenty. Fig. 284 represents the upper and lower permanent teeth in their alveoli or sockets, the external alveolar plate having been removed to show the mode in which

they are articulated. Fig. 285 represents the same teeth when removed from the socket.



a, a. Central incisors. b, b. Lateral incisors. c, c. Canine teeth. d, d. First bicuspidati. e, e. Second bicuspidati. f, f. First molares. g, g. Second molares. h, h. Third molares or dentes sapientiæ.

While the jaws are becoming furnished with teeth and increasing in size, they undergo a change of form, and the branches become more vertical, so as to favour the exertion of force during mastication. When the teeth issue from the gums, they are most favourably situate for the act of mastication; the incisors are sharp, the canine pointed, and the molares studded with conical asperities; but, in the progress of age, they become worn on the surfaces, which come in constant contact.

During the occurrence of these changes, which embrace the whole of the period we are considering, and extend, at times, into the two next, the animal functions, especially that of sensibility, become surprisingly developed, and the intellectual and moral results of a well adapted system of education are strikingly apparent. The nutritive functions are, likewise, performed with energy, the body not yet having attained its full growth; and, towards the end of the period, the organs of reproduction commence that development, which we have to describe under the next period.

The teeth appear with sufficient regularity to permit some inference to be drawn with regard to the age of the individual, and accordingly it has been proposed by Mr. Saunders to make them a test for age in reference to the children employed in the factories of Great Britain.

### 3. ADOLESCENCE.

The commencement of this age is marked by one of the most extraordinary developments, which the frame experiences, and its \* The Teeth, a Test for Age, considered with reference to the Factory Children. Addressed to the Members of both Houses of Parliament, Lond. 1837.

termination by the attainment of full growth in the longitudinal direction. The period of the former of these changes is termed

puberty; that of the latter the adult age.

The age of adolescence has been considered to extend from fifteen years to twenty-five, in men; and from fifteen to twenty-one, in women; but this is only an approximation, like the other divisions of the ages, all of which are subject to great fluctuations in individual cases.

During the periods we have already considered, no striking difference exists between the appearance of the male and female, except as regards the generative organs; but, about the age of puberty, essential changes occur, that modify the characteristics of the two sexes in a manner, which they maintain through the remainder of existence; and these changes affect the whole of the economy to a greater or less degree. In the male, the skin loses more or less of its delicacy and whiteness; the hair becomes darker, the cellular tissue condensed, and the muscles more bulky, so that they are strongly marked beneath the surface; the beard appears, as well as hair upon the pubes, chest, and in the axillæ. The different parts of the body become developed in such manner that the centre of the frame now falls about the pubes. The encephalon has increased in size, especially at the posterior and inferior part, the cerebellum, and has become firmer. The ossification of the bones, in the direction of their length, terminates towards the end of the period. The muscles become more red and fibrinous, losing the gelatinous character they previously possessed, and, in the animal, exhibiting those striking changes which we see - from veal to beef, from lamb to mutton, &c. The larynx undergoes great augmentation, and the glottis particularly is elongated and widened. The jaws complete their growth, and the dentes sapientiæ appear so as to make up the full complement of sixteen teeth in each jaw. The changes in the nutritive organs are not great, consisting chiefly in their development to correspond with the increased size of the frame. The greatest modification is produced in the organs of reproduction, which are now in a state to exercise their important functions. The testicles, at the period of puberty, suddenly enlarge so as to attain twice the diameter they previously possessed; and the secretion of sperm is accomplished. The penis is also greatly increased in size; and, according to Adelon,a "becomes susceptible of erection." The susceptibility, however, exists long before this age. It may be noticed even in the first period of infancy. The scrotum assumes a deeper colour. Such are the chief changes that supervene in the male.

In the female, they are not quite so striking; — the general habit remaining much the same as during childhood. The skin preserves its primitive whiteness; and, instead of the cellular tissue

Physiologie de l'Homme, 2de édit. iv. 448, Paris, 1829.

526 - AGES.

becoming more condensed, and the muscles more marked, as in the male, fat is deposited in greater quantity between the muscles, so that the form becomes more rotund. New hair appears on the organs of reproduction and in the axillæ, whilst that of the head begins to grow more rapidly. The development of the genital organs is as signal as in the male. The ovaries attain double their previous dimensions; the uterus enlarges; and a secretion takes place from it which has been elsewhere described - the menstrual flux; the mons veneris and labia pudendi are covered with hair; the labia enlarge and the pelvis has its dimensions so modified as to render labour practicable. At an early age, the long diameter of the brim is from before to behind; but it now assumes the opposite direction, or from side to side; and the bosom, which prior to this age, could scarcely be distinguished from that of the male, becomes greatly augmented; fat is deposited so as to give the mammæ their rotundity; the mammary gland is enlarged; and the nipple of greater size; - changes fitting the female for the new duties which she may be called on to exercise.

The functions undergo equally remarkable modifications, under the new and instinctive impulse, which animates every part of animal life. The external senses attain fresh, and peculiar activity; the intellectual faculties become greatly developed, and this is the period during which the mental character is more modified and improved by education than any other. It embraces the whole time of scholastic application to the higher studies. Prior to the end of the period, the male youth enters upon the avocation which is to be his future support, and both sexes may become established in life in the new relations of husband and wife, and of parent and child. It is during this age that an indescribable feeling of interest and affection is experienced between individuals of the two sexes; and that the boldness of the male contrasts so strikingly with the captivating modesty of the tender female,—

"That chastity of look, which seems to hang A veil of purest light o'er all her beauties."

The muscles, having acquired their strength and spring, the severer exercises are now indulged, and mechanical pursuits of all kinds, — military or civil, — are undertaken with full effect. The expressions participate in the altered condition of the mental and moral manifestations, and indicate vivacity, energy, and enthusiasm. The voice of the male acquires a new character, and becomes graver, for reasons assigned elsewhere; whilst that of the female experiences but slight modification. The nutritive functions of digestion, absorption and respiration experience but little change; but nutrition, strictly so called, is evidently modified, from the difference which we notice in the development and structure of the various organs. The muscles contain more fibrin; the blood is thicker and richer in globules; and the excretions mani-

fest a higher degree of animalization. Urea has usurped the place of benzoic acid in the urine; and the cutaneous transpiration has lost its acidulous smell, and become rank and peculiar. Lastly, the sexual functions are now capable of full and active exercise, and appear to be intimately connected with the energy and development of many parts of the economy. If the genital organs do not undergo the due change at puberty, or if the testes of the male or the ovaries of the female be removed prior to that age, considerable modification occurs. This is more manifest in the male, inasmuch as the ordinary changes, that supervene at puberty, are in him more marked than in the female. The removal of the testicles, prior to puberty, arrests those changes. The beard does not appear, nor the hair in the axillæ or on the pubes, as in the entire male; and if those animals in which the males are distinguished by deciduous horns, as the stag, — or by crests and spurs, as the cock, - be castrated before their appearance, such appendages never present themselves. If, however, they be castrated after puberty, they retain these evidences of masculine character. The eunuch, likewise, who becomes such after the appearance of the beard, preserves it, although to a less extent than usual.

The development of the larynx is arrested by castration, so that the voice retains, with more or less change, the treble of the period prior to puberty; and hence this revolting operation has been had

recourse to for the sake of gratifying the lovers of music.

In the course of age, we find that, during the progressive evolution of the organs, one set will be liable to morbid affections at one period, and another set at another. In the early ages, the mucous membranes and the head are peculiarly liable to disease; and, at the period we are now considering, affections of the respiratory organs become more prevalent. It is, indeed, the great age for pulmonary consumption, — that fatal malady, which, it was supposed by Sydenham, destroys two-ninths of mankind. In the female, whose proper feminine functions do not appear at the due time, or are irregularly exercised, the commencement, — indeed the whole of this period, — is apt to be passed in more or less sickness and suffering.

## 4. VIRILITY OR MANHOOD.

Hallé has divided this age into three periods, — crescent, confirmed, and decrescent virility. The first of these extends from the age of twenty-five to that of thirty-five in the male, and from twenty-one to thirty in the female; the second from thirty-five to forty-five in the male, and from thirty to forty in the female. Neither of these will require remark, the whole of the functions throughout this work, — when not otherwise specified, — being described as they are accomplished in manhood. Owing to the particular evolution of organs, however, the tendency is not now so great to morbid affections of the respiratory function. It is more especially the age for cephalic and abdominal hemorrhage;

accordingly, apoplexy and hemorrhoidal affections are more fre-

quent than at any previous period.

In decrescent virility, - in which Hallé comprises the period of life between forty and fifty in the female, and between forty-five and sixty in the male, - signs of decline are manifest. The skin becomes shrivelled and wrinkled; the hair is gray, or white and scanty; the teeth are worn at the top, chipped, loose, and many, perhaps, lost. The external senses, especially the sight, are more obtuse, partly owing to a change in the physical portions of the organ, so that powerful spectacles becomes necessary, and partly owing to blunted nervous sensibility. Owing to the same cause, the intellectual faculties are exerted with less energy and effect, and the moral manifestations are more feeble and less excitable. Locomotion is less active, owing to diminution in the nervous power, as well as probably to physical changes in the muscles, so that the individual begins to stoop, - the tendency of the body to bear forwards being too great for the extensor muscles of the back to counteract. The expressions participate in the condition of the intellectual and moral acts, and are, consequently, less exerted than in former periods. The nutritive functions do not exhibit any very remarkable change, and will even remain active until a good old age. The functions of reproduction show the greatest declension, especially in the female. The male may preserve his procreative capabilities much longer than this period, but in the female the power is, usually, entirely lost, the loss being indicated by the cessation of menstruation. After this, the ovaries shrivel, the uterus diminishes in size, the breasts wither, the skin becomes brown and thick, long hairs appear on the upper lip and chin, and all those feminine points are lost, which were previously so attractive. The period of the cessation of the menses is liable to many different disorders, which are the source of much annoyance, and are, at times, attended with fatal consequences. Prior to their total disappearance, they generally become extremely irregular in their recurrence, sometimes returning every fortnight, debilitating by their frequency, and by the quantity of the fluid lost, and laying the foundation, in many cases, for uterine or other diseases of a serious character. Cancerous affections of the mammæ or labia, which had been previously dormant or not in existence, now arise or become developed, and at times with extreme rapidity. In consequence of the great liability to such affections, this has been called the critical age, critical period or critical time of life or turn of life. The danger to the female is not, however, so "critical" at this period as the epithet might suggest, - the statistical researches of De Chateauneuf and of Lachaise, Finlaisona and others having shown, that between the ages of forty and fifty no more women die than men.b M. Constant Saucerotte has, indeed, attempted to show by statistics on a great scale, that the

<sup>&</sup>lt;sup>a</sup> Reports on the Evidence and Elementary Facts on which the tables on Life Annuities are founded, Lond. 1829.

b Dr. D. Davis, in Principles, &c., of Obstetric Medicine, i. 290, Lond. 1836.

mortality amongst women is greater between the ages of thirty and forty than between forty and sixty; and Muret, in his statistics of the Pays du Vaud, did not find between forty and fifty a more critical age than between ten and twenty.

#### 5. OLD AGE.

This is the age when every thing retrogrades. It is the prelude to the total cessation of the functions, where the individual expires, — which is but rarely the case, — from pure old age. This period, has been divided into three stages: — incipient or green old age, reaching to seventy years; confirmed old age or caducity, to eighty-five years; and decrepitude, from eighty-five years upwards.

In incipient or green old age, the declension, which had occurred in the period of decrescent virility, is now more evident. The intellectual and moral manifestations exhibit more marked signs of feebleness; and the muscular powers totter, and require the aid of a support, — as well to convey a part of the weight of the body to the ground, as to enlarge the base of sustentation. The muscles of the larynx participate in this general vacillation; the

"Big manly voice.
Turning again towards childish treble, pipes
And whistles in the sound,"

and is broken and tremulous.

The appetite is great, and the powers of digestion are considerable; but mastication is largely deteriorated. In the first place, the teeth fall out, in consequence of the constant deposition of fresh layers in the dental cavities, which ultimately close them, and obliterate the vessels that pass to the internal papillæ for their nutrition. As soon as the teeth have fallen out, the alveolar processes, which supported them, waste away by absorption, and the depth of the jaw is thus greatly lessened. On these accounts, the jaws only approach each other at the forepart; the chin projects, and the angle of the jaw is thrown more forward. As the teeth and the sockets disappear, the alveolar margins become thin and sharp, and the gum hardens over them; the chin and nose necessarily approach (Figs. 286 and 287), the lips fall in, and the speech is inarticulate. We can thus understand the peculiarities of the mastication of the aged. They are compelled to bite with the anterior portions of the jaws; for which reason, as well as owing to the greater obliquity of the insertion of the levator muscles of the lower jaw, but little force can be exerted; and owing to the too great size of the lips, the saliva cannot be retained. Respiration is not as readily accomplished, partly owing to the complete ossification of the cartilages of the ribs, but

<sup>\*</sup> Churchill, Outlines on the Principal Diseases of Females; Dunglison's American Medical Library Edit., p. 82, Philad. 1839; or Professor Huston's edit., Philad. 1842.
VOL. II. — 45

chiefly to diminished muscular powers, and many of the bloodvessels, especially of the extremities, become more or less ossified, and the pulse is somewhat slow and intermittent, but generally perhaps faster than in the adult. Of



Skull of the Aged, - (Sir C. Bell.)

The valves of the heart

255 women - between the ages of 60 and 96 examined by MM. Hourmann and Dechambres -the average number of pulsations in the minute was 82.29; of respirations, 21.79. Nutrition is effected to such a degree only as to keep the machine in feeble action; and animal heat is formed to an inadequate extent, so that the individual requires the aid of greater extraneous warmth: in many cases,

the powers of reproduction in the male are completely lost. In confirmed old age, the debility of the various functions goes



Physiognomy of the Aged .- (Sir C. Bell.)

on augmenting. The mental and corporeal powers almost totter to their fall, and often a complete state of dementia or dotage exists. Frequently, however, we are gratified to find full intellectual and moral enjoyment prevailing even after this period, with the possession of considerable corporeal energy. The author has had the honour to enjoy the friendship of two individillustrious uals of this country, who had filled the

highest office in the gift of a free people, both of whom are now no more; each of these gentlemen exhibited, after the lapse of Archiv. Général. de Médec. 1825.

OLD AGE. 531

eighty-three summers, the same commanding intellectual powers

and the same benevolence that ever distinguished them.

In this stage, locomotion becomes more difficult; the appetite is considerable, and the quantity eaten at times prodigious, the digestive powers being incapable of separating the due amount of chyle from a quantity of aliment, which was sufficient in the previous ages. Difficulty, however, sometimes arises in defecation, the muscular powers being insufficient to expel the excrement. From this cause, accumulations occasionally take place in the rectum, which may require the use of mechanical means,—as injections, the introduction of an instrument to break them down, &c. Generation is, usually, entirely impracticable, erection being impossible; and during the whole of this and the next stage, the urinary organs are liable to disorder—irritability about the neck of the bladder, and incontinence of urine, being frequent sources of annoyance.

The density of the lungs — together with the quantity of blood they admit — diminishes with the progress of age; the thorax itself is gradually accommodated to the change; it becomes atrophied as the lungs are atrophied; it contracts as they contract, and the diminution in their vascularity, which is always in a ratio with the diminution in texture, shows the direct proportion between the weakened chemical power and the diminished mechanical forces.

(Magendie, and MM. Hourmann and Dechambre.)

Finally, to this stage succeeds that of decrepitude, so well described by Shakspeare:

"Last scene of all,

That ends this strange, eventful history,
Is second childishness, and mere oblivion;
Sans teeth, sans eyes, sans taste, sans every thing."

As You Like It, ii. 7.


The loss of power, mental and corporeal, becomes progressively greater; and, in addition to the abolition of most of the external senses — especially those of sight and audition — the intellectual faculties are, perhaps, entirely gone; all muscular motion is lost, and paralysis requires constant confinement to the bed, or to the easy chair; the excretions are passed involuntarily; sensibility becomes gradually extinct, and life finally flits away as imperceptibly as the twilight merges in the shades of night.

M. Quetelet, of Brussels, has deduced from extensive observations the relative heights and weights of both sexes at different periods of existence. The results are exhibited in the subjoined diagram. (Fig. 288.) The increase in height is most rapid in the first year, and afterwards diminishes very gradually: between the ages of 5 and 16, the annual increase is very regular. The difference

Annales d'Hygiène publique, vi. 89; and in his work, Sur l'Homme, et le Dévelopment de ses Facultés, Bruxelles, 1835; or translation of the same, p. 63, Edinb. 1842.

between the height of the male and female at birth continues to increase during infancy and growth; but it is not very marked until about the 15th year, after which the female grows at a diminished rate; whilst the male goes on in nearly the same degree until about the age of 19. The female, consequently, attains her full height earlier than the male;—the full height of the latter not being generally attained until about the age of 25. At about 50,

Fig. 288.



Curves indicating the development of the heights and weight of man and woman at different ages,—(Quetelet.)

both sexes experience a diminution of stature, which continues during the latter part of existence. The average heights of the male and female who have obtained their full development are about 31 times those of the new-born infant of the sexes respectively. The relative weight of the sexes corresponds pretty nearly with the height. The preponderance exhibited by the male at birth increases gradually during the first few years; but towards the period of puberty the proportional weight of the female increases; and about the age of 12 there is but little difference between the sexes. After this, however, the weight of the male increases much more rapidly, especially between 15 and 20: after this, there is not much increase on the part of the male, although his maximum is not attained until the age of 40; and there is an absolute diminution on the part of the female, whose weight remains less during nearly the whole period of child-bearing. After this, however, her weight again experiences an increase, and its maximum is attained about 50. In old age, the weight of both sex es undergoes a diminution in nearly the same degree. The average weights of the male and female who have attained their full development are twenty times those of the new-born infant of the sexes respectively.2

Such is a brief description of the chief changes, that befall the body in the different ages. To depict them more at length would be inconsistent with the object and limits of this elementary work. It is clear, that, although the divisions, which we have adopted from Hallé, are entirely arbitrary, must run into each other, and be liable to numerous exceptions; - certain well-marked changes occur about the commencement or termination of many of them, and singular diversity takes place in the successive evolutions of organs: whilst some are predominant at one time, they fall behind others at a previous or subsequent period; and such changes may lay the foundation for morbid affections at one age, in certain organs, which do not prevail at another. The ancients, who believed that great mutations occur at particular intervals, every three, seven or nine years for example, as the particular number might be at the moment in favour, - compared these periods to knots uniting the different stages of life, and giving the economy a new direction. These knots they called the climateric or climacteric years, and they conceived the body to be especially liable to disease at the periods of their occurrence. The majority assigned them to the number seven and its multiples; and the fourteenth and twenty-first years especially were conceived to be replete with danger. Others applied the time climacteric to years resulting from the multiplication of seven with an odd number, and especially with nine; the sixty-third year being regarded, by almost all, as the gland climacteric. The error, with the ancients, lay, in considering that the numbers exerted any agency. Every one admits the influence of particular evolutions on the health; and, at the present day, the word climacteric is generally restricted to certain periods of life, at which great changes supervene, independently of any numerical estimate of years; such as the period of puberty in both sexes; — that of the cessation of the menses, or the critical time of life in the female, &c.

It need hardly be remarked, that the different ages we have described, instead of extending through the protracted period of eighty-five years and upwards, may be varied by original constitution, climate, habits of life, &c., so that the stages may be shorter than usual, and all the signs of decrepitude occur many years earlier; and, on the other hand, the period of decrepitude may, through strength of original conformation, and other causes, be

largely postponed.

<sup>\*</sup> Carpenter, Human Physiology, § 772, Lond. 1842.

## CHAPTER II.

SLEEP.

THE difference between the two classes of animal and nutritive functions is strikingly exhibited in the phenomena we have now to consider. Whilst the former are more or less suspended, the latter continue their action with but little modification. The functions of sensibility, voluntary motion, and expression, cannot be indulged for any length of time, without fatigue being induced, and a necessity arising for the reparation of the nervous energy which has been expended during their action. After a time, -the length of which is somewhat influenced by habit, - the muscles have no longer power to contract, or the external senses to receive impressions; the brain ceases to appreciate; mental and moral manifestations are no longer elicited; the whole of the functions of relation become torpid, and remain in this state until the nervous system has been renovated, and adapted for the repetition of those functions, which, during the previous waking condition, had been exhausted. This state constitutes sleep; which, consequently, may be defined - the periodical and temporary suspension of all, or most, of those functions that connect us with the universe. The suspension occurs in those functions and in those only; and hence the consideration of sleep, in many physiological treatises, has immediately followed that of the functions of relation. The nutritive functions continue regularly in action from the earliest period of fœtal formation; before mental manifestations exist in the embryo, and during sleep. For them there is no cessation, and scarcely any declension of activity, until the decadency of the frame affects them along with the whole of the machinery. Sleep, in the language of poetry, has been compared to death; and Dr. Gooda has stated, that the resemblance between them is not less correct upon the principles of physiology, than it is beautiful among the images of poetry. "Sleep is the death or torpitude of the voluntary, organs, while the involuntary continue their accustomed actions. Death is the sleep or torpitude of the whole." Physiologically, the difference appears to us considerable. During the whole of sleep a process of renovation is probably going on in the organs of animal life, which adapts them for subsequent activity, and contrasts signally with the state of annihilation that constitutes death; hence the important difference between healthy sleep, and the state of coma induced by any morbid cause; from which the patient is aroused languid and exhausted, instead of active and recruited.b The fœtus in utero is also described by some as being in a perpetual sleep, until aroused by the new actions established

<sup>2</sup> Book of Nature, ii. 226, Lond. 1826.

b See the author's Therapeutics, p. 401, Philad. 1836; and his General Therapeutics and Materia Medica, vol. ii. Philad. 1843.

SLEEP. 535

at birth; but it appears to us, that there must be, even in this case, alternations of activity and suspension in the nervous functions. We have seen elsewhere, that they are manifestly more or less exerted during intra-uterine existence; nervous energy must therefore be expended; and renovation,—to a much less extent, it is true, than in the new-born child,—be necessary. Linnæus, under the term somnus plantarum, expresses a peculiar state in the constitution of many plants during the night, as evinced by a change of position, generally a drooping or folding together of their leaves or leaflets; such a change being occasioned by the withdrawal of the stimulus of light, and, probably, it has been conceived, constituting a state of rest to their vital functions; but it is obvious, that there can be no similitude between this condition and that of the sleep of animals, which is confined to the functions of relation,—

functions that do not even exist in the vegetable.

The approach of sleep is indicated by signs, that are unequivocal, and referable to the encephalon. The great nervous centre of animal life, feeling the necessity for rest and renovation, an internal sensation arises in it, as well as in the whole of the nervous system over which it presides, termed sleepiness or the sensation or want or desire of sleep, which, provided the waking state has been protracted, ultimately becomes irresistible, and often draws on sleep in spite of every effort to the contrary. It is affirmed, that boys, exhausted by exertion, dropped asleep amid the tumultuous noise of the battle of the Nile; and the fatigued soldier has often gone to sleep amid discharges of artillery. An engineer has been known to fall asleep within a boiler whilst his fellows were beating it on the outside with their heavy hammers. Noises will at first prevent sleep, but the desire is ultimately so invincible, that they cease to produce any effect. In the noisy inns of large towns, where the perpetual arrivals and departures of travellers keep up an incessant din and confusion, sleep may be for a time withheld, but it ultimately supervenes, although the tumult may be even tenfold; and if the noise should, from any cause, suddenly cease, the individual will probably awake. It is reported of the proprietor of some vast iron-works, who slept close to them, not withstanding the noise of sledge-hammers, forges and blast-furnaces, that he would immediately awake if any interruption occurred during the night. This effect of habit is seen in the infant, which has been accustomed to the cradle. The moment the motion and noise of the cradle, or the sound of the nurse's voice - if she have been in the custom of singing the child to sleep - ceases, it awakes.

When the desire for sleep sets in vigorously, the animal functions become more obtuse, until they progressively fail to be exerted. The cessation does not occur in all simultaneously. The power of volition is gradually lost over the muscles; the eyes cannot be kept open; the upper eyelid falls, and if we attempt to

<sup>\*</sup> Amænitat. Académ. tom. iv ; and Choulant, in art. Schlaf, Pierer's Anat. Phys. Real Wörterb. vii. 257, Altenb. 1827.

raise it again, it appears to be weighed down, the eyeball is directed upwards, and the pupil is contracted; a the arms fall where gravity would take them; the extensor muscles of the back, deprived of volition, cease to contract, and the head falls suddenly forwards, occasioning nodding, which rouses the brain to momentary action, to be again lost, however. If the individual be in the erect attitude, his limbs bend under him; and if in the sitting posture, the head gradually falls upon the chest; the extensors of the trunk no longer contract with sufficient force to obviate its tendency to fall forwards; and the attitude, unsupported, cannot be maintained. The same gradual suspension occurs in the muscular movements, concerned in speech and in the production of the voice, which becomes feeble, confused, broken, and ultimately lost. In short, all the strictly voluntary muscles have their action suspended, if we except the orbicularis palpebrarum muscle, which, according to Broussais, b now contracts to close the eye and shut off the stimulus of light. If we determine to resist the desire for sleep, we yawn and stretch, for the reasons elsewhere assigned, and endeavour to arouse the functions to renewed activity. If the state of wakefulness have not been long protracted, we are successful; but all our endeavours fail, if the nervous system be so far exhausted as to render reparation indispensable. From the commencement of sleepiness, the action of the senses is enfeebled, and gradually suspended. The sight yields first, the closure of the eyelids preventing the organ from being impressed by its special irritant. The smell yields after the taste; the hearing after the smell; and lastly, the touch sleeps; although the appropriate irritants may continue to reach the organs of these senses. All the internal sensations, hunger, thirst, &c., as well as the morbid sensation of pain, are no longer appreciated. The intellectual and moral manifestations exhibit, from the commencement of the feeling of heaviness, the languor which pervades the frame. The will gradually ceases to control the functions that are under its dominion, until ultimately the power of volition is lost. In the less perfect kind of sleep, or in slumber, the ideas flit in a disorderly manner, constituting a kind of delirium; but when sleep is complete, the whole encephalic organ appears to be at rest, and perceptions are no longer accomplished: special irritants may be applied to the external senses, but they excite no sensation. Many physiologists affirm, that the internal functions of nutrition acquire more energy during sleep; but Broussaisd properly disputes the affirmation, and maintains, that the want of action in the senses, muscles, and intellect, must necessarily occasion diminished energy in the nutritive functions. During sleep, circulation and respiration appear to be retarded: perspiration is less active, and diges-

Mojon's Leggi Fisiologiche, &c., translated by Skene, p. 34, Lond. 1827.

b Traité de Physiologie, &c. ; or Drs. Bell and La Roche's translation, 2d Amer. Edit. Philad. 1832.

<sup>&</sup>lt;sup>c</sup> Mojon, op. cit., p. 89, Lond. 1827.

SLEEP. 537

tion more tardy than in the waking condition. The difference in the last respect is so great, that, as Broussais remarks, the appetite recurs many hours before the usual time where long watching is indulged, and an additional meal becomes necessary; proving the truth of the old French proverb, — "qui dort dine" — "who sleeps dines." Secretion, nutrition, and calorification are also less energetically performed than usual. Absorption alone, according to some, is more active; but there seems not to be sufficient reason even for this assertion. This notion of the greater activity of the nutritive organs is as old as Hippocrates, and has been acquiesced in by almost all subsequent writers without examination, especially as it seemed to show a kind of alternation and equipoise between the respective periods of activity of animal and organic life.

During sleep, then, all the animal functions are suspended, and the body generally remains in a state of semiflexion, the one which, as we have elsewhere seen, requires little natural effort. To this, however, there are numerous exceptions depending upon habit. The easiest position for the body is perhaps on the back. It is the one assumed in extreme debility, when the prostration is so great that the individual sinks down in the bed like a dead weight; but the extensor muscles of the thigh and leg, under such circumstances, become fatigued, and relief is obtained by drawing the feet upwards so as to elevate the knees. This is a common attitude in the most debilitating maladies, and is often maintained

until within a short time prior to dissolution.

Sleep can persist with the exercise of certain muscles. Couriers, on long journeys, will nap on horseback; and coachmen on their boxes. The author has seen a servant boy erect and asleep in the intervals between the demand for his services at the table.

During the first sleep, the suspension of the animal functions is the most complete; but, towards morning, some of them become less asleep, or more excitable than others. The intellectual and moral faculties are frequently inordinately active, giving occasion to dreams, which, with some individuals, occupy a great portion of the period allotted to rest. The sense of tact, too, is easily roused. If we lie in a position which is disagreeable, it is soon changed; the limbs are drawn away, if irritated in any manner; the clothes are pulled up, if the air is disagreeably cold, &c. The sense of sight and the voluntary motions are least readily aroused, so that those functions, which fall asleep the last, are most easily awakened, and they gradually resume their activity in the order in which they lost it. After six or eight hours of sleep, - more or less according to circumstances, - the individual awakes, not generally at once, however; a state of slumber, like that which preceded sleep, now succeeds it. The organs, which are the last to resume their activity, require to be excited to the performance of their functions. The eyes are rubbed; stretching is indulged, which recalls the nervous influx to the muscles; and sighing and yawning arouse the muscles of respiration, and compensate, in some measure, for

the minor degree of aëration of the blood accomplished during sleep. The urine is discharged, and the phlegm, which may have collected in the air-passages, is expectorated: these excretions have accumulated during sleep, because, owing to diminished sensibility, the call for their evacuation has not been as urgent. In cases of catarrh, accompanied by copious mucous secretion, and in phthisis pulmonalis, the fluid will collect in surprising quantity in the airpassages during sleep, and it is expectorated as soon as the brain

is sufficiently aroused to respond to the sensation.

When the individual is fully awake, the energy, with which the animal functions are exercised, exhibits that the nervous system must have entirely recruited during its state of comparative inaction. The period of sleep, necessary for this purpose, varies in different individuals, and at different ages. Some require eight or ten hours: others not more than three or four; and others are said to have been contented, throughout the course of a long life, with not more than one or two. Men of active minds, whose attention is engaged in a series of interesting employments, sleep much less than the lazy and the listless. General Pichegru informed Sir Gilbert Blane, that in the course of his active campaigns he had, for a whole year, not more than one hour of sleep, on an average, in the twenty-four hours. The great Frederick of Prussia, and the yet more great Napoleon, are said to have spent a surprisingly short time in rest; but with respect to the latter, the fact is controverted by one, b who had excellent opportunities for observation. It is probable, that in these cases the sleep is more intense, and that such of the animal functions, as require rest indispensably, are completely suspended during the whole period consigned to it. These are the functions of voluntary motion more particularly; the intellectual and moral faculties requiring a much shorter period of repose, as is manifest by their incessant activity during dreaming,a condition, which, with some, continues through almost the whole night. The same individual, too, will spend a shorter time in sleep, when strongly interested in any pursuit, than in the monotonous occurrences of ordinary life, and, when any subject occupies us intently, it will frequently keep us awake in spite of ourselves; but although the period of sleep may be protracted much beyond the accustomed hour by unusual excitation, the effect of the stimulus becomes insufficient, and sleep comes on under circumstances, which appear most unfavourable to it. lunatic affords us a wonderful example of powerful resistance to sleep and fatigue, or rather of the short period, which is necessary for the renovation of the nervous system, kept almost incessantly upon the stretch, as it is, in many of these distressing cases. It has been a common remark, that women require more sleep than men, and Georgete assigns them a couple of hours more, -

Medical Logic, 2d edit. p. 83; see, also, Macnish, Philosophy of Sleep, Amer. Edit. p. 35, New York, 1834; and Elliotson's Human Physiology, p. 602.

Bourrienne, Private Memoirs of Napoleon Bonaparte, Amer. Edit. Philad. 1831.

De la Physiologie du Système Nerveux, &c., Paris, 1821.

DREAMS. 539

allotting to men six or seven hours, and to women eight or nine; but Dr. Macnisha judiciously doubts, whether the female constitution require more sleep than the male: at least, he says, it is certain, that women endure protracted wakefulness better than men, "but whether this may result from custom is a question worthy to be considered." The fact is, however, too general to allow custom to be invoked. It would seem, indeed, that the female frame, although far more excitable than that of the male, is longer in having that excitability exhausted, and that the recuperative powers are greater, so that the excitability, when exhausted, is more readily restored. The notion, that the female needs more rest than the male, appears to be traditionary, and like most traditions, to have been handed down from one individual to another, without due examination. The degree of muscular and mental exertion, to which the male is accustomed, would seem to indicate that a longer period of rest ought to be required by him to admit of the necessary restoration of excitability. In infancy and youth, where the animal functions are extremely active, the necessity for sleep is greatest; in mature age, where time is more valued and the cares are more numerous, it is less indulged; whilst the aged may be affected in two opposite ways; they may be either in a state of almost constant somnolency, or their sleep may be short and light.

Sleep has been divided by the physiologist into complete, and incomplete. The former is characterized by suspension of all the animal functions; a state, the existence of which has been doubted by many. Certain it is, that it can occur but rarely, and only when all the organs have stood in equal need of rest and renovation; and when none have preserved, from the preceding state of waking, a peculiar susceptibility for action. The nearest approach to it occurs in the first hours of repose; after this, it becomes incomplete; some of the functions are not equally sound asleep, and consequently respond to excitants with different degrees of facility; and the various organs do not require the same time for reparation, and therefore awake at different intervals; hence, dreams arise, which occur chiefly towards morning, or after the sleep has become incomplete; that is, when some of the animal functions are more or less actively, but irregularly, exer-

cised.

#### 1. DREAMS.

Anciently, dreams were regarded as supernatural phenomena, under the control of the children of Somnus or Sleep,— Morpheus, Phobetor or Icelos, and Phantasos. These three children, according to Ovid, were capable of transforming themselves into any form; the employment of Morpheus being to counterfeit the

Metamorphos. xi. v. 592 ad 645.

<sup>\*</sup> Op. citat. p. 280.

b See, on the Hygienic Relations of Sleep, the Author's "Elements of Hygiène," from p. 444 to 455 inclusive, Philad. 1835.

forms of men; Phobetor imitated the likeness of brutes and objects of terror; and Phantasos that of inanimate creatures. For a long time dreams were supposed to reveal future events by types and figures; as when Hecuba dreamed she had conceived a firebrand, and Cæsar that he should lie with his mother; which was interpreted, that he should enjoy the empire of the earth, - the common mother of all living creatures. Oneiromancy was an encouraged art, and ministered largely to the credulity and superstition of the people. Strange to say, there are yet those, who look upon dreams to be typical and instructive, and consequently supernatural! Mr. Baxtera and Bishop Newton openly maintained this doctrine. They divide dreams into two kinds, - good and evil, - and conceive, that two kinds of agents, good and evil spirits, are concerned in their production; they consequently account for the one or the other sort of dreams, according as the one or the other kind of agents obtains a predominancy !b It is not necessary to combat these views, - which ought of course to be as applicable to animals as to man, - especially as they are universally discarded. Dreaming is now properly considered to be an irregular action of the brain, in which the great controlling power of the will has suspended its agency, and allowed the memory and imagination unlimited sway, so that the most singular and heterogeneous ideas are formed, - still kept, however, somewhat in train by the force of association. At times, indeed, this influence is so great, that every part of the dream appears to go on in the most natural and consistent manner. We witness scenes that have occurred during our waking hours; and seem to see, hear, walk, talk, and perform all the ordinary offices of life. The mind reasons, judges, wills, and experiences all the various emotions. Generally, the whole process is confined to the brain, but, at times, the muscles are thrown into action, and the expression of the feelings and emotions occurs, as in the waking state. The dreamer moves, speaks, groans, cries, sings, &c., and if the dream concern the generative function, the external organs respond, and emission takes place in the male to such an extent, occasionally, as to constitute a true disease, or to be the cause of such, - the paroniria salax of Good, the gonorrhea dormientium, or night pollution of others. During the prevalence of a passion, too, the nutritive organs, in which its effects are experienced whilst awake, may be equally concerned during sleep. The respiration is short and interrupted, and sighs, groans, or laughter, according to the character of the emotion, are elicited; the heart beats with more or less violence, and this state of excitement often continues after the individual has been completely aroused. The nightmare, ephialtes, or incubus affords us an example of suffering as intense as could well be experienced during our waking moments. A sensation of dis-

An Inquiry into the Nature of the Human Soul, &c., Lond. 1730.

<sup>Good's Book of Nature, ii. 13, Amer. Edit., Bost. 1826.
Physiological System of Nosology, cl. iv. ord. 1, gen. v. sp. 3.</sup> 

DREAMS. 541

tressing weight is felt at the epigastrium, and of impossibility of motion, speech or even respiration; the dreamer fancies that some horrible form, or some ferocious being is approaching him, and that all chance of escape is precluded; or that he is about to fall, or is falling, from a lofty precipice; and the anguish which he suffers is indicated by loud groans, or by such painful feelings, apparently in the organs to which the emotions are referred, that he awakes. The ideas, at these times, are even more vivid than during the waking condition; the perceptions, that predominate, not being detracted from by extraneous impressions. On many of these occasions, when we awake, the dream is fresh upon the memory; and by resigning ourselves again to slumber, we can at times recall it, should it be of an agreeable character, or dispel it

altogether by rousing ourselves thoroughly.a

On account of the greater vividness of the ideas during sleep, and their freedom from all distraction, intellectual operations are sometimes effected in a surprising manner; difficulties being occasionally solved, which have obtained the mastery during waking. To a minor degree, every one must have experienced more or less of this. Composition, poetical or other, is often effected with the greatest facility; and a clue is occasionally afforded, which leads to the solution of previous difficulties. Cardan had a notion that he composed one of his works during sleep. Condillac, who attended greatly to this matter, remarked particularly, that, whilst engaged with his "Cours d'Etude," he frequently broke off a subject, before retiring to rest, which he developed and finished the next morning according to his dreams. Condorcet saw in his dreams the final steps of a difficult calculation, which had puzzled him during the day; and Dr. Gregory, of Edinburgh, composed thoughts, and clothed them in words, which were so just in point of reasoning, and so good in point of language, that he used them in his lectures, and in his written lucubrations. Voltaire, Lafontaine, Franklin, Coleridge, and others, have made similar remarks; and events of the kind must have occurred, in some shape, to almost every one. b Dr. Good relates a singular instance which happened to a friend of his, who, amongst other branches of science, had deeply cultivated that of music, of which he was passionately fond. He was a man of irritable temperament, ardent mind, and most active and brilliant imagination; and "was hence," says Dr. Good, "prepared by nature for energetic and vivid ideas in his dreams." On one occasion, during his sleep, he composed a very beautiful little ode, of about six stanzas, and set the same to very agreeable music, the impression of which was so firmly fixed

b Dendy, The Philosophy of Mystery, p. 232, Lond. 1841.

<sup>\*</sup> See, on this subject, the experiments of M. Girou de Buzareingues, in Magendie's Journal de Physiologie, tom. viii.

Book of Nature, loc. cit. See, also, Abercrombie, Inquiries concerning the Intellectual Powers, Amer. Edit. p. 234, New York, 1832; Macnish, Philosophy of Sleep, Amer. Edit. p. 65, New York, 1834; and Elliotson's Human Physiology, p. 615, Lond. 1840.

542 SLEEP.

in his memory, that, on rising in the morning, he copied from his

recollection both the music and the poetry.

In these cases, the will must direct, more or less, the intellectual process. It is scarcely conceivable, that the train of reasoning could go on so connectedly and effectively by association alone. That the will can, in some degree, be kept awake, or in a condition susceptible of being readily aroused, is shown by the facility with which we awake at a determined hour, and exercise a degree of watchfulness during sleep; as well as by the facts, previously mentioned, regarding the courier who sleeps on his horse, or the coachman on his box.

One curious fact, occasionally observed in dreams and likewise in the Mesmeric condition, which is analogous, in many respects, to what is observed in dreaming, is the calling up of impressions, that have been made at an antecedent period, and that may have been entirely forgotten during the waking state. A well-known case of this kind is recorded in the books. A woman, during the delirium of fever, constantly repeated sentences unknown to her attendants and which proved to be Hebrew and Chaldaic. Of these she knew nothing whatever on her recovery; but on referring to her previous condition, it appeared that she had formerly lived with a clergyman, who had been in the habit of reading aloud sentences in those languages, which had impressed her mind without her knowledge.

Dr. Dewar relates the case of a girl, who, when awake, discovered no knowledge of astronomy, or other sciences; but when asleep could define the rotations of the seasons, using expressions the most apt to the subject; and Mr. Dendy, alludes to the case of an Edinburgh lady, who, "during her somnolent attacks, recited somewhat lengthy poems;" and it was curious, that each line com-

menced with the final letter of the preceding.

There is a kind of dreaming, in which the sleep is less profound than during ordinary dreams; where the body has, consequently, more capability of receiving external impressions, but where the will has a certain degree of power over the muscles of voluntary motion, but imperfectly regulates the thoughts. This is somnambulism or sleep-walking. During the continuance of this state, the individual can apparently see, hear, walk, write, paint, speak, taste, smell, &c., and perform his usual avocations, yet remain so soundly asleep, that it is impossible to awake him without making use of violence. Cases are on record, and of an authentic nature, of individuals who have risen from bed asleep, with their eyes closed, and have not only walked about the room or house, going up or down stairs, finding their way readily and avoiding obstacles, but have passed with safety through very dangerous places, as windows, or on the roofs of houses. They have executed, too, yet more difficult feats; such as dressing themselves, going out of a The Philosophy of Mystery, p. 305, Lond. 1841.

DREAMS. 543

doors, lighting a fire, bathing, saddling and bridling a horse, riding, composing verses, &c., and executing all the actions of life correctly, and even acutely; yet they have been asleep during the whole of these acts. The eyes have been shut, or if open, have been incapable of perceiving the brightest light held before them; and the iris has not exhibited its irritability by contracting, so that it is doubtful whether the ordinary functions of the eyes be generally executed during somnambulism; and the fact of the serious accidents, that occasionally befall the sleep-walker, is in favour of this conclusion. It must be remarked, however, that, in the opinion of some physiologists, the sight is awake and employed, and there are cases which strongly favour the idea. A peculiarity of ordinary somnambulism is, that the train of thoughts is usually directed towards one point, and this so profoundly, that notwithstanding the activity of the imagination, and the firm hold it takes on the mind, no recollection is retained of the occurrences during sleep, after the individual awakes, either spontaneously, or by

being forcibly aroused.

Animal magnetisma would seem to be capable of inducing a peculiar kind of somnambulism, in which new faculties appear to be acquired, and intellectual operations to be executed, which are of the most astonishing character. The records of the Académie Royale de Médecine, of Paris, contain many such instances. A singular case of somnambulism is given by Dr. Belden, of Springfield, Vermont. It occurred in a young female, 17 years of age, and the phenomena were attested by numerous observers. One striking circumstance in this case was the astonishingly developed impressibility of the eye. As an evidence of this, when Dr. Belden, in order to test the sensibility of that organ, took one evening a small concave mirror, and held it so that the rays, proceeding from a lamp, were reflected upon her closed evelid; when the light was so diffused, that the outline of the illuminated space could scarcely be distinguished, it caused, the moment it fell on the eyelid, a shock equal to that produced by an electric battery. This female could see as well, apparently, when the eyes were closed as when they were open. The details of this case - and indeed of every case — of somnambulism are full of interest to the mental philosopher. Of late, the various experiments, at one time so much in vogue, when Mesmerism was in fashion, - have been repeated

5 American Journal of the Medical Sciences, No. xxviii.

<sup>\*</sup> See, on the subject of Animal Magnetism, Elliotson's Blumenbach, &c. p. 292, Lond. 1828; and Elliotson's Human Physiology, p. 660, Lond. 1840; Isis Revelata; An Inquiry into the Origin, Progress, and Present State of Animal Magnetism, by J. C. Colquhoun, Esq., Edinb. 1836; Prichard, Treatise on Insanity, p. 410, Lond. 1835; Erfahrungen über den Lebensmagnetismus und Somnambulismus — Commissions-Bericht an die Königl. Med. Akademie zu Paris, von Husson; und Resultate der Praxis einiger Hamburger Aerzte, so wie des Verfassérs, J. F. Siemers, Hamburg, 1835. See, also, Brit. and For. Med. Rev., April 1839, p. 301; and an interesting summary by Prof. J. K. Mitchell, in Quarterly Summary of the Transactions of the College of Physicians of Philadelphia, for Aug., Sept. and October, 1842:

544 SLEEP.

not only by those who are not in the ranks of the profession, but by certain estimable physicians; and of the reality of certain of the effects ascribed to the manipulations of the animal magnetizer we can entertain no doubt. The whole history of the art exhibits, that impressible individuals may have irregularities of nervous distribution induced through the medium of the senses, especially through those of vision and touch, - and that somnambulism and hysteric sleep, with other phenomena referable to a like condition of the nervous system, may be engendered; but that there is any thing like a magnetic fluid or agent, which may be communicated from the magnetizer to the subject of his experiments, is not only not proved but, in the author's opinion, by no means presumable. One of the most curious of the phenomena presented by this singular condition is the greatly developed sensibility to some irritants, and the total insensibility to others. Thus, the author has seen different persons bear without the slightest muscular action the application of a straw or feather to the conjunctiva; the insertion of pointed bodies into various parts of the cutaneous surface; the extraction of a tooth, &c., and yet start at the least puff of air made to come in contact with the face. Dr. Carpentera considers, that these phenomena must be considered as still doubtful; but they have been so often witnessed as not to admit of disbelief. From what the author has himself seen, he can readily credit the statements affirmed on respectable testimony, that even the major operations of surgery may have been executed, whilst the patient was in this state of Mesmeric sleep — if it may be so termed. As to the Hellsehen, clairvoyance, or "lucidity of vision," said to have been possessed by the magnetized, could we assign our belief to it at all, it would be only on the ground, - " credo quia impossibile est."

The causes of imperfect or incomplete sleep, and hence of dreams, are various. The fact, already referred to, of the different organs of the animal functions having their distinct periods of waking and rest, would induce us to suppose, that it ought not to be always equally profound and durable: yet there are individuals whose sleep is nearly complete throughout; but they are not many. The previous occupation of the sleeper exerts great influence. If it have been of a fatiguing nature, all the faculties rest equally long and soundly; but if the fatigue extend beyond the due point, a degree of excitability of the brain is left which renders it extremely liable to be aroused. In this way we understand why dreams should bear upon subjects that have long occupied the mind in its waking state; the tension of the mind on those subjects having left considerable excitability, as respects them, and a disposition to resume them under the slightest irritation. The presence or absence of irritants - external or internal - exerts likewise a great effect on the soundness of sleep, and the formation of dreams. The stillness

4 Human Physiology, p. 231, Lond. 1842.

of the night and the absence of light are hence favourable to repose: the position, too, must be one devoid of constraint; and the couch soft and equable, and especially such as the individual has been accustomed to use. Sleep is impracticable in a badly made bed; and every one must have experienced the antisoporific influence of a strange bed, the arrangement of which, as to size, pillows, &c., differs from that to which he has been habituated. It is not, however, by external irritants that the sleep is usually disturbed. The state of the system itself may react upon the brain, and give occasion to broken sleep, and to dreams of the most turbulent character. Irritations, existing in the viscera, are frequently the cause of dreams, - in children more especially; and a hearty supper, especially if of materials difficult of digestion, may bring on the whole train of symptoms that characterize nightmare. In like manner, any thing that impedes the action of the functions of respiration, circulation, &c., may occasion the wildest phantasies. All these internal impressions are more vividly perceived for the reasons already stated. The nervous system is no longer excited by the ordinary impressions from the external senses; and if the internal impressions be insufficient to prevent sleep altogether, they may

During this incomplete kind of sleep, the external sensations are not wholly at rest; particularly that of touch or tact, which, as it is the last to sleep, is the first to awake. Impressions, made on it, may excite the most exaggerated representations in the brain, in the shape of dreams. The bite of a flea appeared to Descartes the puncture of a sword: an uneasy position of the neck may excite the idea of strangulation: a loaded stomach may cause the sleeper to feel as if a heavy weight, - a house or castle, or some powerful monster, - were on his stomach. A person, having had a blister applied to his head, dreamed that he was scalped by a party of Indians. Moreau de la Sarthe gives the case of a young female, who, from the application of her cold hand against her breast, when asleep, dreamed that a robber had entered her apartment and had seized hold of her. Galen dreamed that he had a stone leg, and, on waking, found that his own was struck with paralysis. Mr. Dugald Stewarta gives a similar case, to show how an impression made upon the body, during sleep, may call up a train of associated ideas, and thus produce a dream. A gentleman, (Dr. Gregory,) who, during his travels, had ascended a volcano, having occasion, in consequence of indisposition, to apply a bottle of hot water to his feet when he went to bed, dreamed that he was making a journey to the top of Mount Ætna, and that he found the heat of the ground almost insupportable. Sir Walter Scotte mentions an analogous instance, which was told him by the nobleman

<sup>\*</sup> Elements of the Philosophy of the Human Mind, i. 335, 3d edit. Lond. 1808.

b Abercrombie's Inquiries concerning the Intellectual Powers, Amer. Edit. p. 216,

New York, 1832; and Elliotson's Human Physiology, p. 625, Lond. 1840.

<sup>c</sup> Letters on Demonology and Witchcraft, Amer. Edit. p. 49, New York, 1830.

546 SLEEP.

concerned. He had fallen asleep, with some uneasy feelings arising from indigestion, which brought on the usual train of visionary terrors. At length, they were all summed up in the apprehension, that the phantom of a dead man held the sleeper by the wrist, and endeavoured to drag him out of bed. He awoke in horror, and still felt the cold dead grasp of a corpse's hand on his wrist. It was a minute before he discovered that his own left hand was in a state of numbness, and that with it he had accidentally encircled his right arm. If, again, the organ of hearing be wakeful, the dreamer may hear an individual speak to him and may reply; so that occasionally secret thoughts and feelings may be elicited. The author has himself replied several times connectedly in this manner; and he has been able to lead on others, especially children,—whose sleep is often interrupted by the existence of irregular internal impressions, — to answer a few times in the same way.

It would seem, that on the loss of any one sense, the dreams, after a lapse of time, will not be referable to it. Darwin has given many instances of this. After blindness had affected certain persons, they never dreamed that they saw objects in their sleep; and a deaf gentleman, who had talked with his fingers for thirty-years, invariably dreamed of finger-speaking, and never alluded to his having dreamed of friends having conversed orally with him.a

In the explanation of the cause of dreaming, we have the most plausible application of the theory of Gall regarding the plurality of organs in the brain. Every explanation, indeed, takes for granted, that certain faculties are suspended whilst others are active. Gall's viewb is, that, during sleep, particular organs of animal life enter into activity; and hence, that the perceptions and ideas, which depend on these organs, awake; but, in such case, their activity takes place without any influence of the will; - that when one organ only is in activity, the dream is simple: the dreamer caresses the object of his affection; he hears melodious music, or fights his enemies, according as this or that organ is exercising its functions; — that the greater the number of organs in activity at the same time, the more confused or complicated will be the dream, and the greater the number of extravagancies; - that, when the organs are exhausted by watching and labour, we generally do not dream during the first hours of sleep, unless the brain is extremely irritable; but, in proportion as the organs get rid of their fatigue, they are more disposed to enter into activity, and hence, near the time for waking, we dream more and with greater vivacity. "Dreaming, consequently," he concludes, "is only a state of partial waking of animal life; or, in other words, an involuntary activity of certain organs, whilst others are resting."c

In many respects, the state of the mind, during dreaming, resembles that in the delirium of fever, as well as in insanity. The

Dendy, The Philosophy of Mystery, p. 228, Lond. 1841.

Bur les Fonctions du Cerveau, ii. 506, Paris, 1825.

See, also, Carmichael, in Transactions of the College of Physicians in Ireland, ii. 48.

imagination and memory may be acting with unusual vivacity, whilst the perception or the judgment may be most erroneous; at times, the perception being accurate and the judgment suspended, so that the individual may be most incoherent; at others, the perception being inaccurate and the judgment right, so that the individual may reason correctly from false premises. As in dreams, too, the delirious may have their ravings modified by impressions made on the external senses. Sir Walter Scotta cites the case of a lunatic, confined in the Infirmary of Edinburgh, whose malady had assumed a gay turn. The house, in his idea, was his own, and he contrived to account for all that seemed inconsistent with his imaginary right of property: - there were many patients in it, but that was owing to the benevolence of his nature, which made him love to relieve distress. He went little, or rather never, abroad, but then his habits were of a domestic and rather sedentary nature. He did not see much company, but he daily received visits from the first characters in the celebrated medical school of the city, and he could not, therefore, be much in want of society. With so many supposed comforts around him, with so many visions of wealth and splendour, one thing alone disturbed his peace. "He was curious," he said, "in his table, choice in his selection of cooks, had every day a dinner of three regular courses and a dessert, and yet somehow or other, every thing he ate tasted of porridge." The cause of this was, that the lunatic actually ate nothing but this at any of his meals: and the impression made upon his palate was so strong as to modify his delusion.

#### 2. WAKING DREAMS.

Nearly allied to dreams, in its physiology — or more properly, perhaps — pathology, is the subject of hallucinations, spectral illusions, or waking dreams, in which the mind may be completely sound, and yet the cerebral or percipient part of the brain, concerned in the senses, be so deranged as to call up a series of perceptions of objects, which have no existence except in the imagination. Such hallucinations are constant concomitants of insanity, delirium, and dreaming; but they may occur, also, when the individual is wide awake, and in the full possession of reasoning powers: he may see the phantasm, but, at the same time, totally disbelieve in the existence of any extraneous body. The most common illusions of this kind affect the senses of sight and hearing.

It has fallen to the lot of the author to meet with some singular and serious cases of this affection; where, for example, the individual, wide awake, has heard the doors of his house violently slammed, his windows thrown up and down, the bells set a ringing, himself subjected to personal violence; yet there has been no slamming of doors, no throwing up and down of windows, no ringing of bells, no personal violence; the whole has been an illu-

sion, a waking dream, and of this no one has been more entirely aware than the sufferer himself. A few years ago, the author was consulted by a most respectable citizen of Virginia, regarding his state of health as well as an illusion of this nature. He was one of the Board of Visiters at West Point, where his duty called him to inspect the demonstrations by the pupils on the black-board. For months after his return to Virginia, he saw the black-board with its demonstrations constantly before him. He had previously experienced an attack of paralysis, and, when he applied to the author, he was labouring under marked evidences of predisposition to a farther attack of encephalic mischief, of which the illusion in question was doubtless one. One of the most impressive cases, however, is that of Nicolai, the eminent bookseller of Berlin, which has been detailed by Drs. Ferriara and Hibbert, and by Dr. Haslame and Mr. Mayo.d Nicolai laid his case before the Philosophical Society of Berlin. He traced his indisposition, for it was manifestly such, to a series of disagreeable incidents that had befallen him. The depression, thus induced, was aided by the consequences of neglecting a course of periodical bleeding to which he had accustomed himself. This state of health brought on a disposition to spectral illusions, and, for a time, he was regularly haunted by crowds of persons entering his apartment, and addressing him or occupied solely in their own pursuits, until, as his health was restored, they gradually disappeared, and ultimately left him entirely. Yet Nicolai, who was a man of unusually strong intellect, was throughout satisfied, that they were mere hallucinations.

The cases of this kind, now on record, are many and curious. Every one engaged in extensive practice, or in frequent communion with the world, must have seen or heard of them. Some, of a deeply interesting character, are detailed by Sir David Brewster, Dr. Abercrombie, and Dr. Macnish, but there are none more extraordinary than some that have been related by Sir Walter Scott. They are signal examples of the illusions that may occur during even our waking moments; and they may, doubtless, account for some of the stories of apparitions, of which so many are upon record. In the hypochondriac, we meet with all kinds of hallucination, and it is one of the most striking of the symptoms of every variety of insanity; but, in the cases referred, to, notwithstanding the constancy and permanency of the illusion,

\* An Essay towards a Theory of Apparitions, Lond. 1813.

b Sketches of the Philosophy of Apparitions, Edinb. 1825.

d Outlines of Human Physiology, 3d edit. p. 213, Lond. 1833. e Letters on Natural Magic, Amer. Edit. p. 42, New York, 1832.

Medical Jurisprudence as it relates to Insanity, in Cooper's Tracts on Medical Jurisprudence, p. 302, Philad. 1819.

f Inquiries concerning the Intellectual Powers, and the Investigation of Truth, Amer. Edit. p. 282, New York, 1832.

Fhilosophy of Sleep, Amer. Edit. p. 214, New York, 1834.

Letters on Demonology, &c., Amer. Edit. p. 34, New York, 1830.

the individual himself has been satisfied that the whole affair had no real existence. Had he believed in the existence of the phantom, and acted from a conviction of its reality, he might, with propriety, have been deemed insane, quoad hoc. An instance of this kind is told in the Memoirs of the Count Maurepas, of one of the princes of the house of Bourbon, who supposed himself to be a plant, and, having fixed himself in the garden, called upon his servant to come and water him. His belief argued unsoundness of mind, yet, even here, the hallucination, we are told, appeared to be confined to this subject.

In youth, when the imagination is extremely vivid, we can call up images in the mind at pleasure, varying them as we may think proper. In the nervous, the delicate, and the imaginative, uneasy sensations can be experienced, when and where the individual wishes. After sedentary habits, long continued, the author has been able to experience, at will, pain in any part of the system, and to make it shift at pleasure from one organ to another.

In the cases of hallucination, referred to above, as well as in every other kind, the cerebral part of the organ of sense is directly or indirectly excited into action; - often by disease of the brain, or of some distant organ which reacts upon it. Hence it occurs as a precursor of apoplexy, epilepsy or other cerebral affection, or it may accompany, or be aggravated by, disorder of the digestive function. It has been seen, that although the passions or emotions are cerebral phenomena, they are felt in the nutritive organs; and we can understand, how a disordered state of those organs may react upon the brain, and call up all kinds of illusions; generally during sleep, but at times even during our waking moments. In this way, we account for the frightful dreams that follow an overloaded stomach, or that accompany impeded respiration or circulation. One of the most distressing symptoms of hydrothorax or water in the chest, which interferes more or less with both these vital functions, is the disturbed sleep, and the frightful sense of impending danger, which nightly distress the unfortunate sufferer.

It appears, then, that in all cases of hallucination, occurring in those of sound or diseased mind, asleep or awake, the encephalic or percipient part of the organ of the sense concerned is irresistibly affected, so as to call up the memory of objects, or to form others, which have no existence except in the imagination; but all this is accomplished without any impression being made upon the external senses from without, even when these senses appear to be most actively exercised. In dreams, this must manifestly be the case. We see a friend long since dead; we parade the streets of a town, which we have never visited; and see, hear, feel and

<sup>\*</sup> See, on this interesting subject, the valuable work of Dr. Conolly on Insanity,

b Rush's Lecture on Medical Jurisprudence, Philad. 1811; and Cooper's Tracts on Medical Jurisprudence, p. 324, Philad. 1819.

550 SLEEP.

touch the different objects. All this must be encephalic; and not less certainly is it the case in the hallucinations of insanity, or in those that occur in the waking condition. The object which we see is not in existence, yet it is a regularly defined creation; a cat in one instance, a gentleman-usher in another, and a skeleton in a third. It cannot depend upon any depraved condition of the organ of sense, as in such case the representation of the mind would be amorphous, irregular, or confused; not a complete metamorphosis, as is invariably the case. Yet we are surprised Sir Walter Scotta should state, that he thinks "there can be little doubt of the proposition, that the external organs may, from various causes, become so much deranged as to make false representations to the mind; and that, in such cases, men, in the literal sense, really see the empty and false forms, and hear the ideal sounds, which in a more primitive state of society, are naturally enough referred to the action of demons or disembodied spirits. In such unhappy cases, the patient is intellectually in the condition of a general, whose spies have been bribed by the enemy, and who must engage himself in the difficult and delicate task of examining and correcting, by his own powers of argument, the probability of the reports, which are too inconsistent to be trusted to."

The explanation is poetic, but manifestly untenable.

A theory, which has been offered to account for the various spectral illusions, occurring in any of the modes we have mentioned, is - that, in all the organs of sense, the mind possesses the power of retransmitting, through the nervous filaments, to the expansions of the nerves that are acted upon by external objects. impressions, which these nerves have previously transmitted to the brain, and, that the vividness of the retransmission is proportional to the frequency with which the impressions have been previously transmitted; that these reproduced impressions are in general feeble in the healthy state of the body, though perfectly adapted to the purposes for which they are required; but, in other states of the body, they appear with such brilliancy as to create even a belief in the external existence of those objects from which the impressions were originally derived. "When the mind," says a writer on this subject, "acquires a knowledge of visible objects it is by means of luminous impressions, conveyed to the sensorium from each impressed point of the retina, through the corresponding filaments of the optic nerve; and when the memory is subsequently called upon, by an act of the will, to present to us an object, that has been previously seen, it does it by retransmission along the same nervous filaments, to the same points of the retina. In the first case, when the presence of the luminous object keeps up a sustained impression upon the nervous membrane, the filaments, which transmit it to the brain are powerfully excited; but, in the process of retransmission by an effort of enemory, the action of the nervous filaments is comparatively feeble, and the resultant impression on the retina faint or transient. When the memory, however, is powerful, and when the nervous filaments are in a state of high excitability, the impression becomes more vivid; and, as in the case of spectral illusions, it has the same strength and distinctness, as if it were produced by the direct action of luminous rays. In one case, the result of the impression and its retransmission to the retina is a voluntary act of the mind, but, in the other, it is involuntary, the controlling power being modified or removed, or the nerves being thrown into a state of easy excitation by some un-

healthy action of the bodily organs."

According to this view, it is indispensable, that the perception, in every case of illusion, shall be referred to the nerves of the organ by which such perception is ordinarily effected; to the retina, if vision be concerned; to the auditory nerve, if audition; and so on. But this retransmission along the nerves appears to us to be wholly unnecessary. When an impression is made upon a sensitive surface, as we have elsewhere shown, sensation is not accomplished, until the impression has been conveyed to the brain by an appropriate organ, and the brain itself has acted; and if we interfere in any manner with the cerebral part of the function, perception is not effected. From the moment, however, that the action of the brain has taken place, the idea formed can be recalled by the exercise of memory; and we have no doubt, that this could take place although the eyes were extirpated. The memory might call up previous perceptions, when the functions of the retina are entirely destroyed. Were it otherwise, it would be impossible for those, who have lost their sight from paralysis of the retina, of which many cases are constantly occurring, to call up any of the scenes and images, of which the brain took cognizance prior to the supervention of their blindness. In dreams, too, we exert every one of the senses; some with the greatest activity. We see, hear, taste, smell, and feel; and, in addition to this, walk, run, fly, and execute the ordinary acts of life not only without apparent difficulty, but with a facility that surprises us. Yet can we suppose, that, in all these cases, the feeling is actually produced by retransmission along the nerves of the organ to which it is referred? It has been asserted, that when examination is carefully made, it will be found, that the images, recalled by the memory, follow the motions of the head and of the eye; but, that this is not the case during sleep is manifest. The individual may remain precisely in the same position, and yet he will seem to move about in all directions in his dreams; will appear to see objects behind as well as before him; and in situations towards which it is impossible that the motions of his head and eye should be directed. Even in most of the illusions of our waking hours, the remark ought to be reversed. The encephalic action is the first of the links in the chain of phenomena; and the motions of the head and the eye follow the images recalled by the memory. When the unfortunate subject of one of the cases of hallucination given by Sir Walter

552 SLEEP.

Scott, saw the gentleman-usher preceding him into company, and circulating among the assembled guests, — as well as when he observed the skeleton at the foot of his bed, — the perception, owing to disease, had so completely taken possession of a part of the encephalic organ of vision, that the idea was constantly in the mind; and volition being actively exercised, the head and the eye were directed towards the phantasm. Yet the perception was not so powerful, as to preclude the reception of impressions from without, as was shown by the skeleton seeming to be shut off by the body of the physician, so that the skull only was seen peering above his shoulder.<sup>a</sup>

Another fact, which shows, that the whole phenomenon may be entirely encephalic, is the occurrence, familiar to the operative surgeon, of a patient, whose lower limb has been amputated, complaining of an uneasy sensation, as of itching, in a particular toe, and in a particular part of a toe. This is, at times, a symptom of an extremely distressing character. It is obviously impossible, that, in such case, there can be any external impression made on the part to which the feeling is referred; or that any retransmission can occur from the brain; the limb having been removed from the body. Broussais asserts, that if a person tells you he suffers in a limb which he no longer has, it is because he experiences irritation in the extremities of the divided nerve, but this, in no respect, removes the difficulty. The sensation is referred to a part which has

no existence except in the imagination.

But, to return to sleep. We have said, that the object of sleep is to repair the loss, which the nervous system has sustained, during the previous condition of waking. This may, consequently, be regarded as the great exciting cause of sleep; but we have seen, also, that certain states of the mind may postpone the usual period of its recurrence. If, indeed, we allow the attention to flag, and suspend the due exercise of volition, sleep can be indulged at almost any hour of the day. In the same manner, any monotonous impression, or action of the brain in thought; the rocking of a cradle, or the song of the nurse to a restless child; the murmurs of a bubbling brook, &c., may soothe us to rest. A like effect is produced by substances, as narcotics, which, by a specific action on the nervous system, prevent the ordinary sources of irritation from being appreciated, as well as by certain morbid affections of the brain, - compression, concussion, inflammation, &c. In these cases, however, the sleep is morbid, and is an evidence of serious mischief, - often of fatal disease; whilst true sleep is as natural as the waking state, and is always -

"Man's rich restorative; his balmy bath,
That supples, lubricates, and keeps in play
The various movements of that nice machine,
Which asks such frequent periods of repair!"

Young's Night Thoughts.

<sup>·</sup> Sir W. Scott, Letters on Demonology and Witchcraft, Amer. Edit. p. 35, New York, 1830.

Yet Haller, Hartley, and numerous others have supposed, that natural sleep is dependent upon an accumulation of blood or other fluids in the vessels of the head pressing upon the brain, and thus impeding its functions. In support of this opinion, it is asserted, that all the phenomena which attend the sleeping state, seem to prove a determination of blood to the head. The face is flushed; the head is hotter; the skin more moist; and it is generally during the night, or when first awake, that bleeding from the nose and apoplexy take place; the frequency of erection during sleep is affirmed to be owing to the pressure exerted on the cerebellum, which, in the theory of Gall, is the encephalic organ of generation; and, lastly, it is argued, that narcotics, and vinous and spirituous liquors produce sleep by causing a similar congestion of blood within the cranium. The case, by no means unique, of the beggar whose brain was exposed, and in whom a state of drowsiness was induced when the brain was pressed upon, which could be increased by increasing the pressure, until at length he became comatose, has also been cited by Hartley and others. But all these are cases of morbid suspension of the animal functions, and are no more to be assimilated to true sleep, than the drowsiness, which Flourense found to prevail in his experiments on animals when the cerebral lobes were removed.

The believers in the hypothesis, that congestion of the vessels of the brain is the cause of sleep, consider that the heaviness and stupor, observable in those who indulge too much in laziness and sleep, are owing to long-continued pressure injuring the cerebral organs. Other physiologists have assumed the opposite ground, and affirmed, that during sleep the blood is distributed to the brain in less quantity, and is concentrated in the abdomen, to augment the action of the nutritive functions; d whilst Cabanise holds, that during sleep there is a reflux of the nervous power towards their source, and a concentration in the brain of the most active principles of sensibility. On all these topics our ignorance is extreme. We know nothing of the state of the encephalon in sleep. Its essence is as impenetrable as that of every other vital function. Dr. Bostockf asserts, that it is not more beyond our grasp than the other functions of the nervous system. This we admit : he has, indeed, afforded us in his own work indubitable evidences of our utter want of acquaintance with the essence of all those functions. The state of sleep is as natural, as instinctive, as that of waking; both are involved in mystery, and their investigation, as Mr. Du-

Element. Physiolog. xvii. 3. On Man, p. 45, Lond. 1791.

<sup>·</sup> Expériences sur le Système Nerveux, Paris, 1825.

d Blumenbach, Instit. Physiol., and in Elliotson's Human Physiology, p. 610, Lond.

Rapport du Physique et du Moral de l'Homme, Paris, 1802. See, also, Adelon, Physiol. de l'Homme, 2de édit. ii. 292, Paris, 1829; and Gall, Sur les Fonctions du Cerveau, ii. 503, Paris, 1825.

Elementary System of Physiology, 3d edit. p. 815, Lond. 1836.

VOL. II. - 47

gald Stewarta has suggested, is probably beyond the reach of the human faculties.

## 3. REVERY.

Revery has been considered to resemble sleep, and, in its higher grades, to be not far removed from the condition of somnambulism. It is characterized by the attention or volition being directed so intently towards particular topics, during wakefulness, that the impressions of surrounding objects are not appreciated. Various grades of this condition of the mind may be traced, from the slightest degree of absence or brown study to a state of total abstraction in which the attention is entirely wound up, and riveted to a particular subject. Most persons must have experienced more or less of this, when any subject of severe study, or any great gratification, anxiety, or distress has strongly occupied the mind. If engaged in reading, they may follow every line with the eye; turn over leaf after leaf, and at length awake from the revery, which had occupied the imagination, and find, that not the slightest impression has been made on the mind, by the pages, which the eye had perused, and the hand had passed over. If walking in a crowded street, they may have proceeded some way under the influence of revery, moving the limbs as usual, performing various acts of volition, winding safely among the passengers, avoiding the posts and other obstacles, yet so exclusively occupied by the conceptions of the mind, as to be totally unconscious of all these acts of their volition, and of the objects they have passed, which must necessarily have impressed their senses so as to regulate those actions, but, owing to the attention having been bent upon other topics, the perceptions were evanescent. In elucidation of the power of a high degree of revery to render an individual torpid to all around him, the case of Archimedes, at the time of his arrest, has been quoted by writers. When the Roman army had at length taken Syracuse by stratagem, which the tactics of Archimedes had prevented them from taking by force, he was shut up in his closet, and so intent on a geometrical demonstration, that he was equally insensible to the shouts of the victors, and the outcries of the vanquished. He was calmly tracing the lines of a diagram, when a soldier abruptly entered his room, and clapped a sword to his throat. "Hold, friend," says Archimedes, "one moment, and my demonstration will be finished." The soldier, surprised at his unconcern at a time of such extreme peril, resolved to carry him before Marcellus; but as the philosopher put under his arm a small box full of spheres, dials, and other instruments, the soldier, conceiving the box to be filled with gold, could not resist the temptation, and killed him on the spot.a

It is to the capability of indulging to the necessary extent in this kind of mental abstraction, that we are indebted for the solution of

b Liv. l. xxxv. c. 3.

a Elements of the Philosophy of the Human Mind, i. 327, 3d edit., Lond. 1808.

every abstruse problem, relating to science or art, and for some of the most beautiful conceptions of the poet. From indulgence, however, in such abstractions, a habit is often acquired, which may be carried so far as to render the individual unfit for society, and to give him a character for rudeness and ill-breeding, of which he may be by no means deserving. Some most amiable and estimable men have, from long habits of abstraction, contracted the disease (aphelxia), as Gooda has constituted it, and have found the cure tedious and almost impracticable; at times, indeed, it appears to have terminated in mental alienation. The difference between this state and that of sleep is, that the attention and volition are here powerfully directed to one object, so as to be torpid to the impressions of extraneous bodies; whilst sleep is characterized by a suspension or diminished exercise of these faculties.

# CHAPTER III.

#### CORRELATION OF FUNCTIONS.

The wonderful and complicated actions of the frame are variously correlated to accomplish that astonishing harmony, which prevails in the state of health, as well as to produce the varied morbid phenomena,—often at a distance from the part originally diseased,—which characterize different pathological conditions. It is not, therefore, simply as a physiological question, that the study of the correlation of functions interests the medical inquirer. It is important to him in the study of every department, which concerns the doctrine of the healthy or diseased manifestations, and the modes adapted for the removal of the latter.

These correlations may be of various kinds; — mechanical, in which the effect exerted is entirely of a mechanical character; functional, in which the action of one organ is inseparably united to that of another, to accomplish a particular object; and sympathetic, in which there is no physical action or direct catenation of functions; but where an organ, at a distance from one affected, is excited to regular or irregular action in consequence of the condition of the latter.

## 1. MECHANICAL CORRELATIONS.

In the description of the different functions, numerous opportunities occurred for showing the influence which organs, in the immediate vicinity of each other, may mutually exert so as to modify their functions. The action of the muscles, — particularly those that contract the larger cavities, as the abdomen and thorax, — on the parts with which they come in contact, must be entirely mechanical. In this way, the diaphragm and the abdominal muscles

A Physiological System of Nosology, cl. iv. ord. 1, gen. v.

act in vomiting and defecation. During the operation of bloodletting, the flow of blood can be augmented by moving the muscles of the hand; and it is probable, that the constant motion of the muscles of respiration impresses a succussion on different organs, which may aid them in accomplishing their functions, although the effect of this is doubtless exaggerated. Every change of position, either of the whole body or of a part, has likewise some effect in modifying the actions performed by it or by neighbouring organs, although such effect may not be easily appreciable. A similar case of mere mechanical influence, which seems to be important to the proper action of certain organs, is exhibited in the pulsation of the different arteries. It has been seen, that a succussion is in this way given to the brain, which appears to be necessary to it; for, if this source of stimulation be in any manner withdrawn, fainting is induced. Perhaps, however, the strongest case, that can be offered, of modification of function by mechanical causes, is that of the gravid uterus, which, by its pressure, gives rise to numerous symptoms in other organs, that are often the source of annovance during gestation.

## 2. FUNCTIONAL CORRELATIONS.

The functional correlations or synergies are of much more moment to the physiologist and pathologist. Many of these have also been described in the preceding history: a brief notice of them will be all that is now requisite. For the maintenance of the healthy function we know that certain conditions are necessary, and that if these be materially modified, in the whole or in any part of the body, disease and death may be the result, even although the derangement may, in the first instance, concern only an apparently unimportant part of the frame, - the affection, by correlation, spreading gradually to more and more essential organs and functions, until the disorder is ultimately too great to allow of a continuance of the vital movements. In this respect, man differs from an ordinary piece of mechanism, in which the various parts are so adapted to each other as to produce a certain result. If one of these parts be destroyed, the whole machine may have its motion arrested; but the effect is owing to the destruction of one part only, the others remaining sound; whilst death, or the stoppage of the living machine, does not necessarily follow the destruction of any except a few essential organs, and is generally owing to the derangement of many. We shall find, indeed, that except in cases of sudden death, it is extremely difficult to say which of the three truly vital organs has first ceased to act; and that in all such cases death begins in one or other of the organs essential to vitality, and soon extends to the rest.

The essentially vital organs are those of respiration, circulation, and innervation; but the great use of respiration is to change the blood from venous to arterial; in other words, to induce a conversion in it by its passage through the lungs, without which it

would be inadequate for the maintenance of life in any organ; and the object of the circulation is, to distribute it to the various parts of the frame as the grand vivifying and reparatory material. If, also, the organs of innervation be destroyed, the nervous influence is no longer conveyed to the different parts of the frame; and as the presence of this influence every where is indispensable, the functions may cease from this cause; so that we may regard, as essential elements to the existence of the frame and of every part of the frame, the proper supply of arterial blood and of the nervous influence. In the production and distribution, however, of these agencies, a number of functions is concerned, giving rise to the correlation, which is the object of our present inquiry. If, in any manner, the blood do not meet with due aëration, as in ordinary cases of suffocation, death supervenes, in the order elsewhere described; and if a slight degree of aëration be accomplished, but still not enough for the necessities of the system, instead of suffocation, the individual dies more gradually; the functions fail in the same order; dark blood circulates through all the textures; hence lividity, especially of those parts where the cuticle is extremely thin, as of the lips, and wherever the mucous membranes commingle with the skin; and the blood gradually becomes inadequate to keep up the action of the brain and nervous system generally, as well as to stimulate the heart, and the individual gradually expires. If, again, the blood, although properly converted in the lungs, be not duly distributed to the organs, owing to the failure of the circulatory powers, - either from direct or indirect causes, - the organs exhibit their correlation in the same manner, and syncope or fainting, or positive death, may be induced. Often, however, the stoppage of the action of the heart is but for a short time. Owing to some painful impression, sudden emotion, or other cause, the organ ceases to contract, either suddenly, - when the persons falls down as if deprived of life, - or gradually when the connexion of the different functions, and the order in which they fail, are manifest. Of this kind of what the surgeon calls - morbid sympathy or constitutional irritation, we have a good example in the effect of a trifling operation upon a delicate, and often upon a strong, individual. Bleeding will sometimes induce fainting, both directly, by the abstraction of fluid from the vessels, so that the brain may cease to act; and indirectly, when the quantity removed cannot be presumed to have exerted any influence. Some, indeed, will faint from the slightest puncture and loss of blood, or even from the sight of that fluid. In these last cases, if the syncope come on gradually, a feeling of great anxiety and oppression, occasionally of vacuity, exists in the epigastric region; and perceptions become confused, the sight obscured, tinnitus aurium and dizziness supervene, the respiration is embarrassed, the face pale, the extremities cold, and the different parts of the body are covered with a cold, clammy sweat, until, ultimately, loss of sensation and motion supervenes, and the individual is temporarily dead; from which state he soon recovers, in the generality of cases, provided he be kept in the recumbent posture, so that the blood may readily pass to the brain. On other occasions, the heart will not cease its pulsations, but will continue to send blood, in undue quantity, to the brain, so that all the above symptoms may ensue, except the temporary privation of vitality. In consequence of the severe pain induced by a displacement of two of the bones of the wrist, by a fall from a carriage, the author remained a considerable time incapable of sight, and at the same time suffering from great anxiety, yet consciousness and the action of the heart never ceased, as in complete syncope. The third vital function, - that of innervation, - when suspended or diminished, draws on a train of pathological phenomena, in the order described under the head of Death; suspending respiration and circulation suddenly, if the cause applied be sufficient; more gradually, and with the symptoms characterizing apoplexy or compression of the brain, if the cause act in a minor degree. All the three vital functions are consequently correlative, and so intimately associated, that if a malign influence act upon one, the effect is speedily extended to the other.a

Owing to the necessity for the blood possessing certain attributes, the most important of which are obtained by its circulation through the lungs, we can readily understand, that if the functions of nutrition be not properly exerted, the composition of that fluid may be imperfect, and disorder take place in various parts of the frame from this cause. Thus, if digestion or the formation of chyle be not properly executed, the blood is not duly renovated, and may be so far impoverished, that the play of the functions is interfered with. We have elsewhere shown, that if omnivorous man be restricted to one kind of diet he will fall off, and become scorbutic, and the affection will be removed by allowing him diet of another kind; - vegetables, if animal food have induced it, and conversely. Enlarged mesenteric glands, consequent, or not, on inflammation of the mucous membrane of the intestine, and the latter affection itself, are cases which may interfere with chylosis, and consequently with the constitution of the blood. In like manner, if nutrition and the various secretions be not duly performed in the tissue of the organs, and, especially, if the great depurations be obstructed, the blood may suffer, and although the due change from venous to arterial may be effected in the lungs, its character may not be such as to adapt it for the healthy execution of the various functions.

The humorists assigned too much importance to the condition of the humours in the production of disease; the solidists, on the other hand, have denied it almost all agency. The medium between these exclusionists is probably the nearest to nature. The

<sup>\*</sup> Richerand's Elémens de Physiologie, 13ème édit. par Bérard ainé, § exevii. Bruxelles, 1837.

solitary fact of black blood being unfit to maintain the perfect and continued vitality of any organ sufficiently exhibits its influence. How the arterial blood exerts its agency, independently of its action as a fluid of nutrition, is beyond our knowledge. It appears to effect a necessary action of stimulation, but in what manner, or on what element, we know not: probably, however, its chief influence may be on the nervous tissue, as the privation of arterial blood soon occasions the cessation of the brain's action.

In the higher classes of animals, innervation is dispensed from three great centres, - the encephalon, the spinal marrow, and the great sympathetic.a The presidency, however, may be fairly assigned in man and in the higher animals, to the first of these. If it fail, death soon becomes general. This, however, is liable to great variation in different animals, and likewise in different functions. In man, if the nervous supply be cut off from any part, the part dies. Physical integrity, continuity, and a due supply of arterial blood, are necessary to the proper exercise of the nervous power. In the former part of this work, the wonderful resistance to death which characterizes the amphibia, and the comparative independence of each portion of the body, in some of the lower order of animals, were pointed out. The polypus may be divided into numerous pieces, yet each may constitute of itself a distinct animal. The snail, after decapitation reproduces the head; and a similar reparatory power is possessed by other animals. We have elsewhere seen, that volition is seated lower in the inferior than in the superior orders of animals; and that in man it is chiefly, - some say, wholly, - restricted to the encephalon. It appears, likewise, that the dependence of the rest of the nervous system on the great nervous centres is less in young than in old animals. Edwards regarded the new-born child as resembling, in many respects, the cold-blooded animal; and Redi, Rolando and Flourens, and Legallois found, that the tenacity of life, after decapitation, was much greater the nearer to birth. The functions also differ with regard to their dependence upon the encephalon. Disease may attack the animal functions and suspend them for a considerable length of time, - as in apoplexy, - before the organic functions are interfered with. This is a topic, however, which will be discussed under the head of Death.

A gifted preceptor of the author, — M. Béclard, b — has defined life to "consist essentially in the reciprocal action of the circulation of the blood and innervation; death always following the cessation of such reciprocal action." But this conclusion is applicable only to animals; although both circulation and innervation are admitted in the vegetable by some physiologists. Legallois, from his experiments, deduced the unwarrantable inference, that "life

Adelon, Physiologie de l'Homme, iv. 146, 2de édit. Paris, 1829; and art. Innervation, in Dict de Méd. lère, édit. tom. xii.

b Elémens d'Anatomie Générale, 2de édit. Paris, 1827; or Togno's Translation p. 106, Philad. 1830. Cur le Principe de la Vie, Paris, 1812.

is owing to an impression made by arterial blood on the brain and spinal marrow, or to the principle, which results from this impression;"—a definition, which would exclude the numerous animals of the lower classes, as well as vegetables, which are deficient in both brain and spinal marrow. The conclusion of Béclard is, perhaps, the limit to our knowledge on this subject. Yet some have endeavoured to discover which of the two functions,—circulation or innervation,—holds the other in domination. They, who consider the nervous substance to be first formed in the fætus, ascribe the supremacy to it; whilst the believers in the earlier formation of the sanguiferous system look upon it as the prime agent. We know no more than that both

"Maintain,
With the mysterious mind and breathing mould
A co-existence and community."

In every important function of the body we find this correlation or catenation of organs existing; all working to one end, and all requisite for its perfect accomplishment. How many organs, for example, are required to co-operate in the elevated function of sensibility! The encephalon, the seat of thought, receives, by the external senses, the various impressions which act upon them from without, and, by the internal sensations, such as arise in the economy, and are generally the indexes of the physical necessities or wants. The intellectual and affective faculties enable us to appreciate the various objects that occasion our sensations, and indicate our social and moral wants: under their direction, volition is sent out, which acts upon the various muscles, and produces such movements as may be required for carrying into effect the suggestions of the mind. Between all these acts there is the closest catenation. In like manner, we observe the correlation between the animal, and the nutritive, and reproductive functions. The internal sensation of hunger suggests to the mind the necessity for a supply of aliment; the external senses are called into action to discover the proper aliment; when discovered, it is laid hold of by muscular movements under the direction of volition, is subjected to various voluntary processes in the mouth, and then passed on, by a mixed voluntary and involuntary action into the stomach. In like manner, the desire for sexual intercourse may be excited in the mind through the organs of vision or touch; the organs of generation are aroused to action, and the union of the sexes is accomplished by the exertion of muscles thrown into contraction by volition. The same catenation is exhibited after a fecundating copulation; menstruation, which was previously performed with regularity is now arrested: the breasts become developed; milk is formed in them, and, whilst the female suckles her child, unless the period is unusually protracted, the non-existence of the menstrual functions continues.

Almost all the phenomena of disease are connected with this

correlation of functions. Derangement takes place in one organ or structure of the body, and speedily all those, that are correlated with it, participate in the disorder. Hence, in part, arises the combination of disordered nervous, circulatory, and secretory function, which characterizes general fever; and the various associated morbid actions that constitute disease in general.

## 3. SYMPATHY.

There is another kind of connexion, which distinguishes the animal body from a piece of ordinary mechanism yet more than those we have considered. In this, owing to an impression made upon one organ, distant organs become affected, without our being able to refer the transmission to mechanical agency, or to the association of functions, which we have just described. This kind of association is called sympathy. A particle of snuff or other irritating substance, impinging on the Schneiderian membrane, produces itching there, followed by a powerful action of the whole respiratory apparatus, established for its removal. The sneezing, thus induced, is not caused by the transmission of the irritation through the intermediate organs to the respiratory muscles; nor can we explain it by the mechanical or functional connexions of organs. It is produced by this third mode of correlation: - in other words, it is a case of sympathy. Again, a small wound in the foot will produce locked jaw, without our being able to discover, or to imagine, any greater connexion between the foot and the jaw than there is between the foot and other organs of the body. We say, that it is caused by sympathy existing between these organs, and, so long as we use the term to signify the unknown cause of these connexions, it is well. It must be understood, however, that we attach no definite idea to the term; that it is only employed to express our ignorance of the agent or its mode of action; precisely as we apply the epithet vital to a process, which we are incapable of explaining by any physical facts or arguments.

Of sympathetic connexions, we have numerous examples in the body; at times, inservient to accomplishing a particular function; but generally consisting of modifications of function produced by the action of a distant organ. Of the sympathetic connexion between the parts of the same organ, for the execution of a function proper to the organ, we have an example in that between the iris and the retina; the former will contract or dilate according to the degree of stimulation exerted by the light on the latter; and the effect is greater when the light is thrown on the retina than when thrown on the iris itself. A similar kind of sympathy exists between the state of the mammæ and that of the uterus, during pregnancy; although this has been frequently referred to ordinary functional correlation or synergy; but the connexion is sufficiently obscure to entitle it to be placed under this division. A singular example of the sympathy between these two organs, soon after

delivery, is the fact of the sudden and powerful contraction which is excited in the uterus, when in a state of inertness, by the appli-

cation of the child to the breast.a

a. Sympathy of Continuity. - This is such as occurs between various parts of membranes that are continuous. For example, the slightest taste or smell of a nauseous substance may bring on an effort to vomit, - the whole of the first passages being unfavourably disposed for its reception. In disease, we have many examples of this kind of sympathy. During dentition, the child is subject to various gastric and intestinal affections. If a source of irritation exist in any part of the intestinal or other mucous membrane, no uneasy sensation may be experienced in the seat of irritation, yet it may be felt at the commencement of the membrane or where it commingles with the skin : - thus, itching at the nose may indicate irritation of the digestive mucous membrane; - itching or pain of the glans penis, stone in the bladder, &c. These facts prove, that, in disease, a sympathetic bond unites the parts concerned, and such is probably the case in health also. We have the same thing proved in the effect produced on the action of glands by irritating the orifices of their excretory ducts. The presence of food in the mouth excites the secretion of the salivary glands, and that of chyme in the duodenum augments the secretion of the liver. In the same manner, a purgative, as calomel, which acts upon the upper part of the intestinal canal, becomes a cholagogue; and duodenitis occasions a copious biliary sccretion. These cases, have, however, been considered by many, to belong more appropriately to functional correlation, as it is presumable, that the propagation of the irritation from the orifice of the excretory duct takes place directly, and along branches of the same nerves as those that supply the glandular organs. It is by this sympathy of continuity that we explain the action of certain medicines. In bronchial irritation, for example, the cough will frequently be mitigated by smearing the top of the larvnx by a demulcent, - the soothing influence of which extends to the part irritated.

b. Sympathy of Contiguity. — A variety of sympathy, differing somewhat from this, is the sympathy of contiguity or contiguous sympathy, in which an organ is affected by an irritation seated in another immediately contiguous to it. The association in action between the lining membrane of the heart and the muscular tissue of the organ has been adduced as an instance of this kind, and chiefly from the experiments of Bichat and Nysten, which showed, that any direct irritation of the muscular tissue of the heart has not as much influence as that of the membrane which lines it. A similar association is presumed to exist between the mucous and muscular coats of the alimentary canal, and the same kind of evidence is adduced to prove that the connexion is sympa-

a Hedenus in art. Brust, (weibliche,) in Encyclop. Wörterb. der Medicin. Wissen schaft, vi. 349, Berlin, 1831; Carus, Lehrbuch der Gynäkologie, Th. ii. 407, Leipz. 1828; and Dr. Rigby, in Lond. Med. Gazette, March, 1831.

thetic.a Other instances of sympathy are, - the convulsive contraction of the diaphragm and abdominal muscles in vomiting consequent on the condition of the stomach, as well as the convulsive action of the respiratory muscles in sneezing, coughing, &c. The general uniformity in the motion of the two eyes has been adduced as an additional instance; but Adelon has judiciously remarked, that the evidence in favour of this view is insufficient. For clearness of vision it is necessary, that the luminous rays should impinge upon corresponding points of the two retines, and should fall as nearly as possible in the direction of the optic axes. For this purpose, the muscles direct the eyes in the proper manner; and subsequently, from habit, the balls move in harmony. We constantly hear, also, a fact adduced from pathology as an instance of sympathy. A molar tooth is lost on one side of the jaw; and it is found, perhaps, that the next tooth which decays is the corresponding molar tooth of the opposite side: - or a tooth has become carious, and we find the one next to it soon afterwards in a course of decay. These have been regarded as evidences of sympathy, - remote and contiguous. This is not probable. The corresponding teeth of the two sides are similarly situate as regards the supply of nerves, vessels, and every anatomical element; and experience teaches us, that the molar teeth - and especially the second great molares - decay sooner than the others. If one, therefore, become carious, we can understand, why its fellow of the opposite side should be more likely to suffer. The opinion, that contiguous teeth are likely to be affected by the presence of a carious tooth, either by sympathy, or by direct contact, is almost universally believed, and promulgated by the dentist. Both views are probably alike erroneous. If the inner side of the second molaris be decayed, we can understand, why the corresponding side of the third should become carious, without having recourse either to the mysterious agency of sympathy, or to the very doubtful hypothesis of communication by contact, - especially as the caries generally begins in-The contiguous sides of the teeth are situate almost identically, as regards their anatomical elements; and, consequently, if a morbid cause affects the one, the other is next likely to suffer, and is very apt to do so. Extracting the diseased tooth prevents this, because it removes a source of irritation, which could not but act in a manner directly injurious on the functions of the tooth next to it.

The fact of the sympathy, which exists between organs of analogous structure and functions, is familiar to every pathologist. That of the skin and mucous membranes is the most intimate. In every exanthematous disease, the danger is more or less dependdent upon the degree of affection of the mucous membranes; and the direct rays of the sun, beaming upon the body in warm climates, induce diarrhæa and dysentery. Acute rheumatism is a disease of the fibrous structures of the joints; but one of its most serious ex-

Adelon, Physiologie de l'Homme, edit. cit., iv. 267, Paris, 1829.

tensions, or metastases, — whichsoever they may be called — is to the fibrous structure of the pericardium. Barthez, a most respectable writer, gives a case of this kind of sympathy from Theden which is inexplicable and incredible. A patient, affected with paralysis of the right arm, applied a blister to it, which produced no effect, but acted on the corresponding part of the other arm. The left becoming afterwards paralyzed, a blister was put upon it, which also acted upon the other arm, not on the one to which it was applied! Owing to this sympathy or consent of parts, Broussais has laid down the pathological law, — that when an irritation exists for a long time in an organ, the textures that are analogous to the one, which is diseased, are apt to contract the

same affection.

c. Remote Sympathies. - As examples of the more distant kinds of sympathies, we may cite the effect produced upon the stomach by distant organs, and conversely. Among the earliest signs of pregnancy are nausea, and vomiting; loathing of food; fastidious appetite, &c. These symptoms are manifestly induced by sympathetic connection between the uterus and stomach; inasmuch as they are not adventitious, but occur more or less in all cases of pregnancy. Their absence, at least, is a rare exception to the rule. Hunger or dyspepsia, again, impresses a degree of langour, mental and corporeal, - which is proverbial; whilst the reception of food and its vigorous digestion give a character of energy and buoyancy, greatly contrasting with opposite circumstances. In disease, too, we find sympathies existing between the most distant portions of the frame, and although these are not apparent to us in health, we are perhaps justified in considering that an occult sympathy exists between them in health, which only becomes largely developed, and obvious to us, when the parts are affected with disease. It is probable, too, that in the successive evolution of organs at different periods of life, new sympathies arise, which did not previously exist or were not observable. The changes that supervene in the whole economy at puberty, strikingly illustrate this; changes which do not occur in those, who, owing to malformation, are not possessed of the essential parts of the reproductive system, or who have had them extracted prior to this period.

d. Imagination. — The effect of the intellectual and moral faculties on the exercise of the functions of other parts is strongly evidenced, especially in disease. The influence of the mind over the body is, indeed, a subject which demands the attention of every pathologist. In health, we notice the powerful effect induced by the affective faculties upon every function. All these are caused by sympathetic association with the brain; the action of the organs being in a state of excitation or depression, according to the precise character of the emotion. The intellectual

\* Nouveaux Elémens de la Science de l'Homme, Paris, 1806.

b Commentaries des Propositions de Pathologie; and Drs. Hays and Griffith's translation, p. 60, Philad. 1832.

manifestations probably exert their influence in a manner less evident, but not the less certain. The effects of one of them, at least, on the bodily functions are remarkable. We allude to the *imagination*; to which we can ascribe most of the cures that are said to have been effected by modes of management, — often of the most disgusting character, — which have been from time to time in vogue, have fretted their hour on the stage, and then sunk into that insignificance from which they ought never to have emerged.

We have had occasion to allude to the excited imagination of the maniac, the hypochondriac, and the nervous, and have remarked, that hallucinations may exist in those of sound mind; - phantoms created by the imagination; pains felt in various bodily organs, &c.; and we can hence understand, that, under particular circumstances, we may have actual disease produced in this manner; and, at other times, the feeling, -- which may be as distressing to the patient, - of disease, which has no existence except in the imagination. It is to the effect produced by the imagination, that we must ascribe the introduction into medicine of magic, sorcery, incantations, Perkinism, and other offsprings of superstition or knavery. The enthusiasm, that has attended the application of these last modes of acting upon the imagination in our own times, is most extraordinary, and their history leads us to be still more impressed with the extensive influence that may be exerted by the mind over the body: they teach the practitioner the importance of having its co-operation, whenever it can be procured; and the disadvantages, which he may expect to ensue, when the imagination is either arrayed against himself personally, or the plan of treatment which he is adopting. The physician, who has the confidence of his patient, will be successful - if he adopt precisely the same plan of treatment that would be pursued by one who has it not - in cases where the latter would totally fail. The applications of this subject are developed by the author elsewhere.b

Again, pathology is invoked as affording us perhaps the best evidences of the existence of extensive sympathetic relations between various parts of the frame, which are supposed to be constantly going on unseen during health, but become developed, and more obvious in disease. The case, we have previously given, of the general effects produced upon the system by local irritation of a part, shows the extent of such association. An insignificant portion of the body may become inflamed, and if the inflammation continue, the function of the stomach may be disordered,—as indicated by loss of appetite, nausea and vomiting; the respiration hurried, as well as the circulation; the senses blunted; the intellectual and moral faculties obscured; and languor and lassitude may indicate the nervous irritation and constraint.

The moral consideration of sympathy does not concern us. It

Demangeon, Du Pouvoir de l'Imagination, &c. chap. ii. p. 39, Paris, 1834.

<sup>&</sup>lt;sup>b</sup> General Therapeutics, p. 56, Philad. 1836; and General Therapeutics and Materia Medica, vol. i. Philad. 1843.

vol. II. - 48

is a subject, - and one of interest to the moral philosopher, - to account not only for these secret causes, which attract individuals towards each other, but which repel them and occasion antipathies. To a certain extent, however, it trends into the province of the physiologist. The tender, susceptible individual, from observing another suffering under pain, feels as if labouring under the same inconvenience, and, by a very rapid, yet complex intellectual process, constituted of numerous associations, may be so strongly impressed as to sink under their influence: - thus, the sight of blood will so powerfully impress the mind, in this sympathetic manner, that fainting may be induced, and the vital functions be for a time suspended. The sight and suffering of a woman in labour may cause abortion in another; and hence the propriety of excluding those, who are pregnant, from the chamber of the parturient female. Hysteric and convulsive paroxysms are induced in a similar way; of which the convulsionnaires of all times must be regarded as affording singular and instructive examples.

Superstitions connected with Sympathy. - Lastly; the mysterious consent, which we observe between various parts of the body, has given rise to some of the most strange and absurd superstitions that can be imagined. It was believed, for instance, almost universally in the fifteenth century, that an intimate sympathy exists, not only between parts of a body forming portions of one whole, but also between any substance that had previously formed part of a body and the body itself: that if, for example, a piece of flesh were sliced from the arm of one person and made to unite to that of another, the grafted portion would accurately sympathize with the body of which it had previously formed part, and undergo decay and death along with it; and it was even proposed to turn this sympathy to account. It was recommended, for instance, that the alphabet should be traced on the ingrafted portion; and it was affirmed, that when any of the letters, so traced, were touched, the party from whom the piece of flesh had been taken would feel similar impressions; so that, in this manner, a correspondence might be maintained. Some went even farther than this, asserting, that such a miraculous sympathy exists between the human body and all that has previously formed part of it, that if a hot iron were run into the excrement of any person, he would feel a sensation of burning in the part whence it had proceeded!

It was also a notion, that grafts of flesh, united to the body of another, die when the person dies from whom they have been taken. In a work on animal magnetism, the case of a man at Brussels is given, who had an artificial nose, formed after the old Taliacotian method, which served every useful purpose, until the person, from whom the graft had been taken, died, when it suddenly became cold and livid, and finally fell off. Tagliacozziahimself lived in an era of superstition, when this belief in the syn-

<sup>\*</sup> Gasparis Taliacotii Bononiensis De Curtorum Chirurgià per insitionem libri duo. Venet. 1597.

chronous death of the parent and graft was universally credited; and the folly has not escaped the notice of Butler : -

> "So learned Taliacotius from The brawny part of porter's bum, Cut supplemental noses, which Would last as long as parent breech; But when the date of nock was out, Off dropped the sympathetic snout." - Hudibras.

Little less singular was the superstition, - that the wounds of a murdered person will bleed afresh, if the body be touched, ever so lightly in any part by the murderer. This idea gave rise to the trial by bier-right, which has been worked up by Sir Walter Scott with so much dramatic skill, in one of his novels, - St. Valentine's Day, or the Fair Maid of Perth. The annals of judicial inquiry furnish us with many instances of this gross superstition.a A case of the kind occurred in our own country. It is contained in the attestation of John Demarest, coroner of Bergen county, New Jersey. The superstition, too, is noticed by many of the older poets. Thus, Shakspeare, — where the Lady Anne reviles Gloster over the corpse of Henry: —

> "O! gentlemen, see, see! dead Henry's wounds Open their congeal'd mouths and bleed afresh! Blush, blush, thou lump of foul deformity; For 'tis thy presence that exhales this blood From cold and empty veins, where no blood dwells. Thy deed, inhuman and unnatural, Provokes this deluge most unnatural."

Richard III., i. 2.

And Webster, in a tragedy published about the middle of the seventeenth century : -

> Her wounds still bleeding at the horrid presence Of yon stern murderer, till she find revenge."

Appius and Virginia.

The belief in these cases of monstrous superstition, which, it need scarcely be said, are usually explicable on purely physical principles, or on the excited imagination of the observer, still exists amongst the benighted inhabitants of many parts of Great Britain and Ireland, and is the main topic of one of the second series of "Traits and Stories of the Irish Peasantry." The superstition, has, indeed, its believers among us. On the trial of Getter, who was executed a few years ago (1833) in Pennsylvania, for the murder of his wife, a female witness deposed on oath, as follows: - " If my throat was to be cut, I could tell, before God Almighty, that the deceased smiled, when he (the murderer) touched her. I swore this before the justice, and that she bled considerably. I was sent for to dress her and lay her out.

\* See the case of Philip Stansfield for the Murder of his Father, Sir James Stansfield, in Hargrave's State Trials, iv. 283; and in Celebrated Trials of all Ages, &c., b Annual Register, for 1767. ii. 566, Lond. 1825.

He touched her twice. He made no hesitation about doing it. I also swore before the justice, that it was observed by other people in the house."

It would be endless to enumerate the various superstitions, which prevailed, a few centuries ago, on topics more or less remotely connected with this subject. We pass on, therefore, to the

interesting, but abstruse, inquiry into the

f. Agents by which Sympathy is accomplished .- The opinions of physiologists have, from time to time, rested chiefly; - on the membranes, the cellular tissue, the bloodvessels, and the nerves: whilst there have been some, who, in the difficulty of the subject, have supposed sympathy to be devoid of all organic connexion: and others, again, have presumed, that all the parts we have mentioned are concerned. The rapidity, however, with which sympathies are evidenced, has led to the abandonment of all those opinions; and the generality of physiologists of the present day look to the nervous system as the great source and medium of communication of the different irradiations, by which distant organs are supposed to react, in this manner, upon each other. The rapidity, indeed, with which the various actions of the nervous system are executed, - the apparent synchronism between the reception of an impression on an organ of sense, and its perception by the brain. as well as between the determination of the will and its effect upon a muscle, — naturally attracted the attention of physiologists to this system as the instrument of sympathy.

The modes, in which it is supposed to be accomplished, are:—
either by the parts that sympathize receiving ramifications from
the same nervous trunks, or from such as are united by nervous
anastomoses; or, by the nervous irradiations, emanating from one
organ, proceeding to the brain, and being thence reflected to every
dependency of the system, but so that certain organs are more
modified by such reflection than others; hence, the distinction
into what have been termed direct sympathies and cerebral

sympathies.

Of the direct sympathies we have already given some examples, — as that between the mucous and muscular coats of the intestines; and if our acquaintance with the precise distribution and connexion of the various parts of the nervous system were more intimate, we might perhaps explain many of the cases that are yet quite obscure to us. The researches of Sir Charles Bell, regarding the nerves concerned in respiration, have thrown light on those associations of organs, which we notice in the active exercise of the respiratory function. It has been elsewhere shown, that although the whole of the nerves, composing his respiratory system, may not be apparently in action during ordinary respiration, yet that when the function has been greatly excited, the association becomes obvious; parts, that are remote in situation, are combined in function, and all the nerves that animate them, he conceives, arise from the same column of the spine. The opinion of Boer-

haave, Meckel, and some others, is, that all sympathies are accomplished in this direct manner. On the other hand, Haller, Whytt, Georget, Broussais, Adelon, and others, make the majority of sympathies to be produced through the medium of the brain. Bostocka indeed affirms, that the facts, adduced by Whytt, are of such a nature as "to prove, that the co-operation of the brain is essential in those actions, which we refer to the operation of sympathy." In many cases, this is doubtless the fact; — as in sneezing and coughing; but there are others in which such co-operation seems improbable and indeed impossible. Something like sympathy exists in the vegetable; in which if we admit, with some naturalists, a rudimental nervous system, we have no reason for presuming, that there is any thing like a centre for the reception or transmission of impressions, and the case of infants, born devoid of brain and spinal cord, affords evidence of a like de-

scription.

We find, that the properties of the vital principle are exemplified by the formation of a body of a certain magnitude, form, structure, composition and duration, and that this applies to all organized bodies, vegetable as well as animal. Where such appearance of design, consequently, exists, we ought to expect, that, in the vegetable, also, a harmony or consent must reign amongst the various functions, tending to the accomplishment of that uniformity, which enables us always to recognise the particular varieties of the vegetable kingdom, and which has kept them as distinct, probably, in their characters, as when first created by Almighty power. The irritation of a single leaflet of the Mimosa pudica or sensitive plant causes the whole leaf, as well as the footstalk, to contract. Dr. John Sims irritated a leaflet of this plant, taking the greatest pains to avoid moving any other part of the leaf; yet the whole contracted, and the footstalk dropped. In order, however, to be sure, that mechanical motion communicated by the irritation had no share in the contraction, he directed a sunbeam, concentrated by a lens, on one of the leaflets, when the leaf again contracted, and the footstalk dropped. Of this kind of vegetable irritability we have many examples, some of which are alluded to under another head.

From these, and other facts of an analogous character, Sir Gilbert Blane<sup>c</sup> concludes, that the functions of living nature, in all its departments, are kept up by a mutual concert and correspondent accordance of every part with every other part, that it would be in vain to waste time in endeavouring to account for them by groping among dark analogies and conjectures; and that it is better to assume them as facts, on which are founded the ultimate and inscrutable principles of the animal economy. We have certainly much to learn regarding the agents of sympathies, and the modes in

\* Elementary System of Physiology, 3d edit. 762, Lond. 1836.

Elements of Medical Logic, 3d edit., Lond. 1819.

b An Essay on the Vital and other Involuntary Functions, § xi. Edinb. 1751.

which they are operated; but still we know enough to infer, that in many cases, in animals, the nerves appear to be the conductors; that the brain is, in others, the centre to which the organ in action transmits its irradiations, and by which they are reflected to the sympathizing organ; and that, in others, again, the effect is caused in the absence of nervous centre, and even of nerves, by vibrations perhaps, but in a manner which, in the present state of our knowledge, is inexplicable, and is, therefore, supposed to be essentially organic and vital, — epithets, however, as we have more than once said, that merely convey a confession of our total ignorance of the processes to which they are appropriated.<sup>a</sup>

# CHAPTER IV.

#### INDIVIDUAL DIFFERENCES AMONGST MANKIND.

The differences, which we observe amongst the individuals of the great human family, are as numerous as the individuals themselves; but this dissimilarity is not confined to man or to the animal kingdom: the vegetable exhibits the same; for, whilst we can readily refer any plant to the species and variety, to which it may have been assigned by the botanist, accurate inspection shows us, that in the precise arrangement of the stalk, branches, leaves, or flowers, no two are exactly alike. We shall not, however, dwell on these trifling points of difference, but restrict ourselves to the broad lines of distinction, that can be easily observed, and an attention to which is of some moment to the physician. Such are the temperaments, constitutions, idiosyncrasies, acquired differences, and the varieties of the human species or the different races of mankind. Of these, the last belongs more especially to the natural historian, and, consequently, will be but briefly noticed.

## 1. TEMPERAMENTS.

The temperaments are defined to be, — those individual differences, which consist in such disproportion of parts, as regards volume and activity, as to sensibly modify the whole organism, but without interfering with the health. The temperament is, consequently, a physiological condition, in which the action of the different functions is so tempered as to communicate certain characteristics, which may be referable to one of a few divisions. These divisions are by no means the same in all physiological treatises. The ancients generally admitted four, — denominated from the respective fluids or humours, the superabundance of which in the economy was supposed to produce them; — the sanguineous,

<sup>&</sup>lt;sup>2</sup> See, on the subject of sympathy, Alison, in Edinb. Med. Chirur. Transact ii. 165; Bostock, op. citat.; Adelon, Physiologie de l'Homme, 2de édit. iv. 260, Paris, 1829; and the author's Therapeutics, p. 72, Philad. 1836; and his General Therapeutics and Mat. Med. Philad. 1843.

caused by a surplus of blood; the bilious or choleric, produced by a surplus of yellow bile; the phlegmatic, caused by a surplus of phlegm, lymph, or fine watery fluid, derived from the brain; and the atrabiliary or melancholic, produced by a surplus of black bile,—the supposed secretion of the atrabiliary capsules and spleen.

This division was kept up for ages without modification, and still prevails with one or more additional genera. The epithets have been retained in popular language without our being aware of their parentage. For example, we speak of a sanguine, choleric, phlegmatic, or melancholic individual or turn of mind, with precisely the acceptation given to them by the Hippocratic school, - the possessors of these temperaments being presumed to be, respectively, full of high hope and buoyancy; naturally irascible, dull and sluggish; or gloomy and low-spirited. Metzger admits only two, - the irritable, (reizbare,) and the dull or phlegmatic (träge). Wrisberga eight, - the sanguine, sanguineo-choleric, choleric, hypochondriac, melancholic, Bæotian, meek, (s a n f t m üthige,) and the dull or phlegmatic. Rudolphib also eight, - the strong or normal, the rude, athletic or Baotian, the lively, the restless, the meek, the phlegmatic or dull, the timorous, and the melancholic; — whilst Broussaise enumerates the gastric, bilious, sanguine, lymphatico-sanguineous, anæmic, nervous, bilioso-sanguine, nervoso-sanguine, and melancholic. It is obvious, that if we were to apply an epithet to the possible modifications, caused by every apparatus of organs, the number might be extended much beyond any of these. Perhaps the division most generally adopted is that embraced by Richerand, who has embodied considerable animation, with much that is fanciful, in his description. In this division, the ancient terms have been retained, whilst the erroneous physiological basis, on which they rest, has been discarded. A short account of these temperaments is necessary, rather, for the purpose of exhibiting what has been, and is still, thought by many physiologists, than for attesting the reality of many of the notions that are mixed up with the subject. With this view, the temperaments may be divided into the sanguine, the bilious or choleric, the melancholic, the phlegmatic, and the nervous.

a. The Sanguine Temperament. — This is supposed to be dependent upon a predominance of the circulatory system; and hence is considered to be characterized by strong, frequent, and regular pulse; ruddy complexion; animated countenance; good shape, although distinctly marked; firm flesh; light hair; fair skin; blue eyes; great nervous susceptibility, attended with rapid successibilité, as the French term it; that is, — a facility of being impressed by external objects, and passing rapidly from one idea

a In his edition of Haller's Grundriss der Physiologie.

b Grundriss der Physiologie, Berlin, 1821.

Traité de Physiologie, Drs. Bell and La Roche's translation, 3d edit. p. 561, Philad.

Nouveaux Elémens de Physiologie, 13eme édit. par M. Bérard ainé, § ccxxviii. Bruxelles, 1837.

to another; quick conception; ready memory; lively imagination; addiction to the pleasures of the table; and amorousness. The diseases of the temperament are generally violent; and are chiefly seated in the circulatory system,—as fevers, inflamma-

tions, and hemorrhages.

The physical traits of this temperament, according to Richerand, are to be found in the statues of Antinous and the Apollo Belvidere: the moral physiognomy is depicted in the lives of Mark Antony and Alcibiades. In Bacchus, both the forms and the character are found; and no one in modern times, in M. Richerand's opinion, exhibits a more perfect model of it than the celebrated Duke De Richelieu; — amiable, fortunate and valorous, but light

and inconstant to the termination of his brilliant career.

If individuals of this temperament apply themselves to labours of any kind that cause the muscles to be greatly exerted, these organs become largely developed, and a subdivision of the sanguine temperament is formed, which has been called the *muscular* or athletic. This is characterized by all the outward signs of strength; the head is small; the neck strong; the shoulders broad; the chest large; the hips solid; the muscles prominent, and the interstices well marked. The joints, and parts not covered with muscles, seem small; and the tendons are easily distinguished through the skin, by their prominence. The susceptibility to external impressions is not great; the individual is not easily roused; but, when he is, he is almost indomitable. A combination of the physical powers, implied by this temperament, with strong intellect, is rarely met with.

The Farnesian Hercules is conceived to offer one of the best specimens of the physical attributes of the athletic temperament.

b. The Bilious or Choleric Temperament. — This is presumed to be produced by a predominance of the liver and biliary organs in general. The pulse is strong, hard and frequent; the subcutaneous veins are prominent; the skin is of a brown colour, inclining to yellow; hair dark; body moderately fleshy; muscles firm, and well marked; the passions violent, and easily excited; the temper abrupt, and impetuous; great firmness and inflexibility of character; boldness in the conception of projects, and untiring perseverance in their fulfilment. It is amongst the possessors of this temperament, that the greatest virtues and the greatest crimes are met with. Richerandb enumerates Alexander, Julius Cæsar, Brutus, Mahomet, Charles XII., Peter the Great, Cromwell, Sextus V., and the Cardinal Richelieu. To these, Good has added Attila, Charlemagne, Tamerlane, Richard III., Nadir Shah, and Napoleon.

The moral faculties are early developed; so that vast enterprises may be conceived, and executed at an age when the mind is ordinarily far from being matured. The diseases are generally combined with more or less derangement of the hepatic system. The

Richerand, op. citat. § ccxxix. Dop. cit. § ccxxx. Book of Nature, iii. 310.

whole of the characters, however, indicate, that an excited state of the sanguiferous system accompanies that of the biliary organs; so that the epithet cholerico-sanguine might, with more propriety, be applied to it. Where this vascular predominance does not exist, whilst derangement is present in some of the abdominal organs, or in the nervous system, we have the next genus produced.

c. Melancholic or Atrabilious Temperament.— Here the vital functions are feebly or irregularly performed; the skin assumes a deeper hue; the countenance is sallow and sad; the bowels are torpid, and all the excretions tardy; the pulse is hard and habitually contracted; the imagination is gloomy, and the temper suspicious. The characters of Tiberius and of Louis XI. are considered to be instances of the predominance of this temperament; and, in addition to these, Richeranda has enumerated Tasso, Pascal, Gilbert, Zimmermann, and Jean Jacques Rousseau.

d. The Phlegmatic, Lymphatic or Pituitous Temperament. -In this case, the proportion of the fluids is conceived to be too great for that of the solids; the secretory system appearing to be active, whilst the absorbent system does not act so energetically as to prevent the cellular texture from being filled with the humours. The characteristics of this temperament are: - soft flesh; pale skin; fair hair; weak, slow and soft pulse; figure rounded, but inexpressive; the vital actions more or less languid; the memory by no means tenacious, and the attention vacillating; with aversion to both mental and corporeal exertion. Pomponius Atticus - the friend of Cicero - is offered as an example of this temperament, in ancient times; Montaigne, in more recent history. The latter, however, possessed much of the nervous susceptibility that characterizes the more lively temperaments. Dr. Goodb suggests the Emperor Theodosius as an example in earlier times; and Charles IV. of Spain, who resigned himself almost wholly into the hands of Godoy, - Augustus, King of Saxony, who equally resigned himself into the hands of Napoleon, - and Ferdinand of Sicily, who surrendered for a time the government of his people to the British, - as instances in our own day. It would not be difficult to find, amongst the crowned heads of Europe, others that are equally entitled to be placed amongst these worthies.

e. The Nervous Temperament. — Here the nervous system is greatly predominant; the susceptibility to excitement from external impressions being unusually developed. Like the melancholic temperament, this is, however, seldom natural or primitive. It is morbid or secondary, being induced by sedentary life, sexual indulgence, or morbid excitement of the imagination, from any cause. It is characterized by small, soft, and, as it were, wasted muscles; and generally, although not always, by a slender form; great vividness of sensation; and promptitude and fickleness of resolution and judgment. This temperament is frequently combined with

<sup>\*</sup> Op. citat. ccxxxi. b Op. citat. iii. 314; and Richerand, op. cit. § ccxxxii.

some others. The diseases that are chiefly incident to it, are of the hysterical and convulsive kind; or those to which the epithet nervous is usually appropriated. Voltaire and Frederick the Great are given by Richeranda as examples of this temperament.

Such are the temperaments, described by most writers. slightest attention to their reputed characteristics will show the imperfection of their definition and demarcation; so imperfect, indeed, are they, that it is extremely rare for us to meet with an individual, whom we could unhesitatingly refer to any one of them. They are also susceptible of important modifications by climate, education, &c., and may be so combined as to constitute innumerable shades. The man of the strongest sanguine characteristics may, by misfortume, assume all those that are looked upon as the indexes of the melancholic or atrabilious; and the activity and impetuosity of the bilious temperament may, by slothful indulgence, be converted into the lymphatic or phlegmatic. It is doubtful, and more than doubtful, also, whether any of the mental characteristics, assigned to the temperaments, are dependent upon them. The brain, we have elsewhere seen, is the organ of the mental and moral manifestations; and although we may look upon the temperaments as capable of modifying its activity, they cannot probably affect the degree of perfection of the intellect; - its strength being altogether dependent upon cerebral conformation. It is even doubtful whether the temperaments can interfere with the activity of the cerebral functions. In disease of the hepatic, gastric or other viscera we certainly see a degree of mental depression and diminished power of the whole nervous system; but this is the effect of a morbid condition, and continues only so long as such morbid condition endures. Nor is it probable, that any predominance of the nutritive functions could induce a permanent influence on the cerebral manifestations. Whatever might be the effect for a while, the nervous system would ultimately resume the ordinary action which befitted its primitive organization. Similar reasons to those have induced the author's late friend, M. Georget, b - a young physician of great promise and experience in mental affections, now no more, - to consider the whole doctrine of the temperaments as a superstition connected with the humoral pathology, and to believe, that the brain alone, amongst the organs, has the power, by reason of its predominance or inferiority, to modify the whole economy.

That a difference of organization exists in different individuals is obvious; but that there is an arrangement of the nutritive organs or apparatuses, which impresses upon individuals all those mental and other modifications known under the name of temperaments, is, we think, sufficiently doubtful. The constitution of an individual is the mode of organization proper to him. A man, for example, is said to have a robust, or a delicate, or a good, or a bad constitution, when he is apparently strong or feeble, usually

<sup>\*</sup> Op. cit. ccxxxiii. b De la Physiologie du Système Nerveux, &c., Paris, 1821.

in good health, or liable to frequent attacks of disease. The varieties in constitution are, therefore, as numerous as the individuals themselves. A strong constitution is considered to be dependent upon the due development of the principal organs of the body, on a happy proportion between those organs, and on a fit state of energy of the nervous system; whilst the feeble or weak constitution results from a want of these postulates. Our knowledge, however, of these topics, is extremely limited, and concerns the pathologist more than the physiologist.

## 2. OF IDIOSYNCRASY.

The word idiosyncrasy is used, by many physiologists, synonymously with constitution: but it is generally appropriated to the peculiar disposition, which causes an individual to be affected by extraneous bodies, in a way in which mankind in general are not acted upon by the same agents.a In all cases, perhaps, these peculiarities are dependent upon inappreciable structure, either of the organ concerned, or of the nervous branches distributed to it; at times, derived from progenitors; at others, acquired, - often by association, - in the course of existence. Hence arise many of the antipathies to particular animate and inanimate objects, which we occasionally meet with, and of which Broussais relates a singular instance in a Prussian captain, whom he saw at Paris in 1815. He could not bear the sight of a cat, a thimble, or an old woman, without becoming convulsed, and making frightful grimaces! The associations must have been singularly complicated to occasion an antipathy to objects differing so signally from each other. Wagner, b of Vienna, has collected a multitude of cases of idiosyncrasy; and the observation of every individual, whether of the medical profession or not, must have made him acquainted with those peculiarities, that render a particular article of diet, which is innoxious, and even agreeable and wholesome to the generality of individuals, productive, in some, of the most unpleasant effects. Haller knew a person who was always violently purged by the syrup of roses. A friend of the author is purged by opium, which has an opposite effect on the generality of individuals. Dr. Parise says he knew two cases, in which the odour of ipecacuanha always produced most distressing dyspnæa: the author knew a young apothecary, who could never powder this drug without the supervention of the most violent catarrh. A friend of Tissot could not take sugar without its exciting violent vomiting. Urticaria or nettle-rash is very frequently occasioned, in particular constitutions,

> "Some love not a gaping pig, And others, when the bagpipe sings i' th' nose, Cannot contain their urine for affection."

Pharmacologia, 4th Amer. Edit. by J. B. Beck, p. 189, New York, 1831.

Shakspeare.

b Hufeland and Himly's Journal der Pract. Heilkund., Nov. 1811, § 55. See, also, art. Idiosyncrasie, Dict. des Sciences Médicales, tom. xxii.; and Fletcher's Elements of General Pathology, p. 21, Lond. 1842.

by taking shell-fish. The same effect is produced on two young female friends of the author by eating strawberries; and similar cases are given by Roose. M. Chevalier relates the case of a lady, who could not take powdered rhubarb without an erysipelatous efflorescence showing itself, almost immediately, on the skin; yet she could take it in the form of infusion with perfect impunity.

The above idiosyncrasies apply only to the digestive function. We find equal anomalies in that of the circulation. In some, the pulse is remarkably quick, upwards of one hundred in the minute; in others, it is under thirty. That of Napoleon is said to have beaten only forty-four times in a minute.<sup>a</sup> It may also be unequal, and intermittent, and yet the individual be in a state of health.<sup>b</sup>

The senses offer us some of the most striking cases of this kind of peculiarity. Many strong individuals cannot bear the smell of the apple, cherry, strawberry, or of musk, peppermint, &c. Pope Pius VII. had such an antipathy to musk, that, on one occasion of presentation, an individual of the company having been scented with it, his holiness was obliged to dismiss the party almost immediately. The idiosyncrasies of taste are also numerous: some of these cases of singular and depraved sense we have described under the sense of taste. Dejeangives us the case of an individual of distinguished rank who was fond of eating excrement. Certain animals, again, as the turkey, have an antipathy to the colour of red; and Von Büchner and Tissot cite the case of a boy who was subject to epileptic fits whenever he saw any thing of a red colour.

Occasionally, we meet with similar idiosyncrasies of audition. Sauvages relates the case of a young man labouring under intense headache and fever, which could not be assuaged by any other means than the sound of the drum. The noise of water issuing from a pipe threw Bayle into convulsions. The author has a singular peculiarity of this kind, derived from some accidental association in early life. If a piece of thin biscuit be broken in his presence, — nay, the idea alone is sufficient, — the muscles that raise the left angle of the mouth are contracted, and this irresistibly.

The sense of tact is not free from idiosyncrasies. Wagner cites the case of a person, who felt a sensation of cold along the back, whenever he touched the down of a peach with the point of his finger; or when the down came in contact with any part of his skin. He was remarkably fond of the fruit, yet was unable to indulge his appetite unless a second person previously removed the skin. Prochaska relates the case of a person, who was affected

with nausea whenever he touched this fruit.

b. Adelon, Physiologie de l'Homme, edit. cit. iv. 511.

" Mathew's Diary of an Invalid.

<sup>&</sup>lt;sup>a</sup> See some cases of unusual slowness of the pulse, by Mr. H. Mayo, in Lond. Med. Gazette, May 5, 1838, p. 232; and Dunglison's American Medical Intelligencer, July 2, 1838, p. 103. In one case of fatal disease the pulse beat only nine in the minute!

It is, of course, all important, that the practitioner should be acquainted with these idiosyncrasies, and so far the notion of "knowing the constitution," — which is apt to be used to the prejudice of the young practitioner, or of any except the accustomed medical attendant, — has some reason in it. It is the duty, however, of the patient to put the practitioner in possession of the fact of such peculiarities, so that he may be enabled to guard against them, and not take that for morbid which is the effect of simple idiosyncrasy. This, however, is a topic which belongs rather to therapeutics.<sup>a</sup>

## 3. OF NATURAL AND ACQUIRED DIFFERENCES.

1. NATURAL DIFFERENCES. — The temperaments, constitutions, and idiosyncrasies may, as we have seen, either be dependent upon original conformation, or they may be produced by external influences; hence, they have been divided into the natural and acquired. Under the former head are included all those individual differences, derived from progenitors, which impress upon the individual more or less of resemblance to one or both parents. It has been properly observed, that the individuality of any human being that ever existed was absolutely dependent on the union of one particular man with one particular woman; and that if either the husband or the wife had been different, a different being would have been ushered into existence. For the production of Shakspeare, or Milton, or Newton, it was necessary, that the father should marry the identical woman he did marry. If he had selected any other wife, there would have been no Shakspeare, no Milton, no Newton. Sons might have been born of other women, but they would not have been the same, either in mental or physical qualities. All this, however, enters into the question of the influence of both parents on the fœtus in utero, which we have considered else-

a. Peculiarities of the Female. — Amongst the natural differences, those that relate to sex are the most striking. In a previous part of this volume, we have described the peculiarities of the sexual functions in both male and female, but other important differences have not been detailed. All the descriptions, when not otherwise specified, were presumed to apply to the adult male. At present, it will be only necessary to advert to the peculiarities of the female.

The stature of the female is somewhat less than that of the male, the difference being estimated at about a twelfth. The chief parts of the body have not the same mutual proportions. The head is smaller and rounder; the face shorter; the trunk longer, especially the lumbar portion, and the chest more convex. The lower extremities, especially the thighs, are shorter, so that the half of the body does not fall about the pubes, as in man, but higher. The neck is longer; the abdomen broader, larger, and

See the author's Therapeutics, p. 38, Philad. 1836; or his General Therapeutics and Mat. Med., Philad. 1843.

VOL. II. -49

more prominent; and the pelvis has a greater capacity to adapt it for gestation and parturition. The long diameter of the brim is from side to side, whilst, in the male, it is from before to behind; the arch of the pubis is larger, and the tuberosities of the ischia are more widely separated, so that the outlet of the pelvis is larger than in the male; the hips are broader, and, consequently, the spaces between the heads of the thigh-bone greater; the knees are more turned in, and larger than in the male; the legs shorter, and the feet smaller. The shoulders are round; but the width across them, compared with that of the hips, is not so great as in man; the arms are shorter, fatter, and more rounded; the same is the case with the fore-arm; the hand is smaller, and softer, and the fingers are more delicate.

The whole frame of the female is more slender; the bones are smaller, their tissue is less compact, and the prominences and corresponding depressions are less marked; the subcutaneous cellular tissue is more abundant, and filled with a whiter and firmer fat; a similar adipous tissue fills up the intervals between the muscles, so that the whole surface is rounder, and more equable, than that of the male; the skin is more delicate, whiter, better supplied with capillary vessels, and less covered with hair; the hair of the head, on the other hand, is longer, finer, and more flexible; the nails are softer and of a more red hue; the muscles of the face are less distinctly marked, so that the expression of the eye, and the emotions, which occasion elevation or depression of the angles of the mouth,—laughing and weeping, for example,— are more strongly defined. On the whole, the general texture of the organs is looser and softer.

The above observations apply to what may be termed the standard female,—one whose natural formation has not been interfered with by employments, which are usually assigned to the other sex. It can be readily understood, that if the female have been accustomed to the laborious exercise of her muscles, they may become more and more prominent, the interstices between them more and more marked, the projections and depressions of the bones on which they move more distinct; the whole of the delicacy of structure may be lost; and the skeleton of the female, thus circumstanced, may be scarcely distinguishable from that of the inactive male, except in the proportions of the pelvis, in which the sexual differences are chiefly and characteristically situate.

Many of the functions of the female are no less distinctive than the structure. The senses, as a general rule, are more acute, whether from original delicacy of organization, or from habit, is not certain; — probably both agencies are concerned. The intellectual and moral faculties are also widely different, and this, doubtless, from original conformation, although education may satisfactorily account for many of the differences observable between the sexes. Gall is one of the few anatomists, who have attended to the comparative state of the cerebral system in the sexes; and the results of his investigations lead him to affirm,

that there is a striking difference in the development of different parts of the encephalon in the two, which he thinks may account for the difference observable in their mental and moral manifestations. In the male, the anterior and superior part of the encephalon is more developed; in the female, the posterior and inferior: the former of these he conceives to be the seat of the intellectual faculties; the latter of those feelings of love and affection, which seem to preponderate in the character of the female. We have elsewhere said, that the views of Gall, on this subject, are not yet received as confirmed truths, and that we must wait until farther experience and multitudinous observations shall have exhibited their accuracy, or want of foundation. Independently, however, of all considerations deduced from organization, observation shows, that the female exhibits intellectual and moral dif- . ferences, which are by no means equivocal. The softer feelings predominate in her, whilst the intellectual faculties have the preponderance in man. The evidences and character of the various shades of feeling and susceptibility, and the influence of education and circumstances on these developments, are interesting topics for the consideration of the moral philosopher, but admit of little elucidation from the labours of the physiologist. The only inference, to which he can arrive, is, that the causes of the diversity are laid in organization, and become unfolded and distinctive by education. The precise organization he is unable to depict, and the influence of circumstances on the mind it is scarcely his province to consider.

The function of muscular motion is, owing to organization, more feebly executed in the female. We have already remarked; that the bones are comparatively small, and the muscles more delicately formed. The energy of the nervous system is also less; so that all the elements for strong muscular contraction are by no means in the most favourable condition; and, accordingly, the power the female is capable of developing by muscular contraction is much less than that of the male. Her locomotion is somewhat peculiar, - the wide separation of the hip-joints, owing to the greater width of the pelvis, giving her a characteristic gait. The vocal organs exhibit differences, which account for the difference in the voice. The chest and the lungs are of smaller dimensions: the trachea is of less diameter; the larynx smaller; the glottis shorter and narrower; and the cavities, communicating with the nose, are of smaller size. This arrangement causes the female voice to be weaker, softer and more acute. The muscles of the glottis, and the ligaments of the glottis themselves, are apparently more supple, so as to admit of the production of a greater number of tones, and to favour singing. The phenomena of expression, as we have often remarked, keep pace with the condition of the intellectual and moral faculties, and with the susceptibility of the nervous system. As this last is generally great in the female, the language of the passions, especially of the softer kind, are more marked in her.

The functions of nutrition present, also, some peculiarities, With regard to digestion, less food is generally required; the stomach is less ample; the liver smaller; and frequently, - at least more frequently than in the male, - the dentes sapientiæ do not appear. The desire for food at the stated periods is not so powerful; and it is generally for light and agreeable articles of diet rather than for the very nutritious; but the appetite returns more frequently, and is more fastidious, owing to the greater sensibility of the digestive apparatus. This, however, is greatly an affair of habit, and we have more instances of prolonged abstinence in the female than in the male. The circulation is generally more rapid, the pulse being less full, but quicker. Of the secretions, that of the fat alone requires mention, which is usually more abundant and the product firmer. The cutaneous transpiration is less active, and the humour has a more acidulous odour. The urine is said, by some writers, to be less abundant, and less charged with salts; whence, it is asserted, there is less disposition to calculous affections. So far, however, as we have had an opportunity for judging, it is secreted in greater quantity, and this may partly account for its seeming to have a smaller quantity of salts in any given amount; but the truth is, the freedom of the female from calculous affections is greatly owing to the shortness and size of the urethra, which admits the calculus to be discharged with comparative facility; and it is a common observation, that where the males of a family, hereditarily predisposed to gout, become, owing to their greater exposure to the exciting causes, affected with that disease, the females may be subject to calculous disorders, - the two affections appearing to be, in some respects, congenerous. For the reasons already mentioned, stone rarely forms in the bladder of the female, and the operation of lithotomy is scarcely ever necessary. The desire to evacuate the contents of the bladder occurs more frequently in the female, probably, in part, owing to habit; and, in part, to the greater mobility of the nervous system.

In addition to these differences as regards the secretions, the female has one peculiar to herself, — menstruation, — a function which has already engaged attention. In the progress of life, too, the glandular system undergoes evolutions which render it especially liable to disease. About the period of the cessation of the menses, — sooner or later, — the mammæ frequently take upon themselves a diseased action, and become scirrhous and cancerous

so as to require the organs to be extirpated.

In the treatment of disease, these sexual peculiarities have to be borne in mind. Owing to the greater mobility of the nervous system in the female, she usually requires a much smaller dose of any active medicine than the male; and during the period when the sexual functions are particularly modified, as during menstruation, gestation, and the child-bed state, she becomes liable to various affections, some of which have been referred to elsewhere;

HABIT. 581

others belong more appropriately to works on pathology, therapeutics or obstetrics,a

2. Acquired Differences, - The acquired differences, which

we observe amongst individuals, are extremely numerous. The effect of climate on the physical and mental characteristics is strikingly exhibited. The temperate zone appears to be best adapted for the full development of man, and it is there, that the greatest ornaments of mankind have flourished, and that science and art have bloomed in exuberance; whilst in the hot, enervating regions of the torrid zone, the physical and moral energies are prostrated; and the European or Anglo-American, who has entered them full of life and spirits, has left them after a few years' residence, listless and shorn of his proudest characteristics. Nor is the hyperborean region more favourable to mental and corporeal development; the sensibility being obviously blunted by the rigours of the climate. The effect of locality is, perhaps, most signally exemplified in the Crétin and the Goîtreux, of the Valais, and of the countries at the base of lofty mountains in every part of the globe; as well as in the inhabitants of our low countries, who are constantly exposed to malarious exhalations, and bear the sallow imprint on the countenance.

Not less effective in modifying the character of individuals is the influence of the way of life, education, profession, government, &c. The difference between the cultivated and the uncultivated; between the humble mechanic, who works at the anvil or the lathe, and him whose avocation, like that of the lawyer, and the physician, consists in a perpetual exercise of the organ of intellect; and between the debased subject of a tyrannical govern-

ment, and the independent citizen of a free state, -

"Lord of the lion heart and eagle eye,"

is signala nd impressive.

1. Habit. - To the acquired differences in individuals from extraneous or intrinsic causes we must refer habit, which has been defined, - an acquired disposition in the living body, that has become permanent, and as imperious as any of the primitive dispositions. It is a peculiar state or disposition of the mind, induced by the frequent repetition of the same act. Custom and habit are frequently used synonymously; but they are distinct. Custom is the frequent repetition of the same act; habit is the effect of such repetition. By custom we dine at the same hour every day: the artificial appetite induced is the effect of habit.

The functions of the frame are variously modified by this disposition, - being, at times, greatly developed in energy and rapidity, at others, largely dminished. If a function be over and over again exerted to the utmost extent of which it is capable, both as regards energy and activity, it becomes more and more easy of execution;

<sup>\*</sup> See, on the whole of this subject, Weber's Hildebrandt's Handbuch der Anatomie, iv. 527, Braunschweig, 1832.

the organ is daily better adapted for its production, and is so habituated to it, that it becomes a real want — a second nature. It is in this way, that we accustom the organs of speech, locomotion, &c., to the exercise of their functions, until, ultimately, the most varied combinations of the muscular movements of the tongue and limbs can be executed with surprising facility. If, on the contrary, the organs of any function possess unusual aptitude for accomplishing it, and we accustom ourselves to a minor degree of the same, we ultimately lose a part of the aptitude, and the organs become less inclined, and less adapted to produce it. By custom we may habituate ourselves to receive an unusually small quantity of nutriment into the stomach, so that at length it may become im-

practicable to digest more.

A similar effect occurs as regards the quantity of the special irritant, which we allow to impinge on any of the organs of sense. If we accustom them to be feebly impressed, yet sufficiently so for the performance of their functions, they become incapable of supporting a greater quantity of the special irritant without indicating suffering. The miner can see into the farthest depths of his excavations, when, to the eye of one, who has descended from the bright light of day, all seems enveloped in obscurity. In this case, the sensibility of the organ of sight is developed to such an extent, that if the individual be brought into even a feeble light, the impression is extremely painful. The nyctalope is precisely so situate. His nerves of sight are so irritable, that, although he can see well in the night, he is incapable of accurate discrimination by day. On the other hand, exposure to intense light renders the sensibility of the visual nerves so obtuse, that objects are not so readily perceived in obscurity. The hemeralope, who sees in the day and not in the night, and who is consequently the antitheton of the nyctalope, has the nervous system of vision unusually dull, and incapable of excitement by feeble impressions.

It may be laid down, as a general principle, that if we gradually augment the stimulus applied to any organ of sense, it becomes less susceptible of appreciating minor degrees of the same irritant; so that, in this way, an augmented dose of the irritant is progressively required to produce the same effect. This is daily exemplified by the use of tobacco,—either in the form of chewing, smoking, or snuffing, which becomes a confirmed habit, and can only be abandoned—without doing great violence to the feelings—by attention to the principle deduced from practice,—that by gradually following the opposite course to the one adopted in acquiring the habit—that is, by accustoming the nerve of sense to a progressive diminution in the dose of the stimulus—an opposite habit may be formed, and the evil, in this manner, be removed.

When, by habit, we acquire extreme facility in executing any function, it may be accomplished apparently without the direct agency of volition This is peculiarly applicable to the voluntary emotions. We have elsewhere shown, that, in this case, habit

HABIT. 583

only communicates the facility, and that there is no natural sequence of motions, and consequently, no reason, - as in executing a rapid musical movement, - why one movement of the fingers should follow rather than another, unless volition were the guiding power. Volition, as Dr. Parr has remarked, is not an exertion of the mind, but apparently a simple impulse directed almost necessarily to an end; and it is affected, by custom, nearly like the organs of the body. Thus, a sensation, which excited a perceptible exertion of volition, will, in time, produce it and the correspondent action, without our being sensible of its interference; and so rapid is this process, that we seem to will two ends or objects at the same time, though they are evidently, when examined, distinct operations. But though, by custom, we are no longer sensible of bodily impressions, or of the exercise of volition, the corporeal organs, in their several functions, acquire, like those of the mind, peculiar accuracy of discrimination. The musician is not, for instance, sensible of his willing any one motion; yet with the most exquisite nicety he touches a particular part of the string of the violin, and executes a variety of the nicest and most complicated movements with the most delicate precision.

It is a common remark, that "habit blunts the feeling but improves the judgment." To a certain extent this is true; but the feeling is not blunted unless the stimulant, which acts upon the organ of sense, is too powerful, and too frequently repeated. When moderately exercised, the effect of education, in perfecting all the senses, is strongly exhibited. Sensations, often repeated, cease to be noticed, not because they are not felt, but because they are not heeded; but if the attention be directed to the sensation, custom adds to the power of discrimination. Hence the sailor is able to detect the first appearance of a sail in the distant horizon, when it cannot be perceived by the landsman; and a similar kind of discrimination is attained by the due exercise of the other senses. This greater power of discrimination is doubtless owing to improvement in the cerebral or percipient part of the visual apparatus; but we have no evidence, that the organ of vision has its ac-

tion necessarily blunted.

It has been presumed, by some physiologists and metaphysicians, that the will, by custom and exercise, may acquire a power over certain functions of the body, which were not originally subject to it; nay, some speculatists have gone farther, and affirmed, that all the involuntary functions were originally voluntary, and that they have become involuntary by habit. Stahl and the other animists, who regarded the soul as the formative and organizing agent in animals, asserted, that it excites the constant movements of the heart, and of the respiratory, digestive and other nutritive organs, by habits so protracted and inveterate, and so naturalized within us, that these functions can be effected without the aid of the will, and without the slightest attention being paid to them. Respiration, according to them, is originally voluntary; but, by habit, has

become spontaneous; so that there is no farther occasion to invoke volition. Respiration goes on night and day, when we are asleep as well as awake; and they regard, as a proof that the action was originally dependent upon free will, that we are still able to accelerate or retard it at pleasure, They cite, moreover, the case of Colonel Townshend, related in another part of this work, (p. 124 of this volume,) to show, that the action of the heart is capable of being influenced by the will; and, likewise, the fact that it is ac-

celerated or retarded under the different passions.

Condillac, Dutrochet, and De Lamarck, again, fantastically assert, that the different instincts, observed to prevail so powerfully in animals, are mere products of an acquired power, transmitted through successive generations. The views of the last distinguished naturalist, regarding the effect of habit on organization, which he considers to tend to greater and greater complication, are most singular and fantastic. It is not, he considers, the organs of an animal that have given rise to its habits; on the contrary, its habits, mode of life, and those of its ancestors have, in the course of time, determined the shape of its body, the number and condition of its organs, and the faculties, which it enjoys. Thus the otter, the beaver, the waterfowl, the turtle, and the frog were not made web-footed that they might swim; but their wants having attracted them to the water, in search of prey, they stretched out their toes to strike the water, and move rapidly along its surface. By the repeated stretching of the toes, the skin, which united them at the base, acquired a habit of extension, until, in the course of time, they became completely web-footed. The camelopard, again, was not gifted with a long flexible neck, because it was destined to live in the interior of Africa, where the soil is arid and devoid of herbage; but, being reduced, by the nature of the country, to support itself on the foliage of lofty trees, it contracted a habit of stretching itself up to reach the high boughs, until its forelegs became longer than the hinder, and its neck so elongated that it could raise its head to the height of twenty feet above the ground!

The objections to all these views are, — that the functions in question are as well performed during the first day of existence as at an after period, and are apparently as free from the exercise of all volition. The heart, indeed, beats through fætal existence for months before the new being is ushered into the world; and when, if volition be exerted at all, it can only be so obscurely. The case of Colonel Townshend is strange — passing strange — but it is almost unique, and the power of suspending the heart's action was possessed by him a short time only prior to dissolution. All the functions in question must, indeed, be esteemed natural, and instinctive, inseparably allied to organization; and

<sup>&</sup>lt;sup>2</sup> Philosophie Zoologique, tom. i. p. 218, and tom. ii. p. 451, edit. 1830; Adelon, art. Habitude, Dict. de Medecine, x. 501, Paris, 1824; and Physiologie de l'Homme, 2de édit. iv. 517, Paris, 1829.

hence differing from the results of habit which is always ac-

quired.

The opinion of Bichat, on the other hand, was, that habit influences only the animal functions, and has no bearing on the organic or nutritive. But this is liable to objections. We have seen, under Digestion, that if a bird, essentially carnivorous in its nature, be restricted to vegetable food, the whole digestive economy is modified, and it becomes habituated to the new diet. We know, also, that where drains are established in any part of the body, they become, in time, so much a part of the physiological condition of the frame, that they can only be checked with safety by

degrees.a

In the administration of medicines, habit has always to be attended to. The continued use of a medicine generally diminishes its power; — hence the second dose of a cathartic ought to be larger than the first, if administered within a few days. Certain cathartics are found, however, to be exceptions to this. The Cheltenham water and the different saline cathartics are so. The constitution, so far from becoming reconciled to lead by habit, is rendered more and more sensible to its irritation. Emetics, too, frequently act more powerfully by repetition. Dr. Cullen asserts, that he knew a person so accustomed to excite vomiting on himself, that the one-twentieth part of a grain of tartarized antimony was sufficient to produce a convulsive action of the parts concerned in vomiting. As a general rule, however, medicines lose their effect by habit, and this is particularly the case with tonics; but if another tonic be substituted for a day or two, and then the former be resumed, it will produce all its previous effects.

2. Association. — Association, employed abstractedly, is a principle of the animal economy nearly allied to habit. When two or more impressions of any kind have been made upon the nervous system, and repeated for a certain number of times, they may become associated; and if one of them only be produced it will call up the idea of the others. It is a principle, which is largely invoked by the metaphysician, and by which he explains many interesting phenomena of the human mind, especially those connected with our ideas of beauty, or the contrary; our likes and dislikes, and our sense of moral propriety. Darwine employed it to explain many complicated functions of the economy; and he laid it down as a law, that all animal motions, which have occurred at the same time or in immediate succession, become so associated, that when one of them is reproduced, the other has a tendency to accompany or succeed it. The principle has, doubtless, great agency in the production of many of the physical, as well as mental, phenomena; but its influence has been overrated; and many of the

b See the author's General Therapeutics, p. 42, Philad. 1836.

Adelon, in Dict. de Méd. x. 498; and Physiologie de l'Homme, edit. cit. iv. 525; see, also, Lepelletier, Physiologie Médicale et Philosophique, i. 259, Paris, 1831.

<sup>·</sup> Zoonomia, i. 49, Lond. 1794.

consecutive and simultaneous actions, to which we have referred under the head of Correlation of Functions, take place, apparently as well the first time they are exerted, as subsequently. Sucking and deglutition are good cases of the kind. Soon after birth, the muscles of the lips, cheeks, and tongue are contracted to embrace the nipple, and to diminish the pressure in the interior of the mouth; and, as soon as the milk has flowed to the necessary extent into the mouth, certain voluntary muscles are contracted. These propel the milk into the pharynx, where its farther progress is effected by muscles, associated or connected functionally, but not in the sense we are now employing the epithet; for here one action could not suggest another, according to the definition we have given of association, which requires that the acts should have been executed previously. Many of the cases, in fact, ascribed by Darwin and Hartleya to the agency of this principle, are instinctive actions, in which a correlation - as in the case of deglutition - exists, but without our being able to explain the nature of such correlation, any more than we can explain other complicated actions and connexions of the nervous system, of which this is doubtless one. Some of the most obstinate diseases are kept up by habit, or by accustomed associated motions; and, frequently, the disease will seem to continue from this cause alone. Whenever intermittent fever, epilepsy, asthma, chorea, &c., have been long established, the difficulty of removing the influence of habit. or the tendency to recurrence, is extreme. In such cases, the principle of revulsion can be invoked with much advantage by the therapeutist.b

3. Imitation. — The principle of imitation falls appropriately under this section. It may be defined - that consent of parts, depending on similar organization, which, under the influence of the brain, enables them to execute acts similar to those executed by the same parts in another individual. Imitation, consequently, requires the action of the brain; and differs from those actions that are natural or instinctive to organs. For example, speech requires the action of imitation; whilst the ordinary voice or cry is effected by the new-born, and by the idiot, who are incapable of all observation, and consequently of imitation. The mode in which speech is acquired, offers us one of the best examples of this imitative principle, - if we may so term it. At a very early period, the child hears the sounds addressed to it, and soon attaches ideas to them. It discovers, moreover, that it is capable of producing similar sounds with its own larynx, and that these sounds are understood, and are inservient to the gratification of its wants; and, in this way, speech, as we have elsewhere seen, (vol. i., page 442.) is acquired. The difficulty is to understand in what manner this singular consent is produced. Sir Gilbert Blane has properly remarked, that the imitation of gestures, is, at first sight, less un-

<sup>a</sup> On Man, i. 102, Lond. 1791.

<sup>&</sup>lt;sup>b</sup> The author's General Therapeutics, p. 42; or his General Therapeutics and Mat. Med., Philad. 1843.

accountable than that of sounds; as they are performed by members which are objects of sight, and would seem therefore to be more readily transferable to the corresponding parts of another person: but he probably errs, when, farther on, he remarks, that when children begin to articulate, they first attempt those letters, in the pronunciation of which the motions of the organs are the objects of sight; such as the p and b, among consonants, and the broad a, among the vowels, "giving occasion to a well-known etymology, from the infantile syllables, expressive of father and mother in all languages." We do not think, that this explanation is happy; and have elsewhere attempted to show, that the combination of letters, and the words referred to, are first enunciated, because they are the easiest of all combinations; and that the expressions of mama, papa, &c., are employed long before the child has acquired the power of imitation, and long prior to his attaching the meaning to the words which he is subsequently taught to adopt. It is certainly singular how the child can learn to imitate sounds, where the action of the organs concerned is completely concealed from view. The only possible way of explaining it is to presume, that it makes repeated attempts with its vocal apparatus to produce the same sound which it hears; and that it recollects the sensation produced by the contraction of the muscles when it succeeds, so as to enable it to repeat the contraction of the muscles, and the sensation, at pleasure. This is, however, a case in which volition is actively exerted. We have others, where the action occurs in spite of the individual, as in yawning. We see the action in a second person, and, not withstanding all our attempts to the contrary, the respiratory organs are excited through the brain, and we accomplish the same act. Nay, even thinking of the action will be sufficient to arouse it. Of a like nature to this is the sympathetic contraction of the uterus, which comes on, where a pregnant female is in the lying-in chamber during the accouchement of another, and to which we have referred under the head of Sympathy." Many morbid phenomena are excited in a similar manner: - of these, squinting and stammering are familiar examples. Of 321 cases of squinting, of which the exciting causes were investigated, 61 were found to be produced by imitation.b

## 4. OF THE VARIETIES OF MANKIND.

To determine the number of varieties, into which the great human family may be divided, is a subject which has been considered to belong so completely to the naturalist that we shall pass it over with a brief inquiry.

If we cast our eye over the globe, although we may find, that mankind agree in their general form and organization, there are many points in which they differ materially from each other. With those forms, proportions and colours, which we consider so beautiful in the fine figures of Greece,—to use the language of

<sup>\*</sup> See Bostock's Physiology, ed. cit. p. 759. b Med. Examiner, July 10, 1841, p. 439.

Mr. Lawrence, - contrast the woolly hair, the flat nose, the thick lips, the retreating forehead, advancing jaws, and black skin of the negro; or the broad, square face, narrow oblique eyes, beardless chin, coarse, straight hair, and olive colour of the Calmuck. Compare the ruddy and sanguine European with the jet black African, the red man of America, the yellow Mongolian, or the brown South-Sea Islanders; the gigantic Patagonian, or the dwarfish Laplander; the highly civilized nations of Europe, so conspicuous in arts, science, literature, in all that can strengthen and adorn society, or exalt and dignify human nature, to a troop of naked, shivering, and starved New Hollanders, a horde of filthy Hottentots, or the whole of the more or less barbarous tribes, that cover nearly the entire continent of Africa; and although we must refer them all to the same species, they differ so remarkably from each other as to admit of being classed in a certain number of great varieties; but with regard to the precise number, naturalists

have differed materially.

Whatever changes have been impressed upon mankind can, of course, apply only to the descendants of Noah. The broad distinctions, we now meet with, could not have existed in his immediate family, saved with him at the time of the deluge. They must necessarily have all been of the same race. None of our investigations on this subject can, consequently, be carried back into antediluvian periods. Hence, the region, on which the ark rested, must be looked upon as the cradle of mankind. The question of the original residence of man has frequently engaged the attention of the philologist. It is one, which could be answered positively by the historian only, but unfortunately the evidence we possess of an historical character is scanty in the extreme, and the few remarks, in the sacred volume, are insufficient to lead us to any definite conclusion. As far back as the date of the most remote of our historical records, - which extend to about two thousand years prior to the Christian era, - we find the whole of Asia and a part of Africa, - probably a large part, - peopled by different nations, of various manners, religion, and language; carrying on extensive wars with each other; with here and there, civilized states, possessing important inventions of all kinds which must have required a length of time for discovery, improvement, and diffusion. After the subsidence of the deluge, the waters would first recede from the tops of the highest mountains, which would thus be the earliest habitable; and, in such a situation, the family of Noah probably increased, and thence spread abroad on the gradual recession of the waters. The earliest habitable region was probably the elevated region of middle Asia, - the loftiest in the world, - not the summits, which would be unsuitable, in every respect, for human existence, but some of the lofty plains, such as that, of which the well-known desert Kobi or Schamo forms the highest point, and whence Asia sinks gradually towards the four quarters, and the great mountain chains proceed that intersect Asia in every direction. This has been suggested by Herdera and Adelungb as the cradle of the human race. In the declivities of this elevated region, and of its mountain chains, all the great rivers arise that flow on every side through this division of the globe. After the deluge, it would therefore soon become dry, and project, like an extensive island, above the flood. The cold and barren elevation of Kobi would not itself have been well adapted for the continued residence of our second parents, but immediately on its southern side lies the remarkable country of Thibet, separated by lofty ridges from the rest of the world, and containing within itself every variety of climate. Although on the snow-capt summits the severest cold perpetually prevails, summer eternally reigns in the valleys, and well-watered plains. The rice, too, the vine, pulse, and a variety of other productions of the vegetable kingdom, which man employs for his nutrition, are indigenous there; and those animals are found in a wild state which man has domesticated and taken along with him over the earth; -the ox, horse, ass, sheep, goat, camel, swine, dog, cat, and even the valuable reindeer, - his only friend and companion in the icy deserts of polar countries. Zimmermann, indeed, asserts, that every one of the domesticated animals is originally from Asia. Close to Thibet, and immediately on the declivity of this great central elevation, is the charming region of Kaschemire, the lofty site of which tempers the southern heat into a protracted spring.

The probabilities in favour of the cradle of mankind having been situate to the south of the elevated region of middle Asia are considered to be strengthened by the circumstance of the nations in the vicinity possessing a rude, meagre and imperfect language, such as might be imagined to have existed in the infancy of the human intellect and of the world. Not less than two hundred millions of people are found there, whose language appears to be nearly as simple as it must have been soon after its formation. Kaschemire, by reason of the incessant changes, which it has exexperienced in ancient and modern times, has, indeed, kept pace with the rest of the world in the improvement of its language; but not so, apparently, with Thibet - its neighbour - and with China, and the kingdoms of Ava, Pegu, Siam, Tunkin, and Cotschinschina. All these extensive countries and these alone in the known world, according to Adelung, betray the imperfection of a newly-formed or primitive language. As the earliest attempt of the child is a stammering of monosyllabic notes, so, - says that eminent philologist, - must have been that of the original child of nature; and, accordingly, the Thibetans, the Chinese, and their two neighbours to the south continue to stammer monosyllabically, as they must have been taught, thousands of years ago, in the infancy of their race. "No separation of ideas into certain

Ideen zur Philosophie der Geschichte der Menschheit, Riga und Leipz. 1785-1792.
 Mithridates oder Allgemeine Sprachenkunde, Berlin, 1806-1817, Erster Theil.
 Einleitung.
 Geograph. Geschichte der Menschen, u. s. w. Leipz. 1778.

VOL. II. - 50

classes, whence arose the parts of speech in cultivated languages. The same sound, which denotes joyful, signifies joy and to gladden, and this in every person, number and tense. No art, connexion, or subordinate ideas are united to the rude, monosyllabic root, thereby communicating richness, clearness and euphony to their meagre tongue. The rude, monosyllabic, radical ideas are placed, perhaps broken, and detached from each other, the hearer being left to supply the intermediate ideas. As the monosyllable admits of no inflection, the speaker either makes no distinction between cases and numbers, or he seeks for aid, in cases of great necessity, in circumlocution. The plural he forms, like the child, either by repetition, — tree, tree, — or by the addition of the words much or more, as tree much, tree more. I much, or I more is the same to him as we." From these and other circumstances, Adelung infers, that these monosyllabic languages are primitive and the honourable ancestors of all others; but the argument is more plausible perhaps than sound. Is has been correctly remarked by the author's distinguished friend, Mr. Duponceau, that, in all languages, there is a strong tendency to preserve their original struc. ture, and that from the most remote period, to which the memory of man can reach, a monosyllabic language has never been known to become polysyllabic, or conversely. Adelung farther infers, that the immediate descendants of Noah originally occupied the favoured region which has been described, and, as population increased, spread into the neighbouring districts, selecting, by preference, the near and charming regions of the south, east, and west. Hence we find, in the countries immediately bordering on Thibet, the earliest formed states, and the oldest civilization. History refers us to the east, for the primordial germs of most of our ideas, arts and sciences, whence they subsequently spread to the countries farther to the west - to Media, Persia, and western

1. Division of the Races. - It is probable that from western Asia, the sons of Noah, - Shem, Ham and Japheth, - branched off in various directions, so as to constitute the three distinct stocks which are found to have divided the old world from time immemorial. These three are-1, the White, Caucasian, Arabico-European, or European; 2, the Olive, Mongolian, Chinese, Kalmuck, or Asiatic; and 3, the Negro, Ethiopian, Africian, Hottentot, &c., each of which has its own principal habitat; - the white being found chiefly in Europe and Asia Minor, Arabia, Persia, and India as far as the Ganges, and in North Africa; the Mongol occupying the rest of Asia, and having its focus on the plateaux of Great Tartary and Thibet; and the negro race covering almost the whole of Africa, and some of the Isles of New Guinea, the country of the Papuas, &c. The white or Caucasian variety are supposed to be the descendants of Japheth ("audax Japeti genus," Horace); the Asiatic the descendants of Shem; whilst Ham is regarded as the parent of the African. These three races. - the

Caucasian, Negro, and Mongolian, - are alone admitted by Cuvier. Blumenbach, whose classification will serve our purpose as well as any of the others to which reference will be made presently, admits four, - the Caucasian, Ethiopian, Mongolian, American and Malay.

a. Caucasian Race. - The Caucasian race is chiefly distin-

guished by the elegant form of the head, which approximates to a perfect oval. It is also remarkable for variations in the shades of the complexion and colour of the hair. From this variety, the most civilized nations have sprung. The name Caucasian was given to it from the group of mountains between the Caspian and the Black Sea, - tradition seeming to refer the origin of this race to that part of Asia. Even at the present day, the peculiar characteristics of the races are found in the highest perfection amongst the people who dwell in the vicinity of Mount Caucasus, - the Georgians and Circassians, -- who are esteemed the handsomest natives of the earth.



Caucasian Variety.

The marginal figure is given by Blumenbach as a specimen of the Caucasian race, near the original residence whence the epithet is derived. It represents Jusuf Aguiah Efendi, formerly ambassador from the Porte to London.

The Caucasian race has been subdivided into several great nations or families: - 1. The Arabs, comprising the Arabs of the desert or the Bedouins, the Hebrews, the Druses and other inbabitants of Libanus, the Syrians, Chaldwans, Egyptians, Phoenicians, Abyssinians, Moors, &c. 2. The Hindoos on the European side of the Ganges ; - as the inhabitants of Bengal, of the coasts of Coromandel and Malabar, the ancient Persians, &c. Scythians and European Tartars, comprising also the Circassians, Georgians, &c. 4. The Kelts or Celts, a dark-haired race, whose precise origin is unknown, but presumed to be Indian. The descendants of this race are the Gauls, Welsh, Rhætiaus, &c., &c.;

<sup>2</sup> Règne animal; also, Elémens de Zoologie, par H. Milne Edwards, p. 259, Paris,

1834 b Handbuch der Naturgeschichtes, § 52, Götting. 1791; and De Generis Humani Varietat. Nativ., Götting. 1777. See, also, Buffon, Hist. Nat. iii. 371; Beclard, Elemens d'Anatomie Générale, p. 110, 2de édit. Paris, 1827; Dumeril, Zoologie Analytique, Paris, 1806; and Lawrence's Lectures on Physiology, &c., Lond. 1839. and lastly, the Goths, a fair-haired race, the ancestors of the Ger

mans, Dutch, Swedes, Danes, &c.

That the time of the first peopling of the European countries must have been very remote is exhibited by the fact, that at the dawn of history, the whole of Europe, from the Don to the mouth of the Tagus, was filled with nations of various physical characters and languages, bearing striking marks of intermixture and modification. At this period, there were, in Europe, at least six great nations. 1st. The Iberians with the Cantabri, in Spain, in a part of Gaul, and on the coasts of the Mediterranean as far as Italy. 2dly. The Kelts, in Gaul, in the British Isles, between the Danube and the Alps, and in a part of Italy. 3dly. The Germani or Goths, between the Rhine, the Danube and the Vistula. 4thly. The Thracians, with the Illyrians, in the southeast of Europe, and in western Asia. 5thly. The Sclavi, in the north: and 6thly. The Fins in the northeast. It is not improbable, that these different races migrated from Asia in the above order - such at least is the theory of certain historians and philologers, and there is some reason for adopting it. They who migrated first would probably extend their wanderings until they were arrested by some invincible obstacle, or until the arrival of fresh tribes would drive them onwards farther and farther towards the west. In this way, they would ultimately reach the ocean, which would effectually arrest their farther progress, unless towards the south and the north. The descendants of the ancient Iberians do now actually occupy the west of Spain, - the residence probably of their forefathers. Nearly about the same time, perhaps, as the Iberians undertook their migration, the Kelts, a populous tribe, migrated from some part of Asia, and occupied a considerable portion of middle Europe. To these succeeded the Goths, to the north, and the Thracians to the south; whilst the Sclavi, the last of the Asiatic emigrants, wandered still farther north. It is not easy to determine the precise link occupied by the Fins in this vast chain of nations, They were first known to history as a peculiar people in the north of Europe; but whence they proceeded, or whether they occupied their position to the north of the Germani from choice, or were urged onwards by their more powerful neighbours, we know not. So long as there was sufficient space for the nations to occupy, without disturbing the possessions of their neighbours, they probably kept themselves distinct; but as soon as the land was filled, a contest arose for the possession of more extensive or more eligible regions; wars were, consequently, undertaken, and the weaker gradually yielded their possessions, or their sovereignty, to the stronger. Hence, at the very dawn of history, numerous nations were met with, amalgamated both in blood and language; - for example, the Kelto-Iberians of Spain; the Belgæ or Kymbri of Gaul and Britain; the Latins, and other nations of Italy, and probably many, whose manners, characters, and language had become so melted into each other as to leave little or no trace of the original constituents. The Letti, Wallachians, Hungarians, and Albanians of eastern Europe, are supposed to afford examples of such amalgamation, whilst the mighty Sclavonic nation has swallowed up numbers of less powerful tribes, and annihilated even their names for ever. This it is, which frequently embarrasses the philological historian; and prevents him, without other evidence, from deducing with accuracy the parent stocks or the most important components in ethnical admixtures.

Dr. Morton, in his splendid and valuable contribution to the Natural History of Man, subdivides the Caucasian race into seven families: — the Caucasian, the Germanic, the Celtic, the Arabian,

the Libyan, the Nilotic and the Indostanic.

b. Ethiopian Race. — The Negro, African, Ethiopian or Black man of Gmelin occupies a less extensive surface of the globe, embracing the country of Africa which extends from the southern side of Mount Atlas to the Cape of Good Hope. This race is evidently of a less perfect organization than the last, and has some characteristics, which approximate it more to the monkey kind. The forehead is flattened and retiring; the skull is smaller, and holds from four to nine ounces of water less than that of the European. On the other hand, the face, which contains the organs of sense, is more developed, and projects more like a snout. The lips are large; the cheek bones prominent; the temporal fossæ hollower: the muscles of mastication stronger; and the facial angle is smaller; —the head of the negro, in this respect, holding a middle place between the Caucasian and the orang-outang. The nose is expanded; the hair short and woolly, very black and frizzled. Skin black. This colour is not, however, characteristic of the race, as the Hottentots and Caffres are yellow.

The marginal figure is the head of J. J. E. Capitein, selected by Blumenbach as the representative of his race. He was an intel-

ligent negro, and published several sermons and other works in Latin and Dutch. His portrait was taken by Van Dyk. This case of great intelligence in the negro is almost unique; and it exhibits what may be expected from him under favourable circumstances. almost all situations in which he is found, it is in a state of slavery and degradation, and no inference can be deduced regarding his original grundkraft — as the Germans call it - or intellectual capability. Hayti has afforded numerous



Negro Variety.

examples of the sound judgment, and even distinguished ability
Crania Americana, p. 5, Philad. 1839.

with which her sable inhabitants are capable of conducting, not only the municipal, but the foreign concerns of a considerable



Mongolian Variety.

community. It must be admitted, however, that from organization, this race would seem to be, cæteris paribus, less fitted for intellectual distinction than the Caucasian, and singular differences have been observed between them and the Caucasian race, when exposed to similar mental and physical influences.

The Ethiopian race is subdivided by Dr. Morton<sup>c</sup> into six families: the Negro; the Caffrarian; the Hottentot; the Oceanic Negro; the Aus-

tralian, and the Alforian.d

c. Mongolian Race. — The Mongolian, Asiatic, Kalmuck or Chinese race, the brown man of Gmelin, is recognised by prominent and wide cheek bones; flat, square visage; small and oblique eyes; straight and black hair; scanty beard, and olive complexion.

The marginal head is from Blumenbach. It is that of Feodor Ivanowitsch, a Kalmuck, given by the Empress of Russia to the hereditary Princess of Baden. He was educated at Carlsruhe, and was a most distinguished painter at Rome. The portrait was

sketched by Feodor himself.

The Mongols are spread over the central and eastern parts of Asia, with the exception of the peninsula of Malacca. They likewise stretch along the whole of the Arctic regions, from Russia and Lapland to Greenland, and the northern parts of the American continent, as far as Behring's Straits, — the Laplanders and Esquimaux being evidently of the same race as the Koriaks, Kamtschadales, Japanese, &c., of the Asiatic continent.

Dr. Morton includes in this race five families: -- the Mongol-Tartar; the Turkish; the Chinese; the Indo-Chinese, and the

Polar.

a See vol. i., p. 295.

b See abstract of a memoir by Dr. B. H. Coates, "on the effects of secluded and gloomy imprisonment on individuals of the African variety of mankind in the production of disease," in Proceedings of the American Philosophical Society, vol. 3, p. 143, Philad. 1843.

d The Alfores or Horaforas are considered aboriginal to many islands of the Indian Archipelago. They are most numerous in New Guinea, the Moluccas, and Magindano: in Celebes, they are said to be sometimes as fair as the Malays, and the savage Dyaks of Borneo appear to belong to the same family. Dr. Morton thinks it not improbable, as suggested by Dr. Prichard, that the Alfores are but a branch of the Australian stock. — Crania Americana, p. 95.

d. American Race. — The American race, red man of Gmelin, or more properly brown man, differs greatly in stature, colour, and physiognomy in various parts of the continent; but his medium



height corresponds with that of the European. His colour is from a cinnamon-brown to a deep copper. The hair is almost always black, straight and stiff. The features are large and strongly marked, except the eyes, which are commonly deep-seated, or sunk in large sockets. The forehead is generally low, somewhat compressed at the sides, and slightly retreating. Facial angle about 80°. Nose generally considerably raised from the face, sometimes arched; cheek bones high, and widely separated; angle of the jaw broad, and chin square. The accompanying head is that of Ongpatonga, (Big Elk,) chief of the Omawhaw Indians.

Dr. Morton divides the American race into two families: — the American and the Toltecan; the latter embracing the civilized

nations of Mexico, Peru, and Bogota.b

e. Malay Race. — The Malay or Australian race, the Tawny man of Gmelin, is admitted by many naturalists, owing to the difficulty of referring it either to the Caucasian Indian, or to the Chinese Mongolian, situate in its vicinity. This Malay variety extends from Malacca to the most remote islands of the great Indian and Pacific Ocean, from Madagascar to the Maldives, in-

From Godman's American Natural History, Philad. 1826-1828.

b Crania Americana, p. 83, Philad. 1839.

clusive; inhabits Sumatra, Java, Borneo, Celebes, and the adjacent islands; the Molucca, Ladrone, Philippine, Marian, and Caroline groups; New Holland, Van Diemen's Land, New Guinea, New Zealand, and the various islands scattered through the South Sea. It is termed Malay, because supposed to have proceeded originally from the Peninsula of Malacca, and to have spread thence over the adjacent islands, - a supposition, which is not confirmed by history: on the contrary, according to Mr. Marsden, it is clearly demonstrated, that the Malays went from Sumatra to Malacca in the twelfth century. No well-marked common characters can be assigned to this variety; for, under the term Malay, races are included, which seem to differ materially from each other; so much so, indeed, as to induce many naturalists to refuse the admission of the Malay as a distinct variety. Their colour may be said to be brown, in various shades, from a light tawny, to almost a black: the forehead is low and round; the nose full and broad; nostrils wide; mouth large; hair thick, crisp, and always black, as well as the iris. Fig. 183 exhibits an individual of this race: it is the head of a New Zealand chief. Cuvier, Rudolphi, Virey and others consider the Malay variety to be a mixture of the Mongol of Asia and the negro of Africa.

Dr. Mortone divides the Malay race into two families - the

Malay, and the Polynesian.

In New Guinea, and the small islands around, the *Papuas* are found, who resemble the negroes yet more strongly; and similar races are met with in the Archipelago of the Holy Ghost, and in the isles of Andaman and Formosa. They are presumed to belong really to the negro race, and to have descended perhaps from individuals of that variety, who have wandered, or been driven, from their original settlements. Some of them resemble the Guinea negro in every particular.<sup>4</sup>

Of late years, many other races have been added to those admitted by Blumenbach; especially by Messrs. Virey, Desmoulins, Malte-brun, and Bory de Saint-Vincent. Gerdy and Broc, — like their predecessors, — dissatisfied with the divisions that have been adopted by naturalists, have taken colour as the basis of theirs. For race they substitute sub-genus; of which they admit

four, — the white, yellow, negro or black, and red.

The various classifications exhibit the vacillation, which yet exists regarding the precise number of races that should be admitted. Every division must necessarily be arbitrary, and the indi-

\* Op. cit. b Grundriss der Physiologie, th. 1, Berlin, 1821. c Op. cit. p. 6.

d Morton, op. cit. p. 92.

· Histoire Naturelle du Genre Humain. 2de édit. Paris, 1824.

Hist. Naturelle des Races Humaines. Paris, 1826.

s Geographie Universelle, Paris, 1816.

h L'Homme, Essai Zoologique sur le Genre Humain. 2de édit. Paris, 1827. i Physiologie Médicale, tom. i.

Essai sur les Races Humaines, p. 25, Paris, 1836.

viduals composing each variety are far from being alike. We find the greatest diversity, for example, amongst the nations of the Caucasian variety, and even amongst any of its subdivisions. The Frenchman can be distinguished from the German, the Spaniard from the English, &c., and if we were to push the system of subdividing, which appears at present to be fashionable, we might constitute almost every nation of the globe into a distinct variety.

2. Origin of the Different Races. - It has been an oft agitated question, whether all the varieties amongst mankind must be regarded as belonging to the same species, - the differences which we observe having been accomplished by extraneous circumstances, acting through a long succession of ages; or whether they must not be regarded as distinct species, ab origine. By many, the discussion of this subject has been esteemed not only unnecessary but profane, inasmuch as the sacred historian has unequivocally declared, that all mankind had a common origin. We have already remarked, however, that this is not a question, which concerns our first parents, but belongs exclusively to the family of Noah; for, in his descendants, all these varieties must necessarily have occurred. From the part of Asia, previously described, his immediate descendants probably spread abroad to the north and the south, the east and the west; Europe being peopled by the migratory hordes, which proceeded towards the northwest, and Africa by those from southwestern Asia. These migrations probably all took place by land, except in the case of our own continent, where a slight sea-voyage, of not more than thirty-nine miles, across Behring's Straits, even in frail vessels, would be sufficient to transport the emigrants without much risk of misadventure; and even this short voyage would be rendered unnecessary during the winter season, the strait being solidified into a continuous mass of ice. Europe probably received its inhabitants long before navigation existed to any extent. Subsequently, when a coasting trade was first established, - to which the enterprise of nations would necessarily be limited in the first instance, until by improved vessels and a better system of management they were enabled to brave the terrors of the ocean, and undertake their adventurous voyages of discovery, - many of the coasts, especially of the Mediterranean, received swarms of emigrants, a circumstance, which accounts for the motley population, observable, at an early period, in these regions. Carthage, we know, was settled by the Phænicians, and Southern Italy and Spain, in this manner, received their Greek colonies. Dr. Coplande has even

Art. Menschen-Varietaten, in Pierer's Anat. Phys. Real Wörterbuch, v. 153.

b Lawrence, op. citat.; Weber's Hildebrandt's Handbuch der Anatomie, iv. 529,
Braunschweig, 1832; J. Cloquet, Anatomie de l'Homme, tom. i. pl. 29, Paris, 1821;

Braunschweig, 1832; J. Cloquet, Anatomie de l'Homme, tom. i. pl. 29, Paris, 1821; Lacepède, art. Homme, in Dict. des Sciences Naturel. xxi. 382; J. E. Doornik, Wysgeerig Natuurkundig Onderzoek aangande den Oorsprongliken Menschen de Oorspronglike Stammen van deszelfs Geslacht. Amsterd. 1828; Elliotson's Human Physiology, 1062, Philad. 1840; Prichard, Researches into the Physical History of Mankind, 3d edit. vol. i. Lond. 1836; and Nat. History of Man, Lond. 1843.

Notes to translation of Richerand's Physiology.

expressed his belief in the view, that this continent was visited "by Phænician navigators, the greater part of whom settled in it, particularly in Mexico; and that the imperfect navigation of that era prevented many of the adventurers, if not all of them, from returning." The notion is, however, altogether hypothetical.

The greatest difficulty has been,—to comprehend how the Caucasian and Ethiopian varieties could have originated from the same source. The other varieties of mankind, if we exclude the negro, could be referred, without much hesitancy, to the same primitive stock,—the changes being caused by adventitious circumstances operating for an immense period,—but it has seemed to many naturalists impossible to suppose, that the characters of the negro could, by any process, become converted into those of the European, or conversely.

The people of antediluvian times probably possessed but few physical differences, — constituting one large family, modified, perhaps, to a certain extent, by circumstances, but not materially; the two antithetical races, — the white and the black, — first arising in postdiluvian periods. If we adopt this view, the question, regarding the difference of species between the white and the black, will require no agitation. But how are we to explain the essential differences as to form and colour, which we notice amongst the

nations of the earth?

In the infancy of anthropology, it was asserted, that the white races inhabit the cold and temperate regions of the earth, whilst the tawny and darker races are situate under a more vertical sun. Within certain limits, the sun is doubtless possessed of the power of modifying the colour. The difference between one, who has been for some time exposed to the rays of a tropical sun, and his brethren of the more temperate climates, is a matter of universal observation. The inhabitant of Spain is, in this way, distinguishable from the French, German, English, &c.; and hence we can understand, why the Southern Asiatic and African women of the Arab race, when confined within the walls of the seraglio, may be as white as the fairest Europeans. There are many exceptions, however, to the notion which has prevailed, that there is an exact ratio between the heat of the climate and the blackness of the skin. For example, at the extreme north of Europe, Asia, and America, we find the Laplanders, Samoiedes, Esquimaux, &c., with the skin very brown, and the hair and iris black; whilst, in the vicinity of the Laplanders, are the Fins, — a people of large stature compared with the Laplanders, with fair skins and bluishgray eyes. In the same manner, to the south of the Greenlander, - of short stature, brown skin, and dark hair, - is the tall and fair Icelander. The Kelt of Wales, and of the western coast of Ireland, of the north of Scotland, and of the west of Bretagne, is still distinguished, by his dark hair and eyes, from the light-haired descendants of the Goth, - the German and the Scandinavian. Many distinct tribes exist in the interior of Africa, having a red or copper hue, with lank black hair, and in the midst of the black varieties of their species. A similar fact was observed by Humboldt in different parts of South America. Again, the negro race is not always found in the torrid zone. On our own continent, none have ever been met with, except what have been imported: and these, after repeated descents, have still retained their original character; whilst, as we have seen, negroes are met with in Australia under a climate as cold as that of Washington. The fact of the slight mutation, effected by ages on the character of a race, is strikingly shown by the circumstance to which we have before referred, - that in some of the monuments of Egypt, visited by Belzoni and Champollion, representations of the negro, presumed to be upwards of three thousand years old, exhibit the features to be almost identical with those of the negro of the present day. The Jew affords an example of the same immutability, as well as the Esquimaux, who strikingly retain the evidence of their Kalmuck origin. Complexion, and, to a certain extent, the figure are doubtless modified by climate, but the essential characters of the organization remain little if at all changed. Volney has fancifully supposed, that the elongated visage of the negro is owing to the wry face habitually made under exposure to the rays of the sun. Independently, however, of the objection, that this would be wholly insufficient to account for the striking peculiarities of the negro head, it has already been remarked, that these peculiarities do not exist among other races, inhabiting equally hot climes; and that the negro himself is not confined to those climates, and ought, consequently, to lose the museau or snout, when the country is so cool as to render the wry face or moue unnecessary.2

It may then, we think, be concluded, that the evidence in favour of the colour of the negro, of the red man, or of the tawny, being produced directly or indirectly by the solar rays, is insufficient to establish the point. One important argument in the negative is the fact, that in all cases, the children are born fair, and would continue so, if not exposed to the degree of solar heat, which had

produced the change in their progenitors.b

In addition to the influence of temperature, and climate, that of food, and of different manners and customs has been frequently invoked, but without any precise results being deduced. The effect of difference in manners and customs is shown in the results of domestication on animals, — as in the case of the wild and the disciplined horse; of the bison and the ox, — which last is regarded as the bison in a state of tameness. The precise cause of such modification we know not. It is not confined to the animal, but is signally evidenced in the vegetable. The flower of the forest, when received into the parterre and carefully nurtured, will develope itself in such a manner as to be with difficulty recognizable. The change seems to be produced by variation of climate and nutrition, but in what precise manner we know not.

The powerful modifying influence of locality on the develop-

Art cit. in Dict. des Sciences Médicales.
Fletcher's Rudiments of Physiology, part iii. p. 69, Edinb. 1837.

ment of the moral and physical powers has been more than once referred to. Perhaps the most remarkable examples are met with at the base of lofty mountains, particularly of the Alps, and in some of the unhealthy districts of France especially. One of these is cretinism, a singular case of malformation, with which we are happily unacquainted in the United States. This is a state of idiocy, which is remarkable in its subjects being always more or less deformed, and in its appearing to originate from local influences. The crétin has every characteristic of the idiot; and, in addition, is often distinguished by a large goître or swelling of the thyroid gland; by soft flabby flesh; and by shrivelled, yellowish, or pale and cadaverous skin, covered, at times, with filthy cutaneous eruptions. The tongue is thick and pendant; the eyelids large and projecting; the eyes gummy, red and prominent; the nose flat; the mouth gaping and drivelling; the face puffy, and at times, violet-coloured, and the lower jaw elongated. In several, the forehead is broad inferiorly, and flattened and retreating above, giving the cranium the shape of a cone rounded towards its smaller extremity. The stature of the cretin is generally small, scarcely ever exceeding four feet and a few inches; the limbs are frequently malformed, and almost always kept in a state of flexion. All the cretins are not affected with goître. Some have large and short, whilst others have thin, and long necks. Like the idiot, the cretin does not generally live long, scarcely ever surviving the thirtieth year.

Authors have differed in opinion as to the causes of this deplorable condition. It is observed almost exclusively in the deep and narrow valleys at the foot of lofty mountains, and in mountain gorges. Hence, it is common in that part of the Alps called the Valais or Wallais; in the valley of Aost, La Maurienne, &c. It is met with, too, at the foot of the mountains of Auvergne, the Pyrenees, the Tyrol, &c. De Saussure, Esquirol, Fodéré, Rambuteau,—and all who have had an opportunity of observing these miserable wrecks of humanity,—believe, that the great cause is the concentrated, moist, and warm air, which prevails throughout almost the whole of the year, in the valleys and mountain gorges where it is found to exist.<sup>a</sup> That it is dependent upon the locality is obvious, but how this acts, we know no more than we do of the causes of other endemic affections,—intermittent fever, pellagra,

beriberi, &c., &c.b

After all, perhaps, the strongest arguments, — in favour of extraneous circumstances occasioning, in the lapse of ages, the different varieties, which we observe in the great human family, — are those derived from the changes that must have occurred amongst many of the inferior animals. The dog, in its wild state, has always

b See the author's Practice of Medicine, 2d edit., vol. i., Philad. 1844.

<sup>&</sup>lt;sup>a</sup> Dr. James Johnson, Change of Air, Amer. Edit. p. 61, New York, 1831; and Assistant Surgeon J. M'Clelland, in Dublin Journal of Medical and Chemical Sciences, May, 1837, p. 293; De Saussure, Voyages dans les Alpes, ch. xlvii. § 1050; Reeve, in Philos. Transact. for 1808, p. 111; Prichard on Insanity, p. 318, Lond. 1835; and Hodgkin, Catalogue of Guy's Museum, §v. part ii. Lond. 1829.

pretty nearly the same characters; being covered with hair of the same colour; the ears and tail, and limbs, having the same shape; and it exhibits, apparently, the same powers and instincts; but, on this matter, our knowledge, derived from observation, is necessarily limited. Yet what a number of varieties are observed in this animal when it becomes domesticated; and how different from each other, in shape, colour, character of skin, and instincts, are the spaniel, hound, greyhound, pointer, mastiff, terrier, cur, pug, lapdog, &c.; differences certainly as great as between the varieties of mankind. These differences, it is presumable, may have been produced partly by the occurrence of accidental varieties, affecting perhaps a whole litter, - male and female; so that if these, again, were to be coupled, the variety, thus accidentally caused, might have become permanent. Such accidental varieties occasionally occur in the human species, but they are soon lost, in consequence of the wise law that prevents individuals, within certain degrees of consanguinity, from marrying. It is by no means uncommon, for example, for different children of the same family, from some accidental cause, to be born with six fingers. The author has met with two families in each of which more than one individual was thus circumstanced; and Sir Anthony Carlisle has detailed the remarkable case of a family from this continent, where the superfluity extended, in the case of a female, to two thumbs on each hand, and to six toes on each foot. She married and had several children, who, in their turn, became parents, and transmitted the peculiarity to the children to the fourth generation. Now, if the members of this family had continued to marry in and in, a new race of individuals might have been perpetuated, possessing the unnecessary additions in question. Under existing laws and customs, it must always happen, that where such peculiarity exists in one parent only, it must soon become extinct; yet, as we have seen, it may be pertinacious enough to persist for some generations. Fortunately, also, it happens, that, as a general rule, no change, which occurs accidentally in the parent after birth, is liable to be extended to the progeny.b Were the rule other than it is, it will be at once seen, the most strange and innumerable varieties of races would exist. Where a limb had become distorted or amputated, a stock of onelimbed animals would be formed; the docked horse would propagate a mutilated colt; the operation of circumcision, performed on one parent, ought to be sufficient for the whole of his descendants, &c., &c.

<sup>a</sup> Philosoph. Transact. for 1814. See, also, Prichard's Researches into the Physical History of Mankind, i. 245 and 352, Lond. 1836; and his Nat. History of Man. Lond. 1843; art. Generation, Cyclop. of Anat. and Physiol. P. xiii. p. 473, Feb. 1838; Morand, Mémoir. de l'Académ. pour 1780, p. 137; Isidore St. Hilaire, Histoire Générale et Particulière des Anomalies de l'Organisation, p. 681, Paris, 1832; and Walker, on Intermarriage, Amer. Edit., New York, 1839.

The author has been informed by one who has had ample opportunity for observation, that when two deaf and dumb individuals marry, the defect is rarely, if ever, observed in their progeny: but that the contrary is the case with the born blind. In addition to this mode of accounting for the great number of varieties in animals of the same species, the influence of difference in manners and customs, which we have already considered, has been adduced; and it has been conceived, that the effect of civilization and refinement on the human race may be analogous to that of domestication on the inferior animals. This kind of influence is said to be particularly observable amongst the inhabitants of Hindusthan, where, in consequence of the division into castes, the same condition of life, and the same occupation are continued without change through many successive generations. The artisans, who are a superior class, are of a manifestly lighter complexion than the tillers of the soil; and in many of the islands of Polynesia, the same difference exists between the classes as in Hindusthan.

The believers, then, in the Mosaic account of the creation, and the deluge, must regard all the varieties of mankind to have descended from the same family, - that of Noah, - and the different changes, which have been impressed upon their descendants, to be results of extraneous influences acting through a long succession of ages, added to the production perhaps of accidental varieties, which may have occurred in the very infancy of postdiluvian existence, when the intermarriage of near relations was unavoidable, and when such varieties would necessarily be perpetuated. The race of Ham appears to have been separated, if not wholly, at least in part, from their brethren by the malediction of Noah; and, whether we consider that a physical alteration was comprised in the malediction, or that such alteration might occur accidentally, as in the cases of those with supernumerary toes and fingers, the very fact of intermarriage with the descendants of the other sons of Noah, being prevented by the curse pronounced on Ham (for many commentators read Ham for Canaan,) would necessarily lead to a perpetuation of the adventitious modification.

But, it has been asked, if all mankind have descended from one family, which of the varieties now extant, must be regarded as their representative? On this we have nothing but conjecture to guide us. It has been supposed, by some, to be more probable, that the changes, induced upon mankind, have been in consequence of the progress from a state of barbarism to one of refinement, than the reverse; and hence, it has been conceived, that the variety ought to be considered primary, which, through all the vicissitudes of human affairs, has remained in the most degraded condition, and which, in its structure, differs most materially from the variety that has uniformly enjoyed the greatest degree of civilization. Upon this principle, the Ethiopian would have to be regarded as the type of our first ancestors, and such is the opinion of Prichard, and of Bostock. Blumenbach, however, maintains the converse view. Bishop Heber, again, suggests, whether the hue of the Hindoo, which is a brownish-yellow, may not have been that of our first parents, whence the transition, he thinks, to the white and black varieties, might be more easy and comprehensible. Philology occasionally aids us in our historical deductions, but the evidence, afforded by it, has to be received with caution. The Hebrew names, like all original appellations, in perhaps all languages, are generally expressive, and therefore worthy of consideration in questions of this nature. The Hebrew word Adam, ( ) is not only the name of the first man, but it signifies man in the abstract, corresponding to the Greek, ardpares, and the Latin, Homo. We are told, in the sacred volume, that, "in the day that God created man, in the likeness of God made he him; male and female created he them; and blessed them, and called their name Adam, in the day when they were created. The word Adam is derived from a Hebrew root, ( ) signifying "to be red," and, accordingly, it is not improbable, that the original hue of the first man was of that character.

The remarks already made, render it unnecessary to inquire into the mode, in which, according to the notions of Blumenbach,2 of Dr. S. S. Smith, or of Dr. Rush, the black colour of the Ethiopian has been produced. Blumenbach imagined, that the heat of the climate gives rise to an excessive secretion of bile; that in consequence of the connexion, which exists between the action of the liver and the skin, an accumulation of carbonaceous matter takes place in the cutaneous vessels, and that this process, being continued for a succession of ages, the black colour of the skin becomes habitual. Dr. Smith had a similar opinion; he thought, that the complexion in any climate will be changed towards black, in proportion to the degree of heat in the atmosphere, and to the quantity of bile in the skin; and, lastly, Dr. Rush, in one of the strangest of the strange views, which have emanated from that distinguished, but too enthusiastic, individual, has attempted to prove, "that the colour and figure of that part of our fellow-creatures, who are known by the epithet of negroes, are derived from a modification of that disease which is known by the name of leprosy."

The following are his deductions from "facts and principles" urged in a communication, read before the American Philosophical Society in 1792:—"1. That all the claims of superiority of the whites over the blacks, on account of their colour, are founded alike in ignorance and inhumanity. If the colour of negroes be the effect of a disease, instead of inviting us to tyrannize over them, it should entitle them to a double portion of our humanity, for disease all over the world has always been the signal for immediate and universal compassion. 2. The facts and principles which have been delivered, should teach white people the necessity of keeping up that prejudice against such connexions with them, as would tend to infect posterity with any portion of their disorder. This may be done upon the ground I have mentioned without offering violence to humanity, or calling in question the

De Gener. Human. Variet. Nat., edit. 3tia, p. 66, Gotting. 1795.

An Essay on the Causes of the Variety of Complexion and Figure in the Human Species, Philad. 1787.

Transact. of the American Philosoph. Society, vol. iv.

sameness of descent, or natural equality of mankind. 3. Is the colour of the negroes a disease? Then let science and humanity combine their efforts, and endeavour to discover a remedy for it. Nature has lately unfurled a banner upon this subject. She has begun spontaneous cures of this disease in several black people in this country. In a certain Henry Moss, who lately travelled through this city, and was exhibited as a show for money, the cure was nearly complete. The change from black to a natural white flesh colour began about five years ago at the ends of his fingers, and has extended gradually over the greatest part of his body. The wool, which formerly perforated the cuticle, has been changed into hair. No change in the diet, drinks, dress, employments, or situation of this man had taken place previously to this change in his skin. But this fact does not militate against artificial attempts to dislodge the colour in negroes, any more than the spontaneous cures of many other diseases militate against the use of medicine in the practice of physic. To direct our experiments upon this subject I shall throw out the following facts: - 1. In Henry Moss the colour was first discharged from the skin in those places, on which there was most pressure from clothing, and most attrition from labour, as on the trunk of his body, and on his fingers. The destruction of the black colour was probably occasioned by the absorption of the colouring matter of the rete mucosum, or perhaps of the rete mucosum itself, for pressure and friction, it is well known, aid the absorbing action of the lymphatics in every part of the body. It is from the latter cause, that the palms of the hands of negro women, who spend their lives at a washing-tub, are generally as fair as the palms of the hands in labouring white people. 2. Depletion, whether by bleeding, purging, or abstinence, has been often observed to lessen the black colour in negroes. The effects of the above remedies in curing the common leprosy satisfy me that they might be used, with advantage, in that state of leprosy, which I conceive to exist in the skin of the negroes. 3. A similar change in the colour of the negroes, though of a more temporary nature, has often been observed in them from the influence of fear. 4. Dr. Beddoes tells us that he has discharged the colour in the black wool of a negro by infusing it in the oxygenated muriatic acid, and lessened it by the same means in the hand of a negro man. The land-cloud of Africa, called by the Portuguese Ferrino, Mr. Hawkins tells us, has a peculiar action upon the negroes in changing the black colour of their skins to dusky gray. Its action is accompanied, he says, with an itching and prickling sensation upon every part of the body, which increases with the length of exposure to it so as to be almost intolerable. It is probably air of the carbonic kind, for it uniformly extinguishes fire. 5. A citizen of Philadelphia, upon whose veracity I have perfect reliance, a assured me that he had once seen the skin of one side of a "Mr. Thomas Harrison."

LIFE. 605

the cheek inclining to the chin, and of part of the hand in a negro boy, changed to a white colour by the juice of unripe peaches, (of which he ate a large quantity every year,) falling,

and resting frequently upon those parts of his body.

"To encourage attempts to cure this disease of the skin in negroes, let us recollect that, by succeeding in them, we shall produce a large portion of happiness in the world. We shall in the first place destroy one of the arguments in favour of enslaving the negroes, for their colour has been supposed by the ignorant to mark them as objects of divine judgment, and by the learned to qualify them for labour in hot, and unwholesome climates. Secondly, We shall add greatly to their happiness, for however well they appear to be satisfied with their colour, there are many proofs of their preferring that of the white people. Thirdly, We shall render the belief of the whole human race being descended from one pair, easy and universal, and thereby not only add weight to the Christian revelation, but remove a material obstacle to the exercise of that universal benevolence which is inculcated by it."

## CHAPTER V.

OF LIFE.

THE knowledge of the mode in which the various functions of the body are exercised constitutes the science of life. The manifestations of life have consequently been considered already. We have seen, that animal and vegetable substances possess the ordinary properties of matter, but that these properties are singularly controlled, so that they are prevented from undergoing those changes, that inevitably occur so soon as they become deprived of vitality. The human body is prone to decomposition. It is formed of substances extremely liable to undergo putrefaction, and is kept at a temperature the most favourable for such change; yet, so long as life exists, the play of the ordinary affinities is prevented. and this constant resistance to the general forces of matter prevails throughout the whole of existence, even to an advanced old age, when it might be supposed the vital forces must be enfeebled almost to annihilation. The case of solution of the stomach after death, described in the first volume of this work, is an additional and forcible evidence of such resistance. So long as life continues in the stomach, the gastric secretions exert no action on the organ, but, when life becomes extinct, these secretions act upon it in the same manner as they do upon other dead animal matter. What, then, is this mysterious power, possessed of such astonishing, such incomprehensible properties? Our knowledge is limited to the fact above stated, that organized matter, in addition to the general physical and chemical forces, possesses one other, - the vital force

606 LIFE.

or principle, vitality or life. This principle exists, not only in the whole, but in every part, of a living body; and its existence is evidenced by the unequivocal signs afforded by the various functions, which we have considered, as well as by others to be presently described. Yet it is not equally evinced in all organs; some appearing to be possessed of more vitality than others,—a result probably produced by diversity of texture, as it would seem irrational for us to admit a different kind of vital principle, where-

ever its manifestations appear to be modified.

Admitting the existence of this controlling principle, what, it may be asked, are the functions through which it immediately acts in keeping up the play of the living machine? It has been elsewhere seen, that, in animals, the reciprocal action of innervation and circulation is indispensable, and that if one of these functions be arrested, the other quickly ceases. This is only applicable, however, to animals; and it has been doubted, whether it applies to all and to every part of them, whilst to the vegetable it is altogether inapplicable, unless we regard it, with some physiologists, to possess a rudimental nervous system. The function of sensibility exhibits to us the mode in which the nervous system acts in connecting man with the objects around him, through the agency of volition; but numerous other acts take place within him, altogether uninfluenced by volition, and yet indispensable for the maintenance of existence. These last acts are equally met with in the animal and the vegetable; and hence a division has been made by Bichat, b into animal life and organic life: - the former evidenced by those functions, that are peculiar to animals - sensibility and voluntary motion - which require the presence of a great nervous centre, that may receive from, and transmit to, the different parts of the body, the nervous irradiations, - the necessary excitant of the different functions: - the latter evidenced by those functions that are common to animals and vegetables, and are inservient to the nutrition of the frame, - as digestion, absorption, respiration, &c., all of which go on without any direct exercise of volition; and occasionally, it has been believed, independently of all nervous influence.

Physiologists may, on this point, be divided into two classes;—they who consider, that the whole of the organic functions are under the government of the nervous influence; and they who think, that the nervous influence does not extend to all the organic functions, but only to the principal of them. The supporters of the first opinion believe, that the agents, or conductors of the nervous influence, are less and less dependent upon the nervous centres, when such exist, the lower the animal is situate in the animal kingdom, and the lower the function; but they consider the nervous influence

b Recherches Physiologiques sur la Vie et la Mort, Paris, 1800.

Fletcher's Rudiments of Physiology, part ii. a, p. 16, Edinb. 1836. See, also, Prichard, on the Vital Principle, Lond. 1829; Dr. W. B. Carpenter, in Edinb. New Philosophical Journal, April, 1838; Principles of General and Comparative Physiology, Lond. 1839; and art. Life, Cyclop. of Anat. and Physiol. Lond. Sept. 1840.

LIFE. 607

to be indispensable to every living being, and to every part of such being. In support of this opinion, they are of course compelled to believe, either that a nervous system exists in the vegetable, or that there is a system, which appears to exert over every part of it an influence necessary for its life, and which is, consequently, analogous to the nervous system of animals. The organ of this influence is, by some botanists, considered to be the medulla or pith; a - whence medullary appendages set out, to be distributed to every part of the vegetable, and which are particularly abundant in such parts as are charged with very active functions, - as the flower. Brachet maintains this idea, and compares the knots of the pith to the ganglions of the nervous system, - destruction of the pith, and especially of these knots occasioning the death of the parts, that receive their filaments from them. Dutrochet, again, considers, that nervous corpuscles exist in the pith of vegetables, which constitute the rudiments of a nervous system; but in the vegetable, this system is diffused, instead of being collected in a mass, as in the animal. The believers in the earlier formation of the nervous system in the fœtus will necessarily be in favour of the first opinion.

The supporters of the second opinion, — that the nervous influence does not extend to all the organic functions, - assert, that it is chiefly exerted on those functions, which are of the highest moment, - the most elevated in animality; that it is less and less in the inferior functions, and ultimately ceases in the lowest acts, - those that immediately accomplish nutrition and reproduction; and the arguments they adduce in favour of their views are, that these lowest acts exist in every living being - vegetable as well as animal; and that in the superior animal, and in man, there are many parts which do not appear to contain nerves. They, moreover, consider the nervous system as one superadded to living beings, not only for life, nutrition and reproduction, but also, where necessary, for sensation, motion, &c., and hence the prolongations or extensions of this system ought to be sent to the organs of the internal or nutritive functions, for the purpose of connecting them with the organs of the external or sensorial functions; and that it is in these connexions only that innervation consists. In this view, consequently, the nervous influence arises only from the necessity of connecting the organs; is but an indirect condition of life; exists in the upper animals only, and can in no way be invoked to account for vegetable life. The last is, in our view, the most accurate opinion. We cannot, in the present state of knowledge, admit the existence of nerves in the vegetable : certainly no such thing as a nervous centre is discoverable, and yet we find the most complicated acts of nutrition and reproduction exercised by it, and the principle of instinct as strikingly evidenced as in many animals. We are, therefore, irresistibly led to the

Sir J. E. Smith's Introduction to Botany, p. 40.
Recherches Expérimentales sur les Fonctions du Système Nerveux Ganglionaire,
&c. Paris, 1830.

608 LIFE.

conclusion, that the manifestations of vitality are but little, if at all, connected with nervous influence, and that the nerves are added, in the upper animals and functions, for other purposes than that of directly communicating vital properties to the part. This deduction will be found confirmed by the facts to be hereafter mentioned, connected with the independence of the vital

property of irritability of the nervous influence.

In the introductory remarks to the first volume of this work, the characters, which distinguish organized from inorganic bodies, were pointed out. All the characters of the former result from the influence of the vital principle, which produces a body of a definite magnitude, shape, structure, composition and duration. There is, moreover, a power, possessed by bodies endowed with the living principle, of being acted upon by certain stimuli, and of being thrown into movement without the participation of the will. This has, indeed, by some physiologists, been considered to be the sole vital property, — with what truth we shall see hereafter. An inquiry into its manifestations will aid us materially in determining whether or not the vital principle be effected directly through the medium of the nerves, and will tend to confirm an opinion, which

we have already expressed on this subject.

Prior to the time of Haller, the nervous system was looked to as the great source of power in the body; and the contractile power of the muscles, - described at length under the head of Muscular Motion, - was considered to be wholly derived from the nerves, which were supposed to transmit the power to the muscular fibre as it was called for, — accurately regulating the quantity supplied. Haller contended for a vis insita, a power of irritability or contractility, essentially residing in the muscles themselves, independently of any condition of the nervous system, and called into action by stimuli, of which, in the case of the voluntary muscles, the nervous influence is one, - contributing, however, like all other stimuli, to exhaust it, instead of furnishing any fresh supply. We have elsewhere shown, that a muscle is capable of being thrown into contraction after a limb has been removed from the body, and for a considerable period after the cessation of respiration, circulation, and consequently of innervation, provided the appropriate stimuli be applied, so as to excite the vis insita, which remains in the muscle for some time after dissolution; and if all the nerves, supplying the limbs of a frog, be divided, and cut out close to the place where they enter the muscles, the muscles will still retain their contractility in as great a degree as when the nerves were entire. They, who believe that the contractility of muscles is wholly derived from the nervous system, maintain, however, that in such case, the stimulus may still act, through the medium of the portions of nerves that must always remain attached to the muscle, however carefully attempts may have been made to remove them; and some have supposed, that these nervous fibres

may even constitute an essential part of the muscular fibre." The most satisfactory reply, that has been made to this argument, is the following experiment of Dr. Wilson Philip. All the nerves, supplying one of the hind legs of a frog were divided, so that it became completely paralytic. The skin was removed from the muscles of the leg, and salt sprinkled upon them, which, being renewed from time to time, excited contractions in them for twelve minutes: at the end of this time, they were found no longer capable of being excited. The corresponding muscles of the other limb, in which the nerves were entire, and of which, consequently, the animal had a perfect command, were then laid bare, and the salt applied to them in the same way. In ten minutes, they ceased to contract, and the animal had lost the command of them. The nerves of this limb were now divided, as those of the other had been, but the excitability of the muscles to which the salt had been applied was gone. Its application excited no contraction in them. After the experiment, the muscles of the thighs in both limbs were found to contract forcibly on the application of salt. It excited equally strong contraction on both sides. In this experiment, the excitability of the muscles, whose nerves were entire, was soonest exhausted; and hence Dr. Philip properly concluded, that the nervous influence, far from bestowing excitability on the muscles, exhausts it like other stimuli; and that the excitability or irritability is a property of the muscle itself.° This is confirmed by the fact, that, when the vital properties of nerves are destroyed by the application of narcotic substances, the irritability of the muscle to which they are distributed may remain for some time longer, so that they must be independent of each other.d

The opinion of J. Müller is, that if muscular irritability be not dependent upon the brain and spinal cord, they supply some influence essential to its exercise, and in confirmation of it he lays much stress on the loss of irritability of muscles within a few weeks after the section of their nerves. This, however, has been shown by Dr. J. Reide to be owing to the alteration in their nutrition consequent on their disuse. He divided the spinal nerves as they lie in the lower part of the spinal canal in four frogs, and both posterior extremities were thus insulated from their nervous connections with the spinal cord. The muscles of the paralyzed limbs were daily exercised by a weak galvanic battery;

See J. W. Earle's New Exposition of the Functions of the Nerves, P. i. Lond.

b An Experimental Inquiry into the Laws of the Vital Functions, &c. p. 100, Lond. 1817.

c See, on the powers of Life, Dr. W. Philip, in Lond. Med. Gazette, for March 18 and 25, 1837; also, a Notice of some Experiments on the Connexion between the Nervous System, and the Irritability of Muscles in Living Animals, by Dr. J. Reid, in the Fourth Report of the British Association, and in Amer. Journ. of Med. Sciences, p. 181, Nov. 1835.

Carpenter, Human Physiology, § 377, Lond. 1842.
 Edinburgh Monthly Journal of Med. Science, May, 1841; and Longet, Comptes Rendus, Juillet, 1841.

610 LIFE.

whilst those of the other limb were permitted to remain quiescent. This was continued for two months; at the end of which time, the muscles of the exercised limb retained their original size and firmness, and contracted vigorously, whilst those of the quiescent limb had shrunk to at least one half of their former bulk, and presented a marked contrast with those of the exercised limb. The muscles of the quiescent limb still retained their contractility, even at the end of two months; but Dr. Reid thought there could be little doubt, that, from their imperfect nutrition, and the progressing changes in their physical structure, this would in no long time have disappeared, had circumstances permitted the prolongation

of the experiment.

It seems that this essential characteristic of living bodies is a distinct vital property, not confined, as Haller supposed, to the muscular structure, but existing over the whole body, In favour of its not being dependent upon the nerves, we have the fact of its presence in the vegetable as well as in the animal. Many plants exhibit this power in a remarkable manner. The barberry bush is one of these. In this flower, the six stamens, spreading moderately, are sheltered under the concave tips of the petals, till some extraneous body, as the feet or trunk of an insect in search of honey, touches the inner part of each filament near the bottom. The irritability of that part is such, that the filament immediately contracts there, and consequently strikes its anther, full of pollen, against the stigma. Any other part of the filament may be touched without this effect, provided no concussion be given to the whole. After a while, the filament retires gradually, and may be again stimulated; and when each petal, with its annexed filament, has fallen to the ground, the latter, on being touched, shows as much irritability as ever. In another plant, — the Cistus helianthemum, dwarf cistus or lesser sunflower, — the filaments, when touched, execute a motion, the reverse of that of the barberry. They retire from the style and lie down, in a spreading form, upon the petals. Owing to the possession of this property, the Apocynum androsæmifolium or dogsbane is extremely destructive to insect life. Attracted by the honey on the nectary of the expanded blossom, the instant the trunk of the fly is protruded to feed on it, the filaments close, and, catching the fly by the extremity of its proboscis, they detain the insect until its struggles end in death, occasioned apparently by exhaustion alone. The filaments then relax, and the body falls to the ground.a

These are only evidences, however, of particular parts possessing an unusual degree of irritability. The property exists in every part of the plant, and, as in the animal, is the essential charac-

teristic of the principle of life.

Irritability or contractility forms a medium of communication between the various parts of the living machine, and is excited to action by extraneous influences. All its movements, however,

<sup>\*</sup> Sir J. E. Smith's Introduction to Botany, p. 211.

appear to be dependent upon the action of appropriate stimuli,

and are, consequently, passively exercised.

1. Instinct. - There is a power, which has been conceived, to be nearly allied to irritability, and is highly characteristic of organized bodies, - vegetable as well as animal, - whose movements or impulsions are active, and most varied. To this power, the term instinct has been appropriated by Virey, Fleming, Good, Good, and others, It is an extension of the ordinary acceptation of the term, but it enables us to understand the phenomena better than where we restrict it to those manifestations of man, or animals, that bear the semblance of reason. It is this power, which, according to those gentlemen, regulates the movements, that are requisite to obtain a supply of food, to remove or counteract opposing obstacles, and to fly from impending danger, or repair injuries. In every organized system," says Dr. Good, "whether animal or vegetable, and in every part of such system, whether solid or fluid, we trace an evident proof of that controlling, and identifying power, which physiologists have denominated, and with much propriety, the principle of life. Of its cause and nature we know no more than we do of the cause and nature of gravitation, or magnetism. It is neither essential mind nor essential matter; it is neither passion nor sensation; but though unquestionably distinct from all these, is capable of combining with any of them; it is possessed of its own book of laws, to which, under the same circumstances, it adheres without the smallest deviation; and its sole and uniform aim, whether acting generally or locally, is that of health, preservation, or reproduction. The agency, by which it operates, is that which we denominate or should denominate instinct, and the actions, by which its sole and uniform aim is accomplished, are what we mean, or should mean, by instinctive actions; or to speak somewhat more precisely, instinct is the operation of the living principle, whenever manifestly directing its operations to the health, preservation or reproduction of a living frame, or any part of such frame. The law of instinct, then, is the law of the living principle; instinctive actions are the actions of the living principle; and either is that power, which characteristically distinguishes organized from unorganized matter, and pervades and regulates the former, uniformly operating by definite means in definite circumstances to the general welfare of the individual system or of its separate organs, advancing them to perfection, preserving them in it, or laying a foundation for their reproduction, as the nature of the case may require. It applies equally to plants and to animals, and to every part of the plant, as well as to every part of the animal, so long as such part continues alive. It is this, which maintains, from age to age, with so much nicety and precision, the dis-

<sup>·</sup> Art. Instinct, in Dict. des Sciences Médicales, xxv. 367.

Philosophy of Zoology, vol. i. 14, Edinb. 1822.

tinctive characters of different kinds and species, which carries off the waste or worn out matter, supplies it with new, and in a thousand instances, suggests the mode of cure, or even effects the cure itself, in cases of injury or disease. It is 'the divinity that stirs within us' of Stahl, the vis medicatrix naturæ of Hoffmann

and Cullen and the physicians of our day, &c., &c."

Of the existence of this instinctive principle we shall adduce a few examples from both the vegetable and the animal kingdom. When the seed of a plant is deposited in the ground, under circumstances favourable for its development, it expands, and the root and stem are evolved. The root descends into the ground, manifestly not from the laws of gravitation, but owing to some inherent force, inasmuch as it penetrates the earth, which is of much greater specific gravity than itself. The stem, too, bursts through the earth, and rises into the air, notwithstanding that the air is of much less specific gravity, until having attained the height to which the action of the vital principle limits it, its upwards development ceases. It rarely happens, however, that the root is capable of procuring nourishment sufficient for its future development in immediate contact with it. It, therefore, sends out numerous filamentous radicles in all directions to search after food, and to convey it to the proper organs. The number and direction of these filaments, and the distance to which they extend, are regulated by the necessities of the plant, and the supply of the soil. A strawberry offset, planted in sand, will send out almost all its runners in the direction in which the proper soil lies nearest; and few, and sometimes none, in the direction in which it lies most remote." When a tree, which requires much moisture, has sprung up, or been planted in a dry soil, in the vicinity of water, it has been observed, that a much larger portion of its roots has been directed towards the water, and that, when a tree of a different species, and which requires a dry soil, has been placed in a similar situation, it has appeared, in the direction given to its roots, to have avoided the water, and moist soil. When a tree, too, happens to grow from seed on a wall, it has been seen, on arriving at a certain size, to stop for a while, and to send down a root to the ground. As soon as this root has been established in the soil, the tree has continued increasing to a large magnitude. The fact has been often noticed with respect to the ash, - a tree, which in consequence of the profusion of its seed, is found more often scattered in wild and singular places, than in any other not propagated by the agency of birds, or conveyed by the winds.b

We find, in all cases, that if the roots of a plant, spreading in search of nourishment, meet with interruption in their course, they do not arrest their progress, but either attempt to penetrate the

Fleming, op. citat. p. 16; and Mayo's Outlines of Human Physiology, p. 14, edit, 4th, 1837.

<sup>&</sup>lt;sup>b</sup> Smith's Introduction to Botany, p. 114; and the Sacred History of the World, by Sharon Turner, F. S. A., Amer. Edit. i. 168, New York, 1832.

opposing body, or to avoid it by altering their direction. Dr. Fleming states, that he has repeatedly seen the creeping root of the Triticum repens or couch grass piercing a potato, which has obstructed its course. It is well known, too, that roots will pass under a stone wall or a ditch, and rise up on the opposite side. A striking case of this nature was communicated to the author, by his venerable friend, the late Ex-President Madison. The wooden pipes, for the conveyance of water to Mr. Madison's establishment, having become obstructed, they were carefully examined, when it was found, that the roots of a honeysuckle, growing immediately above a plug, made of the wood of the Liriodendron tulipifera or American poplar, which is of soft consistence, had penetrated the plug in various places to reach the water, and formed an agglomerated mass in the pipe so as to completely

preclude the passage of the water along it.

The nearest approximation to these manifestations of instinct, in the animal, occurs in the formation of the new being, and in the first actions that take place after birth. From the moment of the admixture of the substances, furnished by the parents at a fecundating copulation, there must be a principle existing in the embryo, which directs the construction and arrangement of its organs after a definite manner, and always according to that peculiar to the species. In the egg, this is seen in the most distinct manner. The germ of the chick is surrounded by the nourishment requisite for its formation. Organ after organ becomes successively evolved, until the full period of incubation is accomplished, when the young animal breaks the shell. At this time, it has within it a portion of nutriment derived from the yolk drawn into the body. This supplies its wants for a short period; but it soon becomes necessary that it should select and collect food for itself, and we observe it throwing its various organs into action for the prehension, mastication, deglutition, &c. of the food, as if it had been long accustomed to the execution of these functions. In the formation of the human fætus in utero the same instinctive action is observable in the successive evolution of organs, and in the limitation of the body to a determinate shape, size, structure, &c.; and when these requisites have been attained, the child bursts the membraneus envelope, and is extruded, to maintain thenceforth an existence independent of the mother. More helpless, however, than the young of the animal kingdom in general, the infant requires the fostering care of the parent for the purpose of supplying it with the necessary nutriment, but as soon as food is conveyed to the lips, the whole of the complicated process of deglutition is effected for the first time, with the same facility as after long practice. As we descend in the animal kingdom, we find these inward actions constituting instinct more and more largely exhibited. In the quadruped, it is not necessary, that the nipple should be applied by the a Op. citat. p. 18.

614 LIFE.

· mother to the mouth of the new-born animal. It is sought for by the latter, invariably discovered, and as invariably seized hold of, by the appropriate organ of prehension, the mouth. The lips are applied; the air is exhausted; and the milk flows according to exact principles of hydrostatics, but without the animal having the least knowledge of the physical process which it accomplishes. Naturalists, indeed, assert, that before the calf has been more than half extruded from the mother, it has been seen to turn around, embrace and suck the maternal teat. As we descend still farther in the scale of creation, we discover the manifestations of instinct yet more signally developed; until, ultimately, in the very lowest classes of animals, the functions are exercised much in the same manner as in the vegetable; and appear to be wholly instinctive, without the slightest evidence of that intelligence, which we observe in the upper classes of the animal kingdom, and pre-eminently in man. This, however, applies only to the very lowest classes; for, a short way higher up the scale, we meet with apparent intelligence, united with instinct, in a manner that is truly surprising

and mysterious.

Again, the similarity of the actions of the instinctive principle, in the animal and vegetable, is exhibited by the reparatory power which both possess when injuries are inflicted upon them. If a branch be forcibly torn from a tree, the bark gradually accumulates around the wound, and cicatrization is at length accomplished. The great utility of many of our garden vegetables, - as spinach, parsley, cress, &c. - depends upon the possession of a power to repair injuries, so that new shoots speedily take the place of the leaves that have been removed: similar to this is the reparatory process, instituted in the lobster that has lost its claw, in the waternewt that has lost an extremity, or an eye; in the serpent deprived of its tail, and in the snail, that has lost its head. These parts are reproduced as the leaves are in the spinach or the parsley. Few animals, however, possess the power of restoring lost parts; whilst all are capable of repairing their own wounds when not excessive, and of exerting a sanative power, when labouring under disease. If a limb be torn from the body, provided the animal should not die from hemorrhage, a reparatory effort is established, and if the severity of the injury does not induce too much irritation in the system, the wound will gradually fill up, and the skin form over it. To a lesser extent we see this power exerted in the healing of ordinary wounds, and in cementing broken bones; and although it may answer the purpose of the surgeon to have it supposed, that he is possessed of healing salves, &c., he is well aware, that the great art, in these cases, is to keep the part entirely at rest, whilst his salves are applied simply for the purpose of keeping the wound moist; the edges in due apposition, where such is necessary, and extraneous bodies from having access to it; - his trust being altogether placed in the sanative influence of the instinctive power situate in the injured part, as in every part of the frame.

It is to this power, that we must ascribe all the properties, assigned to the famous sympathetic powder of Sir Kenelm Digby, which was supposed to have the wonderful property of healing wounds, when merely applied to the bloody clothes of the wounded person, or to the weapon that had inflicted the mischief; - a powder, which, at one time, enjoyed the most astonishing reputation. The wound was, however, always carefully defended from irritation by extraneous substances; and it has been suggested. that the result furnished the first hint, which led surgeons to the improved practice of healing wounds by what is technically called the first intention. It is to this instinctive principle, so clearly evinced in surgical or external affections, but at times, not less actively exerted in cases of internal mischief, that the term vis medicatrix natura has been assigned; and, whatever may be the objections to the views entertained regarding its manifestations in disease, that such a power exists can no more be denied than that organized bodies are possessed of the vital principle. We have too many instances of recovery from injuries, not only without the aid of the practitioner, but even in spite of it, to doubt for a moment, that there is, within every living body, a principle, whose operations are manifestly directed to the health and preservation of the frame, and of every part of such frame.a So far, then, it is manifest, that the instinctive actions of the animal and vegetable are exerted according to the same laws, and probably through similar organs. This, at least, applies to the lowest of all animated beings, where the difference between them and the vegetable is small indeed. It applies equally to the human fætus, which can be considered but to vegetate during the greater part of utero-gestation; and even for some time after birth, its actions are purely instinctive, and differ but little from those of the vegetable, except that, owing to the organization of its nervous system, the acts are of a more complicated character. It is only when the brain has become duly developed, and the external senses fully so, that it exhibits so decidedly the difference between those acts, which it had previously accomplished instinctively, and the elevated phenomena of sensibility, which man enjoys so pre-eminently, but which are likewise possessed, to a greater or less extent, by the whole animal creation.

The cells of the ordinary honeycomb are intended for the larvæ of the different varieties of the occupants of the hive. These cells are usually placed horizontally, with their mouths opening towards the sides of the hive. The bottom of the cells, instead of forming one flat square, is composed of three lozenge-shaped pieces, so united as to make the cell end in a point; consequently, the whole forms an hexagonal tube, terminating in a pyramidal cavity. If the two cells had been a single hexagonal tube, intersected in the middle by a flat, instead of a pyramidal, division, not only would the shape

<sup>\*</sup> See the author's "General Therapeutics," chap. i. Philad. 1837; art. Autocratia, in Most's Encyclopädie, Zweite Auflage, Band i. s. 242, Leipz. 1836; and Dr. Alison, art. Instinct, Cyclopædia of Anat. and Physiol. part xix., July, 1840.

616 LIFE.

not have answered the purpose of the bees, but more wax would have been expended in its construction. Hence, it would seem, that both the body and the base of the tube are adapted for their object; that the greatest strength and the greatest capacity are obtained with the least expenditure of wax in an hexagonal tube with a pyramidal base. Réaumur, when inquiring into the habitudes of these industrious animals, requested König, an able mathematician, to solve the following question: - among all the hexagonal tubes with pyramidal bases, composed of three similar and equal rhombs, to determine that which, having the same capacity, can be constructed with the least possible quantity of matter? König not aware of the precise object of Réaumur's inquiry, solved the problem, and found, - that if three rhombs or lozenges were so inclined to each other that the great angles measured 109° 26', and the little angles 70° 34', the smallest possible quantity of matter would be needed. Maraldi measured the angles actually formed at the bottom of a cell, and found that the great angles gave 109° 28', and the little 70° 32'. All this, however, may be ascribed to blind instinct, proceeding uniformly in the same track. without any evidence of the admixture of reason; but we have innumerable instances, in the same insects, to show, that their operations are varied according to circumstances, and that intelligence is manifestly expended in the adaptation of their means to definite purposes. Of this we shall give but one example. Huber, whose inquiries into this part of entomology have been singularly minute and accurate, having had great ravages committed on his hives by the sphinx atropos or death's head moth, determined to construct a grating, which should admit the bee but not the moth. He did so, and the devastation ceased. He found, however, that in other hives, not protected by his agency, the bees had adopted a similar expedient for their defence; and these defences were variously constructed in different hives. "Here, was a single wall whose opening arcades were disposed at its higher part; there, were several bulwarks behind each other, like the bastions of our citadels: gateways, masked by walls in front, opened on the face of the second rows, while they did not correspond with the apertures of the first. Sometimes, a series of intersecting arcades permitted free egress to the bees, but refused admittance to their enemies. These fortifications were massy, and their substance firm and compact, being composed of propolis and wax." It would be endless, however, and beyond the design of this work, to enumerate the various evidences of intelligence exhibited by the insect tribe, in fulfilling the ends for which they have been destined by the Great Author of nature."

In all our reasonings on the subject of instinct, we must be compelled to admit, in the case of most animals at least, a degree of

<sup>&</sup>lt;sup>2</sup> For numerous examples of the kind, see Alison, art. Instinct. Cyclop. Anat. and Physiol. July, 1840. See, also, Swainson, On the Habits and Instincts of Animals, in Cabinet Cyclopædia, Lond. 1840.

intelligence that strikingly modifies those actions - the impulse to which is doubtless laid in organization. The precise line of demarcation between instinctive acts and reason cannot, however, be established, and this has led some philosophers to call in question the existence of the former. It is owing to this union of intelligence with instinct, that we find animals accommodating themselves to circumstances, so that if prevented from adopting the habits that belong to the species, they have recourse to others as similar as possible. Thus, if a bird be prevented from building its nest in a particular situation, or from obtaining the material, which birds of its own species employ, it has recourse to other materials and to another situation, as like those that are appropriate to it as is practicable. The rook usually and instinctively builds its nest on the summit of the tallest trees: but Dr. Darwin, - who is one of those that call in question the influence of instinct, - asserts, that in Welbourn churchyard, a rookery was formed on the outside of the spire, and on the tops of the loftiest windows. There had formerly been a row or grove of high trees in the neighbourhood, which had been cut down, and, in consequence, the birds exhibited the union of intelligence with instinct, by building on the lofty spire and windows. In like manner, the jackdaws of Selbourne, according to Mr. White, not finding a sufficiency of steeples and lofty houses, on which to hang their nests in that village, accommodated themselves to circumstances, and built them in forsaken rabbit burrows.a

By Stahl, and the animists in general, as well as by more recent philosophers, all the phenomena of instinct have been referred to experience, so obscure as not to be easily traceable, but not the less certainly existent. The insect tribes, however, furnish us with many cases where the young being can never see the parents, and can, of course, derive no benefit from the experience of its progenitors; yet their habits are precisely what they have probably ever been; so uniform, indeed, as to compel us to refer them to some constant impulse connected with their special organization, and, consequently, instinctive. In support of the existence of these natural impulsions, the common occurrence of a brood of young ducks, brought up under a hen, has been adduced.c These little beings, soon after they have broken the shell, and contrary to all the feelings and instincts of the foster-mother, will seek the water, and suddenly plunge into it, whilst the hen herself does not dare to follow them. By what kind of experience or observation, - it has been asked, - by what train of thought or reasoning has the scarcely fledged brood been able to discern that a web-foot adapts them for swimming? Any experience they can have derived must have taught them to shun the water; yet, notwithstanding this,

<sup>&</sup>lt;sup>2</sup> Natural History of Selbourne, with additions, by Sir W. Jardine, Amer. Edit, p. 82, Philad. 1832: Alison, art. Instinct, Cyclop. Anat. and Phys. July, 1840.

b Theoria Vera Medica. Hal. 1737.
CGood's Book of Nature, ii. 118, Lond. 1826.

618 LIFE.

instinct points out to them the habitudes to which they are adapted, and its indications are obeyed in spite of every kind of counter-experience. Attempts have occasionally been made to domesticate the wild turkey of this continent, by bringing the young up under the common turkey, but they have always resumed the way of life to which instinct has directed them, when opportunity offered; in accordance with the Horatian maxim:

"Naturam expellas furcâ, tamen usque recurret."

Mr. Madison reared, with great care, a young hawk, which, for a long time, associated with the young of the poultry, without exhibiting the slightest carnivorous or migratory propensity, until, on one occasion, whilst some of his friends were admiring its state of domestication, it suddenly arose in the air, darted down, and seized a chicken, with which it flew to a neighbouring tree, and, after it had finished its repast, took flight, and was never seen afterwards.

Instinct, then, is possessed by every organized body, animal and vegetable; whilst intelligence is the attribute of those only, that are endowed with a certain nervous development. They are, therefore, manifestly distinct; - the former predominating over the latter in the lower classes of animals; whilst, in the upper classes, intelligence becomes more and more predominant, until, ultimately, in man, it is so ascendant as to appear to be the main regulator of the functions: indeed, some have altogether denied the existence of instinct in him. Instinct is seated in every part of a living body; is totally independent of the nervous system; occurs in the vegetable and the zoophyte unprovided with nerves, or at least in which nerves have never been discovered; whilst intelligence is always accompanied by a nervous system, without which, indeed, its existence is incomprehensible. How can we, consequently, accord with those physiologists who place the seat of instinct in the organic nervous system; and that of intelligence in the brain? Where is the organic nervous system of the zoophyte, and à fortiori of the vegetable? Or how can we admit the seat of the various instincts, with Gall, to be in the brain, seeing that we have them exhibited where there is neither brain nor any thing resembling one. The acephalous fœtus undergoes its full development in other respects in utero, with the same regularity, as to shape and size, as the perfect fœtus, and can we deny it the existence of instinct? Yet, in the upper classes, of animals especially, many of the manifestations of instinct are effected through the nervous system, which, in them, as we have elsewhere seen, seems to hold in control the various functions of the frame, and to be one of the two great requisites for the existence of vitality. The instinctive action in the appropriate organ, which gives rise to the internal sensations of hunger, thirst, &c., is communicated to the great nervous centres by the nerves, and the brain responds to the impression, and excites, through the

medium of the nerves, the various organs into action which are

calculated to accomplish the monitions of the instinct.

What is the nature of this instinctive property? Of this we know no more than we do of the principle of life, of which it is one of the manifestations. It is equally inscrutable with the imponderable agents, light, caloric, electricity, or magnetism, or with the mode of existence of the immaterial principle within us, which gives rise to the mental phenomena: we see it only in its results, which are, in many cases, as unequivocal as those produced by the agents just referred to. All, perhaps, that we are justified in concluding is - with Dr. Good - that instinct is the operation of the principle of organized life, by the exertion of certain natural powers, directed to the present or future good of the individual, whilst reason is the operation of the principle of intellectual life, by the exercise of certain acquired powers directed to the same object; that the former appertains to the whole organized mass as gravitation does to the whole unorganized; actuating alike the smallest and the largest portions; the minutest particles and the bulkiest systems; and every organ, and every part of every organ, whether solid or fluid, so long as it continues alive; that, like gravitation, it exhibits, under particular circumstances, different modifications, different powers, and different effects; but that, like gravitation, too, it is subject to its own division of laws, to which, under definite circumstances, it adheres without the slightest deviation; and that its sole and uniform aim, whether acting generally or locally, is that of perfection, preservation or reproduction.

In this view, reason demands discipline, and attains maturity; instinct, on the contrary, neither requires the one, nor is capable of attaining the other. It is mature from the first, and equally so

in the infant as in the adult.a

2. VITAL PROPERTIES. — The great cause of all those mysterious phenomena, which characterize living bodies, and distinguish them by such broad demarcations from the dead, has been a theme of anxious inquiry in all ages; and has ever ended in the supposition of some special abstract force, to which the epithet vital has been assigned, and which has received various appellations. Hippocrates designated it by the terms outles, and order ; Aristotle styled it the animating or motive and generative principle; Van Helmont, the archwus; Stahl, anima; Barthez and Hunter, vital principle, &c., &c. Yet as Dr. Barclayb has correctly observed, all physiological writers — ancient and modern — seem to be agreed, that the causes of life and organization are utterly

See, also, on this subject, Fletcher's Rudiments of Physiology, part ii. b, p. 16, Edinb. 1836; J. S. Bushnan, The Philosophy of Instinct and Reason, Edinb. 1837; Brit. and For. Med. Rev., Jan. 1841; and Flourens, Eloge de F. Cuvier, Paris, 1840.

An Inquiry into the Opinions, ancient and modern, concerning Life and Organization, Edinb. 1822; and Rullier, art. Force, in Dict. de Méd. ix. 318, Paris, 1824. See, also, Todd and Bowman, Physiological Anatomy and Physiology of Man, Part i. p. 17, Lond. 1842.

LIFE. 620

invisible, whether they pass under the name of animating principles, (Aristotle, Harvey, &c.,) vital principles, (Barthez,) indivisible atoms, spermatic powers, organic particles of organic germs, (Buffon,) formative appetencies or formative propensities, (Darwin,) formative forces, (Needham,) formative nisus or Bildungstrieb, (Blumenbach,) pre-existing monads, (Leibnitz,) semina rerum, (Lucretius,) plastic natures, (Cudworth,) occult qualities, or certain unknown chemical affinities. "All seem agreed, that whatever they be, they have been operating since the world began, and throughout the world operating regularly, without intermission, in various places at the same time. All seem agreed, that their modes of operation are strictly methodical; that they seem to act on definite plans, and actually exhibit specific varieties of chemical combination, and mechanical structure, which human intelligence cannot comprehend, much less explain. From their mutual dependence, and other relations subsisting between them, all seem to speak as if they were subject to one great cause, which regulates and harmonizes the whole.a All seem to speak of this great cause as if it were eternal, omnipotent, omnipresent; whether it be the element of fire, of air, or of water, or whether it be fate,

nature, necessity, or a God."

By virtue of this principle of life, every organized tissue is possessed of certain properties, to which the term vital has been assigned. Regarding the precise number of these properties, physiologists are not agreed. Whilst some have reckoned many; others have admitted but one. All the functions, which we have hitherto considered, are under the influence of life, and are products of the vital properties seated in the tissues; but we do not consider them to be directly caused by these properties. Digestion, for example, is executed by a series of organs, all of which are conducive to a certain result, the aggregate constituting the function of digestion. The result of the action of the salivary gland is very different from that of the liver, yet both operations are vital, but modified by the different organization of the two glands. We do not ascribe the difference to a difference in the vital properties of the glands. These are probably the same in both; and are seated in the primary tissues, of which all the more compound textures and organs are built up. They are primary or fundamental properties of living matter.

Stahl, having observed obscure, oscillatory movements, alternate contraction and expansion in certain parts of the body, either during the exercise of a function, or on the application of some external agent, conceived, that every part of the frame is, at all times, more or less susceptible of similar movements. These movements he called tonic, their effect upon the organs tone, and the property by which they were induced he esteemed peculiar to organization, and termed it tonicity. This vital property, he conceived, influences the progression of the fluids in the vessels; the pheno-

A. F. Hembel, Einleitung in die Physiologie und Pathologie der Menschlichen Organismus, p. 93, Götting. 1828.

mena of exhalation and absorption, and is totally distinct from the

properties possessed by inorganic bodies.

Hallera admitted two vital properties, very different from each other, which seemed to him to be equally elementary. The one of these is that by which a living part exhibits itself to be sensible, or transmits to the sensorium an impression made upon it, either by an extraneous body, or by its own internal and organic action; the other, that by which a part contracts in a manner appreciable to the senses, either by the influence of the will, or of some external or internal stimulus. The first of these he considered to be a special vital property, which he termed sensibility; and the second to be another property, which he called *irritability*. Prior to his time the word irritability had been adopted by Glisson<sup>b</sup> who had noticed the fact, that living matter was acted upon by irritants of various kinds in a mode no wise analogous to physical and chemical motions, and hence he concluded, that every organ of the human frame possesses an inherent and peculiar force, which presides over it movements, and is requisite for the exercise of its functions. This force he called irritability. De Gorter subsequently extended the views of Glisson, and applied them to the vegetable, affirming irritability to be the sole vital property of all organized bodies, vegetable as well as animal. The acceptation, given to the term by Haller, was consequently more limited. He restricted it to those motions of parts which fall under the observation of the senses; such as the contraction of the voluntary muscles, heart, &c. He made numerous experiments on living animals, for the purpose of discovering what parts are possessed or not of the true properties of sensibility or irritability, and he concluded, that the former resides exclusively in the nervous, the latter in the muscular, system. Dr. Marshall Halld still employs the term in the restricted sense of Haller.

This celebrated theory, which formed so large a part of physiological science at one time, and is still an interesting topic to the physiologist, has been referred to in so many parts of this work, as to require but few comments in this place. We have seen that many of the parts, regarded by Haller as insensible, are acutely sensible in disease, and that we cannot pronounce a part to be positively insensible, until we have applied every kind of irritant to it without effect. We have elsewhere defined sensibility to be an exclusive property of the nervous system; and have attempted to show, that irritability is a property of the muscular tissue—a vis insita—totally independent of the nerves, but of which the nervous fluid is an appropriate excitant. As, however, the vital properties of sensibility and irritability were restricted by

b De Ventriculo, in Manget. Bibl. Anatom. i. 80, Genev. 1699.

Medicin. Compendium, Lugd. Bat. 1742.

<sup>\*</sup> Element. Physiol.; and Mémoir. sur la Nature Sensible et Irritable des parties du Corps, Lausan. 1756.

<sup>4</sup> Art. Irritability, Cyclop. of Anat. and Physiol. July, 1840.

Haller to the nervous and muscular systems, they were regarded to be insufficient for the explanation of the various living actions of the frame: the next step was to extend them to every part and to every tissue. It was found, for example, that on investigating the most minute movements of parts, these movements were always preceded by an impression, to which they seemed sensible, and which appeared to excite their actions. This general property, common to every living part, of receiving an impression, was called sensibility; - thus generalizing the property, which Haller had restricted to perceptivity by the mind. Every part was said to be sensible to the blood sent to it for its nutrition. Again, every part was observed to move in consequence of the impression it received, sometimes in an apparent manner, as the heart; at others, too slightly for its movements to be recognised otherwise than by the results, - as in the case of the glandular organs; but always in a manner special to organized matter, and not analogous to any physical or chemical process. This motion was, therefore, referred to another force, called motility, which was nothing more than irritability generalized. These two properties are alone admitted by most modern writers. Every organ is said to feel and to move, after its manner, in the performance of its function; - the stomach in digestion; - the heart in propelling the blood; the muscle in contracting, and the nerve in transmitting sensitive impressions to the brain.

Many modern physiologists, whilst they admit the properties of sensibility and motility, have reckoned a greater number of vital properties: this is owing to their having observed that each part has its own peculiar mode of sensibility and motility, and when these modes have seemed to differ largely from each other, they have elevated them into so many special vital properties. The chief modern theories on the vital properties are those of Barthez, Blumenbach, Chaussier, Dumas, and Bichat. Bartheza admitted five, which we can do no more than enumerate, - sensibility. force of contraction, force of expansion or active dilatation, force of fixed situation, and tonicity. Blumenbach, b also admitted five ; -- sensibility, irritability, contractility, vita propria or proper force of life nisus formativus, force of formation or Bildungstrieb. Dumase referred all the living phenomena to four vital properties; sensibility, motility, force of assimilation, and force of vital resistance. The theory of Bichatd on this subject requires a more detailed notice. He, also, admitted five vital properties; organic sensibility, insensible organic contractility, sensible organic contractility, animal sensibility and animal contractility. First. Organic sensibility is the faculty, possessed by every living fibre of receiving an impression, or of being modi-

<sup>&</sup>lt;sup>a</sup> Nouveaux Elémens de la Science de l'Homme, Paris, 1806.

b Institutiones Physiologicæ, Gotting. 1786; or Elliotson's translation.

<sup>·</sup> Principes de Physiologie, 2de édit. Paris, 1806.

d Anatomie Générale, tom. i.; and Recherches Physiologiques sur la Vie et la Mort, Paris, 1800.

fied by contact, so that the modification is restricted to the part that experiences it, and is not transmitted to the brain. The term sensibility was adopted by Bichat, because already established, and the epithet organic was added, to affirm, that it is the exclusive attribute of organized bodies, and common to all. This property is not only modified in each organ - as the difference in their nutrition and functions demonstrates - but it adapts each organ to its appropriate external stimulant, so that the salivary gland shall be specially influenced by mercury; the upper part of the small intestine by calomel; the lower by aloes, &c., &c. Its exercise is continuous, involuntary, known only by its results, and is more marked as we descend in the scale of animal life; whilst animal sensibility is the contrary. Secondly. Insensible organic contractility is the faculty, possessed by every living part, of moving in an imperceptible manner, in consequence of an impression immediately received, without either the mind having consciousness of the motion, the will participating, or the brain in any manner directing it. We have an example of this in the action of the stomach during digestion; and of every part of the body on the blood sent to it for its nutrition. Bichat applied the term insensible organic contractility to this property, for the following reasons: - contractility, because contraction is the kind of motion which constitutes it; organic, because it is common to all living beings; and insensible, because the brain has no consciousness of it. Like organic sensibility, it is modified in each organ. Its exercise is likewise continuous and involuntary; and it also exhibits itself more intensely as we descend in the scale of beings. It always co-exists with organic sensibility. Thirdly. Sensible organic contractility is the same motive faculty as the last, with this difference, that the movements induced by it fall under the senses and are recognised independently of their results. This property is likewise modified in each organ; its exercise is also involuntary, and it only differs from the last in degree, - the movement that constitutes it being apparent. Thus, the heart contracts independently of the will, but its motions are not imperceptible, as in the cases which belong to the second vital property or insensible organic contractility. Fourthly. Animal sensibility is the property possessed by certain organs of transmitting to the mind, through the medium of the brain, the consciousness of impressions, which they have received. It is sensibility in the restricted acceptation of Haller. The epithet animal was given to it by Bichat to distinguish it from the other variety of sensibility, which belongs to all organized bodies, whilst this is exclusively possessed by animals. The whole of the attributes of this property have been detailed, at much length, in the first volume of this work. Fifthly. Bichat admitted a fifth vital property, under the name animal contractility, which comprised voluntary muscular contraction; treated of else where as one of the functions of the body. It differs from organic contractility, in its exciting causes not being seated

in the organ in which it is developed, — that is, in the muscles, — but in the brain; and, moreover, whilst the other varieties of contractility are irresistibly connected with, and proportioned to, the kind of sensibility correspondent to them, this is not the case with

animal sensibility, and its play is never continuous.

From the distinction we have endeavoured to draw between the fundamental vital properties and the functions, it will be obvious, that the ingenious division of Bichat is susceptible of farther curtailment by analysis. A vital property must be one possessed by all living bodies; it is fundamental in the tissues, and differs according to the precise structure of the tissue. It is found in the vegetable, as well as in the animal. Neither of the two last properties of Bichat, however, corresponds with this definition. They do not exist in the vegetable. They require not only a nervous system, but a brain, that can conceive and will. They are both, indeed, complicated functions, and, as such, have been considered at great length elsewhere. By ultimate analysis, therefore, the five vital properties of Bichat may be reduced to the two we have previously mentioned,—sensibility and motility. Perhaps we ought to rest satisfied with the admission, that every primary tissue is capable of being acted upon by appropriate stimuli, or is sensible; and that it possesses the additional property of moving, in consequence of such impression. Physiologists have, however, attempted to simplify the subject still farther, and fo reduce the vital properties to one only. Such is the view of Broussais, who considers contractility to be the fundamental vital property of all the tissues. Adelon considers, that sensibility is the only living property, that should be admitted, which must carry with it the idea of motion, and is the active, motive faculty of living matter. The term sensibility is, however, unfortunate, in consequence of its conveying the notion of mental perception, and of such acceptation having been received into physiology to designate a function. It has, consequently, been proposed to substitute the term excitability, incitability or irritability, but with the same signification. Rudolphia prefers incitability, (Erregbarkeit,) as not liable to the objection that may be urged against the others, of having been employed in other significations. This incitability differs in the different organs and tissues: in the muscles he terms it irritability (Muskelkraft, Reizbarkeit); in the nerves, sensibility (Nervenkraft, Empfindlichkeit); and by some physiologists, in the membranous parts, it is called contractility (Spannkraft, Zusammenziehungskraft.)b

3. LIFE OF THE BLOOD. — Such are the phenomena which indicate the existence of the vital principle, and such the laws by which

<sup>\*</sup> Grundriss der Physiologie.

b For various views on the subject of irritability, see Alison, art. Contractility, Cyclop. of Anat. and Physiol. i. 717, London, 1836; Adelon, Physiologie de l'Homme, 2de édit. iv. 547, Paris, 1829; and Dr. Hall, art. Irritation, Cyclop. of Anat. and Physiol., July, 1840.

it seems to be governed. By certain physiologists, it is considered to influence solids only; by others, it has been considered to reside in the fluids also, and especially in the blood. The notion of the vitality of this fluid was espoused by the celebrated John Hunter," and to him we are indebted for many of the facts and arguments adduced in its favour, which have impelled the generality of modern physiologists to admit its existence. The analogy of the egg had demonstrated, that life is not restricted to substances, which are solid and visibly organized. The fresh egg, like other living bodies, possesses the ordinary counteracting powers communicated by vitality, and resists those agents, which act upon the dead egg as on other animal substances deprived of the living influence. The fresh egg may be exposed for weeks, with impunity, to a degree of heat, which would inevitably occasion the putrefaction of the dead egg. During the time of incubation, the egg of the hen is kept, for three weeks, at a heat of 105°; yet when the chick is hatched, the remaining yolk is perfectly sweet. The power of resisting cold is equally great. Dr. Hunter performed several experiments, which show the power of the vital principle in resisting cold, and the influence of cold in diminishing the energy of the principle. He exposed an egg to the temperature of 17° and of 15° of Fahrenheit, and found that it took about half an hour to freeze it. When thawed, and again exposed to a temperature of 25°, it was frozen in one-half the time. He then put a fresh egg, and one that had previously been frozen and again thawed, into a cold mixture at 15°; the dead egg was frozen twenty-five minutes sooner than the fresh. These experiments led to the legitimate inference, that the egg possessed the principle of life, and, although fluid, must have enjoyed the properties, which we have described to be characteristic of vitality,—of being acted upon by an appropriate irritant, and of moving responsive to the irritation.

Similar results to those obtained with the egg followed analogous experiments with the blood. On ascertaining the degree of cold, and the length of time necessary to freeze blood taken immediately from the vessel, he found that, as in the egg, a much shorter period, and a much less degree of cold, were requisite to freeze blood that had been previously frozen and thawed, than blood recently taken from the vessel. The inference, deduced from this, was, that the vitality of recent blood being comparatively unimpaired, it was enabled to resist the cold longer than blood whose vital energy had already been partly exhausted by previous exposure. It would appear, however, that the vital principle in fishes can resist the action of frost. Those that were caught by Captain Franklin's party in Winter Lake froze as they were taken out of the nets, and became in a short time a solid

mass of ice; yet they were alive when thawed."

Treatise on the Blood, &c., p. i. ch. i. Philosoph. Transact. 1778, pp. 29, 30.

c See page 194 of this volume; and Dr. W. B. Carpenter, art. Life, Cyclop. of Anat. and Physiol. Sept. 1840.

VOL. II. - 53

The fluidity of the blood, whilst circulating in the vessels, has been regarded as an additional evidence of its vitality. It is obvious, that such fluidity is indispensable, seeing that it has to circulate through the minute vessels of the capillary system, and that the slightest coagulum, forming in them, would lead to morbid derangements. Yet the blood is peculiarly liable to become solid by its constitution, and whenever it is removed from its vessels it coagulates. This is not owing simply to the cessation of its circulation, for if it be kept at the same temperature as in the living body, and be made to circulate with equal rapidity through a dead tube, it equally becomes solid. The cause, consequently, that maintains its fluidity, is the vital agency; or, as J. Müller remarks, the proper combination of its elements is maintained so long only as the blood is under the influence of living surfaces, - that is, of the vessels. The experiments of Shröder van der Kolka show, that coagulation takes place with extraordinary rapidity after the brain and spinal marrow have been broken down: even in a few minutes after the operation, coagula were found in the great vessels. Mayer observed, that after the application of a ligature to the pneumogastric nerve the blood coagulated in the vessels and death was thus produced. Sir Astley Cooper, on repeating the experiment, found that the conversion of venous into arterial blood in the lungs was prevented. Of four experiments, however, which were performed under the direction of J. Müller, b - two on dogs and two on rabbits - although the animals were examined immediately after death, which resulted from the ligature of the pneumogastrics, in two cases only was a small coagulum, of the size of a pea, discovered in the left side of the heart - none in the pulmonary vessels. Another argument in favour of the vitality of the blood is drawn from the facts connected with its coagulation, - facts, which show, that the process is but little influenced by physical agents, and which have induced Magendies to infer, with many other physiologists, who are but little disposed to invoke the vital agency, "that the coagulation of the blood cannot be ascribed to any physical influence, but must be esteemed essentially vital, and as affording a demonstrative proof that the blood is endowed with life." It has, indeed, been attempted to show, that there are certain phenomena, which demonstrate that the vitality of this fluid increases or diminishes with the vitality of other parts of the body. When blood is drawn from a vessel it does not instantly coagulate or die; and, by observing the length of time consumed in the process, it has been thought, that we might, in some measure, be able to estimate the degree of vital energy it possesses. In diseases, where the vital action is exalted, - as in inflammation, - the blood is found to

<sup>\*</sup> Comment. de Sanguin. Coagulat. Groning. 1820; and Diss. sist. Sanguin. Coagulat. Groning. 1820; also, J. Müller's Handbuch, u. s. w., and Baly's translat. p. 97, Lond. 1838. Op. cit. p. 98.

<sup>·</sup> Précis de Physiologie, 2de édit. ii. 234, Paris, 1825.

coagulate much more slowly than in a state of health, and the coagulation itself is more perfect, whilst in diseases, that are dependent upon a diminution of the vital energy, the opposite is the fact; because, in the first case, it is presumed, the blood possesses the vital principle in a higher degree than natural, and consequently resists, for a longer period, the influence of the physical agents to which it is exposed; whilst, in the second case, it possesses the vital principle to a less degree than natural, and therefore yields sooner to the influence of those agents, —the coagula tion, in all instances, being analogous to the rigidity of the muscles, which takes place after dissolution, and indicates the final

cessation of vitality or the last act of life.a

The buffy coat, or inflammatory crust of the blood, called, also, corium phlogisticum, and crusta pleuretica - the nature of which has been investigated before (vol. ii., p. 108,) - is a circumstance connected with the blood's life, which has been invoked by the supporters of this view of the subject. These terms are applied to an appearance of the crassamentum, which is dependent upon its upper portion containing no red particles, but exhibiting a layer of a buff-coloured coriaceous substance, lying at the top, owing to the red particles, during coagulation, sinking to the lower portion of the clot, before coagulation is completed; hence, the colourless state of the upper surface. At the same time, the whole of the coagulated portion is much firmer than usual. The red particles, in such case, have time to subside before the coagulation is complete, which takes place more slowly than in health; and this is conceived to be owing to the blood's possessing a higher degree of vitality, - a view which is confirmed by some experiments of Mr. Thackrah. These consisted in receiving blood, taken from the vessels of a living animal in a full and uninterrupted stream. into different cups, and noting the time at which coagulation commenced in each. Blood, for example, was taken from a horse at four periods, about a minute and a half being allowed to intervene between the filling of each cup. In the first cup, coagulation began in eleven minutes and ten seconds; in the second cup, in ten minutes and four seconds; in the third cup, in nine minutes and thirty-five seconds; and in the fourth cup, in three minutes and twenty seconds. In another experiment, blood was drawn into three separate cups, from the veins of a slaughtered ox. the first of which was filled in the first flow; the second, about three minutes afterwards; and the third, a short time before the death of the animal. Coagulation commenced, in the first cup, in two minutes and thirty seconds; in the second, in one minute and thirty-five seconds; and in the third, in one minute and ten seconds. In a similar experiment, coagulation commenced

See, on the Evidences of the Life of the Blood, from its self-motion, p. 151, of this volume.

b An Inquiry into the Nature and Properties of the Blood, in Health and in Disease, Lond. 1819.

in the first cup, in two minutes and ten seconds; in the second, in one minute and forty-five seconds; and in the third, in thirty-five seconds. Similar phenomena are found to occur in the human subject. Blood, to the amount of about a pint and a half, was taken from the arm of a female labouring under fever. A portion of this, received into a cup on its first effusion, remained fluid seven minutes; a similar quantity, taken immediately before tying up the arm, was coagulated in three minutes and thirty seconds. Of blood, taken as in the last experiment, from the arm of a man, the first portion began to coagulate in seven minutes; the last in four. The vitality of the system, and with it the vitality of the blood, being diminished by each successive abstraction of that fluid, it coagulated or died sooner and sooner in proportion as it was previously more and more enfeebled. It is proper to observe, however, that the blood may remain fluid in the vessels and coagulate when removed from them, long after the death of the body. In a case observed by the author, it flowed freely from the vessels of the brain and coagulated fifteen hours after the total cessation of respiration and circulation; b and many such cases have been observed by others.c They would seem to show, that the phenomenon of coagulation is wholly physical in its nature.d It has been elsewhere shown, that each part of the body as regards the cells that compose it may be considered to have a lifeof its own, - a cellular life; hence, a minute part may die and be reproduced without the general life of the individual suffering; and in certain of the lower animals, so little is the organism in general affected by injuries to a part, that when the animal is cut in pieces, each piece may undergo a distinct development, so as to form so many separate beings. In the higher animals, however, this is not the case. In them, the death and reproduction of every part of the frame is taking place in the function of nutrition; and it is only when organs, that are intimately associated with each other, and whose association is essential to the life of the whole, have their functions interrupted, that the interruption of other functions, and general death follows. The death which takes place in minute parts has been called molecular, that of the whole body somatic.e

But granting that some of the above and other arguments lead to a belief in the vitality of the blood, they are equally favourable,

b Proceedings of the American Philosophical Society, for May, June, and July, 1840,

p. 216; and Amer. Med. Intelligencer, Aug. 1, 1840.

J. Davy, Researches, Physiological and Anatomical, ii. 190, Lond. 1839; or Dunglison's Amer. Med. Lib. Edit., Philad. 1840.

d It was repeatedly noticed by Professor S. Jackson, in those who died in Phila-

delphia of the yellow fever of 1820.

<sup>6</sup> Carpenter, art. Death, in Cyclop. of Anat. and Physiol., vol. i., and in Human Physiology, § 645, Lond. 1842.

a See, on this subject, Fletcher's Rudiments of Physiology, part. ii. a, p. 45, Edinb. 1836; Mandl, Archiv. Génér. de Méd. Nov. 1840; or Encycl. des Sciences Méd., Dec. 1840, p. 248; and Manuel d'Anatomie générale, p. 224, Paris, 1843; and Ancell, Lancet, Oct. 26, p. 145, and Dec. 21, 1839, p. 460.

- many of them at least, - to the life of the chyle, which we have seen, accurately resembles the blood in every property, except in that of coloration; and if we admit the blood to be possessed of life, a question arises, respecting the part at which the nutritive substances, taken into the system, become converted into the nature of the being they are destined to nourish, and receive the principle of life. This must be either through the admixture of the fluids poured out from the supra-diaphragmatic portions of the alimentary canal, with those of the stomach or small intestine, or owing to the mysterious and inappreciable agency of the chyliferous radicles themselves, which separate the same fluid, chyle, from every substance that may be submitted to their action. A reference to what has been said, on these topics, under the heads of Digestion and Absorption, will lead to the opinion, that no vitalizing influence is exerted on the food in the stomach and intestines, and therefore that the infusion of vitality - if the expression may be allowed - must take place in the chyliferous or sanguiferous vessels. As to the mode in which the blood obtains its vitality, great doubt must necessarily exist. The general opinion, perhaps, is, that it is obtained from the organic nerves, distributed to the inner coats of the vessels, and this idea is confirmed by an experiment of the late Mr. Thackrah, which showed, that blood, received into a dead vessel, is always more speedily coagulated than when it is retained by ligature in a living vessel; and thence he inferred, that the vitality of the vessel affects the blood, and retards its coagulation. Mr. Thackrah denies, indeed, the life of the blood, and ascribes all the evidences of it, which it exhibits, to the influence exerted by the living vessels on their contents. These are the only fluids that have been suspected to be endowed. with vitality. None of the others exhibit analogous phenomena, when exposed to similar agencies.

On the whole, we are led to the conclusion, that the vital principle animates both solids and fluids, but all that we seem to know regarding it is — in the language of Dr. Barclay<sup>a</sup>— "that all the organisms of animals and plants are formed out of fluids, and that in a certain species of fluid, secreted from the parent, and afterwards inclosed in a very thin and transparent vesicle, there is a living organizing principle, which also acts upon the fluid in a way which we know not, forming out of it a regularly organized system of solids, and forming not only the rudiments of that system, but causing it afterwards to be nourished, and to grow, through the medium of fluids, which are moved and distributed under the influence of this organizing animating principle." Our knowledge being limited to this category, we are compelled to study life in its results or manifestations. These, as we have seen, constitute the science of Biology or Physiology.

Inquiry into Life and Organization, Edinb. 1822.

## CHAPTER VI.

## OF DEATH.

It has wisely entered into the views of Providence, that the existence of all organized bodies should be temporary. Yet we find considerable difference amongst them in this respect. Whilst some of the lower classes of animals and vegetables are no sooner ushered into being than a process of decay appears to commence; others require the lapse of ages for their various developments and declensions; and, as a general rule, those, in which the attainment of growth has been slow, have the period of decrease proportionably postponed; whilst, where maturity

has been rapidly attained, decay as rapidly supervenes.

The ages of man are numerous and protracted. For a time, the parts of the frame, that are concerned in his development, unceasingly deposit the necessary particles, by a process as beautiful and as systematic as it is mysterious, until ultimately the growth, peculiar to the species and the individual, is attained. At this point, the preponderance, which previously existed in the action of the exhalants over the absorbents, appears to cease. All is equality; but, ere long, the exhalants fall off in their wonted activity; the fluids decrease in quantity; the solids become more rigid; and all those changes supervene, which we have described as characterizing the decline of life, and the approach of the phenomenon, which has now to be considered.

Death is the necessary, total, and permanent cessation of those functions, by which the presence of life is characterized. This cessation may happen at all ages, from accident or disease; a few only ceasing gradually to live by the effects of age alone. Hence, a distinction has been made into that kind of death which is produced by the gradual wear and tear of the organs, and that which cuts off the being prematurely from existence. The former has been termed, by some physiologists, senile or natural, the latter accidental. These differ considerably in their physiology; and

will, therefore, require a distinct consideration.

1. Death from Old Age. — The natural period of life is different in different individuals. It varies according to numerous appreciable and inappreciable circumstances; — the original constitution of the individual; the habits of life; the locality in which he may be situate, &c. Whilst some countries are remarkable for the longevity of their inhabitants, others surprise us by the short period allotted for the natural duration of life. Blumenbach asserts, that by an accurate examination of numerous bills of mortality, he

<sup>&</sup>lt;sup>2</sup> See, on this subject, Bellefroid, Bull. Méd. Belge, Août et Nov. 1839; Lemuel Shattuck, Amer. Journ. Med. Sciences, April, 1841, p. 369; and Traill, Outlines of Med. Jurisprudence, Amer. Edit. with notes by the author of this work, p. 30, Philad. 1841.

has ascertained the fact, that a considerable proportion of Europeans reach their 84th year, but that a few exceed it; whilst, according to Fodéré, a in the insalubrious region of Brenne, in France; nature begins to retrograde at from 20 to 30; and 50 years is the usual term of existence. Hallerb noted one thousand cases of centenarians; sixty-two of from 110 to 120 years; twenty-nine of from 120 to 130; and fifteen who had attained from 130 to 140 years. Beyond this advanced age examples of longevity are much more rare and less sufficiently attested; yet we have some well authenticated cases of the kind. Thomas Parr was born in 1635; married when at the age of 120; retained his vigour till 140; and died at the age of 152, from plethora — it was supposed — induced by change of diet. Harvey dissected him and found no appearance of decay in any organ.c Henry Jenkins, who died in Yorkshire, in 1670, is an authentic instance of the greatest longevity on record. He lived 169 years.d

The following list of instances of very advanced ages has been

given e

					Lived.		Age.
Apollonius of Tyana, -	-	-	-	A.	D. 99	-	130
St. Patrick,	-	-	-	-	491		122
Attila,		1	-	1 -17	500	-	124
Llywarch Hên	-	0.401	-	11/2/	500	100	150
St. Coemgene,	-	1-	-	111-11	618	4	120
Piastus, King of Poland	-	-	-	-	861		120
Thomas Parr,		-	-	-	1635		152
Henry Jenkins,	-	12.3		112	1670	-	169
Countess of Desmond,	-111	-10	121	102	1612	12	145
Thomas Damme, -	2	1300	100	030	1648	111-5	154
Peter Torton,	-	-	-	-	1724		185
Margaret Patten -	-	-	-		1739	-	137
John Rovin and Wife,	-	100	-	-	1741	-	172 and 164
St. Mongah or Kentigen,	-	-	-	-	1781	114	185
					The Court	No. of Contract of	and the second second

It would not seem that the natural period of life has differed much in postdiluvian periods. The Psalmist writes:—

"The days of our years are threescore years and ten; and if by reason of strength they be fourscore years, yet is their strength labour and sorrow, for it is soon cut off, and we fly away."

And when Barzillai excused himself for not visiting the royal palace at Jerusalem, he observed to the king:—

"I am this day fourscore years old, and can I discern between good and evil? can thy servant taste what I eat or what I drink? can I hear any more the voice of singing men or singing women? wherefore then should thy servant be yet a burden unto my lord the king?"

The census of the United States has strikingly exhibited the influ-

- \* Traité de Médecine Légale, et d'Hygiène Publique, tom. v. p. 537, Paris, 1813.
- b Element. Physiol. xxx. 3. c Philos. Transact. iii. 1699. d Art. Lebensdauer, in Pierer's Anat. Physiol. Real Wörterbuch, iv. 697, Leipz. 1821.
- Prichard, Researches into the Physical History of Mankind, 2d edit. i. 421, Lond. 1836.
- r 2 Samuel, xix. 35. See art. Longevity, by the author, in Amer. Quarterly Review, viii. 380, Philad. 1835; and the author's Elements of Hygiène, p. 129, Philad. 1835; Quetelet, Sur l'Homme, &c., Paris, 1835, or English Edit. Edinb. 1842; Haller, Element. Physiol. xxx. 3; and Rullier, art. Longevité, in Dict. de Méd. xiii. 217.

ence of races on longevity in the same country. In 1830, according to Professor Tucker, the proportion of whites over 100 years of age, was 1 in 19,529; of free coloured, 1 in 487; and of slaves, 1 in 1410. The census of 1840 confirms this immense difference,—the whites, over 100, were in proportion of 1 in 17,938; the free coloured of 1 in 597; and the slaves of 1 in 1,866. It is not easy to indicate the character of organization, which is most conducive to longevity and to health. It has been supposed, however, and with some probability, that the state of the nervous system is greatly concerned; for the pathologist looks to this part of the frame as the commencement of most if not all fatal maladies.

Generally, the aged individual sinks silently to death, in the manner described under Decrepture, (p. 529,) totally unconscious of all that surrounds him. At other times, however, he preserves his sensorial powers to the last, and may be capable of locomotion; until, owing to some oppression of one or other of the vital functions during sleep, it becomes the sleep of death, - the elasticity of the organs being insufficient to throw off the oppression and resume their functions. At other times, a slight febrile irritation will be the prelude to dissolution. The great characteristic of this kind of death — as pointed out by Bichat in one of the best of his excellent productions - is, that animal life terminates long before organic life. Death takes place in detail, - the animal functions, which connect the aged with the objects around him being annihilated, long before those that are concerned in his nutrition. Death, in other words, takes place from the circumference towards the centre, whilst in accidental or premature death, the annihilation of the functions begins in the centre and extends to the circumference. As vitality gradually recedes in the aged from the exterior, one of the great centres of vitality - brain, heart or lungs - stops for an instant. The powers are insufficient to restore the action, and total death necessarily ensues.

It has been an interesting inquiry with physiologists to determine the cause of death, thus naturally occurring. Opinions have been various, but such causes as affect the three great vital functions seem to be most entitled to consideration. These have been supposed to be; — First, ossification of the arteries, occasioning an obstacle to the free circulation of blood in the parts; Secondly, ossification of the cartilages of the ribs, and diminution of the capillary system of the lungs, preventing sanguification; and Thirdly, shrivelling and gradual induration of the nervous system, rendering it ultimately unfit for innervation, &c. These are the physical circumstances or changes, which may give occasion to the final cessation of the vital phenomena; but, after all, the difficulty remains,

b Recherches Physiologiques sur la Vie et la Mort, Paris, 1800.

<sup>\*</sup> Progress of the United States in Population and Wealth in fifty years, as exhibited by the Decennial Census, p. 72, New York, 1843.

- and one that is insolvable, - to explain the cause why these changes themselves occur in the organs essential to vitality. We say it is insolvable, for, until we have learned the nature of life, which seems far beyond our comprehension in the present state of our knowledge, it is obviously impracticable to understand the phenomena that arise from its gradual declension and final extinction. This kind of death, produced by the gradual declension of the powers of life, is regarded by Dr. W. Philip, as only the last sleep, characterized by no peculiarity, in which the powers, partly from their own decay, and partly from the lessened sensibility increasing the difficulty of restoring the sensitive system, become incapable of the office, and the individual, therefore, wakes no more. We have before remarked, that there appears to us to be a difference between sleep and death, although they may trend closely on the confines of each other.b It is not common, however, for death to occur in this quiet and gradual manner. Man is liable to numerous diseases, from the earliest to the latest period of existence, many of which are of a fatal character. It was admitted by Sydenham, whose estimate cannot be regarded as more than an approximation, that two-thirds of mankind die of acute diseases; and that of the remaining one-third, twothirds, or two-ninths of the whole die of consumption, leaving, consequently, only one-ninth to perish from other chronic maladies, and from pure old age. How small, then, must be the number of those that expire from decrepitude simply!

2. Accidental Death. — This term has been used, by many physiologists, to include all kinds of death that befall man in the course of his career, and before the natural term; the cause consisting in the supervention of some accidental organic lesion, which arrests the vital movements before they would cease of themselves. This kind of death differs essentially from that we have been considering. The individual is here, perhaps, in the full possession of all his faculties; his organs have been previously, to all appearance, in the most favourable condition for the prolongation of life, and his death, instead of being natural, and unperceived in its approaches by the individual himself, is usually forced

and violent.

Every species of sudden death commences by the interruption of the circulation, the respiration, or the action of the encephalon. One of these three functions first ceases, and the others die in

succession. Each will demand a few remarks.

a. Death beginning in the Heart.—When — owing to fatal syncope, to wounds of the heart or great vessels, or to the rupture of an aneurism—the heart is struck with death, the cessation of the functions is speedy. Sensation and motion are lost; respiration is arrested, and death occurs,—if the cause of the cessation of the heart's action be suddenly and sufficiently applied,—almost

<sup>\*</sup> Philosophical Transactions for 1834; and an Inquiry into the Nature of Sleep and Death, p. 166, Lond. 1834.

instantaneously. The order, in which death takes place in the different organs, is as follows:— The heart failing to propel its blood, the brain no longer receives the necessary impulse for the continuance of its functions; it therefore ceases to act; the consequence of this is the death of all those organs that receive their nervous influx from it; all voluntary motion is annihilated, as well as the action of the respiratory muscles; the mechanical phenomena of respiration are, consequently, arrested; and the air is no longer received into the chest. From this cause, then, the chemical phenomena of respiration would cease, were they not previously rendered unnecessary by the cessation of the heart's action. The phenomena of nutrition, secretion and calorification,— functions of

the capillaries, - yield last.

b. Death beginning in the Brain. — In this case, owing to the loss of innervation, — as in severe injury done to the head, or the worst cases of apoplexy, — the sensorial functions first cease, and the individual lies deprived of all sensation, volition, and mental and moral manifestation. Respiration becomes progressively more irregular and laborious, and ultimately ceases. The order of death is here as follows: — the interruption of the brain's action destroys that of the voluntary and mixed muscles; the mechanical phenomena of respiration therefore cease, and then the chemical. This is followed by cessation of the heart's action, owing to the united loss of nervous influx from the brain, and the want of a due supply of blood. To the cessation of the heart's action succeeds the loss of the general circulation; and lastly, that of the functions of nu-

trition, secretion and calorification.

c. Death beginning in the Lungs. — The action of the lungs may be destroyed in two ways: either the mechanical phenomena of respiration may first cease, as in hanging, strangulation, &c., where the air is prevented from reaching the lungs; or the chemical phenomena may be first arrested, as when air is breathed, which does not contain oxygen, but yet can be respired for a time. In the first case, the order of death is as follows: - the mechanical phenomena cease; to this succeeds cessation of the chemical phenomena, owing to the supply of air being cut off; the blood, not experiencing the necessary conversion in the lungs, soon stagnates in the pulmonary capillaries; for a time, however, the heart continues to beat, owing to the aëration effected by the residuary air in the minute bronchial ramifications; but this soon ceases in consequence of the want of supply of blood; the brain dies, and the other parts in succession.a Where the chemical phenomena first cease, the suspension of the action of the brain follows for the cause already assigned; and the mechanical phenomena of respiration are not arrested, until the nervous influx is cut off by the death of that organ.

The immediate phenomena of death and the order of their suc-

<sup>\*</sup> See the article Asphyxia, by the author, in the American Cyclopedia of Practical Medicine and Surgery, part x., Philad. 1836; and in his Practice of Medicine, 2d edit. vol. i., p. 408, Philad. 1844.

cession are easily undersood, when one of the great centres of vitality is suddenly destroyed, either from accident or disease; but when death does not follow immediately, and time is allowed for a series of morbid phenomena to be established, the problem becomes much more complicated. Some organ or structure is first deranged; and, owing to the intimate connexion, which we have elsewhere seen to exist between the various functions, general derangement or irritation follows, and the individual dies, worn out by such irritation, but without our being exactly able to understand on which of the great centres that dispense vitality the malign influence has been exerted, or whether it may not have affected all equally. In inflammation of the brain, heart, or lungs, we may presume, that the functions of these organs have been respectively annihilated by the diseased action; and that as such functions are essential to the existence of vitality, death may arise in the manner we have already described; but we frequently find the bowels affected with inflammation, or the peritoneum lining the interior of the abdomen; and the case, if neglected, is as surely attended with fatal consequences as the same morbid affection of the organs termed vital; and this in a space of time so short, as not to enable us to understand the nature or the mode of action of the lethiferous agent: but that it must exert its influence on one or more of the great centres of vitality is manifest. In many cases, the heart seems to yield first, not suddenly but gradually; the brain, failing to receive its due impulse, becomes progressively unfit for transmitting the nervous influence to the muscles; insensibility gradually supervenes, until it has attained such an extent, that no nervous influence is sent to the respiratory muscles, when cessation of their action naturally ensues. Of the nature, however, of the morbid condition of the heart, thus induced by disease, we are totally ignorant. It is fashionable to say, that death is produced by irritation, but this is merely concealing our deficiency of knowledge under a term, the explanation of the agency of which comprises the whole difficulty. Adelona thinks, that the brain generally gives way first in these cases; in consequence of which the respiration is disturbed, the lung becomes engorged, the respiration difficult, and death occurs as in a case of gradual asphyxia. There is something extremely obscure in these cases. It often happens, that the intellectual manifestations and the nervous distribution to the muscles of voluntary motion will be executed, even vigorously, until a short time prior to dissolution, whilst the feeble, irregular and intermittent beat of the heart may indicate how greatly its irritability is morbidly implicated.

These remarks are chiefly applicable to death, as it arises from the numerous acute affections, which are so fatal to mankind; but it may occur, also, from those, that persist for a great length of time, and destroy after months or years of morbid irritation, as in cases of calculi of the bladder, engorgements of the viscera, &c. In these cases, likewise, death must ultimately result from the

· Physiologie de l'Homme, 2de édit. iv. 472, Paris, 1829.

destruction of one or other of the vital functions, — respiration, circulation or innervation; but, in a manner so gradual, that it takes place nearly in the same way as in old age; except that, in all cases, it proceeds from the centre to the circumference; the great internal functions first ceasing, and afterwards their dependencies, — a difference, which explains why we are justified in attempting means of resuscitation in sudden death, whilst it would be the height of absurdity to have recourse to them where,

"Like a clock worn out with eating time, The wheels of weary life at last stand still."

The renovation could only be effected by the substitution of new,

for the worn out, machinery.


It has been already shown (vol. i., p. 406) that there are certain causes of death, which affect the two sexes in infancy to a different extent; and the same fact is exhibited when the ratio of deaths of the male and female is taken at all ages. The following table, from the valuable statistical report now annually made by the direction of the British government, shows this in a striking manner.<sup>b</sup>

TO THE STREET,		Number o	f Deaths.		
Causes of Death.	18	38.	1839,		
da grellenthaug de	Males.	Females.	Males.	Females.	
Cancer	620	1828	660	2031	
Hooping-cough -	4036	5071	3683	4482	
Consumption -	27,935	31,090	28,106	31,453	
Child-birth	CONTRACTOR OF STREET	2811	onne de servi	2915	
Violent deaths -	8359	3368	8325	3307	
Hydrocephalus -	4242	3430	4313	3436	
Diabetes	152	55	151	63	
Convulsions -	14,549	11,498	14,245	11,163	
Delirium tremens -	167	15	184	22	
Tetanus	100	29	102	20	
Bronchitis	1193	874	916	747	
Pleurisy	329	253	342	246	
Pneumonia	9887	8112	10,000	8151	
Asthma	3359	2386	3092	2091	
Pericarditis	74	50	83	52	
Aneurism	88	31	- 69	33	
Hernia	318	189	299	175	
Fistula	82	18	81	22	
Stone	282	38	274	25	
Cystitis	103	25	118	20	
Nephritis	113	44	99	82	
Gout	161	46	170	45	
Dropsy	5170	7172	5268	6983	
Intemperance -	125	36	178	40	
Starvation by }_	126	41	85	45	
want, cold, &c. 5	COLUMN TO COLUMN	Trads comats	MON MALT	TODOO WAS	

 See H. Mayo, Outlines of Human Pathology, Dunglison's Amer. Med. Lib. Edit. Introduct. p. 3, Philad. 1839.

b Mr. W. Farr, in Third Annual Report of the Registrar-general of Births, Marriages and Deaths in England, p. 72, Lond. 1841.

The following diagram from M. Quetelet exhibits the relative viability of the two sexes deduced by him from numerous statistical inquiries. The dotted line represents the viability of the



Curve indicating the viability or existibility of male and female at different ages. - Quetelet.

female: the other that of the male. According to this, the maximum of viability is least at the age of 14 in both sexes. After puberty, it diminishes more rapidly in the female than in the male. It is also less during the period of childbearing, from the 27th to the 45th year. The age of shortest viability is immediately after birth; that of longest viability immediately before puberty. The viability of the child after the first month of existence, according to M. Quetelet, is greater than that of the man nearly 100 years old. Towards the 75th year, it is scarcely greater than for the infant about the sixth month after birth.

For some time before dissolution, — both in death from old age and from disease,— the indications of the fatal event become more and more apparent. The speech grows embarrassed; the ideas are incoherent; the hands, if raised by the effort of the will, fall inertly into their former position; the laboured respiration occasions insufficient oxygenation of the blood, and the distress excites an attempt at respiration, which the debility renders nearly

a Op. cit. English edit., p. 32, Edinb. 1843.

ineffectual; distressing yawnings, and gaspings occur to remedy the defective pulmonary action, and the whole respiratory system is in forcible and agitated motion, - the teeth, at times, gnashing, and convulsive contractions occurring at the corners of the mouth. The heart becomes gradually unable to propel the blood with the necessary force into the arteries, so that the fluid ceases to reach the extremities of the body - the hands, feet, nose and ears - which grow cold, and a cold clammy moisture oozes from the vessels. In experiments on animals, the blood is found to be gradually driven no farther than to the feet; then to the groin: afterwards it reaches only to the kidneys, and a kind of reflux occurs through the space along which it had previously been urged forwards. The flux and reflux now reach no farther than the diaphragm, and gradually retreat, until the blood flows back upon the heart itself, which now stops for a time, and then makes an effort to free itself from the contained fluid. The heart's action and respiration are imperfectly performed for a few times at irregular intervals, till at length the contractility of the organ is entirely gone. Respiration ceases by a strong expulsion of air from the chest, - often accompanied with a sigh or a groan, and probably arising, partly from the relaxation of the inspiratory muscles, and still more from the elasticity of the cartilages of the ribs. Hence it is that, in common language, to expire is synonymous with to die. In cases of sudden death, the heart may continue to beat for a time after innervation and respiration have ceased. Under such circumstances, the left ventricle dies first, the obstruction to respiration cutting off its supply of blood.

For some time immediately preceding dissolution, there is usually a peculiar mixed expression of countenance, — a compound of apparent mental and corporeal suffering, - which has given occasion to its being called the agony. It is characterized by facial indications, which were first well described by Hippocrates, and from him called Facies Hippocratica. The nose is pinched, the eyes are sunken, the temples hollow, the ears cold and retracted, the skin of the forehead tense, the lips pendent, relaxed and cold, &c. The eye, during this condition, especially when dissolution approaches, is fixed and slightly elevated, being kept in that position, according to Sir Charles Bell, by the power of the brain over the voluntary muscles of the eve being lost, and the organ being given up to the action of the oblique, which he considers to be involuntary muscles. The word "agony," applied to this condition, in many languages means a violent contest or strife, but its acceptation has been extended so as to embrace what have been termed the "pangs of death" and any violent pain. This agony of death, however, physiologically speaking, instead of being a state of mental and corporeal turmoil and anguish, is one of insensibility. The hurried and laboured breathing, the peculiar sound on inspiration, and the fixed and turned up eyeball, instead of being evidences of suffering, are now admitted to be

signs of the brain having lost all, or almost all, sensibility to impressions. All the indications of mortal strife are such in appearance only: even the convulsive agitations, occasionally perceived, are of the nature of epileptic spasms, which we know to be produced in total insensibility, and to afford no real evidence of cor-

poreal suffering.

Although, from the moment that respiration and circulation permanently cease, the body may be regarded as unquestionably dead, vital properties still remain in some of the organs, the presence of which is an evidence that vitality has previously and recently existed. The functions, which persist after the animal has become dead to surrounding objects, are those that belong to the organic class. Animal heat, for example, may still be elicited for a time, in the internal organs more especially; and it may require several hours, in death caused suddenly or speedily, by accident or disease, before the whole body becomes cold. Absorption is, also, said to have occurred after death, and the beard and hair to have grown; but it is more probable, that, in the last cases, the apparent elongation may have been owing to the shrinking of the integuments. The rectum is very frequently evacuated after dissolution; and cases have occurred where a child has been born by the contraction of the uterus after the death of the mother. The most marked evidence, however, of the continuance of a vital property after dissolution, is in the case of the muscles, which, as we have mentioned in another place, can be made to contract powerfully on the application of an appropriate stimulus, even for an hour or two after death. Nysten, a from his experiments, inferred, that the parts cease to contract in the following order: the left ventricle, the large intestine, the small intestine, the sto mach, the bladder, the right ventricle, the œsophagus, the iris, the different voluntary muscles, and, lastly, the auricles, particularly the right auricle.

The body cools gradually at the surface, and especially towards the extremities, with a rapidity proportionate to the privation of fluids, and the coldness of the atmosphere. Whilst the refrigeration is going on, the blood remains more or less fluid; and owing to the arteries emptying themselves, by virtue of their elasticity, of their contained blood, the fluid generally accumulates in the venæ cavæ, the auricles of the heart, and the vessels of the lungs. By virtue of its gravity, it collects also in the most depending parts, occasioning cadaveric hyperæmiæ, sugillations or livid marks, which might be mistaken for bruises inflicted during life; but may generally be distinguished from them by attention. It will be readily understood, that the situation of the blood in the vessels may differ somewhat according to the vital organ which first ceases its functions. If the action of the right heart stops, the lung may be empty; if the lung or left heart ceases, the lung and the right side of the heart - with the vessels communicating with it - may A Recherches de Physiologie et de Chimie Pathologiques, Paris, 1811.

be surcharged with blood, whilst the organs of the corporeal circulation may be almost empty. During the progress of refrigeration, and especially soon after death, the muscles are soft and relaxed, so that the limbs fall into that position to which the force of gravity would bring them; the eyes are half open; the lips and lower jaw pendent, and the pupil dilated. When the body, however, is cold, the blood is coagulated, and white or yellowish coagula exist, especially in the cavities of the heart, which were at one time supposed to be morbid formations, and termed polypi. They take the shape, more or less, of the cavity in which they are found. Lastly, the muscles become firmly contracted, so that no part can be moved, without the application of considerable force; and, in this state, they continue until the natural progress towards putrefaction again softens their fibres. This has been regarded by physiologists as arising from the last exertion of that residue of vital power, which the body retains after the period of apparent dissolution. With more propriety, perhaps, it may be assigned to physical alterations taking place in the organs, owing to the total loss of those powers, which were previously antagonists to such changes. It has recently been attributed, by M. Brück, a to the coagulation of the liquids in the interior of the tissues. He considers, that the fibrin of the muscle coagulates, when the muscular fibre is deprived of life.

It might seem from the previous enumeration of the signs of death, that no difficulty could possibly arise in discriminating between a living and a dead body. Cases have, however, occurred, where such difficulty has been great and perplexing. Many of the signs may exist, and yet the person be merely in a state of suspended animation; and in certain instances it has even been considered advisable to wait for the manifestations of the putrefactive process, before the body should be consigned to the grave. The following case, given by Dr. Gordon Smith, b strongly exhibits the embarrassment that may occasionally arise. A stout young man had been subject to epilepsy, which became combined with madness. On this account it was necessary to remove him to a private asylum in the neighbourhood of London, where he died suddenly, in a violent epileptic paroxysm. The body was removed to the residence of his friends soon after death, when the necessary preparations for interment were made. On paying attention to the corpse it was found, that the limbs were quite pliable; that the eye was neither collapsed nor glazed; and that the whole features retained their full natural appearance as during life. A surgeon, who, for years, had been in the habit of attending him, was sent for; and although he could find no indications of vitality, he prudently recommended, that the interment should not take place until decomposition had begun to. manifest itself. In the course of two or three days, appearances

<sup>Müller's Archiv. Nov. 1842; cited in Edinb. Med. and Surg. Journal, Oct. 1843,
p. 492.
The Principles of Forensic Medicine, 3d edit., Lond. 1827.</sup> 

still continuing the same, a physician was called in, who concurred in the recommendation that had been already given. Fifteen days from the supposed time of his death had elapsed, when Dr. Smith's informant had an opportunity of inspecting the body. At this time, the countenance retained the appearance described, but the eye seemed beginning to sink, and some degree of lividity had commenced on the surface of the abdomen. The joints were still flexible. At this time, a very eminent professor of anatomy viewed the body, and, considering the hesitation that had prevailed to be altogether groundless, he appointed the following day to examine it internally. The head was accordingly opened, and a considerable extravasation of blood found in the posterior part of the cranium, between the skull and dura mater and between the membranes and substance of the brain. No serum was detected in the ventricles; but the brain itself was remarkably hard. This was sixteen days after death. On the following day, the body was interred. A clamour now arose amongst the neighbours, that he had been prematurely handed over to the anatomist. The body was exhumed; an inquest was held; and the evidence of the medical gentleman demanded. The jury returned a verdict of "apoplexy."

It may hence become a matter of medico-legal inquiry to verify the existence of death, in cases where doubt prevails, owing to the person being in a state of apparent death, — natural or assumed.<sup>a</sup>

Perhaps the most singular case on record of suspension of two of the most important of the vital functions occurred to John Hunter. In the year 1769, being then forty-one years of age, of a sound constitution, and subject to no disease except a casual fit of the gout, he was suddenly attacked with a pain in the stomach, which was speedily succeeded by a total suspension of the action of the heart and of the lungs. By violent exertion of the will he occasionally inflated the lungs, but over the heart he had no control whatever; nor, although he was attended by four of the chief physicians in London from the first, could the action of either be restored by medicine. In about three-quarters of an hour, however, the vital actions began to return of their own accord, and in two hours he was perfectly recovered. "In this attack," says one of his biographers, Sir Everard Home, "there was a suspension of the most material involuntary actions: even involuntary breathing was stopped; while sensation, with its consequences, as thinking and acting, with the will, were perfect, and all the voluntary actions were as strong as ever."b

At one period it was universally credited, that substances could be administered, which might arrest the whole of the vital functions, or cause them to go on so obscurely as to escape detection. This

b See, also, Ottley's Life of John Hunter, Bell's Med. Lib. Edit. p. 38, Philad. 1839; and Hunter, On the Blood, by Palmer, Bell's Edit. p. 189, Philad. 1840.

54\*

<sup>\*</sup> See Dr. J. A. Symonds, in art. Death, Cyclopædia of Anat. and Physiol. i. 791, Lond. 1836.

erroneous popular notion is exhibited in the description of the action of the drug administered by Friar Lawrence to Juliet.a

Death may also be feigned for sinister purposes. The author recollects a body having been brought in a sack to the house of Mr. Brookes, the distinguished anatomist of London, the vitality of which was detected by the warmth of a protruded toe. It was that of a robber, who had chosen this method of obtaining admission within the premises.

The celebrated case of Colonel Townshend exhibits the power occasionally possessed over the vital functions; and Dr. Cleghorn, of Glasgow, knew an individual who could feign death, and had so completely the power of suspending, or at least of diminishing, the action of the heart, that its pulsations were im-

perceptible.b

Lastly, the character of the death, as to violence or gradual extinction, is often exhibited in the physiognomy of the dead. Where it has taken place during a convulsion, or by agents that have forcibly and suddenly arrested respiration or innervation, the countenance may be livid, the jaws clenched, the tongue protruded and caught between the teeth, the eyes forced, as it were, from their sockets; but usually in death from old age, or even from acute and tormenting disease, any distortion or mark of suffering, that may have existed prior to dissolution, subsides after the spirit has passed, and the features exhibit a placidity of expression, singularly contrasting with their previously excited condition. For effect, however, the poet and the painter suit their descriptions of death to the character of the individual whom they are depicting. The tyrant falls convulsed and agonized, whilst the tender and delicate female is described to have progressively withered, till

Without a groan, or sigh, or glance to show
A parting pang, the spirit from her past;
And they who watched her nearest could not know

Romeo and Juliet, iv. 1.

<sup>&</sup>quot;Take thou this phial, being then in bed,
And this distilled liquor drink thou off;
When presently thro' all thy veins shall run
A cold and drowsy humour, which shall seize
Each vital spirit: for no pulse shall keep
His natural progress, but surcease to beat.
No warmth, no breath, shall testify thou liv'st;
The roses in thy lips and cheeks shall fade
To paly ashes; the eyes' windows fall
Like death, when he shuts up the day of life;
Each part, depriv'd of supple government,
Shall stiff, and stark, and cold, appear like death:
And in this borrow'd likeness of shrunk death,
Thou shalt remain full two-and-forty hours,
And then awake as from a pleasant sleep."

b See, also, the case of the Hindoo, mentioned at p. 125 of this volume.

The very instant, till the change that cast
Her sweet face into shadow, dull and slow
Glazed o'er her eyes — the beautiful, the black, —
Oh! to possess such lustre, and then lack."

Byron's Don Juan, canto iv.

643

Warwick's description of the frightful physiognomy of Duke Humphrey, after death from suffocation, exhibits some of this poetical license:—

"But see his face is black and full of blood;
His eyeballs farther out than when he liv'd,
Staring full ghastly like a strangled man:
His hair uprear'd, his nostrils stretch'd with struggling:
His hands abroad display'd, as one that grasp'd
And tugg'd for life, and was by strength subdued.
Look on the sheets; his hair you see is sticking:
His well proportion'd beard made rough and rugged,
Like to the summer's corn by tempest lodged.—
It cannot be but he was murdered here:
The least of all these signs were probable."

King Henry VI., Part ii. Act. 3.

How different is this picture from that of the countenance of the young being, who has gradually sunk to death in the manner above described. The beauty is unextinguished, and the paleness and lividity of death have taken the place of the colours of life; yet the wonted physiognomy may remain:—

> "Hush'd were his Gertrude's lips! but still their bland And beautiful expression seem'd to melt With love that could not die!"

> > Campbell.

Perhaps one of the most beautiful and accurate pictures, drawn by Byron, is his description of the serenity of countenance observable in most fresh corpses; an expression, which, by association is deeply affecting, but not without its consolation to the friends of the departed:—

He who hath bent him o'er the dead Ere the first day of death is fled;

Before decay's effacing fingers Have swept the lines where beauty lingers; And mark'd the mild, angelic air, The rapture of repose that's there; The fix'd yet tender traits, that streak The languor of the placid cheek; And but for that sad, shrouded eye, That fires not, - wins not, - weeps not now; And but for that chill, changeless brow, Where cold obstruction's apathy Appals the gazing mourner's heart, As if to him it could impart The doom he dreads, yet dwells upon: Yes, but for these and these alone, Some moments, ay, one treach'rous hour, He still might doubt the tyrant's power. So fair, so calm, so softly seal'd The first, last look by death reveal'd. Byron's Giaour. 644 DEATH.

An easy death - euthanasia - is what all desire, and fortunately, whatever may have been the previous pangs, the closing scene, in most ailments, is generally of this character. In the beautiful mythology of the ancients, Death was the daughter of Night, and the sister of Sleep. She was the only divinity to whom no sacrifice was made, because it was felt, that no human interference could arrest her arm; yet her approach was contemplated without any physical apprehension. The representation of Death as a skeleton covered merely with skin, on the monument at Cumæ, was not the common allegorical picture of the period. It was generally depicted on tombs as a friendly genius, holding a wreath in his hand, with an inverted torch, - a sleeping child, winged, with an inverted torch resting on his wreath, or as Love, with a melancholy air, his legs crossed, leaning on an inverted torch, the inverted torch being a beautiful emblem of the gradual selfextinguishment of the vital flame.a The disgusting representations of Death from the contents of the charnel-house would not seem to have been common until the austerity of the 14th century, and are beginning to be abandoned. In more recent times, Death has been portrayed as a beautiful youth, and it is under this form that he is represented by Canova, on the monument, erected in St. Peter's at Rome, by George the Fourth of England, in honour of the Stuarts.

Some and the connect that an amount with Leannest will

<sup>&</sup>lt;sup>2</sup> D'Israeli, Curiosities of Literature, 2d Series, Amer. Edit., vol. ii. p. 44, Boston, 1834.

## INDEX.

i. refers to the 1st, and ii. to the 2d volume.

Aberration of refrangibility, i. 179 sphericity, i. 178 Abortion, ii. 428

Absence of mind, ii. 554 Absorption, i. 591

accidental, i. 642 of chyle, i. 591 cutaneous, i. 643 decomposing, ii. 172 digestive, i. 592 of drinks, i. 611

of excrementitial secretions, i. 639

internal, i. 639 interstitial, i. 172 of lymph, i. 619 organic, ii. 172

of recrementitial secretions, i.

of solids, ii. 172 venous, i. 632

Abstinence, deaths from, i. 514 effects of, i. 514

Academia del Cimento, experiments of the, on the gizzards of birds, i. 487

Acid, acetic, where found, i. 33

benzoic, i. 33

where found, i. 33

carbonic, quantity of, consumed in respiration, ii. 46 where met with, i. 33

lactic, where found, i. 33 lithic, where found, i. 30 muriatic, where found, i. 26 oxalic, where met with, i. 33 phosphoric, where met with, i. 25 sulpho-cyanic, where met with, i. 29 uric, where met with, i. 30

xanthic, where met with, i. 31

Adipocire, how formed, i. 335 Adipous exhalation, ii. 235 Admiration, expression of, i. 465

Adolescence, age of, ii, 524

Aëration, ii. 42

Affections, what, i. 266

African race, ii. 593

After-birth, ii. 431

After-pains, ii. 431

Age, critical, ii. 528

Ages, the, ii. 510 Air, atmospheric, properties of, ii. 22

drawn into the veins, i. 618

expulsion of, from the intestines, i.

in the intestines, nature of the, i. 570 in the stomach, nature of the, i. 554

Albino, state of the eyes of the, i. 181

Album græcum, what, i. 573 Albumen, where met with, i. 28

concrete, where met with, i. 28

liquid, where met with, i. 28 Aliments, classification of, i. 507

Allantoid vesicle, ii. 456

Alphabet, how formed, i. 445 Ambiguity, sexual, ii. 341

American race, ii. 594

Amnion, ii. 444

Anastomoses of vessels, effects of, on the

circulation, ii. 155 Anatomie vivante, i. 40

Androgynous being, ii. 351

Angle, facial, i, 296 occipital, 298

Anguish with bodily suffering, expression

of, i. 464

Angus, Mr., trial of, ii. 375

Anhelation, ii. 41

Animalcules, spermatic, ii. 328, 392

Animalculists, ii. 385, 392

Animality, what, i. 19

Animals and vegetables, differences be- Blastema, ii. 181 Blastoderma, ii. 461 tween, i. 19 Blood, ii. 88 cold-blooded, what, ii. 187 aëration of the, ii. 42 warm-blooded, ii. 187 agency of the, in health and dis-Antagonism of nerves, i. 357 ease, ii. 165 Antipathies, ii. 575 analysis of the, ii. 108 Apparatus, what, i. 39 arterialization of the, ii. 42 Appetite, i. 511 atmospherization of the, ii. 42 Arabians, the, on the animal spirits, i. 87 buffy coat of the, 110, 627 Arcus senilis, i. 184 clot of the, ii. 98 Areola, ii. 436 coagulation of the, ii. 103 Arterialization, ii. 42 coloration of the, ii. 57 Arteries, ii. 74 circulation in the, ii. 130 fibre of the, ii. 101 forces that propel the, ii. 144 described, ii. 149 forces that retard the, ii. 153 locomotion of, ii. 154 Articulation, ii. 341 globules of the, ii. 92 halitus of the, ii. 197 Asiatic race, ii. 594 inflammatory crust of the, ii. 110 Association, effects of, ii. 585 infusion of substances into the, ii. Athemprobe, ii. 33 166 Atmospherization, ii. 42 life of the, ii. 624 Atrabiliary capsules, i. 624 quantity of, ii. 89 Attitude, i. 395 red colouring principle of the, ii. 31, erect, i. 395 horizontal, i. 401 specific gravity of the, ii. 91 on one foot, i. 400 transfusion of, ii. 166 on the knees, i. 400 velocity of the, ii. 157 sitting, i. 401 venous, ii. 91 Audition, organs of, i. 138, 152 inspiration of, ii. 145 physiology of, i. 152 weight of, in the body, ii. 90 Aura seminis, what, ii. 362 yellow colouring principle of the, i. insufficient for effecting fecundation, ii. 362, 366 Auscultation, i. 150 corpuscles, red, ii. 92 Australian race, ii. 595 white, ii. 97 Axis, cerebro-spinal, i. 50 Blowing the nose, ii. 37 Body, human, specific gravity of the, i. 408 Azote, sources of, in the food, i. 503 where found, i. 25. See Nitrogen. Bones, i. 36, 337 spongy, use of, in olfaction, i. 132 Borborygmus, what, i. 578 Bosjesman female, generative organs of В. the, ii. 233 nates of the, ii. 240 Ballottement, ii. 422 Beaumont, Dr., case by, i. 558 Brace, Julia, deaf, dumb, and blind, i. 278 Bearing a load, physiology of, i. 412 Brain, i. 51 Beastings, what, ii. 438 analysis of the, i. 75 Belching, i. 582 circulation in the, i. 77 convolutions of the, an index of the Bewegungssinn, i. 372 Bier-right, ii. 567 mind, 300 Biffin, Miss, her case, i. 107 decussation of the, 355 Bildungstrieb, ii. 173, 381 fatty matter of the, i. 32 Bile, ii. 277 insensible, i. 81 colouring principle of the, i. 35 movements of the, i. 79 secretion of the, ii. 265 of the negro, i. 295 use of, in digestion, i. 567 the organ of the mind, i. 268 yellow colouring principle of the, i. 32 a plurality of organs, i. 301 Biliary apparatus, ii. 265 protections of the, i. 51 Bilin, i. 34 ratio of the weight of the, to the Biliverdin, i. 35 other parts, i. 294 Births, legitimate and illegitimate, ii. 404 Breasts, ii. 434 Black man of Gmelin, ii. 593 Bridgman, Laura, case of, i. 280

647 INDEX.

Brown man, ii. 594 Buffy coat of the blood, ii. 109 Burns, Miss, case of, ii. 375 Byron, Admiral, effects of prolonged hunger on, i. 511

Caducity, ii. 529 Calcium, where found, i. 25 Callipædia of C. Quillet, ii. 404 Caloric, laws of, i. 101, ii. 186 Caloricité of Chaussier, ii. 202 Calorification, ii. 187

circumstances influencing ii. 200 seat of, ii. 202 theories of, ii. 202

Capillary circulation, ii. 136 vessels, ii. 77 Carbon, where found, i. 25 Cartilages, i. 36, 342 Casein, where met with, i. 29 Catamenia, ii. 342

Caucasian race, ii. 591 Cell, germinal, i. 42, ii. 180 -life, ii. 183

Cells, epithelial, j. 100 formation of, ii. 180

Cellular membrane, exhalation of, ii. 235 Cerebellum, i. 59

Chabert, M., his resistance to heat, i. 102 Cheselden's case of the boy restored to sight,

Chick in ovo, development of the, ii. 459

Childhood, age of, ii. 520 Chinese race, ii. 594

Chlorine, where found, i. 26

Cholesterin, i. 35 Chondrin, i. 30 Chorion, ii. 444 Chyle, i. 599

where formed, i. 528, 563

Chylification, i. 563

Chyliferous apparatus, i. 592 Chylosis, i. 592

physiology of, i. 602 Chyme, i. 528

Chymification, i. 527 Ciliary motion, i. 472 Circulation, ii. 66, 112

in arteries, ii. 130 in birds, ii. 168 capillary, ii. 136 in fishes, ii. 170 in the fœtus, ii. 479 in the heart, ii. 114 in insects, ii. 170 in mammalia, ii. 168 in reptiles, ii. 169 in veins, ii. 142

Circulation, use of the, ii. 165 Circulatory apparatus, ii. 67 Circumcision in the female, ii. 333 Clairvoyance, ii. 544 Climacteric years, ii. 533 Cœnæsthesis, i. 261 Cold, effects of severe, ii. 191 Colouring matter of organs, exhalation of the, ii. 243 Colostrum, what, ii. 438

Colours, accidental, i. 224 complimentary, i. 224 harmonic, i. 224 insensibility to, i. 245

opposite, i. 224 Combustibility, preternatural, ii. 100 Combustion, spontaneous, ii. 100 Commodus, his feats, i. 344 Composition of man, i. 24. Compounds of organization, i. 27

Conception, ii. 398

at different ages, ii. 400 at different seasons, ii. 400 double, ii. 408 physiology of, 399 signs of, 419

Concord, i. 152 Condiments, i. 509 Consonants, i. 446 Constitution, ii. 574 Contractility, i. 21, ii. 608

Contractilité de tissu, i. 42

par defaut d'extension, ii. 42 Copulation, ii. 313, 356

Cord, jelly of the, ii. 454 umbilical, ii. 453 Corpus luteum, ii. 370 Corium phlogisticum, ii. 109

Corpora Wolffiana, ii. 468 Correlation of functions, ii. 555

Cortical membrane, ii. 447 Cortex ovi, ii. 447.

Coughing, ii. 36 Cowper, Spencer, his case, i. 410 Craniological system of Gall, i. 301

Craniology, i. 307 Cranioscopy, i. 307 Cranology, i. 307

Crassamentum of the blood, ii. 98

Cretinism, ii. 581, 600 Critical age, ii. 528 Cruor of the blood, ii. 98 Crttorin, ii. 101

Crusta pleuretica, ii. 109 Cry, i. 440

Crying, i. 440 expression of, i. 463

of animals, ii. 41 Crypsorchides, ii. 319 Crypts, i. 37, ii. 221 sebaceous, i. 95, ii. 221 Curvatures of vessels, effects of the, on the | Drinks, i. 508 absorption of, i. 511 circulation, ii. 154 Duct, thoracic, i. 598 Cutaneous exhalation, ii. 244 Cutis anserina, i. 468 Cutting, Margaret, her case, i. 438 Cytoblast, ii. 180 E. Cytoblastema, ii. 179 Ear, external physiology of the, i. 153 internal, physiology of the, i. 159 D. middle, physiology of the, i. 154 musical, i. 163 Dazzling, i. 223 trumpet, 150 Deaf-dumb, intelligence of the, i. 277 Echo, i. 149 and blind, ii. 277 Egg, incubation of the, ii. 459 Death, ii. 630 Elasticity of tissue, i. 42 Decapitation, death by, i. 85, 344 Elementary structure of animal substances, Decarbonization, ii. 42 Decidua, ii. 410 Elements, inorganic, i. 15, 25 reflexa, ii. 412 organic, i. 15, 27 Declamation, i. 453 containing azote, i. 27 Decrepitude, ii. 531 not containing azote, Defecation, i. 575 i. 32 Deglutition, i. 523 Emboîtement des germes, ii. 386 of air, i. 527 Embryo, see Fœtus. Dentition, first, ii. 517 Embryology, ii. 443 second, ii. 520 Emotions, i. 266 Depuration, cutaneous, ii. 244 instinctive expressions of the, urinary, ii. 281 i. 469 Derivation, ii. 144 Encasing of germs, ii. 386 Desires, instinctive, i. 263 Encephalon, i. 50 Diastole of the heart, ii. 117 Endosmose, i. 44 Diet, variety of, necessary for man, i. 505 Engastrimism, i. 435 Differences, acquired, amongst mankind, Epigenesis, ii. 381 ii. 581 Epiglottis, use of the, in deglutition, i. 524 individual, ii. 570 Epithelial cells, i. 100 natural, ii. 577 Epithelium ciliated, i. 100 Digestion, i. 474 cylinder, i. 100 buccal, i. 520 pavement, i. 100 in the large intestine, i. 571 tesselated, i. 100 in the small intestine, i. 563 Erectile tissues, ii. 159 oral, i. 520 Erection, ii. 357 of solids, physiology of, i. 510 Eructation, i. 582 of liquids, physiology of, i. 578 Erythroid vesicle, ii. 457 of the stomach after death, i. 548 Ethiopian race, ii. 593 physiology of, i. 510 Evolution, doctrine of, ii. 384 theories of, i. 537 Excitability, ii. 624 theory of, by chemical solution, Excito-motory nerves, i. 71 i. 538 Exhalants, ii. 171 by cection, i. 537 Exhalations, ii. 233 by fermentation, i. 538 adipous, ii. 235 by putrefaction, i. 537 areolar, ii. 244 by trituration, i. 538 colouring, ii. 243 Digestive organs, i. 475 cutaneous, ii. 244 of birds, i. 485 external, ii. 244 of ruminant animals, i. 484 internal, 234 Dislodging a stake, physiology of, i. 413 of the marrow, ii. 241 Diverticula, ii. 158 pulmonary, ii. 254 Docimasia pulmonum, ii. 33 mucous, ii. 258. Dragging a weight, physiology of, i. 413 synovial, ii. 242 Dreams, ii. 539 serous, ii. 234 waking, ii. 547 Exosmose, i. 44

INDEX. 649

Expansibility, vital property of, ii. 139 Expectoration, ii. 38 Expiration, ii. 32 Expression, i. 414 depressing, i. 462 exhilarating, i. 461 Extensibility of tissues, i. 42 Extract of meat, i. 30 Extractive of meat, i. 30 Eye, achromatism of the, i. 202 accessory organs to the, i. 192 accommodation of the, to distances, coats of the, i. 180, 208 diaphanous parts of the, i. 183. dimensions of the, i. 192 insensibility of the, to colours, i. 245 muscles of the, i. 194, 211 phosphorescence of the, i. 223 refracting power of the, i. 198 transparent parts of the, i. 183 Eyes, unequal foci of the, i. 239

#### F.

Face, muscles of the, i. 457 Faculties, affective, i. 266, 290 emotive, i. 266, 290 intellectual, i. 285 intellectual and moral, physiology of the, i. 266 mental, i. 266 moral, i. 286 of the heart, i. 286 Fæces, properties of the, i. 573 Falsetto voice, i. 453 Fat, exhalation of the, ii. 235 Fear, expression of, i. 465 Fecundation, ii. 360 Feeling, common sense of, i. 261 of life, i. 261. Female, characteristics of the, ii. 577 Fibre, what, i. 39 albugineous, i. 39 cellular, i. 38 elementary, i. 38 laminated, i. 38 life, ii. 183 medullary, i. 39. muscular, i. 38, 326 nervous, i. 39 pulpy, i. 39 Fibres, primary, i. 38 Fibrin, where met with, i. 28 Filament, what, i. 39 Flexibility of tissues, i. 42 Flexors, preponderance of the, i. 395 Fluid, nervous, i. 88 Fluids, division of, i. 40 of the human body, 35, 40 Flying, i. 412 VOL. 11. - 55

Fœtal existence, ii. 443 Fœtus, of the, ii. 443, 467 anatomy of the, ii. 443 animal functions of the, ii. 482 calorification of the, ii. 507 circulation of the, ii. 478 dependencies of the, ii. 443 development of the, ii. 458 digestion of the, ii. 470 dimensions of the, it. 471 effect of maternal imagination on the, ii. 504 expression of the, ii . 483 external senses of the, ii. 482 histology of the, ii. 443 increment of the, ii. 467 intellectual and moral faculties of the, ii. 483 internal senses of the, ii. 482 motion of the, ii. 483 nutrition of the, ii. 503 nutritive functions of the, ii. 484 peculiarities of the, ii. 475 physiology of the, ii. 482 position of the, ii. 473 reproductive functions of the, ii. respiration of the, ii. 498 secretions of the, ii. 508 weight of the, ii. 471 Follicle, i. 37, ii. 221 sebaceous, ii. 260 Follicular secretions, ii. 258 Food, of man, i. 499 prehension of, i. 517 Force of formation, ii. 173, 331 nutritive, ii. 173 plastic, ii. 173, 381 of vegetation, ii. 173 vital, ii. 173 Forces, motive, seat of the, i. 347 Foreshortening, i. 248 Free-martin, ii. 354 Friction of the blood retards it, ii. 153 Functions, animal, i. 50 classification of the, i. 48 correlation of, ii. 555 nutritive, ii. 474. of man, i. 47 of relation, i. 50 reproductive, ii. 307 table of the, i. 48

G.

Galactophorous tubes, ii. 435
Gall's craniological system, i. 301
Galvanism, effects of, on the muscles, i. 369
Ganglions, glandiform, i. 37, ii. 300
nervous, i. 37

Hawking, ii. 38 Gaping, ii. 38 Gas animale sanguinis, ii. 97 Gases, permeability of tissues by, i. 45 Gastric juice, i. 541 Gelatin, where met with, i. 28 Gemeingefühl, i. 261 Gemeinsinn, i. 261 Generation, ii. 307 ab animalculo maris, ii. 393 animalcular, theory of, ii. 393 by spontaneous division, ii. equivocal, ii. 307 fissiparous, ii. 312 gemmiparous, ii. 312 marsupial, ii. 314 oviparous, ii. 313 ovo-viviparous, ii. 314 regular, ii. 307 spontaneous, ii. 307 theories of, ii. 380 univocal, ii. 307 viviparous, ii. 314 Generative apparatus, ii. 315 Genital organs of the female, ii. 331 of the male, ii. 315 Germinal cell, i. 180 Germs, dissemination of, ii. 390 encasing of, ii. 386 vital, Darwin's notion of, ii. 386 of animals, ii. 402 Girandelli, Madame, her resistance to heat,

Gestation, ii. 410 Gestures, i. 455 Gland, described, i. 37, ii. 241 Glandiform ganglions, ii. 300 Glandular secretions, ii. 260 Globular elements of tissues, ii. 177 Globulin of the blood, i. 28, ii. 96, 101 Goître, i. 624, ii. 600 Granula seminis, ii. 329 Gras des Cimetières, i. 336 Gravity retards the blood, ii. 184 Greisenbogen, i. 184 Growth of the body, ii. 184 Guillotine, death by the, i. 87, 344 Gustation, i. 111

H.

Habit, ii. 581 Hæmadynamometer, ii. 130 Hæmatosis, ii. 42, 49 Hair, i. 95 Halitus of the blood, ii. 97 Hallucinations, ii. 547 Hand, advantages of the, as an organ of touch, i. 106 Harmony, what, i. 152

Hearing, i. 138 immediate functions of, i. 163 improved by cultivation, i. 169 nerve of, i. 161 organ of, i. 138 Heart, ii. 67 a double organ, ii. 67 beat of the, ii. 121 circulation through the, ii. 114 fœtal, pulsations of the, ii. 423 impulse of the, ii. 121 sounds of the, ii. 117 suction power of the, ii. 144 weight, &c., of the, ii. 71 Heart's action, cause of the, ii. 123 Heat sense of, i. 92, 261 animal, ii. 186 of different animals, ii. 188 Hemachroin, ii. 101 Hematin, ii. 96, 101 Hematosin, ii. 96, 101 Hepar sanguinis, ii. 98 Hermaphrodism, ii. 351 Hermaphrodite, ii. 351 Honeywell, Miss, her case described, i. 108 Humorists, i. 36 Hunger, i. 510 Hunter, Mr., case of, ii. 641 Hybrids, doctrine of, ii. 388 Hydrogen-where found, i. 26. Hygrometric property, i. 46 Hymen, use of, ii. 335

I.

Idiosyncrasy, ii. 575 Illusions, mental, ii. 547 optical, ii. 258 spectral, ii. 547 Imagination, maternal, influence of the, on the fœtus, ii. 397, 504 Imbibition, i. 43, 616 Imitation, effects of, ii. 586 Impulses, cerebral, of motion, i. 353 Incitability, ii. 624 Incubation of the egg, ii. 459 Individualitätssinn, i. 261 Infancy, ii. 510 first period of, ii. 510 second period of, ii. 515 third period of, ii. 520 Inflammatory crust, ii. 627 Infusion of medicines into the blood, ii. 166 Inorganic bodies, i. 14 Insalivation, i. 521 Inspiration, ii 27 of venous blood, ii. 145 first, ii. 510 Instinct, ii. 611

INDEX. 651

Instincts, i. 284
Instinctive signs, i. 470
Insula sanguinis, ii. 98
Intellect, i. 284
Intercellular substance, ii. 179
Intercostal nerve, great, i. 66
Intermediate system of vessels, ii. 79
Iron, where found, i. 26
Irritability, i. 17, 362, ii. 624
Irritation, constitutional, ii. 557
Itching, i. 262

J.

Jelly of the cord, ii. 454 Joints, i. 341 Joy, expression of, i. 465

K.

Kalmuck race, ii. 594
Kiestein, ii. 421
Kidney, royal road to the, ii. 297
Kissing, i. 461
Korpergef ühl, i. 261
Kyestein, ii. 421

L.

Labour, ii. 428 premature, ii. 428 Lactiferous tubes, ii. 435 Lachrymal apparatus, i. 195 Lactation, ii. 434 Lacteals, ii. 592 Language, i. 414 artificial, i. 441 natural, i. 440 origin of, i. 443 Larynx, i. 414 Laughing, i. 440, ii. 40 Laughter, ii. 40 broad, i. 461 of animals, ii, 40 Leaping, i. 406 Lebensgefühl, i. 261 Lebenssinn, i. 261 Lebensturgor, ii. 139 Lenses, various, i. 176 Letters, how divided, i. 445 Life, ii. 605 animal, ii. 606 cell, ii. 183 fibre, ii. 183 of the blood, ii. 629 organic, ii. 606 Ligaments, i. 33, 342 Light, i. 170

Light, colour and decomposition of, i. 177 diffraction of, i. 241 duration of the impression of, on the retina, i. 259 intensity of, i. 172 reflection of, i. 172 refraction of, i. 172 velocity of, i. 171 Likeness of child to parent, remarks on the, ii. 397 Line, facial, i. 296 occipital, i. 298 Liquid, prehension of, i. 580 Liquor amnii, ii. 444 false, ii. 444 sanguinis, ii. 109 seminis, ii. 329 Listening, i. 169 Liver, histology of the, ii. 268 Lochia, ii. 431 Locomotion, nervous system of, i. 348 Locomotility, i. 325 Locomotive influx, i. 347 Longsightedness, i. 234 Lung-proof of infanticide, ii. 34, 477 Lungenprobe, ii. 34, 477 Lymph, i. 624 coagulable, ii. 101 corpuscles, ii. 625 Lymphatic apparatus, ii. 619

M.

Lymphosis, ii. 619, 626

Magnetism, animal, ii. 543 Magnesium, where found, i. 27 Malay race, ii. 595 Mammæ, ii. 434 Manganese, where found, i. 26 Manhood, age of, ii. 527 Mankind, varieties of, ii. 587 Mantle, i. 95 Mariotte, experiment of, i. 218 Marks, mother's, ii. 504 Marrow, exhalation of the, ii. 241 Mastication, i. 476 Matière extractive du bouillon, i, 28 Mechanical principles, i. 379 Meconium, ii. 509 Medulla oblongata, i. 59 spinalis, i. 61 Megalanthropogenesis, ii. 404 Melody, i. 152 Membrana granulosa, ii. 461 . Membrane, i. 37 compound, i. 37 fibrous, i. 37 germinal, ii. 460 mucous, i. 37, 99 nictitating, i. 193

032	
Membrane, serous, i. 37	Natural state of man, i. 502
Menses, ii. 342	Nausea, i. 583
Menstruation, ii. 342	Navel-string, ii. 453
vicarious, ii. 345	Negro race, ii. 593
Mesenteric glands, i. 597	Nerves, i. 36, 61
Milk, ii. 436	composition of the, i. 75
Mind, not proportionate to the state of the	fatty matter of, i. 32
senses, i. 276	Fletcher's division, ii. 71
seat of the, i. 323	Lepelletier's division, ii. 70
Miscarriage, ii. 428	M. Hall's division, ii. 71
Mitchell, the boy, case of, i. 277	pneumogastric, effects of the sec-
Molecules, organic, of Buffon, ii. 383	tion of the, on digestion, i. 549
Mongolian race, ii. 594	on respiration, ii. 61
Monorchides of the Cape of Good Hope,	sensible and insensible, ii. 81
ii. 319	Sir C. Bell's division of the, ii. 66
Monstrosities, ii. 503	Nervi-motion, Dutrochet's views of, i. 43,
Moral acts, i. 290	ii. 607
Motility, ii. 662	Nervous system, i. 50
Motion, ciliary, i. 472	Neurine, i. 75
encephalic nerves of, i. 347	New Zealander, head of, ii. 185
involuntary, i. 346	Nightmare, ii. 545
muscular, i. 325	Nipples, ii. 435
physiology of, i. 343	Nisus formativus, ii. 381
vibratory, i. 472	Nitrogen, quantity of, consumed in respi-
voluntary, i. 325	ration, ii. 44. (See Azote.)
Motive apparatus, i. 325	Norma verticalis of Blumenbach, i. 298
forces, seat of the, i. 347	Nose, blowing the, ii. 37
Mouvement de ballottement, ii. 422	use of the, in smell, i. 131
Movements, locomotive, i. 402, 404	Nuclei, ii. 180
partial, i. 402	Nucleoli, ii, 180
Mucous follicular secretion, ii. 238	Nutrition, ii. 171
exhalation of, ii. 258	force of, ii. 173
membranes, i, 99	Nutritive principle, peculiar, does not ex-
Mucus, where met with, i. 30	ist, i. 502
Muscles, i. 36, 325	Nyctalopes, i. 203
analysis of, i. 335	
colour of, i. 335	0.
contraction of, i. 360	o.
relaxation of, i. 371	Odours, i. 126
simple and compound, i. 334	classification of, i. 129
Muscular contraction, duration of, i. 375	disengagement of, i. 126
extent of, i. 372	divisibility of, i. 128
force of, i. 372	medicinal properties of, i. 130
motion, i. 325, 343	nutritive properties of, i. 130
nervous centre of,	vehicles of, i. 128
i. 347 theories of, i. 360	Oken's bodies, ii. 468
	Old age, ii. 529
velocity of, i. 376 web, i. 95	Olein, where met with, i. 32
Musical tone, i. 150	Olfaction, i. 122
Muskelsinn, i. 372	physiology of, i. 131
Muteosis, i. 455	Onomatopæia, i. 443
Myopy, i. 234	Optic nerves, decussation of the, i. 189
	Organic nerve, i. 66
· 100 months was of comment	Organ, i. 39
N.	Organization, i. 16
The state of the s	compounds of, i. 27
Navi matarni ii 504	Organized bodies, characters of, i. 16
Nævi materni, ii. 504	Organology, i. 307
Nails, i. 99	Oscitation, ii. 38
Natural bodies, i. 13	Oscultation, i. 461
signs, i. 470	Osmazome, where found, i. 30

NDEX. 653

Ovarists, doctrine of the, ii. 384 Ovum, development of the, ii. 458 Oxygen, where found, i. 25 quantity of, consumed in respiration, ii. 44

P.

Pain, i. 265
bodily, expression of, i. 464
Pains, after, ii. 431
Painting, a variety of expression, i. 472
Palpation, i. 92
Palsy, theory of, i. 355
Pancreatic juice, secretion of the, ii. 263
use of, in digestion, i. 567
Pandiculation, ii. 38
Panniculus carnosus, i. 95
Panspermia, ii. 390
Panting, ii. 41
Parturition, ii. 428
Passions, i. 290

expression of the, i. 461 seat of the, i. 274 Pectoriloguy, i. 434

Pepsin, i. 28, 544
Peripheral system of vessels, ii. 77
Peristaltic action, i. 532
Peristole, i. 532
Perceptivity of plants, i. 21
Perspective, i. 249
aërial, i. 257

Perspiration, ii. 244
Phonation, i. 414
Phosphorus, where found, i. 25
Phrenologist, cerebral organs of the, i. 307
Phrenology, i. 307
Physical properties of the tissues, i. 42

Physical properties of the tissues, i. 42 Physiology, general, of man, i. 23 Picromel, where found, i. 34

Placenta, ii. 447

of the blood, ii. 98

Placental souffle, ii. 422 Plasma, ii. 109

Pneumogastric nerves, effect of the section of the, on digestion, i. 552

on respiration, ii. 61
Poetry, a variety of expression, i. 472
Point, visual, i. 225

Point, visual, i. 225
Portal system, ii. 87
Potassium, where found, i. 27
Power, sensorial, i. 324
Pregnancy, ii. 410

duration of, ii. 425 protracted, ii. 427 signs of, ii. 429 tubal, ii. 364

Prehension, organs of, i. 413 of food, i. 517 of liquids, i. 580 Presbyopy, i. 234
Presentations, various, ii. 433
Principle, nutritive peculiar, does not exist,
i. 503
vital, ii. 606
Principles, mechanical, i. 379

proximate of animals, i. 27

Proligerous disc, ii. 341 layer, ii. 341

Propelling a body, how effected, i. 412 Property, hygrometric, of tissues, i. 43 physical, of tissues, i. 42

vital, ii. 620
Prosopose, i. 461
Protein, i. 27
Protogala, ii. 438
Puberty, ii. 525
Pulse, doctrine of the, ii. 160

venous, ii. 115
Purgations, ii. 345

Pylorus, use of the, i. 531

Q.

Quickening, ii. 421

R.

Races of man, ii. 590
origin of the, ii. 597
Racornissement, i. 43
Rage, expression of, i. 465
Red man, ii. 594
Regurgitation, i. 582
Reinigung, ii. 345
Rennet, i. 545.
Reproduction, desire of, ii. 355
functions of, ii. 307
instinct of, ii. 355

Respiration, ii. 13, 26
chemical phenomena of, ii. 42
cutaneous, ii. 60
effects of, on the circulation,
ii. 164
of animals, ii. 64
mechanical phenomena of, ii.
27

Respirations, number of, ii. 34 Respiratory organs, ii. 13 Revery, ii. 554 Rubrin, ii. 101 Rumination, i. 583 Running, i. 408

S.

Saliva, ii. 261 Sapidity, cause of, i. 113 Sanguification, ii. 42

Savours, i. 113	Skin, follicular secretion of the, ii. 259
classification of, i. 114	goose, ii. 468
Schurze of the Bosjesman female, ii. 333	Skull, i. 53
Sea-sickness, i. 584	Sleep, ii. 534
Sebaceous follicles, i. 195	complete, ii. 539
Secretion, ii. 221, 224	incomplete, ii. 539
follicular, ii. 258	walking, ii, 542
glandular, ii. 260	Slumber, ii. 536
Secretory apparatus, ii. 221	Smell, i. 122
Secundines, ii. 431	acuteness of, in animals, i. 137
Selbstgefühl, i. 261	acuteness of, in the blind, i. 138
Self-feeling, i. 261	immediate function of the, i. 134
Semen, secretion of, ii. 325	improved by education, i. 137
properties of, ii. 326	mediate functions of, i. 135
Seminal animalcules, ii. 328	nerves of, i. 124
granules, ii. 329	organs of, i. 123
Seminists, ii. 385	Sneezing, ii. 38
Sensations, i. 50	Sobbing, ii. 41
external, i. 90	Sodium, where found, i. 26
înternal, i. 263	Solander, Dr., effects of severe cold on, ii.
morbid, i. 265	192
organic, i. 263	Solidists, i. 36
Sense of individuality, i. 261	Solids, i. 35
locality, i. 261	compound, i. 39
cold, i. 261	Somnambulism, ii. 542
heat, i. 261	Soul, seat of the, i. 324
hunger, i. 261	Sound, i. 147
thirst, i. 261	acute, malappreciation of, i. 166
life, i. 264	intensity of, i. 151
motion, i. 261,	reflexion of, i. 149
pneumatic, i. 261	sympathetic, i. 148 timbre of, i. 152
sixth, of Buffon, i. 261	tone of, i. 152
geometrical, i. 108 regulating, j. 108	vehicle of, i. 148
Senses, additional, i. 260	velocity of, i. 149
Sensibility, i. 50, 82	Spaying, method of effecting, ii. 364
less in the lower animals, i.	Spectra, ocular, i. 224
271	Speech, i. 442
vital property of, ii. 621	Sperm, ii. 325
Sensorial power, i. 324	Spermatic animalcule, ii. 329, 392
Serosity of the blood, ii. 98	Spermatists, ii. 384, 392
Serous exhalation, ii. 234	Spermatozoa, ii. 329
of the cellular mem-	Spinal marrow, protection of the, i. 80
brane, ii. 235	structure of the, i. 61
Serum of the blood, ii. 98	Spirits, animal, i. 87
Sex, doubtful, ii. 351	Spitting, ii. 37
of the fœtus, ii. 480	Splanchnic nerve, i. 66
Sexes, differences between the, ii. 577	Spleen, ii. 300
proportion of the, born, ii. 408	Spontaneity of plants, i. 21
art of producing the, ii. 403	Spurzheim's craniological system, i. 310
Sexual ambiguity, ii. 351	Squeezing, i. 413
Sheep, fat-buttocked, ii. 240	Squinting, i. 240
Short-sightedness, i. 234	Standing, i. 395
Sighing, ii. 38	Stearin, where met with, i. 32
sound of, i. 440	Stethoscope, i. 150
Sight, sense of, i. 169	Stillborn, ii. 407
Silicium, where found, i. 26	Stomach, digestion of the, after death, i.
Singing voice, i. 453.	548
Sinuses, nasal, use of, in smell, i. 131	and kidney, connexion between
Skeleton, living, exhibited, i. 40	the, ii. 297
Skin, i. 92	Stout, Mrs., her case, i. 410

	Strabismus, i. 240	Temperament, bilious, ii. 572
	Straining, ii. 35	choleric, ii. 572
	Stratum proligerum, ii. 341	influence of, on the mind,
	Stretching, ii. 38	i, 273
	Structure, elementary, of animal substances,	lymphatic, i. 573
	ii. 176	melancholic, i. 573
	Study, brown, ii. 554	muscular, i. 572
	Succus intestinalis, i. 566	nervous, i. 573
	Sucking, i. 581	phlegmatic, i. 573
	Suction power of the heart, ii. 144	sanguine, i. 571
		Temperature, animal, ii. 186
	Sugar of diabetes, i. 34	depressed, effects of, ii. 192
	milk, i. 33	elevated, effects of, ii. 195
	Sulphur, where found, i. 25	of animals, table of, ii. 188
	Superfecundation, ii. 408	of bodies, ii. 186
	Superfectation, ii. 408	Terror, expression of, i. 465
		Testes, descent of the, ii. 481
	Supra-renal capsules, i. 624	Testicondi, ii. 319
	Suspicion, expression of, i. 467	Thaumatrope, of Paris, i. 259
	Sutures, i. 52	Thigh-bone, neck of, advantage of the, i.
	Sweat, what, ii. 252	398
	Swimming, i. 408	
	Sympathetic, great, i. 66	Thirst, i. 578
	Sympathy, ii. 561	sense of, i. 261
	agents of, ii. 568	Thoracic duct, i. 598
	cerebral, ii. 568	Thymus gland, i. 623, ii. 476
	direct, ii. 568	Thyroid gland, i. 623, ii. 417
	morbid, ii. 562	Tickling, i. 262
	of contiguity, ii. 562	Tingling, i. 262
	of continuity, ii. 562	Tissue, i. 39
	remote, ii. 564	albugineous, i. 39
•	superstitions connected with, ii.	cellular, i. 38
	566	compound, i. 39
	Synergies, ii. 556	laminated, i. 38
	Synovia, ii. 242	medullary, i. 39
	System, i. 40	muscular, i. 38
	nervous, i. 50	nervous, i. 39
	nervous, of locomotion, i. 347	primary, i. 38
	Systole of the heart, ii. 117	pulpy, i. 39
		Tissues, permeability of, to gases, i. 46
		physical properties of the, i. 42
	T.	Titillation, i. 262
		Toltecan race, ii. 595
	Tablier of the Bosjesman female, ii. 333	Tone, i. 42
	Tact, i. 92	Tongue, i. 112
	Tapetum, i. 203	Tonicity, i. 42, ii. 620
	Taste, i. 111	Touch, i. 100
	diversity of, in animals, i. 120	immediate functions of, i. 107
	immediate functions of, i. 119	mediate functions of, i. 107
	improvement of, by education, i. 120	regarded the first of the senses, i.
	mediate functions of, i. 119	105
	organs of, i. 111	Townshend, Col., his case, ii. 124
	physiology, i. 116	Transfusion, ii. 166
	Tattooing, ii. 185	Transpiration, cutaneous, ii. 244
	Tawny man, ii. 595	pulmonary, ii. 259
	Tears, i. 214	Transudation, i. 43
	secretion of the, ii. 260	Travail, ii. 429
	use of the, i. 214	Trisplanchnic nerve, i. 66
	Teeth, i. 475	Tunica granulosa, ii. 341
	shedding of the, ii. 520	Turgor vitalis, ii. 139
	Temperament, i. 273, ii. 570	Tutamina cerebri, i. 51
	athletic, ii. 572	oculi, 1, 193
	atrabilious, ii. 573	Twins, proportion of cases of, ii. 401

Umbilical cord, ii. 453 Umbilical vesicle, ii. 454 Understanding, i. 285 Urea, where met with, i. 30 Urinary organs, ii. 281 Urine, ii. 294 secretion of, 381 Utero-gestation, ii. 410 Uvula, use of the, i. 523 V. Varieties of mankind, ii. 587 Vasa vasorum, ii. 87 Vegetables and animals, differences between, i. 19 V egetative nerve, i. 66 Veins, ii. 83 circulation in the, ii. 142 Vena porta, ii. 88 Venous system, ii. 83 blood, ii. 88 Ventrale cutaneum of the Bosjesman female, ii. 333 Ventriloquism, i. 435 Venus, Hottentot, ii. 333 Vesicle, allantoid, ii. 456 erythroid, ii. 457 intestinal, ii. 650 germinal, ii. 341, 460 of Purkinje, ii. 460 umbilical, ii. 454 Vessels, i. 341 Vestiges of the French, what, i. 140 Viability of child, ii. 428 Vibrations, sympathetic, i. 150 Virility, age of, ii. 527 Vis insita, of Haller, ii. 126, 621 mortua, i. 47 medicatrix naturæ, ii. 612 Viscus, i. 37 Visible direction, centre of, i. 220 Vision, i. 169 advantages of, to the mind, i. 646 direct, i. 226 direction of bodies appreciated by, distance appreciated by, i. 247 distinct, point of, i. 225 requisites for, i. 226 double, i. 235 erect, i. 220 immediate function of, i. 244 improved by education, i. 260 indirect, i. 226 magnitude, appreciated by, i. 247 mediate functions of, i. 246

motion appreciated by, i. 253

Vision, multiple, with one eye, i. 242 nerves of, i. 217 nocturnal, i. 223 oblique, i. 225 organs of, i. 180 phenomena of, i. 216 physiology of, i. 196 position, appreciated by, i. 245 seat of, i. 217 single, i. 237 surface of bodies appreciated by, i. 247 Visual angle, i. 247 Vital force, ii. 606 Vital principle, ii. 606 properties, ii. 619 Vitality, ii. 606 of the blood, ii. 456 Vitelline disc, ii. 341 pedicle, pouch, ii. 456 Vitelline fluid, ii. 456 Vitellus, ii. 456 Vocal apparatus, i. 414 Voice, i. 414 intensity of, i. 424 native, i. 440 nerves of, i. 418 physiology of the, i. 420 quality of the, i. 432 timbre of the, i. 432 tone of the, i. 424 Volition, seat of, i. 345 Vomiting, i. 583 at pleasure, i. 582 Vowels, i. 438 W.

Walking, i. 404 Wants, i. 263 Weeping, ii. 40 expression of, i. 463 of animals, ii. 40 Whispering, i. 440 Whistelo's case, ii. 396 Whistling, i. 440 Wolff's bodies, ii. 468 Writing, art of, i. 437

Y.

Yawning, i. 440, ii. 38

Z.

Zoohematin, ii. 101 Zoospermes, ii. 328, 395

# MEDICAL LEXICON, BROUGHT UP TO 1842.

A NEW DICTIONARY OF MEDICAL SCIENCE; Containing a concise account of the various Subjects and Terms, with the French and other Synonymes, and Formulæ for various Officinal and Empirical Preparations, &c. Third Edition, brought up to 1842. BY ROBLEY DUNGLISON, M.D., Professor in the Jefferson Medical College, &c. In One Volume, royal 8vo.

"The present undertaking was suggested by the frequent complaints, made by the author's pupils, that they were unable to meet with information on numerous topics of Professional Inquiry, -especially of

recent introduction,-in the medical dictionaries accessible to them.

"It may, indeed, be correctly affirmed, that we have no dictionary of medical subjects and terms which can be looked upon as adapted to the state of the science. In proof of this the author need but to remark, that he has found occasion to add several thousand Medical Terms, which are not to be met with in the only medical lexicon at this time in circulation in this country.

"The present edition will be found to contain many hundred Terms more than the first, and to have experienced numerous Additions and Medifications."

experienced numerous Additions and Modifications.

"The author's object has not been to make the work a mere lexicon or dictionary of terms, but to afford, under each, a condensed view of its various medical relations, and thus to render the work an epitome of the existing condition of Medical Science."

This New Edition includes, in the body of the work, the Index or Vocabulary of Synonymes that was in the former Editions printed at the end of the Volume, and embraces many Corrections, with the addition of many New Words.

### PEREIRA'S MATERIA MEDICA.

EDITED BY DR. CARSON, WITH NEAR THREE HUNDRED ENGRAVINGS ON WOOD.

ELEMENTS OF MATERIA MEDICA AND THERAPEUTICS; COMPRE-HENDING THE NATURAL HISTORY, PREPARATION, PROPERTIES, COMPOSITION, EFFECTS, AND USES OF MEDICINES, BY JONATHAN PEREIRA, M.D., F.R.S., Assistant Physician to the London Hospital, &c.

Part I, contains the General Action and Classification of Medicines, and the Mineral Materia Medica. Part II, the Vegetable and Animal Kingdoms, and including diagrams explanatory of the Processes of the Pharmacopeias, a Tabular view of the History of the Materia Medica, from the earliest times to the present day, and a very copious index. From the Second London Edition, which has been thoroughly revised, with the Introduction of the Processes of the New Edinburgh Pharmacopeia, and containing additional articles on Mental Remedics, Light, Heat, Cold, Electricity, Magnetism, Exercise, Dietetics, and Climate, and many additional Wood Cuts, illustrative of Pharmaceutical Operations, Crystallography, Shape and Organization of the Feculas of Commerce, and the Natural History of the Materia Medica. The object of the author has been to supply the Medical Student with a Class Book on Materia Medica, containing a faithful outline of this Department of Medicine, which should embrace a concise account of the most important modern discoveries in Natural History, Chemistry, Physiology, and Therapeutics in so far as they pertain to Pharmacology, and treat the subjects in the order of their natural historical relations. Part I, contains the General Action and Classification of Medicines, and the Mineral Materia Medica.

This great Library or Cyclopedia of Materia Medica has been fully revised, the errors corrected, and numerous additions made, by DR. JOSEPH CARSON, Professor of Materia Medica and Pharmacy in the "College of Pharmacy," and forms Two Volumes, octavo, of near 1600 large and closely-printed pages; and it may be fully relied upon as a permanent and standard work for the country,—embodying, as it does, full references to the U.S. Pharmacopea and an account of the Medicinal Plants indigence to the United States. nous to the United States.

# PRINCIPLES AND PRACTICE OF SURGERY, WITH CUTS.

THE PRINCIPLES AND PRACTICE OF MODERN SURGERY, BY RO-BERT DRUITT. From the Second London Edition, illustrated with fifty wood engravings, with notes and comments by Joshua B. Flint, M.D., in one volume 8vo., at a low price.

EXTRACT FROM THE AUTHOR'S PREFACE.

"The arrangement of a work of this kind ought not, as I conceive, to be regarded as a matter of mere indifference, or at most of convenience, but it ought to embody in it something of a principle; and I believe that the arrangement of this work may be useful to the student, by showing him in what order he may best prosecute his researches into the principles of his profession.

"Of the five parts into which it is divided, the first two are more especially devoted to the principles, "Of the five parts into which it is divided, the first two are more especially devoted to the principles, and the three others to the practice of surgery. The first part treats of the disturbances of the constitution at large, that may be produced by injury or disease of a part; beginning with the simple faintness or collapse that follows a blow, and proceeding to consider the varieties of fever and tetanus.

"The second part describes what may be called the elements of local disease; that is to say, those morbid changes of structure or function, which are produced either immediately by external causes, or morbid changes of structure or function, which are produced either immediately by external causes, or morbid changes of the various kinds of injuries, beginning with the simplest mechanical inju"The third part treats of the various kinds of injuries, beginning with the simplest mechanical inju"The third part treats of the various kinds of injuries, beginning with the simplest mechanical inju"The third part treats of the various kinds of injuries, beginning with the simplest mechanical inju-

poisons, &c.

"The fourth part considers the various tissues, organs, and regions of the body in order, and describes the various accidents they are liable to, &c.

"The fifth part describes such of the operations as were not included in the former parts, &c.

"To the whole is appended a collection of formulæ, the number of which is very much increased in this afficien."

this edition."

#### FEVERS OF THE UNITED STATES.

THE HISTORY, DIAGNOSIS AND TREATMENT OF TYPHOID AND TYPHUS FEVER, WITH AN ESSAY ON THE DIAGNOSIS OF BILIOUS REMITTENT AND OF YELLOW FEVER, BY ELISHA BARTLETT, M.D., Professor of the Theory and Practice of Medicine in the Transylvania University. In one volume 8vo; a new work.

Notice has already been given of the appearance of this work: we have become satisfied of its sterling value, and, therefore, without hesitation, feel justified in again recommending it to the immediate notice of practitioners.—Boston Medical and Surgical Journal.

## MÜLLER'S PHYSIOLOGY.

ELEMENTS OF PHYSIOLOGY; BY J. MÜLLER, M.D., Professor of Anatomy and Physiology in the University of Berlin, &c. Translated from the German by WILLIAM BALY, M.D., Graduate in Medicine of the University of Berlin. Arranged from the Second London Edition by John Bell, M.D., Lecturer on Materia Medica and Therapeutics, &c., &c. In One Volume, 8vo. - Just ready.

In arranging the Volume now offered to American readers, from the materials furnished in Muller's Elements of Physiology, the Editor has endeavoured to procure reduction in size, of this latter, without any abstraction of its vitality and mind. With this view he has omitted, for the most part, mere disquisitions, many details of experiments, matters of physics and natural philosophy, including mechanics under the head of locomotion, acoustics and the theories of music under voice and hearing, and of optics under vision,—much of the minutiæ of comparative physiology, and metaphysics or metaphysicophysiology. But, while excluding details on collateral topics, the Editor has been particularly careful to preserve Physiology Proper, which, resting on the basis of Histogeny and General Anatomy, derives important aid from Organic Chemistry and Microscopical Observations, and in its turn serves to illustrate Hygiene, Pathology and Therapeutics: Thus aided and thus applied, in the manner exhibited by Muller himself, Physiology will invite the attention of the Student in these pages.

It will soon be discovered that, although this volume is an abridgement of the large work of Müller, it may rightfully claim to be considered a complete system of Physiology, exceeding in copiousness and comprehensive details, any other work on the same subject which has yet emanated from the London

comprehensive details, any other work on the same subject which has yet emanated from the London press.

# ELEMENTS OF PHYSICS—WITH WOOD-CUTS.

ELEMENTS OF PHYSICS, OR NATURAL PHILOSOPHY, GENERAL AND MEDICAL. A New Edition, complete in One Volume, written for universal use, in plain and non-technical language, and containing New Disquisitions and Practical Suggestions; comprised in Five Parts: 1. Somatology, Statics and Dynamics. 2. Mechanics. 3. Pneumatics, Hydraulics and Acoustics. 4. Heat and Light. 5. Animal and Medical Physics. By NEIL ARNOTT, M.D., of the Royal College of Physicians. A New Edition, revised and corrected from the last English Edition; with additions by Isaac Hays, M.D., and numerous Wood-cuts.

## PRACTICAL MINERALOGY AND GEOLOGY—WITH CUTS.

A TEXT-BOOK OF GEOLOGY AND MINERALOGY, WITH INSTRUCTIONS FOR THE QUALITATIVE ANALYSIS OF MINERALS. BY JOSHUA TRIMMER, F.G.S., with Two Hundred and Twelve Wood-cuts. A handsome Octavo Volume, bound in embossed cloth.

This is a Systematic Introduction to Mineralogy and Geology, admirably calculated to instruct the Student in those sciences. The Organic Remains of the various Formations are well illustrated by numerous Figures, which are drawn with great accuracy.

## ELLIS'S MEDICAL FORMULARY IMPROVED.

THE MEDICAL FORMULARY OF DR. ELLIS; being a COLLECTION OF PRESCRIPTIONS, derived from the Writings and Practice of many of the most eminent Physicians in America and Europe. To which is added an Appendix, containing the usual Dietetic Preparations and Antidotes for Poisons; the whole accompanied with a few brief Pharmaceutic and Medical Observations. By BENJAMIN ELLIS, M.D. The Seventh Edition, completely revised, with many Additions and Modifications, and brought up to the present improved state of the Science; by SAMUEL George Morton, M.D., Professor in the Pennsylvania College of Medicine, &c., &c. In One Octavo Volume.

## PRINCIPLES OF MEDICINE.

THE FIRST PRINCIPLES OF MEDICINE, BY ARCHIBALD BILLING, M.D., A.M., Member of the Senate of the University of London, Fellow of the Royal College of Physicians, &c., &c. In One Volume, 8vo. First American from the Fourth London Edition.

"We know of no book which contains within the same space so much valuable information, the result not of fanciful theory, nor of idle hypothesis, but of close, persevering Clinical Observation, accompanied with much soundness of judgment, and extraordinary clinical tact."—Medico-Chirurgical Review.

A TREATISE ON FEVER. By Southwood Smith, M.D., Physician to the London Fever Hospital, fourth American edition. In one volume octavo.

COATES'S POPULAR MEDICINE, OR FAMILY ADVISER, consisting of Outlines of Anatomy, Physiology, and Hygiene, with such Hints on the Practice of Physic, Surgery, and the Diseases of Women and Children, as may prove useful in families when regular Physicians cannot be procured: being a Companion and Guide for intelligent Principals of Manufactories, Plantations, and Boarding Schools; Heads of Families, Masters of Vessels, Missionaries, or Travellers; and a useful Sketch for Young men about commencing the Study of Medicine. By Reynell Coates, M.D.

Young men about commencing the Study of Medicine. By Reynell Coates, M.D. This work is designed to supply the place of Ewells' Medical Companion, which is

now entirely out of print.

OUTLINES OF A COURSE OF LECTURES ON MEDICAL JURISPRU-DENCE. By Thomas Stewart Traill, M.D., with notes and additions. A small volume.

A PRACTICAL TREATISE ON MEDICAL JURISPRUDENCE, with so much of Anatomy, Physiology, Pathology, and the Practice of Medicine and Surgery, as are Essential to be known by Members of the Bar and Private Gentlemen; and all the Laws relating to Medical Practitioners; with Explanatory Plates. By J. Chitty, Esq., second American edition, with notes and additions adapted to American Works and Judicial Decisions. In One Volume Octavo.

ABERCROMBIE ON THE BRAIN. Pathological and practical Researches on Diseases of the Brain and Spinal Cord. New edition, 1 vol. 8vo.

A PRACTICAL TREATISE ON THE HUMAN TEETH, showing the causes of their destruction and the means of their preservation, by William Robertson. With plates. First American from the second London Edition. In one volume octavo.

ANATOMY, PHYSIOLOGY, AND DISEASES OF THE TEETH. By Thomas Bell, F.R.S., F.L.S., &c. Third American edition. In one volume octavo, with numerous plates.

DISSERTATIONS ON NERVOUS DISEASES. By Drs. James Hope, J. C. Prichard, John Hughes Bennett, Robert H. Taylor and Theophilus Thomson. In one

DISSERTATIONS ON DISEASES OF THE ORGANS OF RESPIRATION. By Drs. Williams, Theophilus Thomson, W. B. Carpenter, and W. Bruce Joy. In one volume octavo.

DISSERTATIONS ON FEVERS, GENERAL PATHOLOGY, INFLAMMA-TION, AND DISEASES OF THE SKIN. By Drs. Symonds, Allison, Christison, &c. &c. In one volume octavo.

DISSERTATIONS ON DISEASES OF THE DIGESTIVE, URINARY AND UTERINE ORGANS. By Drs. Joy, Symonds, Thomson, Ferguson, &c. &c. In one volume octavo.

DISSERTATIONS ON HÆMORRHAGES, DROPSY, RHEUMATISM, GOUT, SCROFULA, &c. &c. By Drs. Burrows, Watson, Shapter, Joy, &c. &c.

In one volume octavo.

The above five volumes are from the Library of Practical Medicine, edited by Dr.

Tweedie, with notes by Dr. Gerhard. Each volume is complete within itself, and is
for sale separately.

THE MEDICAL STUDENT; OR AIDS TO THE STUDY OF MEDICINE. Including a Glossary of the Terms of the Science, and of the Mode of Prescribing; Including a Glossary of Medical Works; the Regulations of the different Medical Bibliographical Notices of Medical Works; the Regulations of the different Medical Bibliographical Notices of Medical Works; the Regulations of the different Medical Colleges of the Union, &c. By Robley Dunglison, M.D., &c. &c. In one vol. Svo.

ESSAYS ON ASTHMA, APHTHÆ, ASPHYXIA, APOPLEXY, ARSENIC, ATROPA, AIR, ABORTION, ANGINA PECTORIS, and other subjects, embraced in the Articles from A to Azote, prepared for the Cyclopædia of Practical Medicine by Dr. Chapman and others. Each article is complete within itself, and embraces the practical experience of its author, and as they are only to be had in this collection, will be found of great value to the profession. The two volumes are now offered at a price so low, as to place them within the reach of every practitioner and student.

OUTLINES OF PHYSIOLOGY; with an Appendix on Phrenology. By P. M. Roget, M.D., Professor of Physiology in the Royal Institute of Great Britain, &c. First American edition revised, with numerous Notes. In one volume octavo.

GEOLOGY AND MINERALOGY, considered with reference to Natural Theology. By the Rev. William Buckland, D.D., Canon of Christ Church, and Reader in Geology and Mineralogy in the University of Oxford. With nearly one hundred copperplates and large coloured maps. A new edition from the late London edition, with supplementary notes and additional plates.

THE BRIDGEWATER TREATISES, complete in seven volumes octavo, em-

I. The Adaptation of External Nature to the Moral and Intellectual Constitution of Man. By the Rev. Thomas Chalmers.

II. The Adaptation of External Nature to the Physical Condition of Man. By John Kidd, M.D.F.R.S.

III. Astronomy and General Physics, considered with reference to Natural Theology. By the Rev. William Whewell.

William Whewell.

IV. The Hand; its Mechanism and vital Endowments as evincing Design. By Sir Charles Bell, K.H., F.R.S. With numerous wood-cuts.

V. Chemistry, Meteorology, and the Function of Digestion. By William Prout, M.D.F.R.S.

VI. The History, Habits, and Instincts of Animals. By the Rev. William Kirby, M.A.F.R.S. Blustrated by numerous engravings on copper.

VII. Animal and Vegetable Physiology, considered with reference to Natural Theology. By Peter Mark Roget, M.D. Blustrated with nearly five hundred wood-cuts.

VIII. Geology and Mineralogy, considered with reference to Natural Theology. By the Rev. William Buckland, D.D. With numerous engravings on copper, and a large coloured map.

The works of Buckland, Kirby and Roget, may be had separate.

A POPULAR TREATISE ON VEGETABLE PHYSIOLOGY, by W. P. Carpenter, Author of Principles of Human Physiology, &c., published under the auspices of the Society for the Promotion of Popular Instruction. With numerous wood-cuts, in one volume 12mo.

A POPULAR TREATISE ON AGRICULTURAL CHEMISTRY; intended for the use of the practical farmer, by Charles Squarry, Chemist. In one vol. 12mo.

ROGET'S ANIMAL AND VEGETABLE PHYSIOLOGY, with nearly five hundred wood-cuts, in two volumes, second American edition.

THE HISTORY, HABITS, AND INSTINCTS OF ANIMALS, by the Rev. William Kirby, M.A.F.R.S. Illustrated by numerous copperplate engravings.

The Ninth Bridgewater Treatise. A FRAGMENT, by Charles Babbage, Esq.

From the second London edition. In one volume octavo.

A PRACTICE OF PHYSIC. Comprising most of the diseases not treated of in Diseases of Females and Diseases of Children, second edition. By W. P. Dewees, M.D., formerly Adjunct Professor in the University of Pennsylvania. In one vol. 8vo.

ELEMENTS OF HYGIENE; on the Influence of Atmosphere and Locality; Change of Air and Climate, Seasons, Food, Clothing, Bathing, Sleep, Corporeal and Intellectual Pursuits, &c., on Human Health, constituting Elements of Hygiene. By Robley Dunglison, M.D. In one volume 8vo.

ABERCROMBIE ON THE STOMACH. Pathological and Practical Researches on Diseases of the Stomach, the Intestinal Canal, the Liver, and other Viscera of the Abdomen. By John Abercrombie, M.D. Third American, from the second London edition, enlarged. In one volume 8vo.

## DISEASES OF THE SKIN.

A NEW WORK.

A PRACTICAL AND THEORETICAL TREATISE ON THE DIAGNOSIS, PATHOLOGY. AND TREATMENT OF DISEASES OF THE SKIN, arranged according to a Natural System of Classification, and preceded by an Outline of the Anatomy and Physiology of the Skin. By Erasmus Wilson, M. D., author of a System of Human Anatomy, &c., in 1 vol. 8vo.

# MEDICAL NEWS

# LIBRARY.

#### SUPPLEMENT TO NO. 17 FOR MAY 1844.

Postage on Medical Journals .- It is of the utmost importance to the profession, and indeed to the whole community, that every facility should be afforded for the diffusion of Medical, Agricultural, and Scientific intelligence. To accomplish this, medical men, agriculturists and others should unite to induce Congress to reduce the postage on such periodicals to the same rate as that on Newspapers.

Duration of the life of Medical Men .- M. Chadwick in his " Sanatory Report" states that in the medical profession examples are not rare of the attainment of extreme old age; yet as a class they bear the visible marks of health below the average. The mortuary registration for the year 1839, gives the following as the average age of death of persons in the three professions asserts that in some cases where the hair falls in England:—Clergymen, 59; Lawyers 50, Meaway from the eyebrows, whiskers, &c. the loss dical men, 45. Only one medical student was included in the registration; had the deaths of those who died in their noviciate been included, the average age of death would have been much lower. This corresponds with the results of the researches of Dr. Caspar of Berlin. Yet the medical profession is notoriously the worst paid and the worst treated of all the three professions -the public for whose good this awful loss of life is sustained, is utterly regardless of the sac-

Prize Essay .- The Academy of Medicine offered a prize of 1500 francs for the best essay on ædema of the glottis, in which should be considered its causes, progress, symptoms, diagnosis and treatment with the advantages and disadvantages of tracheotomy. This prize was awarded to M. Valleix on the 12th December

New Journal,-A new Journal (Journal fuer Kinderkrankheiten) devoted to the diseases of children, edited by Drs. Behrend and Hilderbrand, has recently been commenced at Berlin. The first No. contains memoirs on the intermittent fever of children, on the desquamation of the epithelium of the mucous membranes in acute exanthemata, on the semeiology of new born children, and on pleurisy in infants.

German Medical Diplomas.—Mr. Wait of England says that he has been informed by Profestor Naegele that about 200 forged diplomas purporting to come from the University of Hei-delberg, are supposed to be current in the Bri-tish dominions. How many are there in this

Medical Graduates, 1844. — University of Pennsylvania, 155; Jefferson Medical College, 114; Medical Department of Pennsylvania Col-

lege 7; College of Physicians and Surgeons, N. Y., 32; Harvard University, Boston, 17; University of Maryland, 38; Ohio Medical College 36; Transylvania University, Lexington, 59; Medical College of the State of South Carolina, 82; Louisville Medical Institute, 47; University of the City of New York, 93; Medical College Richmond Va. 24; Jackson Kemper College St. Louis 97; Medical Department of St. lege St. Louis, 27; Medical Department of St. Louis University, 11; Geneva Medical Institution, 44; making a total of 784. Several Institutions are not embraced in the above. The number of young men who have been authorized to practice medicine during the present year must be nearly one thousand.

away from the eyebrows, whiskers, &c. the loss being preceded by intense itching and redness of the skin, it is due to the presence of an insect, which is a parasite of the horse-fly.

Milk-sick Literature .- The Editor of a Western Journal concludes a complimentary notice of an essay on milk sickness with the remark that it "will constitute a valuable addition to the milk-sick literature of our country."!

Adieu to the Year .- A writer in the French Lancet thus bids adieu to the past year:—
"Adieu then, remarkable year 1845! Medicine will long preserve the remembrance of thee,
—year fruitful of events as strange as unexpected -which hath witnessed the death of homocopathy and the birth of tachytomy, which hath seen chiromancy expire and hydropathy flourish."

Naval Medical Bureau. - Dr. Thomas Harris of Philadelphia has been appointed to the Bureau of Medicine and Surgery. Dr. Harris's high reputation as a practitioner and his honourable character as a man, will render this appointment popular in the Navy and of advantage to the ser-

Ligature of the external Iliac Artery.-Mr. B. B. Cooper performed this operation the 5th of March last, on a patient in Guy's Hospital with femoral aneurism.

Sir Henry Hulford .- The London papers recently received, announce the death of this distinguished physician.

Eclipse.-Among the phenomena foretold in a recent No. of the Gazette des Hôpitaux to occur

in 1844 is the following:

"A total eclipse visible in Paris and throughout all France of orthopædic cures,"

#### UNIVERSITY OF PENNSYLVANIA.-MEDICAL DEPARTMENT.-SESSION 1844-5

The Lectures will commence on Monday, the 4th day of November, and be continued, under the following arrangement, to the middle of March ensuing:—Practice and Theory of Medicine, by Nathaniel Chapman, M. D.; Chemistry, Robert Hare, M. D.; Surgery, William Gibson, M. D.; Anatomy, William E. Horner, M. D.; Institutes of Medicine, Samuel Jackson, M. D.; Materia Medica and Pharmacy, George B. Wood, M. D.; Obstetrics and the Diseases of Women and Children, Hugh L. Hodge, M. D. A Course of Clinical Lectures and Demonstrations, in connection with the above, is given at the very extensive and convenient Infirmary called the Philadelphia Hospital.

Clinical Medicine, by W. W. Gerhard, M. D.; Clinical Surgery, Drs. Gibson and Horner.

Clinical Instruction in Medicine is also given from the 1st day of November to the 1st day of

March by Dr. Wood, in the Pennsylvania Hospital, an institution which is well known as one of

the largest and best conducted Infirmaries in the United States.

The rooms for Practical Anatomy will be opened October 1st, and continued so to the end of the Course. They are under the charge of Paul Beck Goddard, M. D., Demonstrator, with a supervision on the part of Dr. Horner.

Extensive cabinets on Anatomy, Materia Medica, Chemistry, Surgery, and Obstetrics exist.

The Professor of Materia Medica, besides his Cabinet, has an extensive and well-furnished Conservatory, from which are exhibited, in the fresh and growing state, the native and exotic Medicinal Plants.

W. E. HORNER, M. D.,

April 1st, 1844.

Dean of the Medical Faculty, 263 Chestnut Street, Philadelphia.

#### JEFFERSON MEDICAL COLLEGE, SESSION OF 1844-45.

The regular Course of Lectures will commence on Monday the 4th of November, and end on the last day of February. Robley Dunglison, M. D., Professor of Institutes of Medicine, & c.; Robert M. Huston, M. D., Professor of Materia Medica and General Therapeutics; Joseph Pancoast, M. D., Professor of General, Descriptive, and Surgical Anatomy; John K. Mitchell, M. D., Professor of Practice of Medicine; Thomas D. Mütter, M. D., Professor of Institutes and Practice of Surgery; Charles D. Meigs, M. D., Professor of Obstetrics and Diseases of Women and Children; Franklin Bache, M. D., Professor of Chemistry. Lectures and Practical Illustrations will be given at the Philadelphia Hospital, one of the most extensive and valuable institutions in the United States, regularly through the Course, by Dr. Dunglison on Clinical Medicine, Dr. Pancoast on Clinical Surgery, and at the Dispensary of the College, by Professors of the Institution. The Dissecting Room will open on the first of October, under the Professor of Anatomy, and Clinical Instruction in Medicine and Surgery will be given at the Dispensary of the College.

R. M. HUSTON, M. D., Dean of the Faculty.

#### MEDICAL INSTITUTE OF PHILADELPHIA, LOCUST STREET ABOVE ELEVENTH.

The Course of Lectures, for 1844, will commence on the first Monday in April, and will continue until the end of the ensuing October, with the exception of a vacation from the 15th of July to the latter part of August.

Clinical Instruction in Medicine and Surgery will be given every afternoon during the season, and in order to make it as practical as possible, those students who desire it will be furnished

with cases.

Clinical Instruction in Midwifery will be given by the lecturer on that branch, and cases of labour provided for students who may wish to attend them.

Practical Instruction in Materia Medica and Pharmacy will also be given, and the Students

taught to compound prescriptions and prepare medicines.

Lectures on Anatomy, by Paul B. Goddard, M. D.; Surgery, W. Poyntell Johnson, M. D.; Obstetrics, M. P. Hutchinson, M. D.; Practice of Medicine, Caspar Morris, M. D.; Materia Medica, Joseph Carson, M. D.; Chemistry, James B. Rogers, M. D.; Clinical Medicine, W. W. Gerhard, M. D.; Clinical Surgery, W. P. Johnson, M. D. Fee for the course, including Clinical Instruction, \$70.

For tickets, apply to

PAUL B. GODDARD, Secretary, No. 7 South Ninth Street.

Lea & Blanchard have published the following new books and new editions, since their last list, for which see the proper heads, in the following pages. Andral on the Blood, Allison's Pathology, Dunglison's Human Physiology, 5th edition. Dunglison's Medical Dictionary, 4th ed.; Dunglison's Practice of Medicine, 8vo., second ed; Chapman on the Viscera; Cooper on Hernia; Cooper on Dislocations; Condie on Children; Smith and Horner's Anatomical Atlas; Prout on the Stomach: Williams's Pathology; Watson's Practice; Cyclopedia of Practical Medicine, part First, Second, and Third. Harrison on the Nerves; Wilson's Dissector, &c. &c.

Messrs. Geo. W. Carpenter and Co., Wholesale Druggists, No. 301 Market Street, Philadelphia, have recently published a large pamphlet of 96 pages, entitled "Carpenter's Annual Medical Advertiser for 1844." It contains an account of the most popular of the new medicines, Surgical Instruments, Anatomical Preparations, Chemical and Philosophical Apparatus, Medical Books, the improvements in Chemistry, Pharmacy, &c., and is distributed gratuitously to the physicians of the United States. Orders for any of the books mentioned in the accompanying list, can be executed by them.

2

## TO THE MEDICAL PROFESSION.

LEA AND BLANCHARD present a condensed list of books published and preparing for publication by them and would refer to the other pages for a more detailed account. The prices and all other information in relation to them will be given on application free of postage. Being extensively engaged in the publication of Medical and Scientific works, it will be their effort to furnish them at prices lower than formerly, and as low as they can be afforded consistent with correct and well executed editions. The latest editions will always be furnished, and to their present extensive list they will add from time to time such other good works as the wants of the profession may call for. Their publications may be found at all of the principal Bookstores throughout the Union. the Union.

Anatomical Atlas, by Smith and Horner, imperial 8vo., Harris on the Maxillary Sinus, I vol. small 8vo , 165 nearly ready.

Arnott's Elements of Physics, in 1 vol. Svo., 520 closely Horner's Special Anatomy, 2 vols. Svo., 6th edition, printed pages.

American Medical Journal, published quarterly at \$5 a Hodge on the Mechanism of Parturition in 1 vol. 4to., year. Fourteen numbers of the new series are now (preparing) with many plates Hope on the Heart, 1 vol 8vo., 572 pages. year. Fourteen numbers of the new series are now published.

Abercrombie on the Stomach, 1 vol. 8vo., 320 pages.

Abercrombie on the Brain, a new edition, 1 vol. 8vo. 324 pages.

Altison's Outlines of Pathology, in 1 vol. 8vo., 420 pages.

Ashwell on the Diseases of Females, complete in 1 vol., 1 Name on Ruptures, 1 vol. 8vo., 480 pages.

Manual Rectantist of Pathology with many plates

(preparing) with many plates

Hope on the Rectanism of Pathology with many plates.

Hope on the Rectanism of Pathology with many plates.

Hope on the Rectanism of Pathology with many plates.

Hope on the Rectanism of Pathology with many plates.

Hope on the Rectanism of Pathology with many plates.

Hope on the Heart, 1 vol. 8vo., 572 pages.

Harrison on the Nervous System, 1 vol. 8vo., 292 pages.

Kirby on Animals, many plates, 1 vol. 8vo., 578 pages.

Lawrence on the Eye, 1 vol. 8vo., 480 pages.

Name of Pathology with many plates. Svo , nearly ready. Andral on the Blood, 120 pages, Svo. Bell on the Teeth, with plates, 1 vol. Svo, 351 pages. Maury's Dental Surgery, with plates, a new work, I vol. 8vo., 285 pages Mütter's Surgery, 2 vols. Svo, now in preparation, with Buckland's Geology and Mineralogy, 2 vols. Svo., with numerous plates and maps.

Müller's Physiology, 1 vol. 8vo., 886 pages.

Berzelius on the Kidneys and Urine, 1 vol. small 8vo., Manual of Ophthalmic Medicine and Surgery, to be 179 pages.
Bridgewater Treatises, with numerous illustrations, 7 vols. 8vo., 3287 pages.
Bartlett on Fevers, &c., 1 vol. 8vo., 393 pages.
Bridgewater Treatises, with numerous illustrations, 7 vols. 8vo., 3287 pages.
Bartlett on Fevers, &c., 1 vol. 8vo., 393 pages.
Bridgewater Treatises, with numerous illustrations, 7 Medical News and Library, published monthly.
Manual of Medical Jurisprudence, in preparation.
Meigs' Translation of Colombat D'Isere on the Diseases of Females, in preparation, 1 vol. 8vo.
Prout on the Stomach and Renal Diseases, 1 vol. 8vo., with coloured plates, 465 pages.
Popular Medicine, by Coates, 1 vol. 8vo., 614 pages.
Philip on Protracted Indigestion, 1 vol. 240 pages.
Principles of Surgery, a new work, in 1 vol., 8vo.
Pereira's Materia Medica, 2 vols 8vo., 1566 very large and closely printed pages. 179 pages. published hereafter. Pereira's Materia Medica, 2 vols 8vo., 1566 very large and closely printed pages.

Carpenter's General and Comparative Physiology, 1 vol. 8vo., at press.

Carpenter's Vegetable Physiology, 1 vol. 12mo., with cuts, 300 pages.

Carpenter's Animal Physiology to be published hereafter.

Cooper, Sir Astley, his work on Hernia, imperial 8vo., with plates, 428 pages.

Cooper on Dislocations and Fractures, 1 vol. 8vo., with cuts, 499 pages.

Costello's Cyclopædia of Practical Surgery, to be published hereafter.

Churchill's Theory and Practice of Midwifery, 1 vol.

Pereira's Materia Medica, 2 vols 8vo., 1566 very large and closely printed pages.

Roget's Animal and Vegetable Physiology, with many cuts, 2 vols 8vo., 871 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Rigby's System of Midwifery, 1 vol 8vo., 256 pages.

Roget's Animal and Vegetable Physiology, with many cuts, 2 vols 8vo., 871 pages.

Rigby's System of Midwifery, 1 vol 8vo., 256 pages.

Roget's Animal and Vegetable Physiology, with many cuts, 2 vols 8vo., 871 pages.

Roget's Animal and Vegetable Physiology, with many cuts, 2 vols 8vo., 871 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 491 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Animal and Vegetable Physiology, 1 vol. 8vo., 516 pages.

Roget's Animal and Vegetable Physiology, 1 vol. 8vo., 516 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Animal and Vegetable Physiology, 1 vol. 8vo., 516 pages.

Roget's Animal and Vegetable Physiology, 1 vol. 8vo., 516 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 256 pages.

Roget's Outlines of Physiology, 1 vol. 8vo., 271 pages.

Roget's Outlines Churchill on Females, 1 vol 8vo., 595 large pages.
Churchill's Theory and Practice of Midwifery, 1 vol.
8vo., 519 pages, with cuts.
Cyclopedia of Practical Medicine by Forbes, &c. Edited by Dunglison, in 4 large super royal vols, at Tweedie on Diseases of the Nervous System, 1 Separate parts of Library of Medic Tweedie on Fevers, Inflammations and Cutavol 8vo. press Tweedie on Diseases of the Organs of Respira-Carson's Medical Formulary, in preparation. Dewees's System of Midwifery, with plates, 10th edition, tion, I vol. 8vo Tweedie on the Digestive, Urinary and Uterine Dewees's System of Midwhery, which of the Dewees's System of Midwhery, which of the Dewees's Practice of Physic, 1 vol. 8vo. 819 pages.

Dewees on Females, with plates, 8th edition, 532 pages.
Dewees's Practice of Physic, 1 vol. 8vo. 819 pages.
Dunglison's Physiology, 5th edition, 2 vols. 8vo., 1304 pages, with 300 cuts
Dunglison's Therapeutics and Materia Medica, a new work, 2 vols 8vo., 1004 pages
Dunglison's Medical Dictionary, 4th edition, 1 vol. 8vo.

Dunglison's Medical Dictionary, 4th edition, 1 vol. 8vo., 727 pages.
Toddi's Cyclopedia of Anatomy and Physiology, to be published hereafter.

Walshe's Diagnosis of the Diseases of the Lungs, 1 vol. 12mo, 310 pages.
Watson's Principles and Practice of Physic, 1 vol. 8vo., 576

Weedie on Rheumatism, Gout, Dropsy, Scott, 1 vol. 8vo., 234 pages.
Trimmer's Geology and Mineralogy, with many cuts, 1 vol. 8vo., 527 pages.
Toddi's Cyclopedia of Anatomy and Physiology, to be published hereafter.

Walshe's Diagnosis of the Diseases of the Lungs, 1 vol. 12mo, 310 pages.
Watson's Principles and Practice of Physic, 1 vol. 8vo., 576 771 very large pages.
Dunglison's New Remedies, 5th edition, 1843, 615 pages
Dunglison on Human Health, in 1 vol. 8vo, at press
Dunglison's Practice of Medicine, 2d edition, 2 vols
8vo, 1322 pages.
Dunglison' Medical Student, 12mo, a new edition, prepages.
Wilson's Dissector, or Practical and Surgical Anatomy,
Wilson's Dissector, or Practical and Surgical Anatomy, paring.

Druitt's Modern Surgery, 1 vol. 8vo., 534 pages.

Ellis's Medical Formulary, 7th edition, 1 vol. 8vo., 262

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages.

Wilson's Dissector, or Practical and Surgical Anatomy, by Goddard, with cuts, 1 vol. 12mo, 444 pages. I vol Svo. Youatt and Clater's Cattle Doctor, 1 vol. 12mo, with Elliotson's Mesmeric Cases, 8vo., 56 pages.
Fergusson's Practical Surgery, 1 vol. 8vo., 629 pages.
Graham's Chemistry, with cuts, 1 vol. 8vo., 750 pages.
Goddard's Dissector's Companion, in preparation.

cuts, 282 pages. Williams' Pathology, 1 vol. 8vo., 383 pages.

## CHAPMAN'S NEW WORK.

## LECTURES ON THE MORE IMPORTANT DISEASES

OF THE

# THORACIC

DELIVERED IN THE UNIVERSITY OF PENNSYLVANIA.

## By N. CHAPMAN, M. D.

Professor of the Theory and Practice of Medicine, &c.

In one volume, octavo.

#### CONTAINING

PHTHISIS PULMONALIS, CYNANCHE LARYNGEA, ASTHMA, ANGINA PECTORIS, GASTRITIS. CHRONIC GASTRITIS, ORGANIC LESIONS OF THE STOMACH, DYSPEPSIA, ENTERITIS, DUODENAL DYSPEPSIA, CHRONIC FLUXES OF THE BOWELS, &c. &c. &c.

CONSTIPATIO, HEPATITIS, CONGESTION OF THE LIVER, HEPATICULA, ICTERUS SPLENITIS. CHRONIC SPLENITIS, ENGORGEMENT OF THE SPLEEN, CHRONIC CONGESTION OF THE SPLEEN,

## MOW READY.

#### THE FIRST PART

OF THE

# CYCLOPÆDIA OF PRACTICAL MEDICINE,

EDITED BY

## DRS. TWEEDIE, FORBES, CONOLLY, AND DUNGLISON.

#### CONTAINING ARTICLES ON

ABDOMEN, EXPLORATION OF THE, Dr. Forbes. | AMAUROSIS, Dr. Lee. ABORTION, Dr. Tweedie. ABSCESS (INTERNAL), ABSTINENCE, ACHOR, ACNE, ACRODYNIA, ACUPUNCTURE, AGE, AIR (CHANGE OF), ALOPECIA. ALTERATIVES, APOPLEXY (CEREBRAL)

AMENORRHŒA, Dr. Tweedie.
Dr. Marshall Hall.
Dr. Todd.
Dr. Todd.
ANASARCA,
ANGINA PECTORIS,
ANODYNES, Dr. Dunglison. ALTHELMINTICS, Dr. Thomson.
Dr. Elliotson. ANTHRACION, Dr. Dunglison.
Dr Roget. ANTIPHLOGISTIC REGISTER. Dr Roget.

Sir James Clark.

ANTIPHLOGISTIC REGIMEN, Dr. Barlow.

ANTISPASMODICS,

Dr. Thomson. Dr. Todd.
Dr. Conolly.
Dr. Clutterbuck.
AORTA, ANEURISM OF THE,
APHONIA,
APHTHÆ,

Dr. Locock. Dr. Marshall Hall. Dr. Darwall. Dr. Forbes. Dr. Whiting. Dr. Dunglison. Dr. Hope. Dr. Robertson.

For further details see the Prospectus, terms, &c., annexed, but the pages are not so well printed there as they are in the work. The publishers are prepared to send a specimen of the work that will exhibit the mechanical execution as well as its professional character. In presenting this as one of the cheapest works that has yet been offered to the profession, the publishers beg leave to state that persons remitting five dollars can have the first ten parts sent them by mail or otherwise, as they may direct.

# LEA AND BLANCHARD,

PHILADELPHIA,

WILL PUBLISH

# THE CYCLOPÆDIA OF PRACTICAL MEDICINE:

COMPRISING

TREATISES ON THE

NATURE AND TREATMENT OF DISEASES,

# MATERIA MEDICA AND THERAPEUTICS,

MEDICAL JURISPRUDENCE, &c. &c.

EDITED BY

JOHN FORBES, M.D., F.R.S.
Physician in Ordinary to her Majesty's Household, &c.

ALEXANDER TWEEDIE, M.D., F.R.S.

Physician to the London Fever Hospital, and to the Foundling Hospital, &c.

JOHN CONOLLY, M.D.

Late Professor of Medicine in the London University, and Physician to the Hanwell Lunatic Asylum, &c.

# THOROUGHLY REVISED, WITH ADDITIONS, BY ROBLEY DUNGLISON, M.D.

Professor of the Institutes of Medicine, &c. in the Jefferson Medical College, Philadelphia,
Lecturer on Clinical Medicine, and Attending Physician at the Philadelphia
Hospital; Secretary of the American Philosophical Society, &c.

# TERMS OF PUBLICATION.

THE WORK WILL BE PRINTED WITH A NEW AND CLEAR TYPE, AND BE COMPRISED IN

TWENTY-FOUR PARTS, AT FIFTY CENTS EACH,

FORMING, WHEN COMPLETE,

FOUR LARGE SUPER-ROYAL OCTAVO VOLUMES,

EMBRACING OVER

THREE THOUSAND UNUSUALLY LARGE PAGES,

IN DOUBLE COLUMNS.

Any person forwarding Twenty Dollars, free of postage, in Current Funds, will be entitled to two copies. The whole work will be completed during the year

# PUBLISHERS' NOTICE

## CYCLOPÆDIA OF PRACTICAL MEDICINE.

This important work consists of a series of Original Essays upon all subjects of Medicine, contributed by no less than SIXTY-SEVEN of the most eminent practical Physicians of Great Britain and Ireland, and among them many of the Professors and Teachers in London, Edinburgh, Dublin and Glasgow, whose reputation conveys a high and just authority to their doctrines. Each subject has been treated by a writer of acknowledged eminence, whose particular studies have eminently fitted him for the task; and all the articles are authenticated with the names of the authors.

The Editors are men of elevated attainments, and in the undertaking have spared no personal pains; in the hope, by uniformity of plan, simplicity of arrangement, and the harmony and consistency of its several portions, to make the Cyclopædia represent, fully and fairly, the state of PRACTICAL MEDICINE at the time of its appearance. From innumerable foreign and domestic sources, the scattered knowledge, which has so fast accumulated since the commencement of the present century, has been gathered together and placed at the command of every reader of the English language; and whilst the great claims of the older cultivators of Medicine have never been forgotten, the labours of the moderns, and more particularly of the French, German, and Italian Pathologists, by which, conjointly with the efforts of British and American Practitioners, the whole face of Practical Medicine has been changed, have attracted the most diligent and thoughtful attention.

The Editors affirm, that if the reader will take the trouble to inspect the mere titles of the articles contained in the work, comprising nearly

#### THREE HUNDRED ORIGINAL ESSAYS

of known and distinguished authors, and will bear in mind either the leading physiological divisions of disease, or consider them with reference to the Head, the Chest, the Abdomen, the Surface, or the general condition of the body, as well as the subjects of OBSTETRICAL MEDICINE, MATERIA MEDICA, or MEDICAL JURISPRUDENCE, he will sufficiently appreciate the care bestowed to make the Cyclopædia satisfactory to all who refer to its pages, and, at the same time, strictly a book of practical reference. No subject, it is believed, immediately practical in its nature or application, has been omitted; although unnecessary disquisition has been, as much as possible, avoided.

It entered consistently and properly into the plan of the Editors to admit a far wider range of subjects than appears heretofore to have been considered necessary in works written professedly on the Practice of Medicine, but a range comprising many new subjects of extreme importance to those

engaged in practice, or preparing for it. Such are the subjects of

ABSTINENCE :	CONTAGION	EXPLORATION OF	PROGNOSIS
ACUPUNCTURE !	CONVALESCENCE	THE CHEST AND	PULSE
AGE	COUNTER-IRRITATION	ABDOMEN	SOFTENING
CHANGE OF AIR	CONGESTION AND DE-	GALVANISM	MEDICAL STATISTICS
ANTIPHLOGISTIC	TERMINATION OF	HEREDITARY TRANS.	STETHOSCOPE
REGIMEN	BLOOD	MISSION OF DISEASE	SUDDEN DEATH
ASPHYXIA	DERIVATION	INDURATION	SYMPTOMATOLOGY
AUSCULTATION	DIETETICS	IRRITATION	TEMPERAMENT
BATHING	DISINFECTION	INFECTION	TOXICOLOGY
BLOOD-LETTING	PHYSICAL EDUCATION	LATENT DISEASES	TRANSFORMATION
MORBID STATES	ELECTRICITY	MALARIA AND MIASMA	TRANSFUSION
OF THE BLOOD	ENDEMIC DISEASES	PERFORATION	TUBERCLE
CLIMATE	EPIDEMICS	PSEUDO-MORBID AP-	VENTILATION
COLD	EXPECTORATION	PEARANCES	MINERAL WATERS

and those of various general articles on the pathology of organs. It will be found, too, that admirable articles from the best sources have been inserted on the important subjects of

DISEASES OF WOMEN AND CHILDREN, AND OF MEDICAL JURISPRUDENCE. In order, however, that the nature and value of the work may be fully understood, a list of the

articles, and the names of the contributors, is appended.

The excellence of this work on every topic connected with Practical Medicine, has been admitted by all who have had the good fortune of being able to consult it. The hope, indeed, expressed by the Editors, has been amply realized - "That they have prepared a work required by the present wants of medical readers, acceptable to the profession in general, and so capable, by its arrangements, of admitting the progressive improvements of time, as long to continue what the general testimony of their medical brethren, as far as it has hitherto been expressed, has already pronounced it to be,

#### "A STANDARD WORK ON THE PRACTICE OF MEDICINE."

Such a work, it is believed, will be most acceptable to the members of the profession throughout the Union, as there exists, at this time, no publication on Practical Medicine, on the extended plan of the one now presented.

To adapt it to the Practice of this country, and to thoroughly revise the various articles, the attention of

## PROFESSOR DUNGLISON

will be directed; whose character and established reputation are a sure guarantee that his portion of the work will be carefully executed.

# CONTENTS OF, AND CONTRIBUTORS TO,

# THE CYCLOPÆDIA OF PRACTICAL MEDICINE.

HI DI CD D	Control De Western
Abdomen, Exploration of, Dr. FORBES.	Counter Irritation Dr. WILLIAMS.
Abortion Dr. LEE.	Croup Dr. CHEYNE.
Abscess Dr. Tweedie.	Cyanosis Dr. CRAMPTON.
Abstinence Dr. M. Hall.	Cystitis Dr. Cumin.
Achor Dr. Topp.	Delirium Dr. PRITCHARD.
Acne Dr. Topp.	Delirium Tremens Dr. CARTER.
Acupuncture : Dr. Elliotson.	Dentition, Disorders of . Dr. Joy.
Acupuncture : Dr. Elliotson.	Derivation Dr. STOKES.
Age Dr. ROGET.	Determination of Blood Dr. BARLOW.
Air, Change of, Dr. CLARK.	Dishetes Dr. Bardsley.
Alopecia Dr. Todd.	Diabotto I I I I I
Alteratives Dr. Conolly.	Diaphoretics Dr. A. T. Thomson.
Amaurosis Dr. Jacob.	Diarrhea SDr. CRAMPTON.
Amenorrhœa Dr. Locock.	(DI. I ORBES.
Anæmia Dr. M. HALL.	Dietetics Dr. PARIS.
Anasarca Dr. Darwall.	Dilatation of the Heart . Dr. Hope.
Angina Pectoris Dr. Forbes.	Disease Dr. Conolly.
Angina rectoris Dr. PORBES.	Disinfection Dr. Brown.
Anodynes Dr. Whiting.	To totti contain a a a a a a a a a a a a a a a a a a
Anthelmintics Dr. A. T. THOMSON.	
Antiphlogistic Regimen Dr. BARLOW.	Dropsy Dr. DAEWALL.
Antispasmodics Dr. A. T. THOMSON.  Aorta, Aneurism of, . Dr. Hope.	Dysentery Dr. Brown.
Aorta, Angurism of, . Dr. Hope.	Dysmenorrhæa Dr. Locock.
Aphonia Dr. Robertson.	Dysphagia Dr. STOKES.
Aphthæ Dr. Robertson.	Dyspnœa Dr. Williams.
Application Company Dr. Common Dr	Dysuria Dr. Cumin.
Apoplexy, Cerebral, . Dr. CLUTTERBUCK.	Diponent
Apoplexy, Pulmonary, . Dr. Townsend.	
Arteritis Dr. Hope.	
Artisans, Diseases of, . Dr. Darwall.	Education, Physical, Dr. BARLOW.
Ascites Dr. DARWALL.	Electricity Dr. APJOHN.
Asphyxia Dr. Roger.	Elephantiasis Arabum . Dr. Scott.
Asthma Dr. Forbes.	Elephantiasis Græcorum Dr. Joy.
Astringents Dr. A. T. Treaseast	Emetics Dr. A. T. THOMSON.
Astringents Dr. A. T. Thomson.	Emmenagogues Dr. A. T. THOMSON.
Atrophy Dr. Townsend.	To an
Auscultation Dr. FORBES.	
Barbiers Dr. Scott.	Emphysema of the Lungs Dr. Townsend.
Bathing Dr. FORBES.	Empyema Dr. Townsend.
Beriberi Dr. Scott.	Endemic Diseases Dr. HANCOCK.
Blood, Morbid States of, Dr. M. HALL.	Enteritis Dr. STOKES.
Bloodletting Dr. M. HALL.	Ephelis Dr. Topp.
CD. Orres	Enidemics Dr. HANCOCK.
Brain, Inflammation of, {Dr. QUAIN. Dr. ADAIR CRAWFORD.	Epilepsy Dr. Cheyne.
Dr. ADAIR CRAWFORD.	Epistaxis Dr. Kerr.
Bronchitis Dr. WILLIAMS.	Epistaxis Dr. Kerr. Erethismus Mercurialis Dr. Burder.
Bronchocele Dr. And. CRAWFORD.	Erethismus Mercurialis Dr. BURDER.
Bullæ Dr. Topp.	Erysipelas Dr. Tweedie.
Calculus Dr. Thos. Thomson.	Erythema Dr. Joy.
Calculous Diseases Dr. CUMIN.	Expectorants Dr. A. T. THOMSON.
Catalepsy Dr. Joy.	Expectoration Dr. Wilson.
Catarrh Dr. WILLIAMS.	Favus Dr. A. T. THOMSON
	CDr. Scott.
	Feigned Diseases 3 Dr. Forbes.
	CDr. MARSHALL.
Chicken Pox Dr. GREGORY.	Fever, General Doctrine of, Dr. TWEEDIE.
Chlorosis , Dr. M. HALL.	1 O and ital
Cholera Dr. Brown.	Modifications 2 11. 1 WEBBIR
Chorea Dr. And. CRAWFORD.	
Climate Dr. CLARK.	a prince
Cold Dr. Whiting.	Epidemie Gustre
CDr. WHITING.	D- Promy
Colic Dr. Tweedle.	Teliminone
777	" Infantile Remittent Dr. Joy.
To 1 - Courses	" Hectic Dr. Brown.
Coma :	
	1 Duarparal . Dr. LEE.
Combustion, Spontane- Dr. APJOHN.	" Puerperal Dr. Gillkrest.
ous Human, ?	" Yellow Dr. Gillerest. "Yellow Dr. Kerr.
ous Human, Dr. Barlow.	" Puerperal Dr. BEE. " Yellow Dr. GILLKREST. Fungus Hœmatodes . Dr. KERR. Galvanism Dr. APJOHN.
Congestion of Blood . Dr. Barlow.	" Puerperal Dr. Bee. " Yellow Dr. Gillerest. Fungus Hæmatodes . Dr. Kerr. Galvanism Dr. Apjohn. Dr. Stokes.
Congestion of Blood . Dr. Barlow.  Constipation Dr. Streeten.	" Puerperal Dr. BEE. " Yellow Dr. GILLKREST. Fungus Hœmatodes . Dr. KERR. Galvanism Dr. APJOHN. Gastritis Dr. Sarrows
Congestion of Blood . Dr. Barlow.  Constipation Dr. Barlow.  Dr. Hastings.  Dr. Streeten.  Dr. Brown.	" Puerperal Dr. BEE. " Yellow Dr. GILLKREST. Fungus Hœmatodes Dr. KERR. Galvanism Dr. APJOHN. Gastritis Dr. STOKES. Gastrodynia Dr. BARLOW.
Congestion of Blood Constipation Contagion Con	" Puerperal Dr. BEE. " Yellow Dr. GILLKREST. Fungus Hœmatodes . Dr. KERR. Galvanism Dr. APJOHN. Gastritis Dr. STOKES. Gastrodynia Dr. STOKES. Gastro-Enteritis . Dr. STOKES. Dr. KERR.
Congestion of Blood Constipation Contagion Convalescence  Or. Barlow.  Dr. Hastings.  Dr. Streeten.  Dr. Brown.  Dr. Brown.  Dr. Tweedie.  Dr. Apair Crawford.	" Puerperal . Dr. DEE. " Yellow . Dr. GILLKREST. Fungus Hæmatodes . Dr. Kerr. Galvanism . Dr. Apjohn. Dr. Stokes. Gastrodynia . Dr. Barlow. Gastro-Enteritis . Dr. Stokes. Glossitis . Dr. Kerr.
ous Human,	" Puerperal Dr. BEE. " Yellow Dr. GILLKREST. Fungus Hœmatodes . Dr. Kerr. Galvanism Dr. Apjohn. Gastritis Dr. Stokes. Gastrodynia Dr. Barlow. Gastro-Enteritis Dr. Stokes. Glossitis Dr. Kerr. Glossitis Dr. Joy.
ous Human,	" Puerperal . Dr. BEE. " Yellow . Dr. GILLKREST. Fungus Hæmatodes . Dr. KERR. Galvanism . Dr. Apjohn. Dr. Stokes. Gastrodynia . Dr. Barlow. Gastro-Enteritis . Dr. Stokes. Glossitis . Dr. KERR. Glossis, Spasm of the, Dr. Joy. Gout . Dr. Barlow.
ous Human,	" Puerperal . Dr. Dr. Dr. GILLKREST.  "Yellow . Dr. GILLKREST.  Fungus Hœmatodes . Dr. Kerr.  Galvanism . Dr. Stokes.  Gastritis . Dr. Barlow.  Gastrodynia . Dr. Stokes.  Glossitis . Dr. Kerr.  Glossitis . Dr. Kerr.  Glossitis . Dr. Kerr.  Glossitis . Dr. Jov.

	D. C
Headach Dr. Burder.	Pityriasis Dr. Cumin. Plague Dr. Brown.
Heart, Diseases of the, Dr. Hope.	
" Displacement of the, Dr. Townsend.	
Hematemesis Dr. Goldie.	
Hemoptysis Dr. Law.	Plica Polonica Dr. Corrigan. Pneumonia Dr. Williams.
Hemorrhage Dr. WATSON.	Pneumothorax Dr. Houghton.
Hereditary Transmis- sion of Disease . { Dr. Brown.	Porrigo Dr. A. T. THOMSON.
Herpes Dr. A. T. Thomson.	Pregnancy, &c., Signs of, Dr. Montgomery.
	Prognosis Dr. Ash.
	Pseudo morbid Annear Co
Hooping-Cough Dr. Johnson. Hydatids Dr. Kerr.	Pseudo-morbid Appear- Dr. R. B. Topp.
Hydatids Dr. Kerr. Hydrocephalus Dr. Joy.	Psoriasis Dr. Cumin.
	Puerperal Diseases Dr. Hall.
Hydrophobia Dr. Darwall.  Hydrophobia Dr. Bardsley.	Pulse Dr. Bostock.
Hydrothorax Dr. Darwall.	Purigo Dr. A. T. THOMSON.
Hypertrophy Dr. Townsend.	Purpura Dr. Goldie.
Hypertrophy of the Heart Dr. Horg.	Pyrosis Dr. KERR.
Hypochondriasis Dr. PRITCHARD.	Rape Dr. BEATTY.
Hysteria Dr. Conolly.	Refrigerants Dr. A. T. THOMSON.
Icthyosis Dr. A. T. THOMSON.	Rheumatism Dr. Barlow.
Identity Dr. MONTGOMERY.	Rickets Dr. Cumin.
Impetigo Dr. A. T. THOMSON.	Roseola Dr. Tweedie.
Impotence Dr. BEATTY.	Rubeola Dr. Montgomery.
Incontinence of Urine . Dr. Cumin.	Rupia Dr. Corrigan.
Incubus Dr. WILLIAMS.	Rupture of the Heart . Dr. Townshend.
Indigestion Dr. Topp.	Scabies Dr. Houghton.
Induration Dr. CARSWELL.	Scarlatina Dr. Tweedie.
Infanticide Dr. Arrowsmith.	Scirrhus Dr. CARSWELL.
Infection Dr. Brown.	Scorbutus Dr. KERR.
Inflammation Dr. A. CRAWFORD, Dr. TWEEDIE.	Scrofula Dr. Cumin.
Influenza Dr. HANCOCK.	Sedatives Dr. A. T. THOMSON.
Insanity Dr. PRITCHARD.	Sex, Doubtful, Dr. BEATTY.
Irritation Dr. WILLIAMS.	Small-Pox Dr. Gregory.
Ischuria Renalis Dr. Carter,	Softening of Organs . Dr. CARSWELL.
Jaundice Dr. Burder.	Somnambulism and Ani- Dr. PRITCHARD.
Kidneys, Diseases of, . Dr. Carter.	
Lactation Dr. Locock.	Soundness, &c., of Mind Dr. PRITCHARD.
Laryngitis Dr. Cheyne.	Spinal Marrow, Dis- Spr. R. B. Topp.
Latent Diseases Dr. Christison.	eases of,
Lepra Dr. Houghton.	Spleen, Diseases of, . Dr. Bigsby.
Leucorrhœa Dr. Locock.	Statistics, Medical, Dr. HAWKINS.
Lichen Dr. Houghton. Liver, Inflammation of, Dr. Stokes.	Stethoscope Dr. WILLIAMS.
Liver, Inflammation of, Dr. Stokes.	Stimulants Dr. A. T. THOMSON.
" Diseases of Dr. VENABLES.	Stomach, Organic Dis- S Dr. Houghman
Malaria and Miasma . Dr. Brown.	
Malformations of the Heart Dr. WILLIAMS.	Succession of Inherit- Dr. Montgomery.
Medicine, Principles and Dr. CONOLLY.	The state of the s
	Suppuration Dr. R. B. Topp.
Melæna Dr. Goldie.	Survivorship Dr. BEATTY.
Melanosis Dr. CARSWELL.	Sycosis Dr. Cumin.
Menorrhagia Dr. Lococк.	Symtomatology Dr. M. Hall.
Menstruation, Pathology of Dr. Locock.	Syncope Dr. Ash.
Miliaria, Dr. Tweedie.  Mortification Dr. Carswell.	Tabes Mesenterica Dr. Joy.
	Temperament Dr. PRITCHARD.
Narcotics Dr. A. T. Thomson.	Tetanus Dr. Symonds.
Nephralgia and Nephritis Dr. CARTER.	Throat, Diseases of the, Dr. Tweedle. Tonics Dr. A. T. Thomson.
Neuralgia Dr. Elliotson.	
Noli me tangere, or Lupus Dr. Houghton.	Toxicology Dr. Apjohn. Transformations Dr. Duesbury.
Nyctalopia Dr. Grant. Obesity Dr. Williams.	Transfusion Dr. Kay.
Œdema Dr. DARWALL.	Tubercle Dr. Carswell.
Ophthalmia Dr. Jacob.	Tubercular Phthisis Dr. CLARK.
Otalgia and Otitis Dr. Burne.	Tympanites Dr. Kerr.
Ovaria, Diseases of the Dr. Lee.	Urine, Morbid States of, Dr. Bostock.
Palpitation Dr. Hope,	Urine, Bloody, Dr. Goldie.
Pancreas, Diseases of the, Dr. Carter.	Urticaria Dr. Houghton.
Paralysis Dr. R. B. Topp.	Uterus, &c. Pathology of, Dr. LEE.
Paralysis Dr. R. B. Todd. Parotitis Dr. Kerr.	Vaccination Dr. Gregory
Pellagra Dr. KERR.	Varicella Dr. GREGORY.
Pemphigus Dr. Corrigan.	Veins, Diseases of, Dr. LEE.
Perforation of the Hol- { Dr. CARSWELL.	Ventilation Dr. Brown.
low Viccore 3 Dr. CARSWELL.	
low viscera	Wakefulness Dr. Cheyne.
Pericarditis and Carditis Dr. Hore.	Wakefulness Dr. CHEYNE. Waters, Mineral, Dr. T. THOMSON.
Pericarditis and Carditis Dr. Hope. Peritonitis Dr. M'ADAM, Dr. STOKES.	Wakefulness Dr. CHEYNE. Waters, Mineral, Dr. T. THOMSON. Worms Dr. Joy.
Pericarditis and Carditis Dr. Hope. Peritonitis Dr. M'ADAM, Dr. STOKES. Persons found Dead . Dr. BEATTY.	Wakefulness Dr. CHEYNE. Waters, Mineral, Dr. T. THOMSON. Worms Dr. Joy. Wounds, Death from, . Dr. Beatty.
Pericarditis and Carditis Dr. Hope. Peritonitis Dr. M'ADAM, Dr. STOKES.	Wakefulness Dr. CHEYNE. Waters, Mineral, Dr. T. THOMSON. Worms Dr. Joy.

# DUNGLISON'S PRACTICE,

SECOND EDITION.

A NEW EDITION, TO 1844,

OF

# THE PRACTICE OF MEDICINE;

ON

# SPECIAL PATHOLOGY AND THERAPEUTICS.

SECOND EDITION.

BY ROBLEY DUNGLISON, M. D.,

Professor of the Institutes of Medicine in the Jefferson Medical College, Lecturer on Clinical Medicine,
Attending Physician at the Philadelphia Hospital, &c.

In two large octavo volumes, of over thirteen hundred pages.

Notwithstanding the short period in which this work has been before the profession, it is so universally known throughout the country, and so extensively used as a text-book in the Medical Colleges, that, in presenting this new and much improved edition, the publishers have only to call attention to the author's preface.

"It is scarcely necessary for the author to say, that the extraordinary sale, which the first edition of this work has met with, is most gratifying to him. It sufficiently satisfies him, that the large amount of labour and reflection, which he bestowed upon it, has been found serviceable to his medical brethren, and more especially so, perhaps, to those who are engaged in the study of their profession.

"Grateful for this result, the author has endeavoured to render the present edition still more useful, by adding whatever of importance has transpired in the short period that has elapsed since the first edition was published, and by supplying omissions, which were almost inevitable in the first impression of a work in which so many subjects are treated of.

"It has been pleasing to the author to observe the favourable manner in which the work has been noticed by almost all the Journals of Medicine, which have received it at home and abroad; and where criticisms have been occasionally made, they have generally been rather against the plan than the manner of execution. That plan was adopted from a full and deliberate conviction of its advantages, and it is, therefore, adhered to in the present edition without modification.

"The solid testimony, which the Profession have afforded in favour of the work will incite the author to untiring exertion to render every succeeding edition more and more valuable."

# THE PUBLISHERS ANNEX A CONDENSED STATEMENT OF THE CONTENTS.

Diseases of the Mouth, Tongue, Teeth, Gums, Velum Palati and Uvula, Pharynx and Esophagus, Stomach, Intestines, Peritoneum, Morbid Productions in the Peritoneum and Intestines.—Diseases of the Larynx and Trachea, Bronchia and Lungs, Pleura, Asphyxia.—Morbid Conditions of the Blood, Diseases of the Heart and Membranes, Arteries, Veins, Intermediate or Capillary Vessels,—Spleen, Thyroid Gland, Thymus Gland, and Supra Renal Capsules, Mesenteric Glands,—Salivary Glands, Pancreas, Biliary Apparatus, Kidney, Ureter, Urinary Bladder.—Diseases of the Skin, Exanthematous, Vesicular, Bullar, Pustular, Papular, Squamous, Tuberculous, Maculæ, Syphilides.—Organic Diseases of the Nervous Centres, Neuroses, Nerves.—Diseases of the Eye, Ear, Nose.—Diseases of the Male and Female Organs of Reproduction—Fever.—Intermittent, Remittent, Continued, Eruptive, Arthritic, Cachex, Scrofulous, Scorbutic, Chlorotic, Rhachitic, Hydropic, and Cancerous.

# WATSON'S PRACTICE OF MEDICINE. LECTURES

ON THE

# PRINCIPLES AND PRACTICE OF PHYSIC.

DELIVERED AT KING'S COLLEGE, LONDON.

BY THOMAS WATSON, M.D.

FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS, PHYSICIAN TO THE MIDDLESEX HOSPITAL, &C. &C.

In one large octavo volume, of over nine hundred unusually large pages, well bound in leather. Containing NINETY LECTURES.

This volume has met with the universal approbation of the medical press, so far as it has yet been expressed, in a manner rarely enjoyed by so comprehensive a work. The publishers submit a few extracts.

"For our parts, we are not only willing that our characters as scientific physicians and skilful practitioners may be deduced from doctrines contained in this volume, but we hesitate not to express our belief, that for sound, trustworthy principles, and good substantial practice, it cannot be paralleled by any similar work in any other country. We would advise no one to set himself down in practice unprovided with a copy."—British and Foreign Medical Review.

"We have here the sterling production of a liberal, well-stored, and truly honest mind, possessed of all that is currently known and established of professional knowledge, and capable of pronouncing a trustworthy and impartial judgment on those numerous points in which truth is yet obscured by false hypotheses."—Provincial Med Journal.

"Open this huge, well furnished volume where we may, the eye immediately rests on something that carries value on its front. We are impressed at once with the strength and depth of the lecturer's views. He gains on our admiration in proportion to the extent of our acquaintance with his profound researches. Whoever owns this book, will have an acknowledged treasure if the combined wisdom of the highest authorities is appreciated." Boston Med. and Surg Journal.

"We know of no other work better calculated for being placed in the hands of the student, and for a text-book, and as such we are sure it will be very extensively adopted. On every important point, the author seems to have posted up his knowledge to the day."—American Medical Journal.

"We know not, indeed, of any work of the same size that contains a greater amount of useful and interesting matter. We are satisfied, indeed, that no physician, well-read and observing as he may be, can rise from its perusal without having added largely to his stock of valuable information."—Medical Examiner.

"It is an admirable digest of general pathology and therapeutics. As a text book for medical schools, it cannot be surpassed, and in no other treatise can practitioners find so concise, and at the same time so complete a summary of the present state of the science of medicine "—Baltimore Patriot.

"One of the most practically useful books that ever was presented to the student—indeed, a more admirable summary of general and special pathology, and of the application of therapeutics to diseases, we are free to say has not appeared for very many years. The lecturer proceeds through the whole classification of human ills, a capite ad calcem, showing at every step an extensive knowledge of his subject, with the ability of communicating his precise ideas, in a style remarkable for its clearness and simplicity "-N. Y Journal of Medicine, \$\psi\_c\$.

"The style is correct and pleasing, and the matter worth the attention of all practitioners, young and old."-Western Lancet.

In order to insure to this volume as wide a circulation as it is entitled to by its merits, the publishers have put it at a price within the reach of all. It recommends itself to practitioners as well as students, to the owners of other works on practice, as well as those without—something new can be gleaned by all from its well-stored pages.

# TWEEDIE'S LIBRARY OF PRACTICAL MEDICINE,

A SYSTEM OF PRACTICAL MEDICINE.

Comprised in a Series of Original Dissertations, arranged and edited by Alexander Tweedie, M. D., F. R. S., &c., &c.

With notes and additions, by W. W. Gerhard, M. D., &c.

SECOND AMERICAN EDITION,

In three volumes octavo.

The design of this work is to supply the want, generally admitted to exist in the medical literature of Great Britain, of a comprehensive System of Medicine, embodying a condensed, yet ample view of the present state of the science. The desideratum is more especially felt by the Medical Student, and by many Members of the Profession, who, from their avocations and other circumstances, have not the opportunity of keeping pace with the more recent improvements in the most interesting and useful branch of human knowledge. To supply this deficiency is the object of The Library of Medicine; and the Editor expresses the hope, that with the assistance with which he has been favoured by Contributors, (many of great eminence, and all favourably known to the Public,) he has been able to produce a work, which will form a Library of General Reference on Theoretical and Practical Medicine, as well as a series of Text Books for the Medical Student.

## DEWEES'S PRACTICE.

A PRACTICE OF PHYSIC. Comprising most of the diseases not treated of in "Diseases of Females" and "Diseases of Children." By William P. Dewees, M. D., &c. &c. Second edition, in one large 8vo. volume.

# HORNER'S ANATOMY.

# SPECIAL ANATOMY AND HISTOLOGY.

BY WILLIAM E. HORNER, M. D.,

Professor of Anatomy in the University of Pennsylvania. Member of the Imperial Medico-Chirurgical Academy of St. Petersburg, of the Am. Philosophical Society, &c. &c.

Sixth edition, in two volumes, 8vo..

"Another edition of this standard work of Professor Horner has made its appearance to which many additions have been made, and upon which much labour has been bestowed by the author. The additions are chiefly in the department of Histology, or Elementary Anatomy, and so important are they that the Professor has added the term to the title of his work. Every part of this edition seems to have undergone the most careful revision, and its readers may rest assured of having the science of Anatomy fully brought up to the present day."—Am Med. Journal.

An Atlas that may accompany this work or be sold separate is now preparing .- See Advertisement.

# WILSON'S HUMAN ANATOMY.

# A SYSTEM OF HUMAN ANATOMY, GENERAL AND SPECIAL,

BY ERASMUS WILSON, M.D.,

Lecturer on Anatomy, in King's College, London.
EDITED BY PAUL B. GODDARD, M.D.,

PROPERCIAL OF ANATOMY IN THE MEDICAL INSTITUTE OF BUILDED BY

## WITH ONE HUNDRED AND SEVENTY ILLUSTRATIONS ON WOOD.

Engraved by Gilbert from designs by Bagg.

In one volume octavo, beautifully printed.

"An elegant edition of one of the most useful and accurate systems of The illustrations are really beautiful, and their execution reflects the highest credit on the able American artist who copied them for this edition of the work. In its style the work is extremely concise and intelligible. Dr. Goddard has added a number of valuable notes, and made some judicious alterations of names. No one can possibly take up this volume without being struck with the great beauty of its mechanical execution, and the clearness of the descriptions which it contains is equally evident. Let students by all means examine the claims of this volume on their notice, before they purchase a text-book of the vitally important science which it so fully and easily unfolds."—Lancet.

# WILSON'S DISSECTOR. THE DISSECTOR,

# PRACTICAL AND SURGICAL ANATOMY.

BY ERASMUS WILSON, M.D.,

AUTHOR OF "A SYSTEM OF HUMAN ANATOMY," &c.

With one hundred and six illustrations, modified and re-arranged by

PAUL B. GODDARD, M.D.,

DEMONSTRATOR OF ANATOMY IN THE UNIVERSITY OF PENNSYLVANIA, &c. In one volume, royal 12mo.

"Those who pursue their labours by this monitor will speedily discover its merits as a plain, safe director; and it is only necessary to use it half an hour to find that we are furnished with an improved edition of an excellent work."—Boston Medical and Surg. Journal.

"The high reputation which Mr. Wilson has acquired, by his valuable system of anatomy, will secure a favourable reception for the present work. As modified and re arranged by Dr. Goddard, it is well suited to favourable reception for the present work. As modified and re arranged by Dr. Goddard, it is well suited to favourable reception for the present work, and is an excellent guide for him in the dissecting room. It the wants of the American student of anatomy, and is an excellent guide for him in the dissecting room. It is very copiously illustrated with wood cuts, which will greatly facilitate the labours of the student in acies very copiously illustrated with wood cuts, which will greatly facilitate the labours of the student in acies were a supplied to the student of anatomy and is an excellent guide for him in the dissecting room. It is very copiously illustrated with wood cuts, which will greatly facilitate the labours of the student in ac-

"It strikes us as being all that a "Dissector" should be. The wood cuts are numerous, and will afford the student the most essential aid in the dissecting room."—Western Journal of Med. & Surgery.

"In this work we have another valuable aid to the student of practical anatomy."-N. Y. Journal of Medicine.

# SMITH AND HORNER'S ANATOMICAL ATLAS.

LEA and BLANCHARD are now Publishing, AN

ANATOMICAL ATLAS,

# ILLUSTRATIVE OF THE STRUCTURE OF THE HUMAN BODY.

By HENRY H. SMITH, M. D.

Fellow of the College of Physicians, Member of the Philadelphia Medical Society, &c., &c.

UNDER THE SUPERVISION OF

#### WILLIAM E. HORNER, M.D.

Professor of Anatomy in the University of Pennsylvania, &c. &c.

This Work is to be in FIVE PARTS, and will, when complete, form one large imperial Octavo Volume. The price will not be over

## SIX DOLLARS, BOUND, And will contain a Complete Series of Plates.

Illustrative of the Bones; Ligaments; Muscular, Dermoid and Nervous Systems; Arteries; Lymphatics; Veins; Viscera; &c.; with between

## Six and Seven Hundred Engravings on Wood

Executed in the very first style of Art.

Parts I. and II. have been published, containing the Bones and Liga-MENTS, and the MUSCULAR and DERMOID SYSTEMS, illustrated by over Two Hundred and Twenty Engravings on Wood, by Gilbert. The greater portion of them after Original Drawings by Pinkerton.

This work possesses novelty both in the design and the execution. It is the first attempt to apply engraving on wood, on a large scale, to the illustration of human anatomy; and the beauty of the two parts already issued, induces the publishers to flatter themselves with the hope of the perfect success of their undertaking. The plan of the work is at once novel and convenient. Each page is perfect in itself, the references being immediately under the figures, so that the eye takes in the whole at a glance, and obviates the necessity of continual reference backwards and forwards. The cuts are selected from the best and most accurate sources; and, where necessary, original drawings have been made from the admirable Anatomical Collection of the University of Pennsylvania. It will also embrace all the late beautiful discoveries arising from the use of the microscope in the investigation of the minute structure of the tissues.

In the getting up of this very complete work, the publishers have spared neither pains nor expense; and they now present it to the Profession, with the full confidence that it will be deemed all that is wanted in a scientific and artistical point of view, while, at the same time, its very low price places it within the reach of all.

It is particularly adapted to supply the place of skeletons or subjects.

It is particularly adapted to supply the place of skeletons or subjects.

"This is an exquisite volume, and a beautiful specimen of art. We have numerous Anatomical Atlases, but we will venture to say that none equal it in cheapness, and none surpass it in faithfulness and spirit. We strongly recommend to our friends, both urban and suburban, the purchase of this excellent work, for which both editor and publisher deserve the thanks of the profession."—Medical Examiner.

"We would strongly recommend it, not only to the student, but also to the working practitioner, who, although grown rusty in the toils of his harness, still has the desire, and often the necessity, of refreshing his knowledge in this fundamental part of the science of medicine."—New York Journal of Medicine.

"The plan of this Atlas is admirable, and its execution superior to any thing of the kind before published in this country. It is a real labour-saving affair, and we regard its publication as the greatest boon that could be conferred on the student of anatomy. It will be equally valuable to the practitioner, by affording him an easy means of recalling the details learned in the dissecting-room, and which are soon forgotten."—American Journ. of the Med Sciences.

"The second part of Drs. Smith and Horner's Anatomical Atlas, in point of interest of subject and execution of the engravings, is even more valuable and attractive than the first, much as that led us to expect. Rich aids to physiological inquiry will be found in the microscopical views of the cellular and adipose tissues, and of the epidermis, rete mucosum, cutis vera, papillæ, the sebaceous and perspiratory organs of the skin, the perspiratory glands and hairs of the skin, the hairs and nails, and the general anatomy of the muscles. The muscles themselves are delineated with all the distinctness of a fine dissection. We cannot, taking the two numbers of the Anatomical Atlas before us as evidence of the execution of the entire work, too strongly recommend it to readers and the medical public at large."—Bulletin of Medical Science.

Specimens exhibiting the plan and execution of the work will be sent by mail or otherwise as may be directed on application free of postage.

# PHYSIOLOGY AND PATHOLOGY.

DUNGLISON'S PHYSIOLOGY. A NEW EDITION, TO 1844.

# HUMAN PHYSIOLOGY: WITH UPWARDS OF THREE HUNDRED ILLUSTRATIONS.

By ROBLEY DUNGLISON, M. D.

Professor of the Institutes of Medicine in Jefferson Medical College, Philada., etc. etc.

FIFTH EDITION,

GREATLY MODIFIED AND IMPROVED.

In Two Volumes of 1304 large Octavo Pages.

Although but a short time has elapsed since the appearance of the last edition of this standard work on Physiology, the labours and discoveries in that branch of science have been so diversified and important, that extensive additions have become necessary in the present edition. Besides the numerous monographs and minor treatises that have appeared from time to time, in the various scientific Journals and Transactions of learned Societies, reference has been made to the large and valuable works of Valentin, Bischoff, Henle, Wilbrand, J. Müller, Wagner, Mandl, Gerber, Liebig, Carpenter, Todd and Bowman, and others, and all the results of their interesting and important investigations will be found embodied in its pages.

The illustrations have also undergone a material change. More than ninety new cuts have been added, chiefly to elucidate subjects now first introduced into this edition; many of the former cuts have been replaced by those much superior, and the rest have been thoroughly retouched and improved, altogether making this the most completely illustrated treatise on Human Physiology that has ever been attempted in this country. In short, the publishers believe that, both in matter and execution, this work will be found fully brought up to the day.

# CARPENTER'S PHYSIOLOGY.

PRINCIPLES OF HUMAN PHYSIOLOGY,

With their chief applications to Pathology, Hygiene, and Forensic Medicine. Especially designed for the use of Students.

With over One Hundred beautiful Illustrations on Wood.

BY WILLIAM B. CARPENTER, M.D.,

LECTURER ON PHYSIOLOGY IN THE BRISTOL MEDICAL SCHOOL. FIRST AMERICAN EDITION, WITH NOTES BY THE AUTHOR, AND NOTES AND ADDITIONS BY MEREDITH CLYMER, M.D.,

In one volume, octavo.

This edition of Carpenter's Physiology has been most carefully prepared by Dr. Clymer, at the request of Professor Jackson, for his lectures at the University of Pennsylvania.

The Author remarks in his Preface—"It has been his object to place the reader in possession of the highest principles that can be regarded as firmly established, in each department of the science; and to explain and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these, by the introduction of as many important facts as could be included within moderate and illustrate these.

# MULLER'S PHYSIOLOGY.

ELEMENTS OF PHYSIOLOGY.

BY JOHN MÜLLER, M.D.,

Professor of Anatomy and Physiology in the University of Berlin.

TRANSLATED FROM THE GERMAN, BY W. BALY, M. D. Arranged from the 2d London edition by John Bell, M. D., &c.

In one volume octavo, of nearly nine hundred pages.

# WILLIAMS, PATHOLOGY.

(A New Work.)

# PRINCIPLES OF MEDICINE,

COMPRISING

GENERAL PATHOLOGY AND THERAPEUTICS, and a general view of ETIOLOGY, NOSOLOGY, SEMEIOLOGY, DIAGNOSIS AND PROGNOSIS.

BY CHARLES J. B. WILLIAMS, M.D., F.R.S.

FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS, ETC.

## WITH ADDITIONS AND NOTES BY MEREDITH CLYMER, M.D.

LECTURER ON THE INSTITUTES OF MEDICINE, ETC. ETC.

In one volume 8vo.

" Such a work, at least so far as English literature is concerned, was clearly demanded, and the importance of the subject, together with the high character of the author, should ensure for it a careful perusal by every practitioner."-Western Lancet.

"Such a work, then, as that of Professor Williams, embodying, as it does, a full account of the recent additions to our knowledge, and bringing the subjects of general pathology and therapeutics up to the present state of medicine, cannot fail to be in the highest degree useful, both as a class book to the student, and as a

work of reference to the practitioner.

"The entire work is one of no ordinary ability, and well sustains the reputation which its author has already acquired."—Provincial Med. Journal.

"This work supplies a deficiency long felt by the profession—the want of an elementary treatise on general pathology. The author has had extensive experience at the bed side, and has drawn chiefly from his own observation of disease, as he has witnessed it in hospitals and private practice during the last twenty years.

"The subject of inflammation occupies many pages of this work. The manner in which it is treated cannot fail to excite the admiration of the intelligent reader. We earnestly recommend this, as every other part of Dr. Williams' excellent Principles of Pathology, to the diligent perusal of every physician who is not familiar with the accessions which have been made to medical science within the last few years."—Western Journal of Medicing and Surgery. of Medicine and Surgery.

### ROGET'S OUTLINES OF PHYSIOLOGY.

WITH AN APPENDIX ON PHRENOLOGY. BY PETER MARK ROGET, M. D., &c. &c. In one volume, octavo.

ROGET'S PHYSIOLOGY.

ANIMAL AND VEGETABLE PHYSIOLOGY, considered with reference to Natural Theology. By PETER MARK ROGET, M.D., second American edition. In two volumes, octavo, with nearly FIVE HUNDRED engravings on wood.

# CARPENTER'S VEGETABLE PHYSIOLOGY.

A POPULAR TREATISE ON VEGETABLE PHYSIOLOGY. Published under the auspices of the Society for the promotion of Popular Instruction. With numerous cuts. By W. B. CARPENTER, M.D. author of Human Physiology, in one volume, royal 12mo.

FIRST PRINCIPLES OF MEDICINE. By ARCHIBALD BILLING, M.D., A.M. &c. Revised from the fourth London edition. In one small 8vo. volume.

# PATHOLOGY, A NEW WORK.

#### OUTLINES OF PATHOLOGY

AND

PRACTICE OF MEDICINE.

BY

#### WILLIAM PULTENEY ALLISON, M. D.,

Professor of the Practice of Medicine in the University of Edinburgh, &c. &c.

In three Parts-Part I - Preliminary Observations-Part II .- Inflammatory and Febrile Diseases, and Part III., Chronic or Non-Febrile Diseases.

In one volume octavo.

# MIDWIFERY, DISEASES OF WOMEN, CHILDREN, &c.

# RAMSBOTHAM'S MIDWIFERY.

THE PRINCIPLES AND PRACTICE

OF

# OBSTETRIC MEDICINE AND SURGERY.

IN REFERENCE TO THE

## PROCESS OF PARTURITION.

ILLUSTRATED BY ONE HUNDRED AND FORTY-TWO FIGURES, ON 52 PLATES.
BY FRANCIS H. RAMSBOTHAM, M. D.

PHYSICIAN TO THE ROYAL MATERNITY CHARITY, &c. &c. second american edition, revised.

In one volume, imperial octavo.

Well and strongly bound in leather, at a low price.

This work has met with almost unexampled approbation from the medical press of England and this country. From among these numerous commendations the publishers append a few, and would particularly call the attention of the medical profession to the numerous plates which form a most important feature in the volume. The great expense they have incurred in its production, calls for an extended sale, which thus far has been commanded, as may be seen from the fact of a second edition having been required within a year after the publication of the first.

"It is the book on Midwifery for students: clear, but not too minute in its details, and sound in its practical instructions—It is so completely illustrated by plates (admirably chosen and executed) that the student must be stupid indeed who does not understand the details of this branch of the science, so far at least as description can make them intelligible."—Dublin Journal of Medical Science.

"We strongly recommend the work of Dr. Ramsbotham to all our obstetrical readers, especially to those who are entering upon practice. It is not only one of the cheapest, but one of the most beautiful works in Midwifery."—British and Foreign Medical Review.

"We feel much pleasure in recommending to the notice of the profession one of the cheapest and most elegant productions of the medical press of the present day. The text is written in a clear, concise, and simple style. We offer our most sincere wishes that the undertaking may enjoy all the success which it so well merits."—Dublin Medical Press.

"This is one of the most beautiful works which has lately issued from the medical press; and is alike creditable to the talents of the Author and the enterprise of the publisher. It is a good and thoroughly practical treatise; the different subjects are laid down in a clear and perspicuous form, and whatever is of importance is illustrated by first rate engravings. A remarkable feature of this work, which ought to be mentioned, is its extraordinary cheapness. As a work conveying good, sound, practical precepts, and clearly demonstrating the doctrines of obstetrical science, we can confidently recommend it either to the student or practitioner."—

Edinburgh Journal of Medical Science.

"Dr. Ramsbotham has treated the subject in a manner worthy of the reputation he possesses, and has succeeded in forming a book of reference for practitioners, and a solid and easy guide for students. Looking at the contents of the volume, and its remarkably low price, we have no hesitation in saying that it has no parallel in the history of publishing."—Provincial Medical and Surgical Journal.

"We most earnestly recommend this work to the student who wishes to acquire knowledge, and to the practitioner who wishes to refresh his memory, as a most faithful picture of practical Midwifery; and we can with justice say, that altogether it is one of the best books we have read on the subject of obstetrical medicine and surgery."

—Medico-Chirurgical Review.

## DEWEES'S SYSTEM OF MIDWIFERY.

## A COMPREHENSIVE SYSTEM OF MIDWIFERY.

CHIEFLY DESIGNED TO FACILITATE THE INQUIRIES OF THOSE WHO MAY BE PURSUING THIS BRANCH OF STUDY.

ILLUSTRATED BY OCCASIONAL CASES AND MANY ENGRAVINGS.
Tenth Edition, with the Author's last Improvements and Corrections.

BY WILLIAM P. DEWEES, M. D.

LATE PROFESSOR OF MIDWIFERY IN THE UNIVERSITY OF PENNSYLVANIA, &c. In one volume, Svo.

# Devees of Females.

A TREATISE ON THE DISEASES OF FEMALES, BY WILLIAM P. DEWEES, M.D., late Professor of Midwifery in the University of Pennsylvania, &c. &c.

Eighth edition, revised and corrected, in one volume, 8vo., with plates.

# CHURCHILL'S MIDWIFERY.

# ON THE THEORY AND PRACTICE OF MIDWIFERY,

BY FLEETWOOD CHURCHILL, M.D., M. R. I. A. PHYSICIAN TO THE WESTERN LYING-IN HOSPITAL, ETC. ETC.

#### WITH NOTES AND ADDITIONS

BY ROBERT HUSTON, M.D.,

Professor in the Jefferson Medical College, &c. &c.

## AND ONE HUNDRED AND SIXTEEN ILLUSTRATIONS,

Engraved by Gilbert from drawings by Bagg and others.

In one volume octavo.

This work commends itself to the notice of the profession from the high reputation of the author and editor, and the number and beauty of its illustrations. Besides accurate directions for THE PRACTICE OF MIDWIFERY,

a portion of the work is also devoted to

#### THE PHYSIOLOGY AND PATHOLOGY

connected with that essential branch of medical knowledge.

"It is impossible to conceive a more useful or elegant manual: the letter-press contains all that the practical man can desire; the illustrations are very numerous, well chosen, and of the most elegant description, and the work has been brought out at a moderate price."—Provincial Med. Journal.

"We expected a first rate production, and we have not been in the least disappointed. Although we have many, very many valuable works on tokology, were we reduced to the necessity of possessing but one, and permitted to choose, we would unhesitatingly take Churchill."—Western Med. and Surg. Journal.

This work is brought out in a style to match Wilson's Anatomy, Fergussons' Surgery, and Carpenter's Physiology, and is extensively used as a Text Book in various Colleges in the Union.

# ASHWELL ON THE DISEASES OF WOMEN.

A NEW WORK, Nearly Ready.

# A PRACTICAL TREATISE

ON THE

# ISEASES OF WOMEN,

Illustrated by cases derived from Hospital and private practice.

BY SAMUEL ASHWELL, M.D.,

Obstetric Physician and Lecturer to Guy's Hospital, &c. &c.

WITH NOTES AND ADDITIONS

#### BY PAUL GODDARD, M.D.

Demonstrator of Anatomy in the University of Pennsylvania, &c. &c. In one vol. octavo, in three parts.

Part I.—Functional Diseases. Part II. & III.—Organic Diseases.

"Situated as is Dr. Ashwell in extensive practice, and at the head of the Obstetric Department of a large Hospital, it could not be but that his work must contain very valuable information—the results of great experience. The book is full of important information and excellent practical description."—Dublin Medical Journal.

"The contributions of Dr. Ashwell coming as they do, armed with the authority of his personal observation and great experience, are eminently entitled to consideration. Throughout every page we have only the result of the author's own observation; and of the work we entertain a high opinion, believing that it is characterized by accurate observation and sound judgment."—Edinburgh Monthly Journal.

## DISEASES OF FEMALES, PREGNANCY AND CHILDBED.

THE PRINCIPAL DISEASES OF FEMALES, TOGETHER WITH THE DISEASES INCIDENT TO PREGNANCY AND CHILD-BED, CHIEFLY FOR THE USE OF STUDENTS. BY FLEETWOOD CHURCHILL, M,D.,

Lecturer on Midwiferyand Diseases of Women and Children, in the Richmond Hospital &c. &c.

WITH NOTES AND ADDITIONS BY R. M. HUSTON, M.D.,

Professor, &c. in the Jefferson Medical College, Philadelphia. Second American edition, in one vol. octavo.

A work extensively introduced as a Text Book.

# A NEW AND IMPORTANT WORK ON MIDWIFERY.

LEA and BLANCHARD have now in Preparation

# A NEW WORK ON THE MECHANISM OF LABOUR.

By HUGH L. HODGE, M. D.

Professor of Obstetrics and the Diseases of Women and Children in the University of Pennsylvania, assisted in the illustrative part

BY PAUL BECK GODDARD, M.D.,

Demonstrator of Anatomy in the University of Pennsylvania.

This important work is intended to exhibit the various OPERATIONS OF DELIVERY, with full instructions for the use of the MOST APPROVED INSTRUMENTS, and means to facilitate child-birth. To illustrate this, plates will be used on an extended scale.

The work will form

# One Large Imperial Quarto Volume,

NUMEROUS MAGNIFICENT PLATES, Executed in Lithography in the best Style of the Art UNDER THE IMMEDIATE INSPECTION OF DR. GODDARD. And will aim to be the most perfect and accurate work yet presented to the Profession in this country.

In the Figures a uniform scale will be adopted, representing about HALF THE SIZE OF NATURE.

The price cannot be precisely arranged, but it is at present not designed to exceed

# EIGHT DOLLARS PER COPY,

A Specimen, with a Plate, will shortly be issued, and sent to such persons as may request it, free of postage.

"Numerous publications have appeared abroad, and much has been attempted, by books and by plates, but almost universally there is a want of accuracy in portraying nature's operations, and embracing fully the science as practised by intelligent obstetricians, in this country, which leads

necessarily to errors in practice.

"The object of the author, therefore, will be to present an accurate history, and, as far as practicable, a delineation of the process of labour, whether natural or preternatural, in the human female; and, also, of the chirurgical means which may be advantageously employed to effect delivery in protracted and difficult labours, so as to diminish the amount of suffering and danger for the mother and her offspring. In other words, to exemplify as minutely as possible, by drawings, the natural 'mechanism of labour,' and the scientific employment of the hand or of instruments in facilitating this painful and frequently dangerous process. No one who is ignorant of the details of labour in every presentation of the child, can advantageously or even safely assist in cases of acknowledged difficulty; an attempt, therefore, to exhibit more fully and minutely the transit of the child through the pelvis in all its various presentations, and the practical results of the knowledge thus acquired, cannot but be advantageous to the practitioner, and, especially, to the student in medicine."

# RIGBY'S MIDWIFERY.

A SYSTEM OF MIDWIFERY, WITH NUMEROUS WOOD CUTS, BY EDWARD RIGBY, M.D., Physician to the General Lying-in Hospital, Lecturer on Midwifery at St. Bartholomew's Hospital, &c. With notes and additional illustrations. In one volume octavo. This volume commends itself for its fulness on the Physiology of Utero-Gestation, with the Diseases

and treatment of Pregnancy-

# CONDIE ON CHILDREN.

A NEW WORK.

# A PRACTICAL TREATISE

ON

#### DISEASES OF CHILDREN.

BY D. FRANCIS CONDIE, M.D.

FELLOW OF THE COLLEGE OF PHYSICIANS; MEMBER OF THE AMERICAN PHILOSOPHICAL SOCIETY, &c. &c.

In one volume, octavo.

The Publishers would particularly call the attention of the Profession to an examination of this work.

THE VOLUME IS METHODICALLY ARRANGED IN TWO PARTS,

#### IN PART I. ARE CONSIDERED

The Hygienic Management of Children. The Peculiarities of Organization and Functions in Infancy and Childhood. Pathology of Infancy and Childhood. And the Semeiology of the Diseases of Infancy and Childoood.

#### IN PART II. ARE EMBRACED

Discases of the Digestive Organs.

The Mouth, Throat, Œsophagus, Stomach, Intestines, Respiratory Organs, Nervous System, the Skin, Eruptive Fevers, Cutaneous Eruptions, Nutritive Functions, and Urinary Organs. And lastly, Congenital Affections, and Accidents occurring soon after Birth.

But this is, however, only an outline of the subjects brought under special notice.

"Dr. Condie has studiously endeavoured to be understood by students, who need to have the elements of Therapeutics presented to them in a comprehensive form."

"An excellent Practical Treatise on the Diseases of Children, and a very safe guide to the juvenile practi-tioner and student."—Med. Examiner.

"For the practical physician who shall turn to its pages to learn all the phenomena which may be presented by the disease which he is treating, and all the means to which he may resort for the cure of that disease, it will offer many and strong attractions, among which may be mentioned completeness, clearness, judgment and good sense. In it the vanity of the author never tempts the compiler into negligence; nor does the laborious care of the compiler weigh down and overlay the original vigour of the author: the two

offices are made to strengthen and illustrate one another.

"Dr. Condie from the very labour which he has evidently bestowed upon this book, is entitled to our respect as an indefatigable and conscientious student; but if we consider the results of his labour, we cannot but

admit his claim to a place in the very first rank of eminent writers on the practice of medicine.

"Regarding his treatise as a whole, it is more complete and accurate in its descriptions, while it is more copious and more judicious in its therapeutical precepts than any of its predecessors, and we feel persuaded that the American medical profession will very soon regard it, not only as a very good, but as the very best 'Practical treatise on the Diseases of Children.'"—Am Med. Journal:

## dewies on children.

A TREATISE ON THE PHYSICAL AND MEDICAL TREATMENT OF CHILDREN. BY WILLIAM P. DEWEES, M.D., late Professor of Midwifery in the University of Pennsylvania, &c. eighth edition, with the author's last improvements and corrections. In one volume 8vo.

The objects of this work are, 1st, to teach those who have the charge of children, either as parent or guardian, the most approved methods of securing and improving their physical powers. This is attempted by pointing out the duties which the parent or the guardian owes for this purpose, to this interesting but helpless class of beings, and the manner by which their duties shall be fulfilled. And 2d, to render available a long experience to those objects of our affection when they become diseased. In attempting this, the author has avoided as much as possible. 'technicality;" and has given, if he does not flatter himself too much, to each disease of which he treats, its appropriate and designating characters, with a fidelity that will prevent any two being confounded together, with the best mode of treating them, that either his own experience or that of others has suggested. Physicians cannot too strongly recommend the use of this book in all families. Physicians cannot too strongly recommend the use of this book in all families.

# SURGERY.

## FERGUSSON'S PRACTICAL SURGERY.

# A SYSTEM OF PRACTICAL SURGERY,

BY WILLIAM FERGUSSON, F. R. S. E.,

Professor of Surgery in King's College, London; Surgeon to King's College Hospital, &c. &c.

WITH TWO HUNDRED AND FORTY-SIX ILLUSTRATIONS.

Engraved by Gilbert, after drawings by Bagg.

#### WITH NOTES AND ADDITIONAL ILLUSTRATIONS

BY GEORGE W. NORRIS, M.D.,

In one volume octavo.

with a clear, sound, and practical treatment of every subject in surgical science. No pains or expense have been spared to present it in a style equal, if not superior to the London edition, and to match the editions of "Wilson's Anatomy," "Churchill's System of Midwifery," and "Carpenter's Human Physiology." It is now extensively used as a text book.

The object and nature of this volume are thus described by the author: "The present work has not been produced to compete with any already before the Profession; the arrangement, the manner in which the subjects have been treated, and the illustrations, are all different from any of the kind in the English language. It is not intended to be placed in comparison with the elementary systems of Cooper, Burns, Liston, Symes, Lizars, and that excellent epitome of Mr. Druitt. It may with more propriety be likened to the Operative Surgery of Sir C. Bell, and that of Mr. Averill, both excellent in their day, or the more modern production of Mr. Hargrave, and the Practical Surgery of Mr Liston. There are subjects treated of in this volume, however, which none of these gentlemen have noticed; and the author is sufficiently sanguine to entertain the idea that this work may in some degree assume that relative position in British Surgery, which the classical volumes of Velpeau and Malzaigne occupy on the Continent." Malgaigne occupy on the Continent."

A volume on the Principles of Surgery to accompany Fergusson will be prepared.

# DRUITT'S MODERN SURGERY.

# THE PRINCIPLES AND PRACTICE OF MODERN SURGERY,

BY ROBERT DRUITT.

Id potissimum agens, ut omissis hypothesibus, in praxi nihil adstruat quod multiplici experientià non sit roboratum.-Act. Erud. Lips., 1722.

FROM THE SECOND LONDON EDITION,

ILLUSTRATED WITH FIFTY WOOD ENGRAVINGS.

WITH NOTES AND COMMENTS BY

JOSHUA B. FLINT, M. D., M. M. S. S., &c. &c.

In one volume, octavo.

It may be said with truth that this work affords a complete, though brief and condensed view, of the entire field of modern surgery. We know of no work on the same subject, having the appearance of a manual, which contains so many topics of interest to the surgeon, and the terse manner in which each has been treated evinces a most enviable quality of mind on the part of the author, who seems to have an innate power of searching out and grasping the leading facts and features of the most elaborate productions of the pen. It is a useful handbook and grasping the leading facts and features of the most elaborate productions of the pen. It is a useful handbook for the practitioner, and we should deem a teacher of surgery unpardonable who did not recommend it to his for the practitioner, and we should deem a teacher of surgery unpardonable who did not recommend it to his pupils. In our own opinion, it is admirably adapted to the wants of the student."—Provincial Med. Journal.

# LAWRENCE ON RUPTURES.

A TREATISE ON RUPTURES, BY W. LAWRENCE, F.R.S. Author of a Treatise on the Diseases of the Eye, &c. &c., from the Fifth London Edition, considerably enlarged. In 1 vol. octavo.

The peculiar advantage of the treatise of Mr. Lawrence is, that he explains his views on the anatomy of Hernia, and the different varieties of the disease, in a manner which renders his book peculiarly useful to the student. It must be superfluous to express our opinion of its value to the surgical practitioner. As a treatise on Hernia, presenting a complete view of the literature of the subject, it stands in the first rank.—

Edinburgh Medical and Surgical Journal.

## SIR ASTLEY COOPER ON HERNIA. With One Hundred and Thirty Figures in Lithography.

# THE ANATOMY AND SURGICAL TREATMENT

# ABDOMINAL HERNIA. By SIR ASTLEY COOPER, BART.

Edited by C. Aston Key, Surgeon to Guy's Hospital, &c.

This important work of Sir Astley is printed from the authorized second edition, published in London, in large super-royal folio, and edited by his nephew, Professor Key. It contains all the Plates and all the Letterpress—there are no omissions, interpolations, or modifications—it is the complete work in

One Large Imperial Octavo Volume,

WITH OVER 130 FIGURES ON 26 PLATES, AND OVER 400 LARGE PAGES OF LETTERPRESS.

The correctness of the Plates is guaranteed by a revision and close examination under the eye of a distinguished Surgeon of this city.

The value of this Work of Sir Astley Cooper is so universally acknowledged by all medical men, that in presenting this edition to the American Profession, the publishers have only to state that they have used their utmost endeavours to render the mechanical execution of the work worthy its exalted reputation, and to put it in such a form and at such a price as to place it within the reach of those who have been prevented from obtaining it by the size and rarity of former editions.

# SIR ASTLEY COOPER'S WORK. ON FRACTURES AND DISLOCATIONS

WITH NUMEROUS WOOD CUTS.

A TREATISE ON DISLOCATIONS AND FRACTURES OF THE JOINTS. By SIR ASTLEY COOPER, BART., F.R.S., Sergeant Surgeon to the King, &c.

A new edition much enlarged; edited by BRANSBY COOPER, F.R.S., Surgeon to Guy's Hospital, with additional Observations from Professor John C. Warren, of Boston. With numerous engravings on wood, after designs by Bugg, a memoir and a splendid portrait of Sir Astley. In one vol. octavo.

The peculiar value of this, as of all Sir Astley Cooper's works, consists in its eminently practical character. His nephew, Bransby B. Cooper, from his own experience, has added a number of cases. Beside this, Sir Astley left behind him very considerable additions in MS. for the express purpose of being introduced into this edition. The volume is embellished with ONE HUNDRED AND THIRTY-THREE WOOD CUTS, and contains the history of no less than three hundred and sixty one cases, thus embodying the records of a life of practice of the Author and his various editors. There are also additional Observations from notes furnished by John C. Warren, M.D., the Professor of Anatomy and Surgery in Harvard University.

# MAURY'S DENTAL SURGERY.

# A TREATISE ON THE DENTAL ART,

FOUNDED ON ACTUAL EXPERIENCE.

ILLUSTRATED BY TWO HUNDRED AND FORTY-ONE FIGURES IN LITHOGRAPHY, AND FIFTY-FOUR WOOD CUTS:

> BY B. F. MAURY, DENTIST OF THE ROYAL POLYTECHNIC SCHOOL. Translated from the French, with Notes and Additions,

BY J. B. SAVIER, DOCTOR OF DENTAL SURGERY.

One volume, octavo.

This work is used as a Text book in the Baltimore College of Dental Surgery, and commends itself to the Profession from the great reputation of the author, and as embracing the latest information on the subject—Its steady demand is the best testimony of the general favour with which the profession has received it. It is distinct the second of the subject is the best testimony of the general favour with which the profession has received it. vided into three parts.

Part I .- Dental Anatomy, Physiology and Pathology.

Part II .- Dental Hygiene and Therapcutics.

Part III.—Mechanical Dentistry in all its various parts and fully illustrated.

#### THE DISEASES OF THE EYE. WITH NUMEROUS CUTS.

#### A TREATISE ON THE DISEASES OF THE EYE. BY W. LAWRENCE,

Surgeon Extraordinary to the Queen, &c., from the last London Edition, with numerous additions, and sixtyseven Illustrations, many of which are from original drawings.

#### BY ISAAC HAYS, M.D.,

Surgeon to the Wills Hospital, &c. &c., in one volume octavo.

The character of this work is too well established to require a word of commendation—it is justly considered the best on the subject. The present is a reprint of the last London Edition, which appeared in 1841, completely revised and greatly enlarged by the author—and to it considerable additions have been made by the editor. Several subjects omitted in the original are treated of in this edition, on which occasion free use has been made of the work of Mackenzie, to which is added the editor's own experience, derived from many years' attention to the subject. tention to the subject.

# MATERIA MEDICA, THERAPEUTICS, PHARMACY, AND CHEMISTRY.

#### PEREIRAIS MATERIA MEDICA.

WITH NEAR THREE HUNDRED ENGRAVINGS ON WOOD.

# THE ELEMENTS OF MATERIA MEDICA AND THERAPEUTICS.

COMPREHENDING THE NATURAL HISTORY, PREPARATION, PROPERTIES, COMPOSITION, EFFECTS, AND USES OF MEDICINES.

BY JONATHAN PEREIRA, M. D., F. R. S. and L. S.

Member of the Society of Pharmacy of Paris; Examiner in Materia Medica and Pharmacy of the University of London; Lecturer on Materia Medica at the London Hospital, &c. &c.

From the Second London Edition, enlarged and improved.

WITH NOTES AND ADDITIONS

## BY JOSEPH CARSON, M. D.

In two volumes, octavo.

Part I. contains the General Action and Classification of Medicines, and the Mineral Materia Medica. Part II, the Vegetable and Animal Kingdoms, and including diagrams explanatory of the Processes of the Pharmacopoias, a tabular view of the History of the Materia Medica, from the earliest times to the present day, and a very copious index. From the Second London Edition, which has been thoroughly revised, with the Introduction of the Processes of the New Edinburgh Pharmacopoia, and containing additional articles on Mental Remedies, Light, Heat, Cold, Electricity, Magnetism, Exercise, Dietetics, and Climate, and many additional Wood Cuts, Illustrative of Pharmaceutical Operations, Crystallography, Shape and Organization of the Feculas of Commerce, and the Natural History of the Materia Medica.

The object of the author has been to supply the Medical Student with a Class Book on Materia Medica, containing a faithful outline of this Department of Medicine, which should embrace a concise account of the most important discoveries in Natural History, Chemistry, Physiology, and Therapeutics, in so far as they pertain to Pharmacology, and treat the subjects in the order of their natural historical relations.

This great Library or Cyclopedia of Materia Medica has been fully revised, the errors corrected, and numerous additions made by DR JOSEPH CARSON, Professor of Materia Medica and Pharmacy in the "College of Pharmacy," and forms Two Volumes, octavo, of near 1600 large and closely-printed pages. It may be fully relied upon as a permanent and standard work for the country,—embodying, as it does, full references to the U.S. Pharmacopoia and an account of the Medical Plants indigenous to the United States.

# ELLIS'S MEDICAL FORMULARY, 7th Edition.

THE MEDICAL FORMULARY; Being a Collection of Prescriptions, derived from the Writings and Practice of many of the most eminent physicians in this country and in Europe. To which is added an Appendix containing the usual Dietetic Preparations and Antidotes for Poisons; the whole accompanied by a few brief Pharmaceutic and Medical Observations.

BY BENJAMIN ELLIS, M. D.

Seventh Edition, revised and extended by Samuel George Morton, M. D. In one volume octavo.

# GRAHAM'S CHEMISTRY. THE ELEMENTS OF CHEMISTRY,

Including the application of the Science to the Arts.

WITH NUMEROUS ILLUSTRATIONS, BY THOMAS GRAHAM, F.R.S., L. and E. D.

Professor of Chemistry in University College, London, &c. &c.

With Notes and Additions, BY ROBERT BRIDGES, M. D., &c. &c.

In one vol. octavo.

The great advancement recently made in all branches of chemical investigation renders necessary a new text book which shall clearly elucidate the numerous discoveries, especially in the department connected with organic Chemistry and Physiology, in which such gigantic strides have been made during the last few years. The present treatise is considered by eminent judges to fulfil all these indications, and to be peculiarly adapted to the wants of the medical student and practitioner. In adapting it to the wants of the American profession, to the wants of the editor has endeavoured to render his portion of the work worthy the exalted reputation of the first chemist, the editor has endeavoured to render his portion of the work worthy the exalted reputation of the first chemist, of England. It is already introduced as a text book many of the Colleges, and has universal approbation. "Professor Graham's work is one of the best, if not the best, of all English text books, and is of such recent "Professor Graham's work is one of the best, if not the best, of all English text books, and is of such recent date as to embrace all the latest discoveries. The appearance of a correct and amended American Edition, under the care of Dr Bridges, will prove an acceptable thing to both teachers and students of chemistry in this country."—Silliman's Journal.

# DUNGLISON'S MATERIA MEDICA, &c. GENERAL THERAPEUTICS AND MATERIA MEDICA. BY ROBLEY DUNGLISON, M.D.,

Professor of the Institutes of Medicine, &c. &c., in the Jefferson Medical College, Philada.

In two volumes octavo.

"A second edition of the work on General Therapeutics, being called for by the publishers, the author has deemed it advisable to incorporate with it an account of the different articles of the Materia Medica. To this he has been led by the circumstance, that the departments of General Therapeutics and Materia Medica are always associated in the Medical Schools. The author's great object has been to prepare a work which may aid the Medical Student in acquiring the main results of modern observation and reflection; and at the same time, be to the Medical Practitioner a trustworthy book of reference.

"Throughout, he has adopted the Nomenclature of the last edition of the Pharmacopæia of the United States, a work which ought to be in the hands of every practitioner as a guide in the preparation of medicines: and he has endeavoured to arrange the articles in each division, as nearly as he could, in the order of their efficacy as Therapeutical agents."

Therapeutical agents.'

"The subject of Materia Medica has been handled by our author with more than usual judgment. He has, very wisely in our opinion given his principal attention to the articles of the Materia Medica as medicines. In conclusion, we strongly recommend these volumes to our readers. No medical student on either side of the Atlantic ought to be without them "-Forbes' British and Fereign Medicat Review, Jan 1844

This work is extensively used as a Text Book in many Colleges.

# DUNGLISON ON NEW REMEDIES, 4th Edition. NEW REMEDIES; PHARMACEUTICALLY AND THERAPEUTICALLY CONSIDERED.

Fourth Edition, with extensive modifications and Additions,

#### BY ROBLEY DUNGLISON, M. D, &c. &c. &c.

In one volume octavo.

This edition will be found more complete than any of the former. It contains all the new remedies introduced since the third edition in 1841, as well as the new matter embodied in the pharmaceutical journals and essays since that period, as well as in the large works of Pereira. Christison, Bouchardat, Lincke and others. The new matter (about sixty or seventy pages) thus introduced through the work can only be appreciated on close investigation. The principal new articles inserted in this edition are Aluminæ Sales, Anthrakekali, Cannabis Indica, Corylus Rostrata, Ferri et Quiniæ Citras, Fucus Amylaceus, Fuligokali, Gentinant Chirage Period (1998). ana Chirayita, Juglans Regia, Matias, Paullinia, and Platini Preparata.

# A new Edition, to 1844,

# DUNGLISON'S MEDICAL DICTIONARY.

# A DIGTIONARY OF MEDICAL SCIENCE:

A concise Account of the various Subjects and Terms, with the French and other Synonymes, Notices of Climates and of Celebrated Mineral Waters, Formulæ for various Officinal and Empirical Preparations, &c. &c.

# FOURTH EDITION, MODIFIED AND IMPROVED. By ROBLEY DUNGLISON, M.D.

Professor in the Jefferson Medical College, etc.

In one Volume, royal Octavo, of near 800 large Pages, double columns.

This may safely be affirmed to be the most complete Medical Dictionary yet published, containing, as it does, upwards of

FORTY THOUSAND WORDS AND SYNONYMES.

It has been the author's anxious desire to render this work a satisfactory and desirable, if not indispensable, Lexicon, in which the student should never be disappointed in his search for any term with which he might meet in the course of his reading. To those unacquainted with the rapid and unceasing progress of medical science, it might seem that no great modification was necessary in the fourth edition of a work like the present, when the third had appeared within two years, and yet it has been found advisable to add no less than

#### TWO THOUSAND WORDS AND TERMS

not included in the preceding editions. Some of these are words of recent formation, while others had previously escaped the author.

# SPECIAL TREATISES ON MEDICINE, SURGERY, &c.

PROUT ON THE STOMACH.

ON THE NATURE AND TREATMENT OF STOMACH AND RENAL DISEASES; Being an Inquiry into the Connection of Diabetes, Calculus, and other affections of the Kidney and Bladder with Indigestion. By William Prout, M. D., F. R. S., &c. From the Fourth Revised London Edition, with Plates, some coloured.

HOPE ON THE HEART.

A TREATISE ON THE DISEASES OF THE HEART AND GREAT VESSELS, and on the affections which may be mistaken for them. By J. Hope, M. D., F. R. S. From the Third London Edition, with notes by C. W. Pennock, M. D., &c. In one volume, 8vo. with

BARTLETT ON FEVERS.

THE HISTORY, DIAGNOSIS, AND TREATMENT OF TYPHOID AND TYPHUS FEVER; With an Essay on the Diagnosis of Bilious Remittent and of Yellow Fever. By ELISHA BARTLETT, M. D., &c. In one volume 8vo.

WILSON ON THE SKIN.

A PRACTICAL AND THEORETICAL TREATISE ON THE DIAGNOSIS, PATHO. LOGY AND TREATMENT OF THE DISEASES OF THE SKIN. Arranged according to a natural system of classification, and preceded by an outline of the Anatomy and Physiology of the Skin. By Erasmus Wilson, author of "A System of Human Anatomy," &c. In one vol. 8vo.

WALSHE ON THE LUNGS.

THE PHYSICAL DIAGNOSIS OF THE DISEASES OF THE LUNGS. By WALTER HAYLE WALSHE, M. D., Professor of Pathological Anatomy in University College London, &c. In one volume, 12mo.

PATHOLOGICAL HÆMATOLOGY.

AN ESSAY ON THE BLOOD IN DISEASE. By G. Andral, Professor of General Pathology and Therapeutics in the University of Paris, &c. &c., translated from the French by J. F. MEIGS, M.D. and ALFRED STILLE, M.D. A small volume.

HARRIS ON MAXILLARY SINUS.

DISSERTATION ON THE DISEASES OF THE MAXILLARY SINUS. By CHAPIN A. HARRIS, M. D., D. D. S. &c. One vol. 8vo.

RICORD ON VENEREAL.

A PRACTICAL TREATISE ON VENEREAL DISEASES; or, Critical and Experimental Researches on Inoculation, applied to the study of these affections; with a Therapeutical Summary, and special Formulary. From the French of J. Ricord, M. D., &c. In one volume Svo.

ABERCROMBIE ON THE BRAIN.

PATHOLOGICAL AND PRACTICAL RESEARCHES ON THE DISEASES OF THE BRAIN AND SPINAL CORD. By JOHN ABERCROMBIE, M. D., &c. A new Edition, enlarged by the author. One vol. 8vo.

ABERCROMBIE ON THE STOMACH.

PATHOLOGICAL AND PRACTICAL RESEARCHES ON THE DISEASES OF THE STOMACH, the Intestinal Canal, the Liver, and other Viscera of the Abdomen. By JOHN ABERCROMBIE, M. D. Third American Edition. One vol. 8vo.

PHILPIP ON RENDRERGERONS. A TREATISE ON PROTRACTED INDIGESTION; and its consequences, by A. P. W.

PHILIP, M. D., F. R. S., &c. One volume, 8vo.

BELL ON THE TEETH.

THE ANATOMY, PHYSIOLOGY, AND DISEASES OF THE TEETH. By Thomas Bell, F. R. S., F. L. S., F. G. S., Member of the Royal College of Surgeons in London, &c. Third American Edition, with numerous plates.

HARRISON ON THE MERVES.

AN ESSAY TOWARDS A CORRECT THEORY OF THE NERVOUS SYSTEM. By JOHN HARRISON, M. D., Professor of Physiology and Pathology in the University of Louisians. In one volume, 8vo.

## ROBERTSON ON THE TEETH.

A PRACTICAL TREATISE ON THE HUMAN TEETH. Showing the causes of their destruction, and the means of their preservation. By WM. ROBERTSON. With Plates. In one vol.

BIBEZIBLIUS ON THE URINIES &C.

THE KIDNEYS AND URINE. By J. J. Berzelius. Translated from the German by J. H. Boyè, and F. Leaming, M. D. In one volume, 8vo.

#### CHITTY'S MEDICAL JURISPRUDENCE.

A PRACTICAL TREATISE ON MEDICAL JURISPRUDENCE, and all the laws relating to Medical Practitioners; with Explanatory Notes. By J. Chitty, Esq., Barrister at Law. Second American Edition, with notes and additions, adapted to American works and judicial decisions. In one vol. 8vo., with plates and cuts.

#### TRAILL'S MEDICAL JURISPRUDENCE.

OUTLINES OF A COURSE OF LECTURES ON MEDICAL JURISPRUDENCE. By THOS. S. TRAILL, M. D., F. R. S. E., Professor of Jurisp. and Med. Police in Univ. of Edinburgh. Revised, with numerous notes. 1 vol. 8vo.

#### BRODIE ON URINARY ORGANS.

LECTURES ON THE DISEASES OF THE URINARY ORGANS. By SIR BENJAMIN C. BRODIE, Bart., F. R. S. &c. &c. From the third London Ed. In one vol. Svo.

# BRODIE ON THE JOINTS.

PATHOLOGICAL AND SURGICAL OBSERVATIONS ON THE DISEASES OF THE JOINTS. By Sir Benj. C. Brodie, Bart., F. R. S., &c. &c. From the Fourth London Edition. In one vol. 8vo.

#### COATES' POPULAR MEDICINE.

POPULAR MEDICINE; or, FAMILY ADVISER, consisting of Outlines of Anatomy, Phy. siology, and Hygiene, with such hints on the Practice of Physic, Surgery, and the Diseases of Women and Children, as may prove useful in families when regular physicians cannot be procured. By REYNELL COATES, M. D., &c. &c. In one volume, 8vo., with cuts.

SELECT MEDICAL ESSATS.

ESSAYS ON PRACTICAL MEDICINE AND SURGERY, By Drs. Chapman, Bache, Horner, DUNGLISON, DEWEES, CONDIE, HAYS, HODGE, &c. &c. In two volumes, 8vo.

ELLIOTSON ON MESMERISM, Price 25 cents.

NUMEROUS CASES OF SURGICAL OPERATIONS WITHOUT PAIN IN THE MES-MERIC STATE; by John Elliotson, M. D., F. R. S., &c. &c.

#### ARNOTT'S PHYSICS.

ELEMENTS OF PHYSICS; or, Natural Philosophy, general and medical, written for universal use, in plain or non-technical language. By Neil Arnott, M. D., &c. Complete in one vol. 8vo., a new edition, edited by Isaac Hays, M. D., &c.

TRIMMER'S GEOLOGY.
PRACTICAL GEOLOGY AND MINERALOGY; with instructions for Qualitative Analysis of Minerals. By J. TRIMMER, F. G. S. In one volume, 8vo. With over 200 wood-cuts.

GEOLOGY AND MINERALOGY, WITH REFERENCE TO NATURAL THEOLOGY. By the Rev. WM. BUCKLAND, D. D. A new edition, in two volumes, with numerous plates and maps.

BRIDGEWATER TREATISES.
THE BRIDGEWATER TREATISES, On the Power, Wisdom, and Goodness of God, as manifested in the Creation. In seven volumes, 8vo. Containing Essays by Roget, Kirby, Buckland, Babbage, Bell, Prout, Chalmers, &c. &c., and illustrated by numerous plates, cuts, and maps, in seven volumes, 8vo.

THE HISTORY, HABITS AND INSTINCTS OF ANIMALS, by the REV. WILLIAM Kirry, M A., F.R.S. Illustrated by numerous copperplate engravings.

THE NINTH BRIDGEWATER TREATISE. A Fragment, by Charles Babbage, Esq. From the second London edition. In one volume octavo.

YOUATT'S TREATISE ON THE DISEASES OF THE HORSE, &c. &c. in one vol. 8vo.

YOUATT AND CLATER'S CATTLE DOCTOR, &c. in one vol. 12mo.

Several of the Treatises in Tweedie's Library, first edition may be had in separate volumes.

# YOUATT AND CLATER'S CATTLE-DOCTOR.

Now Ready.

# EVERY MAN HIS OWN CATTLE DOCTOR:

THE CAUSES, SYMPTOMS, AND TREATMENT OF ALL THE DISEASES INCIDENT TO

OXEN, SHEEP, AND SWINE.

By FRANCIS CLATER.

EDITED, REVISED, AND ALMOST REWRITTEN,

By WILLIAM YOUATT, AUTHOR Of "THE HORSE," &c. Together with numerous Additions, by the American Editor, J. S. Skinner. AMONG WHICH ARE

AN ESSAY ON THE USE OF OXEN, WITH MODES OF BREAKING, FEEDING, GRAZING, ETC. AND A TREATISE ON THE GROWTH, IMPROVEMENT AND BREEDING OF SHEEP, AND THE SOILS ADAPTED TO THEIR RAISING.

> With numerous Cuts and Illustrations. In one volume, 12mo. Price Fifty Cents, in Cloth.

> > ALSO,

THE COMPLETE COOK, edited by Sanderson, price 25 Cents.

THE COMPLETE CONFECTIONER, edited by Parkinson, price 25 Cents.

THE COMPLETE FLORIST, edited by an Experienced Florist of Philadelphia, price 25 Cents.

Persons remitting in current funds, free of postage, ONE DOLLAR, can have the Four Works, by mail, or otherwise, as they may direct, each one sewed in a cover.

THE HORSE.

By WILLIAM YOUATT, a new edition with illustrations; containing a full account of the Diseases of the Horse, with their mode of treatment; his anatomy, and the usual operations performed on him; his breeding, breaking, and management; and hints on his soundness, and the purchase and sale. Together with a General History of the Horse; a dissertation on the American Trotting Horse, how trained and jockeyed, an account of his remarkable performances, and an Essay on the Ass and the Mule, by J. S. SKINNER, Assistant Postmaster General, and Editor of the Turf Register. In one volume, octavo.

ENCYCLOPÆDIA AMERICANA.

A new edition. A Popular Dictionary of Arts, Sciences, Literature, History, Politics and Biography, including a copious collection of original articles in American Biography. On the basis of the seventh edition of the German Conversations-Lexikon. Edited by Francis Lieber, assisted by E. Wigglesworth and T. G. Bradford. In 13 volumes octavo of about 600 pages each.

The publishers offer this valuable work at a very low price in all the various styles of binding.

Over Two Hundred Thousand copies have been sold in Germany.

## THE ENCYCLOPÆDIA OF GEOGRAPHY.

Comprising a complete description of the earth, Physical, Statistical, Commercial and Political; exhibiting its relation to the heavenly bodies—its physical structure—the natural history of each country; and the Industry, Commerce, Political Institutions, and Civil and Social State of all Nations; by Hugh Murray, F.R.S.E., assisted in Astronomy, &c. by Prof. Wallace, Geology, &c. by Prof. Jameson, Botany, &c. by Prof. Hooker, Zoology, &c. by W. Swainson. Illustrated by eighty-two maps, and about eleven hundred other engravings on wood, representing the most remarkable objects of Nature and Art in every region of the Globe; together with a map of the United States, drawn by Drayton, from Tanner's Map, and engraved on copper, in which is embodied the latest information relating to the Internal Improvements of this country. The portion relating to the United States written anew, by T. G. Bradford. Revised, corrected, and brought down to the present period 1844. In three handsome octavo volumes in various bindings.

MISS STRICKLAND'S QUEENS OF ENGLAND.

Lives of the Queens of England, from the Norman Conquest to the reign of Elizabeth; with anecdotes of their Courts, now first published from official records and other authentic documents, private as well as public. A new and improved edition, in six volumes, royal 12mo., bound in cloth, or done up in paper to match.

- SELECT WORKS OF TOBIAS SMOLLETT, with a memoir of his Life and Writings, by Sir Walter Scott. Containing The Adventures of Roderick Random; Price 25 cents.—The Adventures of Peregrine Pickle, Double Number; Price 50 cents.—The Expedition of Humphrey Clinker; Price 25 cents.—The Adventures of Ferdinand Count Fathom; Price 25 cents.—The Adventures of Sir Launcelot Greaves; Price 25 cents;—The History and Adventures of an Atom, and Select Poems, in one part; Price 25 cents;—or the whole done up in one volume.
- SELECT WORKS OF HENRY FIELDING, with a memoir of his Life and Writings, by Sir Walter Scott, and an Essay on his Life and Genius, by Arthur Murphy, Esq., containing Tom Jones, or The History of a Foundling; Double Number; Price 50 cents. The Adventures of Joseph Andrews and his friend Mr. Abraham Adams, Price 25 cents. Amelia; Price 25 cents. The Life of Jonathan Wild, with the Life of Fielding, Essay on his Genius, &c., in one part Price 25 cents;—or the whole done up in one volume.
- COOPER'S NOVELS, AT TWENTY-FIVE CENTS PER VOLUME, embracing Wyandottè, Wing-and-Wing, The Spy, The Pilot, The Water-Witch, Lionel Lincoln, Heidenmauer, The Pathfinder, Precaution, The Wish-Ton-Wish, Homeward Bound, Mercedes of Castile, The Pioneers, The Red Rover, The Two Admirals, The Prairie, The Headsman, The Deerslayer, The Bravo, The Last of the Mohicans, Home as Found, The Monikins, The Travelling Bachelor, and Ned Myers; In all twenty-four different Works, or forty-seven volumes.
- WHITE'S UNIVERSAL HISTORY FOR COLLEGES AND HIGHER SCHOOLS. Elements of Universal History, on a new and systematic plan; from the earliest times to the Treaty of Vienna. To which is added, a summary of the leading events since that period, by H. White, B.A., Trinity College, Cambridge, with additions, by John S. Hart, A.M., Principal of the Philadelphia High School, and Professor of Moral and Mental Science, Member of the American Philosophical Society, &c. in one vol. 12mo., or an Edition without Questions may be had for general use.
- POEMS BY SAMUEL ROGERS, with numerous illustrations, a new edition, revised, with additions by the author, in one volume, imperial octavo, embellished with ten engravings by Turner, Goodall, Wallis, Finden, &c. in cloth extra, or beautiful white calf, gilt edges.
- WALPOLE'S SUPPRESSED LETTERS TO SIR HORACE MANN, from 1760 to 1785, now first published from the original MSS. In two octavo volumes to match the four volumes formerly published, to which they form a continuation.
- MARTIN CHUZZLEWIT, by "Boz," publishing in Nos. with plates. Price 61 cents each.
- THE HISTORY OF IRELAND, commencing with its earliest period to the great expedition against Scotland in 1545. By Thomas Moore. In one volume octavo.
- BROUGHAM'S FRENCH REVOLUTION, price fifty cents, forming a third volume to his Memoirs of Statesmen.
- THE EDUCATION OF MOTHERS; or, the Civilization of Mankind by Woman. By L. Aime Martin. Being the work to which the prize of the French Academy was awarded. 'Translated by Edwin Lee, Esq. Revised from the fourth French edition, done up to match the "Ladies' Cabinet Series." Price seventy-five cents.
- BENTHAMIANA; or, Select Extracts from the Works of Jeremy Bentham. With an outline of his opinions on the principal subjects discussed in his works. Edited by John Hill Burton, Advocate. In one volume, large 12mo., extra cloth.
- THE HISTORY OF ROME, by B. G. Niebuhr. Translated by J. C. Hare, Connop Thirlwall, William Smith, and Leonhard Schmitz. From the third English Edition, revised. The four volumes brought into two large octavo volumes. (Now at press.)
- PICCIOLA, THE PRISONER OF FENESTRELLA; or, Captivity Captive. From the French of M. de Saintine. Third American edition. In one volume, 12mo., price thirty-five cents, in glazed paper.
- THE MILLWRIGHT AND MILLER'S GUIDE. By OLIVER EVANS. The Tenth Edition with Additions and Corrections, by the Professor of Mechanics in the Franklin Institute of Pennsylvania; and a description of an improved Merchant Flour Mill. With Engravings. By C. & O. Evans, Engineers. This is a practical work, and has had a very extended sale.
- AMERICAN ORNITHOLOGY, a supplement to "Wilson," by Prince CHARLES BONA-PARTE. Four volumes, folio, with magnificent coloured plates.
- THE NATIONAL SCHOOL MANUAL. In four volumes, 12mo, with an Atlas and the Clinton Primer as an Introduction.
- A POPULAR TREATISE ON AGRICULTURAL CHEMISTRY. By CHARLES SQUAREY, in one volume 12mo.
- THE DISTRICT SCHOOL; OR, NATIONAL EDUCATION. By J. ORVILLE TAYLOR, Third Edition, in one volume 12mo.
- LIGHTS, SHADOWS, AND REFLECTIONS OF WHIGS AND TORIES. By a Country Gentleman, in one volume 12mo.

## Price Twenty-five cents each.

THE COMPLETE COOK .- Plain and Practical Directions for Cooking and Housekeeping; with upwards of seven hundred receipts: consisting of directions for the choice of meat and poultry; preparations for Cooking, making of broths and soups; boiling, roasting, baking, and frying, of meats, fish, &c. seasonings, colourings, cooking vegetables; preparing salads, clarifying; making of pastry, puddings, gruels, gravies, garnishes, &c. and, with general directions for making wines. With additions and alterations, by J. M. Sanderson, of the Franklin House.

THE COMPLETE CONFECTIONER, PASTRY COOK AND BAKER.—Plain and practical directions for making Confectionary and Pastry, and for baking; with upwards of five hundred receipts; consisting of directions for making all sorts of preserves, sugarboiling, comfits, lozenges, ornamental cakes, ices, liqueurs, waters, gum-paste ornaments, syrups, jellies, marmalades, compotes, bread-baking, artificial yeasts, fancy biscuits, cakes, rolls, muffins, tarts, pies, &c. &c. With additions and alterations, by Parkinson, Practical Confectioner, Chestnut street.

THE COMPLETE FLORIST .- A Manual of Gardening, containing practical instruction for the management of greenhouse plants, and for the cultivation of the shrubbery, the flower garden, and the lawn, with descriptions of those plants and trees most worthy of culture, in each department. With additions and amendments, adapted to the climate of the United States.

Now Preparing to match the Complete Florist, THE KITCHEN AND FRUIT GARDINER.

#### THE LADY'S CABINET SERIES.

L. &. B. are publishing under this title a number of standard works, fitted for the Boudoir, exclusively by female writers, such as Mrs. Jamieson, Mrs. Hemans, Miss Strickland, &c. They are all neatly done up to match in lemon-coloured glazed paper, and will form a beautiful series.

The Poetical Remains of Lucretia M. Davidson, with a Memoir by Miss C. M. Sedgwick. In one volume, price fifty cents.

The Poetical Remains of Margaret M. Davidson, with a Biography by Washington Irving. In one volume, price fifty cents.

The Lives of the Queens of England, by Miss Strickland, first series. In three volumes, price fifty cents each. Second series. In three volumes, price sixty-five cents each.

The Study of the Life of Woman. Translated from the French of Madame Necker de Saussure

of Geneva. In one volume, price seventy-five cents.

The Poetical Works of Mrs. Hemans, to which is prefixed a Memoir by her sister Mrs. Hughes, and an essay on her genius by Mrs. Sigourney. To be completed in seven volumes, price fifty cents each.

The Memoirs of the Loves of the Poets, or the lives of women celebrated in ancient and modern

poetry, by Mrs. Jamieson. In one volume, price seventy-five cents.
Selections from the Writings of Mrs. Margaret M. Davidson, the mother of Lucretia Maria and Margaret M. Davidson, with a preface by Miss C. M. Sedgwick, price fifty cents. Other volumes are in preparation.

## L. & B. are now Publishing

# A CABINET SERIES OF NOVELS IN A FORM FOR PRESERVATION.

It will embody-Destiny; or the Chief's Daughter. By Miss Ferrier, author of the Inheritance, Marriage, &c. In one thick volume, 12mo. price 40 cents.—Now ready.

Marriage. By Miss Ferrier, (at press.)

The Inheritance, by Miss Ferrier, (preparing.)
Other Novels of high character are to be embraced in this series.

# JUST READY.

WHIMS AND ODDITIES. By Thomas Hood. With upwards of eighty spirited wood cuts, In

one volume, 12mo. Price fifty cents.

FAIRY LEGENDS OF THE SOUTH OF IRELAND. By T. Crofton Croker. With fifty-five wood cuts. Second Edition. In one vol. 12mo., price fifty cents.

WALPOLE'S New Letters to Horace Mann, in 2 vols. 8vo., vol. 1 now ready.

THE HUNCHBACK OF NOTRE DAME. By Victor Hugo. With an Illustration. Price 25

TYLNEY HALL. By Thomas Hood. With an Illustration. Price 25 cents. CLATER & YOUATT'S CATTLE DOCTOR, in 1 vol., 12mo. WHITE'S UNIVERSAL HISTORY, a handsome vol. in 12mo.

#### MISCELLANEOUS BOOKS

#### PUBLISHED BY

LEA & BLANCHARD. Brougham, Henry Lord, His Speeches with Historical Introductions, in 2 vols. 8vo.

Do. do. do. Historical Sketches of the Times of George IV., in 2 vols. royal 12mo.

Do. do. do. Critical and Miscellaneous Writings, in 2 vols. royal 12mo.

Butler's Atlas of Ancient Geography, with a complete Index, 1 vol. 8vo.

Do. Geographia Classica, fourth edition, 1 vol. 8vo. to accompany the Atlas. Boz's Works complete, fine edition, with numerous illustrations, 6 vols. royal 8vo. Consisting of Sketches of Every day Life and Every-day People.

Posthumous Papers of the Pickwick Club.
Oliver Twist; or, The Parish Boy's Progress.
Life and Adventures of Nicholas Nickleby.
Old Curiosity Shop.
Barnaby Rudge.
Also, a Cheap Edition, without the plates.
Bulwer's Critical and Miscellaneous Writings, in 2 vols. royal 12mo.
Brewster's Treatise on Optics, a new and improved edition, 1 vol. 12mo.
Bolmar's French Series, Consisting of Perrin's Fables, 1 vol. 12mo. Colloquial Phrases, 1 vol. 18mo.
French Verbs, 1 vol. 12mo. Telemaque, 1 vol. 12mo. Key to do. 1 vol. 12mo.
Cooper's Naval History of the United States, second edition, in 2 vols. 8vo.
Do. Novels and Tales complete, and to match, 44 vols. 12mo. at 25 cts. per volume.
Do. Leather Stocking Tales, 5 vols. 12mo.
Do. Sea Tales, 5 vols. 12mo.
Davidson's, Margaret M. Biography and Poetical Remains, edited by Irving, 1 vol. 12mo. ex. cl.
Davidson's, Lucretia M. Biography and Poetical Remains, edited by Miss Sedgwick. A new and cheap edition, 1 vol. 12mo. price 50 cts.
Ellis, Mrs. Family Secrets; or, Hints to those who would make Home Happy, in 2 vols. 12mo. price Sketches of Every day Life and Every-day People. 50 cts. Encyclopædia of Geography. A new edition, brought up to 1842, publishing in 24 weekly Numbers, royal 8vo. Encyclopædia Americana, a great work, in 13 vols. 8vo. Frederic the Great, His Court and Times, including the Seven Years War, by Thomas Campbell, in 4 vols. 12mo. Fielding's Select Works, with his Life, by A. Murphy, royal 8vo.
Geisler's Text-Book of Ecclesiastical History, 3 vols. 8vo.
Guy Fawkes; An Historical Romance of the Times of James I., by W. H. Ainsworth, 1 vol. royal 8vo. Heber's Poetical Works. Complete in 1 vol. 12mo.

Herschell's Astronomy. A new and improved edition, 1 vol. 12mo.

History of Congress during the Administration of General Washington, in 1 vol. 8vo. a cheap volume. Hemans' Poetical Works, Complete, in 7 vols. 12mo. with a Life by her Sister, and an Essay on her Genius, by Mrs. Sigourney.

Do. Memoirs, by her Sister, Mrs. Hughes, in 1 vol. 12mo.

Horse, (The) by Youatt, with additions by Skinner, 1 vol. 8vo. with many illustrations.

Irving's Select Works, in 2 vols. royal 8vo. with a portrait.

Do. Astoria, in 2 vols. 8vo. Rocky Mountains. A new and cheap edition.

Life and Voyages of Christopher Columbus, in 2 vols. 8vo.

Crayon Miscellany—Containing Abbotsford, Newstead Abbey, Conquest of Spain, &c. Do. Do. Do. vols. 12mo. Jess', Court of England under the House of Nassau and Hanover, 3 vols royal 12mo. Jefferson's Life, by Professor Tucker, in 2 vols. 8vo. Keble's Christian Year, third edition, in one neat 12mo. volume. Keble's Child's Christian Year, third edition, in one neat 12mo. volume.

Keble's Child's Christian Year, a small volume.

Lockhart's Life of Sir Walter Scott, in 7 vols. 12mo. with a portrait.

Language of Flowers, with beautiful coloured plates, a new edition, extra gilt edges.

Madame de Sevigne and her most distinguished Cotemporaries, in 2 vols. royal 12mo.

Moore's, Thomas, History of Ireland, in 1 vol. 8vo.

Millwright's and Miller's Guide, with plates, by Oliver Evans, tenth edition, with improvements, 1 vol. 8vo. 1 vol. 8vo. Madden's Lives and Times of the United Irishmen, 2 vols. 12mo. Mrs. Marcet's Stories for Young Children, small 4to. Porcelain Tower, or Nine Stories of China, by T. T. T. 1 vol. 12mo. with numerous illustrations. Queen of Plowers, or Memoirs of the Rose, 1 vol. coloured plates. Romantic Biography of the Age of Elizabeth, by the Benedictine Brethren of Glendalough, 2 vols. 12mo. Ranke's Ecclesiastical and Political History of the Popes of Rome, during the 16th and 17th centuries, in 2 vols. royal 8vo.

Strickland, Miss, Lives of the Queens of England, 1st series, 3 vols. 12mo. extra cloth.

Do. do. do. do. 2d do. 2 vols. do. do. Do. do. do. do. do. 2d do. 2 vols. do. Stanley Thorn, by Cockton, 1 vol. royal 8vo. with numerous illustrations. Shelley, Mrs. Eminent Literary and Scientific Men of Italy, in 2 vols. 12mo Do. do. Memoirs of the most eminent French Writers, 2 vols. 12mo.

Sentiments of Flowers, or Language of Flora, with coloured plates. Scott's Poetical Works. Complete in 6 vols. royal 12mo. extra cloth. Smollett's Select Works, with a Memoir by Sir Walter Scott, royal 8vo.

Washington Potts, Mrs and Mr. Smith, by Miss Leslie, price 25 cts.
Waleh's Sketches of Conspicuous Living Characters of France, 1 vol. 12mo.
Walpole's Unrivalled Letters. The only complete edition. Containing nearly 300 Letters, with a

portrait, in 4 vols. 8vo. extra cloth.

# THE AMERICAN JOURNAL

OF THE

# MEDICAL SCIENCES

LEA & BLANCHARD are the Publishers of the American Journal of the Medical Sciences, edited by ISAAC HAYS, M.D., Surgeon to Wills Hospital, Physician to the Philadelphia Orphan Asylum, Member of the Am. Philos Soc, &c. &c. &c. assisted by numerous collaborators in every section of the Union. This Journal was commenced TWENTY THREE YEARS AGO, and it is therefore the oldest Medical Journal now existing in the United States, and its permanency may be considered as established. It was originally edited by Dr. Chapman, who has been a constant contributor to its pages; and for the last seventeen years it has been under the editorial direction of Dr. Isaac Hays.

A New Series was commenced in Jan. 1841.

A New Series was commenced in Jan. 1841.

The pages of this Journal contain the records of the experience of the most distinguished members of the Profession in every part of the Union.

#### CONTENTS OF THE NO. FOR APRIL 1844.

MEMORS AND CASES.—Art. I Observations on the Pathological Relations of the Medulla Spinalis. By Austin Flint, M.D. II. Case of Molluscum, associated with Fibro-Cellular Encysted Tumour and Encephaloid Disease. By Washington L. Atlee, M.D. [With a wood cut.] III. On the Pulse of the Insane. By Pliny Earle, M.D. IV. Statistics and Cases of Midwifery; compiled from the Records of the Philadelphia Hospital, Blockley. By Geo. N. Burwell, M.D. V. On the Congestive Fever of Mississippi, with Cases. By R. G. Wharton, M.D. VI. Enteritis—with Cases, exemplifying the decidedly beneficial effects of Blood letting, and the Sedative Treatment. By P. M. Kollock, M.D. VII. Case of Inversion of the Uterus, in which Reposition was effected on the tenth day. By Joseph P. Gazzam, M.D. VIII. Ligature of the external Iliac Artery for Aneurism. By W. P. Boling, M.D. IX Eupatorium Perfoliatum in Epidemic Influenza. By J. F. Peebles, M.D. X. Case of Polypus of the Uterus expelled by the action of Ergot. By Thomas J. Garden, M.D. XI. On the Extraction of Retained Placenta in Abortion. By Henry Bond, M.D. XII. Improved Catheter Bougie. By R. J. Dodd, M.D. Reviews.—XIII. Fenger's Dissertation on Erysipelas Ambulans. XIV. Drs. M William and Pritchett on Catheter Bougie. By R. J. Dodd, M D.

REVIEWS.—XIII. Fenger's Dissertation on Erysipelas Ambulans. XIV. Drs. M William and Pritchett on

African Remittent Fevers.

African Remittent Fevers.

BIBLIOGRAPHICAL NOTICES.—XV. Harrison's Theory of the Nervous System. XVI. Hartshorne's Annual Report of the Eastern State Penitentiary. XVII. Kirkbride's Report of the Pennsylvania Hospital for the Insane. 2 Report of the Ohio Lunatic Asylum. 3. Report of the Managers of the State Lunatic Asylum, (New York) 4. Report of the Superintendent of the Maine Hospital. XVIII Paris's Pharmacologia. XIX. Hare's Statistical Report of one hundred and ninety cases of Insanity admitted into the Retreat near Leeds. XX. White's Address on Insanity. XXI. Report of the Surgeon General U. S. Army. XXII. Dunglison's Human Physiology. XXVIII. Dunglison's Human Physiology. XXVIII. Dunglison's Heman Physiology. XXVII. Weston some of the more important Diseases of Childhood. XXV. Magendie's Elementary Treatise on Human Physiology. XXVII. West's Clinical and Pathological Report on the Pneumonia of Children. XXVIII. Cooper on Hernia. XXVII. West's Clinical and Pathological Report on the Pneumonia of Children. XXVIII. Cooper on Hernia. XXIX. Todd on Rheumatic Gout. Fever, and Chronic Rheumatism of the Joints. XXX. Smith and Horner's Anatomical Atlas. XXXI. Quarterly Summary of the Transactions of the College of Physicians of Philadelphia. XXXII. Chapman's Lectures on the more important Diseases of the Thoracic and Abdominal Viscera.

#### SUMMARY OF THE IMPROVEMENTS AND DISCOVERIES IN THE MEDICAL SCIENCES.

Anatomy and Physiology.—1. Elsæsser on the Period at which the foramen ovale, and the ductus venosus become obliterated. 2. Lacauchie on Absorption. 3. Burrows on Blood in the brain. 4. Erdl on Optic nerves. 5. Spence on Nervus vagus and nervus accessorius. 6. Escape of ova independent of fecundation, and the connection of this with menstruation. 7. Roberton on Age of puberty in girls. 8. Starr on Urachus pervious after birth.

MATERIA MEDICA AND PHARMACY.—9. Galleoti on Species of Veratrum. 10. Martens and Galleoti on Species of Smilax. 11. Formula for fluid extract of Senna. 12. O'Shaughnessy on Gurjum Balsam. 13. King-

don on New preparation of Quinine.

MEDICAL PATHOLOGY AND THERAPBUTICS AND PRACTICAL MEDICINE.—14. Prus on Meningeal Apoplexy.

15. Huss on Communication between the aorta and pulmonary artery without Cyanosis. 16. Huss's remarkable case of Cyanosis. 17. Pritchard on the connection of Insanity with diseases in the organs of physical life. 18 Fioravente on new treatment of Sciatica. 19. Volz on Ulceration of the Appendix Vermiformis. 20. Bredscheider on Rupture of the Trachea. 21. Negrier on Epistaxis. 22. Herberger on blood and urine in chlorotic subjects. 23. Black, on Tubercles and Phthisis. 24. Boudin on Antagonism of typhoid and intermittent Fevers. 25. Taylor on double intussusception. 26. Mackenzie on Epidemic Remittent Fever which provailed in Classon.

which prevailed in Glasgow.

Surgical Pathology and Therapeutics and Operative Surgery.—27. Debency on abortive treatment of Gonorthea, by the Nitrate of Silver, and on the employment of caustic injections at all stages of Urethritis. 28. Leroy d'Etiolles on Cancerous diatheses. 29. Douglas on Dislocation of the Femur—head of that bone on the pubes,—fracture of the neck of femur. 30. Tuton on Ovarian dropsy successfully treated with ioduret of Iron. 31. Mackesy on Amputation of the Leg—ressels so diseased as to preclude the use of ligatures—recovery. 32. Syme on Disarticulation of the ramus of the lower Jaw, without opening the cavity of the mouth. 33. Lepine on Penetrating wound of the Abdomen—Prolapse of the Stomach—Recovery. 34. Fife on Ununited Fracture of the Humerus—excision of the fractured extremities—cure. 35. Purefoy on Trismus following the extraction of a tooth. 36. Cooper on Extirpation of an Ovarian Cyst. 37. Scontetten on Tracheotomy in the last stage of Croup. 38. O'Bryen on Diagnosis of Aneurism of the Aorta. 39. Englehardt on Fibrous tumour mistaken for Aneurism. 40. Rufz on Painful affection of the breast. 41. Danglehardt on Fibrous tumour mistaken for Aneurism. 40. Rufz on Painful affection of the breast. 41. Danglehardt on Suture of the Perineum performed immediately after delivery. 42. Berard on Cure of encysted Tumours. 43. Segalas's remarkable case of Calculus. 44. Lenoir on Dislocation of the os innominatum on the sacrum. 45. Gigon on Polypi of the rectum in young children.

Ophthalmology—46. Mackensie on Postfebrile Ophthalmitis. 47. Lavergne on Amaurosis from concussion of the retina. 48. Ansiaux on Foreign bodies in the Eye. 49. Guepin on Foreign body in the Eye. 50. sion of the retina. 48. Ansiaux on Foreign bodies in the Eye. 49. Guepin on Foreign body in the Eye. 50. sion of the retina. 48. Ansiaux on Foreign bodies in the Eye. 49. Guepin on Foreign body in the condition of the exercises on the excretory ducts of the lachrymal gland. 52. Dornbluth on Cyst in the orbital cavity.

bital cavity.

Midwipery.—53. Chailly Honoré on Artificial rupture of the membranes to accelerate delivery. 54. Elkington on Contagiousness of Puerperal Fever. 55. Guillon's New method of compressing the aorta in utering the morrhage. 56. Wilson on Extra-uterine Foxation. 57. Sherman on Perforation of the bladder by mistake during parturition. 58. Mondini on Absence of the uterus. 59. Kayser's Cosarean operation. 60. take during parturition. 58. Mondini on Absence of the uterus. 59. Kayser's Cosarean operation. 60. Vannoni on Kiestine. 61. Simpson on Uterine Sound.

Vannoni on Kiestine. 61. Simpson on Uterine Sound.

Medical Jurisprudence and Toxicology.—62. Perforation of the intestine by Arsenic. 63. Mialke on Medical Jurisprudence and Toxicology.—62. Perforation of the intestine by Arsenic. 65. Vilation of hydrocyanic Acid and the alkaline cyanurets on Calomel. 64. Professional confidence. 65. Vilaume on Malconformation of the Female Sexual Organs. 66. Sensibility and power of locomotion after laume on Malconformation of the Female Sexual Organs. 66. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who had severe injuries. 67. Death from Calomel and Blue Pill. 68. Vandersavel on Pregnancy in a Female who

31

AMERICAN INTELLIGENCE.—Original Communications.—McClellan on Extirpation of the Parotid Gland.

E. Bissell's Cases of Ovarian Disease, with remarks. Eve on Lithotomy—Bilateral operation, with cases.

Tracy on the Vegetable Acids as correctives of acidity of stomach. Ludlow's Notice of a new Flexible Stethoscope: Dodd's Improved Cupping Apparatus. Crowell's Unusual case of irregularity of the incisors and canine teeth of the superior jaw, remedied in seventy-six days.

Domestic Summary.—Judkins' case of Ovarian Disease. Bedford on Vaginal Hysterotomy.

New Medical Works in preparation.

New Medical Works in preparation.

Each Number contains

ORIGINAL ESSAYS.

BIBLIOGRAPHICAL NOTICES,

And a full Summary of Foreign and Domestic Intelligence.

The publishers invite especial attention to this last department, which constitutes an important feature in this Journal, comprising the fullest

RETROSPECT

of the improvements and discoveries made in the Medical Sciences anywhere to be met with, derived not only from the

BRITISH JOURNALS.

but also from those of

FRANCE, GERMANY, DENMARK, ITALY, THE EAST INDIES, and our own country.

ILLUSTRATIONS ARE GIVEN WITH COLOURED PLATES. WOOD CUTS, &c. &c.

when cases require them, and every effort will be made to improve the appearance of the work that it may continue to command the confidence and patronage of the profession.

TERMS.

This Journal is published quarterly on the first of January, April, July and October, each number containing at least 264 large and closely printed pages. Price FIVE DOLLARS a year, which if forwarded in advance and free of postage, will entitle the subscriber to "The Medical News" without charge.

Early application should be made direct to the publishers. The subscription commences with a

January number.

THE MEDICAL NEWS AND LIBRARY

FOR ONE DOLLAR A YEAR, PAYABLE IN ADVANCE.

LEA & BLANCHARD, Philadelphia, are publishing The MEDICAL NEWS AND LIBRARY, a monthly periodical, of 32 or more large and closely printed pages. It contains in addition to the Miscellaneous Medical Intelligence, reports of Clinical Lectures, accounts of the difference of the diff ferent Medical Schools and Hospitals, with notices of cases and operations in those institutions,

and various other matters of interest to practitioners.

The Library is to comprise a series of Lectures or works on the principal branches of Medical Science, and will constitute a complete Library for the student, and a useful work of reference to

the practitioner.

The price of the News and Library to subscribers is ONLY ONE DOLLAR A YEAR, and it will be sent gratuitously to all subscribers to the Medical Journal who pay their subscription five dollars in advance.

All remittances and orders must be sent free of postage, which can be done by the postmaster

franking the letter.

Any Physician, News Agent or Postmaster, can have a number of the News sent gratis, as a specimen, on application, free of postage.

MOTICE.

With a view of extending the circulation of the American Journal of the Medical Sciences, the publishers offer the following inducement to NEW SUBSCRIBERS. Those who remit TEN DOLLARS shall receive The American Journal of Medical Sciences and Medical News for the Two Years, 1844 and 1845, and in addition The Medical News and Library for the year 1843.

They will thus obtain the first 264 pages of Watson's Practice, which are embraced in the News for 1843, and will have that work completed in the News for 1844. (A new work will commence

the year 1845.

The price of Watson's Lectures on the Practice of Physic, complete, will be itself Three Dol-


lars, and will form a volume of near 900 large pages.

This liberal offer is made solely with a view of extending the circulation of the Journal, and to induce payments in advance, and will not be binding with the publishers after July next, or after the copies of the News and Library for 1843 now on hand are exhausted. To secure the advantages of this offer, remittances should be made early.

As the News for 1844 will embrace about 600 pages of Watson's Lectures and 96 of other matters, subscribers are informed that the amount of Five Dollars for that and the Journal should also be remitted early this year-as the publishers reserve to themselves the right to decline furnishing them under Six Dollars, or the News separate for that year for Two Dollars unless so taken and

paid for.

This paper may be delivered to any physician if declined by the person to whom it is addressed or if they have removed-and Postmasters will particularly oblige the publishers by furnishing a list of the Physicians of their county or neighbourhood In addition to the business it may bring to the office, a copy of "Ned Myers" a Tale by Cooper will be sent gratis for any ten or more names furnished free of cost.



NATIONAL LIBRARY OF MEDICINE

NLM 03206746 0