An essay on the effects of lead: comprising a few experiments on the saccharum saturni, and its application in the cure of diseases / by Thomas Semmes.

Contributors

Semmes, Thomas, 1778-1833.
Barton, Benjamin Smith, 1766-1815.
University of Pennsylvania.
Carr & Smith
National Library of Medicine (U.S.)

Publication/Creation

Philadelphia: Printed Carr and Smith, 1801.

Persistent URL

https://wellcomecollection.org/works/qjbw9ccd

License and attribution


This material has been provided by This material has been provided by the National Library of Medicine (U.S.), through the Medical Heritage Library. The original may be consulted at the National Library of Medicine (U.S.) where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.


You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



UNITED STATES OF AMERICA



FOUNDED 1836

WASHINGTON, D.C.

16-67244-1





ESSAY

ON

THE EFFECTS OF LEAD,

COMPRISING

A FEW EXPERIMENTS

ON

THE SACCHARUM SATURNI,

AND ITS

APPLICATION

In the Cure of Diseases.

BY THOMAS SEMMES.

OF MARYLAND;

Honorary Member of the Philadelphia Medical and Chemical Societies.

"Poisons in small doses are the best medicines; and the best medicines in too large doses are poisonous."

Withering's Botanical Arrangement.

PHILADELPHIA:

PRINTED BY CARR AND SMITH.

1801.

AN

INAUGURAL ESSAY,

FOR

THE DEGREE

OF

DOCTOR OF MEDICINE;

SUBMITTED TO THE EXAMINATION

OF

THE REV. JOHN EWING, S. S. T. P. PROVOST,

THE

TRUSTEES AND MEDICAL FACULTY

OF

The University of Pennsylvania,

ON THE

EIGHTH DAY OF JUNE, 1801.

406386

DOCTOR ELISHA CULLEN DICK.

TO you, in whom are concentered, talents, profeffional eminence, and domestic virtues; under whose patronage and direction, my medical education was commenced and finished, with great pleasure and improvement to myfelf; I beg leave to inscribe the following pages.

Although the connection of preceptor and pupil will now cease between us, permit me, dear sir, to hope for the continuance of those prudent counsels, and valuable instructions, with which, you have been pleased to savor me.

For the many polite and flattering attentions I have received from you, and your amiable family, accept the grateful acknowledgements of

Your fincere Friend,

And affectionate Pupil,

THOMAS SEMMES.

The second of th

BENJAMIN SMITH BARTON, M. D.

Professor of Materia Medica, Natural History and Botany,

IN THE UNIVERSITY OF PENNSYLVANIA;

SIR,

WITH great pleafure, I avail myfelf of the prefent opportunity, to express my thanks for the many particular marks of politeness and friendship, with which you have honored me, during my residence in Philadelphia.

That your happiness, and fuccess in practice, may be commensurate with your talents and eminence in public life, is the wish of

Your fincere Friend,

And obedient Servant,

THE AUTHOR.

INTRODUCTION.

WHEN we contemplate the mischiefs, which have arisen from the application of different preparations of lead to the human body, the objections of those medical writers, who have viewed them in a light so unfavorable, and have so strenuously opposed their internal exhibition, would appear to be just.

That lead has produced many difagreeable confequences will be readily granted; but to suppose that the preparations of this metal are invariably followed by deleterious effects, when admitted into the body, if exhibited within bounds, would, in my opinion, be equally absurd and erroneous. If we take a view of the materia medica, we shall behold, enveloped in poison, many of our most active and useful medicines. The datura stramo-

nium, the conium maculatum, and innumerable metallic fubstances in the hands of the ignorant, or when exhibited in improper quantities, disorganize the vital functions and extinguish life. But if these substances are managed judiciously and administered with discretion, they cure disease, prolong life, and prove highly advantageous to mankind.

The effects of fubflances, ufually called poisons, are relative; and the flightest observation will convince us, they act either as medicines or as poifons, according to their quantity and quality. Animals have a greater or less aptitude to be acted on, by one substance than by another; or possess a peculiar idiofyncrafy of habit, whereby the most baneful effects will be produced by a certain fubstance upon one, while upon another, the fame fubstance would be attended with little or no effect. Hemlock is innoxious to cows; hogs fatten on henbane; and goats may confume large quantities of euphorbium without injury; but each of these vegetables are poifonous to many other animals. The stramonium constitutes an article of food to some animals, while the admission of a few grains into the stomach of a man, proves deleterious. -Opium is a valuable medicine, yet it is known, that two perfons taking equal quantities, may be differently affected. Perfons who labour in quickfilver mines feldom remain free from palfy, or fome other complaint, longer than two, three, or four years; the effects of this mineral are not lefs striking than those which sometimes arise from the effluvia of lead. Gilders, and others who fuffer from mercury, are much afflicted with tremors and debility, particularly in their hands; and when the difease is once confirmed, they have no command over the mufcles, which naturally obey the will.* It alfo, under certain states of the system, disposes wounds of the mildest nature to ulceration, and not unfrequently occasions the true phagedena, a diffreffing and often fatal ulcer. A child, whose fystem was under the influence of this mineral, was bled, the orifice, instead of healing, became inflamed; the wound in three days had confiderably enlarged, destroying the integuments, and leaving bare the blood-veffels; on the fourth day death enfued.+

^{*} Vide Clutterbuck, on the poison of lead and copper.

[†] Professor Barton's M. S. Lectures.

The oxigenated muriate of mercury, in the proportion of an eighth or fixth of a grain, and arfenic in small doses, are innocent and salutary remedies, in certain diseases; but when taken in too large quantities, are sollowed by the most fatal consequences.

The different fystems of the same body are not alike influenced by the same poison. The carbonic acid gas, when inhaled, immediately destroys life, but when taken into the stomach, is not unpleasant to the taste, and in some cases, is useful as a medicine. The venom of the viper, according to the experiments of Fontana, may, without danger, be taken into the stomach, and is said to possess tonic powers; but if it be introduced into the sanguiserous system, it proves rapidly destructive.

Examples, illustrating the relative operation of poisons, might be advanced without number; suffice it to say, that the preparations of copper, arsenic, zinc, and other metals, though the most deleterious articles belonging to the catalogue of medicines, are, when properly managed, important acquisitions to the materia medica.

I have shewn above, that the most valuable medicines may occasionally prove deleterious, yet no one contends, on this account, that these articles should be excluded from the meteria medica. Lead, like all other powerful medicines, when given in too large quantities, becomes a poison; but we have the authority of many respectable physicians for afferting, that its cautious internal exhibition may be practised with safety, and frequently with the greatest advantage to the patient.

It was my intention, when I chose lead as the subject of a differtation, to have instituted an extensive series of experiments, in order to ascertain, as nearly as possible, its operation on the human body; but I regret, that from the difficulty of obtaining a sufficient number of subjects upon which to operate, together with the limited time allowed to prepare a thesis, my design has been in a great measure frustrated.

Lety the many which the party winter about STAR STRAIGHT STEEL STEEL

OF THE

MORBID EFFECTS OF LEAD.

NICANDER, a physician and poet, who lived foon after the time of Hippocrates, feems to have been the first who noticed the poisonous nature of lead; he mentions ceruffe and litharge, as the only metallic poifons known at that time; fince that period much has been written on the fubject. The effects of lead on the conflitution are too obvious to escape our notice; the pallid countenances and frequent indisposition of persons daily occupied in lead mines, are fufficient indications of its unwholesome qualities. I shall, however, notice a few instances, in which its deleterious effects were very plainly marked. The colic, which was fo long epidemic in Devonshire, was owing to the poifon of the leaden utenfils, used by the inhabitants of that part of the country in making cider, which constitutes their principal drink.*

^{*} Vide Medical Transactions, vol. 1.

The celebrated German traveller, Professor Thunberg, on his passage to the Cape of Good-Hope, fupped, together with many others on board the vessel, on some pancakes, in which, through miftake, there was a confiderable quantity of white lead. The confequences of this accident were very diffreffing, although no life was loft; its first effects were fickness at stomach and puking; and most of those, in whom this took place immediately, remained entirely exempt from further inconvenience. The captain, however, who vomited in the commencement, was attacked two days after, with a most violent colic; he was of a confumptive habit, but while the morbid effects of the poison prevailed, he was free from cough and other fymptoms of the pulmonic affection. No one, who fuffered from this unfortunate mistake, sustained greater injury than Dr. Thunberg; he was affected with nausea, violent reachings, colic, a complete falivation and ulceration of the mouth, accompanied with a difagreeable stench; inflammation of the eyes and face, great pains in the ears, difficulty of breathing, violent throbbing of the temples, with pain in the head to fuch a degree, as to warrant the apprehension of apoplexy. The chaplain and cook's mate, were affected nearly in the same manner, except that they had, at times, a perfect iliac passion.

The injurious effects of this mineral are not exclusively confined to the human species; quadrupeds, birds, and even vegetables fometimes fuffer from its noxious influence. Dr. Percival mentions a dog, who from laying frequently on a plate of lead before a fire, was feized with a paralytic affection of his limbs. Cats, he likewife observes, are sometimes driven to madness by it; and we have instances, on his authority, of birds being killed by eating pieces of red wafer, into the composition of which, enters a small quantity of lead. The effluvia that arises from smelting mills, where lead is obtained from the ore by fettling on the adjacent fields, entirely destroys the herbage.

contrate en la contrate de la contra

OF THE

PREMONITORY SYMPTOMS

FROM

The Poison of Lead,

AND THE

PROPHYLAXIS.

AS the prevention of disease ought to be a primary confideration with physicians, a knowledge of the fymptoms, which indicate its approach, are certainly of the greatest importance; from ignorance and inattention to thefe circumstances, many diseases have proved fatal, which might eafily have been remedied, had they been attended to in their commencement. Some of the most formidable, as pulmonary confumption, mania, epilepfey, apoplexy &c. (part of them till lately fupposed to be incurable) are now known to be frequently cured, when attended to in their recent or forming state; but when neglected till they are perfectly formed, or have attained their greatest degree of violence, how often, is even the physician of skill and information, vanquished in his efforts to give

relief; - and how feeble is theory in aiding his attempts. In few difeases is the danger from delay greater than in those induced by the poifon of lead, and the prevention of none can be accomplished with greater ease. Its deleterious effects discover themselves by a flow and almost imperceptible progression; fmelters, miners, painters and potters are feldom affected before they have been exposed to its effluvia for fome months, and frequently not till two, three, or even four years have elapsed. The first remarkable change which is produced in perfons who are exposed to the pernicious influence of this metal, is an unufual yellowness of the complexion, with a dark streak below the eyes. The stomach next begins to be difordered, accompanied with preternatural constipation of the bowels, slight pains about the precordia, cramps in the legs at night, and a fense of weight about the epigastric region. An enlargement of the glands about the throat and fwelling of the gums, with an increased secretion of the faliva, sometimes occur; and these symptoms, if not attended to, frequently terminate in colic and paralysis, confumption, rheumatism and afthma. As foon, therefore, as we are apprifed of their approach, it becomes necessary to

counteract them by every possible means .-With this intention, perfons who are difcovered to be in the least affected, should immediately discontinue all employments, in any manner connected with the metal; obviating, at the same time, costiveness and other urgent fymptoms. Among the various articles which have been proposed as antidotes for this poifon, particular dependence has been placed upon oleagenous and mucilaginous fubstances. I am led, however, from the refult of feveral experiments, to conclude that they are ineffectual, when large quantities of the metal have been received into the stomach. Experience teaches us, nevertheless, that a greafy unctuous diet in some degree protects persons who work in lead ore, from the noxious influence of its effluvia.

Mercury, in fmall doses, has been lately recommended as a preventive against this poison, and there is no doubt it would be serviceable; but the inconveniences arising from its operation, are too considerable ever to allow of its general adoption. Cleanliness is an object of great importance, and it is known, that persons employed in the different manufactories of lead ore, who keep them-

felves clean, are less liable to be affected by it than those who are negligent of their perfons, and who never take the precaution of washing their hands previous to eating.

OF THE

PRIMARY EFFECTS

OF

THE SACCHARUM SATURNI,

ON

The Human Body.

There has been a great change of opinion, concerning the operation of medicines, within the last few years. The doctrine of their fedative operation, was for a long time universally admitted. But latter observation and experience have shewn its absurdity. The justly celebrated Dr. John Brown, was the first to affert, the stimulating qualities of opium; and from this period, succeeding writers were induced to examine more attentively, the primary effects of other substances.

Those, who have written on the subject of lead, have noticed only its remarkable power in lessening the frequency and force of the arterial system, and there are many, who still

adhere to the opinion of its direct fedative action. In order, therefore, to determine whether its primary operation be that of a fedative or incitant, I made the following experiments.

EXPERIMENT 1.

At 9 o'clock in the morning, half an hour after eating a moderate breakfast of light food, I took one grain of faccharum faturni, mixed with a little fyrup, my pulse beating 69 strokes in a minute. In 5 minutes it stood at 69; in ten minutes it was increased 2 strokes; in 15 minutes it beat 73, but was not increased in fullness; in 20 minutes I believed it to be quicker, and it beat 75; in 30 minutes it beat 72; at 40 minutes, 72; in 50 minutes it beat 68, and was fomewhat fmaller; in one hour I felt a flight degree of uneafiness at the stomach, which soon went off; in 70 minutes my pulse beat 66, and was reduced in volume; I attributed this finking of the pulse to the temperature of the room, which was lowered from fuffering the fire to go nearly out; in 80 minutes my pulse was 68, and natural; but continued to vary from 67 to 70, for an hour after.

EXPERIMENT 2.

At 10 o'clock, A. M. about one hour after breakfast, I took 2 grains of the sugar of lead, mixed in a little syrup. My pulse beating 70 strokes in a minute, its natural standard; when the following changes were observed.

In	ıls	e bear	3	6 70	9 72	12 72	15 74	18 73	20 74	23 2 73 7	26 29 70 68	68
30	6	40	45	50	55	60	70	80	90	100	110	120
6	7	69	68	67	1 65	1 65	1 00	1 64	100	1 67	67	69

130 | 140 | 150 | 160 minutes 67 | 68 | 68 | 70 strokes.

In 15 minutes after taking the medicine, my pulse was increased in number, and rather quicker, but no increase of strength was perceptible. In 50 minutes my pulse was slower, and a little diminished in volume; at the expiration of one hour, there was an evident diminution of strength in the pulse—at 80 minutes I had a nausea, which lasted 15 minutes, my pulse a little contracted; in 150 minutes my pulse had resumed its natural action. I continued to examine it for several hours after, when it varied from 68 to 73.

EXPERIMENT 3.

In one hour and 20 minutes, after taking a light breakfast, I took 3 grains of saccharum saturni, dissolved in a little syrup; my pulse beating 72 strokes in a minute:*

In pulse	beati	2	5 68	8 75	11 1 73 7	4 17 76	20 74	23 71	26 2 70 7	9 32 68
35 68	38 70	41 70	45	50 69	60 64	65 64	70 80	0 85	90 62	100 68
110 64	12 6	0	130	140 68	150 69	165 65	180	190	210 70	225

240 minutes 71 strokes.

In 8 minutes my pulse was perceptibly stronger, with a slight heat about the precordia; at the end of 14 minutes my pulse had increased in frequency, and was quicker; when 20 minutes had elapsed, my pulse was considerably reduced in volume; but remained preternaturally quick; in one hour my pulse was still smaller than natural, and in frequency, reduced from 70 to 64, with an uneasy sensation in my head; in 85 minutes my pulse was round and soft; at one

^{*} Dr. Horsefield did me the favour to be present during this experiment.

hour and 20 minutes it had nearly refumed its original fulnefs.

At the expiration of two hours and 40 minutes I had a pain in the head, with naufea, which lasted about 15 or 20 minutes; during the remainder of the day my pulse was natural.

EXPERIMENT 4.

At 40 minutes past 10 o'clock A. M. my friend Mr. George Lee, took 4 grains of the fugar of lead, formed into a pill, with a fmall quantity of conferve of rofes; his pulse beating 60 strokes in a minute, and perfectly natural. In 5 minutes his pulse beat 61; in 10 minutes it beat 62; in 15 minutes his pulse had increased three strokes in frequency, but was not fuller; in 20 minutes, his pulse stood at 63, and was quicker; in 25 minutes it beat 64, but no unufual fenfations were experienced; in 40 minutes it was at 60, and a little contracted; from one to two hours his pulse had lessened in force, and was gradually finking; I was now prevented from further examination, but had every reason to conclude that it would have been reduced feveral strokes more.

EXPERIMENT 5.

At 10 o'clock in the morning, being a little indisposed with a slight head-ach, with some tension and sulness of the arterial system, I took 4 grains of saccharum saturni, combined with a small quantity of conserve of roses; my pulse beating 74 strokes in a minute:

In pulse	beat	3	6 73	9 75	12 13 76 78	18 80	21 80	24 83	28 81	32 82	36
40	50	60	170	90	100	110	120	130) 1	40	150
80	78	80	76	78	76	75	72	68	3	70	67

160 | 170 | 180 | 190 | 200 minutes 69 | 72 | 75 | 70 | 72 strokes.

In 8 minutes I experienced a fensation of heat at stomach; before 20 minutes had elapsed, my pulse was increased in number and fulness, with an increase of heat at stomach; when 40 minutes had expired, my stomach was more affected, but had no disposition to vomit, with a preternatural warmth of the cutaneous system; in 50 minutes had an operation on the bowels; in an hour my skin became moist, which continued for 15 minutes; my pulse at the end of two hours was quicker, though not more frequent than

natural; about this time I had a fecond evacuation from the bowels, which was not fo copious as the first, but more bilious; in three hours my stomach recovered its tone, and my head was relieved from pain.

During the remainder of the day, I felt in no other respect disordered, than a slight disficulty of breathing; my pulse being rather more feeble than usual.

EXPERIMENT 6.

On the tenth day, after experiment the fifth was made, at feven o'clock, A. M. I took on an empty stomach, five grains of the sugar of lead; my pulse beating 66 strokes in a minute:

In puls	e bea	3 t66	68	9 71	12 71	15 73	18 76	20 75	23 26	6 30	35 75
40 73	45 70	50 73	55 72	60 70	70 67	80	90	100 65	110 63	120 60	130 58

^{140 | 150 | 160} minutes 60 | 58 | 57 strokes.

In 6 minutes I felt a warmth at stomach; in 12 minutes my pulse was somewhat stronger, with an increase of heat at stomach; in 20 minutes the sulness of the arterial sys-

tem was rather greater; at the end of 40 minutes my cheeks feemed warmer than usual; when one hour had elapsed, I selt some uneasiness at stomach, my pulse being weak and corded; when two hours had elapsed, considerable nausea occurred, my pulse seeble and not so frequent; at 130 minutes I made several efforts to vomit, but discharged nothing from my stomach; at 140 minutes the nausea returned.

At 160 minutes, feeling extremely unpleafant, with a feeble and tremulous pulse, and not having eat any thing during the morning, I took some toast and coffee, which raised my pulse and greatly composed my stomach.

I now laid down, and after a while a moifture came on, that lasted about half an hour; at one o'clock I arose, being much recovered, and took a little soup. I had throughout the after part of the day, a small degree of tightness across the thorax, my pulse being irregular and seeble; my urine was highly coloured, but my head and bowels remained unaffected. The two following experiments were communicated to me, by Dr. Robert Black, and Mr. Washington, of Alexandria.

EXPERIMENT 7.

Having made, fays Dr. Black, a very light and early breakfast, at 10 o'clock, my pulse beating 88 strokes in a minute, I took two grains of the fugar of lead. In 10 minutes my pulse rose to 96, somewhat contracted and a little tense; my hands and wrists were rather cool, accompanied with a confiderable degree of moisture. In 20 minutes my pulse fell to 90, became rather feeble, and the moisture on my skin increased; in 25 minutes my pulse stood at 88, I felt a slight fensation of heat and uneasiness about the region of the stomach; it was not accompanied with the flightest nausea, but it appeared to be confined principally to the external parts; in 50 minutes my pulse rose to 94, and became irregular. This change I attributed to my approaching the fire, and placing my feet on the projecting plate of a Franklin stove, which was heated. In 75 minutes my pulse beat 90, which is nearly its natural standard, it was foft and regular,

continuing in the same state throughout the day.

EXPERIMENT 8.

At ten o'clock, fays Mr. Washington, after taking a flight breakfast of bread and coffee, I took one grain of faccharum faturni mixed in fugar and water-my pulse beating 80 strokes in a minute, was somewhat fuller and more irregular than natural; in 5 minutes it rose to 84, but not so full or varying; in 10 minutes it was 98, but appeared fmall, tenfe and irregular; in 15 minutes it fell to 86, was foft and even; in 20 minutes there was fome degree of warmth about my stomach, accompanied with a flight naufea, and difpofition to vomit, though an emefis did not take place; in 25 minutes my pulse was at 84, uneven and full, at this time my skin became moist and clammy, imparting to the touch a fenfation not diffimilar to that which is usually experienced, about the close of a paroxysm of fever; in 60 minutes it was regular and foft; and at 76 minutes I felt a difagreeable tafte in my mouth, very like that produced by copper, and refembling that which accompanies an incipient ptyalifm; and in a fhort time after an increased secretion of faliva took place, which continued during the day. At the expiration of 100 minutes my pulse resumed its natural standard.

EXPERIMENT 9.

To a dog, whose pulfation, as well as I could determine, was about 114 in a minute, I gave one drachm of fugar of lead: in 10 minutes he appeared a little uneafy; in 15 minutes his pulse became more frequent and strong with a disposition to vomit. At 20 minutes there was a great irregularity in his movements, with laborious respiration; the diaphram feeming to be much convulfed; in a few minutes after this, he discharged from his stomach some frothy matter, and I judged, nearly all the medicine; at 32 minutes he had an evacuation from the bowels, confifting of fæces and mucus; when 50 minutes had elapsed, his pulse was nearly imperceptible, imparting to the touch a tremulous fenfation; in an hour he became dull, (being much relieved) and fell asleep; in four hours from this time, I injected into his right jugular vein, a faturnine folution; in about 30 minutes he expired, having been previously affected with convulfions, delirium and other distressing symptoms. On examining his body, the appearances were as follow: The whole stomach was contracted, particularly its fuperior orifice; its blood-veffels diftended, with inflammation in feveral parts, especially at the termination of the œfophagus; the duodenum was turgid and inflamed, containing a fmall quantity of a dark fluid; the other intestines had a natural appearance—the bladder, kidneys and liver were not affectedthe gall-bladder was filled with a quantity of bile, of a darker colour than usual, and there were fome marks of congestion in the lungs. On opening the cranium, the external part of the brain appeared natural, but within its fubstance was found a fmall quantity of extravafated blood.

In two fimilar experiments, except that the folution was not injected into the blood-vessel, the result was nearly the same: In the one, however, a preternatural secretion of saliva occurred; and in the other it was remarkable that the rectum was the only part of the intestines which shewed marks of inflammation.

Since the first introduction of lead into the materia medica, it has been uniformly classed with medicines called sedatives, till Professor Barton suggested the probability of its primary operation being that of a stimulant; this opinion is corroborated by the preceding experiments; and the analogy of its effects, without those of other incitants, assords additional proof of the truth of the doctrine: Let us then briefly point out the similarity of the effects of lead, to those of other stimulants.

Do many stimulants, as alkohol, opium, digitalis, stramonium, &c. increase the frequency and force of the arterial system? So does the sugar of lead. In experiments the 4th, 5th, 6th and 7th, the pulse rose from 8 to 12 strokes in the minute, with considerable increase of force tension, and quickness.

Have incitants the effect of ultimately reducing the force and frequency of the arterial fystem? So has the fac. fat. to a great degree.

Do opium, mercury, and fenaka root, occasion an increased secretion of saliva? So likewise do the preparations of lead. This symptom occurred in two of my experiments. Dr. R. Warren says, out of thirty-two cases of colica pictonum, caused by lead, which he had an opportunity of observing, four were salivated, and others complained that their mouths were fore.* The same symptom occurred to a very extensive degree, in Mr. Thunberg and others, as already mentioned.

Are not heat at stomach, nausea and vomiting, consequent upon taking stimulants? Such are the effects of lead. Do not arsenic, opium, and other poisons, produce convulsion and paralysis? The preparations of lead have done the same.

Are not various stimulating substances known to constipate the bowels? Lead has the same effect.

Is the cutaneous fystem influenced by stimuli? So likewise it is by the sugar of lead, causing preternatural heat; this was the effect of lead on myself and several of the subjects of my other experiments.

^{*} Vide Medical Transactions, vol. 1.

Do certain incitants produce diaphoresis? In the experiments communicated from Mr. Washington and Dr. Black, and in one that I made upon myself, the same effect was produced.

Are the different fecretions and excretions affected by stimuli? So likewise they are by the preparations of lead.

Have diffections, discovered inflammation, and other morbid phenomena, to be the effects of stimuli? Such also are the consequences of the sugar of lead. In the examination of two of the dogs, that had taken a drachm of this salt, nearly the whole alimentary canal evinced marks of inflammation and excessive action.

The interesting and minute narrative of its deleterious effects by Mr. Thunberg, as mentioned above, confirms the opinion beyond a doubt; fever, inflammation, and a numerous train of other malignant symptoms, were in this instance its consequences; and no one will contend that these phenomena could have been produced but by powerful stimulants.

We may then fairly conclude, the primary operation of the preparations of lead is that of a stimulant, united to an astringent principle, but that thefe effects are extremely transitory, and are quickly followed by those of a contrary nature. Although we are enabled by experiment, to decide whether the first effects of a substance be fedative or stimulating, still the subject of the modus operandi of medicines is cloathed in much obscurity. To ascertain precisely, upon what principle, or in what manner, the action of fubstances upon the human body is brought about, will be a point as likely to elude the investigation of man, as any connected with the science of medicine. A pill of opium may be taken and exert every active property it possesses, without a diminution of From the rapid improvements which the science of chemistry has lately undergone, fome have been induced to suppose this beautiful and pleafing fludy would lead to an explanation of those mysterious laws and operations of nature. The chemist by analysis, may discover if a substance be composed of a gum, refin, &c. He may, moreover, determine accurately the proportion of each elementary part that is necessary to the formation of a

fubstance; but, can he, by these means, explain the principles and functions of animal or vegetable life; or is this science sufficient to teach us, why two gums, the external characters of which are the same, should in their operation produce such different effects; or will it inform us, wherefore, some substances produce salivation, some vomiting and others sleep?

Whether it depends upon a particular modification of particles, or upon a certain proportion of the incitant, aftringent and fedative principles, or upon what circumstances it does depend, must remain a secret, until phisiologists shall obtain a more perfect knowledge of the laws and economy of organic matter.

All the second of the second o Will be at the second of the second OF THE

INTERNAL USE

OF

THE SUGAR OF LEAD,

IN DISEASES.

Before we enter upon the confideration of the use of lead in diseases, it will be necessary to enquire, to what extent the preparations of this metal have been taken, either intentionally or by accident, without injury.

A girl took eighteen grains of sugar of lead, every day for three weeks, without any injurious consequences. A weak man, forty years old, took (at four doses, in the course of forty-eight hours, in a liniment) one drachm of saccharum saturni, without any other disagreeable effect than a giddiness and heat at stomach. At Salford, in Warwickshire, a perfon swallowed near a bottle of Goulard's extract; and no inconvenience ensued.*

^{*} Johnson's Medical Essays.

A woman took by accident, a scruple of acetate of lead, without inconvenience from it.*

I knew a child of fix years old, to drink a portion of faturnine folution, which contained about four grains of faccharum faturni, and no ill effect followed.

A man for the cure of the gout, was advised by an empiric, to use an oxide of lead, which he did, in such quantities, as to take two pounds in sisteen days; he remained free from disease, till twenty days after he began its use, when he was attacked with dysentery. His recovery was preceded by a very obstinate jaundice.†

I have been informed by a gentleman of Devonshire, says Sir George Baker, that it had long been a custom with him, to restrain the acetous fermentation of his cider, by throwing into it a quantity of sugar of lead; and that he was convinced, no ill effects ever followed the practice.‡

^{*} Professor Barton's M. S. Lectures.

[†] Johnson's Medical Essays, p. 117, 123, 124.

I London Medical Transactions.

Many English Dispensatories, and old systems of medicine, recommend, from seven to twelve grains of saccharum saturni at a dose. And in their former naval and military Dispensatories, a scruple is ordered for a single dose.

Anterior to the time of Paracelfus, we have nothing certain respecting the internal use of the preparations of lead. From the early part of the sixteenth century, till the middle of the seventeenth, Tachenius, Crollius, and others were accustomed to employ them as valuable medicines. In 1741, Hundertmark, published a differtation at Lipsic, wherein he assures us, that he gave the sugar of lead, to the quantity of four grains, with wonderful success, in some desperate cases of phrenetic delirium, and in some cases of peripnumony; * and several differtations also appeared in Europe on the same subject, towards the end of the eighteenth century.

However much, modern physicians have wished to lay aside the internal use of these preparations, it is not the less certain, that the ancients ascribed considerable virtues to them,

^{*} Vide Gmelin.

as powerful and ufeful remedies in many difeases.

The Chinese, very frequently administer internally various preparations of lead, to which they ascribe extraordinary virtues.*

Such efficacy was this mineral supposed to posses in phthis pulmonalis, that one of its preparations acquired the appellation of tinctura antiphthisica. Etmuller, who directed his attention particularly to the use of faturnine preparations in consumption, to prevent the cough and sweating that occur in hectics, recommends a formula, containing ten grains of the sugar of lead, to be taken every night at two doses.

Mr. White, in a letter to Dr. Duncan,† obferves, that he found the action of the heart and the frequency of the pulse, evidently diminished by the use of the tinctura saturnini, and that in phthis pulmonalis, it constantly abated the hectic fever and sweats.

Sir George Baker acknowledges, that there are fcarcely any medicines which pro-

^{*} Vide Medical Transactions.

[†] Medical Com. vol. 3.

duce their effects more immediately, or more certainly, than the preparations of lead, in colliquative fweats, in fluxes, or in hæmorrhages.*

The digitalis has been particularly ferviceable in certain stages of consumption, and from the similarity of its operation to that of sugar of lead, I am led to conclude, the latter will prove a valuable remedy in that stage of the disease in which great morbid action of the arterial system prevails.

In Diabetes,† the acetate of lead has been used with great advantage. Dr. Rush has lately endeavoured to explain the phenomena of this disease, by considering it a state of sever, and the success which attended the practice to which this opinion leads, corroborates the idea of its being a febrile affection. The remedies which he recommends are, repeated blood-lettings in small quantities, salivation, low regimen &c. Considering the power of saccharum saturni, in restraining all morbid secretions and evacuations, and its known influence over the pulse, it appears to

^{*} Medical Transactions, vol. 1.

[†] Vide Etmuller's Practice.

me, that much good would be derived from its application in this difease.

Much likewife has been faid concerning the use of this medicine in dysentery and diarrhœa; and indeed, from their inflammatory nature, and from their being feated in that fystem, on which lead exerts its most immediate and permanent operation, it appears to be particularly adapted to their cure, and accordingly we find that it has been used with advantage by respectable writers. When there is inflammation of the intestines, Revirius observes, it will be necessary to let blood, and give glyfters of the fugar of lead, diffolved in rofe water; and, at the same time fays, ten grains of this falt, mixed with conferve of roses, should be taken by the mouth, every morning. By Etmuller the acetate of lead, is confidered a specific in dyfentery; and when aftringents and abforbents are indicated, Dr. Shaw, confidently informs us, it is a most valuable and useful medicine.

At the royal and univerfal dispensary, there is a formula, containing half a grain of this falt, which is often employed by the physicians of that institution, in cases of hæmorrhages, and obstinate diarrheas with advantage, and no instance has occurred, in which any injury arose from its use.*

The New-York Medical Repository contains an interesting account of the efficacy of the acetate of lead, in relieving feveral cafes of chronic diarrhœa, communicated in a letter from Dr. John Archer, jun. of Maryland, to Dr. John Claiborne of Virginia. The first case which Dr. Archer relates, was of a Mr. S. R. aged 22 years, a man of strong constitution, who, in June 1798, after severe exercife, was attacked with dyfentery, which at the expiration of four weeks terminated in a painful diarrhæa, which continued unrestrained until the latter end of November. His body was much emaciated, and during this time he had violent pains in his bowels, with frequent discharges of blood. After the failure of the usual remedies, faturnine injections were administered, but these also proving ineffectual, recourse was had to the exhibition of this preparation of lead, by the mouth, to the quantity of three grains, three times a day, made into a pill, with fyrup,

^{*} Vide Clutterbuck's Paper, &c.

and instructions to the patient to discontinue the medicine, as foon as the purging moderated. In two days the diarrhœa was much abated; he continued however, to take the medicine, the confequence was costiveness, and extreme pain of the bowels; which was speedily removed, by a dose of Glauber's falts. His difease again returned, but by the use of the faturnine pills, reduced to two grains of the falt, his diarrhœa was kept under, and in the course of two weeks, the disease was perfectly cured. The other cases related by this gentleman are equally interesting; in none of them were there any permanent difagreeable confequences from the use of the medicine: The flight inconvenience which occurred in the case noticed, was obviated by a fingle dose of falts. I regret that the pulse was not more attended to in the relation of thefe cafes.

IN EPILEPSY.

THERE is no disease, the pathology of which is involved in more obscurity than this, and till theory and experience shall elucidate its nature, every medicine that can be prescribed for its cure, must be in a great measure bordering on empiricism.

Dr. Rush, to whom we are much indebted for the explanation of the phenomena of fome difeases, is of opinion, that to epilepfy, no determinate feat can be affigned. He fupposes it to be exclusively confined, neither to the arterial, nor nervous, nor any other fystem, but to extend its morbid influence over the whole animal oconomy. He admits it to be a nervous or febrile difease, and not unfrequently partaking of the nature of both in the same person, but if we judge of the truth of theory, from the fuccess of its application, in practice, we must acknowledge that this, like every other, ferves only to inform us of our ignorance, and to convince us that we have yet to acquire much important information from the mysterious

laws and operations of nature, before the science of medicine shall become perfect, and man shall cease to die, but from old age.

That epilepfy is often a febrile difeafe, may be inferred from its being induced by the fame causes which produce other fevers, and from its being accompanied with many of the usual symptoms of fever; the pulse is often full and tenfe, pupils of the eyes dilated, face flushed, &c. It moreover attacks perfons at that time of life when the fystem is plethoric, and pre-disposed to febrile diseases; it fometimes also occurs with confiderable violence, without any appearance of fever. Opium, camphor, mercury, and lately, the nitrate of filver, have all been ufeful, but the fugar of lead will probably prove to be equal, if not fuperior, to any of them in this difeafe; it has frequently been known to fuspend the fits, and has fometimes cured the difease, but admitting that it only protracts the intervals between the fits, this effect, alone, would be fufficient to recommend it to the attention of physicians. A person, not well versed in the treatment of epilepfy, will find it difficult to point out the particular stages and circumstances of this disease, under which the acebut from the known power of this medicine in lowering the pulfe, I should suppose it particularly adapted to the cure of those cases of the disease which are accompanied with an inflammatory diathesis, and which have regular periods of occurrence; but as this medicine, to be effectual, requires to be taken for some time before the fit, it might be objected to, in cases where the fits occur at distant or irregular intervals, or where the paroxysm was not ushered in by some characteristic symptom, as injurious effects sometimes result from its long continued use.

Near two hundred years ago it was used with success in the epilepsy of children, by Agricola; and a boy was lately effectually cured of this formidable disease by taking two grains of the acetate of lead three times a day; and in two instances likewise in the Pennsylvania Hospital, it suspended the fits for several weeks.*

In Fluor Albus and Gleets, the acetate of lead has been administered internally, with success. In the year 1799, I had an oppor-

^{*} Dr. Rush, M. S. Notes.

tunity of witnessing its efficacy in a case of leucorrhæa, in a poor woman aged about 27 years-fhe had been subject to the complaint for near twenty months, her pulse was preternaturally quick and tenfe, and beat about 85 or 86 in a minute, she had been several times bled, falivated, and had observed an antiphlogistic regimen, without any benefit. In this fituation she was directed to take half a grain of fugar of lead, combined with two grains of gum kino, three times a day; on the third day after the use of the medicine, the pulse was diminished in frequency, three or four strokes, but the discharge and other fymptoms remained as before. The quantity of the falt was now increased to one grain, to be taken at the fame intervals as before; on the fecond day after the use of the increased dose, her pulse continued to decrease in frequency, and the discharge was somewhat abated; fhe was now costive, and being apprehenfive that difagreeable fymptoms might enfue from the too long retention of the medicine in the fystem, it was intermitted, and a dose of castor oil exhibited; in a few days after, the medicine was again had recourse to, which effectually relieved the patient, without causing the most trifling unpleasant fymptom.

IN HÆMORRHAGES.

In the cure of no difease will the sugar of lead be found more advantageous than in hæmorrhagy; and with Dr. Heberden, I believe, there is scarcely any medicine that could with more propriety be called an internal specific, than saccharum saturni, in hæmorrhages of the prima viæ.

Dr. Reynolds of London, has experienced in a particular manner, the efficacy of the preparations of this mineral, in restraining hæmoptysis, epistaxis and memorrhagia. To a young man, labouring under hæmoptysis, was given the following formula.

> R. Sac. Sat. gr. 1. Con. ros. rub, grs. 2. Laud. gtt. v. m.

In a day or two the dose was increased to one grain and a half; he had been bled several times, and had taken saline medicines, and observed the lowest diet without advantage. His pulse beat 108 strokes in a minute, full and hard; on the fourth day after taking the medicine, the spitting ceased entirely, his

pulse reduced to 90, foft and even; he had regularly two motions a day.*

If the spitting of blood is violent, Shaw advises, half a scruple of saccharum saturni to be taken every fifth hour, the patient at the same time to drink freely of diluted tincture of red roses.

In June 1800, a poor man aged 30, of flender make and healthy habit, was attacked with a haemoptyfis, which was occasioned by the too great exercise of the lungs, from blowing the fife, his pulse beat 89 in a minute, quick and tense, breathing laborious, and indicating considerable congestion. He was directed to take the following formula, every fourth hour:

R. Sac. Sat. 2 grs.
Theriac. And. 3 grs. m.

In 15 hours after he commenced taking this medicine, the spitting had entirely ceased, the pulse much reduced in force and lessened in frequency; the congestion being not effectually relieved, he took 15 grains of calo-

^{*} Medical Com. vol. 3.

mel, which operated moderately on the bowels, at the same time slightly affecting the mouth; nothing else after this was necessary.

- "A woman between forty and fifty years old, labouring under uterine hæmorrhage, was relieved by the faturnine tincture."
- "Another woman had a flooding terminated, which had lasted two months, unchecked by all the most powerful astringents in common use, by taking four grains of saccharum saturni."*

Equal quantity of white vitriol and fugar of lead, in the proportion of half a grain, has also been given with the happiest effects, in menorrhagia.

In epistaxis, the success attending the internal use of this salt, has been alike evident; but I suppose it might be employed externally in this variety of bleeding, with every advantage. A physician in this city, was in the habit of using it as snuff, in bleeding from the nose, which had every desired effect.

^{*} Med. Tranf. vol. 2.

In Tetanic Affections, might not the faccharum faturni, be employed with great advantage? The use of this remedy in tetanus, is fanctioned by the authority of Mr. John Hunter, and at all events deserves a trial. The following interesting relation of facts and observations, was communicated to me, from Dr. Barton:

" DEAR SIR,

- "I exceedingly regret that my late fevere illness has prevented me from furnishing you with the particulars of my practice with the acetite or fugar of lead, in the treatment of different diseases. At present, I can give you but little on the subject that is worthy of your attention.
- "In the course of the last five years, I have employed this preparation of lead in cases of hæmorrhages; in hæmoptysis, hæmorrhages from the stomach and intestines, and in uterine hæmorrhages, both of pregnant and of non-pregnant women. Of late, I have also employed it, very largely, in a case of obstinate gonorrhæa. But my practice having been principally confined to the treatment of hæmorrhages, I shall content myself with making a few observations on the use of the medicine in these diseases.
- "In a case of hæmoptysis, which threatened the speedy death of the patient, I gave the sugar of lead, in doses of six grains every two hours, for several hours together. I ascribe the complete recovery of my patient solely to this treatment. This was one of the first cases in which I exhibited the medicine, and I should not have ventured upon such doses (for I had been taught to believe, that the preparations of lead ought always to be exhibited with the greatest circumspection), had I not thought the patient irrecoverable by any other means. I had no cause, however, to regret the employment of the lead; for it produced not the least inconvenience.

"But it is chiefly in uterine hæmorrhages that I have administered the sugar of lead. I have now exhibited it in many cases; I think not less than eighteen or twenty. I have ever given it with some advantage; generally to the complete cure of the patient, and, I have never known it to produce any serious inconvenience. It has sometimes seemed to occasion a slight sickness at stomach, but this, which could not always be ascribed to the lead, was nowise injurious to the patient. In one case, it seemed to occasion a slight degree of tenesmus. But this was soon removed by a dose of castor-oil. I am certain, that it sometimes operates, even in small doses, as a laxative.

"In the administration of the fugar of lead, I have almost always combined it with a portion of opium. I still continue this practice, because I have been induced to believe, that the lead may be given in much larger doses with than without the opium. Perhaps I may be mistaken.

"With respect to the dose of the fugar of lead, this, I need not tell you, must vary according to the age, the strength, and the case of the patient. Hitherto, my practice with this medicine, has been almost folely confined to adults. In uterine hæmorrhages, I have feldom given lefs than two grains at a dofe (in combination with a grain of opium), twice a day. But I have often given larger dofes. Even fmall doses, however, fometimes produce an early, and I was going to fay, an immediate effect. This, I have particularly remarked, in the case of a lady, who had laboured under a menorrhagia for fourteen days, before I faw her. I found her with a quick and irritated pulse, flushings, fevere pain in the region of the loins, and with much affection of the head: I had no hesitation in advising her to lose some blood from her arm; to this she objected. She was immediately put upon the use of the sugar of lead; in

less than one hour, the hæmorrhage was considerably diminished; she took a second dose, about four hours after the first, and was so completely recovered that there was no necessity to continue the medicine.

"I shall not trouble you with any speculations concerning the modus operandi of this medicine in curing hæmorrhages. I have touched upon this subject in my lectures, which you have done me the honour to attend; and, I doubt not, that you have, in the course of your experimental inquiry, been able, in some measure, to satisfy yourself as to the manner in which the lead is of use in these cases. I will only observe, that I am persuaded, that the medicine does not produce all its good effects, merely by its quality of lowering the pulse. It appears to be useful both in the active, and in the passive hæmorrhages. I am not ashamed to mention these old names.

"I have had fome experience with the use of the digitalis in the treatment of uterine hæmorrhages. This is certainly, a valuable medicine. I can readily believe Dr. Ferriar, that it has often done much good. In my hands, however, it has appeared to be much less efficacious than the sugar of lead. Of one thing, I am persuaded, that the digitalis does not, in general, produce so immediate an effect upon the disease as the lead does.

"I am fure that you, who know with what caution I always speak of medicine in my lectures, will not suppose, that I recommend the sugar of lead as a specific for the cure of uterine or other hæmorrhages. I considently recommend it, however, as a powerful and valuable medicine; and I cannot but suppose that it may generally, if not always, be exhibited with perfect safety. Yet it may be necessary, in cases in which we have not already given the me-

dicine, to watch its effects with a nice attention. I believe, that there are some persons who are liable to be injured by very small portions of lead. I believe it is a fact, that lead, whether it be taken into the stomach, or applied in the shape of vapour, externally to the body, does not, in general, produce its bad effects until some days after it has been applied.

"These facts should lead us to observe some degree of circumspection in the use of the medicine, in the treatment of diseases. It is not impossible that one or two grains of the sugar of lead, may now and then occasion some inconvenience. But in the exhibition of this medicine, I have been more governed by the other circumstance which I have mentioned: I have been fearful lest the patient might suffer from an accumulation of the lead in the system. When, therefore, in the treatment of hæmorrhages, I have not sound evident benefit from the medicine in two or three days, I have laid it aside for a short time (three or sour days); and, when necessary, have resumed the use of it again.

"I fear that this very imperfect, but faithful statement of my practice, can be of little use to you. Such as it is, I beg you to receive it with my best wishes for your prosperity and usefulness in life.

I am,

My dear Sir,

Your affectionate friend,

And humble fervant,

BENJAMIN SMITH BARTON.

Philadelphia, June 4th, 1801.

Hi motus arteniamme atque hoccordamma tunta = escent



Med. Hist. WZ 270 5472e 1801

