Every man his own physician : being a complete collection of efficacious and approved remedies, for every disease incident to the human body ; with plain instructions for their common use / by John Theobald, M.D. author of the Medulla medicine ; compiled at the command of His Royal Highness the Duke of Cumberland.

Contributors

Theobald, John, -1760. Babcock, John, 1764-1843 National Library of Medicine (U.S.)

Publication/Creation

Hartford : Printed by J. Babcock, 1800.

Persistent URL

https://wellcomecollection.org/works/f5txa8vx

License and attribution


This material has been provided by This material has been provided by the National Library of Medicine (U.S.), through the Medical Heritage Library. The original may be consulted at the National Library of Medicine (U.S.) where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.


You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



NATIONAL LIBRARY OF MEDICINE Washington





EVERY MAN

HIS OWN

PHYSICIAN.

BEING

A complete collection of efficacious and APPROVED REMEDIES, For every DISEASE incident to the HUMAN BODY.

WITH

Plain Instructions for their common Ufe.

BY JOHN THEOBALD, M. D. Author of the Medulla Medicinæ.


Tompiled at the command of his Royal highnels the Duke of Tumberland,

EIGHTH EDI F

1840

HARTFORD : PRINTED BY J. BABCOCK.

-- 1800.---



PREFACE.

ALTHOUGH publishers of receipts have been always very numerous, yet it has so happened, that the least able have hitherto undertaken this task, as the most easy and likely to answer any mercenary views ; which seems to have been the reason, that more able perfons have neglected it as an under part of their profession. Nevertheless, I have persuaded myself, that I should perform a work, neither intirely woid of use, nor foreign to the duties of my profession, if I made the public partakers of the principal helps against most diseases, which I had either learned by long experience, or selected from the writings of the most eminent physicians. As these receipts are published chiefly for the use of persons residing in the country, whose convenience or abilities, will not allow of the attendance of a physician or apothecary, I have studied plainness in my stile, and in my directions, with regard to the treatment of the difeases, that I might render this treatife as useful as possible.

T. THEOBALD.

PREFACE TO THE SECOND EDITION.

THE favorable reception this work has met with, as is sufficiently shewn by the remarkable quick fale of the first large impreffion, has induced the editor (to support the credit of what is really beneficial to the public, and add weight to the receipts) to publish the authorities from whence those receipts were taken; and to make it still more useful, he has also added the signs of difeases, and some plain and concise directions for bleeding. He begs leave to remark, that although Dr. Theobald hath in general given no more than one receipt for the cure of one diforder, yet he has been careful to select such only, as, upon long experience, were found to be the best : and that although receipts are given for difeases which are feldom cured, (as an apoplexy, confumption, &c.) yet, if it is in the power of medicine to remove any of these complaints, there is the greatest reason to imagine, that the remedies directed in this volume will effect a cure.

EVERY MAN HISOWN PHYSICIAN.

ACHES AND PAINS.

R UB a little opodeldoch upon the part affected, two or three times a day, and wear a flannel upon it; if this does not give relief, take twenty drops of volatile tincture of guaicum, (prepared as directed in the Appendix,) every night and morning, in a glafs of fpring water. *Heifter*.

AFTER-PAINS.

Take one fcruple of fpermaceti, five grains of volatile falt of hartfhorn, five drops of balfam of Peru, and half a drachm of Venice treacle; mix them into a bolus, with as much fyrup of white poppies as is fufficient to make a bolus;

Every Man

6

to be repeated every fix hours till the pains abate. Cover the woman up warm, and let her drink plentifully of caudle.

Lying-inn women fhould on no account be permitted to quit their beds before the end of the first fortnight after delivery, fince nothing fo effectually prevents a milk fever and other dangerous diforders (at the fame time that the neceffary cleanfings are thereby encouraged) as warmth and a fpare diet, with drinking plentifully of warm diluting liquors. Getting up too foon is always productive of bad confequences.

Heister & Smellie.

AGUE, OR INTERMITTING FEVER.

Signs of.

A violent head-ache, a wearinefs of the limbs, a pain in the loins, a coldnefs of the external parts, a fhivering and fhaking, fometimes fo much as to make the very bed fhake under them; a great thirft and burning heat, and frequent inclination to vomit. Afterwards the fymptoms begin to abate, the fkin becomes moift, the urine is of a flame col-

7

our, but without a fediment, and a fweat breaking out terminates the fit.

Method of Cure.

First vomit the fick perfon, by giving half a drachm of the powder of ipecacuanha, and work it off with chamomile tea: then let the fick perfon take the following powder:

Of the beft Peruvian bark, powdered, one ounce; of Virginia fnake-root, and falt of wormwood, each one drachm; mix these well together, and divide them into eight doses, one paper to be taken every two hours in a glass of red wine, or any other liquid. This is a certain and infallible cure; but care must be taken to administer it only in the intervals of the fits; and it must be repeated for two or three days, about ten days after the first cure, or else the diforder will frequently return. In obstinate cafes, removing into a drier air has been found. of great fervice for perfons of robuft constitutions after taking the ipecacuanha vomit. Dr. Mead directs a drachm of powder of myrrh to be taken an hour

Ebery Ban

8

before the ague first comes on, in a glass of fack. Huxbam.

SAINT ANTHONY'S FIRE.

Signs of.

This difeafe affects every part of the body, but most frequently the face, and it happens at all times of the year. But whatever part is affected, a chillness and shivering generally attend the diforder, with great thirst, restless, and other figns of a fever; the face fwells of a fudden, with great pain and redness, and abundance of small pimples appear, which often rife up into small blifters, and spread over the forehead and head, the eyes being quite closed by the largeness of the swelling. This, in the country, is usually called a Blast.

Method of Cure.

Let the fick perfon lofe eight or ten ounces of blood, and repeat the bleeding more than once if the fymptoms run high; apply to the part a poultice of white bread and milk with a little hog's lard in it; let the poultice be changed twice in a day; but flannels wrung out of a ftrong decoction of elder flowers applied warm

9

afford the fpeedieft eafe and relief; and every other morning take the following purge, till the diforder is cured: viz.

Glauber's falt one ounce, manna half an ounce : mix and diffolve it in warm water for one dofe.

The diet in this difeafe must be very low, chiefly water-gruel, or at most weak broth; all strong liquors and flesh meat must be avoided as poison. Sharp.

APOPLEXY.

The immediate preceding Signs are,

Tremblings, ftaggering, a giddinefs in the head, dimnefs of fight, fleepinefs, lofs of memory, noife in the ears, and deep and laborious breathing. The difeafe itfelf is a total deprivation of fenfe and motion, except that of refpiration, which is performed with difficulty and fnoring. It ufually ends in a paralytic diforder, and is feldom curable, but always leaves behind it a great defect of memory, judgment and motion.

Method of Cure.

Cupping in the nape and fides of the neck is always ufeful, provided the fcarifications are deep enough to give a free

Ebery Pan

paffage to the blood ; ftimulating clyfters and warm purges are alfo of fervice, as is the following electuary. Take half an ounce of powdered heath valerian, and one ounce and an half of conferve of orange peel, and mix them together ; the dofe is the quantity of a nutmeg every four hours, diffolved in a cup-full of rofemary tea. Apply a ftrong blifter to the back and the legs. The diet muft be very fparing.

Mead, Sydenham and Heister. APPETITE, Want of.

Drink chamomile tea every day an hour before dinner, or take ten drops of acid elixir of vitriol in a glafs of water, about two hours before, and about two hours after dinner every day.— A gentle puke fhould precede the ufe of thefe remedies, for which purpofe nothing is preferable to the powder of ipecacuanha, a fcruple of which is fufficient for a dofe, and fhould be worked off with weak chamomile tea. Mead. ASTHMA OR PHTHISICK.

Signs of. An obstructed and very laborious

10

breathing, attended with unfpeakable anxiety, and a straitness about the breast. Method of Cure.

Diffolve two drachms of gum ammoniacum in half a pint of pennyroyal water, and add an ounce of oxymel of fquills. Three large fpoonfulls of this mixture may be taken frequently; or from twenty to thirty drops of the paregoric elixir may be taken in pennyroyal water, two or three times in a day. Bleeding is generally proper, as is a large blifter applied to the back or the legs, and gentle vomits. The diet fhould be flender, and malt liquors muft be avoided, being very pernicious. Mead.

BALDNESS.

Rub the part frequently with an onion till it looks red : the ufe of bear's greafe is alfo recommended as particularly ferviceable in this cafe, being more fubtle and penetrating than any other animal fat hitherto known.

BITE OF A MAD DOG.

Dr. Mead's powder, and the Tomquin remedy; for both which receipts fee the appendix.

Ebery Man

BITE OF AN ADDER OR VIPER.

The fat of this animal rubbed well into the part bitten, prevents the ill confequences of fuch a wound. Where this fat cannot be procured, the fame good effects will be produced by bathing the part well with warm fallad oil.

> Philof phical Transactions and Mead. SPITTING OF BLOOD.

Take red rofe leaves dried half an ounce, twenty drops of oil of vitriol, one ounce and an half of refined fugar, and pour two pints and an half of boiling water on thefe ingredients in an earthern veffel; let it ftand to be cold, and take half a gill frequently. In this diforder, frequent bleeding in fmall quantities is proper, not exceeding four, or at the most, fix ounces at each time, according to the ftrength of the fick perfon. Hoffman and Mead.

BLEEDING AT THE NOSE.

Apply to the back part and the fides of the neck, a linen cloth dipt in cold water, in which falt prunella has been diffolved. In very obstinate cases, bleeding in the foot is useful. Internally, the

quantity of a nutmeg of the following electuary may be taken three or four times in a day. Take the feeds of white henbane, and white poppies, each half an ounce; conferve of rofes three ounces; and mix them into an electuary with fyrup of diacodion. Sydenham & Boyle. BLOODY FLUX.

Signs of.

Generally begins with coldnefs and fhivering, fucceeded by a quick pulfe, and intense thirst. The stools are greafy, and fometimes frothy mixed with blood, with filaments intermixt, which have the appearance of melted fuet, and are attended with intolerable gripings, and a painful descent as it were of the bowels.

Method of Cure.

Bleed first, then give the following vomit; half a drachm of powder of ipecacuanha; work it off with chamomile tea; repeat this vomit every other day, for three or four times. On the intermediate days, between each vomit, let the fick perfon take a large fpoonful of the following mucilage, warm, every hour : diffolve half an ounce of gum

Ebery Dan

tragocanth, in a pint of barley water, over a gentle fire. Clifters made of fat mutton broth are of great fervice. The fick perfon must abstain from malt and spirituous liquors. Mead.

BOILS.

Apply a plaifter of diachylon with the gums, once every day, till they are cured. To prevent their return, a few dofes of cooling phyfic are proper.

Wiseman.

BRUISES, INTERNAL.

Take a large fpoonful of cold drawn linfeed oil, two or three times in a day. The patient must also be blooded to the quantity of eight or ten ounces; and if the fymptoms are violent, the bleeding must be repeated at difcretion. Sydenham.

BRUISES, EXTERNAL.

Bathe the part with a little fpirits of wine and camphire, which in flight cafes will effect a cure; but if that fails it will be neceffary to apply a poultice of ftale beer grounds and oatmeal, with a little hog's lard, which must be applied fresh every day till the bruise is entirely cured. Heister,

14

BURNS AND SCALDS.

Take May butter unfalted, and white wax, of each fix ounces; oil of olives, half a pint; lapis calaminaris one ounce and an half; melt the wax and butter with the oil, and ftir in the lapis calaminaris finely powdered, till it is too hard to let it fettle. This is an excellent ointment for the above purpofe, and is to be applied once a day fpread on a fine linen rag. *Turner*.

CANCER.

Signs of.

Begins at the first with a small tumor, about the fize of a nut, which does not change the color of the skin, and sometimes remains for several years without increasing. But as soon as the virulent humor becomes more active, the small hard swelling becomes, all of a sudden, a large, round, livid, unequal tumor, and is attended with an intense shooting pain : at length it breaks, and turns into a sharp stinking fanies, which eating away the found parts, the lips of the wound becomes offensive to the soft, and the patient being worn out with in-

Ebery Man

tolerable pain, at last ensues death. This diforder may infect any part of the body, but most frequently the breast, arm-pits, behind the ears, the lips, nose, and private parts.

Method of Cure.

Boil a quarter of a pound of guaicum, fhavings in fix pints of water, till it comes to four pints : drink a pint of this decoction twice a day, milk warm. External applications moft proper, are poultices of white bread and milk, and fomentations of milk and water ; and the cancer fhould be defended from the cold air, by a piece of fine linen, on which a little of the fpermaceti ointment (the composition of which you will fee in the Appendix) is fpread, and fhould be renewed two or three times in a day. Dr. Love of Greenock.

Dr. Stork of Vienna, greatly recommends the ufe of hemlock in cancerous cafes, and gives feveral furprifing inftances of its fuccefs. The part affected is to be fomented every night and morning with hot flannels wrung out of a decoction of this plant, prepared by boil-

ing four ounces of dried flinking hemlock in a gallon and an half of water, till one third part is boiled away, and is to be afterwards covered with the following poultice. Boil a pint of the hemlock fomentation with as much oatmeal as is neceffary to give it a proper confiftence for a poultice; during the ufe of thefe external applications, the patient must take the extract or infpiffated juice of this plant inwardly, beginning with two grains only night and morning : the quantity is to be increasfed by degrees, till it amounts to the quantity of thirty grains in the twenty-four hours.

CHILBLAINS,

Are fwellings on the hands or feet from violent cold, accompanied with heat, rednefs, pricking pains, and intolerable itching. On the first appearance, bathe them with fnow-water, or hold them over the steam of boiling vinegar; but when they break and become fores, they must be fomented with a fomentation, the receipt of making whereof is given in the Appendix, and a dreffing applied of yellow baselicon mixed with

Ebery Ban

a few drops of fpirits of turpentine and fpread on fine lint; the limb must be kept warm; and, if they are obstinate, a sparing diet must be observed.

Heister.

CHOLICK.

Take two ounces of Daffy's elixir, and repeat it as occasion may require; or half a drachm of powder of rhubarb toasted a little before the fire. Quincey.

COLDS.

Colds may be cured by lying much in bed, by drinking plentifully of warm fack-whey, with a few drops of fpirits of hartshorn in it, or any other warm fmall liquor; living upon puddings, fpoon meats, chickens, &c. and drinking every thing warm. In fhort, it must be treated as a finall fever, with gentle diaphoretics; fuch as half a drachm of the compound powder of contrayerva, taken night and morning; or half an ounce of Mindereus' spirit may be given every night going to reft, drinking a plentiful draught of fack-whey after it. If any cough fhould remain, after using this method a few days, the medicines

directed under the article of Coughs, must be taken.

This is a much more eafy, natural and certain method, than the common practice by balfamics, linctuffes, and the like, which fpoil the ftomach, and deftroy the appetite, and hurt the conftitution. Dr. Cheyne.

CONSUMPTION.

· Signs of.

A dry cough, a difpolition to vomit after eating, an unealy ftraitnefs of the breaft, fpitting of blood, a quicknefs of the pulfe after meals, and flufhing of the cheeks. As the difeafe advances, the matter brought up by the cough, if fpit into the fire, yields an offenfive fmell; if into a glafs of water, it finks to the bottom; profuse night-fweats, loofenefs and wasting away of the whole frame.

Method of Cure.

A milk diet, riding on horfeback, country air, and bleeding frequently in fmall quantities, at each time taking away not more than fix ounces of blood, are the most efficacious remedies in this diftemper; fnails boiled in milk have fometimes been of fervice, as is alfo the Peruvian bark, when it does not occasion a purging.

> Sydenham, Boerhaave and Pringle. CONVULSIONS IN CHILDREN.

Signs of.

The mouth is drawn awry, the eyes are difforted, and the face turns blackifh; the child's fift is clenched, and the globes of the eyes feem immoveable. When thefe fymptoms give way, the child is fleepy till another fit comes on.

Method of Cure.

Give three or four drops of the tincture of wood-foot, or a few hartfhorn drops in a tea-fpoonful of water frequently. A blifter applied to the nape of the neck is alfo ferviceable. Sydenham.

CORNS.

After foaking them a confiderable time in warm water, pare away carefully with a pen-knife the uppermost and hardest furface: then apply a plaister of green wax, or diachylon with the gums, spread on thin leather; repeat this method (which is perfectly fafe) a few times, and it will feldom fail to exitrpate them entirely. Heister.

COSTIVENESS.

Take the fize of a nutmeg of lenitive electuary every morning, or as often as occasion requires. Shaw.

COUGH.

Take oil of fweet almonds and fyrup of balfam of each two ounces; four ounces of barley water, and thirty drops of fpirits of fal volatile; fhake them well together, and take two large fpoonfuls when the cough is troublefome. If this medicine does not remove the cough in a few days, it will be abfolutely neceffary to be blooded. *Pringle*.

DECAYS, SEE CONSUMPTIONS.

DEAFNESS.

Syringe the ears well with fome warm milk and oil : then take a quarter of an ounce of liquid opodeldoch, and as much of oil of almonds; mix them well, and drop a few drops into each ear, ftopping them with a little cotton or wool; repeat this every night going to reft.

Ebery Ban

DELIVERY, to Promote.

Divide one grain of purified opium into two pills, to be taken one fix hours after the other. Or take a fcruple of borax with a few grains of toafted nutmeg made in a bolus with conferve of rofes. *Mead.*

DIABETES.

Signs of.

Is an exceffive difcharge of urine, of the tafte, fmell and colour of honey, attended with an intenfe thirst, and a wasting away of the whole body.

Method of Cure.

Take of the fhavings of faffafras two ounces; guaiacum one ounce; liquorice root three ounces; coriander feeds, bruifed, fix drachms; infufe them cold in one gallon of lime-water for two or three days. The dofe is half a pint three or four times in a day. Or four ounces of alum whey, (which is prepared by boiling four pints of milk over a flow fire, with three drachms of alum, till it is turned into whey) may be taken three times in a day. Mead.

N. B. Lime-water is made by pouring twelve pints of boiling water on a pound of unflacked lime; when it is cold it is fit for ufe.

DROPSY.

Signs of.

Difficulty of breathing, making very fmall quantities of urine, coftivenefs, great thirft, fwelling of the feet and ancles, which, when preft with the fingers, will pit; fwelling of the belly, and falling away of the other parts.

Method of Cure.

Take powder of jalap, cream of tarter and Florentine itis, of each a quarter of an ounce: mix them: the dofe is from thirty to forty grains every other day. On the intermediate days take the quantity of a large nutmeg, every night and morning, of the following electuary. Take two drachms of powdered chamomile flowers, as much ginger, and half the quantity of prepared fteel; make it into an electuary with two ounces of conferve of orange peels. Diuretics fhould alfo be made ufe of. The following is excellent: Take broom afhes half a

Ebery Ban

pound, and steep them in a quart of Rhinish wine cold, strain off the liquor, and take a quarter of a pint three times in a day. Dr. Dover.

Assistances for Drowned Persons. Directions with respect to drowned perfons.

As feveral circumftances may happen to have continued life in fuch an unfortunate fituation, beyond the ufual term, we should always endeavour to afford them the most effectual relief, and not give them up as irrecoverable too foon; fince it has been often known that after the expiration of feveral hours, fuch bodies have fhewn fome apparent tokens of life. First, the wet clothes should be ftripped off, and the body ftrongly rubbed with dry coarfe cloths, and put as foon as poffible into a well heated bed; and this rubbing must be continued a confiderable time together. Some perfon fhould force his own warm breath into the drowned perfon's lungs, and alfo the fmoke of tobacco by means of a pipe or funnel introduced into the mouth, ftopping the fufferer's noftrils close at the same time. If a surgeon can be procur-

25

ed, the jugular vein, or any large vein in the neck, fhould be opened, and ten or twelve ounces of blood taken away. This vein is to be preferred, because it is seldom that any other vein will afford any quantity of blood under these circumstances. The fmoke of tobacco fhould alfo be thrown up into the bowels by the fundament, by introducing the fmall end of a tobacco pipe well lighted up, the bowl of it wrapped up in a paper in which feveral holes are pricked, and through thefe the breath is to be ftrongly forced. Two pipes may be thus lighted and applied, one to the anus and the other to themouth, and at the fame time. The ftrongest volatiles should also be applied to the patient's nostrils. It is useles, and even dangerous, to pour much liquid of any kind into the mouth, as long as the patient fhews no figns of life; but as foon as he difcovers any motion, he should take five or fix spoonfuls of a strong decoction of cardus benedictus fweetened with honey; and, if nothing elfe can be had, fome warm water, with the addition of a little common falt,

Laftly, notwithftanding we difcover fome tokens of life, we fhould not ceafe to continue our affiftance, fince they fometimes irrecoverably expire after thefe first appearances of recovering.

Having pointed out the most effectual means of recovering drowned perfons, I shall just mention those which it is the general cuftom to use in the first hurry. Drowned perfons are fometimes wrapped up in a fheep's fkin immediately flayed off, but this is lefs efficacious than a bed well warmed and long continued rubbing with hot flannels. The method of rolling them in an empty hogshead is dangerous, and lofes a deal of important time. The hanging them up by the feet is alfo dangerous, and fhould be difcontinued; fince the froth, which is one of the causes of their death, is too thick and vifcid to difcharge itself by its own weight; and this posture tends to increafe the over fulnefs of the head and lungs. The covering the perfon with hot ashes, hot fand, or hot falt, has often fucceeded, and may be tried, but fhould not make the 'affiftants lefs attentive to

26

the other methods already directed to be used for the recovery of the drowned perfon. Haller and Tiffot.

DRY BELLY-ACHE, OR NERVOUS CHOLIC. Signs of.

Extreme coftivenefs, a most violent pain in the bowels, coldnefs of the hands and feet, trembling, extreme anxiety, and a difposition to fainting, which is frequently fucceeded by the palfy.

Method of Cure.

Give frequently the following clyfter. Take dryed mallow leaves an ounce; chamomile flowers, and fweet fennel feeds, of each half an ounce; water, a pint; boil it for ufe. Take half a pint of this decoction, and add two fpoonfuls of fweet-oil, and half an ounce of Epfom falt; mix it for a clyfter to be repeated frequently. The warm bath is of the utmost fervice in this diforder, as is alfo balfam o Peru given inwardly from twenty to forty drops in a fpoonful of powdered loaf fugar, three or four times in a day.

Riverius and Sydenham.

Ebery Man.

EAR-ACHE.

The fmoke of tobacco blown into the ear is excellent. Hoffman.

EXCORIATIONS in Children.

Diffolve a little white vitriol in fpring water, and dab the part with it, which will heal them prefently. Heister,

DISORDERS OF THE EYES.

An excellent eye-water. Take two grains of fugar of lead, diffolve it in a quarter of a pint of fpring water.

AN OINTMENT for the EYES.

Dip a feather in a little ointment of tutty, and gently rub it across the eyes every night going to fleep. *Turner*.

FAINTING.

Apply to the noftrils and temples fome fpirits of falarmoniac, an give a few drops in a wine glafs of water inwardly. Hoffman.

FALLING SICKNESS.

Signs of.

A wearinefs of the whole body, an heavy pain of the head, unquiet fleep, dimnefs of fight, a noife in the ears, a violent palpitation of the heart, a coldnefs of the joints, and a fenfe as it were, of

a cold air afcending from the extreme parts to the heart and brain; then they fall fuddenly on the ground, the thumbs are fhut up clofe in the palms of the hands, the eyes are difforted and all fenfation is loft, fo that by no noife, nor even by pinching the body, can they be brought to themfelves: they alfo froth at the mouth.

Method of Cure.

Take of Peruvian bark powdered one ounce, of wild valerian root half an ounce, of fyrup of orange peel a fufficient quantity; make an electuary of this, and let the patient take the quantity of a nutmeg, (after proper evacuations, fuch as bleeding and purging) morning and evening, for three months together, and then repeat it three or four days before the new and full moon. Mead.

FEVERS INFLAMMATORY.

Signs of.

The breath is very hot, and there is a drynefs of the whole fkin, and fenfe of heat; the refpiration is thick, difficult and quick; the tongue is dry, yellow, parched and rough; the thirft is un-

Ebery Man

quenchable; a vaft anxiety, reftlefsnefs, and wearinefs, frequent inclinations to vomit, and fometimes light-headednefs and convultions.

Method of Cure.

Bleed to the amount of ten or twelve ounces, vomit with half a drachm of ipecacuanha, work it off with chamomile tea; if coftive, inject, as often as occafion requires, the clyfter directed under the article of the Dry Belly-Ache, and give inwardly the following mixture. Take of falt of wormwood half a drachm, lemon-juice three quarters of an ounce, falt of prunella ten grains, fpring water one ounce; mix them together for one dofe, and repeat it every fix hours.

Huxman.

Nervous Fever. Signs of.

Slight chilineffes often in a day, with uncertein flufhes of heat, a vaft dejection and anxiety of the fpirits, a giddinefs and pain of the head, an inclination to yawn and dofe, a drynefs of the lips and tongue without any confiderable thirft; the countenance heavy, pale and deject-

ed ; frequent fick fits ; the urine is pale and made often and fuddenly ; the pulfe is low, quick and unequal.

Method of Cure.

A vomit is neceffary when the ficknefs and load of the ftomach is urgent; if the body is coffive, clifters of milk, fugar and falt, may be injected every fecond or third day. Blifters must be applied to the nape of the neck, head and legs: the fick perfon must be kept quiet in body and mind; opiates are commonly hurtful; a little wine whey is proper for common drink. Give the following draught every fix hours. Take falt of hartshorn ten grains, lemon juice half an ounce, fimple wine water one ounce and an half, compound fpirits of lavender and fyrup of faffron, of each one drachm and an half, mix them together-for one dofe. Huxbam.

PUTRID OR SPOTTED FEVERS.

Signs of.

The head aches and is hot, dull, and attended with a dejection of mind from the beginning ; a conftant watchfulnefs; the countenance is dejected ; the pulfe is languid, fmall and low. A pain in the back and loins, a great load at the pit of the ftomach, a perpetual vomiting of black bile: the thirft is commonly very great, and all drinks feem bitter and maukifh : the tongue at the beginning is white, but grows daily more dark and dry, with a kind of dark bubble on the top, and livid or brown fpots appear over the whole furface of the fkin.

Method of Cure.

Gentle vomits are neceffary in the beginning; and, if the body is too coftive, a clyfter of milk, fugar and falt, may be given as often as occafion requires; wine diluted with water, and acidulated with the juice of Seville oranges, may be ufed for common drink; and the following has been found of greater efficacy in this diforder than any other medicine.

Boil three drachms of Peruvian bark in powder, and three drachms of Virginia fnake-root in powder, in a pint of water, till half a pint is boiled away; then add a quarter of a pint of good

red Port wine; the dofe is a tea-cup full every three or four hours. Dr. Pringle. FISTULA.

Take a quarter of a pound of elecampane root, three quarters of a pound of fennel feeds, and a quarter of a pound of black pepper ; pound thefe feparately and fift them through a fine fieve ; take half a pound of honey, and half a pound of powder fugar, melt the honey and the fugar together over the fire, fcumming them continually till they become bright as amber ; when they are cool, mix and knead them into your powder, in the form of a foft pafte. The dofe is the fize of a nutmeg, morning, noon and night, drinking a glafs of wine or water after it.

This is Dr. Ward's receipt for curing Fiftulas; and, if it does not fucceed, immediate recourfe must be had to a skilful furgeon.

GIDDINESS.

Take twenty drops of tincture caftor in a glafs of water two or three times in a day; or from a fcruple to a drachm of powder of valerian root, three or four times in a day: when this diforder proceeds from too great a fulnefs of the veffels of the head, bleeding will certainly give relief. Mead.

Gout.

- Signs of.

A pain refembling that of a diflocated bone, fucceeded by a chillinefs, fhivering and a flight fever, ufually feizes the great toe, heel and calf of the leg or ancle, which becomes fo exquifitely painful as not to endure even the weight of the bed cloaths; the part looks red and is very much fwelled; the urine is high colored, and lets fall a kind of red gravelly fediment.

Method of Cure.

Take a tea-fpoonful of volatile tincture of guaiacum, every night going to reft, in a glafs of water, be covered warm, and drink plentifully of weak fack whey. In the intervals of the fits, exercife, fuch as walking and riding, is abfolutely neceffary to prevent a return; and the duke of Portland's gout powder, the receipt of which we have given in

our Appendix, should be taken according to the directions annexed thereto.

Dr. Cheyne

GRAVEL AND STONE. Signs of.

A pain in the lions, bloody urine, a numbnefs of the thy or leg on the fide affected, a ficknefs at the ftomach, and frequent vomitings.

Method of Cure.

Take one pound of calcined oyfterfhells, and pour thereon twelve pints of boiling water; ftrain it when cold; and take half a pint mixed with a little new milk, three times in a day, gradually increafing the quantity, till it amounts to four pints a day. If the patient is coftive, two ounces of manna diffolved in a quart of whey, fhould be taken for one dofe, once or twice every week.

Dr. Whytt and Sydenham.

GREEN SICKNESS. Signs of.

A pale complexion, fwelling of the ancles, wearinefs of the whole body, difficulty of breathing, a palpitation of the heart, drowfinefs, a defire of eating coals,

Ebery Man

chalk, &c. and a suppression of the monthly courses.

Method of Cure.

Take thirty drops of the tincture of black hellebore, two or three times in a day, in a glafs of water or wine, ufing moderate exercife; or the pills directed under the article of *Obstructions*. Chalybeate waters are also of fervice, and bleeding in the foot about the time of menstruation. *Mead*.

GRIPES.

Take half a drachm of powdered rhubarb, and toaft it a little before the fire, then add a little powder of ginger to it, and mix it for one dofe, to be repeated as occasion requires; or take a wineglafs of Daffy's elixir. Radcliffe.

HEAD-ACH.

Apply leeches behind the ears, and take twenty drops of tincture of cafter in a glafs of water frequently; if this fails, take a fcruple of pil. Rufi every night going to reft, for a week or ten days. Haller.

HEART-BURN.

. Take a little chalk fcraped in a glafs of water, or a tea-cup full of chamomile tea.

36

HOARSENESS.

Take the medicines directed under the article of *Coughs*, or as much as will lie on a fhilling of the following powder, three or four times in a day : take fpermaceti and fugar-candy, of each equal parts, and make them into a fine powder, or a tea-fpoonful of Barbadoes tar in a glafs of old rum, every night going to reft. Three or four Wellfleet oyfters fwallowed early in the morning fafting, afford furprifing relief in this complaint. *Shaw*.

HYSTERICS.

Signs of.

An intercepted breathing, almost to fuffocation, fainting, loss of speech, great drowfines, exceeding costiveness, and making large quantities of urine, almost as clear as water.

Method of Cure.

Take fimple pepper-mint water, twelve ounces; one ounce of powdered valerian, and half an ounce of lavender drops; mix them together, and take three large fpoonfuls two or three times in a day, and also apply to the navel a large galbanum plaister.

Hoffman and Mead.

HARD SWELLINGS.

Apply a plaister of diachylon with the gums, spread thick on leather, or a mercurial plaister. Parey.

HIP-GOUT. Signs of.

A violent pain in the joint of the thigh and lower part of the loins, which fometimes reaches to the leg, and even the extremity of the foot, without any fwelling or change of colour of the fkin.

Method of Cure.

Take a drachm of ætherial oil of terpentine, and three times as much honey mixt with it, every morning, for fix or feven days at fartheft, drinking plentifully of warm fack whey after it.

Cheyne.

HOOPING-COUGH.

Boil a good handful of dried coltsfoot leaves cut fmall, in a pint of fpring water till about one half is boiled away, ftrain the liquor through a cloth and fqueeze

the herbs as dry as you can; diffolve in the liquor an ounce of brown fugarcandy, and give the child (if it be but three or four years old and fo in proportion) a spoonful of it cold or warm, according to the feafon of the year, three or four times in a day, or oftener, till the violence of the diftemper is abated; or the child may take from five to twenty drops, according to its age, of elixir afthmaticum three or four times a day; a gentle vomit should precede the use of either of these remedies; and, in very bad cafes, a blifter applied to the nape of the neck will be neceffary; bleeding is highly improper in this diforder.

Haller.

JAUNDICE.

Signs of.

A yellowithnefs of the whites of the eyes: and of the whole body, bitternefs of the tongue, heavinefs and laffitude, vomiting of galls, the ftools almost white, and the urine of a faffron colour, tinging linen dipt therein yellow.

Method of Cure. Take Venice soap half an ounce, oil

Ebery Ban.

of annifeeds fixteen drops; mix them well together, and make it into middle fized pills. The dofe is three or four, two or three times in a day: if coftive, half a drachm of rhubarb must be taken in the morning twice a week.

Huxbam.

INDIGESTION. Signs of.

Pain and fenfe of weight in the ftomach, attended with frequent belchings, heartburn, &c.

Method of Cure.

Take a large fpoonful of tincture of hiera picra every day, an hour before dinne; or from ten to twenty drops of acid elixir of vitriol, in a glafs of water, two or three times in a day. Pyrmont and fpaw waters are alfo very efficacious in removing this complaint.

Cheyne and Heister.

INFLAMMATIONS.

Take away ten or twelve ounces of blood, and repeat it if neceffary : give cooling purges, and apply to the part a poultice of bread and milk, with fome eintment of elder in it. Sharpe.

Ітсн.

Signs of.

Watery pimples, attended with intolerable itching between the fingers, under the hams, on the arms, and on the thighs.

Method of Cure.

Anoint the parts which brake out every night with the following ointment : take hog's lard, half a pound; flour of brimstone and fulphur vivum, of each two ounces; powdered cloves one ounce; mix them together : two or three weeks is as little time as can be depended upon, and the fame linen must be wore the whole time. This is a never failing remedy. For perfons who are too delicate to bear the fmell of the brimstone. the following ointment is contrived; take an ounce and a half of pomatum; precipitated sulphur, a quarter of an ounce; white precipitate, two scruples; mix them together. During the use of this ointment a dofe of cooling phyfic should be taken every third day, and the linen should be frequently changed.

D 2

Ebery Man

42

High living, rich fauces, &c. must be carefully abstained from.

INFLAMMATIONS OF THE BOWELS.

Signs of.

A burning pain in the belly, attended with a fever, vomiting, and frequently an obftinate coffivenes.

Method of Cure.

Take away ten ounces of blood, and repeat the bleeding according to the urgency of the complaint; give the clyfter directed in our appendix, twice every day, till the patient has had two or three ftools; and, to appeafe the pain, and ftop the exceffive vomiting, give ten drops of liquid laudanum, in a little broth, every fix hours. The diet fhould confift entirely of weak broths.

Juice of lemons taken inwardly in obftinate cafes, has fometimes afforded furprifing relief. Arbuthnot.

INTERMITTING FEVER, SEE AGUE.

KING'S EVIL.

Signs of.

Hard fwellings in the neck, arm-pits and groin, and tumours on the joints and fingers, attended with a fwelling,

rottennefs of the bones, and forenefs of the eye-lids. In fhort, there is no part of the human body which may not be affected with tumours, abfceffes, or ulcers, by this difeafe.

Method of Cure.

Take a fcruple of burnt fponge, and four grains of rhubarb; mix them together for one dofe, which is to be taken every night and morning, with a draught of whey. Drinking fea-water is very efficacious in curing this difeafe.

Sir Edward Hulfe.

LEPROSY.

Signs of.

Red pimples, or poftules, commonly firft break out in the elbows and knees, which gradually increase in fuch a manner, that the whole body is covered with a leprous fcurf.

Method of Cure.

Take fharp pointed dock roots, half a pound : monks rhubarb, a quarter of a pound ; ftick licorice and corinder feeds, of each half an ounce ; boil them in a gallon of fpring water till it comes to three quarts, and to the ftrained liquor add two ounces of diophoretic antimony. The dofe is a pint every morning and evening. Or take twenty drops of the antimonial wine three times a day.

Banyer.

LETHARGY, SEE APOPLEXY. LIMBS FROZEN.

In very fharp winters, it is too common for perfons to be pierced with fo vlolent a degree of cold, that their hands or feet, or fometimes both, are frozen at once, like a piece of flesh meat exposed to the air. If a perfon in this fituation attempts to warm the frozen parts before a fire, his cafe proves irrecoverable; intolerable pains, and at length a mortification, are the confequence of his imprudence. The only certain remedy is to convey the perfon affected in fome place where it does not freeze, but where, however, it is but very moderately warm, and there continually to apply to the frozen parts fnow, if it can be procured; if not, to keep washing them incessantly; but very gently, with ice-water, as the ice thaws in the chamber : by this application the patient will be very fenfible

of his feelings returning very gradually to the part, and of their beginning to recover their motion. In this ftate the perfon may be fafely removed into a place a little warmer, and drink fome balm tea, or any other mild diluting liquid. Many people have been revived, who had remained in the fnow, or expofed to the freezing air for feveral days, and who have difcovered no one fign of life for feveral hours : we fhould therefore, ufe our utmost endeavours for the recovery of perfons in the like fituation, by using the above method.

LOOSENESS.

Take half a drachm of ipecacuanha for a vomit; and work it off with chamomile tea. Abstain from malt liquors. If this does not entirely remove the loofenefs, take half a drachm of powder of rhubarb, made into a bolus, with a little diacordium, and repeat it every day, till the loofeness is intirely stopped.

For common drink, the fick perfon may take hartfhorn drink, or rice boiled in water with a little cinnamon.

45

Towne.

MADNESS. Signs of.

A rednefs of the eyes, grinding of the teeth, ftrange malice to particular perfons, want of fleep, finging in the ears, incredible ftrength, infenfibility of cold, and an exceffive rage when provoked to anger.

Method of Cure.

Bleeding is ufeful in this diforder, and take forty drops of tincture of black hellebore in a glafs of water, two or three times a day; or half a drachm of champhire may be taken every night going to reft. Mead and Kinneir.

MEASLES.

Signs of.

A chillnefs and fhivering, which is fucceeded by a fever, accompanied with great ficknefs, thirft, a continual drowfinefs, flight cough, and an effufion of tears, which is the most certain fign of the measure. About the fourth day little red fpots like flea-bites appear over the whole body, which do not rife above the furface of the skin.

Method of Cure.

First, bleed the fick perfon, then let him or her drink plentifully of the following decoction : take pearl-barley, raifins and figs, of each two ounces; ftick licorice bruifed, half an ounce; boil them in four quarts of water till it comes to two quarts; ftrain it for ufe, and add a quarter of an ounce of falt prunella. You must purge often after this diforder, and the diet and management must be the fame as in the fmall pox. Sydenbam.

MENSTRUAL DISCHARGE, Excessive.

Signs of. Lofs of ftrength and appetite, bad habit of body, fallow complexion, and fwelling of the feet.

Method of Cure.

Bleed according to the ftrength of the patient, and give thirty grains of the following powder in a glafs of red wine three times in a day. Take three quarters of an ounce of rock allum, and a quarter of an ounce of dragon's blood, and mix them together. Mead.

MENSTRUAL DISCHARGE, Suppression of. Signs of.

A fwelling of the belly, a pain in the loins and groin, difficulty of breathing, cold fweats, frequent faintings, and fometimes hyfteric fits.

Method of Cure.

Bleed in the foot, ufe exercife, and take the quantity of a nutmeg, three times a day, of this electuary: Take conferve of dried orange-peel, a quarter of a pound; candied ginger, half an ounce; powder of fteel, two ounces; mix them into the confiftence of an electuary with any fyrup; or a teafpoonful of tincture of black hellebore may be taken in a glafs of warm water twice in a day. Mead.

MISCARRIAGE.

To prevent mifcarriage, bleeding is ufeful and neceffary, about the third month of pregnancy, more or lefs, according to the conftitution of the woman with child. The body fhould alfo be kept open with manna or rhubarb, in the first months especially; and violent exercise must be shunned, and the passions must be kept under. Hamilton.

49

MORTIFICATION, OF GANGRENE.

Signs of.

The pain and inflammation abate, the parts which were before fwelled and tight, become foft and flaccid; and upon preffing with the finger, its impreffion remains: blifters rife up, like those of burns, filled with a yellowish, reddish, or black humour. The limb loses its feeling and power of motion; the colour of the part turns black, and becomes intolerably offensive.

Method of Cure.

Foment the part every night and morning with hot flannels wrung out of the following fomentation; take limewater a pint, and diffolve it in half an ounce of crude fal armoniac; then add three ounces of camphorated fpirits of wine. Afterwards apply a poultice of ftale beer grounds and oatmeal, moiftened with a little hog's lard: when the part begins to fuppurate, apply under the poultice a dreffing of black bafilicon. Inwardly take a drachm of the beft

Every Man

Peruvian bark in fine powder, every four hours, in a gill of mountain wine. Profeffor Monroe and Heifter.

NERVOUS DISORDERS.

Country air, exercife and the cold bath, with the mixture directed under the article of hyfterics, will prove a certain cure for every fpecies of these diforders. *Fuller*.

Noise in the Ears,

May be cured by the vapour of a decoction of rofemary flowers prepared with wine, being conveyed to the ear by a funnel. *Heister*.

A SAFE OPIATE FOR A GROWN PERSON.

Take twenty drops of liquid laudanum in a glafs of wine and water going to reft. The quantity muft be proportionably leffened for children, according to their ages. Boerbaave.

OBSTRUCTIONS IN EITHER SEX.

Take fpecies hiera picra half an ounce, falt of steel one drachm, made into pills of a moderate size with a little syrup of faffron: the dose is five pills twice a day. Shaw.

OLD ULCERS.

When an ulcer is of long standing, it is dangerous to dry it up, without subftituting in the place of a difcharge, which has become almost natural, fome other ; fuch as purging from time to time, or cutting an iffue near the difeafed part. To forward the cure, falt meats, fpices and ftrong liquors muft be most avoided; the usual quantity of flesh meat should be leffened, and the body kept moderately open by a vegetable or milk diet; and if the ulcers are in the legs, it is of great importance to keep in a lying posture; for negligence in this material point, changes the flightest wounds into ulcers, and the most trifling ulcers into obstinate and incurable ones.

Method of Cure.

Take a quarter of a pound of bafilicon, and an ounce and an half of oil of olives, and mix therewith half an ounce of verdigreafe; drefs the fore with this ointment, fpread upon a little tow, after fomenting it well with a decoction made of chamomile flowers and mallow leaves; or the fomentaion directed in the Appendix. I repeat again, take frequently a dofe of cooling physic, and live regularly. Sharpe.

DISORDERS OF THE BREASTS.

If there is a tumour and inflammation of the breafts after delivery, from the ftagnation of the milk, it is proper to apply brandy or rum, hot, with linen cloths; or a white diachylon plaifter, with a little champhire mixed with it; and the milk fhould be drawn by the nurfe.

If the ftagnation of the milk does not occafion great heat, the breafts may be bathed with warm red wine, fpirits of wine and camphire, or a poultice of white bread and red wine, to which add a little fpirits of wine and champhire.

If the breaft feems likely to break, apply a poultice of white bread and milk, or a plaifter of diachylon with the gums, fpread thick on leather. When the fwelling breaks, the wound muft be dreffed with yellow bafilicon, every night and morning, till the diforder is cured. *Theobald*.

53

CHAPT NIPPLES.

Anoint them with oil and bees-wax, equal parts of each; or dab them with a litle Hungary water. Turner.

MILK FEVER.

This diforder happens about the third or fourth day after delivery, and is owing to the congestion of milk in the breasts; it is attended with shivering fits like an ague, tension and swelling in the breasts, that extends to the arm-pits, which parts are often extremely painful. This fever continues a day or two, and sometimes goes off of itself, by the benefit of nature, in plentiful sweats proportionably to the preceding cold fits.

Method of Cure.

The patient fhould drink plentifully of warm, thin diluting liquors, as balmtea, barley-water, &c. Her diet fhould be very flender, and the child fhould be often put to the breaft. Internally the following powder fhould be taken every fix hours, in a cup of balm-tea: take compound powder of contrayerva, twenty-five grains, and mix them together for one dofe.

Ebery Man

54

But if the woman does not intend to give fuck, she must rigourously adhere to a very flender diet, and the foregoing medicine : the breafts should always be drawn by fome proper perfon; and to prevent the milk from coagulating, the breafts must be imbrocated with warm fallad-oil, or the leaf of a red cabbage may be applied thereto. If there is reafon to apprehend an inflammation of the breafts, it will be abfolutely neceffary to lofe eight ounces of blood from the Folded cloths dipt in brandy or arm. vinegar, placed under the arm-pits, are alfo of fervice to drive back the milk. Aftruc and Smellie.

PAIN IN THE STOMACH, SEE HEART-BURN. PAINS IN THE JOINTS, SEE HIP-GOUT.

PILES. Signs of.

A violent pain at the time of going to ftool, the excrement ftreaked with blood, and fmall fwellings like warts on the verge of the anus.

Method of Cure. .

Take of linitive electuary one ounce and an half, flour of brimftone half an

ounce, mix them together; the quantity of a nutmeg to be taken every night and morning; apply outwardly a little of this ointment; take two ounces of white diachylon, two ounces of fweet oil, and half an ounce of vinegar, mix them together. Leeches applied as near the piles as poffible, or on the pile itfelf, afford furprifing relief. Sydenham,

PALSY.

Perpetual blifters are ferviceable and the following drops have frequently afforded great benefit : Take fal volatile drops half an ounce ; lavender drops and tincture of cafter, a quarter of an ounce each ; mix them together : the dofe is forty drops frequently in a glafs of wine and water ; or half a drachm of wild valerian root in powder, may be taken three times in a day. *Allen*.

The diet fhould be warm and attenuating.

PLEURISY. Signs of.

A violent pain in the fide, accompanied with heat, thirst, and other usual symptoms of a fever, and a troublesome

Every Pan

56

cough; if the fick perfon is bled, the blood, when cold, looks like melted tallow.

Method of Cure.

Bleed frequently till the pain abates; apply a blifter to the fide, and take half a pint of the following infusion twice a day: Take fresh horse-dung fix ounces, and pour on it a quart of boiling pennyroyal water; strain it when cold, and add a quarter of an ounce of Venice treacle; mix it for use.

Sydenham and Quincey.

POISON TO EXPEL.

Milk mixed with fallad oil, or either of them alone, drank in large quantities, fo as to occafion a plentiful difcharge by vomit, yield the most certain relief, and therefore should be administered to perfons who are fuspected to have taken any kind of poison, as foon as possible, and must be continued to be drank plentifully, till the violence of the fymptoms are abated, and the fooner it is given, the

QUINSEY OR SORE THROAT. Signs of.

A fwelling in the parts concerned in deglution, accompanied with great pain inflammation, and a fever, fo as to hinder the fwallowing any folid meats, and almost ftop breathing.

Method of Cure.

Bleeding is fometimes neceffary, and cooling phyfic, but often jelly of black currants, fwallowed down leifurly in fmall quantities, effects a cure, without the affiftance of any other medicine.

> Philosophical Transactions. RHEUMATISM.

Signs of.

It begins with chillnefs and fhivering, heavinefs of the joints, and coldnefs of the extreme parts. The appetite is loft, the body is coftive and feverifh, and a racking pain is felt fometimes in one part, fometimes in another; frequently fhifting from place to place, and leaving a rednefs and fwelling in the part vifited laft.

Method of Cure. Take half a drachm of powder of gum

guaicum in a draught of warm ale going to reft, and be covered with a larger quantity of clothes than ufual; perfift in this method a few days, and you will find relief. Bleeding is of fervice in phlethoric conftitutions. *Pringle*.

RICKETS IN CHILDREN.

Signs of.

A fwelling of the belly, and conftant defire of fitting ftill; the bones are crooked, and the joints feem very large, as if knotted; the head is over large, and the child's underftanding exceeds its years.

Method of Cure.

Give the child two grains of ens veneris, diffolved in a fpoonful of wine and water, every night; to this must be joined cold bathing, frictions of the back, exercise, and a strengthening diet.

Boerhaave.

RETENTION OF URINE.

Take a quarter of an ounce of nitre, a drachm of volatile falt of amber, a quarter of an ounce of powder of egg-fhells, and half an ounce of fine fugar; mix them together: the dofe is as much as

will lie on a shilling, two or three times . a day, drinking a draught of marsh-mallow tea after each dose. *Turner*.

RUPTURES,

Are cured in children and young perfons by applying a ftrengthening-plaifter fpread on leather, (the composition of which is fet down in our appendix) and a trufs) which may be had at any trufsmaker's.

SCALD-HEAD.

First shave, then cover the head with a pitch plaister spread on leather, and give inwardly one grain of calomel, made into a pill, with a little conferve of roses, every night going to rest, and a dose of cooling physic twice in a week.

Wifeman.

STRANGURY.

Take half a drachm of camphire, a fcruple of powdered gum arabic, liquid laudanum five drops, with a fufficient quantity of turpentine to make it into a bolus, to be taken night and morning, drinking a draught of marsh-mallow tea after each dose.

Turner.

Ebery Dan

60

STITCH IN THE SIDE, SEE PLEURISY.

Scurvy. Signs of.

A fallow complexion, difficulty of breathing after exercife, fpongy gums fubject to bleed with the flighteft touch, frequent bleeding at the nofe, a fwelling of the legs, accompanied with livid fpots, and unufual lazinefs, and pains over the whole body, refembling the rheumatifm.

Method of Cure.

Take one ounce of acid elixir of vitriol, and one ounce of tincture of Peruvian bark, mix them together; a teafpoonful is to be taken twice in a day, in a glafs of water. A decoction of the tops of the fpruce fir may be taken, the quantity of half a pint, twice in a day. Bathing in the fea, and drinking feawater, are of infinite fervice.

Heifter and Ruffel.

SLOW FEVER, SEE NERVOUS FEVER.

SMALL-POX.

Signs of. A chillnefs, fhivering, and other ufual fymptoms of a fever, attended with a vi-

olent pain in the head and back, and an inclination to vomit, and great propenfity to fweat. In children, a dulnefs and drowfinefs, and fometimes epileptic fits; about the third or fourth day red fpots about the fize of a pin's head appear on the breaft, face, and whole body, which increafe in fize every day, till they are of the fize of a large pea, and are filled with matter.

Method of Cure.

The fick perfon must be kept in bed, taking care to defend him or her from the inclemency of the winter, by proper means, and to moderate the exceffive heat in fummer by cool air, for the patient ought not to be ftified with heat and cloaths, nor fhould the eruption and perfpiration be checked by cold. With regard to diet, it ought to be very flender, moiftening and cooling; fuch as watergruel, pipping-liquor and milk pottage : and in the beginning the best regimen is that which keeps the body open, and promotes urine; as figs, damascene prunes and tamarinds; and by giving fubacid liquors for drink, as finall beer

Ð

fharpened with orange or lemon-juice; whey made with apples boiled in milk, emulfions made with barley-water and almonds, or Rhenifh wine plentifully lowered with water. Repeated purging is neceffary after this diforder as well as after the meafles. Mead & Sydenham.

SHINGLES. Signs of.

An eruption of watery puftules, about the fize of millet feeds, which encircle the body like a belt, of an hand's breath, and occafion an intolerable itching and forenefs.

Method of Cure.

Take two ounces of white diachylon, two ounces of fweet oil, and half an ounce of vinegar; mix them together for a liniment, which fpread on a piece of fine linnen and apply to the part affected, repeating it as occasion requires; this with two or three doses of Glauber's falt, and a cooling and spare diet, will entirely remove this complaint. *Turner*.

STRAINS.

After fomenting with warm vinegar, apply a poultice of stale beer grounds,

and oatmeal, with a little hog's lard, every day till the pain and fwelling are abated; then apply the strengthening plaister directed in our Appendix. Obferving the following rules, will much shorten the cure : let the perfon ftand three or four minutes on both his feet, and fometimes move the strained foot; and when fitting with his foot on a low ftool, let him move it this way or that as he can bear it; let the strained part be gently rubbed with a warm hand feveral times in a day, which will contribute very much to contract the over-stretched veffels, and recover a due circulation of their fluids through them. Sharpe.

TEETH, Difficulty Cutting of.

If the child is coftive, a few grains of rhubarb mixed with an equal qantity of magnefia alba, according to its age and ftrength, muft be given every night in a little pap; hartfhorn-drops in the quantity of three or four drops in a fpoonful of water, are proper three or four times in a day: if this method does not give relief, the gums muft be lanced to give

Ebery Ban

64

opportunity, for the tooth to make its way. Sydenham & Harris.

STONE, SEE GRAVEL.

SWEATING, Exceffive.

Take twenty drops of acid elixir of vitriol, in a glafs of water, two or three times a day. Shaw.

Swelled Legs.

Take the fame medicines which are advifed under the article of the dropfy. TOOTH-ACHE.

The root of yellow water flower-deluce rubbed on the tooth which is painful, or chewed in the mouth, in an inftant, as if by a charm, drives away the pains of the teeth, arifing from what caufe foever. Or take champhire and opium of each as much as a large pin's head, and put it in the hollow tooth ; a blifter may alfo be appled behind the ear. Allen.

THROAT SORE, SEE QUINSEY.

PUTRID SORE THROAT.

Signs of.

A giddinefs, chillnefs, and fhivering, like that of an ague fit, followed by an intenfe heat, a violent head-ach, heat and

forencfs in the throat, fliffnefs of the neck, great ficknefs, vomiting and purging, or both; the face looks red and fwelled, the eyes inflamed and watery, with reftleffnefs, anxiety and faintnefs: frequently a great number of fmall pimples appear on the neck, breaft, and hands, which are fenfibly fwelled; the infide of the mouth and throat are fwelled, and appear of a florid red colour, fometimes of a pale white, furrounded with red.

Method of Cure.

All evacuations which leffen the ftrength, particularly bleeding and purging, and cooling medicines, are highly prejudicial. The hot fteam of a boiling mixture of vinegar, myrrh and honey, is to be received into the throat, through an inverted funnel; this cannot be ufed too frequently. The decoction of the Peruvian bark, made by boiling an ounce of bark in a pint and an half of water till it comes to half a pint, and adding a drachm of acid elixir of vitriol, is the

Ebery Man

best medicine in this diforder, if given, a tea cup full, evey four hours.

Huxham & Forthergill. TETTARS.

Take four ounces of fweet oil, one ounce of white wax, one ounce fpermaceti, and twenty-five grains of camphire; mix them together for a liniment, with which gently touch the tettars three or four times in a day.

THIRST, Ungenchable.

A quart of water mixed with an ounce or two of white wine vinegar will make an agreeable drink, and has been known to extinguish the most violent thirst after other liquors have in vain been tried for that purpose. James.

THRUSH.

Signs of.

Little white ulcers affect the lips, gums, cheeks, tongue, palate, and the whole infide of the mouth.

Method of Cure.

Rub the child's mouth with a linen rag dipt in the following mixture; take honey of rofes an ounce, oil of vitriol fix drops, mix them together; or the

child's mouth may be frequently washed with a decoction of elm bark.

Shaw & Allen.

TYMPANY.

Signs of.

A great fwelling of the belly, which being occasioned by air pent up in the cavity of the belly, gives an hollow found when struck by the hand.

Method of Cure.

The fame medicines which are directed for the dropfy are also of fervice in this diforder. Boerbaave.

VIOLENT PAINS AFTER DELIVERY, See After-Pains.

VOMITING.

Take mint water one ounce, lemonjuice half an ounce, a fcruple of falt of wormwood, mix them for one dofe, to be repeated as occasion may require. Mead.

ULCER OF THE LUNGS,

Requires the fame treatment as is directed under the article of Confumptions. Fuller, Pringle, &c. ULCER OF THE KIDNEYS.

Signs of.

Fleshy excrescences are voided by

urine. The heat and pain in making water comes by intervals; the matter that comes from the kidneys is more plentiful, white and thin, than in an ulcer of the bladder, and is not fetid; the urine looks like milk when first made, but after standing fome hours, the matter feparates from it, and falls to the bottom.

Method of Cure.

Butter-milk not very four, is extremely beneficial, as is alfo an emulfion made of barley-water and fweet almonds. Chalybeate waters have been fometimes of fervice. *Arbuthnot.*

ULCER OF THE BLADDER.

Signs of.

A voiding of finking matter or blood, and fometimes fcales, or a membranous fkin along with the urine, and a continual heat and pains in the urinary paffages.

Method of Cure.

Selter's mineral water mixed with milk are excellent, as is also the constant use of spaw-water. Hoffman.

ULCER OF THE WOMB.

Sings of.

A flux of purulent matter, or matter

mixt wih blood, which is more vifeid and fetid than the whites, and attended with a fixed pain.

Method of Cure.

An injection, composed of barleywater, a pint, and honey of roses two ounces, must be used to the part affected, by the help of a syringe, several times in a day. You may add a small quantity of tincture of myrrh. *Astrue*.

WARTS.

May be removed by rubbing them with the juice of celandine, or milk of fpurge; but when they are fituated about the eye-lids to prevent hurting the eyes, it is proper to furround the wart with a wring of wax, or a piece of plafter with a mole in the middle fo that the wart may mome through, by which means the warts will be deftroyed, without hurting my other part. Heifter.

WATERY GRIPES IN CHILDREN.

Take half a drachm of magnefia alba, and half a drachm of rhubarb, mix them sogether, and give the child three or four rains in its pap every morning and vening. Or take a fpoonful of hemp feed,

Ebery Ban

and boil it in a pint of water, fweetened with fugar. This will also cure the cholic in grown perfons. Cadogan.

WHITES. Signs of.

A flux of matter from the womb, attended with a pain and weight in the loins, lofs of appetite, pale complexion, difficulty of breathing, a liablenefs to mifcarriage, and fwelling of the eyes and feet.

Method of Cure.

Apply a large ftrengthening plaister to the small of the back, and take a quarter of a pint of this decoction, every night and morning : Take cow's milk half a pint, and boil it in one handful of archangel flowers, and a bit of cinnamon: strain it for use. In obstinate cases, bathing in the sea, and drinking sea water, is far preferable to any thing elfe. Russed

WIND.

Take of the powder of angelica feeds, carraway feeds and parfley feeds, of each a quarter of an ounce; make them into an electuary with the fyrup of clove

71

gilliflowers; the dose is the quantity of . a nutmeg at discretion. Hoffman.

WHITE SWELLINGS OF THE JOINTS.

Anoint the part with Barbadoes tar, before a good fire, two or three times in a day, covering it with a bladder, and drink half a pint of fea water very morning fafting. Guy's Hospital.

WASTING AWAY OF CHILDREN.

Take falt of tartar, nitre and arcanum duplicatum, of each a quarter of an ounce; falarmoniac thee drachms; mix them together; a very little of this muft be put into the child's drink, according to its age. *Heister*.

WHITLOES.

This diforder begins with a flow heavy pain; but in a little time, the pain, heat and throbbing becomes intolerable; the part fwells and grows red, and fometimes the whole hand is inflamed and fwelled: on its first appearance the finger should be plunged in hot water or held over the steam of boiling water; and by doing this frequently for the first day, a cure has been obtained; but

Every Man

when the diforder is further advanced, a poultice of white bread and milk muft be applied.

WORMS.

Signs of.

Palenefs of the countenance, itching of the nofe, voracioufnefs, ftartings, and grinding the teeth in fleep, loofeneffes, ftinking breath, hard fwelled belly, and fometimes epileptic fits.

Method of Cure.

Take burnt hartihorn, and the beft feamony in powder, of each a quarter of an ounce; calomel one drachm; powder, of tin a quarter of an ounce; mix them together, and let the child take eight or ten grains according to its age, every other morning, in a tea-fpoonful of roafted apple. *Alfon*.

GREEN WOUNDS.

Drefs them every day, with yellow bafilicon fpread on fine lint, after fomenting them well with the fomentation, the receipt for which is inferted in the appendix. *Turner*.

Directions for preferving Health, and attaining Long Life.

EALTH has been ever effeemed the first of bleffings, and confequently every endeavour towards its prefervation, deferves encouragement. This confideration induces me to publish the following remarks, from a thorough conviction of the truth they contain; which ought with every honest man, to outweigh every timidity of its reception, and give him courage to offer, at all risques, what by well grounded experience, appears to him, (from its nature) of general utility to mankind.

The air, which is a fluid elaftic fubstance that furrounds us on all fides, penetrates our bodies, and yet is fo fine, that it efcapes the fight, is rarified by heat, and condenfed by cold : it is fo neceffary that an animal cannot live a moment without it ; it ferves for refpiration or breathing, and is fufceptible of different qualities; it may be hot, moift, cold, dry, ferene, pure, and temperate. It is fubject to variations more or lefs fudden, and to be mixed with impure, corrupted, infectious vapours, which are prejudicial to health. The fudden changes of the air are dangerous, whence proceed a great number of difeafes which reign in the fpring and autumn. Towards the approach of winter, hofpitals, prifons, places where armies are encamped, places where led is melted and the earth just

thrown up near dunghills, &c. are unhealthy, on account of bad exhalations. Too hot an air occasions malignant and putrid fevers : a cold and moift conftitution of the air produces coughs, pleurifies, rheumatifms, agues, &c. for inftance, agues are common in the Fens of Cambridge and Linconshire, and the Hundreds of Effex, on account of their marshy fituation, and the moist vapours which weaken the fibres, of the body and obstruct the pores of the fkin, and confequently diminish the discharge of the skin, called infensible perspiration. Hence it appears, that to preferve health, dunghills should never be placed too near the dwelling-house, fince the corrupted vapours which are continually exhaling from them, cannot fail being in time prejudicial, and caufing malignant fevers; for though those who are used to them, do not perceive their offensiveness, yet the cause does not cease its unwholesome activity.

From what has been obferved with refpect to hofpitals and jails, the neceffity of daily opening the windows of a bed chamber muft clearly appeor, to prevent the bad confequences which arife from foul and confined air; and why houfes fhould be built rather raifed above the level of the earth, than funk beneath its furface; and the exposure of the front fhould be of the fouth-east, this being the most wholefome afpect.

Our conftitution, from the loss it daily fuftains, requires to be repaired by aliment and

bis own Johylician.

drink ; hence a knowledge of their general kinds and qualities is neceffary, in order to make a proper choice thereof.

The principal and most general food is bread, made of wheat, barley, rye, or oats: wheaten bread is most nourishing; barley is dry; rye and oats laxative. The 'crust is most easy of digestion; the crumb more oily and heavy; though other mealy substances, beans, peas, potatoes, &c. nourish much, but are windy, heavy and viscid, and their constant use is apt to cause obstructions, unless a great deal of exercise is used. Rice is emollient and nourishing; but nuts, almonds and chefnuts although they abound with nutritious particles, are hard of digestion.

Pulpy, tart fruits, which abound with juice, eaten ripe, are refreshing, cooling, quench thirst, and are easy of digestion ; fuch as strawberries, rasberries, currants, mulberries, gooseberries, cherries, apples, pears, apricots, peaches, nectarines, &c. Remark, contrary to the common prejudice which generally prevails, that fruits are hurtful in the bloody flux, and even occasion this diforder ; they may be eaten with great fafety in the bloody flux, and are indeed the real prefervatives against it, as is confirmed by the experience of the most eminent physicians ; the cause of this diforder being an excels of flesh meats, too moilt a state of the air, a fuccession of cold showers to violent heats; and uncleanlinefs. In fact, this diforder has been observ-

Ebery Man

ed lefs frequent, and lefs dangerous, when fruits were plenty and cheap; fo that to efcape this diftemper when it is rife, ripe fruits are to be eaten in plenty, and the quantity of flefh meats are to be leffened confiderably.

Pot herbs and roots are lefs nourifhing than the mealy fubftances. Lettuce, fuccory, endive, forrel, purflain, &c. refrefh, moiften, and are laxative. Artichokes, cellery, creffes, afparagus, parfley, &c. are a little heating. Truffles, mufhrooms, oinions, garlic, pepper, muftard, and the other fpices, heat very much, and are therefore lefs wholefome in particular conftitutions.

Animal food differs very much with regard to its kind, age, manner of living, and fubftance. Fifh nourifh the leaft of all animals. Young animals have the greateft plenty of foft nourifhing juice, but that of the older is most fpirituous and nourifhing. Yet though the juices of old animals are most gelatinous, and agreeable to the tafte, their flefh is the hardeft, and most difficult of digestion, wild animals are light and easier of digestion than tame; their white parts contain a very juicy fubstance, of tender fibres, yield a fost food, and are easy of digestion.

Liquid aliments, are milk, eggs, chocolate, foups, broths, &c. Milk requiring but little preparation in the ftomach, is a good aliment for perfons whofe ftomachs are weak, and children; new-laid eggs are very nourifhing, and eafy of digestion, therefore agree with

exhaufted and old perfons. Chocolate nourifhes greatly, ftrengthen the ftomach, helps digeftion, and foften fharp humours; whence it is proper for weak ftomachs and confumptive perfons. Broths and foups abound with a foft, moiftening, nutritious jelly, whence they are great reftoratives. Meat that is roafted, contains an excellent nourifhing juice the moift parts being diffipated by the fire. Things that are fried, are only proper for ftrong ftomachs.

Drink restores the fluid parts of the body, and helps digeftion; the principal and most falutary is water, of which the foftest is best, which is known by its lathering eafily with foap : but taken in too great quantities it relaxes and weakens the ftomach, and caufes many diforders. Winein moderation strengthens the folids, and promotes digestion; but if used in excess, as well as brandy, rum, and all other fpiritous liquors, hardens the fibres, weakens the nerves, deftroys the appetite, and causes gout, stone, and other chronic difeafes. That malt liquor is the wholefomeft which is the lighteft and cleareft; for then it does not offend the ftomach, but paffes readily through the emunctories or strainers of the body, and particularly by urine ; whence the best beer does not make the head ache nor grows four on the ftomach, nor caufes wind. The making wholefome beer depends greatly on the foftnefs of the water, the proper boiling of the ingredients, and a due working

Every Ban

of the wort; for all thick, muddy, or stale beer, not fufficiently boiled, diforders the head, caufes wind in the ftomach, obstructions, the strangury, asthmas, cholics, and ulcers in the legs. Tea, used in moderation, promotes perfpiration, or the discharge by the fkin, ftrengthens the ftomach, and helps digestion; but that kind of tea which is called hyfon, is improper for perfons who have weak nerves or fubject to hysteric diforders. Coffee, drank after dinner, is thought to quicken digestion, and allay the fumes of wine; but if used in excess, it agitates the blood, caufes watching, and promotes hemorrhages, or bleedings from the nofe or other parts.

Spices, pepper, &c. are pernicious, when ufed to create an appetite; whence made diffes are bad; for the appetite caufed by the quality and difference of the victuals, incites perfons to eat more than the ftomach can well digeft; which caufes indigeftion, and frequently dangerous and fatal diforders. With refpect to diet, the furest method of preferving health, is to live on plain, fimple food, lightly feafoned, in that quantity which by experience nature has been found to require. Perfect digestion is the best rule for regulating a meal, which is known from perfons being more lively and brifk after a meal than before. History furnishes us with many examples of perfons, who by temperance, have lived to a very advanced age, though,

of weakly conftitutions naturally; wherefore those that are defirous of life and health, should imitate their manner of living; fince excession in eating and drinking are both extremely dangerous.

Strong, robust, young people, who use much exercife, ought to eat more than others, and may be free with the groffer kinds of food ; for their ftomachs being ftrong, the lighter foods would digest too eafily, and be diffipated too foon. Perfons of a weak constitution, or who are just recovered from a disease, should use soft, light foods, agreeable to the ftomach. Children, whofe ftomachs are weak, and veffels fine, ought to use a light, thin, foft food, eafy of digestion ; wherefore infants should be fed with a fluid milk, to avoid causing obstructions in their fine, delicate veffels ; confequently the milk of a nurfe newly brought to bed, is more proper for a new-born infant, than the milk of a nurfe who has been delivered five or fix months, becaufe her milk begins to have too great a confiftence. Nurfes should observe an exact diet, and shun violent passions of all kinds, for they difturb digeftion, and communicate their bad effects to the child. When children are weaned, they fhould not be accuftomed to spirituous liquors, or strong food, especially falt or fmoked provisions, becaufe they are hard of digeftion, and yield bad nourishment. Their diet should config of light animal food, taken in a fmall quantity at a time, but often. At all times of life, but efpecially in old age, the conftant ufe of falt and fmoked meats tend to harden and fliffen the folid parts of the body, inftead of affording good nourifhment ; being hard of digeftion ; in old age the fluids are more thick, fecretious, more flow, and the folids harder than in youth ; therefore they require a more foft, moistening diet, eafy of digeftion, and not too much at a time, efpecially of a night.

All great changes ought to be brought about infenfibly, for cuftom is a fecond nature, and an acquired habit is hard to be left off. Many persons enjoy a good state of health when their meat and drink are very indifferent, by being accustomed thereto; and are apt to be fick, when they attempt to change their manner of life. Cuftom confines us to certain hours, but hunger points out the best time of eating. In age, where strength is wanting, and in youth, wherein there is a great diffipation, when much is not eaten at a time, something taken between the fet meals is not improper. It is necessary to observe, that when the ftomach is bad, perfons should not begin to eat again, till the last meal is digested.

Perfons who are much fatigued should rest before eating; and in cases of distress and forrow, the food should be light, and small in quantity, because the stomach is weak at those times.

The ftomach will admit of groffer food in winter than in fummer. Those who eat in a hurry, without much chewing, are fubject to indigestion; for digestion depends in part on well chewing the food, and thereby intimately mixing it with the spittle; for this reason, light foods are most proper for children and old perfons who have lost their teeth.

Vegetables are more difficult of digestion than animal food, therefore improper for weak stomachs. On this account, light food, fuch as veal, lamb, chicken, fish, are the food . which best agree with delicate constitutions. On the contrary, stronger habits are more fubject to be furfeited with tender and young meats, than with beef and mutton; becaufe the degree of heat which converts beef and mutton into the true state it ought to have in the ftomach, carries the tender aliment or lamb, pig, chickens, &c. into a stercorous or excremental state, before it leaves the stomach ; whence fluxes, &c. In fuch ftrong ftomachs, it is neceffary to mix vegetables with the animal food, that the fermentation of the one may oppose the too precipitate digestion of the other by its acidity. Hence it is obvious, that weak ftomachs fhould abstain from vegetables, which require a greater degree of vital heat than even beef or mutton, to be converted into proper nourifhment. This, I hope, will fatisfactorily explain the terms of eafy and difficult digeftion. The

I all all all all all all and and

ftomach, which can bear beer and water, cannot bear roafted pig and water, from too great a power of digestion, not from any deficiency in those faculties; therefore beef and mutton are more eafy of digeftion than lamb, or any of the white meats; and the contrary in weak ftomachs. Surfeits from lobsters, crabs, &c. are of the kind which rife from a too fudden change of these fubstances in the stomach ; therefore they should be always eaten with vinegar or lemon-juice. In fact, butter is no bad preservative against furfeits in this inftance; the oil blending with the alcalious falts, forms a foap of that which would otherwife be a more acrimonious, inflammatory, and offenfive fubstance.

Sleep reftores the firength, and repairs and replaces the wafte which is made by the labour and exercife of the day. The proper time for fleep is the night, when darknefs and filence invite and bring it on; day fleep is lefs refrefhing, exercife and cuftom fhould regulate its duration; fix or feven hours at a time is generally thought fufficient, for exceffive fleeping is attended with great inconveniences; it blunts the fenfes, and renders them lefs fit for the duties of life.

It is beneficial to vary the scenes of life; to be sometimes in the country, sometimes in the town; go to sea, to hunt, to be at rest now and then; but more frequently to use exercise, because a sedentary life brings on

many indifpolitions, and renders the hody weak and unactive ; while, on the other hand, exercise and labor strengthen it. But moderation is to be observed in all these things, and too much fatigue to be avoided ; for too frequent and violent excercise overpowers the natural strength, and wastes the body. Of all kinds of exercise, riding on horseback is the most falutary. I have known many instrength of performs recovering thereby from the most deplorable state, in confumptions, dropsies, cholics, and nervous diforders.

In old age there is feldom fufficient ftrength to use bodily exercise, though so very requifite for health; wherefore frictions with the flesh-brush are necessary, at this time of life, to promote perspiration, which should be done by the person himself if possible.

I have already taken notice, that cold flops the pores of the fkin, and diminifhes both fweat and perfpiration. To avoid this inconvenience, the winter cloathing fhould be put on pretty early in the feafon, and be left off late; befides, care must be taken not to pass too fuddenly from hot into a cold air, and to forbear drinking any thing cold, when the body has been violently heated.

The paffions and affections of the mind, viz. joy, fear, anger, &c. produce very fenfible effects, and, when too much given way to, have a very bad effect on health, for they affect the ftomach, hinder digeftion, and chylification; whence arife many terrible diforders : wherefore it is beft to keep them in bounds as much as poffible, and to preferve an inward ferenity, calmnefs, and tranquility.

Exceffive venery must be avoided, fince the action of coition is very impetuous, and comes near to a convultion. The animal heat is greatly leffened thereby, the habit of body weakened wonderfully, and the whole nervous fystem largely injured.

RULES FOR NURSING SICK PERSONS.

It is a great miftake, to fuppofe that all diftempers are cured by fweating; and that, to procure fweat, fick perfons must take hot medicines, and keep themfelves very hot; for fweating carries off the thinner part of the blood, leaving the remainder more dry, thick and inflamed, which must evidently increase the diforder; for instead of forcing out the watery part of the blood, we fhould rather endeavour to increase it, by drinking freely of barley-water, balm tea, lemonade, or any other diluting liquor made luke warm. What has been already faid on the head of foul confined air, shews the absurdity of stifling the fick perfon with the heat of a close appartment, and a load of bed-cloaths; for these two causes are sufficient alone to produce a fever, even in a healthy perfon. By letting in a little fresh air now and then into a fick perfon's room, and leffening the bedcloaths, you will almost always perceive the fever and oppression in some measure abate.

Inftead of Venice-treacle, faffron, gafcoignpowder, and other heating medicines, in all feverifh diforders, the belly fhould be kept moderately open; whilft those medicines just mentioned render the body costive, and must neceffarily have a bad effect.

Fevers are aggravated by giving the fick perfons food through fear of their dying of weaknefs; which food increafes the diforder, and renders it fatal. This fear is groundlefs; perfons in fevers may be fupported, even for fome weeks, with liquids only, and are ftronger at the expiration of that time than if they had taken more folid nourifhment; for, from the first attack of a fever, whatever folid food is taken, even foup, eggs, bifcuit, &c. corrupts in the ftomach. If a man in perfect health was to eat flinking meat, rotten eggs, four broths, &c. he would be feized with vomiting, load at the ftomach, a purging, fever, and eruptions on the fkin.

The fame articles, even in their foundeft flate, given to a perfon in a fever, are quickly putrified, by the heat and difeafed matter already in his flomach, and in a few hours produce the fame effects. Is it then possible to expect the least fervice from them? No: as long as a fick perfon has a bad humour in his flomach, his weakness increases, in proportion to the food he receives; for this food being corrupted by the infected matter already there, proves incapable of affording the least nourifhment; on the contrary, it

becomes an additional caufe of the diftem-Befides, to heat and cram the fick perper. fon, is wholly opposite to what nature herfelf indicates : the burning heat of which they complain, the dryness of the lips, tongue, and throat, the high colour of their urine, their earnest longing after cooling things, the pleafure and benefit they receive from the admission of fresh air into their chamber, are fo many proofs that we ought to cool them moderately, by refreshing and diluting liquors, fuch as balm-tea, lemonade, &c. to promote an eafy difcharge of the vitiated humours. Those who have the care of fick perfons, fhould particularly attend to this observation, that as long as there is any taste of bitterness, fickness, or defire of reaching; bad breath, heat and feverifhnefs, with offenfive stools, and high-coloured urine made in a small quantity only, fo long all flesh meats, foups, eggs, and all kind of food composed of any of them, and all heating medicines, wine, &c. are fo many abfolute poifons.

If the fick perfon has not two motions for ftool in the twenty-four hours; if the urine is high coloured, the fever runs high, the pain of the head and loins confiderable, a glyfter of warm water, with fweet oil, and a little common falt fhould be given once in a day. As long as the patient has ftrength for it, he fhould fit up out of bed an hour daily, and longer if he can bear it; but he fhould not be raifed whilft in a fweat. His linen fhould

be changed every other day, taking care that the clean linen is well aired; for nothing conduces more to continue the fever and light-headednefs than confining the fick conftantly to their bed, and preventing their changing their foul linen.

Persons recovering from diftempers, require great care and attention ; in proportion to the abatement and decline of the fever, their quantity of food may be gradually increafed ; and when the fever is entirely gone, the fick perfon may venture on a little white. meat, fuch as chicken, rabbit, whitings, flounders, or other flat fish; but falmon; eels, carp, skait, haddock, and the like, are not to be ventured on till the recovery is abfolutely perfect. Soups, new-laid eggs, and a little wine diluted with water; but thefe are to be used with great moderation, because the ftomach, being extremely weakened by the difease, is capable only as yet of a small degree of digestion; and if the quantity of nurishment exceeds its power ever fo little, it will not digest, but become putrid, and delay the recovery.

To procure a compleat and perfect termination of acute difeafes, observe the following rules.

Let perfons recovering, as well as those who are fick, take very little food at a time, and take it often.

Let their meal confift of one kind of food only, and let them chew their food well.

Ebery Man

Leffen their quantity of drink ; the best in general is wine and water ; three parts water, to one part wine ; for too great a quantity of liquids prevents the stomach from recovering its tone, and increases the tendency to a swelling of the legs.

Riding on horfeback, as often as they are able, is abfolutely neceffary : the propereft time for this exercife is in the forenoon.

They should eat nothing, or at most but very little, in the evening, as perfons in this state are feldom quite fo well towards night. Their sleep will be the lefs disturbed for this caution: seven or eight hours, at most, are as much as should be allotted for lying in bed.

A ftool is not abfolutely neceffary every day; but if the coftiveness exceeds the second day, a glyster should be admistered; or sooner, if the person seels uneasy, is restless, or has the head-ache.

If after fome time they should continue very weak, and their stomach is difordered, and they have, from time to time, a little irregular fever, they should take a tea-cup full of the decoction of the Peruvian bark, three or four times in a day, which may be prepared by boiling an ounce of the best bark in powder, in a quart of water, till two thirds are wasted away, and then adding to the remainder a gill of red wine.

Labouting men mult by no means return to work too foon after their recovery, lest it

his emn Phylician.

prevent their ever getting perfectly well, and entirely recovering their loft strength.

DIRECTIONS CONCERNING BLEEDING.

Though bleeding has been directed in many diforders in the course of this work, yet, as the improper use of it is attended frequently with the most fatal confequences, it has been judged neceffary to bring into one point of view all those cases in which bleeding may be administered with advantage, as well as those in which it is highly prejudicial. In all inflammatory difeafes, it should by no means be omitted, as in the pleurify, and peripneumony, during the first days; but so foon as the symptoms of suppuration appear, expectorating medicines are the most proper means from which to expect relief, and the use of the lancet must be totally forbid. In the apoplexy, epilepfy, bloody-flux and inflammation of the bowels, bleeding must be repeated according to the exigence of the fymptoms : it is also useful, sometimes to promote suppuration in large absceffes, where nature is too much opprest by the violence of the inflammation. In inflammations of the eyes, bladder, or womb, ischiatic pains, rheumatifms, coughs, pleurify, head-aches, quinfeys, afthmas, hemorrhages, and nephritic complaints, bleeding is of the utmost fervice # but in every diforder proceeding from a relaxed state of the folid parts, and impoverished state of the blood, attended with a

H

Ebery Man

chachectic habit of body, fuch as dropfies, jaundice, gout, &c. bleeding must be abstained from with the utmost caution, as it will, instead of relieving, greatly aggravate those distempers.

APPENDIX.

A choice Receipt to make Opodeldoc.

AKE of Hungary-water a pint ; Caftile-foap fliced, three ounces; camphire an ounce; let them stand together in a glass clofely stopped, till the soap and camphire are entirely diffolved in the Hungary-water.

The Clyfter Decoction.

Take of dried mallow-leaves an ounce: chamomile flowers, and fennel feeds, of each half an ounce; boil them in a fufficient quantity of water to strain off about half a pint, then add about two ounces of fweet oil, and it is fit for use.

Hartshorn-Drink.

Take burnt hartshorn two ounces, gumarabic two drachms; boil them in three pints of water, till one pint is wasted away, then strain it, and it is fit for use.

Barley-Water.

Take two ounces of pearl barley, wash it well in cold water, and then boil it in half a pint of water for a very little while ; this water will look redifh, and it is to be thrown away; then add four pints of water, and boil it away to one half : the remainder is fit for ule.

Viper Broth.

Take a middle fized viper freed from the head, skin and bowels, and two pints of water, boil them to a pint and an half, then remove the veffel from the fire, and when the liquor is cold, let the fat which congeals upon the furface be taken off. Into this broth put a pullet of a moderate fize, after having fkinned it, and taken off all the fat carefully; fet the veffel on the fire again, that the liquor may boil; then remove it from the fire, take out the chicken, and chop its flefh into little pieces; put thefe into the liquor again, fet it over the fire, and, as foon as it boils up, pour out the broth, firft carefully taking off the fcum.

Broths taken frequently, and in a small quantity at a time, are excellent reftoratives, and of infinite fervice in decays : on this account viper broth may be reckoned a restorative, though I efteem the chicken the principal ingredient to be depended on; for I am convinced by experience, that vipers have no one virtue to recommend them that can be depended on : but it is usual to overlook the efficacy of things we are daily conversant with, and afcribe their good effects to others that are not to common, though lefs to be depended upon : and I am certain that whoever tries chicken broth, with and without the viper, will find as much fervice from the first as the latter. Dr. James.

An excellent Fomentation.

Takefouthernwood and worm-wood dried, and chamomile flowers, of each an ounce; bay leaves dried, half an ounce; boil them

gently in fix pints of water, and strain it off for use: all green wounds, and old fores, should be fomented with this every day before they are dreffed.

The infusion of Senna.

Take three quarters of an ounce of fenna, cream of tatar three drachms, carraway feeds bruifed, two drachms; boil the cream of tartar in half a pint of water till it is diffolved, then pour the boiling water upon the reft of the ingredients, and let it stand until it is cold, then strain it off and it is fit for use.

A Purging Draught.

Take of the infusion of senna, as above directed, two ounces; syrup of buckthorn one ounce; mix them together for one dose, which may be taken in the morning fasting, three times a week, and it is a safe and sure purge : it may be taken in all cases where purging is proper.

A Dofe of Cooling Physic.

Take Glauber's falts an ounce, manna half an ounce, diffolve them in a little boiling water, for one dofe, to be taken as often as occafion requires.

Hiera Picra.

Take of fuccotrine aloes finely powdered, a quarter of a pound; winter's bark finely powdered three quarters of an ounce; mix them together.

Tincture of Hiera Picra.

Steep an ounce of hiera picra, made as above directed, in a pint of mountain wine, for a week or ten days, by which time it will be fit for ufe.

Mindererus's Spirit.

Take a quarter of an ounce of volatile falarmoniac, and add to it by degrees, distilled vinegar, till the effervesence entirely ceases.

An excellent Bitter Tincture.

Take gentian root, and orange peel dried, each two ounces; cochineal bruifed, half a drachm; infufe them in a quart of French brandy for three or four days, then strain off the liquor for use. This is a very pleafant and wholefome bitter, and may be used occafionally to provoke an appetite, and affist digestion.

Duke of Portland's Gout Powder.

Take of round birthwort root and gentian root, of the tops and leaves of germander, ground pine, and centaury : Take of all these, well dried, powdered and fifted as fine as possible, an equal quantity; mix them well together, and take one drachm of this mixed powder every morning fasting in a cup of wine and water, broth, tea, or any other vehicle you like best : keep fasting an hour and an half after it, continue this for three months without interruption, then diminish the dose to three quarters of a drachm for three months longer, then to half a drachm for fix months more, taking it regularly every morning if poffible. After the first year, it will be sufficient to take half a drachm every other day.

As this medicine operates infenfibly, it will take prehaps two years before you receive much benefit; fo you must not be difcouraged, though you do not perceive at first any great amendment; it works flow but fure: it doth not confine the patient to any particular diet, fo one lives foberly and abstains from those meats and liquors that have always been accounted pernicious in the gout; as champaign, drams, high fauces, &c.

N. B. In the rheumatism, that is only accidental and not habitual, a few of the drachm doses may do; but if habitual, or has been of long duration, then you may take it as for the gout. The remedy requires patience, as it operates but low in both diftempers.

Sir Hans Sloane's Oitment for Sore Eyes,

Take prepared tutty one ounce; blood ftone finely prepared two fcruples; fuccotrine aloes in fine powder, twelve grains; prepared pearl, four grains; mix them together in a marble mortar, with as much viper's fat as is fufficient to give it the confiftence of an ointment

This øintment is to be applied with a hair pencil, the eyes winking or a little opened.

It must be observed, that purging medicines are hurtful to the diseases of the eyes, which are cured by this medicine.

Dr. Mead's Remedy for the Bite of a Mad Dog.

Let the patient be bled in the arm, to the amount of nine or ten ounces. Take ash coloured ground liverwort four drachms, black pepper two drachms, mix them together into a fine powder.

This is to be divided into four dofes, whereof one is to be taken in warm milk, in a morning fafting, for four mornings fucceffively; after this the perfon must be put into a cold bath, pond, or river, for thirty days together, early in the morning, and before breakfast; he is to remain in it, with his head above water, not longer than half a minute. The wound should be continually fomented with a pickle made with vinegar and falt, as warm as it can be borne.

Tonquin Remedy for the Bite of a Mad Dog.

Take of native cinnabar, and common vermillion (both ground to a very fine powder) each twenty-four grains; muſk ſixteen grains; rub theſe together, till the muſk is alſo become very fine, and give it all for one doſe in a tea cup full of arrack, or brandy, as foon as poſſible after the perſon is bit; and another doſe thirty days after : but if the perſon has the ſymptoms of madneſs before he has kad the medicine, he muſt take two doſes in an hour and a halſ. This medicine is perſectly ſaſe and innocent, as appears by the great number of perſons who have tried it, none of whom have felt any ill conſequences, or been diſordered ſince through its uſe.

Its only visible effect, is producing a remarkable drowfines, which generally ends

in a perfect cure. This dose is intended for a grown perfon; for children it must be lesfened in proportion to their age.

Mrs. Stephen's famous Gure for the Stone and Gravel, as amen ded by 'Dr. Hartly.

Take two fcruples of calcined egg-shells, three times in a day, in a glafs of any convenient liquor, drinking, after each dofe, a third part of the following decoction : Take two ounces of Castile soap, dissolve it in a quart of foft water, and fweeten it according to your tafte, with honey or loaf fugar. If this decoction is made in a copper veffel, care must be taken that it is well tinned. The taking of these medicines must be continued for fome time after the complaint ceafes, left any part of the flone fhould remain, which being rough and unequal, might occasion exquisite pain. It is common after a few days use of these medicines, to have an increase of pain in making water ; at which time a foft diet, emollient drinks, and reft, are proper. For common drink, milk and water, or a decoction of marsh-mallow roots, parsley, and liquorice, may be used ; but if the perfon has been used to strong liquors, fmall punch, made without any acid, may be drank sparingly. Artichoakes, asparagus, spinnage, letuces, fuccory, parfley, turnips, carrots, potatoes, radifhes, peas, &c. may be fafely ufed; but onions, leeks and cellery are to be preferred to all other vegetables.

The patient ought to drink no more of any liquot than is fufficient to quench his thirft, and he fhould hold his water, as long as he can without great uncafinefs, that it may have the longer time to act on the ftone. If thefe medicines occafion coftivenefs, it will be neceffary now and then, at difcretion, to take a dofe of Glauber's falt and manna.

A Warm Purge, for the Palfy and Apoplexy.

Take tincture of hiera picra two ounces; fyrup of buckthorn three quarters of an ounce; lavender drops a quarter of an ounce; mix them together for one dofe, which may be given in the morning fasting, twice a week, or as often as occasion may require.

Daffy's Elixir.

Take raifins ftoned four ounces, fenna three ounces, carraway-feeds bruifed one ounce, fteep thefe ingredients in a quart of brandy, for three weeks or a month, then ftrain it off for ufe, and keep it in a bottle clofe ftopped.

Tincture of Rhubarb.

Take of rhubarb fliced thin, one ounce; leffer cardamon feeds bruifed, a quarter of an ounce; faffron, a dram; brandy, one pint: steep these ingredients together for three weeks or a month, then strain off the liquor for use.

N. B. Mountain wine may be used instead of the brandy.

Volatile Tincture of Guaicum.

Take an ounce of gum guaicum in grofs powder, and steep it for a week or ten days in fix ounces of fal volatile drops, in a bottle closely stopped.

Fryer's Balfam.

Take of gum benjamin one ounce and an half, of strained storax one ounce, of balfam of Tolu half an ounce, of fuccotrine aloes a quarter of an ounce, rectified spirit of wine a pint. Digest them together till the gums are diffolved as much as possible, then strain off the tincture for use.

Diachylon with the Gums.

Take three quarters of a pound of white diachylon, two ounces of strained galbanum, turpentine and frankincense of each three quarters of an ounce; melt them together over a flow fire.

White Diachylon Plaifter.

Take litharge finely powdered a pound and a quarter, fweet oil a quart, boil them together with a quart of water, till they are thoroughly mixed, and are of a proper confiftence for a plaifter, and look quite white; if the water fhould be entirely wafted away, you must add fome more to prevent its turning black.

An excellent Strengthening Plaister.

Take white diachylon half a pound, frankincenfe two ounces, and draggon's blood three quarters of an ounce; melt the diachylon over a flow fire, and then add the other ingredients finely powdered, and mix them well together, by stirring them continually till the plaister is quite cold.

Ointment of Elder.

Is made by boiling the young leaves of elder in mutton-fuet, till they are quite crifp, and the fuet is of a deep green colour.

Spermaceti Ointment.

Pake a quarter of a pint of the best fallad oil, a quarter of a pound of white wax, and half an ounce of spermaceti; melt these ingredients together, over a gentle fire, and keep them continually stirring, until the ointment is quite cold. This ointment is remarkably efficacious in preventing pits on the face after the small-pox.

Ointment of Marshmallows.

Take half a pound of marshmallow-roots; of linfeeds and fænugreek-feeds, each three ounces; bruise them, and boil them half an hour gently in a quart of water; then add two quarts of sweet oil, and boil them together till the water is quite wasted away; then strain off the oil, and add to it a pound of bees-wax, half a pound of yellow rosin, and two ounces of common turpentine; melt them together over a flow fire, and keep them continually stirring, till the ointment is cold.

Turner's Cerate. See Burns and Scalds. Yellow Basilicon.

Take fweet oil a quarter of a pint; beeswax, yellow rofin, and burgundy pitch, of

each a quarter of a pound; Venice turpentine three quarters of an ounce; mix them together over a flow fire.

An excellent Powder to clean the Teeth.

Take cream of tartar, three quarters of an ounce; choice myrrh, three drachms; dragon's blood, one drachm; mix them for a powder.

An excellent Lip Salve,

Is made by adding a quarter of an ounce of alkanet root to the spermaceti ointment, and simmering them together a few minutes over a gentle fire.

Lime-Water.

Pour a gallon of boiling water on a pound of unflacked lime; ftir them well together, and let the water remain on the lime for twenty-four hous; then ftrain off the clear liquor for ufe.

Champhorated Spirits of Wine.

Diffolve an ounce of camphire in a pint of rectified spirits of wine.

Hungary Water.

Take of fresh rosemary tops a pound and an half; proof spirit of molasses, a gallon, and distil off about five pints.

Lavender Water.

Take of fresh lavender flowers a pound and an half; of proof molaffes spirit, a gallon, and distil off five pints.

Ebery Man his own IPhylician. 102

Lavender Drops. Take of lavender water a pint and an half; of Hungary water, half a pint; of cinnamon and nutmegs, each a quarter of an ounce, of red fanders powdered, a drachm and an half; let them infuse together in a glass bottle well stopped for a month : then strain off the liquor for use.

INDEX OF DISEASES,

For which Remedies are proposed in this Treatife.

tore they wanted half a want a set of the	Page.
CHES, from colds	5
After pains	ib.
Agues, or Intermitting Fevers	6
Apoplexy	9
Apppetite, want of	10
Althma	ib.
Saint Anthony's Fire	8
Baldnefs	II
Bite of a Mad Dog	ib.
of a Viper or Adder	12
Bleeding at the Nofe	ib.
Blood, fpitting of	ib.
Bloody Flux	13
Boils	14
Breasts, Disorders of the	52
Bruises, External	14
Internal	ib.
Burns and Scalds	15
Cancer	ib.
Chapt Nipples	53
Colds	18
Cholic	ib.
Chilblains '	17
Confumption	19
Convultions	20
Corns	ib.
Costiveness, to prevent	21
Cough	ib.
Hooping	38

INDEX.

and the second	Page.
Decays, see Consumptions	21
Deafnefs	ib.
Delivery, to promote	22
Diabetes, and piffing of Blood	ib.
Directions for preferving Health	73
for bleeding	89
Dropfy	23
Dry Belly-Ache	27
Drowning	24
Ear-Ache	
	28
Excoriations	ib.
Eyes, Diforders of the	ib.
Fainting	28
Falling Sicknefs	ib.
Fevers, Inflammatory	29
Fevers, Nervous	30
Spotted	31
Fits, see Hysterics	* 37
Fiftula	33
Giddinefs	
Gout	ib.
Gravel and Stone	34
Green Sicknefs	35
Gripes	ib.
	36
Head-Ache	ib.
Heart-burn	ib.
Hoarfenefs	37
Hysterics	37 ib.
Hard Swellings	38
Hip-Gout	ib.
Hooping-Cough	ib.
Jaundice	39

INDEX.	105	
T 1 0.	- Page.	
Indigeftion	40	
Inflammations Itch	ib.	
Inflammations of the Bowels	41	
Intermitting Fever, see Agues	42	
Anternating Lever, ice 119405	10 4 1 P	
King's Evil	42	
Leprofy	43	
Lethargy, fee Apoplexy	9	
Limbs frozen	44	
Loofenels	45	-
4		
Madnefs	46	
Meafles Marfas investor Electro	ib.	1
Menfes, immoderate Flow of	47	
Suppression of Miscarriage	48 ib,	
Mortifications		
Milk Fever, to prevent	49	
and a croit, to pretent	53	
Nervous Diforders	50	
Nipples, Chapt	53	
Noife in the Ears	50	
	1111	
Opiate	50	
Obstructions	ib.	
Old Ulcers	51	
Ointment for the Eyes	28	
Paine in the Lainte for His Cout	1 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
Pains in the Joints, see Hip-Gout Pain in the Stomach, see Heart-burn	38	
Palfy	36	
town in the second of the	55	

106 INDEX.	
100 INDLA.	Page.
Piles	54
Pleurify	55
Poifon, to expel	56
Quinfey	57
Rheumatifm	ib.
Retention of Urine	58
Rickets in Children	58 ib.
Ruptures	59
Rules for nurfing fick Perfons	84
CB	The second
Scald Head	59
Scurvy	бо
Small-Pox	ib.
Sore Throat, see Quinsey	57
Strains	62
Sweating, Exceflive	64
Slow Fever, see Nervous Fever	30
Stranguary	59
Stone, see Gravel	35
Stitch in the Side, see Pleurify	55
Shingles	62
Swelled Legs.	64
Teeth, difficult cutting of	63
Tooth-Ach	64
Tettars	66
Thirst, excessive	ib.
Thrush	ib.
Throat Sore, see Quinsey	57
Putrid	64
Tympany	67

INDEX.	107
Violent Pains after Delivery see After Pains	Page.
Vomiting	67
Ulcers of the Lungs	ib.
Kidneys	ib.
Bladder.	68
Ulgers in the Womb	ib.
old ·	51
Warts	69
Whites	70
White Swellings	71
Wind	70
Watery Gripes	69
Wasting away of Children	71
Whitloe	ib.
Worms	72
Wounds, green	ib.

INDEX TO THE APPENDIX.

RECEIPT for making Opodeldock ·	91
	ib.
Hartshorn Drink	ib.
-Barley Water	ib.
Viper Broth	ib.
an excellent Fomentation	\$ 92
Infusion of Senna	93 ib.
a Purging Draught	
Hiera Picra	ib.
Tincture of ditto	ib.
and the second	

INDEX.

the same with the service of the	Page.
Mindererus's Spirit	94
the Bitter Tincture	ib.
-Duke of Portland's Gout	Powder ib.
Sir Hans Sloane's Ointme	ent for 7
fore Eyes	\$ 95
-Dr. Mead's Remeby fo	r the 7 .
Bite of a Mad Dog	ib.
the Tonquin Remedy,	96
Steven's Stone Medicine	97
a Warm Purge	. 98
for making Daffy's Elixir	ib.
Tincture of Rhubarb	ib.
Volatile Tincture	99
Friar's Balfam	ib.
Diachylon with the Gums	,ib.
White Diachylon	ib.
Strengthening Plaifter	ib.
Ointment of Elder	100
Spermaceti Ointment	ib.
Ointment of Marshmallow	ib.
Ointment for burns and Sc	alds 15
Yellow Bafilicon	100
a Powder for the Teeth	IOI
-a Lip Salve	ib.
Lime Water	ib.
Camphorated Spirits of W	ine ib.
Hungary Water	ib.
-Lavender Water	ib.
Lavender Drops	102
The second s	

THE END.







