

**A synopsis of the diseases of the eye, and their treatment : to which are prefixed a short anatomical description and a sketch of the physiology of that organ / by Benjamin Travers ; with notes and additions by Edward Delafield.**

### **Contributors**

Travers, Benjamin, 1783-1858.  
Delafield, Edward, 1794-1875.  
National Library of Medicine (U.S.)

### **Publication/Creation**

New York : E. Bliss and E. White, 1825.

### **Persistent URL**

<https://wellcomecollection.org/works/kxqpw967>

### **License and attribution**


This material has been provided by This material has been provided by the National Library of Medicine (U.S.), through the Medical Heritage Library. The original may be consulted at the National Library of Medicine (U.S.) where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>













AN 29 1962









R. J. Bond

A

## SYNOPSIS

OF THE

# DISEASES OF THE EYE,

AND THEIR TREATMENT:


TO WHICH ARE PREFIXED

A SHORT ANATOMICAL DESCRIPTION

AND

A SKETCH OF THE PHYSIOLOGY

OF THAT ORGAN.



BY BENJAMIN TRAVERS, F. R. S.

SURGEON TO ST. THOMAS'S HOSPITAL.

WITH NOTES AND ADDITIONS

BY EDWARD DELAFIELD, M. D.

*Surgeon to the New-York Eye Infirmary, and Lecturer on Diseases of the Eye.*

FIRST AMERICAN FROM THE THIRD LONDON EDITION.

NEW-YORK:

PUBLISHED BY E. BLISS AND E. WHITE, AND H. C. CAREY  
AND I. LEA, PHILADELPHIA.

1825.

*Southern District of New-York, ss.*

**BE IT REMEMBERED**, That on the twenty-seventh day of September, in the fiftieth year of the Independence of the United States of America, E. Bliss and E. White, of the said District, have deposited in this office the title of a Book, the right whereof they claim as proprietors, in the words following, to wit :

" A Synopsis of the Diseases of the Eye, and their treatment : to which are prefixed a short Anatomical Description and a Sketch of the Physiology of that Organ. By Benjamin Travers, F. R. S. Surgeon to St. Thomas's Hospital. With Notes and Additions by Edward Delafield, Surgeon to the New-York Eye Infirmary, and Lecturer on Diseases of the Eye. -- First American from the Third London Edition."

In conformity to the Act of Congress of the United States, entitled "An Act for the encouragement of Learning, by securing the copies of Maps, Charts, and Books, to the authors and proprietors of such copies, during the time therein mentioned." And also to an Act, entitled "an Act, supplementary to an Act, entitled an Act for the encouragement of Learning, by securing the copies of Maps, Charts, and Books, to the authors and proprietors of such copies, during the times therein mentioned, and extending the benefits thereof to the arts of designing, engraving, and etching historical and other prints."

JAMES DILL,

*Clerk of the Southern District of New York.*

WW  
T779s  
1825

C  
102-1

TO

**WRIGHT POST, M.D.**

President of the College of Physicians and Surgeons of the University of the State of  
New-York, and Professor of Anatomy and Physiology, Consulting Surgeon  
to the New-York Hospital and New-York Eye Infirmary,

**THIS WORK IS DEDICATED,**

AS AN

**Evidence of Respect**

FOR HIS

**PROFESSIONAL EMINENCE AND PRIVATE WORTH,**

AND A

**TRIBUTE OF GRATITUDE**

FOR THE

**ADVANTAGES DERIVED FROM HIS INSTRUCTIONS,**

BY HIS SINCERE FRIEND,

**THE EDITOR.**



WRIGHT PORT M.D.

THIS WORK IS DEDICATED

TO THE

PROFESSOR OF

THE

AND

BY HIS SINCERE FRIEND

THE EDITOR

## PREFACE.

---

THE Volume which I now present to the Public, is the result of a more ample opportunity of observing the diseases of the important organ of which it treats, than commonly falls to the lot of Hospital Surgeons. This opportunity, originally derived from my situation during a period of seven years, as surgeon to the London Infirmary for Diseases of the Eye, has been considerably augmented by private practice in the same branch of the Profession, contingent to that appointment.

I have always thought that the advantages obtained by the subdivision of professional talent and labor, are infinitely overbalanced by those which arise from the general and undivided application of these instruments of knowledge. No fact more

strikingly illustrates the truth of the doctrine, that the confinement of any branch of the Profession to the hands of a few, operates prejudicially to science, than the state of information in this country, concerning the Diseases of the Eye. Mr. Samuel Cooper has spoken so precisely, as I think, upon this subject, that I cannot do better than quote his words.

“ The disorders of the eye and its appendages are far more numerous and diversified than those of any other individual part of the body, and some of the requisite operations for their relief ought to be done with the nicest combination of skill and delicacy. These circumstances, strangely enough, have had the effect of inducing an erroneous supposition, that such cases do not properly enter into the department of ordinary surgery ; but ought to be consigned to the care of a man, who makes them exclusively the object of his attention, and disregards disease in every other form. The morbid affections of the eye, it is true, like all other surgical cases, must be studied, in order to be understood.



They have no peculiarity, however, except what depends upon their number, and the tenderness and functions of the organ affected. In their nature they are swayed by the same laws which influence all common diseases, for which the practice of surgery is instituted; and their treatment is regulated by general principles, which prevail throughout the whole of this indispensable art.

“No one, except the thorough surgeon, can make the complete oculist; by which last term is not meant any body who can merely manage to extract the cataract better than the generality of surgeons, but a man whose science leads him to recognise the analogy betwixt the diseases of the eye and those of other parts, and whose knowledge of the latter, while it qualifies him in a great measure for the treatment of the former, gives him a decided superiority over the bare oculist.”

“On a cru faussemment (says the intelligent M. Louis, in adverting to the diseases of the eye), que le sçavoir nécessaire pour discerner le caractère de ces diverses af-



fections contre-nature, et pour y remédier, faisoit en quelque sorte un art particulier. Mais quels fruits pourroit porter cette branche, étant séparée du tronc? Il est bien prouvé, par les faits, que les progrès de cette partie de la chirurgie ne sont dûs qu'aux grands Maîtres qui ont pratiqué l'art dans toute sa plénitude, et dont l'expérience, relative aux maladies des yeux, a été éclairée par les lumières que leur avoient données les principes qui constituent indivisiblement la science, sans laquelle on ne peut exercer aucune partie avec connoissance de cause."—First Lines of the Practice of Surgery, Vol. I. p. 433, 4th edit.

In this country. I believe no one before myself, who designed to practise general surgery, ventured to give more than a cursory attention to the diseases of the eye. A fear of being disqualified in public opinion, by a reputation acquired in these, for the treatment of other diseases, was a motive, however groundless, sufficient to deter surgeons from the cultivation of a large and legitimate field of observation and practice.

It was with a public avowal of the sentiments so well expressed by the writers just quoted, that I accepted the situation of Surgeon to the Eye Infirmary, in the year 1810; and from these I have never swerved.

At the commencement of the following year, the students of surgery were first invited to attend the practice of the Infirmary; an opportunity eagerly embraced, and which many hundreds have since enjoyed.

Among the gentlemen who with ardor and diligence entered upon this new and interesting study, during my connexion with the Infirmary, I have the pleasure of including many of the best educated and most rising men in the Profession; and, if they will permit me to say so, some of my most estimable friends.

Upon occasion of electing a second surgeon to that Establishment in 1814, my friend, Mr. Lawrence, became my colleague. I consider it to be no ordinary sanction of my views, that they were thus seconded by the



co-operation of a gentleman, so highly distinguished as the present senior surgeon of the Infirmary.

But whether my example or my services have been in any degree instrumental in promoting so desirable an object as that of recalling to the notice of the profession at large the neglected subject of these diseases, is a matter of no public interest, and which I am content to leave to the candor of the Profession.

In offering these observations, I entreat it may be understood, that it is far from my meaning to insinuate the slightest derogation from the merits of some truly respectable Members of the Profession, who confine their attention to this class of diseases. We take a different view of the subject; but no man of ingenuous feelings will arrogate to himself that he alone walks in the right path.

In this country, the want of a comprehensive treatise on the diseases of the eye has long been felt and acknowledged. The work of Professor Scarpa was the only



book of reference for English students before the publication of Mr. Saunders's Treatise, which from its intrinsic evidence of a strong and original talent for observation, leaves us to regret as a national calamity, the premature termination of his labors. Mr. Wardrop's ingenious Essays on the Morbid Anatomy of the Eye, have since contributed to instruct and gratify the Profession. But the object is yet unaccomplished, and I can only flatter myself with having advanced a step or two nearer to its completion.

In Germany a merited share of attention has long been devoted to the diseases of the eye. The elaborate work of Professor Beer of Vienna, who has devoted a life to the subject, is said by those of our countrymen who read the German language, and are competent to appreciate its merit, to evince a familiarity with these diseases—a comprehensiveness of arrangement—a depth and minuteness of observation—a promptitude and fidelity of diagnosis—unattained in any other department of Pathology. Walther of Landshut, Schmidt of Vienna, Himly and Langenbeck of Gottingen, and others,

have distinguished themselves in the same field of research.

I ardently hope that the example of German industry may operate as an incentive to our exertions, and should it prove so—without meaning to question the extent or value of their researches in ophthalmology—I may be permitted to express my belief that they can lay us under no deeper obligation.

Nations, like individuals, are distinguished by a peculiar character of mind, to whatever causes attributable, evinced in their respective modes of observing, reflecting, and acting; and the sentiment of Phædrus is as strictly applicable to the one as to the other:

*Sua cuique quum sit animi cogitatio*

*Colorque privus*——\*

I should be sorry to see the sober sense of my countrymen perverted by a taste for fastidious distinctions. Simplicity is the characteristic feature of English Surgery, which is neither more nor less than the

\* Prol. Lib. V.



application of the principles of inflammation, as illustrated by the genius of JOHN HUNTER.

I cannot but fear that the condensed and compendious plan of this Volume may occasion disappointment to persons advanced in the study of eye diseases. But I must remind them that it makes no pretension to the character of an elaborate systematic treatise, and is designed more particularly for the information of surgeons and students of surgery. It is the result of personal observation, not a compilation from the works of others. Neither is it in any degree critical. I have not time enough at my command to be an historian, nor ambition to be a controversialist. Facts, whether new or old, derive a value of authenticity from personal observation, which is not enhanced by the custom of collating materials from other sources. Such a custom is likewise inconsistent with another valuable quality, in practical matters especially—I mean conciseness.

It was at the suggestion of a friend that I prefixed the sketches of anatomy and



physiology, for the purpose of bringing the entire subject before the reader in the compass of a volume. In these short pieces, I have aimed at a simple and perspicuous exposition of all that appears essential to a right understanding of the principal diseases of the organ. For the more subtle anatomical points, or rather questions, which I have purposely avoided, I refer my reader to the Appendix.

Against an overweening spirit of anatomical discovery, the pathologist should be jealously upon his guard, if he would observe honestly and reason accurately on the phenomena of disease. Some ingenious disquisitions have occasionally been given to the Public on the morbid affections of textures, of which the existence is problematical. I would by no means speak with disrespect of the pursuit of minute anatomy, nor presume to question its utility when conducted after the manner of Harvey and Ruysch ; but I must be allowed to remark that such points as are too subtle to admit of demonstration, are not to be cleared by the helps of analogy and conjecture, and indeed are never satisfactorily

proved but by the anatomy of disease, from which they derive their importance.

The second part of the work, containing the Pathology was drawn up and intended for publication in the last Volume of the Surgical Essays published jointly by Sir Astley Cooper and myself. It is unnecessary to state the reasons of its postponement; a principal motive was the belief that it would be more useful and acceptable to the junior part of the Profession, as it now stands, connected with the remaining contents of this volume. I am induced to mention this circumstance by way of apology, for what might appear to be occasional infringements upon the plan of considering the diseases and their treatment distinctly. To a certain extent the blending of these subjects is unavoidable, but the Pathology having been designed for a separate and distinct Essay without any direct view to the subject of treatment, the latter is introduced more freely in the way of illustration, than it would otherwise have been; and, consequently, anticipations and repetitions are more frequent. I think, however, that this fault is mainly in the



plan. I should not therefore *a priori* have adopted it, but I was unwilling to disturb the arrangement of the piece first written, and to some readers it is probable that the division may not be unacceptable.

Some highly important topics, in the third division of the volume, my experience, as the reader may conclude, would have enabled me to treat much more in detail, than the plan of this work permitted. It is difficult to speak in so small a compass upon subjects like those of cataract and artificial pupil, either of which is sufficient from its extent and importance, to furnish materials for a distinct volume. I have endeavoured how successfully I know not, to guard against material omissions, in studying to preserve conciseness.

As I have made little reference to the writings of others, the omission of the synonyma forming the crabbed vocabulary of Ophthalmologists, ancient and modern, will, I hope, need no apology. The nomenclaturing mania appears to me an evil of increasing magnitude.

New Broad Street,

Oct. 28, 1820.



## PREFACE

TO

### THE THIRD EDITION.

IN a careful review of this Treatise in its passage through the press, I have not seen occasion to make any material alteration. The compendious method in which the subject is treated, which forms probably one of its principal recommendations to practical readers, precludes me from adding the details of a more extended observation—a restraint to which I the more readily conform, from the persuasion that they would tend rather to confirm than to correct the pathological views which the work presents; and would so considerably increase the size, as to change the character, and augment the cost of the volume.

I have the pleasure to acknowledge many liberal and handsome testimonies to its value from individuals of great and deserved eminence in the profession ; and the public confirmation of these, I take the liberty to infer from the rapid sale of the work, which has been out of print for many months—but a circumstance which has afforded me the sincerest gratification, is the appearance, during the last year, of a spirited and faithful translation of it into the Italian language by a physician of Pisa, whose name I would gladly announce had his modesty permitted me to do so.

To Dr. T. I. Todd, of Nice, I beg thus publicly to express my obligations for his spontaneous recommendation of the original to the notice of so able and intelligent a translator. To the latter gentleman I offer my congratulations upon the public approval of his translation, evinced, as I am informed, by a very ready sale.

*New Broad Street,*

*April 15, 1824.*

# CONTENTS.

---

## PART I.

	Page
Anatomical Description of the Eye and its Appendages	1
Sketch of the Physiology of the Eye and its Appendages	45

## PART II.

### CHAPTER I.

#### PATHOLOGY OF THE MEMBRANES.

SECTION I. Conjunctiva	87
II. Cornea	106
III. Sclerotica	126
IV. Choroid and Iris	131
V. Retina	138

### CHAPTER II.

#### PATHOLOGY OF THE HUMORS.

SECTION I. Aqueous Humor	203
II. Vitreous Humor	206
III. Crystalline Humor	218
IV. Diseases affecting the Eyeball	227



## CHAPTER III.

## PATHOLOGY OF THE APPENDAGES.

	Page
SECTION I. Orbital Appendages.....	240
II. Facial Appendages .....	247

## PART III.

## CHAPTER I.

## TREATMENT OF THE DISEASES OF THE EYE.

SECTION I. Simple Inflammation of the Conjunctiva...	264
II. Inflammation modified by Struma .....	277
III. Acute suppurative Inflammation of the Con- junctiva.....	284
IV. Secondary Diseases of the Conjunctiva....	292

## CHAPTER II.

SECTION I. Diseases of the Cornea .....	301
II. Scleritis, Choroiditis, and Iritis .....	313
III. Amaurosis .....	318
IV. Diseases affecting the Eyeball.....	332

## CHAPTER III.

SECTION I. On the Operations for the Cataract .....	337
II. Of the Operations for artificial Pupil ....	360

## CHAPTER IV.

## DISEASES OF THE APPENDAGES.

	Page
SECTION I. Diseases of the Eyelids.....	377
II. Obstruction of the Lacrymal Passages....	386

---

APPENDIX.....	411
EXPLANATION OF THE PLATES.....	421
NOTES.....	447

CHAPTER IV

CONTENTS OF THE APPENDIX

Section I. Diseases of the Eye	375
Section II. Diseases of the Ear	380

APPENDIX	411
EXPLANATION OF THE PLATES	411
NOTES	417

A. APPENDIX

THE ART OF READING AND WRITING

Section I. The Art of Reading	411
Section II. The Art of Writing	417

B. APPENDIX

Section I. The Art of Reading	411
Section II. The Art of Writing	417

C. APPENDIX

Section I. The Art of Reading	411
Section II. The Art of Writing	417



PART I  
ANATOMICAL DESCRIPTION  
OF  
THE EYE  
AND ITS APPENDAGES  
ANATOMY AND PHYSIOLOGY

OF

**THE EYE**

AND ITS APPENDAGES.





## PART I.

### ANATOMICAL DESCRIPTION

OF

### THE EYE

#### AND ITS APPENDAGES.

---

THE orbits are two funnel-shaped cavities situated under the arch of the forehead, on either side of the root of the nose. The roof of the orbit is formed by the frontal and sphenoid bones: the floor is contributed by the superior maxillary and malar bones; the malar and sphenoid bones make up the temporal side; the lacrymal, æthmoid, sphenoid, and palate bones compose the nasal. The nasal sides are plane, and nearly parallel; the temporal are considerably divergent, so that the axis of the orbit is an oblique line. A line drawn horizontally across the base of the cavity is also oblique; the nasal being more advanced than the temporal angle. This configuration of the orbits prevents us from commanding the parietes of both cavities in a front view of the cranium, and greatly extends the field of vision. The globe of the eye is considerably

Orbits:

smaller than the receptacle in which it is contained, to allow of its free motion on all sides. The capaciousness of the orbit provides for the lodgment of the adipose substance upon which the eyeball is cushioned; the muscles which move and adapt it to vision in all directions; the vessels which nourish its membranes and secrete its humors; the nerves which supply these several parts with energy; and the gland for preparing the lubricating fluid which is essential to its economy. These being situated posterior to that membrane which shuts up the cell of the orbit, may be termed, for the convenience of description, the orbital appendages, in contradistinction to those which the organ presents on its facial aspect, viz. the supercilium or eyebrow, palpebræ or eyelids, and the lacrymal passages; which parts, together with the tunica conjunctiva, I shall call the facial appendages of the globe.

Their holes  
and depres-  
sions.

The foramina and depressions of the orbit, as of all the bony cavities, form a natural introduction to the anatomy of the parts therein contained. The nasal and superior sides, for example, are terminated by the optic hole for the introduction into the orbit of the optic nerve. The temporal and superior sides are bounded by an irregular slit or fissure, denominated foramen lacerum orbitale superius; the temporal and inferior are separated by a similar fissure, termed foramen lacerum orbitale inferius vel spheno-maxillare.



The numerous nerves, the optic nerve excepted, and the principal vein of the eye and its appendages pass through the foramen lacerum. Two minute foramina left in the suture, connecting the frontal and æthmoid bones, by which a direct vascular and nervous communication is maintained between the nares and the orbit, are termed foramina orbitalia interna, anterius et posterius. A small hole in the orbital portion of the malar bone establishes a similar communication with the cheek. A depression in the orbital plate of the os frontis, next its external angular process, receives the lacrymal gland which is fastened to it by a particular ligament. A minute pit behind and above the internal angular process of the same bone, gives attachment to a ligament and cartilaginous trochlea, in which the tendon of a muscle plays in its passage to the globe. A notch or foramen is observed in the orbital ridge of the os frontis, which permits vessels and nerves to pass from the orbit to the eyebrow, glabella, and forehead. The infra-orbital canal, which opens obliquely at the back of the orbital floor, is continued under it to the cheek, transmits vessels freely communicating with the ophthalmic, and a nerve from which the inferior palpebra derives its chief supply.

THE GLOBE OR BALL OF THE EYE is not exactly Eyeball.  
spherical, the line forming the visual axis ex-

ceeding its transverse diameter. This line is parallel in the two eyes.

The figure of the orbit demonstrates that a part only of the ball is contained within it. A needle placed upon the temporal angle of the orbit, and pushed horizontally across the globe, perforates the orbital plate of the æthmoid bone, and measures its greatest transverse diameter.

The difference in the degree of projection of the eye in different individuals is determined by the relative volume of the ball and its socket; but the figure and the ordinary interspace of the eyelids are subject to variations, which convey a delusive idea of the magnitude of the globe. When a paralysis affects the palpebral muscle of one eye, the organ, compared with its fellow, has the appearance of being diminished in bulk.

The eye of the female is commonly smaller than that of the male; and the fissure of the eyelids, which are rounder, broader, and more delicate in texture, is generally less.

Humors  
and mem-  
branes.

The eyeball is composed of the following parts:—

- |                           |                                    |
|---------------------------|------------------------------------|
| 1. The vitreous humor,    | } united by their<br>common tunic. |
| 2. The crystalline humor, |                                    |



3. The aqueous humor.
4. The retina.
5. The choroid and its appendages, the annulus and processus ciliares.
6. The iris.
7. The sclerotic.
8. The cornea.

The humors give shape to the eyeball, and support to its tunics.

The crystalline is set in the vitreous humor, and washed in front by the aqueous.

The retina is the membranous expansion of the optic nerve, upon which the images of external objects are painted.

The choroid is the bed of the vessels of the eye, and the dark screen which confines and condenses the rays of light. Its appendages are auxiliary to this purpose, and to other parts of the economy of vision.

The iris is the colored membrane in which the aperture termed 'the pupil' is formed.

The sclerotic is the external opaque investiture of the choroid.

The cornea is the anterior transparent membrane which first converges the rays of light.

Nearly in the order in which these several parts have been named, I proceed to describe them.

Vitreous  
humor.

THE VITREOUS HUMOR is the basis upon which the larger tunics are expanded, and fills a space somewhat exceeding three quarters of the volume of the globe. Upon its anterior surface it is rather abruptly flattened, and presents a central cup-like depression; the dimensions of which exactly correspond to the posterior segment of the crystalline humor, which is imbedded therein. Its substance is a glairy fluid heavier than water, perfectly pellucid, and contained within cells formed by processes of its tunic, arranged in horizontal planes. Towards the back and sides of the humor these cells are larger than in the interior, adjacent to the crystalline fossula; the septa are likewise thicker and stronger towards the circumference of the humor. After a careful section of the frozen humor, its substance may be picked out in solid wedge-like flakes from the interstices of the septa. The continuous covering, though of great tenuity and perfect transparency, is of much strength, and resists, owing to the support it receives from the numerous septiform productions of its internal surface, a considerable pressure. When lacerated or wounded, the humor of the corresponding cell or interstice is instantly evacuated; but if the wound is superficial, the humor does not escape in quantity, while supported by the other parts of the globe, or if removed from the globe, while suspended

Hyaloid  
tunic.



in a fluid. But if in any way compressed after a wound, a dribbling of the humor goes slowly on, until the cells, which communicate with each other, are emptied.

The tunica hyaloidea is covered by the retina in the whole extent of that membrane, but is connected with it only at the entrance of the optic nerve. The substance of the humor is penetrated by a branch of the arteria centralis retinæ, which contributes a few very delicate vessels to its containing membrane. In the fœtus they have been displayed ramifying on the capsule at the back of the lens.

THE CRYSTALLINE HUMOR is a double convex lens, its breadth about four lines, its thickness about two. The posterior and most convex face of the lens is exactly fitted to the cup in the fore-part of the vitreous humor; the anterior is opposed to the iris, and the circumference to the canal of Petit. The axis of the lens is that of the pupil, a little to the inner side of the axis of the eye. This humor is of perfect transparency in its healthy state. In the fœtus and new-born infant, it is spherical, semi-fluid, and has a slightly reddish tint. In the adult, it is gelatinous in consistency, its external lamellæ easily broken down between the fingers, but a nucleus of greater firmness is found in the centre, which in some degree resists this pressure.

Crystalline  
humor.



In advanced age, the lens becomes more close and compact in texture, and the nucleus acquires a yellow or topaz color.

The texture of the lens is lamellated; the lamellæ concentric and connected by a very delicate fibrous tissue. After maceration, the crystalline breaks into triangular pieces composed of concentric scales, of which the apices meet in the centre. The anterior may sometimes be separated from the posterior part of the lens, at the line of its circumference, as if it were composed of two segments of spheres of unequal size, applied face to face. The crystalline discovers no vascular organization.

Canal of  
Petit.

The tunic of the vitreous humor, called tunica hyaloidea, has also, upon its exterior surface, a process or duplicature, membranula coronæ ciliaris of Zinn, who considered it a distinct texture. It is produced at the distance of a line's breadth from the circumference of the cup which receives the crystalline humor. At the verge of the cup the duplicatures coalesce, and thus an annular space is included between them, which has been named, after its describer, canalis Petitianus. Inflation of the canal shows that it is not of uniform dimensions; like the intestine colon, it is tacked up into cells or pouches by short transverse septa, whence the name given by Petit, canal gauderonné, or godronné. In



the grooves corresponding to these septa, the posterior edges of the ciliary processes are inserted. The intervening looser portions of the membrane correspond to the interstices of the processes; and the black radiated lines, which appear upon the membrane of the canal, are stains left by the pigment which fills them. Like the corpus ciliare, the canal is broader on the temporal than on the nasal side.

After the condensation of the lamellæ at the margin of the crystalline, the proper tunic is continued over the concave face of the vitreous humor, posterior to the crystalline lens; and a continuous transparent membrane, produced anteriorly, passes before the crystalline, so as to retain it in its place. This portion of the membrane covering the crystalline is termed capsule of the crystalline, or tunica aranea, and is considerably more dense and elastic than the proper tunic of the vitreous humor. Independent of the membranous enclosure now described, the existence of a distinct and proper capsule of the crystalline is generally assumed; but its demonstration is not altogether satisfactory. A small quantity of aqueous fluid contained in the capsule enclosing the crystalline humor, is called after its discoverer, humor Morgagnii.

Capsule of  
the crystal-  
line.

THE RETINA. The optic nerve having perforated the sclerotic and choroid coats at the in-

Retina.



terial and posterior part of the globe, terminates abruptly in a little white conical eminence or papilla. From the base of this papilla proceeds the very delicate membranous expansion termed 'retina.' It encompasses the vitreous humor, the front part only excepted. Its anterior termination is also abruptly defined, and corresponds to that of the choroid tunic which lies exterior to it. It is of exceeding delicacy, and, on dissection, resembles, in semi-transparency and in color, the ground glass of which ornamental lamps are constructed. During life it is of perfect transparency. Without caution it cannot be preserved entire in dissection; and if, when the sclerotic and choroid are divided, the parts of the globe are separated by their weight, by its strict adhesion to the other coats at its origin, it is drawn off the vitreous tunic in the form of a fine medullary rope, which expands and re-assumes its proper form in water. The arteria centralis, emerging from the axis of the optic nerve, distributes a few delicate branches upon it, which do not in the healthy adult convey red blood.

Its central  
foramen.

A minute foramen in the retina is seen on the temporal side of the optic nerve, having a yellow border, around which the arteria and vena centralis, after a delicate injection, display a vascular corona. This appearance first described by Soëmmering, "foramen centrale cum limbo



luteo," is seen only in the recent state of the eye. Its situation corresponds to the extremity of the visual axis. The membranous surface of the retina is opposed to the tunica hyaloidea, the medullary to the choroid. Its attachment, at its insertion into the ciliary body, is very slight, as it commonly yields at that part, if recent and uninjured, rather than tears by the force exerted to separate it entire. The retina is uniformly expanded over the tunica hyaloidea, but has no demonstrable connexion with that membrane.

THE TUNICA CHOROIDES extends from the circumference of the optic nerve to the margin of the exterior or flattened surface of the vitreous humor; there it terminates, together with the retina, in a greyish colored substance, termed ganglion, or ligamentum ciliare, or, better, annulus ciliaris, and which is the common centre of union for the interior membranes of the eye.

Choroid  
tunic.

The choroid is of a dusky brown color in the adult, reddish in infants, and adhering by an abundant and lax cellular tissue, which may be readily inflated, to the sclerotic coat, and by the numerous ciliary vessels and nerves, which perforate the latter to take their course upon the choroid. This cellular substance is more plentiful in the infant than in the adult, and is most abundant in the track of the principal vessels and nerves. The vessels terminating upon it

Its pigment. are extremely numerous, and secrete a dark pigment or varnish, which stains the contiguous adhering surface of the sclerotic; it likewise communicates its stain to the finger, or a piece of white paper, but the texture of the membrane is permanently dark, and is not bleached by maceration.

The interior surface of the choroid is also covered with a black varnish, thicker and deeper colored in the infant than in the adult; but, having no connexion by texture with the retina, its stain is not communicated to this tunic. Around the insertion of the optic nerve, the choroid is destitute of this dye. Residence for some time in alcohol discovers a fine white flocculent substance coating the interior of the choroid, formerly described by Ruysch as a distinct tunic (*tunica Ruyschiana*), but not regarded in this light by modern anatomists. The pigment, there can be no doubt, is secreted into a fine cellulous tissue, flakes of which are detached, in some diseased states of the organ, from the ciliary processes and back of the iris, forming to all appearance a real *membrana nigra*.

Its nerves  
and vessels.

The ciliary nerves run in parallel lines, at equal distances, upon the choroid; and from their size and whiteness are particularly conspicuous. The long ciliary arteries appear, one on either side of the globe, in their course to the *annulus ciliaris*. Beneath these the membrane



presents, on its opposite sides, vessels arranged in form of trees with weeping branches, or of the figure of a jet d'eau; these, which have been named *vasa vorticosa*, are veins returning the blood distributed to the ciliary processes, and are collected into three or four distinct venous trunks. The short posterior ciliary arteries pass under the ciliary veins, in the intervals of the trunks, to the interior of the choroid; and uniting with the anterior at the fore part of the globe, their extremities form a very intricate and beautiful net-work upon its interior surface. The adhesion of the choroid to the sclerotic is most strict, adjacent to the optic nerve behind, and the ciliary ring before, owing to the introduction of the ciliary vessels at these parts.

THE ANNULUS CILIARIS is an elastic ring Ciliary ring. composed of a short and dense pulpy texture, closely adherent to the inner border of the sclerotic, at the distance of a line and a half from the external circumference of the cornea. It is of greater breadth on the temporal than on the nasal side. The choroid and retina adjoin its greater, the cornea and iris its lesser circumference. Anteriorly it adheres firmly to the sclerotic, as before observed, and the ciliary processes are attached to its posterior surface, so that it forms a common centre of union for these tunics. Its color is observed to correspond to that of the iris.



Ciliary  
plaits or  
processes.

**THE PROCESSUS CILIARES.** On the internal surface of the choroid, at the root of the annulus ciliaris, the plicæ or processus ciliares arise in delicate striæ, and, advancing a little anterior to the circumference of the crystalline lens, terminate in a circle of fine grey points at the base of the iris. They appear to be radiated folds of the choroid tunic, from sixty to seventy in number, long and short alternately, and gathered at their origin like the plaits of a shirt at the wristband. Viewed collectively through the vitreous humor they have some resemblance to a radiated flower; a small white circle appears within a large dark one. The white lines represent the edges of the plicæ; the black, their interstices coated with pigment. These edges of the plicæ are engrooved in the duplicature of the vitreous capsule, which assists in forming the canal of Petit.

The extremities of the processes projecting from the interior border of the annulus ciliaris interdigitate with the radical fibres of the iris. To obtain a view of them, let the cornea be accurately removed at its junction with the sclerotic, and the iris be torn away, entire, from its ciliary attachment. The points of the processes will then appear, projecting like the teeth of a comb from behind the annulus ciliaris; and the ciliary border of the iris, upon floating it in water, will be found to present a corresponding arrangement.

The processes having their edges thus inlaid in the tunica hyaloidea at the margin of the crystalline fossula, and their points or anterior extremities, interlaced with the radical fibres of the iris, form a posterior iris, the aperture of which is exactly occupied by the crystalline lens and its capsule. From their origin to their insertion, they are supported exteriorly by the annulus ciliaris, with which substance they are in fact incorporated. The figure of each plica ciliaris is triangular, the internal obtuse angle being opposed to the circumference of the crystalline lens; the posterior, elongated, loses itself in the choroid; the anterior is inserted into the iris. The anterior edge is attached to the annulus ciliaris and root of the iris, the posterior to the tunica hyaloidea, and the internal and shortest measures the space between the verge of the crystalline lens and the basis of the iris; or, in other words, forms the outer boundary of the posterior chamber.

THE IRIS. This is the colored membrane <sup>Iris.</sup> which presents a plane surface traversing the globe horizontally, and dividing the corneal from the sclerotic segment. It is rendered imperfect as a septum by the pupilla or round hole in its centre. The pupil is not, however, quite central in relation to the iris, the breadth of the iris being always somewhat less on the nasal than on the temporal side. It is



divided into a ciliary and pupillary portion. Its attachment is, as already observed, by indenture with the extremities of the plicæ choroïdæ, at the inner margin of the annulus ciliaris, from which it originates. The ciliary portion of the iris is the larger one, and is composed of a delicate fibrous and vascular tissue, in which grey serpentine lines or striæ are seen proceeding like radii from the annulus ciliaris: from this the smaller pupillary portion is distinguished by a darker shade of color, and a gently elevated circular line, most conspicuous on the posterior surface of the membrane. The fibres of this portion have a similar tortuous direction, and are convergent towards the pupillar aperture. The pupillary margin is thin and defined, and presents the appearance of a dark circular line when placed upon a white ground, as e. g. the opaque capsule of the crystalline lens. The iris diminishes in thickness from its base to the margin of the pupil. Its anterior surface is richly colored of different hues in different individuals. It is thickly coated on its posterior surface by the pigmentum nigrum.

Its vessels  
and nerves.

The ciliary vessels, entering the anterior part of the globe, unite with the other detachments, and form arches at the basis of the iris and processes. From the zone thus produced (zona major) the branches run in straight lines upon the iris. In the dilated state of the pupil these

radiated vessels are tortuous ; by its contraction they become straight. At the distance of rather less than half its diameter from the pupil, another zone is formed by their anastomosis, from which branches are detached to the margin of the pupil. The zona minor gives the appearance of the undulating circular line, distinguishing the pupillary from the ciliary portion of the membrane. The two long ciliary arteries chiefly contribute to the formation of these zones and the supply of the iris. The short ciliaries, seen upon the interior of the choroid, detach numerous fasciculi to each ciliary process, which pursue a serpentine course along the fixed edge of the fold, and are inverted to form concentric arches upon its opposite free margin.

The membrana pupillaris, a delicate membrane occupying the pupil of the fœtus, and which is supplied by the vessels of the iris, disappears before birth.

Membrane  
of the pupil.

Of the proper structure of the corpus ciliare nothing is with certainty known. The notion that it is wholly constituted of vascular and nervous tissue, having no proper fibrous texture for its base, which has also been conceived of the iris, is absurdly contrary to observation and analogy. The annulus appears to be a gangliform or bulbous termination of the choroid coat, and the processes resemble plaits or doublings of



this membrane laid back to back, to accommodate it to the area of the posterior chamber. Similar uncertainty prevails as to the structure of the iris, the different opinions of its texture being founded rather upon inference from its functions than upon demonstration. If the former species of evidence be regarded, it is in part unquestionably a muscular texture; the phenomena of its action can be best explained upon the supposition that it is both muscular and elastic, and that these forces act alternately.

Sclerotic  
tunic.

THE TUNICA SCLEROTICA is the external covering of the ball, with the exception of one-fifth part, bearing a proportion to the cornea somewhat similar to that which the vitreous bears to the aqueous humor. It is a dense compact fibrous membrane, of a blueish white color; its fibres appear reticulated on maceration. It has few nutrient vessels, and no traceable nerves; its texture is both extensile and elastic. In the foetus and infant it admits of separation into two plates, but these are inseparably connected in the adult. For the entrance of the optic nerve, with the sheath of which it is intimately connected, it is cribrated or perforated with many small holes, by which the fibres of the nerve enter and terminate in the conical protuberance before described. The choroid and retina adhere firmly to the margin of this cribriform plate. In other parts the connexion between the sclerotic and

choroid is by the medium of blood-vessels and cellular tissue. The sclerotic around the entrance of the nerve, and likewise around the margin of the cornea, has many small oblique passages, of which the apertures on its internal surface are conspicuous, when separated from the choroid, for the entrance and exit of the ciliary vessels and nerves. Adjoining the cornea, the choroid and conjunctival vessels communicate through the foraminula of the sclerotic. On its inner surface it has furrows in right lines, in which the long ciliary vessels and nerves are lodged. The sclerotic is of greatest density in the vicinity of the nerve; it gradually diminishes in thickness towards the middle of the globe, where it is fortified by the tendons of the several muscles. The opening in front of the sclerotic is nearly circular, having its inner edge sloped for the broad insertion of the cornea between its anterior and posterior margins.

THE CORNEA is of a horny texture, less extensive than the sclerotic, and perfectly transparent. It is, onion-like, composed of concentric lamellæ or pellicles, connected by a delicate cellular tissue containing a transparent fluid, in which exhalant and absorbent vessels are abundantly distributed. This tissue is more lax or copious between the anterior than between the posterior lamellæ. The transparent conjunctiva upon the cornea gives a polish and brilliancy to the surface, which the lamellæ of the

Cornea.



cornea do not possess, and which is lost at the approach of death, by the transudation of the aqueous humor. They are scabrous from the adhesion of the cellular membrane connecting them, and void of lustre. The cornea is externally rather elliptical than circular, being of greater length in the transverse than the vertical diameter.

The cornea is of greater thickness than the sclerotic, in infants especially, in whom its posterior surface is contiguous to the iris. The internal surface is likewise half a line broader than the outer, the margin being obliquely extended from without inwards, to correspond with the sloped edge of the sclerotic. After maceration it may be detached from the sclerotic, to which it is connected by cellular substance ; this separation is most readily effected by plunging the macerated eye into boiling water. A fine transparent humor is secreted by colorless exhalant vessels in the areolæ of the cellular membrane between the lamellæ of the cornea. The interstitial substance of the cornea receives no colored vessels. Numerous lines have been observed to form figures of many sides between the plates of the cornea in the eye of the negro, and supposed, from a reddish tinge, to be blood vessels. The existence of nerves has never been demonstrated, and it is much to be doubted if it possess any. On its interior surface the cornea is smooth, and washed by the aqueous humor.

According to modern anatomists it is lined by a tunic proper to the humor, which is reflected from it upon the face of the iris, and advances even to the margin of the pupil. Its tenuity, if it exist, is such as very rarely to allow of its demonstration, at least in the human eye. The convexity of the cornea is greater than that of the sclerotic, being the segment of a sphere seven lines and a half in diameter.

Membrane  
of the  
aqueous  
humor,

**THE AQUEOUS HUMOR.** The name of anterior chamber is given to that space comprised between the cornea and the iris, ordinarily about one line and a half in depth. The posterior chamber, not exceeding a quarter of a line, is the space between the iris and the crystalline lens. They communicate by the aperture of the pupil, and both are occupied by the aqueous humor. This is a transparent fluid, evaporates on exposure to heat, and is uncoagulable by heat, acids, or alkalies; it is in quantity about five grains; in quality, viscous and slightly saline. It gives figure and tension to the cornea, keeps the pupil properly dilated, and supports the parts forming the parietes of both chambers. When discharged by the puncture of the cornea, the pupil contracts, and the chambers are obliterated by the collapse of their parietes: it is however reproduced in a few hours. The aqueous humor in fœtuses and new-born infants is turbid, and sometimes of a reddish tint.

Aqueous  
humor, and  
chambers of  
the eye.



Veins of the  
globe.

The ciliary veins and vasa vorticosa of the choroid jointly return the blood distributed by the ciliary arteries.

They perforate in like manner the sclerotic coat, and terminate in the infra-orbital branch and trunk of the ophthalmic vein, which also receives the vena centralis retinae.

### ORBITAR APPENDAGES.

Periosteum  
and fat.

THE PERIOSTEUM AND ADEPS OF THE ORBIT.  
The dura mater, which is the internal periosteum of the cranium, lines the orbit, and is continuous at all its openings with the periosteum of the head and face; hence the extensive sympathetic pains in the inflammatory affections of the bones of the face and cranium, and their common membrane. Hence also probably, the suppurative inflammation of the dura mater after extensive fractures and injuries of the orbit.

The fat, which in health is secreted abundantly in the orbit, surrounds the optic nerve, and invests the posterior surface and sides of the globe, forming for it a soft bed, and defending the vessels and nerves from compression in its motions. In emaciating diseases its diminution by absorption produces that characteristic sinking of the globe in its socket, and loss of convexity in the eyelid, which is familiarly expressed by the term "hollow eyed." On the other hand,

its secretion in excess, as in morbid obesity, protrudes, compresses, and thus induces congestion in the vessels of the eye.

I proceed to describe the muscles, vessels, and nerves contained in the orbit.

THE MUSCLES are seven in number ; viz. the levator palpebræ ; the rectus superior, inferior, internus, and externus ; obliquus superior and inferior. Muscles.

The levator palpebræ has an acute origin from the periosteum above the foramen opticum ; its fibres spread in their course, giving it a fan-like shape, and they are inserted by a broad aponeurosis in a condensed cellular substance, which connects the upper tarsus to the orbital ridge, between the conjunctiva and the fibres of the orbicularis palpebrarum. From the nature and extent of its connexion with the eyelid, it results that the partial division of the tarsal ligament, or even the removal of the cartilage, does not take away the power of elevating the lid as the paralysis of this muscle does ; the elevation, however, under these circumstances, is imperfectly performed. Elevator of the upper eyelid.

The rectus *superior* lies beneath this muscle, arising from the border of the foramen opticum and the partition between it and the foramen lacerum. Superior straight muscle.



Internal and  
inferior.

The rectus *internus* and rectus *inferior* arise in common from a ligament which in part surrounds the optic foramen, and fills up the foramen lacerum.

External.

The rectus *externus* arises by two distinct heads: the inferior having a common origin with the last named muscles, from the ligament which occupies the inferior angle of the foramen lacerum; the superior from an arch of ligament crossing the foramen above. It is important to note this bicipital origin of the rectus externus, as some of the nerves of the orbit pass through the interspace between its heads, and others through the top of the foramen. The ligament of the foramen spheno-maxillare forks into three intermuscular slips, which give origin and support to the external, inferior, and internal recti muscles, in the manner of the intermuscular ligaments of the extremities. The four recti muscles, varying in length and direction as the sides of the orbit to which they are adjacent, pass over the great circumference of the bulb, between which and the cornea they are inserted, at equal distances, by straight tendinous fibres, into the substance of the sclerotic coat.

Superior  
oblique.

The superior *oblique* muscle rising from the periosteum between, and a little anterior to the origins of the superior and internal recti, passes its slender rope of tendon through a half ring of

cartilage which is affixed by a ligament to the os frontis, a little above and behind its internal angular process. The trochlea is provided with a sacculus mucosus, and the tendon emerging from it is enclosed in a ligamentous sheath to its insertion in the sclerotic coat, at the posterior and upper surface of the globe, beneath the superior rectus muscle.

The inferior *oblique* rises from the orbital plate of the superior maxillary bone, behind the lacrymal fossa, and takes an oblique direction between the globe and rectus inferior, to its posterior and outer surface, where it is likewise inserted into the sclerotic.

Inferior  
oblique.

The single actions of the recti are expressed by the terms levator, depressor, adductor, and abductor. Their co-operation retracts the globe in its socket. The oblique muscles, acting singly, roll or rotate the eye in contrary directions. Their co-operation antagonises that of the recti, which power is demonstrated by the course of the superior oblique, the origin of the inferior, and their posterior insertions.

Their ac-  
tions.

THE ARTERIES of the eye are principally derived from the ophthalmic artery, which has a short but sharp curve at its origin from the internal carotid, before it enters the orbit. This it does through the foramen opticum, upon the temporal side of the nerve.

Arteries.



The *arteria centralis retinae* which runs in the centre of the optic nerve, and the *long ciliary* arteries which pass upon either side of it, are its first branches; next the *lacrymal* artery, which contributes a ciliary branch, a branch to the rectus externus, and another which runs beneath the globe to the obliquus inferior muscle and reaches the inferior palpebra. The lacrymal branch then divides into two: one, a branch of communication with the deep temporal branch of the internal maxillary artery at the outer margin of the orbit; another, which is dispersed in the substance of the lacrymal gland and superior palpebra.

The trunk of the ophthalmic artery then crosses obliquely beneath the optic nerve, and on the nasal side of the nerve sends branches to the superior oblique and levator palpebrae, rectus superior and inferior muscles, and commonly a ciliary artery. The remaining branches of the ophthalmic artery, which is here tortuous, are the *frontal*, through the supra-orbital foramen; one or two to the rectus internus, the *nasal* branch which passes by the anterior æthmoid foramen into the nose, and the *infra-trochlear* branch. The ophthalmic artery at length emerges upon the inner canthus, furnishing the *superciliary* and *palpebral* branches, and anastomoses with the nasal branch of the facial artery from the external carotid. The muscular branches penetrate between the fibres, and run-

ning in the same direction, appear beneath the conjunctiva on the sclerotic coat. Here they subdivide and ramify upon the conjunctiva; the fasciculi inosculating so as to form a faint circulus arteriosus around the cornea, when filled with colored blood. Those of the rectus internus are most numerous.

**THE VEINS.** The ophthalmic and nasal Veins. branches of the anterior division of the facial vein, freely communicate at the inner angle of the orbit with the *ophthalmic vein*; and the anterior and posterior *æthmoidal* or *nasal*, the *lacrimal*, all the *ciliary veins* from the globe, the *vena centralis retinæ*, the *infra-orbital*, the several muscular, periosteal, and adipose branches, are all collected into this trunk in its passage through the orbit. It takes a serpentine course over the optic nerve, through the foramen lacerum, to terminate in the anterior part of the cavernous sinus of the dura mater.

**THE NERVES** of the orbit, exclusive of the optic, are the third pair, or *motores*; the fourth, or *pathetici*; the first division of the fifth, or *trigmini*; and the sixth, or *abducentes*. Nerves.

The third pair enter the orbit between the heads of the rectus externus muscle, in company with the nasal branch of the fifth and the sixth pair. Its lesser and superior branch rising



before its entry into the orbit, joins a twig of the fifth pair, to assist in forming the *ophthalmic* or *lenticular ganglion*, and then divides to supply the rectus superior and levator palpebræ muscles. Its larger and inferior branch passes under the optic nerve towards the nasal side of the orbit; and while covered by that nerve, is divided into a branch to the rectus internus, a short thick stalk to the *ophthalmic ganglion*, and a long slender filament to the inferior oblique muscle.

From the *ophthalmic ganglion* lying concealed in fat, on the temporal side of the optic nerve, a superior and inferior fasciculus of *ciliary* nerves arise and creep along its side in a serpentine direction to the bulb.

The fourth pair of nerves, with the lacrymal and frontal branch of the fifth, pass through the upper part of the foramen lacerum. It sometimes receives a branch of augmentation from the fifth pair, and always increases in size towards its termination in the central fibres of the superior oblique muscle.

The first or *ophthalmic* division of the fifth pair gives off,

1st. The *supra-orbital*, which is subdivided into the supra and infra trochlear ramuli, and

the proper frontal nerve; which last, running in an external and internal branch upon the levator palpebræ, is distributed upon the forehead.

2dly. The *lacrymal* nerve, which, taking a direction to the outer canthus, splits into an external and internal branch. The internal supplies with filaments the glomera of the lacrymal gland; twigs from the external likewise enter the gland, and together they are dispersed upon the superior palpebra.

3d. The *nasal* nerve, which gives a branch to unite with the short stalk of the third pair to form the ophthalmic ganglion, and contributes two *long ciliary* nerves to the globe, then passes obliquely under the superior oblique muscle to the æthmoid foramen, by which it enters the nose, furnishing an infra-trochlear filament to the nasal region of the orbit.

The sixth pair, having parted with the filaments supposed to be the roots of the great sympathetic upon the canalis caroticus, enters the orbit with the nasal of the fifth and the third pair, to be spent upon the rectus externus muscle.

THE LACRYMAL GLAND is of the conglomerate kind, of a flattened oval form, divided by a cleft into two lobes, of which the superior and internal is the smaller and thinner, the inferior and exter-

Lacrymal  
gland.



nal the large extremity of the gland. Its position is oblique; the inferior and internal surface hollowed to suit the convexity of the globe; the superior convex to fit the corresponding surface of the orbit to which the gland is attached, by a ligament passing transversely beneath it. It measures, in length, about ten lines; in breadth five or six. The structure of the gland resembles that of the salivary, its lobules connected by a dense cellular tissue, upon which its vessels and nerves subdivide, to supply the granules of which they are composed. The vessels enter the gland at its posterior margin; and from the anterior, its ducts, five or six in number, pass out in straight lines, and pierce the conjunctiva at the orbital edge of the superior tarsus.

We have now briefly described the orbital appendages, or those parts situated behind the tunica conjunctiva, and proceed to the

### FACIAL APPENDAGES.

*Eyebrows*

**THE SUPERCILIUM, OR EYEBROW.** The arch of the eyebrow corresponds to that of the superciliary ridge upon which it is planted. It extends from the tuberosity of the frontal sinus to the external angle of the orbit. It consists of a thick row of strong short hairs, which have a disposition almost erect at the commencement of the brow, and are then arched obliquely outward, and gradually reduced in number so as to

terminate the arch acutely. The few erect hairs correspond to the fibres of the corrugator supercilii muscle, the crescentic to the fibres of the orbicularis palpebrarum.

The extent and fulness of the brows vary greatly in different persons. In some, especially persons of dark complexion and black hair, they have little if any interspace at their origin, and are long, prominent, and bushy in the centre of the arch. Among the ancients these were esteemed points of female beauty. The fibres of the occipito-frontalis, or epicranial muscle, terminate beneath the skin of the supercilium, blending with those of the orbicular muscle of the palpebra. The former elevates the brow, wrinkling the integument of the forehead horizontally: the latter depresses it, and closes the eyelids, being the sphincter palpebrarum.

The corrugatores approximate the heads of the supercilia, drawing the integuments over the root of the nose into deep longitudinal rugæ: they co-operate with the orbicularis in the act of frowning. The action of the subjacent muscles renders the brow an important feature in regulating the quantity of light, contracting the field of vision, and in assisting the expression of the sterner passions. It would not be a useless ornament if it were insusceptible of motion, the hair being advantageously placed upon the



projecting ridge of the orbit to entangle and arrest particles, solid and fluid, which might otherwise fall or trickle upon the eye. The habitual depression of the brow is usually a concomitant of a weak or morbid retina; it is a characteristic of strumous inflammation, and is observable in all cases where light is offensive, and in those central circumscribed opacities of the cornea and lens, in which the dilated state of the pupil is necessary to vision.

Eyelids.

THE PALPEBRÆ, OR EYELIDS, are those semi-oval curtains which cover the great aperture of the orbit, and graduate the light falling upon the eye by the degree of their separation, or exclude it by their apposition. The skin covering the palpebræ is thin, and loosely connected to the subjacent parts by a fine lax cellular texture, which abounds at the orbital margins of the palpebræ. The frequent œdema of the eyelids, so disfiguring to the countenance, is owing to the abundance of this tissue void of fat, and subject therefore to serous infiltration.

The superior is broader than the inferior palpebra, covering two-thirds of the surface of the globe by its descent. It is also more moveable, the inferior palpebra being inconsiderably elevated to meet it in shutting the eye. The superior palpebra, when drawn up, makes a doubling or deep crescent-shaped fold in the skin

under the orbital arch, which is effaced when the palpebra falls. Upon the skin of the lower eyelid narrow and gently curved rugæ are seen; these, which are signs of the unequal contractility of the skin and the muscular fibres beneath it, are more strongly marked in persons of advanced years, in whom the muscles have been longer and more vigorously employed, and whose skin is loose or redundant from the absorption of the adeps beneath it.

When the eyelids are forcibly closed by the contraction of the sphincter fibres of the orbicularis palpebrarum, the tendon of this muscle starts forward, and the rugæ are extensively radiated from the nasal angle over the skin of the cheek. On removing the skin and the subjacent cellular tissue of the palpebræ, the thinly spread fibres of the orbicularis muscle are seen. The tendon with which these fibres are connected is a little round cord, distinctly seen and felt beneath the tegument, implanted in the nasal process of the maxillary bone, in the great transverse diameter of the orbit. The fibres which lie upon the palpebræ are the interior fibres of the muscle, the fissure of the lids being the axis of the oval formed by it. The inferior external fibres, from the round tendon and contiguous parts of the maxillary bone, take an extensive sweep over the orbital ridge upon the cheek, towards the temple, where they become thin and scattered. The

Orbicular  
muscle of  
the eyelids.



superior from the round tendon and contiguous part of the frontal bone, take the direction of the superciliary arch; being at their commencement connected with the fibres of the corrugator, and in their course blended with those of the frontal muscle. The integument of the palpebræ is adherent to the tendon of the orbicularis, which has been described as the ligament of the palpebræ or tarsi; and the angle of union between them is larger in consequence of this adhesion, than the external. The inner canthus, formed wholly of the doubling of the integument, is that notch, or triangular sinus, formed between the tarsi and the tendon of the orbicularis.

Tarsal cartilages.

THE TARSI are two elliptical cartilages which give figure and firmness to the palpebræ, and afford a basis for the attachment of their several parts. The superior is broader than the inferior. Their opposite edges are broad and sloped from without inward; their orbital edges are thin and continuous with a condensed cellular membrane, which is ligamentous where it is inserted into the orbital circumference, and especially at the temporal side. Their temporal extremities are angular, the nasal rounded. The former terminate the fissure of the palpebræ at the temporal angle; the latter, which are opposed to each other, and brought into contact when the lids are closed, are situated at the distance of two lines from the nasal angle to which they are

connected by the doublings of integuments forming the borders of the inner canthus. The convex surface of the tarsi is covered by the fibres of the orbicularis and the cellular membrane connecting them with the integument; the concave, which is exactly moulded to the face of the globe, is covered by the *membrana conjunctiva*.

**THE CILIA, OR EYELASHES.** The exterior borders of the sloped edges of the tarsi, which are opposed to each other, are furnished with cilia, or eyelashes, disposed in three or four rows: these we may therefore call the ciliary borders of the tarsi. The apertures, in which their bulbs are contained, are seen in the integument when the cilia are extracted: they are more numerous and longer upon the centre than the extremities of the tarsal arch, and fewer and shorter on the lower than the upper tarsus. Their direction is curved, those from the upper being arched upwards, from the lower downwards. The length and fulness of the eyelashes vary in different individuals. They are commonly of the same color as the eyebrows.

**THE MEIBOMIAN FOLLICLES.** Upon the interior border of the tarsus the mouths of a row of follicles, seated vertically on the concave surface of the tarsus, form a slightly eminent line. These follicles, when magnified by a glass, ap-

Eyelashes.

Follicles of  
Meibomius.



pear to be small knotted tubes, resembling studs of the smallest pearls, arranged for the most part in parallel lines, and communicating with each other at their origin from the orbital edges of the tarsi, but terminating by distinct orifices upon their interior borders, which we may distinguish from the ciliary as the meibomian borders of the tarsi. In their length, connexion, and arrangement, they present considerable variety. The fluid, which they secrete, may be expressed in a condensed form in diseased states of the follicles, or, after death, in the shape of small white worms. It is an unctuous fluid lubricating the tarsal edges, preventing the effects of attrition from their frequent contact, and facilitating their motions over the contiguous surface of the globe.

Lacrymal  
conduits.

THE PUNCTA LACRYMALIA, SACCULUS, AND DUCTUS LYCRYMALIS. The obliquity of the tarsal edges, which are opposed to each other, leaves a groove, or sulcus, between the meibomian borders and the surface of the globe, when the ciliary borders are in contact. This increases in breadth towards the nasal angles of the tarsi, where the puncta, or orifices of the lacrymal excretories, are placed upon two small conical eminences accurately opposed, and terminating the meibomian borders: they are two pinholes formed in the cartilaginous substance, and thus preserved permanently open. Their course is at

first perpendicular to the tarsi, in which direction they severally form a short cul de sac ; then they are turned at right angles towards the nose, and lie in the doubling of the skin, forming the borders of the inner canthus. They are, if we except their orifices, purely membranous canals leading to the oblong membranous sac situated in the lacrymal fossa ; they terminate side by side, or more frequently, in a common duct, in the upper and anterior part of this sac, under the tendon of the orbicularis.

The position and direction of the lacrymal sac correspond to the fossa formed by the anterior concave portion of the os unguis, and the nasal process of the superior maxillary bone, upon which it is seated ; it rises a little higher than the termination of the lacrymal duct or ducts, and the transverse tendon of the orbicularis crosses the upper part of it. Below it contracts into a duct, which occupies the canal formed by the nasal process of the maxillary bone, and the spoon-shaped process of the os spongiosum applied to it. The course of this duct is downward, outward, and gently slanting backward ; it opens by an oblique fissure under the convex scroll of the spongy bone, in the side and near the floor of the nostril. A probe, introduced from the nostril into the nasal duct, must have a sharp curvature to enter it with facility.



The membrane of the sac and duct is intimately adherent to the bony parietes. The anterior and exposed surface of the sac is defended by a fibrous or ligamentous expansion, derived from the circumference of the lacrymal fossa. The orbital is distinguished from the nasal portion of the duct by a fold, or duplicature of its lining membrane, and another similar fold sometimes occurs in the nasal part of the duct. The membrane of the lacrymal canals and nasal duct is abundantly furnished with mucous follicles, or lacunæ. The fulness of the membrane, where it is reflected from the nasal extremity of the duct, greatly diminishes the diameter of the bony aperture, and gives it a valve-like form. The diameter of the lacrymal and nasal canals exceeds that of their orifices. The puncta absorb the tears, which have been conducted from the lacrymal ducts by the tarsi, and convey them into the sac to pass off by the nasal duct.

Conjunctive  
membrane.

**THE TUNICA CONJUNCTIVA.** The integument of the eyelids is inflected at the edge of the tarsi, and lines the whole of the concave surfaces of the palpebræ; is reflected upon the visible face of the globe, enters into the puncta, lines the lacrymal sac, and at the nasal extremity of the duct is continuous with the common mucous membrane of the nostrils, fauces, and alimentary canal.

The conjunctiva, having lined the interior surfaces of the tarsi, is connected to the ligaments of the tarsi and palpebral muscles, and thence reflected upon the globe, so as to form an oblong sac or pouch. Its attachment to the sclerotic is such as to prevent its forming folds in the motions of the globe, to the freedom of which it offers no impediment. As it approaches the cornea, its attachment becomes more strict, and at the margin of that membrane it is inseparable from it. Its continuity is ascertained by dissection, but its tenuity and transparency are increased, and when held to the light it has a nearer resemblance to a very delicate lamella of the cornea than to the conjunctiva of the sclerotic. After maceration, the separation is more readily effected.

The character of this membrane is so materially modified by its several relations with the integument, the tarsi, the sclerotic, and the cornea, that its continuity alone establishes its identity. The fact of continuity is, however, corroborated by some pathological phenomena, which so often illustrate problematical points in anatomy and physiology. For example, the conjunctiva furnishes the matrix for the adventitious vessels, which are created to repair breaches of the corneal texture. These vessels, whether formed by the healing process, or open-

Its continuity demonstrated by disease.



ed by long continued diseased action, as in chronic ophthalmia, are obviously superficial. The circumstances by which they are produced, are characterised by different appearances, as I shall hereafter point out. Again, when a small portion of the conjunctiva is abraded by an extraneous particle, the scabrous surface of the cornea is exposed, and ulceration of this surface ensues. The deficiency of the conjunctiva is exactly depicted by the margin of the abrasure, and the contrast of the surfaces. This is very dissimilar to the interstitial ulcer of the cornea. The pterygium, a rare disease, exhibits the continuity in a very striking manner. It has a full broad base next the canthus, where the conjunctiva lies loose, and is gradually flattened and drawn to a point, so as to have a wedge-like form as it approaches the cornea. But although the deposition is beneath the conjunctiva, it does not stop at the cornea, but slowly travels across it. The strictness of the adhesion alters its appearance; the lymph shed between the conjunctiva and cornea, presenting only a progressive dense opacity, instead of the fleshy elevation which it exhibits upon the sclerotic. The continuity of the superjacent texture is demonstrable.

Upon the tarsi the membrane is closely adherent, and although transparent, appears of a pale red tint; upon the sclerotic and cornea it

is colorless. The sclerotic conjunctiva, however destitute of red vessels in the tranquil state of the organ, becomes conspicuously vascular and acquires a deep red color by inflammation, its minutest capillaries appearing to convey red blood, in the vehement acute ophthalmia. Those of the corneal conjunctiva are only to be seen, when, by continued distention, the connection is loosened between the conjunctiva and cornea. In this case, the cornea exhibits red vessels freely inosculating from its opposite sides, and anastomosing with each other. The increase in number and extent of these vessels is a gradual process, demonstrable to observation, and the inflammatory action which precedes this state, is ordinarily of considerable duration. The incapacity of the vessels of the corneal conjunctiva to receive red blood, seems to depend upon the strictness of its adhesion.

THE VALVULA SEMILUNARIS, AND CARUNCULA LACRYMALIS. The conjunctiva is attached to the canthi of the eyelids, and, at the internal canthus, forms a semi-lunar duplicature in shape of a valve. The horns of this crescentic fold are lost in the sinus palpebralis, or angular fold of the conjunctiva.

Semi-lunar  
fold and lacrymal  
caruncle.

On the fore-part of this valve, a small red glandular body, caruncula lacrymalis, is seen, occupying the hollow of the canthus. The ca-



runcula is a granulated substance, of a conical form and a deep red color. The base of the cone is next the orbit, the apex towards the eye. A few fine hairs are scattered over its surface. It is made up of a congeries of minute follicles, secreting that mucus which accumulates during sleep in the form of a gummy matter, at the inner corner of the eye; and appears to perform a similar office to that of the meibomian glands, which are confined to the tarsi.

From the above description it will be understood, that the palpebra, the anterior hemisphere of the eyeball, and the lacrymal passages, are every where covered by the reflected integument, modified in its disposition and qualities as its economy requires, which invests the organs of sense, the hollow viscera, and forms the external covering of the body. It is by the continuity of this membrane that the sympathy is established between these surfaces, healthy and morbid, remote and contiguous, and that the diseases with which they are affected have for the most part a common character. It remains only that I should point out the origin and disposition of the superficial vessels and nerves, by which the palpebræ are supplied.

Vessels and  
nerves of  
the eyelids

THE ARTERIES, VEINS, AND NERVES OF THE PALPEBRÆ. A superior and inferior branch, de-

rived from the ophthalmic artery, at its egress from the orbit, course along the orbital edges of the tarsi, and form by inosculation at the external angle, a complete arcus palpebralis. A superciliary arch is also formed by the union of the superciliary artery, from the ophthalmic, with the temporal. The nasal branch of the facial artery assists in forming these arches, and freely communicates with the frontal branch of the ophthalmic. The superior coronary, transverse facial, infra-orbital and temporal artery, participate in the supply of the palpebræ.

THE VEINS, beginning by small radicles from the opposite margins of the tarsi, form an intricate plexus beneath the skin of the palpebræ, and are collected into the facial, supra-orbital, and deep temporal vein. The arteries pass in the direction of the orbicular fibres, the veins cross them at right angles; their direction according to the breadth of the palpebræ.

THE NERVES take a direction similar to the veins, the frontal branch of the fifth pair supplies the superciliary and superior palpebral branches; and the infra-orbital, or first branch of the superior maxillary nerve, gives off three principal branches, which turn round the trunk of the facial vein to be dispersed upon the lower eyelid.



For the simple and general view which I have taken of the subject of the foregoing description, if any apology be necessary, I know of none more appropriate than that contained in the following paragraph.

“The study of anatomy, as it leads to the knowledge of nature and the art of healing, needs not many descriptions nor minute dissections; what is most worth knowing is soonest learned, and least the subject of dispute; while, dividing and describing the parts, more than the knowledge of their uses requires, perplexes the learner, and makes the science tedious, dry, and difficult.” *Cheselden's Preface to his Anatomy.*

## SKETCH

OF THE

## PHYSIOLOGY OF THE EYE,

## AND ITS APPENDAGES.

---

It is not my intention to enter into an abstract discourse on the phenomena of vision, a subject more allied to philosophy than medicine; but the preceding sketch would be incomplete without some account of the functions of the organ, and the history of its diseases would want the illustration which a competent knowledge of its economy conveys.

I shall suppose the reader acquainted with the prevailing opinions concerning the origin and nature of light; the velocity of its movement; the meaning of the terms direct, reflected, and refracted rays; the equality between the angle of reflection and the angle of incidence; and the facts that refraction is in-

Preliminary  
positions.



creased according to the relative density of bodies, and that the convergence of rays after refraction is proportionate to the curvature of the surface through which they pass: further, the decomposition of light by the prism into seven elementary colors, which differ in their refrangibility; the reflection of all the rays together producing the appearance of white; while their partial reflection occasions the various diversities of color, and their total absorption the sensation of black, which is in fact but the absence of color: lastly, the emission of the rays of light from every visible point of the surface of a luminous body, and their divergence thence so as to form a cone, of which the apex corresponds to the point from which they emanate, and the base to the surface upon which they impinge.

Influence of  
the several  
textures upon  
the rays  
of light.

The operation of the cornea upon the rays of light is to render them convergent towards the retina, by reason of the sphericity of its surface, and its greater density compared with the atmospheric medium through which they pass. The rays which fall within an angle of  $48^{\circ}$ , or thereabouts, measured on the surface of the cornea pass through it, and are refracted in their passage. Those which are not included within this angle are reflected by the verge of the cornea and the sclerotic coat. The aqueous humor, being of inferior density to the cornea,

diminishes in some degree the convergence of the rays which proceed through it, so that the total effect is nearly the same as that which would result from the refractive power of the aqueous humor alone, if the cornea had not existed. The rays which lie remote from the visual axis, are not transmitted through the pupillar aperture, but are reflected by the iris, and in part absorbed by the pigment coating its posterior surface, without which pigment it would be diaphanous, as in the albino. The superior density of the crystalline co-operates with its curvature to increase the convergence of the rays which are admitted within the pupil; and by their passage through its posterior surface, this convergence is increased, because they arrive at the vitreous humor, which is a medium of inferior density. By the operation of all these causes the rays are collected into foci upon the retina, and that part of the object from which the rays proceed is painted upon this membrane.

The result, then, it appears, of a series of refractions of the rays of light in passing through the humors of the eye, is their collection into foci upon the retina, so as to form a complete picture of the external scene.

From this account it will be perceived that each pencil will consist of a double cone of rays,

Inversion  
of the image  
on the  
retina.



the axes of which are right lines, their bases meeting in the crystalline and their apices being situated in the object and in the retina respectively. The rays from the top of the object are deflected to the bottom of the eye, and those from the side of the object to the right of the observer, are deflected to the left side of the eye, and vice versa; hence the inversion of the picture upon the retina. The following simple experiment, demonstrating this fact, is well known. A portion of the coats being removed from the back of the eye, and their place supplied by a piece of oiled or tracing paper, the flame of a candle placed before the cornea is exhibited of diminished size and inverted. We infer that this image excites the perception of the object, because distinct vision is enjoyed only in such conformations and conditions of the eye, as allow of its being accurately formed and impressed.

Correction  
of aberration from  
unequal refraction.

The necessary effect of the spherical figure of the cornea is to occasion an unequal refraction of the rays which permeate it, and hence to create a degree of aberration which would confuse vision. This is corrected in two ways:—first, by the gradually increasing density of the lens from the circumference to the centre, and its consequently refracting with less power those rays which arrive at it with a considerable

obliquity; and secondly, by the mobility of the iris, which adapting the size of the pupil to the circumstances of the case, excludes more or less those rays which would produce aberration.

I have stated that the iris serves to arrest those rays which are denied admission through the pupil: they would be unequally refracted **by** those points of the lens through which, if uninterrupted, they must pass, or would fall so obliquely on the cornea as to be subjected to too great a refraction. This is its passive function; but by its power of dilatation and contraction, in obedience to the stimulus of light upon the retina, it determines the quantity necessary for the purpose of distinct vision. In regulating the quantity of light the iris assists materially in accommodating the eye to different distances; in viewing a distant object the pupil dilates, and in viewing a near one it contracts. It is true that viewing the sun occasions a contraction of the pupil, and the steadfast vision of a near object in deficient light, its dilatation. These are confirmations of the statement that its motions are in obedience to the impression of light upon the retina, because the direct emanation of light from its source in the one case, and the insufficient light in the other, render these objects analogous in this respect to the nearest and the remotest visible objects. But under ordinary circumstances, the illumination of ob-

Office of the  
iris.



jects being conformable to the distance, the pupil, in viewing a distant object, is dilated so as to admit as many rays of the enfeebled light as is necessary to the distinct perception of the object; and on the other hand contracts, to exclude the superfluous rays, which coming from a near object, would otherwise create confusion. Let a person survey the sun whilst the pupil is fully dilated by belladonna, or under the same circumstances, the flame of a candle, brought near to the eye, and in either case he will find his vision confused to dimness. But the fullest permanent dilatation of the pupil will not injure the clearness of his vision of any other remote object; though that of all near objects will be in a degree confused, and the confusion be increased in proportion to the degree of their illumination. Where the iris is from any cause motionless, the power of adapting the eye to distances is lost. I conclude, therefore, that the adaptation of the eye to light co-operates with its adaptation to distance.

Corrective  
power of  
the lens.

By the peculiar constitution of the crystalline lens, before adverted to, its refractive power is so adjusted to that of the contiguous aqueous and vitreous humors, as to correct the aberrations which the figure of the cornea would occasion, and to throw the most oblique pencils of rays with sufficient accuracy upon the concave face of the retina.

Notwithstanding that man, compared with animals, requires the largest quantity of light for vision, the images of objects on his retina are undisturbed by reflection, owing to the absorbing quality of the dark pigment: which being spread over the whole interior of the globe, renders the eye a most perfect camera obscura. Animals, in whom this pigment is a brilliant reflecting surface, have the advantage of seeing in feebler light, and this power is in proportion to the whiteness of the pigment; but the accuracy of their vision, it may be presumed, is in the same proportion defective.

and use of  
the pigment.

Not only is the clearness of the image undisturbed by superfluous light, but it is also destitute of color, the decomposition of light by irregular refraction, being in ordinary vision prevented or corrected by the structure and curvature of the crystalline lens. Light, artificially separated, either by refraction, reflection, or inflection, produces color; but the light which arrives at the eye in its natural combination of elementary rays, undergoes no such decomposition in its passage through the humors.

Achromatic  
power of  
the lens.

The retina is equally expanded over the vitreous humor, but the field of vision is limited. This is not, however, confined to the axis of vision; for in certain positions of the eye, ar-

Field of  
vision.



tificially induced, we have a clear perception of an object from which the rays pass so obliquely as to fall upon the retina not in the axis of vision. It has been long observed, that if an image fall upon a certain spot of the retina, the perception of it is obscured. This spot, about one thirtieth of an inch in diameter, corresponds to the entrance of the optic nerve.

Magnitude  
of the  
image.

The magnitude of the image formed upon the retina, is proportional to the angle which the two extremities of the object viewed subtend with the centre of the eye. Hence, the more remote the object the smaller the image.

Duration of  
the impres-  
sion.

The duration of the impression made upon the retina is in proportion to the strength of the impression; this is illustrated by the appearance of a fiery circle produced by the rapid revolution of a lighted stick. The principal phenomena of ocular spectra admit of an explanation in some degree similar; as for example, the appearance of a luminous halo after looking intently at a colored object, remaining even after the eyelids are closed.

Distance,  
size, and  
position of  
objects.

The eye possesses no absolute power of determining the actual distance, magnitude, and position of objects. Such knowledge is relative, and results from the experience derived from

the combined agency of the senses of sight and touch.

It remains that I should advert to what may be termed the problems of vision, but as I have little from my own observation to offer upon these subjects, and as their investigation is in a considerable degree connected with the department of physical optics, or of metaphysical speculation, I shall be excused for touching them lightly.

The images of objects are inverted upon the retina, yet we see them, as they are in nature, erect.

*Inversion of  
the image.*

If we look in a concave mirror, objects appear inverted. The image formed upon the retina is in this case erect, and we see the object in the same relative position to the image, as all other objects. Of this fact any one may convince himself, by preparing an eye, as before mentioned, and placing beside and a little behind the flame of the candle a spoon, the hollow of which reflects it inverted, when he will observe, on the opposite side of the oiled paper, the images of the real and the reflected object, the first inverted, the second erect.

It has been generally supposed that we actually see objects inverted, and that this error of the sight is corrected by experience. Some, on

*Common  
theories.*



the contrary, have supposed that the mind acquires the perception of objects, not from the picture upon the retina, but from the object itself, by retracing the direction of the pencils to their points of radiation. Others assert, that a decussation of the fibres of the optic nerve corrects the erroneous impression before it is presented to the sensorium.

Berkley's  
theory.

The celebrated explanation of Berkley, in so far as it admits of an abridged exposition, is as follows. Visible and tangible ideas occupy distinct provinces, and have originally no affinity to each other. It is only by experience that they become connected. The impressions on the organ of sight suggest by association the ideas of objects acquired by the sense of touch, just in the same way as the word used to denote an object immediately suggests the idea of that object, to a person who is familiar with the language. The image on the retina is merely the instrument, not the object of vision. Its position has originally no influence on the ideas we form of the situation of external objects; and the supposed difficulty in the case of the inverted images arises from confounding ideas derived from the sense of touch with those derived from the sense of sight\*.

\* A person born blind and suddenly restored to sight, is the case supposed by Berkley and other writers, and so happily exemplified by Cheselden. Such a person, it is clear, would gain nothing by the aid of sight, until the connection

The association of ideas, derived as they are from the external senses, operates imperceptibly to an extent that we have no means of ascertaining, because the original and absolute negation of each sense in succession, so that each should be in turn insulated, is an impossible condition, notwithstanding the seemingly possible independence, in a state of society at least, of the animal and vital functions. Touch, in the extended sense of physical feeling, is the basis of all; sight, hearing, smell, and taste, like the sense of touch itself in its strict and limited import, are but modifications of it. That either

Intercourse  
of the  
senses ne-  
cessary to  
their deve-  
lopement.

between touch and sight grew up and established itself in his mind.

“L’objet propre et immédiat de la vûe n’est autre chose que la lumière colorée : tout le reste, nous ne le sentons qu’à la longue et par expérience. Nous apprenons à voir précisément comme nous apprenons à parler et à lire.—*Voltaire, Physique Newtonienne, Chap. 7.*

Our Shakespeare who ‘needed not the spectacles of books to read nature, but looked inwards and found her there,’ puts this distinction with admirable force and shrewdness, in the dialogue between Gloster and the fellow who feigned to be cured of his native blindness at St. Alban’s shrine.

GLOSTER.

Saunder, sit there, the lying’st knave in Christendom

If thou hadst been born blind,

Thou might’st as well know all our names, as thus

To name the several colours we do wear.

Sight may distinguish colours :

But *suddenly* to nominate them all,

It is *impossible*.

2d Part of KING HENRY VI. Act 2.



or all of these therefore should be wanting, is not incompatible with their constitution; but the sense of contact is so essentially and indivisibly incorporate with the organic nervous system, that its negation would be paramount to acephalous monstrosity. Hence its influence as a substitute and corrector in relation to the rest, when wanting or imperfect, can never be fully appreciated, because it cannot, like them, be subjected to analytical test. But from what we see of the effects of privation of one or more of the external senses, and of their reciprocity in general towards each other in cases of malformation and disease, is it not in the highest degree probable that their natural intercourse and co-operation are essential to the developement of each respectively? To illustrate my meaning—If it be possible to suppose a case in which the eye was the only external organ of sense, would the unfortunate possessor have any distinct idea of visible objects; or, *mutatis mutandis*, the ear of sounds? Certainly not. Dumbness is in most cases only a consequence of the absence of hearing; the organs of speech are perfect: so the loss of visual perception (not of light more than of unharmonized articulation) would result in the case supposed, from the absence of the associated sensations and ideas thence derived.

Single  
vision.

How it happens that impressions made upon our two eyes at one and the same time are represented single to the sensorium, we know as

little as why we hear one sound with two ears, and smell one scent with two nostrils. The mind is incapable of receiving two distinct impressions at the same instant. The interval is too small to be measureable, but the simple experiment of Haller affords unexceptionable evidence of the fact, that we employ our eyes severally, and not at the same instant, in distinct vision.\*

But we know that if the direction of the two eyes is in conformity, each with the other, objects appear as they are, singly; and that when a certain divergence or derangement of the visual axes exists, objects appear double. If the image, for example, is thrown upon a point of the retina of one eye, not in correspondence with the spot impressed in the other, this effect is produced. The double image of a candle is seen when gentle pressure is made on the globe of one eye; and it is either on the horizontal or vertical direction, according as the finger is applied to the side of the cornea, or below it. It is observed that a more considerable inclination of the optical axis is required to produce a double image in the transverse than in the vertical plane.

Double vision.

We are not, however, to conclude, that a double image is formed only, when the obliquity

Double vision with one eye.

\*Elem. Phys. Vol. V. Sec. IV. 9.



of the optic axis is such, as to throw the image beyond the area of the points of correspondence in the retina. It may happen when but one eye is employed, from a partial compression of the retina, optic nerve, or cerebrum, or some peculiarity of figure, or opaque streaks in the humors intercepting the radiant pencils, so as to produce a double refraction.

Double vision from strabismus.

Neither is double vision a common result of strabismus where the distortion of one eye is obvious and permanent; for in squinting, whether congenital or acquired, the distorted eye is weak in comparison with its fellow, and in the majority of cases the loss of association is the consequence of its weakness. It is, in fact, wholly unemployed in intent vision, which it would only tend to confuse. Where double vision occurs, it is seldom, if ever, a permanent symptom, although the squint becomes confirmed, or even increases. The disappearance of this symptom might be accounted for by the very probable accommodation of the deranged eyes, and the substitution of new points of correspondence in the distorted eye under the influence of habit; but in every instance of deviation, I believe it will be found that the averted eye is unfit, in respect of power, and therefore ceases to associate with its fellow. Even in the cast or leer the affected eye is unemployed in vision. The focal distance of the two eyes is in such cases so much at variance, that confu-



sion would necessarily result from their simultaneous employment, if that were possible. This question is not affected by the arguments which go to prove, that for the purpose of intent vision, one eye only is or can be employed at the same time. The points of correspondence are essential to preserve the unity of vision, because an indistinct or confused perception, or a double image, would be produced in the state of indolent vision, when it is admitted that both are employed. The defective eye, it is true, extends the field of sight; but if the sound eye be closed, the person discovers that he is indebted to it for little more; and therefore, if it correspond in direction with its fellow, he finds an advantage in closing it for the purpose of accurate vision; if it is permanently averted, it is as much unemployed, as if it were closed. Cases have occurred in which the exclusive employment of the averted eye has at length restored its tone and direction. The squinting eye recovers its position when the sound eye is closed, but relapses when the latter is again opened, because its employment ceases. Cases of squint arising from mechanical causes are of course excepted in this observation, being incapable of even temporary rectification; but in these the turned eye is idle.

It must be evident to all persons who consider the subject, that the rays of light which issue

Adjustment  
of the eye to  
objects at



different  
distances.

from an object at some distance from the eye, and those issuing from a much nearer object, cannot be collected into foci at the same given distance behind the crystalline lens, unless the eye have a power of altering its focal distance. It must do for itself what a convex glass does for those, who by reason of a certain configuration cannot see near objects distinctly, or a concave glass for those who have no distinct sight of objects beyond a moderate distance. In the first, owing to a defective refracting power, the rays cannot be brought to a focus soon enough; in the second, owing to too great a refractive power, they are brought into a focus too soon. The picture in the one, without the aid of the glass, would be formed behind the retina, and in the other anterior to it. The point of perfect indolent vision, or the extreme focal distance of the eye, and the range or space through which it has the power of preserving distinct vision nearer to the eye, varies in different individuals, and very often, as before stated, in the eyes of the same individual.

Various  
hypotheses.

I shall content myself with a very brief mention of the principal hypotheses to explain the adjusting mechanism. To enumerate all with barely intelligible conciseness, would occupy a large portion of this volume; such is the interest which this subject has excited. It has been ascribed to a change of figure in the cornea, to



the variations in the diameter of the pupil, to a change of figure of the globe by the action of its muscles, to a change of figure of the lens by an action proper to itself, to a change of place of the lens by the contraction of the ciliary processes, and the compression of the vitreous humor at its circumference.

The first supposes a close aponeurotic expansion derived from the tendons of the recti muscles, bracing the anterior segment of the globe; the second assumes the muscularity of the iris, or the extension of its texture, by the sudden injection of its vessels, and vice versâ, its abridgment by their contraction; the third, a power in the muscles of the globe either to shorten or elongate its axis; the fourth attributes muscularity to the crystalline; and the fifth a similar structure to the ciliary processes.

I shall not enter into a discussion of the merits of these hypotheses, because no one, I believe, disputes the force of the objections to which they are more or less exposed.

A healthy state of the retina, of the crystalline lens, and of the iris and ciliary apparatus, are conditions indispensable to the perfection of this mechanism. It is impaired in proportion to the debility of the retina in the various forms of amaurosis; it is suspended during the per-

Conditions  
requisite.



manent contraction or dilatation of the pupil, and it is lost after the removal of the crystalline under the most favourable circumstances; but the failure of any one of these conditions, exclusively, is destructive to it; as for example, though the retina and crystalline be healthy, if the iris be motionless, or though the retina be sound and the iris active, if the crystalline be absorbed. I know that very different statements have been given to the public, so different indeed as to be almost the converse of these. I am ready to admit that the results are subject to modifications, as the cases vary, and no two are exactly similar; but these are the general results of my experience. I have already said that the iris, by regulating the quantity of light, assists in the office of adjustment to distance, and that these functions are in a degree consentaneous. I am disposed to consider adjustment as the result of a change of figure in the lens, such as we may coarsely imitate by gentle pressure of the crystalline of the horse or ox, held in a vertical position between the thumb and fore finger. Its form and lamellated texture render it peculiarly susceptible of such a change, and the absence of a connecting medium between its plates, indeed of any vascular organization, prevents the possibility of a nebulous obscurity resulting from pressure so applied. A very slight increase of its curvature, we have been informed, on competent authority, would



be sufficient to explain the phenomenon of adjustment, assuming its quiescent state, which its elasticity tends incessantly to restore while subjected to compression, as that fitted for perfect indolent vision.

Notwithstanding the absence of anatomical proof, I cannot but regard the motions of the iris as muscular motions, and the pupillary portion an orbicular sphincter, such as environs the several outlets or apertures of the body. To this structure I attribute its uniformity under varying magnitudes—its incapacity of contraction, when having a fixed point, as happens in some malformations; when confined by adhesion at any point of the circle to the capsule of the lens, or when its texture has been the subject of adhesive inflammation—its recovery of a prolapse through a section of the cornea, and resuming its circular figure when overstretched, as in extraction, by a gentle friction of the eyelid—the extreme velocity of its contraction, and the comparative slowness of its relaxation—its ordinary preservation of a mean or middle state, between the spasmodic contraction induced by acute inflammation, and the dilatation we must from ascertained phenomena presume to be induced, by absolute darkness long continued—its inferior power of contraction in children, and the increase of its power by exercise, as in artisans incessantly employed upon minute objects, in whom it is apt

Iris in part  
muscular,



to acquire a rigidity which scarcely admits of dilatation—its obedience, in all respects, to the laws which regulate the muscular system—its contractility in proportion to the strength and perfection of the nerve of sense with which it is associated—its incapacity of perfect contraction when tremulous, and its spastic contraction, even to the resistance of the influence of belladonna, in tetanus—its relaxation when the sphincters are relaxed, as in syncope, asphyxia, apoplexy, or compression of the brain, and after the use of alcohol in excess—its complete dilatation when under the influence of the sedative poisons, as opium, hyoscyamus, belladonna, &c. to which its proper nerves are in a peculiar manner irritable.

and partly  
elastic.

The ciliary portion of the iris I regard as an elastic structure. It is by virtue of its elasticity that the extraordinary dilatation of the pupil, such as we see under the use of belladonna, is produced. Here, as in other parts, elasticity is opposed to muscular motion; hence when the latter is paralysed or from any cause diminished, the former strikingly predominates; when the nervous supply is intercepted, the pupil gapes widely, the action of elasticity being independent of the sensorium.

Pupils of  
animals.

All animals which have a moveable iris, have the pupil circular, oblong or elliptical,



forms favourable to the arrangement of marginal fibres.\* In fish the iris is evidently a prolongation of the choroid without interruption of continuity; it is therefore motionless. I concentrated the sun's rays in the focus of a pocket lens, and threw them upon the pupil of a perch, at the moment of drawing it from the water; it underwent no change. In other animals it contracted to a line, vertical (cat) or horizontal (adder, toad,) according to the figure of the pupil; or to a small pin's head aperture, where it was of a circular form, as in the common snake.

If we look through the vitreous humor exposed for a small space on its posterior aspect, we observe the plicæ advancing upon its anterior surface, beyond the margin of the lens, like a circular fan or screen; if the lens is pressed evenly backward, the plicæ separate and extend the sacculated circle of Petit, to which their edges are affixed. On remission of the pressure the lens springs forward, and the leaves of the fan are closed. The circumferential compression of the globe increases the closeness of their application. In the dead body, only the most coarse and remote analogies can be obtained to the functions of the living. But I cannot believe so obvious and yet so exquisite

Theory of  
the author.

\* It is worthy of notice, that fish in which the iris is without motion, furnish the remarkable exceptions; viz. the dolphin, the skate, the cuttle, &c.



a contrivance for changing the site and figure of the apparatus, as this view affords, can be without necessity or occasion. Looking then at the posterior origin of the processes from the choroid, and their attachment externally to the ciliary ring; their insertion into the vitreous capsule to the edge of the fossula, their encroachment upon the anterior segment of the crystalline, and their termination by distinct prolongations in the substance of the iris at its great circumference; assuming the choroid and annulus as fixed points, and the iris and processes as the moveable parts of the apparatus, it follows that the plicæ will be unbraced and partially open in the state of mean dilatation of the pupil, belonging to passive or atonic vision, and in the state of extreme dilatation of the pupil accompanied with blindness to near objects, totally relaxed and floating. On the contrary, by the steadily contracted state of the pupil suited to the nearest extremity of the focal range, they will be closed and braced together; and, bearing upon the circumference of the crystalline at every point, will necessarily elongate the axis of the lens. These being the extreme states, so, in proportion, the intermediate degrees of adaptation will be accounted for. Hence the actions of the pupil, however excited, will extend their influence to the lens, and by this catenation of motions the general conformity of adjustment to light and adaptation

to distance are to be explained. And this forms no objection to the hypothesis ; because it is only in the voluntary and steadily preserved contraction of the pupil that the latter object is or can be required ; for blindness would as surely ensue from gazing on the sun, as death from suspending the actions of the respiratory muscles, were it in our power to do either ; and therefore the involuntary has the ascendancy over the voluntary action in both these cases, as it has in all cases of mixed muscles.

Radiated fibres are described by Zinn and Haller as raised on the posterior face of the iris, and advancing even to the margin of the pupil. They are distinct from those seen upon its anterior surface, and regarded as continuations of the ciliary processes. In man no such fibres are distinguishable by the naked eye ; but if the observation, however obtained, be correct, it affords a strong presumption in favour of the power of the iris to change the figure of the lens by the instrumentality of the plicæ. The capsule, it is true, is fixed by the processes, but this opposes no impediment to the change supposed ; for the membrane of Petit, to which alone the processes are affixed, is relaxed when they are closed, and extended when they are separated, and thus permits the capsule to yield only in the degree required for the change of figure of the lens ; or, in other words, pre-

Uveal prolongations of the ciliary processes.



serves its exact adaptation to the face of the lens in its opposite and varying states. This I take to be the use of the membranous circle of Petit, that it gives the processes the complete command of the continuous capsule.

Some cases of dilated pupil are accompanied by a bulging of the lens. This is not the effect but the cause of the dilatation, for it never follows the application of belladonna, provided the capsule be entire; but if from any cause the lens be protruded so as to bear down the natural resistance of the processes, the pupil becomes dilated by its pressure.

Faculty of  
adaptation,  
how far en-  
joyed by  
animals.

It would require a more intimate acquaintance than we possess with the economy of the various classes of animals, to determine the several degrees in which they enjoy or require the power of adaptation to distances; but although many unquestionably have a great reach of sight, it is highly improbable that any animal approaches to man in minuteness or accuracy of near vision. The curvatures of the cornea and lens, which are inverse to each other, and the corresponding variations in the quantity of the aqueous humor, are obviously appropriated to the different densities of the media in which they habitually dwell.

The crystalline of man, compared with ani-

mals, is of the softest consistence, and occupies the smallest portion of the volume of the eyeball. The firmness of the crystalline is always in proportion to its convexity.

The absence of the processes in fish; their very slight indication (being close and delicate striæ instead of folds) in birds, and such reptiles as possess them; the absence in all of the membrane and circle of Petit, and the insertion of the processes into the capsule of the lens in the latter classes, offer a marked contrast to the appearances observed in man and quadrupeds, in whom they are full and strongly marked, and especially at the salient angle opposite to the crystalline, where they are unadhering and free to move. In fish, as we have said, the pupil is without motion; in birds and reptiles, as in man and quadrupeds, its motions are vivacious; in some we are told, voluntary, which, whether it be so or not, must be stated on pure conjecture. The adjusting power in fish and birds has been attributed to other mechanism, on account of certain peculiarities of structure, which seemed, *primâ facie*, to be adapted to that purpose, as the choroid gland in fish, and the pecten in birds; or of such deviations from the human structure as rendered the several hypotheses inapplicable.

I believe that the motions of the iris in animals are single, and obedient solely to the sti-



musculus of light, and that they have no control over the pupil by volition; a property which pertains exclusively to the adjusting power, and which is exerted independently of the variations of light. It is probable that they possess it in so far only as it results from the adaptation to light.

The iris is a mixed muscle; its motions are regulated in part by the stimulus of light upon the retina, and in part by an effort of the will.

Involuntary,

That the motions of the iris which take place upon the sudden changes of light are involuntary, there can be no doubt, for they are observed even in sleep, when the will cannot be exerted, and in the earliest infancy. There is another proof that these motions are involuntary, viz. that they occur in some forms of perfect amaurosis. I have seen the pupil act briskly, where the person has been totally devoid of the perception of light from bright sunshine, or the flame of a candle held before the eye.

The sympathy of the iris with the retina must be ascribable to a communication between the retina and the ciliary nerves which supply the iris. The small lenticular ganglion from which these nerves are derived, lies upon the optic nerve, and is probably the medium of communication.

On the other hand, every one may satisfy himself of a power which the will is capable of exercising over the iris, in viewing alternately near and distant objects; the state of relaxation or moderately dilated pupil being suited to the remote, and its tonic or relatively contracted state to the near object. It is seldom that this change is sufficient to be obvious to a bystander where the light remains unchanged, because the faculty is seldom exercised in these circumstances; and still more rare for the state of accommodation to be preserved in defiance of the changes of light, because it is an unnatural effort. I have several times observed, in persons whose eyes were steadily fixed upon an object at some yards' distance, that the approach of a candle towards the eye did not stimulate the pupil to contraction, until it was so placed as that its image should fall upon the most sensible part of the retina, when the pupil instantly contracted. So that the voluntary is in subordination to the involuntary power, where they are opposed; that is, when the stimulus of light opposes the adaptation of the eye to distance. But by continued application, the mind is capable of acquiring over the motions of the iris an extraordinary power, as is well known to be the case with other muscles subjected in any degree to volition. Of this I have seen two or three remarkable instances, but none so striking as that of my ingenious and learned friend Dr. P. M.

and voluntary motion  
of the iris.



Roget, in whom, I may be permitted to say, profound scientific knowledge is accompanied by a characteristic aversion to ostentatious display. It affords me much pleasure to lay before my reader the peculiarity to which I have alluded in the person of Dr. R. as described by himself, at my request.

*" Bernard Street, Russel Square,  
Feb. 21, 1820.*

" MY DEAR SIR,

" I am much pleased with the view you design to take, in your intended work, of the subserviency of the motions of the iris to the changes which accompany distinct vision at different distances, and of these motions being subordinate to the effect of light on the retina; and the more so as they accord with a circumstance relative to my own eyes which I have often made the subject of experiment, and which you will probably recollect my showing to you some years ago.

" When I have stated that I possessed the power of dilating and contracting at pleasure the iris, the fibres of which are usually considered as no more under the dominion of the will than the heart or blood-vessels, my assertion has, in general, excited much astonishment. Such, however, is strictly the fact. I can easily satisfy any person who witnesses the movements

I can produce in them, that this power is totally independent of the influence of light; since I can effectually exert it, although the position of my eye with regard to the window or candle, as well as the direction of the optic axis, continue unchanged. However singular this power may appear, it admits, I conceive, of a very natural explanation. The effort of which I am conscious, when performing the voluntary contraction of the pupil, is the same as that which accompanies the adaptation of the eye to the vision of near objects, and is of course productive of an increase of its refractive power. This very same power of moving the iris is in fact possessed, in a greater or less degree, by every person who enjoys the faculty of distinct vision at different distances. It is accordingly well known, that if a person after looking at a distant object, transfers his attention to a near object, the pupil always contracts. But this change, it is supposed, can never be effected, unless some real object or image, from which light radiates, be present to direct the sight. I have never, indeed, met with any person besides myself, who, while steadily directing his eye to a distant object, and while no other object intervened, could, by a mere effort of volition exerted on the eye, augment its refractive power so as to adapt it to the vision of near objects. That I have acquired such a power I can ascribe to no other cause, than to my hav-



ing from my childhood, been much in the habit of observing optical phenomena, and of practising various experiments relating to vision, a subject which I early took great delight in cultivating.

“It is still more easy for me, while an object is placed near my eye and distinctly seen, immediately to relax the organ so as to fit it for the distinct vision of the most distant objects; and these changes I can effect in succession, with considerable rapidity, each change being accompanied with a corresponding enlargement or diminution of the pupil. The increasing the refractive power of the eye, is always the change that constitutes the effort; the state of vision adapted to parallel rays being that of complete relaxation. The effort which attends this voluntary contraction of the pupil, when there is no object before the eye to call for such a change, is followed by a sense of fatigue; and if often repeated or too long continued, it becomes painful, and continues so for some time afterwards. The fatigue is felt almost exclusively in the eye to which my attention had been directed during the experiment, although the same change in the refractive power takes place, and I believe to the same extent, in the other eye. It is also remarkable, that when there exists a real object of sight which is looked at, and which requires an equal change

in the eye for distinct vision, as in the former case, no sense of fatigue, or hardly any, is experienced.

“ I need scarcely add, that while I thus alter the refractive power of my eye from that which adapts it to the distance of the objects I look at, those objects appear indistinct, from their images either forming before the rays reach the retina, or tending to form beyond it.

“ I am,

“ DEAR SIR,

“ very faithfully yours,

“ P. M. ROGET.”

“ B. TRAVERS, ESQ.”

---

In addition to the nerves derived from the lenticular ganglion, the iris receives two or more branches from the nasal nerve, (5th pair), and its actions may possibly be subjected to the will by virtue of the influence which these nerves convey, for from the same source is derived the nerve which supplies the levator palpebræ, which is purely a voluntary muscle.

Ciliary  
nerves from  
the nasal:



The limited motion which the pupil has when the retina is for the most part insensible, may be considered as an involuntary or automatic motion, similar to that which in a healthy eye affords protection to the retina ; and if, as sometimes happens, the iris contracts in a state of blindness, this likewise must be regarded as its involuntary action, for volition cannot precede sensation. It is probable that those motions of the iris, which are in conformity to the impressions of light upon the retina, are purely involuntary ; and that those which are in conformity to the situation of objects, and are therefore directly subservient to vision, are under the influence of the will. Hence the dependence of the adjusting faculty upon the perfection of the retina.

Lenticular  
ganglion.

The ganglia have been conjectured by an ingenious author to be bars or stops upon volition, and this case of the iris, which he assumed to be purely involuntary, was incorrectly cited in support of the hypothesis. By others ganglia have been supposed to be small sensories or cerebral receptacles, capable of rendering a supply of nervous energy to their filaments, by which they are in a measure independent of the brain and its appendages. The theory, which I have ventured to suggest, attributes the voluntary motions of the iris to nerves unconnected with ganglia, the involuntary to those

derived from the lenticular ganglion, which I regard as a direct medium of communication between these nerves and the retina.

The phenomena of ocular spectra, or images of luminous objects remaining upon the retina after the external impression is withdrawn, are highly interesting and curious. Luminous sparks and flashes, halos or variously colored rings, it is well known, are produced at will by friction or pressure of the closed eyelids, and the first are an instant effect of concussions of the brain. The red is that color called up by the rudest artificial pressure; the violet by the slightest; and the gentlest impulse is the natural one, in which the light suffers no decomposition. Are these appearances really retinal impressions, or illusory mental phantasms, founded on the feeble and obscure analogy subsisting between mechanical pressure and the impression of light? Although blind persons see such appearances, I doubt if they ever present themselves in cases where the retina is disorganised, or after the extirpation of the eyeball, as the mutilated feel their fingers and toes. They seem to me therefore to establish the essential connexion between the retina and the faculty of perception, or the connexion between the corporeal and mental impressions; and this is confirmed by what we observe of morbid spectra, which are

Ocular  
spectra.



symptoms of various disordered states of the retina, of which I shall speak hereafter.

Direct.

I have said that the duration of an impression is in proportion to its intensity. The experiment of the revolving fire-stick demonstrates that the impressions upon single points of the retina, although successive, become blended or confused by vividness and consequent proportionate duration; for if it were not luminous, the appearance of a continuous circle would scarcely be produced. Spectra are direct or reverse. The first is the impression of a luminous object, the shadow of that upon which the eye has for some time dwelt, although with no peculiar degree of intentness, and presenting the outline of the object in color. This either vanishes at once, or it presents a circle of the primary colors, variously associated or successively exhibited, in the order of their relation. The latter is especially the case after looking at the sun or a very bright light. This is the simple effect of a temporary over-excitement of the retina, analagous to the echo of a noise in our ears, by which the auditory nerve has been over-excited. The reverse

Reverse.

spectrum is produced, when a color, occupying a certain space, has been so intensely impressed (as when for example we make the experiment), as to exhaust the irritability of the retina, and

render it inexcitable by any and every combination of the rays of light, in which that color is a constituent. Hence the color of the spectrum is that which results from the abstraction of the offensive ray from white light, or the reverse of that of the object. The stimulus most remote from that which has excited the distress, is the only one to which the retina is alive, and in this it finds the relief of contrast.

In another case the reverse spectrum seems to depend on the excess of susceptibility in the retina, as when the eye has been fixed on a black ground; here the spectrum is white. Hence it is that the ash grey wall, presented to the eye on arriving at the mouth of a cavern, has a silvery brightness. The contrast of white light is alone capable of exciting the retina which has been altogether deprived of its natural stimulus.

Thus we have two states or degrees of nervous excitement; the first, simple over-excitement to the extent of continued or renovated impression. The second, excitement to exhaustion, leaving only a negative sensibility. A third state is that of non-excitement, and consequently accumulated excitability. The two last are directly opposed, being minus and plus states of sensibility, and thus the contem-

Explanation  
of these  
phenomena,



plation of a white surface gives a black spectrum, as the black gives a white one.

As they are presented to the healthy eye, spectra need no further explanation. They are to be referred to the preternatural duration or intensity, or to the privation of the natural stimulus. They are accordingly produced artificially and at will in all persons. The organ is passive. The hypothesis of spasmodic action in the retina is altogether gratuitous.

---

In the "ANATOMICAL DESCRIPTION," I have unavoidably embodied the chief of what it is needful to say on the economy of the appendages. A few additional circumstances, however, come properly to be noticed under this head.

The actions  
of the palpe-  
bral muscles.

The levator palpebræ being purely a voluntary muscle, the simple suspension of its action effects the closure of the lids, as its contraction opens them in the act of waking. Hence the disposition in the upper lid to fall announces the approach of sleep. In febrile and exhausted states of the system, its impaired energy occasions the

drooping expressed by the term 'heavy-eyed', one of the most characteristic symptoms in the physiognomy of disease. A similar state belongs to some morbid affections of vision, of which I shall have occasion to speak hereafter. A voluntary closure of the eyelids, as when the eye is from any cause irritable to light, is performed by the orbicularis palpebrarum, which in some casualties and morbid states contracts spasmodically, and the relaxation of this muscle assists the opening of the closed lids. In going to sleep and awaking from it, the lower lid is therefore passive; in a voluntary shutting and opening of the eye it participates, although inconsiderably, in both actions. Winking is an alternation of the actions of the levator and orbicularis, and therefore a seasonable relief to the former, and a means of preserving the moist and clear condition of the cornea. It is performed by a very slight contraction of the palpebral portion of the orbicularis. The combination of the action of the corrugator and orbicularis is seen in the strained closure of the lids to resist their separation by external force, knitting and depressing the eyebrows, and throwing the nose and forehead into folds; and the equipoise of the actions of the orbicularis and levator is evinced in the approximation or screwing of the eyelids, and peering, as is customary in short sighted persons. When they are both in full action, the



corrugator acts as a moderator to the levator; the orbicularis is the antagonist of the latter.

Actions of  
the muscles  
of the globe.

The actions of the straight muscles in various combinations, and in succession, explain the several intermediate motions to those which they singly perform, and the revolving motions of the ball round its socket. The rotation on its axis inward is performed by the superior oblique, that in the opposite direction by the inferior. The combined actions of the whole preserve the relative position of the eye to the object, independent equally of the motions of the object and the head. The motions of the eyes are in perfect correspondence, and the will cannot place them in opposition.

Superciliary  
arch and  
muscles.

The prominence of the superciliary ridge, as well as the fulness of the brow, is subject to great variety, and is sometimes an impediment to the facility of operations. On the physiognomy I need not say its influence is most marked. The elevation of the eyebrow performed by the occipito-frontalis co-operating with the levator, in staring, and its depression and approximation to the nose by the orbicularis and corrugator in frowning, are habitually employed to a manifest advantage in the opposite states of deficient or feeble, and of superabundant or dazzling light. The eye's 'mute eloquence which passes speech,'

belongs chiefly to its appendages. An opera dancer would be as effective with a divided tendo achillis, as a tragic actor who had lost the moveable apparel of the orbits.

The closed eyelids are penetrated by a full light, so as in ordinary circumstances to occasion waking, and distress to persons whose eyes are inflamed. The superior tarsus, when drawn up, slides under the arch of the orbit, but retains its apposition to the globe, owing to the laxity of its attachment with the integument of the palpebra.

Tarsi permeable to light.

The origin of some fibres of the orbicularis from the ligamentous expansion which supports and protects the lacrymal sac, gives it a power of compressing the sac in its contraction, and thus assists in the excretion of the tears. This is in part proved by the epiphora which accompanies a fixed state of the lower palpebra from injuries, and the paralysis of the orbicularis, which states also prevent the due apposition of the puncta. Hence too, people wink often and forcibly, whose eyes are disposed to water, and after shedding tears.

Muscular compression of the lacrymal sac.

The conjunctiva has been supposed to secrete a lubricating fluid, which serves the purpose of keeping the cornea clear, and facilitating the motions of the lids. This forms, according to

Conjunctiva a non-secreting surface.



that opinion, the habitual lacrymal discharge, and the secretion of the lacrymal gland is but an occasional one, as when from mental emotion or irritation of the conjunctiva the profuse and palpable discharge which we call 'tears' is poured out. I believe the notion rests solely upon the supposed relation of the conjunctiva to the class of mucous membranes. But this need be no fetter upon our conception of the matter; for not only do we see from the varieties of its surface that its economy is not throughout the same, but anatomists describe its continuity with the cutis as much as with the membrana narium. Again, the capillaries of mucous membranes carry red blood, which is not the case in the conjunctiva of the globe in health. But there is no evidence of such a secretion; upon the cornea it is not assumed to exist, yet the difference between the corneal and sclerotic portion is only in the strictness of its adhesion. The follicles and caruncula are specifically provided for preventing the effects of friction, and the incessant although insensible escape of the tears from the lacrymal ducts, unavoidable under the act of winking, in which the upper lid sweeps over and preserves the polish of the cornea, renders such a provision superfluous and therefore improbable. In disease, the sclerotic conjunctiva secretes a mucus which is immediately obvious (the corneal surface is excepted because its vessels do not admit red blood) and this is in conformity

with what we see of the mucous membrane properly so called, as of the urethra and intestinal canal, which continually show that the secretion can be set up by disease upon a whole surface, while in the healthy state this function is confined to its follicles and lacunæ.

A young woman who had never shed tears, and was incapable of doing so, had a shrivelled, opaque and cuticular conjunctiva.

The puncta lacrymalia absorb the tears, not by any capillary attraction, but by a vital action as absorbent mouths. They are often spasmodically contracted, and afford a resistance to the introduction of Anel's probe, but yield to the point of a pin, so as afterwards readily to admit a probe of much larger dimensions. When over-dilated, they lose for a time their absorbing power, and the epiphora is increased. When they are morbidly patulous and atonic, as sometimes happens in age, the epiphora is permanent; and their function is frequently arrested by inflammation of the sac, for we often find the epiphora altogether independent of obstruction.

Excretion  
of tears.

The direction of the superior duct varies a little in relation to the sac, according to the degree of elevation of the upper lid. By drawing



the lid upwards and towards the nose, it is brought nearly into a line with the axis of the sac.

The area of the sac and nasal portion of the duct, exceeding that of their orifices, facilitates the passage of the tears; the slight elevations of the lining membrane and the narrowness and obliquity of the nasal opening probably retard the excretion, which would be inconvenient if constantly taking place.

---

## PART II.

---

### PATHOLOGY

OF

## THE MEMBRANES.

---

---

### CHAPTER I.

---

#### SECTION I.

#### CONJUNCTIVA.

THE vessels of the conjunctiva of the globe are derived from two sources:—1. The palpebral arteries and veins; 2. The ophthalmic muscular branches, and accompanying veins. The first, creeping upon the sclerotic conjunctiva in the sinus palpebrales, and at the angles of the palpebræ, have a reticular distribution upon the loose portion of the conjunctiva. The second, after penetrating the tendons of the straight muscles, advance in four distinct fasciculi. These

Blood vessels of the conjunctiva.



spreading, as they advance upon the opposite sides of the globe, form numerous lateral anastomoses with each other, so as to present a faint circulus arteriosus upon the sclerotic coat. In a state of congestion a free communication is conspicuous between these two orders of vessels, and their distinctive character is preserved under very high degrees of inflammation. In a complete section of the conjunctiva at a short distance from the cornea, both sets may be divided without injury to the sclerotic. It is by infinitely frequent and minute subdivisions of these vessels that the tissue which unites the conjunctiva to the sclerotic is supplied; and hence in inflammation, the areolar distribution of these vessels is most conspicuously demonstrated where this texture is most abundant. At the verge of the cornea both orders of vessels are rectilinear, and never present the reticular arrangement; for the looseness of the conjunctiva upon the sclerotic gradually diminishes from the point of its reflection to the verge of the cornea, and its adhesion to the cornea is so strict, as to render an artificial separation impracticable. This remarkable difference in strictness of adhesion subsisting between the conjunctiva and subjacent textures, suited to their difference of economy, explains some varieties in the morbid affections of the conjunctiva covering these parts respectively.

Independent of the accompanying veins of the muscular arterial fasciculi, long and tortuous branches are seen to proceed from the meshes of the conjunctiva, distinguished from the arteries by their purple hue, their isolated course, and more superficial seat. Their sharp spiral curves are larger as they descend towards the base of the visible hemisphere of the globe, where their attachment to the sclerotic is least strict. These are the palpebral veins of the conjunctiva.

Although in the tonic and tranquil state of the eye, little, if any, red blood is admitted into the superficial vessels, yet under a very temporary excitement, colored blood has a ready admission into the vessels of the sclerotic conjunctiva. Such, however, is the condensation of the connecting texture upon the cornea, as to prevent the admission of red blood into its vessels under a very high degree of inflammation, as even where the white ground of the sclerotic is extinguished\*. The susceptibility of parts permeable to red blood to increased vascularity under excitement, is in proportion to the quantity of cellular texture entering into their composition, or connecting them with subjacent parts. Compare, in this view, the membrane of the fauces and the trachea

Characteristic difference of the conjunctiva of the sclerotic and cornea, under inflammation.

\* I believe further, that it is inadmissible, except a morbid change has previously taken possession of this texture. But of this hereafter.



—pleura pulmonalis and costalis—periosteum and perichondrium—the lining membrane of veins and arteries.

Since the easy admission of red blood into the vessels of the sclerotic conjunctiva discovers a distribution of vessels, not otherwise known, it affords opportunities of observing the accession and progress of inflammation; and this portion of the membrane presents, when inflamed, appearances different from those of the inflamed corneal conjunctiva. But although the colorless circulation of the latter shows the rise and progress of inflammation obscurely, its perfect transparency in health makes the results of it more conspicuous.

Cornea obscured by inflammation.

The first effect of inflammation upon the cornea is haze or dimness, which depends upon the loaded state of its (serous) vessels. The dimness is immediately removed by the recovery of the circulation, as after the removal of an irritant; for example, a foreign particle on the cornea; or after a free evacuation, where the cause is less obvious. Hence transient dimness is merely a condition of congestion. This state, however, continued, produces a deeper and more permanent opacity; viz. effusion into the connecting texture, and thickening of the conjunctiva upon the cornea. This is the progress of simple ophthalmia in the generic sense of the term. Contrast the epidermis in the state of blush (conges-

tion) and of incipient vesication (effusion). The peritoneal coat of the liver, the arachnoid and synovial membranes doubtless exhibit, in the distinct stages of congestion and effusion, the transient and permanent opacity.

The conjunctiva is to the cornea, what the periosteum is to the bone. It nourishes the superficial lamellæ; wherever it is completely detached, the exposed surface of the cornea ulcerates, and its vessels repair the breach. To pursue the analogy, the interlamellar texture of the cornea may represent the medullary membrane; gangrene therefore does not ensue but from a permanent destruction of both textures, as by blows and explosions, which mechanically disorganize; by the action of lime, gunpowder, strong acids, and other chemically destructive agents; or by the strangulation of the vessels of both textures, as in the excessive chemosis, which destroys on the same principle as the paraphymosis, or the strangulated hernia.

Relation of  
the conjunc-  
tiva to the  
cornea.

The forms of inflammation of the conjunctiva which I shall now proceed to refer to, I consider as specific variations from the simple acute inflammation,\* of which the ordinary signs are familiar to every practitioner, originating in a healthy subject from an obvious occasional or accidental cause, as an extraneous particle, or a blow not injuring the texture, or a blast of cold air. An

\* See Plate I. fig. 1.



inflammation purely local, uninfluenced by constitution, both from the nature of its origin, its recent existence, and the health of the subject in whom it occurs. Such is the simplest example we can suggest of the morbid disposition natural to this organ, or to any organ; it is frequent, for all are liable to it, if all are not equally exposed: it is in fact an instance of what may be termed, without a misnomer, the inflammation of health.

Inflam-  
mation of the  
conjunctiva,  
modified by  
scrofula.

The inflammation of the conjunctiva, termed 'strumous,' where it has not proceeded to a change of texture, is not marked by any prominent local character. The vascularity is inconsiderable. This inflammation sometimes accompanies pustule of the sclerotic conjunctiva, in which case the vascularity is diffused instead of being partial, as in pure pustular inflammation, and the intolerance of light characteristic of the strumous inflammation is present in a greater or less degree. It accompanies also the morbid secretion of the lids when the eyeball becomes affected by the acuteness and duration of that disease, and the pustule on the cornea, especially the variolous pustule. In its simplest form it is almost peculiar to young children, stationary, marked by a very slight redness of the sclerotic conjunctiva, and the greatest possible degree of intolerance. I have known it in more than one instance of such severity and duration as to occasion a distortion of the spine from

the habitual depression of the head, and the obstinate maintenance of an awkward and unnatural posture of the body to screen the eyes from light.

The disproportionate degree of inflammation makes it difficult to account for this excessive morbid sensibility. It is purely a disorder of function; for although it far exceeds that which accompanies the acutest inflammation to which the organ is liable, it never in my experience impairs the faculty of vision. I attribute it to a morbid sympathy of the retina with the secreting surfaces of the primæ viæ and the skin, for neither of these organs perform their healthy functions during its existence. The tongue, the index of the former, shows by various signs gastric irritation or disordered digestion, and the cutaneous surface is remarkably dry and harsh. Accordingly it is cured by diaphoretics, as tartar emetic to nausea, James's powder, or calomel combined with opium in small doses; by the warm bath; and materially corrected, if not removed, by a preternatural secretion in the vicinity, as by an open blister on the nape of the neck. I have often seen an aggravated intolerance removed in twelve hours by the application of a blister.

This state of intolerance bears an analogy to cases of depraved, or rather of painfully acute sensibility in other organs of sense. The senses



of hearing and of smell are in some rare cases rendered morbidly acute, independent of the slightest organic affection, so that the ordinarily agreeable stimuli of these organs in a state of health, viz. an acute sound and a pungent odor, become causes of distress. An amateur of music, when labouring under an occasional disorder of the auditory passages, compared his sensations, during a fine performance of instrumental music, to those of Hogarth's enraged musician; all was jar and discord. Every snuff-taker knows the effect of a catarrh to spoil his enjoyment.

The nebula and the pustule of the corneal conjunctiva are the terminations of this inflammation when it affects the texture of the organ, to which may be added the small herpetic ulcers, reddish brown points, giving to the cornea a scabrous appearance. The healing action is always remarkably languid and protracted, as if the state of excessive irritability checked its progress, and prevented its completion.

Apthous or  
pustular in-  
flammation  
of the con-  
junctiva.

The conjunctiva of the sclerotic is disposed to form apthæ or pustules at the verge of the cornea, or near to it. In the former situation, where the more lax adhesion becomes abruptly strict, the pustule is elevated or cone-like, and is the termination of a distinct pencil of vessels; which arrangement sometimes precedes and announces the disposition to pustule. When it is

situated at a distance of a line or two from the corneal margin, it is broad and flattened. It is a small speck or patch of lymph, and seldom advances to suppuration. It is common to see one on either side of the cornea, in the transverse axis of the globe.\* Sometimes they appear in detached clusters, or a zone of pustules environs the cornea. This resembles the aphtha of the mouth and fauces and intestinal canal.

The pustules of the corneal conjunctiva which are less frequent, except in children, are generally situated near to the margin of the cornea, where one or more pustules of the sclerotic portion appear. Like the aphtha of the glans penis and the stricter parts of fine cutaneous texture, the pustule on the cornea usually forms an ulcer.

The ophthalmia with puriform discharge is a disease of the palpebræ. The secretion is supplied by the meibomian follicles and the conjunctiva bordering them, and likewise by the caruncula lacrymalis.

Indamm-  
tion of the  
follicles with  
puriform dis-  
charge.

A puriform discharge is furnished by the cryptæ of the tonsils, the lacunæ of the urethra, and the mucous glands of the nares, fauces, rectum and vagina, in mild inflammation of these parts. But in the vehement acute form of inflammation,

\* See Plate I. fig. 6.



the matter of suppuration is furnished by the tumid and villous surfaces of these membranes.

Acute suppurative inflammation of the conjunctiva.

The sclerotic conjunctiva in acute suppurative ophthalmia presents the following states: 1st, Serous effusion (œdema) which is common to other inflammations, and especially those of a less vigorous kind. 2d, Effusion of lymph (chemosis) peculiar to this form of inflammation, by which it acquires a solid augmentation of bulk. 3d, Villosity, or a subsequent prolongation of the extreme vessels in the form of villi, which secrete pus. The strict adhesion of the conjunctiva to the cornea prevents these changes from taking place upon that membrane. Upon the tarsi the conjunctiva thus affected becomes preternaturally vascular, thickened, and scabrous, or forms fleshy eminences. That the vascular villi of the conjunctiva secrete pus, may be ascertained by the aid of a lens. The pus, when formed, collects in the interstices of the villous texture. We have no evidence, as I have before observed, that the conjunctiva is a secreting surface in the healthy state.

Mild acute suppurative inflammation of the conjunctiva.

A form of disease intermediate to these in extent and severity, a modification of the suppurative ophthalmia, is the villosity and puriform secretion of the conjunctiva palpebralis, as seen upon eversion of the lids, while the membrane upon the globe is simply intu-

mescent, giving it a more rounded figure, and moderately vascular. This is the mild acute suppurative ophthalmia, which seldom injures the cornea, but frequently leaves after it the same fungous or granulated state of the conjunctiva palpebralis which so often follows the most acute form.

The difference then between the inflammation of the meibomian follicles and caruncula (the disease which if neglected terminates in lippitudo), and the suppurative inflammation of the conjunctiva, whether of the palpebræ or the globe, is a difference in kind as well as in seat; the one is the conversion of a transparent and bland secretion proper to the organ (meibomian) into a viscid and irritating mucus, puriform in appearance; the other is the de novo production of a true suppurative surface by inflammation.

The highly contagious nature of the suppurative ophthalmia, whether in the mild or vehement acute form, is sufficiently proved. For one person affected with this disease above three months old, I should think at least twenty are subject to it under that age. The mother is the subject of fluor albus or gonorrhœa, and the discharge is usually perceived about the third day. In new-born infants the disease begins in the conjunctiva palpebralis, and is often confined to that portion of the membrane. Where by

Suppurative  
ophthalmia  
communi-  
cated by  
contact.



neglect or improper treatment, it extends to the conjunctiva of the globe, it often destroys the cornea. I have repeatedly seen the most virulent form of the disease produced by an accidental translation of the matter of gonorrhœa from its source in the same subject, and from one to another, as from the husband to his wife. I have also known it set up by the fluid injected into the eye of a patient, spirting into the eye of the medical attendant, and by the use of a sponge which had been recently employed to cleanse the eyes of an infant affected with the disease. The mild as well as the acute form of the disease, it is well known, runs through armies, schools, and families. There is much reason to believe it epidemic as well as contagious, but the former is a point less easy of decision.

• Chemosis.

The chemosis, as must appear by the description above given, is an affection widely differing from ecchymosis, with which it has been by some writers confounded.\* It is after the existence

\* In the forming stage of chemosis, it is frequently mistaken for a mere œdematous effusion of serum. The apparent softness of the swelling at this early period may readily lead the practitioner into this error. I have not unfrequently known the conjunctiva of the eye-ball freely scarified, when in this condition, with the expectation of discharging the fluid supposed to distend it; and, as a necessary consequence, the inflammation thereby materially increased.—*Editor.*

of this morbid condition which is characteristic of the suppurative ophthalmia, that the conjunctiva forms fungous excrescences, pendulous flaps, or hard callous rolls protruding between the palpebræ and globe, and everting the former (ectropeon), or if not protruding, causing the turning of the lid over against the globe (entropeon). The tarsal portion takes on from the same cause the hard granulated surface, which keeps up incessant irritation of the sclerotic conjunctiva, and at length renders the cornea opaque. These eminences, sometimes cone-like and sometimes flattened, are not granulations in reality, i. e. adventitious glands secreting pus. Granulations, I believe, are never formed without breach of texture.

Preternatural elongations and excrescences of the conjunctiva, concealed in the hollows of the palpebræ, are a sequel of the protracted mild suppurative ophthalmia, where the palpebral conjunctiva has been the principal seat of disease. They are similar to those of the membrane lining the rectum, and the fine skin at the verge of the anus after inflamed piles, and the pudendum muliebre in acute gonorrhœa. All such membranous growths are, I believe, referrible to irritation of inflamed parts by the diseased and confined secretion, as the warts in external gonorrhœa.

Elongations and excrescences.



Simple fungus of the conjunctiva.

A firm fleshy fungus, which sometimes attains considerable bulk, so as to project from between the eyelids and globe in an orbicular figure, even to the circumference of the orbit, I have extirpated. Such fungi are exclusively formed of the conjunctiva, and usually originate from injury.

Carcinomatous fungus of the conjunctiva.

There is a malignant fungus of the conjunctiva, for, like the mucous membrane of other parts, this is sometimes the seat of carcinoma; and excepting the lacrymal gland, I believe no other texture related to the organ of vision is ever primarily so affected. I have removed the contents of the orbit for a painful tubercular fungus, with ulcerated depressions containing an ichorous discharge. The coats and humors of the eye were for the most part absorbed, the lacrymal gland scirrhus. The disease afterwards returned upon the palpebræ, and destroyed the patient. I have at this time a similar case under my observation. The fleshy tubercles grow from the conjunctiva, both on the cornea and sclerotic, and the inferior palpebra is extensively ulcerated.\* It is accompanied by lancinating pains in the supra-orbital region, and an unhealthy discharge.

Pannus.

The pannus is a chronic thickening and opacity of the conjunctiva of the sclerotic, generally

\*See Plate II. fig. 1.

unaccompanied by inflammation. By relaxation of the connecting tissue the membrane becomes redundant in extent, and forms folds or duplicatures, on one, or on all sides of the cornea, which encroach upon it considerably in the motions of the globe. The elongated uvula is the only analogy that occurs to my mind. This is often unpreceded by inflammation, and the extension is purely membranous.

The membranous pterygium is a true nebula of the sclerotic conjunctiva; the fleshy is an adipose or sarcomatous growth beneath the sclerotic conjunctiva. It extends from either canthus or sinus palpebralis, most commonly from behind the caruncula lacrymalis; and by its increase forcibly detaches the conjunctiva from the cornea. In its progress it occasions a permanent and indelible opacity by the thickening of the conjunctiva, and the deposition of lymph in the interspace of these membranes, in the form of a little tongue-shaped process. The wedge-like figure of the fleshy pteryx, and its gradual extension upon the cornea, afford the best pathological demonstration of the continuity of the conjunctiva; and the spread fan-like figure of the membranous, its semi-transparency as well as its termination in simple nebula of the corneal conjunctiva, shows the difference in the nature of the two diseases. Both this and the disease last mentioned, like other morbid growths of the

Pterygium,  
membranous  
and fleshy.



cellular texture or beneath it, are most prevalent in warm climates.

Conjunctival tumors on the sclerotic,

Adipose, steatomatous, and even cartilaginous tumors form in the cellular tissue of the sclerotic conjunctiva, and produce the same change when situated in the vicinity of the cornea as the disease last mentioned, viz. a marginal interstitial deposition.

and cornea.

Circumscribed tumors of a dense and firm texture are sometimes formed upon the surface of the cornea, and attain a considerable magnitude; but such cases are rare. I excised the anterior hemisphere of the eyeball in an elderly lady, in whom the cornea was concealed by a tumor, of a dark purple color, protruding to such an extent between the eyelids, as to occasion great inconvenience and deformity. It had the appearance of being disposed in lobes, somewhat resembling a bunch of currants of unequal size. On dissection, the cornea and sclerotic proved to be entire, and the morbid growth, lying upon and adhering to the corneal and a small portion of the sclerotic surface, had acquired the lobulated appearance, as if by degeneration of the covering conjunctiva; for delicate white bands, the only vestiges of this membrane, were seen intersecting the lobules at irregular distances, in the form of septa. The substance, on section, was firm, of a dark color, here and there mottled with white, and measured

a quarter of an inch in thickness from the external surface of the cornea.\*

The encanthis is a morbid enlargement of the lacrymal caruncle, in the form of a granular tumor, involving the valvula semilunaris. Sometimes the short down growing upon this gland takes on a morbid growth and harshness. The disease is extremely irritating, and occasions epiphora by a forcible diversion of the lacrymal puncta from each other, and from the surface of the globe. I have never known it assume the malignant character ascribed to it by some writers.

Encanthis.

The elongated valvula semilunaris retains its crescentic figure even to the margin of the cornea, where it has a loose and thin edge. By this and other parts of the conjunctiva, fringes or clusters of soft red caruncles are sometimes produced, resembling those occasionally seen in the site of the carunculæ myrtiformes, and about the os externum vaginæ.

Elongated  
valvula se-  
milunaris.

The frena or frenula connecting the conjunctiva palpebrarum and conjunctiva scleroticæ, a troublesome, and often irremediable deformity, follows burns and wounds of the conjunctiva tarsi, and the excision of tumors connected with this portion of the membrane. They are mem-

Frena.

Plate II. fig. 2. and fig. 4.



branous bands formed by adhesive inflammation of the opposed and contiguous surface (pleura costalis and pulmonalis—peritoneum of the intestines and parietes.) It is not necessary that both surfaces should be wounded, if the position be by a mistake of treatment preserved, as by the application of a bandage. “The opposite uninflamed surface,” as Mr Hunter observes, “accepts of the union.” I have seen these frena produced by a slit eyelid from a fall, and trifling as the inconvenience might seem, it so restricted the motions of the globe, and the disease was so materially aggravated by operations to relieve it, i. e. by the multiplication of frenula, that the patient became disturbed in his intellects, from an exaggerated sense of his misfortune.

Co-adhering  
tarsi.

The co-adhesion or concretion of the tarsi by organized adhesion of the conjunctiva tarsorum is rare. I saw a remarkable case of it in a full grown boy, whose eye was found perfect after the division, though he had been thus blind from his infancy. It is similar to the co-adhesion of the nymphæ or labia pudendi, and the closed anus in new-born infants.

Indisposi-  
tion of the  
conjunctiva  
to ulcerative  
inflamma-  
tion.

The conjunctiva is not prone to ulcerate, whilst the substance of the cornea readily assumes that action; hence the frequency of ulcers not opening externally, and of ulcers penetrating into the anterior chamber. Its readiness to assume the adhesive inflammation is evinced

by the rapid formation of a superficial speck where it has been scratched or abraded, and the assistance it affords in healing open ulcers of the cornea. There is a marked disposition in these cicatrices to ulcerate in subsequent attacks of inflammation, which are in the same degree as in other parts slower to heal than the original texture. The synovial membrane is as much less disposed to ulceration than the cartilage, as the conjunctiva is than the cornea, or the periosteum than the bone, or the peritoneum than the mucous coat of the bowel. All these external close membranes accord in their disposition to adhesive inflammation.

The conjunctiva, viewing its compound pathological character, bears in its respective relations to the sclerotic and the cornea, an affinity to the two distinct classes of membranes: viz. the mucous upon the sclerotic, and the serous upon the cornea; hence the frequency of pustule and the tendency to suppurative inflammation of the sclerotic portion, and the indisposition to ulcerate and proneness to adhesive inflammation of the corneal.

Pathological  
relations of  
the conjunc-  
tiva.



## SECTION II.

### CORNEA.

Ulcers of  
the cornea,

THE cornea is disposed to adhesive inflammation, ulceration, and sloughing. It rarely suppurates. The ulcer of the cornea begins, not in abscess, but in a circumscribed deposit of lymph, or in pure ulcerative absorption without pus. In most instances, when of spontaneous origin, it begins in the interlamellar texture. When the conjunctiva has been detached, and the cornea deeply injured, as by a large spiculum, or by an ulcerated pustule of the corneal conjunctiva, the ulcer is filled by an inspissated mucus, or a little dirty white slough which may be picked or washed out, leaving a clear but rough fovea. The ulcerative process is unaccompanied by any appearance of colored vessels, and the adhesive process is, in many instances, conducted by colorless vessels. That the proper vessels of the cornea are capable of secreting adhesive matter, is proved by indelible opacities both with and without breach of its texture, and the healing of interstitial ulcers, without any appearance of a colored vessel.

and their  
modes of  
healing.

The organizing process is, however, in some

instances, performed by colored vessels. Where interstitial ulcers open externally, or pustules of the conjunctiva terminate in ulcers of the cornea, a narrow pencil of vessels is produced from the sclerotic conjunctiva to the breach, which organizes the lymph deposited by the proper vessels of the corneal texture. I have sometimes observed, that the fasciculus of red vessels produced to complete the healing of an ulcer, opening externally, instead of coming from the nearest point of the sclerotic, crosses the greater diameter of the cornea. I have never seen such a production of vessels without a narrow and very delicate substratum of recent lymph in their track; nor have I observed this peculiar deposit distinct from the production of vessels, prior to their appearance, but the vestige of it is discernible for a short time after the vessels have disappeared. In superficial lesions of the cornea, as from the insinuation of a foreign particle, the part is at once restored by adhesion, and marked by a superficial speck; it is only where the ulcerative process has supervened, and the conjunctiva is at the same time destroyed at the mouth of the fovea, that nature adopts the mode of healing by colored vessels. Their office is, I think, limited to the organization of lymph deposited by the transparent vessels of the interlamellar texture; because when the ulcerative process is checked, the ulcer presents the signs of the adhesive action, viz. the



marginal halo of lymph and the contraction and filling up of the fovea, before the vessels of the sclerotic conjunctiva are produced. In this state there is often a pause in the restorative process, when the stimulus of a single injection is followed by the appearance of the red vessels, the effect of which is speedily manifested by a reddish brown tint communicated to the deposit, which soon becomes distinctly vascular. In other words, the adhesive process is already commenced before this phenomenon is observed. I have never seen the conjunctival fasciculus running to a transparent fovea. When the conjunctiva is entire, as in the interstitial ulcer, no red vessel appears, nor can be made to appear by stimulant injections, although they certainly quicken the adhesive process in such cases.

From analogy we are led to conclude that the vessels which secrete are distinct from those which organize the deposit, in open breaches of all textures. An insulated pellicle formed upon the bed of an ulcer never becomes skin, but if it be connected by never so narrow an isthmus with the circumference, its organization is perfected. After the ligature of an artery, the barrier of lymph is deposited from the vessels opening upon the divided cellular membrane, as is evident from its origination within the fissure, but injections show that it is organized by the vessels of the lining membrane. The same

fact is still more clearly demonstrated after the division of the sclerotic conjunctiva, for we may observe a distinct interval to transpire between the deposition of new matter in the breach, and its vascularization by ramuli from the divided trunks. This is in opposition to the opinion of Mr. Hunter, who cites the same example to prove the re-union of divided trunks, or inosculation, an hypothesis which has been totally abandoned, since it has been ascertained that the permanent obliteration of arteries is the invariable consequence of a solution of their continuity, whether occasioned by wound or ulceration. I believe that in all parts the capillaries terminating upon the newly exposed surface furnish the deposit, and that this becomes the matrix of new vessels opened from its ramifications, under the extraordinary impulse consequent upon the obstruction of the trunk.

The appearance of colored vessels upon the conjunctiva of the cornea is to be referred to one or other of the following states, and may be classed accordingly:—1st, To the presence of adhesive inflammation excited by a pustular ulcer of the cornea,\* as in the instance last described, and in that of acute interstitial ulcer opening externally, described at page 114.—2d. To the duration of acute strumous ophthalmia, in which the serous vessels of the cornea are

Colored  
vessels upon  
the cornea.

\* See Plate I. fig. 3.



opened to red blood upon its entire circumference, in the form of radii converging to a centre, to an equal extent of from one to two lines.\* In this case the cornea is more or less obscured.—3d, To a state of chronic inflammation, in which straggling solitary vessels, having a varicose appearance, run to one or more specks, or proceeding from opposite sides of the sclerotic conjunctiva, course over the opaque cornea, and freely anastomose upon it.† This state is a common sequel of the suppurative ophthalmia, whether accompanied or not with the granular conjunctiva tarsi, and which I have been accustomed to designate “chronic inflammation with vascular cornea.” It is commonly seen in cases of disorganized globe and spoiled cornea, from whatever cause.

The first of these conditions I consider adventitious to the adhesive inflammation excited by the open ulcer of the cornea. It is proved to be so by a temporary deposition of a narrow layer of lymph; the direct course of vessels along it to the breach; their passage not always by the nearest route, i. e. from the nearest point of the sclerotic; the entire freedom of the cornea from blood-vessels in other parts; and the spontaneous disappearance of the vessels and the lymph track after the healing of the ulcer. It is the most striking and beautiful instance we

\* See Plate I. fig. 4.

† See Plate I. fig. 5.

have of the handicraft of nature, of the production of vessels in inflammation for a specific purpose, and their gradual contraction to obliteration; the determination ceasing when that purpose is accomplished.

The second and third description of appearances I consider to be produced by the continued vis a tergo overcoming the resistance opposed by the enfeebled tone of the vessels. The second is combined with recent and diffused nebula of the cornea, capable of removal by absorption. The third is as often present with ulcerated specks as with simple opacities, which, whether partial or complete, are seated in the corneal texture, and have usually existed prior to the appearance of the red vessels; and neither the vessels nor the opacities spontaneously disappear. After repeated circular sections of the conjunctiva near the cornea, these vessels undergo contraction, especially under the use of styptic applications, as the liq. plumb. acetatis, and solutions of copper, alum, &c.; that portion of the opaque matter which has been recently deposited, is at the same time absorbed. This operation seems to act beneficially in two ways; viz. by throwing up a barrier of lymph to impede the direct course of the vessels, and by diverting them to the purpose of its organization.

When I speak of the 'production' of vessels,

Production  
of vessels.



I am not unaware of the ambiguity or impreciseness, at least, of the term, which is so different in a mathematical and a general sense. It would be wandering widely from the subject of this treatise, to enter into a discussion of the question, whether the phenomenon of organization is to be referred to an elongation of vessels by virtue of a force operating upon their elasticity, or an occasional formation of ramusculi from contiguous branches. To divided vessels, the former supposition cannot apply, as their orifices, as before observed, become permanently closed. I may observe also, that elongation of vessels implies a looseness of cohesion in the textures in which they appear, which does not belong to the corneal conjunctiva in its healthy state; and the example which I have given in illustration of the formation of vessels, is strikingly opposed, in this respect, to the instance adduced of original vessels acquiring increased capacity, and being rendered, by the color of their contents, conspicuous. In the first, the corneal surface is otherwise healthy; in both the second and third, it is raised, thickened, and more or less disorganized.

We are led by analogy to conclude that the effect of vascular congestion from altered or interrupted texture is only less in degree than that from complete obstruction in larger vessels; viz.

the extension and enlargement of collateral branches. Observe the vessels of the sclerotic conjunctiva in organic amaurosis; of the skin covering indurated tumors; and lastly, the communicating branches after the ligature of the trunk in aneurism. Mr. Hunter thought that the vessels might be formed in a patch of lymph independent of the circulation. "I think," says he, "I have been able to inject what I suspected to be the beginning of a vascular formation in a coagulum, when it could not derive any vessels from the surrounding parts. By injecting the crural artery of a stump above the knee where there was a small pyramidal coagulum, I have filled this coagulum with my injection as if it had been cellular, but there was no regular structure of vessels." Then, likening extravasation under inflammation, and the vascularization of the membranes of the chick to this appearance of a self-organized coagulum, he adds, "I conceive that these parts have a power of forming vessels within themselves."—"But where this coagulum can form an immediate union with the surrounding parts, it either receives vessels first at this surface, or forms vessels first at this union, which communicate with those of the surrounding surface."\*

The infinitely more probable theory of the derivation of vessels from parent branches for

\* Hunter on the Blood, p. 92 et seq.



the organization of deposits, is supported by daily observation, and satisfactorily explains the phenomena.

That vessels are capable of producing their kind is as certain as the reproduction of bone, tendon, and other elementary textures. The dependence of one order of vessels upon another is shown by the existence of *vasa vasorum*; and it seems to me most probable that the vessels first seen in a patch of lymph are fabricated by the *vasa vasorum*, blood-vessels and absorbents, of the terminal vessels of the wounded surface.

Acute interstitial ulcer of the cornea.

The acute interstitial ulcer sometimes opens externally, by absorption of the conjunctiva, as well as of that portion of the lamellæ superjacent to it. Its figure and extent are determined by the deposit of adhesive matter. This is frequently crescentic, and traverses a part or the whole of the diameter of the cornea. It is most frequently situated near the upper or lower margin of the cornea, but occasionally crosses the centre. Upon close examination, the conjunctiva will be found to be absorbed at the part opposite to the ulcer, and the exposed scabrous surface of the cornea renders the motions of the upper lid acutely painful. The deposition of the adhesive track precedes the appearance of red vessels, which are derived to it in one or more fasciculi from the sclerotic conjunctiva, and by which its

healing is perfected, as in the ulcer opening from the surface, before described.

The terms onyx and unguis have been indiscriminately applied to extensive collections of lymph and pus between the layers of the cornea, and to similar collections situated in the anterior chamber. They are applicable only to the crescentic interlamellar depositions above described. The acute interstitial ulcer in debilitated habits of body, or when it is produced by considerable violence done to the cornea, instead of opening upon either of its surfaces, spreads between its lamellæ, and in this case a considerable quantity of puriform matter is secreted. If it occupy a large and central portion of the cornea, it usually terminates by slough of the entire membrane.

Onyx and unguis.

When, as more frequently happens, the interstitial ulcer opens into the anterior chamber, it produces the hypopion, which is a mixed secretion of lymph and pus; the former flaky and inorganizable, and situated exterior to the fluid. The soft lymph pendulous from the mouth of the ulcer is often observed connecting the hypopion with it. I never recollect to have seen the puriform hypopion unaccompanied with ulcer penetrating the interior lamella of the cornea. In these cases, the integrity of the chamber is preserved, and the iris has no share in the restoration.

Hypopion.



*Procidencia  
iridis.*

When the external ulcer opens into the anterior chamber, so that the cornea is perforated, the iris falls into the breach and becomes united to it by adhesion. When the opening is small, as from simple ulceration, the iris presents a small black point; if large, as from sloughing, the opposed portion of the iris is protruded in the form of a little sac; and when this happens, the adhesive process is sometimes set up at once, by its pressure on the margin of the aperture in the cornea, and by the ready disposition of the iris to take on a corresponding action. But the adhesive process is not so promptly set up in this case as in the *procidencia* following a wound, and the prolapsus often increases in size until a stimulant is employed. The healing action is marked by a dusky white line at the verge of the opening. The contraction of the chamber and the disfiguration of the pupil are proportioned to the extent of the prolapsus.

*Procidencia  
corneæ.*

In the progress of an external ulcer to the interior of the cornea, and before it penetrates into the chamber, a remarkable appearance is occasionally presented, viz. a transparent vesicle, which fills the aperture, and is supposed to be the membrane of the aqueous humor. I have never seen this state maintained; the prolapsus *iridis* follows in a few hours, notwithstanding the use of the lunar caustic and other stimulants. This has led me to question its being a distinct texture, and

its appearance corresponds accurately to that of the innermost lamella of the cornea, which after losing its support yields to the pressure of the humor, and assumes the vesicular form. The falling-in, or cup-like depression of the continuous surface of the cornea, where the circumscribed interstitial ulcer opens into the anterior chamber, serves to confirm this conjecture. I have never seen the appearance which I have heard others describe as demonstrating the adhesive inflammation of this tunic, viz. colored vessels arborescent upon a deposit of lymph coating the interior of the cornea. If the observation be accurate, the case is very rare.

Chronic interstitial ulcers from pure ulcerative absorption, succeed to acute inflammation, in which large quantities of blood have been lost, and occur frequently in children imperfectly nourished, or adults much debilitated from various causes. The cornea is perfectly transparent, but indented, like a bonce when struck upon a marble hearth, or pitted, according as the ulcers are diffused, or circumscribed; the vision is very slightly affected. Under nutritive diet, effective tonics, and moderate topical stimuli (vin. opii. sulph. zinci) they become hazy; and this denotes the commencement of the adhesive action.

Chronic interstitial ulcer of the cornea.

Opacities are of three kinds:

Opacities: their kinds.

1. Thickening of the conjunctiva and effusion



of adhesive matter between it and the cornea, or between the lamellæ of the latter. This is commonly the product of acute strumous ophthalmia. The corneal surface loses its smoothness and polish as well as its transparency, when the conjunctiva is affected; when the opacity is in the cornea this is not the case. This is that form of opacity which, while recent, admits of removal by excitement of the absorbents more or less completely, and especially by that which mercury produces, the corneal texture being sound. It is what I have been accustomed to call nebula, and though presenting great varieties of shade, it has a soft, diffused, semi-transparent character.

2. A slow change of texture without breach, similar to that by which the pleura, or choroid, or capsule of the lens is converted into bone. The yellow pearly opacity, resembling the inside of an oyster-shell, is of this kind. It is the result of continued, or frequently relapsing (strumous) rather than violent inflammation, and is deep-seated. In this case the layers of the cornea become opaque, indurated, and condensed, so as not to admit of separation by the knife or maceration; and if such opacities are in any degree relieved, it is by an absorption of the interlamellar deposit in their vicinity.

3. New matter, supplying an absolute loss of substance of the cornea, from ulceration or gangrene. This differs from the second chiefly in

its figure being more abruptly circumscribed, and bearing more resemblance to a cicatrix. In point of density the second often exceeds it, as when the cicatrix does not penetrate the cornea. Both these species of opacity are white in the recent state, and in general the more conspicuously their color is contrasted with this appearance, as yellow or brown, the less is the probability of reducing them.

The peculiar hue and loss of tension, as well as lustre, of the *dead* cornea in acute suppurative ophthalmia, has been aptly pictured by Mr. Saunders, by the terms 'cindery, ragged, flocculent.' It is important, because I have satisfied myself that the first change of the cornea in this disease is purely nebulous, produced by the deposition of adhesive matter; and if the inflammation be arrested even on the verge of gangrene, the cornea is susceptible of restoration by absorption. This fact I had lately an opportunity of establishing, in the case of a lady who was rendered blind by acute suppurative inflammation of the conjunctiva: so inevitable to all appearance was the destruction of the cornea, which had sloughed in a deep sulcus at its junction with the sclerotic above, that the most experienced practitioner of my acquaintance in this branch of surgery pronounced the case hopeless and irremediable, and took his leave. The highest tonic regimen, bark, wine, and opium, followed close upon a very active and bold depletion, and the

Gangrene preceded by adhesive deposition in the acute suppurative ophthalmia.



anterior chamber was fortunately and unexpectedly preserved. No sooner was a sign of the arrest of sloughing ulceration obtained, than I commenced a mercurial course; in three days the system was affected; the recovery of the figure and transparency of the cornea was rapid and complete beyond all expectation, and an equally perfect state of vision was restored and established.

The gangrenous opacities of the cornea produced by lime or other substances destroying its texture, are sometimes superficial and defined in extent, and a process resembling exfoliation ensues. More frequently this disorganization is integral and complete. The cornea, disorganized by acids, is rendered instantly opaque, shrivelled, and of a yellow color, almost resembling a piece of wash leather.

In general, opacities which have a recent diffused semi-transparent character (nebulous) admit of absorption; not so those in which the interstitial deposition has been abundant and of long standing, and the lamellæ are compacted, owing to the entire obliteration of the cellular texture; or in which a new portion of cornea is formed.

I have seen several cases of the conversion of the conjunctiva into a skin, rugous and opaque,

knitting the lids close to the globe, so as to obliterate the sinus palpebrales. I have called it cuticular conjunctiva. In these cases there is no secretion of tears. I have had occasion to observe the accession and progress of this disease, in early and advanced life, among the sequelæ of chronic inflammation of the conjunctiva, and am disposed to consider it depending on an obliteration of the ducts of the lacrymal gland.

All stimulant substances, not escharotic, applied to remove opacities of the cornea, act in the same manner as rubefacients upon the skin; they excite a temporary vascular action, which is followed by a corresponding excitement of the absorbents. I have often seen an opaque portion of the cornea cleared by a puncture with the couching needle. If the point of salutary excitement is exceeded, the increased vascularity is permanent, and occasions increased deposition. Injections applied to ulcers do not excite the absorbent action in the same ratio, but occasion a permanent increase of the vascular action, which is here below the ordinary standard. This instance of the adaptation of the same means to different ends, according to the state of the part, is perhaps the best practical illustration of Mr. Hunter's quaint but expressive phrase, "stimulus of necessity."

Action of  
topical  
stimuli on  
opacities

Opaque specks, even cicatrices, are obliterated

Absorption



during  
growth.

during the period of growth, and, as is observed of cicatrices in other parts of the body, change their relative position remarkably in the years of growth. Thus, a conspicuous speck, encroaching upon the pupil in the infant, becomes a small and scarce discernible speck in the grown child, situated near the verge of the cornea. In other instances, if originally small, it has disappeared altogether. Like cicatrices of other parts, they are always considerably smaller than the ulcers of which they are the vestiges.

Staphyloma;  
its kinds.

The staphyloma is of two kinds, viz. from dilatation and from breach. In the first case the corneal lamellæ have not completely given way, but are only bulged, the membrane having been so far weakened by ulceration as to have lost its due resistance, and the staphyloma consequently affects the whole diameter of the cornea (spheroidal). In the second the staphyloma is circumscribed, the recent lymph, corresponding to the breach, yielding at one or more points to the pressure of the humor (conoidal).

In the spheroidal staphyloma the effect of pressure is to thicken the remaining lamellæ by a deposition of adhesive matter, as in the aneurismal and herniary sac; so that the transverse section of it greatly exceeds in diameter that of the healthy cornea. In the conoidal staphyloma the recently deposited matter yields to the

pressure a tergo before its organization is complete. Sometimes the two forms are combined, and one or more conoidal protrusions are formed upon a spheroidal staphyloma. When, as in the first case, the corneal texture has not actually given way, the anterior chamber is in most instances preserved, and the iris is free. In the other case the iris is usually complicated with the staphyloma. The remediableness of the deformity occasioned by the disease, by means of an operation, depends upon a sufficient portion of the iris being left. The iris is kindly disposed to granulate, as we see in sloughs of the entire cornea from suppurative ophthalmia.\*

Three or four days after the operation for staphyloma, the iris is seen coalescing with the conjunctiva, and throwing up fleshy pullulations, which contracting into a little button-like eminence, seal up and permanently secure the crystalline and vitreous humors; thus the spherical figure of the globe is preserved to support the lids. But if the section be posterior to the plane of the iris, the vitreous humor escapes, and the globe collapses and sinks in the socket.

\* I have seen in such cases a permanent staphyloma of the iris, the pupil closed, and the exposed iris retaining its character, occupying the situation and presenting the figure of the cornea. The case is not common, as the cicatrization of the pupil must precede the fall of the cornea.



Conical  
cornea.

The cornea is occasionally subject to a process of thinning, or an absorption of its interlamellar texture, and in consequence, loses its natural tonic resistance to the pressure of the contents of the globe. It usually assumes a conoidal figure, but this is not always the case; the projection of the cornea is sometimes uniform, describing the segment of a larger sphere. The apex of the cone corresponding to the centre of the cornea, when this figure is assumed, exhibits a degree of tenuity and brilliancy which gives it the appearance of a pellucid fluid, like a dew-drop suspended. The patient's vision becomes so inconveniently short as to render objects confused at a very moderate distance; the change is sometimes slow, occupying months, and even years; and on the contrary, I have seen it produced in its greatest extent in the short space of eight weeks: both eyes are generally affected, though not always in the same degree. The disease is unprecedented by inflammation, or any obvious assignable cause; it is more frequent in women than in men, and in my experience affects the periods of youth and middle life. I have never seen it commencing in infancy or old age. It is as much the disease of the robust as of the weakly constitution and frame of body. If inflammation is excited by stimulants, the apex of the cornea turns opaque; if left to itself, the cornea does not give way, but remains in the condition de-

scribed. No remedy yet proposed has been followed by a beneficial result; but a pupillar aperture set in a black ring frame, about a quarter of an inch or more in depth, greatly assists the patient by lessening the confusedness of his vision.

It seems that the presence of adhesive inflammation is the chief distinction between the staphyloma from dilatation and the conical cornea: the absorption is therefore in the one case the ulcerative, and in the other the interstitial.

The last stage of healing in external ulcers of the cornea exhibits beautifully the third kind of absorption, viz. the modelling, as described by Mr. Hunter. We see it in the lowering, rounding, and smoothing of the jagged edges of the cup of the ulcer, a compromise in some sort with the full and complete finishing of the adhesive process, viz. indelible opacity: and in some cases this state of imperfect restoration is perpetuated to a manifest advantage; for a transparent indentation occasions little, if any impediment to vision.

Finishing  
process in  
ulcers.



### SECTION III.

---

#### SCLEROTICA.

Vessels of  
the sclerotic.

BRANCHES from the straight vessels of the conjunctiva penetrate the sclerotic obliquely towards the margin of the cornea, and the long ciliary vessels pass in sulci of this membrane to the plexus ciliaris at the root of the iris. At the interior border of the sclerotic, where the annulus ciliaris is adhering closely to this tunic, the ciliary communicate with the muscular branches, and being in deep-seated inflammation fully injected with red blood, the condensation of color gives the well known and remarkable appearance of a vascular zone at the margin of the cornea. Injections do not demonstrate this anastomosis; for the communicating vessels, like those which are continued upon the cornea, are too delicate to admit of artificial injection, and only admit red blood after a strong and steadily supported inflammatory action.\* When once they have received red blood, they very slowly recover their healthy

\* A very successful injection of an eye in the state of acute iritis could alone demonstrate this fact to the entire satisfaction of anatomists.

calibre, as is proved by the faint appearance of the zone long after the inflammation has ceased, and the almost instantaneous reproduction of the state of congestion, on forcibly separating the lids. When an inflammation at first affecting only the conjunctiva is allowed to progress, the ciliary vessels partake of the action, and this sign of the extension of it to the interior tunics makes its appearance. But the sclerotic from its situation and texture serves as a shield to the finer tunics, from external inflammation as well as from external violence. By the interposition of the sclerotic the vascular communication of the choroid and conjunctiva is rendered extremely minute and anastomotic; and for this reason inflammation of the conjunctiva may and often does reach to a considerable height, without any indication of its extending to the parts beneath the sclerotic. An acute and obstinate inflammation of the conjunctiva, not threatening injury to the cornea, as the purulent, and that with puriform discharge, does not in any degree affect the choroid and iris. On the other hand, when inflammation has extended to these tunics, the vision is affected in a much greater degree than appearances would often lead us to expect. I only mean to remark that if the transmission of blood to the deeper seated tunics had followed readily to that of the conjunctiva, the consequences of every severe

Situation  
and texture,  
their effect.



superficial ophthalmia would have been mischievous.

Ordinary  
inflammation of the  
sclerotic,  
secondary.

When the sclerotic partakes of the inflammation of the conjunctiva, for it is only as intermediate to the conjunctiva and the other tunics that it is usually affected, the vessels which pursue a straight course to the margin of the cornea are strongly distinguished. They have a somewhat brighter hue than the areolar vessels upon the loose portion of the conjunctiva. I have exhibited this difference of arrangement of the vessels in the sclerotic and conjunctival inflammation, as seen in a singular case of inflammation accompanied with pustules.\* This appearance is always observed, as sclerotic inflammation is always present, during the morbid changes upon the cornea, viz. interlamellar deposits of lymph and ulcerations.

Sclerotitis;

I have occasionally observed in a recent ophthalmia this turgescence of the straight vessels, unaccompanied by any affection of the cornea or iris, and with so slight a vascularity of the loose conjunctiva, as to give reason to consider it a primary sclerotitis. The inflammation is more obstinate than acute; the motions of the ball are painful. By continuance the cornea becomes

\* See Plate 1. fig. 6.

nebulous, and the surface roughened, from effusion beneath the conjunctiva.

The inflammation of the sclerotic sometimes accompanies, and is sometimes vicarious with rheumatic inflammation. This is not surprising, as its texture is of the same class with the ligaments of the joints. The rheumatic ophthalmia presents the zonular arrangement of the vessels, more or less cloudiness of the aqueous humor, and a pupil displaced or drawn a little to one side. It is often seen in company with, or following gonorrhœa, eruptions, or sore throat of a pseudo-syphilitic character; and the pains to which, in my experience, it is generally allied, are those which succeed to the exhibition of mercury.\*

and rheumat-  
ic ophthal-  
mia.

The sclerotic, although a firm texture, possesses in certain persons such a degree of tenuity and consequent transparency, as to convey an obscure tint of the subjacent choroid in the vicinity of the cornea. This is most observable in those of light-colored hair and iris, and in persons of lax and weakly habits. It is relatively thinner in such persons. It is evidently slenderer in its texture adjoining the cornea, than elsewhere. A morbid discoloration of the sclerotic is usually combined with an organic amaurosis, whether congenital or induced by inflammation or its consequences. The sclerotic sometimes yields in the spheroidal staphyloma, a disease

Staphyloma  
of the scler-  
otic.

\* See note A.



proper to the cornea, so much as greatly to increase the conspicuousness and deformity of the disease. This happens in hydropic and other degenerations of the humors. It also frequently becomes attenuated or bulged near its junction with the cornea, in the amaurosis which follows inflammation of the choroid. This protrusion, larger or smaller, is sometimes circumscribed, and in other instances diffused over a large portion of the ball.\* It is often seen encircling the cornea, and presenting a sacculated or pouched appearance. It has a blueish grey tint, and the globe is of course misshapen in proportion to its size. An increase in size of the whole globe, or hydrophos oculi, is often joined with it.

Persons who are not aware that it is the result of a chronic process, viz. an interstitial absorption of the sclerotic, sometimes mistake it, from some faint resemblance to the black fungoid tumor, for a malignant disease. I have heard it pronounced a fungus hæmatodes, and the extirpation of the organ advised. I think it may not improperly be designated *staphyloma scleroticæ*.

Staphyloma  
of the cho-  
roid.

I have met with one or two instances of the actual protrusion of the choroid at the margin of the cornea, which had the appearance of resulting from the separation of the sclerotic; like the *staphyloma iridis* from the fall of the cornea, described page 123.

\* See Plate I. fig. 7.

## SECTION IV.

---

### CHOROID AND IRIS.

THE appearance of the vascular zone at the margin of the cornea, which, taken by itself, is a sign of the inflammation having extended to the sclerotic,—if accompanied with dulness of the humors, a spastic contraction, or a very sluggish and limited motion of the pupil, an impatience of light, and a considerable dimness of vision,—demonstrates that the choroid and iris participate in the inflammation. We ought to consider, that the local and vascular relations of the choroid and iris, distinct as they certainly are both in texture and properties, are such as to make it exceedingly improbable that the one should not, in all cases, participate more or less in the inflammation of the other. We are permitted to see the primary changes induced by disease in the living organ, upon the iris only, and we have not as yet any precise marks by which we can ascertain the commencement of inflammation in the one or the other texture. It is probable, however, as the iritis presents considerable varieties in its form, its access and progress, relatively to the superficial inflammation, and the kind and degree of pain and dimness

Choroiditis.



which accompany it, that the choroid is the seat of the primary inflammation in those cases, in which the changes upon the iris take place later than the other signs of internal inflammation, viz. the arrangement of the vessels, the pain, and the obscurity of vision. I have often seen cases of this description which I have felt disposed to denominate "Choroiditis."

Iritis.

The indications above mentioned, are still further confirmed by the presence of an habitual aching pain affecting the globe of the eye, forehead, and region of the orbit, and by certain appearances of inflammation upon the iris, as hair-like red vessels and specks of extravasated blood in its substance. Adhesive inflammation takes place between the fibres of this muscle; the pupil loses its thin flowing edge, and becomes thick, stunted, and gibbous. Iritis of moderate acuteness is often unaccompanied by any other appearance of inflammation; there is no distinct deposit of lymph, and it is rather inferred from the fixedness or slight change of figure of the pupil, than demonstrated.\* I believe the adhesive matter, in this case, is deposited on the posterior surface, formerly called uvea, for in the course of a few days, the opacity of the capsule of the crystalline, and the co-adhesion with it of the pupillary margin, becomes evident, provided the inflammation be unchecked. In this form of in-

\* See Plate I. fig. 2.

flammation the pain is often augmented in the evening, or at an early hour of the morning, to such intensity as to compel the patient to rise, and even totally to deprive him of rest. Sometimes the pain affects the whole corresponding side of the head. In other instances, it is confined to the eyeball and its immediate vicinity, as the forehead and temple, and bones of the cheek. The sensation is sometimes that of pulsatile pain, marking every injection of the ophthalmic artery, as of the radial artery in a whitloe. A sense of continued pressure or constriction, as from extreme distention of the vessels, is the more common character of the patient's sufferings. In the vehement acute iritis, lymph is variously deposited upon the face of the membrane, in small tufts here and there, or large tubercular masses. The pupil, in this case, is usually much misshapen, being rendered angular at those points of the circle at which the deposit has taken place, or is most abundant. Its aperture is sometimes partially covered, and sometimes completely blocked by a mass of lymph. The pain, in this state, is not always augmented in proportion. It affects more the head than the organ. The vision is nearly, if not quite extinguished. The appearance of a stratum of lymph, coating the face of the iris, with a turbid state of the aqueous humor, belongs to chronic inflammation, which tends to opacity of the capsule of the lens, and constriction of the pupil.



## Primary.

A primary inflammation of the iris, as for example, from syphilis, or from mercury, is distinguished from the secondary, or that by extension from the conjunctiva, by the more sparing vascularity of the conjunctiva, and consequently more distinct and conspicuous appearance of the vascular zone.\* The attack is more sudden, the pain in the region of the orbit and head, commences with the inflammation, and is more severe; the vision is more quickly and completely bedimmed. The effusion of lymph is *en masse*, and the disfiguration of the pupil greater.

## Secondary.

In the inflammation of the iris by continuity, the conjunctival vascularity is more conspicuous and diffused, and the cornea is so much clouded, as partially to obscure the view of the iris; the albuminous deposit is wanting, or if any has taken place, it is small in quantity, white, flocculent, and partially diffused in the aqueous humor, or is deposited at the ciliary margin of the iris, forming a lymphatic hypopion; the pupil is little, if at all, misshapen. The pain in the secondary iritis is usually confined to the ball, and is comparatively inconsiderable. Although the vision is much bedimmed, there is greater susceptibility to the painful impression of light. This state I have heard others describe as the adhesive inflammation of the anterior chamber.

\* See note B.

The terminations of iritis, if unsubdued, are, Terminations of iritis.  
 1st, constricted or closed pupil, with opaque capsule; 2d, co-adhesion of the iris and cornea, partial or entire, the former assuming the convexity of the latter; 3d, organic amaurosis, followed by disfiguration of the globe, and often by protrusions of the choroid and sclerotic.

Iritis, as I have formerly observed, is very frequently in company with, or succeeding to syphilis, and the symptoms called mercurial, as peculiar eruptions, sore throat, and pains of a rheumatic character. Primary iritis is rarely seen unaccompanied or unpreceded by such symptoms. I have never said or thought that it could not exist independently of these symptoms, and their supposed causes, having seen such instances. But I have since had additional opportunities of confirming the facts before advanced, that where mercury had been used in various ways before the iris was affected, and before the other symptoms appeared which were referred to its use;—where the primary affection was either altogether questionable, or at most a gonorrhœa, or a superficial sore, which healed by a simple topical application—the iritis has yielded to the steadily supported influence of mercury upon the system, in a manner the most satisfactory; and that no other remedy with which I am acquainted, was competent to this effect.\* Mercurial iritis.

\* I think it right to state, that the salutary effect of mercury



Morbid  
changes of  
the iris.

The iris undergoes a change of color as well as texture, by a continuance of inflammation. This is owing to the loss of its transparency, and the interruption given to its proper secretion by the lymph deposited upon its posterior surface. The healthy iris is transparent, as may be seen in the albino, white rabbit, and ferret; hence the use of the pigmentum nigrum.

It suffers a loss of mobility from the agglutination of its fibres, and ultimately of its posterior surface to the tunica hyaloidea, by which the posterior chamber is annihilated. It is from this morbid condition extending to the plicæ ciliares, that the loss of figure of the globe, or the staphyloma of the sclerotic and choroid, results. A notable thickening and rigidity, a leather-like toughness of the iris, and a varicose state of its vessels, are changes accompanying

in iritis, unassociated with any specific action, was an observation made at the same time by Dr. Farre and myself, at the 'London Infirmary for Diseases of the Eye.' It was first given to the public in the second edition of Mr. Saunders's work, then in the press. I am quite satisfied that the observation was original, notwithstanding all the attempts of the German scholars to convince us, that at Vienna, and elsewhere, it was a matter "*lippiis et tonsoribus notum*."

I shall further add, that I am unacquainted with any fact in Medical Surgery which ranks with this in point of importance; whether we consider the urgency and frequency of the occasion, or the indispensable necessity, and almost unerring efficacy of the remedy.

the state of chronic closed pupil, after reiterated attacks of inflammation.

A morbid change, sometimes, but rarely witnessed, is the conversion of the choroid into a shell of osseous matter.

I shall have occasion to speak of others under another head.\*

\* There is a mild chronic form of iritis, so little resembling the acute disease, and so insidious in its progress, that the greatest attention is necessary in its diagnosis. The following case is a good specimen of this affection.

An eminent upholsterer of this city applied to me, some time since, complaining that he was gradually losing the sight of one of his eyes; and that the loss of vision had been slowly progressing for several months. The only other local symptom which attended the case, was a constant deep-seated pain in the back part of the head, on the affected side, together with a continual sense of coldness there; and these sensations had existed since the patient first observed his vision beginning to fail. He had been labouring under dyspepsia for several years; and as not the least trace of redness, or other ordinary evidence of inflammation, appeared in the eye at the time, or ever had done so, the case was at once suspected to be amaurosis, dependent upon disorder of the digestive functions. On a closer examination of the organ, however, the lower edge of the pupil was observed to be irregular, inverted, and adherent to the capsule of the lens, but not sufficiently so to strike the observer without very accurate attention. No other symptom of disease of the eye existed; but this last became much more evident upon the application of extract of stramonium, and consequent dilatation of the pupil. This, however, taken in conjunction



with the loss of sight, and the pain in the occiput, was a decided proof that the disease was chronic iritis. Accordingly, the patient was cupped, and immediately commenced the use of mercury. The impediment to vision quickly disappeared, the pain subsided, and in a few weeks the patient was cured.

Two months afterwards, the other eye was attacked in a similar manner, and cured by the same remedies.

Other cases of the same character have occurred to me, but I have noticed that they all, like this, were connected with a disordered state of the digestive organs.

I have remarked one fact as connected with syphilitic iritis, which I do not remember to have seen noticed elsewhere. The disease is very frequently accompanied with tinnitus aurium and deafness; the organ of hearing apparently taking on the same congested condition as the eye. I have observed this symptom in protracted and badly treated cases alone, never in recent ones, nor in those which had not been preceded by syphilis.—*Editor.*

## SECTION V.

---

### RETINA.

THE retina is sometimes, though rarely, the seat of inflammation; but it is an error to suppose that intolerance of light is a sign of this affection, as is clearly proved in the strumous ophthalmia, in which, although the intolerance is in excess, the retina is uninjured; and, secondly, because the effect of inflammation upon a nerve of sense is to produce direct palsy, not increased excitability. Inflammation of the passages and auxiliary textures of the organs of sense may render the impression of their natural stimuli painful; but here, as in the case before referred to, and probably in all cases, the increased acuteness of the sense is sympathetic. The organic sensibility, we may conclude, is increased in the sentient, as in other organs, by inflammation. This is probable from the first and predominant symptom of inflamed retina, viz. a sudden attack of vehement dashing pain of the most distracting kind, which is described to extend from the bottom of the eyeball to the occiput, or in the reverse direction, and the supervention, within a few hours, of total blindness, with occasional sparks and flashes of vivid

Inflammation of the retina.



light. The pupil, upon inspection, is gaping and motionless, as in confirmed amaurosis, and the humors are thick and muddy. The external signs of inflammation are in the commencement disproportionate, and quite insufficient to account for the symptoms.

Accompanied with inflammation of the other tunics.

In some cases, however, the signs of choroid inflammation are present with the attack of pain and the loss of sight. The pupil is not thrown open, but it is without motion. In addition to diffused vascularity of the conjunctiva, the straight ciliary vessels are remarkably loaded, so as to give a livid red hue to the sclerotic around the cornea. The pupil becomes in a few days plugged with lymph, or the whole iris bulges forward, changes color, and the crystalline turns opaque; or instead of this, the same splendid tapetum-like appearance presents itself which is observed in the commencement of the medullary fungus, upon looking obliquely through the pupil. The pain in this attack is accompanied with a sense of confusion so alarming, that the patient apprehends the loss of his intellects. I once saw the disease marked throughout with so much disturbance of the nervous system, e. g. vigilance, temporary wanderings, catches of the muscles of the face, startings and frightful dreams, in the short intervals of repose from exhaustion,—coupled with a sense of heat, constriction, and tenderness of the whole scalp,—that I was at first disposed to consider the

ophthalmia as secondary, and subordinate to inflammation of the brain or its membranes. When the internal signs of inflammation are less obvious, and the humors and internal tunics undergo a slow but complete disorganization in the progress of the disease, meteoric flashes are frequent even after the inflammation has run its course; and I have known patients gratified with this *ignis fatuus*, although conscious that it was no more. I have seldom seen an example of this inflammation, which seemed to afford time for the beneficial operation of a remedy. I have in more than one instance given a full trial to the lancet, and the immediate operation of mercury; but though both were carried as far as could be permitted with safety, the vision was lost. In others the external inflammation has been subdued, and the vision has been recovered so far as to enable the person to distinguish surrounding objects with tolerable precision; but the gaping and motionless pupil, the discolored humors, and the superficial congestion, which remained, afforded little hope of its continuance. One lamentable instance occurred under my observation, of its destroying both eyes in a middle-aged lady within the short interval of a fortnight. She expressed, in the agony of her suffering, a conviction that she must either lose her sight or her senses.

Amaurotic affections, as is well known, differ infinitely in degree, but they differ also in kind; and this affords a more scientific basis of classi-

Amaurosis,  
organic and  
functional.



fication. I divide them into two classes, the organic and the functional. The first comprehends alterations, however induced, in the texture or position of the retina, optic nerve, or thalamus. The second includes suspension or loss of function of the retina and optic organ, depending upon a change either in the action of the vessels, or in the tone of the sentient apparatus.

As causes of the first, we may enumerate,

1. Lesion, extravasation of blood, inflammatory deposition upon either of its surfaces, and loss of transparency of the retina.

2. Morbid growths within the eyeball, dropsy, atrophy, and all such disorganizations as directly oppress or derange the texture of the retina.

3. The state of apoplexy, hydrocephalus, tumors or abscesses in or upon the brain, the optic nerve, or its sheath; and thickening, extenuation, absorption, or ossification of the latter.

As causes of the second,

1. Temporary determination; vascular congestion or vacuity, as from visceral and cerebral irritation; suppressed or deranged, or excessive secretions, as of the liver, kidneys, uterus,

mammæ, and testes ; various forms of injury and disease ; and sudden translations of remote morbid actions.

2. Paralysis idiopathica,\* suspension, or exhaustion of sensorial power from various constitutional and local causes ; from undue excitement or exertion of the visual faculty ; and from the deleterious action of poisons on the nervous system, as lead, mercury, &c.

From this description it will be understood, that organic, and many forms of functional amaurosis are incurable ; and the functional, by continuance, lapses into the organic disease.

\* To apply the term *paralysis* to a nervous tissue, is, to say the least of it, a misnomer. A muscle may be paralysed from pressure or injury of the nerve which supplies it, or the part of the brain or spinal marrow whence that nerve is derived ; but there is certainly nothing in the nerve itself sufficiently analogous to muscular structure or function to apply the term 'paralysis' to both. We have indeed no evidence that a nervous tissue ever loses its function independently of some lesion either of itself or its origin, except where it is affected by sympathy with some other part. To attribute amaurosis, therefore, to idiopathic palsy is entirely gratuitous ; and the result of practice directed to the removal of such a condition of the parts is never sufficiently encouraging to countenance the theory upon which it is predicated. Most, if not all the cases of amaurosis, which are attributed to idiopathic palsy of the retina, are more probably dependent upon some antecedent congestion of the organ, or some other of the causes enumerated by our author.—*Editor.*



Even under the continued suspension of function, much more the duration of a state of excitement, the power of the retina, as of other parts, gradually fades, and is at length exhausted. Thus the removal of a cataract from the eye of a person who had been the subject of the disease for thirty years, was unsuccessful in restoring useful vision. This was a sensorial defect, for the eye had every appearance of health, both before and after the operation.

I am aware of the objections to which this, like most other attempts at a scientific arrangement of such subjects, consistent with practical views, is exposed. Thus the comprehending under the same heads the states of temporary and permanent congestion of the vessels of the retina and brain, and the disordered actions of the vascular and proper texture of the retina, may, *primâ facie*, appear to be examples of incongruity. But for the purpose of descriptive arrangement, a line of division must be somewhere drawn; and opposed to the gradual and often imperceptible transition from functional to organic disease, this division must appear more or less forced and artificial. I conclude that the difference between the disposition to apoplexy and the state of apoplexy, may be acknowledged in the eye as well as in the brain, and that the purely functional irregularities of the former organ as a whole, may be classed with

as much propriety as those of the heart or the stomach.

In treating of the disorders of any sense or function, I deem it an essential character of a scientific arrangement, to include in one view the entire organ, philosophically speaking, subservient to that sense or function: hence, the brain and retina should not be considered separately, but in conjunction. Secondly, to regard the locality and demonstrableness of diseased states, as affording the best ground of division; because the presence or absence of certain external characters affords a stronger distinction between functional and organic disorders, than we could hope to obtain from any analysis of the symptoms characterising the varieties of disordered function, in the present state of our knowledge.

The history and concomitant appearances or morbid states associated with amaurosis, usually indicate to which class it belongs: as for example, diseased changes in the situation or texture of the eyeball, or in the brain. A hemiplegia, or partial paralysis, with other signs of apoplectic or hydrocephalic pressure, whether resulting from an injury of the head or otherwise; or an acute deep-seated inflammation, whether accompanied by a visible opacity or not, point out the organic nature of the affection. I have seen

History, and  
concurrent  
diseases.



such an amaurosis produced by abscess in the cerebral substance, and by the medullary fungus of the cerebrum. On the other hand, I have known the following distinct sources of irritation operating to produce functional amaurosis, viz. wound of the scalp,\* caries of the skull, abscess and caries of the antrum maxillare, with excessive œdema of the integuments of the lids and cheek, a large abscess under the masseter and muscles of the cheek, and an abscess at the extremity of a molar tooth, while the crown of the tooth was sound. In all these cases it is to be understood that the eye was sound, and the orbit was untouched by the disease of the parts in the vicinity, to which the amaurosis was clearly attributable. In like manner, an excessive use, or rather abuse of the visual faculty, the disordered functions of the stomach, liver, uterus, &c., sudden and alarming depletion, excessive or obstinately suppressed secretions, difficult dentition, the presence of worms in the intestinal canal, and the deleterious effects of noxious agents upon the organ or the system, are sufficiently obvious causes of the functional amaurosis.

The professions, circumstances, and habits of patients throw much light on the origin and na-

\* A lesion of the frontal nerve is mentioned among the occasional causes of an amaurosis from Hippocrates downwards. A striking example is reported by Sabatier. *Traité d'Anatomie*, Tom. 3, p. 228.

ture of amaurotic affections. Such as have a direct influence are, sedentary occupations disposing to torpid liver and bowels, combined with the continued exercise of the eye in a depending position of the head upon minute objects;\* in too strong or insufficient light; upon polished reflecting surfaces;† habitual exposure of the organ to a high degree of heat;‡ to acrid fumes and vapours;§ and the customary employment of optical glasses.|| Immoderate grief, excessive indulgence in venery, protracted suckling, continued diarrhœa, repeated hæmorrhages, profuse salivation; and, on the other hand, obstinate amenorrhœa, or constipation of bowels, with determination of blood to the head in a full habit, are ordinary predisposing or constitutional

\* Needle-workers, writers, draughtsmen.

† Inspectors of linen and scarlet cloths, and of new bank-notes; money-counters. It is a curious fact, that several persons so employed at the Bank, at the issue of a new coinage of silver, were affected with symptoms of amaurosis. Color-manufacturers, burnishers, landscape-painters.

‡ Smiths, stokers in iron furnaces and glass-houses, tavern cooks, &c.

§ A wholesale manufacturer of blacking became the subject of gutta serena. He had been a constant superintendant of the process upon a large scale. The mixture of sulphuric acid, with the several ingredients, disengages a pungent and offensive vapor, by which the eyes are very painfully affected.

|| Watch-makers and engravers, philosophical instrument-makers, sea officers.



causes of this disease, as I have had abundant opportunities of learning.\*

Amaurosis, of whichever class, is either perfect or imperfect. The first is marked by total insensibility to light; the second, by defect of vision, infinitely varied in kind and degree. I need scarcely remark, that not only the appearances and symptoms vary, but the essential character of the disease varies in its stages. Thus, an

\* There are strong shades of difference in the cases of these unfortunate persons, as regards the intensity of their feelings under the hopeless privation of sight. The man of pure life has the support of the best philosophy. The literary man has not enjoyed his '*Noctes Atticæ*' in vain; they have provided him with resources. Even the aged voluptuary rises with some degree of complacency as a '*conviva satur*' from the banquet of nature; and contemplating the various evils of the common lot in the circle of his friends, meets his calamity with somewhat of martyr fortitude. The most pitiable is the amaurosis of early life, from excess of sexual indulgence, and especially of solitary vice. The following are strong examples:—A country lad, of robust constitution, became the alternately favoured paramour of two females, his fellow-servants, under the same roof. He was the subject of gutta serena in less than a twelvemonth. Another, at an early period of puberty, suddenly fell into despondency, and shunned society. He never left his chamber but when the shade of night concealed him from observation, and then selected an unfrequented path. It was not discovered until too late, that, in addition to other signs of nervous exhaustion, a palsy of the retina was the consequence of habitual masturbation.

affection, purely functional in its origin, by duration becomes an organic disease.

#### ORGANIC AMAUROSIS.

When the eyeball is the seat of organic amaurosis, it commonly presents some, or all of the following appearances:

Signs of organic change in the eyeball.

1. A pupil fully or preternaturally dilated, contracting feebly, in the first case, on the sudden admission of light, and absolutely motionless in the second. This appearance is not peculiar but common to both classes, though by no means invariable in either.

2. A congestion of the superficial vessels, especially of the long fasciculi of conjunctival veins.

3. A peculiar bluish grey tint of the sclerotic coat; sometimes a degree of bulging or protrusion on one or more sides of the globe; or simply a loss of sphericity, its sides appearing flattened.

4. A diffused turbidity or milkiness, apparently of the vitreous humor, strikingly observable when contrasted with the jetty brightness of a healthy eye. It is little more than the healthy appearance of the humors in the eye of the horse. This state, which the ancients termed glaucoma, is very often mistaken for incipient



cataract; and I have known it called a black cataract, and the operation of extracting the transparent lens performed. It appears deep-seated, diffused, and of uniform density; and in examining some such cases at long intervals, I have not found the appearance vary. The lens remains transparent. There are, however, some cases of a deep-seated opacity so closely resembling that of incipient cataract, that it becomes next to impossible to decide the actual state of the lens. I have seen the latter, upon an experimental extraction in such a case, semi-transparent, and of a bright yellowish tint throughout, and the sight of the patient has been considerably improved. The vision is in general defective in a much greater degree than the visible opacity explains; and this, combined with the depth of the opacity, a dilated and sluggish pupil, and some other symptoms of amaurosis, makes for the opinion that it belongs to the latter class. But where other signs of impaired retina are wanting, and the states of dimness and opacity correspond, the operation would be warrantable, although the site of the opacity should be disputable, if it were the express desire of a patient properly in possession of the circumstances.

5. Another yet more common appearance is that of a white or greenish yellow spot, apparently in the fundus of the eye, a little to one side of the visual axis; sometimes it has a disc of

such breadth and splendor, as to look like the tapetum of sheep, or the colored choroid of fish; but more commonly it occupies a circumscribed annular space, and is seen only in a strong light, and in particular directions of it. Although this appearance is commonly associated with impaired vision, I have now and then seen it in persons who made little, if any complaint of their sight. This appearance has been referred to a circumscribed opacity of the retina, and the central spot supposed to correspond to the porus opticus, or axis of the optic nerve. It has also been conjectured to be the macula lutea of Soemmering.\* It is probably with more propriety to be attributed to a deficient secretion of the choroid pigment, a preternatural adhesion betwixt the choroid and retina, and a discoloration or resplendent appearance of the retina from that cause. I have been led to this opinion from observing it combined with that form of amaurosis in which the vision is confused to dimness in the broad light of day, and is tolerably clear and agreeable after

\* A point of opacity within the reflecting mirror of the eye must, of necessity, create so much delusion, that we can scarcely venture, by inspection, to determine its precise seat. Even the opacities of the cornea and crystalline capsule are liable, in some instances, to be confounded. It is highly doubtful whether anatomy would determine the point in question; except by a nice comparison with the sound organ at a very early period after death, the morbid opacity of the retina would certainly pass undetected.



sunset; and also with that which is disturbed by the partial illumination of objects.

I have also been enabled to make the following observations regarding this appearance. It is neither suddenly induced nor preceded by any signs of inflammation; there is often a degree of blindness joined with it greater than its extent could account for; and a recovery of vision, to which I have been witness under such circumstances, is not accompanied by any perceptible change in the appearance. I am therefore disposed to regard its connection with amaurosis, whatever it may be, as a casual coincidence, a change incidental to age, like the arcus senilis of the cornea; for it is by no means a constant appearance in that disease, nor is it incompatible with useful vision.

Opacity of  
the retina.

I have more than once seen a condensed and palpable opacity at the fundus of the eye succeeding to inflammation of the choroid, which had destroyed vision; and this I have considered to be produced by a change in the texture of the retina. What adds to the probability is, that the crystalline in this case afterwards, as if progressively, becomes opaque; a common sequel of amaurosis induced by inflammation.

Amaurosis  
from deep-  
seated in-  
flammation.

In the amaurosis from inflammation of the choroid or retina, where the diseased action has entirely subsided, the veins of the conjunctiva

are varicose, the iris is discolored, thick, tough, inelastic, and preternaturally vascular; the substance of the crystalline is more or less absorbed, or converted into a fluid and discolored; the vitreous humor is opaque and of a deep yellow color. The retina, like the other transparent textures, becomes opaque under inflammation, and it is probable that under these circumstances, adhesive matter is effused upon the interior of the choroid; this supposition I have never had an opportunity of verifying by dissection, in cases of which the history was known.

I some time ago dissected the eyes of a man who had cataracts with amaurosis. The cataracts had been formed ten years prior to his death; one of them fell down behind the pupil, and he was spontaneously restored to sight, as by a natural couching. Gradually he lost his sight, the eye still remaining plump, and the pupil clear of any opaque substance. The change which had taken place was an interstitial absorption of the vitreous humor, which was proved by the immediate discharge of an unusual quantity of watery humor, on opening the tunics, and the appearance of the vitreous capsule collapsed into a little opaque bag, and adhering to the ciliary body at the inferior margin of the iris. A remnant of the crystalline was involved in it, not exceeding in size a large pin's

From absorption of the vitreous humor and collapse of the retina.



head. The remains of the retina were a mere film or string extending from its attachment, at the back of the globe, to this bag or net of the vitreous capsule. The pupil of the other eye, in which the cataract had retained its place, was closed, and the lens adhered firmly to the iris by its capsule, which was involved with the collapsed tunic of the vitreous lying behind it; for the humor was as completely absorbed in this as in the eye first examined, and the texture of the retina as nearly obliterated. Thus the amaurosis resulted from collapse of the retina, owing to absorption of the vitreous humor. The fall of the lens in one eye resulted from the absorption of the vitreous humor, and would have occurred from the same cause in the other eye, but for the adhesion previously existing betwixt its capsule and the iris. The eyes preserved their figure by the increased secretion of the aqueous humor, which is always in proportion to the default of the vitreous: and could the retina have retained its position, the vision, which was, for a time, recovered, would probably have been retained.

From cerebral tumor.

A young gentleman, the subject of amaurosis in his left eye, was affected with symptoms of a diseased action in the brain; as, deep-seated pain in the fore part of the head on the same side, disposition to sleep, and inability to employ his mind as heretofore. He was repeatedly

blooded and blistered without relief. The digestive functions were much disordered, and he was put upon a course of alterative medicine. The disease, however, advanced; the eyelid became paralytic, and a slight degree of strabismus was accompanied with occasional double vision. The lethargy and the derangement of the secreting organs, and consequent emaciation and debility, increased, and his death soon followed. The eye had no unhealthy appearance; the pupil was regular, and moderately active. On examination, a firm lardaceous tumor, of the size of a garden bean, was found compressing the optic ganglion and nerve at its origin thence, of the same side.

I have seen several cases of amaurosis from concussion, as by a blow on the temple, or the eye. Of these some were attended with signs of disorganization—some were superficially inflamed—and others presented no external appearance of injury. One was the case of a captain of artillery, who was struck by what is called the *wind of a ball*, on the right side of the head. He received no wound, but lost, instantly and irrecoverably, the sight of his right eye.

Amaurosis  
from con-  
cussion.

In another case, a young gentleman received a blow on the eye, by which it was inflamed; the inflammation was superficial and easily subdued, but the vision was so much impaired, that



a surgeon was consulted, who observed the pupil to be dilated and without motion. The iris recovered its activity by depletion, but useful vision was lost. It is not always the eye on the same side of the head which has received the blow that is affected. One remarkable case I shall take the liberty to mention.

A man who had good vision of the right eye, and was nearly dark from a cataract in the left, received a violent blow on the left temple. From that time he lost totally the sight of his right (or perfect) eye, which has now the faded appearance of an incurable organic amaurosis, and owing to the rupture of the lens, which was at the time undergoing absorption, recovered sight with his left eye, which he still enjoys. Frequently the amaurosis from concussion is purely functional, and is cured by a full blood-letting, blisters, and purgatives.

Congenital  
organic  
amaurosis.

I have observed several forms of congenital organic amaurosis: one, in which the organ is preternaturally small and soft, and even flaccid to the touch, as if from deficiency of the vitreous humor; the iris tremulous, and not influenced by the belladonna; the globe affected with an incessant tremor, and not subject to the control of the will. I have often seen this motion of the globe uncombined with the tremulous iris in

cases where there was little more than a natural feebleness of the retina.

A second depends on a deficiency of the pigmentum nigrum; here the tremulous motion of the globe is present; strong light produces uneasiness, and vision is dazzled and confused; the vessels of the choroid give the interior of the eye a deep red tinge, but not the bright scarlet of the albino, or the white rabbit. I have seen several of the children of two families thus affected; they are considerably aided by cylindrical shades, such as are used by connoisseurs in pictures; goggles; glasses covered with black gauze, and every other contrivance to absorb light; even a coat of black varnish besmeared around the eye. I may observe in general, that an amaurotic disposition, greater or less, exists in all persons whose hair and eye-brows approach to white.\*

A third form of congenital amaurosis is that in which the sclerotica so encroaches upon the cornea, that the latter scarcely exceeds the dia-

\* Professor Beer states, that dark colored eyes are more inclined to become amaurotic than those of lighter color. This observation is not at variance with my own, since the above remark refers only to white-haired persons, the characteristic of whose amaurosis is photophobia, or intolerance of light, and in whom the disposition depends on a defective pigment.



meter of the pupil, while the volume of the globe appears somewhat greater than natural. Various malformations of the pupil, an extreme diminutiveness, and even a total deficiency of the iris, are not in my experience ordinarily associated with an imperfectly organized retina.

A fourth kind of congenital amaurosis is unaccompanied by any appearance of organic derangement. The eyes move in concert as if attracted by a faint perception of light, in an oblique direction; but the infant is too certainly blind. A diseased state of the thalami or optic nerves would, I apprehend, be discovered by inspection of these cases after death. It is probable that the opaque retina is sometimes congenital as well as the opaque cornea: of the latter I have seen instances; others are related by the late Mr. Ware.

#### FUNCTIONAL AMAUROSIS.

The functional amaurosis admits of the following subdivisions.

1st. *The symptomatic*, or that which is only a symptom of some general disease or disorder of the system, as for example, general plethora, general debility, &c.

2nd. *The metastatic*, or that produced by the sudden transference of the morbid action from another organ of the body ; as for example, from the skin, the testicle, &c.

3rd. *The proper*, or that which immediately depends upon a peculiar condition of the retina ; as for example, the visus nebulosus, muscæ volitantes, &c.

I proceed to treat of them in succession.

### 1. *Symptomatic.*

Like nervous deafness, amaurosis sometimes follows typhus and scarlet fever and the various forms of acute constitutional disease. This I have several times met with as a result of infantile fevers. It is also sometimes a consequence of chronic wasting diseases, in which organic changes interrupt the nutrition of the system. I have seen a rapid and severe salivation instituted for a remote affection, and where no disease had previously affected the eyes, terminate in gutta serena of both. The same has been observed of the sense of hearing. The state of the circulation has a marked influence upon imperfect amaurosis.\* I know patients whose vision

Amaurosis from constitutional disease and general debility.

\* The effect of fever upon a nervous deafness, was strongly evinced in the case of a lady whom I knew. She had been incurably deaf many years, when, during the existence of a



is benefited in a high degree, and others in whom it is as much deteriorated by the quickened circulation following a full meal and a few glasses of wine. The former are persons of spare and meagre habits; the latter plethoric.

The influence of mental emotion in producing this disease is most frequently seen in the instance of grief. Young widows are peculiarly liable to amaurotic affections; and cases are

puerperal fever, it was remarked, that she had recovered distinct and even acute hearing, which again left her, after the febrile action had subsided.

I was once consulted by a gentleman who was the subject of this species of deafness to a painful degree. He informed me that he was in possession of a remedy for the disease, but unfortunately it was available only while in actual use, and too severe to be employed incessantly. His object was to ascertain if a less objectionable one were known. It consisted in occasional drastic purging, abstemious diet, and the hard daily exercise of a man training to walk against time, or fight, or ride a race. Under this alterative plan he had so excited the action of the extreme vessels as to restore the sentient tone of the nerve. By adopting it, he had more than once recovered a perfect state of hearing, which remained while he had resolution to pursue it. But he thought, as would most others, '*le jeu ne vaut pas la chandelle.*'

While upon this subject, I shall take the liberty of still further extending this digression by adding a much more remarkable proof of the influence of vascular action upon the brain and its more immediate functions, and leave the reader to form his own conclusions.

A young woman, who was employed as a domestic servant

lated, in which what the poet calls 'a short madness' has been productive of the same unhappy consequence.

That form of amaurosis common to protracted suckling, in which the infant may be said to prey upon its mother; and that in which the impaired energy of the whole nervous system, occasioned by various states of physical disorder, shows itself especially in this organ, afford fami-

by the father of the relator when he was a boy, became insane, and at length sunk into a state of perfect idiocy. In this condition she remained for many years, when she was attacked by a typhus fever; and my friend having then practised some time, attended her. He was surprised to observe as the fever advanced, a developement of the mental powers. During that period of the fever, when others were delirious, this patient was entirely rational. She recognised in the face of her medical attendant, the son of her old master, whom she had known so many years before, and she related many circumstances respecting his family, and others which had happened to herself in her earlier days. But alas! it was only the gleam of reason. As the fever abated, clouds again enveloped the mind; she sunk into her former deplorable state, and remained in it until her death, which happened a few years afterwards."—Tuke's Description of the retreat for Insane Persons of the Society of Friends, p. 137.

A man labouring under recent concussion of the brain, and from this cause comatose, was freely let blood, and afterwards took a full dose of emetic tartar. After vomiting, he became immediately sensible.



Amaurosis  
from pletho-  
ric conges-  
tion.

liar instances of the amaurosis from constitutional debility.

Amaurosis depending on vascular congestion is marked by some or all of the following symptoms, viz. dilated and sluggish or immoveable pupil, ptosis or strabismus, and oblique or double vision of the affected eye; a preternatural action of the carotids, flushed face, sense of weight, pain, or stricture of the scalp, lethargy, occasional tinnitus aurium, with greatly disordered and irritable stomach. The patient frequently complains, particularly in straining, stooping, or on first lying down, of seeing luminous sparks or flashes,\* and a reflection of one or more of the choroidal vessels, the visible pulsation of which is a cause of much distress to him. A person thus affected accurately described to me the zona minor iridis, as distinctly presented to his view.

From deter-  
mination of  
blood to the  
head.

A loss of balance in the sanguiferous system, occasioning an undue determination of blood to the head, often exists, distinct from general plethora, and is aggravated by loss of blood. The following case is an example :

\* Persons labouring under dyspepsia are often troubled with this symptom on first closing the eye to sleep; and in the progress of amaurosis from nervous exhaustion, it is sometimes accompanied with the sensation of a crackling or snapping noise.

A young medical man came to me one morning from the country in extreme anxiety, with an earnest solicitation that I would instantly apply a ligature to his carotid artery. This gentleman, aged 25, was of short stature, and constitutionally healthy. His pupils were large, and his countenance was suffused and bore the appearance of preternatural determination of blood to the head. He had been the subject of two attacks of inflammation; one in April, the other in October of the same year; during which he lost upwards of an hundred ounces of blood. He had now a constant heavy pain in the head chiefly over the coronal suture, and in the direction of the sinuses, with tinnitus of the left ear. After stooping, the giddiness was extreme, and a golden colored spot, edged with black, appeared floating before the eye. He had been troubled with *muscæ* in excess, for a year and a half past; he had now fire sparks flashing before the sight, and saw a pulse in the choroid synchronous with that of the wrist.\* When looking at near objects he was not troubled with *muscæ*, but they were always numerous, in proportion as the object was remote. He did not complain of much dimness. His complaints

\* The subjects of chronic iritis, and in whom the pupil is fixed, and the capsule of the lens more or less opaque, are often distressed with this or other signs of undue determination of blood to the head, which is not in any degree relieved by drawing blood.



were not relieved by topical blood-letting. He recovered gradually but perfectly, under a regulated diet, and a course of the blue pill with saline aperients.

Amaurosis  
from inanition.

The amaurosis, from depletion, is sometimes mistaken for its opposite, viz. that from plethoric congestion; this is owing to the coincidence of a dilated and immoveable pupil, *muscæ*, and a deep-seated pain in the head, with occasional vertigo; and its occurrence often in a corpulent habit. It succeeds somewhat abruptly to uterine floodings, and large and sudden depletion for acute diseases. The pain is not confined to the region of the orbit, though it affects chiefly, if not exclusively, the same side of the head; it is that peculiar nervous pain to which women are subject after uterine hæmorrhage, attended with a sense of defined pressure, as of an iron finger on the brain; and sometimes a distressing jarring noise like that of a mill or threshing-floor, or the rattling of the shingles as a heavy wave of the sea recedes. It is perhaps connected with an imperfect injection of the medullary substance. By a cautious use of tonics it is relieved; by whatever lowers or stimulates, whether diet or medicine, it is decidedly aggravated. The vision in this form of amaurosis is further enfeebled by the loss of as much blood as flows from two or three leech-bites. This is not imaginary; I have seen distinctly

marked cases of it, in which large and copious venesection was still urged as the only resource of art. This I consider to be a fatal mistake.

As the causes of amaurosis, to whichever class it belongs, present infinite gradations, so do the imperfections of vision. An unhealthy secretion of the meibomian follicles and caruncula lacrymalis, a very inconsiderable obstruction in the excretories of the tears, or a preternatural excitability of the conjunctival surface from any cause, will give occasion to such a degree of weakness and dimness, as greatly to interfere with, if not to interrupt, the transaction of business. The improvement of the vision in strength and clearness, during the use of astringent lotions and stimulant ointments, is continually observed. Persons affected with an irritable state of the conjunctiva from any cause, frequently complain, not that their sight is indistinct, but weak; by which nevertheless they mean, that they cannot maintain distinct vision for any considerable time together. The retina appears to be sympathetically affected in these cases. They are often accompanied with muscæ, and remarkably benefited by blisters, as well as by applications, which improve the condition of the conjunctiva and eyelids.\*

Amaurosis  
sympathetic  
with irri-  
table con-  
junctiva.

\* See note C.



## FUNCTIONAL AMAUROSIS.

2. *Metastatic.*

Amaurosis  
by metasta-  
sis from the  
chest.

Amaurosis by metastasis is not unfrequent. I have seen it from the state threatening effusion into the chest, from gout in the foot, and swelled testicle ; in all which cases the oppressed organs were suddenly relieved, and the eye as suddenly affected. Thus a person goes to bed with good vision and rises blind.

A lady, above the middle age, who had long been subject to occasional attacks of pulmonary congestion, after one of unusual severity, threatening hydrothorax, was suddenly affected with paralysis of the upper eyelid of the left eye ; the sight was slightly, if at all, impaired. On the following day however, she had totally lost the sight of that eye, the pupil of which was dilated and motionless. On the morning of the third day, the upper eyelid of the right eye was paralysed, and the vision of that eye was also much impaired. On the fourth day the ptosis on the left side had disappeared, and the eyelid on the right side resumed its position. She was now in a state nearly approaching to complete blindness ; both pupils dilated, although the left most so, and nearly, if not quite inactive. The attack was accompanied by a marked simultaneous re-

lief from the threatening symptoms of the original complaint, and occasional fugitive pains across the front and top of the head. Upon inspecting the eyes opposite the light, the appearance described 5, page 150, was conspicuous in both eyes, and led her physicians, who were of the first eminence, to apprehend an incipient opacity in some interior texture. Under the frequent administration of calomel and rhubarb, in moderate doses, and the alternate repeated application of blisters behind the ear, and to the nape of the neck, with abstemious diet, the vision of both eyes was gradually restored.

A man, the subject of hernia humoralis, lost an unusual quantity of blood by the bites of leeches applied to the part. The testes were suddenly and greatly reduced, and he complained of uneasy sensations in his head; to use his own phrase, a pain like opening and shutting.\* Without any other visible sign of amaurosis than a dilated and sluggish state of the pupils, he described a dark screen seeming to rise gradually from below upward, and at length totally ob-

From the  
testis.

\* I once saw an attack of hemiplegia originate under similar circumstances, viz. a sudden reduction of enormously enlarged testes by leeches and cold lotions; and since the first edition of this work was published, I have known a case of fatal apoplexy succeed to a gangrenous inflammation of the scrotum, from extravasation of the fluid injected for the cure of hydrocele, within three days from the operation.



scuring the sight. He lost blood from the head repeatedly, and underwent a full course of mercury without benefit. A state of perfect amaurosis ensued.

The gout attacks the eye through the medium of the stomach. Vomiting occurs with pain in that organ, on the subsidence of an inflammation in the extremities, and is succeeded by violent pain in the head. The loss of sight is sudden and permanent.

From the  
font.

A gentleman, after an attack of gouty inflammation in the foot which suddenly ceased, was attacked with pain in the stomach and vomiting; this in the course of the day subsided, and in the night, a violent pain in the head was succeeded by an almost total loss of sight, which was never afterwards in any degree restored, although, by steeping the feet in a hot infusion of mustard, the great toe inflamed so much as entirely to relieve the head and stomach.

To this class belongs the cases of amaurosis consequent upon the sudden suspension of the catamenia, and of habitual hemorrhoidal discharges; the rapid healing of large ulcers of long standing, and the sudden retrocession of cutaneous eruptions.\*

\* See note D.

## FUNCTIONAL AMAUROSIS.

3. *Proper.*

A short case or two will best explain what I would call a temporary palsy of the retina from over excitement.

Amaurosis  
from over-  
excitement.

The following account is that of a young gentleman who was ardently engaged in the study of the profession when thus interrupted :

“ Having habituated myself for the preceding twelve months to intense study, reading and writing to a very late hour, which had been only interrupted for a few days by a slight inflammation of my right eye, I quitted London to recruit my health in the pure air of ———. This daily improved, but I found a growing imperfection in the vision of my left eye, which advanced, unaccompanied by inflammation, pain, or any other external symptom of disease. It seemed at first a film before the sight, but at length amounted to a total loss of vision. On examination, I found the pupil greatly dilated, and learned that the iris had little or no action. By the advice of Mr. T. whom I now consulted, I applied a blister, extending from the centre of the forehead round the eye to the root of the nose. This drew well, and I continued it open for ten days, closing the



eye from light during that period. I took at the same time a calomel and opium pill thrice a-day. In the space of a few days my mouth became sore; the pupil acted, though unequally, and I experienced a gradual recovery of vision. In the course of six weeks, I was enabled to resume my studies, and could perceive no defect of vision. I had gradually reduced the dose of calomel, and now discontinued it, drinking the decoction of sarsaparilla. At the distance of four months from this occurrence, the pupil is regular and active, and the sight unimpaired."

One of our most eminent and indefatigable artists in landscape was the subject of a superficial irritable ophthalmia, accompanied with much dimness and confusion of vision. This continued after the inflammation had subsided under the ordinary treatment. He became seriously alarmed to find, upon attempting to renew his occupations, that he was unable to discriminate the shades of color from each other, and that in fact he had lost the visual tact, if I might so express myself, essential to his pursuit. He submitted, by my advice, to a gentle course of mercury, and has since enjoyed his former accuracy of vision.

Examples of this species of amaurosis frequently occur among sea officers and others, suddenly, and without any preceding inflamma-

tion. I have repeatedly seen it of sudden accession with no other external sign than an inactive pupil.

A captain in the navy had made much use of his right eye for many years in observations with telescopes and sextants. About a week before he applied to me, he observed a mist before this eye, which increased until it was so dense, that he could neither distinguish the features of his friends, nor the large letters of a title page. The eye was free from inflammation, the pupil large and sluggish; he had no pain either in the eye or the head. He was bled copiously from the arm and temple, and briskly purged with calomel and jalap at short intervals. Blisters were applied to the temples. He then rubbed in a drachm of the strong mercurial ointment for several nights in succession; this produced a copious flow of saliva and violent diarrhœa, so that no benefit was obtained. By the calomel and opium pill taken night and morning his gums were immediately made sore. In three days the mist began to clear, and he was delighted to find that he could tell the hour by his watch. He continued improving so rapidly that, at the expiration of ten days, he could read an ordinary print with perfect facility. The pupil had recovered its natural magnitude and activity.

In one instance this form of the disease fol-



lowed a long exposure to the heat of the sun with such suddenness as to lead the patient to attribute it to a '*coup de soleil*;' and in another it was referred to the habit of reading by fire-light. It is seldom that both eyes are affected, and probable in some cases that the discovery of the amaurosis, and not the disease itself, is of recent occurrence. This point, however, can generally be ascertained.

---

Symptoms of  
amaurosis.

I now proceed to mention the leading symptoms of amaurosis. A great source of difficulty, in the arrangement of this extensive and complicated subject, is the circumstance of many symptoms being common to both classes. I shall appropriate them, as far as I am able, in my description. Pain affecting the forehead and temples is a precursory symptom of amaurosis, diminishing in proportion as the dimness increases; when the amaurosis is perfect, it usually ceases altogether if the disease has its seat in the eyeball. We must judge by its situation and extent, but especially by its association with other symptoms, if the pain be characteristic of organic amaurosis. Pain affecting the parts before mentioned, occasionally inconsiderable, and declining as the dimness increases, is common in some forms of functional amaurosis. If it be severe, remitting imperfectly, immediately increased by exercise, whether diffused

Pain.

over the entire side of the head, or circumscribed to a small space of the anterior cerebral lobes, it is usually connected with an organic cerebral change; but in this case, derangement and torpor of the *primæ viæ*, loss of strength and flesh, disposition to stupor, occasional confusion of intellect, inaptitude to exertion, and paralysis of one or more muscles, will be concomitant symptoms.

I have met with cases of amaurosis clearly depending on cerebral disease or irritation, in which the scalp was universally tender even to soreness.

There is an intermittent spasmodic pain accompanying some cases of amaurosis, shooting through the orbit into the head, of the most acute and distressing severity; it makes a periodic attack at or about the same hour, every night, or every second night, and continues for several hours; it is accompanied with convulsive quivering of the muscles of the eye and eyelids, and profuse lacrymation; there is nothing in the appearance of the organ to explain its nature and origin. What has been described as an intermittent ophthalmia, is I think improperly so termed—the pain, not the inflammation, is intermittent. The pain of tooth-ache and ear-ache, according to the state of the vascular system, is subject to intermissions more or less complete,

Spasmodic  
pain.



but the periodic pain to which I refer, is independent of any visible sign of inflammation. I believe it is a *tic douloureux* affecting one or more of the orbital branches of the fifth pair. I have cured it in two cases by arsenic, where opium failed to prevent the paroxysm.\* I have known one instance of a similar affection, without any defect of vision.

Paralysis.

Paralysis of the levator palpebræ is a sign of cerebral pressure, and always accompanied with some degree of imperfect vision; this, however, varies considerably. Paralysis of the orbicularis palpebrarum is less frequent. It is generally connected with a paralysis of the muscles on the same side of the face. A palsy of either of these muscles is sometimes attended by a degree of vertigo so considerable as to make the patient in danger of falling, if the eye be uncovered. In these cases near objects appear remote, and much diminished in size, as if seen through an inverted opera-glass. The vertigo seems to be excited by this illusion, as it happens to most persons in looking from such an eminence as renders objects dwarfish, for it ceases as soon as the affected eye is closed. This symptom is usually removed by depletion.

Vertigo.

\* Since the publication of the second edition of this work, I have seen and treated several cases of this description, some of them complicated with superficial inflammation, with the same remedy and equal success.

In other cases of fallen eyelid the affection of vision is so slight as scarcely to occasion inconvenience, so that a person may read or write while he supports the lid; but the pupil is invariably over dilated, and I have observed that the pupil does not recover its activity proportionably with the recovery of vision; on the other hand, I have seen both palpebræ affected with paralysis in succession, each for a day or thereabouts, and in this case the blindness supervened upon the removal of the ptosis.

Ptosis.

Ptosis follows injuries of the head and top of the spine: I have known a permanent ptosis and dilated pupil follow an injury of the cervical vertebræ after an interval of some months, without any other symptom of palsy. It is remarkable that there was no paralytic affection below the seat of injury. But paralysis from irritation may happen in any direction. In this case vision was in great measure restored by a course of mercury. The use of strong lead washes will produce a ptosis. I lately saw a temporary paralysis of the muscles on one side of the face, from frequently touching an ulcer of the fauces with the oxymel æruginis.

In proportion as vision fails, the eye affected with amaurosis loses its fellowship with the sound eye, and this loss of correspondence becomes a conspicuous character of the disease.

Loss of association and direction.



It is owing to this loss of correspondence that persons, affected with an imperfect amaurosis of one eye, often mistake the relative position and distance of objects, and frequently see them reflected.

In perfect amaurosis, or gutta serena, as it is absurdly called, the peculiar inexpressiveness or vacancy of countenance, depending on the non-convergence of the optic axes, is too remarkable to escape an ordinary observer, especially if both organs are affected. The patient has either a fixed unmeaning stare, or a constant rolling motion of his eyes. The loss of association in strabismus results from a relative debility of one of the straight muscles, and, if both eyes are employed, is generally accompanied with double vision; but that which I have just described is in no degree depending upon loss of muscular energy, partial or general, direct or indirect; but on the total failure of sensation, by which the actions of all muscles subject to the will are directed.

Strabismus.

Strabismus is either congenital, or from the debility of scrofula, like the paralysis of the lower limbs in children; or from a morbid association accidentally contracted and impressed by habit in childhood; or from a wound of the frontal nerve; or a speck upon the cornea rendering the vision oblique; or from violence

done to the affected straight muscle. I have seen a complete internal strabismus, the effect of a blow on the temple, which a school-boy received in fighting. From its suddenness and the ecchymosis of the conjunctiva on the temporal side, I concluded the abductor was lacerated or palsied ; and so it proved ; for in a few weeks the distortion was entirely removed.\* It is also a symptom of irritation arising from difficult dentition, worms, &c.; or of pressure at the origin, or in the course of the nerve proper to the affected muscle.†

\* Mr. Cheselden relates the case of a gentleman who had strabismus, with double vision, produced by a blow on the head. By degrees, the most familiar objects came to appear single again, and in time all objects did so, without any amendment of the distortion.

† A remarkable case of strabismus, with double vision, occurred in a patient who consulted me, about twelve months since.

J. G. was attacked, six years ago, with a remittent fever of extremely tedious and obstinate character, leaving him in a state of debility from which he very slowly recovered. During his convalescence, the right eye began to be affected with strabismus and double vision, which increased, until it was impossible for him to look with both eyes without producing pain in the head and confusion of thought. It distressed the patient exceedingly, and kept him in a nervous, irritable condition, which impaired his health, and rendered life a burden to him. Four years after the accession of the disease, he applied at the New-York Eye Infirmary, with a request that the affected eye might be destroyed, in order to



There is a complaint of cross sight occasionally made by persons who have no perceptible strabismus. It happens in looking downwards, as in ordinary reading. In looking at objects

put an end to his distress. It was then ascertained, that he saw perfectly well with either eye singly, when the other was covered, but the double vision was complete when both were used.

As his request could not properly be complied with, he would not consent to any other plan of treatment, and successively applied to different surgeons in the city, who uniformly refused to perform the operation for which he was so anxious. Eventually, however, he found an empiric who undertook to sink the eye. He made an incision through the sclerotica, posterior to the cornea, at its lower edge, and thrusting in a probe, succeeded in removing the lens, but did not reduce materially the size of the organ. Severe inflammation followed the operation, which soon also attacked the internal textures of the other eye. After three weeks had elapsed, I was called and found him in a most deplorable situation; the one eye ruined, the other excessively inflamed, the iris adherent, and the pupil filled with lymph; the patient suffering continually intolerable pain. The most active treatment was pursued to subdue the inflammation, but only with partial success. After six weeks of acute suffering, the inflammation began slowly to subside, and the sight gradually to return in the left eye. The patient then went into the country with the hope of advantage from change of air. But here again, his constitutional irritability getting the better of his judgment, he suddenly set off for another of our principal cities. There, before the inflammation was completely reduced, several operations were performed to clear the pupil and remove the lens. The result was, destruction to the patient's sight, and an eye gradually falling into a state of atrophy.

on the same horizontal plane with the eyes, it is not perceived. It is a relaxation of one of the inferior straight muscles.

Although incidental instruction may be gained from the bad practice pursued in this case, the point intended to be illustrated is, the duration of the double vision, as accompanying the strabismus. No change took place in this respect during five years. In general, it is well known that double vision gradually subsides, and the squinting eye becomes accommodated to the new direction of its axis. The degree of distress which the patient suffered was also sufficiently remarkable, when it could induce him to desire the destruction of one of his eyes. Covering the affected eye gave him but partial relief, as the slightest ray of light coming to it brought on the confusion and pain under which he continually suffered.

Another case of double vision has since occurred to me, not unworthy of record. A man received a violent kick upon the cheek, just below the right eye, from a maniac, whom he was endeavouring to secure. The immediate consequence was double vision, but in a vertical instead of a horizontal direction. Every object at which the patient looked appeared double, but the image of one *above* the other. On examination, a firm swelling, caused by extravasated blood, was felt below the eye, deep in the orbit, and it was evident that the organ was somewhat pushed up out of its natural direction. On placing a finger beneath the other eye and pressing into the orbit, the double vision was immediately corrected, by the axis of the sound eye being made to correspond with that of the injured one ; while the symptom returned as soon as the finger was removed. The extravasated blood was gradually absorbed, and the patient after a few weeks regained correct vision.—*Editor.*



Hemiopsia  
or partial  
impairment  
of the retina.

The retina is often partially affected in organic amaurosis. Thus some persons describe a horizontal, others a vertical screen, eclipsing one half of the object viewed, or even of the field of vision. In order to see a given object entire, which is upon a level with the eye, they are compelled to move the eye or the head, in the direction which the obliquity of their vision requires. Again, cases now and then occur in which persons have lost their lateral vision, while they see objects in the direct line of the axis of vision. In the greater number of cases however, the vision of amaurotics is oblique, and in perfect amaurosis, the perception of light, if any, is also oblique. The gradual return of vision is generally first perceived in this direction. The sensibility of the retina, whether more or less, seems to be greatest at the part farthest removed from the axis of vision, or rather from the entrance of the nerve, for it is more frequently observed on the temporal than on the nasal side. I am unable to offer a satisfactory conjecture in explanation of this fact.

A gentleman, who died of apoplexy in his 36th year, and who lost eight pounds of blood in the three days preceding his death, had consulted me for depraved sight of one eye, six months before that event. The pupil was permanently dilated. He had double vision whilst looking

directly forward ; if he looked obliquely to either side, his vision was single. A slight strabismus was perceptible. Many cases analogous to this have fallen under my notice, in which no opportunity was afforded of ascertaining the cause of disease.\* The vision is sometimes perfect or nearly so with the eye, which, by loss of correspondence with its fellow, occasions the duplicity of objects viewed with both. I have known it equal in power to that of the other eye when employed singly. The double vision and giddiness cease, when either eye is employed alone, or is closed. The paralysis affects the nervi motores oculorum directly in such cases, and the optic nerve suffers by sympathy. Where the vision of the affected eye is materially bedimmed, the ptosis or strabismus is more marked, and the symptoms of double vision and vertigo in proportion.

A distorted relative position of objects is also not an uncommon symptom of organic amaurosis. Thus, the lines of a printed page seem zig-zag, and the two eyes of a face appear in different planes, whether one or both eyes are affected. If one object be seen in its proper place, the situation of a contiguous object is erroneous, both as to distance and parallel. If only one eye be affected, the employment of the sound eye exclusively, corrects the error. In the case

Distorted  
position of  
objects.

\* See note E.



of simple non-correspondence, it is corrected by either eye indifferently. This is the reverse of what happens when the eyes are perfect, in which case, the delusions arising from the use of one eye only, are corrected by employing both. Both this and the preceding are for the most part, not always, symptoms of an organic affection. I have been led to this conclusion from observing the morbid appearances, and the inefficacy of remedies, where such signs of the disease were established. But I have known instances of their removal.

Muscae,  
fixed and  
floating.

Muscae are either fixed, when they are usually allied to spectra, and belong to organic amaurosis ; or floating, when they are, I believe, characteristic of functional amaurosis, sympathetic or proper. I have known the fixed musca, permanent, without variation, for years, darkening a certain defined portion of the field of vision. In some rare instances, it precedes acute choroid inflammation ; more frequently it is unconnected with any particular morbid state. Around the opaque spot persons have sufficiently distinct vision. The spot varies in density in different individuals, and under a long but gentle mercurial course, I have known it become considerably less dense, so as not to intercept bright light. Its circumference sometimes reflects a tinge of color or a luminous halo. The fixed musca seldom presents the fantastic

shapes of the floating,\* but it is not uncommon for the two forms to co-exist.

The *musca volitans* is sometimes solitary, following the eye at a fixed angle as it passes along a line; sometimes two, three, or more are presented; more frequently an immense assemblage, descending in a cloud as the eye is raised, and ascending as it is depressed. They are obvious to so many analogies, and apprehension of impending blindness makes patients so minute in their observation and description of them, that it is scarcely possible to do justice to our experience in attempting to describe them. Sometimes they are represented as globular, sometimes annular and flat like a piece of money. Portions of flue, of soot, insects' wings, transparent vesicles, or minute globules of quicksilver, connected like the links of a chain, or short hairs with their bulbs attached to them, are ordinary resemblances. They occupy the air with some persons, and are seen upon looking at the sky, or upon a white sheet of paper, and especially in shifting the eye from one object to another; to some they appear in the fire or candle only, and with others they seem to cover the ground, so that they walk in them knee-

\* To this remark there are exceptions. A gentleman, who a short time ago consulted me, compared the spectrum constantly before each eye to a large dragon fly, darkening the field of sight.



deep. Almost every person has, at some time or other, seen these appearances, but especially those subject to dyspepsia, and disordered function of the stomach and liver. At the moment of approaching deliquium, they appear in one vast cloud, and they are harbingers of the intense bilious headache. At the instant of their appearance, the sentient extremities upon the fingers and tongue are so benumbed, that objects of touch and taste convey a very indistinct impression, as if some muffle were interposed. These sensations I am describing *ad vivum*, for I was formerly often the subject of this attack, which was followed by a certain degree of confusion of intellect, and temporary suspension of memory, so as greatly to embarrass, if not to take away, the power of intelligible expression. I mention these opposite and transitory states of emptiness and plethora concomitant with the floating *muscæ*, to show the purely functional origin of the affection. The one (deliquium) is an uninjected, the other (sick headache) an over-injected or congested state of the nervous texture; or suspension from vacuity, and suspension from plethora. An analogy is plainly to be perceived between the corresponding states of the sentient and visual extremities, described in the last affection, to that of a temporary incomplete paralysis.

The fixed *musca* is generally an organic affec-

tion, probably a deposit or extravasation between the choroid and retina, compressing to a certain space the papillæ of the retina, to which the musca corresponds in figure. In other instances, it is independent of deranged structure, and may be presumed to be only an insensible point on the retina. The single muscæ sometimes coalesce, and form a larger spot. The floating muscæ are altogether a functional affection, not interfering with useful vision, and sometimes, though not often, removed. To some persons they disappear upon looking through glasses, and others see them only upon remote objects. Their magnitude diminishes as the distance is increased, to those who see them at all distances. They are simply a disordered circulation in the vessels of the retina,\* and occur oftener in nervous and spare than in plethoric and robust persons; they are generally induced by overstraining the organ, almost always accompanied with a weak and irregular digestion, and varying with, if not depending upon, the condition of the stomach.

Colored spectra, or luminous impressions of objects remaining upon the retina, are usually preceded by the fixed muscæ, and may then be regarded as a more advanced stage of the com-

Morbid  
spectra.

\* In some cases they have been supposed to depend on floating particles in the humors or minute scabrous points in the cornea.



plaint. This is not always the case: I have known them to be symptoms of functional derangement, and to disappear as the vision recovered. In this affection, a halo of light encircles the opaque spot during the exercise of vision; and if a bright luminous object be contemplated, a colored image or the reflection of it is presented to the mind for a time, greater or less, after the eye is withdrawn or closed. Of the same species are the various morbid refractions of luminous bodies, presenting the object double to one eye, or curiously divided and distorted, as in looking through a crinkled pane of glass;\* and the appearance of prismatic colors in the forms of circles, rainbows, cones, &c. Sometimes ordinary objects are imperfectly represented, or even dark, so that they would not be known but from their outline being illuminated. Thus a man, a tree, or a house, appears fringed with a glory; and on the other hand, it is not uncommon for the outline of objects to be lost in shade, while the centre is clearly discerned.

I might illustrate these observations by a multitude of cases in my possession, which I omit for the sake of conciseness, and in conformity to the plan of this work. On this subject, however, I shall take the liberty of introducing the

\* I have known cases in which interrupted vision was produced by several apertures made with a needle in a capsular cataract, so that the moon appeared as if cut in pieces.

case of an intelligent young gentleman, very accurately drawn up by himself, for the purpose of showing the gradations of this disease in its progress, as well as its origin.

“About a year and a half ago the first symptoms appeared, which gave me any uneasiness with respect to my sight. For several months I read incessantly, not only throughout the day, but also for five or six hours each night by candle-light; and I now perceived numerous circular motes, which combining, formed clouds of irregular figures before my eyes. These motes always appear when I look at the sky or any light-colored object in a strong light; they move with the eye, retaining for some time the same position with relation to each other and to the centre of vision: each consists of a slightly opaque circumference and a central spot, the diameter being, as well as I can judge, about four or five minutes of the circle of vision. Sometimes films appear curved or twisted like hairs, and of the same degree of opacity as the motes. There is a collection of these films always before the right eye, but at such a distance from the centre of vision as not to disturb sight. The number of the motes seems increased by violent exercise as well as by close reading, or a disordered state of stomach. Sometimes for a moment a small circular black spot appears near

Case of  
muscae and  
spectra.



the centre of vision, and sometimes, though not so frequently one faintly luminous.

“The candle next appeared surrounded with a faint halo, which became more vivid as I continued this severe exertion of my sight. When my eyes are unusually weak, or a light is presented to them after I have been some time in darkness, instead of the halo a globular appearance of a muddy yellow color surrounds the flame.

“About six months ago, I began to be annoyed by the retina retaining impressions made upon it. After looking at any white or bright metallic object, on turning away my eyes I distinctly perceive its outline in a darker shade, on any surface to which I may direct my view; the impression lasting from two or three seconds to half a minute, according to the strength of light, the brightness of the object, and the length of time for which I have viewed it. The flame of a candle leaves its image impressed on the retina frequently for a couple of minutes; the sun for a still longer time; the image in both instances being of a muddy yellow color.

“A kind of penumbra surrounds light-colored objects in a strong light, and prevents me from accurately distinguishing their outline. When the object is under a sufficiently small angle to

be seen entire without moving the eye, it seems double, one image being such as would appear to a healthy eye, the other much fainter ; thus is the moon seen, a piece of money, or the gilt letters over shop windows. These appearances take place indifferently, whether I use either eye or both.

“In a few instances, a very severe exertion of my eyes, produced the appearance of innumerable black particles dancing before them.

“When I read for any considerable time, I have a disagreeable sense of heat in my eyes, with pain in the eyeballs, extending to the lower part of the forehead. I am not constantly subject to headaches, though occasionally afflicted by them, especially if I delay breakfasting for any length of time after rising. My tongue is frequently foul for weeks together, my digestion seems weak, and I seldom enjoy a good appetite.

“I ought to observe, that most of the above-mentioned symptoms seem to have been mitigated since I came to London. Since the application of the blisters, the halo round the flame of the candle has nearly disappeared.”

A very frequent and characteristic symptom of functional amaurosis is a thin mist, fog, smoke, or gauze, or, as I have heard some pa-

Amblyopia: nebulous or misty vision.



tients represent it, an indefinable something, as if vision required a peculiar atmosphere, intervening between the eye and the object, which takes off the '*acies oculorum acer claraque*,' the sharp edge of clear vision. Letters of a book run together, and the outline of all minute objects is indistinct. In some cases this indistinctness is constant and unvarying; in others it is the result of exercise of the organ, for a period, varying in different persons from ten minutes, or even less, to half an hour. Repose of the organ, whether obtained by closing the lids, or looking vacantly on distant objects, or gentle friction of the lids, or a slightly stimulant application, enables the person to resume for a short time; but the hindrance returns, and if the employment be persisted in, the dimness becomes little short of blindness, and sometimes occasions pain, always a painful sense of weakness. This affection is unaccompanied with irritable conjunctiva; there is no tendency to suffusion. It is seldom relieved by glasses, and never permanently. It is sometimes combined with *muscæ*, but more frequently distinct. The iris appears irritable and unsteady; it contracts often quickly, but vacillates between contraction and dilatation without a change of the light.

Oscillatory  
vision.

Another functional affection is an oscillation or wavering of objects, so that the want of steadiness occasions a dazzling and confused per-

ception. This may be the result of simple congestion; but I have known it unaccompanied by any sign of this state, and in persons of a frame and temperament distinctly opposed to it. With this is often combined a delusion of something waving or flapping in an oblique relation to the eye, as towards the temple, or pendulous from the eyelash, or brow, unaccompanied by any distinct perception of figure.

An occasional symptom of functional amaurosis is a loss of the faculty of distinct vision at different distances. A gradual abridgment of the focal range at its near extremity, occurring in advanced life, and requiring the aid of convex glasses, has been supposed to depend upon a permanent change in the figure of the globe; I should rather refer it to a loss of power of the retina incidental to age, and a consequent imperfection of function in those parts, which execute the office of adjustment. It is a change similar to this, taking place in early or middle life, and with more abruptness (the indistinctness sometimes pervading near objects exclusively, and in other cases, remote ones), which I consider to be a symptom of amaurosis. It cannot be admitted, that the distinct vision of an object, at a permanent focal distance, proves the power of the retina to be unimpaired. If the organ be originally perfect as an optical instrument, so that the rays of light flowing from near, as well as from remote objects, form images upon

Loss of the  
adjusting  
power.



the retina sufficiently perfect for distinct vision; and if the range of distinct vision be, whether suddenly or gradually, so abridged, that the eye is incapable of relieving itself by a change of focus, the feebleness of the retina is invariably demonstrated by other signs, for the imperfection of adjustment seems in all cases to be in proportion to the loss of vigor of the retina. Ordinary observation proves that the effect of wear and tear is to allow of good distant vision, in which the parallelism of the rays of light supersedes the necessity of adjustment, while the near sight, which requires the active or tonic state of the adjusting faculty, is impaired or lost. But if, as sometimes happens, the vision of near objects remains good, while the distant is obscured, the evidence of the faultiness of the retina is direct. The correction of a defective adjustment by the use of glasses, in either case, proves no more than that the retina is not organically affected, while the failure of this corrective, which is frequent in the cases referred to, demonstrates the functional debility of the retina. In most of these cases the use of glasses is of temporary benefit, but if continued, it is followed by uneasiness or pain in the eyeball.

Many phenomena of impaired adjustment correspond to the degree of mobility of the iris; for in some persons it is quick to contract, but unable to preserve its contraction, and falls open or fluctuates in the same quantity of light, and

I have observed that the point of clear vision shifts accordingly; in others it contracts slowly and imperfectly; in others again it is permanently contracted or dilated, and this, independent of any other defect of sight than an abridgment of the original range of distinct vision, at one or other extremity.\*

I have met with different reports, as to the time of day in which persons affected with imperfect amaurosis, enjoy the best sight. Some see clearest on first waking in the morning, whereas others are particularly dim for an hour or two after rising. In these cases, the state of the stomach has an obvious influence. Emptiness will produce muscæ, and a temporary blindness. Some see only in a full light, others in a weak one, as after sunset. Candle-light, though generally least favorable to such persons, is not invariably so. In a considerable proportion of cases, amaurotic patients see clearest in the evening, and their vision seems to gain strength by exercise. They see better for example on retiring to rest, than they have seen at any period of the day.

Influence of  
light on  
amaurotics,

These differences are, in most cases, referrible to the varieties in susceptibility of the retina,

according to  
the state of  
the retina,

\* See note F.



determining the requisite degree of illumination of objects for vision, and the adaptation of the pupil to that purpose. But in other instances, as I have before hinted, a reflection takes place within the eye, owing to some change in the quality or quantity of the pigment, which renders a screen or colored glasses, or a dark day or twilight, in their several degrees, favorable or even essential to vision. This partakes of the nature of an organic disease.

and of the  
pupil.

I have known persons absolutely blind for two or three minutes, upon going into a darkened room, owing to the imperfect sensibility of the retina, and consequent slow dilatation of the pupil; and they made no complaint, as persons usually do, of dimness from the opposite transition. Such persons, indeed, require a full strong light to see at all. But that such a defect is not directly or exclusively depending on the state of the pupil, is proved by the occasional coincidence of good vision with a permanently contracted pupil. I have met with cases of remarkably small and habitually contracted pupil, in which the glare of light was even painful, and where the inconvenience was at once removed by moderating the light, without any perceptible alteration in the diameter of the pupil. A lady of quality, in whom I made this observation, told me that it had equally attracted the notice of Mr. Hunter, whom she had formerly consulted. The

cases of day and night blindness present the opposite extremes of variation in susceptibility of the retina, and these must be regarded as cases of proper functional amaurosis. The remarkable efficacy of blisters upon the temples in these cases confirms this fact.

I had abundant opportunities formerly of observing the influence of trades, in aggravating as well as producing amaurotic affections: it was a common remark with tailors and shoemakers, that they never saw so well as upon Monday morning, which they justly attributed to the repose of the organ during Sunday.

The activity of the iris, evinced in the motions of the pupil, is, generally speaking, the surest indication of the health of the retina. The contraction is slow, or it is incomplete, or both, upon the sudden admission of light to the eye, where the retina is defective in sensibility. The mean state of the pupil is that of dilatation where an amaurotic affection exists, and this will sometimes discover to an attentive observer, which of the two eyes is affected. When the perception of light fails altogether, the pupil is generally fully dilated, and absolutely motionless. It is in other cases not perfectly a circle.

Action of the iris an index of the state of the retina.

The activity of the iris requires the free and uncompressed state of the retina, iris, and ci-

Phenomenon of active irids in per-



fect amaurosis.

liary nerves. In the various forms of amaurosis, its activity is proportioned to the degree of integrity which these several parts retain, and the intensity of the stimulus. If the retina be opaque, compressed, or unsupported, the iris mechanically disordered, or the ciliary nerves palsied, the pupil is inactive, independently of the state of vision. In the first of these cases, it is evident the vision will be lost; but we continually see useful vision combined with the second and third, as after operations in which the iris has been half destroyed, or has become permanently adherent, or in malformations where it is half wanting; and in paralysis of the ciliary nerves accompanying the state of ptosis. But how shall we explain the activity of the iris in a state of absolute blindness? a case by no means uncommon. We can only explain it by concluding the organ to be sound, and the cause of the amaurosis remote, or at least external to it. Its motions in such a case are purely involuntary; the mental perception being suspended or annihilated. All that is required to excite them is the impingement of the ordinary stimulus upon the unchanged retina, the white sheet upon which the images of objects are impressed, the instrument, not the organ of perception. The iris, in such a case, acts by a sympathy independent of the brain.

Thus in a case of circumscribed tumor com-

pressing the left optic nerve, immediately behind the ganglion opticum, although the blindness was complete, the iris was active. In two young ladies, in whom the eyes, as in the former case, were perfect, and the blindness complete, the iris was even vivacious; and there was the strongest presumptive evidence, from the symptoms, that the amaurosis was in the cerebral portion of the nerve.\*

If this theory be correct, the activity of the pupil in complete blindness proves, that the retina and orbitar portion of the optic nerve are unaltered, and that the disease has its seat in the cerebrum, or cerebral portion of the nerve; while, at the same time, the ciliary nerves are unaffected by it in their origin and course. On the contrary, the fully dilated and motionless pupil shows, that these nerves are paralysed—the disease may be cerebral or orbital, or both. We see this exemplified in hydrocephalus, and in orbitar tumors compressing the nerve and globe. In cases of perfect amaurosis, in which the pupil, of its ordinary size, is absolutely without motion, a case by no means uncommon, the retina has most probably undergone a change of texture. The ciliary nerves are uncompressed, as may be inferred from the undilated state of the pupil, but the source of their

\* Janin relates two cases of lively pupil in a state of total blindness.—*Mem. et Observ. sur l'Œil*, p. 426.



excitement, sympathy with the retina, is destroyed. The symptoms of the disease, in its early stage, will point out its seat, where its locality is definable.\*

Blindness  
with dilated  
pupil, from  
fractured  
basis.

A man was trepanned for a fracture with depression of the right frontal and parietal bones. After the operation he became sensible, but it was discovered that he was totally blind, being unable to perceive a lighted candle held close before his eyes. The pupils were fully dilated and insensible to light. On the fifth day he died of inflammation of the membranes of the brain, having continued until his death in total darkness. On dissection, a fracture of the frontal and parietal bones was discovered on the left side, corresponding to that on the right, but without depression; and these two fractures were connected by a transverse fracture, extending across the basis of the cranium, i. e. through the orbital plates of the frontal bone anterior to the junction of the optic nerves. This transverse fracture extended into the bones of the face, so as to separate them from those of the cranium, and there was displacement of the bones sufficient to occasion a considerable pressure upon the optic nerves.

\* It is to be regretted that in the many excellent observations of Morgagni and others, of diseased states of the optic nerve, the opportunity of learning the signs of the disease during life was so seldom enjoyed.

In this interesting observation, given to me by my able and excellent friend Mr. Brodie, it is evident that all the nerves entering the orbit must have been compressed. Hence the paralysis of the iris concomitant with that of the retina.

In hydrocephalus, the pupils are invariably fully dilated and motionless. In apoplexy generally, but with exceptions; sometimes contracted, but still immoveable; which Dr. Cooke, in his late Treatise on that subject, considers a fatal prognostic. In injuries of the head, with symptoms of depression, the pupils are generally dilated and motionless, but sometimes considerably enlarged, and slightly moveable; in other instances contracted; and very frequently one is permanently dilated, and the other contracted. These varieties depend on the situation in which pressure is applied, and the extent which it occupies, and in some degree on the nature of the compressing cause, whether bone driven in or fluid effused; in hydrocephalus the result is uniform. But it is not the loss of the sight which occasions the dilatation of the pupil, in injuries where these circumstances coincide, but compression of the ciliary nerves, or of those from which they are derived; the loss of motion in the iris is a gradual and not a sudden effect of the loss of sight, and it seldom happens that the vision is lost in those casualties in which the pupil is permanently dilated.



Among the morbid changes of the retina, ossification has been rarely met with.\*

It is remarkable that in the thickened, attenuated, softened, ossified, or otherwise morbid states of the optic nerve or its sheath, the diseased appearance has seldom extended beyond the ganglion opticum. The eyeball has frequently been free from disease. The blindness has probably in as many instances proved a cause, as an effect, of the degenerations of the nerve. Cases indeed are related, of a considerable and very obvious change in the structure of the nerve, where the sight of the corresponding eye has been unaffected.

Case of  
Milton.

It has occurred to me, in concluding the subject of amaurosis, that the case of our great Epic Poet, drawn up by himself for the purpose of its being submitted to Thevenot, a celebrated French oculist, may not be uninteresting to my readers. I subjoin it as the best account that I know of the symptoms of amaurosis, in its progress from the state of functional debility, to the confirmed, perhaps organic, gutta serena. I have preserved his own words for the sake of accuracy.

\* Vide Morgagni, Lett. 52, Art. 30.

“Decennium, opinor, plus minus est, ex quo debilitari atque hebescere visum sensi, eodemque tempore lienem, visceraque omnia gravari, flatibusque vexari; et mane quidem, si quid pro more legere cœpissem, oculi statim penitus dolere, lectionemque refugere, post mediocrem deinde corporis exercitationem recreari: quam aspexissem lucernam, Iris quædam visa est redimere: haud ita multo post sinistrâ in parte oculi sinistri (is enim oculus aliquot annis prius altera nubilavit) caligo oborta, quæ ad latus illud sita erant, omnia eripiebat. Anteriora quoque, si dexterum forte oculum clausissem, minora visa sunt. Deficiente per hoc fere triennium sensim atque paulatim altero quoque lumine, aliquot ante mensibus quam visus omnis aboleretur, quæ immotus ipse cernerem, visa sunt omnia nunc dextrorsum, nunc sinistrorsum natare; frontem totam atque tempora inveterati quidem vapores videntur insedissee; qui somnolentâ quâdam gravitate oculos, a cibo præsertim usque ad vesperam, plerumque urgent atque deprimunt; ut mihi haud raro veniat in mentem Salmydessii vatis Phinei in Argonauticis:

———— κάρος δέ μιν ἀμφεκάλυψεν  
Πορφύρεος, γαῖαν δὲ πῆριξ ἐδόκησε φέρεσθαι  
Νειόθεν, ἀβληχρῶ δ' ἐπὶ κώματι κέκλυτ' ἀναυδός.\*

\* ————Vertigo vero ipsum circumdedit

Atra, et terram opinatus est circumagi

Ab imo, in languidum vero soporem delapsus est elinguis.

BECK'S APPOLLONIUS RHODIUS. Lib. 2, v. 203.



Sed neque illud omiserim, dum adhuc visûs aliquantulum supererat, ut primum in lecto decubuissem, meque in alterutrum latus reclinassem, consuevisse copiosum lumen clausis oculis emicare; deinde, imminuto indies visu, colores perinde obscuriores cum impetu et fragore quodam intimo exilire; nunc autem, quasi extincto lucido, merus nigror, aut cineraceo distinctus, et quasi intextus solet se affundere: caligo tamen quæ perpetuo observatur, tam noctu, quam interdiu, albenti semper quam nigricanti propior videtur; et volvente se oculo aliquantulum lucis quasi per rimulam admittit."

LEONARDO PHILARÆ, ATHENIENSI:

Septemb. 28, 1654.

*Miltoni Opera Amstelodami, 1698. p. 330.*

# PATHOLOGY

OF THE

# HUMORS.

---

## CHAPTER II.

### SECTION I.

#### AQUEOUS HUMOR.

THE simple redundancy of the aqueous humor is a sequel of chronic inflammation, affecting the internal texture of the globe. Its figure is preserved, but the distended sclerotic has a dark blue tinge; the cornea is extended and prominent, the pupil dilated and inactive, and the vision is inconsiderable, if not extinct. In other instances, the state of hydrophthalmia is accompanied with loss of figure of the globe, and staphylomatous enlargement of the cornea, which is specked or exulcerated, and frequently presents fasciculi of red vessels on its surface. This state is the result of a disorganizing inflammation.

Hydroph-  
thalmia.



The bulged and transparent cornea, whether spheroidal or conical, gives the appearance or the idea of a redundant aqueous humôr; but this is only the consequence of the increased capacity of the chamber. The distinction is important; for the treatment of the hydrophthalia and the conical cornea proceeds, as it seems to me, on opposite principles.

Rapid reproduction.

The aqueous humor is always rendered turbid by inflammation of the choroid and iris, but resumes its transparency when the inflammation is subdued. When discharged by accident or operation, it is reproduced in a period of from eight to twelve hours. It is regenerated in all states of the organ, in which the anterior chamber is even in part preserved, in quantity sufficient to give plumpness and figure to the globe, and to refract the light with accuracy enough for the distinct vision of large objects.

Effusions of lymph, from inflammation of the iris, of puriform matter from internal ulcer of the cornea and abscess of the eyeball, and of blood from concussions and wounds of the organ, are frequently observed in the chamber of the aqueous humor.

Solvent power inconsiderable.

The solvent action of the aqueous humor over the exposed fragments of the crystalline lens is not, in my belief, superior to that of water,

#### AQUEOUS HUMOR.

which I have found by experiment to be exceedingly slow. A knife or needle, too highly tempered, has occasionally been broken in the operation for cataract, and the point has been left in the anterior chamber. This accident once happened in my own hands. The rusted appearance of the aqueous humor, and the gradual disappearance of the fragment, have led to a conclusion that it underwent a chemical solution. The rapid removal of the fluid and flocculent cataract, when dissipated in the chamber, has been explained in the same way. I believe that the aqueous humor has no greater solvent property than common water, and that this would be quite insufficient to explain the very quick restoration of its clearness, which we often witness, where it has been loaded with opaque matter. The fragments of the lens have no more power of resisting absorption than an extraneous substance, and the process improperly termed solution, is essentially referrible to the operation of the absorbents. The secreting function of the chamber is evidently a powerful one, from the reproduction of the humor in the course of a single night. That the absorbent function is nearly equal to it, is proved by the facts above mentioned, but still more strikingly, by the rapid diminution and removal of the matter effused under inflammation, when quickened by the excitement of mercury.



## SECTION II.

---

### VITREOUS HUMOR.

THE absorption of the vitreous humor is evident in cases of floating cataract, and in some forms of organic amaurosis, marked by preternatural flaccidity, even without a diminution of volume; also in cases of absorbed crystalline and membranous cataract, with adhesions to the iris.

Diseased  
eyes of  
horses.

I examined the decayed eyes of horses, and found in a considerable number the opaque lens sunk in the vitreous chamber, and this humor almost entirely absorbed. The eye was filled by a morbid accumulation of aqueous humor, so that on a single puncture the whole fluid contents of the globe, which was larger and more tense than usual, escaped uninterruptedly. The sclerotic and choroid in these cases, although often thickened, and even the opposite sides co-adhering, were entire. The retina was usually drawn into folds, and partially absorbed. In some cases the globe was greatly enlarged and flaccid, resembling an undistended dropsical cyst; the crystalline opaque, of its natural size

and firm, and sometimes its capsule thickened and scabrous, bearing marks of inflammation, but in place of the healthy vitreous humor, a gelatinous fluid, of a deep yellow or amber color, filled the globe. In other cases, where the tunics were disorganized, and thickened from injuries, the space occupied by the humors was either exceedingly reduced or obliterated altogether, by the co-adhesion of the opposite sides of the globe. In most cases where the lens was cataractous, and not impacted in the anterior chamber, nor fastened by adhesions of its capsule to the iris, the substance of the vitreous humor was shrunk as before described; and the crystalline, partially absorbed, had receded and sunk in proportion. The globe was supported in its figure by a morbid collection of aqueous humor.

The tremulous iris is, I believe, always connected with a relative disproportion in volume of the vitreous humor, whether congenital or the result of operations and injuries. Couching and the operation by absorption, if roughly performed, break down a portion of the vitreous cells, which become obliterated; hence the frequency of floating cataract and tremulous iris after these operations. The loss of a very considerable proportion of the vitreous humor may take place without permanently impairing the vision, except of minute objects, as is proved

Tremulous  
iris.



by the successful issue of some cases of extraction, in which this accident has happened.\*

\* Disorganization of the vitreous humor, with tremulous iris, although coming on spontaneously, is not necessarily productive of blindness. A young lady, from Vermont, arrived in this city in the early part of last summer, to obtain advice in such a case. The iris was tremulous, the eye-ball soft and readily yielding to pressure, and the lens, having lost its support in the vitreous humor, had fallen through the pupil into the anterior chamber. No severe inflammation accompanied this unnatural situation of the lens, but it was productive of occasional pain, and almost completely destroyed vision by lying in front of the pupil. The lens was still transparent, except at its edges, and although it had remained in this situation several months, was very little diminished by absorption. The other eye having been lost in childhood, from accident, the patient was thus deprived of all useful vision. A section was made in the cornea, through which it was intended to extract the dislocated lens, but immediately upon withdrawing the knife, it passed back through the pupil, and fell below the axis of vision into the bottom of the eye. No inflammation followed the operation, and the patient recovered in a few days. The sight was rendered so perfect, that, with a proper glass, she could read and sew as well as any patient who had undergone the operation for cataract.

In a similar case the result was different.

W. D. — a shoemaker of this city, wounded his left eye with an awl, about sixteen years before he applied at the Infirmary. The instrument pierced the cornea and lens, which in consequence became opaque, and the eye lost its power of vision. It gave him little or no inconvenience, until the day before he came to the Infirmary, when on rising in the morning, he found the eye beginning to be inflamed, with gradually increasing pain, and the lens lying in the anterior chamber. On examining it, in addition to these symptoms, the iris was

I have suspected a diseased state of the vitreous humor in some cases of cataract in elderly people, accompanied by a preternatural convexity of the globe, in which a slightly glairy fluid has distilled in quantity from the eye during the operation, of a consistency between that of the vitreous and aqueous humors.

A diseased state of this humor, frequent in organic amaurosis, with or without cataract, and especially accompanying diseased changes of the

Discolored  
vitreous hu-  
mor.

observed to be tremulous, and the whole organ preternaturally soft. Sight was entirely extinct; the eye not being sensible in the slightest degree to light. It was evident here, that in consequence of the accident, a process of disorganization had been set up in the vitreous humor, which eventually depriving the lens of its support, had allowed it to escape into the anterior chamber. To prevent the dangerous inflammation likely to follow this state of the parts, the lens was extracted through a section of the cornea. The wound healed well, and the eye regained its healthy appearance, but of course, remained perfectly blind.

Our author remarks, that a tremulous state of the iris may follow the rough performance of the operations of couching, and that by absorption. It is equally true that it may sometimes follow them when performed in the most skilful manner, and be productive of no ill consequences to the patient. In such cases the tremulous state of the iris is often not permanent, but subsides after the lapse of a few weeks; rather disproving our author's position that it is "*always* connected with a relative disproportion of the vitreous humor," for such a condition of the parts must be unchangeable, and its effect in producing tremor of the iris permanent.—*Editor.*



iris, is that in which it assumes a deep yellow or a chocolate brown color. From its rapid and uninterrupted egress, in this and the former case, even to a partial collapse of the globe, there is reason to infer that the cellular contexture is broken down; for although the cells of the healthy humor communicate, a gush only of the fluid ensues from wound or rupture of the capsule; the support afforded by the closed lids prevents its further escape, and the obliteration, by inflammation of the ruptured cells, speedily follows.

Hemorrhage  
into the vi-  
treous cells.

I have known blood effused into the cells of the vitreous humor within twelve hours after the operation of extraction, in consequence of straining upon the night-chair, which was instantly followed by severe pain darting towards the occiput. The coagulum was visible both to the patient and to the surgeon; the former described it as a central circular spot, intercepting the light which was strongly reflected from its circumference; in the sunshine it had a bright scarlet hue, and was liver-colored in the shade; it was in the course of time absorbed, so that the patient gradually recovered tolerable vision.

Another case was one of active and continued hemorrhage; it was not occasioned, as far as could be ascertained, by any improper exertion. It produced an excessive distention of the globe, and was attended with exquisite pain. These

symptoms commenced in the evening of the day of operation, and, on the day following, the humor, loaded with an enormous coagulum of blood, protruded at the section.

I have met with other cases, in which hemorrhage into the vitreous cells occurred in consequence of a blow. Inflammation and swelling of the globe ensued, and the cornea, yielding to the pressure, sloughed, when the humor protruded gradually in the form of a large spongy mass, loaded with coagula of blood, so as forcibly to separate and distend the lids, and occupy the entire circle of the orbit. In these cases a severe pain is felt in the head two or three inches above the orbit, and in the temple. The occasional hemorrhage is profuse. The pain is relieved by opiates, and the eyeball ultimately sinks with a total loss of figure.

I have reason to believe that this disease, which at one period assumes much of the aspect and character of a fungus, has sometimes been mistaken for one of a malignant character; a mistake not very unlikely to occur, judging from first appearances, where disorganization is complete.

The vitreous humor is subject to a complete change of consistence and a total loss of transparency, the texture of its cells and its volume and figure remaining; the secretion is converted

Change of  
substance of  
the vitreous  
humor.



from a transparent albumen into an opaque substance resembling curd. In one case it was like ground rice boiled. Although the opacity is visible, the appearance differs widely from that of cataract. While the crystalline remains transparent, the same bright-colored appearance is seen at the bottom or sides of the eye, which is supposed to announce the incipient medullary fungus. In the progress of the disease also, as in the malignant disease, the lens appears to become opaque, and is protruded so as forcibly to dilate the pupil; this becomes fixed, its edge roughened by detached pigment, and the iris convex, so as to give a conoidal figure to the globe.

Several years ago I extirpated the eye of a fine infant, eight months old, in whom this disease was concluded from the appearance described, to be the malignant fungus in its nascent state. The child has since grown to be a fine healthy boy; the other eye has remained sound. Upon section of the eye, the vitreous humor presented the appearance which I have described; the tunics were all entire.\* As the loss of vision had been but recently discovered, and the appearances had in the interval sensibly changed, so as to denote the progress of the disease, this case could not be considered as a congenital mal-organization.

\* See Plate III. fig. 4.

I have since seen several cases of a convex and permanently dilated pupil, with a deep-seated opacity of a splendid yellow tint in children, and doubting, from the preceding history and the child's freedom from indisposition, that such appearances indicated a malignant disease, I have abstained from operating. To my surprise, the appearances have continued stationary for years, unaccompanied with any disorder of the health. One, a child of four years old, I have very recently examined, having seen it at intervals during that period, since the first notice of the disease at the age of three months. I can discover no difference in the appearance of the eye at this time, from that which it then assumed. The bright yellow tint occupies the temporal hemisphere of the globe, supposing it were bisected in a vertical direction; the figure of the globe is slightly conoidal, the pupil much dilated as if from pressure, not perfectly circular, and its edge apparently everted, forming a narrow white line, while small detached flakes of the pigment lie behind it next the lens. The pupil of the other eye is constricted, and closed by an opaque capsule. The child is well grown and in perfect health.

I therefore regard these cases as a simple and uniform conversion of the substance of the vitreous humor, by an altered action of the secreting vessels, wholly independent of a malignant cha-



racter. Unfortunately we have no accurate signs by which to distinguish, in their incipient state, the malignant fungus and the disease above described; nor do we know that the disease may not sooner or later take on an active and malignant character, as is certainly now and then occurring in the testicle, female breast, and other textures.

The peculiar tint and splendor of the opaque substance (a mother described it to me as resembling, when first perceived, the scale of a tench,) is not to be depended upon as a sign of malignity. It evidently depends upon an opaque reflecting surface at the fundus of the globe; and the appearance is produced equally by an opaque retina, as by a morbid growth, except that in the latter it is somewhat more lustrous, from its greater density and projection; the convexity of the iris, the immobility of the pupil, and the apparent opacity of the crystalline, are secondary signs, and common to both. There are, however, two marks of distinction sufficiently strong, between the malignant fungus and this disease of the vitreous humor; viz. the progressive or stationary condition of the disease, denoted by the state of the tunics and the eyeball generally, and secondly, the presence or absence of pain and constitutional irritation. To these I might add, especially as regards children, the affection of one or both organs, as affording a

strong presumption that the disease is harmless in the first case; in the second, a conclusion that it is malignant.

It is remarkable, that both the fungus hæmatodes, or malignant fungus of the eyeball, and the disease which I have just described, are of most frequent occurrence in infancy; I am not aware of having seen the latter in the adult. I have now under my observation a very remarkable case of *congenital* malignant fungus. The child is eight months old. At its birth the eyeball was of the size of a walnut, it is now more than three times that size.\*

\* There is another affection of the vitreous humor, which ought not to be omitted in the description of the diseases of that part. The gradual conversion of this texture into a yellowish green substance, with a partial or complete abolition of vision, constitutes the disease known by the name of *Glaucoma*.

By the German surgeons, this disease is considered as always the consequence of a more or less evident gouty inflammation of the eyeball, although they state that sometimes the inflammatory arthritic affection is so slight, and proceeds in so lingering a manner, that only a very acute observer readily discovers the true disease. It is, however, by no means proved, that this affection can fairly be considered as necessarily belonging to a gouty diathesis; as in several cases of it, which I have seen, there was no other symptom present to warrant the belief that the patient laboured under gout.

It appears to be more consistent with sound pathology, to consider the disease as essentially an inflammation of the vi-



treous humor, and all the morbid changes which occur, as dependent upon this inflammation for their cause.

When the disease commences, the patient suffers pain across the eyebrows, sometimes slight, at others excessively severe, and accompanied with a sense of rending and extreme distension. If we now examine the eye, we perceive that there is a change of color, not immediately behind the iris, as in cataract, but deep-seated in the globe. This, perhaps, is not at first easily distinguished, but in the progress of the disease, becomes extremely manifest. By dilating the pupil by means of belladonna or stramonium, and exposing the eye to a bright light, we can see the change of color very early in the disease. The patient very soon begins to be troubled with specks and motes floating before the eyes, and finds his vision beginning to be seriously impaired. As the color of the vitreous humor deepens, the sight is more and more injured, although not always in an equally increasing ratio. The pupil gradually becomes more and more sluggish in its motions, until eventually it remains fixed. After a time the lens frequently undergoes the same change in appearance, and assumes also a greenish color; *cataracta glaucomatosa*, as it is called. It increases also in bulk, fills the posterior chamber, presses upon the iris, and at last pushing itself into the dilated pupil, partly enters the anterior chamber. Eventually the pain subsides; the eyeball begins to diminish in size, becomes atrophic, and sinking into the orbit, the lids entirely close over it.

When glaucoma has commenced in one eye, it generally, if not always, extends also to the other, if not arrested by remedies; and we will often see the disease in its different stages in the two eyes.

In its fully formed stage, glaucoma is absolutely incurable; but it may often be checked in its progress; and when only one eye is yet affected, it may sometimes be prevented extending to the other. We cannot restore the transparency of the vitreous humor, but may occasionally arrest the disease, and even improve the impaired vision.

*Treatment.*—This must be purely antiphlogistic, and in general, similar to what is proper for inflammation of the deep-seated textures of the eye. Local blood-letting, by means of cupping or leeches, frequently repeated, purging, and counter-irritation, perseveringly persisted in, are the proper remedies in the first stages of the disease; together with regulated diet, and rest both of the organ, and the body generally. This treatment may be advantageously followed or accompanied by the moderate use of mercury, in its milder forms, such as Plummer's or the blue pill. Either of these may be used until they slightly affect the system, but not too freely or too rapidly; as the attendant state of the constitution is generally such as not to bear the remedy in large quantities. Under the use of this all-powerful medicine, we are assured that after the change of colour had evidently taken place, the transparency of the eye has been completely restored, and the patient regained perfect vision.

This disease has proved so rare in my practice, that I have not confined myself, in describing it, to the results of my own observation, but made use of the descriptions given of it by the German authors.—*Editor.*



### SECTION III.

---

#### CRYSTALLINE HUMOR.

Abscess of  
the capsule.

I AM not aware of having witnessed any other result of inflammation of the crystalline and its capsule, than opacity, with the exception of one case. This was that of a lad, who, after a severe blow on the globe of the eye, which produced acute inflammation, had a suppuration within the crystalline capsule, which projected through the pupil in a globular form, and was filled with pus. There was no deposit upon the iris. Under the action of mercury the pus and the lens were absorbed together. The continued application of belladonna did not prevent the gradual constriction of the pupil, and the case terminated in a capsular opacity, with constricted and mis-shapen pupil, and co-adhesion of the iris and cornea.

Capsular  
opacities.

The capsule readily unites by adhesion, when simply incised. This is of course prevented by the intervention of any portion of lens. The capsule, when adhering to the iris, receives delicate red vessels, which run in small brown peduncles or foot stalks of lymph, produced from the interior border of the pupil; small flaky

portions of the pigment are also frequently detached, and conspicuous upon the margin of the capsule. This is an appearance commonly seen in the constricted pupil with partially transparent capsule, after chronic iritis. It is often the result of repeated attacks of inflammation at short intervals, to which a constricted state of the pupil certainly predisposes. The iris is much thickened by repeated depositions of lymph, until its texture becomes quite altered. There is a very imperfect and deranged state of vision, according to the degree and extent of opacity of the capsule, which admits of no improvement by the direction of the light; and sometimes a marked and painful determination of blood to the head. Except in this case, and in punctured wounds, the capsule is seldom partially opaque, but though its opacity is diffused, it is often not of uniform density, so that it has a dotted or mottled appearance. When calcareous matter is deposited, it is in small flakes or scales, which have a brighter tint than the opaque membranous portion. The opacity of the capsule, as of the cornea, varies in degree, from the slightest nebula to the opacity from change or conversion of texture. The incipient nebulosity is often, as before observed, difficult of discernment. Where the capsule is completely opaque, the lens undergoes a slow absorption; the capsule, however, remains transparent in most cases of senile cataract, not pre-



ceded by inflammation. The capsule, like all other textures of the body, undergoes absorption when detached.

Amaurosis  
mistaken for  
cataract.

The cataract, from opacity of the humor Morgagni, is in my belief purely hypothetical. There is an appearance of semi-opacity and yellowness in the lens, in some cases of defective vision, insufficient to account for the degree of dimness. I believe that this is not the cause of it, and that the disease is amaurosis. This I judge from the appearance being stationary, and the symptoms being those of the latter disease. I have known it to be confidently pronounced an incipient cataract, and the patient to remain in anxious expectation of an operation for years, without any sensible change of its density, although in the interim the person had lost his vision. On the contrary I may remark, that the degree of opacity of some soft caseous cataracts, when held up to the light immediately after extraction, appears quite insufficient to explain the degree of blindness, although the sight has been restored by the operation.

Species of  
lenticular  
cataract.

The fluid, flocculent, caseous, and hard cataract, are the four principal and easily distinguished degrees of density of the opaque crystalline. The caseous admits of division into soft and hard, as it approaches nearer in consistence to the second or fourth species. The

nucleated and mixed cataract—the first an opacity confined to the centre, the circumference and superficies transparent, the second a soft caseous or fluid superficies upon a firm centre—are well marked varieties. I refer the reader to my Papers on this subject in the fourth and fifth volumes of the Medico-Chirurgical Transactions.\* The opacity of the posterior capsule i. e. the tunica hyaloidea, is very rare, which it would not be if the lens were invested, as most persons suppose, in a capsula propria, especially after the operation of extraction, in which the anterior capsule only is lacerated, and the lens alone escapes. Where it is met with, the lens and anterior capsule are usually transparent, and when this is not the case, and the cataract escapes with a posterior fold of opaque capsule, it is always in my experience accompanied with a considerable discharge of vitreous humor, for it is owing to a detachment of the tunica hyaloidea, beyond the angle of union with the crystalline membrane, or a separation, beyond the margin of the lens, of the opaque from the transparent portion of that tunic.†

\* Vol. IV. p. 278, and Vol. V. p. 391.

† Until the perfect capsule of the crystalline lens can be exhibited detached from the eye, and the vitreous capsule and Petitian canal at the same time demonstrated entire, and capable of complete inflation, I shall continue to believe, through evil and good report, that the appearances actually and uniformly presented, and supported by all the phenomena of the pathology of these parts, warrant the opinion which I



What has been called 'black cataract,' when this term is used as applied to the lens, and not synonymously for amaurosis, is a modification of the fourth or hard species. A yellowish brown is the ordinary tint, but I have seen it occasionally of a blackish brown color.

Formation  
of cataract.

Some peculiar circumstances relative to the formation of cataract deserve to be briefly noticed.\*

Although the period of its completion from its first appearance is very variable, and can never with any certainty be predicted, it is usually slow; and sometimes a clouded or semi-opaque state or a distinct nucleated opacity remains stationary for years, or even for life; yet it occasionally forms with rapidity although no inflammation is present.

The rapid formation of cataract is generally attended by inflammation, or preceded by diseases of other textures. The result of iritis has been already explained. I have seen a yellow-colored opacity posterior to the lens preceding the rapid formation of cataract. The eye became the subject of a superficial but violent in-

have elsewhere given, that the lens is incased betwixt the strong membranous production, commonly called the anterior capsule, and the proper capsule of the vitreous humor.

\* See note G.

flammation, from sympathy with its fellow which was acutely inflamed. The cataract was completely formed in two days. The pain was of the most acute kind, affecting the eyeball, temple, and cheek. In the former state some useful vision remained, which was now completely extinguished. The previous opacity was a morbid state, I believe, of the vitreous humor, and the lens transparent until the attack of inflammation.

The residence of a perfect cataract in the eye is injurious, or at least attended with much risk of destructive inflammation. The vitreous humor undergoes a partial absorption, and the lens, losing to the same extent its support, bulges forward and presses upon its capsule and the iris. What operators call a narrow anterior chamber, arises more or less from this cause. Where the capsule yields from a blow or by absorption, and the pupil is dilated by the protruding lens, a violent inflammation, attended with very acute pain, is the invariable consequence. I have known this happen suddenly and independent of external injury, where the formation of the cataract has been gradual, and unattended with pain; and the spontaneous occurrence, though not so frequent, is precisely similar in its effects to that produced by the too free laceration of the capsule with the needle.

Partial displacement.

I have heard of the complete spontaneous ab-

Spontaneous absorption.



sorption of the congenital cataract at an early age, and although the eyes had a constant tremulous motion, the person was in after-life enabled to follow useful occupations. I have never seen such a case, but I refer to an instance which is well authenticated.\* The tremulous motion of the globe is not incompatible with a perfect state of vision.

\* A well marked case of this description fell under my notice some years ago. A gentleman from Charleston, S. C. brought his child, a little girl seven years old, to this city, for advice, in consequence of the imperfection of her sight. He described the pupils as having had a white appearance a few days after birth, which after several months, began gradually to diminish, until, before the child was two years old, no trace of it could be seen. In the mean time, the child's sight had gradually improved. When brought to me, she could see sufficiently for all the ordinary purposes of life, but not so accurately as to learn either to read or sew. The eyes had the tremulous motion referred to by our author. The pupils looked perfectly clear, and were of a jet black. The case had been repeatedly pronounced amaurosis, and if merely examined independently of its history, might be readily mistaken for it. The history of the case, however, proved decidedly that the child had congenital cataract, which had been gradually absorbed, and thus the power of vision had been gained. Extract of stramonium was applied, and the pupils widely dilated. All doubt was now removed : slight spots of opacity of a silvery brightness, but very small and with considerable spaces between them, were seen scattered over the capsules of each lens ; and sufficiently accounted for the imperfection of the sight. The parents of the child would not consent to an operation at the time, and the result of the case has not been ascertained.—  
*Editor.*

The absorption of the matter of the opaque lens is quick in proportion to the looseness of its texture, and its complete exposure, by breaking up and detaching its fragments, to the operation of the aqueous humor; and also to the plentiful secretion of healthy aqueous humor.\* In confirmation of the latter statement I have observed, that in all cases of narrowed anterior chamber, by the partial co-adhesion of the iris and cornea, consequent upon the injury which produced the cataract, or inflammation from any other cause, it is slow; and that absorption does not take place during the existence of inflammation, in which state, the aqueous humor is in a morbid condition, and if the inflammation be deep-seated and protracted, the vitreous humor partakes of it.

Progress of  
absorption.

I have seen a case of dislocated lens occupying the anterior chamber, and producing inflammation of the iris; from which membrane it has derived an adventitious capsule of lymph, organized by colored vessels.†

\* I do not consider this fact as militating against the opinion given at page 204. The turbid state of the humor indicates the arrest of its secretion; and its copiousness and transparency, the unabridged extent and health of the secreting surface.

† The hereditary disposition to cataract incident to certain families has long ago been observed by authors. A very remarkable instance of it exists in a neighbouring state, New



Jersey. A girl, of sixteen years of age, with congenital cataracts is now at the Infirmary, and has been operated upon once by my colleague, Dr. Rodgers. Her grandfather, father, and two uncles, with a number of her brothers and sisters, and cousins, amounting altogether to twenty-one, are more or less affected with some degree of congenital opacity of the lens. In several of them it is so slight as not to prevent reading, but in all, the involuntary motion of the eye, so common in congenital cataract, is present.—*Editor.*

## SECTION IV.

---

### DISEASES AFFECTING THE EYEBALL.

UNDER this head might properly be classed the morbid changes which result from the various forms of injury, to which, as all parts are exposed, all are liable in common. These, however, it is impossible to enumerate, and indeed would be, for any practical purpose, unnecessary, after describing the peculiar morbid disposition of each texture, and the characters of inflammatory action which they severally exhibit. The mode and extent of the injury, and the instrument with which it is inflicted, will determine its importance, as regards the preservation of the organ and its functions. Concussion and extravasation of blood are, both of them, frequent causes of a sudden loss of vision, which is, in many instances, gradually restored. The lesions of the internal tunics are to be seriously apprehended, from the probability of the retina being included in the mischief—the case of foreign bodies penetrating deeply into the globe, from the probable disorganization and dissolution of the vitreous humor, and the consequent

Effects of  
injuries.



suppuration, or state of atrophy of the globe. Sometimes the organ is at once destroyed by rupture of the nerve and muscles, and a partial evulsion of the globe from its socket; or by so free a lacerated wound, as to occasion large staphylomatous protrusions of the choroid, or the escape of the humors in toto, and instant collapse. More frequently, however, the cornea, or lens, or iris, are so far severally or exclusively affected, as to leave the figure of the ball uninjured, and to make the secondary changes which may be expected to supervene, an object worthy of the most interesting and anxious attention; and, I may add, to afford considerable scope for the salutary exercise of art.

Suppuration  
of the eye-  
ball.

From a long continued and exasperated inflammation of the interior tunics, but more frequently from injury, the eyeball suppurates, and its texture is totally destroyed. The globe becomes rapidly enlarged, greatly protruded, and exceedingly tense. The conjunctiva, highly tumid and vascular, is rolled out upon the cheek, so as completely to evert the lower eyelid. The pain is very acute, lancinating through the eyeball and head, and continues day and night without intermission. The patient's health is greatly disturbed, and the symptomatic or irritative fever, as in the thecal abscess, or acute paronychia, is considerable. The anterior chamber is at first filled by soft white lymph,

then pus collects in quantity; the clouded cornea turns opaque and slowly yields by ulceration, or dies and sloughs off, when the contents of the globe are more or less discharged, and the pain and symptoms of irritation gradually subside. The eyeball afterwards shrinks up, and the cornea is obliterated. The hypopion or purulent secretion filling the anterior chamber, originating from internal ulcer of the cornea, is not accompanied by the enlargement, or the acute pain and high irritative fever which mark the abscess of the globe; these are referrible to the extreme distention and corresponding resistance of an unyielding texture like the sclerotica, as in the abscesses of tendinous sacs and thecæ. The termination is the same in both, viz. the perishing of the cornea.

I had formerly been led to suppose, that the malignant disease termed cancer, affected the bulb or globe of the eye. Such is the doctrine of most writers on the subject. I have, however, satisfied myself that, as regards the eye, this disease is peculiar to the lacrymal gland, conjunctiva, and eyelids; and I have classed it accordingly. On the contrary, I had believed that the disease termed soft cancer, medullary fungus, or fungus hæmatodes—for these latter terms are descriptive of the appearances of the same disease in its several varieties—had its origin in some individual and peculiar texture;

Malignant  
diseases.



some writers giving it to the retina and medullary substance, some to the vascular tissue of the choroid, and others to the fibrous texture of the sclerotic. I am, upon the evidence of many cases and dissections, assured, that this is not the fact as regards the eye; that, on the contrary, each and every texture, if we except the crystalline and cornea, is capable of generating it, and is occasionally its proper nidus. The disease, therefore, comes properly to be noticed in this section.

Malignant  
fungus.

The early appearances of this formidable disease, of which it has fallen to my lot to see numerous instances, have been accurately described by Mr. Saunders and Mr. Wardrop. The disease, in my experience, has proved speedily destructive, when arrived at that stage in which the visible enlargement and loss of figure of the ball, the consequent livid blue tint of the sclerotic, and the distended vessels of the conjunctiva and eyelids present themselves. The character of the disease is by these decided; they are proofs of its rapid and destructive progress, and taken together with the primary appearances, must be regarded as fatal prognostics. The staphylomatous protrusions of the sclerotic and choroid coats, may, without the exercise of a careful discrimination, be confounded with this disease. I speak from a distinct remembrance of two cases in particular, in which the exist-



ence of the malignant fungus was a matter of dispute between very competent persons, prior to the extirpation of the organ, by which its existence was instantly demonstrated; and from the recollection of several in which the proposal of an operation was overruled, and as the event proved, judiciously, the patients having remained well, and the organ tranquil. Abscess of the globe, chronic enlargement and disfiguration of it from choroidal inflammation, proptosis from enlargement of the appendages, and tumors within the orbit, hydrops oculi, &c. are, on the contrary, sufficiently easy of distinction from the malignant disease, by the absence of the characteristic signs of the latter. Nevertheless I have known some of these mistaken for it. I have before noted a case, viz. protrusion of the vitreous humor from the eyeball, with which the fungus protruding through the slough of the cornea might, *primâ facie*, be confounded by a person unacquainted with the history of the case. But it rarely happens that the origin of the disease has not been accurately noted by the patient or his friends, or that the germ of the disease, in the infant at least, is not apparent upon inspection of the other eye. The complexion, as the disease proceeds, acquires the leaden paleness of cancer, and the rest is broken by deep and lancinating pain. If a child is the subject of the disease, it is heavy-headed and lethargic, as one affected by hydrocephalus; disturbed by occasional convulsive starts; the



stomach often rejects food, the frame emaciates rapidly, and the highest possible degree of irritability and fretfulness is present. The child usually expires in convulsions. The adult suffers from spasmodic shoots of pain through the ball and head, and simultaneous startings in going to rest; but the constitutional disturbance is inconsiderable previous to the protrusion of the fungus; and the hæmorrhage, which usually comes on at this period, is exceedingly distressing.

The metallic appearance at the fundus of the eye, sometimes presents colored blood-vessels, branches of the arteria centralis, which penetrate the vitreous humor; the opacity seems to advance towards the pupil, and might be mistaken for a protruding soft cataract, an appearance which, as Mr. Saunders has remarked, is altogether delusive. In a case in which Mr. Hunter was consulted, the operation for cataract was actually undertaken, and the lens being found transparent, the eye was immediately extirpated, from a conclusion that the disease was malignant.

A dissection of the eye, after the destruction of the cornea, furnishes a very indistinct clue to the original seat of this disease; it is in this state a firm semi-organized mass of lymph intermixed with and surmounted by coagula of blood;\* all

\* It is seldom that the blood-colored fungus acquires that enormous size, and protrudes to such a deforming extent as

the textures of the eye are broken up, but here and there the vestiges of one or more may be traced. If examined at an earlier period, that is, before it fungates, the section presents the real character of the disease, the medullary or soft brain-like substance, which we see in other organs of the body, commonly arranged in small lobes or *moleculæ*. Sometimes one, and sometimes another of the proper textures, appears to be the matrix of the disease. The fungus, in some instances, adheres intimately to the sclerotica, and detaching the choroid and retina, throws these and the vitreous humor to the opposite side of the globe. In others, it splits the sclerotic into two *lamellæ*, distinctly originating in the interstitial substance of that coat. Its progress, by absorption of a part of the sclerotic, gives occasion to a rapid growth of the diseased texture, external to it, within the orbit; but more frequently the diseased growth is luxuriant upon the outer surface of the sclerotic, to which it is as firmly attached as on the inner, while that tunic retains its integrity, and forms a perfect septum between the diseased masses. Hence, fibrous membrane has been supposed to give origin to it. But in other cases, the disease unquestionably begins in the choroid, and that tunic gradually

is seen in Saunders, Pl. III. It is to be regretted that an examination was refused in that case, although there can be little doubt that it would have confirmed this description of the ultimate state of the disease.



degenerates into the diseased mass which, occupying a large portion of the globe, is throughout deeply tinged with the black pigment. Sometimes the sclerotica has a morbid growth externally, and the choroid on the interior. Occasionally, these tunics seem to be only secondarily affected, and are removed by partial absorption in the progress of the disease; the septiform productions, extended, displaced, and broken down, of the tunica hyaloidea, enclose the fungus, and although no vestige of the vitreous humor remain, it is plain that it has formed the nidus of the diseased growth. The iris and corpus ciliare evidently degenerate in the same manner as the choroid. Again, the disease has seemed to originate at the point of entrance of the optic nerve into the globe, pushing the humors before it; and the nerve itself has upon section been found diseased, both contiguous to the sclerotic, and at the distance of three quarters of an inch from that tunic, where the intermediate portion has preserved its healthy aspect. Nay, the optic ganglion, tractus opticus, and thalamus, have been repeatedly found, one or other or all diseased, and the surrounding adipose substance in the orbit has exhibited the disease in its genuine character, and to a considerable extent, where it had no direct communication with the diseased contents of the globe.\* So also, the lymphatic

\* I have a colored drawing which accurately exhibits this state, taken at the moment of inspection.

glands, at the angle of the jaw, frequently take on the same character of diseased structure; and in some instances the bones of the cranium present periosteal tumors resembling soft nodes, which upon section are found to contain the medullary substance, the external table of the bone being to the same extent carious. It appears to me, therefore, that this is not a disease of this or of that texture, as writers would insinuate, but of all the textures, the crystalline and cornea excepted, which yield to its progress, but never exhibit a specific change of texture. The process of suppuration, ulceration, and sloughing, have no share in the diseased action, and are only seen in the cornea. The other tunics and humors yield by progressive absorption, without pus. Its appearance, in comparing the sections before fungating, is evidently modified by its situation, as for example, in the sclerotica and choroid, adipose substance, and vitreous humor; the phenomena of color, of hemorrhage, &c. depend much upon situation, also the preservation of figure, degree of enlargement, and the rapidity of growth, according as it is confined to the interior, or originally affects the exterior of the globe, or communicates with the orbit. We see these modifications of appearance, upon comparing it in the different viscera, and in the cellular membrane, tendinous aponeuroses, and the nerves. The deep blue and black tubera characterise the choroid fungus; the medullary, the substantia



alba of the optic nerve or brain; a more dense fibrous brown tuber, clear of stain, is the production of the sclerotic. These several parts are found to be directly and peculiarly involved, upon dissection.

I have a preparation exhibiting a genuine example of the disease affecting the anterior right lobe of the cerebrum, and protruding the eye from its socket, while the eye itself was perfectly free from disease. This shows the progress of the disease by absorption, and not by contiguity, conformable to our opinion of its nature.

---

A proptosis or protrusion of the ball conveys a delusive idea of the increased magnitude of the organ. I have occasionally met with cases of proptosis to such an extent as to occasion a morbid change upon the cornea, with a varicose state of the vessels of the conjunctiva; and with others, in which, although the cornea remained clear, the vision was materially deranged, where the cause of the protrusion was altogether obscure, and the progress of the disease had been so slow as to occupy a period of several years.

In these cases both eyes are equally affected; they are attended with a distressing degree of deformity, obtuse pain in the forehead, and other occasional signs of determination of blood to the

head. It is probably a morbid increase of the adeps contained in the orbit, obstructing its circulation, as well as protruding the eyeball. The disease, when acute, is of a different and more formidable nature. The following short history is an example:—

Highland, a lighterman, stout and healthy, aged twenty-nine, after frequent bleedings from the right nostril, with an obstructed and snuffling respiration, for which he was unable to account, was attacked with a severe pain over the whole front of the head, with a sense of weight in that part, and extreme lethargy. Although naturally of an active cheerful disposition, he became morose, indolent, and fond of solitude; at intervals he was attacked with tremors, cold perspirations, and syncope. These symptoms had become established, when the right eye began to protrude from its socket; his pain was at this time more severe, and a copious glairy discharge was set up from the nostril. As the disease advanced, his manner to his relations became strange, his intellect confused, and his gait unsteady. The protrusion steadily increased for several weeks without abatement of his pain, except for a few hours after occasional blood-letting. Convulsions at length ensued, and terminated his existence about three months after the commencement of the proptosis. It is remarkable that he retained the vision of the affected eye up to the period of its



protrusion; and before that was obvious to his friends, he described the sensation of something pushing the eye out of its socket. It is also remarkable, that through the whole period of his disease, although his bowels were extremely torpid, he had a good appetite, and little, if any, febrile irritation. These particulars I learned from his surgeon and relatives, having myself seen him only a few days before his death. Upon dissection I found the following appearances:—behind the cavity of the right orbit lay a tumor, which had the appearance of an oblong polypous cyst, and anterior to this was a blood-colored fungus filling the orbit and extruding the globe. The cyst lay anterior to the dura mater, adhering to its surface, and so situated as to make the right hemisphere of the cerebrum appear truncated of its anterior lobe. The æthmoid bone, frontal sinus, and orbitar plate of the os frontis on the same side, were in a state of caries, so that the finger passed readily from the orbit into the cavity of the cranium and posterior nares. A large quantity of yellow viscid matter occupied the frontal sinus, such as had been discharged during life by the nostril. The os frontis in front of the sinus and above the orbit was denuded, and presented numerous worm-hole ulcerations. The anterior lobe of the brain was discolored and softened; there was an extra quantity of water in the left ventricle, and some fluid blood in the right. On a transverse section of the right hemi-

sphere of the brain, it was found broken down in its texture, and the dura mater partially absorbed at its basis, the tumor having opened into the ventricle. The right thalamus was much diminished in bulk, though entire. The hæmatoid fungus in the orbit was mingled with spicula of bone, and distinct from the tumor. The dura mater to which the cyst adhered was continuous behind the cyst, except at the lower part, where it was destroyed. The disease appeared, therefore, to be connected with the external surface of the dura mater, by its increase to have occasioned absorption of the bones and displacement of the eye (which was sound, as were also the optic nerve and muscles of the globe), and ultimately to have ulcerated through the dura mater and anterior cerebral lobe, and discharged itself into the right ventricle. All the nerves were sound except the olfactory; this had disappeared, together with the æthmoid bone, on the right side.

---

SECOND EDITION.—I have two cases at present under my observation, of malignant proptosis from some such formidable and inaccessible disease as that above detailed. In one, a polypous tumor plugs and distends the nostril on the same side; in the other, the temporal wall of the orbit has given way, and the disease is extending to the integuments.



# PATHOLOGY

OF THE

## APPENDAGES.

---

### CHAPTER III.

---

#### SECTION I.

##### ORBITAR APPENDAGES.

Abscess.

ABSCCESS sometimes forms within the orbit, and previous to its discharge occasions an equal protrusion of the globe, with eversion of the palpebra, dilated pupil, and suspended vision. Its situation and effect upon the eye give much pain and apprehension to the patient, as well as considerable disturbance to the system at large. The sight is sometimes permanently extinguished. In other cases it returns after the discharge of the abscess, and consequent removal of pressure.

Adipose and  
encysted tu-  
mors.

Adipose and steatomatous tumors are occasionally formed in the cellular and fatty tex-

ture cushioning the globe. They occupy the interspace of the recti muscles, and in their progress emerge between the globe and the orbital circumference. They have therefore an oblong figure. The globe of the eye is turned and fixed in an opposite direction, and so compressed as to be rendered dim. I have removed them when projecting over the top or on one side of the globe, in several instances. When the conjunctiva is freely divided, the fatty tumor is easily hooked forward, and dissected out by a few touches of the narrow bistoury. The cyst containing a fluid, which is usually transparent, it is not so easy to remove entire, owing to its tenuity and the instant escape of its contents if it be accidentally torn or wounded. In the empty or collapsed state it is scarce possible to extirpate it completely, and even though the integrity of the cyst be preserved, its extent backwards within the orbit renders this in some instances a matter of much difficulty. If the cyst be cut in half, although the lips of the wound heal kindly, it suppurates periodically and discharges many times at the cicatrix, which closes in the intervals. The encysted tumor, although it extend to the bottom of the orbit, seldom occasions the distortion of the globe. A disagreeable sense of numbness and coldness affects the integument of the glabella and forehead, after the division of the frontal and supratrochlear nerves; these therefore should be avoided in the operation.



The tumor sometimes projects exterior to the tarsus, so as to rise upon the palpebra, but more commonly it is beneath the tarsus and contiguous to the globe. In the former case the cyst lies upon the periosteum of the orbit and is adherent to it; in the latter it is adherent to the globe.

Hydatid tumor.

In the last annual report of the London Infirmary for diseases of the eye, is the notice of a singular case of hydatid cyst protruding the globe, with the following remark: "One of these cases was a protrusion of the eye from the orbit, by a cyst containing hydatids deeply seated in the cavity. The hydatids were evacuated by a puncture in the cyst; the eye returned into its natural situation, and the patient was completely cured. This is the only instance of such an affection that has occurred since the opening of the Infirmary."

Aneurismal tumor.

The looseness of the connecting texture in the orbit, and the number and tortuosity of the vessels, seem to predispose to that disease of the arterial and venal extremities, which gives origin to those peculiar vasculo-cellular tumors, the precise nature of which is not yet satisfactorily ascertained; which add to a structure most resembling that of *nævus maternus*, the formidable character of aneurism. See my case of aneurism by anastomosis in the orbit, in the *Medico-Chirurgical Transactions*, Vol. II. Art. I., and

another by Mr. Dalrymple, the able and ingenious surgeon to the Norfolk and Norwich hospital, Vol. VI. Art. 7.

Tumors sometimes form beneath the perosteum of the orbit, giving to the touch a firm resistance. I have seen several cases of this description where the tumor appeared to extend the depth of the orbit, and was presenting on the nasal side. Their anterior edge is thin, being bound down to the orbital circumference, but when they protrude and compress the globe to blindness, as is sometimes the case, it is to be inferred that they have attained considerable bulk posterior to the globe. I once removed one on the abductor side of the globe, by scraping it clean away from the bone; it was of the hardness of cartilage and of great extent. I am unable to say whether the disease returned, having soon afterwards lost sight of the patient. The impression I had of the case was unfavourable, from the character, as well as the extent and connections of the tumor.

Sarcomatous  
and cartila-  
ginous tu-  
mor.

Exostoses of the orbit are not common. I have never seen them in the living subject of a size to create deformity or material inconvenience.

Exostoses.

Polypi of the frontal, sphenoid, and ethmoid sinuses in their progress burst through the eth-

Polypi.



moid and lacrymal bones, and sometimes extrude the eyes, so as to occasion the most horrible deformity. If when they first appear at the inner canthus, having an elastic feel, we puncture them, a thick glairy fluid is discharged, but from the extent of the disease when it has advanced thus far, the swelling does not subside. I know of no disease which presents so truly formidable an appearance as the polypus of these parts, when it has arrived at such an extent as to break up the bony structure of the nose or antrum, and show itself in the orbits.

Diseases of  
the lacrymal  
gland.

The lacrymal gland is subject to simple interstitial enlargement, to suppuration, and to scirrhus, like other glands of similar structure; its enlargement is known by the lobulated appearance of the tumor, on further stretching the skin of the projected eyelid. It often suppurates in children, and occasions an excessive swelling above the upper lid, depressing the tarsus upon the globe so as completely to conceal it. The abscess may be conveniently opened and discharged beneath the lid, with a narrow curved bistoury. I removed the lacrymal gland greatly enlarged and in a state of true scirrhus, from the orbit of a middle-aged man, a merchant's clerk in this city.\* The vision of that eye had suffered considerably during the growth of the

\* Represented by Fig. 6. Plate II,

tumor; in other respects he continued quite well when I last saw him, after an interval of some years. There was no other deformity than a slight drooping of the lid at the outer angle. All these operations should be performed beneath the eyelid when the circumstances admit of it.\*

\* There is a variety of disease, accompanied by protrusion of the eyeball, different from any of those mentioned by Mr. Travers, and depending upon a different cause. In this case, we observe a gradually increasing and uniform protrusion of the globe going on for a certain length of time, without any change in the texture of the eye, or the appearance of any tumor at its side, but apparently depending upon a simple increase in volume of the cellular and adipose substance at the bottom of the orbit. It occurs only, as far as I have remarked, in scrophulous children, whose general health is much impaired. My attention was first called to this affection by a paper of M. Louis in the Memoirs of the French Royal Academy of Surgery.†

He there quotes a case from Theophilus Bonetus, of a child of three years of age, whose right eye was almost entirely protruded from its socket. The child was evidently scrophulous, and a tense and swelled abdomen showed the extent of the strumous diathesis. The patient, after having been purged, was for a month put upon the use of tincture of rhubarb. The eye gradually sunk in the orbit, in proportion as the abdomen decreased in size and tension; and when a healthy state of the abdominal viscera was re-established, the organ had regained its natural situation, without any other treatment.

No modern author, that I have consulted, notices this curious disease; but I have met with more than one instance of

† Vol. 5. page 214.



it. One case closely resembled that just quoted. It occurred in a negro child, of about six years of age, in whom the strumous diathesis was strongly marked. Both eyes were considerably protruded from their orbits. A mode of treatment similar to that adopted by Bonetus was pursued; and by a long course of purgatives, with other remedies calculated to improve the general health, I eventually succeeded in curing the disease.—*Editor.*

## SECTION II.

---

### FACIAL APPENDAGES.

**PALPEBRÆ.** The little abscess called a sty<sup>e</sup>, Hordeolum. situated on either edge of the palpebra, commonly has its origin in an obstruction of one or more of the meibomian follicles, like the mammary abscess, which is an obstruction of one or more of the lactiferous tubes. Obstruction is followed by inflammation of the follicle and surrounding cellular membrane, and it terminates in a little painful abscess. Another description of sty<sup>e</sup> is an abscess forming around the bulbs of the eye-lashes from inflammation of the ciliary foveolæ, which sometimes has its origin in a disease of the hair-bulb. After the discharge of the abscess, and before the falling off or evulsion of the hair, the foveola appears exulcerated within and at its mouth, and continues for a long time to discharge a thin ichor by which the edges of the tarsi are denuded of conjunctiva, or this membrane is kept in a state of exulceration; the discharge forms a scab, by which the lids are firmly agglutinated during sleep, but it is detached, and the diseased secretion renewed by their separation. By the recovery of the foveola the healthy growth of the



cilia is restored. Sometimes the affected foveolæ become obliterated by the adhesive action, when the hairs of course are not reproduced. An habitual production of diseased cilia as well as nails is not uncommon, and for this state plucking in either case is only a palliative remedy. The disease is in the glandular structure which forms them, and its obliteration or removal, by a natural or artificial process, is in most cases the only method by which it can be cured. Independent of the abscess of the meibomian follicle and ciliary foveola, the conjunctiva upon the margins of the tarsi, and the continuous cutis, are liable to furuncular inflammation. The term sty is I believe indiscriminately applied to all. The disposition to form sty is generally a mark of scrofula, and from one or other of these causes is continually recurring as the system is affected.

Warts and  
vesicles.

Transparent vesicles and skin warts are not uncommon upon the margin of the tarsi. The former occurs upon the meibomian border more especially, the latter upon the ciliary.

Lippitudo.

The first stage of lippitudo is a simple excoriation ; the second, an ulceration of the borders of the palpebræ. It is the result of inflammation of the palpebral conjunctiva, aggravated by the acrimonious quality of the vitiated meibomian

secretion. In the chronic form of the disease in strumous subjects, the conjunctiva is greatly thickened, indurated, and altered in its texture; the ciliary glands are destroyed, together with the fine cuticle of the lid, to some extent beyond the ciliary margin; and a partial eversion of the lids, owing to the tumefied state of the conjunctiva, increases the deformity. The mouths of the meibomian glands are obliterated, and the ducts plugged by their inspissated secretion; sometimes the conjunctiva fungates so as to render the eversion complete, and a process of cicatrization makes it permanent. Thus the ectropeon may result from a neglected lippitudo.

The eyelashes are subject to become morbidly dry, harsh, and variously distorted, instead of having an equal curve, and their natural softness and pliancy. The inversion of one or more cilia upon the conjunctiva produces, by continued irritation, a painful degree of intolerance with an undue secretion of mucus and tears, and an opacity of the conjunctiva of the cornea with the prolongation of the colored vessels upon this membrane. In this way the marginal opacity extends over the surface of the cornea and occasions blindness. This state frequently exists independent of entropion. The disordered state of the cilia, above described, is owing to a disease of the glands which secrete

Disease of  
cilia.



and nourish the cilia, as mentioned in speaking of the hordeolum.

*Tinea ciliaris.*

The *tinea ciliaris* is a disease of the ciliary foveola, and hence is commonly combined with *lippitudo*. In the slightest form a branny crust surrounds the roots of the cilia, the skin of the lid being very partially, if at all abraded; in the more advanced stage, a mucus is secreted by the inflamed and excoriated ciliary border, and the thinner part of it evaporating leaves a scab. In the most inveterate form, the disease is the same with the *lippitudo* which has advanced to ulceration; the cilia fall off, and either diseased cilia are produced, so that the pore and the hair mutually react and keep up the disease, or the hair gland is permanently destroyed and the foveola obliterated. The *porrigo* or *crusta lactea* overspreading the eyelids and cheeks, with chaps and exulcerations behind the ears, and within the *meatus auditorius*, are concomitant affections, especially in children.

*Trichiasis or entropion.*

The *trichiasis* is a morbid incurvation of the tarsus, affecting either a part or the whole of it, from the cicatrization and consequent contraction of wounds, burns, and *lippitudinous* ulcers upon its *meibomian* edge, or the palpebral fold of the conjunctiva. Whether the entire cartilaginous border or only a part is in-

verted, depends upon the situation and extent of the disease which has produced it. But it very frequently arises, especially in aged persons, from a simple loss of elasticity in the cartilage, or a redundancy of the integument of the lid and cheek, or these causes combined. There is a case less frequent, in which trichiasis depends on a thickened and callous roll of palpebral conjunctiva, over which the lid turns. From whatever cause the disease originates, the eye becomes irritable, the motion of the lids occasions pain and watering, and from the incessant friction of the tarsus upon the globe, results the inveterate chronic inflammation with opacities, fed by vessels overshooting the cornea. It is very rare, with the exception of the case of protruded conjunctiva last mentioned, that the excision of an elliptical fold of skin at the basis of the lid, is not an efficient remedy. In that case, the roll of conjunctiva must be excised. The excision of the entire cartilage I think superfluous, and a remedy far more severe than the disease.

The ectropeon is the result of injury to the eyelid, as wound, burn, herpetic ulcer, or the sequela of chronic lippitudo. The tarsus of the lower palpebra sometimes falls outward from an apparent loss of elasticity, or the unequal action of the orbicularis muscle. The lid receding from the globe, suffers the tears to collect in a pool

Ectropeon.



between them. An unhealthy state of the conjunctiva is, if not the cause, as when villous and redundant, a certain consequence of its eversion and exposure. The case is much aggravated when coadhesion after burns or neglected wounds, ulceration from any cause, or long enduring eversion, takes place between the skin of the eyelid and cheek. This case admits of palliation, but not of cure. I have much improved several cases by first detaching the fastened lid and forcing it to heal by granulation, and afterwards, removing a triangular portion of the cartilage, according to the proposal of a modern author, for the correction of the eversion, which is the best remedy for such eversions as do not depend upon the protruded conjunctiva.\* In this, which is the simplest case of ectropeon, the excision of the diseased conjunctiva is sufficient. Where the everted lid is adherent to the bone, there is a deficiency of cellular substance to produce granulations, and the case is, generally speaking, slightly if at all benefited by operation.

\* This remedy results from considering the relative condition of the tarsus and skin at the base of the eyelid, in the two diseases, entropeon and ectropeon. In the first the integument is elongated, in the second the tarsus. As in the first case by removing a portion of the redundant skin, we turn out the inverted tarsus, so in the latter by removing a portion of the elongated tarsus, we turn in the everted. It will be understood, that it is only relatively that we speak of the elongation of the tarsus. It is everted, and strictured in the state of eversion, by the skin.

Tumors of the eyelids are encysted, varying in size, and containing a thin yellow fluid, or a fluid of the consistence of honey, or a white and dense caseous substance; or like the common steatom. They are adhering to the tarsi or moveable. In the first case, they give a diffused elevation to the skin of the lid, are circumscribed, and tense to the touch. Upon examination on the interior of the tarsus, a white hollow spot is discerned, surrounded by a blush, which corresponds to the point of their intimate adhesion to and partial absorption of the cartilage. Upon a free incision through the cartilage at this point, the entire cyst is easily expressed through the section. A thickening of the membrane covering the cartilage, will give the sensation of a tumor which does not exist. The adhesion of these cysts to the tarsus is sometimes so intimate, that if their removal were attempted by dissection externally, it would be scarcely possible to detach them, without removing a portion of the cartilage. They are frequently two or three in number, and their fluid contents are not often absorbed.

Tumors of  
the palpe-  
bræ.

The steatomatous tumors form in the cellular substance beneath the cutis, and are freely moveable, and easily turned out through a free incision of the skin. The atheromatous and lardaceous tumors form on or near the edges of the tarsi, and are very common in weakly children,



in whom they acquire a considerable bulk. If left, the skin ulcerates, and a scab forms upon the top, when they may be readily expressed entire between the nails of the thumbs. They resemble the sebaceous tumors behind the ears, and on other follicular parts of the skin, arising from obstruction of the follicles, which, being dilated and their sides condensed by inflammation with the surrounding texture, form these cysts.

Abscess of  
the upper  
lid.

Suppuration of the upper eyelid occurs from slight causes of irritation. The matter should be early discharged, for the cellular membrane, owing to its laxity and abundance, is subject to so rapid and excessive a distention, that if unrelieved, it sloughs out, and an ugly puckering or even a permanent eversion of the tarsus ensues, from deficiency of substance for granulations. I have known this deformity produced in an aggravated degree, by the suppuration going on insidiously and unsuspectedly, under the mask of œdema. The origin of it was snipping off a small skin wart upon the palpebra with a pair of scissors, and the subsequent irritation of the wound by the application of court plaster.\*

\* The eyelids are subject to a peculiar species of ulceration, apparently of a syphilitic character; strongly resembling chancres in appearance, yet probably a symptom of secondary syphilis. These ulcers are very foul, of a white color, deeply excavated, with hard edges, and having no disposition to granulate. They commence on the ciliary margin,

**LACRYMAL PASSAGES.** The puncta are sometimes much constricted, sometimes obliterated by preceding chronic inflammation; hence epiphora from imperfect or non-absorption of the tears, proportionate suffusion from excitement, and dimness. The constricted or closed punctum is always best opened with the point of a middle-sized pin; it afterwards readily admits the dilating probe; but if no vestige of the punctum remains, it is useless to attempt to form an artificial canal. I have seen a congenital deficiency

Constricted  
and closed  
puncta.

and extend to both the internal and external surfaces of the lid. I have seen them involve the puncta; and in one instance, ulcers of precisely the same character were situated on the cheek and ala of the nose.

Whatever may be our opinion of the nature and origin of these ulcers, whether syphilitic or not, there is no doubt as to the appropriate remedy. They always require and readily yield under the use of mercury. Any of the ordinary forms of the remedy will generally succeed, but the most appropriate is the corrosive muriate in doses of gr.  $\frac{1}{4}$  three times a day; the patient at the same time making free use of decoct. sarsaparillæ. The ulcers may be dressed with the ung. nitric. oxyd. hydr. Under this treatment, they soon assume a healthy aspect, and granulating surface. The result is generally much more favourable than was anticipated; for little or no deformity is left after the healing of ulcers of the lids, which appeared to have destroyed a considerable portion of their ciliary margin.

Almost all the cases of this disease which have occurred at the New-York Eye Infirmary have been in females, and the greater proportion of these negroes.—*Editor.*



of the puncta, but the case is very rare; obliteration is much less so.

Patulous puncta.

The over dilated or patulous puncta occur in old people, with more or less separation of the lower lid from the globe. They are so large as apparently to have lost their contractile as well as absorbing power; the conjunctiva of the palpebra is tumid and slightly villous, and the meibomian secretion is morbidly increased.

Wounds of the lacrymal ducts.

The lacrymal conduits are subject to be wounded or divided, hence incurable fistulæ. I have met with several such cases, and have tried in vain to heal them. Small abscesses occur in or adjoining the lacrymal conduits, which are broken and discharged by the passage of the probe. In more than one instance, I have turned out a considerable quantity of calcareous matter wedged in these ducts, like the calculi of the salivary ducts.

Stricture at the mouth of the sac.

With the constricted punctum a stricture of the lacrymal conduit, at the entrance of the sacculus lacrymalis, is often combined. This stricture is readily ascertained, and easily yields to the dilating probe. The tears regurgitate, and cause suffusion, but there is no evidence of any affection of the sac. The most frequent situation of stricture is at the point of termination of

the sacculus lacrymalis in the ductus nasalis. The tumid state of the lining membrane during acute inflammation of the palpebral conjunctiva, occasions a temporary obstruction, and this will continue for a time after the subsidence of the inflammation, but in a less degree. The canal in this state is exquisitely sensible, and the use of probes is improper. The obstruction is gradually removed under the treatment adapted to the inflamed palpebral conjunctiva, with which the lining membrane is continuous.

Acute.

A more considerable and permanent obstruction arises from continued vascularity and slow thickening of the lining membrane, the sac becomes slightly elevated from habitual distention, and a little mucus on pressure is returned upon the eye. Yet there is no discoloration of the skin, or sign of inflammation of the sac, and the epiphora is partial, that is, the tears are only impeded, and the epiphora only occurs when the eye is employed or in any way excited, and the secretion quickened.

Chronic.

In the state of incomplete obstruction, if the tarsi are unaffected, no mucus is discharged on pressure, nor is the sac perceptibly enlarged, nevertheless the suffusion upon reading, writing, working, or exposure to cold air, is exceedingly troublesome. Water injected by the puncta will find its way into the nostril, but slowly

Obstruction partial.



or complete. and only in part; so it is with the tears. When from long continuance of this state the obstruction is confirmed, the epiphora is incessant, the sac becomes sensibly dilated into a tumor, and upon pressure a very considerable discharge of purulent mucus takes place, so as to flood the eye. The sac, when once habituated to a state of over distention and a secretion of purulent mucus, will not contract upon its contents. Thus a very moderate degree of obstruction, or no obstruction at all may co-exist with this state; it may continue, even though the bony part of the canal should be destroyed, and the matter on pressure flow readily into the nostril.

Abscess of  
the sac.

The lacrymal sac is liable to acute inflammation and abscess, a very troublesome, painful and disfiguring disease, the signs and progress of which are well known. The surrounding cellular membrane becomes œdematous, and the cheek and side of the face are enormously swollen, so as to obliterate the orbital fossa and fold of the lower eyelid. When the abscess is chronic, the sac having been previously distended owing to obstruction, the contiguous cellular membrane passes into the state of adhesive inflammation, so that the swelling is not dropsical, but firm and hard. This occasions considerable embarrassment to the young surgeon, from the obliteration of the points, the infra-orbital edge especially, by which the operator directs his in-

cision, and the depth of the sac from the surface. If the disease be left to pursue its course, the skin discolours, the sac ulcerates, and its contents are diffused in the cellular substance; or, as more commonly happens where a previous obstruction has existed, the skin and sac, being condensed by the adhesive inflammation, yield together, and the discharge is external. Hence the fistula lacrymalis properly so called, a term incorrectly applied to all stages of the disease, of which it is but the last. It must not be supposed, however, that abscess of the lacrymal sac is always preceded by obstruction to the tears, any more than that abscess of the prostate is always preceded by obstruction to the passage of urine. It is frequently a sudden and rapid disease unpreceded by epiphora. In other cases it is slow and obviously progressive from the state of imperfect obstruction and retention of mucus and tears. The termination of acute abscess is more speedy and favourable for this reason, than of the chronic. In the former case, upon introducing a common sized probe after opening the sac, it passes readily into the nose.

Fistula.

The lining membrane of the lacrymal sac is liable to take on a morbid action. It forms a hard and dense tumor, which slowly ulcerates and destroys the skin to the extent of the sac. An irritable button-like fungus, of a malignant aspect, is then protruded. A fungus of a looser

Fungus of the sac.



texture sometimes follows the abscess of the sac, or, in other words, the sac laid open by ulceration, throws up luxuriant granulations.

Dropsy of  
the sac.

The sac is also subject to dropsy, in which state it acquires the size of a pigeon's egg, projecting the lower lid next the nose; the tumor is perfectly transparent, containing a fluid like that of hydrocele. Its natural openings are closed, for it does not admit of evacuation by pressure; it is very considerably extended within the orbit on the nasal side. This has been termed the hydatid tumor of the lacrymal sac.

Diseases of  
the canal  
and sur-  
rounding  
parts.

Injuries, as blows flattening the nose in early life, or occasioning exostosis of the ossa nasi, or unguis, produce incurable disease, or permanent destruction of these passages. To these may be added polypi and fungous tumors so situated, or of such magnitude as to compress the sac or nasal extremity of the canal, or occasion absorption of its bony parietes; and caries of the spongy, lacrymal, æthmoid, or maxillary bones. The ulcerative absorption or exfoliation of the bones renders the opening into the nose preternaturally large. But in obstinate chronic fistulæ, accompanied with erysipelatous or herpetic inflammation, or ulceration of the surrounding integument, it is not uncommon to find the bones denuded of the periosteum on the orbital side, still preserved by the pituitary membrane which

remains attached to the nasal. In such cases the proper canal is usually obliterated.

There is a sponge-like fungoid growth, sometimes affecting the interior chambers of the face, which speedily fills and obliterates the nasal duct ; it is firm and almost brittle, much disposed to profuse bleeding, and very quickly regenerated when broken down in attempts to remove it entire. It occasions considerable deformity by forcibly expanding the nasal cartilages. I have long had a case of this description under my care, in a middle-aged woman. Many severe operations have been only available to keep it in check. The fungus is of five years growth.

I have often found the canal completely obliterated by ossific inflammation at its upper orifice in skulls ; and I know cases of enlargement of the ossa nasi, and of periosteal inflammation and thickening, marked by habitual overflowing of the tears and occasional erysipelatous inflammation of the surface, in which the canal is evidently destroyed. Malignant herpetic ulcers of the lupous class, not unfrequently occurring at this part, expose and destroy the whole lacrymal apparatus.



---

## PART III.

---

### TREATMENT

OF THE

## DISEASES OF THE EYE.

---

### CHAPTER I.

IT will be necessary in this department of my work to refer to the states of disease described in the Pathology, but I shall endeavour to avoid repetition, and to seize upon the principles of treatment, to the exclusion of over-minute practical details. The leading object of my undertaking, I have already accomplished, however imperfectly; namely, the description of the principal phenomena of disease in the several textures of which the organ is composed. For the purpose of illustration I have unavoidably anticipated, in some instances, the subject of treatment, and the remarks which I have yet to offer will lie in a small compass; for it would be idle to suppose that general principles of treat-

ment require to be enforced, after the nature of a disease is clearly pointed out. The maxims and modes of successful practice, so far as they are hitherto known, are accessible to all enquirers of ordinary capacity ; and that man is unworthy of his profession who seeks to mystify them, for the purpose of being esteemed wiser than his neighbours. The innumerable modifications and varieties of disease render it impossible to lay down rules that admit of universal application, and the general intelligence of the profession in the present day, forbids such an attempt, if it were in the contemplation of any individual to make it.



## SECTION I.

---

### SIMPLE INFLAMMATION.

**Causes.** THE causes of ophthalmia, like those which lead up to inflammation in other organs, refer first, to the state of the system; secondly, to the direct operation of external agents upon the organ. A person whose general health is disordered, or who is recently convalescent from some other malady, frequently becomes the subject of ophthalmia. The extreme states and sudden changes of temperature; the prevalence of easterly winds, of fog and damp, and peculiar conditions of atmosphere; exposure to draughts of cold air; concentrated heat and light; extraneous particles, and other less obvious circumstances, are regarded as occasional causes.

Whatever is the exciting cause of inflammation of the conjunctiva, the first visible phenomenon is a state of congestion, owing to an increased influx of blood into the capillaries: it may pass away, but the continuance and increase of this, with certain other phenomena,

determine that the act of inflammation is set up.

The sensation of a foreign particle within the lids, whether real or delusive, commencing with the state of congestion, gives occasion to a spasmodic contraction of the orbicularis palpebrarum. A pungent pricking pain in the organ creates a copious secretion of tears, which collecting within the palpebræ, gush out at intervals, and their discharge affords a temporary relief. The increased temperature, volume, and sensibility of parts under inflammation, explain the following symptoms, viz. the sensation of burning heat and scalding tears, the constriction or girthing of the eyelids, and a sense of weight upon the globe; the involuntary exclusion of air and light, and sympathetic pains in the region of the orbit.

Symptoms

The simple inflammation of the conjunctiva, i. e. an inflammation not sympathetic with injury to the organ, nor depending upon any established disorder of the system, nor modified by a scrofulous diathesis, is easily and speedily reduced by the ordinary means adapted to this end. In its acutest form the loss of a few ounces of blood, and some brisk doses of purgative medicine are sufficient to subdue it. Even when it arises from superficial injury to the cornea, if treated in the commencement, it is scarcely less manageable.

Treatment  
of simple  
ophthalmia.



Febrile irritation.

It is rarely that any sensible febrile irritation is present in simple acute ophthalmia, but if there be any, and in certain irritable habits the constitution sympathises with the smallest local malady, it yields to the means above mentioned; repose of the organ, soothing applications, suspension of ordinary employments, a light vegetable diet, and diaphoretic diluents.

Blood-letting.

In many inflammations it is unnecessary to draw blood; the organ recovers speedily on the removal of excitement, the use of soothing applications, and the operation of cathartics. In some, general blood-letting is contra-indicated, both by the character of the inflammation, and the habit of the patient; while on the other hand, the degree of congestion makes it desirable to assist the recovery of the organ by unloading the vessels in the vicinity. In others, a question may arise as to the mode of proceeding to be adopted, in which the patient's convenience or preference may be consulted, or a disposition habitual to not a few persons, to a troublesome erysipelatous inflammation after leech-bites, may be admitted as an objection;\* but there are cases in which the indication is peremptory, both as to the use of topical and general blood-letting. If it be important to make the system

\* The swelling and discoloration which so often follow the application of leeches to the eyelids, especially the lower, make the remedy little less an evil than the disease.

sustain and feel a reduction of power, blood must be taken by the lancet, either from a vein or the temporal artery.\* Cupping has a very decided superiority over leeches; both are well adapted to relieve local congestion. But these modes of depletion are obviously too indirect, however extensively employed, to be used with the first-mentioned view, as a substitute for the lancet. Bleeding from the angular vein, and scarification of the conjunctiva are other means adopted for the relief of the turgid vessels. The latter practice is, in most cases, objectionable in the acute stage of inflammation; in the chronic it is highly beneficial, as in the thickened and over-vascular state of the palpebral conjunctiva; and a considerable discharge of blood may be thus obtained if it be briskly performed with a sharp lancet, the lower lid kept everted, and continually fomented with hot water.†

On the subject of local applications in acute ophthalmia, there is considerable variety of pro-

Topical applications,

\* See note H.

† Little good is derived from scarifications of the conjunctiva in the mode in which that operation is very commonly performed. If the scarification be not made exactly in the proper situation, so little blood is obtained as to be of no advantage to the patient. On everting the lower eyelid, we may observe about a line within the margin of the lid, a narrow strip of the membrane, exactly parallel to the tarsal border,



fessional opinion, but little in the evidence of patients. Dr. Johnson, whose opinion in these

more vascular than the rest ; the vascularity still more conspicuous if inflammation be present. A single stroke of a sharp scalpel drawn along this line will furnish a large quantity of blood, if the lid be kept everted as long as the bleeding continues. If the scarification be made even a single line nearer the globe of the eye, little or no blood is obtained. The situation meant to be designated is that in which the conjunctiva is firmly adherent to the tarsus ; as opposed to the loose portion of it passing from this part to the eyeball.

In this country, scarifications of the conjunctiva are so constantly practised in acute inflammation of that membrane, that I cannot avoid bearing my testimony against it. In the early part of my practice, I extensively used it, and must frankly acknowledge, often did much harm by it. The authority of Mr. Saunders, however, sanctioned by the remark on the subject by our author, ought to be decisive. "Scarifications," observes Mr. Saunders, "as far as I have seen them employed in the active state of the inflammation, are certainly injurious ; they have manifestly aggravated the symptoms ; and I conceive it will appear highly improbable that the infliction of mechanical injury on a part already actively inflamed, can be advantageous—a similar practice does not obtain in surgery on other parts of the body.—When the activity of the inflammation is gone, the vessels of the conjunctiva that have been engaged in the process remain preternaturally enlarged. At such time, the division of a great number of them might cause the whole series to contract, and thus accelerate the diminution of the vascularity, and the adhesive inflammation might only produce a degree of reaction, which would then be immaterial. But the condition of the part is widely different during the progressive state of inflammation. At this stage, the wounds of the lancet are only ad-

matters there can be no presumption in criticising, had a saying, "that there is little virtue in a lotion." In the main, and speaking of them comparatively, if it refers, as I conclude, to medicated lotions, I am much of his opinion; but warm and cold baths, whether employed for a part or the whole, have very unequivocal and sensible effects. These effects are likewise very

warm and  
cold.

ditional stimuli co-operating with the disease, and consequently exasperating the state of the eye."

Of the same character, as a remedy, as scarifications, we may consider the application of leeches to the palpebral conjunctiva; a practice attributed in England to Mr. Crampton, but long ago recommended in France, by Demours. Mr. Crampton insists upon the utility of the remedy in every stage and species of inflammation which attacks the eye. During the first stages of acute conjunctival inflammation, I have seen it do much mischief, and therefore would think it safer to limit its application to cases of chronic ophthalmia, or at least to those in which the severity of the symptoms had first been reduced by other means. With this caution, leeches thus applied constitute a safe and excellent remedy against ophthalmic inflammation.

The mode of applying the leech is sufficiently simple. Having selected one rather less than the medium size, the surgeon everts the lower eyelid with the fore finger of his left hand, and places the animal in such a manner as to fix itself upon the conjunctiva lining the lid, being careful to prevent its attaching itself either to the eyeball or the tarsal edge. Where the leech fixes itself to the eyeball, it sometimes produces severe inflammation, although no permanent injury is the consequence: on the edge of the lid, it gives great pain, and occasionally several days elapse before the irritation produced by it subsides.—*Editor.*



different, as might be expected. Although the sensation of cold is most agreeable to an organ under acute inflammation at the moment of its application, it is generally followed by increase of heat and pain; and in familiar instances, the pulsatile action of the vessels leading to an inflamed part, is so increased as to evince its stimulating effect, and the re-action thereby induced. When, however, the acuteness of inflammation has subsided, and the sensibility of the part is in proportion diminished, the effect of cold is only tonic, and has a salutary tendency to restore the balance of circulation. I therefore decidedly prefer, as a general practice, a tepid application in the painfully acute stage of inflammation, and I appeal to general observation in proof of its efficacy in promoting a grateful sense of coolness, and a more permanent relief from pain. It is objected to as being in the common phrase, "relaxing," which term exactly expresses its recommendation at the period of which I speak. We see its relaxing and resolving effect in incipient acute inflammations of the skin, the lymphatic glands, absorbents, &c. Moisture is a condition almost necessary to inflamed organs, and when the application is continued for some hours, as in poultices, it partakes so soon of the temperature of the surface, that this question is of less importance, but the indication is the same with very few exceptions. To conclude these remarks on what may be regarded as too trifling

to excuse prolixity, I prefer tepid water to all applications in the painfully acute stage of inflammation.

It is remarkable that even the weaker forms of medicated lotions irritate, and none more than that which is esteemed of all the most sedative, I mean opium. The relief afforded by anodyne fomentations in general, is very various. I have often known them objected to as painful, and patients to inquire if they might not substitute warm water for the aqueous solution of opium, and infusions of poppy and hemlock. The same observation applies especially to painful herpetic cutaneous affections, and acutely irritable ulcers. Upon these a solution of opium often acts as a stimulant and augments pain, while the lunar caustic solution as often assuages it. I do not deny that there are occasional exceptions to this remark. I have met with cases in which no other application than the aqueous solution of opium could be borne. I have also known the vapor of laudanum afford the most marked relief to the characteristic symptom of the strumous ophthalmia, viz. irritability to light.

Anodyne  
lotions.

Although during the state of morbidly elevated sensibility accompanying the outset of acute inflammation, warm applications are most soothing, and therefore most eligible; the continuance



of them beyond their necessity is a loss of time, if not injurious.\* When the extreme vascular

\* To avoid doing our patient injury is certainly of as much importance as to do him good. I shall therefore take occasion, whenever necessary, to warn the student of medicine and young practitioner against the use of remedies, which I know to be common in our country, where they are evidently prejudicial. Of these one of the worst, in inflammation of the conjunctiva, either simple or purulent, is warm poultices applied to the eyes. I have repeatedly seen young infants brought to the New York Eye Infirmary, with the corneæ of both eyes sloughed, and the organs entirely ruined, after having been kept under poultices even as long as six weeks without intermission: and the practitioner, perhaps, has not, during that whole period, once examined the patient's eyes. I have now under my care a gentleman, whose eyes were kept constantly under poultices more than a month, where the corneæ are both completely opaque and vascular, and the lids loaded with granulations. I need not say that he is totally blind, and many months will elapse before he recovers his sight. Such cases are by no means uncommon, and their result is a sufficient proof of the impropriety of the practice.

There is also another common practice which is equally prejudicial; the unnecessary confinement of patients suffering under ophthalmic inflammation, in dark chambers. A high degree of morbid sensibility to light is thus induced, which does not belong to the disease. Inflammation of the conjunctiva, when not modified by struma, is attended with very little intolerance, and hardly even requires the use of a shade; confinement in the dark is never necessary. No better guide can be found than the patient's own feelings; and he should be kept in just that quantity of light which is pleasant and agreeable to him. By an opposite practice, patients are often thus confined even months, until they cannot bear the slightest ray of light in their apartments; and long after the inflammation has subsided, the eyes are still useless from their want of power to bear its admission.

congestion and excessive sensibility are reduced, and the inflammation tends to become chronic, the use of cold lotions of a slightly tonic quality is substituted with great advantage for ablutions of warm water. The sulphates of alum and zinc are the best. The smearing of the tarsal edges with cetaceous ointment or cold cream at bedtime is useful in the acute stage, and as it subsides, the tutty or lead, or very dilute citrine ointment may be advantageously substituted.

The œdematous elevation of the conjunctiva is significant of a feeble action, and is by some regarded as erysipelatous. A more than ordinary fulness of the sclerotic conjunctiva is often combined with the nausea, foul tongue, and præcordial oppression, which manifest disorder of

œdematous  
ophthalmia.

When the inflammation is more deeply seated, and the superficial vessels of the sclerotic becoming enlarged, shoot over the margin of the cornea, and involve that membrane in the disease; then indeed, intolerance of light becomes a marked and distressing symptom; and it is necessary so far to diminish the light of the patient's chamber, as to make him comfortable. But even in this case, absolute darkness is prejudicial. A patient deprived of all light in his room, suffers acute and distressing pain at the opening of a door, or any other unavoidable admission of light; whereas if the room be only partially obscured, he suffers little inconvenience from any temporary admission of light.

Observations so minute as these, may perhaps appear trivial; but they are of so much consequence to the comfort of our patients, that they cannot be deemed useless.—*Editor*



the stomach and liver in cutaneous erysipelas; and the solution of emetic tartar given at short intervals, operates very beneficially in reducing it.

Atonic ophthalmia.

There are inflammations which assume a chronic character in their commencement, evidently depending on a state of atony, of very partial extent, void of pain, and scarcely possessing any sign of inflammation, except the congestion of vessels, or if any, so feebly marked as to encourage us to disregard them in treatment. In such cases, a single stimulus will often restore the healthy action at once. The vinous tincture of opium has acquired a nostrum-like importance, from its restorative operation in such cases; a virtue I believe not proper to it. A drop or two of the zinc or the lunar caustic solution, of water impregnated with calomel, or a minute portion of the citrine ointment, or any other stimulant introduced within the palpebræ would do as much. Some old women use their morning's urine with admirable effect in these cases. It is the character of the morbid action, not the application, that explains this sudden recovery. The re-excited or increased momentum of the arterial action clears the stagnant capillaries, and the unloaded vessels recover their tone. Such cases are frequently relapsing, unless means similar to those of cure are continued as prophylactics.

In certain habits, or states of the system—whether the ophthalmia arises from constitutional disorder or local injury—bleeding, purging, and blistering, the ordinary means of arresting inflammation, are employed without apparent benefit, or at least with a very disproportionate degree of advantage; and if the plan is persevered in, it soon becomes injurious; the irritability by which it is marked, increasing as the strength fails. These are cases in which opium, if we so combine it as to countervail its tendency to check the natural secretions, has an admirable effect, viz. with calomel, antimony, or ipecacuanha.

Irritable  
ophthalmia.

In the treatment of simple acute ophthalmia, the object to be kept in view is the soundness of the cornea; the organ is in no danger of deeper injury. The main indication for an activity of treatment beyond that successfully adopted in ordinary cases, is furnished by the state of this membrane. Where the sclerotic conjunctiva is much raised, and the surface of the cornea has in any degree lost its polish, and still more when lymph is effused in or upon the cornea, so as to obscure vision, the anti-inflammatory measures must be as vigorous and decided as the integrity of the organ is important. Blood-letting and blisters, calomel, antimony, and the neutral salts comprise all the requisite means.

Inflamma-  
tion threat-  
ening the  
cornea.



Chronic simple inflammation.

It is seldom that the simple inflammation becomes chronic. Blisters, and issues or setons, the zinc and acetous acid washes, and tonics, especially pure air and exercise, are the most efficacious remedies; but the red and thickened state of the conjunctiva at the margins of the lids, is an occasional and not unfrequent termination of it. Here scarification and the diluted mercurial ointments are employed with obvious advantage; but with some persons all greasy applications inflame so much as to aggravate the complaint, and in such instances moderately stimulant washes, of which a portion is to be admitted within the lids, may be substituted.

## SECTION II.

---

### INFLAMMATION MODIFIED BY STRUMA.

THE apthous inflammation, the inflammation of the follicles, and that characterized by intolerance of light in excess, and commonly denominated strumous ophthalmia, are almost always of an atonic character; and although obstinate when to a certain degree established, are easily subdued in the early stage, or at least prevented from arriving at such a height as to do permanent mischief to the cornea. They are, with very few exceptions, constitutional diseases; and the same remark applies to many instances of the mild acute suppurative ophthalmia. This is proved by the disorder prevailing in the system of nutrition, by the general debility of the habit, and by the concurrence of local affections in other parts referrible to the same source. The habit, age, and sex of the patient frequently contribute to the predisposition. The sphere and mode of life have also a decided influence in the production of these diseases. Children are most frequently affected by them, and those



especially, subject from infancy to glandular enlargements, chilblains, cutaneous eruptions and chaps, psoriasis, tinea, and porrigo. Imperfect nutriment, whether from the nature or deficient quantity of their food or defect in their powers of assimilation and absorption, contributes to them. To this may be added an impure atmosphere and want of cleanliness. The tendency to such diseases is demonstrated before they exist, and to prevent their recurrence is often more difficult than to remove them. It is common for a parent to say, "I know what will remove the complaint, but I cannot prevent its return." This however arises from neglecting to follow up the cure to its completion, and properly to employ the interval of the attacks.

A gentleman determined to relinquish animal food, and lived wholly upon vegetables and water. From the enjoyment of good ordinary health, he was in the course of six months reduced to a lamentable state of disease. The whole mucous surface became affected successively, after a severe and obstinate attack of mild acute suppurative ophthalmia. His system was so alarmingly debilitated by the protraction of his disease, owing to the prostration of his restorative powers rather than to the violence of the morbid action, that a residence in the south of Europe became necessary for the final re-establishment of his health. The dispo-

sition of such a class of diseases to fasten on the organ when once seated, in other words, to become chronic, is as characteristic as their tendency to re-appear when for the time removed. They come slowly, and so depart. Violent means fail to cure them. They are, to the surprise of persons who mistake their character, unaffected by such measures; which if persisted in, produce a change for the worse. If the remedies employed increase the debility of the system at large, it must follow that the part suffers, if this account of their constitutional origin be correct. Hence it is not uncommon for those who treat all inflammations alike, to express their surprise at the obstinacy of these affections, after going through and through again the routine of an active antiphlogistic treatment.

When the inflammation is of a sthenic character, as is more frequent where rapid changes are taking place upon the cornea, as a diffused opacity, or the formation of pustule and its passing into ulcer on that membrane, especially where the deeper-seated tunics are partaking by continuance of the inflammation, the necessity of a more active practice is sufficiently demonstrated. But, as a general observation, blood-letting is not salutary in these inflammations. They are rarely attended with any very acute pain. Rough and depressing purgatives, either



from quantity or quality, are injurious. Warm applications are of no advantage, if not injurious. Blisters on the nape of the neck and behind the ears, are for the most part of very great utility, where the severity or permanency of the inflammation calls for them. When the corneal surface is affected, and the sensibility is from this cause painfully augmented, this is especially the case. Blisters, when required in such cases, should be kept open as long as they do not irritate the system. Where the morbid appearances upon the cornea are notwithstanding stationary, or slowly progressive, issues and setons are of great avail.

The principle of treatment indicated in such cases is, to lessen the irritability without materially depressing the power of the system. The selection of medicine and applications, the regulation of diet, the degree of relief proper for the organ from its natural stimulus, when painful, must of course be determined by the circumstances of the case under consideration. The arrangement of the vessels at the verge of the cornea, and the condition of that membrane, are the special points for observation.

If the cornea be opaque, calomel, or the blue pill, or the oxymuriate of mercury should be exhibited in combination with opium, slightly to affect the system. The efficacy of the mercu-

rial preparation mainly depends on its combination with opium ; it irritates too much if administered alone in quantity sufficient for the purpose.

The following may serve as a synoptical sketch of the treatment for each form.

1. *Strumous inflammation without change of texture, vascularity more or less, intolerance excessive.*

Calomel and opium, or hydr. c. cretâ and Dover's powder at night; emetic tartar to nausea ; gentle alvine evacuants ; diaphoretic drinks ; large open blister on the nape of the neck ; leeches ; tepid bath ; tepid or cold water washes, as most agreeable ; vapor of opium ; large bonnet shade ; no bandages ;\* spacious airy apartments and light bed clothing.

2. *With recent diffused opacity of corneal conjunctiva, and vessels raised upon and over-shooting the corneal margin.*

Calomel and antimony, or opium, or any other more appropriate exhibition of mercury, to ptyalism ; occasional purgatives ; leeches ; blisters

\* Close bandages, I would observe, are always prejudicial. They create a morbid sensibility where it had not before existed, and greatly add to it when present.



alternated behind the ears and on the nape of the neck and temples.

3. *With herpetic ulcers of the cornea.*

The same; blisters on the temples; as the inflammation yields, sol. argent. nitrat.; vin. opii; sol. cup. sulph.; dilute zinc lotion.

4. *With pustules.*

If partial, weak zinc or alum lotion; ung. hydr. nitr.; occasional brisk purgatives; infusion of roses with additional acid; tonic bitters; calumba, gentian, &c.: blisters behind the ears, repeated if necessary. If the vascularity is diffused by the multiplication of pustules or the duration of inflammation with irritability to light; treatment as in strumous inflammation without breach. Ung. zinci. et sub-acet. plumbi.

5. *With inflammation of the follicles and puriform discharge.*

Active measures at first, but not long continued. Blisters, when becoming chronic, if with thickened lids, scarifications; zinc, alum, or copper wash, dilute; ung. hydr. nitr.; hydr. nitr. oxid.; sub-acet. cupri; tonics and sedatives. If obstinate, issue or seton.

## 6. *Convalescent state.*

Infusion of roses; cascarilla; calumba; decoction of bark, with dilute sulphuric or nitric acid; steel: rhubarb and soda, or magnesia, as aperients. Tonic collyria and gently stimulant ointments; nutritive diet; country air; shower or sea bath in the warm months.



### SECTION III.

---

#### ACUTE SUPPURATIVE INFLAMMATION OF THE CONJUNCTIVA.

THE suppurative inflammation is of all the most dangerous to the organ; and its sequelæ, even under a favorable termination, lingering, and sometimes difficult of removal. It is in its nature acute, but this acuteness is either mild or vehement. The former has been confounded with the inflammation of the follicles with puriform discharge, known also by the name of psorophthalmia, ophthalmia tarsi, mucosa, &c. I believe, as I have before explained,\* that the diseases are in their seat and nature distinct, and that the discharge is the only symptom common to both. From the occasional presence of diffused vascularity of the conjunctiva with the inflammation of the palpebra, the misconception has probably arisen. The flakes of mucus lying in the palpebral sinuses are not indicative of the suppurative inflammation, for these are often seen in the inflammation of the follicles, where

\* See page 97.

the palpebral conjunctiva is very slightly affected, and the secretion is not that of suppurative inflammation, but of an irritated mucous surface. (Fluor albus and gonorrhœa.) The intumescence and elevation of the palpebral conjunctiva (chemosis palpebrarum), its villosity, and the fluid and truly puriform nature of the secretion, are characteristic of the mild acute form of suppurative ophthalmia, in which the conjunctiva of the globe is also tumid and vascular; but I do not deny that under aggravation, the inflammation of the meibomian border and follicles may be followed up by the mild suppurative inflammation of the conjunctiva.

In the mild form of the complaint the cornea Mild. is not endangered, unless the disease be neglected or exasperated by stimulants. A very slight haze of the cornea is the worst direct result of it. There is not that excessive swelling of the lids, that intense pain, nor that profuse secretion, which characterises the vehement acute form of the disease; but these symptoms exist in a degree sufficient to require immediate and active treatment, and to this the acuteness of the inflammation speedily yields. The alum solution should be early substituted for the emollient fomentations, which, during the acute period should be freely used; and this should be directed in a gentle stream over the conjunctival surface, from a syringe furnished with



an ivory pipe, introduced at the temporal angle of the lids, without forcibly separating them. Simple purging and abstinence are generally sufficient to allay the febrile irritation, which is moderate. Topical bleedings and a suppurating surface opened by blistering the back of the neck, are of great efficacy. When the pain and irritability to light subside, and the discharge becomes gleety, the conjunctiva pale and flaccid, tonics, especially the extract of bark and the acids, do great good. As an application, the liquor plumbi acetatis may now be advantageously employed. While we are permitted to see the cornea, and to see it clear and bright, for this is the index by which we are guided, we need be under no apprehension.

**Vehement**

The vehement acute suppurative inflammation is sudden in its attack, accompanied with most severe darting pains; the upper lid is in a few hours prolonged upon the cheek, owing to the infiltration and enormous swelling of the tissue connecting the conjunctiva to the tarsus. The cornea is nearly concealed by the fold of conjunctiva which overlaps it all around, and the corneal surface is dusky. The system sympathises, chilliness is succeeded by a hot and dry skin, and the pulse is frequent and hard. The instant relief of a large venesection is indescribable. The pain is mitigated, if not removed; the pulse softened, and the patient sinks into a

sound sleep, and perspires freely. Upon inspection we observe the high scarlet hue and bulk of the chemosis sensibly reduced, and the cornea has a brighter aspect.

But it is rarely that a single blow suffices to vanquish the disease, especially where it arises, as is most frequently the case, from the contact of morbid matter. The most violent cases in my experience have been those produced by the matter of gonorrhœa applied to the eyes, of which I have seen several unequivocal examples. With large blood-lettings repeated, subject to the discretion of the practitioner, until the inflammation yields, a brisk catharsis should be combined, and this followed by a tea-spoonful of a solution of emetic tartar every hour, so as to keep up a state of nausea, perspiration, and faintness. The discharge, at first ropy, viscid, and sparing in quantity, becomes thin, gleety, and more abundant; as the swollen lid subsides, the conjunctiva sinks and becomes pale and flabby; and if at this period, the pain and febrile irritation being past, the cornea retains its tone and brightness, all is well; the disease has given way, and a careful but prompt exhibition of tonics, with the use of cooling astringent lotions, will prevent its lapsing into a chronic form. But if, when the lowering practice has been pushed to the extent of arresting acute inflammation, the patient being at the same time sunk and exhausted, the cornea



shows a lack-lustre and raggedness of its whole surface, as if shrunk by immersion in an acid, or a grey patch in the centre, or a line encircling or half encircling its base, assuming a similar appearance, the portion so marked out will infallibly be detached by a rapid slough, unless by a successful rally of the patient's powers we can set up the adhesive action so as to preserve in situ that which may remain transparent.

To know how far to go and not outstep the boundary; to know when to venture upon a short and sudden reverse of treatment, is the great difficulty of this highly important case. It is a fatal mistake to consider the first change, which is a true adhesive nebula, as the sign of gangrene or death, and thus to temporise, or even under this delusion to support the diseased action.\* Another and scarcely less mischievous error, is to treat the discharge as the disease, which is in fact but an inconsiderable sign of it as regards its importance, and to stimulate by strong astringent injections in its commencement. But the pathology which attributed the destruction of the cornea to the corroding quality of the matter secreted, was so lamentably erroneous, that we cannot be surprised at any effects, however mischievous, which resulted from the treatment thence deduced.

In closing my observations on the treatment

\* See page 119.

of inflammation of this organ, I shall take the liberty of making one or two general remarks. When inflammations in their nature destructive are arrested by the vigor of the means employed, the system stands in great need of the power thus lost for its recovery; to restore parts partially injured, and to supply the place of those which are destroyed. We see this fact exemplified in many instances both of disease and injury. A patient labouring under pneumonia is relieved by excessive bleedings of his attack, and dies a month afterwards of dropsy. A person threatened with apoplexy, who by the advice of his physicians is cupped once a month, soon falls a victim to erysipelas.

If much blood is lost in severe injuries, especially of aged people, the healing powers are prostrate and gangrene ensues. I mention this as a caution against that inconsiderate detraction of blood (and it applies as forcibly to the abuse of mercury) which proceeds without proportioning the quantity to the absolute necessity of the case, and, secondly, without balancing the effect upon the system at large against the importance of the organ. When I hear, as I often have heard, of sixty and seventy ounces of blood taken at one time for an ophthalmia, and this followed by repeated smaller bleedings, I must protest against the necessity of such a practice,



and say with Falstaff, "the better part of valor is discretion."

One of the great errors, it appears to me, in the treatment of inflammations of the eye, though of late years much corrected, has been the irritation of the inflamed organ by stimulant drops and ointments. The advantage of them is fully admitted at a proper season; but during the presence of active inflammation their use is as revolting to common sense as it is injurious.\* I am satisfied many eyes have been thus destroyed. An anomalous species of ophthalmia, or a pseudo-ophthalmia is produced by it, which differs as much from the real character of the disease in either of its forms, and may be as readily distinguished from it, as an artificial from a natural flower. Thus, to mention one of many cases, I

\* I have mentioned certain cases in which stimulants act beneficially. Even mustard has been applied to inflamed eyes, with some real or supposed benefit. The temporary relief which follows pungent applications is to be attributed to the copious secretion and flow of tears which they occasion, which is nature's own mode of relieving the distention of the vessels, quickened by additional excitement. The pain of every inflamed organ is augmented by the retention of its secretion, and in proportion relieved by its discharge. But the means employed to promote this end should not be such as are likely to support and increase the morbid action. Pain is only an effect of this morbid action, and to assuage it by measures calculated to perpetuate the cause, is, to say the least, a most unscientific method of proceeding.

have seen the star-like arrangement of the vessels around the margin of the cornea, the cornea and remaining portion of the conjunctiva clear, and the choroid and iris perfectly free from inflammation, the sequel of an inflammation of the follicles which had been incessantly stimulated; and I know cases of permanent and excessive congestion, or rather varices of all the veins of the conjunctiva with an actual discoloration of the sclerotica, such as would lead to the belief that the person laboured under confirmed organic amaurosis, in which however the sight is perfect. Here the plan of irritative applications had been unremittingly pursued by several practitioners in succession. The anomaly consists in the existence of such appearances unallied with the states of which, by their habitual association, we consider them characteristic. An apprehension suggests itself to my mind, when I see such cases, that the external character may be the prototype of internal disease, or, at least, that the confirmed existence of the one may predispose to the production of the other.



## SECTION IV.

## SECONDARY DISEASES OF THE CONJUNCTIVA.

Graular  
conjunctiva.

THE granular state of the tarsal conjunctiva is a very common result of the mild suppurative ophthalmia. It is characterised by a gleety discharge, irritability to light, drooping of the upper lid, a pricking sensation as of sand in the eye, and a preternaturally irritable and vascular state of the sclerotic conjunctiva; with these are frequently combined, opacities of the cornea. The lid should be everted, and the projecting granules shaved off from the surface and orbital edges of the tarsus, with a keen-edged lancet, or, if peduncular and prominent, they will be more conveniently snipped off with the flat scissors. In doing this, care should be taken to avoid injuring the continuous membrane.

With vascular  
cornea.

When in addition to the state above described, vessels are ramifying over the cornea, opacity of its covering conjunctiva being a contemporaneous result of the inflammation, or a consequence of the irritation excited by the granulations, a section of the membrane should be made at one line's distance from the margin of the cornea.

For this purpose, the globe should be thrown forward and fixed in a state of tension by depressing the edges of the palpebræ with the fingers. The membrane yields instantly to a light hand, and its edges gape asunder; in aggravated cases, the operation, which is painful, requires to be repeated, and some adroitness in exposing and fixing the globe is requisite to its complete performance. After the excision of the granulations and the division of the conjunctiva, a solution of the sulphate of copper, or some astringent, is very advantageously employed in the way of injection. A few drops of the liq. plumb. acetatis, or the tinct. opii vinos. are often highly effective. It should be observed that the two states above described often exist apart, but the treatment adapted to them respectively is equally essential. The application of the blue stone, or of the lunar caustic, is useful in preventing the regeneration of the granulations after their excision.\* †

Another consequence of the disease above described, are folds and flap-like elongations of the conjunctiva filling the palpebral sinuses, and occasioning such a fulness of the lids as to prevent the patient from more than half opening the eye. Upon eversion of the lids they roll out upon the cornea. Another state ensuing

Fungous  
conjunctiva,  
elongations,  
excrescences,  
pannus,  
&c.

\* See note I.

† See note at the end of the Chapter.



upon the excessive chemosis, is a fungoid protrusion of the conjunctiva in a thickened and indurated state. The conjunctiva also, at the point of its reflection from the lid upon the globe, occasionally forms a tumor of considerable magnitude. I have seen it projecting from beneath the upper lid equal in bulk to a middle-sized walnut, producing great distortion and inconvenience, and rapidly increasing so as completely to cover the eye. Such states more frequently result from injuries, as falls and blows. The treatment of all these cases consists simply in the excision of the tumors, which is most conveniently done with a lancet-shaped knife, cutting on both sides. The same may be said of the disease which I have denominated pannus,\* the elongated valvula semilunaris, and the caruncular excrescences which sometimes form in clusters between the tarsus and the globe. In the first named disease, a circular excision of the redundant opaque membrane should be made with the curved scissors at a short distance from the margin of the cornea, and the scissors will be found most convenient in the removal of elongations and excrescences, while such parts are raised by a pair of small forceps. The broad

\* This term is differently appropriated. In the disease to which I have applied it, the conjunctiva clothes and covers the cornea from that side to which the eye is directed, but it has no affinity to the membranous pterygium, or any form of nebulous opacity of that membrane.

or ring-ended forceps are often convenient on these occasions.

The fleshy pterygium is sometimes a chronic and even a stationary condition of disease producing no inconvenience, nor threatening to interfere with vision. Whenever this is the case, I am decidedly of opinion that it should be let alone. When, by its progress, it is encroaching upon the sight, it should be raised by dissection as close as possible to the margin of the cornea, and the relaxed portion of the membrane removed by an incision mid-way between the base of the pterygium and the cornea, and concentric to that membrane. I have experienced the inconvenience pointed out by Professor Scarpa, of carrying the excision to the caruncula, viz. the deposit of lymph in the site of the cicatrix becoming united with the caruncula, and forming a hard frenum or cord which prevents the abduction of the eye. I am also satisfied that the disease is permanently arrested when the connection with the cornea is dissevered. In this operation I prefer the cornea knife to the scissors. It is inadmissible to interfere with any portion of the pterygium that may have encroached upon the cornea. It may be necessary to repress the tendency to reproduction by the application of the caustic pencil to the section of the tumor; but the frequent or diffused application of escharotics is objectionable, as a

Pterygium  
and encan-  
this.



morbidly thickened and tubercular state of the membrane is the consequence of the irritation thus excited. The treatment of the membranous pterygium consists in nipping up a crescentic portion of the opaque membrane as near as convenient to the cornea, and freely excising it with a pair of curved scissors. The extremities of the line of excision both in this and the former species, should extend beyond the diseased part.

The encanthis, when it attains any considerable bulk, becomes condensed with the valvula semilunaris, and presents appendices corresponding to the cornua of this fold. The treatment consists in simple excision.

Frena.

The membranous bands connecting the lid to the globe should be divided, with the precaution to avoid wounding the palpebral conjunctiva. No bandage should be employed, and during the day the patient should not be suffered to keep the eyelids closed. In my experience, escharotics only exasperate the evil.

Tumors.

Tumors upon the globe, unconnected with the palpebra, should be dissected from the sclerotica; and this is the more important, in proportion as they are seated near to the cornea. Where the tumor is tied by angular folds to the eyelid, considerable attention is necessary to

prevent the adhesion of the conjunctival surfaces during the stage of healing, as in the case of frena. The best mode of preventing the approximation of the opposed surfaces, is to produce a partial eversion of the lower lid, by a strip of plaster carried from its margin in an oblique direction across the cheek, and frequently renewed.\*

\* *Granular conjunctiva* of the eyelids, with vascular and opaque cornea, is an extremely common disease in every part of the state of New York; but more particularly in our northwestern counties: and not a few of the inhabitants of those counties are permanently and irremediably blind, from the long continuance of this most unpleasant consequence of purulent ophthalmia. Such patients are constantly presenting themselves at the New York Eye Infirmary; and in a large portion of the cases, the result may be fairly attributed to improper treatment of the acute stage of the inflammation. The disease generally proves extremely difficult of cure, and is constantly liable to severe relapses during the whole progress of the treatment. From the very concise directions for its treatment given by Mr. Travers, the student might be misled, and imagine that it might always be readily and quickly cured. The operations of removing the granulations by the knife or scissars, and dividing the vessels which run upon the cornea, are by no means sufficient in themselves to eradicate the disease; the latter is generally unnecessary as well as inefficient. Cutting off the granulations removes them indeed for the time, but does not remove that condition of the conjunctival membrane which gives rise to them; and if we confine our treatment to this remedy, we will generally be disappointed.

When such cases present themselves, they are always ac-



accompanied with more or less of inflammation. Our first object must be the removal of this inflammation, and before this is effected, all operations are improper. For this purpose, the usual means in common use are sufficient, cupping or leeches, keeping the bowels open by proper medicines, regulating the diet, and removing all sources of irritation. Counter-irritation is, however, our most efficient agent in fulfilling this indication, and should be kept up during the whole course of the treatment. Blisters behind the ears, kept open by ung. sabin. or ung. canthar. may be employed; or still more conveniently, the tartar emetic ointment may be applied to the same situation, or to the nape of the neck. In the recent state of the disease, a due perseverance in these means will often be sufficient to subdue it; and in all cases they must be continued as long as inflammation is present.

If they do not succeed, it becomes necessary to use means for the direct removal of the granulated surface, and the restoration of the conjunctiva to a healthy condition. For this purpose, in mild cases, we may succeed merely by the use of astringent applications to the granular surface; and of these, one of the mildest and most efficacious is the liq. plumb. acet. undiluted, dropped into the eye once or twice a day, or smeared over the diseased surface with a camel's hair pencil. Other astringents are useful with the same view, such as solutions of alum and of nitrate of silver; but neither of them is so efficacious as the liq. plumbi. The ung. hydr. nitrat., diluted with three or four parts of lard, may at the same time be applied to the edges of the lids every day, to prevent their agglutination, and act as a beneficial stimulant to the parts. Counter-irritation should still be kept up either by means of ung. tartr. antim., setons, or issues.

If the case be more severe, or these means fail, still more active measures must be resorted to. If the granular bodies be large and hard, they should be removed by the knife, as directed by our author, and the operation should be repeated, if they grow again, until they are removed as far as possible.

As the whole of the disease, however, can hardly be eradicated in this manner, escharotic substances are employed to complete their removal ; and of these the sulph. cupri arg. nitrat., and the mineral acids are the best ; the first preferable to the other two. In applying them, the lids must be everted, and if the sulph. cupri be used, it should be lightly rubbed over the granulated surface ; if the arg. nitrat., the projecting points may be slightly touched with it. The part should then be freely washed off with water to prevent the application of the caustic to the eyeball. When the acids are employed, they should be diluted with three parts of water, and applied with a camel's hair pencil. Sometimes the use of any of these means excites active inflammation in the eye ; and then all irritating applications must be avoided, and antiphlogistic and soothing treatment employed.

The use of the knife or scissors to remove the granulations is very rarely necessary, and I believe, falling much into disuse in Europe. The sulph. cupri is a better remedy, and generally adequate to the object in view, if perseveringly employed. It may be applied every other day with perfect safety.

Having succeeded in removing the diseased state of the palpebral conjunctiva, the vascularity and opacity of the cornea generally subside, without the use of any direct remedies for their cure. They are excited and kept up by the irritation of the granular lids, and disappear with their cause. They are removed more rapidly, however, by still persisting in the use of the ung. tartr. antim. The circular incisions around the cornea recommended by Mr. Travers will sometimes quicken the process, but should never be used while any inflammation is present, nor until the palpebral lining is restored to a healthy condition. When employed under these circumstances, they sometimes excite severe inflammation.

This last remedy, the division of the enlarged vessels which overshoot the cornea, is so frequently employed in this coun-



try, that it is necessary to make some further remarks with regard to it. Another modification of the same remedy, as recommended by Mr. Ware, the excision of a portion of the enlarged vessels is also much in use among us, and when improperly made use of, is still worse than the first: indeed, if either be necessary, a simple division of the vessels is adequate to every purpose which the remedy is capable of effecting, and is liable to do much less mischief. Mr. Ware's method, therefore, is never necessary; and the other very rarely. Our object should rather be to remove the inflammation which enlarges the vessels, than to diminish their diameters by any operation on themselves. An opposite practice is very often productive of serious injury to the organ, by increasing the inflammation and aggravating all the symptoms. The common error is, to divide the vessels while inflammation still subsists: very often it is done in its very commencement. In severe acute inflammation of the cornea, in ulcers of that membrane with red vessels running to them, and even in common pustular inflammation of the conjunctiva, this operation is very commonly performed, and in every such case, if it does any thing, it does harm. No arguments are necessary to prove the impropriety of the practice in such cases; it is repugnant to the very first principles of surgery, and its employment cannot be justified on the slightest ground. Were I not a daily witness of the mischief arising from the improper use of this remedy, I could not have supposed that what is so evidently wrong, should so often be done. But with this evidence constantly before me, I feel it absolutely necessary to caution the student of this branch of surgery against an operation he will often be advised to perform.—*Editor.*

TREATMENT  
OF THE  
DISEASES OF THE EYE.

---

CHAPTER II.

---

SECTION I.

DISEASES OF THE CORNEA.

THE term, inflammation of the cornea, must be understood as applied to the compound texture so denominated, and not to the lamellæ of horny substance, which has no vessels proper to itself, but derives them from the covering and connecting cellular tissue. These vessels nourish and preserve it in the condition essential to its economy. The crystalline humor is a simpler texture, being wholly dependant on its capsule—as the nails, like the horse's hoof, are sustained by the lamellæ of the cutis in which they are implanted—or the hair, by the bulb alone to which it is attached. It is rarely that red vessels are seen in the inter-lamellar texture of the cornea. Deposits of adhesive matter and of pus are frequent; the former most so; those



of blood are rare, being only a result of severe injury, superadded to a state of inflammation. The cornea is rendered turbid by a congestion in the vessels of its covering or connecting texture; and in this, and the case of interstitial inflammatory secretions, may, if in any, be said to be inflamed. But its subserviency in these processes to the conjunctiva and sclerotica, makes the strict propriety of the term questionable as applied to the corneal lamella. It would be as incorrect to speak of an inflamed crystalline, hair, or nail.

Nebula and  
onyx.

It is only necessary to observe, that the practice employed to reduce inflammation is then most strongly indicated, when the cornea is rendered opaque, or presents an onyx of adhesive matter.

Superficial  
external  
ulcer.

The superficial ulcer is commonly attended with much inflammation of the conjunctiva, and, by continuance, of the sclerotica. The eye is very irritable to light, and the sensation of a foreign particle in the motions of the lids acutely painful. The pain is often spasmodic, and relieved by profuse lacrymation at intervals. Opium should be so combined as to operate on the skin, and the bowels must be kept freely open.

Touching the ulcer with a fine pointed caustic pencil, or the solution of argenti nitratum,

is the best local treatment ; much superior, as an anodyne, to sedative lotions. Warm fomentations afford temporary relief. It will be found advantageous, if not indispensable to prevent relapse, to affect the system with mercury where the inflammation of the sclerotica is intense. The cicatrix being confined to the superficial lamellæ is of very inconsiderable density, so as in time to be scarcely perceptible, and in children to wear quite away.

The indolent and the deep sloughing ulcer may be touched once, or oftener, with the caustic pencil, or washed once a day, or oftener, with the solution. The cleansing of the ulcer, and the opaque adhesive circle is the sign for a less frequent use of it, and the deposition of new matter, undergoing a vascular organization, renders its further use hazardous. The occasional use of leeches is often a necessary accompaniment to this treatment. The administration of tonics and sedatives is at the same time essential.

Indolent and deep sloughing ulcer.

The acute interstitial ulcer cannot be treated distinctly from the adhesive inflammation ; it is a sign only of the inflammation which constitutes the disease. In proportion as this is reduced, its disposition to extend is checked, or we are enabled to employ auxiliaries to that end. But in favorable circumstances of constitution they are not wanted. Healing is a spontaneous ac-

Acute interstitial ulcer.



tion, vicarious with destruction, and commences on the arrest of inflammation.

Abscess.

A large collection of matter in the cornea, whether the puriform onyx, or central abscess, requires, at the same time, a supporting constitutional treatment, mild cathartics, and the application of blisters; calomel should be avoided, as in most instances where ulceration is present. The puncture of the cornea is seldom practised with advantage. By the means above-named, I have seen large effusions absorbed, and no trace left of their existence.

Hypopion.

When the hypopion is so large as to rise towards the pupil, and the ulceration of the cornea is extending, I think its discharge by section near its margin advisable. If not too long delayed, the ulcerative process is checked by it, which would otherwise run into sloughing, and the cornea recovers with only partial opacity and disfigurement.

Procidencia  
iris.

The prolapsus iridis from ulcer should, if small, be touched with the caustic pencil, ground to a fine point. If large and extending, it should be snipped off with a pair of curved scissars, and the caustic pencil immediately applied to the cut surface and margin of the ulcer. In this way I have seen many cases recover with good though abridged vision. This circumstance depends on

the site of the ulcer and the relation of the prolapsed portion of the iris to the pupil. The same treatment is best adapted to prolapsus from wound, as after extraction. The inflammation accompanying these states requires the occasional application of leeches, gentle purgatives, sedatives, light tonics, and mild nutritive diet.

The chronic interstitial ulcer requires only stimulant and astringent injections; blisters in the neighbourhood of the eye, bark and opium, pure air and good diet, with a due attention to the secretions. Rhubarb and aloë are the best aperients.

Chronic interstitial ulcer.

The opacities, in their nature removable, are the nebulous, which depend on a loss of transparency from recent inflammation, or recent interstitial deposition without breach of texture. Cicatrices are only so far benefited as the surrounding deposit is of this description, and susceptible of absorption. The actually changed texture of the entire cornea depending on an obliteration of the interstitial texture, like the cicatrix itself, undergoes no change from the use of stimulant applications. The most effective injections are the lunar caustic and the oxymuriate of mercury, one or two grains to one ounce of water; the former may be used in the decline of the inflammation; the latter, not until after its disappearance. Levigated glass, calomel, loaf-

Opacities.



sugar, are by some coarse practitioners blown into the eye for this purpose. The principle of their operation is the same. In the use of applications to remove opacities, the points of importance to be determined are the time and the frequency of their use. They are mischievous when inflammation is excited or increased by them; their effects as excitants should be temporary. The mercurial ointments are less effective, in my experience, than the injections. Where the internal use of mercury is indicated by the character and duration of the inflammation which has given rise to opacity, its effect upon the latter is more marked than that of any local remedy.\*

\* The extent to which opacities of the cornea are removable in very young patients is truly surprising, and should always prevent an unfavourable prognosis in such cases. The purulent ophthalmia of new-born infants often leaves the cornea universally and densely opaque; and an inexperienced observer seeing the organ in this state, would pronounce its usefulness irremediably destroyed: while in these very cases, the lapse of a few months, and with no other remedy than a weak solution of nitrate of silver, will exhibit the cornea entirely restored to its transparency and the vision perfect.

In the treatment of opacities of the cornea generally, it must never be forgotten, that until all inflammation be removed, stimulant applications can only do harm. Nor are such applications always necessary. The means which are calculated to subdue the inflammation, and of these mercury is one of the principal, will, in many cases, remove the opacity, if perseveringly employed; while the incautious and precipitate use of a stimulant collyrium will, as often, re-induce in-

Under the head of strumous nebula with vessels overshooting the cornea, I have advised ptyalism, upon the strength of several decided proofs of its efficacy; but no form of recent opacity is so intractable; and I should be uncandid not to state that I have seen it increase from day to day under the mercurial action. The vessels which shoot in radii upon the cornea and at length meet in the centre of the membrane, if the disease is unchecked, are situated beneath the conjunctiva and belong to the sclerotica, as may be easily ascertained by close inspection; and the uncontrollable nature of sclerotic inflammation, of which I shall speak presently, is well known to those who have seen much of it. The deposition is interstitial. The oxymuriate or the hydr. cum cretâ, in small but frequent doses, will sometimes succeed better in this case than the other forms of mercury; and the combination of blue pill or calomel with antimony, better than that with opium.

Strumous  
nebula of the  
cornea.

In constitutions which discover an insusceptibility to be affected by the mercurial pill, or in which its exhibition in sufficient doses is attended with inflammation, and extend the opacity. Such collyria are indeed extremely useful in their proper place, but even more injurious if employed too soon.

Slight scarifications of the corneal conjunctiva covering opacities, assist materially in removing them, if made use of after the complete subsidence of inflammation.—*Editor.*



ed with griping pain and diarrhœa, friction should be employed; and indeed in all cases in which the saving of time and strength is peculiarly an object, this is the more certain and efficacious proceeding. I know that the prejudice often existing against the use of the remedy in any shape, is most strongly opposed to this, its best form. But among persons otherwise intelligent, such a prejudice soon gives way to the more rational feeling of confidence in the practitioner. It is necessary, however, that he should support this feeling by a proper confidence in himself—by a steady perseverance in his design: having, therefore, upon mature deliberation decided, that the mercurial action should be set up, nothing but the clearest demonstration of the patient's inability to support it should interfere with the full and fair execution of the plan. A character notoriously abused by indiscriminate excess, is in much danger of being further injured by half measures. This, I think, has been the case of mercury. It is not the most delicate frame which is most ready to admit, or least able to support it; and it is not the quantity consumed, but the quantity absorbed, which is to be taken into account by the practitioner. The progress of disease during its exhibition is no argument against its continued employment; in this view, unless the system be fairly under its influence, all that has been given goes for nothing; nay, I have had occasion to see many cases in which, after all the

signs of absorption were manifest, its operation upon the disease was for a time unobserved, or was null, and was yet ultimately all that could be wished. I venture upon these remarks from having myself felt "afraid to go forward lest I should go wrong," in some very obstinate cases of strumous nebula in young and very delicate subjects, the issue of which gave me no reason to regret that my confidence had triumphed over my fears. And therefore the stationary condition, nay, the natural progress of a disease during the period occupied by the introduction of mercury, or even after its introduction, for a time to be limited, would not deter me from prosecuting it in a case wherein I placed my dependence upon its power; but an alarming degree of arterial excitement, or certain morbid appearances of the organ, not looked for in the natural and ordinary progress of the disease, would, as a matter of course, determine me to withhold it. These remarks are not confined to the case under notice; they are of general application.

I shall take this opportunity of briefly advertising to another point of the ordinary treatment of this case, of some importance. The division of the conjunctival vessels on the decline of the inflammation is injurious during the acute stage of the disease, and at any period its effect upon the vessels, by which the nebula is secreted and maintained, is from their situation indirect; so



that it stands upon the same ground as scarification, and no other.

Staphyloma.

The staphyloma, if purely corneal, and of such size as to occasion deformity, and expose the organ to further injury, or if producing habitual irritation and inflammation of the tarsal borders, should be excised; the ligature passed through and including two-thirds of the diseased cornea, by means of a curved needle, assists the operator, by steadying the globe. If the staphyloma is from dilatation, the iris will be left; if from breach, it is compacted, and removed with the cornea. This circumstance makes no material difference in the healing, unless the section be made much posterior to the ciliary ring, when the globe collapses from the escape of the vitreous humor; which is not the case when the section is at the base of the cornea, although the iris should be included in it, for the vitreous humor is in a considerable degree sunk by absorption in the staphyloma, and the aqueous as much superabundant. A flat double-edged knife is the most convenient instrument for a circumcision of two-thirds of the staphyloma, which is executed in its passage across the globe; the remaining portion may be finished by one stroke of the scissars. A compress of soft linen should be laid upon the closed lids, and retained by a roller. Where the staphyloma is partial and conical, the

section is corneal, and its edges should be touched with the argentum nitratum to prevent a corneal fistula. But when, as in many cases, the protrusion is not such as to prevent the easy motion of the lids, or occasion inconvenience, it should be left, screened or not, at the option of the patient. No benefit results from tapping the globe of the aqueous humor, either in this case or in the hydrops oculi.

Fungous tumors of the cornea must be treated as the staphyloma. They are of very rare occurrence.

The discharge of the aqueous humor is useless in this case, and all attempts to remove the disease have hitherto proved ineffectual. I have found repeated blisters, and the more powerful tonics, as steel or arsenic, decidedly serviceable. To these may be added, cold bathing, and the practice of often opening the eyes in cold spring water. I am unable to say, whether a section of the cornea, as in extraction, would be productive of benefit. It has occurred to me as not improbable. The disease, however, is constitutional, and must be so treated. The tubular spectacle frame with a pupillar aperture, affords more aid in correcting the vision, than any form of lens.\*

Conical  
cornea.

\* A young man from Rhode-Island applied at the New-York Eye Infirmary about eighteen months since, with both corneæ conical and a cataract in one eye. His near-sighted-



ness was so extreme, that he could read with great difficulty. The cataract was removed from the one eye, and with much advantage to his vision. He could see without any glass much more distinctly with the eye operated upon, than with the other assisted by a concave glass of any degree.

This case tends somewhat to prove the propriety of the plan recommended by some surgeons of removing the lens in cases of conical cornea, in order to compensate for the increased refracting power of the cornea. I have had no opportunity, however, of ascertaining the degree of permanency of the improved vision in this case, as the patient immediately returned home, and I have learned nothing from him since.—*Editor.*

## SECTION II.

---

### SCLEROTITIS, CHOROIDITIS, AND IRITIS.

THE signs of inflammation extending to the sclerotica have been described. It is seldom, if ever, the cornea being the seat of diseased actions, that the sclerotica does not participate. Inflammation cannot pass from the surface of the eye to the interior tunics, without involving this membrane, and the impediment, which is happily opposed to its progress, the slowness with which it is in consequence propagated, is accounted for by the texture and properties of the sclerotica, and the minuteness of the vascular communication through its medium between the conjunctiva and the choroid. The structure and properties of the sclerotica also explain why the primary scleratitis is a rare disease. In the commencement of this disease, the cornea is slightly, if at all clouded, and the activity of the iris is but little impaired. An obtuse pain in the eyeball is materially relieved by blood-letting, and by antimony and ipecacuanha with opiates.\* It is by no means so decidedly

\* The pain attendant upon these forms of inflammation, which generally attacks in paroxysms at regular hours, may



influenced by mercury as the iritis, and its obstinacy and disposition to relapse render the case often difficult of treatment. The subject of it is usually reduced and irritable in a high degree, from suffering with rheumatic inflammation in the elbow, knee, or ankle joints.\* I have generally observed that the previous use of mercury has more or less contributed to this state. I have also noticed the frequent accompaniment of gonorrheal inflammation with this disease, or its existence a short time previous. Though it is necessary to use mercury with more reserve than in other forms of inflammation, to suspend its operation at intervals, and allow the system to recover from its immediate effects, yet its exhibition will be found in the majority of cases, indispensable. The rude and profuse employment of it hurries on the disease, and the extension of the inflammation to the interior tunics ultimately destroys the organ. The nitric acid may often be exhibited with marked benefit, in the intervals of the mercurial action.

The Plummer's pill, the oxymuriate in doses of one-twelfth to one-eighth of a grain, or in feeble subjects, the hydr. cum cretâ, five grains

be materially alleviated by rubbing upon the eyebrow and forehead, a short time before the expected paroxysm, an ointment composed of strong mercurial ointment and powdered opium, in the proportion of six grains of the former to two of the latter.—*Editor.*

\* See note K.

to ten, twice or thrice a day, are most available and beneficial forms of the remedy in these cases. As auxiliaries, soothing and allaying irritation, I should mention the Dover's powder, hemlock, and hyoscyamus, and the extract of sarsaparilla, either dissolved in the decoction or taken freely in the solid form. I have seen an obstinate chronic inflammation yield before these latter remedies, in which mercury had been productive of no benefit. In the motley diseases now known by the cant term of pseudo-syphilis, their efficacy is admitted by the most competent authorities.

On the treatment of deep-seated inflammation whether affecting the choroid or iris, I shall not now dwell, having in an essay on this subject, published three years ago, pretty fully stated my opinion; and when treating of the signs of these diseases in the present volume, having repeated my conviction of the remarkable efficacy of mercury, and of the comparative insignificance of every other remedy. One full blood-letting or more should be premised in the acute stage of the disease; and topical blood-lettings are generally required at short intervals during its exhibition. I have now and then found that the incipient inflammation, where it has extended from the conjunctiva, yields to a copious venesection, and two or three brisk doses of calomel and rhubarb, followed up by the infusion of senna; but, generally speaking, the system must



be made to feel the influence of mercury before the disease is permanently subdued. The inflammation which has proceeded to the effusion of adhesive matter, never, in my experience, yields either to the lancet, continued nausea, or full purging ; and it is remarkable that the cases presenting this termination of inflammation are always most sensibly and immediately benefited by the remedy in question, whether the cornea or the iris be affected, or any other texture of the body.

That in many instances, however, the deposition takes place notwithstanding, or immediately succeeding to the action of mercury, and is most prone to do so (I do not say from that cause), I am as sure, as that it seldom fails to yield to its continuance or renewal. But when the mercury arrests inflammation previous to this event of it, there is reason to infer that it prevents such termination, and its less rapid and decided influence under these circumstances, is not a reason why, if the inflammation resists the ordinary antiphlogistic measures, it should not be employed. I believe that the mode of action of the remedy varies according to the degree of its influence, which again varies according to the habit of the patient, the form or stage of the disease, and the quantity of the remedy which is received into the system. But if any two facts are well established in modern medicine,

I apprehend they are these :—first, the power of mercury to arrest acute membranous inflammation, both prior to and after the effusion of adhesive matter; and second, its power rapidly to remove, by an excitement of the absorbing system peculiar to itself, the newly effused adhesive matter. If these facts are admitted, then the propriety of its use is indicated in iritis, as in carditis, pleuritis, peritonitis, and the only practical question that can arise respecting it is, how far the patient's strength is equal to support the remedy. There are, I admit, states of the organ as well as of the constitution, in which it cannot be borne, and no sooner is its influence felt, than the inflammation threatens disorganization, and if the plan is persevered in, quickly runs on to it. The globe becomes enlarged or mis-shapen, the sclerotica assumes a livid hue, and the veins a state of varicose congestion; sometimes the eye-ball suppurates, and the little remaining vision is completely extinguished. In cases where age, or the existence of other diseases, or the already excessive use of mercury, has greatly enfeebled the powers of the system, it must be used, if ventured upon at all, very sparingly, or with intermissions, and the system must be supported by every admissible means, both of nourishment and medicine, during its employment.



### SECTION III.

---

#### AMAUROSIS.

THE term amaurosis comprehends all those imperfections of vision which depend upon a morbid condition, whether affecting structure or function, of the sentient apparatus proper to this organ. That the term is not so defined according to its etymological import is well known, but it is thus employed by pathologists, if I have rightly understood its meaning.

Organic  
amaurosis  
from inflam-  
mation.

The diseases of the other coats and humors of the eye which are present in a considerable number of these cases, are effects of an inflammation which has destroyed the retina. Such are especially, discoloration and absorption of the vitreous humor, or a bright yellow opacity of the crystalline lens, which is indurated—its capsule condensed with it, and firmly adhering to the constricted and perhaps irregular pupil, with peduncles of lymph or detached flakes of the black pigment projecting from its posterior border—or a capsule containing calcareous concretions with an absorbed lens, and a concave and tremulous iris, or an obliterated pupil, or a staphyloma of the sclerotica or choroid.

But there are cases in which a change in the structure of the retina is to be inferred, of a description less conspicuous indeed, but not less fatal to vision. This is the result of a slow and insidious morbid action, and although sometimes accompanied with superficial inflammation, is more frequently altogether independent of inflammation. The congestion of the superficial vessels, the extenuation and consequent blue tint of the sclerotica, the appearances supposed to indicate a caligo of the vitreous humor, or an opacity of the retina, or a deficiency of the pigmentum nigrum, seen upon looking towards the fundus of the eye, are signs of this change.

From a change of texture independent of inflammation.

When the eyeball has the appearance of health, and the loss of vivacity in the motions of the pupil, is the only sign of an amaurosis obtained from inspection of the organ, we are scarcely warranted to suppose any disease of structure. I have called such cases functional, and my object in doing so is to discriminate them from the organic, in the belief that much practical advantage may be gained from the distinction. Diagnosis is a study interesting in a scientific view, but it is awfully important as it affects practice and character. An amaurosis depending on a change of structure in the brain or eyeball, is an irremediable case. The same may too often be said of that which presents no evidence of structural disease, of which

Functional amaurosis.



I shall presently mention examples. It is creditable to a practitioner to know such cases; and if his ingenuousness is equal to his knowledge, he will be a gainer in reputation in every way.

Functional amaurosis I have arranged under three heads: the symptomatic, the metastatic, and the proper. The first includes a class of diseases so large and diversified, that to consider them and the rationale of their treatment in detail, would occupy a volume. Suffice it therefore to say, that the amaurosis being subservient to the disease which affects the system at large, or some one important organ, the latter is the proper object of medical treatment. I may instance the morbid states and actions of the vascular system, the disorder of the digestive organs in its several degrees from impaired appetite to confirmed hypochondriasis, the interruption to the healthy functions of the uterus, the excess, or deficiency, or accumulation of the wonted secretions and excretions, the presence of local irritation, as wounds and abscesses, caries, worms, &c., and the influence of strong mental emotion producing a morbid irritability. These co-existing with an amaurosis must be regarded as the original and substantive disease, the removal of which is the aim and end of treatment. It must be obvious that it is rather the degree, than the nature and origin of the

functional disease, that should in most cases influence our prognosis, yet the latter circumstances, it is equally clear, afford more or less encouragement, in proportion as the pre-existing states of diseases ordinarily admit of relief or otherwise. Thus, for the sake of illustration, I may observe, that the amaurosis from gastric diseases, from plethora, from irritation, are all of them relievable, and, if treated at an early period, remediable. Whereas paralysis, the sequel of fever or of epilepsy,\* or severe constitutional diseases, whether acute or chronic, or depending upon habitual cerebral congestion combined with organic visceral disease, or induced by the operation of noxious agents on the system, is a hopeless form of the malady. It resembles in appearance and character the ordinary gutta serena, or idiopathic palsy of the retina, which occurs in early as often as in advanced life, in which, excepting the gaping and motionless pupil, and the absence of physiognomical expression, no defect appears; on the contrary, the fine, large, well-opened, and singularly brilliant eye, often excites admiration of its beauty as an organ, though unilluminated by the mind.

\* I know a family of several well-formed children, three of whom have dark hair and eyes, the others light hair and blue eyes. Towards puberty, all the dark-haired children have become epileptics, and gradually lost their sight; the eyes, except in the expansion and immobility of the pupils, retaining every appearance of health.



The metastatic amaurosis is rare but well marked. The restoration of the original malady, if it be practicable without involving the patient's safety, or the substitution of an artificial excitement or discharge, which may serve as an equivalent, appears to be the natural indication, and such a practice has been attended with success.\* But the prognosis is necessarily one of great uncertainty.

The proper functional amaurosis presents great variety; but if treated at an early period is very often cured. The extreme states of light and temperature, and the over-exertion of the organ, are the chief causes of it. The remission or removal of these hurtful circumstances even of itself does much towards the cure. The continuance of them frustrates the end of treatment, and the amaurosis becomes confirmed, and ultimately passes into the organic form. The hemeralopia,† and many other cases not assuming

\* See a remarkable case of "Amaurosis from suppressed purulent discharge," successfully treated by Professor Beer, in the "Analecta" of the "Quarterly Journal of Foreign Medicine and Surgery, No. IV."

Although the metastasis of gout, of which I have known two marked instances, has been fatal to vision; yet in three cases, in which I extracted the cataract from gouty subjects, and a smart attack of the disease followed the operation, the eyes were unaffected and the sight was well recovered.

† See an excellent paper on this disease as it affects seamen in tropical climates, by Mr. R. W. Bampffield, Surgeon of the Royal Navy, in the 5th Vol. of the Medico-Chirurgical Transactions.

this precise character, are essentially depending on the injurious influence of the extremes of temperature, and light, and intense colors. I call to mind several cases distinctly referrible to each of the above-named causes. During the correction of this sheet I have been consulted for an amaurosis immediately succeeding to exposure, during several hours, to cold and incessant rain; and which I have the pleasure to say is advancing towards recovery. I have exemplified the treatment of these cases in the pathology;\* if active measures are taken without delay, they generally warrant a favourable prognosis, but only on this condition.

The functional amaurosis varies in its rate of progress as well as in its ultimate extent. Some are sudden in their accession and perfect, as many instances of the metastatic and the proper; others advance steadily but sensibly to a point little short of blindness, at which they begin to assume an organic character; and of others again the actual progress is scarcely perceptible for months in succession, fluctuating from day to day between better and worse. I should say that the slow and the steadily progressive amaurosis is more to be apprehended in the result, that is, is less tractable than either the sudden or the rapidly advancing disease, supposing all to be alike free from the unequivocal signs of organic change.

\* Page 169 et seq.



The removal of an irritating or oppressing cause will often effect a sudden and marked relief, as by clearing the intestinal canal of vitiated secretions therein accumulated, by restoring the digestive functions labouring under manifest derangement, or by taking away blood where the necessity is indicated. I have seen an incipient amaurosis directly arrested by the extraction of a diseased tooth, when the delay of a similar operation had occasioned gutta serena on the opposite side two years before.

The floating muscæ, when the disease is once established, are seldom if ever removed, yet patients retain good sight who have been troubled with them during half their lives. When the mind becomes indifferent about them, they are no longer observed, except in states of anxiety, irritation, or bodily weakness; and the subjects of them are usually aware of this fact.

It appears then upon this representation that certain cases purely functional, whether the affection originates in the organ, or in a remote part of the system, are, in their character, so nearly approaching to the organic class, as at once to convey the impression of their irremediable nature. In common with these they have many symptoms, hence the difficulty of diagnosis; and some of them quickly tend to altered structure, although the external signs of this

change are faintly indicated. On the other hand, cases are of frequent occurrence, more frequent than has been generally supposed, which admit of material and decided improvement, and even of complete recovery ; and I would repeat, with the exception of those above adverted to, that it is rather the degree than the nature and origin of the functional disease that should influence our prognosis.

The treatment of amaurosis is almost exclusively constitutional. To the various forms of external remedies, such as stimulant vapours, drops, and ointments ; spirituous, ethereal, and aromatic embrocations ; sternutatories, &c. &c. my experience leads me to attach no value. The faith yielded to such applications is a relic of the not very remote superstition, which ascribed miraculous powers to the hand of a living king, or a dead culprit. Ophthalmic surgery has been more degraded by manual officiousness, and the confidence placed in externals has been more injurious to its improvement, than to that of any other branch of the profession. It is fortunately not now necessary to do something when nothing can avail. I am quite aware that the transient effect of stimuli is in many of these cases grateful, and seems partially to remove the obscurity of vision, but the patient soon discovers that this is but a fillip. When, indeed, a disordered state of the conjunctiva and

External  
remedies.



Blisters.

eyelids exists in conjunction with an imperfect amaurosis, a more permanent benefit is often obtained by the rectification of this state, sufficient not only to afford encouragement in the use of topical remedies, but to induce a belief that the affection of the retina is, in a degree at least, sympathetic with that of the surface.\* I should make a reserve of cupping, issues, or setons, in certain cases which it is unnecessary to specify, and of blisters in almost all. These, if managed as the case directs, are a remedy of great value: in some, as temporary irritants only; in others, as irritants and drains. With the former view either the vesicle should be preserved by simply puncturing it, or the cuticle should be removed on dressing the blister, and the simple ointment applied. The process should be frequently repeated, and alternately over the superciliary ridge, upon the temple, upon the mastoid process, or the nape of the neck, as most eligible. Or if a more extended surface of irritation is desired in the immediate vicinity of the eye, the blister should take the shape of a chemist's retort, reaching from the zygoma to the glabella. It should be borne in mind, that the operation of blisters is very different in different individuals as regards their susceptibility. The irritation and discharge of an efficient blister, as

\* See page 165.

big as a crown piece, will sink the powers of a delicate female for days, and this effect will be especially felt in a weak retina. I have often known the obscurity of vision decidedly increased for a time by the application. Such cases are yet more affected by the direct loss of blood, even in the smallest quantity, and the permanent blister would be injurious. How is it that a blister is as useful in a proper nervous or paralytic amaurosis, as in one depending upon the congestion of the blood-vessels? I have been asked this question by intelligent persons. The fact is unquestionable and the answer obvious, that the simple and temporary irritation is the object in one case, and a permanent irritation and derivation of blood in the other; and the blister is to be managed accordingly. Hence in a very susceptible subject, or a very irritable skin, a mustard plaster applied for ten minutes, and repeated now and then, may answer the first purpose more conveniently. It is in this particular view that the moxa is used in this and other diseases with so much advantage on the Continent. The eschar, if left to itself and not converted into an issue, is superficial, but the irritation is of the severest kind.

I have heard and read of the effects of elec-

Electricity.

tricity and galvanism in amaurosis. Some nar-



edly entitled to credit. I have had recourse to them in many cases, some of a very favourable description, but have never witnessed a single instance of benefit arising from the application of these powers.

General  
treatment.

The degree of constitutional power which enters into the disease forms the first and most important question in the general treatment. Extremes, it is said, meet, and it is certain that a strong and delusive similarity often prevails between the signs of diseases, which result from conditions diametrically opposite. The treatment in cases of general plethora and of cerebral compression I need not point out. But I have mentioned cases of undue determination of blood to the organ, which are especially common after deep-seated chronic inflammation, or distress from over excitement, by which its vessels have lost their tone; an effect decidedly increased by depletion. Such cases are not difficult of discrimination from the former, if a due attention is given to the history.

All the cases of direct debility and proper paralysis of the retina are aggravated by loss of blood, and the great prevailing mistake in the treatment of amaurosis is the indiscriminate detraction of blood. The same observation, it appears to me, applies to the treatment of cases of general palsy. The practical idea of compres-

sion derived from the demonstrated instance of apoplexy, prevails over and puts aside the theoretic idea derived from the admitted condition of nervous debility or exhaustion.

I have never known any real benefit derived from what are called antispasmodic and antinervous medicines, camphor, assafœtida, valerian, &c. Neither do I recollect an instance of decided benefit from the emetic practice, although in respect to high authority, I have tried it fairly in many instances. The cases of gastric disorder to which it is especially applicable, are most benefited by a long continued course of the blue pill, with gentle saline purgatives and tonic bitters.

In most of these cases we must depend, first, on the regulation of the visceral functions; and secondly, on the employment of such restoratives as the system requires and can bear. The blue pill, with colocynth, rhubarb, or aloes, and the combination of soda with rhubarb and calumba or gentian, are best adapted to the former purpose. The exhibition of general tonics is often strongly indicated, and I have seen much benefit derived from the mineral acids, bark, steel, when admissible, and arsenic, after a due regulation of the digestive functions. I know of no article of the class of stimulants that has any direct claim to notice, or any approach to a specific



virtue, such as has been ascribed to the *arnica montana*, *aconite*, &c.\*

Mercury.

When the amaurosis is recent and sudden, and either the signs of an obscure inflammation are present, or only the amplitude and inactivity of the pupil correspond to the patient's history, the indication is less simple ; mercury should be introduced with all convenient rapidity into the system, I mean so as to ruffle it in the least possible degree. No advantage is obtained by salivation; on the contrary, I think it hurtful; when mercury is beneficial, its efficacy is perceived as soon as the mouth is sore.† I have seen it tried, and have myself tried it in many cases of perfect amaurosis, without the smallest advantage; but in cases of recent occurrence, imperfect, but rapidly progressive from bad to worse, I have been witness to its power in suddenly arresting the disease in too many instances, not to entertain a far higher opinion of it than of any other article of the *materia medica*. I shall not again discuss the knotty question of its *modus operandi*; "*causa latet: vis est notissima*." The form of its administration must be regulated by the circumstances of the case.

\* Some oculists still adhere to the practice of the archangel, and

— "purge with *euphrasy* and *rue*

"The visual nerve."

† See note L.

Superadded to the entire repose of the organ, Dietetic.  
the natural tonics, viz. a pure, dry atmosphere,  
the cold bath, horse exercise, nutritious diet,  
early and sufficient rest, agreeable society, and  
a mind as much as possible diverted from the  
object upon which it is unfortunately and perti-  
naciously prone to dwell—these are of greater  
avail than drugs; and some lighter forms of  
sympathetic amaurosis are as effectually cured  
by them as by the blue pill and rhubarb, and  
upon the self-same principle.



## SECTION IV.

---

### DISEASES AFFECTING THE EYEBALL.

Suppuration  
of the ball.

THE treatment of those injuries to which the organ is subject from external violence, is comprehended in the directions given for the treatment of inflammation and its consequences. In the suppuration of the ball, when the patient's suffering is acute and the constitutional irritation severe, and the part notwithstanding its great tension affords no immediate prospect of relief by a natural opening, the cornea, iris, ciliary ring, and some extent of the sclerotica should be so divided by a deep transverse incision, as to evacuate the globe of its contents. In this manner about a tea-spoonful of pus, more or less, is discharged. The section of the cornea alone, effects this object so imperfectly as to afford little if any relief. The eye should afterwards be lightly covered with a soft poultice confined in a cambric bag.

Extirpation  
of the ball.

The extirpation of the eye, when that operation is determined upon, is most conveniently performed with a straight double-edged knife,

which is to be employed for the purpose of freely dividing the septum of the conjunctiva and oblique muscles, so as to separate the globe and lacrymal gland from the palpebræ and base of the orbit. When this is done, the globe admits of being drawn gently forward by a ligature previously passed through its anterior segment. A double-edged knife, curved breadth-wise, should then be introduced at the temporal commissure of the lids, for the purpose of dividing the muscles, vessels, and nerves, by which the globe remains attached, with greater convenience and despatch. The hemorrhage is repressed by means of a small portion of fine sponge introduced into the orbit, and a light compress of linen should then be laid upon the lid supported by a roller. The sponge should not be suffered to remain longer than the following day, when a soft poultice in a muslin bag may be substituted for the compress. An opiate should be given at bed-time.

The practice of cramming the orbit with lint or charpie, and leaving it to be discharged by suppuration, is objectionable. I knew one case in which this measure was followed by a most extensive suppuration within the cavity, and by abscesses in the neighbouring integument of the lids and forehead; and another has been communicated to me, in which its ill effects were evident.



I once lost a patient, a middle-aged countryman, otherwise in health, within a fortnight after this operation, owing to a suppuration of the dura mater, on the same side of the head. The attack of inflammation was sudden and rapid, commencing about a week after the operation, and ushered in by a severe rigor after exposure to cold, in the square of the Hospital; an imprudence quite unauthorized. There was no continuity of inflamed surface to account for this, although the morbid appearances were confined to the membranes of the corresponding hemisphere. I have performed the operation many times without any serious after-symptom.

Glandular  
enlarge-  
ments and  
tubercles.

The propriety of this measure, from its severity and the uncertainty of its preserving life, should be always matter of very deliberate consultation. I will only observe, that if but one eye is affected with the disease, and the patient free from any material disorder of health, we should be slow to reject the operation on account of glandular enlargements in the vicinity of the orbit, or of tubercles of a suspicious character in other parts of the body. Such affections, supposed to be of the same morbid character, have disappeared in more than one case of malignant fungus, after the removal of the diseased part.

A gentleman, whose case, a fungoid tumor involving the knee-joint, was considered desperate, and was absolutely abandoned as hopeless by the ablest surgeons in this town, on account of many tubercular swellings on other parts of his body, in addition to extreme weakness and emaciation, put himself under the care of an older practitioner, who considered the objections theoretical; the limb was amputated by this gentleman, and the patient recovered; all the tumors subsiding as he regained his health.

The recent enlargement without induration of one or more lymphatic glands in the track of absorption, in cases of scirrhus of the mamma, or the testis, or the lower lip, has not unfrequently proved to be the result of simple irritation, by speedily subsiding after the healing of the wound. I have known cases in which swellings of a more suspicious nature remained stationary after the extirpation of a malignant fungus, and the patient has sunk under visceral disease of a character totally dissimilar. It is right however to state, that the period of survival has been short in these cases. An extensive abdominal abscess proved destructive to a man whose leg had been amputated three months before for a genuine fungus hæmatodes; another died suddenly of apoplexy, several months after the removal of a fungoid testicle of great size. In both



these cases the glandular swellings of the inguen and iliac region of the same side had subsided in the interval.

Admitting therefore the malignant character of a disease, if the symptoms do not indicate the affection of vital organs, it is possible that the removal of the principal source of irritation, may admit of a change in favour of the constitution, and put a stop to its progress. We know not how far the multiplied production of tubercles may be a result of sympathetic irritation, and when doubts of this description arise, cases like that above mentioned should be borne in mind, that the patient's chance of recovery may not be forfeited by indecision.

**TREATMENT**  
**OF THE**  
**DISEASES OF THE EYE.**

---

**CHAPTER III.**

---

**SECTION I.**

**ON THE OPERATIONS FOR THE  
CATARACT.**

BEFORE speaking of the operations, I shall offer a few preliminary considerations.

The extent and importance of the subjects of this and the following chapter, the deep professional interest which they have of late years excited, and the copiousness of the historical details connected with them, render it impossible for me to go minutely into them, consistently with the plan of the present work. Such an undertaking, if it were not superfluous, would of itself furnish materials for a volume of ordinary size; but so much has been ably written upon



the subject, that the inquiring student can be at no loss for all requisite elementary information. This premised, I shall take the liberty of confining myself to those general results of my personal observation, which appear to me worthy of communication.

Cataract in  
one eye,

It has been a custom with oculists where a person has a full formed cataract in one eye, and retains the vision of the other, to advise the postponement of the operation until that also is dark—this advice I think erroneous. I am satisfied that the cataractous eye, if it become the subject of an accidental inflammation, is strongly disposed to go into amaurosis; and further, that the retina loses its vigor by the permanent exclusion of light. I speak from repeated observation of the fact. The objection to the operation on the ground of inconvenience arising from the difference of focus of the two eyes, when one only is the subject of the disease, is trivial, and a consideration altogether subordinate; such a defect may always be remedied by glasses properly adjusted. In several cases of amaurosis ensuing upon cataract, I have been disposed to regard the change in consistence and volume of the lens, as productive of a destroying inflammation, in others of a partial absorption of the vitreous humor.

with amaurosis;

The cataractous eye is not unfrequently amauro-

rotic; nor is it always possible to determine the presence of amaurosis, when the opaque lens is so dense as to account for complete darkness.

It often happens that a patient has a full formed cataract in one eye which presents the signs of amaurosis, and an incipient cataract, or one as much advanced in the other which is at present free from these symptoms. In this case the cataract of the latter should be removed without delay.

In cases of congenital or infantile cataract, which become the subject of operation at an adult age, the lens is found to have undergone absorption, and the capsule alone remains. This absorption does not take place, except in cases of injury, in the cataract formed during adult life. When the lens has early undergone a natural absorption, and the cataract is simply capsular, the organ is always imperfect; the operation therefore seldom increases the distinctness although it may enlarge the field of sight. The fluid cataract of the adult is always joined to a weak, often an insensible retina.

with absorbed lens.

It would be incorrect to say that the operation was unadvisable in all cases of cataract in which the patient has no sense of light, for it is possible that the density of the lens may be

Degree of vision.



such as absolutely to exclude the light, and that the motions of the iris may be therefore suspended, or from some degree of pressure of the lens, or adhesion of the uvea to the capsule, that the pupil may be undilated, and the circumference of the lens permanently covered. But undoubtedly a case of this description is unpromising. A strong sense of light by which at least to know the direction in which it enters the apartment—to be sensible of its falling on the eye, and of a shade, as the hand, for example, intercepting it, with a corresponding freedom of motion in the pupil, is the most favourable state for the operation. There is in this case perception enough to determine the sensibility of the retina, and not enough to occasion the unsteadiness of the globe. If a patient has vision, the eye is irritable to light, and involuntarily rolls as far as possible towards the nose on the introduction of the instrument, one of the greatest perplexities in the operation. Another reason why an operator may naturally prefer an eye in which vision is interrupted, is this : patients are practical and not speculative philosophers, and estimate operations by the amount of the good conferred, not by the magnitude of the evil averted. A blind person restored to sight, is thus gratified in a much greater degree than one whose partially obscured vision is rendered clear. These however are not reasons for deferring the operation

beyond the period at which useful vision ceases, nor do I think any reason can be given for delay, tantamount to the risk of injury from incidental inflammation, or impotence from a continued suspension of the function of the organ.

There are several circumstances in the structure and condition of the organ which have an important influence on the facility and success of operations.\* These are, shortly, the degree of projection of the orbit, or the relative volume of the eyeball and its socket; the absolute size, prominence, and tension of the ball; the dimensions of the anterior chamber; the clear and healthy, or thickened and partially opaque state of the cornea; the absence or presence and encroachment of the arcus senilis; the proportional diameter of the cornea to the globe, and its actual diameter; the healthy state and proportion of the aqueous and vitreous humors; the free and active, or adhering and constricted state of the pupil. A small cornea, a narrow or partially obliterated chamber, and a contracted pupil, are circumstances decidedly unfavorable to any and all operations.

Local circumstances.

The operations have been so repeatedly and

Coupling.

\* To these might be added circumstances of temperament, viz. a calm and steady, or an irritable and very moveable eye.



minutely described, that I shall not fatigue my reader with a prolix detail of them. They are three in number : couching or depression, extraction, and absorption. The first and most ancient is now seldom performed in this country. The cases to which it is alone applicable are those cataracts of firm consistence, the circumstances of which offer a serious impediment to the much more eligible method of extraction. These circumstances may be inferred from the enumeration above given. The needle employed by Scarpa is best adapted to the purpose.\* The couching-needle may be passed through the sclerotica at a line's breadth from the cornea, and a little below the horizontal diameter, so as to avoid the long ciliary artery; or through the inferior part of the cornea and pupil; and the lens may be depressed vertically or horizontally. The term 'reclination' has been applied to the latter method. In both cases the lens must be hitched into a breach of the vitreous humor below the border of the pupil. Its anterior capsule, and the capsule of the

\* The needle employed by the Editor is somewhat different from Scarpa's. Instead of a triangular point like his, it has a lancet point, cutting on both edges, flat on one side, convex on the other, and curved at the point like that of the Italian surgeon. A needle of this form enters the sclerotic more readily, makes an incised instead of a punctured wound, and may be more easily made perfectly sharp by the cutler. See Plate 5, Fig. 1.

vitreous humor, must be divided or torn through, to render the operation effective. The lens corresponds in diameter to the iris, and there is therefore no natural space into which it can be depressed. The posterior capsule, identical with that of the vitreous humor, must be lacerated, to admit of its dislocation backwards and downwards; and if its anterior capsule was left entire, it would become a secondary capsular cataract, and require a subsequent operation.

The operation of couching through the cornea has of late years been warmly advocated, as being unattended by the injurious consequences ascribed to the perforation of the sclerotica and choroid. It is reasonable to prefer that operation which inflicts the smallest injury, and to conclude that it is least liable to be followed by severe inflammation. But the subtilties of theory have little weight in the scale against experience, and no one who has seen much of these operations considers the puncture of the tunics to form a material objection, if it is executed with a proper instrument, and agreeably to the directions which are furnished by a knowledge of the organ. I have so often seen the eye perfect in its aspect and function, after several such penetrations of the larger tunics, and the inflammation immediately resulting from each operation so slight and transitory, as to be convinced that the

Kerato-  
nyxis.



objection is either hypothetical, or is drawn from unskilful and rash procedures.

The real objection to couching is the ultimate step of the operation, viz. the breaking up of the fine texture that fills the globe by the forcible depression of the lens. Whether it be depressed edge-ways or breadth-ways, makes no difference in the result; it must still occupy a breach in the cells of the vitreous humor, and must derange and disorder that delicate texture and those connected with it. A slow insidious inflammation marked by a gradual development of the symptoms of disorganization, viz. congestion of vessels, turbid humors, flaccid tunics, and palsied iris, is too often the consequence. The sight, instead of improving when the immediate effects of the injury are passed away, remains habitually weak and dim, or declines and fades altogether. The advocates for reclination seem to forget that the principle, which is the same in both operations, is the real ground of objection. As to the position of the lens, I suspect less mischief is done by the old method of depression, as less force is required to break a space for the vertical than the horizontal lens, provided the depression be carried to no greater extent than is necessary to clear the inferior border of the pupil. After all, the argument is mere trifling about the position of the lens, absolute or relative; it can

occupy no place but that made for it without serious injury to the organ. It is not fair to bring against any operation objections that apply only to the unskilful performance of it; and this, it is easy to see, is the secret of the frightful catalogue of disasters which the spirit of controversy promulgates, and which those who practise these operations declare to be totally unauthorized in their experience. For example, what has the wound of the retina, of the ciliary body and processes, of the iris, &c. to do with the operation of couching properly performed? Yet all these are marshalled in formidable array as objections, for the obvious purpose of demonstrating the superior advantages of couching through the cornea.

If the operation just described is attempted upon a flocculent cataract, the lens, instead of descending solidly, breaks into pieces, which undergo a gradual absorption; the perfection of which however consists in making the free central aperture by laceration of the anterior capsule, the preliminary step. The needle is introduced either through the cornea or the sclerotica. In most instances the anterior operation is to be preferred, as I have elsewhere stated. The more minutely the lens is broken up and divided in its texture, and the more its fragments are dissipated in the anterior chamber, the quicker the progress of absorption, and the softer the tex-

Absorption.



ture of the lens, the more readily and safely is this object accomplished. If the substance of the lens is dense and compact, this division is not accomplished without considerable force, and the inflammation which follows is hazardous; if the fragments are bulky and press upon the iris, such a result is still more to be apprehended. If therefore this operation is resorted to in a case of firm cataract, it must of necessity be several times repeated if we would preserve the organ uninjured, and this forms an insuperable objection to it in all such cases.\* I would add, that even the utmost caution is inadequate to prevent an internal inflammation from the bulging of the lens after its capsule is freely rent, or its separation, during the absorbing process, into fragments of such a size as to oppress the iris. On this account an amaurosis is not unfrequently the result of this operation, although the slower and milder method is decidedly attended with the least risk. These objections are not applicable to the soft caseous and flocculent cataract, but even in this case the

\* I pass over the description of an operation which consists in the introduction of a knife, whether through the cornea or sclerotica, for the purpose of cutting up the hard crystalline in situ, and throwing the slices into the anterior chamber; and I mention it only by way of caution, if caution be necessary against a measure so desperate and ill-advised. It levels with the proposal to extract through the sclerotica.

cure is often lingering, and subject to be interrupted by inflammation. It is especially to the cataract of infancy that the operation of absorption is applicable. Here indeed there is no alternative, it is fluid, or flocculent; often so far absorbed, that only a thin scale or flake of lenticular substance remains betwixt the capsules, so that its consistence does not allow of depression, and the eye is too unsteady to admit of extraction with safety, if this operation were otherwise as eligible, which in fact it is not. It is impossible to conceive a more simple, sufficient, or gratifying operation than that of Mr. Saunders, if the intention is perfectly executed. I have now enjoyed extensive opportunities of ascertaining its value; having operated, during a period of ten years, upon children of all ages, from four months upwards, and I do not hesitate to affirm that it ranks, in my estimation, as one of the finest discoveries of modern science.

The primary cataract adhering to the iris is for the most part capsular; but whether it be so or not, the needle is best adapted to it. Its toughness and the firmness of its attachment, and the difficulty of couching it, when detached, are circumstances which often render the operation imperfect. The aperture by laceration of the capsule in the centre, and its extension as much as possible by the varied movements of the needle, should be the object of the surgeon,

Cataract  
with adhe-  
sion to the  
iris.



rather than the detachment of the membrane entire. In this, and in all cases in which the needle is used, the employment of the extract of belladonna, softened by the addition of a little water, is a point of the first importance, both prior and subsequent to the operation. The skin above and below the eyebrow should be thickly painted with the solution once or oftener in the twenty-four hours, and this varnish should be preserved moist for a period of half an hour, in order to admit of its absorption. The frequency of the application must be determined by its effect upon the pupil. The preternatural dilatation should not be incessantly maintained; for if it be, the pupil will, in all probability, be misshapen when the application is suspended, and the iris recovers its power.

Extraction.

The operation of extraction is by far the most perfect ever devised for the cure of cataract; but it is one of considerable difficulty, and the several modifications which have been at various times suggested, owe their origin to the disappointments and defeats which operators meet with in learning to execute it with success. The preference entertained for couching rests on no better ground than its greater facility, and therefore less risk. No operation in surgery, I am well satisfied, requires an equal degree of temper and experience for its accurate and successful performance.

The Baron de Wenzel is reported to have said that he had 'spoiled a hat-full of eyes' before he had learned to extract. This was doubtless a figure of speech, but it serves to show the appreciation of its difficulty by a great master of the art. Excellent directions for the operation have been given by Wenzel in his treatise, translated by the late Mr. Ware; and the essay of the latter gentleman, who was in no respect inferior to the Baron as an operator, upon the impediments to the success of the operation, is a work of much merit, and should be diligently studied by all who undertake it. It is objected to this operation, that it is one of which the result is a matter of hazard. I reply, not more so, in the hands of qualified persons, than hernia, lithotomy, aneurism, and other important operations. Secondly, if it fails, it fails beyond recovery. This I contend is rarely the case in the hands of competent persons. That it sometimes is the case I do not deny, nor would I believe that man on his oath who ventured for himself to deny it; but I may be allowed to ask, is not this exception to the general issue of the operation, a condition of every human work? What operation, I should be glad to learn, is not impugnable by such an argument? Thirdly, it is followed by a higher degree of inflammation, and one of a less manageable kind. This again, as the former and all the objections in detail, admit of this general answer. They



apply to the performance, not to the principle of the operation. Thus, if the corneal section be clean, and situated midway between the pupillar edge and the margin of the cornea, or a little nearer to the latter; if it be of such extent as to allow of the perfectly easy escape of the lens; if the sclerotic conjunctiva, sclerotica, and especially the iris be untouched, and the capsule freely lacerated, without læsion of the vitreous capsule, then the operation is perfect. But although all these points should be imperfectly, that is, not strictly fulfilled, yet the result of the operation is ordinarily successful with a due attention on the part of the surgeon. There is not one of them that I have not repeatedly seen reversed, and yet the patient has recovered excellent vision. So that the failure of the operation is by no means a necessary consequence of the casualties that may attend it, even although they should be such as to excite a considerable degree of alarm in the mind of the operator. The deviation of the section from the course intended; the wound, or removal, or prolapsus of a portion of the iris; the escape of a part of the vitreous humor, these, I grant, are derogations from the perfection of the operation. But if the lens be extracted with tolerable facility, such accidents are seldom, with the aid of care and time, permanently injurious to vision. Nevertheless, they are such departures from the fair procedure of the operation, as ought in

common candor to clear it of all imputation from an imperfect result.

The main impediment to the success of this very valuable operation is, as I have elsewhere stated, a section of insufficient magnitude. The easy extraction of a cataract, like the easy extraction of a stone, almost invariably does well, and the difficult and forcible removal of either as certainly augurs unfavorably. The enlargement of the section, if too short, is difficult, and always dangerous to the iris in the collapsed state of the cornea; it is attended, moreover, with imminent risk of a laceration from the want of due support of the vitreous capsule, the loss of a portion of this humor, and the consequent sinking of the lens behind the iris. Thus one difficulty leads on another.

Inadequate  
section.

Again, the protrusion of the iris before the lens in its exit, which only happens from a too narrow section, is almost always followed by the falling of that membrane into the wound, a protracted healing of the wound, and a loss of figure of the cornea.

The free escape of the vitreous humor, owing to an imperfect section, undue pressure, &c. occasions the sinking of the lens in the globe. Such a circumstance may embarrass an operator, and induce him to leave the cataract, in the

Escape of  
vitreous hu-  
mor



hope of its absorption, or of removing it at a future time, rather than hazard the further loss of vitreous humor. This should never be done. For as soon as the wound closes, the cataract is raised by the renewal of the aqueous humor and pressed forward upon the iris. I have seen an inflammation supervene in such a case which speedily went on to suppuration, and destroyed the eye. If, upon making the section, an inordinate quantity of aqueous humor escape, and the lens sinks from this cause, the case is different; here the vitreous humor is already partially absorbed, and the lens is supported by the aqueous. The same ill consequences will not follow in this case, if the surgeon prefers to relinquish the operation. The cornea heals kindly, and he may afterwards operate with the needle. However, in most instances, the lens may be supported by gentle pressure on the inferior part of the globe, and extracted or rather turned out with the hook or spoon end of the curette; and I should always pursue this method under such circumstances, as long as the globe retained its figure.

Section  
verging on  
the sclero-  
tica.

It is a point of considerable importance that the section should be purely corneal. I mean that it should not be carried so low as to verge upon the sclerotica, and thus to leave the corneal margin of an insufficient breadth for

union.\* Two ill consequences arise from this: first, the iris, unsupported at its base, commonly falls into and prolapses at the wound, even though the section be ample in extent, and the escape of the lens perfectly easy; and secondly, from the defect or narrowness of the corneal margin and the non-opposition of homogeneous parts, the healing is always remarkably slow, even though no prolapsus should take place. An oozing of humor is continually occurring, and I have sometimes seen a portion of the capsule, now turned opaque, protrude, and subsequently slough out at the section. When a prolapsus, of whatever kind, prevents the healing of the wound, it should be completely snipped off with a pair of iris scissors, and the surface and edges of the wound touched with the caustic pencil. This practice I have repeatedly adopted with the best effect, in prolapsus from wound as well as from ulcer.

Soft and semi-transparent and unadhering capsular cataracts may all be conveniently ex-

Extraction  
of the soft  
cataract.

\* I have stated elsewhere an additional reason for making the section not too distant from the pupil, viz. the easier escape of the lens. See "Observations on the Cataract." *Med. Chir. Trans.* Vol. V. I prefer it midway between the pupil and margin of the cornea. If it be higher than this, the lower margin of the pupil insinuates itself into the section, and the cicatrix not only disfigures the cornea in the greatest degree, but encroaches very disadvantageously upon the pupil.



tracted. They pass through a smaller section. The capsule is easily laid hold of with a hook or forceps. The semi-transparent, by which I mean the cataract with an opaque nucleus and transparent circumference and capsule, forms in most cases a secondary cataract; that is, a portion of the transparent lamellæ and capsule become opaque, and occupy the pupil or a part of it. It is rent and detached with the greatest ease by a touch of the needle passed through the cornea after the healing of the section.

Instru-  
ments.

The construction of instruments employed in operations is a point which every man must decide for himself. The knife of the eminent Professor Beer of Vienna is that which I am in the habit of using, thinking it on the whole better adapted than either Richter's or Wenzel's, to make a safe and expeditious section.\*

Preparative  
and after  
treatment.

The more or less inflammation which follows the operation is of course depending in great measure upon the habit of body. The patient should be well purged, and live abstemiously for a short time previous to it. If disposed to fulness in the vessels of the head, cupping may be premised the day before the operation. It is a matter of some importance to examine the section, and adjust it accurately before finally closing the eye. I think it useful to let the

\* See Plate V, fig. 4.

patient rest for a few minutes with his eye closed, and then to direct him to open it two or three times successively : a slight friction of the lids assists the pupil to recover its figure, and dissipates any small floating particles of lens. The sitting posture in an easy chair is most favourable after the operation, until the patient feels fatigued and desires to go to bed. Confinement to bed produces great restlessness, and is of no advantage to persons not constitutionally ill. If the patient complains of pain on the evening of the day of operation, a full blood-letting removes it, and should not be omitted.\* I never give opiates. A light bandage passed round the night-cap and fastened to it is a sufficient covering for the eyes. Compresses on the eyelids are generally better omitted ; the bandage may in most cases be laid aside on the second or third day, and a deep black shade substituted for it. During the night, however, the bandage should be applied for the first week to prevent the accident of rubbing the eyes in sleep.

I know of no peculiarities requiring a distinct notice in the treatment of inflammation after extraction. Topical blood-letting and blisters are sometimes necessary, and a strict antiphlogistic regimen should always be enforced. There is

\* It is the custom of M. Roux, of Paris, a dexterous and successful operator, to apply a blister to the nape of the neck after the operation.



often an irritability to light, and aversion to open the eye, which is removed by two or three brisk doses of calomel. When the section protrudes only in a small degree, it soon levels down, so as to restore the figure of the membrane. When the protrusion is more considerable, the patient is afterwards subject to repeated irritable ophthalmia, and a troublesome exulceration of the cicatrix sometimes occurs. I have in one case seen small transparent vesicles form on the line of the cicatrix at intervals, which occasioned much intolerance and distress to the patient for a long time subsequent to the healing of the section.

Unfavorable  
results of  
the operation,

The inflammation of the iris, the interstitial ulceration and opacity of the cornea, the separation of the edges of the section by the intervention of another texture, the redundant deposit of lymph in the section, or the ulceration of its edges, are the mischiefs which occur after unfavorable extractions. Blood shed by a wound of the iris in the anterior chamber is quickly absorbed. Where it has even filled the entire chamber, I have found the aqueous humor clear on the succeeding day.

The coalition of the iris and cornea adjoining the section, occasioning a disfigured pupil, is the result of a prolapsus or a læsion of the iris. Although the process of healing is painfully re-

tarded, and the corneal cicatrix is ever after conspicuous from its breadth and prominence, excellent vision is often obtained under these circumstances. The iritis may, however, be vehement, and proceed to amaurosis, or it may terminate favorably in constricted pupil. This state is seldom uncombined with a secondary cataract, i. e. a portion of continuous capsule, which having become opaque, knits up the pupil by adhesion to its margin. The dimness of the cornea, if any, is slight and transient, except an interstitial herpetic ulcerative action, connected with a bad condition of the edges of the section, be present, when the cornea takes on an opacity of a very intractable kind. The sclerotica is in this case inflamed, and very minute depressions appear on the surface of the cornea, which undergoes a total loss of brilliancy, although it remains obscurely transparent. The restoration of smoothness to the surface does not diminish the lack-lustre appearance of the membrane. The patient has a perception of light, but no vision of objects. In fact, the cornea precisely resembles that of the dead subject. Mercury is of uncertain efficacy in this case, which fortunately is very rare. Time and tonics do most for it.

---

I have now adverted to the principal miscarriages of the operation, both during and consequent upon its performance, which I have met



with, or witnessed in the practice of others. When the operation is perfect, its pre-eminence is too conspicuous to require illustration. It exacts the homage of admiration beyond any effort of the art.

The adjustment of the light, its exclusion from the other eye by the compress and bandage, the regulation of the seats of the patient and the surgeon, the light and firm support of the upper lid, the calm and easy penetration, and the quick and steady passage of the knife across the chamber, and without a pause, through the opposite border of the membrane, so as to anticipate the escape of the humor, and preserve the iris in situ; the deliberate completion of the section, all pressure being removed, either by the progress of the knife, or by a clean back stroke, or by the aid of the finger-nail dividing the cornea upon its edge, as may be most expedient, are, in brief, the material points of the operation. I will only add, the capsule should be freely lacerated in the centre, not incised concentrically to the lower border of the pupil.

It would scarcely be credited by a by-stander who saw the operation happily executed upon a steady and well-formed eye, that it presented any difficulty; a conclusion applied to every thing well done, whether warranted or not. But the incidental embarrassments are too frequent and

numerous to admit of being always anticipated, and the only security against them is the constant habit of practising the operation, and the confidence thence acquired, tempered with a due sense of responsibility.

Habit will make any man ambi-dexter; and the rest for the elbow, so much insisted upon, having been once laid aside, would prove a hindrance rather than a help to the operator. After thoroughly understanding the minutiae of the operation, habit will also render it unnecessary for him to rehearse the several steps and stages of the performance before advancing to it. The memory of a successful operator is altogether technical, and his mementos are carried, to use a homely figure, "*sur le bout du doigt.*"



## SECTION II.

---

### OF THE OPERATIONS FOR ARTIFICIAL PUPIL.

FROM morbid alterations of the cornea, or iris, or both, result those several states of the organ which suggest the formation of an artificial pupil. The disease may be simple, that is, affecting exclusively the cornea or the iris, or it may be complicated, and involve both textures.

#### *Simple states :*

1. An indelible central opacity of the cornea, more or less eclipsing the healthy pupil.

2. A closed or obliterated pupil; the crystalline capsule opaque, and adhering to the iris; and probably the lens also opaque, unless the disease is consequent upon the operation of couching or extraction.

The first supposes a healthy state of the iris and anterior chamber, the second a transparent cornea.

*Complicated states :*

1. A closed pupil, with or without concealed cataract, combined with a partially opaque cornea.
2. A central opacity of the cornea combined with a constricted pupil and cataract.
3. In addition to the closed pupil, a partial adhesion of the iris to the opposite surface of the opaque cornea.
4. A permanent prolapse of the iris through an ancient breach or section of the cornea, involving more or less of the pupillary margin.

In both the latter cases, the anterior chamber is reduced in dimensions in proportion to the extent of the adhesion or prolapse.

These are the chief circumstances—it would be endless and useless to detail all their possible modifications,—which have given rise to the proposal of an operation, having for its principle, the formation of a permanent aperture in the iris opposed to a transparent portion of cornea.

Many very ingenious methods of accomplishing this object have been devised both by foreign and British surgeons and oculists. The



tracts of the late Mr. Gibson of Manchester, and of my friend Mr. Guthrie, surgeon to the Westminster Eye Infirmary, are well entitled to the diligent perusal of students investigating this subject. A remark, which I had early occasion to make in reference to the treatment of cataract, is pointedly applicable to this subject, to wit, that no one method of operating commands an exclusive preference, and that the eligibility of either can only be determined by the peculiar character and bearing of the case before us. It follows, therefore, that a particular description of the various modes of operating, without reference to the precise conditions to which they are applicable, although interesting as an exposé of professional ingenuity, can have no other tendency than to embarrass by a multiplicity of materials, and that it is quite impossible to arrive at a fair estimate of the respective merits of each operation by such a mode of proceeding. Infinitely varied as are the states which call for the operation, a sufficient fertility of invention has been exhibited to admit of an unhesitating preference of some one or other method well suited to every case that can occur; and so far as regards this point, the judgment of the young surgeon requires only to be assisted in its decision by the experience of those who have enjoyed repeated opportunities of putting the merits, general and comparative, of each manœuvre to the test.

In the hope of simplifying a subject which has been in some degree complicated by useless amplification and over refinement, I shall content myself with setting down the operative method which I regard as best adapted to each particular state described.

*Simple states :*

1. *A central opacity eclipsing the pupil.* A section is to be made, with the cornea knife, in the transparent portion of the cornea, from two to three lines in length. This being done, by a gentle pressure upon the opposite side of the eyeball, the iris protrudes in the form of a little sack or bag at the wound, which is to be snipped off with a fine pair of scissors. The iris then recedes, and presents a permanent aperture more or less circular. The corneal section may be made on either side of the pupil as required. The merit of this highly ingenious operation is due to the late Mr. Gibson. It is applicable only to the above state, for if the pupil is closed and in adhesion with the capsule, or if adhesions exist between the iris and cornea, no protrusion will take place. In making the section, it is very important that the direction of the knife should be perpendicular to the cornea, for if its passage is oblique, the cicatrix will be so much extended as to obscure the new pupil. On this account, the practice of Professor Beer to draw out with



forceps and excise the pupillary margin, more completely answers the intention.

There is another operation which I have performed successfully in this case. A narrow bladed and finely pointed knife, cutting on one edge, such as was used by Mr. Cheselden for dividing the iris,\* is introduced through the sclerotic coat and ciliary margin of the iris into the anterior chamber, with its cutting edge opposed to the face of the iris; that membrane is then to be completely divided on the temporal side, including the border of the pupil, by repeated gentle nicks. The iris being sound, and consequently elastic, the section immediately assumes the figure of an equilateral triangle. The superiority of the operations before-mentioned consists in their less liability to be followed by cataract. If cataract ensues, as is pretty uniformly the case in the simple division, it must be broken up for absorption in a subsequent operation with the needle, provided that measure should ultimately be necessary.

2. *A closed pupil, the crystalline capsule opaque and adhering, and probably the lens also opaque, if not previously extracted or depressed.* It should be observed, that in the majority of instances the capsule and lens have undergone the change here presumed; but this is a matter not always pos-

\*.See Plate IV. fig. 9.

sible to be ascertained, and one which affords no ground for varying the plan of operation. It is therefore best to provide for it. The section should be made precisely as in extraction. The centre of the iris should then be raised under the flap of the cornea with the forceps, and as large a piece of the membrane as can be embraced by the convex scissars, should be clipped off. Through such an opening there will be no impediment to the passage of the lens. I have repeatedly performed this operation with perfect success.

Wenzel, finding the mere division of the closed fibres then only useful when they are on the stretch, included a central portion of the iris in the section of the cornea, and removed the triangular flap thus formed, with a stroke of the scissars; a method perfectly efficacious in the state of closed pupil, whether the cataract remains or not. It is only necessary to bear in mind, that a sufficient portion of the iris must be excised to allow of the easy escape of the lens, a point to which Wenzel's directions do not extend, since his operation supposed the previous depression or removal of the cataract.

In the seventh volume of the Medico-Chirurgical Transactions is an interesting paper by Professor Maunoir, of Geneva, relating three successful operations for artificial pupil performed



with his scissars in the manner recommended by him. In two of these cases the pupil was constricted upon an opaque capsule and lens. To this paper is appended a letter from Scarpa, justly complimenting M. Maunoir on his skill and success, and adding his opinion that an operation compounded of Wenzel's and Maunoir's, is the most appropriate to the case of constricted or closed pupil with opaque lens.\*

*Complicated states :*

1. *Closed pupil, combined with a partially opaque cornea.* To this state either of the operations last described may be applicable; this must however depend upon the extent of the opacity. A section of the opaque part of the cornea is unadvisable, as it is strongly disposed to ulcerate instead of healing kindly. I should therefore avoid it, if a portion of the cornea remained sound, sufficient to allow of the unavoidable encroachment of the cicatrix. But in all cases of closed pupil, the iris retaining its position, I hold it to be expedient, if possible, to remove a portion of that membrane; and to this end the section of the cornea is indispensable. The mere division of the fibres of the iris is ineffective, by reason of the loss of its elasticity. An opacity of the cornea however may be of such extent as to leave no choice of operation, as the cicatrix of the section

\* See Explanation of Plate IV.

of that membrane would obscure the small portion which remains transparent. In such a case, the separation of the ciliary margin of the iris, after the manner of Scarpa or Schmidt, is the only practicable proceeding.

2. *Central opacity of the cornea, combined with constricted pupil and cataract.* In this case we need only deviate from the usual method of extracting, by dipping the point of the knife behind the pupillary margin on the nasal side, so as to include a sufficient portion of the iris. If after the removal of the lens, the incised portion should not be completely detached, the flap may be easily snipped off with the iris scissors. It may be useful to remark, that pressure should, as much as possible, be avoided during the extraction of the lens. Other modes of procedure in this case are as follow :

1. An opening is to be made in the cornea, distant about three lines from the point at which the new pupil is to be formed. Through this aperture the ciliary margin of the iris, detached by the double hook of the ingenious M. Reisinger of Augsburg, is to be drawn out and excised.\* If the cataract should impede vision, it must be treated in a subsequent operation.

2. Dividing the iris horizontally on the nasal

\* See Explanation of Plate IV.



side, and couching or breaking up the cataract before withdrawing the needle, which must be adapted to that purpose.

I do not hesitate to prefer to either of these modes the extraction of the lens by excising a portion of the iris, a thing so easy to be done that it is often difficult to avoid it. The distinct performance of the operation for artificial pupil, and the treatment of the lens by the absorbing process, after the lapse of an interval, is for obvious reasons objectionable, when it is possible to do otherwise.

3. *Closed pupil and partial adhesion of the iris to the opaque cornea.* The separation of the cornea and iris when in adhesion, is an attempt altogether unadvisable. It is only at that part of the transparent cornea, which is free from adhesion to the iris, that the attempt to make an artificial pupil can be of any avail. The removal of a portion of the fixed iris, by hooking it through a limited section of the cornea, for the purpose of strangulation or excision, is an operation of much delicacy. It is however in high repute with the German surgeons. If the iris is at any part rendered tense by adhesion to the cornea, an extensive vertical or oblique division of its fibres with the iris knife, or Maunoir's scissors, is best adapted to this case. The simple incision of the iris is enough, if it

either retain its natural and healthy texture, or be placed upon the stretch; and if not, it is inefficient, as before observed, owing to its inelasticity. But in this, as in all cases, if the opacity of the cornea is of such extent as not to allow of incision, the operation of Scarpa must be resorted to.

4. *Permanent prolapse of the iris through a breach of the cornea, involving more or less of the pupillary margin.* This is the state for which the operation was first devised and practised, and when it is consequent upon the removal of the lens, is the most favorable condition for the artificial pupil. The appropriate procedure is that of Mr. Cheselden, viz. the transverse division of the stretched fibres of the iris; and which, if the section be made in front of the membrane, i. e. from before backwards, admits of no improvement. The edges of the section instantly recede, and form an excellent pupil. If the lens has been previously extracted, the intention is completed at once; otherwise the lens must be couched or broken up, and in the latter case may require a subsequent operation. Sharp, Woolhouse, and others, who followed Cheselden's plan indiscriminately, naturally complained of their want of success. Wenzel discovered the cause, and demonstrated the necessity and the mode of removing a portion of the membrane. This was a very important step in advance, and



is the principle of the most successful operation since performed for the artificial pupil.

It may happen that a partial adhesion of the pupillary margin to the cornea may be combined with a healthy lens. In this case, the removal of the free border of the pupil, drawn by a pair of forceps through an incision of the cornea, is preferable, on account of preserving the transparency of the lens.

I am fully aware, not only that other states than those which I have mentioned, may frequently occur; but that even in these, circumstances may possibly arise, to render other modes of operation more expedient than those which I have recommended. In fine, I consider it quite impossible to reduce a subject in its nature so purely circumstantial, and therefore discretionary, to the rule and line. The rationale of such operations is simple and intelligible, and it may safely be left to the genius of the surgeon to meet the exigencies of each individual case, in this as in many other instances. Manual adroitness is, in an especial degree, required in these operations, not only because the space within which the instrument is to move is so confined as to endanger contiguous parts, but because the division of textures in a morbid state is opposed—especially of the iris, on account of its inelasticity and over-vascularity—

by a greater degree of resistance and a larger effusion of blood, than the division of the same parts in health. Hence the operation must be executed as quickly as is consistent with gentleness. For another reason this is important; the organ upon which we are operating is an unsound one, and having been recently and perhaps repeatedly the subject of inflammation, is in too irritable and delicate a state to admit of violent or long continued manual efforts. For the latter reason, it is also highly desirable to finish the operation at one sitting, and I therefore decidedly disapprove of coupling with the formation of an artificial pupil, the displacement and breaking up of the lens to undergo the tedious process of absorption, whenever it is possible to extract or even to couch it. The organ has already suffered too much, and the operator should either not interfere with the transparent lens, or should remove that which is already opaque; and as the latter may almost always be presumed, while it remains *in situ* behind a closed pupil, that operation is preferable, which provides for its removal.

So much for the mechanism of the operation; it now only remains for me to add a remark or two on the more important question of its practical application and value, and the circumstances which should influence our prognosis and determine our practice. There are two



main questions for consideration when these cases present themselves to our notice, both of which should be determined as far as we are able to decide them, before the operation is undertaken. The first in importance as in order is, whether the retina retains its sensibility, and the vitreous humor its texture and transparency. The second, whether the manipulation required is so far practicable as to afford a fair prospect of success. The circumstances under which vision has been suspended are more or less favorable, according to the seat, extent, duration, and character of the inflammation. If the disease is confined to the cornea, the case affords the best encouragement. In the prolapsus iridis from ulceration of the cornea, the iris is only passively concerned, and here it seldom happens that the retina has suffered. If the closure of the pupil has supervened upon wound or injury, as the operation of couching or extraction, it is in general more favorable than when it results from idiopathic inflammation of the choroid and iris of long standing, or from repeated attacks of inflammation. In the latter cases, the organ usually presents certain signs of an organic change in its altered volume and figure, in the yielding and crumpling of the flaccid sclerotica under the impression of the finger, the permanent congestion of the sclerotic and choroideal vessels, the great convexity and discoloration of the iris, and the yellowness of the opaque lens.

if any part of it is visible, the partial detachment of the pigmentum nigrum, &c. An extensive adhesion of the iris to the cornea denotes the greatest degree of organic derangement, and constitutes the most unfavorable state for operation.

In a great proportion of cases in which the operation might, in a manual view, be executed with perfect success, no advantage would result from it; as in cataract, complicated with amaurosis, the extraction proves to be of no avail.

Difficult as the operation may be and often is, it is the state of the eye by reason of the disorganizing inflammation which has preceded it, that forms the most frequent impediment to its success. When the signs of organic change in the eyeball are present, it should never be attempted; without some healthy perception of light, I should think it offered a very small chance of usefulness.\* In the most favorable circumstances, the patient should be led to entertain a moderate degree of expectation; and as persons in a state of blindness are invariably over-solicitous to submit to any experiment for the recovery of vision, the surgeon, for his credit's sake,

\* I say, healthy, because the morbid perception, as of gleams and flashes of blue or variegated light, stars, and sparks, so common in these cases, is decisive of the disorganization of the retina.



should be careful not to limit his inquiry to the mere practicability of an operation; nor, when taking a larger and more deliberate view, to be drawn into it against his judgment. If this is undecided, let him so state the circumstances as to make himself responsible only for the mechanical process. Let it, for example, be expressly understood that without it the case is hopeless, and that if unhappily nothing should be gained, nothing will be lost by the operation. It is seldom, if ever, wise to do the operation upon one eye, the other remaining sound.

The ensuing inflammation is in general moderate, particularly so, if the lens has been previously removed. The success of the operation, however, is sometimes foiled by the train of morbid actions which ensues from the thoroughly diseased condition of the cornea and iris, where the instant result had inspired a hope of better fortune.\*

\* Previously to the performance of any operation for artificial pupil, we should always ascertain whether any improvement to vision may not be gained by the use of belladonna or stramonium. The dilatation of the constricted pupil to the size of a small pin's head, or even less, will often give the patient a degree of vision, which will render any operation unnecessary. This precaution should in no case be neglected; for we will frequently in this manner restore good sight, when we least expect it. A very remarkable instance of this fact occurred in the following case.

J. Steinberg was received March 12th, 1824, at the New York Eye Infirmary, by my colleague, Dr. J. Kearny Rodgers; blind, in consequence of closure of the pupils from inflammation of the iris. The constriction of the pupils appeared to be so complete, and the original inflammation so severe, that the patient was assured that he could expect no relief, except from an operation for artificial pupil, and that even in that case, success was extremely doubtful. The sight remaining to the patient was totally insufficient for any useful purpose. He could not discover the form of any object, and, indeed, could do little more than distinguish light from darkness. Before performing an operation, Dr. Rodgers took the precaution to order the application of extract of stramonium. At the next visit, the patient had gained a degree of vision which made it evident that no operation would be necessary. He was therefore directed to continue the stramonium. Under this application, the sight regularly improved, until at length the patient could read without difficulty type of very moderate size. The most surprising circumstance attending this case is, that notwithstanding a slight enlargement of the pupils could be observed after the application of the stramonium, yet they were still so excessively small, that it was altogether unaccountable how the patient could see so accurately. The improvement in vision was permanent after the discontinuance of the stramonium; and this man now gains a comfortable subsistence by his occupation of comb-making, which he performs to the perfect satisfaction of his employer. The appearance of the pupils is shown in Plate 4, Figures 5 and 6.

This case is taken from a Paper on Artificial Pupil by the Editor, in the New York Medical and Physical Journal, No. 14. He has also annexed the plate from the same paper, with delineations of the results of some operations for artificial pupil, and given an account of the cases in the explanation of the plate.—*Editor.*



## TREATMENT

OF THE

## DISEASES OF THE EYE.

### CHAPTER IV.

#### DISEASES OF THE APPENDAGES.

OF the treatment of diseases of the orbital cavity and appendages much need not be said. I have spoken of the excision of adipose tumors and cysts. Such cases are sufficiently within the observation of the general surgeon to render a fuller discussion of them unnecessary in a work, the more particular object of which is to communicate information upon a department of pathology, which, unfortunately for mankind, he has too much neglected to cultivate.

## SECTION I.

---

### DISEASES OF THE EYELIDS.

STYES, if large and painful from inflaming the eyelids, should be discharged with the point of a lancet, and poulticed, or bathed with a slightly astringent wash, according to circumstances. When phlegmonous, indurated, and slow to suppurate, occasional friction will often promote absorption of these little swellings, as we may conjecture was known of old, from the reputed specific effects of a wedding-ring, or the tail of a black cat. Hordeolum.

The disposition to stye is not only very troublesome, but very injurious to beauty. The permanently conspicuous redness of the borders of the tarsi, a slight degree of thickening and elevation of the conjunctiva, and small cuticular denudations, are the results of their frequent formation, and the loss or scantiness of the cilia greatly increases the deformity. The nitrated, or red or white precipitate ointment of mercury, diluted so as to give a momentary smart upon closing the lids, should be used daily or thrice a week at bed-time, till the part acquires a healthy aspect; and the lids should be occasionally smeared with any soft and unirritating salve at intervening periods. Alum and zinc



washes assist this object. The chronic indurated styte, if not dispersed by the stimulant ointment, should be excised. An appropriate attention to the habit of patients is essential, for this is always faulty.

Lippitudo.

The acute lippitudo generally yields to a single stimulant application. Some exceptions however occur. The lead and zinc ointment, or one combined with opium, will often agree with those slight, but very irritable lippitudoes sometimes met with;\* but there are persons to whose sensations even cetaceous ointment is painful, and who derive no benefit whatever from unctuous applications. In this case hot water affords most relief, and it is a remarkable fact, that it is soothing and refreshing to the eye, at a temperature which is not endurable to the hand.

The chronic lippitudo is a very deforming disease, and often very intractable. It is accompanied with much intolerance. The vessels of the palpebral conjunctiva are turgid, and at length varicose, the membrane a little overlaps the thickened tarsal border; this is partially if not quite denuded of cilia, and small surfaces of the adjoining cutis are excoriated. The follicles are plugged, and here and there is one so

\* The addition of ten or fifteen drops of the liq. plumbi acet. to two drachms of the ung. zinci, I have found to answer completely in such cases.

much distended by inspissated mucus, as to occasion acute inflammation. These should be opened with the point of the lancet, and the white consolidated secretion removed, the conjunctiva should be occasionally scarified, and the meibomian borders stimulated by one of the ointments above-named. The tarsal edges should also be frequently bathed with an astringent lotion. In the aggravated and obstinate cases of lippitudo, where the conjunctiva is altered in its texture, the sulphate of copper lightly carried over the thickened conjunctiva and ulcerated border of the tarsus is highly useful; and stimulant solutions of copper, zinc, lunar caustic, or sublimate, applied with a camel-hair brush to the tarsal edges before smearing them with the ointment, are likewise advantageous.

The degree of strength in which the ointment should be used, the quantity to be used, and the mode of applying it, are points of no small importance in the treatment of these diseases. Unless it excites a pretty smart irritation and provokes a copious flow of tears, it does little if any good; but the irritability of the conjunctiva varies so much according to the stage of the disease, the time of making the application, and even the general habit, that it is impossible to fix a standard of strength. As to the manner, patients should be instructed in the intention; and for the strength and quantity, in the desired effect of



the remedy. The degree of irritation should be such as to prevent the patient from keeping his eyes open for some succeeding minutes ; but no increased congestion should be apparent on the following day. The patient therefore should feel his way, and measure the irritability of the conjunctiva, by advancing gradually from a lower to a higher stimulus. In acute lippitudo a little cold cream or spermaceti will occasion a severe smarting and profuse flow of tears, whereas in the chronic form, the strong mercurial ointment is often necessary to produce this effect.\*

\* The 'golden ointment,' as it is called, is an excellent remedy. I shall be accused of heterodoxy : but I must in justice assert, that the inventor of this arcanum deserves well of his country, for if his patriotism be equivocal, the virtue of his nostrum is at least certain, when judiciously prescribed. If it did not unfortunately aspire to be a panacea, its beneficial operation would be without exception ; and indeed the greatest evil of quack, as of regular medicines, is their abuse. Few things capable of doing much good, are not also capable of doing mischief. An old lady of Paris, whose husband had become famous for an eye-water, had the misfortune to lose her spouse and his secret together. In this dilemma, harassed by applications for the nostrum, she had recourse to the water of the Seine, and was not more gratified than surprised to find that the collyrium had lost nothing of its virtue. After having enriched herself by a successful traffic, it so chanced that she fell sick, and conscience-stricken at the prospect of death, she applied to an eminent professor of surgery, instead of a priest, to relieve herself of the burthen of sin with which her soul was encumbered. "Soyez tranquille, mon amie," said the professor, "de tous les Medecins vous êtes le plus innocent ; vos remedes n'ont fait du mal à personne !"

The correction of the diseased states and secretions of the ciliary apparatus proceeds upon the same principle as the cure of lippitudo, with which, in a greater or less degree, they are very generally combined. The re-establishment of a healthy conjunctival surface and a healthy meibomian secretion, is the object to be attempted. In general the mercurial ointments are the best remedies for this disease.

*Tinea ciliaris.*

Cleanliness is a point of the first importance, and it is the more necessary to mention it, because the disease is often set up, and is always aggravated by neglect of it. The margins of the lids, and the roots of the cilia should be thoroughly cleansed from loose scabs and branny incrustations before anointing them; the ointment, liquefied by a gentle heat, should be applied upon both edges of the cartilage with a camel-hair brush; keeping them through the day slightly besmeared with a mild ointment, as the tutty, prevents the formation of fresh crusts. It is needless to say, that more depends upon the patient than the surgeon in the cure of these complaints, and that the incurable states, those which admit only of palliation, are invariably the consequence of neglect, and might therefore have been prevented.

When cilia are inverted from a diseased growth, they must be kept plucked until by the improved

*Trichiasis.*



condition of the hair-gland, under the means used, the disease is removed. If a case occurs in which the vitiated site or incurvation of one or more cilia does not admit of correction, the corresponding follicle should be obliterated by repeatedly touching it with a fine caustic pencil. When the incurvation is depending on a disease of the tarsus, the case must be treated as entropion.

Entropion.

The treatment of the inverted eyelid, upon the plan recommended by Scarpa, will, according to my experience, be effective in nine cases out of ten; that is, by the removal of a fold of skin with a pair of scissars from the surface of the eyelid. The relaxation of the integument operating as a cause of the disease, is more frequent upon the lower than the upper lid, but the operation is applicable to cases originating from other causes. The surgeon should be careful to adapt the position and extent of the wound to the site and degree of the inversion.\* I have now and then

\* A more recent method of treatment adapted to entropion, and in some respects superior to the one recommended by our author, consists in the application of concentrated sulphuric acid to the skin of the affected eyelid, in such a manner as to produce the necessary degree of eversion, by the contraction thus effected in the skin.

For this purpose, a small quantity of the acid is rubbed upon the skin of the eyelid, either with the blunt end of a probe, or a piece of smooth wood, opposite to the inverted

met with a case in which the inversion was clearly depending upon a callous roll of conjunctiva at the orbital edge of the tarsus, in which case the disease was removed by the excision of this roll; of itself a disease requiring such a remedy, as before noticed. In cases of a circumscribed inversion produced by cicatrix from burn or wound, I have found an operation similar to that of Dr. Crampton, an effective remedy; sometimes the complete division of the conjunctiva and tarsal cartilage, including the inverted portion, and parallel to its border, with the aid of sticking-plaster, has proved sufficient. I should see no objection to the entire removal of that portion of the tarsal edge which was incorrigibly inverted from such a cause, especially when combined with distichiasis, by which is meant a preternatural growth of cilia from the meibomian border of the tarsus; but

cilia, and including a space, extending a little beyond the disease on each side; and from a quarter to half an inch wide, according to the degree of inversion. After remaining a few seconds, the acid must be wiped off, to prevent its spreading over parts which we do not wish to injure.

The application is then to be repeated a second and third time, or even oftener, until the contraction thus induced in the skin shall have everted the lid sufficiently to prevent the cilia any longer rubbing upon the eyeball. A detailed account of this mode of treatment, and of this disease generally, may be found in Guthrie's Operative Surgery of the Eye, recently published.—*Editor.*



the removal of the entire cartilage, which I have twice performed in aggravated cases of this disease, although by no means difficult of execution, is an operation of great and needless severity, and one which, in my experience, is not warranted by the degree of relief which it affords; to say nothing of the permanent deformity which it occasions.

Ectropeon.

The ordinary ectropeon is cured by the excision of a portion of the thickened or redundant conjunctiva which occasions it.\* The perpendicular division, or the removal, as the circumstances require, of a triangular portion of the tarsal border by a double incision, as mentioned p. 252, is the operation best adapted to the long-established and aggravated forms of the disease; an adhesion of the everted eyelid to the cheek, adds much to the difficulty of the case, and renders all modes of treatment merely palliations of the deformity. I have, however, succeeded in greatly lessening the deformity in some such cases.

Palpebral tumors.

Concerning tumors of the eyelids I have little to add to what will be found in the pathology. Often the encysted tumor is seated superficially

\* Simply rubbing the everted and redundant conjunctiva with sulphate of copper, on alternate days, is sufficient to cure cases of very considerable ectropeon. This mode of treatment, however, requires perseverance.—*Editor.*



and loosely connected to the tarsus, in which case it must be removed on the outside of the eyelid, by just separating and turning to either side the fibres of the orbicularis muscle. It is only when an intimate adhesion subsists, (the cyst is often formed betwixt the cartilage and the ligamentary membrane which covers it,) and the appearance of a white circumscribed indentation is seen upon everting the tarsus, that the excision is to be made from the interior by division of the cartilage; in which case it is always executed with perfect facility. If the cyst be not extirpated, but merely incised and its contents expressed, the tumor speedily re-appears.\* If its adhesion to the tarsus be such as to prevent the entire removal of it, the remnant of the cyst should be touched with the caustic pencil.

For the permanently drooping lid, if an operation be ever desirable,—which, as an unclosed state of the lids is a pretty certain fore-runner of diseased conjunctiva, amounts with me to a question—a fold of integument may be removed by the knife, or by pencilling out a portion of it with the strong nitric acid. It is unnecessary to describe an operation for dividing the cohering tarsi. When complicated with adhesion to the globe, the case is incurable.

Lagophthalmus,

and cohering tarsi.

\* My experience does not, in this instance, correspond with that of Mr. Travers, as I have repeatedly succeeded in permanently removing these little tumors by merely puncturing them from the inside of the eyelid, and evacuating their contents.—*Editor.*



## SECTION II.

---

### OBSTRUCTION OF THE LACRYMAL PASSAGES.

Stricture of  
the lacrymal  
and nasal  
ducts.

THE treatment of the disease improperly termed "fistula lacrymalis," has occupied a large share of the labor and talent of the profession;\* yet, notwithstanding this advantage, the practice is to this day unsettled and unsatisfactory.†

\* See the numerous papers in the Memoirs of the French Academy, and the works of Sharpe, Pott, Desault, and other eminent writers.

† In proof of this remark, I may observe that nearly all the schemes hitherto suggested have been executed within my knowledge, by different surgeons, viz. the small probe and injecting syringe of Anel, the sound and syringe for the nasal duct, the seton of silk or catgut, the bougie or nail-headed style, the metallic tube, &c. In Paris, M. Dubois employs the silk seton of Mejan, M. Dupuytren the permanent tube of Wathen, M. Roux the mesh seton, introduced by means of a watch-spring from the sac. M. Beer, of Vienna, uses, for a seton, a coil of catgut, such as is used for fiddle-strings. Among the surgeons of London, Mr. Ware's style is chiefly in use, although the practice is evidently losing credit.

Mr. Pott was in error when he concluded in his criticism of Anel's practice, that the stricture of the lacrymal ducts was very rare, because the mucus of the sac was habitually, or upon pressure, discharged at their orifices; the stricture of the ducts is, on the contrary, frequent; and although this morbid secretion of the sac is often present with a free state of the lacrymal ducts, the cases of obstruction on the nasal side are very frequent, in which no such symptom exists.

The absorption of the meibomian mucus by the puncta lacrymalia, and its regurgitation on pressure, as described and considered by Scarpa to constitute the first stage of the disease, independent of a permanent stricture, is, I think, hypothetical; for, if founded in fact, this symptom would be present in every severe lippitudo or ophthalmia with puriform discharge, which every body knows is not the case. Besides, if the duct were open, there is no reason why the fluid, once admitted, should be arrested or regurgitate instead of passing into the nose; but the office of the puncta is the absorption of the lacrymal fluid, as that of the lacteals is the absorption of chyle, and absorbent mouths are distinguished from capillary tubes by the selection of their proper fluid. It by no means follows because a purulent secretion is discharged from these orifices that they have derived it

Mucous discharge.  
Professor Scarpa's hypothesis.



from the eye. Further, there is every reason to believe that the fluid so discharged is the proper secretion of the sac, and cases are frequent in which it is retained and cannot be expressed, owing to strictures both of the lacrymal and nasal ducts. As to the proof of the meibomian border of the lid of the affected eye being more vascular than the other, I need only remark that the irritation of an obstructed sac naturally produces this appearance upon a continuous and highly sensible membrane; if the redness prove any thing, it proves the presence of irritation, and stricture is as probable a source of it as any. But it is demonstrable that the "*flusso palpebrale*" is as seldom present with the symptoms of obstruction in a degree sufficient to support the hypothesis of obstruction from that cause, as the actual obstruction is rare in those cases of its excess which are of ordinary occurrence.

Origin of  
stricture.

The effect of a severe cold in the head to produce a coryza and troublesome watering of the eye, may enable us to form a pretty accurate idea of the cause of a permanent stillicidium; for although an over-excitability state of the conjunctival surface may occasion a more plentiful secretion of the lacrymal fluid, yet it cannot be questioned that the same temporary condition may prevail in the membrane lining the sac and duct, as in the other parts to which it is distri-

buted; I mean a state of vascular congestion and intumescence. This continued, would lead to a permanent thickening of this membrane, and from that cause a diminution of the calibre of the canal. An adhesive process, whether primary or consecutive to the states of suppuration and ulceration, finally closes the duct, so as to render it absolutely impervious. The actual obliteration of the canal by the degeneration of the membrane into a texture resembling cartilage, is a secondary morbid change, and only the result of long continued obstruction.

I have had occasion before to observe that the inflammation and abscess of the sac, although frequently preceded, and in great measure produced by obstruction of the nasal duct, is by no means always referrible to that cause; a considerable degree of obstruction endures for years without a tendency to excite inflammation of the sac; and on the other hand, the inflammation and abscess of the sac in its acutest form, as from exposure to cold and other exciting causes, and sometimes from injury, often exists without any degree of obstruction. In proof of this, I may remark that the incipient inflammation of the sac often admits of resolution by the use of leeches, &c. without further inconvenience to the patient; and the whole treatment required for the abscess in many cases, is simply the discharge of the

Abscess in-  
dependent of  
stricture.



matter, or, in other words, the treatment applicable to a common abscess.

A free opening of the sac for the purpose of discharging its contents, shortens the sufferings of the patient, and saves the skin; but unless the previous existence of symptoms demonstrates the presence of a stricture, an abscess of the lacrymal sac is by no means a sufficient proof of it, to warrant the employment of any further measures. The existence of an abscess is of itself a cause of temporary obstruction, and the stage immediately preceding the formation of abscess may have been the cause of a temporary stricture, of which the abscess and its discharge are the termination. Of this I am well satisfied: that the supposed invariable connection of an abscess of the lacrymal sac with a stricture of the duct, enforced in the writings of the French Academicians, Mr. Pott, M. Desault, and others, have led to an officious and often injurious treatment of this painful and sufficiently distressing malady; and at all events, the first indication is simply that of giving issue to the matter, by a free incision of the sac, and applying a soft poultice in a bag to the inner angle of the orbit.

Treatment  
of abscess.

Supposing, therefore, the case of abscess so far advanced, instead of introducing a style into the ductus nasalis, after opening such abscess, I recommend simply the examination of the duct

with a fine probe ; if the probe pass without resistance into the nose, the case requires no further operative treatment, the integument recovers its healthy condition under an emollient application, the discharge gradually diminishes, and the wound heals. If, on the other hand, upon examination with the probe introduced through the wound into the sac, resistance be offered to its passage into the nose, no more favorable opportunity will be presented for overcoming such resistance. This therefore should be accomplished, but to this the operative process should be limited, and the wound should be suffered to heal without further disturbance.\*

\* Abscess of the lacrymal sac, terminating in fistulous openings through the integuments, is a disease of no uncommon occurrence in children, and for the most part, in such cases, is a scrofulous affection. It is unattended with stricture, and simply a strumous inflammation of the membrane lining the sac, furnishing a muco-purulent discharge. Accordingly in these young subjects, all operations for the purpose of dilating the nasal duct are unnecessary ; fortunately so, for they would be extremely difficult at this early age. The principal object of the surgeon is to correct the state of the constitution which gives rise to the disease, by means which it is unnecessary here to point out. Little local treatment is necessary. An open blister behind the ear is as useful here as counter-irritation generally is in scrofulous diseases : and a weak astringent wash to the part, together with the application of some stimulant ointment, if the sac have burst, such as the ung. hydr. nitrico oxyd. constitute all the topical remedies which are required.—*Editor.*



Erysipelatous inflammation.

An erysipelatous inflammation of the integuments of the face in the vicinity of the lacrymal sac, in which the peculiar characters of that inflammation seem to indicate that the cutis is primarily affected, however this may be in fact the case, requires to be narrowly watched, especially if it extend to suppuration; for then there will be reason to apprehend a sloughing ulceration of the cellular texture, and probably, the destruction of a portion of the sac from this cause; so that a lacrymal fistula will be the termination of the disease. But a more immediate cause for apprehension exists, if inflammation has been preceded by any sign of obstruction to the passage of the tears, and this, I believe, will be found to be most frequently the case. The inflammation symptomatic of deep-seated suppuration is erysipelatous, as is well known to every practical surgeon, and this is more especially the case where any fascial or ligamentous texture intervenes, like that which covers and supports the sacculus lacrymalis. Therefore, although the incision of the sac is manifestly improper and unadvisable, unless it is distinctly ascertained to be the seat of suppuration, the erysipelatous nature of the inflammation must not be considered to indicate the confinement of the disease to the integument, and the previous healthiness of the sac. I do not hesitate to say that the evil of an uncalled-for incision of the sac is, in every point of view, less than the opening

formed by a process of disease under the circumstances supposed, for in the latter case it will be difficult to prevent a permanent fistula.

The tear falling over the cheek is a sign of complete obstruction, except under strong excitement, when indeed it happens even though the duct is free; but the moist or watery eye indicating the retardation, not the arrest of the tear, is by much the more frequent case. This depends upon an imperfect obstruction of the nasal duct. This state often exists without any other external symptom of disease, and it is a source of considerable inconvenience from the continual suffusion which it occasions, and the necessity it imposes of incessantly wiping the eye.

Moist or  
watery eye.

The epiphora, or, more properly, the stillicidium lacrymarum, resulting from the constricted puncta, or obstructed lacrymal ducts, is relieved by the introduction of a small silver probe into the sac once, or oftener, as may be required. Frequently the obstruction is confined to the saccular extremity of the duct. The case is common, the inconvenience considerable, and the relief complete. The obliterated punctum or canal is a case which admits of no remedy.

Constricted  
puncta and  
obstructed  
ducts.

The patulous puncta are usually combined

Patulous



puncta and  
atony of the  
sac.

with a swollen and atonic state of the canals and sac ; there is no contraction of the orifice on contact with the probe, no obstruction to its passage into the nose, nor any excretion of mucus upon the eye or from the nostril; the situation and figure of the sac are conspicuous from its prominence. Such a state, most frequent in elderly persons, may in part depend upon a feebleness or paralysis of the orbicularis muscle, and a redundant fulness of the skin producing a partial eversion, or an enlargement of the semi-lunar fold of the conjunctiva, displacing the puncta, or disturbing their relative position. It results also from long continued distention of the sac, owing to a morbid increase and retention of its secretion; and under such circumstances, it may be partially relieved by astringent washes, but it is not, in my experience, curable.

Stricture  
symptomatic  
of conjunctival  
inflammation.

The stricture which is occasioned by the extension of conjunctival inflammation to the lacrymal excretories, should be treated only as a sign of that disease. When under the influence of a treatment purely antiphlogistic, and soothing applications, the inflammation subsides, the temporary interruption to the function of these organs ceases; or if it should not cease altogether, the use of a gentle stimulant collyrium will put an end to it. When, however, the state of

distended sac, the regurgitation of mucus upon the eye, and the gathering of the tears in the lacus lacrymalis, are altogether chronic, there can be little doubt of the existence of a stricture, partial or complete; and I am unacquainted with any other mode of treating this disease than such as is adapted to the removal of the stricture, and the restoration of the canal. I am satisfied that the practice of introducing stimulant liquids into the hollow of the inner canthus, in the supine position of the head, and of injecting the sac with astringent lotions, has no other effect than that of aggravating the symptoms of the complaint; how indeed can any other result be rationally expected?

When from the duration of this state the overflow of tears becomes continual, the distention of the sac and the discharge of mucus excessive—the conjunctiva towards the inner angle has a preternatural vascularity, the outline of the sac assumes a circumscribed phlegmonous hardness, and a blush begins to appear upon the skin covering it—when the eminences of the puncta lacrymalia are shrunk and absorbed in the swelling, and in short, the mucous is about to pass into the purulent secretion—even at this period I have repeatedly averted the formation of abscess by re-opening the nasal duct. But when the disease has advanced another

Permanent  
distention  
threatening  
suppuration.

Fluctuation.



stage, and the pointing of the tumor and sense of fluctuation are perceptible, no advantage could be expected from the introduction of the probe through the lacrymal canal, if it were possible; we have now a disease requiring a distinct treatment, and to prevent a complicated fistula—such as results from the yielding of the sac, the diffusion of its contents into the cellular substance, and sinuses spreading in various directions beneath the integument,—the free incision of the sac should be made without delay, and the treatment of the original disease postponed.

The nail-headed style.

When the integrity of the parts is restored, it will be necessary in this case to have recourse to the same mode of proceeding which is adapted to the other stages of the disease, and which appears to me to be the only method of treatment applicable to it upon a rational principle; for I confess myself at a loss to understand how relief can be afforded by the practice of introducing a style to remain in the duct, and I am strongly disposed to doubt whether any permanent benefit was ever derived from such a practice. I am quite aware that the mere opening of the sac affords an immediate and considerable degree of relief; and if a stricture has existed in the nasal duct, which is at the same time overcome by the introduction of the probe, the relief will be still more complete; but the style

which occupies the sac and duct can have no conceivable beneficial influence until it is withdrawn. The disease is transferred from the eye to the cheek, and the oozing of the tears through a small fistulous aperture in the sac, is substituted for their overflow of the natural channel. The state is, upon the whole, less irksome to patients, and in so far the practice must be considered as palliative; but I may fairly say that I have scarcely seen one instance in which this practice had been adopted, and the style was still retained, whether at the expiration of three weeks, or three months, or three years, or double the latter period, in which the disease did not exist in undiminished force, under the modifications which I have just described, that is, as an established lacrymal fistula. Patients are reluctant to part with the style, because, as is natural, they ascribe the degree of relief they have obtained from the opening of the sac, and the diversion of the excretion, to what appears to them to be an essential part of the process.

I have recommended the introduction of a probe into the nose, when such an opportunity is presented, from unwillingness to lose one so favorable for the restoration of the canal; and the only case in which the permanent dilatation is required, is when the passage of the instrument in the direction of the duct is so firmly resisted as to compel the forcible renovation of

Dilatation  
gradual or  
immediate.



the canal. In this case it is obvious that some means must be used to preserve it, and for this purpose two modes of practice are employed. Some surgeons having introduced a dilator into the duct, so far as the obstruction permits, fasten it there, and from day to day renewing the attempt to overcome the obstruction, gain upon it by little and little, until at length the instrument enters the nostril; here it is left for some days, when either an instrument of larger dimensions or a seton is substituted for it, which is not finally withdrawn until the object is accomplished.—Others forcibly overcome the obstruction at once, and afterwards place a style or tube in the newly formed passage. I am not now speaking of the penetration of the *os unguis* or the breaking through of the bony parietes, but of the re-opening of the original passage; which being obliterated by a morbid structure of the lining membrane, of such firmness as to require the employment of force, and to occasion a free hemorrhage from the nostril, is, in fact, the same thing as an artificial channel. Of the two practices, I decidedly prefer the latter; the former is drivelling, tedious, and painful to a degree.

Averse to any and every permanent tent, I formerly introduced a probe into the nose for many days in succession, but the daily increasing facility with which it passed was not a compensation for the pain it inflicted, and the ulceration

of the wound by the repeated interruption of the cicatrizing process.

The practice which I have long employed, and which I adopted as the most successful, after a trial of the several methods of which I have made mention, the tube only excepted, is too obvious to have the merit of novelty. In a large proportion of cases, it has proved successful in curing the disease, both in slight and aggravated forms, in early and advanced stages, without entailing the inconvenience and deformity inseparable from the various contrivances for permanent dilatation, and avoiding altogether a fistulous aperture. Contented with accomplishing the passage of a moderate sized probe into the nose, after the incision of the sac, my attention is exclusively directed to the reduction of the inflammation, and the restoration of the soft parts, with which, be it expressly understood, I never interfere, except in the case of abscess discoloring the skin, and threatening fistula. Thus, with this single exception of abscess, the treatment of the obstruction is one and the same, so far as the point of obstruction is concerned; and it is a point always important to be ascertained; the more so, as it is by no means of uniform occurrence.

The author's practice.

It can hardly be required that I should occupy the time of the reader in showing that the



practice of restoring a passage partially closed, or even establishing an artificial passage, as nearly as possible in the same direction, when the natural channel is obliterated, commands a decided superiority over the practice of making an artificial opening. This applies to the treatment of the urethra, as well as of the ductus nasalis, and it is only in case of abscess, in which the distended and inflamed integument threatens to give way by ulceration, that in either case it becomes necessary to deviate from it.

Lacrymal  
probes.

A set of silver probes, of about five inches long, varying in size, flattened at one end, and slightly bulbous at the point, are the instruments I use for the purpose of restoring the passage. The probe is introduced with perfect facility by one who is familiarly acquainted with the anatomy of the part, from either of the puncta lacrymalia into the corresponding nostril, when no obstruction is offered to its passage. If the punctum be constricted, it is readily entered and dilated by a common pin; and upon withdrawing it, by one of the smaller probes; the direction and relative situation of the lacrymal ducts, the sac, and the nasal canal, point out the proper course of the instrument. It is confirmed by its advance without the employment of force, and the sensation conveyed by the free and unencumbered motion of its point; until the point is fairly within the sac, it

is necessary to keep the eyelid gently stretched and slightly everted; the upper lid being drawn a little upward toward the brow, the lower as much downward toward the zygoma. The point carried home to the sac and touching lightly its nasal side, the lids may be left at liberty, while a half circular motion is performed by the instrument; the surgeon neither suffering the point to recede, nor, on the other hand, allowing it to become entangled in the membrane.

The probe now rests in a perpendicular direction upon the eyebrow towards its inner angle, and in this direction it is to be gently depressed until it strikes upon the floor of the nostril, where its presence is readily ascertained by a common probe, passed beneath the inferior turbinated bone. The probe of smallest dimensions is of sufficient firmness to preserve its figure in its passage through the healthy duct, but it is too flexible to oppose any considerable obstruction, without danger of a change of figure: for the stricture of the lacrymal ducts it is of sufficient strength.

Very many cases of recent origin, and in which the stricture has no great degree of firmness, are completely cured by three or four introductions of the probe into the nostril, at intervals of one or two days.



Lacrymal  
styles.

I have seldom met with a stricture so firm as not to yield to the full-sized probe. I am fully aware of the objection that immediately presents itself, viz. that a passage so obtained is not permanent; by several repetitions of the operation it is often rendered so; but if the resistance is not altogether removed, after a trial of the experiment for some days in succession, I introduce a style having a small flat head, a little sloped, through the punctum lacrymale into the nose, and leave it for a period of twenty-four hours in the duct. If worn longer, as for two days, it ulcerates the orifice; but I have never seen it injure the punctum in the smallest degree, when worn for the full period first named. A day or two should be suffered to elapse before the style is again introduced, and it should then be passed through the other lacrymal duct. The injection of tepid water should be made on the intervening days with Anel's syringe. The plan requires perseverance, as may be said of all plans by which so difficult an object is sought to be effected. In many cases the resistance, in the first instance opposed, is inconsiderable, yet it is sufficient to maintain the disease. The probe passes daily with increasing facility, and after a very few repetitions, with as much ease as through the healthy canal; yet the stillicidium, and even the mucous discharge do not immediately subside, because, although the obstruction is removed by which these symptoms were originally set up,

the parts have not yet recovered the loss of tone which the state of habitual obstruction and inaction has induced; and here the use of the probe is unavailing, if not injurious, as in all cases in which the full-sized probe passes without impediment. It is important that operators should consider this, and not lose sight of the vital function of the parts, in treating the morbid alterations of structure which have interrupted and deranged them. For this mitigated, but not recovered state, time alone, with attention to prevent distention by occasional gentle pressure of the sac if accompanied with mucous discharge, is often sufficient; but the injection of a solution of alum, or even of cold spring water, and the use of astringent washes, will assist. Sniffing a stimulant vapour, as of vinegar, or diluted nitric acid, into the nostril, I have also found useful. It is of course unnecessary to pass a probe, when the fluid injected by the punctum drips in a stream through the nostril or into the throat, as the head of the patient is inclined forwards or backwards; but this test of the freedom of the passage should be had, before the use of the probe is laid aside.

If it be objected to this operation, that it is always painful, and often tedious, I can only reply, that there is too much truth in the objection. I shall be happy to be instructed in one equally effective, and free from these objections. I am



far from assuming that all the other measures employed might not be crowned with success in favorable cases; this, like other complaints, sometimes recovers, not so much in consequence, as in despite of treatment.

Seton.

Of all the other modes, the seton of Mejan alone appears to me to be a rational practice; it too is tedious, and, during its use, deforming.

Tube.

The tube seems, on the other hand, to be of all the most objectionable. I have seen cases of its employment, in which it very speedily became plugged with mucus, the sac habitually loaded, the nostril dry, and the stillicidium permanent. But how a metallic tube can be expected to form a substitute for the natural duct, an inorganic to serve in lieu of an organized part in perpetuity, the functions of the puncta and sac to be restored, and as it were in consent with it, I confess myself at a loss to conceive. I have more than once heard patients sorely regret that they had submitted to its introduction, having received no degree of permanent benefit from the operation, and I have been called upon to remove it, which is not easily done, in one case from the disease which it had set up in the contiguous soft parts, and in others from an exasperation of the symptoms of the complaint. I am now speaking from what has incidentally passed under my notice of the practice of others; but I ought in candor to add, that some very intelligent surgeons, both

English and Foreign, have lately assured me of the general success of the practice in the able hands of M. Dupuytren, at Paris.\*

In the use of the lacrymal probes, caution is requisite; they must be passed with great gentleness, and if the extremity becomes confined, a little withdrawn, so as to prevent their hitching in the membrane, and passing beneath it; the size should never be such as to distend the lacrymal canal, lest it should injure the texture and destroy the tone of the part; and no considerable degree of pressure should be made with one so slight as to be in danger of becoming curved. A probe of sufficient dimensions and firmness to preserve its straightness, is quite within the measure of the lacrymal orifices and ducts, and has strength enough to overcome an ordinary stricture. Injections should be frequently employed to ascertain the progress of the case towards recovery. They are of great use in almost all stages of the disease.

There are undoubtedly many cases of slight epiphora not depending upon mechanical obstruction of the lacrymal excretories.† The zinc and the lunar caustic solutions, the thebaic tincture, the mercurial ointments introduced between the lids, will remove such forms of the complaint, if the patient is so far inconvenienced

Epiphora independent of obstruction.

\* See note M.

† See note N.



as to apply to his surgeon for relief, which is not always the case. Again, the gleety discharge of the sac, as it is not always present with stricture, is sometimes, though not often, present without it. Stimulant collyria, cold water, alum injections, and constitutional tonics, must be employed to cure it. The chronic, thickened, and hardened state of the sac after abscess, and in the state of fistula, is removed by leeches, cold poultices, and saturnine lotions; and if the skin be free from inflammation, it is reduced by the mercurial ointment. A small fistulous aperture, where the disease has been left to nature, is a common and troublesome case. The aperture should be freely dilated, the open state of the canal ascertained by the pocket probe, and the part afterwards treated as above directed.

Abscess with  
caries.

No peculiar treatment is required in the abscess with caries of the bones; a case much less frequent than would be imagined from its description as a stage of the disease. In this case, as in that where the sac has yielded to ulceration before the integument, the skin should be freely divided, that is, beyond the confines of the sac. I never met with a case requiring the use of a trocar, nor do I believe the perforation of the os unguis is ever really required.

Relaxed sac.

I am not practically acquainted with the effect of pressure upon the relaxed or hernial lacrymal

sac. It is an old, and, I believe for a better reason, an obsolete practice, viz. that it is difficult of application and inefficient.

For the opportunity of seeing and treating an interesting case of dropsy of the sac, (the disease described at page 260,) I am indebted to the kindness of Mr. Alexander, to whose extensive observation of this class of diseases it was new, as it was to mine. I exposed the distended and transparent sac by dissection, and removed by the scissars the two anterior thirds of it. A process of suppuration which ensued was for some time troublesome, but the wound at length healed soundly, and the complaint was cured by this treatment.

Dropsy of  
the sac.

With the disease in which the sac is said to acquire the size of a pigeon's egg from an accumulation of inspissated, or rather indurated secretion, having a cartilaginous hardness, and a livid color, I am entirely unacquainted. But a confinement and inspissation of its proper mucous secretion in such accumulated quantity as to distend and morbidly enlarge the sac, giving it the appearance of a firm incompressible globular tumor, I have repeatedly seen and removed by persevering in the use of the probe and syringe. In this case the lacrymal and nasal ducts are equally obstructed. It may be properly

Mucocele.



termed 'mucocoele,' and should not be permitted to continue.

[*Third Edition.*].—On the general treatment of this disease I have only to observe, that additional experience has fully confirmed the efficacy of the practice which I have here recommended. I have frequently anticipated the impending supuration of the sac by the early and gentle introduction of the probe, by which the disease has been arrested; and it is only when the skin partakes of the inflammation as a consequence of distention, that I have recourse to an external opening. After the relief of the sac, a poultice reduces the inflammation; and in a few days the use of the probe may be resumed. In recent obstructions, the disease yields to a very limited course of treatment. In the chronic, perseverance is required, and the early alternate employment of the syringe greatly promotes the object in view. The lacrymal style I have discontinued, finding the benefit resulting from its use disproportionate to the inconvenience.

When the sac has been long over distended, its contents should be gently expressed by the point of the finger from time to time during the injection; which should in this case be made per saltum, and always slowly and with very moderate force; much counter resistance rends the sac, and extravasates the fluid; and although

this accident is of no ultimate importance, it had better be avoided.

Some little time is required for the recovery of the tone of the parts after the canal is re-established. In some cases the symptoms of obstruction only in part disappear during the process of cure; but completely, a short time afterwards. In most cases of inveterate or bony obstruction, the patient finds great relief from the open state of the lacrymal ducts and the occasional ablution of the sac, and I believe makes a compromise on better terms with the disease than by wearing the permanent tube or style.



the accident is of no ultimate importance, it  
had better be avoided.

Some little time is required for the recovery  
of the tone of the eye after the canal is re-  
closed. In some cases the expansion of the  
tonic only in part disappears during the process of  
cure; but completely a short time afterwards.  
In most cases of moderate or heavy obstruction  
the patient finds great relief from the open state  
of the lacrimal ducts and the occasional ab-  
sence of the eye and I believe makes a compari-  
son on better terms with the disease than has  
usually been the case.

The following cases are given as examples of the  
effect of the treatment. In the first case the  
patient had been blind for several years. The  
treatment was successful in restoring vision.  
In the second case the patient had been blind  
for several years. The treatment was successful  
in restoring vision. In the third case the patient  
had been blind for several years. The treatment  
was successful in restoring vision. In the fourth  
case the patient had been blind for several years.  
The treatment was successful in restoring vision.  
In the fifth case the patient had been blind for  
several years. The treatment was successful in  
restoring vision. In the sixth case the patient  
had been blind for several years. The treatment  
was successful in restoring vision. In the seventh  
case the patient had been blind for several years.  
The treatment was successful in restoring vision.  
In the eighth case the patient had been blind  
for several years. The treatment was successful  
in restoring vision. In the ninth case the patient  
had been blind for several years. The treatment  
was successful in restoring vision. In the tenth  
case the patient had been blind for several years.  
The treatment was successful in restoring vision.

## APPENDIX.

---

DR. CHRISTIAN SALAMON, of the Medical and Chirurgical Academy of St. Petersburg, one of ten gentlemen deputed three years ago, by his Majesty the Emperor Alexander, to visit foreign schools for the purpose of acquainting themselves with the state of medical science, an appointment equally honorable to both parties, has politely favored me with the following anatomical sketch, which is the result of some very delicate and laborious dissections. Some of these I have derived much instruction and pleasure from being permitted to witness, and although I do not feel satisfied in all points of the accuracy of Dr. Salamon's conclusions, I have nevertheless seen enough of his general professional intelligence, his talent for minute investigation, and his method of conducting it, to feel assured that the following brief memoir, in his own words, will make no inconsiderable addition to the interest of this work. I have the pleasure, at the same time, to announce, that those who take an interest in these researches, will shortly be gratified by the appearance of an anatomico-physiological dissertation on this subject, upon which Dr. Salamon is at present employed.



OBSERVATIONS  
ON  
SOME POINTS  
OF THE  
ANATOMY OF THE EYE.

---

*SCLEROTICA*, before reaching the cornea, increases in thickness, and divides into two layers ; the outer advances beyond the inner, and is connected with the exterior layers of the cornea ; the inner corresponds to the interior layers of the cornea. On the inside of the inner layer of the sclerotica, near its termination, is an annular groove in the whole circumference, which receives a tendinous ring (*annulus tendinosus\**) ; this ring is situated without the choroid, and firmly adherent to the sclerotic coat ; on its inside it is connected with the origin of the venous layer of the choroid coat.

*Cornea* consists of layers which are more firmly united at its centre than at the circumference. The inner surface of it is covered with a serous membrane (*membrana humoris aquei*, *Wrisbergi*), the existence of which is easiest shown in eyes of aged persons ; this membrane can be separated from the innermost layer of the cornea, and differs from it in its greater tenacity and transparency. The latter quality it preserves in spirit ; the greater firmness and more express character of a serous membrane distinguish it from the layers of the cornea ; it is not so liable as these to ulceration, or to be destroyed by an ulcerative process, and, therefore, sometimes protrudes and forms the disease called by Professor Beer, '*Keratokele*.'

\* *Doellingeri descriptio oculi humani. Wurceburgi.*

*Iris.* Having subjected the iris to maceration, as I knew that by such a process the choroid coat might easily be divided into two layers, I succeeded in doing the same with this membrane. To such a division of the iris into two layers I was led by the observation of some writers on the anatomy of the eyeball, that the membrane of the aqueous humor is continued over the iris into the posterior chamber ; but with them it has been a mere supposition, and not proved by dissection. If there is such a continuation over the iris, this membrane must be divided into two layers ; in ascertaining this I succeeded, and shall now endeavour to give a description of my dissection. I performed the division more easily from the pupillar margin of the iris, where this membrane is thicker, and at this place I could evidently distinguish the turn which is formed by the anterior layer of the iris continued into the uvea ; betwixt these two membranes I saw distinctly the nerves and vessels distributed in a tortuous manner. Both membranes appeared somewhat transparent. The anterior layer, constituting the forepart of the iris, secretes on its inside and between the two layers a pigmentum, which exhibited itself to me in dark eyes, darker than in light ones. From the remarkable difference of this pigmentum in its color, I am inclined to think, that the different color of the iris particularly depends upon it, which then only can appear evident, when the uvea secretes its pigmentum. The anterior layer is afterwards continued to the tendinous ring, where it unites with a serous membrane, which I consider the origin of the venous layer of the choroid coat, under an acute angle. The posterior layer or uvea secretes on its back part the pigmentum nigrum ; when this pigmentum is removed, there appear small white processes going off from the ciliary processes to the uvea, being continued from the ciliary towards the pupillar margin, but not quite reaching the latter ; these processes are, like the ciliary processes, more distinctly seen in dark eyes, and differ from them only in their smallness.

*Choroidea* is easily divided into two layers, after maceration of several days. 1. The outer layer, or *choroidea, stricte sic*



*dicta*, is the thinner serous membrane, in which the ciliary veins are distributed to form the vasa vorticosa; it appears more distinct at its origin on the inside of the tendinous ring, where it unites with the anterior layer of the iris, and exhibits here evidently in its transparency the nature of a serous membrane. I think that this origin of the venous layer of the choroid coat has been described by Duverney as a peculiar serous membrane, covering the choroid coat. This venous layer appears more pallid at its beginning, on account of the ciliary ligament situated under it, and the deficiency of pigment; just behind the ciliary ligament it is perforated by the ciliary nerves and vessels of the iris. 2. The inner layer, or ruyschiana, is firmer, and secretes its pigmentum nigrum in the back part of the eyeball on its outside; as soon as it reaches the origin of the zonula ciliaris, it forms the ciliary body (*corpus ciliare*), which begins with a dentated margin, and secretes here its pigmentum nigrum on the inside; hence the impression of it appears on the zonula ciliaris. Professor Beer distinguishes the posterior part of it as the '*pars non plicata corporis ciliaris*,' which is larger on the temporal than on the nasal side of the globe, on account of the retina advancing more forward on the nasal side, as the optic nerve enters more on that side of the globe. This *pars non plicata* is the very part through which the needle is brought into the vitreous humor in operations through the sclerotic coat, and it is the part which is united by cellular tissue with the zonula ciliaris, to which the *processus ciliares* have no adhesion in the human eye. Professor Beer calls the anterior part of this body, *pars plicata*, to the formation of which the ciliary processes contribute. The ruyschiana, after having formed this body, continues forward—having to its inner surface firmly united the ciliary processes, and to its outer the ciliary ligament—to the back part of the iris into the uvea; so that I consider the ruyschiana as the mere continuation of the uvea. This continuity is not disturbed, after having separated the ciliary processes with their origins; and the appearance of both membranes is completely the same, each exhibiting the nature of a thin serous membrane.

Having now described the choroid coat and the iris, and the connexion of them anterior to the tendinous ring, I must here remark, that when this tendinous ring is separated from the sclerotic coat, the venous layer of the choroid and the anterior lamina of the iris form one membrane turning towards the cornea for the space of about one line ; which appearance makes me believe, that it is the membrane of the aqueous humor, though I could not follow this membrane further by dissection, so as to show clearly its continuation. Yet it is evident that this membrane divides before the tendinous ring into two, the outer and posterior forming the venous layer ; the inner and anterior, forming the iris, continues, uvea and ruyschiana. That it is a division of this membrane in these two different directions, I conclude from the finer structure of each of these two membranes than of that before the tendinous ring. The expressed character of a serous membrane in the latter, and its similarity in structure with that of the aqueous humor, make me believe, that it is really the membrane of the aqueous humor itself ; pathological observations prove also such a contiguity to the iris, i. e. the corneitis so quickly followed by iritis, and *vice versa* ; so that the primary inflammation of the cornea is denied by eminent pathologists.

As to the nature of the iris, choroidea, and ruyschiana, I am inclined to think that they are of a serous kind, from the appearance which they exhibit in their natural state, and the more so, from the morbid alterations to which they are subjected during inflammation, which is most evident in the iris, viz. the disposition to throw out coagulable lymph even in the slightest degrees of inflammation, and thus to produce an adhesion of the uvea to the capsule of the lens, or to close the pupil entirely ; in other cases to form partial or total adhesions of the iris to the cornea. The tubercles in the syphilitic iritis, which Beer calls condylomata, and appear at the ciliary or pupillar margin of the iris, are of a more or less brownish and red color, which variety of color depends on the smaller or greater organization of the coaguable lymph.



In a higher degree of iritis, though more rarely, there may be formed an abscess, which occupies, as Beer observes, the middle of the iris ; in these respects, as to the liability of adhesive inflammation, and rare occurrence of suppuration, in the substance of the iris, it coincides in its nature with that of serous membranes. The same liability to adhesion we may observe in the choroid coat, when a dissection of the eyeball is performed after a deep-seated inflammation, when we shall find not rarely adhesions of the venous membrane to the sclerotic coat, but more frequently of the ruyschiana to the retina ; or after the operation for cataract, adhesion of the zonula ciliaris to the ciliary processes. The difference of function in the iris and ruyschiana, though they are the continuation of a serous membrane, I explain from the addition of other parts, as in the iris, of the ciliary nerves and vessels, to which I think the motion of the iris is to be attributed ; and in the uvea and ruyschiana, from the other ciliary arteries, which secrete the pigmentum nigrum.

*Ligamentum ciliare*, is a cellular substance of a conical shape, situated between the choroidea and ruyschiana, just above the corpus ciliare and ciliary processes, its basis turned to the iris, its apex backwards. It is of various colors ; in light eyes it is lighter, looser, and larger. As to the use of it, I think it is for defending the vessels and nerves of the iris, which go through it. These vessels and nerves are situated at the back part of the globe without the venous layer of the choroid coat, and as soon as they reach the ligament perforate this membrane.

*Processus ciliares*, are situated on the inside of the ruyschiana, occupying nearly the anterior half of the ciliary body ; they originate by their bases in the angle formed by the uvea, continuing into the ruyschiana ; they are situated outwards and backwards ; their attenuated termination is where the connexion of the zonula ciliaris with the ciliary body begins. They may be separated from the corpus ciliare by the help of a needle, and elevated. There appear some larger and some

smaller processes, and two commonly arise together. The origins of these processes form the boundary of the posterior chamber in its entire circumference. From this description, it is evident that the depression of the crystalline lens into the posterior chamber (as described by some oculists) cannot be done without a violent injury to the ruyschiana or iris. From these processes arise smaller ones, continuing to the back part of the uvea. As to these processes, I have still to remark, that they are described by some anatomists as connected with the choroid coat, but I could not find such a connexion. That they are formed by the ruyschiana, I can also not admit, because, 1. they may be separated and raised from the corpus ciliare without injuring the continuity of the ruyschiana; 2d. they are different in their structure, endowed with greater firmness and elasticity; 3d. they do not secrete pigmentum nigrum in the human eye. I consider these processes as consisting in themselves of a different structure from that of the membranes of the eyeball; they appear to me very elastic, and of a substance which I might call one betwixt tendon and ligament. These processes are covered with a cellular tissue, which I have seen fully injected; this cellular tissue seems to be the secreting apparatus of the aqueous humor in the posterior chamber.

*Retina*, covers the corpus hyaloideum, and is connected with it at its back part, after having entered the eyeball, but much firmer in its connexion at the anterior part, where the zonula ciliaris originates at the dentated margin of the ciliary body; here it forms the ora serrata, which is considered the end of the retina. Professor Doellinger describes a thinner continuation under the zonula ciliaris as far as the canal of Petit. I repeated these dissections, and found that there exists a substance exhibiting itself as cellular texture, destitute of medullary substance; whether this is mere cellular texture, or a continuation of the inner layer of the retina, is uncertain. As to the membrane described by Jacob in a late



number of the Philosophical Transactions, I have seen it in brutes ; it appears evidently of a serous nature.

*Zonula Ciliaris.* An accurate description of this part has been given by Professor Doellinger (über das Strahlen-blättchen), of which I shall mention shortly the anatomical facts. The zonula ciliaris is situated under the corpus ciliare, and the serrated margin of that body denotes its commencement, from which place it goes to the anterior capsule, and unites with it intimately. The ruyschiana is easily separated from the retina until it reaches the zonula ciliaris ; it is united with this by firm cellular texture. The zonula Zinnii has about three lines of breadth, has an anterior and posterior margin, an outer and inner surface ; the outer one is connected with the corpus ciliare ; the inner surface is connected with the attenuated continuation of the retina ; the anterior part of the zonula is free. The canalis Petiti is formed by the zonula Zinnii and hyaloid membrane ; the hyaloid membrane is connected with the posterior capsule of the lens more backward, and the zonula Zinnii more forward with the anterior capsule, so as to leave a triangular space. The zonula consists of fascicles, which appear more evident when the canal is filled with air (Zinn) ; these fascicles are connected with the serrate prominences of the retina. Since Winslow's time, this zonula is derived from a splitting of the hyaloid membrane ; but the following observations are opposed to this opinion : 1. The hyaloid membrane has such a thinness, that it cannot be divided into two layers, as Zinn had already observed. 2. The zonula has a quite different structure ; the hyaloid membrane has nothing fibrous. 3. The hyaloid membrane is at the posterior convexity of the lens not thinner than in the other regions of the vitreous humor. 4. Between the corona ciliaris and hyaloid membrane is situated the continuation of the retina under the zonula. I have to mention last, the distribution of vessels in the zonula, which also proves that it is not a continuation of the hyaloid membrane ; the anterior part of the zonula obtains its vessels with the anterior capsule

of the lens ; the posterior part of the zonula from the arteria centralis after its division upon the posterior capsule of the crystalline lens, and forming here an anastomosis with the other vessels ; and finally the zonula has been injected with the capsule, but the hyaloid membrane has not shown the least trace of injection.

*Lens crystallina*, is included in a proper capsule ; the anterior is firmer than the posterior, which depends on the connexion with the zonula ciliaris, and obtains its vessels from the corpus ciliare. In the posterior capsule is distributed the arteria centralis, and it is connected with the hyaloid membrane by loose cellular tissue.

*Humor vitreus*, consists of cells, filled with a fluid like aqueous humor. It obtains its vessels from the retina, and the central artery going to the capsule gives off the arteria lateralis Albini. When the cells are destroyed, they are not restored, but the space is filled by a fluid which resembles aqueous humor.



of the testis, the formation part of the vessels from the external  
 testis after its division upon the posterior capsule of the  
 crystalline lens, and forming here an anastomosis with the  
 other vessels; and finally the vessels has been injected with  
 the capsule, but the hyaline membrane has not shown the  
 least tendency to rupture, and remains as a firm membrane  
 and, within the capsule, the vessels are seen to be in a state of  
 tension, and is included in a proper capsule; the an-  
 stomosis is distant from the posterior capsule, which is separate from the  
 testis with the vessels, and obtains its vessels from  
 the dorsal artery, in the posterior capsule is distinguished  
 the anterior capsule, and it is connected with the dorsal  
 artery by some cellular tissue.

When injected, consists of cells, filled with a fluid, the  
 aqueous humor. It obtains its vessels from the testis, and  
 the central artery going to the capsule, and the vessels  
 of the testis. When the cells are destroyed, they are  
 not restored, but the space is filled by a fluid which remains  
 the aqueous humor.

The capsule is a firm membrane, and is not easily ruptured, and  
 when it is ruptured, the fluid is not restored, and the space is  
 filled by a fluid which remains the aqueous humor. The capsule  
 is a firm membrane, and is not easily ruptured, and when it is  
 ruptured, the fluid is not restored, and the space is filled by a  
 fluid which remains the aqueous humor. The capsule is a firm  
 membrane, and is not easily ruptured, and when it is ruptured,  
 the fluid is not restored, and the space is filled by a fluid which  
 remains the aqueous humor. The capsule is a firm membrane, and  
 is not easily ruptured, and when it is ruptured, the fluid is not  
 restored, and the space is filled by a fluid which remains the  
 aqueous humor. The capsule is a firm membrane, and is not easily  
 ruptured, and when it is ruptured, the fluid is not restored, and  
 the space is filled by a fluid which remains the aqueous humor.






Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.



Fig. 6.





Fig. 7.









## EXPLANATION OF THE PLATES.

---

### PLATE I.

*Fig. 1.* Simple acute inflammation of the conjunctiva. See page 91.

*Fig. 2.* (*Inserted by the Editor.*) Inflammation of the iris, strongly contrasted in character with that of the conjunctiva. In the former, the white line around the cornea, the arrangement of the inflamed vessels in straight lines, running in radii from the cornea, the intensity of the inflammation near the cornea, and its gradual diminution as you recede from it, together with the evidently deeper seat of the enlarged vessels, are easily distinguished from conjunctival inflammation, in which the redness is most intense upon the eyelids, and diminishes as you approach the cornea; the vessels are superficial; evidently in the conjunctiva and not moving with the motions of the eyeball, and not arranged in right lines like those of the sclerotica, which are enlarged in iritis.

*Fig. 3.* A plan to show the cicatrization of an ulcer of the cornea, whether pustular or interstitial, and communicating with the surface by a



fasciculus of vessels carrying red blood. See page 109.

*Fig. 4.* The state of strumous nebula with vessels in the form of radii overshooting the cornea. These are not continuous with, but distinct from the vessels of the conjunctiva, and beneath it. See page 110.

I have seen cases of the prolongation of these vessels even to the centre of the cornea, organizing the nebulous deposit, affecting both eyes of the same patient, and occasioning total opacity and blindness, perfectly restored by continued ptyalism. The first change was the breaking and clearing of the dense deposit of lymph, then the fading away of the rose-colored zone at the verge of the cornea, and lastly, the gradual disappearance of the vessels on the cornea. See page 306 and seq.\*

*Fig. 5.* A plan of the anastomosis of colored vessels upon an opaque cornea; 'chronic inflammation with vascular cornea.' See page 110.

\* The converse of this statement has been repeatedly observed by the Editor in severe corneal inflammation; although, at the same time, he does not deny the accuracy of the author's description in a certain proportion of cases. Where the vessels have been prolonged over the margin of the cornea, and have produced general opacity of that membrane, in general, the vessels first begin to diminish in size

*Fig. 6.* Aphthous or pustular inflammation of the conjunctiva. Two large aphthæ are seen, one on either side of the cornea, situated nearly in the transverse axis of the globe. This figure was selected for the purpose of showing the peculiarity of two distinct orders of vessels bounded by the pustules. The upper segment of the hemisphere presents the ordinary superficial vessels of the conjunctiva, having the areolar distribution. The lower are the straight vessels penetrating the sclerotica, and appearing through the transparent conjunctiva. This two-fold arrangement is seldom so distinctly seen. See pages 95 and 128.

*Fig. 7.* The blue tumor of the sclerotica, 'staphyloma scleroticæ,' accompanying various disorganized states of the globe. See page 130.

The figures 1, 4, 6, and 7, in this plate, are from nature. The three former have an appearance of unnatural magnitude, from the artificial elevation of the upper and the depression of the lower lid, for the purpose of exhibiting the conjunctival surface around the cornea.

and gradually disappear, leaving the deposited lymph in the centre of the cornea, to be slowly removed by a process which may continue many months. In such cases, even in forty-eight hours, the whole of the circumference of the cornea may recover its transparency, and the red vessels completely disappear, while the centre still remains densely opaque.—*Editor.*



PLATE II.

*Fig. 1.* Carcinoma of the conjunctiva affecting the globe and eyelids.


This very accurate representation of the disease was taken from the eye of a patient of mine in St. Thomas's Hospital, by Dr. Zuckerbecker, a very intelligent physician now settled at Moscow. See page 100.

*Fig. 2.* A section of the fungous growth represented in situ, *Fig. 4.*

*Fig. 3.* The state of central slough of the cornea in the last stage of acute suppurative inflammation.

*Fig. 4.* A peculiar fungous growth interstitial to the conjunctiva and cornea. See page 102.

This figure represents the appearance of the disease, as it was presented to me in the person of an elderly lady from Somersetshire, and the idea I then formed of it was that the fungus had originated from the iris, or choroid tunics, consequent to a slough of the cornea. Finding that the sclerotic tunic was sound, I proposed the excision of the tumor by an operation similar to that for staphyloma. This was executed with





## PLATE II.

*Fig. 1.* Carcinoma of the conjunctiva affecting the globe and eyelids.

This very accurate representation of the disease was taken from the eye of a patient of mine in St. Thomas's Hospital, by Dr. Zuckerbecker, a very intelligent physician now settled at Moscow. See page 100.

*Fig. 2.* A section of the fungous growth represented in situ, *Fig. 4.*

*Fig. 3.* The state of central slough of the cornea in the last stage of acute suppurative inflammation.

*Fig. 4.* A peculiar fungous growth interstitial to the conjunctiva and cornea. See page 102.

This figure represents the appearance of the disease, as it was presented to me in the person of an elderly lady from Somersetshire, and the idea I then formed of it was that the fungus had originated from the iris, or choroid tunics, consequent to a slough of the cornea. Finding that the sclerotic tunic was sound, I proposed the excision of the tumor by an operation similar to that for staphyloma. This was executed with

Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.




Fig. 5.



Fig. 6.







facility, and the patient quickly recovered, the remaining portion of the globe collapsing in the orbit. The surface had the deep blue tint, and the currant or berry-like appearance denoted, likening it to what is termed by the Germans, 'Staphyloma racemosum,' but it proved to be totally different in its nature from that disease.


The section *Fig. 2.* is shrunk from immersion in spirit; it is vari-colored and of unequal consistence; in some parts pulpy, in others firm; the sclerotica is not altered in its structure; the iris is thickened by inflammation, the pupil closed, and the capsule of the lens, which is shrunk and opaque, adherent to the uvea. The red line is the section of the iris, and the yellow substance beneath and adhering to it represents the shrunk crystalline. These parts were quite free from the disease, and changed only by compression. The cornea which is seen at the base of the fungus, has lost its lamellated structure, and hence appears firmer; its surface is rough and has a brownish tint, as if beginning to degenerate into the morbid mass which lies above it. This not only covers the cornea, but at one part a little overlaps the sclerotica. The section discovers a subdivision of the larger into lesser lobes, separated by whitish lines intersecting the mass perpendicularly. The lobes differ in structure and in color, as if originating in cells and distinct from each other. In one part adjoining the sur-



face a whitish spot is conspicuous, of a cartilaginous hardness. The fungus is covered by a membrane easily torn; if this be not the conjunctiva, no trace of that membrane remains; but I am disposed to think from its relation to the tumor, and the continuity of the conjunctiva scleroticæ with this membrane, at that part of the circumference of the cornea where the pulpy mass has encroached upon the sclerotica, that the covering membrane is formed of the conjunctiva in an altered state. The intimate adhesion of the fungus to the covering membrane, the total change in texture of the conjunctiva, and the fuller evolution of the disease next the surface, lead me to suppose that it originates from that membrane; and from its lobular arrangement I conclude, that the morbid growth occupies the cells of the connecting membrane indicated by the white lines intersecting it.

*Fig. 5.* Abscess of the eyeball terminating in ulceration and death of the cornea and disorganization of the globe. See page 228.

*Fig. 6.* Scirrhus of the lacrymal gland. See page 244. The deformity was removed by extirpation of the gland.





*Fig. 1.*



*Fig. 2.*



*Fig. 3.*



*Fig. 5.*



*Fig. 6.*



*Fig. 7.*



*Fig. 4.*



## PLATE III.

*Fig. 1. and 3.* represent in different positions the eye of a child affected with the malignant fungus, prior to the opacity of the cornea; in which are observable the change of figure of the globe and annihilation of the anterior chamber, by the bulging of the opaque lens and iris.

*Fig. 2.* is a section of the same eye.

These drawings having been made immediately upon the removal of the eyes after death, represent accurately the recent appearances, but by a minute dissection, after hardening in spirit, I have been enabled to trace some vestiges of the several textures which the mere sections do not exhibit, in a manner which would have given the representation, if that had been possible, additional interest, as it throws much light on the origin and progress of the diseased growth.

A substance answering the description of the 'sarcoma medullare,' occupies the upper and back part of the eyeball, by which the vitreous humor, which has undergone a similar degeneration, is compressed and pushed forward and downward. The scleretica is in a perfectly



Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.



*PLATE III.*

*Fig. 1.* and *3.* represent in different positions the eye of a child affected with the malignant fungus, prior to the opacity of the cornea; in which are observable the change of figure of the globe and annihilation of the anterior chamber, by the bulging of the opaque lens and iris.

*Fig. 2.* is a section of the same eye.

These drawings having been made immediately upon the removal of the eyes after death, represent accurately the recent appearances, but by a minute dissection after hardening in spirit, I have been enabled to trace some vestiges of the several textures which the mere sections do not exhibit, in a manner which would have given the representation, if that had been possible, additional interest, as it throws much light on the origin and progress of the diseased growth.

A substance answering the description of the 'sarcoma medullare,' occupies the upper and back part of the eyeball, by which the vitreous humor, which has undergone a similar degeneration, is compressed and pushed forward and downward. The sclerotica is in a perfectly



healthy state, as are also the cornea, iris, and a considerable part of the choroid; the lens absorbed, its capsule firmly adherent to the uvea. The diseased mass is evidently deposited between the venous and arterial layers of the thickened choroid coat; the former adherent to the sclerotica, being situated external to the morbid mass; the latter (which has been described as the tunica ruyschiana), separated and protruded by it, is denoted by a line crossing the section obliquely. Here no pigmentum nigrum is secreted. The corresponding posterior part of the retina is destroyed, the anterior is adherent to the choroid. At the back of the eyeball is a medullary tumor contained in a capsule, formed by the surrounding cellular texture. A similar tumor is formed by the optic nerve at its entrance. The texture of these is somewhat firmer than that within the ball, and between these tubera and the internal diseased mass there is no communication.

*Fig. 4.* represents the other eye of the same subject, when by the progress of the disease the cornea had perished by ulceration, and the fungus, represented *Fig. 5.*, was just about to protrude. The leaden tint of the extenuated sclerotica is characteristic.

*Fig. 5.* represents the contents of the orbit on removal after death.

*Fig. 6.* is a section of the fungous mass.

The optic nerve filled with the morbid deposition has its neurilema thickened; and a similar substance occupies the cellular texture within the orbit. The white lines bifurcating from the extremity of the nerve, represent the sclerotica thickened. On both sides of the entrance of the optic nerve, a mass is formed on the interior of the sclerotica, resembling that on its outside, except that its texture is firmer. Forwards, on the right side of the figure, the sclerotica is distinctly split into two layers by the pulpy substance which has insinuated itself between them. This corresponds to the upper half of the globe.

The plate, which represents only a recent section, does not illustrate the remaining points of the description. They have been since made out by subdivision and careful separation of the several parts of the mass from each other. As they are highly curious I subjoin them.

The interstitial deposit is seen in the whole anterior circumference of the sclerotica, but the layers of this membrane are less widely separated from it at the inferior part of the ball. At the posterior part, where the sclerotica is single, the membrane is much altered, but it has nowhere given way, so that the diseased masses upon its opposite surfaces have no direct com-



munication. By the mass formed on its inside, the choroid coat is distinctly seen to be protruded forwards. At the anterior part, the sclerotica is firmly connected with the thickened choroid coat. The entire centre of the cornea has perished by an ulcerative process, but a fragment of corneal lamella remains to denote its place, and behind it is a portion of membrane adhering, and resembling the iris much altered in its structure. The softest part of the fungus is connected with this membrane, and between it and the choroid no trace appears of the humors or retina. This anterior mass is distinguished from the posterior growth of the sclerotica, from which it is separated by the choroid, by its softer texture and darker color. Hence it appears that the morbid substance within the ball consists of two different formations; first, of the sclerotica degenerated on its exterior and interior surfaces, and likewise in its substance; secondly, of the choroid degenerated on its interior surface.

*Fig. 7.* This figure represents the section of the eye of a child, aged eight months, which I extirpated several years ago. The subject of the operation has since enjoyed perfect health. See page 212, where *Fig. 4.* is referred to by a misprint.

The cells of the vitreous humor are filled

with an opaque lardaceous substance, by which the lens was slightly protruded, and the iris rendered convex. The eyeball was but little increased in size or altered in figure. The sclerótica was in some parts thinner than usual, and had a bluish hue from the preternaturally firm adhesion of the choroid coat. The retina was for the most part absorbed, the other tunics perfect, and the optic nerve free from disease.

There is no evidence to prove this change of structure malignant, although the external appearances closely resembled those of *Fig. 3*.



## PLATE IV.

(BY THE EDITOR.)

*Fig. 1. and 2. are delineations of the eye before and after the operation for artificial pupil in the following case.*

In the month of November, 1821, Mr. R. Wortendyke, aged sixty-three years, a farmer from Sussex county, New Jersey, applied to me, blind of both eyes. When he was three years of age, he received a wound in the right eye from a fork, which penetrated the cornea near its lower margin, pierced the iris, and wounded the crystalline lens. The consequence was, opacity and subsequent absorption of the lens, and adhesion of the iris to the lower edge of the cornea. In this situation I first saw the eye, sixty years after the accident happened. The pupil was extremely small, and drawn down to the edge of the cornea; but adhering only at this point, and free at its upper edge. The lens had disappeared and the capsule remained behind the pupil, much thickened and opaque. The patient could see light with this eye, but was totally unable to distinguish objects.

The left eye became blind from an injury of much later date. At the age of fifty-nine, four years previously to my seeing him, Mr. Wortendyke received a severe blow upon this eye, from an ear of wheat. In consequence violent inflammation, sloughing of the cornea, protrusion of the iris, and total destruction of vision took place. The eye recovered from all the acute symptoms, but the greater part of the cornea remained densely opaque, and the iris partially adherent to it.

Fig. 10




Fig. 9



Fig. 8



Fig. 4



Fig. 3



Fig. 5



Fig. 6



Fig. 7



Fig. 8






## PLATE IV.

(BY THE EDITOR.)

Fig. 1. and 2. are delineations of the eye before and after the operation for artificial pupil in the following case.

In the month of November, 1831, Mr. J. F. Ward, Jr. aged sixty-three years, a farmer from Falmouth, Me., came to the office of the Editor, with a complaint of blindness. He stated that three years of age he received a wound in the right eye from a flock, which penetrated the cornea and entered the iris, pierced the iris, and wounded the crystalline lens. The consequence was, opacity and subsequent absorption of the lens, and adhesion of the iris to the lower edge of the cornea. In this situation I first saw the eye, thirty years after the accident happened. The pupil was extremely small, and drawn down to the edge of the cornea; but adhering only at this point, and free at its upper edge. The lens had disappeared and the capsule remained behind the pupil, much thickened and opaque. The patient could see light with the eye, but was totally unable to distinguish objects.

The left eye became blind from an injury of much later date. At the age of fifty-nine, four years previously to my seeing him, Mr. Worleudyke received a severe blow upon this eye, from an ear of wheat. In consequence violent inflammation, sloughing of the cornea, protrusion of the iris, and total destruction of vision took place. The eye recovered from all the acute symptoms, but the greater part of the cornea remained densely opaque, and the iris partially adherent to it.







The patient applied to me with the expectation of having some operation performed upon this eye, as he had long since given up all hope of receiving any benefit to the other. The right eye, however was evidently in the best situation for successful operation, and the only unfavorable circumstance about it was, the great length of time which it had remained blind. I determined, however, to make an attempt upon this eye, although it was not without difficulty that the patient could be induced to consent to it; so impossible did he consider it, that any operation upon it should succeed.

The operations which it required were sufficiently simple. It was first necessary to remove the opaque capsule. This was effected without difficulty by means of Saunders' needle. When the eye had recovered from this operation, the patient had some degree of vision, but the pupil was so small and distant from the centre of the cornea, that the sight was extremely imperfect. It was necessary, therefore, to perform some operation for artificial pupil. For this purpose an iris knife was introduced, about two lines behind the cornea, and after puncturing the coats of the eye, its point was pushed through the iris into the anterior chamber, about two lines from its ciliary margin: the point of the instrument was then directed towards the small pupil already described, and all that part of the iris was divided, which lay between this pupil and the point at which the instrument was introduced. The iris immediately retracted, and left a large oval pupil, with its transverse axis running obliquely across the eye. The patient recovered, with very little supervening inflammation, and the pupil remained full as large as the natural one. By degrees he gained so much sight, that he was enabled to distinguish the countenances of his friends, and attend to his usual avocations on his farm. When I last saw him, he had still further improved; but although his sight was sufficiently good for all ordinary purposes, he was



unable to read. The success of the operation, however, was fully as great as could have been expected, under the circumstances.

*Fig. 1.* shows the eye before the operation for artificial pupil, but after the removal of the opaque capsule. The original pupil is seen drawn down and adhering to an old cicatrix, and so small as to be useless. *Fig. 2.* is the same eye after the operation.

*Fig. 3.* A central artificial pupil made in the eye of J. Thomson.

This patient was admitted into the New York Eye Infirmary, May 25th, 1824. He had been the subject of severe iritic inflammation, which had closed both pupils. In the right eye the blindness was complete, the only remaining aperture in the iris being filled with a mass of lymph. In the left, a small portion of the pupil remained free, and gave the patient sufficient sight to enable him to find his way without a guide, about places to which he was accustomed. The situation of the right eye was most unfavorable to the success of an operation, but as the other was of some use, it was deemed most prudent not to risk any operation upon it, until after an experiment upon the worst eye.

The right eye, therefore, was selected for operation, which was performed in the mode originally practised by Cheselden, but now revived and improved by Sir W. Adams. The iris knife was introduced through the sclerotic and iris, as in the preceding case, and a section made across about two-thirds of the diameter of the iris, leaving a large clear pupil in its centre of an elliptical form.

But although the operation succeeded perfectly, as regarded the formation of a good pupil, it was of no advantage to the

patient. The eye proved entirely amaurotic, and remained perfectly blind. The case, however, was a striking illustration of the excellency of this mode of operating. The pupil was made with great facility, and remained afterwards permanent, without the least disposition to contraction. The drawing, made from recollection, will give a better idea of the appearance of the eye after operation, than any description I can give.

*Fig. 4.* The result of a somewhat similar operation upon the eye of a patient, whose case follows :

Peter Simpson, aged forty-two years, in the month of July, 1822, received a slight blow from a hook attached to a pulley, upon the right eye, which caused immediate blindness. He had lost the left eye several years before. In consequence of the last accident, he became a patient of the New York Hospital, whence he was discharged, after the expiration of two months, having recovered sufficient vision only to distinguish light from darkness. During this period he had suffered severely from the violent inflammation which followed the accident.

In September, 1823, he was admitted a patient into the New York Eye Infirmary, with his eye in the following condition. In consequence, apparently, of a rupture of the cornea, from the blow he had received, the iris had prolapsed, was drawn over to the inner side of the eye, and adhered to that edge of the cornea, totally obliterating the natural pupil ; but leaving an open space through the iris, at the inner edge of the cornea, which would have been sufficient for good vision, if that membrane had been transparent. Unfortunately, however, the injury had so far rendered this part opaque, as nearly to make the pupil useless, and it was still further impaired by the presence of an opaque lens behind it. In consequence, the patient's vision was totally insufficient to



enable him to gain a livelihood, although he could, without much, difficulty find his way in the streets. From this inability to work, he had for the previous year been a tenant at the Alms-House. From this description it will be evident that the fibres of the iris were on the stretch, drawn towards the inner canthus, and also upwards and downwards, by their adhesion to corresponding points of the cornea. The case was, therefore, evidently favorable for the operation of simply dividing the fibres of the membrane ; and as the cornea was transparent in its centre, the most advantageous situation was afforded for a new pupil.

Accordingly, the operation was performed upon that principle. The iris knife was introduced through the sclerotic, at the usual situation for the introduction of the couching needle, the back of the knife being held towards the operator. When its point had arrived about two lines from the external edge of the iris, it was pushed forward through the membrane, and then carried on in the anterior chamber, as far as possible, towards the inner canthus of the eye. The iris was next divided by a back stroke of the instrument, in the motion of withdrawing it ; the fibres immediately retracted, and left a large pupil of a triangular form, opposite to the centre of the cornea.

It was now discovered, through the new pupil, that the lens had been partially dislocated by the accident, and thrown towards the inner canthus of the eye, remaining attached opposite to the junction of the cornea and sclerotic ; but its external edge projecting so far behind the new pupil, as to render it useless. It was, however, deemed most prudent to defer the operation for its removal to another time.

Little inflammation followed the operation ; and in a fortnight, the patient was able to bear the full light with no other screen than a common shade.

In the month of November following, another operation was performed, for the removal of the opaque lens. A puncture was made through the cornea, near its inner inferior margin, and a section made, corresponding to about one-fourth its circumference. The lens, already softened in texture, and somewhat reduced in size by its exposure to the aqueous humor, immediately glided through the opening, and left the new pupil perfectly clear and bright. This slight operation was immediately recovered from without any untoward accident. After the expiration of a few weeks, the patient returned to his occupation as a sawyer, and has ever since gained his own subsistence by his daily labour. He can now see so well without glasses as to pick a pin from the floor. Having never learned to read, the accuracy of his sight cannot be tested in that particular; but he can distinguish the difference of small letters, and trace them accurately.

*Fig. 5.* The appearance of the pupil in the best eye of Steinberg, alluded to in the note, page 374, after it had been dilated to its fullest extent by extract of stramonium. The very minute dark line, between the upper edge of the pupil, and a deposite of white lymph lying within it, appears to be the only situation in which light can pass to the retina.

*Fig. 6.* The same eye, before the application of the stramonium.

*Fig. 7.* J. M'Gilvra's eye before the operation: a dense cicatrix across the centre of the cornea, and the only remains of the pupil shown by a few dark lines in its original situation.

*Fig. 8.* The same eye, with an artificial pupil made in the lower half of the iris.



This patient was received into the Infirmary, October 27th, 1824. His right eye was completely blind from organic amaurosis, and had been in that situation about six years. The left had been deprived of vision by an accident which had happened sixteen years before. The eye had been wounded by a thorn, which had torn the cornea entirely across, in its transverse diameter, wounding at the same time the iris and capsule of the lens. In consequence, when these wounds healed the pupil was closed, and the cornea, iris, and capsule united in one firm cicatrix. The upper and lower portions of the cornea were perfectly transparent, and the iris apparently sound. These appearances are more fully shown by the annexed drawing. The patient was deprived of all useful vision, and could only perceive the motion of objects before him, or their interposition between him and the light,

Three operations had been performed upon the left eye, by an eminent surgeon in the northern part of this state, the exact nature of which I could not learn ; but they had the effect, as was afterwards proved, of destroying the lens, and thus rendering the subsequent operation less complicated.\*

In this case, it was evident, that a very favorable situation was offered for a new pupil, in the lower half of the iris. The fibres of this membrane, being adherent at opposite points, viz. at its ciliary margin below, and to the firm cicatrix above, it might fairly be calculated, that upon a simple incision, they would retract sufficiently to leave a good and permanent pupil. Accordingly, on the 1st November, the

\* It is perhaps a matter of doubt, whether the removal of the lens was the effect of absorption, excited by these operations, or by the original injury ; either being competent to this effect.

operation was performed, in a manner similar to that described in the last case. The iris knife was introduced through the sclerotic, somewhat below the transverse diameter of the eye, its point pushed through the iris about a line from its ciliary junction, and after carrying it forward to a corresponding point on the opposite side, the membrane was divided by a gentle back stroke of the instrument, as it was withdrawn. The fibres of the iris immediately retracted, and left a large pupil, such as is shown by the drawing.

The patient soon recovered from the operation, but although his sight was somewhat improved, it was still insufficient to answer any useful purpose. On very close examination, after the organ could bear exposure to a full light, I perceived some minute portions of the capsule of the lens, which obstructed the passage of the light through the new pupil. On the 20th November, these were removed by a second operation, so simple as not to require description. Still, the degree of vision gained was not very material, for several days after the eye could bear full exposure to light ; and I had almost despaired of eventual success in the case. About two weeks after the last operation, however, the sight suddenly began to improve, and gained so rapidly in the course of the following fortnight, that with proper glasses the patient could read with facility the title page of a book. The long disuse of the organ in this case, seemed at first to have rendered it incapable of performing its functions, after the obstacles to vision had been removed ; but, by degrees, it regained its powers. The patient was discharged from the Infirmary early in December.

In March following, four months after the last operation, I received from him the following letter.



Princeton, March 4th, 1825.

Dear Friend,

With much pleasure I inform you that my eyesight has gained so, that I can see to read small print ; as small as that of a pocket bible. My health is good, and I hope you enjoy the same blessing. I return you my humble thanks for your kind attention to me.

Yours with respect,

JOHN M'GILVRA.

DR. DELAFIELD.

The success experienced in this case, was certainly more than I expected. It was not the first instance, in which I learned not to despair of eventual success, when sight was not immediately recovered, after operations on the eye. A more striking one occurred in the case of a young lady, upon whom I operated for cataract about eighteen months since.

The cataract, which proved to be soft, was removed by lacerating the capsule, loosening the texture of the lens, and placing a portion of it in the anterior chamber. In two months after the operation, the whole of the lens and capsule were absorbed, and the pupil perfectly clear and bright ; but the eye was useless, and the patient could distinguish the form of no object with it. It continued in this situation, with very little change, more than eight months ; after which, it began rapidly to improve, and when I last heard from the patient, the sight was so perfect, that she could read and sew without glasses. I do not recollect any author who notices this important fact ; although it can hardly have escaped the observation of those who have had extensive opportunities of operating on the eye.

*Fig. 9.* The iris knife used in the foregoing operations, resembling a very small scalpel, cutting only on one edge,

and extremely sharp at its point. It is the instrument alluded to by Mr. Travers, page 364.

*Fig. 10. a.* Reisinger's double hooked forceps, noticed page 367.

*b.* A side view of a single limb of the instrument.

Reisinger's operation consists in separating the iris from the ciliary body, and strangulating it afterwards between the edges of the cornea; to effect which, he uses a very fine double hook, which by a slight pressure of the finger and thumb, is made to resemble a single one.

An incision is first made in the cornea near its outer edge, one and a half, or at most two lines in length, and if possible, three lines (one quarter of an inch) distant from that part of the iris which is to be separated.

“After the incision in the cornea is completed, the eye should be allowed to close for a moment to afford it rest, and prepare it for the subsequent proceedings, unless it has been previously fixed by an instrument.

“The operator should hold the hooked forceps nearly in the same manner as the cataract knife, the points of the hooks downwards, the thumb resting on the flat side of the shank which is facing the operator, and the first and second finger on the flat side of the opposite shank, so that the point of the second middle finger may reach the end of it. The handle must rest against the *radial* side of the first joint of the first finger; the little finger serving to steady the hand.

“The instrument is now drawn along the edge of the cornea, pressing gently and steadily with the convexity of the united hooks against the small incision, in order that the edges of the wound may be gently opened, and the closed



forceps insinuated into the anterior chamber ; then glide the instrument with the convexity of the hooks against the inner surface of the cornea, as far as the spot where the iris is to be separated ; taking care, however, to bring the hooks as near as possible to the ciliary edge of the iris. The forceps are now to be turned, so that the point of the hooks may be directed towards the iris ; then let the instrument be opened, so that the two hooks may be at least one line, and not more than two lines asunder ; press the convexity of the hooks against the ciliary ligament, sink the points into the iris, and close the forceps, at the same time gently drawing them towards you ; by these motions, which must be almost simultaneous, the iris is steadily seized and easily separated. The closed forceps are now to be further drawn out of the eye, the convex edge of the hooks being carefully turned towards the inner surface of the cornea to avoid pricking it. In this manner a considerable part of the iris will be separated, and a triangular pupil formed, of the size of at least one quarter of the iris, and which will reach the middle of the eye.

“ The handle of the forceps is now to be depressed, and as the convexity of the hooks glides out at the upper angle of the incision, withdraw the forceps, together with that part of the iris which has been laid hold of, so as to produce a pretty considerable procidentia iridis. The eye should now be immediately closed to assist the strangulation of the prolapsed iris by the pressure of the eyelid.

“ If the prolapsed portion of the iris should recede, it will generally happen from the incision having been made too large. In this case, it may probably be advisable to cut off the part of the iris which has been separated, and which must again be drawn out ; and so combine *Coredialysis* with *Coretomia*, to insure an artificial pupil of a proper size.\*”

\* Reisinger, in his description of this operation, part of which I have quoted, gives many more minute directions concerning it. I have extracted sufficient to show the nature of the operation, and to do more would unreasonably lengthen this article.

Maunoir's operation with the scissors is noticed by our author without any particular description of it. I have introduced it as described by Scarpa.

The scissors are made so small and delicate, as to be easily introduced through an incision in the cornea, with their blades forming a large angle with the handles.\* The upper blade, or that which is intended to pass through the anterior chamber between the cornea and iris, terminates in a small button. The lower one, with which the iris is to be pierced, has a very sharp lancet point.

“An incision is made in the cornea at its lower or lateral segment, as may be most convenient, of *half the extent* of that which is usually made for the extraction of the crystalline lens. Through this small opening in the cornea, the scissors are to be introduced closed, with the flat part in a line parallel to the transverse diameter of the iris; and as soon as the point of the instrument has advanced near to the great margin of the iris, that is to say, nearly opposite the small incision made in the cornea, it is gently opened and inclined in such a manner that the inferior pointed blade may perforate the iris, and run along the posterior surface of that membrane, until the small button of the upper blade reach the part where the cornea and sclerotica unite. The iris is then to be divided in its transverse diameter, by a single stroke passing as nearly as possible through its centre. This incision being executed, another is to be expeditiously made, so far diverging from the first, that the two incisions may form in the centre of the iris a triangular flap of the figure of the letter V, the apex being precisely in the centre of the iris, and the base near its greater margin.—On opening the eye operated on, five or six days after, the apex of the triangular shred is found to have retracted towards its base, leaving in the middle of the iris an artificial pupil of the form of a parallelogram.”

\* See Plate V. Fig. 5.



By means of the scissors, also, an operation may be effected, according to Scarpa, without injury to the transparent lens or its capsule, in the case of a "contraction of the natural pupil occasioned by the iris and pupil being stretched towards some point in the cornea. This happens, in general, in consequence of prolapsus of the iris through ulcers of the cornea." In this case, the lens and capsule often preserve their transparency, and the cornea is only opaque immediately around the adhering point.

For this operation, it is necessary to have the scissors of Maunoir, made with the point of each blade terminating in a button. Then the operation having been commenced as in the first instance, "one of the blades, by means of the small button, is introduced within the contracted natural pupil, and conducted behind the posterior surface of the iris, until the other blade, defended in the same manner, has reached the confines of the cornea and sclerotica. The iris is then to be divided in the form of the letter V, without at all injuring either the capsule or the lens, both of which have preserved their transparency."

How far this operation is practicable without injury to the lens or capsule, I cannot say, as I have never performed it; but I confess, I somewhat doubt the possibility of carrying an instrument between the iris and capsule, without lacerating the latter membrane. If practicable, the operation is certainly very beautiful.

If a hard lens be supposed to be present, Maunoir operates with his scissors in a manner similar to the first case, but divides, with one stroke of the instrument, the iris, capsule, and lens, which is afterwards extracted through the incision in the cornea, either by means of a forceps or scoop, or simply by slight pressure on the eyeball.





*Fig. 1.*




*Fig. 2.*



*Fig. 3.*



*Fig. 4.*



*Fig. 5.*

## PLATE I

(BY THE EDITOR.)

*Fig. 1.* Front and side views of the searching needle employed by the author. The only material difference between this instrument and Scarpa's needle is that it has a blunt point, instead of a sharp one, and the other needle is pointed, instead of being flattened behind, and being bent back wider, it has a slightly greater length. The line is the size of depression.

*Fig. 2.* Scarpa's couching needle. This instrument is generally used in this country of a much larger size than recommended by Scarpa. I have seen it at least four times the proper size, and of course, unnecessary violence committed by it upon the eye. The delineation in the plate shows the precise size recommended by the inventor.

*Fig. 3.* Saunders' needle, for the anterior operation, more particularly useful in congenital cataract in infants.

*Fig. 4.* Beer's cornea knife for extracting the cataract.

*Fig. 5.* Maunoir's iris scissors, described in the explanation of Plate 4.



Fig. 1.



Fig. 2.




Fig. 3.




Fig. 4.





Fig. 5.

*PLATE V.*

(BY THE EDITOR.)

*Fig. 1.* Front and side views of the couching needle employed by the editor. The only material difference between this instrument and Scarpa's needle is, that it has a lancet point, instead of a triangular one, and therefore makes an incised, instead of a punctured wound, and being somewhat wider, it less readily passes through the lens in the act of depression.

*Fig. 2.* Scarpa's couching needle. This instrument is generally used in this country of a much larger size than recommended by Scarpa. I have seen it at least four times the proper size, and of course, unnecessary violence committed by it upon the eye. The delineation in the plate shows the precise size recommended by the inventor.

*Fig. 3.* Saunders' needle, for the anterior operation, more particularly useful in congenital cataract in infants.

*Fig. 4.* Beer's cornea knife for extracting the cataract.

*Fig. 5.* Maunoir's iris scissors, described in the explanation of Plate 4.



## PLATE I.

BY THE AUTHOR.

Fig. 1. Front and side views of the compound microscope as proposed by the author. The only material difference between the instrument and Hooke's is, that it has a barrel joint, instead of a hinge joint, and thereby it makes an inclined motion of a quadrant, which is not a disadvantage, when it is used to pass through the lens in the end of depression.

Fig. 2. Another compound microscope. This instrument is generally used in the country of a more larger size than recommended by Hooke. I have seen it at least four times the proper size, and of course, unnecessary violence is done by it upon the eye. The objection to the plate shows the proper way of using it for the purpose.

Fig. 3. Another, which is the simplest of the three, and is particularly useful in examining small objects.

Fig. 4. Hooke's camera lucida for extending the object.

Fig. 5. Hooke's camera lucida, described in the explanation of Plate 4.

# NOTES.

---

## NOTE A.

THE relations of structure in every organ are such as to render it impossible that any texture can long be singly affected. But it seems reasonable to consider that texture as the proper seat of the morbid action which presents the earliest, or strongest, or exclusive signs of it. I do not therefore adopt the term "*Ophthalmitis Scleroticæ*," as applied to all the deeper seated inflammations; nor do I consider the corneal and iritic inflammations merely symptomatic, and as having their proper seat and base in the sclerotica (although in rare instances it may be so), because from the affinity and vascular connexion of the parts, that membrane presents the ordinary appearances of inflammation during their existence. However inflammation may deviate from the ordinary course and the order of local relation, and, however complicated its results may be, it is to be referred to the texture primarily and principally affected, and is thence properly denominated; as in the instances of scleritis to which this note refers.

Intolerance of light is considered by some respectable authors to be diagnostic of sclerotic inflammation. This is not my opinion, having in numberless instances witnessed that symptom in the most aggravated degree in the absence of every external sign of inflammation. But on the other hand. I am willing to admit, on the ground above stated, that sclerotic inflammation is often present in such instances.

The sympathy of the sclerotica and the ligamentous capsules with the urethra in gonorrhœa, is as unquestionable as that which has been more generally observed, because more



frequently occurring, between the latter and the synovial membrane. I have seen cases so nearly resembling that described by Dr. Vetch in his late valuable treatise,\* under the head of 'gonorrheal ophthalmia' (page 243), that I can vouch for the accuracy of the description. This phenomenon it is not so easy to explain as the coincidence of the suppurative inflammation of the conjunctiva with acute gonorrhea, which, notwithstanding the apparent contradiction of some experiments, I am convinced originates from contact, as is indeed proved by the history of the ophthalmia of infants, and by the fatal effects which have unfortunately fallen more than once under my observation, of an accidental application of morbid matter to the sound organ.

The disposition between remote parts to be reciprocally affected, admits of two modes of explanation. First, by a partial sympathy depending on identity of structure (serous, synovial, mucous surfaces.) Second, by a mode of the universal sympathy which prevails throughout the system, independent of the alliances of structure and organization, which disposes parts different in properties to be affected reciprocally by the same form of diseased action (joints of the hand and foot, stomach and retina, in gout; muscular fibre and ligamentous capsules in rheumatism; skin, mucous membrane, and iris, in syphilis). To the numerous, extensive, and complicated sympathies of the latter class belong many cases of metastasis, in which the secondary often differ from the primary forms of diseased action. Inflammation in one organ occasions in another congestion, and *vice versa*; and the cessation of an habitual secretion in one, whether natural or morbid, gives occasion to inflammation in another. Preternatural irritability, swelling, pain, spasm, are thus excited by this reciprocal sympathy, in remote organs; or a metastasis, a change of place, strictly speaking, occurs; the morbid action abruptly quitting one part before it appears in the other.

\* See "A Practical Treatise on the Diseases of the Eye," by John Vetch, M.D. F.R.S.E. London, 1821.

But not only does the reflected action often differ in its nature from the original, but this, in many instances, continues in undiminished activity, so that the metastasis is only an incidental feature of the association. Now, if the sclerotica and the ligamentous capsule are liable to be reciprocally affected, we may refer it to the first stated sympathy of kindred textures, as in the case of hernia humoralis and gonorrhea, whether simply propagated or metastatic; and of rheumatic or scrofulous inflammation passing from joint to joint. But if the above mentioned parts are, as I believe, subject to be consensually affected during gonorrheal inflammation,\* I should explain it by reference to the second and more diffused sympathy which connects all organs and textures through the medium of the common sensory, with various degrees of affinity, according to their vital powers and properties.

---

#### NOTE B.

IN the history of the idiopathic iritis, by Dr. Schmidt of Vienna, a yellowish red tubercle is described as forming upon the surface of the membrane, which enlarges, and at length bursts and discharges its contents into the chamber. This he denominates an abscess, of which the cyst remains visible for some time afterwards. But for the general accuracy of this author's descriptions, I should decide against the correctness of this observation. I have never been able to discover any thing resembling an abscess in the iris, though perfectly familiar with the appearance of a tubercle like that described; and it is highly improbable that lymph should be effused in tubercles, which in some parts become rapidly organized and absorbed, and in other parts form cysts of abscesses, upon the same texture, at one and the same time. Such abscesses

\* I speak here of true sclerotic inflammation independent of any mixture of puriform conjunctival ophthalmia.



would degenerate into ulcers, an appearance never witnessed. Pus, as I believe, is never formed in pure iritis.

A species of iritis is described by the same author, which he names the arthritic, an inflammation either primary or secondary in a gouty habit to a common ophthalmia. It is attended with excruciating pain. Its diagnostic signs are the appearance of a narrow white ring at the verge of the cornea, and a varicose disposition of the vessels of the conjunctiva. In spare and irritable individuals the pupil becomes contracted as in the idiopathic iritis, the blood-vessels of the iris are varicose, and the disease terminates in a diminution of volume of the eyeball. In persons of full habit and relaxed fibre, the pupil, on the contrary, becomes remarkably contracted, but not uniformly, being transversely oval. No lymph is effused, but a greenish yellow appearance of the humors is observed, and the lens bulges forward, of a sea-green color. This pain, which is periodical, and announced by a burning sensation around the organ, and a profuse flow of tears, is of the most severe description. The vessels of the choroid assume the same varicose state as those of the conjunctiva, and the transparent sclerotica presents a dark ring in the situation of the corpus ciliare. Total blindness accompanies this state, and atrophy of the globe ensues.

In this country we have not been accustomed to distinguish gouty inflammations in this organ. If an inflammation, characterised as above, is peculiar to the arthritic diathesis, the distinction is borne out; if not, the division is frivolous. I am unable to decide upon the value of the diagnostic sign first mentioned; not that it has altogether escaped my observation, as being in some instances of deep-seated inflammation more strongly marked than in others; but that no evidence has been conclusive to my mind of the existence of a distinct species of iritis affecting gouty subjects. A varicose state of the vessels is the ordinary result of continued inflammation, whatever be the texture affected. The same may be said of the ultimate state of atrophy of the entire organ from interstitial absorption. But to speak my opinion candidly, the two

states which are here described as varieties of iritis, according to the different habits of body in which it appears, are essentially different forms of disease. The fully dilated and transversely oval pupil, the collapsed and disorganized iris, the varicose vessels of the choroid and conjunctiva, the attenuation of the sclerotica and bulging of the ciliary ring, all indicate not an existing inflammation, but a disease gone by; they are slow after-changes which ensue upon acute destructive inflammation of the choroid and retina; and the agonizing attacks of pain which I have often witnessed in this precise condition, are the result of an universal congestion of the vessels and consequent distension of the eyeball, for every loss of blood gives temporary relief, and when the loss of volume is observed to commence, these attacks cease and return no more. That the angularities of the pupil are in one form of the iritis (syphilitic) observed to be upward and downward, and in another (arthritic) from side to side; that the sclerotic vessels, in one case, advance boldly to the cornea; and, in another, are separated by a line which is only visible through a magnifier, are circumstances too much of a contingent and casual nature to be admitted, as affording a ground for invariable and specific distinctions; and to look at them as types of corresponding constitutional states, is almost ludicrous.

---

#### NOTE C.

THIS head and that of amblyopsia, page 139, include a very considerable proportion of the cases of functional amaurosis. The amblyopsia may be regarded as the advanced stage of the symptomatic amaurosis, and borders upon paralysis. The disorder of the conjunctival surface, in many instances of pure weakness of sight, unattended by any degree of dimness, is so slight, as either to induce a belief that it has no share in the disease, or that it stands in the relation of an indirect effect, rather than of an exciting cause of the complaint. We must therefore consider it in such cases as a primary nervous



affection ; or as originating from sympathy with some other organ, or some peculiar state of the system. The existence of such causes, in many instances, is too obvious to escape notice : in others, I have looked for it in vain. Neither has the organ been over-exerted nor oppressed ; nor have the functions of the stomach, liver, uterus, &c. varied from a state of health. Excepting the absolute suspension of all such employments as demand the exercise of what may be called active vision, I know of no remedy for the disease. It is more frequent in early and middle, than in advanced life ; and in females than in males. There is no unusual intolerance of light, nothing amounting to actual pain, no defect of vision ; in short, the sum and substance of the complaint is the sensation of an effort to see, and the want of power to continue it. The removal of the exciting cause, the first and most important step in the treatment of all diseases, although so negative as almost to amount, in the patient's estimation, to an abandonment of the case, is in this more efficient than active measures, as abstinence may on some occasions be advantageously substituted for cathartics. Accordingly I have known the complaint materially mitigated by perseverance in this system for six or twelve months, with a scrupulous attention to regimen ; and perhaps by extension of the principle, a state of protracted sleep, if it were possible, might be the readiest mode of cure.

A young gentleman who consulted me three years ago for *muscæ*, a weakness of sight, and inability to exercise his sight in any way requiring continued attention, a painful sensation from looking at pointed objects, as pins, needles, or the corner of a chimney-piece,\* I have lately seen. There is not now, nor ever has been, the slightest deviation from a healthy

\* This is the only example I have met with of this very expressive symptom of a tender, or highly irritable retina. It seems to me to resemble those disagreeable but natural sensations, viz. the teeth on edge, or the *culis anserina*, when amounting, as in some individuals, to a morbid excess, and excited by impressions of which they alone are susceptible.

appearance of the eyes; the pupils contract freely, and there is no intolerance of light, nor does his uneasiness, though it compels him to desist from employment, ever amount to pain. He has a frame rather spare than plethoric, a healthy complexion, good appetite, animated disposition, and spirits corresponding to his health, if some apprehension about his sight did not occasionally depress him. His education has been from this cause interrupted, and he is unable to indulge his inclination in the choice of a profession.

Before I saw him he had been subjected to different plans of treatment. First, undue determination of blood without any sufficient reason was presumed; for this hypothesis he suffered leeching, cupping, and blistering. Secondly, it was regarded as a nervous affection; evacuations of blood were countermanded, and a particular attention to the bowels, sea-bathing, exercise, and various tonics substituted. I gave my opinion in writing three years ago, that the disorder was purely functional, that the systematic regulation of the bowels, attention to diet and exercise, with as much indulgence of the organ as was possible, comprised all the means of treatment which my experience suggested. That depletion on the one hand and the higher tonics on the other, would not amend, if they did not aggravate the complaint. At the same time, I recommended a trial to be given to the blue pill, the bitter infusion, and blisters. The disease is as nearly as possible the same at this moment, as when I was first consulted. His own observation is, that his sight is evidently more affected by the state of his stomach than any other circumstance, that the more observant he has been of the plans laid down for the regulation of diet, &c. the more tranquil and comfortable has been the state of his eyes. Reading a few pages of a book is infinitely more distressing than the glare of a theatre or a ball-room; and the sense of debility is greater during the morning than at any other time. But even though the bowels are perfectly regular, and the digestion strong, the same feeling prevails; and on the other hand, when these functions



have been very irregular, it has often happened, that the state of vision has been as good as at the best.

---

#### NOTE D.

PROFESSOR BEER refers to many cases of this description, as from suppression of febrile diseases of the skin, scarlatina, variola, &c. in the first period of the eruptions. The pupil in these cases is much contracted and immoveable; the prognosis favourable, if treated early. He has also seen the disease after suppressed chronic diseases of the skin, as psora; after the amputation of the plica polonica, and very many in consequence of an abrupt healing of old leg ulcers. The prognosis in these cases is unfavourable. In that arising from suppressed catarrhus narium, in which the pupil is angular and drawn toward the outer canthus, he considers the prognosis encouraging. He mentions as rare and unfavourable forms of the metastatic amaurosis, those arising from suppressed secretion of milk in lying-in women, and from suppressed passions of the mind.

---

#### NOTE E.

AN elderly gentleman, the subject of confirmed organic amaurosis, whom I had seen at intervals, died lately, and his friends kindly afforded me an opportunity of examining his head after death. In the year 1816 he first complained of dimness—was unable to mend his pen as usual, changed his spectacles repeatedly—and from writing a small neat hand wrote large and straggling. In walking he imagined that he saw objects on the ground which intercepted his path, and endeavoured to avoid them by taking long and high steps. The flames of the candle at night appeared multiplied and undefined. There was little if any difference between the two eyes. About the time that his sight grew dim, he complained of uneasiness and oppression in the head. He

often described the horrible sensation of passing under an archway, with the fear of being crushed by its falling.\* His habit was plethoric, and he was apparently in the full vigor of health. As these symptoms, together with much lethargy, were considered to threaten apoplexy, he was freely and repeatedly bled, and in other respects treated accordingly. He became totally blind and soon afterwards dark. His eyes were perfect in appearance. They had the unmeaning roll characteristic of the disease in its last stage. Pupils a good deal dilated and motionless. A full course of mercury and electricity were employed without any effect. The latter was persisted in for six months.

In the progress of the case he was attacked with fits of a mixed kind, partly apoplectic, with temporary hemiplegia, and in part epileptic; his mind and speech failed him. Great torpor of bowels and indigestion, scantiness of urine and pain in voiding it, coldness and œdematous swelling of the lower extremities, with frequent and severe convulsions of his whole frame, were the symptoms most remarkable towards the close of his life.

On inspection of the head, the ventricles of the brain were found greatly surcharged with serous fluid, and the optic nerves to and from the ganglion opticum shrunk, or rather absorbed; so that they appeared flat instead of cylindrical, and of a straw color instead of a silvery whiteness. In slitting, and cutting them across, it was evident that only the sheath of the nerve remained, the medullary substance had entirely disappeared. The eyes nevertheless were in all respects sound, and had the plumpness and clearness of health. There was no vestige of an apoplectic effusion.

The following case exemplifies the exclusive paralysis of the *nervi motores* referred to in the paragraph to which this note belongs.

\* Another morbid horror, somewhat resembling this, I have heard described, viz. the approximation of the walls of the apartment, so as to give the patient the impression of being in a closet just large enough to contain his person. I need scarcely observe that all such delusions have their origin in the sensorium.



Mrs. W. a healthy woman, aged 28, suckling an infant of five months, was attacked, in June, 1820, with severe pain in the head on first rising in the morning, which in an hour or two subsided, but after some time it continued during the whole day, affecting chiefly the left side. She, of her own accord applied leeches to her temples and a blister behind each ear, but without relief. In August following finding the pain almost insupportable, her family surgeon was consulted. The bowels, he informed me, were so obstinately costive as to be with difficulty acted upon by powerful cathartics. Being still unrelieved after brisk and effective purgative medicine, she concluded it to be rheumatism, and wrapped her head in flannel. In crossing the road on the 8th of November following, she felt a sudden smart shock between the orbital processes, like a pea striking her forehead. From this moment the pain in the head ceased; but she found the vision imperfect—that is, she saw objects in unnatural positions, and although she could see distinctly with either eye, she could not with both, and therefore tied up one, the left, which was rather the weakest when engaged in business. Both eyes appeared perfectly healthy, and the pupils were equally active. On the third of December following, she first saw objects double, and the strabismus was so marked as very sensibly to disfigure a pretty and pleasing countenance. Both eyes were turned towards the nose, the left most so. Both pupils nevertheless acted freely. She had suffered no return of pain since the 18th ultimo. She was now cupped to twenty ounces from the nape of the neck and temples. Her bowels had been kept in action by the pil. hydrarg. and an occasional purgative draught. Since the double vision began, these were directed to be continued. A large blister was applied to the occiput after the bleeding, and three days afterwards no improvement appearing, each temple was bled with six leeches, and these parts were also blistered. The dose of blue pill was gradually raised to ten grains twice a day, and in a fortnight the mercurial action was established.

Previous to this event, it was noticed that her vision had

been less confused. She was cupped twice in this period to six ounces, and blisters applied behind the ears were kept open for some days. Her mouth continued sore about a month, during which time she gradually recovered single and perfect vision, and the strabismus was corrected. During her recovery she saw but one object on looking steadfastly in a straight direction; but upon turning her eyes to either side, or upwards, she still saw objects double; and even now that her vision in all directions is ordinarily single, whilst in a supine posture, and especially whenever her mind recurs to the subject, her vision is occasionally double.

During her treatment, she was seldom free from a sense of heaviness about the forehead and occiput, and vertigo. The pulse was quick and feeble, countenance pale and haggard, mind irritable and anxious; and she labored evidently under great debility, both muscular and nervous. The recovery of the eyes was very gradual; no sensible acceleration of the rate of progress was observed during the mercurial influence. She did not however lose what had been gained in sight, though she lost strength. Even since the recovery of the sight, and in a great measure of her flesh and strength by country air, the continuance of weight and uneasy sensations in the head, led me to recommend a seton in the nape of the neck, which she adopted with advantage.

This young woman, I should observe, the mother of several children, acted as her husband's book-keeper, and whilst pursuing this sedentary and anxious occupation, had been accustomed to drink freely of potent home-brewed beer.

This case is contra-distinguished to that in which the affection of the retina precedes the strabismus. Here, it is the symptom—there, the cause of the strabismus. The sympathetic affection of the retina in the case just related, is the slightest possible. In the majority of such cases, it is more marked, so that the vision of one eye is much stronger and clearer than that of the other, and *one* eye may be said to



be in fault. When strabismus ensues upon amaurosis, this difference is still more conspicuous. But the prognosis is not least serious when the retina is least affected; a squint from blindness adds little to the case but confirmation that the retina is insensible; but strabismus coming first connects the origin of the disease with the cerebrum, and what alarm we feel in one case for the sense of vision, we feel in the other for life, or, what is of yet more value, for intellect.

I must beg my reader's excuse for still farther lengthening this note. A few days ago I was desired to visit a gentleman between thirty and forty years of age, who had just arrived from the West Indies, on account of a large and hard tumor seated in the abdomen, about the nature of which his medical attendants were in doubt, from its external character, its apparent insulation, and frequent change of place, being sometimes in the epigastrium or beneath the umbilicus, and at other times distinctly felt in the right iliac region, in the position of the caput coli. Sometimes it was concealed and could not be felt any where. The disease was of seven months' standing, attended with marked symptoms of stricture in some part of the intestinal canal: a very imperfect and disturbed state of the alimentary functions, frequent hiccough, sharp pains in the belly, and great emaciation.

On the day of my seeing him, he was suddenly seized with a new symptom; viz. convulsions and total blindness; and in the interval of the fits, which were protracted and severe, he complained of pain across the top and front of the head. His pulse was regular, but compressed; his skin was covered with a cold perspiration; except during the convulsive paroxysm, his mind was perfect, but his manner was changed from anxiety to apathy. The pupils were dilated to the utmost, as in hydrocephalus; he had no sense of light, no perception of its interruption. I directed his head to be immediately shaved, and twelve ounces of blood to be taken by cupping from the temples, and above each mastoid process. Afterwards a blister to be applied in the direction of the coronal and sagittal sutures, sinapisms to his feet, and a pill to

be taken, composed of five grains of calomel and half a grain of opium. Two hours afterwards, the convulsions recurring, and his skin being still cold and clammy, he was ordered a cordial draught and a small quantity of brandy in gruel occasionally. At the same time a turpentine glyster was administered; this was soon followed by a very plentiful discharge from the bowels, a warm and copious perspiration, and a sound sleep of several hours. The next morning he awoke free from pain, and his vision was as perfect as ever. He had no return of convulsion or cerebral disorder of any kind. No material change occurred in the symptoms of the original malady during this short but truly alarming attack.

This amaurosis, there can be no doubt, was an example of sympathetic irritation and congestion. The cupping was a precautionary measure indicated by the amaurosis; there was pain, a firm pulse, and this unequivocal symptom of compression; effusion was instantly threatened; a stimulus at this moment appeared hazardous. We expected to see him expire in each fit. The pill, followed by the stimulant glyster, was remedial, and for the time saved him.

P. S. Since the above was written, this gentleman is deceased. On inspection, a firm, very irregular, fungoid tumor, of the size of a man's double fist, knobulated, fissured, and of a truly malignant aspect, was found occupying the head of the colon; and, by its origin from an extensive surface of the mucous membrane, partially inverting and concealing the termination of the ileon and the cæcum, of which only the appendix vermiformis was seen. He had no other organic disease.

---

#### NOTE F.

THE different forms of amblyopsia amaurotica enumerated by Professor Beer are as follow:—  
Visus interruptus—the person in reading sees only single words or letters.



- Visus dimidiatus—S. hemiopsia.  
 Visus muscorum—S. myodesopsia.  
 Visus reticulatus—a higher degree of the former.  
 Visus lucidus—Photopsia. Marmaryge. (*Hipp.*) Sparks and flashes of lightning perceived by the patient.  
 Photophobia—light painful.  
 Oxiopsia—a state of vision which enables the patient to see with perfect accuracy the smallest point when deprived of light.  
 Visus nebulosus.  
 Visus duplicatus—S. Diplopia, Luscitas, et Strabismus.  
 Visus coloratus—S. Crupsia, all objects seem colored, green, blue, yellow, &c.  
 Visus defiguratus—S. Metamorphosia.  
 Myopia and Presbyopia.  
 Nyctalopia and Emeralopia.  
 Amblyopia vaga—periodica—intermittens.

---

#### NOTE G.

PROFESSOR WALTHER, in an essay on this subject, has stated some very original and curious notions. He thinks that cataract is the primitive and natural state of the lens, and that congenital cataract is therefore not an altered but an unaltered condition, in consequence of a check given to the development of the embryo. Like other malformations, it is not owing to the influence of any active or formative cause, but having been originally present in every embryo at certain periods of its existence, does not disappear in its progress to a more perfect state, as it does where this progress is unchecked. The three months' fœtus has a hare lip, with but one cavity for the mouth and nostrils. The iris is imperforate, and so are all the apertures of the perfect body. The eyelids are fastened together over the naked eye, and the cavity of the um-

bilical cord being one with the abdomen, exomphalos is the natural and original state.

Walther considers cataract to be always the result of inflammation of the capsule, acute or chronic. By a powerful magnifier he has discovered a wreath of vessels about a quarter of a line distant from the pupillary edge of the iris, forming a concentric circle with the pupil. To this vascular wreath vessels pass in radii from the circumference of the capsule, and into the posterior surface of the iris. Nay, a net-work of more delicate vessels is described to have been seen deeper seated in the lens itself, "the larger trunks of which are not always derived from the circumference of that body, but evidently come from its posterior surface directly forwards, and then divide into branches."\* This is an appearance entirely morbid, the same authority deciding that there is no organized connexion between the lens and its capsule in health, and that the lens is nourished by imbibition or absorption of the humor Morgagnii, secreted by the vessels of the capsule, into which it again deposits its waste, being merely furnished with absorbent and exhalent vessels. Hence inflammation of the lens is always secondary to that of the capsule, in the same manner as inflammation of the capsule is secondary to that of the iris. Spots seen in the capsule, whether gray or brown, are, we are told, deposits of lymph, in which the prolonged vessels are seen terminating.

The inflammation of the capsule and of the lens are described as diseases marked by certain signs and appearances. The latter is always chronic like that of the bones, cartilages, and fibrous textures. When the disease is established, the blood vessels of the lens and capsule become varicose. The firm cataract is the termination of inflammation in induration. The milky exemplifies suppuration. The dry siliquous or shrunk cataract is a dry gangrene. The hard cataract when occurring without inflammation is a scirrhus, and the purulent

\* I quote the words of the Analysis in the Quarterly Journal of Foreign Medicine and Surgery.



may sometimes be the effect of ulceration of the lens. Other cataracts are considered to be sarcomatous!

The first part of these observations, namely, that referring to the appearances exhibited in inflammation of the capsule, from its consistency and analogy with the phenomena that are open to observation, has been anticipated in the way of hypothesis, and may be admitted with proper allowance for the chances of optical delusion in the employment of a sextuple magnifier. But unfortunately the enthusiastic devotion to system which the author betrays in his pathological notions (which are to my seeming pure nonsense), gives an air of marvel to the whole story.

Professor Beer divides cataracts into true and spurious. The true is within the capsule; the spurious is placed between it and the iris. The principal kinds are,

*True.* 1. Lenticular. 2. Anterior capsular. 3. Posterior capsular. 4. Morgagnian. 5. Capsulo-lenticular. 6. Cystic. 7. Siliquous. 8. Cataract with a cyst or sac containing pus. 9. Trabecular.

*Spurious.* 10. Lymphatic. 11. Purulent. 12. Sanguineous. 13. Pigmentous.

The distinction and dignity of a name given to each variety, is the only novelty of this list. The cystic is the floating cataract, the capsule opaque and thickened, and the lens more or less absorbed. The siliquous is the capsular cataract, the lens being absorbed, as after wound or rupture of the capsule.\* The cyst containing pus is rare. The trabecular is probably the radiated. As to the four last, they are results of iritic inflammation, not cataracts; nor are they in this country confounded with them. They are seldom if ever met with but after blows, wounds, and operations.

\* Do the travelled pedants, who deal such heavy blows among their ignorant and besotted countrymen, suppose that these every day forms have escaped our notice? In England, as in Germany, the same things are seen, but their importance is differently estimated; they are differently explained, arranged, and reasoned upon. I am content it should be so.

## NOTE H.

THE often mooted questions—first, in what cases topical blood-letting commands an advantage, if in any, over phlebotomy, as usually practised at the arm ;—second, whether drawing blood from an artery is of greater efficacy than from a vein, the quantity being the same—may be answered, I think; in a few words. To the first I should reply—if the system is inflamed, i. e. if the pulse indicates that the action of the heart is excited by the state of the organ, or if the activity or rapid progress of the inflammation, however local, threatens the safety of the organ, we ought not to trust to topical blood-letting. In such cases it may be employed subserviently with great advantage, but not principally. If, on the other hand, no such sympathy is evinced, and the inflammation, though acute, is in its nature weak and slow of progress, the local blood-letting may commonly suffice. But is it in such cases to be preferred? Generally I think it is, because the inflamed vessels are relieved from the state of congestion and tension, and are enabled to recover their contractile tone, at a smaller expense to the system. To pale the inflamed conjunctiva by opening a vein in the arm, supposing there is no disposition to syncope, will require a loss of from twelve to sixteen ounces of blood ; whereas this effect will often be produced by a loss of one third of that quantity drawn as quickly from the immediate neighbourhood of the inflamed organ. Syncope from dread of the lancet will produce the best effect of a topical bleeding, and may stand in its stead. It will be no substitute for general bleeding where that is indicated, for the same reason that topical bleeding is no adequate substitute for general. Thus, though a general bleeding will answer the main purpose of a local one, it is least economical when neither the character of the inflammation, nor the danger of its duration, calls for it ; and, therefore, in strumous and other weakly habits, in atonic and sluggish inflammations, it is least eligible.



Nevertheless, when the inflammation and the part affected are such as to require general blood-letting, and the patient owes his safety to its employment, topical is continually had recourse to with remarkable advantage, and the latter is more eligible as an auxiliary in such cases, than the former as a substitute in those of an opposite description. The difference of the inflammatory action according to the texture of the organ affected, is comprehended in the above general statement.

Leeches are the least effectual mode of topical blood-letting. In many instances, the blood derived by them is not sufficient to counterbalance the irritation caused by the wound, and they act as irritants, augmenting the vascularity, swelling, and pain.

To the second question my observation leads me to reply in the negative. The effect upon the heart's action will be determined by the quantity lost, and by that alone—taking the distance from the heart and the size of the current—in any artery which it is usual or would be discreet to bleed from.

---

#### NOTE I.

THE gratitude of the public, the highest, if not the only reward for public benefit to which an honorable mind aspires, is unquestionably due to Dr. Vetch, for the successful treatment of this formidable disease, especially in its first and most formidable stage. He reprobates the excision of the granulations and the division of the conjunctiva. Although I have seen cases of the absorption of large and even pendulous granulations, and believe that such absorption almost invariably takes place where the disease is left to run its course, abundant experience has convinced me that the employment of the scissars is highly important to the favorable issue of the case, since the preservation of the cornea depends chiefly upon the restoration of the lining membrane of the lids, which this practice essentially promotes. Yet so highly do I

appreciate the importance of caustic and astringent applications judiciously employed, that if the joint use of these remedies were in any case proscribed, I should prefer, as a single measure, the use of the lunar caustic or the blue vitriol, to the employment of the knife.

Dr. Vetch objects to the use of these substances in solution as too stimulant, and contends that when lightly applied in the solid form they act most beneficially. There is much truth in this observation. The fact is, that in this, as in all chronic morbid changes, the treatment rests not upon one, but on a variety, or perhaps a combination of measures; not upon this or that form or mode of application, but upon a form and mode suited to the existing circumstances. A topical application made with advantage to-day would probably be hurtful to-morrow. To watch the caprices of the case, the moment of excitement and of relaxation, to subdue, to soothe, or to support promptly, and thus to break the force of each successive relapse, varying the means as the circumstances permit, are points indispensable to conduct the case to a favorable termination, and are those which distinguish the man of science from the empiric. But the local treatment is by no means the only material part of the 'Therapeia' of these cases.



## NOTE K.

WHERE these circumstances are not present, the subjects of this inflammation are, in my experience, weakly and of a scrofulous habit ; the tame, indolent, shifting character of the disease, viz. the pink-colored zone at the verge of the cornea, vivid one day, and scarcely perceptible the next, renders it more difficult to subdue than a fixed and vigorous action ; and the practitioner is insensibly betrayed into irresolution by the seeming inertness of the disease and the obvious delicacy of the patient. But diseases, like other dangers, are formidable in proportion as they are disguised, and it would often prevent mischief, if a bolder practice were instituted in the outset than circumstances appear to the inexperienced to warrant.

---

## NOTE L.

THIS statement requires some qualification. What I mean is this ; where the mercurial action being fairly established is productive of no sensible improvement, its continued and freer use is attended with no advantage, and is, therefore, constitutionally injurious ; but I do not mean to say that the full advantage will always be obtained by a short and gentle course ; on the contrary, where in the commencement of the mercurial action the improvement is only sufficient to give encouragement to persevere, a course of three or even four months is often necessary to accomplish the end in view.

In amaurosis supervening on inflammation, especially recent inflammation, the remedy promises most ; but even here, if the inflammation has induced perfect amaurosis, it will often restore the iris to its color and activity without materially benefiting vision ; for mercury is not a remedy for

paralysis. If however from inflammation—whether the result of injury or spontaneous, whether from the operation of a direct or a remote cause, the state of congestion or atony, the state of serous or lymphatic, perhaps of partial sanguineous effusion—the sensorial function be interrupted, our first hope is topical blood-letting and counter irritation—the second, the action of mercury.

At the distance of four years from a fall on the occiput, followed by severe symptoms of cerebral injury, I have known sight restored to the eye which had ever since been deprived of it, by a full mercurial course. Again and again I have seen the same effect produced by the same agent, where neither injury nor any other cause nor symptom of inflammation had ever existed, and where only a slow moving pupil corresponded to the patient's complaint of dimness, to such an extent as to render indistinguishable the features of a person standing before him: and on the other hand I have never known it to be of any efficacy in cases, ushered in by severe frontal pains, in which blindness was already complete, and the pupil largely and permanently dilated, where the greenish cast of the humors was strongly marked, and the visus lucidus was complained of.

---

#### NOTE M.

I CONFESS that I have seen no reason to alter the opinion here expressed, in several trials of this plan of treatment with the tube of M. Dupuytren, since the publication of this work. Through a learned member of the University of Paris, a patient of mine, and also of M. Dupuytren, I requested the professor to favor me with a case that might serve as a report in detail of the treatment which he had adopted. Upon this gentleman the tube operation, though performed by the Baron, was unsuccessful; but as he had previously worn for two or three years a nail-headed style with as little



advantage, the case certainly did not afford a fair chance of success, and the failure was attributed to this circumstance. M. Dupuytren, with the liberal and courteous spirit of a true friend to science, immediately transmitted to me the subjoined report of an interesting case which had very recently occurred ; and I have great satisfaction in presenting it to my readers.

“ Madame Daive, âgée de quarante deux ans, demeurant à Sarre-Louis, vint à Paris dans le mois de Mai 1821, pour consulter Mons. le Professeur Dupuytren pour une tumeur lacrymale d'un côté, et une fistule de l'autre côté.

“ Il y a six ans que la malade s'aperçut pour la première fois, que l'œil du côté gauche étoit larmoyant, qu'il se formoit souvent à son grand angle une petite tumeur qui se vidoit par la pression, et qui ne tarδοit pas à se reproduire. Cette dame avoit un écoulement involontaire de larmes sur la joue ; l'œil de ce côté étoit toujours chassieux, larmoyant, la narine toujours sèche. Cette petite tumeur augmenta bientôt de volume, la peau qui la recouvroit, s'enflamma, s'aminçait ; les paupières se tuméfièrent. Ces accidens la firent recourir aux soins d'un chirurgien, qui ouvrit de suite cette tumeur : du pus, des larmes, du mucus, et du sang s'écoulèrent ; la malade fût soulagée, l'inflammation tomba ; mais à la tumeur succéda une fistule qu'il falloit guérir, et voici les moyens qui furent employés.

“ On fit d'abord des injections. Quoique continuées pendant un temps assez long, elles n'eurent aucun résultat heureux. Ce premier moyen ayant échoué, on en essaya un second ; c'est-à-dire, qu'on tenta de faire passer dans les fosses nasales un ressort de montre. On fit des essais pendant plusieurs jours, leur inutilité fit adopter le moyen suivant, dans l'intention de désobstruer par cautérisation les voies lacrymales. Un stilet rougi au feu fut introduit entre les lèvres de la fistule, et les cautérisa : du gonflement survint ; au bout de quelques jours il diminua ; bientôt les escarres formées se détachèrent ; des bourgeons charnus s'élevèrent ; mais le chirurgien s'appliqua chaque jour à introduire, pendant quelques instans, un cylindre de nitrate d'argent poudré.

Cette manière d'agir eut pour résultat l'aggrandissement de la fistule, l'adhérence de ses bords aux os ; les bords prirent une organisation cutanée : enfin on introduisit pendant longtemps une espèce de broche en plomb—Tous ces moyens loin de guérir la maladie, l'avoient rendue presque incurable. Désirant trouver un remède à son infirmité, Madame Daive se présenta chez M. le Baron Dupuytren ; elle se trouvoit alors dans l'état suivant.

“ Au grand angle de l'œil gauche existoit une ouverture de trois lignes de diamètre ; les bords tapissés par la peau amincie, avoient pris l'organisation cutanée dont s'emparent toujours les ouvertures fistuleuses, qui donnent passage à des corps étrangers. Par là s'écouloient sans cesse des larmes qui venoient irriter l'œil, enflammer et excorier la peau de la joue.

“ Au grand angle de l'autre œil existoit depuis quatre ans une petite tumeur, plus grosse pendant le temps humide, plus aussi le matin que le soir. Cette tumeur pouvoit être facilement vidée par la pression ; alors il s'écouloit par les points lacrymaux une matière purulente, muqueuse, mêlée à l'humour des larmes ; la narine de l'un et l'autre côté étoit sèche.

“ La guérison de la tumeur lacrymale étoit certaine, celle de la fistule pouvoit être douteuse ; la malade en fut prévenue, et l'opération pratiquée le 4 Mai 1821, de la manière suivante.

“ La malade étoit assise sur un chaise placée vis-a-vis d'une fenêtre, la tête appuyée sur la poitrine d'une aide : Monsieur Dupuytren tend alors avec le *medium* et le doigt indicateur de la main gauche, la peau des paupières de l'œil droit, en la portant un peu en dehors, tandis qu'avec la main droite armée d'un bistouri à lame étroite, il fait à la peau qui recouvre la tumeur, une incision perpendiculaire, qui la divise, ainsi que le sac lacrymal. On vit bien que l'instrument n'avoit pas dévié, par la profondeur à laquelle il pénétra sans difficulté, et à la sortie de mucosité purulente. Changeant alors de main, M. Dupuytren saisit avec la droite le bistouri, et avec le gauche le mandrin, revêtu de sa canule en or.\* Le bistouri est un peu retiré pour permettre à

\* Voyez la description à la fin de l'observation.



l'extrémité du mandrin qui est glissé sur sa lame, d'être introduite à mesure qu'on fait entrer le mandrin ; enfin lorsqu'on est entré à la hauteur du canal nasal, il ne reste plus qu'à l'enfoncer. On est averti qu'il a pénétré assez avant, par la résistance qu'on éprouve à l'enforcer davantage ; ce qui provient du contact de la canule sur le rebord de la gouttière lacrymale. Voulant s'assurer que la communication existoit entre le sac lacrymal et le fossé nasal, Monsieur Dupuytren ferma l'ouverture antérieure des fosses nasales, et ordonna à la malade de faire des efforts comme pour se moucher, aussitôt on vit de l'air mêlé à du pus et à des mucosités sanguinolentes s'échapper par la petite ouverture ; on y présenta la flamme d'une bougie, elle fut éteinte.

“ De l'autre côté l'ouverture fistuleuse permit l'introduction de la canule ; elle fut facile, et chose étonnante, mais qui arrive toujours, c'est que la malade ne sentant nullement la canule, avoit peine à croire qu'on l'eût introduite.

“ Restoit à savoir l'issue qu'auroient ces deux opérations. Au bout de vingt quatre heures la petit plaie du côté droit fut cicatrisée, la tumeur n'existoit plus, le cours des larmes étoit parfaitement rétabli, et la narine de ce côté avoit repris son humidité naturelle.

“ Plusieurs jours après l'opération, l'ouverture fistuleuse du côté gauche parut un peu rétrécie ; cependant la malade éprouvoit de ce côté la même incommodité.

“ Que pouvoit-on faire pour cicatriser cette ouverture ? Devoit-on détruire les adhérences de la peau, enlever les bords de la fistule ? Mais en agissant ainsi, on pouvoit craindre de ne pas réussir, et d'aggraver au contraire l'état de la malade ; aussi ce parti ne fut-il pas adopté.

“ Les succès brillans que Monsieur le Professeur Dupuytren venoit d'obtenir dans la guérison de fistules recto-vesicales, uretro-vaginales, par le cautère, lui suggérèrent l'idée d'employer ce moyen. En effet quinze jours après l'opération, l'ouverture fistuleuse n'ayant fait aucun progrès vers la cicatrisation, Monsieur Dupuytren la toucha avec un petit pinceau de charpie trempée dans du nitrate de mercure, avec excès d'acide nitrique ; par-dessus il mit encore de la charpie

hachée, également imprégnée de ce caustique. Du gonflement, de la douleur survinrent ; ils furent calmés par quelques lavemens, des pédiluves (bains de pieds) sinapisés, du petit lait, &c.

“ Au bout de quatre jours l'escarre tomba, et Monsieur Dupuytren vit avec plaisir que la plaie s'étoit un peu rétrécie. Enhardi par ce premier succès, il fit une seconde, troisième, quatrième, et jusqu'à une septième cautérisation ; toutes furent pratiquées à quatre ou cinq jours d'intervalle, chaque fois on trouva l'ouverture rétrécie. Enfin le 3 Juillet, deux mois depuis l'opération pratiquée, cette ouverture fistuleuse à parois cutanées, organisée depuis plusieurs années, et que plusieurs médecins avoient jugé incurable, étoit parfaitement cicatrisée, le cours des larmes rétabli ; en un mot, il étoit difficile de pouvoir assurer que cette malade avoit eu une tumeur lacrymale d'un côté, et une fistule de l'autre, tant elle étoit bien guérie. On ne pouvoit voir à l'œil que la malade avoit dans le nez deux canules en or, et leur présence se faisoit si peu sentir, que la malade avoit peine à croire qu'on les lui eut laissées. Enfin elle quitta Paris parfaitement guérie, et heureuse d'avoir été délivrée d'une infirmité qu'on avoit jugé incurable.”

---

*Description de la canule d'or de Monsieur le Professeur Dupuytren.*

“ Cette canule doit être aussi longue que le conduit nasal ; par conséquent elle est de 10 à 14 lignes, sur une, ou une et demie de diamètre, cependant sa partie supérieure est un peu plus large ; elle offre une légère courbure pour s'accommoder à celle du canal. Un rebord renflé, en forme de bourrelet saillant en dehors, fait le contour de l'ouverture, qui doit aboutir au sac lacrymal ; il est destiné à s'opposer à la chute de la canule dans le nez, chute qui permettroit à la maladie de se reproduire ; l'autre extrémité de la canule est taillée en bec de flûte, afin qu'un de ces côtés moins long, ne




by the absence of such a symptom during the palpable existence of such conditions ; the infrequency of this case, and the frequency of the cases proposed ; others are altogether hypothetical, and I will not add to the '*obscurum per obscurius*.'

## NOTE X.

Though their general complexion is light, the countenance is sometimes pale and sometimes flushed, and the eyes are sometimes watery and sometimes dry. The conjunctival surface is highly irritable, and a profuse gush of scalding tears overflows the cheeks, at the moment that the eye is opened and exposed to the light. The skin of the lower lid and cheek is excoriated, and the cornea becomes rough and scaly from the quantity and density of the lachrymal discharge. The patient's only relief consists in keeping the eyelids closed. The repeated pressure of the probe, if some slight obstruction should have existed, and the flowing of water injected by the puncta in a stream from the nostril, afford sufficient freedom of the canal. THE END. A preternatural sensibility, but no relief to the epiphora. A preternatural sensibility of the membrane may be inferred from the excessive pain and dread of the operation. I have seen such a case in a delicate young female of many months' duration ; in which neither soothing nor anæsthetic applications, neither anæsthetics nor alterative, tonic nor sedative medicine, produced the smallest benefit. The congestion of the superciliary vessels is inconsiderable : no morbid appearance is seen on examining the upper lid ; the orifices of the canaliculi lachrymales are natural ; there is no fulness nor unevenness felt on pressure in the region of the lachrymal gland. What is the disease ? Is it an affection of the lachrymal gland ? and if so, does it result from any undue excitement, or any morbid irritability, like a 'tic sensitivus' of the face ? NEW YORK : PRINTED BY J. SEYMOUR, JOHN STREET.


3281













NATIONAL LIBRARY OF MEDICINE



NLM 02093951 2