

**A study of the surface tension of blood serum by the drop weight method /
By Harold E. Woodward.**

Contributors

Woodward, Harold Edward, 1888-
Augustus Long Health Sciences Library

Publication/Creation

Worcester, Mass. : C.W. Burbank & Co., Printers, 1912.

Persistent URL

<https://wellcomecollection.org/works/gea82ygg>

License and attribution

This material has been provided by This material has been provided by the Augustus C. Long Health Sciences Library at Columbia University and Columbia University Libraries/Information Services, through the Medical Heritage Library. The original may be consulted at the the Augustus C. Long Health Sciences Library at Columbia University and Columbia University. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

COLUMBIA LIBRARIES OFFSITE
HEALTH SCIENCES STANDARD



HX64100898

QP99 .W87 1912 A study of the surfa

1912

Woodward.

A Study of the Surface Tension of Blood Serum
by the Drop Weight Method.


RECAP


Columbia University
in the City of New York

THE LIBRARIES



Medical Library





Digitized by the Internet Archive
in 2010 with funding from
Open Knowledge Commons (for the Medical Heritage Library project)

150
28
RECEIVED
A Study of the Surface Tension
of Blood Serum by the
Drop Weight Method

DISSERTATION

SUBMITTED IN PARTIAL FULFILMENT OF THE REQUIRE-
MENTS FOR THE DEGREE OF DOCTOR OF PHILOSOPHY
IN THE FACULTY OF PURE SCIENCE IN COLUMBIA
UNIVERSITY IN THE CITY OF NEW YORK

BY

HAROLD E. WOODWARD, B. A.


NEW YORK CITY

1912

WORCESTER, MASS.

CHARLES W. BURBANK & Co., Printers

1912



A Study of the Surface Tension
of Blood Serum by the
Drop Weight Method

DISSERTATION

SUBMITTED IN PARTIAL FULFILMENT OF THE REQUIRE-
MENTS FOR THE DEGREE OF DOCTOR OF PHILOSOPHY
IN THE FACULTY OF PURE SCIENCE IN COLUMBIA
UNIVERSITY IN THE CITY OF NEW YORK

BY

HAROLD E. WOODWARD, B. A.

NEW YORK CITY

1912

WORCESTER, MASS.

CHARLES W. BURBANK & Co., Printers

1912

QP 99

W 87

1912

ACKNOWLEDGMENT

This investigation was suggested by Professor J. L. R. Morgan, and carried out under his directions. I wish to express my appreciation of the assistance and encouragement I have received from him.

Professor W. J. Gies directed the experiments on dog serum, and I am very grateful to him for the interest he has taken in the work and the assistance he has given me.

I am also deeply indebted to Dr. Bailey, Dr. Hopkins and Dr. Smith, of St. Lukes Hospital, for valuable material aid and advice, and to Dr. Butterfield and Dr. Bronfenbrenner, of the Rockefeller Institute, and Dr. Warren, of Roosevelt Hospital.

HAROLD E. WOODWARD.

LABORATORY OF PHYSICAL CHEMISTRY,
HAVEMEYER HALL, COLUMBIA UNIVERSITY.

CONTENTS

Introduction and Object of the Investigation.....	5
Apparatus and Method.....	6
Dog Blood Serum.....	11
Human Serum.....	15
Comparison of Animal Sera.....	17
Meiostagmin Reaction.....	20
Summary.....	27
Vita.....	28

A Study of the Surface Tension of Blood Serum by the Drop Weight Method

Ascoli and Izar* have shown that the immunizing reaction in pathological blood serum is accompanied by a lowering of the surface tension of the serum. They used Traube's stalagmometer, working at room temperature, and observed the change in the number of drops in a given volume of the serum after the immunity reaction had taken place. It seemed advisable to confirm this change, which they called the Meistagmin reaction, by the more accurate drop weight apparatus of Morgan †. At the same time it was thought that an accurate study of the surface tension of normal and pathological blood serum might be valuable, and that there might be a normal value for the surface tension of the blood serum of all mammals.

In measuring surface tension by such methods as capillary rise or the drop method (stalagmometer), the density of the liquid must be used to change the height of the liquid in the capillary tube or the drop volume to surface tension. But since drop weights from any one tip are proportional to surface tensions, for any liquid and any temperature, it is only necessary to get the weight of the falling drop, thus avoiding the error and difficulty of determining densities. From drop weights then we have

$$\gamma = \text{weight} \times \text{constant}$$

This work, which is necessarily rather fragmentary, is divided into the following parts,—

*Juhnke, *Interstate Medical Journal*, 18, 233, (Feb. 1911)

†Morgan, *Jour. Amer. Chem. Soc.* 32, 349 (1911)

- I Surface tension of blood serum.
 - A— Dog serum; the condition of the dog being controlled.
 - B— Human serum, normal and pathological.
 - C— Comparison of different animal sera.
- II The Meistagmin reaction; the decrease in surface tension accompanying the union of antibody and antigen.

APPARATUS AND METHOD

The apparatus used is the same as described by Morgan.* For measuring the drop weight of serum the apparatus was allowed to remain in a constant temperature bath regulated at 37°C. (body temperature) for nearly half an hour, then a drop of the serum was pulled over and left hanging for five minutes in order that the air in the weighing vessel should be saturated with its vapor. This drop was then carefully forced back into the supply vessel, which contained about 5 cc. of serum, and a fresh drop was slowly pulled over and allowed to fall of its own weight. The weighing vessel was then weighed as usual after condensing the vapor in it by water at room temperature for one minute. Next a blank was run in the same way, except that the drop was not allowed to fall, and by subtracting this weight from the other the correct weight of one drop of the serum was obtained.

The determinations in the Meistagmin reaction were made at 0°C. as there is almost no evaporation at that temperature and in order that no reaction should go on in the solution during the determination. Five drops were taken and weighed, and no blanks were necessary.

As in the work on water and solutions, the tip had to be cleaned after each determination or the results would be

**Loc. cit.*

too low. The cleaning was best done by alkalin permanganate solution followed by chromic acid in diluted sulfuric acid, then the tip was washed with distilled water and dried by suction.

STANDARDIZATION OF TIP

One tip was standardized on benzene at 30°. The constant is found by means of a modified Ramsay and Shields

equation,
$$K = \frac{W \left(\frac{M}{d} \right)^{\frac{2}{3}}}{t_c - t - 6}$$

in which W = drop weight in milligrams, $M = 78$, $d = .86824$, and $t_c = 288.5$. Surface tensions may be found from the ratio

$$W : \gamma :: K : K' \quad \text{or} \quad \gamma = \frac{K'}{K} W$$

γ will be in dynes per centimeter, and K' is the surface tension constant from the Ramsay and Shields formula, $K' = 2.1148$.*

Tip I gave the following results with benzene at 30°.

Vessel + 30 drops	Vessel + 5 drops	
11.7112	10.9695	
11.7113	10.9694	25 drops = 0.74182 gram.
11.7114	10.9697	
11.7113	10.9695	1 drop = 0.0296728
11.7114	10.9693	
11.7112		= 29.67 mg.
11.71130	10.96948	

$$K = 2.3572$$

$$\gamma = \frac{2.1148}{2.3572} W = 0.8972 W$$

*Morgan and McAfee, *Jour. Amer. Chem. Soc.* 33, 1275. (1911)

$$\text{diameter of tip}^* = \frac{2.3572}{.4224} = 5.580 \text{ mm.}$$

Two other tips were standardized on distilled water at 37°. The surface tension of water, from

$$\gamma = 75.872 - 0.1547t - 0.000222t^2 \text{ is } 69.844 \text{ at } 37^\circ$$

Tip II gave results as follows,

Vessel + 30 drops	Vessel + 5 drops	
13.2202	11.3082	25 drops = 1.91208 grams.
13.2207	11.3085	
13.2199	11.3078	1 drop = 0.0764832
<u>13.2202</u>	<u>11.3082</u>	
13.22025	11.30817	= 76.483 mg.

$$\gamma = \frac{69.844}{76.483} W = 0.91319 W$$

$$K = \frac{2.1148}{.91319} = 2.3158$$

$$\text{diameter of tip} = \frac{2.3158}{.4224} = 5.483 \text{ mm.}$$

Results on Tip III were

Vessel + 30 drops	Vessel + 5 drops	
12.9084	11.0109	25 drops = 1.8976
12.9085	11.0108	
<u>12.9082</u>	<u>11.0106</u>	1 drop = 0.075904
12.90837	11.01077	= 75.904 mg.

$$\gamma = \frac{69.844}{75.904} W = 0.92016 W$$

*Morgan and Cann, *Jour. Amer. Chem. Soc.* 33, 1060 (1911)

$$K = \frac{2.1148}{.92016} = 2.2983$$

$$\text{diameter of tip} = \frac{2.2983}{.4224} = 5.441 \text{ mm.}$$

In order to test the tips with thick viscous liquids somewhat like serum, the drop weights of glycol and sugar solutions were first tried.

Glycol has practically no vapor tension at ordinary temperatures, (b. p. 200°) so 10 drops were weighed without any blank. At 0° glycol is so viscous that it took over thirty minutes to get the ten drops, while the same number of drops of water are generally dropt in less than five minutes.

Results on glycol (CH_2OH)₂

Temp.	Tip	10 drops	Average grams	1 drop mgs.	Crit. Temp.	γ
0	I	0.5265	0.52683	52.683	329.5	47.26
		0.5266				
		0.5273				
		0.5269				
		0.5270				
		0.5268				
30	I	0.5051	0.50504	50.504	350.0	45.31
		0.5053				
		0.5047				
		0.5051				
		0.5050				
55	III	0.4728	0.4731	47.310	365.9	43.53
		0.4733				
		0.4734				
		0.4729				

The values for surface tension give, by least squares, the formula, $\gamma = 47.277 - 0.0675t$. Walden's* formula, from capillary rise is

$$\gamma = 48.48 (1 - 0.00205t), \text{ or } \gamma = 48.48 - 0.0994t.$$

*Walden, *Zeits. Phys. Chem.* 65, 143.

The large difference at low temperatures is probably due to the fact that the liquid is so viscous at 0° that the rise in a capillary tube can not be accurately measured.

Comparison of results

	$\gamma = 47.277 - 0.0675t$	$\gamma = 48.48 - 0.0994t$
0	47.28	48.48
30	45.25	45.50
55	43.57	43.01

Sugar solutions at 37°

Tip	5 drops .5 molar	1 drop	W	γ
III	0.3819	0.0770	76.50	70.30
	0.3826	<u>0.0768</u>	<u>76.30</u>	
	<u>0.3829</u>	-.0006		
	0.3825		76.40	
1 molar				
II	0.3881	0.0780	87.63	70.86
	<u>0.3882</u>	<u>0.0780</u>	<u>77.50</u>	
	0.38815	-.0005	77.56	
1.5 molar				
III	0.3887	0.0785	77.81	71.67
	<u>0.3894</u>	<u>0.0784</u>	<u>77.95</u>	
	0.38905	-.0005	77.88	
2 molar				
II	0.3983	0.0804	79.81	72.90
	<u>0.3998</u>	<u>0.0803</u>	<u>79.85</u>	
	0.39905	-.0005	79.83	

The method of least squares applied to these results, with 69.84 as the surface tension of water, gives

$$\gamma = 69.878 + 0.456C + 0.5202C^2$$

for the surface tension of different concentrations of sugar at 37°. The calculated results from this agree very well with the observed results.

	obs.	calc.	diff.
Water	69.84	69.88	+0.04
0.5 molar sugar	70.30	70.24	-0.06
1 " "	70.86	70.85	-0.01
1.5 " "	71.67	71.73	+0.06
2 " "	72.90	72.87	-0.03
			<u>0.00</u>

DOG BLOOD SERUM

Except for the first three which were used for practice, the dogs were kept in special cages and fed the regular diet (15 gr. meat, 4 gr. cracker meal, 3 gr. lard and 35 cc. water per kilogram of body weight) until they were in normal condition. They were then subjected to various conditions which a person might undergo, such as missing meals, having extra food, loss of blood, action of cathartic, etc., and they were then bled in order to see if the surface tension of the blood serum would change under such conditions.

At first a few measurements made on defibrinated blood and centrifuged serum, to see if the surface tension was the same in either case. There seemed to be almost no difference, altho a French investigator reported a difference of over 2%. Results on one dog were, with Tip I.

Blood	W		γ	Serum	W		γ
5 drops				2 drops			
0.2487	49.66		44.6	0.0991			
0.2479				0.0992	49.56		44.5
				0.0993			
				0.0989			

As the serum was easier to use, and the blood decomposed rather easily, the serum was used alone.

A comparison was then tried between centrifuged and clotted serum, to see if the method of getting the serum made any difference Serum from one dog gave on Tip III.

Centrifuged			Clotted		
1 drop	W	γ	1 drop	W	γ
0.0502			0.0499		
<u>0.0502</u>	49.7	45.7	<u>0.0497</u>	49.3	45.4
-.0005			-.0005		

There is thus a little more than half of one percent difference, but as the clotted serum remained clear longer than the centrifuged, the blood was allowed to clot after No. 7.

EXPERIMENTS ON DOGS

No.	Tip	Dog	1 drop	W	γ	Remarks
1	I	1 Fem.	0.0991			
			0.0992	49.56	44.5	
			0.0993			
			0.0989			
2	I	1 "	0.2560			
			0.2559	51.19	45.9	
			0.2560			
			0.2558			
3	I	2 Male	0.2757			
			0.2765	55.2	49.5	Dog did not eat or urinate for several days.
			0.2760			
			0.2759			
4	II	3 "	0.0502	49.6	45.7	
			0.0499			
5	III	4 Fem.	0.0503			
			0.0500	49.6	45.7	
			0.0503			

6	III	5	Male	0.0494 0.0493	48.9	45.0	
7	III	5	"	0.0502 0.0502	49.7	45.7	Centrifuged
				0.0499 0.0497	49.3	45.4	Clotted
8	III	6	"	0.0500 0.0498	49.4	45.5	
9	III	6	"	0.0498 0.0498	49.3	45.4	Same dog, next day
10	II III	4	Fem.	0.0496 0.0490 0.0491	49.1 48.0	44.8 44.7	Extra meal
11	II III	4	"	0.0496 0.0492 0.0493 0.0491	49.1 48.7	44.8 44.8	Extra meal
12	III	5	Male	0.0499 0.0499	49.4	45.5	No food day before bleeding
13	III	6	"	0.0505 0.0503	49.9	45.9	No food for two days
14	III	6	"	0.0499 0.0497	49.3	54.4	Extra water
15	II III	7	"	0.0508 0.0504	50.3 49.9	45.9 45.9	Did not eat for several days
16	III	8	"	0.0500 0.0502 0.0502	49.7	45.7	10 gr. salt
17	III	9	Fem.	0.0493 ¹ 0.0493 ⁰ 0.04924	48.8	44.9	10 gr. sugar

18	III	9	"	0.04980			
				0.04984	49.3	45.4	Dose of Magnesium sulfate
				0.04968			
19	III	10	"	0.0502	49.6	45.6	
				0.0501			
				0.0501			
20	III	10	"	0.0504	49.8	45.8	Same dog next day
				0.0503			
				0.0503			

NOTE:—In No. 1 two drops were weighed, and in No. 2 and 3 five drops were weighed in a vessel containing some serum. In the others one drop was weighed in an empty vessel and the weight of the vapor, as a result of several determinations was taken as 0.5 mg.

The first eight experiments show that the normal surface tension of dog blood is about 45.5 dynes per centimeter. The first three vary from this more than the others because the dogs had not been kept in condition.

Experiment 9 was to see if the loss of blood had any effect on the surface tension. The dog was bled about as far as was safe each day (No. 8 and 9), and the surface tension was about the same each day. This experiment was repeated in No. 19 and 20, except that the two bleedings were about 19 hours apart instead of 26 hours. In this shorter time the surface tension had not quite reached its normal value, for the blood was still rather diluted.

In No. 10 and 11 the dog was given an extra meal several hours before the blood was taken. The decrease of about 1.5% shows the effect of the products of digestion, especially fat, on the surface tension of the blood.

In experiments 12 and 13 the dog starved one and two days, respectively, before the blood was taken. The result in No. 13 shows a rise of about 1% in surface tension as the blood becomes poorer in the products of digestion. The dog in No. 15 refused to eat for several days, and the surface tension in this case confirms the other result.

500 cc. of water given two hours before the bleeding in No. 14 seemed to have no effect on the surface tension, and a good dose of magnesium sulfate, in No. 18, which would remove water from the system, did not seem to cause any change.

In experiment 16 the last meal before bleeding contained a large amount of salt, and the surface tension was raised a little, as it should be if much salt was taken up by the blood. The effect of sugar in the diet is shown in No. 17, where there is a decided decrease. This may be because glucose lowers the surface tension of solutions.

HUMAN SERUM

The results on human serum are not as satisfactory as those on dog serum, as it has been shown that the amount and kind of food eaten has an effect on the surface tension of the serum, and the conditions governing the human subjects were not regulated, as was the case with the dogs. Most of this serum was obtained from St. Luke's and Sloan Hospitals, and some was given by fellow workers in the laboratory.

Results on human serum.

No.	Tip	1 drop gr.	W mg.	γ	Remarks
1	I	0.0499			
		0.0500	49.5	44.5	
		0.0500			
2	I	0.2577	51.5	46.2	Placental
		0.2578			
3	II	0.0507	50.3	45.9	"
4	II	0.0511			
		0.0514	50.8	46.4	"
		0.0512			

5	II	0.0502 0.0503	49.8	45.5	"
6	II	0.0520 0.0521 0.0520	51.6	47.1	Mict
7	II	0.0530 0.0532	52.6	48.0	Kidney
8	III	0.0563 0.0565	55.9	51.4	Chronic nephritis
9	III	0.0517 0.0519	51.3	47.1	Same, later
10	I	0.0499 0.0501	49.5	45.1	" "
11	III	0.0535 0.0535	53.0	48.8	Loc. atax.
12	III II	0.0492 0.0495	48.7 49.0	44.8 44.7	Apoplexy
13	III	0.0489 0.0491 0.0492	48.6	44.7	J. S. B.
14	III	0.04876 0.04856 0.04863	48.1	44.3	"
15	III	0.0499 0.0498	49.35	45.4	F. R. E.
16	III	0.0506 0.0506 0.05068	50.1	46.1	H. E. W.
17	III	0.04968 0.04952 0.04964	49.1	45.2	"

NOTE:—In all of these, except No. 2 in which five drops were weighed, a single drop was weighed in the empty vessel and the weight of the vapor, as a result of several determinations, was taken as 0.5 mg.

Of these, No. 1—5 and 13—17 are probably quite normal, and give an average value of 45.4 dynes as the surface tension of human serum, but most of them vary considerably from this because the conditions were not controlled as was the case with the dogs. The placental sera (No. 2—5) are rather high (average = 46.0), probably because the patients had not been eating as much as the others (average = 45.0). The blood in No. 13 and 14 was obtained in the afternoon while it was taking up the digested food, which is the reason for the low value. No. 15 was obtained about an hour after eating, before the blood could absorb any food. No. 16 was taken at noon—about eighteen hours after eating, since nothing was eaten that morning—while a few days later No. 17 was taken about four hours after eating, thus showing the great variation in one individual at different times of day.

Several of the pathological sera show a higher surface tension than would be expected even from a low diet. In the case of the kidney trouble and nephritis this is probably due to retention of salt, and in the latter it accompanies a high blood pressure.

COMPARISON OF DIFFERENT ANIMAL SERA

The blood serum of horse, rabbit, guinea-pig and sheep was obtained from Rockefeller Institute and St. Luke's Hospital, for comparison with dog and human serum. The results do not agree as closely as they did on dog serum, since the condition of these animals was not governed quite as carefully as with the dogs.

Results on animal serum

No.	Tip	Animal	1 drop	W	γ
1	III	Horse	0.0516 0.0514 0.0514	51.0	46.9
2		"	0.0488 0.0488 0.0488	48.3	44.5
3		Rabbit	0.0496 0.0495	49.1	45.2
4		"	0.0506 0.0505	50.1	46.1
5		"	0.0521 0.0521 0.0521	51.6	47.5
6		Guinea pig	0.0503 0.0503	49.8	45.3
7		Sheep	0.05223 0.05229	51.7	47.6
8		"	0.04922 0.04941	48.8	44.9

Comparison of surface tensions

Human	from	44.3	to	46.4	av.	45.4
Dog		44.7		45.9		45.3
Horse		44.5		46.9		45.7
Rabbit		45.2		47.5		46.3
Guinea pig		45.3				45.3
Sheep		44.9		47.6		46.2

The surface tension of blood serum is much below that of water, but this is not surprising when the number of substances in solution is considered. The list includes

neucleoprotein, serum globulin, serum albumin, dextrose, fat, enzymes, lecethin, cholesterol and esters, gases, coloring matter and inorganic salts such as chlorides, phosphates, carbonates and sulfates of sodium potassium and magnesium. The only ones of this list which raise the surface tension of water are the inorganic salts, while the proteins, being colloids, would lower it somewhat and emulsions of fat, lecethin and chloesterol lower it still more.

The surface tension of serum has been mesured by a French investigator* using a stalagmometer. In this method the number of drops of serum in a certain volume is counted and compared with the number of drops of water in the same volume of water as follows—

$$\frac{\text{No. of drops of water} \times \text{density of serum}}{\text{No. of drops of serum}} \times 75 = \text{surface tension}$$

He found it necessary to determine the water "constant" several times a day, so the method can not be very accurate. The use of 75 as the surface tension of water shows that he probably workt at low temperatures (about 5°C), instead of body temperature. His values vary from 69 to 74 (almost equal to pure water), while a few determinations by drop weights at 0° gave about 65. He gave results for different dilutions of serum with water, which do not agree very well. In some cases the very dilute solutions gave results above water, and in others they were nearer the value for undiluted serum. The drop weights of a few solutions of serum were taken, but no irregular curves like those were found. The drop weight work on solutions was not continued, as these did not seem to be any constant cure for dilution, and the very dilute solutions decompose too easily. A knowledge of the surface

*Iscovesco, *Comp. Rend. Soc. Biol.* 69, 353, 70, 1166, 93, etc.

tension of serum may be of importance in the case of injection of salt solution after loss of blood, as this is generally followed by a fever, since the surface tension of the injected salt solution is higher than that of the serum. It may be that a study of serum diluted with salt solution may be of importance in this connection.

Some Italian investigators* have used the capillary rise method on blood serum, but they worked at nearly all temperatures except body temperature, so the results can not be compared. Both the French and Italian workers have said that the surface tension of the serum is less than that of distilled water or isotonic salt solution, and varies according to the species and the individual. The results of this investigation show that the variation in any one individual may be as great as between different individuals, and the value for any individual is about 45 or 46 dynes.

MEIOSTAGMIN REACTION

Ascoli and Izar, of the University of Pavia, who originated the Meiostagmin reaction, showed that when a specific antigen is added to a specific serum, the surface tension of the liquid is lowered after the antibody, present in the pathological serum, has bound the antigen, by incubation for two hours at 37°. The reaction has been tried on typhoid, syphilis, tuberculosis and cancer serum, being positive in about 90% of the cases which are positive by standard methods, and nearly always negative in negative cases.

This reaction was tried on the drop weight apparatus in a series of syphilis cases—about 25 being positive by the Wassermann and 5 negative. The serum came from St.

*Fano and Mayer, *Arch. di Fisiol.* 4, 165

Luke's and Roosevelt Hospitals, and the Wassermann tests were made in their laboratories. It seemed best to make the determinations at 0° instead of at room temperature, so that no reaction would be going on while the determinations were being made, and water was used for diluting since there were no corpuscles present,—otherwise the reaction was carried out according to directions. The reaction does not seem to be as quick as many such reactions. When red blood corpuscles, for example, are treated at 0° with an antibody (from the serum of an animal which has been immunized against these corpuscles), and the corpuscles are immediately centrifuged away from the solution, it is found that they have combined with all the antibody in the few seconds that they were together. In the Meiostragmin reaction it takes about two hours for complete union of the antibody and antigen at 37° and there seems to be almost no union at 0° .

Izar used alcoholic ethereal splenic extract from a syphilitic fetus for an antigen, but in this work the antigens were mostly from beef heart. The following were used,—

- Antigen I The acetone insoluble part of an alcoholic and ethereal extract of beef heart, in alcohol solution. Titrated strength is .04 cc. to .1 cc. of serum.
- “ II An alcoholic extract of guinea pig heart. Strength is .06 cc. .1 cc. serum.
- “ III Same as I. Strength is .01 cc. to .1 cc. serum.

The solutions were made up so that the serum was 1 to 25.

Antigen	I	0.2 cc.	II	0.3 cc.	III	0.05 cc.
Water		11.8		11.7		11.95
Serum		<u>0.5</u>		<u>0.5</u>		<u>0.5</u>
Total		12.5		12.5		12.5

Half of the solution was put in the incubator, while the other half was put in the apparatus and its surface tension determined at once. The weight of five drops was found and the determination was repeated to get a check within 1 mg.—this allowed an error of about 0.1% and most of the changes were over 0.5%.

It seemed to make no difference whether the serum was inactivated (by heating to 55° for 30 minutes) or not, but most of the samples were inactivated, as they had been used for the Wassermann test in which the serum must contain no complement. In one case the Meistagmin reaction was tried both before and after inactivation of the serum. Inactivation destroys the complement (which is present in all serum), but leaves the immune body, and it is the latter which is necessary for this reaction.

Tip III			Fresh Serum			Antigen III		
Before incubation			After incubation					
5 drops	W	γ	5 drops	W	γ			
0.3976			0.3955					
<u>0.3981</u>	79.57	73.22	<u>0.3965</u>	79.20	72.88			
0.39785			0.3960					

Same Serum after 30 minutes heating at 55°

0.4072			0.4057					
<u>0.4070</u>	81.42	74.35	<u>0.4047</u>	81.04	74.01			
0.4071			0.4052					

Change in both cases is -0.5%

The Meistagmin reaction works therefore, whether the serum is fresh and contains complement or whether it has lost its complement either by keeping it three or four days or by heating it.

Five non-syphilitic cases were tried, four being negative by the Wassermann test, and the fifth being blood serum of the author which was used as a control.

Tip III Results on non-syphilitic cases

		Before incubation			After incubation			Change
No.	Ant.	5 drops	W	γ	5 drops	W	γ	%
1	I	0.3753	75.02	68.51	0.3757	74.99	68.49	0.
		<u>0.3749</u>			<u>0.3742</u>			
		0.3751			0.37495			
2	II	0.3577	71.62	65.36	0.3593	71.93	65.66	+0.3
		<u>0.3586</u>			<u>0.3600</u>			
		0.3581			0.35965			
3	III	0.4056	81.07	74.60	0.4048	80.90	74.45	-0.2
		<u>0.4051</u>			<u>0.4043</u>			
		0.40535			0.40455			
4	III	0.4006	80.16	73.76	0.4008	80.16	73.76	0.
		<u>0.4010</u>						
		0.4008						
5	III	0.3997	80.04	73.65	0.3999	80.04	73.65	0
		<u>0.4007</u>			<u>0.4005</u>			
		0.4002			0.4002			

The negatives do not show more than 0.2% decrease, and that is about the limit of accuracy with these solutions, as it means a check of 0.8 mg. on the weight of five drops before and after incubations. The one in which Antigen II was used, probably showed an increase on account of the large amount of the alcohol solution which had to be used.

Results on positive syphilis

		Before incubation			After incubation			Change
No.	Ant.	5 drops gr.	W mg.	γ	5 drops gr.	W mg.	γ	%
6	I	0.7507	75.03	69.04	0.7412	74.14	67.30	-2.5
		<u>0.7499</u>			<u>0.7416</u>			
		0.7503			0.7414			

7	<u>0.3337</u> 0.3343 0.3340	66.80	61.00	<u>0.3261</u> 0.3262	65.23	59.54	-2.4
8	<u>0.3635</u> 0.3632 0.36335	72.67	66.87	<u>0.3565</u> 0.3571 0.3568	71.36	65.67	-1.8
9	<u>0.7927</u> 0.7932 0.79295	79.30	72.42	<u>0.3969</u> 0.3959 0.3964	79.28	72.40	0
10 II	<u>0.3537</u> 0.3547 0.3542	70.84	65.15	<u>0.3343</u> 0.3350 0.3346	66.92	61.60	-5.5
11	<u>0.3591</u> 0.3591	71.82	66.07	<u>0.3523</u>	70.46	64.87	-1.8
12	<u>0.3575</u> 0.3567 0.3571	71.42	65.72	<u>0.3550</u> 0.3539 0.3544	70.89	65.23	-0.8
13 III	<u>0.3783</u> 0.3797 0.3790	75.80	69.75	<u>0.3710</u> 0.3716 0.3713	74.26	68.33	-2.
14	<u>0.3963</u> 0.3974 0.3968	79.37	73.04	<u>0.3885</u> 0.3881 0.3883	77.76	71.55	-2.
15	<u>0.3881</u> 0.3872 0.3876	77.53	71.34	<u>0.3818</u> 0.3810 0.3814	76.28	70.19	-1.6
16	<u>0.3955</u> 0.3951 0.3953	79.06	72.75	<u>0.3914</u> 0.3915	78.29	72.04	-1.
17	<u>0.3976</u> 0.3981 0.3978	79.57	73.22	<u>0.3955</u> 0.3965 0.3960	79.20	72.88	-0.5

18	0.4070 <u>0.4064</u> 0.4057	81.34	74.85	0.4053 <u>0.4048</u> 0.4050	81.01	74.55	-0.4
19	0.4033 <u>0.4031</u> 0.4032	80.64	74.20	0.4025 <u>0.4017</u> 0.4021	80.42	74.00	-0.3
20	0.3965 <u>0.3961</u> 0.3963	79.26	72.94	0.3951 <u>0.3955</u> 0.3953	79.06	72.75	-0.26
21	0.4046 <u>0.4050</u> 0.4048	80.96	74.50	0.4039 <u>0.4045</u> 0.4042	80.84	74.39	-0.15
22	0.4027 <u>0.4033</u> 0.4030	80.60	74.17	0.4027 <u>0.4027</u>	80.54	74.11	-0.1
23	0.3909 <u>0.3919</u> 0.3914	78.28	72.03	0.3721 <u>0.3731</u> 0.3726	74.52	68.57	-4.8
24	0.3949 <u>0.3954</u> 0.3951	79.03	72.72	0.3873 <u>0.3883</u> 0.3878	77.56	71.37	-1.9
25	0.3985 <u>0.3979</u> 0.3982	79.64	73.28	0.3918 <u>0.3910</u> 0.3914	78.28	72.03	-1.7
26	0.3992 <u>0.3992</u>	79.84	73.47	0.3944 <u>0.3933</u> 0.3938	78.77	72.49	-1.3
27	0.4024 <u>0.4030</u> 0.4027	80.54	74.10	0.3999 <u>0.4005</u> 0.4002	80.04	73.65	-0.6

28	0.3997	79.97	73.59	0.3974	79.58	73.23	-0.3
	<u>0.3990</u>			<u>0.3984</u>			
	0.3993			0.3979			

NOTE:—In No. 6 and the first half of 9, ten drops were weighed, instead of five. No. 7 and 9 were on Tip II, all the others on Tip III. In No. 23—28, the solutions were made up so that the serum was 1 volume in 10, instead of 1 in 25, with apparently no difference in results. Results on four other positive cases which gave a decided decrease in surface tension are omitted as the results were not checked very closely on account of the tip not being cleaned well enough.

In all but three of these there is a decided decrease in surface tension after incubation, but in No. 9, 21 and 22 the change is no greater than might be found in a negative case, so these can not be called positive by the Meios-tagmin reaction. The history of these three cases, however offers an explanation. No. 9 was a case in which the primary occurred twenty-five years ago, and it was cured at that time by the regular treatment. The Wassermann test was a very weak positive, as titration showed only two units of antibody while an active case contains about eight or ten times as much. No. 22 was a case of tertiary syphilis in which the primary was ten years ago. It gave a positive Wassermann with one antigen and negative with another. The Wassermann test was not tried with the antigen which was used in the Meios-tagmin. No. 21 was a case of early tabes probably on syphilitic basis. The Meios-tagmin reaction is almost positive, but is very doubtful. No. 20, which gives a weak positive Meios-tagmin, gave a doubtful positive Wassermann test. The case had been treated for about eight months with mercury and potassium iodide.

The Meios-tagmin reaction is evidently not quite as delicate as the Wassermann at present, tho it probably would be more delicate if the average of three or four determinations were taken instead of two, or if more drops

were weighed. It is a simpler test to make, for it only requires the serum and antigen, while the Wassermann requires inactivated serum, antigen, guinea pig serum for complement, sheep blood corpuscles and rabbit serum which has been injected with sheep corpuscles, all of which must be tested to see if they can be used together.

SUMMARY

1. The surface tension of the blood serum in an individual may change during the day, depending on the food that is being absorbed by the blood.

2. The surface tension of the blood serum of different individuals and different species is approximately the same, if account is taken of the daily change in any one individual.


3. The surface tension of the blood serum seems to be abnormally high in certain diseases, especially those in which the kidneys are affected.


4. The Meistagmin reaction was found to be specific in about 87% of a series of clinically positive syphilis cases.


5. The clinically positive cases of syphilis in which the Meistagmin reaction was not positive were those in which the Wassermann test was weak or rather doubtful.

VITA

Harold Edward Woodward was born in Worcester, Massachusetts, July 5, 1888. He received the degree of Bachelor of Arts from Amherst College, in June, 1910. Since then he has been studying Chemistry at Columbia University, under the Faculty of Pure Science, for the degree of Doctor of Philosophy. During the year 1910-1911 he taught Chemistry in Horace Mann High School, and he was Samuel Anthony Goldschmidt Fellow in Chemistry for 1911-1912.







Woodward

QP99
W87
1912

A study of the surface tension
of blood serum.

QP99
W87
1912
Anne

BOUND

NOV 29 1956

