## Infant mortality and milk stations / Edited by Philip Van Ingen and Paul Emmons Taylor.

#### **Contributors**

Van Ingen, Philip. Taylor, Paul Emmons. Augustus Long Health Sciences Library

#### **Publication/Creation**

New York : New York Milk Committee, 1912.

#### **Persistent URL**

https://wellcomecollection.org/works/hz272rwm

#### License and attribution

This material has been provided by This material has been provided by the Augustus C. Long Health Sciences Library at Columbia University and Columbia University Libraries/Information Services, through the Medical Heritage Library. The original may be consulted at the the Augustus C. Long Health Sciences Library at Columbia University and Columbia University. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



# RECAP


Ingen, P. V. and Taylor, Emmons
... Infant Mortality and Milk Stations

RA610-V26

# Columbia University in the City of New York

College of Physicians and Surgeons Library







## INFANT MORTALITY AND MILK STATIONS

## SPECIAL REPORT

OF THE COMMITTEE FOR THE REDUCTION
OF INFANT MORTALITY OF THE

NEW YORK MILK COMMITTEE

1912

111

THE NEW YORK MILK COMMITTEE

105 EAST TWENTY-SECOND STREET

NEW YORK

RAGIO

V26

Columbia University in the City of New York

College of Physicians and Surgeons




Reference Library



Digitized by the Internet Archive in 2010 with funding from Open Knowledge Commons

# INFANT MORTALITY AND MILK STATIONS





CONSULTATION DAY AT THE MILK STATION

## INFANT MORTALITY AND MILK STATIONS

SPECIAL REPORT DEALING WITH THE PROBLEM OF REDUCING INFANT MORTALITY. WORK CARRIED ON IN TEN LARGEST CITIES OF THE UNITED STATES

#### TOGETHER WITH

DETAILS OF A DEMONSTRATION BY PUBLIC AND PRIVATE AGENCIES IN NEW YORK CITY DURING 1911 TO DETERMINE THE VALUE OF MILK STATION WORK AS A PRACTICAL MEANS OF REDUCING INFANT MORTALITY

### Edited by

## PHILIP VAN INGEN, M.D.

Member of the New York Milk Committee and Medical Director of the Committee for the Reduction of Infant Mortality

and


### PAUL EMMONS TAYLOR

Secretary of the New York Milk Committee and Director of the Committee for the Reduction of Infant Mortality

PUBLISHED BY
THE NEW YORK MILK COMMITTEE
1912

Infant mortality statistics presented in this report were prepared By Philip Van Ingen, M.D.

PRICE, ONE DOLLAR



## The New York Milk Committee

ORGANIZED BY THE NEW YORK ASSOCIATION FOR IMPROVING THE CONDITION OF THE POOR IN 1906 **INCORPORATED IN 1910** 

#### Purpose

Create through education and publicity a public demand for a clean and safe milk supply for all. Prevent unnecessary loss of infant life.

#### OFFICERS

STEPHEN G. WILLIAMS, Chairman THOMAS S. McLane, Vice-Chairman and Treasurer PAUL E. TAYLOR, Secretary

#### ACTIVE MEMBERS

THE OFFICERS AND HAVEN EMMERSON, M.D. STEPHEN FRANCISCO Mrs. J. Borden Harriman Philip Van Ingen, M.D. MISS ALICE LAKEY CHARLES E. NORTH, M.D. Mrs. A. M. Palmer GODFREY R. PISEK, M.D. MARSHALL L. WARRIN IRA S. WILE, M.D.

#### ADVISORY COUNCIL

E. H. BARTLEY, M.D. HENRY DWIGHT CHAPIN, M.D. V. EVERIT MACY HENRY L. COIT, M.D. HENRY IVES COBB LEWIS A. CONNER, M.D. Mrs. Arthur M. Dodge Rt. Rev. David H. Greer A. Jacobi, M.D. REV. WALTER W. LAIDLAW

WALTER W. LAW FRANKLIN D. ROOSEVELT JOHN G. SAXE DEWITT J. SELIGMAN JOHN SPARGO Mrs. Henry Villard ARTHUR E. WAKEMAN GEORGE W. WICKERSHAM

## Committee for the Reduction of Infant Mortality

GODFREY R. PISEK, M.D., Chairman Mrs. J. Borden Harriman, Vice-Chairman PAUL E. TAYLOR, Director

#### EXECUTIVE COMMITTEE

THE OFFICERS AND S. Josephine Baker, M.D. ELI LONG, M.D. Mrs. J. J. Moorhead Mrs. A. Gordon Norrie E. B. Phelps Mrs. Nathaniel Bowditch Potter Rev. A. W. McCurdy MRS. BELMONT TIFFANY IRA S. WILE, M.D. PHILIP VAN INGEN, M.D.

#### GENERAL COMMITTEE

MRS. FELIX ADLER REUEL A. BENSON, M.D. HOMER FOLKS MISS ALICE P. GANNETT ROBBINS GILMAN JOHN A. KINGSBURY THOMAS M. MULRY

PROF. M. I. PUPIN Mrs. V. G. Simkhovitch MISS M. DEG. TRENHOLM MORRIS D. WALDMAN MISS MARIANNA WHEELER GAYLORD S. WHITE EGERTON L. WINTHROP, JR.

#### MEDICAL COUNCIL

ABRAHAM JACOBI, M. D., Chairman WALTER LESTER CARR, M.D. HENRY DWIGHT CHAPIN, M.D. FLOYD M. CRANDALL, M.D. ROWLAND G. FREEMAN, M.D. S. S. GOLDWATER, M.D.

L. EMMETT HOLT, M.D. L. E. LAFETRA, M.D. C. G. KERLEY, M.D. WILLIAM P. NORTHRUP, M.D. N. R. NORTON, M.D. JOHN A. WYETH, M.D.



## Table of Contents

Introduction	PAGE
CHAPTER I.—THE PROBLEM  Statistics of Infant Mortality  (a) In Foreign Countries.  (b) In the Registration Area of the United States.  Conditions Affecting Infant Mortality.  Poverty and Ignorance, Resulting in—  (a) Unsanitary Housing.  (b) Neglect of Babies and Expectant Mothers.  (c) Impure Milk.  (d) Bad Methods of Feeding.  (e) Crime and Illegitimacy.  Relation of Meteorological Conditions to Infant Mortality.	17
CHAPTER II.—THE CAMPAIGN AGAINST INFANT MORTALITY IN NEW YORK CITY (PRIOR TO 1911).  Beginning and Growth of the Movement. Work of Various Charitable Organizations. Work of the Health Department. Co-operation Between Organizations. Development of Prevention.	22
CHAPTER III.—THE CAMPAIGN IN NEW YORK CITY IN 1911  Work of the New York Milk Committee.  (a) Establishment of Thirty-one Milk Stations.  (b) Plan of the Campaign.  (c) Tremendous Growth of Movement.  (d) Appropriation Granted by Board of Estimate and Apportionment for Fifty-five Milk Stations in 1912.  Work of the Health Department.  Work of Other Organizations.	27
CHAPTER IV.—RESULTS OF THE CAMPAIGN IN NEW YORK CITY AS SHOWN BY COMPARATIVE TABLES	36
CHAPTER V.—RESULTS AT THE MILK STATIONS OF THE NEW YORK MILK COMMITTEE. A STATISTICAL STUDY OF THE RECORDS	52

VIII	TABLE OF CONTENTS	
Сн	APTER VI.—CAMPAIGNS IN OTHER CITIES	PAGE 84
	Statistical Study of Infant Mortality.	
	Weather Conditions. Efforts at Reducing Infant Mortality.	
	Chicago.	
	Philadelphia.	
	St. Louis. Boston.	
	Cleveland.	
	Baltimore. Pittsburgh.	
	Detroit.	
	Buffalo.	
Сн	APTER VII.—GENERAL CONCLUSIONS	122
CII	Summary of Infant Mortality in Various Cities.	122
	Comparison of Weather Conditions in Various Cities.	
	Comparative Summary of Campaigns Against Infant Mortality.  General Deductions.	
	General Deductions.	
Сн	APTER VIII.—THE EXTENSION OF RELIEF THROUGH MILK STATIONS	131
	Method of Handling Free Milk Problems.	
	Statistical Study of Milk Relief Applications. Other Forms of Relief.	
	Ice.	
	Hospital Care. Outings.	
	Conclusions.	
_		
Сн	APTER IX.—ESTABLISHMENT AND OPERATION OF MILK AND HEALTH STATIONS	147
	Survey of Needs.	141
	Cost and Equipment of Station Plants.	
	Organization and Operation.	
	Cost of Operation. Physicians.	
	Nurses.	
	Matrons.	
Сн	APTER X.—Extension of Milk Stations' Activities	165
	Education of Expectant Mothers.	
	Control of Midwives. Supervision of Boarded-out Babies.	
	Care of Children Under School Age.	
Δрі	PENDLY	174
AF	Tables Showing the Quality of Milk Dispensed by the Committee's Stations and that	1/4
	Dispensed by City Stations.	
	Summary of Milk Committee's Work for Season.	
Ex	HIBITS	

ILLUSTRATIONS

## Introduction

The purpose of this report is to make available to every one having an interest in the campaign against infant mortality the information acquired by the Committee for the Reduction of Infant Mortality of the New York Milk Committee during the summer of 1911, when it undertook a demonstration to determine the value of infants' milk stations as a means of reducing that mortality.

It is not intended as a general report of the work done by the Committee, which is contained in the Fifth Annual Report of the New York Milk Committee. Statistics were carefully kept and tabulated. They are here presented, with the deductions drawn from them, together with the opinions derived from the experience of those who organized and directed and carried on the work.

In the first chapter is given a very brief summary of the problem, to help in the solution of which the milk station was called upon. A frightful tragedy, costing the lives of some 1600 men and women, horrifies the entire world, and yet every year many thousands of babies die unnecessarily. Tables are given showing that in the civilized world from one in every ten to one in every three babies born die during the first year of life.

The various factors concerned in this waste of life are very briefly discussed. In the next chapter is given a résumé of what has been done in New York city during recent years for the prevention of this high infant mortality. An attempt is made to trace the development of the activities working together today toward this end.

Chapter IV deals with the results obtained in New York city as a whole, and its different subdivisions. Tabulations of the official figures of the Health Department for a number of years are given. These tables show a most remarkable reduction in infant mortality for the year, and chiefly during the summer months. They show that, with a low total infant mortality, only 28 per cent. occurred during the three summer months in Manhattan, as against an average for the preceding ten years of 33.8 per cent.

A tabulation and brief discussion of weather conditions in New York is also given, which shows that in the first half of the summer conditions were very unfavorable for a low infant mortality. The first part of July was the most unfavorable since 1901, and yet the deaths during that month were the lowest since 1901, not even taking into consideration the increase in population.

Chapter V deals with the results obtained at the milk stations. Careful records were kept containing all data which could be used to any advantage. The feeding records of 3182 babies under two years are analyzed and studied in some detail. Of these babies, only 158 died, and of them, 58 were never treated in any way by the station physician or nurses, being immediately sent to a hospital

or private physician. The milk stations were carried on to prevent sickness among babies. Many facilities exist for treating sick ones. Every baby who died, however, if it once came inside the station, was charged against that station, in order to avoid possible charges of selecting cases. The results show the entire feasibility of teaching the mother to modify the milk at home.

Detailed studies of mortality in a large portion of the borough of Manhattan were made by one of the Committee, in order to try to show the relation of the mortality among the station babies to that in the blocks surrounding the stations and from which the babies came. The results are rather striking.

In order to determine whether results in New York, especially the boroughs of Manhattan, where the Committee was working, and Brooklyn, where the Health Department was most active, were better than in other cities, as complete records of their vital statistics as possible were made, together with an attempt to find out what was being done there. The nine largest cities after New York were selected. More could not be attempted, owing to the difficulty of securing information. The lack of published statistical records makes it very difficult to make a thorough study of conditions. Comparatively few cities outside of the largest ones publish monthly bulletins of vital statistics. Of those that do, the data printed in regard to infant deaths are often scanty and classified entirely differently in different cities. Even the Census Bureau, in its splendid yearly reports on mortality, does not tabulate *infant deaths* by months of occurrence, though it does adults. For many of the data the Committee is indebted to health officers, who supplied them on written request.

In Chapter VI the various data collected are tabulated and summarized, and an attempt is made to compare results fairly and impartially. The Committee believes the value of the milk station as a means to reduce infant mortality through educational prevention is proved.

In the chapters that follow, the organization of the stations and the central office is shown for the benefit of any communities planning to organize such a work. The cost of maintenance is most carefully and fully worked out. The problem of relief and the details of its management are given fully. A chapter is devoted to the future of the milk station and its possibilities of usefulness.

The work of the Committee was carried on to help the Department of Health demonstrate the value of milk stations in reducing infant mortality. The city appropriated funds in 1910 to maintain 15 municipal milk stations. When the Board of Estimate and Apportionment met to pass on the appropriations for 1912, the Committee submitted a report to each member stating what had been accomplished, and illustrating results with maps, charts, and tables. It also was represented at the public hearing on the budget, and that the city increased the previous appropriation so that the Department of Health could maintain 55 in 1912, they feel sets the official seal of approval on the work of milk stations,—not theirs,—but what is far more important, milk stations as a means for the prevention of infant mortality through education.

The Committee wishes in this way to acknowledge the help and coöperation of many—it is impossible to mention them by name, they were so many. The

various relief organizations, settlements, and hospitals gave much assistance; without it the work would have been impossible. The coöperation of the Health Department and the Bureau of Municipal Research was much appreciated. The ice companies did their share in generously supplying free ice to those unable to buy it.

Our thanks are due to the many health officers who provided statistical material and information. Many physicians and private individuals also furnished valuable information.


And, finally, we wish to express our thanks and appreciation to the staff of physicians, nurses, matrons, and the office force, who worked in season and out of season, early and late, under most trying and almost impossible conditions, giving their very best to their work. Without such a spirit on the part of all we could not have accomplished what we did.



DR. P

SIO, TO Commis Addi Kees REVIEN





#### CHAPTER I

## The Problem

"A baby who comes into the world has less chance to live one week than an old man of ninety; and less chance to live a year than one of eighty."

So said Bergeron some years ago, and it is practically true today. In 1900, 2.2 per cent. of the total population of the Registration Area\* were under one year of age. Assuming the same proportion to hold good in 1910, among 2.2 per cent. of the population occurred† 19.2 per cent., or nearly one-fifth, of all the deaths. The infantile mortality rate‡ in eight States having fairly complete birth registration varied in 1910 from 108 to 168 (†); that is, from one in every six to one in every ten, babies died before they were one year old.

These figures are not exceptionally high, as will be seen from the following table:§

TABLE 1.—SHOWING GENERAL AND INFANT DEATH-RATES IN FOREIGN COUNTRIES IN 1909||

	Course Double Day	T D
n	GENERAL DEATH-RATE	INFANT DEATH-BATE
England and Wales	14.5	109
Scotland		121**
Ireland		92
New South Wales	9.6	74
Victoria		71
Queensland		72
S. Australia	9.3	61
W. Australia		78
Tasmania		65
New Zealand		62
Denmark		123**
Norway		76**
Sweden		85**
Prussia		164
		212
Hungary	22.2**	
Austria		204††
Servia		158
Netherlands		99
Belgium	16.5**	147**
France		135††
Switzerland	16.2**	108**
Spain	23.4	173tt
Italy		153**
Chili		320**

<sup>\*</sup> The registration area in 1900 comprised 40.5 per cent. of the population of continental United States. It embraces those States and cities which have adequate laws for registration of vital statistics.

<sup>†</sup> Census Bureau Bulletin, 109.

<sup>‡</sup> Infant mortality rate is deaths of infants per 1000 living births for same period.

<sup>§</sup> From Seventy-second Annual Report of Registrar General of Births, Deaths, and Marriages in England and Wales (1909).

<sup>||</sup> General death-rate is number of deaths at all ages per 1000 of population. Infant death-rate is number of deaths under one year per 1000 births.

<sup>\*\* 1908</sup> figures.

Ireland, Norway, Sweden, The Netherlands, and the great Australian colonies, New Zealand and Tasmania, alone fall below 100; while in Prussia, roughly, one in every six; Austria and Hungary, one in every five; and Chili, one in every three, babies die before they are a year old.

In order to understand the problem we must analyze these figues still further. It is necessary to know when the babies die, and from what causes. The following tables show the age at death and the cause of death (general groups of diseases) in England and Wales in 1909:

TABLE 2.—INFANT MORTALITY IN ENGLAND AND WALES. NUMBER OF DEATHS PER 1000 BIRTHS AT VARIOUS AGES AND FROM VARIOUS GROUPS OF DISEASES, BASED ON FIGURES FOR 1909\*

	UNDER ONE DAY	ONE DAY TO ONE WEEK	ONE	Two	THREE	UNDER ONE MONTH	Under One Year		
I. Common infectious diseases II. Diarrheal diseases III. Wasting diseases IV. Tuberculous diseases V. Miscellaneous	10.14	10.68	3.82	3.16	2.08	29.88	41.69, or 38.3%	infant " " "	deaths
All causes	11.55	13.18	5.70	5.33	3.99	39.20	108.73		

#### By Months.

	1	1-2	2-3	3-4	4-5	5-6	6-7	7-8	8-9	9-10	10-11	11-12	UNDER 1 YEAR
I. II. III. IV. V.	0.11 0.80 29.88 0.06 8.90	0.36 1.47 4.30 0.20 5.26	0.36 1.63 2.35 0.32 4.12	0.33 1.50 1.46 0.34 3.44	0.32 1.47 1.02 0.44 3.18	0.42 1.23 0.71 0.41 2.93	0.54 1.02 0.52 0.38 2.89	0.74 0.89 0.40 0.36 2.76	0.85 0.81 0.32 0.40 2.72	0.93 0.72 0.30 0.39 2.58	0.97 0.56 0.23 0.35 2.35	1.04 0.54 0.20 0.35 2.30	6.97 12.64 41.69 4.00 43.40
All		11.59	8.78	7.07	6.43	5.70	5.35	5.15	5.10	4.92	4.46		108.73

- I. Includes smallpox, chicken-pox, measles, scarlet fever, diphtheria, whooping-cough.
- II. All gastro-intestinal diseases except gastritis and gastric catarrh.
- III. Includes premature births, congenital defects, want of breast milk, starvation, atrophy, debility, marasmus.

<sup>\*</sup> Compiled from Annual Report of Registrar General, England and Wales, 1909. It should be noted here that in Group V are included "meningitis, not tuberculous," and "convulsions," which amount to 10.43 under one year, and "gastritis, gastric catarrh," amounting to 1.83. The latter, in our classification, would probably be included in Group II, and the former are probably largely digestive in character. This should be remembered in comparing the above table with Table 4.

TABLE 3.—INFANT MORTALITY IN ENGLAND AND WALES. PERCENTAGE OF TOTAL DEATHS UNDER ONE YEAR OCCURRING AT VARIOUS AGES, BASED ON FIGURES FOR 1909

Of the infant death	1s-						
10.6% oc	curre	d under	1 day.				
12.1%	"	1 day			n 1 w	eek.	
5.2%	**	1 week	and	less t	han 2	week	rs.
4.9%	44	2 week		**	" 3	- 11	
3.6%	**	3 "	"	**	" 1	mon	th.
36.5%	44	under	1 mon	th of		mon	cii.
	**					2	
10.6%					than		onths.
8.0%	"	2 mor	iths '		"	3	"
6.5%	44	3			"	4	"
5.9%	"	4	11 11	1 11	44	4 5	11
5.2%	44	4 5	14 6	11	**	6	11
4 90%	44			11	**	7	**
4 70%	44	6 7			**	8	**
4 60%	**				**	0	**
4.5%	"	9			**	10	**
4.570						10	
4.1%		10			"	11	"
4.0%	"	11		"	"	12	**

Of these deaths, 10.6 per cent. occur under one day old, 22.7 per cent. under one week old, and over a third during the first month of life. From then on the proportion steadily decreases. The above tables also show that 38.3 per cent. of these deaths under one year occur from the so-called "wasting diseases," and 11.6 per cent. from diarrheal diseases. As noted above, the proportion of the latter is probably too low, but practically 50 per cent. are due to these two classes.

The following table shows the proportion of deaths under one year in the Registration Area of the United States due to these causes:

TABLE 4.—PERCENTAGE OF DEATHS UNDER ONE YEAR OF AGE IN THE REGISTRATION AREA OF THE UNITED STATES DUE TO DIARRHEAL AND WASTING DISEASE\*

	1900-4	1905-9	1900-9
Diarrhea and enteritis (II)	.23.72	26.41	25.23
Wasting disease (III)		27.79	26.83
	49.33	54.20	52.06

It will be seen that in the United States also a little over one-half of all the deaths under one year are due to these two classes of disease. Further, the proportion has shown a steady increase from year to year, as will be seen from the following figures:

TABLE 5.—PERCENTAGE OF ALL DEATHS UNDER ONE YEAR DUE TO DIARRHEAL AND WASTING DISEASES IN THE REGISTRATION AREA OF THE UNITED STATES\*

1900	190552.46 per cent.
190148.40 "	190653.21 "
190247.57 "	190754.23 "
190349.96 "	190855.59 "
190451.84 "	190955.03 "

Such, then, in brief is the problem which confronts us. When we look still

<sup>\* &</sup>quot;A Statistical Survey of Infant Mortality's Urgent Call for Action," Phelps. Amer. Assoc. for Study and Prevention of Infant Mortality, Annual Meeting, 1910.

further and try to discover the causes of this appalling state of affairs, we are brought face to face with practically the entire social problem of today.

#### CONDITIONS AFFECTING INFANT MORTALITY

Ignorance and poverty are written all over the situation. As has been said, one-quarter of all the deaths in babies are due to the so-called "wasting diseases." Poverty forces married women to be wage-earners. It forces the expectant mother to work long hours, often in the factory, under unsanitary conditions, with poor food and with her family at home to take care of after working hours. This is a poor preparation for the ordeal through which she must go. She is less likely to give birth to a baby who has a fair start in the world. Further, her own physical condition is such that she is less likely to be able to give the baby the food that nature intended it to have. Even if she can give the baby the breast, poverty again drives her to return to work immediately after her confinement. This forces her not only to feed the baby artificially, but to leave it to the care and tender mercies of a "Little Mother," often a mere child herself, or to confide it to the care of a day nursery.


Poverty forces the poor to live in overcrowded, unsanitary tenements. Exhibit 2 shows graphically the distribution of infant deaths along the East Side of Manhattan—New York city. In some of these blocks as many as 5000 people are herded together, and it is in these districts that the infant deaths occur in such large numbers. Compare the black dots near the river with those near Fifth Avenue. We are here confronted with the problem of congestion of population in all its phases.

Another quarter of the deaths among babies is due to diarrheal disease. This means bad food and bad methods of feeding. Deprived of its mother's milk, the baby must be artificially fed, and milk should be the chief article of diet during the first year. This brings us face to face with the entire milk problem. How is the city to be insured a clean milk supply? About 2,000,000 quarts of milk a day are brought into New York city from 44,000 farms in seven different States. To the average mother milk is milk. She is ignorant of the necessity for clean, pure milk. Poverty forces her to buy the cheapest milk she can get. Pure milk costs more to produce than dirty milk, but if buying food for the older members of the family, she would not willingly buy decomposing meat or vegetables. She is ignorant of the fact that impure milk is a poison to her baby, just as decomposing food is to the rest of the family.

Bad methods of feeding, even when the milk used is a good milk, are responsible for a vast number of deaths. The mother is ignorant of how to feed her baby, and it is not only the mother, but unfortunately often the physician as well, who is ignorant. When her baby is sickly, in desperation she tries anything her neighbors suggest—condensed milk, proprietary foods, or even solid food. Artificial feeding diminishes tenfold the baby's chance to survive the first year.

Finally, crime and illegitimacy come in as causes. The figures for England and Wales for 1909 show the death-rate among illegitimate children under one year of age to be 211.1, as against 104.8 for legitimate children—more than twice as high a mortality.





STUDY OF DEATH CERTIFICATES SHOWS THE HIGH INFANT MORTALITY AREAS-MANHATTAN-EAST SIDE (Canal to 75th Street) 4





EXHIBIT 2
A STUDY OF DEATH CERTIFICATES SHOWS THE HIGH INFANT MORTALITY AREAS (Each dot represents an infant's death in 1910)



### WHERE THE BABIES DIE IN BROOKLYN

(Each dot represents a baby's death in 1910)





EXHIBIT 3

IN LESS THAN ONE-HALF OF BROOKLYN'S TOTAL AREA OCCURS NINETY-FIVE PER CENT. OF THE BOROUGH'S INFANT MORTALITY



In general, then, the two great fundamental causes of infant mortality are poverty and ignorance.

Among the more remote causes meteorologic conditions are considered to play a large part in producing infant deaths. From one-third to almost one-half of the baby deaths in some cities are recorded during the third quarter of the year—July-September.

In a recent careful study of conditions in Berlin Lieffmann and Lindemann have drawn certain conclusions as to the effect of atmospheric temperature. They have constructed charts showing the daily deaths in Berlin, together with the course of the temperature for a number of years. They show that for the winter and spring months there is a sort of normal mean daily curve for deaths. In early summer, with the advent of hot days, there is a sharp rise in the number of deaths; with the fall of temperature there is an almost immediate fall in the deaths to the so-called normal. In midsummer and in late summer, while the sharp rises occur with each hot day, the fall does not recur to the "normal," but the basal curve tends to remain high. It is only after a considerable period of cool weather that it does so return.

They conclude, from their studies, that atmospheric heat above 23° C. (73.4° F.) has a directly injurious effect upon infants. They consider that a great number of deaths in hot weather are due to "heat-stroke" ("Hitzschlag"), just as in adults. While admitting that many babies, during the hot weather, do suffer from diarrheal disease, they consider that this is not the real cause of death. Artificial feeding, to their mind, is usually faulty in quality and quantity in hot weather. Cow's milk under these conditions causes an injury to the intestinal tract which makes such babies particularly susceptible to the effects of heat. Certain experiments on animals seem to bear out this theory.

This fact seems to explain the failure of the death-curve to return to the "normal" in later summer during cooler days. Many babies die from diarrhea each day, regardless of the temperature, and on each very hot day the "heat cases" are added.

Whether this is a true explanation or not; whether the theory that the heat diminishes the baby's resistance, acts unfavorably on the milk or food it gets, and so renders it an easy prey to the diarrheal disease brought on by faulty feeding or filthy milk, is accepted—the fact is not altered that the daily deaths do rise sharply in hot weather, and tend to remain high throughout the late summer. Exhibits 4 and 5, showing the daily deaths under one year in Manhattan in 1909 and 1910, and Exhibit 7, showing the same for Brooklyn for 1910, illustrate this very clearly.

Relative humidity has been less studied, and it is somewhat difficult to give its proper place. It is to be noted that humidity in dwellings is probably quite different from that recorded at weather bureaus.

Rainfall—the amount and distribution of the rain throughout the summer is thought to have an influence. Dry summers and long periods without rain, even when the total rainfall is large, are supposed to increase the infant mortality.

All these factors will be discussed more in detail in studying the conditions and results in 1911 compared with other years.

### CHAPTER II

### History of the Campaign Against Infant Mortality in Greater New York

Before describing in some detail the story of the campaign in New York in 1911, it will be well to take a brief glance at what was done in previous years. The New York Milk Committee is convinced that the phenomenal decrease in infant mortality in 1911 can in large part be laid to improved methods, and a view of the development of these methods will not be out of place.

In 1873 the New York Diet Kitchen Association was founded, its object being to distribute nourishing food to the sick among the very poor. "Good milk" was, of course, distributed to old as well as young.

In 1891, at the Good Samaritan Dispensary, milk was dispensed in summer to sick babies. It was modified to a set formula, suited to a definite age, and pasteurized. For several years this part of the work was carried on only in summer. Since 1898 it has distributed milk all the year.


In 1892 Mr. Nathan Strauss founded his well-known Pasteurized Milk Laboratories. Any baby in need of milk could obtain the day's supply, already prepared, and needing only to be warmed before using. Mr. Strauss's depots were situated in various parts of the city. As his plan grew summer stations were added to meet the demand. A safe, well-prepared milk was provided, the formulæ being provided by the best medical authorities in the city. This was a great step in advance, and paved the way for most later developments.


In Brooklyn, in 1901, the Children's Aid Society entered the field, following the lines of the Strauss stations. Their depots were open only a couple of months the first year and were in charge of a matron. The Health Department provided a physician to supervise the preparation of the milk. Eleven stations were opened, chiefly in settlements or hospitals, and 1087 babies were fed the first summer. In 1889 and 1890 the Brooklyn Bureau of Charities had started this work, which was later taken over by the above-mentioned society.


Before 1901 the work of the Health Department had been small in the field of infant mortality. As far as the annual reports show, until 1902 it consisted in a small summer corps who visited sick babies in the tenements.

In 1902, however, the work was reorganized, and a house-to-house canvass of the worst tenement districts was carried on. Nurses followed up the cases and carried out orders. Dispensaries were established at recreation piers; 206,121 families were visited. The inspection of the milk supply was undertaken—at the farm, in transit, and in the city. Funds were not available for a very extensive campaign, but the work was begun.

The Brooklyn Children's Aid Society continued their work, adding another







Daily Maximum and Mean Temperatures Compared With Daily Deaths of Infants-1910, 1911, Brooklyn



station, and caring for 1197 babies. This year they were open from June 24th to September 5th. The same general principle was followed, dispensing milk according to age, already modified. The Henry Street Nurses' Settlement began this year to dispense milk to the poor in their neighborhood in connection with their visiting work.

In 1903 the Health Department broadened out its work considerably. The summer corps of physicians and nurses began visiting the families of every baby whose birth was registered with the department between August 1, 1902, and June 30, 1903. A circular on "How to Keep the Baby Well" was mailed to each address. After three days it was found that 30 per cent. of the families had moved, and so many unregistered babies were found that a house-to-house canvass was started, and a report made on every child under one year found. Literature was distributed, and, when necessary, sick babies were treated by inspectors and nurses. Six inspectors chosen for their fitness were specially assigned to this. In Brooklyn the Children's Aid Society increased their stations to 14, feeding 1547 babies from June 17th to September 5th. In Manhattan the Strauss laboratories, the Good Samaritan, and the Diet Kitchen continued their work as before.

In 1904 the department continued its work along the same lines, visiting the tenement districts, tabulating all cases, and treating the sick children. The physicians visited 159,726 families and treated 1428 sick babies. Thirty-three nurses were doing district work, and visited 56,271 families and cared for 437 sick babies. They distributed orders for milk and ice as needed. In Brooklyn the Children's Aid Society improved on their method by attempting seriously to carry on some instruction. The matrons visited among the sick babies, and a corps of volunteer physicians (nine) followed up the sick cases. Again, 14 stations were maintained and 1233 babies fed from June 27th to September 3d.

In 1905 the Health Department increased its activities, employing 31 nurses in Manhattan, 14 in Brooklyn, 1 in the Bronx, and 2 each in Queens and Richmond. The families visited numbered 289,114, and 1985 sick babies were treated. The New York Diet Kitchen Association had been paying more and more attention to babies and to tuberculosis work, and in their reports only milk and eggs are mentioned as being distributed. In Brooklyn 15 stations were maintained by the Children's Aid Society from June 26th to September 9th, feeding 1649 babies.

In 1905 the milk problem was energetically handled. The New York Association for Improving the Condition of the Poor carried on a very extensive investigation with the coöperation of the Health Department. As a result of this investigation many violations of the law were discovered and offenders punished. It was the beginning of a vigorous campaign.

In 1906 a new agency came into the field. Mr. John D. Rockefeller accepted the suggestion made by the A. I. C. P. that the land adjoining the Rockefeller Institute be utilized for a summer hospital for sick babies, where mothers could be taught that they can "save their own babies, in their own tenement homes, if they will give them clean milk, clean air, and clean bodies."

The old Schermerhorn mansion, in a state of apparently hopeless decay, was rejuvenated by soap and water and plenty of whitewash. Five open-air shacks were provided, capable of accommodating 60 babies. Everything was of the simplest. A large shelter tent was provided, and "Junior Sea Breeze" was ready for work. Mothers were allowed as free access as possible to the shacks in order that they might learn that the simplest articles were sufficient to care for the baby; also how much had to be done when a baby was sick. With the hospital work proper was combined an educational campaign. In the shelter tent there were given, twice a week, simple lectures and demonstrations to the mothers in the care of their babies, the importance of maternal nursing, "clean milk, clean air, clean bodies." A nurse did follow-up and district work.

The Health Department continued its work on much the same lines, always increasing in amount: 350,618 families were visited, and 3331 sick children treated. The Children's Aid Society in Brooklyn continued as before, feeding 1620 babies at its 15 stations from June 25th to September 8th. There was little change other than Junior Sea Breeze. In 1906 the appropriation for milk inspectors for the Health Department was doubled, and a vigorous inspection was carried on.

In 1906 the New York Milk Committee came into existence as a result of a conference called by Dr. Darlington, then Commissioner of Health. This committee was formed to investigate the milk problem, especially in its relation to infant mortality. The Mayor also appointed a committee to investigate the problem of insuring a safe milk supply.

In 1907 the work of the Health Department, Diet Kitchen, Strauss laboratories, Good Samaritan, and Children's Aid Society of Brooklyn was continued with little change. The chief feature of the summer work was the enlarging of the scope of the work of Junior Sea Breeze. A very vigorous educational campaign was carried on. Nineteen nurses were employed in district work. This work was confined to the nineteenth ward, extending from Fortieth Street to Eighty-sixth Street, and from Sixth Avenue to the East River. A house-to-house canvass of the district was made; mothers were urged to feed their babies on wholesome milk if nursing was impossible, and watch was kept on the babies of the district. It was a piece of intensive work of great importance. While the summer deaths were higher in the rest of the city, in the nineteenth ward they were lower than the previous year.

The Association for Improving the Condition of the Poor also started this summer its Caroline Rest, where mothers and babies were sent as soon after the confinement as possible. Here the mother had a chance to recuperate and fit herself to nurse her baby. She was given the most careful instruction in the care of babies and older children.

The year 1908 marks the beginning of a very vigorous campaign, with emphasis on the educational factor. Up to this year the Department of Health had carried on its summer corps work as part of the Division of Contagious Diseases. In August, 1908, the Division of Child Hygiene was formed, in charge of a chief and 5 assistant chiefs, 16 supervising inspectors, 1 superintendent of nurses, 5 supervising nurses, and 195 district nurses. This was a very great step

in advance. The summer work was carried on by 83 nurses and 20 doctors. Visits were made to all new-born babies as the births were reported. Lectures were given at various places by Health Department inspectors, and it is noticeable that this year there was a very marked decrease in infant deaths and in the infant death-rate.

Early in the spring the Commissioner of Health issued an invitation to all organizations interested in any way in the campaign against infant mortality to attend a conference at the Health Department. This conference was held and was largely attended, and the coöperation of all agencies was earnestly asked for. In Brooklyn the department increased its coöperation by detailing inspectors and nurses to the stations of the Brooklyn Children's Aid Society. The work of this organization was, therefore, greatly increased, and the educational factor became very pronounced. The physicians and nurses were provided with simple medicines which were given to any sick children. From their 16 stations 2296 babies were fed. Seventy mothers' meetings and clinics were held, in charge of a physician and nurse. Literature was distributed in the homes, and free icetickets, provided by the Herald Free Ice Fund, were given when needed.

On June 17, 1908, the New York Milk Committee opened seven stations, from which they dispensed milk modified at the stations according to set formulæ. Volunteer physicians had charge of all the consultations, and trained nurses were employed to prepare the milk and supervise the babies in their homes. No mother was permitted to obtain milk from the station until it was proved that she was unable to nurse her baby, and she was allowed to obtain milk then only on condition that she would bring her baby once a week to be weighed during consultation hours. A careful record of all cases was kept as a basis for further work. From three of the depots pasteurized milk was dispensed, and from four, raw milk. Consultations were also held at the Union Settlement in East One Hundred and Fifth Street and in West Ninety-third Street, though milk was not dispensed from these places. Junior Sea Breeze continued its work as in the past year, although not so many nurses were employed in visiting work as in the previous year.

In 1909 the work of the Division of Child Hygiene was increased. It was made to include the supervision of midwives, boarded-out children, nurseries, and other institutions caring for children. From April 15th to September 1st the summer campaign was carried on. A great step in advance was made by beginning the campaign early. The summer corps visited 57,059 mothers of new-born infants.

Another important step in advance was the formation of the so-called "Little Mothers' League." There were 54 centers for this league, which enrolled the little mothers in the public schools, giving them lessons and demonstrations in the care of the baby and general hygiene. An inspector and a nurse held these meetings once or twice weekly in each center. The New York Milk Committee maintained seven stations throughout the year. This organization was originally a special committee of the New York Association for Improving the Condition of the Poor. In October of this year it became an independent organ-

ization, in accordance with the policy of the Association in allowing its offspring to become independent as soon as they were able to care for themselves.

In Brooklyn the Children's Aid Society continued its coöperation with the Health Department. Sixteen stations were maintained, caring for 2142 babies. Physicians and nurses were provided by the Health Department, and there was also coöperation by the District Nursing Committee. Increased emphasis was laid on educational work, maternal nursing, and the care and preparation of milk. Three hundred and twenty meetings were held during the

year.

In 1910 the work continued as before. The New York Milk Committee was able to maintain only four stations, owing to lack of funds. These were conducted also along different lines. The distribution of already modified milk was discontinued, and the policy of teaching mothers to prepare the baby's food themselves, under the supervision of the nurse, according to the doctor's direction, was carried out. It was with considerable hesitation that this work was undertaken, but the results soon proved its value. The mothers were more regular in their attendance, and took greater personal interest and pride in the progress of their babies when they themselves prepared the food. During this year the Health Department coöperated with the New York Diet Kitchen, providing physicians and nurses for eight of their nine stations. The ninth station was managed by the association itself. Physicians and nurses held at first weekly clinics, then semi-weekly, and, finally, during the summer, daily clinics. Expectant mothers were also cared for. In Brooklyn the Children's Aid Society continued as before. The importance of small classes and individual instruction was brought home to them, and greater attention was paid to this detail. From 18 stations 2182 babies were fed, of whom half were sick on admission, and only 32 deaths occurred.

It will be noticed that of late years, especially since 1908, prevention has been playing a greater and greater part in the attempts at the reduction of infant mortality. What has been given above is merely a very brief outline of certain phases of the work which has been carried on. It is impossible to give in detail everything that has been done. One of the most important agencies in the prevention of the waste of infant lives has been the adoption of the so-called social service work of Dr. Cabot, of Boston. Most of the hospitals and dispensaries of this city now have a social service department. It interests itself, among other things, with the care of babies and the prevention of disease among them. Another very important step has been the increasing cooperation developing among various agencies. The Health Department has been ready and willing to make the work which is being done in every way more efficient. In 1907 the department turned over practically all the work of visiting new-born babies in the nineteenth ward to Junior Sea Breeze Hospital. A list of births was provided to the Junior Sea Breeze nurses, just as to their own. This cooperation increased in a most encouraging way, as will be seen in the report of what has been done in 1911.




EXHIBIT 9

ORGANIZATION CHART OF THE COMMITTEE FOR THE REDUCTION OF INFANT MORTALITY OF THE NEW YORK MILK COMMITTEE



### CHAPTER III

### The Campaign in New York City in 1911

The year 1911 marks the beginning of the most extensive and efficient campaign for the reduction of infant mortality ever undertaken in New York city. The lessons of past years had apparently been learned. Here were a number of organizations, each working independently, with practically no general coöperation, though many worked well with the Health Department.

### CAMPAIGN OF THE NEW YORK MILK COMMITTEE

In the fall of 1910 the Board of Estimate and Apportionment appropriated sufficient funds to maintain, during the entire year, 15 infants' milk depots under city control. These stations were put under the control of the Division of Child Hygiene. The New York Milk Committee realized that this was a new venture, and that the extension of this system of milk depots depended on the success of the year's work. They also realized that the summer months were the ones when the greatest reduction ought to be obtained. To make the experiment a strongly convincing one, a sufficiently large number of stations must be provided to cover thoroughly a large portion of the city. The Health Department decided to place only five in Manhattan, one in the Bronx, and nine in Brooklyn. It was felt that the mortality of the city could not be demonstrably affected by the existing stations, so a subcommittee of the New York Milk Committee, known as the Committee for the Reduction of Infant Mortality, was formed. They were authorized to plan a campaign, raise money, and conduct their demonstration, subject to the approval of the Executive Committee of the parent body.

The new subcommittee selected an Executive Committee and a Finance Committee. The former committee thoroughly canvassed the city, studying the available records of population and mortality, and the presence of other similar organizations in given districts. As a result of this study it was decided to start, as soon as funds could be raised, 22 stations, in addition to the 4 already maintained by the Committee. It was thought that these stations could care for most of the worst portions of the city.

The Finance Committee started in on a whirlwind campaign. The city was divided into districts, each district being in charge of a leader. Mrs. J. Borden Harriman was chairman of the Committee, and she associated with her a number of prominent and active women as district leaders. Each district leader divided her district into streets, and appointed a street leader for each. These latter approached the residents in their vicinity with a personal letter of appeal. The result was quite startling. The financial campaign did not start

until April, but enough money was obtained to open 11 stations of the 22 on the second of May. By the twentieth of May 10 more were ready to open, and on May 22d a meeting was held at Seward Park, officially opening the campaign. At this meeting Borough President McAneny, the Rt. Rev. David H. Greer, the Rt. Rev. Monsigneur McMahon, the Rev. Rabbi Wise, and Commissioner of Health Lederle spoke.

On May 31st the last 2 of the 23 stations were opened; with the original 4

stations, this made 27.

Realizing that the problem was a large one, and that every energy of all who were interested was needed to do the necessary work, a scheme of coöperation was devised to prevent unnecessary duplication of work. The Association of Infants' Milk Stations was organized in May. Its objects were briefly:

- 1. To prevent duplication of work by districting the city, and giving each milk station a distinct area for its activities.
- To establish some uniform system of record keeping, so that the work of the various agencies could be put together at the end of the season.
- 3. To render the information in the possession of one group available to all.
- 4. To conduct a publicity campaign in the interests of the reduction of infant mortality.

At the invitation of Commissioner Lederle a meeting was held, and such an association was formed May 24, 1911. Eight organizations, maintaining 79 milk stations, comprised the Association, and an Executive Committee, composed of a representative of each, was organized, as follows:


S. JOSEPHINE BAKER, M.D., Chief Divisi	on of
Child Hygiene	Department of Health.
REUEL A. BENSON, M.D	Babies' Dairies Assn.
MARIA L. DANIELS, R.N	
ARTHUR R. GREEN, M.D	Nathan Strauss Pasteurized Milk Labora-
	tories.
JANE E. HITCHCOCK, R.N	Nurses' Settlement.
G. N. McCurdy	Morningside Milk Dispensary.
PAUL E. TAYLOR, Director	
ARTHUR E. WAKEMAN	

Through the courtesy of Dr. Baker office room, with use of a stenographer and clerk, was provided at the headquarters of the Division of Child Hygiene. Dr. H. H. Hart, of the Child-Helping Department of the Russell Sage Foundation, provided funds for the salary of a secretary. Mr. Charles J. Storey was appointed to fill this position.

By a committee of the Association the city was districted, thus preventing two nurses from visiting the same family and wasting time and effort. Whenever possible, a baby applying at a station living outside the district was referred to the station located in that district, but, owing to the price of the milk varying in different sets of stations, this was not always possible to bring about. No canvassing was done, however, outside the district assigned to each particular station.

Each station sent in a weekly report in detail showing its enrolment and

### EDUCATION THROUGH PRINTED MATTER





all details of importance, which were tabulated and made accessible to the other organizations.

The 27 stations founded by the Milk Committee were situated in the most congested parts of the city. The object of the Committee's campaign was to make a demonstration of the value of milk stations in reducing infant mortality; to show that these stations should be situated in the most densely populated districts of the city; and that the ordinary store was suited for a milk station.

Many of the situations selected were far from ideal; vacant stores had to be utilized, and they were sometimes not situated just where desired, and, as was afterward proved, were often too small. The Union Settlement in East One Hundred and Fourth St., the Vanderbilt Clinic, the Wilson Industrial School, and the Alfred Corning Clark House gave room rent free for the stations.

The plan of campaign was as follows: Each station was in charge of a graduate nurse specially chosen for her knowledge of infant hygiene, for her interest in the problem, and her willingness to give herself, as well as her time, to the work. She was assisted by a matron. In certain districts where the population was almost entirely foreign, and where English was spoken and understood as rarely as in Italy or Russia, a matron was selected, when possible, who was able to talk the language of the district. Where two or more languages must be spoken, an interpreter was provided in addition.

Each nurse canvassed her district from house to house, looking for babies, leaving the folder of the station with the mother, and offering to help her keep her baby well. That was the great thing the stations tried to teach—to prevent sickness. The windows of each station were adorned with placards in various languages inviting mothers to make use of it. The outside of each station was painted a light, bright blue, and the "Blue Fronts" became a regular expression of the district, being incorporated into many foreign tongues.

The Milk Committee stations tried first and foremost to encourage maternal nursing. Mothers were told how necessary it was and how it would save the baby many of the dangers of the hot weather. Not only were they urged to nurse their babies, but they were taught to care for themselves so that they could nurse them. They were told what to eat and, more important, what not to eat and drink. When breast-milk seemed to be failing, they were provided with milk, and, if necessary, nourishing food, to try to increase the ability to nurse. When nursing was impossible, artificial feeding was ordered by the doctor in attendance at the station. Each baby was treated as an individual, and the food ordered according to its individual needs.

Let us trace a baby through its whole progress at the station. The doctor was in attendance at definite hours twice or three times a week. If a new baby was brought to the clinic, it was seen by the physician, and its food ordered by him. If brought in at another time, after a preliminary talk from the nurse in which the whole system was explained, the mother was asked if she wanted to enroll her baby. If so, it was stripped and weighed by the nurse and the weight recorded on the individual chart. The nurse's record and the history card were then filled out, and the nurse ordered a temporary feeding for the

child, according to instructions very carefully prepared by the supervising physician and indorsed by the Medical Council. If the baby was sick, it was sent to a station where a doctor was in attendance that day, or the station doctor was communicated with and arranged to see the baby.

Its food having been ordered, the mother procured her supply of milk, bottles, barley-flour, etc., and returned to her home, whither the nurse speedily followed her. At this visit she was taught how to prepare the food; and in her own home, where existing conditions could be most readily appreciated, she was given a lesson in general hygiene. The formulæ ordered for the babies were of the simplest kind possible. Whole milk mixtures were used almost exclusively. She was then told to bring the baby back to the station on the next clinic day, and to come every morning for her supply of milk. She was also urged, at the first sign of illness, however slight, to report at once to the station. On clinic days every baby was stripped, weighed, and examined by the station physician, its progress discussed with the mother, any necessary changes ordered, and the nurse instructed what to do. Very sick babies were referred to hospitals or to private physicians.

If the mother was unable to buy milk, through an arrangement with the Charity Organization Society, the Association for Improving the Condition of the Poor, and the United Hebrew Charities, such cases were immediately reported to them, and were at once investigated. In almost every case—probably in every deserving case—milk relief was promptly ordered by these organizations.

Regularity in attendance at clinics was demanded from the mothers. If they refused to attend, they were not allowed to obtain milk.

It was hoped that as many as 75 babies might be enrolled at each station. By the fifteenth of July many of the stations began to cry for help. They were so overcrowded they could not properly carry on their work. So many babies were brought to the clinics that they could not all get in and blocked the sidewalk. The nurses were unable to do the follow-up visiting essential to a successful campaign. The situation was carefully considered, and, although July is not a favorable month to try to raise money, funds were procured, and 4 new stations were opened to relieve those most urgently in need of assistance. The full list was as follows:

OLD STATIONS.....306 E. 21st St.
73 Cannon St.
244 Mulberry St.
438 W. 48th St.

OPENED MAY 2D....248 E. 105th St.
722 E. 9th St.
1391 Avenue A.
229 E. 4th St.
114 Thompson St.
307 Tenth Ave.
35 Norfolk St.
174 Eldridge St.
325 E. 5th St.
239 W. 69th St.
56 Leroy St.

### EDUCATION THROUGH PRINTED MATTER




EXHIBIT 11 PUBLICITY AND INFORMATION



OPENED MAY 20TH . . . . 37 Washington St. 326 E. 11th St. 316 E. 56th St. 512 Second Ave. 873 Second Ave. 1705 Second Ave. 235 E. 81st St. 315 E. 112th St. 3 E. 115th St.

OPENED MAY 31ST ....165 Ludlow St. 200 E. 97th St.

OPENED JULY 8TH....331 Stanton St., to relieve 73 Cannon St. OPENED JULY 10TH....223 E. 75th St., to relieve 1391 Avenue A.

Vanderbilt Clinic.

OPENED JULY 12TH . . . . 78 Ninth Ave., a new station where the summer deaths were high.

OPENED JULY 14TH....1457 Madison Ave., to relieve 200 E. 97th St.

The dates given above are those on which *milk* was first distributed. The nurse and matron had been at work in the district for several days.

Not only was the scope of work increased by adding four new stations, but every station whose enrolment passed the 100 mark was provided with an extra nurse. This was essential if the purpose of the station—instruction—was to be carried out.

In August, therefore, 31 stations were in operation. Doctors were on duty, holding clinics, and often visiting the sick babies in their homes; nurses were daily at the stations and visiting in the homes, watching, advising, encouraging, scolding, teaching, as the case demanded. As the nurse went about from house to house, from family to family, she found many expectant mothers. Part of the campaign was to get in touch with these women and to try to advise and teach them so that they might pass successfully through their pregnancy and be in physical condition to nurse their babies. It was soon found that this was a large undertaking, and in August a special nurse was delegated to take charge of this work. Her salary was kindly provided by Dr. Hastings H. Hart, of the Russell Sage Foundation. During the summer, 964 expectant mothers were under observation.

The coöperation existing between the Milk Committee and the various charitable organizations, especially those already mentioned, was of the closest kind. Whenever tired-out mothers were found whose babies were doing well enough to be sent away, through the cordial relations existing an outing was easily arranged.

The Association for Improving the Condition of the Poor offered to receive any baby needing hospital treatment at Junior Sea Breeze Hospital. This offer was gladly accepted, and during the summer months 43 babies were admitted there. At discharge or death a report of the case was made to the Committee. Discharged babies were immediately referred back to the milk station from which they came.

Figures showing that a large percentage of deaths among babies occur in foundling institutions, the Committeeoffered the services of its milk stations to the New York Foundling Hospital and the New York Infant Asylum. They explained to those in authority at these institutions the purpose of the stations. The offer was gladly accepted, and during the summer 110 babies were enrolled from these institutions. The fact that so many of the foster mothers lived in the outlying boroughs prevented a greater number from being able to make use of the stations.

The milk which was used during the summer was provided by the New York Dairy Demonstration Company.\* It came from tuberculin-tested herds and was of the highest standard. It was sold for seven cents a quart. The contract for this milk was awarded after bids had been asked for from the chief milk-dealers of the city. A constant watch was kept on the quality of this milk. Bacterial counts were made daily from samples taken at various stations, and control bacterial counts were made frequently by the Research Laboratory of the Health Department, through the courtesy of Dr. W. H. Park. The milk was used raw. These counts were compared with those made by the Health Department of the milk dispensed at their stations and compared favorably. The table showing the bacterial counts will be found at the end of this report.

The central office of the Committee was in charge of the Director, Mr. Paul E. Taylor, who was selected by the committee because of his proved executive and constructive ability and his knowledge of the problem from his active participation in the Philadelphia campaign of 1910. He was associated with the Philadelphia Bureau of Municipal Research, who kindly granted him leave of absence to undertake the work here.

The general supervision of the whole work was in the hands of the Director; a supervising nurse was at the office constantly; station nurses were instructed to report by telephone to headquarters every difficulty or complication which arose; an assistant to the supervising nurse was constantly in the field, visiting and inspecting the stations; and a special assistant was appointed during the summer to keep track of the relief cases.

The work of the physicians was supervised by a volunteer supervising physician, who visited the various stations as often as possible, and especially difficult cases when asked, and was general adviser on all strictly medical work. Much of the supervising work was done by the assistant supervising physician.

The entire scope of the work was submitted to the Medical Council in considerable detail and was indorsed by them.

### WORK OF THE HEALTH DEPARTMENT

The Department of Health conducted a very vigorous campaign. Fifteen stations were maintained by them at the following locations:

MANHATTAN	108 Cherry St.	BROOKLYN185 Bedford Ave.
	31 Roosevelt St.	296 Bushwick Ave.
	209 Stanton St.	128 Dupont St.
	2287 First Ave.	994 Flushing Ave.
	207 Division St.	698 Henry St.
-		651 Manhattan Ave.
BRONX	511 East 149th St.	176 Nassau St.
		129 Osborn St.
		303 Williams Ave.

<sup>\*</sup>This company was organized by the New York Milk Committee to demonstrate that a high-grade milk from healthy herds can be produced by the ordinary farmer with ordinary equipment, and sold at a price usually charged for bottled milk of uncertain quality. (See Fifth Annual Report of New York Milk Committee for details of this experiment.)




EXHIBIT 12 Number of Babies Registered at Milk Stations September, 1911



The same general policy was carried out in the stations of the Health Department as in those of the New York Milk Committee. Individual formulæ were ordered,—almost always whole milk formulæ,—and the food was prepared at home under supervision of the nurse. Careful records were kept and the cases were followed up. The work of the summer corps was coördinated with that of the milk stations. The district nurses and inspectors received lists of sick babies from the stations each day, and visited and cared for the sick children of those unable to pay a physician. The department coöperated with the other organizations, sending their inspectors to see any sick child in the care of a milk station on request.

The "Little Mothers' League" continued its active and most valuable work in teaching the future mothers of the city how to care for the little brothers and sisters left in their charge.

The efficient coöperation of the Health Department was further shown by the assignment of physicians and nurses to the stations of the Brooklyn Children's Aid Society and the New York Diet Kitchen stations.

The milk used at the stations was the Sheffield Farms Slawson-Decker pasteurized milk, and was sold in quart and pint bottles, unmodified, for seven cents per quart.

### WORK OF OTHER ORGANIZATIONS

The New York Diet Kitchen maintained 8 stations in Manhattan and one in the Bronx, as follows:

MANHATTAN	146 E. 7th St.
	205 E. 66th St.
	451 E. 86th St.
	209 E. 103d St.
	169 Mott St.
	437 W. 41st St.
	205 W. 62d St.
	152 W. 100th St.
Bronx	583 Courtlandt Ave.

Daily clinics were held at their depots in charge of the Health Department physicians and nurses. Formulæ were ordered according to each baby's individual needs, and mothers were taught by the nurses to prepare the food at their homes.

Certified milk in bulk was dispensed, the daily quantity being "dipped." It was sold at six cents a quart. The same records and system were employed as at the Milk Committee and Health Department stations.

The Nurses' Settlement station was continued as before. Two physicians were in charge of the consultations, and the settlement nurses did the home visiting. Milk was tuberculin tested and shipped to the settlement in cans, where it was bottled and sold in quart and pint bottles at eight cents a quart. Modification was carried out at the homes according to individual prescription under the supervision of visiting nurses.

All the above-mentioned organizations worked along similar lines, laying

special emphasis on instruction, in the belief that permanent results are more likely to follow teaching a mother to care for her baby and its food herself, than by simply giving her the food already prepared. The campaign was also participated in by other organizations, who did their share along slightly different lines.

The Nathan Strauss Pasteurized Milk laboratories maintained eight stations:

322 E. 59th St. 303 E. 111th St. 38 Macdougal St. 45 Monroe St. 348 E. 32d St. 402 W. 37th St. Tompkins Square Park Educational Alliance Roof Garden, 197 East Broadway

From these stations pasteurized milk, already modified to set formulæ, was dispensed in 3-, 6-, and 8-ounce individual feeding-bottles, at ten cents a quart modified, or for older children at eight cents a quart unmodified. There was no home instruction or visiting and no compulsory medical supervision.

The Babies' Dairies provided tuberculin-tested milk in individual feeding-bottles, modified at the station on individual prescription, at ten cents for a day's feeding. Home supervision was not carried out. Medical supervision at the station was provided. The stations were located at—

416 E. 65th St. 523 E. 78th St. 511 W. 41st St. 117 W. 63d St. (St. Cyprian's, colored)

The Good Samaritan Dispensary, at the corner of Broome and Essex Streets, continued its work, dispensing pasteurized milk modified to set formulæ. The Morningside Milk Dispensary, at Morningside Avenue and 122d Street, dispensed pasteurized milk, modified to set formulæ, in individual bottles of 4, 6, and 8 ounces, at two and three cents the bottle.

In Brooklyn the Children's Aid Society maintained 14 stations. Through the active coöperation of the Health Department in providing the physicians and nurses, follow-up work at the home was carried on, and consultations were held. Considerable attention was paid to the educational side. Pasteurized milk, modified to set formulæ, was sold in 3-, 6-, and 8-ounce bottles, at one and two cents a bottle; unmodified milk in 8-ounce bottles at two cents. The stations were located as follows:

105 Fleet St. 608 Fourth Ave. 159 Wyona St. 817 Park Ave. 106 S. 3d St. 27 Columbia Place 146 Union St. 190 Fourth Ave. Pitkin, cor. of Watkins St. 148 Jackson Ave. 165 Johnson Ave. 85 Java St. 201 Hoyt St. 15 Garnet St.

These stations were open from June 15th to September 15th.

The various dispensaries and hospitals continued their follow-up work through visiting nurses. Bellevue Hospital, Presbyterian, Roosevelt, St. Luke's, Mt. Sinai, The Babies', University and Bellevue Medical School Dispensary,





EXHIBIT 13
MILK—ITS VALUE, USE, AND CARE





Ехнізіт 14

THE COMMITTEE PROVIDED THE BOARD OF ESTIMATE AND APPORTIONMENT WITH THE FACTS—
THE NUMBER OF MUNICIPAL STATIONS WAS INCREASED FROM FIFTEEN TO FIFTY-FIVE



Vanderbilt Clinic, and many others have visiting nurses who instruct mothers in the care of their babies, and all are factors in the campaign. There was, however, no particular change from previous years in their methods. The same applies to the St. John's Guild and Junior Sea Breeze, except as already noted in regard to coöperation with the milk stations.

Coöperation was earnestly sought from all the special agencies. The New York Milk Committee sent out an information card, giving the list of all milk stations and information regarding kind of milk, method of preparation, cost, etc. This card was sent to hospitals, dispensaries, social workers, nurses' settlements, and many physicians.

Police Commissioner Waldo distributed lists of the milk stations, provided by the Milk Committee in size suitable for carrying in the pocket, to all patrolmen of the police force, and issued instructions to give the address of the nearest station to any mother with a sick baby seen in the parks or on the recreation piers. The coöperation of the press was active. Articles appeared frequently in many newspapers in New York and throughout the country. Editorial help was also frequently given.

Toward the latter part of the summer the committee prepared to back up the Commissioner of Health in his request for funds to maintain 75 milk stations in New York in 1912. All summer special statistics had been compiled through the courtesy of Dr. Guilfoy, Registrar of the Health Department, who allowed access to the records. These statistics were put together in as striking a manner as possible, with charts and tables. A detailed study of the location of the high mortality centers was made by a member of the Executive Committee, and the results of this study were embodied in a report which was delivered to each member of the Board of Estimate and Apportionment before the hearings.

The Committee was represented at the public hearings; only one protesting voice was raised there against the appropriation, and in this case it was evident that the protester did not understand what the money was to be used for. The Board finally appropriated money to maintain 55 stations, which, although 16 less than the number urged by the Milk Committee, was a very great advance over the 15 provided for the previous year; and there is hope that if these are efficiently carried on in 1912, the full number will be provided for in 1913.

### CHAPTER IV

### Results of the Campaign in New York City

In the pages that follow an honest attempt is made to discuss fairly the facts shown by vital statistics. As already mentioned, the factors in infant mortality are many and complicated, and at the present moment many statistics are still unavailable. Only the most obvious facts can be more than briefly touched upon.

In studying the mortality among infants in this country we are sadly handicapped by lax registration ordinances—lax themselves or laxly enforced. The recognized method of rating yearly infant mortality is to record the number of deaths in infants under one year of age per 1000 births during the same year. In New York the registration of births has been incomplete up to the last few years; therefore comparisons of infant mortality rates (deaths per 1000 births) cannot be made between widely separated periods. The years 1901 and 1911, for instance, cannot be compared. The greater the number of unreported births, the higher will be the rate, of course. A glance at Table 6 will show that in 1901 the rate was 191.5, while in 1911 it was 111.6. Nobody believes this really happened. So inaccurate is this method that the Census Bureau has only in 1910 begun to record these figures, and then (Bulletin No. 109) only for 35 large cities. Only 5 of the 10 largest cities in the country have sufficiently complete registration of births to be included. Chicago admits that not more than one-third of the births are registered, but steps are being taken to remedy this state of affairs.

Another method of rating infant mortality is on the estimated population under one year, based on the percentage at that age according to the last United States Census, and assuming this percentage to remain constant. The results are, of course, only approximate.

Again, the number of deaths under one year of age in proportion to all deaths is an approximate method of estimate.

Tables 6, 7, 8, and 9 show in some detail the infant mortality by months and groups of months, in the city of New York as a whole, the boroughs of Manhattan and Brooklyn, and the "Rest of the City" (comprising Bronx, Queens, and Richmond), for the years 1901 to 1911 inclusive, with five- and ten-year averages. It also shows the relation to total mortality and to births.

It will be noted that there is a steady tendency to increase from 1901 through 1907, with the exception of 1903. The year 1903 shows the lowest gross total mortality of any year in the decade 1901–10.

Again it will be noted that in 1908 there was a sharp fall of 1206 deaths in New York city from the record of 1907; a further decrease of 255 in 1909; a rise of 239 in 1910; followed in 1911 by a fall of 1198. In other words, in 1908 there was a sharp fall, a small change in 1909 and 1910, followed by another sharp fall in 1911.

# TABLE 6.—MORTALITY UNDER ONE YEAR OF AGE, 1901-1911

### CITY OF NEW YORK

AVERAGE 1901-10	1,174 1,139	_					2,263 1,530		1,534 1,332	1,227 1,189	925 993	1,651 1,104	16,067 15,017	73,780 75,423	71.7			30.1 30.5	18,509 18,048	31.3 25.4	107,597 134,477	149.3 111.6
AVERAGE 1906-10	1,204	1,150	1,318	1,266	1,191	1,193	2,167	2,102	1,580	1,311	973	1,111	16,609	75,867	2.18	1	9,890	29.7	18,789	31.7	122,281	135.8
1910	1,261	1,081	1,320	1,228	1,160	1,297	2,253	1,840	1,481	1,344	903	1,047	16,215	76,750	21.1	1	4/c'c	34.3	18,785	29.6	129,080	125.6
1909	1,234	1,076	1,322	1,272	1,216	1,102	1,759	1,939	1,456	1,308	1,089	1,208	15,976	74,105	6.12	1	901,6	2.20	17,512	29.4	122,975	129.9
1908	1,180	1,147	1,283	1,259	1,218	1,170	2,317	1,950	1,472	1,232	953	1,050	16,231	73,072	2.52	1	9,139	99.9	18,081	31.7	126,862	127.9
1907	1,307	1,209	1,336	1,285	1,175	1,184	2,292	2,560	1,650	1,315	926	1,168	17,437	79,205	22.0		7000	21.2	20,139	32.2	92,914 111,772 120,720 126,862 122,975 129,080 122,281	144.4
1906	1,221	1,240	1,330	1,290	1,189	1,215	2,218	2,224	1,843	1,364	896	1,086	17,188	76,203	6.55		0,285	36.5	19,430	32.3	111,772	153.7
AVERAGE 1901-05	1,107	1,071	1,186	1,138	1,048	1,209	2,359	1,905	1,489	1,144	877	966	15,526	71,694	21.0	-	9,793	37.0	18,230	31.5	92,914	167.1
1905	1,169	1,092	1,259	1,281	1,126	1,352	2,802	1,963	1,415	1,180	876	1,007	16,522	73,714	22.4		0,180	37.4	160,61	32.3	555 103,881	159.0
1904	1,133	1,138	-	1,222	1,095	1,267	2,552	1,861	1,542	1,077	698	1,048	16,125	78,060	20.0	-	900	36.9	18,435	32.3	99,555	161.9
1903	1 038	1,051	1,122	972	1,021	1,188	1,920	1,711	1,391	1,148	928	923	14,413	67,864	21.2		_	34.8	16,637	30.1	94,755	152.1
1902	1,183	1,104	1,110	1,113	1,057	1,204	2,117	1,837	1,380	1,145	863	066	15,103	68,112	1.52			99.9	17,353	30.7	85,644	176.3
1901	1,016	971	1,121	1,105	941	1,035	2,404	2,154	1,718	1,170	849	983	15,467	70,720	21.5	10	0,270	40.9	19,635	31.9	80,735	191.5
TOTAL DEATHS UNDER ONE YEAR	January	February	March	April	May	June	July	August	September	October	November	December	Year	All ages 70,720 68,112 67,864 78,060	Per cent, under one year	July-August-September:	Under one year	Per cent. total year	All ages	Per cent. under one year	Births for year 80,735 85,644 94,755 99,	Infant deaths per 1000 births 191.5 176.3 152.1

TABLE 7.—MORTALITY UNDER ONE YEAR OF AGE, 1901-1911

## BOROUGH OF MANHATTAN

1161							764						8,192		21.3		9 2,294				3 66,527	_
AVERAGE 1901-10	685	652	747	737	999	829	1,119	1,072	847	069	518	594	8,989		23.1		3,039	33.8	9,48	32.(	60,336	_
AVERAGE 1906-10	705	670	780	773	969	674	1,043	1,118	857	729	535	612	9,196	39.099	23.5		3,019	32.8	9,417	32.0	65,121	141.2
1910	989	639	756	741	651	723	1,126	1,004	822	720	481	605	8,954	38 660	23.1		2,952	32.9	9,376	31.4	66,357	134.9
1909	689	647	794	811	739	630	810	1,071	759	733	597	634	8,914	37 963	23.4		2,640	29.6	8,519	30.9	63,599	140.1
1908	692	621	768	771	756	644	1,185	1,059	199	677	529	547	9,048	37,723	23.9		3,043	33.6	9,181	33.1	66,875	135.3
1907	763	694	780	755	663	738	1,069	1,313	968	740	540	651	9,602	41.310	23.2		3,278	34.1	10,120	32.4	65,771	145.9
1906	669	751	803	787	675	635	1,025	1,144	1,013	778	529	625	9,464	39.831	23.7		3,182	33.6	068'6	32.1	63,005	150.2
AVERAGE 1901-05	665	634	715	701	636	642	1,196	1,027	837	651	501	576	8,782	38.857	22.6		3,060	34.8	9,556	32.0	55,551	158.0
1905	711	658	757	800	707	711	1,443	1,037	814	658	523	573	9,401	39.671	23.6		3,294	35.0	9,945	33.3	60,203	126.1
1904	654	189	852	751	629	672	1,290	1,014	880	629	490	615	9,207	42.326	21.7		3,184	34.5	199'6	32.9	59,196	155.5
1903	644	614	662	611	640	637	066	919	802	643	498	521	8,181	37.013	22.1 23.3 22.1 21.7		.,	-	~	30.7	49,990 52,291 56,078 59,196	145.8
1902	739	657	662	169	622	588	-	666	807	889	494	577	8,529 8,594	36.769	23.3		.,	33.6	9,040	31.9	52,291	164.3
1901	580	564	645	639	535	605	1,175	1,167	883	641	501	594	8,529	38.507	22.1		3,225	37.8	10,315	32.2	49,990	170.6
TOTAL DEATHS UNDER ONE YEAR	January	February	March	April	May	June	July	August	September	October	November	December	Year	All ages.	under one year	July-August-September:	Under one year	Per cent. total year		Per cent. under one year	Births for year	Infant deaths per 1000 births 170.6 164.3

# TABLE 8.—MORTALITY UNDER ONE YEAR OF AGE, 1901-1911

### BOROUGH OF BROOKLYN

1161	321	352	362	372	300	328	506	545	448	415	326	354	4,629	24,481	10.0		1,499	32.3	5,795	25.8	45,699	101.3
AVERAGE 1901-10	361	334	366	341	333	414	818	929	491	383	293	331	5,097	24,161	0.17		1,937	38.0	6,182	31.3	33,209	153.5
AVERAGE 1906-10	389	340	381	357	357	379	784	640	496	401	307	354	5,190	25,009	7.07		1,920	36.9	6,250	30.7	39,855	130.2
1910	414	313	402	365	363	426	782	543	443	430	278	300	5,059	25,676	1.0.1		1,768	34.9	6,197	28.5	42,708	118.4
1909	384	300	367	328	335	341	682	562	489	385	324	426	4,923	24,365	*0.0		1,733	35.2	5,973	29.0	41,494	118.6
1908	355	366	360	355	330	379	694	592	474	371	315	346	5,012	23,938	6.0%		1,835	36.6	5,922	30.9	41,906	119.6
1907	394	365	400	383	374	325	875	804	512	401	304	366	5,503	26,043	W.1.1		2,191	39.8	6,789	32.2	38,632	142.4
1906	400	360	379	358	386	425	812	200	562	422	316	333	5,453	25,024	,		2,074	38.0	6,373	32.5	34,538	157.8
AVERAGE 1901-05	333	329	350	326	308	449	852	613	487	365	279	309	5,004	23,314	4.1.7		1,953	39.0	6,115	31.9	26,562	188.4
1905	338	317	370	346	282	521	944	635	425	387	257	328	5,150	23,935	61.0		2,004	38.9	6,328	31.6	30,972	166.2
1904	342	340	341	350	326	451	915	575	472	311	273	319	5,015	24,831	1.0%		1,962	39.1	6,145	31.9	28,859	173.7
1903	290	329	331	260	291	441	662	543	540	384	325	295	4,601	23,271 22,344 22,192 2			1,655	35.9	5,562	29.7	27,292	168.5
1902	364	355	343	323	336	507	808	632	451	348	272	320	5,059	22,344			1,891	37.3	5,845	32.3	23,507	215.2
1901	333	304	368	351	307	327	934	683	640	396	270	286	5,197	23,271	2000		2,257	43.4	6,695	33.7	22,182	243.3
TOTAL DEATHS UNDER ONE YEAR	January	February	March	April	May	June	July	August	September	October	November	December	Year	All ages	cente unidei one year	July-August-September:	Under one year	Per cent. total year	All ages	Per cent, under one year	Births for year 22,182 23,507 27,292	Infant deaths per 1000 births 243.3 215.2

TABLE 9.—MORTALITY UNDER ONE YEAR OF AGE, 1901-1911
REST OF CITY (BRONX, QUEENS, RICHMOND)

January. 103  February. 103  March. 108  April. 115  May. 103  June. 295  August. 295  September 105	107 125 148 113 119								-		-	İ	-
	125 148 113 119 140	104	137	120	114	121	150	133	191	161	145	129	160
	148 113 119 140	108	137	1117	118	138	150	160	129	129	141	129	147
	113	129	128	132	129	148	156	155	191	162	156	143	168
	119	101	121	126	115	146	147	133	133	122	136	125	158
	140	06	06	137	107	128	138	132	142	146	137	122	169
		110	144	120	123	156	120	147	131	148	140	131	134
	323	268	347	415	329	381	348	363	267	345	340	335	260
	289	249	272	291	281	380	443	299	306	293	344	312	327
-	165	139	190	176	173	268	243	199	208	216	226	199	213
	125	121	137	135	130	164	174	184	185	194	180	155	147
	132	105	106	96	103	123	112	109	168	144	131	1117	139
December	46	107	114	106	105	128	151	157	148	142	145	125	174
Year	1,883	1,631	*1,923	1,961	1,827	2,281	2,332	2,171	2,139	2,202	2,221	2,021	2,196
All ages. 8.942	8,999	8,659	10,903	10,108	:	11,148	11,843	11,411	11,777	12,406	11,717	:	12,526
nder one year		18.8	17.6	19.5	:	20.4	19.6	19.0	18.1	17.7	18.9	:	17.5
July-August-September:													
Under one year 794	777	929	800	882	783	1,029	1,034	861	781	854	910	847	800
Per cent. total year 45.6	41.3	40.3	42.1	45.0	42.8	45.1	44.3	39.6	36.5	38.7	41.0	41.8	36.4
All ages 2,622	2,368	2,255	2,629	2,818	2,538	3,167	3,229	2,978	3,030	3,213	3,123	2,830	3,287
H	32.5	29.1	30.7	31.3	30.8	32.4	32.0	28.9	25.7	26.5	29.2	29.9	24.3
			11,500	12,706	10,800	14,229	16,317	18,081	17,882	20,015	17,304	14,052	22,251
Infant deaths per 1000 births 203.3		190.2 143.2	165.4	155.1	168.8	159.6	142.9	120.0	119.6	110.0	128.4	143.9	98.6

\* Note: The Slocum disaster, in which 918 lives were lost, occurred in June, 1904. Deducting these from the total, the number of deaths for the year would be 9985. The per cent. under one year was 19.0.




EXHIBIT 15 Infant Deaths Occurring in New York City by Weeks (1909, 1910, 1911)



As already mentioned, in 1908 there was a very great increase in activity in the campaign in New York. That was the year of the formation of the Division of Child Hygiene in the Health Department; the coöperation of the Health Department with the Children's Aid Society; the effort to bring about general coöperation, as shown by the first conference on the summer care of babies; the opening of seven milk depots by the New York Milk Committee; in other words, a great extension of the campaign and the coming to the front of educational prevention.

The year 1911 was another one of great extension in the work, especially

along educational lines.

It must be remembered that the special efforts were confined largely to the summer months; also it must be remembered that the problem is very different in different boroughs, owing to their different physical characteristics. The following tables illustrate the changes that have occurred since 1908:

TABLE 10.—MORTALITY UNDER ONE YEAR IN NEW YORK CITY OCCURRING DURING THREE SUMMER MONTHS

	1908	1909	1910	1911
Actual	5739	5154	5574	4593
	35.3	32.2	34.3	30.5

It will be noticed, from Table 10, that while the total year's mortality in 1908 was low, the summer mortality was quite high—higher than either 1909 or 1910—35.35 per cent. in 1908, as against 32.26 per cent. in 1909, and 34.37 per cent. in 1910. It was, however, below previous years, except 1903.

In 1911 the summer mortality was low, being only 30.58 per cent. of the year's total. Table 11 shows the difference between these years and 1911.

TABLE 11.—DIFFERENCE BETWEEN MORTALITY UNDER ONE YEAR IN THREE SUMMER MONTHS OF 1911 AND CERTAIN OTHER YEARS

	FROM 1908	FROM 1909	FROM 1910
Actual	—1146	—561	—981
Percentage	—19.9	—10.8	—17.6

The summer mortality in 1911, compared with the three previous years in the different boroughs, as shown in Tables 12 and 13, is worth noticing.

TABLE 12.—MORTALITY UNDER ONE YEAR DURING THREE SUMMER MONTHS IN DIFFERENT SECTIONS OF NEW YORK CITY

		1908	1909	1910	1911
	Actual Percentage of year Actual Percentage of year	3043 33.63 1835 36.6	2640 29.61 1733 35.2	2952 <b>32.96</b> 1768 <b>34.9</b>	2294 28.00 1499 32.3
REST OF CITY		861 <b>37.9</b>	781 <b>36.5</b>	854 <b>38.7</b>	800 <b>34.8</b>

TABLE 13.—DIFFERENCE BETWEEN MORTALITY UNDER ONE YEAR IN THREE SUMMER MONTHS OF 1911 AND CERTAIN OTHER YEARS

	FROM 1908	FROM 1909	FROM 1910
MANHATTAN Actual Per cent Actual Per cent Actual Per cent Per cent Per cent Per cent Per cent Per cent	-24.6 -336 -18.3 -61	-346 -13.1 -234 -13.5 +19 +2.4	658 22.2 269 37.8 54 6.3

It will be seen that while there was a very great actual and percentage reduction in 1911 from the figures in 1908 to 1910 in both Manhattan and Brooklyn, in the Rest of the City this was not so marked. There was an actual increase over 1909 figures.

The proportion of deaths occurring in the third quarter of the year was very low. Table 14 shows the distribution of the mortality by trimesters for the years 1901–1911, with five- and ten-year averages for the boroughs of Manhattan, Brooklyn, and the "Rest of the City."

It will be noted that in Manhattan only 28 per cent. of the year's mortality occurred during the third quarter, while the ten-year average is 33.8 per cent., and the five-year average, 1906–1910, is 32.8 per cent. In no single year was so low a figure reached, and in only one, 1909, did it fall below 30 per cent. This occurred in connection with a very low total mortality for the whole year.

The figure for Brooklyn is 32.4 per cent., as against the ten-year average of 38 per cent. It is also the lowest of any single year by 2.5 per cent.

The "Rest of the City" shows 36.4 per cent., the ten-year average being 41.9 per cent. But in 1909 the proportion was 36.5 per cent. The lower figure and the proportionately greater reduction seem to make the inference fair that the more extensive and active campaign in Manhattan was responsible, in part, at any rate.

The New York Milk Committee opened its stations in 1911 in the latter part of May. The campaign did not really come into full swing until June. So far as the Milk Committee was concerned, it ceased its work on the first day of November, when the Health Department took over 22 of the 31 stations which it had maintained during the summer. In order to estimate the effect of the campaign during these five months, tabulations have been made of the deaths in the various parts of the city for a number of years and the births occurring during corresponding twelve-month periods ending October 1st.

In the summer of 1911 there were in Manhattan 57 milk stations; in Brooklyn, 23; and in the Bronx, 1. These tabulations have been made for Manhattan, Brooklyn, and the "Rest of the City." This last area has been taken because in all three boroughs there was practically no milk station work. The regular summer corps work of the Health Department was carried on as heretofore. Table 15 shows the results.

14	
NEW YORK	
~	
0	
>	
3	
(+)	
=	
4	
~	
GREATER	
=	
4	
(1)	
$\overline{z}$	
1	
0	
1-	
=	
0	
m	
5	
9	
2	
IS IN VARIOUS SECTIONS OF C	
2	
H	
S	
in	
5	
=	
$\simeq$	
$\sim$	
4	
>	
-	
Z	
-	
S	
~	
巴	
STE	
STE	
FESTE	
MESTE	
IMESTE	
RIMESTE	
TRIMESTE	
TRIMESTE	
Y TRIMESTE	
BY TRIMESTE	
BY TRIMESTE	
R BY TRIMESTE	
AR BY TRIMESTE	
EAR BY TRIMESTE	
YEAR BY TRIMESTE	
YEAR BY TRIMESTERS 1	
E YEAR BY TRIMESTE	
VE YEAR BY TRIMESTE	
ONE YEAR BY TRIMESTE	
ONE YEAR BY TRIMESTE	
ONE YEAR BY TRIMESTE	
OR ONE YEAR BY TRIMESTE	
IER ONE YEAR BY TRIMESTE	
DER ONE YEAR BY TRIMESTE	
NDER ONE YEAR BY TRIMESTE	
UNDER ONE YEAR BY TRIMESTE	
UNDER ONE YEAR BY TRIMESTE	
S UNDER ONE YEAR BY TRIMESTE	
HS UNDER ONE YEAR BY TRIMESTE	
THS UNDER ONE YEAR BY TRIMESTE	
VITHS UNDER ONE YEAR BY TRIMESTE	
TATHS UNDER ONE YEAR BY TRIMESTE	
EATHS UNDER ONE YEAR BY TRIMESTE	
DEATHS UNDER ONE YEAR BY TRIMESTE	
-DEATHS UNDER ONE YEAR BY TRIMESTE	
.—DEATHS UNDER ONE YEAR BY TRIMESTE	
4.—DEATHS UNDER ONE YEAR BY TRIMESTE	
14.—DEATHS UNDER ONE YEAR BY TRIMESTE	
3 14.—DEATHS UNDER ONE YEAR BY TRIMESTE	
E 14.—DEATHS UNDER ONE YEAR BY TRIMESTE	
3LE 14.—DEATHS UNDER ONE YEAR BY TRIMESTE	
ABLE 14.—DEATHS UNDER ONE YEAR BY TRIMESTE	
TABLE 14.—DEATHS UNDER ONE YEAR BY TRIMESTE	
TABLE 14.—DEATHS UNDER ONE YEAR BY TRIMESTE	
TABLE 14.—DEATHS UNDER ONE YEAR BY TRIMESTE	
TABLE 14.—DEATHS UNDER ONE YEAR BY TRIMESTE	

2171     2180     2115     2144       24.0     24.5     23.6     23.3       3043     2640     2962     3021       33.6     29.6     33.1     32.8       1753     1964     1806     1877       19.4     22.0     20.1     20.4
2180 24.5 2640 29.6 1964 22.0
3043 33.6 1753 19.4
1932 1931 <b>20.4 20.1</b>
1932 <b>20.4</b>
1729
18.7
200
-

TABLE 15.—MORTALITY UNDER ONE YEAR DURING MONTHS JUNE 1ST TO NOVEMBER 1ST IN VARIOUS SECTIONS OF GREATER NEW YORK

	1908	1909	1910	1911
MANHATTAN:				
	644	630	723	572
June July	1185	810	1126	764
August	1059	1071	1004	859
September	799	759	822	671
October	677	733	720	627
Total	4364	4003	4395	3493
BROOKLYN:				
June. 3	379	341	426	328
July	769	682	782	506
August	592	562	543	545
September	474	489	443	448
October	371	385	430	415
Total	2585	2459	2624	2242
REST OF CITY":				
June	147	131	148	134
July	363	267	345	260
August	299	306	293	327
September	199	208	216	213
October	184	185	194	147
Total	1192	1097	1196	1081

It will be noted in Table 15 that the deaths under one year in 1908 and 1910 in all three areas were nearly the same. In 1909 the total was lower than in the years preceding and following. The year 1911 shows a very great drop in Manhattan and a very distinct drop in Brooklyn, and almost no drop in the Rest of the City compared with 1909. Table 16 shows the differences in actual numbers and in percentages. From the gross figures it appears that the Rest of the City did not do nearly so well as Manhattan and Brooklyn.

TABLE 16.—DIFFERENCE BETWEEN MORTALITY UNDER ONE YEAR IN MONTHS JUNE 1ST TO NOVEMBER 1ST, 1911, AND CERTAIN OTHER YEARS IN DIFFER-ENT SECTIONS OF GREATER NEW YORK

	FROM 1908	From 1909	FROM 1910
MANHATTAN:			
Actual	-871	-510	902
Percentage	-19.9	-12.7	-20.5
BROOKLYN:			
Actual	-343	-217	-382
Percentage	-13.2	-8.8	-14.5
"REST OF CITY":	10.2	0.0	11.0
Actual	-111	-16	—115
Percentage	-9.3	-1.4	-9.6

Table 17 shows the actual and percentage changes in the gross infant mor-

tality for these three areas for each of the years 1909, 1910, and 1911, compared with previous years back to 1908.

#### TABLE 17.—DIFFERENCE IN MORTALITY UNDER ONE YEAR DURING MONTHS JUNE 1ST TO NOVEMBER 1ST BETWEEN CERTAIN YEARS AND YEARS PRE-CEDING

1909 as Compared With 1908					
	TUAL DIFF	ERENCE		PERCENTAG	E DIFFERENCE
MANHATTAN	361			_	8.27
BROOKLYN					4.87
"REST OF CITY"	— 95	5			7.97
1910 as Compared With 1908 and 1909					
Аст	UAL DIFFI	ERENCE	PERCE	NTAGE DIFFE	RENCE
1908		1909	1908		1909
MANHATTAN+31		+392	+0.71	+	9.79
BROOKLYN+39		+165	+1.50		6.71
"Rest of City" + 4		+ 99	+0.33		9.02
1911 as Compared With 1908, 1909, an	d 1910				
Асти	AL DIFFER	RENCE	PERCE	NTAGE DIFFE	RENCE
1908	1909	1910	1908	1909	1910
MANHATTAN —871	-510	-902	-19.9	-12.7	-20.5
Brooklyn —343	-217	-383	-13.2	- 8.8	-14.5
"REST OF CITY"111	- 16		- 9.3	- 1.4	— 9.6

Table 18 compares the percentage difference in actual deaths under one year during this period for each area in the years 1909, 1910, 1911, and with the year immediately preceding. It will be seen that in Manhattan and Brooklyn the percentage decrease in deaths during these months in 1911 was greater than in either 1910 or 1909, while for the rest of the city the change is very much less.

## TABLE 18.—PERCENTAGE DIFFERENCE BETWEEN THE MORTALITY UNDER ONE YEAR DURING MONTHS JUNE 1ST TO NOVEMBER 1ST IN 1909, 1910, AND 1911, AND THE YEAR IMMEDIATELY PRECEDING

1909	1910	1911
MANHATTAN8.27%	+9.79%	-20.5%
BROOKLYN	+6.71	-14.5
"REST OF CITY" -7.97	+9.02	<b>—</b> 9.6

These gross figures, as given above, make no allowance for the increase in population. Taking the census figures of 1910 as a basis, the percentage increase in population for 1900–1910 for the various areas is:

Manhattan, ten years,	26.020
Yearly average,	2.602
Brooklyn, ten years4	4.010
Yearly average	00.380
Yearly average	9.038

If we apply this increase to the number of deaths occurring in 1909, the low year of the five preceding 1911, and compare with these figures the actual deaths which did occur in these years during this same period, we find some interesting points, as shown by the following table:

# TABLE 19.—MORTALITY UNDER ONE YEAR, WHICH SHOULD HAVE OCCURRED DURING THE MONTHS JUNE TO NOVEMBER, 1910 AND 1911, ON BASIS OF 1909 FIGURES, ALLOWING FOR INCREASE IN POPULATION

	1910	1911
Manhattan	4107	4211
Brooklyn	2557	2656
"Rest of City"	1196	1295

The actual and percentage differences from these estimated mortalities are:

Act	UAL	Perc	ENTAGES
1910	1911	1910	1911
MANHATTAN+288	-718	+7.0	-17.0
BROOKLYN+ 67	-414	+2.6	-15.5
"REST OF CITY" + 0	-214	+0.0	-16.5

There has been in the "Rest of the City" a steady tendency downward during the last three years. In 1910, when Manhattan and Brooklyn were in excess of the estimated mortality, the "Rest of the City" maintained the same proportion. In 1911 it fell more than Brooklyn, and almost as much as Manhattan.

It is possible that the influence of the campaign in New York city may have had something to do with this result. Conditions in these outlying boroughs are very different from Manhattan and Brooklyn.

If, now, we look at the infant mortality rates for these years, we find that, based on the total births for the year, the proportion of infant deaths in the months of June to November 1st was:

TABLE 20.—INFANT MORTALITY RATE FOR MONTHS JUNE TO NOVEMBER FOR DIFFERENT SECTIONS OF GREATER NEW YORK

1908	1909	1910	1911
Manhattan	62.9	66.2	52.5
Brooklyn	59.2	61.4	49.0
"Rest of City"65.9	61.3	59.7	48.5

Table 21 shows the actual and percentage difference in these rates compared with the years preceding, during the period discussed:

TABLE 21.—DIFFERENCE IN INFANT MORTALITY RATE FOR MONTHS JUNE TO NOVEMBER, BETWEEN 1909, 1910, AND 1911, AND YEARS PRECEDING, FOR DIFFERENT SECTIONS OF GREATER NEW YORK

#### 1909 as Compared With 1908

	ACTUAL	PER CENT.
Manhattan	2.3	-3.52
Brooklyn	2.4	-3.89
"Rest of City"	4.6	-6.98

#### 1910 as Compared With 1908 and 1909

	ACTUAL	PER	CENT.
190	8 1909	1908	1909
MANHATTAN +1.		+1.53	+5.24
Brooklyn	2 +2.2	-0.32	+3.71
"REST OF CITY"6.	2 —1.6	-8 16	-26

#### 1911 as Compared With 1908, 1909, and 1910

		ACTUAL			PER CENT	
	1908	1909	1910	1908	1909	1910
MANHATTAN	-12.7	-10.4	-13.7	-19.4	-16.5	-20.7
BROOKLYN	-12.6	-10.2	-12.4	-20.4	-17.2	-20.1
"REST OF CITY"	-17.4	-12.8	-11.2	-26.4	-20.8	-18.7




EXHIBIT 16
DEATHS OF INFANTS, 1909, 1910, 1911, BY WEEKS
UPPER CHART—MANHATTAN; LOWER CHART—BROOKLYN



Comparing the percentage change in the infant mortality rates in consecutive years we find:

TABLE 22		
1909	1910	1911
Manhattan	+5.2	-20.0
Brooklyn	+3.7	-20.1
"REST OF CITY" -6.9	-2.6	-18.7

The mortality rate of the "Rest of the City" fell in 1909 much more than that of Manhattan and Brooklyn. In 1910 it fell, while that of the other two boroughs rose. In 1911, in Manhattan and Brooklyn, the mortality fell more than in the "Rest of the City."

The tables given above would seem to show that-

- (1) In the city of New York during the summer and early fall of 1911 there was a very great diminution in infant mortality.
- (2) That the diminution is demonstrable both in the gross infant mortality and in the infant mortality rate.
- (3) That the diminution is slightly but distinctly more marked in the boroughs of Manhattan and Brooklyn, where the greatest efforts were put forth, than in the "Rest of the City."

It is a well-known fact that infant mortality tends to vary from year to year in given localities. This is usually explained on the ground of varying weather conditions. Summers of intense heat and scant rainfall have usually been followed by a high infant mortality. It remains now to investigate weather conditions which have existed in New York during certain years, and to see how they compare with the mortality rates.

Table 23 shows the number of days on which the maximum temperature and mean temperature reached or passed certain figures.

TABLE 23.\*—NUMBER OF DAYS MAXIMUM AND MEAN TEMPERATURES IN NEW YORK REACHED CERTAIN FIGURES

TIL W	OICH RELIE	TILD CL	CIMIN I IOC	ILLO	
	MAXIMU	M TEMPERATU	JRE WAS	MEAN TEMPE	RATURE WAS
	95° or +	90° or +	85° or +	85° or +	80° or +
June	TOTAL STREET	100000000000000000000000000000000000000			
1901	2	5	10	1	5
1903		0	0	0	0
1908		1	8	0	4
1909		4	9	0	6
1910	0	2	6	0	3
1911		0	2	0	0
July					
1901	2	9	19	2	12
1903		6	12	0	5
1908	2	7	21	2	18
1909	0	2	10	0	3
1910	0	4	19	0	10
1911		10	18	6	9
August					
1901	0	1	7	0	2
1903		0	3	0	0
1908		4	10	3	6
1909	Δ :	4	7	0	2
1910		0	1	0	0
1911		5	12	0	3

<sup>\*</sup> New York Meteorological Observatory Reports.

Maxi	MUM TEMPERATUR	RE WAS	MEAN TEMPE	RATURE WAS
95° or +	- 90° or +	85° or +	85° or +	80° or +
September				
1901	0	4	0	0
19030	0	3	0	0
1908	0	0	0	0
1909	0	0	0	0
1910	1	2	0	1
19110	0	3	0	0
June-September				
19014	15	40	3	19
19031	6	18	0	5
19082	12	39	5	28
1909	10	26	0	11
1910	7	28	0	14
19116	15	35	6	12

The years 1901 and 1903 are included in the above table, because the latter showed the lowest year's mortality on record since 1901. The annual meteorologic summary for 1911, issued by the Weather Bureau (New York), contains the following statement: "July, period—— hot weather exceeded only by that of July, 1901, in duration and intensity, prevailed during the first thirteen days of the month." The highest temperature was only one degree below the highest ever recorded at the station, and for thirteen days the average temperature was 81°. These records were taken 400 feet above the street.

Not only was the number of days when the temperature reached the high point greater in 1911 than in these other years, but the duration of the heat was greater, as will be seen in Table 24.

TABLE 24.\*—NUMBER OF HOURS TEMPERATURE WAS AT OR ABOVE CERTAIN FIGURES

FIGUR	CES		
		TEMPERATURE WAS	
	95° or +	90° or +	85° or +
June	,, ,,	, o or 1	00 01 1
1901	3	23	"
4003			66
1903		0	0
1908	0	5	38
1909	0	14	65
1910		5	34
1911		ő	10
		· ·	10
July			
1901	10	40	134
1903	1	26	80
1908		33	149
1909		6	41
1010			
1011		20	100
1911	25	67	151
August			
1901	0	3	35
1903			
		0	11
4000		24	76
	0	10	34
1910	0	0	5
1911	0	15	59
September		-	
1001			
****		0	12
		0	11
1908	0	0	0
1909	0	0	0
1910	0	5	14
1911		0	
	0	U	8

<sup>\*</sup> New York Meteorological Observatory Reports.

		TEMPERATURE WAS	
June-September	95° or +	90° or +	85° or +
1901	13	86	247
1903	1	26	102
1908	3	62	263
1909	0	30	140
1910	0	30	153
1911	25	82	228

This table shows a very much greater duration of heat than in any other of the years tabulated, especially in regard to temperatures of 90° or over. In 1908 a temperature of 85° or over was maintained longer than in 1911, but in 1909 and 1910 weather conditions were much less severe. The years 1901 and 1903 will be discussed later. The mean temperature for the month taken alone is deceptive in making comparisons, a long period of hot weather being often balanced by a long cool period, or short hot and cool periods following one another. The following table (Table 25), taken in conjunction with Table 24, shows the records for the four months of June, July, August, and September, and the average monthly mean over a period of forty-two years:

TABLE 25.—MEAN MONTHLY TEMPERATURES AND INFANT MORTALITY RATES, JUNE-SEPTEMBER\*

Average Mean Temperature for Month for Forty-two Years	1901	1903	1908	1909	1910	1911
June       70.3         July       75.3         August       73.3         September       67.1	70.5	64.2	72.9	72.1	68.7	69.8
	77.7	74.6	79.8	73.5	77.2	77.5
	74.9	69.0	75.7	72.2	72.0	72.8
	68.1	67.0	68.3	66.8	69.4	67.6
Infant mortality	7311	6210	6909	6256	6871	5627
	(90.5)	(65.5)	54.4	50.8	53.2	41.8

Table 25 shows that the mean monthly temperature in 1908 for all four months was distinctly above the average for the forty-two years. In this year the summer mortality was high, being 54.4. In 1909 June alone was above the average mean, July and August being considerably below, and September slightly below; the infant mortality was low, the rate being 50.8. The records for 1910 and 1911 were very similar. In June both were below the average mean, especially 1910. July was nearly two degrees above the mean in both years; August was below in both, 1911 being the higher of the two. September shows considerable excess in 1910 and a slight excess in 1911. In the death-rates for the two years there is a difference of 1.4 in favor of 1911. So far as temperature was concerned, it would seem that 1911 was a bad year.

The year 1903 was the remarkable one of the decade as to low infant mortality, both for the year and for the four months June, July, August, and September. Its death-rate, based upon births for the entire year and for the summer months, was also much below the years preceding and following; in fact, the yearly rate was not equaled again until 1907, when there was a very great increase in the reported births. In 1903 there can be no question as to the remarkable weather

conditions that prevailed. June was over 6° below the mean; July, 0.7° below; August, over 4° below; and September, about the average. On only one day of the four months did the temperature reach 95° or over, and it remained there for only one hour. For only nine hours did it reach 90° or over, and for only 102 hours was it 85°+. The mean temperature never reached 85° and was only 80° or over on five days. There has been no such record as this in any year since 1900, the nearest approach being 1909, as already mentioned.

The years 1901 and 1911 each had a prolonged hot spell in July. In the former year, at 6 A.M. on July 1st, the temperature was 81° and did not fall below 80° until 2 P.M. on the third, a period of fifty-six hours, when between 1 P.M. and 6 P.M. there was a hard storm, the thermometer falling from 94° to 77° in one hour. From 11 A.M. to 9 P.M. on the first, from 9 A.M. to 6 P.M. on the second, and from 10 A.M. to 1 P.M. on the third, the temperature registered 90 or over and reached 100° on the first and second. Then followed cooler weather. There was a period of high temperature from July 14th to 17th inclusive; August 21st to 24th inclusive, and August 29th and 30th, but the temperature did not reach such great heights, nor did it remain so high during the entire twenty-four hours. During the month of July, 1901, there were 2404 deaths under one year of age. In the week ending July 6, 1901, 617.

In July, 1911, during the first thirteen days, there was a period of extreme heat. From 10 A.M. on the morning of the second to 7 P.M. on the evening of the seventh, a total of one hundred and twenty-nine hours, the temperature was recorded below 80° for only twenty-six hours. From 3 P.M. to 9 P.M. on the second, 9 A.M. to 9 P.M. on the third, 11 A.M. to 8 P.M. on the fourth, 12 M. to 9 P.M. on the fifth, 11 A.M. to 8 P.M. on the sixth, it was continually at 90° or over; then came thirty-nine hours of relief, to be followed by another period of extreme heat. From 10 A.M. on the ninth, to 10 P.M. on the thirteenth, a total of one hundred and eight hours, the temperature was below 80° only twenty-three hours. From 3 P.M. to 8 P.M. on the ninth, 11 A.M. to 5 P.M. on the tenth, 12 M. to 9 P.M. on the eleventh, 4 P.M. to 9 P.M. on the twelfth, it was continually 90° or over, and yet during the month of July, 1911, only 1530 babies died, compared with 2404 in 1901. In the week ending July 8, 1911, only 347 babies died, compared with 617 in 1901. This remarkable failure of the infant deaths to rise promptly and markedly during the hot weather is even more strikingly shown in Exhibits 6 and 8, which show the daily deaths in Manhattan and Brooklyn during this hot period. On no day in the entire summer did the deaths in Manhattan reach 40. In 1909, on the other hand, 57, and in 1910, 65, deaths were recorded in a single day. The hot period in 1901, while it lasted a shorter time, was a little more intense in that the temperature did not fall below 80° for fifty-six hours. In 1911, in the early mornings, between 2 and 7 or 8 A.M., the thermometer registered 77° to 80°, but its effect on the thoroughly baked tenements could not have been great.

When we come to consider the subject of rainfall and the number of rainy days occurring during these years, several points must be emphasized. The total rainfall for the month, so far as its bearing upon health conditions may be concerned, must be considered as to its distribution throughout the month, as well as its total amount. For instance, August, 1911, had a total rainfall of 9.28 inches, as against 1.07 in 1910, but up to and including the twenty-third only 1.95 inches had fallen in 1911, as against 0.89 in 1910.

Table 26 shows the monthly rainfall and the number of rainy days in each month for the years we have been considering, and also the rainfall from July 1st to August 23d inclusive.

TABLE 26.—RAINFALL AND RAINY DAYS, NEW YORK CITY RAINFALL, INCHES

	KAINTA	LL, INCHE	n.)			
Average, Forty-two Years	1901	1903	1908	1909	1910	1911
JUNE	1.00 7.64	9.78 3.93	1.27 3.80	3.02 2.17	6.03 0.49	5.67 2.16
AUGUST	6.55 2.42	7.85 4.31	6.01 1.91	8.50 2.65	1.07	9.28 1.70
TOTAL	17.61	25.87	12.99	16.34	9.01	18.81
N	UMBER O	F RAINY I	DAYS			
JUNE       9         JULY       10         AUGUST       9         SEPTEMBER       8	5 16 9 12	18 10 11 6	3 7 10 3	10 5 8 7	12 6 11 5	14 11 12 9
Тотац	42	45	23	30	34	46
RAINFALL,*	JULY 1ST	TO AUGUS	ST 23D, IN	CHES		
JULYAUGUST	7.64 4.05	3.93 3.66	3.80 2.59	2.17 8.50	0.49 0.89	2.16 1.95
TOTAL	11.69	7.59	6.39	10.67	1.38	4.11

It will be seen that during the entire period, 1911 had a greater precipitation than any of the three previous years, slightly greater than in 1901, and less than in 1903. The number of rainy days was greater than in any of the five other years considered, although during the period July 1st to August 23d the rainfall was lower than in any of the five years except 1910. We must, therefore, admit that in 1911 the rainfall and number of rainy days were in favor of low mortality, as compared with 1910, during the latter part of the summer. During July and the first part of August climatic conditions were unfavorable.

In a general way summers having bad climatic conditions had a high infant mortality. During 1911 these conditions were unfavorable, especially during July. This Committee believes that the reduction in the mortality in the summer of 1911 was not due to "chance" variation in mortality, but to increased efficiency in the methods used to combat it.

<sup>\*</sup> New York Meteorological Observatory figures.

#### CHAPTER V

## The Results at the Milk Stations of the New York Milk Committee


The results of the work of the Milk Stations may be gauged to a certain extent by the response of the communities in which the stations were located; by the mortality among the babies within their spheres of influence; and by the mortality among the station babies.

In order to obtain these figures a vast amount of statistical work had to be done, and it was impossible to carry it out for all the stations in Greater New York. The work was, therefore, limited to the stations maintained by the Milk Committee.


It has already been stated that the response to the work was so great at most of the stations that extra nurses and sometimes extra matrons had to be provided. In 20 stations an extra nurse was provided, and in 15 an extra assistant. Table 27 shows the enrolment of babies actually under supervision at the end of each week from May 20th to the end of the season.

In order to estimate the mortality among the babies in the areas under the influence of the milk stations, the following method was adopted: A map was made and plotted out showing the location of the station and the number of babies enrolled in each block surrounding the station. In this way the actual sphere of influence of the station was determined. A few babies came from longer distances, and, therefore, from outside the district. In order to determine the mortality in the district thus established a search of the records at the Health Department and a tabulation of all deaths occurring in the months of June to November were made for the years 1910-1911. In order to determine the infant mortality rates it was necessary to tabulate the births by blocks in the same areas. The Committee originally intended to close its work on October 1st, so that the births were tabulated for the years October 1, 1909, to September 30, 1910; and October 1, 1910, to September 30, 1911. The deaths were originally tabulated from January 1st to September 30th, but it seemed more accurate to take only the five months of active work, and this tabulation was made a second time. It was, however, impossible to go through the birth figures again.

The following table shows the deaths under one year for the "Station Areas" in the months June to October inclusive in 1910, before, and in 1911, after, stations were maintained. It should be remembered that the stations at Mulberry, Cannon, East 21st, and West 48th Streets had been running for several years. This table also shows the births for the twelve months ending September 30th, and the infant mortality rate based thereon. For comparison the death-rate



WHERE MILK STATIONS ARE NEEDED. STREET SCENE ON LOWER EAST SIDE, NEW YORK



-	
$\sim$	
0)	
100	
T	
1-3	
-	
(BY WEE)	
>	
037	
N	
20	
=	
The state of	
1	
[7]	
_	
2	
_	
_	
-	
2	
~	
0	
(1)	
-	
4.0	
20	
-	
-	
_	
-	
6	
14	
-	
N	
-	
0	
~	
~	
377	
-	
5	
5	
-	
4	
1-1	
=	
C D	
_	
=	
1-	
-	
0	
01	
SZ	
SNO	
ONS	
IONS	
LIONS	
TIONS	
ATIONS	
FATIONS	
TATIONS	
STATIONS	
STATIONS	
K STATIONS	
K STATIONS	
LK STATIONS	
ILK STATIONS	
IILK STATIONS	
MILK STATIONS	
MILK STATIONS	
MILK STATIONS	
T MILK STATIONS	
AT MILK STATIONS	
AT MILK STATIONS	
AT MILK STATIONS	
T AT MILK STATIONS	
IT AT MILK STATIONS	
NT AT MILK STATIONS OF THE NEW YORK MILK COMMITTEE	
ENT AT MILK STATIONS	
ENT AT MILK STATIONS	
MENT AT MILK STATIONS	
MENT AT MILK STATIONS	
LMENT AT MILK STATIONS	
LMENT AT MILK STATIONS	
OLMENT AT MILK STATIONS	
ROLMENT AT MILK STATIONS	
ROLMENT AT MILK STATIONS	
NROLMENT AT MILK STATIONS	
NROLMENT AT MILK STATIONS	
ENROLMENT AT MILK STATIONS	
ENROLMENT AT MILK STATIONS	
-ENROLMENT AT MILK STATIONS	
-ENROLMENT AT MILK STATIONS	
1.—ENROLMENT AT MILK STATIONS	
7.—ENROLMEI	
TABLE 27.—ENROLMENT AT MILK STATIONS	
7.—ENROLMEI	

Oct. 31	123   123   124   125	4937
OCT.	198 1125 1125 1125 125 125 127 120 120 120 120 120 120 120 120 120 120	5070
Ост.	195 110 110 110 110 110 1111 1111 1111 1	5019
Oct. 14	196 106 106 107 107 107 107 107 107 108 108 108 108 108 108 108 108 108 108	5016
Oct.	188 107 1154 1138 1138 1138 1138 1138 1138 1138 113	4971
SEPT.	193 1111 153 1111 1121 1131 1132 1145 1145 1145 1145 1145 1145 1145 114	4923
SEPT.	194 143 160 160 160 160 160 160 160 160 160 160	4903
SEPT.	188 1139 1141 1142 1150 1160 1173 1173 1173 1173 1173 1173 1173 117	4796
SEPT.	184 1118 1123 1134 1140 1140 1140 1140 1140 1140 114	4684
SEPT.	185 195 195 195 195 196 197 198 198 198 198 198 198 198 198 198 198	4457
Aug. 26	175 175 175 175 175 175 175 175 175 175	4320
Aug.	166 168 168 105 105 107 107 107 108 108 108 108 108 108 108 108 108 108	4016
Aug.	151 177 177 177 177 178 188 188 188 188 18	3674
Aug.	150 177 178 178 178 178 178 178 178 178 178	3361
July 29	156 175 175 175 175 175 175 175 175 175 175	2395
July 22	113 113 113 114 115 115 115 115 115 115 115 115 115	2816
July 15	441 156 156 156 157 167 167 168 177 177 177 177 177 177 177 177 177 17	2483
JULY	21 24 1881 1881 1881 1881 1881 1881 1881 1882 1883 1883 1884 1985	2025
July 1	21 04 04 15 16 16 16 16 16 16 16 16 16 16 16 16 16	
JUNE 24	126 159 159 159 159 159 159 159 159 159 159	1477
JUNE 17	110 110 110 110 110 110 110 110 110 110	967 1058 1245 1477 1715
JUNE 10	112 152 153 153 154 171 171 172 173 173 174 175 175 175 175 175 175 175 175 175 175	1058
JUNE 3	100 100 100 100 100 100 100 100 100 100	196
May 27	1529 1529 1539 1539 1539 1539 1539 1539 1539 153	834
MAY 20	105 1135 1135 1135 1135 1135 1135 1135 1	634
STATION NUMBER	EAST  1 2 3 3 4 4 5 6 6 7 7 11 11 12 12 13 14 14 15 16 16 17 18 18 22 23 23 24 25 25 26 66 7 7	TOTAL

among babies under one year, based on the enrolment, is given for each station; also the totals for East and West Side stations and for the total area.

TABLE 28.—STATISTICS OF STATION AREAS AND STATIONS

1910

1911

STATION MORTALITY

STATION AREA EAST SIDE	Deaths, June- Oct.	Births, Oct Sept.	Infant Mortal- ity	Deaths, June- Oct.	Births, Oct Sept.	Infant Mor- tality	Deaths, June- Oct.	Enrol- ment	Death- rate
1. Mulberry	42	1307	32.1	54	1243	43.4	12	256	46.8
2. Norfolk	16	601	26.6	11	580	18.9	0	123	0.0
3. Eldridge	39	1305	29.8	34	1274	26.6	2	207	9.6
4. Cannon and Stanton (21)	78	1909	40.8	52	1777	29.2	6	430	13.9
5. Ludlow	65	1130	57.5	30	1012	29.6	0	155	0.0
6. East 3d	58	1219	47.5	40	1199	33.3	1	153	6.5
7. East 5th	33	882	37.4	29	820	35.3	7	142	49.3
8. East 9th	55	1011	54.4	36	911	39.5	3	158	18.9
9. East 21st	53	598	88.6	34	561	60.6	5	187	26.7
10. Avenue A	53	831	63.7	50	786	63.6	6	205	29.2
11. East 105th	90	1678	53.6	83	1663	49.9	14	299	46.8
12. East 11th	92	1568	58.6	70	1556	44.9	5	283	17.6
13. East 56th	20	358	55.8	14	366	38.2	1	111	9.0
14. 512 2d Ave	47	554	84.8	31	553	56.0	4 9	176 170	22.7 52.9
15. 873 2d Ave	37	568	65.1	39	605	64.4			37.0
16. E. 81st St	30	505	59.4	23	471	48.8	5	135 135	44.4
17. 1705 2d Ave	24 53	378	63.4	21	397 783	52.8 51.0	6		22.4
18. E. 97th		834	63.5	40			6	267	100
19. E. 112th	110	1593	69.0	84	1631	51.5	8	198	40.4 14.9
20. E. 115th	13	349	37.2	10	338	29.5	2	134	14.9
21. Combined with 4	27	979	72.3	14	900	38.2	2	101	19.8
23. Madison Ave	27	373		14	366 776			189	0.0
		738	36.5	25	110	32.2	0	109	0.0
TOTAL EAST SIDE	1062	20289	52.3	824	19668	41.8	104	4214	24.6
		1910			1911		STATIO	ON MORT	ALITY
STATION AREA WEST SIDE	Deaths.	Births,	Infant	Deaths.	Births.	Infant	Deaths.		Infant
11231 0102	June-	Oct	Mortal-	June-	Oct	Mor-	June-	Enrol-	Mor-
	Oct.	Sept.	ity	Oct.	Sept.	tality	Oct.	ment	tality
			7.5	-					
1. Thompson	42	917	45.8	47	861	54.5	4	236	16.9
2. Leroy	40	550	72.7	32	555	57.6	10	129	77.5
3. 10th Ave	54	629	85.8	46	524	87.7	8	117	68.3
4. W. 48th	127	1360	93.3	101	1409	71.7	7	198	35.3
5. W. 69th	32	375	85.3	25	378	66.1	2	91	21.9
6. Washington	33	222	148.5	24	242	99.1	5	121	41.3
7. Vanderbilt Clinic	59	545	108.2	44	481	91.4	2	139	14.3
8. 9th Ave	44	373	117.9	19	365	52.0	3	134	22.3

It will be seen that in the station areas on the East Side there was a reduction of 10.5 over 1910, and on the West Side, 16.4. The infant population affected on the West Side, however, was about one-quarter that of the East Side. This

86.6

52.3

86.6

59.0

338

824

338

1162

4815

19668

4815

24483

70.2

41.8

70.2

47.4

1165

4214

1165

5379

35.1

24.6

35.1

26.9

41

104

41

145

TOTAL WEST SIDE .

ALL AREAS.....

EAST SIDE.....

West Side....

431

1062

1493

431

4971

20289

25260

would seem to point either to less efficient station technic or to a poorer class of patients, *i. e.*, needing the stations more. Comparing the station rate and the district rate, the latter conclusion is borne out, for on the West Side the station rate was just half the district rate. The testimony of the nurses and physicians in charge of the stations also points to this being the case.

Another noticeable fact is that in the Mulberry Street, East 5th Street, and Leroy Street stations the station mortality is in excess of the district mortality. This is explained by the fact that the nurses in these districts were particularly active in seeking out sick babies; also by the particularly bad character of the neighborhood in which at least two of the stations (East 5th Street and Leroy Street) were situated. Although many of these babies were seen only once and cannot be fairly charged against the station, the Committee adopted the policy of accepting the responsibility for the death of every baby enrolled, even if seen only once, in order to avoid criticism, and it stands on these records as given.

The West Side figures show the great need of stations in this part of the city. The areas of intensely congested population are more scattered; there is not the long stretch that is seen on the East Side, and this part of the city has been neglected. In only 4 of the 23 East Side station areas does the death-rate equal or exceed the lowest one on the West Side. In 4 of the West Side districts the rates are higher than those of the worst East Side district.

Some of the figures are very striking. In the Washington Street district, among the Syrian population, the death-rate in 1910 was 148.5, and fell in 1911 to 99.1, a net difference of 49.4. This little strip of densely populated territory extends from Battery Place to Cedar Street, from West Side to Greenwich. The people rarely speak English, and an interpreter was needed almost all the time. Two hundred and forty-two babies were born in the year, and the enrolment was 121, or exactly 50 per cent., while the station death-rate was 41.3, or 41.6 per cent. of the district mortality.

Table 29, while not strictly accurate, is suggestive for purposes of comparison. If the enrolment be deducted from the district births, and the station deaths from the district deaths, the comparison is even more striking.

This is not strictly accurate, because the enrolment was not absolutely confined to the station area of influence. For instance, at the East 105th Street station several enrolled babies lived in the Bronx. The station where the figures did most closely correspond was probably Washington Street, which shows a mortality of 41.3 as against the "corrected" rate of 157 for the district.

Another suggestive table, again not strictly accurate for the same reason, but not more than 1 or 2 per cent. out, is Table 30, showing the percentage of births enrolled at the stations and the percentage of the district deaths among babies enrolled.

One of the objects of the demonstration carried on by the New York Milk Committee in coöperation with other organizations was to show that home modification was practicable among the very poor and ignorant, among whom the work was done. The Committee felt very strongly that not only was this a more practical way, because less expensive, but that it accomplished more in that it taught something.

TABLE 29.—STATISTICS FOR STATION AREAS, DEDUCTING STATION ENROL-MENT FROM DISTRICT BIRTHS, AND STATION DEATHS FROM DISTRICT DEATHS, COMPARED WITH STATION FIGURES

	5	STATION DISTRIC	CTS		STATIONS	
	DEATHS	BIRTHS	RATE	DEATHS	ENROLMENT	RATE
1 E	42	987	42.5	12	256	46.8
2 E	11	457	24.0	0	123	0.0
3 E	32	1067	29.9	2	207	9.6
4 and 21 E	46	1347	34.1	6	430	13.9
5 E	30	857	35.0	l ő	155	0.0
6 E	39	1046	27.2	1	153	6.5
7 E	22	678	32.4		142	49.3
7 E	33			2		
8 E		753	43.8	7 3 5	158	18.9
9 E	29	374	77.5		187	26.7
10 E	44	581	75.7	6	205	29.2
11 E	69	1364	50.5	14	299	46.8
12 E	65	1273	51.0	5	283	17.6
13 E	13	255	50.9	1	111	9.0
14 E	27	377	71.6	4	176	22.7
15 E	30	435	68.9	9 5	170	52.9
16 E	18	336	53.5	5	135	37.0
17 E	15	262	57.2	6	135	44.4
8 E	34	516	65.8	6	267	22.4
9 E	76	1433	53.0	8	198	40.4
20 E	8	204	39.2	2	134	14.9
22 E	12	265	45.2	2 2	101	19.8
	25	587	42.5	0	189	0.0
23 E	23	901	42.3	0	100	0.0
Γotal East Side	720	15454	46.5	104	4214	24.6
1 W	43	625	68.8	4	236	16.9
2 W	22	426	51.6	10	129	77.5
3 W	38	407	93.3	8	117	68.3
4 W	94	1211	77.6	7	198	35.3
5 W	23	287	80.1		91	21.9
6 W	19	121	157.0	5	121	41.3
7 117	42		122.8	2 5 2 3	139	14.3
7 W		342		2		
8 W	16	231	69.2	3	134	22.3
Total West Side	297	3650	81.3	41	1165	35.1
SUMMARY						
EAST SIDE	720	15454	16.5	104	4214	24.6
		15454	46.5	41	1165	35.1
West Side	297	3650	81.3	41	1100	33.1
Готац	1017	19104	53.2	145	5379	26.9

With this point in view, careful records were kept of each case, and from the individual records the following tables (Tables 31 to 42) have been compiled, showing in some detail the feeding history of 3182 babies. The period of observation extended from June 1st to September 15th. The latter date was selected in order that something definite might be compiled for the Board of Estimate and Apportionment. If the full time—June 1st to November 1st—could have been utilized for these records, the results might have been even better, as many babies were recorded as well at the end who showed no gain in weight.

TABLE 30.—RELATION OF ENROLMENT AT STATION TO BIRTHS IN DISTRICT AND STATION DEATHS TO DISTRICT DEATHS

STATION DISTRICTS	PERCENTAGE OF BIRTHS ENROLLED	PERCENTAGE OF DISTRICT DEATHS AMONG ENROLLED
EAST SIDE		
1	20.5	22.2
2	21.2	0.0
3	16.2	5.8
4 and 21	24.2	11.5
5	15.3	0.0
6		2.5
7	17.3	24.1
8	17.3	8.3
9		14.7
10	26.0	12.0
11	17.9	16.8
12	18.1	. 7.1
13	30.3	6.8
14	31.8	12.9
15	28.0	23.0
16	28.6	21.7
17		28.5
18		15.0
19	12.1	9.5
20	39.6	20.0
22	27.5	14.2
23	24.3	0.0
	21.4	12.6
WEST SIDE		
1	27.4	8.5
2		31.2
3		17.3
4		6.9
5	24.0	8.0
6	50.0	20.8
7	28.9	4.5
8	36.7	15.7
	24.2	12.1
GRAND TOTAL	21.9	12.4

With reference to the condition of "well" or "sick" at the close of the period, the criticism may be made that overenthusiasm of the physicians and nurses may have made the well too many, and the sick too few. This criticism cannot be disproved, but even leaving out of all consideration the statement of "well" or "sick," the weights, which show pretty conclusively the success or failure of feeding, cannot be questioned. They must be regarded as accurate and as free from error due to the personal equation.

In considering the following tables, noting the "condition at close," it must be remembered that the period of observation was short,—an average of six weeks,—and many babies classed as well did not have a chance to gain in weight. Every baby enrolled more than a week is included in this record. Fatal cases are included in the tables, although no data are given. These cases are considered in detail elsewhere. As no babies who did not attend a week or over are included in the table, the number of deaths is only 74 instead of 119, as 45 babies were seen only once and immediately referred to a hospital or private physician, or else the mother refused to attend.

TABLE 31.—SHOWING FOR WELL BABIES, BREAST-FED ON ADMISSION AND THROUGHOUT PERIOD OF OBSERVATION, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

	DIED		:	-	-	:	-		: :	:	:	:	:	:	;	:	:	:	:	:	:				:	:	3
	TION	Sick	:	:	:	:	:	-	2	-	:	:	:	-	:	:	:	:	:	:	:				:	:	S
188	CONDITION AT CLOSE	Well	-	4	:	-	-	-	2	4	3	9	-	3	:	-	7	2	2	:	-				: -	1	36
	WK.	Ozs.	10.00	6.20	:	10.00	9.00	38	3.06	7.38	1.90	1.69	10.00	3.79	:	0.21	4.75	3.25	1.00	:	3.00				1.86		3.48
	05	Lbs.	:	:	:	-	:		: :	:	;	:	:	:	:	:	:	:	:	:	:				:	:	:
ЕІСНТ	Av. Loss Case   Pe	Ozs.	14.00	15.50	:	10.00	00.6	1 50	13.00	11.80	6.33	7.33	0.10	13.25	:	3.00	9.50	6.50	1.50		00.6				13.00	200	11.37
N NI	PER (	Lbs.	-	:	:	-	:		: :	:	:	:	:	:	:	:	•		:	:	:				-000	:	:
SHOWING LOSS IN WEIGHT	VT.	Ozs. 1	00.9	3.75	:	8.00	15.00	12 00	10.75	11.00	12.00	00.6	2.00	12.75	:	00.6	8.00	9.50	12.00	:	:		:	:	. 6	3	15.07
OWING	AV. WT.	Lbs.	ın		:		14 1	10 1			17 1	19	20	18 1	:	-	18	20	_	:	19				33	-	16 1.
SH		Ozs. I	4.00	3.25	:	2.00	8.00	13 50			2.33		12.00	10.00	:		1.50		1.00		00.6			:	14.00		10.44
	AV, WT. AT ENTRANCE	Lbs. C	1	10		=======================================	15 8		15	-	18	20 (	20 12	19 10			19	21		-	9 61		:		23 14	- 1	17 10
	ERVED	OBSI	3.00	2.50		1.00	1.00	4 00 1			3.33 1	4.33 2	1.00	3.50	:		2.00 1	2.00			3.00 1				. 00 2	- 1	3.27 1
	CASES.	OE (	1 3	4 2		1 1	1 1	2			3	6 4	1 1	4 3	:	1 14			2 1		1 3					-	41 3
	WBEE MRE			4	2			_	. 2		_	_	_		-	2		- 8		•					_		-10
	CONDITION AT CLOSE	1 Sick	:			:	:		_		_	:	~				:	-	-	:	:	2			:	:	1 13
	CON	Well	68	135	71	99 1	9 57	42		-	7 14	3 18	00	8	4	-	3		9		0						6 611
	WK.	Ozs.	7.14	5.50	5.57	4.21	4.49	3.43	4.54	3.00	3.37	2.83	1.98	2.09	5.74	2.34	8.08	10.50	3.00	2.00	8.50	2.10	4.50	10	2 73	5	4.86
_	GAIN	Lbs.	:	:	:	:	:		: :	:	:	:	:	:	:	:	:		:	:	:		:	:		:	:
VEIGH	Av. Case	Ozs.	6.35	10.63	14.15	8.69	9.14	3 36	2.89	2.06	0.40	12.72	68.6	11.25	10.40	12.00	3.00	10.00	13.50	00.9	2.02	10.50	2.00	2 50	2.50	0.4	9.35
NI .	PER	Lbs.	2	-	-	-	-	-		-	-	:	:	:	-	:	7	2	:	:	:		-			1	1
SHOWING GAIN IN WEIGH	WT.	Ozs.	0.64	9.79	3.05	4.21	3.35	14.05	14.43	9.62	8.20	8.78	13.55	7.38	3.60	6.93	10.66	11.00	8.25	12.00	00.6	00 9	00.9	200	12 33	14.33	5.03
OWING	AV. WT.	Lbs.	Ξ	=	13	14	15	Ť.			18	17	19	18	17	36	19	22	330	1/2	LANCE OF	22	23	200	103	1	14
SH	T. AT	Ozs.	10.29	15.16	4.90	11.52	10.21	11 50	11.54	7.56	7.80	12.06	3.66	12.13	9.20	10.93	2.66	1.00	10.75	00.9	2.00	11 50	4.00		06.1	00.0	11.68
	AV. WT. AT ENTRANCE	Lbs.	00		=			141	-	0.5	17	130	19	75133		18	17	20	100	07/05	21	21	500	10	17	2	12
	WKS.	OBS	5.37	4.84	5.41	5.86	2.60	5 64	4.16	6.03	4.87	4.50	5.00	5.38	4.60	5.13	4.33	4.00	4.50	3.00	4.00	2 00	4.00	4 00	200	30.0	5.22
	CASES	OE (	68	139	73		57	42			15		6		_	15			4	2		2	-		7 "	-	624
ac	BABIES UMBER	N	06	144	74	29	28	45	41	37	18	24	10	12	S	16	S	3	9	7	2	2	-		7 4	+	899
	AGE BY MONTHS		1		3	4	S	9	1	8	6	10	11	12	13	14	15	16	17	18	19	20	21	33	23	2	TOTAL

TABLE 32.—SHOWING FOR WELL BABIES, BKEASI-FED ON ADMISSION, LAIER CHANGED 10 MIAED FEEDING, GAIN, OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

	DIED		:	:	:	:	:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	0
	TION	Sick	:	:	:	-	:		:	:	:		:	:	-	-	:	:	:	:	:	:	:	:	:	:	3
	CONDITION AT CLOSE	Well	3	:	-	:	3	c	7 .	-	:	-	-	-	:	-	:	:	:	:	:	:	:	:	:	:	14
	WK.	Ozs.	0.89	:	0.75	1.38	2.43	,	00.1	3.33	:	2.17	1.86	2.50	11.00	2.13	:	:		:		:	:	:	:	:	2.00
	24	Lbs.	:	:	:	:	:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
EIGHT	Av. Loss Case PE	Ozs.	2.66	:	3.00	11.00	2.67	9	3.00	4.00	:	13.00	13.00	5.00	11.00	8.50	:	:	1			:	:	:	:	:	7.29
IN W	PER C	Lbs.	:	:	:	-:	:		: '	-	:	-:	-:	:	:	:	:	:	:			:	:	:	:	:	:
SHOWING LOSS IN WEIGHT		Ozs. I	15.66	:	8.00	2.00	8.33	8	00.0	14.00	:	3.00	3.00	00.9	00.6	4.00	:					:	:	:	:	:	1.88
WING	AV. WT.	Lbs. C	7 15	:		11 5	13 8			14 14		17 3	13 3	1000		17 4							:	:	:	:	13
Sно		Ozs. L	2.33	-	1.00		14.00 1			2.00		-	-	1.00		12.50											0.17
	Av. Wt. at Entrance			•	-		1		S. IV		•			-			-	•						-	•		_
	Walter State	Lbs.	8	-	6 00		13			00 10	330	00 18	00 14	00 16		00 17	•	•				•	•	•	•	•	55 13
	WKS.	.vA	3.00	:	4.00	8.00	2.33			0.00	:	00.9	7.00	2.00		4.00	:	:					:	:	:	:	3.65
	MBER	NU OF	3	:	1	-	3		7 .	_	:	_	-	-	_	2	:	:				:	:	:	:	:	17
	CONDITION AT CLOSE	Sick	2	3	2	:	1		:		:	-	1	:	:	:	:	:				:	:	:	:	:	10
	COND	Well	13	12	==	15	14	,	10	00	6	9	9	9	00	:	4	7	2		-	-	2	:	:		136
	WK.	Ozs.	3.56	4.45	4.68	4.09	3.96		3.21	2.48	2.27	3.16	2.38	3.05	2.29		1.27	2.67	4.54		1.60	1.38	0.81	:	:	:	3.29
	GAIN	Lbs.	:	:	:	:	:		:	:	:	:	:	:	:	:	:	:					:	:	:	:	
VEIGHT	Av. C	Ozs.	5.33	10.67	8.46	8.80	10.13	1	2.73	3.25	15.89	4.29	1.71	11.17	1.75	:	9.50	12.00	13.50		8 00	11.00	6.50	:	:	:	5.01
NI	PER	Lbs.	1	-	-	-	-		٦,	-	:	-	-	:	-	:	:	:	-			:	:	:	:	:	-
SHOWING GAIN IN WEIGH	Wr.	Ozs.	2.40	10.80	14.23	5.53	8.06		0.75	2.13	3.11	8.86	13.85	13.00	6.50	:	0.25	1.00	8 00		:		00.9	:	:	:	9.07
OWING	AV. WT.	Lbs.	6		10	12	13	;	0 :	16	17	17	18		19	:	18	20	20	1	10	21	17	:	:	:	14
SH	AV. WT. AT ENTRANCE	Ozs.	13.07	0.13	5.77	12.73	13.93	9	11.00	14.88	3.22	4.57	12.14	1.83	4.75	:	6.75	2.00	10.50		8 00	5.00	15.50	:	:	:	4.06
	AV. W Entr	Lbs.	7		6		=				16	16		17	18	:	17	19	00		2		100	:	:	:	13
	WKS.	SHO	0.00	00.9	5.23	6.07	90.9	;	0.03	7.75	2.00	6.43	7.43	3.66	7.75	:	7.50	4.50	05 9	2	200	8.00	8.00	:	:	:	6.38
	CVSES	OB (	15	15	- 22		- 22	,	10	00	6	7	7	9	00	:	4	7	0	,	-	-	2	:	:	:	146
ac	BABIES BABIES		18	15	14	16	18		18	6	6	00	00	7	6	2	4	2	2	•	-	-	2	:	:	:	163
	AGE BY MONTHS		1	2	3	4	S	,	0	1	00	6	10	=======================================	12	13	14	15	16	17	. %	19	20	21	22	23	Total

TABLE 33.—SHOWING FOR WELL BABIES, BREAST-FED ON ADMISSION, LATER CHANGED TO ARTIFICIAL FEEDING, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

	DIED		:	:	:	:	:	:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	0
	TION	Sick	:	:	:	:	:	:	:	:	:	:	:	:		:	:	:	:		:	:	:	:	:	
	CONDITION AT CLOSE	Well	:	-	:	:	:	:	:	-	-	-	:	2	-	3	-	:	:	-	:	:	:	:	:	12
	WK.	Ozs.	:	4.00	:	:	:	:		0.33	0.50	4.50	:	1.80	0.57	1.56	0.30	:		1.57	:	:	:	:	:	1.32
	~	Lbs.	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	
нон	Av. Loss Case Pe	Ozs.	:	4.00	:	:	:	:	:	2.00	3.00	00.6	:	3.50	4.00	8.33	1.00	:		1.00	:	:	:	:	:	7.17
IN WI	PER C	Lbs.	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:			-	:	:	:	:	:	
SHOWING LOSS IN WEIGHT		Ozs. I	:	4.00	:	:	:	:	:	8.00	4.00	00.7	:	1.00	1.00	9.33	3.00	:		8.00	:	:	:	:	:	6.25
OWING	AV. WT.	Lbs.	:	10	:	:	:	:	-:	13	14	81	:	19	16		20 1			18	:	:	:	:	:	17
SH		Ozs. L	:	0	:	:		:	:	10.00	7.00	:	:	14.50			14.00			0	:	:	:	:		13.42
	AV. WT. AT ENTRANCE	Lbs.	:	10	:	:	•	:		13 10	14	19	:	19 14	16		20 14			19	:	:	:	:		17 13
	WKS.	OBS		0	:	:	-	:	:	00.9	00.9	2.00	:	7.50	7.00		5.00			00.7	:	:	:	:	:	5.42
	VSES OF	ro l	:	-	:	:	:	:	:	-	-	-	:		_	8				-	:	:	:	:	:	12
			1			-				-				-									:	:	:	2
	CONDITION AT CLOSE	ell Sick	7	2		7		9	3	-			-	4		. 9	-	-		_	2	-			. 2	11
		. Well	2	95	9,	88	6,	96.1	23	23	02	.65	0.0	1.73	8	4	98	9		0.5	8	22				1
	PER WK.	Ozs.	3.77	4.56	3.76	6.58	3.79	=	3.53	3.72	1.30	-	2.50	=	0.78	2.44	3.86	3.00		3.50	3.08	1.67	:	:	1.55	3.40
-	GAIN	Lbs.	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:				:	:	:	:	:	
VEIGH	Av. CASE	Ozs.	15.13	9.00	12.60	6.57	14.33	14.67	3.33	1.50	5.20	12.38	15.00	13.00	2.80	13.00	11.00	2.00		5.00	2.50	10.00	:	:	8.50	4.68
N IN	PER	Lbs.	-		-	2	-	0	2	2	:	:	:	:	:	:	-	-		-	-	:	:	:	:	-
SHOWING GAIN IN WEIGHT	Av. Wr.	Ozs.	1.13	14.50	8.20	2.71	2.00	12.67	9.00	1.00	5.60	8.13	5.00	5.75	1.20	4.33	4.00	14.00		0006	7.00	:		:	:	3.76
номп	AV.	Lbs.	6	11	12	14	15	14	15	17	17	18	20	17	18	19	23	20		26	22	24	:	:	21	16
S	Av. Wt. at Entrance	Ozs.	2.00	5.50	11.60	12.14	3.67	14.00	5.67	15.50	0.40	11.75	00.9	8.75	14.40	7.33	00.6	12.00		4.00	4.50	00.9	:	:	7.50	15.08
	AV. W Entr	Lbs.	1	6	10	11	13	13	13	14	17	600	19	16	17	18	21	10		25	21	23	:	:	20	14
	WKS.	OBS	8.25	00.6	09.7	5.86	8.00	7.50	10.00	00.6	4.00	7.50	00.9	7.50	3.60	5.33	7.00	00.9	2	00.9	00.9	00.9	:	:	5.50	80.9
	ZEE OE	CAUMI	00	2	S	7	3	9		2	25	00	1	4	2	9	1	-	1	: -	2	-	:	:	2	73
ac	BABIES BABIES		00	3	S	7	3	9	3	3	9	6	-	9	9	6	2	-		. 2	2	1	:	:	2	85
	AGE BY MONTHS		1	2	3	4	25	9	7	00	6	10	==	12	13	14	15	16	17	18	19	20	21	22	23	TOTAL

TABLE 34.—SHOWING FOR WELL BABIES, MIXED FED ON ADMISSION AND THROUGHOUT PERIOD OF OBSERVATION, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911 ED

	DIE		2	2	:	-	-	:	-	:	:	:	:		:	:	:	-		:				:	:	:	00	
	TION	Sick	-	3	:	:	:	-	:	:	:	:	-	-	:	:	:			:	-			:	:	-	6	
	CONDITION AT CLOSE	Well	-	-	-	-	-	3	3	-	3	S	2	2	2	3	3	-	-		-	2		:	:	1	39	
	WK.	Ozs.	1.14	2.36	2.00	0.14	12.00	2.18	2.44	0.29	2.17	2.71	2.92	1.42	3.88	2.67	1.12	2.20	3.00	0.57	3.67	2.28		:	:	5.67	2.13	
	04	Lbs.	:	:	:	:	:	:	:	:	:	:	:		:	:	:			: 1	: :			:	:	:	:	
IN WEIGHT	Av. Loss Case Pe	Ozs.	4.00	8.25	00.9	1.00	2.00	9.25	7.33	2.00	4.33	09.7	1.66	5.66	5.50	8.00	6.33	1.00	3 00	4 00	5.50	8.00		;	:	8.50	7.50	
W NI	PER C	Lbs.	:	:	:	:	:	:	:	:	:	:	:		-	:	:	-		:	:			:	:	:	:	
SHOWING LOSS		Ozs. I	1.00	2.75	14.00	2.00	8.00	1.25	1.33	12.00	11.67	08.9	8.67	13.67	2.00	:	3.33	5.00	2 00	4 00	10.00	10.50		:		2.00	13.42	
WING	AV. WT.	Lbs. C	6		9 1		13	13 1		16 1	18 1		18	16	-	22		21		-				:	:	_	16 1	
SHO		Ozs. L	5.00		4.00	8.00	4.00	4.50		14.00	:	14.40	4.38	3.33	2010		99.6		10 00			2.50		:	_	7.50	4.92	
	Av. Wt. at Entrance	1	_	_	_		_	4	_		-		-			-	_				_	_	-					
	1007.02	Lbs.	6 0	-	0 10	0 13	0 14	-	0 13	0 16	0 19	0 18	0 19	17			71 17	0 22						:		0 25	17	
	WKS.		3.50	3.50	3.00	7.00	1.00	4.25	3.00	7.00	2.00	2.80	4.00	4 00	4.00	3.00	5.67	5.00	1.00	7.00	1.50		:	:		1.50	3.52	
	CASES	NUI OF (	2	4	1	-	-	4	3	-	3	S	3	"	2	3	3	_		-	, ,		1	:		. 2	48	
	TION	Sick	:	2	:	-	-			:	2	-	-		: :	:	-		:	:	:	:	:	:		: -	10	
	CONDITION AT CLOSE	Well	30	47	30	32	16	10	24	21	21	20	26	1,	15	10	7	9	u	o 11			,	:	-	2 2	361	
	WK.	Ozs.	4.75	4.94	5.47	3.23	3.10	3.52	3.19	2.92	3.41	4.26	89.2	9 55	4.13	3.03	2.83	17.	1 0	9 6	000	64.9	2	:	1.71	3.84	3.81	
	GAIN	Lbs.		:	:	:	:			:	:	:	:		: :		:		:			:	:	:		: :	:	-
WEIGHT	Av. CPER CASE	Ozs.	1.87	9.94	0.13	10.63	11.53	13.05	12.46	1.67	1.04	12.81	1.29	15 00	10.40	10.60	1.62		7 60	1000	6.00	0000	0.00	:			10.58	
N NI	PER	Lbs.	2	2	2	1	130	-	1000		-	1	1		: -		-	-		1333			1	:		: -	-	-
SHOWING GAIN IN	NT.	Ozs.	4.27	1.27	3.03	7.24	12.23	8 05	12.13	10.81	14.56	12.57	10.59	15 76	0.93	5.40	15.00	12 33	000	2.00	14 00	4 22	4.55	:	10 00	12.67	14.92	
OWING	AV. WT.	Lbs.	10	12	12	13	14	15	17		2000	11600	177	10	3314	- 33	99.	00	-	07			27	:	24	NO 2015	15	-
SH	A CONTRACT OF	Ozs. 1	2.40	7.33	2.90	12.61	0.70	11 00	15.67	9.14	14.52	15.76	9.30	94.0	6.53	4.80	13.38	12 33	000	7.60	00.0	11 67	70.1	:	14.00	12.67	4.34	-
	AV. WT. AT ENTRANCE	Lbs.	00	6	10		13	13	1000	(100)	Up las	Value of	1000	10	17	-	_	101	10.0	Section 1	010	11		:	23 1	(15 × 10 ×	14	1
	CHARD	OBSE	7.13	8.49	5.87	8.24	88.8	8 26	8 92	6.05	5.00	92.9	6.44	00 4	6.40			0 33	200	00.4	1.20	1.00 6 22	00.0	:	00 4	4.17	86.9	
	WKS.	OF C	30 7				17 8	10		_			27 6	1,							0 +	1000				1 9	371 (	
-	BABIES	1	34					23		9704			30	00			==	o	0 4	0 4	2 0	) W	0	:	-	- 00	428 3	
d	AGE BY MONTHS	-200	-		1 60		100								13			71	10	77	0 0	61	07	21	23	23	TOTAL 4	

TABLE 35.—SHOWING FOR WELL BABIES, MIXED FED ON ADMISSION, LATER CHANGED TO ARTIFICIAL FEEDING, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

	DIE		-	2	:	:	:	-	:	2         2         10.00         15         10.00         18         2.50         2         8.50         4.05         2		4														
	TION	Sick	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	-		:	:	1
	CONDITION AT CLOSE	Well	:	:	:	:	:	:	:	:	:	-	-	-	-	:	:		-	:	:	:		-	:	9
		Ozs.	:	:	:	:	:	:	:	:	:	80.0	5.67	1.25	0.75	:	:	:	0.67	:	:	0.27		2.67	:	86.0
	OSS PER WK.	Lbs.	:	:	:	:	:	:	:	:	:	:	:		:	:	:		:	:	:	:		:	:	
HOIS	Av. Loss Case   Pe	Ozs.	:	:	:	:	:	:	:	:	:	1.00	1.00	0.00	3.00	:	:		4.00	:	:	3.00		8.00	:	6.57
IN WI	PER C	Lbs.		:	:	:	:	:	:	:	:	:	-			:	:		:	:	:	:			:	
SHOWING LOSS IN WEIGHT	- 3300	Ozs. I		:	:	:	:	:	:		:	00.	00.9	00.	00.	:	:		00.8	:	:	00.0			:	4.86
OWING	AV. WT.	Lbs. C	:		-	:	:	:		:		_		1			:							1		18
SH		Ozs. L																					-			
	AV. WT. AT ENTRANCE	Lbs. O										7 13	36	-	10000				_			_		777		I SURPLINE
	ERVED				•	•	•	•		•		00				•	•				•					
	WKS.	.vA		•	•	•	•	-	•		•	12.0	3.	×	1.4	:	:		1 6.0	:	:					1
-	MBER			:	:	:	:	:	:	:	:	_	_			:	:		_	:	:	_			-:	
	CONDITION AT CLOSE	Sick	1	:	-	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:		:	:	2
	CON	Well	:	2		3	:	:	2					1					:	-	:	-	-	-	:	32
	VIN PER WK.	Ozs.	6.00	3.43	0.32	4.67	:	:	0.86	4.05	0.13	4.29	1.05	4.00	5.00	5.62	2.65	1.79	:	1.50	:	12.00	4.17	1.00	:	3.24
	GAIN	Lbs.	:	:	:		:	:	:	:	:	:	:	-	:	:	:		:	:	:	:	-	:	:	
VEIGHT	Av. Case	Ozs.	14.00	7.50	1.00	14.67	:	:	9.50	8.50	1.00	14.00	8.60	4.00	10.00	4.50	1.25	1.00		9.00	:	4.00	0.00	10.00	:	8.62
4 IN W	PER	Lbs.	1	2	:	2	:	:	:	2	:	-	:	1		2	-	-	:	:	:	7	7		:	-
SHOWING GAIN IN WEIGH	Vr.	Ozs.	1.00	00.6	6.50	00.6	:	:	1.00	2.50	1.00	0.50	2.80	9.00	:	0.50	1.75	3.50	:	00.6	:	:	2.00	0.00	:	89.6
HOWIN	Av. Wr.	Lbs.	12	13	6	14	:	:	15	18				16	23	21	20	19	:	22	:	23			:	17
S		Ozs.	3.00	1.50	5.50	10.33	:	:	7.50	0.00	0.00	2.50	4.20	5.00	00.9	2.00	0.50	2.50	:	:	:	2.00	3.00	:	:	1.06
	Av. Wt. at Entrance	Lbs.	10	11	6	==	:		14				-	15				18	:	22			20	18	:	16
	WKS.	OBSI	5.00	11.50	4.50	10.00	:	:	0								6.50		:		:				:	7.59
	CASES	OE (	-	-1-1	2		:	:	4760			-			_					-		-	-		:	34
ac	BABIES UMBER		2	4	2	3	:	-	2	2	-	3	9	2	2	2	4	2	-	1		2	-	2	:	45
	AGE BY MONTHS		1	2	3	4	s	9	7	00	6	10	11	12	13	14	15	16	17	18	19	20	21	22	23	TOTAL

TABLE 36.—SHOWING FOR WELL BABIES, ARTIFICIALLY FED ON ALMISSION, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

	Drei		-	2	:	-	:	-	:	-	:	2	:		-	'	-		: '	-	-	:	:		:		:	12	
	TION	Sick	-	2	:	:	:		-	-	-	:	:		:	. 2	:			:		-	:		: '	-	:	10	
	CONDITION AT CLOSE	Well	ıs	S	2	S	7	0	9	3	S	S	00	0	1	12	S	0	2 '	2	S	S	6	4	۲,	-	7	134	
	WK.	Ozs.	1.88	2.94	0.38	2.39	0.35	1.13	1.75	2.61	2.48	0.74	2.34	4.63	2.61	2.15	1.69	9 63		2.24	2.94	2.46	1.31	1 61	100	2.60	1.18	2.13	
	Loss	Lbs.	:	:	:	:	:				:	:	:		:	: :			:	:	:	:	:		:	:	:	:	
WEIGHT	Av. L	Ozs.	7.83	6.29	3.00	11.00	3.00	4.78	2.00	2.25	10.33	5.20	1.00		13 43	14.57	8.80	3 00	0.50	09.7	9.40	9.83	2.67	20 0	07.6	14.00	4.71	10.49	
IN W	PER C	Lbs.	:	:	:	-	:				-	:	-	-		:	: :		:	:	:	:	:		:	:	:	:	
SHOWING LOSS IN		Ozs. I	4.50	2.28	11.00	5.60	00.9	0.78	9.57	0.25	1.50	00.9	7.50	14 00	0 43	2.00		11 70	0.10	2.80	0.40	6.33	15.89	. 75	0.10	8.00	12.29	13.19	
WING	Av. Wr.	Lbs. (	6	6		_	13	13	_	_	_	15	-	17 1		_	•	185 To 110				21 (	20 13	21	-		21 13	16 1	
SHO		Ozs. L	12.33	8.57	14.00	10.60	00.6	5.50					20	3 11							9.80	0.16	5.56				1.00	7.68	
	AV. WT. AT ENTRANCE	Lbs. 0	9 12	8 6	1 14	13 10	13 9	13			_	-	16 8.	10		_							21 5	,,			22 1	17 7	
	ERVED		4.17	2.14	8.00	4.60 1	8.50 1	4 22 1						4 56 1									4.33 2				4.00 2	4.93	
	Wks.	.vA		82		5 4.			_		6 4.	_	-	_		1.555				-				_					
	NBER	inN		_							_				1	-	100	, ,	=	S		9	6			2		144	
	CONDITION AT CLOSE	Sick	:	:	:	:	-		:	: :	-	-	:	-	-	-	1 1/2	, -	-	-	2	1	:		-	_	:	18	
	CONI	Well	25	29	28	34	33	30					47	2.							29	30	37				55	806	İ
	WK.	Ozs.	4.05	4.68	4.12	4.28	3.26	3.11	3.83	2.99	3.56	2.67	2.97	2 20	9 99	4.33	2.66	60 6	6.00	2.35	2.52	2.66	2.34	9 6	9.00	3.88	3.03	3.29	
	GAIN	Lbs.	:	:	:	:	:			: :	:	:	:		:	:	:		:	:		:	:		:		:	:	
VEIGHT	Av. C	Ozs.	11.68	14.83	11.50	0.88	9.91	0 50	10.52	6.39	80.9	2.18	4.81	0 27	2 16	12 01	1.96	74 74	10.74	15.88	0.45	0.03	15.40		40.7	8.04	14.32	5.84	
NI V	PER	Lbs.	1			2	-	-			-	-	-	-		1576			:		-	6491	:	,	-	Evo	:	1	-
SHOWING GAIN IN WEIGH	Wr.	Ozs.	10.12	14.55	96.0	13.29	62.6	5 67	7 92	3.67	8.88	3.91	3.26	2 4	77.7	2.10	6 78	63	00.11	14.98	3.61	2.22	12.29	000	00.0	5.56	5.45	8.18	1
OWING	Av. Wt.	Lbs.	10	13		13 1	1000	14.1	1007	16	17	17	18	0	10	12-11	Marie .	92	201		400	-0.00	22 1		17	23	23 1	18	1
SH	L. AT	Ozs.	14.44	15.72	5.46	2.41	15.88	80.9	13.40	13.28	2.80	1.73	14.45	13 33	2 50	0.02	4 82	9	11.89	15.10	3.16	2.19	12.89	700	0.00	3.52	1.13	2.34	
	Av. Wt. at Entrance	Lbs.	8		10	11	12 1	-		10.00			-	7			-			19	21	21		6	4	100	23	17	1
	ERVED WKS.	ESEO	6.84	30.5	89.9	89.7		8 23	6 03	7.50	6.20	6.82	2.00	1 22					10.0	6.85	6.52	6.03					4.73	6.64	-
	YSES	OB C	25			34	-	30						_	27			_	_		_		37	-	-	-	55	824	-
-	BABIES OF		32	38	30	40	36	40	40	41	41	40	55	;	14	24	30	5 1	45	47	37	37	46	1	17	27	62	8086	-
	AGE BY MONTHS		1	2	3	4	25	4	10	- 00	0 0	10	=		77	5 5	1 1	2	10	17	18	19	20	,	77	22	23	TOTAL	

TABLE 37.—SHOWING FOR SICK BABIES, BREAST-FED ON ADMISSION AND THROUGHOUT PERIOD OF OBSERVATION, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

ED

	DIE				-	:	:		:	:	:	1	:	:		: :	:	:		:	:	:	:	:		:	:	:	2
	TION	Sick		-	2	-	1		:	:	-	:	:	-		-	:	1				7	:	:		:	:	-	12
	CONDITION AT CLOSE	Well	:		: :	:	:		:	1	-	:	:	-	-	-	:	:		:	:	:	-	:		:		:	9
	WK.	Ozs.	;	2.00	6.71	8.00	3.50		:	3.00	2.14	:	:	1.13	77	0.88	:	3.00		:	. 0	0.00	90.0	:		:		9.00	3.19
	000	Lbs.			: :	:	:		:	:	:	:	:	:		: :	:	:		:	:	:	:	:		:	:	-	
IGHT	7-	Ozs. I	:	2.00	7.50	8.00	00.7		:	2.00	7.50	:	:	4.50	2 00	5.00	:	3.00		:		7.50	3.00	:		:	:	11.00	10.72
IN WE	AV. PER CASE	Lbs. C				:	:		:	:	:	:	:	:		: :	:	:		:	:	:	:	:		:	:	4	:
Loss		Ozs. L		2.00	2.00	:	00.9			2.00	3.50	:	:	00.9	10 00	1.00	:	2.00		:	. 0	7.00	2.00	:		:		2.00	14.22
SHOWING LOSS IN WEIGHT	Av. Wr.			-	_	_	_		_	-		-	-		15 10	•	-	17 5			-		25 5	:		:		23 5	14 14
SHO		. Lbs.		9 00		_					00 15	-	-	50 15	1 0			0	-				8.00 2	_				•	8.94
	AV. WT. AT ENTRANCE	Ozs.		4.00	-	8.00	13.00		:	4.00	11.00	-	:	10.50						:	-		_	:		:	:		
	Av. Ent	Lbs.	:	9			80				15	:		15	16		:	11		:			72	:		:		28	15
	WKS.	Av.	:	1.00	3.50	1.00	2.00			1.00	3.50	:	:	4.00	4.00	5.50	:	1.00			. 2	3.0	2.00	:		:	:	3.00	3.39
	CASES	OB (	:	-	2	-	-		:	-	2	:	:	2	-	7	:	1		:	: "	7	-	:			:	1	18
	TON	Sick	:	3	00	4	2	c	7	3	7	3	7	2		: :	-	:				-	:	:		:	:	:	34
	CONDITION AT CLOSE	Well	S	9	S	7	9	v	0 0	7	4	3	-	-		-	-	:	:	-	, ,	7	:	:		-	:	:	46
		Ozs.	6.89	7.74	5.59	2.00	3.47	9	3.18	3.04	4.58	2.48	4.51	2.46		1.60	7.22	:		9 11	1 6	2.31	:	:		3.00	:	:	4.80
	AIN PER WK.		9								-		-										:	:		7	:	:	-
HT	0	s. Lbs.	. 02	. 19	46	. 19	. 55		4.43	1.00	. 19	17	00	2.00		8.00	0.50				200	. /0	-	_	_	00.9		100	8.19
WEIG	AV. PER CASE	Ozs.	5.2	0.67	8.46	10.67	5.25	-	4.		7.	11.1	12.00	2.0		8.0	0.	:				0.0	:	:				:	∞i
N IN		Lbs.	- 20		1						3 1	:	:	1		: :		:		: "		:	:	:			:	:	0
SHOWING GAIN IN WEIGHT	AV. WT.	Ozs.	7.00	9.56	15.00	6.17	1.75	0	8.00	11.20	13.83	4.17	10.33	5.66		11.00	13.50	:			0.0	14.33	:	:		:		:	14.30
номп	AV.	Lbs.	10	Ξ	==	14	15		12	14	17	16	14	19		18	18	:		: 6		13	:	:		25	:	:	14
S	AV. WT. AT ENTRANCE	Ozs.	1.80	8.89	6.54	11.50	12.50		3.57	10.20	6.16	9.00	14.33	3.66		3.00	13.00	:		3.	3.	2.00	:	:		10.00	:	:	6.13
	AV. W Entr	Lbs.	00	6	10		2000	:			16	15	13	18		18		:		: :	1 5	19	:	:		21	:	:	13
	Wks.	OBSI	5.40	4.22	4.38	5.33	6.13	67.7	6.43	5.60	5.17	4.50	2.66	7.33		5.00	4.50	:			3.50	3.00	:	:		2.00	:	:	5.04
	NBER	NUI NUI	ıs	6	13	9	00	1	7	S	9	9	3	3		: -	2	:			7 0	3	:	:	1	-	:	:	80
			100	-	9	1	6		1	9	00	7	3	S	-	4 10	2	-			4 1	0	-	:		-		-	100
A	BABIES BABIES	IN	I/O	10	16																								

TABLE 38.—SHOWING FOR SICK BABIES, BREAST-FED ON ADMISSION, LATER CHANGED TO MIXED FEEDING, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

		DIE		:	:	:	:	:		:				: :		:		: :		: :	:	:	:		:		:	0
-		TION	Sick	:		:		:		: :	-			1		:	: :				:	:	:		:	:	:	2
		CONDITION AT CLOSE	Well	:	:	:	:	:		: :				:		:	: :	:		: :	:	:	:		:	:	:	:
		WK.	Ozs.	:	:	:	:	:		: :	2.00			0.30		:	: :	:		: :	:	:	:	. 1	:	:	:	0.36
		OSS PER WK.	Lbs.	:	:	:	:	:		: :	:	- :		:		:	: :	:		: :	:	:	:		:	:	:	:
	SHOWING LOSS IN WEIGHT	Av. Loss Per Case   Pe	Ozs.	:	:	:	:	:		: :	2.00	:		2.00		:	: :	:		: :	:	:	:		:	:	:	2.00
	IN N	PER (	Lbs.	:	:	:	:	:		: :	:	:		:		:	: :	:		: :	:	:	:		:	:	:	:
	Loss	/T.	Ozs.	:	:	:	:	:		: :	00.6	:		0.00		:		:		: :	:	:	:					4.50
	OWING	AV. WT.	Lbs. (	:	:	:	:	:		: :	15			17 (		:												16 4
	SH		Ozs. L						-		1.00			2.00														6.50 1
		AV. WT. AT ENTRANCE	Lbs. 0		•						5 11															-		
		БЕУЕВ			-		-	-			-	•		17		•						•	•			•		09 10
		WKS.	.vA		-	:	<u>:</u>	•			-	-	:	10				:			:	:	:		:	:	:	5.50
=		MBER		:	:	:	:	:				:	:	_		: :		:		:	:	:	:		:	:	:	2
		CONDITION AT CLOSE	Sick	2	:	:	:	:	-	-	:	-	-	1		: :	:	:	-	:	:	:	:		:	:	:	∞
		CONT	Well	9	:	:	7	:	-	2	:	3	1	:		: :	-	:	-	:	:	:	:		:	:		17
		NIN PER WK.	Ozs.	3.74	:	:	2.00	:	1.60	2.77	:	9.07	0.41	0.64		: :	10.80	:	6.71	:	:	:	:		:	:	:	3.40
	_	GAIN	Lbs.	:	:	:	:	:		:	:	:	:	:		: :	:	:			:	:	:		:	:	:	
	SHOWING GAIN IN WEIGHT	AV. OPER CASE	Ozs.	7.38	:	:	14.00	:	8.00	12.00	:	15.75	5.50	7.00		: :	00.9	:	7.50	:	:	:	:		:	:	:	4.52
	NI Z	PER	Lbs.	-	:	:	:	:			:		:	:		: :	3	:	1	:	:	:	:		:	:	:	1
	G GAIR	WT.	Ozs.	7.38	:	:	4.00	:	8.00	3.00	:	14.75	1.50	2.00			11.00	:	11.00	:	:	:	:		:	:	:	3.72
	OWIN	AV. WT.	Lbs.	6	:	:	=	:	16	15	:	18	18	17		:	1	:	18		:	:	:			:	:	14
0	מ	AV. WT. AT ENTRANCE	Ozs.	:	:	:	00.9	:	:	00.7	:	15.00	12.00	11.00		:	5.00	:	3.50	:	:	:	:			:	:	15.20
		Av. W Entr	Lbs.	00	:	: :	10	:	16	14	- 143	10117	100	16 1	:	:	6	:	17	:	:	:	:	:		:	:	12 1
		WKS.	OBS	6.25	:	: 1	7.00	:	5.00	4.33	:	3.50	13.50	11.00	:	:	2.00	:	3.50	:	:	:	:	:	-		:	6.04
		CASES	NO N	00	:	: '	7	:	2	3	:	_	2 1	-	:	:	-	:	2	:	:	:	:	:		:	:	25
	OE	BABIES	N	00	:	: '	7	:	2	3	-	4	7	7	:	:	1	:	7	:	:	:	:	:		:	:	27
-		AGE BY MONTHS		-	7	ю.	4 -	S	9	7	00	6	10	=	12	13	14	15	16	17	18	19	07	21	22	23	2	TOTAL

TABLE 39.—SHOWING FOR SICK BABIES, BREAST-FED ON ADMISSION, LATER CHANGED TO ARTIFICIAL FEEDING, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF

						0	OBSERVATION	LVAT	TON.	PE	PERIOD JUNE	JUL	VE 1	TO	SEPT	EMB	TO SEPTEMBER 15, 1911	5, 19	=							1
	a					SHOW	SHOWING GAIN IN WEIGHT	N IN	VEIGHT								S	HOWIE	SHOWING LOSS IN WEIGHT	IN W	BIGHT				1	
AGE BY MONTHS	MBER O	GREE CASES	WKS.	Av. Ent	AV. WT. AT ENTRANCE		AV. WT.	PER	AV. GAIN PER CASE   PE	GAIN	PER WK.	CONDITION AT CLOSE	LOSE	MBER	WKS.	Av. W Entr	AV. WT. AT ENTRANCE	AV.	AV. WT.	AV. PER CASE	Av. Loss	OSS PER WK.		CONDITION AT CLOSE		Dir
	UN	MUN	.vA	Lbs.	Ozs.	Lbs.	Ozs.	Lbs.	Ozs.	Lbs.	Ozs.	Well	Sick	N N OF		Lbs.	Ozs.	Lbs.	Ozs.	Lbs.	Ozs.	Lbs.	Ozs.	Well 8	Sick	1
-			13.00	0	1.33	10	11.33	-	10.00	:	2.00	2	1	:	:	:	:	:	:	:	:	:	:	:	:	:
2	) 4				0.50		15.00	-	14.50	:	6.78	2	2	:	:	:	:	:	:	:	:	:	:	:	:	:
3	:		:	:	:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
4	3		200		14.33			:	10.33	:	1.94	7	-	:	:	**	:	:	:	:	:	:	:	:	:	:
S	-		2.00	6	9.00	6	15.00	:	00.9	:	1.20	-	:	:	:	:	:	:	:	:	:	:	:	:	:	:
9	-		00.9	13	2.00	14	7.00	-	5.00	:	3.50	-	:	:	:	:	:	:	:	:	:	:	:	:	:	:
1	:	-			:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
80	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
66	:	:	:	:	:	:	:	:	:	:	:	:	:		:	:	: !	: :	: !	:	: :	:	: ;	:	: '	:
10	-		:	:	:	:	:		:		:	:	:	-	3	14	8.00	13	13.00	:	11.00	:	3.67	: '	-	:
11	3	2	8.00		15.00	14	15.00	3	:	:	00.9	2	:	-	S	17	:	15	00.6	-	7.00	:	4.60	-	:	:
12	-		9.00	=	3.00	12	4.00	-	1.00	:	1.89	-	:	:	:	:	:	:	:	:	:	:	:	:	:	:
13	:				:		:	:	:	:	:	:	:	:	:	:	:	:	:		:	:	:	:	:	:
14	:	_	:	:	:		:	:	:	:	:	:	:	:	:		:	:	:	:	:	:	:	:	:	:
15	-	-	8.00	15	00.9	17	:	-	10.00	:	3.25	-	:	:	:	:	:	:	:	:	:	:	:	:	:	:
16																				:		:		:		:
17	: :	: :	: :	: :	: :	: :	: :	: :	: :	: :	: :	:		:	:	:	:	:	:	:	:	:	:	:	:	:
18	:	:	:	:	:	:	*	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
19	:				:	:			:	:	:	:	:				:	:	:	:	:	:	:	:	:	
20	-	_	3.00	19	8.00	20	:	:	8.00	:	2.67	-	:	:	:	:	:	:	:	:	:	:	:	:	:	:
21	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
22	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
23	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
TOTAL	. 19	17	7.06	6	11.00	11	2.82	1	7.82	:	3.37	13	4	2	4	15	12.00	14	11.00	-	1.00	:	4.25	-	-	0
The state of the s																			1	-		ı	1		Ī	-

IABLE 40. - SHUWING FUR SICK BABIES, MIXED FED ON ADMISSION AND THROUGHOUT PERIOD OF OBSERVATION, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

		Dig		3	4	:	7	:	:	-	_	:	:	:	-	:	:	:	:	:		:	:	:	:		:	:	:	=	
F		CONDITION AT CLOSE	Sick	-	7	-	-	2	0	2	:	-	:	-	,	4 -	- (	7	-	3	0		- 1	7			:		:	26	
		COND AT C	Well	:	-	-	:	:	:	:	:	-	:	-		: '	-	:	:			:	:	:	:		:	:	:	5	
			Ozs.	09.0	2.84	1.33	0.50	1.50	1.38	3.43	:	1.13	:	1.67	-		4.17	2.90	3.00	2.70	0 67		0.33	2.30	:		:	:		2.13	
		Loss Per Wk.	Lbs.	:	:	:	:	:	:	:	:	:	:	:		:	:	:	:	:		:	:	:	:		:	:	:	:	1
	SHOWING LOSS IN WEIGHT	Av. I	Ozs.	3.00	2.00	8.00	1.00	3.00	11.00	8.00	:	4.50	:	7.50	2	00.0	12.50	14.50	3.00	00.0	3 00	900	1.00	11.50			:		:	9.16	1
	N N	PER	Lbs.	:	-	:	:	:	:	-	:	:	:	:		:	:	:	:			:	:	:	:		:	:	:	:	1
	ross	WT.	Ozs.	5.00	11.67	9.50	00.9	9.50	4.00	7.50	:	11.00	:	7.50	000	2.00	0.00	7.50	1.00	6.67	12 50	00.0	:	4.50	:		:	:	:	8.90	
	IOWING	AV. WT.	Lbs.	9		6	6	11	6	10	:	14	:	15	,	07	0 !	17	18	22	100.0		70	19	:	T	:	:	:	14	-
	Š	Av. Wt. at Entrance	Ozs.	8.00	13.67	1.50	7.00	12.50	15.00	15.50	:	15.50	:	15.00	i.	0000	2.50	00.9	4.00	5 67	0 50	00.0	1.00	:	:		:	:	:	2.06	
		Av. W Entr	Lbs.	9	8	10	100	=======================================	6	=======================================	:	14	:	15	,	0 !	11	18	18	22	0	01	07	20	:		:	:	:	15	
		WKS.	OBS	5.00	6.33	00.9	2.00	2.00	8.00	7.00	:	4.00	:	4.50	000	2.00	3.00	2.00	1.00	3 33	4 50	4.30	3.00	2.00	:		:	:	:	4.32	-
		CASES		-	3	2	-	7	-	2	:	2	:	2		7	7	7	-	"		7	-	2	:		:		:	31	-
		TION	Sick	S	6	3	S	4	3	-	:	S	-	4	-	- ,	0	9	-		,	:	7	-	-		:	-	3	11	=
200		CONDITION AT CLOSE	Well	2	4	6	4	S	3	2	7	3	3	-	,	0	-	7	:			7	-	:	:		:	:	3	55	
201			Ozs.	2.35	3.65	6.31	3.31	89.2	3.57	3.34	2.09	4.52	3.71	4.33	0	08.2	4.33	3.96	4.57	32.2	200	2.11	2.17	12.00	2.13		:	1.57	89.2	3.64	-
1		GAIN PER WK.	Lbs.	:	:	:	:	:	:	:	:	:	:	:		:	:	:	:		:	:	:	:	:		:	:	:	:	-
:	WEIGHT		Ozs.	10.43	13.23	10.58	10.44	3.33	00.9	2.50	11.50	14.00	13.00	15.60	- 1	2.00	0.71	1.00	6.85	12 22	4.00	4.00	13.00	8.00	1.00		:	11.00	13.33	7.23	-
100	Z	Av. Per Case	Lbs.			_	-	-	-	2	:	-	:	:		-	-	-	-		: 0			-	-		:	:	:	1	
OBSERVATION	SHOWING GAIN	WT.	Ozs.	1.71	0.46	14.50	5.33	3.44	8.67	7.67	0.50	0.88	1.50	1.20		7.07	10.57	4.63	10.71	2,66	200	00.0	00.9	8.00	10.00		:	13.00	14.00	13.06	-
5	OWING	Av. WT.	Lbs.	1	==	1000	1000	==	13	12	17	16	16	15				1545	17	1	1 5	19	20	16	16			1000	23	14	-
	SH	AV. WT. AT ENTRANCE	Ozs. 1	7.29	3.23	3.92	10.89	0.11	2.67	5.17	5.00	2.88	4.50	1.60		13.67	98.6	3.63	3.86	6 22	0.00	4.50	00.6	:	00.6		:	2.00	0.67	5.83	-
		Av. W Entr	Lbs.	9	6	11		10	12	10	16	14	15	14			10	15	16	4	2 5	1/	19	15	15		:	18	23	13	
		WKS.	OBSI	4.43	8.00	6.75	8.00	7.22	6.17	10.33	5.50	6.63	3.50	3.60	-	7.50	3.86	5.75	2.00	2 67	2000	13.00	00.9	2.00	8.00		:	00.7	5.17	6.38	-
		YSES	OB (	7	13		6		_	6 1			_	2	_		7			"		_	_	-			:	_	9	126	
-	B	BABIES UMBER C		=	20	14	12	11	7	6	3	10	4	7		00	6	10	00	4	,	4	4	3	-		:	-	9	168	10000
		AGE BY MONTHS		1	7	3	4	5	9	1	80	6	10	11		12	13	14	15	16	07,	17	18	19	20		21	22	23	TOTAL	

TABLE 41.—SHOWING FOR SICK BABIES, MIXED FED ON ADMISSION, LATER CHANGED TO ARTIFICIAL FEEDING, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

	DIE		1	:	:	:	:	-	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:		: :		2
	TION	Sick	:	:	:	:	:	:	:	:	:	:	:	:	:	2	:	:			:	:	:		: :	1	2
	CONDITION AT CLOSE	Well	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	-	:	:	:	:	:		: :	İ	-
		Ozs.	:	•	:	:	:	:	:	:	:	:	:	:	:	1.30	:	12.00	:	:	:	:	:		: :	İ	3.08
	DER WK.	Lbs.	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:		: :	T	
ICHT	Av. Loss	Ozs. I	:	:	:	:	:	:	:	:	:	:		:	:	0.50	:	8.00	:	:	:	:	:		: :	1	12.33
N WE	AV. PER CASE	Lbs. C																1 8	•						: :	1	
7088 I					·	-			-	·					-	5.00	-	10.00		-	-	<u>.</u>			700	1	3.00
SHOWING LOSS IN WEIGHT	AV. WT.	s. Ozs.		•	•		•	•	•		•			-	:	_	•	1000	•	•	•	•		_		+	_
SHOV		Lbs.	:	•	•	•	:	:	•	:	:	:	:	:		0 18	:	0 19	:	•	:	:	-		: :	-	3 19
	AV. WT. AT ENTRANCE	Ozs	:	:	:	:	:	:	:	:	:	:	:	:		00.9	:	2.00	:	:	:	:	-		: :		15.33
	Av. Ent	Lbs.	:	:	:	:	:	:	:	:	:	:	:	:	:	19	:	21	:	:	:	•	:		: :		19
	WKS,		:	:	:	:	:	:	:	:	:	:	:	:	:	5.00	:	2.00	:	:	:	:	:		: :		4.00
	CASES	NU TO	:	:	:	:	:	:	:	:	:	:	:	:	:	2	:	-	:	:	:	:			: :		3
	ION	Sick	:	:	:	:	:	:	:	:	:	:	:	-	:	:	:	2	:	:	:	:			: :	İ	3
	CONDITION AT CLOSE	Well	3	:	:	:	-	:	:	:	-	-	-	-	2	1	:	2	:		:	:	:	_	: -	-	14
		Ozs. V	3.63				2.33				80.2	1.20	10.83	2.87	3.56	7.75		2.90							1.40	1	3.69
	PER WK.	-					• • •					-	9				•	6%	-	-		-				+	
нт	0	Lbs.		•	•	•											•		•	•	•	:	-	_	: 2		
SHOWING GAIN IN WEIGHT	AV. PER CASE	Ozs.	3.3	:	:	:	7.00	:	:	:	9.00	12.00	2.00	5.50	:	15.00	:	15.2	:	:	:	:	- :		7.00		9.18
NI NI		Lbs.	1	:	:	:	:	:	:	:	1	:	8		2		:	:	:	:	:	:			: :	310	1 1
NG GA	AV. WT.	Ozs.	12.00	:	:	:	7.00	:	:	:	2.00	11.00	8.00	12.00	7.00	15.00	:	13.50	:	:		:			7.00		1.06
номг	AV.	Lbs.	10	:	:	:	==	:	:	:	15	12	18	17	16	17	:	18	:	:	:	:			21		16
53	AV. WT. AT ENTRANCE	gozs.	8.67	:	:	:	:	:	:	:	9.00	15.00	00.9	6.50	7.00	10.00	:	14.25	:	:	:	:			:	:	7.88
	AV. V ENTR	Lbs.	6	:	:	:	11	:	:	:	13	Ξ	10	16	14	16	:	17	:	:	:	:			21	1	14
	WKS.	OBSI	5.33	:	:	:	3.00	:	:	:	12.00	10.00	12.00	7.50	00.6	4.00	:	5.25	:	:	:	:			2 00		6.82
	CASES	Ob (	3	:	:	:	-	:	:	:	1 1	No.	1 1	1000	2		:	4	:	:	:	:			: -	-	17
ac	BABIES DABIES		4	:	:	:	-		:	:	-	-	-	2	2	3	:	r.	:	:	:	:			: -	1	22
	AGE BY MONTHS		1	2	3	4	S	9	7	80	6	10	=======================================	12	13	14	15	16	17	18	19	50	21	20	23		TOTAL

TABLE 42.—SHOWING FOR SICK BABIES, ARTIFICIALLY FED ON ADMISSION AND THROUGHOUT PERIOD OF OBSERVATION, GAIN OR LOSS IN WEIGHT FOR EACH AGE GROUP, AND CONDITION AT CLOSE OF PERIOD OF OBSERVATION. PERIOD JUNE 1 TO SEPTEMBER 15, 1911

	DIED		3	4	S	4	2	3	4	-	2	7	-		: :		:	-		:	:	:	:			:	:	32	1
	TION	Sick	-	3	2	4	9	ıc	S	3	9	3	4		2	-	2	4			3	-	2	2		-	4	89	
	CONDITION AT CLOSE	Well	-	:	:	2	-			-	-	3	2	-		:	:			7	:	:	7	:		:	-	18	
		Ozs.	1.00	3.00	2.60	1.86	1.77	5.06	0.89	2.00	1.55	1.76	3.10	9 54	4.58	1.55	6.80	8.94	000	2.30	2.72	2.00	1.25	2.63	-	0.50	1.86	2.48	
	OSS PER WK	Lbs.	:	:	:	:	:	:			:	:	:		:	: :	:		:	:	:	:	:			:	:	:	
BIGHT	Av. Loss Case Pe	Ozs.	5.00	15.00	6.50	6.83	6.57	0.20	7.80	14.00	6.43	7.33	5.50	7 00	2.33	1.00	1.00	00 9	1 61	1.0.1	6.33	7.00	6.25	10.50	000	2.00	8.20	10.90	
IN W	PER C	Lbs.	:	-:	:	:	:	-		-	:	:	:		-		-	,	1	:	:	:	:				:	:	1
SHOWING LOSS IN WEIGHT	VT.	Ozs.	11.00	5.67	10.50	1.50	2.71	0.80	3.60	9.75	98.0	0.17	8.17	200	0.33	5.00	10.00	15 75	2.00	00./	4.67	00.9	4.75	10 00	20.0		10.40	10.38	-
OWING	AV. WT.	Lbs.	8	1000	11 1	1000	12	=	=				15	1	10000		alie i	18	-			18	19 1	22 1		21	19 1	14 1	-
SH		Ozs. L	:	4.67	1.00	8.33	9.28	1.00						20 0				77				13.00	2.00	4 50			2.60	5.28	-
	Av. Wt. at Entrance	Lbs.	6			-	12	12		1			16		11					2000		18 1	20	23		21	20	15	-
	WKS.	OBS	5.00	5.00	2.50	3.67	3.71	3.20	8.80	7.00	4.14	4.17	5.00	27.0	4 00	1.00	2.50	A 25	27.7	3.33	2.33	1.00	2.00	4 00	200.1	4.00	4.40	4.40	-
	CASES	OE (				9				-	7	-	No. 17 to	-		-	7	-		3		-		-			S	98	-
		Sick	2	ro.	7	-	00	=	14	00	7	3	4		) W		. 00	u	0 '	9	3	2	6	-	۲		9	128	=
	CONDITION AT CLOSE	Well Si	2	11	10	16	16	13			=	16	13	4	0 0	2 1	9		CI.	9	00	==	00	-	1	6	6	232 1	-
		Ozs. W	6.23				3.43	4.30						200	4 70	3.59	2.67	9 6	00:	3.45	5.37	10.1	3.13	4 13		4.15	88.2	3.85 2	-
	NIN PER WK.		9	60	4	4	e6	4	. 4	. +	4	65	€.	•			-	•	_	-	3	-	_		-			1	-
THE	0	. Lbs.	, in	. 0	-	4		0	. 0		9	00		-			4	-	:	86	. 00	60			:	68		91	-
SHOWING GAIN IN WEIGH	Av.	Ozs.	14.75	8.50	12.71	6.64	00.9	12 70	14.30	14.75	10.16	11.00	7.23	1	12 20	7.33	13.14	-		7.58	9.90	8.69	0.76	200	13.0	8.89	14.80	8.16	-
N IN	PER	Lbs.				3 2		-					7					,				3 1				1	:	1	-
ig GA	Av. Wt.	Ozs.	12.50	5.00	7.71	6.23	4.00	3 87	13 80	10.63	5.88	13.89	12.47	1	11 20	3.62	7.85	0 67	0.0	5.91	5.63	14.23	12.70	09 0	5.6	15.89	10.53	1.08	
HOWIN	AV.	Lbs.	10	12	10	12	=	13			15	15	15	1	17		17	10	10	21	20	20	20	10			21	16	
S	AV. WT. AT ENTRANCE	Ozs.	13.75	12.50	11.00	15.59	14.00	7 17	15 50	11 88	11.72	2.89	5.24	2 22	14 00	12 20	10.71	7 5	00.1	14.33	11.73	5.54	11.94	10 60	10.00	7.00	11.73	8.92	
	AV. W Entr	Lbs.	7			6		10				15	14	,	1 10			1	1	19	18	19	19	1	1	21	20	14	
	WKS.	OBSI	7.50	6.75	6.41	88.8	6.42	6.67	6.36	6 63	5.89	5.32	8.00	,,	6.00	6.63	4.93	ti ti	0.00	6.83	4.82	6.15	5.35	2 60	3.00	00.9	5.13	6.28	
	CASES		4	16	17	17	24	24	38	16	18	19	17	(	2 2	24	14	0	01	12	11	13	17	U	0	6	15	360	
ac.	BABIES UMBER C	N	6	23	24	27	33	33	37	21	27	27	24	,	15	25	16	22	67	15	14	14	21	1	-	10	20	478	
	AGE BY MONTHS		-	2	3	4	25	9	10	- 00	0	10	=	,	12	C1 1	15	2	10	17	18	19	20		17	22	23	TOTAL	

# TABLE 43.—SUMMARY.

		DIED		6		0		-	=	27				-			4	7	7.
=	1	AT SE	Sick	N	3	:	6	-	10	28	12	2	-	26	2	88	Ξ	139	
		CONDI-	Well	36	14	12	39	9	134	241	9	:	1	S	-	18	31	272	:
		WK.	Ozs.	3.48	2.00	1.32	2.13	96.0	2.13	2.19	3.19	0.36	4.25	2.13	3.08	2.48	2.47	2.28	:
		Loss	Lbs.	:	:	:	:	:	:	:	:	;	:	:	:	:	:	:	
	SIGHT	Av. Lo	Ozs. I	11.37	7.29	7.17	7.50	6.57	10.49	9.64	10.72	2.00	1.00	9.16	12.33	10.90	10.48	9.93	
1	×	PER C	Lbs.		:	:	:	:	-:	:	:	:	-	:	:	:	:	:	:
	SHOWING LOSS IN WEIGHT		Ozs. L	15.07	1.88	6.25	13.42	4.86	13.19	10.79	14.22	4.50	11.00	8.90	3.00	10.38	12.46	0.31	
	ING	AV. WT.		16 15	3 1	17 6	16 13	18 4	16 13	16 10	4	16	14 11	14	19	14 10	14 17	16	-
1	SHOW		. Lbs.		-				7.68	- 11	8.94	6.50		2.06	5.33	5.28 1	6.94		
1	"	AV. WT. AT ENTRANCE	Ozs.	10.44	9.17	13.42	4.92	11.43	7.0	4.43	8.		12.00	2.0	_	S.	0.0	10.24	
		Av. Ent	Lbs.	17	13	17	17	18	17	17	15	16	15	15	19	15	15	16	
		ERVED	Av.	3.27	3.65	5.42	3.52	6.71	4.93	4.41	3.39	5.50	4.00	4.32	4.00	4.40	4.25	4.36	
		CASES	OB O	41	17	12	48	7	144	269	18	2	2	31	3	86	142	411	
=		AT AT	Sick	13	10	2	10	2	18	55	34	00	4	71	c.	128	248	303	442
		CONDI-	Well	611	136	71	361	32	806	2017	46	17	13	55	14	232	377	2394	3.21 2666 442
		WK.	Ozs.	4.86	3.29	3.40	3.81	3.11	3.29	3.77	4.80	3.40	3.37	3.64	3.69	3.85	3.87	3.80	3.21
		GAIN B PER	Lbs.	:	:	:	:5		:	:	:		:	:	:	:	:	:	
	WEIGHT	Av. G	Ozs.	9.35	5.01	4.68	10.58	8.62	5.84	7.69	8.19	4.52	7.82	7.23	9.18	8.16	7.85	7.72	3.27
	IN W	PER	Lbs.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-
	GAIN IN	AV. WT.	Ozs.	5.03	9.07	3.76	14.92	9.68	8.18	16.60	14.30	3.72	2.82	13.06	1.06	1.08	7.34	3.06	2.70
	SHOWING	AV.	Lbs.	14	14	16	15	17	18	16	1	14	==	14	16	16	15	16	16
	SHO	AV. WT. AT ENTRANCE	Ozs. I	11.68	4.06 14	15.08 16	4.34 15	1.06 17	2.34	14.91	6.13 14	15.20 14	9 11.00 11	5.83	7.88	8.92	15.49	11.34	5.43
		IV. W	Lbs.		13		14	16			13		9 1	13	14	14	13 1	141	41
-		ERVED	OBSI	.22	.38	73 6.08 14	86.	.59	.64	.28	.04	.04	90.						8.
		CASES WKS.		624 5.22 12	146 6.38 13	73 6	371 6.98 14	34 7.59 16	824 6.64 17	2369 2072 6.28 14	80 5.04 13	25 6.04 12	17 7.06	126 6.38	17 6.82	360 6.28	625 6.17	9 46	9 80
-		BABIES	UM	9 899	163 1	82	428 3	45	8 086	20 50	100	27	19	168	22	478 3	814 6	32 26	31
_	A	UMBER O	N		. 16				36	-	1				2000			318	318
		CONDITION AT ENTRANCE AND FEEDING		Well babies breast-fed.  No change in feeding	eding	Well babies breast-fed. Changed to artificial feeding	Well babies mixed fed.  No change in feeding	Well babies mixed fed. Changed to artificial feeding	Well babies artificially fed. No change in feeding	O TOTAL FOR WELL BABIES	Sick babies breast-fed.  No change in feeding	Sick babies breast-fed.  Changed to mixed feeding	Sick babies breast-fed.  Changed to artificial feeding	No change in feeding	Sick babies mixed fed.  Changed to artificial feeding	No change in feeding	TOTAL FOR SICK BABIES	TOTAL WELL AND SICK BABIES 3182 2697 6.25	TOTAL AND AVERAGES WELL AND SICK: Gain and loss

TABLE 44.—SUMMARY.—RESULTS OF FEEDING OF 3182 BABIES UNDER TWO YEARS OF AGE AS TO GAIN OR LOSS IN WEIGHT AND DEATH AT END OF PERIOD

1	CENT.	10	0	3	1	2	5	~	_	_	1	_		_		_	_	~	_						
	Рев	4.5	5.	3.	3.	=	3.	3,	2.	2.	2.7	0	:	0.0	•	=	2.1	1.2	1.4	:	:	:	:	:	2.3
OSE	DIED	10	16	1	∞	3	-	9	3	3	4	-	:	-	:	-	2	-	-	:	:	:	:	:	74
CONDITION AT CLOSE	PER CENT.	6.7	7.3	4.8	7.3	0.6	12.0	12.3	12.2	15.4	17.6	17.2	23.5	18.7	10.1	16.9	21.6	17.3	18.1	20.6	19.7	16.2	9.3	15.4	12.9
NDITION	Loss	15	23	10	16	17	23	23	18	23	26	26	27	20	26	15	21	14	13	13	16	9	4	16	411
	PER CENT.	88.8	87.7	6.16	0.68	89.4	84.4	84.5	85.7	82.5	7.67	82.1	76.5	80.4	83.9	82.0	76.3	81.5	80.5	19.4	80.3	84.8	7.06	84.6	84.8
ALL BABIES.	GAIN	198	278	193	195	169	162	158	126	123	118	124	88	98	135	73	74	99	28	20	65	31	39	88	2697
	TOTAL	223	317	210	219	189	192	187	147	149	148	151	115	107	191	68	97	81	72	63	81	37	43	104	3182
	PER CENT.	15.4	15.5	11.11	11.8	3.6	08	9.1	5.1	6.1	5.3	2.4	:	:	:	:	2.8	:	:	:	:			: :	5.8
TOSE	DIED	9	6	9	9	2	4	. 10	7	65	7	-	:	:	:	:	-	:	:	:	:			: :	47
CONDITION AT CLOSE	PER CENT.	7.7	12.1	11.1	15.7	18.2	12.0	14.5	21.2	18.4	18.4	28.6	28.0	23.3	12.2	15.4	22.2	23.8	26.1	22.2	17.4	25.0	0.1	21.4	17.4
CONDITI	Loss	3	7	9	00	10	9	000	1	0	1	12	7	7	2	4	00	S	9	4	4	,		9	142
SICK BABIES.	PER CENT.	76.9	72.4	77.8	72.5	78.2	0 08	76.4	72.7	75.5	76.3	0.69	72.0	7.97	87.8	34.6	75.0	76.2	73.9	77.8	82.6	75.0	000	78.6	76.8
SICK I	GAIN	30	42	42	37	43	40	42	24	37	29	59	18	23	36	22	27	16	17	14	19	4	,	22	625
	TOTAL	39	58	54	51	55	S	200	33	40	38	42	25	30	41	26	36	21	23	18	23	٥	-	28	814
	Рев. Свит.	2.2	2.7	9.0	1.2	8.0	,	1.7	0.0	:	. ~	:		1.3		1.6	1.7	1.7	2.0		:		:	: :	1:1
M	DIED	4	7	-	2	-			-	•		:	;	-		-	-	-	-		:		:	: :	27
CONDITION AT CLOSE	PER CENT.	6.5	6.2	2.6	4.7	5.2	120	11 3	0.6	14.0	17.3	12.8	22.2	16.9	17.5	17.5	21.3	150	14.3	20.0	20.7	120	0.01	13.2	11.4
DITION	Loss	12	16	4	00	7	1	1 2	1 :	14	101	14	20	13	21	11	13	0	1	6	12	,	+ 6	10	269
	PER CENT.	91.3	91.1	8.96	94.1	94.0	0 20	02.0	80.5	0.40	80.0	87.2	77.8	81.8	82.5	80.9	77.0	83.3	83.7	80.0	79.3	06.7	7.00	86.8	87.5
WELL BABIES.	GAIN	168	236	151	158	126	100	771	100	707	000	95	70	63	66	51	47	9	41	36	46	30	3 6	67 99	2072
WEI	TOTAL	184	250	156	168	134	140	147	114	100	110	109	06	77	120	63	19	9	40	45	58	00	67	76	2368
	AGE	-	,		4	2	,	0 1	- 0	0 0	10	= =	12	13	14	15	16	11	18	10	20		17	23	TOTAL
												71													

TABLE 45.—SUMMARY.—RESULTS OF FEEDING OF 3182 BABIES UNDER TWO YEARS OF AGE AS TO WELL, SICK, OR DEAD AT END OF PERIOD

1	PER CENT.	4.5	2.0	3.3	3.7	1.6	3.6	3.2	2.1	2.1	2.7	0.7		: ;	6.0	:	1.1	2.1	1.2	1.4		:		:	:	:	2.3
SE	Digo	10	16	7	00	3	7	9	3	3	4	-		:	-	:	-	2	1	1	:	:		:	:	:	74
CONDITION AT CLOSE	PER CENT.	8.1	12.3	13.3	8.6	13.8	13.1	16.6	12.2	19.4	9.5	13.9		12.7	16.8	14.9	28.1	19.6	13.6	19.4	14.3	10.1	18.0	10.7	9.3	15.4	13.9
NDITION	SICK	18	39	28	19	26	25	31	18	29	14	21	;	14	18	24	25	19	11	14	6	13	1		4	16	442
	PER CENT.	87.4	82.7	83.4	87.7	84.6	83.3	80.2	85.7	78.5	87.8	85.4	1	87.8	82.3	85.1	8.02	78.3	85.2	79.2	85.7	83.9	1 10	1.10	200	84.6	83.8
ALL BABIES,	WELL	195	262	175	192	160	160	150	126	1117	130	129		101	88	137	63	20	69	57	54	89	30	200	39	88	2666
V	TOTAL	223	317	210	219	189	192	187	147	149	148	151		115	107	191	68	97	81	72	63	81	27	10	43	104	3182
	PER CENT.	15.4	15.5	11.1	8.11	3.6	8.0	9.1	5.1	6.2	5.3	2.4		:	:	:	:	2.8	:	:	:	:			:	:	5.8
LOSE	DIED	9	6	9	9	2	4	N	2	3	2	-		:	:	:	:	-		:	:	:		:		:	47
CONDITION AT CLOSE	PER CENT.	30.8	43.1	42.6	33.3	41.8	46.0	47.3	46.4	46.9	28.9	42.8		40.0	20.0	46.4	73.1	50.0	47.6	52.2	33.3	52.2	75.0	0.01	18.2	50.0	44.1
CONDIT	SICK	12	25	23	17	23	23	26	15	23	11	18		10	15	19	19	- 82	10	12	9	12	4	0	2	14	359
SICK BABIES.	PER CENT.	53.8	41.4	46.3	54.9	54.6	46.0	43.6	48.5	46.9	65.8	54.8	4	0.09	20.0	53.6	26.9	47.2	52.4	47.8	66.7	8.74	0 20	0.07	81.8	50.0	50.1
SICK I	WELL	21	24	25	28	30	23	24	16	23	25	23	1	15	15	22	7	17	=	=	12	==	·	7	6	14	408
	ТотоТ	39	58	54	51	55	50	55	33	49	38	42	1	25	30	41	26	36	21	23	18	23	0	0	==	28	814
	PER CENT.	2.2	2.7	9.0	1.2	8.0	2.1	8.0	6.0	0.0	1.8	:		:	1.3	:	1.6	1.7	1.7	2.0	:	:		:	:	:	1.1
SE	DIED	4	7	-	2	-	60	-	-	:	2	:		:	-	:	-	-		-	:	:		:	:	:	27
CONDITION AT CLOSE	PER CENT.	3.2	5.4	3.2	1.2	2.7	1.4	3.7	2.6	0.9	2.7	2.8		4.5	3.9	4.2	9.5	1	1.7	4.1	6.7	1.7		0.0	6.3	2.6	3.5
NDITION	SICK	9	14	S	2	3	2	ı,	3	9	3	3		4	3	S	9	-	-	2	3	-	•	1	2	2	83
	PER CENT.	94.6	91.9	96.2	97.6	0.76	96.5	95.5	96.5	94.0	95.5	97.2		95.5	8.46	95.8	6.88	2 90	196	93.9	93.3	98.3	200	6.06	93.7	97.4	95.4
WELL BABIES.	WELL	174	238	150	164	130	137	126	110	94	105	106		98	73	115	56	50	000	46	42	57	00	07	30	74	2258
H	TOTAL	184	259	156	168	134	142	132	114	100	110	109		06	77	120	63	61	09	49	45	58	00	67	32	92	2368
	AGE	1	2	3	4	10	9	7	00	6	_	=======================================		12	13	14	15	16	17	18	19	20		17	22	23	TOTAL

It is also to be remembered, in studying these tables, that the period of obervation was during the hottest summer weather. The gain in weight is very triking. This is most marked among the breast-fed babies, as would be exected. It is also interesting to note the average weights of the babies under one 'ear. In the group over one year it will be noted that they were practically all very much under normal weight when admitted, and that the gains were coniderable.

Summary Tables 44 and 45 show the condition of babies at the close of the period of observation, grouped according to age, sick or well on admission, and otal, expressed in gross and percentage figures.

As would be expected, the results for babies well on admission are much better than for sick babies. In the former group, 87.5 per cent. showed gain in weight, as against 76.8 per cent. in the latter, the average being 84.8 per cent. Again, 95.4 per cent. of the well babies were well at the close, 3.5 per cent. were sick, and 1.1 per cent. had died.

Of the sick babies, 50.1 per cent. were well, 44.1 per cent. were still sick, and only 5.8 per cent. had died. That over-optimism did not make the "well" too nigh may be gathered from the relation of gain in weight and "well" among the sick babies.

A more detailed discussion of the deaths will be given later on in this chapter.

The following table shows the condition of babies on admission, grouped according to feeding history, and further divided into groups of babies under one year and over one year of age.

TABLE 46.—SUMMARY OF FEEDING HISTORIES AND CONDITION ON ADMISSION

FEEDING	Ui	NDER O	NE	(	OVER O	NE		TOTAL	
	Well	Sick	TOTAL	Well	Sick	Total	Well	Sick	Total
Breast to mixed	608 140 54	83 24 16	691 164 70	60 23 31	17 3 3	77 26 34	668 163 85	100 27 19	768 190 104
Total breast on admission	802	123	925	114	23	137	916	146	1062
Mixed	329 26	108	437 35	98 19	60 13	158 32	427 45	168 22	595 67
Total mixed on admission	355	117	472	117	73	190	472	190	662
Artificial	451	284	735	529	194	723	980	478	1458
Total	1608	524	2132	760	290	1050	2368	814	3182

Of 3182 babies at end of period-

2697, or 84.87 per cent., showed gain in weight.

2666, or 83.8 per cent., were well. 74, or 2.3 per cent., had died.

Of 3182 babies on admission-

2368, or 74.4 per cent., were well. 814, or 25.6 per cent., were sick.

Of 2368 babies well on admission, at end of period-

2072, or 87.5 per cent., showed gain in weight.

2258, or 95.4 per cent., were well. 27, or 1.1 per cent., had died.

Of 814 babies sick on admission, at end of period-

625, or 76.8 per cent., showed gain in weight.

408, or 50.1 per cent., were well. 47, or 5.8 per cent., had died.

TABLE 47.—NUMBER OF BABIES GROUPED ACCORDING TO FEEDING HISTORY, NUMBER SICK AND WELL ON ADMISSION, AND PERCENTAGES OF TOTAL ENROLMENT FOR EACH GROUP AND CONDITION

	BREA	ST-FED	TO N	ST-FED LIXED ED	то А	ST-FED RTIFI- Y FED		ED FED	то А	ED FED RTIFI- Y FED		TIFI- Y FED	Тот	ALS
	No. Per Cent.		No.	Per Cent.	No.	Per Cent.	No.	Per Cent.	No.	Per Cent.	No.	Per Cent.	No.	Per Cent.
Well	668 100	21.0 3.1	163 27	5.1 0.8	85 19	2.7 0.6	427 168	13.4 5.3		1.4 0.7	980 478		2368 814	74.4 25.6
TOTAL	768	24.1	190	5.9	104	3.3	595	18.7	67	2.1	1458	45.8	3182	99.9

TABLE 48.—FEEDING HISTORY AND PROPORTION OF EACH GROUP, SICK OR WELL ON ADMISSION

	BREA	ST-FED	TO M	ST-FED IXED ED	то А	ST-FED RTIFI- Y FED		ED FED	TO A	ED FED ARTIFI- LY FED		TIFI- Y FED	Ton	TALS
	No. Per Cent.		No.	Per Cent.	No.	Per Cent.	No.	Per Cent.	No.	Per Cent.	No.	Per Cent	No.	Per Cent.
Well	668 100	87.0 13.0	163 27	85.8 14.2			427 168		45 22	67.2 32.8	980 478	67.2 32.8	2368 814	74.4 25.6
TOTAL	768	100.0	190	100.0	104	100.0	595	100.0	67	100.0	1458	100.0	3182	100.0

Nearly one-quarter of the entire enrolment, therefore, were breast-fed throughout, and 30 per cent. were entirely or partially nursed. The ratio of sick to well increases from breast-fed to mixed fed to artificially fed.

# ABLE 49.—PROPORTION OF TOTAL BABIES BREAST-FED, MIXED FED, AND ARTIFICIALLY FED ON ADMISSION

Br	EAST	M	IXED	Art	IFICIAL		
No.	Per cent.	No.	Per cent.	No.	Per cent.		
1062	33.4	662	20.8	1458	45.8		

Of 1062 babies breast-fed on admission-

768, or 72.3 per cent., were breast-fed throughout; at end of period-704, or 91.7 per cent., showed gain in weight.

699, or 91.0 per cent., were well. 5, or 0.47 per cent., had died.

190, or 17.9 per cent., were changed to mixed feeding; at end of period— 171, or 90.0 per cent., showed gain in weight.

167, or 87.9 per cent., were well.

None had died.

104, or 9.8 per cent., were changed to artificial feeding; at end of period— 90, or 86.5 per cent., showed gain in weight.

97, or 93.3 per cent., were well.

None had died.

## Of 662 babies mixed fed on admission—

595, or 89.9 per cent., were mixed fed throughout; at end of period— 497, or 83.5 per cent., showed gain in weight.

460, or 77.3 per cent., were well. 19, or 3.2 per cent., had died.

67, or 10.1 per cent., were changed to artificial feeding; at end of period— 51, or 76.1 per cent., showed gain in weight.

53, or 79.1 per cent., were well. 6, or 9.0 per cent., had died.

Of 1458 babies artificially fed on admission; at end of period-

1184, or 81.2 per cent., showed gain in weight.

1190, or 81.6 per cent., were well. 44, or 3.0 per cent., had died.

## TABLE 50.—PROPORTION OF TOTAL BABIES WELL OR SICK ON ADMISSION AC-CORDING TO FEEDING ON ADMISSION

	BREA	AST-FED	Mn	KED FED	ARTIFICIALLY FED			
	No.	Per Cent.	No.	Per Cent.	No.	Per Cent.		
Well	. 916 86.3 146 13.7		472 190	71.3 28.7	980 478	67.2 32.8		
[OTAL	1062	100.0	662	100.0	1458	100.0		

# Of 916 babies well and breast-fed on admission-

668, or 72.9 per cent., were breast-fed throughout; at end of period-624, or 93.4 per cent., showed gain in weight.

647, or 96.8 per cent., were well.

3, or 0.45 per cent., had died.

163, or 17.8 per cent., were changed to mixed feeding; at end of period—146, or 89.6 per cent., showed gain in weight.

150, or 92.0 per cent., were well.

None had died.

85, or 9.3 per cent., were changed to artificial feeding; at end of period—73, or 85.0 per cent., showed gain in weight.
83, or 97.6 per cent., were well.

None had died.

Of 472 babies well and mixed fed on admission-

427, or 90.5 per cent., were mixed fed throughout; at end of period—371, or 86.9 per cent., showed gain in weight.

8, or 1.9 per cent., were well.

45, or 9.5 per cent., were changed to artificial feeding; at end of period—34, or 75.5 per cent., showed gain in weight.

38, or 84.4 per cent., were well. 4, or 9.1 per cent., had died.

Of 980 babies well on admission and artificially fed throughout; at end of period—

824, or 84.1 per cent., showed gain in weight.

940, or 95.9 per cent., were well. 12, or 1.2 per cent., had died.

Of 146 babies sick and breast-fed on admission-

100, or 68.5 per cent., were breast-fed throughout; at end of period—80, or 80 per cent., showed gain in weight.

52, or 52 per cent., were well. 2, or 2 per cent., had died.

27, or 18.5 per cent., were changed to mixed feeding; at end of period—25, or 92.6 per cent., showed gain in weight.

17, or 63.0 per cent., were well.

None had died.

19, or 13 per cent., were changed to artificial feeding; at end of period—
17, or 89.5 per cent., showed gain in weight.

14, or 73.7 per cent., were well.

None had died.

Of 190 babies sick and mixed fed on admission-

168, or 88.4 per cent., were mixed fed throughout; at end of period—126, or 75.0 per cent., showed gain in weight.

60, or 35.7 per cent., were well. 11, or 6.6 per cent., had died.

22, or 11.6 per cent., were changed to artificial feeding; at end of period—
17, or 77.3 per cent., showed gain in weight.

15, or 68.2 per cent., were well. 2, or 9.1 per cent., had died.

Of 478 babies sick on admission and artificially fed throughout; at end of period—

360, or 75.3 per cent., showed gain in weight.

250, or 52.3 per cent., were well. 32, or 6.7 per cent., had died.

The above figures are for babies under two years of age. It may be well to study the results of the feeding in the 2132 babies under one year of age.

Of 2132 babies under one year of age, at end of period-

1844, or 86.5 per cent., showed gain in weight.

1796, or 84.2 per cent., were well. 68, or 3.2 per cent., had died.

Of 2132 babies on admission-

1608, or 75.4 per cent., were well. 524, or 24.6 per cent., were sick.

Of 1608 babies well on admission, at end of period-

1449, or 90.1 per cent., showed gain in weight.

1534, or 95.4 per cent., were well. 22, or 1.4 per cent., had died.

Of 524 babies sick on admission, at end of period-

395, or 75.4 per cent., showed gain in weight.

262, or 50 per cent., were well. 46, or 8.8 per cent., had died.

# TABLE 51.—PROPORTION OF TOTAL BABIES UNDER ONE YEAR WHO WERE BREAST-FED, MIXED FED, OR ARTIFICIALLY FED ON ADMISSION

BREAS	I-FED	Mi	KED FED	ARTIFICIALLY FED				
No.	Per cent.	No.	Per cent.	No.	Per cent.			
925	43.4	472	22.1	735	34.5			

## Of 925 babies breast-fed on admission-

691, or 74.7 per cent., were breast-fed throughout; at end of period—648, or 93.8 per cent., showed gain in weight.

634, or 91.8 per cent., were well. 5, or 0.72 per cent., had died.

164, or 17.7 per cent., were changed to mixed feeding; at end of period—148, or 90.2 per cent., showed gain in weight.

144, or 87.8 per cent., were well.

None had died.

70, or 7.6 per cent., were changed to artificial feeding; at end of period—64, or 91.4 per cent., showed gain in weight.
63, or 90.0 per cent., were well.

None had died.

#### Of 472 babies mixed fed on admission—

437, or 92.6 per cent., were mixed fed throughout; at end of period—375, or 85.8 per cent., showed gain in weight.

353, or 80.8 per cent., were well. 18, or 4.1 per cent., had died.

35, or 7.4 per cent., were changed to artificial feeding; at end of period—27, or 77.1 per cent., showed gain in weight.

27, or 77.1 per cent., were well. 6, or 17.1 per cent., had died.

Of 735 babies artificially fed on admission, at end of period-

582, or 79.2 per cent., showed gain in weight.

575, or 78.2 per cent., were well. 39, or 5.3 per cent., had died.

# TABLE 52.—PROPORTION OF BABIES UNDER ONE YEAR WHO WERE WELL OR SICK ON ADMISSION ACCORDING TO FEEDING ON ADMISSION

В	REAST-FED		Mix	ED FED	ARTIFICIALLY FED				
	No.	Per cent.	No.	Per Cent.	No.	Per cent.			
Well	802 123	86.7 13.3	355 117	75.2 24.8	541 284	61 4 38.6			
TOTAL	925	100.0	472	100.0	735	100.0			

Of 802 babies well and breast-fed on admission-

608, or 75.8 per cent., were breast-fed throughout; at end of period-577, or 94.9 per cent., showed gain in weight. 591, or 97.4 per cent., were well.

3, or 0.49 per cent., had died.

140, or 17.5 per cent., were changed to mixed feeding; at end of period— 126, or 90.0 per cent., showed gain in weight.

129, or 92.1 per cent., were well.

None had died.

54, or 6.7 per cent., were changed to artificial feeding; at end of period— 50, or 92.6 per cent., showed gain in weight. 52, or 96.3 per cent., were well. None had died.

Of 355 babies well and mixed fed on admission—

329, or 92.7 per cent., were mixed fed throughout; at end of period— 294, or 89.4 per cent., showed gain in weight.

308, or 93.6 per cent., were well. 7, or 2.1 per cent., had died.

26, or 7.3 per cent., were changed to artificial feeding; at end of period-20, or 76.9 per cent., showed gain in weight.

20, or 76.9 per cent., were well. 4, or 15.4 per cent., had died.

Of 451 babies well and artificially fed on admission; at end of period-

382, or 84.7 per cent., showed gain in weight.

434, or 96.2 per cent., were well. 8, or 1.8 per cent., had died.

Of 123 babies sick and breast-fed on admission-

83, or 67.5 per cent., were breast-fed throughout; at end of period-

71, or 85.5 per cent., showed gain in weight.

43, or 51.8 per cent., were well. 2, or 2.4 per cent., had died.

24, or 19.5 per cent., were changed to mixed feeding; at end of period-22, or 19.7 per cent., showed gain in weight.

15, or 62.5 per cent., were well.

None had died.

16, or 13 per cent., were changed to artificial feeding; at end of period-14, or 87.5 per cent., showed gain in weight. 11, or 68.7 per cent., were well.

None had died.

Of 117 babies sick and mixed fed on admission-

108, or 92.3 per cent., were mixed fed throughout; at end of period—81, or 75.0 per cent., showed gain in weight.

45, or 41.7 per cent., were well. 11, or 10.2 per cent., had died.

9, or 7.6 per cent., were changed to artificial feeding; at end of period—7, or 77.8 per cent., showed gain in weight.

7, or 77.8 per cent., were well. 2, or 22.2 per cent., had died.

Of the 284 babies sick and artificially fed on admission, at end of period-

200, or 70.4 per cent., showed gain in weight.

141, or 49.6 per cent., were well.

31, or 10.9 per cent., had died.

The above figures prove pretty conclusively that the home modification of nilk is possible among the very poor and ignorant, and that the results, when this seding is carefully ordered and careful supervision insures the instructions being arried out, are as good as when modified milk is dispensed.

As has been already stated, every fatal case which was seen once by the hysician or nurse at the station has been charged against that station. This obviously unfair, for in practically every case the mother was referred to a hysician or advised to take the child to a hospital, arrangements usually being tade at once by the nurse, and a card given to the mother. However, in order hat no charge of selecting cases could be made, all were credited to the stations, and the general mortality rates for each station and for the stations as a whole aclude these deaths.

Of the 158\* deaths occurring among the station babies during the entire eriod June to November, the method of feeding, condition on admission, disosition of the case, place of death, average interval elapsing between last visit nd date of death, and age at death, are shown in the following detailed tables, rouped according to diagnosis and number of clinics attended.

In studying these tables it must be remembered that the diagnoses are those iven on the death-certificates by the reporting physician. It will be noted that iarrheal disease is credited with 79, or exactly 50 per cent. of the total deaths; nalnutrition, 21, or 13.3 per cent.; respiratory diseases, 17, or 10.8 per cent.; ontagious diseases, 14, or 8.9 per cent.; all other diseases, 27, or 17 per cent. In his last group there are included:

Diseases of cerebrospinal system	
Congenital debility	
Tuberculosis	
Congenital syphilis 1	
Mastoiditis 1	
"Choked to death" 1	
Unknown 5	
Total	

<sup>\*</sup> The deaths given in connection with the feeding tables are for the period June 1st to Sepember 15th.

TABLE 53.—DATA CONCERNING 158 FATAL CASES

	12-24	7000	2	0001	1	1001	2	-0	3	6211	7
HIS	6-12	9 4 1 1 2	27	0014	5	2015	11	420%	6	2005	7
AGE IN MONTHS	3-6	12 12 3	29	5500	4	1501	4	0000	0	8004	7
AGE	1-3	r-400	17	1335	6	0000	0	0000	0	8101	5
	Under One Month	0000	4	7000	2	0000	0	4400	2	4000	1
AVERAGE IN-	TERVAL IN DAYS	9.86 13.83 27.20 15.41	14.50	3.75 6.00 12.50 23.12	13.95	20.40 0.0 16.30 21.50	20.20	25.10 15.30 1.00 14.50	18.20	20.90 21.00 25.00 6.60	15.20
AT	Hosp,	10 5 6 12	33	0084	7	7100	3	7017	5	4116	6
DIED	Ноше	13 10 10	46	4004	14	2507	14	4602	6	8008	18
RE-	ATTEND	2139	12	0011	2	0100	3	0000	4	1017	4
UED TO	Hosp.	10 7 12	34	00%%	9	0000	4	7017	4	2112	6
REFERRED	Doctor	13 10 8 8	33	4004	13	2002	10	2013	9	810%	14
ADMISSION	Sick	26 13 7 19	65	4608	21	2024	11	1050	6	11808	19
ON AD	Well	0 10 m m	14	0000	0	3102	9	0110	5	1019	00
	Artifi- cial	23 9 3 15	50	2045	11	000	6	4102	7	0215	17
FEEDING	Mixed	N 7- 4 6	19	1535	8	0014	S	1001	S	0107	3
	Breast	-284	10	7000	2	1011	3	0011	2	£004	7
NUM-	BER	29 10 22	79	4000	21	73007	17	9814	14	11 32 11	27
		DIARRHEAL 1 clinic 2 clinics 3 clinics 4 or more clinics	TOTAL	MALNUTRITION 1 clinic 2 clinics 3 clinics 4 or more clinics	TOTAL	RESPIRATORY 1 clinic 2 clinics 3 clinics 4 or more clinics	TOTAL	Contagious 1 clinic	TOTAL	OTHER CAUSES 1 clinic 2 clinics 3 clinics	TOTAL

	r004	15	79090	15
	17 7 6 29	59	27 11 9	59
	91 - 7 - 41	44	29 4 4 0 7	44
	12 8 5 6 8	31	17 0 0 0 0 0 0	31
	9800	6	40001	6
	14.50 13.90 20.10 15.50	15.50	14.50 13.95 20.20 18.20 15.20	0.0
	23 25 6 15	57	33 7 8 9 9	57
	23 29 29	101	46 14 14 9 18	101
	12 6 8 4	25	22 8 4 4	25
IMAKX	13 6 14 24	57	34 0 4 9 9	57
SOF	33 15 4 24	92	33 13 10 6 14	92
	52 21 15 37	125	65 21 11 9 19	125
	6 6 15	33	41 0 0 8 8	33
	42° 62	94	50 111 9 7	94
	13 11 11	40	19 8 3 3	40
	5 12 12	24	10 2 2 2 7	24
	58 27 21 52	158	79 17 14 27	158
	ALL CAUSES 1 clinic	TOTAL	ALL CLINICS Diarrheal Malnutrition Respiratory Contagious All others	TOTAL

It is possible that, as only four died in the hospital, some of the "cerebrospinal diseases" may have been diarrheal in nature.

The above figures give a death-rate from the different groups of causes as follows, based on the enrolment of 7919 babies:

# TABLE 54.—DEATH-RATE FOR EACH GROUP OF DISEASES

Diarrhe	al			 							 			. :	8.0	per	thousa	nd
Malnut	rition	١		 			 								2.6			
Respira	tory.			 			 								2.1		"	
Contag	ious.			 			 		٠						1.8		"	
Other c	auses			 			 						٠		3.4			

In order to get a really accurate view of what was accomplished it is necessary to deduct the deaths of babies seen only once and referred to hospitals or physicians, numbering 58, or 26.7 per cent. This leaves exactly 100 deaths to be credited to the stations.

We, therefore, have 100 deaths among 7861 babies (deducting 58 from enrolment totals under two years of age), a death-rate of 1.27 per thousand.

There were charged against the stations 145 deaths among babies under one year, and 13 deaths among babies between one and two years. Of these cases there attended only one clinic, or were seen but once, 51 under one year and 7 over one year and under two years.

Deducting these figures from the deaths and enrolment at the different age periods we have—

94 deaths among 5328 babies under one year, or 17.6 per thousand.

6 deaths among 2533 babies over one year, and under two, or 2.4 per thousand. The deaths under one year in Manhattan for the same months, based on the births of the preceding twelve months, were 52.5 per thousand—almost exactly three times as many.

If, in addition to the deaths among babies attending only one clinic, we exclude everything except diarrheal and malnutrition deaths, we shall have the startling total of 66 deaths among 5287 babies, or 12.5 per thousand. This is, of course, putting the matter in its most favorable light, and is mentioned only as an interesting statement.

It is of interest to study the percentage of the deaths in each group in relation to the number of clinics attended:

TABLE 55.—PERCENTAGE OF DEATHS IN RELATION TO DIAGNOSIS AND NUM-BER OF CLINICS ATTENDED

	1 CLINIC	2 CLINICS	3 CLINICS	4 OR MORE
Diarrheal	36.7 19.0 41.2 42.9 44.4	22.8 14.3 0.0 21.4 11.1	12.7 28.6 17.6 7.1 3.7	27.8 38.1 41.2 28.6 40.7
	36.7	17.1	13.3	32.9

The interval elapsing between the time when the baby passed from under the observation of the station to date of death is shown by the following table:

TABLE 56.—PERCENTAGE OF DEATHS IN RELATION TO NUMBER OF CLINICS AND INTERVAL BETWEEN LAST ATTENDANCE AND DATE OF DEATH

DAYS ELAPSING	1-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80	81-90	TOTAL
One clinic	48.0	24.0	17.0	2.0	3.0	4.0	2.0	0.0	0.0	100.0%

From the above the following conclusions seem to be warranted:

First: That milk stations did have a distinct influence in diminishing the mortality among babies in the districts in which they were situated.

Second: That the milk stations did not indirectly encourage artificial feeding—32.4 per cent. of all babies under one year of age were entirely breast-fed throughout; 60.6 per cent. were partly breast-fed.

*Third:* That home modification, even among the very poor and ignorant, is possible.

Fourth: That the results, as shown by the mortality and by the condition of the surviving babies at the end of the period of demonstration, prove that as good results can be obtained as when already modified milk is distributed.

#### CHAPTER VI

# Campaigns in Other Cities

While the figures that have been given for New York prove conclusively that there was a greater reduction in infant mortality in 1911 than in any previous year; that the weather conditions were not such as to explain this low mortality, especially in the early summer; and that the diminution in Manhattan and Brooklyn, where a very vigorous campaign was carried out, exceeded that of the rest of the city, the question arises, Was the mortality in New York reduced more than in other large cities?

With this question to answer, a careful tabulation of mortality statistics has been made for the nine largest cities in the country after New York. Study was also made of the weather conditions, and, as far as data could be secured from published records and articles, and in many cases from personal letters from health officers and lay and medical workers in those cities, a comparison was made of the

extent and efficiency of the campaign against infant mortality.

It must be admitted that most of the efforts at reducing infant mortality in this country have been carried on spasmodically, being almost always confined entirely to the hot summer months. It is, therefore, these three hot months, generally July, August, and September, that we shall discuss most fully, although the figures for the entire year also give important information as showing the effects of the educational campaign carried on. The various cities will be discussed in the order of their population.

#### CHICAGO

The details of the mortality under one year of age in Chicago are shown in Table 57.

If we analyze these figures, we find that the total deaths in Chicago under one year rise steadily from 1906 to 1908, and that the percentage of total deaths at all ages occurring during the first year of life rises from 21.05 in 1906 to 22.67 in 1908. In 1909 there was a diminution of 504 in the infant deaths for the year, the proportion of total deaths being 20.4. In 1910 the total rose almost to the figure of 1908, but the percentage of the total mortality remained about the same. In 1911 the gross total reached the lowest figure since 1906, although it was only 75 below 1909.

A similar condition is noted in the deaths occurring during the three summer months. Again 1909 was a low year, 1911 being 60 less than 1909, and the proportion of the year's deaths under one year of age occurring during these months is smaller than in any of the five preceding years. The percentage of total deaths



# AROUSING GENERAL INTEREST IN





EXHIBIT 17


IMPORTANT FACTS BROUGHT TO THE ATTENTION OF

# THE PROBLEM OF SAVING BABIES



Ехнівіт 17

MAYORS, HEALTH OFFICERS, AND CIVIC ORGANIZATIONS



at all ages is in excess of 1910, but less than any of the other four years. The Assistant Registrar of Vital Statistics of Chicago states that about one-third of the births are reported, thanks to an inefficient ordinance, so that infant mortality rates cannot be compared.

TABLE 57.-MORTALITY UNDER ONE YEAR IN CHICAGO

	1906	1907	1908	1909	1910	AVERAGE FIVE YEARS	1911
January	484	568	606	588	521	553	576
February	454	579	698	537	497	553	506
March	504	611	627	557	663	592	543
April	494	546	547	480	524	518	495
May	497	536	466	490	526	503	540
June	344	404	389	407	478	404	393
July	585	487	671	435	811	597	614
August	749	967	863	830	758	833	739
September	589	676	620	699	640	644	551
October	480	457	525	510	504	495	434
November	414	390	425	399	445	414	421
December	520	478	451	452	474	475	497
YEAR Per cent. total mortality	6114 21.05	6699 <b>20.81</b>	6888 22.67	6384 <b>20.40</b>	6841 <b>20.58</b>	<b>6581</b> 21.08	6309 <b>19.31</b>
July, August, and September	1923	2130	2154	1964	2209	2074	1904
Per cent. year's infant mortality .	31.45	31.79	31.27	30.76	32.29	31.52	30.18
Per cent. of total deaths	26.64	27.81	29.23	26.13	23.82	27.55	24.69

For the year 1911, then, in relation to 1910, there was a reduction of 7.7 per cent. of actual deaths and a reduction of 5.8 per cent. in the percentage of total deaths. Of deaths for July, August, and September, there was a reduction of 13.8 per cent. in actual deaths, but an increase of 3.3 per cent. in the proportion of total deaths. The proportion of the infant mortality for the year occurring during the summer months is less by 6.5 per cent.

If, now, we estimate what the mortality would have been in the years following 1906 if the infant deaths had increased at the same rate as the population (2.87 per cent. yearly), and then compare the actual mortality in these years, we find the conditions which are set forth in the following table:

TABLE 58.—MORTALITY WHICH SHOULD HAVE OCCURRED ON BASIS OF 1906 FIGURES, ALLOWING FOR INCREASE IN POPULATION, COMPARED WITH ACTUAL MORTALITY, IN CHICAGO

	1907	1908	1909	1910	1911
Estimated mortality Actual mortality Difference	6289 6699 +410	6464 6888 +424	6640 6384 —256	6815 6841 +26	6991 6309 —682
Percentage difference	+6.5%	+6.5%	-3.8%	+0.3%	-10.1%

The deaths in each year from 1907 on were in excess of this estimated rate by from 0.3 to 6.5 per cent., except in 1909, when they were 3.8 per cent. below the estimated figure. In 1911 there was nearly three times as great a reduction from the estimate as in 1909.

The proportion of deaths occurring during the third trimester is the lowest of any of the cities considered except Manhattan and Buffalo. It may be of interest to tabulate these figures here for reference.

TABLE 59.—DEATHS UNDER ONE YEAR BY TRIMESTERS, CHICAGO

	1906	1907	1908	1909	1910	Average 1906-10	1911
First quarter: Actual	1442	1758	1931	1682	1681	1698	1625
	23.6	26.2	28.0	26.3	24.6	25.8	25.8
	1335	1486	1402	1377	1528	1425	1428
	21.8	22.2	20.4	21.6	22.3	21.7	22.6
	1923	2130	2154	1964	2209	2074	1904
	31.5	31.8	31.3	30.8	32.3	31.5	30.2
	1414	1325	1401	1361	1423	1384	1352
	23.1	19.8	20.3	21.3	20.8	21.0	21.4

### WEATHER CONDITIONS

The weather conditions in Chicago during the last three years may be compared in the following tables:

TABLE 60.—WEATHER CONDITIONS: NUMBER OF DAYS MAXIMUM AND MEAN TEMPERATURES REACHED OR EXCEEDED CERTAIN FIGURES, CHICAGO\*

17	.,,

1000				
MAXIMUM TEMPERATURE WAS-	-		MEAN TEMPER	ATURE WAS-
95° or -	+ 90° or +	85° or +	85° or +	80° or +
June0	0	5	0	1
July0	3	11	0	2
August	2	8	0	4
September 0	0	2	0	1
_			_	_
1910	5	26	0	8
June0	3	7	0	6
July2	6	13	0	9
August	0	10	0	4
September0	0	0	0	0
_	_	-		
1911	9	30	0	19
June	6	9	2	5
July	9	15	6	10
August0	4	7	0	4
September0	0	1	0	0
_				
8	19	32	8	19

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Chicago, and Monthly Weather Review, Washington.

TABLE 61.—MONTHLY MEAN TEMPERATURE, PRECIPITATION, AND NUMBER OF RAINY DAYS, CHICAGO\*

	Average 39 Years	1909	1910	1911
June				
Mean	66.4°	67°	68°	72°
Rainfall	3.56 in.	5.09 in.	0.91 in.	2.54 in.
Rainy days		13+5	4+1	10+2
July				200000000000000000000000000000000000000
Mean	72°	72°	76°	76°
Rainfall	3.57	1.77	1.79	2.65
Rainy days		8+6	8+5	11+2
August				
Mean	71°	75°	73°	72°
Rainfall	3.10	6.20	3.08	3.72
Rainy days		10+3	6+5	12+2
September		2010		
Mean	65°	64°	65°	67°
Rainfall	3.07	3.60	3.90	4.03
Rainy days		8+1	9+4	16+4
Total		012		
Mean	68.6°	69.5°	70.5°	71.7°
Rainfall	13.30	16.66	9.68	12.94
Rainy days	10.00	39+15	27+15	49+10

Note.—Two figures are given for number of rainy days—the first, in front of the plus sign, is the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is number of days there was a trace of precipitation, but less than 0.01 inch.

If we study these tables, we find that the maximum temperature for the four months, taken as a whole, reached 85°, 90°, and 95° on more days than in either 1909 and 1910. The thermometer never reached 95° in 1909, and on only five days did it reach 90°. In 1910 it registered 95° or + on only two days, and 90° on only nine days, as against eight and nineteen days respectively in 1911. Similarly, the mean temperature never reached 85° in either 1909 or 1910, while it did so on eight days in 1911.

Only the month of August fell below 1909 and 1910 as to mean temperature for the whole month. The average for four months was decidedly higher than either year, and 3° higher than the average for thirty-nine years. In each month, however, the total rainfall was higher than in 1910, though it was above the thirty-nine-year average only in August and September. In July and September it exceeded the figures for 1909. For the four months the rainfall was well above that of 1910 and well below that of 1909. In the number of rainy days 1911 is largely in excess of either year.

The month of July is particularly interesting. The mean temperature for the month was 4° higher in 1911 than in 1909, and the deaths were 170 more than in 1909. There was the same mean for 1910 and 1911, and the deaths in 1911 were 197 less than in 1910. Also the deaths in 1910 were 376 more (86 per cent.) than in 1909, as against 170 more (39 per cent.) in 1911.

In 1910 and 1911 August was 2° and 3° below 1909, and the deaths were 72

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Chicago, and Monthly Weather Review, Washington.

and 91 respectively below those of 1909. In September, on the other hand, 1910 was 1° and 1911 was 3° above 1909, and yet the deaths were respectively 59 and 148 below those of 1909.

In July the rainfall was practically the same for 1909 and 1910, and 0.86 inch more in 1911. August showed less rainfall by 3.12 and 2.48 inches in 1910 and 1911 than in 1909, in both of the latter years the deaths being less. In September the rainfall was nearly the same, +0.30 inch in 1910 and +0.43 inch in 1911.

#### THE CAMPAIGN

For a number of years Chicago had been getting ready for an organized effort for combating infant mortality. The United Charities, then consisting of two separate organizations, attempted to get mothers and babies out of the city, to care for the sick babies, and to educate the mothers in the city itself by means of public tents, in charge of a trained nurse and visited by a physician. The Children's Hospital Society established a Milk Commission to provide a modified and pasteurized milk for babies at a reasonable price. Milk was dispensed already modified to set formulæ.


In 1908 the Health Department first entered the field, putting 100 physicians at work visiting from house to house, instructing the mothers, during July and August. The work was not organized, and very little came of it; but a more comprehensive plan was worked out during the ensuing winter.

In 1909 the United Charities, the Health Department, and the Visiting Nurses' Association coöperated. The Health Department mapped out the areas of the highest death-rate, furnished 20 nurses and 14 doctors, and spent \$10,000 on the campaign. The United Charities furnished nurses, tents, and workers. The Visiting Nurses' Association supervised the tents, furnished nurses in the tents, and worked in the field.

The general plan was a house-to-house canvass by nurses in the most crowded and unsanitary districts. Wherever a small child was found, instruction in feeding and care was given. At first the Health Department confined its activities to locating and referring cases, the actual care of the sick babies being turned over to the Visiting Nurses' Association, the settlements, and the United Charities. During July so much work had to be done that the Health Department had to take up part of the work of caring for the sick. Careful records were kept of all the cases, and information was acquired which was tabulated during the following months for use in the campaign of the next year.

It will be noted that 1909 was the first year with a lowered mortality since 1906. The percentage of the total mortality was also low.

Armed with the information obtained in 1909, the work was more active in 1910 and better organized. The activities of the Health Department were confined to one entire ward and parts of three others, as far as house-to-house visiting was concerned; and in 13 other wards they visited the homes of every baby that could be discovered from the records of the United Charities or the County Clerk's register of births. Ten public tents were located in centers having the highest





Garbage and Waste Forty-eight Hours in the Street; One of the Handicaps of a Summer Baby-saving Campaign



death-rates. These were maintained by the United Charities. Dr. Caroline Hedger spent her whole time in the work. An Infant Welfare Society was formed, consisting of representatives of all the various charities in Chicago.

During the summer the mothers of 23,984 babies were instructed. Ice tickets were distributed, and relief was provided through the United Charities, Jewish Aid, etc. The nurses were on the lookout for violations of the Sanitary Code, and 500 complaints were made.

The Civic Federation placarded the city fences and walls with a telling poster, "Don't Kill your Baby!"—with information how not to, and this was most effective. The well-known Bulletins of the Health Department featured the campaign, and its "Health Grams" and striking cartoons went far and wide.

In 1911 there was much greater coöperation among the nurses of the Department and other organizations. The work of the Health Department became more definite, and in every way the campaign was more efficient.

The Infant Welfare Society maintained 10 milk stations, open all the year, each in charge of a nurse. Volunteer physicians carried on the consultations, of which 806 were held. Two thousand one hundred and twenty-nine babies attended these conferences. Home modification was taught "when possible"; when impossible, modified milk was dispensed. Six hundred and twenty mothers were taught. Certified milk only was used.\*

It will be noted that there was an increase in mortality in 1910 over 1909 nearly up to the figures of 1908, although this is only 0.3 per cent. greater than the mortality estimated on the 1906 figures. The work of the nurses, while more efficient, was not nearly so wide-spread as in 1909. In 1911, due to increased experience and better coöperation, a great diminution was again obtained, especially marked during the summer months, in spite of unfavorable weather conditions. Dr. Hedger, who has had charge of the work in large part during the last three years, says: "The campaign should be a year-long affair; a few weeks of instruction in the summer is not sufficient. It must be line upon line and precept upon precept, year in and year out."† Chicago realizes the importance of continuous instruction to obtain permanent results, and not an ineffectual, short campaign.

Since 1910 Chicago has been attempting to enforce the laws pertaining to a safe milk supply. Seventy-five per cent. of the city's milk is pasteurized. All the rest must be from tuberculin-tested herds. As thorough an inspection of pasteurization plants is carried out as available funds will permit. There is also inspection of dealers and depots, platform inspection, and a certain amount of country farm, creamery, and bottling plant inspection. It is interesting to note that, excepting that provided to restaurants, 95 per cent. of the city's milk is bottled.

<sup>\*</sup> The above facts are obtained from the report of the Department of Health of the City of Chicago for the years 1907–1910, and from personal letters. In 1912 we are informed that practically all modification is done at the home. Through coöperation with certain milk dealers certified milk is delivered direct to the homes at 10 cents a quart.

<sup>†</sup> Report of the Campaign Against Summer Diarrhea, Chicago, 1909, etc.

# PHILADELPHIA

We have been unable to obtain reports or records of infant mortality in Philadelphia by months prior to 1909. The following table gives the figures as far as obtainable:\*

	1906	1907	1908	1909	1910	1911
January				387	398	286
February				438	404	289
March				433	497	427
April				330	349	332
May				289	313	344
June				396	540	401
July				659 490	799 552	587 556
August				496	426	394
September				364	315	313
October				238	284	330
December			::	350	355	357
Total Per cent. total mortality	5781 21.1	5508 <b>20.0</b>	5291 <b>20.1</b>	4870 <b>19.4</b>	5232 <b>19.3</b>	4616 17.6
July, August, and September Per cent. year's infant mortality Per cent. total mortality	<b>5</b> ::	::		1645 33.7 28.4	1777 33.9 27.9	1537 33.2 25.0
BIRTHSINFANT MORTALITY	34,771 166.2	35,142 136.7	36,641 144.1	37,540 <b>129.7</b>	38,676 135.3	41,000† 112.5†

In 1911, therefore, Philadelphia had less deaths by 254 than in 1909, the low record of the five-year period 1906–1910; 1165 less than in 1906, the high year; and 616 less than in 1910. The proportion of the total mortality for the year is 17.6 per cent., a fall of 1.8 from 1909; 3.5 from 1906; and 1.7 from 1910.

The summer infant mortality was practically the same proportion of the year's infant mortality as in 1909 and 1910; but of the total deaths it was 3.4 per cent. and 2.9 per cent. lower.

The infant mortality rate also shows a marked fall, although it is only estimated. There is a reduction of 22.8 and 17.2 per thousand from 1909 and 1910 respectively. In tabular form this appears as follows:

TABLE 63.—DIFFERENCE BETWEEN MORTALITY UNDER ONE YEAR IN 1911 AND OTHER YEARS

	ACTUAL D	IFFERENCE	PERCENTAGE DIFFERENCE		
Year's mortality	1909	1910	1909	1910	
	-254	616	- 5.2	-11.7	
	-1.8	1.7	- <b>9.2</b>	- 8.8	
	-108	240	- 6.5	-13.5	
	-0.5	0.7	- 1.4	- 2.0	
	-3.4	2.9	- <b>11.9</b>	-10.3	

<sup>\*</sup> From figures supplied by Dr. S. W. Newmeyer, Philadelphia.

In 1903 there were 4597 deaths under one year of age in Philadelphia, which was 17.7 per cent. of the total mortality. This is the low record year of the decade in both respects. The following table shows the deaths which would have occurred had they increased in proportion to the population increase, and the actual and percentage differences from these figures:

TABLE 64.—MORTALITY UNDER ONE YEAR, WHICH SHOULD HAVE OCCURRED IN DIFFERENT YEARS, BASED ON 1903 FIGURES, COMPARED WITH ACTUAL MORTALITY, PHILADELPHIA

	1906	1907	1908	1909	1910	1911
Estimated deaths	+ 913	4959 5508 + 549 + <b>11.0</b>	5049 5291 +242 + <b>4.7</b>	5140 4870 —270 — <b>5.2</b>	5230 5232 + 2 + <b>0.0</b>	5321 4616 — 705 — <b>13.2</b>

There is, therefore, a very great reduction in 1911 from every point of view.

#### WEATHER CONDITIONS

The weather conditions during the last three years are set forth in the following tables:

TABLE 65.—NUMBER OF DAYS ON WHICH MAXIMUM AND MEAN TEMPERA-TURES REACHED CERTAIN FIGURES, PHILADELPHIA\*

	MA	XIMUM TEMPERATU	MEAN TEMPERATURE		
	95° or +	90° or +	85° or +	85° or +	80° or +
909 June	0	8	10	1	8
July	0 1 1 0	8 5 3 0	17 13	1	8 8 3 0
September	Ô	0	0	Ō	0
	2	16	40	3	19
910					
Tune	1 0	5	8	1 1	5 17 0 2
July	1 0 0 0	5 10 0 1	8 19 5 3	0	0
September	_	_	_	_	_
	1	15	35	2	24
911					
June July August	0	2	9	0	12
August	0 5 0	10 3 0	19 13 2	6 0 0	3 12 5 0
September	_	_	_	_	_
	5	15	43	6	20

<sup>\*</sup> Monthly Meteorological Summary, Philadelphia; and Monthly Weather Review, Washington, D. C.

TABLE 66.—MONTHLY MEAN TEMPERATURE, RAINFALL, AND NUMBER OF RAINY DAYS, PHILADELPHIA\*

	Normal	1909	1910	1911
June				
Mean temperature	71.2°	72.4°	69.6°	71.6°
Rainfall	3.30 in.	2.26 in.	5.40 in.	5.10 in.
Rainy days		13	14	14+3
JULY				
Mean temperature	75.8°	75.3°	78.2°	78.5°
Rainfall	4.33 in.	2.19 in.	1.84 in.	4.19 in.
Rainy days		4+4	5	9+2
August				
Mean temperature	73.8°	73.0°	73.2°	74.5°
Rainfall	4.61 in.	1.95 in.	5.70 in.	12.10 in.
Rainy days		4+2	11	14+3
SEPTEMBER	**			
Mean temperature	67.4°	67.2°	70.0°	68.6°
Rainfall	3.38 in.	3.55 in.	3.05 in.	2.52 in.
		8	5	7+6
Rainy days		0		170
TOTALS	72.05°	71.9°	72.75°	73.3°
Mean temperature		9.95 in.	15.99 in.	23.91 in.
Rainfall	15.62 in.			The state of the s
Rainy days		29+6	35	44+14

Note.—Two figures are given above for number of rainy days—the first, in front of the plus sign, is the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is number of days there was a trace of precipitation, but less than 0.01 inch.

Except for extremely hot weather, that is, the number of days when very high temperatures were reached, there was very little difference in the temperature records for the three years.

The mean temperatures for the month show 1911 above the normal in all four months, especially in July; 1910 was lower than 1911 in each month except September, although there was only 0.3° difference in July. July and September were above the normal; 1909 was below the normal in all months except June, although very slightly. In 1909 the heat was more uniformly distributed throughout the summer.

In 1911 rainfall for the month and the number of rainy days were considerably in excess of 1910, especially in July and August. June and September, 1910, had a little more rain.

During June, July, and August temperature conditions may be said to have been distinctly less favorable to a low infant mortality than in 1909 and 1910. In respect to rainfall and its distribution during the same period, conditions were more favorable.

#### THE CAMPAIGN

Before 1909 there was very little effort made toward the reduction of infant mortality outside the ordinary work of the various charitable organizations. The Modified Milk Society was about the only special agency at work. It pro-

<sup>\*</sup> Monthly Meteorological Summary, Philadelphia; and Monthly Weather Review, Washington, D. C.



ANY PLACE WILL DO FOR A MILK STATION



ided milk modified at a central laboratory and distributed to infants from several joints in the most congested sections of the city.

In 1909 Dr. Neff, Director of the Department of Public Health and Charities, rought about a revision of the milk ordinances after considerable publicity had seen given to the statement that 12 per cent. of the milk furnished the city was rom cattle infected with tuberculosis.

A conference was held early in the summer, at which plans were considered or reducing infant mortality. As a result, the Modified Milk Society, various ettlements, and a number of women's clubs conducted work along independent ines. The Health Department placed its medical inspectors in the most concested parts of the city to canvass for sick babies and to instruct mothers in their are. Milk dealers were provided with bags upon which were printed simple lirections as to the care of milk and feeding of the baby. Bottled milk was sold in these bags.

In 1910 a conference was called by the Mayor. This was attended by representatives of some 200 agencies, the object being to bring about a better working relation between the various organizations engaged in summer work for nothers and children. As a result of this conference a Bureau of Registration and Information was established at the City Hall. A directory was published and distributed, giving the names of agencies and the kind of service each was prepared to render.

The Modified Milk Society established 10 distributing stations and later opened 8 more. The City Council appropriated funds to maintain a corps of 8 nurses, and, by utilizing the medical inspectors and district physicians, a temporary Division of Child Hygiene was established, with Dr. S. W. Newmayer in charge. This division began intensive work in June in four of the most congested wards. Other organizations assisted by furnishing nurses who reported daily to the Health Department and worked with their nurses. Two nurses each were provided by the Bureau of Municipal Research, the Visiting Nurses' Society, Starr Centre, and the Light House, and one each by the Phipps Institute and the Baby Alliance.

Previous to the closing of the schools in the four wards selected, demonstrations were given to girls in the grammar grades on the care of babies. This work was done by the medical inspectors and school nurses through the coöperation of the school authorities. A house-to-house canvass was carried on in the district by the Health Department nurses to instruct mothers in their homes.

At the central office a careful system of record keeping for each case was installed. Day and night telephone service was established for receiving requests for aid in emergency cases. The Police and Fire Departments coöperated by instructing all police stations and fire houses to forward, by telephone, any request for medical or nursing aid. A large number of bulletins, posters, and circulars were distributed, chiefly through the police department.

The Visiting Nurses' Society assumed supervision and care of sick babies. Piers were utilized for receiving and caring for babies and mothers. Special

hospital facilities were made available, the hospital at Redbank, N. J., coöperating by receiving babies.

Free ice was distributed, midwives were checked up, and several baby farms were closed.

The Philadelphia Association of Retail Druggists passed resolutions indorsing the stand taken by the Health Department in regard to soothing syrup and comforters, and discouraging their sale by its members.

There was a reduction of 40 per cent. in the deaths under two years in the four wards where the work was concentrated. In the city itself, during the three summer months, there was an increase of 132 deaths over those in 1909, though the proportion of the total deaths which occurred under one year fell 0.5 per cent.

In 1911 a still more vigorous campaign was waged. A "Milk Show" was held May 20th-27th, which was attended by 110,681 people. Considerable interest in the problem was aroused.

The City Council voted \$5,000 for the summer work, and the nursing staff was increased. The work was carried on in four wards. Sick babies were cared for at the recreation piers by night as well as by day.

The Mayor appointed a strong commission to investigate and report upon the whole subject of the milk supply. The refrigeration of milk in transit was required for the first time.

The experience of previous years bore fruit in a better organized campaign and a great reduction in the summer mortality, both from 1910 and 1909 figures. In the four wards where the nurses were working there was a reduction of 11.3 per cent. in all deaths under one year from the 1910 figures, and of 34.6 per cent. in diarrheal deaths under two years.

#### ST. LOUIS

Table 67 shows the infant mortality statistics for St. Louis for the five-year period 1906–1910, the average for the five years, and the figures for 1911.

TABLE 67.-MORTALITY UNDER ONE YEAR, ST. LOUIS

	1906	1907	1908	1909	1910	Average Five Years	1911
anuary	143	127	146	136	129	136	166
ebruary	112	121	102	151	139	125	107
March	122	144	117	128	146	131	102
April	132	121	120	116	93	116	92
May	100	76	116	121	115	105	96
une	147	122	189	151	177	157	160
uly	264	315	229	229	216	250	228
August	204	181	204	196	201	197	170
september	195	160	115	137	149	151	141
October	132	107	125	127	133	124	111
November	125	112	148	128	105	123	109
December	138	129	123	103	104	119	97
YEAR	1814	1715	1734	1723	1707	1738	1579
Percentage total mortality	17.91	16.50	17.72	16.04	15.67	16.74	14.55
une-September	810	778	737	713	743	756	699
Percentage year's infant mortality	36.5	38.2	31.6	32.6	33.2	34.5	34.1
Percentage deaths all ages	24.3	23.4	23.8	20.8	21.4	22.7	20.5
nfant mortality rate*				113.3	112.1		101.6

It will be seen that during the four years 1907–1910 there was very slight variation in the total year's mortality under one year of age. The year 1906 was a high one, the low year being 1910. In comparing the proportion of total deaths which occurred under one year of age there is no very great difference between the first and third years of the five-year period. In 1909 and 1910, however, there is a distinct fall. The year 1911 gives the lowest gross total for the year, as well as the lowest percentage of all deaths. In comparing the summer mortality it will be noticed that the four months, June, July, August, and September, are given instead of July, August, and September. St. Louis mortality begins to increase earlier than the more northern cities, and in many years June is much higher than September. If we compare these figures, we will see that there has been a slight general decrease since 1906, and that this decrease was greater in 1911 than in any other year.

<sup>\*</sup> From figures provided by Mrs. Minnie D. Weiss.

The following table shows what the deaths would have been had they increased from 1907, the low year up to 1910, at the same rate as the population, 1.94 per cent. per year.

TABLE 68.—MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED,
BASED ON 1907 FIGURES, COMPARED WITH ACTUAL MORTALITY,
ST. LOUIS

	1908	1909	1910	1911
Estimated mortality	1748	1781	1814	1848
	1734	1723	1707	1579
	— 14	— 58	—107	— 269
	— <b>0.8</b>	— <b>3.2</b>	— <b>5.</b> 8	— <b>14.5</b>

Yearly rate of increase in population, 1.94 per cent.

Allowing for the increase in population, then, there has been a steadily decreasing mortality rate, and the percentage difference from the estimate has been greater each year up to and including 1911.

#### WEATHER CONDITIONS

Comparing the weather conditions which existed in St. Louis during these years we find the facts set forth in the following tables:

TABLE 69.—NUMBER OF DAYS MAXIMUM AND MEAN TEMPERATURE REACHED OR EXCEEDED CERTAIN FIGURES, ST. LOUIS\*

	Max	KIMUM TEMPERA	MEAN TEMPERATURE		
	95° or +	90° or +	85° or +	85° or +	80° or +
1909					
June	0	5 6 18 2	14	0	6 9 23 1
July	1	6	15	1	9
August	1 4 0	18	14 15 27 8	6	23
September	0	2	8	0	1
	-	_	_	_	
	5	31	64	7	39
1910					
June	0	7	14	0	8
July	0	7 5 5	15 13	0	8 11 6 1
August	0	5	13	0	0
September	0	0	6	0	1
	0	17	48	0	26
1911	U	17	48	U	20
	0	13	22	5	15
June	8 6 2 1	13	21	5 7 3 0	15 15 11 4
August	2	13 8 5	15	3	11
September	1	5	15	0	4
Deptember	_		13		
	17	39	73	15	45

<sup>\*</sup> From Monthly Meteorological Summary, Local Weather Bureau, St. Louis, and Monthly Weather Review, Washington, D. C.



AFTER A CONSULTATION—FUTURE MOTHERS ARE BEING TAUGHT




TABLE 70.—MONTHLY MEAN TEMPERATURE, PRECIPITATION, AND NUMBER OF RAINY DAYS, ST. LOUIS\*

	AVERAGE, THIRTY- EIGHT YEARS	1909	1910	1911
une				
Mean	75°	75°	72°	79°
Rain	4.49 in.	2.63 in.	4.24 in.	1.34 in.
Rainy days		10+5	11+0	8+1
Tuly				
Mean	79°	77°	77°	79°
Rain	3.63 in.	7.34 in.	4.21 in.	0.64 in.
Rainy days		13+4	14+0	5+5
lugust				
Mean	78°	81°	75°	76°
Rain	2.48 in.	0.66 in.	1.90 in.	3.51 in.
Rainy days		4+3	7+3	11+1
September				
Mean	70°	68°	70°	74°
Rain		4.22 in.	6.09 in.	7.09 in.
Rainy days		00+0	10+2	14+5
TOTAL				
Average mean		75.2°	73.5°	77°
Rain		14.85 in.	16.44 in.	12.58 in.
Rainy days		00+0	42+5	38+12

Note.—Two figures are given for number of rainy days—the first, in front of the plus sign, s the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is he number of days there was a trace of precipitation, but less than 0.01 inch.

It will be seen that June, 1911, averaged for the month higher than either of the other years or than the thirty-eight-year average. The same is true for July.

August, on the contrary, is 5° below 1909 and 2° below the mean, but August and some very hot weather, the thermometer registering 95° on two days and the nean for the day being 85°+ 'on three days. September was hot, the mean being well above either year and the thirty-eight-year average. The rainfall also in 1911 was very scant through June and July, but above the average for August and September, especially in the latter month. Extremely high temperatures were reached during the four summer months of 1911, remained high for onger periods, and the hot weather was continued throughout the summer more evenly than in any of the other years considered.

In spite of very much worse conditions in June and July, the mortality in June was 17 below 1910 and 9 above 1909. In July it was 1 below 1909 and 12 above 1910.

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, St. Louis, and Monthly Weather Review, Washington, D. C.

# THE CAMPAIGN

In 1904 the St. Louis Pure Milk Commission was incorporated for the purpose of distributing modified milk to poor babies. After two years of work Dr. A. S. Bleyer started consultations at one of the stations of the Commission. This work was founded with the idea of instructing mothers in the care of their babies, and modified milk was dispensed. In the spring of 1911 the "Save the Babies" League was formed, to coöperate with existing agencies for the prevention of sickness and death among the babies of St. Louis. In the previous summer the Pure Milk Commission had carried on some nursing work, but was financially unable to continue it. In 1911 the Visiting Nurses' Association had one nurse in the field. The object of this League was the maintenance of conferences where mothers were advised and instructed by the doctors in charge in the care and feeding of their babies; follow-up work in the homes by the nurses, especially in the case of sick babies; and coöperation with the St. Louis Pure Milk Commission in furnishing safe modified milk for babies.

On June 1st one nurse started in on her work, the number being rapidly increased to 9. Thirteen stations were in operation, 6 of them being opened for five months,—from May 1st to October 1st,—and 729 babies being enrolled. There were 17 doctors, 9 nurses, and 6 assistant workers engaged in the work. Particular stress was laid upon the home work, and during the latter part of the season a considerable number of the mothers were taught home modification of milk. This was the first year that any extensive campaign was carried out. Coöperation between various organizations was extremely good, and plans are in hand for a more extensive campaign in 1912.

The Health Department did very little special work in connection with the campaign. Milk inspection in St. Louis is still far from thorough, owing to lack of available funds.

### BOSTON

The following table gives the facts in regard to infant mortality in Boston for the years 1906-1911:

TABLE 71.-MORTALITY UNDER ONE YEAR, BOSTON

TABLE /IMORTALI	11 0	VDER	ONE	I LAIN,	DOST	711	
	1906	1907	1908	1909	1910	Average Five Years	1911
Tanuary	187	177	197	175	163	179.8	163
JanuaryFebruary	168	138	198	157	183	168.8	175
March	186	96	208	192	189	174.2	196
April	201	156	198	161	163	175.8	191
May	197	185	182	146	140	170.0	180
June	152	151	150	117	168	147.6	129
July	240	172	305	188	225	226.0	280
August	357	356	405	313	254	337.0	260
September	271	282	300	240	270	272.6	224
October	186	229	222	159	225	204.2	172
November	165	147	186	124	148	154.0	138
December.	158	187	191	153	121	162.0	137
YEAR	2468 21.62	2276 <b>19.47</b>	2742 23.28	2125 <b>19.20</b>	2249 <b>19.45</b>	<b>2372.0</b> 20.62	2245 <b>19.08</b>
July, August, September		810 <b>35.58</b> <b>29.88</b>	1010 36.83 34.04	741 34.87 29.25	749 33.30 27.18	835.6 35.22 30.27	764 33.58 25.10
Births		18,436 123.4	18,521 4148.0	17,728 119.8	17,741 126.7	17,942 132.2	1258 126.0

It will be noted that, excepting 1908, a very high year, there has been little striking change in the gross figures. In percentages of total mortality 1907, 1909, 1910, and 1911 remain practically the same, differing only in a small fraction of 1 per cent.

In the totals for summer mortality the last three years have been very nearly the same, being slightly higher each year. The infant mortality rate shows an increase in 1910 and 1911 over both 1907 and 1909. The rate for 1911, as furnished by the Registrar of Vital Statistics for Boston, Dr. Davis, is given at 126, practically the same as in 1910.

If we estimate the deaths which would have occurred if they had increased in the same proportion as the population, 1.96 per cent., yearly, taking 1909, the low year for the five-year period, as a basis, and then compare these figures with actual deaths, we find the following:

TABLE 72.—MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED ON BASIS OF 1909 FIGURES, COMPARED WITH ACTUAL MORTALITY,

BOSTON	
1910	1911
Estimated mortality	2208
Actual mortality	2245
Difference+83	+37
Percentage difference+3.8	+1.7

Estimating for the three summer months, taking 34.87 per cent. as the proportion of deaths occurring during these months in 1909, we have the figures shown in the following table:

TABLE 73.—MORTALITY UNDER ONE YEAR IN THREE SUMMER MONTHS WHICH SHOULD HAVE OCCURRED ON BASIS OF 1909 FIGURES, COMPARED WITH ACTUAL MORTALITY, BOSTON

	1910	1911
Estimated mortality	755	769
Actual mortality	749	764
Difference		<b>—</b> 5
Percentage difference	—0.79	-0.65

Boston, therefore, shows almost no improvement in infant mortality for the year compared with three of the five preceding years. The summer mortality has not been reduced in the last three years. The percentage of the year's infant deaths occurring during the three summer months is still fairly high, as is the proportion of all deaths during this period.

Taking the *high* figure, 2742, in 1908, and estimating the deaths which should have occurred on the basis of a yearly increase of 1.96 per cent., there is only a decrease of 22.7 per cent., while New York shows for the year 1911 a decrease of 20.4 per cent. from the *lowest*, allowing for increase in population.


# WEATHER CONDITIONS

The weather conditions in Boston during the last three years are shown in the following tables:

TABLE 74.—NUMBER OF DAYS ON WHICH MAXIMUM AND MEAN TEMPERA-TURES REACHED CERTAIN FIGURES, BOSTON\*

		MAXIMUM	Mi	EAN	
	95° or +	90° or +	85° or +	85° or +	80° or +
1909					
June	0	4	9	0	6
July	1	4	8	0	4
August	0	4 3 0	6	0	4 4 0
September	0	0	0	0	0
	-	-	-	-	-
TOTAL	1	11	23	0	14
910					
June	0	1	6	0	2 5 0
July	0	6	16	0	5
August	0	0	4	0	0
September	0	0	0	0	0
Tomas	0	7	-	0	7
1911 TOTAL	0	,	26	0	,
June	0	0	2	0	0
July	0 8		14	7	10
August	0	9 3 0	6	0	1
September	0	0	1	0	Ô
	_	_	_	-	_
TOTAL	8	12	23	7	11

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Boston, and Monthly Weather Review, Washington, D. C.



DISTRIBUTED TO 3500 STUDENTS ATTENDING UNIVERSITY SUMMER SCHOOL IN NEW YORK CITY EXHIBIT 18




TABLE 75.—MONTHLY MEAN TEMPERATURE, RAINFALL, AND NUMBER OF RAINY DAYS, BOSTON\*

	Normal	1909	1910	1911
June				
Mean temperature	66°	69°	65°	66°
Rainfall	3.08 in.	4.45 in.	4.89 in.	3.67 in.
Rainy days		8	12+2	8+7
July				
Mean temperature	72°	72°	75°	77°
Rainfall	3.18 in.	0.97 in.	1.15 in.	4.65 in.
Rainy days		5+7	6+1	4+3
August				
Mean temperature	70°	69°	69°	69°
Rainfall	3.84 in.	3.55 in.	0.98 in.	4.18 in.
Rainy days		8+3	7+2	9+4
September				
Mean temperature	63°	63°	63°	63°
Rainfall	3.22 in.	5.15 in.	2.14 in.	2.95 in.
Rainy days		12+1	11+1	10+1
TOTALS				
Mean temperature	68.7°	68.2°	68.0°	68.7°
Rainfall	13.32 in.	14.12 in.	9.16 in.	15.45 in.
Rainy days		32+11	36+6	31+15

Note.—Two figures are given for number of rainy days—the first, in front of the plus sign, is the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is number of days there was a trace of precipitation, but less than 0.01 inch.

Except for the month of July, the conditions are very similar. In this month the temperature reached 95° or over on eight days in 1911, and the mean temperature was 85° or over on seven days. The high figures are all crowded into the first two weeks of July; August was cool. In summing up the monthly figures compared with the normal, June was exactly the same. July was 5° over; August was below; September was the same. The rainfall the first three months is in excess of the normal; September, somewhat below.

Compared with 1910, June was 1° hotter; July, 2° hotter; August and September were the same. In 1909 June was 3° hotter; July, 5° cooler; August and September, the same.

The rainfall in 1911 was less in June, much more in July and August, and less than 1909 in September, but more than in 1910. It was less evenly distributed throughout the summer than either year.

#### THE CAMPAIGN

In Boston practically the only organization doing purely baby work, and laying particular stress on prevention, is the Boston Milk and Baby Hygiene Association. In 1905 a group of public-spirited citizens organized a "poor people's" modified milk service. They distributed milk already modified to

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Boston, and Monthly Weather Review, Washington, D. C.

various settlements, day nurseries, etc. In 1907 a committee on modified milk stations was organized to take charge of the work. After a year this committee came to the conclusion that its policy was unintentionally encouraging weaning, and a new method was adopted. In 1909 they opened 10 milk stations, under the control of the Committee on Milk and Baby Hygiene. A director was appointed, and consultations were inaugurated, under volunteer physicians, a medical director, and trained nurses.

In 1910 the association was incorporated, and has been carrying on the work since. Twenty volunteer physicians have charge of the 10 stations; consultations are held from once to three times a week; modified milk is dispensed; earnest efforts are made to encourage breast-feeding; mothers receive home instruction; all sick babies are referred to physicians or dispensaries; sick babies are referred also to the Instructive Visiting Nurses' Association, whose nurses care for the babies while sick.

In 1911, 9 stations were maintained; 10 nurses were in the field; 13 volunteer physicians had charge of consultations. Home modification was taught "where possible."

In 1909 and 1910 the Women's Municipal League carried on a demonstration of prenatal and postnatal instruction to mothers. They have since discontinued their postnatal work, which was planned as a demonstration.

The work in Boston in 1911 showed no increase except in efficiency, perhaps, over that done in the three previous years. The work of the Milk and Baby Hygiene Association has been excellent, but it cannot reach more than a small proportion of those who need its aid.

The Health Department had two nurses in the field doing prophylactic work, including prenatal instruction.

# CLEVELAND

In considering the mortality returns for Cleveland, it must be remembered that, with the exception of Detroit and the three boroughs of Bronx, Queens, and Richmond in New York, the percentage increase in population during the past decade has been greater than in any of the other cities considered. The following table shows the details of the infant mortality for the years 1906 to 1911 inclusive:

TABLE 76.-MORTALITY UNDER ONE YEAR, CLEVELAND

	1906	1907	1908	1909	1910	AVER- AGE 1906-10	1911
January	108	166	153	133	134	139	163
February	145	162	144	148	141	148	147
March	180	170	171	170	194	177	147
April	194	150	135	142	132	150	148
May	123	137	119	117	147	128	140
June	215	124	104	99	157	140	137
July	272	164	240	204	322	140	225
August	176	304	209	213	278	136	199
September	145	188	142	137	174	157	145
October	122	175	142	126	128	138	124
November	121	113	128	108	127	119	92
December	124	135	114	114	123	122	121
YEAR Percentage of all deaths	1925 <b>26.1</b>	1988 <b>25.9</b>	1801 <b>25.7</b>	1711 24.5	2057 <b>25.6</b>	1896 25.5	1788 22.4
July, August, and September	593 <b>30.8</b> <b>30.2</b>	656 <b>33.0</b> <b>30.1</b>	591 32.8 33.2	554 32.4 30.3	774 37.6 35.3	634 33.4 32.7	569 <b>32.0</b> <b>28.5</b>
Births	11,201 171.9	10,700 185.8	12,010 150.0	13,100 130.6	13,456 152.9	12,093 156.8	12,589 142.0

During the years 1906-09 inclusive there was a steady diminution in the total number of deaths under one year and in the proportion of all deaths. The year 1910 showed a sharp rise both in the year's total and during the summer months, the proportion of the mortality occurring during the summer months being the highest in any of the years considered.

On the other hand, 1911 had a lower total mortality than any year except 1909. The same is true of the summer mortality, and the percentage of all deaths both for the year and for the summer months; the proportion of the year's deaths occurring during the three summer months is also lower than any year except 1906. The greatest reduction, therefore, occurred during the summer months, but there is a marked fall for the entire year.

In the following table is shown the comparison of 1911's figures with those of other years:

TABLE 77.—DIFFERENCE BETWEEN MORTALITY UNDER ONE YEAR IN 1911 AND OTHER YEARS, CLEVELAND

	1906	1907	1908	1909	1910
Year  Percentage all deaths  July, August, and September  Percentage of year's infant mortality .  Percentage all deaths	$-24 \\ +1.2$	-210 -3.5 -87 -1.0 - <b>1.6</b>	-13 -3.3 -22 -0.8 -4.7	+77 -2.1 +15 -0.4 -1.8	-269 -3.2 -205 -5.6 - <b>6.</b> 8

If, now, we compare the results in 1911 with those in 1909, allowing for the increase in population, we find results shown in the following table:

TABLE 78.—MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED, BASED ON 1909 FIGURES, COMPARED WITH ACTUAL MORTALITY, CLEVE-LAND

	1910	1911
Estimated mortality Actual mortality Difference, actual Per cent. difference	1791 2057 +266 + <b>14.8</b>	1871 1788 83 <b>4.4</b>

Yearly rate of increase in population, 4.69 per cent.

The year 1910 showed 266, or 14.8 per cent., above 1909's figures, allowing for the increase in population; whereas 1911 was 83, or 4.4 per cent., less.

There was, therefore, a distinct improvement in the infant mortality conditions in Cleveland, although if the registration of births is anywhere near complete, the death-rate is still very high, and there is not the same degree of improvement noted as in a number of other cities.

## WEATHER CONDITIONS

The weather conditions in Cleveland are shown in Tables 79 and 80.

In 1909 in no month was the temperature above the normal. On only one day did it reach 90°, and the mean daily temperature reached 80° only on one day. In 1910 the temperature for the month was 1° higher than the normal in September. The mean for the four months was 0.5° below the normal; and again there was only one day when 90° or over was recorded. The mean was 80° or + on four days.

In 1911 all four months were above the normal, and the mean for four months was 1.3° above. Individual days of high temperature also were more numerous; 95° or over was reached once, 90° or over seven times. Twice the mean daily temperature was 85° or over and 9 times 80° or over. The high temperatures were scattered through the summer, except in July, when on four consecutive days 90° or over was reached (July 2d to 5th). The rest of July was cool.

TABLE 79.—NUMBER OF DAYS ON WHICH THE MAXIMUM AND MEAN TEM-PERATURE REACHED CERTAIN FIGURES, CLEVELAND\*

		MAXIMUM	MEAN		
	95° or +	90° or +	85° or +	85° or +	80° or +
909					ine in
June	0	0	3	0	0
July	0	1	4	0	0
August	0	0	3	0	1
September	0	0	1	0	0
910 TOTAL	0	1	11	0	1
June	0	0	0	0	0
July	0	1	7	0	2
August	0	0	8	0	2
September	0	0	1	0	0
911 TOTAL	0	1	16	0	4
June	0	2	6	0	2
July	1	4	7	2	5
August	0	1	6	0	2
September	0	0	1	0 -	0
TOTAL	1	7	20	2	9

TABLE 80.—MONTHLY MEAN TEMPERATURE, RAINFALL, AND NUMBER OF RAINY DAYS, CLEVELAND\*

	NORMAL	1909	1910	1911
June				
Mean temperature	67°	67°	64°	69°
Rainfall	3.54 in.	3.72 in.	2.01 in.	2.88 in.
Rainy days		13	9+3	10+5
July		10	710	10 10
Mean temperature	72°	70°	72°	73°
Rainfall	3.56 in.	3.12 in.	0.94 in.	1.71 in.
Rainy days	0.00 m.	9+5	9+3	7+5
August		110	10	
Mean temperature	70°	70°	70°	71°
Rainfall	2.87 in.	2.31 in.	1.08 in.	5.18 in.
Rainy days		10+3	6+0	11+5
September		1010		
Mean temperature	64.1°	62°	65°	65°
Rainfall	3.33 in.	3.40 in.	4.09 in.	4.67 in.
Rainy days		10+3	11+2	11+3
ramy days		10   0		
TOTALS				
	68.2°	67.2°	67.7°	69.5°
Mean temperature Rainfall	13.30 in.	12.55 in.	8.12 in.	14.44 in.
Rainy days	19.90 III.	42+11	35+8	39+18

Note.—Two figures are given for number of rainy days—the first, in front of the plus sign, is the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is the number of days there was a trace of precipitation, but less than 0.01 inch.

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Cleveland, and Monthly Weather Review, Washington, D. C.

The weather conditions were less favorable for a low infant mortality in 1911 than in 1909 or 1910, except in regard to rainfall, the most marked advantage in this respect in 1911 being in August.

# THE CAMPAIGN

In 1899 a Milk Fund Association was formed, which distributed clean milk in bulk to any one who desired to buy it. In 1906 the Infants' Clinic was established by the Milk Fund Association and the Visiting Nurse Association, to provide, during the summer, medical counsel, nursing care, and instruction in the homes and clean milk for sick babies in its neighborhood. The clinic was so successful that it became evident that a permanent organization was needed. The value of the home nursing was demonstrated beyond question.

In December, 1906, the Babies' Dispensary and Hospital of Cleveland was incorporated. A medical director was appointed, and Dr. Gerstenberger has continued from its founding to devote his time and thought to the undertaking. It is largely due to him and his clear conception of the problem that the work has succeeded as it has. In May, 1907, the dispensary and the milk laboratory were opened. Three visiting nurses were provided, and two dispensary physicians appointed. Milk was modified at the central plant and distributed by hand.


In 1908 an out-physician was added to the staff to visit cases too sick to come to the dispensary. Four nurses were employed, who made 8503 visits. Milk was still delivered to the homes with a wagon. Considerable modification of milk was done at the home by mothers; those incapable of doing so received already modified milk.

Particular stress was laid on the educational side and to the encouragement of maternal nursing.


In 1909 four branch dispensaries were formed. Great coöperation existed between the Milk Fund Association, the Visiting Nurses' Association, the Health Department, and various settlement houses. An outdoor ward was established where sick babies were treated—at first from 8 A. M. to 6 P. M., and later day and night. The Milk Fund Association became amalgamated with the dispensary, thus putting the work on a more rational basis.

In 1910 there were 6 branch dispensaries. A vigorous campaign of education for reduction of infant mortality was carried out, and was participated in by a large number of civic and philanthropic organizations.

In 1911 the New Central Dispensary was built and in operation. The City Council appropriated \$10,000 for the reduction of infant mortality, and 5 new stations were opened, making a total of 11 with 60 distributing stations. Two were managed by the Health Department, but, owing to the wonderful coöperation which existed, the Babies' Dispensary and the Health Department worked together. In October, 5 of the branch dispensaries were taken over by the city. A Bureau of Child Hygiene was formed, and in September Dr. Gerstenberger became its head. Cleveland is a fortunate city in the complete, broad coöperation that has marked its infant mortality work, especially in recent years.



EVERY STATION BABY EXAMINED BY A CAPABLE PHYSICIAN



# BALTIMORE

In the following tables is set forth the infant mortality by months and groups of months, with the percentage of all deaths occurring under one year of age for the years 1907 to 1911:

TABLE 81.-MORTALITY UNDER ONE YEAR, BALTIMORE

	1907	1908	1909	1910	1911
January	167	180	155	176	143
February	153	160	139	170	138
March	182	166	161	148	165
April	134	157	162	132	141
May	139	157	173	114	152
June	140	201	168	183	125
July	397	399	317	306	280
August	370	233	264	266	254
September	239	184	215	180	182
October	215	138	172	190	132
November	145	123	157	138	109
December	162	117	144	145	137
YEAR	2443 21.84	2215 21.26	2227 <b>21.43</b>	2148 <b>19.97</b>	1958 18.82
July, August, September	1006 41.10 34.01	816 <b>36.84</b> <b>31.14</b>	796 35.74 29.35	752 <b>35.01</b> <b>28.05</b>	716 36.57 28.23

From this table it will be seen that the gross figures for the years have had a steady tendency to diminish, and that this tendency has been increasingly marked in 1910 and 1911. The relation of the deaths under one year to total deaths remained practically constant during the first three years shown in the table, but has steadily fallen in 1910 and 1911.

Comparing the summer mortality, there is the same steady decrease in the gross figures. The decrease, however, has been slow. The proportion of the year's deaths under one occurring during the three summer months in 1911 was higher than in either 1909 or 1910. There is also a slight increase in the proportion of all deaths during the three summer months which are under one year of age. In other words, there was a greater proportionate decrease during the rest of the year than during the summer months.

The population of Baltimore increased 9.7 per cent. in the decade 1900–10, or 0.97 per cent. yearly. If we estimate the deaths which would have occurred had the total increased 0.97 per cent. yearly from 1908, which was the low figure up to 1910, and then note the difference between this estimate and the deaths that did occur, we shall have the results shown in the following table:

TABLE 82.—MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED, BASED ON 1908 FIGURES, COMPARED WITH ACTUAL MORTALITY, BAL-TIMORE

	1909	1910	1911
Estimated mortality	2236	2257	2279
	2227	2148	1958
	—9	—109	—321
	— <b>0.4</b>	— <b>4.8</b>	— <b>14.0</b>

Yearly average increase in population, 0.97 per cent.

The results, as shown by the yearly figures, are therefore good. The reduction in the summer mortality, however, is slight, and there is still a very high proportion of the year's infant deaths occurring during the summer months.

Birth statistics for Baltimore are of no value.

# WEATHER CONDITIONS

The following tables show weather conditions in Baltimore:

TABLE 83.—NUMBER OF DAYS ON WHICH MAXIMUM AND MEAN TEMPERA-TURES REACHED CERTAIN FIGURES, BALTIMORE\*

	MA	XIMUM TEMPERAT	MEAN TEMPERATURE		
	95° or +	90° or +	85° or +	85° or +	80° or +
909					
June	0	9 7	13	1	9
July	0 1 0	7	16	1	9 9 6 0
August	1	6	13	1	6
September	0	0	0	0	0
_	_			_	T.
TOTAL	1	22	42	3	24
910					
June	0	4	10	1	6
[uly	0	15	23	1 2	16
August	0	1	9 5	0	3
September	1	2	5	1	16 3 3
	_	-	-	-	-
TOTAL	2	22	47	4	28
911					
June	1	7	10	1	4
July	1 7 2 0	11	20	1 7 2 0	14
August	2	6	14	2	9
September	0	0	6	0	9
	-	-	_	_	_
TOTAL	10	24	50	10	28

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Baltimore, and Monthly Weather Review, Washington.

TABLE 84.—MONTHLY MEAN TEMPERATURE, RAINFALL, AND NUMBER OF RAINY DAYS, BALTIMORE\*

	NORMAL	1909	1910	1911
Tune				
Mean temperature	73°	74.1°	70.4°	73.4°
Rainfall	3.84 in.	4.38 in.	5.3 in.	5.52 in.
Rainy days		13+8	14+1	16
uly	**	1370	14+1	10
Mean temperature	77.3°	76°	78.8°	79.2°
Rainfall	4.82 in.	1.31 in.	0.95 in.	3.53 in.
		5+7		
Rainy days		3+1	7+2	6+4
August	74.7°	74.4°	74 20	76.00
Mean temperature			74.3°	76.2°
Rainfall	4.21 in.	0.86 in.	1.37 in.	12.28 in.
Rainy days		5+3	11+3	15+4
September	00.00	CM CO	B	
Mean temperature	68.6°	67.6°	71.6°	70.4°
Rainfall	3.85 in.	3.97 in.	2.13 in.	1.6 in.
Rainy days		5+5	4+7	7+1
TOTALS		AND LABOR.		
Mean temperature	73.4°	73.02°	73.7°	74.8°
Rainfall	16.72 in.	10.52 in.	9.75 in.	22.93 in.
Rainy days		28+23	36+13	44+9

Note.—Two figures are given for number of rainy days—the first, in front of the plus sign, is the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is the number of days there was a trace of precipitation, but less than 0.01 inch.

For maximum temperatures below 95° and mean temperatures below 85° there is a very close resemblance between the three years. But temperatures of 95° or over were recorded 10 times in 1911, as against twice in 1910 and once in 1909. Mean temperatures of 85° or over were also recorded 10 times in 1911 as against 4 times in 1910 and 3 times in 1909.

The monthly mean temperatures in 1911 were well above the normal in each month, while in 1909 only June, and in 1910 only July and September, exceeded the normal. The year 1911 had a higher monthly mean than 1909 in each month except June, and higher than 1910 in each month except September.

Rainfall, on the other hand, was much greater in 1911 than in either 1909 or 1910, except in September, and exceeded the normal in the months of June and August. August had the remarkable rainfall of 12.28 inches, which is 2.79 inches greater than any recorded in August since the establishment of the Weather Bureau.

#### THE CAMPAIGN

The chief agencies engaged in work for the reduction of infant mortality in Baltimore are the Babies' Milk Fund Association and, to a much less extent, the Mothers' Relief Society. The latter society was founded in the early nineties, its purpose being the assistance of the mothers during and after confinement.

The Babies' Milk Fund was started in 1904, when the Thomas Wilson Sanitarium established four milk stations in distant parts of the city. The work was

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Baltimore, and Monthly Weather Review, Washington, D. C.

shown to be of such value that, after a few years, the Babies' Milk Fund Association was incorporated. Funds were raised by popular subscription, the Thomas Wilson Sanitarium being the largest contributor.

Babies were referred to the stations from hospitals, dispensaries, and private physicians. Milk modified to set formulæ was dispensed, and special formulæ could be secured. Milk was dispensed by trained nurses, who gave the greater part of the day to visiting at the homes, instructing and demonstrating.

In 1910 the two above-mentioned organizations united to form the Maryland Association for the Study and Prevention of Infant Mortality. It was managed by a committee composed of representatives of both organizations. After one

year the Mothers' Relief Society withdrew.

In 1911 the work done increased considerably in efficiency and also in extent. Nine stations are now maintained, open all the year round. They are in charge of a trained nurse. Babies are referred by physicians and organizations, and there is little or no house-to-house canvassing. Milk already modified is supplied from six of the stations. From the other three home modification is practised and taught, tuberculin-tested milk being delivered directly to the mothers' homes, and the modification taught by the nurse. At the first two visits she modifies for the mother, and at the last visit watches the mother do it. This latter work—home modification—was started in 1911.

A weekly clinic is held at each station, when babies are weighed and mothers instructed. Great emphasis is laid on breast-feeding. Instruction is individual, not in classes. Attendance is not compulsory; milk is delivered whether they attend or not. During the year Dr. Henrietta Thomas has been conducting follow-up work for cases discharged from the Johns Hopkins Maternity services. She has had 500 babies under her care. They are also visited by the association nurses from time to time. Milk is not dispensed to these babies.

The work of the association is being widened in its scope by the establishment of welfare clinics at six of the stations, and by prenatal work among expectant mothers referred by three hospitals in Baltimore.

The number of babies supplied with milk by the stations has been as follows:

TABLE 85.—NUMBER OF BABIES FED AT MILK STATIONS IN BALTIMORE AND MORTALITY AMONG SAME

				DEATHS	RATE
February	, 1908,	to February,	19091073	103	96.0 per 1000
"			19101217	105	86.3 " "
11	1910,	to "	19111356	188	138.7 " "
**	1911,	to "	19121194	50	41.9 " "

Of these 1194 babies in 1911–12, 927 were supplied with modified milk; 267 with whole milk modified at home.

The Department of Health has done little active special work, owing to lack of funds. During the last two years considerable interest in the problem has been aroused by the annual meeting of the American Association for the Study and Prevention of Infant Mortality in 1910, and by general exhibits on Baby Hygiene, etc.

#### PITTSBURGH

The following table shows the infant mortality statistics for the years 1907-11 n Pittsburgh:

TABLE 86.-MORTALITY UNDER ONE YEAR, PITTSBURGH

	1907	1908	1909	1910	1911
[anuary	129	223	177	162	171
February	146	259	157	176	134
March	137	183	120	185	151
April	139	172	131	169	142
May	148	142	140	148	131
[une	148	160	153	166	116
July	244	251	273	338	250
August	189	214	184	231	197
September	127	176	164	178	144
October	126	144	143	191	132
November	112	143	134	157	118
December	159	127	147	150	124
YEAR	1804	2194	1923	2251	1810
Percentage all deaths	24.45	24.29	23.04	23.43	22.33
I.I. A S	560	611	621	747	501
July, August, September	560 <b>31.04</b>	641 29.20	621 32.29	747 33.18	591 <b>32.65</b>
Percentage year's infant mortality  Percentage all deaths	29.82	31.08	31.19	33.33	27.75

It will be seen that in 1911 the year's total is the lowest since 1907, which year was 6 below 1911's figures. The percentage of the total deaths occurring under one year of age is lower than in any year, including 1907.

During the summer months the deaths were 591, the lowest total of the five years except 1907. The percentage of the year's mortality which occurred during these three months was lower than in 1910, but higher than in 1907, 1908, or 1909.

If, now, taking as a basis the figures of 1907, the lowest year up to 1911, we estimate the deaths as increasing at the same rate as the population, and compare the actual deaths with these figures, we find the results shown in the following table:

TABLE 87.—MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED, BASED ON 1907 FIGURES, COMPARED WITH THE ACTUAL MORTALITY, PITTSBURGH

	111151	JUNGII		
	1908	1909	1910	1911
Estimated mortality	1836 2194 +358 + <b>19.5</b>	1869 1923 +54 + <b>2.8</b>	1902 2251 +349 + <b>18.3</b>	1935 1810 —125 — <b>6.4</b>

Average yearly increase in population, 1.82 per cent.

These figures show that in each of the years 1908, 1909, and 1910 more deaths occurred than the estimate, while 1911 alone shows a decrease of 6.4 per cent. from the estimate.

There has, therefore, been a very decided improvement this year over previous years. The proportion of infant deaths occurring during the three summer months has not, however, fallen as much as in other places.

# WEATHER CONDITIONS

Weather conditions are shown in Tables 88 and 89.

TABLE 88.- NUMBER OF DAYS ON WHICH MAXIMUM AND MEAN TEMPERA-TURES REACHED CERTAIN FIGURES, PITTSBURGH\*

	MA	XIMUM TEMPERA	MEAN TEMPERATURE		
	95° or +	90° or +	85° or +	85° or +	80° or +
1909					
June	0	0	6	0	0
July	0	0	7	0	0
August	0	2	8	0	1
September	Ö	0 2 0	7 8 3	0	Ô
September	_	_	_	_	_
	0	2	24	0	1
1910	0	-	21		
	0	2	10	0	0
June	Ö	1	13	0	2
July	0	1		0	3 0
August	0	1	10	0	0
September	U	0	1	0	U
	_	_	24	_	3
1011	0	4	34	0	3
1911					
June	0 5 1	4 8 6 0	5 17	0	2 9 7
July	5	8	17	4	9
August	1	6	18 3	0	7
September	0	0	3	0	0
	_	7.	_	_	_
	6	18	43	4	18

Extremely high temperatures were reached more often in 1911 than in either 1910 or 1909, especially as compared with the latter year. In neither 1909 nor 1910 was 95° recorded. The same general relation is true for the daily means. The hot weather was especially severe in July and August, 1911 (during forty-eight days beginning July 1st the thermometer reached 85° or over on thirty-two days). In neither 1909 nor 1910 did the mean reach 85°, against six and four days in July and August, 1911.

The monthly mean temperatures are very much the same in 1910 and 1911, differing only in August, when in 1911 it was 1° hotter, and June, when it was 4° hotter. All four months of 1911 are above the normal averages. June, 1909 and 1911, were the same, but all other months were much below in 1909.

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Pittsburgh, and Monthly Weather Review, Washington, D. C.



THE STATION NURSE INSISTS UPON MATERNAL NURSING WHEREVER POSSIBLE



TABLE 89.—MONTHLY MEAN TEMPERATURE, RAINFALL, AND NUMBER OF RAINY DAYS, PITTSBURGH\*

	Normal	1909	1910	1911
une				
Mean	70.7°	71°	67°	71°
Rainfall	3.71 in.	4.92 in.	1.94 in.	2.63 in.
Rainy days		16	13+3	9+6
uly			20,0	210
Mean	74.8°	71°	75°	75°
Rainfall	4.39 in.	1.22 in.	1.26 in.	2.17 in.
Rainy days		9+6	9+4	7+6
ugust		710	212	7-10
Mean	72.8°	72°	73°	74°
Rainfall	3.07 in.	3.33 in.	2.47 in.	6.3 in.
Rainy days		9+5	8+1	
eptember		713	071	13+4
Mean	66.7°	65°	68°	68°
Rainfall	2.48 in.	0.76 in.	5.5 in.	
		8 8		6.36 in.
Rainy days	••	0	10+4	13+4
COTALS				
Mean	71.2°	69.7°	70.7°	72°
Rainfall	13.65 in.	10.23 in.	11.17 in.	17.46 in.
Rainy days	20.00 1	42+11	40+12	42+20

Note.—Two figures are given for number of rainy days—the first, in front of the plus sign, is the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is number of days there was a trace of precipitation, but less than 0.01 inch.

The rainfall in August and September, 1911, was above the normal, and also above either 1909 or 1910, especially in August. June and July were below the normal, but in excess of 1910 in June and in excess of both years in July.

It has been impossible to obtain information first hand in regard to the work being done in Pittsburgh.

\* Monthly Meteorological Summary, Local Weather Bureau, Pittsburgh, and Monthly Weather Review, Washington, D. C.

# DETROIT

In studying the mortality figures for Detroit it must be remembered that the population has increased 63 per cent. in ten years. The following tables give the statistics of infant mortality for the years 1906 to 1911:

	1906	1907	1908	1909	1910	AVERAGE 1906-10	1911
January	97	124	126	130	148	125	160
February	113	107	141	126	132	124	154
March	137	153	165	149	200	161	138
April	147	114	180	150	196	157	104
May	145	119	119	140	141	133	150
June	134	103	117	114	198	133	133
July	221	186	206	211	275	220	244
August	172	198	207	220	251	209	215
September	154	166	162	171	182	167	179
October	136	133	131	149	125	135	130
November	116	100	91	114	118	108	121
December	107	95	89	127	129	109	134
YEAR Percentage of all deaths	1679 <b>28.0</b>	1598 <b>26.4</b>	1734 29.5	1801 28.5	2095 <b>28.1</b>	1781 28.1	1862 <b>26.1</b>
uly, August, and September	547	550	575	602	708	596	638
Percentage year's infant mortality	32.6	34.4	33.6	33.4	33.8	33.5	34.3
Percentage all deaths	33.3	35.8	38.5	36.0	35.3	35.7	34.0
Births	7752	8341	9572	9617	11,509	9358	13,373
nfant mortality rate	216.5	191.5	181.1	187.2	182.0	190.3	139.2

TABLE 91.—DIFFERENCE BETWEEN MORTALITY UNDER ONE YEAR IN 1911 AND PREVIOUS YEARS, DETROIT

	1906	1907	1908	1909	1910
Year	+183	+264	+128	+61	-233
	-1.85	0.24	-3.36	-2.42	-1.99
	+91	+88	+63	+36	-70
	+1.32	0.16	+1.10	+0.84	+0.47
	+ <b>0.74</b>	<b>1.82</b>	-4.51	-2.02	- <b>1.26</b>

It will be seen that from 1907 through 1910 there has been a slowly increasing gross total infant mortality. The proportion of total deaths which occurred under one year of age was higher than in 1907 during the three years following it, although it has slowly fallen each year after 1908. The year 1911, however, shows the lowest percentage of all deaths of any of the six years, being even lower than 1907.

Comparing the summer mortality, the total has slowly increased, but not in

proportion to the population. The year 1911 shows a lower total than 1910, but higher than the four previous years. The percentage of the year's mortality which occurred during these three months is also higher than in any year except 1907. In other words, the summer mortality is not reduced quite so much as that for the rest of the year. The percentage of all deaths occurring during the summer months is, however, lower than any since 1906. If we take deaths in 1907, the low year, as a basis, and estimate the increase on the same basis as increase in population, the mortality should have been:

TABLE 92.—MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED ON BASIS OF 1907 FIGURES, COMPARED WITH ACTUAL MORTALITY, DETROIT

	1908	1909	1910	1911
Estimated mortality	1698	1799	1900	2000
	1734	1801	2095	1862
	+36	+2	+195	—138
	+ <b>2.1</b>	+ <b>0.1</b>	+ <b>9.3</b>	— <b>6.9</b>

Average yearly increase in population, 6.3 per cent.

Applying the same method to the summer mortality we find:

TABLE 93.—MORTALITY UNDER ONE YEAR DURING THREE SUMMER MONTHS WHICH SHOULD HAVE OCCURRED ON BASIS OF 1907 FIGURES, COM-PARED WITH ACTUAL MORTALITY, DETROIT

	1908	1909	1910	1911
Estimated mortality	584	619	654	688
	575	602	708	638
	—9	—17	+54	—50
	— <b>0.15</b>	— <b>2.7</b>	+8.2	— <b>7.3</b>

The summer reduction has, therefore, kept pace with the yearly reduction, allowing for increase in population. The infant mortality rate also shows a very marked fall to 139.2, but it must be remembered that the great increase in births during 1910 and 1911 is probably partly due to a more efficient registration.

## WEATHER CONDITIONS

Weather conditions in Detroit are shown in Tables 94 and 95.

The most striking differences occur in 1909 and 1911. August was hotter in 1909 than in either 1910 or 1911. The mean for the month was higher, and also above the normal. In the other three months 1911 was in excess of 1909 by from 2° to 3°. Ninety-five degrees was never reached in 1909, while it was reached three times in 1910 and five times in 1911. The daily mean was never 85°, and only 80° or over three times in 1909, while in 1911 it was 85° three times and 80° or over thirteen times.

TABLE 94.—NUMBER OF DAYS ON WHICH THE MAXIMUM AND MEAN TEM-PERATURE REACHED CERTAIN FIGURES, DETROIT\*

		MAXIMUM	Mean		
	95° or +	90° or +	85° or +	85° or +	80° or +
1909					
June	0	0	3	0	0
July		1	5	0	0
August	0 0	2 0	3 5 8 2	0	0 3 0
September	0	0	2	0	0
September	_	_			_
	0	3	18	0	3
1910			10		
June	1	3	8	0	3
July	1 2 0 0	3 3 1	8 15 7	2	3 3 1 0
August	Õ	1	7	ō	1
August	ů.	Ô	Ó	ő	ô
September	_	_	_	_	_
	. 3	7	30	2	7
1011			30	-	
1911	2	2	6	0	2
June	2 3 0	3 8	12	0 3	3 8 2 0
July	0	1	12	0	0
August	0	0	12 7 2	0	2
September	U	0	2	U	0
	-	10	27		12
	5	12	27	3	13

TABLE 95.—MONTHLY MEAN TEMPERATURE, RAINFALL, AND NUMBER OF RAINY DAYS, DETROIT\*

	NORMAL	1909	1910	1911
[une				
Mean	67.4°	67°	66°	69°
Rainfall	3.80 in.	4.95 in.	1.47 in.	1.48 in.
Rainy days		13	9+0	9+7
uly				
Mean	72.1°	71°	74°	74°
Rainfall	3.50 in.	5.94 in.	1.48 in.	0.79 in.
Rainy days		9+2	6+5	9+0
August				
Mean	70.1°	72°	71°	71°
Rainfall	2.73 in.	3.75 in.	1.11 in.	1.31 in.
Rainy days		9+5	7+3	9+6
eptember				
Mean	63.5°	61°	63°	64°
Rainfall	2.55 in.	0.75 in.	2.02 in.	4.32 in.
Rainy days		8+2	10+2	14+4
COTALS				
Mean	68.2°	67.7°	68.5°	68.5°
Rainfall	12.58 in.	15.39 in.	6.08 in.	7.9 in.
Rainy days		39+9	32+10	41+17

Note.—Two figures are given for number of rainy days—the first, in front of the plus sign, is the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is number of days there was a trace of precipitation, but less than 0.01 inch.

<sup>\*</sup> Monthly Meteorological Summary, Local Weather Bureau, Detroit, and Monthly Weather Review, Washington.

June, 1910, was cooler than 1911 by 3°, and September by 1°. The other two months had the same mean, 1911 being above the normal in all four months.

Except in September the rainfall in 1911 was scanty, being well below the normal in each month. The first three months were drier than in 1909. July was much drier than 1910, and had very slightly greater rainfall in June and August.

It is in the months of July and August that the greatest reduction in infant

deaths occurred.

# THE CAMPAIGN

Until 1910 very little was done in Detroit for preventing infant mortality outside of the ordinary relief work done by charitable organizations. The Babies' Milk Fund distributed good milk, whole or modified, to patients referred by the various clinics and hospitals. No attempt was made to follow up these babies, nor was any home instruction given.

In 1910 the Health Department began a "mothers' clinic" under the charge of Dr. Duffield. The Visiting Nurses' Association assigned one nurse to this clinic, which attempted to secure proper feeding for babies and instruction for their mothers. The milk was distributed unmodified, and was prepared at home under supervision.

The "Children's Bureau" was organized this same year, composed of representatives of 18 different organizations; a trained social worker was employed, and coöperation and greater efficiency resulted.

In 1911 the Babies' Milk Fund started two milk stations in charge of a Medical Director, Dr. Cooley, with nurses in attendance. The Milk Fund gave up the modification of milk except in very exceptional cases, teaching home modification instead. Home visiting by the nurses is made a great feature of the work, and coöperation exists with the Visiting Nurses' Association. Consultations are held at the stations, attendance by the mothers being demanded. Classes have been started for mothers and young girls in infant hygiene, making of clothes, etc. Obstetric clinics are being started in connection with the stations, and promise to be valuable additions to the work.

The Mothers' Clinic of the Health Department increased its activities very largely in 1911. The school nurses were detailed to its work in the summer months. Births reported by midwives were visited, and the mother instructed.

A campaign of publicity was started to try to improve the milk supply. A milk exhibit was held under the auspices of the Board of Commerce and the Health Department, which resulted in greatly improving the supply through the raising of the standards required.

A considerable amount of work has been done, and very nearly all of it during the years 1910 and 1911. The educational *prevention* of infant mortality is being emphasized.

BUFFALO

The mortality under one year for Buffalo is shown in the following table:

TABLE 96.—MORTALITY UNDER ONE YEAR, BUFFALO\*

	1906	1907	1908	1909	1910	AVERAGE 1906-10	1911
January	84	39	107	110	109	90	96
February	85	118	110	101	76	96	82
March	92	114	96	77	111	98	94
April	97	109	100	70	135	102	97
May	80	100	77	89	123	94	88
June	77	79	73	64	67	72	104
July	130	290	154	136	139	170	97
August	221	242	174	192	254	216	103
September	67	152	141	117	173	130	90
October	99	92	144	98	109	108	70
November	93	73	77	85	91	84	63
December	121	87	74	114	95	98	66
YEAR Percentage all deaths	1246 <b>19.6</b>	1495 22.5	1317 21.1	1253 19.8	1482 21.3	1358 20.9	1050 20.7
July, August, and September  Percentage of year  Percentage all deaths	418 33.5 23.7	684 45.7 38.8	469 <b>35.6</b> <b>29.9</b>	445 35.5 27.5	566 38.2 32.0	516 38.0 30.5	290 27.6 20.0
Births	8075 154.3	8891 168.1	9171 143.6	9027 138.8	10,008 148.1	9272 146.5	10,546 99.5

The difference between 1911 and the other years is shown in the following table:

TABLE 97.—DIFFERENCE BETWEEN MORTALITY UNDER ONE YEAR IN 1911 AND PREVIOUS YEARS, BUFFALO

	1906	1907	1908	1909	1910
Year	-128 -5.9 -3.7	-445 -1.8 -394 -18.1 -13.8 -67.6	-267 -0.4 -179 -8.0 -9.9 -44.1	-203 +0.9 -155 -7.9 -7.5 -39.3	-432 -0.6 -276 -10.6 -12.0 -48.6

It will be seen at once that there was a most phenomenal decrease in the infant mortality in 1911. Only January and June are above the five-year average. The decrease is all in the second half of the year. The figures for the first and second half-years are:

<sup>\*</sup> These figures are taken from the monthly bulletin of the State Department of Health. They will be found to differ considerably in some cases from the Census Bureau figures, and from those of the Buffalo Department of Health Annual Reports, owing apparently to different methods in tabulating still-births.

TABLE 98.—MORTALITY UNDER ONE YEAR BY HALF-YEARLY PERIODS, BUFFALO

	1906	1907	1908	1909	1910	AVERAGE 1906-10	1911
First half-year		559 936	553 764	511 742	621 861	552 806	561 489

The striking situation exists of the second half of the year having a lower mortality than the first half; even more striking are the figures by trimesters:

TABLE 99.-MORTALITY UNDER ONE YEAR BY TRIMESTERS, BUFFALO

	1906	1907	1908	1909	1910	AVERAGE 1906-10	1911
First trimester	261 254 418	271 288 684	303 250 469	288 223 445	296 325 566	284 268 516	272 289 290
Fourth trimester	313	252	295	297	295	290	199

The percentages of the year's total deaths occurring in each of the four trimesters in 1911, as compared with the average for five years, are:

TABLE 100.—MORTALITY UNDER ONE YEAR BY TRIMESTERS, IN PERCENT-AGES, BUFFALO

	FIRST	SECOND	THIRD	Fourth
1911	25.91	27.52	27.62	18.95
	<b>20.91</b>	19.73	<b>38.00</b>	<b>21.36</b>

A difference of one-tenth of 1 per cent. is shown between the second and third quarters in 1911, as against 18.27 in the average. The third quarter in 1911 is 10.38 less than the average.

The percentage of all deaths for the year, however, is only 0.6 lower than 1910, and is 1.1 per cent. higher than 1906. For the summer months it is 12 per cent. lower than 1910, and 3.7 per cent. lower than 1906.

The infant mortality rate is from 39.3 to 67.6 lower than the preceding years, being 99.5, a very low figure.

If we estimate the deaths that would have occurred if the mortality had increased at the same rate as the population, based on the figures of 1906, the low year, we find:

TABLE 101.—YEAR'S MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED, BASED ON 1906 FIGURES, COMPARED WITH ACTUAL MOR-

TAI	1907	FFALO 1908	1909	1910	1911
Estimated mortality	$^{1495}_{+224}$	1296 1317 +21 +1.6	1321 1253 —68 —3.1	1347 1482 +135 +10.0	1372 1050 322 23.5

Average yearly increase in population, 2.02 per cent.

Estimating in the same way the summer mortality we find:

TABLE 102.—MORTALITY UNDER ONE YEAR, DURING THREE SUMMER MONTHS, WHICH SHOULD HAVE OCCURRED, BASED ON 1906 FIGURES, COM-PARED WITH ACTUAL MORTALITY, BUFFALO

	1907	1908	1909	1910	1911
Estimated mortality	426	435	443	452	460
	684	469	445	566	290
	+258	+34	+2	+114	—170
	+ <b>60.6</b>	+ <b>7.8</b>	+ <b>0.5</b>	+25.2	— <b>36.9</b>

There has been a most phenomenal decrease in infant deaths in Buffalo. The infant mortality rate has fallen to 99.5, and the actual figures for births are only 546 more than 1910, so that there cannot have been a very greatly increased reporting of births, in all probability.


# WEATHER CONDITIONS

Weather conditions in Buffalo are shown in the following tables:

TABLE 103.—NUMBER OF DAYS ON WHICH MAXIMUM AND MEAN TEMPERA TURES REACHED CERTAIN FIGURES, BUFFALO\*

		MAXIMUM	TEMPERATURE		MEAN TEMPERATURE			
	95° or +	90° or +	85° or +	80° or +	85° or +	80° or +	75° or +	
1909		<b>F</b>						
June	0	0	1	2	0	0	1	
July	0 0 0	0 0	0 2 1	2 7	0 0	0	1 3 6 1	
August	0	0	2	10	0	0	6	
September	0	0	1	3	0	0	1	
TOTAL	0	0	4	22	0	0	11	
1910								
June	0	0	1	6	0	0	1	
July	0 0 0	0	1 2 5 0	6 9	0 0	0	1 3 7 0	
August	0	0	5	12	0	0	7	
September	0	0	0	0	0	0	0	
Total	0	0	8	27	0	0	11	
1911								
	0	0	0	2	0	0	0	
June July	0	0 2 2 0	0 6 6	11	0	0 3 3	10	
August	Ô	2	6	11	Ô	3	10 9 0	
September	0	0	0	1	0	0	0	
TOTAL	1	4	12	25	1	6	19	

<sup>\*</sup> Monthly Meteorological Summary, Buffalo, and Weather Review, Washington, D. C.



Where Home Instruction is Needed



TABLE 104.—MONTHLY MEAN TEMPERATURE, RAINFALL, AND NUMBER OF RAINY DAYS, BUFFALO\*

	NORMAL	1909	1910	1911
June				
Mean	65°	64.6°	62.8°	66.3°
Rainfall	3.03 in.	1.05 in.	1.05 in.	2.23 in.
Rainy days		5+7	8+5	12+6
July		011	013	12+0
Mean	70°	68.6°	70.1°	71.6°
Rainfall	3.32 in.	2.49 in.	5.33 in.	4.41 in.
Rainy days	0.0.4 1111	10+3	8+4	8+7
August		10   0	011	0+1
Mean	69°	69.4°	69.4°	69.8°
Rainfall	2.95 in.	1.69 in.	2.72 in.	4.61 in.
Rainy days		8	8+3	11+4
September		0	073	1174
Mean	63°	61.7°	62.0°	62.2°
Rainfall	3.09 in.	2.61 in.	2.16 in.	3.81 in.
Rainy days		11+4	10+4	11+6
TOTAL		11 11	10 1.4	11+0
Mean	66.7°	66.07°	66.07°	67.4°
Rainfall	12.39 in.	7.84 in.	11.26 in.	15.06 in.
Rainy days	14.60 III.	34+14	34+16	42+23

Note.—Two figures are given for number of rainy days—the first, in front of the plus sign, is the number of days precipitation was 0.01 inch or more; the second, after the plus sign, is number of days there was a trace of precipitation, but less than 0.01 inch.

Temperatures above 90° are unusual in Buffalo. They were not recorded during the years 1909 and 1910. Figures for a maximum of 80° and a mean of 75° are, therefore, included in the table.

The year 1911, therefore, stands out as a year when there were periods of extreme heat, 90° or more being recorded on two days in both July and August. A mean temperature of 80° or more was also recorded in both July and August, which did not occur in either 1909 or 1910. A mean temperature of 75° was reached nineteen times in 1911 as against eleven times in the two previous years. The monthly means show 1911 above the normal in each month except September, and higher than either 1909 or 1910 for each month.

The rainfall except in June was above the normal in each month, and was greater than the corresponding months in 1909 and 1910, except in July, when less rain fell than in 1910.

Except in rainfall, weather conditions were unfavorable for a low infant mortality.

#### THE CAMPAIGN

It has been difficult to obtain exact information in regard to the activities being carried on in Buffalo. In a general way everybody seems to be agreed that during the last year some very active work has been done by the Health Department and social organizations. The problem of the milk supply is being very actively handled by the Health Commissioner, Dr. Fronczak. The Children's Welfare Society and the Babies' Milk Dispensary are both very active in their campaign. Considerable coöperation exists between the various agencies, and the Health Department is doing all in its power to increase this coöperation.

\* Monthly Meteorological Summary, Buffalo, and Weather Review, Washington, D. C.

# CHAPTER VII

# General Summary

For convenience in reference the mortality in the various localities studied is summarized in the following tables.

Table 105 shows the highest, lowest, and difference between same, for the years 1906–10 inclusive, in the year's actual mortality under one year; the percentage of all deaths, and the infantile mortality rate; the actual deaths, percentage of all deaths, and proportion occurring in this period for the third quarter; and the relation of the 1911 figures to the highest, lowest, and 1910 figures in each of these groups.

Table 106 shows the same thing expressed in percentages. By "Extremes" is meant the difference between highest and lowest in the period 1906–10 expressed in percentage of the highest. The yearly average increase in population is also noted.

A consideration of these tables shows that:

- 1. In actual yearly deaths-
  - Manhattan, Brooklyn, City of New York, Philadelphia, St. Louis, Baltimore, and Buffalo show a decrease in 1911 from the lowest actual mortality recorded in the previous five-year period.
  - In the amount of this decrease from lowest figures, expressed in percentages, the figures for Manhattan are exceeded only by Baltimore and Buffalo.
  - "Rest of City," Chicago, Boston, Cleveland, Pittsburgh, and Detroit show an increase in 1911 from this lowest figure.
- 2. In the ratio of infant deaths to total deaths for the year, in 1911—
  - All but Buffalo show a decrease from the lowest figure recorded. In actual figures the proportion of decrease is greater in Manhattan than in any of the other localities except Cleveland, Pittsburgh, and Detroit.
  - In the percentage decrease from the lowest previous figure, Manhattan is exceeded only by Philadelphia and Cleveland (8.8 and 8.6 respectively, as against 7.8 for Manhattan).
- 3. In infant mortality rate for 1911-
  - Manhattan is higher than any other city in actual figures except
    - Boston, Cleveland, and Detroit.
  - A decrease from the lowest previous rate is noted in all except Boston, which is 6.2 higher than its lowest previous figure, and Cleveland, which is 11.4 higher. These figures, however, are of little real value in most cases, as birth registration has only within the last year or two received any attention from the authorities.

TABLE 105.—INFANT MORTALITY. COMPARATIVE TABLE BASED ON STATISTICS FOR YEARS 1906-11.—

ACTUAL CHANGES

BUFFALO	1495 1246 249	22.6 19.6 3.0	168.1 138.8 29.3	684 418 266	38.8 23.7 15.1	45.7 33.5 12.2
TIORIZA	2095 1598 497	29.5 26.4 3.1	216.5 181.1 35.4	708 547 161	38.5	34.4 32.6 1.8
Ріттѕвивсн	2251‡ 1804‡ 447‡	24.4† 23.0† 1.4‡	100100100	747‡ 560‡ 187‡	33.3‡ 29.8‡ 3.5‡	33.2‡ 29.2‡ 4.0‡
ВАГТІМОВЕ	2443‡ 2148‡ 295‡	21.8‡ 19.9‡ 1.9‡	100100100	1006‡ 752‡ 254‡	34.0‡ 28.0‡ 6.0‡	41.1‡ 35.0‡ 6.1‡
CLEVELAND	2057 1711 346	26.11 24.5 1.6	185.8 130.6 55.2	774 554 220	35.3 30.1 5.2	37.6 30.8 6.8
возгои	2742 2125 617	23.3 19.2 4.1	148.0 119.8 28.2	1010 741 269	34.0 27.2 6.8	25.9 21.3 4.6
Sr. Louis	1814 1707 107	17.9 15.7 2.2	100100100	663 548 115	25.7 21.4 4.3	38.2
Ригловгрніл	5781 4870 911	21.1 19.3 1.8	166.2 129.7 36.5	1777† 1645† 132†	28.4† 27.9† 0.5†	33.9† 33.7† 0.2†
Сніслео	6888 6114 774	22.6 20.5 2.1	100100100	2209 1923 286	29.2 23.8 5.4	32.2
CITY OF NEW	17437 15976 1461	22.5 21.1 1.4	153.7 125.6 28.1	6502 5154 1348	32.3 29.4 2.9	37.3 32.3 5.0
"REST OF	2332 2139 193	20.4 17.7 2.7	159.6 110.0 49.6	1034 781 253	32.4 25.7 6.7	45.1 36.5 8.6
Вкооктии	5503 4923 580	21.8 19.7 2.1	157.8 118.4 39.4	2191 1733 448	32.5 28.5 4.0	39.8 34.9 4.9
МАТТАНИАМ	9602 8914 688	23.9 23.1 0.8	150.2 134.9 15.3	3278 2640 638	33.1 30.9 2.2	34.1 29.6 4.5
EXTREMES, 1906-10	Year Actual: Highest Lowest	Percentage all deaths: Highest Lowest. Difference.	Infant mortality rate: Highest. Lowest. Difference.	JULY-AUGUST-SEPTEMBER Actual: Highest. Lowest. Difference.	Percentage all deaths: Highest Lowest Difference	Percentage years deaths under one: Highest

\* "Rest of City" means Boroughs of Bronx, Queens, and Richmond combined. † Based on figures 1909–11.

§ Not accurate enough for any use.

TABLE 105.—INFANT MORTALITY. COMPARATIVE TABLE BASED ON STATISTICS FOR YEARS 1906-11.— ACTUAL CHANGES.—(Continued.)

BUFFALO	1050 445 196 432	$^{20.7}_{-1.9}$	99.5 68.6 39.3 48.6	290 -394 -128 -276	20.0 -18.8 -3.7 -12.0	27.6 -18.1 -5.9 -10.5
тюятаП	1862 -233 +264 -233	26.1 -3.4 -0.3 -2.0	139.2 -77.3 -41.9 -42.8	638 -70 +91 -70	34.0 +4.5 +0.7 -1.3	34.3 -0.1 +1.7 +0.5
Ритевивен	1810 —4411 +6‡ —441	22.3 -2.1‡ -0.7‡ -1.1	LOT LOT LOT	591 -156 +31 -156	27.7 —5.6‡ —2.1‡ —5.6	32.6 -0.6‡ +3.4‡ +0.4
ВАСТІМОВЕ	1958 -485‡ -190‡ -190	18.8 -3.0 -1.1 -1.1	conconcon	716 —290‡ —36‡ —36	28.2 -5.8‡ +0.2‡ +0.2	36.6 -4.5‡ +1.6‡ +1.6
Сгелетуир	1788 —269 +77 —269	22.4 -3.7 -2.1	142.0 -43.8 +11.4 -10.9	569 —205 +15 —205	28.5 -6.8 -6.8	31.8 -5.8 -1.0 -5.8
возгом -	2245 	19.1	126.0 -22.0 +6.2 -0.7	764 —246 +23 +15	25.1 -8.9 -2.1	33.6 -1.9 +2.7 +2.7
smod .rs	1579 —235 —128 —128	14.5 -3.4 -1.2	101.6	699 —124 —27	20.5 -4.6 -0.3	34.1 -4.1 +2.5 +0.9
Ригьрегриіл	4616 1265 254 616	17.6 -3.5 -1.7	112.5 -53.7 -17.2 -23.8	1537 240† 108† 240	25.0 -3.4† -2.9† -2.9	33.2 -0.7† -0.5† -0.5
Сніслео	6309 -579 +195 -532	19.3 -3.3 -1.2	101101101101	1904 305 305	24.7 4.6 +0.8 +0.8	30.2 -2.1 -0.6 -2.1
CITY OF NEW YORK	15017 -2420 -959 -1198	19.9 -2.6 -1.2	111.6 -42.1 -14.0 -14.0	4593 -1909 561 981	25.4 -6.9 -4.0 -4.3	30.6
"REST OF	2196 —136 +57 —6	17.5 -2.9 -0.2 -0.2	98.6 -61.0 -11.4	800 234 +19 54	24.3 -8.1 -1.4 -2.2	36.4 -8.7 -0.1 -2.3
BROOKLYN	4629 —874 —294 —430	18.9 -2.9 -0.8	101.3 —56.5 —17.1 —17.1	1499 —692 —234 —269	25.9 -6.5 -2.5	32.3 —7.4 —2.5 —2.5
илтинимМ	8192 —1410 —722 —762	21.3 -2.6 -1.8 -1.8	123.1 —27.1 —11.8 —11.8	2294 —984 —346 —658	25.7 -7.5 -5.3 -5.8	28.0 —6.1 —1.6 —4.9
EXTREMES, 1906-10	YEAR Actual, 1911. Difference from highest. Difference from lowest. Difference from 1910.	Percentage all deaths, 1911  Difference from highest Difference from lowest  Difference from 1910	Infant mortality rate, 1911  Difference from highest  Difference from lowest  Difference from 1910	JULY-AUGUST-SEPTEMBER Actual, 1911. Difference from highest Difference from lowest. Difference from 1910.	Percentage all deaths, 1911.  Difference from highest.  Difference from lowest.  Difference from 1910.	Percentage years deaths under one, 1911 Difference from highest Difference from 1910

\*"Rest of City" means Boroughs of Bronx, Queens, and Richmond combined.

† Based on figures 1909-11.

124

BUFFALO	16.6 -29.8 -15.7 -29.2	13.3 +5.6 -2.8	17.4 40.8 -28.3 -32.8	38.9 -57.6 -30.9 -38.8	38.9 -48.4 -15.6 -37.5	26.7 -39.6 -17.6 -27.6 2.02
DETROIT	23.7 -11.1 +16.5 -11.1	10.5 -11.5 -1.1	16.3 —35.7 —23.1 —23.5	22.7 —9.9 +16.6 —9.9	13.5 -11.7 +2.1 -3.7	
нэялязтич	19.8‡ -19.6‡ +0.3‡ -19.6	8.77 -8.64 -3.04	conconcon	25.0‡ —20.9‡ +5.5‡ —20.9	10.5‡ -16.8‡ -7.0‡ -16.8	12.0 -0.2 +11.6 +1.2 1.82
BALTIMORE	12.11	8.77 -13.85 -5.11	conconcon	25.21 -28.81 -4.81 -4.8	17.6‡ -17.0‡ +0.7‡ +0.7	14.8‡ +4.6‡ +4.6 0.97
Стелегуир	16.8 -13.1 +4.5 -13.1	6.1 -14.2 -8.6 -12.5	29.7 -23.5 +8.7 -7.1	28.4 -26.5 +2.7 -26.5	14.7 -19.2 -5.3 -19.2	18.1 +3.2 +3.2 -15.4 4.69
котгоя	22.5 -18.1 +5.6 -0.2	17.6 -18.0 -0.5 -1.5	19.0 +5.2 -0.5	26.6 -24.3 +3.1 +2.0	20.0 -26.2 -7.7 -7.7	17.7 -7.3 +12.7 +12.7 1.96
St. Louis	5.9 12.9 7.5	12.3 —19.0 —7.6 —7.6	101101101101	17.3 -18.7 -1.6 -4.7	16.7 -17.9 -1.4 -5.4	17.3 +7.9 +2.7 1.94
Ригловерніл	15.7 —21.8 —5.2 —11.7	8.5 -16.5 -8.8 -8.8	21.9 -32.3 -13.2 -17.5	7.5† —13.5† —6.5† 13.5	-11.9 $-10.3$ $-10.3$	-2.0 -2.0 -1.4 -2.0 1.97
Сніслео	11.0 -8.4 +3.1 -7.7	9.2 -14.6 -5.8 -5.8	100100100100	12.9 -13.8 -0.9 -13.8	18.4 +3.3 +3.3	4.6 -6.5 -1.9 -6.5 2.87
CITY OF NEW YORK	8.3 13.9 6.0 7.4	9.3 -11.5 -5.7	18.3 -27.4 -11.1 -11.1	20.7 -29.3 -10.9 -17.6	9.0 -21.3 -13.6 -14.5	13.4 —17.9 —5.3 —11.0 3.87
" REST OF	8.3 + 2.6 - 0.3	12.8 -13.8 -1.1	31.1 -38.2 -10.4 -10.4	24.6 -22.8 +2.4 -6.3	20.7 -25.0 -5.4 -8.3	19.2 -19.5 -0.3 -5.9 9.038
BROOKLYN	10.5 -15.9 -6.0 -8.5	9.9 -0.4 -0.4	25.0 -35.8 -14.4 -14.4	20.4 -31.6 -13.5 -15.2	12.3 -20.0 -8.8 -8.8	12.3 -18.6 -7.2 -7.2 4.01
МАПТАНИАМ	7.1 -14.6 -8.1 -8.5	3.3 -10.9 -7.8 -7.8	10.2 -18.0 -8.7 -8.7	19.4 -30.0 -13.1 -22.3	6.6 -22.6 -17.1 -18.4	13.1 —17.8 —5.4 —14.9 2.602
	Actual: Extremes	Extremes. 1911 from highest. I lowest. I lowest.	Extremes	Extremes 1911 from highest lowest 1915	Extremes.  1911 from highest.  " lowest.  " 1910.	Extremes

\* "Rest of City" means Boroughs of Bronx, Queens, and Richmond combined. † Based on figures 1909–11.

§ Not accurate enough for any use.

The percentage decrease in Manhattan is the smallest of any which show a decrease. The real value of this is also small, as stated above.

4. In actual deaths during the third quarter of the year 1911—

Manhattan, Brooklyn, New York City, Chicago, Philadelphia, St. Louis, Baltimore, and Buffalo show a decrease from the lowest previous year.

"Rest of City," Boston, Cleveland, Pittsburgh, and Detroit show

an increase from the lowest previous figure.

Manhattan shows a greater percentage of decrease from the lowest previous figures than any locality except Brooklyn and Buffalo.

- 5. In ratio of infant deaths to total deaths occurring in third quarter of 1911—
  - There is a decrease from the lowest previous figures in all save Chicago, Baltimore, and Detroit.

In actual decrease from lowest previous figure Manhattan is exceeded only by Cleveland, Baltimore, Pittsburgh, and Detroit.

- Manhattan shows the greatest percentage decrease from lowest previous figure—17.1 per cent. as against 15.6 per cent. for Buffalo.
- In percentage of year's infant mortality occurring in third quarter of 1911—
  - There is a reduction from the lowest previous figures in Manhattan, Brooklyn, "Rest of City," New York city, Chicago, Philadelphia, and Buffalo.

An increase is noted in St. Louis, Boston, Cleveland, Baltimore,

Pittsburgh, and Detroit.

Manhattan shows the lowest actual figure of any locality except Buffalo—28 as against 27.6 per cent.

The percentage of decrease from lowest previous figure in Man-

hattan is exceeded only by Brooklyn and Buffalo.

7. Leaving out of consideration the infant mortality rate which is absolutely unreliable for comparative study over a period of years—

Manhattan shows a greater percentage reduction than Chicago, St. Louis, Boston, Pittsburgh, and Detroit in every respect. Philadelphia and Cleveland have a better showing only in the ratio of infant deaths to total deaths for the year. Baltimore exceeds only in the percentage decrease of actual deaths for the year.

Buffalo has a better showing in each respect except in the ratio of infant deaths to total deaths for the year and for the third

trimester.

The percentage reduction in mortality for the third trimester is greater in Manhattan than in any other region studied. Compared with the "Rest of the City," Manhattan makes a better showing in reduction in every point.

There is still another method of estimating results based upon the method of comparison with previous low records, allowing for increase of population. The following tables are constructed to put the 1911 mortality figures to as severe a test as possible.

Taking the low record total for the year in each city and allowing for increase in population, we have the following:

TABLE 107.—MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED IN 1911, BASED ON LOW FIGURE PREVIOUS TO 1910, ALLOWING FOR INCREASE IN POPULATION, COMPARED WITH ACTUAL MORTALITY IN 1911 IN VARIOUS LOCALITIES

	LOWEST PREVIOUS YEAR'S TOTAL	YEAR	INCREASE IN POPU- LATION	ESTIMATED MORTALITY	MORTALITY, 1911	DIFFERENCE	PER CENT. DIFFERENCE
Buffalo	1246	1906	10.10	1371	1050	- 321	-23.5
St. Louis	1715	1907	7.76	1848	1579	- 269	-14.5
Baltimore	2215	1908	2.91	2279	1958	- 321	-14.0
Philadelphia	4597	1903	15.76	5321	4616	- 705	-13.2
Chicago	6114	1906	14.35	6991	6309	- 682	-10.1
Detroit	1598	1907	25.20	2000	1862	- 138	- 6.9
Pittsburgh	1804	1907	7.28	1935	1810	— 125	- 6.4
Cleveland	1711	1909	9.38	1871	1788	— 83	- 4.4
Boston	2125	1909	3.92	2208	2245	+ 37	+ 1.7
Brooklyn	4601	1903	32.08	6077	4629	-1448	-23.8
"Rest of City"	1631	1903	72.30	2810	2196	- 614	-21.8
New York City	14413	1903	28.56	18529	15017	-3512	-20.4
Manhattan	8184	1903	20.81	9884	8912	-1692	-17.1

Of the first nine cities in the list, Buffalo alone has a greater reduction from this estimated mortality than Manhattan. Brooklyn, the "Rest of the City," and New York as a whole had a greater reduction than Manhattan.

If, now, we compare the mortality during the three summer months in the same way, we see the following results:

TABLE 108.—MORTALITY UNDER ONE YEAR WHICH SHOULD HAVE OCCURRED DURING THE THREE SUMMER MONTHS OF 1911, BASED ON PREVIOUS LOW FIGURES, AND ALLOWING FOR INCREASE IN POPULATION, COMPARED WITH ACTUAL MORTALITY IN 1911 IN VARIOUS LOCALITIES

	ESTIMATED MORTALITY	MORTALITY IN 1911	DIFFERENCE	PER CENT. DIFFERENCE
Buffalo	502	290	- 312	-63.9
Chicago	2198	1904	- 294	-13.4
Detroit	719	638	- 81	-11.3
Philadelphia	1709	1537	- 172	-10.1
Cleveland	606	569	- 37	- 6.1
Baltimore	759	716	<b>—</b> 43	<b>—</b> 5.7
St. Louis	740	699	<b>—</b> 41	- 5.5
Pittsburgh	600	591	- 9	- 1.5
Boston	770	764	- 6	- 0.8
Brooklyn	2185	1499	- 686	-31.4
City of New York	6576	4593	-1983	-30.1
"Rest of City"	1120	800	- 320	-28.6
Manhattan	2787	2294	<b>— 493</b>	-17.9

From this estimated summer mortality, Buffalo again alone has a greater percentage difference than Manhattan, except for the city of New York as a

whole, Brooklyn, and the "Rest of the City." It is this summer mortality which has been most influenced by the campaign against infant mortality.

Another even more severe test of results may be applied. If we take the lowest recorded proportion of the year's deaths occurring in the third quarter and apply it to the lowest year's total, we should in most cases have still lower figures for a basis. For example in 1903 in Manhattan the year's total under one year was 8181, the lowest recorded in the ten-year period 1901–1910. In 1909 the proportion of the year's mortality falling in the third quarter was 29.6, the lowest recorded. Taking then 29.6 per cent. of 8181, we get the optimum figures for comparison. The following table shows these figures:


TABLE 109.—INFANT MORTALITY IN THE THIRD QUARTER OF 1911 COM-PARED WITH ESTIMATED MORTALITY BASED ON OPTIMUM FIGURES FOR PERCENTAGE OF YEAR'S MORTALITY AND TOTAL YEAR'S MORTALITY

	ESTIMATED MORTALITY	1911 MORTALITY	DIFFERENCE	PER CENT. DIFFERENCE
Buffalo	459	290	- 179	-39.0
Philadelphia	1793	1537	- 256	-14.3
Chicago	2146	1904	- 242	-11.3
Baltimore	797	716	- 81	-10.2
Detroit	652	638	- 14	- 2.1
Cleveland	576	569	<b>—</b> 7	- 1.2
Pittsburgh	565	591	+ 26	+ 4.6
St. Louis	583	699	+ 116	+19.8
Boston	470	764	+ 294	+62.5
Brooklyn	2120	1499	- 621	-29.2
New York City	6096	4593	-1503	-24.6
"Rest of City"	1025	800	— 225	-21.9
Manhattan	2925	2294	<b>— 631</b>	-21.5


Only six of the cities outside of New York can show a decrease; Pittsburgh, St. Louis, and Boston show an increase—Boston, of 62.5 per cent. Manhattan again exceeds all but Buffalo, which again ranks first.

A study of charts shown in Exhibits 3 to 7, comparing daily deaths with temperatures in Manhattan and Brooklyn, is of interest in connection with the theories of Liefmann and Lindemann, as mentioned in Chapter I. Their theory is that the summer mortality among babies is due to two chief causes. The first of these is the directly fatal effect of high atmospheric temperature, which results in sharp rises in the mortality curve following twenty-four to forty-eight hours after the rise in temperature. These rises, followed by sharp falls to the "normal," occur chiefly in early summer. In late summer another factor is added. The basal line, so to speak, is continually high, and represents practically the diarrheal deaths. On this basal curve occur the rapid rises and falls in the mortality curve immediately following hot periods.

These conditions are very plainly seen in plates showing the temperature and mortality in Manhattan in 1909 and 1910 and Brooklyn in 1910. In 1911, however, the mortality curve is a very different one. Sharp rises and falls do occur, but the diarrheal element is markedly diminished. Heat deaths occur, but the



TEACHING HOME MODIFICATION OF MILK



high basal line representing diarrhea is down nearer the normal. Also the sharp rises are not so great, which tends to show the absence of the predisposing element resulting in great susceptibility to high temperature.

Briefly summarizing weather conditions in 1911 in a general way, temperature was decidedly unfavorable for a low infant mortality throughout the country in July and early August. In St. Louis heat was excessive and continued throughout the summer. Detroit and Buffalo especially are unused to such temperatures as occurred in 1911.

Rainfall was not so uniformly unfavorable as temperature. In Chicago, Philadelphia, and Baltimore, while below the normal, it was not nearly so scanty as in other places. In Boston and Buffalo it was above the normal. In New York and the other cities studied it was very scanty during July and early August, especially in St. Louis.

It is rather difficult to gage the activities of health departments and private organizations. In a general way all the cities discussed, except Boston, were more active in 1911 than ever before. In Greater New York, Manhattan and Brooklyn were very decidedly more active. Never in their history has so great and general an effort been put forth, and these boroughs were fairly well cared for. Chicago and Philadelphia profited by previous experience, and their campaigns were more efficient and extensive. St. Louis also extended its campaign, as did Detroit. Baltimore added home instruction as an important feature of its work.

Buffalo made the most remarkable showing of any city studied. The Health Department had begun a vigorous campaign for pure milk, and the milk stations of the Babies' Milk Dispensary were active.

Boston is the striking exception to the general rule of reduction. Considerable effort has been made to find out conditions existing there in 1911, and apparently there has been absolutely no increase in effort nor expansion of the work over past years. The weather conditions were no worse comparatively than in other places, and in rainfall they were better than most others. With these conditions went a stationary infant mortality compared with 1910, and a poorer showing compared with previous years than most of the other cities record.

#### CONCLUSIONS

From the information gained through experience and study of the work of reducing infant mortality the following conclusions seem to be justified:

- 1. The reduction of infant mortality in 1911 in the cities studied was very marked, especially during the summer months. During the first part of the summer meteorological conditions was not favorable to a low mortality.
- 2. This reduction was not due to chance, but bears a close relation to the activity of the campaign for the reduction of infant mortality. The difference between cities in infant mortality rates is probably more a matter of public conscience and quality of official endeavor, than weather and character of population.
- 3. The full effects of any campaign will not be seen immediately. Dr. Robertson, Health Officer of Birmingham, England, said at the conference at Caxton Hall that he would be satisfied if he got results after ten years' work.

- 4. In the education of the mother in the care of herself and her baby we have the strongest weapon for fighting infant mortality.
- 5. The results in the milk stations, as shown by the detailed histories of 3182 babies, has proved the usefulness of the milk station in the reduction of infant mortality. It is one of the most efficient forces in the educational prevention of sickness and deaths among babies.
- 6. Its field of usefulness can be very greatly extended into other branches of infant and child welfare, such as prenatal work; the care of children under school age; follow-up work in inspection of school-children; supervision of boarded-out babies and midwives, etc. Stations should be maintained all the year round in order to make their value cumulative.
- 7. The sphere of influence of a milk station is limited, so that many are needed in a large city.
- One nurse for every 100 babies is the limit for the maximum of efficiency, and probably 300 babies the limit for any one station.
- Compulsory attendance under penalty of stopping milk supply should be insisted on.
- 10. When efficiently managed, the milk station is an economical means of forwarding the movement for infant welfare.
- 11. The results of the methods employed at the Committee's milk stations prove conclusively the entire feasibility of teaching home modification of milk, thus not only doing away with expensive laboratory plants, but making the mother eventually independent of the station.
- 12. The milk station can and should be a valuable means of encouraging and making possible maternal nursing. It need not be an encouragement to artificial feeding.
- 13. Private agencies are warranted in initiating milk station work, but their program should provide for having the municipality assume charge as soon as practical. The activities which are possible for the stations are proper functions for the municipality. Close coöperation between health authorities and private organizations is entirely feasible and necessary.
- 14. The education of the "little mothers" in the care of the baby will insure not only immediate results, but will pave the way for better motherhood.

#### CHAPTER VIII

# Extension of Relief Through Milk Stations

Because ignorance, poverty, squalor, and sickness in the home lessen the chance of the baby to survive a heated summer, it is obvious that a well-organized campaign to reduce infant mortality must provide ways and means for doing a great deal more than giving advice in the homes where these conditions exist. If the campaign is confined to babies living under average home conditions with mothers anxious for instruction, only a moderate reduction in the death-rate can be expected, because it is not the baby in the average home who is most likely to die. The mother who comes to a milk station of her own accord with the well baby to learn how to keep it well is not the mother most likely to lose her baby. The mother most needing the influence of the milk station is the one with whom the nurse has to work hard to have her baby enrolled because of ignorance and bad home conditions. When these mothers are visited, the station nurse must be prepared to solve the problem of the home; she must be able to suggest the proper solution for each problem presented to her. To do this, the nurse must know the facilities available, be they private or public, and act. If she acts wisely, she gains the mother's confidence and cooperation in giving the baby a chance to live; visits and advice where substantial relief is needed avail nothing if the relief is not to be obtained. If the father needs hospital care, it must be provided; if out of work, he must be brought in touch with the proper agency dealing with such cases; if the children are undernourished and sickly, an agency prepared to provide the necessities must be found.\* While the station nurse may be particularly solicitous for the welfare of the infant member of the family, yet she must of necessity deal with the whole situation. Parents so destitute that they are unable to buy milk for the baby are not cases for free milk alone, but for general relief.

Provision for extending relief to needy cases coming under the supervision of the milk station was recognized as an important factor in the demonstration. Relief in the form of milk, medical attendance, nurse service, ice for properly keeping the milk, hospital accommodations, and outings for sick mothers and babies are needed in addition to the general relief; of these, free milk for babies of poor mothers was considered most vital to the success of the demonstration.

At first it was planned to raise a large relief fund to supply free milk to all worthy cases. This was abandoned because it was thought best to make the demonstration under conditions which might be expected to exist another year if the work of conducting a large number of milk stations should be undertaken by

<sup>\*</sup> Attention is called to Table 116, showing the living conditions of 2156 families having babies enrolled at the Committee's stations.

the city. As the city could hardly be expected to give free milk, its stations would have to rely upon the organized relief agencies to provide milk for the poor. Another reason for abandoning the original plan of supplying free milk was the attitude of many of the Committee's larger contributors, who urged that they were also substantial contributors to established relief agencies better prepared not only to investigate the needs of applicants, but to extend all the relief necessary.

A From	APPL	ICATION FOR RELIEF	то	5435
		Date		1911
В			ETURNED BY RELIE ARTERS OF REFERR	
		Date		1911
C From	APPL	ICATION FOR RELIEF	то	5435
		Date		1911
Surname	Address	Floor	No. Rooms	Rent
FIRST NAME	AGE	OCCUPATION	INCOME	HEALTH
Woman				
Children				
Amount of milk now purchased	by family		Price	
Relief needed				
Form No. 114 K.		(Signed)		Nurse

Ехнівіт 19

In April letters were addressed to the superintendents of the various established relief organizations, in which the position of the Committee and its contributors was made known. This brought about several conferences between relief-giving organizations and the agencies conducting milk stations. The relief organizations agreed to make special provision for the prompt handling of cases brought to their attention and furnish free milk to babies wherever necessary. A

# EXTENSION OF RELIEF THROUGH MILK STATIONS 133

uniform system of reporting cases was adopted in order to avoid delay in extending milk relief and to eliminate duplication of effort in the investigation of home conditions of applicants.

Milk relief cases were divided into two classes:

- 1. Mothers having babies registered at the station who were unable to pay the difference in price between loose milk sold at the grocery and the bottled milk dispensed at the station; in such cases the organization extending the relief agreed to pay the difference in price of the two grades of milk—these were termed "partial relief" cases.
- Mothers having babies registered at the station and unable to bear any part of the cost of the milk required by the baby—these were designated "whole relief" cases.

It was agreed in both instances that the period during which relief would be extended should depend upon the emergency. Exhibit 19 for reporting applications for milk relief was adopted.

This record and reporting form is divided into three parts, each part bearing the serial number of the case. Whenever an application for relief was made through the milk station, the part marked "C" was filled out at the station from the information obtained by the nurse; wherever possible the information reported on this form was verified by a personal visit of the nurse to the home of the applicant. If the mother applying for relief had already registered her baby at

YOU ARE AUTHORIZED TO CHARGE
FORQUARTS OF MILK DAILY FROM
FOR
NEW YORK MILK COMMITTEE Date:

the station, the information recorded upon this application was verified by consulting the case history card described in a previous chapter. After completing this record the nurse forwarded the form to the Committee's central office; here the application was handled by a special relief agent formerly connected with one of the organized relief societies. If, in the opinion of the central office, the case was one requiring relief, form "A" was made out and forwarded, together with the form "B," to the agency which was expected to furnish the relief. Applications from Hebrew families were invariably sent to the United Hebrew Charities, and all other applications reported upon this form were forwarded to the Joint Registration Bureau, maintained by the Association for Improving the Condition of the Poor and the Charity Organization Society, where it was determined to which society the case properly belonged. The action taken by the agency was recorded on form "B" and returned to the central office of the Committee, where it was filed with the copy of the application, form "C." The station nurse was then notified as to extending relief to the cases on the postal card form shown in Exhibit 20.

#### CHARACTER OF CASES REFERRED

The character of cases referred to organized agencies for relief were those principally of mothers having ill-nourished or sick babies, due largely to their inability to pay the price of a pure, clean milk, and who, because of their poverty, were endeavoring to rear their babies on either condensed milk or milk purchased from the "corner grocery," which, through an investigation made by the Committee, was found to be entirely unfit for use in the feeding of infants. In some instances families were already receiving relief in other forms, but nothing had been done toward supplying the baby with the right kind of food.

Even where the family income was sufficient to supply the baby with the cheap "dipped" milk of uncertain quality, it was difficult for the nurse immediately to bring the mother to recognize the danger in giving her baby this milk; however, once an improvement was shown in the baby's health, it was easier to persuade the mother to purchase the milk from the station. If the milk relief could be extended for a certain period in such cases, to the extent of paying the difference between the price of loose milk (5 cents a quart) and the price of the milk dispensed at the station (7 cents a quart), the baby would be brought under the influence of the station, and upon its showing improvement the mother would consent to economize in other ways so as to continue giving good milk to the baby. No cases of this character were referred for relief excepting where extreme ignorance existed in the homes, and the baby, undernourished or sick, required immediate action by the nurse in order to save its life; these were termed "emergency relief" cases and were given either partial relief or entire milk relief.

Cases requiring the attention of a relief organization, where the home conditions were deplorable, but where the baby, in spite of a bad milk supply, was in no immediate danger, were referred, and the report of the agency awaited before any relief in the form of milk was given. It might be stated that in the "emergency relief" cases the nurse was authorized to extend temporary relief until the applica-

tion had been acted upon by the relief agency. If, however, the relief agency refused to act, the station was reimbursed by several individuals particularly interested in emergency cases found by the milk station.

Cases which the relief organizations rejected were thoroughly investigated by the special relief agent of the Committee, and if new facts were brought to light, the cases were again presented to the relief agency with a full statement of the conditions. If, after reopening a case, no aid could be obtained and the facts were still found to justify the extending of relief, the case was brought to the attention of persons sufficiently interested to supply it. Our only purpose in doing this was to save the baby even at the expense of breaking some of the established rules of dealing with the extension of relief in case of unworthy parents.

Owing to the most excellent cooperation existing between the milk stations and the relief agencies, comparatively few cases considered worthy by the Committee were refused aid. The following is an example of a case refused relief in the form of milk for the baby, although considered worthy by the Committee:

Case No. 766.—An Italian family consisting of father, mother, and two children, one aged twelve years and the other four months, came to the notice of the Mulberry Street milk station nurse; it was regularly reported through the central office to the Charity Organization Society. The father, a laborer, earned enough to cover the rent of three rooms shared with another family; rental, fourteen dollars; mother sews on pants whenever able; had been ill four months from heart trouble; father able to obtain employment only two days a week, and under medical treatment for skin disease. The mother receives six cents a pair for finishing trousers—a maximum earning of two dollars per week; father's earnings two and a half dollars per week; baby in poor physical condition, suffering from malnutrition and marasmus, artificially fed, and using grocery milk.

An investigator of the relief agency reported that the father was earning six to eight dollars a week and the mother three dollars a week, and there seemed no reason why the family should not manage without outside help; but further investigation by the Committee confirmed the report rendered by its station nurse, and

milk relief was furnished through other sources.

On the other hand, we have a station nurse forwarding an application (Case No. 781) from a woman who had been deserted by her husband and who had her only baby registered at the station. Investigated by the relief agency, it was discovered that the woman was already receiving free milk from a diet kitchen, and she acknowledged that in wanting another free quart from the milk station she had misrepresented her circumstances.

To illustrate the effectiveness of close supervision over relief work done this summer the case of a Bohemian family who begged for milk relief but two weeks after their admittance to the country might be cited. At that time the man was earning four dollars a week. The case was reported in the usual way to a relief agency, which brought the matter to the attention of the Commissioner of Immigration. Investigation from that quarter aroused the head of the family to greater efforts. He soon began to earn twelve dollars a week, and the family was made self-supporting.


The care and good judgment exercised by the station nurse in investigating home conditions in the case of applicants for relief are shown by the fact that out of 648 cases referred to three relief agencies during the period May 1st to November 1, 1911, 482, or 74.3 per cent., were accepted by them as proper cases for relief.

Naturally, it might be expected that, by establishing a large number of milk stations throughout the city, many mothers would take advantage of the situation to obtain a free supply of milk for their babies during the summer, even though they were able to pay for it. However, such was not the case; in spite of continued canvassing by nurses who repeatedly urged the mothers to bring their babies to the station for supervision, the number of babies receiving free milk in relation to the enrolment was even less during the demonstration than had been the experience of the Committee's stations during the previous year. In three stations operated by the Committee during the year 1910 the number of registered babies receiving free milk amounted to 16.3 per cent. of the total enrolment, while during the period of the demonstration,-May 1 to November 1, 1911,-out of a total registration of 7919 different babies, only 724, or 9.1 per cent., received free The comparatively small number receiving milk relief is due mainly to the care exercised by the nurse in handling applications for relief in a quiet manner, and not letting it become generally known among the mothers that some were being aided. The willingness of relief organizations to grant relief in worthy cases was 'not taken advantage of as a means of increasing the station enrolment.

Table 110, giving by stations and nationalities the number of cases applying for free milk, in relation to enrolment, shows that the number of applicants for free milk at the different stations varied from 1.5 to 35.8 per cent. of the total enrolment.

This statement also shows that the percentage of cases applying for free milk varied considerably among the different nationalities. Of 1149 American parents having babies enrolled, 15.4 per cent. applied for relief; of 2422 Italian parents having babies enrolled, 12.5 per cent. applied; while the applicants of 1995 Hebrew, 256 Bohemian, 666 Irish, and 164 German babies enrolled represent 9.9, 17.7, 12.8, and 4.2 per cent. respectively. It is also clear that the percentage of enrolment applying for milk relief varied considerably among the different stations dealing with babies of the same nationality. Taking, for instance, Station No. 24, with a total enrolment of 310 Italian babies, 15.1 per cent. applied for free milk, while in Station No. 12, with an enrolment of 358 Italian babies, only 6.9 applied. The same variation is true as regards other nationalities. This would seem to show either a more liberal policy to be enforced by the relief agency working in those particular districts, or a difference in the activity of the station nurse in seeking out babies of families in destitute circumstances and bringing them into the station.

Table 111 shows the monthly increase in enrolment as compared with the new application for free milk.



THE NURSE HAS MANY FRIENDS IN EACH TENEMENT



TABLE 110.—SHOWING BY STATIONS AND NATIONALITIES COMPARISON OF NUMBER	R OF CASES AF	PLYING	FOR MILK RELIE	S.F.
TO ENROLMENT.—PERIOD MAY 1 TO NOVEMBER	1911			

-	LITY	P.	:::::	100.0	:::::	20.0	. : : 100.0	100.0	:	17.4	
	NATIONALITY NOT GIVEN	A.	:::::	:::04	:::::	-::::	::::=	:":::	:	11	
	ZX	E.	ve2 ::	:::79	w: 7: w	n : :00	2 :	~ ≈ + : :	:	63	
	TIES	P.	.: 8.3 15.7	1.5 9.6 18.7	16.6 7.5 6.4 6.4	3.5	6.0	42.8 10.0 18.1 7.2 11.1	:	4.2	
	OTHER	A.	::000	:0 :00	22421	დ := ::	-::::	11253	:	55	
	NATI	E.	. 42 19 17	16 131 182 52 52 16	87 12 53 31 35	55 33 28 21 16	101 27 27 15	50 11 151 9	20	316	
-	z	ъ.	:::::	83.3	25.0	: : : : : :	50.0	55.5	20.0	12.8 1	elief.
	GERMAN	A.	:::::	:::ייי	-::-:	:= : : :	:∾::म	:2 :	-	21	for 1
	0	E.	:: 14	12631	4 :1,4	22 118 19	.980	96168	r.	164	ying
17	H	P.	:::::	100.0	50.0 15.0 31.2 18.4	25.0 10.5 56.2	 100.0 50.0 13.4	22.3 13.6 38.8 6.0 31.5	9.4	17.7	enrolment applying for relief
	IRISH	A.	:::::	. 1 12 ::	1 :027	200 ::	:: " " " "	118 17 2 9	11	118	men
		E.	:::::	: :4	40 16 38 38	12 16 16 17	18 3 3 5 2 5 2	67 132 18 33 33	116	900 118	nrol
200	NNS	P.	:::::	7.5	:::::	3.2	:::::	:::::	:	7.0	o jo
2	BOHEMIANS	A.	:::::	::::	:::::	<b>-</b> ::::	:::::	:::::	:	18	Per cent. of
	m	Ei .	:::::		:::::	31	:::::	:::::	:	256	= Per
1	S	P.	8.0 8.0 7.0 12.2	5.2 31.5 32.8 40.0	37.7	14.2 11.1 7.5	16.3 44.4 4.3 	20.0	:	6.6	F=
3	HEBREWS	A.	22 25 25 25 25 25 25 25 25 25 25 25 25 2	119 22 2	10 3	5 :: 2 ::	0411::	7::::	:	199	lief.
	HB	Э	202 149 371 180	210 60 67 31 5	24 88 13 13	35 9 158 6 126	55 9 254 1 3	. 125	:	1995	or re
	SZ	P.	18.1 20.7 73.6	100.0 .: 13.3 50.0	10.3 6.9 9.3 6.8	5.2	27.2 27.2 15.1 10.2	36.3 16.1 13.1 14.3	33.3	12.5	=Applicants for relief.
	ITALIAN	Ą.	58 222 14 	1 : : 15	31 25 1 14 6	:: 25:	£ : 47 12	. 1058	2	304	oplic
	7	Э.	322 106 19	8 112 4	301 358 39 150 88	3 245 245	310	22 31 76 9	9	15.4 2422 304	A-A
PININGENIEN	SN	P.	:::::	.: 16.6 20.0	15.7 7.7 12.8 28.9 8.9	22.2 11.2 4.2 7.6	5.2 7.7 100.0	59.3 14.2 34.3 11.5	15.5	15.4	100
2	AMERICANS	A.	:::::	:::%	3 20 7	3: 62	:4=01	35	12	177	Ime
	AM	¥	: 17.60	12 30 35	35835	45 107 140 7 39	3 13 13 9	59 63 32 32 147	77	1149	= Enrolment.
	APPLYING FOR RELIEF	Per Cent.	17.6 13.8 12.4 10.4 8.9	8.2 10.2 8.4 15.0 11.7	10.9 7.8 7.9 15.2 8.4	8.2 8.2 8.0 6.6 1.5	7.5 9.1 4.8 15.7 14.0	35.8 15.5 21.4 7.5	11.6	11.2	# E=
	APPI	No.	58 29 36 43 24	222 222 36 36	48 31 16 42 21	23 16 19 33 33	13 14 15 28 28	45 30 15 25	26	903	
	TOTAL	ENROLMENT	329 210 289 412 268	229 215 261 280 306	440 393 202 276 248	205 193 411 285 195	172 153 307 330 199	290 290 140 199 197	224	8031	
	STATION	NUMBER	-1264s	0 1 8 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 1 0 0 1	112111111111111111111111111111111111111	110 113 119 20	21 22 23 24 25	26 27 28 30	31	TOTAL	

TABLE 111

Монтн	ADDED TO ENROLMENT	Number of Appli- cants for Relief	PER CENT.
May (enrolled May 30, 1911)	894	66	7.3
JuneJuly	1263 1892	124 230	9.8 12.1
August	1949 1242 791	268 126 89	13.7 10.1 11.2
	• • • • • • • • • • • • • • • • • • • •		11.0
3	8031	903	11.2

From the above it will be seen that out of an enrolment of 894 babies in May, 66, or 7.3 per cent., applied for free milk, and out of 1263 new babies added to the enrolment during the month of June, 124, or 9.8 per cent., applied for free milk. The percentage of the new babies added to the enrolment applying for free milk reached its highest point in August, when 1949 babies were added to the enrolment, and 268, or 13.7 per cent., applied for free milk.

The following tabulation gives the number of instances in which whole or partial relief was extended and the authorizing agency. In the partial relief cases the agency paid the difference between the price of the store milk and the price of the bottled milk supplied by the station:

TABLE 112.—CASES GIVEN WHOLE AND PARTIAL MILK RELIEF

Relief Agency	CASES GIVEN RELIEF		WHOLE RELIEF		PARTIAL RELIEF	
	Number	Per Cent. of Total	Number	Per Cent.	Number	Per Cent.
A. I. C. P. C. O. S. U. H. C. Hudson Guild Union Settlement N. Y. Foundling Hospital Other agencies Individuals (through Mr. Thos. S. McLane)	335 157 119 58 6 19 12 243*	35.4 16.5 12.5 6.1 0.6 2.0 1.2 25.7	288 153 117 40 6 1 12 209	85.9 97.4 98.3 68.9 100.0 5.2 100.0 86.0	47 4 2 18  18  34	14.0 2.5 1.6 31.0 94.7  13.9

The above statement shows that out of 949 cases in which relief was extended, 826, or 87 per cent., were given entire relief, and 123, or 12.9 per cent., were given partial relief. Of the 949 cases furnished milk relief 335, or 35.4 per cent., were aided by the Association for Improving the Condition of the Poor; 157, or 16.5 per cent., by the Charity Organization Society; 119, or 12.5 per cent., by the United Hebrew Charities, and the remainder received relief through other sources.

<sup>\*</sup> Includes 225 cases given temporary relief until they were taken over by relief agencies.

# EXTENSION OF RELIEF THROUGH MILK STATIONS 139

The following tabulation shows the number of cases in which relief was extended on the initiative of the workers of the different relief agencies:

TABLE 113.—ORGANIZATIONS EXTENDING MILK RELIEF ON THEIR OWN INITIATIVE

Relief Agency	TOTAL NUMBER OF CASES GIVEN	GIVEN RELIEF ON THE INITIATIVE OF THE AGENCY		
	Relief	Number	Per Cent.	
A. I. C. P	335	103	30.7	
C. O. S	157	26	16.5	
J. H. C	119	20	16.8	
Hudson Guild	58	58	100.0	
N. Y. Foundling Hospital	19	19	100.0	
Jnion Settlement	0	6	100.0	
Other agencies	12	12	100.0	
ndividuals	243	12	4.9	
Total	949	256	26.9	

In 30.7 per cent. of the cases granted relief by the Association for Improving the Condition of the Poor the initiative was taken by that organization, while cases given relief by the Charity Organization Society and the United Hebrew Charities, 16.5 and 16.8 per cent. respectively were given relief on the initiative of these societies. Of a total of 949 cases receiving milk relief during the demonstration, 256, or 26.9 per cent., were given relief upon the initiative of the relief agency.

The following tabulation shows, for the three main relief agencies, the number of cases referred by the Committee for milk relief and the action taken:

TABLE 114.—CASES REFERRED FOR MILK RELIEF AND RESULT

Society	A. I. C. P.	C. O. S.	U. H. C.	TOTAL
Number of cases referred	275	209	164	648
Number given milk relief	232	131	99	462
Per cent	84.4	62.6	60.4	71.2
Number aided in other ways	7	8	5	20
Per cent	2.5	3.8	3.0	3.1
Total number aided	239	139	104	482
Per cent	86.9	66.4	63.4	74.3
Number refused relief	28	64	54	146
Per cent	10.1	30.6	32.9	22.6
Number of cases withdrawn	8	6	6	20
Per cent	2.9	2.8	3.6	3.0

From the foregoing it appears that out of 275 cases referred to the Association for Improving the Condition of the Poor, 232, or 84.4 per cent., were granted milk relief, and 7, or 2.5 per cent., were aided in other ways, giving favorable action in 86.9 per cent. of the total cases referred; of 209 cases referred to the

Charity Organization Society, 131, or 66.4 per cent., were given milk relief, and 8, or 3.8 per cent., were aided in other ways, while out of 164 cases referred to the United Hebrew Charities 99, or 63.4 per cent., were given milk relief, and 5, or 3 per cent., were aided in other ways, showing that out of a total of 648 cases referred to the three relief agencies through the central office of the Committee, favorable action was obtained in 482, or 74.3 per cent., of the total cases referred.

Below is given a tabulation showing the number of cases receiving milk relief through the different agencies and the amount of money expended during the demonstration on account of relief given in the form of milk and nursing supplies, such as bottles, nipples, barley, etc.


NAME OF SOCIETY, ETC.	AMOUNT EXPENDED	Amount Expended Number of Cases Given Relief	
A. I. C. P.	\$1645.55	335	\$4.91
C. O. S	1004.99 525.17	157 119	6.40 4.41
Other agencies	372.84 487.44	95 243	3.92 2.01
Individuals	\$4035.99	949	\$4.25

TABLE 115.—COST OF RELIEF


The total amount of relief extended by the Association for Improving the Condition of the Poor amounted to \$1645.55; \$1004.99 was extended by the Charity Organization Society; \$525.17 by the United Hebrew Charities, and \$860.28 by all other agencies. The average amount of money expended per case by all the agencies was \$4.25. During the period of the demonstration the total value of milk and nursing supplies dispensed amounted to \$30,860.98, of which, as shown by the foregoing statement, \$4035.99, or 13.0 per cent., was given as relief. This does not include a loss sustained by the Committee of one-half cent upon each quart of milk dispensed at its stations.

Relief in other forms was extended to needy families through the efforts of the station nurses independently of those formally reported for milk relief. No attempt was made to maintain complete records where the station nurse acted on her own initiative in helping poor mothers to solve their economic problems, but numerous cases came to the attention of the central office which evidenced the earnestness and the good sense of the nurse. The efforts to save some of the babies' lives and to straighten out home conditions frequently required a great deal of personal sacrifice on the part of the nurses; an example of a case by no means uncommon was reported by the nurse at Station No. 29 as follows; in this case what was done by the station nurse is best described in her own words:

"On the evening of May 10th, just as I was closing the station at 5 p. m., a poor, frail little girl, about six years old, came to the door of the station and said: 'My mother wants a bottle of milk.' I immediately went to the home, and there



A Lesson in Bathing



found five children, of which this little girl was the eldest. There were twin babies, about three months old, in the family. The poverty and squalor are almost indescribable. The mother was a tall, sickly looking woman, sitting on the floor with the twins in her arms; there was absolutely no food in the home. I returned to the station, sent them some milk, and reported the case to the central office, requesting that the relief be extended for a time. The father, I afterward learned, was a longshoreman out of work and addicted to drink. The babies were under-nourished and suffering from marasmus—one of them was developing pneumonia. Through the generosity of a member of the committee who visited the station a nurse, medicine, and doctor were obtained, but the baby having pneumonia died. I obtained sufficient money in the neighborhood and from my friends to bury the little mite. Conditions in the home did not improve, and the family was soon dispossessed. The mother and children came to the milk station, where they stayed until night, when the father returned. He obtained a room on Clarkson Street and removed the furniture there.

"I continued to visit them in their new home and did everything possible for the other baby, instructing the mother how to care for it; but she, like her husband, had become discouraged, and I not infrequently found her intoxicated and the children sitting around her on the floor, stupid from hunger. The baby was very frail, weighing only about four pounds, and it looked as if she too would die; but by visiting the home every day, making the milk formula, and not trusting the mother, the baby started a slow but steady gain in weight. During all this time the other three children came to the station twice daily, where I gave them a glass of milk and shared with them the lunches intended for the matron

and myself.

"The family had been reported to various relief agencies, but with unsatisfactory results, as they claimed the father refused to comply with the requirements under which aid would be extended. Finally, I brought the case to the attention of the parish priest, and through him the children were placed in an


institution.

"Of all the cases with which I have had to deal this summer, this family required more thought and time to accomplish results than any other, but I hold that not only the life of the remaining baby, but the lives of the three other children were saved through the efforts of the milk station."

The citing of another case will show how the station nurse helped the mother solve a problem which she was required to face. This mother had a baby eighteen months old, suffering from gastro-intestinal trouble, and a nursing baby ten days old. Her husband had died ten days before the baby was born, leaving the mother, a girl of seven, and two little babies without any support. The case was referred to the Charity Organization Society, which advised putting the children in an institution and obtaining employment for the mother; but the station nurse felt that every endeavor should first be made to prevent the breaking up of the home and the separating of the mother from her children, especially from the nursing baby. After the decision of the relief agency the nurse called at a maternity hospital, where applications for wet-nurses are often received, and was given the address of a lady who wanted a wet-nurse, with the result that the mother was engaged and secured a good home in the country for herself and her nursing baby. The problem of providing for the remaining children was solved by the station nurse's hunting up relatives of the family who were willing to take the children until the mother could otherwise make provision for them.

Case after case could be cited, showing the many ways in which the statton nurse helped distressed mothers with whom she came in contact. Too much cannot be said of the enthusiasm and keen interest manifested by the nurses during the demonstration, and it was their untiring efforts that made the milk stations real live social service organizations for the districts in which they were located. Mothers soon learned to appreciate the character of the service and the spirit manifested by the station nurses and their assistants. That they came to the station for advice and help is an indication of the possibilities of such centers and the decided influence which they exert.

The milk stations cooperated with various other agencies furnishing help in forms other than milk relief. Those milk stations located in the vicinity of distributing points for free ice through the Free Ice Fund of the "New York Herald" were provided with tickets in the form shown below:



Ехнівіт 21

The numbers in the squares indicate the days of the week the ticket has been used by the holder. These tickets were given out only to mothers unable to purchase a sufficient amount of ice to insure the keeping of the daily supply of milk at a low temperature. The name, address, agency issuing the ticket, and the date of issuance were reported on the perforated stub. The stub was detached by the agent dispensing the ice when the ticket was first presented at the distributing place.

It has formerly been the custom of the "New York Herald" to distribute free ice during the months of July and August. During last year the distribution began early in June, and the distributing points were selected after a careful study of the locations of milk stations. An early morning scene, showing mothers waiting at one of the depots for ice, is shown in this report.

#### EXTENSION OF RELIEF THROUGH MILK STATIONS 143

In addition to the coöperation of the "New York Herald" in furnishing ice the Knickerbocker Ice Company issued coupon books containing 31 coupons, each good for 10 pounds of ice if presented to any of their 14 ice depots; the form of this coupon is shown below:

ш	Book No. 1144 New York, 1911
SFERABLE	The Bearer,
TRANSFE	is entitled to receive Ten Pounds of Ice for each of the attached coupons when this book is presented to the Weighmaster of any of the Ice Depots named on back of book, or to the Driver of any retail ice wagon of the Knickerbocker Ice Company when countersigned by an authorized agent of the
NOT	NEW YORK MILK COMMITTEE.
-	ByAgent
	Only one coupon will be honored each week day and two Saturday.

Ехнівіт 22

Book No. 1144 Coupon No. 1

This Coupon when not detached is good for

### TEN POUNDS OF ICE

according to terms stated on front cover of this book.

**NEW YORK MILK COMMITTEE.** 

#### Ехнівіт 23

The ice-box problem has already been solved, as very few homes visited by the station nurse during the demonstration were found not to have facilities for keeping the ice. The education of mothers as to the importance of keeping the babies' milk cold, and the established custom of the agencies, already mentioned, of distributing free ice, has caused the poor mothers to solve the ice-box problem satisfactorily. To the ice-distributing agencies much credit is due for their efforts to aid the mothers in keeping the babies' milk supply sweet and wholesome.

#### RELIEF IN THE FORM OF HOSPITAL CARE

The relief extended in the form of hospital and dispensary care of the sick babies during the period of this demonstration contributed in no small way to its success. However, the difficulties in placing sick babies in hospitals seriously lessen the effectiveness of the hospitals. During the summer a careful canvass of the hospitals in New York city was made, and the result shows that, for the care of infants under two years suffering from general diseases, there are approximately 800 beds which are not restricted by requiring a legal commitment or a permit from the Department of Charities. These beds are not all available for the use of infants under two years because of the requirements for admissionnone will receive cases complicated by mild contagion; some do not receive babies under one year of age; some require mothers to accompany babies; others have no accommodation for nursing mothers, while some receive no cases during the summer and others receive cases only during the summer; most of these beds are for what are known as pay or part-pay patients. Only 21 per cent. are reserved exclusively for free patients. Again, these 800 beds are not available at all times, as often a hospital is in quarantine with some contagious disease.


The existing facilities, though inadequate, cannot be properly utilized under the present system, as no one knows the total number of available beds at any given time, and to place a case one must telephone from hospital to hospital until accommodations are found. A central hospital bureau prepared to give prompt information as to available beds at any time would make possible a better utilization of the present hospital facilities, which are generally known to be woefully inadequate.

Because of the difficulty in placing sick babies under proper hospital care comparatively few of those registered at the Committee's stations were sent to hospitals. Junior Sea Breeze, which has special facilities for the treatment of gastro-intestinal diseases, responded promptly and satisfactorily in every case brought to its attention.


#### RELIEF IN THE FORM OF OUTINGS

Through the coöperation of various agencies maintaining seaside and country homes for sick mothers and children the station nurses were liberally provided with tickets to distribute to those in need of this kind of relief. In the case of the St. John's Guild Floating Hospital, tickets were good only for sick children or mothers recommended by either the family or station physician.

In very few instances did the station nurses encourage mothers to take their babies to seashore and country homes if the baby was well and gaining in weight under the station supervision. This is particularly true of babies artificially fed,



GIVING ADVICE AS TO THE DIET OF THE OLDER CHILDREN. THE NURSE COMES IN CONTACT WITH THE WHOLE FAMILY



# EXTENSION OF RELIEF THROUGH MILK STATIONS 145

because the change in milk and feeding formula had been found in a number of instances to prove detrimental, at least for a time.

However, special effort was made by the station nurses to obtain for the mother a rest in her own home by having relief agencies provide a country or seashore outing for the older children. This plan proved far more beneficial to all concerned, and was keenly appreciated by the overworked mother.

TABLE 116.—SHOWING FOR 2156 FAMILIES HAVING BABIES REGISTERED AT MILK STATIONS, RENTAL, INCOME, AND LIVING CONDITIONS

Number of Persons IN FAMILY	TOTAL FAMILIES	TOTAL NUMBER OF PERSONS	TOTAL NUMBER OF ROOMS	TOTAL NUMBER OF DARK ROOMS	AVERAGE YEARLY RENTAL PER FAMILY	AVERAGE YEARLY INCOME PER FAMILY
2 3 4 5 6 7 8	1 541	2 1623	1 1627	408	\$108 145	\$260 641
4	379	1516	1182	355	145	621
6	429 304	2145 1824	1398 1019	372 300	147 151	633 631
7 8	205 143	1435 1144	720 499	250 137	156 164	599
	76	684	270	92	151	626 546
0ver 10	45 33	450 396	177 141	39 50	183 196	678 740
TOTAL	2156	11219	7034	2003	\$150	\$628

Number of Persons in Family	Number of Families Living in								
TOMBER OF TENSORS IN PAREL	1 Room	2 Rooms	3 Rooms	4 Rooms	5 Rooms	6 Rooms	7 Rooms	8 Rooms	
2 3 4 5 6 7 8 9 10 Over 10	1 7 6 8 4  1 	118 70 74 39 15 10 4 2	297 199 195 131 100 71 38 12 6	104 82 111 112 67 45 25 22 20	13 21 35 13 18 13 6 7	2 1 4 4 3 1 3 1	 2 1 2 2 2 		
Total families Total number of persons Total number of rooms Total number of dark rooms Average yearly rental per family* Average yearly income per family*	27 119 27 10 \$82 \$433	332 1479 664 227 \$124 \$516	1049 5250 3147 920 \$138 \$602	588 3381 2352 640 \$171 \$678	130 778 540 160 \$211 \$852	19 120 114 19 \$258 \$918	8 58 56 19 \$229 \$921	3 34 24 8 \$384 \$1716	

<sup>\*</sup> Cents omitted in amounts given.

#### In conclusion:

- 1. The milk stations can be successfully conducted without making the distribution of free milk a prominent part of the work.
- 2. The organized relief agencies should be called upon to handle every case needing free milk. If a family is too poor to buy milk for the baby, the case is not one for free milk alone, but for general relief, and as such should be dealt with by an agency prepared to deal with the situation.
- Whatever relief is extended in the way of free milk should be given in a
  quiet way. The desire for free milk is contagious with milk-station
  mothers.
- 4. Arrangements should be made to have some agency undertake to furnish needy cases with the necessary ice to keep the baby's milk sweet and wholesome. The mothers can and will solve the icebox problems.
- 5. Hospitals with many so-called free or part free beds do not mean available hospital service for sick babies. The standardizing of requirements for admission and the establishment of a Central Hospital Bureau for information will insure a better utilization of exisiting hospital facilities.
- The older children may be sent away for a country or seashore outing, but the baby should be left at home with its mother. Travel and change of milk and feeding formula may be disastrous to the bottle-fed baby.
- 7. The milk station should be made a place where mothers will come with their troubles. The nurse should be prepared to know what to advise; if she shows good judgment, she will win the confidence and friendship of the mother.

#### CHAPTER IX

# Establishment and Operation of Milk and Health Stations

SURVEY TO DETERMINE THE NEEDS OF EACH LOCALITY

A preliminary survey was made in 1910 to determine how many milk stations would be necessary to carry on an intensive demonstration in Manhattan. Information and experience obtained in operating six milk stations during 1909 and census figures served as the basis for this study. The records of these milk stations were analyzed, a map of the district in which the stations were located was made, and in each block was indicated the number of babies registered at the station. The number of babies was then considered in relation to area and density of baby population.

As the six stations were located in different sections of Manhattan it was thought that their experience could be taken as an index of what might be expected if additional stations were opened in similar districts. Accordingly, it was estimated that the average sphere of influence in acres of a station on the lower East Side was 16; on the upper East Side, 24, and on the West Side, 22.

By considering station enrolment in relation to baby population it was estimated that in districts with more than 25 children per acre 25 per cent. might be expected to come under the supervision of milk stations; in districts with 15 to 25 children per acre, 10 per cent.; and in districts with less than 15 children per acre, 5 per cent.—might be expected to come under the supervision of stations. Allowing 100 babies as a maximum enrolment for a milk station, it was estimated that, to provide for the entire number of babies needing milk-station supervision in Manhattan, 60 stations in addition to those already in operation would be required.

In April, 1911, after the Committee had secured sufficient funds to begin a demonstration, another survey was made to determine the number of babies under two years who might be brought under the influence of the milk stations. This study was made by wards or parts of wards, in which the block population was used as the basis for estimating the number of children under two years of age. From this, together with the experience of the stations already in operation, it was estimated that a minimum of 13,137 babies under two years could be brought under the influence of milk stations if these were properly located. Information from organizations already conducting milk stations showed that 4500 of the total estimated number were already under the supervision of established stations, leaving 8637 babies to be provided for. After a thorough canvass of the districts not covered by milk stations it was decided that if 28 stations were established by this Committee in addition to the 5 to be maintained in Manhattan by the Department of Health and 22 by other agencies, it would be pos-

sible, by employing extra nurses, to care for the remaining 8637 babies needing supervision.

With the tentative locations for new stations indicated on a map of Manhattan, a careful inspection of each district was made. The clergy of all denominations, as well as social workers and representatives of hospitals and dispensaries in the locality, were interviewed, and many valuable suggestions were obtained relative to the proposed location of the station in their district.

Once having determined the vicinity in which the station was to be opened, a canvass was begun for suitable quarters; in selecting these, an endeavor was made to have the stations placed on streets which were most used by the mothers of the neighborhood.

#### KIND OF QUARTERS SELECTED

In all but five instances stores were selected for quarters for the milk station, and an effort was made in each case to obtain at least 600 square feet of floor space, although in certain districts this was not possible, as nearly all the stores were the so-called single stores, having a width of 10 feet and extending back 40 to 45 feet. Wherever possible stores were obtained having partitions so arranged as to give one large room in front with two smaller rooms in the rear. (See Exhibit 24.)

Free quarters were provided for five of the stations by different organizations—two of these were in settlement houses; one in an industrial school; one in a dispensary, and one in the room adjoining the vestibule of a chapel. It became necessary, however, to remove the station from the chapel after occupying the quarters for six weeks, owing to the refusal of mothers to bring their babies to a station located in a church not of their religion. No difficulty was encountered in obtaining a large enrolment at stations located in settlements; in fact, with one exception, the settlement stations had the largest enrolment throughout the summer of any of the Committee's stations.

#### EQUIPMENT AND COST

A careful list of the equipment necessary for the station was compiled, taking into consideration the economy of floor space and the amount of money available. The following is a complete list of the furniture and equipment purchased for each station, as well as the cost of same:

#### LIST OF EQUIPMENT OF STATIONS

Folding chairs (24) wooden			\$15.50
Stool (1), white enamel, wooden			.87
Cabinet (1), white enamel, wooden			
Mirror (1), white, size 12½ x 16 inches			
Blackboard (1), size 24 x 36 inches			.32
Chairs (2), white enamel			
Table (1), size 24 x 30 inches, white enamel, wooden			
Table (1), size 20 x 48 inches, white enamel, wooden, with draw			
x 19 inches			6.50
Table (1), size 24 x 60 inches, white enamel, wooden, 5 drawers-	-4 5	side.	
1 long			
Desk (1), size 24 x 40 inches, white enamel, wooden, 5 drawers-			
1 long		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	10 25
A IUME			10.20

Wash-stand (1)	3.50
Refrigerator (1), size 27 inches x 5 feet, oak	34.00
Linoleum (to measure), 65 cents per vd., made and laid	
Pan for refrigerator	.55
Pan for refrigerator	
inches	5.00
Howe scales with special pan (1)	9.00
Pad for scales	.22
Double boiler (1), 1 quart, enamel	.57
Strainer (1), large, to fit quart measure, enamel	.25
Quart measure (1), enamel	.30
Spoon (1), enamel, white, size 14 inches	.10
Jars (4), glass, I quart, with tops	.88
Funnel (1), glass	.09
Bath-tub (1), baby's	1.05
Graduate glass (1), 16 ounce	.20
Kettle (1), 1 quart, enamel.	.80
Tea-kettle (1), 4 quarts, enamel.	.78
Pitcher (1), gallon, enamel	.65
Saucepan (1), gallon, white enamel	.80
Dishpan (1), size 16 x 5 inches, enamel  Mop (1) and handle	.60
Mop (1) and handle	.30
Boiler (1), No. 6 Kramer	1.12
Irons (2), 6 pounds, each	.44
Ironing-board with cover, size 4½ feet	.80
Scrub-pail (1) and wringer (1)	1.00
Washboard (1)	.32
Bottle-cleaners (2) for small nursing bottles	.50
Bottle-cleaner (1), large, with wooden handle	.25
Nail-brush (1)	.04
Scrub-brush (1)	.11
Soap-dish (1), enamel	.15
Towel-rack (1), nickel	.42
Clothes-line (1)	.45
Broom (1)	.30
Ice-pick (1) and hammer (1)	.25
Dust-pan (1) and brush (1)	.31
Slop-pail (1), enamel	.63
Hand-basin (1), enamel	.24
Bowl and pitcher for wash-stand, enamel	.97
Window-screening	.25
White dotted muslin (20 yards)	1.90
Towels (12), hand, 1 yard each, hemmed	1.38
Towels (12), dish, 1 yard each, hemmed	1.32
Floor-cloths (6), ½ yard each, hemmed	.66
Scrap-basket (1), wire	.35
Clock (1), New Haven alarm	.60
Inkstand (1), with 2 bottles and pen-holder	.75
Cash-box (1), tin, size 7 x 9 x 2 inches, 6 compartments	.55
Thermometer for refrigerator	.48
Dairy thermometer	.17
Tongue depressors.	.60
Clinical thermometer (two-minute)	.50
Breast-pump (1)	.30
Proceeds (1)	.45
Pyoscope (1)	2.80
Card-drawer files (daily record cards), at 80 cents	1.80
Card-drawer mes (daily record cards), at oo cents	1.00

The average cost of equipment was \$184.72 per station. All the furniture was white enamel, with the exception of the folding chairs, and was made to order for the Committee. With the exception of the refrigerators, the equipment selected proved well adapted to the needs of the milk station. The refrigerators, although of well-known make, did not prove satisfactory, as it was impossible to keep the temperature of the milk below 55° F. unless the bottles were placed

in direct contact with the ice. It is doubtful if any refrigerator will give as satisfactory results in keeping milk at a low temperature as an ice-chest in which the bottles can be buried in cracked ice.

#### ARRANGEMENT OF INTERIOR OF STATION

Wherever possible, as has been said, stores with three rooms were obtained for milk stations—the front room, being the largest, was used for a class-room and for the dispensing of milk. In the exhibit showing floor plan of a three-room station the milk-dispensing table is in the fore part of the front room. In some places a rail and gate, together with the table, separated the remaining part of the room used for class purposes from the milk-dispensing portion. A three-paneled screen is utilized to separate the class-room from the view of the street. The refrigerator, with a capacity of 200 quart bottles of milk, is placed alongside of the cabinets. These cabinets contained nursing bottles of various sizes, barley, milk-sugar, nipples, borax, and other supplies which were sold to the mothers at cost. A portion of the front room utilized for class work or as a waiting-room for mothers on consultation day with the physician is provided with 24 to 36 folding campchairs and a table for the use of the nurse or physician conducting class work.

The adjoining room is known as the "demonstrating room"—this is the smallest of the three; here mothers were instructed in the modification of milk, bathing and dressing of babies, etc. It is provided with a large demonstrating table, a sink with running water, wall shelves on which were kept the utensils used in demonstrating the modification of milk, a cabinet for supplies, and a cloak closet.

The third or rear room is nearly twice as large as the demonstrating room, and is known as the "consultation room," in that it is used almost exclusively for individual consultation work with mothers. The station physician and nurse are each provided with a desk. At the side of the room nearest the nurse's desk is a table holding the scales used in weighing babies, and on the opposite side is the cabinet containing medical supplies used in the station work.

The arrangement, as per the plan outlined, makes it possible for a station with apparently small quarters to handle quite conveniently a great many cases. The use of that part of the front room nearest the street door for receiving mothers or older children coming for milk, without interfering with the work carried on in the other part of the station, is particularly advantageous. The class-room makes it possible for the general lessons in hygiene and care of the baby by the station nurse and the special talks by the station physician to be given without interruption from persons coming in for the sole purpose of obtaining the day's supply of milk. By having the consultation room in the rear, a good light is assured for the examination of babies by the physician. The room affords a certain amount of privacy, and the mother feels that the station is giving her baby individual attention and that she is being granted a personal consultation, rather than being instructed as one of a group. The character and amount of individual attention given to each baby and mother by the station physician and nurse determine largely the influence which the station is to have upon that mother.





EXHIBIT 24
FLOOR PLAN OF MILK STATION
STORES ARE USED FOR MILK STATIONS



All records with the exception of the ledger for recording the milk dispensed are kept in the consultation room. On the walls of the front room of the station are placards giving advice or warnings to mothers. A blackboard in the milk-dispensing portion of the station affords a means of notifying mothers of changes in clinic hours and other important notices, such as announcements or reminders of clinics and classes to be held, excursions to parks or seashore, "little mothers'" meetings, etc.

#### PAINTING AND DECORATING THE STATION

A blue-and-white color scheme for both the city milk stations and the Committee milk stations was adopted. The exterior of the station was a light blue

# THE BLUE FRONTS

# **BABIES' PURE MILK STATIONS**

Every Summer thousands of Babies die We can save your Baby Ask us how

Committee for Reduction of Infant Mortality

New York Milk Committee

#### Ехнівіт 25

PLACARDS USED TO ADVERTISE THE STATIONS (Printed in Five Languages; Size, 22 x 28 inches)

with white trimmings around the windows. White calcimine was used for the ceiling, and a white-enamel paint for the side walls—in some cases white-enamel paint for the side walls was used up to a height of 5 feet, and the remaining portion of the wall calcimined. In some of the city stations blue was used in marking borders in the interior. Having the exterior of the station a striking blue, this being an unusual color for fronts of buildings, served at once to attract the attention of the neighborhood, and it was soon known as the "House with the

Blue Front," and greatly aided the nurses in directing the mothers to the milk station, by simply giving the street upon which it was located and telling them to look for the "House with the Blue Front." In most of the stations there were signs, either at right angles to the front of the station or painted upon the large plate-glass window. The interior of the station being in white, its white-enamel furnishings and linoleum floor-covering gave the visitor the impression of clean, sanitary surroundings.

#### BUILDING UP THE STATION ENROLMENT

The location of the station was determined, quarters secured, and the premises renovated and painted, furniture and equipment installed, and the organization was completed by engaging a nurse, matron, and a physician for carrying on the work of the station.

With this personnel, work was begun immediately to build up a station enrolment. The nurse visited the dispensaries, hospitals, settlements, clergy, social workers, private practitioners, and druggists of the district to make known to them the aims and purpose of the station, and to obtain their coöperation, urging them to refer babies to the station for feeding supervision. The following postal-card form was distributed to facilitate referring cases:

	Date
We have	this day temporarily referred for feeding to Milk Station
No	Street
	of the NEW YORK MILK COMMITTEE
Name	
Address	
Remark :	
	Physician

#### Ехнівіт 26

Wall-cards and convenient pocket-cards, giving the locations of all the milk stations of the city and the organization conducting the work, were left with those who were willing to coöperate. Large printed signs were placed in the window of the station, to announce the work that was to be carried on, and to notify mothers to bring their babies for examination by the station physician. These were also placed in store windows, and not infrequently were to be seen in corner saloons. Hand leaflets, showing contrasts between sick and well babies, and urging mothers to bring their babies to the stations, were distributed by druggists and shopkeepers, with good results.

After making this announcement of the opening of the stations, the nurse began canvassing systematically the tenements on each street of her district. From the janitors she obtained information as to the families having babies under two years of age, and these homes were visited and the mothers urged to come to the stations with their babies. They were told the purpose of the station and the kind and price of the milk dispensed. As the babies became registered the mothers were given circulars to distribute among their neighbors and friends; they were asked to bring with them other mothers whose babies were sickly, to get them well; and those whose babies were well, to learn how to keep them well.

Each home having a baby under two years of age needing supervision was visited, and the following card form was filled out:

	Born.	4/1/11	ELIGIBLE BABY Flight 2 frt. beld ;		
	Name	K	Address VI Cornel	ia Si	£.
EE			PARTLY BREAST FED ARTIFICIAL	LLY FE	ED V
TTIM			LOOSE L'CONDENSED Hovery	PAT. FOOD	TABLE FOOD
- OM	DATE	Condition	ACTION TAKEN—REMARKS		N.VISIT
K CC	8/1	Por	Terged to tetens Clinio - Instructe	7	8/19
MIL.	8/19	n	no improvement - will attend clinics		1/20
Form	8/43	4	not as well - Instructed		8/27
W	The	sec	advised to see dotor - Will register &	aly a	8/31
NEW	0/3/	4	Called to bring mother & Clinic		

Ехнівіт 27

This is a salmon-colored card, 3 x 5 inches, punched so as to carry in a loose-leaf cover, and is known as the "Eligible Baby" record. The term "eligible babies" refers to those babies who live within the district which the station is supposed to serve, and who, in the judgment of the nurse on visiting the homes, need the care and supervision of the milk station. In other words, every home visited in which conditions were unfavorable, due either to poverty, ignorance, or neglect, the baby was considered a candidate for registration at the milk station.

The above record shows that this particular baby lives on Cornelia Street, second flight, back hall, and apartment to the right. The baby was born on April 1, 1911, and artificially fed on "loose" (dipped) milk purchased at a grocery store. The date of the first visit to the home was August 15th; the condition of the baby indicated was "poor." The nurse urged the mother to attend the clinic, and instructed her in the care of bottles, nipples, and milk. The nurse revisited the home on August 19th, 23d, 27th, and 31st; the last was for the pur-

pose of bringing the mother to the station to have the baby examined by the station physician.

#### A REGISTERED BABY

Once a mother brings a baby to the station and the baby is examined by the station physician, and the mother promises to follow the instructions given to her by the physician and nurse and attend the consultations as directed, the baby becomes a "Registered Baby," and the following card record is made:

		,	BAE	3Y	Flight 2 frt. bok. ry lft. r	r. house
	0	L	A	ddŕess	- 0 . /	St.
		ACTION	TION	DATE	HOME VISITS ACTION TAKEN	NEXT VISIT
8/31	I11	Castor oil 3	hrs	8/31	Instructed mother	r
9/7	Imbrove	ed .				9/1
9/14				91		9/2
9/21	Very 111	Saline 80	2	01		
	1		,	. 1		
on	ce do	ily			and B.W. diet	
				9/22	Saline.	
	Name ATTEN DATE 8/31 9/7 9/14 9/21 Muy	Relief Case V.  Name S  ATTENDANCE AT DATE CONDITION  8/31 III  9/7 Improve  9/14 "  9/21 Very III  Murphy in	Relief Case Name S L  Attendance at Medical Consultate Condition Castor oil 3. B. W. diet 36  9/7 Improved  9/14 "	Relief Case	Name S L Address  Attendance at Medical Consultation  Date Condition Action 11 Date  8/31 III B.W. diet 36 hrs 8/31  9/7 Improved  9/14 " 9/1  9/21 Very III Saline 8 03 9/2  Murphy installation 9/19  once daily	RELIEF CASE

ATTEN	NDANCE AT	MEDICAL CONSULTATION		Home Vi	SITS	NEXT	
DATE	CONDITION	Action	DATE	Action	TAKEN	Visit	
9/18	Imbroved	Saline discontini	9/23 e0	Saline ir	rigation	9/24	
10/5	acod	None	9/24		,	9/25	
10/12		4	9/25			9/26	
19/19	u	Changed Formula	9/26		91	9/27	
10/26	ч		9/27				
			19/19	Taught	milk.		
				modifi	cation		
					Lange Ope		

Ехнівіт 28

Pert. Excellent

Good

EXHIBIT 29 A SEPARATE MEDICAL RECORD FOR EACH BABY



This card contains the important information shown on the "Eligible" card of the baby, and the current information as to the clinic consultation, the action taken by the station physician, the home visits, and notes of the station nurse. It forms the working memorandum which can be conveniently used by the nurse in exercising supervision and control over the case. This card form also has punched holes for carrying in the loose-leaf binder whenever the nurse makes her home visits upon the case.

Both the "Eligible Baby" and the "Registered Baby" card forms are filed at the station in a tickler file, under the date on which the case is to receive further attention. When the "Eligible Baby" becomes a "Registered Baby," the "Eligible" card record is then filed alphabetically, in a terminated record file, and the "Registered" baby card becomes the active record. When the registered baby is attending clinic regularly, with no unusual conditions prevailing which would require home visiting and special work on the part of the nurse, this record is filed by street and numerically by the house number. If home visiting, etc., is necessary, the card is filed in a tickler file according to the date on which further action regarding the case is to be taken. Upon termination of a case by removal, transfer to hospital, or other agency, etc., the record is then filed alphabetically in a file for terminated cases. The registration card gives information as to the name of the baby, the address, date of birth, whether the case is one in which relief is extended, and the case number. The form also provides for recording the attendance of the baby at the medical consultations, its conditions, and the action taken as a result of the medical examination. It further provides for recording the home visits made by the nurse by giving the date of the visit, the action taken, and the date of the next visit, if such is necessary. The date indicated for the next visit determines the date under which the card is to be filed in the tickler file by the nurse upon her return to the station.

#### INDIVIDUAL WEIGHT CHART AND MEDICAL RECORD OF BABY

In addition to the foregoing card record maintained for each registered baby, a weight chart and medical record is kept in each case. This record is a  $9\frac{3}{8} \times 12\frac{1}{4}$  inch card, printed on both sides. One side is used for recording information relating to the baby if under one year of age; the other side, if over one year of age. This record is made up at the time the baby is first examined by the station physician, and the mother agreed to follow the instructions of the physician and nurse. (See Exhibit 29.)

This record provides for recording the name and address of the case, the date of birth, the number of the case, the station at which the baby is registered, sex, whether full-term or premature birth, reported weight at birth, length of time breast-fed, length of time mixed fed, the reason for artificial feeding, the kind of food given, nationality of mother, and the names of the station physician and nurse. All this information relates to conditions existing at the time the baby is registered.

The major part of the form provides for recording information relating to the case, as noted by the station physician, and the kind of action taken. In the

upper part of this space provision is made for recording the month and the date the case comes under the observation of the station physician, the kind of formula for home modification if artificially fed, instructions regarding the number and intervals of feeding if breast-fed, and the weight of the baby in pounds and ounces for the age in weeks.

The lower part of the chart records the general condition upon medical examination at entry, the physician's notes regarding the case, the average weight curve for normal babies, and the weight curve of the particular baby under observation. The reverse side of the card records similar information for the baby in its second year, the main difference being in the normal weight curve. These chart records are filed alphabetically, according to the clinic or the consultation day on which the mother is instructed to bring her baby to the station.

Whenever a case is transferred to another station, the chart record, together with the other records pertaining to the case, is forwarded to the station to which the baby is being transferred. If the case is terminated for whatever cause, the chart is filed alphabetically under terminated cases, and the reason for termination is reported to the central office.

#### CASE HISTORY CARD

In addition to the registration card and weight chart and medical record, a case history card is made out for each baby registered. This gives information concerning conditions which prevailed previous to admittance to the station, and takes the character of a social record of the case. This is a 5 x 8 inch card, printed on one side; the following shows the form of this record:

NEW YORK MILK COMMITTEE STATION NUMBER \$5 CASE NUMBER \$41  WARD NUMBER NAME  SABY'S NAME  NEW YORK MILK COMMITTEE STATION NUMBER \$5 CASE NUMBER \$41  CASE HISTORY CARD  (CASE HISTORY CARD  (CASE HISTORY CARD  (CASE HISTORY CARD  (CASE NUMBER STATION NUMBER \$5 CONNELIS \$5.											
	Tenement	1	BASY	M	1	Cowis' milk	1	MOTHER'S NAME Mary			
	Floor	2	( Number	3		Raw		Nationality Italian			
	Pront		Birth Normal	1		Bolled	1	Occupation Housework + Embroidery			
	Rear	1	(Abnormal			Pasteurized		Born in U. S			
Residence	No. of rooms .	4	(Hospital		Milk	Certified		Color White V Can read V			
	No. of light	2	Home	V	used	Condensed		Illack Cannot read			
	Rental	17.00	Physician			Loose	1	Number of days before Number tel days after			
	Home		Attendant Midwife	1		Bottled		not work O sumed work 21			
	Boarding		at Birth Nurse			Modified (Home) .		Kind of work Emberdery Kind of work House			
Average we	eekly income .	12. 10	Other			Already modified .	1000	General health: Good Fair Poor			
	Father	1	Mother	1		/Patent food		FATRER'S NAME Frank			
	Mother	1	Cared for Other adult			Table food		Nationality Italian			
Family	Adults	21	. by Older child	1		Rave fruit		Occupation Pastry-work			
consists of	Children	3	Neighbor		Other Foods	Coffee		Born in U. S Mo Intemperate			
	Bourder		Present (Hospital		Beverag	Tea		Color White V Can read V			
	Roomers	1	Medical Dispensery .		1000000	Soda water		Elack Cannot read			
	(Clean	V	Supervision (Private	V		Beer		Children living 3 Ages 7473. 5473 4 70475.			
Cleanliness			Good			Wine		Remarks:			
of Premises	Very dirty .		Health Fair	-		Hours	1	Baby never breast Led.			
	(Good		Condition (Foor	V	Outing	Regular		Thin and ill orderished			
Ventilation	Fair	V	(Regular			(Irregular	V	Feeding irregular as			
	Poor		Condition Irregular		Baby car	riage used		to or antitu and interval			
Buth tub .			Sleeps alone .	1	Faby car	riage not used	V				
Eath tub us	red	1	Breast fed Mother		Name as deale	d address of milk					
("	Tash	1	Wet-eurse		13	esser		~ ~ 0 0 1. 11			
Tollet - N	lo flash		Partially breast fed	,	310	ornelia		Nurse 12. 6. Schofield			
(P	rivy		Artificially fed	V				Date compiled 9 - 11 - 11			

The information relates largely to a history of the baby and its home conditions. The record is compiled as soon as possible after the baby becomes registered, generally on the first visit of the nurse to the house after registration. The arrangement of items provides a convenient way for recording data by checking the items upon which information is desired. The purpose of this history card was to ascertain the character of the homes which the milk station was able to reach, and the previous history of cases coming under their influence. It greatly facilitated matters in looking up a case whenever application for relief was made.

#### ARTIFICIALLY FED BABIES

When artificial feeding is prescribed for a registered baby, the mother receives the daily supply of milk from the station. If it is not a case of extreme poverty, she is required to pay seven cents for a quart bottle and four cents for a pint bottle of milk. If the case deserves relief, milk is furnished upon instruction from the central office after the case has been investigated and some organization or individual has agreed to reimburse the station for the amount of the value of the milk dispensed in the form of relief.

For the purpose of control over mothers coming to the station for their milk supply a ledger, providing a page for recording information relating to each case, is maintained; the case is identified by the case number given the registered baby. The quantity of milk dispensed each day, also the feeding formula, as prescribed by the station physician, is here recorded. All this affords an important means of control over mothers having babies registered at the station and artificially fed, as the matron dispensing the milk promptly reports to the nurse cases where the mother fails to come for her daily supply. These are immediately followed up by the station nurse, to determine the cause; usually, if the baby is well, mothers are apt to be negligent or careless in making their station purchase, and, through motives of economy, substitute cheap grocery milk.

The ledger has column-rulings with the following headings:

Date Milk Dispensed Quarts Pints Cash Received Relief charged to Formula prescribed (oz.) Whole Milk Oatmeal Water Lime Water Barley Water Boiled Water Plain Water Milk Sugar Cane Sugar Feeding Intervals

By having recorded the feeding formula of the baby receiving the milk, the nurse and the matron have conveniently at hand information as to the feeding and the quantity of milk the mother should buy for her baby. It is further used for questioning the mother in regard to the feeding formula prescribed. If no change has been made in this for several weeks, it is an additional means of bringing to the nurse's attention the failure of the mother to bring the baby to the consultations for the purpose of having any necessary changes made in the feeding formula, as no formulas are supposed to be written for a longer period than three weeks.

#### CHARACTER OF REPORTS RENDERED BY STATION EMPLOYEES

During the demonstration daily reports were made out at the close of each day's work, by both physician and station nurse, and mailed to the central office, where daily tabulations were made and the results used for general supervision and control.

#### NURSE'S DAILY REPORT CARD

The nurse made her daily reports on the card form, 4 x 6 inches, as shown in Exhibit 31.

This report provides for reporting the name of the nurse, the date, station number, hour of arrival, and hour of her departure; information relating to changes in the station enrolment by giving the cases brought forward from the previous day; the new babies received, and from where they were received; the registered cases terminated, and the reason for termination, with the total number of active cases carried forward to the next day; the first visit, and revisits of well babies and sick babies to the station; the number of first visits and revisits made by the station nurse to homes of well babies, sick babies, babies not found; the number of visits and revisits of coöperation to dispensaries, hospitals, settlements, private practitioners, charitable organizations, etc., and the total number of first visits and revisits, and total for all visits.

The form also provides for recording the number of mothers instructed individually, once or more than once, in the modification of milk or other matters relating to the care of the baby; the number of mothers instructed in classes for the first time and for two or more times, and the total number instructed.

In the lower right-hand corner of the report is recorded the number of babies fed, the number of mothers fed, and the others fed, and the total individuals to whom milk was dispensed; the number of requests for relief, the number of cases where instruction had been received to extend entire relief, partial relief, or to refuse or to discontinue relief.

Space is also given to the daily accounting of milk received and dispensed by the station, recording the number of quarts and pints carried forward from the previous day's supply; quantity received on the date to which report relates, and the total amount to be accounted for; the quantity dispensed to babies, mothers, and to others; quantity transferred to other stations, spoiled, or left over, and the total quantity accounted for.

The reverse side of the report provides for recording the name, address, time of visit, and recent action taken, and the date of the next visit for each home. Information is also recorded relating to cases transferred to hospitals by giving the name, home address, reason for transferring, name of hospital, and the date on which the case entered the hospital. For deaths among station babies the

form provides for the name, the home address, the cause, the place where the death occurred, and the date of death.

		MIL S DAI			иітт	EE		ATTENDAN	CE	First Visit	Re- visit	То	na1
		5 DAI	Mothers		4	9	1	3					
830-14			Babies, well .		4	2	1	/					
ARRIVAL O		/	Rabies, sick .			2	2	V					
DEPARTURE JA	<u>.</u>		1. 7		ale	٠.	NURSE	Visits to well be	bies	/	4	•	-
CA	SES BR	OUGHT I		D			159	Visits to sick ba	bies		4	1	1
CASES		RECE				ERMINAT	-	Visits to babies	not fou	nd /		1	1
	Ref.	Trans.	Other	Total	Trans.	Other	Total	Visits of co-oper	ration				
Re-admitted		-			-		-	Total visits				1	'n
Hos. or Disp		-			v		2	Mothers Instruc	t Ind	i.e.	-		-
Board of Health											2	-	2_
Other Organization		1		1				Mothers Instruc	t.Clas	sses			
Other N. V. M.C. Station								Babies Fed	138	MILK	Qt	s. 1	Pts.
Private Practitioner	1			1			1	Mothers Fed .	3	Carried for			
Canvass		Laure D	4	4				Gibers Fed	2	Received . Total to	ha	-	_
Removed								Total Fed	143	accounted	for /4	-	
Refused to Attend								Request for Relief		Disp. to bai	-	1	_
Dead								Entire Relief .		mothers		3	
				Partial Relief .		Disp. to ot		3.	-				
		-	-	Refused Relief	Spoiled .		-	-					
Total	/	/	4	6	2		2	Discontinue	Left Over		4		
		CA	SES CAR	RIED FO	RWARD		163	Relief		Total acc'd		43	

!					NEXT
NAME	ADDRESS	TIME	REASON	ACTION TAKEN	VISIT
1 Walter Wengels	511 6.78 St	1000	Siet	Advised	8/8
2 Jerome monio	425671.	1130	Follow up	out	8/8
3. Florence Fredle	4216,74 .	1040	Sick	advised	8/8
1 Junge Koska	503 6.73 "	110	Inus Clinic	Moth Sick wie Come	_
5 Lik Ruse	436 6.73 .	730	Follow_	we allas cline	-
6 Jenne Loka	414 6.73 "	200	Sick	adired Buy lotter	8/10
7 Franks Larsdop	420 6. 70 1	35	Kew	Sustrates - Will come	_
8 Her Lenkonds	506 6.70 .	345	Followy	lives attend clarice	_
9 Jos. Jahora	1372 1# ace	415	change	Newsonstate per Franke	_
10 Frank Jara	1431 and a	445	Sect	By letter to Tunde	8/10
SENT TO HOSPITAL	HOME ADDRESS	REA	SON	NAME OF HOSPITAL	DATE
Theo. Nazer	404 8. 7384.	Enteri	tie	Jamor La Presge	8/2
Jos. Barturer	424 6.75			]	8/3
DEATHS	HOME ADDRESS	CA	USE.	PLACE	DATE
	-			Marine Marine	

EXHIBIT 31 Nurse's Daily Report

#### DAILY REPORT OF PHYSICIAN

The stations' physicians were required to make a report each day on which services were rendered; the form of this report is shown below:

Form P-6-1-11-1500					MITTEE
DISTRICT West	PH	IYSICIAI	N'S DAIL		DATE SUBTEL 1911
Station Number	IX W.		IW		REMARKS
Hours	10 12.		2 4.39		0
INDIVIDUAL WORK				The same of	Austin Coal
Mothers Instructed	33		150		- Jung
Well Babies Examined	30		45		lack awal for
Sick Babies Examined	3		5		Topens, for
Total Babies Examined	33		50		as a rol of
CLASS WORK					
No. of Classes Held					tomsils.
No. Mothera Present					
No. Little M. Present					
CASES REFERRED			Sele In Nichola		
Hospital Surgical			2		
Dispensary   Surgical   Medical					
Board of Health				· ·	0
Outing			3		1 1274
Central Office	And the second				do Wilhomos
Total	2		5		Signature of Physician

Ехнівіт 32

It gives the district served by the physician and the date of the report. The column rulings at the right of the items shown in the first column are provided for recording separate information for each station to which services were rendered by the physician. The information called for relates to work performed by the physician, showing the hour of arrival and departure, the individual work performed by recording the number of mothers instructed and babies examined; the number of mothers and "little mothers" or older girls present at consultations. The physician also reports the action taken in referring cases, by giving the number of cases referred to hospitals and dispensaries and the reason for referring; the number referred to the Department of Health, to private practitioners, and to the central office, together with the total number of cases referred. To the right of the column rulings space is provided for remarks intended to be brought to the attention of the central office.

In addition to the foregoing, special reports were rendered by physicians and nurses when requested. No other regular report except a weekly financial statement as to the milk and supply sales was required.

#### OF COST OF OPERATING THE COMMITTEE'S MILK STATIONS DURINGCINE

	awin			Tell (					No.	U.M.	
	2000.										
					85.00.						
		3:005									
	Mino.										
						00.7					
					0690						
								10073			
				10001					32.10		
			0000.		2100						
	08300.										
	10000									10010	
	Jones									in to	

# TABLE 117.—SHOWING STATEMENT OF COST OF OPERATING THE COMMITTEE'S MILK STATIONS DURING THE MONTH OF SEPTEMBER, 1911 NETAL ADMINISTRATION EXPENSES STATION OPERATION EXPENSES EDU

	100			8	UPERVE	SION A	ND CEN	THAL	LDMINI	STRATI	DON EX	PRODUC										arne.	1004 000	m-max ru		mesma											EDUCA	HUSANL	EXPE	19363			CENTRAL	Atomic.
NUMBER	Executes tor Barr	Days	Salabi	es	STENOGR	LAPHY	REST	AND	PAINTIN STATIO	FO, AND	Iscne Exec	NTAL NEED	Tors		SALARIES TROS		Res		GA		Ice		CLEAN	eteg	POST. PRINTIN STATIO	PG, AND	THERP	BONE	CARP	ARE	I SCHOOL EXPER	NTAL ISSES	Ton		SALARIES SICIA		Salasis Nuss		Posts Bulla Circus	STIME.	Total	a	THOSE EN	N STA-
SOLLY.	Scarce B	ď	Amount	Average per Raby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Raby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Buly Day	Amount	Average per Baby Day	Amount	Average per Balty Day	Amount	Average per Baby Day	America	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per Baby Day	Amount	Average per liaby Day
1 2 3 4 5	201 6 150 4 193 5	5,790 5,030 5,500 5,790 5,170	\$41.38 43.04 32.15 41.38 29.77	.0071	5.79 4.32 5.57	\$.0010 ,0010 ,0010 ,0010 ,0010	5,36 4,01 5,16	.0009 .0009 .0009	3.04 2.27 2.92	\$.0003 .0003 .0003 .0005 .0005	7.51 5.61 7.22	.0012 .0012 .0012	\$62.25 64.74 48.36 62.25 44.77	.0107 .0107 .0107	45.00 51.00 45.00	\$.0137 .0074 .0113 .0077 .0107	38.09 45.09 32.59	,0063 ,0100 ,0056	1.20 7.52 1.66	.0002 .0016 .0003	6.83 4.41 21.80	.0011 .0010 .0038	5.00 4.75 9.20	.0008 .0011 .0016	.56 .46 .45	.0000 .0001 .0000	3.00 3.00 3.00	.0005 .0007 .0005	1.40 .50 1.00	.0002 .0001 .0002	2.22 8.53 1.92	.0004 .0018 .0003	103,30 123,26 116,62	.0171 .0278 .0201	22.00 25.00 22.00	.0036 .0035 .0038	75.00 153.63 150.00	.0124 .0341 .0259	.58 1.25 .38	.0000 .0002 .0000	97.58 179.88 172.38	.0161 .0399 .0297		.0140 .0786 .0607
7	143 4 186 5 174 5	3,270 1,290 5,580 5,220 5,850	23.34 30.60 39.83 37.23 41.80	.0071	4.12 5.36 5.01	,0010 ,010 ,010 ,010 ,010	3.81 4.96 4.64	.0009 .0009 .0009	2.16 2.81 2.63		5.34 6.94 6.49	.0012 .0012 .0012 .0012 .0012	46.03 59.90	.0107 .0107 .0107	45.00 58.50 91.00	.0243 .0104 .0104 .0174 .0076	23.09 27.09	.0077 .0054 .0049 .0067 .0043	.72 1.36 .80	.0002	4.25 4.80 5.20	.0013 .0010 .0009 .0010 .0015	5.75 3.75 .20	.0012 .0013 .0007 .0000 .0008	.54 .97 1.02		3.00 3.30 2.95	,0007 ,0006 ,0006	.40 .20 1.30	.0000 .0000 .0002	3.49 .05 1.21	.0008 .0000 .0002	138.77	.0201 .0179 .0266	26.00 25.00 32.00	.0060 .0044 .0061	75.00 125.00 150.00	.0174 .0224 .0287	1.97 .41 1.00	.0004 .0000 .0001	100.45 102.97 150.41 183.00 183.85	.0260 .0269 .0350	235.24 310.33	.0548 .0556 .0724
11 12 13 14 15	242 7 116 3 157 4	7,200 7,260 1,480 1,710 1,410	51.86 24.79 33.60	.0071	6.97 3.33 4.52	,0010 ,0010 ,0010 ,0010 ,0010	6.46 3.09 4.19	,0009 ,0009 ,0009	3.66 1.75 2.37	.0003	9,04 4,32 5,86	.0012 .0012 .0012 .0012 .0012	77.99 37.28	.0107 .0107 .0107	45,00 45,00 45,00	.0093 .0055 .0129 .0096 .0102	40.10 25.10 45.10	.0072	1.20 .72 2.00	.0002 .0002 .0004	5.10 3.75	.0006 .0015 .0008	4.00	.0011	1.58 1.00 1.54	.0000 .0002 .0003 .0003	3.05 3.00 3.00	.0004 .0009 .0006	3.75	.0005 .0000 .0001	12.64 1.22 .09	.0017 .0003 .0000	114.72 85.44 105.08	.0158 .0246 .0223	28.50 22.00 23.00	.0039 .0063 .0048	162.42	.0201 .0215 .0344	.56 1.08 .36	.0000 .0003 .0000	175.68 98.08 185.78	.0241 .0281 .0394	371.97 368.39 220.80 341.40 280.37	.0507 · .0634 .0728
16 17 18 19 20	112 3 252 7 194 5	3,510 3,360 7,560 5,820 3,570	23,96 53,93 41,49	.0071	3.22 7.25 5.58	,0010 ,0010 ,0010 ,0010 ,0010	2.98 6.72 5.17	.0009 .0009 .0009	1.69 3.81 2.93	,0003 ,0003 ,0003 ,0005 ,0005	4.18 9.40 7.23	.0012 .0012 .0012 .0012 .0012	81.11 62.40	.0107 .0107 .0107	40.50 48.00 45.00	.0092 .0121 .0063 .0077 .0119	37.10 14.10 35.10	.0111 .0018 .0060	.88 .72 3.84	.0003 .0000 .0007	10.30 6.20 18.48 5.20 12.90	.0018 .0024 .0009	7.00 5.50 6.00	.0021 .0007 .0010	.62 .80 .20	.0003 .0002 .0001 .0000 .0001	3.00 3.00 3.00	.0009 .0004 .0005	1.40 1.75 .10	.0002	2.63 6.50 6.40	.0007 .0008 .0019	99,33 98,83 104,84	.0296 .0131 .0180	25.00 33.00 25.50	.0074 .0043 .0043	112.50 80.00 147.50 150.00 75.00	.0238 .0195 .0257	.86 .50 1.40	.0002 .0000 .0002	105.86 181.00 176.90	.0315 .0239 .0303		.0717 .0477 .0591
21 22 23 24 25	104 3 184 5	4,110 3,120 5,520 5,370 3,390	22.30 39.41 38.37	.0071 .0071 .0071 .0071 .0071	3.00 5.30 5.16	.0010 .0010 .0010 .0010 .0010	2.78 4.91 4.78	.0009 .0009	1.58 2.79 2.71	.0005 .0005 .0005 .0005 .0005	3.89 6.87 6.69	.0012 .0012 .0012 .0012 .0012	59.28 57.71	.0107 .0107 .0107	45.00 48.87 67.00	.0097 .0144 .0089 .0123 .0193	25.10 40.10 32.10	.0073 .0060	3.84 1.12 .72	.0012 .0002 .0001	14.10 4.40 8.92 4.28 5.15	.0014 .0016 .0008	4.75 5.25 .70	.0010 .0015 .0010 .0001 .0006	1.32 .30 .80	.0001 .0004 .0000 .0001 .0000	3.00 3.00 3.00	.0010 .0005 .0005	.20	.0000 .0000 .0000 .0002	1.05 2.50 .14	.0005	88.76 110.26	.0284 .0200 .0203	20.50	.0104 .0042 .0038	75.00 75.00 131.86 125.00 80.00	.0240 .0238 .0232	1.01	.0000	96.31 108.12 155.84 146.51 101.31	.0282 .0272	230.43 325.38 313.16	.0738 .0590 .0583
29		3,180 3,420	41.48 22.30 22.71 27.37		5.58 3.00 3.05 3.28		5.17 2.78 2.53 3.03	,0009 ,0009 ,0009	2.93 1.58 1.61 1.72		7.23 3.89 3.96 4.25	.0012	62.39 33.53 34.16 36.65	.0107 .0107 .0107 .0107	62.50 45.00 45.00 65.00	.0190	40.10 32.50 42.60 32.50	.0095	.80 1.65 1.68 1.66	.0003 .0005 .0005	5.01	.0022 .0037 .0016 .0015	3.25 1.50 3.00	.0003 .0006 .0005 .0009	.46 .73 .68 1.34	.0002 .0000 .0002 .0002 .0004	3.00 3.00 3.00 3.00	.0005 .0010 .0009 .0009	.60 1.10 1.03 .10	.0000	,90 1,25 ,82 ,63	,0001 ,0004 ,0002 ,0001	98.32 102.99 109.26	.0214 .0315 .0324 .0319	32.50 15.75 23.58 23.00	.0055 .0050 .0074 .0067		.0257 .0279 .0385 .0387	.40 .51 .61 1.01	.0000 .0001 .0001 .0002	103.44 146.69 156.51	.0314 .0331 .0461 .0457	369.70 235.31 283.84 302.42	.0635 .0754 .0592 .0584
-	113												1	-								_			-	-																	319.35 99227.35	-

#### THE COST OF OPERATING MILK STATIONS

The cost of conducting the milk-station work may be divided into four divisions of expenditures as follows:

- 1. Supervision and central administration.
- 2. Maintenance of milk-station plants.
- 3. Medical and educational work.
- 4. Loss on milk sales and cost of relief.

Under expenditures on account of "supervision and central administration" may be properly included the salaries of the administrating official or officials, the salaries of supervisors, and the salaries of the central office force, including chief clerk, relief agent and clerks, the cost of stenographic service, rent of central office, telephone service, postage, printing, stationery, office supplies, and incidentals.

The "maintenance of milk-station plants" would also include wages of matrons and interpreters, rental of stations, gas, ice, cleaning, when not performed by station matron, postage, printing, stationery, carfare, supplies, and incidental expenses.

The "medical and educational work" carried on at the station would include salaries of nurses and physicians; cost of printing and circulating posters, bulletins, and circulars; medical and educational supplies, carfare, and incidental expenses.

"Loss on milk sales and cost of relief" represents the difference between the purchasing and the selling price of the milk if it is sold below cost, and the cost of milk dispensed in the form of relief.

Table No.117 is a statement giving the distribution of the expense of operating 31 of the Committee's milk stations during the month of September, 1911; this month is taken because it represents very nearly the average monthly expenditures during the period of demonstration. This analysis will give an idea of the system employed by the Committee in arriving at the cost of operating the various stations.

The statement is divided into three parts—i. e.: (a) Supervision and administration; (b) maintenance of milk station plants; (c) medical and educational expenses. Inasmuch as the "loss on milk sales" is a constant factor, and the relief in the form of milk was extended through other agencies, these expenditures are not shown in this statement.

Expenses on account of "supervision and central administration" are charged to the different stations on the basis of their enrolment, consequently the amount shown as the average expenditure per baby day under this item is the same for all the stations, although the actual amounts charged against the stations vary by reason of the difference in the average daily enrolment of babies.

The total amount expended during the month of September, as shown by the statement on account of "supervision and central administration," was \$1559.99, or an expenditure of \$0.0107 per baby day. These amounts are made up as follows:

	TOTAL	AVERAGE COST PER BABY DAY
Salaries	\$1037.14	\$0.0071
Stenography		0.0010
Rental and telephone		0.0009
Postage, printing, and stationery		0.0005
Incidental expenses	180.86	0.0012

The expense of "maintenance of milk station plants," exclusive of the cost of "medical and educational work" during this month, was \$3274.35, or an average of \$0.0225 per baby day, made up as follows:

	TOTAL	AVERAGE COST PER BABY DAY
Salaries of matrons\$	1609.34	\$0.0111
Rental	1014.20	0.0069
Gas	51.52	0.0004
Ice	238.70	0.0016
Cleaning	140.95	0.0010
Postage, printing, and stationery	23.06	0.0001
Telephone	92.30	0.0006
Carfare	22.45	0.0001
Incidental expenses	81.83	0.0006

#### MAINTENANCE OF MILK-STATION PLANTS


The cost varied considerably among the different stations. The amount spent for rent varied from \$14.10 to \$45 per month. The expense incurred for cleaning the stations varied because at some of the stations the matrons did this work, while at other stations it was deemed advisable to employ outside help, either because the matron could be used to better advantage in other ways, or because she was not strong enough to do scrubbing and cleaning. The amount expended for wages of matrons varied from \$32.27 to \$91 per station. In some stations only one matron was required, while in others the character and size of the enrolment necessitated two and sometimes three matrons.

The cost for "medical and educational work" represents expenditures of maintaining the staff of physicians and nurses and expenditures incidental to their work. The amount expended under this heading during the month of September was \$4393.01, or an average of \$0.0302 per baby day, made up as follows:


	TOTAL	PER BABY DAY
Salaries of physicians	\$772.20	\$0.0053
Salaries of nurses		0.0247
Posters, bulletins, and circulars for dis-		
tribution	23.62	0.0001

The expenditures, therefore, for the month of September may be summarized as follows:

TOTAL	AVERAGE COST PER BABY DAY
Supervision and central administration .\$1559.99	\$0.0107
Maintenance of milk station plants 3274.35	0.0225
Medical and educational work 4393.01	0.0302
\$9227.35	\$0.0634



FORMS USED IN THE BUSINESS ADMINISTRATION OF MILK STATIONS



Upon the basis of the foregoing statement the yearly cost for each baby supervised by the milk stations may be estimated as \$23.14, made up as follows:

Supervision and administration	8.22
\$	23.14

#### PHYSICIANS

Ten physicians were placed in charge of the medical consultation work of the milk stations, two of whom served as volunteers at one consultation per week; the others were paid according to the number of consultations held, their salaries varying from \$50 to \$100 a month.

The Committee was fortunate in that it could pay for this service, as it placed its relations to the physicians upon a business basis. Their work was inspected by the supervising physician, and they were required to conform to a uniform scheme of procedure in discharging their duties. Daily reports were required, and punctuality in attendance was made obligatory. Failure to conform to instructions and inaptitude for the consultation work demanded several changes in the staff during the demonstration.

The milk station work calls for physicians possessing not only the technic of infant feeding, but the ability and personality which will enable them to reach, benefit, and hold the most ignorant mother. It also requires patience to do the systematic routine and painstaking tasks incidental to intelligent and scientific record keeping.

The quality of the service rendered by the medical staff is evidenced by the facts set forth in the chapter dealing with the results achieved.

#### NURSES

All the nurses appointed, with one exception, were graduates from recognized training-schools for nurses. The nurse appointed without training-school experience proved unsatisfactory after several weeks' trial. Of the 63 nurses appointed, 17 gave up the work during the six months' demonstration, either because they received appointments to civil-service positions or could not adapt themselves to the milk-station work. Only six of the nurses appointed were familiar with any language other than English, although a large proportion of the station enrolment was drawn from non-speaking English homes. It has been the experience of the Committee that, with an interpreter or a matron familiar with the language of the district, better results could be obtained by employing a station nurse of a different nationality than that comprising the station enrolment. Both the Italian and the Jewish mothers seem to have more confidence in the American nurse than in one of their own nationality. A case in point is our experience at the Third Street station. This station was opened during the month of July, in a locality largely made up of Polish Jews. A nurse of the same nationality was appointed, and began an active canvass to build up the station enrolment. For nearly four weeks comparatively little progress was made,

although the district is densely populated. The nurse reported that in spite of all she could do the mothers would not consent to bring their babies to the station. As the experience of this nurse was unusual, special aid was given her by the central office in advertising the station in the neighborhood. At the end of four weeks a change was decided upon, and an American nurse with no knowledge of the neighborhood or its language took charge of the work, with the result that within two weeks a full enrolment of babies was obtained, and this became one of our most successful stations. Her personality, enthusiasm, and tact were sufficient to more than overcome any handicap caused by lack of familiarity with the language of the neighborhood. In this case the new nurse soon found several interpreters from among the "little mothers" of the neighborhood, and one of them accompanied her on her daily rounds.

Out of a total of 6684 nurse days served, only sixty days' absences were due to illness. When it is considered that this service was rendered during the six months of the most tedious and trying season of the year, the absence on account of illness is remarkably low, and can be attributed only to the keen interest and sense of responsibility of the nurses, which made them feel that absences from their stations might mean disaster to some of the lives intrusted to their care. Their devotion to the work is further manifested by the fact that the time record shows the nurses to have given 2441 hours, or 349 nurse days, in overtime and Sundays, without extra remuneration.

#### MATRONS

The matrons or nurses' assistants were chosen from a list of 185 applicants. Of these, 68 appointments were made during the period of the demonstration. Preference was given to capable applicants residing in the locality of the station and speaking the language of the prospective station enrolment.

The duties of the matron were so varied that it was a problem to obtain the right kind of women. They were required to be able to keep a record of milk sales, to attract mothers to the stations, to be neat in their work, which included mopping, dusting, and keeping ice-box and utensils scrupulously clean, and to render general assistance to the physician and nurse in the instruction of mothers. Some of the older matrons with personal maternal experience proved unsatisfactory because they persisted in instructing the mothers in their own language according to their own ideas, and otherwise undermining the influence of the physician and nurse.

Only 15 of the 68 appointments made for matrons were given to Americanborn women; the other appointments were given to Italian, Hebrew, German, Hungarian, Polish, and Bohemian applicants.

Out of 6614 days served by matrons, only 49 absences were due to illness. As a rule, the same devotion and enthusiasm which prevailed among the nurses marked the work of the matrons.

#### CHAPTER X

### Extension of Milk Station Activities

The milk station has demonstrated what can be done in preventing sickness and useless waste of infant life through the education of mothers in the feeding of their babies. As the station organization and methods become perfected greater results may reasonably be expected.

This chapter will show how the milk station can be made the basis for-

- The prevention of deaths of early infancy due to the lack of proper prenatal care.
- 2. The official supervision and control of midwives.
- The official supervision and control of foster mothers having boardedout babies.
- 4. The education of mothers in the care of children under school age.

All these activities may be properly carried on by the milk stations established and maintained by the city, while stations conducted by private agencies may eventually be expected to make the education of expectant mothers and the supervision of children under school age a regular feature of their work.

The following table shows the main causes of deaths among infants before they reached one year of age, and the rate per thousand in each case:

#### TABLE 118.—INFANT DEATHS, CAUSES, AND PROPORTION (NEW YORK CITY, 1909)

Congenital debility, malformations, and ill-				
defined causes of early infancy	Out	of	1000	deaths
Diseases of digestive system	**	44	"	"
Diseases of respiratory system			**	"
Infection, contagion, and tuberculosis 94	**	"	**	"
Diseases of nervous system	**	"	44	**
All other causes	"	"	**	"

The first three main causes account for 85 per cent. of the total infant mortality in New York city. It is not comforting to know that these deaths are largely preventable; that out of every 1000 infant deaths 326 are due in a great measure to the lack of prenatal care as a result of ignorance and social sins; 282 are largely due to ignorance and bad food, and 232 are mainly due to ignorance and bad air. In all these causes ignorance must be given the first place; hence the solution of the problem lies in education.

The systematic instruction by which the expectant mother will come to act intelligently and wisely in the care of herself will not only enable her to produce a better and healthier baby, but will help her to nurse the baby as nature intended. The fact, therefore, of starting the education of the mother before the baby arrives will not only reduce the infant death-rate from congenital diseases, but will afford an additional means of reducing the death-rate from enteritis.

The following table shows the reasons for terminating breast-feeding in the case of 1656 milk-station babies artificially fed before they reached the age of nine months:

#### TABLE 119.—REASONS FOR ARTIFICIAL FEEDING

	Number	PER CENT.
Artificially fed under nine months	1656	100.0
Insufficient milk		19.9
Mother ill		15.5
Weaned, mother not willing to nurse.	210	12.8
Mother working		12.3
Abnormalities of breasts	176	10.6
Mother dead		10.2
Mother's milk "does not agree"	159	9.7
Mother pregnant		9.8

The resorting to artificial feeding because of lack of milk, ill health, and abnormalities of breasts constituted 46 per cent. of the reasons for depriving the baby of its right to nurse, and thereby lessening its chances to live through the first year of life.

The milk station offers a means of not only reaching the expectant mother several months prior to confinement, but affords facilities by which she may be successfully instructed in the hygiene of pregnancy.


This Committee began an experiment during the early part of last summer which would cover several thousand expectant mother cases, to determine the best working method by which the milk station could be utilized in educating the expectant mother. The nurses were enthusiastic over this feature of the work, and within a few weeks nearly 1000 cases were regularly registered at the stations and were receiving instructions. The milk station gives the nurse and physician the best and most delicate means of approaching the mother on the subject. Nearly 15 per cent. of the expectant mothers registered was made up of primipara cases, showing that even the young married woman was interested and came to confide her secret to the station nurse and receive advice. The following popular little letter was distributed with good effect to 10,000 mothers, and read by them in their native tongues. (Exhibit 34.)

The leaflet containing this letter gave a list of milk stations, as also a list of hospitals receiving maternity cases and hospitals and dispensaries maintaining outdoor departments. Furthermore, 5000 circulars on the "Prevention of Blindness" (published by the Society for the Prevention of Blindness) were distributed through the milk station in order to warn mothers of the danger of improper care at confinement.

#### EXPECTANT MOTHERS REGISTERED AT MILK STATIONS

The 31 stations of the Committee were used as centers where expectant mothers were registered, many of them already having a baby under the supervision of the milk station. Those not having babies registered at the station were brought under its influence by the nurse while visiting the home, and by being referred to the station by mothers already registered. For each expectant mother with whom the nurse came in contact a special card was made out. (Exhibit 35.)

WAITING FOR ICE



#### A LETTER TO EXPECTANT MOTHERS

Dear Mothers:

Last year fully five thousand babies died in New York before they were one month old. They died because they were born too weak to combat the diseases of babyhood. They were born weak because their mothers did not properly care for themselves during the period of pregnancy; overwork, lack of exercise, insufficient or improper nourishment, irregularity in eating and sleeping, worry and nervousness were some of the many causes.

Every mother desires to have her baby born healthy and strong. Every baby has a right to be so born. You can do much at this time to insure your baby being born well and strong. Prepare yourself in advance—be regular in your habits; build yourself up by eating wholesome food; avoid beer, whiskey and drugs; talk to your doctor about yourself, or go to any BABIES' PURE MILK AND HEALTH STATION for advice.

When the time comes for the baby to be born, you must use special care so as not to have your own health or that of your baby impaired. Remember that one-half of all the blindness in this world is due to im-

proper care of the baby's eyes immediately after birth.

To avoid having your own health and that of your baby impaired at the time of birth, you will need special care. Obtain the best. It will be worth while. If you can go to the hospital, do so by all means; there you will receive proper medical and nursing care for yourself and baby. Select from the list of hospitals given in this circular the one to which you desire to apply for admission, and make arrangements at least two weeks in advance of the date of confinement. If you cannot leave your home because of other children, go to a "Blue Front" BABIES' PURE MILK AND HEALTH STATION for advice; there the nurse will be glad to tell you of hospitals and settlements which will provide medical treatment and nurse services in your own home. Many agencies will furnish women physicians if desired. These services can be obtained at a small charge, or free if you cannot afford to pay.

Do not forget you owe it to your unborn babe to bring it into the world well and strong. This you can do if only you are willing to learn how.

#### Ехнівіт 34

This card is a distinctive blue color, and was filed at the station under the month in which the confinement was expected to take place.

Two nurses were employed especially for following up these cases and giving the expectant mothers the proper instruction—this was in addition to the general instruction given by the station nurse. When the special nurse visited the home, a more detailed history of the case was obtained. The form of this record is shown in Exhibit 36.

These were then forwarded to the central office, where they were classified as to the necessary attention which the case would require, and filed under date of next visit for the purpose of instruction or for the completion of the record.

	EXPECTANT MOTHER
	Name
E	Address
O	By Whom Referred
-4-28-12-2000 O MILK COMMITTEE	Date Expected Confinement
13-20 K CO	Medical SupervisionOther Supervision
Form K-4-28-12-2000	Family consists of HusbandChildrenAges
rm K-	Home Conditions: GoodFairBadMother's Health:GoodFairBad
O Form K NEW YORK	Referred to
NEW	Remarks
	Dates VisitedBy Whom

Ехнівіт 35

COMMITTEE FOR THE REDUCTION OF INFANT MORTALITY OF THE NEW YORK MILK COMMITTEE EXPECTANT MOTHER							
Name of Mother				Address		Age	
Where Born	Kind of Wor	k			Place	Management of the Control of the Con	
					(1	actory, Store, I	(ome, etc.)
History of Previous Indicated by M-Miscarriage P-Premature	Pregnancies  Age at Death	1st	2nd	3rd	4th	5th	6th
N-Normal Birth S-Still Birth TTwins	Age at Death	7th	Sth	9th	10th	11th	12th
Date Stopped Work be	fore present Confi	nement	Dat	e Began Work	after present (	Confinement	
Date of Birth	Normal	Place	Atten	dant at Birth	(Check) Physic	ian Midwife	Nurse
Date of Death	(Yes	or No) (H Cause	espital or Hom	The second second second	h after 1 Mont	h old. Goo	d Poor
Breast Fed	Bottle F	ed	Dat	e Registered at	Station		
Date of Report							
							NURSE

A careful canvass was made to ascertain the requirements for admittance at lying-in hospitals, and facilities available for rendering service at confinement in the homes. Dispensaries were visited and arrangements were made for sending patients for preliminary examinations. The card form shown below was used in referring cases to the various agencies rendering services to pregnant women:

	REDUCTION OF INFANT MORTALITY OF THE ORK MILK COMMITTEE
То	191
	is hereby referred to you
(Station stamp here)	NURSE

Ехнівіт 37

Much valuable assistance was given by district nurses from various settlements. Relief agencies came to the rescue where the home conditions were deplorable and mothers sick or undernourished by reason of poverty.

#### CHARACTER OF INSTRUCTION TO EXPECTANT MOTHERS

Every endeavor was made to obtain the enrolment of the cases before the sixth month of pregnancy; this was not always possible, yet essential if the instruction was to be thorough and effective.

The character of the instruction varied according to existing conditions. The improperly nourished were especially advised as to diet, while the sick were placed under medical care. The health of the mother was given first consideration, and all possible means were employed to improve her general health; this included instruction as to her habits, daily routine, exercise, diet, condition of bowels, clothing, etc. In every case the kind and amount of instruction were dependent upon the difficulties which the case presented. This of necessity left much to the discretion of the nurse, and in order to have the cases handled as

uniformly as possible, the special nurses visited each case in the home after the preliminary instruction had been given by the station nurse.

The special nurses were selected for their tact and ability to instruct the mothers. They continued the visiting and instructing until the baby had reached one month of age, when the case was referred to the nearest milk station. The results obtained in this work with 500 expectant mothers are encouraging. They are not given because the number is too small to prove conclusively the value of prenatal instruction in reducing infant mortality in New York. Furthermore, this work was merely preliminary to determine the best working organization methods and records for an intensive study and experiment along this line during 1912.

#### THE MILK STATION AS A BASIS FOR OFFICIAL CONTROL OVER MIDWIVES


The midwife has been condemned as incompetent and dangerous to both mother and baby by her slovenly and ignorant ways. She has had many accusers and few loud defenders. Attempts have been made to bring her under rigid official supervision. She is required to have a permit to practise, and she must comply with certain regulations, one of which is to have a proper outfit for caring for her cases. She is visited occasionally in her home by an inspector from the Health Department, just for what purpose it is difficult to understand, unless it is to see if she still has her license and outfit. At first these official visits caused her not a little concern, but she soon began to see that they were merely formal; the possession of a clean outfit, license, and ready answers to routine questions are now rarely lacking.


The milk station offers a medium for exerting adequate control over the midwife. Her license or permit should be issued through the milk station nearest her home; she should be made to feel that a renewal of the permit depends largely upon her ability to demonstrate to the station nurse and physician that she has not only performed her own work well, but has actively coöperated in advancing the municipal milk station work. The mutual advantage in bringing the midwife in touch with the station nurse and her work is apparent.

THE MILK STATION AS A BASIS FOR SUPERVISION OF BOARDED-OUT BABIES

The relation of the foundling and boarded-out baby to the infant mortality rate is an important one. The official supervision of these babies is intrusted to the Health Department, and is part of the work of the Division of Child Hygiene. Foster mothers are required to take out permits from the Department, which are issued after inspection of the premises, and visits are supposed to be made at regular intervals by inspectors. At best these inspections can be only at considerable intervals, and it is doubtful if the supervision can be adequate. It is also undoubtedly true that a considerable number of babies are living with foster mothers of whom the Health Department has no record, and over whom it can exert no supervision.

The majority of these babies come from two institutions in this city which maintain their own inspection service in addition to that maintained by the





Under School Age. A Problem to be Solved



Health Department. That the mortality among the charges of these institutions is high may be gathered from the following figures, published by one of them. In 1908, among 2190 admissions under one year, there were 1097 deaths, or a mortality of 49.2 per cent.; in 1909, 2237 admissions, with 1253 deaths, or 56.5 per cent. mortality. In fairness it must be remembered that a very large proportion of these babies are illegitimate, and have already been deprived of the breast on admission. Figures recently published show that in certain European institutions of a similar character the mortality is high, and in some cases higher,\* but this does not mean necessarily that it cannot be lowered.

The establishment of municipal milk stations throughout the city opens the way for bringing boarded-out babies under a more thorough and effective system of supervision, by making the milk station the headquarters in each district for officially dealing with such babies and their foster mothers. A record of foster mothers in the district could be kept at each milk station, and the home supervision carried on by the nurse. It could also be made mandatory for the foster mother to bring her baby to the milk station at regular intervals for examination by the station physician. The control could be made even more effective if the agency at whose expense the baby is being boarded out were to demand a record of attendance before payment for board is made.

During the summer of 1911 a number of babies placed out by the two large institutions were referred to the milk stations for supervision, and the results showed that this can be carried on without interfering with the authority of the institutions over their charges. Both institutions require that babies who become sick shall be returned for treatment, and this relation with the milk station should insure their return before they become desperately sick, and should also give the institution valuable information as to the care being given its charges. It would save the Health Department much time and labor expended by its inspectors in following up the cases all over the city, and make the supervision more thorough and economical.

# THE MILK STATION AS A BASIS OF SUPERVISION FOR CHILDREN UNDER SCHOOL AGE

Once a child reaches school age and is enrolled in a public school, its mental and physical development becomes a matter of public concern. Modern school plants with every facility for safeguarding its life are provided. Trained teachers undertake its education; physicians employed by the city examine the child at regular intervals to discover any physical defect that may endanger its health or retard its progress in school. Whenever necessary, municipal nurses go to the home to advise the parent as to its care and to bring about the proper treatment to remove any physical defect. Municipal playgrounds and public baths are maintained for the children's use and enjoyment; but little, if anything, has been done by the municipality to safeguard the lives and promote the health among

<sup>\*&</sup>quot;Der gegenwärtige Stand der Frage über die Kindersterblichkeit in Russland und deren Bekämpfung," Hubert, "Archiv f. Kinderheilkunde," 1912, Bd. lvii, p. 351.

that portion of our population represented by children between the second and sixth years of life.

A beginning has already been made to exert official supervision of the infant population by establishing municipal milk stations with physicians and nurses to educate mothers in the care and feeding of their babies, but no comprehensive plan has been adopted by the city to make certain that the child coming to the public school for the first time is not handicapped by unnecessary physical and mental defects. Wherever careful physical examinations have been made of the first-year pupils in our public schools, one child out of every three has been found to have some physical defect needing attention. In many instances these defects have already impaired the general health of the child and have acted as a serious handicap to its proper development. It is certain, therefore, that if the condition of the child had been made known to the parents before it reached school age and the defect treated or removed, it would not only make possible a better first-year pupil, but in all probability would insure against its becoming a repeater in the grades.

Milk stations offer the basis for officially reaching and supervising the child under school age, not only to bring about the correction of physical defects, but to educate the mothers in those things which will enable her to rear a healthy and well-developed child. During the past year a start was made to ascertain what could be done through the milk stations with this class of children, but no attempt was made to make this a prominent feature of the work of the milk station. According to information obtained by the Association of Infant Milk Stations, about 1200 of these older children came under the supervision of the milk station physicians and nurses, largely from homes having babies registered at the stations. In most of the stations the older children given attention were those in poor health or seriously in need of special care. In such cases suggestions were given to the mothers as to their care and treatment. Nearly 150,000 quarts of milk were dispensed by the various milk stations to this class of children.

In the Committee's stations children between two and six years of age, who were in poor physical health, were carried on a separate register from the station babies. The total number of cases which came under the attention of the station nurse were 541. Of these, the diet of 432 was supplemented by milk furnished by the station and paid for by the mothers or by relief agencies to whose attention some of the cases were brought. The medical examination of these children has shown that 248, or 46 per cent., of the number registered were in need of medical attention, and were referred to the proper agency for treatment; 6 cases were sent to hospitals for surgical operations; 18 entered hospitals for medical care. In most instances children receiving hospital or dispensary care were later referred by the station nurse to agencies prepared to furnish country outings. Of the 541 children registered, 284 were sent to the country through the coöperation of these agencies.

The mothers were carefully instructed as to the care of the older children during the visit of the nurse to the home. A special diet card was prepared by Dr. Godfrey R. Pisek during the early part of the summer, and a large number

#### DIET FOR CHILD FROM 12TH TO 18TH MONTH

FIRST MEAL-ON RISING.

(1) 1 to 2 ounces juice of a sweet crange or Pulp of 6 stewed prunes

1 ounce pineapple juice.

(2) 8 ounces milk with either swieback, or toasted biscuits or stale toasted bread.

NOTE: Fruit must be given either 1/2 hour before or 1/5 hour after milk.

SECOND MEAL-DURING FORENOON. Milk alone or with zwieback.

NOON MEAL.

(I) 6 ounces soup

3 ounces beef juice.

NOTE: Soup may be made of chicken, beef or motton.

(2) Stale bread may be added to the above.

FOURTH MEAL-AFTERNOON.

Milk or toasted bread and milk.

EVENING MEAL.

(1) 4 ounces thick gruel mixed with 4 ounces top half milk.

Taken with zwieback.

NOTE: Grael may be made of outmeat, farina, barley, bonney, wheating, or rice.

(2) Apple sauce

Prone telly.

Total milk in 24 hours. I to 11, quarts.

NOTE. A sunces is equal to a half pint.

COMMITTEE FOR THE PEDUCTION OF INFANT MORTALITY

NEW YORK MILK COMMITTEE

#### DIET FOR CHILD FROM 18TH TO 24TH MONTH

BREAKFAST.

Juice of I sweet orange

Pulp of 6 stewed prunes

Pineapple juice (fresh or bottled) I ounce.

A cereal such as cream of wheat, oatmeal, farina, or hominy preparations with top milk (top 16 ounces) sweetened or salted. A glass of milk, bread and butter.

NOTE: It constiguted give the fruits ½ hour before breakfast with water; if not, they may be given during the

foresoon.

Raw fruit juice must be given either 3/2 hour before or
1/2 hour after milk.

#### FORENOON.

A glass of milk with two toasted biscuits or zwieback or graham crackers.

#### DINNER.

I Broth or soup made of beef, mutton, or chicken, and thickened with peas, farina, sago or rice

Beef 'juice with stale bread crumbs; or clear vegetable

soop with yolk of egg

Egg soft boiled, with bread crumbs, or the egg poached, with a glass of milk.

(2) Dessert: apple sauce, prime pulp, with stale lady-fingers or graham wafers

Plain puddings; rice, bread, tapioca, blanc-mange, junket or baked custard.

SUPPER.

Glass of milk, warm or cold: swieback and custard or stewed

Total milk in 24 hours, 11, quarts.

COMMITTEE FOR THE REDUCTION OF INFANT MORTALITY NEW YORK MILK COMMITTEE

#### DIET FOR CHILD

#### FROM TWO TO THREE YEARS

#### BREAKFAST.

Juice of I sweet orange

Pulp of 6 stewed prunes

I ounce pineapple juice (fresh or bottled)

Apple sauce.

(2) A cereal such as oatmeal, farina, cream of wheat, hom or rice, slightly sweetened or salted as preferred, with the addition of top milk (top 16 ounces)

A soft boiled or poached egg with stale bread or toast.

(3) A plans of milk.

NOTE: If constipated give the fruits 15 hour before break-last with water; if not, they may be given during the

forences.

Milk and raw fruit juice must not be given at same meal.

#### DINNER.

(f) Broth or soup made of chicken, mutton or beef, thickened with arrowroot, split peas, rice, or with addition of the yolk of an egg or toast squares.

(2) Scraped beef or white meat of chicken or broiled fish (small amount)

Mashed or baked potato with fresh peas or spinach or

(3) Dessert: apple sauce, baked apple, rice pudding, junket or custard.

SUPPER.

(1) A cereal or egg (if egg is not taken with breakfast) with stale bread or toast

Bread and milk or bread and cocoa or bread and custard.

(2) Stewed fruit.

COMMITTEE FOR THE REDUCTION OF INFANT MORTALITY NEW YORK MILK COMMITTEE

#### DIET FOR CHILD

#### FROM THREE TO SIX YEARS

#### BREAKFAST.

- (1) Fruits: an orange, apple, pear or stewed prunes.
- (2) Cereal: oatmeal, hominy, rice or wheat preparations, well cooked and salted, with thin cream and sugar

Egg: soft boiled, poached, omelet or scrambled.

(3) Milk or cocoa.

#### DINNER.

- (1) Soup: beef, chicken or mutton.
- (2) Meat: chicken or beefsteak or roast beef or lamb chops
- (3) Vegetables: spinach or carrots or string-beans, peas, cauliflower tops, mashed or baked polato, beets or lettuce (without vinegar).

Bread and butter not fresh bread or rolls.

(4) Dessert: custard, rice or bread or tapioca pudding, ice cream (once a week), cornstarch pudding (chocolate or other flavor), stewed prunes, or baked apple.


#### SUPPER

(1) Milk toast or graham crackers and milk

A thick soup, as pea, or cream of celery with bread and butter

A cereal and thin cream with bread and butter.

2: Stewed fruit: custard or a plain pudding: jam or jelly.



of these were distributed by the milk station nurses. (See Exhibit 38.) These special diet cards gave suggestions for each meal for children of different ages; for example, the diet card relating to children from twelve to eighteen months gave suggestions for the first meal on rising; the second meal during the forenoon; the noon meal; the fourth meal during the afternoon, and the evening meal, and the total amount of milk which should be given during the twentyfour hours. Similarly, the diet is prescribed for children from the eighteenth to the twenty-fourth month, two to three and three to six years of age. For the child of two to three years the mother is told that the breakfast should consist of fruit, cereal, and milk, or fruit with soft-boiled or poached egg, with stale bread or toast and milk. She is told that if the child is constipated, to give fruits one-half hour before breakfast with water, and if not, the raw fruit may be given during the forenoon, and that milk and raw fruit-juice must not be given at the same meal. That the diet of the midday meal should consist of broth or soup, meat or fish, vegetable, and dessert. The kinds of soup or broths are given, as is also suggested the kind of meat, vegetable and dessert. For the evening meal, a diet of cereal or eggs with stale bread or toast; or bread and milk; or bread and cocoa; or bread and custard and stewed fruit are prescribed. It is difficult to estimate the amount of good the distribution of these diet-cards did, but the mothers seemed glad to receive them, and in many instances followed the directions explicitly, particularly in those cases where the children were in delicate health. The influence of the milk station in promoting the health and welfare of the older children cannot be overestimated.

## Appendix

# COMPARISON BETWEEN THE BACTERIA CONTENT OF MILK DISPENSED BY THE MUNICIPAL MILK STATIONS AND THAT DISPENSED BY THE NEW YORK MILK COMMITTEE STATIONS

PERIOD, AUGUST 22 TO SEPTEMBER 9, 1911

The municipal stations dispensed pasteurized milk and the New York Milk Committee dispensed raw milk from tuberculin-tested herds. In both cases samples were collected and examined by the same laboratory.

	New York Milk Committee Stations	MUNICIPAL MILK STATIONS
Date	Bacteria content per c.c.	Bacteria content per c.c.
	22 28,500	
	22	
11	22	
"	22	
11	22	
Sept.	1	
ocpt.	1	
11	1	
"	1	
11	2	22.000
**	2	** ***
44	2	14,000
**	2	9,500
**	5	
**	5	
**	5	
"	5	
**	5	
11	5	
11	6	42.000
**	6	40,000
**	6	15,000
**	6	12,000
**	6	125 000
11	7	8,300
**	7	
**	7	
11	7	
11	7	
**	8	
- 11	8	
11	8	
**	8	
**	8	
"	9	11,000
11	9	12.000
**	9	
**	9	
16	9	

#### STATEMENT SHOWING BACTERIA CONTENT OF SAMPLES OF MILK TAKEN FROM NEW YORK MILK COMMITTEE STATIONS AND EXAMINED BY DR. NORTH'S LABORATORY AND DEPARTMENT OF HEALTH

(Samples taken four to ten hours after delivery of supply to station and two to three hours after the mothers had received their day's supply.)

S M			BACTERIA CONTENT PER C.C.						
1911	Number of Samples	UNDER 5000	5000 TO 10,00 0	10,000 TO 20,000	20,000 TO 30,000	30,000 TO 50,000	50,000 TO 75,000	75,000 TO 100,000	OVER 100,000
May	11	6	4	1					
June	23	10	6	4	1	1	1		
July	23		3	7	7	2	1	2	1
August	35	1	12	5	5	2	1		9
September	37	1	14	12	5	1	2		2
October	9	2	3	2	1				1
	138	20	42	31	19	6	5	2	13

#### STATEMENT SHOWING A SUMMARY OF THE WORK OF NEW YORK MILK COM-MITTEE STATIONS, AS PER TABULATION OF DAILY REPORTS

#### PERIOD, MAY 20 TO OCTOBER 31, 1911

Number of behice on hand May 20th			634
Number of babies on hand May 20th			7,951
			1,551
Received by transfer	24	291	
Hospital or dispensary	25		
Board of Health stations.			
Other New York Milk Committee stations			
Other organizations	52	121	
Received by reference		636	
Hospital or dispensary	80		
Board of Health stations			
Private practitioners	193		
Other New York Milk Committee stations			
Other organizations	120	November 1	
Received by canvass		6,538	
Readmitted		486	0.000
Number of cases terminated			3,648
Refused to continue to attend	1,657		
Removed	1,053		
Transferred	780		
To other New York Milk Committee stations367			
To Board of Health stations			
To hospital or dispensary			
To other organizations			
To private practitioners			
Died	158		
Gastro-intestinal diseases			
Other causes			
Total number of babies on hand October 31st			4.937
Total number of public on hand october officers.			2,701


STATEMENT SHOWING A SUMMARY OF THE WORK OF NEW YORK MILK COMMITTEE STATIONS,


AS PER TABULATION OF DAILY REPORTS.—(Continued.)		
Attendance at milk stations for advice		43,739
Mothers' first visits for advice		
Well babies' first visits		
Well babies' revisits		
Sick babies' first visits		
Sick babies' revisits		39,586
Home visits	7 462	
To well babies' first visits		
To sick babies' first visits.	1,551	
To sick babies' revisits		
To babies' (not found) first visits	1,572	
To babies' (not found) revisits	. 2,060	29,227
Visits of cooperation, first visits	318	
Visits of cooperation, revisits		524
Number of mothers instructed first time	12,401	
Number of mothers instructed two or more times	9,964	
Total number of mothers instructed individually		
Total number of mothers instructed in class	894	22,365
Total new requests for free milk (relief)	903	
Number given free milk (relief)		
Milk dispensed (quarts)		.420,451
To babies. To mothers.		
To others.		
*Milk dispensed (quarts) from May 20th to May 31st		
Total number of physicians' consultations		
Total number of babies examined		. 27,095
Well babies examined (cases)		
Sick babies examined (cases)		. 562
Surgical treatment		. 302
Medical treatment		

<sup>\*</sup> Not itemized.











DATE DUE				BS
p MAR	0 2001	MAR 3 1 2	<del>01</del>	t the g, as with
- DA	7.6.20	MAR SU	72000	DUE
//	10V % 6 20	2		==
		-	+	
	-			
				_
DEMCO	38-296			
C28 (251) 100M	1		1	



RAGIO
Van Ingen
Infant Mortalizy and Milk Stations
JAN 2 8 1948

U. BINDERY

ON PERSONAL RESERVE SHELF

RA 610

V26

Annex

