Essentials of diagnosis : arranged in the form of questions and answers / prepared especially for students of medicine by Solomon Solis-Cohen and Augustus A. Eshner.

Contributors

Solis-Cohen, Solomon, 1857-1948.
Eshner, Augustus Adolph, 1862-
Augustus Long Health Sciences Library

Publication/Creation

Philadelphia : W.B. Saunders, 1900.

Persistent URL

https://wellcomecollection.org/works/u2hg7ts8

License and attribution

This material has been provided by This material has been provided by the Augustus C. Long Health Sciences Library at Columbia University and Columbia University Libraries/Information Services, through the Medical Heritage Library. The original may be consulted at the the Augustus C. Long Health Sciences Library at Columbia University and Columbia University. where the originals may be consulted.
This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
https://wellcomecollection.org

DUPLICATE
HX00014346

tion Compends

Essentials of Medical Diagnosis

Cohen & Eshner



RC71 C66 1900

Columbia University
in the City of New York

College of Physicians and Surgeons

Reference Library

N ALS

Th nd-books
on th ct amply
demc rs New
Seri students
and

Th sent liter-
ature d practi-
tione American
c lle xhaustive
in de uction of
"cas ok.
Th advanced
lectu ctitioner:
to th s for what
he w e; to the
former they will afford safe guides to the essential points of study.

The **Saunders New Series of Manuals** are conceded to be superior to any similar books now on the market. No other manuals afford so much information in such a concise and available form.

Any of these Manuals will be mailed on receipt of price (see List on page 3 of Cover).

W. B. SAUNDERS, Publisher,

925 WALNUT STREET, PHILADELPHIA.

Saunders' New Series of Manuals.

PHYSIOLOGY. By JOSEPH HOWARD RAYMOND, A. M., M. D., Professor of Physiology and Hygiene and Lecturer on Gynecology in the Long Island College Hospital, etc. Price, $1.25 net.

SURGERY, General and Operative. By J. CHALMERS DACOSTA, M. D., Clinical Professor of Surgery, Jefferson Medical College, Philadelphia; Surgeon to the Philadelphia Hospital. New edition, thoroughly revised and greatly enlarged. Handsome octavo volume of 900 pages, with 386 illustrations. Cloth, $4.00 net; Sheep or Half Morocco, $5.00 net.

DOSE-BOOK AND MANUAL OF PRESCRIPTION-WRITING. By E. Q. THORNTON, M. D., Demonstrator of Therapeutics, Jefferson Medical College, Philadelphia. Price, $1.25 net.

MEDICAL JURISPRUDENCE. By HENRY C. CHAPMAN, M. D., Professor of Institutes of Medicine and Medical Jurisprudence in the Jefferson Medical College of Philadelphia, etc. Price, $1.50 net.

SURGICAL ASEPSIS. By CARL BECK, M. D., Surgeon to St. Mark's Hospital and to the German Poliklinik; Instructor in Surgery, New York Post-Graduate Medical School, etc. Price, $1.25 net.

MANUAL OF ANATOMY. By IRVING S. HAYNES, M. D., Adjunct Professor of Anatomy and Demonstrator of Anatomy, Medical Department of the New York University, etc. (Double number.) Price, $2.50 net.

SYPHILIS AND THE VENEREAL DISEASES. By JAMES NEVINS HYDE, M. D., Professor of Skin and Venereal Diseases, and FRANK H. MONTGOMERY, M. D., Lecturer on Dermatology and Genito-urinary Diseases, in Rush Medical College, Chicago. (Double number.) Price, $2.50 net.

PRACTICE OF MEDICINE. By GEORGE ROE LOCKWOOD, M. D., Professor of Practice in the Woman's Medical College of the New York Infirmary, etc. (Double number.) Price, $2.50 net.

OBSTETRICS. By W. A. NEWMAN DORLAND, M. D., Assistant Obstetrician to the Hospital of the University of Pennsylvania, etc. (Double number.) Price, $2.50 net.

DISEASES OF WOMEN. By J. BLAND SUTTON, F. R. C. S., Assistant Surgeon to the Middlesex Hospital, and Surgeon to the Chelsea Hospital for Women, London; and ARTHUR E. GILES, M. D., B. Sc. Lond., F. R. C. S. Edin., Assistant Surgeon to the Chelsea Hospital for Women, London. (Double number.) Price, $2.50 net.

VOLUMES IN PREPARATION.

NERVOUS DISEASES. By CHARLES W. BURR, M. D., Clinical Professor of Nervous Diseases, Medico-Chirurgical College, Philadelphia, etc.

*** There will be published in the same series, at short intervals, carefully-prepared works on various subjects, by prominent specialists.

N

Th
on th
demo
Seri
and

Th
ature
tione
c lle
in de
" cas
Th
lectu
to th
he w
forme
Th
be su
manu
able

FRONTISPIECE.



GEO. S. HARRIS & SONS, LITH. PHILA.

THE RELATIONS OF THE THORACIC AND ABDOMINAL VISCERA,

THE ANTERIOR THORACIC AND ABDOMINAL WALLS REMOVED.

A.—Upper bone of the sternum.

B.B.*—Two first ribs.

C.C.*—Second pair of ribs.

D.D.*—Right and left lungs.

E.—Pericardium, enveloping the heart—the right ventricle.

F.—Lower extremity of the sternum.

G.G.*—Lobes of the liver.

H.H.*—Right and left halves of the diaphragm, in section; the right half separating the right lung from the liver, the left half separating the left lung from the cardiac extremity of the stomach.

I.I.*—Eighth pair of ribs.

K.K.*—Ninth pair of ribs.

L.L.*—Tenth pair of ribs.

M.M.*—The stomach.

N.—The umbilicus.

O.O.*—The transverse colon.

P.P.*—The omentum, covering the transverse colon and small intestines.

Q.—The gall-bladder.

R.R.*—The right and left pectoral prominences.

S.S.*—The small intestines.

SINCE the issue of the first volume of the **Saunders Question-Compends,**

OVER 175,000 COPIES

of these unrivalled publications have been sold. This enormous sale is indisputable evidence of the value of these self-helps to students and physicians.

Saunders' Question Compends, No. 17.

ESSENTIALS OF DIAGNOSIS

ARRANGED IN THE FORM OF

QUESTIONS AND ANSWERS

PREPARED ESPECIALLY FOR

STUDENTS OF MEDICINE.

BY

SOLOMON SOLIS-COHEN, M. D.,

Professor of Clinical Medicine and Therapeutics in the Philadelphia Polyclinic; Lecturer on Clinical Medicine in Jefferson Medical College; Physician to the Philadelphia Hospital and to the Rush Hospital for Consumptives, etc.,

AND

AUGUSTUS A. ESHNER, M. D.,

Professor of Clinical Medicine in the Philadelphia Polyclinic; Physician to the Philadelphia Hospital, etc.

ILLUSTRATED.

SECOND EDITION, REVISED AND ENLARGED.

PHILADELPHIA:

W. B. SAUNDERS,

925 Walnut Street.

1900.

GIFT

COPYRIGHT, 1900.
BY W. B. SAUNDERS.

Dr. Janeway
10/15/12

TO

J. M. DA COSTA, M.D., LL.D.,

EMERITUS PROFESSOR OF THE PRACTICE OF MEDICINE AND OF CLINICAL MEDICINE IN JEFFERSON MEDICAL COLLEGE,

THE MASTER DIAGNOSTICIAN,

THE BRILLIANT TEACHER, THE DISTINGUISHED CLINICIAN,

THIS LITTLE BOOK

IS AFFECTIONATELY INSCRIBED

BY

TWO GRATEFUL PUPILS.

PREFACE TO THE SECOND EDITION.

THIS book has been thoroughly revised and considerably enlarged. Much new material has been added, and recent additions to knowledge incorporated. Designed primarily for students, no attempt has been made to render the work encyclopedic, yet it is believed that the essentials of the subject have been set forth with sufficient fulness to serve even for reference by the practising physician. The book is not a mere compilation, but, while based upon the common knowledge and its records, represents in no small measure the results of personal observation. Care has been taken to omit mere opinions, and, as before, it has been deemed wisest in matters of doubt to set forth most prominently the prevailing view. The kind words of correspondents, the cordial reception accorded the first edition by the medical press of England and America, and the comparatively large sale of the book in Great Britain and the British colonies, have been to us sources of pleasure, for which we herewith make grateful acknowledgment.

S. S. C.
A. A. E.

PREFACE.

THIS book is intended to meet a popular demand. While the endeavor has been to make it reliable and helpful, the student is advised not to depend upon it to the exclusion of standard and more elaborate works. It is elementary in character, devoid of detail, and represents but an outline of the subject with which it has to deal. This outline must be filled from observation and further reading.

Being written especially for students, everything has been sacrificed to accuracy and brevity; references to authorities consulted have been omitted; and it has been deemed wisest to conform with prevailing views in matters concerning which there may be differences of professional opinion.

In the arrangement of material, systematic classification has often been departed from, to secure the benefit of association of ideas.

Diagnosis must be studied from patients, not from books; even the best of books can only direct the student what to look for at the bedside, and warn him against probable errors.

If this book facilitates the true methods of study, it will have accomplished its purpose.

CONTENTS.

PAGE

The Principles and Methods of Diagnosis 17
Fever 27
Simple Continued Fever—Ephemeral Fever—Febricula 31
Ardent Fever 31
Catarrhal Fever—Influenza—Epidemic Catarrh—La Grippe 32
Typhoid Fever—Enteric Fever 35
Typhus Fever 44
Cerebro-spinal Fever—Epidemic Cerebro-spinal Meningitis 46
Asiatic Cholera—Cholera Infectiosa 50
Relapsing Fever 52
Malarial Diseases 54
Malta Fever 61
Bubonic Plague 62
Beriberi 63
Yellow Fever 64
Weil's Disease 67
Leprosy 68
The Exanthemata 68
Morbilli—Measles—Rubeola 69
Scarlatina—Scarlet Fever 70
Rubella—Rötheln—Roseola—German Measles—French Measles . 74
Variola—Smallpox 75
Varicella—Chickenpox 79
Erysipelas 80
Glandular Fever 82
Miliaria—Miliary Fever—Sweating Disease 82
Dengue—Break-bone Fever 83
Diphtheria 84
Glanders—Farcy—Equinia 87
Anthrax—Wool-sorters' Disease—Charbon—Malignant Pustule—Splenic Fever 89
Actinomycosis 90

PAGE
Foot-and-mouth Disease 91
Milk-sickness 91
Hydatid (Echinococcus) Disease 92
Trichiniasis 93
Filariasis 95
Dracontiasis 95
Acute Rheumatism—Rheumatic Fever 95
Gonorrheal Synovitis 97
Syphilitic Arthritis 97
Subacute Rheumatism 98
Myalgia 98
Chronic Rheumatism 99
Acute Gout 99
Chronic Gout 101
Lithemia 101
Rheumatoid Arthritis—Arthritis Deformans 102
The Blood 104
Anemia 106
Chlorosis 107
Pernicious Anemia 107
Hyperleukocytosis (Leukocytosis) 108
Leukemia 109
Pseudo-leukemia—Hodgkin's Disease—Lymphatic Anemia—Malignant Lymphoma or Lymphadenoma 111
Infantile Pseudo-leukemic Anemia 112
Scorbutus—Scurvy 113
Purpura 114
Hemophilia 116
Addison's Disease 116
Rachitis 117
Mollities Ossium 118
The Heart 118
Inspection 118
Palpation 119
Percussion 119
Auscultation 120
Malformation 120
Dextrocardia 125
Functional Disturbance of the Heart 125
Tachycardia 126
Brachycardia 127
Irritable Heart 127

PAGE

Angina Pectoris 128
Hypertrophy of the Heart 128
Dilatation of the Heart 129
Valvular Disease of the Heart 131
Mitral Incompetency—Mitral Regurgitation 132
Mitral Obstruction 132
Aortic Obstruction 133
Aortic Incompetency—Aortic Regurgitation 133
Tricuspid Incompetency—Tricuspid Regurgitation 134
Tricuspid Obstruction 135
Pulmonary Obstruction 135
Pulmonary Incompetency—Pulmonary Regurgitation 135
Acute Pericarditis 135
Diseases of the Myocardium 138
Acute Endocarditis 139
Heart-clot 140
Diseases of the Mediastinum 141
Thoracic Aneurism 141
Arterio-capillary Fibrosis 143
Local Syncope—Local Asphyxia—Local or Symmetrical Gangrene . 144
Erythromelalgia 145
Angioneurotic Edema 146
The Respiratory System 146
Coryza—Acute Nasal Catarrh 148
Hay-fever—Hay-asthma—Rag-weed Fever—Autumnal Catarrh—June-cold—Rose-cold—Idiosyncratic Coryza—Periodic Vasomotor Coryza 149
Acute Laryngitis 150
Edema of the Larynx 151
Acute Tuberculous Laryngitis 152
Laryngismus Stridulus 153
Laryngeal Vertigo 153
Catarrhal Croup—Spasmodic Croup 154
Membranous Croup 155
Whooping-cough—Pertussis 157
Chronic Laryngitis 158
Chronic Tuberculosis of the Larynx 159
Physical Diagnosis 159
Acute Pleurisy 165
Chronic Pleurisy 169
Acute Bronchitis 170
Chronic Bronchitis 171

PAGE

Plastic Bronchitis—Fibrinous Bronchitis 172
Putrid Bronchitis 172
Bronchiectasis 173
Capillary Bronchitis 174
Catarrhal Pneumonia—Broncho-pneumonia—Lobular Pneumonia 175
Acute Croupous Pneumonia—Lobar Pneumonia 176
Pulmonary Gangrene 182
Pulmonary Tuberculosis 182
Acute Miliary Tuberculosis 191
Interstitial Pneumonitis 193
Pulmonary Emphysema 194
Pneumothorax 195
Asthma 197
New-growths in the Lungs 199
The Digestive System—The Mouth 199
Catarrhal Stomatitis 199
Aphthous Stomatitis 200
Thrush—Muguet—Parasitic Stomatitis 200
Ulcerative Stomatitis 200
Mercurial Stomatitis 201
Gangrenous Stomatitis—Noma—Cancrum Oris 201
The Tongue 202
Glossitis 202
Leukoplakia Lingualis—Leukoplakia Buccalis 202
Geographical Tongue 203
Glossanthrax 203
Nigrities—Black Tongue—Hairy Tongue 203
Mumps—Parotiditis 204
The Pharynx 204
Pharyngitis 204
Tonsillitis 206
Herpetic Sore-throat—Herpetic Tonsillitis—Herpes of the Pharynx—Common Membranous Sore-throat—Ulcero-membranous Angina—Diphtheroid Throat 208
Gangrenous Pharyngitis—Putrid Sore-throat 209
Retro-pharyngeal Abscess 210
The Esophagus 211
Esophagitis 211
Stricture of the Esophagus 211
The Stomach 212
Neuroses of the Stomach 214

PAGE

Splanchnoptosis 215
Gastralgia 215
Acute Gastritis 215
Chronic Gastritis—Chronic Gastric Catarrh 217
Dilatation of the Stomach—Gastrectasis 218
Gastric Ulcer 218
Carcinoma of the Stomach 220
The Intestines 223
Acute Enteritis 223
Membranous Enteritis 224
Cholera Morbus—Cholera Nostras 225
Cholera Infantum 225
Chronic Enteritis—Chronic Intestinal Catarrh 225
Acute Dysentery 226
Chronic Dysentery 227
Typhlitis—Appendicitis—Perityphlitis 228
Intestinal Obstruction 231
Intussusception—Invagination 232
Carcinoma of the Intestine 233
Intestinal Parasites 234
Tænia Solium 235
Tænia Mediocanellata 236
Bothriocephalus Latus 236
Ascaris Lumbricoides—Round-worms 237
Ankylostomiasis 238
Distomiasis 239
Oxyuris Vermicularis — Seat-worms — Thread-worms — Spool-worms 239
Occlusion of the Mesenteric Vessels 240
Acute Peritonitis 241
Chronic Peritonitis 243
Tabes Mesenterica 244
The Liver 244
Floating Liver 245
Congestion of the Liver 246
Acute Hepatitis 247
Acute Yellow Atrophy of the Liver 248
Abscess of the Liver 250
Interstitial Hepatitis—Cirrhosis of the Liver 252
Fatty Liver 253
Amyloid Disease of the Liver 254
Carcinoma of the Liver 254

PAGE

Hydatid Cyst of the Liver 256
Perihepatitis 258
Cholangitis—Cholecystitis 258
Biliary Calculi 261
The Spleen 263
Floating Spleen 264
Splenitis 264
Enlargement of the Spleen—Splenic Tumor 264
Splenomegaly (Banti's Disease) 265
The Pancreas 265
Acute Pancreatitis 265
Carcinoma of the Pancreas 265
Pancreatic Cysts 266
Pancreatic Hemorrhage 266
The Genito-urinary Apparatus 267
Indicanuria 273
Oxaluria 273
Pyuria 274
Albuminuria 275
Chyluria 278
Lipuria 279
Hematuria 279
Hemoglobinuria 280
Paroxysmal Hemoglobinuria 281
Endemic Hematuria 282
Diabetes Insipidus 282
Glycosuria 283
Diabetes Mellitus 283
Cystitis 288
Vesical Calculus 289
Neoplasms of the Bladder 289
Pyelitis 289
Renal Inadequacy 290
Displacements of Kidney 290
Renal Calculus 291
Acute Nephritis 292
Chronic Parenchymatous Nephritis 292
Amyloid Kidney 293
Chronic Interstitial Nephritis 293
Abscess of the Kidney 295
Perirenal Abscess 296
Tuberculosis of the Kidney 296

PAGE

Malignant Disease of the Kidney 297
Hydronephrosis 297
Hydatid Cyst of the Kidney 298
The Nervous System 299
Neuritis 302
Multiple Neuritis 303
Sciatica 305
Facial Hemiatrophy 306
Paralysis of the Facial Nerve 306
Paralysis of the Phrenic Nerve 308
Paralysis of the Musculo-spiral Nerve 309
Neuromata 310
Neuralgia 310
Migraine 311
Spinal Meningitis 312
Cervical Pachymeningitis 313
Hemorrhage into the Spinal Membranes 314
Anemia of the Spinal Cord 314
Hyperemia of the Spinal Cord 315
Myelitis 315
Chronic Myelitis 316
Acute Anterior Poliomyelitis 318
Acute Ascending Paralysis 319
Amyotrophic Lateral Sclerosis—Progressive Muscular Atrophy—Glosso-labio-laryngeal Palsy 320
Acute Bulbar Palsy 323
Pseudo-bulbar Palsy 324
Asthenic Bulbar Paralysis 324
Progressive Muscular Dystrophy 324
Arthritic Muscular Atrophy 326
Thomsen's Disease—Myotonia Congenita 327
Posterior Spinal Sclerosis—Locomotor Ataxia—Tabes Dorsalis . 327
Primary Lateral Sclerosis—Spastic Paraplegia 329
Postero-lateral Sclerosis—Ataxic Paraplegia 329
Friedreich's Ataxia—Hereditary Ataxic Paraplegia 331
Cerebro-spinal Sclerosis—Insular Sclerosis—Multiple Sclerosis—Disseminated Sclerosis 332
Paralysis Agitans—Shaking Palsy 333
Spinal Hemorrhage 335
Spinal Compression 336
Tumor of the Spinal Cord 337

PAGE

Syringomyelia 339
Morvan's Disease—Analgesic Panaris 340
Cerebral Meningitis 342
Hydrocephalus 345
Hemorrhage into the Cerebral Membranes 345
Congenital Spastic Paraplegia 346
Cerebral Anemia 347
Cerebral Hyperemia 348
Cerebritis 349
Cerebral Abscess 349
Cerebral Hemorrhage 350
Cerebral Softening 354
Cerebral Embolism 355
Cerebral Thrombosis 356
Cerebral Tumor 357
Intracranial Aneurism 359
General Paralysis of the Insane—Paretic Dementia 359
Sunstroke — Heat-stroke — Insolation — Heat-fever — Thermic Fever—Heat-exhaustion 361
Delirium Tremens 363
Plumbism 364
Torticollis 366
Occupation-neuroses 366
Writers' Cramp 367
Chorea 368
Hereditary Chorea 369
Spasmodic Tic 369
Epilepsy 370
Hysteria 372
Neurasthenia 375
Family Periodic Paralysis 377
Tetanus 377
Tetany 379
Hydrophobia 380
Aural Vertigo—Labyrinthine Vertigo—Ménière's Disease . . . 381
Exophthalmic Goiter—Graves's Disease—Basedow's Disease . . 381
Vasomotor Ataxia 384
Cretinism 384
Myxedema 385
Akromegaly 387
Rhizomelic Spondylosis 392

ESSENTIALS

OF

MEDICAL DIAGNOSIS.

ESSENTIALS OF DIAGNOSIS.

THE PRINCIPLES AND METHODS OF DIAGNOSIS.

DIAGNOSIS, in medicine, is the art and science of observing and of discriminatingly interpreting the phenomena of disease. In its study, a knowledge of the phenomena of health is an essential prerequisite. Unless familiar with the sounds elicited upon percussion of the normal chest, one cannot decide whether or not the sound heard in a given case is indicative of abnormal conditions. Unless one knows the characteristics of normal urine, he cannot hope to gain from urinalysis a clue as to the nature of a case of disease. Unless one knows the appearance of a healthy brain, he cannot determine whether the brain seen at a necropsy is or is not the seat of morbid change. Unless one knows the function of a normal joint, he cannot affirm that a joint under investigation has had its function impaired.

Having learned, by observations upon the healthy, to recognize when departures from health have taken place, one must further learn by observation of the sick to appreciate the significance of such departures. No opportunity should be lost to examine *post mortem* the organs and tissues that have been altered by disease.

A knowledge of the effects of drugs upon healthy and unhealthy persons and organs may likewise render easy a diagnosis that might otherwise be difficult.

The phenomena indicative of the existence of disease are in a general way termed *symptoms*. These may be either *subjective*—known only to the patient by his sensations; or they may be *objective*—capable of investigation by the senses of the observer,

aided, it may be, by instruments of precision. Thus pain, vertigo, nausea, ringing in the ears, are subjective symptoms, while high temperature, vomiting, unsteadiness of motion, loss of voice, are objective symptoms.

The objective manifestations of disease may be further divided into *symptoms* (in a restricted sense) and *signs*. While the two are not rigidly separable, *symptoms* may be defined as manifestations of disordered function, *signs* as manifestations of altered structure. Thus, cough is a symptom; the laryngeal congestion discovered by inspection with the mirror, the bronchial râles heard upon auscultation, are signs. Difficulty in deglutition is a symptom; a pulsating tumor in the chest, indicative of aneurism pressing upon the esophagus, is a sign. Dyspnea is a symptom; distension of the abdomen with gas (tympanites) or with fluid (ascites), causing pressure upon the diaphragm and restricting its movements, is a sign.

Thus, signs are in many cases directly explanatory of symptoms, and their discovery is one step further in the diagnosis. It must not be forgotten, however, that signs are not always to be found; that, when found, a sign may not account *in toto* for the symptom with which it is associated; and that even when sufficient to explain the symptom, the sign itself remains to be explained before the diagnosis is complete. Like all other morbid phenomena, signs must, therefore, be considered in relation with all the evidence presented in a given case.

Certain signs that are not at once manifest, but that require for their study special means of exploration, are termed *physical signs*. In its restricted sense, the term "physical signs" is applied to the phenomena elicited by special methods (*inspection*, *mensuration*, *palpation*, *percussion*, and *auscultation*), used chiefly in examination of the chest and abdomen, though often applied elsewhere; while the signs elicited by examination of the blood, the urine, the sputum, the feces, or by laryngoscopy, ophthalmoscopy, cystoscopy and the like, are not given other qualifying designation than the special names describing the respective processes. They might appropriately be termed *instrumental signs* and *signs of research*.

Phenomena corresponding with logical deductions as to the direct results of deranged function, including subjective and

objective symptoms and signs, are called *rational signs*, in contradistinction from physical signs, which denote the mechanical condition of the structures examined. Thus, in a case of valvular disease of the heart, weakness, vertigo, shortness of breath, pallor and dropsy would be termed rational signs, while the area of cardiac percussion-dulness and the character of the sounds heard upon auscultation would be termed physical signs.

Symptoms may also he divided into *general* or *constitutional* symptoms and *local* symptoms. General symptoms are those, like fever, depression, delirium, that may result from unbalancing of the organism as a whole, and are common to affections of many kinds; while local symptoms are those, like swelling or discoloration of a part and circumscribed pain or tenderness, that are confined to a certain locality and result from localized morbid conditions, usually that of the organ or tissue diseased.

Symptoms caused by local disease not at the seat of manifestation and that are not dependent upon mere mechanical influences or upon interference with related function, but that arise indirectly, as a result of nervous irritation, are termed *reflex symptoms*. Thus, the difficulty of breathing occasioned by the pressure of a mediastinal tumor upon the trachea is a local symptom, due to a mere mechanical influence; dyspnea from deranged action of the heart is a symptom dependent upon interference with related function; while an asthmatoid condition dependent upon disease of the nose is a reflex symptom. It must not be forgotten, however, that the same symptom may at one time be local; at another time, part of a general process; at another time, reflex. Thus, vomiting may indicate local disease of the stomach; or it may be part of the general disturbance caused by certain febrile diseases; or it may reflexly indicate disease in the abdomen or in the brain.

Having, by careful observation and interrogation, ascertained the *present condition* of the patient and having, by inquiry, learned his *family history* (in order to judge of the probable influence of hereditary disease, diathesis or liability) and his *previous history* (anamnesis) of health or disease (including a knowledge of his mode of life and of his surroundings, as well as the mode of invasion of the disease under investigation and

its course up to the moment of examination), it becomes necessary to interpret the information thus gained—in other words, to *make a diagnosis.*

In making a diagnosis one has to consider not only the bearing of the signs and symptoms individually and collectively, but also their relations with the phenomena of health and with each other. Further, one has to consider (and this is what may, to some extent, be learned from reading) the historical experience of the medical profession as to the significance of certain symptoms and groups of symptoms, and as to the relation of certain symptoms and groups of symptoms with lesions observed *post mortem.*

A diagnosis may thus be made (1) by the *inductive method*, reasoning, upon anatomical (structural) and physiological (functional) data, from the character of the disturbance to the organ affected and the nature of the affection; (2) by the *historical* or *empirical* method, relying upon the recorded experience of other observers and upon one's own experience that certain symptoms manifested under certain circumstances indicate the existence of a definite malady; or (3) by the method of *pathological association*, which is based upon the fact that when certain symptoms have been observed during life, definite lesions have been discovered after death. By analogy, the lessons of pathological association may be applied in affections not necessarily of a fatal character. It is obvious, therefore, that a knowledge of the various gross and minute morbid changes occurring in the body generally or in special organs, tissues or cells and of the circumstances under which special changes are likely to occur is essential for precision in diagnosis. The most satisfactory results are to be obtained when all of the methods indicated can be concurrently availed of.

Diagnosis may further be *direct*, *differential* or by *exclusion.* Diagnosis is said to be *direct* when one or more of the signs or symptoms, independently of or in relation with other symptoms or with the age, sex, physical and mental characteristics, residence or occupation of the patient, or with his family history, enable direct affirmation to be made of the nature of the malady. Thus, a paroxysm of chill, fever and sweating, in association

with the presence in the blood of characteristic parasites, permits a direct diagnosis of malarial fever to be made.

Diagnosis is said to be *discriminative* or *differential* when the signs or symptoms are suggestive of more than one disease and a decision is reached by comparison and contrast. It is necessary to compare the ideal pictures of various diseases in turn with the actual picture presented, in order to establish the resemblance or unlikeness; finally affirming the nature of the case with more or less certainty, according to the completeness and definiteness of the observations made and the degree of knowledge on the part of the clinician as to the conditions present in the respective affections under review. Diagnosis is usually differential, and as one is often compelled to balance probabilities, with incomplete evidence before him, differential diagnosis calls for the greatest knowledge and skill. Numerous examples of differential diagnosis will be found throughout this book. Reference may here be made to malarial fever, in case hematozoa are not demonstrable, and the fever of hepatic suppuration or of pulmonary tuberculosis. Careful observation of the temperature-course and painstaking physical examination may be required to establish the points of difference.

In diagnosis by *exclusion* one is unable to affirm the nature of the affection directly; and even after comparison of the evident phenomena with the phenomena of the respective diseases suggested, the points of resemblance are not sufficiently great in number or in character to warrant an affirmative conclusion in any one instance. It then becomes necessary to prove a negative; to do which, reliance must be placed entirely upon points of unlikeness. One endeavors to recall in the ideal picture of a certain malady some symptom or association of symptoms so necessary that its absence from the actual case may warrant a decided negative; or, on the other hand, to discover in the case before him some symptom or association of symptoms so incompatible with the ideal picture of the malady under consideration as likewise to warrant a negative conclusion. Thus, from the absence of what ought to be present and from the presence of what ought to be absent, one after another of the conditions discussed is set aside, until finally one remains that

cannot be so excluded; and there is reached a more or less probable diagnosis.

Diagnosis by exclusion is the least satisfactory, as one can never be sure that he has passed in review and has excluded all of the conditions that ought to be excluded. Nevertheless, it sometimes affords most brilliant results. Thus, paralysis of the left vocal band, associated with recurring cough and occasional dyspnea and dysphagia, every other suggested cause for which has been excluded, has led to a correct diagnosis of aneurism of the arch of the aorta, not discoverable by the most careful physical exploration.

As a rule, that diagnosis that best and most readily accounts for all of the symptoms is the most likely to be correct. When any symptom is unaccounted for, the diagnosis is at least incomplete, if not doubtful. Diagnosis of a common disease, other things being equal, is more likely to be correct than diagnosis of a rare disease. On the other hand, statistics have no bearing upon the individual case; the rare disease may be present and be overlooked, from want of knowledge or in consequence of superficial examination.

More than one disease may be present in the same patient at the same time; but certain diseases are held to be antagonistic and thus not likely to coëxist. Data bearing on this point are defective and are undergoing revision and correction. Not a few cases have been recorded contradictory of some of the teachings of the past. Hence, in this work, little stress has been laid upon these antagonisms. On the other hand, there are many lesions and morbid processes that are frequently found associated. Such associations are specifically mentioned wherever it seemed likely that a knowledge of their existence would be of service.

One disease sometimes strongly predisposes to another. A knowledge of facts of this nature is often of assistance in diagnosis, as well as in preventive treatment.

A knowledge of the *complications* and *sequelæ* that ordinarily or exceptionally attend or follow certain diseases, independently of its bearing upon therapeutics, is an important equipment for the diagnostician. This can be acquired only by extensive

reading or by prolonged experience. In anticipation of the latter source of information, the former must be sedulously cultivated. In default of such knowledge, the mistake may be made of diagnosticating a single well-marked condition, as, for instance, pleuritis, as the whole of the disease; when, in reality, the condition may be but a comparatively unimportant complication in the course of typhoid fever or of an hepatic abscess. Similarly, a sequela to some acute disease, remote in time or obscure in symptoms, may be diagnosticated, prognosticated and treated as an independent affection, often to the grave detriment of the patient. Thus, a temporary loss of knee-jerk following unrecognized diphtheria has led to an incorrect diagnosis of locomotor ataxia. Sometimes the initial phenomena of disease escape observation (a chancre may be concealed, especially in the female, or scarlatina may occur without eruption or appreciable fever, thermometry not being resorted to), and when the later phenomena (such as syphilitic fever or scarlatinal dropsy) develop, the case may be misinterpreted, unless this possibility of masked beginnings be borne in mind.

The so-called *negative signs* of disease should not be underrated. The absence of headache may assist in the exclusion of brain-tumor; the absence of albumin and casts from the urine, after sufficiently careful and extended observation, may be of great importance in discriminating among the causes of a train of symptoms including vertigo, optic neuritis, vomiting and paroxysmal dyspnea; the absence of rose-rash and of splenic enlargement may determine the diagnosis between enteric influenza and typhoid fever. In recording cases it is necessary to note negative points in order to show that the inquiry or search has been made.

No less important is a knowledge of the morbid phenomena that may be caused by certain drugs—not merely by the narcotic agents, opium, belladonna, and the like, but by such poisons as mercury, arsenic and lead, which may produce symptoms closely resembling acute and chronic diseases of common occurrence or may give rise to nervous and other phenomena puzzling in the extreme. Similarly, common or unusual drug-effects, perhaps due to idiosyncrasy, occurring in the course of treat-

ment, may mask the symptoms of disease, or give rise to additional phenomena that, unless caution be exercised, may lead to error in diagnosis.

The *data* for diagnosis are obtained by *observation* and *inquiry*.

Inquiry may often have to be made of those about the patient, the latter being unable or incompetent to answer; and not rarely the same questions will have to be repeated in various ways and be controlled by questions requiring opposite answers, in order that the clinician be not deceived, intentionally or otherwise.

Observation can be made while the questioning is proceeding; sometimes the manner, method and form of a patient's answers to questions are in themselves part of the data acquired by observation. For example, in cases of aphasia, the patient is unable to find words in which to express himself, though he may understand the questions put and may even believe that in his monotonous repetition of certain words and phrases he has given an intelligent answer.

Observation includes (1) a more or less rapid survey of the *general condition* of the patient, and (2) a careful examination into the condition of *special structures*, the performance of *special functions* and the constitution, chemic and microscopic, of the *blood*, the *secretions* and the *excretions*, both as to normal and abnormal constituents.

The *whereabouts* of the patient may afford information as to the acuteness or severity of the attack. He may be attending to his work as usual, or he may be confined to his house, his room, his bed.

Position and movement are next noted. Thus, if the patient paces restlessly about, a condition of excitement, of physical or mental causation, is indicated. If, whether seated or recumbent, his position is easy and unconstrained, it indicates that, whatever the disease may be, the attack is, for the time being, mild and that there is no serious respiratory or circulatory trouble, or inflammation of an important part. If the patient lie passively upon his back, or huddled up in a heap, a prostrating sickness, usually an acute infectious disease, is indicated. If he lie persistently on one side, affection of the lung or pleura of that

side is likely to exist; because this position restrains the movement of that side of the thorax, lessens the pain of acute inflammation and, in addition, permits the healthy lung to better perform its work. Sometimes, however, patients with pneumonia lie on the sound side, having less pain in that posture.

If, whether the patient be in an easy-chair, or in bed, the thorax is propped into an upright or semi-upright posture, there is great difficulty in respiration (*orthopnea*), which may be be due to cardiac or pulmonary disease, to obstruction in the upper air-passages, or to abdominal or general effusions. If the patient lie upon his back with the legs drawn up, there is likely to be peritonitis, and if on the right side with the right leg drawn up, appendicitis should be suspected.

If the head is retracted, or fixed in opisthotonos, disease of the cerebrum or of the meninges is to be suspected. So, too, deviation of the head to one or the other side, picking at the bedclothes, helplessness of one or more extremities, general restlessness, jactitation, local spasms or general convulsions, incoördination, tremor, temporary or persistent, sometimes throw considerable light on the condition of the nervous system.

The *expression of the countenance* should be observed. It may be indicative of comparative comfort or of intense suffering; it may be dull and apathetic, as in typhus fever; sunken and anxious, as in cholera; eager and brilliant, as in some cases of pulmonary tuberculosis; indeed, a careful study of the physiognomy of patients may often directly reveal much, or indirectly serve to lead the attention of the examiner to matters that might otherwise be overlooked. Thus, there is in many cases of chronic nephritis a peculiar pallor that can hardly be mistaken by the experienced observer. Connected with the expression of the face are the *general nutrition*, the condition of the *skin*, and the condition of the *mind*, all of which must be included in the general examination.

Concerning the *mental condition*, without special effort it can be learned whether the patient is interested in his surroundings or is oblivious to them, whether he is conscious or unconscious, whether he is delirious or is aware of his sayings and doings, whether he is able to fix his attention or is continually wander-

ing. To the facts developed on special inquiry no attention is now being paid.

The *skin* may be of normal hue, pallid from impoverishment of blood, flushed with fever, discolored by inflammations, bruises, extravasations, eruptions, jaundice, the various cachexias. It may be swollen, or give evidence of swelling or effusion beneath. It may be abnormally smooth or rough, dry or moist, hot or cold.

The *state of the nutrition* is usually evident in the preservation of the full, rounded outlines, or in the sharpness and hollows of emaciation. For accuracy and completeness of knowledge the weight should be determined by the scales.

Abnormal prominences of various kinds and situation may indicate the existence of *tumors*, *effusions*, *articular*, *osseous*, *glandular* and *visceral enlargements* of inflammatory or other origin, to be determined by further examination.

Pulsations may be visible that, by their abnormal situation or abnormal character, are indicative of abnormal circulatory conditions, or of tumors or collections of fluid external to the heart or vessels. The tortuosity of visible arteries, or the localized or general turgescence of superficial veins, may indicate disease or obstruction in or affecting the circulatory apparatus.

The manner in which *breathing* is performed, the character of the *voice*, the nature of a *cough*, afford information as to the condition of the respiratory apparatus.

Examination of the *pulse*, the *tongue*, and the *temperature*, while affording general indications, yet mark the transition from general to special examinations.

Except in a few instances, the plan of this book does not contemplate descriptions of methods of special examination, or enumeration of the conditions indicated by special symptoms and signs (*semeiology*, *inductive diagnosis*), but rather an account of the observed clinical groupings of signs and symptoms in the recognized and named affections of nosology (*historical diagnosis*, *pathological association diagnosis*), and an elucidation of the points of contrast between diseases most likely to be confounded (*differential diagnosis*).

FEVER.

What is fever?

Fever is a complex morbid process, of which the most striking feature is elevation of temperature. Sometimes the term fever is restricted to the abnormal heat, but this is more correctly termed *pyrexia*. In addition to pyrexia, fever is attended with acceleration of the pulse and of the respiration, with thirst and with disturbance of digestion. There are likewise increased and perverted tissue-changes, as manifested by alterations of secretions and excretions, which may be completely arrested or diminished in quantity and changed in quality. As a result of the deprivation of the normal products of metabolism and of the retention in the blood of abnormal products of metabolism, as well as of the waste-products of normal metabolism, other symptoms are often caused, such as malaise, depression, headache, insomnia, delirium, etc. When fever is long-continued, wasting takes place; partly as a result of the abnormal heat, both directly and indirectly; partly from other causes. Many febrile processes are at some period in their course attended with subnormal temperature.

What is the average normal temperature of man?

The *average normal temperature* of the adult is 98.4° or 98.6° F. (37° C.); in health it may vary from this a little more than a degree; from 97.3° F. (36.25° C.) to 99.5° F. (37.5° C.). It is higher in children than in adults, and in the aged it is sometimes a few tenths higher or lower.

The temperature is increased after meals by the activity of digestion; it is increased by exertion, and in children and hysterical persons by emotion. Weather has but slight influence upon the temperature of the healthy; it often has a marked influence upon that of the sick. There are daily periodic variations. The *daily minimum* occurs in the early morning between two and five o'clock; the *daily maximum* is reached between five and eight o'clock in the evening. The *average difference* between maximum and minimum is about 1.5° F.; it may be much

more. In the tropics the average body-heat is said to be slightly higher, and the daily range greater than in temperate climates.

In sickness the diurnal variations (morning remission, evening exacerbation) become quite marked.

What is subfebrile temperature?

The term *subfebrile temperature* is not to be confounded with *subnormal* temperature. It is applied to a moderate morbid elevation of temperature not exceeding 100.4° F. (38° C.).

What is the ordinary range of febrile temperature?

A temperature of from 100.4° F. (38° C.) to 101.2° F. (38.4° C.) is considered an indication of *slight fever*.

A temperature of from 101.3° F. (38.5° C.) to 102.2° F. (39° C.) in the morning, and of 103° F. (39.5° C.) in the evening, is indicative of *moderate fever*.

A temperature of from 103° F. (39.5° C.) to 104° F. (40° C.) in the morning, and of 105° F. (40.5° C.) in the evening, indicates *high fever*.

What is hyperpyrexia?

Febrile temperature exhibiting a tendency to remain above 107° F. (41.7° C.) is called *hyperpyrexia*. This is ordinarily directly dangerous to life.

Very high temperatures have been observed in hysterical cases in which, apparently, deception has been guarded against; a case of injury to the spine, in which recovery took place, is reported to have exhibited on many occasions a temperature of 122° F. (50° C.).

What is meant by inverse temperature?

Ordinarily the course pursued by morbid temperature is like that of normal temperature in having its maximum towards evening, and its minimum in the early morning. Sometimes, however, the maximum may be much earlier or much later than usual, even at noon or at midnight. Sometimes there are great fluctuations during day and night. When a complete reversal takes place, so that the *morning* temperature exhibits the *maxi-*

mum, and the *evening* temperature the *minimum*, the condition is said to be one of "inverse temperature."

Inverse temperature not infrequently occurs in acute tuberculous affections. It is of rather rare occurrence in typhoid fever.

What is the significance of a sudden fall of temperature?

A sudden fall of temperature may be part of the usual phenomena of a disease, as in intermittent fever; or it may indicate termination by crisis, as in pneumonia; or when it is not part of the usual course, and is not brought about by obvious loss of blood (as by venesection or menstruation), by drugs or cold applications, or other extraneous influence, it may indicate the development of albuminuria, or the occurrence of internal hemorrhage.

What is subnormal temperature?

A temperature lower than 97.3° F. (36.25° C.) is considered *subnormal*.

The temperature may be subnormal in influenza and in acute yellow atrophy of the liver as part of the essential course of the disease.

Subnormal temperature occurs, usually associated with profuse perspiration, at periods of critical recovery, as in pneumonia. There is simultaneous decrease in the frequency of pulse and of respiration and improvement in the general condition and sensations of the patient. In many acute diseases that terminate by lysis, such as typhoid fever, the temperature falls below the normal with the setting in of convalescence.

A subnormal temperature develops abruptly in conditions of collapse. It is then associated with sudden failure of the heart, as indicated by a feeble, rapid and irregular pulse, a fluttering and feeble cardiac impulse, pallor, coldness of the skin, with or without sweating, faintness or syncope, and profound prostration and weakness. The pulse becomes feebler, less rapid, and may finally cease. If recovery take place, the temperature again rises; or, if the termination be fatal, the temperature may rise *post mortem*.

The temperature becomes subnormal in some cases of severe hemorrhage, and at times, temporarily or persistently, in chronic

wasting diseases, such as pulmonary tuberculosis and carcinoma, in chronic Bright's disease and in some diseases of the brain.

What is an essential fever?

An *idiopathic* or *essential fever* is one in which the pyrexia and its concomitants do not result secondarily from an anatomical lesion, but arise primarily from the action of a specific poison or a similar cause. Essential fevers may or may not present invariable lesions as a part of their phenomena.

What is a symptomatic fever?

A *symptomatic* or *deuteropathic fever* is one that arises secondarily as the result of irritation, intoxication or perverted function, caused by disease or injury of a special organ or tissue. The causative lesion is the essential element; the fever is but one of the symptoms.

What is a specific fever?

A *specific fever* is an essential fever due to a specific pathogenetic agent. Typhoid fever and yellow fever are examples of specific fevers.

What is a continued fever?

A *continued fever* is one in which the temperature pursues an uninterrupted course, without sudden variation from beginning to end. The diurnal range will not exceed 1.8° F. (1° C.). The first rise may be sudden or gradual; there may be steady maintenance of a maximum, or continuous increase; there may be gradual or sudden final defervescence; but there is not a decided fall and a renewed rise during the progress of the case.

What is a remittent fever?

A *remittent fever* is one the course of which is interrupted once or repeatedly by a marked decline in temperature, not reaching the normal, and followed by renewed exacerbation. The daily difference will exceed 1° C. (1.8° F.).

What is an intermittent fever?

An *intermittent fever* is one in the course of which a complete intermission takes place once or repeatedly; that is to say, in

the progress of the case the temperature becomes normal (or even subnormal), remains normal for an appreciable time, of greater or less duration, and subsequently rises to about its previous level. The maximum may be high.

Simple Continued Fever.

What is the clinical course of simple continued fever?

Simple continued fever, *ephemeral fever* or *febricula* is a non-specific fever, sometimes apparently idiopathic, but often symptomatic of digestive aberration or of some local irritation. It may be *caused* by error in diet, by fatigue, anxiety, shock, unusual mental or physical exertion, exposure to cold and damp, to a foul atmosphere, to the sun, or to other source of metabolic perversion giving rise to intoxication.

The disease *sets in* with a feeling of indisposition, which is followed by chilliness, fever, with heated skin, headache, thirst, anorexia and pain in the back and limbs. The *pulse* is accelerated. The *tongue* is coated. There is usually *constipation*, less frequently *diarrhea*. The *urine* is scanty and high-colored. The *pyrexia* is slight or moderate; the diurnal variations are small, the rise rapid, the fall gradual or by critical defervescence, with profuse sweating or watery diarrhea. The *duration* is commonly less than a week. It may be less than twenty-four hours. Recovery is invariable and convalescence is rapid.

Graver forms of the affection, induced by the graver causes, may be attended with considerable depression, and running a more protracted course, in duration from ten to fourteen days, may simulate typhoid fever. Especially is this resemblance marked in cases of septic intoxication.

Ardent Fever.

What is ardent fever?

Ardent fever is a non-specific, continued fever, occurring in hot countries, in which the ordinary symptoms of febricula are exaggerated. The *temperature* reaches or exceeds 103° F.; there is throbbing of the temporal arteries, with severe headache and

even delirium. The symptoms resemble those of an inflammatory fever, so that the utmost care must be observed to avoid error in diagnosis.

Catarrhal Fever—Influenza.

What is catarrhal fever?

Catarrhal fever, epidemic catarrh, influenza or *la grippe*, now commonly called "grip" in the United States, is a widespread contagious disease that occurs epidemically, endemically or pandemically, more rarely sporadically, and exhibits a protean capability of variation in its symptoms. It is dependent upon the activity of a small bacillus with rounded extremities that is found in the nasal and bronchial secretions. The period of *incubation* is from one to four days.

There are four prominent *varieties* of the disease: (1) the *catarrhal*, in which the predominant features are those of catarrhal inflammation of the respiratory tract; (2) the *thoracic*, in which, with or without the ordinary catarrhal manifestations, there are symptoms of profound involvement of the thoracic viscera; (3) the *abdominal* or *gastro-intestinal*, in which disturbances of the digestive tract are most marked; (4) the *nervous* or *cerebral*, in which neural phenomena attract attention, and in which, as in the abdominal variety, the clinical course of typhoid fever is often closely simulated. Much greater refinement in subdivision might be made, as the symptoms of two or more types are frequently associated in one case. Some epidemics are characterized by greater or even exclusive prevalence of one or two types; but in other epidemics, half a dozen patients in one household may exhibit six varieties of the disease.

What is the clinical course of influenza?

The *invasion* of influenza is commonly sudden, sometimes with a chill, usually with more or less irritation or inflammation of the mucous membranes of the nose and throat, often associated with catarrhal conjunctivitis and blepharitis (pink-eye). Epistaxis may occur at this time or later. There is always a feeling of

lassitude, passing into unaccountable depression, physical and mental, with soreness in the back and limbs, aching of the muscles, more or less stiffness of the neck, cutaneous hyperesthesia, headache, dyspnea and anorexia.

The *temperature* is decidedly irregular; it may rise high; it is often subnormal for more or less protracted periods, and is subject to extreme fluctuations. It may be periodically or irregularly remittent or intermittent. The *pulse* is not much accelerated; it ordinarily is weak, the heart sharing in the general depression. The *tongue* is coated; the *bowels* are often constipated; the *urine* is scanty and high-colored, or profuse and light-colored. The *skin* is hot and dry and often peculiarly sensitive to the touch. Sometimes there are irregular perspirations.

In the *catarrhal type*, sneezing, coryza, watering of the eyes, odynphagia, painful respiration and a dry, harassing, irritative cough are early and prominent symptoms. The difficulty and distress in breathing are out of all proportion to the phenomena elicited by auscultation and percussion. Later, the cough becomes freer, and associated with expectoration of a thick, glairy mucus, sometimes blood-streaked or discolored; the nasal discharge becomes thicker and even purulent and hemorrhagic. Involvement of the sinuses (frontal, ethmoid, maxillary) may cause excruciating pain. With recovery, profuse, watery diarrhea occurs. In some cases, symptoms of tonsillitis or of catarrhal pharyngitis are the earliest, those of laryngitis and of bronchitis following. Coryza may be a late manifestation. In some cases, there is a peculiar edematoid condition of the tonsil, uvula, and neighboring structures that has been termed "solid edema," as puncture gives exit not to serum, but to a sanious, stringy, lymphoid material. Suppurative otitis media, preceded by excruciating earache and usually affecting both ears in succession, may be the only manifestation in the upper air-tract. This otitis is quite common in children and usually runs through a household.

In the *thoracic type*, pleuritis, pericarditis, endocarditis, pneumonia—catarrhal, croupous, or hemorrhagic—or edematous or

hemorrhagic infiltration of the lung may cause grave symptoms or even death.

In the *abdominal* or *gastro-intestinal type*, vomiting and purging are common, and there may be great pain and tenderness in the epigastrium or over the entire abdomen. Catarrhal jaundice may occur.

In the *cerebral* or *nervous type*, the headache is intense; there may be insomnia, photophobia, tinnitus aurium, talkativeness or even mild delirium. There is decided hyperesthesia; there may be restlessness, tremor, muscular twitchings or jactitations.

In the *typhoid type*, the course of the disease may be protracted, and the temperature may remain continuously elevated. The prostration may be extreme; the mental phenomena may be characterized by depression. There may be epistaxis, pain in the splenic region, special tenderness in the right iliac fossa and considerable diarrhea. There may be anomalous eruptions —papular, herpetic or erythematous. The action of the heart may be exceedingly feeble. Heart-failure is always a threatening danger.

The *duration* of influenza is as variable as are the symptoms. It may be less than forty-eight hours; it may be several weeks. Commonly, the acute symptoms last from three or four days to a week, but the weakness and depression continue much longer. *Convalescence* may be tardy, prolonged subnormal temperature and annoying perspiration being notable features.

In addition to the *complications* mentioned, there may be hemorrhages from various organs, meningitis—cerebral and spinal—multiple neuritis, arthritis and nephritis. Among the *sequelæ* are tuberculosis, paralyses of various kinds, hemicrania and melancholia, and other psychoses. Of itself influenza is not often fatal or directly provocative of serious complications or sequelæ; but it aggravates existing lesions or morbid processes, reawakens latent disease or searches out the weak point in the organism and renders this liable to the action of exciting causes. Hence, the great variability in its clinical course, and hence, too, the high mortality it occasions among the previously sick and debilitated, among infants and the aged.

With what diseases may influenza be confounded?

The *diagnosis* of influenza may be extremely easy or extremely difficult. According to its type, it may be mistaken for simple catarrhal inflammation of the mucous membrane of the eye, ear, nose, throat, bronchi, stomach or intestines; for measles; for cerebro-spinal meningitis; for acute articular rheumatism; for dengue; for ordinary types of pleurisy or pneumonia; for malarial fever; for acute tuberculosis; for typhoid fever.

Upon what does the discrimination depend?

In times of prevalence of epidemic influenza, or of the analogous epizoöty, the knowledge of that fact will cause one to be on the lookout for the disease, and he may, perhaps, even call other affections by its name. Characteristics upon which stress should be placed are the sudden onset, the great depression, the cutaneous hyperesthesia, the lumbar and muscular pains and the excessive respiratory distress. These serve to distinguish it from indigestion, gastro-intestinal catarrh, bronchitis, coryza and "colds." In contradistinction from typhoid fever or typhus fever, the common occurrence of some form of catarrhal symptoms, the irregularity of the temperature, the shorter duration and the absence of the characteristic symptoms to be described, are additional discriminating points. From ordinary pneumonia influenza differs in evolution, course and temperature, in the marked depression, in the irregular distribution of the pulmonary lesions, and in the presence in the expectorated matters of distinctive bacilli. The differential diagnosis from the other diseases mentioned will be successively developed.

Typhoid Fever—Enteric Fever.

What is typhoid fever?

Typhoid or *enteric fever* is an acute, infectious disease, dependent upon a specific microörganism, and presenting inflammation, swelling, softening and ulceration of the intestinal lymphatic structures, enlargement and softening of the mesenteric glands, tumefaction of the spleen, and changes in the parenchyma of other organs. Rarely the intestines escape.

The disease runs a course of about twenty-four days, beginning gradually and terminating by lysis. It is most common in young adults and in the autumn.

The *typhoid-bacillus* (Fig. 1) is a plump, motile organism, with rounded extremities. It is about one-third as long as the diameter of a red blood-corpuscle and one-third as wide as it is long. It presents a characteristic growth upon potatoes and does not liquefy gelatin. It is best stained by means of an alkaline solu-

FIG. 1.

Bacilli of typhoid fever. (Von Jaksch.)

tion of methylene-blue or by carbol-fuchsin or anilin-water fuchsin. It differs from other organisms found in the stools in not generating indol. The bacillus has been found during life in the blood, in the urine, in the stools, in the sputum and in the sweat; and, after death, in the intestinal wall, in the mesenteric glands, in the spleen, in the bone-marrow, in the liver, in the bile, and in the midst of complicating lesions.

How is the disease acquired?

Typhoid fever is most commonly transmitted through polluted water, though direct contagion and conveyance through air or food may take place.

What is the clinical course of typhoid fever?

The *period of incubation* of enteric fever varies between eight and twenty-three days. The *onset* is usually insidious, occasion-

ally abrupt, with a chill. The *prodromal period* is characterized by headache, pains in the back, epistaxis, general malaise, disturbed sleep, loss of appetite and coated tongue. These symptoms become aggravated, while to them are added relaxation of the bowels, abdominal tenderness, especially in the right iliac fossa, intestinal gurgling and elevation of temperature.

The *temperature* pursues a characteristic course. (Fig. 2.) There is a period of gradual ascent (first week), a period of maintained height (second week and half of third week), and a period of gradual fall (last half of third week and fourth week). For from five to seven days the temperature rises two or three degrees each evening, to recede a degree or a degree and a half on the following morning, until it has reached a level of from 102.5° to 104° F. (in grave cases even higher), at which, with

FIG. 2.



Temperature-chart of a case of typhoid fever. (Wunderlich.)

slight evening remissions and morning exacerbations, it lingers for from ten to fourteen days, then to decline inversely pretty much as it rose, until, with the setting in of convalescence, it may fall below the normal. Rarely, the temperature does not rise above the normal.

The *pulse* is accelerated, but not, as a rule, in proportion to the rise of temperature. It may not exceed 90. It is peculiarly soft and rebounding, giving rise to an apparent duplication known as *dicrotism.* The *tongue* is coated in the middle, but red at the margins and tip. The coating is thick and, at first,

white. The red, uncoated portion at the tip occupies a characteristic wedge-shaped area. The lips, gums and teeth become covered with sordes. The patient often exhales a characteristic odor. The *pupils* are dilated. A limited flush colors the cheek.

Toward the end of the first week or early in the second, a varying number of small, slightly elevated, *rose-colored spots* that disappear upon pressure or upon stretching the skin may be observed upon the trunk—upon the abdomen or chest, anteriorly or posteriorly. These spots appear in successive crops, each of which lasts for several days. A diffuse, scarlatiniform erythema has been observed in some cases, and an eruption resembling that of measles or typhus in other cases.

The *urine* presents a characteristic reaction. If to one part of a (one-half per cent.) solution of sodium nitrite in distilled water (1 : 40) and forty parts of a saturated solution of sulphanilic acid in dilute hydrochloric acid (1 : 20) is added an equal bulk of urine and the whole is rendered alkaline with ammonia-water, a deep-red color is produced.

The *blood* also yields a distinctive reaction after the first week, causing agglutination and sedimentation of typhoid bacilli, with loss of motility, when added in dilution of, say, 1 to 20 to a bouillon-culture—Widal's reaction. The number of red blood-corpuscles and the percentage of hemoglobin undergo reduction, while the number of colorless corpuscles is but little changed. Generally the large mononuclear and transitional leukocytes are increased, while the polynuclear neutrophiles are diminished.

The palms of the hands and the soles of the feet sometimes present a peculiar yellowish discoloration.

In classically typical cases, there is *diarrhea;* but in almost an equal number, however, the bowels are constipated. The diarrheal stools present a characteristic appearance, being thin and yellowish, "ochrey," or like pea-soup; they have a peculiar, fetid odor; sometimes they contain blood, independently of a formal hemorrhage. Tympanites is common. The area of *splenic percussion-dulness* is *increased.* In the second week, if not earlier, *mental dulness* and listlessness are manifest. The patient pays little heed to his surroundings, but usually re-

sponds when spoken to. Deafness and visual disturbances are not uncommon, and there is frequently low delirium. In the third week, the first sound of the heart is observed to be feeble; *emaciation* is decided, and the *tongue* often becomes dry, fissured and coated with a heavy, brown fur. Intestinal hemorrhage may occur. Perforation of the bowel, with consecutive peritonitis, is among the dangers.

Usually, towards the end of the third or the beginning of the fourth week, progressive improvement is manifested, coincidently with the decline of temperature. The morning remissions exceed the evening rises. Commonly by the twenty-fourth day, but often much later, the temperature has fallen to the norm. Convalescence is slow.

Death may take place in or after the second week, from exhaustion, toxemia, fever or the accidents of the disease.

Other *complications* and *sequelæ* than those mentioned are bedsores, phlebitis, thrombosis, pericarditis, endocarditis, parotiditis, edema, inflammation or ulceration of the larynx, bronchitis, pleuritis, pneumonia, pulmonary tuberculosis, osteomyelitis, meningitis, peripheral neuritis, nephritis, orchitis, rupture of the spleen, cholecystitis, cholangitis, abscess of the liver, the formation of abscesses and gangrene.

Sometimes the symptoms are mild, constituting *ambulatory* or *walking typhoid fever*. At other times, the attack, while perhaps severe, terminates at the end of a week, or of two weeks, constituting *abortive typhoid fever*. A few days after convalescence has set in, the temperature may again rise, and the attack be repeated, though curtailed in duration, constituting a *relapse*. A relapse may likewise interrupt the declining course of the disease. Sometimes, when convalescence is apparently about to set in, the temperature reascends and remains elevated for an uncertain period, constituting a *recrudescence*.

What is the distinction between a relapse and a recrudescence in typhoid fever?

A typical *relapse* in typhoid fever includes a redevelopment of the entire group of morbid phenomena of the primary disease, as indicated by a characteristic temperature-curve, splenic enlargement and rose-rash, though the duration may be shortened.

A *recrudescence* refers only to a reappearance of fever. The temperature of recrudescence does not pursue a typical course; it may fall as suddenly as it rose. A recrudescence may depend upon some complication or some accidental source of irritation, *e. g.*, peritonitis, or constipation or the premature or injudicious taking of food. A relapse indicates renewed activity of the specific cause of the disease.

FIG. 3.



Temperature-chart of relapse in typhoid fever. Convalescence from the relapse interrupted by recrudescence. (From a case at the Philadelphia Hospital.)

How is perforation of the bowel in typhoid fever to be recognized?

Perforation of the bowel in the course of typhoid fever may or may not be preceded by intestinal hemorrhage. When it occurs, the patient experiences a sudden and intense pain, localized at one spot in the abdomen, but soon extending. There may also be rigors. Tympanites, if absent, develops; if present, it increases. Vomiting may occur. There is exquisite abdominal tenderness; the patient lies upon his back, with the legs drawn up; the *face* is pale, pinched and anxious; the *pulse* is small, hard and rapid; the *breathing* is shallow and thoracic. *Shock* and *collapse* are common. With the fall of temperature, the patient's mind may become clear. *Death* may take place in

a few hours; but more commonly the temperature again rises and the symptoms of *peritonitis* become predominant, death occurring in the course of a few days.

Fatal perforation may, however, occur without decided symptoms either of shock or of peritonitis; or there may be a sudden fall or a sudden rise of temperature, a sudden increase in the pulse-rate or in the intensity of the prostration, or there may be sudden vomiting; or there may be simply persistent and rebellious tympanites, with comparatively slight abdominal tenderness and, perhaps, marked depression in the general state of the patient. While perforation is usually fatal, recovery has occurred in well-authenticated instances.

How are typhoid fever and pyemia to be differentiated?

Pyemia may be attended with typhoid symptoms, diarrhea and cerebral manifestations. The temperature (Fig. 4), however,

FIG. 4.



Temperature-chart of a case of pyemia. (Wunderlich.)

pursues a different course from that of typhoid fever; it is irregular and presents wide variations in range, often declining below the normal. The morning temperature may be normal, that of noon several degrees above normal, that of evening lower than that of noon. There may be great and sudden changes from day to day. In addition, there may be recurrent chills and

sweats. The rose-rash, the diazo-reaction and the Widal reaction are wanting. The colorless blood-corpuscles, and especially the polynuclear leukocytes, are increased in number. The detection of a primary focus of suppuration and the results of metastasis point to the cause of the equivocal symptoms.

How are typhoid fever and a typhoid condition to be differentiated?

A condition of asthenia and low vitality developing in the course of certain febrile conditions, surgical affections, septicemia and pyemia, possibly attended with diarrhea, is to be distinguished from typhoid fever by the previous history: on the one hand, there are the evidences of some constitutional or local disease; on the other, epistaxis, headache, continuous fever of gradual invasion. The age of the patient may have weight: typhoid fever being infrequent after thirty-five and rare after fifty, while in the aged many diseases, especially pneumonia, commonly assume a typhoid type. A rose-rash, the diazo-reaction and the Widal reaction are all significant of typhoid fever.

How are typhoid fever and yellow fever to be differentiated?

Initial headache and pains in the loins attend both typhoid fever and yellow fever; but, in the latter, epistaxis and diarrhea are wanting, the onset is abrupt and the symptoms remit on the second or third day. The discoloration of the skin that gives its name to yellow fever is wanting in typhoid. Yellow fever is a disease of hot climates; typhoid fever, one rather of temperate climates. The characteristic facies of yellow fever is wanting in typhoid, while albuminuria is less common and the slowing of the pulse, in proportion to the temperature, is less marked in the latter than in the former.

How are typhoid fever and variola to be differentiated?

For several days it may be impossible to distinguish typhoid fever and variola from one another. Both present headache and pains in the back. Epistaxis and diarrhea may be wanting in typhoid fever. The onset of variola, however, is likely to be

abrupt; that of typhoid fever insidious and gradual. On the third or fourth day, there appears in variola a characteristic eruption; simultaneously, the temperature declines. The eruption of typhoid fever is unlike that of variola and rarely appears before the fifth day. The subsequent course of the two diseases is sufficiently diverse to remove any possibility of confusion. The presence or absence of vaccine protection and the existence of other cases of one or of the other disease may have some weight in the early diagnosis.

How may one avoid confounding pneumonia with typhoid fever?

While pneumonia may present symptoms of a typhoid character and a temperature-course not unlike that of typhoid fever, the respiratory frequency is out of all proportion to the pulse-rate, while careful physical examination will reveal the signs of pulmonary consolidation (dulness on percussion, bronchial breathing, crepitant râles, increased vocal fremitus and vocal resonance) and perhaps the friction-sound of an associated pleurisy. Pleurisy, however, is by no means rare early in the course of typhoid fever. The appearance of rusty sputum clinches the diagnosis of pneumonia. It must, however, be borne in mind that pneumonia may occur as a complication of typhoid fever. In such a case, the association must be recognized by the rose-rash, the splenic enlargement, the diarrhea and the protracted course of the disease.

How are typhoid fever and trichiniasis to be differentiated?

Trichiniasis may present many of the manifestations of typhoid fever, but the epistaxis, the severe headache, the enlargement of the spleen, the characteristic stools, the typical temperature-curve and the rose-spots are usually wanting. The knowledge that there is such a condition as trichiniasis, with inquiry as to the food taken and the detection of nodules in the painful muscles, ought to be sufficient to prevent mistake. The increased number of eosinophile leukocytes is a point strongly in favor of trichiniasis.

Typhus Fever.

What are the symptoms of typhus fever?

Typhus fever, also called *famine-fever*, *ship-fever* and *jail-fever*, is an acute, infectious disease of sudden onset, with a papular eruption and pronounced nervous symptoms, usually terminating by crisis at the close of the second week. It is highly contagious and develops in crowded and unwholesome places, among the poor and wretched, although it may be communicated to any. There are no distinctive lesions. The disease has a *period of incubation* of variable duration—from a few hours to two weeks, during which the patient is comparatively comfortable, or there may be a brief stage of preliminary depression.

With the *onset* of the disease there are general malaise, headache, perhaps a chill, pains in the back and a heavily coated tongue, perhaps with nausea. The *temperature* rises to between 104° and 106° F.; the *pulse* is frequent, at first full, but early becoming feeble. Stupor soon develops. The *bowels* are usually constipated. The *expression* is dull. The *conjunctivæ* are injected. The *pupils* are usually contracted. The *face* appears livid. A *musty odor* is manifest. The *body* may be covered by a diffuse, red rash. Between the fourth and the sixth day a *coarse, papular eruption* appears, usually on the trunk and extremities, exceptionally on the face. Intermingled with this are many petechial spots. For two or three days new papules appear, to recede gradually and disappear.

At the close of the first week or early in the second, low, muttering delirium, or coma-vigil, without great restlessness or with ceaseless tossing, muscular twitching and jactitation, appears; the mental depression is profound. In the *cerebral type* there is a wild, fighting delirium, with intolerance of light and illusions of sight and hearing. Excitement is soon succeeded by weakness and prostration, perhaps by fatal coma. In some cases the *respiration* is shallow, irregular and noisy, though no change in the lungs can be detected. The *heart-sounds* are feeble, though the beat may be excited. Often, there develops an endocardial murmur, due to the depraved state of the blood.

The *tongue* is brown and cracked, the teeth and gums covered with sordes. The *urine* is scanty, high-colored, deficient in chlorides and may contain albumin.

In cases that recover the temperature gradually subsides at the end of the second week, a decided decline taking place on the fourteenth or sixteenth day, accompanied, perhaps, with profuse perspiration, diarrhea or a copious discharge of urine. *Relapses* are rare. An attack confers subsequent *immunity.* Pulmonary *complications* are the most common. Others may be meningitis, phlebitis, gangrene, erysipelas, parotiditis, edema of the larynx. During the last stages, or after convalescence, acute tuberculosis may develop.

How are typhoid fever and typhus fever to be differentiated?

Typhus fever is contagious; typhoid is not. Typhus is the more likely to be epidemic. Prodromata are more common and of longer duration in typhoid. The onset is acute in typhus, insidious in typhoid. Typhus lasts about two weeks; typhoid not less than three. The eruption of typhoid consists of small rose-spots, usually confined to the abdomen and chest, and appearing in successive crops; that of typhus is coarse, macular and petechial and of more extensive distribution. The skin is usually moist in typhoid; it is dry in typhus. In typhus, the body exhales a characteristic musty odor. The bowels are often loose in typhoid fever; they are usually constipated in typhus. Nervous prostration is the more profound in typhus. The course of the temperature is different in each. Epistaxis is common in typhoid; uncommon in typhus. The Widal reaction and the diazo-reaction belong to typhoid and not to typhus fever. In typhoid fever, one finds on post-mortem examination intestinal ulceration and enlargement of the spleen and of the mesenteric glands; in typhus, no constant lesions; though the spleen is likely to be diffluent. Typhoid fever and typhus fever may, though rarely, coëxist in the same patient.

How are variola and typhus fever to be differentiated?

Both variola and typhus fever are in a high degree contagious. Should both be simultaneously epidemic, the diagnosis may be difficult during the first few days of the attack. The eruption

of variola, however, appears from twenty-four to thirty-six hours earlier than that of typhus. The former is usually situated on the face, as well as on the trunk, and passes successively through the stages of papule, vesicle and pustule, the pustules rupturing and leaving cicatrices. The eruptionof typhus fever rarely or never appears on the face ; it remains largely papular, though in part it becomes petechial. In typhus fever, the temperature becomes high at the onset and continues high ; in variola, the temperature declines with the appearance of the eruption. Vaccination commonly protects against variola ; it affords no protection from typhus fever. Finally, typhus is a disease of about two weeks' duration ; variola, of quite three.

Cerebro-spinal Fever—Epidemic Cerebro-spinal Meningitis.

What are the clinical features of cerebro-spinal fever?

Cerebro-spinal fever or *epidemic cerebro-spinal memingitis* is an acute infectious disease, dependent upon the activity of the meningococcus or diplococcus intracellularis meningitidis, which is present in the blood, in the exudate, and in fluid obtained by lumbar puncture. Epidemics occur most commonly in winter and spring. Overcrowding, filth and other unsanitary conditions are predisposing influences. The disease varies much in its clinical manifestations, but it is usually characterized by decided disturbances of the cerebro-spinal functions. In some cases cerebral, in others spinal symptoms predominate. Not infrequently, respiratory phenomena or blood-changes assume great prominence. A characteristic eruption is usually a marked feature.

The attack may develop gradually, but more often it *sets in* suddenly, with a rigor followed by fever ; malaise ; nausea ; great thirst and vomiting, often with a clean tongue and no gastric derangement ; vertigo ; excruciating headache, remitting, but never entirely ceasing, and attended with paroxysmal exacerbations ; rigidity of the head and neck, sometimes passing into opisthotonos ; muscular twitchings or convulsions ; dry-

ness of the skin, with hyperesthesia and paresthesia. There may also be photophobia and tinnitus aurium. Prostration soon becomes profound, though restlessness may continue. Delirium may set in and be followed by stupor and coma. The *expression* is anxious. The *pulse* is rapid and extremely irregular. The *temperature* fluctuates between wide limits. Hyperpyrexia is not rare. It may develop suddenly and persist until death. The temperature may continue to rise after death. A sudden fall of temperature may usher in collapse and death. A gradual fall of temperature precedes recovery. When the thigh is placed at a right angle with the trunk the leg can be only partially flexed. The action of the sphincters is often deranged, so that there may be incontinence of urine or feces, or retention of urine, or constipation. As a rule, retention is an early, incontinence a late, symptom. The *urine* is often albuminous and contains an excess of urates. The number of *colorless blood-corpuscles* is increased, particularly the polynuclear leukocytes.

Between the first and the third day, purpuric spots, or an erythematous eruption that quickly becomes petechial, may appear upon the trunk and extremities. The disease is sometimes called "spotted fever," from the character of this eruption. Between the third and the sixth day, herpetic vesicles may appear on the face about the lips.

In the further progress of the case, the *pupils*, at first contracted, become dilated; paralysis and anesthesia of irregular distribution appear; disturbances of sight and hearing, perhaps also blindness and deafness develop. The *respiration* may be profoundly disturbed. As death approaches there is difficulty in swallowing and the breathing may assume the Cheyne-Stokes type. Short remissions in the general severity or in individual symptoms may occur, to be followed by renewed exacerbations. The *duration* of the disease is variable. The *fastigium* is commonly reached on the sixth day. In protracted cases, profound emaciation occurs. *Death* may take place early or late, in coma, by exhaustion, or by apnea. If recovery ensues, *convalescence* is tardy, and sometimes protracted, while permanent loss of special senses is common. Pneumonia is a common, arthritis

a rare, *complication*. Palsies, headache and epileptiform convulsions may be additional *sequelæ*.

In addition to the ordinary type of epidemic cerebro-spinal meningitis, there may be *fulminant* cases (death occurring within twelve hours), *mild* or *abortive* cases and *protracted* or *typhoid* cases. *Sporadic* cases are rare.

Instances of contagion (direct transference from the sick to the well) and of portagion (conveyance by the person or belongings of those that have been in contact with the sick) of cerebro-spinal fever appear to have been authenticated, but are extremely uncommon.

How are cerebro-spinal fever and tetanus to be differentiated?

Cerebro-spinal fever appears in epidemics, while tetanus usually occurs sporadically, as a result of the infection of a wound by soil. Trismus, an early symptom of tetanus, is the less common in cerebro-spinal fever. Opisthotonos, general rigidity and spasm are more marked in tetanus than in cerebro-spinal fever. The presence of the meningococcus in the fluid obtained by lumbar puncture is distinctive of the latter. Recovery from tetanus is exceptional. Death is not the invariable termination of cerebro-spinal fever.

Tetanus is wanting in the palsies, the eruption, the leukocytosis, the derangement of intellection and sensation and the febrile symptoms of cerebro-spinal fever, though toward the fatal termination the temperature may rise inordinately high.

How are cerebro-spinal fever and typhus fever to be differentiated?

While both cerebro-spinal fever and typhus fever occur in epidemics, and both may be sudden in onset and attended with profound nervous phenomena and petechial eruption, cerebro-spinal fever has not the dusky, stupid facies of typhus, while the herpes of the face, the retraction of the head, the fixed spinal pain, the muscular rigidity and the heightened sensibility of cerebro-spinal fever are not observed in typhus; nor is typhus, as a rule, accompanied with the great impairment of special senses or followed by the paralytic sequelæ of cerebro-spinal fever. The general course of the two diseases, the fever and

the eruption may be discriminated on careful observation. The greatest difficulty occurs in cases of malignant cerebral typhus.

How are cerebro-spinal fever and torticollis to be differentiated?

The muscular contraction that gives rise to torticollis is usually unilateral and limited, while in cerebro-spinal fever the contraction is symmetrical and not confined to the muscles of the head and neck. The symptoms of an acute, febrile disease, with disturbances of the sensorium and paralytic concomitants and sequelæ, are not present in torticollis, but are characteristic of cerebro-spinal fever. Even mild cases of cerebro-spinal fever, lacking the characteristic febrile course and without eruption, will present severe headache.

What are the distinctions between cerebro-spinal fever and smallpox?

Headache, vertigo, nausea, vomiting, pain in the back and fever attend both cerebro-spinal fever and smallpox; but retraction of the head, muscular rigidity and paralysis, hyperesthesia and anesthesia are wanting in smallpox, and the peculiar temperature-record and the characteristic eruption of smallpox are not seen in cerebro-spinal fever.

What are the distinctions between cerebro-spinal fever and yellow fever?

Yellow fever is especially a disease of hot climates; when found elsewhere its importation may be traced. If cerebro-spinal fever display any susceptibility to climatic conditions, it is most common in temperate zones. Characteristic symptoms of motor and sensory derangement, observed in the course of cerebro-spinal fever, are wanting in yellow fever, which is a disease of brief duration, in contrast to cerebro-spinal fever, the duration of which may be protracted. Petechial and herpetic eruptions appear during the progress of cerebro-spinal fever, while yellow fever is characterized by a saffron-yellow color of the skin. The black vomit often seen in yellow fever is entirely wanting in cerebro-spinal fever. Although remissions in the intensity of special symptoms may occur in the course of cerebro-spinal fever, the characteristic "lull" of yellow fever is absent.

How are cerebro-spinal fever and typhoid fever to be differentiated?

In cerebro-spinal fever, the onset is usually abrupt; in typhoid it is insidious. In typhoid fever the temperature pursues a typical course; in cerebro-spinal fever there is no regularity. The eruption of cerebro-spinal fever is petechial or herpetic and appears early—before the fourth day; that of typhoid is roseolous and appears not earlier than the fifth or sixth day. Constipation is the rule in cerebro-spinal fever; diarrhea often attends typhoid. Nausea and vomiting occur in cerebro-spinal fever, but not usually in typhoid. The retraction of the head, the paresthesiæ and the paralyses of cerebro-spinal fever are all wanting in typhoid fever. The headache is more intense in cerebro-spinal than in typhoid fever; in the latter it disappears when delirium sets in; in the former, delirium and headache coëxist. Leukocytosis attends cerebro-spinal fever, not uncomplicated typhoid. Lumbar puncture may disclose the presence in the arachnoid fluid of the specific diplococci of the former. The knowledge of an epidemic assists in the diagnosis. The discovery ophthalmoscopically of recent papillitis (choked disc) or optic neuritis would be diagnostic of cerebro-spinal meningitis in contradistinction from uncomplicated typhoid fever. In rare cases, typhoid fever is complicated by meningitis.

Asiatic Cholera.

What are the symptoms of Asiatic cholera?

Asiatic cholera, also called *cholera infectiosa*, is an acute, infectious disease, having its home in tropical climates and occurring in epidemics. It is dependent upon a specific organism, which is found in the alvine discharges of the patient and is principally transmitted by means of milk and drinking-water.

The disease has a *period of incubation* of from half a day to three or four days. It may *set in* suddenly, with a chill, but more usually the attack proper is preceded by a moderate diarrhea, to which the name *cholerine* has been given. This con-

stitutes the *first* or *premonitory stage*. The course of the disease may be arrested at this stage.

In the *second stage* (*stage of spasm*, or *stage of serous diarrhea*), there occur violent cramps in the abdomen and legs, and the intestinal flux increases in severity. There is often obstinate

FIG. 5.

Comma-bacillus of cholera. (Vierordt.)

vomiting. The patient complains of thirst, is restless and anxious ; prostration is marked ; the pulse is weak and thready ; the skin is cold and shrunken ; the eyeballs are sunken. The *temperature*, taken in the rectum or with a thermometer carefully applied and allowed to remain at least ten minutes in the axilla, will be found to be elevated. The *stools* are almost liquid and colorless and contain large quantities of epithelium, constituting the so-called "rice-water" discharges, in which the *comma-bacillus* of Koch (Fig. 5) is to be found.

Soon, a *third* or *algid stage*, or *stage of collapse*, sets in. The *circulation* fails, and there is marked depression of the vital powers ; the *respiration* is shallow and accelerated ; the *skin* becomes as cold as marble ; the *breath* may be chilling ; the *voice* is lost. Suppression of urine often occurs. The *urine* that is secreted is albuminous and contains casts.

In this condition of collapse the patient may die, or he may enter upon a *fourth stage*, or *stage of reaction*, convalescence setting in or a low, typhoid condition developing, with fever and delirium and possibly with suppression of urine. This stage may terminate in death or in convalescence. Convalescence may be *complicated* by ulceration of the cornea and by parotiditis.

How are cholera nostras and cholera Asiatica to be differentiated?

Cholera nostras occurs sporadically; cholera Asiatica, endemically or epidemically, and, in Europe and America, by importation. Asiatic cholera is by far the graver affection; the stools present a characteristic "rice-water" appearance, and in them a specific bacillus is to be found. If an apparently similar bacillus be found in the stools of cholera nostras, its morphology and culture will prove it to be different.

How is arsenical poisoning to be distinguished from cholera?

Poisoning by arsenic occasions vomiting, cramps in the abdomen and legs, and diarrhea with stools of a "rice-water" character. Local evidences of the ingestion of arsenic may be present in the mouth; vomiting precedes diarrhea; the stools are bloody and do not contain the specific comma-bacilli.

Relapsing Fever.

What are the distinguishing features of relapsing fever?

Relapsing fever is a specific, infectious disease of intermittent type, dependent upon the presence in the blood of a specific

FIG. 6.



Spirochetæ of relapsing fever. (V. Jaksch.)

organism—the *spirocheta Obermeierii.* (Fig. 6.) Transmission is thought to take place through the intermediation of mosqui-

toes. The disease has commonly prevailed amid unfavorable hygienic conditions.

The growth and development of the parasite give rise to the periodic paroxysms that characterize the disease and give it its name. After a *period of incubation*, varying from several hours to two weeks, the disease *sets in* suddenly with a chill, followed by fever, with decided elevation of *temperature*, muscular pains, vertigo, headache, nausea and vomiting. The spleen is enlarged. Often there is jaundice.

In the course of from five to seven days the attack abates with the suddenness with which it set in, the temperature falling to the normal and profuse diaphoresis occurring.

For about a week the patient is free from symptoms. At the end of this time, the paroxysm is repeated, to again subside; and this may happen a number of times.

Convalescence is tardy and protracted. During the paroxysms, spirochetæ in large numbers may be found in the blood. They are not to be found during the intermissions.

How is relapsing fever to be diagnosticated from typhoid fever?

In typhoid fever, both onset and subsidence are gradual; in relapsing fever, they are sudden. The former is a continued fever; the latter intermittent and periodic. Relapsing fever does not present the rose-rash of typhoid fever. Jaundice, often present in relapsing fever, is rare in typhoid fever. Spirochetæ are not found in the blood in typhoid fever, while the Widal reaction and diazo-reaction are distinctive of this disease.

How does relapsing fever differ from typhus fever?

The course of relapsing fever is interrupted by intermissions, giving rise to a disease of periodic type; typhus is a continued fever.

The marked nervous symptoms, as well as the exanthem, of typhus are wanting in relapsing fever.

Spirochetæ are found in the blood only in relapsing fever.

Upon what does the differential diagnosis between relapsing fever and yellow fever depend?

Yellow fever is a disease of not more than three stages; there

may be more or less in relapsing fever. The primary acute stage, as well as the period of intermission or remission, is longer in relapsing fever than in yellow fever. Vomiting occurs late in yellow fever, early, if at all, in relapsing fever. Vertigo is a more marked symptom in relapsing than in yellow fever. The occurrence of one or the other in an epidemic might afford a clue in diagnosis. Spirochetæ are present in the blood in relapsing fever; they are not found in yellow fever.

Malarial Diseases.

What are the characteristics of malarial disease?

The group of diseases known as *malarial* is characterized by paroxysmal periodicity, enlargement of the spleen and liver, melanemia and the presence in the blood, free or within the red corpuscles, of parasites that exert a deleterious influence upon the red cells. (Fig. 7.) Tertian, quotidian and estivo-autumnal parasites have been described. The parasites of tertian and quartan fever are ameboid bodies that enter the red blood-corpuscles, at whose expense they develop. They are at first unpigmented, but in the course of development pigment-granules appear. The parasites complete their cycle of development and break up into spores at the end of forty-eight and seventy-two hours respectively. The occurrence of sporulation coincides with, or rather, precedes, the appearance of the paroxysm. The estivo-autumnal parasite appears to be much more irregular in development. Sometimes the parasites assume a crescentic shape. More than one generation of the same parasite, or more than a single variety of parasite, may be present, and give rise to corresponding variations in the type of fever. It is believed that the disease is conveyed through the agency of mosquitoes.

During the paroxysm, the urine is said to be irritant, the proportion of both water and solids being increased. It often contains albumin, and there is sometimes a complicating nephritis.

Malarial diseases prevail principally in the lowlands of warm climates, with marshy soils.

FIG. 7.



Organisms of malaria—intracorpuscular.

FIG. 8.



Organisms of malaria—extracorpuscular. (Macé.)

What are the varieties of malarial fever?

Malarial fevers are said to be *intermittent* when between two paroxysms there intervenes a period of freedom from symptoms, with a restoration of the temperature to the normal; *remittent*, when between two paroxysms the symptoms moderate and the temperature falls (but not to the normal level).

When the paroxysm is repeated daily, the fever is designated *quotidian;* if repeated on alternate days, *tertian;* if with an interval of two days, *quartan.* If two paroxysms occur daily, the fever is called a *duplicated quotidian.* There may be a

double tertian, in which occur daily paroxysms, of which only those of alternate days are alike; a *double quartan*, and other combinations. When the paroxysms succeed one another so closely that the cold stage of one begins before the sweating stage of its predecessor ends, the fever is called *subintrant*.

Morphologic and biologic differences among the organisms present in the various types of malarial fever have been recorded.

What are the features of a malarial paroxysm?

A typical malarial paroxysm consists of a *cold stage*, a *hot stage* and a *sweating stage*. The disease is therefore called "chills and fever." It is also known as *ague*.

What are the characteristics of each stage?

The *cold stage* or *chill* sets in with malaise, nausea, vertigo, shivering; as the *rigor* becomes more pronounced, the patient may be severely shaken; his teeth chatter; the *skin* is cold and rough; the *breathing* is shallow and hurried; the *pulse* is small and rapid; the *temperature* of internal parts, however, is febrile.

Gradually, the feeling of coldness subsides and gives way to a sense of warmth, the temperature of the rectum or of the mouth continuing to rise; the surface of the body becomes flushed and the eyes are brilliant. The patient has now subjective sensations of *fever*. After the lapse of a variable time, a more or less copious *perspiration* sets in, with a decline in the temperature and an amelioration or disappearance of the symptoms. The paroxysm is at an end.

What are the clinical features of malarial intermittent fever?

In *intermittent fever*, as the name indicates, there is in the interval between two paroxysms a complete intermission of the symptoms, the temperature becoming normal or subnormal.

The *cold stage* lasts from fifteen minutes to an hour, the *hot stage* and the *sweating stage*, respectively, a varying number of hours. The beginning of each successive paroxysm anticipates in time of the clock that of the preceding paroxysm. The spleen is enlarged; herpes is common.

Untreated, the paroxysms of intermittent fever lose their regularity.

The disease may gradually and spontaneously subside, or the

paroxysms may become remittent or pass into the malarial cachexia.

After apparent recovery, a tendency to a return of the paroxysms on the fifth, seventh, ninth or fourteenth day is sometimes observed.

Fig. 9.



Temperature-chart of quotidian intermittent fever. (Wunderlich.)

Fig. 10.



Temperature-chart of tertian intermittent fever. (Wunderlich.)

What are the clinical features of malarial remittent fever?

Remittent fever represents a more profound degree of intoxication than does intermittent fever.

The *chill* is usually severe and protracted. In addition, there is gastric irritability, perhaps vomiting, sometimes jaundice. The *temperature* attains a high degree. The *hot stage* may last for from six to eighteen hours and is followed by profuse *perspiration.*

In the *interval* between two paroxysms, the symptoms moderate and the temperature declines, but does not reach the normal.

After the occurrence of several paroxysms, the chill may be wanting; or there may be but one—the initial chill. Without medicinal intervention, the remissions may gradually become less decided and a typhoid condition may develop.

What are the symptoms of hemorrhagic malarial fever?

Hemorrhagic malarial fever is a grave form of malarial intoxication in which the height of the paroxysm is marked by headache, severe pain in the back, nausea, vomiting, decided jaundice and hemorrhages from various mucous surfaces, particularly from the kidneys.

What is pernicious malarial fever?

In certain localities in which the malarial organisms are exceedingly numerous or virulent, the attack manifests a *pernicious* tendency. The clinical picture depends upon the system that bears the brunt of the disease. There may thus be a *cerebral form*, characterized either by delirium and excitement or by coma and depression; or there may be a *thoracic form*, in which the respiration is accelerated, and there is an urgent sense of the need of air; or there may be a *gastro-intestinal variety*, attended with nausea, vomiting, jaundice and diarrhea; or there may be an *asthenic* or an *algid variety*, in which there is a condition of marked debility and a striking coldness of the surface and of the breath. Pernicious malarial fever usually manifests its character only after a preliminary paroxysm of apparently ordinary intermittent or remittent fever. Unless promptly and vigorously treated, it is likely to be fatal.

What are the symptoms of the malarial cachexia?

After protracted residence in a malarious district or following untreated or rebellious intermittent or remittent fever, there may occur irregular chilly sensations, with some tendency to periodicity, an occasional sense of feverishness, headache, mental torpor, drowsiness, a sallow complexion, constipation or diarrhea, impaired appetite, enlargement of the spleen and liver—a complex of symptoms that may not yield to medicinal treatment, but which improves on removal to a non-malarious climate.

What is meant by "Dumb Ague?"

There is a variety of irregular manifestations of malarial or paludal poisoning, sometimes acute, but more frequently subacute or chronic, which do not at any time present the classical picture of chill, fever and sweat. These attacks of "dumb

ague," "masked malaria," "larval paludism," as they are variously called, comprise chilly sensations, irregular fever, or flushes, or subjective sensations of heat, joint-pains and muscle-pains, headache and neuralgias of various kinds, cough, with or without bronchial or laryngeal signs, gastric and other visceral disturbances, sometimes taking the form of crises; in severe or protracted cases, anemia, anasarca and albuminuria, hematuria and hemoglobinuria have been observed. A common variety is the so-called "brow-ague," a form of frontal headache frequently associated with tenderness of the nerves at the supra-orbital and infra-orbital foramina and sometimes with an intensely painful sensation of pressure or constriction referred to the nasal bones; the manifestations usually exhibiting an irregular periodicity. Appearing in the morning and disappearing at night, or when aggravated by sunlight, it is called "sun-pain."

Enlargement of the spleen is sometimes demonstrable. A careful study of the phenomena in these cases will usually elicit some periodicity in their recurrence, or while the prominent symptoms, gastric or neuralgic, or whatever they may be, may not be periodic, careful temperature-observations will show an unsuspected periodic rise. Sometimes, if quinine be administered for a short time, and then withheld, a distinct periodicity in the symptoms will be developed upon withdrawal of the drug. The discovery of characteristic plasmodia in the blood would establish the diagnosis. The organisms are usually comparatively few in number and of the *crescentic* or *sickle-shaped* variety. In some patients there is a tendency to *annual recurrence* of critical manifestations.

What is meant by Ague-cake?

The enlargement of the spleen in a case of chronic malaria or of malarial cachexia is sometimes quite manifest to ordinary inspection. When not so readily manifest to sight, it may easily be detected by palpation. Its connection with malaria being well known, the enlarged spleen has in vulgar parlance acquired the name of *ague-cake.*

What are the distinctions between malarial fever and the fever of suppuration or of septic infection?

Suppuration and septic infection usually give rise to fever of

a remittent or intermittent type. When an obvious cause exists, the recognition of the nature of the fever is easy. When, however, the fever is induced by deep-seated suppuration, as when abscesses form or in case of pulmonary tuberculosis or of occlusion of the hepatic or of the common bile-duct, the connection may be obscure. The distinguishing features, however, are that the symptoms of suppurative or of septic fever are rarely of regular periodicity or typical in course; that they are attended with leukocytosis; that they are often uninfluenced by the administration of quinine, which acts specifically in the malarial diseases; and that the hematozoa characteristic of malarial fever are wanting.

With what conditions may pernicious malarial fever be confounded?

The *gastro-intestinal* type of the disease may simulate ordinary gastro-intestinal catarrh, but the one is a grave condition, while the other is not; the one is febrile the other afebrile; the one submits to treatment by quinine, while the other does not; in the one the blood contains characteristic hematozoa and blood-pigment, while in the other it does not; in the one the spleen is enlarged, in the other it is unchanged in size.

The pulmonary type may be confounded with pleurisy or with pneumonia, but the physical signs and many symptoms will clear up any doubt. In doubtful cases, examination of the blood and treatment with quinine will furnish irrefutable evidence.

The *algid* variety resembles cholera, but it does not occur in epidemics, the general symptoms of cholera are wanting, and treatment decides the result.

Jaundice attends the *hemorrhagic form;* hemorrhages take place from various mucous surfaces and blood is found in the urine. Hemorrhagic malarial fever differs from paroxysmal hemoglobinuria in not being dependent upon exposure to cold, while the urine contains red corpuscles and not merely hemoglobin. Quinine cures the one, but fails to influence the other.

The cerebral type is to be distinguished from those conditions that give rise to apoplexy and from profound intoxications of various kinds. The essential, distinguishing features are the occurrence of the symptoms in the course of an attack of

malarial fever, the absence of palsies and localizing symptoms, the tendency to recovery under treatment, the enlargement of the spleen, the presence of hematozoa in the blood.

How are typhoid fever and malarial fever to be differentiated?

Typical cases of intermittent and remittent fever are not likely to be confounded with typhoid fever, but if an intermittent or remittent has existed for some time, uninfluenced by medication, a typhoid condition develops, and the symptoms may occasion some doubt in diagnosis. Under such circumstances, the previous history must be considered.

The diarrhea, the rose-spots, the temperature-course of typhoid fever are all different from what is seen in malarial fevers.

The reaction of the urine to sulphanilic acid and sodium nitrite, described by Ehrlich, and a characteristic bacillus are not found in malaria. The detection of the plasmodia of malaria in the blood or the reaction of typhoid bacilli to the blood dissipates even the remotest doubt.

Malarial fever and typhoid fever may coëxist as so-called typho-malarial fever.

How does syphilitic fever differ from malarial fever?

When secondary syphilis is marked by fever, the elevation of temperature usually occurs at night and is associated with bone-pains, cutaneous eruption and other evidences of syphilis. Cerebral and meningeal syphilis may also give rise to febrile movement. In many cases, the discovery of the plasmodia in the blood and the results of treatment by quinine on the one hand, and the results of treatment by mercury and iodides on the other hand, must make the diagnosis.

Malta Fever.

What is Malta fever?

Malta fever, also known as *Mediterranean fever*, *rock-fever*, *Neapolitan fever*, *undulant fever*, is an acute infectious disease of warm climates dependent upon the activity of a specific micrococcus which has been isolated from the spleen.

What are the clinical features of Malta fever?

The disease prevails especially, though not exclusively, in countries bordering on the Mediterranean. It is most common in summer, and it may be epidemic. The period of *incubation* is from six to ten days. The disorder presents an irregular *temperature-curve*, characterized by undulations of from one to three weeks, with intermissions of two or more days. There are, beside, obstinate constipation, profuse perspiration, neuralgic pains, arthritic effusions, anemia and debility; there may be orchitis. The *duration* is uncertain, if not indefinite, without change of climate. A malignant type, of short duration, and an intermittent type, with daily pyrexia, are described. The *blood* is capable of arresting the motility of the micrococcus melitensis.

How are Malta fever and malarial fever to be differentiated?

Each has a distinctive temperature-curve; while the blood of the one arrests the motility of the micrococcus melitensis and that of the other contains distinctive plasmodia; quinine is without effect upon the one and of almost specific therapeutic value in the other.

How are Malta fever and typhoid fever to be differentiated?

Malta fever is unattended with the epistaxis, the diarrhea, the rose-spots of typhoid fever; the temperature of typhoid fever is of continued type, that of Malta fever peculiarly undulating; the blood of the one arrests the motility of the typhoid bacillus, that of the other the motility of the micrococcus melitensis; splenic puncture will in the one instance disclose the presence of the bacillus, in the other that of the micrococcus.

Bubonic Plague.

What are the clinical features of the bubonic plague?

The *plague* is an acute contagious disease of great virulence, characterized especially by lymphadenitis, and occurring in epidemics, especially in the far east. It is dependent upon the activity of a small, motile bacillus that gains entrance into the blood and the organs of the body. Insects and domestic animals may convey the infection. The disease is most common in the

hot season and is predisposed to by unfavorable hygienic conditions. The period of *incubation* is three or four days. The attack sets in with headache, pain in the back, stiffness in the extremities, a sense of anxiety and restlessness and depression of spirits. *Respiration* is accelerated and hemorrhage may take place, especially from the air-passages. After from twelve to thirty-six hours the *temperature* rises and the *pulse* becomes accelerated. Symptoms of collapse may appear and death ensue. In less grave cases the lymphatic glands, axillary, cervical, popliteal, become enlarged, and they may undergo suppuration or gangrene. Carbuncles, as well as petechiæ, may appear on various parts of the body. The *mortality* is high.

How do plague and typhus fever differ?

Typhus fever is wanting in the hemorrhages, the buboes, the carbuncles of plague, and its duration is ordinarily longer. The discovery of the specific bacillus in the blood and in pus from the buboes would leave no doubt as to the existence of plague.

How are plague and yellow fever to be differentiated?

Yellow fever is unattended with buboes and carbuncles, while it presents jaundice, black vomit, a peculiar facies and disproportion between temperature and pulse-rate. The detection of the bacillus pestis in the blood or the pus from the buboes, or the arrest of the motility of the bacillus icteroides by the blood, will remove any doubt in diagnosis.

Beriberi.

What is beriberi?

Beriberi, or *kakké*, is a disease of undetermined origin, occurring in warm climates and attended with symptoms of peripheral neuritis, together with anasarca.

What are the clinical features of beriberi?

The disease has occurred in epidemics and has been attributed to both infection and food-intoxication. It is *predisposed* to by unhygienic conditions. The *symptoms* include paresthesia, pain, anesthesia, weakness and wasting of varying degree and extent, with reaction of degeneration and loss of reflexes. There occur, besides, palpitation of the heart, shortness of breath, albuminuria, edema and effusion into serous cavities.

Yellow Fever.

What are the characteristics of yellow fever?

Yellow fever is a specific, epidemic disease of hot climates, occurring in a single paroxysm of three stages: the *first*, a *febrile stage*, lasting from thirty-six to forty-eight hours, which sets in with a chill, followed by fever, with capillary congestion, especially of the face and eyes, pains in the head, the back and the calves of the legs, restlessness and anxiety, irritability of the stomach, vomiting, thirst, constipation; the *second*, a *stage of remission* or *lull*, of less than six hours, in which the fever subsides and the skin assumes a deep-yellow or bronze hue; the *third*, a *stage of renewal*, in which the symptoms reappear, prostration becomes pronounced and hemorrhages take place from various mucous surfaces; the vomited matters present a characteristic *black* appearance. The urine usually contains albumin and often casts. Suppression of urine may occur. The flushed and somewhat swollen *face* and the injected *eyes* are considered characteristic. The *pulse* may not be accelerated even though the temperature be high.

The *mind* is usually clear almost up to the moment of death, but in some cases delirium and stupor develop. A slender, motile bacillus has been found in the blood and the tissues, and in a number of cases the blood of persons suffering from yellow fever has caused agglutination and arrested the motility of the specific microörganisms in culture. The period of *incubation* is three or four days.

Death may result from collapse or with convulsions and the symptoms of uremia. If recovery take place, *convalescence* is often gradual, and may occasionally be interrupted by *relapse*. Some cases are quite *mild*, recovery taking place at the end of the first stage. Even grave cases may be so mild in the first stage as to be unrecognized; the patient *walking* about, to be suddenly seized with prostration, quickly followed by *black vomit* and death.

An attack protects against subsequent infection.

What are the distinguishing features between yellow fever and malarial remittent fever with jaundice?

Yellow fever is epidemic; remittent fever, endemic. Yellow fever is a disease of a single paroxysm, not lasting more than a week; remittent fever is a disease of repeated paroxysms, of periodic recurrence, and lasts more than a week.

In yellow fever, the eyes become injected and watery, the expression anxious or fierce. In remittent fever, there is no especial change in the eyes or in the expression.

Prostration and muscular pains are decided in yellow fever and are not so prominent in malarial fever.

Delirium is common in bilious remittent fever, and the mind is always dull. Delirium is not common in yellow fever, and the mind is usually clear.

The pulse may become very slow in yellow fever; it is always quick in remittent fever.

Hemorrhages from mucous surfaces take place in yellow fever; not in ordinary remittent fever.

The urine of yellow fever contains albumin, and suppression may take place; the urine of remittent fever contains no albumin and suppression does not commonly occur. Bile-pigment gradually disappears from the urine of yellow fever and increases in the urine of bilious remittent fever.

An attack of yellow fever confers immunity from subsequent infection; one attack of remittent fever predisposes to other attacks. Yellow fever is commonly fatal, remittent fever rarely fatal.

The treatment of yellow fever is uncertain; remittent fever yields to quinine.

Plasmodia malariæ are not found in the blood in uncomplicated yellow fever; they are diagnostic of malarial fever. The blood in yellow fever causes, further, agglutination and arrests the motility of cultures of the bacillus icteroides.

How are hemorrhagic malarial fever and yellow fever to be differentiated?

Both hemorrhagic malarial fever and yellow fever occur in hot climates and are attended with jaundice, hematemesis and other hemorrhages.

Yellow fever, however, is epidemic; hemorrhagic malarial fever, endemic. The former consists of but a single paroxysm, of three stages, including a remission; the latter is marked by a series of paroxysms, each followed by a remission.

Black vomit is the more characteristic of yellow fever; hemorrhage from the kidneys, of hemorrhagic malarial fever. Albumin and casts are commonly found in the urine in yellow fever; not in malarial fever. An attack of yellow fever confers immunity from subsequent infection; an attack of malarial fever predisposes to the occurrence of other attacks. The detection of the plasmodia of malaria in the blood or of the specific reaction in yellow fever establishes the diagnosis.

How is yellow fever to be distinguished from acute yellow atrophy of the liver?

Yellow fever is epidemic; acute yellow atrophy is sporadic. In acute yellow atrophy, the area of hepatic dulness becomes rapidly and decidedly diminished; in yellow fever, there is either enlargement or no demonstrable change.

Yellow fever is, and acute yellow atrophy is not, attended with a distinct remission in the severity of the attack. Yellow fever is sometimes followed by recovery; acute yellow atrophy but rarely.

The injection of the eyes, the pains in the back and extremities, found in yellow fever, are wanting in acute yellow atrophy of the liver.

In acute yellow atrophy, leucin and tyrosin are found in the urine, and while cerebral symptoms are more pronounced than in yellow fever, the temperature never rises so high and may even be subnormal.

What are the clinical differences between yellow fever and dengue?

When yellow fever and dengue prevail synchronously, the differentiation may be exceedingly difficult. Dengue is, however, likely to be unattended with the peculiar facies of yellow fever, the albuminuria, the relative slowness of pulse, the hemorrhages and the jaundice; while the agglutinating and sedimenting influence of the blood on the bacillus icteroides removes any doubt.

Weil's Disease.

What are the symptoms of Weil's disease?

Weil's disease, also called *acute infective jaundice*, is an intermittent febrile affection, usually exhibiting two periods of activity separated by an uncertain interval; the first of a little more, the second of a little less, than a week's duration. The disease may *set in* abruptly with nausea and vomiting. The *temperature* at once rises to a considerable height, but falls decidedly on about the night of the fifth day; subsequently declining gradually until the normal level is reached. After an afebrile period of from twenty-four hours to a week, there is a return of fever lasting a few days or a week.

The *attack* is characterized by headache, vertigo, malaise, debility, somnolence, and, sometimes, nocturnal fever and restlessness, hyperesthesia, diarrhea, muscular pains and jaundice. The *pulse* is small and frequent and sometimes dicrotic. The *respiration* is accelerated. The areas of splenic and hepatic percussion-dulness are increased. The *urine* passed is diminished in quantity and contains bile-pigment, bile-acids, albumin and casts. Hemorrhages from various mucous surfaces may take place—epistaxis, hematemesis, hemoptysis and intestinal hemorrhage. Petechial spots may appear in the skin.

The disease has been observed most commonly in summer and in vigorous young men, butchers and soldiers seeming to display a peculiar proclivity. Similar manifestations have followed poisoned wounds. In fatal cases, degeneration of the liver and kidneys and spleen has been found. Parotiditis, pneumonia, iridocyclitis and motor weakness have been *sequelæ*. In cases that recover, the *convalescence* is protracted.

How are Weil's disease and yellow fever to be differentiated?

Weil's disease and yellow fever probably exhibit a closer resemblance in description than in actuality.

Weil's disease shows a special predisposition for young adults, especially butchers and soldiers; yellow fever occurs in epidemics and does not confine itself to any class of individuals.

Diarrhea is the rule in Weil's disease; constipation in yellow fever. The initial stage of yellow fever is of shorter duration; it is earlier attended with a remission; the remission is less complete, and both the remission and final stage are shorter than is the case in Weil's disease. Black vomit is not common in Weil's disease; the injection and excitement are less than in yellow fever. Weil's disease does not present the peculiar facies of yellow fever, nor the relative slowing of pulse.

Leprosy.

What is leprosy?

Leprosy is a chronic infectious disease dependent upon the activity of a specific bacillus resembling that of tubercle.

What are the symptoms of leprosy?

The disease occurs in two forms, the *tubercular* and the *anesthetic*. In the former, areas of cutaneous erythema appear, which subsequently become pigmented and finally the seat of nodules. The hair falls out and the mucous membranes may be involved. Anesthetic leprosy is attended with pains and hyperesthesia, followed by anesthesia and trophic changes. The nerves are at first tender, but subsequently thickened and nodular. Blisters may form and rupture, leaving ulcers. The digits may undergo contractures and necrosis.

The Exanthemata.

What are the exanthemata?

The term *exanthemata* or *eruptive fevers* is applied to a group of contagious, epidemic diseases, each depending upon a specific infection and having as prominent signs, fever and specifically characteristic eruptions on the skin and often on the visible mucous membranes. As exanthemata are commonly described scarlatina or scarlet fever; morbilli or measles; rubella, also called German measles, French measles or Rötheln; variola or smallpox and its modification, varioloid; vaccinia—the usual consequence of vaccination; varicella or chicken-pox. Diphtheria and erysipelas might also be included.

Morbilli—Measles.

Upon what does the diagnosis of morbilli depend?

Morbilli or *measles*, also called *rubeola*, is an acute, contagious disease, common in children, attended with catarrhal symptoms (coryza, rhinitis, pharyngitis, laryngitis, bronchitis, conjunctivitis), febrile elevation of temperature and a characteristic exanthem.

The *period of incubation* is from seven to fourteen days. The *onset* is somewhat abrupt, with a quick rise of temperature to from 102° to 104° F., more or less headache, restlessness, injection and watering of the eyes, sneezing, running from the nose, often swelling of the nose and lip, perhaps cough and slight sore-throat. *Digestion* is commonly disturbed, and the *urine* may be scanty. The *temperature* (Fig. 11) undergoes a considerable ele-

Fig. 11.



Temperature-chart of measles. (Strümpell.)

vation with the onset of the attack, declining on the second or third day, to rise again on the fourth, with the appearance of the eruption. This consists of coarse, pink papules, primarily dis-

crete, then becoming surrounded by a somewhat paler border and soon coalescing, to form slightly elevated patches arranged in crescentic form, with intervening healthy skin. The eruption first appears upon the face and neck, then upon the body. The fever declines and the eruption begins to disappear between the fifth and the seventh day. Branny desquamation follows as the attack comes to an end on the ninth day. The eruption may usually be detected in the throat a day or two in advance of its appearance on the skin. Small, round, bluish-white spots, surrounded by an area of congestion or upon a diffuse red background, may be seen on the buccal mucous membrane, especially opposite the lower molar teeth. Sometimes a sense of rubbing or crackling may be elicited on pressure upon the abdomen, from the presence of the eruption on the peritoneum.

An attack of measles commonly protects against subsequent infection; though second attacks are not rare and third attacks not remarkable. The disease is ordinarily mild and benign; sometimes, however, it is malignant and hemorrhagic (*black measles*). Lobular pneumonia and catarrhal otitis media are not uncommon *complications*. An attack of measles is likely to precipitate the development of tuberculosis in one predisposed and to accelerate the course of the disease when it already exists.

How does morbilli differ from typhus fever?

Typhus fever is of longer duration and decidedly more grave than measles, than which it is relatively less common in children. While the eruption of measles in some degree resembles that of typhus, it appears earlier and, as a rule, has no tendency to become petechial; it begins on the face, while in typhus the face escapes. The catarrhal symptoms of measles are wanting in typhus; the profound nervous depression of typhus is not seen in measles. Typhus in North America is rare and as a rule imported, most frequently by emigrant-ships.

Scarlatina—Scarlet Fever.

What are the symptoms of scarlatina?

Scarlatina is an acute, contagious disease, to which children

and young persons evince a special predisposition. An attack confers relative immunity from subsequent infection.

The *period of incubation* may be short. It varies from twenty-four hours to a week, rarely ten days. The *onset* is usually abrupt, perhaps attended with vomiting or convulsions. The *temperature* at once rises to a considerable height (104° or 105° F.) and the *pulse* attains a striking frequency. In the first twenty-four hours, or sometimes a little later, a diffuse, fine, punctiform, *red rash* appears, at first on the neck and breast and in the flexures of the joints, soon spreading as a uniform scarlet flush over the greater part of the body. Pressure causes temporary dissipation of the redness. There are intense subjective burning and itching of the skin.

The *throat* is usually sore and swallowing is painful. The fauces and palate are reddened, the tonsils and uvula and the adjacent cervical glands are enlarged and there is stiffness of the muscles of the neck. The scarlet discoloration may sometimes be detected in the throat, especially on the free border of the soft palate and on the uvula, in advance of its appearance on the skin. The throat-affection is often ulcerative or pseudo-membranous in character. The larynx is rarely invaded. The nose is usually involved and there is more or less serous or sero-purulent discharge. The inflammation may extend into the Eustachian tube and involve the auditory apparatus. Suppurative otitis is not an uncommon complication, and perforation and deafness may result.

The *tongue* is at first heavily coated, but, in the course of a little while, the dense fur is cast off, exposing the surface of the reddened organ, with its enlarged and prominent papillæ—an appearance characteristically described by the designation "strawberry tongue." Thirst is often great. *Digestion* is deranged. The *urine* is scanty. The number of *colorless blood-corpuscles* is increased.

Severe *nervous symptoms* may occur, twitchings or convulsions, restlessness, insomnia, delirium, stupor, fatal coma; or the nervous disturbances may be very mild and cease with the setting in of convalescence.

In favorable cases, the eruption fades by the fourth or fifth day.

The temperature, which has remained high, begins to decline; *defervescence* taking place by somewhat rapid *lysis* (Fig. 12). At the end of a week or nine days, the skin undergoes a furfuraceous or membranaceous *desquamation*, the temperature falls to the normal and convalescence may set in. Persistence of high temperature into the second week is not common and usually denotes a complication. So, too, a sudden recrudescence of pyrexia, after decline has begun, indicates suppuration or other accident. Convalescence may be interrupted by the appearance of symptoms of an *acute nephritis;* edema of the

FIG. 12.



Temperature-chart of scarlatina. (Strümpell.)

face and body, with diminished elimination of a dense, high-colored or smoky-looking urine, containing considerable albumin and blood, as well as blood-casts and epithelial casts of the uriniferous tubules. Not rarely, albuminuria and other evidences of renal congestion or inflammation may be detected prior to the appearance of edema. With or without suppression of urine, there may occur convulsions, delirium, stupor, coma and death.

Another rather frequent *complication* is arthritis, which may involve a single joint or several joints. Sometimes, with or without joint-symptoms, there is endocarditis, pericarditis or pleuritis. Permanent valvular lesions of the heart may be *sequelæ*.

Various *types* of scarlatina are observed in addition to the ordinary form. When throat-symptoms predominate, the attack is called "anginose." Sometimes the disease is *mild* or *abortive*. At other times it is *malignant* in virulence. In malignant cases, the rash may be delayed; it may be pale and indistinct or dark and livid. Cases in which the rash is wanting may be mild or severe. This form is called "larval" or "scarlatina sine exanthemate." In these cases, dropsy or suppression of urine may be the first symptom to attract attention. Anasarca may occur without nephritis, but its dissociated occurrence is not common. Scarlatina is said occasionally to be *hemorrhagic*.

How are scarlatina and measles to be differentiated?

Scarlatina usually sets in with vomiting or convulsions; measles rarely so begins. Rather severe sore-throat and glandular enlargement characterize scarlatina; catarrhal symptoms are present in measles. The great rapidity of pulse and elevation of temperature commonly observed in scarlatina are wanting in measles The eruption of scarlatina appears on the first or second day of the disease and is finely punctate, occasioning an appearance of diffuse redness; the eruption of measles appears not before the third day and is commonly papular, arranged crescentically, with areas of intervening healthy skin. After a preliminary elevation, the temperature of measles falls on the second or third day, to rise again with the appearance of the eruption, then to subside rapidly; the temperature of scarlatina at once mounts high and, after a few days, declines gradually. Nervous symptoms are decided in scarlatina, wanting in measles.

What are the features that differentiate acute exfoliative dermatitis and scarlet fever?

Acute exfoliative dermatitis may set in suddenly with febrile symptoms, and the exanthem persist for five or six days and be followed by desquamation. It is, however, unattended with the peculiar appearance of the tongue, the angina, the grave constitutional symptoms and complications and the leukocytosis of scarlet fever.

How is scarlet fever to be differentiated from toxic dermatitis?

Inflammation of the skin of varying intensity and extent

may arise from many toxic influences. The exanthem of scarlet fever may itself be considered to be of this nature, and numerous other infectious diseases and processes, as, for instance, rubella, variola, diphtheria, septicemia, typhoid fever, are capable of giving rise to a similar manifestation. Cutaneous erythema has been observed also after the use of medicaments, as, for instance, quinine, belladonna, atropin, bromids, and of antitoxin; and as a result of direct irritation of the skin, as, for instance, by rhus toxicodendron and turpentine. The diagnosis will be based upon the history, upon a recognition of the etiologic factors and upon the course of the respective affections. High fever, rapid pulse, profound constitutional disturbance, sore throat, albuminuria, anasarca, desquamation, when present, make up a clinical picture that is not readily mistaken.

Rubella.

What are the symptoms of rubella?

Rubella, also called *Rötheln*, *roseola*, *German measles* and *French measles*, is an acute, contagious exanthem, presenting an eruption resembling that of morbilli and throat-symptoms like those of scarlatina. An attack protects from subsequent attacks, but not from measles or scarlet fever; neither does an attack of scarlet fever or one of measles confer immunity from rubella. The *period of incubation* of rubella is from five to twenty-one days. The *onset* is usually abrupt. The *temperature* is moderate; the *pulse* is not very rapid. In the course of a day or two there appears, first upon the face and then progressively invading the trunk and the extremities, an *eruption* of small pinkish papules, usually separated from one another by skin of normal appearance. Sometimes the intervening skin is erythematous. The eruption lasts for from four days to a week, is attended with itching and is sometimes followed by slight desquamation. The *throat* is usually sore, the fauces reddened and the cervical glands enlarged. *Catarrhal symptoms* are common. The *course* of the disease is usually mild and uncomplicated.

How are rubella and scarlatina to be differentiated?

Rubella is inherently a mild disease; scarlatina is never without gravity. Rubella lacks the rapid pulse, the high temperature, the "strawberry tongue" and the grave complications of scarlatina. The eruption of rubella more nearly resembles that of morbilli than that of scarlatina. Slight edema of the hands, sometimes present in scarlatina, is not found in rubella. Neither rubella nor scarlatina is protective against the other.

How are rubella and morbilli to be differentiated?

Rubella resembles morbilli in its rash, in its mildness and in the catarrhal symptoms it presents. The eruption of Rötheln, however, is brighter-hued and does not show a tendency to crescentic arrangement, while it appears earlier and the individual papules are smaller than is the case in morbilli. Rubella is milder than morbilli, even as relates to the catarrhal symptoms. An attack of either confers no immunity from the other.

Variola—Smallpox.

Upon what does the diagnosis of smallpox depend?

Smallpox or *variola* is an acute, contagious, epidemic disease, *setting in* with a chill, pains in the back, head and extremities, nausea and vomiting, elevation of *temperature* to 102° or 103° F.,

Fig. 13.



Temperature-chart of smallpox. (Strümpell.)

with marked exacerbations, increased rapidity of pulse, and a

diffuse, *red, scarlatiniform rash* or a *macular measly rash* that is followed on the third or fourth day by the appearance of *papules;* when the temperature declines. The papules commonly appear first on the lips and forehead. The preceding red or macular rash usually appears first on the abdomen, the arms, and in the neighborhood of the groin. The fauces are usually reddened. There may be decided catarrhal symptoms. In the course of the succeeding four or five days, the papules become *vesicles* and the vesicles, in turn, *pustules.* The surface of the pustule is depressed at the middle—*umbilicated.* Each pustule is surrounded by an area of redness, constituting an *areola.* On the eighth or ninth day the pustules rupture and discharge their contents, and the *temperature again rises* (secondary or pyemic fever). (Fig. 13.) Secondary fever is sometimes announced by rigor, and its temperature-course is remittent. The evening temperature reaches from 103° to 105° F. The period of maturation and discharge lasts from three to five days, when crusts form and the temperature declines; in the course of a week the scabs fall off, leaving red cicatrices, which in the course of time become whiter and contracted, leaving "pits." The *spleen* is enlarged, and active *leukocytosis* is present.

Ulceration of the larynx and trachea, bronchitis, pneumonia, pleurisy or orchitis may *complicate* variola. Secondary inflammations are most likely to occur coincidently with the secondary fever. The eye frequently suffers permanent injury as the result of an attack of smallpox. The *period of incubation* of variola is about twelve days.

What are the varieties of smallpox?

Smallpox may be *simple* or *discrete, confluent* or *hemorrhagic;* it may assume a *malignant* character.

What are the characteristics of discrete smallpox?

Discrete smallpox is the mildest type of smallpox; the eruption is least profuse, the pustules occurring isolated. Sometimes the pustules are in contact at their periphery, when the disease is said to be *coherent.*

What are the characteristics of confluent smallpox?

In *confluent smallpox,* the pustules are numerous and run into

one another. The temperature fails to decline with the appearance of the eruption and is apt to be decidedly high during the period of maturation. The gravity of the case is greater than in discrete smallpox. Typhoid symptoms, delirium, stupor and fatal coma may develop; or death may be brought about by diarrhea, ulceration of the larynx or trachea, endocarditis or other complication.

What are the characteristics of hemorrhagic variola?

In *hemorrhagic variola* the exanthem may be from the first constituted of ecchymoses, while hemorrhages may take place from any of the mucous membranes. This is the gravest form. Few cases recover. Sometimes hemorrhage does not take place until the vesicular or pustular stage is reached.

What is malignant smallpox?

The epithet *malignant* is sometimes applied to a variety of smallpox, most frequently encountered at the beginning of an epidemic, in which death may occur early in stupor, following delirium, the eruption being ill-defined and perhaps only developed *post-mortem*.

What is varioloid?

Varioloid is smallpox modified by vaccination or by a previous attack of smallpox. The symptoms of the disease are similar to those of smallpox, but milder in degree and shorter in duration; secondary fever is absent. The eruption of varioloid commonly appears on the second or third day; that of variola on the fourth day. The course of varioloid is completed in about fourteen days; that of variola in about twenty-one days.

What is vaccination?

Vaccination consists in the introduction of the virus of cow-pock[1] into the lymphatic system of man, usually through the skin

[1] **What is meant by humanized virus?**

Humanized virus is vaccine-lymph that is not taken directly from the cow (or calf), but from the vaccine-pustule of a human being, usually a child, who may have been inoculated with matter obtained from another child, or from the cow. Unless one is sure as to the purity of the antecedents of the source of the humanized virus, bovine virus is to be preferred.

denuded of its epithelium, as a protection against variola. The protective influence continues for a period of about seven years, at the end of which time vaccination should be repeated.

What is vaccinia?

Vaccinia is the result of the inoculation of cowpock in human beings, and is protective against smallpox. For two or three days following vaccination, little is to be observed, locally or constitutionally. At the end of this time, the site of inoculation presents an appearance of slight redness, which in the next few days becomes intensified, as a vesicle forms and becomes transformed into a pustule, umbilicated and surrounded by a distinct areola. This process goes on for four or five days, when the pustule ruptures and the intensity of the inflammation gradually subsides. In the course of a week or ten days, the crust falls off and leaves a reddish cicatrix, which subsequently becomes white and depressed.

How does smallpox differ from measles?

The eruption of measles is coarsely papular throughout, with a tendency to crescentic arrangement, and is followed by branny desquamation; that of smallpox is irregular in arrangement and passes from the papular into a vesicular and then into a pustular stage, sometimes leaving disfiguring cicatrices. In measles, *after a previous decline in temperature, the appearance of the eruption is attended with renewed elevation;* in smallpox, *the temperature declines with the appearance of the rash* and reascends with the occurrence of pustulation. Smallpox is a grave disease, lasting three weeks; measles is a mild disease, lasting less than two weeks.

How are variola and scarlatina to be differentiated?

Smallpox may be attended with a primary, diffuse red rash, not unlike that of scarlatina, but, at the end of three or four days, papules appear, in turn to be succeeded by vesicles and pustules, finally leaving disfiguring cicatrices. With the appearance of the secondary rash, the temperature declines. The eruption of scarlatina undergoes no change and terminates in desquamation. The temperature is high from the outset and is continuous. Variola is not characterized by the same rapidity

of pulse or by the typical "strawberry tongue" of scarlatina. Uncomplicated scarlatina is a disease of less than two weeks' duration; variola rarely lasts less than three.

Varicella—Chickenpox.

What are the characteristics of varicella?

Varicella or *chickenpox* is a mild, contagious disease of childhood, attended with moderate elevation of temperature and the appearance on the first or second day of an eruption of papules, which in turn become transformed into vesicles. These may occasionally form bullæ, or undergo suppuration or gangrene. The *eruption* appears on the trunk and extremities, on the scalp and face. It comes out in crops and continues for three or four days, the vesicles desiccating and falling off, occasionally leaving cicatrices. The period of *incubation* is from two to three weeks.

How does varicella differ from smallpox?

Varicella is a mild disease; variola a grave disease. The eruption of smallpox appears on the third or fourth day and passes through papular and vesicular stages to become pustular; that of varicella appears on the first or second day and does not pass beyond a vesicular stage. The cutaneous lesion of variola has, further, a peculiar, shot-like hardness; it is surrounded by an inflammatory areola and the pustules are umbilicated. The appearance of the eruption in smallpox is attended with a fall in the temperature; in varicella, the temperature, rarely high, is uninfluenced by the appearance of the rash. Varicella is a disease of scarcely a week; smallpox is a disease of three weeks. Neither is protective against the other.

How are varicella and varioloid to be differentiated?

There may be a close similarity between the manifestations of varicella and those of varioloid. Varicella is rare in adults because of the immunity conferred by an attack in childhood. Varioloid does not respect age; it is likely to appear in the course of an epidemic of variola in those that have been protected by vaccination. Varicella is the milder affection, and is of the

shorter duration. Varioloid is but an attenuation and abbreviation of variola. Both varicella and varioloid are contagious. The existence of parallel cases may be decisive in diagnosis. An attack of the one does not protect against invasion by the other.

How are varicella and morbilli to be differentiated?

The eruption of measles is coarsely papular; it appears on the third or fourth day and displays a special proclivity to invade the face, with a tendency to crescentic arrangement. The eruption of chickenpox is at first papular and subsequently vesicular; it appears on the first or second day and is rather less than more abundant on the face than on the trunk. The catarrhal symptoms of measles are wanting in varicella. The temperature-course is not characteristic in varicella, as it is in measles.

Erysipelas.

What are the symptoms of erysipelas?

Erysipelas is an acute contagious disease, depending upon the activity of a streptococcus, and occurring most commonly in the spring. The period of *incubation* is from three to seven days. The disease usually *sets in suddenly* with a *chill*, sometimes with nausea and vomiting, or with convulsions, followed by considerable *elevation of temperature*, and the appearance, usually at some part of the face, and in most instances near one or other ear, or on the bridge of the nose, of an area of *red, brawny induration*. The redness progressively increases in extent, is *definitely circumscribed* by an elevated line of demarcation, and is attended with swelling.

The *appetite* is impaired. The *tongue* is coated.

The *urine* is usually albuminous, and often contains tube-casts, sometimes leucin and tyrosin. The number of *colorless blood-corpuscles* is increased.

There is often some soreness of the throat, and in some cases the disease may extend through the nasal passages into the pharynx and larynx, causing grave complications, and, perhaps, fatal edema. Sometimes the disease begins in the throat.

Involvement of a cerebral sinus may occur, giving rise to

severe symptoms and eventually causing death. Suppuration is not uncommon.

Usually, in the course of a week or ten days, the redness and swelling subside, desquamation sets in, and the temperature gradually reaches the normal.

In some cases the disease displays a *migratory* tendency (*erysipelas migrans*), continuing for a long period of time, and in turn appearing at various parts of the body. In children, erysipelas sometimes appears first in the neighborhood of the *anus*. The disease often attacks *wounds*. Recurrence is not uncommon.

What are the differential features between erysipelas and scarlatina?

Scarlatina is especially a disease of childhood. Erysipelas is more common in adults than in children.

The temperature is high in both, but the rapidity of the pulse is the more characteristic of scarlatina.

The rash of erysipelas is circumscribed in extent, limited in area, homogeneous in color; it usually appears upon the face, and presents a peculiar brawny induration, with well-defined borders; while the eruption of scarlatina is extensive in distribution and punctate in character.

An attack of scarlatina confers immunity from subsequent attacks; one attack of erysipelas rather predisposes to the occurrence of subsequent attacks.

Of the two, scarlet fever is the longer in duration.

How does erysipelas differ from simple erythema?

Simple erythema presents a diffuse redness, of transitory character, without febrile concomitants or sequelæ.

Erysipelas is a serious affection, with considerable elevation of temperature and characteristic rapidity of pulse. It lasts a week or more, is followed by desquamation, and is intimately related with nephritis.

How are facial erysipelas and herpes zoster of the forehead and face to be differentiated?

Facial erysipelas and herpes zoster of the forehead and face present a number of symptoms in common. That which dis-

tinguishes the latter, however, is that the eruption begins as a number of vesicles and does not extend beyond the middle line. The pain is much greater in herpes than in erysipelas. The constitutional symptoms are more profound in erysipelas than in herpes.

How are variola and erysipelas to be differentiated?

When variola is attended with a primary roseola, the disease may, for several days, simulate erysipelas. The redness of erysipelas, however, is distinctly circumscribed, although it may slowly spread, and is attended with brawny induration, while that of smallpox rapidly spreads from the face to the trunk and extremities. On the third or fourth day, if the disease is variola, papules appear, progressively passing through the stages of vesicles and pustules. The eruption of erysipelas undergoes little change, unless large blebs form, until on about the fifth or seventh day, when it may subside with desquamation.

Glandular Fever.

What is glandular fever?

Glandular fever is probably an infectious disease, occurring especially in children, and attended with redness of the throat and enlargement of the cervical lymph-glands.

What is the symptomatology of glandular fever?

The disease *sets in* suddenly, with pain on movement of the head and neck. The temperature is slightly elevated, and there may be nausea, vomiting and abdominal pain. Slight redness of the throat is visible and sometimes also the axillary and the cervical, the inguinal and the mesenteric lymphatic glands become enlarged and somewhat tender. In many cases liver and spleen also are enlarged. The adenitis may persist for several weeks. Suppuration may take place.

Miliaria—Miliary Fever—Sweating Disease.

What is miliary fever?

Miliaria is probably an infectious disease characterized by a

vesicular eruption and excessive sweating. There may be, besides, gastric disturbance and general malaise, and in severe cases high fever, delirium, prostration and hemorrhage. The *eruption* appears on about the fourth day, and is attended with itching and an aggravation of existing symptoms.

Dengue.

What are the symptoms of dengue or "break-bone fever?"

Dengue is an exanthematous and arthritic disease of hot climates, occurring in *epidemics*, and having a *period of incubation* of about four days. It is characterized by a peculiar intermittent temperature-course and by severe pains in the muscles and joints, which latter may be stiff and swollen. The knees are especially prone to be affected, so that the gait presents a peculiar character. A micrococcus has been described as present in the blood. The disease may be gradual in *onset*, with anorexia, headache, vertigo, drowsiness, or it may set in suddenly with a chill, moderate elevation of temperature and the appearance of an erythematous rash. There are stiffness of the neck and pain along the spine and in the lumbar region. The *fever* reaches its acme within twenty-four hours. The *temperature* fluctuates, rising and falling, but in the course of two or three days subsides nearly or quite to the normal; the eruption disappears, though constitutional depression and more or less pain continue. As a rule, nausea and vomiting do not occur at this stage, though the tongue may be heavily coated, and there may be other symptoms of gastric irritability.

After an intermittence of from forty-eight to seventy-two hours or more, the fever returns and a new eruption appears, as a rule resembling the eruption of scarlatina. Sometimes the eruption is more like that of measles, or it may be urticarious or vesicular. It is attended with heat and itching. Nausea and vomiting are usually manifested with this renewal of the fever.

In the course of a few days, desquamation occurs; convalescence sets in, but is tardy and protracted; there is weakness and more or less rheumatoid muscular pain; lymphatic swellings in the neck, groin or axilla often appear during the

febrile period or during convalescence. When the disease invades a community, few escape.

How does dengue differ from scarlatina with arthritic manifestations?

In dengue the fever is not continuous and pursues a course different from that of scarlatina; the arthritic symptoms are of earlier occurrence, and the pain is of a characteristic nature, giving the name "break-bone" fever. The eruption develops much later, and is sometimes quite different in appearance from that of scarlatina. The erythematous rash of the period of invasion is slight, inconstant and disappears without desquamation when the remission or intermission occurs. Throat-symptoms are not common.

In what respect does dengue differ from influenza?

Influenza is the graver disease; its symptoms are the more intense.

Catarrhal symptoms are frequent in influenza, infrequent in dengue.

Eruptions are exceptional in influenza, the rule in dengue.

Joint-pains are more decided in dengue. Hyperesthesia of the cutaneous surface is more common in influenza.

The course of dengue is interrupted by a remission. The course of influenza is usually though not invariably continuous to its termination.

Influenza is independent of climate. Dengue prevails only in certain localities. Convalescence is even more tedious and protracted from dengue than from influenza.

Diphtheria.

What are the symptoms of diphtheria?

Diphtheria is an acute, contagious disease affecting children especially, but adults as well, and characterized by superficial coagulation-necrosis of mucous membranes, especially of those of the pharynx, larynx and nares, with symptoms of constitutional intoxication. It is dependent upon the activity of the Klebs-Löffler bacillus, which is found in the false membrane

and at the site of the disease, and which varies considerably in virulence. The conjunctiva, the ear and the external integument also may be attacked. The period of *incubation* is from two to seven days.

The attack may *begin* insidiously, or *set in* suddenly with a chill, followed by considerable fever.

Pain in the throat in swallowing may or may not be complained of. The fauces will be seen to be livid, the tonsils usually swollen.

Soon, there appear, in greater or less extent and rapidly spreading over the tonsils, the half-arches, the uvula and the posterior wall of the pharynx, grayish or yellowish patches of false membrane, the forcible detachment of which is followed by bleeding. There are enlargement of the submaxillary and cervical glands and tumefaction of the soft tissues of the neck, externally. The number of *colorless blood-corpuscles* is increased.

The *constitutional disturbance* becomes profound. Albuminuria is common. From the pharynx the inflammation and necrosis may extend to the larynx and to the nose, and to the difficulty of swallowing are added croupy cough, aphonia, difficulty of breathing and a nasal discharge. The diphtheritic process is sometimes primary in the nose, and may thus escape detection, unless careful examination be made. From the larynx the false membrane may invade the trachea and bronchi. Pneumonia may occur. The action of the *heart* becomes weak and often intermittent. Septicemia, heart-failure and suffocation from obstruction of the larynx or bronchi are the common causes of death.

The *fatality* of the disease varies in different epidemics. Even in times of grave epidemics, there are many mild cases that become foci of infection. *Chronic diphtheria* of the throat is not so rare as it is commonly considered to be and is likewise a focus of infection. Paralysis from peripheral neuritis or cerebral thrombosis may be a *sequel*.

The *palsy* of diphtheria may appear in the course of the disease, but it is more common after the acute attack is at an end. Adults are rather more prone to suffer than children. The most common manifestations are *paralysis of the palate*, permit-

ting regurgitation of fluids and giving rise to nasal speech; *paralysis of the ciliary muscle* (cycloplegia), causing loss of power of accommodation; loss of *knee-jerks*. There may be more general palsy, with deranged sensation, ataxia and trophic changes.

How are diphtheria and scarlatina to be differentiated?

At the onset, the diseases may be indistinguishable. Both present the evidences of constitutional disturbance, with localized throat-symptoms. Possibly, the pulse may be relatively more rapid in scarlatina than in diphtheria. In from twenty-four to thirty-six hours, however, the appearance of a scarlet rash, as well as the subsequent course of the disease, dispels all doubt. In diphtheria, the symptoms centralize themselves about the throat; in scarlatina the throat-disturbance represents but a part of the general derangement. The detection of diphtheria-bacilli in the false membrane removes all doubt. The paralyses commonly seen after diphtheria are rare after scarlatina. More common during the course of scarlet fever or subsequently are suppurative ear-disease, nephritis and glandular enlargement.

Diphtheria and scarlatina may coëxist in the same patient.

How is tonsillitis to be distinguished from diphtheria?

Deposits on the tonsil may appear diphtheritic. They show little or no tendency to spread, however. Extension is characteristic of the diphtheritic membrane. In lacunal tonsillitis, the discreteness of the plugs and their situation at the orifices of the ducts are characteristic, and their creamy color is different from that of the diphtheritic pseudomembrane. Microscopically, they will be seen to be made up of desquamated epithelium, of sebaceous material and of ordinary fungi. The diphtheric membrane is constituted of meshes of fibrin including necrotic tissue, and it contains diphtheria-bacilli. In herpetic tonsillitis the eruption first appears as papules that soon become vesicles; but it is rarely seen at this stage; when ulcers and fibrinous deposits form and become confluent, the discrimination is difficult. Still, the herpetic patch is quite superficial, and more readily detached, leaving less erosion and causing less

hemorrhage in its separation than does the diphtheric patch. The former is usually the less extensive, and here and there, perhaps, the circular form of an isolated ulcer may give evidence of its origin. If necessary, inoculation-experiments and bacteriologic investigation will also help to discriminate. The constitutional symptoms of tonsillitis are less profound than those of diphtheria; local subjective symptoms, such as soreness, odynphagia and burning, are usually the more intense in tonsillitis, which is not, as a rule, followed by paralysis of the palate. Albuminura is not usual.

How is membranous croup to be distinguished from diphtheria?

Until the physician acquires sufficient experience to warrant a personal opinion, he had best consider all cases of membranous croup diphtheritic. The discrimination is difficult and disputed.

How are diphtheria and stomatitis to be differentiated?

The deposits in stomatitis are seated upon the mucous surface of the lips and cheeks and upon the tongue, while the membrane of diphtheria is usually seated in the pharynx, from which, as a center, it is distributed. The constitutional derangement is not as profound in stomatitis as in diphtheria. Stomatitis readily yields to mild general and local measures, while diphtheria is more rebellious to treatment. The fatality and the severe sequelæ of diphtheria are wanting in stomatitis.

Glanders—Farcy—Equinia.

What are the clinical features of glanders?

Glanders, *farcy* or *equinia* is an infectious disease, especially peculiar to horses, asses, and mules, from which it is transmitted to man through abrasions of the skin and through the mucous surfaces of those that come in contact with the diseased animals.

The site of inoculation displays evidence of active inflammation; especially is this marked in the nasal passages. There are also malaise, headache, elevation of temperature and pains in

the limbs; the urine may be albuminous. Soon there appears a macular eruption, which becomes vesicular, then pustular and finally umbilicated. The pustules may rupture and leave ugly ulcers. It is dependent upon the activity of a short, non-motile bacillus. The disease may appear in an acute or a chronic form. The period of *incubation* is three or four days. In addition, nodules form beneath the skin; these also soften and may rupture, discharging sanious pus and detritus. Lymphatic glands are enlarged. Another characteristic symptom of glanders is ozena. There is at first a moderate, thin discharge from the nostrils, soon, however, becoming profuse and purulent. The mucous membrane of the nares and contiguous structures is involved in intense inflammation and may become ulcerated. Catarrhal pneumonia and purulent arthritis are occasional complications. Glanders may be transmitted from man to the lower animals by inoculation.

How are glanders and variola to be differentiated?

In glanders, there may be a history with the local evidences of inoculation with the specific virus of the disease. The eruption of variola does not appear until the third or fourth day; with its appearance the temperature falls. The eruption of glanders may appear within the first twenty-four or forty-eight hours of the disease; it reaches a pustular stage much earlier than that of variola; there is no decline of temperature with its appearance. The ozena and the subcutaneous nodules of glanders are wanting in variola.

How are the ozena of glanders and that of syphilitic disease to be differentiated?

Ozena is dependent upon destruction of the nasal structures and putrefactive decomposition of the secretions. It is thus not distinctive of a single disease. Occurring in syphilis, it is a late manifestation, and probably has been preceded by well-defined symptoms. As seen in glanders, it occurs at the height of the disease, and is associated with a pustular eruption and the presence of nodules beneath the skin. The mode of infection differs in the two diseases.

Anthrax—Wool-sorters' Disease.

What are the clinical features of anthrax?

Anthrax, *wool-sorters'* or *brush-makers' disease*, *charbon*, *malignant pustule* or *splenic fever* is an infectious disease, due to inoculation with the bacillus anthracis. (Fig. 14.) It develops in butchers, wool-sorters, workers in hides, stevedores, and others that, with cut or wounded or abraded hands, manipulate the wool, hair or skins of animals that have died of splenic fever or charbon. The infection sometimes gains entrance through a scratch on the cheek or an abrasion of the lips. In butchers, the tongue is sometimes infected from a knife taken between the teeth. In those that carry hides upon their shoulders, the neck may be the site of local infection; these cases are likely to be more than ordinarily dangerous. The disease may apparently result from eating the flesh or from drinking the milk of infected animals.

FIG. 14.



Anthrax bacilli in blood. (Vierordt.)

At the site of inoculation, a pimple appears; the skin in its neighborhood becomes red and infiltrated; the papule becomes vesicular and pustular, with subsequent gangrene; other vesicles or pustules form and also become gangrenous; there results a characteristic eschar, which presents the appearance of an elevated patch, consisting of a zone of low, whitish vesicles

surrounding a depressed brownish, purplish, or black center, with an outer zone of red induration. Beyond this is usually a region of swelling and edema of variable extent. The spleen and the lymphatic glands in communication with the infected regions enlarge. There is often considerable local tenderness. The *constitutional symptoms* are those of septicopyemia: malaise, headache, depression, fever. According to the mode of introduction of the poison, or the direction in which infection spreads, other manifestations appear. Sometimes the gastro-intestinal tract appears to bear the brunt of the disease and there are nausea, vomiting, abdominal pains and diarrhea, the stools being bloody. Death may take place from exhaustion or from septicemia. At other times, thoracic symptoms predominate. There are then dyspnea, a sense of oppression of breathing, hemoptysis and cyanosis. Death may take place from edema of the larynx or of the mediastinum. Characteristic bacilli may often be found in the blood, pus, sputum, feces, or urine. Under proper treatment recovery frequently takes place.

Actinomycosis.

What is actinomycosis?

Actinomycosis is a condition dependent upon the presence of ray-fungi: *actinomyces bovis.* (Fig. 15.) The disease is more fre-

FIG. 15.



Actinomyces. (Ziegler.)

quent in drovers and in those that have to do with cattle, from which the parasite, as found in man, is usually derived. The

cattle become infected through their food. The fungus gains entrance through a breach in continuity of the surface and, finding its way to a suitable nidus, gives rise to the formation of a sero-purulent collection ; this manifests itself as a tumor that usually finds vent externally. In the matter discharged, yellowish miliary nodules, composed of fungi, can be detected. The lower jaw seems to be a favorite seat of the disease, infection taking place through decayed teeth ; sometimes extensive destruction of bone results. At other times, purulent collections form in internal viscera. The intestinal tract, the lungs, the skin, the brain, may be the seat of the disease. When the pleura is infected the ribs may suffer severely. The symptoms vary with the localization of the morbid process.

Foot-and-Mouth Disease.

What is foot-and-mouth disease?

Foot-and-mouth disease is a rare affection that occurs in sheep, cows, pigs and horses, and that occasionally seems to be transmitted to man. It manifests itself by the appearance of vesicles and bullæ in the mouth and on the feet at the margins of the hoofs and, in cows, on the udder and teats. The disease may be transferred directly to man by inoculation—thus to the butcher or to the veterinary surgeon, or it may be conveyed by milk. In man, vesicles form in the mouth, on the face, on the hands and on the feet. In the course of two or three days, the vesicles rupture, discharging opaque, yellowish fluid, and leaving dark-red ulcers. There are also fever, loss of appetite, pain in eating, swelling of the tongue, fetor of the breath, salivation and derangement of digestion. In children the disease may prove fatal.

Milk-Sickness.

What is milk-sickness?

This affection is probably a form of *food-poisoning*, arising in human beings who have partaken of the flesh or the milk (or

its products, butter and cheese) derived from animals suffering from a disorder known as trembles.

What are the symptoms of milk-sickness?

There is complaint of a sense of fatigue and languor, with headache, thirst, loss of appetite, nausea, vomiting, pyrosis, epigastric pain and constipation. The breath is peculiarly offensive; the skin is dry, the tongue moist and coated. Respiration is labored and sighing; the temperature is not elevated and may be subnormal; the pulse is not accelerated. Prostration may progress to coma and death within a few days. Convalescence is tardy and may prolong the illness for several weeks.

Hydatid (Echinococcus) Disease.

What is hydatid disease?

An *hydatid cyst* is a parasitic formation due to ingestion of the ova of the *tenia echinococcus*, the tape-worm of dogs, in which it is derived from the flesh of sheep, or pigs, or less frequently, kine, suffering from hydatid disease.

FIG. 16.



Tenia Echinococcus—vesicle, scolex and hooks. (After Heller.)

When the ovum enters the stomach of man, its capsule is dissolved, and the immature embryo or scolex (Fig. 16) is set free to continue its migrations. Arriving at its destination, the irritation to which it gives rise results in the formation of a membranous envelop, in which the parasite continues its development. This capsule, and its contents together constitute

an *hydatid cyst.* An hydatid cyst contains within the capsule, a vesicle or mother-sac, consisting of concentric layers of a gelatinous material, inclosing the embryo and more or less fluid. The fluid is clear, opalescent, and faintly alkaline. Within this develop other similar sacs, so-called daughter-vesicles, and within these again, granddaughter-vesicles; the mother-sac and its investing membrane continue to enlarge, if in a favorable situation, until ultimately the cyst attains an enormous size. Multiple cysts may form. The daughter-vesicles contain a germinating layer that produces new scolices. These consist of a head, four suckers and a row of hooklets. The discovery of the hooklets (Fig. 16) in fluid removed from the cyst is diagnostic. *Sterile echinococci* or *acephalocysts* do not produce scolices. The cysts may perforate into adjacent organs, rupture externally or undergo suppuration.

Hydatids may develop in various viscera, but are most common in the liver; then in order of frequency follow the genito-urinary system, the intestinal canal, the lungs and pleura, the brain and spinal canal.

The symptoms are those of a cystic tumor and vary with the size and situation of the formation. When superficial the cyst may yield on palpation and percussion a peculiar vibratile tremor or fremitus. When perforation or rupture takes place, or on exploratory puncture, the characteristic hooklets may be detected in the fluid obtained. In the event of suppuration chills, fever, and other symptoms may appear.

Trichiniasis.

What is trichiniasis?

Trichiniasis is a disease set up by the *trichina spiralis* (Fig. 17), a small roundworm that finds its way into the intestine with meat obtained from diseased swine.

In the intestine the mature female throws off embryos, which pass through the walls of the intestine and into the blood-current, finding their way into the voluntary muscles in different parts of the body. Here the embryos occasion irritation and inflammation, so that about each a capsule forms in which

lime-salts are in time deposited; in this way the death of the embryo may be brought about.

What are the symptoms of trichiniasis?

FIG. 17.



Trichina. *a*, Male; *b*, Female; *c*, Muscle-Trichina. (V. Jaksch)

The symptoms of trichiniasis present themselves in three *stages*. In the *first*, which lasts about a week, the trichinæ are undergoing development in the alimentary canal, as a result of which the symptoms of gastro-intestinal derangement appear. In the *second stage*, lasting two or three weeks, the embryos pass from the intestine into the muscular tissue. Finally, retrogressive changes take place in and around the trichinae encapsulated in the muscles.

In the *first stage*, the appetite is impaired; the tongue is coated, the breath is foul; there are malaise, nausea, a bad taste, diarrhea, abdominal pain and slight fever. Leukocytosis is marked, the esinophile cells being especially increased in number.

In the *second stage*, there is edema, apparent in the face and sometimes extending downward; there are also pains in the muscles, which are swollen; defects of the ocular muscles and of accommodation sometimes appear; the senses and various functions may be affected; sleep is disturbed and there is moderate fever. The muscles are sensitive to touch, which may detect the minute nodules. Death may result from exhaustion, pneumonia or ulceration of the bowel. In favorable cases, the symptoms gradually subside and the patient enters upon the *third stage* of the disease. Active symptoms are now in abeyance, but there may be some stiffness of the muscles, while noduless in the muscles may be detectable.

Ultimately, however, the patient may be restored to a fair degree of health.

Filariasis.

What is filariasis?

Filariasis is a condition arising from the presence in the lymph or the blood of the *filaria sanguinis hominis*, a nematode parasite, from 3 to 4 inches in length, that periodically discharges its ova into the circulation. By obstruction there may result *hematochyluria*, *lymph-scrotum* and *elephantiasis*.

Dracontiasis.

What is dracontiasis?

Dracontiasis is a condition arising from the invasion of the *filaria* or *dracunculus medinensis* (Guinea-worm), a parasite from 20 to 40 inches in length, that gains entrance through the stomach, the impregnated female penetrating the intestine and finding its way to the subcutaneous and intermuscular connective tissue, where it gives rise to the formation of a vesicle that ruptures and leaves an ulcer.

Acute Rheumatism—Rheumatic Fever.

What are the symptoms of acute rheumatism?

Acute rheumatism or *rheumatic fever* usually follows exposure to cold. An almost identical condition is sometimes observed in the course of puerperal fever and as a sequel of scarlatina. It has been thought to be dependent upon the presence of lactic acid in the blood. It is probable that the disorder is of bacterial origin, though the evidence is not yet conclusive. The disorder prevails most in temperate and humid climates, and young adult males are most commonly affected.

The *onset* is generally abrupt, one of the larger *joints* becoming painful, enlarged, hot and reddened. It is evidently inflamed; sometimes the presence of fluid can be detected. Soon, another joint, probably the corresponding joint on the opposite side, or the next contiguous joint, becomes similarly involved; and in this way the process may extend, until most or all of the large

joints are in turn attacked. On account of the pain, the patient is immovably helpless. From the surface of the body exudes an acid sweat. The *temperature* ordinarily ranges from 102° F. to 104° F., rising with each fresh access of joint-symptoms, and declining gradually with the termination of the disease. The pulse is disproportionately frequent ; it may be full and bounding. The *urine* is scanty, high-colored, and quite acid ; it may contain a trace of albumin. Leukocytosis is marked.

In many cases, to the articular manifestations *endocarditis* is added, as a result of which permanent valvular lesions may be established. Among other *complications* are inflammations of the pericardium, the pleura, the peritoneum, the kidney, the iris, and, though rarely, the cerebro-spinal meninges. As a rule, delirium is due to the toxemia and not to meningitis.

Sometimes the temperature rises even to *hyperpyrexia*, and there are decided cerebral symptoms—delirium, convulsions, coma, death.

Untreated, the *duration* of acute rheumatism is about six weeks ; sometimes much longer. Relapses are not uncommon. An attack predisposes to subsequent attacks.

Purpura is sometimes seen in the course of rheumatism ; sometimes tuberculated cutaneous nodules.

In some cases of rheumatism, hereditary influences can be traced. Some authorities consider the disease a dyscrasia, others a neurosis. There appears to be an indefinable, yet close relation between acute rheumatism and chorea.

Articular rheumatism not uncommonly accompanies, precedes or follows acute tonsillitis. In cases in which joint-involvement is insignificant or absent, pericarditis and endocarditis may sometimes be detected if carefully searched for.

How are pyemia and acute rheumatism to be differentiated ?

Both pyemia and acute rheumatism occasion arthritis, sweats, cardiac complications and cerebral symptoms. Periodicity and rigors, as well as metastatic invasion of internal structures remote from the primary seat of disease, which are common in pyemia, are wanting in rheumatism. The constitutional depression is more profound in pyemia than in rheumatism. In one, an obvious or obscure focus of suppuration exists ; in the

other, there is a history of rather abrupt onset following exposure to cold or wet.

How are acute synovitis and acute rheumatism to be differentiated?

Acute synovitis usually involves but a single joint; it is characteristic of acute rheumatism to progressively attack many joints. The constitutional phenomena are more profound in acute rheumatism than in acute synovitis.

The peculiar, acid sweats, as well as the cardiac complications of rheumatism, are not seen in synovitis.

In duration, synovitis is the shorter disease.

Gonorrheal Synovitis.

What is gonorrheal synovitis?

Occasionally, in the course of an attack of gonorrhea, a large joint—and usually but one joint, such as the knee, the elbow, the wrist, the ankle or the shoulder, becomes tumid, painful, tender and hot, and the adjacent textures may be edematous. The urethral discharge often ceases with the appearance of the synovitis. In successive attacks of gonorrhea, different joints may be involved. Permanent stiffness and impaired mobility constitute a common sequel. Cardiac complications have been observed in some cases of gonorrheal synovitis. The diagnosis depends upon a knowledge of the existence of a specific urethritis. Sometimes, the course of the temperature is suggestive of a pyemic or of a septic condition.

Syphilitic Arthritis.

What is syphilitic arthritis?

Every now and then, in the course of syphilis, one or more joints become involved in inflammation, with all of the characteristics of an arthritis. The discrimination of the condition depends upon a recognition of its association with syphilis. As a rule, cardiac complications are wanting in syphilis, and the arthritis is more strictly limited to one, or at most, two joints, the migratory tendency of acute articular rheumatism being absent.

Subacute Rheumatism.

What is subacute rheumatism?

At times, as a result of exposure to cold and wet, muscular movement becomes painful, in consequence of a rheumatic involvement of the muscle-sheaths or the tendons. The patient sometimes mistakes for paralytic weakness the restraint of motion by pain. The pain is also in some degree spontaneous and influenced by meteorologic conditions. Affecting the muscles of the lumbar region, the condition is termed *lumbago*. Affecting the muscles of the neck it may give rise to *torticollis* or *wry-neck*.

How are subacute rheumatism and neuralgia to be differentiated?

Both subacute rheumatism and neuralgia occur in paroxysms, superinduced by suitable meteorologic conditions. Rheumatism is more common in men; neuralgia in women. The pain of the former is rather dull and diffused; that of the latter sharp and confined to the distribution of an affected nerve, in the course of which may be found several tender points. Rheumatic pain more commonly than neuralgic pain is aggravated by movement.

How are subacute rheumatism and trichiniasis to be differentiated?

In trichiniasis, in addition to symptoms simulating those of subacute rheumatism, there are evidences of a cachexia, wasting, debility and symptoms of gastro-intestinal derangement, with a history of the ingestion of diseased meat. At an advanced stage of the disease, it may be possible to detect the nodules to which the encapsulated parasites give rise. Leukocytosis with eosinophilia is characteristic of trichiniasis.

Myalgia.

What is myalgia?

As a result of muscular strain, or of traumatism, groups of muscles become painful to touch and on movement, in association with some degree of cutaneous hyperesthesia.

Chronic Rheumatism.

What are the symptoms of chronic rheumatism?

As a result of an attack or of repeated attacks of acute rheumatism, numerous joints in different parts of the body remain enlarged, stiff and painful. Sometimes the condition is insidious, progressive and chronic from the outset. However produced, the functions of the various joints are impaired; attacks of pain occur and are apparently influenced by meteorologic conditions. Wasting of the muscular structures adjacent to the diseased joints takes place.

Chronic rheumatism may affect both joints and muscles, or the muscle sheaths or tendons only (*muscular rheumatism*), or it may attack the nerve-sheaths (*rheumatic neuralgia*). The principal symptoms of muscular and of nerve-rheumatism are pain, spontaneous and on motion, with accompanying tenderness, usually localized.

How are the enlarged joints of chronic rheumatism to be distinguished from those of chronic spinal disease?

Trophic changes in the large joints—enlargement, effusion, subluxation, arthropathies—take place in the course of some chronic spinal affections, notably posterior spinal sclerosis. As a rule, but one or a few joints are involved. In chronic rheumatism, many joints are involved. In case of disease of the spinal cord ordinary scrutiny should detect the existence of symptoms indicative of such a condition.

Acute Gout.

What are the symptoms of acute gout?

Acute gout is a recurrent paroxysmal affection thought to be dependent upon the presence of an excess of uric acid in the blood. It occurs chiefly in those of a sedentary or inactive mode of life, who indulge excessively in the luxuries of the table, more especially in meats, sweets, sweet wines and malt liquors. Those that have been active in out-door sports and afterwards, while diminishing their exercise, maintain the heavy diet formerly

appropriate, are extremely liable to gout. The tendency to gout is distinctly hereditary, and in some cases of marked gouty diathesis the attacks may occur despite personal abstemiousness.

The *paroxysm* may be brought on by an unusual excess, by a fit of anger, by worry or anxiety or by exhaustion. Its *advent* is sometimes unannounced; at other times, it is preceded by symptoms of indigestion, by mental irritability or depression. The attack usually *sets in* suddenly at night, the patient being awakened by intense pain most commonly referred to the metatarso-phalangeal joint of the great toe. There is *fever* in proportion to the intensity of the local affection. The pain moderates somewhat towards morning, when the patient falls into a gentle perspiration and is again able to sleep. Towards night the pain returns. The joint is now noticed to be tender, red, swollen and edematous; finally desquamation takes place. Other joints are successively involved, the morbid process showing an affinity for the smaller articulations. The attack gradually subsides, leaving the affected joints a little stiffened and swollen. At the height of the attack the proportion of uric acid in the blood is increased, while that excreted in the urine is diminished. When the paroxysm is over the quantity of uric acid in the urine is increased. In an attack of acute gout, the joint-symptoms may suddenly subside, and gastric, cardiac or even cerebro-spinal symptoms be substituted.

Sometimes the attack is manifested from the first by visceral rather than by articular crises. Visceral crises are more likely to occur late in the history of the case than early in its course, and they sometimes prove fatal.

How are acute rheumatism and acute gout to be differentiated?

Gout is an hereditary affection, occurring in paroxysms, in which the first metatarso-phalangeal articulation and other small joints are involved. In acute rheumatism, a history of heredity is frequently wanting; the large joints are especially involved. The duration of an attack of rheumatism is many weeks; an attack of gout subsides in the course of a week or two. The uratic deposits of gout are wanting in rheumatism. The sweats of rheumatism are absent from gout. Cardiac complications are common in acute rheumatism; gout never occasions

endocarditis, but chronic interstitial nephritis is a common sequel.

Chronic Gout.

What is chronic gout?

In those that have had a number of paroxysms of acute gout, or sometimes chronically from the first, deposits of urates take place around the diseased joints, in the articular cartilages and elsewhere, as in the lobe of the ear, in the kidneys and in the spleen. As a result there is painful thickening of the affected articulations, which are stiff and finally become deformed. Sometimes distinct "chalk-stones" may be felt, and in extreme cases these may cause ulceration and appear externally. Gout is a potent cause of arterio-capillary fibrosis. An excess of fibrous tissue develops in the viscera and in the walls of the blood-vessels, with secondary contraction. Chronic interstitial nephritis is a common sequel.

Lead-poisoning may give rise to lesions exactly resembling those of chronic or of subacute gout.

Lithemia.

What are the clinical features of lithemia?

Lithemia is modified gout—a manifestation of the uric-acid diathesis. It is caused by defective oxidation within the body and is dependent upon imperfect tissue-metabolism.

Lithemia manifests itself by varied symptoms, among which are sallowness or abnormal redness of complexion, impaired or perverted appetite, a metallic taste, deranged digestion, constipation, at times alternating with diarrhea, in some cases with the passage of mucous casts of the bowel, headache, vertigo, irritability of temper, a tendency to neurasthenia or melancholia, abnormal drowsiness or sleeplessness, palpitation of the heart, precordial distress, irritative cough, disturbance of vision, noises in the ears, anomalous cutaneous eruptions, transient localized edema, and cramps in the calves of the legs. *Micturition* may be frequent and burning, the *urine* usually being diminished in

quantity, of high specific gravity, and containing an excess of uric acid and urates; phosphates and calcium oxalate are likewise frequently in excess; albumin and tube-casts are sometimes found, and in cases attended with paroxysmal flushing red blood-corpuscles likewise. Chronic catarrhal hepatitis and functional inactivity of the liver are frequently—perhaps causally—associated with lithemia. Fibrous degeneration of the kidneys and of the walls of the smaller arteries, with cardiac hypertrophy, may be an ultimate *sequel*.

For what affections may lithemia be mistaken?

Unless in a given case one bears in mind the possibility of the existence of lithemia, and is on the alert for its detection, the condition may be mistaken for almost any functional disorder, or even for serious organic disease of the heart, brain, stomach, intestine, or other organ.

The discrimination depends partly upon the exclusion of visceral lesions, and partly upon the results of urinalysis: the finding of an excess of urates or free uric acid pointing to the existence of lithemia. An hereditary tendency to gout or rheumatism, or the existence of gout, rheumatism, or diabetes in other members of the patient's family, or the fact that the patient's habits of life are such as are likely to lead to the development of gout, should direct attention to the probability of the existence of lithemia.

It must not be forgotten that fibroid changes in the blood-vessels and kidneys are frequent concomitants of the uric-acid diathesis.

Rheumatoid Arthritis—Arthritis Deformans.

What are the clinical features of rheumatoid arthritis?

Rheumatoid arthritis, *arthritis deformans*, often incorrectly called *rheumatic gout*, is a morbid condition in which destructive changes take place in one or more joints of the body, resulting in thickening, impairment of mobility, deformity and pain. The articular cartilages undergo softening and absorption; the ends of the bones become enlarged and sclerotic, while the apposed

surfaces become smooth from mutual pressure; the subjacent bone, however, becomes rarefied and brittle; lime-salts are deposited in the remains of the articular cartilages. A peculiar crepitus due to the apposition of roughened surfaces can often be elicited on manipulation of the affected joint. At a later stage, eburnation takes place, and the apposed bony surfaces slide over one another with abnormal facility.

The disease manifests a tendency to symmetrical involvement. When the hands are involved, a peculiar deformity results—the fingers being deflected towards the ulnar side. Occasionally, hard fibrous nodules are found in the muscles at a short distance from the affected joints.

In some cases nodes form at the sides of the terminal phalanges. Less commonly a single joint is affected.

The *onset* of the disease is usually insidious; occasionally it is acute and attended with febrile symptoms, with pain, with swelling and with redness of the affected joints.

Arthritis deformans occurs in those exposed to unfavorable hygienic influences, in the weak and ill-fed, in those exhausted by frequent childbearing, by prolonged lactation, by grief or by anxiety.

The disease is not directly fatal. It occasions no cardiac complication. When the larger joints (especially the hip, knees and elbows) are involved there often results decided *muscular atrophy*.

How are chronic rheumatism and rheumatoid arthritis to be differentiated?

In rheumatism, the larger joints of the body are especially involved; rheumatoid arthritis involves the smaller joints as well. The deformity of rheumatism is essentially dependent upon a hyperplasia of the fibrous structures that enter into the formation of the articulation; the joint-lesions of rheumatoid arthritis are partly destructive in character and occasion peculiar deformities of the hands and feet, while irregular exostoses form on the articular extremities of the bones. The tendency to symmetrical invasion is more conspicuous in rheumatoid arthritis than in chronic rheumatism. The latter is usually a disease of advanced life; the former may appear in early adult life.

How are gout and rheumatoid arthritis to be differentiated?

Rheumatoid arthritis lacks the paroxysmal character of gout. Unlike gout, it is observed in the underfed rather than in the overfed. In rheumatoid arthritis the deposits of uric acid and of urates in various structures, characteristic of gout, are wanting. Gout never presents the peculiar deformities of the hands and feet seen in rheumatoid arthritis. The latter does not lead to the fibroid condition of the kidneys, heart and vessels to which gout gives rise.

THE BLOOD.

What are the methods of studying the constitution of the blood?

For purposes of diagnosis especially, the blood is frequently examined as to its *corpuscular richness*, as to the *proportion of hemoglobin* it contains, and as to the presence of *abnormal* bodies.

The *corpuscular richness* of the blood is determined by means of an instrument called a *hemocytometer* or *blood-cell counter*, which consists of a shallow cell of known capacity, mounted on a glass slide, in microscopic divisions of which the numbers of red and white blood-corpuscles contained in a centesimal dilution of blood are respectively counted. For enumerating the white cells alone a decimal dilution is to be preferred. The number of cells can also be determined by the use of the *hematokrit*, a centrifugal apparatus provided with graduated capillary tubes of known calibration. In healthy men the blood contains about five million red corpuscles to the cubic millimeter; in women, about four and a half millions. The number of white cells is held not to permanently exceed under normal conditions, 10,000 per cubic millimeter. The normal ratio of white cells to red cells varies between 1 : 400 and 1 : 800.

Differential studies of the leukocytes are made with the aid of selective stains. The blood, spread in a thin layer on a cover-slip or slide, is fixed by heat or otherwise and exposed to the action of eosin and methylene-blue or hematoxylin, etc. Several varieties of leukocytes are distinguished, in accordance with their morphology and their staining affinities: (1) small

mononuclear, or lymphocytes, from 15 to 25% ; (2) large mononuclear, from 3 to 6% ; (3) polymorphonuclear or neutrophiles, from 65 to 75% ; or (*a*) acidophile or eosinophile, from 1 to 7% ; (*b*) basophile, from 20 to 25% ; (*c*) neutrophile, from 65 to 75%.

The *proportion of hemoglobin* in the blood is determined by comparing the color of the diluted blood with a standard, called a *hemoglobinometer* or hemometer, the result being expressed in percentages.

The percentage of hemoglobin is sometimes spoken of as *absolute* or *total;* sometimes as *relative.* By the term *absolute percentage* is meant the comparative richness in hemoglobin of the whole volume of blood, as shown by the direct reading of the hemoglobinometer scale. By *relative percentage* is meant the relation or ratio of the absolute hemoglobin percentage to the percentage of red corpuscles. Thus, if in a given case, the number of red corpuscles to the cubic millimeter is estimated at 3,000,000, or 60% (5,000,000 being taken as the standard or 100 per cent.), and the hemiglobinometer reading is 54 per cent., the latter figure (54%) would represent the absolute hemoglobin percentage, while the relative hemoglobin percentage would be $\frac{54}{60}$ or 90 per cent.

Deficiency of blood is known as *anemia* or *oligemia* ; deficiency of corpuscles, as *oligocythemia;* deficiency of hemoglobin, as *oligochromemia;* deficiency in solid constituents of the plasma, especially albumin, as *hydremia.* Excess of white corpuscles may appropriately be designated *hyperleukocytosis;* deficiency of white corpuscles, *hypoleukocytosis.* The morbid condition, of which persistent excess of white corpuscles is often the most conspicuous and is perhaps the essential feature, is known as *leukemia* or *leukocythemia.* The normal diameter of the red blood-corpuscle is 7μ. Smaller red corpuscles are called *microcytes;* larger, *megalocytes.* Irregularly-shaped red corpuscles are called *poikilocytes.*

The blood sometimes contains *parasites,* such as the hematozoa of malaria, the spirilla of relapsing fever, the bacilli of anthrax, the embryos of filaria sanguinis hominis, etc.

Anemia.

What are the symptoms of anemia?

Simple anemia may be a *result* of hemorrhage, of long-continued discharges, of syphilis, malaria, fevers and wasting diseases, of mal-assimilation, of impaired nutrition, of the presence of parasites or of poisons in the system.

The *blood* is *deficient* in *quantity* and in *quality*. The actual volume of the circulating fluid, as well as its corpuscular richness, is diminished. The number of *red cells* and the number of *white cells* are less than normal; so is the absolute quantity of *hemoglobin*, while the relative proportion per corpuscle may be scarcely altered. Some of the red cells are ill-shaped and diminutive, and some nucleated.

Anemia long continued gives rise to fatty degeneration of various structures, notably of the walls of the bloodvessels, of the heart and of other viscera.

The *countenance* and visible mucous membranes are usually pale; though in exceptional instances of great vascularity of the face the complexion may be rosy. The eyeball has a bluish tint. There are shortness of breath, an undue readiness of fatigue, and a disinclination to mental or physical effort. The patient complains of neuralgia, of headache, of vertigo and of sleeplessness. The *appetite* and *digestion* are impaired; constipation is the rule. The *urine* is pale and may be of low specific gravity, from diminution in urea. Emaciation is not always evident. The action of the *heart* is enfeebled. Palpitation is common. The *pulse* is soft, compressible, and usually small. A soft, blowing murmur is often to be heard at the base of the heart and in the vessels of the neck; in the arteries less constantly than in the veins (*venous hum*, "*bruit de diable*"). Edema ultimately develops and hemorrhages from various surfaces may take place.

When anemia is associated with enlargement of the spleen it is termed *splenic anemia*. By some authorities splenic anemia is considered to be a variety of pseudo-leukemia. In the yellow, waxen countenance of its subjects, in its clinical course, which is sometimes remittent or intermittent, in its intensity and its fatality, the disease closely resembles pernicious anemia. Un-

like the latter affection, splenic anemia presents a definite visceral lesion, a relative increase of white cells, and a diminution in the relative percentage of hemoglobin.

Chlorosis.

What are the symptoms of chlorosis?

Chlorosis or *green-sickness* is a depraved condition of the blood, seen especially at about the time of puberty in young women, with derangement of menstruation. In addition to the symptoms of simple *anemia*, the *complexion* and the conjunctivæ present a yellowish-green hue. The number of *red corpuscles* is not diminished in the same degree as is the percentage of *hemoglobin*, while the number of *leukocytes* is not appreciably altered.

Pernicious Anemia.

What are the symptoms of pernicious or idiopathic anemia?

There is a form of anemia in which the impoverishment of the blood is marked, and which pursues a progressive and usually fatal course. It is more common in men than in women. No constant visceral lesion has been found associated with the disease. In some instances, there has been atrophy of the gastric glands; in others, disease of the medulla of the bones; in others, increase of iron-containing pigment in the liver; in still others, no lesion except that of the blood has been detected. The disease sometimes appears in connection with pregnancy.

The *symptoms* are those of intense anemia, with irregular outbreaks of febrile temperature. The *complexion* is pallid and often assumes a strikingly yellowish (lemon) hue. The lips and palpebral conjunctivæ may be white. The bones, especially the sternum, exhibit tenderness on pressure. The *pulse* is usually rapid. The *urine* may be notably dark-colored, and contains an excess of nitrogenous matters.

Sometimes deceptive remissions in the symptoms occur. The layer of subcutaneous fat is often well-preserved. Hemorrhages take place from the mucous surfaces, beneath the skin, into the

retina and elsewhere, as a result of fatty degeneration of the coats of the arteries. The *heart* is also likely to undergo fatty degeneration.

The volume of *blood* is small; the number of *red corpuscles* is considerably diminished (less than 2,000,000 per c. mm.), as is necessarily the absolute quantity of *hemoglobin*, of which, however, the relative proportion per corpuscle is increased. The red corpuscles are poorly developed and ill-shaped; some are nucleated. Megalocytes preponderate; microcytes and poikilocytes are likewise found. In some cases minute, highly-colored globules resembling "small red-tinged, fat globules" are seen. The number of white corpuscles may remain unaltered; in some cases, it has been increased; in others, diminished.

What special precautions must be taken in arriving at a diagnosis of the various forms of anemia?

When a condition of anemia is discovered, a searching inquiry into the history and symptoms and a careful physical examination must be made, to determine whether or not there be a coëxistent morbid condition, such as carcinoma, tuberculosis, hemorrhoids or other source of hemorrhage, nephritis, malarial infection, intestinal parasites, defective or insufficient nutrition, mal-assimilation, arsenical, plumbic, mercurial or other form of poisoning, and conditions that occasion jaundice.

Hyperleukocytosis (Leukocytosis).

What is leukocytosis?

Leukocytosis is a condition of the blood in which, without alteration in the number of red corpuscles, without enlargement of the spleen, of the liver or of the lymphatic glands, the number of white corpuscles undergoes an intermittent or transitory increase, involving especially the polymorphonuclear cells. It may be *physiologic*, as when it occurs after meals, during pregnancy or in the newborn; or *pathologic*, as when it occurs in association with suppuration, the various diseases of the blood and lymphatic glands, as chlorosis, pernicious anemia, leukemia; inflammatory, suppurative and exudative conditions (including infections), as pleurisy, pericarditis, meningitis, nephri-

tis, acute articular rheumatism, diphtheria, erysipelas, scarlet fever, variola, cholera, tetanus, trichiniasis; malignant disease, carcinoma and sarcoma; after the use of certain medicinal agents, as pilocarpin, antipyrin, nuclein, camphor, ethereal oils, bitters, etc.; or it may be a *transitional* stage of leukemia or pseudo-leukemia.

Leukemia.

What is leukemia or leukocythemia?

Leukemia is a morbid condition in which, in association with enlargement of the spleen, of the liver, of lymphatic glands, or with alterations in the medulla of bones, the blood contains a permanent excess of *leukocytes* and is deficient in *red cells* (Fig. 18).

FIG. 18.



Appearance of the blood in leukemia. (Funke.)

Three varieties of leukemia are recognized: *lienal*, *lymphatic* and *medullary*, as the spleen and liver, the lymphatic system, and the medulla of bones, respectively, are involved.

Early in the course of the disease, the *spleen* becomes conspicuously enlarged; sometimes friction-fremitus may be detected on palpation, and a blowing murmur on auscultation;

ascites may develop. In the further course of the disease, the liver and the lymphatic glands may also become enlarged; the bones may be painful and tender; changes in the kidneys and in the intestinal glands are sometimes noted.

The number of *white corpuscles* in the blood is increased beyond 10,000 to the cubic millimeter, sometimes in extraordinary degree. The number of *red cells* and the quantity of *hemoglobin* are diminished. Through increase in the number of the white, and diminution in that of the red, there ensues a great change in the ratio of white cells to red cells. The ratio is, as a rule, not less than 1 : 50, and may reach or exceed 1 : 1. Masses of leukocytes may occasion infarcts in the spleen or lungs. It is said that in lienal leukemia large leukocytes predominate; in lymphatic leukemia, small leukocytes; and in myelogenous leukemia, large leukocytes, having large nuclei, together with transition-types and many eosinophile cells.

The depravity of the blood occasions dyspnea and edema; conjoined with the secondary changes in the bloodvessels, it predisposes to extravasation of the blood. There are thus breathlessness on exertion, epistaxis and other hemorrhages from mucous surfaces and into the retina and subcutaneous connective tissue. Diarrhea is common. The *urine* contains an excess of uric acid. Rarely, disturbances of vision occur—as a result of hemorrhage, of leukemic deposit, or of leukemic retinitis. From time to time febrile periods, lasting from a few days to a week, in which the temperature may rise to 102° F., are observed. Remissions in all of the symptoms sometimes take place during the progress of the case.

Leukemia is most common between the ages of twenty and fifty; it is more common in men than in women. It is usually insidious in *onset*, the early clinical phenomena being those of anemia in general. Sometimes the disease pursues a rapidly fatal course. Epistaxis is often the first significant symptom. The *etiology* of leukemia is obscure. In some cases there has been a history of antecedent malarial infection; in others, of traumatism of the spleen; in others, of traumatism of bones. Privation and exposure are cited among predisposing and exciting causes.

Pseudo-leukemia.

What is pseudo-leukemia?

Pseudo-leukemia, Hodgkin's disease, lymphatic anemia, malignant lymphoma or *lymphadenoma*, is an affection characterized by hyperplasia of adenoid tissues, and especially by enlargement of the lymphatic glands in various parts of the body, and by changes in the blood. *Anemia* prevails, the number of *red corpuscles* and the relative proportion of *hemoglobin* being diminished, the number of *white corpuscles* remaining unchanged or not uncommonly being slightly increased. An apparent excess of white cells, due to decrease in the number of red cells, with consequent moderate elevation of the ratio, must not be mistaken for an actual increase in the number of leukocytes. The superficial glands, especially those of the neck and axilla, are usually chiefly involved. Sometimes the glandular enlargement is first noticed in the groin. The spleen, liver and other viscera may also be enlarged. In addition to the symptoms of anemia and malnutrition, pressure-phenomena are observed. Enlargement of the tonsils, of the lingual, postnasal, cervical, peritracheal and peri-bronchial glands gives rise to cough, dyspnea and dysphagia. The cerebral circulation may be impeded from pressure upon the vessels of the neck. Pressure upon nerves may cause paralysis, pain or edema. Jaundice and ascites are sometimes observed. There are, in addition, a tendency to effusions, hemorrhages and petechiæ, and from time to time febrile periods.

How are leukemia and pseudo-leukemia to be differentiated?

It is possible that leukemia and pseudo-leukemia represent stages of a single condition. Individual cases present symptoms of both affections. The differentiation of well-marked cases, however, is not difficult.

In leukemia, the spleen enlarges early; the lymph-glands are not necessarily involved; when they are, however, it is late in the disease. In pseudo-leukemia, the lymph-glands are primarily enlarged; the spleen and liver may escape entirely; should they become involved, it is late in the disease. In leukemia, the number of white blood-cells is always increased—

often in decided degree. The alteration of relation of white cells to red cells in pseudo-leukemia is due to decrease in the number of the red cells; actual excess of white cells constitutes no essential part of the disease; though not rarely observed, it never reaches the high figures of leukemia.

How are tuberculosis of the lymph-glands ("scrofula") and pseudo-leukemia to be differentiated?

The lymphatic glands over a wide distribution may become tuberculous. Such a condition will also be associated with anemia—so that a clinical picture will be presented simulating that of pseudo-leukemia. Microscopic examination of a portion of one of the growths will disclose the presence of tubercle-bacilli in the one instance and a condition of simple hyperplasia in the other.

In pseudo-leukemia the enlarged glands are not inflamed, and do not give rise to inflammation in the surrounding connective tissue. They are not hot or tender to the touch; they are not adherent to the skin, and are, as a rule, freely movable upon each other. Tuberculous glands are usually adherent to the skin, and to each other, and manifest a tendency to chronic inflammation, with caseation and suppuration. There are usually other evidences of tuberculosis present, in the bones, joints, or elsewhere; and the *facies* is often significant. Glandular tuberculosis and pseudo-leukemia may, however, coëxist; or inflammatory changes may be set up by traumatism in the glands of pseudo-leukemia.

Infantile Pseudo-leukemic Anemia.

What is infantile pseudo-leukemic anemia?

Jaksch has described in infants a form of anemia in the last stage of which there are a deficiency of *red corpuscles*, a deficiency of *hemoglobin*, and an increased number of *white corpuscles*.

The condition differs from leukemia in that the increase of white corpuscles is not so great, that eosinophile leukocytes are never present, that, while the spleen is enlarged, the liver is not, and that the prognosis is favorable.

Table showing alterations in the constitution of the blood in disease.

	Number of red cells.	Number of white cells.	Proportion of hemoglobin.	
			Total.	Per red cell.
Anemia . . .	Diminished.	Normal or diminished.	Diminished.	Normal or diminished.
Chlorosis . . .	Diminished.	Normal or diminished.	Greatly diminished.	Diminished.
Pernicious anemia .	Greatly diminished.	Variable.	Greatly diminished.	Increased.
Splenic anemia . .	Diminished.	Increased.	Diminished.	Diminished.
Pseudo-leukemia .	Diminished.	Normal or increased.	Diminished.	Normal or diminished.
Leukocytosis . .	Normal.	Increased.	Normal.	Normal.
Leukemia . . .	Diminished.	Greatly increased.	Diminished.	Normal or diminished.

Scorbutus.

What are the symptoms of scorbutus?

Scorbutus or *scurvy* is a disease dependent upon dietetic errors, especially a deficiency in certain substances usually contained in the juices of fresh vegetables and fruits. It was formerly common at sea, and in jails and poorhouses. Outbreaks still occasionally occur. Isolated cases sometimes appear under unexpected circumstances, as in adults upon restricted diet and in children on artificial food.

The disease is characterized by a depraved condition of the blood, with degeneration of the walls of the vessels and consequent hemorrhages. The *gums* are soft and spongy, and bleed on slight provocation or spontaneously. The *breath* is defiled by the fetor of the decomposing necrotic tissue. Hemorrhages take place from other mucous surfaces, while inflammation of serous membranes, with hemorrhagic effusions, may occur. Extravasations of blood take place into the subcutaneous tissues, especially in the course of the superficial veins of the dependent parts of the body, giving a mottled, bluish or purplish appear-

ance to the skin in more or less extensive areas. The discoloration frequently resembles that of a bruise, and, as it slowly fades, assumes a greenish tint. It is often quite persistent. Extravasation into the muscles may likewise occur. With this, edema is often associated. Brawny induration of the connective tissue in various parts of the body may develop. Diffuse, dull pains are felt. The *skin* is dry and rough. There is mental apathy, with a sense of lassitude and an undue readiness of fatigue. Fungated hemorrhagic ulcers form. Existing ulcers assume an unhealthy, spongy appearance. Fractured bones fail to unite, while the union of broken bones may be dissolved. The *complexion* is sallow; anemia becomes evident, with shortness of breath and rapidity and feebleness of pulse. In children pain and swelling of the extremities, with loss of mobility, and elevation of temperature are prominent symptoms. In some epidemics, dysentery has been a *complication*. Sometimes hemeralopia or night-blindness has been observed. Without proper treatment death takes place. With proper treatment recovery is slow.

Purpura.

What are the clinical features of purpura?

Purpura is a manifestation of a deteriorated condition of the blood and vessels, as a result of which petechiæ form, and hemorrhages from the mucous membranes take place. Sometimes blood is extravasated into the serous cavities; and occasionally intracranial hemorrhage occurs. The disease sometimes develops in the midst of apparent health; at other times, in association with infectious, toxic, cachectic and neurotic states and as a result of mechanical conditions, such as venous stasis and violent muscular strain. Sometimes articular symptoms are present. Occasionally, transient febrile periods are observed. Sometimes the disease *begins* acutely, with a chill followed by pain in the back or limbs, but without fever. The purpuric spots may be large or small, of limited or of extensive distribution. Usually they appear in successive crops. At first reddish, they soon become of a deep purple, gradually fading to

brown, and then to yellow, as they disappear. The varieties of purpura commonly recognized are *purpura simplex*, *purpura hæmorrhagica* and *purpura* (or *peliosis*) *rheumatica*. The names are sufficiently descriptive. Henoch has described a recurring variety, observed especially in children and characterized by an erythematous exanthem, attacks of pain, vomiting and diarrhea, slight articular pain or swelling and hemorrhages from mucous membranes.

How are scorbutus and purpura to be differentiated?

Scorbutus occurs among those that are massed together and are insufficiently supplied with fresh vegetables; purpura may develop amid circumstances apparently the most favorable and in the face of an abundant supply of fruits and vegetables. The gums may bleed in purpura, but they are wanting in the sponginess and lividity of scorbutus. The breath is fetid in scurvy, not necessarily so in purpura. The petechiæ of scorbutus usually develop about the hair-follicles; those of purpura are indifferent in distribution. The individual discolored patches are more extensive in scorbutus than in purpura. In the latter, large blotches can usually be resolved into a number of smaller spots aggregated. The hue in the two cases differs to the experienced eye. Scorbutus in children may closely simulate purpura.

How is purpura to be distinguished from measles?

The eruption of purpura has in certain instances presented sufficient superficial resemblance to that of morbilli to occasion mistake—the youth of the patients and the coincidence of febrile symptoms increasing the difficulty. The course of the eruption is, however, quite different in the two affections. The distribution of the purpuric spots is not so general as is that of the eruption of measles. The spots of purpura often appear first on the legs, while in measles the rash begins on the face and trunk. The purpuric spots change color and fade, new spots appearing while others are receding; no such phenomenon occurs in measles. Catarrhal symptoms do not occur in purpura; hemorrhages are not usual in measles. The characteristic temperature-curve of measles is absent from the course of purpura.

Hemophilia.

What are the clinical features of hemophilia?

Hemophilia is a morbid condition manifested by an abnormal tendency to the occurrence of hemorrhages, spontaneously or upon slight provocation. Individuals so affected are called "bleeders." The disease occurs in families; it is much the more common in males, but is mostly transmitted through females, in whom, when it occurs, its manifestations are usually mild. Slight wounds, a scratch, the extraction of a tooth, may be followed by alarming or even fatal hemorrhage. Petechiæ form, while sometimes large extravasations of blood take place into the subcutaneous textures and into the joints. The disease usually first makes its appearance early in life. The *diagnosis* depends essentially upon the history of hereditary transmission, the unusual proneness to the occurrence of alarming hemorrhage, the formation of petechiæ, the development of subcutaneous and articular extravasations of blood, and the appearance of the first symptoms early in life.

Addison's Disease.

What are the clinical features of Addison's disease?

Addison observed that in certain cases, in which after death the *supra-renal bodies* were found tuberculous, there had existed during life a peculiar pigmentation of the skin, with remarkable asthenia and nausea and vomiting. The discoloration is usually of a brownish hue, like that which develops in one exposed to a tropical sun; it resembles the pigmentation of the dark-skinned races. It occurs in plaques, on exposed surfaces, at parts that have been compressed or constricted, in the flexures of joints, and about the genitalia and the nipples. The mucous membrane of the mouth and tongue may be pigmented. Progressive weakness is manifested; and prostration finally develops. The subcutaneous fat, however, may be preserved. The *heart's action* is feeble; the *pulse* is small and compressible. The *appetite* is impaired; nausea and vomiting are common; there may be diarrhea or constipation. *Death*

may take place gradually, from exhaustion, or with unexpected suddenness.

Tuberculous disease of the supra-renal bodies has been found when there was no bronzing of the skin. In such cases it has been thought that the disease had not progressed sufficiently to have occasioned symptoms. In other cases the skin has been pigmented, but no changes were found in the supra-renal bodies. Pigmentation of the skin may, however, be a result of other conditions than supra-renal disease. By some, Addison's disease is thought to depend upon loss of an internal secretion of the adrenals; by others upon changes in the semilunar ganglia and branches of the sympathetic or visceral system of nerves.

Rachitis.

What are the clinical features of rachitis?

Rachitis is a disease dependent upon defective and perverted development of the osseous structures of growing children, probably as a consequence of faulty nutrition. There is doubtless congenital predisposition to its occurrence. The *bones* are soft and yielding, and wanting in their natural firmness and stability, so that various deformities result. The sides of the chest become flattened, and the sternum projects, giving rise to the "pigeon-breast." Nodules or "beads" form at the junction of the ribs and their cartilages. The long, supporting bones become bent and their epiphyseal extremities enlarged. The soft and deformed bones are especially liable to green-stick fractures. The *lower jaw* is narrow and dentition is delayed. The teeth may decay and fall out soon after their appearance. The *head* appears large; its summit is flat; the fontanels close late. Imperfect ossification of the cranial bones gives rise to so-called cranio-tabes, with "parchment-crackling." The deformities are maintained by the ultimate hardening of the affected bones.

Rachitic children present a pasty complexion and pearly conjunctivæ; they are undersized and poorly resist disease. They are restless, and display a tendency to excessive sweating of the head. Often there are diffuse soreness of the body and slight elevation of temperature. *Digestion* is impaired; the *abdomen*

is often protuberant. Laryngismus stridulus and convulsions are common occurrences. Internal viscera may be enlarged as a result of hyperplasia of the interstitial connective tissue.

Mollities Ossium.

What are the clinical manifestations of mollities ossium?

Mollities ossium is a morbid condition that develops in adults amid unfavorable hygienic surroundings and in women that have borne many children. It manifests itself by both softening and rarefaction of the bones, so that progression is impossible, and fractures are common. Occasionally febrile symptoms are present. *Death* is the common result, either from exhaustion, or from mechanical interference with respiration.

How are rachitis and mollities ossium to be differentiated?

Mollities ossium is a disease of adult life, attended by changes in developed bone, and usually of fatal termination. Rachitis is essentially a disease of childhood, dependent upon abnormalities in developing bone, and from which recovery usually takes place, only the sequels of the disease remaining.

THE HEART.

Inspection.

What can be learned of the heart by inspection?

On *inspecting* the *normal chest*, one perceives in the left fifth intercostal space, two inches below and an inch within the nipple, in an area of perhaps an inch in diameter, a gentle rise and fall—the *cardiac impulse* or *apex-beat*. It is less distinctly visible in fat than in lean persons.

The position of the apex-beat varies slightly with the respiratory movements, with posture, and with the state of the abdominal viscera. It may be *displaced*, as to the left by an effusion in the right pleural cavity, or by adhesions of the left pleura; or to the right by an effusion in the left pleural cavity, or by adhesions of the right pleura. It may be *increased* in

extent, as when the heart is enlarged, or when the pericardium is distended with fluid. Under varying conditions an impulse is seen in the epigastrium.

The impulse may be *strong*, as in cases of cardiac hypertrophy; *feeble*, as in cases of dilatation; *wavy*, as when the pericardial cavity is occupied by fluid; when the pericardium is adherent "systolic dimpling" occurs.

Palpation.

What is to be learned of the heart by palpation?

Palpation confirms and reinforces what is learned by inspection. An impulse that cannot be seen can sometimes be felt. A *feeble* impulse indicates that the action of the heart has been embarrased by disease or by an effusion in the pericardial cavity; a *strong* impulse is indicative of over-action, of hypertrophy. A pericardial friction-rub may-sometimes be felt. In most cases of mitral valvular obstruction, and in some of aortic insufficiency, a *purring tremor* is perceived on palpation.

Percussion.

What is to be learned of the heart by percussion?

By *percussion*—which is best practised with the patient recumbent—the approximate *size* of the heart can be learned. A considerable portion of the organ is covered by lung-tissue. By *superficial percussion* of a normal chest the cardiac dulness is found to be represented by a triangle included between a point at the lower margin of the fourth left costal cartilage at its junction with the sternum, another at the apex-beat, and a third at the lower extremity of the sternum at its left border. *Deep percussion* defines a somewhat more extended area.

The area of cardiac percussion-dulness is *increased* when the heart is enlarged, or the pericardial sac is distended by fluid, when the heart is uncovered by retraction of the lung, or in full expiration. The area is *diminished* when the heart is covered by emphysematous lung-tissue, or by the lungs in full inspiration.

Auscultation.

What is to be learned of the heart by auscultation?

Auscultation constitutes the most important method of physical examination of the heart. It reveals the frequency, the rhythm, the quality, and the purity of the heart-sounds.

In health, the heart of an adult at rest beats about seventy-two times to the minute. The *frequency* of action varies with the degree of bodily exertion, and with posture. It is *increased* by exertion and by excitement, after meals, in febrile affections and in many diseases of the heart; it is greater in the upright than in the horizontal posture; it is *diminished* by rest and by jaundice; in aortic obstruction, in fatty degeneration and during convalescence from acute disease. It is affected by various drugs and by nervous influences.

The action of the heart in health is *rhythmical* and *regular*. There is a prolonged, dull, *first sound*, and a shorter, sharper *second sound*, followed by an *interval of silence*—with rhythmical repetitions. Occasionally a cycle is omitted—the heart *intermits*. A normal first sound may be followed by two second sounds—*duplication*. The first sound may likewise be duplicated. The normal rhythm of the heart may be disturbed by organic disease of the heart, such as degeneration by functional or nervous disorder, and by derangement of other organs, as of the stomach.

The *first sound* of the normal adult heart is a dull, but well-defined thud; the *second* is a shorter, sharper, snapping or ringing sound; it is followed by an *interval of silence*.[1]

Three elements enter into the production of the first sound, which is synchronous with the contraction of the ventricles (*systole*) and the closure of the auriculo-ventricular (*mitral* and *tricuspid*) valves. These are: the *closure of the valves*, the *muscular contraction*, and the *impact of the heart* against the chest-

[1] There can likewise be detected by the trained ear a short interval of silence between the first and second sounds. The fact is mentioned here to avoid misleading the student in his further studies; but for practical purposes this "minor silence," as it is called, may be entirely ignored.

wall. The second sound, which is synchronous with the beginning of the *diastole*, is valvular ; it is due to the quick approximation of the semilunar flaps, preventing return of blood through the arterial (*aortic* and *pulmonary*) orifices.

The first sound of the heart may be *altered* in volume, in tone, in duration and in strength. These are usually *increased* permanently when the heart is hypertrophied, and may be *increased* temporarily under excitement and after the administration of certain drugs. They are *diminished* when from any cause the action of the heart is enfeebled—among such causes are dilatation, degeneration, pericardial effusion. The character of the second sound is largely dependent upon the tension in the arteries; the higher the tension the sharper the sound.

The *purity* of the heart-sounds depends upon the condition of the heart-muscle, upon the state of the blood and upon the functional efficiency of the various valves and orifices.

The functional *efficiency* of the valves and orifices of the heart can be determined by a study of the sounds of the heart as heard over the respective parts ; but these are so close together that points in the course of the blood-stream are selected for auscultation. That for the mitral sound corresponds to the situation of the apex-beat ; that for the aortic, to the junction of the right second costal cartilage with the sternum ; that for the tricuspid, to the ensiform cartilage ; and that for the pulmonary, to the left second intercostal space close to the sternum.

Alteration of the structure and derangement of the function of the valves and orifices of the heart are revealed by adventitious sounds, called *murmurs*, that accompany or replace the normal sounds, or occur in the interval between them. Murmurs are often blowing ; sometimes they are soft, sometimes harsh, sometimes musical.

The first sound occupies the ventricular *systole ;* the *diastole* is taken up by the second sound and the period of *silence.* During the systole, the blood pases from the ventricles into the aorta and pulmonary artery ; the arterial valves should be freely open ; the auriculo-ventricular valves should be perfectly closed : murmurs generated at the arterial orifices indicate obstruction to the outflow of blood from the ventricles ; murmurs generated at

the auriculo-ventricular orifices indicate reflux of blood into the auricles.

At the conclusion of the systole, there is a brief pause, followed by the diastole, during which the blood flows from the auricles into the ventricles; the auriculo-ventricular valves should be freely open; the arterial valves should be perfectly closed: murmurs generated at the auriculo-ventricular orifices indicate obstruction to the onflow of blood into the ventricles; murmurs generated at the arterial orifices indicate reflux of blood from the arteries.

Incompetency or insufficiency of a valve permits of regurgitation.

Constriction or occlusion of an orifice occasions obstruction.

The valves and orifices may be so altered by adhesion as to prevent perfect closure. Under such conditions both *regurgitant* and *obstructive* murmurs may be audible.

The seat and character of an endocardial murmur are deter-

FIG. 19.



Auscultation of the heart-sounds. The small letters indicate the situation of the valves; the large letters, the points for *auscultation*. *aA*, aortic; *mM*, mitral; *tT*, tricuspid; *pP*, pulmonary. (Vierordt.)

mined from the *place* at which it is best heard (site of *maximum intensity*), in association with the *time* of its occurrence and the direction in which it is *transmitted*.

Analogous conditions of the right and left heart necessarily give

rise to murmurs at the same time; analogous conditions of the auriculo-ventricular and arterial valves give rise to murmurs at different times; and conversely. Thus, for example, the murmurs of mitral insufficiency and of tricuspid insufficiency coincide in time with each other and with the murmurs of aortic obstruction and of pulmonary obstruction. Mitral and tricuspid regurgitant murmurs and aortic and pulmonary obstructive murmurs are heard during the systole—with or in place of the first sound. Mitral and tricuspid obstructive murmurs and aortic and pulmonary regurgitant murmurs are heard during the diastole; the latter two with or in place of the second sound; the first two, however, in what is normally the period of silence immediately preceding the first sound—hence *presystolic*.

Mitral regurgitant murmurs are heard best at the apex of the heart; they are transmitted in the course of the fifth and sixth ribs to the axilla, and may be heard below the posterior inferior angle of the scapula. *Mitral obstructive* murmurs are best heard at the apex of the heart, or a little above the apex; they are but feebly transmitted. *Aortic regurgitant* and *aortic obstructive* murmurs are most distinctly heard at the junction of the second costal cartilage on the right with the sternum; the former are transmitted downwards in the course of the sternum and may often be heard all over the chest; the latter, upwards in the course of the great vessels, especially the carotid. Exceptionally the murmur of mitral regurgitation may be so transmitted as to be heard at the aortic cartilage; or the murmur of aortic obstruction be transmitted downwards towards the apex of the heart. Murmurs generated at the orifices and by the valves of the right side of the heart are exceedingly rare. *Tricuspid regurgitant and obstructive* murmurs are best heard at the ensiform cartilage, or a little farther to the right. *Pulmonary* murmurs should be best heard in the second intercostal space on the left, close to the margin of the sternum. In anemia, soft blowing murmurs, dependent upon the condition of the blood, are heard in the same situation.

All murmurs heard in the precordial region are not endocardial. Auscultation reveals as well the sounds of pericardial friction as of adjacent pleural friction.

How are murmurs due to organic disease to be distinguished from so-called functional murmurs?

In addition to adventitious sounds generated at the orifices of the heart as a result of structural changes, murmurs are heard when the condition of the blood is deteriorated, or when, from disturbed action of the heart or other cause, abnormal currents are generated in the blood-stream. These so-called *functional murmurs* are distinguished by their inconstancy and their softness; they are usually heard only at the base of the heart and over the body of the organ; they are not transmitted; they are intensified by pressure with the stethoscope; and they disappear with the removal of the conditions upon which they depend. Organic murmurs are usually harsher, more constant, and vary comparatively little in character and intensity.

What is the sphygmograph?

The sphygmograph is an instrument by which an artery is made to record certain of the characters of its pulsation. The *sphygmogram* (Fig. 20) is an important aid in diagnosis, but

FIG. 20.



Normal pulse-tracing. (After Eichhorst.)

cannot be relied upon apart from the ordinary rational and physical signs. The use of the sphygmograph and the significance of its tracings must be learned by experience.

Malformation.

What are the most common malformations of the heart?

The most common malformations of the heart consist in an imperforate interventricular septum and a failure of the foramen ovale to close.

To what symptoms do malformations of the heart give rise?

Individuals in whom there exist serious abnormal communications between the lateral halves of the heart rarely reach adult life. Cyanosis is the most common symptom. Systolic murmurs are heard practically indistinguishable from those occasioned by valvular derangement.

Dextrocardia.

What is dextrocardia?

Dextrocardia is a congenital displacement of the heart on the right side, commonly associated with displacement of the liver on the left and of the spleen on the right.

How is dextrocardia to be recognized?

In case of dextrocardia the impulse of the heart is wanting in its usual situation, and is seen to the right of the sternum. The area of cardiac percussion-dulness occupies on the right an extent corresponding to that which it normally occupies on the left. The sounds of the heart are heard on the right side instead of on the left. The hepatic percussion-dulness is not found in its usual situation, but in a corresponding position on the left. The splenic dulness is found on the right instead of on the left.

Functional Disturbance of the Heart.

What is meant by functional disturbance of the heart?

Under various conditions, as when the nutrition is impaired or the digestion is deranged, as a result of overwork, or of dissipation, or of the excessive use of tobacco, or tea, or coffee, and in connection with gout or lithemia, with hysteria or hypochondriasis, the action of the heart may be deranged without recognizable structural change. There will be present such symptoms as pain, palpitation, anxiety, headache, vertigo and breathlessness, sometimes with irregularity and increased frequency of the heart's action. The diagnosis depends upon the recognition of the primary condition and upon the absence of the physical signs of a cardiac lesion. Functional disorder,

great in degree and long continued, may lead to structural change.

Tachycardia.

What is tachycardia?

Tachycardia is a term applied to a somewhat rare condition of excessive rapidity of the action of the heart, accompanied with palpitation, the rhythm of the heart sometimes remaining unaffected. The pulse may exceed 200 beats a minute. Occurring in paroxysms, the qualification *paroxysmal* is applied. When no etiologic lesion is discoverable, the condition is termed *essential paroxysmal tachycardia*. The paroxysm usually begins suddenly, with or without warning; at times without apparent exciting cause; at other times seeming to result from some such condition as overdistention of the stomach. Tachycardia may be *due* to temporary paralysis of the vagus or stimulation of the cardiac accelerator nerve. Increased cardiac dulness and indefinite murmurs may be observed during the attack, and disappear with subsidence of the symptoms. The condition may be unattended with other symptoms, and ordinarily does not shorten life. It may be a part of other neuroses. It has been observed in women at the menopause.

How does tachycardia differ from angina pectoris?

Tachycardia is wanting in the threatening symptoms of angina pectoris—the anxiety, the pain, the cardiac failure. Rapidity of the heart's action and palpitation are the essential features of tachycardia; while in angina pectoris the pulse is of variable frequency.

How does tachycardia differ from exophthalmic goiter?

Palpitation of the heart and increased frequency of the pulse are among the earliest phenomena of exophthalmic goiter. Before the thyroid gland has become enlarged or the eyeballs protrude, the distinction from tachycardia is not possible. There is no difficulty in the diagnosis, however, when not only the exophthalmos and the goiter, but the array of other symptoms that characterize exophthalmic goiter, have also developed.

Brachycardia.

What is brachycardia?

Unusual infrequency of heart-beat is normal to some individuals. In others it may be related to disease, febrile, metabolic, toxic, hemic, digestive, respiratory, circulatory, cutaneous, genito-urinary or nervous. Sometimes there is merely a discrepancy between the number of heart-beats and the number of pulse-beats.

Irritable Heart.

What are the phenomena of irritable heart?

Irritable heart is a condition originally observed in soldiers in active service, in which there are in addition to increased frequency of the action of the heart, often with disturbed rhythm, recurring attacks of palpitation and pain in the precordia. There are usually headache and vertigo, especially during the paroxysms. The general health may suffer little or not at all. The first sound may be short and sharp or may be barely audible; the second sound is accentuated. There is no constant murmur. The pulse is compressible and easily influenced by position. Respiration is but little if at all accelerated. A similar condition may develop in civil life in those unaccustomed to arduous labor called upon to perform unusual tasks. It has also been found in athletes and others who have committed excesses in physical exertion, and in masturbators. Under proper regimen, restoration to the normal results; under other circumstances, hypertrophy of the heart develops.

How does irritable heart differ from tachycardia?

Irritable heart results from well-recognized causes that are not concerned in tachycardia. In irritable heart the frequency is less than in tachycardia, and is habitual; in tachycardia the increased frequency is extraordinary and usually occurs in paroxysms. It is pain and palpitation rather than increased frequency of action that marks the paroxysmal seizures of irritable heart. Tachycardia is also wanting in the distressing subjective sensations and the grave issue of irritable heart.

Angina Pectoris.

What are the characteristics of angina pectoris?

Angina pectoris is a paroxysmal disorder for which no definite structural lesion has been found. Perhaps the most common condition associated with it is atheroma of the coronary arteries.

The attack *sets in* suddenly, with a sense of oppression, dyspnea, and pain in the precordia, rising to a high pitch of intensity, and attended with a sense of impending dissolution—and not uncommonly death does occur in the paroxysm. There is often a sensation as of throttling. The pain in the heart is described as "clutching," "squeezing," and "tearing." The pain radiates in various directions from the heart, especially to the left shoulder, and extends down the left arm. The *face* is pale, the features drawn, the *pulse* variable. The attacks may occur spontaneously, but are usually brought on by excitement or exertion, or by gastro-intestinal derangement. They recur with varying frequency, sometimes over a long period of years. The disease is most common in men in advanced middle life.

How does intercostal neuralgia differ from angina pectoris?

The pain of intercostal neuralgia never attains the intensity or presents the peculiar character of that of angina pectoris; nor are the general manifestations ever so portentous. The tender points of Valleix, found in intercostal neuralgia, are absent in angina pectoris. An herpetic eruption on the chest, following the course of an intercostal nerve, is diagnostic of neuralgia.

Hypertrophy.

What are the symptoms of hypertrophy of the heart?

The size of the heart is proportionate to the demands made upon it, or to the stimulation that it receives. Valvular disease and other conditions impeding the circulation may cause one or all of the cardiac chambers to enlarge and the walls to thicken. Hypertrophy, uncomplicated by valvular disease, results when the heart is called upon to perform excessive labor, or is stimu-

lated by abnormal nervous influences. Symptoms arise only when the action of the heart is in excess of the requirements of the system. Under such conditions there will be a sense of discomfort in the precordial region, palpitation, paroxysmal cough, shortness of breath, headache, vertigo, ringing in the ears, disturbed sleep, deranged digestion, a florid complexion, and a tendency to hemorrhage.

What are the physical signs of hypertrophy of the heart?

When the heart is *hypertrophied*, its *impulse* is decided and extended, and usually displaced to the left. The area of *percussion-dulness* is increased. As the enlargement, in most cases, principally involves the left ventricle, the area of dulness increases to the left, though enlargement of the left ventricle is likely to be followed in turn by hypertrophy of the remaining cavities. Owing to the position of the heart in the chest, the left ventricle may be moderately enlarged without giving rise to appreciable percutory abnormality. Enlargement of the right heart may be the first cause of extension of the area of dulness. The *action* of the heart is moderately accelerated, regular and rhythmical. The *first sound* is strong and booming, the *second* accentuated. The *pulse* is correspondingly full and strong.

Dilatation.

What are the symptoms of dilatation of the heart?

Dilatation of the heart occurs under pretty much the same conditions as give rise to hypertrophy, except that the organ is unable to meet the demands made upon it. Dilatation may thus be a sequel of hypertrophy. As dilatation is frequently an ultimate result of obstruction to the pulmonary circulation, the right heart usually suffers the more. The symptoms are those of *failing circulation:* precordial anxiety, palpitation, headache, vertigo, syncope, pallor, cough, dyspnea, venous congestion, and dropsy.

What are the physical signs of dilatation of the heart?

The cardiac *impulse* is feeble and diffused, and usually displaced to the left. The area of cardiac *percussion-dulness* is in-

creased. The *first sound* is weakened in correspondence with the disproportion between the enlargement of the cardiac chambers and the thickness of their walls; the *second* is little changed. If the dilatation has been great enough to so enlarge the orifices or weaken the muscles that the valves are no longer competent to effect complete closure, regurgitant murmurs may be developed, even in the absence of structural alterations in the valves. Such murmurs are usually rather soft and may be inconstant. The *action* of the heart may be rapid and irregular. The *pulse* is small and soft.

What are the distinctions between hypertrophy and dilatation of the heart?

HYPERTROPHY.	DILATATION.
Face florid; ringing in ears; rushing of blood to the head.	Face pallid; tendency to syncope; dropsy.
Cardiac impulse strong, extensive.	Impulse feeble, diffused, often wavy.
Increased area of percussion-dulness.	Increased area of percussion-dulness.
Action rapid, regular, rhythmical.	Action rapid, irregular.
First sound strong.	First sound relatively enfeebled.
Pulse full, strong.	Pulse small, yielding.

How does dilatation of the heart differ from fatty degeneration of the heart?

Fatty degeneration of the heart may be recognized when the skin has a peculiar, greasy appearance, when an arcus senilis is present, when the sounds of a heart not enlarged are exceedingly feeble, and attended with the symptoms of a failing circulation. It differs from dilatation in the unchanged or diminished size of the heart and the association with such other evidences of fatty degeneration as may be present. The pulse is usually, but not invariably, slow in fatty degeneration, while it is rapid in dilatation.

How is a pericardial effusion to be distinguished from dilatation of the heart?

Fluid may collect in the pericardium in connection with pericarditis, or as a part of a general dropsy. In dilatation of the heart the cardiac impulse is feeble and diffuse, but not as

feeble or as wavy as in effusion. When dilatation exists, the percussion-dulness in the precordia is not as extensive as when there is an effusion, nor is it peculiarly triangular in outline. While the sounds of the heart are enfeebled over the organ when dilatation exists, they are almost obliterated in case of effusion, except at the base, where, in case of pericarditis, friction-sounds may likewise be detected. The recognition of a condition that gives rise to pericardial effusion is an aid in diagnosis.

Valvular Disease.

What are the symptoms common to valvular lesions of the heart?

The functional deficiency of a diseased heart-valve may be compensated for by an improved condition of the heart-muscle. The presence or absence of rational signs will depend upon the extent and constancy of the structural and functional alteration of the muscle. If the alteration be too great, the symptoms will be those of an hypertrophied or overacting heart. If it be too little, the symptoms will be those of an enfeebled heart. If it be inconstant, symptoms of excitement, of overaction or of enfeeblement may appear in paroxysms, or for prolonged periods at irregular intervals. Even when the compensatory change is apparently constant and sufficient, there may be at times or continuously, with or without pallor and weakness, a sense of precordial discomfort, palpitation of the heart, shortness of breath, perhaps headache and vertigo—all aggravated by exertion. Often, when compensation is but slightly imperfect, there are evidences of gastric and intestinal derangement, and a sense of fulness in the hypochondria. Should the heart fail, and compensation be disturbed, the condition becomes virtually or actually that of dilatation of the heart; the phenomena are those of insufficient *vis a tergo* in the circulation. As a result of the stagnation of blood in the veins, there is engorgement of various organs, especially of the lungs, liver, kidneys and spleen; the existing symptoms become intensified, and in addition there appears dropsy, first manifesting itself in the lower extremities, and extending upwards.

When two or more lesions coëxist, the compensation is less readily established and more easily ruptured than in the case of single lesions. Pulmonary congestion and hemoptysis occur not infrequently.

Mitral Incompetency—Mitral Regurgitation.

What are the signs of incompetency of the mitral valve (mitral regurgitation)?

As a result of endocarditis, or of fibrous or calcareous degeneration, the bicuspid leaflets, or the chordæ tendinea, become thickened and contracted, interfering with the accurate apposition of the segments of the mitral valve during the systole. Sometimes a similar result is brought about by dilatation of the left side of the heart and of the mitral orifice. In consequence, blood is abnormally diverted backward through the mitral orifice, giving rise to a blowing (systolic) sound, heard with greatest distinctness at the apex, and transmitted, in the course of the fifth and sixth ribs, to the axilla and to the inferior angle of scapula. The heart is usually enlarged—under favorable conditions, hypertrophied; under unfavorable conditions, dilated. The pulmonary second sound is usually accentuated.

Mitral Obstruction.

What are the physical signs of obstruction at the mitral orifice?

The mitral orifice becomes contracted as a result of endocarditis or of degenerative changes in the segments of the valve. The characteristics of mitral obstruction are: a rumbling murmur heard just before the systole, with greatest intensity over the left ventricle, and a purring tremor or thrill perceived by the hand placed over the precordia. The first sound is short and sharp, often resembling the second sound; and the pulmonary second sound accentuated. The left ventricle diminishes in size, while the left auricle becomes enormously enlarged. In turn the right side of the heart becomes enlarged. The pulse is small and feeble. Not rarely mitral incompetency is associated with obstruction.

Aortic Obstruction.

What are the physical signs of aortic obstruction?

Aortic obstruction is a result of structural changes in the semilunar valves or in the aorta, induced by inflammation or degeneration. Aortic obstruction and incompetency are often associated. The action of the heart is strong; the left ventricle becomes hypertrophied; the pulse may be full and strong or small and tardy. Occasionally a systolic thrill can be felt at the base of the heart on the right. At midsternum and on the right side over the junction of the second costal cartilage with the sternum, a coarse, blowing, systolic murmur is heard, transmitted into the great vessels. The murmur of aortic obstruction is sometimes to be heard over the right carotid artery, when at midsternum and at the aortic cartilage merely an obscuration of the first sound can be detected.

Aortic obstruction, apart from insufficiency, is one of the less common lesions. A systolic murmur heard at the aortic cartilage and transmitted into the neck may be due to dilatation or roughening of the aorta, without valvular or orificial change, and may give rise to no circulatory disturbance.

Aortic Incompetency—Aortic Regurgitation.

What are the physical signs of incompetency of the aortic valve (aortic regurgitation)?

Aortic incompetency arises from conditions similar to those that give rise to imperfections at the other orifices of the heart, as well as from atheromatous changes. It is often associated with aortic obstruction. It is characterized by a blowing, diastolic murmur, replacing or accompanying the second sound of the heart, heard with greatest intensity at the aortic cartilage, and transmitted downwards in the course of the sternum. The action of the heart is powerful; the left ventricle becomes enormously dilated and hypertrophied; vascular pulsation is common and may be evident even in parts remote from the center of circulation, as the dorsal artery of the foot; the pulsa-

tion in the carotids may be so strong that the head is shaken; the pulse comes up well with the systole of the heart, but immediately recedes—constituting the gaseous pulse, also called the water-hammer pulse, or the pulse of Corrigan. This char-

FIG. 21.



Pulse-tracing of aortic insufficiency. (After Strümpell.)

acter is well shown in the sphygmographic tracing. (Fig. 21.) Sometimes a thrill can be felt and a presystolic murmur heard over the heart.

Tricuspid Incompetency—Tricuspid Regurgitation.

What are the signs of tricuspid regurgitation (incompetency of the tricuspid valve)?

Lesions of the valves and orifices of the right side of the heart are uncommon. They may be congenital, or due to endocarditis, although they more commonly follow disease of the left side of the heart and disease of the lungs.

If the tricuspid valve is incompetent, blood regurgitates with the contraction of the ventricle; in consequence, a blowing systolic murmur is heard at the ensiform cartilage, and systolic pulsation is visible in the veins of the neck. In some cases the liver is seen to pulsate.

Tricuspid Obstruction.

What are the signs of obstruction at the tricuspid orifice?

When the tricuspid orifice is obstructed the blood must accumulate in the peripheral veins; a presystolic murmur is heard at the ensiform cartilage.

Pulmonary Obstruction.

What are the signs of obstruction at the pulmonary orifice?

When the pulmonary orifice is obstructed a systolic murmur is heard in the third intercostal space on the left, close to the sternum.

Pulmonary Incompetency—Pulmonary Regurgitation.

What are the signs of incompetency of the pulmonary valve (pulmonary regurgitation)?

When the pulmonary valve is incompetent regurgitation occurs with the dilatation of the ventricle; as a result a diastolic murmur is heard at the junction of the third costal cartilage on the left with the sternum; the murmur may be transmitted downwards in the course of the sternum. The vascular phenomena attending aortic insufficiency are wanting.

Acute Pericarditis.

What are the distinguishing features of acute pericarditis?

Acute pericarditis may develop in association with rheumatism, infectious diseases, nephritis, or pleurisy, endocarditis, or other adjacent disease; it may also result from exposure to cold or from blows upon the chest. It is attended with pain in the precordial region, and sometimes in the epigastrium; considerable elevation of temperature and other febrile symptoms; a sense of anxiety; dyspnea; irregularity and increased rapidity of the heart; irritable cough; possibly headache, vertigo, deli-

Endocardial Murmurs.

Time of occurrence.	Site of greatest intensity.	Direction of Transmission.	Seat of lesion.	Nature of lesion.
Systolic.	At cardiac apex.	Along left fifth and sixth ribs—in left axilla—in the back, at inferior angle of left scapula.	Mitral orifice.	Incompetency—Regurgitation.
Systolic.	At junction of right second costal cartilage with sternum.	To junction of right clavicle with sternum—in course of right carotid.	Aortic orifice.	Narrowing—Obstruction.
Systolic.	At ensiform cartilage.	Feebly transmitted.	Tricuspid orifice.	Incompetency—Regurgitation.
Systolic.	At left second intercostal space close to sternum.	Feebly transmitted.	Pulmonary orifice.	Narrowing—Obstruction.
Diastolic.	At junction of right second costal cartilage with sternum.	To midsternum—in course of sternum.	Aortic orifice.	Incompetency—Regurgitation.
Diastolic.	At left second intercostal space close to sternum.	In course of sternum.	Pulmonary orifice.	Incompetency—Regurgitation.
(Diastolic) presystolic.	Over body of heart.	To apex of heart.	Mitral orifice.	Narrowing—Obstruction.
(Diastolic) presystolic.	At ensiform cartilage.	Feebly transmitted.	Tricuspid orifice.	Narrowing—Obstruction.

rium, nausea, and vomiting. The gastric phenomena may obscure the actual condition, or the cerebral symptoms may suggest meningitis ; careful inquiry and examination will, however, reveal the inflammation of the pericardium.

What are the physical signs of acute pericarditis?

In the *first stage*, the characteristic sign is a rubbing *friction-sound* heard close to the ear at one or more points in the precordia, synchronously with one or both sounds of the heart. The sound is to be distinguished from an endocardial murmur by its quality, its superficiality and by increase of distinctness caused by pressure with the stethoscope.

In the *second stage*—that of *effusion*, a murmur is no longer heard, except perhaps above the fluid at the origin of the great vessels. The exudation may be sero-fibrinous, hemorrhagic or purulent. The cardiac *impulse* is feeble, extended and wavy. The area of *percussion-dulness* is increased, and presents a peculiar outline—being *triangular*, with the base below. At the apex and over the body of the heart, the *sounds* are heard but feebly or not at all, while at the base they may be heard with ordinary distinctness.

In the *third stage*, the fluid has disappeared, and murmurs may again be heard. In the course of time, however, the opposite surfaces of the inflamed membrane become adherent, so that the pericardial cavity is in whole, or in part, obliterated. Adhesions to the chest-wall give rise to localized retraction, and to *systolic dimpling*. The heart in turn becomes enlarged.

Diaphragmatic adhesions may cause visible systolic tugging at points where the diaphragm is attached.

How are acute pericarditis and acute pleurisy to be differentiated?

Pericarditis and pleuritis arise under similar conditions. The two may be associated. The physical phenomena of both are analogous, differing, however, in degree, extent and situation. The friction-sound of pleurisy is heard synchronously with the respiratory movements, that of pericarditis synchronously with the cardiac movements—the latter is thus the more frequent. If doubt arise, the breathing should be suspended

for a short time: a pleural friction disappears. The percussion-dulness occasioned by an effusion into the pericardium is situated in the precordia, and has a characteristic triangular outline; the percussion-dulness of an effusion into the pleural cavity involves the lateral and posterior aspects, as well as the anterior aspect of the chest.

Diseases of the Myocardium.

What are the most common affections of the myocardium?

The myocardium may undergo *degenerative change* in consequence of interference with the blood-supply through the coronary arteries; of alterations in the quality of the blood from deficiency of red cells or of coloring matter, or from the presence of toxic substances in association with or independently of infectious diseases; of adjacent disease; and of high bodily temperature. In obese persons there may also be an *increased deposit of fat* upon or in the myocardium. Interference with the supply of blood through the coronary arteries may give rise to *anemic necrosis*, possibly with *aneurismal formation* and *rupture*, or even the development of an *abscess*, and to *fibrosis* of the myocardium. General anemia may cause *fatty degeneration* of the muscular structure of the heart; as may also some toxic processes, while others cause fibrosis. *Parenchymatous degeneration* is usually the result of some toxic process or of infectious disease, and myocarditis is usually present with endocarditis and pericarditis. *Fragmentation* and *segmentation* may attend other forms of myocardial disease. Among other morbid processes that may attack the myocardium are *brown atrophy*, *amyloid degeneration*, *hyaline transformation* and *calcareous degeneration*. The heart may, further, be the seat of *syphilis*, of *tuberculosis* and of *new-growths*.

What are the most conspicuous symptoms of disease of the myocardium?

Disease of the myocardium may be unattended with symptoms, and sudden death may occur. During life the action of the heart may be enfeebled and slow, the rhythm disturbed, even to intermittence, and precordial distress, palpitation, dyspnea, vertigo and edema be present.

Acute Endocarditis.

What are the characteristics of acute endocarditis?

Acute endocarditis develops in the course of various infectious diseases and of nephritis and chorea, but more especially in connection with acute rheumatism and other varieties of arthritis. It is not rare in connection with tonsillitis and specific urethritis. It may be engrafted upon pre-existing chronic endocarditis. It may be *simple* or *vegetative*, or *malignant* or *ulcerative*. The valves of the heart are more often affected than its walls. The disease is characterized by a sense of distress in the precordia, palpitation and increased frequency of action of the heart, sometimes with irregularity, elevation of temperature, dyspnea, slight cough and an anxious expression. In addition, the face may be flushed; there may be chills, headache, vertigo, delirium, icterus, irritability of the stomach, and diarrhea. Finally, symptoms of embolism in various parts of the body may appear. The characteristic sign of acute endocarditis is a soft, blowing, systolic murmur, though it is possible for endocarditis to exist without the detection of a murmur. The occurrence of intermittent fever in a case of chronic endocarditis should awaken suspicion of its ulcerative character.

How is the murmur of an acute endocarditis to be distinguished from a murmur the result of a past endocarditis?

The murmur of acute endocarditis is soft, inconstant in its seat, unattended with enlargement of the heart and associated with the fever, precordial anxiety and other symptoms of acute endocarditis; a murmur dependent upon a past endocarditis is harsher, fixed in seat, unattended with fever, but associated with cardiac enlargement and a history of one of the conditions that may cause endocarditis or of a remote attack of endocarditis.

ACUTE ENDOCARDITIS.	CHRONIC VALVULAR DISEASE.
The murmur is soft and inconstant in seat and occurrence.	The murmur is harsher, fixed in seat and constant in occurrence.
Unattended with enlargement of the heart.	Attended with enlargement of the heart.

ACUTE ENDOCARDITIS.	CHRONIC VALVULAR DISEASE.
Associated with fever, precordial anxiety and other symptoms of acute endocarditis.	Not associated with fever or other symptoms of an acute condition.
No history of a remote acute endocarditis.	Remote history of acute endocarditis.
Associated with rheumatism or other cause of acute endocarditis.	Absence of an immediate cause of acute endocarditis.

How is acute endocarditis to be distinguished from acute pericarditis?

The pain of pericarditis is likely to be more severe than that of endocarditis. The murmur of pericarditis appears closer to the ear than that of endocarditis; the former is a friction-sound, diastolic as well as systolic; the latter is blowing in character and only systolic. When a pericardial effusion exists, it presents a peculiar area of dulness on percussion, the heart-sounds at the apex are enfeebled or they may be absent, and there is a diffused, feeble and wavy impulse, not found in endocarditis.

How are a pericarditic and a pericardial effusion to be differentiated?

An effusion of fluid into the pericardium may take place as part of a general dropsy. It presents physical phenomena indistinguishable from those of pericarditis in the stage of effusion, though without the febrile symptoms, and the friction-murmurs of the first and third stages. Other evidences of general dropsy, such as edema, pleural and peritoneal effusions, and symptoms or signs of the causative disease—nephritis, for example—will be found if looked for.

Heart-Clot.

What are the symptoms of heart-clot?

The blood may coagulate in the cavities of the heart in the course of endocarditis, of diseases in which the coagulability of the blood is increased, such as pneumonia, and when from any cause the action of the heart has become enfeebled. Under such circumstances the surface of the body becomes cold and livid,

there is intense dyspnea, the action of the heart becomes rapid, feeble and irregular, a faint murmur may be heard over the organ and the area of cardiac percussion-dulness is increased. There is great anxiety, nausea, vomiting, nervous excitement, delirium, venous turgidity and attacks of syncope.

Diseases of the Mediastinum.

What are the most common disorders of the mediastinum?

There may be simple or suppurative lymphadenitis, new-growths and emphysema.

What are the general clinical features of mediastinal disease?

Apart from the constitutional symptoms there may be pain, dyspnea, cough, hoarseness, distention of veins, edema, dysphagia, variations in the size of the pupils, with protrusion of the sternum, dulness on percussion, enfeeblement of the breath-sounds, diminished vocal resonance and fremitus, and displacement of the heart. Emphysema, when it appears externally, gives rise to a peculiar doughy swelling, with crackling on palpation.

Thoracic Aneurism.

What are the symptoms of thoracic aneurism?

The symptoms and physical signs of *thoracic aneurism* vary with the location of the aneurism. Aneurisms of the aorta are the most common. The frequency of aneurisms of the aorta diminishes with the distance from the heart. The *characteristic manifestations* of an aneurism are the existence in the course of a bloodvessel of a *pulsating expansile tumor*, attended with a *thrill* and *bruit* and diastolic shock; on percussion a circumscribed *area of dulness* will be found to correspond with the area of expansile pulsation, over which a thrill can be perceived with the palpating hand, and a systolic bruit can be heard on auscultation. Other *symptoms* depend upon the compression of adjacent structures. There may thus be pain, a peculiar metallic cough, with or without expectoration; dyspnea, dysphagia; irregularity of the action of the heart; venous stagna-

tion and edema; manifestations of derangement of the cerebral circulation; localized sweating; dilatation or contraction of one pupil; paralysis of one or both vocal bands; enfeeblement of respiratory murmur over certain areas, from obstruction of a large bronchus; inequality of the radial pulses or obliteration of pulsation in peripheral arteries; tracheal tugging; neuralgic pain and erosion of bone. Rupture may take place, externally or into an adjacent viscus, with gradual oozing or fatal hemorrhage. Examination with the fluoroscope will usually reveal the presence of an abnormal shadow, which may be seen to contract and expand with the pulsation of the aneurism. The shadow will be more definitely located by skiagraphy.

How is an intra-thoracic tumor to be distinguished from an aortic aneurism?

An aneurism develops only in the course of a bloodvessl; a tumor of any other sort is not so restricted. The former presents pulsation, expansion, and usually thrill and bruit; the latter may rise and fall if seated over an artery, but does not expand or present a thrill or bruit. The sphygmographic tracing (taken *in loco*) of a solid tumor situated above a normal artery is that of the normal artery; the curve given by an aneurism is abnormal and often characteristic. Intra-thoracic tumors are usually either secondary or give rise to metastases. If the tumor were a gumma, the diagnosis would depend upon a history of syphilis or upon other manifestations of syphilis, together with a recession of the symptoms on appropriate treatment. Tuberculosis of the mediastinal glands would be but one manifestation of a general tuberculosis.

How is an abscess of the mediastinum to be distinguished from an aneurism of the aorta?

In addition to the features that distinguish all other thoracic tumors from aneurism, an abscess presents fluctuation if accessible, rigors, elevation of temperature, and sweating. There is a history of traumatism or of pyemia.

How is incompetency of the aortic valve to be distinguished from an aneurism of the aorta?

Incompetency of the aortic valve may be attended with marked pulsation in the course of the aorta, with a thrill and

bruit, but the evidences of tumor and the consequences of compression are wanting; the radial pulses are of the Corrigan or water-hammer type, but are equal; while the second sound of the heart at the base is accompanied or replaced by a murmur transmitted downwards in the course of the sternum.

How is a dilated auricle to be distinguished from an aortic aneurism?

An enlarged auricle may give rise to pulsation at the base of the heart, but not to the compression-phenomena of an aneurism, not to the thrill or bruit, beyond the sounds generated in the heart itself as a result of the lesion that has led to the enlargement of the auricle: usually obstruction at the mitral orifice with a presystolic murmur.

How are a thoracic aneurism and pulsating pleurisy to be differentiated?

While left-sided empyema is sometimes attended with pulsation transmitted through the heart, it is unattended with the auscultatory signs and the expansion of aneurism, and it is, besides, indicated by the physical signs of a pleural effusion.

Arterio-Capillary Fibrosis.

What is arterio-capillary fibrosis?

Arterio-capillary fibrosis is a degenerative process of the walls of the smaller bloodvessels, due to the long-continued circulation in the blood of irritants, such as alcohol, lead, the poisons of syphilis, of gout, of rheumatism, and of infectious diseases. It may be a result also of constant or frequent increase in arterial pressure from whatever cause. Sometimes an inherited predisposition exists. Vaso-motor spasm and hyperplasia of the connective tissue of the arteries and of the intercellular elements of parenchymatous organs take place. The coats of the vessels become thickened and the size of organs becomes increased. From secondary contraction of the newly-formed fibrous tissue, the lumen of the arteries becomes narrowed and the organs become reduced in size. The circulation is curtailed and nutritive processes are interfered with; atrophy may

result; thrombi may form *in situ*. Sclerotic vessels are prone to rupture, in the brain, in the retina, and elsewhere. It has been suggested that the condition is really one of hyperplasia of the muscular layer of the arteries—*arterial hypermyotrophy*.

What are the symptoms of arterio-capillary fibrosis?

The diagnostic symptoms of arterio-capillary fibrosis or arterio-sclerosis are those of vascular spasm, impeded circulation, and impaired nutrition: shortness of breath, vertigo, mental impairment, nervous derangement, irregularity of cardiac action, resistant radial arteries, prominent, tortuous temporals, increased arterial tension. Edema may be present. The *urine* becomes increased in quantity and may contain a trace of albumin and an occasional tube-cast. Numbness and coldness of the extremities are common. The *nails* are sometimes bluish from impeded circulation. The *knee-jerk* is often exaggerated. Transient local palsies sometimes occur. Angina pectoris and pseudo-angina are not uncommon. The action of the *heart* may be arrhythmic and the sounds duplicated; the second or arterial sound is usually accentuated. An hypertrophied left ventricle, without valvular disease of the heart, should be suggestive of the existence of vascular disease. Chronic interstitial nephritis is a frequent result of arterio-capillary fibrosis, or of the same causes that give rise to the vascular changes. Cerebral and retinal thrombosis or hemorrhage may be *complications*.

Local Syncope—Local Asphyxia—Local or Symmetrical Gangrene.

What are the symptoms of Raynaud's disease?

Raynaud's disease is a vaso-motor neurosis characterized by pallor, lividity or gangrene of peripheral portions of the body, as fingers, toes, nose and ears, especially induced by cold, and sometimes attended with hemoglobinuria. The disorder is believed to be dependent upon spasm of arterioles, with dilatation of small veins. In accordance with the intensity of the morbid process and the clinical manifestations the condition is designated *local syncope, local asphyxia* or *local gangrene*. The affected

parts may be the seat of severe pain. When gangrene takes place cicatrization follows as a rule. In rare instances extensive areas are involved, sometimes even parts of the trunk, and death may result.

How are the symptoms of Raynaud's disease and those resulting from frost-bite to be differentiated?

The appearances presented by Raynaud's disease and those presented by frost-bite may be almost indistinguishable. Frost-bite is not likely, however, to exhibit the symmetry of distribution observed in cases of Raynaud's disease. The symptoms of frost-bite, moreover, usually recede when cold weather departs, while the manifestations of Raynaud's disease persist, though, perhaps, also aggravated by cold. Raynaud's disease is not necessarily dependent upon severe cold as a cause; while frost-bite is essentially so dependent.

Erythromelalgia.

What is erythromelalgia?

Erythromelalgia is a morbid condition characterized by pain and redness of one or more extremities, aggravated by the dependent position and by heat, and attended with local elevation of temperature. It is thought to be due to inflammation of nerve-end filaments. Peripheral nerves have been found degenerated and peripheral arteries sclerotic. The condition has been observed also in association with central nervous disease.

How are Raynaud's disease and erythromelalgia to be differentiated?

Raynaud's disease is more common in females, erythromelalgia in males. In the former the affected parts become white or cyanotic or gangrenous and cold, while in the latter they become dusky-red or violaceous and hot. Erythromelalgia is by far the more painful affection and is aggravated by heat and the dependent position, while Raynaud's disease is made worse by cold.

Angioneurotic Edema.

What is angioneurotic edema?

Angioneurotic edema is a vaso-motor disorder characterized by the appearance suddenly of more or less circumscribed swellings in one or other portion of the body, with equally sudden disappearance. The attack is sometimes attended with colicky pain and symptoms of gastro-intestinal disturbance. Involvement of the larynx may cause death by asphyxia. Some cases exhibit periodicity of recurrence. The predisposition to the affection is often inherited.

THE RESPIRATORY SYSTEM.

How are affections of the upper air-passages (nose, pharynx, larynx and trachea) to be diagnosticated?

The diagnosis of affections of the upper air-passages is based, in part, upon the symptoms; but the only certain methods are those of *direct inspection*, or of *indirect inspection* by means of a suitable mirror. For both direct and indirect inspection, proper illumination is necessary. When direct daylight does not suffice, a reflector is employed, either with daylight or with artificial light. There are also apparatus for direct illumination by artificial light. Palpation with probe or finger is sometimes additionally necessary.

It is sometimes necessary to remove from the line of vision, by means of suitable retractors, certain structures, as the anterior palatine folds, the uvula, soft palate or epiglottis.

By *translumination*, or transmission of light through the tissues, information is sometimes gained as to the comparative density of structure.

Direct inspection of the nasal passages by means of a speculum inserted by way of the nostrils is called *anterior rhinoscopy.*

Indirect inspection of the nasal passages by means of a mirror so placed in the pharynx, behind and below the soft palate, as to reflect an image of the posterior choanæ and neighboring structures, is called *posterior rhinoscopy.* The

vault of the pharynx and the mouths of the Eustachian tubes are examined by the methods of posterior rhinoscopy.

Direct inspection of the pharynx is termed *direct pharyngoscopy*. To carry it out, only light to illuminate and a spatula to depress the tongue are required. Those parts of the pharynx inaccessible to direct inspection are examined by means of a mirror appropriately placed (*indirect pharyngoscopy*).

The laryngeal structures are rarely accessible to direct inspection. Sometimes a part of the epiglottis may be seen, and in exceptional instances a glimpse of the supra-arytenoid structures has been obtained. With the aid of a special tongue-depressor (Kiastew's autoscope), however, the structures may be dragged forward and the vocal bands and posterior laryngeal wall brought into view.

Laryngoscopy is practised by means of a mirror so placed in the pharynx, with the uvula lifted out of the way upon the back of the mirror, as to reflect an image of the deeper parts. In favorable cases, the interior of the trachea, to its bifurcation, may be seen. When the patient phonates, the superior face of the vocal bands and the structures above are reflected. When the patient inspires deeply, the internal face of the vocal bands and the structures below are additionally revealed. One endeavors to observe the contour, the color and the motility of the various parts, and to determine the presence or absence of secretion, ulceration, cicatrices, abnormal growths or foreign bodies.

To what conditions may difficulty of nasal respiration be due?

Difficulty of nasal respiration may be dependent upon enlargement, engorgement or distortion of the septum or of the turbinate bodies, upon the presence of foreign bodies, or of polypi or other neoplasms, or, especially in children, upon overgrowth of the glands of the vault of the pharynx (adenoid vegetations). The *diagnosis* can be made only by rhinoscopy, anterior and posterior, sometimes in conjunction with palpation.

What symptoms other than those directly referable to the nose may arise from the conditions that occasion difficulty in nasal respiration?

In addition to the local symptoms, the conditions that occa-

sion difficulty of nasal respiration may be attended by anemia from deficient oxygenation, together with persistent cough, facial tic, headache, asthmatoid seizures, epileptiform convulsions, chorea, exopthalmic goiter, or various other reflex disturbances. The *diagnosis* is to be based upon a careful study of all of the phenomena and conditions, general and local.

What is a frequent, unsuspected cause of epistaxis?

An ulceration or erosion of the mucous membrane of the nasal septum, usually situated near the anterior inferior angle of the triangular cartilage, is a not uncommon cause of epistaxis. Malignant growth, especially sarcoma, which may involve the frontal sinuses and even be hidden from rhinoscopy is another condition worthy of mention. The bleeding may be insignificant in quantity, and might easily be considered of little moment.

Coryza—Acute Nasal Catarrh.

What is coryza?

Coryza or *acute nasal catarrh* is a superficial inflammation of the nasal mucous membrane, attended by sneezing and a profuse watery discharge from the nostrils. There is usually headache and sometimes more or less fever. The swollen membrane and more especially the engorged turbinate bodies impede respiration and modify articulation. In the course of a few days, the discharge becomes more viscid, often muco-purulent. Usually the attack subsides completely in a week. Conjunctivitis, pharyngitis, laryngitis, bronchitis and otitis media are among the *complications* and *sequelæ* sometimes manifested.

How is coryza to be differentiated from influenza?

Some authorities refuse to make a distinction between coryza and influenza. Coryza, with conjunctivitis or bronchitis, is sometimes epidemic. From fully-developed influenza, simple coryza, whether due to cold, to non-specific irritation or to infection, differs by the absence of the constitutional symptoms and of the profound depression characterizing the former disease. Characteristic bacilli are present in the secretions from a case of influenza.

Hay-Fever.

What is hay-fever?

Hay-fever, *hay-asthma*, *rag-weed fever*, *autumnal catarrh*, *June cold*, *rose-cold*, *idiosyncratic coryza*, *periodic vasomotor coryza*, are names applied to a group of symptoms developed in susceptible individuals as the effect of special irritants upon the mucous membranes of the eyes and air-passages, more especially of the nose. The *manifestations* vary much in severity and in constancy in different individuals, and in the same individual at different times, and comprise conjunctivitis, coryza, pharyngitis, laryngitis, bronchitis, asthma and gastric, enteric and renal crises. There is commonly intolerable itching of the eyelids and of the palate. Fever is not common; but when it occurs, it is irregular.

In some individuals the syndrome occurs when any irritating substance gains entrance to the nasal passages. In others the powder of certain drugs, such as ipecacuanha, or the pollen of certain plants, such as grasses, rag-weed, roses, is the *exciting cause*. The *predisposing cause* is usually a neurotic constitution, perhaps induced by excessive mental exertion or by undue indulgences; sometimes the presence of nasal abnormities aggravates the symptoms.

In those in whom the attacks depend on special pollens, the manifestations necessarily recur *periodically*. In North America, the rag-weed is the most common provocative agent, and with most sufferers the attacks begin about the middle of August, and last until the cause has disappeared. June-colds are dependent on hay and roses.

How is hay-fever to be distinguished from simple coryza?

The distinguishing features of hay-fever are the invariability of its causation, and the severity and association of its symptoms; together with the rapid disappearance of symptoms on removal to a locality (usually the mountains or the seashore) where the provocative agency is not present.

Acute Laryngitis.

What are the symptoms of acute laryngitis?

Acute laryngitis usually results from exposure to cold, or from the inhalation of irritating fumes. It occurs deuteropathically in the course of many inflammatory and infectious diseases, whether general or of the respiratory or digestive organs. With great variations in the degree of severity, it manifests itself by laryngeal irritation, hoarseness or painful aphonia, painful deglutition, dyspnea and slight hawking cough, with mucous expectoration.

There may also be slight elevation of temperature.

Laryngoscopically, the laryngeal structures will be seen to be more or less reddened and swollen. If edema occur, the breathing will be labored and stridulous, swallowing excessively painful and finally impossible, and, as a result of the impediment to respiration, cyanosis may develop. Laryngoscopy or palpation will reveal the cause.

How is laryngitis to be distinguished from pharyngitis?

The only reliable method of discrimination is by inspection, both directly and by means of the mirror, as symptoms are very likely to be misleading, and the two affections not infrequently coexist.

In pharyngitis the alteration in voice—aphonia or hoarseness, of laryngitis is wanting. The difficulty and pain of swallowing are likely to be the less in laryngitis, and inspection will reveal the seat of the inflammation.

How does parotiditis differ from laryngitis?

In parotiditis the voice remains unaffected. In laryngitis there is no swelling about the face.

How does tonsillitis differ from laryngitis?

Inspection discloses the swelling of the tonsils and the absence of involvement of the larynx.

In tonsillitis the voice may be nasal, but it is not hoarse or lost; cough and expectoration are absent, while the difficulty of breathing may be great.

The difficulty and pain of swallowing are greater in tonsillitis than in laryngitis.

How does hysterical aphonia differ from acute laryngitis?

Hysterical aphonia, as a rule, sets in more or less suddenly; often in association with emotional disturbance, in a person, usually a female, presenting other manifestations of hysteria; and is unattended with pain, difficulty of breathing or elevation of temperature. Laryngoscopic examination will reveal the motor impairment of the vocal bands, and the non-existence of an inflammatory process. Recovery may take place as suddenly as did the onset. The attack is likely to be repeated.

Edema of the Larynx.

What are the symptoms of acute edema of the larynx?

Acute edema of the larynx may occur so insidiously as to produce death without giving rise to appreciable symptoms, or the symptoms may be sudden and overwhelming.

When edema of the larynx occurs in the course of other affections, or as a sequel, there may be the usual prodromes of inflammatory fever, but as a rule the onset is sudden, and the severity of the attack rapidly increases. There are local tenderness, dryness and heat in the throat, with a sensation as of the presence of a foreign body; more or less ineffectual cough; muffling or extinction of voice; difficulty of swallowing; difficulty of inspiration, with harsh stridor, occurring in paroxysms that increase in frequency and severity, and unless relief be afforded, result in expiratory difficulty and apnea. *Respiration* is hurried and spasmodic; the *pulse* is small, frequent and irregular; the *temperature* is elevated; the *eyes* are prominent; the *face* is flushed and anxious, finally cyanotic.

When dependent upon acute disease, the suffocative attacks will be abrupt and violent, and will recur at intervals of a few hours.

When dependent upon chronic disease, the paroxysms may pass off, to recur irregularly, at progressively shorter intervals. Fever is absent, unless the underlying disease itself be febrile.

Simple inspection of the pharynx may reveal the swollen epiglottis projecting up behind the base of the tongue. *Palpation* with the finger may detect a soft, elastic, bladder-like swelling of the epiglottis, or of the ary-epiglottic folds; but unless cautiously practised it may induce a paroxysm of suffocation from the additional obstacle or irritation. *Laryngoscopic examination* at once reveals the cause of the symptoms.

What are the symptoms of chronic edema of the larynx?

Chronic edema of the larynx may be due to laryngeal disease, or to the usual causes of effusions. It gives rise eventually to progressive difficulty of breathing, paroxysmally aggravated, with perhaps some impairment of voice, and pain or difficulty in swallowing. The diagnosis is only to be made by laryngoscopy, in default of which palpation may perhaps be of service.

How is chronic edema of the larynx to be distinguished from asthma?

Principally by laryngoscopic examination. The paroxysms may exactly simulate asthmatic attacks. Suffocative paroxysms, due to laryngeal neoplasm, may likewise be mistaken for asthma if laryngoscopic examination be not made. Before the days of laryngoscopy many cases of sudden death were found to be due to laryngeal neoplasm.

Acute Tuberculous Laryngitis.

What are the characteristics of acute tuberculous laryngitis?

Acute tuberculous laryngitis is characterized by the *constitutional* symptoms of acute miliary tuberculosis, and *locally* by great pain in swallowing, by cough, dysphonia or hoarseness, and sometimes pain in respiration. *Laryngoscopic inspection* reveals peculiar thickenings of the epiglottis and other laryngeal structures, which are quickly followed by ulceration; or there may be ulceration from the first. Tubercle-bacilli are sometimes to be detected in the secretions or in the débris of ulcers.

Laryngeal manifestations are sometimes observed before evidences of pulmonary tuberculosis can be detected; but the latter, sooner or later, become manifest. A somewhat rapid fatal

termination is the rule, but recognized instances of recovery are multiplying.

Laryngismus Stridulus.

What is laryngismus stridulus?

Laryngismus stridulus is a spasmodic affection characterized by contraction of the constrictor muscles of the larynx, giving rise to dyspnea, inspiratory stridor and a ringing, croupy cough. It occurs most frequently in rachitic children. The attacks are paroxysmal and usually nocturnal. The child may be awakened from sleep with a sense of suffocation and an appearance of lividity. The paroxysm lasts for a few minutes and terminates with a deep inspiration, attended with a crowing sound. Occasionally death results in the paroxysm ; perhaps, from incarceration of the epiglottis.

How does laryngismus stridulus differ from asthma?

Laryngismus stridulus is a disease of childhood ; asthma is rare in children.

Rickets predisposes to laryngismus stridulus ; asthma may be apparently idiopathic, or some recognized irritation, direct or reflex, may be discoverable. The wheezing and râles of asthma differ from the cough and stridor of laryngismus.

The asthmatic paroxysm terminates in abundant expectoration ; laryngismus stridulus subsides without critical phenomenon.

Laryngeal Vertigo.

What is laryngeal vertigo?

The name *laryngeal vertigo* is applied to a group of symptoms of rare occurrence that may appear in connection with recognized nervous diseases, as posterior spinal sclerosis or epilepsy, or without obvious explanation. The symptoms are not always the same, but the essential elements in association are pain or spasm of the larynx, vertigo and perhaps syncope. The occurrence of laryngeal vertigo should lead to careful search for symptoms or signs of the causative disorder.

Catarrhal Croup—Spasmodic Croup.

What is catarrhal or spasmodic croup?

Catarrhal or *spasmodic croup* is essentially a catarrhal laryngitis, associated with a tendency to spasm of the constrictor muscles of the larynx, causing paroxysms of suffocation. It is almost exclusively a disease of children. It may *follow* exposure, may be simultaneous with the prevalence of epidemics of influenza, of measles or of scarlatina. It may *begin* insidiously with slight hoarseness, or suddenly, with chill. It usually attracts attention at first toward evening; the voice will be hoarse, with a slight cough. In the middle of the night, the child will suddenly awaken, with crying and paroxysms of suffocation, and with a peculiar, ringing, metallic cough, which is termed croupy. There is slight elevation of temperature. The spasm may quickly pass away, or it may be repeated several times during the night and for three or four successive nights, the symptoms of ordinary laryngeal or laryngo-bronchial catarrh being manifested during the day. *Recovery* may take place, or high fever may set in, with a firm, bounding pulse, flushing of the cheek, abnormal brilliance of the eye, and development of the obstructive symptoms of pseudo-membranous laryngitis.

How does catarrhal croup differ from laryngismus stridulus?

Laryngismus stridulus is a disorder of repeated occurrence, and comparatively chronic duration, occurring in rachitic children, and not associated with fever, cough or other symptom of inflammation. Spasmodic croup begins acutely, is of short duration, attacks healthy as well as rachitic children, and is associated with slight fever, cough and symptoms of catarrhal inflammation.

How are croup and edema of the larynx to be distinguished?

Apart from the revelations of laryngoscopy, croup is a disease of childhood principally—edema usually occurs in adults. In febrile cases the history of exposure to irritating fumes, or of other exciting cause of acute edema, the pain in swallowing, and the local tenderness; in other cases the absence of fever,

and the knowledge of the existence of causative conditions; together with the absence of the croupal cough, and the purely inspiratory character of the stridor, are additional discriminative features of edema.

Membranous Croup.

What are the symptoms of membranous croup?

Membranous croup or *pseudo-membranous laryngitis*, whether an idiopathic disease or due to the extension or original formation of diphtheritic membrane in the larynx and trachea, is manifested by symptoms of mechanical and spasmodic or paralytic obstruction of the larynx and trachea, associated with more or less fever and the constitutional symptoms of diphtheria, if that disease be present. In some instances, the palate, tonsils and pharynx are involved in simple, exudative or diphtheritic inflammation. In severe cases, the process extends into the bronchi, and in some cases there is pneumonitis. There is a peculiar ringing cough, which gradually becomes muffled, with progressive muffling or extinction of the voice; and great difficulty in breathing, intensified in paroxysms, which may progress to suffocation. Evidences of the severity of the obstruction are the recession upon inspiration of the soft tissues above and below the sternum and the use by the child of the auxiliary muscles of respiration, in its efforts to obtain breath. There is extreme restlessness and agitation, with paroxysmal exacerbations, attended by protrusion of the eyeballs, distention of the nostrils, flushing of the cheek, grasping at supports.

Sometimes a child will sit up straight in bed and clutch at its neck, as if trying to pull away some obstruction.

Suffocative paroxysms, being partly spasmodic, are sometimes relaxed by the effects of the resulting carbonic acid poisoning, and with the production of cyanosis there comes a deep sighing expiration, followed by a deep inspiration, and a period of comparative quiet.

Flakes of membrane or even membranous casts of the trachea and bronchi may be expectorated. The *duration* may be short, from twenty-four to forty-eight hours; it may be prolonged to

two or three weeks; usually, however, asphyxia or recovery occurs in from five to eight days. Convulsions often precede death.

How are membranous croup and retropharyngeal abscess to be distinguished?

Retropharyngeal abscess may be detected by inspection and palpation. The evidence of tuberculosis or syphilis, the history of previous infectious disease, may be suggestive of abscess. The voice is toneless but distinct in croup; in abscess it is nasal and indistinct. In abscess there are pain and difficulty in swallowing; these symptoms are absent from croup. In abscess there is stiffness of the neck, with tumefaction and pain on pressure externally; this is not the case in croup. In croup there is a peculiar cough and sometimes expectoration of false membrane; these are not present in abscess.

How do membranous croup and catarrhal croup differ?

In catarrhal croup fever and constitutional symptoms are slight. In membranous croup, fever is severe; and in diphtheria, toxemic symptoms are evident.

In catarrhal croup the obstruction to breathing is purely spasmodic, and disappears with relaxation of the spasm.

In membranous croup there is mechanical obstruction, continuing when the spasm has passed.

In catarrhal croup the voice remains clear, or at most becomes hoarse; the cough preserves its ringing, "croupal" character. In membranous croup voice and cough become toneless.

In membranous croup shreds of membrane or casts may be expectorated, and diphtheritic membrane is sometimes visible in the pharynx. Laryngoscopic inspection, when possible, will settle the diagnosis.

How is obstruction of the larynx, trachea or a bronchus by a foreign body to be distinguished from croup?

In the absence of the history, which would at once prevent error, the lack of fever and its concomitants, the absence of the peculiar cough, the variation in the symptoms, and finally laryngoscopic examination or the evidences of bronchial obstruction discovered on auscultation and percussion, permit the diagnosis of foreign body to be made.

Whooping-Cough—Pertussis.

What are the symptoms of whooping-cough?

Pertussis, or *whooping-cough*, is a contagious disease of childhood, attended with catarrhal symptoms, to which is added a peculiar cough, occurring in paroxysms and attended with a characteristic whoop. For several days there is coryza, accompanied with an acrid discharge, conjunctivitis, laryngitis and bronchitis. The cough then becomes paroxysmally explosive, a series of expiratory efforts being followed by a peculiar, ringing, inspiratory whoop. During these attacks the child may become livid and appear as if about to suffocate; sometimes vomiting is induced. The violence of the cough may cause ulceration of the under surface of the tongue on either side of the frenum; or there may be hemoptysis or epistaxis. Emphysema, interstitial or vesicular, may be induced. The paroxysms are repeated with variable frequency. The number of colorless blood-corpuscles, especially the lymphocytes, is increased. The disease may last many weeks, the severity and number of attacks progressively diminishing. It may be *complicated* by catarrhal pneumonia, which sometimes proves fatal. Bronchitis may long persist as a *sequel*. The period of *incubation* is from seven to ten days. The disease is supposed to be dependent upon a small bacillus with rounded extremities.

What are points of discrimination between membranous croup and whooping-cough?

In whooping-cough there is little or no fever, no continuous dyspnea, no alteration of voice, and the child is usually up and about, unless there is pulmonary complication. In croup there is not the characteristic whoop.

What are the differential features of laryngismus stridulus and whooping-cough?

Laryngismus stridulus is a paroxysmal neurosis, most common in rickety children; whooping-cough is a contagious disease that attacks all children alike.

The cough of laryngismus stridulus is croupy; that of whooping-cough occurs as a series of explosive expiratory sounds, fol-

lowed by a distinctive inspiratory whoop. During the intervals of freedom, whooping-cough presents the symptoms of bronchitis ; laryngismus stridulus does not.

How does tuberculosis of the bronchial glands differ from whooping-cough ?

Whooping-cough is a self-limited disease ; tuberculosis is not. Grave constitutional symptoms attend the latter ; recovery from whooping-cough is comparatively rapid. While the cough induced by tuberculosis of the bronchial glands is ringing, it does not possess the peculiar character of the cough of pertussis. Cases of whooping-cough do not occur isolated. By pressure on adjacent structures, enlarged bronchial glands may occasion dyspnea, cough and cyanosis, and edema of the face and neck.

Chronic Laryngitis.

What are the symptoms of chronic laryngitis ?

Chronic laryngitis may be a result of repeated attacks of acute laryngitis ; it may depend on nasal disease, gastric catarrh, dilated heart, constipation, or kidney disease, or other near or remote affection. It is common in those who use their voices improperly or immoderately, and in those who smoke to excess. It is characterized by hoarseness, persistent, irritable cough and discomfort in phonation and deglutition. Laryngoscopically the structures within the larynx are seen to be thickened, congested, covered by tough secretion ; the vocal bands have lost their pearly, glistening appearance, and their motility may be impaired. In many cases chronic laryngitis is of tuberculous or of syphilitic origin. In such cases there may be characteristic ulceration, and in the case of tuberculosis, tumefaction of the epiglottis and aryteno-epiglottic folds ; vegetations in the meso-arytenoid fold are quite common.

How is chronic laryngitis to be distinguished from laryngeal neoplasms ?

The only safe method of diagnosis is by means of laryngoscopy, and its neglect is criminal.

Chronic Tuberculosis of the Larynx.

What are the characteristics of chronic tuberculosis of the larynx?

In case of *chronic tuberculosis of the larynx* the subjective and objective symptoms may be simply those of ordinary chronic laryngitis, or there may be characteristic tumefactions, ulcerations or vegetations in the larynx, or all combined; with more or less pain and difficulty in swallowing and in respiration, and alteration of the voice. The signs of pulmonary tuberculosis are usually well advanced. Cases of recovery from the local disease in the larynx, or of its prolonged abeyance, are not rare.

How are paralyses, ulcerations, abscess, stricture and other conditions affecting the larynx and trachea, and giving rise to alterations in voice and difficulty in respiration, to be diagnosticated?

The only reliable method is laryngoscopy.

PHYSICAL DIAGNOSIS.

What are the methods of physical exploration?

In a *physical exploration*, systematically conducted, five means of investigation are employed: *inspection*, *mensuration*, *palpation*, *percussion* and *auscultation*.

What does physical exploration teach?

By physical exploration information is gained of physical or mechanical conditions, so far as these may influence the size, contour, movement, temperature, density, elasticity and acoustic relations of the structures examined. The nature of pathologic conditions cannot be directly determined from physical signs, but is to be inferred from the latter in association with all other phenomena in a given case.

What is to be learned by inspection?

By *inspection* the general appearance of an individual is noted—the height, the apparent relative nutrition, the color,

the configuration, the movement, the state of the pupils, the expression.

Examining the chest more especially, its symmetry or asymmetry, its fulness or retraction; the frequency and character of the respiratory movements; the extent of the respiratory excursion; the situation, extent and vigor of the cardiac impulse are observed, and abnormal manifestations are looked for.

The chest is *distended* symmetrically in emphysema and usually in case of hydrothorax; pleural effusions and pneumothorax commonly occasion unilateral bulging.

The chest is *retracted* when the parietal and visceral pleuræ are adherent, and when the lung is shrunken, as in interstitial pneumonitis. The retraction is well marked after some forms of operative treatment of empyema, and when the lung is collapsed from any cause.

A healthy adult breathes from eighteen to twenty times in the minute. The normal respiratory excursion is curtailed and the frequency of respiratory action is accelerated by almost all diseases of the lung and pleura and also in case of peritonitis. The respiratory frequency is also accelerated when the pulse is quickened, as in febrile conditions. In case of pleurisy, the patient lies on the affected side, so as to reduce to a minimum functional movement of that side. Rapidity of breathing may be hysterical. Women breathe mostly with the upper portion of the chest; men with the lower.

What is to be learned by mensuration?

Mensuration determines with precision what inspection does approximately: the size, configuration, symmetry or asymmetry and respiratory excursion of the chest.

What is to be learned by palpation?

By *touch* or *palpation* one distinguishes elasticity from rigidity and resistance, investigates the existence of pain or tenderness, of edema, of moisture or dryness, of heat or cold. When the palm of the hand is applied to the chest of a speaking person a diffuse, vibratory sensation is perceived, usually slightly greater in intensity on the right side than on the left. This is known as the *vocal* or *tactile fremitus.*

It is increased in conditions of condensation of the lung, and diminished or lost in the presence of thickening and adhesion of the surfaces of the pleura and of collections of fluid in the pleural cavity. Variations in vocal fremitus correspond pretty closely with variations in vocal resonance. Sometimes a pleuretic friction-rub can be felt.

How is percussion performed?

Percussion may be *immediate* or *mediate*. In the former a sharp blow is struck with the tips of the fingers bunched together, or with the palmar surface of the extended fingers.

Mediate percussion is performed by means of a thin, flat plate of ivory or of hard rubber (called a plexímeter), applied over the part to be examined, and a small rubber-tipped hammer (called a plexor).

For many purposes it is preferable to use the extended finger of one hand as a plexímeter, and the index or middle finger (or both) of the other hand as a plexor. Percussion is said to be *weak* or *strong*, *superficial* or *deep*, according to the energy of the blow of the plexor.

What is to be learned by percussion?

Percussion gives information as to the relative distribution of gases (usually air), fluids and solids in the structures examined. Attention is paid to the *quality* (or *timbre*), the *pitch* and the *intensity* of sounds.

Percussion of the healthy chest elicits a sound called *clear*, representing the *normal*, *pulmonary*, *vesicular resonance*. This may be *impaired* by increased density of the pulmonary tissues or of superjacent structures—as in interstitial pneumonitis, in pleural thickening or when the chest-wall is thickened.

Percussion-dulness is dependent upon a high degree of condensation—as in the solidification of pneumonia, or at the late stage of tubercle-formation.

The sound elicited over solid viscera, as the liver, heart or spleen, or over serous effusions—as in hydrothorax or hydropericardium, or in ascites, is *flat*.

The sound occasioned by air (or gas) in inclosed spaces larger than the alveoli of the normal lung may be *hyper-resonant* or *tympanitic*—as in vesicular emphysema or over the intestines.

An *amphoric* or *metallic* sound is elicited by percussion over large closed cavities, with tense walls, containing air—as a distended stomach or a large vomica in the lungs.

A *cracked-pot* or *cracked-metal* sound is occasioned by percussing over a cavity of some size, with a small opening, through which air escapes—as in the case of a pulmonary cavity communicating with a bronchial tube.

The percussion-sounds may display alterations in *degree* or *pitch*, and in *intensity* or *volume*.

Increased density gives heightened pitch—hence a dull sound is higher in pitch than a clear one. The pitch of tympanitic sounds varies. As a rule, it is higher than that of the normal pulmonary resonance. Other things being equal, the greater the volume of matter set in vibration the greater the intensity of the sound.

By mediate percussion with the fingers a sense of *elasticity*, or of *resistance*, can be appreciated.

Sometimes auscultation is practised by one while a second at the same time practises percussion—so-called *auscultatory percussion*. With a double stethoscope or phonendoscope one may practise auscultatory percussion without assistance. Some prefer simply to tap lightly over the part to be examined, gradually receding from or approaching to the point of auscultation.

The percussion sounds elicited during full-held inspiration differ from those elicited during full expiration—so-called *respiratory percussion*.

How is auscultation practised?

Auscultation, like percussion, may be *immediate* or *mediate*. In the former, the ear is directly applied to the part to be auscultated; in the latter, auscultation is performed through the mediation of a stethoscope. Stethoscopes are monaural or binaural. Each has its advantages. The phonendoscope also may be employed. It is especially useful for auscultatory percussion. The student should become familiar with all of the methods of auscultation.

What is to be learned by auscultation of the lungs?

Auscultation gives information as to the movement of air and

fluids, as to the comparative calibers and lengths of the tubes through which the air passes, and as to the presence in the path of the air-current of matters capable of acting the part of reeds in the production of musical tones.

Listening to the normal respiratory sounds one hears a soft, breezy inspiration—the *normal vesicular murmur*, followed by a scarcely audible, briefer expiration.

These sounds may be *exaggerated*—as they normally are in children; hence they are then called *puerile*.

The respiratory sounds are also *intensified* in a lung, or in a portion of lung, performing an excess of function—as after violent exercise, or, compensatorily, when the function of the other lung, or of a portion of the same lung, is interfered with by condensation or compression.

The respiratory murmur is *enfeebled* or wanting when there is an obstruction to the circulation of air in the lung—as in incipient pulmonary tuberculosis, and in occlusion of the air-passages—as by intrathoracic aneurism, or when the air vesicles have lost their elasticity—as in emphysema.

The respiratory murmur is rendered *harsh* and is *heightened in pitch* when the bronchial tubes are thickened, and their caliber is narrowed—as in bronchitis. Under similar conditions, *rhonchi* or *dry râles* may be heard—*sibilant*, if generated in the small tubes; *sonorous*, if generated in the large tubes. Sometimes the sounds are *high-pitched* and *wheezing*. *Isolated sibilant râles* are usually indicative of pulmonary tuberculosis.

If the vesicular murmur is but partially obliterated, the breathing is called *vesiculo-bronchial*; or if the bronchial element predominates, *broncho-vesicular*. When occlusion of the alveoli entirely obliterates the vesicular quality, the breathing is said to be *bronchial* or *tubular* or *blowing;* these terms respectively indicating progressive encroachment upon the integrity of the smaller air-tubes. In case of a cavity in the lung, the breathing may be *cavernous;* if the cavity have a tense wall and a small orifice of communication with a bronchus, the sound transmitted to the ear is *amphoric*.

The presence of secretion in the bronchial tubes occasions

râles, usually qualified as *moist: subcrepitant*, if in the smaller tubes, *mucous* if in the larger.

The presence of fluid in a vomica may give rise to *bubbling* or *gurgling*.

The separation of the adherent surfaces of pulmonary alveoli lined by viscid secretion, as in pneumonia, gives rise to the *crepitant râle*. When isolated softening of pulmonary tubercles takes place, a sound is generated comparable to that produced when salt is thrown upon fire, or when a few hairs are rubbed together between the fingers—so-called *crackling*.

When moist râles are heard through solidified pulmonary tissues they are transmitted with a peculiar clearness and with a sort of ringing character. They are then called *consonating*.

In the first stage of pleurisy, before effusion has taken place, and in the third stage, after the fluid poured out has been absorbed, a *coarse crackling*, or *creaking*, or *rasping* sound of pleural *friction* may be heard.

Metallic tinkling, resembling the splashing of water in a pool, may be heard when, in the presence of a collection of air in a cavity large enough to act as a resonating chamber, bubbles of air escape through fluid, fluid drops into fluid, or air enters through a valve-like orifice. These conditions are fulfilled in *pneumothorax, hydropneumothorax*, and *vomicæ of large size*, especially those containing a moderate quantity of fluid. When air and fluid are both present, shaking of the patient, with the ear applied to the chest, may elicit the sound of splashing—the so-called Hippocratic *succussion sound*.

When the ear is applied to the normal chest of a speaking individual a confused sound is heard—the *vocal resonance*.

If fluid is effused into the pleural cavity the transmission of the voice is intercepted below the upper level of the fluid. If the pulmonary tissues are solidified, as in pneumonia and tuberculosis, the resonance is increased, constituting *bronchophony*. When spoken language and whispers can be distinguished, the phenomena are termed *pectoriloquy* and *whispering pectoriloquy* respectively. The latter, if circumscribed, is usually indicative of the existence of a cavity in the lung.

When, in the course of pleurisy, a small quantity of fluid has

been poured out the voice is transmitted above the level of the fluid with a peculiar bleating character, constituting *egophony*.

Not only may the character of the breath-sounds be altered, but their *rhythm* may deviate from the normal. Thus, expiration may be *prolonged*, when, as in emphysema, the elasticity of the walls of the alveoli is diminished or when the alveoli are compressed by infiltration around them, as in incipient tuberculosis; or the expiratory sound may be *prolonged* or *jerking*, when, as in the early stages of pulmonary tuberculosis, there is some obstruction to the exit of air.

Acute Pleurisy.

What are the symptoms of acute pleurisy?

Acute pleurisy may set in with a chill and sharp pain in the side, aggravated by the respiratory movements. The dyspnea may be slight or considerable. The *temperature* rises moderately high; the *breathing* is feeble, shallow and rapid. There is slight irritative cough, and scanty, frothy expectoration. In the course of a few days the symptoms subside, and at the end of a week or ten days the patient is well. Acute pleurisy may occur as a primary condition. It may follow traumatism or exposure to cold. Quite commonly it is secondary to inflammation of adjacent structures, especially of the lungs. It also occurs in the course of various infectious diseases, of nephritis and of rheumatism. Pleurisy may be "dry" or attended with effusion. The exudation may be serous, sero-fibrinous, purulent or hemorrhagic. Effusion gives rise to respiratory and circulatory embarrassment proportionate to the volume of fluid poured out. It may be encysted.

The fluid poured out may not be readily absorbed. On the contrary, it may remain obstinately persistent. In the course of time, it may become purulent; under other circumstances, it is purulent from the outset; in either case the condition is known as *empyema*. More commonly, the fluid is absorbed and the two layers of the pleura become adherent, with obliteration of the pleural cavity; in the progress of the case considerable thickening takes place, followed in turn by retraction of the

chest. When the effusion is purulent there are repeated rigors, fever, sweats, leukocytosis, edema of the chest and emaciation; otherwise the health may be preserved, unless the chronic pleurisy is tuberculous. By the action of the heart, pulsation may be imparted to a collection of fluid in the left pleural cavity, so that an aneurism may be simulated. When the collection is purulent, the condition is designated *pulsatory empyema*. Pleurisy may be confined to the diaphragmatic layer or to the folds between the lobes of the lungs.

What are the physical signs of acute pleurisy?

For convenience of study, the physical signs of acute pleurisy may be considered in *three stages*. In the *first*, or *plastic stage*, as the surfaces of the two layers of the pleura, roughened by exudation, slide over one another in inspiration and expiration, a harsh, creaking sound is heard, the vibration occasioning which may sometimes also be appreciable on palpation—the so-called *friction-sound* or *rub*. The *breathing* is shallow, feeble and rapid; the movements of the affected side being restrained in the greater degree in order to mitigate the pain.

In the *second stage*, or *stage of effusion*, serum in variable quantity, possibly mixed with blood, is poured out into the pleural cavity. The corresponding lung is pushed upward and backward, and the heart, liver, spleen and diaphragm may be displaced. One side of the chest appears larger than the other, and the interspaces corresponding to the situation of the effusion are abnormally prominent. The *cardiac impulse* is visibly displaced. The *respiratory excursion* is limited. Below the upper level of the fluid the *percussion-note* is flat; above, it is subtympanitic. Through the effusion the *breath-sounds* are heard feebly and indistinctly; the *voice-vibrations* are not transmitted to the palpating hand; nor, as a rule, to the auscultating ear, though exceptionally bronchophony may exist. To the ear applied to the chest above the level of the fluid the *voice* may be transmitted with a peculiar bleating quality—constituting egophony. Exploratory puncture may be necessary to determine the presence of fluid and its character.

In the *third stage*, the effusion has been absorbed and the apposed layers of pleura again come in contact. The friction-

sound of the first stage returns. Ultimately the pleura may be restored to its original condition or its apposed surfaces may become adherent and thickened, giving rise to retraction of the chest and percussion-dulness.

How does acute pleurisy differ from croupous pneumonia?

Pleurisy usually accompanies pneumonia. The signs of the former, however, should not be permitted to obscure the existence of the latter. Neither the local nor the general symptoms of pleurisy are so profound as those of pneumonia. While pneumonia may then present the symptoms of pleurisy, pleurisy does not present the blood-streaked expectoration, the crepitant râle, the blowing breathing, the deficiency of the chlorides in the urine, or the critical termination of pneumonia.

The percussion-note of pneumonia is dull; that of pleurisy, when an effusion exists, is flat. The dulness of pneumonia is usually over a lower lobe; that of pleurisy is universally at the base. In pleurisy, the breath-sounds are heard feebly, or not at all, through the effusion; in pneumonia the breathing is bronchial. Vocal resonance and fremitus are increased in pneumonia; they are diminished or absent in pleural effusion; above the fluid, however, a bleating sound is transmitted to the auscultating ear. Simple, uncomplicated pneumonia is unattended with friction-sounds and occasions no displacement of adjacent viscera.

How are intercostal neuralgia and acute pleurisy to be differentiated?

The pain of intercostal neuralgia may simulate that of acute pleurisy, and give rise to rapid, shallow, feeble respiratory movements. Intercostal neuralgia is paroxysmal and unattended with friction-sound, dulness on percussion or fever; it occurs in anemic individuals with neurotic tendencies, and may be attended with a unilateral herpetic eruption in the course of the nerve affected. In addition there are a number of tender points.

How are a pleuritic effusion and an hydatid cyst of the liver to be differentiated?

When an hydatid cyst of the liver attains proportions suf-

ficient to give rise to definite physical phenomena, these will appear in a region beyond that in which the signs of a pleuritic effusion on the right side are found. The history of an acute attack is wanting, while it is present in pleuritic effusion. Neither in case of pleuritic effusion, nor in case of hydatid of the liver, is the percussion-dulness confined to the right hypochondrium ; in the one it extends rather upwards, in the other downwards ; in the latter it is associated with fluctuation. In a case of hydatid cyst it may by percussion be possible to elicit the characteristic thrill or fremitus. In a case of pleuritic effusion the breath-sounds and the vibrations of the voice are feebly transmitted ; breath-sounds, vocal resonance, and fremitus are unaltered in hydatid cyst of the liver. Egophony is characteristic of the presence of fluid in the pleura ; it is thus not present in case of hydatid cyst of the liver. Exploratory puncture may determine the presence of hydatid hooklets, the detection of which places the diagnosis beyond doubt.

How are a pleuritic effusion and abscess of the liver to be differentiated ?

Symptoms of respiratory interference are naturally less conspicuous in case of abscess of the liver than in case of pleuritic effusion. Hepatic abscess may occasion tumefaction in the right hypochondrium ; pleural effusion renders the chest asymmetrical from unilateral bulging. Pleuritic effusion impairs the transmission of the breath-sounds and of the vibrations of the voice, which is unaltered by an hepatic abscess. Egophony may be elicited in case of pleuritic effusion, but not in case of hepatic abscess. Rigors commonly attend an abscess of the liver ; they only take place in cases of pleuritic effusion when suppuration has occurred. In case of hepatic abscess, there is usually a history of gallstone, of ulceration of the bowel or of pyemia ; in case of pleuritic effusion, there is a history of an acute attack of pleurisy. Hepatic abscess and perihepatitis may, however, give rise to pleural effusion by extension.

How are pleuritic effusion and abscess of the spleen to be differentiated ?

Abscess of the spleen is most common as a manifestation of py-

emia; pleuritic effusion is a sequel of an acute pleurisy. The symptoms of respiratory derangement, a necessary part of pleuritic effusion, are subordinate in case of splenic abscess. Repeated rigors occur in abscess of the spleen; rigors occur in case of empyema, but not when the effusion is not purulent. Egophony may be developed when fluid is present in the pleural cavity, but not in case of splenic abscess.

How is an intra-thoracic tumor to be distinguished from chronic pleurisy?

Tumors in the chest may arise from the pleura or from the lung; they may develop in the mediastinum; or they may be aneurismal. Malignant tumors are secondary to growths elsewhere—an element in diagnosis. They give rise to circumscribed areas of dulness on percussion, not necessarily limited to one side; to enfeebled breathing and perhaps friction-sounds; in their physical signs, they more closely resemble encysted pleurisy than ordinary chronic pleurisy. When actually complicated by pleurisy, the diagnosis may not be possible. Aneurisms occur in the course of the aorta and large vessels, and, in addition to the evidences of tumor, are accompanied by thrill, bruit and pulsation.

How are a pleuritic effusion and hydrothorax to be differentiated?

Hydrothorax or passive pleural effusion occurs as a result of cardiac insufficiency or as a part of a general dropsy, from nephritis, for instance.

The effusion of hydrothorax is usually bilateral; that of pleurisy is almost invariably unilateral. In the latter there has probably been antecedent pain and friction-sounds and also fever; in the former there are other evidences of cardiac failure or of a general dropsy. The fluid obtained on exploratory puncture in case of inflammatory effusion is usually richer in salts and albumin than the simple serous effusion of non-inflammatory origin.

How is a pericardial effusion to be distinguished from a pleural effusion?

A pericardial effusion occurs under circumstances similar to

those that give rise to a pleural effusion—as a result of inflammation or as a part of a general dropsy. The position and outline of the percussion-dulness to which a pericardial effusion gives rise, however, are entirely different from what is found in a case of pleural effusion ; nor are the breath-sounds notably interfered with, while the circulation is embarrassed and the heart-sounds are almost obliterated at the cardiac apex.

Chronic Pleurisy.

What are the clinical features of chronic pleurisy?

The exudation of acute pleurisy may not be absorbed, and, even without acute symptoms, adhesions may form between the two layers of pleura, with secondary thickening. As a result one side of the chest may undergo retraction and flattening, while the percussion-resonance is impaired, the breath-sounds distant or enfeebled, the vocal resonance and fremitus diminished. Breathing may be interfered with, the heart displaced and health may suffer in varying degree. Sometimes the extremities of the fingers and toes become enlarged and bulbous and the finger-nails unduly curved and beak-shaped.

Acute Bronchitis.

What are the symptoms of acute bronchitis?

Acute bronchitis results from exposure to cold, the inhalation of irritating fumes, or as a secondary disorder in the course of fevers, rheumatism or heart-disease. There is irritative cough, and, at first, scanty mucous expectoration, which subsequently becomes more copious and muco-purulent; slight elevation of temperature ; increased frequency of respiration ; some dyspnea ; retro-sternal pain, and mild constitutional symptoms.

What are the physical signs of acute bronchitis?

The *chest* is not deformed and expands well. The *percussion-resonance* is vesicular. The *breathing* is harsh at first ; dry râles,

sonorous and sibilant, are heard ; subsequently large and small moist râles. *Vocal resonance* and *fremitus* are not perceptibly altered.

How does acute bronchitis differ from acute miliary tuberculosis?

In acute miliary tuberculosis the dyspnea is greater, the temperature is higher, with greater oscillations, the breathing is more rapid, and the symptoms are more profound than in acute bronchitis. The further progress of the case clears up any possible doubt. Recovery from acute bronchitis is the invariable rule. Pulmonary consolidation and softening, percussion-dulness and fine moist râles, emaciation, hectic fever, gradual failure of the vital powers and ultimately death mark the usual course of acute miliary tuberculosis.

Chronic Bronchitis.

What are the symptoms of chronic bronchitis?

Chronic bronchitis is usually a result of repeated attacks of acute bronchitis ; it may manifest itself as a special susceptibility to acute bronchitis ; at first it appears as a winter cough, subsequently becoming continuous. It may obstinately resist treatment ; it is attended with a good deal of cough, copious muco-purulent expectoration, marked shortness of breath, and may in time give rise to emphysema, or to bronchiectasis. It is often attended with loss of flesh and strength.

What are the physical signs of chronic bronchitis?

If emphysema coexists the *chest* may be enlarged ; otherwise it is not abnormal in size or form ; the *respiratory excursion* is somewhat diminished ; the *percussion-resonance* is little or not at all impaired ; the *breathing* is harsh and may be feeble, the bronchial element preponderating. Coarse râles, moist and dry, are heard at all parts of the chest. *Vocal fremitus* and *resonance* are rather increased.

How is chronic bronchitis to be distinguished from interstitial pneumonitis?

A certain degree of bronchitis commonly attends interstitial pneumonitis; but the chest is likely to undergo retraction, which is not the case in bronchitis. If impairment of resonance attend chronic bronchitis, it is general and not well defined, while the dulness of interstitial pneumonitis is the more circumscribed and the more decided.

How does chronic bronchitis differ from pulmonary tuberculosis?

While chronic bronchitis may be attended with obstinate cough, muco-purulent, sometimes blood-streaked, expectoration, loss of flesh and strength, the physical signs are general in their distribution and not localized as in tuberculosis. The impairment of resonance is not so great in chronic bronchitis as it is in tuberculosis. The elevation of temperature observed in tuberculosis is wanting in bronchitis. Tubercle-bacilli are not found in the sputum of cases of simple chronic bronchitis.

Plastic Bronchitis—Fibrinous Bronchitis.

What are the characteristics of plastic or fibrinous bronchitis?

In addition to the phenomena of ordinary bronchitis, there are present in *plastic bronchitis* decided dyspnea and cyanosis, together with the expectoration of tough, fibrinous casts of the smaller bronchial tubes; there occur lancinating pains in the chest; and there may be bleeding from the nose and mouth.

Putrid Bronchitis.

What are the clinical features of putrid bronchitis?

In case of bronchitis with bronchial dilatation, accumulation of secretion may take place, with ulceration and inflammation of the bronchial mucous membrane, as a consequence of which expectoration is augmented, the breath and sputa possessing an offensive, fetid odor. To these phenomena fever and a typhoid condition may be added—and even pulmonary gangrene may supervene.

Bronchiectasis.

What is bronchiectasis?

Bronchiectasis consists in a cylindrical or saccular dilatation of the bronchial tubes, usually developed in the course of some condition attended with powerful or sustained expiratory efforts—such as chronic bronchitis. It may also result from contraction of the peribronchial tissues. In addition to the symptoms of the causative affection, the characteristic feature of bronchiectasis is the periodical occurrence of paroxysms of cough, attended with the expectoration of large quantities of muco-purulent secretion of offensive odor. The tubes filled with fluid may give rise in small areas of irregular distribution to *percussion-dulness* that disappears with the evacuation of the fluid. On *auscultation* large, coarse, moist râles and gurgling may be heard. If of moderately large size, bronchiectatic cavities may yield a hyper-resonant percussion-note, with blowing breathing and whispering pectoriloquy.

How does bronchiectasis differ from pulmonary gangrene?

The grave constitutional symptoms of gangrene are lacking in bronchiectasis. The expectoration of both is offensive, but that in gangrene is actually fetid. The sputum of gangrene contains shreds of pulmonary tissue; that of bronchiectasis does not.

How does bronchiectasis differ from pulmonary abscess?

A sacculated dilatation of a bronchial tube containing fluid fulfils the physical conditions of an abscess communicating with a bronchial tube. The expectoration of a case of bronchial dilatation, however, has an offensive odor not present in abscess. The fever and sweats of abscess are wanting in bronchiectasis. Abscess is usually a sequel or complication of pulmonary inflammation or some general pyemic process.

How are bronchiectasis and catarrhal pneumonia to be differentiated?

In the small areas of dulness of irregular distribution, bronchiectasis may simulate catarrhal pneumonia, but the former is wanting in the febrile phenomena and the constitutional depression of the latter disease. Catarrhal pneumonia is a disease

of acute onset, though sometimes of protracted duration; while bronchiectasis is essentially a chronic disease. The one is most common in childhood, the other later in life. The offensive sputa of bronchiectasis are entirely wanting in catarrhal pneumonia.

Capillary Bronchitis.

What are the symptoms of capillary bronchitis?

Capillary bronchitis, sometimes called *suffocative catarrh*, is an inflammation of the smallest or "capillary" bronchi, most common in children and in old persons, and usually secondary to bronchitis of the larger tubes. In addition to the symptoms of the latter affection, capillary bronchitis is attended with decided constitutional depression; the dyspnea is more profound, and cyanosis may be marked.

Cough is severe, while expectoration is scanty; the *pulse* is rapid, the *respirations* hurried, the *countenance* discolored, the expression anxious; fever is moderately high.

What are the physical signs of capillary bronchitis?

The *chest* is symmetrical. The *breathing* is rapid and shallow. The *percussion-resonance* is vesicular, though in spots it may be slightly impaired. When it exists, the impairment of resonance does not depend upon exudation, but upon atelectasis. The phenomenon may, therefore, be inconstant, and its location shifting. The *breath-sounds* are harsh, and in many places in both lungs small, mucous râles may be heard.

Vocal resonance and *fremitus* are slightly increased. The deviations from the normal are most marked at the bases of the lungs posteriorly.

How are capillary bronchitis and acute miliary tuberculosis to be differentiated?

Capillary bronchitis is a quite brief disease of the very young and the very old; acute miliary tuberculosis is a more prolonged disease of the young and of adults. Dyspnea, cyanosis and constitutional depression are early more decided in capillary bronchitis than in acute miliary tuberculosis. The fever of tuberculosis is more intense and vacillating than that of bronchitis. Localized areas of persistent percussion-dulness, followed

by the moist râles of softening, may develop in the course of acute miliary tuberculosis, which is progressively fatal; recovery from capillary bronchitis is not uncommon.

Catarrhal Pneumonia—Broncho-pneumonia.

What are the symptoms of catarrhal pneumonia?

Catarrhal pneumonia, also called *lobular pneumonia*, or *broncho-pneumonitis*, may be apparently primary, but is often secondary to capillary bronchitis, as also to measles, influenza and other constitutional affections; it may result from the inspiration of infectious material, whether contained in the food or from septic surfaces in the respiratory tract.

It is attended with cough, muco-purulent expectoration, dyspnea, increased frequency of respiration, decided febrile phenomena that may be hectic, and constitutional depression. The course of the disease may be protracted.

What are the physical signs of catarrhal pneumonia?

The *chest* is symmetrical; *respiration* is shallow.

The *range of movement* is sometimes curtailed on one side.

Small, irregularly distributed patches of *percussion-dulness* may be found here and there over both sides of the chest, anteriorly and posteriorly; the *breathing* is harsh; fine and large moist râles may be heard; the *vocal resonance* and *fremitus* are intensified in patches corresponding, as a rule, to the areas of percussion-dulness.

How does catarrhal pneumonia differ from capillary bronchitis?

The gravity of the symptoms in both depends upon the extent of the disease. Other things being equal, capillary bronchitis is the more acutely dangerous affection, the depression sometimes being profound. The duration of catarrhal pneumonia is the longer. Its febrile phenomena are the more decided, and its temperature-range is marked by great oscillations. The impairment of percussion-resonance is more decided in catarrhal pneumonia than in capillary bronchitis; it is constant, and does not change its seat.

What are the means of distinguishing atelectasis from catarrhal pneumonia?

Atelectasis is the pre-natal condition of the lungs, before they have been inflated with air. It may also arise in the course of affections attended with inspiratory obstruction of the bronchial tubes, as in whooping-cough or capillary bronchitis, for instance, a plug of mucus acting as a ball-valve, allowing the expiration of air but preventing inspiration. It is said also to occur in parts of the lung functionally inactive. The condition is not an inflammatory one and is not necessarily attended with elevation of temperature, as is catarrhal pneumonia. The collapse is not persistent, but may successively involve different parts of the lung; so that the physical signs are not constant, but transitory as well as migratory.

How does catarrhal pneumonia differ from pulmonary infarction?

Catarrhal pneumonia does not set in with the acuteness and the severe pain of pulmonary infarction; nor is the expectoration as persistently blood-streaked in the former as in the latter. Catarrhal pneumonia is attended with decided febrile manifestations and is of considerable duration, while pulmonary infarction scarcely occasions fever and its symptoms soon subside. The detection of a source of an embolus would be strongly in favor of the existence of infarction.

Acute Croupous Pneumonia.

What are the symptoms of croupous pneumonia?

Croupous pneumonia, also called *lobar pneumonia*, and at one time *lung fever*, is an acute, infectious disease dependent upon the invasion of microörganisms, of which the most common is the encapsulated lanceolate diplococcus. (Fig. 22.) It is the most widely distributed and the most fatal acute disease. The susceptibility and the mortality increase progressively after the fifteenth year. Males suffer in greater number than females, and the disease is more common in the city than in the country. Intemperance and other debilitating influences are pre-

disposing causes. The disease often follows exposure to cold, *setting in* suddenly with a chill, sometimes with convulsions, the *temperature* rising at once to 103° or 104° F., the *respiratory frequency* reaching 35, 40, 50 or more, while the *pulse* is not proportionately accelerated.

There is at first slight cough, with scanty expectoration of a viscid sputum, which soon becomes blood-streaked (rusty), and usually contains the causative microörganisms.

There is considerable dyspnea; the alæ nasi are retracted in breathing; and one or both cheeks are flushed. As a rule the number of *colorless blood-corpuscles* is increased until the crisis occurs.

The cough increases; the expectoration becomes more free, more hemorrhagic, less viscid, and presents the appearance of prune-juice. There is pain in the chest, often referred to the nipple of the affected side; usually involving the lateral or axillary region in addition. The pain is increased on attempting to make prolonged inspiration. The *lips* and *cheeks* may be cya-

FIG. 22.



Micrococci of croupous pneumonia. (Vierordt.)

notic. The *tongue* is coated; the *appetite* is lost; *thirst* is increased; there may be jaundice, and, at the height of the attack, delirium.

Between the second and fifth days, herpes of the lips may appear. The symptoms continue for from five to seven or nine days, to subside by *crisis*. Profuse perspiration occurs; the temperature declines; and convalescence sets in.

The cough is still free; the expectoration copious and muco-purulent. During the attack the urine is deficient or wanting in chlorides, which return as resolution proceeds. Headache, delirium and other toxic symptoms may be present. Resolution may fail to take place or it may be but partial; so that suppuration or gangrene may ensue or chronic induration be a sequel. Death happens in a variable proportion of cases, not rarely from heart-clot or heart-failure. In the aged and in alcoholics the disease may be insidious in onset and latent in course.

Pleurisy is a common accompaniment of pneumonia. Occasionally large effusions are poured into the pleural cavity. Pericarditis, endocarditis and meningitis may be complications.

What are the physical signs of croupous pneumonia?

For convenience of description, pneumonia is divided into three stages: a stage of congestion, a stage of exudation, a stage of resolution.

What are the physical signs of the first stage of pneumonia, or the stage of congestion?

The aspect of the *chest* is unchanged.

The *breathing* is accelerated, becoming shallow and labored.

The *percussion-resonance* is little or not at all impaired.

The *respiratory murmur* is roughened over a circumscribed area, usually corresponding to a lower lobe. At the end of inspiration a fine, moist, crepitating râle may be heard.

Vocal resonance and *fremitus* are not perceptibly altered.

What are the physical signs of the second stage of pneumonia, or the stage of exudation?

The *respiratory excursion* is curtailed; the *breathing* is hurried.

The lower part of the *chest* on one side, corresponding to the lower or middle lobe of the lung, may be a little fuller than elsewhere. Here the *percussion-note* is dull. Above, the note is hyper-resonant. The right lung appears to suffer more commonly than the left; not rarely both become involved. Sometimes the upper lobe is attacked, particularly in children. When the disease is deep-seated the physical signs may be ill-defined.

At the dull area, the *breathing* is bronchial in character. At other parts of the chest the breathing is puerile. The crepitant râle has disappeared. Small mucous râles may be heard.

Vocal resonance and *fremitus* are increased at the area of percussion-dulness. A pleuritic friction-sound may be heard and the signs of effusion may develop.

What are the physical signs of the third stage of pneumonia, or the stage of resolution?

The localized fulness of the *chest* may remain.

The *respiratory excursion* and the *expansion* have somewhat augmented.

Some degree of impairment of the *percussion-resonance* persists.

The *breathing* is yet bronchial, though not in the same degree as it was. A small subcrepitant râle, termed the *crepitus redux*, can now be heard; other moist râles also become plainly discernible.

Vocal resonance and *fremitus* are still increased.

How is edema of the lungs to be distinguished from pneumonia?

Pulmonary edema may develop in the course of heart-disease, if the compensation be disturbed; in the course of nephritis or in the course of general asthenia; in subjects of these affections, it may follow exposure. It is not confined to one lung, but involves both alike throughout their entire extent.

Pulmonary edema may be uncomplicated, but is likely to be associated with effusions into the serous cavities and dropsy elsewhere. The sputum is frothy and abundant, but not rusty or like prune-juice. The cheeks are flushed in pneumonia; the face is pale or livid in pulmonary edema. Edema usually does not present febrile symptoms; the percussion-resonance scarcely suffers; large and small moist râles are heard all over the chest; the proportion of chlorides in the urine remains unaltered; while the brief duration of the condition, speedily terminating, as it does, in recovery or in death, and the knowledge of the existence of a condition that may occasion it, make the diagnosis clear.

In what respects do catarrhal and croupous pneumonia differ?

Croupous pneumonia is sudden in onset; catarrhal pneumonia is rather commonly secondary to some other condition. Lobar pneumonia usually attacks robust adults; broncho-pneumonia, children, the aged, the debilitated. The sputum of catarrhal pneumonia is not often blood-streaked, as is usually that of croupous pneumonia. Croupous pneumonia terminates in the course of seven or nine days by crisis; catarrhal pneumonia is a disease of long duration, with tardy convalescence. The physical signs of croupous pneumonia are well-defined, usually limited to one lung and to a lower lobe; the signs of catarrhal pneumonia are ill-defined and irregularly disseminated.

How is passive or hypostatic congestion of the lungs to be distinguished from pneumonia?

Passive or hypostatic congestion of the lungs may develop in the course of acute or chronic debilitating diseases or as a result of heart-disease; it usually affects the bases and most dependent portions of the lungs. It is attended with increased frequency of breathing, impaired percussion-resonance, and the presence of fine râles. It differs from pneumonia in its association with the conditions that give rise to it; in the absence of fever, unless arising in the course of a febrile disorder; in its bilateral distribution; in the absence of blowing breathing; in its course and in its submission to appropriate management. If expectoration attend congestion of the lungs it is not hemorrhagic.

How is pulmonary infarction to be distinguished from pneumonia?

Coagulation of the circulating blood may take place in the right side of the heart as a result of increased coagulability or of retarded circulation, or a thrombus may form in a distant vein, and detached fragments of clot be swept into the lungs. The occurrence is attended with pain in the chest, dyspnea, and blood-streaked expectoration. Small areas of dulness on percussion can be detected, with enfeebled or bronchial breathing. There may be slight fever, from secondary inflammation. The condition differs from pneumonia in being secondary to a pre-

existent condition, in the circumscribed distribution of the physical phenomena on the part of the lungs, in the comparative absence of fever, and in the course of the disease. Pulmonary infarction is not of itself fatal. The danger is from the condition of which the infarction is but one result.

How is pulmonary infarction to be distinguished from pulmonary abscess?

Pulmonary embolism may result in either infarction or abscess. The outcome depends upon the presence or absence of pyogenic microörganisms. The immediate phenomena are the same in both instances. If suppuration take place, however, there are repeated chills, decided fever, with purulent expectoration. Pulmonary abscess is obviously the graver condition. The existence of disseminated suppuration is an aid in the diagnosis.

How is pneumonia to be discriminated from pulmonary tuberculosis?

Pneumonia is an acute disease, terminating by crisis in from five to nine days; the tendency of pulmonary tuberculosis is to be chronic; its duration is indefinite. The sputum of pneumonia may be profusely admixed with blood, but actual hemorrhage does not occur except in the peculiar form of pneumonia associated with influenza. Characteristic diplococci are present; while tubercle-bacilli are found when pulmonary tuberculosis exists. The emaciation, the anemia and the sweats of pulmonary tuberculosis are not present in uncomplicated croupous pneumonia. The temperature of tuberculosis commonly displays wide variations; the temperature of pneumonia is of a continued type. Tuberculosis usually involves the upper portion of the lung, successively invading the remainder of the lung and the opposite lung as well; pneumonia is usually unilateral and situated in a lower lobe. The physical signs differ in accordance with this peculiarity of distribution. So-called tuberculous pneumonia may be indistinguishable from the ordinary type of croupous pneumonia until tubercle-bacilli appear in the sputum; it may be suspected if diplococci are absent.

Pulmonary Gangrene.

What are the clinical features of pulmonary gangrene?

Gangrene of the lung may result in the course of pulmonary tuberculosis, or of pneumonia; it may depend upon ulcerative communication of the lung with an adjacent septic process; it may be occasioned by the inspiration of septic matters from some portion of the respiratory tract; finally it may have its origin in a septic embolus. The symptom characteristic of pulmonary gangrene is the expectoration of a horribly offensive, stinking sputum, in association with a typhoid condition. Unless cavitation occurs, the physical signs are obscure; at best, they may be masked by those of the primary condition.

How is pulmonary gangrene to be distinguished from pulmonary abscess?

The physical signs and the local phenomena of pulmonary gangrene and of pulmonary abscess may be similar; but the expectoration in a case of abscess is copious and purulent, while that of a case of gangrene is brownish and horribly offensive and more likely to contain fragments of pulmonary structure. The signs of a cavity are more likely to attend abscess than to attend gangrene. Gangrene is more nearly certain than abscess to be fatal.

How are pulmonary tuberculosis and pulmonary gangrene to be differentiated?

Gangrene may complicate tuberculosis of the lung. It differs from uncomplicated tuberculosis by the characteristic odor of the breath and sputum and the absence of tubercle-bacilli from the matters expectorated.

Pulmonary Tuberculosis.

What are the varieties of pulmonary tuberculosis?

In the present state of knowledge pulmonary tuberculosis may, for practical purposes, be considered under two forms: *confluent* and *disseminated*. The lesion in both is essentially the

same, the process in the one instance having a tendency to localization and chronicity, with slow extension; in the other to diffusion and acuteness. In both, also, the process may be acute or chronic. Acute miliary tuberculosis is rarely a pulmonary affection alone, but most frequently involves several organs from the first, and is comparable, in many respects, to an acute infectious fever, such as one of the exanthemata. Indeed, the lungs may be less involved than, for example, the intestines or the peritoneum. Localized tuberculosis may undergo retrogression and cicatrization by fibroid substitution, sometimes associated with calcareous deposits. On the other hand, it may terminate by softening (cheesy necrosis or caseation), leading to excavation or the formation of vomicæ. Calcification, however, may occur even after caseation has taken place.

Galloping consumption or *florid phthisis* is either an extremely rapid form of confluent tuberculosis of the lung or a rapidly caseating subacute or chronic broncho-pneumonitis, on which tuberculosis has supervened.

Fibroid phthisis is a slow, localized tuberculosis in which cicatrization keeps almost even pace with tuberculous ulceration.

The fibroid conditions of the lungs found in miners, in steel-grinders, and in others exposed to the inhalation of fine particles of dust, will be considered as forms of interstitial pneumonitis, comparable to interstitial hepatitis and interstitial nephritis. Tuberculosis may supervene, running then the ordinary course.

What are the symptoms of pulmonary tuberculosis?

The course of ordinary chronic pulmonary tuberculosis may pathologically be divided into *three stages:* a *first* or *incipient stage*, in which the formation of tubercles begins; isolated, miliary nodules are localized at some part of the lung, most commonly at an apex; a *second stage*, of *consolidation*, in which the formation of tubercles has increased in number and density, but not correspondingly in extent, the tendency to localization continuing; and a *third stage*, of *softening*, in which, as a result of caseation and breaking down of the older tuberculous formations, cavities develop; while at the same time invasion of new areas takes place.

The *physical signs* in the different stages differ in accordance with the lesions. The *symptoms* of one stage, however, pass by imperceptible gradations into those of another, and it must be remembered that while in certain portions of the lung the process may have reached the third stage, in other portions tuberculous infiltration may be just beginning.

The *onset* of the disease is usually insidious. The individual complains of a sense of lassitude and of an unusual readiness of fatigue. Exertion induces shortness of breath. There is a gradual failure of nutrition. The *color* fades; the *digestion* is deranged. Perhaps now a slight, irritating cough, attended with little or no expectoration, sets in. Sometimes, however, the first symptoms are observed after a neglected "cold," or supervene upon an attack of catarrhal pneumonia (so-called caseous phthisis). An occasional sense of chilliness in the back may be perceived. The *body-weight* diminishes. Careful observations may detect a slight rise of temperature at noon or within an hour afterward, and perhaps again toward five o'clock. Perhaps at this time, unprovoked, the patient feels a tickling in the throat, perceives a salty taste, and before he realizes what is about to transpire he ejects a mouthful of bright-red, frothy blood. The *hemorrhage* may be repeated; at this stage it is not likely to be fatal. The cough becomes more aggravated; it occurs in paroxysms, the severity of which may induce vomiting, especially on rising in the morning. The *sputum* is abundant and muco-purulent; it contains elastic fibers and tubercle-bacilli. The dyspnea has become decided. Emaciation progresses; recognizable evening fever appears; debilitating night-sweats occur. Hemorrhages become more common and more profuse. The *cheek* is flushed, the *eye* bright, the *intelligence* quickened, the *mind* hopeful. The *hair* becomes straight and prematurely gray. *Digestion* fails. Diarrhea becomes troublesome. Atrocious pains in various parts of the chest indicate the existence of accompanying pleuritic processes. The ends of the fingers become enlarged and bulbous, the nails incurvated. The anemia is profound; the *lower extremities* become edematous. The patient must take to his bed. In the last stages of the

disease, the patient may suddenly be seized with agonizing pain in the side, increased dyspnea, and a sense of impending death. One side of the chest becomes enlarged; the percussion-resonance is tympanitic; the breathing is but feebly transmitted; and metallic tinkling may be developed. *Pneumothorax* has occurred.

Death ultimately results from exhaustion, and is frequently preceded by the supervention of disseminated tuberculosis, giving rise to the febrile and toxemic symptoms of that condition. Occasionally a large hemorrhage brings about a fatal issue by suffocation, from inspiration of a portion of the fluid.

Pulmonary tuberculosis is not rarely complicated by or associated with other tuberculous affections, especially laryngeal tuberculosis and intestinal tuberculosis, the symptoms of which are then superadded.

What are the physical signs of the first stage of pulmonary tuberculosis?

The *chest* may be full and symmetrical. The *respiratory excursion* may be sufficient; perhaps, by standing behind the patient and placing the hands on the front of his chest, a slight deficiency of expansion at an apex may be detected. The *percussion-resonance* is vesicular, except over a small area at the uppermost portion of one lung, where there is slight relative impairment. At this point, too, the vesicular murmur is altered. The inspiratory murmur is less soft and breezy than normal, while expiration is prolonged and heard more distinctly than usual; or inspiration is jerking and arrhythmic. Careful *auscultation* by an acute observer may detect fine moist râles at the end of inspiration. *Vocal resonance* and *fremitus* are slightly increased. If there is any expectoration, careful search may detect a small number of tubercle-bacilli in the *sputum*. With the *fluoroscope* a faint shadow may be seen to replace the normal pulmonary transparency in the affected area, and this may be demonstrated by *skiagraphy*. The excursion of the diaphragm may also be seen with the fluoroscope to be limited on the affected side.

What are the physical signs of the second stage of pulmonary tuberculosis?

The *respiratory frequency* is slightly accelerated. The upper part of the chest is less full on one side than on the other, and the *expansion* in this area is deficient. The *resonance* here is impaired; the *percussion-note* may be dull. The vesicular element of the respiratory murmur is wanting. Inspiration is harsh; expiration is prolonged and blowing; the breathing may be bronchial. Fine crackling sounds are heard. *Vocal resonance* and *fremitus* are notably increased. Blowing sounds may be heard on stethoscopic auscultation over the subclavian vessels. The *sputum* contains many bacilli. Fluoroscopic shadows are marked.

What are the physical signs of the third stage of pulmonary tuberculosis?

The general emaciation is striking. Decided depressions exist above and below the clavicle, on one or on both sides. The ribs are prominent, the interspaces narrowed. The *chest* appears rigid; it may heave, but it scarcely expands in respiration. The *percussion-dulness* is marked, perhaps in different degree over the upper lobes of the two sides. Here and there, in irregular areas, are heard the dull sounds of a thickened pleura. In the midst of the dulness a tympanitic or amphoric or cracked-pot sound is elicited. Coarse and fine râles, gurgling, bronchial or cavernous breathing are heard. Bronchophony or whispering pectoriloquy is present. *Tactile fremitus* is increased. The *sputum* contains bacilli and elastic fibers.

What are the best methods for detecting the presence of tubercle-bacilli in sputum?

In all bacteriologic investigations, the instruments and appliances used must be rigidly sterile. Sputa for examination should be collected in glass or porcelain vessels that have been made clean by boiling, and finally washed with a 1 : 1000 solution of mercuric chloride. The examination should not be too long deferred. In the interval the vessel should be covered.

The sputum should be poured from its receptacle on a sterilized plate of glass having a black background. With a platinum wire, first heated to redness in the flame of a lamp or

Bunsen burner and permitted to cool, a small quantity of purulent matter is taken up and spread in a thin layer upon a cover-glass that has been washed with alcohol, ether, and bichloride solution and carefully wiped dry. The thin layer of sputum is permitted to dry at ordinary temperature, or the process is accelerated by gentle heat. When perfectly dry, the cover-glass preparation is drawn three or four times through the flame so as to fix the albuminoid matters.

The preparation is now ready to be stained. For the staining, a filtered solution composed of one part of fuchsin (or gentian violet), four of carbolic acid, ten of alcohol and one hundred of sterilized distilled water, may be used ; or ten or fifteen drops of anilin oil and about a dram of sterilized distilled water are shaken together and filtered, and then sufficient of a saturated alcoholic solution of gentian-violet (or fuchsin) addded until a deep coloration is produced.

The cover-glass preparation may now be floated on the surface of one of the solutions indicated, for twenty-four hours, at the ordinary temperature, or the process may be accomplished in

FIG. 23.



Tubercle-bacilli in sputum. (Ziegler.)

fifteen minutes by the application of moderate heat until the vapor of steam arises ; or face upward, a few drops of either of the solutions indicated are placed upon the surface of the cover-glass preparation, which is gently heated until the vapors of steam arise.

The excess of free stain is removed with water. The preparation is decolorized in a solution of nitric acid (1 : 3), or in one of sulphuric acid (1 : 4). It is then briefly passed through seventy per cent. alcohol. It may at once be dried and mounted and examined, but it is better to stain again with an aqueous solution of a color contrasting with that first used—fuchsin and methylene-blue, gentian-violet and vesuvian or Bismarck-brown.

The search for bacilli in sputum may be facilitated by adding to about four ounces of sputum, one ounce of sterilized distilled water, and from four to eight drops of liquor sodæ and heating; from two to three ounces more of water are added and the mixture is put aside in a conical glass for from twenty-four to forty-eight hours, when the supernatant fluid is decanted and cover-glass preparations made from the sediment.

Centrifugation may be practised when bacilli are not readily found otherwise.

What is the clinical course of "galloping consumption"?

The rapid form of confluent pulmonary tuberculosis, known as "quick consumption" or *phthisis florida*, as distinguished from the infectious fever known as acute miliary tuberculosis, not infrequently supervenes upon a neglected "cold," commonly a catarrhal pneumonia of the apex; or it may suddenly become manifest after localized tuberculosis of the apex, so slight as to be unsuspected, has existed for a longer or shorter time; it may closely or remotely follow an attack of influenza; or its immediate antecedents may be untraceable. Frequently, the subjects of phthisis florida present an hereditary liability to tuberculosis. Careful inquiry may elicit a history pointing to tuberculous infection of the bronchial glands in childhood, as shown by recurrent attacks of cough and fever, or to actual "scrofulous" manifestations.

Clinically the disease is marked by high temperature of a continued or, more frequently, a remittent type; profuse night-sweats; profound digestive and circulatory disturbance; rapid wasting, with accompanying weakness; and pulmonary symptoms of progressive severity, with corresponding physical signs. Pathologically it is characterized by rapid caseation, with for-

mation of extensive cavities; severe laryngeal and intestinal complications are frequent. It may apparently begin as an acute laryngitis of ordinary or of tuberculous type. Sometimes persistent diarrhea is the earliest symptom to attract attention. The duration of the disease is from a few weeks to about eighteen months.

How is a pulmonary hemorrhage to be discriminated from a gastric hemorrhage?

The blood of pulmonary hemorrhage has a sweetish or saltish taste, is usually fluid, bright-red, alkaline and more or less frothy. There is a previous history or evidence of disease of the lungs or heart. The expectoration continues blood-streaked for several days. In gastric hemorrhage the blood is usually acid, dark and clotted; its taste is masked as a result of admixture with the contents of the stomach; the stools that follow are tarry.

Pulmonary hemorrhage may occur with or without cough, and is not provoked by taking food. Gastric hemorrhage usually occurs with vomiting, and is provoked by taking food.

How is the distinction to be made between pulmonary tuberculosis and chronic pleurisy?

Pleural adhesion and thickening are extremely common. They occasion retraction of the chest-wall, often more or less displacement of the heart, and give rise to dulness on percussion, enfeebled transmission of the breath-sounds and diminished vocal resonance and fremitus; but they are not associated with the ascultatory and constitutional phenomena of pulmonary tuberculosis, or with the presence of tubercle-bacilliin the sputum. If a pleural effusion is present, there are bulging of the lower part of the chest, flatness on percussion, feebleness or absence of breath-sounds, and diminished vocal resonance and fremitus. If the fluid is purulent, there are chills, fever, sweats and emaciation.

How is pulmonary carcinoma to be distinguished from pulmonary tuberculosis?

Carcinoma of the lung is more commonly secondary than

primary. The metastatic growths give rise to irregularly distributed, multiple areas of percussion-dulness, without corresponding change in the auscultatory phenomena. The temperature is likely to be sub-normal rather than febrile, unless inflammation of the lung or pleura is excited. The duration of pulmonary carcinoma is limited; pulmonary tuberculosis may be indefinitely protracted. Tubercle-bacilli are not found in the sputum in pulmonary carcinoma. The presence of carcinomatous new-growths elsewhere is significant in doubtful cases.

How is pulmonary syphilis to be distinguished from pulmonary tuberculosis?

Syphilis of the lung appears either in the form of gummata or as fibroid induration. It is to be distinguished from tuberculosis by the absence of bacilli from the sputum, by the involvement of the middle and lower parts rather than the upper part of the lung, by the diffuse and irregular rather than concentrated localization of physical signs, by the absence of the constitutional phenomena of tuberculosis, and by the history or the knowledge of the existence of other syphilitic manifestations. The therapeutic test is sometimes available, but should be cautiously applied, as cases of tuberculosis are usually injured by potassium iodide.

Syphilis and tuberculosis may coëxist.

How are bronchiectasis and pulmonary tuberculosis to be differentiated?

Bronchiectasis is attended with copious muco-purulent expectoration, and perhaps with decided emaciation; the pulmonary resonance may be impaired, and large, moist râles may be heard; but the physical phenomena are often bilateral in distribution and most decided at the bases of the lungs; pulmonary tuberculosis usually begins at the apex of one lung, whence it extends. When, as is not uncommonly the case, bronchial dilatation is found in the upper portion of one lung, its discrimination from a tuberculous vomica is difficult, and must be based rather on the results of microscopic investigation and the general phenomena than on the physical signs. Bronchiectasis

is slowly if at all progressive; tuberculosis less commonly stops short of a fatal termination. The sputum of a case of bronchiectasis does not contain tubercle-bacilli.

How is an abscess of the lung to be diagnosticated from pulmonary tuberculosis?

An abscess of the lung may develop in the course of a pneumonia or an empyema, as a result of traumatism, or it may constitute a manifestation of a general pyemia. It is to be distinguished from tuberculosis by a knowledge of its possible origin, by the appearance of the phenomena in a lower rather than in an upper lobe and upon one side only. The condition is acute rather than chronic, and the sputum does not contain tubercle-bacilli. The physical phenomena are peculiarly circumscribed.

By what means are malarial fever and pulmonary tuberculosis to be differentiated?

When caseation and suppuration are taking place in a lung, the temperature is high in the evening, declining towards morning. In addition, however, there are persistent dyspnea, increased frequency of breathing, cough, expectoration containing tubercle-bacilli, loss of flesh, night-sweats, dulness on percussion, and mucous râles on auscultation. The symptoms last enumerated are wanting in intermittent fever, while the plasmodia of malaria are wanting in the blood in pulmonary tuberculosis. Possible mistakes in diagnosis are to be avoided by a physical examination.

Acute Miliary Tuberculosis.

What are the symptoms of acute miliary tuberculosis?

Acute miliary tuberculosis sets in insidiously. The victim may be a member of a tuberculous family. He is taken ill with slight cough, scanty or no expectoration, marked dyspnea, elevation of temperature, rapidity of pulse, delirium, and other symptoms of a grave intoxication. Tuberculous involvement of the bowel will cause diarrhea; of the cerebral meninges, the signs of meningitis. The course of the disease may be rapid or protracted. Death may take place from exhaustion, or, in

rarer instances, the acute symptoms subside, and those of chronic tuberculosis appear.

What are the physical signs of acute miliary tuberculosis?

The physical signs of acute miliary tuberculosis of the lungs are at first practically those of an acute bronchitis: a full chest; rapid breathing; harsh, vesiculo-bronchial respiratory sounds; unaltered percussion-resonance; unchanged vocal resonance and fremitus. The formation of subpleural tubercles gives rise to a peculiar friction-fremitus appreciable on palpation. Later, dulness at one apex or at both apices, or just below, and fine crackling râles are heard; and still later, evidences of caseation are found on auscultation and percussion. The detection of tubercle-bacilli may be very late. Some cases early develop indications of apex-pneumonia.

How are acute miliary tuberculosis and typhoid fever to be distinguished from one another?

The prostration and the general condition are much alike in both; but in acute miliary tuberculosis the epistaxis, the rose-spots, the characteristic temperature-course, urine-reaction and blood-reaction, and the peculiar stools of typhoid fever are wanting. The temperature of typhoid fever pursues a definite course, that of acute miliary tuberculosis presents considerable oscillations on the same day; without therapeutic interference, it may be normal, subnormal and very high. Bronchitis may attend typhoid fever, but the dyspnea is never so marked as it is in acute miliary tuberculosis. The progress of the case determines the diagnosis. Not only do the marked physicial signs of destructive change in the lungs develop in the one disease and not in the other, but typhoid fever is a self-limited disease of known duration, with a tendency to recovery, while acute miliary tuberculosis is a disease of uncertain duration, and sooner or later almost invariably fatal.

Tubercle-bacilli may sometimes be discovered in the sputum of acute miliary tuberculosis, in the blood or in the spleen.

The knowledge of previous scrofulous or tuberculous disease in the patient, or a family history of scrofula or tuber-

culosis, should excite suspicion of acute tuberculosis; and in susceptible subjects the latter disease may even quickly follow typhoid fever.

Interstitial Pneumonitis.

What are the characteristics of interstitial pneumonitis?

As a result of the irritation occasioned by the constant inhalation of fine particles of dust by miners and grinders, and others, a chronic bronchitis with hyperplasia of the interstitial pulmonary tissue develops. The two layers of pleura are often adherent and thickened. As time progresses, contraction takes place, terminating in condensation of the lung and dilatation of the bronchial tubes. Interstitial pneumonitis may develop as a part of a general fibroid degeneration, it may follow acute pneumonia or other disease of the lung; or it may be associated with pleural thickening.

In coal miners the condition is known as *anthracosis;* in iron-workers as *siderosis;* in grinders as *chalicosis.*

What are the symptoms of interstitial pneumonitis?

There is obstinate cough and abundant muco-purulent expectoration, which sometimes contains particles of the dust inhaled. There may be considerable wasting and dyspnea.

What are the physical signs of interstitial pneumonitis?

The *chest* is diminished in size, rather flattened anteriorly, from retraction. The *respiratory excursion* is small, the chest expanding but little in inspiration. The *percussion-resonance* is impaired over a large area of pulmonary surface. The *breathing* is bronchial. Large moist and dry râles are heard. The *vocal resonance* and *fremitus* are increased.

How does interstitial pneumonitis differ from chronic pleurisy?

Interstitial pneumonitis and chronic adhesive pleurisy are allied conditions and are often associated. Occurring alone the former is wider in its distribution and is more likely than the latter to be bilateral. Blowing breathing, râles, cough and expectoration attend the pulmonary, but not the pleural, condition.

Pulmonary Emphysema.

What are the characteristics of pulmonary emphysema?

Pulmonary emphysema is usually a result of long-continued excessive expiratory effort with closed glottis—as in protracted cough or in the habitual blowing of wind instruments. It is often associated with chronic bronchitis. The air-vesicles become dilated into large sacs, and the bloodvessels in the interlobular septa are obliterated. The increased work thrown upon the right heart in turn gives rise to dilatation.

Emphysema manifests itself by dyspnea, expiratory in character; shortness of breath, aggravated in paroxysms, and attended with distressing cough and scanty expectoration; and cyanosis. The cardiac insufficiency adds to the dyspnea and may be the cause of dropsy.

What are the physical signs of emphysema?

The *chest* is large, "barrel-shaped;" the circumference being increased in greater degree than the vertical diameter. The *respiratory excursion* is slight and may be scarcely visible; although the chest may rise and fall as a whole, the ribs being fixed in the position of full inspiration. The *percussion-resonance* is heightened, almost tympanitic. The *breath-sounds* are feeble, being almost continuous and without intermission, expiration being prolonged and attended with a succession of puffs. The *vocal resonance* and *fremitus* are diminished. The area of superficial cardiac dulness is diminished, but deep percussion will reveal enlargement of the right ventricle.

How is emphysema to be distinguished from pneumothorax?

Pneumothorax usually results from the breaking through the pleura of a destructive process in the lung, such as the caseation and suppuration of tuberculosis. The symptoms to which it gives rise are abrupt in onset and distressing in character. The patient perceives a sudden, severe pain in the side, and is seized with great dyspnea and a sense of impending death. The chest is seen to be enlarged, the breathing rapid and shallow, the respiratory excursion small; the percussion-resonance is tympanitic and the breath-sounds are feeble. In pneumo-

thorax, the dilatation of the chest and the tympany are, however, unilateral; in emphysema, they are symmetrical. In pneumothorax, an amphoric blowing sound is heard on inspiration, and if pleurisy with effusion have set in, metallic tinkling, succussion sounds and egophony as well. The recognition of a condition that may give rise to perforation and pneumothorax aids in the diagnosis.

How is emphysema to be distinguished from a pleural effusion?

An effusion into the pleura may take place in the course of a pleurisy, or as a result of disease of the heart or kidney. Adjacent organs are displaced in proportion to the quantity of fluid poured out. If the percussion-note be hyper-resonant at the upper part of the chest, above the level of the fluid, it is flat below. Pleuritic effusions are usually unilateral; when the effusion is part of a general dropsy it is usually bilateral. Emphysema is rarely unilateral. In emphysema, the breath-sounds are everywhere feebly heard; in pleural effusion the breathing may be puerile, or even bronchial, above the level of the fluid, but the breath-sounds are not well heard through the fluid. Vocal resonance and fremitus are diminished in emphysema; absent below the level of a pleural effusion. If the effusion becomes purulent, rigors, hectic fever, sweats and marked emaciation occur.

Pneumothorax.

What are the clinical features of pneumothorax?

Pneumothorax may result from traumatism that causes fracture of a rib and perforation of the lung; from a perforating wound of the chest; from the rupture of an emphysematous pulmonary alveolus; from communication between the pleural cavity and an adjacent hollow viscus; from the softening of a septic infarct or of an area of pneumonic consolidation; from the development of gas in the pleural cavity; but the most common cause is ulcerative perforation of a tuberculous cavity in the lung. The condition usually *sets in* suddenly, with sharp pain in the side, intense dyspnea and a sense of great oppres-

sion. The *breathing* is hurried and shallow. The *chest* on the affected side becomes bulging; the *percussion-note* is tympanitic; the *breath-sounds* are enfeebled or distant, perhaps amphoric; as fluid is poured out, metallic tinkling may be heard, and by shaking the patient, with the ear applied to his chest, a succussion-sound may be elicited; the *voice-sounds* and *vibrations* are poorly transmitted. Voice and cough assume a metallic tone in transmission, and a peculiar bell-like resonance is given the transmitted percussion-note when metallic plexor and plexímeter (two coins) are used. The *heart*, as well as other viscera, may be displaced.

How are pneumothorax and diaphragmatic hernia to be differentiated?

The stomach may be dragged upwards beneath the ribs by a contracting lung, or the stomach or the large intestine may bulge through a yielding portion of the diaphragm, so that there may be unilateral percussion-tympany, with displacement of the heart; vibrations of the fluid in the hollow viscus may give rise to metallic tinkling, which, however, occurs independently of respiration and is associated with rumbling sounds peculiar to the gastro-intestinal tract. The interference with the respiratory functions that results is, moreover, not only acute in onset, as it is in pneumothorax (of which the etiologic elements are wanting), but it may be equally sudden in disappearance. Diaphragmatic hernia may have a history dating from birth; it bodes no danger unless strangulation occurs, of which the symptoms are characteristic. The outcome of pneumothorax is in the nature of things often rapidly fatal.

How are pneumothorax and a pulmonary cavity to be differentiated?

A large cavity in the lung may yield a tympanitic percussion-sound, amphoric respiration, metallic râles or tinkling, and succussion-sounds, but the phenomena are usually circumscribed; there is no bulging, no acute exacerbation of pain, dyspnea and oppression, and the percussion-flatness of an effusion into the pleural sac is wanting.

How is pneumothorax to be distinguished from a pleural effusion?

Pneumothorax sets in suddenly, with acute pain and dyspnea. A pleural effusion usually takes place insidiously. Pleurisy with effusion usually attends pneumothorax, so that the signs of both may be present; but when the pneumothorax is the primary condition, the percussion-note above the level of the effusion is more pronouncedly tympanitic and the voice more distinctly egophonic. Succussion-phenomena and tinkling are not heard unless both air and fluid are present in the pleural sac. Finally, the recognition of a condition that gives rise to pneumothorax may decide the diagnosis.

How are pneumothorax and subphrenic abscess to be differentiated?

Sometimes an accumulation of pus and gas takes place beneath the diaphragm in the sequence of perforation of the stomach or bowel. It may occur on either side and give rise to symptoms closely resembling those of pneumothorax. The differentiation will depend essentially upon a knowledge of the previous existence of disease of the lung on the one hand or of the stomach or bowel on the other.

Asthma.

What are the symptoms of asthma?

Asthma is a paroxysmal affection, the symptoms of which may be dependent upon spasmodic narrowing or exudative inflammation of the smaller bronchial tubes, turgescence of the bronchial vessels or an urticaria-like affection of the bronchial mucous membrane. It is predisposed to by a neurotic state. The attacks recur at irregular intervals, with or without apparent exciting cause. Among provocative conditions are changes of climate and weather, overeating, indigestion, emanations from various sources, possibly reflex disturbances and sexual excitement.

The seizure is sudden in *onset*, usually occurring at night. The patient is awakened with a sense of oppression and distressing dyspnea, inspiratory in character. Orthopnea is common. Dur-

ing the *paroxysm*, which may last for several hours, the chest heaves spasmodically, but the *pulmonary expansion* is slight; the *face* is pale or livid, and the distress is evident. Loud, wheezing sounds are heard. On auscultation, the *breath-sounds* are feeble, or the vesicular murmur may be obscured by the wheezing or replaced by sonorous and sibilant râles. The *percutory phenomena* are unaltered. The *crisis* takes place with a profuse expectoration of mucus, perhaps, also, with a copious discharge of limpid urine. This may terminate the attack, or the same phenomena may be repeated on several successive nights or even invade the day. The sputum consists of thin mucus, and round gelatinous masses, which are found, microscopically, to have a spiral arrangement. Pointed octahedral cystals also are present. The number of esinophilous blood-corpuscles is increased.

Asthma may be primary or secondary. It may result from reflex influences, such as disease of the nose. It may apparently replace the convulsive seizure of epilepsy. Asthmatoid seizures are common accompaniments of chronic bronchitis and emphysema, of cardiac insufficiency, and of chronic nephritis.

How is asthma to be distinguished from an asthmatoid condition?

In true asthma, no cause for the disease may be found or else some local or reflex irritation may be discovered. When so-called asthmatoid attacks occur, their discrimination depends upon the recognition of an organic pulmonary affection, such as emphysema or bronchitis, or of cardiac incompetency or of chronic nephritis. Such attacks do not pursue the typical course of true asthmatic seizures and do not terminate suddenly with profuse expectoration.

How does asthma differ from whooping-cough?

Whooping-cough is a disease of children; asthma, a disease of adults. The characteristic whoop of pertussis is wanting in asthma. An attack of whooping-cough does not last more than a couple of months, at most; asthma may continue indefinitely.

What are the points of differentiation between asthma and paralysis of the diaphragm?

The dyspnea resulting from paralysis of the diaphragm is not

paroxysmal, as is that of asthma. Paralysis of the diaphragm is not characterized by an absence of the breath-sounds, followed by wheezing and high-pitched râles. When the diaphragm is paralyzed, inspiration is attended with expansion of the chest and depression of the abdominal wall; in expiration the chest collapses and the abdominal wall is elevated; stimulation of the phrenic nerves restores the normal harmony of action; attempts at bearing down are futile or ineffective.

New-growths in the Lungs.

What are the clinical features of new-growths in the lungs?

New-growths in the lungs are usually secondary, rarely primary. The most common varieties are epithelioma, encephaloid and scirrhus. Sarcoma is less common. Primary carcinoma usually involves one lung only, secondary growths both. Among the principal symptoms are pain, dyspnea, cough and prune-juice expectoration. There may be also cyanosis, edema and dilatation of the veins from pressure. The percussion-resonance is impaired or lost, the breathing enfeebled, possibly bronchial; vocal resonance and fremitus are diminished. External lymphatic glands may be enlarged. Sometimes there is elevation of temperature. Emaciation is moderate.

THE DIGESTIVE SYSTEM—THE MOUTH.

Catarrhal Stomatitis.

What are the clinical characteristics of catarrhal stomatitis?

Catarrhal inflammation of the mucous membrane of the mouth may *result* from the ingestion of irritating substances, from the presence of carious teeth, or by extension from adjacent disease; it may also develop in conjunction with morbid dentition, or derangement of digestion, or in the course of the exanthemata. It *manifests* itself by redness, tumidity, and increased heat of the structures within the mouth, and by increased secretion. The taking of food is attended with discomfort or

with pain ; taste is impaired ; and the breath is offensive. As a rule, there is little or no constitutional disturbance.

Aphthous Stomatitis.

What are the clinical features of aphthous stomatitis?

In individuals exposed to unfavorable hygienic conditions, and in those debilitated by disease, small vesicles, surrounded by reddened areolæ, appear and rupture, leaving ulcers with grayish bases at various parts of the mucous membrane of the mouth. Mastication in adults and nursing in infants are difficult and painful; the secretions of the mouth are increased; the appetite may be impaired; digestion may be deranged; and there may be diarrhea.

Thrush.

What are the clinical manifestations of thrush?

Thrush, *muguet* or *parasitic stomatitis* is a mycotic inflammation of the mucous membrane of the mouth and throat, to which children are especially prone. It is dependent upon the presence of a fungus, termed the *mycoderma* (or *oïdium*) *albicans*. An acid reaction of the secretions of the mouth is an essential condition for the development of the affection. The growth of the fungus and the resulting irritation give rise to the formation of minute curd-like masses upon various parts of the mucous membrane. Forcible detachment of the masses occasions bleeding. The *symptoms* are those of the ordinary form of stomatitis, plus the flaky deposits containing the characteristic fungus. There may also be diarrhea and a varying degree of constitutional disturbance. In nurslings, the inability to suckle may result in inanition and death.

Ulcerative Stomatitis.

What are the symptoms of ulcerative stomatitis?

Ulcerative stomatitis is an aggravated form of inflammation of the mouth, attended with ulceration, which it is stated is

usually unilateral. The affection *arises* amid conditions of crowding and filth, and in those supplied with insufficient and inappropriate food ; it may also be a sequel of other forms of stomatitis, of caries of the teeth, or of the maxillary bones. It is *manifested* by impaired appetite, fetid breath, increased salivation, pain in eating, and constitutional symptoms of varying intensity. Sometimes adjacent lymphatic glands undergo enlargement.

Mercurial Stomatitis.

What are the clinical manifestations of mercurial stomatitis?

Stomatitis sometimes *results* from the medicinal ingestion of large quantities of *mercury*, or of small quantities by persons possessing an idiosyncrasy, or as a manifestation of mercurial intoxication by means of articles of food or drink, or from exposure to the metal in certain occupations. The *symptoms* vary greatly in severity. The gums especially become swollen, reddened, tender, and sometimes ulcerated. The *teeth* may fall out, and the maxillary bones become carious. The *breath* is fetid. The *saliva* contains mercury ; its secretion and discharge are inordinately increased, giving rise to the term "salivation," as descriptive of the affection.

Gangrenous Stomatitis—Noma.

What are the clinical features of gangrenous stomatitis?

Gangrenous stomatitis, *noma* or *cancrum oris* is essentially an affection of childhood, rare, and almost invariably fatal, which *develops* in those of depraved constitution, often at the termination of one of the exanthemata, particularly measles. It is probably of bacterial origin. The disorder is *manifested* by a brawny induration of one cheek, the structures of which undergo disintegration, with resulting ulceration of the mucous and cutaneous surfaces, and, not rarely, perforation. Adjacent portions of the gums may by contiguity become involved in the process. The teeth may fall out and the maxillary bones be-

come carious. The early symptoms may be obscured by those of the antecedent condition. Soon, however, the breath becomes fetid, and, in addition to the local manifestations, the symptoms of septic intoxication may appear, in the midst of which the child may die. *Recovery* may take place, with hideous deformity of the face. Pneumonia, pulmonary gangrene, and entero-colitis may be *complications.*

THE TONGUE.

Glossitis.

What are the symptoms of glossitis?

When the tongue is inflamed, from whatever cause, the organ becomes enlarged, tumid, reddened, painful; speech, deglutition and mastication, sometimes respiration, are interfered with, and the secretions of the mouth are increased. The swelling may be so great that suffocation results, unless relief be given by incision. Glossitis may be *superficial* or *parenchymatous*, *acute* or *chronic;* the intensity and character of the symptoms being modified accordingly. It may be due to erysipelas, either primarily or secondarily. Epidemics of erysipelatous glossitis have occurred, and the name "black tongue" has been applied to this affection. Parenchymatous glossitis sometimes proceeds to suppuration.

Leukoplakia Lingualis.

What is leukoplakia of the tongue?

Leukoplakia lingualis, or *leukoplakia buccalis*, is a name applied to a peculiar chronic affection of the tongue or of the tongue and buccal mucous membrane, characterized by the formation of persistent, horny, whitish patches upon the surface, sometimes extending entirely through the epithelial layer. Untreated, it is said to lead at times to the development of carcinoma. Sometimes it appears to be related with gout. Syphilis, tobacco and alcohol are also enumerated among its causes.

Geographical Tongue.

What is geographical tongue?

This is a disorder characterized by desquamation of the superficial epithelium of the tongue in circular patches that extend from the periphery. It is attended with a sense of itching and heat. It has been observed in association with digestive derangement and it has been considered of gouty origin. Relapse or recurrence is common.

Glossanthrax.

What is glossanthrax?

Glossanthrax is a term applied to the localization of malignant pustule upon the tongue. It is to be differentiated from carcinoma, tuberculosis, syphilis and other affections leading to suppuration or ulceration. The local and constitutional symptoms are those of anthrax in general. The appearance of the eschar is characteristic and the presence of anthrax-bacilli is diagnostic.

Nigrities.

What is nigrities?

Nigrities, also called *black tongue* and *hairy tongue*, is an affection of the filiform papillæ of the tongue, supposed to be due to the irritation of a special fungus. The papillæ in various situations become discolored, thickened, and elongated, giving the appearance of a scattered or compact hairy growth upon the dorsum of the tongue. Desquamation takes place, after which the tongue may remain comparatively clean for a longer or shorter period; then the growth recurs. The teeth also may be black. The affection is to be differentiated from staining of the tongue by tobacco, medicines, and the like.

MUMPS—PAROTIDITIS.

What are the symptoms of parotiditis?

Parotiditis or *mumps* may be primary, or secondary in the course of infectious diseases. It may be epidemic or endemic. It is *characterized* by pain at the angle of the jaw, followed by tumefaction of the parotid gland, at first on one side and then on the other. Movement of the jaw, as in mastication, is difficult and painful. Deglutition is not interfered with; hearing may be deranged. The secretion of saliva is usually excessive; it may be diminished. There are febrile manifestations of moderate severity. The period of *incubation* is from fourteen to twenty-five days. The *duration* of the disease is from seven to ten days.

Orchitis or ovaritis is a peculiar *complication* of parotiditis. It is likely to occur as the parotid swelling subsides.

THE PHARYNX.

Pharyngitis.

What are the symptoms of pharyngitis?

In *acute catarrhal pharyngitis*, or *angina*, the symptoms vary with the intensity and extent of involvement. Ordinarily, there are "sore throat," irritable cough, pain or difficulty in deglutition, interference with respiration, enlargement of the tonsils and of the glands of the neck. On *inspection*, the soft palate, the uvula, the tonsils, the posterior and lateral walls of the pharynx, or the palatine arches may, one or more, be seen to be reddened and tumefied, and often coated with glairy mucus. A moderate degree of *fever* attends acute pharyngitis.

Acute phlegmonous pharyngitis is a much more serious affection, involving not only the mucous membrane, but also the submucous connective tissues and even at times the sheaths of the muscles. The constitutional symptoms are in accordance with the severity of the process, and may be those of pyemia. The pus may gravitate to the cellular tissues of the neck, manifesting

as an external swelling and causing dyspnea or even suffocation from compression of the trachea. The inflamed tissues of the throat, especially the soft palate and uvula, may be greatly swollen and edematous.

Chronic pharyngitis may be a sequel of repeated acute attacks. It is common in those who smoke, or drink alcohol excessively, or use their voices a good deal. The pharynx presents a granular and sometimes a glazed appearance. There is increased secretion of tenacious, adherent mucus.

What is acute tuberculous pharyngitis?

Acute tuberculous pharyngitis is a form of acute miliary tuberculosis, apparently beginning in the pharynx.

The *constitutional manifestations* are those of an acute febrile process of grave type, sometimes simulating typhoid fever.

Locally, deposits of tubercle may be observed beneath the mucous membrane as little semi-transparent, grayish nodules, resembling in size and form vermicelli-seeds or fish-eggs. These are collected into little patches more or less confluent, which eventually undergo ulceration.

The uvula is sometimes thickened into a somewhat characteristic gelatinous-looking, reddened, club-shaped mass. This gelatinous thickening may likewise take place in other portions of the pharynx.

The disease may extend to the epiglottis, tongue, and larynx, or to the vault of the pharynx and the nasal passages.

The ulcerative process usually begins on a palatine fold or on a lateral wall of the pharynx, whence it rapidly extends. Ulceration may penetrate the submucous tissues, and the muscles may undergo tuberculous or fatty degeneration. Pus is usually absent from the surface of the ulcers and the bacillus tuberculosis is sometimes found in the detritus.

The chief and most distinctive *local subjective symptom* is intense pain in swallowing, often more than can be accounted for by the extent of visible disease. The pain may extend into the ears.

As the disease progresses, cough is superadded, emaciation becomes rapid, and signs of pulmonary disease, and perhaps of other complications, become manifest.

Death may result within a few weeks, and is rarely postponed beyond two or three months.

Recovery is the exception.

How is tuberculous pharyngitis to be distinguished from syphilitic sore-throat?

The intense pain in swallowing, the characteristic deposit and gelatinous infiltration, the absence of pus from the ulcers, the history of the attack, the personal history, the family history of the patient, and the febrile symptoms; the discovery of the tubercle-bacillus in the detritus of the ulcers, or in the sputum, together with the detection of the evidence of pulmonary tuberculosis as the case proceeds, are the points upon which the diagnosis from syphilis must depend.

Syphilitic and tuberculous disease may, however, coëxist.

The supervention of typhoid symptoms in a case of supposed syphilitic ulceration of the throat should excite suspicion of the existence of tuberculosis.

How is tuberculous pharyngitis to be distinguished from typhoid fever?

Typhoid fever is sometimes accompanied by ulcerative pharyngitis and laryngitis, though the cases in which this occurs are much rarer in North America than they appear to be in Europe. The characteristic fish-egg-looking infiltration of tuberculosis and the non-purulent character of the ulceration would make the discrimination locally; while, constitutionally, there would be absence of the characteristic temperature-course and blood-reaction, of the rose-spots, and of the peculiar stools of typhoid fever. Discovery of the tubercle-bacillus would be conclusive.

Tonsillitis.

What are the symptoms of tonsillitis?

When the tonsils are inflamed, adjacent parts usually participate in the process. *Parenchymatous tonsillitis* or *quinsy* may be primary or secondary to various infectious diseases. It sometimes *sets in* suddenly, with a chill, followed by decided elevation of temperature and other febrile manifestations. Usually, one tonsil only is affected, or first one and then the other, but bilateral involvement may occur and the glands become so intensely swollen

as to meet in the middle line, practically obstructing deglutition and respiration. In such cases the voice is nasal, and fluids attempted to be swallowed may return through the nose. There is always some interference with swallowing. The pain is atrocious, and may extend into the ear on the affected side. There is increased secretion of saliva; swallowing is increased in frequency, and aggravates the pain. Sometimes the patient lies with open mouth, making labored and noisy efforts at respiration, while the saliva dribbles. On examination, the tonsils are seen to be enlarged and angry, adjacent parts in some degree participating in the inflammatory process.

The enlargement of the glands can, sometimes, be distinctly detected from without. Tenderness on pressure beneath the angle of the jaw is common.

An inflamed tonsil or the peritonsillar tissues may suppurate, and grave complications, such as ulceration into the carotid artery and suffocation from rupture into the larynx, have occurred.

Lacunal or *follicular tonsillitis* is a much less serious affection. It may follow exposure to cold or wet, or it may arise amid unhygienic surroundings. The process is superficial, the lining membrane of the lacunæ or ducts being involved rather than the substance of the gland. Scattered over the surface of the inflamed and enlarged tonsils are a number of yellowish points or patches, indicative of accumulations of sebaceous matter, desquamated epithelium and fungi, at the orifices of the ducts. These plugs may be readily removed by means of a scoop, sometimes by syringing—a point of some importance in the discrimination from diphtheria. The streptococcus is the microörganism found most commonly in the secretion. Parenchymatous tonsillitis sometimes follows lacunal tonsilitis, and tonsillar or peritonsillar abscess may then develop. Untreated, the duration of tonsillitis is from two to ten days or more.

Some cases of tonsillitis are associated with pain and other rheumatic manifestations in the muscles or joints, and endocarditis and pericarditis have been noted. An endocardial murmur is not uncommon. Swelling of the joints may appear in apparent metastasis as a tonsillar inflammation declines.

Anomalous eruptions and albuminuria are among the less common concomitants of tonsillitis. Paralyses are very rare sequelæ.

Chronic tonsillitis is most common in young persons and is characterized by hyperplasia, causing obstruction of the pharynx, with inability to breathe through the nares, in consequence of which there result deformities of the chest, changes in facial expression, mental impairment and bodily stunting. Asthmatoid seizures may occur by day, and snoring, with "terrors" at night. The breath is fetid. There are also headache, altered nasal voice, imperfect articulation; and hearing, taste and smell may be impaired. Similar conditions may attend hyperplasia of the adenoid tissues of the vault of the pharynx, the so-called pharyngeal tonsil.

Herpetic Sore-Throat.

What is herpetic sore-throat?

Herpetic sore-throat, *herpetic tonsillitis*, *herpes of the pharynx*, *common membranous sore-throat*, *ulcero-membranous angina*, *diphtheroid throat*, are names applied to a disease often mistaken for diphtheria, but which is, in reality, a form of inflammation of the mucous membrane of the palate, tonsils, uvula and pharynx, characterized by the eruption of herpetic vesicles, which soon rupture, leaving little circular ulcers that coalesce and become covered with a fibrinous exudation. It is sometimes associated with herpes of the lips.

Constitutional symptoms may be absent, but, when present, are usually of a mild febrile type. There may, however, be high fever, preceded by malaise or chill.

The pain in deglutition (odynphagia), and the dryness and heat of the throat are often much greater than in ordinary forms of pharyngitis. In rare instances, in children, the false membrane has extended into the larynx, causing suffocation. Usually the disease terminates in recovery in about a week or ten days.

Chronic or *recurrent herpes of the throat* is encountered in rare instances.

How is common membranous sore-throat to be distinguished from diphtheria?

The diagnosis is sometimes very difficult, and, when in doubt, the safer plan is to consider the case one of diphtheria until bacteriologic investigation has been made. As a rule, however, the islet-like distribution of the patches of fibrinous exudation covering the ulcers left by rupture of the vesicles of herpes is quite different from the appearance presented by the coherent, continuous mass of thick, yellowish or grayish membrane observed in diphtheria. Herpes is more frequent upon the palate and tonsils. Diphtheria usually involves the pharynx extensively. The constitutional symptoms of diphtheria are, as a rule, much more profound than those of herpetic sore-throat. Herpes of the lips often coëxists with herpetic sore-throat, rarely with diphtheria. Diphtheria is contagious. Herpes is non-contagious. A specific bacillus causes the one and not the other.

Gangrenous Pharyngitis.

What is gangrenous pharyngitis?

Gangrenous pharyngitis, or *putrid sore-throat*, may *originate* independently of any other malady or may follow ordinary forms of pharyngitis, or the sore-throat of the exanthemata, or of dysentery, or of typhus or of typhoid fever. It sometimes occurs in cases of tuberculosis. *Constitutional symptoms* are typhoid in type. The *local symptoms* are those of violent inflammation of the mucous membrane of the tonsils, palatine folds, and walls of the pharynx, which soon become covered with gangrenous patches. The destructive process rapidly extends, sometimes into the esophagus, the larynx, and the nares. Sometimes the process is extremely limited, as to the tonsils. Erosion of blood-vessels may cause fatal hemorrhage.

How is gangrenous sore-throat to be distinguished from diphtheria?

In gangrene, the patches are grayish-black in color from the outset, while the pseudo-membrane of diphtheria becomes dark only as the disease progresses.

Swelling of the cervical glands is unusual in putrid sore-throat, and the characteristic odor of gangrene is almost unmistakable.

Diphtheria presents a specific bacillus, while the organism active in gangrenous sore-throat, if specific, has not been identified.

Retro-pharyngeal Abscess.

What are the symptoms of retro-pharyngeal abscess?

Suppuration in the retro-pharyngeal tissues is most commonly a result of destructive disease of the cervical vertebræ; it may also be due to inflammation of the lymphatic glands or of the connective tissues, resulting from traumatism or developed in the course of infectious disease, or by extension from adjacent disease. The affection is more common in children than in adults, and in tuberculous or syphilitic than in other subjects. There may be an initial chill, with nausea and vomiting, followed by considerable elevation of temperature and acceleration of pulse. In some cases, especially in adults, the disease is insidious in onset and course, and febrile movement is absent.

There are soreness of the throat, with pain and difficulty of swallowing; orthopnea; suffocative paroxysms; and noisy breathing; the voice is muffled or nasal. The head is often thrown back. There is little or no cough. There may be regurgitation of fluids through the nose.

The neck may be swollen and tender to touch, especially behind the angle of the jaw, in front of the sterno-mastoid muscle. The submaxillary glands may suppurate and fluctuation become evident. The diagnosis is to be made by inspection and palpation.

On *inspection* of the throat the posterior wall of the pharynx may appear tumid, or a distinctly circumscribed projection may be seen, the mucous membrane over and around the swelling being reddened, perhaps ecchymotic.

The abscess may be so hidden that the use of the mirror, or digital exploration, may be necessary for its detection.

On *palpation* fluctuation may be elicited.

THE ESOPHAGUS.

Esophagitis.

What are the clinical features of esophagitis?

Inflammation of the esophagus may arise from the ingestion of irritating food, corrosive substances or foreign bodies, or as a secondary manifestation of some primary disorder, local or constitutional. The mucous membrane is thickened, perhaps eroded, sometimes the seat of false membrane. Ulceration may take place. The condition is attended with pain in deglutition and dull pain behind the sternum.

In connection with cardiac insufficiency and cirrhosis of the liver the esophageal veins may be enlarged and varicose, and give rise to hemorrhage, which may be fatal.

Stricture of the Esophagus.

What are the symptoms of stricture of the esophagus?

The esophagus may be narrowed by new-growths in its walls; by cicatrices resulting from previous ulceration or syphilitic or tuberculous disease or from the ingestion of corrosive substances; by pressure from without, as by an aneurism or a new-growth; or as a congenital malformation. The constriction is most commonly near the upper or the lower extremity of the tube. Deglutition is interfered with, so that it may be possible to swallow only liquids. If the narrowing is decided, a pouch forms above the seat of constriction, in which considerable quantities of food accumulate, to be periodically rejected. Esophageal growths, if favorably situated, can sometimes be detected by the methods of laryngoscopy. *Esophagoscopy* has not yet been sufficiently developed for diagnostic purposes. On auscultation to the left of the spinal column the sound generated by swallowed water may be found retarded. Conclusive evidence of the existence of a stricture of the esophagus is furnished by the resistance encountered in the introduction and withdrawl of a bulbous bougie; if aneurism be suspected, this means of exploration is not permissible.

How are functional and organic strictures of the esophagus to be differentiated?

Spasmodic contraction of the esophagus is apt to result from the presence of a foreign body, as a bougie, in the gullet. Spasm of the esophagus (*esophagismus*), in a degree sufficient to give rise to symptoms, sometimes occurs in hysterical persons. Under the conditions last named, food may be obstinately rejected and a considerable degree of emaciation result. The introduction of a bougie may be met with some resistance, which, however, slowly yields to gentle pressure. Careful observation will disclose the fact that not all the food taken is rejected. A powerful impression, judiciously made, may at once cause the disappearance of the symptoms, which occur, without discoverable cause, in a person with other hysterical attributes.

THE STOMACH.

How is the acidity of the gastric contents determined?

If the gastric contents are *acid* from any cause they will redden blue litmus-paper. In the presence of *free acid* they will cause in Congo-red a change to blue, which disappears when heat is applied if the acid be *organic* and persists if it be *hydrochloric*.

What are the most available tests for hydrochloric acid?

Töpfer's test consists in the use of a 0.5% solution of dimethyl-amido-azobenzol, a cherry-red color developing in the presence of free hydrochloric acid.

Günzburg's test consists in the use of a solution of vanillin 1 part, phloroglucin 2 parts, absolute alcohol 30 parts, a carmine-red color resulting on application of gentle heat when free hydrochloric acid is present.

Boas' test consists in the use of a solution of 5 parts of resublimed resorcin, 3 parts of cane-sugar, 100 parts of 94% alcohol, a rose-red or vermilion-red color developing in the presence of free hydrochloric acid when gentle heat is applied.

How is the total acidity of the gastric contents determined?

A known quantity of the gastric contents is titrated with a decinormal solution of sodium hydroxid, phenolphthalein being

used as an indicator, the number of cu.cm. of the solution employed being converted into terms of hydrochloric acid.

How is the presence of lactic acid in the gastric contents determined?

Uffelmann's test consists in the use of a solution of 5 drops of a strong solution of carbolic acid, 5 drams of water, and 2 drops of a solution of ferric chlorid, which gives rise to a canary-yellow color in the presence of lactic acid.

How are the functions of the stomach studied?

By means of an investigation of its *digestive activity*, of its *motor activity* and of its *absorptive activity*. For these purposes test-meals are employed. Ewald and Boas' *test-breakfast* consists of from an ounce to an ounce and a half of wheat-bread and from ten to twelve ounces of water or weak tea without milk or sugar. The gastric contents are examined after an hour. Riegel's *test-dinner* consists of twelve ounces of soup, six ounces of beefsteak, an ounce and a half of wheat-bread and six ounces of water. The contents of the stomach are examined after four hours.

How is the digestive activity of the stomach determined?

A tablet of three or four grains of potassium iodid is enclosed in a bit of thin vulcanized rubber tubing, the ends of which are carefully folded over, and the whole is tied with three or four strands of fibrin. This is swallowed after a test-breakfast and the saliva is tested for iodin, which normally should appear within two hours.

How is the presence of pepsin in the gastric contents demonstrated?

The digestion within a few hours of coagulated egg-albumin, fibrin or serum-albumin by the gastric contents kept at body-temperature indicates the presence of pepsin.

How is the presence of lab-ferment or rennet-ferment in the gastric contents demonstrated?

The coagulation within fifteen minutes of neutralized milk by an equal quantity of neutralized gastric contents kept at body-temperature indicates the presence of lab-ferment or rennet-ferment.

How is the motor activity of the stomach determined?

By the time that elapses after the ingestion of fifteen grains of salol in a gelatin capsule at the height of digestion before the urine yields a violet color on the addition of a drop or two of a neutral solution of ferric chlorid. Normally this takes place within from half an hour to an hour and a quarter.

How is the absorptive activity of the gastric mucous membrane determined?

By the time that elapses after ingestion of two or three grains of potassium iodid carefully enclosed in a gelatin capsule before iodin appears in the saliva, as shown by starch-paper being made blue. Normally this should take place within ten or fifteen minutes.

Neuroses of the Stomach.

What are the neuroses of the stomach?

The functional activity of the stomach may be deranged in the absence of appreciable structural alteration. The disorder may be *motor*, *sensory* or *secretory*.

Thus, there may be *increased motor activity* of the stomach (*hyperkinesis*, *supermobility*), causing premature discharge of the chyle into the duodenum. *Peristaltic unrest* consists in unduly active movement of the stomach (and sometimes also of the intestine) after the taking of food, with borborygmi and gurgling. *Nervous eructations* may occur paroxysmally and be induced by emotional disturbance. *Nervous vomiting* may take place independently of the ingestion of food and be unattended with nausea. It consists rather in regurgitation than in forcible ejection of the contents of the stomach. *Rumination* or *merycism* consists in regurgitation of food, which may then be rechewed. There may further be *spasm or insufficiency of the cardia*, *spasm or insufficiency of the pylorus* and general *atony of the stomach*.

Among the sensory neuroses may be mentioned *hyperesthesia*, *gastralgia* or *gastrodynia*, *bulimia* (excessive hunger), *akoria* (absence of a sense of satiety) and *nervous anorexia*.

The secretory neuroses include *excessive acidity* of the gastric

juice (*hyperacidity, hyperchlorhydria*); *excessive secretion* of the gastric juice (*supersecretion*), which may be *intermittent* or *continuous;* and *deficient acidity* of the gastric juice (*subacidity, anacidity*). With the last there may be associated also *deficiency of the digestive ferments* (*achylia gastrica*).

Splanchnoptosis.

What is splanchnoptosis?

One or more of the abdominal viscera may be unduly free or mobile or variously displaced in consequence of excessive length or relaxation of the supporting ligaments. In addition to the physical signs of visceral displacement and manifestations of digestive derangement neurasthenic symptoms are often present.

Gastralgia.

What is gastralgia or gastrodynia?

Gastric pain is usually symptomatic of inflammation, ulceration or neoplasm. It may, however, occur independently of recognizable structural disease, and is then considered a neurosis—*neuralgia of the stomach.* The pain of neuralgia is spasmodic, of a cutting character, usually rather brief in duration, shooting and shifting in seat, and occurs spontaneously or immediately after the ingestion of food. It may induce and be relieved by vomiting or eructation. It is relieved by heat or pressure. It is not accompanied by vomiting of blood or by the usual manifestations of disordered digestion. Other neurotic symptoms may coëxist.

Acute Gastritis.

What are the symptoms of acute gastritis?

Acute gastritis results from the entrance of irritating matters into the stomach. It varies in degree in accordance with the intensity of the causative irritant. The milder attacks are sometimes called "*acute gastric catarrh*" or "*acute indigestion.*"

Acute gastric catarrh may complicate infectious diseases, or its exciting cause may escape detection. Phlegmonous or suppurative, membranous and mycotic forms of gastritis have also been described.

The *symptoms* are anorexia, nausea, vomiting, often epigastric pain and tenderness; in a severe attack, the *face* is pale and anxious, the *pulse* small and firm; the *skin* may be cold and clammy; there may be moderate elevation of *temperature;* syncope and collapse may occur. The vomited matters consist of the contents of the stomach, mucus, perhaps bile, and sometimes blood. There is often violent retching, without expulsion of the contents of the stomach, or occurring when the organ is empty. In addition there are commonly headache and thirst. The tongue may be red and angry. Perforation of the walls of the stomach may take place. Peritonitis may develop. The intestines are not likely to entirely escape.

How are the gastric symptoms occurring at the onset of acute febrile disorders or in the course of cerebral disease to be distinguished from the symptoms of acute gastritis?

The vomiting of acute, febrile disorders, or of cerebral disease, is not dependent upon the ingestion of food; nor is it necessarily attended with nausea, coated tongue, or pain in the epigastrium. Tenderness is not common. If symptoms indicative of a constitutional affection have not been obtrusive, careful investigation will succeed in detecting them at once, or after the lapse of a variable period. No single symptom, but an association of symptoms establishes the diagnosis.

How are acute gastritis and intestinal obstruction to be differentiated?

The vomiting of intestinal obstruction usually, but not invariably, becomes fecal; the rolling of the obstructed intestines may be apparent through the abdominal walls. The matter vomited in acute gastritis consists largely of mucus, perhaps of some blood, and shreds of mucous membrane; while there is a history of irritation, poisoning, or indiscretion in diet. Obstinate constipation attends obstruction; acute gastritis is rather likely to be associated with diarrhea. If an

obstruction of the bowel be not speedily removed, death results. Recovery from the milder forms of acute gastritis may take place spontaneously.

Chronic Gastritis.

What are the symptoms of chronic gastritis?

Chronic gastritis or *chronic gastric catarrh* is the principal factor in the production of dyspepsia. The symptoms differ from those of acute gastritis not only in degree, but also in character. There are many varieties of the affection but the general phenomena are the same in all. The more or less persistent ingestion of food or drink, improper in quality or excessive in quantity, is the principal cause of chronic gastric catarrh. The disorder manifests itself by nausea, vomiting, impaired appetite, coated tongue, epigastric discomfort, aggravated by the ingestion of food, flatulence, eructations, pyrosis, acidity, bad taste, offensive breath, and palpitation of the heart. There is dull, diffuse abdominal pain, relieved by vomiting. There is also diffuse epigastric tenderness. The vomiting, as a rule, takes place from one-and-a-half to two hours after the ingestion of food. Constipation is common; sometimes there is diarrhea, with undigested food in the stools. Usually there is increased thirst. The complexion is often sallow. Headache is frequent and vertigo not uncommon. There may be loss of flesh and anemia. Mental operations are in many cases temporarily sluggish, depression ("blues") is not rare, and sleeplessness or disturbed sleep and disquieting dreams are quite common.

If much accumulation of undigested and decomposing food takes place, or impairment of the motor activity of the stomach becomes decided, dilatation of the viscus (*gastrectasia*) may result. A succession of meals may be taken before vomiting occurs; when it does, astounding quantities of liquid, containing the macerated and fermenting remains of food, and various fungi (especially the yeast-fungi and sarcinæ), are ejected. The quantity of hydrochloric acid in the gastric juice may or may not be changed. Organic acids of fermentation are usually present.

Dilatation of the Stomach—Gastrectasis.

What are the clinical features of gastrectasis?

Dilatation of the stomach may be acute or chronic. It may follow ingestion of large quantities of food or drink, obstruction at the pylorus or atony of the muscular coat from any cause. In addition to dyspeptic symptoms there occurs at intervals vomiting of large quantities of offensive fluid containing perhaps particles of food taken long before. The upper part of the abdomen may be prominent and the enlargement of the viscus may be appreciable on inspection. The area of gastric percussion-tympany is increased when the stomach is empty and that of dulness when it is filled. Splashing can sometimes be felt and heard. The diagnosis will sometimes be facilitated by inflation of the stomach.

Gastrectasis is to be distinguished from enlargement of the stomach—*megagastria*—and from displacement of the stomach —*gastroptosis*.

Gastric Ulcer.

What are the symptoms of gastric ulcer?

Destruction of the mucous coat of the stomach may be a result of acute gastritis. More commonly, however, gastric ulceration develops in the course of chronic gastric catarrh, especially in anemic women with impaired nutrition. Long-continued hyperacidity may be a contributing cause. In some cases, ulcers of the stomach result from occlusion or obstruction of gastric vessels. Gastric ulcer is not infrequent in cases of cirrhosis of the liver. The symptoms are sometimes obscure, and the disease may go unrecognized until suddenly perforation and death result. Symptoms of gastric catarrh are usually present, with vomiting, impaired appetite, discomfort after meals, flatulence, acidity, coated tongue. There are acute pain and marked tenderness in the epigastrium or hypochondrium, also sometimes in the lower dorsal or upper lumbar region. Pain and tenderness over or near the spinal column opposite the site of epigastric pain are quite characteristic of gastric ulcer. The

pain and accompanying tenderness, anteriorly as well as posteriorly, are most frequently distinctly circumscribed. Pain is aggravated by the ingestion of food, especially solid food, and relieved by vomiting, which usually occurs soon after food is taken. The vomited matters are often blood-streaked; or a considerable quantity of blood, bright-red in hue, or discolored by the gastric juice, is vomited. Following hematemesis the stools contain more or less black, tarry matter. The acidity of the gastric juice, and especially the quantity of free hydrochloric acid, is usually increased. Anemia is a common attendant upon gastric ulcer and there may be considerable loss of weight. Dyspnea and palpitation are its concomitants. There are often great emaciation and profound prostration. Occasionally perforation of the walls of the stomach results; if adhesive inflammation have occurred, adjacent structures, as liver or pancreas, may constitute the floor of the ulcer; otherwise fatal peritonitis ensues. Cicatrization may cause deformity of the stomach and obstruction.

Ulceration of the duodenum arises under much the same conditions as gastric ulceration, although more common in males, and in some instances being related especially to burns of the surface of the body. The pain after eating occurs usually later than with gastric ulcer, the tenderness is prone to be further to the right and lower, and hematemesis is less likely, and melena more likely to take place.

How is chronic gastritis to be distinguished from gastric ulcer?

With ulceration of the stomach are associated the symptoms of chronic gastritis, but in addition there are the exquisite localized epigastric pain and tenderness, which differ from the diffuse soreness of simple gastritis. Hematemesis is usual in ulcer of the stomach and less usual in chronic gastritis of other than alcoholic origin; and the blood in the latter case is not likely to be bright in color or of frequent appearance in the vomit. In gastritis vomiting occurs not only after taking food, but not uncommonly on an empty stomach as well. The vomiting of ulceration is usually brought on by eating. Chronic gastritis responds more readily than gastric ulcer to judicious treatment. The age and sex of the patient sometimes help the

diagnosis, as ulcer is most frequent in young persons and especially in anemic girls.

How are gastric ulceration and gastralgia to be differentiated?

The one is and the other is not attended with hematemesis and possibly melena. The pain of gastralgia is more likely to be relieved and that of ulcer to be aggravated by the taking of food and by pressure. Dyspeptic symptoms, anemia, impaired nutrition and increased acidity of the gastric juice are all more common with ulcer than with gastralgia.

Carcinoma of the Stomach.

What are the symptoms of carcinoma of the stomach?

Carcinoma of the stomach appears principally in two forms. In the one the cellular element predominates; in the other, the fibrous. The former involves the coats of the body of the stomach; the latter, almost exclusively the pylorus. Either may, however, be circumscribed or diffuse. The *symptoms* occasioned differ in each case. When the body of the stomach is involved the symptoms are those of an aggravated chronic gastritis: impaired appetite, pain after meals, sometimes vomiting, with slow emaciation. The hydrochloric acid of the gastric juice is usually diminished and sometimes wanting; and lactic acid is often present. Among the many bacteria found in the gastric contents a long, non-motile bacillus has been isolated and is believed to be responsible for the production of lactic acid. Often there is febrile elevation of temperature. As time goes on, ulceration takes place in the neoplasm, with the addition of the symptoms of this condition: severe, deep-seated pain in the region of the epigastrium, aggravated by the ingestion of food, vomiting of blood-streaked matter and of discolored blood presenting an appearance of coffee-grounds. The stools contain tarry matter, from disorganized blood. With the infiltration of the walls of the stomach, adhesions are formed with adjacent organs, which may be progressively invaded. Perforation may take place.

When the new-growth involves the pylorus, a characteristic feature is dilatation of the stomach. The organ may be considerably displaced by the weight of the tumor. Emaciation ensues; while the development of the carcinomatous cachexia is of diagnostic significance. The feature without the establishment of which the diagnosis of carcinoma of the stomach is doubtful is the presence of a tumor in the epigastrium. In scirrhus of the pylorus death results from exhaustion, as a result of inanition; in cellular carcinoma metastasis plays a prominent part in determining a fatal issue.

How are carcinoma and ulceration of the stomach to be differentiated?

The similitude between the symptoms of ulcer of the stomach and those of carcinoma of the stomach is sometimes so great that the diagnosis is extremely difficult. The detection of a tumor in the area occupied by the stomach is strong evidence, although a cicatrized gastric ulcer may present like signs. In its absence, a cachexia or the detection of enlarged glands or new-growths at other parts of the body may afford corroborative evidence. While the symptoms of ulceration may set in acutely, and rapidly assume a grave aspect, they may on the other hand be comparatively mild; recovery is the rule. The symptoms of carcinoma are more slowly developed, more persistent and more profound than those of ulcer, and the progress of the case reveals its malignancy. If the pylorus is involved, vomiting does not take place for some time after food has been taken, while both in ulceration and in cellular carcinoma of the body of the stomach vomiting, when it occurs, takes place early. Obstruction of the pylorus by carcinoma occasions dilatation of the stomach. Before the fortieth year of life ulceration is the more common; after forty, carcinoma. In ulcer, the gastric ulcer hydrochloric acid is often excessive; in carcinoma, often wanting. In simple ulcer, sudden, more or less profuse hemorrhages occur; hence the blood is often bright, and when discolored it is a viscid fluid or coagulated in coherent clots. In carcinomatous ulcer there is a slight but more or less continuous oozing; hence vomiting of blood rarely occurs apart from the

admixture with food of the disorganized "coffee-grounds" sediment.

How is carcinoma or sarcoma of the omentum to be distinguished from carcinoma of the stomach?

The detection of a tumor in the region of the epigastrium is not exclusively indicative of carcinoma of the stomach. The omentum may be the seat of carcinoma or sarcoma. In such a case the symptoms of gastritis are not necessarily present; in particular the vomiting of matter resembling coffee-grounds is wanting.

How is carcinoma of the stomach to be distinguished from carcinoma of the pancreas?

It may be impossible to decide from anatomic considerations, especially in view of the displacement that often occurs, whether a tumor in the epigastrium is gastric or pancreatic. If the former, definite symptoms of gastric derangement are present; the visceral symptoms of the latter are ill-defined. The existence of diabetes and the imperfect digestion of fats would point to involvement of the pancreas. Jaundice sometimes accompanies carcinoma of the pancreas from pressure upon the bile-ducts. In gastric carcinoma without hepatic complication this does not occur.

How is chronic gastritis to be distinguished from carcinoma of the stomach?

It is not sufficient to make a diagnosis of chronic gastritis. The existence of carcinoma should, if possible, always be excluded. If, in addition to the symptoms of gastritis, there is severe and persistent pain in the epigastrium, or pain increased or developed after eating, with vomiting shortly after meals or after a variable interval, the ejected matters resembling coffee-grounds; if there exist cachexia, new-growths in various parts of the body, and a tumor can be detected in the region of the epigastrium, the gastritis is but a concomitant of the malignant disease, which, in the course of a year or eighteen months, is almost necessarily fatal. Carcinoma is uncommon before forty; gastritis may occur at any time of life. The gastric juice is

likely to contain free hydrochloric acid in nearly all cases of chronic gastritis, but there is usually none in cases of gastric carcinoma.

THE INTESTINES.

Acute Enteritis.

What are the symptoms of acute enteritis?

Acute inflammation of the small intestine may vary greatly in severity, from a simple catarrh of the mucous lining to an intense inflammation involving the submucous and muscular tunics and even the peritoneal investment.

Acute intestinal catarrh or *mucous enteritis* is, in most cases, dependent upon the presence of irritating matters in the bowel, introduced from without or generated within. As a result, there occur diarrhea, with colicky pains, and often considerable tenderness in the abdomen, and febrile and other constitutional manifestations of varying intensity. The *stools* may number from three to six or more in twenty-four hours; they are thin and liquid, containing undigested food and considerable mucus, and may be streaked with blood. The stomach is likely to share in the inflammatory process, and the symptoms of a more or less intense gastritis are superadded.

When *gastro-enteritis* is due to the ingestion of corrosive and poisonous substances or to the ingestion or development of ptomaïnes, the symptoms become more severe, and collapse and death may occur.

In the severe cases of *acute enteritis*, involving the serous, muscular and submucous coats, constipation, or constipation alternating with slight, irritative diarrhea, is the rule; actual obstruction of the bowel, from inflammation, paralysis, or incarceration by bands of lymph, is not uncommon. The constitutional and local symptoms are correspondingly intense. There may be an initial chill. While the pain may be colicky at first, it soon becomes constant, subject, however, to paroxysmal exacerbations. It is increased by pressure, tenderness at times being exquisite. As in peritonitis, the patient lies upon his back

with flexed thighs, to relax the abdominal muscles. There is not uncommonly a marked and distressing pulsation to the right of the umbilicus. Thirst, nausea, vomiting and retching may be decided, even in the absence of gastric involvement. The *fever* becomes high and does not remit, as it does in intestinal catarrh. The *pulse* is rapid ; at first tense and full ; afterward, small and wiry. Following a copious stool, amelioration and recovery may ensue ; or symptoms of failing circulation, with distention of the abdomen, hiccough, incessant retching, sweating, anuria and exhaustion, may precede death.

How are acute catarrhal enteritis and typhoid fever to be differentiated ?

Some degree of inflammation of the small intestine necessarily attends typhoid fever. Occurring as an independent affection, however, enteritis is wanting in the epistaxis, the severe headache, the rose-spots, the characteristic temperature-curve, the peculiar character of the stools, the diazo-reaction, the agglutinating reaction, the gravity and the typical course and duration of the general infectious disease.

Membranous Enteritis.

What is membranous enteritis ?

Membranous enteritis, tubular diarrhea or mucous colic, is a somewhat uncommon disease, characterized by the discharge of mucoid casts of the bowel, or of flakes of false membrane. These are usually expelled, after paroxysms of colic, with painful straining, in watery stools containing mucus, sometimes blood, but, as a rule, little fecal matter. The paroxysms are often preceded by constipation, and associated with or followed by diarrhea. Nervous symptoms are common. The affection is obstinately recurrent. It occurs chiefly, if not exclusively, in hysterical or hypochondriacal subjects in early adult or middle life, and principally in women. Enteroptosis has been noted as an associated condition, and gastric achylia has been present in some cases.

Cholera Morbus.

What are the symptoms of cholera morbus or cholera nostras?

Cholera morbus or *cholera nostras* is an acute disease, most prevalent during the summer months and characterized by inflammation of the stomach and intestines. It is commonly a result of the ingestion of unsuitable and irritating articles of diet. The affection is manifested by colicky pains in the abdomen, by nausea, vomiting and diarrhea, by cramps in the legs, by increased thirst, by headache, vertigo and debility, by coldness of the extremities, by prostration and rapid wasting. Despite the severity of the symptoms, recovery is almost invariable.

How are cholera morbus and cholera Asiatica to be differentiated?

The symptoms of cholera morbus and those of cholera Asiatica differ principally in degree. Cholera Asiatica, however, occurs in epidemics; isolated cases are rare. Cholera morbus, on the other hand, is not an epidemic disease. The mortality from cholera Asiatica is high; recovery from cholera morbus is the rule. The detection of characteristic comma bacilli in the feces or in the vomit confirms a diagnosis of cholera Asiatica.

Cholera Infantum.

What are the symptoms of cholera infantum?

Cholera infantum is practically gastro-enteritis in children. It is a disease of the summer months, and is intimately related with heat, foul air, uncleanliness and fermentation of food. It is manifested by vomiting, profuse watery, often fetid diarrhea, fever, rapid wasting, depressed fontanel, convulsions and coma. Among the poor the fatality of the disease is great.

Chronic Enteritis.

What are the symptoms of chronic enteritis?

As *chronic enteritis* or *intestinal catarrh* most frequently results

from persistent errors in diet, the symptoms depend upon retarded and defective intestinal digestion. Pain and oppression occur at a time after the ingestion of food that varies with the seat of the morbid process. In duodenal catarrh the symptoms appear earlier than when the jejunum or ileum is involved. When the lower portion of the intestine is affected, colicky pains are frequent. There may be persistent diarrhea, or obstinate constipation, or diarrhea alternating with constipation. The stools are slimy and may contain undigested food. The *abdomen* is distended; flatulence and belching are usual. The *complexion* is sallow. The *nutrition* is impaired. Headache is common. *Sleep* is disturbed by dreams, while there is unusual drowsiness.

Acute Dysentery.

What are the symptoms of acute dysentery?

Acute dysentery is practically an inflammation of the large intestine, probably dependent upon a specific infection; rarely the lower portion of the small intestine is involved. The inflammation may be of varying intensity. It may be catarrhal, ulcerative or diphtheritic. There are slight fever, abdominal pains of a griping character (tormina), terminating with frequent, small, slimy stools, streaked with blood, each evacuation being attended with bearing-down, burning pain and muscular spasm (tenesmus). The stools sometimes contain pus, sometimes shreds of membrane; in many cases the ameba coli is found. In addition there are headache, vertigo, weakness, thirst and perhaps nausea and vomiting. Dysentery is a prolific source of hepatic abscess. Death may result from exhaustion, or from perforation of the bowel.

How are intussusception of the bowel and acute dysentery to be differentiated?

In cases of intussusception of the bowel the stools may be small, frequent, mucous, blood-streaked, and attended with tenesmus. Inquiry, however, will elicit a history of abrupt onset; examination will disclose the presence of a sausage-shaped abdominal tumor, perhaps also protrusion of the bowel

at the anus. Intussusception is the more common in children; dysentery, the more common in adults.

How are acute enteritis and acute dysentery to be differentiated?

The pain of acute intestinal catarrh is colicky, but the peculiar tormina and tenesmus of dysentery are wanting. The stools of dysentery are more frequent than those of enteritis, are smaller in quantity, contain more blood, and are in less degree fecal. Dysentery is more likely than is enteritis to be epidemic.

How are acute dysentery and typhoid fever to be differentiated?

While both acute dysentery and typhoid fever are attended with diarrhea, the stools of dysentery are frequent and small, perhaps ineffectual, and are composed principally of blood-streaked mucus; the stools of typhoid fever are not necessarily frequent, are larger, often thin and like pea-soup. The epistaxis, the rose-spots, the diazo-reaction, the agglutinating reaction and the characteristic temperature-curve of typhoid fever are wanting in acute dysentery. The duration of dysentery is briefer than that of typhoid fever.

Chronic Dysentery.

What are the symptoms of chronic dysentery?

Dysentery may from the outset manifest a tendency to chronicity; or an acute dysentery may become chronic. The symptoms of chronic dysentery differ in several respects from those of the acute disease. Tormina and tenesmus are uncommon, except with exacerbations. The bowels are loose, the stools containing mucus, but little blood and membrane. Sometimes diarrhea and constipation alternate. Febrile symptoms are generally wanting; intermissions occur; wasting takes place; anemia develops; the complexion becomes sallow; hepatic abscess may form.

Typhlitis—Appendicitis—Perityphlitis.

What are the symptoms of typhlitis?

Inflammation occurs in the cecum or the vermiform appendix as a result of accumulation and impaction of fecal matters; or from irritation by a calculus, the nucleus of which may be inspissated mucus from catarrh of the appendix, or by a foreign body, such as a cherry-stone or a grape-seed. Traumatism and straining are occasional exciting causes. It is probable that bacteria play an important exciting part. The process may be catarrhal, obliterative, ulcerative or gangrenous. The premonitory symptoms are vague and are frequently mistaken for simple colic. Diarrhea may alternate with constipation, still further misleading the inattentive observer. If the cecum be involved, it becomes paralyzed and distended with accumulated feces; thus, the condition declares itself primarily by the symptoms of intestinal obstruction, by pain, often severe, and increased by motion, in the right iliac fossa and right hip, with tenderness, a sense of doughy induration, and dulness on percussion. The tumor is superficial and sausage-shaped, its long axis pointing inwards and downwards. It is slightly movable; gurgling may sometimes be developed. The pain may be paroxysmal or paroxysmally aggravated, and is frequently of an agonizing character. There is usually some fever, and at times the temperature may reach from 102° to 104° F. Peritonitis may develop, even without ulceration and perforation of the bowel, and its symptoms then predominate. It may remain localized or become diffused. With relief to the obstruction, recovery may ensue, or collapse may suddenly occur, recovery or death following.

If *appendicitis* alone exist there may be no interference with the passage of the intestinal contents, though constipation often exists; and the evidences of a tumor in the iliac fossa are wanting, unless the appendix have undergone great distention. There are, besides, colicky pain in the right iliac fossa, with localized tenderness most commonly at a point where a line from the anterior superior iliac spine on the right to the umbilicus crosses the outer border of the rectus muscle; usually

elevation of temperature; and nausea and vomiting. The right rectus muscle is usually rigid, "on guard," especially in the iliac region, and sometimes the enlarged appendix can be felt through the abdominal wall. Sometimes epigastric pain and tenderness, with or without vomiting, are the initial and most prominent symptoms. In ulcerative appendicitis, perforation not rarely results, giving rise to a general or localized purulent peritonitis. This event may be announced by sudden pain, shock, chill and rise of temperature, followed by the development of a fluctuating tumor indicative of abscess; or the condition may develop insidiously and be difficult of recognition. Sometimes, after initial shock and pain, deceptive improvement is manifested. Sudden cessation of pain, and of fever may denote the occurrence of gangrene. In cases suggestive of appendicitis surgical exploration may be a diagnostic necessity. Attacks of appendicitis are prone to be repeated.

Perityphlitis, or inflammation of the tissues surrounding the cecum and its appendix, usually occurs in the course of typhlitis or of appendicitis. The fibrous structures and the peritoneum are involved alone or in association. If perforation takes place, an abscess may form, or general peritonitis result, or both complications may be present. As a rule, the pus is shut off from the general peritoneal cavity by a capsule. The symptoms vary with the pathologic association. To those of the primary condition are added an acute exacerbation of pain and tenderness, as well as of the general symptoms, perhaps preceded by a chill. The pain is deep-seated and is increased by flexing the right thigh upon the abdomen. Sometimes the patient is unable to lift the right leg. He usually lies upon the right side, with the thigh semiflexed. If an abscess forms, there may be repeated rigors and a fluctuating tumor in the right iliac fossa. The tumor is not superficial and sausage-shaped like that of cecitis, but is deep-seated and irregular. A pericecal abscess may sometimes be detected by rectal exploration. If peritonitis develop, death may ensue, from septicemia or gradual exhaustion, or suddenly, with manifestations of collapse.

With what conditions may appendicitis or perityphlitis be confounded?

Inflammations of the cecum and its appendix or inflammation and abscess of the surrounding tissue have been mistaken for typhoid fever, and for "idiopathic peritonitis."

The mistake is more likely to be made when the inflammation of the appendix has been slow and the symptoms indistinct, until perhaps perforation occurs, causing a limited abscess or a general septic peritonitis. Deep-seated but limited abscess may for a time be concealed from other than surgical exploration, and through septic poisoning give rise to the "typhoid state."

Before localizing symptoms, such as induration, or the presence of a fluctuating tumor in the iliac fossa, render the case clear, the occurrence of one or more rigors, the absence of distinctive characteristic symptoms of typhoid fever, the course of the temperature, and sometimes the location of the tenderness, should prevent mistake. The location of the pain and tenderness, however, may be misleading, as the appendix varies greatly in its position; and appendicitis has even been known to simulate hepatic disease. Sometimes only an exploratory incision can settle the diagnosis.

The exploring needle or aspirator should never be used.

How are perityphlitis and typhlitis to be distinguished from a lumbar abscess?

Destructive disease of a lumbar vertebra is usually followed by suppuration, the pus following in the course of the psoas muscle and seeking exit below Poupart's ligament. A collection of pus forming in this way differs from typhlitis or perityphlitis by the absence of symptoms of intestinal derangement. The situation of the tumor is different in each case. In the one case, examination will reveal a deformity of the spinal column, with pain and tenderness in the lumbar region. The symptoms are slow and progressive. They may be associated with visceral tuberculosis.

How are typhlitis and perityphlitis in a woman to be distinguished from an abscess of the right ovary?

An abscess of the right ovary is situated nearer the middle line than is the swelling of typhlitis or perityphlitis. With typhlitis

or perityphlitis is associated gastro-intestinal derangement; with abscess of the ovary, uterine and menstrual derangement. Vaginal and rectal examination may clear up any doubt.

How are typhlitis and perityphlitis to be distinguished from carcinoma of the cecum?

The symptoms of typhlitis and perityphlitis are likely to appear suddenly; those of carcinoma insidiously and progressively. Typhlitis and perityphlitis are affections of comparatively brief duration; carcinoma of the cecum will probably continue for a number of months after its detection. The inflammatory processes are usually attended with fever; carcinoma is not. When typhlitis or perityphlitis has existed for some time, fluctuation—indicative of the occurrence of suppuration—can be detected in the tumor; the carcinomatous new-growth retains its original density. Perityphlitis or typhlitis may be attended with emaciation and sallowness of skin, but not with the cachexia of carcinoma. When the cecum is the seat of carcinoma, like new-growths are usually found in other parts of the body.

Intestinal Obstruction.

What are the symptoms of intestinal obstruction?

The lumen of the bowel may be obliterated by an accumulation of feces, by a large gall-stone, by an intestinal calculus (enterolith) or other foreign body, by organic narrowing of the bowel, by stricture or neoplasm, by a twist or volvulus, by external or internal constriction and by incarcerated or strangulated hernia, internal or external.

The condition may arise in a subject of habitual constipation or of hernia. It may follow a violent physical effort. It may be due to acute enteritis. From the onset, or after a variable period during which no stool has been passed, abdominal pain and rumbling set in. Vomiting ensues; at first of the contents of the stomach, then of yellowish-green fluid and mucus; finally the vomiting becomes stercoraceous. The apparent constipation does not submit to ordinary measures. The abdomen becomes distended. The expression of the face is drawn and

anxious, the pulse small, rapid and feeble, the surface cold and clammy, and if the condition be not relieved by medicine or operative intervention, death is the inevitable result.

When the obstruction is not complete, as in some cases of fecal impaction, there may be more or less frequent passages of liquid matters somewhat fecal, which the patient will describe as diarrhea. Inspection of the stools, and the evidences given by palpation and percussion, of the presence of a hard mass in the course of the bowel, usually in the transverse or descending colon, will prevent error. The nature of the obstruction in any case is to be determined principally by physical examination. In cases of strangulated hernia the knowledge of the existence of hernia may assist the diagnosis. In the absence of such history, examination must none the less be made, if only to exclude that condition from among the possibilities in the case. Incarceration may result from unusual exertion or indiscretion in diet. Often the pain is colicky and referred to the umbilicus.

Intussusception.

What are the symptoms of invagination or intussusception of the bowel?

Under certain circumstances, not definitely recognized, one portion of the bowel becomes invaginated in another portion. The small intestine or the large intestine, respectively, may be alone involved; but most commonly the small intestine enters the large at the ileo-cecal orifice. The occurrence of the accident is announced by a sudden attack of pain, repeated in paroxysms, followed by the presence of a sausage-shaped tumor in the abdomen, and stools of a dysenteric character. Sometimes no fecal matter is passed, and there are frequent discharges of blood-stained mucus. The pain is intense, and the child (for the affection is most common in children) often draws up its legs close to the belly. Pressure and manipulation relieve the pain, and quiet the excruciating cries of agony. In the course of a variable period of time, blood is passed by the bowel, the pain becomes continuous and vomiting occurs, with the symptoms of intestinal obstruction. The invaginated bowel

may be accessible to rectal examination; it may even protrude from the anus; it may slough and be detached, and recovery ensue; or it may occasion stenosis of the bowel. Intussusception is more common in children than in adults, and in boys than in girls.

How are intussusception and obstruction of the bowel to be differentiated?

Intussusception is the more common in children; obstruction, in adults. The symptoms of intussusception appear abruptly; those of obstruction are often abrupt in onset, though sometimes of long standing. A sausage-shaped tumor is characteristic of intussusception; certain varieties of obstruction are attended with abdominal tumors of irregular shape. Constipation is not so absolute in intussusception as in obstruction; and stercoraceous vomiting is less common in the former than in the latter. A discharge of blood and the protrusion of a portion of bowel from the anus are diagnostic of intussusception. Digital exploration of the rectum will often assist in discrimination.

How are typhlitis and intussusception of the bowel to be differentiated?

Both typhlitis and intussusception may present a sausage-shaped tumor in the right iliac fossa and be attended with severe pain and the evidences of intestinal obstruction; but intussusception, unlike typhlitis, is sudden in onset, is uncommon in adults, is usually afebrile, is likely to be attended with ineffectual or bloody stools, and perhaps to be accompanied by protrusion of the bowel at the anus. The tumor of intussusception is not necessarily confined to the right iliac fossa.

Carcinoma of the Intestine.

What are the clinical manifestations of carcinoma of the intestine?

Carcinoma of the intestine is most commonly situated in the rectum, the sigmoid flexure, the cecum, the vermiform appendix or the duodenum. When in the duodenum, the papilla of the pancreatic duct and common bile-duct is usually involved and

jaundice is apt to result. When other parts of the bowel are involved, in addition to the pain and constitutional phenomena occasioned by the malignant growth, symptoms of partial intestinal obstruction develop. There is obstinate constipation, and when the bowels are moved, the stools appear as thin, flat bands, often streaked with pus and blood. On physical examination a tumor may be detected.

Intestinal Parasites.

What are the most common varieties of intestinal parasites?

The most common intestinal parasites belong to the order of vermes, of which there are two important classes—*cestodes*, or tape-worms, and *nematodes*, or round-worms. The distomata are *trematodes*, or sucking-worms.

Of the former the more important are the *tænia solium*, the *tænia mediocanellata*, and the *bothriocephalus latus*.

Of round-worms the most common are the *ascaris lumbricoides*, the *oxyuris vermicularis*, the *trichina spiralis*, the *filaria sanguinis hominis*, the *dracunculus medinensis*, and the *ankylostoma duodenale*.

The oxyuris vermicularis inhabits the large intestine, the tape-worms, the lumbricoides and the ankylostoma the small intestine; the trichina migrates from the stomach and small intestine into the muscles, setting up an irritative fever with special symptoms, the condition being known as *trichiniasis*.

What symptoms are occasioned by the presence of animal parasites in the intestinal canal?

The presence of worms in the intestinal canal of an otherwise healthy individual may occasion no appreciable disturbance. In other cases, however, there are evidences of gastro-intestinal derangement, capricious appetite, abdominal uneasiness, colicky pains, possibly diarrhea, nausea, vomiting, loss of flesh, debility, cachexia, irregular fever, disturbed sleep, gritting of the teeth, itching of the nose and anus, nervous manifestations, even epileptiform convulsions. The filaria gains entrance to the lymph or the blood, where it causes obstruction and sets its embryos free. The dracunculus wanders from the intestine to the sub-

cutaneous and intermuscular connective tissue. The sucking-worms cause profound anemia by abstraction of blood.

The diagnosis of intestinal worms depends upon the discovery of the parasites or of their ova in the stools.

Tænia Solium.

What are the characteristics of the tænia solium?

The *tænia solium* (Fig. 24) is a tape-worm having a small head, or scolex, and a slender neck. The head is of the size of a pin-head, and is surmounted by a circle of twenty-six hooklets, around which are four suckers. From the neck pass off segments, or proglottides, that progressively increase in size. The entire worm, or strobila, may be from seven to ten feet long and composed of from four hundred to six hundred segments. The tænia

FIG. 24.



Head. Mature segment. Ovum.

Tænia Solium, magnified (Heller).

solium usually develops in man from the ingestion of "measly" pork or the flesh obtained from swine infected with cysticerci cellulosæ, which in turn develops in animals that have swallowed the ova of the tænia solium. When ova of the tænia solium gain entrance into the stomach of man their capsules are dissolved and the embryo parasites invade the muscles, the brain, the eye and other parts, where they become encapsulated or encysted, constituting *cysticerci cellulosæ*.

Tænia Mediocanellata.

What are the characteristics of the tænia mediocanellata?

The *tænia mediocanellata*, or *unarmed tape-worm* (Fig. 25) differs from the tænia solium in that the head though surmounted by four suckers, is without hooklets. The tænia mediocanellata may attain a length of from ten to twenty feet, and be composed of from eight hundred to more than one thousand segments, which are longer and broader than those of the tænia solium.

The tænia mediocanellata is transmitted to man by the raw flesh of sheep or cows, in which hosts the larvæ develop from the ova of the tænia mediocanellata.

Tape-worms are usually present in small numbers; there is often but a single worm; sometimes there are two or three. About three months are requisite for the development of a tape-worm.

FIG. 25.



Head. Mature segment. Ovum.

Tænia mediocanellata, magnified (Heller).

Bothriocephalus Latus.

What are the characteristics of the bothriocephalus latus?

The *bothriocephalus latus* (Fig. 26) sometimes called the *tænia*

lata, is a cestode worm, with a club-shaped head and a filament-

Fig. 26.



Head. Mature segment. Ovum. Ovum: embryo developed.

Bothriocephalus latus, magnified (Heller).

ous neck. On either side of the head is a longitudinal sucker. The mature segments present a characteristic stellate appearance, dependent upon the distention of the uterus with ova. The worm may be from fifteen to twenty-five feet in length, and constituted of from three thousand to four thousand segments. It is thought to be derived from fish or fresh-water molluscs.

Ascaris Lumbricoides.

What the characteristics of the ascaris lumbricoides?

Lumbricoid or *round worms* inhabit the small intestine. Small worms, embryos, or ova, are supposed to gain access to the alimentary canal of man through drinking water, and in some instances to be conveyed by the fingers to the mouths of those engaged in cleansing privies or otherwise handling excrement, but the exact history of their development is unknown.

Mature worms are from eight to fifteen inches in length, are attenuated at both extremities, and resemble common earthworms. Lumbricoid worms may be present in varying numbers, from one to a dozen or more. They are usually multiple, and in rare instances may be so multitudinous as to occlude the intestine. The worms may find their way into the stomach, and

be expelled by vomiting. They may pass from the esophagus into the larynx and trachea. They have in this way caused suffocation in children. They may cause occlusion of the bile-

FIG. 27.



a, natural size; *b*, head, magnified; *c*, ovum, magnified.
Ascaris lumbricoides (v. Jaksch).

duct or pancreatic duct, and, though rarely, suppuration of the liver or of the pancreas. They may also leave the intestine by way of a perforation, causing peritonitis or fecal abscess.

Ankylostomiasis.

What is ankylostomiasis?

Ankylostomiasis is a condition arising from the presence in the intestial tract of the *ankylostoma duodenale*, a nematode worm from ¼ to ¾ inch long, with a mouth provided with tooth-like

hooks, by means of which the parasite attaches itself to the mucous membrane of the upper portion of the small bowel, and thus withdraws blood by suction. In addition to anemia, with dyspnea, edema, wasting, and debility, there may be colicky abdominal pains and diarrhea. The oval eggs appear in the stools. *Infection* takes place through drinking-water.

Distomiasis.

What is distomiasis?

Distomiasis is a condition due to the presence in various structures of the different varieties of distomata, trematode sucking-worms.

The *distoma hepaticum* is a liver-fluke, about 1 inch long and ½ inch wide, that finds its way into the biliary passages, the inferior vena cava and the intestine especially in ruminants, and rarely in man.

The *distoma lanceolatum* is about ⅓ inch long and ¼ as wide. It also is found in the biliary passages, but is not common in man.

The *distoma hæmatobium* is from ½ to ⅔ inch long, with a tendency to a cylindrical shape. It is found in the trunk and branches of the portal vein, the splenic vein, the mesenteric veins and the vessels of the rectum and the bladder. The ova penetrate the mucous and submucous membranes of the tubes, the bladder, the rectum, at times the liver, the lungs, the kidneys and the prostate, giving rise to inflammation, ulceration, etc. The disease is common in Egypt and Abyssinia in both monkeys and man.

Occlusion of the Mesenteric Vessels.

What are the clinical features of occlusion of the mesenteric vessels?

Thrombosis or embolism of the mesenteric arteries or veins

may arise from various causes, such as valvular disease of the heart, atheroma or inflammation of the vessels, arteriosclerosis, etc. The resulting symptoms vary somewhat with the rapidity of occurrence of the disturbance. They may simulate intestinal obstruction and consist in severe abdominal pain, with vomiting that may become stercoraceous or bloody, tympanites and signs of collapse. There may, however, be diarrhea with bloody stools. A tumor may sometimes be detected upon palpation. The bowel undergoes ulceration or gangrene.

Oxyuris Vermicularis.

What are the characteristics of the oxyuris vermicularis?

Oxyures vermiculares, *seat-worms*, *thread-worms*, or *spool-worms*, are from one-eighth to one-half of an inch long. They infest the large intestine, especially the rectum. The worms find their way out of the anus, and give rise to intense itching.

FIG. 28.



1, Female; 2, males. Ovum, magnified.
Oxyuris vermicularis, natural size (Vierordt).

Sometimes they gain access to the vagina, and occasion unpleasant symptoms. The parasites may be present in the bowel in large numbers; they are often found in the stools in tangled masses, resembling bunches of thread. Infection takes place probably through drinking-water or green vegetables.

Acute Peritonitis.

What are the symptoms of acute peritonitis?

Acute peritonitis arises from the activity of irritants either conveyed through the blood or resulting from local disease. It may thus be caused by nephritis, rheumatism, tuberculosis, other infectious diseases, inflammatory, ulcerative or suppurative processes in any of the abdominal viscera, traumatism and operative procedures.

Acute peritonitis may be *local* or *general;* it may be *fibrinous, sero-fibrinous, purulent, putrid, hemorrhagic* or *dry.* It is attended with acute abdominal pain, tympanites, nausea, vomiting, constipation, considerable elevation of temperature and other febrile manifestations. The *pulse* is small, rapid and tense—"wiry." The *face* is drawn; the *expression* anxious.

The *breathing* is shallow, rapid, and thoracic. The abdominal pain and tenderness are so intense that the patient shrinks from the slightest movement, and complains of the lightest covering. The legs and thighs are drawn up in flexion, to relax the abdominal parietes. The surface of the body may be covered with a cold sweat and collapse may ensue. Effusion may take place or adhesions form among the structures in the peritoneal cavity. Should pus form, there are repeated rigors and hectic fever. If the inflammation is putrid, death speedily takes place amid the symptoms of profound intoxication.

How are acute gastritis and acute peritonitis to be distinguished from one another?

Nausea, vomiting, constipation, headache, abdominal pain and tenderness, and febrile symptoms attend both acute gastritis and acute peritonitis. In the former, however, the vomiting occurs earlier and is more aggravated than in the latter, and the vomited matter may contain blood. In peritonitis, the abdominal pain and tenderness are not confined to the epigastrium, but are more extensive and more intense than in gastritis, while abdominal distention is more decided. A cause will be obvious for an acute gastritis sufficiently intense to simulate peritonitis, the symptoms of which are relatively the more profound.

How are acute peritonitis and acute enteritis to be differentiated?

The pain of enteritis is colicky; that of peritonitis, lancinating. Tenderness is greater and more general in peritonitis than in enteritis. Diarrhea is common in enteritis; constipation is the rule in peritonitis. Nausea and vomiting are more decided in peritonitis than in enteritis. Effusion occurs in peritonitis, not in enteritis. The constitutional disturbance of peritonitis is comparatively more profound than that of enteritis. Rigors and fluctuating temperature are suggestive of peritonitis. The cause of peritonitis is sometimes obvious in the history of the case, or discoverable upon vaginal or other examination.

How are acute intestinal obstruction and acute peritonitis to be differentiated?

Constipation attends both acute peritonitis and acute intestinal obstruction; it may yield in the one, but it is insuperable in the other. Prior to the acute symptoms of obstruction there may have been small, liquid evacuations, but, when the symptoms of obstruction have set in, constipation is absolute. The vomiting of acute peritonitis does not present any unusual features; that of intestinal obstruction soon becomes stercoraceous. The pain of obstruction is colicky; that of peritonitis is sharp and lancinating. The exquisite abdominal tenderness of peritonitis is not encountered in obstruction. When obstruction exists, the rolling of the intestines may be apparent to the eye or to the palpating hand. Peritonitis occasions paralysis of the bowel. The febrile symptoms of peritonitis are wanting in uncomplicated obstruction.

Inquiry and physical examination may elicit one of the known causes of peritonitis or of obstruction respectively.

How are acute peritonitis and intestinal colic to be differentiated?

Acute peritonitis and intestinal colic have pain in common; the latter, however, is unattended with febrile manifestations.

The pain of colic is inconstant and is relieved by frictions; in peritonitis the pain persists and is intensified by the slightest touch.

Colic, as a rule, is symptomatic, and its cause is to be ascertained upon careful search. A blue line on the gums would indicate lead-poisoning.

How is subacute rheumatism involving the muscles of the wall of the abdomen to be distinguished from acute peritonitis?

The pain of rheumatism is not so severe as that of peritonitis; nor is it as constant; nor does it give rise to symptoms of gastro-intestinal derangement; and it is not attended with febrile manifestations; while in addition there are other evidences and a history of exposure to influences productive of subacute rheumatism.

Chronic Peritonitis.

What are the symptoms of chronic peritonitis?

Chronic inflammation of the peritoneum may be the sequel of an acute attack; it may be insidious in development. It is commonly a result of persistent irritation, such as may depend upon the presence of new-growths, carcinomatous, sarcomatous or tuberculous, or upon chronic inflammation of the abdominal or pelvic viscera.

Adhesions form between adjacent viscera and fluid collects in the abdominal cavity. Paroxysmal attacks of pain occur. The abdomen is distended. There may be nausea and vomiting.

The action of the intestines is interfered with and constipation usually results. In tuberculous cases, with concomitant ulceration of the intestine, there may be intercurrent or continuous diarrhea. In proportion to the chronicity of the attack the omentum is found indurated and rolled up close to its attachment to the stomach, the mesentery is shortened and the lumen of the bowel is narrowed, producing visible and palpable distortions and prominences.

There is commonly more or less fever, sometimes hectic, especially if the effusion be purulent. Sometimes, particularly in tuberculous disease, there are recurrent febrile exacerbations. In any case there occur gradual emaciation and loss of strength,

breathlessness, edema of the lower extremities; spontaneous recovery sometimes happens, but as a rule, unless relief be given, surgically or medicinally, death ensues. The cases are often protracted.

Malignant disease and tuberculosis of the peritoneum are most frequently associated with similar conditions in other structures.

How are chronic peritonitis and malignant disease of the liver to be differentiated?

Malignant disease of the liver, like chronic peritonitis, may give rise to a tumor in the upper portion of the abdomen, and to ascites; but the physical signs of enlargement fuse with those that normally belong to the liver; while the percussion-dulness dependent upon an omentum rolled up and contracted by chronic disease is separated from the hepatic dulness by an area of tympanitic resonance.

Tabes Mesenterica.

What are the symptoms of tabes mesenterica?

In predisposed children the glands of the mesentery sometimes become tuberculous, and there occur derangement of health, anemia, and wasting. On palpation it may be possible to detect the enlarged glands. In other respects, the symptoms are like those of subacute peritonitis and chronic enteritis. There is much confusion and dispute concerning the existence of a non-tuberculous tabes mesenterica, of which the principal objective symptom is the tumid abdomen.

THE LIVER.

What are the normal limits of the liver as determined by physical examination?

Under normal conditions the area of hepatic percussion-dulness (Figs. 29 and 30) is included between the sixth rib on the right, in the nipple line, the lower margin of the sixth rib on the right in the axilla, and the tenth rib posteriorly on the right, on the one hand, and the inferior border of the right

Fig. 29.



The relations of the heart, lungs, liver, stomach and spleen, as seen from the front (Weil). The deeply-shaded areas represent the portions of the heart, liver and spleen not covered by the lungs; the lightly-shaded areas represent portions covered by the lungs. *b*, *c*, *d*, boundary between lung and heart; *e f*, lower boundaries of lungs; *g h*, upper boundaries of lungs; *l*, lower limit of hepatic dulness; *m*, area of splenic dulness; *n*, greater curvature of stomach; *p*, upper limit of deep hepatic dulness.

costal arch on the other. The left lobe extends into the left hypochondrium and the dulness to which it gives rise is practically inseparable from the cardiac percussion-dulness.

Floating Liver.

To what symptoms does a floating liver give rise?

Occasionally the coronary ligament of the liver becomes lengthened and the organ acquires an abnormal freedom of movement.

Fig. 30.



Showing the relations of the lungs, liver, spleen and kidneys, as seen from behind (Weil-Luschka). The deeply-shaded areas represent portions of the liver and spleen not covered by the lungs; the lightly-shaded areas represent portions covered by the lungs. *a b*, lower border of the lungs; *c d*, boundaries of complementary pleural spaces; *e f g*, divisions between lobes of lungs; *i*, lower margin of liver.

That such an unusual tumor in the abdomen is the liver is determined by the size and conformation of the organ, by the percussion-resonance in the normal region of the liver and by the absence of the phenomena of malignant disease.

Congestion of the Liver.

What are the symptoms of congestion of the liver?

Congestion of the liver results from derangement of the circulation dependent upon disease of the heart or lungs, or upon com-

pression of the hepatic vessels; it may result from constant irritation of the hepatic cells by improper, and especially stimulating, articles of diet; it is wont to occur in those of inactive and sedentary habits. The condition is manifested by a sense of weight and dull pain in the right hypochondrium, pain at the right shoulder, headache and vertigo. The *appetite* is impaired; the *tongue* is coated; there are nausea and vomiting; the digestion is deranged; the *bowels* are constipated; the *skin* and conjunctiva may present a yellowish tinge; hemorrhoids frequently develop; there is, sometimes, mental depression or irritability of temper; the *heart* may be irritable, as manifested by palpitation. The *urine* may contain bile-pigment and an excess of urates.

Acute Hepatitis.

What are the symptoms of acute hepatitis?

Acute hepatitis is essentially a disease of tropical climates. It may also occur in association with dysentery and other infectious diseases.

It is manifested by pain in the right hypochondrium, febrile symptoms and possibly slight jaundice. The *appetite* is impaired, and there may be nausea and vomiting. Abscess is a not uncommon sequel of inflammation of the liver.

How is the malarial cachexia to be distinguished from acute hepatitis?

As a sequel of chronic malarial poisoning, the liver and the spleen become crowded with pigment and undergo enlargement. There may be slight intermittent fever; the fever of hepatitis is continuous and not periodic. In the malarial cachexia the complexion is sallow, not jaundiced, as it may be in hepatitis. In the malarial cachexia, too, plasmodia are to be found in the blood.

How are acute hepatitis and portal phlebitis to be differentiated?

Portal phlebitis may develop in the course of infectious diseases or of pyemia; it may also result by extension from adjacent disease. It is attended by pain in the right hypochondrium, by enlargement of the liver, by distention of the

veins of the abdominal wall, by ascites, by enlargement of the spleen and by diarrhea. Gastric or intestinal hemorrhage may occur. When suppuration takes place, there are recurring chills and fever, with wasting and debility. Enlargement of the spleen, distention of the abdominal veins and gastric and intestinal hemorrhage are not a part of acute hepatitis.

Acute Yellow Atrophy of the Liver.

What are the symptoms of acute yellow atrophy of the liver?

The etiology of *acute yellow atrophy of the liver* is obscure. The disease is more common in females than in males, and in young adults than in others; it sometimes appears in the course of pregnancy; at other times, in conjunction with profound emotion. After death, the liver is found to be much reduced in size and weight; in places it presents areas of reddish, purplish and yellowish discoloration, in the midst of which the hepatic cells are replaced by granules and oil-globules. By some the condition is said to be a parenchymatous inflammation of the liver.

The onset of the more grave symptoms may, for a few days or weeks, have been preceded by jaundice. This has even been associated in rare instances with enlargement of the area of hepatic percussion-flatness. Commonly, there are soon added headache, intolerance of light, delirium, stupor, convulsions and coma. The tongue is dry and coated; there are nausea and vomiting. Hemorrhages from the mucous surfaces often occur, and subcutaneous petechiæ appear. The area of hepatic percussion-dulness progressively diminishes; the abdomen becomes distended. The urine may be scarcely discolored; it contains leucin and tyrosin, and sometimes albumin; it is deficient in urea, uric acid, chlorids, phosphates and sulphates; it may contain an excess of ammonia. There may be chills and irregular fever, sometimes high; usually, however, there is but slight elevation of temperature; and in most cases the temperature is at some time subnormal. The disease rarely lasts longer than a week; it usually terminates fatally.

How are acute yellow atrophy of the liver and typhoid fever to be differentiated?

Typhoid fever is a disease lasting three or four weeks; recovery is common. The active stage of acute yellow atrophy rarely lasts longer than a week; the termination is usually fatal. There is but slight or very irregular fever in acute yellow atrophy and nearly always a tendency to subnormal temperature; the course of typhoid fever is decidedly febrile and typical. Diarrhea is common in typhoid fever; constipation, in acute yellow atrophy. The size of the liver becomes much reduced in acute atrophy; it is unchanged or increased in typhoid fever. Rose-spots are wanting in acute atrophy, in which there may be petechiæ. Jaundice is invariable in acute atrophy; rare in typhoid fever. The cerebral symptoms are more decided in the hepatic disease than in the infective fever. The presence of leucin and tyrosin and the deficiency of urea, uric acid, chlorides, phosphates and sulphates in the urine are characteristic of acute yellow atrophy; while the blood and the urine of typhoid fever exhibit distinctive reactions.

How are acute yellow atrophy of the liver and phosphorus-poisoning to be differentiated?

The symptoms occasioned by poisoning with phosphorus closely resemble those of acute yellow atrophy of the liver. Phosphorus-poisoning, however, usually sets in with vomiting and purging, to which the subsequent manifestations succeed; there is sometimes necrotic disease of the inferior maxillary bone. The alkalinity of the blood may be diminished and the number of red cells increased. The first symptom of acute yellow atrophy of the liver is jaundice. In phosphorus-poisoning the liver is enlarged prior to becoming smaller; the diminution in the size of the liver in acute yellow atrophy is progressive from the onset of active symptoms, though the organ may have been enlarged previously. Colicky abdominal pains usually attend phosphorus-poisoning, of which it may be possible to elicit a history. The anatomic lesion of acute yellow atrophy is a parenchymatous hepatitis; of phosphorus-poisoning, a fatty degeneration of the liver.

Abscess of the Liver.

What are the symptoms of abscess of the liver?

Abscess of the liver may follow traumatism or acute hepatitis; it may result from the activity of pyogenic organisms introduced into the structure of the organ in the course of inflammation of the biliary ducts, or of ulcerative processes in the distribution of the tributaries of the portal vein, or as a part of a general pyemia. When the suppuration is dependent upon traumatism, hepatitis or ulceration, but a single abscess usually develops; in cases of pyemia, there may be multiple abscesses, not only in the liver, but also elsewhere. Dysentery is a common cause of hepatic abscess. When the two are associated, the amœba coli may be found both in the stools and in the abscess or its walls. The right lobe suffers more commonly than the left. As a result of the inflammation associated with the development of an abscess, the liver becomes enlarged and tender, the enlargement tending upward and being detectable by palpation and percussion. There is pain over the liver and at the right shoulder. The tissues over the liver are often edematous. The right hypochondrium may be exquisitely tender. Repeated rigors may occur; there may be periodic exacerbations of temperature, followed by copious sweats. Slight jaundice may develop, but it is often absent. There may be nausea, vomiting and diarrhea. The number of colorless blood-corpuscles is increased. Hiccough, cough and dyspnea may result from pressure upon the diaphragm, and pleurisy, with the symptoms and physical signs denoting a rapid effusion, may result from extension of the inflammation or conveyance of the infection. When the abscess is superficial, a fluctuating tumor may be detectable in the right hypochondrium. The abscess may rupture into the pleural cavity or into the peritoneum, with a fatal termination; or the pus may make its exit through the abdominal wall, or even into the intestines. A hepatic abscess may be evacuated through the bronchial tubes, the fluid evacuated presenting the reddish-brown appearance of anchovy sauce. Death may result from septic poisoning or from exhaustion.

How are abscess of the liver and occlusion of the biliary passages to be differentiated?

If the common bile-duct is obstructed, bile accumulates in the gall-bladder and in the radicles of the hepatic ducts, the liver becomes enlarged, and jaundice, rigors, fever and sweats appear. The stools are pale.

If the cystic duct is occluded, there is no jaundice and the stools are not discolored. In both instances, the gall-bladder becomes distended with fluid and constitutes a fluctuating tumor, which may simulate an abscess. The differentiation depends upon the fact of the tumor occupying the situation of the gall-bladder, upon a knowledge of the existence of a cause of biliary obstruction, such as a calculus or a neoplasm, and upon the absence of a cause of abscess; nor is an abscess likely to occasion jaundice.

How are abscess of the liver and carcinoma of the liver to be differentiated?

Medullary carcinoma may give rise to a sort of fluctuation on palpation, but beyond this and the enlargement of the liver, the symptoms differ radically from those of abscess. In cases of carcinoma, chills, fever and sweats do not occur; but there is distinct cachexia, perhaps with nodules in other situations.

How is actinomycosis of the liver to be distinguished from abscess of the liver?

Actinomycosis is dependent upon the presence of a special fungus, which usually gains entrance through an abrasion of the surface, and may be disseminated by the blood-stream.

The irritation caused by the presence of the fungus, or perhaps by some product of its metabolism, is followed by inflammation and suppuration, so that the conditions of abscess are reproduced, and in that sense there is no discrimination to be made.

The diagnosis depends upon a knowledge of infection or of the existence of the disease at the original site of entrance, or the discovery of the ray-fungus (actinomyces) in the pus.

Interstitial Hepatitis—Cirrhosis of the Liver.

What are the symptoms of interstitial hepatitis, also called sclerosis or cirrhosis of the liver?

When irritating matters are in constant circulation in the blood-current, especially in the portal stream, the interstitial connective-tissue of the liver slowly undergoes hyperplasia, to the consequent detriment of the parenchymatous structures. The rapidity of the process depends upon the intensity of the irritation. When the increase of connective tissue has reached its height, contraction sets in and the liver-cells suffer still more. The organ, at first irregularly, perhaps nodularly, enlarged, now becomes diminished in size. This is the condition of cirrhosis. The different stages are sometimes respectively called *hypertrophic cirrhosis* and *atrophic cirrhosis.* In many cases the enlargement persists, notwithstanding extensive destruction of the secreting structures. In other cases the hepatic percussion-dulness and palpation-area are apparently diminished from the first. In the early stages of the disease the symptoms are inconspicuous; there may be manifestations of derangement of the gastro-intestinal system. As the contraction becomes more marked, however, there appear evidences of interference with the functions of the liver. Ascites develops; the superficial veins of the abdomen become enlarged and prominent; this may be especially marked in the neighborhood of the umbilicus, giving rise to the so-called *Caput Medusæ;* small networks of venules appear at various parts of the surface of the body; hemorrhoids form; hemorrhages may take place from the nose and stomach; the skin assumes a pallid, clayey hue; and ultimately jaundice appears. The spleen is often enlarged. Vertigo is not rare. Late in the disease, drowsiness and coma may develop. Excessive indulgence in alcohol is the most common cause of cirrhosis of the liver. This variety of cirrhosis has been designated *portal.*

Under other circumstances the hyperplasia involves more particularly the distribution of the biliary radicles, and the liver maintains its enlarged size. In this variety of cirrhosis jaundice appears early, while ascites occurs late if at all. The affection

is of long duration, and attended with pain in the right hypochondrium, recurring in attacks. The spleen is enlarged and hard. This variety of cirrhosis has been designated *biliary*.

Fatty Liver.

What are the symptoms of fatty liver?

The liver may undergo *fatty infiltration* or *fatty metamorphosis*. A temporary accumulation of fat in the liver occurs under normal conditions after the ingestion of food rich in hydrocarbons. When, from any cause, oxidation becomes deficient, this fatty infiltration may become permanent and excessive, the more especially if, at the same time, immoderate eating or drinking is indulged in. Thus the condition is found in the indolent, in gourmands, in the obese, in the subjects of pulmonary tuberculosis and in drunkards.

Drunkards, however, are more liable to true *fatty metamorphosis*, the liver-cells in circumscribed or extensive areas undergoing oily degeneration, a condition that may likewise occur in wasting diseases, or as an apparent result of protracted discharges, or of blood-changes, in pyemia, infectious diseases, or other morbid states accompanied by protracted high temperature. It is sometimes a part of the degenerative processes of old age. It may accompany carcinoma, cirrhosis and amyloid degeneration of the liver.

The symptoms are not early obtrusive. The area of hepatic percussion-dulness is increased, and to palpation the organ seems smooth and rounded, perhaps soft and doughy. The skin may feel greasy or velvety, and is sometimes smooth and glistening; it may be pale or flushed. Hemorrhoids are not infrequent. Jaundice is uncommon; there is no pain; ascites is rare. Diarrhea is the most constant symptom, and the least indiscretion in diet may provoke gastro-intestinal catarrh. In the more advanced stages anemia, hydremia, dyspnea and patent signs of failure of hepatic function appear.

Amyloid Disease of the Liver.

What are the symptoms of amyloid, waxy or lardaceous disease of the liver?

When the liver undergoes amyloid degeneration, other organs, such as the spleen, the kidneys, perhaps, also, the stomach and the intestines, are likewise involved, either simultaneously or consecutively. The liver is smoothly and often enormously enlarged; digestion is impaired; there may be persistent diarrhea. Pain, jaundice and ascites are uncommon. Isolated disease of the liver usually escapes detection. The process is chronic. Amyloid degeneration is a sequel of syphilis, of tuberculosis, of suppuration, of bone-disease.

How are amyloid disease and cirrhosis of the liver to be differentiated?

The liver is enlarged in both amyloid disease and cirrhosis of the liver; the enlargement is persistent in the one, replaced by contraction in the other. The amyloid liver is smooth, the cirrhotic liver usually irregular. Cirrhosis is attended with ascites and jaundice, while amyloid disease, as a rule, is not. In amyloid disease, there is a history of syphilis, of tuberculosis, of suppuration or of bone-disease; in cirrhosis, there is in most cases a history of alcoholism. With amyloid disease of the liver is usually associated amyloid disease of the kidney, occasioning the presence of albumin and tube-casts in the urine. The tube-casts will sometimes give a characteristic reaction with iodine.

Carcinoma of the Liver.

What are the symptoms of carcinoma of the liver?

The liver is a frequent seat of carcinoma, which may be primary or secondary. Primary carcinoma of the liver may be massive, nodular or cirrhotic; secondary carcinoma is usually nodular. In its incipiency carcinoma of the liver may escape detection. Soon, however, there is unaccountable loss of flesh, with the development of the characteristic cachexia. The liver is noted to be enlarged, and on palpation adventitious nodules

may be felt through the abdominal wall, in the region of the right hypochondrium. Jaundice is uncommon, unless the biliary passages are compressed. The abdomen becomes distended and fluid collects in the peritoneal cavity. The digestive derangement becomes marked. Emaciation progresses and the patient becomes reduced to the lowest degree. There is often excruciating lancinating pain. Tenderness in the right hypochondrium is a fairly constant attendant of carcinoma of the liver.

How is carcinoma of the liver to be distinguished from amyloid disease of the liver?

Emaciation and anemia attend both carcinoma and amyloid disease of the liver; but the straw-colored appearance presented by a patient with the carcinomatous cachexia is wanting in the conditions that give rise to amyloid disease. Cases of carcinoma do not present the peculiar distribution of amyloid disease in liver, spleen, kidneys and gastro-intestinal tract. The liver is often nodulated when carcinomatous; an amyloid liver is always smooth. Pain is present in carcinoma, absent in amyloid disease. A history of congenital, or of acquired syphilis, of tuberculosis, or the presence of a suppurating focus anywhere in the body would weigh in favor of amyloid disease. Amyloid disease itself is not immediately fatal. Life is not likely to be prolonged for more than a year after carcinoma of the liver has been discovered.

How are carcinoma of the omentum and carcinoma of the liver to be differentiated.

When carcinoma develops in the omentum, this structure becomes shortened and rolled up beneath its attachments. As a consequence, there is dulness on percussion, and evidences of a new-growth on palpation, not alone in the right hypochondrium, but in the epigastric and umbilical regions, and in the left hypochondrium.

Carcinoma of the omentum is more commonly than carcinoma of the liver associated with ascites. In carcinoma of the omentum the symptoms of derangement of the functions of the liver are wanting. A new-growth of the liver rises and falls with this

organ in expiration and inspiration; when the omentum is involved the mass is fixed.

How are carcinoma of the stomach and carcinoma of the liver to be differentiated?

Carcinoma of the stomach and carcinoma of the liver present many symptoms in common: a tumor in the right hypochondrium, progressive emaciation, cachexia and secondary growths; but when the stomach alone is involved there is obstinate vomiting, with absence of evidence of hepatic derangement; while if the liver only is involved ascites is common, and vomiting of coffee-ground material is lacking. A new-growth of the liver will rise and fall with this organ in respiration; a gastric tumor is more fixed. The percussion-dulness yielded by a tumor of the liver merges with the dulness of the liver; while the dulness of a tumor of the stomach may be separated from the hepatic dulness by an interval of tympanitic resonance. Hepatic and gastric carcinoma not infrequently coëxist.

Hydatid Cyst of the Liver.

To what symptoms does an hydatid cyst of the liver give rise?

A small, deeply-seated hydatid cyst of the liver, not interfering with the function of other parts, may give no sign. When the liver is the seat of an hydatid cyst in an accessible situation, the condition may be recognized by the presence of a soft, elastic, resistant, fluctuating tumor, the viscid contents of which may, on percussion and palpation, transmit a peculiar thrill or fremitus. This "purring tremor," recalling the trembling of a bowl of jelly, is present in about one-half of the cases. In addition, there may be pain in the right hypochondrium. The presence of such a new-growth, together with the absence of symptoms of profound constitutional disturbance, distinguishes an hydatid cyst from a malignant growth in the liver and its neighborhood.

Pressure-signs vary with the location of the tumor. A cyst situated near the hepatic duct or common bile-duct may cause

obstruction and fatal jaundice; pressure on the portal vein may cause ascites; jaundice and dropsy, however, are not usual. If on the upper surface of the right lobe, the cyst will push up the diaphragm, giving rise to dyspnea and cough. The heart may be displaced. If the enlargement takes a different direction, the abdominal viscera may be encroached upon. There may be multiple cysts.

Retrogressive changes may take place in an hydatid cyst, attended with a diminution in size. The contents of the cyst may be evacuated through the stomach, bowel, bronchial tubes or abdominal wall, or into the abdominal or pleural cavity.

Spontaneous evacuation may lead to recovery or it may cause inflammation and suppuration of the invaded organ and death result. Sudden death may happen from invasion of the vena cava or of the pericardium. When retrogression or evacuation is not brought about, spontaneously or therapeutically, gradual failure and death may occur from exhaustion, or from septicemia.

How are abscess of the liver and hydatid cyst of the liver to be differentiated?

An hydatid cyst of the liver is ordinarily wanting in the constitutional phenomena of hepatic abscess: rigors, fever, sweats, emaciation. Should suppuration take place an abscess virtually results. Hydatid cysts are insidious in onset, slow in development, protracted in duration, and may undergo spontaneous disappearance; when accessible they occasion extensive percussion-dulness and a peculiar thrill on palpation. An hepatic abscess may be insidious in onset, but it soon gives rise to decided symptoms; spontaneous resolution never occurs, and thrill or fremitus is wanting. The detection of echinococcus hooklets in the contents of an hydatid cyst, and of amœbæ coli in the pus from an hepatic abscess may in each case be regarded as diagnostic.

There is a growing tendency to regard exploratory puncture as inconclusive and dangerous, and to substitute exploratory incision—the latter to be followed if necessary by immediate operation.

How are an hydatid cyst of the liver and a distended gall-bladder to be differentiated?

Distention of the gall-bladder is usually dependent upon occlusion of the common bile-duct, or of the cystic duct, and is commonly associated with intense jaundice and clay-colored stools. Jaundice is exceptional in the case of an hydatid cyst, and the action of the bowels is not necessarily deranged. A distended gall-bladder occupies a definite situation; an hydatid cyst may be seated in any part of the liver. The "purring tremor" is not given by a distended gall-blader.

Perihepatitis.

What are the clinical features of chronic inflammation of the capsule of the liver, or perihepatitis?

The capsule of the liver may become thickened as a result of contiguous inflammatory processes, such as pleuritis and hepatitis, or as may be occasioned by a gastric ulcer. Perihepatitis may also be but a part of a chronic peritonitis. The liver itself is deformed, and may be diminished in size. The symptoms are ill-defined. Pain and tenderness in the region of the liver are usually present. Ascites develops. The kidneys are said usually to be diseased, so that the urine contains albumin.

How are chronic perihepatitis and cirrhosis of the liver to be differentiated?

Cirrhosis of the liver and perihepatitis may be associated. Occurring independently, the enlargement or the diminution in the size of the liver and the jaundice of cirrhosis are likely to be wanting in perihepatitis; while the presence of albumin in the urine points to the existence of perihepatitis.

Cholangitis—Cholecystitis.

What are the symptoms of inflammation of the bile-ducts and gall-bladder?

Inflammation or catarrh of the bile-ducts and gall-bladder, the most common cause of jaundice, is a result of the extension of

inflammation or catarrh or of the invasion of microbes from the duodenum or of the presence of biliary calculi.

In consequence of the swelling of the mucous membrane, the flow of bile is interfered with and jaundice results. The surface of the body and the visible mucous membranes assume a yellowish or greenish hue ; the urine is similarly discolored ; the stools become scanty and clay-colored. The saliva may be discolored. It may be possible to palpate the enlarged gall-bladder.

There is pain in the right hypochondrium ; the liver is enlarged. The action of the heart is retarded, and the respiratory frequency is lessened. There is itching of the skin. The appetite is usually impaired ; the tongue is coated ; digestion is enfeebled ; the bowels are constipated. Chills and irregular fever or transient rise of temperature may occur.

How is catarrhal jaundice to be distinguished from other forms of jaundice ?

Many conditions occasion jaundice. In some, as in catarrhal jaundice, there is an interference with the discharge of bile into the intestine ; in others, no such obstruction can be detected. In the first group of cases belong such conditions as occlusion of the common bile-duct by gall-stones, or parasites, stricture or obliteration or new-growths of the duct, compression of the common bile-duct by enlarged glands or by neoplasms or as a result of inflammation and thrombosis of the portal vein. In the second group belong yellow fever, Weil's disease, the intense forms of malarial fever, acute yellow atrophy of the liver and cirrhosis of the liver.

When the common bile-duct is occluded by a *calculus*, repeated attacks of colic are apt to occur, attended with excruciating pain, referred to the right hypochondrium ; and occasionally one or more calculi are expelled into and from the intestine, so as to be found in the stools. If the calculus is large, or if more than one is present, they may be detected by palpation, and fremitus and friction-sound may be appreciable. Colic and severe pain are wanting in catarrhal jaundice, the icterus of which is apt to be less intense and less protracted than that of obstructive disease.

In *compression* of the common bile-duct, the portal vein is likely also to be compressed, so that ascites develops. When such compression is dependent upon an *enlarged gland* this may be palpable from without.

Pylephlebitis is likely to be associated with cirrhosis or syphilis of the liver, a new-growth of the portal vein, adjacent proliferative peritonitis, perforation by gall-stones or by a suppurating hydatid cyst, and may be attended with the sudden onset of symptoms of engorgement of the portal system, with hematemesis, melena, ascites and splenic enlargement. It may be secondary to ulcerative disease of the bowel, as appendicitis or dysentery; or due to the presence of a foreign body, as a pin. In addition to febrile phenomena there may be evidences of pyemia or septicemia.

Malignant disease may involve the biliary passages or the surrounding structures. A malignant growth occasions emaciation and a peculiar cachexia manifested by a straw-colored complexion. Ascites, tumor, repeated rigors, decided fever, emaciation and cachexia are wanting in catarrhal jaundice.

Yellow fever is an epidemic, infectious disease, characterized by a train of grave symptoms, including headache, pains, fever, vomiting and prostration, that is wanting in catarrhal jaundice.

Weil's disease also is accompanied by pronounced constitutional manifestations, including fever, headache, vertigo, malaise, albuminuria, hemorrhages.

When *malarial fever* is attended with jaundice, there are commonly, in addition, chills, fever, sweats and perhaps hematuria.

The symptoms of *acute yellow atrophy of the liver* may supervene upon those of an apparently catarrhal jaundice. In such a case the area of hepatic percussion-dulness rapidly becomes diminished, cerebral symptoms appear, the urine contains lucin and tyrosin, and is deficient in urea, uric acid, chlorids, phosphates and sulphates, and death is the usual issue.

When jaundice attends *portal cirrhosis of the liver*, the discoloration is gradual in onset and slight in degree. There are, besides, enlargement or shrinkage of the liver, enlargement of the abdominal veins, and ascites. In *biliary cirrhosis* jaundice may appear early, be pronounced and be unattended with

ascites, but associated with enlargement of the spleen as well as of the liver.

Biliary Calculi.

What are the clinical manifestations of biliary calculi?

Biliary calculi are more common in women than in men, and late in life than early. They result from the precipitation and agglomeration of the less soluble elements of the bile. They may be single or multiple. They occasion symptoms when they give rise to obstruction. When large or more than one is present they may be detected by palpation, and a peculiar fremitus may be appreciable, and on auscultation a characteristic friction-sound. The presence of biliary calculi may be detected by means of fluoroscopy or skiagraphy.

The passage of a gall-stone through the common bile-duct is attended with excruciating colicky pain in the right hypochondrium, radiating at times to the shoulder, with shivering, elevation of temperature, nausea, vomiting and hiccough. The face is pale and the surface of the body is covered with a cold sweat. Jaundice soon develops—transient, if the calculus passes into the bowel; persistent, if the calculus remains impacted in the common duct. The spleen may be enlarged.

By ulceration and perforation an impacted gall-stone may find its way into the stomach, the bowel or the peritoneal cavity, or it may be evacuated externally through the walls of the abdomen.

Attacks of hepatic colic are prone to be repeated. Occlusion of the common or of the hepatic duct is followed by stagnation of bile, increased secretion of mucus, and ultimately by the formation of pus. The gall-bladder may become distended into a great sac, and the accumulation advance into the biliary radicles. Under such circumstances, jaundice persists and a type of fever develops that has been designated hepatic intermittent. Periodic elevations of temperature occur, preceded by rigors and followed by sweats. Unless relief is afforded, death ultimately results from cholemia or from acholia.

Biliary calculi and carcinoma of the biliary passages are not rarely associated.

How are hepatic intermittent fever and malarial intermittent fever to be differentiated?

The intermittent fever that occurs as a result of obstruction of the biliary passages, with or without subsequent suppuration, closely resembles malarial intermittent fever; but in the former, there is jaundice, intense and persistent; the distended gall-bladder or the obstructing calculi may be detectable by palpation; parasites are not present in the blood; and the fever does not yield to quinin.

How are hepatic colic, renal colic and intestinal colic to be differentiated?

The pain of hepatic colic is referred especially to the right hypochondrium, whence it may radiate to the right shoulder; that of renal colic is referred to one loin, not necessarily the right, whence it radiates in the course of the corresponding ureter; the pain of intestinal colic is especially umbilical, whence it may radiate over the entire abdomen. Hepatic colic is soon followed by jaundice; the urine contains biliary pigment, and is greenish or brownish in color. In cases of renal colic, the urine may contain pus-corpuscles, blood-corpuscles and crystalline matters; jaundice is wanting. Intestinal colic depends upon intestinal derangement, sometimes upon lead-poisoning; the urine remains unaltered; there is no jaundice.

How are carcinoma of the biliary passages and obstruction of the bile-duct by calculi to be differentiated?

Obstruction of the biliary passages either by gall-stones or by carcinoma occasions persistent and intense jaundice; but in case of malignant disease there is also progressive emaciation, death usually occurring within a year of the discovery of the existence of the disease. Attacks of hepatic colic attend the presence of gall-stones in the biliary passages.

The obstruction dependent upon calculi may be overcome, spontaneously or therapeutically. Unrelieved, death ultimately results from acholia or cholemia.

The diagnosis is often extremely difficult. The uncertainty is increased by the fact that calcular obstruction and malignant disease are sometimes associated. The detection of primary

carcinoma elsewhere, or of metastatic nodules, indicates the existence of malignant disease. It was at one time thought that fever was wanting in malignant disease, but this is not an invariable rule.

THE SPLEEN.

What is the normal situation of the spleen, as determinable by physical examination?

The spleen (Fig. 31) is placed obliquely beneath the ribs, from

FIG. 31.



Relations of the spleen (Weil). *M*, middle line of back; *A*, *B*, *C*, axillary lines; *Sc*, line of scapula; *a b c d*, limits of spleen; *a b c' d*, limits of rhomboidal spleen; *e f g*, limits of kidney; *l b c*, angle between lung and spleen; *d h g*, angle between spleen and kidney; *n m*, lower margin of liver.

the ninth to the twelfth, on the left side below the axilla. The organ is about four inches long by three inches wide; its axis

passes downwards and forwards. Posteriorly, the dulness of the spleen merges with that of the left kidney.

Floating Spleen.

To what symptoms does a floating spleen give rise?

The attachments of the spleen may become relaxed and prolonged, so that the mobility of the organ is abnormally increased, and its displacement is facilitated. The essential points in the recognition of a floating spleen are the presence of a solid body in an unusual situation, the absence of the spleen from its normal seat, the possible facility of replacement and the absence of pronounced symptoms.

Splenitis.

What are the symptoms of splenitis?

Inflammation of the spleen is a condition clinically obscure. It may occur in the course of infectious diseases or may be set up by emboli swept into the splenic vessels. It has in cases been attributed to traumatism. Under such circumstances the organ becomes enlarged. There are pain and tenderness in the left hypochondrium, or rather at the lower lateral aspect of the left chest, aggravated by respiration; there are also nausea, vomiting and elevation of temperature. Inflammation of the spleen may terminate in suppuration. Then chills, fever and sweats are superadded. The greatest danger is from the rupture of an abscess into the peritoneal cavity. Symptoms resembling those of exophthalmic goiter have been noted in a case of splenic abscess.

Enlargement of the Spleen.—Splenic Tumor.

Under what conditions does enlargement of the spleen take place?

Enlargement of the spleen may occur in conjunction with infectious diseases, particularly typhoid fever, malarial fever,

pyemia, septicemia; cirrhosis of the liver; venous stasis; amyloid degeneration; hemorrhagic infarction, and leukemia.

Splenomegaly (Banti's Disease).

What is splenomegaly?

This is a morbid condition characterized by enlargement of the spleen, with anemia, diminution in the size of the liver, jaundice and ascites. The etiology is undetermined. The disease occurs most commonly in young persons and adolescents. Its duration may cover years, and death finally results.

THE PANCREAS.

Acute Pancreatitis.

What are the symptoms of acute pancreatitis?

The frequency of acute pancreatitis is not known, as the condition is difficult of recognition. It may result from extension of adjacent inflammation, from the irritation of matters contained in the blood, from traumatism and from hemorrhage into the pancreas. It is characterized by acute, deeply-seated pain and tenderness in the epigastric region, or perhaps colicky pain, shooting to the back and shoulders; by anorexia, thirst, nausea, vomiting of a thin viscid liquid, sometimes of bile; by tympanites, dyspnea, restlessness, anxiety and depression; by febrile symptoms; and usually by constipation—phenomena similar to those presented by peritonitis. In some cases there is diarrhea, the stools being thin, watery, and resembling saliva. Suppuration may take place in the pancreas and an abscess form. Gangrene of the organ may be a sequel. Death is the usual termination. It may be sudden, with the phenomena of collapse. Fat-necrosis is a common accompaniment of all forms of pancreatic disease.

Carcinoma of the Pancreas.

What are the symptoms of carcinoma of the pancreas?

New-growths of the pancreas are not common, but of those that

occur carcinoma is the commonest. The disease may be primary or secondary. The head of the organ is involved in most instances. Symptoms may long be latent. Among the most constant are epigastric pain, jaundice and distention of the gall-bladder from pressure on the biliary passages, the presence of a tumor, wasting and cachexia. Pressure on the pylorus may give rise to dilatation of the stomach, with nausea and vomiting. There may be fatty diarrhea.

Pancreatic Cysts.

What are the clinical features of cysts of the pancreas?

Cysts of the pancreas may arise from retention and accumulation of fluid in an occluded or obstructed pancreatic duct, from the proliferation of pancreatic tissue, or as a result of traumatism or inflammatory processes, leading to hemorrhage and dropsy of the lesser peritoneal cavity. Among the most common symptoms are colicky pains in the abdomen, with nausea and vomiting and the appearance of a comparatively fixed tumor in the upper half of the abdomen. The bowels may be loose and the stools contain undigested fat. Jaundice and ascites may result from pressure. The urine may contain albumin or sugar, and the secretion of saliva may be increased. Loss of weight and failure in strength usually take place. Fluid obtained on puncture may contain blood-corpuscles or hemoglobin, granular detritus, fat globules and cholesterin, and it may exhibit the fermentative activity of pancreatic juice.

Pancreatic Hemorrhage.

What are the symptoms of hemorrhage into the pancreas?

Hemorrhage into the pancreas is characterized by the occurrence of severe pain of sudden onset in the upper part of the abdomen, with nausea, vomiting, a sense of anxiety, restlessness, depression, coldness of the surface, sweating, a weak, rapid pulse, tympanites, constipation.

THE GENITO-URINARY APPARATUS.

Examination of the urine is important not only when disease in the genito-urinary tract is suspected, but also in many other varied conditions.

The *constitution of the urine* may be quantitatively or qualitatively altered. The proportions of normal ingredients may be increased or diminished. Substances not normally found in the urine may under certain conditions be present.

The *quantity of urine* excreted by a healthy adult in twenty-four hours varies between forty and fifty ounces. The quantity is physiologically *increased* after the ingestion of large quantities of fluid and when the cutaneous, pulmonary and intestinal transpiration is diminished. The quantity is physiologically *diminished* under the reverse conditions.

The quantity of urine excreted is altered in many conditions of disease. It is increased in diabetes insipidus and diabetes mellitus, in chronic nephritis and, temporarily, in emotional states, including hysteria. The quantity of urine is diminished in acute nephritis, in the last stages of chronic nephritis, in lithemia, in renal insufficiency, in conditions characterized by feebleness of the heart, in Asiatic cholera, in yellow fever, in acute intestinal obstruction and in febrile conditions generally.

Urine is normally clear, transparent, of an amber color, without obtrusive odor, of acid reaction and a specific gravity varying from 1010 to 1030. The color is paler or darker, and the specific gravity lower or higher, as the quantity is increased or diminished.

The *transparency of the urine* is interfered with by the presence of matters rendered insoluble by an altered reaction of the urine (such as phosphates), or by a change of temperature (such as urates); by the presence of matters foreign to normal urine (such as pus, chyle or blood); and sometimes as a result of the action of bacteria.

The *color of urine* is intensified when the quantity is diminished, as in fevers, or when the action of the skin is increased. The color is altered by the ingestion of some medicaments and foods and by the presence of other abnormal ingredients. Thus,

rhubarb imparts a gamboge-yellow color to acid urine, a violet-red to alkaline urine; santonin imparts a golden-yellow to acid urine, an orange-yellow to alkaline urine; senna imparts a brownish, logwood a reddish, carbolic and gallic acids each a blackish tinge. Urates may color the urine orange-red. The presence of blood gives rise to a smoky, reddish or chocolate color; the presence of biliary pigment to a brownish or greenish hue; the presence of chyle to a milky appearance. In cases of melanotic tumors the urine may be dark or black; the color is likewise dark in some cases of pernicious anemia. The urine is pale when it is excreted in excess, as in hysteria, in diabetes and in chronic nephritis.

The *odor of normal urine* is typical, but not obtrusive. When fermentation takes place the urine becomes ammoniacal. The ingestion of turpentine imparts to the urine an odor of violets. Other aromatic oils impart odors abnormal to urine. The urine of diabetes mellitus has a sweetish odor. If it contain acetone, the odor resembles that of chloroform.

The total twenty-four hours' urine of a healthy person is of acid *reaction*. The urine may be alkaline from fixed alkali a few hours after a meal, especially if large quantities of alkalies have been taken; in cases in which the stomach is washed out; and in cases of obstinate vomiting. The urine is alkaline from volatile alkali when ammoniacal fermentation has taken place.

The *average specific gravity* of normal urine is 1018 or 1020. Except in diabetes mellitus, the specific gravity bears in general an inverse relation to the quantity of urine. It is heightened after the ingestion of large quantities of nitrogenous food, in diabetes mellitus and in acute nephritis and other febrile conditions. It is lowered in diabetes insipidus, in chronic nephritis, in hysteria, when the activity of the skin is lowered and after the ingestion of large quantities of fluid.

The normal average quantity of *urea* excreted in twenty-four hours is about five hundred grains. The excretion of urea is increased on a generous nitrogenous diet and in fevers in which the urine is not suppressed. The quantity of urea in the urine is diminished on a vegetable diet; when oxidation is interfered with, as in pulmonary or cardiac disease; in cases of

grave hepatic disease, renal insufficiency or suppression of urine; and when the urea is retained in the circulation.

If urine or any fluid containing urea be concentrated to a syrupy consistence, by evaporation in a water-bath, and nitric

FIG. 32.



Uric acid and urates. (Funke.)

acid be added, a crystalline precipitate of the rhombic plates of urea nitrate will be thrown down.

The quantity of *uric acid* excreted in the urine in twenty-four hours varies from eight to sixteen grains. It is increased when the diet is nitrogenous; when oxidation is defective, as in pulmonary or cardiac disease; after an attack of gout or lithemia; and in febrile processes. It is diminished on a diet of carbohydrates; in chronic diseases after hemorrhage; during an attack of gout or lithemia.

If ten parts of hydrochloric acid be added to one hundred parts of urine and the mixture be permitted to stand for forty-eight hours, a sediment of fine, red crystals of uric acid will form.

The quantity of *hippuric acid* excreted in the urine in twenty-four hours is between eight and thirty grains. It is increased on a vegetable, and diminished on an animal diet. It is only precipitated from acid urine, when it appears as rhombic crystals.

The quantity of *sodium chloride* excreted in the urine in

twenty-four hours is about half an ounce. It depends upon the quantity ingested. The excretion of chlorides is diminished in febrile disorders, and especially when exudation or transudation occurs.

The presence of chlorides in the urine may be determined by first acidulating the urine with nitric acid, and then adding a solution of silver nitrate. If the precipitate that forms is dense

FIG. 33.



Calcium phosphate. (Laache.)

and curdy, the quantity of chlorides is normal; if the precipitate is milky, the chlorides are diminished.

Between thirty and eighty grains of *phosphoric acid* are eliminated in the urine in twenty-four hours, two-thirds as alkaline (acid sodium and potassium) phosphates, and one-third as earthy (calcium and magnesium) phosphates. The quantity of phosphates excreted is diminished at the beginning of febrile processes, and increases with defervescence and convalescence.

The earthy phosphates are precipitated when the reaction of the urine is alkaline. The alkaline phosphates are precipitated by the addition of a solution containing one part each of magnesium sulphate, ammonium chloride and liquor ammoniæ, and eight parts of distilled water.

About thirty grains of *sulphuric acid* are eliminated as sulphates in the urine in twenty-four hours. The quantity varies with the character of the food and the degree of activity.

The presence of sulphates in the urine is determined by the addition of a solution of barium chloride, which gives a cloudy precipitate, insoluble in water or acids.

Biliary coloring matter appears in the urine when jaundice exists. The urine is yellowish, greenish or olive-brown in

FIG. 34.



Triple phosphates and ammonium urate. (Laache.)

color, and stains the linen. When the jaundice is not due to biliary obstruction it is said that the biliary acids are not found in the urine.

The presence of biliary coloring matter in the urine may be detected by overlaying nitric acid in a test-tube with urine, or by permitting the fusion of a drop each of nitric acid and of urine on a white plate. A play of color from green, blue, violet, red to yellow, results.

To detect the presence of the biliary acids in the urine a small quantity of cane-sugar is added to the urine, which is introduced into a test-tube, so as to overlay some sulphuric acid. A purple color is formed at the line of contact. If a bit of filter-paper is

dipped in the saccharated urine and touched with a drop of sulphuric acid a purple color results.

What are leucin and tyrosin?

Leucin and *tyrosin* are products of the destructive metamorphosis of proteids, and are among the waste-products of

Fig. 35.



Leucin and tyrosin. (Laache.)

pancreatic digestion. They do not occur in normal urine, but are found in the urine in cases of phosphorus-poisoning and of acute yellow atrophy of the liver, for which they may have diagnostic significance. They have also been exceptionally noted in variola, typhus, pernicious anemia, and in a case of obstructive jaundice caused by hydatid cyst of the gall-duct.

To obtain leucin and tyrosin in the sediment, the urine should be evaporated in a water-bath to syrupy consistence, or a drop may be boiled down on an object-glass. Leucin appears in the form of faintly shining spheres of variable size, the larger ones sometimes exhibiting radiation or concentric rings. Tyrosin crystallizes in very fine needles, commonly aggregated into sheaves or bundles.

Indicanuria.

What is indicanuria?

The presence in the urine of potassium indoxyl-sulphate is known as indicanuria. Indican results from the decomposition by bacteria of albumin in the intestines, and it occurs in normal urine in only small amounts. The quantity is increased in wasting diseases and when excessive albuminous disintegration is taking place. Its presence is disclosed by the bluish-black discoloration that takes place when 2 or 3 drops of solution of chlorinated soda are added to equal parts of urine and strong hydrochloric acid. On shaking with chloroform, indigo is dissolved out and settles at the bottom.

Oxaluria.

What is oxaluria?

The presence of oxalates in the urine is sometimes associated with a complex of symptoms, to which the designation oxaluria has been applied. Individuals so affected complain of languor, of dull pains in the loins, are irritable or dejected, have boils or

FIG. 36.



Calcium oxalate. (Laache.)

carbuncles; and their nutrition is impaired. The *urine*, the

specific gravity of which is increased, contains an excess of urea and of oxalates, and occasionally a trace of albumin. The condition has been observed in persons in whom the nervous system has been severely put to task. The oxalates in the urine may be increased by certain foods, such as tomatoes and rhubarb. By accretion they may form calculi.

Pyuria.

What is pyuria?

The presence of pus in the urine constitutes *pyuria*. Pus appears in the urine when there is suppuration in any portion of the genito-urinary tract, as in urethritis, cystitis, ureteritis or pyelitis. Rupture of an abscess into the genito-urinary passages gives rise to the sudden discharge of a large quantity of pus. When, in women, leucorrhea exists, some of the pus may find its way into the receptacle for urine. The addition of caustic alkalies converts pus into a gelatinous viscid mass. Hydrogen dioxide gives a characteristic foaminess, from liberation of oxygen. The microscope is the most certain test. Urine containing pus responds to chemic tests for albumin.

When *urethritis* exists, micturition is burning. The first urine passed is turbid; that which follows may be clear. The reaction of the urine remains unchanged. In the vast majority of cases, urethritis is gonorrheal.

When there is *cystitis*, there may be vesical tenesmus, with frequent passage of small quantities of urine that is usually of alkaline reaction. Cystitis may be secondary to urethritis; it may be a result of obstruction, as from stricture of the urethra or an enlarged prostate; it may be associated with a calculus in the bladder; it may have been caused by the introduction of septic matters into the bladder in the course of some surgical procedure, such as the employment of an unclean catheter, sound or bougie; it may be tuberculous; it is often seen in association with diseases of the brain and spinal cord. The urine is not only turbid, but it is usually also alkaline in reaction and offensive in odor, from ammoniacal fermentation.

Ureteritis and *pyelitis* are usually associated. Both are com-

monly secondary to cystitis, though they may result from the presence of calculi. In uncomplicated cases the urine is acid in reaction.

When an *abscess* ruptures into the genito-urinary tract, a certain sense of relief of tension is perceived, followed by the evacuation of a large quantity of pus in the urine, which had previously presented no abnormal characters.

Albuminuria.

What is albuminuria?

Albumin is found in the urine under many varied conditions; thus, urine containing pus or blood will respond to the tests for albumin. Under other circumstances, however, the albumin is derived directly from the circulating blood, as a result of defective action of the epithelium of the secretory apparatus of the kidney, or of changes in the blood or in vascular tension. Such a condition is present in the various morbid states of the kidney, in the course of fevers and toxic and infectious processes and in association with violent convulsions, obstructive disease of the heart, and interference with the respiratory function. The ordinary form of albumin found in the urine is serum-albumin; less commonly paraglobulin is present.

How is the presence of albumin in the urine to be determined?

The best test for the detection of albumin in the urine is that by heat. The urine should be clear and of acid reaction. If not clear, it should be filtered. If not acid, a drop or two of acetic acid—sufficient to impart an acid reaction, should be added. The upper half of the urine, in a test-tube, is heated. A resulting cloudiness is due to the presence of albumin or of phosphates. If the urine have been acid, the cloudiness is not likely to be due to phosphates. The addition of a few drops of acetic acid removes any doubt; if the cloudiness is dependent upon the presence of phosphates, it at once clears; if due to the presence of albumin, it persists. To the filtered urine about one-sixth its quantity of saturated solution of sodium chlorid and 5 or 10 drops of dilute acetic acid (50%) may be added before heat is applied.

A simpler test for the presence of albumin consists in overlaying a quantity of cold nitric acid in a test-tube with the urine. A white ring at the line of contact of the two fluids is indicative of the presence of albumin. Its appearance may be delayed.

FIG. 37.



Red blood-corpuscles and a blood-cast of a uriniferous tubule. (Eichhorst.)

FIG. 38.

Epithelial cast of a uriniferous tubule. (V. Jaksch.)

FIG. 39.



Granular casts of the uriniferous tubules. (V. Jaksch.)

FIG. 40.



Hyaline casts of the uriniferous tubules. (Vierordt.)

A faint ring of urates that forms above the line of contact is dissipated by the application of heat. A mixture of one part of strong nitric acid and five of saturated solution of magnesium sulphate may be employed in the same way.

Another test consists in adding to half a test-tubeful of urine a dram or more of solution of potassium ferrocyanid (1 : 20); and after thorough admixture, about 10 drops of dilute acetic acid (50%). The occurrence of turbidity indicates the presence of albumin.

A saturated solution of picric acid used by the contact-method in the same way as nitric acid, the urine, however, being over-

FIG. 41.



a

Waxy casts of the uriniferous tubules. *a*, a waxy cast containing crystals of calcium oxalate. (V. Jaksch.)

laid by the acid, constitutes a most delicate test for the detection of albumin in the urine, but it is equivocal, as it reacts with peptones, urates and with the alkaloids. Heat, it is true, dissipates the ring formed with these, but it may also cause a diffu-

sion of the turbidity produced by albumin, so that the latter may escape detection.

What is the significance of the presence of tube-casts in the urine?

The presence in the urine of casts of the uriniferous tubules is indicative of nephritis. The presence of albumin is usually associated with that of tube-casts. Albumin and tube-casts may, however, be absent, or at least escape detection, at times when nephritis exists; and they may occasionally be present unassociated with inflammation of the kidney. Centrifugation should be practised when tube-casts are not otherwise readily found.

The character of the casts varies according to the form of nephritis. *Blood-casts* and *epithelial casts* are indicative of an acute process; *granular casts* of a chronic process; *hyaline casts* may be present in many conditions. *Fatty casts* bespeak the late stage of a chronic parenchymatous nephritis—fatty kidney; *waxy casts* may be found in acute and chronic nephritis and in amyloid disease of the kidney.

Chyluria.

What is chyluria?

The presence in the urine of chyle or its molecular base, fat, constitutes *chyluria*. The urine presents a whitish, milky appearance and displays a tendency to spontaneous coagulation.

FIG. 42.



Filaria sanguinis hominis. (V. Jaksch.)

Albumin is present; in cases due to filaria, blood may be found. Under the microscope lymph-corpuscles and a finely granular

basis are recognized. Chyle finds its way into the urine as a result of abnormal communication between the lymphatic system and the genito-urinary tract, in consequence of rupture following obstruction in the lymph-channels, most commonly by *filaria sanguinis hominis*. The scrotum and the lower extremities are sometimes infiltrated, and may be enlarged so as to create a resemblance to elephantiasis. *Filariasis* is a chronic affection, as a rule irregularly intermittent in its manifestations. When filariæ are present, ova or embryos should be discovered in the lymph or in the blood.

Filaria nocturna appears during the period of sleep or rest, therefore usually at night; *filaria diurna*, during the period of activity, therefore usually by day; while *filaria pustano* may be found at any time.

Lipuria.

What is lipuria?

The presence of fat in the urine constitutes *lipuria*. It may occur in health after excessive ingestion of fatty food. It has been observed in connection with fatty degeneration of the kidney, pyonephrosis, diabetes mellitus, phosphorus-poisoning and pregnancy. Urine containing chyle responds to the tests for fat. The condition may be recognized by the addition to the urine of a small quantity of potassium hydrate and then shaking with ether. The ether is permitted to evaporate and the fat collects as globules.

Hematuria.

What is hematuria?

The presence of blood in the urine constitutes *hematuria*. The *urine* is turbid and darker in color than normal; the specific gravity is increased; and albumin is present in considerable quantity. With the microscope there is no difficulty in distinguishing the red corpuscles.

Blood may be present in the urine as a result of hemorrhage from traumatism, stone, neoplasm, tuberculosis, inflammation or ulceration in the genito-urinary tract; as a symptom of certain nervous diseases; as a manifestation of a hemorrhagic dia-

thesis, as in leukemia, scurvy, purpura, hemophilia, exophthalmic goiter, and the like; as a manifestation of influenza or malaria, or as the result of drug-action. Like chyluria, hematuria may be due to *Filaria sanguinis hominis*. *Distoma hematobium* and *strongylus gigas* are additional parasites that may cause hematuria.

It may sometimes be important to determine whether the blood present in urine comes from the bladder or from the kidneys. A comparatively small quantity of blood; its uniform admixture with the urine; the presence of blood-casts of the tubules, or of the ureter; small size, discoloration, or so-called "ringing" of the blood-cells, point to renal hemorrhage. The passage of pure blood or of clots unmixed with urine, or with but slight admixture, or only at the beginning or close of urination, points to hemorrhage from the bladder or urethra. Exploration or cystoscopic examination may detect the source of hemorrhage and perhaps its cause, as a calculus or a neoplasm.

Hemoglobinuria.

What is hemoglobinuria?

When the coloring-matter of the blood appears in the urine the condition is designated *hemoglobinuria*. The coloring-matter of the blood may appear free in the urine as a result of conditions attended with disorganization of the blood, such as scurvy, purpura, pyemia, typhus, insolation, extensive burns, and poisoning by potassium chlorate, hydrogen arsenide, phosphorus, carbolic acid and chloral. Spontaneous hemoglobinuria has been observed in the new-born infant.

The *urine* is dark-red or chocolate-brown in color, of high specific gravity, and contains considerable albumin. Microscopically, few or no blood-cells are found. A drop of such urine treated on a slide with sodium chloride, acetic acid, and heat yields microscopically the flat, rhombic, prismatic tables known as Teichmann's crystals. The addition of a few drops of fresh tincture of guaiac to a small quantity of the same urine gives rise to a whitish or greenish coloration; shaken well, the addition of a few drops of a solution of hydrogen dioxide or of

old oil of turpentine causes a change of color to blue. Spectroscopic examination reveals characteristic striæ.

Paroxysmal Hemoglobinuria.

What is paroxysmal hemoglobinuria?

Hemoglobinuria may be *intermittent* and *paroxysmal*. In some cases no cause for the condition can be elicited; in most, however, the exciting factor has been exposure to unusual cold. Attacks may be induced by mental or physical exhaustion. The disease is most common in young adult males. The onset of a paroxysm is indicated by a sense of languor and fatigue, to which is soon added a feeling of chilliness. The fingers and toes and tips of the ears become numb, cold and cyanotic. The paroxysm ends by the restoration of a comfortable sense of warmth. During the attack, or a short time subsequently, the symptom that gives the disease its name appears. The urine, however, soon assumes a normal condition. Paroxysmal hemoglobinuria may be part of the syndrome of Raynaud's disease, and in a mild form may be associated with other varieties of vasomotor ataxia.

How are paroxysmal hemoglobinuria and malarial hematuria to be differentiated?

Paroxysmal hemoglobinuria is not known to be related to malarial infection; other possible manifestations of malaria are wanting; the periodicity of the attack is not rhythmical; malarial hematuria is a manifestation of profound malarial intoxication, associated with other distinct characteristics of malaria. In hematuria, with the microscope, many red blood-corpuscles are to be seen in the urine; the absence of blood-corpuscles in the urine is characteristic of hemoglobinuria. The one condition yields to treatment with quinine; the other does not. Examination of the blood in a case of paroxysmal hemoglobinuria discloses nothing characteristic; an examination of the blood in cases of malarial hematuria reveals the presence of characteristic parasites.

Endemic Hematuria.

What is endemic hematuria?

In tropical climates the presence of a nematode worm, the *distoma hæmatobium* or *Bilharzia hæmatobia*, in the veins about the bladder, occasions the appearance of blood in the urine. The ova of the parasite find their way into the genito-urinary passages and there give rise to irritation and inflammation and extravasation of blood. The disease is most common in youth; it shows a tendency to subside after the age of puberty. The condition predisposes to the development of urinary calculi. The *diagnosis* depends upon the endemic character of the affection and the detection of ova, which are often concealed in shreds of tissue contained in the urine.

Diabetes Insipidus.

What are the clinical features of diabetes insipidus or polyuria?

Diabetes insipidus or *polyuria* is a morbid condition characterized by the elimination of an excessive quantity of feebly acid urine, of low specific gravity, containing neither albumin nor sugar. The disease is most common in young persons, and males suffer oftener than females. The progress is usually gradual, at times rapid. There is increased thirst; an abnormal quantity of fluids may be ingested, of which, however, the quantity of urine excreted is usually in excess. The *skin* and visible mucous membranes may be rough and dry. In some cases the *appetite* is excessive. The general *nutrition* may be maintained, but is rather more likely to suffer deterioration, as manifested by wasting and debility. Diabetes insipidus is not directly fatal. Its *etiology* and *pathology* are obscure. In many cases, there is a history of injury to the head, or disease of the brain is found post mortem. It may depend upon disease of the pancreas. Several cases have been found in a single family; in other instances, some members of a family present diabetes insipidus, others diabetes mellitus.

How are diabetes insipidus and chronic interstitial nephritis to be differentiated?

In cases of chronic interstitial nephritis the quantity of urine eliminated may be abnormally large, but it does not attain the proportions encountered in diabetes insipidus; in addition, careful investigation will detect the presence of albumin and casts in the urine, as well as the existence of other symptoms indicating a degenerative lesion of the kidney: heightened arterial tension, hypertrophy of the heart, breathlessness, edema, headache, diarrhea.

How are hysterical polyuria and diabetes insipidus to be differentiated?

The excessive discharge of urine that is sometimes observed in hysterical patients is paroxysmal rather than persistent; and does not constitute the essential feature of the disease, but is rather an incidental symptom of a malady of protean feature.

Glycosuria.

What is glycosuria?

By *glycosuria* is meant the presence of *grape-sugar* or *glucose* in the urine. The condition may be *transient*, *intermittent* or *persistent*. Thus, it may follow the ingestion of excessive amounts of sugar or starch or of chloral or other drugs, inhalation of chloroform, or traumatism, or attend infectious diseases and nutritional diseases, *e. g.*, chronic malaria, exophthalmic goiter, akromegaly, myxedema, acute and chronic suppurative appendicitis. It is the distinguishing feature of diabetes mellitus.

Diabetes Mellitus.

What are the clinical features of diabetes mellitus?

Diabetes mellitus is a morbid condition attended by the discharge of large quantities of urine characterized by the persistent presence of a varying quantity of glucose. It is a metabolic disorder of adult life, though at times observed in the young. Males suffer rather more commonly than fe-

males. The disease occurs sometimes in connection with cerebral disease, sometimes with disease of the pancreas or of the liver, sometimes in women at the menopause and sometimes as a sequel of infectious diseases or of exhausting labors; it sometimes attends or follows pregnancy; but it is commonly unassociated with any evident visceral disease. In many cases there is a racial or family predisposition or a gouty or rheumatic diathesis. In not a few instances the advent of the disease has been preceded by grief, anxiety or profound emotion. The course of the affection is ordinarily slowly progressive; exceptionally it is acute and rapidly fatal.

Among the earliest *symptoms* is a gradual failure of health, with a sense of fatigue disproportionately great. Often there is mental alteration, especially irritability of temper. It may be observed that the patient is passing an excessive quantity of *urine*, examination of which shows it to be pale in color, of a sweetish, fragrant odor, usually of high specific gravity, and chemically containing glucose. The proportion of sugar varies, being increased by indulgence in, and diminished by abstinence from, sugars and starches. At the same time, there is increased thirst, together with the ingestion of large quantities of fluids. The urine dissolves gentian-violet. The *appetite* is also wont to be excessive, and large quantities of food are taken. Constipation is the rule, though exhausting diarrhea may occur. The *skin*, mouth and throat are dry. A sweetish taste is perceived. The *breath* may exhale a fragrant, a balsamic, an ethereal odor, or an odor of chloroform. The *teeth* become carious, the gums spongy and loose. Boils and carbuncles are common. Persistent furunculosis, without obvious cause, should always awaken a suspicion of diabetes. In women, pruritus vulvæ, and, in men, excoriations about the meatus urinarius have been observed. The blood, dried and fixed on glass slides, does not stain with 1% aqueous solution of Congo-red. It changes the color of a weak alkaline solution of methylene-blue to a yellowish-green or yellow. Symptoms of peripheral neuritis, with abolition of the knee-jerks and ataxia may appear. Bilateral sciatic neuritis sometimes develops. Failure of sexual vigor is sometimes an early symptom. Emaciation and debility may

ultimately become extreme. Exceptionally an appearance of perfect health is preserved. Some have distinguished two groups of cases, those with a tendency to emaciation, and those with a tendency to obesity.

Sometimes, there is sense of chilliness and the temperature is subnormal. Gangrene from vascular occlusion may result and bring about a fatal issue. After death the bladder is often found hypertrophied. Cataract not rarely develops in the course of diabetes. Retinitis also may occur and retinal hemorrhage take place. Nephritis is sometimes superadded. *Death* commonly results from an intercurrent affection, such as pulmonary tuberculosis or pneumonia. Sometimes coma appears, with or without delirium and convulsions. Coma is in many cases preceded for a few days by the appearance in the urine of acetone and diacetic acid, concurrently, it may be, with diminution in the amount of urine and in the percentage of sugar. Obstinate constipation also may be a forerunner.

What is Moore's test for the presence of glucose in the urine?

The simplest test for the presence of sugar in the urine is Moore's test. To a moderate quantity of urine in a test-tube half as much liquor potassæ is added. The mixture is heated. If sugar be present in greater proportion than 0.3 per cent., a sherry-wine color results. If a drop or two of nitric acid be added, the brownish color disappears and the odor of molasses becomes perceptible. Other substances than sugar respond similarly in color; so that the test cannot be infallibly relied upon.

What is Boettger's test?

To a small quantity of urine in a test-tube is added about an equal quantity of liquor potassæ, and then a small pinch of bismuth subnitrate. The mixture is heated; if sugar be present a black precipitate of metallic bismuth is thrown down. Sulphur compounds precipitate black sulphides, so that albumin, if present, should first be removed by precipitation and filtration. If the reaction fails to take place, the presence of sugar in significant quantity can safely be excluded.

What is Trommer's test?

To a moderate quantity of urine in a test-tube a smaller quantity of liquor potassæ, and then a few drops of a ten per cent. solution of cupric sulphate are added. If heat be applied, an orange-red precipitate of cuprous oxide is thrown down if sugar be present.

What is Fehling's test?

Two solutions are used. One consists of an ounce of cupric sulphate and sufficient water to make fifteen ounces; the other of five ounces of crystallized Rochelle salts, three ounces of a solution of sodium hydrate having a specific gravity of 1.34, and sufficient water to make fifteen ounces. When the test is to be made, equal quantities of the two test-solutions are heated in a test-tube. The urine is then slowly added. Should no orange-red precipitate result, the presence of sugar can be excluded.

Fehling's test can be applied for quantitative analysis by employing solutions of which a known volume will be reduced by a definite quantity of glucose 10 cu. cm. of Fehling's solution will be reduced by 5 mgm. of glucose.

What is the fermentation-test?

If a bit of baker's yeast be added to urine containing sugar, fermentation takes place, carbon dioxide and alcohol being generated. In making a quantitative examination the specific gravity is taken and recorded; eight ounces of urine are introduced into a pint bottle, a piece of compressed yeast about the size of a walnut added, and the bottle closed by a perforated cork, and placed in a warm place (from 86° to 90° F.) for four or five hours or longer. The urine is then permitted to cool, and its specific gravity is again taken. Each degree of loss of specific gravity represents about one grain of sugar to the ounce, or the total number of degrees lost multiplied by 0.23 will give the percentage of sugar. Einhorn's fermentation-saccharimeter is so graduated as to indicate the approximate percentage of sugar by the volume of carbon dioxid given off from a measured quantity of urine.

What are the tests for acetone?

1. To 4 or 5 cu.cm. of urine are added a few drops of a fresh

solution of sodium nitroprussid, and then a strong solution of sodium hydroxid. If acetone be present a red color appears in from five to ten minutes, giving place to a purple or violet on addition of acetic acid. 2. About 4 cu.cm. (1 dram) of solution of potassic hydrate, containing 1 gram (15 grains) of potassium iodid, are placed in a test-tube and boiled. An equal volume of urine is then cautiously poured in, so as to float on the surface of the alkaline liquid. At the point of contact a ring of phosphates will form, which, if acetone be present, will in a few minutes be colored yellow and studded with crystals of iodoform. 3. Tincture of iodin or compound solution of iodin is added to the urine or a distillate thereof and then sufficient ammonia to occasion a deep-black precipitate. This disappears gradually if acetone be present and is replaced by a yellowish precipitate of iodoform, which may be recognized by its odor and by the appearance microscopically of six-sided tables or stars.

What is a convenient test for diacetic acid?

Solution of ferric chlorid is slowly added to the fresh urine; the phosphates thus precipitated are removed by filtration, and more ferric chlorid is added. If aceto-acetic acid is present, the urine becomes wine-red. The test must then be repeated with urine previously boiled. If the boiled urine gives no reaction with the iron-solution, or only a slight coloration, and if acetone is likewise found to be present, it may be concluded that *diaceturia*, a forerunner of diabetic coma, exists.

How are diabetes insipidus and diabetes mellitus to be differentiated?

The diagnostic distinction between diabetes mellitus and diabetes insipidus is the presence or absence of sugar in the urine, of which the specific gravity is lower in the latter than in the former. The occurrence of boils, of itching of the skin, of cataract and of gangrene, the presence of a sweetish taste and of a fragrant breath, encountered in saccharine diabetes, are wanting in insipid diabetes. The prognosis in the case of diabetes insipidus is more favorable than in that of diabetes mellitus.

Cystitis.

What are the symptoms of cystitis?

The mucous membrane of the bladder may undergo inflammation in the course of constitutional disorders, or as the result of local irritation or infection. Cystitis gives rise to pain in the hypogastrium, a frequent desire to urinate, followed by the evacuation of small quantities of urine, attended with burning pain. The last drops of urine passed are opaque and viscid. The quantity of urine voided does not in the aggregate exceed the normal. When the cystitis is acute the reaction of the secretion is acid; in chronic cystitis the urine is alkaline; the urine is turbid and precipitates a dense, copious sediment; it is albuminous in a degree proportionate to the quantity of pus contained. Examined microscopically the sediment is found to be made up of pus-corpuscles, pavement epithelial cells and mucus. In the earliest stages of acute cystitis the urine may contain blood; in chronic cystitis, crystals of the triple ammonio-magnesium phosphate may be found. The walls of the bladder become thickened, its cavity reduced in size, its lining membrane ribbed. The dangerous sequelæ of cystitis are ureteritis, pyelitis, hydronephrosis and suppurative nephritis, dependent upon extension of the inflammatory process.

How are cystitis and nephritis to be differentiated?

Pain in the hypogastrium is a feature when there is acute cystitis. If pain attend acute nephritis, it is referred to the loins. Urine is passed less frequently and in smaller quantities when acute nephritis exists than is the case in acute cystitis. The urine of cystitis is the more distinctly purulent; that of nephritis, the more distinctly albuminous. In cases of nephritis, the urine contains casts of the uriniferous tubules. The urine of simple cystitis never contains tube-casts. The alkalinity of the urine, the presence of pus, and the absence of tube-casts, in addition to the absence of other phenomena, such as edema, dyspnea, anemia, cardiac hypertrophy, distinguish chronic cystitis from chronic nephritis.

Vesical Calculus.

To what symptoms does a vesical calculus give rise?

The presence of a calculus in the bladder gives rise to irritation and inflammation. In addition to the signs of cystitis—frequent, burning discharge of small quantities of urine of alkaline or acid reaction—there occur attacks of vesical tenesmus, with complete obstruction to the discharge of urine, with pain referred to the glans penis, followed by the presence of blood in the urine. Careful examination, by means of a sound or the cystoscope, may furnish conclusive evidence. A vesical calculus may not only excite cystitis, but it may also give rise to inflammation and dilatation of the ureters and pelves of the kidneys and to hydronephrosis.

Neoplasms of the Bladder.

To what symptoms do neoplasms of the bladder give rise?

When the lining of the bladder is the seat of new-growths, hemorrhage is of frequent occurrence, together with paroxysms of vesical tenesmus. The *urine* usually contains cellular evidence of the presence of a neoplasm, while cystoscopic examination may afford ocular demonstration of the fact.

Pyelitis.

What are the symptoms of pyelitis?

Inflammation of the mucous membrane of the pelvis of the kidney may arise in the course of acute infectious diseases, may result by extension from the urethra, through the bladder and ureters, or from the presence of a calculus or a new-growth. Pyelitis may be tuberculous. There is pain in the loin. The *urine*, the reaction of which remains acid, precipitates a copious sediment principally constituted of pus, with some epithelial cells. From time to time the urine may contain blood. If suppuration be pronounced there may be chills and fever, with anemia and emaciation.

In cases of tuberculosis, tubercle-bacilli may be found in the urine, and there may be evidences of tuberculous disease elsewhere. It is said that bacilli morphologically indistinguishable from tubercle-bacilli and presenting the same color-reactions are sometimes found in the urine of healthy persons. The numbers are smaller, and the organisms are less constantly present than in cases of tuberculosis, and the inoculation-test proves them to be innocuous.

How are pyelitis and cystitis to be differentiated?

The pain of pyelitis is referred to the lumbar region; that of cystitis to the hypogastric region. Small quantities of urine are frequently evacuated in case of cystitis; neither quantity nor frequency is altered in uncomplicated pyelitis. When there is cystitis, the urine is most likely to be alkaline in reaction; when there is pyelitis, it is usually acid.

Renal Inadequacy.

What is renal inadequacy?

Some individuals never pass more than a normal or slightly less than normal quantity of urine of low specific gravity and containing abnormally little urea, even though they should partake of an excess of fluids. The condition has been termed *renal inadequacy.* The symptoms are vague and indefinite. There is a sense of ill-being, an undue readiness of fatigue, and a lowered resistance to disease. Patients so affected bear operative interference badly. There may be slight edema, but the urine contains neither albumin nor casts.

Displacements of the Kidney.

To what displacements is the kidney subject?

The kidney may undergo several degrees of displacement and in several ways. It may be congenitally fixed in some unusual situation, as behind the umbilicus, in front of the vertebral column, in the iliac fossa, at the sacro-iliac junction—*displaced kidney.* Secondly, the kidney may acquire greater or less

freedom of movement, though practically confined within its normal situation—*movable kidney.* Finally, provided with a mesonephron, it may possess a wide range of movement—*floating kidney.*

Congenital displacement of the kidney usually is unattended with symptoms and escapes recognition during life. Undue mobility of the kidney is much more common in women than in men, and on the right side than on the left. It may be induced by constriction of the waist, alternately repeated abdominal distention and relaxation, traumatism, heavy lifting, and adjacent disease, and it may occur as part of a general splanchnoptosis.

The condition may be unattended with symptoms, but frequently there is pain in the loin, with a sense of dragging and discomfort. Neurotic manifestations, neurasthenia, hysteria, hypochondriasis, are not uncommon accompaniments. The organ may be merely palpable, or it may be found quite out of place, although susceptible of reposition. Floating kidney may be attended with crises of abdominal pain, with chills, fever, nausea, vomiting, collapse. By twisting of the ureter, urine may accumulate in the renal pelvis, to be evacuated when the quantity has reached a certain limit.

Renal Calculus.

To what symptoms does a renal calculus give rise?

A calculus in the pelvis of the kidney occasions constant dull pain in the loin, aggravated into great intensity in paroxysms, superinduced by active physical exercise or by concussion. Following the paroxysm the urine contains blood. In the interval the urine contains pus. If the calculus enter the ureter, efforts at its expulsion give rise to excruciating pain in the course of the ureter, with retraction of the corresponding testicle, numbness of the thigh, nausea and vomiting. When a renal calculus is a cause of obstruction to the flow of urine, hydronephrosis or atrophy of the kidney may result. Renal calculi may be constituted of uric acid, of oxalates, of phosphates and less commonly of cystin and xanthin.

Acute Nephritis.

What are the clinical features of acute nephritis?

Acute inflammation of the kidney is common in association with acute infectious diseases—especially scarlatina, erysipelas, small-pox and yellow fever. It may follow exposure to cold, or result from the ingestion of toxic substances. It sometimes develops in the course of pregnancy, and occasionally in conjunction with cutaneous lesions, such as burns. The kidneys are large and smooth and reddened. The inflammation particularly involves the parenchyma of the organs. The occurrence of acute nephritis may be manifested by a chill, which is followed by fever, by pains in the loins, by headache, perhaps with nausea, vomiting and diarrhea. The *face* (notably the eyelids) is puffy; the extremities are swollen and edematous. The excretion of *urine* is diminished to a minimum; there may be actual suppression. The urine presents a smoky, reddish, turbid aspect; its specific gravity is high; it is deficient in urea, but it contains a decided quantity of albumin and hyaline and epithelial casts, and blood-casts, as well as free blood-corpuscles. The symptoms of *uremia* may supervene; to the headache, nausea and vomiting, are added derangement of vision, delirium, convulsions and coma. In the course of several weeks, the symptoms gradually subside, though the accompanying anemia yields but obstinately, while the urine may continue albuminous for a long time, perhaps permanently. Inflammations of the serous membranes are frequent in the course of acute nephritis.

Chronic Parenchymatous Nephritis.

What are the symptoms of chronic parenchymatous nephritis?

Recovery from acute inflammation of the kidney may be imperfect; repeated attacks may leave a permanently damaged organ; finally an inflammation of the parenchyma of the kidney may be insidious and essentially chronic from the outset. However produced, the *symptoms* are variable. The essential characteristic is the persistent presence of a considerable quantity of albumin in the urine, together with granular, hyaline

or fatty tube-casts. The quantity of urine may scarcely deviate from the normal. A characteristic variety of retinitis may be detected with the ophthalmoscope. Edema of the face and extremities occurs, anemia is manifest, assimilation is impaired, the arterial tension is heightened, and the heart becomes enlarged. Ascites is common. Edema of the larynx or lungs, or effusions into the pleuræ or pericardium may take place suddenly or gradually. Headache, impaired vision, somnolence, convulsions and coma, from uremia, may develop. Individuals affected with chronic parenchymatous nephritis badly resist disease. The probabilities are that the so-called fatty kidney is but a modification of the kidney of chronic parenchymatous nephritis, dependent upon secondary degeneration. The symptoms of both conditions are much alike. In the secondary degeneration, the presence of fatty casts and of oil-globules is characteristic.

Amyloid Kidney.

What are the clinical features of amyloid disease of the kidney?

Amyloid, lardaceous or *waxy disease of the kidney* depends upon a peculiar hyaline degeneration of the coats of the renal capillaries, which is often associated with some degree of chronic parenchymatous nephritis, and with amyloid degeneration in other organs, such as the liver, the spleen and the intestine. The *symptoms* are practically those of the associated condition. In addition to edema, albuminuria, the presence of tube-casts in the urine, heightened arterial tension and cardiac hypertrophy, there may be a peculiarly waxy complexion, diarrhea, and painless enlargement of the liver, without ascites and jaundice. Amyloid degeneration results usually in consequence of protracted suppuration or ulceration, or of syphilis.

Chronic Interstitial Nephritis.

What are the symptoms of chronic interstitial nephritis?

The interstitial structure of the kidney participitates, in some degree, in every inflammation of the organ. The involvement

becomes especially obvious in the late stages of all forms of nephritis. In some cases, however, the interstitial connective tissue undergoes hyperplasia from the outset, ultimately contracting and giving rise to a diminution in the size of the organ, which becomes small, red, granular ; the capsule is thickened ; on attempting to remove it the adherent cortex tears. This chronic interstitial nephritis is often but one part of a widely distributed arterio-capillary fibrosis. The principal *symptom* referable to the kidneys is frequent micturition, the urine being considerably increased in quantity, of a low specific gravity, and containing a trace of albumin and hyaline and granular tube-casts. Albumin and casts are sometimes only to be found by repeated, careful examination. There is little or no edema, unless dilatation of the heart develop. There is shortness of breath. The arterial tension is high and the heart is hypertrophied. The second sound of the heart is accentuated. The appetite is impaired, digestion is deranged and vomiting may occur. There is sometimes persistent diarrhea. Retinitis, papillitis, and hemorrhages may occur. *Death* usually results from uremia or cerebral hemorrhage. The commonest causes of chronic interstitial nephritis are alcoholism, gout and lead-poisoning.

What are the symptoms of uremia?

Uremia is an auto-intoxication, which sometimes arises in the course of disease of the kidneys and is dependent upon the retention in the fluids of the body of certain products that it is the function of the kidneys to remove. It may be acute or chronic. Uremia gives rise to headache, vertigo, nausea, vomiting, diarrhea, motor palsy, tremulousness, dyspnea, dilatation of the pupil, altered vision, delirium, somnolence, convulsions and coma. Effusions and pulmonary edema frequently accompany the condition. The temperature may be subnormal. It has exceptionally been observed to be febrile. The body and breath exhale a musty odor. Uremia is one of the gravest complications of disease of the kidneys.

How are a uremic seizure and a paroxysm of epilepsy to be differentiated?

Probably dependent upon analogous conditions of irritation,

there is little difference between the convulsions. The onset of an epileptic attack is sudden, attended with a cry and biting of the tongue, and possibly preceded by an aura; the convulsions of uremia are usually preceded by headache, vertigo and nausea, and occur without sound or injury to the tongue. The urine may contain albumin during or after an epileptic paroxysm, but casts are present as well when the convulsions are uremic. Epileptic seizures are far more likely than uremic convulsions to have been and to be repeated. A knowledge of the previous history of the case simplifies the diagnosis.

How are uremia and insolation to be differentiated?

The failure on the part of the skin as well as on the part of the kidneys to eliminate certain excrementitious products may in cases of insolation give rise to phenomena resembling those presented by uremia; but in the former the temperature attains a height never seen in the latter, while the season of the year at which the accident happens, as well as the attendant circumstances, makes the diagnosis clear.

Abscess of the Kidney.

What are the symptoms of abscess of the kidney?

Suppuration may take place in the kidney as the result of an acute infectious nephritis, as a part of a general pyemia, or in consequence of traumatism. When pus has formed, the *symptoms* occasioned are those to which imprisoned pus usually gives rise: rigors, hectic fever, sweats, emaciation. The *urinary secretion* will be altered in accordance with the intensity and extent of the causative or associated nephritis. In the urine may be found albumin, pus, tube-casts and blood-corpuscles. There may be pain and tenderness in the region of the kidney. If the accumulation of pus attains considerable proportions it may give rise to a palpable tumor, with fluctuation, in the loin. The abscess may rupture into the pelvis of the kidney, into an adjacent viscus, into the abdominal cavity; or the pus may make its way out through the abdominal wall.

Perinephric Abscess.

What are the clinical features of perinephric abscess?

Suppuration may take place in the perinephric connective tissue as a result of an extension of inflammation in the pelvis or in the structure of the kidney or as a result of blows or injuries. The condition is to be recognized by the existence of pain and tenderness in connection with a boggy, fluctuating tumor in the loin, associated with chills, fever, sweats and emaciation. The tumor is not movable, and cannot be separated from the kidney. Exploratory puncture may detect the presence of pus. Such an abscess may burrow and find an exit above or below Poupart's ligament, in the loin, into the kidney, ureter or bladder, into the peritoneum, intestines or into a lung.

How are a perinephric and a renal abscess to be differentiated?

An abscess in the loin developing in the course of pyemia is more likely to be renal than perinephric; an abscess following traumatism is more likely to be perinephric than renal; while disease of the pelvis or of the kidney may give rise to one or the other. The pus of a perinephric abscess is more likely than that of a renal abscess to burrow and point at some distant spot. In cases of renal abscess evidences of disease are likely to appear in the urine; there may be pus with albumin and tube-casts.

Tuberculosis of the Kidney.

What are the clinical features of tuberculosis of the kidney?

Miliary tubercles may be developed in the kidneys in common with other organs, as a part of an acute miliary tuberculosis. More usually, when the kidney is tuberculous, the foci are more extensive, and exhibit a tendency to undergo caseation. The kidney is rarely the exclusive seat of tuberculosis. As a rule, other portions of the genito-urinary tract are likewise involved, and not uncommonly the lungs as well. Miliary tuberculosis of the kidney does not give rise to symptoms apart from those of

the general condition of which it is but an incident. In the more localized lesion, the symptoms are vague. There may be pain in the loin, with tenderness; exceptionally a tumor is discoverable on physical examination; the urine may contain pus and detritus, in which tubercle-bacilli can, perhaps, be found; sometimes there is hematuria; there may be hectic fever. Ultimately emaciation sets in, exhausting sweats take place and the vitality fails. Tuberculosis of the kidney is more frequent in the young than in the old, in males than in females.

Malignant Disease of the Kidney.

What are the clinical manifestations of malignant disease of the kidney?

Malignant disease of the kidney is sarcomatous or carcinomatous. Sarcoma may be congenital or it may appear during childhood. Carcinoma rarely develops before forty. The classical symptoms are tumor, pain and hematuria.

The swelling occupies one of the loins, and is separated from the anterior abdominal wall by the large intestine. The pain is dull and heavy, with acute exacerbations. Hematuria is not constant. Its absence is of less significance than would be its presence. In addition, the appetite fails; there may be nausea or vomiting, constipation or diarrhea; emaciation sets in; and cachexia develops. Edema of the lower extremities and ascites may result from pressure on the veins.

Hydronephrosis.

What is hydronephrosis?

When there exists at any part of the genito-urinary tract an obstruction to the flow of urine, such as may be occasioned by a calculus, a neoplasm, an enlarged prostate gland, a kink or a twist in, or pressure from without upon, a ureter, fluid accumulates behind the point of obstruction, and causes dilatation. The condition is designated as hydronephrosis. It is sometimes congenital and it may be either unilateral or bilateral. If the obstruction be situated in the urethra or at the orifice of the bladder, the bladder, the ureters, and the pelves of

the kidneys all become thickened and dilated. If but one ureter is obstructed the corresponding pelvis suffers. The dilatation of the pelvis gives rise to compression and atrophy of the renal parenchyma. Sometimes the fluid resembles pale urine of low specific gravity; at other times it contains pus. In the latter case the condition is designated *pyonephrosis*. The symptoms are ill-defined. There may be dull pain, a sense of fulness, and a fluctuant tumor in one or the other loin, attended perhaps with the periodic discharge of large quantities of urine, as the obstruction is temporarily relieved, with a corresponding recession of the symptoms. There may, in addition, be rigors, hectic fever, sweats and emaciation. Constipation due to pressure upon the bowel, is not uncommon in cases of hydronephrosis. The urine may be normal, except at such times as the periodic copious discharges take place, when pus or other matters may be found present. In cases of double hydronephrosis there is great liability to the sudden development of uremia.

Hydatid Cyst of the Kidney.

What are the symptoms occasioned by an hydatid cyst of the kidney?

The kidneys are relatively a common seat of hydatid cysts. Such a cyst in connection with the kidney may give rise to a fluctuating tumor in the loin or in the lateral region of the abdomen, yielding the characteristic thrill on percussion, and occasioning symptoms dependent upon pressure; or prior to surgical exploration, it may not give rise to definite symptoms and may elude physical exploration. Under any circumstances a diagnosis may not be possible, unless the contents of the cyst find their way into the genito-urinary tract and are evacuated; when the detection of the hooklets in the urine leaves no room for doubt. The recognition of an additional cyst in another situation may facilitate the diagnosis.

How are an hydatid cyst of the kidney and hydronephrosis to be differentiated?

An hydatid cyst of the kidney and hydronephrosis both occasion a fluctuating renal tumor, but the peculiar thrill on

percussion that characterizes many hydatid cysts is wanting in all cases of water-logged kidney. Hydronephrosis is dependent upon some form of genito-urinary obstruction ; such a condition bears no relation to the development of an hydatid cyst. The periodic discharge of a large quantity of fluid, with a temporary disappearance of the tumor, is characteristic of hydronephrosis ; should a discharge of fluid take place, in connection with an hydatid cyst, microscopic examination will reveal the characteristic hooklets in the urine.

THE NERVOUS SYSTEM.

How is a study of diseases of the nervous system to be approached?

By an investigation of motion, sensation, reflex action, nutrition, electric reaction and cerebral functions.

How may motion be affected?

The power of muscular movement may be impaired or lost ; motility may be exaggerated ; or coördination may be impaired. Complete loss of muscular power is called *paralysis;* partial loss, *paresis.* Motor over-action is seen in spasm and convulsion. Incoördination gives rise to *ataxia.*

The power of voluntary motion is impaired or lost in destructive disease at any part of the motor tract, from the cerebral cortex to the end-plates. Movement may be interfered with by disease of the muscle itself, or of an adjacent joint ; or abstained from on account of the pain that it causes in a joint or muscles.

Loss of motor power confined to one side of the body is called *hemiplegia ;* if bilateral, *diplegia ;* limited to the lower extremities, *paraplegia ;* localized to a member, *monoplegia.*

Hemiplegia is usually dependent upon a unilateral cerebral lesion ; it may possibly be caused by a unilateral lesion seated in the upper portion of the cord.

Diplegia may be a result of symmetrical bilateral cerebral lesions, or of a lesion in a situation where the motor tracts are close together ; like hemiplegia, it may possibly be caused by a lesion in the upper portion of the spinal cord.

Paraplegia is almost invariably a result of disease of the cord.

Monoplegia is a result of a circumscribed lesion of the brain, usually in or near the motor cortex. Any form of paralysis may be simulated by hysteria.

Spasm or *convulsion* is indicative of irritation at any part of the motor tract. It may occur in acute disease, cerebral (hemorrhage, embolism, thrombosis) or constitutional (tetanus, hydrophobia); in chronic irritative disease (cortical softening, tumor); or it may be a symptom of nutritional disease (epilepsy), or of functional disease (hysteria); it may also occur in disease not strictly nervous (uremia).

Ataxia is the result of derangement of the conducting fibers or the coördinating centers in the cord or cerebellum. Loss of tactile sensation also gives rise to ataxia. Defective equilibration may result from disease of the internal auditory apparatus.

What are the varieties of sensation?

Common or tactile sensibility, the pain-sense, the temperature-sense, the muscular sense, the stereognostic sense; special sensation: smell, vision, taste and hearing.

How may sensation be affected?

Sensation may be impaired, retarded, lost, perverted or heightened. Impairment of sensation is *hypesthesia;* loss of sensation is *anesthesia;* heightening of sensation, *hyperesthesia;* perversion of sensation, *paresthesia.* Alterations in sensation are dependent upon derangement in the course of the sensory path. As the lesion may be destructive or irritative, so may the symptoms be those of diminished sensation, of supersensibility, or of perverted sensation. One or the other variety of sensation may be affected without involvement of all.

What are the varieties of reflexes?

Superficial or cutaneous, and muscular or deep.

What are the principal superficial reflexes

The plantar, the gluteal, the cremasteric, the abdominal, the epigastric, the conjunctival, the reaction of the iris to light, and to cutaneous irritation.

What are the principal varieties of deep reflexes?

The knee-jerk, ankle-clonus, the tendo-Achillis jerk, the biceps-jerk, and the elbow-jerk.

How are the reflexes altered in disease?

They may be exaggerated, enfeebled or entirely lost.

The superficial reflexes are exaggerated when the influence of the inhibitory apparatus is removed, or when the activity of the reflex apparatus of the spinal cord is heightened; the deep reflexes are exaggerated under similar conditions, and likewise when degeneration has taken place in the cerebro-spinal motor tract. Both superficial and deep reflexes are lost when the reflex arc is interrupted at any point in its course—sensory nerve, reflex center in the cord, and motor nerve.

How is the nutrition affected by disease of the nervous system?

Trophic changes often attend disease of the nervous system. Bedsores sometimes form. Muscles may waste. A paralyzed limb is edematous and its temperature is subnormal; in children the growth of paralyzed parts is retarded. Decided alterations in joints may take place. The skin sometimes becomes glossy. Trophic changes are most marked in degenerative disease of the nerves and of the gray matter of the cord, especially of the anterior horns. The nutrition of a palsied member slowly suffers.

How are the electric reactions affected by disease?

The normal response of muscle and nerve to both the voltaic and faradic currents may be enfeebled or exaggerated—a *quantitative change;* or the response of nerve to both faradism and voltaism, and of muscle to faradism may be enfeebled or lost, while the response of muscle to voltaism persists and may be exaggerated and is altered in its polar relations—a *qualitative change*, the *reaction of degeneration.*

Normally, the response of muscle to voltaic stimulation is most ready upon application of the kathode and making (closure) of the current; least ready upon application of the kathode and breaking (opening) of the current. Response upon anodal closure and anodal opening is less ready than that of kathodal closure, more ready than that of kathodal opening.

The sequence: (1) K.Cl.C.; (2) $\begin{matrix}\text{An.O.C.};\\ \text{An.Cl.C.};\end{matrix}$ (3) K.O.C.; is

called the normal formula. A deviation from this sequence gives a formula expressive of the *reaction of degeneration*—a *qualitative* change. Quantitative changes take place in simple wasting dependent upon disuse or disease of the muscle or of an adjacent joint, or upon interruption in the motor path between the cerebral cortex and the ganglion-cells of the anterior horns of the spinal cord; qualitative changes take place in degenerative muscular wasting dependent upon lesions of nerve or cord, between the motor end-plates and the ganglion-cells of the anterior horns inclusive.

How are the cerebral functions affected by disease of the nervous system?

Delirium, headache and vertigo may appear. Vision, hearing, taste or smell may be perverted, excessively acute, impaired or lost; memory may be enfeebled; speech may be altered or lost; articulation may be defective; the emotions may be affected; consciousness may be deranged; the higher mental powers may suffer deterioration.

Neuritis.

How may neuritis be classified?

Into perineuritis, interstitial neuritis and parenchymatous neuritis. Neuritis may be acute or chronic; it may affect one nerve or many.

What are the causes of neuritis?

Inflammation of a nerve may result from local injury or compression, from adjacent inflammation, from exposure to cold, or from the influence of irritating matters circulating in the blood, such as alcohol, metallic poisons (lead, arsenic, mercury), the toxic products of infectious diseases (diphtheria, syphilis, influenza), and of perverted metabolism (diabetes, nephritis, gout). Injury, compression, adjacent inflammation and exposure to cold give rise to circumscribed neuritis; irritants in the blood, as a rule, to multiple neuritis.

What are the symptoms of neuritis?

When a nerve is inflamed, hyperesthesia, paresthesia and

anesthesia, pain, tingling, numbness in the peripheral distribution and tenderness in the course of the nerve-trunk in turn appear, with impairment or loss of motion. If the exciting conditions be maintained, nerves degenerate and muscles waste and the skin becomes glossy. Exposure to cold is a cause of inflammation of the facial nerve; rheumatism, gout and diabetes, of inflammation of the sciatic; lead-poisoning, of inflammation of branches of the radial, median and ulnar; pressure, of inflammation of the musculo-spiral; syphilis and alcoholism, of inflammation of the peripheral nerves of the extremities; alcohol, tobacco, syphilis, of inflammation of the optic nerve. Obstinate neuralgias are sometimes dependent upon neuritis. Acute neuritis is attended with fever and other constitutional phenomena.

With what conditions may neuritis be confused?

With subacute rheumatism, with neuralgia and with disease of bone. The pain occurring in the course of diseases of the spinal cord may be mistaken for an evidence of neuritis.

How is the diagnosis of neuritis to be made?

The symptoms are limited to the distribution of one or more nerves and there is tenderness on pressure. When neuralgia is not dependent on neuritis, there are certain tender points, the pain shoots and intermits, and loss of sensation does not result. Pain of spinal origin is not local in distribution or associated with tenderness.

Multiple Neuritis.

What are the causes of multiple neuritis?

Multiple or *disseminated neuritis*, or *polyneuritis*, may be a complication or a sequel of infectious diseases, such as syphilis, diphtheria and influenza; it may develop in the course of posterior spinal sclerosis or it may appear as a part of anesthetic leprosy; it may be due to metallic poisoning, as with lead, mercury, arsenic; it may occur endemically, as the disease known as *kak-ké* or *beri-beri;* the most common causes, however, are chronic alcoholism and exposure to wet and cold.

What are the symptoms of multiple neuritis?

If multiple neuritis set in suddenly, it may present rigor, elevation of temperature and other febrile symptoms.

It is attended with numbness, tingling, pain, tenderness and loss of power in the distribution of the nerves affected, usually those of the extremities. Paralysis, wasting and impairment of sensation soon follow. There may be wrist-drop, foot-drop, abolition of reflexes and ataxia.

Trophic changes occur in the skin, nails and hair. Edema is not uncommon. The wasted muscles present reactions of degeneration. Secondary contractions may take place in the unopposed muscles.

FIG. 43.



Multiple neuritis, wrist-drop and foot drop. (Gowers.)

Nerves of special sense, as the olfactory, the optic, the auditory and the gustatory, may suffer.

How are spinal pachymeningitis and multiple neuritis to be differentiated?

Spinal pachymeningitis is usually cervical; the symptoms are thus especially referable to the upper extremities. There is also an absence of tenderness in the course of the nerves, while pain radiating from the spinal column is the rule.

How does acute myelitis differ from multiple neuritis?

Myelitis usually involves a limited area of the spinal cord in the dorsal or lumbar region. The symptoms are thus manifested in the lower half of the body and constitute the type of paraplegia. The action of the sphincters is impaired. There

is anesthesia, with a zone of hyperesthesia and but little pain. The paralyzed muscles waste rapidly. The reflexes are mostly exaggerated in the parts supplied by the nerves that arise below the affected area of the cord.

Sciatica.

What are the causes of sciatica?

Sciatica is usually dependent upon inflammation of the sciatic nerve.

It is more common in males than in females and in middle life than at any other period. It may be primary, developing in gouty, diabetic, or rheumatic persons or after exposure to wet and cold, or following injury, or as the result of pressure from sitting on the edge of a chair; or secondary, in consequence of compression by tumors in the course of the nerve, within or without the pelvis, or as a result of extension from adjacent disease, as of the hip-joint.

What are the symptoms of sciatica?

Sciatica is attended with pain and tenderness in the course and distribution of the sciatic nerve, increased by movement. There are certain tender points: 1, at the posterior inferior spine of the ilium; 2, at the sciatic notch; 3, at the middle of the thigh; 4, on the posterior aspect of the knee; 5, below the head of the fibula; 6, behind the external malleolus; and 7, on the dorsum of the foot. There are also paresthesia, tingling, formication and numbness.

In aggravated cases, the affected muscles waste and present degenerative reactions. There may also be fibrillary contractions and muscular cramps.

Primary sciatica is rarely bilateral.

With what conditions may sciatica be confounded?

With sciatic neuralgia, with disease of the hip-joint, and with disease of the spinal cord or cauda equina.

How is the differential diagnosis to be made?

Sciatic neuralgia occurs in debilitated and anemic persons, with neuralgia elsewhere; the pain is intermittent, and in the

distribution rather than in the course of the nerve; it may be induced by movement, but is not otherwise aggravated thereby; there is spontaneous pain rather than tenderness.

Disease of the hip-joint may give rise to pain about the hip-joint and knee; but there is no tenderness in the course of the nerve; and investigation will disclose the obvious cause of the symptoms.

The pain to which disease of the spinal cord or of the cauda equina gives rise is bilateral in distribution; it is not attended with tenderness in the course of the nerve; but it is associated with additional symptoms indicative of its origin.

How are primary and secondary sciatica to be differentiated?

The pain and tenderness in the course of the nerve in primary sciatica are wanting when the sciatica is secondary; the pain is then rather in the distribution of the nerve; other symptoms, and careful examination per rectum, if need be, may lead to the detection of an adequate cause for the pain.

Facial Hemiatrophy.

What is facial hemiatrophy?

Atrophy of one side of the face may begin in childhood or it may follow blows or injuries later in life. It is believed to be dependent upon disease of the trigeminal nerve. Both the soft parts and the bone are involved, including sometimes also the tongue, and the hair and the teeth on the affected side may fall out. The disease sometimes extends beyond the median line, and sometimes appears also in other parts of the body. The condition must be discriminated from *enlargement* of one side of the face.

Paralysis of the Facial Nerve.

What are the causes of paralysis of the facial nerve?

The facial nerve may be paralyzed by lesions above its nucleus (supra-nuclear), by lesions of the nucleus (nuclear), or by lesions in the course of the nerve (infra-nuclear). The first is repre-

sented by the paralysis of hemiplegia. The nucleus and the root-fibers may be damaged as a result of hemorrhage, softening, degeneration, or inflammation, or by a new-growth. The nerve may suffer from compression by tumors, from traumatism, from neuritis (primary or secondary). The most common cause of facial paralysis is exposure to cold.

What are the symptoms of paralysis of the facial nerve?

In facial paralysis of nuclear or infra-nuclear origin all of the muscles of one side of the face are paralyzed; the face is rendered asymmetrical by the unantagonized action of the muscles of the opposite side; the eye upon one side remains open and cannot be closed; the lower lid is relaxed and epiphora results; the forehead on that side is smooth and cannot be wrinkled; the angle of the mouth droops; saliva dribbles from the corner of the mouth; the normal furrows are obliterated; the lips cannot be puckered, as in whistling, nor elevated, as in displaying the teeth; associated and emotional movements are

FIG. 44.



Facial paralysis. The figure on the right represents an attempt to close the eyes. (Gowers.)

wanting on the paralyzed side. Numbness and tingling indicate involvement of filaments of the fifth nerve.

The affected muscles waste and reactions of degeneration develop. If complete recovery do not occur, secondary contraction in the paralyzed and wasted muscles takes place.

If the seventh nerve is injured between the geniculate ganglion and the origin of the chorda tympani, the sense of taste is lost in the anterior portion of the tongue.

How are facial palsy of supra-nuclear origin and that of nuclear or infra-nuclear origin to be differentiated?

Facial palsy of supra-nuclear origin is but part of a hemiplegia, of which other manifestations will be evident; unlike what takes place in nuclear and infra-nuclear disease, the orbicularis palpebrarum and the occipito-frontal muscles escape. In nuclear and infra-nuclear palsy, muscular wasting takes place, with degenerative reaction, which is not the case in palsy of supra-nuclear origin. In supra-nuclear paralysis, associated and emotional movements of the face are symmetrically performed; in nuclear or infra-nuclear palsy, they are not.

Paralysis of the Phrenic Nerve.

What are the causes of paralysis of the phrenic nerve?

The function of the phrenic nerve may be interfered with by disease of the spinal cord or of the vertebræ, by the pressure of a tumor or of an aneurism, as a result of a neuritis or of a deep wound of the neck.

To what symptoms does paralysis of the phrenic nerve give rise?

When the phrenic nerve is paralyzed, the movements of the diaphragm are enfeebled or abolished. If the nerve on one side is affected, the loss of power is not marked; but if both are paralyzed, the diaphragm does not actively descend in inspiration; instead of the protusion of the upper portion of the abdominal wall, there is retraction; dyspnea is induced by exertion; cough becomes difficult.

From what conditions is paralysis of the phrenic nerves to be differentiated?

From superior intercostal breathing, from inflammation of the diaphragm and from degeneration of the diaphragm. In the first condition, inspiratory contraction of the diaphragm must

be carefully looked for; in the second, there has been an adjacent inflammation, most probably of the pleura or peritoneum; the third has only been recognized after death.

If the paralysis of the phrenic nerves is dependent upon disease of the cord, there are present other symptoms than inactivity of the diaphragm.

Paralysis of the Musculo-spiral Nerve.

What are the causes of paralysis of the musculo-spiral nerve?

The musculo-spiral nerve may be paralyzed as a result of traumatism, as from a blow or when the humerus is fractured; the nerve may be compressed by callus, by contraction of the triceps muscle, or by the head of a crutch in the axilla; neuritis may result from exposure to cold. The most common cause of musculo-spiral paralysis is pressure by the head on the arm during sleep.

How is paralysis of the musculo-spiral nerve to be recognized?

The characteristic symptom of paralysis of the musculo-spiral nerve is loss of power in the extensors of the elbow and wrist, in the long extensors of the fingers and thumb, and in the supinators. There is wrist-drop and the fingers and thumb cannot be extended. From the want of antagonism the power of flexion is also enfeebled. Subsequently, the palsied muscles may present reactions of degeneration.

How does the neuritis of lead-poisoning differ from the neuritis of the musculo-spiral nerve resulting from compression or traumatism?

Both are attended by wrist-drop, which in lead-poisoning is bilateral, in compression-neuritis unilateral. In the former, the onset is slow and gradual and the action of the supinators is unimpaired; in the latter, the onset is acute and supination cannot be performed. A blue line on the gums attends lead-poisoning; careful chemic examination during the administration of potassium iodide may detect lead in the urine; while in musculo-spiral palsy a cause of a different kind is readily ascertainable.

Neuromata.

To what symptoms do neuromata give rise?

Tumors in the course of nerves may be composed of nervous structure or be heterologous. They may be hereditary, congenital, or a result of traumatism; they are not uncommonly seen on the stumps of amputated members. The symptoms that such growths occasion necessarily depend upon their situation. At first they give rise to irritation and heightened function, as evidenced by pain and spasm; subsequently they cause abolition of function and paralysis.

Upon what does the diagnosis of neuromata depend?

Upon the detection of a tumor and the obstinacy of symptoms referable to the distribution of one or more nerves.

Neuralgia.

What are the clinical features of neuralgia?

Pain in the course of a nerve, independent of neuritis, appears most commonly in adult life, in anemic and ill-nourished women of emotional temperament. In some instances, an hereditary predisposition can be traced. Among the causes of neuralgia are exposure to cold, rheumatism, gout, malaria, alcoholism, plumbism, traumatism and peripheral irritation.

The pain occurs in paroxysms, in the intervals between which some sensitiveness persists. In the course of the nerve are certain tender points — *points douloureux* of Valleix. Muscular spasm sometimes takes place in the distribution of the motor nerve corresponding to the sensory nerve involved. The paroxysm is sometimes attended with vomiting. Vaso-motor derangement and trophic changes may manifest themselves in the course of the disease. Neuralgia may be localized to a single nerve; it may progress in a radiating manner, or it may change its seat from nerve to nerve.

How are neuritis and neuralgia to be differentiated?

The pain of neuralgia is intermittent and paroxysmal; that of neuritis is continuous; in neuritis there is, in addition, ten-

derness in the course of the nerve, with swelling. Neuritis is not limited to sensory nerves; in consequence, muscular weakness and, later, wasting and alterations in the electrical reactions are manifest.

What are the symptoms of trigeminal neuralgia?

The symptoms of *neuralgia of the fifth nerve*, *trigeminal* or *trifacial neuralgia*, or *tic douloureux*, depend upon its distribution. One or several branches of the nerve may be involved.

Of the first division, the supra-orbital branch is most commonly affected. Pain is felt on the forehead, on the eyelid, in the eye and on the side of the nose. There are supra-orbital, palpebral, nasal and ocular tender points.

Of the second division of the fifth nerve, the infra-orbital branch is most commonly attacked by neuralgia. Pain is referred to the cheek and ala nasi, between the orbit and the mouth. There are infra-orbital, nasal, malar and gingival tender points.

The pain of neuralgia of the inferior maxillary nerve is referred to the parietal eminence, the temple, the lower jaw, the ear and the tongue. There are inferior dental, temporal and parietal tender points. Sometimes the inferior dental and the auriculotemporal branches are alone involved.

Migraine.

What are the clinical features of migraine?

Migraine, *hemicrania* or *sick headache* is a paroxysmal neurosis characterized by unilateral headache, associated with nausea, vomiting and derangement of vision and sensation. It is more common in females than in males and in the first half of life than at any other time. Frequently a neurotic heredity can be traced, some members of the same family presenting migraine, epilepsy, neuralgia, insanity or some other neurosis.

The paroxysm often begins with some sensory disturbance, such as tingling or numbness; or with a perversion of vision, such as the appearance of a luminous or brightly colored object; or with an impairment of vision usually presenting the char-

acters of hemianopsia or with auditory manifestations, such as tinnitus or a sudden explosive sound. When the headache reaches a considerable degree of intensity nausea and vomiting occur. In extreme cases motor weakness and aphasia are present. The headache, at the beginning of the disease unilateral, may subsequently become bilateral. It is often periodic, and not rarely in some fixed relation with menstruation, which it may accompany, or precede or follow at a certain interval.

It may sometimes be traced to reflex influences, such as may arise from gastric disturbance or eye-strain. These, however, are to be regarded merely as excitants, acting upon a predisposed nervous system, and would not alone be effective.

How are migraine and ordinary headache to be differentiated?

Headache may arise from a multiplicity of conditions—among others from neuralgia, from rheumatism, from gastro-intestinal derangement, from toxemia, from eye-strain, from anemia, from hyperemia and from organic disease of the brain. It is unassociated with other subjective or with visual manifestations; it is bilateral; it is not paroxysmal; it is dependent upon other influences than is migraine.

Spinal Meningitis.

What are the varieties of spinal meningitis?

Spinal meningitis may be *acute* or *chronic;* it may involve the dura, arachnoid or pia, giving rise respectively to *pachymeningitis*, *arachnitis* or *leptomeningitis;* it may be *simple*, *purulent*, *tuberculous* or *hemorrhagic.*

What are the causes of spinal meningitis?

Inflammation of the spinal meninges may result from an extension of adjacent inflammation, from traumatism, from exposure to cold and wet; it may develop in the course of infectious diseases, or as a manifestation of syphilis, of tuberculosis, of pyemia or of alcoholism.

What are the symptoms of acute spinal meningitis?

Meningitis may set in abruptly with a chill, followed by elevation of temperature, pain in the back and radiating in the course

of the nerves, hyperesthesia, stiffness and spasmodic contraction of various muscles, exaggerated reflexes, retention of urine and constipation. The pain is intensified in paroxysms. The muscular spasms are induced by efforts at movement. The finger drawn over the skin leaves a red streak.

With the subsidence of the acute symptoms, the pain may yet remain, but the manifestations of irritation give way to those of paralysis: muscular weakness and wasting, anesthesia, abolition of reflexes and enfeeblement of the sphincters. The extent and distribution of the symptoms depend upon the seat of the meningeal inflammation.

How are rheumatism of the muscles of the back and spinal meningitis to be differentiated?

Pain may accompany dorsal or lumbar rheumatism, but it is not radiating, nor is it attended with muscular rigidity, or followed by anesthesia, loss of power and wasting.

Cervical Pachymeningitis.

What are the symptoms of cervical pachymeningitis?

Inflammation of the dura mater in the cervical region is recognized by pain and stiffness in that region, with shooting pains in the arms, and numbness, and impaired sensibility, and loss of power, and wasting. The over-extension of the hand, with flexion of the fingers, that results from the paralysis of the long flexors of the wrist and fingers, and of the interossei gives rise to characteristic deformity. There may also be weakness of the lower extremities.

How is cervical pachymeningitis to be distinguished from progressive muscular atrophy and subacute anterior poliomyelitis?

All three may give rise to wasting and weakness in the upper extremities, and weakness in the lower; but in pachymeningitis sensation is deranged, the wasting is irregular in distribution, and there have at some time been stiffness in the muscles of the neck, and pains in the back and radiating down the arms; while spasm and sensory symptoms are wanting in progressive muscular atrophy and poliomyelitis.

Hemorrhage into the Spinal Membranes.

What are the varieties of hemorrhage into the spinal meninges?

Hemorrhage may take place between the dura and the vertebræ, being then *extrameningeal;* or within the dura, being then *intrameningeal.* Intrameningeal hemorrhage may take place between the dura and the arachnoid, being then *subdural;* or between the arachnoid and the pia, being then *subarachnoid.*

What are the causes of hemorrhage into the spinal meninges?

Hemorrhage into the membranes of the cord is usually a result of traumatism, as from falls or blows, or of violent muscular activity, as during convulsions or labor; it may occur in the course of acute infectious diseases or as a manifestation of a hemorrhagic diathesis; or an aneurism that has eroded the vertebræ may rupture and empty its contents into the spinal canal.

What are the symptoms to which hemorrhage into the spinal membranes give rise?

Hemorrhage into the spinal meninges is marked by sudden, severe pain in the back, to which are soon added rigidity, muscular spasm, radiating pains, hyperesthesia, retention of urine and constipation; should death not occur, these symptoms in turn give way to muscular weakness, anesthesia and derangement of the sphincters.

How are spinal meningitis and hemorrhage into the spinal meninges to be differentiated?

The onset of hemorrhage is sudden, that of meningitis gradual. Meningitis is from the outset attended with elevation of temperature; in hemorrhage, the temperature rises only when meningitis sets in.

Anemia of the Spinal Cord.

What are the manifestations of anemia of the spinal cord?

Anemia of the spinal cord may be part of a general anemia. There is no definite symptom by which it can be recognized.

Whatever the symptoms to which it may give rise, they are comprised in those of the condition of which the spinal anemia is a part. There is aching in the legs and the patient fatigues readily. There may be wasting of the extremities and paraplegia.

Hyperemia of the Spinal Cord.

What are the symptoms of hyperemia of the spinal cord?

The clinical recognition of hyperemia, other than as a condition antecedent to inflammation of the spinal cord, is as doubtful as is that of spinal anemia. A diagnosis of hyperemia is justifiable when symptoms apparently indicative of a beginning myelitis disappear in the course of a few days.

Myelitis.

What are the varieties of myelitis?

Myelitis may be *acute*, *subacute* or *chronic;* it may be *transverse*, *focal*, *disseminated*, *diffuse* or *central;* or it may involve only the anterior horns of the gray matter.

What are the causes of myelitis?

Myelitis is more common in males than in females. It may result from traumatism, from exposure to cold and wet, from compression, by extension from adjacent disease and in the course of acute infectious diseases; syphilis may be a cause of myelitis.

What are the symptoms of acute myelitis?

The symptoms of acute myelitis *set in* with variable abruptness and with febrile manifestations. There may at first be pains of moderate severity in the back. Soon, there is loss of motion and sensation in the parts supplied by the nerves arising from the cord below the level of the inflammation, with a girdle-sensation and a zone of hyperesthesia corresponding to the distribution of the nerves given off at the upper limit of the inflammation. Grave trophic changes, as wasting and the formation of bed-sores, may take place, with the development of the reactions of degeneration. The reflexes within the dis-

tribution of the nerves arising from the inflamed area are lost; those below are exaggerated. Control of the sphincters is lost.

After the subsidence of the acute symptoms, some improvement slowly takes place, greater in respect to sensation than in respect to motion. Cystitis and consecutive pyelonephritis may develop. Contractures may take place. The inflammation of the cord may extend.

How are acute spinal meningitis and acute myelitis to be distinguished from one another?

The two are likely to be associated; the symptoms of the one or the other, however, predominating. Pain, hyperesthesia and muscular spasm, followed by anesthesia and palsy, characterize acute meningitis. In myelitis, pain and hyperesthesia are slight, transient and circumscribed; muscular spasm is wanting; and wasting, palsy, anesthesia and derangement of the sphincters appear early.

How are acute myelitis and spinal hemorrhage to be differentiated?

The onset of hemorrhage is more abrupt than that of myelitis. Initial fever indicates myelitis rather than hemorrhage. Hemorrhage into the spinal cord is soon followed by myelitis, and the differentiation then becomes impossible.

Chronic Myelitis

What are the causes of chronic myelitis?

Chronic myelitis is more common in males than in females, and in early and middle adult life than at any other period. The same causes that occasion acute myelitis may also give rise to *chronic myelitis.* Thus chronic myelitis may follow syphilis, repeated exposure to cold and wet, repeated over-exertion, compression of the cord, chronic alcoholism and lead-poisoning. Chronic myelitis may be a sequel of acute myelitis or of chronic meningitis.

What are the symptoms of chronic myelitis?

The symptoms of chronic myelitis vary in distribution with

the localization of the disease. There is at first an undue readiness of muscular fatigue, soon progressing to weakness and followed by actual palsy. *Sensation* is deranged ; paresthesiæ are common ; girdle-pain may appear. The *reflexes* are exaggerated. A tendency to spasm develops. The *muscles* waste and present quantitatively altered electric reactions, which, in the later stages of the disease, suffer qualitative change. The *sphincters* usually participate in the loss of motor power.

How does chronic differ from acute myelitis ?

The symptoms of chronic myelitis, unlike those of acute myelitis, are slow and gradual in onset, progressive in course, irregular in distribution and unattended with febrile reaction.

What are the distinctions between chronic myelitis and progressive muscular atrophy ?

The symptoms of chronic myelitis are irregular in distribution ; the manifestations of progressive muscular atrophy appear somewhat symmetrically. Sensory symptoms constitute a distinctive feature of chronic myelitis, but are inconsiderable or wanting in progressive muscular atrophy. Exaggeration of reflexes and spasm characterize myelitis rather than muscular atrophy. Fibrillary muscular contractions and heightened mechanical irritability mark progressive muscular atrophy.

In what respects does chronic myelitis differ from lateral sclerosis ?

In lateral sclerosis, sensation is not deranged ; there is no girdle-sensation ; and the action of the sphincters remains unimpaired.

What is the distinction between chronic myelitis and spinal pachymeningitis ?

Spinal pachymeningitis is most commonly cervical and the symptoms are limited to the upper extremities ; myelitis irregularly involves all four extremities. Pain is a more constant manifestation of pachymeningitis than of myelitis.

Acute Anterior Poliomyelitis.

What are the causes of acute anterior poliomyelitis?

Acute inflammation of the anterior horns of the gray matter of the spinal cord is more common in males than in females, in late infancy than at any other period of life, and in summer than in winter. It follows sudden chilling of the heated body; injuries to the spine; acute infectious diseases; over-exertion.

What are the symptoms of acute anterior poliomyelitis?

Acute anterior poliomyelitis may set in abruptly, with convulsions, vomiting, diarrhea, delirium, headache, dull, heavy pains in the back and in the extremities, and febrile symptoms. Soon extensive palsy appears, followed, after a variable period, by wasting of certain groups of muscles. There may have been incontinence of urine and of feces.

In the course of several weeks the original extent of paralysis gradually diminishes, until ultimately the loss of power and the wasting are limited to certain parts that remain permanently deficient in nutrition, growth and function. Sensibility is unaffected.

Degenerative reactions of nerve and muscle develop and the reflexes are enfeebled or lost.

What are the distinctions between myelitis and poliomyelitis?

The permanent palsy is more extensive in myelitis than in poliomyelitis. In the former the reflexes are exaggerated; in the latter, enfeebled or lost. Sensory symptoms characterize myelitis; not poliomyelitis. Bedsores are common in the former; not in the latter.

How are the paralysis of poliomyelitis and that of cerebral origin to be differentiated?

The paralysis of poliomyelitis may be limited to one side; but it is not characteristically hemiplegic. It is attended with wasting and the reaction of degeneration; cerebral paralysis may be attended with wasting, but not with the reaction of degeneration. Athetoid movements appear in the course of palsy of cerebral origin; not in the course of the palsy of poliomyelitis.

The reflexes are exaggerated in cerebral disease; enfeebled or lost in poliomyelitis.

How are multiple neuritis and anterior poliomyelitis to be differentiated?

Sensory symptoms—tingling, pain, tenderness, numbness and anesthesia—as seen in neuritis, are wanting in poliomyelitis. Multiple neuritis is rather symmetrical, poliomyelitis irregular in distribution. The one is the more common in adults; the other, in children.

Acute Ascending Paralysis.

What are the symptoms of acute ascending paralysis or Landry's paralysis?

The nature of acute ascending paralysis is not yet known. The disease is thought to depend upon a toxic influence exerted upon the nervous system. It arises under conditions similar to those that precede acute myelitis. It is characterized by progressive motor paralysis, beginning in the lower extremities and gradually extending upwards. The power of movement is lost; respiration, deglutition, and articulation are interfered with. Numbness and tingling are sometimes present at the onset and followed by impairment of sensation. The reflexes are enfeebled. Trophic changes are wanting. The sphincters are usually uninvolved. The spleen is enlarged. Most cases terminate fatally.

How are acute myelitis and acute ascending paralysis to be differentiated?

Febrile symptoms, anesthesia, trophic changes and involvement of the sphincters do not attend acute ascending paralysis (Landry's disease). Recovery from acute ascending paralysis is less common than is recovery from myelitis.

How is acute ascending paralysis to be distinguished from multiple neuritis?

The view has been expressed that acute ascending paralysis is dependent upon multiple neuritis. Acute ascending paralysis, however, does not appear in all four extremities at once, and

is said not to present the alterations of sensation, the muscular wasting and the elevation of temperature of multiple neuritis.

What is the distinction between acute ascending paralysis and anterior poliomyelitis?

The onset of acute anterior poliomyelitis is, while that of acute ascending paralysis is not, attended with febrile manifestations. In poliomyelitis, the paralyzed muscles waste; in ascending paralysis, they do not. Poliomyelitis is rarely fatal; ascending paralysis is usually fatal.

Amyotrophic Lateral Sclerosis — Progressive Muscular Atrophy—Glosso-Labio-Laryngeal Palsy.

There are three affections intimately related to one another: *amyotrophic lateral sclerosis*, *progressive muscular atrophy* and *glosso-labio-laryngeal palsy*. The pathologic lesions are similar in all: a degenerative process in the gray matter and in the conducting paths. The symptoms of the three affections differ according to the situation of the disease in the gray matter, and according to whether the change begins first in the gray matter and then involves the white or *vice versa*.

If the white matter of the lateral columns is first involved, with partial or with subsequent invasion of the gray matter of the anterior horns, amyotrophic lateral sclerosis results. If the gray matter of the cord be first involved, the symptoms of progressive muscular atrophy appear; while if the process extend to or originate in the gray matter of the medulla the symptoms are those of bulbar palsy, superadded or occurring isolated.

What are the causes of amyotrophic lateral sclerosis, progressive muscular atrophy and chronic bulbar palsy?

In many cases, no cause can be determined; in others the symptoms have been preceded by mental strain, by exposure to cold and wet, by concussion, by syphilis, by metallic poisoning, as with lead, mercury or arsenic, and by acute diseases; in some there is an indirect neurotic heredity. All three affections are

more common in males than in females, and in middle life than at any other period.

What are the symptoms of amyotrophic lateral sclerosis?

Amyotrophic lateral sclerosis begins with symptoms of lateral sclerosis: weakness; a stiff, awkward, spastic gait; muscular spasm; exaggerated reflexes; spinal epilepsy; to which in turn may be superadded the symptoms of degeneration of the anterior horns of the gray matter: muscular wasting, paralysis, loss of reflexes, reaction of degeneration.

What are the symptoms of progressive muscular atrophy?

Progressive muscular atrophy is usually insidious in onset; sometimes slow, sometimes rapid, but always progressive in course. Muscular wasting, preceded by pain, is first observed in one portion of the body, usually in an upper extremity, and followed by weakness; both wasting and weakness in turn successively invade all four extremities and the trunk as well; respiration may be embarrassed. If the duration of the case be long enough, symptoms of bulbar paralysis are superadded.

The wasting is especially manifest in the thenar and hypothenar eminences and in the interosseous spaces; as a result, a peculiar deformity—the claw-like hand—develops. The wasting muscles are the seat of spontaneous fibrillary contractions. Mechanical muscular irritability is heightened. The deep reflexes are enfeebled in a degree proportional to the muscular wasting. The electric reactions are also dependent upon the nutrition of the muscles and nerves. At first, they present quantitative, subsequently qualitative changes.

What are the symptoms of chronic or progressive bulbar paralysis—glosso-labio-laryngeal palsy?

The symptoms of *glosso-labio-laryngeal paralysis* are analogous to those of progressive muscular atrophy, the degenerative process, however, involving the nuclei of the cerebral rather than those of the spinal nerves, especially the facial, glosso-pharyngeal, pneumogastric, spinal accessory and hypoglossal. The disease is marked by difficulty in mastication, in deglutition, in respiration, in phonation and articulation, and by wasting of the

muscles concerned in these functions. The affected muscles display spontaneous fibrillary contractions. Speech becomes pro-

Fig. 45.



Progressive Muscular Atrophy. (Gowers.)

1. Wasting of the muscles of the back and arms.
2. Wasting of the trapezii and deltoids.
3. Wasting of the muscles of the neck; A, habitual posture of the head; B, position into which the head falls if it be not inclined backward.

gressively more difficult, indistinct, and finally nasal. Swallowing is difficult and fluids regurgitate through the nose. Saliva

dribbles from the mouth. Food may find its way into the larynx. Reflex action in the throat may be lost. Rarely the ocular and the facial muscles participate in the palsy.

How is progressive muscular atrophy to be distinguished from acute anterior poliomyelitis?

Progressive muscular atrophy is especially a disease of adult life. It is usually gradual in onset and progressive in course, while acute anterior poliomyelitis is especially a disease of childhood, and is acute in onset and retrogressive in course. Febrile symptoms attend acute poliomyelitis, but not progressive atrophy. The fibrillary twitching of the latter is not present in the former. Terminal bulbar symptoms attend progressive muscular atrophy, but not acute poliomyelitis.

How are progressive muscular atrophy and progressive neural muscular atrophy to be differentiated?

The condition that has been described as the peroneal type of progressive muscular atrophy is dependent upon neuritis and, in addition to weakness, wasting, degenerative electric reactions and deformities, presents sensory disturbances. Besides, it usually sets in early in life and attacks first the feet and the legs, sometimes the hands and the forearms, gradually extending. Several members of the same family are sometimes attacked.

Acute Bulbar Palsy.

What is acute bulbar palsy?

Symptoms almost identical with those of progressive glosso-labio-laryngeal paralysis may set in acutely. They may subsequently undergo some improvement and then remain stationary. The condition is dependent upon an acute lesion—softening or hemorrhage or inflammation of the medulla, involving the nuclei of the glosso-pharyngeal, pneumogastric, spinal accessory and hypoglossal nerves or the fibers passing to or from them.

How is acute to be distinguished from chronic bulbar palsy?

The symptoms of chronic bulbar palsy are gradual in onset, progressive in course and symmetrical in distribution; those

of acute bulbar palsy are sudden in onset, regressive or stationary and are apt to display slight irregularities of distribution.

Pseudo-Bulbar Palsy.

What is pseudo-bulbar palsy?

A bilateral lesion, such as softening or hemorrhage, involving the lower portion of the ascending frontal convolution of both hemispheres, or the corresponding motor tracts in symmetrical situations, will give rise to the symptoms of bulbar palsy: paralysis of the lips, tongue and throat.

How is pseudo-bulbar palsy to be distinguished from true bulbar palsy?

Pseudo-bulbar palsy, in contradistinction to true bulbar palsy, is likely to be characterized by two distinct attacks, by slight asymmetry of distribution of the symptoms, and by the absence of wasting.

Asthenic Bulbar Paralysis.

What is asthenic bulbar paralysis?

Asthenic bulbar paralysis is a disorder of unknown, though possibly toxic origin, characterized by weakness of voluntary muscles, and especially those controlled by the bulbar nerves, or by undue readiness of fatigue after ordinary functional activity, without wasting or changes in reflexes or in sensibility. Remissions and exacerbations are common and often abrupt. The affected muscles are readily exhausted by tetanizing electric currents.

Progressive Muscular Dystrophy.

What are the symptoms of progressive muscular dystrophy?

Progressive muscular dystrophy is a congenital or family-affection, transmitted through females, but occurring more commonly in males, usually appearing in childhood, and characterized by weakness and wasting of muscles, the fibers of which may first undergo hypertrophy and then atrophy, while the interstitial and fatty tissue increases. Several varieties have been described

—*idiopathic, pseudo-hypertrophic, juvenile* or *scapulo-humeral, infantile* or *facio-scapulo-humeral, hereditary*—in accordance with

Fig. 46.



Progressive muscular dystrophy of pseudo-hypertrophic type in two brothers. (Gowers.)

the distribution, the time of appearance and the presence or absence of primary enlargement.

The disease usually makes its appearance early in life and is progressive in course.

The gait has a peculiar oscillating character; ascending stairs is difficult, and the manner of rising from the floor is characteristic (Fig. 47): the patient first gets on his hands and knees; then extending the legs he stands upon his feet; finally, by supporting his hands upon his thighs he manages to reach the erect posture. Ultimately the muscles become reduced in size; the electric reactions suffer quantitative changes, indica-

Fig. 47.



Progressive muscular dystrophy. Mode of rising from the ground. (Gowers.)

tive of diminished excitability; the deep reflexes are enfeebled and lost. Talipes equinus, spinal curvature and muscular contractions occur at a late stage of the disease. Intelligence does not suffer, and sensibility and the function of the sphincters remain unimpaired.

Arthritic Muscular Atrophy.

What is arthritic muscular atrophy

It has been observed that in the course of inflammation of a joint, acute or chronic, spontaneous or traumatic, the related muscles, especially the extensors, undergo a varying degree of atrophy. With the disappearance of the arthritis, the affected muscles gradually return to their normal condition. If the joint-disorder continue for a long period, the deep reflexes in the region involved are heightened and the muscles present quantitative electrical changes; contracture of the opponent muscles may also occur.

Thomsen's Disease—Myotonia Congenita.

What are the symptoms of myotonia congenita—Thomsen's disease?

The disorder described by Thomsen is one that occurs as a congenital affection in families. It is characterized by tonic muscular spasm on voluntary movement following a period of rest. If movement is persevered in, the spasm relaxes. The spasm is intensified by emotion or by the fear of its occurrence. Muscular hypertrophy is the ultimate result. A peculiar electric reaction, *myotonia electrica*, is developed, the muscular contractions induced attaining their maximum slowly and subsequently relaxing slowly.

Posterior Spinal Sclerosis—Locomotor Ataxia.

What are the causes of posterior spinal sclerosis?

Posterior spinal sclerosis, *locomotor ataxia*, or *tabes dorsalis* is more common in males than in females, and in middle adult life than at any other period. Its most common cause is syphilis; other causes are exposure to cold and wet, concussion, overexertion and sexual and alcoholic excesses; the disease has been observed to follow acute infectious diseases; it is sometimes secondary to other forms of spinal disease; in some cases an hereditary neurotic influence can be traced. It has been attributed to metallic poisoning. The etiology is sometimes obscure.

What are the symptoms of posterior spinal sclerosis?

Posterior spinal sclerosis is characterized by impairment of coördination, giving rise to difficulty in performing delicate movements, to unsteadiness of gait and of station, particularly when the guidance of vision is removed; by shooting pains in the extremities; by a girdle-sensation; by other abnormalities of sensation, particularly anesthesia, and especially in the soles of the feet; by abolition of the deep reflexes; by primary exaggeration and secondary loss of the superficial reflexes; by primary increase of the sexual propensity and secondary impairment of the sexual power; by derangement of the sphincters, manifested by retention or incontinence; by narrow pupils that

act in accommodation, but do not respond to the stimulus of light; by paralysis of ocular muscles, giving rise to strabismus and diplopia; by atrophy of the optic and auditory nerves, occasioning loss of vision and of hearing; by laryngeal, gastric, intestinal and other visceral crises, manifested by paroxysms of intense distress; by trophic changes in the joints, giving rise to enlargement and subluxation, rendering the bones brittle and liable to spontaneous fracture; by perforating ulcers of the foot; sometimes by peripheral neuritis; by hyperextensibility at the joints, particularly the knee and the hip; and in the last stages by muscular wasting.

How is multiple neuritis to be distinguished from posterior spinal sclerosis?

If abolition of the reflexes, unsteady gait and station, loss of sensation and sharp pains attend neuritis, they may recede and ultimately disappear; once present in the course of posterior spinal sclerosis they persist. Posterior sclerosis is progressive in course, and unyielding in treatment. With appropriate treatment neuritis is retrogressive. The girdle-sensation of sclerosis is wanting in neuritis. In neuritis, the muscles undergo degeneration and waste; there is palsy; the electric reactions are qualitatively changed. In sclerosis, muscular wasting and weakness result only after long-continued inactivity; if the electric reactions undergo any change, it is quantitative. In neuritis there are pains and points of special tenderness in the course and in the peripheral distribution of various nerves; in posterior sclerosis the muscles are not tender. In walking the foot is raised high in posterior sclerosis and brought down on the heel or the sole; in neuritis foot-drop necessitates raising the foot only so high as to avoid scraping the floor.

How are lumbar pachymeningitis and posterior spinal sclerosis to be differentiated?

Pachymeningitis in the lumbar region may be attended with shooting pains in the thighs and with abolition of the knee-jerks, but not with manifestations indicative of involvement of cerebral nerves. In meningitis the muscles waste early; in posterior sclerosis, not at all or only late. Impairment of co-ordination characterizes posterior sclerosis.

What are the distinctions between posterior spinal sclerosis and cerebellar tumor?

A tumor in or compressing the middle lobe of the cerebellum may give rise to unstable equilibrium and to a staggering gait; it gives rise in addition to occipital headache, to vertigo, vomiting, nystagmus, optic neuritis and to other evidences of compression; while lightning-pains and alterations of sensation are wanting.

Primary Lateral Sclerosis—Spastic Paraplegia.

What are the causes of primary lateral sclerosis?

Primary lateral sclerosis or *spastic paraplegia* occurs with equal frequency in both sexes, and a little earlier in life than posterior sclerosis.

In many cases, no etiologic element can be discovered; in others, there is a history of concussion, of exposure to cold and wet, of syphilis, of excesses or of neurotic heredity.

What are the symptoms of primary lateral sclerosis?

Primary lateral sclerosis, or spastic paraplegia, is characterized by muscular weakness and spasm; the latter usually involves the extensors of the lower extremities occasioning a peculiar spastic gait and so-called "clasp-knife" rigidity; attacks of "spinal epilepsy" occur; the reflexes are exaggerated; ankle-clonus is present. There is usually muscular wasting, occasionally hypertrophy. The arms are affected less commonly and in less degree than the legs. There may be abnormal sensations, but no anesthesia. The sphincters may be involved.

The electric reactions undergo quantitative changes.

How is lateral sclerosis to be distinguished from myelitis?

While lateral sclerosis presents the symptoms of degeneration of the lateral columns, as seen in myelitis, it is always gradual in onset and unattended with febrile manifestations, with girdle-pain, or with impairment of sensation. Typical paraplegia, with muscular wasting and degeneration and impairment of the sphincters is less characteristic of lateral sclerosis than of myelitis.

How may lateral sclerosis simulate and be distinguished from cerebral hemiplegia?

The manifestations of primary lateral sclerosis may be unilateral; but the face is not involved, as it usually is in hemiplegia of cerebral origin; and careful examination will detect exaggeration of the reflexes of the upper as well as of the lower extremity on the apparently uninvolved side, unlike what is found in cerebral hemiplegia.

Postero-lateral Sclerosis—Ataxic Paraplegia.

What are the causes of ataxic paraplegia or postero-lateral sclerosis?

Postero-lateral sclerosis, or *ataxic paraplegia*, is more common in males than in females and in middle adult life than at any other period. Its etiology is often obscure. In some cases a neurotic heredity can be traced; in others there has been exposure to cold and wet; in still others concussion seems to have been the cause; a history of syphilis is uncommon.

What are the symptoms of postero-lateral sclerosis, or ataxic paraplegia?

The symptoms of ataxic paraplegia are dependent upon sclerosis of the posterior and lateral columns of the spinal cord. The disease is gradual in onset and marked by muscular weakness, spasm and incoördination. There may be pain in the sacral region, but lightning-pains are absent. Articulation may be defective. Tremor of the muscles of the face may occur. The deep reflexes are exaggerated. The iris usually reacts to light. Atrophy of the optic nerve is rare. Sexual power may be lost. The sphincters become enfeebled. Ultimately, contractures develop.

How is posterior spinal sclerosis to be distinguished from postero-lateral sclerosis?

POSTERIOR SCLEROSIS.	POSTERO-LATERAL SCLEROSIS.
Syphilis a common cause.	Syphilis a rare cause.
Weakness a late symptom.	Weakness an early symptom.
Knee-jerk lost early.	Knee-jerk exaggerated.

POSTERIOR SCLEROSIS.	POSTERO-LATERAL SCLEROSIS.
No ankle-clonus.	Ankle-clonus.
Never muscular spasm.	Characteristic muscular spasm.
Argyll-Robertson pupil.	Pupil responds to light.
Lightning-pains.	Dull sacral pains.
Sensory impairment.	Sensation unimpaired.
Girdle-sensation.	No girdle-sensation.
Optic-nerve atrophy common.	Optic-nerve atrophy rare.
Visceral crises.	No crises.

How is postero-lateral sclerosis to be distinguished from primary lateral sclerosis?

Ataxic paraplegia is to be distinguished from spastic paraplegia by the presence of symptoms of muscular incoördination, dependent upon involvement of the posterior columns of the cord.

What are the distinctions between a tumor of the cerebellum and postero-lateral sclerosis?

A growth involving the middle lobe of the cerebellum may give rise to weakness, ataxia, spasm, and heightened reflexes; there are, besides, occipital headache, optic neuritis, vertigo, vomiting and nystagmus; other evidences of pressure may be present.

Friedreich's Ataxia—Hereditary Ataxic Paraplegia.

What are the symptoms of Friedreich's ataxia?

The symptoms of *Friedreich's ataxia* are dependent upon sclerosis of the lateral and posterior columns of the spinal cord. The disease occurs in families and appears early in life; it attacks both sexes alike. It is attended with an ataxic gait, impairment of coördination and equilibration, muscular weakness, abolition of knee-jerks, early talipes equinus, lateral curvature of the spine, nystagmus and defective speech; sensation is usually unimpaired.

What are the distinctions between Friedreich's ataxia and ataxic paraplegia?

The early period of life at which the symptoms appear, its occurrence in families, the absence of knee-jerks and of ankle-

clonus and the presence of nystagmus distinguish Friedreich's disease from ataxic paraplegia.

Cerebro-Spinal Sclerosis—Insular Sclerosis.

What are the causes of cerebro-spinal sclerosis?

The more commonly-recognized causes of *disseminated, multiple, insular*, or *cerebro-spinal sclerosis* are exposure to cold and wet, traumatism, nervous shock and acute febrile diseases. Occasionally a neuropathic heredity can be traced.

What are the symptoms of cerebro-spinal sclerosis?

The symptoms of disseminated, multiple, insular, or cerebro-spinal sclerosis vary with the distribution of the islets of sclerosis. As a rule, there is a peculiar jerky incoördination or tremor, most marked in the upper extremities, and sometimes involving the head and the tongue, aggravated by effort, emotion or observation; there is commonly nystagmus, usually lateral, sometimes vertical, sometimes rotatory; speech is often slow, scanning, syllabic, parts of words being dropped; the reflexes are usually exaggerated. In addition there are certain mental changes, often manifested by a sense of complacency, contentment and self-satisfaction, quite at variance with the patient's condition. Evidences of muscular weakness are not rarely present. There may also be headache, vertigo and optic neuritis.

How is cerebro-spinal sclerosis to be distinguished from postero-lateral sclerosis?

There may sometimes be considerable difficulty in differentiating cerebro-spinal sclerosis and postero-lateral sclerosis (ataxic paraplegia) and Friederich's (hereditary) ataxia. In postero-lateral sclerosis, the defect of coördination involves the lower extremities primarily; in multiple sclerosis, there is coarse tremor rather than true incoördination, and the upper extremities are especially involved. Nystagmus occurs in Friedreich's ataxia and in multiple sclerosis, but not in ataxic paraplegia. The knee-jerks are exaggerated in multiple sclerosis and in ataxic paraplegia, but are wanting in Friedreich's ataxia. Friedreich's ataxia is a family-disease and appears early in life. Multiple

sclerosis and ataxic paraplegia appear in middle life. Cerebral symptoms and mental phenomena belong to multiple sclerosis rather than to postero-lateral sclerosis. Speech is slow and scanning in cerebro-spinal sclerosis; in Friedreich's ataxia there are merely elision and occasional separation of syllables.

Paralysis Agitans—Shaking Palsy.

What conditions favor the development of paralysis agitans?

Paralysis agitans or *shaking palsy* begins most commonly at about fifty years of age. It is more common in males than in females. In some cases, an hereditary influence can be traced in an etiologic connection. In others, the onset of the disease has been preceded by decided emotional disturbance, by physical injury or by acute disease.

What are the symptoms of paralysis agitans?

Paralysis agitans, shaking palsy or Parkinson's disease is characterized by tremor, weakness and rigidity. Tremor is usually first observed; weakness and rigidity subsequently. The manifestations first appear in one extremity, and gradually extend to the others. The tremor is fine and rhythmic; it is increased by observation and by emotional disturbance, and diminished or restrained temporarily by active and by passive movement, and sometimes spontaneously. As a rule, the head does not participate in the tremor; exceptionally it does. The expression of the face is fixed and immobile. The shoulders are bent forward, giving rise to the phenomena of propulsion—a tendency to run forward. Occasionally there is retropulsion, or there may be a tendency to lateral movement. The hands assume a peculiar, semi-flexed attitude, the fingers performing movements as if rolling a small object between their tips; or the fingers are held as if grasping a pen. The voice is monotonous; words are uttered rapidly, with a tendency to confluence of syllables. The reflexes are usually unaltered; exceptionally the knee-jerks are exaggerated and ankle-clonus can be elicited. A sensation of abnormal heat, sometimes with perspiration, is often present.

What are the differential features between paralysis agitans and cerebro-spinal sclerosis?

The diagnosis between these two conditions, ordinarily simple, may under some circumstances be extremely difficult. The tremor of cerebro-spinal sclerosis is coarse and irregular, and is

FIG. 48.



Paralysis agitans. (After St. Leger.)

induced and aggravated by voluntary effort, by emotion and by observation; that of paralysis agitans is fine and regular, and is constant, at least during the waking hours. In cerebro-spinal sclerosis, the head participates in the movements; in paralysis agitans, on the contrary, the face is fixed, immobile, expressionless. Paralysis agitans does not present the slow, scanning speech and nystagmus of multiple sclerosis; while the latter does not present the tendency to forward or backward or even lateral movement of the former. The characteristic attitude of the hand, as if holding a pen in writing, or as if rolling pills between the fingers, seen in paralysis agitans, is not seen in multiple sclerosis.

Spinal Hemorrhage.

What are the causes of hemorrhage into the spinal cord?

Spinal hemorrhage is more common in males than in females, and in adult life than at any other period. It may be *primary*, dependent upon disease of the bloodvessels, or as a result of exposure to cold, or of over-exertion, or of sexual excess; *secondary*, in the course of inflammation, tumors and cavities in the cord; *accessory*, occurring towards the close of convulsive disorders; and *traumatic*, following blows, falls and other injuries.

What are the symptoms of hemorrhage into the spinal cord?

The occurrence of hemorrhage into the spinal cord is indicated by sudden, severe pain in the back, with loss of motion and sensation. Consciousness may be lost, but is likely to be preserved. A girdle-sensation exists at the level of the lesion, and loss of motion and sensation below, on alternate sides, or on both sides of the body, according to the seat of the lesion. The sphincters are likely to be deranged and trophic changes to occur. Respiration will be interfered with if the lesion is in the cervical or dorsal region of the cord.

Some degree of myelitis and meningitis are usually developed in the progress of the case, giving rise to fever and irritative symptoms. Subsequently, the symptoms become paralytic in type. Some degree of paraplegia remains permanently. The paralyzed muscles waste. The deep reflexes are exaggerated.

What are the distinctions between spinal hemorrhage and spinal meningitis?

The abruptness of onset is more decided in hemorrhage than in meningitis. Febrile manifestations attend meningitis from the beginning; they only appear in hemorrhage when myelitis is established. Motor and sensory impairment is more decided in hemorrhage than in meningitis; in the latter, there is a preliminary stage of spasm and pain.

How are hemorrhage into the cord and hemorrhage into the meninges to be distinguished from one another?

Both the local and the radiating pains are less severe in spinal than in meningeal hemorrhage, while the subsequent

anesthesia is more decided in the former than in the latter. In the case of spinal hemorrhage the paralytic symptoms are more decided from the outset; while in meningeal hemorrhage these are preceded by muscular spasm. Trophic changes characterize spinal hemorrhage, and are wanting in meningeal hemorrhage.

Spinal Compression.

What are the causes of compression of the spinal cord?

The cord may be compressed by tumors of the spinal canal; by dislocation of the vertebræ, independently, or as a result of caries, or of fracture; by an exostosis; by an aneurism that has eroded the bones; or by an hydatid cyst.

How can the causes of spinal compression be differentiated?

The recognition of a tumor in the spinal canal depends upon its detection from without, or the detection of new-growths in other parts of the body.

Simple dislocation and fracture of the vertebræ follow traumatism; the symptoms to which dislocation gives rise set in suddenly and are profound in degree; a deformity of the spinal column may be detectable.

The diagnosis of caries depends upon the knowledge of a history of syphilis or of tuberculosis and the detection of a painful deformity in the back.

An hydatid cyst of the spinal canal may be diagnosticated by the detection externally of a fluctuating tumor, upon puncture of which the characteristic hooklets may be found.

It may be impossible to diagnosticate the cause of compression of the cord dependent upon an exostosis or upon an aneurism.

To what symptoms does compression of the spinal cord give rise?

The rapidity with which the symptoms of compression appear depends somewhat upon the cause. Dislocation is apt to occasion manifestations of immediate gravity. In other cases, the symptoms are gradual in appearance and progressive in course.

Compression gives rise to two groups of phenomena, referable to the nerve-roots and to the cord, respectively. There is local pain in the back, aggravated by movement, as well as pains of a radiating character, with girdle-sensation; ultimately anesthesia develops. There is loss of motion in the parts supplied by the nerves from the cord below the seat of compression; with exaggerated reflexes and involvement of the sphincters. The palsied muscles slowly waste and degenerative reactions set in. Contractures may develop.

How is compression-myelitis to be distinguished from hemorrhage into the spinal cord?

When a vertebra softened by destructive disease suddenly gives way, the resulting displacement may be followed by compression of the cord, occasioning symptoms with which those produced by hemorrhage into the cord may be identical. The symptoms of compression, however, are usually more extensive and more absolute than those of hemorrhage; the existence of a deformity of the spine makes the diagnosis certain.

What are the distinctions between compression of the cord and chronic myelitis?

Symptoms of irritation referable to the nerve-roots are wanting in myelitis. The recognition of a cause of compression determines the diagnosis.

Tumor of the Spinal Cord.

What are the symptoms of tumor of the spinal cord?

Tumors of the spinal canal may be situated without or within the dura mater, or within the structure of the cord itself. They are most diverse in character. The symptoms will depend upon the situation of the tumor and upon the degree of mechanical interference with the functions of the cord that it occasions. Pressure on the nerve-roots gives rise to pain, to abnormal sensations, to girdle-pain and to muscular spasm and rigidity. Pressure on the cord or myelitis occasions anesthesia, paralysis and exaggerated reflexes below the level of the lesion and

abolition of the reflexes within the area innervated from the seat of the growth. Involvement of the anterior horns of the gray matter is attended with wasting and other trophic disorders; involvement of the lumbar enlargement causes abolition of the knee-jerk, loss of control of the sphincters and wasting and palsy in the lower extremities; involvement of the cervical enlargement occasions wasting and palsy in the upper extremities. The unilateral appearance of symptoms of spinal disease, or evidence of sensory derangement on one side and motor derangement on the other, is strongly suggestive of tumor of the spinal canal. With the growth of the tumor the symptoms become bilateral.

From what conditions is spinal tumor to be distinguished?

The diagnosis of spinal tumor includes the determination of the nature of the tumor and its distinction from other conditions presenting similar symptoms.

In the first connection, a history of syphilis, of tuberculosis, or of tumors situated elsewhere is to be considered.

The differentiation from caries of the vertebræ depends upon the recognition of disease of the bone, upon the deformity that results and upon the greater degree of pain on movement that attends caries.

From hypertrophic pachymeningitis spinal tumor differs in course; being more rapidly progressive, while symptoms of irritation precede those of paralysis and wasting. The symptoms of pachymeningitis are bilateral from the outset and more circumscribed in distribution than are those of tumor.

The symptoms of tumor differ from those of myelitis, in being irritative in character—attended by pain and spasm, rather than paralytic. If tumor give rise to myelitis, the recognition of the condition depends upon a knowledge of the previous symptoms.

The persistence of obstinate neuralgia, especially if bilateral, should excite suspicion of spinal tumor.

Syringomyelia.

What are the symptoms of syringomyelia?

Cavities in the spinal cord may be a result of defective apposition of the lateral halves of the cord in the course of development; of occlusion of the central canal, with accumulative distention by cerebro-spinal fluid; of the disintegration of gliomata; or they may appear subsequently to myelitis. The course of the disease is slow and chronic, sometimes covering many years. The symptoms vary somewhat with the situation of the cavity. They are usually most conspicuous in the upper extremities. They may be largely bulbar. There are muscular wasting and weakness, preceded by alterations of sensation. Common sensibility is usually preserved, while the perception of pain and of heat and cold is enfeebled or lost. In some cases severe pains occur. A spastic condition may be present in the lower extremities. The sphincters may escape or be involved. Trophic changes are common, and arthropathies are occasionally observed. There may be cutaneous eruptions, as eczema or herpes. The skin may be thin and glossy or thick and horny. Undue sweating may take place. Vaso-motor disturbance may be manifested by coldness and lividity.

From what conditions is syringomyelia to be differentiated?

Myelitis, hypertrophic pachymeningitis and progressive muscular atrophy occasion certain symptoms in common with syringomyelia.

Myelitis is recognized by the much more profound palsy and loss of sensory power, without involvement of the pain-sense and the temperature-sense; hypertrophic pachymeningitis by the attendant pain and the less extensive anesthesia; and chronic muscular atrophy by the absence of conspicuous sensory symptoms.

How are syringomyelia and leprosy to be differentiated?

Many of the symptoms of leprosy are dependent upon peripheral neuritis, so that nerves may be swollen and tender, and all forms of sensibility suffer equally. Spastic symptoms are wanting and the sphincters are not deranged. The detection

of lepra-bacilli in the serum of blisters or in the discharges would remove any doubt in diagnosis.

Morvan's Disease—Analgesic Panaris.

What are the symptoms of Morvan's disease?

Under the name of *Morvan's disease* has been described a syndrome of symptoms, including the development of a painless inflammation at the extremities of the fingers, followed by necrotic sequestration of the phalanges. At the beginning of the disease, the affected parts may be the seat of pain. Subsequently, analgesia develops, together with the destructive process in the fingers. In most cases abnormal curvature of the spinal column has been observed. After death, hyperplasia of the connective tissue of the peripheral nerves and in the posterior horns, posterior columns, and the gray matter of the cervical segment of the spinal cord has been found. The disorder is considered a variety of syringomyelia.

What is the distinction between Morvan's disease and scleroderma?

Scleroderma is a morbid condition in which, as a result of inflammatory changes in the subcutaneous arteries of a varying distribution, hyperplasia of the connective tissue takes place, with hardness, swelling and fulness, which in turn is succeeded by contraction and atrophy. The condition may be circumscribed or diffuse. It is sometimes attended with pigmentation and sometimes with desquamation. The etiology of the affection is obscure.

A condition in which the extremities of the fingers close to the nails undergo discoloration, followed by the formation of bullæ, with perhaps loss of the nails and shortening of the fingers, has been described as a variety of scleroderma or *sclerodactyly*. When nose and ears and face, as well as hands and feet, became reduced in size, the name *akromikria* has been employed. In scleroderma there may be anesthesia, but there is not analgesia. Trophic changes may take place in the affected parts, but they are not of a necrotic character. The nails may be lost, but there is no exfoliation of bone.

How are Morvan's disease and anesthetic leprosy to be differentiated?

In anesthetic leprosy, a destructive process may be set up in the extremities, attended sometimes with the loss of fingers or toes, but the condition is usually to be recognized by the association of other definite symptoms of leprosy, such as patches of anesthesia and leprous nodules in various parts of the body.

FIG. 49.



Appearance of the hand in Morvan's Disease. (After Charcot.)

From what other conditions is Morvan's disease to be differentiated?

Raynaud's disease is attended with cyanosis of the extremities, sometimes terminating in gangrene; but sensibility remains unimpaired; and there are other characteristic symptoms.

Syphilis may occasion a destructive dactylitis. The diagnosis will depend upon the absence of sensory symptoms and upon

a knowledge or a history of infection or of other manifestations.

Necrosis of bone may attend *diabetes*, but sensory symptoms are wanting, distinctive symptoms are present and examination of the urine will make clear the clinical association.

Cerebral Meningitis.

What are the varieties of inflammation of the membranes of the brain?

Inflammation of the dura, pia or arachnoid is known as *pachymeningitis*, *leptomeningitis* or *arachnitis*, respectively; most commonly the pia and arachnoid are involved together—*pia-arachnitis* or *leptomeningitis*. Inflammation of the dura mater may be attended with the extravasation of a membranous hemorrhagic exudate—*hemorrhagic pachymeningitis*. Meningitis may especially involve the *convexity* or the *base*; it may be *acute* or *chronic*, *primary* or *secondary*; it may be *simple*, *purulent*, *tuberculous*, *syphilitic*.

What are the causes of cerebral meningitis?

Inflammation of the pia and arachnoid may be a result of traumatism or of contiguous inflammation; it may develop in the course of infectious disease or of pyemia; it may depend upon a deposit of tubercles in the membranes.

What are the symptoms of cerebral meningitis?

The symptoms of meningitis vary somewhat, as the base or the convexity of the brain is involved. Tuberculous meningitis is usually basilar. Meningitis dependent upon causes other than tubercles, usually, though not exclusively, affects the convexity. Certain symptoms are common to meningitis in any situation. In addition to those of the associated condition, there are, together with febrile manifestations, headache, delirium, vomiting, convulsions, retraction of the head, derangement of the sphincters, palsies, vaso-motor and trophic disturbances, inequality of the pupils and irregularity of the pulse and respiration. When the base is involved, there are optic neuritis followed by atrophy, irritation followed by paralysis of cranial nerves and more decided alteration of pulse and respiration,

trophic and vaso-motor disturbances and retraction of the head. When the inflammation is tuberculous, tubercles may, on ophthalmoscopic examination, be detected in the choroid.

What are the symptoms of cerebral pachymeningitis?

Inflammation of the dura mater may be a result of traumatism or of adjacent disease. In addition to the symptoms of the primary condition, there may be headache, delirium, convulsions, febrile manifestations. A collection of pus may occasion paralysis.

What are the symptoms of hemorrhagic pachymeningitis?

Hemorrhagic pachymeningitis is more common in males than in females, and late in life than at any other period. It is usually preceded by a history of alcoholism, of insolation, of traumatism, or of insanity; it may also develop in the course of blood-diseases. It is attended with headache, contracted pupils, intellectual torpor, apoplectiform convulsions, unnatural drowsiness and mental wandering. The condition is rather obscure, and the symptoms not well defined. It is often unexpectedly found *post mortem*.

How is simple cerebral meningitis to be distinguished from tuberculous meningitis?

Ordinary cerebral meningitis sets in rather acutely; tuberculous meningitis usually more insidiously. The ordinary form of inflammation is associated with well-known causes: traumatism, adjacent inflammation, infectious diseases and pyemia; tuberculous meningitis is usually secondary to tuberculous processes in other parts of the body. The symptoms of the former especially indicate involvement of the convexity and freedom of the base, while in the latter the conditions are reversed, and choked disc and paralysis of cranial muscles are more common. Tubercles are never found in the choroid in ordinary meningitis. Simple meningitis is not uncommonly attended with recovery; tuberculous meningitis but rarely.

How are the cerebral symptoms of acute general disease to be distinguished from the symptoms of cerebral meningitis?

Cerebral symptoms may appear at any time in the course of

acute general diseases, such as scarlatina, smallpox, yellow fever, typhoid fever, rheumatism, pneumonia and acute miliary tuberculosis. Investigation should be directed to the determination of the association, so that the primary condition may not be obscured. Sufficient cause for the symptoms may be found in the high temperature, in the state of inanition, or in the poisoned condition of the blood. When the delirium of febrile affections appears, the headache that has been present disappears. Headache and delirium continue together in meningitis. The manifestations of irritation and paralysis of cranial nerves, including optic neuritis and atrophy, seen in meningitis, are wanting in simple febrile disorders.

How is cerebral meningitis to be distinguished from cerebro-spinal fever?

The differences are of degree, extent and etiology. When the membranes of the brain are inflamed, the spinal membranes are usually in some degree also involved. Cerebro-spinal fever is considered a constitutional affection, cerebral meningitis a local affection. In the former, however, the spinal involvement stands boldly out, as evidenced by the tendency to opisthotonos, the symptoms on the part of the trunk, the sphincters, and the lower extremities; while the latter is especially marked by the involvement of cerebral structures. Cutaneous eruptions mark cerebro-spinal fever, but are no part of cerebral meningitis. The recognition of a cause of cerebral meningitis, or a knowledge of the existence of other cases of cerebro-spinal fever may determine the diagnosis. The detection of the specific bacteria of cerebro-spinal fever in the fluid obtained by puncture of the membranes in the lumbar region affords conclusive evidence.

How is cerebral meningitis to be distinguished from acute mania?

The mental wandering of meningitis does not usually attain the high degree seen in mania; and in the former it is attended with febrile manifestations, in the latter not. The symptoms on the part of the cranial nerves, the headache, the convulsions of meningitis are wanting in mania.

Hydrocephalus.

What are the varieties of hydrocephalus?

An excessive accumulation of cerebro-spinal fluid may take place in the subdural space or within the ventricles of the brain. The one condition is called *external;* the other *internal hydrocephalus.* Either may be *acute* or *chronic*, *primary* or *secondary*, *congenital* or *acquired.*

What are the causes of hydrocephalus?

Hydrocephalus may be a result of meningitis, of obstruction of the orifices of communication between the ventricles and between the fourth ventricle and the subarachnoid space, or of pressure on the veins of Galen. Occasionally no causative factor is recognizable.

To what symptoms does hydrocephalus give rise?

Occurring in a child prior to the closure of the sutures and the complete ossification of the bones, hydrocephalus causes a separation and a thinning of the cranial bones. When the fluid distends the ventricles, the surrounding cerebral tissue becomes attenuated. Under such conditions the head is enlarged; the fontanels may be open; the eye-balls roll; the mental condition is defective; there is muscular weakness; there may be convulsions; and various cranial nerves may be paralyzed.

In the more aggravated cases idiocy exists; there is blindness, with choked discs; and life may be terminated by convulsions and coma.

Analogous symptoms also occur when hydrocephalus develops later in life.

Hemorrhage into the Cerebral Membranes.

What are the varieties of hemorrhage in the cerebral meninges?

The extravasation of blood without the dura mater, between the dura and the bone, constitutes *extradural* hemorrhage; hemorrhage within the dura mater, between the dura and arach-

noid is called *subdural;* and between the arachnoid and pia mater, *subarachnoid.*

Under what conditions does hemorrhage into the meninges of the brain occur?

Hemorrhage into the cerebral membranes may be a result of traumatism ; of the rupture of an aneurism of a meningeal vessel ; of the rupture of a hemorrhage in the brain ; of the same conditions as occasion cerebral hemorrhage ; it may occur in the insane ; rarely it occurs spontaneously.

To what symptoms does hemorrhage into the cerebral membranes give rise?

The symptoms of hemorrhage into the cerebral membranes, depend upon the volume and the extent of the extravasation. The symptoms are those of meningitis, plus those of cerebral hemorrhage, and may be preceded by headache, vertigo and vomiting.

The infantile meningeal hemorrhages that result during labor occasion the so-called birth-palsies, with the symptoms of congenital spastic paraplegia or hemiplegia.

Congenital Spastic Paraplegia.

What is infantile or congenital spastic paraplegia?

When, as a result of injury to the child during birth, bilateral meningeal hemorrhage takes place over the central convolutions, the pyramidal tracts fail to develop or they degenerate. The child is late in learning to walk. It presents the symptoms of spastic paraplegia. The gait is peculiar, one foot being placed over or in front of the other, or a swinging oscillation taking place. Growth and development are retarded. Athetoid movements in the hands are common. Initial convulsions may occur and be repeated, subsequently to cease. Should the hemorrhage be circumscribed or unilateral, the distribution of the symptoms will vary accordingly. There is usually defective mental development.

What are the distinctions between congenital spastic paraplegia and pseudo-hypertrophic paralysis?

The reflexes are exaggerated in congenital spastic paraplegia, enfeebled or lost in pseudo-hypertrophic paralysis. The gait of congenital spastic paraplegia is spastic; that of pseudo-hyper-

FIG. 50.



Case of congenital spastic diplegia. (Philadelphia Hospital.)

trophic paralysis is oscillating; the manner in which the child rises from the floor in the latter is characteristic. In congenital spastic paraplegia the contractures are yielding; in pseudo-hypertrophic paralysis unyielding; the former is retrogressive, the latter progressive. Should there be decided enlargement of some of the muscles, this will constitute a diagnostic feature.

Cerebral Anemia.

What are the conditions that lead to cerebral anemia?

Anemia of the brain may be general: as a part of a systemic anemia, from cardiac insufficiency, from accumulation of blood elsewhere, from pressure on the large vessels to the head, from pressure on the brain; or partial: from obstruction of the circulation by vascular occlusion or by pressure from without. Cerebral anemia may develop gradually or suddenly.

To what symptoms does cerebral anemia give rise?

General anemia of the brain, suddenly induced, occasions the

symptoms of syncope: failure of vision, ringing in the ears, vertigo, nausea, shallow, sighing respiration, contracted pupils, pallor of the face, a cool, moist skin and loss of consciousness. Nystagmus and convulsions may occur. The loss of consciousness may pass into coma, and coma into death.

General anemia of the brain, gradually induced, occasions enfeeblement with irritability of function. There are headache, vertigo, impaired intellection, motor weakness, drowsiness or insomnia, hallucinations, mania or melancholia. The optic disc may be pale. The symptoms are aggravated by the erect posture and may be mitigated by recumbency or inhalation of amyl nitrite.

Partial anemia of the brain is followed by impaired nutrition and loss of function in the affected area. Occurring suddenly, it is attended with loss of consciousness and convulsions; when of gradual development, it occasions headache, vertigo, numbness, tingling and weakness.

Cerebral Hyperemia.

What conditions lead to cerebral hyperemia?

Hyperemia of the brain may be active or passive. *Active hyperemia* may be caused by an overacting left ventricle; by sudden contraction of the vessels elsewhere; by insolation; it may be part of a general plethoric condition; and it occurs as the first stage of the inflammatory process.

Passive hyperemia is a result of cardiac insufficiency, of venous obstruction as a result of pressure from without or of pulmonary disease.

To what symptoms does cerebral hyperemia give rise?

The symptomatology of cerebral hyperemia is somewhat obscure. The recognition of the condition is not always easy. Usually there are dull headache, a sense of fulness of the head, vertigo, mental torpor, derangement of sleep, a disinclination to activity, flashes of light and tinnitus aurium. The countenance may be flushed; the vessels of the eyeground injected. There may be transient loss of consciousness. Convulsions are uncommon.

Cerebritis.

What are the clinical features of cerebritis?

Inflammation of the structure of the brain may be acute or chronic. It is usually a result of traumatism, or it may arise by extension from adjacent disease. Some degree of cerebritis is coincident with inflammation of the membranes. The symptoms are rather obscure and ill-defined. There are headache, vertigo, delirium, convulsions and febrile symptoms. Cerebritis may be followed by abscess.

How is inflammation of the membranes to be distinguished from inflammation of the structure of the brain?

Inflammation of the membranes and inflammation of the structure of the brain are to some degree always associated, the symptoms of either condition respectively predominating. Inflammation of brain-tissue, however, is usually attended with symptoms more profound and more depressing than are those of meningitis; while the symptoms of meningitis are usually more widely distributed and more irritative in character.

Cerebral Abscess.

What are the causes of abscess of the brain?

The development of cerebral abscesses is usually dependent upon traumatism, suppurative disease of adjacent structures or pyemia. Distant suppuration may give rise to embolic abscess. Inflammation of the brain may terminate in suppuration and the formation of an abscess. Cerebral abscess may be acute or chronic in its course; it may be single or multiple.

What are the symptoms of abscess of the brain?

The symptoms of *abscess of the brain* present themselves in three stages: a *primary acute stage*, in which the symptoms of the associated condition may obscure the cerebral symptoms; a *secondary stage of lull*, in which the symptoms are latent and in abeyance; and a *terminal stage*, in which rupture of the abscess gives rise to the abrupt appearance of the signs of meningitis,

of cerebritis or of apoplexy, followed by stupor, coma and death. These stages are not always distinct. Commonly the one passes by gradations into the other.

The more obvious symptoms are fever, recurrent rigors, headache (which may be localized), vertigo, vomiting and optic neuritis (which are most marked in cerebellar abscess), epileptiform convulsions and paralysis (when the abscess is seated in the motor area). There may be, in addition, unilateral ptosis, defect of speech, impairment of articulation, difficulty of deglutition and mental disturbance.

From what conditions is abscess of the brain to be differentiated?

Abscess of the brain gives rise to many of the symptoms of tumor of the brain. In the terminal stage it occasions the symptoms of meningitis, of cerebritis or of apoplexy. The distinction from these several conditions depends upon the recognition of a cause of abscess, upon the recurrence of chills, upon the presence of febrile symptoms. The headache and optic neuritis of tumor are more intense than those of abscess. In meningitis there is greater involvement of cranial nerves than in abscess. The course of the symptoms of abscess is of longer duration than is the case in cerebritis. Optic neuritis is wanting in apoplexy. The symptoms of the terminal stage of abscess are to be distinguished from meningitis, cerebritis and apoplexy by a knowledge of the previous existence of obscure symptoms of cerebral disease.

Cerebral Hemorrhage.

What are the causes of cerebral hemorrhage?

The ultimate cause of cerebral hemorrhage, not traumatic in origin, is disease of the bloodvessels. The event may be induced by over-action of the heart. In some cases of cerebral hemorrhage, an hereditary tendency can be elicited. Hemorrhage in the brain is most common at the degenerative period of life, and its freqency increases with age; the condition occurs more commonly in males than in females; in temperate than in

tropical climates; in winter than in summer. The antecedent disease of the vessels may depend upon alcoholism, gout, rheumatism, syphilis, nephritis, lead-poisoning, infectious diseases and blood-diseases.

What is the pathology of cerebral hemorrhage?

The symptoms that follow the occurrence of cerebral hemorrhage depend upon laceration of the brain-tissue, and upon the compression and irritation of adjacent structure to which the extravasation gives rise. In the course of time, the blood, as well as the products of disintegration, are absorbed, until there remains to mark the previous hemorrhage nothing but a fibrous cicatrix or a cyst. The conducting motor fibers related to the structures destroyed undergo secondary degeneration.

FIG. 51.



Cerebral miliary aneurisms. (Eichhorst.)

The most frequent seat of cerebral hemorrhage is the region of the central ganglia; next in frequency is the centrum ovale; then the cortex, pons and cerebellum.

What are the symptoms of cerebral hemorrhage?

The occurrence of hemorrhage in the brain is indicated by sudden, complete loss of consciousness, with general muscular relaxation, and depression of temperature. True prodromata are wanting. The breathing is labored and stertorous; an initial convulsion may occur; the pulse is apt to be slow, full and hard; the pupils unequal; the reflexes abolished or unequal;

and the head and eyes deviated to one side (conjugate deviation). Urine and feces may be passed involuntarily. The urine may contain albumin, or even sugar. Vomiting may occur. As the shock of the seizure passes off consciousness slowly returns and the evidences of hemiplegia become apparent. The leg, the arm, perhaps the face, are paralyzed on the side opposite to that on which the lesion is situated. The tongue, protruded, deviates to the paralyzed side. There is difficulty in deglutition, in phonation, in articulation. There may be partial or complete anesthesia of the paralyzed members. Secondary fever appears. Gradual improvement slowly takes place, in greatest degree in those parts that habitually act in association with corresponding parts of the opposite side.

If the hemorrhage occur in the cortex, epileptiform convulsions take place ; if in the posterior part of the third frontal convolution of the left side, aphasia results ; if in the posterior part of the first temporal convolution, word-deafness ; if in the angular gyrus and occipital lobe, hemianopsia and word-blindness ; if in the anterior part of the uncinate convolution, loss of the sense of taste. If the hemorrhage is large and extensive the patient may never emerge from the primary coma, but death may speedily or slowly take place. Implication of the pons or medulla is indicated by abnormal elevation of temperature. In most cases the hemorrhage involves the region of the central ganglia and the internal capsule.

Athetosis, post-hemiplegic chorea and other spasmodic conditions may be sequelæ of cerebral hemorrhage. The reflexes become exaggerated, especially upon the paralyzed side, as a result of descending degeneration of the pyramidal tracts. The memory is impaired. Emotional mobility is increased. Various vasomotor and trophic changes may take place in the affected parts, which may also become fixed by contractures.

In what respects do the symptoms of cerebral hemorrhage differ from those of cerebral congestion ?

Cerebral congestion may be attended with transient loss of consciousness, not so profound, however, as is the case in hemorrhage. The event is prone to be repeated, but the condition does not give rise to permanent paralytic manifestations.

What are the distinctions between the loss of consciousness of syncope and that of cerebral hemorrhage?

Syncope is usually a result of cerebral anemia, induced by insufficiency of the circulation; it may also result from nervous influences that stimulate the vagus. The loss of consciousness is of but brief duration. The pulse is feeble, the face pale, the breathing shallow and sighing. The reflexes are preserved. The sphincters are continent. The return to consciouness is not attended with evidences of paralysis. The attack is prone to be repeated.

How is the coma of cerebral hemorrhage to be distinguished from the coma of alcoholic intoxication?

The symptoms of cerebral hemorrhage come on suddenly; those of alcoholic coma, less suddenly, as a direct consequence of the imbibition of a toxic quantity of alcohol. The coma of alcoholism is not so profound as that of hemorrhage. It may be possible by appropriate stimuli to rouse an individual overcome by alcohol, but there is no response from the coma of hemorrhage. The odor of alcohol in the breath, and the detection of alcohol in the urine would be doubtful evidence; but the absence of these conditions would exclude alcoholism. Conjugate deviation of the head and eyes is a common feature of hemorrhage; alcoholism will not cause it. The pupils are usually dilated in alcoholism, not contracted and unequal as in hemorrhage. The temperature-alterations of hemorrhage are wanting in alcoholic intoxication. On the return of consciousness after alcoholic coma, there is no palsy; palsy is characteristic of hemorrhage.

How is the coma of cerebral hemorrhage to be distinguished from that of opium-poisoning?

In the absence of the previous history, cerebral hemorrhage is to be distinguished from opium-poisoning by the primary depression and secondary elevation of temperature, by the conjugate deviation of the head and eyes, by the evidences of paralysis.

The small pupils of opium-poisoning are equal; if hemorrhage causes contraction of the pupils, one is likely to be smaller than

the other. The noisy breathing of hemorrhage does not become so infrequent as the breathing in opium-poisoning.

How is uremia to be distinguished from cerebral hemorrhage?

The conditions under which uremia develops predispose to the occurrence of cerebral hemorrhage. The recognition of disease of the kidney does not help in diagnosis. The coma of hemorrhage, however, is likely to set in more suddenly and to be more profound than is that of uremia.

Uremia is more likely than hemorrhage to appear in young persons. The convulsions that attend uremia are, as a rule, general; unilateral convulsions are more likely to be indicative of cerebral hemorrhage. The primary depression of temperature is followed by a secondary elevation in hemorrhage, but usually not in uremia. Inequality of the pupils and conjugate deviation of the head and eyes are indicative of hemorrhage rather than of uremia.

If loss of muscular power follow uremia, it is usually circumscribed and transitory; in hemorrhage it is usually hemiplegic and persistent.

In uremia the tongue is dry and covered with a thick brownish fur; there is often vomiting; the body often exhales a peculiar, musty odor, not observed in hemorrhage.

How is asphyxia to be distinguished from cerebral hemorrhage?

When an individual is asphyxiated from whatever cause, the appearance is cyanotic. Respiration is interfered with, but is not noisy. There is no conjugate deviation of the head and eyes. Measures of resuscitation may, in a little while, prove successful, while the paralytic phenomena attendant upon cerebral hemorrhage are absent.

Cerebral Softening.

What are the varieties of softening of the brain?

Softening of the brain may be acute or chronic. *Acute softening of the brain* may be a result of inflammation; it may follow vascular occlusion, thrombotic or embolic, arterial or venous.

Chronic softening of the brain is an affection of the degenerative period of life. It usually involves the white matter of the hemispheres. The symptoms to which it gives rise are not well defined. There is usually progressive loss of motor power and impairment of sensation, sometimes with rigidity. Mental deterioration may be slight or considerable. Life may be prolonged for from two months to two years, death usually resulting from bedsores, pneumonia, or some other intercurrent malady.

Cerebral Embolism.

Under what conditions does cerebral embolism occur?

A vegetation from a diseased cardiac valve may be washed into one of the vessels of the brain; or the plug may have its source in a thrombus that has formed in one of the cavities of the heart, or in a calcareous fragment from the wall of an atheromatous aorta.

To what symptoms does cerebral embolism give rise?

The sudden plugging of one of the vessels of the brain is usually manifested by transient loss of consciousness of varying duration, often accompanied by convulsions. On the return to consciousness, the symptoms of hemiplegia are evident, often right-sided, and associated with aphasia. In the course of a few days, secondary fever appears. Unilateral convulsions may be repeated. Choreoid, athetoid, and other disorders of movement are likely to develop.

How do the symptoms of cerebral hemorrhage differ from those of cerebral embolism?

Cerebral embolism is a disease rather of early life; hemorrhage is one of advanced life. Hemorrhage occurs in association with disease of the vessels; embolism with valvular disease of the heart or other source of emboli. Consciousness is invariably lost when hemorrhage takes place; it is more likely not to be in cases of embolism. Embolism is more likely to be cortical and to give rise to initial and consecutive epileptiform convulsions, to localized palsies and to aphasia; hemorrhage is more

likely to be central, to be unattended with convulsions, and to present hemiplegia. Both direct (local) and indirect (general) symptoms are more profound and more extensive in hemorrhage than in embolism. The detection of emboli in other organs is an important element in diagnosis.

Cerebral Thrombosis.

Under what conditions does cerebral thrombosis occur?

The most common cause of cerebral thrombosis is vascular disease, whether atheromatous, syphilitic or fibroid. The blood may also coagulate in the vessels in conditions of lowered vitality and enfeebled circulation. Increased coagulability of the blood predisposes to the occurrence of thrombosis. A thrombus may form from an embolus by extension.

To what symptoms does cerebral thrombosis give rise?

The actual occurrence of thrombosis in a cerebral vessel may for a long time be preceded by distressing headache, vertigo, numbness, tingling, muscular weakness, impairment of memory and mental failure. The formation of the thrombus is announced by loss of consciousness, of varying degree and duration, perhaps attended with convulsions or with delirium. In the course of a few days, decided secondary fever may appear. Subsequently, the evidences of hemiplegia become manifest; while recurring localized convulsions may take place. As in the case of embolism, involuntary and spasmodic disorders of movement in the affected parts are common. Thrombosis of the sinuses gives rise to localized edema of the face and scalp.

How are cerebral hemorrhage and cerebral thrombosis to be differentiated?

Thrombosis, unlike hemorrhage, is of not rare occurrence in infancy and extreme old age. The symptoms of hemorrhage set in suddenly, with loss of consciousness; those of thrombosis are preceded by prodromata and develop gradually, with headache, delirium, perhaps convulsions. The convulsions of thrombosis are wont to be localized or unilateral; if convulsions attend

hemorrhage, they are usually general. Hemorrhage is the more likely to occur under conditions of excitement; thrombosis, amid conditions of depression. In hemorrhage the circulation is active, the arterial tension high; thrombosis is not uncommonly associated with a stagnant circulation in a debilitated individual. The symptoms are more profound and extensive in hemorrhage than in thrombosis. Mobile spasm, athetosis and other disorders of movement more commonly follow thrombosis than hemorrhage. Recurrent convulsions also point to thrombosis rather than to hemorrhage.

Cerebral Tumor.

What are the more common varieties of tumor of the brain?

Cerebral tumors are most commonly gliomata, tuberculomata, sarcomata, gummata, myxomata, carcinomata or psammomata. They may develop in the cerebral mass, in the membranes, or in the bone.

Not rarely, cysticerci or echinococcus cysts develop in the brain. Gliomata sometimes follow injuries to the head.

Tuberculomata appear in young persons, and are usually associated with tuberculosis elsewhere than in the brain. Carcinomata occur later in life, give rise to a cachexia, and are associated with similar growths elsewhere. Gummata are present with other manifestations and a history of syphilis.

What are the symptoms of tumor of the brain?

The symptoms vary somewhat with the size and situation of the new-growth. Certain phenomena, however, characterize the presence of a neoplasm in any part of the brain. There is persistent headache, intensely aggravated in paroxysms; optic neuritis followed by atrophy; mental failure; impaired memory; emotional mobility; vomiting; vertigo; sometimes forced movements; slowness of speech; separation of syllables; confluence of articulation; sometimes aphasia; paralysis of varying distribution, associated with or followed by contractions; sometimes choreoid or athetoid movements, or a jerky incoöordination; convulsions resembling those of major or of minor

epilepsy, or of Jacksonian epilepsy, or attacks of transient convulsions; alterations of sensation; palsy of cranial nerves; sometimes conjugate deviation of the head and eyes; abnormalities of the pulse; interference with the respiration. The quantity of urine evacuated may be increased; there may be albuminuria or glycosuria. Finally, stupor and coma may be the precursors of death.

How may chronic nephritis simulate cerebral tumor; how are the two to be differentiated?

Chronic nephritis may be attended with headache, vertigo, vomiting, optic neuritis and convulsions, but the urine contains casts as well as albumin, edema exists, the arterial tension is heightened, the heart is hypertrophied, and the palsies of tumor are wanting.

What symptoms does anemia have in common with tumor of brain, and how are the two to be differentiated?

Both anemia and cerebral tumor may occasion headache, vertigo, optic neuritis and albuminuria, but the headache of anemia is not so intense as that of tumor; the optic neuritis of anemia is more rapid in development than is that of tumor; while the pallor, the breathlessness, the edema, the submissiveness to judicious treatment of the one, and the convulsions, the palsies, the progressiveness, and the resistance to treatment of the other are distinctive.

How are the epileptiform convulsions of cerebral tumor to be distinguished from the convulsions of essential epilepsy?

In themselves the convulsions may be indistinguishable, though those of tumor are the more likely to be local or unilateral. Epilepsy, however, fails to present the distinctive symptoms of tumor: headache, vertigo, vomiting, optic neuritis and palsies.

How may the symptoms of cerebral tumor be mistaken for those of cerebro-spinal sclerosis; how is the diagnosis to be made?

A tumor in the brain may occasion a peculiar jerky incoördination, but it does not occasion the nystagmus, the peculiarity

of speech, the exaggeration of reflexes of cerebro-spinal sclerosis, while headache and optic neuritis are more common and more intense.

Intracranial Aneurism.

To what symptoms does an intracranial aneurism give rise?

Aneurism of the cerebral vessels may develop as a result of vascular disease, syphilitic or otherwise, of traumatism and of embolism.

The symptoms of intracranial aneurism necessarily vary with the situation of the aneurism. The more constant symptoms are those of an intracranial tumor: headache, vertigo, vomiting, convulsions, palsies, optic neuritis. The detection of a bruit or the demonstration of symptoms dependent upon the presence of a tumor in the course of a cerebral vessel would be diagnostic.

The occurrence of rupture is indicated by the symptoms of cerebral hemorrhage.

General Paralysis of the Insane.

What are the symptoms of general paralysis of the insane?

General or *progressive paralysis of the insane*, or *paretic dementia*, is a fatal disease of insidious invasion, occurring in those who have abused their mental and physical energies, and characterized by psychic alteration and failure, with progressive palsy.

Syphilis also is a most important etiologic factor. The disease is much more common in men than in women, and in early middle life than at any other period.

Among the first symptoms are an alteration of disposition, and a change of character. An individual, previously more or less refined, affectionate, attentive, scrupulous, methodical, precise and neat, becomes careless, indifferent, negligent, coarse, indecent, irritable and forgetful. He fails to keep appointments. He makes ill-judged investments and extravagant expenditures. He eats irregularly and hastily, and sometimes eats and drinks excessively, and sleeps poorly.

The palsy is not of the ordinary type. There is at first loss of coördinating power for delicate tasks, such as writing or

painting, while general muscular strength, as for lifting, is but slightly impaired. The paralysis is associated with tremor. The pupils may be contracted, sluggish and unequal.

To these symptoms are added the so-called delusions of grandeur, which vary from time to time. The patient believes himself fabulously wealthy. He projects colossal enterprises, or announces wonderful discoveries. He is, however, contented with his surroundings. In a milder form, the delusions take the less obtrusive shape of an invincible, unreasoning optimism, and may be overlooked. In other cases there are alternations of depression and exaltation. Sleeplessness becomes more obstinate. Speech becomes defective, indistinct and stuttering. In writing, words, syllables, letters are omitted or misplaced or repeated. There is a peculiar tremor of the lips. The protruded tongue presents both rhythmical tremor and fibrillary contractions. The gait becomes shuffling. Control over the limbs is progressively lost. At this stage, apoplectiform and epileptiform seizures may take place, and paroxysms of rage and fury may occur. Sensibility is not impaired until late, when it may be almost abolished.

Deceptive lulls and remissions may interrupt the progress of the disease, but finally a condition of terminal dementia is reached. Nutrition gradually fails. Paralysis, mental and physical, becomes so great that death may result from the entrance of food into the air-passages. Coma, convulsions, painful contractions, obstinate diarrhea, pulmonary affections, especially pneumonia, are among the phenomena preceding death. At the autopsy chronic congestive, inflammatory and degenerative changes may be found in the brain and membranes and in the posterior and lateral columns of the spinal cord.

How are progressive paralysis of the insane and posterior spinal sclerosis to be differentiated?

As the symptoms of posterior spinal sclerosis may be superadded to those of progressive paralysis of the insane, the differentiation of uncomplicated cases from locomotor ataxia, as from cerebro-spinal sclerosis and paralysis agitans, essentially depends upon a recognition of the presence or absence

of those manifestations of central dissolutional processes that characterize progressive paralysis: the changes in character and disposition, the emotional disturbances, the impairment of memory, the maniacal outbursts, the delusions of grandeur, the peculiar tremor of lips and tongue, the inequality of the pupils, the derangement of the language-faculty in speech and writing, the progressive failure of voluntary motion and intellection, the epileptiform and apoplectiform attacks.

Acute development of the peculiar loss of coördinating power, mental and physical, without obvious cause, in a man of middle life, especially one given to mental, emotional, venereal or alcoholic excess, or with a history of syphilis, should at once excite a suspicion of the development of paretic dementia.

Sunstroke.

What are the symptoms of sunstroke?

Sunstroke, *heat-stroke*, *insolation*, of which there are at least two principal varieties, *heat-fever* or *thermic fever* and *heat-exhaustion*—is an affection of the heated term. It occurs in persons exposed to intense heat, solar or artificial, in whom the respiratory and cutaneous transpiration is checked.

Crowding and defective ventilation predispose to the occurrence of heat-stroke. The *onset* may be abrupt or gradual, the symptoms severe or mild (incomplete). While most often coming on during active muscular exertion, as in the case of laborers and marching soldiers, the attack not infrequently follows the mid-day meal. It usually but not invariably happens during the period of maximum heat of the day.

There occur increased frequency of micturition, headache, vertigo, nausea, vomiting, delirium, loss of consciousness, dysphagia, stertorous breathing and coma, associated in the febrile form with a hot, often flushed, skin, frequent pulse and high temperature. The pupils may be dilated or contracted, or contraction and dilatation may be present at different times in the same case. There may be rectal and vesical incontinence, sometimes partial suppression of urine. The duration of the symptoms varies from a few minutes to several hours.

Death may occur early or late; *recovery* may be speedy or tardy. Persistent headache, vesical irritability, choreoid movements of the upper extremities, mental impairment, epilepsy, and in rare instances hemiplegia, or other paralysis, usually transient, may supervene as *sequelæ*. Anhydrosis, with headache and elevation of temperature, annually recurring near the anniversary of the attack, has been observed.

General paralysis of the insane (paretic dementia) has been attributed to sunstroke.

What are the principal points of differentiation between heat-exhaustion and heat-fever, or sunstroke proper?

In heat-exhaustion the skin is pale, cool or cold, moist or clammy; the temperature is low, even subnormal; there is syncope rather than coma; the pupils are usually dilated; the pulse is feeble; convulsions are absent; and rapid relief follows the use of warmth and stimulants.

How does cerebral hemorrhage differ from sunstroke?

The coma of sunstroke, unlike that of cerebral hemorrhage, does not set in unannounced; nor is it as lasting or as profound; while dysphagia may be greater, stertor is less.

The symptoms of sunstroke are symmetrical; the pupils and the reflexes are equal; hemiplegia is not an ordinary sequel.

The temperature rises much higher in insolation than at any time in hemorrhage. The pulse is frequent, often feeble; that of cerebral hemorrhage is commonly slow and full.

It must, however, not be forgotten that cerebral hemorrhage may occur amid circumstances favorable for the development of sunstroke.

With what other conditions might sunstroke be confounded?

Sunstroke might be mistaken for acute alcoholism, meningitis, uremia or narcotic poisoning. Alcoholic excess may bring on an attack, so that the phenomena of both might be intermingled. The history and the symptoms detailed should prevent error in other cases.

In uremia, too, convulsions usually precede coma.

Delirium Tremens.

What are the symptoms of delirium tremens?

The continued ingestion of excessive quantities of alcohol sometimes gives rise to a condition in which with impaired appetite and gastric irritability are associated inability to sleep, tremor and delirium. The temperature is likely to be elevated. The urine may contain albumin. The patient is restless and evinces a tendency to talkativeness; he is always busily engaged with his delusions. Sometimes the delirium is more violent. The victim is terror-stricken by hideous illusions and hallucinations. The figures of the wall-paper, and articles of furniture are converted into crawling reptiles; attendants, into demons. The patient is unable to sleep. Hypnotics may prove futile. There is little desire for food. That which is taken may be rejected. If recovery is to take place, sleep is gradually restored; else the patient is worn out and succumbs to exhaustion. Pneumonia, especially of the apices, is not an uncommon complication of alcoholism. The use of tobacco during convalescence is sufficient to renew the attack.

How is delirium tremens to be distinguished from acute mania?

The delirium of alcoholic intoxication may be maniacal, but it is associated with a history of alcoholism, while acute mania usually develops in a person with a psychopathic family history, and has probably been preceded by a prodromal period characterized by a change in manner, in disposition or in behavior. The character of the illusions and hallucinations differs in the two disorders. Delirium tremens usually subsides after sound sleep for a number of hours consecutively, and recovery is complete. Evidences of insanity persist after the attack of acute mania is at an end.

How are delirium tremens and cerebral meningitis to be differentiated?

The distinction between delirium tremens and cerebral meningitis depends upon a knowledge of the cause, the history, the

course, the symptoms and the termination of the two affections respectively. Delirium tremens is preceded by a history of alcoholism; meningitis, by a history of traumatism or infection. Delirium tremens lasts from a few days to a week; meningitis for a much longer time. In meningitis there are, and in delirium tremens there are not, muscular spasm, convulsions and sensory disturbance, followed by paralysis. The delirium and tremor of alcoholism are peculiar and characteristic. The temperature may be slightly elevated above the normal in delirium tremens, but meningitis is distinctly a febrile disease.

Plumbism.

What are the clinical manifestations of lead-poisoning?

Lead may gain entrance into the system and give rise to toxic manifestations under diverse circumstances. Thus saturnine intoxication occurs in those who work with the metal, as miners, color-grinders, painters, plumbers, type-founders, compositors. Lead may also be introduced into the system by means of drinking-water conveyed through lead-pipes, or with food prepared in leaden vessels, or otherwise adulterated, or by the use of hair-dyes, of cosmetics, or of snuff packed in lead-paper, or by the medicinal administration of lead in some form. Next to colic, the most common manifestation of plumbism is multiple neuritis.

The symptoms vary in kind, degree, distribution and acuteness. The more common and the more characteristic are obstinate constipation, abdominal colic, wrist-drop, and a blue line on the gums. In addition, the nutrition is impaired and there is anemia. Among other symptoms are cramps in the legs, tremor, headache, convulsions, delirium and coma. The paralysis of the extensors of the wrist and fingers upon which the wrist-drop depends is bilateral, but does not involve the supinator longus. Sometimes the scapulo-humeral muscles are affected; sometimes the intrinsic muscles of the hand atrophy; at times, too, the lower extremities are paralyzed. The knee-jerks may be enfeebled or lost; station may be unsteady; coör-

dination may be deranged. Occasionally, symptoms of sensory derangement, laryngeal palsy, mental failure, melancholia, inequality of the pupils and impairment of vision are observed. Hysterical symptoms also have been noted. The pupils may be small and fail to react to light. The optic nerve may undergo

FIG. 52.



Wrist-drop from lead poisoning. (Gowers.)

atrophy. Paralyzed muscles display the reaction of degeneration. Cardio-vascular-renal changes are of strikingly common occurrence. Arthritic manifestations and deposits indistinguishable from those of gout are not rare. If potassium iodid be administered, the urine may respond to chemic tests for lead.

How may lead-poisoning be confounded with cerebral tumor, and how is the differentiation to be made?

When the system has been impregnated with lead, there may be optic neuritis, headache, convulsions, delirium and palsy; but there are also a blue line on the gums; a history of exposure to the toxic metal; of intestinal colic; of obstinate constipation; and the palsy gives rise to wrist-drop.

Torticollis.

What is torticollis?

Torticollis or *wry-neck* is a condition in which, from shortening or from spasm of one or more muscles of the neck, the head is maintained in an abnormal position.

What are the varieties of torticollis?

There are two types of wry-neck. In one, as a result of traumatism, or of defective development, the muscle, usually the sterno-mastoid upon one side, is atrophied and shortened.

The head is rotated to the side opposite to that of the affected muscle, which is conspicuous for its prominence. In the second variety of wry-neck, the muscles involved, most commonly the sterno-mastoid, the trapezius and the splenius, are in a state of active contraction, tonic or clonic, or alternately both. As a result, the head may be rotated or inclined to one side or retracted in over-extension, in accordance with the muscle or combination of muscles that participates in the spasm. Not infrequently, with the spasm of the muscles of the neck is also associated spasm of muscles of the arm or of the face.

Occupation-Neuroses.

What are the occupation-neuroses?

As the result of persistent and long-continued movement of a part in a constrained position, involving the activity of certain groups of muscles, spasmodic interference with the performance of the same movement may develop.

Thus, as a type of the disease, there is produced writers' cramp or scriveners' palsy. An analogous condition may be produced in telegraphers, shoemakers, piano-players, violinists, zither-players, seamstresses, smiths, painters, turners, watch-makers, knitters, engravers, masons, compositors, cigarette-makers, milkers, money-counters, typewriters, motormen, and others.

Writers' Cramp.

What are the symptoms of writers' cramp?

In those who write much, especially in such a way that the burden of the work falls upon the small muscles concerned, there sometimes develops a condition that renders writing impossible. Under such circumstances, as soon as an attempt to write is made, the fingers contract in spasm that may be painful, so that the act, if possible, is imperfectly accomplished. At other times, there may be tremor, and rarely weakness. In some cases, neuralgic pain is the most conspicuous symptom. Sometimes vaso-motor manifestations are present, such as local heat, discoloration, glossiness of skin, undue sweating. There may be slight muscular wasting and changes in electric reactions. In aggravated cases the manifestations are not confined to the

FIG. 53.



Mode of holding the pen favorable to the development of writers' cramp. (Gowers.)



Mode of holding the pen when writing becomes difficult. (Gowers.)

act of writing, but may appear upon attempts to perform other movements. The onset of the symptoms is usually gradual, but occasionally it is sudden.

Chorea.

What are the etiologic elements concerned in the determination of an attack of chorea?

Chorea is most common at or about the age of puberty, and

more so in girls than in boys. At Philadelphia, attacks are said to be more common in the spring of the year than at any other time. It has been observed that chorea is more common in white than in colored children. In many cases, an hereditary influence can be traced—directly, as chorea, or, more commonly, in the nature of other diseases of the nervous system, as epilepsy or insanity, or of rheumatism. Fright is frequently an exciting cause of the disease. Occasionally traumatism has preceded an attack. An obscure but well-determined causal relation exists between acute rheumatism, heart-disease and pregnancy, on the one hand, and chorea on the other.

What are the symptoms of chorea?

Chorea is a spasmodic neurosis, manifested by irregular, involuntary, incoördinated muscular movements, aggravated by excitement or observation. The movements cease during sleep, but they may be sufficiently violent to prevent sleep. They are often more decided on one side of the body than on the other. Speech is often interfered with. A slight degree of muscular weakness exists. The electric irritability is usually increased; sometimes there are slight qualitative changes. Occasionally pain is felt in the parts affected. The temperature is, as a rule, slightly elevated. There is mental irritability or dulness of intellect. Anemia is often present.

At times, hemic heart-murmurs are heard; at other times, murmurs dependent upon organic disease, either antecedent or coincident, are heard; endocarditis is not an uncommon complication of chorea.

Chorea is a self-limited affection, the average duration of which is about ten weeks. It is said sometimes to involve the diaphragm, and to give rise to obstinate hiccough. It is thought that the irregularity of the action of the heart, sometimes observed in choreic children, is dependent upon involvement of the cardiac muscle; chorea of the larynx has also been seen.

So-called choreoid movement, athetosis, and mobile spasm, are often observed in cases of paralysis of cerebral origin.

Hereditary Chorea.

What is hereditary chorea or Huntingdon's chorea?

A disorder resembling chorea sometimes appears in families, developing at the middle period of life, and possibly continuing until death, in hastening which it may play some part. It is attended with irregular, incoördinate movements in the hands and face. Speech is slow, difficult, and indistinct, and the gait is somewhat staggering. Evidences of mental impairment are, however, often present, and after death degenerative changes have been found in the cerebral cortex.

Spasmodic Tic.

What is spasmodic tic?

Spasmodic tic, habit-spasm or *habit-chorea*, is a disorder characterized by twitching or jerking of one or more muscles or groups of muscles, and manifested by blinking of the eyes, facial grimaces, shrugging of the shoulders, sniffing of the nose and the like. The movements may disappear from one part to appear in another. The condition may arise from local irritation, as from eye-strain or ocular or nasal disease, or as a result of imitation. It may also attend trigeminal neuralgia. It is sometimes associated with explosive utterances.

How are chorea and spasmodic tic to be differentiated?

The two affections differ, first of all, in etiology. Then the movements in each differ in character. In chorea they are incoördinate and purposeless, and widely distributed, being general or confined to one-half of the body. The movements of spasmodic tic, on the other hand, are apparently purposive and, as a rule, less widely distributed; further, they vary from time to time, one set alternating with another. Spasmodic tic is usually the more protracted disorder, sometimes lasting for years, and it yields less readily to treatment, especially to arsenic.

Epilepsy.

What is the etiology of epilepsy?

Epilepsy is immediately dependent upon an irritable condition of the nervous system that must be referred to changes in the nerve-cells, beyond our present means of recognition. The disease is more common in females than in males. In a considerable proportion, a neurotic heredity can be traced. In many cases, no etiologic factor can be determined. The first attack usually occurs in early life, frequently in conjunction with dentition, or in association with rachitis. In other cases, it may be excited by mental influences, especially anxiety, by falls or blows, by acute disease, by the presence of parasites in the intestine, by menstrual derangement, by nasal polypi, by phimosis, by cicatrices, and by other sources of peripheral irritation.

When no source of irritation, direct or reflex, can be found, the disease is designated *essential* or *idiopathic*.

What are the symptoms of epilepsy?

Epilepsy is a paroxysmal neurosis, characterized by attacks of varying frequency, duration and severity, which may appear as transient loss of consciousness (*petit mal*, *minor epilepsy*) or may be preceded by a premonitory sensation or aura, followed successively by a cry, loss of consciousness, frothing at the mouth, tonic followed by clonic convulsions, headache, biting of the tongue and somnolence (*grand mal*, *major epilepsy*).

The attacks of petit mal may be so inconspicuous as to escape observation. There may be simply a vertiginous or other abnormal sensation, or an abrupt interruption in speech; or the grasp of an object held in the hand may be relaxed; or there may be a twitching of the muscles of the face.

When the seizure is preceded by an aura, this may be motor, sensory, visceral or psychic. In the attack the face becomes at first pallid, then flushed, and finally cyanotic; the pupil is dilated and insensible to light; urine and feces may be expelled involuntarily. The urine may contain a trace of albumin.

Mental deterioration ensues in a degree corresponding to the

frequency of the attacks. Paroxysms may be succeeded by transient hemiplegia and aphasia, by the automatic performance of some act or by mania.

Sometimes a series of convulsions occurs, one attack scarcely ceasing before another begins, constituting the *status epilepticus*.

Nothing is known of the pathologic anatomy of epilepsy.

How are the epileptiform convulsions of cerebral disease to be distinguished from attacks of true epilepsy?

Epileptiform convulsions usually result from lesions involving the cortex cerebri. The most common of these lesions are softening from vascular occlusion, hemorrhage and new-growths. The convulsions occasioned are likely to be local in character and not at the onset attended with loss of consciousness; while the convulsions of epilepsy are usually general and simultaneous with loss of consciousness. In the intervals between epileptiform convulsions, some evidences of cerebral disease—hemiplegia, athetoid movements, choked discs—are likely to be present.

What are the distinctions between chorea and epilepsy?

Epilepsy occurs in paroxysms with intermissions, and may persist during the greater part of life; chorea is but a temporary, self-limited condition, the symptoms of which are constant during the continuance of the disease. The movements of chorea do not possess the violence of the convulsions of epilepsy; nor in chorea is consciousness lost or the tongue bitten.

How are the attacks of syncope and those of petit mal to be differentiated?

Syncope occurs under well-known circumstances that occasion temporary cerebral anemia; the pulse at the wrist is almost obliterated; loss of consciousness is of brief duration; the return to the normal condition takes place slowly. In case of *petit mal*, there have been previous attacks of a similar character or perhaps of major epilepsy; the loss of consciousness is sudden and transitory; the normal condition is soon resumed; there may have been slight muscular twitchings; or urine may have been passed involuntarily; or some automatic action is performed.

How is eclampsia to be distinguished from epilepsy?

By eclampsia is meant convulsions occurring incidentally and temporarily, as during dentition, or in connection with pregnancy and parturition. The relation of eclampsia to epilepsy and epileptiform convulsions is uncertain. The recognition of eclampsia and its distinction from epilepsy and epileptiform convulsions depend upon its incidental and transient occurrence in connection with one of the conditions that give rise to it.

How are the convulsions of uremia and those of epilepsy to be differentiated?

The convulsions of uremia are not unlike those of epilepsy. Their significance, however, depends upon their association with edema, a peculiar mustiness of the breath, the presence of albumin and tube-casts in the urine, and the existence of degenerative changes in the retina. If obtainable, the previous history may determine the diagnosis.

Hysteria.

What are the symptoms of hysteria?

Hysteria is a disease of the nervous system, more common in females than in males, and in early adult life than at any other period. In many cases a neuropathic heredity can be traced. The manifestations of the disease may be induced by mental emotion, anxiety, visceral disease and other disease. As a rule, they are sudden in onset and as sudden in disappearance. They may be dissipated by a profound mental impression. The symptoms are protean; some are continuous, others paroxysmal. Among the former are derangements of sensibility and motility, visceral, vaso-motor and cerebral disturbances; the paroxysmal symptoms include convulsive seizures. That which fundamentally characterizes the disease is a loss of self-control, a deficiency of will-power, a heightened self-consciousness, a tendency to exaggeration and a morbid desire for sympathy. The patient is moved to laughter or tears with undue readiness and without adequate cause. Duplicity and deception are practised, and there is an irresistible tendency to imitation. Sensation may

be lost or heightened, or otherwise abnormal, locally, or upon one-half of the body, or in irregular distribution, with unsteady station and gait when the eyes are closed. Spinal tenderness is common. There is frequently a recurrent sensation of obstruction in the throat, as of a ball rising from the stomach, or of a sense of pharyngeal constriction—the so-called *globus hystericus*. At times, there is a feeling of intense pain in the head, such as might be occasioned by the driving in of a nail—the *clavus hystericus*. The special senses may be variously impaired. Motor palsy of diverse distribution may exist, occasioning paraplegia, hemiplegia or monoplegia, unattended with muscular wasting or alteration in the electric reactions or impairment of activity of the sphincters. Sometimes there is inability to stand or to sit—*astasia-abasia*. The reflexes are usually unaltered; in a number of cases, they are exaggerated. There may be aphonia, loss of the power of articulation, retention of urine, constipation, stammering. Contractures may take place and give rise to troublesome deformity. Laryngeal spasm may occasion distressing dyspnea, or a peculiar barking cough. Independently of any spasm, there may be increased frequency of respiration, with shortness of breath. Spasm of the pharynx or of the esophagus may entail the rejection of all food ingested. There may in addition be such repugnance to food that little is taken. On the other hand, the appetite may be strangely, even disgustingly perverted. There may be local flushing and perspiration. In females, menstruation may be deranged; ovarian tenderness is common. The paroxysmal attacks of convulsions are marked by tonic and clonic spasm, without loss of consciousness or biting of the tongue. The arms and legs are wildly thrown about by coördinated muscular activity—purposive movement. The patient may shriek and bite the lips. Opisthotonos and trismus may be developed. At times, there are evidences of profound psychic disturbance, as manifested by hallucinations and delusions, perhaps of an erotic nature. Lethargy and catalepsy are sometimes observed. The temperature may display decided irregularity.

How are hysterical paralysis and the cerebral paralysis of organic origin to be differentiated?

The palsy of hysteria, hemiplegic, paraplegic, diplegic or monoplegic, is usually atypical.

The face is rarely involved. Sensory derangement is common. The evidences of secondary descending degeneration: contractures and exaggeration of reflexes may be wanting. There may be constipation and retention of urine, but not incontinence.

While the onset may have been sudden, it will not have been apoplectic. Other symptoms of hysteria may make the character of the palsy clear, but the mere existence of the symptoms of hysteria should not obscure the possible simultaneous existence of organic disease.

How are the convulsions of hysteria and those of epilepsy to be differentiated?

An epileptic paroxysm takes place without special excitation, and sets in abruptly, often with a cry; an hysterical attack may be brought on by emotion, sets in gradually, and is attended with screaming. The convulsions of epilepsy pursue a regular sequence, and are associated with cyanosis, biting of the tongue, and insensibility of the iris to light; in the hysterical attack, the members are wildly thrown about, the pupil responds to stimulation, while the patient may bite the lips or hands or other persons or things. Involuntary extrusion of urine and feces may take place in epilepsy; but does not occur in hysteria.

Loss of consciousness is complete in epilepsy; consciousness is retained or perverted in hysteria, the paroxysm of which is characterized by talkativeness.

The epileptic attack lasts but a few minutes; the hysterical, ten minutes or longer. Opisthotonos is common in hysteria, exceptional in epilepsy.

An epileptic seizure may occur at any time and under any circumstances; hysterical attacks take place only in the presence of a second person. The hysterical patient is careful in the paroxysm to suffer no injury; the epileptic falls, whatever the attendant dangers, powerless to avert them.

An hysterical patient may, however, also be epileptic.

What are the distinctions between spastic paraplegia and hysterical paraplegia?

Spastic paraplegia develops at an age when hysteria is common; but in hysteria the reflexes may not be exaggerated and the peculiar muscular spasm of lateral sclerosis does not occur, while sensory and emotional symptoms cannot escape observation.

How are acute myelitis and hysterical paraplegia to be differentiated?

Myelitis should not be overlooked because it occurs under circumstances amid which hysteria is to be expected. Exaggerated reflexes, trophic changes, wasting, derangement of sphincters and girdle-pain indicate the existence of more than hysteria.

How are the symptoms of hysteria and those of meningitis to be differentiated?

If symptoms of meningitis appear in an hysterical individual, their true significance may be overlooked. Convergent strabismus may be hysterical; but divergent strabismus always depends upon organic disease. There may be retention of urine in hysteria, but there is never incontinence. Trophic changes and continued elevation of temperature are evidences of the existence of something more than functional disease.

How are hysteria and cerebral tumor to be differentiated?

Hysteria may present any of the imitable symptoms of cerebral tumor, but in addition there will always be indubitable evidence of their nature. Optic neuritis or ptosis, of course, is not to be duplicated. Hysterical manifestations in association with cerebral symptoms should not be permitted to obscure the existence of organic disease.

Neurasthenia.

What is neurasthenia?

Neurasthenia is a condition characterized by undue readiness of fatigue from ordinary mental or physical activity, with enfeebled power of recuperation, probably dependent upon nu-

tritive disturbance of the nervous system and manifested by irritable weakness.

What are the causes of neurasthenia?

Neurasthenia is more common in women than in men, and during active adult life than at any other period. The predisposition is transmitted by heredity. The disorder is superinduced by overwork, with inadequate rest, by worry, anxiety, grief, mental shock, or other emotional disturbance, by excessive use of tea, coffee, tobacco, and alcohol, by drug-addiction, by sexual excesses, by traumatism, by protracted lactation, by wasting discharges, by debilitating diseases, by impaired nutrition from whatever cause.

What are the symptoms of neurasthenia?

Among the most common clinical manifestations of neurasthenia are a sense of tire or fatigue, spontaneous or readily induced, with indisposition to activity; a feeling of distress in the head, or of fulness, or of weight or pressure; noises in the ears, especially in the recumbent posture; disturbed and unrefreshing sleep at night, with perhaps undue drowsiness by day, tremulousness or agitation or fright on slight or on no provocation; morbid fears of all kind; mental depression, with a tendency to weep; irritability of temper; pain in the back and elsewhere. The reflexes are often sensitive and irritable, and station uncertain. Digestion is commonly impaired and constipation is the rule, though sometimes the bowels are loose. There may be increased frequency of micturition, but, as a rule, the urine is not increased in amount, nor does it contain abnormal ingredients. In women menstruation is likely to be deranged and flushing is complained of. Palpitation of the heart may be present, together with pulsation of the aorta and a sense of beating in various situations.

How are hysteria and neurasthenia to be differentiated?

The one is a psycho-neurosis characterized especially by want of functional coördination and attended by varied disturbances of cerebro-spinal activity and of motility and general and special sensibility; the other is essentially a condition of irritable weakness, a fatigue psycho-neurosis, and wanting in the character-

istic motor, sensory and psychic phenomena of the former. Hysteria is often of abrupt origin, due to a powerful cause, and terminating abruptly; while neurasthenia, as a rule, is due to a long-acting cause and of gradual development and decline. The symptoms of hysteria are often local in distribution; those of neurasthenia usually general. Both diseases may be present in the same individual.

Family Periodic Paralysis.

What is family periodic paralysis?

Under this name there has been described a form of flaccid motor palsy, of varying degree and distribution, with loss of reflexes and electric irritability, but without sensory derangement. The condition recurs periodically, the patient being free from symptoms in the intervals. In many cases a family history of the disorder can be elicited.

Tetanus.

What are the symptoms of tetanus?

Tetanus is a spasmodic disorder dependent upon a specific bacillus contained in soil and introduced into the system through wounds or abrasions. It is manifested by painful rigidity of the head and jaw, soon progressing to trismus, and by stiffness of the tongue. In turn, the rigidity involves the muscles of the face (resulting in the *risus sardonicus*), the muscles of the trunk, the respiratory muscles, and the diaphragm. To the tonic spasm of the muscles are added frequently recurring clonic exacerbations, which may be induced by external irritation. Sometimes there are paralysis of the facial muscles and difficulty in swallowing.

The body may be arched in strong extension and supported only on the head and heels, constituting *opisthotonos*; it may be strongly arched forward (*emprosthotonos*), or laterally (*pleurosthotonos*), or it may be rigidly straight (*orthotonos*). The symptoms of tetanus set in at a variable period after inoculation—from a few hours to several days. The duration of the disease likewise varies from a few days to several weeks. Recovery is

exceptional. Towards the close of life or even after death, the temperature may rise to an extraordinary height.

How are hysterical trismus and opisthotonos to be distinguished from tetanus?

Hysteria is an unruly disease, the symptoms of which are disorderly in appearance, while the symptoms of tetanus appear in fairly regular succession. Should trismus or opisthotonos develop as a manifestation of hysteria, it is likely to be associated with other symptoms of hysteria. The paroxysms remit and recur and do not go on to a fatal termination.

How is strychnine-poisoning to be distinguished from tetanus?

Strychnine-poisoning gives rise to some of the manifestations of tetanus. If trismus develops in strychnine-poisoning, it does so late; in tetanus, it is one of the first symptoms.

The convulsions of strychnine-poisoning are intermittent, but may be induced by external irritation; those of tetanus are continuous, with paroxysmal exacerbations; inquiry into the history of the case may elicit important evidence.

How are hemorrhage into the spinal membranes and tetanus to be differentiated?

Meningeal hemorrhage is sudden, tetanus gradual, in onset. Pain is a more prominent symptom in hemorrhage than in tetanus. Trismus is wanting in hemorrhage; it is characteristic of tetanus. Spasm is constant in tetanus, with exacerbations; intermittent in meningeal hemorrhage.

How are acute spinal meningitis and tetanus to be differentiated?

Meningitis sets in abruptly with a chill, followed by elevation of temperature; tetanus is of gradual development, usually after an injury, and is at the outset unattended with elevation of temperature. Trismus is characteristic of tetanus, but exceptional in meningitis.

The convulsions of tetanus are excited by slight peripheral irritation; the muscular contractions of meningitis are induced by efforts at movement.

Tetanus is far more commonly fatal than is meningitis. Motor and sensory impairment are common sequelæ of meningitis.

Tetany.

What is the etiology of tetany?

Tetany is most common in infancy and early adult life, when males are more prone to the disease than females. Occurring later in life, females are affected in larger proportion. In some cases, an hereditary influence can be made out. In many, diarrhea is an exciting cause. In others, the affection has been preceded by one of the acute infectious diseases. Pregnant or nursing women seem especially predisposed. The disease has been observed to develop in a considerable number of cases following removal of the thyroid gland. It has also been seen in association with dilatation of the stomach. In children it is often associated with rachitis, laryngismus stridulus, carpo-pedal spasm and convulsions. Other possible causes are exposure to cold and blows and injuries. Epidemics of the disease have been observed.

What are the symptoms of tetany?

Tetany is an affection characterized by muscular spasm, of symmetrical distribution, which usually begins and is most marked in the extremities.

The spasm may be continuous, remittent or intermittent.

A paroxysm may be induced by compression of the vessels and nerves of a part.

At the onset of the attack, there may be headache, vomiting, spinal pain, numbness and tingling. There may be moderate elevation of temperature.

In the intervals between paroxysms, the mechanical excitability and electric irritability of nerve and muscle are heightened. The disease may continue for a period ranging from a few days to several months.

Cases in which the spasm is intermittent are longer in duration than those in which the spasm is continuous.

How are tetany and tetanus to be distinguished?

The spasm of tetany is more likely than that of tetanus to be intermittent. Trismus is an early symptom of tetanus; if

it occurs at all in tetany, it appears late. The participation of the extremities in the spasm is a feature of tetany.

Hydrophobia.

What are the clinical features of hydrophobia?

Hydrophobia results from the inoculation of man with rabies of animals. The infection is usually transmitted by the saliva of a rabid beast, as a dog, a cat, a fox, or a wolf, through a bite or a preëxisting wound. The period of incubation of hydrophobia is extremely variable, but is on an average from six to ten weeks. During this time, the primary wound may have healed, and no symptom have been present. Local pain may be perceived preceding the development of the disease proper, with the onset of which there are a sense of malaise, mental depression and slight difficulty in swallowing. Sleep is impaired and there may be respiratory spasm. The stomach now becomes intolerant, rejecting everything introduced. The muscular spasm extends and cutaneous hyperesthesia becomes manifest. External stimuli readily induce convulsions. The mental distress becomes intense. The temperature is elevated. Priapism may occur. Albumin and sugar are sometimes found in the urine. Ultimately, paralytic phenomena may supervene. Death may result from exhaustion, by reason of the inability to retain food, and the violence of the convulsions; from suffocation, as a result of respiratory spasm; or from heart-failure.

How are lyssophobia or pseudo-hydrophobia and true hydrophobia to be differentiated?

Persons who have been bitten by animals, rabid or not, develop a state of fear and dread, sometimes difficult to distinguish from true hydrophobia. Usually, however, the laryngeal and pharyngeal spasm characteristic of hydrophobia is wanting, while judicious moral assurance may cause a dissipation of the symptoms.

How are hydrophobia and tetanus to be differentiated?

In hydrophobia there is the history of a bite, with a long period of incubation; in tetanus of wound-infection with earth, and a short period of incubation. In hydrophobia, respiratory and pharyngeal spasm is an early manifestation; in tetanus, trismus is among the first symptoms. In the former all food is rejected; in the latter, if food can be introduced into the mouth, there is no difficulty of retention. Tetanus does not present the intensity of mental distress and disturbance encountered in hydrophobia.

Aural Vertigo—Labyrinthine Vertigo.

What is aural vertigo?

Menière's disease, *aural* or *labyrinthine vertigo*, is an affection dependent upon a pathologic condition of the terminal fibers of the auditory nerve in the labyrinth. The disease of the labyrinth may be inflammatory, gouty, syphilitic or degenerative. The symptoms occasioned are impairment of hearing, tinnitus aurium and vertigo. The last is aggravated in paroxysms, in which in addition there are nausea and vomiting, with pallor of the face and cold sweats. The patient may fall to the ground and the vertigo be so intense that he is temporarily unable to arise.

How are epilepsy and aural vertigo to be differentiated?

Epilepsy is, and aural vertigo is not, attended with muscular spasms and loss of consciousness. In the intervals between the paroxysms of aural vertigo some degree of dizziness persists and there are also impairment of hearing and tinnitus aurium, which are not accompaniments of epilepsy.

Exophthalmic Goiter.

What are the symptoms of exophthalmic goiter?

Exophthalmic goiter, also called *Graves's disease* and *Basedow's disease*, is an affection characterized by increased frequency of action and palpitation of the heart, protrusion of the eyeballs,

enlargement of the thyroid gland and tremor. It is clinically a vaso-motor disorder of autotoxic origin, and its manifestations are probably dependent upon excessive functional activity of the thyroid gland.

The disease is more common in females than in males, and in adult life than at any other period. In some cases a neuro-

FIG. 54.



Case of exophthalmic goiter. (Personal observation.)

pathic heredity can be traced. The exciting cause is usually mental emotion, shock, grief, or anxiety. Traumatism is sometimes the determining factor, and the disease has developed after operation followed by infection. By some observers in-

toxication from the intestinal tract is believed to be a potent cause.

The onset may be either insidious or abrupt, and the course gradual or rapid.

Palpitation of the heart, associated with or independent of an organic lesion, is one of the earliest symptoms. The action of the heart is increased both in frequency and in energy, and is often extremely irregular. The tachycardia and the palpitation may reach a high degree of intensity, the pulse sometimes exceeding 140 in the minute. There may also be evident pulsation in the arteries and a bruit in the vessels of the neck.

Anemia is often an early symptom. In the progress of the case, the eyes protrude and the thyroid gland becomes enlarged.

The exophthalmos and the goiter may each be symmetrical or unilateral; the protrusion and the enlargement, respectively, are not uncommonly greater upon the right side. In some cases, the upper lids are retracted, displaying the sclerotic, and do not follow the eyes when the glance is depressed. Infrequency of winking and insufficiency of convergence have been observed.

Over the enlarged thyroid gland a blowing systolic bruit is usually to be heard. The goiter frequently pulsates; it is at first usually elastic; ultimately it may become firm. The appearance of the goiter may be intermittent; when permanent, the enlargement may undergo spontaneous variation.

Among other symptoms are nystagmus, tremor of the extremities, dermagraphism, an abnormal sense of heat, flushing and perspiration. It has been stated that the bodily electric resistance is diminished. Various cutaneous affections have been noted. In women, menstruation is commonly deranged.

In some cases, albuminuria, in others, diabetes has been observed. Hemoptysis, hematuria, hematemesis and other hemorrhages, and sometimes purpura, occur. Mental changes are not uncommon. Mania is an occasional complication.

Exophthalmic goiter has been followed by myxedema; it is not directly fatal. Spontaneous recession or recovery is possible. Temporary exacerbations take place.

What are the features distinguishing exophthalmic goiter from simple goiter?

Occasionally, the enlargement of the thyroid gland may be the first symptom of exophthalmic goiter observed. The bruit heard over the gland in exophthalmic goiter is wanting in the simple enlargement. Then, investigation will disclose the existence of palpitation and examination will reveal increased rapidity of the action of the heart, with the development, in the progress of the case, of protrusion of the eyes and the other symptoms of exophthalmic goiter. Even before the heart is much disturbed, there may be a tendency to flushing of the face, and occasionally red blood-cells may be found in the urine.

Vasomotor Ataxia.

What is vasomotor ataxia?

In predisposed persons, especially those of neurotic inheritance, there occur under various exciting influences, principally emotion, temperature, weather, drugs, and other toxic ingesta or disturbed metabolism, and in most women during the climacteric, symptoms of deranged circulation, spasmodic, paretic, or mixed. These vary much in intensity from mere "flushes of heat" and "chilly sensations," urticarious and other eruptions, to hemorrhagic discharges, vertigo, and transient blindness, tachycardia, angina, etc. Exophthalmic goiter, hay-fever, angioneurotic edema, certain forms of asthma and renal leakage, and other forms of neuro-vascular disorder may all be exaggerated types of this affection.

Cretinism.

What is cretinism?

Cretinism is a morbid condition, occurring endemically or sporadically and manifested by characteristic mental and physical abnormalities. The body is undersized; the head is broad, but shallow; the eyes are far apart and the nose is flat; the lips are thick; the tongue is sometimes enlarged; the hands are

broad and spade-like; the skin is dry, rough and wrinkled; the aspect is that of age; the hair is straight and stiff; the intelligence is feeble, even to the degree of idiocy; masses of fat sometimes appear in the posterior triangles of the neck; the thyroid gland is often enlarged; it is sometimes wanting. In a number of cases premature ossification of the occipito-sphenoidal suture has been found. Some members of families in which cretinism existed have presented goiters. The condition is closely related to, if not identical with, myxedema.

Myxedema.

What are the clinical features of myxedema?

Myxedema is a peculiar condition, apparently more common in women than in men, in which a mucoid substance is found in the subcutaneous tissue in various parts of the body, in association with anemia, atrophy of the thyroid gland and mental impairment. The condition has been termed *sporadic cretinoidism* in distinction from sporadic cretinism.

The patient becomes dull, stupid and irritable; *speech* thick and hesitant; the *memory* enfeebled. *Movement* is awkward and clumsy, and there is undue readiness of fatigue. *Sensibility* is impaired, while abnormal sensations of heat and of chilliness are perceived. There may be visible flushing.

The *heart* is slow and feeble, the *respiration* sluggish, with breathlessness on exertion; the *temperature* is, as a rule, subnormal; the *appetite* is impaired, and taste is deranged; the *digestion* is enfeebled; the *nutrition* is poor, the *urine* is reduced in quantity, and late in the disease may be albuminous. In women menorrhagia is common.

The *face* is pallid and puffy, and of roundish ("full-moon") outline; the expression is dull and stupid; the *thyroid gland* is diminutive or absent. The tongue is enlarged, and swellings form above the clavicles. The teeth may become carious and fall out. Speech is slow, thick, monotonous, measured. There may be choking sensations and difficulty in swallowing. The *skin* is thickened, dry, rough, sometimes scaly, somewhat translucent in appearance, of a doughy consistence, but with a

certain degree of elasticity; there is no edema and no pitting on pressure; perspiration is diminished. The *hands*, if affected, become square or "spade-shaped"; the *fingers* clubbed; and the nails brittle and distorted. The *hair* is dry and harsh and brittle, and may fall out.

FIG. 55.



Myxedema. (Personal observation.)

In the progress of the case mental and physical failure increases, irritability becomes marked, hallucinations develop, stupor sets in, and death may take place in coma, from exhaustion or from uremia.

A condition resembling myxedema, occurring in circumscribed areas in various situations—*cachexia strumipriva*—has been induced in man and in animals by removal of the thyroid gland. Myxedema has followed exophthalmic goiter.

How are myxedema and scleroderma to be differentiated?

In a given case the diagnosis between myxedema and scleroderma may be exceedingly difficult. In scleroderma, however, the skin is brawny and indurated; not elastic and doughy, as in myxedema. In scleroderma, too, the thyroid gland remains unaffected, while the mental phenomena of myxedema are wanting. The histologic changes in the two conditions differ: in scleroderma there is hyperplasia of the subcutaneous connective tissue; in myxedema there is a mucoid deposit or degeneration.

How are myxedema and obesity or adiposis to be differentiated?

The various forms of fat-deposition and fat-accumulation bear only a remote resemblance to myxedema. They may be diffuse or circumscribed in various situations. The nature of the tissues is, however, different, and, accordingly, while myxedema yields a somewhat brawny feeling, adiposis yields a lobulated or worm-like feeling. Mental and trophic alterations are more pronounced in myxedema than in adiposis. Thyroid

FIG. 55.



Dolorose adiposis. (Personal observation.)

atrophy is distinctive of myxedema; the thyroid gland may apparently be unchanged in adiposis. A dolorose form of adiposis has been described.

Akromegaly.

What is akromegaly?

Akromegaly is a morbid condition characterized by increase of the distal parts of the extremities in thickness, but not corre-

spondingly in length, the bones of the hands and feet and face being especially affected. Some of the fibro-cartilages, as those of the ear and larynx, also become enlarged. The related soft parts undergo corresponding enlargement. There develop decided weakness and slight muscular atrophy. The interosseous spaces become exaggerated. Owing to the enlargement of the inferior maxillary and the frontal bones the face assumes a peculiarly elongated, elliptical outline. The hypertrophy of the nasal bones gives the nose a thickened appearance. The enlargement of the malar bones increases the normal temporal fossæ. The enlargement of the frontal sinuses and the projection of the superciliary ridges give the forehead a retreating appearance. The chin is prominent, and the lower teeth project beyond the plane of the upper teeth. The tongue is enlarged and thickened. The lips and eyelids may also be thickened. There are increased thirst, usually polyuria, and sometimes glycosuria. The appearance may be cachectic, the conjunctivæ icteric. There is commonly spinal curvature, usually a cervico-dorsal kyphosis, with a compensatory lumbar lordosis. The abdomen may be protuberant, but later it may be diminished. The stature is at first increased, The skin is thickened and may be pigmented. The nails are excessively developed and may pre-

FIG. 57.



Akromegaly. (Personal observation.)

sent longitudinal striation. There may be severe spontaneous pain; headache is common. There may be somnolence. In many cases varicose veins and hemorrhoids have been observed. In individual cases there has been hemianopsia, limitation of the visual fields, blindness, or deafness. In women menstruation is usually deranged. The thyroid gland may be atrophied or hypertrophied. The thymus is often present, sometimes enlarged. The disease appears most commonly in young adults, and is chronic in course.

How does akromegaly differ from osteitis deformans, Paget's disease?

Tbe disease called by Sir James Paget *osteitis deformans* more especially involves the long bones, which, while they become

FIG. 58.



Outline of the face, 1, in myxedema; 2, in akromegaly; 3, in osteitis deformans. (Marie.)

enlarged, also become distorted, with resulting deformities; akromegaly displays a predilection for the small bones of the hands and feet and face, and is unattended with distortion and deformity. In osteitis deformans, as a result of the *enlargement of the cranial bones*, the face presents a triangular outline, with the base above and the apex below; in akromegaly the *enlargement of the bones of the face* occasions a characteristic elongated, elliptical outline. In akromegaly there may be an increase in stature; in osteitis deformans there is more likely to be a diminution. The former appears early, between fifteen and thirty-five; the latter, later, after forty. Akromegaly may appear in

several members of one family; osteitis deformans does not. The invasion of akromegaly is symmetrical; that of osteitis deformans is indiscriminate.

How may the differential diagnosis between akromegaly and myxedema be made?

The enlargement that appears in myxedema is dependent upon changes in the skin and subcutaneous soft structures;

FIG. 59.

Hand from a case of empyema with pulmonary hypertrophic osteo-arthropathy. (Personal observation.)

while the enlargement of akromegaly is due to a hyperplasia of bone. In the latter the skin is healthy and mobile; in the former it is thickened and adherent. In myxedema, in contrast to akromegaly, the hands and feet may be entirely uninvolved. In myxedema the face is almost round, "moon-shaped;" in akromegaly it is elliptical and elongated, with a prominent lower jaw, prominent malar processes and prominent nasal bones.

How is akromegaly to be discriminated from pulmonary hypertrophic osteo-arthropathy?

Under a name signifying "*hypertrophic disease of bones and joints in association with pulmonary lesion*," Marie has described a group of cases presenting, in connection with empyema, pleurisy, chronic bronchitis, tuberculous disease of the lungs, and possibly other disease of the lungs and their appendages, skeletal changes somewhat resembling those of akromegaly.

There are kyphosis, enlargement of the distal epiphyses of the bones of the forearm and of the leg, and, most characteristic, an enlargement in all directions of the distal phalanges of both fingers and toes, with the peculiar curvation of the nails that gives rise to the so-called Hippocratic finger. The lower jaw projects, but not so markedly as in akromegaly, and the other facial bones are not enlarged; on the contrary, the features are usually sharpened.

There is commonly dulness on percussion over the manubrium sterni, supposably dependent upon the persistence of the thymus gland. The thyroid gland may be wanting.

Akromegaly.	Pulmonary osteo-arthropathy.
Stature heightened.	Stature not heightened.
Cervico-dorsal kyphosis.	Dorso-lumbar kyphosis.
Projection of abdomen.	No projection of abdomen.
Enlargement of face.	No enlargement of face.
Macroglossia.	Tongue not enlarged.
Hands broadened—spade-like.	Hands lengthened—not spade-like.
Digits uniformly enlarged.	Terminal phalanges enlarged.
Nail short, broad, flat, not reaching end of finger.	Hippocratic nail—curved, longitudinally striated, overlapping finger, "parrot-beak-like."
Epiphyseal enlargements uncommon.	Epiphyseal enlargements the rule.
Polyuria and disturbance of special senses.	Intra-thoracic disease.

FIG. 60.



Rhizomelic spondylosis. (Personal observation.)

Rhizomelic Spondylosis.

What is rhizomelic spondylosis?

This is a disorder characterized by hyperplastic alterations in the joints of the spinal column and of those between the trunk and the extremities, leading to deformity and impaired mobility. Sometimes spinal-nerve roots become included in the morbid process, with the development of corresponding sensory or motor symptoms.

INDEX.

ABORTIVE typhoid fever, 39
Abscess, cerebral, 349
Abscess, cerebral, differentiated from—
cerebral meningitis, 350
cerebritis, 350
tumor of the brain, 350
Abscess, genito-urinary, 274
Abscess, lumbar, differentiated from—
perityphlitis, 230
typhlitis, 230
Abscess of the kidney, 295
Abscess of the kidney differentiated from perinephric abscess, 296
Abscess of the liver, 250
Abscess of the liver differentiated from—
actinomycosis of the liver, 251
carcinoma of the liver, 251
hydatid cyst of the liver, 257
occlusion of the biliary passages, 251
pleuritic effusion, 168
Abscess of the mediastinum differentiated from aneurism of the aorta, 142
Abscess of the right ovary differentiated from—
perityphlitis, 230
typhlitis, 230
Abscess of the spleen differentiated from pleuritic effusion, 168
Abscess, perinephric, 296
Abscess, perinephric, differentiated from abscess of the kidney, 296
Abscess, pulmonary, differentiated from—
bronchiectasis, 173
pulmonary gangrene, 182
pulmonary infarction, 181
pulmonary tuberculosis, 190
Abscess, retro-pharyngeal, 210
Abscess, retro-pharyngeal, differentiated from membranous croup, 156
Abscess, subphrenic, differentiated from pneumothorax, 197
Absorptive activity of gastric mucous membrane, 214
Acetone in the urine, tests for the presence of, 286
Achylia gastrica, 215
Acidity of gastric contents, 212
Acidity, total, of the gastric contents, 212
Acids, biliary, in the urine, 271
Actinomycosis, 90
Actinomycosis of the liver differentiated from abscess of the liver, 251
Action of the heart, 120
Acute alcoholism, 362
Acute anterior poliomyelitis, 318
Acute ascending paralysis, 319
Acute bronchitis, 170
Acute bulbar palsy, 323
Acute catarrhal pharyngitis, 204
Acute croupous pneumonia, 176
Acute dysentery, 226
Acute edema of the larynx, 151
Acute endocarditis, 139
Acute enteritis, 223
Acute gastric catarrh, 215
Acute gastritis, 215
Acute general disease differentiated from cerebral meningitis, 343
Acute gout, 99
Acute hepatitis, 247
Acute indigestion, 215
Acute infective jaundice, 67
Acute intestinal catarrh, 223
Acute laryngitis, 150
Acute mania, 344, 363
Acute miliary tuberculosis, 191
Acute myelitis, 315

Acute nasal catarrh, 148
Acute nephritis, 292
Acute pancreatitis, 265
Acute pericarditis, 235
Acute peritonitis, 241
Acute phlegmonous pharyngitis, 204
Acute pleurisy, 165
Acute rheumatism, 95
Acute synovitis, 97
Acute tuberculous laryngitis, 152
Acute tuberculous pharyngitis, 205
Acute yellow atrophy of the liver, 248
Addison's disease, 116
Adiposis, 387
Ague, dumb, 58
Ague-cake, 59
Air-passages, obstruction of, differentiated from croup, 155
Air-passages, upper, diseases of, 146
Akoria, 214
Akromegaly, 387
Akromegaly differentiated from—
myxedema, 390
osteitis deformans, or Paget's disease, 389
pulmonary hypertrophic osteo-arthropathy, 391
Akromikria, 340
Albumin in urine, tests for, 275
Albuminuria, 275
Alcoholic intoxication differentiated from cerebral hemorrhage, 353
Alcoholism, acute, differentiated from sunstroke, 362
Ambulatory typhoid fever, 39
Amyloid disease of the liver, 254
Amyloid disease of the liver differentiated from—
carcinoma of the liver, 255
cirrhosis of the liver, 254
Amyloid kidney, 293
Amyotrophic lateral sclerosis, 320, 321
Anacidity, 215
Analgesic panaris, 340
Anamnesis, 19
Anemia, 106
Anemia, cerebral, 347
Anemia, diagnosis of, 108
Anemia differentiated from cerebral tumor, 358
Anemia, idiopathic, 107
Anemia, infantile pseudo-leukemic, 112
Anemia, lymphatic, 111
Anemia of the spinal cord, 314
Anemia, pernicious, 107
Anemia, simple, 106
Anemia, splenic, 106
Anesthesia, 300
Anesthetic leprosy, 340
Aneurism, intracranial, 358
Aneurism of the aorta differentiated from—
abscess of the mediastinum, 142
aortic incompetency, 142
dilatation of the auricle, 143
intra-thoracic tumor, 142
pulsating pleurisy, 143
Aneurism, thoracic, 141
Angina, 204
Angina pectoris, 128
Angina pectoris differentiated from—
intercostal neuralgia, 128
tachycardia, 126
Angina, ulcero-membranous, 208
Angioneurotic edema, 146
Ankylostomiasis, 239
Anorexia, nervous, 214
Anthrax, 89
Aorta, aneurism of, 141, 142
Aortic incompetency or regurgitation, 133
Aortic incompetency or regurgitation differentiated from aneurism of the aorta, 142
Aortic obstruction, 133
Apex-beat of the heart, 118
Aphonia, hysterical, 151
Aphthous stomatitis, 200
Appendicitis, 228
Appendicitis, differential diagnosis of, 230
Arachnitis, 342
Ardent fever, 31
Arsenical poisoning differentiated from Asiatic cholera, 52
Arterial hypermyotrophy, 144
Arterio-capillary fibrosis, 143
Arthritic muscular atrophy, 326
Arthritis deformans, 102
Arthritis, rheumatoid, 102
Arthritis, syphilitic, 97
Ascaris lumbricoides, 237
Asiatic cholera, 50

Asphyxia differentiated from cerebral hemorrhage, 354
Asphyxia, local, 144
Astasia-abasia, 373
Asthenic bulbar paralysis, 324
Asthma, 199
Asthma differentiated from—
an asthmatoid condition, 198
chronic edema of the larynx, 152
laryngismus stridulus, 153
paralysis of the diaphragm, 198
whooping-cough, 198
Ataxia, 299, 300
Ataxia, Friedreich's, 331
Ataxia, locomotor, 327
Ataxia, vasomotor, 384
Ataxic paraplegia, 330
Ataxic paraplegia, hereditary, 331
Atelectasis differentiated from catarrhal pneumonia, 176
Atrophy, muscular, arthritic, 326
Atrophy, muscular, progressive, 320, 321
Atrophy, muscular, progressive, differentiated from—
acute anterior poliomyelitis, 323
cervical pachymeningitis, 313
chronic myelitis, 317
progressive neural muscular atrophy, 323
syringomyelia, 339
Atrophy, progressive neural muscular, 323
Atrophy, yellow, acute, of the liver, 248
Atrophy, yellow, acute, of the liver differentiated from—
catarrhal jaundice, 260
phosphorus-poisoning, 249
typhoid fever, 249
yellow fever, 66
Aural vertigo, 381
Auricle, dilatation of, 143
Auscultation, 162
Auscultation of the heart-sounds, 122
Auscultatory percussion, 162
Autumnal catarrh, 149

Banti's disease, 265
Basedow's disease, 381
Beriberi, 63, 303
Bile-duct, common, compression of, 260
Biliary acids in the urine, 271
Biliary calculi, 261
Biliary coloring-matter in urine, 271
Biliary passages, carcinoma of, 262
Biliary passages, occlusion of, 251
Black tongue, 203
Bladder, neoplasms of, 289
Blood, 104
Blood, constitution of, 104
Blood in disease, 113
Blood-casts of the uriniferous tubules, 278
Boas's test for hydrochloric acid, 212
Boettger's test for the presence of sugar in the urine, 285
Bone, disease of, differentiated from neuritis, 303
Bothriocephalus latus, 236
Bowel, invagination or intussusception of, 232
Bowel, perforation of, in typhoid fever, 40
Brachycardia, 127
Break-bone fever, 83
Breathing, superior intercostal, differentiated from paralysis of the phrenic nerve, 308
Bronchial glands, tuberculosis of, 158
Bronchiectasis, 173
Bronchiectasis differentiated from—
catarrhal pneumonia, 173
pulmonary abscess, 173
pulmonary gangrene, 173
pulmonary tuberculosis, 190
Bronchitis, acute, 170
Bronchitis, acute, differentiated from acute miliary tuberculosis, 171
Bronchitis, capillary, 174
Bronchitis, capillary, differentiated from—
acute miliary tuberculosis, 174
catarrhal pneumonia, 175
Bronchitis, chronic, differentiated from—
interstitial pneumonitis, 172
pulmonary tuberculosis, 172
Bronchitis, plastic or fibrinous, 172
Bronchitis, putrid, 172
Bronchophony, 164
Broncho-pneumonia, 175
Brow-ague, 59

Brushmakers' disease, 89
Bubonic plague, 62
Bulbar palsy, acute, 323
Bulbar paralysis, asthenic, 324
Bulbar paralysis, progressive, 321
Bulimia, 214

Cachexia, malarial, 58
Cachexia strumipriva, 386
Calculi, biliary, 261
Calculi, biliary, differentiated from—
carcinoma of the biliary passages, 262
catarrhal jaundice, 259
Calculus, renal, 291
Calculus, vesical, 289
Cancrum oris, 201
Capillary bronchitis, 174
Caput Medusæ, 252
Carcinoma of the biliary passages differentiated from biliary calculi, 262
Carcinoma of the cecum differentiated from—
perityphlitis, 231
typhlitis, 231
Carcinoma of the intestine, 233
Carcinoma of the liver, 254
Carcinoma of the liver differentiated from—
abscess of the liver, 251
amyloid disease of the liver, 255
carcinoma of the omentum, 255
carcinoma of the stomach, 256
Carcinoma of the omentum differentiated from—
carcinoma of the liver, 255
carcinoma of the stomach, 222
Carcinoma of the pancreas, 265
Carcinoma of the pancreas differentiated from carcinoma of the stomach, 222
Carcinoma of the stomach, 220
Carcinoma of the stomach differentiated from—
carcinoma of the liver, 256
carcinoma of the omentum, 222
carcinoma of the pancreas, 222
chronic gastritis, 222
sarcoma of the omentum, 222
ulceration of the stomach, 221
Carcinoma, pulmonary, differentiated from pulmonary tuberculosis, 180
Cardia, insufficiency of, 214
Cardia, spasm of, 214
Cardiac impulse, 118
Caries of the vertebræ differentiated from tumor of the spinal cord, 338
Casts of the uriniferous tubules, 278
Catarrh, acute nasal, 148
Catarrh, autumnal, 149
Catarrh, epidemic, 32
Catarrh, gastric, acute, 215
Catarrh, gastric, chronic, 217
Catarrh, gastro-intestinal, differentiated from pernicious malarial fever, 60
Catarrh, intestinal, acute, 223
Catarrh, intestinal, chronic, 225
Catarrhal croup, 154
Catarrhal fever, 32
Catarrhal jaundice, 259
Catarrhal pharyngitis, acute, 204
Catarrhal pneumonia, 175
Catarrhal stomatitis, 199
Cavity, pulmonary, differentiated from pneumothorax, 196
Cecum, carcinoma of, 231
Cerebellar tumor, 329, 331
Cerebral abscess, 349
Cerebral anemia, 347
Cerebral congestion, 352
Cerebral disease differentiated from epilepsy, 371
Cerebral disease, paralysis of, differentiated from the paralysis of hysteria, 374
Cerebral embolism, 355
Cerebral functions, alterations in, in disease of the nervous system, 302
Cerebral hemorrhage, 350
Cerebral hyperemia, 348
Cerebral membranes, hemorrhage into, 345
Cerebral meningitis, 342
Cerebral softening, 354
Cerebral thrombosis, 356
Cerebral tumor, 357
Cerebritis, 349
Cerebritis differentiated from—
cerebral abscess, 350
cerebral meningitis, 349
Cerebro-spinal fever, 46
Cerebro-spinal fever differentiated from—
cerebral meningitis, 344
tetanus, 48
torticollis, 49

Cerebro-spinal fever differentiated from—
typhoid fever, 50
typhus fever, 48
variola, 49
yellow fever, 49
Cerebro-spinal sclerosis, 332
Cervical pachymeningitis, 313
Chalk-stones, 101
Charbon, 89
Chickenpox, 79
Chickenpox differentiated from—
measles, 80
smallpox, 79
varioloid, 79
Chlorosis, 107
Cholangitis, 258
Cholecystitis, 258
Cholera Asiatica, 50
Cholera Asiatica differentiated from—
arsenical poisoning, 52
cholera morbus, 52, 225
Cholera differentiated from pernicious malarial fever, 60
Cholera infantum, 225
Cholera infectiosa, 50
Cholera morbus, 225
Cholera morbus differentiated from cholera Asiatica, 52, 225
Cholera nostras, 225
Chorea, 367
Chorea differentiated from—
epilepsy, 371
spasmodic tic, 369
Chorea, hereditary, 369
Chorea, Huntingdon's, 369
Chronic bronchitis, 171
Chronic dysentery, 227
Chronic edema of the larynx, 152
Chronic enteritis, 225
Chronic gastric catarrh, 217
Chronic gastritis, 217
Chronic gout, 101
Chronic herpes of throat, 200
Chronic interstitial nephritis, 293
Chronic intestinal catarrh, 225
Chronic laryngitis, 158
Chronic myelitis, 316
Chronic nephritis, 358
Chronic parenchymatous nephritis, 292
Chronic peritonitis, 243
Chronic pharyngitis, 205
Chronic pleurisy, 169
Chronic rheumatism, 99
Chronic tonsillitis, 208
Chronic tuberculosis of the larynx, 159
Chyluria, 278
Cirrhosis of the liver, 252
Cirrhosis of the liver differentiated from—
amyloid disease of the liver, 254
catarrhal jaundice, 260
chronic perihepatitis, 258
Clavus hystericus, 373
Colic, hepatic, intestinal and renal, differential diagnosis of, 262
Colic, intestinal, differentiated from acute peritonitis, 242
Colic, mucous, 224
Color of urine, 267
Coloring-matter, biliary, in urine, 271
Common membranous sore-throat, 208
Complications in disease, 22
Compression of the common bile-duct differentiated from catarrhal jaundice, 260
Compression, spinal, 336
Compression, spinal, differentiated from chronic myelitis, 337
Compression-myelitis differentiated from hemorrhage into the spinal cord, 337
Confluent smallpox, 76
Congenital spastic paraplegia, 346
Congestion, cerebral, differentiated from cerebral hemorrhage, 352
Congestion of the liver, 246
Congestion, pulmonary, differentiated from croupous pneumonia, 180
Constitution of urine, 267
Constitutional symptoms, 19
Consumption, galloping, 183, 188
Continued fever, 30
Convulsion, 300
Cord, spinal, anemia of, 314
Cord, spinal, hyperemia of, 315
Coryza, 148
Coryza differentiated from—
hay-fever, 149
influenza, 148
Coryza, idiosyncratic, 149
Coryza, periodic vasomotor, 149
Cramp, writers', 367
Cretinism, 384

Cretinoidism, sporadic, 385
Croup, catarrhal, 154
Croup, catarrhal, differentiated from laryngismus stridulus, 154
Croup differentiated from—
edema of the larynx, 154
obstruction of the air-passages, 155
Croup, membranous, 155
Croup, membranous, differentiated from—
catarrhal croup, 156
diphtheria, 87
retro-pharyngeal abscess, 156
spasmodic croup, 156
whooping-cough, 157
Croup, spasmodic, 154
Croup, spasmodic, differentiated from—
edema of the larynx, 154
laryngismus stridulus, 154
membranous croup, 156
Croupous pneumonia, 176
Cystitis, 274, 288
Cystitis differentiated from—
nephritis, 288
pyelitis, 290
Cysts, pancreatic, 266

DATA for diagnosis, 24
Deep reflexes, 301
Degeneration, fatty, of the heart, 130
Degeneration of the diaphragm, 308
Degeneration, reaction of, 301
Delirium tremens, 363
Delirium tremens differentiated from—
acute mania, 363
cerebral meningitis, 363
Dementia, paretic, 359
Dengue, 83
Dengue differentiated from—
influenza, 84
scarlet fever, 84
yellow fever, 66
Dermatitis, acute exfoliative, differentiated from scarlet fever, 73
Dermatitis, toxic, differentiated from scarlet fever, 73
Deuteropathic fever, 30
Dextrocardia, 125
Diabetes insipidus, 282
Diabetes insipidus differentiated from—
chronic interstitial nephritis, 283
diabetes mellitus, 287
hysterical polyuria, 283
Diabetes mellitus, 283
Diabetes mellitus differentiated from—
diabetes insipidus, 287
Morvan's disease, 342
Diacetic acid in the urine, tests for the presence of, 287
Diaceturia, 287
Diagnosis, 17
Diagnosis by exclusion, 21
Diagnosis by the historical or empirical method, 20
Diagnosis by the inductive method, 20
Diagnosis by the method of pathological association, 20
Diagnosis, data for, 24
Diagnosis, direct, 20
Diagnosis, discriminative or differential, 21
Diagnosis, physical, 159
Diaphragm, degeneration of, differentiated from paralysis of the phrenic nerve, 308
Diaphragm, inflammation of, differentiated from paralysis of the phrenic nerve, 308
Diaphragm, paralysis of, 198
Diaphragmatic hernia, 196
Diarrhea, tubular, 224
Differential diagnosis, 21
Digestive activity of the stomach, 213
Digestive system, 199
Dilatation of the auricle differentiated from aneurism of the aorta, 143
Dilatation of the heart, 129
Dilatation of the heart differentiated from—
fatty degeneration of the heart, 130
hypertrophy of the heart, 130
pericardial effusion, 130
Dilatation of the stomach, 218
Diphtheria, 84
Diphtheria differentiated from common membranous sore-throat, 209

Diphtheria differentiated from—
gangrenous sore-throat, 209
membranous croup, 87
scarlet fever, 86
stomatitis, 87
tonsillitis, 86
Diphtheroid throat, 208
Diplegia, 299
Direct diagnosis, 20
Discrete smallpox, 76
Discriminative diagnosis, 21
Displacements of the kidney, 290
Disseminated neuritis, 303
Disseminated sclerosis, 332
Distention of the gall-bladder differentiated from hydatid cyst of the liver, 258
Distomiasis, 239
Dracontiasis, 95
Drug-effects, 23
Dumb ague, 58
Duplication of the heart-sounds, 120
Dysentery, acute, 226
Dysentery, acute, differentiated from—
acute enteritis, 227
intussusception of the bowel, 226
typhoid fever, 227
Dysentery, chronic, 227
Dystrophy, progressive muscular, 324

Echinococcus disease, 92
Eclampsia differentiated from epilepsy, 372
Edema, angioneurotic, 146
Edema of the larynx, acute, 151
Edema of the larynx, chronic, 152
Edema of the larynx, chronic, differentiated from asthma, 152
Edema of the larynx differentiated from spasmodic croup, 154
Edema, pulmonary, differentiated from pneumonia, 179
Efficiency of the heart, 121
Effusion, pericardial, differentiated from—
dilatation of the heart, 130
pericarditic effusion, 145
pleuritic effusion, 162
Effusion, pleural, differentiated from—
pericardial effusion, 169
pneumothorax, 197
Effusion, pleural, differentiated from pulmonary emphysema, 195
Effusion, pleuritic, differentiated from—
abscess of the liver, 168
abscess of the spleen, 168
hydatid cyst of the liver, 167
hydrothorax, 169
Egophony, 165
Electric reactions, alterations in, in disease, 301
Embolism, cerebral, 355
Embolism, cerebral, differentiated from cerebral hemorrhage, 355
Emphysema, pulmonary, 194
Emphysema, pulmonary, differentiated from—
pleural effusion, 195
pneumothorax, 194
Empirical method, diagnosis by, 20
Emprosthotonos, 377
Empyema, 165
Empyema, pulsatory, 166
Endemic hematuria, 282
Endocardial murmurs, 136
Endocarditis, 139
Endocarditis, acute, differentiated from—
acute pericarditis, 140
valvular disease of the heart, 139
Endocarditis, malignant or ulcerative, 139
Endocarditis, simple or vegetative, 139
Enlargement of the spleen, 264
Enteric fever, 35
Enteritis, acute, 223
Enteritis, acute, differentiated from—
acute dysentery, 227
acute peritonitis, 242
Enteritis, catarrhal, acute, differentiated from typhoid fever, 224
Enteritis, chronic, 225
Enteritis, membranous, 224
Enteritis, mucous, acute, 223
Ephemeral fever, 31
Epidemic catarrh, 32
Edidemic cerebro-spinal meningitis, 46
Epilepsy, 370
Epilepsy differentiated from aural vertigo, 381

Epilepsy differentiated from—
cerebral disease, 371
cerebral tumor, 358
chorea, 371
eclampsia, 372
hysteria, 374
syncope, 371
uremia, 294, 372
Epilepsy, essential, 370
Epilepsy, idiopathic, 370
Epilepsy, major, 370
Epilepsy, minor, 370
Epistaxis, 148
Epithelial casts of the uriniferous tubules, 278
Equinia, 87
Eructations, nervous, 214
Eruptive fevers, 68
Erysipelas, 80
Erysipelas differentiated from—
erythema, 81
herpes zoster, 81
scarlet fever, 81
smallpox, 82
Erythema differentiated from erysipelas, 81
Erythromelalgia, 145
Erythromelalgia differentiated from Raynaud's disease, 145
Esophagismus, 212
Esophagitis, 211
Esophagus, 211
Esophagus, inflammation of, 211
Esophagus, stricture of, 211
Essential epilepsy, 370
Essential fever, 30
Examination of urine, 267
Exanthemata, 68
Exclusion, diagnosis by, 21
Exophthalmic goiter, 381
Exophthalmic goiter differentiated from—
simple goiter, 384
tachycardia, 126

Facial hemiatrophy, 306
Facial nerve, paralysis of, 306
Family history, 19
Family periodic paralysis, 377
Famine-fever, 44
Farcy, 87
Fatty casts of the uriniferous tubules, 278
Fatty degeneration of the heart differentiated from dilatation of the heart, 130
Fatty infiltration of the liver, 253
Fatty liver, 253
Fatty metamorphosis of the liver, 253
Febricula, 31
Febrile temperature, 28
Fehling's test for the presence of sugar in the urine, 286
Fermentation-test for the presence of sugar in the urine, 286
Fever, 27
Fever, ardent, 31
Fever, break-bone, 83
Fever, catarrhal, 32
Fever, cerebro-spinal, 46
Fever, continued, 30
Fever, deuteropathic, 30
Fever, enteric, 35
Fever, ephemeral, 31
Fever, essential, 30
Fever, glandular, 82
Fever, hemorrhagic malarial, 58
Fever, hepatic intermittent, 262
Fever, idiopathic, 30
Fever, intermittent, 30
Fever, malarial intermittent, 56
Fever, malarial remittent, 57
Fever, Malta, 61
Fever, Mediterranean, 61
Fever, miliary, 82
Fever, Neapolitan, 61
Fever of suppuration, 59
Fever, pernicious malarial, 58
Fever, rag-weed, 149
Fever, relapsing, 52
Fever, remittent, 30
Fever, rheumatic, 95
Fever, scarlet, 70
Fever, simple continued, 31
Fever, specific, 30
Fever, splenic, 89
Fever, spotted, 47
Fever, symptomatic, 30
Fever, syphilitic, 61
Fever, thermic, 361
Fever, typhoid, 35
Fever, typhus, 44
Fever, undulant, 61
Fever, yellow, 64
Fevers, eruptive, 68
Fibrinous bronchitis, 172

Fibroid phthisis, 183
Fibrosis, arterio-capillary, 143
Filariasis, 95, 279
Floating kidney, 291
Floating liver, 245
Floating spleen, 264
Florid phthisis, 183, 188
Follicular tonsillitis, 207
Foot-and-mouth disease, 91
Fremitus, vocal or tactile, 160
French measles, 74
Frequency of action of the heart, 120
Friedreich's ataxia, 331
Frost-bite differentiated from Raynaud's disease, 145
Functional heart-murmurs, 124
Functional stricture of the esophagus, 212

GALL-BLADDER, distention of, 258
Galloping consumption, 183, 188
Gangrene, local or symmetrical, 144
Gangrene, pulmonary, 182
Gangrene, pulmonary, differentiated from—
bronchiectasis, 173
pulmonary abscess, 182
pulmonary tuberculosis, 182
Gangrene, symmetrical, 144
Gangrenous pharyngitis, 209
Gangrenous stomatitis, 201
Gastralgia, 215
Gastralgia differentiated from gastric ulceration, 220
Gastrectasis, 218
Gastric catarrh, acute, 215
Gastric catarrh, chronic, 217
Gastric contents, acidity of, 212
Gastric contents, hydrochloric acid in, 212
Gastric contents, lab-ferment in, 213
Gastric contents, lactic acid in, 213
Gastric contents, pepsin in, 213
Gastric contents, rennet-ferment in, 213
Gastric contents, total acidity of, 212
Gastric hemorrhage, 189
Gastric mucous membrane, absorptive activity of, 214
Gastric ulcer, 218
Gastric ulceration differentiated from gastralgia, 220
Gastritis, acute, 215
Gastritis, acute, differentiated from—
acute peritonitis, 241
intestinal obstruction, 216
the gastric symptoms of acute febrile disease or of cerebral disease, 216
Gastritis, chronic, 217
Gastritis, chronic, differentiated from—
carcinoma of the stomach, 222
gastric ulcer, 219
Gastrodynia, 214
Gastro-enteritis, 223
Gastro-intestinal catarrh, 60
Gastroptosis, 218
General paralysis of the insane, 359
General symptoms, 19
Genito-urinary apparatus, 267
Geographical tongue, 203
German measles, 74
Glanders, 87
Glanders differentiated from—
smallpox, 88
syphilis, 88
Glandular fever, 82
Globus hystericus, 373
Glossanthrax, 203
Glossitis, 202
Glosso-labio-laryngeal paralysis, 320, 321
Glycosuria, 283
Goiter, exophthalmic, 381
Goiter, simple, differentiated from exophthalmic goiter, 334
Gonorrheal synovitis, 97
Gout, acute, 99
Gout, acute, differentiated from acute rheumatism, 100
Gout, chronic, 102
Gout differentiated from rheumatoid arthritis, 104
Grand mal, 370
Granular casts of the uriniferous tubules, 278
Graves's disease, 381
Green-sickness, 107
Grippe, la, 32
Günzburg's test for hydrochloric acid, 212

HABIT-CHOREA, 369
Habit-spasm, 369
Hairy tongue, 203

Hay-asthma, 149
Hay-fever, 149
Hay-fever differentiated from coryza, 149
Headache differentiated from migraine, 312
Headache, sick, 311
Heart, 118
Heart, action of, 120
Heart, action of, intermission of, 120
Heart, apex-beat of, 118
Heart, dilatation of, 129
Heart, efficiency of, 121
Heart, fatty degeneration of, 130
Heart, functional disturbance of, 125
Heart, hypertrophy of, 128
Heart, irritable, 127
Heart, malformation of, 124
Heart, valvular disease of, 131
Heart-clot, 140
Heart-murmurs, 121
Heart-murmurs, organic, differentiated from functional, 124
Heart-sounds, auscultation of, 120
Heat-exhaustion, 36
Heat-exhaustion differentiatedfrom heat-fever or sunstroke, 362
Heat-fever, 361
Heat-stroke, 361
Hematokrit, 104
Hematuria, 279
Hematuria, endemic, 282
Hematuria, malarial, differentiated from paroxysmal hemoglobinuria, 281
Hemiatrophy, facial, 306
Hemicrania, 311
Hemiplegia, 299
Hemocytometry, 104
Hemoglobinometry, 105
Hemoglobinuria, 280
Hemoglobinuria, paroxysmal, 281
Hemoglobinuria, paroxysmal, differentiated from malarial hematuria, 281
Hemophilia, 116
Hemorrhage, cerebral, 353
Hemorrhage, cerebral, differentiated from—
alcoholic intoxication, 353
asphyxia, 354
cerebral congestion, 352
cerebral embolism, 355
cerebral thrombosis, 356
Hemorrhage, cerebral, differentiated from—
opium-poisoning, 353
primary lateral sclerosis, 330
sunstroke, 362
syncope, 353
uremia 354
Hemorrhage, gastric, differentiated from pulmonary hemorrhage, 189
Hemorrhage into the cerebral membranes, 345
Hemorrhage into the spinal membranes, 314
Hemorrhage into the spinal membranes differentiated from—
spinal hemorrhage, 335
spinal meningitis, 314
tetanus, 378
Hemorrhage, pancreatic, 266
Hemorrhage, pulmonary, differentiated from gastric hemorrhage, 189
Hemorrhage, spinal, 335
Hemorrhage, spinal, differentiated from—
acute myelitis, 316
compression-myelitis, 337
meningeal hemorrhage, 335
spinal meningitis, 335
Hemorrhagic malarial fever, 58
Hemorrhagic pachymeningitis, 383
Hemorrhagic smallpox, 77
Hepatic colic, 262
Hepatic intermittent fever differentiated from malarial intermittent fever, 262
Hepatitis, acute, 247
Hepatitis, acute, differentiated from—
malarial cachexia, 247
portal phlebitis, 247
Hepatitis, interstitial, 252
Hereditary ataxic paraplegia, 331
Hereditary chorea, 369
Hernia, diaphragmatic, differentiated from pneumothorax, 196
Herpes, chronic or recurrent, of throat, 208
Herpes of the pharynx, 208
Herpes zoster differentiated from erysipelas, 81
Herpetic sore-throat, 208
Herpetic tonsillitis, 208
Hip-joint, disease of, differentiated from sciatica, 305
Hippuric acid in urine, 269

Historical method, diagnosis by, 20
History, family, 19
History, previous, 19
Hodgkin's disease, 111
Humanized virus, 77
Huntingdon's chorea, 369
Hyaline casts of the uriniferous tubules, 278
Hydatid cyst, 93
Hydatid cyst of the kidney, 298
Hydatid cyst of the kidney differentiated from hydronephrosis, 298
Hydatid cyst of the liver, 256
Hydatid cyst of the liver differentiated from—
abscess of the liver, 257
distention of the gall-bladder, 258
pleuritic effusion, 167
Hydatid disease, 92
Hydremia, 105
Hydrocephalus, 345
Hydrochloric acid in the gastric contents, 212
Hydrochloric acid, tests for, 212
Boas's test for, 212
Günzburg's test for, 212
Töpfer's test for, 212
Hydronephrosis, 297
Hydronephrosis differentiated from cyst of the kidney, 298
Hydrophobia, 380
Hydrophobia differentiated from—
pseudo-hydrophobia, 380
tetanus, 381
Hydrothorax differentiated from pleuritic effusion, 169
Hyperacidity, 215
Hyperchlorhydria, 215
Hyperemia, cerebral, 348
Hyperemia of the spinal cord, 315
Hyperesthesia, 300
Hyperkinesis, 214
Hyperleukocytosis, 108
Hypermyotrophy, arterial, 144
Hyperpyrexia, 28
Hypertrophic pachymeningitis, 338, 339
Hypertrophy of the heart, 128
Hypertrophy of the heart differentiated from dilatation of the heart, 130
Hypesthesia, 300
Hypoleukocytosis, 105
Hysteria, 372
Hysteria differentiated from—
acute meningitis, 375
cerebral tumor, 375
epilepsy, 374
neurasthenia, 376
tetanus, 378
Hysterical aphonia differentiated from acute laryngitis, 151
Hysterical paralysis differentiated from the paralysis of cerebral disease, 374
Hysterical paraplegia differentiated from—
acute myelitis, 375
spastic paraplegia, 375
Hysterical polyuria, 283

Idiopathic anemia, 107
Idiopathic epilepsy, 370
Idiopathic fever, 30
Idiosyncratic coryza, 149
Immediate percussion, 161
Impulse, cardiac, 118
Inadequacy, renal, 290
Incompetency, aortic, 133
Incompetency, aortic, differentiated from aneurism of the aorta, 142
Incompetency, mitral, 132
Incompetency, pulmonary, 135
Incompetency, tricuspid, 134
Indicanuria, 273
Indigestion, acute, 215
Inductive method, diagnosis by, 20
Infantile pseudo-leukemic anemia, 112
Infantile spastic paraplegia, 346
Infarction, pulmonary, differentiated from—
catarrhal pneumonia, 176
croupous pneumonia, 180
pulmonary abscess, 181
Inflammation of the diaphragm, 306
Inflammation of the esophagus, 211
Influenza, 32
Influenza, abdominal type of, 34
Influenza, catarrhal type of, 33
Influenza, cerebral type of, 34
Influenza, diagnosis of, 35
Influenza differentiated from—
coryza, 148
dengue, 84
Influenza, gastrointestinal type of, 34

Influenza, nervous type of, 34
Influenza, thoracic type of, 33
Influenza, typhoid type of, 34
Inquiry, 24
Insane, general paralysis of, 359
Insolation, 361
Inspection, 118, 146, 159
Instrumental signs, 18
Insufficiency of the cardia, 214
Insufficiency of the pylorus, 214
Insular sclerosis, 332
Intercostal neuralgia, 128, 167
Intermission of the heart's action, 120
Intermittent fever, 30
Intermittent fever, hepatic, 262
Intermittent fever, malarial, 56
Interstitial hepatitis, 252
Interstitial nephritis, chronic, 293
Interstitial pneumonitis, 193
Intestinal catarrh, acute, 223
Intestinal catarrh, chronic, 225
Intestinal colic, 242, 262
Intestinal obstruction, 216, 231
Intestinal parasites, 234
Intestines, 223
Intestines, carcinoma of, 233
Intoxication, alcoholic, 353
Intracranial aneurism, 359
Intra-thoracic tumor, 142, 170
Intussusception of the bowel, 232
Intussusception of the bowel differentiated from—
acute dysentery, 226
obstruction of the bowel, 233
typhlitis, 233
Invagination of the bowel, 232
Inverse temperature, 28
Irritable heart, 127
Irritable heart differentiated from tachycardia, 127

JAIL-FEVER, 44
Jaundice, acute infective, 67
Jaundice, catarrhal, differential diagnosis of, 259
acute yellow atrophy of the liver, 260
Jaundice, catarrhal, differentiated from—
biliary calculus, 259
cirrhosis of the liver, 260
compression of the common bile-duct, 260
malarial fever, 260
Jaundice, catarrhal, differentiated from—
malignant disease, 260
pylephlebitis, 260
Weil's disease, 260
yellow fever, 260
June-cold, 149

KAK-KE, 63, 303
Kidney, abscess of, 295
Kidney, amyloid, lardaceous or waxy, 293
Kidney, displacements of, 290
Kidney, floating, 291
Kidney, hydatid cyst of, 298
Kidney, malignant disease of, 297
Kidney, movable, 291
Kidney, tuberculosis of, 296

LAB-FERMENT in gastric contents, 213
Labyrinthine vertigo, 381
Lactic acid, test for, 213
Lacunal tonsillitis, 207
Landry's paralysis, 319
Lardaceous kidney, 293
Larval paludism, 59
Laryngeal neoplasms differentiated from chronic laryngitis, 158
Laryngeal vertigo, 153
Laryngismus stridulus, 153
Laryngismus stridulus differentiated from—
asthma, 153
catarrhal croup, 154
whooping-cough, 157
Laryngitis, acute, 150
Laryngitis, acute, differentiated from hysterical aphonia, 150
Laryngitis, acute tuberculous, 152
Laryngitis, chronic, 158
Laryngitis, chronic, differentiated from laryngeal neoplasms, 158
Laryngitis differentiated from—
parotiditis, 150
pharyngitis, 150
tonsillitis, 150
Laryngitis, pseudo-membranous, 155
Laryngitis, tuberculous, acute, 152
Laryngoscopy, 147
Larynx, edema of, 151
Larynx, tuberculosis of, chronic, 159

Lateral sclerosis, amyotrophic, 320, 321
Lateral sclerosis, primary, 329
Lead-poisoning, 364
Lead-poisoning differentiated from cerebral tumor, 365
Lead-poisoning, neuritis of, differentiated from neuritis of the musculo-spiral nerve, 309
Leprosy, 63
Leprosy, anesthetic, differentiated from Morvan's disease, 341
Leprosy differentiated from syringomyelia, 339
Leptomeningitis, 312, 342
Leucin, 272
Leukemia, 105, 109
Leukemia differentiated from pseudo-leukemia, 111
Leukocythemia, 105, 109
Leukocytosis, 108
Leukoplakia lingualis s. buccalis, 202
Lipuria, 279
Lithemia, 101
Lithemia, diagnosis of, 101
Liver, 244
Liver, abscess of, 168, 250
Liver, actinomycosis of, 251
Liver, amyloid disease of, 254
Liver, atrophy of, acute yellow, 248
Liver, carcinoma of, 254
Liver, cirrhosis of, 252
Liver, congestion of, 246
Liver, fatty, 253
Liver, floating, 245
Liver, hydatid cyst of, 167, 256
Liver, malignant disease of, differentiated from chronic peritonitis, 244
Lobar pneumonia, 176
Lobular pneumonia, 175
Local asphyxia, 144
Local gangrene, 144
Local symptoms, 19
Locomotor ataxia, 327
Lumbago, 98
Lumbar pachymeningitis, 328
Lumbricoid worms, 237
Lung-fever, 176
Lungs, new-growths in, 199
Lymphadenoma, 111
Lymphatic anemia, 111
Lymph-glands, tuberculosis of, 111
Lymphoma, malignant, 111
Lyssophobia, 380

Major epilepsy, 370
Malaria, masked, 59
Malarial cachexia, 58
Malarial cachexia differentiated from acute hepatitis, 244
Malarial diseases, 54
Malarial fever differentiated from—
  catarrhal jaundice, 260
  Malta fever, 62
  pulmonary tuberculosis, 191
  syphilitic fever, 61
  the fever of suppuration or of septic infection, 59
  typhoid fever, 61
Malarial fever, hemorrhagic, 58
Malarial fever, hemorrhagic, differentiated from—
  paroxysmal hemoglobinuria, 60
  yellow fever, 65
Malarial fever, pernicious, 58
Malarial fever, pernicious, diagnosis of, 60
Malarial fever, varieties of, 55
Malarial hematuria, 281
Malarial intermittent fever, 56
Malarial intermittent fever differentiated from hepatic intermittent fever, 262
Malarial paroxysm, 56
Malarial remittent fever, 57
Malarial remittent fever differentiated from yellow fever, 65
Malformation of the heart, 124
Malignant disease differentiated from catarrhal jaundice, 260
Malignant disease of the kidney, 297
Malignant disease of the liver, 244
Malignant endocarditis, 139
Malignant lymphoma, 111
Malignant pustule, 89
Malignant smallpox, 77
Mania, acute, differentiated from—
  cerebral meningitis, 344
  delirium tremens, 363
Masked malaria, 59
Malta fever, 61
Malta fever differentiated from—
  malarial fever, 62
  typhoid fever, 62
Measles, 69
Measles differentiated from chicken-pox, 80

Measles differentiated from—
purpura, 115
rubella, 75
scarlet fever, 73
smallpox, 78
typhus fever, 70
Measles, French, 74
Measles, German, 74
Mediastinum, abscess of, 142
Mediastinum, abscess of, differentiated from aneurism of the aorta, 144
Mediastinum, diseases of, 141
Mediate percussion, 161
Mediterranean fever, 61
Megagastria, 218
Megalocytes, 105
Membranes, spinal, hemorrhage into, 314
Membranous croup, 155
Membranous enteritis, 224
Membranous sore-throat, common, 208
Ménière's disease, 381
Meningitis, acute, differentiated from—
hysteria, 375
sunstroke, 362
Meningitis, cerebral, 342
Meningitis, cerebral, differentiated from—
acute general disease, 343
acute mania, 344
cerebral abscess, 350
cerebritis, 349
cerebro-spinal fever, 344
delirium tremens, 363
Meningitis, cerebral, simple, differentiated from tuberculous meningitis, 343
Meningitis, epidemic cerebro-spinal, 46
Meningitis, spinal, 312
Meningitis, spinal, acute, differentiated from—
acute myelitis, 314
tetanus, 378
Meningitis, spinal, differentiated from—
hemorrhage into the spinal meninges, 314
rheumatism of the muscles of the back, 313
spinal hemorrhage, 335
Meningitis, tuberculous, differentiated from simple cerebral meningitis, 343
Mensuration, 160
Mercurial stomatitis, 201
Merycism, 214
Mesenteric vessels, occlusion of, 240
Microcytes, 105
Migraine, 311
Migraine differentiated from ordinary headache, 312
Miliaria, 82
Miliary fever, 82
Miliary tuberculosis, acute, 191
Milk-sickness, 91
Minor epilepsy, 370
Mitral incompetency or regurgitation, 132
Mitral obstruction, 132
Mollities ossium, 118
Mollities ossium differentiated from rachitis, 118
Monoplegia, 299, 300
Moore's test for the presence of sugar in the urine, 285
Morbilli, 69
Morvan's disease, 340
Morvan's disease differentiated from—
anesthetic leprosy, 341
diabetes, 342
Raynaud's disease, 341
scleroderma, 340
syphilis, 341
Motion, 299
Motor activity of the stomach, 214
Mouth, 199
Movable kidney, 291
Mucous colic, 224
Mucous enteritis, 223
Muguet, 200
Multiple neuritis, 303
Multiple sclerosis, 332
Mumps, 204
Murmurs, endocardial, 136
Murmurs, organic, differentiated from functional heart-murmurs, 124
Muscular atrophy, arthritic, 326
Muscular atrophy, progressive, 320, 321
Muscular atrophy, progressive neural, 323
Muscular dystrophy, progressive, 324
Muscular rheumatism, 99

Musculo-spiral nerve, paralysis of, 309
Musculo-spiral neuritis differentiated from the neuritis of lead-poisoning, 309
Myalgia, 98
Myelitis, 315
Myelitis, acute, differentiated from—
acute anterior poliomyelitis, 318
acute ascending paralysis, 319
acute spinal meningitis, 316
chronic myelitis, 317
hysterical paraplegia, 375
multiple neuritis, 304
spinal hemorrhage, 316
Myelitis, chronic, 316
Myelitis, chronic, differentiated from—
acute myelitis, 317
compression of the cord, 337
lateral sclerosis, 317
primary lateral sclerosis, 329
progressive muscular atrophy, 317
spinal pachymeningitis, 317
syringomyelia, 339
tumor of the spinal cord, 338
Myocardium, diseases of, 138
Myotonia congenita, 327
Myxedema, 385
Myxedema differentiated from—
adiposis, 387
akromegaly, 389
obesity, 387
scleroderma, 386

Nasal catarrh, acute, 148
Nasal respiration, difficulty of, 147
Nasal septum, ulceration or erosion of, 148
Neapolitan fever, 61
Negative signs, 23
Neoplasms, laryngeal, 158
Neoplasms of the bladder, 289
Nephritis, acute, 292
Nephritis chronic, differentiated from cerebral tumor, 358
Nephritis, chronic interstitial, 293
Nephritis, chronic interstitial, differentiated from diabetes insipidus, 283
Nephritis, chronic parenchymatous, 292
Nephritis differentiated from cystitis, 288
Nervous anorexia, 214
Nervous eructations, 214
Nervous vomiting, 214
Neural muscular atrophy, progressive, 323
Neuralgia, 310
Neuralgia differentiated from—
neuritis, 303, 310
subacute rheumatism, 98
Neuralgia, intercostal, differentiated from—
acute pleurisy, 167
angina pectoris, 128
Neuralgia of the stomach, 215
Neuralgia, rheumatic, 99
Neuralgia, sciatic, differentiated from sciatica, 305
Neuralgia, trifacial or trigeminal, 311
Neurasthenia, 375
Neurasthenia differentiated from hysteria, 376
Neuritis, 302
Neuritis, diagnosis of, 303
Neuritis differentiated from—
disease of bone, 203
neuralgia, 303, 310
subacute rheumatism, 303
Neuritis, multiple or disseminated, 303
Neuritis, multiple, differentiated from—
acute anterior poliomyelitis, 319
acute ascending paralysis, 319
acute myelitis, 304
posterior spinal sclerosis, 328
spinal pachymeningitis, 304
Neuritis, musculo-spiral, 809
Neuromata, 310
Neuromata, diagnosis of, 310
Neuroses of the stomach, 214
New-growths in the lungs, 199
Nigrities, 203
Noma, 201
Nutrition, alterations in, in disease of the nervous system, 301

Obesity, 387
Objective symptoms, 17
Observation, 24
Obstruction, aortic, 133
Obstruction, intestinal, 231
Obstruction, intestinal, acute, dif-

ferentiated from acute peritonitis, 242
Obstruction, intestinal, differentiated from acute gastritis, 216
Obstruction, mitral, 132
Obstruction of the air-passages, 155
Obstruction of the bowel differentiated from intussusception, 233
Obstruction, pulmonary, 135
Obstruction, tricuspid, 135
Occlusion of the biliary passages differentiated from abscess of the liver, 251
Occlusion of the mesenteric vessels, 240
Occupation-neuroses, 366
Odor of urine, 268
Oligemia, 105
Oligochromemia, 105
Oligocythemia, 105
Omentum, carcinoma of, 222, 255
Omentum, sarcoma of, 222
Opisthotonos, 377
Opium-poisoning differentiated from cerebral hemorrhage, 353
Organic heart-murmurs, 124
Organic stricture of the esophagus, 212
Orthotonos, 377
Osteitis deformans, 389
Osteo-arthropathy, hypertrophic, pulmonary, 391
Ovary, right, abscess of, 230
Oxaluria, 273
Oxyuris vermicularis, 239

Pachymeningitis, cerebral, 342, 343
Pachymeningitis, cervical, 313
Pachymeningitis, cervical, differentiated from—
progressive muscular atrophy, 313
subacute anterior poliomyelitis, 313
Pachymeningitis, hemorrhagic, 342, 343
Pachymeningitis, hypertrophic, differentiated from—
syringomyelia, 339
tumor of the spinal cord, 338
Pachymeningitis, lumbar, differentiated from posterior spinal sclerosis, 328
Pachymeningitis, spinal, differentiated from—
chronic myelitis, 317
multiple neuritis, 304
Paget's disease, 389
Palpation, 119, 160
Palsy, bulbar, acute, 323
Palsy, bulbar, acute, distinguished from chronic bulbar palsy, 323
Palsy, bulbar, differentiated from pseudo-bulbar palsy, 324
Palsy, pseudo-bulbar, 324
Palsy, pseudo-bulbar, differentiated from true bulbar palsy, 324
Palsy, shaking, 333
Paludism, larval, 59
Panaris, analgesic, 340
Pancreas, 265
Pancreas, carcinoma of, 222, 265
Pancreatic cyst, 266
Pancreatic hemorrhage, 266
Pancreatitis, acute, 265
Paralysis, 299
Paralysis agitans, 333
Paralysis agitans differentiated from cerebro-spinal sclerosis, 334
Paralysis, ascending, acute, 319
Paralysis, ascending, acute, differentiated from—
acute anterior poliomyelitis, 320
acute myelitis, 319
multiple neuritis, 319
Paralysis, asthenic, bulbar, 324
Paralysis, bulbar, progressive, 321
Paralysis, cerebral, differentiated from acute anterior poliomyelitis, 318
Paralysis, family periodic, 377
Paralysis, general, of the insane, 359
Paralysis, general, of the insane, differentiated from posterior spinal sclerosis, 360
Paralysis, glosso-labio-laryngeal, 320, 321
Paralysis, hysterical, 374
Paralysis, Landry's, 319
Paralysis of the diaphragm differentiated from—
asthma, 198
paralysis of the phrenic nerve, 309
Paralysis of the facial nerve, 306
Paralysis of the musculo-spiral nerve, 309
Paralysis of the phrenic nerve, 308

Paralysis, pseudo-hypertrophic, differentiated from congenital spastic paraplegia, 347
Paraplegia, 299, 300
Paraplegia, ataxic, 330
Paraplegia, ataxic, hereditary, 331
Paraplegia, ataxic, hereditary, differentiated from postero-lateral sclerosis, 331
Paraplegia, hysterical, 375
Paraplegia, spastic, 329
Paraplegia, spastic, congenital, 346
Paraplegia, spastic, congenital, differentiated from pseudo-hypertrophic paralysis, 347
Paraplegia, spastic, differentiated from hysterical paraplegia, 375
Parasites, intestinal, 234
Parasitic stomatitis, 200
Parenchymatous tonsillitis, 206
Paresis, 299
Paresthesia, 300
Paretic dementia, 359
Parkinson's disease, 333
Parotiditis, 204
Parotiditis differentiated from laryngitis, 150
Paroxysmal hemoglobinuria, 281
Paroxysmal hemoglobinuria, differentiated from hemorrhagic malarial fever, 60
Pathological association, diagnosis by the method of, 20
Pectoriloquy, 164
Pectoriloquy, whispering, 164
Pepsin in gastric contents, 213
Percussion, 119, 161
Percussion, auscultatory, 162
Percussion, immediate, 161
Percussion, mediate, 161
Percussion, respiratory, 162
Perforation of the bowel in typhoid fever, 40
Pericardial effusion differentiated from—
dilatation of the heart, 130
pericarditic effusion, 140
pleural effusion, 169
Pericarditis, acute, 135
Pericarditis, acute, differentiated from—
acute endocarditis, 140
acute pleurisy, 137
Perihepatitis, 258
Perihepatitis, chronic, differentiated from cirrhosis of the liver, 258
Perinephric abscess, 296
Periodic paralysis, family, 377
Periodic vasomotor coryza, 149
Peristaltic unrest, 214
Peritonitis, acute, 241
Peritonitis, acute, differentiated from—
acute enteritis, 242
acute gastritis, 241
acute intestinal obstruction, 242
intestinal colic, 242
subacute rheumatism, 243
Peritonitis, chronic, 243
Peritonitis, chronic, differentiated from malignant disease of the liver, 244
Perityphlitis, 229
Perityphlitis, differential diagnosis of, 230
Perityphlitis differentiated from—
abscess of the right ovary, 230
carcinoma of the cecum, 231
lumbar abscess, 230
Pernicious anemia, 107
Pernicious malarial fever, 58
Pernicious malarial fever, diagnosis of, 58
Pertussis, 157
Petit mal, 370
Pharyngitis, 204
Pharyngitis, acute catarrhal, 204
Pharyngitis, acute phlegmonous, 204
Pharyngitis, acute tuberculous, 205
Pharyngitis, chronic, 205
Pharyngitis differentiated from laryngitis, 150
Pharyngitis, gangrenous, 209
Pharyngitis, tuberculous, differentiated from—
syphilitic sore-throat, 206
typhoid fever, 206
Pharyngoscopy, 147
Pharynx, 204
Pharynx, herpes of, 208
Phlebitis, portal, differentiated from acute hepatitis, 247
Phlegmonous pharyngitis, acute, 204
Phosphoric acid in urine, 270
Phosphorus-poisoning differentiated from acute yellow atrophy of the liver, 249

Phrenic nerve, paralysis of, 308
Phrenic nerve, paralysis of, differentiated from—
degeneration of the diaphragm, 308
inflammation of the diaphragm, 308
superior intercostal breathing, 308
Phthisis, fibroid, 183
Phthisis, florid, 183, 188
Physical diagnosis, 159
Physical signs, 18
Pia-arachnitis, 342
Plague, bubonic, 62
Plague differentiated from—
typhus fever, 63
yellow fever, 63
Plastic bronchitis, 172
Pleural effusion, 169, 195, 197
Pleurisy, acute, 165
Pleurisy, acute, differentiated from—
acute pericarditis, 137
acute pneumonia, 167
intercostal neuralgia, 167
Pleurisy, chronic, 170
Pleurisy, chronic, differentiated from—
interstitial pneumonitis, 193
intrathoracic tumor, 170
pulmonary tuberculosis, 189
Pleurisy differentiated from pernicious malarial fever, 60
Pleurisy, pulsating, differentiated from thoracic aneurism, 143
Pleuritic effusion, 167, 168, 169
Pleuritic effusion differentiated from—
abscess of liver, 168
abscess of spleen, 168
hydatid cyst of liver, 167
hydrothorax, 169
Pleurosthotonos, 377
Plumbism, 364
Pneumonia, acute croupous or lobar, 176
Pneumonia, catarrhal, 175
Pneumonia, catarrhal, differentiated from—
atelectasis, 176
bronchiectasis, 173
capillary bronchitis, 175
croupous pneumonia, 180
pulmonary infarction, 176
Pneumonia, croupous, differentiated from—
acute pleurisy, 167
catarrhal pneumonia, 180
pulmonary congestion, 180
pulmonary edema, 179
pulmonary infarction, 180
pulmonary tuberculosis, 181
Pneumonia differentiated from—
pernicious malarial fever, 60
typhoid fever, 43
Pneumonia, lobular, 175
Pneumonitis, interstitial, 193
Pneumonitis, interstitial, differentiated from—
chronic bronchitis, 172
chronic pleurisy, 193
Pneumothorax, 195
Pneumothorax differentiated from—
diaphragmatic hernia, 196
pleural effusion, 197
pulmonary cavity, 196
pulmonary emphysema, 194
subphrenic abscess, 197
Poikilocytes, 105
Poisoning, arsenical, 52
Poisoning, narcotic, differentiated from sunstroke, 362
Poliomyelitis, anterior, acute, 318
Poliomyelitis, anterior, acute, differentiated from—
acute ascending paralysis, 320
acute myelitis, 318
cerebral paralysis, 318
multiple neuritis, 319
progressive muscular atrophy, 323
Poliomyelitis, anterior, subacute, differentiated from cervical pachymeningitis, 313
Polyneuritis, 303
Polyuria, 282
Polyuria, hysterical, differentiated from diabetes insipidus, 283
Portal phlebitis, 247
Posterior spinal sclerosis, 327
Postero-lateral sclerosis, 330
Previous history, 19
Primary lateral sclerosis, 329
Primary sciatica, 306
Progressive bulbar paralysis, 321
Progressive muscular atrophy, 320, 321

Progressive muscular dystrophy, 324
Progressive neural muscular atrophy, 323
Progressive paralysis of the insane, 359
Pseudo-bulbar palsy, 324
Pseudo-hydrophobia, 380
Pseudo-leukemia, 121
Pseudo-leukemia differentiated from—
leukemia, 111
tuberculosis of the lymph-glands, 112
Pseudo-leukemic anemia, infantile, 112
Pseudo-membranous laryngitis, 155
Pulmonary abscess, 173, 181, 182, 191
Pulmonary carcinoma, 189
Pulmonary cavity, 196
Pulmonary congestion, 180
Pulmonary edema, 179
Pulmonary emphysema, 194
Pulmonary gangrene, 182
Pulmonary hemorrhage, 189
Pulmonary hypertrophic osteo-arthropathy, 391
Pulmonary incompetency or regurgitation, 135
Pulmonary infarction, 176, 180
Pulmonary obstruction, 135
Pulmonary syphilis, 190
Pulmonary tuberculosis, 182
Pulsating pleurisy, 143
Pulsatory empyema, 166
Purity of the heart-sounds, 121
Purpura, 114
Purpura differentiated from—
measles, 115
scorbutus, 115
Purring tremor, 119
Putrid bronchitis, 172
Putrid sore-throat, 209
Pyelitis, 274, 289
Pyelitis differentiated from cystitis, 289
Pyemia differentiated from—
acute rheumatism, 96
typhoid fever, 41
Pylephlebitis differentiated from catarrhal jaundice, 260
Pylorus, insufficiency of, 214
Pylorus, spasm of, 214
Pyonephrosis, 298
Pyrexia, 27
Pyuria, 274

Quantity of urine, 267
Quinsy, 206

Rachitis, 117
Rachitis differentiated from mollities ossium, 118
Rag-weed fever, 149
Râles, 63
Rational signs, 19
Raynaud's disease, 144
Raynaud's disease differentiated from—
frost-bite, 145
Morvan's disease, 341
Reaction of degeneration, 301
Reaction of urine, 268
Recrudescence in typhoid fever, 39
Recurrent herpes of the throat, 208
Reflex symptoms, 19
Reflexes, 300
Reflexes, alterations in, in disease, 301
Reflexes, deep, 301
Reflexes, superficial, 300
Regurgitation, aortic, 133
Regurgitation, mitral, 132
Regurgitation, pulmonary, 135
Regurgitation, tricuspid, 134
Relapse in typhoid fever, 39
Relapsing fever, 52
Relapsing fever differentiated from—
typhoid fever, 53
typhus fever, 52
yellow fever, 53
Remittent fever, 30
Remittent fever, malarial, 57
Renal calculus, 291
Renal colic, 262
Renal inadequacy, 290
Rennet-ferment in gastric contents, 213
Resonance, vocal, 164
Respiratory percussion, 162
Respiratory system, 146
Retro-pharyngeal abscess, 156, 210
Rheumatic fever, 95
Rheumatic neuralgia, 99
Rheumatism, acute, 95
Rheumatism, acute, differentiated, from acute gout, 100

Rheumatism, acute, differentiated from
acute synovitis, 97
pyemia, 96
Rheumatism, chronic, 99
Rheumatism, chronic, differentiated from—
chronic spinal disease, 99
rheumatoid arthritis, 103
Rheumatism, muscular, 99
Rheumatism of the muscles of the back differentiated from spinal meningitis, 313
Rheumatism, subacute, 98
Rheumatism, subacute, differentiated from—
acute peritonitis, 243
neuralgia, 98
neuritis, 303
trichiniasis, 98
Rheumatoid arthritis, 102
Rheumatoid arthritis differentiated from—
chronic rheumatism, 103
gout, 104
Rhinoscopy, 146
Rhizomelic spondylosis, 392
Risus sardonicus, 377
Rock fever, 61
Rose-cold, 149
Roseola, 74
Rötheln, 74
Round-worms, 237
Rubella, 74
Rubella differentiated from—
morbilli, 75
scarlatina, 75
Rumination, 214

Sarcoma of the omentum differentiated from carcinoma of the stomach, 222
Scarlatina, 70
Scarlet fever, 70
Scarlet fever differentiated from—
acute exfoliative dermatitis, 73
dengue, 84
diphtheria, 86
erysipelas, 81
measles, 73
rubella, 75
smallpox, 78
toxic dermatitis, 73
Sciatic neuralgia, 305
Sciatica, 305
Sciatica differentiated from—
disease of the hip-joint, 305
sciatic neuralgia, 305
Sciatica, primary, differentiated from secondary sciatica, 306
Sciatica, secondary, differentiated from primary sciatica, 306
Sclerodactyly, 340
Scleroderma differentiated from—
Morvan's disease, 340
myxedema, 386
Sclerosis, cerebro-spinal, 332
Sclerosis, cerebro-spinal, differentiated from—
cerebral tumor, 358
paralysis agitans, 334
postero-lateral sclerosis, 332
Sclerosis, disseminated, 332
Sclerosis, insular, 332
Sclerosis, lateral, amyotrophic, 320, 321
Sclerosis, lateral, differentiated from chronic myelitis, 317
Sclerosis, lateral, primary, 329
Sclerosis, lateral, primary, differentiated from—
cerebral hemiplegia, 330
chronic myelitis, 329
postero-lateral sclerosis, 331
Sclerosis, multiple, 332
Sclerosis, postero-lateral, 330
Sclerosis, postero-lateral, differentiated from—
cerebellar tumor, 331
cerebro-spinal sclerosis, 332
hereditary ataxic paraplegia, 331
posterior spinal sclerosis, 330
primary lateral sclerosis, 330
Sclerosis, spinal, posterior, 327
Sclerosis, spinal, posterior, differentiated from—
cerebellar tumor, 329
general paralysis of the insane, 360
lumbar pachymeningitis, 328
multiple neuritis, 328
postero-lateral sclerosis, 330
Scorbutus, 113
Scorbutus differentiated from purpura, 115
Scurvy, 113
Seat-worms, 239
Secondary sciatica, 306
Sensation, 300

Septic infection, fever of, differentiated from malarial fever, 59
Sequelæ, 22
Shaking palsy, 333
Ship-fever, 44
Sick headache, 311
Signs, 18
Signs, instrumental, 18
Signs, negative, 23
Signs, physical, 18
Signs, rational, 19
Simple continued fever, 31
Simple endocarditis, 139
Smallpox, 75
Smallpox, confluent, 76
Smallpox differentiated from—
cerebro-spinal fever, 49
chickenpox, 79
erysipelas, 82
glanders, 88
measles, 78
scarlet fever, 78
typhoid fever, 42
typhus fever, 45
Smallpox, discrete, 76
Smallpox, hemorrhagic, 77
Smallpox, malignant, 77
Smallpox, varieties of, 76
Sodium chloride in urine, 269
Softening, cerebral, 354
Sore-throat, common membranous, 208
Sore-throat, common membranous, differentiated from diphtheria, 209
Sore-throat, gangrenous, differentiated from diphtheria, 209
Sore-throat, herpetic, 208
Sore-throat, putrid, 209
Sore-throat, syphilitic, differentiated from tuberculous pharyngitis, 206
Spasm, 300
Spasm of the cardia, 214
Spasm of the pylorus, 214
Spasmodic croup, 154
Spasmodic tic, 369
Spastic paraplegia, 329
Spastic paraplegia, congenital, 346
Specific fever, 30
Specific gravity of urine, 268
Sphygmogram, 124
Sphygmograph, 124
Spinal compression, 336
Spinal disease, chronic, differentiated from chronic rheumatism, 99
Spinal hemorrhage, 335
Spinal membranes, hemorrhage into, 314
Spinal meningitis, 312
Spinal pachymeningitis, 304, 317
Spinal tumor, 337
Splanchnoptosis, 215
Spleen, 263
Spleen, abscess of, 68
Spleen, enlargement of, 264
Spleen, floating, 264
Splenic fever, 89
Splenic tumor, 264
Splenitis, 264
Splenomegaly, 265
Spondylosis, rhizomelic, 392
Spool-worms, 239
Sporadic cretinism, 385
Sporadic cretinoidism, 385
Spotted fever, 47
Sputum, tubercle-bacilli in, 186
Status epilepticus, 371
Stomach, 212
Stomach, atony of, 214
Stomach, digestive activity of, 213
Stomach, dilatation of, 218
Stomach, functions of, 213
Stomach, hyperesthesia of, 214
Stomach, motor activity of, 214
Stomach, neuralgia of, 215
Stomach, neuroses of, 214
Stomatitis, aphthous, 200
Stomatitis, catarrhal, 199
Stomatitis differentiated from diphtheria, 87
Stomatitis, gangrenous, 201
Stomatitis, mercurial, 201
Stomatitis, parasitic, 200
Stomatitis, ulcerative, 200
Stricture, functional, of the esophagus differentiated from organic stricture, 212
Stricture of the esophagus, 211
Strychnine-poisoning differentiated from tetanus, 378
Subacidity, 215
Subacute rheumatism, 98
Subfebrile temperature, 28
Subjective symptoms, 17
Subnormal temperature, 29
Subphrenic abscess, 197
Sugar in the urine, Boettger's test for the presence of, 285

Sugar in the urine, Fehling's test for the presence of, 286
Sugar in the urine, fermentation-test for the presence of, 286
Sugar in the urine, Moore's test for the presence of, 285
Sugar in the urine, Trommer's test for the presence of, 286
Sulphuric acid in urine, 271
Sun-pain, 59
Sunstroke, 361
Sunstroke differentiated from—
acute alcoholism, 362
acute meningitis, 362
cerebral hemorrhage, 362
heat-exhaustion, 362
narcotic poisoning, 362
uremia, 295, 362
Superficial reflexes, 273
Supermotility, 214
Supersecretion, 215
Suppuration, fever of, differentiated from malarial fever, 59
Suprarenal bodies, tuberculosis of, 116
Sweating disease, 82
Symmetrical gangrene, 144
Symptomatic fever, 30
Symptoms, 17
Symptoms, constitutional, 19
Symptoms, general, 19
Symptoms, local, 19
Symptoms, objective, 17
Symptoms, reflex, 19
Symptoms, subjective, 17
Syncope differentiated from—
cerebral hemorrhage, 353
epilepsy, 371
Syncope, local, 144
Synovitis, acute, differentiated from acute rheumatism, 97
Synovitis, gonorrheal, 97
Syphilis differentiated from—
glanders, 88
Morvan's disease, 341
Syphilis, pulmonary, differentiated from pulmonary tuberculosis, 190
Syphilitic arthritis, 97
Syphilitic fever differentiated from malarial fever, 61
Syphilitic sore-throat, 206
Syringomyelia, 339
Syringomyelia differentiated from chronic myelitis, 339
Syringomyelia differentiated from—
hypertrophic pachymeningitis, 339
leprosy, 339
progressive muscular atrophy, 339

Tabes dorsalis, 327
Tabes mesenterica, 244
Tachycardia, 126
Tachycardia differentiated from—
angina pectoris, 126
exophthalmic goiter, 126
irritable heart, 127
Tactile fremitus, 160
Tænia mediocanellata, 236
Tænia solium, 235
Tape-worm, 235
Temperature, 27
Temperature, febrile, 28
Temperature, inverse, 28
Temperature, subfebrile, 28
Temperature, subnormal, 29
Temperature, sudden fall of, 29
Test-breakfast, 213
Test-dinner, 213
Tetanus, 377
Tetanus differentiated from—
acute spinal meningitis, 378
cerebro-spinal fever, 48
hemorrhage into the spinal membranes, 378
hydrophobia, 381
hysteria, 378
strychnine-poisoning, 378
tetany, 379
Tetany, 379
Tetany differentiated from tetanus, 379
Thermic fever, 361
Thomsen's disease, 327
Thoracic aneurism, 141
Thread-worms, 239
Throat, chronic or recurrent herpes of, 208
Throat, diphtheroid, 208
Thrombosis, cerebral, 356
Thrombosis, cerebral, differentiated from cerebral hemorrhage, 356
Thrush, 200
Tic douloureux, 311
Tic, spasmodic, 369
Tongue, 202
Tongue, black or hairy, 203
Tongue geographical, 203

Tonsillitis, 206
Tonsillitis, chronic, 208
Tonsillitis differentiated from—
diphtheria, 86
laryngitis, 150
Tonsillitis, herpetic, 208
Tonsillitis, lacunal or follicular, 207
Tonsillitis, parenchymatous, 206
Töpfer's test for hydrochloric acid, 212
Torticollis, 98, 366
Torticollis differentiated from cerebro-spinal fever, 49
Touch, 160
Translumination, 146
Transparency of urine, 267
Tremor, purring, 119
Trichiniasis, 92, 234
Trichiniasis differentiated from—
subacute rheumatism, 98
typhoid fever, 43
Tricuspid incompetency or regurgitation, 134
Tricuspid obstruction, 135
Trifacial neuralgia, 311
Trigeminal neuralgia, 311
Trommer's test for the presence of sugar in the urine, 286
Tube-casts in urine, 278
Tubercle-bacilli in sputum, 186
Tuberculosis, acute miliary, 191
Tuberculosis, acute miliary, differentiated from—
acute bronchitis, 171
capillary bronchitis, 174
typhoid fever, 192
Tuberculosis, chronic, of the larynx, 159
Tuberculosis of the bronchial glands differentiated from whooping-cough, 158
Tuberculosis of the kidney, 296
Tuberculosis of the lymph-glands differentiated from pseudo-leukemia, 111
Tuberculosis of the suprarenal bodies, 116
Tuberculosis, pulmonary, 182
Tuberculosis, pulmonary, differentiated from—
bronchiectasis, 190
chronic bronchitis, 172
chronic pleurisy, 189
croupous pneumonia, 181
malarial fever, 191
Tuberculosis, pulmonary, differentiated from—
pulmonary abscess, 191
pulmonary carcinoma, 189
pulmonary gangrene, 182
pulmonary syphilis, 190
Tuberculous laryngitis, acute, 152
Tuberculous meningitis, 343
Tuberculous pharyngitis, 206
Tuberculous pharyngitis, acute, 205
Tubular diarrhea, 224
Tumor, cerebellar, differentiated from—
posterior spinal sclerosis, 329
postero-lateral sclerosis, 331
Tumor, cerebral, 357
Tumor, cerebral, differentiated from—
anemia, 358
cerebral abscess, 350
cerebro-spinal sclerosis, 358
chronic nephritis, 358
epilepsy, 358
hysteria, 375
lead-poisoning, 365
Tumor, intra-thoracic, differentiated from—
aneurism of the aorta, 142
chronic pleurisy, 170
Tumor of the spinal cord, 337
Tumor of the spinal cord differentiated from—
caries of the vertebræ, 338
chronic myelitis, 338
hypertrophic pachymeningitis, 338
Tumor, splenic, 264
Typhlitis, 228
Typhlitis differentiated from—
abscess of the right ovary, 230
carcinoma of the cecum, 231
intussusception of bowel, 233
lumbar abscess, 230
Typhoid condition differentiated from typhoid fever, 42
Typhoid fever, 35
Typhoid fever, abortive, 39
Typhoid fever, ambulatory or walking, 39
Typhoid fever differentiated from—
acute catarrhal enteritis, 224
acute dysentery, 227
acute miliary tuberculosis, 192
acute yellow atrophy of the liver, 249

Typhoid fever differentiated from—
cerebro-spinal fever, 50
malarial fever, 61
Malta fever, 62
pneumonia, 43
pyemia, 41
relapsing fever, 53
trichiniasis, 43
tuberculous pharyngitis, 206
typhoid condition, 42
typhus fever, 45
variola, 42
yellow fever, 42
Typhoid fever, perforation of the bowel in, 40
Typhoid fever, recrudescence in, 40
Typhoid fever, relapse in, 39
Typhus fever, 44
Typhus fever differentiated from—
cerebro-spinal fever, 48
morbilli, 70
plague, 63
relapsing fever, 53
typhoid fever, 45
variola, 45
Tyrosin, 272

Uffelmann's test for lactic acid, 213
Ulcer, gastric, 218
Ulcer, gastric, differentiated from chronic gastritis, 219
Ulceration of the stomach differentiated from carcinoma of the stomach, 221
Ulcerative stomatitis, 200
Ulcero-membranous angina, 208
Undulant fever, 61
Unrest, peristaltic, 214
Urea in urine, 268
Uremia, 293
Uremia differentiated from—
cerebral hemorrhage, 354
epilepsy, 294, 372
sunstroke, 295, 362
Ureteritis, 274
Urethritis, 274
Uric acid in urine, 269
Urine, acetone in, tests for the presence of, 286
Urine, albumin in, 275
Urine, biliary acids in, 271
Urine, biliary coloring-matter in, 271
Urine, color of, 267
Urine, constitution of, 267
Urine, diacetic acid in, tests for the presence of, 287
Urine, examination of, 267
Urine, hippuric acid in, 269
Urine, odor of, 268
Urine, phosphoric acid in, 271
Urine, quantity of, 267
Urine, reaction of, 268
Urine, sodium chloride in, 269
Urine, specific gravity of, 268
Urine, sugar in, Boettger's test for the presence of, 285
Urine, sugar in, Fehling's test for the presence of, 286
Urine, sugar in, fermentation-test for the presence of, 286
Urine, sugar in, Moore's test for the presence of, 285
Urine, sugar in, Trommer's test for the presence of, 286
Urine, sulphuric acid in, 271
Urine, transparency of, 267
Urine, tube-casts in, 278
Urine, urea in, 268
Urine, uric acid in, 269

Vaccination, 77
Vaccinia, 78
Valvular disease of the heart, 131
Valvular disease of the heart differentiated from acute endocarditis, 139
Varicella, 79
Variola, 75
Varioloid, 77
Varioloid differentiated from chickenpox, 79
Vasomotor ataxia, 384
Vegetative endocarditis, 139
Vertebræ, caries of, 338
Vertigo, aural 381
Vertigo, aural, differentiated from epilepsy, 381
Vertigo, labyrinthine, 381
Vertigo, laryngeal, 153
Vesical calculus, 289
Virus, humanized, 77
Vocal fremitus, 160
Vocal resonance, 164
Vomiting, nervous, 214

Walking typhoid fever, 139

Waxy casts of the uriniferous tubules, 278
Waxy kidney, 293
Weil's disease, 67
Weil's disease differentiated from—
  catarrhal jaundice, 260
  yellow fever, 67
Whispering pectoriloquy, 164
Whooping-cough, 157
Whooping-cough differentiated from—
  asthma, 198
  laryngismus stridulus, 157
  membranous croup, 157
  tuberculosis of the bronchial glands, 157
Wool-sorters' disease, 89
Writers' cramp, 367
Wry-neck, 98, 366

YELLOW fever, 64
Yellow fever differentiated from—
  acute yellow atrophy of the liver, 66
  catarrhal jaundice, 260
  cerebro-spinal fever, 49
  dengue, 66
  hemorrhagic malarial fever, 65
  malarial remittent fever, 65
  plague, 63
  relapsing fever, 53
  typhoid fever, 42
  Weil's disease, 67

Medical and Surgical Works

PUBLISHED BY

W. B. SAUNDERS, 925 Walnut Street, Philadelphia, Pa.

	PAGE
Abbott on Transmissible Diseases	18
American Pocket Medical Dictionary	35
*American Text-Book of Applied Therapeutics	8
*American Text-Book of Dis. of Children	13
*An American Text-Book of Diseases of the Eye, Ear, Nose, and Throat	15
*An American Text-Book of Genito-Urinary and Skin Diseases	14
*American Text-Book of Gynecology	12
*American Text-Book of Legal Medicine	44
*American Text-Book of Obstetrics	9
*American Text-Book of Pathology	44
*American Text-Book of Physiology	7
*American Text-Book of Practice	10
*American Text-Book of Surgery	11
Anders' Theory and Practice of Medicine	21
Ashton's Obstetrics	43
Atlas of Skin Diseases	28
Ball's Bacteriology	43
Bastin's Laboratory Exercises in Botany	36
Beck's Surgical Asepsis	41
Boislinière's Obstetric Accidents	39
Brockway's Physics	43
Burr's Nervous Diseases	41
Butler's Materia Medica and Therapeutics	24
Cerna's Notes on the Newer Remedies	32
Chapin's Compendium of Insanity	35
Chapman's Medical Jurisprudence	41
Church and Peterson's Nervous and Mental Diseases	17
Clarkson's Histology	33
Cohen and Eshner's Diagnosis	43
Corwin's Diagnosis of the Thorax	37
Cragin's Gynæcology	43
Crookshank's Text-Book of Bacteriology	27
DaCosta's Manual of Surgery	23
De Schweinitz's Diseases of the Eye	29
Dorland's Pocket Medical Dictionary	35
Dorland's Obstetrics	41
Frothingham's Bacteriological Guide	30
Garrigues' Diseases of Women	34
Gleason's Diseases of the Ear	43
*Gould and Pyle's Curiosities of Medicine	17
Grafstrom's Massage	28
Griffith's Care of the Baby	38
Griffith's Infant's Weight Chart	39
Gross's Autobiography	26
Hampton's Nursing	39
Hare's Physiology	43
Hart's Diet in Sickness and in Health	36
Haynes' Manual of Anatomy	41
Heisler's Embryology	19
Hirst's Obstetrics	20
Hyde's Syphilis and Venereal Diseases	41
International Text-Book of Surgery	6
Jackson's Diseases of the Eye	19
Jackson and Gleason's Diseases of the Eye, Nose, and Throat	43
Keating's Pronouncing Dictionary	26
Keating's Life Insurance	39
Keen's Operation Blanks	36
Keen's Surgery of Typhoid Fever	22
Kyle's Diseases of Nose and Throat	18
Lainé's Temperature Charts	32
Levy & Klemperer's Clinical Bacteriology	44
Lockwood's Practice of Medicine	41
Long's Syllabus of Gynecology	34
Macdonald's Surgical Diagnosis and Treatment	22
McFarland's Pathogenic Bacteria	30
Mallory and Wright's Pathological Technique	22
Martin's Surgery	43
Martin's Minor Surgery, Bandaging, and Venereal Diseases	43
Meigs' Feeding in Early Infancy	30
Moore's Orthopedic Surgery	23
Morris' Materia Medica and Therapeutics	43
Morris' Practice of Medicine	43
Morten's Nurses' Dictionary	38
Nancrede's Anatomy and Dissection	31
Nancrede's Anatomy	43
Nancrede's Principles of Surgery	19
Norris' Syllabus of Obstetrical Lectures	37
Penrose's Diseases of Women	24
Powell's Diseases of Children	43
Pryor's Pelvic Inflammations	33
Pye's Bandaging and Surgical Dressing	23
Raymond's Physiology	41
Saundby's Renal and Urinary Diseases	25
*Saunders' American Year-Book of Medicine and Surgery	16
Saunders' Medical Hand-Atlases	3, 4, 5
Saunders' Pocket Medical Formulary	35
Saunders' New Series of Manuals	40, 41
Saunders' Series of Question Compends	42, 43
Sayre's Practice of Pharmacy	43
Semple's Pathology and Morbid Anatomy	43
Semple's Legal Medicine and Toxicology	43
Senn's Genito-Urinary Tuberculosis	24
Senn's Tumors	25
Senn's Syllabus of Lectures on Surgery	37
Shaw's Nervous Diseases and Insanity	43
Starr's Diet-Lists for Children	38
Stelwagon's Diseases of the Skin	43
Stengel's Pathology	20
Stevens' Materia Medica and Therapeutics	32
Stevens' Practice of Medicine	31
Stewart's Manual of Physiology	37
Stewart and Lawrance's Medical Electricity	43
Stoney's Materia Medica for Nurses	31
Stoney's Practical Points in Nursing	27
Sutton and Giles' Diseases of Women	29, 41
Thomas's Diet-List and Sick-Room	38
Thornton's Dose-Book and Manual of Prescription-Writing	41
Van Valzah and Nisbet's Diseases of the Stomach	21
Vecki's Sexual Impotence	33
Vierordt and Stuart's Medical Diagnosis	28
Warren's Surgical Pathology	25
Watson's Handbook for Nurses	26
Wolff's Chemistry	43
Wolff's Examination of Urine	43

GENERAL INFORMATION.

One Price. One price absolutely without deviation. No discounts allowed, regardless of the number of books purchased at one time. Prices on all works have been fixed extremely low, with the view to selling them strictly net and for cash.

Orders. An order accompanied by remittance will receive prompt attention, books being sent to any address in the United States, by mail or express, all charges prepaid. We prefer to send books **by express** when possible.

Cash or Credit. To physicians of approved credit who furnish satisfactory references our books will be sent free of C. O. D. **One volume or two** on thirty days' time if credit is desired; larger purchases on monthly payment plan. See offer below.

How to Send Money by Mail. There are four ways by which money can be sent at our risk, namely: a post-office money order, an express money order, a bank-check (draft), and in a registered letter. Money sent in any other way is at the sender's risk. Silver should not be sent through the mail.

Shipments. All books, being packed in patent metal-edged boxes, necessarily reach our patrons by mail or express in excellent condition.

Subscription Books. **Books in this catalogue** marked with a star (*) are for sale by subscription only, and may be secured by ordering them through any of our authorized travelling salesmen, or direct from the Philadelphia office: they are **not** for sale by booksellers. All other books in our catalogue can be procured of any bookseller at the advertised price, or directly from us.

Miscellaneous Books. We carry in stock only our own publications, but can supply the publications of other houses (except subscription books) on receipt of publisher's price.

Latest Editions. In every instance the latest revised edition is sent.

Bindings. In ordering, be careful to state the style of binding desired—Cloth, Sheep, or Half Morocco.

Special Offer. Monthly Payment Plan. To physicians of approved credit who furnish satisfactory references books will be sent express prepaid; terms, $5.00 cash upon delivery of books, and monthly payments of $5 00 thereafter until full amount is paid. Any of the publications of W. B. Saunders (100 titles to select from) may be had in this way at catalogue price, including the American Text-Book Series, the Medical Hand-Atlases, etc. All payments to be made by mail or otherwise, free of all expense to us.

SAUNDERS'

MEDICAL HAND-ATLASES.

THE series of books included under this title consists of authorized translations into English of the world-famous **Lehmann Medicinische Handatlanten,** which for **scientific accuracy, pictorial beauty, compactness,** and **cheapness** surpass any similar volumes ever published. Each volume contains from **50 to 100 colored plates,** executed by the most skilful German lithographers, besides numerous illustrations in the text. There is a full and appropriate **description,** and each book contains a condensed but adequate **outline of the subject** to which it is devoted.

In planning this series arrangements were made with representative publishers in the chief medical centers of the world for the publication of translations of the atlases into nine different languages, the lithographic plates for all being made in Germany, where work of this kind has been brought to the greatest perfection. The enormous expense of making the plates being shared by the various publishers, the cost to each one was reduced to practically one-tenth. Thus by reason of their **universal translation** and reproduction, affording international distribution, the publishers have been enabled to secure for these atlases the **best artistic and professional talent,** to produce them in the **most elegant style,** and yet to offer them at a **price heretofore unapproached in cheapness.** The great success of the undertaking is demonstrated by the fact that the volumes have already appeared in **thirteen different languages**—German, English, French, Italian, Russian, Spanish, Japanese, Dutch, Danish, Swedish, Roumanian, Bohemian, and Hungarian.

In view of the unprecedented success of these works, Mr. Saunders has contracted with the publisher of the original German edition for **one hundred thousand copies** of the atlases. In consideration of this enormous undertaking, the publisher has been enabled to prepare and furnish special additional colored plates, making the series even **handsomer and more complete** than was originally intended.

As an indication of the great practical value of the atlases and of the immense favor with which they have been received, it should be noted that the **Medical Department of the U. S. Army** has adopted the "Atlas of Operative Surgery," as its standard, and has ordered the book in large quantities for distribution to the various regiments and army posts.

The same careful and competent **editorial supervision** has been secured in the English edition as in the originals. The translations have been edited by the **leading American specialists** in the different subjects.

(*For List of Volumes in this Series, see next two pages.*)

SAUNDERS' MEDICAL HAND-ATLASES.

VOLUMES NOW READY.

Atlas and Epitome of Internal Medicine and Clinical Diagnosis. By DR. CHR. JAKOB, of Erlangen. Edited by AUGUSTUS A. ESHNER, M. D., Professor of Clinical Medicine, Philadelphia Polyclinic. With 68 colored plates, 64 text-illustrations, and 259 pages of text. Cloth, $3.00 net.

"The charm of the book is its clearness, conciseness, and the accuracy and beauty of its illustrations. It deals with facts. It vividly illustrates those facts. It is a scientific work put together for ready reference."—*Brooklyn Medical Journal.*

Atlas of Legal Medicine. By DR. E. R. VON HOFMANN, of Vienna. Edited by FREDERICK PETERSON, M. D., Chief of Clinic, Nervous Dept., College of Physicians and Surgeons, New York. With 120 colored figures on 56 plates, and 193 beautiful half-tone illustrations. Cloth, $3.50 net.

"Hofmann's 'Atlas of Legal Medicine' is a unique work. This immense field finds in this book a pictorial presentation that far excels anything with which we are familiar in any other work."—*Philadelphia Medical Journal.*

Atlas and Epitome of Diseases of the Larynx. By DR. L. GRUNWALD, of Munich. Edited by CHARLES P. GRAYSON, M. D., Physician-in-Charge, Throat and Nose Department, Hospital of the University of Pennsylvania. With 107 colored figures on 44 plates, 25 text-illustrations, and 103 pages of text. Cloth, $2.50 net.

"Aided as it is by magnificently executed illustrations in color, it cannot fail of being of the greatest advantage to students, general practitioners, and expert laryngologists."—*St. Louis Medical and Surgical Journal.*

Atlas and Epitome of Operative Surgery. By DR. O. ZUCKERKANDL, of Vienna. Edited by J. CHALMERS DACOSTA, M. D., Clinical Professor of Surgery, Jefferson Medical College, Philadelphia. With 24 colored plates, 217 text-illustrations, and 395 pages of text. Cloth, $3.00 net.

"We know of no other work that combines such a wealth of beautiful illustrations with clearness and conciseness of language, that is so entirely abreast of the latest achievements, and so useful both for the beginner and for one who wishes to increase his knowledge of operative surgery."—*Münchener medicinische Wochenschrift.*

Atlas and Epitome of Syphilis and the Venereal Diseases. By PROF. DR. FRANZ MRACEK, of Vienna. Edited by L. BOLTON BANGS, M. D., Professor of Genito-Urinary Surgery, University and Bellevue Hospital Medical College, New York. With 71 colored plates, 16 black-and-white illustrations, and 122 pages of text. Cloth, $3.50 net.

"A glance through the book is almost like actual attendance upon a famous clinic."—*Journal of the American Medical Association.*

Atlas and Epitome of External Diseases of the Eye. By DR. O. HAAB, of Zurich. Edited by G. E. DE SCHWEINITZ, M. D., Professor of Ophthalmology, Jefferson Medical College, Philadelphia. With 76 colored illustrations on 40 plates, and 228 pages of text. Cloth, $3.00 net.

"It is always difficult to represent pathological appearances in colored plates, but this work seems to have overcome these difficulties, and the plates, with one or two exceptions, are absolutely satisfactory."—*Boston Medical and Surgical Journal.*

Atlas and Epitome of Skin Diseases. By PROF. DR. FRANZ MRACEK, of Vienna. Edited by HENRY W. STELWAGON, M. D., Clinical Professor of Dermatology, Jefferson Medical College, Philadelphia. With 63 colored plates, 39 half-tone illustrations, and 200 pages of text. Cloth, $3.50 net.

"The importance of personal inspection of cases in the study of cutaneous diseases is readily appreciated, and next to the living subjects are pictures which will show the appearance of the disease under consideration. Altogether the work will be found of very great value to the general practitioner."—*Journal of the American Medical Association.*

SAUNDERS' MEDICAL HAND-ATLASES.

VOLUMES IN PRESS FOR EARLY PUBLICATION.

Atlas and Epitome of Diseases Caused by Accidents. By Dr. Ed. Golebiewski, of Berlin. Translated and edited with additions by Pearce Bailey, M.D., Attending Physician to the Department of Corrections and to the Almshouse and Incurable Hospitals, New York. With 40 colored plates, 143 text-illustrations, and 600 pages of text.

Atlas and Epitome of Special Pathological Histology. By Dr. H. Dürck, of Munich. Edited by Ludvig Hektoen, M.D., Professor of Pathology, Rush Medical College, Chicago. Two volumes, with about 120 colored plates, numerous text-illustrations, and copious text.

Atlas and Epitome of General Pathological Histology. With an Appendix on Patho-histological Technic. By Dr. H. Dürck, of Munich. Edited by Ludvig Hektoen, M.D., Professor of Pathology, Rush Medical College, Chicago. With 80 colored plates, numerous text-illustrations, and copious text.

Atlas and Epitome of Gynecology. By Dr. O. Schäffer, of the University of Heidelberg. With 90 colored plates, 65 text-illustrations, and 308 pages of text. Edited by Richard C. Norris, A. M., M. D., Gynecologist to the Philadelphia and the Methodist Episcopal Hospitals.

IN PREPARATION.

Atlas and Epitome of Orthopedic Surgery. By Dr. Schultess and Dr. Lüning, of Zurich. About 100 colored illustrations.

Atlas and Epitome of Operative Gynecology. By Dr. O. Schäffer, of Heidelberg. With 40 colored plates and numerous illustrations in black and white from original paintings.

Atlas and Epitome of Diseases of the Ear. Edited by Prof. Dr. Politzer, of Vienna, and Dr. G. Brühl, of Berlin. With 120 colored illustrations and about 200 pages of text.

Atlas and Epitome of General Surgery. Edited by Dr. Marwedel, with the coöperation of Prof. Dr. Czerny. With about 200 colored illustrations.

Atlas and Epitome of Psychiatry. By Dr. Wilh. Weygandt, of Würzburg. With about 120 colored illustrations.

Atlas and Epitome of Normal Histology. By Dr. Johannes Sobotta, of Würzburg. With 80 colored plates and numerous illustrations.

Atlas and Epitome of Topographical Anatomy. By Prof. Dr. Schultze, of Würzburg. About 100 colored illustrations and a very copious text.

* **THE INTERNATIONAL TEXT-BOOK OF SURGERY. In two volumes. By American and British authors.** Edited by J. COLLINS WARREN, M.D., LL.D., Professor of Surgery, Harvard Medical School, Boston; Surgeon to the Massachusetts General Hospital; and A. PEARCE GOULD, M. S., F. R. C. S., Eng., Lecturer on Practical Surgery and Teacher of Operative Surgery, Middlesex Hospital Medical School; Surgeon to the Middlesex Hospital, London, England. Vol. I.—**General and Operative Surgery.**—Handsome octavo volume of 947 pages, with 458 beautiful illustrations, and 9 lithographic plates. Vol. II.—**Special or Regional Surgery.**—Handsome octavo volume of 1050 pages, with over 500 wood-cuts and half-tones, and 8 lithographic plates. Prices per volume: Cloth, $5.00 net; Half-Morocco, $6.00 net.

Just Issued.

In presenting a new work on surgery to the medical profession the publisher feels that he need offer no apology for making an addition to the list of excellent works already in existence. Modern surgery is still in the transition stage of its development. The art and science of surgery are advancing rapidly, and the number of workers is now so great and so widely spread through the whole of the civilized world that there is certainly room for another work of reference which shall be untrammelled by many of the traditions of the past, and shall at the same time present with due discrimination the results of modern progress. There is a real need among practitioners and advanced students for a work on surgery encyclopedic in scope, yet so condensed in style and arrangement that the matter usually diffused through four or five volumes shall be given in one-half the space and at a correspondingly moderate cost.

The ever-widening-field of surgery has been developed largely by special work, and this method of progress has made it practically impossible for one man to write authoritatively on the vast range of subjects embraced in a modern text-book of surgery. In order, therefore, to accomplish their object, the editors have sought the aid of men of wide experience and established reputation in the various departments of surgery.

CONTRIBUTORS:

Dr. Robert W. Abbe.
C. H. Golding Bird.
E. H. Bradford.
W. T. Bull.
T. G. A. Burns.
Herbert L. Burrell.
R. C. Cabot.
I. H. Cameron.
James Cantlie.
W. Watson Cheyne.
William B. Clarke.
William B. Coley.
Edw. Treacher Collins.
H. Holbrook Curtis.
J. Chalmers Da Costa.
N. P. Dandridge.
John B. Deaver.
J. W. Elliot.
Harold Ernst.

Dr. Christian Fenger.
W. H. Forwood.
George R. Fowler.
George W. Gay.
A. Pearce Gould.
J. Orne Green.
John B. Hamilton.
M. L. Harris.
Fernand Henrotin.
G. H. Makins.
Rudolph Matas.
Charles McBurney.
A. J. McCosh.
L. S. McMurtry.
J. Ewing Mears.
George H. Monks.
John Murray.
Robert W. Parker.

Dr. Rushton Parker.
George A. Peters.
Franz Pfaff.
Lewis S. Pilcher.
James J. Putnam.
M. H. Richardson.
A. W. Mayo Robson.
W. L. Rodman.
C. A. Siegfried.
G. B. Smith.
W. G. Spencer.
J. Bland Sutton.
L. McLane Tiffany.
H. Tuholske.
Weller Van Hook.
James P. Warbasse.
J. Collins Warren.
De Forest Willard.

***AN AMERICAN TEXT-BOOK OF PHYSIOLOGY.** Edited by WILLIAM H. HOWELL, PH. D., M. D., Professor of Physiology in the Johns Hopkins University, Baltimore, Md. One handsome octavo volume of 1052 pages, fully illustrated. Prices: Cloth, $6.00 net; Sheep or Half-Morocco, $7.00 net.

This work is the most notable attempt yet made in America to combine in one volume the entire subject of Human Physiology by well-known teachers who have given especial study to that part of the subject upon which they write. The completed work represents the present status of the science of Physiology, particularly from the standpoint of the student of medicine and of the medical practitioner.

The collaboration of several teachers in the preparation of an elementary text-book of physiology is unusual, the almost invariable rule heretofore having been for a single author to write the entire book. One of the advantages to be derived from this collaboration method is that the more limited literature necessary for consultation by each author has enabled him to base his elementary account upon a comprehensive knowledge of the subject assigned to him; another, and perhaps the most important, advantage is that the student gains the point of view of a number of teachers. In a measure he reaps the same benefit as would be obtained by following courses of instruction under different teachers. The different standpoints assumed, and the differences in emphasis laid upon the various lines of procedure, chemical, physical, and anatomical, should give the student a better insight into the methods of the science as it exists to-day. The work will also be found useful to many medical practitioners who may wish to keep in touch with the development of modern physiology.

CONTRIBUTORS:

HENRY P. BOWDITCH, M. D.,
Professor of Physiology, Harvard Medical School.

JOHN G. CURTIS, M. D.,
Professor of Physiology, Columbia University, N. Y. (College of Physicians and Surgeons).

HENRY H. DONALDSON, Ph. D.,
Head-Professor of Neurology, University of Chicago.

W. H. HOWELL, Ph. D., M. D.,
Professor of Physiology, Johns Hopkins University.

FREDERIC S. LEE, Ph. D.,
Adjunct Professor of Physiology, Columbia University, N. Y. (College of Physicians and Surgeons).

WARREN P. LOMBARD, M. D.,
Professor of Physiology, University of Michigan.

GRAHAM LUSK, Ph. D.,
Professor of Physiology, Yale Medical School.

W. T. PORTER, M. D.,
Assistant Professor of Physiology, Harvard Medical School.

EDWARD T. REICHERT, M. D.,
Professor of Physiology, University of Pennsylvania.

HENRY SEWALL, Ph. D., M. D.,
Professor of Physiology, Medical Department, University of Denver.

"We can commend it most heartily, not only to all students of physiology, but to every physician and pathologist, as a valuable and comprehensive work of reference, written by men who are of eminent authority in their own special subjects."—*London Lancet.*

"To the practitioner of medicine and to the advanced student this volume constitutes, we believe, the best exposition of the present status of the science of physiology in the English language."—*American Journal of the Medical Sciences.*

***AN AMERICAN TEXT-BOOK OF APPLIED THERAPEUTICS. For the Use of Practitioners and Students.** Edited by JAMES C. WILSON, M. D., Professor of the Practice of Medicine and of Clinical Medicine in the Jefferson Medical College. One handsome octavo volume of 1326 pages. Illustrated. Prices: Cloth, $7.00 net; Sheep or Half-Morocco, $8.00 net.

The arrangement of this volume has been based, so far as possible, upon modern pathologic doctrines, beginning with the intoxications, and following with infections, diseases due to internal parasites, diseases of undetermined origin, and finally the disorders of the several bodily systems—digestive, respiratory, circulatory, renal, nervous, and cutaneous. It was thought proper to include also a consideration of the disorders of pregnancy.

The articles, with two exceptions, are the contributions of American writers. Written from the standpoint of the practitioner, the aim of the work is to facilitate the application of knowledge to the prevention, the cure, and the alleviation of disease. The endeavor throughout has been to conform to the title of the book—Applied Therapeutics—to indicate the course of treatment to be pursued at the bedside, rather than to name a list of drugs that have been used at one time or another.

The list of contributors comprises the names of many who have acquired distinction as practitioners and teachers of practice, of clinical medicine, and of the specialties.

CONTRIBUTORS:

Dr. I. E. Atkinson, Baltimore, Md.
Sanger Brown, Chicago, Ill.
John B. Chapin, Philadelphia, Pa.
William C. Dabney, Charlottesville, Va.
John Chalmers DaCosta, Philada., Pa.
I. N. Danforth, Chicago, Ill.
John L. Dawson, Jr., Charleston, S. C.
F. X. Dercum, Philadelphia, Pa.
George Dock, Ann Arbor, Mich.
Robert T. Edes, Jamaica Plain, Mass.
Augustus A. Eshner, Philadelphia, Pa.
J. T. Eskridge, Denver, Col.
F. Forchheimer, Cincinnati, O.
Carl Frese, Philadelphia, Pa.
Edwin E. Graham, Philadelphia, Pa.
John Guitéras, Philadelphia, Pa.
Frederick P. Henry, Philadelphia, Pa.
Guy Hinsdale, Philadelphia, Pa.
Orville Horwitz, Philadelphia, Pa.
W. W. Johnston, Washington, D. C.
Ernest Laplace, Philadelphia, Pa.
A. Laveran, Paris, France.
Dr. James Hendrie Lloyd, Philadelphia, Pa.
John Noland Mackenzie, Baltimore, Md.
J. W. McLaughlin, Austin, Texas.
A. Lawrence Mason, Boston, Mass.
Charles K. Mills, Philadelphia, Pa.
John K. Mitchell, Philadelphia, Pa.
W. P. Northrup, New York City.
William Osler, Baltimore, Md.
Frederick A. Packard, Philadelphia, Pa.
Theophilus Parvin, Philadelphia, Pa.
Beaven Rake, London, England.
E. O. Shakespeare, Philadelphia, Pa.
Wharton Sinkler, Philadelphia, Pa.
Louis Starr, Philadelphia, Pa.
Henry W. Stelwagon, Philadelphia, Pa.
James Stewart, Montreal, Canada.
Charles G. Stockton, Buffalo, N. Y.
James Tyson, Philadelphia, Pa.
Victor C. Vaughan, Ann Arbor, Mich.
James T. Whittaker, Cincinnati, O.
J. C. Wilson, Philadelphia, Pa.

"As a work either for study or reference it will be of great value to the practitioner, as it is virtually an exposition of such clinical therapeutics as experience has taught to be of the most value. Taking it all in all, no recent publication on therapeutics can be compared with this one in practical value to the working physician."—*Chicago Clinical Review.*

"The whole field of medicine has been well covered. The work is thoroughly practical, and while it is intended for practitioners and students, it is a better book for the general practitioner than for the student. The young practitioner especially will find it extremely suggestive and helpful."—*The Indian Lancet.*

***AN AMERICAN TEXT-BOOK OF OBSTETRICS.** Edited by RICHARD C. NORRIS, M. D.; Art Editor, ROBERT L. DICKINSON, M. D. One handsome octavo volume of over 1000 pages, with nearly 900 colored and half-tone illustrations. Prices: Cloth, $7.00 net; Sheep or Half Morocco, $8.00 net.

The advent of each successive volume of the *series* of the AMERICAN TEXT-BOOKS has been signalized by the most flattering comment from both the Press and the Profession. The high consideration received by these text-books, and their attainment to an authoritative position in current medical literature, have been matters of deep *international* interest, which finds its fullest expression in the demand for these publications from all parts of the civilized world.

In the preparation of the "AMERICAN TEXT-BOOK OF OBSTETRICS" the editor has called to his aid proficient collaborators whose professional prominence entitles them to recognition, and whose disquisitions exemplify **Practical Obstetrics.** While these writers were each assigned special themes for discussion, the correlation of the subject-matter is, nevertheless, such as ensures logical connection in treatment, the deductions of which thoroughly represent the latest advances in the science, and which elucidate *the best modern methods of procedure.*

The more conspicuous feature of the treatise is its wealth of illustrative matter. The production of the illustrations had been in progress for several years, under the personal supervision of Robert L. Dickinson, M. D., to whose artistic judgment and professional experience is due the **most sumptuously illustrated work of the period.** By means of the photographic art, combined with the skill of the artist and draughtsman, conventional illustration is superseded by rational methods of delineation.

Furthermore, the volume is a revelation as to the possibilities that may be reached in mechanical execution, through the unsparing hand of its publisher.

CONTRIBUTORS:

Dr. James C. Cameron.
Edward P. Davis.
Robert L. Dickinson.
Charles Warrington Earle.
James H. Etheridge.
Henry J. Garrigues.
Barton Cooke Hirst.
Charles Jewett.
Dr. Howard A. Kelly.
Richard C. Norris.
Chauncey D. Palmer.
Theophilus Parvin.
George A. Piersol.
Edward Reynolds.
Henry Schwarz.

"At first glance we are overwhelmed by the magnitude of this work in several respects, viz.: First, by the size of the volume, then by the array of eminent teachers in this department who have taken part in its production, then by the profuseness and character of the illustrations, and last, but not least, the conciseness and clearness with which the text is rendered. This is an entirely new composition, embodying the highest knowledge of the art as it stands to-day by authors who occupy the front rank in their specialty, and there are many of them. We cannot turn over these pages without being struck by the superb illustrations which adorn so many of them. We are confident that this most practical work will find instant appreciation by practitioners as well as students."—*New York Medical Times.*

Permit me to say that your American Text-Book of Obstetrics is the most magnificent medical work that I have ever seen. I congratulate you and thank you for this superb work, which alone is sufficient to place you first in the ranks of medical publishers.

With profound respect I am sincerely yours, ALEX. J. C. SKENE.

*AN AMERICAN TEXT-BOOK OF THE THEORY AND PRACTICE OF MEDICINE. By American Teachers. Edited by WILLIAM PEPPER, M. D., LL.D., Provost and Professor of the Theory and Practice of Medicine and of Clinical Medicine in the University of Pennsylvania. Complete in two handsome royal-octavo volumes of about 1000 pages each, with illustrations to elucidate the text wherever necessary. Price per Volume: Cloth, $5.00 net; Sheep or Half-Morocco, $6.00 net.

VOLUME I. CONTAINS:

Hygiene.—Fevers (Ephemeral, Simple Continued, Typhus, Typhoid, Epidemic Cerebro-spinal Meningitis, and Relapsing).—Scarlatina, Measles, Rötheln, Variola, Varioloid, Vaccinia, Varicella, Mumps, Whooping-cough, Anthrax, Hydrophobia, Trichinosis, Actinomycosis, Glanders, and Tetanus.—Tuberculosis, Scrofula, Syphilis, Diphtheria, Erysipelas, Malaria, Cholera, and Yellow Fever.—Nervous, Muscular, and Mental Diseases etc.

VOLUME II. CONTAINS:

Urine (Chemistry and Microscopy).—Kidney and Lungs.—Air-passages (Larynx and Bronchi) and Pleura.—Pharynx, Œsophagus, Stomach and Intestines (including Intestinal Parasites), Heart, Aorta, Arteries and Veins.—Peritoneum, Liver, and Pancreas.—Diathetic Diseases (Rheumatism, Rheumatoid Arthritis, Gout, Lithæmia, and Diabetes.)—Blood and Spleen.—Inflammation, Embolism, Thrombosis, Fever, and Bacteriology.

The articles are not written as though addressed to students in lectures, but are exhaustive descriptions of diseases, with the newest facts as regards Causation, Symptomatology, Diagnosis, Prognosis, and Treatment, including a large number of approved formulæ. The recent advances made in the study of the bacterial origin of various diseases are fully described, as well as the bearing of the knowledge so gained upon prevention and cure. The subjects of Bacteriology as a whole and of Immunity are fully considered in a separate section.

Methods of diagnosis are given the most minute and careful attention, thus enabling the reader to learn the very latest methods of investigation without consulting works specially devoted to the subject.

CONTRIBUTORS:

Dr. J. S. Billings, Philadelphia.
Francis Delafield, New York.
Reginald H. Fitz, Boston.
James W. Holland, Philadelphia.
Henry M. Lyman, Chicago.
William Osler, Baltimore.
Dr. William Pepper, Philadelphia.
W. Gilman Thompson, New York.
W. H. Welch, Baltimore.
James T. Whittaker, Cincinnati.
James C. Wilson, Philadelphia.
Horatio C. Wood, Philadelphia.

"We reviewed the first volume of this work, and said: 'It is undoubtedly one of the best text-books on the practice of medicine which we possess.' A consideration of the second and last volume leads us to modify that verdict and to say that the completed work is, in our opinion, THE BEST of its kind it has ever been our fortune to see. It is complete, thorough, accurate, and clear. It is well written, well arranged, well printed, well illustrated, and well bound. It is a model of what the modern text-book should be."—*New York Medical Journal.*

"A library upon modern medical art. The work must promote the wider diffusion of sound knowledge."—*American Lancet.*

"A trusty counsellor for the practitioner or senior student, on which he may implicitly rely."—*Edinburgh Medical Journal.*

***AN AMERICAN TEXT-BOOK OF SURGERY.** Edited by WILLIAM W. KEEN, M. D., LL.D., and J. WILLIAM WHITE, M. D., PH. D. Forming one handsome royal octavo volume of 1230 pages (10×7 inches), with 496 wood-cuts in text, and 37 colored and half-tone plates, many of them engraved from original photographs and drawings furnished by the authors. Price: Cloth, $7.00 net; Sheep or Half Morocco, $8.00 net.

THIRD EDITION. THOROUGHLY REVISED.

In the present edition, among the new topics introduced are a full consideration of serum-therapy; leucocytosis; post-operative insanity; the use of dry heat at high temperatures; Krönlein's method of locating the cerebral fissures; Hoffa's and Lorenz's operations of congenital dislocations of the hip; Allis's researches on dislocations of the hip-joint; lumbar puncture; the forcible reposition of the spine in Pott's disease; the treatment of exophthalmic goiter; the surgery of typhoid fever; gastrectomy and other operations on the stomach; new methods of operating upon the intestines; the use of Kelly's rectal specula; the surgery of the ureter; Schleich's infiltration-method and the use of eucain for local anesthesia; Krause's method of skin-grafting; the newer methods of disinfecting the hands; the use of gloves, etc. The sections on Appendicitis, on Fractures, and on Gynecological Operations have been revised and enlarged. A considerable number of new illustrations have been added, and enhance the value of the work.

The text of the entire book has been submitted to all the authors for their mutual criticism and revision—an idea in book-making that is entirely new and original. The book as a whole, therefore, expresses on all the important surgical topics of the day the consensus of opinion of the eminent surgeons who have joined in its preparation.

One of the most attractive features of the book is its illustrations. Very many of them are original and faithful reproductions of photographs taken directly from patients or from specimens.

CONTRIBUTORS:

Dr. Phineas S. Conner, Cincinnati.
Frederic S. Dennis, New York.
William W. Keen, Philadelphia.
Charles B. Nancrede, Ann Arbor, Mich.
Roswell Park, Buffalo, New York.
Lewis S. Pilcher, New York.
Dr. Nicholas Senn, Chicago.
Francis J. Shepherd, Montreal, Canada.
Lewis A. Stimson, New York.
J. Collins Warren, Boston.
J. William White, Philadelphia.

"If this text-book is a fair reflex of the present position of American surgery, we must admit it is of a very high order of merit, and that English surgeons will have to look very carefully to their laurels if they are to preserve a position in the van of surgical practice."—*London Lancet.*

Personally, I should not mind it being called THE TEXT-BOOK (instead of A TEXT-BOOK), for I know of no single volume which contains so readable and complete an account of the science and art of Surgery as this does."—EDMUND OWEN, F. R. C. S., *Member of the Board of Examiners of the Royal College of Surgeons, England.*

*AN AMERICAN TEXT-BOOK OF GYNECOLOGY, MEDICAL AND SURGICAL, for the use of Students and Practitioners.

Edited by J. M. BALDY, M. D. Forming a handsome royal-octavo volume of 718 pages, with 341 illustrations in the text and 38 colored and half-tone plates. Prices: Cloth, $6.00 net; Sheep or Half-Morocco, $7.00 net.

SECOND EDITION, THOROUGHLY REVISED.

In this volume all anatomical descriptions, excepting those essential to a clear understanding of the text, have been omitted, the illustrations being largely depended upon to elucidate the anatomy of the parts. This work, which is thoroughly practical in its teachings, is intended, as its title implies, to be a working text-book for physicians and students. A clear line of treatment has been laid down in every case, and although no attempt has been made to discuss mooted points, still the most important of these have been noted and explained. The operations recommended are fully illustrated, so that the reader, having a picture of the procedure described in the text under his eye, cannot fail to grasp the idea. All extraneous matter and discussions have been carefully excluded, the attempt being made to allow no unnecessary details to cumber the text. The subject-matter is brought up to date at every point, and the work is as nearly as possible the combined opinions of the ten specialists who figure as the authors.

In the revised edition much new material has been added, and some of the old eliminated or modified. More than forty of the old illustrations have been replaced by new ones, which add very materially to the elucidation of the text, as they picture methods, not specimens. The chapters on technique and after-treatment have been considerably enlarged, and the portions devoted to plastic work have been so greatly improved as to be practically new. Hysterectomy has been rewritten, and all the descriptions of operative procedures have been carefully revised and fully illustrated.

CONTRIBUTORS:

Dr. Henry T. Byford.
John M. Baldy.
Edwin Cragin.
J. H. Etheridge.
William Goodell.
Dr. Howard A. Kelly.
Florian Krug.
E. E. Montgomery.
William R. Pryor.
George M. Tuttle.

"The most notable contribution to gynecological literature since 1887, ... and the most complete exponent of gynecology which we have. No subject seems to have been neglected, ... and the gynecologist and surgeon, and the general practitioner who has any desire to practise diseases of women, will find it of practical value. In the matter of illustrations and plates the book surpasses anything we have seen."—*Boston Medical and Surgical Journal.*

"A thoroughly modern text-book, and gives reliable and well-tempered advice and instruction."—*Edinburgh Medical Journal.*

"The harmony of its conclusions and the homogeneity of its style give it an individuality which suggests a single rather than a multiple authorship."—*Annals of Surgery.*

"It must command attention and respect as a worthy representation of our advanced clinical teaching."—*American Journal of Medical Sciences.*

***AN AMERICAN TEXT-BOOK OF THE DISEASES OF CHILDREN. By American Teachers.** Edited by LOUIS STARR, M. D., assisted by THOMPSON S. WESTCOTT, M. D. In one handsome royal-8vo volume of 1244 pages, profusely illustrated with wood-cuts, half-tone and colored plates. Net Prices: Cloth, $7.00; Sheep or Half-Morocco, $8.00.

SECOND EDITION, REVISED AND ENLARGED.

The plan of this work embraces a series of original articles written by some sixty well-known pædiatrists, representing collectively the teachings of the most prominent medical schools and colleges of America. The work is intended to be a PRACTICAL book, suitable for constant and handy reference by the practitioner and the advanced student.

Especial attention has been given to the latest accepted teachings upon the etiology, symptoms, pathology, diagnosis, and treatment of the disorders of children, with the introduction of many special formulæ and therapeutic procedures.

In this new edition the whole subject matter has been carefully revised, new articles added, some original papers emended, and a number entirely rewritten. The new articles include "Modified Milk and Percentage Milk-Mixtures," "Lithemia," and a section on "Orthopedics." Those rewritten are "Typhoid Fever," "Rubella," "Chicken-pox," "Tuberculous Meningitis," "Hydrocephalus," and "Scurvy;" while extensive revision has been made in "Infant Feeding," "Measles," "Diphtheria," and "Cretinism." The volume has thus been much increased in size by the introduction of fresh material.

CONTRIBUTORS:

Dr. S. S. Adams, Washington.
John Ashhurst, Jr., Philadelphia.
A. D. Blackader, Montreal, Canada.
David Bovaird, New York.
Dillon Brown, New York.
Edward M. Buckingham, Boston.
Charles W. Burr, Philadelphia.
W. E. Casselberry, Chicago.
Henry Dwight Chapin, New York.
W. S. Christopher, Chicago.
Archibald Church, Chicago.
Floyd M. Crandall, New York.
Andrew F. Currier, New York.
Roland G. Curtin, Philadelphia
J. M. DaCosta, Philadelphia.
I. N. Danforth, Chicago.
Edward P. Davis, Philadelphia.
John B. Deaver, Philadelphia.
G. E. de Schweinitz, Philadelphia.
John Dorning, New York.
Charles Warrington Earle, Chicago.
Wm. A. Edwards, San Diego, Cal.
F. Forchheimer, Cincinnati.
J. Henry Fruitnight, New York.
J. P. Crozer Griffith, Philadelphia.
W. A. Hardaway. St. Louis.
M. P Hatfield, Chicago.
Barton Cooke Hirst, Philadelphia.
H. Illoway, Cincinnati.
Henry Jackson, Boston.
Charles G. Jennings, Detroit.
Henry Koplik, New York.

Dr. Thomas S. Latimer, Baltimore.
Albert R. Leeds, Hoboken, N. J.
J. Hendrie Lloyd, Philadelphia.
George Roe Lockwood, New York.
Henry M. Lyman, Chicago.
Francis T. Miles, Baltimore.
Charles K Mills, Philadelphia.
James E. Moore, Minneapolis.
F. Gordon Morrill, Boston.
John H. Musser, Philadelphia.
Thomas R. Neilson, Philadelphia.
W. P. Northrup, New York.
William Osler, Baltimore.
Frederick A. Packard, Philadelphia.
William Pepper, Philadelphia.
Frederick Peterson, New York.
W. T. Plant, Syracuse, New York.
William M. Powell, Atlantic City.
B. K. Rachford, Cincinnati.
B. Alexander Randall, Philadelphia.
Edward O. Shakespeare, Philadelphia
F. C. Shattuck, Boston.
J. Lewis Smith, New York.
Louis Starr, Philadelphia.
M. Allen Starr, New York.
Charles W. Townsend, Boston.
James Tyson, Philadelphia.
W. S. Thayer, Baltimore.
Victor C. Vaughan, Ann Arbor, Mich
Thompson S. Westcott, Philadelphia.
Henry R. Wharton, Philadelphia.
J. William White, Philadelphia.
J. C. Wilson, Philadelphia.

*** AN AMERICAN TEXT-BOOK OF GENITO-URINARY AND SKIN DISEASES.** By 47 Eminent Specialists and Teachers. Edited by L. BOLTON BANGS, M. D., Professor of Genito-Urinary Surgery, University and Bellevue Hospital Medical College, New York; and W. A. HARDAWAY, M. D., Professor of Diseases of the Skin, Missouri Medical College. Imperial octavo volume of 1229 pages, with 300 engravings and 20 full-page colored plates. Cloth, $7.00 net; Sheep or Half Morocco, $8.00 net.

This addition to the series of "American Text-Books," it is confidently believed, will meet the requirements of both students and practitioners, giving, as it does, a comprehensive and detailed presentation of the Diseases of the Genito-Urinary Organs, of the Venereal Diseases, and of the Affections of the Skin.

Having secured the collaboration of well-known authorities in the branches represented in the undertaking, the editors have not restricted the contributors in regard to the particular views set forth, but have offered every facility for the free expression of their individual opinions. The work will therefore be found to be original, yet homogeneous and fully representative of the several departments of medical science with which it is concerned.

CONTRIBUTORS:

Dr. Chas. W. Allen, New York.
I. E. Atkinson, Baltimore.
L. Bolton Bangs, New York.
P. R. Bolton, New York.
Lewis C. Bosher, Richmond, Va.
John T. Bowen, Boston.
J. Abbott Cantrell, Philadelphia.
William T. Corlett, Cleveland, Ohio.
B. Farquhar Curtis, New York.
Condict W. Cutler, New York.
Isadore Dyer, New Orleans.
Christian Fenger, Chicago.
John A. Fordyce, New York.
Eugene Fuller, New York.
R. H. Greene, New York.
Joseph Grindon, St. Louis.
Graeme M. Hammond, New York.
W. A. Hardaway, St. Louis.
M. B. Hartzell, Philadelphia.
Louis Heitzmann, New York.
James S. Howe, Boston.
George T. Jackson, New York.
Abraham Jacobi, New York.
James C. Johnston, New York.
Dr. Hermann G. Klotz, New York.
J. H. Linsley, Burlington, Vt.
G. F. Lydston, Chicago.
Hartwell N. Lyon, St. Louis.
Edward Martin, Philadelphia.
D. G. Montgomery, San Francisco.
James Pedersen, New York.
S. Pollitzer, New York.
Thomas R. Pooley, New York.
A. R. Robinson, New York.
A. E. Regensburger, San Francisco.
Francis J. Shepherd, Montreal, Can.
S. C. Stanton, Chicago, Ill.
Emmanuel J. Stout, Philadelphia.
Alonzo E. Taylor Philadelphia.
Robert W. Taylor, New York.
Paul Thorndike, Boston.
H. Tuholske, St. Louis.
Arthur Van Harlingen, Philadelphia.
Francis S. Watson, Boston.
J. William White, Philadelphia.
J. McF. Winfield, Brooklyn.
Alfred C. Wood, Philadelphia.

"This voluminous work is thoroughly up to date, and the chapters on genito-urinary diseases are especially valuable. The illustrations are fine and are mostly original. The section on dermatology is concise and in every way admirable."—*Journal of the American Medical Association.*

"This volume is one of the best yet issued of the publisher's series of 'American Text-Books.' The list of contributors represents an extraordinary array of talent and extended experience. The book will easily take the place in comprehensiveness and value of the half dozen or more costly works on these subjects which have hitherto been necessary to a well-equipped library."—*New York Polyclinic.*

*** AN AMERICAN TEXT-BOOK OF DISEASES OF THE EYE, EAR, NOSE, AND THROAT.** Edited by GEORGE E. DE SCHWEINITZ, A. M., M. D., Professor of Ophthalmology, Jefferson Medical College; and B. ALEXANDER RANDALL, A. M., M. D., Clinical Professor of Diseases of the Ear, University of Pennsylvania. One handsome imperial octavo volume of 1251 pages; 766 illustrations, 59 of them colored. Prices: Cloth, $7.00 net; Sheep or Half-Morocco, $8.00 net.

Just Issued.

The present work is the only book ever published embracing diseases of the intimately related organs of the eye, ear, nose, and throat. Its special claim to favor is based on encyclopedic, authoritative, and practical treatment of the subjects.

Each section of the book has been entrusted to an author who is specially identified with the subject on which he writes, and who therefore presents his case in the manner of an expert. Uniformity is secured and overlapping prevented by careful editing and by a system of cross-references which forms a special feature of the volume, enabling the reader to come into touch with all that is said on any subject in different portions of the book.

Particular emphasis is laid on the most approved methods of treatment, so that the book shall be one to which the student and practitioner can refer for information in practical work. Anatomical and physiological problems, also, are fully discussed for the benefit of those who desire to investigate the more abstruse problems of the subject.

CONTRIBUTORS:

Dr. Henry A. Alderton, Brooklyn.
Harrison Allen, Philadelphia.
Frank Allport, Chicago.
Morris J. Asch, New York.
S. C. Ayres, Cincinnati.
R. O. Beard, Minneapolis.
Clarence J. Blake, Boston.
Arthur A. Bliss, Philadelphia.
Albert P. Brubaker, Philadelphia.
J. H. Bryan, Washington, D. C.
Albert H. Buck, New York.
F. Buller, Montreal, Can.
Swan M. Burnett, Washington, D. C.
Flemming Carrow, Ann Arbor, Mich.
W. E. Casselberry, Chicago.
Colman W. Cutler, New York.
Edward B. Dench, New York.
William S. Dennett, New York.
George E. de Schweinitz, Philadelphia.
Alexander Duane, New York.
John W. Farlow, Boston, Mass.
Walter J. Freeman, Philadelphia.
H. Gifford, Omaha, Neb.
W. C. Glasgow, St. Louis.
J. Orne Green, Boston.
Ward A. Holden, New York.
Christian R. Holmes, Cincinnati.
William E. Hopkins, San Francisco.
F. C. Hotz, Chicago.
Lucien Howe, Buffalo, N. Y.
Dr. Alvin A. Hubbell, Buffalo, N. Y.
Edward Jackson, Philadelphia.
J. Ellis Jennings, St. Louis.
Herman Knapp, New York.
Chas. W. Kollock, Charleston, S. C.
G. A. Leland, Boston.
J. A. Lippincott, Pittsburg, Pa.
G. Hudson Makuen, Philadelphia.
John H. McCollom, Boston.
H. G. Miller, Providence, R. I.
B. L. Milliken, Cleveland, Ohio.
Robert C. Myles, New York.
James E. Newcomb, New York.
R. J. Phillips, Philadelphia.
George A. Piersol, Philadelphia.
W. P. Porcher, Charleston, S. C.
B. Alex. Randall, Philadelphia.
Robert L. Randolph, Baltimore.
John O. Roe, Rochester, N. Y.
Charles E. de M. Sajous, Philadelphia.
J. E. Sheppard, Brooklyn, N. Y.
E. L. Shurly, Detroit, Mich.
William M. Sweet, Philadelphia.
Samuel Theobald, Baltimore, Md.
A. G. Thomson, Philadelphia.
Clarence A. Veasey, Philadelphia.
John E. Weeks, New York.
Casey A. Wood, Chicago, Ill.
Jonathan Wright, Brooklyn.
H. V. Würdemann, Milwaukee, Wis.

*AN AMERICAN YEAR-BOOK OF MEDICINE AND SURGERY. A Yearly Digest of Scientific Progress and Authoritative Opinion in all branches of Medicine and Surgery, drawn from journals, monographs, and text-books of the leading American and Foreign authors and investigators. Collected and arranged, with critical editorial comments, by eminent American specialists and teachers, under the general editorial charge of GEORGE M. GOULD, M. D. Volumes for 1896, '97, '98, and '99 each a handsome imperial octavo volume of about 1200 pages. Prices: Cloth, $6.50 net; Half-Morocco, $7.50 net. Year-Book for 1900 in two octavo volumes of about 600 pages each. Prices per volume: Cloth, $3.00 net; Half-Morocco, $3.75 net.

In Two Volumes. No Increase in Price.

In response to a widespread demand from the medical profession, the publisher of the "American Year-Book of Medicine and Surgery" has decided to issue that well-known work in two volumes, Vol. I. treating of **General Medicine**, Vol. II. of **General Surgery**. Each volume is complete in itself, and the work is sold either separately or in sets.

This division is made in such a way as to appeal to physicians from a class standpoint, one volume being distinctly medical, and the other distinctly surgical. This arrangement has a two-fold advantage. To the physician who uses the entire book, it offers an increased amount of matter in the most convenient form for easy consultation, and without any increase in price; while the man who wants either the medical or the surgical section alone secures the complete consideration of his branch without the necessity of purchasing matter for which he has no use.

CONTRIBUTORS:

VOL. I.

Dr. Samuel W. Abbott, Boston.
Archibald Church, Chicago.
Louis A. Duhring, Philadelphia.
D. L. Edsall, Philadelphia.
Alfred Hand, Jr., Philadelphia.
M. B. Hartzell, Philadelphia.
Reid Hunt, Baltimore.
Wyatt Johnston, Montreal.
Walter Jones, Baltimore.
David Riesman, Philadelphia.
Louis Starr, Philadelphia.
Alfred Stengel, Philadelphia.
A. A. Stevens, Philadelphia.
G. N. Stewart, Cleveland.
Reynold W. Wilcox, New York City.

VOL. II.

Dr. J. Montgomery Baldy, Philadelphia.
Charles H. Burnett, Philadelphia.
J. Chalmers DaCosta, Philadelphia.
W. A. N. Dorland, Philadelphia.
Virgil P. Gibney, New York City.
C. H. Hamann, Cleveland.
Howard F. Hansell, Philadelphia.
Barton Cooke Hirst, Philadelphia.
E. Fletcher Ingals, Chicago.
W. W. Keen, Philadelphia.
Henry G. Ohls, Chicago.
Wendell Reber, Philadelphia.
J. Hilton Waterman, New York City.

"It is difficult to know which to admire most—the research and industry of the distinguished band of experts whom Dr. Gould has enlisted in the service of the Year-Book, or the wealth and abundance of the contributions to every department of science that have been deemed worthy of analysis. . . . It is much more than a mere compilation of abstracts, for, as each section is entrusted to experienced and able contributors, the reader has the advantage of certain critical commentaries and expositions . . . proceeding from writers fully qualified to perform these tasks. . . . It is emphatically a book which should find a place in every medical library, and is in several respects more useful than the famous 'Jahrbucher' of Germany."—*London Lancet.*

*** ANOMALIES AND CURIOSITIES OF MEDICINE.** By GEORGE M. GOULD, M.D., and WALTER L. PYLE, M.D. An encyclopedic collection of are and extraordinary cases and of the most striking instances of abnormality in all branches of Medicine and Surgery, derived from an exhaustive research of medical literature from its origin to the present day, abstracted, classified, annotated, and indexed. Handsome imperial octavo volume of 968 pages, with 295 engravings in the text, and 12 full-page plates. Cloth, $3.00 net; Half-Morocco, $4.00 net.

POPULAR EDITION REDUCED FROM $6.00 to $3.00.

In view of the great success of this magnificent work, the publisher has decided to issue a "Popular Edition" at a price so low that it may be procured by every student and practitioner of medicine. Notwithstanding the great reduction in price, there will be no depreciation in the excellence of typography, paper, and binding that characterized the earlier editions.

Several years of exhaustive research have been spent by the authors in the great medical libraries of the United States and Europe in collecting the material for this work. **Medical literature of all ages and all languages** has been carefully searched, as a glance at the Bibliographic Index will show. The facts, which will be of **extreme value to the author and lecturer,** have been arranged and annotated, and full reference footnotes given.

"One of the most valuable contributions ever made to medical literature. It is, so far as we know, absolutely unique, and every page is as fascinating as a novel. Not alone for the medical profession has this volume value: it will serve as a book of reference for all who are interested in general scientific, sociologic, or medico-legal topics."—*Brooklyn Medical Journal.*

NERVOUS AND MENTAL DISEASES. By ARCHIBALD CHURCH, M. D., Professor of Clinical Neurology, Mental Diseases, and Medical Jurisprudence, Northwestern University Medical School; and FREDERICK PETERSON, M. D., Clinical Professor of Mental Diseases, Woman's Medical College, New York. Handsome octavo volume of 843 pages, with over 300 illustrations. Prices: Cloth, $5.00 net; Half-Morocco, $6.00 net.

Second Edition.

This book is intended to furnish students and practitioners with a practical, working knowledge of nervous and mental diseases. Written by men of wide experience and authority, it presents the many recent additions to the subject. The book is not filled with an extended dissertation on anatomy and pathology, but, treating these points in connection with special conditions, it lays particular stress on methods of examination, diagnosis, and treatment. In this respect the work is unusually complete and valuable, laying down the definite courses of procedure which the authors have found to be most generally satisfactory.

"The work is an epitome of what is to-day known of nervous diseases prepared for the student and practitioner in the light of the author's experience . . . We believe that no work presents the difficult subject of insanity in such a reasonable and readable way."—*Chicago Medical Recorder.*

DISEASES OF THE NOSE AND THROAT. By D. Braden Kyle, M. D., Clinical Professor of Laryngology and Rhinology, Jefferson Medical College, Philadelphia; Consulting Laryngologist, Rhinologist, and Otologist, St. Agnes' Hospital. Octavo volume of 646 pages, with over 150 illustrations and 6 lithographic plates. Cloth, $4.00 net; Half-Morocco, $5.00 net.

Just Issued.

This book presents the subject of Diseases of the Nose and Throat in as concise a manner as is consistent with clearness, keeping in mind the needs of the student and general practitioner as well as those of the specialist. The arrangement and classification are based on modern pathology, and the pathological views advanced are supported by drawings of microscopical sections made in the author's own laboratory. These and the other illustrations are particularly fine, being chiefly original. With the practical purpose of the book in mind, extended consideration has been given to details of treatment, each disease being considered in full, and definite courses being laid down to meet special conditions and symptoms.

"It is a thorough, full, and systematic treatise, so classified and arranged as greatly to facilitate the teaching of laryngology and rhinology to classes, and must prove most convenient and satisfactory as a reference book, both for students and practitioners."—*International Medical Magazine.*

THE HYGIENE OF TRANSMISSIBLE DISEASES; their Causation, Modes of Dissemination, and Methods of Prevention. By A. C. Abbott, M. D., Professor of Hygiene in the University of Pennsylvania; Director of the Laboratory of Hygiene. Octavo volume of 311 pages, with charts and maps, and numerous illustrations. Cloth, $2.00 net.

Just Issued.

It is not the purpose of this work to present the subject of Hygiene in the comprehensive sense ordinarily implied by the word, but rather to deal directly with but a section, certainly not the least important, of the subject—viz., that embracing a knowledge of the preventable specific diseases. The book aims to furnish information concerning the detailed management of transmissible diseases. Incidentally there are discussed those numerous and varied factors that have not only a direct bearing upon the incidence and suppression of such diseases, but are of general sanitary importance as well.

"The work is admirable in conception and no less so in execution. It is a practical work, simply and lucidly written, and it should prove a most helpful aid in that department of medicine which is becoming daily of increasing importance and application—namely, prophylaxis."—*Philadelphia Medical Journal.*

"It is scientific, but not too technical; it is as complete as our present-day knowledge of hygiene and sanitation allows, and it is in harmony with the efforts of the profession, which are tending more and more to methods of prophylaxis. For the student and for the practitioner it is well nigh indispensable."—*Medical News,* New York.

A TEXT-BOOK OF EMBRYOLOGY. By JOHN C. HEISLER, M. D., Professor of Anatomy in the Medico-Chirurgical College, Philadelphia. Octavo volume of 405 pages, with 190 illustrations, 26 in colors. Cloth, $2.50 net.

Just Issued.

The facts of embryology having acquired in recent years such great interest in connection with the teaching and with the proper comprehension of human anatomy, it is of first importance to the student of medicine that a concise and yet sufficiently full text-book upon the subject be available. It was with the aim of presenting such a book that this volume was written, the author, in his experience as a teacher of anatomy, having been impressed with the fact that students were seriously handicapped in their study of the subject of embryology by the lack of a text-book full enough to be intelligible, and yet without that minuteness of detail which characterizes the larger treatises, and which so often serves only to confuse and discourage the beginner.

"In short, the book is written to fill a want which has distinctly existed and which it definitely meets; commendation greater than this it is not possible to give to anything."—*Medical News*, New York.

A MANUAL OF DISEASES OF THE EYE. By EDWARD JACKSON, A. M., M. D., sometime Professor of Diseases of the Eye in the Philadelphia Polyclinic and College for Graduates in Medicine. 12mo, 604 pages, with 178 illustrations from drawings by the author. Cloth, $2.50 net.

Just Issued.

This book is intended to meet the needs of the general practitioner of medicine and the beginner in ophthalmology. More attention is given to the conditions that must be met and dealt with early in ophthalmic practice than to the rarer diseases and more difficult operations that may come later.

It is designed to furnish efficient aid in the actual work of dealing with disease, and therefore gives the place of first importance to the recognition and management of the conditions that present themselves in actual clinical work.

LECTURES ON THE PRINCIPLES OF SURGERY. By CHARLES B. NANCREDE, M. D., LL.D., Professor of Surgery and of Clinical Surgery, University of Michigan, Ann Arbor. Handsome octavo, 398 pages, illustrated. Cloth, $2.50 net.

Just Issued.

The present book is based on the lectures delivered by Dr. Nancrede to his undergraduate classes, and is intended as a text-book for students and a practical help for teachers. By the careful elimination of unnecessary details of pathology, bacteriology, etc., which are amply provided for in other courses of study, space is gained for a more extended consideration of the Principles of Surgery in themselves, and of the application of these principles to methods of practice.

A TEXT-BOOK OF PATHOLOGY. By ALFRED STENGEL, M. D., Professor of Clinical Medicine in the University of Pennsylvania; Physician to the Philadelphia Hospital; Physician to the Children's Hospital, Philadelphia. Handsome octavo volume of 848 pages, with 362 illustrations, many of which are in colors. Prices: Cloth, $4.00 net; Half-Morocco, $5.00 net.

Second Edition.

In this work the practical application of pathological facts to clinical medicine is considered more fully than is customary in works on pathology. While the subject of pathology is treated in the broadest way consistent with the size of the book, an effort has been made to present the subject from the point of view of the clinician. The general relations of bacteriology to pathology are discussed at considerable length, as the importance of these branches deserves. It will be found that the recent knowledge is fully considered, as well as older and more widely-known facts.

"I consider the work abreast of modern pathology, and useful to both students and practitioners. It presents in a concise and well-considered form the essential facts of general and special pathological anatomy, with more than usual emphasis upon pathological physiology." —WILLIAM H. WELCH, *Professor of Pathology, Johns Hopkins University, Baltimore, Md.*

"I regard it as the most serviceable text-book for students on this subject yet written by an American author."—L. HEKTOEN, *Professor of Pathology, Rush Medical College, Chicago, Ill.*

A TEXT-BOOK OF OBSTETRICS. By BARTON COOKE HIRST, M.D., Professor of Obstetrics in the University of Pennsylvania. Handsome octavo volume of 846 pages, with 618 illustrations and seven colored plates. Prices: Cloth, $5.00 net; Half-Morocco, $6.00 net.

Second Edition.

This work, which has been in course of preparation for several years, is intended as an ideal text-book for the student no less than an advanced treatise for the obstetrician and for general practitioners. It represents the very latest teaching in the practice of obstetrics by a man of extended experience and recognized authority. The book emphasizes especially, as a work on obstetrics should, the practical side of the subject, and to this end presents an unusually large collection of illustrations. A great number of these are new and original, and the whole collection will form a complete atlas of obstetrical practice. An extremely valuable feature of the book is the large number of references to cases, authorities, sources, etc., forming, as it does, a valuable bibliography of the most recent and authoritative literature on the subject of obstetrics. As already stated, this work records the wide practical experience of the author, which fact, combined with the brilliant presentation of the subject, will doubtless render this one of the most notable books on obstetrics that has yet appeared.

"The illustrations are numerous and are works of art, many of them appearing for the first time. The arrangement of the subject-matter, the foot-notes, and index are beyond criticism. The author's style, though condensed, is singularly clear, so that it is never necessary to re-read a sentence in order to grasp its meaning. As a true model of what a modern text-book in obstetrics should be, we feel justified in affirming that Dr. Hirst's book is without a rival."—*New York Medical Record.*

A TEXT-BOOK OF THE PRACTICE OF MEDICINE. By JAMES M. ANDERS, M.D., PH.D., LL.D., Professor of the Practice of Medicine and of Clinical Medicine, Medico-Chirurgical College, Philadelphia. In one handsome octavo volume of 1292 pages, fully illustrated. Cloth, $5.50 net; Sheep or Half-Morocco, $6.50 net.

THIRD EDITION, THOROUGHLY REVISED.

The present edition is the result of a careful and thorough revision. A few new subjects have been introduced: Glandular Fever, Ether-pneumonia, Splenic Anemia, Meralgia Paresthetica, and Periodic Paralysis. The affections that have been substantially rewritten are: Plague, Malta Fever, Diseases of the Thymus Gland, Liver Cirrhoses, and Progressive Spinal Muscular Atrophy. The following articles have been extensively revised: Typhoid Fever, Yellow Fever, Lobar Pneumonia, Dengue, Tuberculosis, Diabetes Mellitus, Gout, Arthritis Deformans, Autumnal Catarrh, Diseases of the Circulatory System, more particularly Hypertrophy and Dilatation of the Heart, Arteriosclerosis and Thoracic Aneurysm, Pancreatic Hemorrhage, Jaundice, Acute Peritonitis, Acute Yellow Atrophy, Hematoma of Dura Mater, and Scleroses of the Brain. The preliminary chapter on Nervous Diseases is new, and deals with the subject of localization and the various methods of investigating nervous affections.

"It is an excellent book—concise, comprehensive, thorough, and up to date. It is a credit to you; but, more than that, it is a credit to the profession of Philadelphia—to us." —JAMES C. WILSON, *Professor of the Practice of Medicine and Clinical Medicine, Jefferson Medical College, Philadelphia.*

"The book can be unreservedly recommended to students and practitioners as a safe, full compendium of the knowledge of internal medicine of the present day . . . It is a work thoroughly modern in every sense."—*Medical News,* New York.

DISEASES OF THE STOMACH. By WILLIAM W. VAN VALZAH, M. D., Professor of General Medicine and Diseases of the Digestive System and the Blood, New York Polyclinic; and J. DOUGLAS NISBET, M. D., Adjunct Professor of General Medicine and Diseases of the Digestive System and the Blood, New York Polyclinic. Octavo volume of 674 pages, illustrated. Cloth, $3.50 net.

An eminently practical book, intended as a guide to the student, an aid to the physician, and a contribution to scientific medicine. It aims to give a complete description of the modern methods of diagnosis and treatment of diseases of the stomach, and to reconstruct the pathology of the stomach in keeping with the revelations of scientific research. The book is clear, practical, and complete, and contains the results of the authors' investigations and of their extensive experience as specialists. Particular attention is given to the important subject of dietetic treatment. The diet-lists are very complete, and are so arranged that selections can readily be made to suit individual cases.

"This is the most satisfactory work on the subject in the English language."—*Chicago Medical Recorder.*

"The article on diet and general medication is one of the most valuable in the book, and should be read by every practising physician."—*New York Medical Journal.*

SURGICAL DIAGNOSIS AND TREATMENT. By J. W. MACDONALD, M. D., Edin., F. R. C. S., Edin., Professor of the Practice of Surgery and of Clinical Surgery in Hamline University; Visiting Surgeon to St. Barnabas' Hospital, Minneapolis, etc. Handsome octavo volume of 800 pages, profusely illustrated. Cloth, $5.00 net; Half-Morocco, $6.00 net.

This work aims in a comprehensive manner to furnish a guide in matters of surgical diagnosis. It sets forth in a systematic way the necessities of examinations and the proper methods of making them. The various portions of the body are then taken up in order and the diseases and injuries thereof succinctly considered and the treatment briefly indicated. Practically all the modern and approved operations are described with thoroughness and clearness. The work concludes with a chapter on the use of the Röntgen rays in surgery.

"The work is brimful of just the kind of practical information that is useful alike to students and practitioners. It is a pleasure to commend the book because of its intrinsic value to the medical practitioner."—*Cincinnati Lancet-Clinic.*

PATHOLOGICAL TECHNIQUE. A Practical Manual for Laboratory Work in Pathology, Bacteriology, and Morbid Anatomy, with chapters on Post-Mortem Technique and the Performance of Autopsies. By FRANK B. MALLORY, A. M., M. D., Assistant Professor of Pathology, Harvard University Medical School, Boston; and JAMES H. WRIGHT, A. M., M. D., Instructor in Pathology, Harvard University Medical School, Boston. Octavo volume of 396 pages, handsomely illustrated. Cloth, $2.50 net.

This book is designed especially for practical use in pathological laboratories, both as a guide to beginners and as a source of reference for the advanced. The book will also meet the wants of practitioners who have opportunity to do general pathological work. Besides the methods of post-mortem examinations and of bacteriological and histological investigations connected with autopsies, the special methods employed in clinical bacteriology and pathology have been fully discussed.

"One of the most complete works on the subject, and one which should be in the library of every physician who hopes to keep pace with the great advances made in pathology."—*Journal of American Medical Association.*

THE SURGICAL COMPLICATIONS AND SEQUELS OF TYPHOID FEVER. By WM. W. KEEN, M. D., LL.D., Professor of the Principles of Surgery and of Clinical Surgery, Jefferson Medical College, Philadelphia. Octavo volume of 386 pages, illustrated. Cloth, $3.00 net.

This monograph is the only one in any language covering the entire subject of the Surgical Complications and Sequels of Typhoid Fever. The work will prove to be of importance and interest not only to the general surgeon and physician, but also to many specialists—laryngologists, ophthalmologists, gynecologists, pathologists, and bacteriologists—as the subject has an important bearing upon each one of their spheres. The author's conclusions are based on reports of over 1700 cases, including practically all those recorded in the last fifty years. Reports of cases have been brought down to date, many having been added while the work was in press.

"This is probably the first and only work in the English language that gives the reader a clear view of what typhoid fever really is, and what it does and can do to the human organism. This book should be in the possession of every medical man in America."—*American Medico-Surgical Bulletin.*

MODERN SURGERY, GENERAL AND OPERATIVE. By JOHN CHALMERS DACOSTA, M.D., Clinical Professor of Surgery, Jefferson Medical College, Philadelphia; Surgeon to the Philadelphia Hospital, etc. Handsome octavo volume of 911 pages, profusely illustrated. Cloth, $4.00 net; Half-Morocco, $5.00 net.

Second Edition, Rewritten and Greatly Enlarged.

The remarkable success attending DaCosta's Manual of Surgery, and the general favor with which it has been received, have led the author in this revision to produce a complete treatise on modern surgery along the same lines that made the former edition so successful. The book has been entirely rewritten and very much enlarged. The old edition has long been a favorite not only with students and teachers, but also with practising physicians and surgeons, and it is believed that the present work will find an even wider field of usefulness.

"We know of no small work on surgery in the English language which so well fulfils the requirements of the modern student."—*Medico-Chirurgical Journal*, Bristol, England.

"The author has presented concisely and accurately the principles of modern surgery. The book is a valuable one which can be recommended to students and is of great value to the general practitioner."—*American Journal of the Medical Sciences.*

A MANUAL OF ORTHOPEDIC SURGERY. By JAMES E. MOORE, M.D., Professor of Orthopedics and Adjunct Professor of Clinical Surgery, University of Minnesota, College of Medicine and Surgery. Octavo volume of 356 pages, with 177 beautiful illustrations from photographs made specially for this work. Cloth, $2.50 net.

A practical book based upon the author's experience, in which special stress is laid upon early diagnosis and treatment such as can be carried out by the general practitioner. The teachings of the author are in accordance with his belief that true conservatism is to be found in the middle course between the surgeon who operates too frequently and the orthopedist who seldom operates.

"A very demonstrative work, every illustration of which conveys a lesson. The work is a most excellent and commendable one, which we can certainly endorse with pleasure."—*St. Louis Medical and Surgical Journal.*

ELEMENTARY BANDAGING AND SURGICAL DRESSING. With Directions concerning the Immediate Treatment of Cases of Emergency. For the use of Dressers and Nurses. By WALTER PYE, F.R.C.S., late Surgeon to St. Mary's Hospital, London. Small 12mo, with over 80 illustrations. Cloth, flexible covers, 75 cents net.

This little book is chiefly a condensation of those portions of Pye's "Surgical Handicraft" which deal with bandaging, splinting, etc., and of those which treat of the management in the first instance of cases of emergency. The directions given are thoroughly practical, and the book will prove extremely useful to students, surgical nurses, and dressers.

"The author writes well, the diagrams are clear, and the book itself is small and portable, although the paper and type are good."—*British Medical Journal.*

A TEXT-BOOK OF MATERIA MEDICA, THERAPEUTICS AND PHARMACOLOGY. By GEORGE F. BUTLER, PH.G., M.D., Professor of Materia Medica and of Clinical Medicine in the College of Physicians and Surgeons, Chicago; Professor of Materia Medica and Therapeutics, Northwestern University, Woman's Medical School, etc. Octavo, 874 pages, illustrated. Cloth, $4.00 net; Sheep, $5.00 net.

Third Edition, Thoroughly Revised.

A clear, concise, and practical text-book, adapted for permanent reference no less than for the requirements of the class-room.

The recent important additions made to our knowledge of the physiological action of drugs are fully discussed in the present edition. The book has been thoroughly revised and many additions have been made.

"Taken as a whole, the book may fairly be considered as one of the most satisfactory of any single-volume works on materia medica in the market."—*Journal of the American Medical Association.*

TUBERCULOSIS OF THE GENITO-URINARY ORGANS, MALE AND FEMALE. By NICHOLAS SENN, M.D., PH.D., LL.D., Professor of the Practice of Surgery and of Clinical Surgery, Rush Medical College, Chicago. Handsome octavo volume of 320 pages, illustrated. Cloth, $3.00 net.

Tuberculosis of the male and female genito-urinary organs is such a frequent, distressing, and fatal affection that a special treatise on the subject appears to fill a gap in medical literature. In the present work the bacteriology of the subject has received due attention, the modern resources employed in the differential diagnosis between tubercular and other inflammatory affections are fully described, and the medical and surgical therapeutics are discussed in detail.

"An important book upon an important subject, and written by a man of mature judgment and wide experience. The author has given us an instructive book upon one of the most important subjects of the day."—*Clinical Reporter.*

"A work which adds another to the many obligations the profession owes the talented author."—*Chicago Medical Recorder.*

A TEXT-BOOK OF DISEASES OF WOMEN. By CHARLES B. PENROSE, M.D., PH.D., Professor of Gynecology in the University of Pennsylvania; Surgeon to the Gynecean Hospital, Philadelphia. Octavo volume of 531 pages, with 217 illustrations, nearly all from drawings made for this work. Cloth, $3.75 net.

Third Edition, Revised.

In this work, which has been written for both the student of gynecology and the general practitioner, the author presents the best teaching of modern gynecology untrammelled by antiquated theories or methods of treatment. In most instances but one plan of treatment is recommended, to avoid confusing the student or the physician who consults the book for practical guidance.

"I shall value very highly the copy of Penrose's 'Diseases of Women' received. I have already recommended it to my class as THE BEST book."—HOWARD A. KELLY, *Professor of Gynecology and Obstetrics, Johns Hopkins University, Baltimore, Md.*

"The book is to be commended without reserve, not only to the student but to the general practitioner who wishes to have the latest and best modes of treatment explained with absolute clearness."—*Therapeutic Gazette.*

SURGICAL PATHOLOGY AND THERAPEUTICS. By JOHN COLLINS WARREN, M. D., LL.D., Professor of Surgery, Medical Department Harvard University. Handsome octavo, 832 pages, with 136 relief and lithographic illustrations, 33 of which are printed in colors.

Second Edition,

with an Appendix devoted to the Scientific Aids to Surgical Diagnosis, and a series of articles on Regional Bacteriology. Cloth, $5.00 net; Half-Morocco, $6.00 net.

Without Exception, the Illustrations are the Best ever Seen in a Work of this Kind.

"A most striking and very excellent feature of this book is its illustrations. Without exception, from the point of accuracy and artistic merit, they are the best ever seen in a work of this kind. * * * Many of those representing microscopic pictures are so perfect in their coloring and detail as almost to give the beholder the impression that he is looking down the barrel of a microscope at a well-mounted section."—*Annals of Surgery*, Philadelphia.

"It is the handsomest specimen of book-making * * * that has ever been issued from the American medical press."—*American Journal of the Medical Sciences*, Philadelphia.

PATHOLOGY AND SURGICAL TREATMENT OF TUMORS. By N. SENN, M. D., Ph. D., LL. D., Professor of Practice of Surgery and of Clinical Surgery, Rush Medical College; Professor of Surgery, Chicago Polyclinic; Attending Surgeon to Presbyterian Hospital; Surgeon-in-Chief, St. Joseph's Hospital, Chicago. One volume of 710 pages, with 515 engravings, including full-page colored plates. New and enlarged Edition in Preparation.

Books specially devoted to this subject are few, and in our text-books and systems of surgery this part of surgical pathology is usually condensed to a degree incompatible with its scientific and clinical importance. The author spent many years in collecting the material for this work, and has taken great pains to present it in a manner that should prove useful as a text-book for the student, a work of reference for the practitioner, and a reliable guide for the surgeon.

"The most exhaustive of any recent book in English on this subject. It is well illustrated, and will doubtless remain as the principal monograph on the subject in our language for some years. The book is handsomely illustrated and printed, ... and the author has given a notable and lasting contribution to surgery."—*Journal of the American Medical Association*, Chicago.

LECTURES ON RENAL AND URINARY DISEASES. By ROBERT SAUNDBY, M. D., Edin., Fellow of the Royal College of Physicians, London, and of the Royal Medico-Chirurgical Society; Physician to the General Hospital. Octavo volume of 434 pages, with numerous illustrations and 4 colored plates. Cloth, $2.50 net.

"The volume makes a favorable impression at once. The style is clear and succinct. We cannot find any part of the subject in which the views expressed are not carefully thought out and fortified by evidence drawn from the most recent sources. The book may be cordially recommended."—*British Medical Journal.*

A HANDBOOK FOR NURSES. By J. K. WATSON, M. D., Edin., Assistant House-Surgeon, Sheffield Royal Hospital. American Edition, under the supervision of A. A. STEVENS, A. M., M. D., Professor of Pathology, Woman's Medical College, Philadelphia. 12mo, 413 pages, 73 illustrations. Cloth, $1.50 net.

This work aims to supply in one volume that information which so many nurses at the present time are trying to extract from various medical works, and to present that information in a suitable form. Nurses must necessarily acquire a certain amount of medical knowledge, and the author of this book has aimed judiciously to cater to this need with the object of directing the nurses' pursuit of medical information in proper and legitimate channels. The book represents an entirely new departure in nursing literature, insomuch as it contains useful information on medical and surgical matters hitherto only to be obtained from expensive works written expressly for medical men.

A NEW PRONOUNCING DICTIONARY OF MEDICINE, with Phonetic Pronunciation, Accentuation, Etymology, etc. By JOHN M. KEATING, M. D., LL.D., Fellow of the College of Physicians of Philadelphia; Editor "Cyclopædia of the Diseases of Children," etc.; and HENRY HAMILTON, with the Collaboration of J. CHALMERS DACOSTA, M. D., and FREDERICK A. PACKARD, M. D. One very attractive volume of over 800 pages. Second Revised Edition. Prices: Cloth, $5.00 net; Sheep or Half-Morocco, $6.00 net; with Denison's Patent Ready-Reference Index; without patent index, Cloth, $4.00 net; Sheep or Half-Morocco, $5.00 net.

PROFESSIONAL OPINIONS.

"I am much pleased with Keating's Dictionary, and shall take pleasure in recommending it to my classes."

HENRY M. LYMAN, M. D.,
Professor of Principles and Practice of Medicine, Rush Medical College, Chicago, Ill.

"I am convinced that it will be a very valuable adjunct to my study-table, convenient in size and sufficiently full for ordinary use."

C. A. LINDSLEY, M. D.,
Professor of Theory and Practice of Medicine, Medical Dept. Yale University; Secretary Connecticut State Board of Health, New Haven, Conn.

AUTOBIOGRAPHY OF SAMUEL D. GROSS, M. D., Emeritus Professor of Surgery in the Jefferson Medical College of Philadelphia, with Reminiscences of His Times and Contemporaries. Edited by his sons, SAMUEL W. GROSS, M. D., LL.D., and A. HALLER GROSS, A. M., of the Philadelphia Bar. Preceded by a Memoir of Dr. Gross, by the late Austin Flint, M. D., LL.D. In two handsome volumes, each containing over 400 pages, demy 8vo, extra cloth, gilt tops, with fine Frontispiece engraved on steel. Price per Volume, $2.50 net.

PRACTICAL POINTS IN NURSING. For Nurses in Private Practice. By EMILY A. M. STONEY, Graduate of the Training-School for Nurses, Lawrence, Mass.; Superintendent of the Training-School for Nurses, Carney Hospital, South Boston, Mass. 456 pages, handsomely illustrated with 73 engravings in the text, and 9 colored and half-tone plates. Cloth. Price, $1.75 net.

SECOND EDITION, THOROUGHLY REVISED.

In this volume the author explains, in popular language and in the shortest possible form, the entire range of *private* nursing as distinguished from *hospital* nursing, and the nurse is instructed how best to meet the various emergencies of medical and surgical cases when distant from medical or surgical aid or when thrown on her own resources.

An especially valuable feature of the work will be found in the directions to the nurse how to *improvise* everything ordinarily needed in the sick-room, where the embarrassment of the nurse, owing to the want of proper appliances, is frequently extreme.

The work has been logically divided into the following sections:

I. The Nurse: her responsibilities, qualifications, equipment, etc.
II. The Sick-Room: its selection, preparation, and management.
III. The Patient: duties of the nurse in medical, surgical, obstetric, and gynecologic cases.
IV. Nursing in Accidents and Emergencies.
V. Nursing in Special Medical Cases.
VI. Nursing of the New-born and Sick Children.
VII. Physiology and Descriptive Anatomy.

The APPENDIX contains much information in compact form that will be found of great value to the nurse, including Rules for Feeding the Sick; Recipes for Invalid Foods and Beverages; Tables of Weights and Measures; Table for Computing the Date of Labor; List of Abbreviations; Dose-List; and a full and complete Glossary of Medical Terms and Nursing Treatment.

"This is a well-written, eminently practical volume, which covers the entire range of private nursing as distinguished from hospital nursing, and instructs the nurse how best to meet the various emergencies which may arise and how to prepare everything ordinarily needed in the illness of her patient."—*American Journal of Obstetrics and Diseases of Women and Children*, Aug., 1896.

A TEXT-BOOK OF BACTERIOLOGY, including the Etiology and Prevention of Infective Diseases and an account of Yeasts and Moulds, Hæmatozoa, and Psorosperms. By EDGAR M. CROOKSHANK, M. B., Professor of Comparative Pathology and Bacteriology, King's College, London. A handsome octavo volume of 700 pages, with 273 engravings in the text, and 22 original and colored plates. Price, $6.50 net.

This book, though nominally a Fourth Edition of Professor Crookshank's "MANUAL OF BACTERIOLOGY," is practically a new work, the old one having been reconstructed, greatly enlarged, revised throughout, and largely rewritten, forming a text-book for the Bacteriological Laboratory, for Medical Officers of Health, and for Veterinary Inspectors.

MEDICAL DIAGNOSIS. By Dr. OSWALD VIERORDT, Professor of Medicine at the University of Heidelberg. Translated, with additions, from the Fifth Enlarged German Edition, with the author's permission, by FRANCIS H. STUART, A. M., M. D. In one handsome royal-octavo volume of 600 pages. 194 fine wood-cuts in the text, many of them in colors. Prices: Cloth, $4.00 net; Sheep or Half-Morocco, $5.00 net.

FOURTH AMERICAN EDITION, FROM THE FIFTH REVISED AND ENLARGED GERMAN EDITION.

In this work, as in no other hitherto published, are given full and accurate explanations of the phenomena observed at the bedside. It is distinctly a clinical work by a master teacher, characterized by thoroughness, fulness, and accuracy. It is a mine of information upon the points that are so often passed over without explanation. Especial attention has been given to the germ-theory as a factor in the origin of disease.

The present edition of this highly successful work has been translated from the fifth German edition. Many alterations have been made throughout the book, but especially in the sections on Gastric Digestion and the Nervous System.

It will be found that all the qualities which served to make the earlier editions so acceptable have been developed with the evolution of the work to its present form.

THE PICTORIAL ATLAS OF SKIN DISEASES AND SYPHILITIC AFFECTIONS. (American Edition.) Translation from the French. Edited by J. J. PRINGLE, M. B., F. R. C. P., Assistant Physician to, and Physician to the department for Diseases of the Skin at, the Middlesex Hospital, London. Photo-lithochromes from the famous models of dermatological and syphilitic cases in the Museum of the Saint-Louis Hospital, Paris, with explanatory wood-cuts and letter-press. In 12 Parts, at $3.00 per Part.

"Of all the atlases of skin diseases which have been published in recent years, the present one promises to be of greatest interest and value, especially from the standpoint of the general practitioner."—*American Medico-Surgical Bulletin*, Feb. 22, 1896.

"The introduction of explanatory wood-cuts in the text is a novel and most important feature which greatly furthers the easier understanding of the excellent plates, than which nothing, we venture to say, has been seen better in point of correctness, beauty, and general merit."—*New York Medical Journal*, Feb. 15, 1896.

"An interesting feature of the Atlas is the descriptive text, which is written for each picture by the physician who treated the case or at whose instigation the models have been made. We predict for this truly beautiful work a large circulation in all parts of the medical world where the names *St. Louis* and *Baretta* have preceded it."—*Medical Record*, N. Y., Feb. 1, 1896.

A TEXT-BOOK OF MECHANO-THERAPY (MASSAGE AND MEDICAL GYMNASTICS). By AXEL V. GRAFSTROM, B. Sc., M. D., late Lieutenant in the Royal Swedish Army; late House Physician, City Hospital, Blackwell's Island, New York. 12mo, 139 pages, illustrated. Cloth, $1.00 net.

DISEASES OF THE EYE. A Hand-Book of Ophthalmic Practice. By G. E. DE SCHWEINITZ, M. D., Professor of Ophthalmology in the Jefferson Medical College, Philadelphia, etc. A handsome royal-octavo volume of 696 pages, with 255 fine illustrations, many of which are original, and 2 chromo-lithographic plates. Prices: Cloth, $4.00 net; Sheep or Half-Morocco, $5.00 net.

THIRD EDITION, THOROUGHLY REVISED.

In the third edition of this text-book, destined, it is hoped, to meet the favorable reception which has been accorded to its predecessors, the work has been revised thoroughly, and much new matter has been introduced. Particular attention has been given to the important relations which micro-organisms bear to many ocular diseases. A number of special paragraphs on new subjects have been introduced, and certain articles, including a portion of the chapter on Operations, have been largely rewritten, or at least materially changed. A number of new illustrations have been added. The Appendix contains a full description of the method of determining the corneal astigmatism with the ophthalmometer of Javal and Schiötz, and the rotation of the eyes with the tropometer of Stevens.

"A work that will meet the requirements not only of the specialist, but of the general practitioner in a rare degree. I am satisfied that unusual success awaits it."

WILLIAM PEPPER, M. D.
Provost and Professor of Theory and Practice of Medicine and Clinical Medicine in the University of Pennsylvania.

"A clearly written, comprehensive manual. . . . One which we can commend to students as a reliable text-book, written with an evident knowledge of the wants of those entering upon the study of this special branch of medical science."—*British Medical Journal.*

"It is hardly too much to say that for the student and practitioner beginning the study of Ophthalmology, it is the best single volume at present published."—*Medical News.*

"It is a very useful, satisfactory, and safe guide for the student and the practitioner, and one of the best works of this scope in the English language."—*Annals of Ophthalmology.*

DISEASES OF WOMEN. By J. BLAND SUTTON, F. R. C. S., Assistant Surgeon to Middlesex Hospital, and Surgeon to Chelsea Hospital, London; and ARTHUR E. GILES, M. D., B. Sc., Lond., F. R. C. S., Edin., Assistant Surgeon to Chelsea Hospital, London. 436 pages, handsomely illustrated. Cloth, $2.50 net.

The authors have placed in the hands of the physician and student a concise yet comprehensive guide to the study of gynecology in its most modern development. It has been their aim to relate facts and describe methods belonging to the science and art of gynecology in a way that will prove useful to students for examination purposes, and which will also enable the general physician to practice this important department of surgery with advantage to his patients and with satisfaction to himself.

"The book is very well prepared, and is certain to be well received by the medical public."—*British Medical Journal.*

"The text has been carefully prepared. Nothing essential has been omitted, and its teachings are those recommended by the leading authorities of the day."—*Journal of the American Medical Association.*

TEXT-BOOK UPON THE PATHOGENIC BACTERIA. Specially written for Students of Medicine. By JOSEPH MCFARLAND, M. D., Professor of Pathology and Bacteriology in the Medico-Chirurgical College of Philadelphia, etc. 497 pages, finely illustrated. Price, Cloth, $2.50 net.

SECOND EDITION, REVISED AND GREATLY ENLARGED.

The work is intended to be a text-book for the medical student and for the practitioner who has had no recent laboratory training in this department of medical science. The instructions given as to needed apparatus, cultures, stainings, microscopic examinations, etc. are ample for the student's needs, and will afford to the physician much information that will interest and profit him relative to a subject which modern science shows to go far in explaining the etiology of many diseased conditions.

In this second edition the work has been brought up to date in all departments of the subject, and numerous additions have been made to the technique in the endeavor to make the book fulfil the double purpose of a systematic work upon bacteria and a laboratory guide.

"It is excellently adapted for the medical students and practitioners for whom it is avowedly written. . . . The descriptions given are accurate and readable, and the book should prove useful to those for whom it is written.—*London Lancet*, Aug. 29, 1896.

"The author has succeded admirably in presenting the essential details of bacteriological technics, together with a judiciously chosen summary of our present knowledge of pathogenic bacteria. . . . The work, we think, should have a wide circulation among English-speaking students of medicine."—*N. Y. Medical Journal*, April 4, 1896.

"The book will be found of considerable use by medical men who have not had a special bacteriological training, and who desire to understand this important branch of medical science."—*Edinburgh Medical Journal*, July, 1896.

LABORATORY GUIDE FOR THE BACTERIOLOGIST. By LANGDON FROTHINGHAM, M. D. V., Assistant in Bacteriology and Veterinary Science, Sheffield Scientific School, Yale University. Illustrated. Price, Cloth, 75 cents.

The technical methods involved in bacteria-culture, methods of staining, and microscopical study are fully described and arranged as simply and concisely as possible. The book is especially intended for use in laboratory work.

"It is a convenient and useful little work, and will more than repay the outlay necessary for its purchase in the saving of time which would otherwise be consumed in looking up the various points of technique so clearly and concisely laid down in its pages."—*American Med.-Surg. Bulletin.*

FEEDING IN EARLY INFANCY. By ARTHUR V. MEIGS, M. D. Bound in limp cloth, flush edges. Price, 25 cents net.

SYNOPSIS: Analyses of Milk—Importance of the Subject of Feeding in Early Infancy—Proportion of Casein and Sugar in Human Milk—Time to Begin Artificial Feeding of Infants—Amount of Food to be Administered at Each Feeding—Intervals between Feedings—Increase in Amount of Food at Different Periods of Infant Development—Unsuitableness of Condensed Milk as a Substitute for Mother's Milk—Objections to Sterilization or "Pasteurization" of Milk—Advances made in the Method of Artificial Feeding of Infants.

MATERIA MEDICA FOR NURSES. By EMILY A. M. STONEY, Graduate of the Training-school for Nurses, Lawrence, Mass.; late Superintendent of the Training-school for Nurses, Carney Hospital, South Boston, Mass. Handsome octavo, 300 pages. Cloth, $1.50 net.

The present book differs from other similar works in several features, all of which are introduced to render it more practical and generally useful. The general plan of contents follows the lines laid down in training-schools for nurses, but the book contains much useful matter not usually included in works of this character, such as Poison-emergencies, Ready Dose-list, Weights and Measures, etc., as well as a Glossary, defining all the terms in Materia Medica, and describing all the latest drugs and remedies, which have been generally neglected by other books of the kind.

ESSENTIALS OF ANATOMY AND MANUAL OF PRACTICAL DISSECTION, containing "Hints on Dissection." By CHARLES B. NANCREDE, M. D., Professor of Surgery and Clinical Surgery in the University of Michigan, Ann Arbor; Corresponding Member of the Royal Academy of Medicine, Rome, Italy; late Surgeon Jefferson Medical College, etc. Fourth and revised edition. Post 8vo, over 500 pages, with handsome full-page lithographic plates in colors, and over 200 illustrations. Price: Extra Cloth or Oilcloth for the dissection-room, $2.00 net.

Neither pains nor expense has been spared to make this work the most exhaustive yet concise Student's Manual of Anatomy and Dissection ever published, either in America or in Europe.

The colored plates are designed to aid the student in dissecting the muscles, arteries, veins, and nerves. The wood-cuts have all been specially drawn and engraved, and an Appendix added containing 60 illustrations representing the structure of the entire human skeleton, the whole being based on the eleventh edition of Gray's *Anatomy*.

A MANUAL OF PRACTICE OF MEDICINE. By A. A. STEVENS, A. M., M. D., Instructor in Physical Diagnosis in the University of Pennsylvania, and Professor of Pathology in the Woman's Medical College of Pennsylvania. Specially intended for students preparing for graduation and hospital examinations. Post 8vo, 519 pages. Numerous illustrations and selected formulæ. Price, bound in flexible leather, $2.00 net.

FIFTH EDITION, REVISED AND ENLARGED.

Contributions to the science of medicine have poured in so rapidly during the last quarter of a century that it is well-nigh impossible for the student, with the limited time at his disposal, to master elaborate treatises or to cull from them that knowledge which is absolutely essential. From an extended experience in teaching, the author has been enabled, by classification, to group allied symptoms, and by the judicious elimination of theories and redundant explanations to bring within a comparatively small compass a complete outline of the practice of medicine.

MANUAL OF MATERIA MEDICA AND THERAPEUTICS.

By A. A. STEVENS, A. M., M. D., Instructor of Physical Diagnosis in the University of Pennsylvania, and Professor of Pathology in the Woman's Medical College of Pennsylvania. 445 pages. Price, bound in flexible leather, $2.25.

SECOND EDITION, REVISED.

This wholly new volume, which is based on the last edition of the *Pharmacopœia*, comprehends the following sections: Physiological Action of Drugs; Drugs; Remedial Measures other than Drugs; Applied Therapeutics; Incompatibility in Prescriptions; Table of Doses; Index of Drugs; and Index of Diseases; the treatment being elucidated by more than two hundred formulæ.

"The author is to be congratulated upon having presented the medical student with as accurate a manual of therapeutics as it is possible to prepare."—*Therapeutic Gazette.*

"Far superior to most of its class; in fact, it is very good. Moreover, the book is reliable and accurate."—*New York Medical Journal.*

"The author has faithfully presented modern therapeutics in a comprehensive work, . . . and it will be found a reliable guide."—*University Medical Magazine.*

NOTES ON THE NEWER REMEDIES: their Therapeutic Applications and Modes of Administration.

By DAVID CERNA, M. D., PH. D., Demonstrator of and Lecturer on Experimental Therapeutics in the University of Pennsylvania. Post-octavo, 253 pages. Price, $1.25.

SECOND EDITION, RE-WRITTEN AND GREATLY ENLARGED.

The work takes up in alphabetical order all the newer remedies, giving their physical properties, solubility, therapeutic applications, administration, and chemical formula.

It thus forms a very valuable addition to the various works on therapeutics now in existence.

Chemists are so multiplying compounds, that, if each compound is to be thoroughly studied, investigations must be carried far enough to determine the practical importance of the new agents.

"Especially valuable because of its completeness, its accuracy, its systematic consideration of the properties and therapy of many remedies of which doctors generally know but little, expressed in a brief yet terse manner."—*Chicago Clinical Review.*

TEMPERATURE CHART.

Prepared by D. T. LAINÉ, M. D. Size 8×13½ inches. Price, per pad of 25 charts, 50 cents.

A conveniently arranged chart for recording Temperature, with columns for daily amounts of Urinary and Fecal Excretions, Food, Remarks, etc. On the back of each chart is given in full the method of Brand in the treatment of Typhoid Fever.

A TEXT-BOOK OF HISTOLOGY, DESCRIPTIVE AND PRACTICAL. For the Use of Students. By ARTHUR CLARKSON, M. B., C. M., Edin., formerly Demonstrator of Physiology in the Owen's College, Manchester; late Demonstrator of Physiology in the Yorkshire College, Leeds. Large 8vo, 554 pages, with 22 engravings in the text, and 174 beautifully colored original illustrations. Price, strongly bound in Cloth, $4.00 net.

The purpose of the writer in this work has been to furnish the student of Histology, in one volume, with both the descriptive and the practical part of the science. The first two chapters are devoted to the consideration of the general methods of Histology; subsequently, in each chapter, the structure of the tissue or organ is first systematically described, the student is then taken tutorially over the specimens illustrating it, and, finally, an appendix affords a short note of the methods of preparation.

"The work must be considered a valuable addition to the list of available text-books, and is to be highly recommended."—*New York Medical Journal.*

"One of the best works for students we have ever noticed. We predict that the book will attain a well-deserved popularity among our students."—*Chicago Medical Recorder.*

THE PATHOLOGY AND TREATMENT OF SEXUAL IMPOTENCE. By VICTOR G. VECKI, M. D. From the second German edition, revised and rewritten. Demi-octavo, about 300 pages. Cloth, $2.00 net.

The subject of impotence has but seldom been treated in this country in the truly scientific spirit that it deserves, and this volume will come to many as a revelation of the possibilities of therapeusis in this important field. Dr. Vecki's work has long been favorably known, and the German book has received the highest consideration. This edition is more than a mere translation, for, although based on the German edition, it has been entirely rewritten by the author in English.

"The work can be recommended as a scholarly treatise on its subject, and it can be read with advantage by many practitioners."—*Journal of the American Medical Association.*

THE TREATMENT OF PELVIC INFLAMMATIONS THROUGH THE VAGINA. By W. R. PRYOR, M. D., Professor of Gynecology in the New York Polyclinic. 12mo, 248 pages, handsomely illustrated. Cloth, $2.00 net.

In this book the author directs the attention of the general practitioner to a surgical treatment of the pelvic diseases of women. There exists the utmost confusion in the profession regarding the most successful methods of treating pelvic inflammations; and inasmuch as inflammatory lesions constitute the majority of all pelvic diseases, the subject is an important one. It has been the endeavor of the author to put down every little detail, no matter how insignificant, which might be of service.

DISEASES OF WOMEN. By HENRY J. GARRIGUES, A.M., M.D., Professor of Gynecology in the New York School of Clinical Medicine; Gynecologist to St. Mark's Hospital and to the German Dispensary, New York City. In one handsome octavo volume of 728 pages, illustrated by 335 engravings and colored plates. Prices: Cloth, $4.00 net; Sheep or Half-Morocco, $5.00 net.

A PRACTICAL work on gynecology for the use of students and practitioners, written in a terse and concise manner. The importance of a thorough knowledge of the anatomy of the female pelvic organs has been fully recognized by the author, and considerable space has been devoted to the subject. The chapters on Operations and on Treatment are thoroughly modern, and are based upon the large hospital and private practice of the author. The text is elucidated by a large number of illustrations and colored plates, many of them being original, and forming a complete atlas for studying *embryology* and the *anatomy* of the *female genitalia*, besides exemplifying, whenever needed, morbid conditions, instruments, apparatus, and operations.

Second Edition, Thoroughly Revised.

The first edition of this work met with a most appreciative reception by the medical press and profession both in this country and abroad, and was adopted as a text-book or recommended as a book of reference by nearly *one hundred* colleges in the United States and Canada. The author has availed himself of the opportunity afforded by this revision to embody the latest approved advances in the treatment employed in this important branch of Medicine. He has also more extensively expressed his own opinion on the comparative value of the different methods of treatment employed.

"One of the best text-books for students and practitioners which has been published in the English language; it is condensed, clear, and comprehensive. The profound learning and great clinical experience of the distinguished author find expression in this book in a most attractive and instructive form. Young practitioners, to whom experienced consultants may not be available, will find in this book invaluable counsel and help."

THAD. A. REAMY, M.D., LL.D.,
Professor of Clinical Gynecology, Medical College of Ohio; Gynecologist to the Good Samaritan and Cincinnati Hospitals.

A SYLLABUS OF GYNECOLOGY, arranged in conformity with "An American Text-Book of Gynecology." By J. W. LONG, M.D., Professor of Diseases of Women and Children, Medical College of Virginia, etc. Price, Cloth (interleaved), $1.00 net.

Based upon the teaching and methods laid down in the larger work, this will not only be useful as a supplementary volume, but to those who do not already possess the text-book it will also have an independent value as an aid to the practitioner in gynecological work, and to the student as a guide in the lecture-room, as the subject is presented in a manner at once systematic, clear, succinct, and practical.

THE AMERICAN POCKET MEDICAL DICTIONARY. Edited by W. A. NEWMAN DORLAND, M. D., Assistant Obstetrician to the Hospital of the University of Pennsylvania; Fellow of the American Academy of Medicine. Containing the pronunciation and definition of all the principal words used in medicine and the kindred sciences, with 64 extensive tables. Handsomely bound in flexible leather, limp, with gold edges and patent thumb index. Price, $1.00 net; with thumb index, $1.25 net.

SECOND EDITION, REVISED.

This is the ideal pocket lexicon. It is an absolutely new book, and not a revision of any old work. It is complete, defining all the terms of modern medicine and forming an unusually complete vocabulary. It gives the pronunciation of all the terms. It makes a special feature of the newer words neglected by other dictionaries. It contains a wealth of anatomical tables of special value to students. It forms a handy volume, indispensable to every medical man.

SAUNDERS' POCKET MEDICAL FORMULARY. By WILLIAM M. POWELL, M. D., Attending Physician to the Mercer House for Invalid Women at Atlantic City. Containing 1800 Formulæ, selected from several hundred of the best-known authorities. Forming a handsome and convenient pocket companion of nearly 300 printed pages, with blank leaves for Additions; with an Appendix containing Posological Table, Formulæ and Doses for Hypodermatic Medication, Poisons and their Antidotes, Diameters of the Pemale Pelvis and Fœtal Head, Obstetrical Table, Diet List for Various Diseases, Materials and Drugs used in Antiseptic Surgery, Treatment of Asphyxia from Drowning, Surgical Remembrancer, Tables of Incompatibles, Eruptive Fevers, Weights and Measures, etc. Handsomely bound in morocco, with side index, wallet, and flap. Price, $1.75 net.

FIFTH EDITION, THOROUGHLY REVISED.

"This little book, that can be conveniently carried in the pocket, contains an immense amount of material. It is very useful, and as the name of the author of each prescription is given, is unusually reliable."—*New York Medical Record.*

A COMPENDIUM OF INSANITY. By JOHN B. CHAPIN, M.D., LL.D., Physician-in-Chief, Pennsylvania Hospital for the Insane; late Physician-Superintendent of the Willard State Hospital, New York; Honorary Member of the Medico-Psychological Society of Great Britain, of the Society of Mental Medicine of Belgium. 12mo, 234 pages, illust. Cloth, $1.25 net.

The author has given, in a condensed and concise form, a compendium of Diseases of the Mind, for the convenient use and aid of physicians and students. It contains a clear, concise statement of the clinical aspects of the various abnormal mental conditions, with directions as to the most approved methods of managing and treating the insane.

"The practical parts of Dr. Chapin's book are what constitute its distinctive merit. We desire especially, however, to call attention to the fact that in the subject of the therapeutics of insanity the work is exceedingly valuable. The author has made a distinct addition to the literature of his specialty."—*Philadelphia Medical Journal.*

AN OPERATION BLANK, with Lists of Instruments, etc. required in Various Operations. Prepared by W. W. KEEN, M. D., LL.D., Professor of Principles of Surgery in the Jefferson Medical College, Philadelphia. Price per Pad, containing Blanks for fifty operations, 50 cents net.

SECOND EDITION, REVISED FORM.

A convenient blank, suitable for all operations, giving complete instructions regarding necessary preparation of patient, etc., with a full list of dressings and medicines to be employed.

On the back of each blank is a list of instruments used—viz. general instruments, etc., required for all operations; and special instruments for surgery of the brain and spine, mouth and throat, abdomen, rectum, male and female genito-urinary organs, the bones, etc.

The whole forming a neat pad, arranged for hanging on the wall of a surgeon's office or in the hospital operating-room.

"Will serve a useful purpose for the surgeon in reminding him of the details of preparation for the patient and the room as well as for the instruments, dressings, and antiseptics needed."—*New York Medical Record*

"Covers about all that can be needed in any operation."—*American Lancet.*

"The plan is a capital one."—*Boston Medical and Surgical Journal.*

LABORATORY EXERCISES IN BOTANY. By EDSON S. BASTIN, M. A., Professor of Materia Medica and Botany in the Philadelphia College of Pharmacy. Octavo volume of 536 pages, 87 full-page plates. Price, Cloth, $2.50.

This work is intended for the beginner and the advanced student, and it fully covers the structure of flowering plants, roots, ordinary stems, rhizomes, tubers, bulbs, leaves, flowers, fruits, and seeds. Particular attention is given to the gross and microscopical structure of plants, and to those used in medicine. Illustrations have freely been used to elucidate the text, and a complete index to facilitate reference has been added.

"There is no work like it in the pharmaceutical or botanical literature of this country, and we predict for it a wide circulation."—*American Journal of Pharmacy.*

DIET IN SICKNESS AND IN HEALTH. By MRS. ERNEST HART, formerly Student of the Faculty of Medicine of Paris and of the London School of Medicine for Women; with an INTRODUCTION by Sir Henry Thompson, F. R. C. S., M. D., London. 220 pages; illustrated. Price, Cloth, $1.50.

Useful to those who have to nurse, feed, and prescribe for the sick. In each case the accepted causation of the disease and the reasons for the special diet prescribed are briefly described. Medical men will find the dietaries and recipes practically useful, and likely to save them trouble in directing the dietetic treatment of patients.

A MANUAL OF PHYSIOLOGY, with Practical Exercises. For Students and Practitioners. By G. N. STEWART, M. A., M. D., D. Sc., lately Examiner in Physiology, University of Aberdeen, and of the New Museums, Cambridge University; Professor of Physiology in the Western Reserve University, Cleveland, Ohio. Handsome octavo volume of 848 pages, with 300 illustrations in the text, and 5 colored plates. Price, Cloth, $3.75 net.

THIRD EDITION, REVISED.

"It will make its way by sheer force of merit, and *amply deserves to do so. It is one of the very best English text-books on the subject.*"—*London Lancet.*

"Of the many text-books of physiology published, we do not know of one that so nearly comes up to the ideal as does Professor Stewart's volume."—*British Medical Journal.*

ESSENTIALS OF PHYSICAL DIAGNOSIS OF THE THORAX. By ARTHUR M. CORWIN, A. M., M. D., Demonstrator of Physical Diagnosis in the Rush Medical College, Chicago; Attending Physician to the Central Free Dispensary, Department of Rhinology, Laryngology, and Diseases of the Chest. 219 pages. Illustrated. Cloth, flexible covers. Price, $1.25 net.

THIRD EDITION, THOROUGHLY REVISED AND ENLARGED.

SYLLABUS OF OBSTETRICAL LECTURES in the Medical Department, University of Pennsylvania. By RICHARD C. NORRIS, A. M., M. D., Lecturer on Clinical and Operative Obstetrics, University of Pennsylvania. Third edition, thoroughly revised and enlarged. Crown 8vo. Price, Cloth, interleaved for notes, $2.00 net.

"This work is so far superior to others on the same subject that we take pleasure in calling attention briefly to its excellent features. It covers the subject thoroughly, and will prove invaluable both to the student and the practitioner. The author has introduced a number of valuable hints which would only occur to one who was himself an experienced teacher of obstetrics. The subject-matter is clear, forcible, and modern. We are especially pleased with the portion devoted to the practical duties of the accoucheur, care of the child, etc. The paragraphs on antiseptics are admirable; there is no doubtful tone in the directions given. No details are regarded as unimportant; no minor matters omitted. We venture to say that even the old practitioner will find useful hints in this direction which he cannot afford to despise."—*New York Medical Record.*

A SYLLABUS OF LECTURES ON THE PRACTICE OF SURGERY, arranged in conformity with "An American Text-Book of Surgery." By N. SENN, M. D., PH. D., Professor of Surgery in Rush Medical College, Chicago, and in the Chicago Polyclinic. Price, $2.00.

This work by so eminent an author, himself one of the contributors to "An American Text-Book of Surgery," will prove of exceptional value to the advanced student who has adopted that work as his text-book. It is not only the syllabus of an unrivalled course of surgical practice, but it is also an epitome of or supplement to the larger work.

"The author has evidently spared no pains in making his Syllabus thoroughly comprehensive, and has added new matter and alluded to the most recent authors and operations. Full references are also given to all requisite details of surgical anatomy and pathology."—*British Medical Journal*, London.

THE CARE OF THE BABY. By J. P. CROZER GRIFFITH, M. D., Clinical Professor of Diseases of Children, University of Pennsylvania; Physician to the Children's Hospital, Philadelphia, etc. 404 pages, with 67 illustrations in the text, and 5 plates. 12mo. Price, $1.50.

SECOND EDITION, REVISED.

A reliable guide not only for mothers, but also for medical students and practitioners whose opportunities for observing children have been limited.

"The whole book is characterized by rare good sense, and is evidently written by a master hand. It can be read with benefit not only by mothers, but by medical students and by any practitioners who have not had large opportunities for observing children."—*American Journal of Obstetrics.*

THE NURSE'S DICTIONARY of Medical Terms and Nursing Treatment, containing Definitions of the Principal Medical and Nursing Terms, Abbreviations, and Physiological Names, and Descriptions of the Instruments, Drugs, Diseases, Accidents, Treatments, Operations, Foods, Appliances, etc. encountered in the ward or the sick-room. By HONNOR MORTEN, author of "How to Become a Nurse," "Sketches of Hospital Life," etc. 16mo, 140 pages. Price, Cloth, $1.00.

This little volume is intended for use merely as a small reference-book which can be consulted at the bedside or in the ward. It gives sufficient explanation to the nurse to enable her to comprehend a case until she has leisure to look up larger and fuller works on the subject.

DIET LISTS AND SICK-ROOM DIETARY. By JEROME B. THOMAS, M. D., Visiting Physician to the Home for Friendless Women and Children and to the Newsboys' Home; Assistant Visiting Physician to the Kings County Hospital; Assistant Bacteriologist, Brooklyn Health Department. Price, Cloth, **$1.50** (Send for specimen List.)

One hundred and sixty detachable (perforated) diet lists for Albuminuria, Anæmia and Debility, Constipation, Diabetes, Diarrhœa, Dyspepsia, Fevers, Gout or Uric-Acid Diathesis, Obesity, and Tuberculosis. Also forty detachable sheets of Sick-Room Dietary, containing full instructions for preparation of easily-digested foods necessary for invalids. Each list is *numbered only*, the disease for which it is to be used in no case being mentioned, an index key being reserved for the physician's private use.

DIETS FOR INFANTS AND CHILDREN IN HEALTH AND IN DISEASE. By LOUIS STARR, M. D., Editor of "An American Text-Book of the Diseases of Children." 230 blanks (pocket-book size), perforated and neatly bound in flexible morocco. Price, $1.25 net.

The first series of blanks are prepared for the first seven months of infant life; each blank indicates the ingredients, but not the *quantities*, of the food, the latter directions being left for the physician. After the seventh month, modifications being less necessary, the diet lists are printed in full. *Formulæ* for the preparation of diluents and foods are appended.

HOW TO EXAMINE FOR LIFE INSURANCE. By JOHN M. KEATING, M. D., Fellow of the College of Physicians and Surgeons of Philadelphia; Vice-President of the American Pædiatric Society; Ex-President of the Association of Life Insurance Medical Directors. Royal 8vo, 211 pages, with two large half-tone illustrations, and a plate prepared by Dr. McClellan from special dissections; also, numerous cuts to elucidate the text. Third edition. Price, Cloth, $2.00 net.

"This is by far the most useful book which has yet appeared on insurance examination, a subject of growing interest and importance. Not the least valuable portion of the volume is Part II., which consists of instructions issued to their examining physicians by twenty-four representative companies of this country. As the proofs of these instructions were corrected by the directors of the companies, they form the latest instructions obtainable. If for these alone, the book should be at the right hand of every physician interested in this special branch of medical science."—*The Medical News*, Philadelphia.

NURSING: ITS PRINCIPLES AND PRACTICE. By ISABEL ADAMS HAMPTON, Graduate of the New York Training School for Nurses attached to Bellevue Hospital; Superintendent of Nurses and Principal of the Training School for Nurses, Johns Hopkins Hospital, Baltimore, Md.; late Superintendent of Nurses, Illinois Training School for Nurses, Chicago, Ill. In one very handsome 12mo volume of 512 pages, illustrated. Price, Cloth, $2.00 net.

SECOND EDITION, REVISED AND ENLARGED.

This original work on the important subject of nursing is at once comprehensive and systematic. It is written in a clear, accurate, and readable style, suitable alike to the student and the lay reader. Such a work has long been a desideratum with those entrusted with the management of hospitals and the instruction of nurses in training-schools. It is also of especial value to the graduated nurse who desires to acquire a practical working knowledge of the care of the sick and the hygiene of the sick-room.

OBSTETRIC ACCIDENTS, EMERGENCIES, AND OPERATIONS. By L. CH. BOISLINIERE, M. D., late Emeritus Professor of Obstetrics in the St. Louis Medical College. 381 pages, handsomely illustrated. Price, $2.00 net.

"For the use of the practitioner who, when away from home, has not the opportunity of consulting a library or of calling a friend in consultation. He then, being thrown upon his own resources, will find this book of benefit in guiding and assisting him in emergencies."

INFANT'S WEIGHT CHART. Designed by J. P. CROZER GRIFFITH, M. D., Clinical Professor of Diseases of Children in the University of Pennsylvania. 25 charts in each pad. Price per pad, 50 cents net.

A convenient blank for keeping a record of the child's weight during the first two years of life. Printed on each chart is a curve representing the average weight of a healthy infant, so that any deviation from the normal can readily be detected.



SAUNDERS' NEW SERIES OF MANUALS

for Students and Practitioners.

THAT there exists a need for thoroughly reliable hand-books on the leading branches of Medicine and Surgery is a fact amply demonstrated by the favor with which the SAUNDERS NEW SERIES OF MANUALS have been received by medical students and practitioners and by the Medical Press. These manuals are not merely condensations from present literature, but are ably written by well-known authors and practitioners, most of them being teachers in representative American colleges. Each volume is concisely and authoritatively written and exhaustive in detail, without being encumbered with the introduction of "cases," which so largely expand the ordinary text-book. These manuals will therefore form an admirable collection of advanced lectures, useful alike to the medical student and the practitioner: to the latter, too busy to search through page after page of elaborate treatises for what he wants to know, they will prove of inestimable value; to the former they will afford safe guides to the essential points of study.

The SAUNDERS NEW SERIES OF MANUALS are conceded to be superior to any similar books now on the market. No other manuals afford so much information in such a concise and available form. A liberal expenditure has enabled the publisher to render the mechanical portion of the work worthy of the high literary standard attained by these books.

Any of these Manuals will be mailed on receipt of price (see next page for List).

SAUNDERS' NEW SERIES OF MANUALS.

VOLUMES PUBLISHED.

PHYSIOLOGY. By JOSEPH HOWARD RAYMOND, A. M., M. D., Professor of Physiology and Hygiene and Lecturer on Gynecology in the Long Island College Hospital, etc. Price, $1.25 net.

SURGERY, General and Operative. By JOHN CHALMERS DACOSTA, M. D., Professor of Clinical Surgery, Jefferson Medical College, Philadelphia. Second edition, revised and greatly enlarged. Octavo, 911 pages, 386 illustrations. Cloth, $4.00 net; Half-Morocco, $5.00 net.

DOSE-BOOK AND MANUAL OF PRESCRIPTION-WRITING. By E. Q. THORNTON, M. D., Demonstrator of Therapeutics, Jefferson Medical College, Philadelphia. Price, $1.25 net.

MEDICAL JURISPRUDENCE. By HENRY C. CHAPMAN, M. D., Professor of Institutes of Medicine and Medical Jurisprudence in the Jefferson Medical College of Philadelphia, etc Price, $1.50 net.

SURGICAL ASEPSIS. By CARL BECK, M. D., Surgeon to St. Mark's Hospital and to the German Poliklinik; Instructor in Surgery, New York Post-Graduate Medical School, etc. Price, $1.25 net.

MANUAL OF ANATOMY. By IRVING S. HAYNES, M. D., Adjunct Professor of Anatomy and Demonstrator of Anatomy, Medical Department of the New York University, etc. Price, $2.50 net.

SYPHILIS AND THE VENEREAL DISEASES. By JAMES NEVINS HYDE, M. D., Professor of Skin and Venereal Diseases, and FRANK H. MONTGOMERY, M. D., Lecturer on Dermatology and Genito-urinary Diseases in Rush Medical College, Chicago. Price, $2.50 net.

PRACTICE OF MEDICINE. By GEORGE ROE LOCKWOOD, M. D., Professor of Practice in the Woman's Medical College of the New York Infirmary, etc. Price, $2.50 net.

OBSTETRICS. By W. A. NEWMAN DORLAND, M. D., Assistant Demonstrator of Obstetrics, University of Pennsylvania; Chief of Gynecological Dispensary, Pennsylvania Hospital. Price, $2.50 net.

DISEASES OF WOMEN. By J. BLAND SUTTON, F. R. C. S., Assistant Surgeon to the Middlesex Hospital, and Surgeon to the Chelsea Hospital for Women, London; and ARTHUR E. GILES, M. D., B. Sc. Lond., F. R. C. S. Edin., Assistant Surgeon to the Chelsea Hospital for Women, London. 436 pages, handsomely illustrated. Price, $2.50 net.

IN PREPARATION.

NERVOUS DISEASES. By CHARLES W. BURR, M. D., Clinical Professor of Nervous Diseases, Medico-Chirurgical College, Philadelphia, etc.

*** There will be published in the same series, at short intervals, carefully prepared works on various subjects, by prominent specialists.

SAUNDERS' QUESTION COMPENDS.

Arranged in Question and Answer Form.

THE LATEST, MOST COMPLETE, and BEST ILLUSTRATED SERIES OF COMPENDS EVER ISSUED.

Now the Standard Authorities in Medical Literature

WITH

Students and Practitioners in every City of the United States and Canada.

THE REASON WHY.

They are the advance guard of "Student's Helps"—that DO HELP; they are the leaders in their special line, *well and authoritatively written by able men, who, as teachers in the large colleges, know exactly what is wanted by a student preparing for his examinations.* The judgment exercised in the selection of authors is fully demonstrated by their professional elevation. Chosen from the ranks of Demonstrators, Quiz-masters, and Assistants, most of them have become Professors and Lecturers in their respective colleges.

Each book is of convenient size (5 × 7 inches), containing on an average 250 pages, profusely illustrated, and elegantly printed in clear, readable type, on fine paper.

The entire series, numbering twenty-four subjects, has been kept thoroughly revised and enlarged when necessary, many of them being in their fourth and fifth editions.

TO SUM UP.

Although there are numerous other Quizzes, Manuals, Aids, etc. in the market, none of them approach the "Blue Series of Question Compends;" and the claim is made for the following points of excellence:

1. Professional distinction and reputation of authors.
2. Conciseness, clearness, and soundness of treatment.
3. Size of type and quality of paper and binding.

*** **Any of these Compends will be mailed on receipt of price (see next page for List).**

SAUNDERS' QUESTION-COMPEND SERIES.

Price, Cloth, $1.00 per copy, except when otherwise noted.

1. **ESSENTIALS OF PHYSIOLOGY.** 4th edition. Illustrated. Revised and enlarged. By H. A. Hare, M. D. (Price, $1.00 net.)

2. **ESSENTIALS OF SURGERY.** 7th edition, with a chapter on Appendicitis. 90 illustrations. By Edward Martin, M. D. (Price, $1.00 net.)

3. **ESSENTIALS OF ANATOMY.** 6th edition, thoroughly revised. 151 illustrations. By Charles B. Nancrede, M. D. (Price, $1.00 net.)

4. **ESSENTIALS OF MEDICAL CHEMISTRY, ORGANIC AND INORGANIC.** 5th edition, revised, with an Appendix. By Lawrence Wolff, M. D. ($1.00 net.)

5. **ESSENTIALS OF OBSTETRICS.** 4th edition, revised and enlarged. 75 illustrations. By W. Easterly Ashton, M. D.

6. **ESSENTIALS OF PATHOLOGY AND MORBID ANATOMY.** 7th thousand. 46 illustrations. By C. E. Armand Semple, M. D.

7. **ESSENTIALS OF MATERIA MEDICA, THERAPEUTICS, AND PRESCRIPTION-WRITING.** 5th edition. By Henry Morris, M. D.

8, 9. **ESSENTIALS OF PRACTICE OF MEDICINE.** By Henry Morris, M. D. An Appendix on Urine Examination. Illustrated. By Lawrence Wolff, M. D. 3d edition, enlarged by some 300 Essential Formulæ, selected from eminent authorities, by Wm. M. Powell, M. D. (Double number, price $2.00.)

10. **ESSENTIALS OF GYNÆCOLOGY.** 4th edition, revised. With 62 illustrations. By Edwin B. Cragin, M. D.

11. **ESSENTIALS OF DISEASES OF THE SKIN.** 4th edition, revised and enlarged. 71 letter-press cuts and 15 half-tone illustrations. By Henry W. Stelwagon, M. D. (Price, $1.00 net.)

12. **ESSENTIALS OF MINOR SURGERY, BANDAGING, AND VENEREAL DISEASES.** 2d edition, revised and enlarged. 78 illustrations. By Edward Martin, M. D.

13. **ESSENTIALS OF LEGAL MEDICINE, TOXICOLOGY, AND HYGIENE.** 130 illustrations. By C. E. Armand Semple, M. D.

14. **ESSENTIALS OF DISEASES OF THE EYE, NOSE, AND THROAT.** 124 illustrations. 2d edition, revised. By Edward Jackson, M. D., and E. Baldwin Gleason, M. D.

15. **ESSENTIALS OF DISEASES OF CHILDREN.** 2d edition. By William M. Powell, M. D.

16. **ESSENTIALS OF EXAMINATION OF URINE.** Colored "Vogel Scale," and numerous illustrations. By Lawrence Wolff, M. D. (Price, 75 cents.)

17. **ESSENTIALS OF DIAGNOSIS.** 2d edition, thoroughly revised. 60 illustrations. By S. Solis-Cohen, M. D., and A. A. Eshner, M. D. (Price, $1.00 net.)

18. **ESSENTIALS OF PRACTICE OF PHARMACY.** 2d edition, revised. By L. E. Sayre.

20. **ESSENTIALS OF BACTERIOLOGY.** 3d edition. 82 illustrations. By M. V. Ball, M. D.

21. **ESSENTIALS OF NERVOUS DISEASES AND INSANITY.** 48 illustrations. 3d edition, revised. By John C. Shaw, M. D.

22. **ESSENTIALS OF MEDICAL PHYSICS.** 155 illustrations. 2d edition, revised. By Fred J. Brockway, M. D. (Price, $1.00 net.)

23. **ESSENTIALS OF MEDICAL ELECTRICITY.** 65 illustrations. By David D. Stewart, M. D., and Edward S. Lawrance, M. D.

24. **ESSENTIALS OF DISEASES OF THE EAR.** 114 illustrations. 2d edition, revised and enlarged. By E. Baldwin Gleason, M. D.

Some of the Books in Preparation for Publication during 1900.

AMERICAN Text-Book of Pathology.

Edited by LUDVIG HEKTOEN, M.D., Professor of Pathology, Rush Medical College, Chicago; and DAVID RIESMAN, M.D., Demonstrator of Pathological Histology, University of Pennsylvania.

AMERICAN Text-Book of Legal Medicine and Toxicology.

Edited by FREDERICK PETERSON, M.D., Chief of Clinic, Nervous Department, College of Physicians and Surgeons, New York City; and WALTER S. HAINES, M.D., Professor of Chemistry, Pharmacy, and Toxicology, Rush Medical College, Chicago.

BECK—Fractures.

By CARL BECK, M.D., Professor of Surgery in the N. Y. School of Clinical Medicine.

BÖHM, DAVIDOFF, and HUBER—A Text-Book of Human Histology.

Including Microscopic Technic. By DR. A. A. BOHM and DR. M. VON DAVIDOFF, of the Anatomical Institute of Munich, and G. C. HUBER, M.D., Junior Professor of Anatomy and Histology, University of Michigan, Ann Arbor.

EICHHORST—A Text-Book of the Practice of Medicine.

By DR. HERMAN EICHHORST, Professor of Special Pathology and Therapeutics and Director of the Medical Clinic, University of Zurich. Translated and edited by AUGUSTUS A. ESHNER, M.D., Professor of Clinical Medicine in the Philadelphia Polyclinic.

FRIEDRICH — Rhinology, Laryngology, and Otology in their Relations to General Medicine.

By DR. E. P. FRIEDRICH, of the University of Leipsig.

LEVY AND KLEMPERER — The Elements of Clinical Bacteriology.

By DR. ERNST LEVY, Professor in the University of Strassburg, and DR. FELIX KLEMPERER, Privat-Docent in the University of Strassburg. Translated and edited by AUGUSTUS A. ESHNER, M.D., Professor of Clinical Medicine in the Philadelphia Polyclinic. *Just Ready.* Cloth, $2 50 net.

McFARLAND—A Text-Book of Pathology.

By JOSEPH MCFARLAND, M.D., Professor of Pathology and Bacteriology, Medico-Chirurgical College, Philadelphia.

OGDEN — Clinical Examination of the Urine.

By J. BERGEN OGDEN, M.D., Assistant in Chemistry, Harvard Medical School.

PYLE—A Manual of Personal Hygiene.

Edited by WALTER L. PYLE, M.D., Assistant Surgeon to Wills' Eye Hospital, Philada.

SCUDDER—The Treatment of Fractures.

By CHARLES L. SCUDDER, M.D., Assistant in Clinical and Operative Surgery, Harvard University.

SENN—Practical Surgery.

By NICHOLAS SENN, M.D., PH.D., LL.D., Professor of the Practice of Surgery and of Clinical Surgery, Rush Medical College, Chicago. Octavo volume of about 800 pages, profusely illustrated.

The Pathology and Treatment of Tumors.

By NICHOLAS SENN, M.D., PH.D., LL.D., Professor of the Practice of Surgery and of Clinical Surgery, Rush Medical College, Chicago. A New and Thoroughly Revised Edition in preparation.

STENGEL AND WHITE — The Blood in its Clinical and Pathological Relations.

By ALFRED STENGEL, M.D., Professor of Clinical Medicine, University of Pennsylvania; and C. Y. WHITE, M.D., Instructor in Clinical Medicine, University of Pennsylvania.

STEVENS—The Physical Diagnosis of Diseases of the Chest.

By A. A. STEVENS, A.M., M.D., Lecturer on Terminology, and Instructor in Physical Diagnosis, University of Pennsylvania.

STONEY — Surgical Technique for Nurses.

By EMILY A. M. STONEY, late Superintendent of the Training Schools for Nurses, Carney Hospital, South Boston, Mass.

SAUNDERS'

MEDICAL HAND-ATLASES.

THE series of books included under this title are authorized translations into English of the world-famous

Lehmann Medicinische Handatlanten,

which for **scientific accuracy, pictorial beauty, compactness,** and **cheapness** surpass any similar volumes ever published.

Each volume contains from **50 to 100 colored plates,** besides numerous illustrations in the text. The colored plates have been executed by the most skilful German lithographers, in some cases more than twenty impressions being required to obtain the desired result. Each plate is accompanied by a full and appropriate **description,** and each book contains a condensed but adequate **outline of the subject** to which it is devoted.

One of the most valuable features of these atlases is that they offer a **ready and satisfactory substitute for clinical observation.** Such observation, of course, is available only to the residents in large medical centers; and even then the requisite variety is seen only after long years of routine hospital work. To those unable to attend important clinics these books will be absolutely indispensable, as presenting in a complete and convenient form the most accurate reproductions of clinical work, interpreted by the most competent of clinical teachers.

While appreciating the value of such colored plates, the profession has heretofore been practically debarred from purchasing similar works because of their extremely high price, made necessary by a limited sale and an enormous expense of production. Now, however, by reason of their projected **universal translation** and reproduction, affording international distribution, the publishers have been enabled to secure for these atlases the **best artistic and professional talent,** to produce them in the **most elegant style,** and yet to offer them at a **price heretofore unapproached in cheapness.** The great success of the undertaking is demonstrated by the fact that the volumes have already appeared in **thirteen different languages**—German, English, French, Italian, Russian, Spanish, Dutch, Japanese, Danish, Swedish, Roumanian, Bohemian, and Hungarian.

The same careful and competent **editorial supervision** has been secured in the English edition as in the originals. The translations have been edited by the **leading American specialists** in the different subjects. The volumes are of a uniform and convenient size (5 × 7½ inches), and are substantially bound in cloth.

(For List of Books, Prices, etc. see next page.)

Pamphlet containing specimens of the Colored Plates sent free on application.

VOLUMES NOW READY.

Atlas and Epitome of Internal Medicine and Clinical Diagnosis. By Dr. Chr. Jakob, of Erlangen. Edited by Augustus A. Eshner, M.D., Professor of Clinical Medicine in the Philadelphia Polyclinic. With 68 colored plates, 64 text-illustrations, and 259 pages of text. Cloth, $3.00 net.

Atlas of Legal Medicine. By Dr. E. von Hofmann, of Vienna. Edited by Frederick Peterson, M. D., Chief of Clinic, Nervous Department, College of Physicians and Surgeons, New York. With 120 colored figures on 56 plates and 193 half-tone illustrations. Cloth, $3.50 net.

Atlas and Epitome of Diseases of the Larynx. By Dr. L. Grunwald, of Munich. Edited by Charles P. Grayson, M. D., Physician-in-Charge, Throat and Nose Department, Hospital of the University of Pennsylvania. With 107 colored figures on 44 plates, 25 text-illustrations, and 103 pages of text. Cloth, $2.50 net.

Atlas and Epitome of Operative Surgery. By Dr. O. Zuckerkandl, of Vienna. Edited by J. Chalmers DaCosta, M. D., Clinical Professor of Surgery, Jefferson Medical College, Philadelphia. With 24 colored plates, 217 illustrations, and 395 pages of text. Cloth, $3.00 net.

Atlas and Epitome of Syphilis and the Venereal Diseases. By Prof. Dr. Franz Mracek, of Vienna. Edited by L. Bolton Bangs, M. D., Professor of Genito-Urinary Surgery, University and Bellevue Hospital Medical College, New York. With 71 colored plates, 66 text-illustrations, and 122 pages of text. Cloth, $3.50 net.

Atlas and Epitome of External Diseases of the Eye. By Dr. O. Haab, of Zurich. Edited by G. E. de Schweinitz, M. D., Professor of Ophthalmology, Jefferson Medical College, Philadelphia. With 76 colored illustrations on 40 plates and 228 pages of text. Cloth, $3.00 net.

Atlas and Epitome of Skin Diseases. By Prof. Dr. Franz Mracek, of Vienna. Edited by Henry W. Stelwagon, M. D., Clinical Professor of Dermatology, Jefferson Medical College, Philadelphia. With 63 colored plates, 39 illustrations, and 200 pages of text. Cloth, $3.50 net.

Atlas and Epitome of Special Pathological Histology. By Dr. H. Durck, of Munich. Edited by Ludvig Hektoen, M. D., Professor of Pathology, Rush Medical College, Chicago. Two volumes, with about 120 colored plates, numerous text-illustrations, and copious text.

Atlas and Epitome of Diseases Caused by Accidents. By Dr. Ed. Golebiewski, of Berlin. Translated and edited with additions by Pearce Bailey, M. D., Attending Physician to the Department of Corrections and to the Almshouse and Incurable Hospital, New York. With 40 colored plates, 143 text-illustrations, and 600 pages of text.

IN PREPARATION.

Atlas of General Pathological Histology.
Atlas of Orthopedic Surgery.
Atlas of General Surgery.
Atlas of Operative Gynecology.
Atlas of Psychiatry.
Atlas of Diseases of the Ear.

THE

AME[illegible]Y.

Th[illegible]vera[illegible]
ideal [illegible]uiren[illegible]
dicti[illegible]ary,[illegible]
ability and[illegible]
bring it.[illegible]
The boo[illegible]

the
ew
rial
can
of

RC71 C66 1900
Solis-Cohen
Essentials of Diagnosis



COLUMBIA UNIVERSITY LIBRARIES
0052113515