

Observations, anatomical, physiological and pathological, on the pulmonary system : with remarks on some of the diseases of the lungs, viz. on haemorrhage, wounds, asthma, catarrh, croup, and consumption : tending to establish a new pathology of the lungs, founded on the anatomy and physiology of the parts : some remarks are introduced on the broken-wind of horses, and to the whole is added an appendix, containing Observations on some of the articles of the materia medica, viz. on the Rosa Rubra, Flores Chamaemeli and Sarsaparilla, as also on the Cicuta, Stramonium, Hyosciamus and Aconitum / by William Davidson.

Contributors

Davidson, William, 1765-
Bartlet, P.
Bristol Royal Infirmary. Library
University of Bristol. Library

Publication/Creation

London : Printed by S. Low, Berwick Street; for T. Egerton, Charing Cross; J. Johnson, St. Paul's Church Yard; and W. Richardson, Royal Exchange, 1795.

Persistent URL

<https://wellcomecollection.org/works/nj42whhn>

Provider

Special Collections of the University of Bristol Library

License and attribution


This material has been provided by This material has been provided by University of Bristol Library. The original may be consulted at University of Bristol Library. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>







D 307. a

151 273 2036

Restr. Med. 18

Bartley

P. Bartlett

. P*Bartlet

RESERVATION
NATIONAL SYSTEM OF MATHEMATICAL

MONARCHY SYSTEM

THE NATIONAL SYSTEM OF MATHEMATICAL

P. Bartlett

OBSERVATIONS,
ANATOMICAL, PHYSIOLOGICAL, and PATHOLOGICAL,
ON THE

PULMONARY SYSTEM:

WITH
REMARKS ON SOME OF THE DISEASES OF THE LUNGS, VIZ.
ON
HÆMORRHAGE, WOUNDS, ASTHMA, CATARRH, CROUP,
AND
CONSUMPTION;

TENDING
TO ESTABLISH A NEW PATHOLOGY OF THE LUNGS,
FOUNDED ON THE
ANATOMY AND PHYSIOLOGY OF THE PARTS.

SOME REMARKS
Are introduced on the BROKEN-WIND of HORSES.

AND TO THE WHOLE IS ADDED AN

APPENDIX,

CONTAINING
OBSERVATIONS ON SOME OF THE ARTICLES OF THE
MATERIA MEDICA, VIZ.

ON THE
ROSA RUBRA, FLORES CHAMÆMELI AND SARSAPARILLA;

AS ALSO ON THE
CICUTA, STRAMONIUM, HYOSCIAMUS AND ACONITUM.

Deo, Optimo, Maximo, duce, in regiones pathologicas tendimus, Anatomia
et Physiologia monstrantibus iter. Passibus autem æquis Chymia fe-
quatur, ut, illis deficientibus, hæc quoque magnas suas opes suppeditet.

— Si quid novisti rectius istis
Candidus imperti; si non, his utere mecum.


HOR.

BY WILLIAM DAVIDSON.

LONDON:

PRINTED BY S. LOW, BERWICK STREET;
FOR T. EGERTON, CHARING CROSS; J. JOHNSON, ST. PAUL'S CHURCH
YARD; AND W. RICHARDSON, ROYAL EXCHANGE.

1795.



OBSERVATIONS

ANATOMICAL, PHYSIOLOGICAL, AND PATHOLOGICAL

ON THE

PULMONARY SYSTEM

WITH

REMARKS ON THE THEORY OF THE LUNGS

BY

MEMORIALS, WOUNDS, & OTHER CASES, REPORTED

PATRICK BARTLET, ESQ.

CONSUMPTION

AND

TO ILLUSTRATE A NEW THEORY OF THE LUNGS

AND

ANATOMY AND PHYSIOLOGY OF THE PARTS
FOLLOWING SHEETS

Are included in the BROOKLYN KIND OF HORSES
AND THE RESPECTIVE ANATOMY

APPENDIX

ON THE TESTIMONY OF ESTHETIC AND RESEARCH

BY

HON. RICHARD, HUGHES CHAMBERLAIN, & BARRISTER AT LAW

AND

CIRCUIT, ATTORNEY AT LAW, AND MEMBER OF THE
AND THE MOST Eminent

OF THE MEDICAL SOCIETY, IN THE CITY OF NEW YORK

WILLIAM DAVIDSON

Author of "The Anatomy of the Human Eye"

1825

BY WILLIAM DAVIDSON

LONDON

PRINTED BY J. JOHNSON, ST. PAUL'S CHURCH

AND W. CLAYTON, ST. MARK'S CHURCH

CONTENTS

TO

PATRICK BARTLET, ESQ.

Ec. Ec.

THE

MUTARE

FOLLOWING SHEETS

Page 68, line 18, for 'I' read 'we'

ARE RESPECTFULLY INSCRIBED,

The Author is sorry that in some other passages a similar inadvertence will be perceived, which, he hopes, the reader will excuse.

SMALL TESTIMONY OF ESTEEM AND REGARD,

BY

HIS MUCH OBLIGED,

AND MOST OBEIENT

HUMBLE SERVANT,

WILLIAM DAVIDSON.

QUEEN ANNE-STREET, EAST.

1795.

CONTENTS

PATRICK BARTLET, ESQ.

ERRATUM.

Page 66, line 18, for 'I,' read 'we.'

The Author is sorry that in some other passages a similar inadvertence will be perceived, which, he hopes, the reader will excuse.

CONTENTS.

	Page
INTRODUCTION, — — —	vii

CHAP. I.

<i>General Anatomy and Physiology of the Lungs, with some preliminary Observations,</i>	1
---	---

CHAP. II.

<i>General Observations, and practical Remarks, on Active Hæmorrhage from the Lungs: together with some Remarks on Wounds pe- netrating their Substance, —</i>	12
<i>Several Cases of Pulmonary Hæmorrhage, speedily and successfully cured by a limited Use of Liquids, — —</i>	29

CHAP. III.

<i>Observations, and practical Remarks on the Asthma, — —</i>	49
<i>A few Thoughts on the Broken-wind in Horses and other Animals, as being a disease ana- logous to the Asthma of the Human Subject,</i>	77

CHAP. IV.

<i>General Remarks on Pulmonary Consumption, with some Observations on Catarrh, and Croup; as also on the Seat, Origin, Formation, and Termination of Tubercle,</i>	84
<i>On Inflammation, Suppuration, and Ulceration of Tubercle,</i>	124
<i>Prevention and Cure of Tubercle,</i>	128

CHAP. V.

<i>Remarks on the Treatment of Ulceration of the Lungs, and Pulmonary Consumption,</i>	139
<i>Two Cases of Consumption, the one greatly relieved, and the other cured, by the limited Use of Liquids,</i>	173

APPENDIX.

<i>Observations on some of the Articles of the Materia Medica,</i>	183
--	-----

CHAP. I.

<i>On the Rosa Rubra, Flores Chamæmeli and Sarsaparilla,</i>	ibid.
--	-------

CHAP. II.

<i>On the Cicuta, Stramonium, Hyosciamus and Aconitum; with some Remarks on Sedatives in general,</i>	190
---	-----

INDEX,	221
--------	-----

INTRO-

INTRODUCTION.

AS diseases of the Lungs are, by far, the most common in this country, and the treatment of them involved in much obscurity and difficulty, any rational attempt to throw light upon the subject will, I doubt not, be examined with that candour and attention which the magnitude of the object and the nature of the undertaking require.

And, when I inform the reader of the extent and importance of the subject I am about to consider, that it comprehends, and influences the treatment of, the whole of

the diseases of some vital organs, and tends to point out a principle in the cure of them hitherto unnoticed; he will, I trust, be induced to peruse these observations with patience, and to forgive some errors which he will undoubtedly meet with. Therefore, in treating a business of so great moment, and attended with so much difficulty, instead of incurring the severity of criticism for presumptive boldness, I hope, I shall receive the approbation of the Public for having had sufficient courage to attempt so arduous an undertaking. For notwithstanding the task is arduous indeed, and although in this field of Pathology I have to contend with a most powerful enemy, I shall, nevertheless, attempt his overthrow. And, animated by the exclamation of Baglivi,

“ O quantum difficile est curare morbos pulmonum ! ”

I boldly

I boldly proceed ; and, under the auspices of Heaven, doubt not of some success.

The business of the following remarks, therefore, is to point out a *principle* hitherto neglected, but of the utmost importance, in the treatment of every morbid affection of the Lungs ; and from which arises a new pathology of all the pulmonary diseases, founded on the best of all possible bases, the anatomy and physiology of the parts. This principle, which, I conceive the reader will find established in the following remarks, is that of a particular and strict *limitation of liquids* during the treatment of every pulmonary disease ; a principle simple and obvious, involved in no obscurity, and easily applied.

It appears wonderful that practitioners have never thought of this principle, but
 have,

have, on the contrary, constantly treated patients, labouring under pulmonary affections, in the same way as if they had laboured under similar diseases of any of the other parts of the body. But it seems probable that this proceeded from their not attending to their peculiar structure, which is different from every other part of the body; for in all the other viscera, and in almost every other soft part of the body, there is some fleshy substance in their composition besides their vessels; but, in the lungs, there is no parenchymatous or fleshy substance, they being entirely composed of vessels of different kinds, of which the blood vessels form a very principal part. Respecting the common treatment of pulmonary diseases, every practitioner knows, that the patient is ordered to drink plentifully of diluting drinks; which appear grateful to the patient, because the dry and
irritable

irritable fauces receive a temporary relief from the passage of these soft liquids. But whoever considers the structure of the lungs, and remembers that they are entirely composed of vessels, and that their natural functions cannot be easily performed, even in health, if much additional liquid is taken into their blood vessels, will immediately see the impropriety of the practice, and be no more astonished at our want of success in the treatment of their various diseases.

To endeavour, therefore, to remove this error, and to establish the almost self-evident principle above-mentioned, constitutes the chief intention, as has been already observed, of publishing these remarks.

This principle of treating diseases of the lungs first occurred to me while attending a patient

a patient affected with hæmorrhage from them; but, since then, I have found a proper limitation of liquids of much benefit in all the other pulmonary diseases. Therefore, instead of ordering my patients to drink many quarts of diluents in the twenty-four hours, as is the common practice, I have only allowed them half a pint, a pint, or a pint and a half, of liquid, including tea and every other kind of fluid taken by the patient, during that period.

And my practice has been crowned with the most flattering success.

And case second of pulmonary hæmorrhage exhibits an example of actual consumption, and where the hæmorrhage was most probably a consequence of ulceration
of

of the lungs, where not only the hæmorrhage, but also all the other pneumonic affections were removed by that method of treatment.

In the cure of asthma I have also found a proper attention to this principle of great importance.

And in the following observations, I hope I have satisfactorily proved that the proximate cause of this morbid affection, when an original disease, does not always consist in constricted air vessels, as has been hitherto supposed, but more generally in over-distended blood vessels, occasioning difficulty of breathing from compression, and not from constriction, of the bronchia.

To

To the remarks on asthma some observations are added on the broken-wind of Horses, which disease in them I conceive to be the same as the asthma of the human subject.

After these remarks on asthma, &c. a few observations are offered on that most horrible of the pulmonary diseases, the Pulmonary Consumption.

In the treatment of consumptive patients, I have observed many advantages to arise from the moderate use of liquids; for even those, who were in the last stage of the disease, found their breathing become thereby easier, and their hectic symptoms also much diminished.

In pointing out the application of the above principle in the cure of the different
pulmonary

pulmonary diseases I have rather chosen, for the present, to deliver the doctrine as it naturally arises from practice and observation, than to confine myself by any systematic arrangement. But when a proper opportunity shall be allowed me I will, with pleasure, resume the subject, and arrange it in a more regular form; adding, at the same time, whatever new observations may occur from future practice.

To my observations and remarks on pulmonary diseases I have added a few observations on some of the articles of the materia medica, from a strong conviction, that, if properly considered, they will in some degree contribute to alleviate the miseries of mankind.

To conclude: I have only to add, that I rely, with confidence, upon that Public, in
 whose

whose service I am most assiduously engaged, for excusing many errors which must unavoidably occur under the circumstances in which I am placed.

OBSERVATIONS,

OBSERVATIONS,

Æc. Æc.

CHAP. I.

*General Anatomy and Physiology of the Lungs,
with some preliminary Observations.*

AS a particular description of the anatomy and physiology of the lungs would far exceed the limits of this publication, a very concise and general view only will be given; it being solely intended to call to the reader's mind some general circumstances which are more immediately connected with the present subject. The reader will be pleased to recollect that the lungs, which are the organs of respiration, are situated in the lateral parts of the thorax, and consti-

tute the chief bulk of its contents; that they nearly furround the heart, with which and their appendages, they form one large mass adapted in figure to, what is commonly called, the cavity of the thorax: that they consist of a congeries of blood vessels, absorbing vessels, nerves, and air vessels, joined loosely together by the cellular membrane, the common connecting medium of the body; and that the whole is enveloped by the pleura, a fine smooth membrane, which, being continued from the lungs over the internal surface of the ribs, intercostal muscles, and diaphragm, forms the internal lining of the chest. There is no fleshy substance in their composition besides these vessels; so that they may be justly considered as two bundles of vessels, right and left, forming the great pulmonary system; of which the blood vessels make a very principal part. These blood vessels, which, in structure, resemble those of the other parts of the body, are the pulmonary artery, arising from the right side of the heart, and branching through the lungs; and the pulmonary veins, arising from the extremities of the arteries and passing on to terminate in the left side of the heart. And by these pulmonary arteries and veins the blood is exposed

exposed to the influence of the air, and transmitted from the right to the left side of the heart. But, besides these great vessels of exposition and transmission, the lungs have another set of blood vessels, in common with every other part of the body, for their nourishment and support; we mean the bronchial arteries with their corresponding veins, whose minute ramifications constitute the *vasa vasorum* of all the other pulmonary vessels. Whether the last-mentioned vessels be the chief source of the internal secretion is not yet clearly determined. Now it will appear evident, that the pulmonary blood vessels, as possessing the same muscular structure as the other blood vessels of the body, are liable to the same diseases. The same increased tone and activity may certainly exist in them, as in the general system; as well as the same debility, and the same disposition to over-distention, from which that contractile power, necessary for a proper circulation, may be either diminished or destroyed. Nor is it improbable to suppose, that aneurism and varix may exist in the blood vessels of the lungs, as well as in those of the other parts of the body.

The system of air vessels consists of the larynx, the trachea arteria, and its branches (the
B 2
bronchia),

bronchia), which terminate in a particular cellular termination, called the air cells.

Here I beg leave to call the recollection of the reader to the muscular and cartilaginous structure, as well as to the irritable internal membrane, of this system.—He will recollect that this internal lining is a continuation of that mucous membrane which covers the fauces; and that, like it, it is also furnished with glands which secrete a slimy fluid, or mucus, for keeping it moist and defending it from the irritation of noxious vapours, or a too sharp atmosphere. For the variety of situation and circumstances, under which man might be accidentally placed, rendered it necessary that his constitution should thus adapt itself to them in the article of air, as well as in that of heat and cold. But besides this secretion of mucus for defence, there is also a considerable secretion or exhalation of watery vapour on every expiration.

The air cells, or the ultimate terminations of the bronchia, have been supposed by some anatomists to communicate with the common cellular membrane of the lungs: but they certainly do not, although in structure they bear a resemblance, inasmuch as the air cells communicate

with

with one another, in the same way as the cells of the cellular membrane.

If that supposed communication existed, dropy of the lungs, or a collection of water in their common cellular membrane, could not exist; but which I have several times seen, even when the lungs have been apparently found in every other respect.

And in the second volume of Medical Communications, p. 471, a case is related by the learned, the ingenious, and accurate Dr. James Carmichael Smyth, of effusion of blood and serum into the cellular connecting membrane of the lungs, which destroyed the patient; yet not a drop of it was expectorated, as it was poured out into the connecting cellular substance, and not into the air cells.—Dr. Smyth likewise mentions his having seen a case of the same kind after peripneumony; and that he had also seen one instance of pure emphysema of the lungs. Monsieur Vitet, of Lyons, in his dissections of the larger animals, as the horse, the ox, &c. could find no communication between the air cells and the connecting cellular membrane. His words are, “ L’existence des vesicules est donc aussi imaginaire que *le passage de l’air dans le tissu cellulaire*, & que sa sortie à travers les membranes

membranes qui convrent la surface externe des poumons.*” Whenever, therefore, this communication has appeared, we may venture to say that it was occasioned either by disease, from rupture of the air cells, or from some perforation of them, or extrication of air after death. The reader will observe we are only speaking of the human subject and quadrupeds; as we know that, in birds, the air pervades almost every part of the body, which answers a good purpose in their œconomy. But, even in them, it has been demonstrated that the aërial system is a system by itself, having no communication with the common cellular membrane †. The reader already knows that absorbing vessels and nerves are sent to the lungs, in common with every other part of the body.

To all the different branches of the system just mentioned, different offices are assigned; which, when regularly and easily performed, constitute its general health, and *e contra*. To the absorbent vessels is assigned the office of counteracting the natural exhalation into the different interstices, of absorbing extravasation, and,

* Vide Médecine Vétérinaire par M. Vitet, Docteur & Professeur en Médecine, à Lyon.

† Vide Philosophical Transactions, 1774.

perhaps,

perhaps, tubercle and induration ; and if they fail to do their duty, disease and consequent oppression will succeed. By its nerves is conveyed that influence, whatever it may be, which is necessary for producing the different actions of the whole. By means of the air vessels the air is constantly passing to and from the lungs, which, either by conveying something vivifying to, or carrying off something noxious from, the constitution (perhaps both) becomes so necessary for health, that we can live but a very short time without it ; and, in order that this particular function may be properly performed, it is necessary that the lungs must be in perpetual motion. By the large and numerous pulmonary arteries and veins, the whole blood of the body is exposed in the lungs, and conveyed from the right to the left side of the heart, as I shall now briefly relate. The blood, collecting together from all the different parts of the body, comes into the right side of the heart, from whence it is circulated by the pulmonary artery and its different ramifications through the lungs ; and the minute branches of this artery terminating in those of the pulmonary veins, the blood is thereby conveyed to the left side of the heart ; from whence

it is thrown into the aorta, or great artery; which carries it all over the body, for the various purposes of the animal œconomy. From this account of the circulation through the lungs, it will clearly appear that it must keep pace with the motion of the heart: so that, if the blood returns in great quantity to the heart, it will be roused to stronger action, and will throw the blood with force and celerity, and in increased quantity into the vessels of the lungs; but if the blood returns in moderate or diminished quantity to the heart, it will be propelled with slowness and regularity into the lungs, and its circulation through them will be gentle and uniform. In the former case the vessels will be distended beyond their healthy state, and rupture of any weak parts be likely to follow; and, when ruptured, may continue to pour out their contents, and an union of the orifices be thereby prevented; thus hæmoptoe, which becomes a very frequent cause of consumption, is produced; and it would appear probable that the distention of the vessels proves one of the chief obstacles to the cure. In the latter there will be no impediment to any of the natural pulmonary functions: so that, if in a state of disease, breath-

ing

ing will become less frequent and easier, ruptured vessels will unite, incipient tubercles may disappear, extravasations be absorbed, and ulcerations have a greater chance of a cure.

To conclude, I trust it will appear evident, from what hath been said, that when the absorbents of the lungs do their duty, when the air passes and repasses with ease, when the blood flows uniformly and easily through them, and when the nervous influence is properly conveyed, that then they are in health.

After this general view of their structure and healthful functions, we shall next beg leave to offer a few other general observations before we enter upon the treatment of their various diseases. It will doubtless be allowed that, in the different diseased states of the several parts of the body, a cure can only be brought about by the proper exercise of the powers naturally inherent in those parts, and that the business of the physician and surgeon is either to excite or restrain those actions, so that their exertions towards recovery may be precisely adapted to the state of the parts, or nature of the disease, under which they labour: and that, for this purpose, it is, in general, necessary that the part affected should be at rest, that those natural

tural

tural efforts may be allowed to proceed, without interruption, in the business of restoration. On this principle, the surgeon having properly placed the ends of a fractured bone, and put the limb in the most easy natural position, leaves the rest to nature; and no plaister, no bandage, can unite these bones, if the natural functions of the part, from inability or want of excitement, are not exerted on the occasion; for by them, and them alone, can the health of the part be restored, or a callus formed. These exertions the surgeon will either excite or restrain, according as they are either too remiss or too active in the performance of their respective functions. In the former case he will give the bark, wine, and other stimuli; and in the latter he will use bleeding, purging, and the other means of lessening action commonly employed. In the different morbid affections of the lungs, the physician will endeavour to adopt the same plan which the surgeon useth in the treatment of a broken limb, viz. he will endeavour to keep them as quiet, and as much at their ease, as possible, and so regulate their natural powers as to give the best chance of a cure: but here he labours under many disadvantages, as the natural offices of the parts require

require them to be in perpetual motion ; and, in cases of inflammation, rupture of the vessels, induration or other disease of any part of them, the variety of local applications, made use of to external parts, cannot be adopted. No fomentation, no poultice, no styptic, can be applied. Inhalation is the only mean we have of local treatment, but which, when properly regulated, may be of considerable service.

However, as no application can form the callus of a bone, so no particular medicine, we know, whether locally applied or internally administered, can, to a certainty, remove a tubercle, or heal an ulcer of the lungs.

The most rational plan, therefore, and which will, most probably, give the greatest chance of recovery, seems to be to retain them, as much as possible, at ease, that their natural powers may be allowed to exert themselves, and be so regulated as to accomplish a cure.

CHAP. II.

*General Observations, and practical Remarks, on
Active Hæmorrhage from the Lungs: together
with some Remarks on Wounds penetrating their
Substance.*

AS the whole of the doctrine to be delivered in these observations, is more particularly applicable to, and originated in, the treatment of bleeding from the lungs, and, as this disease proves a very common cause of pulmonary consumption, it may be most natural to notice it first.

Active hæmorrhage from the lungs may arise from a variety of causes; and may be naturally divided into two different states, viz.

- 1st, When the hæmorrhage alone constitutes the disease;
- 2d, When accompanied with more or less of other morbid affection.

This distinction, although worthy of consideration in forming a prognosis of the disease,

is of little use in the application of the principle we contend for, or in the administration of the other remedies commonly made us of.

The more common causes of the first state of hæmoptysis are, coughing, straining, or any violent exertion, by which considerable determination of the blood is made to the lungs, from which proceeds rupture and consequent hæmorrhage. In this case it is more apt to happen in a plethoric state of the system, and most frequently to persons of narrow chests, whose lungs are therefore compressed, and their vessels confined. Sudden rarefaction of the blood, while the vessels have been constricted from cold, is another very common cause of this disease.

But from whatever cause it arises, if the rupture is considerable, and a cure is not speedily accomplished by a proper union of the ruptured vessel, great bleeding will, most probably, ensue, and ulceration and consumption very generally follow. Whereas, if proper attention is paid to the principles here pointed out, the treatment becomes easy and effectual; and scarcely one patient in a hundred will become consumptive from this cause, providing his constitution is otherwise sound. What proportion of patients

tients become consumptive from ulceration of the lungs, the most common consequence of hæmorrhage from them, I do not exactly know; but when we find the learned and ingenious Dr. Beddoes, and other respectable authors, writing many pages, and in them mentioning it as the chief cause, we may readily conclude it is considerable.

Since the idea of moderate drinking, in diseases of the lungs, occurred to me, I have had many patients under my care in this first state of the disease, and have succeeded far beyond my most sanguine expectations by that mode of treatment.

The second state of pulmonary hæmorrhage is when it is attended with other disease, as tubercle, abscess, or induration. In the first state the ulceration commonly succeeds the hæmorrhage; but, in the second, it more frequently becomes the cause; for being nature's effort, by means of the absorbing vessels to get rid of tubercle, matter or other extraneous substance in the lungs, the blood vessels are thereby often eroded, and so pour forth their contents. And if a portion of the lungs becomes indurated, the circulation will be carried on with more difficulty; and, when any accidental fulness takes place, there will be most probably

probably rupture of the neighbouring vessels and consequent hæmorrhage.

The hæmorrhage in this second state, therefore, is in general a consequence of the other morbid affections. Here the cure becomes more difficult and uncertain; and the success will depend much upon the nature of the concomitant disease: yet under whatever circumstances the patient may be, he will soon be sensible of the great advantages resulting from the due regulation of his drink to be hereafter mentioned. And it may, perhaps, be satisfactory to know, that, from whatever cause the hæmorrhage proceeds, our principle is equally applicable, and will not only tend to fulfil the chief indication of healing the bleeding vessels, but also be a principal mean of removing all the other pulmonary affections, by leaving the lungs less embarrassed in the performance of their natural functions.

This second state of pulmonary hæmorrhage also comprehends all wounds penetrating the lungs, as by a ball, sword, or bayonet; in the cure of which the principle of a moderate use of liquids is of infinite importance.

We now go on to the treatment; which will include both the states of this disease, viz. ac-

tive hæmorrhage from the lungs, either with, or without, other pneumonic affection. Here I beg leave to call the attention of the reader to the plethoric state of the system which generally exists in active hæmorrhage, to the distended state of the blood vessels, and their consequent increased action; from which he will readily perceive, that this distention and increased action are the chief causes of the hæmorrhage, and the chief impediments to a cure.

And if he pays proper attention to the particular structure of the parts affected, to the size and number of their blood vessels, to the great quantity of blood circulated through them, and to their constant motion, he will soon be convinced of the danger of the disease; while the general circumstances just mentioned will intimate to him the most proper mode of relief.

It may be worthy of observation, that the doctrine, now delivered, is, in general, applicable to all cases of active hæmorrhages, or where they are attended with fever, although more particularly so in those of the lungs: and this fever, or increased action of the heart and arteries, the practitioner will attack by every mean in his power. For this purpose,
a strict

a strict adherence to every part of the antiphlogistic regimen, has been recommended: but from this I make an exception of that part of it, which commands plentiful drinking or dilution, for reasons to be hereafter given. The remedies for active hæmorrhage from the lungs may be ranked as follows, which however is rather the order of their exhibition than of their importance. They are bleeding, purging, limitation of liquids, saline nauseating medicines, blisters, ligatures on the extremities, &c. all of which we shall briefly consider in the order in which they are placed; and shall begin with

Bleeding. In all cases of active hæmorrhage, bleeding has been considered as a very sovereign remedy; and it certainly is so when used with moderation: but, like every other active medicine, it requires proper circumspection and regulation in its use.—For, as the blood is the vital fluid which warms, nourishes and supports every part of the body, and as its loss is very difficultly made up, so it ought never to be taken away excepting under the most urgent circumstances. Since the fatal doctrine of lentor was introduced, it may be questioned whether as many of the human race have not fallen sacrifices to the lancet as to the

sword: for it must be allowed that the one is as destructive as the other in improper hands.

I have been led into these observations from having seen practitioners, of great respectability and eminence, resting the chief stress of the cure of pulmonary hæmorrhage on repeated bleedings; in almost all of which cases the patients either became consumptive, or remained invalids for many months after the cure. That bleeding is a powerful and speedy mean of lessening distention, and of diminishing the increased action of the heart and arteries is well known; and it ought to be employed, without delay, on every pressing occasion: but when the cure can be accomplished without a repetition of it, the patient will recover more speedily than if his strength is exhausted by the lancet. When the most urgent symptoms, therefore, are removed, the other means of emptying the vessels and of lessening action, whose debilitating effects are not so permanent as those of bleeding, should be adopted; such as purgatives and saline nauseating medicines, never losing sight of a proper regulation of the liquids taken by the patient. By pursuing this method I have always found one bleeding sufficient: and after the removal of the disease the

patients

patients have generally been in better health and spirits than before their indisposition. The quantity of blood taken away in this one operation I have always regulated according to the urgency of the symptoms.

Purging comes next in order on our list of remedies: and, in active hæmorrhages of the lungs, when the vessels are over-filled, it becomes of infinite service; not only by diminishing the quantity of fluids in the blood vessels in general, but also by determining the blood from the parts affected, and thus giving the ruptured vessels time to unite. In order that purging may have its full effect, scarcely any liquid should be allowed during the operation. For neither your bleeding nor purging will have the desired effects of taking off the distention of the vessels, if these vessels are immediately filled again by a quantity of diluting liquors, which is commonly prescribed on these occasions. It may be necessary to specify what purgatives are most useful. All the saline purgatives, as coinciding with the antiphlogistic plan followed in these cases, are proper. Hoffman gave the preference to the *magnesia vitriolata*, which is the purgative I have, in general, used; and I think it possesses certain antispas-

modic and other powers, as well as its evacuant properties, which may assist in the cure of this disease.

The *limited use of liquids*, which is our *grand principle*, upon which the hinge of success in treating the disease now under consideration must turn, is placed next in order, although of the first importance. As the body, in its healthy state, is continually employing and discharging a particular portion of liquid, it is necessary that a certain quantity should be taken: but it commonly happens that from pleasure, or an evil habit, we drink much more than is required, and so over-distend the vessels, and embarrass nature in many of her salutary operations. In health, the quantity absolutely necessary is very inconsiderable; and, in sickness, we often drink too much. This has constantly been the case in pulmonary diseases; and particularly in hæmorrhages from the lungs, according to the common method of treating them. Practitioners had surely forgotten that the chief cause of the rupture and hæmorrhage, and the chief impediment to the cure, was the distention or too great fulness of the blood vessels; otherwise they would not have added to this fulness and distention by their plentiful dilution. When

no very urgent symptoms of hæmorrhagy are present, a pint of liquid, including tea and every other kind of fluid taken by the patient, is sufficient in twenty-four hours, and cannot safely be increased. But in the watchman's case, hereafter-mentioned, where apoplexy was present, accompanied with strong full pulse, as well as the hæmorrhage, notwithstanding I bled and purged him, I allowed him no drink for the first six hours, and half a pint only for the next twenty-four hours. He drank nothing during the operation of the physic: and the change produced by this regulation of liquid, even in a very short time, was astonishing. His vessels, of course, became emptier; fever and thirst were much abated; the apoplectic symptoms had disappeared; and, in short, all the morbid affections were more favourable.

From what hath been said I conceive it will be allowed that a proper regulation of the liquids taken by the patient is of the greatest importance in the treatment of pulmonary hæmorrhage, and experience enables me to assert, that, if early and proper attention is paid to this principle, the patient will, in general, be speedily restored; whereas, if neglected, and a contrary method pursued, even

all the other means of cure may prove ineffectual.

In a late conversation with a learned and intelligent Foreigner I was informed that almost all the French, who are taken with any considerable bleeding from the lungs, sink under the disease. On enquiring how much liquid they generally drank in twenty-four hours, he assured me the quantity was commonly very considerable; and that, when a purgative was given, the direction constantly was to drink *abondamment*. If so, the efficacy of our principle receives additional support; while the mortality resulting from a very opposite treatment is easily accounted for.

The *Saline Nauseating Medicines* next claim attention: and, in active hæmorrhage, are certainly of considerable use by lessening the increased action of the heart and vessels, and by determining the blood to the surface of the body. And, if we add to them, occasionally, some mild narcotic medicine, which may allay cough or irritation without increasing the action of the sanguiferous system, they will, without doubt, assist the general plan of cure.

For fulfilling the latter intention, opium has been recommended: but I always prefer
the

the fyrupus papaveris albi to the tinctura opii; the former possessing sedative powers sufficient for our purpose, while the latter cannot be so safely used on account of its stimulating properties. On most occasions, however, I have preferred the succus cicutæ spissatus to both.

The following formula I commonly use, varying the quantities of the ingredients according to circumstances.

Rx. Kali præparati scrupulum unum,
 Succu Limonum, q. s. ad saturationem,
 Magnesiæ Albæ scrupulum unum,
 Nitri grana decem,
 Vini Antimonialis guttas viginti,
 Lactis Amygdalarum unciam cum dimidia, &
 Syrupi Papaveris Albi drachmam unam (vel,
 ejus loco, Succu Cicutæ spissati grana quinque).

Misce, ut fiat haustus quatuor quaque horâ, vel pro re natâ, sumendus.

Blisters have generally been supposed useful in hæmorrhages of the lungs by taking off spasm, and assisting the determination to the surface of the body: I have, therefore, often employed them as auxiliaries to the general treatment.

Ligatures on the Extremities have been also used in this disease, particularly by the Ancients:

cients: and they may be of service by retarding the blood in its progress to the heart, and so allowing the circulation through the lungs to be carried on more gently; and, by thus lessening the distention of their blood vessels, give the orifices time to unite. For, as the whole blood brought to the right side of the heart must pass through the lungs in its passage to the left side, it surely follows that if a quantity of that blood, which is in the habit of returning to the right side of the heart, is arrested in the arms or legs, the quantity thrown into the lungs will be diminished in proportion; from which their vessels will become emptier, and thus the ligatures on the extremities will assist in accomplishing a cure. I should have conceived it unnecessary to observe that the nearer these ligatures are to the trunk of the body the better, had I not observed some authors advising to apply them to the wrist.

On Wounds of the Lungs. Before we conclude our remarks on hæmorrhagy, we shall beg leave to say a few words on recent wounds penetrating the lungs; where our principle of the limited use of liquids is particularly concerned, and will greatly assist the Surgeon in performing a cure. If the reader will consider
that,

that, in wounds of the lungs, there will be more or less of extravasation into the cellular membrane, as also into the aërial system, which will produce more or less of irritation, inflammation, cough, and fever, and which must be either absorbed or expectorated; and that the wounded vessels are also to be healed: he will readily believe, that the more quiet the lungs are kept, and the more empty their vessels are retained, the sooner and the easier will nature accomplish these salutary operations.—For this purpose, therefore, a due regulation of liquids is to be particularly enjoined, as nothing will contribute more towards a cure; although moderate bleeding and purging, together with the saline nauseating medicines, and blisters will also assist in removing the disease.

The following case came under the care of a very respectable medical friend; a man of correct judgement, and great veracity, who related it to me. In the year 1781, a soldier, in America, received a shot above the left breast, and the ball passed through the lungs, and also the scapula: from the uneasiness of moving, and some other accidental circumstances, he lived for four days without drinking any thing, but a very small quantity of barley water, or
weak

weak chicken broth. This foldier not only recovered speedily, but marched afterwards through almost the whole of North America, and is perhaps alive at this moment.

I recollect a case of a wound of the lungs, inflicted by a bayonet, which happened above nine years ago, and by a thrust from below upwards, and passed into the chest three or four inches; where the patient, finding pain when he moved or swallowed, would scarcely taste any thing for two days. He recovered in the course of a few weeks, and is now alive. Thus, from the uneasiness of moving or swallowing, have the patients been prevented from drinking; and thus our principle of keeping the lungs at ease hath been accidentally adopted, and a cure thereby accomplished. If wounds of the lungs were always treated in this way, I conceive all the patients would recover, excepting in cases of mortal wounds occasioned by the destruction of some great blood vessel.

From henceforward, therefore, I trust the proper limitation of liquids will be more particularly attended to: which may be the happy mean of saving the lives of many of the brave British sailors and foldiers, who are now gloriously employed in our defence.

Having

Having already demonstrated that, in active hæmorrhage, the vessels are in general full, distended, and acting too vigorously, it will be superfluous to observe that bark and elixir of vitriol, and every other tonic medicine, as keeping up that action, will be highly improper, and tend to increase the disease. For it has ever been, and ever will be found, that the more these medicines are given in active hæmorrhage, the longer will the bleeding continue, and the more obstinate will be the cure.

When the fever is gone, and the hæmorrhage becomes what is called passive, where the vessels are in general deficient in their contractile power; then, and then only, can astringent and tonic medicines be allowed.

The following is a pleasant, elegant, and effectual preparation of this kind, and has scarcely ever failed me in those cases of passive hæmorrhage where I have employed it.

Decoctum Rosarum *.

℞. Rosarum Rubrarum exsiccatarum drachmas tres,
 Aquæ puræ libram unam : coque ad libram dimidiam,
 & cola.

* Vide the Appendix.

℞. Decocti

℞. Decocti Rosarum supra-præscripti uncias duas,
 Tincturæ Opii guttas tres, vel q. f.
 Syrupi Croci drachmam unam,
 Elixir Vitrioli (Ph. Vet. Lond.) guttas octo, vel q. f.
 Mife, & fiat haustus quartâ, 6tâ, vel 8vâ, quâque horâ sumendus.

Here also, as well as in active hæmorrhage, a due attention is to be paid to the quantity of liquids taken by the patient, and more or less to be allowed according to the degree of the hæmorrhage, and state of the vascular system. Having, in the third and fourth volumes of Medical Facts and Observations, related several cases of active hæmorrhage successfully treated in the manner before-mentioned, and as the general principle I have adopted is new, and will be by them farther illustrated, it may not be improper to add them here: they therefore follow, as they were then published, together with the reflections and observations which then occurred.

Several

Several Cases of Pulmonary Hæmorrhage, speedily and successfully cured by a limited Use of Liquids.*

C A S E I.

On the 6th day of March, 1792, I was requested to visit Mr. S——, a man of a florid complexion, full habit of body, and about forty-five years of age. He had been much affected with head-ach, and hard dry cough, for three or four weeks. His pulse was now full, frequent, hard, and quick; and the veins upon his hands and arms were so much distended, that they appeared as if ready to burst. The cough was almost incessant, attended with considerable expectoration of florid frothy blood, which made its appearance this morning, after a severe fit of coughing, and his head-ach still continued.

The plan I followed was the following:

I took, from a large orifice in the arm, twelve ounces of blood, which from the long time it remained fluid after being taken from the arm, and the consequent appearance of (what is com-

* Vide Medical Facts, &c. Vol. III. p. 68.

monly

monly called) inflammation, both indicating the great action of the sanguiferous system, seemed to point out the necessity of employing the most vigorous antiphlogistic treatment. Much danger was also to be apprehended from the enlargement of the opening of the ruptured vessel.

Accordingly, I ordered him a saline draught, with antimonial wine, to be taken every three hours, adding to the night draught some syrup of white poppies, and an opening saline draught to be taken the morning following, and repeated every other morning.

This course of medicine, together with abstinence from animal food, and a strict adherence to a light cooling diet, was regularly pursued for three days; during which time the bleeding, although moderated, still continued, but the cough was much better.

9th. He was directed to continue the same diet, and to avoid much exercise; and the turgid state of the veins of his hands admonishing me that his vessels were still too full, it occurred to me to advise, instead of a second bleeding, that he should drink as sparingly as possible; from which I thought the vessels would become less full, and the ruptured vessel

fel have a greater chance of uniting than when constantly distended by drinking; and that, if I could avoid taking away more blood, my patient would recover from his indisposition much sooner than if I rested the chief stress of the cure upon this operation. He was, therefore, allowed a pint of liquid only, including tea and every other kind of drink, (all of which were given cold) in the twenty-four hours. When thirsty, I recommended it to him to suck an orange or lemon, instead of drinking. On former occasions of this kind, viz. in active hæmorrhages, I have prescribed (as is the common practice) cooling emulsions, milk whey and other diluents, in considerable quantities, with a view of relaxing the vascular system, and thereby lessening its increased action, not considering that the stimulus of distention kept up this action, and was, therefore, one of the chief things I had to guard against. But as there is now little to be dreaded from the Boerhaavian lentor, so there is no particular occasion for the great dilution commonly practised, and which seems to have been founded upon this doctrine. The medicines prescribed this day were similar to the former.

10th. I found him very cool, and without cough or expectoration of any kind. The pulse was softer, less frequent, and in every respect better. The appearance of the cutaneous veins also was so different, that I was convinced this great alteration for the better was chiefly to be attributed to his having avoided much drinking during the preceding day and night. The draughts kept the body regularly open once or twice a day, and induced a soft skin and comfortable sleep. They were, therefore, continued for three days, four every day; and three days more, two every day, still observing the same rule as to drinking. They always produced the same salutary effects. From this time the patient was perfectly well, and has remained so ever since.

In this case it would appear probable that no particular pneumonic affection existed, excepting the bleeding, which was most probably occasioned by a plethoric state of the constitution and particular determination to the lungs by the cough.

CASE

CASE II.

Soon after my attendance on the above patient, another case of hæmoptysis occurred, but which differed from the former in being attended with considerable pneumonic affection besides the hæmorrhage. The patient was a tall, thin man, about thirty years of age, of a pale complexion, narrow chest, and high shoulders, and had been affected with a severe cough for nearly four months previously to his application to me, accompanied with much yellow expectoration, and was supposed by his friends to be in a deep decline. He had no night sweats; but for the last three weeks had been affected with a continual pain of the right side; which, as far as I could discover, did not originate from any rheumatic affection of the external muscles, but from some internal disease of the thorax, and which I conceived to be a slow inflammation of the lungs, from which, and the violence of the cough, the hæmorrhage proceeded. He applied to me in the beginning of April, when he was coughing violently, and bringing up blood in mouthfuls. He had considerable fever, with a full hard pulse. I took from him ten ounces

of blood, and prescribed in every respect as in the foregoing case, enjoining to him great attention not to drink more than a pint of liquid in twenty-four hours. This, and every other rule directed, he regularly observed for about three weeks, when the bleeding had ceased for three or four days, and also the pain in the side. But returning imprudently to his former diet, and drinking the usual quantity as when in health, previously, as I suppose, to the obstruction or inflammation of the lungs being removed, his cough returned, with some little appearance of bloody expectoration, mixed with that kind of yellow mucus, which is commonly discharged by mucous secreting surfaces when inflamed. These symptoms, however, were entirely removed in the course of ten days, by a steady attention to the spare diet, and abstinence from liquids, formerly recommended, and the medicines before used.

Since then he has been, and now is, in the most perfect health, without cough, pain in the side, or any other thoracic or pneumonic affection. It occurs to me that this second attack, and the success of the subsequent treatment, point out the delicate situation of the lungs, and also the efficacy of this method of cure.

Having

Having related the above two cases with every necessary precision, I shall beg leave to offer some few observations on active hæmorrhage in general, and on that of the lungs in particular. In all active hæmorrhages a plethoric state of the system generally exists: all the blood vessels of the body are full, distended, and acting vigorously; and hence, very commonly, rupture and consequent hæmorrhage.

Therefore the chief proximate cause seems to be distention and consequent increased action of the vessels: Dr. Cullen, indeed, adds congestion of blood, which certainly may happen either from accidental determination of blood to a part, or some particular fault in the original conformation, or acquired relaxation, of the coats of the vessels of certain parts. But it is well known that hæmorrhages may arise from general distention, without any particular congestion; and, in this case, will happen wherever the vascular system is weakest or least supported. The proximate cause being clearly ascertained, the method of cure will appear obvious. Remove the preternatural distention of the vessels, and their action will soon diminish; then nature, with very little assistance, will do the rest. Although this is evidently the case, it appears sin-

gular that, hitherto, almost all practitioners have neglected the most effectual method of accomplishing this desirable purpose, viz. by a due abstinence from liquids. In Dr. Moffatt's translation of Aretæus, page 347, are the following words: "The drink ought to be very sparingly
 " exhibited, for moisture is disadvantageous in
 " a dry diet." But, although this was written when treating of hæmorrhage, the intentions of Aretæus were only that the astringency (upon which he seemed to place his chief hope) of his diet might not be weakened by drinking.

The idea of moderate drinking is adopted by Dr. Rowley, in his treatise on "Female nervous diseases," published in 1788. When treating of the "Immoderate flow of the
 " menses," page 32, he observes, "as hæmor-
 " rhages seldom happen, unless there be a suffi-
 " cient quantity of blood in the body to rup-
 " ture the vessels, one principal part of the cure
 " consists in not only obtaining, but preserving
 " a diminished quantity of blood, by a great
 " *abstinence* from liquids; for by this means, the
 " very sources of supply are cut off. If little
 " be drank, the blood vessels which are, or have
 " been, distended beyond their proper dimen-
 " sions, will gradually contract themselves to
 " their

“ their original size, acquire strength daily, and
 “ not having so large a column of blood to cir-
 “ culate, they will resist the morbid disposition
 “ of nature to evacuate so violently the cata-
 “ menia.” But the late celebrated Dr. Cullen,
 when treating on hæmoptysis, particularly re-
 commends, that “ every part of the antiphlo-
 “ gistic regimen be strictly enjoined *,” which
 includes “ taking in large quantities of mild
 antiseptic liquors † :” and says, that the phlo-
 gistic diathesis is to be taken off by bleeding,
 more or less, according to circumstances. If,
 however, the stimulus of distention is kept up
 by filling the vessels with liquids, the good ef-
 fects of the bleeding are counteracted, and a
 frequent repetition rendered necessary—Where-
 as, if abstinence from liquids be particularly at-
 tended to, one bleeding will have more effect,
 than three or four, if accompanied with that part
 of the antiphlogistic regimen, and the loss of
 blood be thereby prevented; which, considering
 its importance in the constitution, and the diffi-
 culty with which its loss is made up, should be
 at all times avoided when possible.

* First Lines of the Practice of Physic, Vol. II. p. 353,
4th Edition.

† Ibid. Vol. I. p. 132.

Of all cafes of hæmorrhagy, that from the lungs is the moft dangerous in its nature, and moft difficult of cure. This will appear evident if we recollect their particular ftructure, their large and numerous veffels, their constant motion, &c.

As to their ftructure, anatomy demonstrates that they are compofed of a congeries of blood veffels, abforbents, and nerves, together with the air cells; and that all thefe are only connected by the cellular membrane, the common connecting medium of the body: for I do not mention their pleuritic covering, as I am only fpeaking of their fubftance. The blood veffels, with which alone our prefent fubject is connected, are very large, and in greater number than in any other part of the body of the fame fize. This was abfolutely neceffary to circulate the very large quantity of blood generally fent to them. Haller obferves*, that the quantity of blood which enters into the lungs is equal to, or even perhaps greater than, that which is fent in the fame time throughout the reft of the body. And, as the chief bufinefs of the lungs is for refpiration, by which

* Prim. Lin. Phyfiol. § 246:

they are kept constantly in action, so it will appear evident why hæmorrhages here are more dangerous, as well as more obstinate to cure, than in any other part, as their constant motion counteracts and prevents the union of the ruptured vessel.

*A Case of Pulmonary Hæmorrhage, with Remarks on Tubercle, Induration, Consumption, &c.**

C A S E III.

Having, in the third volume of *Medical Facts and Observations*, related two cases of pulmonary hæmorrhage, and from them endeavoured to demonstrate that the proximate cause of active hæmorrhage often consists in distention and consequent increased action of the blood vessels, and to point out that abstinence from liquids is a principal mean of removing this distention; I shall now beg leave to add another, which lately occurred, in farther confirmation of that doctrine, and of the advantages resulting from such a method of treatment. The patient was a robust man, of a sanguineous temperament, and about sixty-

* Vide *Medical Facts, &c.* Vol. IV. p. 129.

four years of age. He had been affected with a severe cough for near four months before the present attack; and during the last seven or eight weeks had been spitting blood, mixed with a yellow expectoration; but without any pain in or about the chest.

Being sent for on the fourth day of October, 1792, I found him in an insensible state, as if from oppression of the brain, with occasional strong contractions or convulsions of the right arm. His pulse was strong, frequent and full; his tongue was furred, and his breathing laborious. He had been just brought into the house from a Stone-mason's yard, where he was employed in sawing.

The persons about him informed me, that, while at work, he was seized with a fit of coughing, and brought up about three or four pints of blood; that he soon after became insensible, and was immediately brought home. Considering the great hæmorrhage which had taken place, and the apoplectic symptoms now present, I conceived my patient to be in considerable danger, and that the most active method of relieving him should be adopted: accordingly sixteen ounces of blood were instantly taken from a large orifice in the arm. The blood,
when

when coagulated, was covered with the buff coat, as it is called. A few minutes after the operation he became sensible, and complained of great pain in the anterior part of the chest; which, he said, he had first perceived that morning. About half an hour after the bleeding, he took a purging draught, chiefly composed of magnesia vitriolata. A large blister was also applied to the breast. He was particularly directed to refrain from drink during the operation of the purgative medicine; and, if thirsty, only to moisten his mouth and throat with a little barley water.

In the evening (six hours after my former visit) I found him sensible, with less fever, his cough quiet, his breast easier, and he had not brought up much blood. His medicine had purged him several times.

A saline draught containing twenty drops of antimonial wine was now directed to be taken every six hours; he was strictly enjoined to drink about half a pint only of liquid during the first twenty-four hours; and in every other respect to adhere rigidly to the antiphlogistic regimen.

October 5th, he had rested pretty well, and expectorated about an ounce or two only of
blood,

blood, which was chiefly in coagula: his breast was easier, but still a little tight; his pulse was much improved, and his skin was cool and moist. He had little thirst, and his tongue was less furred. The use of the saline draught was continued, and the opening draught was directed to be repeated in the morning. Being so much better he was now allowed a pint of liquid (including tea, &c.) in the twenty-four hours, and the same quantity only was permitted every day during the whole of his illness.

October 6th, he was still much better: he had rested well, had less cough, less fever, little bloody expectoration, and his pulse was nearly natural: his chest was much easier.

From this time to the 12th he continued gradually to recover. He had no expectoration of blood after the 8th, but the saline draught, and likewise the purgative medicine, were occasionally repeated, and he persevered in the limited use of liquids till the 12th, when I thought it unnecessary to visit him any longer. His pulse was then sixty-eight in a minute, and he was apparently in good health, only a little weak.

I afterwards learned that, contrary to my directions, he went upon duty, as patrol,

on

on the Monday following, the 15th day of October.

Considering, therefore, the nature of this office, the season of the year, the age of the patient, and the short time since his recovery, it cannot seem surprising that the disease was reproduced; accordingly, on the 25th, he was again seized with fever, difficult breathing, cough, and hæmorrhage. He continued, notwithstanding this return of the complaint, to attend his duty regularly until Sunday the 28th, when he was again taken with considerable bleeding, while on the patrol, and instantly expired.

On Tuesday the 30th, having an opportunity of inspecting the body, the following appearances presented themselves: the thorax and abdomen being laid open, we observed on the anterior surface of the right lung an incipient inflammation, which, however, could not account for the patient's death, for, on farther examination, it seemed evidently to be occasioned by the hæmorrhage. There were also some adhesions, apparently rather vestiges of former than of any recent inflammation. There were no tubercles. A small portion of the aorta was ossified. All the abdominal viscera

were

were found. In the stomach there was some coagulated blood, which had been swallowed; but there was not the smallest erosion of its coats.

In the two former cases of hæmoptysis, I have noticed the great difficulty of curing a ruptured vessel in the lungs, on account of their constant motion, and the great quantity of blood circulated through them; but that this difficulty might be, in general, overcome by a steady adherence to the plan of cure there recommended, viz. moderate bleeding and purging, but particularly a due abstinence from liquids. The success attending the treatment of the present case must evidently establish the superiority of that method of cure over every other hitherto recommended. Here a blood vessel, of considerable magnitude, was ruptured in a part of the body which, from its natural office, must be in perpetual motion, and where no local application could be made; yet this rupture was healed in almost as short a time as the most experienced Surgeon can heal an external accident of the same nature, even with the assistance of compresses and bandages. For example, I have seen a rupture of some superficial vessels require these applications for many
 days.

days. It may be said that the bleeding, which has been more or less plentifully used (I mean as to quantity, for it was never used more than once in each case) according to the urgency of the symptoms, was the chief mean of cure. But a practitioner, who has seen a patient blooded twelve or thirteen times for an hæmorrhage from the lungs, and still sink under the disease, will not readily subscribe to this opinion.

It may be necessary to observe that the patient to whom I allude was allowed to take, and actually did drink, several quarts of diluents in the twenty-four hours. But supposing he had recovered, after such loss of blood he must have remained infirm for many months: whereas this patient, who was so soon relieved by abstinence from liquids, had he been in easy circumstances, and could he have kept from labour and improper exposure to the night air, for another week or two, might have obtained a perfect and permanent cure, without any particular diminution of bodily strength.

Respecting the other medicines, they were doubtless of service, and conspired to effect a cure, which, had the usual quantity of diluents been used, I am convinced, would, notwithstanding,

standing, have been much more tedious. For in vain do practitioners attempt to lessen distention by emptying the vessels, either by purging or bleeding, if they are immediately filled again by plentiful drinking. The spare use of liquids, therefore, may justly be considered as one of the greatest improvements in the modern treatment of hæmorrhage: and particularly in hæmorrhages from the lungs. And why should not the idea be carried farther? Indeed, from some cases I have lately attended, I think I may venture to assert, that, in all diseases of the lungs, moderate drinking will be of service. For seeing they are a congeries of vessels, if these vessels are overfilled, or kept in a continued state of distention, they may so press upon one another that their healthy actions shall be either prevented or greatly impeded, particularly the actions of the absorbent system: whereas, if they are but moderately filled, the different systems of vessels are left more at liberty to exercise their respective functions, either in the business of health, or in the removal of disease. When tubercles are formed in the lungs, why should they not be absorbed? We know that the most solid tumors in other parts of the body frequently disappear; and that

that even bone itself is capable of being absorbed, as is clearly demonstrated by the different changes which take place in it as well in health as in disease. And in the lungs there are many absorbent vessels, which, if their actions were not lessened or prevented, might soon remove the most confirmed induration of their substance. As emetics are powerful promoters of absorption, is it not on this principle that many patients, seemingly labouring under tubercles of the lungs, have been cured by vomits, particularly of the stronger kind? I hope the time is not far distant when practitioners, being better acquainted with the laws and functions of this important system, shall be enabled to direct its actions with more certainty, either in removing a tubercle or the most scirrhous tumor. But when this happy period arrives it can only be carried into effect by a proper regulation of the quantity of liquids; and, in general, a diminution of the usual prescribed quantities. Perhaps the advantages arising to consumptive patients from a warm climate and the use of flannel, are principally from their doing the same thing as abstinence from liquids, viz. determining the tide of circulation to the surface of the body, and thus leaving the vessels

of

of the lungs more empty, and, therefore, more ready to recover themselves when under the influence of disease.

From the above cases I hope the reader is convinced that the moderate use of liquids is not only of infinite importance in the treatment of hæmorrhages in general, but is also more particularly so in that of active hæmorrhage from the lungs.

After the successful treatment of case second, where considerable pneumonic affection existed, besides the hæmorrhage, and which was removed, together with it, by our plan of cure, I was induced to view the spare use of liquids as a principle of greater importance than I had at first imagined: and, having recollected that the lungs were chiefly composed of vessels, among which inflammation, tubercle, induration, and all the other morbid affections must arise, I conceived that the less these vessels were distended, the less would the parts affected be embarrassed, and the more readily would a cure be accomplished. It therefore appeared applicable, not only in the treatment of hæmorrhage, but also in that of all the other diseases of the lungs. Accordingly many successful trials have confirmed me in the opinion, that,

that, in the treatment of every pulmonary complaint, a proper limitation of liquids will be productive of great advantages to the patient. For, in recent cases, a cure may be, thereby, more speedily accomplished; and, in even the most deplorable of these melancholy affections, the sick will derive much ease and comfort from this regulation of their drink, as the following cases, and those already related, will sufficiently demonstrate.

In attempting an explanation of this doctrine and its application in the other morbid affections of the lungs, we shall begin with Asthma previously to our remarks on Consumption, as the former disease is often a prelude to the latter.

C H A P. III.

Observations, and practical Remarks on the Asthma.

WE shall now proceed, in the same cursory way in which we have hitherto prosecuted our remarks, to point out the application of our principle in the cure of Asthma. Since the

E idea

idea of employing this principle in the treatment of this disease occurred to me, several opportunities of trying the efficacy of a moderate use of liquids in this most distressing of the pulmonary affections have presented themselves; and they have, in general, been crowned with the same success which attended my trials in active hæmorrhage. This indeed might have been naturally expected, as the same turgescence and distention of vessels are generally present in both, as will appear from the following quotations from Sir John Floyer and Dr. Cullen, which I have extracted from their works since the success of my own practice had determined me to publish the result for the public use.

But, before I mention the opinions of these respectable physicians, it may not be improper to relate one or two of the cases which I attended and treated successfully, and where the principle of moderate drinking was particularly attended to; introducing, at the same time, a few remarks, as they naturally arise from actual practice. These cases seem to me to demonstrate that the pathology of asthma, when considered as an original disease, has been in general erroneous; and that, therefore, the treatment

ment recommended has not been accompanied with the wished-for success.

C A S E I.

The patient, whose case I am now to relate, was a lady of a middling stature, of a pale complexion, of a full and relaxed habit of body, and about fifty-three years of age. She had been subject to repeated attacks of asthma for eight years; although, in every other respect, she had enjoyed very tolerable health. Her fits, which were always occasioned by any great hurry or fatigue, or severe cold, came on very irregularly: but, when they did, they generally continued for several months. In endeavouring to trace the cause, she informed me, that, for five years previously to a most violent attack, from which the eight years above-mentioned are dated, she felt a gradual difficulty of breathing coming on, but which only particularly affected her on going up stairs, or on being hurried; but never in any great or permanent degree. At the beginning of the eight years, having been a considerable distance from her house, and there threatened by a violent storm, she was obliged to run home as

fast as possible; upon which such great difficulty of breathing immediately followed, that her life was supposed to be in the most imminent danger. Two physicians being called to her assistance, the one advised immediate bleeding, and the other to try some medicine first. The latter plan was adopted, and the medicine (the composition of which she knows not) vomited and purged her severely; but relieved her so much, that there was no occasion for the bleeding. After being indisposed for many weeks, she recovered; but has, ever since, been subject to repeated attacks of the disease; particularly from the beginning of autumn to the end of spring, during which time she has very little interval of ease. It may be necessary to observe, that menstruation still continues, and has never been particularly interrupted, even when her asthma has been the most violent.

After having been indisposed about a week she sent for me, on the 25th of December, 1792, when I found her Asthma very severe; so much so that she could scarcely speak to me. She had also much cough, and some expectoration, but no pain about the chest. Her tongue was white and dry; and her pulse was frequent;

frequent, but neither full nor strong. I prescribed as follows, viz.

Capiat haustum emeticum ex antimonio tartarizato & vino ipecacuanhæ statim : et pilulam anodynam diaphoreticam, horâ fomni.

Primo mane sumat haustum catharticum e tinctura jalapii, &c. non bibendo inter operationem.

December 26th. Her emetic has answered very well, and she has had a comfortable night. Her physic has operated powerfully, and she breathes easier, and feels much relieved in every respect. I now allowed her a pint of liquid only for the next twenty-four hours, and sent her a mixture, composed of lac ammoniacum, oxymel simplex, &c. to be taken every three or four hours.

27th. She has slept tolerably well, and is getting better. The same rule respecting liquids to be continued; only, during the operation of the following pills, half a pint of weak tea may be taken.

Capiat haustum ex æthere vitriolico, &c. horâ fomni; & mane, sumat pilulas duas catharticas e refina jalapii, sapone & calce antimonii illotâ.

28th. She has had a very good night: her medicine has operated seven or eight times;

and she feels much better. I prescribed the same æther draught to be taken every night at bed-time, and advised the same limitation of liquids.

30th. She is exceedingly well; her pulse is nearly natural, and she has scarcely any difficulty of breathing. Her expectoration is now almost gone, having gradually diminished as her breathing became better. I therefore recommended her to take the æther draught for two nights, and to take another dose of her pills; and also to observe great moderation in drinking for several days.

On the 2d of January, 1793, I found her perfectly well. She had another severe attack about the end of January, and got well in six days by a similar treatment.

In June following she was seized again; and, after being ill for several days, she sent for me on the 14th, when her asthma was extremely severe. By a like treatment she was well on the 19th of the same month.

I saw her towards the end of February, 1794, and she was then well; and told me, with seeming astonishment, that since June, 1793, her asthma had not returned: which was the only autumn and winter she had missed it for the last

last eight years. She appeared lively and was not so corpulent as she had been when I attended her. It may be proper to observe, that in this patient the asthma was not hereditary.

Now the history of the commencement of this disease, in the case just now related, seems to me to throw considerable light on the nature of the asthma, as an original disease, and to point out the proper mode of relief. For I conceive that, during the first five years, when the difficulty of breathing was gradually coming on, that the blood vessels of the lungs were then, as gradually, losing their contractile power; so that the blood was not very regularly propelled through them. In this case, congestion of blood must take place in the lungs, whenever, by hurry or any extraordinary exercise, the blood was sent into them more quickly, or in greater quantity, than usual; and consequent difficulty of breathing must ensue. And, at the beginning of the eight years, when confirmed asthma took place, it seems probable that, from the great exertion used in running home, the vessels were so much distended as to lose, in great measure, their contractile power; from which such con-

E 4 gestion

gestion of blood in the lungs was produced, as might have proved fatal, had not timely assistance been procured.

CASE II.

The subject of this case, Mrs. Rose's cook, is a full young woman, of a relaxed habit of body, and about thirty-one years of age. She has been of this full habit of body from her infancy: and her father, after having been tormented with the asthma for many years, died of the disease in the beginning of February, 1794, soon after my attendance on his daughter. Her father's fits generally continued for many months at a time, without receiving any relief from a variety of medicines administered by his medical attendant.

January the 20th, 1794. This young woman has, in general, enjoyed a good state of health; and has never before had any asthmatic affection, although she has been extremely so for the last eight days; and is now scarcely able to breathe, notwithstanding she has taken some oily emulsions and nitrous medicines prescribed her by an apothecary in the neighbourhood. She is hot and thirsty, and her pulse is frequent

quent, and rather strong; and her tongue white and furred. Her face is full and nearly livid. She has also a troublesome cough, but which is not attended with any particular expectoration. Last night, she took, by the advice of her mistress, a table-spoonful of oleum ricini, which has operated five or six times. She is a little relieved, but still very ill. I prescribed for her a mixture composed of æther, simple oxymel and camphor, to be taken every four hours; and particularly directed her to drink a pint of liquid only during the next twenty-four hours.

21st. She breathes easier, and has had a tolerable night. Her pulse is less frequent, and her tongue moister.

Repetatur mistura, & cras mane, sumat haustum catharticum e magnesia vitriolata, infusione et tinctura sennæ, et cum tincturæ jalapii drachma, non bibendo inter operationem.

22d. She has had still a better night, and her breathing is more comfortable, with less fever and less cough. Her physic has operated five or six times; but she informs me, that, during the operation, she drank about half a pint of tea. But as, from her account, the
evacuations

evacuations were considerable, we cannot suppose that the disease was in the least increased by this little deviation.

Capiat, horâ somni, haustum æthereum cum camphorâ, &c.

23d. She has rested exceedingly well, breathes easy, and her pulse is natural; having also very little cough. I ordered the draught to be continued; and sent her also a mixture of the same kind, to be taken three or four times a day.

24th. Excepting a little cough, she is now perfectly well: and can run up stairs with the greatest ease. But, lest a relapse should take place, I directed her to continue her night draught for two nights, to repeat her purging draught once more, and to observe the directions above given, as to liquids, for several days.

On the 28th she informed me, she had strictly observed my directions, and had been well ever since the twenty-fourth.

This second case exhibits an instance of a very rapid recovery from one of the most distressing of the pulmonary diseases, and under the most disadvantageous circumstances.

Her asthma was hereditary, and the patient was ill eight days before I saw her; yet she is perfectly

perfectly cured of a disease, which sometimes continues for many months, and often proves fatal, in the short space of four days. It may reasonably be questioned, whether the annals of medicine can produce such an example. For the asthma has always been mentioned as a disease almost incurable by art, and generally considered as one of the *opprobria medicorum*. At any rate, the observations naturally arising from case first, and the method of cure followed in both, together with the experience I have had in treating the other cases which have come under my care, seem to me to prove that this disease, when idiopathic, most commonly originates from too great fulness or over-distention of the blood vessels of the lungs: which, in these cases, may be justly considered as the proximate cause of the disease. For here I consider the difficulty of breathing to arise, not from constriction, but from compression, of the air vessels by means of turgid or over-distended blood vessels: nor do I conceive it necessary there should be constriction of the bronchia, as the compression above mentioned, by hindering the ingress of the air into the more minute branches of the air vessels, will account for the *respiratio alta*, without having recourse to spasmodic constriction,

striction, which seems to have done much mischief in the treatment of this complaint. I would, therefore, rather consider the proximate cause of the disease to exist more generally in the blood vessels, than in the air vessels: and the constriction of the bronchia, when it happens, rather as a symptom than as the cause of the disease. And, although spasmodic constriction may occasionally be produced from vapours, or airs of different kinds, as well as from other causes, and then become the sole original disease; it will, however, in general, either speedily destroy, or be speedily removed.

Dr. Withers, who treats of the asthma as a convulsive disease, observes, page 28, "all convulsive disorders are sudden in their attack, and they are often as sudden in their termination." But this is not commonly the case in asthma; for notwithstanding that the severe exacerbations may soon disappear, yet the disease still exists, and will often continue for many weeks: sometimes it will exist in a moderate degree, at other times great exacerbations will come on and continue, with very little remission, for several days. From the two cases above related the reader will be convinced, that they are not histories of a spasmodic disease,

case, which comes on suddenly, and goes off in a moment; but rather a detail of morbid affections, arising from full and weakened pulmonary blood vessels, and which disappear gradually as these causes are removed. That the asthma is more generally occasioned by over-distended blood vessels compressing the bronchia, agreeable to the notions already hinted, seems acknowledged by the practice of the most respectable physicians who have written on this disease, although they were wholly unacquainted with the principle upon which their occasional success depended. Our principles therefore are particularly applicable in asthma, and will in general produce the most salutary effects. Indeed since I adopted the idea of moderate drinking in affections of the lungs, and have paid particular attention to their diseases, several cases of periodical asthma, as already observed, have come under my care, which speedily yielded to the above plan of cure.

Here I wish to observe, that I am well aware that Nosologists have mentioned a species of asthma under the name of asthma plethoricum: but the intelligent reader must perceive that the description now given, as well respecting the precise state of the vessels, as the method of treatment,

treatment, has not yet been properly observed by any author.

In the treatment of the asthma we have been describing, where no other disease of the lungs has given rise to it, the following two general indications of cure naturally present themselves, viz. 1st, to diminish the quantity of fluids in the blood vessels, by which their over-distention is kept up. 2d, To restore the contractile power to these vessels, after the turgescence and distention have been removed.

These indications we shall notice hereafter; whilst, in the mean time, we proceed to point out the passages of those eminent physicians, whose writings confirm the practice we have adopted, although they were by no means acquainted with our general principles or their mode of application.

The first of the respectable physicians just mentioned is Sir John Floyer; whose treatise on this disease has been deservedly held in high estimation, on account of the many excellent practical observations therein contained. This respectable practitioner, after a long unpleasant experience in his own person, and an attentive observation of this complaint in others, seems clearly of opinion that fulness and distention,
arising

arising from effervescence, (as he expresses himself) of the blood of the lungs, are the chief causes of the asthma; as the following passages from his treatise will clearly demonstrate.

In his dedication* he observes, “and though
 “it still be a dispute, whether the motion of
 “the heart in a fever be by an irritation of the
 “fermenting blood, or the disordered spirits,
 “yet it will be the same thing as to practice:
 “for by whichsoever of them the *rarefaction*
 “happens in the asthma, I must level my me-
 “thod against the *effervescence*, if I will cure the
 “fit or prevent it:” and in the 29th page of
 the same treatise, he says, “The blood of asth-
 “matics is very subject to *effervescencies*; and
 “whatsoever produces that, occasions the fits.”

Also, page 30, “the nature of the asthma
 “consists in a slow effervescence, or ebullition
 “of our blood, on which the several symptoms
 “of that disease depend.” Sir John accordingly found that heated rooms, hot weather, the heat of the bed, volatile and all other heating medicines, tended to produce and keep up his complaints; while cool air, cold weather, and cooling medicines never failed to give relief.

* Vide a Treatise of the Asthma, 3d edition, ded. p. iv.

The diet which agreed best was the cool and temperate. From which he was of opinion, that Hippocrates's observation, "if a man eats and drinks little, he shall have no disease," might be applied with propriety to asthmatics. That the latter part of the venerable old man's observation is here strictly applicable, will, I trust, appear evident from what we have already observed. And had Sir John known how to apply it in the manner mentioned in these observations, his asthma, which continued about thirty years, might have doubtless been cured in a short time, and a relapse thereby prevented.

This sagacious physician, indeed, found, from actual experience, that much drinking was hurtful; and, therefore, recommended moderation. But, although practice had thus convinced him of the good effects of moderate drinking, he still remained unacquainted with its particular application; as will appear from what follows. For, page 78, he observes, "no distemper requires more orderly diet than the asthma; but especially a moderation in drinking, that the serum of the blood may not abound too much, and a constant use of those liquors that are moderate neither too hot nor too cool; but

“ but suited to the constitutions and age. I know
 “ a fat asthmatic who was much relieved by
 “ drinking very little of any liquor.” Now,
 although Sir John thus recommends moderation
 in drinking, yet, when he tells you, in the same
 page, that he himself generally drank a quart
 of broom small beer at dinner, you will imme-
 diately perceive he had no very correct idea
 of the nature of this principle or of its proper
 regulation.

The next author, whose authority we shall
 adduce in support of our doctrine, is the justly
 celebrated Dr. Cullen: who, when treating of
 the asthma, observes, “ From the whole of the
 “ history of asthma now delivered, I think it
 “ will readily appear, that the proximate cause
 “ of this disease is a preternatural, and in some
 “ measure a spasmodic constriction of the mus-
 “ cular fibres of the bronchia; which not only
 “ prevents the dilatation of the bronchia ne-
 “ cessary to a free and full inspiration, but
 “ gives also a rigidity which prevents a full and
 “ free expiration. This preternatural constrict-
 “ tion, like many other convulsive and spaf-
 “ modic affections, is readily excited by a tur-
 “ gescence of the blood, or other cause of any
 “ unusual

“unusual fulness and distention of the vessels
“of the lungs.”

The intelligent reader will perceive that, although Dr. Cullen, in the former part of this quotation, from not being able in any other way to account for the different phenomena of the disease, coincides in opinion with the most respectable practitioners who had gone before him in supposing the asthma a spasmodic disease; yet, towards the latter end, is obliged to acknowledge the unusual fulness and distention of the vessels which we contend for; although, even there, he does not mention the over-distention, which may certainly happen to every hollow muscular structure.

The Cure.

We shall now beg leave to point out a method of cure founded on the principles which, I trust, we have fully established. Here, as we have before observed, two principal indications present themselves, viz. 1st, To take off the too great fulness and over-distention of the blood vessels. 2dly, To restore their contractility

tractility and tone after they have been sufficiently emptied.

Respecting the former of these indications, we have, when speaking of hæmorrhage from the lungs, considered the different methods commonly employed for emptying the blood vessels and taking off their too great fulness and distention; to which the reader will be pleased to refer.

He will there see, that, together with our *limitation of liquids*, moderate bleeding and purging have been recommended; both which may be also used in asthma, although with a little more circumspection towards the latter stages of the disease.

We shall just say a few words on these separately, and notice any peculiarity which we think worthy of notice in the treatment of this disease.

Bleeding may be occasionally necessary in the cure of asthma; but the intelligent practitioner will recollect many reasons, why it should be avoided, if possible, in the treatment of patients of this description. In some urgent cases, however, where immediate suffocation threatens the life of the patient, this operation must be performed without delay. But when

the situation of your patient will allow you to begin with vomiting and purging, that is, with emptying the vessels without at the same time filling them again by the usual dilution, the lancet will commonly be unnecessary—I have never used it in any of the cases above mentioned. And, in the very advanced stages of the disease, when the patient's strength is nearly exhausted, great caution is necessary in recommending it.

Vomiting and purging. In the early stage of asthma, when the patient's strength is good, a medicine which operates speedily and plentifully in this way will be of infinite service, as appears evident from the history of case first, and from the case of an asthmatic lady mentioned by Sir John Floyer, page 22. But, when the symptoms are moderate, the vomiting will not be required, providing an active purgative is used and no liquid allowed during the operation. And, in asthma, if the disease arises from over-distended blood vessels, active purgatives, which, while they empty the vessels, tend also to excite their contractile power, will be preferable to the milder ones. Here I must beg leave to observe, that my practice has uniformly contradicted the idea commonly entertained

tained respecting purging in diseases of the chest, viz. that it seldom relieves the vessels of the thorax; for I have constantly found great advantage from their use, and think them absolutely necessary in order that our principle of the moderate use of liquids may have its full and speedy effects. That idea is particularly expressed by Dr. Cullen, and (perhaps from his authority) is but too commonly adopted by practitioners in general; from which one of the most powerful means of relieving the lungs, when oppressed, has been neglected. That purging has often cured the asthma must have been seen by almost all practitioners, although not attended to so particularly as a fact of such importance deserved; for surely every one engaged in the practice of medicine must have, occasionally, seen his asthmatic patient cured by an accidental diarrhoea. I have often observed it happen, before I thought seriously of the application of this natural cure to practice. Sir John Floyer, page 91, says, “ a gentle-
 “ woman about 60 years old, being always asth-
 “ matic, fell into a diarrhoea, by which she was
 “ freed from stone, cholic, and *asthma*; but
 “ that being permitted too long, run into a con-
 “ sumptive state, and extremely wasted her fat
 “ body,

“ body, and disposed her to surfeit upon every
 “ occasion: I cured her by rhubarb-purges,
 “ steel and bitters.” Here is a cure accomplish-
 ed by nature and Sir John Floyer exactly on our
 principle. The diarrhœa and rhubarb-purges
 emptied the vessels, and Sir John’s steel and
 bitters restored their contractility, and tone.
 Notwithstanding this remarkable case, Sir John
 repeatedly finds fault with purging, as occasion-
 ing an effervescence of the blood: but here
 his theory outruns his practical observations,
 almost all of which tend to confirm our practice.
 Sir John farther observes, “ I remember an
 “ asthmatic who took some quack-pills, by
 “ which he had twenty or thirty stools; this
 “ very much relieved him.” And, page 176,
 he relates a case from Dr. Willis, cured by
 vomiting and purging. And, according to my
 observation, a natural cure of asthma frequently
 takes place by the vessels being emptied by an
 accidental purging, and the patient afterwards
 going into a cool constricting air, which re-
 stores tone and energy to the whole pulmonary
 system. There are some observations of the
 ancients which give additional support to this
 method of cure.

Ætius says, “ maximum est remedium purgatio fortior per pharmaca fortiora.” And Bellonius “ In difficultate spirandi non est formidanda frequens et magna purgatio.” And Dr. Withers*, notwithstanding the great stress he places on the flowers of zinc, acknowledges, in a case of asthma, even complicated with “ general weakness, relaxation of the stomach and bowels, indigestion, dropical swellings and rheumatism,” where several doses of physic, composed of senna, jalap, &c. were given, that purging gave considerable relief. His own words are, “ She says that her physic always relieves her.” And again, “ The purging physic relieves her materially.” And this case is cured by this plan and the flores zinci. Is not this an ample confirmation of the doctrine we have been endeavouring to establish? For, by purging, the vessels are rendered less full; and by the flowers of zinc, their tone is restored.

In the very last stages, however, of this disease, when the *vis vitæ* is nearly exhausted, and there is considerable secretion of mucus into the bronchia, if the strength is farther

* Vide a Treatise on the Asthma, by T. Withers, M.D. p. 207.

weakened by either bleeding or purging, suffocation will almost instantly follow.

In this respect, the *spare use of liquids* claims a superior consideration; because in no one state of the disease can it be adopted improperly. For, supposing this turgescence of the blood vessels in asthma arises from some previous over-distention or over-stretching of these vessels, from which they have in some degree lost their contractile power, and are thereby disabled from carrying on the circulation with that ease necessary for the healthy actions of the lungs; in the same way as the bladder, when over-stretched or over-distended by a suppression of urine, loses its expulsive power from the same loss of tone and contractility of its muscular coat. In the one case, as well as in the other, these vessels must be kept pretty empty, while, by various means, you endeavour to restore their contractile power. A moderate use of liquids, therefore, is here of infinite importance; and, if properly attended to, the vessels will soon recover their contractility and tone, in the same manner as an over-distended bladder recovers its natural powers of contraction, by the application of proper means, after the distention has been removed. For
the

the same muscular structure exists in the blood vessels of the lungs (as well as in all the other vessels of the body) as in the bladder: and, by leaving these muscular tubes more empty, they recover their contractility and elasticity (which the blood vessels also possess in a certain degree) so as to carry on the circulation through the lungs with that regularity and ease which prevents embarrassment of the air vessels, as well as of the different other branches of the pulmonary system. Thus, in pulmonary hæmorrhage, are the ruptured vessels allowed to unite; and thus, also, in asthma, the compression and constriction of the bronchia are removed; and, with these, the proximate cause of the disease. Whereas, in asthma, if the distention is kept up by plentiful drinking, which is very commonly recommended, the disease will become worse every hour, and may become at last incurable. For, independent of the present difficulty of breathing from the compression of the bronchia by over-distended blood vessels, if this compression is long continued obliteration of the more minute cells of the bronchial terminations must take place; and, in consequence, a permanent and perhaps destructive asthma,

By

By this compression also the mucus in the most minute cells of the bronchia will, most probably, be there arrested, and its different thickened particles become the nuclei of future tubercles, as we shall have occasion to mention by and by. And when these are formed a more permanent cause of difficulty of breathing commences; where, notwithstanding, the limited use of liquids will be of much importance.

Blisters have been, also, recommended in this disease; and, from their occasioning a determination of the blood to the surface of the body, they may prove useful; although, if the foregoing plan is properly pursued, they will be seldom necessary.

Diaphoretics, or those medicines which determine to the surface of the body, without heating or rarifying the blood, may also be, occasionally, used with advantage.

Paregorics, by taking off the irritability of the lungs, will prove serviceable on some occasions.

In the asthma we have just described, unattended with inflammation, if the principles we have mentioned are attended to, the neutral salts and common saline medicines become unnecessary; as well as the long list of antispasmodics

modics commonly employed in the treatment of this disease.

We now come to consider the most probable means of fulfilling our second indication, viz. *to restore the tone and contractility of the blood vessels of the lungs after their too great fulness and distention have been removed.*

For this purpose I have generally found the *vitriolic æther, simple oxymel, &c.* sufficient, and these certainly excite the vessels to exertion, and thus assist them in recovering their contractile power; while the spasmodic constriction of the bronchia (if it exists) will also, most probably, be thereby removed. Dr. Cullen observes, "the vitriolic æther has been found to give relief, but its effects are not lasting." If, however, it is seasonably given, after the vessels have been properly emptied, it will be found an useful remedy: as also the acids, such as vinegar, &c. in so far as they also excite the contractile power of the vessels. And considering the flowers of zinc, so highly extolled by Dr. Withers, as possessing considerable tonic powers, we think they may also be of service in fulfilling the present indication.

But should the patient, notwithstanding the above treatment, remain weak and have occasional

fional shortness of breathing on using exercise, or after any particular exertion, I would recommend the steel and myrrh medicine of the late Dr. Griffiths, and other more powerful tonics, together with cold bathing. The cold bath has been recommended by several practitioners as an effectual remedy in this disease; but cannot be used with safety until the vessels of the lungs have become tolerably empty: and, even during its use, a proper limitation of liquids is of much consequence. The authors I allude to, who recommend cold bathing in this disease, are Cælius Aurelianus, Dr. Baynard, Dr. Millar, and particularly Dr. Ryan, in whose treatise the reader will find a full account of cold bathing in the asthma*.

Respecting the air most proper for asthmatic patients, this must be varied according to the circumstances of irritability of the lungs, and of the precise state of their blood vessels. Hence some breath more comfortable in a moist air, of moderate temperature, while others delight in a dry and keen one. But after the vessels have been sufficiently emptied by the plan of

* Vide Obs. on the History and Cure of the Asthma, by M. Ryan, M. D.

cure above recommended, a cool atmosphere, with the addition of an increased proportion of oxygen, or vital, air, will invigorate the system in general, and tend to restore the proper contractility to the pulmonary vessels.

As the following observations on that disease of horses, commonly called, broken-wind, may tend to elucidate those above, on the asthma, I hope the reader will allow me to add them in this place. Here indeed I may claim the privilege of precedent, as Sir John Floyer has, to his valuable treatise on the asthma, added some observations on a broken-winded mare: which, by the by, would seem to intimate, that he also believed that a resemblance existed between the two diseases.

A few Thoughts on the Broken-wind in Horses and other Animals, as being a disease analogous to the Asthma of the Human Subject.*

Thinking that some analogy existed between the asthma of the human subject, and the disease

* The term broken-wind is the most common English name for this disease; although an unnatural and improper one, as tending to convey a wrong notion of the disease. It is called *pouffe* by the French, which, in English, is *short-wind*; which

ease of horses, commonly called *broken-wind*, and eagerly wishing to afford relief to that most noble and useful animal, I was induced to look into some of the most eminent of the veterinary authors, to see if any of their observations tended to confirm my ideas: and, I am happy to find, that many of their practical observations, as well as their dissections after death, seem to confirm the general doctrine respecting the cause and cure; as, I presume, will clearly appear from the following general abstract of their observations.

The authors I allude to are M. Vitet, *Docteur et Professeur en Médecine à Lyon*; and our own countryman, Mr. Taplin. I shall begin with the latter, interspersing the observations of the former.

Mr. Taplin says, that the broken-wind in horses is a defect of the lungs, and laments that all the dissections and minute investigations, which have taken place, have but little improved his knowledge of this disease, or of the method of cure. He adds, that the most eminent au-

which is a much more natural appellation. An American Negro, when asthmatic, naturally observes, "his wind is short." And this is a very natural description of the *respiratio alta* of the asthmatic.

thors,

thors, who have gone before him, have only either believed or suspected, that the lungs were, by some means, too large for the cavity of the chest*. He himself conjectures, that a viscid, tough phlegmatic matter may gradually accumulate in the lungs, and obstruct respiration; and that, after this obstruction has formed, considerable increase of pulmonary affection may take place from any sudden, hasty and long-continued exertions†; justly observing, that, when obstructions once form, the mischief generally increases. In another place he thinks it may arise from “viscidty of the blood from *coarse, full, and foul* feeding.” And upon this theory Mr. T. founds his method of cure. He advises to remove these obstructions of the lungs by bleeding, and the mild purging balls, slightly impregnated with mercury. He then

* M. Vitet, p. 689, tom. ii. observes, “enfin on observe que la plupart ont les poumons trop volumineux respectivement aux cavité où ils sont renfermés.” May not this be from congestion of blood?

† That Mr. Taplin’s conjectures may be right in some cases cannot be doubted; but that these circumstances do not constitute the general causes of the disease seems proved by the various phænomena of the disease, as well as by the dissections of M. Vitet.

orders

orders soap, gum ammoniac, &c. During this course of medicine he recommends "to be strictly observed, that hay and water are to be dispensed with a sparing hand, so as to prevent too great an accumulation in the stomach or intestines."

Indeed Mr. Taplin, Monsieur Vitet, and all those conversant in the practice of farriery, have generally noticed that much water commonly oppressed the lungs, and aggravated their diseases, particularly that disease now under consideration.

They have, therefore, ordered drink to be given sparingly*, mashes to be left off gradually, and recommended a dry diet. The oppression of breathing was self-evident: it was a practical fact noticed by all; but, in my opinion, satisfactorily accounted for by none.

They have, in general, attributed it to the horizontal position of the animal, and the consequent pressure of a full stomach on the diaphragm: but, had they attended to another fact, that the animal will be often more oppressed some hours after drinking a quantity of liquid,

* M. Vitet, p. 692, vol. ii. mentions, as one of his directions for the cure, "à faire boire le moins qu'il est possible."

when

when that liquid must have left the stomach, and got into the blood vessels, they would have endeavoured to trace the affection to some other source. Mr. T. from the actual observation of the good effects of avoiding much liquid, observes, "For, whether as a preventive or cure, mashes should occasionally be given, and gradually declined, till the food becomes regularly dry."

M. Vitet (p. 693, tom. ii.) quotes a very striking example of an accidental cure by abstinence from liquids. His words are "Les marechaux font attentifs à faire boire les chevaux pouffifs le moins qu'il est possible, étant fondés sur une observation de *Soleysel*, qui constate qu'un cheval pouffif abandonné dans une grange à foin pendant six semaines sans boire, fut parfaitement guéri de la pouffe." I shall take the liberty to add another remarkable passage from the same author which evidently confirms the analogy which we think results from the history of both diseases, and throws considerable light upon our notions of the asthma.

Vol. 2d, page 689, when treating of "*difficulté de respirer sans fièvre*," (pouffe, or short-wind), amongst a variety of different appearances of the lungs of the horse, and of other animals

animals subject to this disease, after death, he says "chez le plus grand nombre de ces animaux on voit les vaisseaux sanguins des poumons, dilatés par beaucoup de sang."

Therefore, respecting the cause of asthma in the human subject, and that of the short-wind in horses, the analogy appears very striking. A horse, from hard exercise, becomes short-winded: the human subject, from any severe straining, running, or any other exertion, by which the blood is determined in very considerable quantity to the lungs, becomes asthmatic.

In both, I consider the disease arises from over-distention of the blood vessels of the lungs. Nor does it lessen the probability of this doctrine to know, that the minister's horse, who is supposed to lead a very retired and easy life, also becomes short-winded. For is it not as well known, that a lazy, inactive life, produces relaxation of the vessels; and that the blood vessels of the lungs possess the same structure, and the same disposition to relaxation, to a varicose or aneurismal state, and consequent congestion, as the blood vessels in other parts of the body?

If,

If, therefore, our analogy is well-founded, and the short-wind in horses be a disease similar to, and arising from the same cause as, the asthma in the human subject, the method of cure should, of course, be the same: which is, first to empty the blood vessels by active purgatives, allowing them scarcely any drink for several days, and by occasional bleeding: and, lastly, to restore their contractile power; for unless the proper tone is given to the over-distended vessels, the disease will return and be continued. The best purgative, I should conceive, would be a bolus compounded of the resin of jalap, soap, and the unwashed calx of antimony, or some calomel. Respecting the other pulmonary diseases of this, and the many other useful animals employed in the service of man, it will appear probable, from the foregoing reasoning and the similar structure of their lungs, that moderate drinking, as well as other parts of the above doctrine, will be equally useful to them, while they are indisposed.

CHAP. IV.

*General Remarks on Pulmonary Consumption,
with some Observations on Catarrh, and Croup;
as also on the Seat, Origin, Formation, and
Termination of Tubercle.*

WE now come to say a few words on that most horrible of the pulmonary diseases, the Pulmonary Consumption: of which also, as already of hæmorrhage and asthma, we do not mean to give a systematic account; but, adhering strictly to our title, only beg leave to offer a few remarks.

With much propriety hath it been observed that man is an inconsistent creature, always fond of extremes. And in no instance is the observation more fully confirmed than in that of the different doctrines which have been delivered on pulmonary consumption. For various and contradictory have been the opinions
of

of authors on the nature of this disease, and as various have been their methods of cure. For while some have contended it was inflammatory, and pursued their bleedings even to the 50th or 60th time*; others have held a contrary opinion, declaring the pulmonary consumption a putrid disease, and contending that the bark was the only remedy. Nor have their sentiments been less discordant respecting the precise state of the lungs which constitutes the disease. For while one endeavours to convince you that it consists in scrophulous tubercles; another, with equal ingenuity, proves to you, that the genuine phthisis cannot exist without ulceration of the lungs.

Yet it is well known that this disease appears in different forms, and takes origin from all the different circumstances we have mentioned; and it has also been sufficiently ascertained that, although active inflammation may sometimes exist, it does not always exist; and that, therefore, notwithstanding blood-letting and the antiphlogistic regimen are often extremely proper, yet they are not always so: nor does the idea of putrefaction or debility appear so generally

* Dr. Dover.

as to warrant an indiscriminate use of the bark. But the practitioner who founds his practice upon the rational basis of the anatomy, physiology, and pathology of the body, avoiding all extremes, and unbiaſſed by theory, will vary his plan of operations according to circumstances; ſometimes uſing the lancet, and ſometimes the bark. And, by an attentive conſideration of theſe authors, he will find in every one of them ſome valuable facts, which he will adopt as circumſtances ſhall require.

In theſe remarks, therefore, we ſhall endeavour to ſteer a middle courſe; and direct the attention of the reader to principles, which we hope are rational, and conſiſtent with the laws of the animal œconomy: by which we ſhall attempt to demonſtrate, that the hitherto declared *opprobrium medicorum*, the hitherto ſuppoſed incurable conſumption, may be attacked ſucceſsfully ſo as, in general, to bring about a cure, if early application is made. And although we have no famous baſam, no ſpecific remedy to propoſe, and no infallible cure for conſumption; yet, we ſhall endeavour to point out a certain plan of treatment, which, when properly adapted to the conſtitution of the patient, and the particular circumſtances of his
caſe,

case, will cure in the beginning, and give infinite relief in the advanced stages of the disease.

And although, respecting the nature of this disease, many doctrines have been delivered, yet one of the principles which we contend for, and which we have already successfully adopted in the cure of the foregoing diseases, has an advantage over every other kind of remedy; that, under whatever circumstances the disease may appear, and upon whatever foundation you attempt a cure, it is not only safe, but absolutely necessary in order that the plan of cure, whatever it may be, may have a fair chance of success. This principle is the *limited use of liquids*.

Now we have before observed, that any part of the body, when diseased, can only recover itself by the natural powers inherent in that part; and that, as no medicine, no application we know, can form the callus of a broken bone, so no particular medicine, we are acquainted with, can remove a tubercle or heal an ulcer of the lungs, if the natural powers of the parts do not exert themselves on the occasion.

But a broken limb has an advantage over diseased lungs, in as much as the former can be placed and retained at perfect ease, while the

natural offices of the latter keep them in perpetual motion; which motion all authors have considered as a great obstacle to the cure. And indeed every practitioner knows that if this natural motion is by any means increased, or kept up, while they are affected by disease, the cure is thereby much impeded, if not entirely prevented.

For, in attempting the recovery of any great affection, either of the constitution in general, or of any particular part, both the body and the part must be, if possible, at rest. For example, a man affected with a fever will recover sooner, if kept quiet in bed, than if permitted to go abroad: and every body knows, that no man can be cured of a fracture, unless he allow the parts to be at rest. Now I trust those principles which tend to place the lungs, when diseased, in the same situation as to ease as a broken limb, when fractured, will be supposed the most rational in attempting the cure of pulmonary consumption. For if the lungs are retained in the same easy quiet state, as a broken leg, the powers of restoration will often, of themselves, accomplish a cure; in the way that the same powers bring about the union of a broken bone. To lessen their continual motion,

tion, therefore, or to *keep the lungs as much as possible at ease*, must constitute one of our chief indications, in attempting the cure of pulmonary consumption, from whatever cause it may arise.

As pulmonary consumption is a general decay or wasting of the body, arising from some previous morbid affection of the lungs, we shall first endeavour to ascertain wherein this morbid affection consists, and upon this found our method of cure. And we will begin by tracing this terrible malady from its most early periods; from the incipient catarrh, which, when neglected, often becomes the fatal source of its direful termination.

Catarrh. This morbid affection, which is commonly known by the term *a cold*, and which has been denominated “a defluxion of sharp serum from the glands about the head and throat,” consists in a considerable determination of blood to the mucous membrane which lines the fauces, the larynx, and trachea arteria, attended with more or less of inflammation, irritation, increased, and often changed secretion, and cough.

The chief exciting cause of this disease has been allowed by all physicians to be a check
of

of perspiration: and they have therefore employed, in order to remove it, all those remedies which have been supposed efficacious in restoring and promoting this cutaneous discharge; among which they have ranked plentiful drinking or dilution as the chief. The cause cannot be denied: but how far the plan of cure, just mentioned, is either rational or successful we shall now proceed to enquire.—First let us see what will be the natural consequences of this obstructed perspiration. When a check of perspiration takes place so as to terminate in catarrh, the cutaneous vessels are constricted, and a quantity of fluid, which should have naturally passed off by the skin, is retained in the habit. To make up for this deficiency of perspiration nature endeavours to rouse the kidneys to exertion, and to increase the exhalation by the lungs: but, unfortunately, the pulmonary vessels themselves are often constricted, and the kidneys are not always faithful to their office; hence there is, in a short time, a considerable increase of the circulating fluids; which, from the external constriction, are either detained in, or determined upon, the different viscera and internal parts: and hence the lungs, as being entirely composed of vessels, and being

more

more exposed than the rest of the viscera, become more or less oppressed; while the above-mentioned mucous membrane, as being exposed to the air in respiration, and as sharing in the general internal determination, becomes more and more irritable and inflamed; and, at the same time, its mucous secretion is more or less altered and increased, and is often so acrid as to irritate the surface extremely, producing perpetual cough. And if that portion of the same membrane which lines the nose be affected in like manner, sneezing will, in the same way, be produced.

The inflammation of this tracheal lining will assume different shapes, and will be more or less acute, according to the state of the constitution in general, or the existing tone and activity of the arterial system: and the state of the secretion will entirely depend upon the state of the irritability and inflammation, although it may be in some degree affected by the atmosphere. In one state of inflammation it will be more or less mucous; in another it will be thin and acrid; and, in another, there will be an exudation of coagulating lymph, which constitutes the most serious of these affections. By this exudation the disease, called croup, is produced;

duced; from it also, when it happens in the air cells, do we suppose that tubercles may arise.

As farther confirmation that the fluids exist in an increased quantity in catarrh, it has been found that blood, taken from the veins of patients labouring under this affection, has generally contained an unusually large proportion of serum.

Doth not, therefore, the increased fulness of the vessels, which we have above-mentioned, point out a contrary method of cure to that commonly made use of, viz. a spare use, instead of a plentiful use, of diluents? We think so: and that the following indications of cure naturally arise from what hath been said, viz.

1st, To lessen the quantity of the circulating fluids.

2dly, To restore the proper determination to the skin.

3dly, To lessen the inflammation and irritability of the pulmonary system in general, and of the internal mucous membrane in particular.

These we shall briefly consider in their order: but shall first notice, in a very few words, some of the usual modes of treating this disease; from which the reader will be better able

able to judge of the different methods of treatment. When a person gets a severe catarrhus affection, he is immediately advised to keep in a warm room, and to drink plentifully of warm diluting liquors: and if a plentiful perspiration is procured in this way, the vessels become thereby less full, and the general determination to the surface allows the inflammation and irritability of the affected membrane to go off, and thus is a cure sometimes obtained. But this method of treatment is liable to some objections. For a patient, treated in this way, is not only frequently weakened by such copious dilution with hot liquids; but, from the increase of perspiration, thereby occasioned, is subject to a relapse on the least exposure to cool air; from which he perhaps get a fresh cold the moment he leaves his chamber, and indeed often to an alarming degree.

Ladies of delicate frames suffer particularly from this kind of treatment, and the consequent increased disposition to relapse. If, therefore, the first indication of cure above-mentioned is proper, the great dilution usually adopted, as being quite contrary to that indication, must be improper; as, by it, the vessels, instead of being emptied, are filled fuller, unless counterbalanced by

by considerable evacuations; and the pulmonary system becomes thereby much oppressed.

All the patients which I have attended, for some time past, have been treated agreeable to the three above-mentioned indications of cure, which are now to be considered; and they have, in general, speedily recovered.

In fulfilling, therefore, the first indication, the reader will recollect that the quantity of the circulating fluids may be lessened in two ways, viz. by diminishing the quantity of liquids taken into the body, and by increasing the different secretions. And he will adopt either the one or the other of these methods, according to circumstances; and, on many occasions, both of them will be worthy of his attention.

When speaking of hæmorrhage, we have there fully pointed out the good effects of moderate drinking and purging in diminishing the fulness and over-distention of the vascular system; we have found them equally useful in the treatment of catarrh, and of incipient pulmonary affection. The reader will, however, perceive, that the plan of drinking little in this disease is contrary to the usual practice, which, as we have already remarked, constantly en-
joins

joins to drink plentifully of warm diluting liquors.

But, although the irritable fauces are much comforted by the frequent passage of soft warm liquids, yet, when much of them is taken, the patient will feel his breathing oppressed, from the pulmonary vessels being over-filled; and, if this plan is continued for several days, the constitution becomes, in general, relaxed; and, even although a cure should be thereby accomplished, (which is not always the case) the patient becomes much more liable to a relapse.

Whereas, if a cure is obtained by evacuations, and a proper regulation, and limited use, of liquids, the vessels will become sufficiently empty, while the proper action of the skin will naturally take place, without leaving that irritability of the body, and openness of the pores, (if I may be allowed the expression) which dispose to a relapse. The quantity of liquids, which I have found generally to answer best with my catarrhal patients, has hitherto been about a pint, or a pint and an half in the twenty-four hours; and the most grateful and best adapted liquid is the almond emulsion, which
 possesses

possesses some nourishing as well as demulcent properties.

On some occasions, I have advised about half a pint of some soft weak liquid to be taken at night, when in bed, a little warm: which has been sufficient to promote the proper temporary relaxation of the vascular system, and determination to the surface of the body, without inducing that permanent weakness and disposition to relapse, commonly produced by the great quantity of warm diluents usually employed. And indeed the common drink may be tepid, but not hot; and never in great quantity, which always, more or less, embarrasses the pulmonary system. We have already, when treating on hæmorrhagy, had occasion to notice the good effects of purging in emptying the vascular system, and in determining from the lungs: and wherever the vessels are full, and emptying is indicated, as is the case in catarrh, according to our notions, I never allow my patients above half a pint of liquid during the operation of the purge; for it appears a very inconsistent practice, when you wish to empty the vessels by purging, to pour into the stomach much more liquid than the quantity carried off by the physic.

The

The purgative may be given every other morning, for two or three times; but which, like every other medicine or plan of cure, must be regulated according to circumstances. The cathartic medicines, preferable in this disease, seem to be those which will resist inflammation, and carry off most of the superfluous water from the blood; such as those composed of magnesia vitriolata, radix jalapii, &c. &c.

We now come to the *second indication* of cure; which is, to restore the proper action of the skin: but we might rather say, allow it to take place; as, when the vessels are sufficiently empty, the natural perspiration will, in general, proceed as usual. Practitioners, forgetting some of the consequences of the first constriction of the skin, viz. the fulness of the vessels, &c. have, therefore, in endeavouring to remove it, paid no attention to remove this fulness, otherwise they would not have attempted the removal by adding to it by their plentiful dilution.

Indeed the cure by increased perspiration is of all others the most uncertain in this changeable climate, and renders the patient very liable to a relapse, as we have before mentioned. And all heating medicines and drinks, given with this view, as increasing and keeping up the inflammation

of the affected membrane, and as increasing the disposition to the phlogistic diathesis, which often exists in this disease, are hurtful: whereas a cooling regimen, and the saline antimonial, and sedative medicines, not only tend to remove the phlogistic diathesis, and relax sufficiently the vascular system, but also to restore the natural cutaneous discharge, without that danger of relapse attendant on the use of the warm diluting regimen. And, during the above and following treatment, there is no necessity of confinement or nursing in a warm room, from which the inflammation and irritability of the affected membrane will be rather increased and supported: but, on the contrary, the patient is advised, either to remain in a room of very moderate temperature, or to go out when the weather will permit; from which the cure is accelerated, and the probability of a relapse rendered less.

Respecting the air, it is of much consequence to regulate its temperature according to the state of the disease. In the earlier and middle states, a moderate temperature, inclining to warm, will be the best; as the blood will be thereby solicited to the surface of the body, and the irritability of the mucous membrane will be thereby diminished:

diminished: whereas a cold constricting air would increase the causes of the disease, as would likewise a too high temperature.

Hydrogen (inflammable) air, in a certain proportion, might also be useful from its sedative properties. In the latter part of the treatment, when the vessels have been sufficiently emptied, and the feverishness, if any existed, is removed; a moderately cool, dry, and pure air, by giving energy to the pulmonary system, as well as to the whole body, will tend to the recovery of the patient.

The most effectual medicine we know for promoting the natural cutaneous discharge, and which will also assist in fulfilling the third indication, is the sedative antimonial draught recommended when speaking of hæmorrhagy; only that, when I have used it for catarrh, I have, in general, increased the quantity of the antimonial wine *ad vomitum usque*; after which I have, generally, found the patient recover very fast. This saline antimonial draught I have commonly given every four, six, or eight hours, according to the urgency of the symptoms, in the intervals of purging; and with the wished-for success.

The *third indication* of cure requires our next consideration: and the attentive practitioner will perceive, that all the various means mentioned, when speaking of the two former indications, are equally applicable in the present; as tending to diminish the internal determination, irritation and inflammation, as well as to restore regularity to the natural functions. The mild opiate or narcotic in the antimonial draught will here be of great use, in taking off the irritability of the parts affected, and may be increased according to circumstances. Indeed, after the vessels have been sufficiently emptied, and the disposition to the phlogistic diathesis removed, the remaining irritability of the mucous membrane, and attendant cough, may often be removed in one night by adding to the night draught some drops of the *tinctura opii*, and removing the patient next day into a dry and moderately cool air. Thus will the increased secretion be diminished; and, losing its irritating properties, by which the cough is kept up, it will return to its natural bland and mucous state; and thus will the disease disappear. I have said nothing of blood-letting, which, although not often required, may, notwithstanding,

ing,

ing, be necessary on some occasions. In an incipient, or recent catarrh, Mr. Mudge has strenuously recommended the inhalation of the vapour of warm water: and it may, without doubt, be useful on some occasions; although, if the plan of cure, above recommended, be properly pursued, it will generally be found unnecessary. When I have thought this local application adviseable I have found more advantage from a warm decoction of the white poppy-heads, or from an infusion of the cicuta, than from pure warm water: which might readily be imagined, as some volatile sedative property will be conveyed to the parts affected together with the vapour.

Vomiting is also of great advantage in every stage of this disease.

Having briefly noticed this catarrhal affection of the larynx and trachea, we are naturally led to another, viz. the disease commonly called the croup: a disease of the most dangerous nature, and which often, by a very rapid progress, goes on to a most fatal termination.

On the Croup.

That the croup*, when attended with that membranous or pulpy substance, sufficiently described by authors†, is a disease arising from active inflammation of the internal mucous membrane which lines the larynx, the trachea arteria and its branches, and of a very opposite nature to the spasmodic affection which, as resembling the croup in some of its symptoms, has been often mistaken for it, has been now fully ascertained. And, if we consider that the one is a disease of high inflammation, and the other of considerable debility, it will appear evident that the medicines which do good in the one case will do manifest injury in the other; and that it will, therefore, be of infinite importance to make the proper distinction. But that the croup, attended with high inflammation, and consequent exudation is by far the most common in this country, is now, I believe, generally allowed. It will therefore follow, that all medicines of a stimulating nature, which increase active inflammation in general, will be

* *Cynanche Trachealis Cullenii.*

† Vide Dr. Baillie's *Morbid Anatomy*, Dr. Home, &c.

here

here highly improper and tend to keep up the disease. And that, therefore, the stimulating antispasmodics, which may be occasionally proper in a spasmodic affection of the larynx or trachea, will also help to increase this internal inflammation. The antiphlogistic plan of cure, which is now commonly practised, is that which I have seen the most successful in this active state of the disease. In using topical bleeding, I prefer six or eight small leeches, to three or four in proportion larger; because the former, while they occasion as much evacuation of blood as the latter, will leave three or four additional points of external determination: and, as this determination should be increased and kept up as much as possible, I endeavour to promote it by applying a blister immediately after the bleeding, which I allow to remain, if necessary, until a certain degree of ulceration takes place. But, in the early state of the disease, I have generally found that the emetic tartar, given so as to produce severe and speedy vomiting, has immediately stopped the disease.

A few days ago, I gave a solution, containing half a grain of the emetic tartar in each dose, every ten minutes, to a child about a year old, until it produced considerable vomiting:

and the croup was, by this emetic and a saline antimonial mixture, entirely removed in twenty-four hours. It may be worthy of observation, that all the children of this family, consisting of several sons and daughters, born of robust and healthy parents, were subject to inflammatory diseases, particularly pleurisy and croup. And, some years past one child died of the croup, although attended by a very eminent physician: but, it should also be observed, that the disease had existed for three days before any assistance was called.

Respecting the internal exudation generally found in croup, physicians speak of it as an inorganic, membranous or pulpy substance. Now dissection has demonstrated, that exudation of coagulating lymph is one of the effects or terminations of active inflammation in other parts of the body; and that, by this lymph, a vascular connection between the part affected and the neighbouring parts is frequently formed. And it is also believed that the exudation found in croup is formed of the same coagulating lymph which is found on other inflamed surfaces, only with the addition of a certain proportion of mucus, which, in the early stage of the inflammation, is secreted in an increased quantity.

quantity. This exudation appears more pulpy, as being mixed with mucus as well as exuded coagulating lymph; but seems to possess the same property of uniting itself to the surface upon which it is exuded. And although, from being situated in a cavity where air is alternately passing in and out, it may, in general, adhere more loosely to the inflamed surface than other exudations differently circumstanced; yet that, like them, it may also become vascular, I had two very particular demonstrations, which I shall now mention.

C A S E I.

In the year 1788, I was requested to visit a child, between two and three years of age, who had laboured under the croup for the four preceding days. As the disease was well marked, I informed the parents of the great danger the child was in, and expressed my sorrow that it had been allowed to proceed thus far without any medical assistance. I immediately sent a strong solution of the emetic tartar, to be given *gradatim* until it produced plentiful vomiting; intending afterwards to apply blisters, and use other means for the recovery of the patient.

The

The emetic answered very well : but, in the action of vomiting, the adventitious membrane or substance was separated, and soon after brought into the mouth. The mother attempted to remove it; but was unsuccessful. It was drawn back into the throat; and, a portion of it shutting up the aperture of the larynx, the child was thereby suffocated.

Soon after the child's death the membranous substance was perceived, and having been removed, was destroyed; so that I never saw it. But having obtained permission just to look at the inside of the larynx and trachea, I found a raw tender surface, upon which appeared different spots of blood, like as when the dura mater is torn from a fresh cranium. These appearances seemed to me to point out a connection by means of vessels; and which is evidently confirmed by the following case.

CASE II.

In November, 1791, the niece of Mrs. Day, who now lives in this neighbourhood, a young woman about eighteen years of age, while under the small pox, was, on the third day of the eruption, suddenly seized with the croup. This
circumstance

circumstance was mentioned, on the same day, to the practitioner who attended her: who being, as I presume, unacquainted with the nature of the disease, told her aunt that she had only a common cold which was of no consequence. That he really thought so was evident, as he prescribed nothing for her, and did not visit her again until after I had been called to her in the act of suffocation from this new-acquired disease.

I saw this young woman late in the evening, on the sixth day of the eruption, when I found her sitting up in her bed, almost black in the face, and nearly in a state of suffocation. I was informed, that she had been three days affected with what evidently appeared, from the description of the symptoms, to be the croup; and that her breathing had been extremely difficult for the greatest part of that day; during which time she was only able, on account of her breathing, to swallow a tea-spoonful of liquid at a time. While she was exerting herself in endeavouring to speak to me, she brought up a large portion of the adventitious pulpy substance above mentioned, in one piece, which was about two inches long; and was thereby much relieved. Notwithstanding, I sent her an
 emetic

emetic draught, composed of emetic tartar and ipecacuanha wine, which operated very well; and brought up a considerable quantity of the same substance, in detached pieces. After this, she became greatly better, and could breath with ease, although very hoarse. One surface of the largest portion of this adventitious substance, which was evidently that next to the larynx and trachea, was studded with bloody specks, similar to those mentioned in the first case; so that there appeared clearly to have been a continuation of vessels between the internal surface of the larynx and trachea, and the adventitious substance formed in consequence of the inflammation.

Do not these appearances confirm the active nature of this inflammation, and also direct us to studiously avoid all warm stimulating medicines in this disease? The asafœtida, having been found occasionally of advantage in the spasmodic *suffocatio stridula*, has been also recommended by some practitioners, in every state of this disease, without discrimination; and even when attended with the most active inflammation. But it will, in general, be found, that, even in the spasmodic state of this disease, if the asafœtida does not occasion vomiting, or some other

other evacuation, it will by no means produce those antispasmodic effects generally expected from it. Dr. Home observes*, when speaking of *asafoetida*, “as it heats and quickens the pulse, “it must always be improper in inflammatory “cases.” This young woman remained hoarse for five or six days; during which time she coughed occasionally, and therefore took some paregoric and gentle diaphoretic medicines. The small pox proceeded as usual, and she got perfectly well.

*On the Seat, Origin, Formation, and Termination
of Tubercle.*

From what has been said on catarrh and croup, it will be readily perceived that, when either of these diseases, both of which consist in certain inflammatory affections of the internal lining of the larynx, trachea, and bronchia, is neglected, the inflammation may at last extend to the lungs themselves, and give rise to the affections we are now to consider, which often terminate in the pulmonary consumption.

* Vide Clinical Experiments.

Therefore,

Therefore, from the trachea arteria, and its branches, the bronchia, we naturally come to the morbid affections of the air cells, which constitute the ultimate terminations of these branches, where the fatal source of pulmonary consumption often commences.

Practitioners have generally mentioned two morbid states of the lungs which they have considered as chiefly giving rise to pulmonary consumption, viz. tubercle and ulceration. The former, as it constitutes that affection of the air cells we have just mentioned, we shall first endeavour to investigate; and afterwards proceed to say a few words on ulceration: a very common consequence, or termination, of tubercle.

Tubercle, therefore, is a circumscribed, and generally hard body, bearing some analogy to an absorbent gland; and sometimes many of them, of different sizes, viz. from the smallest particle to near an inch in diameter, will be found more or less dispersed throughout the lungs of consumptive patients*. But it may be

* This account of the size of tubercle agrees with that given by Dr. Stark; who, when speaking of their cavities, observes, that they are "from the smallest perceptible, to half an inch, or three quarters of an inch, in diameter."

proper

proper to observe, that, although this is the more common appearance, yet it will be seen hereafter that they may get to almost any size from extended exudation, adhesion, &c. &c.

As the nature of these seemingly glandular bodies is acknowledged by the most respectable authors* to be but little understood, I shall, with every due submission, beg leave to offer a few observations on their origin and formation; as well as respecting their situation, which appears to me to be erroneously stated even by the best authors who have written on this disease.

As the phthisis pulmonalis happens most frequently to scrophulous constitutions, and tubercles have generally been found in the lungs of those patients, resembling the scrophulous absorbent glands in other parts of the body, they have concluded that these tubercles were morbid enlargements of some little, almost invisible glands, which they have described as existing in the connecting cellular membrane of the lungs, and which they have said opened at last into the bronchia.

Now as these glandular bodies have never yet been seen in the natural healthy state of the

* Dr. Simmons, &c.

parts, we have, therefore, no reason to suppose that any such exist; nor is their existence necessary in order to account for the formation of tubercle; which we will endeavour to prove to be a disease arising in the air cells, and not in the cellular connecting membrane, as has been commonly supposed. We have already said, that the different bronchia terminated, each in a particular cellular termination, consisting of a number of minute cells communicating with one another; but which we believe to be a distinct aërial apparatus, and to have no communication with the common cellular membrane*.

Now into these cells, as well as into the whole internal surface of the bronchia, is secreted more or less of mucous for their defence; which is liable to many alterations, according to the state of the atmosphere, or of the secreting surface.

In some particular states of atmosphere and surface, this soft lubricating mucus will become thickened, and adhere so firmly as to be brought up with the greatest difficulty.

Now, we trust it will not be thought improbable to suppose, that a particle of this inspissated mucus, stagnating in one of these minute

* Vide page 4.

air cells, may give rise to tubercle : and, as, in an inflamed state of this internal surface, there will often happen an exudation of coagulating lymph, which is well known to adhere frequently with firmness to the surface where it is exuded, we may also conceive that some particles of this coagulating lymph, lodging in the air cells, may likewise become the nuclei of tubercles. That tubercles generally arise in the air cells seems farther confirmed by the observations of the very respectable and learned Dr. Simmons* : his words are, “ and it is not unusual for millers, “ stone-cutters, and others, to die consumptive, “ from their being so constantly exposed to dust, “ which in these cases probably acts by pro- “ ducing similar concretions (viz. tubercles). “ I have seen two instances of this sort in mil- “ lers ;” and Dr. Kirkland observes, “ that “ scythe-grinders are subject to a disease of the “ lungs, from particles of sand mixing with iron “ dust, which among themselves they call the “ grinder’s rot. Many instances in this way may “ be met with in Ramazzini, Morgagni, and other “ writers.” Here it is evident that these extra-

* Vide Practical Observations on the Treatment of Consumptions,

neous substances were drawn into the air cells by inspiration, and *there* (and not in the connecting cellular membrane) gave rise to the tubercles which destroyed the patients and were found in their lungs after death: and had each tubercle been as large as to have included the whole of the air cells, up to the bronchial branch to which the cellular termination was attached, they would have put on the same appearance as those described by Dr. Stark, as we shall hereafter mention. Do not the circumstances which attended the experiment on the dog, as mentioned by Dr. Saunders*, give additional support to this opinion? His words are “Two
 “ drachms of crude mercury were injected into
 “ the crural vein of a dog: after a short time,
 “ he became feverish, with dyspnœa, cough, and
 “ daily increasing symptoms of diseased lungs,
 “ of which he died. His lungs were found full
 “ of tubercles, each of which contained a glo-
 “ bule of mercury, forming, as it were, its nu-
 “ cleus.” For I presume it will be readily believed that the mercury was secreted into the different cellular terminations of the bronchia, and that, being there confined, its particles be-

* Vide Treatise on the Liver, p. 302.

came the nuclei of different tubercles, as by him related. And had it been secreted into the common cellular membrane, would it not, by its specific gravity, have all fallen to the most depending parts of the lungs; and there formed only one, or two tubercles, instead of the many which were said to have been produced, and which we presume were not confined to any particular depending parts of the lungs, as no mention is made of that circumstance?

From the account of tubercle given by the late ingenious Dr. Stark, it appears that he was of opinion, that tubercle originated in the connecting cellular membrane, and that it had no communication with the air cells or bronchia until it had gotten to a certain size: but his own words seem to me to prove, that the chief seat of tubercle is in the air cells, agreeable to our notions. For he observes that, when tubercles have arrived at a size exceeding half an inch, they “ have constantly a round opening made by a branch of the trachea.”

Now, how could this constantly happen if tubercles originated, as is commonly supposed, in the lateral connecting cellular structure? Are not the air cells the ultimate terminations of the tracheal or bronchial branches? They certainly

are : and it hence appears evident that our opinion receives additional confirmation from this pathological fact. It may be said that tubercle, according to our ideas of its formation, may also arise in the common cellular membrane. It certainly may, as well as the different scrophulous tumours which happen in the common cellular membrane of other parts of the body : but it must be granted that it will be much more apt to happen, and more easily produced, in a part where mucus is secreted, and where solid and heavy substances may by inhalation be received and become nuclei, than where there is only the fine interstitial vapour.

Supposing then our ideas, respecting the seat and origin of tubercle, to be well founded ; let us next see how we can account for the formation and appearances of this apparently glandular body.

If a particle of thickened mucus, of coagulating lymph, or of some heavy extraneous body, should stagnate in the air cells, what will naturally follow? First, we presume there will be more or less of irritation, and consequent increased secretion of mucus, or exudation of lymph ; from which the original particle, whether mucus, coagulating lymph, or extraneous substance, will
receive

receive an additional covering, together with, perhaps, adhesion of one or more of the air cells; from which, additional irritation, secretion, and still farther addition and adhesion will take place; until at last this enlargement so irritates and so interferes with the œconomy of the lungs, as to produce inflammation and perhaps ulceration and hæmorrhage.

When the exudation or stagnation, which constitutes the tubercular nucleus, happens, there will be naturally absorption, or exhalation, of the thinner parts, and the tubercle will be formed of a substance composed of mucus, lymph and air cells or extraneous substance, alternately, and almost intimately united: and as we know that the coagulating lymph unites parts, we can easily conceive that the air cells will be, by this intervening substance, firmly united, so that the whole will form the little tumour called tubercle. Yet we have every reason to believe that the particular vessels of the air cells are only included, and not firmly united, with the tubercular mass; for, from different changes which dissection points out to have taken place in some tubercles, it would appear probable that some vascular action continued to be occasionally exerted within them. But there may perhaps be

another sort of tubercle, of a solid, hard, and an inorganised substance; but which, most likely, is formed in the way which we have just mentioned, and which would differ from the former in as much as there would be a complete obliteration of every cell and vessel. That some tubercles are organised, there can be no doubt from the circumstances already pointed out: but if there are others inorganised, they may then form two distinct species, of which the latter may be more compact in its structure, when cut into; and may therefore bear more resemblance to an absorbent gland than the former, in which some interstices may be perceived when submitted to this mode of examination.

Tubercles have been found of different colours, some appearing of a white, and others of a brown, colour. Now may not this variation of colour be merely accidental, and depend upon the nature of the secretion, or exudation; or perhaps be owing to a little blood, or to something inhaled into the lungs at the time of the stagnation or formation of the tubercular nucleus? Some tubercles contain fluids of different kinds, as pus, a watery fluid, &c. &c. these fluids must be secreted or exuded. Is not this a proof of arterial connection, either by vessels

vessels carrying red blood, or by lymphatic arteries? We also often find ulcerated little orifices, which we presume are the efforts of nature to get rid of the contained fluid; and which will, I doubt not, now be allowed to be from the action of the absorbing vessels. Dr. Stark observes that there are no vessels to be seen in tubercles, even when examined with a microscope, after injecting the pulmonary artery and vein: but, with all due deference to this ingenious author, I am of opinion that he ought to have endeavoured to discover their vascularity, not by injecting the pulmonary artery and vein, but by injecting the bronchial arteries, which, as being the proper blood vessels of the bronchia and air cells, must certainly communicate more or less with tubercle. For, as we have already observed, the chief business of the pulmonary arteries and veins, is for the exposition, of the blood in the lungs, and for transmitting it from the right to the left side of the heart; whereas that of the bronchial vessels is for the nourishment and support of the whole pulmonary system, including even the vessels of exposition and transmission. So that although no vascularity appeared in tubercle on injecting the pulmonary artery and vein, we must not therefore conclude

that tubercle has no vessels: for the pulmonary artery and veins, as being only for performing the functions above mentioned, have no communication with tubercle whether it be seated in the connecting cellular membrane or in the air cells.

Had Dr. Stark, therefore, injected the bronchial vessels with fine injection, he would have, without doubt, discovered vascularity in some tubercles; although, as we have already noticed, there may, perhaps, be some wholly without vessels passing into their substance.

Whether the orifices of the absorbents of tubercle, which are the chief agents of ulceration, be large enough to admit mercury, in the way that the absorbents are every now and then filled, by plunging a tube filled with mercury into an absorbent gland, I am not able to determine: although I shall take the first opportunity of investigating that circumstance. But whether they are, or not, the different changes which take place in tubercle sufficiently shew their existence.

Hitherto I have endeavoured to give a general idea of the seat, origin and formation of tubercle: but, for a more minute account of their progress, and the various circumstances or
changes

changes which take place in them, together with the different other appearances they assume, I beg leave to refer the reader to the following pathological authors, viz. Morgagni, Dr. Stark, my learned and ingenious friend Dr. Baillie and others; while I go on to offer a few remarks on their more common termination, viz. in inflammation, abscess, and ulceration. But it may not be altogether useless first to inquire what relation exists between tubercle and scrophula, and whether the former be really a scrophulous disease, as is generally imagined. As tubercles bear very strong analogy to the absorbent glands, whenever the latter have been diseased, and the lungs at the same time affected, practitioners have said that the same disease existed in the lungs, and that their absorbent glands were also enlarged: although it is well known that no absorbent glands exist in the substance of the lungs. Now we have already endeavoured to convince the reader that, whatever analogy tubercles may bear to absorbent glands, which are the chief seat of scrophula, they are notwithstanding very different substances: therefore, although scrophula may exist in the body, and the whole absorbent glands be thereby affected, it is still no certain proof that
tubercles,

tubercles, found in the lungs of these patients, are the same disease. But, as it must still be acknowledged that tubercles and consumption more frequently occur in scrophulous constitutions than in any other, we shall now endeavour to ascertain how this happens.

In the first place it is generally agreed that the scrophulous constitution is a weak one; and perhaps debility alone, or laxity of the vascular systems, will account for all the phenomena of scrophula: for from this cause may arise diminished absorption, and a too languid circulation from want of a sufficient power in the vessels to propel forward their contents; from which stagnation in the smaller vessels in the glands, in the air cells, and other internal surfaces, may ultimately take place; and thus produce all the appearances commonly called scrophulous. It will also be found, that in all weak constitutions the mucous secretions are more abundant than in the stronger ones: it therefore follows that, as the internal surface of the aërial system is a mucous secreting surface, and as there will be more of that secretion in a scrophulous patient than in any other, stagnation in the air cells, and the other circumstances which give rise to tubercle, are more likely to happen
to

to them than to other more robust habits, where less of this mucus is secreted. It may also be observed that consumption may be more apt to happen in scrophulous constitutions, as in them the commencement is insidious; and steals on by such slow and imperceptible degrees, that the disease has often arrived at a very advanced state before you are aware.

That the mucous secretions are much increased in weak constitutions must be evident to every practitioner who has attended to the great quantities of slime or mucus which is in general discharged from the bowels of weak, rickety, or scrophulous children; and where the other mucous discharges, as from the nose, the lungs, &c. are also in considerable quantity. And this disposition to increased secretion of mucus will often continue to the most advanced age: hence some people have a great and constant discharge of this kind from the lungs, which often accompanies them through life, and frequently terminates, at last, in pulmonary consumption. For when this secretion is not carried off by expectoration, stagnation, and consequent mischief will follow: and it will be generally found, as we have just now observed, that these people will die consumptive, unless

unless they are speedily carried off by sudden death, or some acute disease.

If, therefore, our notions of the nature of tubercle are well founded, it will appear probable, that tubercles have no farther connection with scrophula than that, from the circumstances above mentioned, they are more likely to be produced in a scrophulous patient than in any other: and a very respectable and ingenious author has observed, "I am fully of opinion, that at least nine in ten of those who die of consumption are scrophulous subjects*."

On Inflammation, Suppuration, and Ulceration of Tubercle.

When a tubercle has acquired a certain size, it begins to disturb the natural offices of the lungs. The blood is prevented or impeded in its passage through that portion of them, and respiration is more or less embarrassed.

Mr. Mudge, in his treatise on the catarrhus cough, p. 45, observes, when speaking of tubercles, "for by obstructing the regular circulation of the blood through the small, and in-

* Dr. Hamilton's Obs. on Scrophulous Affections, p. 27.

deed sometimes larger branches of the pulmonary vessels, they produce a distention of their sides, which by this means becoming thin and weak, frequently burst during the violent exertions of the cough; the consequence of which is an hæmorrhage, always alarming, and sometimes fatal." But supposing this hæmorrhage does not take place, yet the whole pulmonary system will become irritated, and particularly the part or parts affected: hence will arise a severe cough, and by and by inflammation will be produced, which, if not early attended to, will often terminate in an abscess, which may soon occasion ulceration, and all its dreadful consequences. The tubercular inflammation, which every now and then is confined to the tubercle itself, and always includes it, will often so affect the neighbouring parts as to form that disease of very considerable extent which is particularly described by Dr. Baillie*. For according to the nature of the stagnation, the irritation and extent of the consequent secretion or exudation, will either tubercles be formed or this more extended disease. In the organised tubercle, when the inflammation is confined to

* Morbid Anatomy, p. 49.

the tubercle itself, and matter has been formed in it, ulceration generally takes place, producing one or more openings by which the matter is poured into the bronchia, by the natural motion of the lungs, from whence it is expectorated. And when this communication with the cavity of the bronchial branch, is once formed, the air received into the lungs by inspiration will have more or less of access to the internal surface of this tubercular abscess, and often occasion additional irritation and cough. On opening into these abscesses, their internal surface is in general an unequal, ulcerated surface, although, every now and then, it is lined with a smooth membranous cyst.

The termination of tubercle in ulceration, and which is the most dangerous, is now generally allowed to be the work of the absorbing vessels; and is considered as nature's effort either to remove, or get out of the way of, an irritating substance; for whatever irritates a part to a certain degree, excites to this kind of action of the absorbing vessels. Whenever, therefore, a certain irritation from tubercle takes place, either the surrounding absorbents, or those in the tubercle itself, are thereby stimulated, and set to work to remove the irritating body;

body; either by the complete absorption or removal of the whole of it, or by endeavouring to get out of its way by ulceration, which is also accomplished by the same vessels. These circumstances clearly point out the two different actions of the absorbing vessels, viz. the one complete absorption, or absorption without ulceration; and the other absorption, with ulceration. May not the former be an increased exertion of their natural healthful functions; and the latter, an angry action, which they never adopt but when obliged?

For it is observable that whatever teazes them excites to this kind of action, as poisons, too great friction, &c. And perhaps this irritable action of the absorbing vessels in carrying on ulceration may be kept up by an acrid secretion from the blood vessels. Or does the matter, or ichor produced, and the ulceration depend upon a certain weakness or morbid irritability of both the sanguiferous and absorbing systems? It may be questioned, whether the two different actions of the absorbing vessels just mentioned depend upon different states of the vessels themselves, or upon the diversity of stimulus impressed by the different stimulating or irritating causes. We suspect the latter: and conceive

ceive that one kind of stimulus excites to healthy vigorous action; while another excites to a more feeble, irritable, or angry action, if, in the language of a late celebrated physiologist, I may be allowed to use the expression. After these general remarks, we proceed to say a few words on the

Prevention and Cure of Tubercle.

Vomiting. If tubercles take origin from different substances lodged in the air cells, agreeable to our notions of them, one chief indication, by way of prevention, will be to dislodge these substances; and vomiting will constitute the principal mean. In scrophulous constitutions, therefore, where there is great secretion of mucus in the aërial system, and where the patient is thereby more liable to the disease, a vitriolic emetic, exhibited occasionally, will prove an efficacious preventive: and, either a vitriolic or antimonial emetic should never be omitted on the first attack.

The vitriolic emetic, which I have generally used, has been the white vitriol, in doses of twenty or thirty grains.

While

While the practitioner is thus endeavouring to prevent the stagnation of mucus in the air cells, he will also endeavour to lessen the secretion, by attempting to strengthen the pulmonary system by every method he is acquainted with, viz. by riding, sailing, and particularly by an early and well-directed use of the cold bath; while he at the same time guards against too great fulness of the blood vessels.

When we suppose tubercle is formed, even then the more active emetic medicines, as being very powerful promoters of absorption, and as it is not improbable that tubercles may be absorbed, will prove very valuable remedies.

In the early state of the disease, when active inflammation is present, the emetic tartar is preferable to the vitriolic emetics, as being more adapted for lessening the action of the sanguiferous system; while it at the same time promotes evacuations and determinations, tending to the removal of the disease. But after the disease has existed some time, and the fever is become a fever of irritation or debility, where the action of the sanguiferous system is weak; then the active and tonic emetic medicines, such as the white or blue vitriol, should be adopted.

In my remarks on case third, of pulmonary hæmorrhage, I have observed, why should not tubercles be absorbed? Now it is well known that the absorbing vessels are distributed in great numbers through the lungs, as well as through every other part of the body; and that not only collections of fluids, but solid tumors, and even bone itself, are frequently removed, in different parts of the body, by their means, without any ulceration. And why should not tubercles also be removed in the same way; as likewise scirrhus, or induration of the lungs; for even through the substance of scirrhus there generally remain some vessels capable of their usual offices, as is clearly demonstrated by the increase and diminution, and other changes, which every now and then take place in enlarged and indurated parts? We certainly believe they may, did we but know how to excite or command the vigorous and healthy action of the absorbing vessels. When we have acquired this knowledge sufficiently, we shall then be able to command the processes of complete absorption, and of ulceration with greater certainty. Then, I trust, we shall be enabled to say unto them, remove this tubercle, and they will do it: and,

when

when engaged in the process of ulceration, stop; and we shall be obeyed.

The learned Dr. Simmons was of opinion, long ago, that the increase of tubercle might be prevented by absorption. He observes*, “medicines that operate in a general manner upon the system may, by promoting absorption and diminishing the determination to the lungs, tend to disperse tubercles or to prevent their formation;” and after, “if any remedy is capable of dispersing a tubercle, I believe it to be vomits.” I have generally used the white vitriol, as already mentioned, and given it in the above-mentioned doses, once a week or oftener, according to circumstances. Dr. Simmons recommends the blue vitriol, which may, perhaps, be more effectual. The resolution of tubercles in this way is most desirable; as when they terminate in abscess and ulceration, the situation of the patient becomes more dangerous, although the natural powers are often capable of remedying these morbid affections when properly regulated; and when, by the moderate use of liquids, and the other means recom-

* Practical Observations on the Treatment of Consumptions, p. 29.

mended in these remarks, the different systems are left unembarrassed, and at liberty to exert their powers of restoration.

Now it is well known that ipecacuanha will vomit, and that all emetics will promote absorption to a certain extent; but we conceive that the metallic emetics possess the latter property in a more eminent degree. Here it may not be improper to notice some of the different effects of the emetic substances usually employed, respecting which there seems to exist a considerable difference in the effects produced by their operation. One class of emetics, when given in doses to produce vomiting, proves emetic without having much disposition to purge, viz. ipecacuanha, &c. Another vomits, and often purges also, as the emetic tartar: and a third, while it only vomits without any risk of purging, gives, at the same time, a more considerable and more permanent contractile stimulus to every contractile fibre in the body. The two former are preferable when too great vascular action exists; and the latter, when you wish to excite other actions in the body besides vomiting. By the latter, therefore, you may clear the stomach, promote expectoration, and work other salutary changes in the lungs, without

out weakening your patient so much as by the former. And, in case of vomica, where matter has been formed, and is pouring out into the bronchia, the vitriolic emetic will assist in emptying it, and also tend to contract the cavity so as to prevent fresh formation: it may also, perhaps, excite new and more healthy actions of the absorbing and sanguiferous systems.

Purgatives may also be occasionally adopted in the treatment of tubercle, not only on account of their determining from, and helping to keep sufficiently empty, the vessels of the lungs; but as also tending to promote the action of the absorbents.

The magnesia vitriolata, and different preparations of jalap, are great promoters of absorption, as also evacuants; and, as such, I use them on this occasion. I have found calomel, joined with antimonial powder, produce many good effects.

Moderate drinking. While you are thus endeavouring to prevent the formation of tubercle by vomiting, or, when formed, to promote its absorption by vomiting, purging, and other remedies; a due abstinence from liquids, by allowing greater freedom of circulation through the lungs, and the absorbents more liberty to

act, as well as permitting the proper expectoration to take place, will be of considerable service, and should be constantly kept in view.

The absolute necessity of attending to this principle will appear more forcible if we recollect that, in consumption arising from tubercles, the disease now under consideration, as well as when that disease is accompanied with vomica and ulceration, it often happens that the vessels of the lungs become gradually more and more impervious; until, from fresh inflammation, adhesion and obstruction, the greatest part of the vessels are obliterated, and the patient, at last, thereby destroyed. This is particularly demonstrated by the dissections of Dr. Stark; who observes, "The pulmonary arteries and veins, as they approach the larger vomicæ are suddenly contracted; a blood vessel, which, at its beginning, measured nearly half an inch in circumference, sometimes (although it had sent off no considerable branch) could not be cut up farther than an inch; and when, outward, they are of a larger size, yet, internally, they have a very small canal, being almost filled up by a fibrous substance." Thus by this fibrous substance, or coagulating lymph, are the cavities of the larger blood vessels gradually diminished;

minished; while by it also, in the same way, as well as in the manner before-mentioned, when speaking of the formation of tubercle, are the lesser blood vessels also obliterated, forming such impediments to the circulation that it will often be impossible for a fourth part of the quantity of blood, usually circulated through the lungs, when in health, to pass through them. How then is it possible that many additional quarts of liquids, which, if drunk, as is the common practice, in those cases, must pass through the lungs before they are expelled the body (unless they are carried through the bowels by purging), can find a passage, from the right to the left side of the heart, through so small a number of remaining vessels?

Anodynes, and other remedies. While these principal operations are attending to, it will be of infinite consequence to take off irritation, allay cough and fever, by occasional blood-letting, blistering, saline antimonial medicines, a cool spare diet, and gentle anodynes. Respecting the latter I shall just beg leave to make a few observations. In treating diseases of the lungs I have tried opium in every form, and have often found inconvenience from its use: for it is well known that opium, as being a very

acrid stimulating substance, whatever ultimate sedative effects it may produce, tends, in its first operation, to increase the phlogistic diathesis of the system; and I cannot help believing that, in this way, it often keeps up pulmonary inflammation. I have found the following the most effectual sedative in the treatment of pulmonary complaints, viz. the Succus Cicutæ Spissatus Ph. Edinb. which is made, by mixing with the inspissated juice, when reduced to a certain consistence, a quantity of the powder. Of this preparation I generally begin with five grains, increasing the dose and frequency of exhibition according to the state of the disease, and the existing irritability of the patient: it seems to me to possess considerable sedative powers, without any stimulating property; and indeed the chief good effects of the cicuta seem to depend upon this direct sedative property*.

Hence it may justly be considered as a medicine worthy of attention wherever you wish to quiet or diminish action, either of a particular part or of the system in general, without increasing or keeping up the phlogistic diathesis or disposition to inflammation.

* Vide the Appendix.

And

And it seems particularly adapted to answer our present indications, viz. to diminish the cough, and suspend the irritations of the pulmonary system, until we, by various plans of treatment, endeavour to remove the existing causes of the disease. By this sedative property, therefore, it may stop or suspend the increase of tubercle, or other morbid enlargements, and consequent ulceration; either by taking off the irritation naturally produced by them, and thus preventing farther determination, or by lessening the irritability of the absorbing vessels, and thus preventing ulceration.

The syrupus papaveris albi, although an opiate, is much less stimulating than the pure opium or its tincture; and is, therefore, in many cases, preferable to the other preparations of this narcotic medicine. This will readily occur to us, if we recollect, that, in this syrup, the stimulating ingredient is sheathed by the mucilaginous part of the poppy-head, which, in a greater or lesser proportion, is always extracted in the preparation.

When we have reason to believe that a patient has tubercles, and have tried the methods above recommended, or others, in order to promote their resolution, without effect; the

next

next indication which naturally presents itself, is, to endeavour to prevent a disposition to inflammation and ulceration, by making nature contented with them. On this principle balls, and other extraneous substances, may remain quietly in different parts of the body for many years: and stones formed in the kidneys, of long standing, and whose size may be presumed to be far beyond the reach of expulsion, can be made to continue quiet, and the patient enjoy tolerable health for a long series of years. For this purpose, a strict sedative plan, as giving the cicuta, swinging*, sailing, &c. is to be pursued; and every thing which can tend, either to occasion or keep up irritation, is to be avoided. The proper determinations from the lungs and other general circumstances are to be attended to; by which, and a particular attention to moderate drinking, the vessels of the lungs will be retained sufficiently empty, until the constitution becomes reconciled to the newly-formed impediments to the circulation through the lungs, and to the other offices of the pulmonary system.

* Vid. An Account of the Effects of Swinging, employed as a remedy in the Pulmonary Consumption, by James C. Smyth, M. D. F. R. S. and Physician Extraordinary to his Majesty.

It may, however, happen that, notwithstanding every effort to prevent them, tubercles will go on to inflammation, suppuration and ulceration: and thus end in the ulcerative consumption, which we next proceed to notice.

CHAP. V.

Remarks on the Treatment of Ulceration of the Lungs, and Pulmonary Consumption.

IN the progress of our remarks we have ventured to state our ideas on the rise, progress and termination of tubercle; and we have seen that a common termination of it was in ulceration, which some Physicians have considered as the sole cause of every true consumption: and indeed almost all the morbid affections of the lungs, which produce that long lingering disease, such as hæmoptœ, long-continued asthma, tubercle, inflammation, &c. generally terminate at last in ulceration; which commonly produces a consumption, more or less rapid. We have

have already observed, that great difference of opinion existed among practitioners, respecting the cause of this disease: and that many, having considered it as an inflammatory disease, had attempted to cure it by bleeding, and a rigid antiphlogistic plan; while others, imagining that it originated from scrophula, had pursued a very different treatment.

It may be worthy of notice, that, in a state of debility, as well as in scrophula, the blood vessels are weak, and seem often to want that contractile power necessary for a free and proper circulation, and hence that plethoric state of the lungs, and of the whole body, frequently present in those states of the constitution. Dr. Simmons, in his book on consumption, already mentioned, p. 8, says, "The genuine phthisis is usually the effect of a certain pre-disposition of body which is very often hereditary."

Now, may not this pre-disposition consist in the relaxation, and consequent plethoric state, of the vessels of the lungs above mentioned; occasioning over-distention of the blood vessels, and hence rupture of them, compression of the ultimate branches of the bronchia, stagnation of mucus or lymph, tubercle, ulceration, and consequent consumption.

This

This would seem probable from what we have said of the asthma, which frequently takes origin from the causes above mentioned: for, in our remarks on asthma, p. 74, we have particularly observed, that, from the distention of the blood vessels, and the consequent pressure upon the bronchia, mucus may be arrested in the air cells, which may produce tubercles, and ultimately consumption. And dissection has shown that all asthmatics, who have had the disease for any length of time, have had more or less of other pulmonary disease: and that tubercles, which we believe may be occasioned in this way, are frequently found in the lungs of asthmatic patients. If, therefore, our ideas of the cause of asthma be allowed, and that tubercle and consumption may be thereby produced, it will be allowed that weak and over-distended pulmonary blood vessels constitute, at least, one cause of the disease. Now, we have already stated, what can scarcely be denied, that, when any part of the body is diseased, that part can only be restored by the proper exertion of the powers naturally inherent in that part; and it will not be disputed, that the lungs possess the same powers of restoration, as any other part of the body: it will, therefore, naturally follow, that

that every impediment to the proper exertion of the natural powers of the lungs will constitute a cause, why ulceration and other morbid affections of the lungs may be the more difficult to cure. Why, therefore, are not ulcers of the lungs as readily healed as ulcers of other parts of the body? This we presume arises from the following causes, viz. 1st, From their constant motion, from which the ulcerated surface is kept in a continued state of irritation. 2d, From a constant exposure to atmospheric air, which is in general hurtful to ulcerations of external parts exposed to its influence, and which may, perhaps, on some occasions, proceed from its containing an increased proportion of oxygen. 3d, From embarrassment of the different pulmonary systems, viz. either from over-distended, or too constantly, or improperly acting blood vessels. From a due consideration of these causes, therefore, do we deduce our present indications of cure, in pulmonary consumption arising from ulceration of the lungs; which we conceive, may be divided into the three following general heads, viz.

1st, To place the lungs, as much as possible, in that easy, quiet state, in which you would place an external part affected with an irritable
ulcer,

ulcer, or where certain circumstances exposed that part to irritation; by removing every impediment to the proper and easy exercise of their natural functions.

2d, To regulate the application, or inhalation of the atmospheric, or other airs, or vapours, so that the actions of the internal ulcerated surface may be either increased or diminished, according as circumstances shall require.

3d, To direct properly the natural powers of the whole pulmonary system in the due performance of its natural functions: to excite the actions of its different branches, if too languid; or to restrain them, if too active.

We shall here just observe, that, in fulfilling these three general indications of cure, it will be perceived by every intelligent practitioner, that the plan of treatment may be so blended, that either two, or perhaps the whole, may be fulfilled at the same time.

The 1st *indication of cure*, we attempt to fulfil by lessening the quantity of the circulating fluids, and by regulating the circulating system so, that the blood may be determined, as much as possible, to the other parts of the body. These purposes we accomplish by flannel, a moderately warm atmosphere, vomiting, purg-
ing,

ing, the moderate use of liquids, and occasional bleeding; by giving sulphur, tar water, and mild volatile demulcent medicines; by rube-facients, dry cupping, blisters, caustics, setons, and other external applications and remedies, which will readily occur to the sagacious well-informed practitioner.

Now we have already seen, that the two great impediments to the removal of any morbid affection of the lungs are, their constant motion in respiration, and the too great fulness of their blood vessels; and that the ease of respiration depends much upon the state of these vessels. We have, also, when treating on hæmorrhage, asthma, and catarrh, pointed out the various modes of removing this fulness: to which the reader will be pleased to refer. We have there clearly demonstrated that the plentiful dilution usually employed is inconsistent and improper; and have pointed out an opposite principle, viz. *the spare use of liquids*; the good effects of which will be readily acknowledged by every candid practitioner who will make the trial. Indeed, in no one pulmonary disease ought this principle to be neglected: we therefore beg leave to enforce it in the treatment of pulmonary consumption as one of the chief means of keeping
the

lungs at ease. Having lately had occasion to consult the justly-admired Aretæus, whose practical acuteness and accuracy has scarcely been equalled in any age or country, I found, in his treatise on the cure of Peripneumony, the following very remarkable passage, viz. "but on the whole, drink should be exhibited very moderately, for moisture is pernicious to the lungs, as they naturally attract, both from the gullet and ventricle." Here his penetration discovered, or rather his practice taught him, that little drinking was proper for the patient; although, being unacquainted with the nature of the principle we contend for, he accounts for it by a false theory. This, however, does not diminish the practical fact; and, therefore, Aretæus's practice evidently confirms the propriety of our principle in the treatment of pulmonary diseases. The quantity of liquid, which I have generally found sufficient in the twenty-four hours, has been from half a pint to a pint and an half. By this regulation of drink the breathing has soon become easier, and the hectic symptoms greatly diminished: indeed the night sweats and colliquative diarrhœa seldom continue during this mode of cure.

One great difficulty occurs in the cure of consumption, which is that many different states of the lungs often exist at the same time, viz. in one portion shall exist a tubercle, in a state of suppuration; in another, tubercles may be forming; and, in a third, the process of ulceration may be going on. In the treatment, therefore, the above general principles can only be adopted. In an ulcerous state of the lungs, how far the exhibition of emetics, as being considerable promoters of absorption, and ulceration being itself an action of the absorbing vessels, is proper, I am at present unable to determine. But I can conceive that emetics may suspend the present existing morbid action, and perhaps induce a mode of absorption more adapted to the recovery of the patient.

Dr. Cullen found that, in many ulcerations, the blue vitriol was more useful than any other kind of emetic. Was this from its giving the contractile stimulus to the ulcerating surface, and thereby constricting, or shutting up the mouths of the absorbing vessels; or was the tone of the general systems of the body thereby increased, and more firm, or more salutary actions both of the sanguiferous and absorbing systems thereby induced? I believe the latter: and
think

think that a vitriolic *emetic* (either of white or blue vitriol), increases the healthy action, both of the absorbing, and sanguiferous systems. But the reader will be pleased to attend, that I do not mean that the blue vitriol may be given in small doses, as a tonic: on the contrary, I believe that this preparation of copper, as well as all the preparations of this metal, act as poisons, when given in this way. I have seen it given in mortifications of the extremities, but with the same effects: it produced debility, anxiety, cold sweats, &c. The white vitriol I have often given, in small doses, as a tonic; and with evident good effects.

Now, if we examine all the cures which have been performed, whether by nature or art, of both of which there are many well-authenticated instances, it will appear probable, that they have chiefly been accomplished by different circumstances which tended to fulfil the present indication, and to put the lungs in that quiet, and easy state, upon which in a great measure depends the proper exertion of the powers of restoration, by which alone recovery is to be brought about.

Bleeding has been the chief anchor of hope to many physicians in the treatment of this disease:

and Dr. Dover, as we have before noticed, has prescribed this operation even to the fiftieth or sixtieth time. And the late Sir William For-
 dyce rested his chief hope upon blood-letting. The quantity taken at one time was certainly small, which used to be the common direction, viz. a frequent repetition of venæsection in small quantities. Feeble indeed must this practice have been, which was only calculated to relieve a symptom, which would recur in a few hours, or perhaps, minutes, after the operation. For it could only relieve, for a moment, some difficulty of breathing, which was perhaps occasioned (or if it was not, would certainly be increased), by the plentiful dilution so improperly recommended in diseases of the lungs. Would it not have been much more rational to have used greater moderation in drinking, and, at the same time, the different means of determination, and thus saved the strength of the patient? For by these bleedings the patient was drained of that vital power upon which alone rested his recovery. The loss of blood relieved him for the moment, in the same way as a loose motion or gentle diaphoresis, each tending to relieve the lungs: but, how much more preferable than bleeding? When we recommend this
 operation,

operation, it is generally under some urgent circumstances, where no room is left for adopting the other means of cure.

Moderate Drinking. Here we shall beg leave to remark that the principle of moderate drinking, which we contend for, has been often adopted without being attended to, and a cure thereby accomplished. In this way I would account for the recovery of most of those consumptive patients, who have been cured by a voyage to Lisbon, to the West Indies, &c. which happens in this way. In the first place, they are generally sick on first going to sea; and if they do not vomit, they are at least several days without drinking much liquid; and indeed the water is in general so indifferent, that they drink but little during the voyage. And if they are also affected with almost continual vomiting, which often happens, it will be readily conceived, that the blood vessels will be left sufficiently empty. Thus, therefore, are the lungs left at ease, in the same way as by moderate drinking; while the change of air, and the sedative effects of the ship's motion, perhaps assist in accomplishing a cure.

On the same principle do I believe that grapes, and other fruits, have cured a consump-

tion, in a warm climate, by purging; while the blood was, at the same time, determined to the surface of the body, and there retained by the congenial warmth of the atmosphere.

Indeed all the accounts of consumption, cured in the confirmed state of the disease, seem to me to be so many histories of particular determinations from the lungs; while, at the same time, the strength was supported, or salutary changes induced, by good air, change of scene, agreeable company, &c.

By the one, the vessels of the lungs were kept sufficiently empty; and by the other, an end was put to the diseased actions, and new and healthy ones produced in their place.

Purging. Although all purgatives, by their stimulus upon the alimentary canal during their action, will in some degree call the blood off from the lungs; yet some purgatives are preferable to others, in attempting to evacuate the superfluous water of the blood by the bowels.

But whenever they are taken for this purpose, little drink should certainly be taken during the operation. Authors have in general recommended to diminish the determination to the lungs, but while they have been using purging and the other means proper for this purpose, their

their plentiful drinking or dilution has ruined every attempt to succeed, by counteracting their endeavours. In the latter stages of the disease much circumspection is necessary in administering purgative medicines. Then the vital powers are too low for attempting this mode of determination; which however has been accidentally successful: and, as we believe, on the same principle of keeping the lungs at ease, by retaining their blood vessels in a moderately distended state. We have before mentioned Dr. Cullen's idea, that purging did not relieve the vessels of the thorax. That the Doctor was mistaken, is very evident from what we have said of asthma, hæmorrhage, &c. But, many more examples could be given, where consumptive patients have been cured by considerable purging; which has happened sometimes, by accident, and, at other times from medicines given with that view. In May, 1793, a striking example of this kind occurred to myself. I was then sent for by a lady, as respectable for her rank, as venerable for her years and virtues, who had laboured under pulmonary affection for some time. She was then bringing up blood and matter, and was confined to her bed. Her pulse in the morning was about one hun-

dred and ten, and possessed some degree of firmness; and her bowels were much confined. As I was not her apothecary, she only sent for me to give her a little opening medicine, to remove costiveness. I, accordingly, sent her two draughts, each composed of half an ounce of magnesia vitriolata, with a little tincture of senna and rhubarb to quicken the operation; with directions to take one immediately, and repeat the other after four or five hours, should the first fail of success. As no effect was produced by the first, she took the second: and they both purged her about twenty times.

She gave me this account next day; and added, angrily, that the physic I had given her was fitter for a horse than for a christian. However she became better from this moment: and the morbid affections of the lungs gradually disappearing, she soon recovered her usual health. I now attend her whole family, and am happy to say that she enjoys at this time the most perfect health, although at a very advanced period of life.

Vomiting is useful, not only by determining to the surface of the body, and promoting expectoration, but also by stopping, or suspending, the existing morbid actions of the lungs.

We

We have already noticed the different effects of the different emetic substances commonly employed; so that it remains with the practitioner to judge when the one, or when the other, is to be adopted. Although, when ulceration has actually commenced, the vitriolic appears to claim a decided preference. In an active inflammatory state, the emetic tartar is preferable to any other; as, besides the property of promoting evacuation, and determining to the surface of the body, it also possesses the power of relaxing, and thus diminishing the action of, the sanguiferous system.

Sulphur has been recommended, by some physicians, in the pulmonary consumption, but, being considered as a very heating substance, and as more or less of fever is generally present in this disease, it has seldom been used. But Dr. James Sims, in his observations on epidemic disorders, observes, page 124, “Sulphur is almost the only laxative which does not diminish perspiration; neither does it raise any degree of heat.” From his recommendation I have often used it as one of the means of determining from the lungs, and of keeping the body open; and with considerable advantage. And the most particular attention to the pulse
never

never pointed out any heating effects, or increased action of the blood vessels in consequence, although given in considerable doses.

Tar-Water may also be given as another auxiliary in fulfilling this indication. There is a cordial, exhilarating property in tar-water, which seems to determine from the centre to the circumference, without heating, and which has appeared well adapted for our present purpose. But when it was used, the other liquids allowed were diminished in proportion. A quarter of a pint of strong tar-water was given twice a day. Tar-pills have also been found useful on the same principle.

Mild Volatile Medicines seem to act in the same way as the tar-water, viz. by determining to the surface of the body.

Rubefacients, Dry Cupping, Blisters, Caustics, and Setons also determine outwardly, while they, at the same time, produce a set of external actions which may, in some measure, call off the nervous influence from the internal ones; and, thus, either stop, or suspend them.

Flannel and Fleecy Hosiery prove of much advantage to the consumptive of this country; by keeping up a regular and uniform determination to the skin, and thus preventing those internal determinations

determinations which, otherwise, regularly attend the sudden changes of this variable climate.

Gum Ammoniac and Squill Vinegar. Pure gum ammoniac, triturated with squill vinegar, so as to form a kind of ointment, has been spread on leather and applied to the outside of the chest, for the purpose of external determination: and often with good effects.

Turpentine Ointment. The following I have used with advantage in endeavouring to keep up the external determination, viz.

R. Terebinthinæ Venetæ,
Mellis, singulorum drachmas duas,
Spiritus Ammoniacæ Compositi drachmas tres,
Axungia Porcinæ unciam unam: misce et fiat unguentum, partibus thoracis externis, bis vel ter die, applicandum.

But an objection may justly be started to the frequent use of external applications, particularly in cold weather; as determination to the lungs is apt to happen, at the moment of using them: the bad consequences of which could not be counterbalanced by their good effects.

In order therefore to keep up a regular determination to the surface of the body, I have often thought that a flannel vest, whose internal surface was thinly besmeared with tar, and re-
newed

newed after three or four days, would answer very well for this purpose without being liable to the objection just mentioned. But I have not yet had an opportunity of trying it. Would a mixture of honey and tar; or would any other substance be better?

We know that the ancients anointed the external parts of the chest with oil with a similar intention.

Here I need not remind the reader, that a certain degree of strength is necessary to bring about a proper determination for the lungs; for he well knows that, under whatever disease the body labours, it must be more or less supported, according to the nature of the disease, and constitution of the patient: nor can the natural functions of diseased lungs be properly performed, unless they possess adequate powers. With a view, therefore, of giving strength, and promoting external determination, particularly in the latter stages of the disease, some white, or light, wines, will be found of considerable service; given either diluted, or alone, according as the practitioner shall judge proper.

Climate. It has been observed, that many insular situations have proved unfavourable to the consumptive. Whether this is from a too cold,

or too changeable atmosphere, or from any thing noxious in the proximity of the sea, I leave to the determination of the more experienced physician. It is generally supposed that the changeableness of our climate renders the inhabitants of Great Britain particularly liable to pulmonary complaints; and that, when these are once formed, the same cause operates as one of the greatest obstacles to recovery.

It is, therefore, usual for the English physicians to order their consumptive patients into a warmer climate, and to one less changeable than our own. And it is observable, that a certain degree of warmth of climate, when accompanied with a certain steadiness of temperature, by keeping up a regular continued determination to the surface of the body, is highly conducive to the cure of pulmonary complaints. Yet it has been found, that, beyond a certain temperature, the fate of the consumptive was accelerated, as appears by the following quotation from a very respectable author.

He observes, page 301, "Pulmonary consumptions rarely originate in the island (Jamaica), but those who come from England, with that complaint already begun, are not benefited by the warmth of the climate; on
" the

“ the contrary, the disease is precipitated, and
 “ proves sooner fatal than it would have done
 “ in a more temperate air. Of this we had re-
 “ peated examples among the soldiers, several
 “ of whom arrived in the island with beginning
 “ consumptions, and were all quickly carried
 “ off by that disease*.”

The same author observes, that in Jamaica, during the hotter months, the thermometer ranges from 85° to 90° . Whether the above effects arise from the heat producing general debility, or from its effect upon the internal surface of the lungs, I do not pretend to determine: perhaps both. The climate, therefore, should neither be very hot nor very cold, nor liable to frequent changes of temperature.

Pregnancy produces another determination from the lungs, which has been found to suspend the fate of the consumptive: and, I believe, that, during pregnancy, the disease may be frequently cured, if proper methods are pursued.

The second Indication. We have already noticed that inhalation, either of air or vapour, is the only mode of local treatment, when the

* Observations on the diseases of the army in Jamaica, &c. by John Hunter, M. D. F. R. S. and Physician to the army.

lungs are diseased. And it has been observed by authors, that one great reason why ulcerations of the lungs are so difficult to cure is their being so constantly exposed to the action of the atmosphere. We shall therefore enquire into some general circumstances respecting the effects which are likely to be produced upon an ulcerated surface by some of the different kinds of air.

In treating ulceration of the lungs, or even of any other part of the body, great respect is to be paid to the temperature as well as to the chemical properties of the air: cold air is in general more unfriendly to all, than a kindly warm air; and cold air is, in general, hurtful to the consumptive, as, besides its effects upon the internal surface, it will also determine the blood inwardly upon the lungs, and thus occasion more or less increase of the existing morbid affections. It is now well known, that atmospheric air is a compound, consisting, according to the more modern accounts, of oxygen (pure) air, one part; and azotic air, or air unfit for animal and vegetable life, and combustion, three parts: and that, according to the proportions of these different ingredients, the common air is more or less stimulating, and will be therefore

more

more adapted at one time than at another for favouring the recovery of ulcers exposed to its influence. As every experiment hitherto made with oxygen air, respecting the lungs, proves it to possess stimulating properties to so great a degree as to be incompatible with health even in their sound state. It will, therefore, be readily believed, that the constant application of any considerable proportion of this stimulating substance to their tender, irritable, and ulcerated internal surface, must be highly detrimental in attempting a cure. Therefore, in an atmosphere containing a super-abundant quantity of oxygen air, the ulcerated and tender surface will be irritated, and the patient's cough increased in consequence, as well as perhaps all the other existing pulmonary affections. But, notwithstanding that a certain quantity of oxygen air is so absolutely necessary for life that we cannot live but a very short time without it: yet, as a very small quantity of it will support life for some time, its proportion may, therefore, be lessened, if too stimulating, by reducing it with hydrogen air, which seems to be the best adapted for that purpose; as, besides the property of rendering the oxygen less stimulating, by its chemical combination, and decomposition,

the

the hydrogen air seems to possess sedative properties, which, together with the abstraction of the stimulant ones of the oxygen, will produce considerable good effects upon an irritable ulcerated surface, commonly exposed to an atmosphere superabounding with oxygen.

The ingenious Dr. Beddoes conceiving that consumption originated from a superabundant quantity of oxygen air in the blood, has recommended to inspire the common atmospheric air, mixed with a certain proportion of hydrogen air; which he imagines will neutralize the superabundant oxygen, and thus prove a valuable remedy in the cure of consumption. Whether his conjecture be, or be not, well founded remains yet to be determined. The sedative properties, however, of the hydrogen air above-mentioned seem to me to account for the good effects produced in the experiments made by this indefatigable physician; but which, respecting consumption, I imagine, are chiefly confined to the internal surfaces with which it comes in contact. For it is well known that hydrogen air is so highly sedative that, if increased to a certain degree, it will destroy life, in the same way as a too increased dose of cicuta*, and

* Vide Observations on sedatives in the Appendix.

other sedative substances of that class, viz. by gradually diminishing all the actions of the body, and at last destroying them, together with that of the heart itself. By this sedative property, therefore, do I conceive that a certain quantity of hydrogen air, mixed with an atmosphere abounding with a too-increased proportion of oxygen air, may prove useful in ulceration of the lungs. It may imitate, in a certain degree, that soft and congenial atmosphere so eagerly desired by the consumptive.

And as it is probable that, to an over-proportion of oxygen air existing in atmospheric air, is owing the irritation of wounds exposed to the air; so the hydrogen air, as combining with the oxygen, will thus deprive the atmospheric air of its irritating property: while the superabundant portion of hydrogen air will add sedative properties of infinite importance to an irritable sore. Dr. Ewart observes, “ and the
 “ accurate Mr. Watt assures me that this air
 “ (hydrogen) has a powerful effect in allaying
 “ the pain of external inflammation and sores,” which gives additional support to the idea of good effects being produced by it on the internal surface of the lungs, when ulcerated. While you thus attend to the chemical properties, it is
 also

also necessary to pay attention to the temperature, which, if not exactly adapted to the constitution and nature of the case, will more or less disagree with the patient. For it is well known, that many pulmonic affections are either relieved or cured by removing the patient from a cold and bleak situation into a warmer, although less pure. But, if the pneumonic affections arise from debility, as is often the case in asthma, a cool, keen air invigorates the whole pulmonary system, and often gives almost instantaneous relief.

On the above principles I have frequently relieved asthmatic patients, according to the various causes which produced their complaints, according to the season of the year, or according to the particular state of the vessels of the lungs of each patient, by sending some of them from Mary-le-Bone to Charing Cross or to the city, and others to Hampstead: and, in my election, I always attended as much to temperature as to chemical properties.

The inhalation of sedative vapour from decoctions or infusions of the cicuta, or white-poppy heads, I have, occasionally, used with good effects. And, when I have thought that an inactive state of the internal surface existed,

I have added to the liquor, the vapour of which was to be inhaled, a small quantity of distilled vinegar, in order to increase its activity. And, when the lungs have so far recovered themselves, that the general symptoms indicate rather an inactive and relaxed state of the whole pulmonary system, than an irritable or inflamed one; then an increased proportion of oxygen air, added to the usual atmosphere, will doubtless tend to the recovery of the patient, by invigorating the constitution in general, and the pulmonary system in particular. Whenever this plan is indicated, it may be a question, whether it will tend as much to the benefit of the consumptive patient to seek for it modified in nature's own way, viz. in the midst of luxuriant verdure and foliage, when the sun has properly performed that natural operation sufficiently demonstrated by the experiments of Dr. Ingen-Houfz; as when it is obtained from manganese, or nitre, &c. and mixed with atmospheric air in the way most approved by Dr. Beddoes, and the other gentlemen who have made this branch of medicine their particular study. The latter plan indeed must be adopted when the patient is unable to leave his chamber. As a certain proper mixture of oxygen, or pure air, is of
great

great importance to the consumptive in every stage of the disease, and as plants, whether in leaves, flowers, or fruit, have more or less influence upon the air as to its purity, it may not be improper to lay before the reader some observations on this subject, by the above-mentioned respectable philosopher, Dr. Ingen-Houfz. They therefore follow.

The Doctor, in his preface, page 64, says,
 “ * J’observai que les plantes n’avoient pas
 “ seulement la faculté de corriger l’air impur
 “ dans l’espace de six jours ou plus, comme les
 “ expériences de M. Priestley semblent l’indi-
 “ quer ; mais qu’elles s’acquittent de ce devoir
 “ important dans peu d’heures, de la manière la
 “ plus complete ; que cette opération marveil-
 “ leuse n’est aucunement due à la végétation,
 “ mais à l’influence de la lumière du soleil sur
 “ les plantes. Je trouvai que les plantes possè-
 “ dent en outre l’étonnante faculté de purifier
 “ l’air qu’elles contiennent dans leur substance,
 “ & qu’elles ont sans doute absorbé de l’atmos-
 “ phère, & de le changer en un air des plus
 “ purs, véritablement déphlogistiqué ; qu’elles
 “ versent une espèce de pluie abondante (s’il

* Vide Expériences sur les Végétaux, Vol. I.

“ est permis de s’exprimer ainsi) de cet air vital
 “ & dépuré, qui, en se répandant dans la masse
 “ de l’atmosphère, contribue réellement à en
 “ entretenir la salubrité, & à la rendre plus ca-
 “ pable d’entretenir la vie des animaux—qu’il
 “ s’en faut beaucoup que cette opération soit
 “ continuelle, mais qu’elle commence seulement
 “ quelque temps après que le soleil s’est élevé
 “ sur l’horizon, après qu’il a, par l’influence de
 “ sa lumière, éveillé les plantes engourdies pen-
 “ dant la nuit, & après qu’il les a préparées &
 “ rendues capables des reprendre leur opération
 “ salutaire sur l’air, & ainsi sur le règne animal:
 “ opération suspendue entièrement pendant l’ob-
 “ scurité de la nuit; que cette opération des
 “ plantes est plus ou moins vigoureuse, en rai-
 “ son de la clarté du jour, & de la situation de
 “ la plante plus ou moins à portée de recevoir
 “ l’influence directe du soleil; que les plantes
 “ ombragées par des bâtimens élevés ou par
 “ d’autres plantes, ne s’acquittent pas de ce de-
 “ voir, c’est-à-dire, n’améliorent pas l’air, mais,
 “ au contraire, exhalent un air mal-faisant &
 “ nuisible aux animaux qui le respirent, & ré-
 “ pandent un vrai poison dans l’air qui les en-
 “ vironne; que la production du bon air com-
 “ mence à languir vers la fin du jour, & cesse
 entièrement

“ entièrement au coucher du soleil ; mais qu’il
 “ faut en excepter un petit nombre de plantes,
 “ qui continuent leur action salutaire un peu
 “ plus long-temps que le reste ; que toutes les
 “ parties de la plante ne s’occupent pas de cet
 “ ouvrage, mais seulement les feuilles, les tiges
 “ & rameaux verts qui les supportent ; que les
 “ plantes âcres, puantes, & même les vénéne-
 “ uses, s’acquittent de ce devoir comme celles
 “ qui répandent l’odeur la plus suave, & qui
 “ sont les plus salutaires ; que la plupart des
 “ feuilles, sur-tout celles des arbres, versent cet
 “ air déphlogistiqué en plus grande abondance
 “ de leur surface inférieure ; que les feuilles
 “ nouvelles, & celles qui n’ont pas encore ac-
 “ quis tout leur accroissement, ne répandent
 “ point autant d’air déphlogistiqué, ni d’aussi
 “ bonne qualité, que celles qui sont parvenues
 “ à leur grandeur naturelle, ou déjà vieilles ;
 “ que quelques plantes préparent un air déphlo-
 “ gistiqué, d’une meilleure qualité que d’autres ;
 “ que quelques plantes, sur-tout parmi les aqua-
 “ tiques, excellent dans cette opération ; que
 “ toutes en général corrompent l’air environ-
 “ nant pendant la nuit, & même au milieu du
 “ jour, dans l’ombre ; que quelques plantes ce-
 “ pendant, qui ne cèdent à aucune autre dans

“ leur opération diurne à préparer l’air déphlo-
 “ gistique, surpassent néanmoins les autres dans
 “ leur pouvoir d’infecter l’air commun pendant
 “ la nuit & dans l’ombre, jusqu’au point même
 “ de rendre en peu d’heures une grande masse
 “ d’air tellement corrompue, qu’un animal plongé
 “ dans cet air y périt en quelques secondes ; que
 “ toutes les fleurs exhalent constamment un air
 “ mortel, & gâtent l’air environnant pendant le
 “ jour & pendant la nuit, à la lumière & à l’om-
 “ bre ; & qu’elles répandent un poison réel &
 “ des plus terribles dans une masse considérable
 “ d’air, où elles se trouvent enfermées ; que les
 “ racines récemment tirées de la terre ont la
 “ même influence mal-faisante sur l’air qui les
 “ environne, que les fleurs, à l’exception cepen-
 “ dant de quelques racines ; que les fruits en
 “ général conservent cette influence pernicieuse
 “ en tout temps, sur-tout dans l’obscurité, & que
 “ cette qualité vénéneuse des fruits est si grande,
 “ que quelques-uns, même des plus délicieux,
 “ tels que les pêches, peuvent, dans une seule
 “ nuit, rendre l’air tellement empoisonné, que
 “ nous serions en danger de périr, si nous cou-
 “ chions une seule nuit dans une petite chambre,
 “ dont la porte & les fenêtres seroient exacte-
 “ ment fermées, & où se trouveroit une grande
 “ quantité

“ quantité de ce fruit ; que le soleil, semble n’a-
 “ voir pas le pouvoir d’arrêter l’influence per-
 “ nicieuse des fleurs, est cependant capable de
 “ modérer les exhalaisons nuisibles de quelques-
 “ uns des fruits ; que le soleil lui même n’a pas
 “ le pouvoir de rendre l’air commun d’une meil-
 “ leure qualité, sans la concurrence des plantes ;
 “ mais qu’au contraire, il est plutôt capable de
 “ le corrompre, s’il agit seul, &c.”

The *third indication* of cure will be fulfilled by the proper exhibition and regulation of the different substances we have mentioned, when treating of the two former indications, whether applied internally to the stomach, or to the internal surface of the lungs by means of inhalation, in such a way as the proper knowledge of the anatomy, physiology and pathology of the lungs will point out. Indeed it will in general be found, that, in fulfilling the first and second indications, you also accomplish this at the same time.

For example, it will often happen, that medicines will promote a particular determination from the lungs, and, at the same time, lessen or increase the action of their blood vessels : for instance, the emetic tartar, and saline nauseating medicines will more or less determine to the surface of the body, while they will, at the same time,

ime, diminish the increased action of the sanguiferous system; and white wine, volatile and other medicines, while they determine outwardly, will, at the same time, promote a more vigorous action of the blood vessels. And a medicine which determines to the bowels, to the kidneys, or to the skin, may also increase the action of the absorbing vessels; as the magnesia vitriolata, jalap, calomel, antimonial powder, calomel and nitre, and a variety of other medicines which will readily occur to the attentive and well-informed practitioner.

We shall conclude our present remarks, by relating two cases of consumption; the one of the most deplorable kind, where the patient's sufferings were greatly alleviated and her life evidently prolonged by our method of treatment; and the other, where the patient was completely cured in the course of five weeks.

CASE

C A S E I.

Of Consumption, where the Lungs were so much diseased that the Patient had no Chance of Recovery, much relieved by a limited Use of Liquids, and the Life of the Patient thereby prolonged: together with some general Remarks.

Mrs. Todd, wife of — Todd, journeyman blacksmith, then living in Mary-le-Bone Street, and aged between thirty and forty years, applied to me on the 30th of May, 1793, after having been in a consumption about eighteen months.

May 30th. This poor woman looks a most deplorable object: she is wasted to a skeleton, with large drops of sweat standing on her forehead, and cannot raise or turn herself in her bed without assistance. She is now spitting blood and matter, and has done so for many months. She has considerable pain in the chest, attended with short breathing and incessant cough; and brings up about a pint of bloody expectoration in the twenty-four hours. She cannot lie upon either side. Her pulse is extremely weak, and beats about 130 in a minute:
and

and she has considerable thirst, and profuse colliquative sweats, both night and day.

Her bowels are disposed to constipation. She is generally thirsty, and drinks a considerable quantity of weak liquids every day. About six weeks ago she was able to walk out, when she was condemned as incurable by a very respectable Physician; who advised her to go home, and pray to God Almighty to take or relieve her, as medicine could afford her no farther relief.

A few days after this sentence was pronounced, she began to keep her bed, gradually getting worse until now; when, conceiving herself dying, she requested my assistance. From an attentive consideration of the history of the disease, and state of the patient, I was but too well convinced of the sagacity of the physician, and of the justness of his prognostic; and had, therefore, very little hopes of success. Yet, having no doubt but that the limited use of liquids would afford her, at least, some temporary comfort, I immediately set about the application of this principle; while I, at the same time, watched its effects with the most anxious and careful attention.

For

For the first three days, therefore, a due abstinence from liquids was enjoined, viz. she was allowed a pint of liquid only in the twenty-four hours, including tea, &c. and no medicine was given during that time which could interfere with the proper effects of this principle.

June 2d. She has adhered strictly to the limitation of liquids prescribed: and now looks cheerful and animated, and says that she is better, and that her perspiration is very much diminished. Her pulse is a little stronger, and also less frequent. Although, from this new trial, I was, more and more, convinced of the efficacy of my principle, yet I did not think myself warranted in trusting to it entirely; I, therefore, commenced the exhibition of every other medicine (together with a moderate use of liquids), which I thought was likely to forward the recovery of my patient.

And thus, by the moderate use of liquids, did I attempt to keep the lungs, as much as possible, at ease; while, by the diligent application of other means, I endeavoured to stop the existing morbid actions, and bring about healthy ones in their place.

As it would be too tedious to give the particulars of this case, and as they would, perhaps,
 answer

answer no good purpose, I shall only notice the more material circumstances which occurred during the treatment.

She took the flores sulphuris for some days, in doses of a drachm each, every four or six hours, and with evident advantage: for, besides retaining the body sufficiently open, without heating, or increasing the frequency of the pulse, it seemed to relieve the general pulmonary affections.

The cicuta was also of service, by allaying irritation and cough. But, having given the calx antimonii illota for several days, from an idea that, by removing some obstructions of the lungs, or fever, it might relieve the difficulty of breathing, and thus contribute to the benefit of the patient. I found I was mistaken; for the general weakness, as well as the fever and night sweats were thereby increased.

I advised a nourishing diet, and did not restrain her from a little animal food, when she was inclined for it.

June 14. Having become progressively better, her pulse is now tolerably firm and equal, and about 104. She has had no perspiration for the last three nights, and is now able to sit up in a chair. The expectoration is much bet-

ter,

ter, and in a diminished quantity, and she has one regular motion every day.

June 29. Her pulse is now only 92. She has little expectoration, little cough, and scarcely any perspiration.

For eight or ten days she continued, in every respect, so much better, that I began to entertain hopes of a recovery; and although her breathing was still short, I was inclined to impute part of that inconvenience to weakness. In the course of that time she was so well as to be able to go down stairs, with some assistance, and drink tea with the lodgers below. And after this she discharged her nurse, and was able to cook her husband's dinner, and do some other necessary offices in her apartment, for about a week: when feeling weak and exhausted, and becoming feverish, she complained of cough, short breathing, &c. And all these symptoms increased rapidly, notwithstanding every attempt to remove them, until the 18th, when she died.

She never had any colliquative diarrhœa, not even when she died: and as her strength was so much exhausted when I saw her, and as her bowels continued very regular, I did not venture to try that mode of determination.

Appearances

Appearances on Dissection. Having, on the 20th of July, opened the body, the following appearances presented themselves. In the abdomen every thing appeared cool and natural, nor were there scarcely any marks of putrefaction. The small intestines, in which there appeared no extrication of air, were much contracted and lying in a very small compass, but perfectly natural. In the pelvis was about half a pint of a clear brownish fluid; which was, perhaps, from transudation after death. All the other abdominal viscera were found: nor was there any inflammation on the peritoneum, or about the uterus.

Having next, after turning aside the integuments and external muscles, cut through the cartilaginous extremities of the ribs, I endeavoured to get into the chest as usual; but found the pericardium so closely connected with the sternum, that I could not raise the latter without cutting a small portion of the former, which appeared a thick, tough, leathery kind of substance. The sternum having been raised, a small vomica appeared, on the fore part of the left lung, opposite to the cartilaginous extremity of the third rib, containing about an ounce
of

of pure pus, which had no communication with any of the bronchial branches, and, therefore, could not be expectorated.

All the contents of the thorax appeared as if glued together; and adhered so firmly all round, that the sternum, ribs, intercostal muscles, pleura, diaphragm, and the contents of the thorax might have been justly considered as one great mass firmly united together. The heart seemed natural, but pale; and the right auricle and ventricle were full of blood: and in the pericardium was contained several ounces of the same kind of fluid as that found in the pelvis.

On examining the lungs, they appeared wholly diseased, excepting a small portion of the lower anterior edge of the right lung. And, on cutting into their substance, they were found full of small suppurations, some containing pus, others a cheesy matter; together with adhesion and obliteration of the greater number of the smaller blood vessels, air cells, and the smaller bronchial branches: the whole exhibiting a remarkable instance of what great disease may exist in the lungs, and yet life go on.

Now, as this poor woman had been confined to her bed for near six weeks before I saw her:

and as, when I did see her, she appeared to me in a dying state; and considering that she, upon the whole, gradually recovered from my first attendance until a few days before her death; I trust it will not appear improbable to suppose, that the whole of the morbid affections, which appeared on dissection, were actually present when I first saw her.—If so, the inference is obvious. For seeing that, in the above state of the lungs, even a fourth part of the quantity of blood, which was commonly circulated through them in health, could not now pass through them, on account of the great obstruction and obliteration of vessels which had taken place; I, therefore, conceive, that the only chance this poor woman had of having her life prolonged was, to withdraw her liquids; and thus, by diminishing the quantity of fluids, which, if taken into the body, must necessarily pass through the lungs, allow her to live until either the blood could find no passage through the remaining vessels, or her strength was unequal to the task of propelling it through the lungs. If, therefore, this patient got so well as to be able to do the necessary offices for herself and husband, although the lungs were at

that

that moment almost impervious to the circulating blood, can this be attributed to any other principle than that which was adopted and strictly adhered to, viz. the limited use of liquids? Respecting the above-mentioned useful medicines, and a variety of others which were given her, none of them produced any very permanent good effects; as, after a few days, I was, generally, on that account, obliged to try something new.

CASE II.

Of Pulmonary Consumption, speedily cured by our Method of Treatment.*

The patient, whose case I am about to relate, is a little man, of a dark complexion, sharp nose, high cheek bones, and about thirty years of age.

May 5th, 1793. He has had a severe cough for about six months, attended with considerable expectoration, short breathing, and pain in the side.

* Vide Dr. Duncan's Medical Commentaries for 1793.

During the last three or four weeks, he has brought up a considerable quantity of blood, and yellow expectoration; and the spitting is now purulent and bloody. His countenance is ghastly and desponding, being impressed with an idea (not generally entertained by patients of this description), that he shall not recover. He has now a pain in one side, violent night sweats, a dry furred tongue; is restless, and his pulse is hard and frequent. He lives several miles from London, where he has been attended by his own apothecary, who has bled and blistered him repeatedly, and used other means for his recovery, but without success. He has been in the habit of drinking many quarts of diluents every day. I ordered him a light, cooling, vegetable diet, and the following medicines; enjoining him particularly not to exceed a pint of liquid in the twenty-four hours, including tea, &c.

Rx. Extract. Cicut. ʒj. divide in Pil. xvij. quarum sumat ij. omni nocte.

Capiat etiam Haust. Cath. e Magnes. Vitriol. omni altero mane, non bibendo inter operationem.

May 12th. He has taken his medicines regularly, and observed the directions in regard to liquids.

liquids. His pills quieted the cough, which is now greatly better. He has had no night-sweats since he took his first draught, and has seen no blood for four days past. The expectoration is much diminished, and is now mixed with a frothy phlegm or mucus.

He eats his vegetable diet with pleasure, and sleeps well; his tongue is moist, with scarcely any fur upon it, and he is not thirsty. His opening draughts generally operated about three or four times. The pulse is much softer, and less frequent; and the pain of the side is gone.

The medicines and regimen to be continued.

May 19th. He has now, to my great astonishment, scarcely any complaint. He has no expectoration, no fever, and no cough: he sleeps well, and is acquiring flesh and strength. He sometimes feels his breathing a little short; yet he takes a deep inspiration without pain or coughing.

Capiat pilulas, ut antea; et haustum Cath. bis in hebdomada tantum.

I allowed him a little more freedom as to liquids, but still recommended moderation.

He was to return in a fortnight; but, being quite free from complaints, I only saw him

about a month after, when he was in perfect health, and is so at this moment, without having had any return of his pulmonary complaints.

In the above case, it is evident that great pneumonic affection existed, and that the patient was marching, with hasty strides, towards the other world. The lungs were overwhelmed with disease, while they were at the same time oppressed by the quantity of drink taken by the patient. Their morbid affections were thereby increased, while their healthy efforts were either lessened or prevented.

APPENDIX.

OBSERVATIONS ON SOME OF THE ARTICLES OF THE
MATERIA MEDICA.

CHAP. I.

*On the Rosa Rubra, Flores Chamæmeli and
Sarsaparilla.*

THE Rosa Rubra*, or Red Rose, has been allowed, both by the ancients and moderns, to possess considerable astringent powers; which no one, who has ever made the conserva rosarum and accidentally used an iron spatula, will dispute. Dr. Ruddy observes, "maxime adstringens est, præcipue in gemmis:" and, notwithstanding that every experiment which

* Rosa Gallica Linnæi.

can be made will prove this fact, no practitioner has yet pointed out how these astringent powers are to be obtained. Yet it will appear, on a moment's consideration, that the chief astringent properties of the rose can be obtained, for common use, in no other way than by decoction, although some of its astringency may be extracted by infusion. For it is well known that, wherever the virtues of a simple reside in its more fixed parts, decoction is preferable to infusion, for extracting them; and vice versa, when they reside in the more volatile parts. Now, as the chief virtues of the red rose are entirely confined to its more fixed parts, it must appear evident that decoction is the most proper mode of extracting them. And it may also be observed, that, when you give a preparation of this rose for stopping a bleeding vessel, you have no particular wish to preserve its fine volatile odour, which cannot, in the least, add to its astringent or tonic properties. Respecting the different effects of decoction and infusion in extricating the astringent virtues of the rose, the reader will be convinced by the following experiments:

After having made the *infusum rosæ*, as prescribed in the present *Pharmacopœia*, I boiled the
the

the same rose leaves in a double quantity of water to one half (which remainder precisely equalled the quantity produced by infusion), and I found that the decoction was not only as strong, but, by every appearance and trial, one half stronger than the infusion. So that the infusum rosæ extracts one part, while the decoctum extracts two parts and an half.

And, if you add to this the additional proportion of rose leaves in our decoctum, which is, as 48 to 180 of the Pharmacopœia*, this decoction will doubtless be considered as a medicine worthy of some attention: and, as such, we beg leave to recommend it. If to half a pint of this decoction you add six drachms of the mel rosæ, and a proportion of mineral or vegetable acid adapted to the state of the parts affected, you have a very effectual gargle for ulcerated sore throats; and one much preferable to any which can be prepared from the infusion or tincture.

What we have now said of the red rose is applicable to many other simples of the materia medica. As, for example, the decoctions of wormwood, and chamomile flowers, are, on

* Vide page 25.

the same principles, much more powerful stomachics than their infusions; as their stomachic properties reside in their fixed parts alone. The following decoction of chamomile I have found of great use in some affections of the stomach, after many other medicines had failed.

Decoctum Florum Chamæmeli*.

R. Florum Chamæmeli unciam unam,

Aquæ puræ, vel distillatæ, libram unam; coque ad libram dimidiam, et ab igne remotis adde

Pulveris crassi Radicis Zingiberi Albi drachmas duas: fiat infusio donec frigescat; deinde valide exprimens cola. Liquoris colati sumat æger cyathum vinosum, bis vel ter quotidie.

The *Sarsaparilla* † is another article of the materia medica, whose fixed parts contain all its medicinal properties; and which, therefore, can only be extracted by decoction.

It seems surprising that Dr. Cullen, although he allows that certain acrimony may exist in the blood and become the cause of disease, should, notwithstanding, contend that the sarsaparilla, which evidently contains a bland, insipid, mucilaginous property, which may be taken into

* *Anthemis Nobilis* Linnæi.

† *Smilax Sarsaparilla* Linnæi.

the blood vessels and sheath this acrimony, is good for nothing.

But, from its sensible properties, and from my own experience of its effects, I am convinced that the sarsaparilla possesses some bland, mucilaginous and sheathing properties, which, when properly extracted by decoction, may prove assisting in the cure of many obstinate diseases. That common or ammoniacal salt, may exist in the blood, in an increased quantity, as well as many other stimulating substances, such as sublimate mercury, &c. and thus give rise to many morbid affections, every one acquainted with the nature of these substances and of the animal œconomy must allow. And, that the decoction of sarsaparilla is a substance well adapted for being absorbed and carried into the blood vessels, and there sheathing every kind of acrimony or stimulating substance it may meet with, cannot be denied.

Many diaphoretic properties have been ascribed to the sarsaparilla: but this idea appears to have chiefly arisen from the mode of its administration. For, as the sarsaparilla has been more commonly given in the form of decoction, and that weak, and in considerable quantities; it may not be improbable to imagine that the
 diaphoretic

diaphoretic effects generally succeeding its exhibition proceeded more from the warm water in the decoction, than from any particular diaphoretic property naturally existing in the sarsaparilla.

Dr. Cullen* places the Sarsaparilla in his chapter on Stimulantia: but, would it not have been more natural to have placed it among the Demulcentia?

The London and Edinburgh colleges, attentive to the proper mode of extracting its virtues, have given very proper directions for obtaining them. I have, however, met with patients who could not take it in the form prescribed in sufficient quantity to produce any very beneficial effects, without having the stomach and whole constitution very much relaxed by the large draughts necessary to be taken. To obviate this inconvenience I have, for some time past, generally boiled the quantity of decoction which commonly made a quart down to a pint; of which the patient took a quarter of a pint, instead of half a pint, twice or three times a day. And, as it frequently happens that, together with the sarsaparilla, some tonic or

* Vide Mat. Med.

strengthening

strengthening medicine is also indicated, I have, on these occasions, commonly added some of the Peruvian bark, and prepared my decoction in the following manner; by which the virtues of the sarsaparilla appear to be wholly extracted, while a watery vapour only is lost by the long boiling.

Decoctum Sarsaparillæ cum Cortice Peruviano.

R. Radicis sarsaparillæ incisæ uncias tres,

Liquoritæ incisæ drachmas duas,

Pulveris Craffi Corticis Peruviani unciam dimidiam,

Aquæ puræ libras quatuor : decoque ad libram unam,
et cola.

Bibat æger liquoris colati uncias quatuor bis vel ter die.

During this demulcent and gently tonic plan, the patient's appetite and general health commonly mend; while all the secretions are performed regularly, and in proper quantity. I have seen this medicine, which, in general, rather strengthens than weakens the stomach, have very good effects in the leprosy, and other cutaneous affections. Was this from the decoction entering the blood vessels, and there sheathing acrimony so as to allow it to pass off by the different excretories; or were the good effects

effects entirely produced by the action of this medicine upon the muscular fibres of the stomach? I am inclined to believe, from both.

CHAP. II.

On the Cicuta, Stramonium, Hyosciamus and Aconitum; with some Remarks on Sedatives in general.

THE *Cicuta**, or *Hemlock*. Having for many years past entertained an idea that the good effects produced by the *cicuta*, in the cure of the different diseases for which it was administered, depended chiefly upon its sedative powers: and finding that the same is confirmed, not only by my own practice, but even by that of the celebrated Dr. Storck and others, I think it a duty incumbent on me to submit the result of my reflections and observations to public examination. And if, in so doing, I shall be

* *Conium Maculatum* Linnæi.

found to have assisted in establishing a fixed principle, by which practitioners may be directed in the exhibition of this remedy, I shall feel much satisfaction.

That the cicuta possesses *direct sedative properties*, and by them alone has cured, or assisted in the cure of, the various diseases for which it has been given, will, I think, appear evident from almost all the cases published on the subject, where the particular effects are properly related. And although the learned Dr. Storck, of Vienna, to whom the world is much indebted for the pains he bestowed in the investigation of this medicine, seems, in some parts of his cases, inclined to impute the good effects of the cicuta to some property of purifying the blood; yet, upon the whole, he seems to consider its action as insensible. That is to say, he could not account for it upon any fixed or visible principle. Therefore, when summing up the result of his experiments, he observes, "*Agit modo insensibili, nec alvum, nec vomitum, nec urinam, nec sudorem movet**." And notwithstanding, in the 3d Corollary to his *supplimentum necessarium*, he allows that some evacua-

* Vide Antonii Storck libellum primum de cicuta.

tions may take place by the skin and kidneys during its use, yet he there adds, "In plurimis tamen ægris nullam excretionem sensibiliter auget." The justly-admired and venerable Cullen says, "I am still at a loss to say what are truly the powers and virtues of this plant;" although he allows it capable of producing very considerable effects on the human constitution*. It appears to me wonderful that these two eminent physicians should not have been able to discriminate upon what principle it acted; although its direct sedative effects are particularly visible through the greatest number of Dr. Storck's cases, in his first, second and third publications, as I shall now endeavour to demonstrate. But shall first beg leave to state what is generally meant by a sedative. Dr. Cullen, in his chapter on *Sedantia*, says, "These are the medicines which directly, and without evacuation, diminish the motion and powers of the human system."

By this definition of sedatives the Dr. seems only to consider, as such, those sedative medicines which promote no particular evacuation. But may not evacuations of different kinds be

* Treatise of the Mat. Med. Vol. II.

by them produced merely from their operation as sedatives? I conceive there may: as, for example, by the skin, from relaxing a spasmodic constriction of the extreme vessels; or by the kidneys, from taking off constriction also of their secretory vessels. And in this way the cicuta may, every now and then, occasion a flow of thick urine, or a general perspiration; according to the particular state of the cuticular or renal vessels.

Notwithstanding, the cicuta more generally corresponds with Dr. Cullen's description; as will appear from the above quotations from the writings of Dr. Storck, as also from the writings of every other author on the same subject.

Now it is well known to practitioners, that, since the stimulating powers of opium and other substances, generally accounted sedative, but which only prove so indirectly, or after they have first produced some stimulating effects, have been particularly attended to, it has been a *desideratum* to find out a medicine which would immediately produce sedative effects, without occasioning any stimulating operation. And, indeed, some practitioners have contended that no such direct sedative existed in nature. But that the cicuta is one, I trust the reader will soon be convinced by the following remarks.

*Proofs of the direct sedative Powers of the Cicuta,
from Dr. Storck's Libellus Primus de Cicuta.*

In Case 4th, an account is given of a virgin, eighteen years of age, where, after severe affection of one breast, the disease at last terminates in an open cancer. The Dr. begins his treatment thus: "Hinc externe fomentum ex foliis cicutæ adplicui, et interne dedi mane & vesperi tres pilulas, quarum quælibet grana duo pondere habuit;" and immediately observes, "*Eodem adhuc die his remediis dolores multum imminuebantur.*" Case 15th exhibits a dreadful case of cancer of the scrotum, and of fungous excrescencies on the penis, also become cancerous, which originated from a venereal cause, and where that disease still existed, nothing having been taken to counteract or remove it. After describing the miserable state of the patient, that "nec quiete jacere, nec præ dolore dormire, minus autem ambulare potuit," he says, "In principio statim ter de die sex pilulas (a gr. ij) præbui, & partes affectas fomento diligentissime feci foveri. *Eadem vespera remiserunt dolores, & miser sponte cepit dormire.*" Are not direct sedative powers

powers evidently demonstrated by the two foregoing quotations? In the last both cancerous and venereal irritation are immediately counteracted by this medicine.

Case 16th, is a remarkable one of many fistulous sinuses, where the patient, exhausted with pain and a long continuance of the disease, and after the administration of many remedies without effect, was at last cured by the cicuta internally and externally administered. The Dr. observes, "*Primo statim die dolores mitigabantur, & ægra sine opio dormivit, quod ante nunquam factum erat.*" Were not the morbid actions from this moment diminished by the sedative effects of the cicuta, and the patient gradually restored by a return of healthy ones?

Other Proofs from Dr. Storck's Libellus Secundus de Cicuta.

By Case 2d, are clearly demonstrated the sedative powers of the cicuta, producing direct debility by continued application. In it the learned Baron gives an account of a great disease of the breast, in a woman, thirty-six years of age, which is at first large, hard, and painful;

ful; and then becomes open, discharging ichorous matter from an ill-conditioned and troublesome ulceration. A cure is at last accomplished, chiefly by the cicuta and the Peruvian bark. Now, if the reader will particularly attend to the effects of these medicines on the patient, during the treatment, he will perceive them always sedative or tonic according to the mode of their administration. For when the cicuta is continued by itself for any length of time, or when the dose is much increased, even when accompanied with a little bark, sedative and debilitating effects soon become apparent. In short, the whole history exhibits a succession of sedative effects producing debility, shivering, &c. which are always removed by the increased administration of the bark, and the diminution or omission of the cicuta. In one part of this case, after having continued the cicuta alone for twenty-eight days, in considerable doses, he says, "*Verum ægræ vires prosternebantur, & nova iterum febricula orta est, et sæpius advertentebantur horripilationes:*" and adds, "*Hinc dedi iterum corticis Peruviani nuciam dimidiam per diem.*" And he diminished the quantity of the cicuta from a drachm and an half to a scruple in the day. It will be also observable, that, when
the

the sedative power of the cicuta is applied so long as to produce debility, then the discharge becomes more ichorous, and every thing assumes a more unfavourable aspect; until, by withdrawing the different causes of debility and giving some tonic medicines, the strength is again restored.

Case 3d. Here a man, twenty-seven years of age, afflicted with a large, foul, spreading cancerous ulcer of the side, which had, for a long time, resisted many different modes of treatment, and even the bark itself, although given in large doses and for a considerable time, was at last cured by the inward and outward application of the cicuta.

In this case the sedative effects of the medicine soon became apparent, by diminishing the hectic fever of irritation, and thus procuring quiet sleep and a diminution of all the other unfavourable appearances of the disease. Now, as it may sometimes happen that, in order to cure ulceration, nothing is wanting but to diminish the irritable action of the absorbing vessels; it cannot, therefore, appear improbable that the cicuta, by its sedative powers, should accomplish this desirable purpose, and thus allow a cure to take place without the assistance

of the bark. This sedative power is even evident by the external application alone ; for by only bathing an irritable cancerous sore with an infusion of the cicuta the irritability is immediately diminished, and the thin fœtid ichor is gradually changed into a discharge of a more favourable kind.

Case 7th produces an example of the sedative powers of the cicuta when thrown even into the cavity of the chest. It was in the dreadful case of a virgin, of twenty-one years of age, where, from a severe and tedious cancerous affection, the ribs were become carious, and an opening made into the cavity of the chest, through which fœtid ichor was discharged. A variety of injections were thrown into the cavity of the thorax in order to cleanse it, many of which the patient could not bear : that of barley water and honey was the easiest, but the fœtid ichor still continued to be discharged, attended with heat and pain. An injection of the infusion of cicuta was therefore adopted : upon which the matter became almost immediately less fœtid, and the heat and pain in some degree decreased. His words are, “ *Altero die jam materies ex*
“ *cavo pectoris prodiens non amplius adeo fœ-*
“ *tuit, et dolor, ardorque paulisper imminuti*
“ *sunt.*”

“ sunt.” And a little farther he adds, “ Lene
 “ ab initio fiebat infusum cicutæ pro injectione,
 “ quoniam autem vidimus id nec pulmones irri-
 “ tare, nec augere anxietates, & bonum præstare
 “ effectum, id saturatum dein adhibuimus.” Un-
 der the same title of casus 7mus. is related an
 account of a man, affected with empyema, where
 the operation was performed, the matter was
 discharged, and the ribs found carious. Here,
 after mentioning that the surgeon had tried many
 injections, he adds, “ Nullam autem magis uti-
 “ lem invenit, quam eam, quæ infuso cicutæ facta
 “ fuit, hæc enim pus laudabile fecit.” And a
 little farther, “ Mirum! quod saturatum infu-
 “ sum cicutæ in cavum pectoris, et ad nudum
 “ pulmonem injectum nullas turbas, nullas anx-
 “ ietates, &c. excitaverit.”

Now had a solution of opium been injected,
 much pain would have been at first produced, as
 is the case when applied to the eye or to any
 tender surface.

In this second publication of Dr. Storck there
 are many of his medical correspondents who
 confirm his good opinion of the cicuta. One
 of them, Mr. Ferdinand Leber, writes thus,
 “ Fœmina, 34 annorum, post terrorem habuit
 “ mammam induratum, dolentem, moles duritiei

“semper increvit, & aucti sunt dolores, Data
 “cicuta mox melius se habuit, & intra binos &
 “dimidium menses fuit penitus sanata”—which
 I trust will be allowed as another instance of the
 sedative powers of the cicuta. Here the ir-
 ritability of the part was so much increased that
 it was about to break out into an open cancer,
 but was happily prevented by the sedative
 powers just mentioned. In the first corol-
 lary of this book Dr. Storck, when speak-
 ing of the effects of the cicuta upon the blood,
 observes that the blood of many people, who
 had taken this plant in large doses and for a
 long time, was of proper colour and consistence:
 and says that he and Dr. Restler found the blood
 of a lady, in the last stage of pregnancy, and
 who had taken cicuta during several months of
 her time, resembling that of persons in good
 health. Here he does not even mention that
 the buff coat (commonly so called), which is
 generally seen in the latter months of pregnancy,
 appeared on this occasion. In the same corol-
 lary the Doctor adds another fact, which is of
 great importance towards establishing the direct
 sedative principle I have been endeavouring to
 demonstrate. His words are, “Dum hæc scribo,
 “sanguinem, iterum mihi alii comiti, quæ cicuta
 “a binis

“ a binis mensibus utitur, & video eundem op-
 “ time ligatum, & longe saniozem, quam fuit
 “ ante usum cicutæ, tunc enim semper sanguis
 “ erat tenaci, mucosa, & *variegata crusta tectus,*
 “ fibrosus.” Doth it not appear evident from
 this quotation that the cicuta, instead of increas-
 ing the action of the heart and arteries, and thus
 also the phlogistic diathesis of the system, rather
 tends to diminish both? It certainly does: and
 I have often used it for this purpose in the acute
 rheumatism, and with the greatest success *.

So

* The salutary effects of the cicuta in the very last stage
 of pregnancy is also confirmed by Dr. Butter, to whom the
 publick is much obliged for the trouble he has taken in re-
 commending this remedy in the kinkcough; the efficacy of
 which in that disease I have often experienced. The Doctor's
 judicious rules for making its different preparations are well
 worthy of being perused by all who wish to have them in
 perfection, particularly the extract †. In making the latter
 preparation I particularly adhere to the Doctor's directions;
 only that, as I imagine that much of the sedative effects of the
 cicuta depend upon its volatile parts, I, therefore, endeavour,
 as much as possible, to preserve them: and, for this purpose,
 while the evaporation and coagulation are going on, I care-
 fully collect the coagulated portions, or clots, into a pot
 until the end of the process; when I mix the whole together,
 and reduce them to the consistence of an extract. And if I
 wish to make the extract according to the Edinburgh Pharma-

† Vide a Treatise on the Kinkcough.

So far do the two first publications of Dr. Storck seem to confirm the idea of the direct sedative powers of the cicuta. I shall beg leave to mention another proof from his *Supplimentum Necessarium*, or 3d publication, de cicuta.

In case 9th of this supplement is related an account of a cancer of the nose and upper lip, with which the patient was affected for two years. The parts were livid, hard, swelled, and so painful that no sleep could be procured. After having used purgatives, antiscorbutics, mercurials and purifiers of the blood, from which the disease seemed rather to increase, he begins the use of the cicuta; and with the happiest effects. His words are, “*Adhibita cicuta ichor copiosissimus, sed longe blandior, effluxit, mox partes omnes detumuerunt, dolor evanuit, lividus color mutatus est in naturalem.*”

Further proofs of the powers of the cicuta from Dr. Andree†, with some other remarks on the different sedatives.

copœia, which I have always found to be the most effectual, I mix the powder at the same time; which renders less evaporation necessary.

† Vide Observations upon a treatise on the virtues of Hemlock, &c. by John Andree, M. D.

Having

Having above endeavoured to point out the sedative property of the cicuta from the writings of Dr. Storck, I shall now proceed to give some extracts from his opponent, Dr. Andree, in further confirmation of this idea. Dr. Andree, in his observations upon Dr. Storck's treatise, labours to refute the Doctor's ideas respecting the innocence of this remedy, and very properly cautions against trusting entirely to it in the cure of cancer, and the other diseases in which Dr. Storck recommends it. But, leaving their dispute to themselves, I shall only adduce a few of the facts which tend to establish the sedative powers of the cicuta.

Dr. Andree, page 6th, mentions the case of Elizabeth Web, where, after taking " six grains
 " of the extract night and morning, for two or
 " three days, she was obliged to leave it off;
 " for it brought on such dizziness of the head,
 " and dimness of the eyes, that she thought she
 " should lose her sight. Her speech was also
 " affected, and she felt a numbness in her limbs,
 " especially the arms and hands; so that she
 " was afraid she should lose the use of them."
 Are not these the effects of a direct sedative power applied so as to bring on direct debility? I believe so: and think that, if it had been
 given

given in a smaller dose, these symptoms of debility would not have appeared.

At page 8th is given the case of a Lady, of a bad habit of body, who also took the cicuta for a cancer. She took two pills every day. "After the second dose," says he, "she was taken with a dizziness in the head, and sickness. On taking the fourth dose, she became paralytic all over, lost her speech, and for several days seemed to be dying. By the assistance of cardiac, &c. medicines, she recovered from this dangerous situation." This is still a more striking case of debility induced by the application of a sedative power, and which is removed by the use of cordial medicines. Several other cases of the same kind are related by Dr. Andree: who, afterwards, observes, respecting the action of the cicuta, that "It seems principally to attack the primæ viæ, and debilitate the nervous system."

Upon the whole, it appears that he considers it as either dangerous or useless.

But as the virtues of all the articles of the materia medica are only relative, and as they become either poisonous, medicinal, or good for nothing, according to the mode of exhibition; so also may the cicuta be given so as to
be

be either salutary, poisonous or of none effect. For by the same sedative property by which it allays irritation and pain, and is capable of suspending the different morbid actions, may it also suspend all the actions of the body, and even that of the heart itself. This effect of the cicuta has not been denied by the modern physicians; and is particularly acknowledged by the ancients, who were therefore very fearful of using it as a medicine.

Indeed, in greatly debilitated constitutions, some caution is necessary in proportioning the dose to the strength of the patient's constitution. For I once saw an unfortunate case, which intimated to me this kind of circumspection. It was in that of a patient who had been long afflicted with chronic rheumatism, by which she was considerably weakened; where, from taking some pills of the extractum cicutæ, in a quantity beyond that ordered by the gentleman who attended her, and, I presume, in a quantity beyond the strength of her vital powers, all the actions of the body were gradually diminished, and at last entirely suspended; and the patient thus delivered, quietly and without a groan, into the arms of death. Nor, in this case, would it have been any good objection to the sedative power

of

of the cicuta, if the *vis medicatrix naturæ* should have been roused, and, in its last struggles for the defence of the constitution, produced some convulsive efforts.

Now we have above seen, that the cicuta, when given in a moderate dose, or in one properly adapted to the strength of the patient and nature of his disease, produces the most beneficial sedative effects: as also, that, when carried a little farther, debility ensues; and that, in a still increased proportion, life at last may be in danger. But that the cicuta is perfectly innocent in proper hands, is very well ascertained; and, that its effects arise chiefly or entirely from its sedative powers, I trust, have now been sufficiently demonstrated. When, therefore, a fever of irritation exists, whether it arises from a tubercle about to inflame and suppurate, from a scirrhous tumour about to break open or form a cancerous or other irritable ulcer, or even when cancer is formed; by its sedative powers it allays the irritation, quiets pain, and diminishes the frequency of the pulse; and thus allows the healthy operations, when properly directed towards the removal of the disease, to proceed without interruption. In cases where several pieces of carious bone were to be removed, the

cicuta

cicuta has been, in the same way, found of service; and, by diminishing the irritability of the neighbouring parts, it has allowed them to come away without much pain or irritation. Seeing, therefore, that direct sedative powers exist, and are the only demonstrable properties, in the cicuta, we ought not to wonder if, after having allayed all the symptoms of irritation, the sedative powers can extend no farther for the benefit of the patient*. And, as the taking off of irritation is not always the only treatment necessary for the cure of scirrhus or cancer, so the practitioner, who, in those cases, trusts entirely to the cicuta, will be often disappointed. Indeed we see that Dr. Storck, and also his correspondents, every now and then, used other remedies together with the cicuta; for example, in case 18th of his *Libellus 2dus*, he ordered a purgative of rhubarb and sal polychrest to be taken every fortnight, together with the cicuta; and the patient recovered perfectly. And the learned Dr. Kaifin, in a Letter to Dr. Storck, says, “Etenim fere omni septimana dabam meis ægris purgans, & eis conducebat.” Might not these

* *—* sunt certi denique fines,
Quos ultra citraque nequit consistere rectum.

Hor. Sat. 1. lib. 11.

purgatives,

purgatives, or some invigorating or different air, or some other application to the system, such as good news, love, an extraordinary glass, &c. bring about new and healthy actions in the parts affected, while the morbid actions or dispositions to inflammation were either lessened or suspended by the cicuta?

In considering sedatives according to their effects, may they not be naturally divided into two kinds, viz. *Direct* and *Indirect*?

By a *Direct Sedative*, I mean a medicine which operates more or less immediately as a sedative, without producing any stimulating effects; such as the cicuta, hydrogen air, and perhaps many other substances.

By an *Indirect Sedative*, I mean a medicine which, although it ultimately produces sedative effects, yet has some other previous stimulating operation; such as opium, paregoric elixir, &c.

Now, although opium has generally been ranked as the chief of the sedantia, yet its stimulating power is at present very well ascertained: and every practitioner knows (what we have already mentioned), that, if applied to the eye, or to a tender surface, it will produce more or less of irritation and pain, whatever ultimate sedative effects it may occasion. From this stimulating

mulating property, which is always more or less discoverable on its first exhibition, are we prevented from employing it in cases of strong active inflammation? for, in them, if opium is given alone, the phlogistic diathesis of the system is in general thereby increased, and the disease thus rendered more difficult of cure. But it is not so with the direct sedative we have been treating of: for in the most acute rheumatism, or in the most violent pleurisy, it may be given with advantage; and, instead of increasing, it will rather help to diminish, the phlogistic diathesis*. Nor will the cicuta, if applied to the most irritable surface, or (as we have seen above) if even thrown into the cavity of the chest itself, produce pain or irritation: on the contrary, it will immediately relieve both, and in so doing will clearly demonstrate its direct sedative powers. Although this fact is so clearly proved by the writings of Dr. Storck, and others, as to be beyond the possibility of doubt, it seems strange that no practitioner has yet pointed it out so as to be kept in view in general practice.

* *Cicuta nec majorem motum, nec in sanguinis circulo turbam excitat.*

Supplim. necess. de cicuta, corol. 2dum.

The reader has already seen the candid acknowledgement of Dr. Cullen: and he must also know, that no particular principle has yet been settled among practitioners respecting its administration. For, while one condemns it as either dangerous or useless; another gives it in all cases, whatever they may be, or in whatever state they may exist, provided they happen in a scrophulous constitution. And even the celebrated Baron Storck, notwithstanding his great success in the treatment of many scrophulous, cancerous, and other complaints, by this remedy, was still at a loss to account for its mode of operation; and had, therefore, no particular principle to direct him*.

From what hath been said, may we not give the following definition of the medicinal properties of the *cicuta*?

The *cicuta*† is an innocent, but powerful, remedy;

* Being of opinion that the foregoing reasoning, on the facts related by Dr. Storck and Dr. Andree, sufficiently establishes the direct sedative property of the *cicuta*, I shall, for the present, omit some particular facts of my own, as being unnecessary, and as they are in general connected with other subjects which are intended for publication.

† It may be necessary to inform the reader that the preparations, from this herb, without the root, are those meant

medy; possessing sedative properties, by which it is capable of diminishing the morbid irritability of the body, from whatever cause it may arise, without producing any stimulating operation: from which, therefore, it may either lessen, or entirely suspend, all diseased actions which arise from, or depend upon, morbid irritability.

A few Observations on Dr. Storck's Treatise on the Stramonium, Hyosciamus, and Aconitum.*

I have already made mention of three publications, by Dr. Storck, on the cicuta: soon after these he published another on the stramonium, hyosciamus, and aconitum; which seem to possess sedative properties, more or less resembling those of the cicuta.

Of each of these the Doctor makes several trials, chiefly to prove that these simples, which had hitherto been accounted highly poisonous, might be given with safety in the cure of disease. After a variety of experiments on himself and others, he proceeds to exhibit them in every obstinate disease, or where no other remedy proved in this definition, viz. the infusion, decoction, extract, and powder.

* Vide Libellum de Stramonio, Hyosciamo et Aconito.

of service; and often with success, although he had no fixed principle to guide him.

*Stramonium**, or *Thorn-apple*. After mentioning that authors had generally spoken of this plant as highly hurtful to man and brutes, viz. “*stramonium turbare mentem, adferre insaniam, delere ideas et memoriam, producere convulsiones;*” he prepares an extract from it, and with it he cautiously tries some experiments respecting its effects on the human body. He put upon his tongue a grain and an half of the extract, from which he found no particular inconvenience or affection; and, having dissolved it in the mouth, he only perceived a disagreeable, nauseous taste; but does not mention any stimulating effects, such as redness, inflammation or pain. He swallowed it without observing any particular effect. Therefore he, afterwards, ventures to give it in several cases of mania and epilepsy; in some of which it seemed to produce good effects. But, as the cases are related, no decided opinion can be formed whether or not this plant possesses direct sedative powers, similar to the *cicuta*, although the Doctor’s experiment upon himself seems to say so. It may therefore deserve farther trial.

* *Datura*, Linn. Sp. Plant.

*Hyoscyamus**, or *Henbane*. Having prepared an extract of the leaves and stalks of this plant, he tries it first on a dog and then on himself; and being, thereby, convinced that a moderate dose might be given with safety, he goes on to try it on his patients.

Experiment 1st. A woman, thirty-seven years of age, had been afflicted with dreadful convulsions, for above a year and an half, every day. Many medicines had been given without effect: and she only received a temporary relief from opium, given in large doses. She was at last cured by the extractum hyosciami, which was first given in the quantity of a grain, three times a day, and gradually increased to nine grains a day. The convulsions soon disappeared: and the body was kept open by the medicine.

Experiment 2d. A convulsive tremor of the foot was cured by this remedy, in three weeks. The patient had copious stools.

Experiment 5th. A man, about thirty years of age, affected with mania, having tried blood-letting, purging and other medicines without effect, took also paregorics and opiates, in large

* *Hyoscyamus niger*, Linnæi Syst. Veg.

doses; but, instead of procuring sleep, he says, “noctes inde magis inquietæ reddebantur, mens fiebat anxia, oriebaturque febris.” He therefore tried the extract of the hyosciamus, in the dose of one grain three times a day, and adds, “nox prima jam magis tranquilla fuit.” And by increasing the dose, the rigors, delirium, &c. entirely left him, and the body became more open.

Experiment 6th. A man, thirty-three years of age, afflicted with a severe tickling cough, and expectoration a little streaked with blood, together with anxious nights, and having tried many things in vain, took one grain of the extractum hyosciami in the morning, and one in the evening. The Dr. observes, “Nox fuit longe tranquillior, & æger sibi opium datum esse, putavit.” The dose was gradually increased to nine grains a day, viz. three in the morning, at noon, and in the evening: and all the morbid affections gradually disappeared. He had generally three or four stools a day.

Experiment 9th. A woman, thirty years of age, from having been accused of theft, became disturbed in her mind, and so violent as to be obliged to be bound with cords. From the

11th to the 18th of September every thing which the art of physic could suggest was tried without any advantage; and two grains of opium having been given, they only procured her a short sleep, from which she awoke worse than ever. The extractum hyosciami was then given; of which she took two pills every day for three days, and afterwards three; and left the hospital, about the end of October, perfectly restored. The only effects, mentioned to have been produced by the hyosciamus, were evidently sedative, viz. “ægra pacatior fieri videbatur,” and “his (pilulis) sensim rediit quies menti.”

Case 10th. A woman, eighteen years of age, is cured of the epilepsy by the hyosciamus. The greatest dose was two pills (each a grain) three times a day. Here purgatives were occasionally thrown in, as she was costive.

Cases 11th and 12th. Are cases of fits, cured by this remedy.

Upon the whole, it appears sedative with the power of keeping the body open.

*Aconitum**, or *Wolfsbane*.

Dr. Storck having powdered the leaves and stalks of this plant, which had been reckoned amongst the strongest poisons, he put a little of the powder upon his tongue, and immediately felt a heat, which continued for some time; together with some momentary, vague and lancinating pains, which often pervaded the tongue, but produced no unpleasant consequence: and the powder, being left on the tongue for two minutes, neither brought on inflammation nor redness. As long as the heat remained in the tongue, so long he had a copious flow of saliva; but in other respects he perceived no kind of inconvenience. Doth not the absence of redness and inflammation demonstrate the absence of stimulus; and therefore prove, that the aconitum has no stimulating properties? And may not the heat, momentary pains, &c. be referred to sensation, which may not necessarily imply increased action or stimulating effects? The above powder the Dr. also sprinkled upon a cancerous, fungous sore; and he observes,

* *Aconitum Napellus*, Linn. S. Veg.

“ Primo

“ Primo die levis oriebatur suppuratio, et æger nullum *dolorem*, nec *ardorem* conquestus est, secundo, 3tio. quarto & 5to. die idem observatum fuit.” Is not this a farther proof of direct sedative powers? Having prepared an extract in the usual manner, and put some upon the tongue, it only caused a very slight titillation. And having put a grain within the lower eye-lid, he observed no particular burning heat, but only the usual effects as from any other heterogeneous body*. He then proceeds cautiously to administer it inwardly; first to himself, and then to others. The first effect he observed was, that “ *extrema totumque corpus præter consuetudinem toto die multum transpirare, madere;*” which was constantly the case while he took the medicine: but, intermitting it for one day, these effects by the skin disappeared, although they returned immediately on returning to the aconitum. Whether the action of the sanguiferous system was thereby increased, we are not informed; although, from the experiment on himself, we may conclude it

* “ *Nec inde afficiebar aliter ac a quocunque corpore heterogeneo.*”

* Aconitum Napellus, Linn. S. Veg.

was

was not, as he says, "Nulla inde actio corporis turbabatur."

After these experiments on himself, and many upon others, he concludes, that the extract of aconitum is an innocent and very efficacious medicine. And adds, "Acre, quod circa articulos, tendines, & ossa hæret, irritat nervos, excitatque summos dolores, solvitur inde & agitur in motum, & per urinam, vel alvi fluxum, vel per sudorem, vel per insensibilem transpirationem ex corpore ejicitur." And, a little farther, he observes that sometimes the aconitum cures when the cicuta either disagrees or fails. Notwithstanding, he immediately acknowledges that the cicuta often relieves and cures after the aconitum has failed. "Hinc," he says, "de cicuta semper verum manet: esse eam medicamentum summe efficax in morbis curatu difficillimis."

Now, having clearly pointed out that the stramonium, the hyosciamus and aconitum, possess certain sedative powers, resembling those of the cicuta; I leave for the determination of the learned physician, whether the evacuations above-mentioned, whether the purgative effect of the hyosciamus, or the sudorific of the aconitum,

nitum, are to be ascribed to sedative or stimulating powers. But, that neither the stramonium, the hyosciamus, nor the aconitum possess the stimulating properties of opium, is evidently demonstrated as above.

THE END.

INDEX.

INDEX.

Page

A

ABSORBING Vessels, their two distinct actions, 127
 ————, their irritable action, perhaps the
 cause of ulceration, ———— *ibid.* and 127
 ————, conjectures and queries respecting
ibid.
 Aconitum, its properties considered, 216
 Aerial System has no communication with the connecting cel-
 lular membrane, ————
 Air Atmospheric, its component parts, 128
 — Hydrogen, or inflammable, its use, ————
 ————, is a direct sedative, ———— *ibid.*
 — Oxygen, or pure, is highly stimulating, ————
 ————, irritates the ulcerated internal surface of the
 Lungs, ———— *ibid.*
 ————, its stimulating power destroyed by combination
 with Hydrogen ———— *ibid.*
 ————, is obtained, naturally, from vegetables, 128
 ————, artificially, from manganese, nitre,
ibid.
 &c. ————
 Anatomy and Physiology of the Lungs, a general view, 1
 Anodynes, observations on, ———— 132
 Astringens

I N D E X.

A.	Page
A BSORBING Vessels, their two different actions,	127
—————, their irritable action, perhaps the cause of ulceration, — — — <i>ibid.</i> and	197
—————, conjectures and queries respecting, <i>ibid.</i>	
Aconitum, its properties considered, — — —	216
Aërial System has no communication with the connecting cel- lular membrane, — — —	4
Air Atmospheric, its component parts, — — —	158
— Hydrogen, or Inflammable, its use, — — —	161
—————, is a direct sedative, — — —	<i>ibid.</i>
— Oxygen, or pure, is highly stimulating, — — —	163
—————, irritates the ulcerated internal surface of the Lungs, — — —	<i>ibid.</i>
—————, its stimulating power destroyed by combination with Hydrogen — — —	<i>ibid.</i>
—————, is obtained, naturally, from vegetables,	165
—————, artificially, from manganese, nitre, &c. — — —	<i>ibid.</i>
Anatomy and Physiology of the Lungs, a general view	1
Anodynes, observations on, — — —	135

Aretæus recommends an astringent diet in hæmorrhages,	36
———'s practice confirms the utility of moderate drinking in diseases of the lungs,	145
Asthma, observations, and practical remarks on,	49
———, the proximate cause is often over-distention of the blood vessels of the lungs,	59
———, difficulty of breathing in, more commonly from compression than from constriction of the bronchia, <i>ibid.</i>	
———, our doctrines confirmed by the practice of Sir John Floyer,	62, & seq.
———, and by the writings of Dr. Cullen,	65
———, Cases where the limited use of liquids was used with advantage,	51
———, the cure,	66
———, a natural cure frequently produced,	70
———, a spare use of liquids particularly recommended,	72
———, great analogy between it and the broken-wind in horses, &c.	82

B.

Broken-wind in Horses, an improper name,	77
———, a disease analogous to the asthma, and, like it, seems to arise from over-distended blood vessels of the lungs,	77, and 82
———, the cure,	83
———, their pulmonary blood vessels found particularly full of blood after death,	82

C.

Catarrh,	89
———, limitation of liquids recommended,	94
	Cicuta,

Cicuta, or Hemlock,	—	—	190
——, is a direct sedative,	—	—	191, 207
Circulation of the blood through the lungs,	—	—	7
Climate, the best for consumptive patients,	—	—	158
Croup,	—	—	102
——, an active inflammatory disease,	—	—	108
——, heating antispasmodics improper,	—	—	<i>ibid.</i>
Cullen, Dr. his practice in hæmoptysis objected to,	—	—	37
——, Remarks on his definition of sedatives,	—	—	192

D.

Decoctum Rosarum, recommended in passive hæmorrhage,	—	—	27
—— Sarfaparillæ cum cortice Peruviano,	—	—	189
——, recommended in leprosy and other cutaneous diseases,	—	—	<i>ibid.</i>

E.

Emetics, their use in the prevention and cure of tubercle,	—	—	139
——, different kinds, and effects,	—	—	132
——, exhibition in ulceration considered,	—	—	146
——, recommended in consumption,	—	—	152

F.

Flores Chamæmeli, decoction preferable to infusion as a stomachic,	—	—	186
—— Sulphuris, useful in consumption,	—	—	153

H.

Hæmorrhage, passive,	—	—	27
——, active pulmonary, general observations and Remarks,	—	—	12
			Hæmorrhage,

Hæmorrhage, treatment of,	— — —	15
————, cases, with remarks,	— — —	29
Hyofciamus, or Henbane, some observations on,		213

I.

Ingen-Houfz, Dr. his observations on vegetables, respecting their effects in spoiling or purifying the air,	—	165
---	---	-----

L.

Lungs, general anatomy and physiology,		1 & seq.
————, general pathology, and preliminary observations,		9
————, Hæmorrhage of, general observations and practical remarks on,	— — —	12
————, treatment of,	— — —	15
————, wounds of	— — —	24
Liquids, the limited use of, recommended in active hæmorrhage of the lungs,	— 20, 29, 33, and 39	
————, recommended in passive hæmorrhage of,		28
————, ————— in the asthma,	— 51, and 72	
————, particularly useful in the short-wind of horses,		81
————, recommended in catarrh,	—	94
————, ————— in the cure of tubercle,	—	133
————, ————— in the ulcerative consumption,	144, 149	
————, ————— in menorrhagia, by Dr. Rowley,		36
Aretæus's practice confirms its utility,	—	145

P.

Physiology and anatomy of the lungs,	—	1
Pulmonary Consumption, remarks on,	—	47, 84
		Pulmonary

Pulmonary Consumption, treatment of,	—	139
—————, the pre-disposition to, considered,		140
—————, indications of cure,	—	142
—————, suspended by pregnancy,		158
—————, a very deplorable case greatly relieved by a limited use of liquids,	—	171
—————, a case cured by the same,		179

R.

Rosa Rubra, or Red Rose, possesses considerable astringent properties,	—	183
—————, decoction preferable to infusion, for extracting them,	—	184
Rowley, Dr. recommends abstinence from liquids in menorrhagia,	—	36

S.

Sarsaparilla, some observations on,	—	187
—————, possesses demulcent properties,	—	<i>ibid.</i>
—————, decoction with Peruvian bark, good in the leprosy and other cutaneous diseases,	—	189
Sedatives in general, remarks on,	—	136, 193, 208
—————, direct and indirect,	—	208
Short-wind in horses,	—	77
—————, moderate drinking of great use,	81, 2, 3	
Sims, Dr. James, recommends sulphur in consumption,		153
—————, tried with good effects,		153 and 174
Strength of the patient must be supported, during the treatment of consumption,	—	156
	℞	Succus

Succus Cicutæ spissatus, a direct sedative,	191, 207
—————, useful in pulmonary complaints,	136, 180
Stramonium, or Thorn-Apple, its properties considered,	212

T.

Tubercle, remarks on,	—	—	46
—————, frequently formed in asthma,	—	—	74
—————, description of,	—	—	110
—————, Origin and seat, in the air cells and not in the connecting cellular membrane	—	—	113
—————, Formation of,	—	—	116
—————, Vascularity, considered,	—	—	119
—————, relation between it and scrophula,	—	—	121
—————, inflammation, suppuration, and ulceration of,	—	—	124
—————, absorption of,	—	—	126
—————, prevention and cure of,	—	—	128

U.

Ulceration, may be sometimes cured by diminishing the irri- table action of the absorbents,	—	—	197
————— of the lungs, the treatment of,	—	—	139
—————, why so difficult to cure,	—	—	142
—————, the effects of oxygen and hydrogen air on,	—	—	159

... of the ...
... of the ...
... of the ...

T

... of the ...
... of the ...
... of the ...
... of the ...
... of the ...
... of the ...
... of the ...
... of the ...
... of the ...
... of the ...

... of the ...
... of the ...
... of the ...
... of the ...
... of the ...

