Aphorisms on the application and use of the forceps and vectis, on preternatural labours, on labours attended with hemorrhage, and with convulsions / by Thomas Denman, M. D. licentiate in midwifery of the College of Physicians, London; and honorary member of the Royal Medical Society at Edinburgh.

Contributors

Denman, Thomas, 1733-1815. University of Bristol. Library

Publication/Creation

London: Printed for J. Johnson, St. Paul's Church Yard, 1793.

Persistent URL

https://wellcomecollection.org/works/ga9w95st

Provider

Special Collections of the University of Bristol Library

License and attribution


This material has been provided by This material has been provided by University of Bristol Library. The original may be consulted at University of Bristol Library. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



1512732107




UNIVERSITY OF BRISTOL


MEDICAL LIBRARY

Reste Med. 18

HELF

.A.





APHORISMS

ON THE

APPLICATION AND USE

OF THE

FORCEPS AND VECTIS;

ON PRETERNATURAL LABOURS,

ON LABOURS ATTENDED WITH HEMORRHAGE,

AND WITH CONVULSIONS.

THE FOURTH EDITION.

By THOMAS DENMAN, M.D.

LICENTIATE IN MIDWIFERY OF THE COLLEGE OF PHY-SICIANS, LONDON; AND HONORARY MEMBER OF THE ROYAL MEDICAL SOCIETY AT EDINBURGH.

LONDON:

PRINTED FOR J. JOHNSON, ST. PAUL'S CHURCH YARD.

1793.





DR. THOMAS SAVAGE,

DEAR SIR,

THESE Aphorisms, designed for the use of students, I request you will permit me to dedicate to you. I wish also to take this opportunity of conveying those sentiments of respect I have ever entertained for the integrity and benevolence which so eminently distinguish your character; and to express my gratitude for that kindness and partiality you have always shewn to,

DEAR SIR,

Your obliged, and affectionate humble fervant,

Dec. 7, 1792. THOMAS DENMAN.

ARRANGEMENT

OF

LABOURS.

FOUR CLASSES.

I. NATURAL.

II. DIFFICULT.

III. PRETERNATURAL.

IV. Anomalous, or Complex.

CLASS I. NATURAL LABOURS.

CHARACTER. Every labour in which the process is completed within twenty-four hours, the head of the child presenting, and no adventitious assistance being required.

VARIETIES.

I. The face inclined towards the facrum.

2. The face inclined towards the offa pubis.

B

3. The

THE RADIO WARRA

4.0

and U o a h

TOUR CLASSES

ARREST AND

Carallela in Fig.

ILL PARTERIAL TRACT

manager De se je udukurum A. . .

Value of Language de Co

edi disidi ni arcidi petali anteria talen umbergaran migilia basel peparai fusione gallimeter philosophia trabali siti pransi englimitat pransitini ambitansaka da base

V KRISTERS.

A. The face inches towards the faceous.
2. The face inchined towards the eff public.
3. The face inchined towards the eff public.
3. The face inchined towards the effective face and the face of the

- 3. The head presenting with one or both arms.
- 4. The face presenting.

That part of a child which descends lowest into the pelvis, is to be esteemed the presenting part.

Circumstances attending Labours.

- 1. Anxiety.
- 2. Rigours.
- 3. Strangury.
- 4. Diarrhœa.
- 5. Mucous discharge, with or without a mixture of blood.
- 6. Pain.

Causes of pain.

- 1. Expulsatory action of the uterus.
- 2. Resistance made to the effect of that action.

Distinctions of pain.

- I. True.
- 2. False.

Causes and figns of false pain. Means of removing them.

B 2

Means

gial price in a golden price in a series

That part of a child which deficant lower -

Giver Cours attacker Labour.

- t, Amilei
 - model is
- andapuring &
- atomic set
- a modern ve dite providelle seconde .
 - Alle I de l'alle

Cuffe of pain.

a Regulator Santon of the interest of that a state of that a state of that a santon of that

Difficultion of paint E

and a

day side to sond hos silve!

Means of comering them.

Since NIL

Means by which true pains are supposed to be regulated, and their effect promoted.

Note. The pains attending labour are subsequent to the action of the uterus, though in common language the word pain, and the action of the uterus, are used synonymously.

Progress of natural labours.

Three periods or stages.

Ist period.

Dilatation of the os uteri.

Rupture of the membranes.

. Discharge of the waters.

2d period.

Descent of the child.

Dilatation of the external parts.


Expulsion of the child.

3d period.

Separation of the placenta.

Expulsion or extraction of the placenta.

Note. It very often happens that the membranes do not break till the head of the child is on the point of being expelled. This



is the natural and most desirable progress of a labour, and it is a negative proof that the labour has been well conducted; that is, not interrupted. But the description given above, will answer the purpose of impressing a clear idea of labours in general.

The two circumstances which principally require attention in natural labours are, to guard the perinæum and to extract the placenta with discretion.

CLASS II. DIFFICULT LABOURS.

CHARACTER. Every labour in which the process is prolonged beyond twenty-four hours, the head of the child presenting.

Note. Some objections may be made to this definition taken from time, but it will be found to apply to practical uses better than if it was taken from circumstances.

It would often be extremely difficult to fay with precision when a labour actually begins, because

is a natural and made defeable progressions in the or and it is a negative proof that the last the last that the last that the factor has been used conducted; that is, not interrupted. But the defealation riven they acted will answer the purpose of imprecising a clear

1 4 1

villagioning dulider round fine unit of cult of a control of contr

Cines II. Dericular Lacoune.

CHEANGTER! Describered in which the

Tops 1. School of Short may be smade to this desirable to the desirable to the short the short that it will be found to apply to profited ules better than if

Ha as studility the extremely difficult to a sale of the sale of t

because of the number of concurrent changes. But in general some progress must be made before we can allow a labour to be commenced.

FOUR ORDERS.

ORDER I.

Labours rendered difficult from the inert or irregular action of the uterus.

CAUSES.

- 1. Too great distention of the uterus.
- 2. Partial action of the uterus.
- 3. Rigidity of the membranes.
- 4. Imperfect discharge or dribbling of the waters.
- 5. Shortness of the funis umbilicatis.
- 6. Weakness of the constitution.
- 7. Fever.
- 8. Want of a due degree of irritability.
- 9. Passions of the mind.
- 10. General deformity.

ORDER

Sherral know Island said lanes in 1981 TOUR ONDERS.

ORDER II.

Labours rendered difficult by the rigidity of the parts to be dilated.

- 1. First child.
- 2. Advancement in age.
- 3. Too early rupture of the membranes.
- 4. Oblique position of the os uteri.
- 5. Fever or local inflammation.
- 6. Extreme rigidity of the os uteri.
- 7. Uncommon rigidity of the external parts.

ORDER III.

Labours rendered difficult from disproportion between the dimensions of the cavity of the pelvis and the head of the child.

- 1. Original smallness of the pelvis.
- 2. Distortion of the pelvis.
- 3. Head of the child unufually large, or too much offified.
- 4. Head of the child enlarged by disease.
- 5. Face inclined towards the offa pubis.
 - 6. Presentation

ti singol

Library resident and the state of the state of the Land and Land a

建冷却是让

al Adenoe neet in open

manufactured to support the over the

A thin the fall on the party and the fall

and the state of t

and to the tagellists amount to

Letting father and harpling a real model.

AM SACRO

Letterer rendered of their franciscoperation have a decision of the carrier of the property of the property of the property of the property of the carrier o

Original finalinule of the polying

To the Land of the

1

A Hand Street Counties and Street by discolor to the country of th

- 6. Presentation of the face.
- 7. Head presenting with one or both arms.

ORDER IV.

Labours rendered difficult by diseases of the soft parts.

- 1. Suppression of urine.
- 2. Stone in the bladder.
- 3. Excrescences of the os uteri.
- 4. Cicatrices in the vagina.
- 5. Adhesion of the vagina.
- 6. Steatomatose tumours.
- 7. Enlargement of the ovaria.
- 8. Rupture of the uterus.

NOTE. The disturbance of the natural progress of labours, more especially the premature rupture of the membranes, is the most general cause of difficulties in parturition.

Women are to be relieved in difficult labours,

- 1. By time and patience.
- 2. By encouragement to hope for a happy event.

3 among the second of and equil our anome you have

- 3. By regulating their general conduct.
- 4. By lessening the obstacles to the effects which should be produced by the pains.
- 5. By the affiftance of instruments.

Intentions in the use of instruments.

- I. To preserve the lives both of the mo-
- 2. To preserve the life of the mother.
- 3. To preferve the life of the child.

Instruments contrived to answer the first intention.

1. Fillets. 2. Forceps. 3. Vectis.

Three things are to be confidered with respect to the Forceps or Vectis, and to the use of instruments in general.

- 1. To make an accurate distinction of those cases which require their use.
- 2. Of those cases which allow their use.
- 3. Of the manner in which they ought to be used.

We are in the first place to speak of the application and use of the forceps.

Directions

0.0

the state of the s

Athenia de de chrai antenda. Le company de chrain antenda. Le company de chrain antenda de company d

2. To present the life of the moulter, y. To preferre the life of the child.

Anglergensen come land in angles of a flag Administra or Fillers in Marcoph. In Million

Firmer things are to be confident with respect to the till of inflataments in grants.

1. To make on accurate difficulting of facility and a second of the cales which allow designally a second of the planter in which they or the facility of the bearing.

- We are in the first place to figure of the


Directions for, and admonitions in, the application and use of the Forceps.

SECTION I.

- rule, that no instruments are to be used in the practice of midwifery; the cases in which they are used are therefore to be considered merely as exceptions to this rule.
- 2. But such cases can very seldom occur in the practice of any one person; and when they do happen, neither the *forceps* or any other instrument is ever to be used in a clandestine manner.
- 3. The first stage of a labour must be completed, that is, the os uteri must be dilated and the membranes broken before we think of applying the forceps.
- 4. The intention in the use of the forceps is, to preserve the lives both of the mother and child, but the necessity for using them must be decided by the circumstances of the mother only.

and the second of the second or to the firms Billian and Autor Stelle expensively and the last

- 5. It is meant, when the forceps are used, to supply with them the insufficiency or want of labour pains; but so long as the pains continue, we have reason to hope they will produce their effect, and shall be justified in waiting.
- 6. Nor doth the cessation of the pains always prove the necessity of using the forceps, as there may be a total or a temporary cessation of the pains.
- 7. In the former, the pulse, the countenance, and the general appearances of the patient indicate extreme debility, and resemble those of a person worn out with disease or fatigue.
- 8. But in the latter there are no alarming fymptoms, and the patient often enjoys short intervals of refreshing sleep.
- 9. A rule for the time of applying the forceps has been formed from this circumstance; that, after the cessation of the pains, the head of the child should have rested for six hours in such a situation as to allow the use of the forceps before they are used.



- to prevent the rash and unnecessary use of the forceps, must be subject to the judgment of the person who may have the management of any individual case.
- 11. Care is also to be taken that we do not, through an aversion to the use of instruments, too long delay that assistance we have the power of affording with them.
- 12. The difficulties which attend the application and use of the forceps are far less than those of deciding upon the proper time when, and the cases in which, they ought to be applied.
- 13. The lower the head of the child has descended, and the longer the use of the forceps is deferred, the easier will in general their application be, the success of the operation more certain, and the hazard of doing mischief less.
- 14. The forceps should always be applied over the ears of the child; it must therefore be improper to apply them when we cannot feel an ear.

(II)

behave it idea tribe grave base side mil .com
off to the complete tribes when the distributed tribes of the
and to the complete tribes which as described as the complete tribes are tribes of the complete tribes and the complete tribes are tribes of the complete tribes and the complete tribes are tribes are tribes and tribes are tribe

ablitude one express on the bear and the first of the bear and the first one of the first o

Applications and the local of the company and the company of the c

self .as

15. But when an ear can be felt by a common examination, the case is always manageable with the *forceps*, if the circumstances of the mother require their use.

16. The ear of the child which can be felt, will be found toward the offa pubis, or under one of the rami of the ischia.

17. The ears are not turned to the sides of the pelvis till part of the hind head has emerged under the arch of the osa pubis, when the use of the forceps can very seldom be required.

18. When we have determined on using the forceps, and explained the necessity of using them to the patient and her friends, she is to be placed in the usual position on her left side, near to the edge of the bed; and the instruments, warmed in water and smeared with some uncluous application, are to be laid conveniently by you.

Note. Women, impelled by their fears and their sufferings in difficult labours, will very generally implore you to deliver them with instruments long before you will be convinced of the necessity of using them. In

 many cases I have found it expedient and encouraging to them to fix upon some distant time when they should be delivered, if the child were not before born; six or eight, or twelve hours, for instance. In some cases of great apprehension I have also shewn them, upon one of my knees, all that I intended to do with the forceps.

The following rules are given on the prefumption that the head of the child presents with the face inclined or verging towards the hollow of the facrum, and that the common short forceps are intended to be used; but if any other kind of forceps should be preserred, the rules must be adapted to the instrument.


SECTION II.

- 1. Carry the fore finger of the right hand to the ear of the child.
- 2. Then take the blade of the forceps to be first introduced by the handle in the left hand, and conduct it between the head of the child and

which is the beautiful to the second property and the or believe to the line of the contract of the

and the finger already introduced, till the point reaches the ear.

- 3. The farther introduction must be made with a motion resembling a slight degree of Temi-rotation, and the point of the blade must be kept close to the head of the child, by gently raising the handle as the instrument is advanced.
- 4. The blade of the forceps must be carried up till the lock reaches the external parts, near the inferior edge of the osa pubis.
- 5. Should any difficulty occur in the introduction of either of the blades, we must withdraw them a little, to discover the obstacle, and never strive to overcome it with violence.
- 6. When the first blade is introduced, it must be held steadily in its situation, as it will be a guide in the introduction and application of the second blade.
- 7. The second blade of the forceps must be conducted upon the fore singer of the left hand, passed between the head of the child and the perinæum, in the same cautious man-



ner as the first, till the lock reaches the perinæum, or even presses it a little backward.

- 8. When the fecond blade is properly introduced, its fituation will be opposite to the first.
- 9. In order to lock the forceps, the handles of which are at a confiderable distance from each other, the blade first introduced must be brought down and carried so far back that it will lock with the second blade, held in its first position.
- to. Care should be taken that nothing be entangled in the lock of the forceps by carrying the finger round it.
- 11. It is convenient to tie the handles of the forceps together, when locked, with force fufficient to keep them from sliding or shifting their position.
- 12. If the blades of the forceps were introduced fo as not to be opposite to each other, they could not be locked.
- 13. Should the handles of the forceps when applied come close together, probably the bulk of the head is not included between them,

er ge the first sinches and an inches entrances and an

of which wis a considerable different at a considerable difference from a such original difference from and original original and considerable fix back make another the back make another the back make another the back make another the back and another the

be offered to the fact of the particular of the profiles.

The offered to the fact of the fact of the particular of the

the furrest engages of the desired with furest

deced in as one to be oppose to the control information of the second control of the sec

the constitution of the second of the second

and therefore when we acted with them they would flip.

- 14. If the handles when locked are at a great distance from each other, they are not well applied, and will probably slip.
- 15. But in these estimations allowance is to be made for the different dimensions of the heads of children.
- 16. The forceps will never slip if judiciously applied, if the case be proper for their use, and we act circumspectly with them.

Note. The difficulties in the application of the forceps arise, from attempting to apply them too soon; from passing them in a hurry, or in a wrong direction; or from entangling the soft parts of the mother between the instrument and the head of the child. Of course, we are always to be guarded against these circumstances.

the second

SECTION III.

- 1. There is no occasion, and it would be hurtful to attempt to change the position of the head, when the forceps are applied, before we began to extract.
- 2. For if the action with the forceps be flow, the head of the child will turn in the same manner, and for the same reasons, as in a natural labour.
- 3. Therefore the forceps being fixed upon the head must also change their position according to its descent, and the handles be gradually turned from the offa pubis and sacrum, where they were first placed, to the sides of the pelvis.
- 4. The handles of the forceps likewise, though originally placed far back towards the facrum, that is, in the direction of the cavity of the pelvis, are to be gradually turned, as the child advances, more and more towards the pubes, that is, in the direction of the vagina.

D . 5. The

- 5. The first action with the forceps must be to bring the handles, firmly grasped in one or both hands, slowly towards the pubes till they come to a full rest.
- 6. After waiting till the pains return, or an imaginary interval if there should be a total want of pain, the handles are to be carried back in the same slow and cautious manner to the perinæum, using at the same time a certain degree of extracting force.
- 7. The subsequent actions must be from handle to handle, or occasionally by simple traction; but the action of that blade which was towards the pubes, must be stronger and more extensive throughout the operation, than the action with the other blade which has no fulcrum to support it.
- 8. By a repetition of these actions, always directed according to the position of the handles, with their force increased, diminished, or continued, according to the exigence of the case, we shall in a short time perceive the head of the child descending.

D 2

g. When

being at of the william for any of the state of the state

The division spiriture and the problem of the probl

And the continue of the contin

Mark to St.

.

- 9. When the head begins to descend, the force of the action with the forceps must be abated, and as that advances, the direction of the handles must change by degrees more and more to each side, and towards the pubes.
- fcends, the more gently we must proceed, in order to prevent any injury or laceration of the perinæum or external parts, which are likewise to be supported in the same manner as in a natural labour.
- casioned by the application of the forceps, or the very expectation of their being applied, will bring on a return or an increase of the pains sufficient to expel the child without their assistance.
- 12. In other cases we are obliged to exert very considerable force, and to continue it for a long time; so that one operation may be safely and easily finished in twenty minutes, or even a less time, and another may require more than an hour for its completion, and

and the death of subset had no in the to that want to be the same with the course to be

,

the repeated exertions of very confiderable force.

- 13. In some cases it happens also, that the obstacle to the delivery exists at one particular part of the pelvis, and when that is surmounted, the remainder of the operation is easy; but in other cases there is some difficulty through the whole course of the pelvis.
- 14. Before the exertion of much force we are always to be convinced that a small or a moderate degree of force is not equal to our purpose.
- 15. In every case in which the forceps have been applied, they are not to be removed before the head is extracted, even though we might have little or no occasion for them.
- 16. When the head of the child is born the forceps are to be removed, and the remaining circumstances are to be managed as if the labour had been natural.

Note. The general arguments against the use of instruments have been drawn from their abuse: it appears, however, that necessity

will

the repeated wastrious of very confidential

and their policy services in heavy and the service of a particular and their contractions and their services and their services are followed to the services and their services are followed to the services are services as a service and their services are services as a service and their services are services as a service and their services are services as a service and the services are services as a service and

and angel during to magnetic additional to a state of the state of the

The state of the s

Rolling the county branch of a county of a

will justify the use of the forceps; that when such necessity exists, their use is not only justifiable, but often highly advantageous; that delay to apply them, and slowness in their application and use, will secure, as far as is possible, both the mother and child from untoward accidents; but that mischief cannot be prevented if they are applied too soon, or the operation with them be performed in a hurry.

It would be a very defirable thing that every student should have an opportunity of seeing the operation with the forceps performed before he goes into practice; but that is not always possible. Yet if he has been properly instructed in the principles of the application and use of the forceps, reslects seriously before he determines on performing the operation, and proceeds slowly but not timidly in it, he can hardly fail to succeed. Hurry, in any operation, is a very common sign both of want of information and of sear; and attention is to be paid to the order of the rule in Celsus, 1. tutò, 2. citò, 3. jucundè.

SECT. IV.

On the application and use of the Vectis.

- I. We shall have a just idea of the vestis by considering it as one blade of the forceps a little lengthened and enlarged, with the handle placed in a direct line with the blade.
- 2. The general condition and circumstances of labours before stated, as requiring and allowing the use of the forceps, will hold equally good when the vectis is intended to be used.
- 3. In the application of the vectis two fingers, or the fore finger of the right hand is to be passed to the ear of the child.
- 4. Then taking the vectis by the handle, or with the blade shortened in the left hand, conduct it slowly till the point of the vectis reaches the ear, however that may be situated.
- 5. The instrument is then to be advanced, as was advised with the forceps, till according to your judgment the extremity of the blade reaches

1 22 J

1.72 22.2

the river at the land and the fact of the course

Aparella de la cidade de la compania de la gracio de la compania del compania del compania de la compania del compania del compania de la compania del compania del compania del compania de la compania del compania

reaches as far, or a little beyond the chin of the child.

- 6. Then grasping the handle of the instrument firmly in the right hand, wait for the accession of a pain.
- 7. During the continuance of the pain raise the handle of the instrument gently but firmly towards the pubes, drawing at the same time with some degree of extracting force.
- 8. When the pain ceases let the instrument rest, and on its return repeat the same kind of action, alternately resting and acting in imitation of the manner of the pains.
- 9. By a repetition of this kind and manner of action the head of the child is usually advanced, and the face turning gradually towards the hollow of the facrum, the position of the handle of the vectis will be altered, and the direction of the action with it of course be changed.
- we must proceed more slowly and carefully, according to the degree of descent, in order to prevent any injury to the external parts, which

To mile alle buoyes almil a to tall a resident

-matter out to offered an element of the settlers.

Acres a la maistre

Services and property of the control of the formation of the control of t

designation of the minutes of the policy of

of action the tales of the caldidate of these of the caldidate of the cald

to prevent any input to the categories of defending to the depose of defending to the part of the categories and a to prevent any input, to the categories and a to prevent any input, to the categories and a to prevent any input, to the categories and a

which is to be prevented, as was directed, when the forceps are used.

- rate force before recommended, the head should not descend, it must be gradually and cautiously increased till it becomes sufficient to bring down the head.
- part of the instrument must rest upon the symphysis of the offa pubis, or upon the ramus of the ischium according to its position, as upon a fulcrum, for its support.
- 13. By passing the flat part of the hand to the back of the blade of the instrument when in action, we shall be occasionally able to lessen or take off this pressure which must otherways be made upon the parts of the mother.
- 14. Some have recommended the vectis to be used when the head of the child was higher up in the pelvis than is before stated, as justifying the use either of this instrument or the forceps.
- 15. They have also recommended the vectis when the head of the child was firmly locked

in the pelvis, and have afferted that by its use there is often obtained a very good chance of preserving the life of a child, which must otherwise be inevitably lost.

- 16. Others have by frequent use acquired such dexterity as to be able to extract the head of a child in the situation first stated, with a single sweep of the instrument.
- 17. Some have also advised the introduction of the vectis between the facrum, or sacroficiatic ligaments, and the head of the child, from a belief that it could be equally or more advantageously used in this position than in that first stated.
- 18. But having ever considered the use of all instruments as a thing to be lamented, and when I did use them, esteemed the safety of using them as my principal object, I cannot deviate from these principles, or enter upon a discussion of points of practice, of which, as far as I am competent to judge, I cannot approve.

Note. Before, and immediately after the publication of my second Essay on Difficult E Labours,

ı and the second

Labours, feveral Gentlemen, with whom I converse, and to whom I ought to pay great respect, reprehended in very decided terms what I have advanced with regard to the forceps and vestis. Some maintained that the forceps is an instrument far superior to the vectis, of which I was accused of speaking too favourably. Others, of equal respectability, accused me of speaking with timidity, or restraint, of those advantages which, they asferted, the vectis had over the forceps. This very strong evidence could only be invalidated by its contradiction, but the very respect which I bear to the witnesses, compelled me to pass over their evidence, and to rely upon my own experience and judgment.

I did not speak of the mechanism of the instruments, or of the operation performed when we had applied, and acted with them, as these have hitherto been very imperfectly and often erroneously explained. The subject came under consideration in the ordinary course of the work, and having frequently used both the instruments, I stated

E 2

tentiled you wanted one road have yourself

T. C.

the matter equitably, according to the best of my abilities, and in such a way that, I thought, students, who were principally concerned in the discussion, being left with the choice of either instrument, according to the doctrines of the particular professors whom they might attend, could not be missed. It is not to be expected that men versed in practice should change their opinions or alter their practice, or, in short, pay much regard to disputes about instruments, if any were disposed to raise them.

It then was, and yet remains my opinion, founded, as I before observed, on my experience with both instruments, that the superior excellence which has been attributed to each of these instruments, ought chiefly to be ascribed to the dexterity which may be acquired by the habit of using either of them. It is also my opinion that we may, in general, either with the forceps or vestis, effectually and conveniently give that assistance which is required in cases of difficult parturition, allowing and justifying their use. In particular cases

9 to fled see or unification assistings inversely Applications of the property of the more and the property THE RESERVE OF THE PARTY OF THE

cases it may perhaps be proved that one inftrument is more commodious than another.

But if the vectis be depreciated by those who have never used it, and are not expert in its use, because they prefer the forceps, or if the known properties of the forceps be not allowed by those who do not use them, because they prefer the vectis, the proper inference would not be, that either of the instruments ought to be condemned; but that we are in possession of two instruments well adapted to answer the same purpose, if they are prudently used.

CLASS III. PRETERNATURAL LABOURS.

CHARACTER.—Labours in which any part of the child prefents, except the head.

TWO ORDERS.

ORDER I.

Presentations of the Breech, or inferior Extremities.

ORDER

Control of the second distance of the second of the second

TWO CASE CITE

ar and the same

Profession of the Break or inferior Phys.

SALGEO.

ORDER II.

Presentations of the Shoulder, or Superior Extremities.

SECTION I.

- of birth may be of three kinds. I. With the head. 2. With the breech, or inferior extremities. 3. With the shoulder, or superior extremities.
- 2. Presentations of the first kind are called natural, those of the second and third kind, preternatural.
- 3. Preternatural prefentations have been fubdivided into a much a greater variety, but without any practical advantage.
- 4. The prefumptive figns of the preternatural prefentation of children are very uncertain, nor can it ever be determined what the prefentation

MOUTOUR !

AND THE RESIDENCE OF THE PARTY OF THE PARTY

Sile-ries

delles see inch first six to consider our see and consider decided and see about decided.

and seed interesting house with the seed of the seed o

contrativity of a significant problem, and if the second problem is the second problem of the second problem in the second problem is the second problem in the second problem in the second problem is the second problem in the second problem in the second problem is the second problem in the second problem in the second problem is the second problem in the second problem in the second problem in the second problem in the second problem is the second problem in the second problem i

presentation is, till we are able to feel the prefenting part.

5. When any part of a child can be felt, we may form our judgment of the presenting part by the following marks.

6. The head may be distinguished by its

roundness, its firmness, and its bulk.

- 7. The breech may be known by the cleft between the buttocks, by the parts of generation, and by the discharge of meconium.
- 8. The foot may be distinguished by its length, by the heel, by the shortness of the toes, and the want of a thumb; and the hand by its slatness, by the thumb, and the length of the singers.

SECTION II.

On the first Order of Preternatural Presentations.

1. In this kind of presentation the breech, one hip, the knees, and one or both legs, are to be included.

a side on the pictories

perpulsion is all we are able to led the pre-

or id both bearing of the Real of a

Asia and his automa and your end of the later of

tion in the second second second second second

The state of the s

- 14 - 110 - 120

Or als part, to be at Proposition to

the best of the bear of the parties and the state of the

of the

- 2. In these presentations it was formerly supposed necessary, as soon as they were discovered, to introduce the hand to bring down the feet, and to extract the child with expedition.
- 3. But, according to the present practice, such labours are not to be interrupted, but allowed to proceed as if the presentation was natural; unless the necessity of giving assistance should arise from some circumstance independent of the presentation.
- 4. By acting on this principle, when the breech of the child is expelled by the pains, the parts are sufficiently distended to allow the body and head to follow without any danger from delay.
- 5. But if the feet of the child were to be brought down in the beginning of labour, the difficulty with which it would be expelled or could be extracted, increasing as it advanced, the child would probably die before the woman was delivered, and she would be in danger of suffering mischief.

windered approved annual angular of the control of

seem lib

All approximates the continue of the process of the continue of the process of the continue of the process of the continue of

call mader adjudants and an accordance of the control of the contr

E A

- 6. In cases of this kind there is also equal reason, when the breech is on the point of being excluded, for our guarding the perinæum from the hazard of laceration as in presentations of the head.
- 7. In first labours, the child, unless it be very small, will not unfrequently be born dead when the breech, or inferior extremities, present; but in subsequent labours they will usually be born living, if there be no other impediment than that which is occasioned by the presentation.
- 8. The injuries which the presenting part of the child, especially the penis and scrotum, may sustain will often be alarming, and appear dangerous, but by soothing and gentle treatment, they are soon recovered.
- 9. Should there be reason to think the child dead, or the powers of the mother insufficient to expel it, we must then give such assistance as may be required.
- 10. This affishance must be given with the hand, or with a blunt hook or crotchet, hitched in the groin of the child; or, which I

prefer,

designation of sends being either to educe of the sends gradules with the sends of the sends of

(22)

The state of the s

E. The inchies which the preferring part
of the child, electricity the principal region
over the child, electricity the principal region
over the child, electricity the manning that the
principal diagraphs had by inothing man grands

Eliterate dans considered and chings at

The same of the sa

prefer, by paffing a ligature round the bent part of the child at the groin, with which we can hardly fail to extract it.

11. But every affistance of this kind must be given with discretion, and we must first be convinced of the necessity before we interfere.

SECTION III.

Of the second Order of Preternatural Presentations.

- 1. In this kind of presentation are included the shoulders, the elbows, and one, or both arms.
- 2. In all these presentations we shall be under the necessity of turning the child, but as they may be attended with circumstances widely different, it is necessary to make the following distinctions.
- 3.—1. When the os uteri is fully dilated, the membranes unbroken, or the waters lately discharged,

 discharged, a superior extremity being perceived to present, before the uterus is contracted.


- 4.—2. When the membranes break in the beginning of labour, the os uteri being little dilated.
 - 5.—3. When the os uteri has been fully dilated, the membranes broken, and the waters long discharged, the uterus being at the same time strongly contracted, and the body of the child jammed at the superior aperture of the pelvis.
 - 6.—4. When, together with any of these circumstances, there is a great disproportion between the size of the head of the child, and the dimensions of the cavity of the pelvis.

SECTION IV.

On the Cases which come under the first Distinction.

1. Whenever there is a necessity of turning a child, the patient is to be placed upon

F 2



her left fide, near the edge of the bed; or fometimes, when we expect or find much difficulty, in a prone position, resting upon her elbows and knees.

- 2. All the advantage to be gained from any particular position of the patient is, to allow us the free and dexterous use of our hands; the situation of the child not being altered by the position of the patient.
- 3. The os externum is then to be dilated with the fingers reduced into a conical form, acting with a femi-rotatory motion of the hand.
- 4. The artificial dilatation of all parts must be made slowly, in imitation of the manner of natural dilatation.
- 5. The os externum should be amply distended before the hand is carried farther, or its contraction round the wrist will be an impediment in the subsequent part of the operation.
- 6. When the hand is passed through the os externum, it must be slowly conducted to the os uteri, which being wholly or sufficiently dilated,

The second secon THE PROPERTY WILLIAM STATES AND ASSESSED. re-tain from a feature of the section of the section

dilated, we must break the membranes by perforating them with a finger, or by grasping them firmly in the hand.

- 7. The hand must then be passed along the fides, thighs, and legs of the child, till we come to the feet.
- 8. If both the feet lie together we must grasp them firmly in our hand; but if they are distant from each other, and we cannot conveniently lay hold of both feet, we may deliver by one foot without much additional difficulty.
- 9. Before we begin to extract we must be assured that we do not mistake a hand for a foot.
- 10. The feet must be brought down, with a slow waving motion, into the pelvis; when we are to rest and wait till the uterus begins to contract, still retaining them in our hand.
- on, the feet are to be brought lower at each return of pain, till they are extracted through the external orifice, and the labour may then be finished, partly by the efforts of the mother, and partly by art.

12. If

- 12. If the toes are turned towards the pubes, the back of the child is towards the back of the mother, which is an unfavourable position.
- 13. But if the toes are towards the facrum, the back of the child is towards the abdomen of the mother, which is proper; and all other positions of the child must be gradually turned to this as the body is extracting.
- 14. Yet this position of the child is only advantageous when the head comes to be extracted.
- 15. Wrap the feet of the child in a cloth, and wait till there is a contraction of the uterus, or a pain, during the continuance of which gently draw down the feet.
- 16. When the pain ceases we must rest, and proceed in this manner through the delivery, assisting the efforts of the patient, but not making the delivery wholly artificial.
- 17. When the breech comes to the os externum, the child must be extracted very slowly through it, and in the proper direction, or there

processes the process of a street of a win has to contract salmy and the first than the second state of the second

there will be danger of lacerating the pe-

- 18. When the child is brought so low that the funis reaches the os externum, a small portion of it is to be drawn out, to slacken it to lessen the chance of compression, or to prevent the separation of it from the body of the child, or of the placenta from the uterus; and from this time the operation should be finished as speedily as it can with safety.
- 19. But if the circulation in the funis be undisturbed there is no occasion for haste, as the child, we are then assured, is in safety.
- 20. The child may be extracted without much difficulty if we act alternately from fide to fide, by making a lever of its body, and fometimes by pressing it from the offa pubis with the singers.
- 21. If the child should stick at the shoulders, the arms must be successively brought down.
- 22. This is to be done by raising the body the opposite way, and by bending them at the elbow very slowly, lest they should be broken, and

will oils with some to refund to their souls

and the hand must be cleared toward the pubes.

- 23. When both the arms are brought down, the body of the child must be supported upon our left hand placed under the breast, and the singers on each side of the neck.
- 24. Then placing the right hand over the shoulders, and pressing with our singers the head towards the facrum, we must ease the head along, gradually turning the body of the child as it advances toward the abdomen of the mother.
- 25. If the head should not come easily away, we must introduce the fore singer of the lest hand into the mouth of the child, by which the position of the head will be rendered more convenient.
- 26. When the head begins to enter the os externum, we must proceed very slowly, and support the perinæum, by spreading the singers of the lest hand over it.
- 27. In some cases there may be a necessity of speedily extracting the child in order to preserve its life, but we must also recollect, that



that the child is often lost by endeavouring to extract it too hastily.


28. When a child has been extracted by the feet, the placenta usually separates very soon and very easily; but in the management of this we are to be guided by the general rules.

SECTION V.

On the Cases which come under the second Diftinction.


- 1. We are first to ascertain the presenting part, and if, together with the arm, the head is perceived by a common examination, there may be no occasion to turn the child, such case only constituting the third variety of natural labour.
- 2. But if the case should be such as to require the child to be turned, it might be doubted whether it were proper to dilate the os uteri by art, or to wait for its spontaneous dilatation.

3. Perhaps



- 3. Perhaps neither of the methods can be constantly followed, but we may generally say, that there is under these circumstances neither danger or increase of difficulty, from waiting for the spontaneous dilatation, which is therefore in general to be preferred.
- 4. But if more speedy dilatation should be required, whatever is done by art should be done slowly, and in imitation of nature.
- 5. The os uteri is always to be confidered as completely dilated when we judge it will allow of the easy introduction of the hand.
- 6. When we have fixed upon the proper time and begin the operation, the os externum must be dilated in the manner before advised.
- 7. The hand must always be introduced into the uterus, or that side of the pelvis where it will pass most conveniently; and there is usually most room at that part which will lead to the feet.
- 8. It is generally most convenient to pass the hand between the body of the child and

G




the offa pubis, the feet being most commonly found lying toward the belly of the mother.

- 9. In cases which come under this distinction the uterus is seldom contracted very strongly upon the body of the child, but always in some degree.
- operation of turning the child, in these cases, will be fully explained under the following distinction.

SECTION VI.

On the Cases which come under the third Distinction.

- 1. The difficulty in the management of these cases depends upon the degree of contraction of the uterus, and upon the distance or awkward position of the feet of the child, but chiefly upon the former circumstance.
- 2. The uterus is in some cases contracted in a globular, and in others in a longitudinal form.



- 3. It is always easier with an equal degree of contraction to turn the child when the uterus is contracted in a globular, than in a longitudinal form.
- 4. When we are called to a case of this kind it is better not to form, or to give a hasty opinion, nor to attempt to deliver the patient immediately, but to deliberate upon it, and then to make a second examination.
- 5. If the fecond examination should confirm our first opinion, we may prepare for the operation.
- 6. We shall be able to judge in what part of the uterus the seet of the child lie, if we confider whether it be the right or left hand which presents, which may be known by the direction of the thumb and of the palm of the hand.
- 7. But the contraction of the uterus is the principal difficulty to be furmounted, and the danger in turning the child is in proportion to the difficulty.
 - 8. The danger in turning a child when there

of the contract of the contrac . there is a strong contraction of the uterus, is a single danger, that of rupturing the uterus.

- 9. The contraction of the uterus is of two kinds; first, the permanent contraction, in consequence of the waters having been long drained off, which may occur when there has been little or no pain.
- arising from the action of the uterus, returning at intervals, and always attended with pain.
- degree of force sufficient gradually to overcome the permanent contraction of the uterus, or the operation could never be performed.
- 12. But if we were to attempt to overcome the extraordinary contraction, it must follow, that we can, or cannot overcome it.
- 13. In the first instance we should be in danger of rupturing the uterus, and in the second the hand would be cramped, and we should be unable to proceed with the operation.
 - 14. The deduction is therefore clear, that

we ought not to proceed in our attempts to turn the child while the uterus is acting with violence.

- 15. The action of the uterus is rendered more frequent and strong by the generally increased irritability of the patient.
- 16. Before we attempt to deliver it will be prudent to endeavour to lessen this irritability, in many cases by bleeding, by clysters, and by an opiate, which, to answer this purpose, should be given in two or three times the usual quantity.
- 17. When the opiate takes effect, and the patient becomes disposed to sleep, we must consider this state as extremely favourable, and proceed without loss of time to the delivery.
- 18. There never can be occasion to separate the arm which presents from the body of the child, and when this has been done, instead of facilitating, it has impeded the operation.
- or left hand, as may be most convenient to ourselves,

Color 7

the grains of memoral states of the second

the against of the same in traderal

The state of the s

position of the opinion takes of the body and the particular of the body and the bo

The control of the co

The same of the sa

ourselves, must be introduced in the manner before directed, and conducted slowly into the uterus if there be sufficient room.

- 20. But if the child be jammed at the fuperior aperture of the pelvis, the hand cannot be introduced.
- 21. We must then fix our foresinger and thumb in the form of a crutch in the armpit of the child, and pushing the shoulders to-wards the head, and towards the fundus of the uterus, we must by degrees raise the body of the child till there be room for the introduction of the hand.
- 22. If while we are introducing our hand we perceive the action of the uterus come on, we must not proceed till that ceases or is abated.
- 23. The hand is also to be laid flat during the continuance of the action of the uterus, lest the uterus be injured by its own action on the knuckles.
- 24. When the action ceases or is abated, we must renew our attempts to carry up our hand to the feet of the child.

and some named of the state of a second second which seads this life because the four ore guinelt and half as of title at heart set? community of the state of the s

25. In this manner we are to proceed, alternately resting and exerting ourselves, till we can lay hold of one or both feet.

26. There is sometimes much difficulty in getting to the seet, and sometimes in extracting them, especially when the uterus is contracted in a longitudinal form.

27. In such cases it is often convenient, when we can reach the knees, to bend them cautiously, and to bring down the legs and feet together.

28. But before we begin to extract we should examine the parts we hold, and be assured they are the feet; and we must extract slowly and steadily.

29. If we hurry to bring down the feet they may flip from us, and return to the place from which they were brought.

30. We must then carry up the hand again, and grasping the foot or feet more firmly, bring them down in the cautious manner before advised.

31. When the feet are brought down, if there be difficulty in extracting them, we must

. The Superior of the Revenue of the state of Hor Foreign and the second of the second . THE RESIDENCE OF THE PARTY OF T the second of the second second second and the second second second second the state of the s

must endeavour to slide a noose, first formed upon our wrist, over the hand to secure the feet, by which the hazard of their return will be prevented, and the succeeding part of the operation much facilitated.

- 32. When the noofe is fixed over the ancles, we must pull by both ends of it with one hand, and grasp the feet with the other.
- 33. When there is afterward much difficulty in extracting the child, it is probably owing to the body of the child being jammed across the superior aperture of the pelvis.
- 34. It will then be proper to pass the finger and thumb as directed at 21, to raise the shoulders and body of the child toward the fundus of the uterus, with one hand, and with the other extract at the same time with the noose.
- 35. When the breech of the child has entered the pelvis, we must proceed with deliberation, but there will be little farther difficulty, except from the smallness of the pelvis, of which we shall speak in the next section.

the same of the sa

SECTION VII.

On those Cases which come under the fourth Distinction.

- 1. The disproportion between the head of the child and the dimensions of the pelvis, may be added to any of the circumstances mentioned under the preceding distinctions.
- 2. But as the management of these has been already directed, there is now occasion to speak only of the peculiar difficulties arising from that cause.
- 3. The degree of difficulty in these cases is greater or less according to the degree of disproportion; but the difficulty of extracting any part of the body of the child is little, compared with that which attends the extraction of the head.
- 4. We will therefore suppose the body of the child to be brought down, but that the head cannot be extracted by any of the methods before recommended.
 - 5. The force with which we endeavour to

the figure of the state of the server the same of the server extract must then be increased, till it is sufficient to overcome the difficulty or resistance.

- 6. But as the necessity of using great force can only be known by the failure of a less degree to produce the desired effect, we must begin our attempts with moderation, and gradually increase our efforts according to the exigence of the case.
- 7. The force exerted should also be uniform, controuled or commanded, and exerted by intervals, in the manner of the natural pains.
- 8. If the head should not descend with the force which we judge can be safely exerted, we must rest, and give it time to collapse.
- 9. We may then renew our attempts, extracting from fide to fide, or backwards and forwards, as may best conduce to ease the head through the distorted pelvis, alternately resting and endeavouring to extract.
- 10. But if the head should descend in ever so small a degree, the force is not to be increased with the view of finishing the delivery

H 2 expeditiously,

.

expeditiously, but we must be satisfied with our success, and proceed circumspectly.

- 11. When the head once begins to descend there is seldom much subsequent difficulty in finishing the delivery, as the cause of the difficulty usually exists at one particular part of the pelvis.
- 12. But should the head rest in this situation for several hours, no additional inconvenience would thence arise to the mother, and the longer it rested the greater advantage we should probably gain when we renewed our attempts to extract it.
- 13. It may be prefumed when the head of the child has been wedged for a long time in the position we are supposing, and great force has been used to extract it, that there is little reason to expect the child should be born alive; yet instances of this are said to have occurred in practice.
- 14. When we can hook a finger on the lower jaw of the child, the direction of the head may be changed to one more favourable, and the delivery thereby facilitated.

15. But

Tellin bullion and them convert the training A STATE OF THE PARTY OF THE PARTY OF THE PARTY OF THE PARTY.

- 15. But we must not extract with so much force as to incur the hazard of breaking or tearing away the jaw.
- offa pubis to the facrum, with the fingers carried up as high as we can reach, will often be of great use in these cases.
- 17. If the difficulty of extracting the head arises from its enormous size, occasioned by some disease, as the hydrocephalus, &c. these methods steadily pursued will answer our intention, as by a prudent use of the force in our power, the integuments will burst, or even the bones be broken.
- in which it feemed expedient to use either one, or both blades of the forceps, or to lessen the head.
- 19. But if such cases should occur, the utmost care must be taken that we do no injury to the mother.
- absolutely necessary to lessen the head of the child, the perforation may be conveniently made

1 The second secon The state of the s

made behind either of the ears, and the general rules of the operation must be followed.

21. By the force used should the neck of the child give way, we are not to separate the body from the head, but we must rest longer


and act moderately.

22. Should the body be separated from the head by the force we have used, or should we be called to a case of this kind, there will be no occasion for this reason alone to all hastily or rashly, as the head may even then be expelled by the pains.

23. But if this should be impossible, or is it be absolutely necessary to extract the head speedily, on account of the state of the mo-

ther;

24. Then the general rules for lessening the head must be accommodated to the exigencies of this particular case, and the head may be confined to a proper situation by compressing the abdomen.



SECTION VIII.

Miscellaneous Observations.

- 1. It fometimes happens that no part of the child can be perceived before the membranes break, though the os uteri be fully dilated.
- 2. In such cases we should not be absent when the membranes break, lest it should prove a preternatural presentation, requiring the child to be turned.
- 3. In some cases even when the os uteri isdilated, the membranes broken, and the waters discharged, no part of the child can be: felt.
- 4. It will then be prudent to introduce the hand into the uterus in the cautious manner before directed, to discover the part which does present.
- 5. If the head be found to present we should withdraw our hand, and suffer the labour to proceed in a natural way.
 - 6. If the inferior extremities should pre-

fent, we may bring down the feet, and then fuffer the labour to go on uninterruptedly.

- 7. But if the shoulder or superior extremities should present, we may proceed to the feet, and turn the child as was before directed.
- 8. By this conduct we shall guard against the danger of turning a child in a contracted uterus.
- 9. If we should be called to a case in which the arm presented and much force had been used to extract the child in that position, the arm having perhaps been mistaken for a leg, and the pains being at the same time violent, it may be impossible to turn the child, or even to introduce the hand into the uterus, the shoulder of the child being pushed low down into the pelvis.

per to attempt to introduce the hand into the uterus, or to turn the child, as it will be expelled by the efforts of the mother.

11. Yet in these cases the body of the child does not come doubled, but the breech

ARE REPORTED FROM THE MEDICAL PROPERTY OF THE SECOND and the second of the second o product of the party plants and a rebide

is the first part delivered, and the head the last, the body turning, as it were, on its own axis.

- gard to a small child coming prematurely, as it will apply to a child of a common size, and when a woman is at her full time, provided the pelvis be well formed.
- 13. This fact, of the possibility of a child being expelled in this position, though originally contradicted with great considence, is now confirmed in the most satisfactory manner by many cases which have been recorded, in some of which the children have even been born living.
- 14. From these it might be inferred that a woman in a state of nature, or in perfect health, would not die undelivered, though the arm of the child might present, supposing that she was not assisted by art.
- 15. Yet it is always requisite and proper to turn children when the superior extremities present, if the operation can be performed without the hazard of injuring the mother,

8

in the first part delivered, and the land the la

the contract of the second of the contract of

reing expelled it, on the continue of the cont

the second the child might be identical that a perfect twenty in the first metals, or in perfect beattles, would not als seconds and also seconds are seconds and also seconds and also seconds and also seconds are seconds are seconds and also seconds are seconds and also seconds are seconds and also seconds are seconds are seconds and also seconds are seconds are seconds are seconds and also seconds are seconds

to the state of the state to produce and produce to the state of the s

and we have generally a better chance of preferving the child.

- 16. But when there is no chance of preferving the child, and yet it cannot be turned without the greatest danger to the mother, knowing the possibility of its being expelled in this position, it is necessary to consider the propriety of the operation before we perform it.
- 17. It remains, however, to be proved by future experience, how far, and in what cases the preceding observation ought to be a guide in practice.
- 18. In cases of presentation of the superior extremities, in which the difficulty of turning the child would be very dangerous, and great or insurmountable, another method has been recommended.
- 19. But of this method, which has been practifed by one gentleman to whose knowledge and experience I pay great respect, I am not a competent judge, having never tried it.
 - 20. I therefore refer to the annexed note

1-17-1

and we have passerally a bitter chapte of gra-

income the call of the common is considered to the constant of the constant of

of hands of a second distance of the product by a second by a second distance of the contract and a second distance of the second distanc

minger and in neither where in a control of a control of the contr

And the second s

The second secon

for an explanation and history of the method to which I allude.

Note. Hoorneus, sæpe laudatus, adhuc peculiarem, novum eumque breviorem modum, sætum mortuum cum brachio arctissimè in vagina uteri hærente extrahendi, invenit atque descripsit, qui in eo consistit, ut quando ad pedes pervenire nequit, collum, utpote quod in sætibus valde adhuc tenerum est, vel scalpello a reliquo trunco resecet, vel unco idoneo quam cautissimè auserat. Hoc enim sacto, vel sponte mox prorumpit ex utero sætus, vel tamen, dum brachium propendens attrahitur, quod medico loco habenæ inservit, quam sacillimè excutitur. Caput vero deinde seorsim mox vel manu, vel aliis propositis artisiciis, si manus parum esset, ejiciendum.

HEISTER. cap. cliii. fect. ix.

The latter part of this description is further explained in the seventh section. the same series and till or of the metallic

The state of the s

The figure of the late of the second second

I AM induced to reprint the following, as they were the very cases which first gave me an opportunity of observing the spontaneous evolution.

CASE I.

In the year 1772, I was called to a poor woman in Oxford Street, who had been in labour all the preceding night, under the care of a midwife. Mr. Kingston now living in Charlotte Street, and Mr. Goodwin, surgeon, at Wirksworth, in Derbyshire, who were at that time students in midwifery, had been sent for some hours before I was called. The arm of the child presenting, they attempted to turn and extract it by the seet, but the pains were so strong as to prevent the introduction of the hand into the uterus. I found the arm much swelled and pushed through the external parts in such a manner,

that

coops all or visio to distance on the the series of th emined by the size has been been been be-AND THE RESIDENCE OF THE STATE The same action with a first and actions: ACCOUNT HAND IN FRANCE OF THE PARTY COLUMN

The woman struggled vehemently with her pains, and during their continuance, I perceived the shoulder of the child to descend. Concluding that the child was small and would pass, doubled, through the pelvis, I defired one of the gentlemen to sit down to receive it, but the friends of the woman would not permit me to move. I remained by the bed-side till the child was expelled, and I was very much surprised to find, that the breech and inferior extremities were expelled before the head, as if the case had originally been a presentation of the inferior extremities.

The child was dead, but the mother recovered as foon and as well as she could have done after the most natural labour.

CASE II.

In the year 1773, I was called to a womanin Castle Street, Oxford Market, who was attended: Control of the placementary belongs to the place of the p

21 2 3 3

The sales of the builts were it has a series of all of a series of the contract of the contrac

attended by a midwife. Many hours after, it was discovered that the arm of the child prefented. Mr. Burosse, surgeon, in Poland Street, was fent for, and I was called into consultation. When I examined, I found the shoulder of the child pressed into the superior aperture of the pelvis. The pains were strong, and returned at short intervals. Having agreed upon the necessity of turning the child, and extracting it by the feet, I fat down and made repeated attempts to raife the shoulder, with all the force which I thought could be fafely used; but the action of the uterus was fo powerful that I was obliged todefift. I then called to mind the circumstances of the case before related, mentioned them to Mr. Buroffe, and proposed that we should wait for the effect, which a continuance of the pains might produce, or till they were abated, when the child might be turned with less difficulty. No further attempts were made to turn the child. Then every pain propelled it lower into the pelvis, and in a little more than one hour the child

was born, the breech being expelled, as in the first case.

This child was also dead, but the mother recovered in the most favourable manner.

Having been prepared for observing the progress of this labour, I understood it more clearly, and attempted to explain both in my lecture on the subject, and in the aphorisms which were printed for the use of the students, my opinion of the manner in which the body of the child turned as it were, upon its own axis. I also pointed out the circumstances, in which, I supposed, the knowledge of the fact might be rendered useful in practice; but with great circumspection.

CASE III.

January the 2d, 1774, I was called to Mrs. Davis, who keeps a Toy-shop, in Crown Court, Windmill Street. She had been a long.

and the state of t

Hartrem that the bush of the best that

long time in labour, and the arm of the child presented.

The late Mr. Eustace had been called on the preceding evening, and had made attempts to turn the child, which he had continued for feveral hours without fuccess. I was fent for about one o'clock in the morning, and on examination found the arm pushed through the external parts, the shoulder pressing firmly upon the perinæum. The exertions of the mother were wonderfully ftrong. I fat down while she had two pains, by the latter of which, the child was doubled and the breech expelled. I extracted the shoulders and head, and left the child in the bed. Mr. Eustace expressed great astonishment at the sudden change, but I affured him that I could claim no other merit on account of this delivery, except that I had not impeded an effect which was wholly produced by the pains.

This child was also dead, but the mother recovered in the most favourable manner.

In all these cases, the women were at the full

this said in this said and that the child

The White the transfer of the more and the set of the set of the set of the transfer of the set of

with the party and the same that the

All of the language and which there is not

Mag.

full period of utero-gestation, and the children were of the usual size.

Many other cases of the same kind have occurred to me, and with the histories of several, varying in the time or manner in which the evolution of the child was made, I have lately been favoured by gentlemen of eminence in the profession, and many others have been published, in different countries. But these are sufficient to prove the sact, that in cases in which children present with the arm, women would not necessarily die undelivered, though they were not assisted by art.

With respect to the benefit we can, in practice, derive from the knowledge of this fact, I may be permitted to repeat, that the custom of turning and delivering by the feet in presentations of the arm, will remain necessary and proper, in all cases, in which the operation can be performed with safety to the mother, or give a chance of preserving the life of the child. But when the child is dead, and when we have no other view but merely to extract the child, to remove the danger

tall periods of area-earlierers will the title-

Minny other rates of the shift of him of him of the carting of the

picture of the control of the contro

danger thence arifing to the mother, it is of great importance to know the child may be turned spotaneously, by the action of the uterus. If we avail ourselves of that knowledge, the pain and danger which fometimes attend the operation of turning a child may be avoided. Nor would any person, fixing upon a case of preternatural presentation, in which he might expect the child to be turned spontaneously, be involved in difficulty, if, from a defect of the pains, or any other cause, he should be disappointed in his expectations. Nor would the fuffering, or chance of danger to the patient be increased by such proceeding, as the usual methods of extracting the child could, under any fuch circumstances, be fafely and fuccefsfully practifed.

TOTAL PROPERTY OF THE PROPERTY OF THE PARTY the state of the s

CLASS IV. Anomalous or Complex Labours.

FOUR ORDERS.

ORDER I.

Labours attended with Hemorrhage.]

ORDER II.

Labours attended with Convulsions.

ORDER III.

Labours with two or more Children.

ORDER IV.

Labours in which the Funis Umbilicalis prefents before the Child.

K 2

On

Times as an annual times and the same and th

es serio sunt

aligno.

of the latest of

THE PERSON

Lateres of the sold south Long several L.

THE HARRY

Laboury coint trans or only will be being transfer.

OFF STREET

And the State of t

On Labours attended with Hemorrhage.

HEMORRHAGE. A discharge of blood from the uterus, inordinate with respect to time or quantity.

VARIETIES.

- 1. In abortions.
- 2. At the full period of utero-gestation.
- 3. After the birth of the child.
- 4. After the expulsion of the placenta.

Note. No general description or character can be given to Anomalous Labours as a class, because the different orders bear no refemblance to each other. They are brought together merely to prevent the multiplication of classes.

ON ABORTIONS.

SECTION I.

- 1. With respect to the time of pregnancy, all expulsions of the fætus may be reduced under two distinctions.
- 2. In the first will be included all those which occur before the uterus is sufficiently distended to allow of any manual operation, and these may be properly called abortions.
- 3. In the fecond may be classed all those which allow of manual assistance, if required, and which are therefore to be esteemed as labours, premature or at the full time.
- 4. But no precise period of pregnancy can be fixed as a line for these distinctions.
- 5. We may, however, in general say that all expulsions of the fætus, before the end of the sixth month, are to be considered as abortions.

6. But

ON A BORELONE

.a. Welfulle

bending and was parent of the annihing as its

the design of the property of the services of

nes you as my to print to be and an analy as

and the contract of property of the confidence to

1

- 6. But all expulsions of the fætus, after the expiration of the fixth month, are to be esteemed as labours, and, if attended with the same circumstances, should be managed upon the same principles.
- 7. Yet expulsions of the fætus sometimes happen so critically, as to make it doubtful to which distinction they should be ascribed.
- 8. When manual affiftance is thought needful, the longer the time wanting to complete the full period of pregnancy, the more difficult must be any operation.

SECTION II.

On the Causes of Abortions.

- 1. The predisposing causes of abortion are; 1st, general indisposition of the constitution; 2d, infirmity of the uterus.
- 2. The general state of women who are disposed to abortion is very different, some being weak and reduced, and others plethoric.

3. Weakly

(200)

The same

A STATE OF THE STA

- 3. Weakly women become more liable to abortion, because they are susceptible of violent impressions from slight external causes.
- 4. Plethoric women are more liable to abortion, from the disposition which the vessels of the uterus have, from structure and habit, to discharge their contents.
- 5. Every action in common life has been affigned as a cause of abortion.
- 6. But it is to the excess of these actions that we are to attribute their effects, for women in health seldom abort, unless from violent external causes.

SECTION III.

On the Prevention of Abortion.

I. As every disease to which women are liable may dispose to abortion, the method instituted to prevent it, must be accommodated to the disease, or to the state of the constitution.

2. In.

mad and all and an income and an income and and

ATT MOTTON

on the Delegation of themsion

the state of the s

all of

- 2. In fome constitutions abortions may be prevented by repeated bleeding in small quantities, by antiphlogistic medicines, and sometimes by warm bathing.
- 3. In others, abortion may be prevented by nourishing and invigorating diet and medicines, by bark, by wine, especially claret, and often by cold bathing.
- 4. But it will be proper, in every case, to avoid all violent exercise, to keep the mind composed, and to rest frequently in an horizontal position.
- 5. Women feldom abort while they have the vomiting which usually attends early pregnancy.
- 6. In women who have no spontaneous vomiting, this may be excited with safety and advantage by frequently giving small doses of *Ipecacuanha*.
- 7. Pregnant women are usually costive, and abortions have been often occasioned by too great assiduity to remove this costiveness, which is a natural and proper state, in the early part of pregnancy.

SECTION

, ,

SECTION IV.

On the Signs of Abortion.

- The figns of abortion are, frequent micturition, tenesmus, pain in the back, abdomen, and groins, with a sense of weight in the region of the uterus.
- 2. But the most certain sign is, a discharge of blood, which proves that some part of the ovum is separated from the uterus.
- 3. It has been supposed when this last sign appears, that there is scarcely a possibility of the patient proceeding in her pregnancy.
- 4. But I have met with an infinite number of cases in practice, in which, notwithstanding this appearance, once or oftener, to a considerable degree, the discharge has ceased, and no ill consequences have followed.
- 5. We are therefore to persevere in the use of those means which are thought reasonable and proper, till the abortion has actually happened.

6. It

the most day

MARINE MARINE CO.

6. It is not always prudent to give a decided opinion of the probable event of those cases which may be attended with the symptoms of abortion, as their termination is very often different from what might have been expected from the symptoms.

SECTION V.

On the Treatment of Women at the Time of Abortion.

- 1. The treatment must vary according to the nature and degree of the symptoms.
- 2. There is an endless variety in the manner in which abortion takes place. Some women abort with sharp and long continued pains, others with little or no pain; some with a profuse and alarming hemorrhage, others with very little discharge. In some the ovum has been soon and perfectly expelled, in others after a long time, in small portions,

L

Car I

.

or very much decayed; but the only alarming symptom is the hemorrhage.

- 3. The hemorrhage in abortions is not always in proportion to the period of pregnancy, this being in some advanced cases very small; and in others, though very early, abundant.
- 4. The hemorrhage usually depends upon the difficulty with which the ovum may be expelled, and upon the state of the constitution of the patient naturally prone to hemorrhage.
- 5. The general principles which should guide us in the treatment of hemorrhages, from any other part of the body, are applicable to those of the uterus, regard being had to the structure of the uterus.
- 6. If the patient be plethoric, some blood should be taken from the arm at the commencement of the hemorrhage, and the saline draughts with nitre, or acids of any kind, may be given in as large a quantity, and as often as the stomach will bear.

L 2

7. Thefe

- 7. These may also be given during its continuance, and cloths wet with cold vinegar may be applied to the abdomen and loins, and renewed as they become warm. The patient should be exposed to, and suffered to breathe, the cold air.
- 8. Every application or medicine, actually or potentially cold, may be used. A large draught of cold water or ice may be given with great propriety, and it is the custom in Italy to sprinkle ice over the body of the patient if the danger of the case be imminent.
- 9. Every medicine or application which has the power of flackening the circulation of the blood, eventually becomes an aftringent, but aftringents, properly so called, can have no power in stopping hemorrhages from the uterus.
- 10. Hemorrhages are stayed by the formation of coagula, plugging up the orifices of the open blood vessels, or by the contraction of the coats of the blood vessels.
- 11. These effects are produced more favourably during a state of faintness, which, though

Thefe may alfa be given during its construction of the constructio

was blue one address.

The constant of the constant o

the name of factoring the curvivino of the blood, executably decemps on all autopa, back adving the called one save to a strangence, properly for called one save to proper in factoring factorings beautiful.

alife his

tion of the contract of the co

and the state of the product of the state, and the state of the state

though occasioned by the loss of blood, becomes a remedy in stopping hemorrhages.

- 12. Cordials are not therefore to be hastily given to those who are faint from loss of blood; unless the faintness should continue so long as to make us apprehensive for the immediate safety of the patient.
- 13. The introduction of lint or any foft substance into the vagina, has been recommended, and sometimes used with advantage, by favouring the formation of coagula.
- 14. Cold or astringent injections into the vagina have also been recommended.
- 15. Opiates have been advised in abortions attended with profuse discharges, and they may sometimes be proper to ease pain, or to quiet the patient, especially when there is a chance of preventing the abortion, or after the accident has happened.
- 16. But when there is no hope of preventing the abortion, the degree of pain proving the degree of action of the uterus, and the action of the uterus producing and favouring the contractile power of the blood vessels, if by opiates

sign decided to the second to a state of the old, being the country of the second to a state of the second to a se

nedwise fator of the patient.

1.3. The introduction of line or envelope
Labitance this the engine, in the recem-

the Cold or eliminated michigandian size

continued in a continue of the continue of the

primitive to sold one production with the control of the control o

opiates the action of the uterus should be prevented or checked, they may contribute to the continuance of the hemorrhage.

17. Hemorrhages in abortions, independent of other complaints, though very alarming, are not dangerous.

18. But if women abort in consequence of acute diseases, there will be very great danger.

19. For they abort because they are already in great danger, and the danger is increased and accelerated by the abortion.

20. The ovum has been fometimes retained in the uterus for many months after the symptoms of abortion had appeared, and when it had lost the principle of increasing.

or proper in abortions, to use any means for bringing away the ovum, or any portion of it which may be retained.

constant and a first point of the sea to depute A RESIDENCE TO THE PARTY OF THE desired to see the second

SECTION VI.

On Hemorrhages at the full Period of Uterogestation.

- I. Under this fection will be included all those hemorrhages which may happen in the three last months of pregnancy.
- 2. These are occasioned first by the attachment of the placenta over the os uteri; secondly, by the separation of a part, or of the whole placenta, which had been attached to some other part of the uterus.
- 3. Hemorrhages arising from the first cause are more dangerous than from the second; but those from the second have sometimes proved fatal.
 - 4. The danger attending hemorrhages is to be estimated from a consideration of the general state of the patient, of their cause, of the quantity of blood discharged, and of the effect of the loss of blood, which will vary in different constitutions.

5. Hemorrhages

AND THE PARTY OF T

- 5. Hemorrhages are infinitely more dangerous with sudden than with slow discharges of blood, even though the quantity lost may be equal.
- 6. The danger arising from hemorrhages is indicated by the weakness or quickness of the pulse, or by its becoming imperceptible, by the paleness of the lips, and a ghastly countenance, by inquietude, by continued fainting, by a high and laborious respiration, and by convulsions.
- 7. The two last symptoms are usually mortal, though when women are extremely reduced, they are liable to hysteric affections of a similar kind, that are not dangerous.
- 8. The vomiting which generally follows violent hemorrhages indicates the injury which the constitution has sustained by the loss of blood, but by the action of vomiting the patient is always relieved, and it contributes to the suppression of hemorrhages.
- 9. Near the full period of utero-gestation, women are always in greater danger in those hemorrhages

1 of electronic of presentations Philippites seithber beiter beiter

hemorrhages which are not accompanied with pain.

of the uterus, and this proving that the strength of the constitution is not exhausted, the danger in hemorrhages may often be estimated by the absence or degree of pain.

SECTION VII.

On those Hemorrhages which are occasioned by the Attachment of the Placenta over the Os Uteri.

- be distinguished from the membranes as soon as the os uteri is a little opened, be attached over the os uteri, the woman usually goes through the early part of pregnancy without any inconvenience, or symptom which denotes the circumstance.
- 2. But when the changes previous to labour come on, there must be an hemorrhage,

8 because

because a separation of a part of the placenta is thereby occasioned, and as the disposition to labour advanceth, the hemorrhage is generally, though not universally, increased.

- 3. With this circumstance very slight external causes are also apt to occasion hemorrhage.
- 4. When a hemorrhage from this cause has once come on, the patient is never free from danger till she is delivered.
- 5. The powers of the constitution are undermined by hemorrhages profuse or often returning, so that no efforts, or only very seeble and insufficient ones, are commonly made for the expulsion of the child.
- 6. We are therefore often obliged to free the patient from the imminent danger she is in by artificial delivery.
- 7. Of the propriety of this delivery, in cases of dangerous hemorrhage, there is no doubt, or can be any dispute, except as to the precise time when the patient ought to be delivered.
 - 8. On the first appearance of the hemorr-

with the state of the state of

hage, unless it be prodigious in quantity or unusually terrifying in its effect, it is seldom either requisite or proper to attempt to deliver by art.

- 9. Nor does it often happen that a fecond or a third return of the discharge compel us to the delivery by art.
- cannot be fecure till she is delivered, and as the delivery is seldom completed by the natural efforts, and as the artificial delivery, though performed before it be absolutely necessary, is not dangerous, if performed with care, we must be on our guard not to delay the delivery too long.
- thought eligible to deliver on account of the hemorrhage, the parts are so unyielding as not to allow of the operation itself without some hazard.
- 12. Yet when the parts requiring dilatation make no resistance to the passage of the hand, the event of the operation is always more pre-

M 2

carious,

Parasi undelli di din produgnam da quantita or nondondiy terajiga in in effect, it is foldom condondiy terajiga in in effect, it is foldom collere require an arrange en delirer

the same and the state of the same of the

to sine delevery his are,

enticlimates of the said that the said and said the said and said the said

work nor street

ed to mission to make a minimum of any of a minimum of the instance of the minimum of the policy of the minimum of the instance of the instanc

,

Louis Bernal

And the state of the parameter of the bank and the bank, being a state bank, being a state bank, being a state bank, being a state bank a state bank

carious, the operation having been deferred too long.

- 13. But though it may be proper in some cases to determine on immediate delivery, the operation must always be performed with the utmost deliberation.
- 14. The first part of the operation has been described under preternatural presentations.
- attached over the os uteri, it is of little confequence whether we perforate the placenta with our fingers, or separate it on one side till we come to the edge, though the latter is generally preferable.
- 16. If the hand be passed through the placenta, we shall come directly to the part of the child which presents.
- 17. But if we separate the placenta to the edge, the hand will be on the outside of the membranes, which must be ruptured before we lay hold of the feet of the child.
 - 18. No regard is to be paid to the part of the

carious, the operation having been deferred teo long.

eq. But abough it may be prepar in fome and the to determine on immediate delivery, the spectament with the attention and always he performed with the attention deliveration.

to The first part of the openium has been described under preterment pretentations.

is go. When the hand is carried to the placental attached over the as sirely, it is in the leader on-frequence whether we performs the placental in with our fingura. Or framero is on one this sall we come to the edge, though the larger is yearenly preferred as

refer of the band he pailed through thought in the case, we that oddered refer to the child tokich undone.

edges, the board will be on the placents so the cityes, the board will be on the confidence of the manner before the beauty which are the children of the children beauty.

to rich with the files and or at the part off . The

the child which may present, as it must always be delivered by the feet.

- 19. The feet of the child being brought flowly into the pelvis, we must wait till the uterus is contracted to the body of the child, which will be indicated by pain, and known by the application of our hand to the abdomen.
- 20. The delivery must then be finished very slowly, to give the uterus time to contract as the child is withdrawn from its cavity; but this part of the operation has likewise been described under preternatural prefentations.
- 21. An affistant-should make a moderate pressure upon the abdomen during the operation, to aid the contraction of the uterus, and to prevent ill consequences from the sudden emptying of the abdomen.
- 22. When the child is born, the hemorrhage will be generally stayed, if the operations has been performed slowly.
 - 23. But if the hemorrhage should continue or

the schild which may protein, as it mate al-

ign The destroiting child being intenglic should being intenglic should be not the second with the life the solid, and the solid, and the solid, while the helicated by print and answer by the the application of our head to the one-steer.

produce to said the condition of a second product of the condition of the

Company of the particular state of the separate of the separat

AND THE RESIDENCE OF THE PARTY OF THE PARTY

or return, the placenta is to be managed as will be afterwards directed.

- 24. Should no uncommon difficulty attend the delivery, children will be often born living in cases of hemorrhage which are attended with the utmost danger to the mother; or, as it has sometimes happened, after the death of the mother.
- 25. Before, during, or after delivery in cases of hemorrhage, the means and applications before recommended, may be occationally used with advantage.

SECTION VII.

On those Hemorrhages which are occasioned by the Separation of a Part, or of the whole Placenta, before or in the Time of Labour.

- 1. Hemorrhages arising from this cause are feldom so alarming or dangerous as the preceding.
 - 2. But if the separation of the placenta be sudden

(30)

to began in it is a service of mineged as

sac Sheeds no unconduct his after his such in a standing she the delivery bettering the delivery bettering the cases to see a standard of a second delivery on as a standard of a second delivery on as the standard of the second delivery of the second de

nit was the man of guitain and the 22.

-saidga man agreement of guitain and the effect of the man of the man of the color and t

SECTION THE P

the second of th

ers dim to a special garden garden send or mile et al.

and standing the particular and when

fudden and extensive, the danger may be equal, and the same mode of proceeding required.

- 3. Our conduct must be guided by a confideration of the degree and effect of the hemorrhage, and of the state of the labour when it occurs.
- 4. Should the hemorrhage from this cause occur in the first period of labour, the action of the uterus will be weakened, but it may be sufficient to dilate the os uteri.
- 5. If the quantity of blood lost in these cases be very considerable when the os uteri is sufficiently dilated, the greater the degree the better, the membranes containing the waters may be ruptured.
- 6. By the discharge of the waters the distention of the uterus will be lessened, and by the consequent contraction, the size of the vessels being diminished, the hemorrhage will of course be abated or removed.
- 7. After the abatement or suppression of the hemorrhage, the action of the uterus will become stronger, so that the delivery will, in general,

fudden and testanfive, the danger may be county and the labe mode of proceeding re-

the control with the paided by a confine control of the degree and allied of the his hisecontrol of the degree and allied of the his winess

come and the homest of them the section of the sect

The transport of the state of t

b. Its the obligates of the waters the difference of the water of the section of

in malifer apput the amended to the A or of the action of

general, be then completed without further affiftance.

- 8. But if the hemorrhage should continue after the discharge of the waters in such a degree as to threaten danger; or if it should commence in the second period of the labour, the interposition on our part must vary according to the circumstances, and chiefly according to the situation of the child.
- 9. It may in some cases be necessary to deliver by art as in the preceding section, and in others to deliver with the forceps or vectis, if the hemorrhage be profuse, and we despair of the child being expelled by the natural efforts.
- 10. The proper management of all such cases may be collected from what will be generally said on the subject, being always on our guard to distinguish between sear and real danger.

afficience.

de flue if the hearthage finder de la soll of a la continue of the difference of the soll of the land of the land

p. It may in tests taken be according to decide in from by art as in the mencering feether, and in extreme with the respective and of decides in the hemotrings be profined and of decides of the instance of

the tree of the contract of th

SECTION VIII.

On those Hemorrhages which occur when the Placenta is retained after the Birth of the Child.

- the spontaneous action of the uterus in a short time after the birth of the child.
- 2. But sometimes the placenta is retained, 1st, from the inaction or insufficient action of the uterus; 2d, by the irregular action of the uterus; 3d, by the scirrhous adhesion of the placenta to the uterus.
- 3. Sometimes there is a profuse discharge of blood, when no action is exerted by the uterus to expel the placenta.
- 4. Whenever there is a hemorrhage, the whole or a portion of the placenta must have been previously separated, and the hemorrhage usually continues, or returns till the placenta is expelled or extracted out of the cavity of the uterus.

THEF WORTZON

of the state of the second state of the second

the plantage of penerally regular lies

the form the investion or emperators without of the property of the interest of the property o

appealate Salvey e si acide criss demail e. Calvel demana el noille de criste de al la

consideration of the state of t

1100

FORTING

SECTION IX.

On the Retention of the Placenta from the Inaction or insufficient Action of the Uterus.

- I. Though the placenta be retained after the birth of the child, if there be no hemorrhage, we are to wait, without any interpofition on our part, in expectation of the action of the uterus.
- 2. The time which it may be proper and expedient to wait will depend upon the state of the patient, and the state of the patient generally depends upon the previous circumstances of the labour.
- 3. But no patient ought to be left before the placenta is brought away, because a dangerous hemorrhage may at any time come on.
- 4. When the patient complains of pain, the expulsion of the placenta may be safely forwarded, by aiding the contraction of the uterus by moderate pressure with the hand

N

upon

MI MOLTON

On the Report of the Property of the action of the United States

i. If in each the concern to religion of the confirm of the confir

The fine to wait to distinct the proper and conceined the called the fine of the callent, and the fine the callent, and the fine the provide the private depends of the highest the provide the callent.

grades administ and the second control beddle and the places and the product and the places and

upon the abdomen, and by pulling gently by the funis.

- 5. But if the first pain, with the aid we think it prudent to give, should not bring down the placenta, we are to wait for a return of the pains, proceeding in the same cautious manner.
- 6. When that part of the placenta into which the funis is inserted can be felt, little danger or difficulty is to be apprehended, and we are to extract it slowly.
- 7. But if a hemorrhage was to come on, the placenta being retained, it would be equally necessary to extract the placenta as it would be to extract the child, provided the degree of hemorrhage was equally profuse or sudden.
- 8. After the birth of the child, the extraction of the placenta is therefore to be confidered as the only method by which an apprehended or present hemorrhage is to be prevented or avoided.
- 9. Yet all discharges of blood do not require a speedy extraction of the placenta, but

 N 2 such

fuch only as by their violence or continuance, or frequent returns, threaten danger.

- 10. If much force be used in pulling by the funis, there will be danger; 1st, of tearing it from the placenta; 2d, of inverting the uterus; 3d, of injuring the uterus by the violence; 4th, of increasing the hemorrhage.
- 11. The danger of these consequences is greater when force is used to extract the placenta by the funis, than by the prudent introduction of the hand into the uterus for that purpose.
- will fometimes be able to bring down the placenta in cases in which the uterus acts insufficiently, just using so much force as will prevent the retrocession of it in the act of inspiration.
- may be brought into the pelvis, it should be suffered to remain there till the action of the uterus comes on, or so long as there is reason to fear a return of the hemorrhage, and it must then be carefully withdrawn.

SECTION.

and successful to the same and produces the self-sent and the service and the state of t and the first spirit the part of the second manners and the state of the second second

SECTION X.

On the Retention of the Placenta from the irregular Action of the Uterus.

- with equivalent force, and at the same time, the combined power will contribute to the expulsion of whatever is contained in its cavity.
- 2. But if the uterus should act irregularly, the contrary effect might be produced.
- 3. If the fundus uteri should not act when the other parts are in action, the longitudinal contraction of the uterus would be produced; but if the central parts should only act, the uterus would then be contracted in the form of an hour glass.
- 4. As the placenta cannot be excluded when the uterus acts in this irregular manner, it must be extracted by introducing the hand into the uterus, provided the state of the hemorrhage should require it, or when it can-

et matematical reported to deliver the state of the s

not be extracted by using the means before mentioned.

- 5. The hand ought never to be introduced into the uterus except in cases of real necessity, and then with the utmost care; and the hand when introduced should not be withdrawn until the placenta is detached and brought into the pelvis.
- 6. If the whole placenta be loosened this is easily effected, but if a portion of it should be found adhering, this must be separated by bending it back from the uterus, or by passing gently the singers between it and the uterus.
- 7. When the uterus is found contracted in the form of an hour-glass, the contracted part must be dilated in the manner recommended for the dilatation of the os uteri, and the contracted part must be amply dilated, or it will immediately contract again round the wrist.
- 8. We must then proceed as is before ad-

Agency had the attempt to transport a second or a A PARTY OF THE PAR AND THE PARTY OF T many and the second THE THE PARTY TO SEE TH

SECTION XI.

On the Retention of the Placenta from the scirrhous Adhesion of it to the Uterus.

- 1. Should there be a degree of hemorrhage fufficient to make it necessary to introduce the hand to extract the placenta, a part of it must be separated, though there may be a scirrhous adhesion of the remainder to the uterus.
- 2. Then the method advised in the last fection must be put in practice, and the sirmer the adhesion the slower the separation ought to be made.
- 3. But if there should be no hemorrhage of importance, and merely a retention of the placenta beyond its due time, we may say, for example, more than four hours, and the means before recommended are insufficient to bring down the placenta;
- 4. It may then be necessary to introduce the hand carefully to separate and extract the placenta,

The state of the s The state of the s placenta, and the difficulty will not be increafed by the delay.

- 5. Following the navel string as our guide, we must then pass the hand to the placenta; and if it should be found wholly adhering, we must begin with great caution to separate at the edge, and gradually proceed as before directed until the separation is completed.
- 6. Then grasping the placenta, we must slowly withdraw our hand, that the uterus may contract accordingly, and the chance of a subsequent hemorrhage be prevented.
- 7. The irritation made by the introduction of the hand, will generally occasion a return of the action of the uterus, before dormant, that will greatly facilitate the separation.
- 8. Yet it is possible that a portion of the placenta may adhere so firmly as to make it unsafe to separate it with our singers.
- 9. Should this circumstance occur notwithstanding the most deliberate and firm proceeding, it may sometimes be more justifiable to leave the adhering part remaining than to use violence in separating it.

6. 20- 11 Manufacture and the second of when the greater portion of placenta is brought away, it is always a defirable thing to bring away the placenta and membranes in a perfect state.

SECTION XII.

On those Hemorrhages which follow the Expulsion or Extraction of the Placenta.

- I. The hemorrhage in these cases may be either a continuation of that which existed before the exclusion of the placenta, or it may only follow the exclusion of the placenta.
- 2. When it is of the former kind, we may presume that it was not within our power to prevent it, but the latter kind may often be attributed to the violence or hurry with which the placenta has been extracted.
- 3. This is not fo dangerous as either of the varieties of hemorrhage of which we have last spoken, though with imprudent manage-

ment,

no. But thingly bearing our allocate in branching when the greater portion of placesta in branching to descript a stay, it is always a dedirable thing to descript away the place on and members a line perfect.

TIK MOLYCE

On their Minnestern was reliable for the figure of the first and the first of the f

as Valuen in it of the Colored kinds will many property of the property of the colored with which will be the property of the colored to the violence of the colored to the places of the places of the been extracted.

evidenteleide le egalementele e electrici ede Enedenteleide le egalementele e electrici ede Enedenteleide icin d'agants prodogli, fiel ment, or under particular circumstances, it has sometimes proved fatal.

- 4. All the cautions given with respect to the management of the placenta, relate to the prevention of this kind of hemorrhage.
- 5. When the strength of women is much reduced by any cause which existed previous to labour, or when they have gone through much fatigue in the course of it, there is usually great heat and a rapid circulation of the blood at the time of delivery.
- 6. While they are in this fituation, if the placenta were to be brought away hastily, an extraordinary quantity of blood must of necessity be discharged.
- 7. The interval of time which passeth between the birth of the child and the expulsion of the placenta, should therefore be employed in cooling the patient and recovering her from her fatigue.
- 8. Even when the placenta is excluded out of the cavity of the uterus, it should be fuffered

ment and analyze the state of the state of the

Tuffered to remain there till all tumult is quieted, and then with the membranes, flowly extracted.

- 9. The quantity of blood discharged in consequence of the separation of the placenta will vary in different women, or in the same women at different labours, independently of the manner in which the placenta may come away.
- ro. The less the quantity of blood discharged the better women in general recover, provided there be no morbid cause of its diminution.
- great discharge of blood after the separation of the placenta, whatever care may be taken in extracting it.
- 12. This may often be prevented by keeping the patient out of bed till the membranes are broken and the waters discharged to the very moment of the child being born.
- 13. In all cases of dangerous hemorrhage after the extraction of the placenta, it is first O 2 necessary

T. John

the state of the s

The state of the s

to the state of the second state of the second

necessary that we should be assured, by an examination per vaginam, that the uterus is not inverted.

- 14. Should there be an alarming hemorrhage after the separation and exclusion of the placenta, notwithstanding all the care which can be taken according to the methods before mentioned,
- 15. The doctrine of hemorrhages before given, and the general treatment already recommended, will enable you to fix upon the line of conduct it will be expedient to purfue, and to restrain or suppress them as far as they are under the influence of art.
- 16. In cases of hemorrhage so very profuse as to occasion frightful faintings, continuing so long as to raise great solicitude for the immediate safety of the patient, it was generally said, that cordials ought not to be given.
- 17. But this requires explanation. When the patient has continued faint so long as to give time, according to our judgment, for the wessels of the uterus to contract, then cordials

the sale of the sa

the Should exceed the an distinction bearing to the configuration of the

stated department to entire of Very and have department of Very and have department of the department of the department of the state of the department of the state of the sta

to eath of a common to eath of the state of

estable and the continues of the continu

dials and nourishment in small quantities, very often repeated, are really needful.

- 18. Other means are also to be used for the purpose of recovering women from this long continued fainting; and one of the most effectual is, sprinkling the face freely with cold water.
- 19. After a profuse hemorrhage the patient will frequently have a disposition to steep, which has generally been considered as dangerous.
- 20. But short sleeps are very refreshing, though long ones in a very weak state are, under every circumstance, found to be injurious.
- 21. When there has been a dangerous hemorrhage, the patient should remain for many
 hours undisturbed, and in an horizontal position; and our attention must be continued aslong as any danger is to be apprehended.

(1001)

Carlo and the second in final questions beauty

the fact of or old surround and control of the fact of

english egales and elike hange that grant and a seek hours that self-reference are reached all political and there exists has grant all marris and this engage.

And same part of a product of a part of the contract of the co

On Labours attended with Convulsions.

- nancy very much refemble the epilepsy, but to the symptoms, which these have in common, may be added, the peculiar hisping noise which women make with their lips during the convulsions.
- 2. When convultions happen to women with child, they are generally, but not univerfally, accompanied or followed with fymptoms of labour.
- 3. These convulsions are indicated by a piercing pain in the head, by giddiness and other vertiginous complaints, by blindness, by vacillation of the mind or a slight delirium, by violent cramp or pain at the stomach, by a fulness or apparent strangulation of the neck and fauces, and other affections of the vascular and nervous system.
 - 4. The means to be used for the preven-

Sempliance of the Control of the Con

Annous Tallespan Indiana, in head In-

CONTRACTOR OF THE STATE OF THE

tion or cure of convulsions when threatened or existing, must be regulated according to the constitution of the patient and the violence of the symptoms.

- 5. In general it will be necessary to take away some blood, or sometimes to repeat the bleeding, and it has been sound particularly serviceable to open the jugular vein. Emetics, when they could be given, have been useful, as has also the warm bath. Clysters may be frequently exhibited. Opiates, joined with nervous medicines, may be given; and the patient is, by all the means in our power, to be soothed and restrained. From violent exertions.
- 6. During the convulsions the means by which contrary irritations may be excited are to be used; and of these the most powerful is, the dashing of cold water in the face, which has been known to prevent, or even to cure, convulsions.
- 7. Some writers have recommended the fpeedy delivery of the patient, as the most eligible,

Construct and another the sup or not see that the second of the second o

and the court of the control of the

Sent de la company de la compa

eligible, and only effectual method of removing puerperal convulsions; but others have infisted that the labour should be uninterrupted.

- 8. From the histories of all the cases of puerperal convulsions which have been recorded, it appears, that a greater number have died of those who were delivered by art, than when the labours were resigned to nature.
 - 9. As far as my experience enables me to judge, we ought not to attempt to deliver women with convulsion's before some progress is made in the labour.
- Tufficiently, or to a certain degree, the patient fafely may, and ought to be delivered by art, if from the urgency of the convultions, and the general danger of the case, delivery should appear necessary.
- these cases, whether the operation be performed with the forceps or vectis, or by turn-

E 801

man although and although and and and a

ACTUAL DE

already been fully explained.

12. The event of the operation, both to the mother and child, will also very much depend upon the skill and circumspection with which it may be performed.


THE END.

Published by Dr. DENMAN,

- Y. An Introduction to the Practice of Midwifery.
 Vol. I. 8vo. Price 6s. in Boards.
- 2. An Essay on Preternatural Labour, 2s.
- 3. An Essay on Hæmorrhages, 2s.
- Price 2s. each.
- Printed for J. Johnson, No. 72, St. Paul's Church Yard.







264)102





