A treatise on medical sympathy, and on the balance and connection of the extreme vessels of the human body : comprehending, I. The nature of sympathy in general; ... II. The nature of febrile sympathy, ... / by Seguin Henry Jackson, M.D.

Contributors

Jackson, Seguin Henry, 1752-1816. Howell, John, 1777-1857 Bristol Royal Infirmary. Library University of Bristol. Library

Publication/Creation

London : Printed for J. Robson and W. Clarke, booksellers, 1787.

Persistent URL

https://wellcomecollection.org/works/sm62xs64

Provider

Special Collections of the University of Bristol Library

License and attribution

This material has been provided by This material has been provided by University of Bristol Library. The original may be consulted at University of Bristol Library. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



100 151293 56 86 PRESENTED BY Dr. John Howell to the Library of the BRISTOL INFURMARY. Rest. Med. 18





TREATISV

ON

A

A FOR STATION

MEDICAL SYMPATHY

AND ON THE

BALANCE AND CONNECTION

TH

EXTREME VESSELS OFTIE UMAN BODY:

COMPREHENDING,

- I. The Nature of Sympathy in general; that of Antipathy; and the Force of Imagination; and their extensive Importance and Relation to the Animal Economy: With fome of Mr. JOHN HUNTER's interesting Observations on partial and universal Sympathy.
- II. The Nature of Febrile Sympathy, and CONSENT OF Ex-TREME VESSELS, illustrated by practical Remarks; and a new Explanation of the various Affections of the Stomach and Skin, in Fevers: In which is attempted a full Refutation of the Doctrine delivered by Dr. CULLEN on the fame Subject, from the Practical Chair at the celebrated University of Edinburgh.

BY SEGUIN HENRY JACKSON, M.D.


Member of the Royal College of Phyficians, London, and of the Royal Medical Society, Edinburgh ; and Phyfician to the Weilminster General Dispensary, and to the Infirmary for the Poor of Saint George's, Hanover-Square.

SECOND EDITION.

LONDON:

Printed for J. ROESON and W. CLARKE, Bookfellers, New Bond-Street.

M DCC LXXXVII.



THIS TREATISE

IS RESPECTFULLY INSCRIBED

TO HIS BRETHREN,

THE MEMBERS OF THE ROYAL MEDICAL SOCIETY OF EDINBURGH.

AS A GRATEFULL TESTIMONY OF THE MANY ADVANTAGES WHICH HE DERIVED FROM THE INSTITUTION, WHEN RESIDENT AMONG THEM.

BYTHEIR

DEVOTED

FRIEND,

AND OBLIGED

HUMBLE

SERVANT,

THE AUTHOR.

Digitized by the Internet Archive in 2015

LTT OF TOINEURGI

THISTREATISE

HTERETHOT

https://archive.org/details/b21441996

To the R E A D E R.

Hanover-Street, Hanover-Square, April 12th, 1787.

AFTER having neglected the Publication of the following Treatife near Six Years, the Author is again induced to offer it, in its original State, to the Attention of the Medical Profession, and in particular the Student ; from a firmer Conviction of the Importance of the Subject, than when it first appeared. The Reafoning he has fometimes fuccefsfully indulged, in difficult Cafes, from his Partiality to the general Doctrine, has made him confident, that where the Art of Medicine can admit of much Improvement, it will be on the Principles laid down in the following

INTRODUCTION.

lowing Sheets, namely, that a VIS INSITA SYMPATHIÆ rules and directs, and at Times diffurbs the whole Functions of the human Body. Her Influence is indelibly flamped on our Nature, by Actions complicated—Affections wonderful—Impreffions innumerable, daily improving, strengthening, varying, and encreasing, both in Sickness and in Health. Let us then, my Brethren, as Slaves to Health, and Enemies to Difeafe, unite in the general Caufe, and by improving, on this new Principle, the Practice of Phyfic, raife the Names of its most fuccelsful Votaries to the highest Pinnacle of Fame.

The Author has read with Pleafure what Mr. J. Hunter has faid on the Subject in a late Publication, and hopes he fhall fucceed in his Attempt on fome future Occafion, to anfwer the Objections made by a chirurgical Commentator.

PREFACE.

it has ever been, to look upon every

the indulgence of a profession, liberal

PREFACE

of its younger THE author of the following sheets is too diffident of his own abilities, as well as too fenfible of the weighty objections there are to the publications of young men, upon professional subjects, not to feel a confiderable degree of anxiety for the light, in which his attempt will be looked upon: all that he has to fay is, that he does think the fubject he writes upon of importance to medicine, and that he has thought upon it long, and attentively; for the reft he throws himfelf with confidence upon the

vi PREFACE.

the indulgence of a profession, liberal in the most extensive sense of the word, and the characteristic of which it has ever been, to look upon every effort of its younger members with complacency, and appreciate them with candour.

Indeed, there is a poffibility of one event happening, which could entitle him to praife, and this would be, if the honorable ambition of diftinguishing himfelf in the profession he has chosen (however premature the attempt it has given birth to) should at length provoke Dr. Cullen to the combat, which he has hitherto declined; declined, it is to be feared, really out of com-

PREFACE. vii

compafiion to his antagonift, but not without circumftances, which have contributed to raife his prefumption.

The author cannot finish without acknowledging his, and his readers' obligations to Mr. John Hunter, for the very liberal manner in which he gave him leave to make use of his observations upon the subject of ME-DICAL SYMPATHY.

April 9th, 1781.

PART THE FIRST.

ON the nature of Sympathy in general; that of Antipathy, and the force of Imagination; and their extensive importance and relation to the animal æconomy: with many interesting observations on medical Sympathy.

CHAP. I.

On the extensive relation of Sympathy to the animal aconomy.

Page. SECT. I. Sympathy, the first and last principle of animal life - - - 3 II. Sympathy, when mental, and when corporeal - - 6 SECT. III.

SECT. III. Sympathy, when inherent, and when acquired, - - 12 IV. How sympathy prevents and cures diseases, 14

II. Difeafed S.H. A. P. A. H. D tial, and how

Sympathy which attends the healthful state of the system.

SECT. I. Sympathy, one of the most extensive principles in the animal acconomy, 18 II. Sympathy, when similar, and when diffimilar, 22 III. How Sympathy is varied in different animals. 25 IV. Sympathy least varied in the most simple V. Sympathy most strongly marked by affections of the stomach, _____ 29 VI. Mental Sympathy often connected with fimple life, and how, venoathy, cohen the p

СНАР.

CHAP.

ix

CHAP. III.

On the Sympathies which attend diseases. SECT. I. Inflammation when Sympathetic and when specific, 35 II. Difeafed Sympathy, how partial, and how universal, III. 41 -----, when particularly partial, 44 IV. _____, when in a special manner universal, 47 V. Universal Sympathy, when immediate, nd when secondary, 55

CHAP. IV.

On the uses of Sympathy.

SECT. I. Natural Sympathies most evidently useful, - - 63
II. Diseased Sympathies less evidently useful, 67
III. What Sympathies may be applied to medical use, and how, - 72
IV. Sympathy, when the province of the physical cian, and when of the surgeon, - 77

CHAP.

XV

deferent ani-

CHAP.

CONTENTS. CHAP. V.

On the fympathetic operation of medicines. SECT. I. Effects of medicines, when local, and when remote, - 78 H. Sympathetic effects of aqueous applications to the fkin, - 83 HI. How unctuous and volatile applications by Sympathy increase motion, - 86 IV. How unctuous and volatile applications procure resolution, - 88

CHAP. VI.

the University of E.

On the Sympathies of the senses, and force of imagination.

SECT. I. On the Sympathies produced by the passions and affections of the mind, - 94
II. On the sympathetic affections of the external fenses, - 103
III. On the secret effects of Sympathy and Antipathy on the child in the womb, - 106
IV. Sympathy sometimes disorders, and throws into confusion, the aconomy of human life, 110.

PART

PART THE SECOND.

xiz

d force of

in of imman tite, IIO

PART

On febrile Sympathy and Confent, and on the balance and connection of extreme veffels, illustrated by practical remarks, and a new explanation of the various affections of the stomach and skin in fever; in which is attempted a full refutation of the doctrine delivered on the same subject from the practical chair at the University of Edinburgh.

CHAP. I.

CHAP. VI.

A concife view of the theory of fever. SECT. I. The author's original letter to Dr. Cullen, with the Doctor's anfwer to it, 119 II. Of the phænomena of fever, - 126. III. Of the proximate caufe of fever, - 137 IV. Of the remote caufes of fever, - 139

CHAP,

CHAP. II.

Of the various febrile affections of the Romach.

SECT. I. Of the physiology of the stomach, 144
II. Of anorexia, as a febrile affection of the
Stomach, 149
III. Of nausea, as ditto, 155
IV. Of vomiting, as ditto, 159

C H A P. III.

On the Sympathy and Consent between the stomach and the skin in fever.

- SECT. I. How the Sympathy and Confent depends upon the balance and connection of extreme veffels, and not upon the condition of the muscular fibres of the stomach, - 173
- 11. How the balance and connection of the extreme veffels, giving the confent between the ftomach and skin, may be illustrated by the operation of medicines, as well as by the general phænomena of fever, - 179

CHAP.

C H A P. IV.

How the author's doctrine is illustrated by the fame arguments, which Dr. Cullen has made use of to illustrate his own.

SECT. I. Remarks on the fact related by Dr. Sydenham concerning the plague, 189
II. On the ceffation of vomiting at the coming on of the hot flage of fever, and very certainly on any fweat appearing, - 200
III. On the effects of vomits in bringing on the hot flage of fever, - 205
IV. On the effects of cold water taken into the flomach, - 214
On the effect of cold applied externally to the body, - 172

CHAP. V.

Some opinions are delivered in a concife point of view, as a fummary to the whole.

SECT. I. Some general conclusions, respecting the febrile affections of the stomach, by way of corollaries, - - 220 II. The

CONTENTS. i.i

II. The author attempts to give a new fummary of the pathology of the fever, - 223
III. Some promifed obfervations on the terms Sympathy and Confent, - 226
Conclusion, - 231

ATREA-

CONTENTS. 1.1 II. The author accompts to give a new fummary of the pathology of the fever, - 223 III. Some promised cifetuations up the terms

TREATISE ON SYMPATHY.

A

and the second second second


PART THE FIRST.

On the Nature of SYMPATHY in general; that of ANTIPATHY; and the force of IMAGINA-TION; and on their extensive Importance and Relation to the ANIMAL ŒCONOMY.

With many interefting Obfervations

0 N

MEDICAL SYMPATHY.



ON THE

NATURE OF SYMPATHY IN GENERAL, &c.

CHAP. I.

On the extensive relation of sympathy to the animal acconomy.

SECT. I.

Sympathy, the first and last principle of animal life.

I.

GENERAL doctrines are premifed in MEDICINE, that we may rationally eftablish a systematical mode of preferving health, and of preventing and curing difeases, and they are called its Institutions.

II.

Hitherto authors and teachers of medicine have delivered the inftitutions under three A 2 general

general heads or divisions; to wit, the following.

1ft. The doctrine of life and health, or phyfiology.

2d. The doctrine of difeafes, or pathology.

3d. The doctrine of the materia medica, or means used in the practice of physic *.

I shall take the liberty of adding a fourth, very lately discovered, but of sufficient importance to claim a place and engage our attention; namely,

4th. The doctrine of reftoring animation, or the vital principle, when apparently loft. Or, * * * * * *.

III.

Sympathy is concerned in each of these doctrines (II. 1, 2, 3, 4.). Life and health exist by it; the theory of pain and diseafe is often built upon it; relief is frequently

* Therapeica.

obtained

5

obtained from medicines, by their fympathetic operation; and though I am unacquainted with the particular doctrine at prefent inculcated by the ingenious Dr. Hawes (one of the inftitutors of the Humane Society) on the recovery of perfons apparently dead, I have not a doubt in my own mind, but that fympathy is the furviving principle in the animal œconomy, through which the means of reftoring life fucceed, applications being generally made to the ftomach, which is the feat and fountain of fympathy.

IV.

Sympathy even gave us life; the breathed into us when born into the world; the preferves our lives while in it, guards us againft the difeafes of it, proves fatal to us when in excefs, and when life has not been too long *apparently* extinct, is capable of reftoring us the world again.

V.

Sympathy does not belong more particularly to the nervous fystem, than to other folids in the body. If it be attached in par-A 3 icula

ticular to the moving extremities of the nerves, commonly called mufcular fibres, independent of the nerves themfelves, (Dr. Cullen has delivered it as his opinion that they are appendages to the nervous fyftem) we muft then confider it as a principle of fimple life, or in itfelf the *living principle*. On the contrary, if we admit Dr. Cullen's idea of their nervous connection to be well founded, and the nerves to be mutually concerned, we muft then lay it down as a principle belonging to the nervous fyftem; or of itfelf the *fenfitive principle*.

VI.

Sympathy, whether owing to a connection with the brain or not, is certainly a quality of theliving folid, and moving fibre.

CHAP. I. SECT. II.

Sympathy, when mental, and when corporeal.

VII.

Sympathy is of two kinds, mental or corporeal. The first arises from a sensation in the mind, determining to particular organs, or particular parts of the body, and raising in them.

7

them certain feelings, actions, and inclinations, fometimes agreable, and fometimes difagreable. Of this fort are longings of various kinds, depreffing paffions, &c. Thefe I would call SYMPATHIES OF CONSCIOUSNESS.

VIII.

The fecond kind depends upon the operation of external bodies, and the condition of the moving and fentient extremities of the nerves, and more generally occurs in difeafed flates of the fystem. These I would name SYMPATHIES OF IMPRESSION.

IX.

Both kinds of fympathies (VII. VIII.) are produced through the medium of the five principal fenfes; but in a more particular manner the fympathies of impression (VIII).

X.

UW SIUMO

A certain force of impression is often neceffary to produce fome of the sympathies of impression, (VIII.) Thus in some difeased states of the system pain becomes the necesfary agent.

A 4

XI. An

- aniloni Las , adoifadXI.

8

An itching is to be confidered as a leffer degree of pain, but it will not raife that fympathy, which ftronger pain in the fame part would do. However, an itching will fometimes of itfelf produce fympathetic actions.

.IIX and the condition of

The operations of fympathy are often proportioned to the ftrength or weaknefs of the parts fympathifing; and are themfelves ftronger or weaker according to the influence of cuftom and habit.

Both kinds of fy.HIX airs (VII. VIII) are

Different fympathies do not always point out a different mode of action in the caufes producing them, but a variety only in different conftitutions. The fame caufe will produce in fome ficknefs at ftomach, in others a pain of the ear, or the tooth-ach.

XIV.

batz in forme divisied

Any particular fympathy is encreafed by the mind's dwelling long upon it. Thus, for

for inftance, LOVE is a fympathy of confcioufnefs, (VII.) to which duration and attention are neceffary to give it its full effect. The mind is induced to yield to thefe from the pleafure it enjoys from them, though this is often mixed with excruciating anxiety.

XV.

When the force of an imprefiion has continued any length of time, with a correfpondent attention of the mind to the impreffion, the fympathy arifing from it will even continue long after the caufe of impreffion, originally producing it, has ceafed to act. Thus the recollection of a difagreable object, or melancholy event, will renew the imprefiion originally felt from them. The remembrance of absent love has also a similar effect.

XVI.

From length of time, the caufe having ceased to act, the impression grows weaker and weaker, until the fympathy at last ceases altogether. odin tann

XVII.

The fame or fimilar caufe of fympathy again operating, does not fo readily produce the

5

the fame impression as before. Hence, all new sympathies are, cæteris paribus, the strongest.

XVIII.

Sympathies often shew themselves, not at the part where the impression was first made; but at a distant part, where those nerves terminate that were originally impressed. Thus, an uneasy fensation has been felt at the fingers; though the cause producing it in fact existed in the substance of the brain.

XIX.

The fympathies attending the healthful ftate of the fyftem are in general agreable and pleafant. Those attendant on difeases are in general difagreable and often painful : fuch, for instance, as proceed from debility, laffitude, difficult respiration, &c.

XX.

No fympathies arife originally in the mind, without being preceded by, or mutually accompanied with, fome change in the actions and affections of fome part of the body.

body. Thus longings are produced probably by fome change of action in the organs concerned.

CHAP. I. SECT. III.

Sympathy when inherent, and when acquired.

XXI.

Sympathies are either inherent or acquired. The inherent ones are those which particularly fall under the direction of the vis confervatrix and vis medicatrix naturæ, and have commonly an evident utility. Of this kind are the fympathies between the ftomach, and the whole fystem, and between the stomach, uterus, and skin. Hunger and thirst are fympathies of this kind.

XXII.

Those may be called *acquired*, which are only brought to light by the occurrence of difease, and do not appear to answer any very apparent end or utility. I may mention, as of this kind, the sympathetic pain between the inflamed liver and shoulder; the

the uneafinefs at the glans penis, from diforders of the urinary paffages, &c.

XXIII.

The inherent fympathies (XXI.) attached to the living folid and moving fibre, remain fome time after life is apparently loft, (III.) and conftitute a vis vitæ reflauratrix.

XXIV.

The acquired fympathies (XXII.) have a given duration, they ceafe on the abatement of the morbid affection, and return not again but with a renewal of the caufe.

XXV.

-off or

Sympathy, as an inherent quality of the moving fibre and living folid, commonly takes place from changes in the affections of the brain and nervous fyftem, which excite or alter the actions of the moving fibres.

XXVI.

Sympathies are called forth, when the balance between the contraction and relaxation of moving fibres in any part is loft, in

in which cafe they conflitute what is called fpafm and convultion.

XXVII.

The fympathies of fpafm and convultion more generally occur, when the lofs of balance (XXVI.) is in any part of the circulating fyftem. By fuch violent exertions of the mufcular fyftem the balance may probably be again reftored.

XXVIII.

The great variety of fpafmodic and convulfive affections may depend upon the difference of the part where the balance in the circulation is diffurbed.

XXIX.

Sympathy (XXI.) is often an imitative faculty, fometimes involuntary, often without confcioufnefs: thus we yawn when we fee others yawn. "So the laughing of " another maketh to laugh."

XXX.

Sympathy directs us to the objects on which the gratification of our appetites and inclina-

inclinations is founded (XXI). It raifes alfo within us propenfities of the inherent kind, which have for their object the removal of fome uneafy and painful fenfation. Of this kind are yawning, ftretching, fneezing, fighing, coughing, hiccuping, vomiting, and the like,

CHAP. I. SECT. IV.

How Sympathy prevents and cures difeases.

XXXI.

Sympathy is nature's hand-maid in the conftitution and government of the animal occonomy.

The Vires confervatrices and medicatrices naturæ, and I may now add the vires vitæ reftauratrices, are every where, and at all times, directed by the influence of fympathy in the conftitution.

XXXII.

Sympathy not only affifts in the operations of those causes, which are by their nature calculated to stir up healthful actions in

in the fystem, but it also guards us against the dangerous confequences, which might arife from other causes, which from their nature, tend to diminish the due energy and activity of those functions by which alone life is supported. Hence many narcotic powers become immediately causes of ftirring up very confiderable sympathetic operations in the animal system. Such are cold, fedative passions, poisons, &c.

XXXIII.

These fympathetic operations so called forth, are always of the inherent kind (XXI). When narcotic powers very violently operate, they produce such a dangerous degree of collapse, that the loss of life is apparently threatened. It is then that the spearently plication of stimulating powers becomes neceffary, in order to fan into life the expiring flame of sympathy (III).

XXXIV.

Sympathy, and, in a more efpecial manner, the inherent, mostly relates to those parts of the system, where connections are established

eftablished between the brain, and certain parts of the body, which have a common function and conffitution. These parts, and the brain, are mutually affected either from fome original difference and alteration in the condition of the brain itfelf, or from the fame first taking place in those parts, which from relation to the brain, and their own fimilarity of organization, are more particularly difpofed to confent. The organs of voluntary motion; the primæ viæ, and more efpecially the ftomach; the heart and circulating fystem, and in particular the cutaneous extreme veffels, with those of the first paffages; the uterine fystem, confisting of its numerous external and internal appendages, are in a peculiar manner under the influence of this mutual and reflex fyinpathy, and are to be confidered as of the first importance in pathology,

XXXV.

What I have faid in XXVI, XXVII, and XXVIII, is agreeable to this mutual fympathetic dependance between parts, that have been

17

been endowed with a common function and conftitution; namely, that a fympathy(XXI.) probably exifts between the organs of voluntary motion, and the circulatory fyftem; for Dr. Cullen has obferved in his Inftitutions of Medicine CLXIV, that "The "motion of the blood in the arteries of any "particular part is promoted by the action " of adjoining mufcles."

XXXVI.

These various leading fympathies in the animal œconomy (XXXIV.) cannot be explained upon any known, or even imaginary, continuity or contiguity in the origin or courfe of the nerves belonging to the parts fo fympathizing; and in my opinion, can only be explained by fuppofing, that, from a law implanted in the animal acconomy by the Supreme Being, these parts being of the first importance to the principal functions of life, are more immediately under the influence of the vis confervatrix and vis medicatrix nature, which in all probability act by a general impression, first made on the brain, and then determined to the particular part, where the neceffary fympathy is required.

B

CHAP.

CHAP. II.

Sympathy, which attends the healthful state of the fystem.

SECT. I.

Sympathy, one of the most extensive principles in the animal æconomy.

XXXVII.

SYMPATHY is a principle in animal bodies fo connected with them ab origine, and fo attached to them even when animation has apparently ceafed, that, in the first place, hardly any actions, affections, or impressions, can be produced in any one part during life, without also producing fecondary actions, affections, and impressions, in other parts.

XXXVIII.

Secondly, for fome time after life is apparently deftroyed, it is impossible to fix any precife time, when we can fay it is abfolutely irrecoverable, and for ever lost to its former,

former, though stationary, existence. (III. XXIII.)

XXXIX.

Sympathy is one of the moft extensive principles in the animal œconomy, and may juftly be confidered as the basis of all its compound actions. (IV.) It is very easy to shew that the most perfect animals are endowed with fensation and the fimple principle of life. Out of these two arises the mind, as a third principle. These three principles then, fimple life, fensation, and the mind, which is a compound of the other two, have each their peculiar affections, which affections produce fimilar fympathies.

XL.

I have already (as in the preceding Chapter) divided fympathy into mental and corporeal, the latter of which includes the two principles of fimple life and fenfation. They in general fo ftrongly co-operate together, that it feems to be difficult, and even unnatural, to diffinguish their separate affections.

B 2

XLI. Sym-

XLI.

Sympathies arife from the readinefs of any part of the body to fall into action, without any politive caufe operating upon it, but folely in confequence of fome action, affection, or imprefision, having taken place in another part of the body. (VII. VIII. XXI. XXII.)

XLII.

Sympathy in fome inftances is an imitative fenfation in the fame body, as well as between different bodies. (XXIX.) It is imitative, in the fame body, when the fympathifer is affected in the fame manner as the part originally affected.

XLIII.

Sympathy is not an original, but a fecondary affection; it acts often as a caufe of other affections; and thefe again may become caufes of other fympathies; each of which may again produce other actions, affections, and imprefisions. So that we have thus a long fucceffion of fympathies.

XLIV.

Sympathy is greater in young than in old people; it is much lefs determinate in the young,

20

young, than in the old, which most probably arises from the parts of a young animal being more fusceptible of irritations.

XLV.

The fenfitive principle is not fo determinate in young animals as in old: thus (fays Mr. J. Hunter) if I prick my finger, the particular part of fenfation belonging to that part is fo beaten already, that no other part will fympathize: but in a child, where the fenfation is not fo determinate, every part will fympathize, and the whole body be thrown into convulfions,

XLVI.

I think that the convultions which attend the teething of infants, are alfo owing to the fame condition of the fenfitive principle (XLV.), which is not equally affected under the fame circumftances, i. e. during the fame procefs, later in life.

XLVII,

Sympathy thus depends upon the different degree of fenfibility in the individual body, and, together with its affections, becomes more or lefs confined. (XLIV.)

B 3

CHAP,

CHAP. II. SECT. II.

Sympathy, when similar, and when distimilar.

XLVIII.

Sympathy is either fimilar to its original caufe, or diffimilar: and its affections or actions may again be fimilar to those, which have already taken place in other parts, or diffimilar according to the affections of the fympathant.

XLIX.

Sympathy is fimilar, when a part fympathizes, i. e. has any affection or action, and another part, not apparently connected, has an affection or action fimilar to the other. We then fay, in the language of Mr. J. Hunter, that the fympathifer has the fame caufe as the fympathant, which had produced the original affection or action,

L.

Sympathy is diffimilar, when the affec, tion or action in the fympathifer is diffimilar to the affection or action in the fympathant,

pathant. We are then to conclude that the caufes producing the affections are different.

LI.

Upon the fame principle of diffimilarity, (L.) affection in the fympathant will often produce action in the fympathifer : and fometimes on the other hand, action in the fympathant will produce affection in the fympathifer.

LII.

To explain this: If you ftrain your ancle, or fuffer much from the dreffing of a wound, either will produce ficknefs and vomiting. The affection of the ancle from the ftrain, or wound, is either a fenfation, or an affection producing a fenfation, to wit, pain, yet this fenfation from affection produces action in the ftomach, where no morbid condition or affection can be fuppofed to be then prefent,

LIII.

Sometimes the action or affection of the fympathifer will be different from that of the fympathant. Thus fyncope has occafionally followed the voiding of coffive B 4 fæces.

fæces. Their labored expulsion is attended with a fenfation producing action from excitement, yet this fenfation caufes an affection of the fenforium from collapfe. Here the head fympathizes with the rectum, tho' the two affections are diffimilar.

LIV.

-mit out in norios

This difference of affection and of action will not in every cafe arife in the fympathifer, from a difference in the nature of the part fympathifing, but often from the nature of the ftimulant.

The affection of the Valle from the fillin, or

eicher will produce fickners and vomitive.

A ftimulant of one kind fhall produce an affection or action of the fame kind in the fympathifer; while a ftimulant of another kind fhall produce a fympathy of a different kind.

LVI.

It is even poffible that a ftimulus of the fame kind fhall produce a fympathy in one part of the body, and yet not in another, though

though the mode of action in the fimulant be the fame in both parts.

CHAP. II. SECT. III.

How sympathy is varied in different animals.

LVII.

Sympathy is fimilar, in the fame animal when the actions, affections, and fenfations are exactly the fame. It is diffimilar, when the actions, affections, and fenfations, of the part fympathifing, are different from the actions, affections, and fenfations of the part imprefied.

LVIII.

Such fympathies as arife from the affection of any one principle, as that of animal life, the fenfitive principle, or the mind, *fingly*, or from affections of two, or all of them, *combined*, if fimilar, are like the unifons of founds in mufic, or the unifons of combipations of founds.

LIX, The

inely make of an eo LIX.

The idea of a fimilar irritation or fympathy is confined either to those animals that are wholly fimilar, (if there are any fuch) or to those animals whose parts are endowed with fimilar actions and affections.

LX.

Animals have not all their original actions fimilar, or produced alike from the fame irritation or ftimulus, but from different ones; nor are all animals confined to the fame actions or affections in different parts.

LXI.

The fame ftimulus shall produce actions or affections, which shall be the cause either of a similar or diffimilar action, or of both kinds of actions, according to the variety of affections of the different parts that sympathize; so that a compound sympathy may take place.

CHAP. II. SECT. IV.

Sympathy least varied in the most simple animal.

LXII.

The moft fimple animal always admits of the moft fimple, and moft fimilar fympathy; but it is alfo probable, that fuch are not only capable of the fame fympathy, but alfo of different fympathies, (XLVIII, LI.)

LXIII,

A polypus may be confidered as an inftance of a fimple animal. It has been fuppofed to have agreeable and difagreeable fenfations or affections; as in the first place, hunger, which is a difagreeable fenfation; fecondly, the propagation of its fpecies, which is an agreeable one. But thefe will not take place without their proper caufes.

LXIV.

Too great a degree of heat, or too great a degree of cold, ftimulate much; both of them must therefore become difagreeable; but what variety of affections they must produce in a simple animal is not yet known. In

In a compound animal they may produce a variety of fympathies.

LXV.

An animal may have an agreeable feeling, yet no fenfation of it; but every part of the body of the moft fimple animal is capable of being ftimulated, and of fympathifing with another part, whatever be the ftimulus that gives the imprefiion. This (Mr. J. Hunter has alledged) will take place in the inverse proportion to perfection,

LXVI.

It has been fup-

A child in the womb has been confidered, as lefs fufceptible of ftimuli than a polypus. May it not have agreeable and difagreeable feelings, without a fenfation of them? I fufpect a child in the womb fympathizes with the mother. Hence the fatal confequence to the child of drunkennefs in the mother, of violent frights, &c. (LXII.)

if cold, ,IIVXL te manning both of

A fimple animal has the feweft fympathies, for if you compound an animal, each part

part has its peculiar part that fympathizes with it, in proportion as you make it more compounded.

CHAP. II. SECT. V.

Sympathy most strongly marked by affections of the stomach.

LXVIII.

of the thing us.

(ciotiliae)

Hunger, in the moft perfect and compounded animal, is a fenfation arifing from fympathy; that is, the ftomach fympathizes with the whole body in a flate of inanition. (XXI.) Here a kind of reflex fympathy is faid to be produced. The ftomach fympathizes with the whole body during hunger, and is thereby excited to action, which action acts again upon the body, fo as to call up other actions in the fyftem different from the former. (XLIII.)

LXIX.

When hunger has ceafed, from the acquifition of a fufficient quantity of aliment to the ftomach, this fympathy is fucceeded by heavinefs

heavinefs to fleep, as if the brain, or mind, confcious of the utility of reft to digeftion, withdrew for a while its powerful influence over the fyftem, that the function of the ftomach might at first proceed undiffurbed.

LXX.

The whole fyftem fympathizes with the ftomach varioufly, according to the nature of the ftimulus. When the ftomach is ftimulated in one way, a fweat fhall break out on different parts of the fkin; when in another way, univerfal debility fhall be produced, together with pain in the head. When the head aches, we have ficknefs at ftomach. Thefe are two different fenfations, though the nature of the ftimulus which produces each, is probably the fame. An external injury often caufes ficknefs, &c.

LXXL

Some fympathies of impression and of confciousness from external objects require a certain degree of warmth, a certain supply of nutriment, and a certain state of body to produce them. Thus a Man, in perfect

fect health and vigor, who lives well, &c. shall be fo captivated by the beauty of one woman, or the good fense and understanding of another, as to excite a *fympathy of impref fion* from the fight of the one, and *fympathy of confcioufnefs* (XIV.) from the company of the other; but, on the contrary, if he be lean, starving, &c. no such impression or affection starving in full possible of the feast of the feast is rendered fusceptible of the feast fimuli, unless the body is properly nouriss.

CHAP. II. SECT. VI.

Mental Sympathy often connected with Simple life, and how.

LXXII.

Three kinds of fympathies (XXXIX.) are fuppofed to exift, which may often be compounded; to wit, those of fimple life, of fensation, or the fensitive principle, and of the mind, each of which is not capable of producing irritations, affections, and actions, without some other irritable part through fympathy being affected in a diffimilar manner. (L. LI.)

LXXIII. A

LXXIII.

A few examples may be mentioned. An injection into the urethra will fometimes produce ficknefs at ftomach.—Great affections of the mind will often produce involuntary motions, even in those muscular parts that are not immediately under the influence of the will.—Fear will make the hand shake; or it will produce laxity of the bowels and diarrhœa; or it will affect the bladder with an involuntary flow of urine; or it will affect the eyes, fo as to occasion a fecretion of tears.

LXXIV.

It is not yet precifely underftood, how far the third principle, the mind, is capable of fympathifing with the first principle, fimple life. But it does appear, on fome occasions, as if the state of the mind became affected by the state and condition of the body.

wallout fome of the irritable part chrough

LXXV. Simple

A.IIIXXII

and in a diffimilar

LXXV.

Simple life belongs more particularly to the natural functions of every animal, fuch as digeftion, and the like. An affection therefore of the mind may be produced by an affection of part of the body only, as the ftomach, &c. There are inftances, in which it appears, that the mind is capable of fympathizing with life, as in the hypochondriac affections.

LXXVI.

On the other hand, the mind will be often light and eafy, without any apparent reafon or caufe, which may rationally be attributed to a fympathy it has with fome part of the body in a ftate of reft, or in a condition of performing its function with uninterrupted eafe and freedom.

LXXVII.

In the hypochondriac affection, fome part of the body, particularly connected with the natural and animal functions, is in all probability difeafed, though we cannot ex-C actly

actly fay what the difeafe is. The body thus interrupted in its more common action, produces affection in the mind. (L. LI.)

LXXVIII.

The full force and vigor of the mind arife from the fenfes being at reft, or at eafe; and its variations, as far as they relate to the greater or leffer degree of its excitement or collapfe, most likely depend upon one or more of the fenfes, or functions of the body, being fome way deranged, and difaffected.

LXXIX.

The mind fympathizes with the first principle, fimple life, through the affections of the fecond, the fensitive principle, which conftitutes what is called inftinct: or in other words, the mind may be faid to fympathize with the living principle in her wants. Mr. J. Hunter observes, that there is a necessfary connection kept up between all three.

CHAP. III.

VIN SACINGS VVINC

On the Sympathies which attend difeafes.

SECT. I.

Inflammation, when sympathetic, and when specific.

LXXX.

IN Chap. I. Sect. IV. I mentioned the relation of fympathy to the animal æconomy, in the prevention and cure of difeafe. There are fome fympathies purely preventative; fuch are fome of the phænomena at the commencement of fever, as naufea and loathing of food. Thefe affections of the ftomach are produced before the fever is formed, in order to remove by re-action the principal and fupporting caufe of fever, whether at the ftomach or fkin, The fickness of pregnant women, from the great fympathy between the ftomach and uterus, often, I believe, prevents abortion and premature labor. A vomiting will even prevent premature death. (XXXIV.) C 2 LXXXI. Some

LXXXI.

Some of the fympathies which attend difeafes contribute towards a cure, many are apparently ufelefs. They may vary much either from the nature of the part fympathizing, or from the nature of the ftimulus belonging to the fpecific quality of the difeafe.

LXXXII.

It is not in all cafes eafy to be determined, how far fympathy is fimilar to its caufe, either in the fpecific effects of the ftimulus, that has fpecific properties, or in the nature of the part which fympathizes, when the whole or fome one part of the body is affected with fpecific difeafes. (XLVIII.)

LXXXIII.

Mr. J. Hunter fufpects that it is fimilar in all local fympathies, fuch as fpreading cancer, fpreading venereal chancre, &c. Thefe are certainly to be confidered as fpecific difeafes, and when parts fympathize with them, and the fympathy is nearly local, then the part that fympathizes feems to be of

of the fame nature with the difeafed part itfelf.

LXXXIV.

When a bubo fpreads Mr. J. Hunter confiders the furrounding parts as venereal, and calls it a local fympathy. But local and connected fympathy in fome of its attributes is not fimilar, either in the action of the ftimulant, or nature of the part fympathizing, fo that when any remote part fympathizes, that fympathy is not fpecific, nor of the fame nature as the difeafe itfelf.

LXXXV.

These differences, it has been imagined, may arise from some real difference in the nature of the stimulant; or, which is most probable, from a real difference in the part sympathizing. (LIV. LV. LVI.)

LXXXVI.

When the urethra is inflamed by a fpecific difeafe, for example, the venereal, the teftis often fympathizes; but the fympathy is not fimilar, and, like the original affec- C_3 tion

tion of the fympathant, fpecific ; it is there fore to be confidered as non-venereal.

LXXXVII.

Two opposite views have been taken of the affection of the glands in the groin, in the cafe of gonorrhœa. 1ft. These glands often swell in this disease; but as it also very often happens, that they are not venereally tainted, they must therefore be supposed to be affected from sympathy, because they are known to sympathize with the urethra.

LXXXVIII.

Secondly. When the glands of the groin fwell, and alfo become painful, in confequence of a venereal gonorrhœa, it may be reafonably fuspected, that it is not fympathy that affects them, but an abforption of the venereal virus; and if it is abforption, we must confider them as venereally tainted.

LXXXIX.

The first view of this affection of the glands (LXXXVII.) is well founded; for there is not a doubt, but that they often fym-

fympathize with the irritation of the urethra in gonorrhœal cafes. When they, on fuch occasions, sympathize in confequence of the venereal difeafe in the urethra, fuch fympathy arifes not from a venereal stimulus at the gland, but is to be confidered as a common inflammation.

XC.

In the cafe of the cancerous breaft, the glands in the axilla will fometimes fwell, and fooner or later again fubfide. This affection of the glands has been alfo confidered as a fympathy, becaufe we know of no inftance of a cancerous affection fubfiding any where. This has led Mr. J. Hunter to imagine, that specific inflammations are not able to give to the fympathifer the fpecific variety.

XCI.

The most cautious investigation has not hitherto been able to determine, how far many other fimilar properties of specific inflamman tions differ from their fympathies. But it has been fupposed by Mr. J. Hunter, that those complaints, which are common to all parts C 4 alike

39.

alike, fuch as common inflammation and common fuppuration, may have the power of fympathy in these common respects. To illustrate this, it may be observed, that the glands of the axilla swell from the inflammation caused by a blister applied to the breast. An inflammation also of one eye will affect the other.

XCII.

An inflammation of the teftis, fays Mr. J. Hunter, whatever be the caufe from which it arifes, or whatever be the confequence which it may produce, will always be attended with a heavy fickly pain. But where the nature of the two parts is fuch, as to caufe a difference in their attributes, there the affections of the fympathant and fympathifer will differ. (LXXXI.)

XCIII.

The fame difference will alfo take place, where the parts themfelves differ in the nature of their difeafes; fo that if any part becomes difeafed in a manner peculiar to its attributes,

attributes, the fympathifer will be alfo affected in a manner agreeable to, and confiftent with, its proper attributes.

XCIV,

Again; when the fympathifer has no peculiar or fpecific mode of action, then it will be affected in the common way, altho' the part ftimulated, or fympathant, has fome fpecific nature, or a peculiarity in its mode of action. (LXXXVI. LXXXIX.)

CHAP, III. SECT. II.

Difeased Sympathy, bow partial, and how universal.

XCV.

Sympathy in various ways becomes partial or univerfal. It may be partial from a local or partial injury, or it may be partial from an univerfal difeafe.

XCVI.

Sympathy is partial from a local difeafe, when the caufes of action in a part become the caufe of action in a diftant part. Exempli

empli gratiâ, when any thing tickles the nofe or the fides, the mufcles of refpiration act by fneezing and laughing. Again; when any thing, particularly inflammation, irritates the liver, the fhoulder fympathizes. This is an inftance of a difference taking place between the action and affection of the fympathant and fympathifer; for the pain of the fympathant is obtufe, but that of the fympathifer acute.

XCVII.

Sympathy may be partial from an univerfal difeafe; for inftance, the flomach fympathizes with the whole body at the beginning of a fever; and the tongue, it is faid, fympathizes immediately with the conflitution on the leaft diffurbance: or the tongue and mouth are affected with thirft, when the conflitution is in want of fluids.

XCVIII.

There feems to be, from the cleareft proof, the fame fympathetic relation between the tongue and mouth, and the conflitution when in want of fluid, as between the

the stomach, and the constitution when in want of folid aliment.

XCIX.

Sympathy often becomes univerfal, from a local or partial injury. Thus when a local injury is done to any part, or an operation of confequence performed, they will each be followed by an inflammation, fuppuration, &c. Thefe are to be looked on as local evils; but from fuch partial affection, fever will arife in the fyftem, which is to be confidered as univerfal fympathy, and is commonly called fymptomatic fever.

C.

Wounds and injuries of joints often produce great difturbance in the fyftem, which is indicated by that univerfal fympathy, commonly called hectic fever, which is fometimes accompanied with partial fympathetic affections. On removing by amputation fuch difeafed parts, this difturbance of the fyftem has immediately fubfided. But it

it will be beft now to fpeak of each fympathy feparately.

CHAP. III. SECT. III.

Difeased Sympathy, when particularly partial.

CI.

Partial fympathy belongs to the fyftem, as well when in health, as when labouring under difeafe; and is therefore natural or difeafed. The natural fympathy more properly belongs to the fecond Chapter, but as far as it is partial, I fhall introduce it here,

CII.

Partial fympathy, when natural, always tends to produce fome falutary effect. There is an inftance of this during the period of utero-gestation, when the breasts of women increase in fize, in proportion commonly with the distension of the uterus. Again, their swelling at the time of menstruation, with darting pains in them, is another instance of natural sympathy.

CIII. To

CIII.

To this head of natural fympathy may be referred, the contracting action of the abdominal mufcles with the actions of the mufcles about the anus and rectum, in the expulsion of the fœces. In a like manner are we to confider the increased fecretion of the lachrymal gland, on ftimulating the eye-lids, or affecting the mind in any particular manner. But of these natural fympathies I need speak no more, as they are not directly to our present purpose.

As etamples of .VID have, we feel that.

The difeafed partial fympathy may be divided into two kinds, viz. that belonging to fenfation only, and that arifing from real difeafe, or difeafed action or affection. The first kind of fympathy is the most common. (XCVI. XCVII.)

CV. CV.

other fevers.

The greater the local mifchief, the greater is the fympathy and confent: for example, If great mifchief is done to the liver by inflammation, &c. the fhoulder fympathizes, in its

its pain in proportion. The more violent an inflammation of the teftis, the greater will be the pain in the back.

s au CVI. aroda antonen o

The greater the difeafe in the conflitution, the greater will be the local or partial fympathy; and the greater confequence the fympathant is of in the conflitution, the greater the affection of the fympathifer. (LXX)

CVII.

As examples of the above, we fee, that the flomach fympathizes with a difeafe of the brain, teftis, &c. A remarkable inftance of it likewife occurs in fever when the affection is proportioned to the fpafmodic condition of the cutaneous extreme veffels: therefore in the plague the flomach is more affected than in other fevers.

CVIII.

In the fecond part of this treatife I shall shew, that there is a wonderful sympathy between the stomach and skin, founded, in a great

a great measure, on a balance and connection between the internal and external extreme veffels. Hence affections of the stomach often produce confiderable sympathetic operations in the skin, so as to throw out fometimes a sweat, at other times an eruption, &c.

CIX.

I have often observed an extraordinary fympathy between the stomach and kidneys. It shews itself both in health and difease. A single glass of wine has often produced considerable diures in a short space of time; and great affections of the stomach often attend nephritic complaints.

CHAP. III. SECT. IV.

Difeased Sympathy, when in a special manner universal.

CX.

Univerfal fympathy takes place, when the whole body in a great meafure fympathizes with the difeafe, which was at first confined to some one part of the body. Thus, fymptomatic fever attends topical inflammations. (XCIX.)

to with by

CXI.

When fuch univerfal fympathy fubfides from a different mode of action taking place, to wit, from fome change for the better which the parts immediately concerned have not been able to bring to a complete cure, the conflictution fympathizes in another way, to wit, by hectic fever. (C.)

CXII. CXII

Thus, when the inflammation of any wounded or injured part is gone, but it ftill remains in an ulcerous flate, the conflitution is teazed; and this teazing Mr. J. Hunter has confidered as the principal part of what is called hectic fever, which by most physicians has been generally attributed to the absorption of purulent matter.

CXIII.

It is a very curious and ufeful obfervation made by Mr. J. Hunter, that a difeafe, which the conftitution can cure, never produces an hectic Of this kind are fanious wounds and ulcers; but if fuch an unhealthy part is removed, even the violence of an operation

operation does not obviate the acquisition of speedy relief to the system.

CXIV.

Mr. J. Hunter has feen a wound in the knee keep a man awake feveral nights, attended with a conftant purging, and the man at the fame time becoming hectic. By amputating the part, the man flept the whole night, the purging ceafed, and a coftivenefs rather enfued. (C.)

CXV.

A man with a wound in the elbow joint was attacked exactly in a fimilar way. His pulfe was hard and quick. Within ten minutes after the removal of the arm, the pulfe became flow and fofter. The conflitution immediately felt fhe had got rid of fomething fhe could not manage, and was quiet. In fuch cafes fymptomatic fever commonly arifes, but they lofe the hectic fymptoms. (C.)

CXVI.

49

2

CXVI.

This univerfal fympathy of the fyftem is greateft, when certain parts are injured. Thus, it will more eafily take place, when an injury has been done to an involuntary part, than if the fame quantity of mifchief had been done to a voluntary part, becaufe the first are more universal in their connection with the constitution than the last, and belong to important functions. (CVI.)

CXVII.

When a vital part is deceafed, there is a greater fympathy between the deceafed part and other vital parts, than between it and other lefs important parts of the fyftem : and there must be a failure, in fuch cafe, in the function of the lefs important parts, befides the fympathy of the vital parts. (CVII.)

CXVIII.

Univerfal fympathy is alfo greater, if the injury is done to a part far remote from the fupport of life, the heart, than if the part injured

injured be nearer, provided the injury be the fame. The animal machine is then more confcious of inability, more alarmed, and hectic fever is more liable to be produced. (C.)

CXIX.

Ex. gr. If an injury be done to the toes, or foot, the conftitution becomes more affected and difturbed, than if a fimilar injury had been done to the fhoulder, even where all other circumftances are the fame. Pulmonary injuries and complaints make however, in my opinion, an exception to this rule. (CXVII.)

CXX.

Univerfal fympathy takes place more readily, or its actions and affections are more ftrongly marked, when the powers of the animal machine are capable of repairing the injury received; and, vice verfa, it takes place more flowly, and is lefs ftrongly marked by its confequences, when the powers of life are more languid. (CVIII.)

CXXI.

CXXI.

The ftate of an individual body is often fuch, from various occafional caufes affecting the predifposition and temperament, that it will more readily fall into fympathy at one time, than at another. Thus people in general, and women more than men, are eafily affected, at particular periods, from flight caufes. (XII.)

CXXII.

Not only the fame conflictution differs at times owing to the change of predifpolition, but different conflictutions vary owing to the difference of temperament. Hence, it is not at all to be wondered at, that fome people's feelings are affected much more readily than those of others. (XLVI.)

CXXIII.

As there is no part of an animal body, that is entirely independent of the conftitution, fo in all material injuries, the whole must fympathize more or lefs with a part material-

ly

ly injured : but as every individual part has a more intimate connection with fome one particular part, than with the whole together; fuch parts, as are particularly connected, must fympathize more readily, and in a greater degree.

CXXIV.

It may be obferved, in illustration of this fympathizing connection, that the ftomach fympathizes more with a difeafed teftis, than the ferotum, or the whole conftitution. So alfo the ftomach fympathizes moft with an affection of the extreme veffels; and yet I greatly fufpect, it is in a fecondary way; viz. through the medium of the fympathizing heart: for it is moft natural to fuppofe that the heart from its connection with them would very readily fympathize with an affection of the extreme veffels. Such a fympathy may perhaps be the foundation of fever, affifted by the operations of the ftomach in vomiting, an idea it may be pro-

per

53

per to inveftigate at fome other time. We have no ready way of knowing that the heart fympathizes with the conftitution, but by the ftate of the pulfe. If the heart can at any time become a fympathifer, it most probably is fuch when hectic fever accompanies, either a part materially injured, or an univerfal difeafe, (C. CXIX.)

CXXV.

Moreover, univerfal fympathy in the confitution, from local or univerfal difeafe, has been faid to have its order of parts. Thus the head, the ftomach, and the back have been faid to be the firft parts which fympathize with the whole conftitution, From what I have faid in the preceding paragraph, *I think the heart must be reckoned* with them. The tongue, fkin, and kidneys have been faid next to fympathize, &c. Thus there feems to be a certain order of local fympathies from an univerfal caufe,

CXXVI.

55

CXXVI.

This priority of fympathetic operations in the fyftem fhews itfelf moft remarkably at the commencement of fever. Thus the head aches; a pain in the back comes on; the ftomach is affected with anorexia and naufea; the pulfe becomes flower and weaker; the fkin becomes dry; the tongue is parched with thirft; the kidneys fecrete an urine of a particular quality, &c. All thefe phænomena feem to arife from fympathy between parts endowed with fimilar properties and functions.

CHAP. III. SECT. V.

Universal Sympathy, when immediate, and when secondary.

CXXVII.

Univerfal fympathy is observed to be either immediate or secondary. The conflitutional fympathic affections may arise in D 4 many

many cafes from the local fympathy, and not from the action and affection of the part first injured. This is observed to be very remarkable, when the fympathizing part happens to be a vital part, whether the first injured part is vital or not. (CXII.)

CXXVIII.

To illuftrate this; if either the ftomach or heart fympathizes, fuch a fympathy produces more morbid effects in the fyftem, than what the original caufe of fuch fympathy was capable of doing; and thus the conftitutional fympathic affections do not all arife from the firft local affection, but fome are caufed, and fome are increased, in a fecondary way, through the medium of either the fympathizing heart, or the fympathizing ftomach. (CXXIV.)

CXXIX.

Many of these fympathizing connections are by nature fixed and regular. Thus we see, that the muscles of respiration constantly

ftantly fympathize with the throat, and nofe, when they are tickled, or otherwife irritated. The muscles of respiration also fympathize with the lungs. They become fecondarily affected with the disease of the lungs, through the medium of the orifice of the trachea, where the fympathy with the diseased part of the lungs first takes place.

CXXX.

The refpiratory mufcles are alfo fpafmodically affected in fympathy with the ftomach, when diforders of it bring on vomiting. Cough alfo, and laborious refpiration accompany any impediment to the free action of the heart, in giving an even circulation. Stated fympathies are alfo produced from affections of the uterus. Thus cough and fever are often caufed by uterine irritation, at the time of abortion, and when the mifcarriage comes away, fuch univerfal fympathies very fhortly go off. The ftomach likewife commonly fympathizes with uterine affections.

CXXXI.

CXXXI.

The glans penis alfo fympathizes with affections of the bladder, as when there is a calculus, or calculi within it, &c. There is hardly an inftance of any one having a difeafe in the bladder, without being troubled with more or lefs of a painful uneafinefs at the glans penis.

CXXXII.

-ordinal oli

CXXXX

Again, the head fympathizes with diforders of the ftomach; and vice verfa, affections of the incephalon often produce ficknefs and vomiting. An affection of the heart, fuch as a difposition to fyncope, will produce both fieknefs, and head-ach. The fame often takes place, upon recovering from a deliquium animi. Of these and other fympathizing connections (CXXIX. CXXX. CXXXI.) fome are natural, fome difeafed, and fome, from accidental circumstances, either natural or difeafed.

CXXXIII.

CXXXIII.

On the other hand, fome parts of the body feem to have no fympathizing connections; at leaft the effects of them are not obfervable. But those difeased fympathies, which are observed regularly and constantly to take place, have been supposed to have the principal of their regularity and constancy founded upon the fympathizing parts having a natural connection with the healthful operations of each other. Such supposition is very rational.

CXXXIV, CXXXIV,

This way of accounting for their regularity and conftancy more immediately relates to the fympathic affections of the vital parts. Thus it holds good with all the parts connected with, and concerned in, refpiration, as the nofe, trachea, &c.; alfo in most of the affections of the head, stomach, and heart; and of the stomach with the intestinal canal and rectum, &c. They are in sickness to be

be confidered as no more than the actions of healthy fympathies, præternaturally increafed from difeafe.

CXXXV.

The difeafed fympathies, between parts that have no connection, are but few; there are however fome examples of them. Such is the fympathizing affection of the right fhoulder with the inflammation, &c. of the liver; the difturbance of the ftomach from affection of the kidney from ftone, &c. the affections of the teftes in the male fex, and breafts in the female fex, from the fwelling of the glands of the neck in the Cynanche parotidœa.

CXXXVI.

be parts con-

Difeafed fympathies, which are immediate, take place but rarely between diftant parts of the body; for which reafon they are very properly termed irregular. The left arm, however, has been known to fympathize with

with a difeafed bladder, and the left hand with an abfcefs in the fubftance of the brain.

CXXXVII.

Mr. J. Hunter relates, at his lectures, a remarkable cafe of the Earl of Clarendon. He was, it feems, fubject to violent fits of the ftone, unaccompanied with the ufual pain at the glans penis or bladder, but the pain was all in the left arm. When he voided urine, and it was commonly in his fits, the pain in his arm would come on : he became then fo ill, that he often thought the fits would kill him, and in fact he was deftroyed by one of thefe fits. This was a very uncommon fympathy, and an irregular one.

CXXXVIII.

It is fomewhat remarkable, that few of the fympathies from natural connection can be reverfed. Thus when a difease attacks the intercostal muscles, we observe no fympathetic

thetic affection of the lungs and throat; but when a difeafe attacks the throat or lungs, the intercoftal muscles more or lefs constantly fympathize.

CXXXIX.

When the glans penis is difeafed from chancre, &c. no irritation paffes to the neck, or any part of the bladder. Nor do difeafes of the inguinal glands or teftes affect in any way the urethra. The mutual fympathy however between the head, ftomach, and heart, is contrary to this general rule.

CHAP.

CHAP. IV.

On the uses of Sympathy.

SECT. I.

Natural Sympathies most evidently useful.

CXL.

SYMPATHIES have undoubtedly their ufes, and anfwer very important purpofes in the animal œconomy. Their utility can be most plainly discovered in the natural and healthful state of the system; or when there is not any material organic injury, but only an irregular morbid action of a found part.

CXLI.

The found parts most likely to produce fympathetic actions and affections, belong to the vital and other functions of the fystem, the constitution of which are somewhat fimilar,

lar, and whole operations are in a great degree dependent on each other. (XXXIV.)

CXLII.

In Section IVth of Chapter Ift, which treats of the extensive relation of fympathy to the animal œconomy, I have observed, that the prevention and cure of disease, and even the restoration of animation when apparently lost, are founded on a VIS INSITA SYMPATHIE, fuited to the feveral purposes, and called forth by the omnifcient will of Providence.

CXLIII.

Natural fympathies eftablifh a connection between the fame principle in different parts of the body; in this way fimple life, and fenfation, or the fenfitive principle, are often preferved in a part fympathant, through the medium of a diftant part fympathizing. Hence, where uncommon exertions of the natural actions of parts are wanting, the ufe of fympathy is often abundantly evident.

CXLIV.

CXLIV.

Some proof and illustration of this may be drawn from the effects of any uncommon irritation on the lungs, throat, or nofe. When fuch occurs the muscles of respiration produce cough, with uncommon force, and the offending matter, whether extraneous or derived from the constitution, is thereby thrown off.

CXLV.

The fympathy becomes fomewhat more complicated, when the irritation, as is moft frequently the cafe, takes place in the lungs; then the fympathizing irritation is firft transferred to the larynx or gullet, denoted by a tickling fenfation of the part, which induces the mufcles of refpiration to fympathize in their turn, whereby coughing is produced, either to get rid of the proximate caufe, whatever that may be, or the difeafe with its confequence, which commonly is an increafed morbid fecretion.

CXLVI.

CXLVI.

An affection of the lungs, producing fuch irritation at the larynx, appears to me to point out the neceffity of a connection fome way or other between the fympathant and fympathifer. The mufcles of refpiration, on fuch an occafion, fympathize indirectly with the lungs, through the medium of the fympathizing larynx, as if the want of continuity between these mufcles and the lungs rendered it impossible for any direct fympathy to take place.

CXLVII.

This complicated fympathetic connection alfo affords us an inftance of a difeafed fympathy producing a natural one; for we muft confider the affection of the larynx as a difeafed fympathy, while the action of the refpiratory mufcles is a natural fympathy : and this compound fympathy at the fame time affords us a proof, that affection in the fympathant can produce action in the fympathifer. (LI. XCVI.)

CHAP.

CHAP.IV. SECT. II.

t entemerated do not feens

Diseased Sympathies less evidently useful.

CXLVIII.

It is a queftion curious in its nature, and interefting in its tendency, why, in real difeafes partial or univerfal, fympathy fhould take place, when the affections or actions of the part fympathizing do not anfwer, at leaft apparently, any visible good to the part difeafed ? Such operations are not eafily accounted for:

CXLIX:

The fympathies between the liver and fhoulder; between the flomach and teftes; and between the flomach and kidneys, tend to no apparent utility. The fame may be faid of the fympathy between the brain and flomach. It feems highly proper, that parts from a fimilarity of conftruction, connection, &c. fhould be affected by fympathy, and occafionally called into action by it; and fuch fympathy may be accounted for. But E 2 the

the fympathies just enumerated do not feem to arife from any organic connection, as far as we anatomically know.

CL.

Though fuch difeafed fympathies can neither be explained on the fame principle as the natural fympathies, nor anfwer any wife purpofe that we can fee, it may however be obferved, that in fuch cafes the fympathifer may fomewhat relieve the fympathant, though no immediate advantage is apparently derived from it; on the contrary the cafe may perhaps feem worfe than before.

CLI.

It is worth our attention, fays Mr. J. Hunter, to confider how far the fenfitive fympathy will admit of a division of abfolute pain; for (continues he) we may affirm, that any given quantity of abfolute pain may, when diffused, become infensible, which might, when concentrated, become intolerable.

CLII.

CLII.

This idea of Mr. J. Hunter's leads me to think, that the fympathy of a found part relieves the fympathant, or difeafed part, by enduring part of that pain, which collected at the difeafed part would have been infupportable, the irritability of it being neceffarily increafed by its morbid condition. The ftomach therefore fympathizes with the teftes, &c. If however it is found that pain is increafed by fympathy, this theory falls to the ground.

CLIII.

Mr. J. Hunter mentions the affection of the glans penis from a difeafe of the bladder, as a difeafed fympathy without apparent utility, and difficult of explanation. May there not however be a fympathetic connection between the glans penis and kidneys, fimilar to that which takes place between the larynx and mufcles of refpiration? To render this analogical opinion plaufible at leaft, I fhall attempt to draw a comparison between an affection of the bladder, and an affection of the lungs.

E 3

CLIV.

CLIV.

I will fuppofe, that the lungs are irritated by an acrid effusion or fecretion, or by infarcted yeffels. The action of coughing may give relief by expelling the one, or deterging the other. To produce therefore this neceffary operation, an antecedent fympathy takes place at the larynx. It is of itfelf an uneafinefs which the conftitution would attempt to get rid of by coughing.

CLV. benorg oritest alleft

In like manner, the bladder may be irritated by gravel, acrid faline urine, &c. An increafed fecretion of urine might relieve by expelling the one, or weakening the other. To this purpofe the kidneys muft be flimulated to action, to which they may probably be excited by the irritation at the glans penis; which irritation may alfo take off the flricture caufed at the neck of the bladder by the irritating matter; and it may be further obferved, in proof of this opinion, that generally an evacuation of urine takes place foon after a fit of the flone and gravel.

CLVI.

CLVI.

The parts concerned in each of the above cafes of compound fympathy (CLIV. CLV.) belong to a feparate and diffinct function in the conftitution, the difeafes of which bear fome analogy, and therefore their fympathies will reafonably admit of a fimilar explanation.

CLVII.

I have given a chapter upon inflammation when fympathetic, and when fpecific; yet I fhall beg to throw in a queftion here, on that division of the fubject. What can poffibly be the ufe of a common fympathic inflammation in one part, from a fpecific difeafe in another part? I fufpect, that fuch inflammation, when it feizes on lymphatic glands, is intended to prevent the abforption of the fpecific virus. This may be illuftrated by the fwelling of the inguinal glands in a venereal gonorrhœa, or of the axillary glands in a cancer of the breaft.

CLVIII.

71

CLVIII.

This opinion is rendered probable from obferving, that the lues venerea very frequently makes its appearance after a gonorhœa has been cured, when there has been no antecedent affection of the glands of the groin; and vice verfa, where the glands have fwelled and been very painful, which is frequent in cafes of gonorhœa, the affection of them abates, as the original complaint goes off, and is not commonly followed by any fyphilitic complaints.

CHAP. IV. SECT. III.

What Sympathies may be applied to medicinal use, and how.

CLIX.

By being acquainted with the natural fympathies in the conftitution, we are enabled to apply our remedies to the healing of a partial affection, on the principle of affecting the fympathant, through the medium of the fympathifer. We are alfo, by fuch

fuch a knowledge of the animal æconomy, guarded against a mistake we might otherwise easily fall into, of confidering a part as difeased, which in fact is only affected from sympathy with some distant discased part,

CLX.

I fhall mention a few fympathies ufeful to be known. An attention to the fympathy between the ftomach and fkin may on fome occafions affift us in our practice. In obftinate vomitings the warm bath may relieve, and effectually put a ftop to them. External applications of theriaca, &c. to the pit of the ftomach have been attended with the fame good effect. In cutaneous difeafes we fhould remember, that the ftomach may only be fympathetically affected; and that fuch diforders may be cured by the operation of medicines on the ftomach. (CVIII.)

CLXI.

Diforders of the head frequently originate from a morbid condition of the ftomach. Thus epileptic and other nervous complaints have followed too liberal an indulgence of the

the appetite in wine, &c. leaving fometimes behind them a failure of the memory, and occafionally paralytic complaints. In fuch cafe, our remedies are to be fuited to the morbid condition of the ftomach.

CLXII,

Int's

A fwelled teftis can be cured on the principle of fympathy between it and the ftomach. Thus a few grains of ipecacoanha have been known to cure fuch an affection. Emetics have also cured the white fwelling of the knee.

div bebasis accended with.

It has long been a received opinion, that cantharides applied to any part of the body, affect the bladder, and (as fome fay) exulcerate it, particularly if they ftay on long. I have alfo read of a kind of ftone, brought out of the Weft Indies, which was faid to have a peculiar property of difcharging gravel, and of diffolving the ftone; infomuch that when laid to the wrift, it has fo forcibly expelled urine and gravel, by its violent manner

ON SYMPATHY IN GENERAL. 75 ner of operating, that the fick perfon has been glad to remove it.

foles of the feet fr. VIXLD leeding pigeons.

tice to onle the head by applying to the

What has long been noticed, is even to this day confirmed by frequent experience, namely, that the foles of the feet have a great affinity of confent with the head, and upper orifice of the stomach. It is generally known, that going barefooted to those that are not accuftomed to it, will fome times affect both, each with its specific complaints. These affections are often removed by hot applications to the feet, to attenuate, as the ancients used to fay, the rheum. Formerly a phyfician, fays a voluminous writer, who wished to appear mystical, prefcribed for the cure of the rheum, that the patient fhould walk continually upon a camomile alley (fuch was the old language) meaning thereby that he fhould put camomile into his focks.

CLXV.

VX12e fick werlon has

Again: it has occasionally been the practice to eafe the head by applying to the foles of the feet fresh flain bleeding pigeons. Sinapisms are to this day frequently applied to the feet to relieve the head. Sleep is also fometimes obtained by the application of foporiferous medicines to the feet.

CLXVI.

-1; general-

VXJO

The wrifts and the hands have been fuppoled by fome to have a fympathy with the heart. Thus, it has been obferved, that the affections and paffions of the heart and fpirits are notably difclofed by the pulfe : and long continued agues, it is faid, have been cured by the application of garlick, and the like medicines, to the hands, and wrifts. Eggs of alabafter, and balls of chryftal have been held in the hand, in order to appeale the fury of a febrile heat. Solutions of nitre applied to the palms of the hands might, perhaps, on the fame principle, anfwer a fimilar purpole.

CHAP.

CHAP. IV. SECT. IV.

Sympathy, when the province of the physician, and when of the Surgeon.

CLXVII.

Difeafes with their fympathies are either local or univerfal. If fympathy be either particular or univerfal, and the caufe be known, it falls under the care of the *medical* furgeon; but if the caufe be not known, or if it be not even known that a caufe exifts, then the fympathy becomes the province of a phyfician, more efpecially if it is univerfal.

CLXVIII,

Local difeafes in all cafes come more properly under the care of the furgeon; when univerfal they fall under the province of the phyfician.

CHAP.

CHAP. V.

CHAP. IV. SEC

On the Sympathetic operation of medicines:

With their lyingath

SECT. I.

Effects of medicines, when local, and when remote.

CLXIX.

MEDICINES produce universal effects by partial operations. The fympathetic effects of medicines may be divided into two kinds; namely, those local or fimilar to the action of the stimulus, and those remote and particular from the nature of the sympathiser.

CLXX.

The fympathetic action will be local, when only the furrounding parts act in concert with the part ftimulated. It is to be confidered as remote, when the remote parts fympa-

fympathizing have fome fpecific connection with the part ftimulated, as the ftomach with the teftis; head with the ftomach, &c.

CLXXI.

The fympathetic operation of a medicine may be either peculiar to the mode of action of the part that fympathizes; or it is poffible that it may be agreeable to the mode of action of the part flimulated.

CLXXII.

That fympathy, whofe influence is fimilar, or which only acts topically, lofes its power by diftance fimply, and may be called truly local; and fuch medicines, as produce thefe effects, may be claffed among the local applications.

CLXXIII.

If fimilar and local fympathy did not affift the operation, and effects of medicines, they could have but little power to do good, as most local difeases extend fome way beyond the furface of exposure, which extension itfelf

SO ON SYMPATHY IN GENERAL:

felf arifes from fympathy; therefore it is probable, that the fympathetic effects of medicines may alfo extend as far.

CLXXIV.

If this was not the cafe, how could a fomentation give eafe to a fwelled teftis? for it only comes in contact with the ferotum; neither could æther, on any other principle but fympathy, cure or eafe a head-ach, by being only applied to the fkin of the forehead.

CLXXV.

It is well known to every one, that by applying medicines fimply to the fkin you may relieve inflammations, that go fome depth: if it was only the furface of contact, which was ftimulated, that could receive eafe, in fuch a cafe the medicine would produce very little effect; but as they produce a healthy action in the furface of contact, the other neighbouring parts fympathize with it, and all are relieved in a proportionate degree.

CLXXVI.

CLXXVI.

Local and fimilar fympathy is certainly effential towards the giving efficacy to all external applications. For if it was neceffary that the medicine fhould come in contact with every part that was difeafed, it would be impoffible for any part, difeafed in this way, to be either relieved or cured.

CLXXVII.

On the other hand it is certain, that all fympathetic effects of medicines are not fimilar to their effects on the part of application; for we know that many medicines, when applied to fome parts, fhall cure, by effects on fuch parts, a diftant part when the fame medicine, applied to the difeafed part, would kill. Such are the effects of many applications to the ftomach, which cure pain, and other complaints of the head; but if you was to apply the fame medicine to the head itfelf (*i. e.* to the brain) it would kill. (CLXX.)

CLXXVIII.

If therefore remote fympathy did not act in many cafes upon the fame principle F

as local, we could not account for the effect of a fomentation, in relieving a pain of the bowels, when applied to the fkin of the belly; or in relieving a pain of the thorax, when applied to the breaft.

CLXXIX.

It is alfo known, that there are many medicines, which cure by remote fympathy, which could have no effect when applied to the part difeafed. Such is the effect of ipecacoanha applied to the ftomach in curing a complaint of the teftis, which, applied to the teftis itfelf, would be of no avail. (CLXII).

CLXXX.

In the cafe just mentioned, the medicine is rendered capable of performing a cure by fympathy, from having produced a peculiar effect on the stomach; fo that the effect which fuch medicine has in the stomach, is not similar to any effect that it would be like to have, if applied to the part itfelf.

CHAP.

CHAP. V. SECT. II. Sympathetic effects of aqueous applications to the skin.

CLXXXI.

We fhall be better able to underftand the foregoing theory, (which is chiefly Mr. J. Hunter's, and most of the arguments in the following Sections of this Chapter are to be received as his) if, by way of illustration, we contrast it with the common received explanation and opinion respecting topical applications. Their mode of operation has been generally supposed to depend upon their entering the pores of the skin, and being, by such channels, conveyed to the parts affected. Hence arose the expression, "Such a medicine is penetrating."

CLXXXII.

Let us in the next place fee, how phyficians have eftablished their mode of practice upon this mechanical idea. To inflammations they order fomentations to be applied as hot as they can be borne, because warmth and moisture penetrate. The pores are thereby opened, and the moisture is supposed to pass in; but if we reflect for a moment, we F_2 supposed to the full

83

fhall know, that the parts beyond the furface of contact cannot be made moifter than they naturally are.

CLXXXIII.

It is upon the fame principle, that people labouring under diflocations, ruptures, &c. are ordered to be put into the warm bath, or have fomentations applied to the injured parts; for there is no doubt, but that the relaxation of the parts morbidly affected would be very falutary and beneficial in fuch cafes, if it could be accomplifhed; and nothing has hitherto been confidered fo effectual, towards attaining fuch an end, as warmth and moifture.

CLXXXIV.

A piece of dried meat, fays Mr. J. Hunter, or dried leather, may be affected in this way, but a dead body, that has as yet undergone no fuch change, cannot; much lefs a living one. " A man, fays he, meets with a diflocation, he is ordered to be put into warm water to be relaxed; now it is morally impoffible (continues he) that the warm

warm water can relax by adding moifture to the body: not a drop of water can poffibly pafs into the joint. The ligaments will not become a bit moifter, though you was to foak him to eternity."

CLXXXV.

It becomes then a queftion; Do thefe applications ever do good ? Mr. J. Hunter allows they do, but upon the principle of fympathy. The furface of contact is by their operation lulled or foothed, and the irritability either of the whole, or part, of the difeafed affection is put to reft. This acquired diminution of partial excitement allows us, where violence is to be ufed, as for inftance in a diflocation, to ufe it more freely than before.

CLXXXVI.

Irritation is often removed upon the principle of remote fympathy. Thus a man is put into a warm bath for a fuppreffion of urine. Surely, warm water, for the purpofe of penetrating fimply, cannot in fuch a cafe be neceffary ! How then does it act ? F 3 It

It acts upon the fame general principle of fympathy. The whole body is put to reft by the irritability of the fkin being foothed, by which means the fuppreffion, from the irritation at the neck of the bladder being taken off, is removed, or in fome meafure relieved.

CHAP. V. SECT. III.

How unctuous and volatile applications by Sympathy increase motion.

CLXXXVII.

Volatile fubftances (fays Mr. J. Hunter) are alfo frequently recommended upon the fame fuppofed principle of their being penetrating. But it is well known, that bodies in vapour are not fo penetrating, as the fame bodies in the fluid form. Neither air, nor the fteam of hot water will pafs, where water alone will pafs readily. Hence the operation of volatile fubftances is affifted by fympathy.

CLXXXVIII.

Oils have been recommended in the way of friction, upon the fame mechanical principle

ple of being penetrating; and two purpofes were meant to be anfwered by them; firft, the giving of motion, and fecondly, their procuring refolution. The mild and foft oils have been recommended for the firft intention : and this idea of the medical utility of oil muft have arifen from its anfwering fo well in making an old hinge go eafy, or an old boot fit eafy.

CLXXXIX.

Unfortunate for the animal body, which labors under the want of any free motion, oil cannot penetrate beyond the furface to affect mechanically the inward parts, even fuppoing in other refpects the mechanical conftruction the fame; for an animal is moift or wet in all its internal parts. Therefore oil cannot poffibly touch any internal part of an animal body. Fortunately for the hinge, oil may penetrate into all its moving apparatus.

CXC.

Animal oils have been commonly most esteemed, and preferred to the vegetable, F 4 because,

becaufe, having come from an animal, it was thought they might more readily enter one again. Good effects are no doubt obtained from the ufe of fuch outward applications, but they most probably arife folely from the mechanical motion ufed in the application of them.

CHAP. V. SECT. IV.

How unctuous and volatile applications procure refolution.

CXCI.

The fecond intention, for which fuch applications have been recomménded, is to procure refolution; to anfwer this indication the warmer and effential oils have been in general ufed. The fame objections hold against their penetrating, as were made against the operation of the mild oils. Therefore whatever may be the medicinal quality they posses, it must depend upon their stimulating power operating upon the part of application, and producing its confequent fympathy,

CXCII.

CXCII,

A greater and more certain benefit may however arife from local applications of the ftimulating kind, than what can be procured either from the ftimulus they produce on the fkin, or from their fympathetic effects: for though we do not allow of their having any mechanical penetration, yet we may be obliged to admit of their operation by abforption, which in many cafes may have confiderable effects.

CXCIII.

If an external remedy be fo applied, that the abforbents paffing through the difeafed part, neceffarily take it up, it may confiderably affect the difeafe in its road to the circulation; for the medicine will be thus applied to the difeafed abforbents; and as experiments have demonstrated, that abforbent veffels are very irritable, fo the fympathy, which arifes from the operation of the medicine on them, may be more immediate, than from its operation fimply on the fkin.

89

CXCIV.

Mercury is the fpecific remedy for the venereal difeafe. When a gland in the groin fwells, it is the common practice to rub the fwelled part with fome mercurial ointment. It is probable however, in fuch a cafe, that the effect is only produced by fympathy, from the ftimulus of the remedy on the fkin: for the abforbents at the part of application in all probability do not pafs through the fwelled gland.

CXCV.

Moreover, the quantity of the medicine abforbed, from fo fmall a furface, can have no very great effect. In order then to produce a more confiderable effect it is proper to apply the mercurial preparation to the leg or infide of the thigh, which feldom fails of procuring the defired effect.

CXCVI.

The fuccefs of fuch applications on difeafed abforbents cannot attend other indurated glandular parts, that are not abforbent. Therefore the advantage derived from a fimilar practice in these cases must arise from

from the application of the remedy to the part affected, by its fympathetic operation, from the ftimulus afforded by it to the fkin.

CXCVII.

With refpect to all outward applications for the purpose either of increasing motion or procuring refolution, it may be observed, that the mechanical effect which arifes from them is different from their mechanical effect on a hinge. Upon fuch their action is immediate, by immediately altering the condition of the parts, which was the caufe of the fliffnefs. But their mechanical operation on a living body acts only as a ftimulus to the parts. By acting thus as a ftimulus to the external parts, they increase their action, and make also the neighbouring and more-deep-feated parts act by confort in a fimilar manner, till by this compound action the parts of themfelves become capable of removing those obstructions, which had existed as the impediment to their free motion, and as the caufe of induration.

CXCVIII.

CXCVIII.

I cannot conclude this chapter without taking fome notice of the operation of internal medicines. Their falutary effects are often fo quickly obtained, at diftant parts of the body, when the medicine cannot poffibly have been directly applied to them, that no other way is left to account for their fudden manner of relieving, but on the principle of the fympathy of the ftomach with the whole of the animal machine. It is in this way that we must explain the fudden good effects of Peruvian bark, in preventing the acceffion of an intermittent paroxyfm : and it is on the fame principle that a ftrong opiate will ftop the further progress of the paroxyfm, though the cold ftage has already commenced. I have frequently made trial of this mode of practice. An increased perfpiration is also caused by nauseating doles of antimonials from fympathy of parts.

CXCIX.

It will however still remain a question, what are the minute parts of the stomach, which

which are immediately acted upon by medicines? The fentient extremities of the nerves may very reafonably be fuppofed to be concerned in their operation; however I fhall have occafion to treat of this fubject more at large in the fecond part of the work, when I fhall endeavour to make it clear, that extreme veffels, from their fenfibility and natural connection, are most likely to become the channels of *Medical Sympathy*, through the medium of the fympathifing heart : and that this vafcular fympathy is particularly founded on a balance and connection between the external and internal extreme veffels, more efpecially those of the STOMACH.

CHAP.

CHAP. VI.

On the Sympathy of the Senses, and force of imagination.

SECT. I.

On the Sympathies produced by the passions and affections of the mind.

CC.

SYMPATHY "relates both to the operations of the affections, and to thole of the imagination :" these are often so much connected, that they have generally been treated of together. It is impossible indeed to confider the affections of all the fenses, without admitting the *force of imagination* to be often materially concerned. I intend however to confine my remarks chiefly to those affections and operations, which are not entirely regulated by its influence, and power.

CCI.

I cannot however proceed with this part of my work, without expressing a hope, that it will not be confidered as foreign and unneceffary to the plan of my prefent undertaking. I view it as a proper medical enquiry, to speak of whatever may concern the qualities and paffions of the mind, and affections of the fenses, in as far as, by their operations, they may affect and alter the corporeal stamina of man. The various temperaments are materially concerned in the ftate and condition of the mind, and therefore I hold it useful, to be well acquainted with the changes that may be produced in the body, through the fenfations and affections of the mind.

,CCII.

The passions and affections of the mind produce in the body different fensations and impressions, and as sympathies of consciousness determine in general the spirits to those parts, which labor most, or are most to be affected. "Thus *fear* and *anger* determine to the

4

the heart; lust to the eyes, &c. joy, pity, wonder, and the like, to the head. This observation may be of great use."

CCIII.

The paffions and affections have been faid to impress, and act upon, the body, in the following manner: " I. FEAR caufeth palenefs, trembling, the standing of the hair upright, starting, and screiching. 2. GRIEF and PAIN caufeth fighing, fobbing, groaning, fcreaming, and roaring: THEY alfo caufe tears, difforting of the face, grinding of the teeth and fweating. 3. Joy caufeth a cheerfulnefs and vigour in the eyes, finging, leaping, dancing, and fometimes tears. 4. ANGER produces palenefs in fome, and the going and coming of the colour in others: alfo trembling in fome, fwelling, foaming at the mouth, ftamping, and bending of the fift. 5. SLIGHT DISPLEASURE OF DISLIKE caufes fhaking of the head, frowning, and knitting of the brows. 6. SHAME caufeth blufhing, and caffing down of the eyes. 7. PITY caufes fometimes tears, and a flexion or caft of

of the eyes afide. 7. WONDER caufeth aftonifhment, and an immoveable pofture of the body, cafting up of the eyes to heaven, and lifting up of the hands. 9. LAUGHING, though hardly to be confidered as a paffion, fince it is produced by an affection of the mind, caufeth a dilatation of the mouth and lips; a continued expulsion of the breath ; with a loud noife, which maketh the interjection of laughing, fhaking of the breaft, and fides, and running of the eyes with water, if it be violent and continued. 10. LUST caufes a flagrancy in the eyes, and priapifm."

CCIV.

The affections of the mind of one perfon will often work upon the fpirits of many. Thus whole companies are fometimes difpofed to be fad and melancholy, or merry and jovial, when any one is prefent much inclined to either of those states of mind; and it has been observed, that old people, who have loved the company of the young, and have been conversant continually with G them,

them, have generally lived long. But young people muft not conclude from this, that the company and converfation of the grave and old will operate upon their *fimple* life, and fenfitive principle, through the affections of their mind, and difpofe them to be fhortlived. On the contrary, by thus improving their understanding, they will be more enabled to fortify their constitution, and refist the ravages of youthful indulgence.

CCV.

It may also be further observed, that those tender sympathetic affections, which lay hold of the mind, at the representation of theatrical performances, originate from the fame principle, while they are to be confidered as the furest test of just execution in the actor, and of the expressive language of the author. Indeed all *stage effect* depends on sympathy.

CCVI.

The affections of the mind make the fpirits more powerful and active, efpecially those which manifest themselves by the eyes.

eyes. Two in particular may be named, LOVE and ENVY. As fympathies of confcioufnefs their operations are more eafily felt than defcribed. Though opposite in their nature, they are equally violent in obtaining their particular ends. The one can no more fuffer indifference and difappointment, than the other contempt and haughtinefs.

CCVII.

It has been faid, that the paffions of the mind are occafionally infective, particularly fome of them. Thus FEAR and SHAME are fometimes very fuddenly fo. We frequently may have occafion to fee, that the ftarting of one will make another ready to ftart. Again, when one man is out of countenance in company, others will often blufh in his behalf. However, the ferious paffions may *furely* be fo under the controul of reafon, as to refift infection, whatever may be the cafe of *temporary mufcular or nervous attraction*.

G 2

CCVIII.

99

CCVIII,

I think there is much reafon to fufpect, that a connection between the affections and fenfations of the female mind and uterus is very materially concerned in the procefs of generation, and probably can alone give efficacy to those actions and impressions subfervient to conception, through the fympathizing affections of the mind. I speak of this generative principle as an universal one, and wish it not to be confidered, as belonging folely to rational beings. Such an influence may have been given to all animated nature.

CCIX.

-of sW.

One of the first medical philosophers of the prefent time is of opinion, that the mother has always the powers and principles of fashioning her child within herself, but that they are not rouzed to action without the stimulating influence of the male. The principles that must be immediately concerned are the sentient and living, but it is through the influence of the mental principle, that the

and image of the embryo is the form It has been fuid and many ftamped: opinion, that the force of in

With refpect to the depravity and force of the imagination, in the production of fympathies, they always operate most upon " weak minds and fpirits, and therefore most on women, fuperstitious and fearful perfons, fick people, children, and young creatures." " Their effects however fometimes fail to appear, becaufe they are encountered and overcome by the mind and fpirit, before they work any manifest effects." or becaufe his fe

not any secult of IX22

Such effects are obviated upon the fame principle, which establishes the prevention. of bodily difease : " for in infection and contagion from body to body (as for example, during the plague) the miasma may be received, but from the strength and good difposition of the body, it is expelled and wrought out, before it has had fufficient time to form the disease."

11111

CCXII.

CCXII.

It has been faid, and many are of the opinion, that the force of imagination doth often forward the end propofed. Thus for inftance, it has been put as a queftion, "whether a man, when he conftantly and ftrongly believes that fuch a thing fhall be (as that fuch a one will love him, and the like) helps any thing to the effecting the thing defired ?" Certainly not in the manner which has been advanced, namely, "by a *fecret operation* on the fpirit of another." If he fucceeds, it is either becaufe he perfevered, or becaufe his *perfeverance and earneftnefs* (and not any *occult operation*) makes him at length be attended to.

CCXIII.

There is not a doubt, but the force of imagination often gives energy to our actions. It may however, unlefs we are much on our guard, eafily delude us afide from reafon. It has been the tree which has yielded the fruits of fuperfition in former times, and which has often fed the human mind with

with the most extravagant notions of fympathy. Sympathies of this kind, *fuch as* the power of charms, and the like, are now pretty generally exploded.

CHAP. VI. SECT. II.

On the Sympathetic affections of the external senses.

CCXIV.

The five principal fenses, hearing, tasting, smelling, feeling, and feeing, are confcious of a fympathetic impreffion from odious objects. " I. A difagreable found will fet the teeth on edge, and make all the body fhiver. 2. The fwallowing of a naufeous medicine will be attended with a shaking of the head and neck. 3. Difagreeable fmells produce nearly the fame effect, which are lefs perceived, becaufe there is a remedy at hand by ftopping the nofe. 4. If you come fuddenly out of the fun into the shade, the sense of feeling is diffurbed by a chillinefs, or shivering of the whole body. 5. And even fudden darkness produces a propensity to fhivering."

G4

CCXV

CCXV. the fiom out dates

There is a very apparent reafon why a fympathy fhould take place between the eyes. Hence their motions are fynchronous. It may be faid, that cuftom and habit difpofe the eyes to move one and the fame way; " for when one eye moveth towards the nofe, the other eye moveth from the nofe."

CCXVI.

Though the eyes are by nature prone to move in concert, cuftom will however deftroy this natural confent, and produce the contrary. Thus fome people will fquint when they will. I would therefore give this caution to mothers and nurfes : " let them not fuffer infants to fit with a candle placed behind them, for both their eyes will difpofe to move outwards, as affecting to fee the light of the candle, which may bring on the .habit of fquinting."

nove baA . 2 . . b CCXVII. of no ganavidt

It appears as a quality in the fenfes of hearing and feeing, "that the inftrument of each

each feparate fenfe has a fympathy and fimilitude to that, which giveth the reflection." Thus it has been obferved, " that the eye will fympathize with a chryftal glafs, or water, and the ear with caves and fuch hollow places as are fuited to report echo."

jects operate in aIHVXOO way, in produc-

Sympathies have been compared to unifons of found in mufic. (LVIII.) Unifons of found produce agreeable fympathetic feelings; the reverfe produce difagreable feelings, (CCXIV.) "All concords and difcords of mufic are (no doubt) fympathies and antipathies of found." Moreover, " they are faid to work as well by report of found as by motion."

CCXIX.

The fenfe of *feeling* may be diffurbed by any uncommon, though apparently flight, irritation. Thus tickling the fides, or foles of the feet, will caufe *laughter* : and again, tickling the noftrils will raife *fneezing*, and on a fudden wonderfully increase the fecretion of tears. Both these *operations*, as fympathies,

thies, tend to remove both caufe and effect, " by producing a fudden emiffion of the fpirits," and the expulsion (if there should be any) of the offending matter.

CCXX.

The moft agreable as well as odious objects operate in a fecondary way, in producing those fympathetic impressions and actions, which they commonly give rise to. An increased fecretion of *faliva* often takes place at the fight of a favourite distribution and the running of water from a bottle, or otherways, will sometimes affect individuals, of a particular idiofyncrafy, with an invofuntary propensity to void urine.

CHAP. VI. SECT. 111.

On the fecret effects of Sympathy and Antipathy on the child in the womb.

CCXXI.

I have frequently had occasion to mention the great fympathy which, in various ftates of the animal machine, is difcovered between

between the uterus and other important parts of the body, fuch as the heart, ftomach, and head. Therefore the fecret effects of fympathy and antipathy on the child in utero fhould not be paffed over unnoticed, notwithftanding it has been faid in a former chapter that a child in utero is to be confidered as a more fimple animal than a polypus. (LXIII. LXVI.)

CCXXII.

I shall first lay before my reader the opinion of the very learned Bacon on this interesting fubject. " It has been obferved (fays he) that the diet of a woman with child doth work much upon the infant; as if the mother eat quinces much, and coriander feed (the nature of both which is to reprefs and stay vapours that ascend to the brain) it will make the child ingenious; and on the contrary fide, if the mother eat (much of) onions, or beans, or fuch vapourish food; or drink wine, or ftrong beer immoderately; or fast much ; or be given to much musing ; (all which fend or draw vapours to the head) it endangereth the child to become lunatic,

lunatic, or of imperfect memory; and I make the fame judgment of tobacco often taken by the mother." Particular cafes may have, no doubt, occurred in fupport of fome of these opinions.

CCXXIII.

There is too good reafon to imagine, that much injury may be done to the child in the womb, by the imprudent conduct of the mother, though fometimes innocently through her ignorance; and if things taken into the flomach can fo materially affect the unborn babe, is it not worth our while ferioufly and minutely to attend to the fubject, in order to difcover what may particularly advantage, or what may particularly prejudice, the child in utero, in order that we may confcientioufly fet about to obtain the one, or guard againft the other ?

CCXXIV.

Unborn children, though confidered as fimple animals, are no doubt liable to both acute and chronic difeafes, which prove often fatal

fatal to them. They are probably brought about by fome defect in the living principle belonging to the circulating and mufcular fyftems, which together conflitute fo confiderable and effential a part of the animal machine. I cannot confider them as under the influence of either the fentient or mental principles. It is on this ground, that I think they are to be looked upon as fimple animals,

CCXXV.

If I take the liberty of fetting afide the neceffity of either a fentient or mental principle before birth, and of thinking the living principle, or fimple life, is alone concerned in the animal œconomy of an unborn babe, it is owing to the following fact; that children in utero, whofe configurations deprive them of nervous influence, namely, fuch as want both brain and fpinal marrow, are as well grown in every other refpect, and have the fame quantity of motions before birth, as perfect and well formed children. This fact clearly proves that the mufcular fyftem has

has certainly a principle of motion, independent of nervous connection.

CHAP. VI. SECT. IV.

Sympathy sometimes diforders, and throws into confusion, the acconomy of human life.

CCXXVI.

Sympathy will often raife actions and affections, in direct contradiction to the eftablifhed law of the animal œconomy, that is, fhe will act against herfelf, and produce diforder in, when she ought to preferve the equilibrium of, the animal functions. She then injudiciously becomes an imitative faculty, from the acuteness of her fensibility, and in this way injures the habitation, which she was appointed particularly to preside over and preferve. Viewing her in this light we must consider her operations as inordinate and difeased.

CCXXVII.

In order to illustrate what I mean, I shall enumerate a few striking instances of such sym-

fympathies, as appear to me to be confiftent with the above fentiment, and which, in my opinion, cannot be viewed in any other light.

CCXXVIII.

The falling into epileptic and hyfteric convultions on any flight alarm, or at the relation of an affecting flory, appear to me to be inconfiftent with *natural fympathy*, and an affront to the dignity of human nature; when the fame takes place at the fight of a diffreffed and fuffering object, I view them as those inordinate actions, which arife from the weakness of human nature, at a time when she ought to command ability and ftrength of reason, to have affisted in the diffress she had just been witness to, and not have exposed herfelf to the necessity of receiving the same, if not greater affistance, from others.

CCXXIX.

I am inclined to place under the fame head of fympathy, as an *irregular imitative faculty*,

culty, those inordinate convulsive actions, which arife in one, or many perfons, on feeing another in the agony of an epileptic paroxysm. Frequent instances occur of knowing that individuals may thus be fympathetically affected; but the instances are rare, where the affections of many are at the fame time fimilarly operated upon by imitation. A noted instance of this once occurred at Haerlem Hospital in Holland, where many of the fame ward were feized with epileptic fits, as soon as the patient, who had been admitted for them, was feized with his.

CCXXX.

Again, it has been already faid, that the force of imagination operates moft on weak minds and fpirits, and therefore moft on women, &c. (CCX.) Such, from a fimilar reafon, are alfo moft readily influenced by tender and fympathetic feelings. It is on this principle, that pregnant women, when they are witneffes to the pangs of labour in another woman, very commonly will

will complain of feeling those pains, which are peculiar to the process of parturition; and I have been further told, and believe it to be true, that premature labour has been brought on, on the fame occasion, when it could only be accounted for, upon the supposed influence and authority of *imitative fympathy*. In short, we are such fympathizing creatures, that we must every now and then be exposed to all the ill confequences of idiopathic difease, owing to the power of fympathy.

CCXXXI.

I cannot conclude the fubject without obferving that the influence of fympathy even extends itfelf to the inanimate part of nature. Plants, metals, &c. fympathize with each other. The whole world was formerly fuppofed by the ancients to fympathize in all its parts. Their doctrines were however carried too far, and built upon fuperfitious principles, though they were occafionally fupported by many operations on the earth, and attractions between certain H parts

parts which conftitute it, and which were fuppofed to make, by confent of the univerfe, part of the diurnal revolution.

CCXXXII.

Human nature herfelf may be confidered as a *microcofm*, and is certainly regulated by the primitive laws and influences of the animal œconomy. Thus we, after the like manner, obferve, that alternate fleep and watching, fatiety and hunger, conftitute, though in part only, and by the confent of human nature, the diurnal revolution of the animal œconomy.

CCXXXIII.

We cannot, in fhort, enquire into any part of animate or inanimate nature, without difcovering the univerfality of fympathy and confent through all her operations. The attraction and cohefion of bodies in general amount to the fame general principle, and point out to us, that through all nature fympathy is the univerfal bond of union: fhe has ftamped her empire indelible on the mind of man, by actions complicated, affections wonderful,

wonderful, impressions innumerable, daily improving, strengthening, varying, and increasing: neither is she less to be admired where she has directed, as by instinct, the œcohomy of the brute creation; and when, by her attractive operations, she affects and alters inanimate matter, she then confirms her universal dominion.


CCXXXIV.

SHAKESPEARE knew well the power of fympathy and antipathy over the conftitution of man, when he made Shylock exclaim,

Some men there are like not a gaping pig;
Some, that are mad, when they behold a cat;
And others, when the bag-pipe fings i'th' nofe,
Cannot contain their urine; for affection,
Mafterlefs paffion! fways it to the mood
Of what it likes or loathes."

MERCHANT OF VENICE.

END of the FIRST PART.



TREATISE on SYMPATHY.

A


PART THE SECOND,

On FEBRILE SYMPATHY and CONSENT; and on BALLANCE and CONNECTION of EXTREME VESSELS, illustrated by *Practical Remarks*, and a new Explanation of the various Affections of the Stomach and Skin in Fever.

IN WHICH IS ATTEMPTED

A full Refutation of the Doctrine delivered on the fame Subject from the Practical Chair at the UNIVERSITY of EDINBURGH.

H 3



FEBRILE SYMPATHY

ON

AND

CONSENT, &c.

CHAP. I.

A concife view of the Theory of Fever.

SECT. I.

The author's original letter to Dr. Cullen, and the Doctor's answer to it.

CCXXXV.

BEFORE I enter upon the confideration of the fubject intended for the following part of the work, I think it proper for feveral reafons, as well as refpectful to my friend and mafter Dr. Cullen, to lay be-H 4 fore

20 ON FEBRILE SYMPATHY, &c.

fore my reader an exact copy of the original letter, which I fent to Dr. Cullen in Nov. 1777, and of his obliging anfwer. I had been induced to transmit my fentiments to him *in writing*, in confequence of his having invited, from the professional chair the preceding morning, his hearers, (among whom I was then one) freely to communicate their thoughts on, and objections to, any part of his newly advanced Doctrine of Fever, he having preferred *that mode* to a perfonal conversation, for which, he faid, he could not command leifure.

(C O P Y.)

"SIR,

Edinburgh, Nov. 30, 1777.

When I heard your very ingenious explanation of the Sympathy and Confent between the mufcular fibres of the ftomach, and the veffels on the furface of the body, which take place in fever, it occurred to me, that fuch might depend upon an atony and fpafm in the fuperficial veffels of the ftomach itfelf."

ON FEBRIL'E SYMPATHY, &c. 121

"It appears to me not lefs confiftent with your doctrines, and equally fo with the laws of the animal œconomy, to fuppofe, that the confent fhould take place between veffels and veffels, rather than between veffels and mufcular fibres. I would therefore refer the atony of the ftomach, corresponding to the atony on the furface of the body, to the minute veffels on its furface corresponding to the flate of the minute veffels terminating at the fkin, from which would arife a mutual and proportionate fpasm, if the debility be confiderable enough to produce it."

" If fuch be probable, I would humbly offer it as my opinion, that the action of vomiting is *folely* the effort of the vis medicatrix naturæ, operating to overcome the fpafm and obftruction in, and increase the action of, its fuperficial and fecretory veffels, to the fame end that the action of the heart and arterious fystem is increased, in order to remove the fpafm at the furface of the body, namely,

namely, the fpafm of the extreme veffels there terminating."

"I further prefume, Sir, to imagine, that the arguments which you have made use of in the XLIII paragraph of your FIRST LINES to prove the confent as you have laid it down, will equally explain and illustrate the above supposed valcular connection."

"I have taken the liberty, Sir, of communicating my thoughts to you in a letter, becaufe I think I fhall be lefs troublefome to you in this manner, than I fhould have been by having done it in perfonal converfation, for which, as you publickly declared from the chair, you have no leifure."

"I have the honour to remain, with the greatest respect,

SIR,

To Dr. Cullen, Your much obliged, Mint, Edinburgh. very humble fervant, SEQUIN HENRY JACKSON."

CCXXXVI.

CCXXXVI.

The foregoing letter of Nov. 30, 1777 remained unnoticed till July 20, 1778, on which day I received the following anfwer to it, but not without having troubled Dr. Cullen, a few days before, with fresh application by letter.

(COPY.)

"Dear Sir, Edinburgh, July, 20, 1778. On the other page I have given you thortly my opinion of your doctrine, and I would have you view it, as I would have all my opinions viewed, as given with great diffidence, and to be fubmitted by me patiently to the judgment of every body elfe."

"Wishing you success in all your ingenious speculations, and every part of your studies, I am with regard,

SIR,

Your faithful and obedient fervant, WILLIAM CULLEN." Second

Second Page of Dr. Cullen's Letter.

"When I fuppofe there is a confent between the flomach and furface of the body, I fuppofe it is a confent between the muscular fibres of the flomach, and the muscular fibres in the veffels on the furface, and fuch confent is not unufual in the animal œconomy."

"I cannot allow that the veffels of the fomach have any concern in this matter, as the phænomena of appetite and vomiting cannot, in my opinion, be referred to any flate of the veffels, but must be to the fibres in the muscular coat of the stomach."

"That the mufcular fibres in the veffels on the furface of the body may have a confent with the mufcular fibres of the flomach, appear to me fufficiently probable from hence, that the mufcular fibres of thefe veffels are affected by various flates of the nervous fyftem, as particularly appears from their

their being affected by the paffions of the mind."

WILLIAM CULLEN."

To Dr. S. Henry Jackson, Broughton, near Edinburgh.

CCXXXVII.

After receiving this letter, I felt myfelf fomewhat discouraged from indulging my fpeculations on the fubject any longer; but my turbulent spirit for enquiry into the philofophy of medicine still kept the afcendency over me, and would not fuffer me to reft long; fo that during the following winter, after having more attentively confidered Dr. Cullen's doctrines, I engaged in preparing the following part of this treatife, which has fince then received very few additions, and has of late only undergone a new arrangement of its parts. I have brought the period of viewing Dr. Cullen's doctrines to a conclusion with great diffidence, having at last, but not without ftruggles, furmounted that timidity, which has hitherto delayed the publication. 126 ON FEBRILE SYMPATHY, &c, of my treatife. I shall now commence the subject of fever.

CHAP. I. SECT. II.

Of the Phanomena of Fevers.

CCXXXVIII.

The phænomena of fevers are fo various and numerous, and at the fame time fo neceffarily connected together, that while I confine my attention to the inveftigating the morbid febrile condition of the function of the ftomach, it will no doubt appear to moft at first view, that I must have passed over many important circumstances connected with the doctrine of fever.

CCXXXIX.

No part of the pathology of fever appears to me fo imperfectly underftood, or more difficult of investigation, than the one I am about to engage in. Most of the other phænomena have been of late years very fatisfactorily explained by Dr. Cullen, our learned professor in

in the Univerfity of Edinburgh, agreeably to his own fyftem; but I have always thought, that his explanation of the febrile ftate of the ftomach, though extremely ingenious, was deficient: in confidering therefore the fubject in hand, I fhall unavoidably be led to offer my objections to that part of his fyftem, which treats of the fympathy and confent of the ftomach and fkin in fever.

CCXL.

When I am obliged to differ from fuch great authority, I do it with fear and diffidence; and yet from a firm perfuation, that any attempt, however feeble, to elucidate a difficult fubject, will be received with candour by the learned, I enter on it with pleafure, and will purfue it with freedom.

CCXLI.

Though I mean to confine my reflections more particularly to an illustration of the more common effects produced on the stomach by the febrile state of the system, yet I think

I think there is a neceffity of being more particular at first, and therefore, before I enter on the particular subject intended for these states intended for these states intended for these states intended for two others as preparatory, and on which the explanation in view will be founded; I mean the pathology of fever, and physiology of the states is the states of the states of the states is the states of the states of the states of which I am to raise my future fabric, and without which, it is probable, fome of my readers might be at a loss to discover the proper application of the reasoning I am hereafter to make use of.

CCXLII.

Fever is a flate of the fystem characterized by the following circumstances.

After an almost imperceptible diminution of strength in the animal functions, there comes on some degree of cold shivering, followed by an increase of heat, an increased frequency of pulse, and increase of the general debility; and though severs, on some occa-

occasions, put on different appearances with respect to the accession and duration of the phænomena, yet I think there is one general cause acting in the production of all.

CCXLIII.

By the modern pathologifts the paroxyfm of a fever, ex. gr. an intermittent, has been (I think juftly) divided into three ftages or periods, to wit, the cold, the bot, and the *fweating*: each of thefe may be faid to be characterized by its peculiar phænomena, which are certain occurrences in the courfe of the difeafe, conflituting when taken together the whole of the paroxyfm. Thefe phænomena I shall briefly enumerate, and mention them as they occur in the feries of caufe and effect, i. e. in the order of the cold, hot, and fweating stages. I begin then by the cold stage.

CCXLIV.

i. At the commencement of the cold ftage, the pulfe may be observed to be flower, and weaker than natural; as it increases, I the

the pulse becomes smaller, very frequent, and often irregular.

2. The refpiration is fmall, frequent, and anxious, and a fhort dry cough is fometimes attendant on it.

3. The function of the flomach is much diffurbed with anorexia, nausea, and vomitio, in other words, want of appetite, fickness, and vomiting.

4. There is much thirst, and the mouth and fauces are dry and clammy.

5. The urine is almost colorless without cloud or fediment.

6. The alvine difcharge is commonly impeded; or if there be an evacuation of fœces during the cold ftage, they are more coffive and dry, than during the health of the fyftem.

7. If there are tumours on the furface of the body, they are obferved, at this time, to be confiderably diminisched; and ulcers have also been dried up during the cold stage. (This last phænomenon there are frequent opportunities of seeing in perfons who have lately applied a blister to any part of the body, or who have issues, sec.)

8. The fenfibility of the body is often greatly impaired.

9. There is a difficulty of recollection more or lefs through the whole of the paroxyfm, often arifing to delirium.

10. Sleep and drowfinefs will fometimes attend this ftage, often to the degree of coma.

11. A head-ach is fometimes felt early in the cold ftage, with pains in the back, and there are more or lefs of general pains in all the flefh and joints; thefe changes in the various functions of the fyftem conftitute the I 2 cold

cold stage, which is of longer or shorter duration according to the nature and type of the fever.

CCXLV.

The cold ftage gradually and indeed fometimes pretty fuddenly changes into the hot ftage; this is characterized by the following phænomena:

1. From being flow, weak, and irregular, the pulfe has become more regular, hard, and full, and in thefe refpects increafes till the fweat flows; (from the increafed action of the heart and arterious fyftem, the pulfe in fever runs further along it than at other times, and hence a pulfation is fometimes to be felt at the tips of the finger. This is very remarkable in child-bed fevers, and when confiderable is a dreadful fymptom.)

2. The refpiration is now more full and free, but still frequent and anxious, with fome continuation of cough.

3. The

3. The anorexia still continues, and the vomiting will fometimes be violent at the beginning of this stage, but abates as it advances.

4. The thirst continues from the heat now prevailing.

5. The urine becomes high colored, but still continues without sediment.

6. The belly still remains bound.

7. Tumours are yet fometimes diminished, and ulcers still dry.

8. The fenfibility is now recovered, and is often confiderably encreafed.

9. The head-ach, though fometimes felt early in the cold stage, more commonly is not felt till the hot ftage be formed, and then it is ufually attended with a throbbing of the temples. When the head-ach comes on, whether fooner or later, there are generally pains in the back or fome of the great joints. This 1 3

This ftage has no limited duration, and by infenfible degrees changes into the following, viz. the fweating ftage.

CCXLVI.

In this stage the changes that occur in the various functions of the body are striking.

1. The pulfe is now become fofter and lefs frequent, and when the fweat ceafes returns to its natural ftandard.

2. The anxiety in the refpiration is relieved, and it is now lefs frequent and freer, and unaccompanied with cough.

3. There is most commonly a total ceffation of the nausea and vomiting.

4. The mouth becomes moifter, and the thirft gradually abates.

5. The urine now deposits a fediment, commonly lateritious, the fweat having flowed freely,

6. Till

6. Till towards the end of the paroxyfm ftools feldom occur, and are commonly after the first evacuation lax.

7. Tumours on the body are now increafed, and ulcers will again discharge matter.

8. And, laftly, there is an abatement of the head-ach, and pains of the back and joints; and when the fweat goes off they commonly ceafe all together.

CCXLVII.

Sympathy, in my opinion, lays the foundation of thefe various changes; (CCXLIV. CCXLV. CCXLVI.) parts fimilar fympathize at one and the fame time with the ftomach, the grand fympathifer, and fountain of fympathy (Part the First, Chap. II. Sect. V.). By the state of this viscus, as a fecreting organ, is regulated the state of all the other fecreting organs in the system, as will, I flatter myself, evidently appear in the course of the subsequent pages. If then

it

it can be made to appear, that fuch influence may be allowed to the flomach over the conftitution at large, there is just room to fuspect, that the proximate cause of the whole phænomena first takes place in the flomach: allowing that some of the principal remote causes of fever are capable of primarily acting, from their nature, on the stomach, which I think they may do. Hence arises its first derangement, of which impaired appetite generally gives the alarm.

CCXLVIII.

Fevers have been divided into the intermittent and continued; and this divifion has a real foundation. There are others called remittent, but thefe are not readily comprehended. For a particular account of thefe varieties, and their origin, I beg leave to refer the reader to Dr. Cullen's First Lines on the Practice of Physic, page 18 to page 24 *. It is unneceffary to the particular fubject in view to dwell longer on this part of the pathology; I shall therefore

* See also his Chapter on the Difference of Fevers and its causes, page 47 of the First Lines. ON FEBRILE SYMPATHY, &c. 137 fore proceed to confider the immediate and efficient caufe of the forementioned chain of phænomena,

CHAP. I. SECT. III.

Of the proximate cause of Fever.

CCXLIX.

Proximate caufes have always been a favourite purfuit with medical philofophers; and though the *hypothefes* founded upon them fometimes want fupport, they may ftill be confidered, in as far as they may tend to illuftrate the fubject, as ufeful delufions. There feems to be, in my opinion, great improvement made in this part of the fubject by the attentive obfervations of Dr. Cullen, who has juftly laid afide the former doctrines of *Lentor* and *Vifcidity*, and of *morbific Matter*, and has adopted a fyftem of his own more confiftent with the phænomena of fever.

CCL.

CCL.

This learned profeffor's idea of fever is as follows; "That a fpafm of the extreme veffels, however induced, may prove an irritation to the heart and arteries," and that fuch is to be confidered as a principal part in the proximate caufe of fever. (Dr. Cullen's Firft Lines, paragraph XL.) But moreover, "he is led to believe, that, together with the fpafm, there is an atony fubfifting in the extreme veffels, and fupporting the fpafm affecting them." (Firft Lines, par. XLII.)

CCLI.

For fome illustration and proof of this doctrine I must refer my reader to his Chap. on the Proximate Caufe of Fever (page 24 of First Lines). Dr. Cullen however observes, " that it will still remain a question, what is the caufe of this spasm, whether it be directly produced by the remote causes of fever, or if it be only a part of the operation of the vis medicatrix naturæ." (F. L. par. XL.) He is disposed to be of the latter opinion. Resting then satisfied with this short account of the

the proximate caufe, I shall next confider the remote causes.

CHAP. I. SECT. IV.

Of the remote causes of Fever.

CC.LII.

The operation of these appears to me to be ftill involved in some obscurity. Dr. Cullen has hinted it as a doubt, whether they act directly in producing the spass. (CCLI.) I cannot but entertain a confused notion, that they may act topically, both on the fentient extremities of the nerves, and the extreme vesses of the serves, and the extreme vesses of the serves in the serves and in this way, by producing the atony, lay the foundation of fever: and as they are not all of the same nature, and do not posfess the same specific properties, so some more readily may produce bad effects externally, others on the contrary, internally. (Last Part of CCLXVII.)

CCLIII.

Dr. Cullen speaks thus of the action of stimulant and sedative powers upon extreme vessels:

veffels: "The tone and action of the arteries, as a mulcular part, may be increased by ftimuli immediately applied to them, or by the increased force of the nervous and animal powers with regard to them; and they may be diminished by fedative powers applied to them, or by weakening the nervous and animal powers." (Cul. Phys. par. CLX.) On such principle fome of the remote causes of fever feem to me to act.

CCLIV.

The principal remote caufes of fever are, contagion, miafmata, and cold: and I beg leave to add, excefs, or imprudent indulgence, of the appetite. Occafionally fear has been known to throw the fyftem into a febrile ftate; they have all been confidered as of a fedative nature, though cold is well known to have both ftimulant and tonic effects on fome occafions. Whatever their operation may be, they (fear excepted) feem to me as likely to act topically as in any other way, and I am inclined to confider both the general debility, and fubfequent rc-action of the

vehicls:

that are the

the fystem, as effects of the operation of the vis medicatrix naturæ, agreeable to fome established law in the animal œconomy.

CCLV.

Fear, confidered as one of the depressing paffions of the mind, has been fuppofed by Dr. Cullen primarily to affect the cutaneous extreme vessels, in confequence of which the ftomach through fympathy is difordered. * But it would be most natural to fuppofe, from the channel through which fuch a caufe must operate, that the brain and nervous fystem are primarily affected, and that the heart and arterious fystem become diffurbed, by that neceflary and mutual dependence exifting between these two fystems. The extreme minute parts of both thefe fystems; namely, the fentient extreme nerves and capillary arteries, neceffarily accompany each other, and are allowed to be highly irritable. No wonder then, from fuch connection and irritability, at the difturbance that takes place.

CCLVI.

* See Dr.: Cullen's letter to the author (par. CCXXXVI).

CCLVI.

This flort account of the pathology of fever I fhall conclude with a paragraph from Dr. Cullen's First Lines, in which the whole of his doctrine is fummed up in a general way, and shews the connection of its different parts.

" The remote caufes of fever are certain fedative powers applied to the nervous fyftem, which diminishing the energy of the brain, thereby produce a debility in the whole of the functions, and particularly in the action of the extreme veffels; fuch, however, is, at the fame time, the nature of the animal occonomy, that this debility proves an indirect ftimulus to the fanguiferous fystem; whence, by the intervention of the cold stage, and spasm connected with it, the action of the heart and larger arteries is increased, and continues fo till it has had the effect of reftoring the energy of the brain, of extending this energy to the extreme veffels, of reftoring therefore their action, and thereby especially overcoming

coming the fpafm affecting them; upon the removal of which, the excretion of fweat, and other marks of the relaxation of veffels take place."

This fummary of the doctrine may be confidered, as an account of the rife and progrefs of a febrile paroxyfm, the whole phænomena of which evidently depend, according to Dr. Cullen's view of the fubject, upon debility, fpafm, and re-action. I next fhall proceed to the phyfiology of the flomach, on which I propose to be very brief.

ATTRACTOR STATISTICS BO

the bit brother () as a star fragment of

CHAP.

144 ON FEBRILE SYMPATHY, Seci

CHÁP. II.

Of the various affections of the Stomach.

SECT. Í.

Of the Physiology of the Stomach:

CCLVII.

THE stomach may be confidered as the distinguishing characteristic between the animal and vegetable kingdoms. The function of it in the conversion of our aliment is certainly involved in much difficulty and darkness, notwithstanding the many ingenious experiments that have of late been made, with the view of throwing light upon the subject. It was subjected to many various opinions among the ancients; I shall fay little more than, in a few words, is necessary to give my own opinion on it, as far as the structure of the parts affists me, and the

the analogy of other operations in the fyftem corresponds.

or such galirie, liquor, the exiftance of

which, I imagin JIIV JOD cely to depend

The flomach is a mulcular bag, the internal furface of which is lined with a very villous and valcular membrane, and furnifhed *alfo* with numerous fimall glands. Many very different hypothefes have been offered to explain its action, which being fo well known, it would be idle fully to enumerate; and I fhall therefore only obferve that, "Some have compared it to a mill, others to a flewing pot, others to a worttrough, when all the while it muft have appeared, that it was neither a mill, nor a flewing pot, nor a wort-trough, nor any thing elfe, but a flomach *."

ovad ware spinler of CCLIX.

The change produced on our aliment, in all probability, depends upon the fecretion of a menftruum peculiar to the ftomach, K and

Dr. Hunter in his Introductory Lecture.

and endowed with properties fuited to that end. Many experiments prove the prefence of fuch gaftric liquor, the exiftence of which, I imagine, most likely to depend on a due tone and action in the vafcular ftructure of the stomach. In what particular way the menstruum acts upon the folid aliment, experiments have not yet quite fo fatisfactorily shewn, as is to be wished; it is probably affisted by some peculiar action in the stomach itself, for we cannot imitate the folutions performed in the stomach, by the fecreted fluids taken out of it.

CCLX.

On the whole, the action of the flomach upon the aliment appears to me principally to depend upon a due flate of action in the vafcular flructure of the villous membrane, which confifts both of an exhalent and fecretory fyftem of veffels. They may have the effect of loofening the fixed air of the alimentary matters, as by fome authors have been faid (Cull. Phyf. CCXXXVI.) and thus prepare them for the putrefactive and acefcent

acefcent fermentation ; but of this, and other opinions, it is not neceffary to fpeak.

CCLXI.

The remarks thus far made, I confider, as pointing out the primary end and function of the ftomach, as life could not long exift without its manfion being daily repaired by frefh fupplies : but its relative and fecondary use is the influence it exerts over the fystem at large. It is the feat of fympathy; the throne of fensibility, to which all the other functions of the fystem look up; its long acknowledged influence is attested by the records and opinion of the ancients.

CCLXII.

The flomach had been by them obferved fo frequently to be affected by fympathy with many other parts of the body, even with the brain, that they even went fo far as to fix the feat of the foul in the *cardia*, or upper orifice of the flomach. Their opinion was further confirmed from obferving, that a blow on the pit of the flomach would fome-

K 2

times,

times, even fuddenly, prove fatal, and this without doing any apparent injury to the heart or its large veffels, or any of the neighbouring parts. This fingular incident feems therefore to argue a very high fenfibility in the ftomach, and a mutual dependance between it and the vital functions. (LXVIII. LXX.)

CCLXIII.

I hope my reader will be fatisfied with thefe few obfervations on the ftomach and dgieftion; I fhall now immediately proceed to confider the febrile affections of the ftomach, to wit, the *anorexia*, *naufea*, and *vomiting*, and I hope to render it probable, that they, as part of the phænomena formerly mentioned, may depend upon the fame debility, fpafm, and re-action, as the other febrile affections.

the brain, that they even went to far as to

furthet coultmed from obletving, that a

he feat of the foul in the cardid, or upper

rea of the flamach. Their opinion was

12012137

-anto bluew donnoft and lo. tig CHAP.

CHAP. II. SECT. II.

Of ANOREXIA as a febrile affection of the stomach.

CCLXIV.

The affections of the ftomach take place in the order in which I have juft now mentioned them, and their violence will depend upon the degree of the aforementioned ftates of the whole fyftem, I mean, the *debility*, *fpafm*, and *re-action*; thus the greater the debility, the ftronger will be the fpafm, and proportioned to the latter will be the fubfequent re-action, and thus to the violence of thefe general principles in the conflitution, the febrile affections of the ftomach bear a relative proportion. That this may appear evident, I fhall mention the connection of the feveral affections of the ftomach with the general febrile ftate of the fyftem,

CCLXV.

On the first approach of fever, but particularly when a fense of cold is felt in any K 3 part,

part, the appetite fails, and generally continues, through the whole of the fever, impaired, and even averfe to all folid food; (CCXLIV.) while the coldness increases, and the patient becomes affected with tremors, and fuccuffions of all the different parts of the body, the anorexia is changed into, or rather accompanied with, nausea, and if the cold stage be violent, fuch ficknefs will end in vomiting, which fometimes, from the violence of the cold ftage, will be of a bilious nature. As the vomiting is in general most fevere at the height of the cold stage, fo it will fometimes continue even at the beginning of the hot ftage, (CCXLV.) but when this is completely formed, the ficknefs and vomiting abate, and will ceafe altogether on the appearance of fweat. (CCXLVI.) debt, I fhail mention the

CCLXVI.

The occurrence of these affections of the ftomach, at the very fame time that all the other fecretions are impeded, namely, the faliva and mucus in the mouth, the urine, fweat,

fweat, and alvine discharge, &c. leads me greatly to sufpect that the fame cause, which obstructs the action of their respective organs, similarly affects the function of the stomach, impeding thereby its operations, and disturbing the digestive process. It will therefore, in the next place, be proper to constider, whether the cause alledged for the impeded perspiration, and glandular secretions in general, is sufficient to explain any of the morbid affections of the stomach.

CCLXVII.

I fhall take it for granted, that a debility of the nervous fyftem lays the foundation of fever (though I have my doubts about it, CCXLVII.) in confequence of which debility the action of the heart, and probably that of the larger arteries, is much weakened, fo that a fpafmodic conftriction takes place on all the extreme veffels, which will very readily account for the fuppreffed perfpiration. But while it is admitted, that fuch fpafm takes place on the fkin, it cannot be denied to exift in many other parts of the K 4 body;

body; the extreme veffels terminate on all the furfaces of the body, both externally and internally, and accordingly are numerous throughout the first paffages as well as on the exterior furface of the body; in the former, as well as the latter, they may be faid to be external with respect to the fystem at large, and are to be confidered as very fingularly numerous on the furface of the stormach, on account of its villous structure, where they may be faid to be exposed to the action of feveral external causes.

CCLXVIII.

Every one will allow, that there is fufficient proof of fuch a fyftem of veffels being prefent in the ftomach, from the frequent inftances of large ferous fecretions, or, more properly fpeaking, effufions, which are collected there, and often evacuated by vomiting. If therefore there are juft grounds for fuppofing a fpafm to be formed on the furface of the body, it is but reafonable to conclude, that the fame takes place in the ftomach, and even through the whole of the inteffinal

testinal tube; and if the former is in the extreme vessels of the skin, I may, from analogy, be allowed to suppose, that the latter depends upon a similar affection of similar vesfels in the first passes.

bnadap Aum Aur un CCLXIX.

A question here naturally presents itself: Will the action of the ftomach upon our aliment be affected by fuch a morbid ftate of its veffels? I should certainly think it would, if there are just grounds for fuppofing digeftion in any way to depend upon a due tone and action in its exhalent and fecretory fystem. (See Dr. Cullen's letter to the author, CCXXXVI.) Befides the effusions of ferum from the usual terminations of the arterious fystem, physiologists have agreed, that there is a fuccus gastricus proprius; that a spasm will affect the former, is, I hope, rendered fufficiently probable, from their immediate connection with the circulation; but will the febrile state of the system disturb the fecretion of the latter ? If we admit the force of analogy, we shall readily allow that it must be disturbed.

CCLXX.

teffinal tube ; an.XX.133 former is in the

In all other parts of the body fecretions are observed to be diminished by the weakness of the fanguiserous system, at the commencement of sever; so that if the secretion of the *fuccus gastricus* does in any way depend upon the state of the arterious system, in giving momentum to the blood, it must, when this is debilitated, suffer the same injury, which affects the other secretions.

would, if there .IXXLOO mds for fuppoling

If then I have fucceeded in my endeavours to make it appear, that appetite depends upon a due tone in the fecreting organs of the ftomach, and due quantity of the *fuccus gaftricus*, we can, on fuch principles as the forementioned, readily account for the *anorexia*, which fo conftantly attends the febrile ftate of the fystem.

Hiw and phoing CCLXXII.

Thus far I have only endeavoured to render it probable, that the primary debility of the fyftem, which must produce an universal constriction

confiriction of the extreme veffels, lays the foundation of an atony, with more or lefs of fpafm, in the exhalents of the ftomach, and caufes at the fame time a diminution in the produce of the gaftric liquor, on which two I have fuppofed the condition of the appetite to depend. I fhall next proceed to confider *naufea*, the fecond morbid effect which arifes in the ftomach, from the febrile condition of the fyftem.

CHAP. II. SECT. III.

hope to they, when liceme more fully to

Of NAUSEA as a febrile affection of the stomach.

CCLXXIII.

Naufea has been defined by three nofologifts, SAUVAGE, LINNÆUS, and VOGEL, " a vain attempt to discharge by the mouth " the contents of the stomach," and is faid to be accompanied with flatulence. This last fymptom however is far from being constant, and perhaps only attends the chronic affections of the stomach, such as for instance, the dyspeptic. I am inclined to consider the nausea

nausea in fever, as little more than a greater degree of the anorexia, arising from an increafe of atony in the excretory and fecretory veffels on the villous membrane, and combined, probably, with more or lefs of a spafmodic constriction in them; I do suppose that this confiriction proves an irritation to the fentient extremities of the nerves, and caufes that uneafy fenfation, which gives rife to the action of vomiting, which action, I hope to fhew, when I come more fully to confider it, may be justly confidered as an effort or exertion of the fystem to produce a falutary end, namely, that of reftoring the morbid condition of the vafcular and villous membrane in the ftomach to its natural healthful ftate, SAUVAGE. LINNAUS, and VOCEL,

VIXXI23 s by the mouth

Naufea and vomiting may arife from caufes acting immediately on the ftomach; thus emetics in general, and acrid ingefta operate; but in the cafe of fever they fhould be confidered as effects fecondary, and dependent upon the ftate of the circulating fyftem. They

They are then produced on the fame principles as when they follow venæfection. When a perfon is bled a *deliquium animi* fometimes becomes the confequence, attended with an univerfal *atony* on the outer furface of the body, which I prefume to be the condition of the veffels of the fkin from the flowing of the perfpirable lymph at that time. A concomitant *naufea* and vomiting take place.

CCLXXV.

It appears to me, that this altered flate of the flomach, upon opening a vein, depends greatly upon the depletion of the fanguiferous fyftem, and debility produced in the action of the heart and arteries : the tenfion in the veffels of the brain before bleeding, being by it removed, produces the *deliquium animi* : the *fudor* and *naufea* are owing to a morbid alteration taking place in the arterious extreme veffels : while the *vomiting* is to be confidered as a falutary effort of the flomach, arifing from its univerfal fympathy with the fyftem, to reftore the due tone and action of the extreme veffels: the good

i misfirit

good effects of which in fuch a cafe I beg leave to explain as follows.

CCLXXVI.

The fystem, from the loss of blood, has been disturbed by the change of tension in the circulation; the action of vomiting, while it roufes the nervous fystem, ferves to fupply the depletion of the fanguiferous fystem, by forwarding the contents of the thoracic duct and lymphatic fystem, as well as of the venous fystem, into the arterious fystem, and thus restoring the tenfion of the whole: and I cannot help throwing in here a reflection, though I want observation sufficient to authorize me, and I acknowledge it is merely conjectural; it is the following. When we mean by venæfection to diminish the volume of the circulating fluids, I think we fhould avoid, if poffible, nausea and vomiting fubsequent to it, as I much fufpect that the latter may have obviated the benefit that was intended, or deftroyed that already effected, by the fupply it is capable of affording to the circulating fystem;

fystem; and therefore I throw it out as a hint, whether it may not be proper, where we intend to employ both, to order the emetic before the bleeding, unless any confiderable *plethara* be prefent.

CCLXXVII. I tall to stimil

So much for nausea, as bringing on the action of vomiting; I shall now proceed to consider the latter more fully, the end it is capable of answering, and the agents by which such end is produced.

CHAP. II. SECT. IV.

provided fuch confequences are obferved to

Of VOMITING as a febrile affection of the stomack.

alore in the cole

CCLXXVIII.

I have feveral times obferved, that I confider vomiting as a falutary effort of nature, a vis medicatrix naturæ: but as this term, though well known in all the fchoolsof phyfick, has not yet been precifely defined, nor its limits marked out, I shall briefly explain

plain the fense in which I mean to use it, before I go on to apply it to the action of vomiting.

- mos vers claim CCLXXIX.

Whatever may be the just definition and limits of that law of the animal œconomy, to which physicians have given the name of vis medicatrix naturæ, I am difpofed to apply it to every change, or exertion of the fyftem, which feems capable of producing good confequences, by removing a previous morbid state in some one part of the system, provided fuch confequences are observed to be frequently, although not regularly, the event of fuch exertion. We have a ftriking inftance of fuch exertion and its confequences pretty conftantly taking place in the cafe of fever, where an interruption to the action of the extreme veffels, or in other words, a constriction of them, is the difease to be overcome; fuch confiriction probably acts as a fimulus to the circulating fystem, in confequence of which its action is increased ; this action finally ferves to remove the affection of

of the extreme veffels, and reftore the healthful balance of the circulation. From hence the *pyrexia* has been confidered as a vis medicatrix nature.

CCLXXX.

I shall now apply this reasoning to the circumstances of vomiting. A caufe which produces the forementioned affection of the external extreme veffels, appears to me capable of affecting alfo the internal extreme veffels at the fame time, without fuppofing any neceffary confent of parts to exift. Now because the internal parts are of more importance to the fystem than the external, and therefore, when morbidly affected, expose to more danger, we may eafily conjecture why, in the cafe of fever, vomiting should be produced. The increased action of the heart and arteries may not alone be fufficient in every cafe, namely, in fuch for instance as are attended with a violent cold stage, to remove the affection of the internal parts fast enough; in order therefore to obviate, or guard against fuch deficiency, the Supreme Being L

Being feems to have implanted in our conflitution an effort to remove this internal affection of extreme veffels by the action of vomiting, which, while it operates immediately in removing fuch affection, ferves alfo to rouze to action the heart and arterious fyftem.

CCLXXXI,

We fee fuch an effect produced in fever, where the whole of the phænomena evidently evince a general spasmodic constriction of the extreme veffels, both on the fkin and primæ viæ; now unlefs this be removed, it must prove fatal, but the only power capable perhaps of removing it, is the increased action of the heart and arteries, aided, nay probably began, by the impulse of vomiting. These falutary efforts appear to me evident and neceffary confequences flowing from the laws of the animal æconomy, and as fuch I would call them the vires medicatrices naturæ, without going fo far as the Stablians did, who fupposed them the operations of an intelligent agent. (XXXVI.) CCLXXXII. 31.10

CCLXXXII.

But fuppofe for a while, that fuch a confriction of veffels does not exift, and that there only takes place an atony in them, let us fee if we can reafonably confider the vomiting as capable of removing it. A very celebrated phyfiologift of the prefent time has delivered it as his opinion, that the ftomach fympathizes with all parts of the body; (LXVIII.) that when an universal debility of the fystem takes place, vomiting is more or lefs prefent, as the confequence of the faid fympathy; and that vomiting is as an index of a general debility, and often the means of relief to the fystem, performing in the first sense of the expression, the office of a vis medicatrix naturæ. (CCXLVII.)

Hence I would conclude, that as it rouzes the circulation, it must also excite to action the fystem of capillary vessels, and thereby remove the atony which becomes the foundation of the constriction.

L2

CCLXXXIII.

CCLXXXIII.

On the above principle long continued hæmorrhages have been fuppofed to have been ftopt by vomiting, which in fuch cafes is frequently observed to occur; a laxity of the bleeding veffels being confidered as a fupporting caufe of the hæmorrhage: for the action of vomiting, by roufing the circulation, and giving the difpofition to contraction in the veffels, ferves to put a ftop to the effusion of blood. The stoppage of uterine floodings may very justly be attributed to fuch a caufe; and I can affirm from experience, what many have before obferved, that vomiting commonly attends thefe when in excefs. If then fuch a pathological conjecture be well founded, we may also confider the vomiting in fever as capable of obviating the fpafm, by removing the previous atony of the veffels.

CCLXXXIV.

The end then which the action of vomiting answers, is that of restoring the healthful

ful ftate of the ftomach and alimentary canal, and the proper balance between the internal and external extreme veffels, all which, as performing the office of excretories and fecretories, muft be fimilarly affected with the other fecerning organs. Sympathy, I contend, bringing about thefe falutary changes; and may not the want of due fympathy, between the ftomach and parts morbidly affected, be a probable caufe, why many difeafes become lingering, and at laft produce *hectic* fever in fome, *leucophlegmafia* in others, *hemiplegia* and *paralyfis* in others, and the like ? But much riper reflection than mine muft eftablifh fuch hypothefes.

CCLXXXV.

The immediate effect of vomiting muft be that of expelling from the flomach its contents, which, if fuffered to remain in the first paffages at the attack of fever, would prove a *febrile fomes*, because they would not undergo the change appointed by nature, by reason of the impeded action of the digestive organs. Hence we see not only the L_3 utility

utility of their expulsion, but of the loathing of food and anorexia which comes on, and which continues more or lefs till the ftomach has recovered its action, by the removal of the impediments to digeftion.

CCLXXXVI.

At the time that this is effected, the ventricular mucous glands are emulged, and the fpafm of the internal extreme arterious veffels probably overcome, by the particular action of the ftomach itfelf, which fome are of opinion is alone fufficient to expel the contents of the ftomach.

CCLXXXVII.

Dr. Hunter has publicly delivered it as his opinion, that during vomiting the contents of the flomach are thrown out by the contraction of the flomach itfelf; and he concludes fo becaufe he has found the bladder fo contracted as to have expelled the laft drop. But I confider vomiting as a more general operation of the fyftem, which gives a flimulus to the remoteft parts of the

the circulation, whereby the conftriction of extreme veffels externally is also often removed.

CCLXXXVIII.

Nature feems to have been particularly anxious and attentive in guarding us againft any permanence of a difordered flate of the ftomach, by implanting in it fuch a fingular fenfibility, that, when affected as in fever, and feveral other diforders, it feems of itfelf capable of removing the morbid condition by its own innate powers. In fuch a light I am inclined to confider vomiting, which is a convultive action of the ftomach, affifted by the exertions of many other mufcles.

CCLXXXIX.

Vomiting in fever has been hitherto fuppofed to depend upon a change of tone, or indeed loft tone, in the mufcular fibres of the ftomach itfelf, and confidered as a fpafmodic affection neceffarily taking place upon fuch previous atony; add to this, that the fuppofed debility of the ftomach, which L4 produces

produces vomiting, has been conjectured to depend upon an atony of the extreme veffels on the external furface of the body. (See Dr. Cullen's letter to the author, parag. CCXXXVI.)

CCXC.

If we reflect but for a moment, we shall find, that the phænomena of vomiting do not at all correspond with the explanation hitherto given, of the febrile affections of the ftomach, not even with the very ingenious conjectures of that celebrated profesior at Edinburgh Dr. Cullen; for, when vomiting takes place, not only the mufcular fibres of the ftomach itfelf are thrown into a convultive action, but all the mufcles of refpiration are fpafmodically affected. and more or lefs every mufcular fibre in the body; therefore to make all the phænomena confistent with Dr. Cullen's doctrine, it should have been laid down, that all these parts fo acting are in fympathy and confent, and confequently each part, and likewife the ON FEBRILE SYMPATHY, &c. 169 the whole, dependent upon the condition of vefiels at the furface of the body.

CCXCI.

That fluggifhnefs of the whole body, and its unwillingness to exert any part of the muscular system at the commencement of fever, feem to me to be indications of the general effort intended to be made by the whole fystem, and are preludes to the action of vomiting, as if nature, incapable of attending to her various functions at one time, withdrew for a while her accustomed influence over the mulcular fyftem as from the least important, to fix her watchful eye on the state of the stomach and heart, and finding them diffurbed in their functions, fat about removing the caufe by rouzing the whole mufcular fabric to action. (CCXC.) This reafoning is supported by the phanomena, which may be observed on the exhibition of an emetic, of which I shall fay more hereafter.

CCXCII.

and youdition

CCXCII.

Sicknefs and vomitings are ufual attendants on the firft months of pregnancy, and are marks of the general irritability of the whole fyftem; but they in a peculiar manner denote a ftrong fympathy between the *interus* and ftomach. They are in a general way removed by the act of quickening, yet when violent they have continued, and even caufed abortion.

CCXCIII.

Quickening has been attributed to the first motion of the child observed by the mother, but by practitioners it is suspected to be brought about by the fudden emersion of the *uterus* from the *pelvis* into the *abdo*men. Why the affections of the standard should then cease altogether, or even abate, it is difficult to explain.

CCXCIV.

Great debility, general and partial spafms, thirst, anasarcous affections, &c. have occured in

in cafes of violent vomitings, of which inftances are related in the IIId Volume of the Medical Obfervations and Enquiries published at *London*, 1767. But be the caufe whatever it may, that diffurbs the ftomach, fuch ftates of it denote its universal influence over the whole fystem.

CCXCV.

To conclude my remarks on vomiting, I beg leave to obferve, that the opinion thrown out in CCXC, refpecting the mufcular fyftem at large, and the general and *united fympathy* of all its parts during vomiting, not having been advanced either by Dr. Cullen, or any body elfe, as far as I know, I therefore do prefume to aver, that the whole may be juftly confidered as a general effort of the fyftem to remove the *fpafm* and *conftriction*, that muft neceffarily affect the veffels of the ftomach, as well as thofe on the exterior furface of the body.

CCXCVI.

CCXCVI.

I am now naturally led to confider a little fully, that fingular *apparent* fympathy and confent, which have been fuppofed to take place between the ftomach and fkin on many occafions.

the dor extreme

CHAP.

CHAP. III.

On the Sympathy and Confent between the flomach and skin in fever.

SECT. I.

How the SYMPATHY and confent depends upon the balance and connection of the extreme vellels, and not upon the condition of the muscular fibres of the stomach.

CCXCVII.

BEFORE I proceed, I think it proper to obferve, that thefe terms, fympathy and confent, in the fenfe here used, seem liable to some objections and restrictions, which I shall particularly point out by and by, and then propose others in their sead, which seem to me more proper. At present I shall use the old language, that I may not be misunderstood.

CCXCVIII.

CCXCVIII.

This fympathy is particularly evident in the cafe of fever, where it is obferved, that the impeded function of the ftomach is feemingly dependent upon the *ftate* or *morbid condition* of the fkin; but there are likewife other occafions on which we find that the active powers of the ftomach feem to be proportioned to a certain degree of tone and energy in the action of the extreme veffels terminating at the fkin. How this is I fhall now attempt to explain.

CCXCIX.

The pathological explanation of the fympathy in queftion, which first particularly claims my attention, is the one advanced by Dr. Cullen in his First Lines on the Practice of Physic, and is as follows. He has fupposed, that the affections of the ftomach corresponded to the atony and spass of the extreme vessels on the outer surface of the body, by a corresponding atony and spass in the muscular fibres of the stomach itself, arising from sympathy and consent. In the first

first place then, let me confider a little, if these muscular fibres of the stomach can be admitted to be in an atonic state.

CCC.

I have already observed, that at the time of vomiting, not only the ftomach, but all the muscles of respiration are likewise spasmodically affected; (CCXC.) this phænomenon alone, which shews the strong exertion of the whole fystem, evinces to me, that there is rather an increased tone and action in the muscular fystem at that time; or at least it be must be allowed that there is a general fympathy between all the mufcles which act in the time of vomiting, and the skin; and though there feems to be a general debility in the body, it may arife from an affection of the fenforium, in confequence of the debile flate of the vital organs; and this view of the general debility will be further rendered plaufible, by what I am hereafter to observe with respect to the effects of vomits.

CCCI.

CCCI.

When I confider the anatomy of the ftomach, together with that of the ikin, I cannot help reprefenting it to myfelf as improbable, that any fuch fympathy and confent should exift between parts fo evidently different in their ftructure and proportion, namely, between invisible veffels (for they may be faid to be fuch) and muscular fibres of confiderable fize; add to this, what Ihave already obferved, that the phænomena of vomiting do not at all agree with fuch fupposition, as vomiting does not fo much depend upon the action of the ftomach, as upon the exertion of the whole fystem, as is evident from the state of respiration, voluntary motion, &c. during vomiting. (CCLXXXII. CCXC.) Having thus flated my objection's to the olddoctrine, I shall proceed to establish a newone.

CCCII.

It is the fame confideration I made above, viz. that of the anatomical ftructure, which leads

leads me to think, that the *apparent* fympathy may be explained on another principle, which will be fupported and made probable, by the *phænomena* being confonant to one another. I have already fuppofed a neceffary affection of internal veffels, from the fame caufe which produces the affection of the external ones; and therefore I need go no farther, in order to explain the occurrence of fymptoms at the fame point of time, than to fuppofe, that the fympathy and confent takes place between veffels and veffels, and thefe too probably in every refpect fimilar, rather than between veffels and the mufcular fibres of a large organ.

CCCIII.

This explanation is firstly confiftent with the general doctrine of fever delivered by Dr. Cullen, at the fame time that it agrees better with the general laws of the animal ceconomy; I am therefore inclined to refer the correspondence of the atony of the stomach with that of the outer surface of the body to the correspondence of the state of M

the minute veffels on its furface with the ftate of the minute veffels terminating at the fkin, from which relation there arifes a mutual and proportionate fpafm, if the debility is confiderable enough to produce it.

CCCIV.

On this fuppolition I am led to confider the inordinate action of the ftomach which excites or rather produces vomiting, in conjunction with other mulcular parts fimilarly affected, as not proceeding from an atony in the mulcular fibres of the ftomach, but as folely depending on the operations of the *vis medicatrix naturæ*, exerted to overcome the fpafm and obftruction in, and increafe the action of, its fuperficial and fecretory veffels, for the fame purpofe, that the action of the heart and arterious fyftem is increafed, in order to remove the fpafm at the furface of the fkin, namely, the fpafm of the extreme veffels there terminating.

СНАР.

CHAP. III. SECT. II.

How the balance and connection of the extreme veffels, giving the confent between the stomach and skin, may be illustrated by the operation of medicines, as well as by the general phanomena of fever.

CCCV.

I shall now attempt some further proof and illustration of the foregoing doctrine; and that I may not conceal any thing from my reader, let me offer one obfervation to his candour, which, I hope, tends rather to support than destroy the hypothefis I have advanced. It is this, that though I am contending for an affection of veffels only, yet it ought to be confidered that these veffels must every where be fupposed furnished with muscular fibres, and that therefore if a fympathy could poffibly exift between mufcular fibres alone (fee Dr. Cullen's letter to the author CCXXXVI.) in diftant parts of the body, it would feem more likely to take place between those fibres belonging to Ma parts

parts fimilar, rather than to parts diffimilar, i. e. rather between mulcular fibres of internal veffels and external veffels, than between those of the ftomach itself, and those of veffels at a diffant part of the body, to wit, at the fkin.

CCCVI.

To proceed however with the further proof of my doctrine, I think, in the first place, it may be observed, that the affections of the ftomach take place more or lefs, according to the degree and extent of the atony prefent, or of the spasm formed in confequence of the atony, which furnishes a prefumptive proof, that the difordered state of the stomach depends upon this vafcular connection, and condition of the cutaneous veffels; for to what can we fo readily attribute the morbid flate of the flomach, as to an affection of its vascular ftructure. when we fee it fo conftantly take place at the time, that the other phænomena of fever evidently evince a general affection of the exhalent and fecretory fystems, and when it feems to particularly proportionate to the extent

extent of the morbid ftate of the veffels at the furface of the body. (CCCII. CCCIII.)

CCCVII.

Secondly, The fudden operation of emetics, as well as the peculiar effects of naufeating dofes, feem further to fupport this affection of veffels. Their operations have been hitherto explained by pathologifts, even by Dr. Cullen, on the principle of their affecting the mufcular fibres of the ftomach. But I cannot fee how an external ftimulus can be particularly applied to the mufcular fibres alone, without first affecting those more irritable parts of the villous membrane, to which they are immediately applied.

CCCVIII.

When the emetic matter is of a fedative nature, it may operate on the *fentient extremities* of the nerves in the flomach, and by fympathy weaken the general circulation, and thus form a fpafm on the extreme arterious veffels every where, which probably would be too permanent, if not M_3 quickly

quickly removed by the exciting efforts of the vis medicatrix in producing vomiting, to expel the noxious emetic matter, and prevent further mifchief.

CCCIX.

Effects nearly fimilar take place when the emetic matter is ftimulant, but not with fuch figns of general debility. The emetic cannot then be fuppofed to produce vomiting by caufing an atony in the mufcular fibres. In either cafe therefore it appears to be the operation of the vis medicatrix, which throws out what is diffurbing the ftomach in its vafcular ftructure, and might prove hurtful to the conflictution.

CCCX.

The *dyspeptic* flate of the flomach (or in other words, want of appetite and fickness) has been fupposed to depend upon a loss of tone in its *muscular* fibres. If it *be* fo, it may be looked upon as an argument against the fupposed fympathy between *these* fibres and the vessels on the skin, unless we are not to suppose the sympathy mutual. If it

15

is mutual, then, in dyfpeptic flates of the flomach, the veffels of the fkin fhould be morbidly affected. But this is not the cafe, and probably the reafon why it is not, is, that this is a chronic, and therefore partial affection, and otherwife brought about.

CCCXI.

The healthful ftate of the ftomach, as I have before obferved, feems to me entirely to depend upon a due tone and action in the exhalent and fecretory veffels of the villous membrane; and I am inclined to confider the mufcular ftructure of the ftomach itfelf, as wholly intended to keep *its contents* in proper contact every where with the villous membrane, that *they* may be advantageoufly acted upon by the digeftive *menftruum*; and *ingefta*, whether fedative or ftimulant, appear to me to act by producing an alteration in the action of the veffels,

CCCXII.

Thus, for inftance, common falt is a *flimulus* much in use, and the strongest that M 4 can

can be applied to the mufcular fibres, as experiments have fatisfactorily fhewn; fothat, if it be allowed that veffels have irritability, (and from the paffage taken from the first lines of Dr. Cullen's Physiology it appears that they have a great deal) we must certainly expect that they will be quickly and confiderably affected.

have before obf. HIX, OOAns to me entirely

The healthful flate of the flohach, as I

The paffage alluded to is the following; " It is probable that the mufcular fibres of " the arteries become more irritable as the " arteries are more diftant from the heart." Inftitutions of Medicine. 1777. Parag. CLIX. page 131--2.

CCCXIV. h vd hong boffs

mombrane, that Very may be advantageout

Notwithstanding this Dr. Cullen supposes the operation of all stimuli (for instance, of the vitriolic acid) to take place on the mufcular fibres of the stomach; but when I consider how well these are defended from *fuch action* by the thickness of the villous membrane, I cannot readily admit the explanation, and must suppose that all stimuli for

fo applied will first, at least, affect the extreme veffels on the furface of the stomach, and produce a change of action in them.

CCCXV.

Befides, the rules given for the administration of medicines in general are contrary to fuch a fuppofed operation. Medicines have been fupposed to act best on a fafting flomach, and accordingly in general have been advifed to be given during fuch a ftate of it. Why? (my answer would be) Becaufe in that ftate they are both lefs diluted, and act more immediately upon the villous membrane. Such exhibition then must be directly contrary to their fuppofed operation on the muscular fibres, because, the ftomach being empty, thefe are very confiderably defended by the puckering together of the villous membrane, from the contracted state of the stomach.

CCCXVI.

From this I am led to observe, that if it is wished that a medicine should operate on the

the mulcular fibres (if fuch action can be of any fervice) it fhould be given in the diftended flate of the flomach, when the villous membrane is greatly expanded; but even in that flate of the flomach, if it can affect the mulcular fibres, it must also materially affect the villous valcular membrane.

CCCXVII.

I shall endeavour to explain my opinion refpecting the operation of medicines in general, by fome observations on the Peruvian bark, &c. The Peruvian bark has not generally been supposed, I believe, to enter the circulating fystem, but to produce its effects by acting on the primæ viæ, by which we can account for its fudden effects in intermittents. Such operation is confiftent with the opinion which has been advanced of the fympathy of the ftomach with all parts of the body. But does it act on the parietal muscular fibres of the primæ viæ? In my opinion it does not; becaufe to do that it must have entered the abforbent fyftem to be in contact with them, as the villous

willous internal membrane is the part to which it is immediately applied. Here it might be alledged, that it affects them by fympathy, from acting on the fentient extremities of the nerves.. I do not doubt but that thefe nerves may be affected, ftill I cannot fee how the extreme veffels can efcape being acted upon, they having been proved, from experiment, to be very irritable.

CCCXVIII.

All medicines feem to me neceffarily to produce their action, firft on the villous membrane, after the manner in which cathartics operate ; thefe are *flimuli*, which from their irritation on the exhalent veffels, and excretory ducts of fmall glands, caufe an increafed effusion from the former, and fecretion in the latter, from which two circumftances arife the liquid ftools. If fuch then be the operation of cathartics, it is to be prefumed, that medicines in general act upon thefe and fimilar parts, through the whole

whole of the first passages, each producing its effects, according to its nature, on this or that particular part.

CCCXIX.

Thus, the bark, for inftance, acts as a tonic in general (from the connection and confent between the extreme veffels of the flomach and all parts of the body); *ipecacoanba* as a vomit; *jalap* as a cathartic, &c., And I greatly fufpect that medicines (bark, for inftance, when it fails of curing the ague) difappoint us in not producing their effects from the total want, or the diminifhed influence, of this principle in the conftitution. Emetics are capable, I imagine, of calling forth this principle, as after the exhibition of them, we often find medicines fucceed, which before had failed.

on in the latter, from which two cite

tes and the liquid finals. If fich

the operation of cathantics, it is to

med, that medicines in ceneral act

CHAP.

CHAP. IV.

How the Author's doctrine is illustrated by the fame arguments, which Dr. Cullen has made use of to illustrate his own.

SECT. I.

Remarks on the fact related by Dr. Sydenham concerning the plague.

CCCXX.

T behoves every man, who means to fubfitute a new doctrine to an old one, to omit no argument in favour of that which he has newly advanced.

I shall therefore prefume to observe, that the vascular connection and confent I wish to establish, to wit, that when there is an atony and constriction of the extreme cutaneous vesfels, there exists also in the stomach a similar affection of similar parts, may be further

ther illustrated by the very fame arguments. which Dr. Cullen has made use of with the view of illustrating his own doctrine, which is briefly the following. He has fuppofed, that the atony of the cuticular extreme veffels is the caufe of the febrile affections of the flomach, by inducing an atonic flate of the muscular fibres of the ftomach, from fympathy of parts. Now I do not fuppofe that fympathy is fo immediately concerned in this matter, but that the affection of external and internal extreme veffels chiefly depends (as I may more than once have already observed) on their balance with each other, and their mutual connection with the circulating fystem.

CCCXXI.

The arguments referred to, which support this connection and mutual affection of veffels, are the following, which are to be found in the XLIII paragraph of Dr. Cullen's *Firft Lines* on his practice.

I. The fact related by Dr. Sydenham concerning the plague.

II. The

- II. The ceffation of vomiting at the coming on of the hot stage of fever, and very certainly on any sweat appearing.
- III. The effects of vomits in bringing on the hot ftage.
- IV. The effects of cold water taken into the ftomach.
- V. The effects of cold applied externally to the body,

I think it quite unneceffary to quote the whole paragraph from the First Lines, as fuch a valuable work is no doubt in every one's possefition. I shall now proceed to confider each of the above arguments at large, in the order they have been mentioned.

Fact concerning the Plague.

CCCXXII.

At the commencement of the pestilential disease known by the name of the plague, there

there comes on a degree of vomiting fo violent, as to hinder any medicines remaining on the ftomach, which prevents altogether the relief which the phyfician in fuch cafe might otherwife afford. This excefs of vomiting appears to me eafily accounted for on the reafoning I have all along made ufe of. The plague has been confidered by the moft eminent authors and teachers of medicine, as no other than a *typhus violentiffima*, (if you except the occurrence of the *bubo*,) the debility of the fyftem being fo confiderable, as foon to bring the miferable fufferers to a fatal diffolution. In every *typhus* the debility creates the danger.

CCCXXIII.

In this place I think it proper to obferve, that the debility, which appears to me to prove fatal in fuch cafes, properly belongs to the various functions of the fyftem, on which life more materially depends; nor am I inclined to confider the concomitant debility of the mufcular fyftem, as in the leaft dangerous independent of the various affections

affections of other more important parts. Indeed I am more inclined to confider the muscular system soon after the approach of fever, as rather acquiring an increased tone, and energy of exertion, in order to obviate the dangerous confequence of a debility in the vital functions, which is rendered fo highly probable from the common occurrence of vomiting at that period.

CCCXXIV.

every view of the fubject, the exertions of

After this view of the apparent debility of the muscular system, it appears to me difficult to explain that occurrence of affections in the ftomach, and on the fkin at the fame time, from any fuppofed fympathy between the muscular fibres of the former, and veffels in the latter. For we might reafonably have expected, that, if there had really exifted a fympathy between these parts, such violent exertions of the ftomach and fystem in general, at the time of vomiting, would have been accompanied with a correspondent increase of action in the vessels on the furface traigent , monopen Nemma out to olgion of

194 ON FEBRILE SYMPATHY, &c. of the body, which form the capillary fyftem.

CCCXXV.

This fympathy therefore, from the action of vomiting, must neceffarily have ended in removing the atony and confiriction of veffels on the cutaneous furface, the fuppofed principal and fupporting caufe of fever. In every view of the fubject, the exertions of the fystem during vomiting feem to me to argue an increased tone or tonic power of the stomach, (for it is difficult to fuppofe that an atony could exist in the muscular fibres of the stomach at that time,) which, if Dr. Cullen's alledged fympathy had existed, must have produced a correspondent tone and exertion in the extreme cutaneous vessels.

CCCXXVI.

Since vomiting is not an exertion of the ftomach alone, but of the fystem at large, the more I reflect on its phænomena, the more reason I have to believe, that it is a principle of the animal œconomy, implanted in

in it by the *fupreme Being*, to reftore the action of the ftomach, when impeded by the febrile ftate of the fyftem(CCLXXX.). Can its violence therefore at the commencement of the plague be accounted for on this principle ?

forface of th. IIVXXXDD Cthis purpole

with the view of relaxing the fpatin on the

I shall take it for granted, that the greater the debility in the function of the system, the greater are the exertions of nature to reflore them to their healthful state; therefore, as the debility in the plague is greater, *exteris paribus*, than in any other febrile affection, the vomiting is of confequence more violent, and yet not always sufficient to answer the intentions of the vis medicatrix: and I cannot help considering the vomiting as hurtful in the end, when it is not in due time followed by a relaxation of vessels, because it may wastefully expend the vigor of the system, and increase the debility of the functions (CV. CVI. CVII.).

IIIVXXOOO functor of the feeretory

ory where, and remove the inter-

CCCXXVIII.

Dr. Sydenham found the vomiting fo confiderable in the plague as not to admit any medicine to ftay upon the ftomach ; he had thereforere courfe to external applications, with the view of relaxing the fpafm on the furface of the body, and for this purpofe made use of the hot bath : as foon as the fweat broke out the vomiting abated, and foon entirely ceased. This phænomenon has been fuppofed by Dr. Cullen to be brought about by reftoring the tone of the mufcular fibres of the ftomach, and removing the fpasm affecting them; but I think, notwithstanding this great authority, that its operation may be explained on my own hypothefis of a supposed mutual affection of veslels.

CCCXXIX.

The hot bath proves a general *flimulus* to the fystem, and of course increases the action of the heart and arteries: this must finally restore the action of the extreme vesfels every where, and remove the interruption in the function of the fecretory organs;

organs; but the water will further tend greatly to relax the confriction of the cutaneous veffels. While then the fweat flows on the furface of the body, the exhalent and fecretory organs in the ftomach, as well as in other parts of the body, recover their action; and this reftoration having been the chief object of vomiting, as an effort of the vis medicatrix, it then ceafes, there being no longer any need of its continuance.

CCCXXX.

That this is the cafe in the plague is well known. The fact has been taken notice of by feveral other writers on the fubject, who have all agreed in the obfervation, that on the breaking out of a fweat the vomiting has ceafed. An obftinacy in vomiting might no doubt take place at the violent attack of any other fever; and I fhould not hefitate to have recourfe to the ufe of the hot bath, at the commencement of moft fevers, if it wasattainable, as I think it more efficacious than a vomit, from its being fo immedia-N 3 ately

ately an universal stimulus to the circulation, on the principle of sympathy between the extreme vessels and the HEART (CXXIV.).

CCCXXXI.

I cannot quit this part of the fubject, without offering a few reflections on the variolous attack. Great nausea and vomiting will fometimes, nay do often, occur during the eruptive fever of the fmall pox, especially if there be any crudities and corrupted humours in the stomach and primæ viæ. In this diforder the greater or leffer quantity of the eruption is probably owing to the ftate of the extreme veffels, at the time of the eruptive fever; and the previous state of the stomach has, in general, been observed to bear a relative proportion to the quantity of the eruption that afterwards does appear. no doubt take piace at the violent

CCCXXXII. IIXXXDD

If then, from the univerfality and permanency of the confiriction of the extreme veffels, the greater quantity of the eruption takes place, and has already been preceded by

by the above-mentioned affections of the ftomach, from there having probably been a fimilar eruption forming on the villous membrane from a fimilar caufe, may we not reafonably expect, that by obviating the conftriction of veffels, we fhall have lefs fever, and afterwards fewer pullules.

CCCXXXIII.

In this way, I should imagine, nauseating dofes may operate advantageoufly at the commencement, and during the period, of the eruptive fever : and vomiting, from whatever caufe produced, whether by nature or art, may be confidered as a means ufed to throw off, by the perspiratory organs, and extreme veffels in general, the variolous matter which forms the puftules, by timely overcoming, if not preventing, the confiriction of veffels, that would retain it in the fkin, and capillary fystem.

The explanation of this will appear evident fitter where I have sheet or faid. It a sign of the pervous fillers insuld by the NA CHAP.

1.14

CHAP. IV. SECT. II.

On the ceffation of vomiting at the coming on of the hot stage of fevers, and very certainly on any sweat appearing.

CCCXXXIV.

I fhall next confider the ceffation of vomiting, which takes place commonly at the coming on of the hot ftage of fever, but very certainly on any fweat breaking out. Dr. Cullen, when fpeaking of this phænomenon, makes the following obfervation: " It is indeed probable, that the " vomiting in the cold ftage of fevers, is " one of the means employed by nature, " for reftoring the determination to the " furface of the body." Firft Lines, paragr. XLIII.

CCCXXXV.

The explanation of this will appear evident from what I have already faid. If a debility of the nervous fystem should lay the foundation of a weaker action in the heart and

and arterious fystem, the momentum of the blood, which depends on the action of the heart chiefly for its velocity, must also be diministred, which diminution again will be chiefly felt in the fystem of the extreme vessels, or *capillary* system; hence, from wanting the distensile power, by which the tone of these small vessels is supported, they will naturally contract, or be constricted.

CCCXXXVI.

I have no doubt but that, in long continued fevers, where the confriction has been permanent, the fides of thefe fmall veffels may have coalefced together, and become imperviable to the momentum of the ferofity ever after, fo that if the fever proves not fatal, there must be an increased determination of ferofity to fome other fecerning organ, that life may be fupported; for I imagine it cannot long continue (or, if it does, it must be in an unhealthful flate) without more or lefs of the infenfible perfpiration continuing, or (as an equivalent to the

the lofs of it) without an increased secretion in some other organ, or effusion internally from other serves arteries.

CCCXXXVII.

ad How more motion will be

To fuch a permanent confiriction may be owing the diabetes, obfinate diarrheas, anafarea, incyfied dropfies, &c. which have been obferved fometimes to come on after long continued fever; and fuch permanent confiriction of the extreme veffels on the furface of the body, particularly on the extremities, is more likely to continue than in the internal parts, as they are more diffant from the heart, and are at the fame time exposed to many external caufes that may fupport the confiriction, and counteract the good effects of the increased momentum of the blood, from the increased action of the heart and arteries (CCCXXXV.).

CCCXXXVIII.

It is on this principle likewife, that I would account for the partial fweats that fometimes occur

occur on the head and breaft, while a con-Ariction exifts on other parts of the furface. I likewife think it probable, that the pluritic, arthritic, and other affections, which fometimes follow intermittent fevers, may be owing to the continuance of the atony and fpasm in the internal extreme veffels of the parts fo affected. Upon the whole I think it fufficiently probable, that a mutual exertion and balance in the extreme veffels is effentially neceffary to perfect health, and that many topical difeases of the fystem may be owing to a total abolition of their function in other parts, though not made known to the phyfician by any apparent change in those parts, where the function of the extreme veffels may have been loft.

CCCXXXIX.

I shall conclude the confideration of the prefent argument with observing, that as the action of vomiting, at the accession of fever, keeps a corresponding proportion with the state of the vessels on the outer surface of the

of the body, i. e. that as its commencement, continuance, violence, and ceffation, are regulated by the flate of action in the cutaneous extreme veffels, it ftrongly argues a dependence on afimilar affection of fimilar veffels at the ftomach. (CCLXIV.) The fecretory minute parts in the ftomach are no doubt fimilarly affected with the other fecerning organs, from the ceffation of the febrile affections of the ftomach, when the other fecretions are returning to their natural state, whether immediately affected by the action of vomiting, or the augmented force of the circulation. The appetite is neceffarily reftored in due time, becaufe the digeftive organs will have then recovered their wonted vigor and free action (CCLXV.).

I fhall conclude the confidention of the preferst argument with obferving, that as the action of vomiting, as the accellent of fever, .9 A H D redponding perportion with the

CCCXXXIX

CHAP. IV. SECT. III.

On the effects of vomits in bringing on the hot stage of fever.

CCCXL.

I am now to confider the effects of vomits in bringing on the hot ftage. Agreeable to the opinion I have all along advanced refpecting vomiting, I am inclined to explain the operation of emetics in the following manner; I do fuppofe, that, from fome specific noxious quality which they poffefs, they in fome way difturb the action of the extreme veffels, and probably alfo the fecreting organs of the fuccus gastricus, (at the fame time may irritate the fentient extremities of the nerves) from which operation the fenfibility of the ftomach, feeling an enemy harraffing its territories, or interrupting its function, is rouzed, and then calls to her aid the fuperior exertions of vomiting, as a law of the vis medicatrix, which happily terminate in the expulsion of the offending matter.

matter. This may be confidered as a primary effect (CCCVII. CCCVIII. CCCIX.):

CCCXLI.

As a fecondary one I would fay, that the fhock, which the whole fystem receives by the action of vomiting, must be attended with a temporary increase of the circulation; that therefore, by giving an emetic before the cold ftage of an intermittent comes on, we often altogether obviate it, and perhaps the fucceeding ftages, by having rouzed the nervous fystem, and put into brisk action the vital organs, fo as not to allow that debility in the circulation to take place, which has been fuppofed to lay the foundation of the cold stage, and all the subsequent fymptoms. If the emetic be exhibited later, namely, when the cold ftage has already commenced, we ftill may fhorten the cold ftage, by haftening the accession of the hot.

CCCXLII.

It is fomewhat fingular, that on the exhibition of a vomit, and as preparatorily to its acting

acting as fuch, the fystem in general should undergo a change, in feveral particulars, very much refembling the accellion of fever. These particulars, as they have been generally unattended to, I shall describe; they are the following: Soon after taking an emetic, or rather, I should fay, fome time before it operates, an evident languor and inactivity come on, accompanied with palenels of the face, and a weaker, fmaller, and rather quicker pulfe than natural; the fick man will totter, and be giddy, as from inebriation, on walking about the room : a moisture may be often observed on the Skin, particularly on the face, and an increase in the falivary fecretions, indicating the relaxation of excretories : I myfelf lately experienced, on taking an emetic, fome of the above, together with aching pains in my hands, arms, and legs, and a fenfe of coldness, particularly on walking about the room. Soon after these fymptoms have come on, the nausea increases, and shortly ends in vomiting, which reftores the fystem to its priftine ftate.

CCCXLIH.

CCCXLIII.

The temporary debility of the voluntary motions, and perhaps the weakness of the circulation, both feem owing to the inattention of nature at that particular time to those functions, the whole of her attention being taken up in obviating the retention of the emetic matter, and effectuating its expulsion by the exertion of vomiting. One would be almost led to fuspect, on attentively confidering the operation of an emetic, that the remote cause of fever may often act principally on the flomach, and bring on the fymptoms of the cold stage in the fame manner, that the emetic matter produces the general languor, and the debility of the circulation, previous to vomiting(CCXLVII. CCLII.) Probably the caufe of fever is more difficultly got rid of, and hence arifes the continuance of it, even though nature has called in the affiftance of vomiting.

CCCXLIV.

As the operation of emetics feems occafionally to vary, as they are of a *flimulant*

or

or *fedative* nature, I shall make fome obfervations on the peculiarities, that may be supposed to belong to each.

CCCXLV.

sure, and fuited to remove

If they are of the first kind, their effects may depend upon their irritating the fentient extremities of the nerves, as well as the extreme veffels, and other minute parts in the villous membrane, which, from caufing an uneafy fenfation, gives rife to the convulfive action of the ftomach; the irritating matter is then expelled, and the organ reftored to its natural state. The operation of fuch emetics is commonly attended with an increafe in the fecretions of, and effusions into, the flomach, as is evident from the great quantity difcharged by vomiting : there will also be an evident glow of heat on the ikin, and an increase of perspiration, which feems a ftrong proof of the particular connection between the external and internal extreme veffels; notwithstanding which the stimulant emetics, such for instance as the mustard seed powdered, have even been fuppofed

pofed to produce an atony in the mufcular fibres of the flomach; but this certainly could not happen, as they are of a tonic nature, and fuited to remove atony.

CCCXLVI.

With regard to the fedative emetics, I think it probable, that when applied to the ftomach, they check the nervous influence in the fentient extremities of the nerves: this diminishes the effusions from the extreme veffels, and perhaps the fecretion of the fuccus gastricus on account of the ceffation in the ofcillatory motion of the capillary veffels. This impeded action of the villous membrane at length produces nausea and rouzes the fystem to exert its timely effort to the removal of the affection. Such emetics are attended with a paleness of the Skin, languor, and inertia liquidi nervofi, all of which denote the fedative effects of the emetic matter on the nervous, and fanguiferous fystems. Notwithstanding this very apparent diminution of irritability, the muscular syftem can fully exert itself in the production of

of vomiting, in obedience to the vis medicatrix naturæ, and thus reftore the vigour of the whole.

CCCXLVII.

In order further to illustrate the effects of an emetic, I shall beg leave to make use of a *fimile* borrowed from the animal æconomy, and which is one, among many other instances that might be mentioned, of the wise operations of nature, who, while she establisted laws for the prefervation of the whole system, under the guidance of the vis infita fympathiæ, did not neglect to screen from injury particular parts of it.

CCCXLVIII.

It was neceffary to diffinct vision, that the anterior furface of the eye, particularly the cornea, fhould be kept bright, and every, the most minute, extraneous body prevented from lodging on it. Nature has accordingly made provision for this purpose, by an appendage to the eye, the glandula lachrymalis, the fource of the tears. It is fituated within O_2 the

the orbit above the outer canthos, and its ducts open near the angle formed by the two eye-lids near to their inner edge. The tears are carried acrofs the eye to the inner canthos, and pafs out to the faccus lachrymalis, to be from thence conveyed to the nofe. Thus they wash every, the most minute, particle from before the fight. Let us fee then the effect of a mote falling into the eye.

CCCXLIX.

The tunica conjunctiva, a reflected membrane covering the anterior furface of the eye-ball, and the internal furface of the eyelids, is highly fenfible, particularly that part of it which lies on the eye-lids, and fympathizes with the lachrymal gland. The mote falling on the eye is generally drawn within the upper eye-lid, and gives confiderable irritation. The confequence of this is, that a very fudden increase in the fecretion of the tears takes place, and will, if the mote is not too large, effectually wash it from the eye, or hurry it to the faccus lachrymalis. If the offending matter should perchance be

an

an acrid fluid, as vinegar, the flow of tears, by diluting it, deftroys its power of further irritating the eye. Such has been the wife providence of nature, in preferving that most useful and important of all our organs.

CCCL.

The attended for the

Let us make the application to our fubject. The villous membrane of the ftomach is irritated in its way by the emetic matter, as the tunica conjunctiva was by the mote, and if it were not expelled might probably fuffer fome injury from it. Nature, attentive to this, raifes a naufea, relaxes the excretories of the ftomach, increafes thereby the quantity of the gaftric fluids, weakens thus the power of the medicine, and then, to prevent further bad effects, expells nearly the whole by vomiting.

the read of all to these to stimp the

has harrough a between the external and

03

ale courran er alte the the derauge sin

and an entropy the second applied

CHAP.

CHAP. IV. SECT. IV.

On the effects of cold water taken into the stomach.

CCCLI.

The next proofs of my dectrine relate to the effects of cold on the human body; and first, I am to consider the effects of cold water taken into the stomach.

The circumftances already mentioned feem me to be fufficient evidence of a connection between the ftomach and the fkin : ftill the prefent argument, and the one next to be confidered, are further proofs of fuch a connection.

CCCLII.

After all that has been faid, I am ready to allow it a very difficult tafk to arrange the facts, and give them their proper application, fo as clearly to fhew to others, why fuch a connection between the external and internal parts fhould exift, and ftill a more difficult

difficult one to fay, in what fystem of minute invisible parts it particularly confist.

It has been, notwithstanding, from frequent observation allowed to take place, and I will venture to offer it as my opinion, that it is to be referred to the capillary system of veffels, (CXCIX.) and may, as I fuspect, be owing to a law of nature, a VIS INSITA SYMPATHIÆ, that when one extremity of the circulating fyftem (confidering the external and internal capillary veffels as the extremes of the faid fystem) is morbidly affected, the other should confent, that the whole system may be rouzed, and the vis medicatrix naturæ exert her feafonable endeavours to remove the affection. In making this reflection I have given great latitude to the influence of fympathy and confent.

CCCLIII.

Though cold water taken into the ftomach may produce hurtful confequences in fome ftates of the fyftem, it would be foreign to our prefent purfuit to confider them O 4 now.

now. What I mean at prefent, is to offer a few remarks on its falutary effects, as thefe will most strikingly prove the connection in view. (CCCLII.)

CCCLIV.

It has been the practice with fome phyficians to give a draught of cold water in fever, in order to procure a fweat; and the practice has been attended with fuccefs. It had been observed by them, that a cold draught taken in warm weather foon after excited a glow over the body: they took advantage of this phænomenon, and made use of it in the cure of fever. I cannot conceive the effects produced in fuch a cafe to depend upon any other operation, than that of the cold applied to the veffels and nerves of the ftomach, and, from a change produced in the tone and action of thefe, affecting, by balance of sympathy and consent, the extreme veffels on the furface of the body.

CCCLV.

As cold moderately applied, when the fyftem is in full vigor, acts by a tonic and ftimulant

ftimulant power, it can be readily fuppofed, in the prefent cafe, that it produces an increafed action in the vafcular ftructure of the ftomach, and (from an *eftablifhed confent*,) that fimilar effects take place on the furface of the fkin. It must be allowed a weighty argument, in fupport of the *favorite doctrine* of fympathy and confent, as I do not fee that we can comprehend, how pure water alone can act by entering the circulation, and being determined to the fkin : the quantity of the water ufed, and the fuddennefs of its effects, are far from being agreeable to fuch a fuppofed modus operandi.

CHAP. IV. SECT. V.

On the effects of cold applied externally to the body.

CCCLVI.

Laftly, I have to confider the effects of cold applied externally to the body. Thefe likewife

likewife very clearly fhew the dependence of the condition of the flomach on that of the fkin, from the changes produced in it by cold applied to the fkin. Frequent inflances occur of the invigorating effects of cold on the human body, when it does not ftop the exhalation of the perfpirable matter: it proves then a very potent mean of increasing the appetite. We have a remarkable inftance of this in its effect on the appetite of those perfons, who, during froft, exercise themselves in skaiting : they will at that time perfpire most freely, and have a most voracious appetite; but both these fymptoms, it has been observed, abate, or altogether go off, when the ftimulating effects of the cold on their bodies have ceafed.

CCCLVII.

Again, when fedative and aftringent effects, and a retention of the perfpirable matter,

matter, proceed from cold applied externally, it commonly diminishes the appetite; which is a further proof of the balance and connection of the parts concerned.

5 E C 1.

CHAP.

matter, proceed them cold applied externalit, is community diminifies the appetices valid, is a further proof of the balance and

CHAP. V.

Some opinions are delivered in a concife point of view, as a fummary to the whole.

SECT. I.

Some general conclusions respecting the fe brile affections of the stomach, by way of COROLLARIES.

CCCLVIII.

HAVE finished, in the preceding chapter, with my arguments in illustration of the doctrine I have ventured to take up. I am apprehensive of having been too tedious and profix in some parts, most probably not altogether clear in others, and of having unavoidably fallen, from the nature of the work, into the too frequent repetition of the fame ideas. I hope however that my reasoning has not wanted facts to support it, and that

that the difficulty of the fubject will be an apology for its deficiencies.

CCCLIX,

I fhall now lay before my reader fome general conclusions, by way of corollaries, and finish with a summary of the pathology of fever, and the *promifed observations* on the terms sympathy and consent. (CCXCVII.)

COROLLARIES.

in confequence of the moreste of Mony in th

CCCLX.)

COR. I. If a debility of the nervous fyftem, from whatever caufe induced, ceafes a weakened action of the heart and arteries, this muft neceffarily and principally be felt in the terminations of the arterious fyftem, namely, the fyftems of capillary arteries, and fecretory veffels; I therefore confider the ANOREXIA, which fo generally attends the beginning of fever, as depending on an ATONY of the extreme veffels, and a diminifhed action of the villous glands of the ftomach.

141010

CCCLXI.

CCCLXI.

COR. II. If it can be reafonably fuppofed, that the continuance and increase of the general debility are attended with a proportionate atony of the extreme veffels, and that the spass of the extreme veffels is more or lefs according to the extent of the atony in them, then I conclude, that the NAUSEA depends upon the CONSTRICTION formed, in confequence of the increase of atony in the veffels of the stomach.

CCCLXII.

COR. III. If the continuance and increafe of the general confriction of veffels, together with the total ceffation of the function of the flomach, can be juftly confidered as a fufficient caufe to rouze the vis medicatrix fympathiæ to her falutary exertions, then I fhould confider the VOMITING as the RE-ACTION of the flomach and mufcular fyftem, completely fuited, in moft cafes, to reftore to its due action, that important organ of digeftion.

CCCLXIII.

CCCLXIII.

COR. IV. and laft. If debility, fpaim, and re-action can fatisfactorily explain the morbid changes which occur in the circulating fystem, and fecretions in general, atony, constriction, and re-action, as above, will equally explain the morbid changes or affections of the ftomach, to wit, the anorexia, that a cor nausea, and vomitio. an obfirection teles plats

CHAP.V. SECT. II.

m the feer shops in

The author attempts to give a new fummary of the pathology of fever.

CCCLXIV.

I shall now attempt to give briefly at one view the doctrine and extent of the vis medicatrix in fever, as a fummary of that part of the treatife in particular, which relates to the fubject of fever, agreeable to the pathology I have all along adopted. I fay then, that the remote causes of fever act with a fedative effect, and induce a debility of the nervous [w]

nervous system, whereby the vital function of the heart and arteries is confiderably and particularly affected :---- that this debility in the circulation will most readily be felt at the extreme veffels, which terminate at every part of the body, but particularly in those at the skin and primæ viæ, from establifhed connection and mutual balance; that a confiriction will then be formed, and an obstruction take place in the fecretions in general, in the exhalation of the perfpirable matter, and effusion of the gastric and inteftinal liquors ;- that this confriction not only proves a general ftimulus to the circulation, in confequence of a fympathy between the heart and veffels, but a particular one to the ftomach itfelf, as the first of the fecreting organs; -and that this ftimulus, at the fame time that it increases the action of the heart and arteries, excites the ftomach to vomiting, both which effects are to be confidered as the efforts of the vis medicatrix naturæ, for the falutary purpose of restoring the condition of the extreme veffels, and fecretory organs in general, to their natural and healthful

ful state, which being fully accomplished the *pyrexia* and vomiting abate. These operations are supposed to be brought about, upon the principle of a VIS INSITA SYMPA-THIÆ (CCCLII.).

The reader is defired to compare the above fummary of fever with the one drawn up by Dr. Cullen, agreeable to his own fystem. It has been mentioned (parag. CCLVI.) in this treatife, as a quotation from his First Lines.

CCCLXV.

I would have the above fummary confidered as the regular train of fymptoms in the progrefs of an intermittent paroxyfm; when there are wanting the complete folution, and perfect intermiffion, this conftitutes the continued and remittent fevers, which I think may be explained by faying, that while the quicknefs of the pulfe, and lofs of appetite continue, they argue either an imperfect folution of the fpafm, or continuance of the atony in the extreme veffels. There P only

only now remains to offer a few remarks on the terms fympathy and confent.

CHAP. V. SECT. III.

Some promised observations on the terms Sympathy and Consent.

CCLXVI.

I observed in a former part of the work, that thefe terms were liable to fome objections and reftrictions, and then proposed only to use them as the old language, that I might not, by the innovation of new terms in their flead, confuse my reader. (CCXCVII.) I still however find it impoffible entirely to fet them afide, as they often are most applicable to the fubject. Thefe terms have however been found ufeful to explain, or account for, occult caufes, and are fometimes the last refource, the ne plus ultra, of a deficient comprehension. I do not think that they are firstly or juftly applicable to that observance in point of time, which

which takes place between the affections of the flomach and fkin in fever, as fuch fympathy could only properly arife from nervous connection.

CCCLXVII.

Agreeable to the explanation I have offered, the affection of the extreme veffels on the fkin and primæ viæ appear to flow neceffarily from the laws of the circulating fyftem alone, the vigor of which, as depending on the heart for its fountain and fupport, being weakened, will naturally produce the confequences arifing from fuch debility in the extreme veffels every where, but more particularly in those of the fkin and primæ viæ for feveral reasons, and above all, from their being exposed to the action of many remote causes, that fupport the debility in them, but cannot affect the other parts.

CCCLXVIII.

In my opinion, therefore, the flate of the extreme veffels in fever, both externally and P 2 internally,

internally, which is a neceffary confequence of the debility in the circulation, cannot be confidered as ftrictly depending either on the nervous system, or any particular sympathy connected with it. The terms were properly used on the supposition delivered by Dr. Cullen, as no immediate or neceffary connection could be fuppofed to take place between the extreme veffels of the ikin, and the muscular fibres of the ftomach, fo that he was obliged to have recourfe to a fuppofed fympathy. But on the fuppolition of my doctrine of extreme vessels being the true one, I would propose, that the terms balance and connection should alone be made use of, when the pathology of fever is the fubject fpoken of, as I think they are fufficient, becaufe fuch veffels are effentially connected with the state of the circulating system, and accordingly are affected, when that fystem is debilitated (CCCXXXV.).

CCCLXIX.

This mutual balance and connection between the cutaneous extreme veffels, and the analogous

analogous veffels of the primæ viæ, which. are fo apparently affected in fever, appear. alfo to take place as neceffary to the healthful state of the fystem. This may be obferved from the peculiar effect of exercise, or any other active caufe, on the state of these vessels, when such a cause increases the general circulation. This balance does not manifest itself in the terminations of the arterious fystem at the fecretory organs; becaufe, whatever increases the action of the heart and arteries, increases, as a necesfary confequence, the quantity of the perfpirable matter and fweat, yet at the fame time does not feem manifeftly to augment any of the other fecretions, as if the terminations of arteries in the fecretory organs were not fimilarly, at least not proportionably, affected.

CCCLXX.

The fecretions in general are often varioufly and confiderably affected by the influence of fympathy. The affections of the mind will often interrupt glandular fecre- P_3 tions.

230 ON FEBRILE SYMPATHY, &c.

tions. Thus we often fee the fecretion of milk fuddenly ftopped by diftrefs of mind, and fometimes it will not return again. A balance is kept up between the fecretions. This is most remarkable between the fkin and kidneys. It also in a great measure takes place between the *breasts and uterus*. But a plethoric state alone of the circulating fystem most remarkably increases the fecretions of perspiration, urine, and *sometimes* milk.

CCCLXXI.

The other fecretions feem neither affected by the quality, or the quantity, of the circulating fluids, while both produce changes in the fecretions I have juft now mentioned; and it appears, that the others are only affected by ftimulants, either immediately applied to their organs, or indirectly, through the affection of the fenforium, and the nervous fyftem, agreeably to the laws of the animal œconomy. When the fecretions are thus indirectly affected, the changes are produced through the power of fympathy, and not from any immediate balance or connection with the circulating fyftem (CCCLXX.).

Ç O Na

CONCLUSION.

CCCLXXII.

I have delivered it as my opinion in feveral parts of the treatife, that the *extreme arterious veffels*, forming a part of the capillary fyftem, are most likely to become the channels of *medical fympathy*, through the medium of the fympathizing heart (CXCIX.) I cannot therefore close the fubject, without informing my reader, that fome very eminent men are of a different opinion.

CCCLXXIII,

The fympathy of an animal body has been explained by the unifons of found produced on the ftrings of a mufical inftrument. (LVIII. CCXVIII.) Such fympathy has been referred by Mr. J. Hunter to the fyftem of *lymphatic veffels*. Mr. Cruickfhanks is of opinion, that it takes place by confent of the *nerves*. And Dr. Cullen is an advocate, in *pyrexia*, *arthritis*, and *dyfpepfia*, at leaft, for fympathy between *mufcu-*

lar

232. ON FEBRILE SYMPATHY, &c.

lar fibres. The reader is left to judge, which of the above four opinions is the most probable, or determine whether they are all mutually concerned.

CCCLXXIV.

It may be obferved, that the minute parts of an animal body are fo immediately connected with each other, that a mutual fympathy may, or may not, take place between them, and yet the *fast* cannot be eafily afcertained. Nerves and veffels may fympathize with each other. This very ftrikingly appeared in an experiment made by the indefatigable Mr. Sheldon, whofe laborious refearches in anatomy and phyfiology will, no doubt, be rewarded with the acquifition of riches and honor.

CCCLXXV.

Mr. Sheldon divided the *carotid* artery in a dog, and afterwards tied it together. The union between the divided ends of the artery did not again take place, but the neighbouring part of the nerve *par vagum* fhrunk feveral

ON FEBRILE SYMPATHY, &c. 233

feveral inches in length. This may be looked upon as a fingular inftance of fympathy between an artery and a nerve.

CCCLXXVI.

The absorbent system has yet to engage the attention of speculative men. It must be left to the mutual labors of ingenious anatomists and physiologists (CCCLXXIII.) to find out, by diffection and ferious reflection, of what confequence the absorbent system is, in the pathology of the animal œconomy. Its importance in the phyfiology of an animal body is now very generally admitted. I do not doubt, but that future investigation will discover, that it bears as important a concern in the pathology of fever, as is at prefent given to the extreme arterious veffels : for, it is certain, that many parts of the body fympathize with the abforbent fystem.

CCCLXXVII.

Mr. J. Hunter has mentioned fuch a cafe as the following, in proof of fuch fympathy.

234 ON FEBRILE SYMPATHY, &c.

thy. A woman pricks her finger with a needle, foon after which a red line appears, extending itfelf at laft up the arm to the abforbent gland in the *axilla*, which then enlarges. Vomiting is fhortly brought on in confequence of this, and is to be confidered as a fympathetic affection.

andel labors

of marin firstant 10 21 40

CUT SSLAID ATTACK

activit fault to toorg

. . . .

In fhort, TIME only can, and will, difcover to us, that life has been fupported and preferved by fympathy alone. HE may not ftop there, but may alfo particularly explain to us *hereafter*, how man not only continued to live, and move, and at laft *died*, but how originally he had his BEING. Till then let us turn to the beft use can the knowledge we already poffers.

END of the SECOND PART.

the entreme arterians reliais: for, 10 is certain.

INDEX.

State failure

INDEX.

15

ALBORENIA IS OWING to

Appearte, cold inc

N.B. The Numbers refer to the Paragraphs, and not to the Pages.

A.

BSOLUTE pain, sensitive sympathy causes a division of, - - 151, 152 Absorbent system, the importance of the, in pathology not yet fufficiently attended to, 376, 377 Abforption, external specific applications, when properly applied, operate by, 192, 193, 195 -----, fympathetic glands alone can be relieved by direct, 196, 197 Acceffion of fever irritated on the exhibition of an emetic, 342 Actions, affections, and impreffions, fympathies are produced by, 37, 41, 43 . Affec-

236 I N D E X.

Affections of the mind of one perfon will often operate upon many, ____ 204,205 ------, make the fpirits more powerfull and active, -----206 Aliment, the expulsion of the, from the stomach ----neceffary in fever, 285 Animal body, effects of heat and cold on an, 64 Animal æconomy, vomiting feems to be an innate principle in the, ---- 326 Anorexia is owing to the impeded fecretion of the fuccus gaftricus, ____ 270, 271 depends upon atony of veffels 360, Cor. i. Appetite, cold increases the, when it does not check perspiration, 356 , cold diminishes the, when it retains the perfpirable matter, ----- 357 Appetites, fympathy feeds our, - 30 Atony and spalm, the affections of the ftomach are proportioned to the general, - 306 Attention, fympathies of confciousness are increased by, 14 Attributes, fympathies vary from difference in the fympathizing, _____ 92,93 Author, his original letter to Dr. Cullen, 235 , Dr. Cullen's letter to the, -236 , his opinion on the fecondary ufe of the ftomach as the grand fympathiler, 261 Author,

Author, his opinion confirmed by the records of 262 the ancients, -, his hypothefis rendered most probable from the anatomical ftructure of the parts concerned, 302 -, his fummary of the pathology of fever, 364 The reader is requefted to compare it with Dr: 250

INDEX.

237

Cullen's,

B

Balance, fympathies are called forth by a lofs of, in the moving fibres, -----26, 27, 28 ----- of extreme veffels is reftored by vomiting, 284 ----- of extreme veffels is effentially neceffary to perfect health, 338 Bare-footed, ill effects of going, 164 Birth, the fentient and mental principles not neceffary before, 255 Body, the ftomach fympathizes in hunger with the whole, 68 -----, the whole, fympathizes with the ftomach during digeftion, 69 fome fympathies require ftrength of, 71 Body, Body, the mind fympathizes with the, in hypochondriac affections, 77 -----, fympathy often raifes actions and affections injurious to her habitation, the, ---- 226 -----, internal medicines through fympathy produce their effects in diftant parts of the, 318, 319 ----- the, does not receive moifture from the warm bath, fomentations, &c. 182, 183, 184 Bodily pain produces fympathies of impression, 10, II Brain, diftant fympathies are produced by the organic affection of the, 18 -----, what parts are particularly difposed mutually to fympathize with the, 34

C esterne vend 29 10

Cantharides, &c. how they operate on the bladder by fympathy, ______ 163 Capillary fyftem of veffels, the author refers fympathy to the, ______ 199, 352 Caufe, different fympathies may be produced by the fame, _______ 13 -____, proximate, of fever, 249. Proximate caufes are ufeful delufions Caufes, remote, of fever, 252. Dr. Cullen's doubt. The author's opinion. Child

I N D E X. 239

Child in utero more fimple than a polipus, - 66 may be affected by the conduct of the mother, _____ 221, 222, 223 Children, unborn, how liable to acute and chronic difeafes, _____ 224 Circulating fystem, a fympathy exists between the organs of voluntary motion, and the, 35 _____, the extreme veffels are the most irritable parts of the, ____ 212, 213 Circulation, vomiting gives a ftimulus to the remoteft parts of the, _____ 286, 287 Coalescence of extreme veffels the cause of diabetes, dirrahœa, anafarca, &c. _____ 337 Cold, effects of, on the animal body, - 64 ----- ftage of fever, the phænomena of the, 244 ----- water operates upon the veffels of the ftomach, 354 ----- increases the appetite, when it does not check perspiration, ---- 356 ----- diminishes the appetite, when it retains the perfpiration, - '357 Common inflammation, specific miasmata produce diftantly, _____ 86, 87, 89, 90 fympathetic inflammation, what is the ufe of, _____ 157, 158 Commencement of fever, the fluggifhnefs of the whole

whole body at the, prognofficates great and unufual exertions in the fystem, -291 Commencement of the plague, the vomiting at the, violent, 322 Connections, many of the fympathizing, are by nature fixed and regular, 129, 130, 131, 132 Connection, fympathies between vital parts are founded on a natural, - 133, 134, 141 -----, difeafed fympathies between parts that have no, are but few, 135, 136, 137 , few fympathies from natural, can be reverfed, -138, 139 , natural fympathies eftablish a, between the fame principle in different parts of the body, - 143, 144, 145, 146 fuppofed to take place between the female mind and uterus, - 208, 209 ----- of the febrile affections of the ftomach, with other febrile phænomena, - 265 ____, Dr. Cullen's explanation of the, between the ftomach and fkin, 299 Confcioufnefs, fympathies of, what, 7 -, are increased by attention, 14 , will remain when the caufe no longer operates, 15 _____, originate from corporeal affections, 20

INDEX.

241

Confent, the eyes are by nature prone to move by, 215,216 Conftitution, vomiting obviates many diforders of the, 288 Constriction of veffels at the stomach causes nausea, 361, Cor. ii. Continuity and contiguity of nerves will not account for fympathies, 37 Convullion, and spasm, depend upon sympathetic muscular affection, _____ 26, 27, 28 Convulsions, hysteric and other, from irritation, inconfiftent with natural fympathy, 228, 229 Cuftom and habit, fympathies vary from, - 12 Cutaneous veffels affected by confent from cold water on the ftomach, 354

D

INDE X. 242 Delusions, proximate causes are useful, - 249 Depravity and force of imagination affect most weak minds, 210,211 Diabetes caufed by coalescence of extreme veffels, 337 Diarrhœa caused by ditto, 337 Digeftion, the whole body fympathizes with the ftomach during, 69 ----- not yet rightly underftood 257 Dr. Hunter's opinion of the various hypothefes on, 258 -depends upon a menftruum peculiar to the ftomach, 259 -is ftopt by the febrile condition of the ftomach, 269 Diminution in the momentum of the blood caufes conftriction, 335 Direction, sympathies under the, of the vis conf. et med. naturæ, 21 Difease, the acquired sympathies are commonly produced by, 22 -----, the greater the, the greater the fympathy 105,106,107 probably founded on the want of fympathy, 284 Difeafed fympathies, without organic connection, are few, _____ 135,136,137 Difeated Difeafed bladder, and glans penis, fympathy between, uleful, _____ 153,154,155,156 Difeafes, which the conftitution can cure, not attended with hectic fever, 113,114,115 of the vital and involuntary parts produce the greatest fympathies, - 116, 117 Diforder of the conflitutions obviated by vomiting, 288 Diffimilar fympathies are produced by diftant irritation, _____ 72,73 Disturbance, local sympathy often creates universal, 127,128 Division of the inftitutions of medicine, - 1, 2 ------fympathy is concerned in every, 3,4 of pain caufed by fenfitive fympathy 151, 152 of fever unnecessary to the subject of the treatife, ____ 248 Dr. Cullen's letter to the author, - 236 _____ arguments enumerated, _____ 321 Doctrine of fever, Dr. Cullen's; confistent with the author's hypothefis, _____ 302,303 -----, the author's, fupported by Dr. Cullen's arguments, _____ 320 Doubts, Dr. Cullen's, on the subject of fevers, 251 Q2 Doubts,

INDÈX.

Doubts, Dr. Cullen's on the remote causes, 252 Dyspepsia owing to an affection of vessels, 310

244

E

Effects of heat and cold on an animal body, 64 Emetics, the fudden operation of, fupport the opinion of a balance and connection of veffels, 307 ----- difference between the fedative and ftimulant, - 308, 309, 345, 346 Emetics, effects proper to the ftimulant, - 345 _____fedative, — 346 Emetic, the exhibition of an, attended with febrile phænomena, ----342 ----- the operation of an, may throw light on the remote caufes of fever, -----343 , effects of an, illustrated by a fimile, 347, 348, 349,350 Exertions, unufual, in the fyftem, how prognofticated, 291 ----- of the fyftem are proportioned to its debility, 327 Expulsion of the aliment in fever, why necessary, 285 External specific applications operate by absorption, 192,193,195 Extreme

I N D E X. 245 Extreme veffels, the tone of, reftored by vomiting, 282 284 Extreme veffels the most irritable parts of the circulating fystem, _____ 312, 313 Extreme veffels, how the conftriction of, takes place, 335 ----- from conftriction may coalefce, 336 _____the condition of, regulates vomiting, 339 ------ univerfally affected by a draught of cold water, 354 ----- affection of, produce anorexia, naufea, and vomiting, - 360, 361, 362, 363 ------ condition of, the caufe of continued fevers. 365 ---- remote causes affect the, of the primæ viæ, ----367 ----- fever, why not influenced by fympathy, -----368 fecretory organs, partial, _____ 369 Eyes, the, are by nature prone to move in concert, 215,216

2 Q om debility in the vita

Fear,

F

Fear, observations respecting, as a remote cause of fever, 255 Febrileaffections of ftomach imperfectly underftood, 239 Feeling, the fenfe of, affected by odious objects, 214 ---- may be difturbed by flight irritation, -219 Female mind, what connection between the, and 208, 209 uterus, Fever, the, accompanying phlegmafiæ fympathetic, 110 ____, phænomena of the three ftages of, 244,245, 246 , Dr. Cullen's idea of, and his doubts, 250, 25I , remote caufes of; Dr. Cullen's doubt; author's opinion, 252,253 -----, the principal remote causes of, - 254 -----, Dr, Cullen's fummary of, _____ 256 , nausea in, is a secondary effect, 274, 275 ____, the danger in, arifes from debility in the vital functions, 323 Fever,

INDEX. 247 Fever, acceffion of, imitated by the operation of a vomit. 342 quested to compare it with Dr. Cullen's, 256 Fevers are different. How divided into ftages 242, 243. -----are to be confidered as universal fympathies. _____ 247 ----- how, become continued, -- 365 Fibres, mufcular, how fympathetically affected, 291, 295, 300 Fomentations, how, relieve inflammations, 174, 175,178 ----------do not relieve by adding moifture to the body, - - 182, 183, 184, -----relieve by local and remote fympathy, - - - 185, 186 Fomes, a febrile, might arife from the retention of ingesta, 285 Force of imagination operates most on weak minds, 210, 211 ---- how the, forwards the end proposed, 212

Q4

Gastricus,

IDDEX.

G

Gastricus, fuccus, anorexia caufed by impeded fecretion of, 270, 271 General phænomena of fever, 242, 243 Glans penis: the utility of its affection with a difeafed bladder, — 153, 154, 155, 156 Grand fympathifer, the ftomach to be confidered as the, 261 Greater, the, that the difeafe is, the greater the fym-105, 106, 107 pathy, -----Greatest sympathies are produced by difeases of the 116, 117 vital part, -, univerfal fympathy is, when the injury is 118, 119 remote,

H

Habit, and cuftom, fympathies vary from, 12 Hæmorrhages; how they are ftopt by vomiting, 283 Health is often preferved by vomiting, 32, 80, 292, 293 is fupported by balance of extreme veffels, 33⁸ Hearing

N D E

T

249

Hearing fympathically affected from odious objects, 214

Heart and hands, fympathy between the, - 166 Heart and arteries are rouzed to action by vomiting, 281

64 Heat and cold, effects of, on an animal body, Hectic fever is universal sympathy, - 111, 112 never attends a difease the constitution can cure, 113, 114, 115 Hot bath, how it ftops vomiting at the beginning of fever, 328 ---- in fever, acts by a general ftimulus and relaxing power, 329 330 Hot stage of fever, phænomena of the, -245 Hunger is a fenfation arifing from fympathy 68 Hunter's opinion of the various hypothefes on digeftion, 258 Hypochondriac affections, the mind fympathizes with the body in, 77 Hypothefis of fympathy between invifible veffels, and large mulcular fibres, improbable, 301 Hypothefis, the author's, of affection of veffels, how probable, 302 Hypothesie,

INDEX.

250

Hypothefis, the author's, ftricily confiftent with general doctrines, ______ 3°3 Hyfteric convultions from imitation inconfiftent with natural fympathy, ______ 228,229

Idea thrown out on the pathology of fever, 124 — of fever, Dr. Cullen's, — - 250 Ill effects of going bare-footed, ____ 164 Imitative faculty, sympathy is an, ____ 29, 42 Imperfect solution of spasm constitutes continued fever. 365 Importance of the abforbent fystem in pathology not yet fufficiently attended to, 376, 377 Impression, fympathies of, what, 8 _____produced by bodily pain, 10, 11 Impressions give rife to sympathies, 37, 41, 43 Increase of anorexia causes nausea, 373 Individual fympathy varies from temperament, 121, 122, Inflammation, use of common sympathetic, 157, 158 how relieved by fomentations, 174, 175, 178 Influence of fympathy when not allowed, 368 Influence

DE X. N I 251 Influence of sympathy often affects the fecretions, 370, 371 Inherent fympathies directed by the vis conf. et vis med. nat. 2I ---- constitute a vis vitæ restauratrix. 23 Injuries partial, may produce universal fympathies, 99, 100 . ____, univerfal fympathies greateft from remote, 118, 119 Inftitutions of medicine confift of four divisions, 1,2 Inftinct, how produced by fympathy, ---79 Infufficiency of violent vomiting argues against Dr. Cullen's doctrine, - - 324, 325 Internal medicines relieve by fympathy, 198, 199 Invisible vessels, fympathy between, and muscular fibres, improbable, 301 Involuntary parts, when difeafed, raife great fympathies, _ _ _ 116, 117 Irritable parts, extreme veffels are the moft, 312, 313 Irritation, distant, produces diffimilar sympathies, 72, 73 flight, will raife fympathetic actions, 219 Kidneys,

252

K

Kidneys, Tympathy between the ftomach and, 109 Known, what fympathies are particularly uleful to be, _____ 160, 161, 162

L

Liable, unborn children, to difeafes, — 224 Local fpecific fympathies produced by fpecific miafmata, — 83 — difeafes produce partial and univerfal fympathies, — 96, 97, 101 — fympathy often creates univerfal difturbance, 127, 128 — fomentations, relieve by, 185, 186 Lofs of balance in moving fibres call up fympathies, — 26, 27, 28

M

INDEX.

Medicines, internal, how they relieve diftant dif------, the author's opinion of their operation, 317 first operate on the villous membrane, 318, 319 Mental principle, not necessary before birth, 225 Menftruum peculiar to the ftomach, - 259 Mercurial ointment, when it operates by fympathy, 194, 195 Miasmata, specific, produce local specific sympathies, 83 distantly common inflammation, — — 86, 87, 89, 90 Mind, the, fympathizes with the principle of fimple life, — 74 & feq. -----, paffions of the, determine to particular - 202, 203 parts, -----, affections of the, of one perfon, operate upon 204, 205 many, _____, make the fpirits more powerful, and active, -----206 ----- fome paffions of the, occafionally infective, 207 ---- the ferious passions of the, not infective, 207 ---- connection between the female, and uterus, 208, 209 Moifture not added to the body by fomentations, 182, 183, 184 Momentum,

254 INDEX.	
Momentum of the blood, veffels are conftrict	ed
from diminished, — 3	35
Mother, the conduct of, may affect the child	in
utero, — — 221, 222, 2	-
Moving fibres, loss of balance in, calls forth fy	m-
pathies, 26, 27,	
Muscular fibres, vomiting hitherto supposed ow	-
	289
invisible vessels	
, belonging to parts fimilar, r	301
	305
of the ftomach confidered as parie	
	311
well defen	
from stimuli, — —	314
not acted upor	h by
medicines, — 315,	316
when they i	may
best be acted on, — 315,	1
Muscular sympathetic affections give rife to sp	
and convulsion, — 26, 27	
fystem, the, may claim the principle	-
	5, 6
of the whole, — —	
at the beginning of fever, why	295
affected with debility, -	300
Mu	-

INDE

Mufic, fympathies are compared to unifons of found in, — 58, 218 Mutual exertion of extreme veffels neceffary to health, — — 338

Χ.

255

N.

Natural connection regulates the fympathy of vital - 122, 124, 141 parts, --- fympathies connect the fame principle in different parts of the body, 143, 144, 145, 146 fympathy, when not confiftent with imitatative fympathy, 208, 209 Nature has made regular many of the fympathizing connections, 129, 130, 131, 132 has eftablished a confent between the eyes, 215, 216 -----, the univerfality of fympathy through all, 231, 232, 233, 234

----- depends upon constriction of vessels, 361, Cor. ii.

Nau-

¥

256

Nauseating doses, the effects of, support the opinion of an affection and balance of veffels, 307 Neceffity of a balance of veffels to perfect health, 338 Nerves, continuity and contiguity of, will not account for fympathies, 36 and veffels may fympathize, - 374, 375 Nervous fystem, the, has no superior claim to sympathy, 5,6 Nurses, caution to, to prevent children from fquinting, 216 Nutriment neceffary to excite fome fympathies, 71

0

Object, the acquifition of an, forwarded by imagination, 212 Observation, on vomiting in fever, 364 Observations respecting fear as a remote cause, 255 are frequent, which establish the universal sympathy of the stomach, 292, 293, 294 Occurrence of the febrile affections argues one ge-266 neral caufe, Oils, animal, preferred to the vegetable, 190 Oily applications do not relieve by penetration, 187; 188, 189 Oily

Oily applications operate by fympathy, - 191 Old people more determinate in their fympathies than the young, - 44, 45, 46, 47 Omnes vitæ vires directed by fympathy, 31 One perfon may influence the fpirits of many, 204, 205 Operation of medicines when local and fimilar to the action of the ftimulus, 169, 170, 172, 173 when remote and particular from the nature of the fympathifer, 177, 179, 180 of vomiting viewed as a vis medicatrix, 295, 304 of emetics, fudden, from connection of veffels, 307 ----- of medicines explained by the author, 317, 318, 319 of an emetic is attended with the phænomena of fever, 342 ----- and effects proper to ftin ulant emetics, 345 -- fedative emetics, 346 -- of cold water on extreme veffels of the ftomach, 354 R Opera-

E ...

258 INDEX Operation of cold water on the fkin, by confent, 354 of cold air on the appetite, 356, 357 Opinion, the author's, on the remote caufes of fever, — — 252, 253 -----, Dr. Hunter's, of the various hypothefes on digeftion, — — 258 ____, the author's, of the fecondary use of the ftomach, - 261 _____, confirmed by the records of ancients, from the datara 262 Opinions, various, respecting medical sympathy, 372, 373 Organs of voluntary motion fympathize with the circulation, 35

P

Pain, abfolute, divided by fenfitive fympathy, 151, 152 Parietal mufcular fibres of the flomach, the ufe of, 311 Partial fympathy may arife from local or univerfal difeafe, ______96, 97, 101 ______ difeafe may give rife to univerfal fympathy, 99, 100 Partial

INDEX. 259 Partial fympathy when natural, always falutary, 102, 103 Parts, what, most readily fympathize with the brain, - 34 _____ ---- vary in their sympathies, how, -- 92, 93 particularly connected fympathize most readily, 123 -----, univerfal fympathy has its order of, 125, 126 vital, fympathize from natural connection, 133, 134, 151 ----- without connection, rarely fympathize in difeafe, _____ 135, 136, 137 Parturient pains, when caufed by fympathy, 230 Paffions of the mind produce particular determinations, ____ 202, 203 _____ fome of the, occasionally infective, _____. 207 Pathology, the abforbent fystem, not fufficiently attended to in, ____ 376, 377 _____ of fever, an idea thrown out on, 124 -----, remarks on, why neceflary, 24I Penetration does not affift the operation of volatile and oily applications, - 187, 188, 189 Penis glans; affection of it, from difease of bladder, how uleful, - 153, 154, 155, 156 R 2 Per-

INDEX. 260 Permanence of confriction may caufe coalescence of veffels, 336 Perspiration produced by sympathy with the stomach, 354, 355, 356 Phænomena of the cold stage of fever, - 244 hot ftage of fever, - 245 fweating flage of fever, 246 -----, the whole of the, of fever, connected with the febrile affections of the ftomach, 264, 265, 266 the, of vomiting do not correspond with Dr. Cullen's doctrine, - 290 Plegmafiæ are attended with fympathetic fever, 110 Phyliology of the ftomach, remarks on, why ne--nalo ceffary, 241 Plague, vomiting most violent at the commencement of the, 322 _____ in the, infufficient, 324, 325 is ftopt by the warm bath, 328 Polypufa, confidered as a fimple animal, - 63 child in utero more fimple than a, 66 Prædifpofition caufes individual fympathy to differ, the ov to nonvisoro and min 121, 122 Premature labour, how brought on by fympathy, 230 Primz 2.5 -75"

INDEX. Primæ viæ, the veffels of the, exposed to remote

caufes. 367 Primary effect of vomits, what, 340 Principal remote caufes of fever, 254 Principal fenfes; the five, confcious of fympathetic impreffions, 214 Principle, fensitive, less determinate in the young than the old, - 44, 45, 46, 47 ----- of fimple life fympathized with by the mind, 74 & feg. Principles of life are connected by natural fympathiesin different parts of the body, 143, 144, 145, 146 Propenfities are raifed in us by fympathy, 30

Proximate caufe of fever, 249

vioalitio vitato a cQia

Questions. May not the absorbent system be much difordered in fever? 376, 377 -----. Do external applications ever do good? 185 . May not permanence of constriction cause coalescence of veffels? - 336, 337 ----. How does fear operate as a remote caufe ? 255

R 3.

Queftions.

Queftions. Through what fystem of minute invifible parts does fympathy take place? 199, 352 Do not the remote caufes of fever often act principally upon the stomach? — 343 May not want of fympathy lay the foundation of difease? — 284

R

Re-action, of universal importance in fever, 264, 265, 266 ----- of the ftomach is the act of vomiting, 362, Cor. iii. the phænomena of fever, 363, Cor. iv. Reader requested to compare CCCLXIV. with CCLVI. 364 Records of the ancients confirm an opinion, 262 Rules for the exhibition of medicines contrary to Dr. Cullen's doctrine, 315, 316 1 Remote injuries produce the greateft univerfal fympathies, 118, 119 -----, operation of medicines when, 177, 179, 180 - fympathy, fomentations, et fim. relieve 185, 186 by, Remote

INDEX.

Remote caufe, observations on fever as a, 255 ----- caufes often fupport the debility of the fkin and primæ viæ, -267 ---- may often act principally upon the ftomach, 343 Remoteft parts of the circulation affected by vomiting, 286, 287 Reftauratrix, fympathy constitutes a vis vitæ, 3, 4, 23, 33, 38 Restoration of animation, apparently lost, by vomiting, 33 Reftriction of veffels from cold diminishes the appetite, 357

S

Secretory organs often influenced by fympahy, 370, 371

Senfes, the five, how affected by fympathy, 214 Sentient principle not necessary before birth, 225 Simile to explain the operation of an emetic, 347, 348, 349, 350 Simple animal, a polypus confidered as a, 63 Simple life when in fympathy with the mind, 74. et seq. R_4

Sinapifms

Sinapifms, &c. often operate by fympathy, 164 Sluggishness denotes future excitement, - 291 Soporiferous effects produced by fympathy, 165 Spafm depends upon fympathetic mufcular affection, 26, 27, 28 Specific miafmata produce local specific sympathies, 83 ----- fympathetic inflammation; 86, 87, 89, 90 Stomach confidered as the grand fympathifer, 261 Stomach, fympathizes in hunger with the whole body, 68 ----- is fympathized with by the whole fyftem, 70 , is excited to action by emetics, - 340 , the phyfiology of the, not rightly underftood, 257, affections of the, proportioned to the general atony and confriction, - 306 , why first acted on by the remote causes of fever, and a second a second second second 343 ---- and fkin how fuppofed in confent by Dr. Cullen, 299 Succus gaftricus neceffary to digeftion and appetite, 269, 270, 271, Summary of fever, Dr. Cullen's, 256 Sweating

Sweating stage of fever, phænomena of the, 246 Sympathy is concerned in every division of theinftitutes 3,4 belongs, not more to the nerves than mufcles, 5,6 directs all the principles of life, 31, 32, 80 _____ is an imitative faculty, _____ 29,42, _____ between voluntary motion and circulation, 35 lefs determinate in the young than the - 44, 45, 46, 47 old, ------- varies from being fimple or compounded, 58, 61, 72 when partial from universal difease, 96, 97, 101 ----- univerfal from partial injury, 99, 100 of individuals how different, 121, 122 ——— between the heart and hands, — 166 causes a division of absolute pain, 151, 152 affifts the operation of external applications, ----169 to 199 of the fenfes a proper medical enquiry, 201 Sympathy

INDEX. 366 Sympathy and confent particularly evident in fever, 298 ---- between invisible vessels and muscular fibres improbable, _____ 301 fimilar veffels and veffels more probable, _____ 302, 305, and confent, of the terms, when not applicable, _____ 297, 366, 368 by experiment between nerves and veffels, 374, 375, 7 ----- impreffion what ? _____ 8 _____ produced by bodily 1.1. - IO, II pain, ____ Sympathies vary from cuftom and habit, 12 _____ differ, though arifing from the fame this wait the first but caufe, 13, ____, of confcioufness increased by attention, La Caufes a divition of abindute main, 151 14 remain, when the caufe has ceased, _____ 15 diftant, produced from disease of the brain, _____ 18 when agreable and when difagreable, 19 Sympathy Sympathies

INDEX.

Sympathies of confcioufness arise from corporeal affections, 20 -----, the inherent, directed by the vis conf. et vis, med. -----, the acquired, commonly produced by difeases, 22 ----, inherent constitute a vis vitæ restauratrix, -----3, 4, 23, 33, 38 -----, how called forth by lofs of balance, in the moving fibres, --- 26, 27, 28 ----, not explained by relation to nerves, 36. -----, how produced, -- 37, 41, 43, varied, as arifing from fimple life, fenfation, or the mind, 39 , diffimilar, produced by diftant irritation, - 72, 73 ____, when greatest, 116, 117, 118, 119, 123, ____, founded on natural connection, 133, I34, 141 -, what, cannot be reverfed, 138, 139 - are most useful, and when, 141, 142 , not always apparently useful, 148, 149, 150 Sympathies

267

INDEX. 268 Sympathies most useful to be known, 160, 161, 162 are compared to unifons of found, 58,218 --- how produced in a fecondary way, 220 -- often injure the body, 226, 228, 229,230 ----- are called forth by the operation of a vomit. DATES DOLESS 319 System affected with fever on exhibition of an emetic, 342

T.

Tafte, when offended, varies fympathetic impreffions, 214
Temperament caufes individual fympathy to differ, 121,122
Term, vis medicatrix naturæ, explained, 279
Terms fympathy and confent when improper, 297, 366, 368
Terminations of the arterious fystem, at the fecretory organs, not in balance and connection, 369
Tone of extreme veffels restored by vomiting, 282

Variety

Volatile applications

269

Variety of Sympathies, how caused, 39, 58, 61, 72, 92,93 Variolous attack, suitable reflections on the, 331, 332, 333 Vascular part of the flomach most immediately af-- 267, 298 fected, Vegetable oils are lefs efteemed for external application, than the animal, _____ 190 Venæfection why attended with naufea and vomiting, 274, 275, 276 Veffels and nerves may fympathize, - 374, 375 Villous membrane of great importance in digeftion, 259, 267, 268, 270, 271, 273, 311 ---- first affected by the operation of medicine. 318, 3'9; Vires confervatrices et medicatrices naturæ directed by inherent fympathy, ---21, 32,80 Vires vitæ omnes directed by fympathy, 31, 32, 80 Vis medicatrix naturæ what, 279, 304 Vis vitæ restauratrix constituted by inherent fympathy, - - - 3, 4, 23, 33, 38 Vital parts produce the greatest fympathies when difeased, in this 116, 117 ---- fympathize from natural connection, 133, 134, 141 Volatile

I N D E X.

270

Volatile applications do not relieve by penetration, 187, 188, 189 Voluntary motion, when inordinately affected by fympathy, 26, 27, 28, 35 Vomiting how useful, when it follows venæfection, 276 is a vis med. naturæ, 278 com mences the operations of the vis med. naturæ, 280 rouzes the hear and arteries to action, 281, 340 removes spasm, and restores tone, 282 _____ how it puts a ftop to hæmorrhages, 283 reftores the balance of extreme veffels, 9, 267, 268, 270, 271, 27 284 ._____ ftimulates the remotest parts, 286, 287, 341 218, 210. _____ obviates many diforders of the fyftem, mpathy, 288 21, 32,80 fuppofed to depend on affections of muf-A Cular fibres, 289 ____, the phænomena of, not agreeable to Dr. Cullen's hypothefes, - -----290 fometimes may produce difeafe, - 294 during pregnancy may caufe abortion, 292 02110011333 Vomiting 124, 141 pibalo.7.

INDEX. 271 Vomiting may fometimes reftore life apparently 33 loft, ---- most violent at the commencement of - 1 322 the plague, ---- obviates the danger from excefs of debility in the vital functions, - 323 - in the plague denotes the ftrong exertions of the muscular system, 323, 324, 325, 326 , when violent, may in the end prove hurtful, 294, 327 how ftopt by the operation of the hot-328, 329 bath, in fever, Dr: Cullen's observation on, 334 regulated by ftate of extreme veffels, 339 muscular fystem, — 362, Cor. iii. Vomits rouze the ftomach to action, - 340 , by their operation, give a flock to the whole system, 341

Ule of a corramon fym U held a

INDEX. 272 Unborn children are liable to difeases, - 224 Unctuous applications operate by fympathy, 191; 194, 195 Unifons of found are fympathies in mufic, 58, 218 Universal disease may cause partial sympathy, 96, 97, 101 --- fympathy may arife from partial injury, 99,100 ---- constitutes hestic fever, III, II2 when greateft, 116, 117, 118, 119 has its order of parts, 125; 126 _____ fympathies, fevers in general are, 247 ---- fympathy of the ftomach great, 292, 293, 294 Universality of sympathy through all nature, 231, 232, 233, 234 Unufual exertions, how indicated by languor, and apparent debility, 291 Use of a common sympathetic inflammation, 157, 158 --- of fympathy not always apparent, 148, 149, 150 Uterus,

INDEX. Uterus connected by fympathy with the female 208, 209 mind, Utility of the fympathy of the glans penis, 153, 351 1551 (451 lefs determinate in their fympathies, re fenfitive principle than the old, 44, 43,

W

4 5 47

Warm bath does not relieve by adding moifture to." the body _____ 182, 183, 184, 185, 186 , why recommended by Dr. Sydenham in the plague, ____ - 328 acts by a general ftimulus, as well as topical relaxation, 329 ----- might be useful at the beginning of most fevers, 330 Warmth is neceffary to produce fome fympathies, 71 Want of fympathy in the ftomach may lay the foundation of difease, 284 Water, draught of cold, does not affect the fkin by having entered the circulation, 355 What sympathies are most useful, - 141, 142 particularly ufeful to be known; 160, 161, 162 When sympathies are most useful, - 141, 142

S

XYZ

273

274 I N D E X.

DE VO SVS

182. 162. 1

XYZ

Young people lefs determinate in their fympathies, and fenfitive principle than the old, 44, 45, 46, 47

FINIS.

E R R A T A.

Par. CXVII. For deceased, read diseased. Par. CCLII. For (last part of CCLXVII. read CCXLVII.) For sympathic, read sympathetic.





