An experimental history of the materia medica, or, of the natural and artificial substances made use of in medicine: containing a compendious view of their natural history, and account of their pharmaceutic properties, and an estimate of their medicinal powers, so far as they can be ascertained by experience, or by rational induction from their sensible qualities / by William Lewis.

Contributors

Lewis, William, 1708-1781. Aikin, John, 1747-1822. University of Glasgow. Library

Publication/Creation

London: printed for J. Johnson; and R. Baldwin, 1784.

Persistent URL

https://wellcomecollection.org/works/cnev3knk

Provider

University of Glasgow


License and attribution

This material has been provided by This material has been provided by The University of Glasgow Library. The original may be consulted at The University of Glasgow Library. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



18th Sprit


-g 31_a.1




Bibliotheca Universitatis Glasquensis

64.3.7.

Q193.16_





EXPERIMENTAL HISTORY

OF THE

MATERIA MEDICA,

OR OF THE

NATURAL AND ARTIFICIAL SUBSTANCES

MADEUSEOFIN

MEDICINE:

CONTAINING

A Compendious View of their NATURAL HISTORY,

AN ACCOUNT OF THEIR PHARMACEUTIC PROPERTIES,

And an Estimate of their MEDICINAL POWERS, so far as they can be ascertained by Experience, or by rational Induction from their sensible Qualities.

By WILLIAM LEWIS, M.B. F.R.S.

THE THIRD EDITION,
WITH NUMEROUS ADDITIONS AND CORRECTIONS
BY JOHN AIKIN.

Rationalem quidem puto Medicinam effe debere: instrui vero ab evidentibus. CELSUS.

LONDON:

Printed for J. Johnson, in St. Paul's Church-Yard; and R. Baldwin, in Pater-nofter-Row.

Digitized by the Internet Archive in 2015

https://archive.org/details/b2147025x

ADVERTISEMENT

TO THE THIRD EDITION.

A Confiderable time having elapsed since the last publication of Dr. Lewis's very valuable work, it was thought necessary, now that a new edition was demanded, to procure such additions and corrections, as later improvements in natural history and medicine had produced, and which the author himself, had he now been living, would undoubtedly have made. For this purpose it was put into the hands of the present editor, who has in the following manner endeavoured to fulfil the intention of the publishers.

To every vegetable article he has subjoined the Linnæan name.

He has corrected all the references to the Edinburgh Pharmacopæia, both in the catalogue of fimples, and the officinal preparations, by the last edition of that work.

He has added feveral entirely new articles; among which are all the new medicines received into the present Edinburgh catalogue, and such others as have been recommended upon apparently

apparently fufficient grounds in various publications, domestic and foreign.*

From the same sources, and also from some marginal notes in Dr. Lewis's own copy of his work, he has derived many additional facts and observations relative to sormer articles. All the additional matter is distinguished by an asterisk prefixed.

At the same time that the editor has so freely added to this work, he has not thought it allowable to omit any thing; though some of the articles are now expunged from the catalogues of materia medica, and the state of medical theory has undergone considerable changes since the author wrote. It is still, in every respect, Dr. Lewis's entire work.

WARRINGTON, October 1, 1783. JOHN AIKIN.

* These are the new Articles.

Aer Fixus.
Cardamine.
Columbo.
Filix Mas.
Flammula Jovis.
Geoffræa Jamaicensis.
Lichen Islandicus.
Lobelia Syphilitica.

Oenanthe Crocata.
Peruvianus Cortex Ruber.
Cinchona Carribæa.
Pulfatilla Nigricans.
Quaffia.
Radix Lopeziana.
Rhododendron Chryfanthemum.
Spigelia.

Stramonium.
Viola Tricolor.
Winteranus Cortex.
Aconitum Napellus.
Cursuta.
Hippocastanum.

ERRATA.

Page 234. It is faid, by mistake, that the extract of the seeds of hemlock, is the only officinal from that plant directed in the last Edinburgh dispensatory. The common extract, made with the juice of the leaves, inspissated, and a fifth part of the powder of the leaves, is ordered under the title of Succus spissatus Cicutæ.

P. 396. Lingua Cervina is not expunged from the Pharm, Edinb.

N. B. See the Poffcript for fome emiffions.

PREFACE.

HE medicinal materials, in the infancy of physic necessarily few, were by degrees exuberantly multiplied; and new ones are still, from time to time, discovered and introduced. Their estimation and use have been variable; substances at one time in high esteem being often at another disregarded, and those which in one age had fallen into neglect being often in another revived; a sluctuation apparently owing, in many articles, to the fancy or caprice which influences other human things, and in many to ignorance or error. The design of the present work is, to examine the several substances which are or have been in repute; with a view to ascertain, as far as possible, their real powers, and to establish this important part of medicine on a just soundation. It will, perhaps, appear from this examination, that there are some materials of little significance among those which are retained in practice, and many of more utility among those which are overlooked.

The materia medica is commonly understood to comprehend, not only the materials afforded by nature, but many of those also which are prepared, produced, or compounded by art; as minium, potash and soap. In pharmacopæias, intended as directories, for the preparation of such medicines only as are made in the shops, all those artificial medicinal substances are referred to the materia medica, which are either brought from abroad, or commonly prepared by particular persons, as articles of commerce. A scientisic distinction

Productions effentially different, in their medical and other properties, from the subject from which they were produced, and which had nothing analogous to them præexisting in the subject; as the fixt alkaline salts of vegetables, and the volatile salts of animals;—compounds resulting from the coalition of opposite ingredients; of ingredients which in mixture lose their specific powers, and form together a substance of new qualities; as neutral salts and soap;—the constituent parts of natural compounds of this kind, separated and purified by art; as magnesia and the mineral acid spirits;—may be considered as distinct medicinal materials, or as articles of the materia medica, wheresoever they are prepared.

It were to be wished, that the several subjects could be methodically arranged, from some qualities subservient to medicinal intentions. This has been attempted by different writers, on different plans, but in my opinion with little success: nor indeed does it seem to be practicable; the qualities of medicines being too intricate, and compounded, and multifariously diversified, to serve for the basis of any useful distribution of them. The division into minerals, animals, and vegetables, and the subdivisions into roots, barks, leaves, slowers, &c. are equally exceptionable; some substances not being clearly reducible to either of the three kingdoms, and different parts of one vegetable being commonly made use of. I know of no practicable method that promises any advantage above the alphabetic one; and what conveniency there is in this regards rather the author than the reader.

Each article may be considered in three points of view; as an object of natural, medicinal, and pharmaceutic bistory.

The office of natural history, so far as it relates to the materia medica, confists, in distinguishing the several substances from one another by criteria drawn from their external form and structure, and in ascertaining their origin and production.

The criteria of natural history are peculiarly adapted to vegetables and animals in their entire and perfect state; the form, structure, and disposition of the several parts, collectively considered, affording here generally sufficient means, and indeed the only means, of distinguishing each particular species from all others. In this branch of knowledge, of late years so diligently cultivated and so remarkably improved as a general science, little new matter

can be expected in a work of the present kind. If, of the accounts given in detail by those who have written professedly on these subjects, the more interesting and useful particulars relative to the medicinal articles, are perspicuously and concifely expressed; if the more obvious and invariable discriminative appearances are justly selected from the writings of others or assigned from my own observation, so as to render the descriptions strictly definitive or characteristic, without regard to the systems of naturalists; I have in this point accomplished my intentions.

Of the distinct parts and productions of vegetables and animals, there are many, which cannot be sufficiently discriminated by any external marks, and which require the assistance of characters drawn from pharmacy or pharmaceutic chemistry, that is, from their intrinsic properties. The criteria of natural history, strictly so called, are still more insufficient, and those of chemistry of consequence more necessary, in regard to the products of the mineral kingdom; where, oftentimes, one and the same matter assumes different forms, and different kinds of matter the same form. In assigning the criteria drawn from this fource, I have endeavoured to determine by experiment those properties, which, at the same time that they are obvious and easily examined, may be fully characteristic of the subject in all its forms.

The medicinal history, or the knowledge of the powers and effects of medicines in the human body, though apparently a most essential branch of the bealing art, has been far more incuriously cultivated, and still, perhaps, continues less cleared from the errors of former ages, than any other science. Even in these later times; after the arbitrary qualities of cold, hot, dry, and moist, and the ridiculous similitudes and conceits which some enthusiasts of the last century relied on as a test of medicinal activity, had been exploded: the advancement of true medicinal bistory has met with many obstructions; partly from the officiousness of compilers in collecting and preserving the fictitious virtues; partly from a fondness, in original writers, of aggrandizing their favourite medicines; partly from a fallacy, in ascribing to a particular ingredient in a composition the effects which more powerful ones had produced, a fallacy which the exuberance of mixture made sometimes unavoidable; partly from the difficulty of distinguishing, in many cases, the real effects of medicines from the operations of nature unassisted; partly from a practice, too common among writers on the materia medica, of barely enumerating the diseases, or even the parts of the body in whose general diseases, a medicine

B 2

had, or was supposed to have, done good, as if diseases of the same parts, or of the same name, were always of the same nature, or were always to be treated by the same remedies. Medicinal history has perhaps suffered also from the misapplication of other sciences.

After botany had been regularly cultivated, and methodized into systems; it was observed that several of the plants, which had been ranged together from their agreement or affinity in botanic characters, agreed or were allied also in medicinal virtue. As nature appeared to have in some instances established a connection of this kind, it was fondly presumed that she had done so in all; that, the virtues of some particular plants being known, those of all the others, ranked in the same botanic class, might also be inferred; and accordingly rules have been drawn up, for judging of the virtues of plants upon this principle, by botanists of the greatest name. But so far is a similarity of virtue from obtaining through the several genera which constitute one class of vegetables, that frequently it does not obtain through the several species of one genus: there are solanums, lettuces, herb-mercuries, cucumbers, mushrooms, &c. esculent and deleterious; and even the same individual often varies, from culture or other circumstances, as much as two plants which have no botanic affinity.

The chemists, in like manner, extending the discoveries of their useful art beyond the proper limits, endeavoured to investigate the virtues of plants from the substances into which they are resoluble by fire; and in this view, the French academicians analysed almost all those made use of in medicine. From their experiments it appears, that the substances thus obtained have no resemblance in quality to the original vegetable, and can afford no foundation for judging of its virtues; that plants the most remote in virtue, purgative and astringent, poisonous and esculent, are changed by force of sire into similar principles. It is matter of concern, that these analyses should have been preserved in the posthumous works of a writer so judicious as Mr. Geoffroy, while the editor was sensible that the author himself, in his later years, disapproved of them.

There are, nevertheless, in most vegetables, certain sensible qualities, either obvious, or easily discoverable; which afford, under due restrictions, an excellent test of their virtues; and in which indeed, oftentimes, their virtues wholly

wholly confist. Aromatics, acrids, fetids, astringents, bitters, sweets, acids, unctuous and mucilaginous substances, which comprehend the greater number. of the articles of the vegetable kingdom, operate generally by such qualities as are the immediate objects of smell and taste; and from the degree of force with which they affect those senses, their degree of medicinal efficacy may be generally inferred. The smells and tastes of the several materials, on which some have already laid considerable stress, but which for the most part have been either wholly neglected, or regarded only as they affect the medicine in point of elegance, I have examined with no little care: and though it is not to be supposed, that the particular degree of each can be precisely determined; or its particular species, especially in regard to smell, fully expressed in words, any otherwise than by comparison with substances more known; or that any exact limits can be always fixed, as between fetid and aromatic, grateful and ungrateful; I nevertheless flatter myself, that the observations of this kind will furnish, in many instances, sufficient data to the physician for judging what may be expected from materials he has not experienced. In this part, as in the descriptive history, great assistance has been drawn from pharmaceutic chemistry: for, in many vegetables, the active matter is so far divided and diluted by the berbaceous inert substance; and in others, different kinds of active matter are so blended together; that they cannot be discovered, or distinguished, till they are extracted, or separated from one another, by the operations of chemistry.

Some other experiments are, in particular cases, very useful auxiliaries in this enquiry. Thus, a solution of vitriol of iron, made in water, is by many vegetables, turned to a black colour; by others, a solution of sulphur, made in alkaline liquors, is rendered milky or turbid, and of a strong setid smell. It is not known, that any vegetable substance produces the first of these effects, but those which have an astringent power; or that any produces the second, but those partaking of an acid, which unites with and neutralizes the alkaline matter in the liquor, and disengages the sulphur which was thereby kept dissolved. By these criteria, lower degrees of astringency and acidity are often discovered, than the taste gives any notice of.

The effects of medicines on the fluid and solid parts of dead animals; as their producing or resolving coagulations, relaxing or contracting the sibres, promoting or retarding putrefaction, or varying the degree and the species

as well as the facility of the resolution; afford likewise, in some cases, considerable light into their medicinal operation. Of these cases, however, the number appears to be much smaller than seems to have been imagined by the generality of those who have prosecuted these enquiries; who have not, perhaps, fufficiently considered, bow different is the operation of medicines on animal substances in a vital and in an inanimate state; and how much the fluids of an animal are influenced by the action of medicines on the folid parts: it is probable, that the operation of most medicines is immediately or principally upon the folids, and that the fluids are in most cases only consequentially affected. The fluids most likely to answer any useful purpose, in these sorts of trials, are those which are secreted into the alimentary canal: experiments on blood seem to be of no medical utility: green vitriol, mixed with the blood drawn from a vein, instead of rendering it more florid, the common medical effect of this chalybeate preparation, changes it grey: mercurials examined in the same manner, discover nothing of that remarkable colliquation, which they produce in the blood of living animals.

There are substances, in which all the foregoing means of investigation fail; and which operate by some latent power, of which they give little or no intimation to the senses. Of this kind are most of the purgative, emetic, and narcotic plants; those, which from their deleterious effects when taken in certain quantities, are called poisonous; and most of the metallic bodies and their preparations. Experiments on brutes are here of use, but of limited use: for if prudence requires us to refrain from substances which are noxious to brutes, it does not always authorize us to venture on fuch as may to them be innocent: experience shews, that the crocus of antimony, of which a grain or two operate on the human body as a virulent cathartic or emetic, may be given to borfes in the quantity of an ounce, without producing any very remarkable effect: that a moderate dose of jalap throws a dog into convulsions, who could well bear a much greater quantity of opium than could be given with safety to a man. The virtues of these kinds of substances can be known only from their effects in the human body itself: and as, of all medicines, they have the most obvious and apparent effects; they are, happily, of all medicines, those which admit of the least deception, and in which, of consequence, we can most avail ourselves of the observations of former writers. Indeed many of them being now received in general practice, their powers have been determined by general experience.

With such assistances as I could draw from these sources or from my own experience, I have endeavoured to point out chiefly the primary effects of the several subjects, or the immediate sensible operation which constitutes their true medical character. I judged it useless to enter into an enumeration of diseases in which a medicine is or is not proper, when the salutary or pernicious effects, which it produces in those diseases, are no other than obvious consequences of its general power: it nevertheless appeared frequently necessary to specify some particular cases, as being either illustrative of the general power, or subservient to its discovery, or where it could not be precisely ascertained.

The pharmaceutic history of simples, closely allied to the medicinal, regards, chiefly, the variations of their qualities in different states and forms naturally or artificially induced; the separability or non-separability of their active principles by different menstrua or different operations; and their miscibility or non-miscibility one with another. In these properties, remarkable diverfities and contrarieties are observed among the different medicinal simples, even among those in which no material disagreement has been generally suspected. Thus, the virtue of some vegetables accompanies the fluid which they yield on being pressed, while that of others remains behind locked up and concentrated in the subject, and that of others is destroyed in the operation: some plants, in being dried, lose all their virtue, some bave their virtue improved, and some bave it changed to another kind: some, by infusion, give out their virtue both to water and to spirit of wine, some to water only, some to spirit only, and some neither to one nor the other. Nor can these diversities be reduced to any general rules, or any otherwise determined than by a separate examination of each particular article.

This province belongs peculiarly to chemistry; but notwithstanding its obvious importance to the practice of medicine, even the medical chemists have been very remiss in the cultivation of it. I know only of two persons, whose labours have been considerable, and whose success may be applauded. Neumann, one of the first who, rejecting the useless analyses of vegetables made by vehemence of sire, endeavoured to separate their component parts, unaltered, by means of menstrua; examined by this method a considerable number of the officinal drugs, not indeed directly in a medicinal view, but in the way of a general chemical enquiry into the products of nature. Cartheuser, consining himself more closely to medicinal considerations, followed nearly Neumann's plan so far as it included

thefe, and made fundry valuable additions.

In the present work, the enquiry is extended to a far greater number of simples, and conducted likewise on somewhat different principles. The quantity of matter, which water or spirit extract from a plant, or which either menfruum extracts after the action of the other, is, medicinally, of little importance to be known; unless it be known also, what are the precise qualities of the several preparations, whether the virtue of the plant refides in the part extracted by one or the other menstruum, or whether in this separation of the parts of the subject, any active matter is discovered which was not perceptible before. And on the other hand, though the qualities of the infusions, extracts, &c. be very carefully and minutely examined; yet if they are described independently of one another, and if no account is taken of the remaining substance of the plant, or of the vapour that exhales in the inspissation of the spirituous tincture; it will be impossible, in many cases, to judge between the dissolving powers of water and spirit, or whether either is a complete menstruum for the active parts, or whether the spirituous extract retains the full virtue of the subject, or whether a part of the virtue exhales or distills with the spirit. Without embarrassing the reader with a minute history of experiments, I have given only their refult, or the general pharmaceutic habitude of the subject deduced from them: it is in trying to make these general deductions from the experiments bitherto published, that their insufficiency, in regard to the greater number of the articles, is most conspicuous. I have nevertheless been obliged, in some cases, by the multiplicity of the labour, and the difficulty of procuring specimens of some few articles, to be satisfied with such information as those experiments

To prevent the necessity of frequent repetitions under the particular subjects,

some observations of a general nature are here premised.

1. Of the collection and curation of Simples.

Vegetables should be gathered chiefly from those soils, in which they naturally delight, or in which they are found most commonly to rise spontaneous; for, though many of them may be raised, and made to grow with vigour, in very different ones, their virtue generally suffers by the change. A variation of seasons occasions also differences considerable enough to require, oftentimes, an allowance to be made in the quantity; plants in general proving weaker, though more luxuriant, in rainy than in dry ones.—Herbs and slowers are to be gathered in a clear dry day, after the morning dew is gone off from them. Leaves, for the most part, are in their greatest perfection, when come to their full growth,

just before the slowers appear: flowers, when moderately expanded: seeds, when they begin to grow dry, before they fall spontaneously: woods and barks, as is supposed, in the winter: annual roots, before the stalks begin to rise: biennial roots, in the autumn of the first year: perennial roots, in the autumn, after the leaves have fallen, or early in the spring before they begin to vegetate. To most of these rules there are some exceptions, which are specified

under the particular subjects.

Of the vegetables which lose their virtue in being dried, the greater number, perhaps all, may be preserved for a considerable length of time, by impeding the exhalation of their native moisture; for so long as they retain this, they seem to retain also their medical activity. Thus roots have their virtue preserved by being buried in sand, which should be dry, that they may not vegetate; leaves and slowers, of a more corruptible nature than roots, by being beaten with about thrice their weight of sine sugar to prevent their

corruption, and kept in a close vessel.

Plants which bear drying, are commonly hung in a warm airy place, defended from the fun. The colours of herbs and flowers are for the most part changed or destroyed, in drying, by the sun's beams; but that their medicinal virtue suffers a like diminution, does not appear. Thus much is certain, that a heat of culinary sire, equal to that of the sun in summer, does them no injury in either respect: and that both slowers and leaves, when thus hastily dried by sire, preserve the liveliness of their colour, and their smell and taste, more perfectly than by slow exsiccation. The leaves of moderately juicy plants are reduced, by drying, to about one fourth of their original weight.

Some roots, and some other parts of vegetables, how thoroughly soever they have been dried, are liable, in keeping, to grow mouldy and carious. This inconvenience might probably be obviated by dipping them, when dried, in boiling spirit of wine, or exposing them to its vapour in a close vessel. It is said, that some of the oriental spices are made less perishable, by being dipt in

a mixture of lime and water.

The pulps of fruits are separated from the seeds and membranous parts, by forcing them through a strong hair sieve. If the fruit is unripe and hard, or if it is dry, it should be previously softened by boiling in a little water; and the pulp, after passing through the sieve, is to be inspissated over a gentle sire, with care to prevent its burning.

The concrete gummy-resinous juices brought from abroad, which have usually a considerable mixture of bits of stalks, leaves, seeds, &c. are purified, by adding so much boiling water, as will so far soften or dissolve them, that they may be pressed, whilst hot, through a strainer; and then inspissating the strained liquid, in a gentle heat, to the original consistence of the gummy-resin: if the quantity of water is considerable, the resinous part commonly separates and subsides, and in this case is to be kept by itself till towards the end of the inspissation of the gummy, at which time they may be easily united again together into an uniform mass. Some of the gummy-resins, exposed to the heat of boiling water, melt thin enough, without any addition, to be pressed through a canvas strainer. In either process, the operator must be careful to prevent, as much as possible, the dissipation of the more volatile parts; an injury which cannot be wholly avoided, especially when the subjects are dissolved by water. The siner tears unpurished are in many cases preferable, for internal use, to those that have been strained.

Pulverable bodies of an earthy texture, or such as are brittle and not dissoluble in water, after being reduced to a powder of moderate fineness, are brought to an impalpable or very subtile state, by grinding them with a little water on some bard smooth instrument: the matter is commodiously dried on a chalk-stone, or rather on a cake of plaster-of-Paris, which equally absorbs the moisture, without adhering to the powder, like substances of the chalky kind. Powders thus levigated are still found to contain a quantity of gross parts; which may be separated by shaking the matter with water, till it is diffused through the fluid, and then suffering it to settle: the grosser parts soon subside; and the turbid liquor, being now poured off, deposites more slowly the finer powder. By this process, powders may be obtained of any degree of fineness; the tenuity being in proportion to the length of time that they remain suspended in the fluid. On the same principle, the bolar earths may be separated from the gritty matter naturally mixed with them, metallic bodies from those of the earthy kind, and the calces of metals from metallic particles uncalcined.

Salts are purified from indisfoluble admixtures, by folution in water and filtration through paper. Water disfolves, in a boiling heat, a much larger quantity of most kinds of falts than it can retain when cold: thus, of nitre, it disfolves when boiling near three times its own weight, but in cooling, a part of the falt gradually separates, till at length, when grown thoroughly cold,

cold, in frosty weather, it does not retain one eighth its own weight, or one twenty-fourth of the quantity of falt at first dissolved. The neutral salts, or those composed of an acid and an alkali; several of those which consist of an acid and an earthy or metallic body; and many of the acid falts of vegetables; in this separation from their solutions, concrete, unless too bastily forced together by sudden cooling, or disturbed by agitation or other causes, into transparent masses, of regular figures peculiar to each particular kind of falt, and thence called crystals .- There are two general methods of recovering falts from their folutions in a crystalline form; one adapted to some falts, and the other to others. The one is, by keeping the folution in a gentle and equable warmth, that the water may gradually exhale, and leave the falt crystallized. The other is, by boiling down the solution, till, on dropping a little of it on a cold glass plate, crystalline filaments appear; then covering the veffel, and suffering it to cool very slowly. Some of the difficultly crystallizable falts are made to shoot more freely, by adding, after sufficient evaporation, a small proportion of rectified spirit of wine, which weakens the disfolving power of water on most kinds of saline bodies .- As different salts require different quantities of water to keep them suspended; when two or more are dissolved together, they begin to concrete at different periods of the evaporation, that, which requires most water for its dissolution, shooting first, and leaving the more soluble dissolved: on this foundation, salts are purified, by crystallization, from admixtures of one another.

2. General pharmaceutic analysis of vegetables by incision and pressure.

The medicinal juices of vegetables commonly refide in distinct vessels; and often exude upon the surface, from a spontaneous rupture, or artificial incisions of them. The juices of herbs and trees issue chiefly during the summer heats; as the gum of the cherry-tree, the resin of the sir, the sweet juice of the manna ash, and the unctuous exudations on the leaves of many plants; though some trees, as the birch and maple, yield early in the spring, on being bored or deeply wounded, a very copious sweetish watery juice, of which, in summer, they yield little or nothing, the watery menstruum being now perhaps consumed. Some roots bleed gummy-resinous juices in the spring, as bryony and angelica among us; others in summer, as the asafetida roots in Persia, and the scammony in Syria. Some fruits, particularly the several varieties of lemons, citrons, and oranges, have numerous vesicles, in their

their outer rind, filled with a fragrant oil; great part of which may be extracted, by rolling the fruit on a plane stuck full of sharp points, which lay open the oily vesicles, or by rubbing it on a mass of sugar, which imbibes the oil.

From succulent berbs and fruits, the different fluid juices they contain are forced out, mixed, by bruifing and pressing them. Vegetables of the fiveet or saline kind, as the summer fruits and the acid herbs; several of the acrid plants, as arum and scurvygrass; and those of the lattescent kind, as dandelion and the spurges; generally give out by this process great part of their active matter along with the watery fluid: but the juices expressed from aromatic berbs, as mint, have, for the most part, little or nothing of the peculiar smell, taste, or virtue, of the subject; and many of the fragrant flowers, as lilies and violets, have their fragrance destroyed by the pressure. The juices of plants, thick, turbid, and very impure when newly expressed, by settling and repeated straining become clear: many, in this depuration, lose nothing considerable of their virtue: from others, the medicinal parts, not dissoluble in watery fluids, separate and subside along with the feculent matter. To the depurated juices, defigned for keeping, a small proportion of " restified (pirit of wine may be added, which, on standing for some time, generally throws down a fresh sediment: the liquor is then to be put in small bottles that have been washed with spirit and dried, a little sweet oil poured on the furface so as nearly to fill the bottles, and the mouths slightly stopt: by this method, most of the juices that bear depuration may be preserved, in a cool cellar, for a year or two; excepting perhaps only the very fermentable sweet ones, which can scarcely be long restrained from fermentation without boiling. Those which are not injured in their virtue by evaporation, may be inspissated to the confistence of a syrup, or of a thick or solid extract: from those of the faline kind, duly depurated and inspissated, the faline part commonly separates, on long keeping, in a crystalline form.

The kernels of fruits, the seeds that on being triturated with water form an emulsion or milky liquor, and some other vegetable substances, yield, on being strongly pressed, an oil; which, of itself, is slavourless and inspid; but which, in some cases, is impregnated with the smell and taste of the subject. The aromatic seeds and kernels, as annifeeds and nutmegs, and some of the purgative ones, as the ricini, are the principal substances that give out with their oils their peculiar virtues, the oils of most of the others having no particular impregnation. There are, however, considerable differences among the unstavoured

unflavoured and insipid oils; in their consistence; in their disposition to congeal by cold; in their disposition to grow rancid by heat; in the degree of heat necessary to make them boil; in their power of dissolving certain bodies; and in their combinability with fixt alkalies into soap. The extraction of the oil from the subject is greatly facilitated by heat; and hence the preparers of these oils for mechanic purposes generally warm to a considerable degree either the subject itself, or the iron plates of the press, or both. Where the product is intended for medicinal use, this practice is generally condemned; heat being apt to impress upon the oil an ungrateful flavour, and increase its disposition to become rancid. Nevertheless, though a great heat has undoubtedly these effects, yet a gentle warmth is in some cases necessary, and not, perhaps, very injurious in any: in winter at least, both the subject and the apparatus may be warmed, with safety, to the greatest degree of heat that obtains in the shade in summer. The heat is never to be continued longer than the expression requires; and the oil, as soon as drawn, is to be set in a cold place.

3. General pharmaceutic analysis of vegetable and animal substances by menstrua.

Water, the proper menstruum of gums, of gellies, and of salts, extracts the gummy and saline parts of vegetables, and the gelatinous matter of animals. By the mediation of these principles, it dissolves others of more activity, oils and refins being made miscible with water by the mediation of gums. All-the substances which water extracts from vegetables or animals, it dissolves almost unlimitedly, so as, by repeated infusion on fresh parcels of the subjects, to become more and more impregnated with their active parts, till so far loaded as to have its further action impeded by the diminution of its fluidity: it generally takes up first the lighter and more grateful matter, and afterwards the groffer and more difagreeable; and hence, by skilful management, it may be richly impregnated with the former, without much admixture of the latter. The subject should be moderately dried (unless it be of such a kind as to lose its virtue in drying) as in this state it communicates a remarkably stronger impregnation than an equivalent quantity when fresh. Most leaves and flowers yield a great share of their more active matter by cold maceration, or more readily by warm infusion: by boiling, the dissolving power of the water is for the most part greatly increased, and the volatile parts, if the subject contained any, are distipated with its steam.

The vapour which exhales in the boiling of odorous substances, and many of the pungent vegetables, in water, collected and cooled in proper veffels, forms a liquor impregnated with their smell and pungency. This impregnation depends on a subtile principle, whose matrix is a volatile oil; of which oil a part often separates in its own form, either floating on the surface or finking to the bottom according to its gravity; and which, from the specific flavour of the subject being concentrated in it, is distinguished by the name of effential. As this subtile oily matter is here separated from the more fixt gummy parts that rendered it before almost unlimitedly dissoluble, the water now can retain only a certain proportion of it, and generally but a small one: if the distilled water, once saturated so that a part of the oil appears distinct, be redistilled from repeated fresh quantities of the subject, the aqueous fluid receives no further impregnation, and the quantity of oil that separates proves proportionably larger than if fresh water had been used. These oils differ from the expressed ones formerly mentioned, in being dissoluble in spirit of wine, and volatile in the heat of boiling water; on either of which foundations, when expressed oils are mixed with essential ones, whether artificially, or in their expression from subjects that contain both, the two oils may be completely separated from one another. The sophistications of the dearer essential oils, commonly practifed, by the admixture of cheaper ones, can be distinguished only by the smell and taste: the smell and taste which they communicate to liquors in certain known quantities, when dissolved in spirit, or, by means of fugar or mucilages, in water, is an useful criterion also of the degree of goodness of the oils when genuine; for the same kinds of vegetables, produced in different foils and seasons, vary not only in regard to the quantity of water they are capable of impregnating by distillation, and in the quantity of oil they afford, but likewise in the strength of the oil itself.

There are some substances whose virtues reside wholly in an essential oil, and are whosly dissipated in boiling: there are others, which have one virtue residing in an essential oil, and another which remains behind in the decoction, which last may be brought to a concentrated state by evaporating the watery menstruum with a gentle heat till the matter becomes thick or solid: there are others, which contain no oil, and whose virtue remains entire in the inspissated extract, provided it has been skilfully prepared. A difference in the quantity of water will in many cases occasion a sensible variation in the qualities of the extract, by requiring more or less heat for its evaporation; for, independently of the dissipation of the volatile parts, even those of the more sixt kind suffer a

confiderable

considerable change from continued beat: by long boiling with water, sweet substances become nauseous, and the drastic purgatives lose their virulence, without any remarkable separation of their parts. Some have endeavoured to avail themselves of this observation, for converting the stronger cathartics and emetics, asarum, tobacco, and others, into medicines of safety and utility; and report that extracts made from these plants, by long boiling with a large quantity of water, were found to act as mild aperients or deobstruents: these kinds of preparations, however, must necessarily be too precarious in strength to be received in general practice; the abatement of the virulence of the medicine depending on what no care can adjust to one standard, the degree and continuance of the beat.

Pure spirit of wine, the appropriated dissolvent of refins and essential oils, and which dissolves also certain saline bodies, as the sweet saccharine salts of vegetables; extracts, for the most part, such virtues of vegetable and animal fubstances, as reside in those principles, or in principles analogous to them. Of the substances, whose virtues reside apparently in these principles, there are many, which give a strong impregnation to water as well as spirit, but few that impart their virtues in an equal degree to the two menstrua: from a compound of pure gum and refin, water, by infusion, extracts directly the gummy matter, and by the intervention of this, a part of the more active refin, leaving great part of the refin undiffolved: whereas, contrariwife, pure spirit extracts directly the refin, and leaves undiffolved greatest part of the inert gum, of which it does not appear to take up so much as water does of the resin. Hence, in the analyses of these kinds of subjects, it is generally observed that spirit loads itself with their active parts much more than water is capable of doing: that the extracts made with spirit are much smaller in quantity, and proportionably stronger, than the watery extracts: and that the Spirituous tinctures, loaded with the refinous parts, grow turbid on the admixture of water, and deposite their pure refin; the gummy matter, that the spirit had taken up, remaining diffolved in the aqueous fluid, and being infufficient in quantity to keep any considerable portion of the resin suspended. Hence saturated resinous tinctures, those especially of the cathartic kind, require, in being diluted for exhibition, an admixture of gummy or faccharine matter, to keep the refin divided, and to prevent its separation: on this foundation may be prepared, from these kinds of tinctures, elegant gummy-refinous extracts; by mixing with them, when inspissated to the confistence of a balfam, a thick solution of any simple gum or mucilage, and continuing the evaporation, with a gentle heat, till the matter becomes

becomes dry. In like manner, the refinous and gummy parts of one subject, or those parts which pure spirit extracts, and which water extracts after spirit, may in some cases be advantageously united into one mass; by separately inspissating the tineture and decoction to a certain thickness, and then mixing them together.

Pure spirit, which exhales or distils with a much less heat than water, carries off with it, for the most part, much less of the essential oils of vegetables. There are many substances, whose active parts are almost wholly dissipated in the preparation of the watery extract, and almost wholly retained in the spirituous. There are some, however, whose oils are so volatile, as to rise with pure spirit as perfectly as with water; and in this case, the distilled spirit proves sometimes stronger than the distilled water; spirit keeping dissolved all the oil that rises with it; whereas, when water is used, a part of the oil frequently separates.

Wines, as being compounds of water and inflammable spirit, take up such parts of vegetables and animals as are soluble in those liquors; but their dissolving power is somewhat weaker than that of purer mixtures of water and spirit, on account partly of their viscous unctuous matter, and partly of their acid. Wines are impregnated with the active parts of medicines, chiefly by maceration in the cold, or with a very gentle warmth; the heat, which is often advantageously applied for expediting or promoting the action of water and spirit, occasioning in wines a disagreeable alteration. Malt liquors are commonly medicated, by macerating the ingredients in them during the fermentation, or boiling or infusing them in the wort.

Acids, both vegetable and mineral, somewhat weaken the dissolving power both of water and spirit on bodies of the resinous and oily kind; and when added to insussions or tinetures, generally precipitate a part of what the menstruum had before taken up: nevertheless, when acids are intimately combined with vinous spirits into what is called a dulcified liquor, the compound proves a more essimated increase the action of water on resinous and oily bodies, and weaken or restrain its action on gummy ones, rendering water incapable of holding pure gums dissolved: they have been supposed to promote the action of spirit on dissincture, the quantity extracted is found to be the same without as with it.—Volatile alkalies precipitate gums from water, like the fixt, but in other cases their effects are more variable.

Expressed

Expressed oils extract the odoriferous, refinous, and oily parts of vegetables; and with these they are sometimes impregnated, both for the purposes of perfumes, and for external medicinal uses. The more fragrant flowers give out their odour by cold maceration. The more fixt refinous parts of the leaves of plants are extracted, by boiling them in the oil till their watery moisture has exhaled, that is, till they are almost crisp, and the oil appears tinged of a green colour. Animal fats, liquefied and boiled with recent berbs, become in like manner impregnated with their refinous parts, and with the green colour residing therein. It does not appear, that the oils or fats receive, by this management, any valuable virtues: the heat, requifite for making them boil, impresses an ungrateful scent, and dissipates the more volatile parts of the subject: they may be impregnated more elegantly with the active parts of vegetables, by an admixture of effential oils or of spirituous extracts. The spirituous tinctures of the leaves of most plants are of a deep green colour; and the inspissated extracts, though often brown or black in their folid or confistent state, give generally a like greenness to fresh spirit, to essential oils, expressed oils and fats.

Air, or its watery moisture, seems to act as a true dissolvent, in the same manner, though not so expeditiously, as water in its grosser form. astringent virtue of the walnut tree, and the purgative of the damask rose, have been observed to be diffused through the air; though they are obviously not of the volatile but of the fixt kind, not exhalable by heat but dissoluble by menstrua. Hence the atmosphere may become impregnated with all those virtues of vegetables, which at least watery menstrua can extract: and bence many medicinal substances are gradually robbed by it of their virtues;

powders the most speedily, as exposing the largest surface to its action.

4. General chemical analysis of vegetable and animal substances, and mineral bitumens, by fire.

Vegetable substances, burnt in the open air, are resolved, partly into smoke, which, condensed, forms soot; and partly into white ashes, which generally give out, on being boiled in water, a fixt alkaline falt. Animal fubstances, and mineral bitumens, are resolved in like manner into soot and ashes; with this difference, that the ashes of these yield rarely any alkaline matter, and that they emit in burning a different kind of smell.

On

On submitting the same subjects to a like degree of heat in close vessels, different products are obtained. From most vegetables there arises a watery and acid liquor; a reddish, empyreumatic, acrimonious oil, which swims on its surface; at length, in the utmost degree of sire, a thicker black oil, which sinks to the bottom; and sometimes a little volatile alkaline salt: from animals, a watery and alkaline liquor, a volatile alkaline salt, and oils of a more fetid kind: from bitumens, an acidulous liquor, an oil approaching to the nature of petroleum, and sometimes a concrete subacid salt: a black inspid coal remaining in all cases behind. About the appearance of the first oil, there is commonly extricated a large quantity of air, or elastic vapour, which, if the sire is hastily urged, and no exit allowed it, either bursts the vessels, or blows off the receiver.

5. Calcination of metallic bodies.

The metals (a) called imperfect or destructible, as tin, lead, bismuth, zinc, regulus of antimony, copper, and iron, on being exposed to the joint action of fire and air, gradually lose their metallic form, and change into a friable or powdery calx; which, urged with a stronger heat, either does not melt, or runs into a vitreous mass, not miscible with metals in their entire state any more than earths and earthy glasses. Some emit slames in their calcination: zinc in particular burns strongly and vividly: from whence it is presumed, that an inflammable substance is one of the constituent principles of these metals, and that the loss of their metallic form and qualities in calcination is owing to the avolation of this principle.

The calcination is greatly promoted by the addition of nitre; which, with most of these metals, visibly destagrates, and is by all of them alkalized in the same manner as by charcoal or other instammable substances. The calx is freed from the saline matter by ablution with water: a part of it commonly dissolves in the water along with the alkalized salt, but either separates spontaneously on standing, or may be precipitated by adding any acid.

⁽a) The term femimetal is throughout this work avoided, as being liable to ambiguity. All the pure metallic bodies I have called by the general appellation of metals: fuch of them as want malleability, are, I prefume, as properly distinguished by the epithet brittle or unmalleable, as by a name which may be imagined to imply that one half of their substance is of an unmetallic nature, and which, in effect, has been often understood in this sense, and accordingly applied to ores, vitriols, and metallic recrements.


A little powdered charcoal, or any other inflammable matter not partaking of a mineral acid, added to the destructible metals in fusion, prevents their calcination: and the calces and glasses, melted with the like additions, recover the principle which they had lost, and are revived or reduced; that is, they resume their metallic appearance, and all their former qualities. But, in order to this reduction, as calces in general melt much more difficultly than the metals themselves, and some of them scarce at all when the sire acts on them through the sides of a vessel; an addition of sixt alkaline salt, borax, or susplied glass, is generally requisite, for bringing them into suspin as well as of inflammable substances, for restoring their metalleity.

All the metals disjolve in acids; some in one acid, and others in another: the disjolution, like that of absorbent earths and alkaline salts, is generally accompanied with an effervescence, heat, and discharge of vapours. In this process, the phlogiston or inslammable principle of the imperfect metals, is absorbed or expelled: hence the vapour, which arises during the disjolution in the vitriolic acid of the metals which abound with this principle, is inslammable and truly sulphureous: and hence the metal, precipitated from the acid by alkaline salts, or by other bodies void of inslammable matter, is found to be a true calx, which, like the calces made by fire, cannot be revived without the

introduction of fresh phlogiston.

The perfect metals, gold, filver, and mercury, fuffer no refolution, or diffipation of any of their principles, from any known power. If changed into the appearance of a calx, by fire or by additions, they are recoverable without loss, either by the simple action of a stronger heat, or by the addition of such substances as may barely absorb from them the matter by which their form had been concealed.

All the WEIGHTS and MEASURES mentioned in this book, are those of the London Pharmacopæia:—the troy pound, divided into twelve ounces, the ounce into eight drams, the dram into three scruples, and the scruple into twenty grains:—the wine gallon, divided into eight pints, the pint into sixteen ounces, and the ounce into eight drams or two spoonfuls.



MATERIA MEDICA.

ABIES.

IR: an evergreen coniferous tree; with numerous, narrow, stiff leaves, standing solitary, or unconnected at the bases with one another.

- I. ABIES Pharm. Parif. Abies conis sursum spectantibus sive mas C. B. Pinus picea Linn. The yew-leaved or silver fir; with a white bark, roundish-pointed leaves somewhat cloven at the tips, and short cones standing upwards: the leaves are marked on the lower side with three green lines and two white depressions.
- 2. PICEA: Abies picea Pharm. Parif. Picea major prima five abies rubra C. B. Pinus abies Linn. The common, or red fir, or pitch tree; with a reddish bark; long slender quadrangular sharp-pointed leaves, and long cones hanging downwards.

These trees are natives of the northern climates: the first grows chiefly on dry, mountainous places; the second in lower and moister grounds. In this kingdom, they are rarely found wild, particularly the first fort: Norway, Switzerland, and some parts of Germany, produce both kinds in abundance.

ALL the parts of these trees contain a resinous juice impregnated with essential oil; in smell not disagreeable, in taste bitterish and moderately

moderately pungent: from incisions made in the trunks, one of the finest of the turpentines is obtained. The red fir appears to be the most refinous; the silver fir is the most grateful: of both forts, the cones are more agreeable than the leaves, the young leaves than the old, and these than the wood. The leaves, though evergreen on the tree, lose of their colour on being dried, and change in keeping to a yellow or brown.

Rectified spirit of wine, digested on the fir, dissolves completely its active parts, along with which it takes up also some portion of the infipid gummy or mucilaginous substance: from the fresh or new dried leaves it gains a yellowish green, from the cones and the wood a brownish or yellowish red tincture. The filtered folutions, mixed largely with water, grow milky and throw off greatest part of their refin with its oil, which may thus be obtained in a state nearly approaching to that of turpentine, the gummy substance being retained by the aqueous fluid. On committing the folutions to distillation, the spirit brings over with it a little of the lighter oil of the fir, so as to be fenfibly impregnated with its fmell; leaving behind an extract, different from the refin separated by water and from the native turpentines, in having an admixture of gummy matter, from which they are free.

Water, though it disfolves little or nothing of the pure turpentines, yet, by the mediation of the gummy matter in the fir itself, extracts part of its refin. In distillation with water, a considerable quantity of effential oil arises: the oil drawn from the wood is nearly similar to Oleum tem- the oil of turpentine: that obtained from the fresh cones is superiour, in fubtility and fragrance, to all the oils of the terebinthinate kind usually met with (a). The decoction remaining after the distillation, inspissated to the consistence of an extract, retains the bitterishness, and some share of the pungency of the fir.

plinum verum Germanorum.

> The tops and cones of the fir tree, by virtue of their balfamic juice, tend moderately to warm and strengthen the habit, and promote perspiration and urine, and the natural fecretions in general. Among us, they are used chiefly by the common people, as an ingredient in diet-drinks; in fome parts of Europe, they are prescribed by physicians in decoctions and spirituous tinctures; for nearly the same intentions as the

exotic woods. Frederic Hoffman the elder relates, that in a fcurvy which raged among the Swedish army, during their wars with the Muscovites, a decoction of the leaves and tops of the fir, made in water or ale, was found an effectual remedy and preservative (a). The Augustan Effentia college joins, to the balfam of the fir, the pungent virtues of cochlearia; by bruifing the cones whilft young, tender, and of a red colour, digesting them for two days in four times their quantity of spirit of scurvygrafs, and then preffing out and filtering the tincture, which is, doubtless, as the authors observe, a medicine of great efficacy. A spirit diffilled from the young leaves is faid to be used in some places as a fuccedaneum to Hungary water.

- 3. ABIES CANADENSIS Pharm. Parif. Abies minor pectinatis foliis, virginiana, conis parvis subrotundis Plukenet. Pinus Balsamea Linn. Virginia or Canada fir; with roundish-pointed leaves, sometimes cloven, standing like the teeth of a comb in two rows on each fide the branches, and variegated underneath with a double line of whitish dots.
- 4. BALSAMEA; Abietis taxi foliis species odore balsami gileadensis Balm-of-gilead fir; fo called from the fragrant smell of the leaves when rubbed. The leaves are roundish-pointed, and slightly cloven, nearly like those of the filver fir: the cones are long and pointed, and stand erect.

THESE foreign firs, now naturalized to our own climate, promife to be superiour, for medicinal uses, to the two preceding; their refinous matter being of a finer and more grateful kind. From the Canada fir is extracted, in America, by wounding it during the fummer heats, an elegant balfam, transparent and almost colourless, which is fometimes brought into Europe under the name of balfamum Canadense. The balm-of-gilead fir has a more agreeable fragrance, approaching to that of the celebrated balfam from which it receives its name: a valuable refin exudes from the cones, in confiderable quantity; and refins nearly of the fame kind may be extracted by spirit of wine. both from the cones and from the leaves.

⁽a) Clavis Schraderian. p. 394.

ABROTANUM MAS.

ABROTANUM MAS Pharm. Edinb. Abrotanum mas angustifolium majus C. B. Artemisia Abrotanum Linn. Southernwood:
a plant, with woody brittle branches; numerous greyish green leaves,
divided into slender segments; and small yellow naked discous flowers,
hanging downwards, in clusters, along the sides of the stalks and
branches. It is a native of open mountainous places, in the warmer
climates: with us, it is raised in gardens, from slips or cuttings;
seldom producing seeds, and not often flowers: the leaves fall off in
the winter: the roots and stalks are perennial.

THE leaves and tops of fouthernwood have a strong smell, to many people agreeable; and a nauseous penetrating bitterish taste: they lose a little by drying both of their taste and smell. The flowers are somewhat weaker than the leaves. The cortical part of the younger roots has a light not ungrateful bitterness, with little or nothing of the peculiar slavour of the herb.

Infusions of the leaves made in water are of a brownish hue, in taste and smell not unpleasant; decoctions are darker coloured and very nauseous. In distillation with water, there arises an essential oil, of a bright yellow colour, in smell exactly resembling the plant. This oil distils slowly, and towards the end of the process proves very soul: rectified, or distilled a second time with fresh water, it leaves behind a considerable quantity of an inodorous and almost insipid resinous matter. From sixteen pounds of the fresh leaves and tops were obtained scarcely three drams of oil, which left in rectification above half a dram of resin.

Tinctures of the leaves, made in rectified spirit, are of a deep green colour, and taste strongly of the southernwood: the smell is covered by the spirit. The spirit, distilled off from the siltered tincture, has very little of the slavour of the herb: the remaining extract retains a considerable share of its smell, and resembles it more perfectly in taste than an extract made by water; though it is much less ungrateful than either that extract or the herb in substance. Rectified spirit appears to dissolve the aromatic part more easily, and the nauseous part more difficultly, than watery menstrua.

This bitterish pungent plant has been employed as a moderately stimulating deobstruent, in different cachectic disorders; as an anthelmintic: mintic; and as possessing some degree of an anodyne or antispassmodic virtue depending on the oil or odorous matter. In the present practice, it is scarcely otherwise made use of than for external purposes, as an ingredient in discutient and antiseptic somentations; in which intention it appears to be of no inconsiderable efficacy. It has likewise been recommended in unquents for promoting the growth of hair; a virtue to which it does not appear to have much claim.

ABROTANUM FEMINA.

ABROTANUM FEMINA Phar. Edinb. (a) Abrotanum femina foliis teretibus C. B. Santolina Chamæcy parissus Linn. Lavender-cotton: a bushy shrubby plant, all over hoary; with oblong slender leaves, composed each of four rows of little knobs set along a middle rib; and naked discous yellow flowers standing solitary on the tops of the stalks. It is a native of the southern parts of Europe; slowers, in our gardens, from June to near the end of summer; and holds its leaves all the winter.

This plant is supposed to agree in virtue with the foregoing abrotanum, and to be the most effectual of the two in hysteric cases, and as an anthelmintic. It has been customary among the common people to use, in this last intention, a decoction of the leaves, made in milk; which receives from them a thick consistence, and a strong, though not very disagreeable taste.

On careful examination, the two abrotana appeared to differ very confiderably in quality. The femina is in smell weaker and less agreeable than the mas; in taste, nauseous and acrid, but void of the penetrating bitterness which prevails in the other. Insusions, tinctures, and extracts, prepared from the femina, are more unpleasant than those of the mas, though not bitter. The essential oils of the two plants, and of consequence their distilled waters and spirits, approach nearer in slavour to one another, though not entirely alike.

These differences, doubtless, affect their virtues as internal medicines. Nevertheless, for fomentations, which is the principal use that either of them is applied to in the present practice, they may be looked upon as very nearly alike: hence the college of physicians of London, under the name of abrotanum, allow either the mas or femina to be taken indifferently.

ABSINTHIUM.

WORMWOOD: a perennial plant; with hoary, divided leaves; firm woody stalks, which die in the winter; and small yellow naked discous flowers, hanging downwards, like little buttons, along the sides of the stalks and branches.

1. ABSINTHIUM VULGARE Pharm. Lond. & Edinb. Absinthium ponticum seu romanum officinarum seu dioscoridis C. B. Artemisia absinthium Linn. Common Wormwood; with large leaves, divided into several deeply indented segments, of a whitish green colour above, and whiter underneath, broader than those of any other species of wormwood. It grows wild about dunghills, and on dry waste grounds; and flowers in June or July.

The leaves of wormwood have a strong offensive smell, and an intensely bitter nauseous taste: the flowers seem to be equally bitter, but somewhat less nauseous *(a): the roots are warm and aromatic, without any thing of the bitterness or offensiveness which prevail in the other parts of the plant. The leaves lose a part of their ill smell, on being dried and kept for some time.

Wormwood leaves give out nearly the whole of their smell and taste both to aqueous and to spirituous menstrua. The watery insusions, prepared without heat, are the least ungrateful. The colour of the insusions made in cold water is a pale brown, in warm water a sooty brown, in proof spirit yellowish. Rectified spirit gains from the fresh leaves a beautiful green, from the dry a reddish or brown tincture.

Rectified spirit elevates little from this plant in distillation: water brings over nearly the whole of its smell and flavour. Along with the aqueous sluid, there arises an essential oil, which smells strongly and tastes nauseously of the wormwood, though not bitter. The oil drawn

Oleum absinthii Pharm. Lond. & Edin.

 ⁽a) The Edinburgh college, in their last edition, direct a tincture of the dried flowering tops
of wormwood, in the proportion of fix ounces to a quart of rectified spirit, under the title of
Tinctura Absinthii.

from

from the fresh herb is commonly of a dark green; from the dry, of a deep yellowish brown colour. The quantity of oil varies greatly, according to the foil and feafon, in which the wormwood is produced: in fome years, ten pounds have afforded upwards of two ounces; in others, twenty pounds have yielded little more than one ounce. Geoffroy obferves, that it is in rainy feafons, and moist foils, that it yields the most oil; that in dry years the oil is accompanied with a refinous matter, and proves of a fine green colour; and that in wet ones it is less refinous, and not green (a).

A decoction of wormwood in water, long boiled, and inspissated to the confistence of an extract, loses the distinguishing smell and ill flavour Extractum of the plant, but retains its bitterness almost entire. An extract made absinthii Ph. with rectified spirit contains, along with the bitter, nearly the whole of the nauseous part; water carrying off, in the evaporation, all the oil, in which the offensive flavour resides, while pure spirit elevates very little of it. The watery extract gives out its simple bitterness, not only to water again, but to rectified spirit.

WORMWOOD is a moderately warm stomachic and corroborant: for these intentions, it was formerly in common use, but has now given place to bitters of a less ungrateful kind. The above experiments, however, point out a method of obtaining from this plant a bitter fufficiently elegant, of little or no particular flavour, and this either in a folid form, or in that of a watery or spirituous solution.

The effential oil is fometimes given, in doses of a drop or two, properly diluted by folution in spirit of wine, as a mild antispasmodic. Its more frequent use is as a vermifuge, for which purpose, it is both applied to the belly, and taken in pills made up with crumb of bread: the spirituous extract, however, promises to be, in this intention, preferable to the pure oil; as it contains, along with the oil, all the bitter matter of the wormwood.

This plant very powerfully refifts putrefaction, and hence is made a principal ingredient in antiseptic fomentations.

Boerhaave commends, in tertian agues, a medicated liquor, prepared by grinding about feven grains of the oil of wormwood with a dram of

fugar, and two drams of the alkaline falt extracted from the ashes of wormwood; and afterwards diffolving the compound in fix ounces of the distilled water of the plant. Two hours before the fit is expected, the patient is to bathe his feet and legs in warm water, and then drink half an ounce of the liquor every quarter of an hour till the two hours are expired: by this means, he fays, cases of this kind are generally cured with eafe and fafety, provided there is no scirrhosity or suppuration (a). The medicine is, doubtless, a very serviceable aperient, where obstructions of the viscera prohibit the immediate use of bark, and in fuch obstructions as the imprudent use of astringents has given rise to. Its virtues might be improved by an addition of the bitter watery extract; though the compound, thus laboriously prepared, would not be at all fuperiour to a fimple infusion of the plant, in pure water, impregnated with a due proportion of fixt alkaline falt.

The roots of wormwood, though not hitherto, that I know of, introduced into medicine, promife to be applicable to some useful purposes; being moderately warm and aromatic, of a flavour sufficiently grateful, and remarkably durable in the mouth. Their virtue refides chiefly in the cortical part, the interiour woody matter being nearly infipid. Rectified spirit extracts their flavour, more perfectly than watery liquors. The spirituous tincture is of a reddish brown colour: inspissated,

it yields an extract more grateful than the root in substance.

2. ABSINTHIUM MARITIMUM nostratibus romanum falso dictum Pharm. Lond. Artemifia maritima Linn. Sea-wormwood, falsely called in our markets Roman wormwood; with finely divided leaves, hoary all over. It grows plentifully about our falt marshes, and in several parts on the fea coast.

This species is in taste and smell considerably less unpleasant than the common wormwood; and hence is preferred by the college as an ingredient in some of the distilled waters. Even the effential oil, which contains the whole of its flavour concentrated, is somewhat less ungrateful; and the watery extract, somewhat less bitter, than those of the common wormwood. Its virtues are the same, differing only in degree;

it is less effectual as an antiseptic and anthelmintic, on account of its being weaker; and more eligible as a stomachic, on account of its being less offensive. A conserve of the tops, made by beating them with thrice their weight of fine sugar, is kept in the shops.

Conferva fummitat. abfinth. maritimi Pharm. Lond.

3. ABSINTHIUM MINUS Pharm. Parif. Absinthium ponticum tenuisolium incanum C. B. Artemisia pontica Linn. Roman wormwood; with more numerous, more finely divided, darker coloured leaves, hoary only underneath. This is a foreign species, but as hardy, and as easily raised, as the others: the roots quickly spread, and send up abundance of new plants. Sea wormwood has been often sold for it in the markets, though the difference betwixt the two, above pointed out, is very obvious on sight.

ROMAN WORMWOOD is confiderably less ungrateful than either of the two foregoing: its smell is weaker, and not unpleasant; and its bitterness is mixed with a kind of aromatic flavour, so as scarce to be disagreeable. It appears to be the most eligible of the three as a stomachic and corroborant; in which intention, a conserve of the tops has been greatly recommended, and is undoubtedly an elegant and useful preparation.

- 4. ABSINTHIUM ALPINUM Pharm. Parif. Absinthium alpinum candidum humile C. B. Artemisia glacialis Linn. Mountain wormwood: procumbent, fine leaved, and covered with a glossy silk-like down (a).
- 5. ABSINTHIUM VALESIACUM: Absinthium seriphium montanum candidum C. B. Herba alba Dod. Mountain wormwood of Valais: erect, fine leaved, and covered with a cotton-like down: the leaves are curled about the edges, so as to appear, with their down, pulpy and of an oblong rounded figure (a).

HALLER informs us, that the first of these plants is frequent in stony grounds on the Alps, and the second by the sides of sandy roads in the territory of Valais in Switzerland; that the former is bitterish, aromatic,

of great estimation among the inhabitants of the Alps, the common remedy against the intermitting fevers which often rage there, and for exciting the menstrual discharges, to which the sudden colds of those countries give frequent checks: that the latter has an acrid aromatic smell and taste, without bitterness, and promises, from its sensible qualities, to be a plant of great virtues. They have not yet been introduced into practice in this country.

ACACIA.

ACACIA Pharm. Lond. Acacia vera Ægyptiaca. ACACIA: a sub-astringent gummy substance, usually of a firm consistence, but not very dry; brought from Egypt, in roundish masses, wrapt up in thin bladders, from sour to eight ounces in weight; outwardly of a deep brown colour inclining to black, inwardly of a reddish or yellowish brown; prepared by inspissating, to a due consistence, the juice expressed from the unripe pods of a large prickly tree called by Caspar Bauhine acacia foliis scorpioidis leguminosæ: the Mimosa nilotica of Linnæus.

Acacia has no manifest smell. Applied to the tongue, it quickly softens, and discovers a moderately rough not ungrateful taste, which is followed by a kind of sweetishness. It dissolves totally in water, except the impurities, which, in the specimens I examined, amounted to a considerable quantity. Proof spirit dissolves a part: rectified spirit extracts from it little or nothing. This juice appears therefore to be truly of the gummy kind; and to differ essentially, in its nature and pharmaceutic properties, from the generality both of astringent juices, as hypocistis and terra japonica, and of astringent vegetables in substance, as bistort and tormentil roots, whose styptic matter is extracted by spirit of wine as well as water.

This mild gummy aftringent may be given to advantage in diforders arifing from laxity and acrimony, as habitual diarrhœas, uterine fluors, and catarrhal coughs. It is used by the Egyptians against spittings of blood, in doses of a dram; and employed in collyria for strengthening the eyes, in gargarisms for quinseys, and in glysters for diarrhœas (a).

⁽a) Alpinus, de plant. Ægypt. cap. 4. & de medicina Ægyptior. lib. iv. cap. 14.

Among us, it is scarcely otherwise made use of than as an ingredient in mithridate and theriaca.

ACANTHUS.

ACANTHUS Pharm. Parif. Branca-ursina: Acanthus sativus vel mollis virgilii C. B. Acanthus mollis Linn. Brankursine or Bearsbreech: a plant with large, elegantly sinuated, soft leaves; among which arises a single stalk, bearing a long spike of irregular monopetalous labiated sless-coloured flowers, the upper lip of each of which is wanting, the stamina standing in its place. This plant is a native of moist warm soils in the southern parts of Europe, and cultivated with us in gardens. It is perennial, and slowers in June and July.

THE roots and leaves of brankurfine abound with a foft, infipid, mucilaginous substance; which is readily extracted by coction or infusion in water, and remains entire upon evaporating the liquid. The roots are the most mucilaginous; and the mucilage, obtained from them, is the most viscous and tenacious.

Rectified spirit, digested on the leaves, extracts from them a fine deep green tincture; which, as the editor of the Wirtemberg pharmacopæia observes, is more durable than the green communicated to spirit by other herbs. The menstruum receives no particular taste or slavour from the plant.

Brankursine has long been a stranger to practice in this country. In those places where it is common, it is employed for the same purposes, to which the althea and other mucilaginous vegetables are applied among us.

ACER.

ACER MAJUS: Acer montanum candidum C. B. Acer Pseudo-Platanus Linn. Great Maple, improperly called Sycamore: a large tree, with pentangular serrated leaves; producing small greenish flowers, and a fruit composed of two capsules, including each a whitish seed at the end where they are joined, and spreading at the opposite end into a membranous wing. It is a native of the mountains of Switzerland and Austria, and now common in England.

ALL the parts of the maple contain a fweet faccharine juice; which, exuding on the furface of the leaves, renders them subject to be preyed on by insects. The roots, trunk, or branches, wounded early in the spring, bleed a large quantity of clear liquor; which, in its dilute state, tastes somewhat sweetish; and being inspissated, yields a brown coloured concrete sugar, with a survey matter resembling malesses.

concrete fugar, with a fyrupy matter refembling melaffes.

The juice, unboiled, has been drank as an antifcorbutic. The fugar and melasses, which are said to be less sweet than those extracted from the sugar cane, and their sweetness to be likewise somewhat different in kind (a), are supposed to be more medicinal in disorders of the breast. Considerable quantities of this sugar, made from a species of maple in Canada, are imported for that use into some parts of Europe, particularly France: the samples which I have seen of it are of a brown colour, and of a more grateful sweetness than the common brown sugars.

Saccharum Canadense Ph. Paris.

ACETOSA.

ACETOSA, Lapathum acetosum, Rumex. Sorrel or Sour-dock: a species of dock with acid leaves.

- I. ACETOSA VULGARIS, OXALIS, Pharm. Edinb. Acetofa pratenfis C. B. Rumex acetofa Linn. Common Sorrel; with the leaves shaped like an arrow-head, and very short or no ears at the bottoms.
- 2. ACETOSA ARVENSIS, minor: Acetofa arvensis lanceolata C. B. Rumex Acetosella Linn. Sheeps Sorrel; with arrow-headed leaves, of which those on the stalk have no ears, those from the root long diverging ones.
- 3. ACETOSA rotunda, P. Paris. Acetosa hortensis rotundisolia C. B. Rumex scutatus Linn. Garden or French forrel; with roundish leaves and ears.

THE leaves of these plants are mildly acid, without any smell or particular flavour: the common forrel is the least, the garden fort the most

agreeable. They were all formerly directed as officinals; and occasionally made use of, for abating heat, quenching thirst, and preventing or correcting a tendency to putrefaction, in febrile and scorbutic disorders; but at present are less regarded, other vegetable acids having in good measure supplied their place.

The leaves yield, upon expression, a large proportion of thick, turbid, green-coloured juice: which, on standing till the seces have subsided, becomes clear and reddish, and in taste more gratefully acid than the herbs in substance. This is one of the most elegant preparations of forrel for medicinal use, and may be advantageously joined, in scorbutic cases, to the juices of the acrid herbs: the inhabitants of Greenland, who are very subject to these distempers, are said to employ, with good success, a mixture of sorrel and cochlearia (a).

Greatest part of the acid matter of sorrel may be obtained also in the form of a concrete salt; by inspissating the depurated juice to a due consistence, and setting it to crystallize. This salt is supposed to approach to the nature of tartar: from which, however, it obviously differs, in being more acid, more easily dissoluble in water (b), and much less, if at all, purgative.

The roots of the forrels have a roughish bitterish taste, without any acidity. They have been looked upon as aperient and diuretic; and, in these intentions, have been sometimes made ingredients in watery infusions and decoctions, to which they communicate a reddish hue: the garden forrel gives the lightest, the common wild fort the deepest red. It is observable, that acid liquors, which in general heighten vegetable reds, destroy this red tincture of forrel roots: alkalies change it to a purplish; chalybeate solutions, to a deep green.

The feeds of forrel are very flightly, if at all, astringent, without acidity or bitterness. They had long a place in the shops, as ingredients in some of the old alexeterial compositions, from which they are now deservedly expunged.

ACETUM.

ACETUM Pharm. Lond. & Edinb. VINEGAR: a vegetable acid liquor, produced by fermentation; either directly from fermentable

(a) Bartholinus, Act. Haffniens. 1671. obs. 9. (b) Neumann, Chemical works, p. 257, 424.

juices or infusions; or from such as have been previously fermented into a vinous state. The more spirituous the wine, the more acid is the vinegar.

Vinegar is not a pure or fimple acid, like those of the mineral kingdom; in open vessels, it grows vapid, ropy, and putrid, while the mineral acids remain unchanged. Distilled by a moderate heat, not exceeding that of boiling water, it yields first a phlegmatic liquor, afterwards a slightly acid one, which is succeeded by stronger and stronger acids; till the matter in the distilling vessel becomes thick and unctuous like honey: the vinegar prepared from malt liquors contains more of this viscous substance than that of wine, and hence is more disposed to become ropy and slimy in keeping. This residuum, urged with a stronger fire, gives over an empyreumatic oil, and a penetrating acid spirit tainted with the ill smell and yellow colour of the oil. There now remains a black coal, which, burnt into white ashes, yields a considerable proportion of fixt alkaline salt.

Acetum difiillatum Pb. Lond. & Edin.

Pure fixt alkaline falt, faturated with the colourless distilled liquors, and afterwards exsiccated, contracts a yellowish or brown tinge; and thus betrays, that the acid still retains a portion of the oil. On gently melting the dry salt, the oily matter burns to a black coal, which separates on dissolution in water: the solution exhaled to dryness, leaves a perfectly white neutral salt, containing the pure acetous acid combined with the alkali. On adding to this compound a little oil of vitriol, the acetous acid is disengaged, and may now be collected by distillation, in a highly concentrated state, and of a very pungent volatile smell.

The quantity of fixt alkaline falt, which vinegar is capable of faturating, is one of the furest criterions of its strength. The best of the German vinegars, according to Stahl, saturate little more than one fortieth of their own weight; the French vinegars, examined by Geoffroy, above one thirty-fifth, and some of them no less than one twelfth; the common distilled vinegar of our shops about one twentieth (a). By con-

⁽a) It cannot be affirmed that the ftrengths of the several vinegars examined were exactly in the proportion of the above numbers, as the alkaline salt, used by different persons or at different times, may have differed in purity or dryness, and as the common way of judging of the saturation is too vague for determining the quantity to any degree of nicety. For all trials of this kind, whether with vinegar or other acids, the alkaline salt (that of tartar is the most eligible) should be previously melted, that all remains of watery moisture may be expelled from it; and the saturation should be determined by means of coloured papers, as mentioned at the end of this article.

gelation, and by distillation from alkalies, as above-mentioned, and from some metallic bodies, particularly copper, the acid may be so far concentrated as to saturate near equal its own weight; a greater degree of strength, than even the mineral acid spirit of sea salt can easily be

brought to.

The acetous acid, however purified or concentrated, differs effentially from all the others: - from the native vegetable acids, in fubtility and volatility; not being obtainable in the form of a concrete falt, which most, perhaps all, of the native ones are; and rising in distillation with a moderate heat, which very few of the native ones have been found to do; most of the acid juices giving over, in the heat of boiling water, only their aqueous fluid, and having greatest part of their acidity destroyed by a stronger heat: - from the mineral acids, in its habitude to different bodies, and the nature of the compounds which it forms with them: thus, whatever alkaline, earthy, or metallic fubstance, the acetous acid be combined with, the addition of any mineral acid will disjoin them, the mineral taking the place of the acetous: neutral falts, composed of the acetous acid and fixt alkalies, dissolve, totally and plentifully, in rectified spirit of wine, whilst those, composed of the same alkalies and mineral acids, are not at all foluble in that menftruum: in this property, the acetous acid differs also from most, perhaps from all, of the acids of its own kingdom: —and from all acids in general, in its peculiar odour.

VINEGAR disfolves the elixated ashes of vegetables, at least in great part; animal earths, purified by incineration, or when naturally blended with but little gelatinous matter, as in shells; the earth of alum; and the mineral calcareous earths. - The folubility of calcareous earth in the acetous acid, and its precipitability by that of vitriol, afford a ready method of discovering the sophistication of vinegar, said to be sometimes practifed, with vitriolic acid. If a faturated folution of any calcareous earth, as chalk, made in strong vinegar, be added to such as is suspected of containing vitriolic acid, no change will enfue if the vinegar was pure; but if it contained even a minute portion of that acid, the mixture will immediately become milky, and on standing for a little time deposite a white fediment: if the calcareous folution be gradually dropt in, fo long as it produces any milkiness or cloudiness, all the vitriolic acid will be absorbed by the chalk, and as this new compound is exceeding F 2 fparingly

fparingly diffoluble, nearly the whole of it will precipitate, so as to leave the vinegar almost pure.

It dissolves, among metallic bodies, zinc, iron, copper, tin, lead, bis-muth, and regulus of antimony; the two last in very small quantity, but sufficient to give a strong impregnation to the vinegar. It dissolves lead more easily when reduced into a calx, than in its metallic state: boiled even with the glass of lead, or in the common glazed earthen vessels, in the glazing of which this metal is a principal ingredient, it extracts so much as to become strongly tainted with the pernicious qualities of the lead.

It dissolves the vegetable inspissated juices, and several of the gummy refins, and extracts the virtues of fundry plants in tolerable perfection; but at the same time its acidity makes a notable alteration in them, or superadds a virtue of a different kind. Some drugs, however, for particular purposes, it excellently assists or coincides with, as garlic, squills, ammoniacum: and in many cases, where this acid itself is principally depended on, it may be advantageously impregnated with the flavour of certain vegetables: most of the odoriferous flowers impart to it their fragrance; and the blue, bright red, and some others, tinge it at the same time of a fine purplish or red colour.

It unites, like the mineral acids, with rectified spirit of wine, into what is called a dulcified liquor, provided the vinegar has been highly concentrated. On distilling the mixture with a boiling heat, a large proportion of a subtile sluid is obtained, similar in its general properties to the æther prepared with the other acids (a).

It mingles equally with blood and its ferum, and with most of the fluids of animals; not thickening or coagulating them, like the acids of the mineral kingdom; but tending rather, as Boerhaave justly observes, to attenuate and resolve coagulations. It is likewise, when taken internally, less stimulating than the mineral acids, and less disposed to affect the kidneys *(b).

THIS mild unctuous acid is a medicine of great use in the different kinds of inflammatory and putrid distempers, both internal and external.

⁽a) M. le Compte de Lauraguais, Hist. Acad. Par. 1759.

^{* (}b) It is less liable to undergo changes in the first passages than the native vegetable acids, which have yet to go through the process of fermentation. Cullen-

It is one of the most certain antiphlogistics and sudorifics in high fevers, and one of the best preservatives against pestilential and other putredinous contagions. Fainting, vomiting, lethargic and hysteric paroxysms, are likewise frequently relieved, by vinegar, applied to the mouth and nose, or received into the stomach: lethargic persons are often found to be excited more effectually by vinegar blown into the nose, than by the far more pungent volatile spirits. Boerhaave observes, that this acid counteracts, in a peculiar manner, the effects of spirituous liquors.

The daily use of vinegar, with food, is falutary in hot bilious difpositions, and where there is a tendency to inflammation or putrefaction. It is prejudicial to children, to aged, hysterical, and hypochondriacal perfons, in cold pale phlegmatic habits, where the veffels are lax, the circulation languid, and the power of digestion weak. It tends in all cases, if used freely, to prevent corpulence; Hoffman (a) suspects that it produces this effect by impeding the formation of chyle, or deftroying the union of the unctuous and ferous fluids of which chyle is composed; an effect common to all acids, as appears from their coagulating milk and artificial emulfions. I have known great corpulence reduced by the liberal use of vinegar, which is the acid commonly employed for this purpose, but not with impunity; diseases succeeding, which eluded the power of medicines, and proved at length fatal.

Combinations of vinegar with different earthy bodies differ in virtue according to the nature of the earth. A folution of the aluminous earth in this acid is strongly styptic; of vegetable earths, or magnesia alba, bitterish and gently purgative: both these solutions are milder, and less ungrateful, than those of the same earths made in the mineral acids, and though as yet unknown in practice, certainly deferve to be introduced. Solutions of different animal and the calcareous mineral earths are bitterish and fubauftere, in various degrees; and supposed to act as mild resolvents,

fubaftringents, or diaphoretics.

Combinations of vinegar with fixt alkaline falts, are useful aperients, diuretics, and cathartics. I have known two drams of the alkali, diffolved in as much vinegar as was fufficient to faturate it, occasion ten or twelve copious watery stools, and a plentiful discharge of urine, without griping or fatiguing the patient. Mixtures of alkali and distilled vinegar,

* Tartarus regeneratus Phar. Edinb. + Sal diureticus Phar. foliata tartari, & arcanum tartari, vulgo.

Spiritus Mindereri

Pb. Edinb.

evaporated to a dry falt, are kept in the shops; either in a brownish oily state, as obtained by fimple evaporation*; or purified to perfect whiteness, by gentle fusion and folution in water +: these preparations are given in doses of ten or twenty grains as mild aperients, and to a dram or two as Lond. Terra purgatives and diuretics.

> It is difficult, in the common way of managing the process, to hit the exact point of faturation between the acid and the alkali. After fourteen parts of distilled vinegar have been gradually poured upon one part of the falt, the addition of a little more of the acid will occasion no further effervescence while the mixture is cold; but if well heated and stirred, the effervescence begins again, and continues till four or five parts of fresh acid have been added: on exhaling the aqueous fluid, the remaining dry falt will generally still raise an effervescence with fresh vinegar, and require two or three parts more of the acid to render-it completely neutral. There is, therefore, this advantage, in reducing the falt to a dry form, that the perfect neutralization is obtained with greater certainty than when the ingredients are barely mixed together. The purification of the dry falt, or separation of its oil, is intended to render it fitter for weak stomachs, on which it would not fit so easily in its common impure state; though the medicine, thus purified, is in some particular cases less to be depended upon than the oily falt. It may be observed, that the imperfection of the oily falt, which the purification is defigned to remedy, does not depend upon the oil as fuch, but on its receiving some degree of burnt taint from the too strong heat commonly employed in the evaporation, and may therefore be effectually prevented by the prudent use of a water-bath.

> Combinations of vinegar with volatile alkaline falts, commonly made with distilled vinegar added gradually to the falt till the effervescence ceases, scarcely yield any solid salt, the saline matter evaporating with the watery fluid, or even before it: on distilling the mixture in a retort, a falt fometimes concretes about the fides of the receiver, but liquefies again as the veffels grow cold. These mixtures have little purgative virtue, but operate powerfully as aperients; by urine, if the patient walks about in the cool air; by perspiration or sweat, if kept warm in bed. They are principally made use of in this last intention, in doses of half an ounce: and as they act without irritation, they have place in inflammatory cases, where the warm sudorifics, if they fail of exciting a fweat, aggravate the diftemper.

> > Great

Great care ought to be taken in the neutralization of this liquor, which is very difficult to be hit exactly by the common method. The best way of judging of the saturation is, by trying the liquor from time to time with certain coloured vegetables juices, or on paper stained with them. A thick writing paper may be stained pale blue on one side with the blue preparation of archil commonly called lacmus; and pale red on the other fide by a mixture of the same infusion with so much dilute spirit of salt as is just sufficient to redden it. If a small slip of this paper be dipt occasionally into the liquor to be tried, or a drop of the liquor applied upon both fides of the paper; the red fide turns blue fo long as any of the alkali remains unfaturated; the blue fide turns red when the acid begins to prevail; and no change at all is produced when the faturation is complete. This way of trial I strongly recommend to the apothecary in making all neutral mixtures, as he may thus at all times find, expeditiously and with certainty, the exact point of neutralization, which is not perhaps possible to be found by the common way of judging from the effervescence: how precarious and indeterminate a mark the ceffation of effervescence is, is apparent from the observations on the preceding preparation. Where lacmus cannot be procured, the paper may be coloured with the juices of certain blue flowers, as violets, iris, cyanus, &c. or with the blue juice pressed out from scrapings of the cortical part of common radish roots: with these juices it is sufficient to stain the paper on one fide, this one colour discovering both acidity and alkalescence, the former changing it red, and the latter green; but the change produced by liquors flightly alkaline is much less conspicuous than that produced by the fame liquors on the red paper above-mentioned, which is therefore to be preferred.

ADIANTHUM.

MAIDENHAIR: an evergreen plant; with flender, smooth, shining blackish stalks; producing no manifest flower. The seeds are a fine dust, lying in roundish specks, about the edges of the backs of the leaves, which curl over and cover them.

1. ADIANTUM Pharm. Parif. Adianthum verum. Capillus veneris:
Adianthum folio coriandri C. B. Adiantum capillus veneris Linn. True
maidenhair:

maidenhair: about half a foot high, with several pinnæ of little roundish sinuated or nearly triangular leaves towards the tops of the stalks.

2. ADIANTUM CANADENSE Pharm. Paris. Adianthum fruticosum brazilianum C. B. Adiantum pedatum Linn. Canada maidenhair: larger, with spreading branches.

The first fort grows wild in Italy and the southern parts of France, from whence the dry leaves are sometimes brought to us. The second, a native of America, is cultivated in some of our gardens.

The leaves of both the maidenhairs have a flight sweetish roughish taste, and a pleasant but weak smell, very perceptible when boiling water is poured on them. They readily give out to the water the whole of their smell, taste, and medicinal virtue: the insussions are not ungrateful; particularly that of the Canada fort, whose slavour is both pleasanter and stronger than that of the other. Insussions or decoctions of them, inspissated, yield a moderately rough, bitterish, mucilaginous extract. Rectified spirit of wine takes up their taste and slavour, and gains from them a deep green colour, but dissolves little of the mucilaginous substance, in which a considerable part of their virtue consists: the extract, obtained by inspissating the tincture, is less in quantity, and stronger in taste, than that made with water.

Maidenhair has long been held in esteem against disorders of the breast; for promoting expectoration, softening recent coughs, and allaying the tickling in the throat occasioned by defluctions of thin rheum. For these purposes, a syrup of the true fort, slavoured with a little orange-flower water, has been usually brought from France; and a syrup of the Canada sort, made with maple sugar, is sometimes received from America. The virtue of the maidenhair is obtainable, however, to much better advantage, by drinking an insusion of the herb as tea, sweetened either with sugar, or by the addition of a little liquorice. The English maidenhair has been commonly substituted in the pectoral syrups and insusions made among us: the Canada species, which appears to be superiour to both, is said to have been long made use of in France, and has lately been introduced into practice in this country.

* AER FIXUS.

IT is not here intended to give a chemical account of the different kinds of air. It is well known to all, in any degree conversant with the subject, that besides the air which we breathe, there have been discovered other species of permanently elastic invisible sluids, possessing very different properties. Of these, one of the best known, and the only one as yet applied to medical purposes, is the Fixed, Fixable, or Mephitic Air, or Gas.

This kind of air is naturally contained in a great variety of substances; and is set free in many processes of art. Every kind of vinous fermentation extricates a large quantity of it, which floats on the surface of the fermenting liquor. It is expelled from alkaline salts, and from absorbent earths, by the action of fire, and of acids. The properties by which it is principally distinguished, are, its extinguishing slame, being destructive to animal life when inspired pure and in large quantity, and being readily absorbed by water, to which it gives a slightly acid taste, accompanied with a degree of briskness and spirit. It has the general properties of a weak acid, of a very volatile nature.

The idea of its medical virtues feems to have arisen chiefly from some experiments of its great antiseptic power when applied pure; and from the discovery of its presence in large proportion in some of the most celebrated mineral waters, and also in other substances very efficacious in

the cure of particular diseases.

At present, fixed air is considerably employed in medicine, chiefly in

the following ways.

The celebrated antiemetic mixture of Riverius, composed of a spoonful of lemon juice and a scruple of salt of wormwood taken in the act of effervescence, is supposed to owe its efficacy chiefly to this principle, which is set free during the combination. On this soundation, its use is extended to many diseases in which a tendency to putridity is suspected. Any other acid may be substituted with equal advantage as far as the fixed air is concerned.

By various contrivances, the air expelled from an absorbent earth by the addition of an acid is received into water, in which it dissolves, communicating to the liquor the qualities above-mentioned. This becomes a very grateful beverage in fevers and putrescent diseases, being cooling, antiseptic, and at the same time causing a temporary glow in the stomach. It should frequently be made fresh, and kept well corked, as the aerial spirit very readily slies off again from the water. As fixed air is in reality an acid, water impregnated with it may be neutralized by an alkali, thus forming an agreeable neutral julep, which is found to be an useful refrigerant and diuretic. If a few clean iron filings are thrown into water to be impregnated with fixed air, so much of the iron is dissolved by its means, as to produce an artificial chalybeate, possibly not inferior in virtue to the most celebrated natural ones.

As the antiseptic powers of fixed air seem to be most considerable when a stream of the pure air itself is thrown upon the matter to be sweetened, some practitioners have injected bladders full of it by way of clyster where the primæ viæ were loaded with highly putrid sæces. On this principle, too, patients with ulcerated lungs have been directed to respire fixed air as it rose from an effervescing mixture; and, notwithstanding the instantly satal effects of breathing this air absolutely pure, it was found to be perfectly innoxious when thus passing through a body of atmospherical air. The same vapour has also been received upon the naked surface of cancerous and other putrid ulcers, with a view to sweeten and correct the discharge.

In these methods has this substance been directly applied to medical purposes. But its application has by some been supposed to be indirectly much more extensive, as they have attributed to it the efficacy of various other remedies. Fermented and fermentable liquors, fruits and fresh vegetables, have been thought useful in putrescent habits and diseases chiefly as affording a large supply of this principle, which might be absorbed by the stomach and intestines. The use of Wort in the Sea Scurvy (see the article Frumentum) was professedly suggested by Dr. Macbride on this supposition.

From what has been above faid, the experienced reader will eafily perceive in what classes of diseases this medicine may be expected to be serviceable. There is one disorder, however, in which its proposed use may not appear so deducible from its obvious qualities: this is, the stone in the kidneys and bladder. It is well known that the medicines from which most of late years has been expected in these cases, are limewater and caustic alkali, the direct chemical opposites to fixed air. But

as the nature of urinary calculi is acknowledged to be very various, it is not unreasonable to propose opposite solvents for them. That calcareous earth may be rendered soluble in water by the medium of fixed air, is not to be doubted; and there are calculi which are certainly calcareous. From indisputable experiments it appears, that pieces of calculi have undergone a solution out of the body in water impregnated with fixed air; but it can scarcely be hoped that the menstruum will reach the bladder in so concentrated a state. Cases have, however, been published, in which manifest relief was obtained from the free use of a beverage of water saturated with fixed air; and it has this advantage above alkaline solvents, that substances abounding in fixed air are more friendly to the general health than alkalescent ones. It has been proposed to throw pure fixed air directly into the bladder by a suitable apparatus; but no experiment of this kind has been made public.

The reader who wishes for further information concerning the medical uses of fixed air, may consult the ingenious Commentary on the subject by Dr. Dobson.

ÆRUGO.

ERUGO Pharm. Lond. & Edinb. Erugo vel viride æris Ph. Parif. Verderis: copper corroded by a fermented vegetable acid into a bluish green substance. The greatest quantities are prepared about Montpelier, by stratifying copper plates with grape stalks, that have been previously soaked in strong wine, and exposed to its vapour during a second fermentation of the wine continued to an acetous state (a): the subtile acid, with which the stalks are thus impregnated, corrodes the surface of the plates, in a sew days, into verdegris; which is afterwards scraped off, moistened, and packed up in skins. The masses, as brought to us, have generally some grape stalks intermixed: these may be separated by pulverization, as being less pulverable than the verdegris itself. The goodness of verdegris is judged of from the deepness and brightness of its colour, its dryness, and its forming, when rubbed on the hand with a little water or saliva, a smooth passe free from grittiness.

This concrete is partially diffoluble in water and in rectified spirit, and almost totally in vinegar: from the acetous solution, well saturated,

Distilled verdegris so called and left to exhale flowly in a warm air, greatest part of the verdegris may be recovered in a crystalline form. The crystals, distilled with a suitable fire, in a retort or other like vessel, give over the acetous acid, in a highly concentrated state, but somewhat altered by the process.

Verdegris is employed externally for deterging foul ulcers, and as an escharotic. Hossman (a) recommends it particularly for destroying the callofities of old fiftulæ: tents of powdered verdegris, made up with faliva, or other liquids not fat or oily, confume, he fays, the hardest callus, in three or four days, fo as to render it completely separable. A detergent ointment is commonly prepared, by gently boiling five parts of verdegris in fine powder, with fixteen of honey and feven of vinegar, till reduced to a due confistence. On keeping this mixture for some time, a thick matter, containing greatest part of the verdegris, falls to the bottom, and a thinner floats on the top: this last is the part made use of, unless where particular occasions require it to be rendered more acrid by shaking up the thick among it. In the last Edinburgh difpensatory, an ointment is directed, composed of white wax and refin each two ounces, olive oil one pint, and verdigris half an ounce. When thefe kind of applications are employed for venereal or other ulcerations in the mouth and tonfils, great caution is requifite, on the part of the patient, to guard against any portion of them passing into the stomach; an accident which is faid to have fometimes happened, particularly in childrens cases, and to have produced very dangerous and even fatal confequences.

Mel Ægyptiacum Pb. Lond.

Unguentum ex Ærugine Ph. Edin.

> Verdegris is rarely or never given internally. Some recommend it, indeed, in the dose of a grain or two, as an emetic, which operates almost as soon as received into the stomach, and which may therefore be of use, where poisonous substances have been taken, to procure their immediate rejection. It appears, however, highly imprudent, to have recourse, on such occasions, to a remedy in itself so dangerous and so virulent; and more especially as a speedy evacuation may generally be obtained, by means of substances, which are not only innocent, but at the same time weaken the force of the poison by diluting and obtunding it; as warm water, milk, oils.

AGARICUS.

AGARIC: a fungus, growing on the trunks of trees, without any pedicle; internally of a simple and uniform structure throughout its whole substance.

(a) Med. rational. De ulceribus.

1. AGARICUS Pharm. Lond. Agaricus sive sungus laricis C. B. Agaric: covered with a brown bark, full of small holes underneath;

internally white.

This fungus is met with on old larch trees, in the Levant, and in different parts of Europe: that produced in the Levant is accounted the best, but from what particular place or country the shops receive it, is not very clear.—It comes forth on the tree in the beginning of spring, and continues to increase till autumn: at this time, it is cut off, the cortical part separated, and the internal part exposed for some weeks to the sun, by which its whiteness is improved. It is brought into the shops in irregular pieces, of different magnitudes, of a chalky whiteness, and very light: the best is easily cut with a knife, friable betwixt the singers, and has no hard, or gritty, or coloured veins.

AGARIC has no remarkable smell: chewed, it impresses first a considerable sweetness, which is followed by a nauseous acrimony and bitterness. It is difficultly reduced to a fine powder in a mortar, on account of its fungous texture: it may be rendered more easily pulverable, by moistening it with a solution of gum tragacanth, and afterwards thoroughly drying it.

It gives out little of its active matter to aqueous menstrua: after long boiling in water, it retains great part of its taste, and proves remarkably viscid and tenacious. The decoction has little taste or colour: inspissated,

it leaves a fmall quantity of a brown coloured nauseous extract.

Rectified spirit takes up nearly the whole of the active matter, leaving the agaric almost insipid. The tincture is of a fine yellow colour, and of an unpleasant sweetness, which continues long in the mouth, and in good measure covers the heat and pungency of the spirit. The extract, remaining on distilling off the spirit, discovers less of the sweet, and more of the offensive bitterness of the agaric. Proof spirit has nearly the same effect as the rectified.

These experiments were made on the internal substance of agaric, as commonly met with in the shops. The cortical part seems to be of a different quality: Mr. Boulduc relates, that a spirituous tincture, drawn from this, had such an abominable taste, that a single drop, laid on the tongue, occasioned vomiting, and a loathing of food for a whole

whole day(a). This fungus appears to differ also greatly in quality, at different periods of its growth: Bellonius informs us, that when full of juice, before it has come to maturity, its offensive effluvia are apt to excite violent symptoms in those who incautiously cut it from the tree (b).

Agaric, taken from a scruple to two drams or more, is said to act weakly, though not very mildly, as a cathartic. It was formerly held in confiderable esteem, and supposed to evacuate peccant humours from the remote parts of the body: but the great flowness of its operation, from which alone that quality appears to have been deduced, its occasioning little evacuation, and being commonly productive of nauseæ, sickness, and gripes; have brought it now deservedly into difuse. Gummy or mucilaginous substances, with which it was formerly made into troches and pills, in some degree correct its ill qualities: aromatics are, in this intention, of very little use. Extracts made from it with vinegar, with wine, and with water in which a little fixt alkaline falt has been dissolved, are faid to purge more effectually, and with less inconvenience, than the agaric in substance; though even these preparations do not appear to be equal to the more common and experienced cathartics. The antients supposed it to be possessed of alexiterial powers, and in consequence of this imaginary virtue made it an ingredient in the theriaca, which is the only officinal composition wherein it is now retained.

2. AGARICUS QUERCINUS, fungus igniarius: Agaricus pedis equini facie Tourn. Fungi arborei ad ellychnia J. B. Boletus igniarius Linn. Agaric of the oak, called by fome, from its readily catching fire, touchwood or fpunk: growing in form of a horse's hoof; externally of an ash colour, internally dusky coloured, soft and tough. Though denominated from the oak, on which the best fort is supposed to be produced, the same sungus is found on several other kinds of old trees, throughout Europe.

THE agaric of the oak has lately come into esteem as an external styptic. It has been said to prevent hæmorrhages after amputations, as

(b) Bellonius, de arborib. conif. &c. p. 26.

⁽a) Boulduc, Mem. de l'acad. roy. de scienc. de Paris, pour l'ann. 1714.

effectually as the painful operation by the needle; and to restrain bleedings in wounds, of several days or weeks standing, where the parts are become so rotten as to become incapable of bearing ligatures. For these purposes, the internal soft part of the sungus, divided into pieces of different sizes, and beaten with a hammer till it may be easily torn with the singers, is applied to the orisices of the vessels, with the usual dressings over it. In a short time the extremities of the vessels are said to be found contracted into a conical shape, and the orisices stopt with plugs of coagulated blood, sufficient to resist the force of the circulation.

Cases have been published, in which this application seemed to answer the character given it (a): in others, and those not a few, it proved ineffectual. Some have remarked, that where it seemed to succeed, the subjects were brought so low before the operation, that little danger was to be apprehended from a hamorrhagy, though no other application

had been made than that of dry lint and flour (b).

Thus much is certain, that the agaric has already lost greatly of its repute, both in France, where it was first introduced, and in England; and that it does not appear, from its sensible qualities, to be possessed of any truly styptic power, at least in any considerable degree. Chewed, in substance, it discovers no taste: boiled in fresh parcels of water, it yielded about one fourteenth its weight of extract, which had only a weak sweetish taste, mixed with a kind of bitterness: treated in the same manner with rectified spirit, it yielded about one eighth its weight of an extract, which had less taste than the other.

It is probable that this fungus acts no otherwise than as a pliable soft substance, adhering to the orifices of the vessels, till they have contracted spontaneously. Some other suggi were employed formerly in the same intention, and there are late accounts, in the Philosophical Transactions, of two having been used with success; namely, the *lycoperdon*, or dusty mushroom; and that found on the casks and walls of wine-vaults, and and thence called suggestions.

AGNUS.

AGNUS CASTUS, Vitex, Pharm. Parif. Agnus folio non serrato J. B. Vitex Agnus Castus Linn. AGNUS-CASTUS, or chaste-tree:

(a) Warner, Cafes in Surgery, &c.

(b) Neale, Observations on the use of agaric, &c.

a small tree or shrub, with tough branches, digitated narrow leaves, and monopetalous purplish flowers standing in spikes on the tops of the branches, followed by oblong whitish seeds. It is a native of the warmer climates, and cultivated in some of our gardens.

The feeds of agnus were formerly celebrated as antaphrodifiacs; but experience does not discover in them any degree of such virtue, and some have ascribed to them an opposite one. From their sensible qualities, their virtues, of whatever kind, do not appear to be very considerable. The seeds in substance, as met with in the shops, have little taste, and scarcely any smell, though described by authors as very hot and biting: extracts made from them, by water or spirit, are weakly bitterish and somewhat pungent. They seem to abound chiesly with a gross insipid oil, of the expressible kind; which is in part taken up by rectified spirit, and separates and falls to the bottom during the inspissation of the tincture: the oily matter is of a deep saffron colour, the inspissated extract somewhat paler.

AGRIMONIA.

AGRIMONIA Pharm. Edinb (a). Eupatorium verum sive agrimoni C. B. Agrimonia Eupatoria Linn. AGRIMONY: a hairy plant; with winged leaves, composed of oblong indented segments, with smaller portions between, set on middle ribs which stand alternately on the stalk: on the top grows a long spike of pentapetalous yellow slowers, followed by little burs, containing, each, one or two seeds. It is perennial, grows wild in hedges and about the sides of fields, and flowers in May.

THE leaves of agrimony have a flightly bitterish roughish taste, accompanied with an agreeable, though very weak, aromatic slavour: the flowers are in smell stronger and more agreeable than the leaves, and in taste somewhat weaker. They readily give out their virtues both to water and to rectified spirit: the leaves impart to the former a greenish yellow, to the latter a deep green colour: the flowers yield their own deep yellow tincture to both menstrua. In distillation with water, there

arises a very small portion of a yellowish essential oil, which smells strongly and agreeably of the herb.

Agrimony is one of the milder corroborants; and in this intention is fometimes employed, especially among the common people, against habitual diarrheas, and cachectic, and other indispositions from a lax state of the solids: insusions of the leaves, which are not ungrateful, may be drank as tea. It is sometimes joined with other ingredients in diet drinks for purifying the blood; and in pectoral apozems.

This plant is often raised in gardens; and does not seem to receive, from culture, any material change in its quality. Another species or variety, of foreign original, common also in our gardens, and differing little in appearance from our indigenous agrimony, promises to be superiour to it in virtue; as its taste is more aromatic, and its smell much stronger and very agreeable: Caspar Bauhine calls it eupatorium odoratum, Fabius Columna eupatorium dioscoridis, odoratum & aromaticum.

ALCHIMILLA.

ALCHIMILLA Pharm. Edinb. (a) Pes leonis five alchimilla J. B. Alchimilla vulgaris Linn. Ladies mantle: an herb, with undivided plaited multangular leaves, and imperfect flowers standing in form of umbels on the tops of the stalks: the cup consists of four larger and four smaller leaves placed alternately. It is perennial, grows wild in dry fields and meadows, and flowers from May to August.

The leaves of alchimilla are weakly aftringent, without any remarkable smell or flavour. They have been recommended, internally, against alvine and uterine fluxes, in which they may doubtless be of some service; and externally, against certain relaxations, which can yield but little to astringents of so mild a kind. Their styptic matter is extracted both by water and by spirit, and when separated from the sluids by inspissation, is still found to be weak: the spirituous extract is the strongest, this menstruum dissolving less, than water does, of the insipid mucilaginous substance of the leaf. The roots of the plant are more astringent than

the leaves, and the extracts made from them, are proportionably stronger. They both strike a black colour with solutions of chalybeate vitriol.

ALKEKENGI.

ALKEKENGI, Halicacabum, Pharm. Edinb. (a) Solanum veficarium C. B. Physalis Alkekengi Linn. Winter-cherry: a low, somewhat hairy plant; with unbranched stalks; large heart-shaped acuminated leaves, standing in pairs at the joints; and whitish bell-shaped flowers, rising in the bosoms of the leaves, divided about the edges into sive segments: the flower-cup changes into a pentagonal capsule or bladder, which, bursting, discovers in its bottom a red fruit like a cherry, containing numerous small seeds with a juicy pulp.—It grows wild in some parts of Europe, and spreads so much in our gardens as not to be easily extirpated. The fruit ripens about the beginning of October, and sometimes continues to near the end of December; after which, the plant dies to the ground.

WINTER-CHERRIES have an acidulous not unpleasant taste, mixed with, or followed by, a slight bitterness: the covering, in which they are inclosed, has a strong pungent bitterness, with which it is apt to impregnate the cherries, unless some care is taken in gathering them. As medical writers in general speak of this fruit as being very bitter, we may presume that it has been often used with this extraneous

bitter impregnation.

These cherries are accounted powerful diuretics, operating without heat or irritation, and which may therefore be ventured on in inflammatory distempers: five or six of the cherries in substance, or an ounce of the expressed juice, are directed for a dose. They are said to be, in some places, eaten, among the common people, by handfuls (b), and with good success, against suppressions of urine, and for promoting the expulsion of mucus and gravel. Mr. Ray tells us of a gouty person, who was cured, and kept free from returns of his disorder, by taking eight of these cherries at each change of the moon; and that the operation of the medicine procured a discharge of extremely fetid matter by

urine(a). The cherries may be dried so as to be pulverable, or the depurated juice inspissated with a gentle heat to the consistence of a rob or extract, and in this state preserved for use.

ALLIARIA.

ALLIARIA Pharm. Edinb. (b) C. B. Erysimum alliaria Linn. SAUCE-ALONE, or jack-by-the-hedge: a plant, with roundish, or heart-shaped, slightly indented leaves; and firm upright stalks; on the tops of which, and in the bosoms of the leaves, come forth clusters of tetrapetalous white flowers; followed by oblong bivalvous pods, full of black feeds. It is biennial, grows wild in hedges, and flowers in May.

THE leaves of alliaria have a moderate acrimony, and a strong flavour greatly refembling that of garlic or onions: they give the fame kind of durable taint to the breath, as those roots; and have been used for the fame culinary purposes. They lose greatest part of their smell, and a little of their tafte, on being moderately dried: after keeping for fome months, the tafte, as well as fmell, feemed to be wholly loft: the leaves, on being chewed, proving merely mucilaginous. The juice, expressed from the fresh leaves, is strongly impregnated with their active matter, but loses greatest part of it on being inspissated to an extract with the gentlest warmth: in its liquid state, duly secured from the air, it may be kept uninjured for many months. On distilling the fresh herb with water, there arises a small portion of essential oil, which tastes and smells exceeding strongly. Great part of the virtue of the plant arises also in evaporation with rectified spirit; an extract, made by this menstruum, having little taste or smell, though the tincture, before the inspissation, has a strong flavour of the alliaria. This herb appears, therefore, to differ from garlic, and agree with onions, in the volatility of its active principles.

Alliaria, taken internally in any confiderable quantity, frequently excites a fweat, which is impregnated with its fmell (c): it stands recommended as a very powerful diaphoretic, and diuretic, as a deobstruent in asthmatic disorders, and externally as an antiseptic, in gangrenes and

putrid ulcers. Boerhaave informs us, that he cured a gangrene of the leg, arifing from a neglected fracture and contusion, by applying the leaves of alliaria bruised with wine (a).

ALLIUM.

ALLIUM Pharm. Lond. & Edinb. Allium fativum C. B. & Linn. Garlic: a plant with long narrow grafs-like leaves; among which arises a single straight hollow stalk; bearing on the top a cluster of small white hexapetalous flowers; each of which is followed by a fruit about the size of a pea, full of dark coloured roundish seeds. The roots are of the bulbous kind, of an irregularly roundish shape, with several sibres at the bottom: each root is composed of a number of smaller bulbs, called cloves of garlic, inclosed in one common membranous coat.—It is said to grow wild in Sicily: with us, it is raised in gardens, from seed, for culinary as well as medicinal uses.

THE roots of garlic have a penetrating highly acrimonious taste, and a strong, offensive disfusive smell. Applied to the skin, they inflame and often vesicate the part. Taken internally, they seem to extend their action, in a short time, through the whole habit; impregnating, with their strong scent, not only the breath, but the urine, the milk of the breast or of the udder, the serum which oozes from sores or issues (b), and the sluid which perspires through the skin. The other parts of the plant possess the same qualities, in a lower degree. In Spain, garlic is said to be equally mild with onion, and is used as food.

Garlic root has been celebrated, by some practical writers, in a variety of disorders; and condemned by others, not only as an offensive, but as a noxious drug. It is certain, that there are many cases, in which it is extremely prejudicial; but that there are many also in which it is of great utility. To warm and stimulate the solids, attenuate thick humours, and resist putrefaction, seem to be its primary virtues. Hence, in hot bilious constitutions, where there is already a degree of irritation, where the juices are thin and acrimonious, or the viscera or intestines

⁽a) Boerhaave, bift. plant. Lugd. Bat. 437.

⁽b) Bennet, [Chrift. Benedictus] tabidorum theatr. exerc. 29. p. 81.

unfound, it is apparently improper, and feldom fails to produce headachs, flatulencies, thirst, febrile heats, and inflammatory symptoms in various shapes. In cold sluggish phlegmatic habits, on the other hand, it proves a falutary and powerful corroborant, expectorant, diuretic, and, if the patient is kept warm, sudorific. In loss of appetite, and humoral asthmas, where the stomach or lungs are oppressed by viscid phlegm, this medicine has generally good effects. It has likewise been found serviceable, as a warm strengthener, in the beginning of dropsies, and for preventing a new accumulation of water after evacuation: Sydenham (a) relates, that he has known the dropsy cured by the use of garlic alone.

Some have held it in great esteem as an antidote against the contagion of pestilential and other putrid disorders; whence it received the name of theriaca rusticorum. It is used also among the common people, slightly boiled in milk, as an anthelmintic; and Hoffman looks upon it as one of the capital medicines of that class.

Garlic is sometimes employed externally, in unguents and lotions, as an antiseptic and discutient; and is frequently made an ingredient in the stimulating epithems, applied to the soles of the feet, in the low stage of acute distempers, for raising the pulse and relieving the head. Sydenham assures us (b), that none of the stimulants operate, in this intention, more powerfully than garlic: he observes, that it sometimes occasions intolerable pain, which may be relieved by a cataplasm of bread and milk. Dr. Cullen remarks that it is not so apt to ulcerate the part as mustard; more capable of being absorbed, and extending its action to remote parts.

This root loses in drying almost nine parts in fifteen of its weight, without suffering any considerable loss of its taste or smell: hence six grains, dry, may be looked upon as equivalent to sifteen grains of the fresh root.

The fresh root yields, upon expression, about one fourth its quantity of a very viscid glutinous juice; which smells strongly of the garlic, and in good measure retains its scent after being inspissated, by a gentle warmth, to the consistence of an extract.

Both the fresh and the dry root give out their virtue to water by warm infusion. A quart of water, poured boiling hot upon a pound of the

Syrupus ex allio Pharm. Lond. fresh root cut in slices, and suffered to stand upon it in a close vessel for twelve hours, becomes strongly impregnated with the smell and taste of the garlic. This infusion, with a proper quantity of sugar, makes the syrup of garlic of the shops.

Vinegar and honey excellently coincide with and improve this medicine, as a detergent and deobstruent, in disorders of the breast. A composition of this kind is prepared by insusing an ounce and a half of the fresh root in half a pint of vinegar, and dissolving in the strained liquor, by the heat of a water-bath, ten ounces of clarified honey: to cover in some degree the ill smell of the garlic, a little carraway and sweet sensel seeds bruised, two drams of each, are boiled for a short time in the vinegar before the garlic is put in.

Oyxmel ex allio Pharm. Lond.

The garlic itself is never to be boiled, either with vinegar or with watery liquors; the virtues of this root residing in an essential oil, which exhales along with the steam of boiling water, leaving, if the decoction be inspissated, an inert mucilaginous extract, which has very little of the taste and nothing of the smell of the garlic. The oil, obtainable by distillation, is of a pale yellowish colour and a thick ropy consistence, in small quantity but of great activity, of an extremely strong smell and a fiery taste: great part of the oil remains dissolved in the distilled water, which is very strongly impregnated with the taste and scent of the garlic.

Rectified spirit of wine, digested on dry garlic root, extracts its virtues more readily and more perfectly than either water or vinegar. With this menstruum, the active matter of the garlic does not easily exhale: the spirit distilled off from the filtered tincture has very little taste or smell of the root, nearly all its virtue remaining in the inspissated extract.

ALNUS.

ALNUS Pharm. Parif. Alnus rotundifolia glutinosa viridis C. B. Betula Alnus Linn. ALDER: a tall coniferous tree, common in watery places; with very brittle branches; roundish, crenated, clammy leaves; a rugged blackish brown bark; and reddish wood.

ALL the parts of the alder tree are more or less astringent and bitter.

The bark is a strong styptic, and might, doubtless, be applied to the

fame purposes as the other substances of that class, though at present rarely or never made use of medicinally, unless sometimes among the

common people in fomentations and epithems.

Tournefort reports (a), that in the Alpine countries, it is customary to procure a plentiful sweat, by covering the patient all over with bags of alder leaves heated; and that by repetitions of this operation, rheumatisms, and sciaticas, are commonly cured. In this way of application, it is apparently the heat and moisture, and not any peculiar quality of the alder leaves, that is the medicine. * Dr. Murray, of Gottingen, recommends from his own experience, the leaves of alder chopt and heated over the fire, as the best remedy with which he is acquainted for dispersing milk in the breasts (b).

ALOE.

ALOES: a bitter, gummy-refinous, inspissated juice; prepared from the leaves of certain thick fleshy-leaved plants of the same name. Three forts of it are distinguished in the shops.

- I. ALOE SOCOTORINA Pharm. Lond. Aloe fuccotrina Ph. Edinb. Socotorine aloes; brought from the island Socotora in the Indian ocean, wrapt in skins; obtained from the Aloe focotrina angustifolia spinosa flore purpureo Breyn: a variety of the Aloe perfoliata Linn. This fort of aloes is of a bright surface, and in some degree pellucid; in the lump, of a yellowish red colour with a purplish cast; when reduced into powder, of a golden colour. It is hard and friable in the winter, somewhat pliable in the summer, and softens betwixt the singers. Its bitter taste is accompanied with an aromatic slavour, but not sufficient to prevent its being disagreeable: the smell is not very unpleasant, and somewhat resembles that of myrrh.
- 2. ALOE HEPATICA Pharm. Edinb. Hepatic, Barbadoes, or common aloes; usually brought from Barbadoes, the best fort in large gourd shells, an inferiour kind in pots, and a still worse in casks; extracted from the Aloe C. B. Aloe dioscoridis & aliorum. Sloan jamaic. This is darker coloured than the foregoing, and not so clear or bright. It is

generally drier and more compact; though fometimes, especially the cask fort, quite soft and clammy. Its smell is much stronger and more disagreeable: the taste intensely bitter and nauseous, with little or nothing of the aromatic slavour of the socotorine.

3. ALOE CABALLINA. Caballine or horse aloes; prepared, probably, from the aloe guineensis caballina vulgari similis sed tota maculata Commel. prælud. not, as is generally supposed, the feces of the hepatic; the difference not being in purity, but in quality. It is easily distinguished from both the foregoing by its strong rank smell: in other respects it agrees pretty much with the hepatic, and is, not unfrequently, fold in its place. Sometimes it is prepared so pure and bright as scarce to be distinguishable by the eye even from the socotorine, but its offensive smell readily betrays it; and if this also should be dissipated by art, its wanting the aromatic slavour of the siner aloes will be a sufficient criterion.

Aloes is a stimulating cathartic bitter. Taken in sufficient doses to purge effectually, as half a dram or two scruples, it occasions commonly a great irritation about the anus, and fometimes a discharge of blood. In fmaller doses, as ten or twelve grains, repeated once or twice a day, it not only unloads the first passages, but attenuates and disfolves viscid humours in the remoter parts, warms the habit, quickens the circulation, and promotes the menstrual and hæmorrhoidal fluxes: its continued use renders the blood sensibly more fluid, as appears on venefection. For a time, in these small doses, it does not act by stool; but at length it produces a gentle looseness, of longer continuance than that occasioned by most other purgatives: hence its utility in habitual costiveness. This stimulating cathartic is particularly adapted to persons of a phlegmatic temperament and sedentary life, to cachectic indispositions, and oppressions of the stomach by viscid crudities contracted from irregularity: in dry bilious habits, it is often injurious, immoderately heating the blood, or inflaming the bowels.

This bitter juice is accounted destructive to worms, or to the matter which favours their production, whether taken internally, or applied in plasters to the umbilical region. It is powerfully antifeptic;

feptic; and commonly made an ingredient in tinctures and balfams for cleanfing and healing wounds or putrid fores.

Aloes confifts of a refinous matter, and a large proportion of a fubstance called gum. By boiling in water, in the proportion for instance of four ounces to a quart, it nearly all disfolves, except the impurities, into a dark coloured liquor; which on standing in the cold for a night, deposites the refin to the bottom, the gummy part continuing disfolved. Refina aloes From this folution (poured off from the precipitated refin, and, if any feculencies appear in it, passed through a strainer) the gum may be recovered in a folid form by evaporation. The coarfer forts of aloes may be purified from their feculencies, without any separation of the gummy and refinous parts, by straining the folution whilst hot, and fetting it directly to evaporate, without fuffering it to fettle.

The hepatic aloes is found to contain more refin and less gum than the focotorine, and this than the caballine. Twelve ounces of caballine aloes yielded two of refin, the fame quantity of focotorine three, of hepatic almost four: of gummy extract, the caballine yielded nine ounces, the focotorine fomewhat less than nine, the hepatic eight. The watery solution of the gummy part of the focotorine, after the separation of the refin, appeared of a bright brown colour, with a cast of red; that of the caballine, deep reddish brown; of the hepatic, brownish yellow, with-

out any tendency to redness.

The refins of all the forts, purified by folution in spirit of wine, (for in their fettling from the watery decoction of the aloes, the impurities of the juice subside along with them) have little smell; that obtained from the focotorine has fearcely any perceptible tafte, that of the hepatic a flight bitterish relish, and that of the caballine a little more of the aloetic flavour. The gummy extracts also are less disagreeable than the crude aloes: the extract of the focotorine has very little fmell, and is in tafte scarcely unpleasant: that of the hepatic is in smell somewhat stronger, but seems to be in taste rather less ungrateful than the extract of the focotorine: the gum of the caballine retains a confiderable share of the peculiar rank smell of this kind of aloes, but its taste is not much more unpleasant than that of the extracts made from the other two.

The purgative virtue of aloes, contrary to that of most of the other cathartic vegetables, refides chiefly in the gummy part; the refin, though taken in confiderable doses, whether divided by testaceous powders, or

Ph. Lond.

Gummi aloes Pharm.

Aloe lota Ph. Edinb. dissolved in spirit of wine, having little or no cathartic power. Socotorine aloes, which contains more gum than the hepatic; purges more, and with greater irritation: the former therefore is to be preferred where a ftimulus is required, as for promoting or exciting the menstrual flux; whilst the latter is better fitted for a common purge. The vulnerary and balfamic virtues, on the other hand, refide principally in the refin; and hence the hepatic, which is more refinous than the focotorine, is found to be more ferviceable in external applications. The caballine aloes, on account of its offensive smell, is very rarely made use of, at least under its own name, either internally or externally.

The purgative aloetic gum dissolves, not only in watery, but likewise in spirituous menstrua; and even more readily in proof spirit and in rectified spirit, than in water or wine. When powdered aloes is macerated, or digested in a gentle warmth, with water, with wine, or with vinous spirits largely diluted, the powder softens, and becomes tenacious, and the folution goes on exceeding flowly: hence in making tinctures or folutions of aloes in these kinds of menstrua, it is of advantage to mix with the powder fome clean dry fand, which by keeping it divided, promotes the diffolution. With rectified and proof spirits, the aloes does not cohere, but continue's powdery till diffolved.

Aloes is fometimes taken by itself, fometimes mixed with saponaceous medicines, warmed with aromatics, acuated with pungent materials, combined with the deobstruent gums, &c. Many of these kinds of compositions have been received as officinals: a pill, for example, composed of equal parts of aloes and extract of gentian, with a fourth part of fal polychrest*: a powder, of eight parts of aloes, with two of canella alba+: a tincture, made by digefting five ounces of this powder in five ! pints of mountain wine; or one ounce of aloes, with one dram each of Jamaica pepper and ginger, in a pint and a half of the same wine | : pills of four parts of aloes, two of myrrh, and two or one ** of faffron, made up with fyrup of faffron or of orange peel: vinous and spirituous tinctures of the aloes with different proportions of the myrrh and faffron 1, &c. Among different aromatic materials made trial of, cloves feemed the best adapted for alleviating the offensiveness of the aloes: the committee appointed by the London college for reforming their pharmacopæia, made choice of canella alba, on account of its not rendering the medicine fo hot as the necessary quantity of the clove itself would do,

* Pil. aloeticæ Pharm. Edinb. +Hierapicra Pharm. Lond. tTinct.facra Pharm. Lond. Il Ph. Edinb. § Pil. Rufi Ph. Lond. ** Pil. communes vulgo Rufi Pharm. Edinb. 4 Elixiria proprietatis varia.

and yet having so much of the clove flavour, as to cover the aloes in a sufficient degree: some commend the casia caryophyllata, or clove bark, as having more of the clove flavour, than canella alba, and yet not being very hot.—Where volatile spirits are to be joined, a solution of the aloes in dulcified spirit of sal ammoniac, or in spirit of sal ammoniac made with quicklime, are very elegant preparations, and require little assistance from aromatics to render them supportable to the palate; the offensiveness of the aloes being greatly abated by the spirit, and the pungency of the spirit sheathed by the aloes: the spirit of sal ammoniac made with fixt alkaline salt does not dissolve near so much of the aloes as the two abovementioned.

ALSINE.

ALSINE Pharm. Edinb. (a) Alfine media C. B. & Linn. CHICK-WEED: a fmall, creeping, juicy herb; annual, common at all times of the year in shady cultivated grounds.

This herb was formerly employed in cataplasms against inflammations; and its expressed juice, or decoction, given also internally, as an aperient, antiscorbutic, antiphlogistic, and as a restorative, that is, perhaps, for abating hectic heats, in atrophies and consumptions. Nor do the virtues ascribed to it appear to be wholly without foundation: experiment discovers, that it is not destitute of active matter, though this matter is so far divided and diluted in the herb, as scarcely to manifest itself till separated from the grosser parts.

The fresh leaves have an herbaceous somewhat saline taste, without any remarkable smell: in distillation, with water or with spirit, they give over nothing. On expression they yield a large quantity of green coloured turbid juice; which dissicultly deposites its seces upon standing, but immediately parts with them on being heated to ebullition, and being now passed through a strainer, looks clear and reddish. The depurated juice, inspissated to the consistence of an extract, discovers to the taste a cool penetrating saline pungency, which quickly goes off, leaving a slight austerity in the mouth.

(a) Expunged.

ALTHEA.

ALTHEA Pharm. Lond. & Edinb. Althæa dioscoridis & plinii C. B. Althæa officin. Linn. Marshmallow: a soft hoary plant: with oblong undivided leaves; and pale flesh-coloured monopetalous flowers, cut deeply into five sections, set in a double cup, the outermost of which is divided into nine parts, the inner into five: the fruit consists of a number of capsules, set in form of a flat disk, containing each a single seed: the roots are long and slender, with several sibres, of a pale yellowish colour on the outside, and white within.—It grows wild in marshes and other moist places, though frequently cultivated in gardens. It is perennial, and flowers from June to near the end of summer.

All the parts of althea abound with a glutinous juice, of scarcely any smell or particular taste. The dry roots, boiled in water, give out near half their weight of gummy matter; which, on evaporating the aqueous sluid, forms a flavourless, yellowish mucilage. The leaves afford scarcely one fourth their weight, and the flowers and seeds still less; though the two latter have been looked upon by some as the most mucilaginous, and accordingly prescribed in less quantity (a) than the other parts of the plant.

Of all the mucilaginous vegetables, marshmallow root is, among us, of the most general use; for obtunding and incrassating acrimonious thin sluids, in tickling coughs from defluxions on the sauces and lungs, in hoarseness, erosions of the stomach and intestines, difficulty and heat of urine; and for lubricating and relaxing the passages in nephritic and

calculous complaints.

The root is sometimes given in powder, from a scruple to a dram or two, either by itself, or in conjunction with other materials of similar intention, as gum tragacanth, starch, &c. It is rather too bulky, however, for this form; and may, in most cases, be taken to better advantage in that of an infusion or decoction, sweetened with a little liquorice: an ounce of the dry root is sufficient for a quart or three pints of water, a larger proportion rendering the liquor disagreeably slimy.—A syrup,

made by boiling a pound of the fresh roots in a gallon of water till half Syrupus ex the liquor is wasted, pressing out the decoction, and after settling for a night, boiling it down with four pounds of fine fugar till the weight of the whole is fix pounds, is kept in the shops, and employed occasionally in fome diforders of the breaft, and for fweetening emollient decoctions in nephritic cases.

ALUMEN.

ALUMEN Pharm. Lond. & Edinb. ALUM: a semitransparent, austere, styptic falt; composed of the vitriolic acid, and a certain earth; which earth is either the pure argillaceous earth, or else is contained, in great quantity, in all the argillaceous fossils that have been examined.

The greatest quantities of this salt are artificially produced from different kinds of minerals, whose nature and composition are little A bluish slate found in the hills near Scarborough and in fome other parts of England (a), and a whitish stone at Tolfa near Rome (b), become richly aluminous by calcination; and a bituminous earth near Hall in Saxony, by exposure to the air: this last, if laid in large heaps, grows hot, like the pyritæ, and at length bursts into flame (c). There are, in Sweden, ferrugineous pyritæ, from which alum, as well as vitriol, is obtained (d); and most, if not all, of the aluminous flates, participate also largely of vitriol: it is probable that in all the matrices of this falt, the part, which becomes alum, differs from that which in the pyritæ becomes vitriol, only in the former having an argillaceous earth in the place of the metallic calx of the latter.

The alum, produced in the mineral, fometimes shoots upon the furface into fibrous efflorescences, called by the antients, from their form, alumen plumofum; though later times have applied that name to a fubstance of a very different kind. The falt is extracted from the earthy matter by elixation with water; and afterwards brought to a crystalline form, by evaporating the folution to a proper pitch, and then fetting it to

⁽a) Colepress, philosoph. transact. No. 142.

⁽b) Mercatus, metallothec. armarium iii. cap. 2.

⁽c) Hoffman, observ. physico-chym. lib. iii. obs. 8.

⁽d) Leopold, relatio de itinere suo suecico, p. m. 84, & segq.

shoot, with the addition of a little alkaline lye or putrefied urine, without which the crystallization does not succeed. Even when the pure earth, separated from alum, is redissolved in the vitriolic acid, the solution does not easily shoot into perfect crystals, till some alkaline salt, fixt or volatile, is added; this acid seeming not to fully satiate itself with the aluminous earth, and the unsatiated part preventing the crystallization of the rest (a). The alkaline liquor is to be dropt in by degrees, till a white precipitation begins to appear; a mark, that all the redundant acid is now saturated, and that a further addition would decompose more and more, proportionably to its quantity of the alum itself.

The English alum is colourless, and commonly in large masses; into which it is formed, by melting the crystals over the fire, with the addition of a little water, and pouring the fluid matter into wooden tubs, in which it concretes and assumes the figure of the vessel: the Roman is of a reddish hue, and in smaller crystallized masses. The name rock or rock alum is applied among us to the English, on account of the hardness and size of its masses; and by foreign writers to the Roman, on account of the hard stone or rock from which it is extracted. The Roman is thought to be somewhat less styptic and less nauseous than the English, and is supposed by some to have for its basis a somewhat different kind of earth.

ALUM is a strong astringent; one of the strongest of the substances of that class. It is in common use for external purposes; against relaxations of the uvula; in gargarisms for spongy scorbutic gums; in epithems and collyria for inflammations and defluxions of the eyes, &c. In this last intention, we have scarcely any application more effectual than the coagulum recommended by Riverius, made by agitating the white of an egg with a lump of alum, till it acquires the consistence of an unguent, which is to be spread on tow, and applied warm to the eyes at bed-time; proper evacuations, if the inflammation is considerable, being premised.

Coagulum aluminofum Pharm. Lond.

Internally, it is given in small doses, of half a grain or less, as a mild corroborant; and in larger ones, as ten, fifteen, and sometimes twenty grains, for restraining immoderate hæmorrhages. These large doses are

never advisable, but in profuse and threatening evacuations; as they are apt to nauseate the stomach, occasion gripes, and leave obstinate constipations of the bowels. The first dose or two sometimes purge a little.

It has been customary to mix alum, for internal use, with an equal * * Pulvis or with half its quantity of dragons blood; which ferves to difguife the alum, and render it, especially when the mixture is made by melting them together*, more flow of folution in the stomach, in consequence of which it fits eatier and may be given with less inconvenience in confiderable doses: this is, perhaps, the only advantage of the addition of dragons blood to alum. The Edinburgh college, in their last edition, Pulvis styphave in place of dragons blood substituted the gum kino, in the pro- ticus I Edinb. portion of three drams to one ounce and a half of alum; an alteration which much improves the medicine; as this aftringent gum is perfectly foluble in watery menstrua. Dr. Thompson, in the medical essays published by a society at Edinburgh, gives an account of the good effects of the former compound in uterine hæmorrhages; and affures us, that he had never found any medicine so much to be depended on, whether for correcting the too frequent return of the menses or their too great abundance, for stopping the floodings which women with child are subject to, or moderating the flow of the lochia. In violent bleedings, he gave half a dram, of a mixture of equal parts of the two, every half hour; and feldom failed to suppress the discharge before three or four drams had been taken. The fuccess of this medicine in these disorders induced him to prescribe it in the fluor albus, and in this also it had excellent effects.

Alum diffolves in twelve times its weight or less of water: on fetting the folution to exhale flowly in a moderately warm air, the falt concretes into crystals of eight or more triangular sides. The solution changes the colour of the blue flowers of plants, or their juices, to a red or purple, as acids do; and like them also, it coagulates milk and the serous humours of animals. The whey obtained by boiling a pint of cow's milk with two drams of powdered alum, is sometimes given in uterine hæmor- Serum alurhages, and recommended also in the diabetes (a), in doses of a quarter Ph. Lond. of a pint, three or four times a day. This liquor, like other aluminous folutions, is not a little ungrateful: nor does this method of obtaining

the folution admit of fo much precision, as could be wished for in a medicine of fuch efficacy in regard to the dose; a considerable part of the alum being retained in the curd, which taftes rather more strongly aluminous than the whey. The whey may be made more elegant by a

proper addition of fugar, and of dried red rofe buds.

Alumen uftum Phar.

This falt, exposed to the fire, easily liquefies, bubbles up in blifters, emits watery vapours amounting to about one fixth of its weight, and then turns to a light spongy unfusible mass, which seems on tasting to be almost infipid, but gradually diffolving in the mouth, discovers at length the same taste as the alum at first. This dried, or burnt alum, as it is called, is sometimes employed for drying foul ulcers, and confuming proud flesh, which it does with great mildness, but it is said to have an inconvenience of leaving a hardness upon the part.

The burnt alum, urged with a strong fire, gives out an acid spirit exactly fimilar to that obtained by the fame means from vitriol; the matter which remains, if the fire has been fufficiently intense and long continued, is the pure earth of the alum, white, light, and infipid. If any of the alum still retains its acid, which a considerable part commonly does though a pretty strong fire had been continued for fome days, this part may be extracted, by boiling in water, from the pure indiffoluble earth.

The earth of alum may be separated also by dissolving the alum in water, and adding a folution of any pure alkaline falt, or rather a volatile spirit: the liquor grows instantly milky on this addition; and on standing for a little time, the aluminous earth falls to the bottom, its acid being absorbed by the alkali. This earth, freed from the faline matter by repeated ablutions with boiling water, diffolves readily in all acids: folutions of it in the nitrous and marine are more flyptic, and more nauseous than alum itself: folutions of it in vegetable acids, though strongly styptic, are of a milder and less ungrateful kind, and promise to be, in many cases, medicines of no small utility.

AMBRAGRISEA.

AMBRAGRISEA Pharm. Edinb. Succinum grifeum. Succinum cinereum. Ambarum. AMBERGRIS: a marine bitumen; very light, fo as to fwim both in water and in rectified spirit of wine; growing soft in a gentle warmth; when warmed, of a fragrant smell; soluble in boiling spirit of wine, from which, if the saturated solution be set in a very cold place, or if a part of the menstruum be exhaled, a proportionable quantity of the ambergris concretes into a whitish unctuous substance.

The greatest quantities of ambergris are met with in the Indian ocean: pieces have likewise been now and then discovered in our own (a) and other northern seas. It is found floating on the surface of the sea, or adhering to rocks, or thrown out upon the shores, and sometimes in the stomachs of large sisses. It is usually in small masses, though there are accounts of very large ones, weighing more than an hundred (b) pounds, opake, rugged, of a greyish or ash colour intermingled with yellowish and blackish specks or veins, of a loose texture, friable in a certain degree like wax, breaking rough and uneven, and frequently containing pieces of shells and other like matters. It is said to be at first soft; and when sound in this state, to be often adulterated by incorporating different substances with it; an abuse which may in good measure be distinguished by the appearance and texture of the mass, and with more certainty by its differences from true ambergris in solubility, volatility and smell.

AMBERGRIS has scarcely any particular taste; and very little smell, unless heated, or much handled; in which circumstances, its smell is very fragrant, and to most people agreeable: set on fire, it smells like burning amber. It softens betwixt the singers, melts in a small degree of heat into the appearance of oil, and in a strong one proves almost totally volatile. Distilled, it yields an aqueous phlegm, a brown coloured acidulous spirit, a deeper coloured oil, at length a thick balsam, and sometimes a small portion of a concrete salt. The spirit, oil, balsam, and salt, are similar to those obtained by the same treatment from amber: except that the oil is of a more grateful smell.

It dissolves in pure spirit of wine, almost totally, but sparingly, and not without the assistance of a boiling heat. Neumann observes, that pure spirit may be made to take up about one twelfth its own weight of the ambergris: that spirits impregnated with a little essential oil, whether

⁽a) Charleton, de animal. append. de fossilib.

⁽b) Chevalier, description de la piece d'ambregris, &c. pesant 182 livres.

by the addition of the oil itself, or by distillation from oily vegetables, diffolve it more readily than pure spirit: that spirits drawn over from fixt alkaline falts, extract a deeper tincture, but diffolve no more, than those which have been rectified without that addition: and that the

dulcified acid and alkaline spirits have very little effect on it.

Ambergris is, in general, one of the most agreeable of the perfumes, and the least apt to disorder weak constitutions, or such as are liable to be offended by fubstances of that class. Taken internally, from two or three grains to a fcruple, it is accounted a high cordial, corroborant, and antispasmodic; in which light it is prescribed by Riverius in hypochondriacal affections. A folution of it made in a very highly rectified spirit distilled from roses, is recommended by Hoffman, in his physicochemical observations, as one of the most effectual corroborants of the nervous fystem. The orientals are faid to look upon it as an aphrodifiac, and suppose that the frequent use of it contributes to longevity.

Tinct. feu essentia ambræ Pharm. Parif.

Tinct. feu effentia regia Pharm. Parif.

The faculty of Paris directs a tincture to be drawn, by digefting two fcruples of ambergris in two ounces of a high rectified spirit impregnated with roses. They have also a compound tincture made from the same quantity of ambergris, with half as much musk, ten grains of civet, fix drops of oil of cinnamon, and four drops of oil of rhodium, digested together in four ounces and a half of a spirit impregnated with roses and orange flowers. This compound tincture is a very high perfume: a few drops of it give a fine fcent to a large proportion of inodorous matters. It is used also for heightening the natural odours of other bodies, as aromatic waters, spirits, &c. the principal secret, for this purpose, confifts in adding the perfume so sparingly, that while it heightens and improves the smell of the substance it is joined to, it may not betray its The most advantageous way of preparing these kinds of tinctures, in regard to the ambergris, appears to be, to make the spirit boil or fimmer with it first, that this ingredient may be completely dissolved before the more foluble ones are added. The vapour, which exhales during the coction, caught and condensed in proper vessels, has little flavour of the ambergris: water, distilled from it in the same manner, proves confiderably impregnated with its fragrance.

AMMI.

AMMI. BISHOPSWEED: an umbelliferous plant; producing small oblong seeds flat on one side, convex and surrowed on the other. The upper leaves are finely divided; the lower narrow, indented, set in pairs along a middle rib, with an odd one at the end.

1. Ammi verum Pharm. Edinb. (a) Ammi alterum semine apii C. B. Ammi odore origani J. B. Sison Ammi Linn. True bishopsweed; with reddish brown seeds: a native of Egypt, from whence the seeds are sometimes, though rarely, brought to us.

The feeds of the true ammi, when in perfection, are an elegant aromatic carminative; of a warm pungent taste, and a pleasant smell approaching to that of origanum. Distilled with water, they yield a confiderable quantity of a yellowish essential oil, containing their whole smell and slavour: the remaining decoction, thus divested of the aromatic part of the feed, is unpleasantly bitterish. Spirit of wine appears also to carry off, in its exhalation, the odorous principle of the ammi; an extract made by this menstruum, though very warm and pungent, and seeming to contain the whole taste of the feeds, having little or nothing of their specific smell.

2. Ammi Pharm. Lond. Ammi majus C.B. & Linn. Ammi vulgare majus latioribus foliis semine minus odorato J.B. Common bishopsweed; with larger and paler seeds: a native of the southern parts of Europe, and propagating itself plentifully in our gardens by the seeds which fall in autumn.

The feeds of this species are weaker both in smell and taste than those of the preceding; nor does their slavour at all resemble that of origanum. The several preparations of them are proportionably different: the essential oil, and the spirituous extract, are both less grateful and less pungent.

AMMONIACUM.

GUMMI AMMONIACUM Pharm. Lond. & Edinb. GUM AMMONIACUM: a concrete gummy-refinous juice; brought from the

(a) Expunged.

East Indies (a); generally in large masses, composed of little lumps or tears, of a milky whiteness: the external parts of the mass are commonly yellowish or brownish, and the white tears change to the same colour on being exposed for some time to the air. Of the plant, from which it is extracted, we have no further knowledge, than what is learnt from the seeds found among the tears; which resemble those of dill, except that they are larger, and apparently belong to a plant of the umbelliferous kind.

Ammoniacum has a strong smell, like that of galbanum, but less ungrateful, and a nauseous sweetish taste which is followed by a bitter one. Its principal virtue is that of resolving obstructions; in which intention, it is frequently made use of in asthmas and difficulty of expectoration, in menstrual suppressions, and cachectic indispositions. In obstructions of the breast, it is accounted the most effectual of the aperient gums: in hysteric cases, some of the others are preferred or joined to it, on account, chiefly, of their more powerful smell. It is most commodiously taken in the form of pills: the dose is a scruple or half a dram, every night or oftener: in larger doses, as a dram, it generally loosens the belly. Applied externally, it is supposed to discuss hard indolent tumours.

Gummi ammoniacum colatum Ph. Lond. It is purified from the feeds, small stones, &c. commonly intermixed among the tears, by softening or dissolving it in a little boiling water, pressing it, whilst hot, through a strainer, and then inspissating it to its former consistence. For internal use, the larger and siner tears, unpurified, are preserable to the common strained gum; for unless the process be very skilfully managed, it loses in the purisication great part of its smell, and not a little of its taste. In the shops, a composition of much inferiour virtues has been often sold in the room of strained ammoniacum.

Lac ammoniaci Phar. Lond. Ammoniacum, triturated with water, dissolves into an emulsion or milky liquor, and in this form acts rather more effectually than in the solid one of a pill. Simple penny-royal water is commonly employed for this purpose, in such proportion, that four spoonfuls (that is, two ounces) of the emulsion contain thirty grains of the ammoniacum. Some

have dissolved it in vinegar of squills, and thus obtained an expectorant

undoubtedly powerful, though more unpalatable.

If the milky folutions are kept fome time, they deposite a considerable quantity of resinous matter, and become clear. Inspissated, they yield an extract, of no smell, and of only a weak bitterish taste. In distillation, no essential oil is obtained, and the distilled water is but slightly impregnated with the slavour of the ammoniacum. In this respect, ammoniacum differs remarkably from most of the other deobstruent gums, as asafetida, galbanum, and sagapenum, which afford not only a strong distilled water, but an actual oil containing the concentrated slavour of the gums.

Rectified spirit of wine dissolves near one half of ammoniacum into a transparent reddish yellow liquor, which tastes strongly of the drug: the undissolved mucilaginous matter is nearly insipid. On distilling the filtered tincture by a gentle heat, the spirit which comes over has hardly any flavour of the ammoniacum: nevertheless the remaining extract proves

weaker, both in fmell and taste, than the juice in substance.

AMOMUM VERUM.

AMOMUM Pharm. Parif. Amomum racemosum C. B. Elettari primum Hort. malabar. True amomum: a cluster of round fruits, or seed vessels, of an oriental plant. Each fruit is about the size of a middling grape; and contains, under a membranous cover, a number of small rough angular seeds, of a blackish brown colour on the outside, and whitish within: the seeds are lodged in three distinct cells, and those in each cell joined closely together, so that the fruit, on being opened, appears to contain three seeds. Ten or twelve of these capsules stand together, without pedicles, upon a woody stalk about an inch long: each single capsule is surrounded with six leaves set in form of a star; and the part of the stalk, void of fruit, is clothed with leafy scales.—Of the other parts of the plant, we have no certain account.

THE feeds of amomum are a strong and grateful aromatic; of a quick penetrating fragrant smell, somewhat like that of lavender, but more agreeable; and of a very warm pungent taste, approaching to that of camphor. They are said to yield in distillation a large portion of a subtile

fubtile effential oil. The husks have the same kind of flavour, in a lower degree.—These seeds have long been a stranger to this country. They are directed as an ingredient in the theriaca, in which they have been commonly supplied by the seeds of the amomum vulgare; and the London college, under the name amomum, have now allowed either the verum or vulgare to be taken indifferently. The college of Edinburgh, while they retained that composition, employed cloves as a succedaneum to the amomum.

AMOMUM VULGARE.

AMOMUM VULGARE. Sifon quod amomum officinis nostris C. B. Sifon amomum Linn. BASTARD STONE PARSLEY: an umbelliferous plant; very much branched; with a firm stalk higher than the branches; deep green, winged, serrated, parsnep-like leaves; upright umbels; and small, narrow, oblong, striated, dark brownish seeds, slat on one side and convex on the other. It is perennial, grows wild under moist hedges and by the sides of ditches, slowers in June and July, and ripens its seeds in August.

THE feeds of the amonum vulgare have a light agreeable smell and a mild warm aromatic taste. They have been sometimes given as carminatives and diuretics, like the other warm seeds, and usually substituted in the shops for those of the amonum verum, from which, however, they are very considerably different, in quality as well as in appearance: they are not near so hot or pungent, nor is their slavour of the same kind.

These seeds, insused in water, give out very little of their virtue: by boiling, their slavour is soon dissipated, and the liquor becomes disagreeably bitterish: in distillation with water, they yield a small portion of a yellowish essential oil, which tastes and smells strongly and agreeably of the seeds.

Rectified spirit readily extracts their virtue, and what is pretty singular, gains from them a green tincture: the spirit, drawn off by distillation from the filtered liquor, brings over with it nothing considerable of the slavour of the seeds: the remaining extract tastes strongly and smells lightly of the amomum, and proves a moderately warm, bitterish, not ungrateful aromatic.

AMYG-

AMYGDALA.

ALMOND: an oblong, flattish, white kernel, covered with a thin brownish skin: produced by a tree which resembles the peach in its leaves and flowers, but differs in the fruit; the stone being covered with a dry tough matter, of a disagreeable taste; and the shell, though wrinkled and cavernulous, yet not rugged.

THE almond tree, Amygdalus communis Linn. is a native of Africa, and cultivated in great plenty in some of the southern parts of Europe. It is now likewise naturalized to our own climate, in which it produces fruit not inferiour to that which we receive from abroad. It flowers earlier in the spring than most other trees, though the fruit does not ripen till autumn.

There are two forts of almonds, one of a foft fweetish taste, the other bitter. The eye distinguishes no difference betwixt the trees which yield the sweet and the bitter fort, nor between the kernels themselves. It is faid, that the same trees, which in a wild state bore bitter almonds, have, when cultivated, afforded the sweet kind; and that the sweet, from want of culture, have degenerated into bitter. The almonds we receive from Barbary, where the tree is indigenous, are bitter; whilst those of Europe, and other parts where it is cultivated, are in general sweet.

Great care is requisite in the choice of these kernels, particularly the sweet fort; as they are very apt to become rancid in keeping, and to be preyed on by an insect, which eats out the internal part, leaving the almond to appearance entire.

I. AMYGDALÆ DULCES Pharm. Lond. & Edinb. Sweet almonds. These are, for most purposes, blanched, or freed from the outer thin acrid skin, by steeping them in hot water till it is softened sufficiently to be peeled off.

Sweet almonds, used in food, are difficult of digestion, and afford very little nourishment, unless extremely well comminuted. As medicines, they contribute, on account of their soft unctuous quality, to blunt

blunt acrimonious humours in the first passages, and thus, sometimes,

give present relief in heart-burns and other like complaints.

On expression, they yield a large quantity, near half their own weight, of oil: which, though it has no particular taste or slavour, is somewhat more agreeable to the palate than oil olive, or most of the other common expressed oils; and hence is employed medicinally, for internal uses in preference to those oils; for obtunding acrid juices, and softening and relaxing the solids; in tickling coughs, hoarseness, costiveness, nephritic pains, &c.

On boiling almonds in water, a part of their oil separates and is gradually collected on the surface: in digestion with rectified spirit, no separation was observed. Very little of the almond is dissolved either by spirit or by water; the decoctions, made in both menstrua, leaving, on evaporation, only a small portion of a somewhat unctuous sweetish matter.

On triturating the almond with water, the oil unites with the aqueous fluid, by the mediation of the mucilaginous and farinaceous matter of the kernel, into an emulsion or milky liquor; a small quantity of powdery matter remaining undissolved: almonds that have undergone the strongest action of the press, retain still so much of their oil, as to communicate a milky hue to water.

These liquors participate of the emollient virtues of the oil, and hence are prescribed in the same intentions as the oil itself; particularly in heat of urine and stranguries, whether arising from a spontaneous acrimony of the humours, or the operation of cantharides or other irritating medicines. They are given also as diluents in acute diseases; and in some cases, for supplying, in some degree, the place of animal milk, with which they have a great analogy.

Emulsio communis. An ounce of almonds forms an emulsion of a due consistence with a quart of water; which is to be gradually poured on, after the almonds have been first thoroughly pounded. A little sugar or other grateful materials are commonly added, the palatableness of the liquor being a point of some importance, as it is in all cases intended to be drank plentifully. For most of the intentions, in which emulsions are generally given, gum arabic is an useful addition: if the water is heated, to hasten the solution of the gum, it must stand till grown cold before it is poured on the almonds, otherwise the emulsion will be imperfect.

The

The pure oil of almonds, exposed for a few days, to a heat equal to that of the human body, becomes rancid and acrimonious. Emulfions, on the other hand, on standing for some hours, throw up a white cream to the furface, and the whey-like liquor underneath grows, not rancid, but four. Hence fome afcribe to emulfions an advantage, in inflammatory distempers, above the pure oil, of not being subject to become acrid and irritating by the heat of the body, but tending rather to a state in which they may contribute to abate inflammation. Acids, mixed with emulfions, promote the feparation of the oily and ferous parts, producing immediately a thick curd, nearly after the fame manner as they do in animal milk.

The pure oil, triturated with a thick mucilage of gum arabic, forms a more permanent emulfion; from which the oil does not feparate on standing for some days, nor on the addition of acids; though it is speedily difengaged by alkalies both fixt and volatile. One part of gum, made into a mucilage with an equal quantity of water, is fufficient for four parts of the oil. The white or yolk of an egg, and a mixture of fyrup with a small quantity of volatile spirit, render the oil also soluble in water, but less perfectly.

Sweet almonds are an useful intermedium for uniting with water substances which of themselves are not miscible with it. Camphor, and the purgative and other refins, whether native or prepared by art, triturated with about fix times their quantity of almonds, diffolve along with them in water into a milky liquor, and are thus excellently fitted for being taken in a liquid form.

2. AMYGDALÆ AMARÆ Pb. Lond. & Edinb. Bitter almonds.

BITTER almonds agree with the fweet in yielding a large quantity of oil, and in being miscible with water into an emulsion. The oil has no perceptible bitterness, and is not in any respect distinguishable from that of the fweet almonds: the college allow, for medicinal use, the oil of either fort to be taken indifferently. The matter remaining after the Oleum expression of the oil, retains all the bitterness, and tastes much stronger amygdalinum Pharm. than the almond did at first.

Lond. & Edinb.

Great part of the bitter matter dissolves, by the assistance of heat, both in water and in rectified spirit: and a part arises also with both menstrua menstrua in distillation. Spirit seems to extract, and water to elevate, the most. It did not appear that the whole is dissolved or elevated by

either, or by the alternate application of both.

Bitter almonds, and emulsions made from them, have been recommended as aperients, resolvents, diuretics, and anthelmintics. They are, doubtless, of some use in the above intentions, but apparently of too dangerous a kind. The almonds in substance, taken freely, occasion sickness and vomiting: to dogs, and some other animals, they are poisonous. A simple water, strongly impregnated with their volatile parts by distillation, has been found also poisonous to brutes; and there are instances of cordial spirits slavoured by them being poisonous to man.

It is probable, that the directly noxious matter of the almond is that in which its bitterness and flavour reside; and that the activity of this matter is increased, by its separation from the gross oil and farinaceous substance, by which it was enveloped and obtunded in the kernel itself. The kernels of other fruits, that have any bitterness or particular flavour, appear to be impregnated with a substance of a similar nature to this poisonous principle of bitter almonds *(a).

ANACARDIUM.

ANACARDIUM: a moderately large kind of nut; whose kernel is covered by two tough rinds; betwixt which is lodged a fungous sub-stance, containing in its cells an extremely acrid matter, in a liquid state when the nut is fresh, though often by long keeping growing dry. It is the produce of certain large Indian trees, of the class of the pruniferæ of Ray.

I. ANACARDIUM Ph. Parif. Anacardium orientale. Anacardium or Malaca bean: externally of a shining black colour, of the shape of a heart flattened, with a very thick pedicle occupying almost the whole basis. The tree, which is found only in the East Indies, is called by Ray arbor indica fructu conoide cortice pulvinato nucleum unicum nullo ossiculo

^{*(}a) Bergius mentions having frequently prescribed with success an emulsion of bitter almonds in intermittents, in the quantity of a pint or two daily during the intermission; and afferts that it had sometimes cured where the bark had failed. Mat. Med. p. 412.

tectum claudente. It is thought to be the Avicennia germinans of Linnæus.

2. Acajou, Cajous, Anacardium occidentale. Occidental anacardium or cashew nut: externally of a greyish or brownish colour, of the shape of a kidney, somewhat convex on one side, and depressed on the other. The tree, a native both of the east and west Indies, is called by Ray pomifera seu potius prunifera indica, nuce reniformi summo pomo innascente, the Indian tree bearing a fruit like an apple, with a kidney-shaped nut growing on the top of the apple; or rather with an apple growing between the nut and its pedicle, for the nut, as he observes, is produced first. It is the Anacardium occidentale of Linnæus.

These nuts have been commended by fome as possessing great medicinal virtues, and condemned by others as very dangerous. The kernels appear to have no hurtful quality: they are said to be eaten by the Indians, have a pleasant sweetish taste, yield an insipid oil upon expression, form an emulsion with water, and are apparently of the same nature with sweet almonds. The acrid juice lodged between the rinds is strongly corrosive, and is said to be used by the Indians for confuming sungous slesh, and for destroying the sensibility of aching hollow teeth. The juice is recommended by some against freckles, and other cutaneous desormities; which it removes only by excoriating the part: Geoffroy cautions against the too free application of this cosmetic, and relates that he has seen erysipelases break out all over the face from the imprudent use of it.

ANAGALLIS.

ANAGALLIS Pharm Parif. Anagallis flore phæniceo & anagallis cæruleo flore C.B. Anagallis arvensis Linn. Pimpernel: a low, creeping, juicy plant, resembling chickweed; from which it differs, in the leaves being spotted underneath, and having no pedicles; in the seed vessel not opening at top, but horizontally; in the slowers being not white, but red or blue. The red slowered pimpernel is called male, and the blue seemale: they are both annual, grow wild in corn-sields and other cultivated L 2 grounds,

grounds, chiefly in fandy ones, and flower from May to August; the first is frequent, the other rare.

The leaves of both the pimpernels have hardly any smell; and when chewed in substance, discover little other than an herbaceous taste. They are not however wholly destitute of medicinal powers: for the expressed juice, on being depurated by settling, and then inspissated to the consistence of an extract, affects the organs of taste with a pungent saline austerity. It appears therefore that these herbs have some claim to the resolvent and detergent virtues ascribed to them by some writers; though neither a decoction or tincture of them, nor their juice in its dilute state, and much less their distilled water, can exert those virtues in any considerable degree.

ANCHUSA.

ANCHUSA, Alcanna, Pharm. Edinb. Anchusa floribus puniceis C. B. Anchusa tinctoria Linn. Alkanet: a rough hairy perennial plant, with unbranched procumbent stalks; of the bugloss kind, and differing from the common buglosses chiefly in the red colour of its roots. It grows wild about Montpelier and in the eastern countries, and is cultivated in some of our gardens; but the roots, produced in this climate, are paler coloured than those which we receive from abroad.

The roots of anchusa, when in perfection, are externally of a deep purplish red colour. The red cortical part, separated from the whitish woody pith, imparts a fine deep red to oils, wax, and all unctuous substances, and to rectified spirit of wine. To water, it gives only a dull brownish hue. The spirituous tincture, on being inspissated to the consistence of an extract, changes its fine red to a dark brown. In these general properties, the deep and pale roots agree with one another, and differ from all the rest of the red drugs we know of: it is not, therefore, probable, that the deep colour of the foreign roots is owing, as some have supposed, to the introduction of an extraneous tincture.

Alkanet root has little or no smell, and scarcely any taste: extracts made from it, by water and by spirit, are bitterish and roughish, but in too low a degree to be regarded as medicines. Its chief use is for colouring oils, plasters, lip-salves, &c. which receive a fine deep red

from

from one fortieth their weight of the root: the confistent uncluous materials are for this purpose to be liquefied in the heat of a water bath, the powdered anchusa added, the mixture stirred now and then, till suffi- Pomatum ciently coloured, and then strained through a linen cloth.

rub. Pharm. Parif.

ANDROSACE.

ANDROSACE sive Acetabulum Pharm. Paris. Acetabulum marinum minus Tourn. Androfaces matthioli, sive fungus petræus marinus, sive umbilicus marinus J. B. Cotyledon marina: A submarine production, found on rocks and on the shells of fishes, about the coasts of Montpelier and elsewhere; consisting of numerous, slender, short, filaments, more or less bent or arched, of a whitish or grey colour, hard and brittle, bearing each upon the top a striated concave body nearly of the figure of an inverted cone.

This fubstance reduced into powder, is used in France, as we are told by the faculty of Paris in the last edition of their codex medicamentarius, for destroying worms, and for dropsies. It does not however promife to be of any fervice in either of these intentions; or to differ materially from the coralline, which has also been used, as a vermifuge, with little fuccess: like that marine body, it is apparently of a stony or testaceous nature, impregnated with a little faline matter, which, when fresh, it discovers to the taste. It is remarked of the dried androsace, that on being held in the flame of a candle, it yields a dazzling brightness, and this repeatedly for several times (a); a phenomenon which I have observed the coralline also to exhibit.

ANETHUM.

ANETHUM Pharm. Lond. & Edinb. Anethum hortense C. B. Anethum graveolens Linn. DILL: an annual umbelliferous plant, with very finely divided leaves' and yellow flowers: producing brownish or dark coloured oval feeds, flatted on one fide, convex and marked with three longitudinal ridges on the other, and furrounded about the edges

with a yellowish leafy margin. It is a native of the warmer climates, cultivated with us in gardens, flowers in July, and in September sheds its seeds, by which the plant is plentifully propagated.

THE feeds of dill have a moderately warm pungent taste, and an aromatic smell, but not of the most agreeable kind: they are given as carminatives, to the quantity of a dram at a time, in flatulent colics and indigestion from a laxity of the organs and viscidity of the humours. The leaves are weaker and less grateful than the seeds: the roots have nothing of their flavour.

Water extracts very little of the virtues of dill feeds by infusion or digestion for many hours. In boiling, their whole slavour exhales along with the watery vapour, and may be collected by distillation: the distilled water, drawn off to the quantity of a gallon from a pound of the seeds, is kept in the shops, and occasionally made the basis of carminative draughts and juleps: its flavour is more agreeable than that of the seeds in substance. The simple water keeps better than any in the shops. A two ounce vial full, corked, after standing on a shelf many years, was clear, without seculence, and retained the slavour of the dill; and might then be looked on as an elegant simple water. Along with the water arises a considerable portion of essential oil, in taste moderately pungent, and smelling strongly of the dill: this is given from one to three or four

Ol. effent. fem. anethi. Pharm. Lond.

Aqua feminum anethi

Pharm. Lond.

Rectified spirit, digested on dill seeds, readily extracts both their smell and taste: the colour of the tincture is a bright yellow: the spirit, gently distilled off from the siltered liquor, brings over very little of its slavour, leaving in the extract nearly all the active parts of the seed.

drops or more, as a carminative, and in hiccups.

ANGELICA.

ANGELICA: a large umbelliferous plant; with hollow, jointed stalks; and indented, oval, pointed leaves, set in pairs along a middle rib with an odd one at the end, containing in their veins a milky juice, which on drying turns yellowish: the ribs of the leaves are channelled on the upper side, and joined to the stalks by large membranous bases or sheaths. The seeds are white or pale coloured, somewhat oval, slat on one side, convex and marked with three longitudinal ridges

ridges on the other, furrounded about the edges with a leafy margin. The roots are long and thick, externally of a dark brown colour, internally white and juicy, and when dry of a fpongy texture.

- 1. Angelica silvestris: Angelica fylvestris major C. B. Angelica sylvestris Linn. Wild angelica; with all the leaves alike, except that the odd one at the end is larger than the rest. This species grows wild, in moist grounds, in several parts of England: it is perennial, and slowers in July. All the parts of this plant are similar in quality to those of the following species, but rather weaker, and hence the medicinal use of this is now superseded by the other.
- 2. Angelica, Pharm. Lond. & Edinb. Angelica fativa C. B. Angelica Archangelica Linn. Angelica, garden angelica; with the odd leaf at the end of each rib, and generally some of the others also, cut into two or three lobes.

This is found by the fides of rivulets in the mountains of Lapland, and cultivated in gardens in the different parts of Europe for medicinal purposes and for the use of the confectioners. Bohemia and Spain are supposed to produce the best; the college of London directs the roots brought from Spain only to be kept in the shops. Linnæus, however, assures us, that it proves most vigorous on its native northern mountains (a). It is naturally a biennial plant; but if the stalks are cut down before they have run to slower, the roots send forth new heads, and may thus be continued for many years. The roots are in greatest perfection in the second spring: they should be thoroughly dried, kept in a very dry place, and frequently aired, otherwise they are apt to grow mouldy, and to be preyed upon by worms.

The roots of angelica are one of the principal aromatics of European growth, though not much regarded in the present practice. They have a fragrant agreeable smell, and a bitterish pungent taste, mixed with a pleasant sweetishness, glowing upon the lips and palate for a long time after they have been chewed. On wounding the fresh root early in the spring, it yields, from the inner part of the bark, an uncluous yellowish odorous juice, which, gently exsiccated, retains its fragrance, and proves

an elegant aromatic gummy-resin. On cutting the dry root longitudinally, the resinous matter, in which the virtue and flavour of the angelica resides, appears concreted into little veins (a). In this state, it is readily and totally dissolved by rectified spirit, and tinges the menstruum of a bright golden colour: on distilling off the spirit from this solution, very little of the slavour of the angelica arises with it, nearly all the active matter of the root remaining concentrated in the extract. Water gains also from this root a pretty deep yellow colour, but extracts little of its taste or smell: in distillation with water, there arises a small portion of essential oil, of an highly pungent taste, and smelling strongly of the angelica: the remaining decoction, thus divested of the aromatic matter of the root, is nauseously sweetish and subscrid.

The other parts of the plant have the same kind of taste and flavour with the roots, but their active principles are far more perishable. The feeds, which come the nearest to the roots, can scarce be kept till the fpring after they have been gathered, without the lofs of their vegetative power, as well as a diminution of their medicinal virtue: the leaves lofe greatest part of their virtue on being barely dried. For some purposes, however, they are well adapted: the fresh leaves, as well as the seeds, on being distilled with water, give over to the liquor the whole of their aromatic matter, which in this form proves sufficiently durable: some of the officinal distilled waters are flavoured with these materials, and the committee of the London college report, that after trial of fundry others, for removing the difagreeable flavour which the addition of vinegar communicates to spirituous waters, angelica was found to answer this end the most effectually. The virtue of the seeds, like that of the roots, is extracted very imperfectly by water, and completely by spirit; and though it rifes totally in distillation with water, is left by spirit, almost entire, in the inspissated extract: the spirituous tincture is of a bright straw colour, the watery infusion of a dark brown.

Caules angelicæ conditæ Pharm. Lond.

The stalks, candied with fugar, make an agreeable sweetmeat.

ANIME.

ANIME Pharm. Edinb. (b) Resina courbaril. ANIME: a transparent amber-coloured resin, exuding from the trunk of a large tree

(a) Grew, idea of philosophic, hist. of plants, §. 41.

(b) Expunged.

growing

growing in Brazil and New Spain; Hymenæa Courbaril Linn. A finer fort is faid to be fometimes brought from the eastern countries; but in the shops, only the American is met with, of different degrees of purity: the small tears are generally the purest; the larger masses being often full of earth, agreeably to Piso's account, that the liquid juice, running down from the tree, sinks into the ground, and is thence afterwards dug up.

Anime has a light pleasant smell, and little or no taste. It is readily friable between the teeth, but on long chewing softens and sticks together. Laid on a red-hot iron, it immediately melts, catches slame, and burns quickly away, with a fragrant smell, leaving only a small quantity of whitish ashes. It gives out little or nothing to aqueous liquors, but dissolves entirely in rectified spirit: the solution is of a yellow colour, smells agreeably of the anime, and has a warm pungent bitterish taste. The fragrance of this resin arises totally in distillation with water, and in part with spirit: on distilling with water a large quantity of anime, a small portion of essential oil is obtained.

The Brazilians are faid to employ anime in fumigations for pains and aches proceeding from cold; and in liniments or plasters for paralytic complaints, bruises, $\mathcal{C}c$. With us, it is rarely, if ever, made use of

for any medicinal purpose.

ANISUM.

ANISUM Pharm. Lond. & Edinb. Apium anisum dictum semine suaveolente Tourn. Pimpinella Anisum Linn. Anise: a small annual umbelliserous herb; producing roundish striated seeds, flatted on one side and pointed at one end, of a pale colour inclining to a green. The upper leaves are divided into sine segments; the lower entire, roundish, and serrated about the edges. This plant, said to be a native of Egypt, Syria, and other places of the east, is cultivated, for medicinal and culinary uses, in the southern parts of Europe: it is raised also in some of our gardens, but seldom brings its seeds to perfection in this climate. The seeds brought from Spain, which are distinguished from those of other countries by being somewhat smaller, are accounted the best.

Aniseeds have an aromatic smell, and a pleasant warm taste accompanied with a degree of sweetness. They are of common use, as a warm M carminative,

carminative, in flatulent colics, in the gripes to which young children are subject, in flatulent pains and obstructions of the breast, in weakness of the stomach and indigestion, in diarrheas, and for strengthening the tone of the vifcera and intestines in general: they are supposed to be in these intentions the most effectual of the warm seeds. They are fometimes taken in powder, from a scruple to a dram; and in some places entire, candied with fugar.

They totally give out their virtue to rectified spirit, the seeds, after the action of this menstruum, proving inodorous and infipid: the tincture is of a bright lemon colour, and taftes very agreeably. The fpirit, distilled off from the filtered tincture, has a light taste of the seeds, but leaves far the greatest part of their virtue behind in the extract, which proves a very pleafant, fweetish, moderately warm, and not very pungent aromatic. In all these preparations made with rectified spirit, the peculiar fmell of the anifeeds, to some persons offensive, is in great measure covered by the spirit.

Of effentiale fem. anifi Ph. Lond. & Edinb.

Infused in water, they impart a little of their smell, but scarcely any tafte: in distillation they give over the whole of their flavour, the remaining decoction having nothing of the peculiar fcent or tafte of the anifeeds. Along with the water arises an effential oil, to the quantity of an ounce or more from three pounds. This oil, in colour yellowish, congeals, even when the air is not fenfibly cold, into a butyraceous white concrete. Its fmell, which exactly refembles that of the anifeeds, is extremely durable and diffusive; its taste milder and less pungent than that of almost any other distilled vegetable oil: twenty drops may be taken for a dose, though common practice rarely goes beyond half that number: it is recommended chiefly in diforders of the breaft, and faid to be less effectual in flatulencies and colics, than the feeds in substance. Geoffroy observes, that milk, drawn from the breast soon after the oil has been taken, is found impregnated with its fmell.

These seeds yield an oil likewise upon expression, of a greenish colour, in tafte grateful, and strongly impregnated with the flavour of the feeds: fixteen ounces, lightly moistened by exposure to the steam of boiling water, are faid to afford one ounce. This oil is composed of a gross, infipid, inodorous one, of the fame nature with the common expressed oils; and of a part of the effential oil of the feed, on which its flavour depends. On digefting the compound in rectified spirit, the odorous oil

is extracted; in distillation with water it is elevated, so as to leave the other by itself inodorous and insipid. The gross oil appears to reside in the kernel of the feed, the effential in the cortical part.

Among the aromatics, of fimilar intention, that have been tried in composition with aniseeds, those of angelica seem the best adapted to improve their flavour. A spirituous water prepared from a mixture of Aqua sem. equal parts of the two, by drawing off a gallon of proof spirit from half a pound of each of the feeds, is commonly kept in the shops, and Lond. proves a fufficiently elegant carminative cordial.

ANISUM STELLATUM.

ANISUM STELLATUM, seu sinense & philippense, & semen badian Pharm. Parif. Fæniculum sinense; Cardamomum siberyense. Zingi. INDIAN or STELLATED ANISE: a fruit or feed veffel; confisting of rufty brown coloured hard wrinkled capfules, half an inch or more in length, joined together by the bases, to the number of fix or more, in the form of a star; each of which includes one feed or kernel, externally gloffy and of the colour of linfeed, internally white. It is the produce of a small tree, growing in Tartary, China, and the Philippine islands, called by Plukenet euonymo affinis philippinarum infularum, anifum spirans, nuculas in capsulis stelliformiter congestis proferens: by Linnæus, Illicium anifatum.

THE capfules or husks of the stellated anise have a fragrant smell, and a fweetish glowing, not fiery, aromatic taste, resembling those of the common anifeeds, or rather of a mixture of anifeeds and fennel-feeds, but stronger, and more agreeable. The feeds are faid by some to have neither tafte nor fmell: of fmell they have very little; but in chewing they fill the mouth with an agreeable aromatic flavour, of the fame kind with that of the husks, but weaker, and accompanied with a greater fweetnefs.

The feeds afford, in distillation with water, the largest quantity of effential oil; and the husks, on being treated with spirit, yield the most acrid refinous extract (a). The oil is more limpid, and fubtile, as well as more fragrant, than that of the common anifeeds (b); and the

⁽a) Cartheuser, fundamenta m. m. ii. 327.

⁽b) Geoffroy, m. m. ii. 470.

fpirituous extract much warmer and more pungent. Infusions of the husks in water, divested of their more volatile parts by evaporation, leave an extract slightly aromatic, amounting to twice the quantity of that obtained by spirit, or half the quantity of the husks themselves (a).

These seeds are employed in the eastern countries, and in some parts of Europe, in preference to the common aniseeds, to which they appear, from their sensible qualities, to be superiour. They have not as yet been received in practice among us, and are very rarely to be met with in the shops.

ANTHORA.

ANTHORA Pharm. Parif. Antithora. Aconitum falutiferum five anthora C. B. Aconitum Anthora Linn. Yellow helmet flower: a plant with divided leaves, and naked flowers confisting of five petals, the uppermost of which is shaped like a hood: each flower is followed by three or more pods, containing wrinkled angular seeds. It is distinguished from the other aconites or wolfsbanes, by the leaves not being glossy, by their being cut quite down to the pedicle, and by the segments being very narrow and of nearly the same width from end to end.

This plant is a native of the Alps and Pyreneans, from whence the dried roots are fometimes brought to us. They are generally of an irregular roundish shape, sometimes a little oblong, of a brown colour on the outside and white within, hard to break, but not tough.

The root of anthora has a faint smell, and an acrid bitter taste, constringing the fauces and throat, accompanied with a kind of nauseous sweetishness. Its medical qualities are doubtful. Some (b) look upon it as a safe anthelmintic, an useful alexipharmac in malignant severs, and even as an antidote to the poisonous aconites, particularly the species called thora, from its supposed efficacy against which it is said to have received its name: others (c) ascribe to it virulent qualities, and relate instances of its occasioning vomiting, purging, great disorders of the stomach, heat, thirst, and anxiety. A competency of experiments, to fully determine this point, is as yet wanting: possibly this root, like

⁽a) Cartheuser. (b) Gesner, epist. p. 66 & 142. Geoffroy, mat. med. ii. 11.
(c) J. Bauhin. bist. plant. p. 94. L'Obel, adv. p. 301. Hoffman, de medicament. insecur. §. 30.
many

many others, may be possessed of noxious qualities when fresh, which are in great measure distipated or destroyed by drying or long keeping. But as all the falutary effects, that can be rationally expected from this drug, are obtainable from medicines of known innocence; common practice has never received the anthora, and the colleges both of London and Edinburgh have now expunged it from their catalogues of officinals.

ANTIMONIUM.

ANTIMONIUM Pharm. Lond. Antimonium, stibium, Pharm. Edinb. ANTIMONY: a ponderous brittle mineral, composed of long shining streaks like needles, intermingled with a dark leaden coloured substance; of no manifest taste or smell. It is usually brought into the shops in the form of conical loaves.

There are feveral mines of antimony in Germany, Hungary, and France, and fome likewife in England. It is fometimes found tolerably pure, but more commonly blended with a hard stone or spar, from which the antimony is feparated by eliquation. The mineral being broken in pieces, put into earthen pots whose bottoms are perforated with small holes, and a moderate fire applied round the veffels, the antimony melts out, and is received in conical moulds placed underneath. In thefe, the lighter and more droffy matter rifes to the furface, while the purer and more ponderous subsides to the bottom: hence the upper broad part of the loaves is confiderably less pure than the lower. The antimony, thus purified, is called crude, in distinction from its officinal preparations.

In some places the native mineral has been employed without purification. The masses which have suffered fusion may be readily distinguished, by the form which they receive from that operation; by their being free from any visible stony matter, pieces of which are generally found adhering to the unwrought ore; and by their striæ being larger. The English antimony appears to be, of all the forts, the most unfit for medicinal use, as having sometimes an admixture of particles of lead ore, of which I have feen specimens.

ANTIMONY was employed by the ancients in collyria against inflammations of the eyes, and for staining the eye-brows black. Its internal use does not seem to have been established till towards the end of the fifteenth

fifteenth century, and even then it was by many looked upon as poisonous. Experience has now fully evinced, that in its crude state, or when duly prepared, it is a medicine of fufficient fafety, and of great efficacy in fundry obstinate disorders, and that though some of its preparations are most violently cathartic and emitic, yet even these, by a slight alteration or addition, lose their virulence and become mild.

Antimonial medicines are principally made use of, as alterants, deobstruents, or gentle evacuants; in cutaneous foulnesses not scorbutic; in rheumatic pains and contractions of the limbs (a); in leucophlegmatic, cachectic, and catarrhous disorders; in intermittent fevers from obstructions of the vifcera, as obstinate quartans; and sometimes in continual fevers, and for promoting expectoration in peripneumonic and afthmatic cases: they generally have better effects in cold serous habits, than in hot bilious dispositions. The more active preparations are employed as emetics in apoplectic and maniacal diforders. It is observable, that even the strongest antimonials, the caustic solutions in mineral acids excepted, are given to horses in large quantity, some ounces a day, without any ill effect: in these animals, both crude antimony and its preparations feem to operate by promoting perspiration.

The virulent effects, which antimony produces in certain circumstances, have been ascribed by many to its participating of an arsenical fubstance (b). But the chemical properties of antimony, alledged in proof of this supposition, are by no means characteristic of that poisonous mineral; and its operation in the human body is extremely different. The most violent antimonials are rendered inactive by means which do not lessen the deleterious quality of arsenic; and some act with violence in far less doses than pure arsenic itself.

timonii.

CRUDE antimony is properly an ore, or a combination of a particular Regulus an- metal with common fulphur. The metallic part, like that of other fulphureous ores, is separated in its proper form, by roasting the powdered mineral over a gentle fire till the fulphureous fumes cease, and then melting the remaining grey calx with inflammable fluxes. The flux

commonly

⁽a) Two remarkable cases of the efficacy of antimony in pains and in inveterate contractions of the limbs, are related by Kunckel in his laboratorium chymicum, 3 theil, 32 capit.

⁽b) Neumann, chym. med. dogmat. experimental. ii. 339. Hoffman, metallurg. morbif. §. 21. Stahl, menf. Decemb. cap. 3. opufc. p. 486. 491.

commonly used for these purposes by the chemists, called from its colour black flux, is composed of two parts of crude tartar and one of nitre, ground together, fet on fire, and burnt in a covered veffel to a blackish alkaline coal. - The fulphur also may be obtained in its pure state, by Sulphur andigesting the powdered mineral in aqua regis, which dissolves the metallic rum. part, leaving the fulphur in form of a greenish yellow substance: this, purified by fublimation, appears, on all trials, the fame with common brimstone. The proportions of sulphur and metal vary in different antimonies; fome forts feem to hold about two parts of metal to one of fulphur, and others nearly equal parts of each.

The pure metal, called regulus of antimony, is of a bright white colour, a plated or leafy texture, very brittle, nearly feven times specifically heavier than water. It melts in a low white heat, and if continued in fusion, in an open vessel, gradually exhales in thick whitish fumes, which condense, on the bodies adjacent, into white flowers. Melted with common brimstone, it becomes similar, both in appearance and quality, to crude antimony. Crude antimony, like most other sulphureous ores, is easier of fusion than its pure metal: it melts before it grows red-hot, though not before the veffel is confiderably fo.

It is in this metallic part of antimony, that its proper medicinal powers refide. The pure metal is a medicine of extreme activity: a quantity too minute to be fensible on the tenderest balance, is capable of producing violent effects if given dissolved or in a soluble state. Acid wines take up so little of it, that the metal, after a number of infusions, feems to have loft nothing of its weight: thefe tinctures, neverthelefs, prove, in moderate doses, strongly emetic or cathartic; and in very small ones, for the most part diaphoretic. It has been cast into the form of fmall pills, which acted as violent cathartics, and after their paffage through the body have operated in the same manner again, and this repeatedly for a great number of times.

THE activity of this metal is abated by calcination, or by the expulsion of the inflammable principle, which makes a constituent part of this as of other metallic bodies: when thoroughly calcined, it appears entirely inert. Thus, if ground with twice or thrice its weight of nitre, and Ceruffa thrown by little and little into a red-hot crucible, it flightly deflagrates, and being now freed from the faline matter by ablution with water, is

found

found changed into a perfect white calx, which though taken in doses of a dram or two, is faid to have no fensible operation. In this deflagration, a part of the nitre is alkalized, and a portion of the calx disfolves in the water along with the alkali, as generally happens in the calcination of other metallic bodies with nitre: acids, added to this folution, precipitate the diffolved calx in form of a fubtile white powder, which is equally inactive with the undiffolved part. These perfect calces, of themselves fixt and unsufible in the fire, melt with saline additions, as fixt alkaline falt and borax, into a pale yellowish glass, inert (a) as the calces at first.

Materia perlata.

> The precise gradations of activity, between the virulence of the metal in its perfect metallic state, and its indolence in that of a perfect calx, are not well known; but thus much is certain, that it continues extremely active till the calcination is almost complete. When crude antimony is roasted over a gentle fire and kept constantly stirring, till the ceffation of the fumes shews the sulphur to be diffipated, the metal remains in form of a greyish-white powder, so far calcined, that on being urged with a strong fire, it melts into a dark yellowish red glass, no part of it refuming its metallic form. Nevertheless, both the calx and the glass are very virulent emetics; differing, however, from the metal itself, in this, that their active parts are soon exhausted by repeated infusion in vegetable acids (b), whereas the metal, so far as the experiment has been carried, feemed to lofe nothing of its power.

Vitrum antimonii Pb. Edinb.

> THE activity of the metal is restrained likewise by the combination of fulphur with it. Crude antimony, a natural mixture of it with fulphur, is altogether mild; doses of half a dram, or a dram, for the most part only gently loofening the belly, or promoting infenfible perspiration: the greater degree of tenuity the powdered mineral is reduced to, the more confiderable are its effects; and the case appears to be the same in regard to all the antimonials that are not totally diffoluble in the animal fluids.

> If a part of the fulphur of the antimony be separated, by such operations as do not calcine the metal, the remaining mass proves proportionably more active. Hence, as different forts of crude antimony, and different parts of one and the same mass, hold manifestly

(a) Malouin, chimie medicinale, part iv. chap. 50. (b) Boerhaave, elem, chem. process. 210.

different

different proportions of fulphur, it is probable that they vary in degree of activity.

The fulphur of antimony is separated by deflagration with nitre: the greater the quantity of nitre, to a certain point, the more of the fulphur is confumed, and the more does the metal, thus divested of its corrector, exert its virulence. An increase of the nitre, beyond the quantity which is fufficient to feparate the fulphur, renders the products, contrariwife, milder and milder; by more and more calcining, or de-

stroying the powers of, the metal itself.

Thus, antimony deflagrated with one eighth its weight of nitre, is faid Crocus antito act chiefly, in doses of fifteen or twenty grains, as an alterative or diaphoretic: with one fixth its weight, it vomits and purges, for the most part very mildly, in doses of eight or ten grains: with balf its weight, it vomits strongly, in the quantity of from one to five or fix grains; and Crocus antiwith equal its own weight +, it proves, in the same doses, a most violent emetic, operating as it were inexhauftibly, till its whole fubstance is Lond. col. expelled. All these preparations are of a dark red or yellowish red colour, and hence perhaps their name of crocus or faffron. The three first are taken from the fire as foon as the deflagration ceases: the last, which is the officinal crocus, is kept for fome time in fusion, during which a whitish faline scoria rifes to the surface, which is separated when the mass grows cold.

The last of the above proportions of nitre, to wit, equal the weight of the antimony, feems to be nearly that by which all the fulphur is destroyed, and the metallic part left bare. If the nitre be increased to twice the weight of the antimony, the metal itself is so far calcined by it, as to appear, after the deflagration, white; and if now freed from the faline matter by ablution with water, proves fo mild, as to occasion only some Emeticum light nausea and gentle vomiting, with a large discharge of saliva and thick urine. If the antimony be treated in the same manner with thrice its weight of the falt, it becomes a perfect indolent calx, the same with monii Phar.

that obtained by calcining the pure metal with nitre.

In this deflagration, a part of the nitre is changed by the fulphur into a neutral falt fimilar to that prepared from pure fulphur and nitre deflagrated together, that is, to the nitrum vitriolatum: this falt may be recovered from the water in which the calx is washed, by filtration, evaporation, and crystallization. A part of the nitre becomes likewise an

monii medi-

monii mitich committ. of +Crocus antimonii Pb. Lond. & Edin. metallorum vulgo: Hepar antimonii quibusdam.

mite antimonii Boer-Calx anti-Lond. Antimon. diaphoret. vulgo. Anodynum minerale Phar. Brand. alkaline falt, which, as formerly observed, renders a part of the metallic calx diffoluble: the crystallized falt is found to retain a little of this calx, but cannot be expected to receive from thence any particular virtues.

Calx antimonii nitrata Ph. Edinb.

Brand.

* The Edinburgh college, in their last pharmacopæia, direct the grey calx of antimony, as prepared for making the glass, to be calcined for an hour with equal its weight of nitre, and the mixture then to be washed with warm water till it becomes tafteless.

Febrifugum craanii Ph.

THE fulphur of antimony is separated also by fusion with fixt alkaline falts, which absorb it, and form with it a scoria on the surface. On melting five parts of antimony, with one of falt of tartar, and four of fea falt, which last does not appear to be of any great use in the process, a ponderous dark reddish mass is obtained, which, separated from the fcoria, is found to be fimilar in quality to the crocus prepared with one eighth of nitre; about as much of the fulphur being here absorbed and fcorified by the alkali, as is there burnt off by the nitre. This preparation is greatly celebrated by Hoffman and others, in fundry obstinate chronical diforders, and esteemed one of the best antimonials that can be given with fafety as alterants: it operates chiefly as a diaphoretic, and fometimes, though rarely, by stool or vomit; the dose is from three or four grains to a fcruple.

If eight parts of antimony, fix of tartar, and three of nitre, be mixed together, deflagrated, and brought into fusion, the alkaline falt, resulting from the nitre and tartar, will absorb the whole of the sulphur of the antimony, and the metallic part will fall pure to the bottom. Only a small quantity, however, of the metal separates in this process; for as foon as the alkali and fulphur are combined together, this compound begins to disfolve and scorify the metal, and scorifies more and more of it in proportion to the continuance of the fire; if the pure metal be melted with a composition of sulphur and alkali, it is in like manner changed into a scoria.

Sulphur antimonii præcipitatum Pharm. Lond.

These alkaline scoriæ dissolve in boiling water; and on adding acids, as spirit of falt, to the filtered folution, the sulphur and metal are precipitated together, in form of a reddish or reddish-yellow powder. - A like folution may be obtained by boiling crude antimony in alkaline lye: which, like the alkaline falts brought into fusion by fire, first disfolves the fulphur, and then, by the mediation of this, takes up a very confide-

rable

rable part of the metal: the college of Edinburgh directs two pounds of powdered antimony to be boiled in two quarts of foap lyes diluted with Sulph. authree pints of water, the matter being kept stirring with an iron spatula, and fresh water occasionally added to supply that which evaporates, for three hours; and the precipitation to be made by dropping diluted spirit of nitre into the strained liquor whilst hot : if the solution is suffered to Kermes micool, a spontaneous precipitation happens. - It is probable, that when the folution is thus procured by boiling in lye, the precipitate will be of more uniform strength, or vary less in the quantity of metal, than when the antimony and alkaline falt have been melted together; and that the precipitate, thrown down by acids, will be less variable than that which is permitted to separate spontaneously. In either case, however, the powder, which falls first, proves darker coloured, contains more of the metal, and operates with more force, than that which fubfides after-The using of the nitrous, marine, or vegetable acids, for the precipitation, is indifferent to the medicine; but the vitriolic might occasion a variation; the neutral falt, resulting from the coalition of this acid with the alkali, being less soluble in water, and not easily separable from the precipitate by washing. - These precipitates, washed from as much of the adhering faline matter as hot water will diffolve, prove gently emetic, in doses of five or fix grains, when taken on an empty stomach. Made into pills with extracts or refins, and taken on a full stomach by a little at a time, they act chiefly as alteratives and deobstruents: with these cautions, I am told, they have been increased to fixteen grains a day, without occasioning any disturbance upwards or downwards.

The alkaline scoriæ of antimony, pulverized whilst hot, and digested for three or four days in rectified spirit of wine, communicate the same colour, taste, and smell, as a mixture of pure sulphur with alkalies. The colleges both of London and Edinburgh judge, contrary to what fome have supposed, that a part of the metal, as well as of the sulphur, is taken up by the spirit : the former direct a tincture to be drawn, with Tinctura a quart of spirit, from fix ounces of powdered antimony melted with antimonii Ph. Lond. & twelve of any fixt alkaline falt; the latter, from four ounces of anti- Edinb. mony and fix of alkali (a). It is faid that these tinctures, taken on an empty stomach, have fometimes proved emetic.

rat. antim. Ph. Edinb.

neral. vulgo.

THE fulphur of antimony is absorbed likewise by most of the metals; most freely by iron. For this purpose, some iron nails, wire, or other like fmall pieces that may lie loofe in the crucible, are heated to a strong red heat, and about twice their quantity, or a little more, of antimony thrown upon them: the fulphur of the antimony immediately acts on the iron, and as fulphur greatly promotes the fusion of that metal, the whole foon melts: a little nitre is then injected, about one part to fix of the antimony, the crucible covered again, and the matter, when brought into thin fusion, poured into a warm greafed cone or mortar. regulus, freed from the fulphureo-ferrugineous fcoriæ, is purified by repeated fusion with one fixth or one eighth its weight of fresh nitre, till the nitre no longer receives from it any yellow or amber colour: if the regulus discovers, by its dull grey colour, sponginess, hardness, and difficulty of fusion, that it retains much of the iron, a little fresh antimony is injected, whose fulphur, absorbing the iron, hastens the purification. If the metal when poured out be in exceeding thin fusion, and the quantity of fcoriæ covering its furface confiderable, it generally assumes on the top a radiated star-like efflorescence.

Regul. antim. martialis.

Regul. antim. stellatus.

This regulus, though venerated by some of the chemists, is not materially different from that obtained by simply calcining the antimony, and reviving the calx with inflammable fluxes.—The scoriæ resulting from the first susion with iron are little other than a sulphurated iron, scarcely retaining any thing of the metallic part of the antimony: exposed to the air, in a shady place, they fall into a black powder, whose since parts, washed off with water, and deslagrated with thrice their weight of nitre, are Stahl's aperient crocust: the grosser part, treated in the same manner, is said to be not aperient, but enormously aftringent \$\frac{1}{2}\$.—The amber coloured scoriæ, arising in the purification of the regulus with nitre, are an excessively strong caustic alkali; powdered and thrown whilst hot into highly rectified spirit of wine, they impart, by digestion and agitation, a deeper or paler red colour \$\frac{1}{2}\$, according as the spirit was more or less oily, together with a penetrating pungency,

+ Crocus martis aperitivus Stahlii. t Crocus martis aftringens Stahlii || Nitrum causticum, Scoriareguli ant. fucci-§ Tinet. antimonii acris Ph. Brand. &c.

THE metallic part of antimony is corroded by the nitrous and vitriolic acids, into a white powder; and totally diffolved by aqua regia, if made

and, as is supposed, a detergent and diuretic virtue (b).

(a) Stahl, menf. Januar. Opufc. p. 523.

(b) Stahl, menf. Decemb. Opufc. p. 505.

with only a small proportion of nitrous acid, into a corrosive liquor. It may likewise be combined with the marine acid into a liquid form, by particular methods of application. If corrofive mercury fublimate (a combination of mercury with the concentrated marine acid) be mixed in powder with half its weight of powdered antimony; the acid of the sublimate begins immediately to act upon the metallic part of the antimony, and fumes, extremely noxious, arife fo copiously, that the utmost circumspection is requisite for avoiding them: the mixture being fet to distil in a wide necked retort, with a fire cautiously increased, the antimonial regulus arises, combined with the acid into a thick caustic liquor, which congeals, in the neck of the retort, in appearance like ice. This concrete, exposed for some time to the air, imbibes moisture and becomes fluid+: it may likewise be melted down from the neck of the + Causticum retort by cautiously applying a live coal, and afterwards rendered permanently fluid by distillation 1 in another retort: when liquefied in the first 1 Butyrum way, it is fomewhat less corrosive than in the other. The use of this butter, as it is called, is for confuming fungous flesh, and the callous lips of ulcers: it acts exceeding quickly, producing an eschar, which, as Boerhaave observes, generally separates the same day it is formed.

antimoniale Pharm. Lond. antimonii Pb. Edinb.

The butter, diluted with a large quantity of water, grows milky, and deposites its metal, intimately combined with a portion of the concentrated acid, in form of an exceeding white powder. The powder, repeatedly washed with water, becomes insipid, but still retains a portion Mercurius of the acid, and operates, in the dose of two or three grains, as a most violent and dangerous emetic.

Spirit of nitre dropt into butter of antimony, fo long as it occasions any effervescence, forms with the marine acid of the butter an aqua regia, which keeps the metal perfectly diffolved. If this folution be committed to distillation, the marine acid comes over first, and a little of the nitrous after it: the rest of the nitrous acid may be totally expelled from the remaining powder, by calcining it in a crucible for half an hour or more Bezoardiwith a strong red heat. Spirit of nitre, poured on the mercurius vitæ, cum minein like manner expels the marine acid, and is itself expelled by fire. The calces thus obtained, though formely looked upon as medicines of great virtue, are equally inactive with those, which are more compendiously prepared by deflagrating crude antimony with thrice its weight of nitre.

After the distillation of the butter of antimony, there remains in the

retort a black powder, composed of the mercury of the sublimate and the sulphur of the antimony. This, like the ethiops made from mercury and sulphur directly, on being urged with a red heat, sublimes into a cinnabarine mass, generally darker than the common cinnabar, and somewhat of a needled structure. It has been supposed that this cinnabar participates of the metallic matter of the antimony; but experiment shews that it does not, and that its difference from common cinnabar consists wholly in its containing a larger proportion of sulphur. Common cinnabar, sublimed with a little fresh sulphur, becomes exactly similar to that of antimony; and cinnabar of antimony, sublimed from a little iron filings, or such other substances as may detain its superstuous

fulphur, becomes the fame with common cinnabar.

Cinnabaris antimonii Pharm. Lond.

Crocus antimonii lotus Ph. Lond.

Tartarum emeticum Ph. Lond. & Edinb. CRYSTALS of tartar, boiled in water with the pure regulus, or crocus, or glass of antimony, dissolve a part of the metallic matter, small indeed, but sufficient to communicate a strong medicinal impregnation: the glass is said to dissolve more easily, and in greater quantity, than the other preparations. The college of London directs half a pound of crocus of antimony (reduced to a subtile powder, and boiled and washed with water till it becomes perfectly insipid) to be boiled with an equal quantity of crystals of tartar, in three pints of water, for half an hour; the liquor to be filtered, and after due evaporation set by to crystallize. The Edinburgh college orders six ounces either of the washed crocus, or powdered glass, and four of creme of tartar, to be boiled in four quarts of water for six hours; the liquor to be filtered when cooled, and either crystallized or evaporated to dryness.

The total evaporation of the fluid appears the best way of securing uniformity of strength to the medicine: for as only a part of the tartar is saturated with the metal, and as the part thus saturated is more soluble than the rest, some of the unsaturated tartar is apt, in crystallization, to shoot by itself. The solubility of the compound affords one of the best means for estimating its strength, or the degree of its impregnation with the antimony. Dr. Saunders relates (a) that an ounce of cold water, about the middle temperature of the air, dissolved, of some of the com-

mon emetic tartars of the shops, not thirty-two grains, or one fifteenth its own weight; whereas, of a well faturated fort, which he had himfelf prepared by long boiling, the same quantity of water dissolved fifty-two grains, or near one ninth its weight. Perhaps the most certain way of obtaining a faturated and uniform preparation of this kind would be, to digest the common emetic tartar in eight times its weight, or less, of cold water, and evaporate the filtered yellow folution to dryness: or to continue the boiling of the glass of antimony and tartar for twelve hours or longer, adding water enough occasionally to keep the tartar always diffolved, and at length to let the water waste so far, as not to exceed eight times the quantity of the tartar employed, after which the liquor is to be fuffered to cool, and then filtered and evaporated.

This preparation is one of the best of the antimonial emetics; as containing the active part of the antimony, made foluble by a mild vegetable acid, which does not, like those of the mineral kingdom, communicate any degree of corrofiveness: the dose is from two or three to fix or eight grains. It may be given also as an alterative or diaphoretic, in doses of a quarter of a grain or half a grain or more; and added, in the quantity of a grain or two, as a stimulus to the milder vegetable cathartics. It is faid that cafia diminishes the power of this medicine, but probably on no

good foundation.

Most forts of vinous liquors contain so much acid, as to extract, in a short time, a strong impregnation from the antimonial metal. The Vinum anticollege of London directs an ounce of the washed crocus to a pint and moniale Ph. a half of mountain; that of Edinburgh, the same quantity of the glass to a pint. It does not appear, that these or much greater differences in the quantities, affect the strength of the preparation; the same crocus being fufficient to impregnate many fresh portions of liquor. tinctures have been chiefly used, in the quantity of half an ounce or an ounce, as ftrong emetics: in fmall dofes, as thirty to fixty drops, they act commonly as diuretics or diaphoretics. A case is related in the Edinburgh Essays, Vol. II. in which the whey made with a gill and a half of antimonial wine produced fleepiness without vomiting. The curd had the fame effect.

The virulence of fome of the antimonials is greatly abated, by intimately mingling them with wax or refins. Powdered glass of antimony, injected into one eighth its weight of melted bees wax, over a gentle

Lond, & Edin.

fire.

Vitrum antimonii ceratum Ph. Edinb. fire, and kept conftantly stirring for half an hour, becomes so mild, that when given from two or three grains to twenty, it occasions for the most part only a few stools, or a slight nausea or sickness, and sometimes produces no sensible evacuation. This preparation has for some time been celebrated in dysenteries: several instances of its good success in these cases are related in the fifth volume of the Edinburgh medical essays.

From the foregoing review of the antimonial medicines, it appears, that the feveral preparations of this mineral, the caustic butter excluded. differ from one another only in degree of activity; and that the greater number must vary in strength, from small and unheeded variations in the manner of preparing them. And indeed, though their real qualities should be always the same, they may nevertheless operate with different degrees of force; from the juices in the first passages, or the food taken during their use, occasioning more or less to be dissolved. Sometimes the milder preparations, and even crude antimony itself, have, from acid foods, proved strongly emetic; and sometimes the more active have lain for a time indolent in the body, and afterwards, on taking the flightest acids, fuddenly exerted unexpected violence. Tinctures of the pure metallic part in mild vegetable acids appear to be the most safe and certain of all the antimonials; and capable of being fo managed, as to answer all the falutary purposes that can be rationally expected from any preparation of this mineral; what is effected in the others, by rendering the metal more or less foluble, being here obtained, with much less uncertainty, by giving actual folutions of it in larger or fmaller doses. Whether the wine, recommended by Huxham, or the tartarous folution, is the most eligible, experience only can determine. It is certain that both of them, as commonly prepared, are very variable in strength; the vinous folution, from differences in the degree of acidity, and confequently in the diffolving power, of the wine itself; the tartarous, from the process being more or less skilfully performed. We ought therefore, in prescribing these preparations, if we are not well assured of their strength, to begin with a small dose, and gradually increase it according to the effect.

APARINE.

APARINE Pharm. Edinb. (a) Aparine vulgaris C. B. Philanthropus. Goosegrass or cleavers: a flender, rough, annual plant, spreading upon bushes and sticking to whatever it touches; with foursquare, brittle, jointed stalks; oblong narrow leaves, set in form of a star, about eight at a joint; and small whitish bell-shaped slowers, followed by little round burs.

THE leaves and stalks of aparine yield upon expression a large quantity of turbid green juice, which when depurated becomes clear and reddish. The leaves in substance have no smell, and very little taste: the juice also, in its dilute state, seems little more than watery and herbaceous; but when inspissated to the consistence of an extract, it affects the organs of taste strongly, though only momentarily, with a pungent saline bitterness.

The juice of this herb has been given, in doses of two or three ounces, as an aperient in obstructions of the viscera, and as a diuretic in hydropic cases and suppressions of urine. * This medicine came into great vogue for scorbutic complaints a few years ago, in consequence of a letter printed in the news-papers. It was found in several cases to have a considerable effect, probably merely as a fresh vegetable juice, as its visible operation manifested very little activity. It seems now to be again fallen into neglect.

APES.

APES Pharm. Edinb. (b) BEES. This infect, dried and powdered, has been given internally as a diuretic, and applied externally (ground with honey or other like substances) for promoting the growth of hair. Some have slightly calcined the bee, in a close vessel, to blackness, and esteemed it, when thus prepared, to be a medicine, in some cases, of more virtue; a saline matter being now in good measure generated by the fire, though not as yet extricated from the other principles.—For my own part, I have had no experience of the bee itself prepared or unpre-

(a) Expunged.

(b) Expunged.

pared, nor is it used in practice: the valuable products, which this insect affords, honey and wax, will be treated of in their places.

APIUM.

APIUM, Eleoselinum, Pharm. Edinb. (a) Apium palustre & apium officinarum C. B. Apium graveolens Linn. SMALLAGE: an umbelliferous plant, with bright green winged leaves, cut flightly into three roundish portions, ferrated about the edges: the feeds are small, oval, plano-convex, furrowed, of a pale brownish or ash colour: the root long, about the thickness of the finger, furnished with a number of fibres, of a pale yellowish colour on the outside, and white within. It is biennial; flowers in August; grows wild in rivulets and watery places; and is frequently cultivated in gardens.

A poisonous plant, the cicuta aquatica or water hemlock, which grows naturally in the fame places with wild fmallage, has been fometimes mistaken for it. This may be distinguished, by its leaves being deeply divided, quite to the pedicle, into three long narrow sharp pointed fegments; whereas those of smallage are only slightly cut into three

roundish obtuse ones.

THE fresh roots of smallage, especially when produced in its native watery places, are supposed to participate, in some degree, of the ill quality of those of the hemlock kind, and to be particularly hurtful to epileptic persons and pregnant women. They have an unpleasant smell, and a bitterish somewhat acrid taste, weaker than those of the roots of the cicuta, but so much of the same kind, as to countenance the suspicion. that the fresh roots of wild smallage, if taken in considerable doses, may not be entirely innocent. By drying, they lofe greatest part of their ill flavour, and become fweetish: the poisonous quality of the cicuta also is faid to be abated by exficcation.

The dry roots of smallage have been employed, in apozems, as aperients and diuretics, in conjunction commonly with the other aperient roots. They give out their virtue, together with a pale yellow colour, both to watery and spirituous menstrua. On evaporating the watery infusion, the flavour of the root exhales, and the remaining extract proves unpleasantly sweetish. The spirituous tincture, inspissated, yields an extract, somewhat sweeter and less ungrateful than that made with water, and of a slight warmth or pungency: the smell of the root, which is pretty strong in the watery insusions, is in good measure covered by spirit both in the tincture and extract.

The feeds of smallage have been sometimes used as carminatives and aperients, and appear to be possessed of greater virtues than the root. They have a moderately strong grateful smell, and a warm bitterish taste. Insused in water, they impart to it very little of their flavour: distilled with water, they yield a small quantity of essential oil, of a very pungent taste, smelling strongly and agreeably of the seeds: the remaining decoction is unpleasantly bitterish. They give out the whole of their taste and smell to rectified spirit, and tinge the menstruum of a yellowish colour: the spirit, distilled off from the siltered tincture, has very little of the slavour of the seeds: the remaining extract is a moderately warm, pungent, bitterish aromatic.

This plant has been greatly improved, by culture, in the fouthern parts of Europe, and thence received in our gardens under the name of celery, apium dulce celeri italorum Tourn. In this state, it is much paler coloured, quite white towards the roots, of a pleasant sweetish somewhat warm taste, without any thing of the ill slavour of the roots and leaves of common smallage. Ray observes, that, if neglected for a few years, it degenerates into smallage again.

The roots of celery lose in drying about two thirds of their weight: the matter which exhales appears to be mere water. The dried roots, digested in rectified spirit, with a heat a little below boiling, soon give out the whole of their active matter, and become insipid. The tincture, which is of a yellow colour, deposites, on standing for some weeks, a considerable quantity of truly saccharine white slakes: inspissated, it yields a whitish extract, of a grateful warm aromatic sweetness. An extract made by water is likewise considerably sweet, but has nothing of the aromatic warmth of the spirituous extract.

The seeds of celery are much inferiour in aromatic flavour to those of smallage; and the several preparations of them are proportionably weaker and less grateful: the essential oil, in which the taste and flavour are

concentrated, is far less pungent than the oil of smallage seeds, and of very little smell. Thus one part of the plant degenerates in its quality, in proportion as the other is improved.

A 2. U A.

AQUÆ COMMUNES. Aqua nivalis, pluvialis, fluvialis, fontana. Common waters: snow, rain, river, spring waters.

It is needless to observe, how much the purity of waters is conducive to health; and how greatly, though by insensible degrees, the human body must necessarily be affected, by minute quantities of insalubrious matters in this universal diluent, and vehicle of all our aliment.

Among the common tests of the purity of water, the least fallacious are, its being perfectly colourless, transparent, and void of smell and taste; its dissolving soap into a smooth lather; boiling pulse tender; not changing the colour of syrup of violets or the juices of other blue flowers; and its mingling with alkaline and with acid liquors, with solution of sulphur in alkalies, and solution of silver in the nitrous acid, without precipitation or change of transparency. These trials serve to distinguish, in most cases, whether waters contain any considerable quantity of foreign matters, but what the particular matters are, they never can discover, different substances exhibiting, in the several experiments, similar phenomena: thus blue juices are changed red by alum as well as by acids, and green by the calcareous marine salt as well as by alkalies.

To determine, with any degree of precision, the contents of waters, a quantity of the water is to be evaporated, in clean glass vessels, with a heat scarcely exceeding that which the hand can support; that the solid contents may be procured by themselves, with as little danger as possible of the extrication or transposition of any of their principles. The dry matter being digested in a little pure distilled water, a faline substance is commonly extracted by the water, and an earthy one is left, no longer dissoluble in aqueous menstrua.

The earthy matter is commonly not one simple earth, but a combination of two or more: 1. aluminous earth, distinguished by its dissolving in the vitriolic acid into an austere liquor; 2. magnesia, dissolving in

the

the same acid into a bitter liquor; 3. calcareous earth, not dissolving at all in the vitriolic acid, but readily in the nitrous and marine, from both which it is precipitated by the vitriolic; 4. felenites, not dissoluble in any acid, till strongly calcined in contact with burning suel, by which process it is reduced to calcareous earth; 5. some of the absolutely indissoluble earths, whose particular species, in the small quantities wherein they are obtained in these kinds of experiments, it is difficult and of little importance to determine. The two first are rarely met with in the residua of waters; the others are frequent, perhaps universal.

The faline substances are; the mineral fixt alkali, natron; the vitriolic acid, combined with this alkali into fal mirabile, or with magnesia into fal catharticus, or with the aluminous earth into alum; the nitrous, acid, combined with the alkali into nitre (a), or with some of the soluble earths into nitrous salts; the marine acid, combined with the alkali into common salt, or with soluble earths into muriatic salts; or a volatile alkali combined with the acids into ammoniacal salts (see the respective salts.) *(b) The common, muriatic and nitrous salts are frequent; nitre, alum, sal mirabile, and ammoniacal salts very rare.

Most of these salts may, by careful crystallization, be separated in their proper form: they may likewise be distinguished, however blended together, by additions. 1. The fixt alkali, unsaturated, is known,

(a) Leigh, natural bift. of Lancasbire, &c. p. 39.

*(b) Another medium by which mineral substances may be rendered soluble in water is fixable air, or gas. This matter is found to be contained in large quantity in many mineral waters, particularly fuch as are diffinguished by a brisk sharp taste, and the property of sparkling when poured out. From some experiments of the Hon. Mr. Cavendish (Philof. Trans. Vol. LVII.) it was found, that calcareous earth could be diffolved in water impregnated with fixable air; and it appeared probable, that this combination actually takes place in the composition of Rathboneplace water. Mr. Lane, in Vol. LIX. of the Philof. Tranf. has proved by experiment, that the same kind of gas will render iron soluble in pure water, without any other addition; whence it may be concluded with fufficient probability, that a part at least of the iron in the chalybeate waters of Spa and Pyrmont, which are known to abound with fixable air, is held in folution by means of this principle. This supposition well explains several of the phenomena of these waters; as particularly their speedy decomposition in the open air, when this very volatile principle is distipated. The above conclusions are confirmed by Dr. Brownrigg's experimental inquiry into the nature of the elastic spirit contained in the Pouhon water and other acidulæ. Philos. Trans. Vol. LV. and LXIV. In order to detect the presence of this matter in waters, some of the water must be boiled in a close vessel, from which a tube is carried, having a bladder tied to its extremity. The air expelled by the heat will enter the bladder, and may then be examined by the proper tests of fixable air. Vid. Aer fixus.

by its raifing an effervescence with spirit of salt. 2. The species of acid is distinguished, by adding to the exsiccated mass a little oil of vitriol: if the acid is the marine, it will be expelled in white, and if the nitrous, in red vapours; but if it is the vitriolic, no change will ensue. The marine acid may likewise be known, by the compound enabling pure aqua fortis to dissolve gold leaf, or a mark made with gold on a touchstone; the nitrous, by its deslagrating, when ignited, on the contact of any inslammable matter; the vitriolic, by its precipitating any solution of calcareous earth, as of chalk in aqua fortis, and the precipitate being a selenites, or not dissoluble in fresh aqua fortis. 3. The basis, or substance combined with the acid in the saline compound, is found, by adding to a solution of the matter, a little solution of salt of tartar, or any other fixt alkaline salt: if the basis is a fixt alkali, no change will ensue; if an earth, it will precipitate; if a volatile alkali, a pungent smell will discover it.

The purest of the common waters is that of snow, carefully collected on the tops of mountains, or on an open plain. A gallon, slowly evaporated or distilled, leaves only two or three grains of solid matter. Distilled water itself leaves nearly as much, upon a second, and upon repeated distillations; but with this difference, that the residuum of snow water, like that of all the other natural ones, is brownish and saline; whereas, that of the distilled is a fine white earth, void of saline matter, partly calcareous, and partly indissoluble. Snow water, kept in a warm place, in clean glass vessels, not closely stopt but covered from dust, &c. becomes in time putrid; though in well-stopt bottles it remains unaltered; I have seen some, which after keeping for many years was perfectly clear and tasteless. Distilled water suffers no alteration in either circumstance. The saline matter of snow water is commonly of the nitrous kind, composed of the acid of nitre united with calcareous earth.

The next in purity is rain water, collected with the same precautions as the foregoing, after the rain has continued for some time, so as to clear the air from insects or other light bodies that may float in it. Neither this water nor the preceding discover any heterogeneity on the common trials with acids, alkalies, soap, blue vegetable juices, or metallic solutions, till great part of the aqueous fluid has been separated by evaporation. Evaporated to dryness, it leaves four or five grains of solid matter

on the gallon. Its falt is often nitrous, and its earth in great part calcareous.

The water of limpid rivers stands next in purity; and proves, though not equally with the two preceding, yet sufficiently soft, and fit for all the purposes of life. Rivers are for the most part purer and softer than the springs from which they are supplied; at a distance from, than near to the source; when their course is rapid, than when slow.

Of spring waters, there are some, which approach in purity to that of rain; but the greater number are of all waters the hardest and most impure. Some, even of those which the eye and palate judge to be good waters, contain above an hundred grains of solid matter on the gallon. The saline part of these waters is most commonly nitrous (a) or muriatic, that is, composed of the nitrous or marine acids united with earths: on adding to them, by little and little, a solution of any alkaline salt, the liquor becomes turbid and milky, more and more, till the acid, completely neutralized by the alkali, parts with all the earth, which on standing settles to the bottom. The water thus corrected, though really no purer than at first, is sound perfectly soft for economical uses, and much less, if at all, detrimental to health; its pungent, austere, earthy salt, being now converted into a mild neutral one.

River waters generally putrefy fooner than those of springs: during the putrefaction, they throw off a part of their heterogeneous matter, and at length become sweet again and purer than at first. Hard waters are remarkably indisposed to corrupt, and even preserve putrescible substances for a considerable length of time: hence, as Dr. Home observes, they seem to be best fitted for keeping at sea, especially as they are so easily softened by a little alkaline salt.

The purest waters soonest freeze; hence ice is purer than the water that remains unfrozen: on this principle, vinous and some saline liquors may be freed from a part of their superfluous water by gentle congelation. Ice, exposed to the open air, loses of its weight, its superficial parts being dissolved or abraded by the motion of the atmosphere. This property of ice was known to Hippocrates; who, imagining not the ice in its whole substance, but some of its finer and lighter parts to be dissipated, was hence led unjustly to condemn both melted ice and snow as the most impure of all waters.

⁽a) Home, experiments on bleaching. Marggraf, mem. de l'acad. des scienc. de Berlin, anno 1751.

WITH regard to the medicinal powers of pure water, little more can be faid, than what is too obvious to require being mentioned. Simple fluidity; universal innocence, or the absence of every quality that can offend the tenderest organ; miscibility with all the animal juices, in a state of perfect health, except sat; unstracts to dilute or mingle with them when greatly thickened, as in some diseases; a disposition to pass off by the cutaneous pores, more speedily and more plentifully than by the kidneys, in consequence perhaps of its total want of irritation; make the principal part of its medical character. To which may be added, that it is the most commodious medium for applying to the human body the powerful agents, heat and cold; of which the one expands and relaxes, the other contracts and constringes, all the sluid and soft parts of the animal machine.

AQUÆ MEDICINALES.

THE medicinal or mineral waters participate more or less of the earthy and saline substances found in common waters; with generally some prevailing ingredient from which they have received their names.

I. AQUÆ ALKALINÆ. Alkaline waters, as that of Tilbury.

The waters of this kind are impregnated chiefly with the mineral alkaline falt, and with calcareous earth; both which readily discover themfelves in the residuum left upon evaporation.

The Tilbury water, one of the strongest perhaps of this class, has been found serviceable, not only in complaints arising from acidities in the first passages, but likewise in obstinate alvine fluxes, some cutaneous defedations, semale weaknesses, and other disorders from a laxity and debility of the fibres: it generally passes off freely by urine or perspiration, and sometimes, on first taking, purges a little. It may be drank to the quantity of a quart a day, or more; cold, or just warmed; by itself, or with the addition, if requisite, of milk, with which it persectly agrees. It does not bear evaporation, or a boiling heat; soon growing milky, and depositing a part of its earth and virtue. In close vessels, it keeps well: some that had been carried to the coast of Guinea and brought back again, appeared to me unaltered in all its properties.

2. AQUÆ CATHARTICÆ AMARÆ. Bitter purging waters.

These waters are distinguished by their bitter taste; and by their depositing, on the addition of alkaline salt, a copious white earth, great part of which is found to be magnesia. The dry matter, left upon evaporating them, consists of the salt catharticus amarus, intermixed with different earths, and often a small proportion of other saline matters. The quantity of salt differs in different waters; some yielding scarcely two drams, and others an ounce or more on the gallon. One of the strongest of the purging waters in this kingdom is that of Jessop's Well near Kingston, of which a pint left on evaporation seventy-two grains; Kilburn and Cheltenham waters yielded sixty-four grains; Acton, not sifty; and that of Epsom, less than forty.

These waters are mild and gentle purgatives, operating with sufficient efficacy, yet in general with ease and safety; rarely occasioning any gripes, nausea, or lowness. They may likewise be so managed, as to promote the animal secretions in general, and prove excellent aperients and attenuants in sundry chronical disorders.

The dose of these waters, as a purgative, is from one to three pints, according to their strength, to be drank by a little at a time. Their virtue may be increased, by dissolving in the water some of the purished salt, or other purgative saline substances, as the artificial salt of Glauber, soluble tartar, Rochelle salt, or manna: additions of this kind are more eligible than boiling down the water, as its strength is augmented in a more certain ratio, and its natural constitution preserved entire. To render the liquor more acceptable to the palate and stomach, some gratefull distilled water, or aromatic tincture, as the tinctura cardamomi, may be added: it has been customary to insuse or boil in the water some aromatic seeds, as those of caraway: but very little of the virtue of the seeds is by this treatment communicated to the water. Their taste is excellently covered by honey. As alterants, the water may be used for common drink; diluted with simple water, milk, whey, wine, or other liquids, so as but just to keep the belly open.

These waters are found to purge more in their natural state, than after they have been boiled, and than the salt obtainable from an equal quantity of them.—They contain, besides the purging salt, no small proportion of calcareous earth. Now, if a solution of calcareous earth,

made either in pure water or in acids, be mixed with a folution of the purging falt, and the liquor evaporated; great part of the falt will be destroyed, its acid being transferred from its own earth into the calcareous earth, and forming, with this, a concrete neither purgative nor dissoluble, namely selenites. As such a concrete is found in the dry residua of the purging waters, we may presume, that it owes its origin, as in this experiment, to the destruction of a part of the purgative ingredient; and that the water holds naturally a greater quantity of falt, than can be extracted from it by art.

3. AQUA MARINA. Sea water.

SEA WATER contains, besides the common alimentary salt, a portion of bitter purging salt, similar to that of the foregoing waters, and which remains dissolved after the common salt has crystallized. After the purging salt also has been separated, there remains a small portion of a pungent saline liquor, which refuses to crystallize, and which appears to be a solution of calcareous earth in the marine acid. The quantity of salt in different seas varies, according to the greater or less evaporation, and accession of fresh water, from about one fiftieth to near one twentieth of the weight of the water.

Sea water has lately come into esteem, against strumous swellings and obstructions of the glands, and different cutaneous soulnesses. Dr. Russel observes, that in the inflammatory state of glandular swellings it is improper; that where the tumor tends to suppuration, it does no good till the pus is discharged; that in other circumstances it is a remedy of great service, whether for resolving the tumor, or preventing a fresh fluxion upon the part; and that it is useful in disorders of the internal glandulous parts, as those of the mesentery, the liver, &c. as well as in those of the external. It has been given also in the true marine scurvy, and found to promote the cure; though incapable of conquering the disease without assistances from the vegetable kingdom.

The dose of sea water is from half a pint to a pint, which may be repeated every morning for some months. In these doses, it gently purges the belly, promotes also the other excretions, and somewhat warms and strengthens the habit: in large ones, it excites vomiting. In many cases, bathing in the water is advantageously joined; both as a general corroborant, and as a topical discutient and antiseptic.

This

This water, at first taking, is apt to occasion great drought; an inconvenience which is seldom much complained of after its use has been continued for some time, and which may in good measure be palliated by sleeping immediately after it is drank. It is apt likewise, in some constitutions, to produce immoderate heat; and even when used only externally, an uneasiness and itching of the skin: it is therefore to be refrained from in all inflammatory cases, and in habits prone to phlogoses. Among all the common saline bodies, to heat and to dry seem to be qualities peculiar to the marine salt.

4. AQUÆ CHALYBEATÆ. Chalybeate or steel waters; as those of Spa, Pyrmont, Tunbridge, Islington, &c.

THESE waters discover their being impregnated with iron, by striking a blue colour with a solution of fixt alkaline salts that have been calcined with animal coals, or with a tincture made by digesting the pigment called Prussian blue in volatile alkaline spirits: this last preparation is preferable to the other, as it may be saturated more completely with that matter which tinges dissolved iron blue.

Iron in waters is discovered also by the purplish, bluish, or blackish colour, which they assume on the addition of certain vegetable astringents, among which powdered galls are the most eligible. This last method of trial, which is that commonly made use of, distinguishes as minute proportions of iron as the first, provided the liquor contains no more acid than is sufficient to keep the metal dissolved; but if the quantity of acid is very large, astringents give no notice of the iron, whereas the tincture of Prussian blue discovers it universally.

Chalybeate waters appear to differ from one another, not only in the

degree, but in the species, of their impregnation.

Some resemble a solution of vitriol of iron made in common spring water. Like that solution, they strike a blue or black colour with galls, deposite on standing some of their iron in an ochery form, but retain great part of it for a length of time, and yield on evaporation a saline matter, which communicates a ferrugineous impregnation to fresh water, and which appears to be a true vitriol of iron. From some waters, as that of Hartfell in Scotland, the vitriol has been crystallized (a) in its proper form.

Others refemble a folution of the same vitriol with an admixture of natron or the mineral fixt alkaline falt. Like fuch a mixture (a), they strike, when fresh, a purple colour with galls, deposite the whole of their iron in a very little time, and yield on evaporation, not a vitriol or chalybeate falt, but a falt composed of the acid of vitriol and alkali: fome, as that of Geronsterre at Spa, yield also a little pure alkali (b), besides what is satiated with acid. It may be observed, that in artificial mixtures of alkalies with folutions of vitriol, or other metallic folutions made in acids, (and possibly something of the same kind may obtain also in the natural,) if the veffel is immediately stopt, so as to have no vacuity after the instant of mixture; the acid and the alkali have no action on one another fo long as they are kept confined, that is, fo long as the extrication of air, the common concomitant of their mutual action, is prevented: but as foon as the veffel is opened, or the contained air has an opportunity of escaping, the alkali begins to absorb the acid, a sparkling or effervescence ensues, greater or less in proportion to the quantities of the two, and the metal, thus divested of its acid folvent, precipitates(c).

It is not however to be prefumed, that the speedy separation of the ferrugineous matter of waters is owing universally to an alkaline precipitant. Solutions of pure vitriol in pure water deposite a part of their iron spontaneously; and if the solutions be so far diluted, as to strike with astringents a colour little more than perceptible, they will lose so much in a few hours as to exhibit with the same astringents no tinge at all.

In general, a blue or black colour produced with galls may be looked upon as a mark of the absence of alkaline salt; and a purple, as a mark that either the water originally contained an alkali, or has become alkalescent or verging to putresaction by standing. On the same principle, a degree of alkalescence, or of tendency to corruption, in common waters, very far too minute to be sensible on any other known trial, may be made conspicuous; viz. by the water, when impregnated with a little vitriol, as a grain or two to a pint, striking a purple colour with galls.

⁽a) That an alkaline addition is necessary, to make solutions of vitriol strike with galls the purple colour that chalybeate waters do, is a discovery of Mr. Reynolds, exper. on a chalybeate water near Bromley in Kent.

⁽b) Rutty, synops. 323. (c) A discovery of Mr. Scheffer, Suenska vetensk. acad. bandl. 1753.

The spontaneous separation of the iron, which happens in many of the chalybeate waters, and which, though it may be retarded, cannot be prevented by any care in stopping the bottles, after the waters have been once exposed to the air, renders them unsit for long keeping or carriage. A small addition of any acid prevents the separation, even in those whose virtues are naturally the most fugitive: it is suspected, that the chalybeates brought from Germany have commonly this artificial impregnation.

These waters are used, like other chalybeates, in debilities and laxities of the stomach, chylopoietic organs, and of the viscera in general; in decays of constitution; in cachectic, chlorotic, and other like indispositions. Where they pass freely, they are accounted more invigorating than the artificial preparations of iron, and less liable to disorder particular constitutions: many of them however are more apt to fail of taking due effect, on account perhaps of the acid solvent being more disposed to quit the metal. Some of them are rarely observed, and some scarcely ever, to give any black tinge to the seces, though drank in large quantity; a phenomenon which may perhaps be ascribed to their depositing their iron in the first passages in an indissoluble and inactive state, rather than to the cause which some have assigned, their carrying

5. AQUÆ CUPREÆ. Cupreous waters; as those of Neusol in Hungary, and Wicklow in Ireland.

it entire into the blood. They are taken to the quantity of two or three pints or more in a day, divided into different doses; and require the

fame caution in their use as the artificial chalybeates.

THESE waters, which are little other than a folution of vitriol of copper, and those which contain a much smaller proportion of that metal blended with vitriol of iron and other ingredients, betray their cupreous impregnation, by staining a polished iron, immersed in them, of a copper colour, and by striking a blue with volatile alkaline spirits. Some of them have been used, like other venereal solutions, as external detergents. Some, more slightly impregnated with the copper, have been taken internally as emetics, purgatives, and deobstruents; a practice which appears much too hazardous to be followed.

ALL the mineral waters we know of, are impregnated with more or fewer of the foregoing ingredients, combined in various proportions. The hot waters called thermæ or baths, have not, as fuch, any peculiar impregnation; their heat depending, not upon an intrinsic, but an external cause: the hot springs of Toplitz in Germany appear, from Hoffman's experiments upon them, to be no other than fimple water. Of the waters called fulphureous, or those which have a fetid smell refembling that of fulphureous folutions, there are feveral which do not feem to contain any actual fulphur: nor is there any actual fulphur in the extremely fetid and diffusive vapour, which arises from folutions of fulphur itself during their precipitation with acids. Analogous to this, perhaps, is the fulphureous impregnation of certain waters. The nature and medicinal effects of this subtile volatile principle are little known; the fulphureous waters containing, at the fame time, other active ingredients. There are likewise waters which appear to be impregnated with fulphur in its whole fubstance, and which may therefore be prefumed to participate of the virtues of the artificial folutions of that concrete. - For the analyses and uses of particular waters, the reader may confult Dr. Rutty's synopsis.

AQUILEGIA.

AQUILEGIA Pharm. Parif. Aquileia. Aquilina. Aquilegia flore simplici J. B. Aquilegia vulgaris Linn. Columbine: a plant with slender reddish stalks, and bluish green leaves, in shape somewhat roundish, with several slight indentations, and one or two deep ones. The slower, commonly blue, sometimes red or white, consists of sive irregular petala, each of which is supposed to resemble a slying eagle or pigeon (aquila or columba) whence the names of the plant: the slower is sollowed by sive pods, full of shining black oval seeds. It is perennial, grows wild in woods, and slowers in June.

THE feeds of columbine have been commended, in substance and in emulsion, as an anthelmintic, as an aperient in the jaundice, and for promoting the eruption of the measles and small pox. Their sensible qualities afford little foundation for these kinds of virtues, as they do not seem to differ materially from those of the cold seeds so called; the columbine

columbine feeds being only fomewhat more mucilaginous, and accompanied with fomewhat of a difagreeable relish.

The virtues ascribed to a tincture of the flowers, as an antiphlogistic, and for strengthening the gums and deterging scorbutic ulcerations in the mouth, appear to be better founded; the tincture being made with an addition of the vitriolic acid, and differing little from our officinal tincture of roses. The flowers themselves, and the conserve and distilled water of them directed in some foreign pharmacopæias, are insignificant.

ARANEARUM TELÆ.

ARANEARUM TELÆ Pharm. Edinb. (a) Cobwebs. These are applied by the common people for stopping the bleeding of wounds; which they effect, not by any styptic power, but by adhering to the part, and closing the orifices of the vessels.

ARGENTINA.

ARGENTINA, Potentilla, Anserina, Pharm. Edinb. (b) Pentaphylloides argenteum alatum seu potentilla Tourn. Potentilla anserina Linn.
Silverweed or wild tansy: a low creeping plant, with winged
leaves, composed of seven or eight pair of oblong indented segments set
along a middle rib, with smaller portions between, green above, and
covered with a silver-coloured down underneath; the slowers, which rise
on long pedicles in the bosoms of the leaves, are composed, each, of sive
gold-coloured petals with a number of threads in the middle, and sollowed by a small cluster of naked seeds. It is perennial; common by the
sides of rivulets and in moist uncultivated places; and slowers in June.

The leaves of argentina have been generally looked upon as strong astringents, and recommended as such in sluxes and hamorrhages. That they have an astringent quality is manifest to the taste, and from their striking a black colour with solutions of chalybeate vitriol: but in the leaves in substance, whether fresh or dry, and in their insusions, decoctions, and expressed juice, the preparations which have been generally made use of, the astringency is very weak; and even the extracts made

from them, by water and by rectified spirit, in which all their active matter is concentrated, are only among the milder styptics or corroborants. The spirituous extract is stronger than the watery, and in proportionably smaller quantity.

ARGENTUM.

ARGENTUM Pharm. Lond. & Edinb. SILVER: a white metal; becoming yellow, and at length black, from the vapour of fulphureous folutions and of putrefying matters; extremely malleable; near eleven times specifically heavier than water; fusible in a bright red heat; fixt and indestructible in the fire; soluble, in the nitrous acid, into a limpid liquor, which stains the solid parts of animals black; not soluble, by moderate digestion, in the marine or vitriolic acid; precipitable by these acids from its solution in the nitrous.

The greatest quantities of this metal are found in the mines of Chili and Peru; commonly in small grains and silaments, embedded in different earths and stones; from which it is separated by pulverization, ablution with water, and amalgamation with mercury. Several mines in England, Germany, and other parts of Europe, afford silver; rarely native, or in distinguishable masses; commonly reduced to a state of ore, of a red, or of a yellow or brown, or of a dusky leaden or black colour, by an intimate admixture of arsenic, or sulphur, or both; from which it cannot be extracted by quick silver, but which are dissipated by calcination, so as to leave the silver separable from the remaining earth by suspenses.

CRUDE filver, however comminuted or attenuated, has not been obferved to produce any medical effect; though abundance of virtues were afcribed to it by the credulity of former times. It is not foluble in any of the fluids of the animal or vegetable kingdom.

It disfolves, by the assistance of a moderate heat, in about twice its weight of pure aqua fortis: the solution, duly exhaled and set in the cold, crystallizes into thin colourless transparent plates. The crystals, or the dry matter left upon inspissating the solution, melt in a moderate fire, and on cooling form a dark coloured caustic mass. This preparation is in common use for consuming warts and callosities; but is less fit for

Causticum lunare Ph. Lond.&Edin.

fuch

fuch purposes as require a considerable quantity to be applied, as the laying open of imposthumations, being apt to liquefy by the moisture of the skin, and spread beyond the limits in which it is intended to operate. For the greater conveniency of using, it is cast into oblong slender pieces, either in iron pipes heated and greafed, or in holes made by fome fmooth instrument in a lump of tempered tobacco-pipe clay: each piece is wiped clean, and wrapt in dry foft paper. The matter is to be poured out as foon as it flows thin: if kept a little too long in fusion, it becomes too thick to run into the mould, and parts with fo much of its acid as not to be fufficiently corrofive: by a longer continuance of the fire, all the

acid is gradually diffipated, and a lump of pure filver remains.

A preparation fomewhat less caustic than the foregoing, is recommended internally by Angelus Sala, Boyle, and others, as an anthelmintic, and as a purgative in hydropic and inveterate ulcerous difeases. For this purpose, the crystals of filver are dissolved in water, and mingled with a folution of equal their weight of nitre: this mixture is evaporated to dryness, and the residuum calcined with a gentle heat, just not sufficient to melt it, and kept continually stirring, till no more fumes arise. Boerhaave affures us, that two grains of this preparation, made into pills Pilula with crumb of bread and a little fugar, and taken on an empty flomach, fome warm water fweetened with a little honey being drank immediately after, purge gently without griping, and bring away a large quantity of water almost without the patient's perceiving it. He nevertheless cautions against the too liberal or continued use of this medicine, and obferves, that by its corrofive quality it weakens the bowels, particularly the stomach, and that therefore proper corroborants, as rob of juniper berries, ought to be interposed. Even with this assistance, however, it is at best a dangerous medicine, and as fuch deservedly stands excluded from practice.

ARGENTUM VIVUM.

ARGENTUM VIVUM Pharm. Lond. Hydrargyrus, Argentum vivum, Mercurius, Pharm. Edinb. MERCURY OF QUICKSILVER: an opake filver-coloured metallic fluid, appearing to the eye like melted lead or tin; about fourteen times heavier than an equal bulk of water;

not congealable by the greatest known degree of natural cold (a); totally exhaling, by a heat below ignition, in subtile sumes, which condense into running mercury again.

Quickfilver is fometimes found in the earth in its fluid form, and is then called virgin mercury; but for the most part it is intimately blended with sulphur or earthy matters into a state of ore. The sulphureous ores are of a more or less beautiful red colour; the earthy or stony ones, grey, yellowish, brown, leaden coloured, &c. From these last, the metallic sluid is extracted by simple distillation: the sulphureous require an addition of quicklime, iron silings, or some other substance that may absorb and keep down the sulphur, which otherwise would rise in conjunction with the mercury. The principal mines of quicksilver, of which we have any account, are in Spain, Hungary, and the province of Friuli in the Venetian territories: considerable quantities are brought also from the East Indies.

This fluid, supposed by the Greeks to be poisonous and corrosive, was introduced into medicine by the Arabians, as an ingredient in external applications, against different cutaneous maladies. This practice was followed by some physicians in Europe towards the end of the thirteenth century, but was not established, or looked upon in general to be safe, till about the beginning of the sixteenth, when the venereal distemper, then lately received from America, was found to yield to mercurial applications alone; and now also the internal use of mercury began to be ventured on, in this and in other diseases.

Pure mercury has no perceptible acrimony, or taste, or smell: there are examples of its having been lodged, for years, in cavities both of the bones and of the slessly parts, without having injured or affected them (b). Taken into the stomach in the quantity of an ounce or two, it soon passes through the intestinal tube, unchanged, and unfelt: hence some have been induced to give a pound or more in violent constipations,

⁽a) It is faid, that in fome late experiments made at Petersburg, with very intense degrees of artificial cold, (produced by mixing snow and spirit of nitre separately brought to great coldness,) pure mercury congealed into a filver-like malleable metal, which quickly melted again on an abatement of the cold; and that in Fahrenheit's thermometer, it sunk, before its congelation, to between three and sour hundred divisions below 0; that is, about as far below the point at which water freezes, as the heat, in which tin melts, is above it.

hoping that this innocent fluid, by its great weight and slipperiness, would force open obstructions, that had resisted the common methods of cure by purgatives, relaxants, and emollients. This practice, so far as I can learn, has not been attended with any remarkable success; nor do the principles, on which mercury has been given in these cases, appear to be just. The slipperiness of this sluid consists only in the mobility of its own parts, not in any power by which it can lubricate the vessels of an animal. Its weight can be of no use, unless where the obstruction lies in some descending part of the tube: and even supposing it to act perpendicularly, to the greatest advantage, there is room to fear, that the pressure of a pound or two will rather distend the superiour part of the intestine, than be able to force a passage through the obstinate obstructions against which it is recommended.

When mercury is refolved into fume, or altered in its form by fire, or combined with a small portion of mineral acids, or otherwise divided into minute particles and prevented from reuniting by the interposition of proper substances; it operates with great power, and extends its action through the whole habit. In these forms, whether taken internally, or introduced into the blood from external application, it seems to liquefy all the juices of the body, and may be so managed as to promote excretion through all the emunctories. If its power is not restrained, or determined by additions, it tends chiefly to affect the mouth; and having sused the humours in the remoter vessels, occasions a plentiful evacuation of them from the salival glands, with considerable swellings, inflammations, and ulcerations of the parts. The salivation is accompanied with a diminution of most of the other discharges, and an increase of these diminishes the salival flux.

The falutary effects of mercurials have, in many cases, very little dependence on the quantity of sensible evacuation. Venereal maladies, and chronical distempers proceeding from a viscidity of the humours and obstruction of the small vessels, are often successfully cured by mercurials taken in such doses as not to produce any remarkable discharge; especially if affisted by diaphoretics and a warm diluent regimen. In this view, camphor, and the resin or extract of guaiacum, are frequently joined to the mercury; and to the more active preparations, a little opium; which not only promotes the diaphoresis, but prevents the mercury from irritating the first passages and running off by the grosser emunctories.

0 2

This appears to be, in general, the most advantageous method of using mercurials; excepting, perhaps, only in venereal maladies of long standing, or such as have arisen to a great height, or have affected the bones; which demand, for the most part, a full ptyalism. In these cases, the disease has been subdued for a time by the alterative method, but afterwards broken out afresh, and been completely cured by salivation: and, on the other hand, some cutaneous soulnesses, after resisting salivation, have yielded to an alterative course.

Though mercurials are found to be falutary in fundry cutaneous defedations, and impurities of the blood and juices vulgarly called fcorbutic; they are always pernicious in the true fcurvy, and dangerous in conftitutions inclining to this difease, where the humours are acrimonious, and colliquated, and disposed to a putrescent state. In such circumstances, mercurial medicines are apt to operate with violence: small doses have occasioned high and lasting salivations. The removal of these accidents is to be attempted by glysters, purgatives, diaphoretics, or such other means, consistent with the patient's strength and the particular symptoms, as may procure a speedy revulsion from the salival ducts.

A long continued use of mercury is in no case free from danger, as it manifestly colliquates the whole mass of blood, and tends to weaken the nerves, so as to bring on tremors and paralyses. The miners, and those who are exposed to the sumes of mercury in extracting it from the ore, are said to be almost always, sooner or later, seized with these kinds of complaints; to become generally in a few years paralytic, and at last to die hectic.

Mercurials are destructive to insects, perhaps of every kind. They are sometimes given internally against worms; and sometimes applied externally, in unguents, for destroying cutaneous animalcula. The itch, now reckoned an animalcular disease, is sometimes cured by mercurial unguents; which, nevertheless, cannot be depended on for this effect, unless in slight cases; as their antipsoric efficacy seems to reach no farther than those parts of the skin to which they are applied, and as they cannot with safety be applied freely, to any great extent of the body, particularly of the trunk.

Mercury has been of late recommended as an effectual antidote against the poison of the mad dog. Several cases are related, by Dr. James, Desault, and Du Choisel, both of brutes and human subjects, bitten by

mad

mad dogs, being preserved from the usual consequences of this bite, by mercurial unctions, and mercurials taken internally. There are some instances given also of a cure being obtained, by the same means, after symptoms of madness had appeared (a).

This fluid diffolves, by the affistance of trituration or heat, most metallic bodies; retaining its own natural colour, but having its confistence increased in proportion to the quantity of the metal: iron is the only one of the common metals, to which it will not easily adhere (b). Bismuth unites with it more intimately (c) than any other metallic body, and remarkably promotes the union of lead with it: mercury, impregnated with a little bismuth, was found to dissolve considerable masses of lead, in a heat no greater than that of the human body. Cases sometimes happen, in which the surgeon may probably avail himself of this property.

From most of the sluid amalgams, or mixtures of mercury with metals, great part of the quicksilver may be separated by pressure through leather; but bismuth, and mixtures of bismuth with lead, are so intimately dissolved, as to pass, in considerable quantity, through the leather with it; and hence, with these metals, it has been frequently adulterated. This abuse may be discovered, by the mercury contracting a dull coloured skin upon the surface, or staining paper blackish; by its not running freely into round globules, but forming tears or vermicular striæ; by its leaving, upon evaporating a little of it, a powdery matter or a coloured spot on the bottom of the vessel; and by its producing a turbid milkiness during its dissolution in aqua fortis.

Quickfilver is commonly purified from these and other like admixtures, by distilling it in a glass or rather iron retort, or in an iron pot, with a head made of one piece, the vessel commonly used for this purpose

⁽a) James, treatise on canine madness.

⁽b) The other metallic bodies, with which it cannot be united by the usual methods, are regulus of arsenic, and two lately discovered metals called nickel and regulus of cobalt. See Neumann's chemical works, p. 152, &c.

⁽c) Though part of the bismuth separates spontaneously from the mercury, it does not follow, as some have concluded, that the union is imperfect or superficial, but that the mercury can retain only a certain quantity of bismuth; and a certain quantity I have sound it to retain after long continued agitation.

Argent. viv. purif. Phar. Lond.

by the refiners, and afterwards washing it with vinegar, or with common falt and water. The chemists, suspecting that some metallic bodies (a) may be carried up by the mercury in distillation, recommend certain additions, particularly fulphur; which, from its rendering the mercury itself less disposed to arise, may be presumed to have this effect in a greater degree on the metals that are naturally not volatile: they fublime the mercury and fulphur together into cinnabar, which fee in the fequel of this article, and adding to this compound fome iron filings to abforb the fulphur, distil off the mercury supposed now to be completely pure. Mr. Malouin recommends uniting the quickfilver with crude antimony instead of fulphur, by leifurely pouring the mercury heated into an equal quantity of antimony made fluid by fire, and then separating the mercury, as from cinnabar, by distillation with iron: the antimony, he fays, by virtue of its reguline parts, detains the other metallic bodies more effectually than fulphur can do, and the mercury is thus brought to a state of purity even greater than that of the animated mercuries of the alchemists; for the processes, by which they imagined it to be animated or exalted in its powers, appear to have done no more than to purify it to a certain degree. He has not, however, communicated the particular facts on which this affertion is built; or given any experimental proofs of the greater purity of mercury distilled from antimony, than of such as has been revived from factitious cinnabar, or even of fuch as has been carefully diffilled without addition.

QUICKSILVER, triturated with powdery or with thick uncluous matters, is gradually divided, and incorporated with them into one uniform compound, in which no particle of the mercury can be distinguished by the eye. It is most difficultly mixed with earthy powders, most easily with thick balsams and mucilages.

Killed or extinguished, that is, ground till the mercurial globules disappear, with one twelfth its weight or more of Venice turpentine, or half that quantity of balsam of sulphur, it is mingled with plasters; which, for this purpose, are to be melted, and taken from the fire, before the mercury is stirred in. The college of London directs the mercury, killed with balsam of sulphur, to be mixed with four times its

quantity of the common plaster+, or of gum ammoniacum 1: that of + Emplastr. Edinburgh uses a pound and a half of common plaster, and two ounces and a half of Venice turpentine, to eight ounces of the quickfilver | . These compositions are applied as resolvents and discutients, against venereal pains, and indurations of the glands; the mercury exerting itself ammon. c. in some degree upon the part, though it is rarely introduced into the Lond. blood in such quantity as to affect the mouth. Astruc observes, that even || Emplastr. by covering all the limbs with mercurial plasters, the method once Ph. Edinb. practifed for raifing falivations, it was difficult to obtain a complete and effectual ptyalism (a).

cum mercurio Ph. 1 Emp. ex merc. Phar. mercuriale

Substances of less consistence, as ointments, leave the mercury at more liberty to act; and are generally and deservedly preferred to the plasters, in the intention of topical refolvents, &c. as well as in that of conveying the mercury into the habit. A dram of quickfilver mixed with unguents, well rubbed into the skin, and repeated every day, or rather every other day, generally produces, foon after the third application, and fometimes after the fecond, appearances of inflammation in the mouth, which are followed by a free and copious ptyalism: those employed in rubbing the ointments on others, have been falivated by the mercury imbibed through the palms. The ptyalism raised by unction is often more effectual, and accompanied with fewer inconveniencies, than that produced by mercurials taken internally; which last are apt, in some constitutions, to run off by the intestines, without affecting the salival glands; and in others, to affect the mouth fo hastily, as to excite a copious falivation without extending their action fufficiently to the remoter parts. The mercurial ointments are commonly prepared, by rubbing the mercury with lard or other fat matters of a due confiftence: three parts of hog's lard, and one of mutton fuet, make a commodious basis, with which may be mixed one part or more of mercury +. As a + Ung. mergood deal of labour is required for thus uniting the quickfilver with fimple fats, fome are accustomed to previously extinguish it with a little turpentine tor balfam of fulphur ||, after which it is mixed, more eafily, tung. cawith twice | or four times ! its weight of lard; these additions, however, particularly the turpentines, are, in this form, accompanied with || Ung. caan inconvenience; being apt, by frequent rubbing, to fret the skin-

rul. mitius Ph. Lond. rul. fortius Ph: Lond.

Merc. alkalazat. vulgo.

Pil. mercurial. Pharm. Lond.

Pil. mercurial. Pharm. Edinb. Mercury is divided also, with different materials, for internal use, and given, as an alterative and as an anthelmintic, from two or three grains to eight or more. Half a dram of quicksilver is ground, for example, with two scruples and a half of prepared crabs-eyes. Sometimes also purgative materials are joined: in this intention, the above quantity of mercury is commonly killed with twelve grains of Strasburg turpentine, and then beaten up with eight grains of the cathartic extract, and fix of powdered rhubarb: the pills called Belloste's, are supposed to be a composition of this kind. The Edinburgh college, in their last dispensatory, have rejected all these compound forms, and direct only a mass of pills made of quicksilver divided by equal its weight of honey, and afterwards beat up with double the weight of bread-crum and a sufficient quantity of water.

A treatife has been published by Mr. Plenck of Vienna, recommending a mixture of quickfilver and gum-arabic as preferable to all the other mercurial preparations hitherto known; being in all cases safe, rarely or never producing a falivation, and acting foon on the venereal virus. He directs one dram of purified quickfilver, and two drams of the gum, to be ground together in a stone mortar, adding by degrees half a spoonful of water, till the quickfilver disappears, and the whole becomes a viscid grey mucilage, which happens in a short time: half an ounce of fyrup, and eight ounces of a fimple water, are then gradually added, and two spoonfuls of the mixture given every morning and evening. Part of the mercury remains suspended in the liquor, part fettles to the bottom, but retains fo much of the gum as to continue divided and form a grey mucous fediment, which readily unites again with the water on shaking the vessel: if the sluid be separated, and the fediment dried by heat, the mucilage then lofes its power, and the mercury runs into globules.

The author made trial of several other substances, both animal and vegetable, for the extinction of mercury, but found none that answered equally with gum-arabic. The mucus expectorated from the throat extinguished it, and kept it divided after the addition of water; but very little of the mercury remained suspended in the liquor, both the mercury and the mucus subsiding. Saliva had much less effect: yolks of eggs, white of eggs, bile, glue of isinglass, gum-tragacanth, mucilage of quince seeds, clarified honey, simple syrup, either did not extinguish the

mercury

mercury at all, or fuffered it, on dilution with water, to run together again. But thick honey, unclarified, kept it divided; and fyrups greatly

promoted the effect of gum-arabic.

The trials of the gummy mercurial that have come to my knowledge, afford little foundation to expect from it any advantages above the common mercurial preparations. I have known it given, without much benefit, in cases which were afterwards cured in a short time by solution of fublimate. I have been informed, that in some of our hospitals even fmaller doses than those directed by the author have on the third day brought on a ptyalism. Dr. Baldinger, in his remarks on the last Edinburgh pharmacopæia, recommends this folution as an efficacious anthelmintic, especially against the lumbrici.

Mercury, thus fimply divided by thefe or other like matters, feems to operate more mildly, not only in the first passages, but after it has been received into the blood, than when combined with mineral acids, or reduced by fire into the form of a calx. At the same time, however, it is more uncertain, or more liable to fail of taking full effect; on account, perhaps, of the substances, by which its particles are disunited, being foon fubdued and separated by the digestive powers, so as to leave part of the mercury to run together again and pass off inactive through the intestines. How easily the union is dissoluble, may be judged from hence, that when mercury is perfectly mixed with turpentine, if the mixture be beaten up with extracts and powders into a mass for pills, a confiderable part of the quickfilver, by this mechanic agitation, is often feparated and fqueezed out in globules.

MERCURY, triturated with equal its weight of fulphur, forms there- Æthiops with a greyish black powder, which grows darker coloured in keeping, mineralis Ph. Lond. or on continuing the triture, and is commonly distinguished by the name & Edinb. of athiops. This compound is one of the mildest of the mercurial preparations; the mercury being far less active in mixture with fulphur, than with any other known species of matter.

The union of the quickfilver with the fulphur, effected by triture, at least by such a degree of triture as the shops are accustomed to bestow upon them, is not intimate any more than that with the fubstances abovementioned. If the æthiops be rubbed on gold, a part of the mercury adheres to the gold fo as to make it white: on mixture with fyrups or

R

other

other like matters into the confistence of an electary or mass for pills, a part of the mercury is generally spued out. The longer the quicksilver and sulphur are ground together, the less they will be disposed to separate; a circumstance which does not appear to obtain in the mixtures of mercury with resinous or earthy bodies.

Æthiops cum igne, vulgo. A more intimate coalition of mercury and fulphur may be fpeedily effected, by melting the fulphur over a gentle fire, and gradually stirring into it the quicksilver, with care to cover the vessel if an ebullition or swelling up of the matter shews it ready to catch slame; an accident which sometimes happens when the quantities are large. This compound gives no whiteness to gold, and suffers no separation of its parts on being made into electaries or other forms. Even the acid solvents of mercury are incapable of extracting it when thus combined with sulphur; and the alkaline solvents of sulphur extract only, by long boiling or digestion, so much of that concrete as is more than sufficient to satiate the mercury.

If mixtures of quickfilver and fulphur, thus intimately united by fire, in due proportions, as twenty-four or twenty-five parts of the former to feven of the latter, be powdered and fet to fublime; the two ingredients rife together without feparation, (except that a part of the fulphur, and generally a very confiderable one, burns away in the process,) and concrete, in the upper part of the fubliming jar, into a red mass called cinnabar or vermilion. This preparation, though containing much more mercury than the æthiops, does not appear to be more active; and is by many looked upon rather as a medicine of the antispasmodic kind, than as possessing the proper virtues of mercurials. Indeed the real virtues, either of cinnabar or of æthiops, cannot perhaps be precisely assigned.

Cinnabaris factitia Ph. Lond & Edin.

When mercury is intimately combined with a certain quantity of fulphur, it feems to operate, though given in confiderable doses, as a scruple or half a dram of the compound, only in an insensible manner; and in many cases, to pass off inactive through the intestinal tube (a). It may be presumed, that an increase of the sulphur, beyond the quantity sufficient for this perfect satiation of the mercury, will not vary its action; but that a diminution of the sulphur will leave the mercury, or a part of it, in a state of more activity; analogously to what has been

⁽a) Cartheuser, rudimenta m. m. p. 481. Malouin, chim. medicinale, part. iv. ch. 34.

before observed to happen in regard to the antimonial regulus. It may be prefumed also, that in perfect cinnabar the mercury is completely faturated, and that in perfect æthiops, it is both faturated, and blended with fome redundant fulphur: but that in fome cinnabars, it is not faturated from a deficiency in the quantity of fulphur; and that in some forts of æthiops it is not faturated, from an imperfection in the mixture. There are examples both of æthiops and cinnabar, one of which, in regard to the latter, has fallen within my own knowledge, having unexpectedly produced falivations. It should seem therefore, that the æthiops made by fire is the most to be depended on, or the most certain and equal in its power, whatever this power may be, of any of the fulphurated mercurials.

When æthiops or cinnabar are thrown on a red-hot iron, their fumes are of great activity. The fumes of cinnabar are fometimes directed, not only to be received on the lower parts; but likewise to be taken into the mouth, against venereal ulcerations in the nose, mouth, and throat. Of all the ways of applying mercury, this last requires the greatest caution.

QUICKSILVER, included in a flat-bottomed glass having a small hole Mercurius open to the air, and kept for several months in a constant heat just not Ph. Lond. strong enough to make it evaporate, calcines by degrees into a red powder. A greater heat, fufficient to make the mercury freely distil, not only does not promote the calcination, but revives fuch part, as has been already calcined, into running mercury again. A weaker heat, as that of the human body, or even of boiling water, though continued for years, changes only a small part of the mercury into a blackish powder: constant triture or agitation produces similar effects to this low degree of heat, and in a much shorter time (a). If the free access of air should be found to influence the calcination of this, as it does that of the metallic bodies called imperfect, the tedious process might be expedited, by using, for the vessel, a glass tube, with both its ends bent upwards, and one of them confiderably higher than the other; through which, a constant stream of fresh air would pass over the surface of the fmall thread of mercury at the bottom.

⁽a) Boerhaave, Philosoph. transact. No. 430, 443, 444, & Mem. de l'acad. des scienc. de Paris, pour l'ann. 1734.

The red powder has been by some greatly esteemed in venereal cases, and supposed to be the most effectual and certain of the mercurials. It is accompanied with one considerable inconvenience, being greatly disposed to irritate the first passages, and occasion gripes; to prevent which, a small quantity of opium, and some warm aromatic material, are commonly joined to it: the antivenereal pills of a late celebrated empyric are supposed to have been a composition of this kind. Even when thus corrected, however, it does not appear, from what I have been able to learn of its effects, to have any advantage above the mercurials in common use and of easier preparation. The dose is from half a grain to two grains: five or six grains are said to vomit and purge violently.

Mifaubin's pills.

Pure aqua fortis, affished by a moderate heat, dissolves equal its weight or more of quicksilver into a limpid corrosive liquor; which, largely diluted with pure water, the common spring waters turning it milky and precipitating a part of the mercury, has been employed in lotions against some kinds of cutaneous defedations, and where mercurial lotions are adviseable, is perhaps one of the best of them. An ointment is likewise prepared, for venereal ulcers, &c. by mixing the corrosive solution with fats: an ounce of quicksilver is dissolved in two ounces of spirit of nitre, the solution poured hot into a pound of lard melted and just beginning to grow stiff, and the whole briskly stirred up till an uniform yellow mixture is procured.

Ung. ci-

On inspissating the mercurial solution over a gentle fire, there remains a white mass highly caustic; which, calcined with a gradual heat, becomes first brown, then yellow, and at length, on increasing the heat, of a deep red colour. If the aqua fortis, used for the dissolution, has been previously drawn over from a small proportion, the hundred and twenty-eighth part of its own weight, of sea salt, the red mass is supposed to assume more readily the sparkling appearance which is looked upon as the characteristic of its goodness. This preparation is employed as an escharotic; and mixed with ointments and cerates, as a digestive; in which intention, Mr. Sharp observes, that it is very effectual.

Merc. corrofiv. ruber, vulgo præcipit. ruber
Ph. Edinh.
Merc. corrofiv. ruber
Ph. Lond.

Sundry methods have been tried for abating the corroliveness of this preparation, so far as to render it safe for internal use. One of the most certain seems to be, digesting it two or three days, with a gentle heat, in about thrice its quantity of rectified spirit of wine, then setting fire to

the

MATERIA MEDICA.

CHARLES SHAPE

the spirit, and keeping the powder constantly stirring till all the spirit is burnt off. In this process, the corrosive is deprived of a little of its acid, Mercurius which is partly perhaps absorbed and dulcified by the spirit during the Ph. Lond. digestion, and partly dislipated by the heat during the burning. medicine, nevertheless, is still a very rough one, operating, generally, in doses of a few grains, both upwards and downwards. Different preparations of this kind have been kept as fecrets in particular hands, but Arcanum it does not appear that any of them are superiour in virtue to some other mercurials of greater fafety and more equal power.

IF oil of vitriol be poured on half, or equal its weight of quickfilver, and gradually heated till the liquor boils and diffils; the more phlegmatic parts arife, while the stronger acid corrodes the mercury into a white caustic mass. On the affusion of warm water, the mass falls into powder, and becomes immediately yellow; a part of it, fatiated with acid, diffolving in the water: the larger the quantity of acid made use of, and the less thoroughly the matter has been exficcated or calcined, the more of it will dissolve. The yellow powder, ground with fresh quantities of water till all the foluble part is extracted, becomes infipid, and Merc. emet. in this state, commonly called turpeth or turbith mineral, it proves, Hav. Pharm. though not corrofive, strongly emetic; operating, in this intention, the flav. vulgo most effectually of all the mercurials that can be given with fafety. It nerale Pb. is used chiefly in virulent gonorrheas, and other venereal cases accompanied with a great flux of humours to the parts: it is faid likewise to have been employed with fuccefs, in robust constitutions, against leprous disorders, and obstinate glandular obstructions. The dose, as an emetic, is from two grains to fix or eight; though fome constitutions, habituated to mercurials, can bear larger quantities: I knew an instance of twenty grains producing no fenfible evacuation or diffurbance. It may be given in smaller doses, as half a grain or a grain, as an alterative, after the fame manner as the red calx of mercury already mentioned: and even when intended as an evacuant, it may perhaps, as Malouin observes, be most adviseable, to give only a small quantity at a time, as one grain, and repeat this dose every hour till the vomiting succeeds.

THE marine acid has no action on mercury, unless either the mercury be previously dissolved in other acids, or the marine spirit be applied in a very concentrated state and in the form of fume.

corallinum, pulvis principis, panacea merc. rubra, &c.

turpeth. mi-

Merc. præcipitat. alb. Boerhaave. On adding a folution of fea falt to a folution of mercury made in aqua fortis, the nitrous acid quits the mercury, and unites with the alkaline basis of the sea salt; and, at the same time, the acid of the sea salt unites with the mercury, and forms with it a compound difficultly and only partially dissoluble, of which, therefore, great part subsides, on standing for some time, in form of a white powder. This powder, washed with fresh quantities of hot water, till the more soluble parts are extracted, becomes nearly insipid. In this state it is recommended by Boerhaave as one of the best of the mercurials, and said, in doses of three grains, to purge and vomit gently. It appears however too corrosive for internal use; being so much so, as to be employed by the farriers for the purposes of an escharotic. The preparation is likewise a very unfrugal one, a considerable part of the mercury remaining unprecipitated, and a considerable part of the precipitate being dissolved and carried off in the ablution.

If the dry white mass, obtained by inspissating a solution of mercury in aqua fortis, be powdered and mixed with equal its weight of dried fea falt, the mixture put into a matras or other like vessel, of which it may fill nearly one half, and fet in a fand-heat gradually increased; the same transposition of the acids will happen, as in the foregoing case, and nearly all the mercury will now be fatiated with the marine acid, and form with it a faline compound, which fubliming into the upper part of the matras, concretes into a white crystalline mass, called corrosive sublimate. If the vitriolic acid be used instead of the nitrous, that is, if the unwashed turbith be taken and mixed with fea falt, the event will be the fame; the mercury fubliming with the acid of the fea falt; while the acid, before combined with it, remains behind united with the fea falt's alkali, forming therewith a nitrum cubicum when the nitrous acid has been used, and a fal catharticus when the vitriolic. In like manner, if forty parts of quickfilver, rubbed with one part or more of sublimate already made, till it is divided into minute grains, be ground with twenty-eight parts of nitre, then with thirty-three of common falt, and afterwards, for a little time longer, with fixty-fix of calcined green vitriol, and the mixture fet to fublime; or if to four parts of mercury disfolved in as much nitrous acid, and evaporated to dryness, five parts each of calcined sea falt and calcined green vitriol be added, and the mixture fubmitted to fublimation; the fame compound will be produced: the acids of the nitre and fea falt

Merc. corrof. fublimat. vel alb. Pb. Lond.
Mercur. fublimat. corrof. Pharm.
Edinb.

are extricated by that of the vitriol: the nitrous acid affifts the marine to corrode the mercury: and the mercury, combined with the marine, fublimes, and, if the process is duly conducted, concretes into a crystalline cake, the form in which this compound is expected in the shops.

This preparation, undiluted, is a most violent corrosive. A solution Aq. phageof it in lime-water, in the proportion of a dram to a quart, and a stronger folution, made by boiling the same quantity of powdered sublimate, with equal its weight of alum, in a pint of common water, till half the liquor Aq. alumiis wasted, are employed for some external purposes, as the cleansing of foul ulcers and suppressing fungosities, and removing obstinate defedations of the skin. The lime-water, like lixivia of fixt alkaline salts, precipitates a part of the mercurial preparation, and hence the impregnation of the liquor cannot be precifely afcertained; for the stronger the limewater, the more of the fublimate will be precipitated, and the less corrofive will the folution be: at the fame time also, the lime in the water changes its nature, by its coalition with the acid which it abforbs from the fublimate. In the aluminous folution, no feparation happens, both the fublimate and the alum retaining their full force: for on mixing together folutions of the two made separately, no precipitation or turbidness ensued.

Small doses of this corrosive preparation, properly diluted, have been ventured upon internally. Boerhaave relates, that if a grain be dissolved in an ounce of water, and a dram of this folution foftened with fyrup of violets, taken twice or thrice a day, it will perform wonders in many reputed incurable distempers. Van Swieten brought it into more general use, for the cure of venereal maladies: he dissolves a grain of the fublimate in two ounces of proof spirit, [rectified spirit dissolves it more perfectly] and gives of this folution from one to two spoonfuls twice a day; continuing the medicine so long as any of the symptoms remain. with a low diet, and plentiful dilution. In the medical observations and inquiries, published by a society of physicians in London, there are many instances of the success of this method: the sublimate operated chiefly by urine and fweat, though fometimes, for the first two or three days, by ftool; and appeared not only fafe, but more to be depended on, for the removal of the fymptoms, than any of the other mercurials used as alteratives. If it be true, as some have presumed, that the completeness of the cure has any dependence on the quantity of mercury introduced

into the blood (a); it would follow, that the cure by fublimate must be less complete than that obtained by any other mercurial preparation, and that those preparations which can be taken without disturbance in considerable doses, as five or fix grains or more, promise the most lasting cures: experience however has now sufficiently shewn, that the cures obtained by sublimate are in general perfect.

Corrosive sublimate consists of mercury, united with so much marine acid, as to be dissoluble in boiling water. If by separating a part of the acid, or adding more mercury, the proportion of acid is rendered so small, as that no part of the compound shall be dissoluble, when sinely powdered, by long boiling in water; its corrosiveness will be destroyed, and it may now be taken with safety in doses of some grains. A little volatile spirit or alkaline lye, dropt into the water after the boiling, will discover if it has taken up any part of the mercury, by turning it cloudy or yellow: rain, snow, or rather distilled water, should be employed for this trial, as the common spring waters are themselves made cloudy by alkalies.

Merc. præcipit. albus Pb. Edinb.

Merc. præcipit. albus Ph. Lond.

Unguent. e merc. præc. Pb. Lond.

Spirit of fal ammoniac, or other volatile alkalies, dropt into a filtered folution of fublimate, abforb a part of the acid; and the mercury, retaining fo little as to be indiffoluble, renders the liquor milky, and fubfides, on standing, in form of a fine white powder, which, washed by repeated affusions of hot water, becomes insipid. Solutions of fixt alkaline falts, fubflituted to the volatile spirit, produce a yellow precipitation: but if an equal weight of crude fal ammoniac be diffolved along with the fublimate, fixt alkalies, added to this folution, extricate the volatile alkali of the fal ammoniac, and the precipitate proves the fame, as if the volatile alkali alone had been added in its pure state. These precipitates are used chiefly on account of the elegance of their colour, in unguents for cutaneous eruptions: one part of the mercurial precipitate, three of precipitated fulphur, and eighteen of the simple ointment or pomatum, moistened with a little strong alkaline lye, make the common mercurial application for these complaints. The precipitates have been given internally; but mercurius dulcis, which differs from them only in being more mild, and more equal and certain in its effects, is in

this intention greatly to be preferred. It does not appear, that a combination of mercury with fo small a proportion of acid, that is, so mild and fafe a mercurial, can be obtained by any kind of precipitation,

as by the process by which mercurius dulcis is prepared.

Mercurius dulcis is sublimate made mild, by combining with it so much fresh mercury, as is sufficient to satiate the redundant acid. Four parts of powdered fublimate are ground with three, or three and a half of quickfilver (an operation in which great caution is necessary, to avoid the lighter corrofive particles that fly off) till they are thoroughly incorporated; or, which is much more commodious, digested together in a gentle heat, by which the union will be performed as effectually. mixture is then fublimed in a glass matras or phial; the sublimed white mass freed from the whitish acrid matter about the mouth of the vessel, and from fuch mercurial globules as may happen to appear distinct, then pulverized, and fublimed again: the college of Edinburgh directs the Merc. dulcis fublimation to be repeated three or four times, that of London fix times. Lond. Edin. By repeated sublimations, if a sufficient quantity of mercury has not been Aquila alba. united at first, the medicine becomes less liable to irritate the first passages and run off by stool; on account of some small part of the acid, or some portion of the compound not fully dulcified in the first operation, being separated or diffipated by the heat. The dulcification depends folely on the combination of fo much fresh mercury with the sublimate, as may fully fatiate the acid: the union of the two is effected by the digestion previous to the sublimation: and the only use of the sublimation itself is to separate such part as may remain undulcified, this part being the most volatile.

Mercurius dulcis appears to be the best and safest of the mercurial preparations that can be taken in a folid form, whether as a fialagogue or as a general alterant; no one of the mercurials, whose transmission into the blood can be depended on, being fo little disposed to affect the first passages. As a sialagogue, sive, ten, and sometimes sifteen grains, made into a bolus or pills, are repeated every night or oftener, till the ptyalism begins. As an alterative, it is given from one to two or three grains. It generally answers best in small doses, which may be repeated, with due caution, every evening, for a confiderable time, without inconvenience.

fubl. Pharm.

Mercury, precipitated from aqua fortis by fixt alkali, dissolves totally, by the assistance of heat, in distilled vinegar: on cooling, the salt crystallizes into fine brilliant plates, which float in the liquor like pieces of silver leaf, and are very dissicult of solution in water. This salt, as appears from the experiments of Hellot and others (a), is the basis of an antivenereal medicine which has lately come into great repute abroad, but which does not seem, from the accounts that have been published of it, to be either more safe, or more effectual, than some of the common officinal mercurials.

Keyfer's pills.

Æthiops antimonialis.

Pil. æthiopicæ Pb. Edinb. In some obstinate desedations of the skin, mercurials and antimonials, joined together, have frequently better effects than either of them unassisted by the other. Some triturate quicksilver with twice its weight of crude antimony, till the mercurial globules disappear, and the mixture becomes an uniform æthiops or black powder: others, instead of the crude antimony, use the medicinal regulus, or the golden or precipitated sulphur; and thus obtain an æthiops of more activity. The college of Edinburgh has given a prescription of pills on this principle, composed of three parts of quicksilver, two parts each of golden sulphur of antimony, gum guaiacum and honey, and so much mucilage of gum-arabic as will reduce them into a mass of a due consistence: if a dram of the mass is made into twenty pills, the dose may be increased from one to six or more, according to the operation.

Some of the mercurial preparations have been faid to be oftentimes for for five in the cinnabar and red corrofive with red lead, the corrofive fublimate and mercurius dulcis with arfenic. The red lead may be readily discovered by fire; the mercurial part evaporating, while the saturnine remains behind. With regard to the other abuse, some have affirmed it to be impracticable (b); for if arsenic be mixed with sublimate, and the mixture set to sublime, the marine acid quits the mercury, and unites with the arsenic; with which it composes, not a solid crystalline, but a soft butyraceous concrete, called by the chemists butter of arsenic. If arsenic should nevertheless, in certain circumstances (c), be combinable with the sublimate into a crystalline cake; and if the pernicious artifice

(a) Memoires de l'acad. roy. des sciences de Paris, 1759. (b) Neumann, Chemical works, p. 142.

should

should be ever practifed, the reports of which we presume to be ground-less; the well known properties of arsenic afford sufficient means for detecting it. If a compound of sublimate and arsenic be mixed with equal its weight or more of fixt alkaline salt, chalk, or vegetable ashes, and exposed to a moderate heat, the arsenic will sublime into the upper part of the glass, and may now be distinguished by its own proper characters. Some recommend alkaline lixivia as a criterion of this abuse: sublimate that contains arsenic being said to give a black colour with the alkali: on what soundation this should happen, I cannot conjecture; for arsenic strikes no blackness with alkalies either fixt or volatile; solutions of it are, on the contrary, by both alkalies made white.

ARISTOLOCHIA.

BIRTHWORT: a plant with heart-shaped leaves set alternately on the stalks; in the bosoms of which come forth irregular tubulous flowers, with a wide mouth, whose lower part is produced into a long flap like a tongue: the seed-vessel is large, roundish, and divided into fix cells.

- vera C. B. Long birthwort, with uncut leaves, standing on pedicles; and oblong roots, not tapering to a point, brownish on the outside and yellow within.
- 2. ARISTOLOCHIA ROTUNDA Linn. Aristolochia rotunda flore ex purpura nigro C. B. Round birthwort: with uncut leaves joined immediately to the stalks, and roundish roots.
- 3. PISTOLOCHIA: Aristolochia tenuis Pharm. Paris. Aristolochia pistolochia dicta C. B. & Linn. Bushy birthwort: with indented leaves set on pedicles; and bushy roots, composed of a number of fibres issuing from one head. In this and the foregoing sorts, the stalks are weak and trailing, and the flowers stand solitary.
- 4. ARISTOLOCHIA TENUIS Pharm. Lond. & Edinb. Aristolochia clematitis recta C. B. Aristolochia clematitis Linn. Creeping birth-

wort: with upright stalks, flowers standing in clusters, and long slender creeping roots rarely exceeding the thickness of a goose quill.

THESE plants are natives of the fouthern parts of Europe, from whence we are supplied with the dry roots. They bear the colds of our own climate; the third fort excepted, which dies in fevere winters. The fourth spreads fast, to a great distance, so as not to be easily extirpated.

ALL the birthwort roots have fomewhat of an aromatic fmell, and a warm bitterish taste. They are represented by authors, as being extremely hot and pungent: Boerhaave fays, they are the hottest of the aromatic plants, and, as it were, burn the tongue and palate, having probably examined the fresh roots; but whatever their qualities may be in that state, such as are usually met with in the shops, have no great pungency. The long and round forts, on first chewing, scarce discover any taste, but in a little time prove nauseously bitterish; the round fomewhat the most so. The other two instantly fill the mouth with a kind of aromatic bitterness, not very ungrateful.

These roots give out their virtue by infusion both to spirituous and watery menstrua, to the first most perfectly: the colour of all the tinctures is brownish or yellowish. In distillation, pure spirit brings over little or nothing: with water there arises, at least from both the flender rooted forts, a small portion of effential oil, possessing the smell and flavour of the roots. The extracts made with spirit smell moderately, and taste strongly, of the birthworts: the watery extracts have nothing of their peculiar flavour, and are much more naufeous in tafte than either the spirituous extracts, or the roots in substance.

The birthwort roots are celebrated as warm attenuants and deobstruents, particularly in suppressions of the uterine purgations, from which virtue they are supposed to have received their name: the dose is from a scruple to a dram and upwards. They have likewise been recommended, particularly the fourth fort, as alterants in the gout: Boerhaave observes, that the pituitous gout, as he calls it, is often relieved by an infusion of these roots in spirit of juniper berries, sweetened with fugar, and taken to the quantity of a spoonful at a time; but that in other kinds of the gout, and in subjects of a tender constitution, this medicine

medicine occasions a loss of appetite, a weakness of the stomach, and a languidness, less supportable than the gout itself. A powder composed of this and other similar materials, which was prescribed by the ancients as an antiarthritic, and has lately come again into esteem, has also produced complaints of the same kind. Externally these roots have been used as discutients, detergents, and antiseptics: Simon Paulli relates, that the long birthwort roots, applied as an epithem or in somentation, were found remarkably serviceable in stubborn ulcers of the legs.

ARSENICUM.

ARSENICUM ALBUM Pharm. Parif. Arsenicum simpliciter dictum. Arsenic or white arsenic: a semitransparent crystalline concrete, assuming an opake milky hue on being exposed for some time to the air: soluble plentifully in alkaline lixivia, more sparingly in oils, and still more so in acids; dissolving also, by the assistance of a boiling heat, in water, but separating and crystallizing in great part as the liquor cools; totally exhaling, by a heat below ignition, in thick sumes, distinguishable from those of all other known mineral substances, by a strong fetid smell resembling that of garlic.

The fumes, caught in veffels, condense, either into a crystalline form again, or into a powdery one, according as the receiver is less or more removed from the action of the heat. If the arsenic be mixed with vegetable or animal coals, or other inflammable substances not sulphureous, that is, not participating of vitriolic acid, and exposed to a moderate heat without communication with the air, it sublimes in form of a bright greyish metallic substance, quickly tarnishing to a black, lighter and less compact than most of the other metallic bodies, scarcely seven times specifically heavier than water, changeable into a calx or white arsenic again by sublimation with the admission of air.

Arfenic is contained, in greater or less quantity, in the ores of most metallic bodies, particularly in those of tin and bismuth, and in the mineral called cobalt, cobaltum, cadmia metallica; from which last, greatest part of the arsenic brought to us is extracted, in Saxony, by a kind of sublimation: the arsenic rises at first into a large horizontal chimney communicating with the surnace, in form of a greyish meal, which, more carefully resublimed, concretes into the crystalline white

arfenic

arfenic of the shops. Henckel observes, that of all the metallic bodies, mercury and the antimonial metal are the only ones which are never found to have any arsenic in their ores: to these perhaps may be added zinc, whose proper ore, calamine, appears to be pure from arsenic.

White arfenic is one of the most violent poisons. Besides the effects, which it produces in common with other poisonous substances, it is said to render the coats of the stomach remarkably thin, to occasion a swelling and sphacelation of the whole body, and a sudden putrefaction after death, particularly of the genitals in men (a). Where the quantity taken has been so small as not to prove fatal, tremors, palsies, and lingering hectics succeed. It has likewise been observed to produce very dangerous, and sometimes mortal symptoms, when applied externally (b), which it was formerly recommended to be, against cancers and scrophulous tumors.

* This dangerous mineral has of late been confidered by some as a real specific against the cancerous virus. Mr. Le Febure has ventured publicly to recommend its internal use, together with a topical application of it to the affected part, in cancerous cases; and positively affirms that he has found it efficacious in more than two hundred instances, without any bad effects. He gives a very dilute solution of white arsenic, mixed with milk and syrup of poppies. Mr. Justamond, who published a treatise on cancers two or three years since, agrees with the above-mentioned author in the idea that arsenic is specific against this disease, but laments that even the most guarded external use of it, while it produces the happiest effects in healing cancerous ulcers, yet occasions such disagreeable symptoms of the paralytic kind, that it cannot be persisted in. The latest trials in London are said to confirm this account.

The remedies against this, as against most other poisons, are, milk and oily liquors, immediately and liberally drank. Hossman tells us of several persons of distinction, who, on tasting food with which white arsenic had been mixed instead of sugar, were all seized with anxiety at the breast, pain at the stomach, tremor of the lips, and reachings: milk

⁽a) Stahl. Mens. Novemb. cap. iv. opusc. p. 454, 455. Lindestolpe, De venenis, edit. Stentzel, cap. xvi. thes. x. p. 755.

⁽b) Degner, Hift. dysenteriæ biliosæ, p. 214. Hildanus, Obs. p. 606. Heimrich, Ad. nat. cur. vol. ii. obs. 10. Fernelius, De methodo medendi, lib. vi. cap. 18.

and oil were poured down, plentifully and repeatedly, so as to keep them strongly vomiting for half a day; some vomited no less than an hundred times: by this simple remedy they all escaped (a), and some instances of the same kind have fallen within my own knowledge. Tachenius relates, that convulsions of the limbs, gripes, bloody urine with inexpressible pain, and a contraction of the whole body, which he had been feized with from exposure to the fumes of arsenic, being relieved by milk and oil, a flow fever succeeded, which continued during the winter, and of which he was at last cured by decoctions of the vulnerary herbs, and by the use of cabbage sprouts with orange juice, oil, and a little falt (b). * Sage, in his Mineralogy, fays that the regulus is far less dangerous than the calx or glass: he gave half an ounce of the regulus to a cat; who grew meager for some time, but afterwards fat again - that acids, particularly vinegar, are the antidotes to this poison. Oils and emulsions do not obtund its effects as acids do; of this he has had experience on brutes - that the regulus is not foluble in water; and that the founders are more afraid of fumes of lead than of arfenic.

SULPHUR, which restrains the power of mercury and the antimonial metal, remarkably abates the virulence of arfenic; compositions of arsenic and sulphur being far less poisonous than the pure white arsenic, and those, in which the quantity of fulphur is considerable, seeming to be almost innocent. Different compositions of this kind are both prepared by art, and found native in the earth.

The bright yellow, fomewhat transparent masses, called yellow arsenic, Arsenicum are prepared, by mixing the arfenical meal, as extracted by the first flavum. fublimation from the ore, with one tenth its weight of fulphur, and fubliming them together: on doubling the quantity of fulphur, the compound proves more opake and compact, of a deep red colour, refem- Arfenicum bling in the mass that of cinnabar, but losing of its beauty on being ground into powder, whilft that of cinnabar is improved by trituration: by varying the proportions of arfenic and fulphur, fublimates may be obtained of a great variety of shades of yellow and red. The fosfil fulphurated arfenics differ remarkably in texture as well as in colour, fome

⁽a) Hoffman, Syft. med. rat. de feb. fect. ii. cap. iii. obf. iii.

⁽b) Tachenius, Hippocrates chymicus, p. 149.

being smooth and uniform like the factitious masses, and others composed of small scales or leaves; the former are commonly distinguished by the name zarnichs, the latter by that of auripigmenta or or piments: the orpiments are the only substances to which the Greeks gave the name arsenicon, the preparation of white arsenic being a discovery of later years: the red zarnichs are the sandarache of the Greeks, and the realgar and risigal of the Arabians and some of the chemical writers.

That these fossils are really sulphurated arsenics is evident from sundry experiments. When set on fire, the arsenical, as well as the sulphureous smell, is plainly distinguishable. If triturated with quicksilver, and exposed to a suitable heat, the sulphur is detained by the mercury, and a pure white arsenic sublimes. A mixture of fixt alkaline salt, with any vegetable or animal substance, as the compound called by the assayers black flux, in like manner keeps down the sulphur, and at the same time revives the arsenic into its reguline or metallic form.

All these compounds are mild, compared to the white arsenic; and several of them are looked upon by many as having no poisonous quality. Some, both of the factitious and native, have been given to dogs in considerable quantities, without producing any ill effect. The native minerals have been used as medicines in the eastern countries, and by some imprudently recommended in our own.

ARTEMISIA.

ARTEMISIA Pharm. Lond. & Edinb. Artemifia vulgaris major C.B. Artemifia vulgaris Linn. Mugwort: a plant, with firm stalks, generally purplish; deeply divided leaves, resembling those of common wormwood, of a dark green colour above, and hoary underneath; and small, purplish, naked, discous flowers, standing erect, in spikes, on the tops of the branches. It is perennial, grows wild in fields and waste grounds, and flowers in June.

This plant has been chiefly recommended for promoting the uterine evacuations, and abating hysteric spasms; for which purposes, insusions of it have been drank as tea and used as a bath. It appears to be one of the mildest of the substances commonly made use of in such intentions; and may, perhaps, be of service, in some cases, where medicines of more activity would be improper.

The

The leaves have a light agreeable smell, especially when rubbed a little; but scarcely any other than an herbaceous taste. An extract made from them by water is likewise almost insipid; and an extract made by spirit has only a weak aromatic bitterishness. Baierus informs us, in a dissertation on this plant, that by fermenting a large quantity of it, and afterwards distilling, and cohobating the distilled water, a fragrant sapid liquor was obtained, with a thin fragrant oil on the surface. The flowery tops are considerably stronger than the leaves, and hence should seem to be preferable for medicinal use.

ARTHANITA.

ARTHANITA, Cyclamen, Pharm. Edinb. (a) Cyclamen orbiculato folio inferne purpurascente C. B. Cyclamen europæum Linn. Sowbread: a low plant, without any other stalk than the slender pedicles of the leaves and flowers: the leaves are pretty large, round, of a green colour above with white specks, and purplish underneath: the flowers purplish, monopetalous, deeply divided into five segments, followed by round seed-vessels: the roots large, somewhat globular, with several sibres, blackish on the outside and white within. It is perennial, a native of the southern parts of Europe, and cultivated in some of our gardens.

The fresh roots of arthanita have a nauseous, acrid, biting taste, and no remarkable smell: by drying, their acrimony is greatly abated; by long keeping, it is almost destroyed; though after they have lost so much as to make very little impression on the organs of taste, they still betray, when taken internally, a great degree of irritating power. Dried and powdered, they have been given in doses of a dram, and found to operate as a strong inflammatory, yet slow cathartic. The juice is said to purge when applied externally to the belly in ointments; and the juice or bruised root to be of great efficacy for softening and discussing indolent hard tumours. The flowers are of a different nature, having a moderately strong and very pleasant smell, and little other than a mucilaginous taste: they have not been used medicinally, and the use of the roots is now, among us, in great measure laid aside.

ARUM.

ARUM Pharm. Lond. & Edinb. Arum maculatum maculis nigris C. B. Arum maculatum Linn. Wake-robin of cuckowfint: a low perennial plant, growing wild under hedges and by the fides of banks. It fends forth, in March, three or four smooth leaves, triangular, or shaped like the head of a spear: these are succeeded by a naked stalk, bearing a purplish pistil inclosed in a long sheath, which is followed, in July, by a bunch of red berries: the root is irregularly roundish, about an inch thick, brownish on the outside and white within. In some plants the leaves are spotted with black, in others with white spots, and in others not spotted at all: the black spotted fort is supposed to be the most efficacious, and hence is expressly directed by the London college. All the forts are said to be stronger, when produced in moist shady soils, than in dry open exposures.

THE fresh roots of arum have an extremely pungent acrimonious taste: slightly chewed, they continue to vellicate and burn the tongue, the part which they principally affect, for many hours, producing at the same time a considerable thirst: these uneasy sensations are somewhat alleviated by milk, butter, or oily liquors. The other parts of the plant are likewise highly acrid, though rather less so than the roots. No part has any smell, except the pistil, which has a faint fetid one.

This root loses greatest part of its acrimony on being dried sufficiently to become pulverable. Kept dry for some time, it seems, on first chewing, to be an insipid farinaceous substance: it still, however, retains a kind of latent pungency, so as when chewed long, in any considerable quantity, to produce a sensation as of a slight excoriation of the tongue. Parkinson observes, that the white farinaceous starch-like powder has been used in washing, and that it has sometimes blistered the hands.

The fresh root, dug up in autumn, yielded upon expression about one sixth its weight of a milky juice; which, on standing, deposited a white fecula and became clear: the clear liquor was insipid: the fecula was considerably pungent, but, like the root in substance, loses its pungency on being dried. The fresh and the moderately dried roots were digested in water, in wine, in proof spirit, and in rectified spirit, with and without heat: the liquors received no colour, and little or no taste. In distillation.

Fœcula ari.

distillation, neither spirit nor water brought over any sensible impregnation from the arum: the watery and spirituous extracts also were nearly The root, nevertheless, loses in these operations almost the

whole of its pungency.

The root may be preserved fresh in fand for several months; and if the fand is fomewhat moift, so as to fuffer it to vegetate, it will be the better fecured both from rotting and from losing of its virtue. It appears to be of equal vigour, or at least of sufficient vigour for medicinal use, at all periods of its growth, and in all feafons of the year. As it has hitherto been used only in a dry state, it has been generally taken up about the time of the plant beginning to dye, as the root is then leaft juicy, and shrinks least in drying.

ARUM root, newly dried and powdered, is given in doses of a scruple and upwards; for stimulating the folids, attenuating viscid juices, and promoting the natural fecretions; in cold, languid, phlegmatic habits; against weakness and relaxations of the stomach, and catarrhous and rheumatic diforders. It has been generally given in conjunction with other materials, and to some of these the effects of the compound have been in great part ascribed; thus, in the officinal powders of arum, fixteen grains of the dry root are accompanied with eight grains of the pungent roots of pimpinella, the same quantity of those of the yellow Pulv. ari water flag, four grains of cinnamon, and two grains of falt of wormwood, with four grains of crabs-eyes. In the medicine recommended by Sydenham against rheumatisms, the acrid antiscorbutic herbs are largely joined to it.

The vinous and spirituous tinctures of this root, by some recommended, appear, from the experiments above related, to be infignificant: but though spirituous liquors are incapable of dissolving or extracting the active matter of the arum, they feem nevertheless, when given along with the dried root as a vehicle, to promote its action: Juncker observes, that a dram of the powder, taken with a spoonful of brandy, procures a very copious fweat, even in persons little disposed to that evacuation; while the powder, by itself, has no such effect (a).

The insupportable pungency on the tongue, which has hitherto prevented this root from being used in a fresh state so as to exert its full virtues, I have found to be effectually covered by unctuous and gummy The fresh root, beaten into a smooth mass, with the addition of a little testaceous powder which promotes the division of it, may either be mixed with about an equal quantity of powdered gum-arabic, and three or four times as much conferve, fo as to make them into an electary; or rubbed with a thick mixture of mucilage of gum-arabic and spermaceti, gradually adding any suitable watery liquors and a little fyrup, fo as to form an emulfion: two parts of the root, two of gum, and one of spermaceti, make an emulsion which scarcely impresses any degree of pungency though kept long in the mouth. In these forms I have given the fresh root from ten grains to upwards of a scruple, three or four times a day: it generally occasioned a sensation of slight warmth, first about the stomach and afterwards in the remoter parts, manifestly promoted perspiration, and frequently produced a plentiful fweat. Several obstinate rheumatic pains were removed by this medicine, which is therefore recommended to further trial.

ASA FOETIDA.

ASA FOETIDA Pharm. Lond. & Edinb. Laser. Laserpitium. Sylphium. ASAFETIDA: the fetid concrete juice of a large plant, growing in the mountains of the provinces of Chorafaan and Laar in Persia, called by Kæmpfer umbellifera levistico affinis, &c. "The um-" belliferous plant akin to lovage, with branched leaves like those of " piony, a very large full stalk, and naked solitary foliaceous seeds like "those of parsnep or cow-parsnep, yielding asafetida from its root." It is the Ferula Asia Fætida of Linnæus. The root of the plant, when grown to a proper age and fize, is bared of earth at the top, skreened from the fun by the leaves that have been pulled off, after fome days cut horizontally, and again carefully skreened: the juice gradually rises, and in a day or two is accumulated on the furface; and being thence collected, the superficial part of the root, that has become dry, is cut off, to allow an exit to the remainder of the juice. A particular detail of this process may be seen in Kæmpfer's amanitates exotica. This juice, as it first issues from the root, is liquid and white like milk: on being exposed to the air, it turns brownish, and gradually acquires different degrees of confistency. It is brought to us in large irregular masses, composed of various shining little lumps or grains, which are partly whitish,

whitish, partly of a brownish or reddish, and partly of a violet hue. Those masses are accounted the best, which are clear, of a pale reddish colour, and variegated with a great number of fine white tears.

THIS juice has a strong fetid smell approaching to that of garlic, and a nauseous bitterish biting taste. It is by much the strongest of the deobstruent warm fetid gums; and is given not only against hysterical complaints, flatulent colics, and obstructions of the breast; but in most of the diforders called nervous, in which it frequently acts as an antispasmodic and an anodyne: in some cases musk, and in some opium, are usefully joined to it. It is sometimes used also as an anthelmintic, and faid by Hoffman to be one of the capital medicines of that class. It is most commodiously taken in the form of pills; from a few grains to a scruple or half a dram. It loses with age of its smell and strength; a circumstance to be attended to in dofing it: Kæmpfer informs us, from his own observation, that a single dram of the recent juice smells more than an hundred pounds of fuch as is commonly fold in Europe.

Asafetida is composed of a gummy and a resinous substance, the first in largest quantity. Its smell and taste reside in the resin; which is readily diffolved and extracted by pure spirit, and, in great part, along with the Tinetura gummy matter, by water. The tincture made in pure spirit is of a foetida Ph. transparent yellow colour; that made in proof spirit, and the watery infusion, is turbid. A tincture in rectified spirit is kept in the shops; whereof two drams contain nearly all the virtue of fifteen grains of the

asafetida.

In distillation with water, it impregnates the aqueous fluid highly with its fcent, and yields a small portion of a pale-coloured essential oil which fmells exceeding strongly: the remaining decoction, inspissated, leaves a weakly nauseous bitterish extract, of very little smell. Rectified fpirit distilled off from the tincture made in that menstruum, proves likewife confiderably impregnated with the flavour of the afafetida, though much less fo than the distilled water: the remaining extract smells moderately, and tastes strongly of the juice.

ASARUM.

ASARUM Pharm. Lond. & Edinb. C. B. Nardus rustica; Vulgago. Afarum europæum Linn. ASARABACCA: a low plant without stalks:

the leaves are stiff, roundish, with two little ears at the bottom, somewhat resembling a kidney, of a dark shining green colour, somewhat hairy, set on pedicles three or sour inches long: the flowers, which rise among the leaves on shorter pedicles, consist of purplish stamina standing in a darker coloured cup, and are followed, each, by a capsule containing six seeds. It is perennial and evergreen, a native of the southern parts of Europe and the warmer climates, and raised with us in gardens. The dried roots have been generally brought from the Levant; those of our own growth being supposed of weaker virtue.

THE roots and leaves of asarum have a moderately strong, not very unpleasant smell, somewhat resembling that of valerian or nard; and a nauseous bitterish, acrid taste. The roots given in substance, in doses of a scruple or more, prove strongly emetic and cathartic. The leaves have the same operation, but their dose or degree of sorce has not been precisely determined: according to some, they are of more activity than the roots.

It is faid, that this emetic plant has been of fervice in ferous diforders, and hurtful in melancholic cases: that in small doses, it promotes perspiration, urine, and the uterine flux: that tinctures made in spirituous liquors posses both the emetic and cathartic virtues of the asarum: but that the extracts, obtained by inspissating these tinctures, act only, and with sufficient mildness, by vomit; requiring to be given in as large doses as the plant in substance, to produce as plentiful evacuations: that insusping in water operate mildly both upwards and downwards: that by coction in water, the emetic power is first destroyed, and asterwards the purgative; the decoction long boiled, or an extract prepared with a large quantity of water not acting at all by stool or vomit, but proving powerfully deobstruent, diuretic, &c. It is obvious, however,, as the activity of the asarum is diminished more and more by boiling, that both the decoction and the extract must be accompanied with one capital inconvenience, precariousness in point of strength.

The principal use of asarum among us is as an errhine. The root is one of the strongest of the vegetable substances commonly employed in this intention: a grain or two, snuffed up the nose, procure a large evacuation of mucus both from the nose and mouth, without provoking sneezing like the white hellebore root, or discovering any remarkable

irritation.

irritation. The leaves, though supposed to be stronger than the roots as emetics and cathartics, appear to be milder as errhines. Geoffroy relates, that after fnuffing a dose of this errhine, he has observed the falutary discharge to continue for three days together, and that he has known a paralysis of the mouth and tongue cured by one dose: he recommends this medicine in stubborn disorders of the head proceeding from viscid matters, in palfies, and in foporific distempers. During its operation, the patient must carefully avoid cold; which is apt to produce pustules, inflammations, and swellings of the face, and sometimes more alarming fymptoms. This herb is a principal ingredient in the cephalic or sternutatory powders of the shops: some take three parts of dried asarum and one of marjoram leaves, + others equal parts of the dried leaves of afarum, + Pulv. cemarjoram, and marum fyriacum, and dried lavender flowers ‡. empyrical herb snuffs have likewise the leaves of asarum for their basis, I Pulv. sterbut often mixed with ingredients of a more dangerous nature.

nutatorius Ph. Lond.

ASPARAGUS.

ASPARAGUS Pharm. Edinb. (a) Asparagus sativa C. B. Asparagus officinalis Linn. ASPARAGUS: a perennial plant, cultivated for culinary use. In the spring appear a number of straight naked shoots; which, rifing to the height of two or three feet, divide into flender, firm, fpreading branches, clothed with foft, green, capillary leaves: the flowers are of a pale greenish colour, and succeeded by shining red berries.

THE young shoots of asparagus, boiled, are supposed to promote appetite, but afford little nourishment. They give a strong fetid smell to the urine in a little time after being eaten, and for this reason have by fome been accounted useful diuretics, by others injurious to the kidneys. It does not appear, from common experience, that they possess either of these qualities in any considerable degree. * (b)

The roots of the plant, which are the part principally employed for medicinal purposes, are less agreeable in taste than the young shoots,

(a) Expunged.

^{*(}b) Bergius afferts that he knew a lady who after eating asparagus generally made bloody urine. Mat. Med. p. 268.

and supposed to be more aperient and diuretic: they appear to be similar, in virtue as in taste, to the roots of sennel, parsley, and the others commonly called aperient, to which they have been sometimes joined in apozems and insusions. It is observable, that neither the roots, nor the stalks when grown up so as to part into branches, give any ill smell to the urine.

ATRIPLEX OLIDA.

ATRIPLEX OLIDA Pharm. Lond. Atriplex fætida Ph. Edinb. & C. B. Chenopodium fætidum Tourn. Chenopodium Vulvaria Linn. Blitum fætidum vulvaria dictum Raii fyn. Garofmum. Stinking orache or arach: a low, procumbent plant, sprinkled all over with a whitish clammy meal: the leaves are small, of a roundish figure, with an obtuse point: on the tops of the branches come forth clusters of imperfect flowers, followed each by a flattish seed. It is annual, grows wild about dunghills, and flowers in August.

This plant has a moderately strong smell, not a little offensive, somewhat akin to that of salt-sish, and which lasts long on the hands after touching the herb: it is sometimes met with among old rubbish; in which situation, its smell proves weaker than when produced in moister places, which it naturally delights in, and is also somewhat of a different kind: in either case, its taste is not very considerable. It gives a strong impregnation to water, both by insusion and distillation: the smell is extracted likewise by rectified spirit, and by this menstruum in some degree covered. In drying, the smell becomes weaker and of a less offensive kind: in keeping it is dissipated, but not soon.

Stinking arach, on account of its strong scent, is reckoned an useful antihysteric; in which intention some recommend a conserve of the leaves, others a watery insusion, and others a spirituous tincture of them. On some occasions, it may perhaps be preferable to the fetids which have been more commonly made use of; as not being accompanied with any pungency or irritation, and seeming to act merely by virtue of its odorous principle.

AURANTIA.

AURANTIA & arantia malus. Citrus Aurantia Linn. TREE: a beautiful evergreen tree or shrub; with numerous, flexible, fomewhat prickly branches; fmooth, firm, broad leaves, having each two heart-like appendages on the pedicle; pentapetalous white flowers, fet thick together among the leaves; and a large round yellow fruit, divided internally into eight cells, filled with a juicy pulp and whitish feeds. It is a native of the warmer climates, and fcarcely bears the winters of ours without artificial shelter.

1. AURANTIA HISPALENSIS, malus aurantia fructu acido Pharm. Lond. Aurantium bispalense Pharm. Edinb. Malus aurantia major C. B. SEVILLE ORANGE: with dark yellow warty fruit, containing an acid juice.

THE flowers of this tree are highly odoriferous; and, on account of their fine fmell, have been used in perfumes, and as a flavouring ingredient in medicinal compositions: their taste is slightly bitterish. They communicate their smell and taste both to water and rectified spirit, most perfectly to the latter: the watery infusion is of a brownish, the fpirituous of a yellow colour. In distillation with water, they impregnate the aqueous fluid strongly with their agreeable odour, and yield a Aq. naphæ fmall quantity of a fragrant effential oil: the distilled water and oil, the preparations principally made use of, are generally brought to us from Ol.feuessen-Italy and France, being rarely prepared in this country on account of the scarcity of the flowers. The watery decoction, inspissated, yields an extract unpleasantly bitterish: an extract made by rectified spirit retains, along with the bitterish matter, a moderate share of the fine flavour of the flowers.

THE leaves also have a pleasant though weak smell, and a bitterish tafte. Viewed against the light, they exhibit numerous transparent fpecks, which appear to be little veficles filled with effential oil. In distillation with water, a small portion of oil separates, of an agreeable flavour, but less so than that of the flowers.

THE yellow rind of the fruit, carefully freed from the fungous white matter underneath, is a grateful warm aromatic bitter, of frequent use as a stomachic and corroborant, and for giving an agreeable flavour to other medicines. It is warmer than the peel of lemons, of a more durable flavour, and abounds more with a light fragrant effential oil; which is lodged in distinct cells on the surface of the peel, and exudes upon wounding it. It may be made into a conserve by beating it into a pulp with triple the weight of double refined sugar.

Conferva aurantiorum Pb. Edinb.

Syrup. e cort. aurantiorum. Pb. Lond. & Edinb. Infused in boiling water, it gives out nearly the whole of its smell and taste, together with a bright yellow tincture: eight ounces of the fresh rind give a strong impregnation to four pints of water; and by dissolving in this infusion a proper quantity of sugar, an agreeable syrup is prepared in the shops. Cold water, on the other hand, extracts chiefly the bitter matter, leaving the aromatic behind: hence when the fresh peel is steeped by the confectioners, for making a sweetmeat, till it has lost its bitterness, it still retains a great share of its peculiar slavour: when large quantities are macerated, a portion of oil is found floating on the surface, from some of the cells having been distended and burst by the aqueous sluid.

In distillation with water, the essential oil, in which the slavour of the peel resides, totally arises, leaving only the bitter matter behind in the decoction. Both the oil and distilled water are very grateful: a spirituous water, moderately impregnated with the slavour of the orange peel, by distilling a gallon of proof spirit from six ounces of the dry rind, is an elegant cordial: and a simple water, more lightly slavoured with it, by drawing over a gallon of water from sour ounces of the dry peel, is an useful diluent in severs, and other diseases, where the stomach and palate are apt to receive quick disgust.

aurant. spirituosa Ph. Lond. Aq. cortic. aurant. smp. Ph. Lond.

Aq. cortic.

Rectified spirit of wine, digested on orange peel, extracts its virtues more perfectly than water, and receives from it a like yellow tincture: after the action of the spirit, the peel remains crisp, after water tough. The spirit, drawn off by distillation, tastes considerably of the peel, but discovers little or nothing of its smell: the remaining extract contains, along with its bitterness, great part of its aromatic flavour, but is less agreeable than the rind in substance.

THE juice of oranges is a grateful acid, of great use in inflammatory and putrid disorders both acute and chronical. Its acid matter differs

in some of its pharmaceutical properties, both from the sermented acid of vinegar, and from the native acid salts of the leaves of plants, at least of such as have been examined;—from the former, in its not being volatile, or not exhaling upon inspissating the juice, nor rising in distillation with the heat of boiling water;—from the latter, in its being soluble in spirit of wine; the inspissated juice, at least all its saline matter, dissolving readily in this menstruum as well as in water, and liquesying also in the air. These properties afford commodious means of preserving the acidity of the orange for many years; either in the form of a thick extract, or of a more dilute spirituous solution. The inspissation of the juice must be performed with a very gentle heat, especially towards the end of the process, when the matter begins to grow thick, as it is then not only liable to contract an empyreuma, but at the same time to have great part of its acidity destroyed.

The young unripe fruit, commonly called Curasso apples, (Aurantia curassaventia Pharm. Edinb. Aurantia enascentia & immatura Pharm. Paris.) is a grateful aromatic bitter, of a flavour different from that of the peel of the ripe fruit, and without acidity; when fresh, it has a little tartness, which in drying is in great measure lost. It readily gives out to rectified spirit the whole of its bitterness and flavour, together with a fine green tincture: water extracts its virtue less perfectly. Distilled with water, it yields a considerable quantity of yellow essential oil, of an agreeable and very fragrant smell. The spirit, distilled from the spirituous tincture, brings over likewise some share of its flavour, leaving however the greatest part concentrated in the extract, which proves an elegant, mild, aromatic bitter.

2. AURANTIA SINENSIS. Aurantia dulcis. Aurantium dulci medulla vulgare Ferrant. besperid. & Pharm. Paris. . China or Sweet orange: with bright yellow smooth fruit, containing a sweet juice.

The rind of this kind of orange has a weak smell, and very little bitterness; and is scarcely ever employed for any medicinal use. The juice, of a grateful subacid sweetness, agrees, in its general qualities, with the fructus horæi of our own climate; and like them, if taken immoderately, produces gripes and fluxes. It is a useful refrigerant in U 2 inflammatory

inflammatory dispositions, and an excellent antiseptic in scorbutic and other putrid disorders.

AURUM.

AURUM vel Sol Pharm. Parif. Gold: a yellow metal, extremely ductile; above nineteen times heavier than water; fufible in a low white heat; fixt and indeftructible in the fire; not foluble by any of the fimple acids, in the common ways of making folutions; eafily diffolving in a mixture of the nitrous and marine acids, called aqua regis,

into a yellow liquor which stains the skin purple.

Effential oils, shaken with this solution, imbibe the gold from the acid, and carrying it up to the surface, keep it there for a time dissolved; but gradually throw it off again, on standing for some hours, in form of bright yellow films, to the sides of the glass. The ether or spiritus vini æthereus takes up the gold more readily and completely, and keeps it permanently dissolved. Rectified spirit of wine mingles uniformly with the acid solution; but on standing for some days, the gold separates from the mixture, and rises in silms to the surface. A piece of tin, placed in the solution largely diluted with water, changes it red or purple, and throws down a precipitate of the same colour. By the appearances resulting from these additions, very minute portions of gold, dissolved in acid liquors, may with certainty be discovered.

This metal is found chiefly native: in small granules or filaments; intermingled among earths, or bedded in stones; in the mines of the Spanish West-Indies, among the sands of some of the African coasts and of some European rivers, and blended with the ores of some other metals. According to the nature of these admixtures, and their degree of union with the gold, the extraction of the metal is differently effected; by ablution with water; by amalgamation with mercury; by bringing the whole matter, that contains the gold, into suspense with the

addition of proper fluxes.

Gold was introduced into medicine by the Arabians, and held to be one of the greatest cordials and comforters of the nerves. As it apparently can have no medicinal effect in its gross state, not being dissoluble by any fluid that can exist in the bodies of animals; the chemists

have

have attempted to fubtilize and refolve it, and to extract what they called an anima or fulphur from it. But as no means have been discovered of feparating the component parts of this metal, their tinctures and aurum potabiles either contained none of the gold, or were no other than diluted folutions of its whole substance. That the aurum potabile of the faculty of Paris, reckoned one of the best of the preparations of this kind, (made by shaking some oil of rosemary with a solution of gold in aqua regis, Tine. auri and afterwards digesting the oil for a month in rectified spirit of wine) retains none of the gold, is obvious from the characters of this metal Parif. above laid down.

Solutions of gold in aqua regis are corrofive: fo far diluted, as that they can be taken with fafety, they are, according to Hoffman, purgative: the dry matter left upon inspissating them, is a strong caustic. The purple precipitate, made by adding pure tin to the folution, is faid to be diaphoretic: a precipitate made by alkalies is strongly purgative and emetic. This last precipitate washed from the adhering saline matter by repeated affusions of water, purges more moderately, though rarely Aurum fulwithout gripes, and fometimes operates by fweat: it has been given, Parif. from half a grain to five or fix grains, in fevers, and in convulfive and other disorders arising from, or supported by, crudities in the first passages: but as its operation is extremely variable, as it has often produced dangerous symptoms, (a) and as its best effects are no other than what may be obtained from medicines of known fafety, it is now, in this country, entirely in difuse; being regarded only as a matter of curiofity, on account of its property of exploding violently when heated or strongly rubbed.*(b)

Some have amalgamated gold with pure quickfilver, and fet the compound to calcine, as directed in page 103 for the calcination of mercury by itself, till it was converted into a red powder: others have melted the gold with twice its weight or more of martial regulus of antimony, and exposed the powdered mixture, in a glass vessel, to a moderate heat,

⁽a) Ludovici, Pharm. moderno seculo applicanda, diss. i. oper. p. 102. Stahl, Digram. de proexeucriseos dignitate, sect. viii. Hostman, Philosophia corp. haman. morbos. par. ii. cap. viii. §. 13.

^{* (}b) Aurum fulminans is still in use in Germany. A late writer (M. A. Plenciz) recommends it in every case where a sure and safe laxative is wanted; alledging that it does not act with the violence many practitioners have afferted.

till the powder became purple. That these kinds of preparations have very considerable medicinal virtues, is not to be questioned; but that those virtues have any dependence upon the gold, is scarcely to be presumed: all that can be rationally expected from this ingredient, is, to obtund the activity of the mercurial calx, and of the not fully calcined antimonial metal. When gold is thus divided by the admixture of other metallic bodies, and in some degree calcined along with them, it proves dissoluble in one of the mineral acids which would not touch it before, to wit, spirit of salt; but the acids of the vegetable and the animal kingdom, it still resists as permanently as sine gold in the mass.

BALAUSTIA.

BALAUSTIA Pharm. Lond. & Edinb. BALAUSTINES: large rofe-like flowers, of a deep red colour, fet in long bell-shaped tough cups. They are the produce of the wild or double-flowered pomegranate tree, (Granata malus Pharm. Edinb. Balaustia flore pleno majore C. B. Punica Granatum Linn.) a low prickly tree or shrub, with long narrow leaves, bearing a brownish acerb fruit about the size of an orange; a native of the southern parts of Europe, and cultivated in some of our gardens on account of the beauty and continuance of its flowers. The shops are usually supplied with the dried flowers from abroad; those of our own growth do not appear to be anywise inferiour to the foreign, but are not to be procured in sufficient quantity.

BALAUSTINE flowers are mildly aftringent and corroborant; of a moderately rough and somewhat bitterish taste, and of little or no smell, or particular flavour. They give out their aftringent matter, together with a pale red colour, both to water and rectified spirit:*(a) the extracts, obtained by inspissating the tinctures, in which the active parts of the flower are concentrated, are pretty strongly styptic. The spirituous tincture is of a paler colour, and the extract in less quantity and proportionably stronger in taste, than the tincture and extract made with water. The spirituous extract, as well as the watery infusion, strikes an inky blackness with solution of chalybeate vitriol; a proof, that the astringent matter of the balaustine, after its separation from greatest part of the mucilaginous and other grosser substances of the flower, is still dissoluble in water.

BALSAMITA.

BALSAMITA: a perennial plant, with undivided indented leaves, and yellow naked discous flowers set in form of umbels on the tops of the stalks; a native of the southern parts of Europe, and cultivated in our gardens.

- 1. Balsamita Mas. Costus bortensis vel tanacetum bortense Pharm. Paris. Mentha saracenica Officinarum Germaniæ. Mentha bortensis corymbisera C. B. Tanacetum Balsamita Linn. Costmary or alecost; with oval leaves.
- 2. AGERATUM, Balsamita femina, Costus hortorum minor, Eupatorium mesues, Herba julia, Mentha corymbisera minor. Ageratum foliis serratis C. B. Achillæa Ageratum Linn. Maudlin; with numerous, small, oblong, narrow leaves.

THESE herbs have been used as mild corroborants and aperients, in weaknesses of the stomach, obstructions of the viscera, and cachectic indispositions; and though at present disregarded, they promise, from their sensible qualities, to be medicines of some utility.

They have a moderately strong pleasant smell, somewhat approaching to that of mint, and a weakly aromatic bitterish taste: the two sorts differ a little in flavour and in taste from one another, the first having the most of the mint smell, and likewise the greatest bitterishness. Infusions of them in water smell pretty strongly, and taste slightly, of the herbs: in the tinctures made with rectified spirit, which are of a deep green colour, the fmell is in good meafure covered by the menstruum. In distillation with water, they yield a small quantity of essential oil, of a pungent tafte, and which smells strongly and agreeably of the balfamitæ; that of the fecond species is the most grateful: the remaining decoction, thus deprived of the aromatic matter, is unpleafantly, though but weakly, rough, bitter, and subsaline. They give over a part of their flavour also in distillation with rectified spirit, particularly the second species, whose odorous matter appears to be of a more volatile kind than that of the first: the extracts, obtained by inspissating the spirituous tinctures, are moderately, and not difagreeably, warm and bitterish.

BALSAMUM COPAIBA.

BALSAMUM COPAIVA Pharm. Lond. Balfamum copaibæ Pharm. Edinb. Balfamum brafiliense. Balsam of copaiba or capivi: a liquid resinous juice, obtained from a large tree of the same name, (Copaiba braziliensibus Marcg. Arbor balfamisera braziliensis fructu monospermo Raii bist. Copaisera officinalis Linn.) which grows spontaneously in the woods of Brazil, and has been lately introduced into some of the British American islands. The balsam is extracted by making deep incisions in the trunk of the tree, in the middle of the summer heats: if no juice flows, the wounds are for a time closed up. It is said, that at the proper season, several pounds of balsam issue in an hour or two; that one tree yields in all sive or six gallons; but that after once bleeding, it never affords more.

The juice, as it issues from the tree, is limpid and colourless, like the distilled oil of turpentine. As brought to us, it is usually of a pale yellowish hue, and about the consistence of oil olive or a little thicker: by long keeping, it grows nearly as thick as honey, but has not been observed, like most of the other resinous juices, to grow solid or dry. In all its states of consistency, it continues clear and transparent.

We fometimes find in the shops, under the name of copaiba, a thick, whitish, almost opake balsam, with a quantity of turbid watery liquor at the bottom. This fort, probably, is either adulterated by the mixture of other substances, or has been extracted, by boiling in water, from the bark or branches of the tree. It is much less grateful than the genuine balsam.

Balsam of copaiba has a moderately agreeable smell, and a bitterish biting taste, not very intense, but durable in the mouth. It has been employed principally, and preferably to the other balsams, in gleets, the fluor albus, and in ulcerations of the urinary passages and the lungs. Fuller says, he has known dry deep coughs, coughing up of blood and pus, voiding of chyle instead of urine, with great pains and weakness, cured by it; and that, notwithstanding the manifest warmth and bitterness of its taste, he has found it to agree in hectic cases: he observes that it gives the urine a bitter taste, but not a violet smell as the turpentines do,

and

and that if taken in doses of two or three drams, it proves like them, purgative. The usual dose is from ten to thirty or forty drops. *(a)

This balfam, agitated with water, in fome degree unites with it, and renders the liquor turbid and milky, but foon separates and rises to the furface on standing. Dropt on sugar, or triturated with thick mucilages, or with whites or yolks of eggs, it becomes more permanently miscible with water into an uniform milky liquid: it is generally taken either in this form, or mixed with powdery and other matters into a bolus or electary. It mingles with oils, eafily with the diffilled, more difficultly with the gross ones obtained by expression. It dissolves in rectified spirit of wine into a transparent liquor, of a fragrant smell, more agreeable than that of the balfam itself.

Distilled with water, it yields nearly half its weight of an essential oil, which when newly drawn is limpid, but by age grows yellowish: the part of the balfam, which remains behind in the still, is a tenacious inodorous refin, of a yellowish colour inclining to green. The refin diffolves in rectified spirit more easily than the entire balfam; the oil more difficultly, requiring, as Hoffman observes, near four times its weight of the menstruum, whereas the balsam will dissolve in twice its weight or less.

The balfam, distilled in a retort, without addition, by a fire gradually raifed, gives over first a light yellow oil, smelling considerably of the juice; then a darker coloured oil, and afterwards a fine blue one, both which are of a very pungent taste, and have little other than an empyreumatic flavour, though not of a very ungrateful kind. In the shops is Ol. copaivæ kept an empyreumatic oil, drawn from a mixture of a quart of the comp. Phar. balfam with four ounces of gum guaiacum.

BALSAMUM PERUVIANUM.

BALSAMUM PERUVIANUM, indicum, mexicanum, americanum. BALSAM OF PERU: a refinous juice, obtained from certain odoriferous trees (Cabureiba Pison. Hoitziloxitl seu arbor balsami indici Hernand.) growing in Peru and the warmer parts of America.

^{*(}a) For a censure of this practice, and of the use of the other balsams and resins in confumptive cases, see a paper of Dr. Fothergill's in Vol. IV. of the Lond. Med. Observ. and Inq.

I. Balsamum peruvianum peruvianum peruvianum nigrum, fuscum, vulgare. Common balsam of Peru; usually about the consistence of thin honey, and of a dark opake reddish brown colour inclining to black. It is said to be extracted by boiling the tops and bark of the tree in water, and to be found floating on the surface when the liquor cools (a). The balsam, however, as brought to us, dropt into cold water, does not float, but sinks immediately to the bottom: if a drop be let fall into water almost boiling hot, it separates into two parts, an oily cuticle, of a very penetrating taste, which spreads upon the surface, and a grosser matter, in larger quantity, which sinks. It may be presumed therefore, that the balsam is extracted by some other method than that above pointed out.

This balsam does not in any degree unite with water, or render it milky or turbid, by agitation. It becomes miscible with water, like that of copaiba, by the intervention of mucilage or yolk of eggs, but not perfectly by sugar: when united with sugar in a dry form, or with thick solutions of it, great part of the balsam separates and subsides on diluting the mixture with water. It dissolves in rectified spirit of wine, a small quantity of impure matter commonly remaining: and likewise, by the assistance of a boiling heat, in alkaline lixivia (b). It unites readily with distilled oils; but not at all with expressed oils or with sluid animal sats, a circumstance in which it differs remarkably from all the other resinous juices that have been examined: after it has been blended, by trituration, with consistent unctuous matters, and with wax, it separates and falls to the bottom as soon as the mixture is made fluid by heat. Nor does it mingle very perfectly with the vegetable juices of its own kind, the native balsams and turpentines.

Distilled with water, it yields about one fixteenth its weight of effential oil, of a reddish colour, a fragrant smell, and a very pungent taste: this oil is remarkably difficult of solution in spirit of wine, requiring, according to Hoffman's experiments, no less than twelve times its own weight of the spirit. The balsam, distilled in a retort, without addition, yields a larger quantity of a yellowish red empyreumatic oil, and com-

⁽a) Monardes, apud Clusium, exoticorum, lib. x.

⁽b) Hoffman, Diff. de balfamo Peruviano, cap. ii. 11 & 23.

monly, as Neumann observes, a small portion of saline matter similar to flowers of benzoine.

This juice has an agreeable aromatic smell, and a very hot pungent taste. It is one of the hottest of the natural balsams, and hence preferred, in cold phlegmatic dispositions, for warming the habit, and strengthening the nervous as well as the vascular system. The dose is from two or three grains to ten or twelve. It is used also, in preference to the other balsams, externally, for wounds and ulcers: Van Swieten observes, that for preventing or abating the terrible symptoms arising from punctures of the nerves or tendons, one of the best remedies is balsam of Peru, dropt warm into the wound, and made to spread and penetrate by applying a warm spatula (a).

2. Balsamum peruvianum album seu Styrax alba Ph. Paris. White balsam of Peru, or white storax; brought over in gourd shells; of a pale yellowish colour, thick and tenacious, becoming by age solid and brittle. It is supposed to be the produce of the same trees which afford the common or black balsam, and to exude from incisions made in their trunks.

This balfam is in taste less hot and pungent than the foregoing, in smell more fragrant and agreeable, somewhat approaching to that of storax. It readily dissolves in rectified spirit, and unites with oils both expressed and distilled, as also with animal sats. Dropt, in its shuid state, into warm water, it spreads totally upon the surface, and forms a pellicle cohesive enough to be taken off entire (b), one of the principal criteria by which the precious balsam of Gilead has been distinguished. It is rarely met with in the shops.

BALSAMUM TOLUTANUM.

BALSAMUM TOLUTANUM Pharm. Lond. & Edinb. BALSAM OF TOLU: a refinous juice; flowing from incisions made in the trunk of a tree said to resemble the pine, (Toluifera Balsamum Linn.) growing in the province of Tolu in the Spanish West Indies (c), from whence

⁽a) Comment. in Boerb. aphorismos, §. 164. vol. i. p. 242.

⁽b) Bartholin, Diff. de theriaca, ii. 26.

⁽c) Monardes, apud Clusium, exoticorum, lib. x.

the balfam is brought to us in little gourd shells. It is of a yellowish brown colour inclining to red, in consistence usually thick and tenacious: by age it grows hard and brittle, without suffering any great loss of its odoriferous parts.

This balfam has an extremely fragrant smell, somewhat resembling that of lemons; and an agreeable, warm, sweetish taste, very slightly pungent, and not accompanied, like that of most of the other balfams, with any nauseous relish. It possesses the same general virtues with those of Peru and copaiba, differing only in being milder, less hot or irritating, and more grateful to the stomach as well as the palate.

Boiled in water for two or three hours, in a circulatory vessel, or in a matras with a very long neck or having a long tube inserted into its

mouth, so as to prevent evaporation, it communicates to the liquor great part of its fragrance: eight ounces of the balsam give a strong impregnation to three pints of water: the decoction strained off from the un-

dissolved refin, forms, with a proper quantity of sugar, an elegant balsamic syrup. The balsam dissolves totally in rectified spirit of wine, and in this form may be mixed in substance with syrups, so as to im-

pregnate a much larger quantity of them with its fine flavour: forty-five grains of the balfam, diffolved in an ounce of pure spirit, are sufficient for two pounds of a simple flavourless syrup made from sugar and

water: the folution is to be stirred gradually into the syrup just warm from the fire, and the mixture kept in the gentle heat of a water bath till the spirit has exhaled. This balsam may likewise be dissolved in

water, into a milky liquor, by trituration with gums or mucilages. It unites readily with distilled oils, more difficultly with expressed oils

and with fats.

In distillation with water, it impregnates the liquor strongly with its fragance; and this, perhaps, is the most advantageous method of obtaining its simple odoriferous matter in the form of a watery solution: if the quantity of balsam, submitted to the distillation, be large, a small proportion of a very fragrant essential oil may be collected. Distilled in a retort, without addition, it yields a pale and a dark coloured empyreumatic oil, and sometimes a small quantity of a kind of saline matter, of the same nature with slowers of benzoine.

Syrup. balfamic. Ph.

Tinct. tolutana Pharm. Edinb.

Syrup. balfamic. Pb. Edinb.

BAMIA.

BAMIA MOSCHATA Pharm. Parif. Ab-el-mosch, i. e. granum moschi. Semen ketmiæ ægyptiacæ. Musk-seed: flat, kidney-shaped striated seeds, about the size of a large pin's head, of a greyish or brownish colour on the outside, and white within: produced by a shrubby plant, of the mallow kind, alcea ægyptiaca villosa C.B. Hibiscus Abelmoschus Linn. a native of Egypt, and of the East and West Indies.

THESE seeds have a fragrant smell, approaching to that of musk, and a slight aromatic bitterish taste. They are used in persumes, and seem to have a claim, as medicines, to the cordial and nervine virtues experienced from most of the other substances of that class. They are ranked in France among the officinals, but are rarely to be met with among us.

BARDANA.

BARDANA MAJOR, Lappa major, Pharm. Edinb. Lappa major, arcium dioscoridis C. B. Personata sive lappa major aut bardana J. B. Arctium Lappa Linn. Burdock: a biennial plant, common by road sides, sufficiently known by its scaly heads or burs; the leaves are very large, shaped somewhat like a heart, of a deep green colour above and whitish underneath: the seeds flatted, nearly oval, somewhat crooked, slightly striated, of a dark brown or blackish colour: the root large, straight, brownish on the outside and white within, composed of a thick cortical part, and a spongy medullary substance, with more or less of a woody septum between them.

The roots of burdock have very little smell, and a weak taste, chiefly sweetish, mixed as it were with a slight bitterishness and roughness. Boiled in water, they impart a brownish colour, and a soft vapid kind of taste: extracts made from them, both by water and by rectified spirit, are weakly sweetish, bitterish, subastringent, and subsaline. These roots are recommended as mild diuretics, diaphoretics, and sweeteners, in scorbutic, rheumatic, gouty, and venereal disorders; and are supposed to be of similar virtue to china and sarsaparilla; to which, in their sensible qualities, they have a considerable resemblance. They are used chiefly,

Decoctum bardanæ Nosocom. Ed. chiefly, and to the best advantage, in the form of decoction: two ounces of the dried root are boiled in three pints of water, till one pint is wasted, and a pint or more of the strained liquor taken warm every day.

The expressed juice of the leaves has been sometimes given, to the quantity of a quarter of a pint or more, in the same intentions. The leaves are bitter and more saline than the roots, and have nothing of their sweetishness: the juice, depurated, and inspissated to the consistence of an extract, discovers a moderately strong penetrating, saline bitterness.

The feeds also are bitterish and slightly aromatic; and have been given, in doses of a dram, as a diuretic, and as an aperient in disorders of the breast. They are said to be purgative. A good deal of care is requisite, to clear them thoroughly from the prickly matter with which they are covered, and which, if swallowed, immediately discovers how much it offends the parts it passes through, by the uneasy sensations it produces in the throat (a).

BDELLIUM.

BDELLIUM Pharm. Edinb. (b) BDELLIUM: a gummy-refinous juice; the produce of an oriental tree, of which we have no particular description; brought from Arabia and the East Indies, in pieces of different magnitudes and figures, externally of a dark reddish brown colour not unlike myrrh, internally clear and somewhat resembling glue.

This gummy-refin has a moderately agreeable smell, and a bitterish slightly pungent taste. It grows soft and tenacious in the mouth, and sticks to the teeth. Laid on a red-hot iron, it readily catches slame, and burns with a crackling noise; during which, little streams of liquid matter ooze out at the surface. Both water and rectified spirit dissolve, each, near one half of the bdellium: the spirituous tincture, of a transparent reddish yellow or orange colour, tastes stronger and smells more agreeably of the bdellium, than the watery insusion, which is turbid and brownish. Geosfroy relates, that its whole substance is dissolved by wine, vinegar, tartarized spirit of wine, and alkaline liquors: the active matter of the bdellium is indeed extracted by all these menstrua,

but the three first were found upon trial to leave a considerable part of its substance undissolved: proof spirit took up nearly the whole. In distillation with water, it impregnates the aqueous sluid weakly with its slavour; nor is there any appearance of essential oil, at least when only small quantities, as three or four ounces, are submitted to the operation. The distilled spirit has very little slavour of the bdellium; nevertheless, the spirituous extract proves weaker, both in smell and taste, than the juice in substance, its active parts being probably enveloped, in this preparation, by the tenacious resin.

This gummy-refin stands recommended, as a corroborant and attenuant, in disorders of the breast, for promoting urine and the menses; and externally for resolving or maturating hard tumours. It appears to be one of the weakest of the deobstruent gums, and is at present rarely

made use of.

BECABUNGA.

BECABUNGA Pharm. Lond. Anagallis aquatica minor folio subrotundo C. B. Veronica aquatica folio subrotundo Morison. bist. Veronica
Becabunga Linn. BROOKLIME OF WATER PIMPERNEL: a low creeping plant; with round thick smooth reddish stalks, naked and procumbent at bottom, erect at top, and clothed with roundish firm juicy
leaves, of a dark shining green colour, slightly indented about the edges,
set in pairs at the joints: from the bosoms of the leaves arise naked
footstalks, bearing spikes of blue flowers which are deeply cut into four
segments and followed by flattish seed-vessels. It is common in rivulets
and ditches, and flowers in June: the leaves remain all the winter, but
are in greatest perfection in the spring.

This herb is ranked among the antiscorbutics, and supposed to possess, in some degree, the virtues of the cochlearia and nasturtium. It is chiefly employed in conjunction with these and the other acrid antiscorbutic herbs, to which it appears to be an useful addition, though not entirely similar to them in quality. It has nothing of the subtile volatile smell of the plants of the scurvygrass kind, and discovers hardly any pungency to the taste; what taste it has being rather subsaline and bitterish than acrid.

BELLIS MAYOR.

BELLIS MAJOR Pharm. Edinb.(a) Bupthalmum majus; Leucanthemum vulgare; Confolida media; Oculus bovis. Bellis sylvestris caule folioso major C. B. Chrysanthemum Leucanthemum Linn. GREATER or OXEYE DAISY; a plant with oblong narrow deeply indented leaves, joined close to the stalks, which are pentagonal, hairy, branched, and bear on the tops pretty large folitary flowers composed of white petala fet round a yellow difk. It is perennial, grows wild in corn fields and dry pasture grounds, and flowers in May and June.

THE leaves of this plant have been recommended in diforders of the breaft, both afthmatical and phthifical, and as diuretics. Geoffroy relates, that the herb, gathered before the flowers have come forth, and boiled in water, imparts an acrid tafte, penetrating and fubtile like pepper; and that this decoction is an excellent diuretic and vulnerary. Either this experiment was made, not with the bellis major, but with the bellis minor; or else the herb loses its pungency when the flowers appear: the bellis minor is manifestly acrid, but in the major, when in flower, no acrimony could be observed: the leaves, whilst fresh, seemed little other than herbaceous; when dried, they discovered to the palate a not ungrateful mucilaginous fweetness.

BELLIS MINOR.

BELLIS MINOR, Consolida minima, Pharm. Edinb. (b) Bellis sylvestris minor C. B. Bellis perennis Linn. Common DAISY: a low fomewhat hairy plant, with oblong leaves lying on the ground, widening from the root to the extremity, which is rounded: among these arise round flender naked pedicles, bearing folitary flowers composed of white or purplish petala set round a yellow disk. It is perennial, common almost every where, and flowers early in the spring.

This plant stands recommended as a vulnerary, detergent, and refolvent; in difeases of the breast, internal bruises, hypochondriacal complaints, and disorders proceeding from the drinking of cold liquors when the body has been much heated. Schræder informs us, that the leaves and flowers loofen the belly.

The leaves, which have been chiefly made use of, are in taste slightly acrid. The roots are considerably stronger, of a subtile penetrating pungency, not hot or siery, but somewhat of the contraverva kind; and though at present disregarded, promise to be a medicine of no small virtue. Their pungent matter is not dissipated in drying, is dissolved both by water and spirit, and on inspissating the solutions is left in great part behind, in the watery, as well as in the spirituous extract. No part of the plant has any remarkable smell.

B E N.

BEN, sive Balanus myrepsica, Pharm. Parisiens. Behen. Glans unguentaria. Ben nut: a whitish nut, about the size of a small silberd, of a roundish triangular shape, including a kernel of the same sigure covered with a white skin. It is the produce of a middle-sized tree, said to resemble the birch, and to grow spontaneously in the East Indies and in America. It is the Guilandina Moringa of Linnæus; and the same the wood of which is called Lignum Nephriticum.

These kernels have a nauseous oily bitter taste; and when taken internally, are said to disorder the stomach, and occasion purging, sickness, and vomiting. On expression, they yield about one fourth their weight of a yellowish oil, of scarcely any particular taste or smell; the nauseous bitter matter remaining behind, not soluble in oily menstrua. This oil differs from most of the others of the expressed kind, in not being subject to grow rancid by long keeping; on account of which property, it is employed as a basis for perfumes and odoriferous unguents. It coagulates in a slight degree of cold. It is impregnated with the fragrance of jasmin and other slowers, by stratifying them with cotton dipt in the oil, and repeating the process with fresh slowers, till the oil becomes sufficiently odorous, after which it is squeezed out from the cotton in a press.

BENZOINUM.

BENZOINUM Pharm. Lond. & Edinb. Benjoinum; Benzoe; Asa dulcis. Benzoine or Benjamin: a concrete refinous juice, obtained Y

from a middle-fized tree, (Croton Bentzoë Linn.) with undivided, fomewhat oval leaves, pointed at both ends, not ribbed, falling off in the winter (a); bearing flattish nuts, about the fize of nutmegs, whose fleshy covering is externally rough and hairy (b). This tree is a native of the East Indies, where the benzoine is extracted by making deep incisions in the upper part of the trunk, about the origin of the first branches: it is said that one tree never yields above three pounds (b). The juice exudes white, and concreting on the tree becomes yellowish, reddish, or brownish; its colour turning darker, the longer it lies exposed to the air. It is brought to us in large brittle masses, composed partly of white, partly of yellowish or light brown, and often also of darker coloured pieces: that which is clearest and contains the most white matter, called by authors benzoe amygdaloides, is accounted the best.

This refin has very little tafte, impressing on the palate only a slight sweetness: its smell, especially when rubbed or heated, is extremely fragrant and agreeable. It totally dissolves in rectified spirit of wine, the impurities excepted, which are generally in very small quantity, into a deep yellowish red liquor; and in this state discovers to the taste a degree of warmth and pungency as well as sweetness. It imparts, by digestion, to water also, a considerable share of its fragrance, and a slight pungency: the filtered liquor, gently exhaled, leaves, not a refinous or mucilaginous extract, but a crystalline matter, seemingly of a saline nature, amounting, as I have found on several trials, to one tenth or one eighth the weight of the benzoine.

Flores benzoini Pharm. Lond. & Ed. Exposed to a gentle heat, in a retort or other proper vessel, it melts, and sends up into the neck white shining flowers, similar to the crystals obtained by water. These are followed by a thin yellowish oil, slightly empyreumatic, intermingled with an acidulous liquor; and at length, by a thick butyraceous matter, which, liquested in boiling water, gives out to it a little more crystalline matter, separable by filtration and proper evaporation: the whole quantity of saline matter obtainable by this method is somewhat greater than that extracted by boiling the benzoine in water. The thin oil, redistilled with water, loses its em-

⁽a) Linnæus, Species plantarum, i. 370.

⁽b) Grimm, Ad. nat. curiof. dec. ii. ann. 2. obf. 152. Rumph. Herbarium amboinenfe.

pyreumatic taint, and in this state fmells agreeably of the benzoine, and appears of the same nature with essential oils: the benzoine itself, diftilled with water, has not been observed, like most of the other refinous

juices, to yield any effential oil.

The flowers or crystals of benzoine have a grateful saline taste, and partake of the fragrance of the refin. They dissolve in spirit of wine; and, by the affistance of heat, in water; but from this last, they separate again, in great part, as the liquor cools, shooting into saline spicula, which unite together into irregular maffes: the addition of fo much fugar, as will reduce the water to the confistence of a fyrup, prevents their feparation, the flowers continuing fuspended in the fyrup after it has grown cold. Distilled with water, they arise entire, concreting into their original form, without communicating any fmell or tafte to the distilled liquor.

These flowers, unless sublimed with a very gentle heat, are apt to be tainted with an empyreumatic fmell, and a yellow colour, on account of a little of the oil being forced up with them. From this they may be purified, by mixing them with fome dry tobacco-pipe clay, and fubliming them afresh; or perhaps more perfectly, by solution in water, filtration, and crystallization. Though purified, however, to a snowy whiteness, they still participate of oil, and hence prove inflammable in the fire, and are fubject, in long keeping, to grow yellow again. They grow fooner yellow in close vessels, than in open ones; the oil, in the latter case, being perhaps carried off by the air in proportion as it is

extricated from the faline matter.

The principal use of this fragrant resin is in perfumes, and as a cosmetic; for which last purpose, a solution of it in spirit of wine is mixed with fo much water as is fufficient to render it milky, as twenty times its Lac virquantity or more. It promises, however, to be applicable to other uses, and to approach in virtue, as in fragrance, to storax and balfam of Tolu. It is faid to be of great service in disorders of the breast, for resolving obstructions of the pulmonary vessels, and promoting expectoration: in which intentions the flowers are fometimes given, from three or four grains to fifteen. The white powder, precipitated by water from folutions of the benzoine in spirit, has been employed by some as similar Magister. and fuperiour to the flowers, but appears to be little other than the pure benzoine in substance: it is not the saline but the resinous matter of

the benzoine, that is most disposed to be precipitated from spirit by water. The flowers, snuffed up the nose, are said to be a powerful errhine.

BERBERIS.

BERBERIS seu Oxyacantha galeni, Pharm. Edinb. (a) Berberis dumetorum C. B. Berberis vulgaris Linn. Spina acida; Crespinus. BARBERRY: a large prickly bush, with brittle branches, covered with an ash-coloured bark, under which lies another of a deep yellow; small, smooth, somewhat oval, pale green leaves, finely serrated about the edges, and yellow monopetalous flowers, standing in clusters on the tops upon naked footstalks, followed by oblong red berries, containing, each, generally two seeds: some of the individuals have no seeds in their berries, and sometimes berries with and without seeds are found on one bush. It grows wild on chalky hills in several parts of England, slowers in May, and ripens its fruit in September.

THE fruit of this shrub is a mild restringent acid, acceptable to the stomach, and of great medicinal efficacy in hot bilious disorders and a colliquative or putrid disposition of the humours. Prosper Alpinus informs us, that the Egyptians employ, in ardent and pestilential fevers and in fluxes, a diluted juice of the berries, prepared by macerating them in about twelve times their quantity of water, for a day or a night, with the addition of a little fennel feed or a piece of bread, and then preffing out and straining the liquor, which is sweetened with sugar, or sugar of roses, or syrup of citrons, and given the patient plentifully to drink: he fays he took this medicine himself, with happy success, in a pestilential fever, accompanied with an immoderate bilious diarrhæa (b). Simon Paulli relates, that he was cured of a like disease, by the use of syrup of barberries diluted with water; and that a concrete falt, which he calls tartar, may be obtained from the juice, by mixing two pounds of it with two ounces of lemon juice, digefting them together in a fand-heat for two days, then gently evaporating the filtered liquor to one half, and fetting it in a cellar for fome days: the tartar, he fays, incrustates

⁽a) Expunged.

the fides and bottom of the glass, proves very grateful both to the palate and stomach, and resists febrile heat and the corruption of the humours; from whence it may be presumed to be the essential acid salt of the fruit; by further inspissating the remaining juice, more of this saline substance separates in the same manner. Among us, these berries are commonly made into a gelly, by boiling them with an equal weight of sine sugar, over a gentle sire, to a due consistence, and then straining the sluid through a woollen cloth. By drying the berries, their acidity is abated, and their astringency improved.

The leaves of the barberry bush have likewise a not ungrateful restringent acid taste, and have sometimes been employed in the same intentions as the fruit, as an ingredient in cooling salads. The inner yellow bark, in taste austere and bitterish, is said to be gently purgative, and to be serviceable in jaundices: Mr. Ray commends, in this disease, from his own experience, a decoction in ale or other liquors, or rather an insusion in white wine, of the yellow bark both of the branches and the roots. It gives a deep yellow tincture both to watery and spirituous menstrua.

BETA.

BETA Pharm. Edinb. (a) Beta vulgaris Linn. BEET: a plant with large, smooth, broad-ribbed, juicy leaves; and slender, striated, branched stalks; bearing spikes of imperfect slowers standing in siveleaved cups, followed each by a roundish, rough, warty seed-vessel. Different sorts of this plant, supposed by Linnæus to be varieties of the wild beet found on some of the sea coasts of England and Holland, are cultivated in our culinary gardens: a whitish-leaved called sicula or cicla, a green-leaved, and a reddish-leaved, all with long thick white roots; and a long-rooted, and a turnep-rooted, all over red. They are all biennial.

BEETS, used as food, are difficult of digestion, and afford little nourishment: taken in quantity, they tend to loosen the belly, and are supposed by some to be prejudicial to the stomach. Their emollient or laxative virtue is extracted by boiling in water, and may be concentrated,

though not without confiderable diminution, by infpiffating the decoction. The red forts give out their colour along with their aqueous juice upon expression, and by infusion tinge rectified spirit as well as water of a deep red. The juice, both of the roots and the leaves, of the red and the white beets, snuffed up the nose, is said to be a powerful errhine, occasioning a copious discharge of mucus without provoking sneezing.

The roots of the beets have, when dry, an agreeable sweetish taste, which is totally extracted by boiling in rectified spirit: the tinctures, on standing some weeks in a cool place, deposite whitish saline concretions, of a saccharine sweetness. Mr. Marggraf observes, that the red beet loses in drying seven eighths of its weight, and the white six eighths; that the dry beet root yields nearly one twenty-sixth its weight of the saccharine salt, and the white fort one sixteenth; and that a good sugar is obtainable from the juice of the fresh roots, by the method practised abroad for preparing it from the sugar cane (a).

BETONICA.

BETONICA Pharm. Edinb.(b) Vetonica Cordi. Betonica purpurea C. B. Betonica officinalis Linn. BETONY, or wood-betony: a low plant, with dark green, oblong, wrinkled, crenated, somewhat hairy leaves, set in pairs; and square unbranched stalks; bearing thick spikes of labiated purplish flowers, each of which is followed by four oblong triangular seeds inclosed in the flower cup. It grows wild in woody and shady places, flowers in June and July, and in winter dies to the ground, the roots continuing.

This herb is recommended as a corroborant and aperient; in obfiructions of the viscera; in catarrhal, vertiginous, paralytic, hysteric, and other disorders both of the nervous and the vascular system. Its virtues have, by many practical writers, been greatly exaggerated; those of the more efficacious medicines to which it was joined, as rue, mint, cloves, guaiacum, and others, being often placed to the account of this favourite ingredient. The leaves and tops of betony have an agreeable but weak smell, which in keeping is soon distipated: to the taste, they discover a slight warmth, roughishness and bitterishness. The powder of the dry leaves, snuffed up the nose, provokes sneezing, and hence is sometimes made

an ingredient in sternutatory compositions.

Infusions of the leaves in boiling water smell and taste lightly, and not ungratefully of the herb: on inspissating them, the specific slavour of the betony is dissipated, and only a weak bitterishness and a kind of saline austerity remain in the extract. The vapour which exhales in the boiling, caught in distilling vessels, is lightly impregnated with the smell of the betony: when large quantities are distilled at once, a very small portion of essential oil separates, in colour yellowish, in taste moderately warm and pungent, and in smell pretty strong, but somewhat less grateful than the herb in substance. Spirituous tinctures, in colour deep green, discover rather less of the smell and taste of the betony than the watery insusions, though the spirit extracts all the active parts of the herb. On inspissating the filtered tinctures, a considerable part of the odorous matter exhales: the remaining extract has little smell, and a weakly pungent, bitterish, aromatic taste.

The roots are faid to be very different in quality from the other parts

of the plant; to be nauseous, bitter, purgative, and emetic(a).

BETULA.

BETULA Pharm. Edinb. (b) & C. B. Betula alba Linn. BIRCH: a tree or shrub, common in moist woods; with numerous very flexible branches; and somewhat oval, sharp pointed, serrated, deep green leaves, hanging on long and weak pedicles; producing small scaly cones, which contain little winged seeds. The bark, which appears externally white and chapt, consists of a thick brittle substance, of a dark brownish red colour, covered with three or four whitish, very thin, smooth, flexible, tough, semitransparent, membranous coats.

On deeply wounding or boring the trunk of the birch tree early in the fpring, there issues by degrees a very large quantity of a limpid, watery,

⁽a) L'Obel, Adversaria, p. 229. Ray, Hist. plant. p. 551. Caspar Hoffman, De med. officinal. lib. ii. cap. xxxvii. §. 6.

fweetish juice. It is said, that one tree will bleed a gallon or two in a day; that the juice extracted near the root is much more watery, and of less taste, than that obtained from the upper part of the trunk or from the branches; and that after the leaves have begun to appear, the juice loses its sweetness and becomes disagreeable. This juice has been drank as an antiscorbutic and deobstruent. It sensibly promotes urine, and if taken freely, loosens the belly. In keeping, it soon turns sour, unless defended from the air, by covering its surface with a little oil: by fermentation, it is converted into a weak vinous liquor. Inspissated to the consistence of a thin syrup, and set in a cool place for some weeks, it yields brownish saline concretions, approaching, as Marggraf observes, to the nature of manna.

The leaves and the bark of the tree have been employed, chiefly externally, as refolvents, detergents, and antifeptics. Simon Paulli relates, that an universal pruriginous scabies, which had been received by infection, was cured by bathing with a decoction of the bark and young branches, in which some nitre and tartar had been dissolved. With regard to the leaves, they discover to the touch a resinous unctuosity, and to the taste an unpleasant bitterness; rubbed a little, they yield a pretty strong, and not disagreeable, smell. The bark has no smell: the thin membranes have no taste; the thicker brittle part has a slight roughish one. This last gives a pale yellowish tincture to rectified spirit, and a deep yellowish red to water: the watery infusion strikes a brownish black colour with solution of chalybeate vitriol, but immediately throws off the colouring matter and becomes limpid, whereas the generality of these kinds of black mixtures retain their blackness for a length of time: on inspissating the infusion, the remaining extract proves moderately austere.

The bark of this tree has been recommended also in fumigations, for correcting contagious air. The membranes are highly inflammable, in burning yield no particular smell, and give out a resinous exudation of no smell or taste. The brittle part is less inflammable, emits a strong acid vapour, and no resin.

BEZOAR.

BEZOAR: a preternatural or morbid concretion, formed in the bodies of land animals. Several of these kinds of substances have been used

used medicinally, and distinguished either by the names of the countries from whence they are brought, or of the animals in which they are generated.

ftone: supposed to be produced in the pylorus, or in a cavity at the bottom of the fourth stomach, of an animal of the goat kind, which inhabits the mountains in different parts of Persia. It is said, that the bezoar is found only in the old animals, only in those which feed on some particular mountains, as the eastern one of the tract called Benna in the province of Laar, and only in a few of these; and that, though of great value in Europe, it is of greater in Persia itself; from whence it has been inferred, that the generality of the stones, sold under this name in Europe, must be of another original (a). Thus much is certain, that artificial compounds have been often substituted in the room of this costly concretion.

The genuine oriental bezoar of the shops is about the size of a kidney-bean (b), of a roundish or oblong rounded figure, of an even smooth surface, and of a shining olive or dark greenish colour: on being broken, it appears composed of a number of concentrical coats, of which the inner are smooth and glossy as the outer: in the middle is either a cavity, or some powdery matter, or some small bits of the leaves or stalks of plants, or other like substances. The common marks of its genuineness are, its striking a yellow or green colour on white paper that has been rubbed with chalk; a red-hot needle not piercing into it, or occasioning any bubbles, but either making no impression at all, or at most taking off only a little scale or crust; and its suffering no diminution of its weight, or disunion of its parts, by steeping in water.

The genuine stone has no manifest smell (c) or taste; and is not fensibly acted on by rectified spirit any more than by water. Reduced

⁽a) Kæmpfer, Amanitates exoticae, p. 398 & feqq. Slare, Differtation on bezoar. Neumann, Chemical works, p. 534.

⁽b) Mercatus (Metallotheca, armar. viii. cap. i. p. 173) describes a stone of this kind (presented by the king of Portugal to cardinal Alexandrinus) weighing sour ounces; so that if equally compact as the common bezoar (whose gravity is to that of water nearly as one and half to one) its volume must have been about five cubic inches.

⁽c) The flight ambergris fmell, perceivable in fome of the oriental bezoars, is supposed to be introduced by art: Cartheuser looks upon those which have this smell as being wholly factitious (Rudimenta m. m. i. 214.)

Lapis bezoar præparatus Pb. Lond. into an impalpable powder, it retains its greenish hue; which, by moistening the powder, in levigation, with a little spirit of wine, is somewhat improved. The powder agitated with water or spirit, subsides uniformly and totally; leaving no greenish matter dissolved in the liquors, as those powders do, in which the bezoar tincture has been imitated by certain vegetable matters. The powdered bezoar dissolves almost totally, and with considerable effervescence, in the acids of nitre and of sea salt; and tinges them of a deep yellow or red colour. The vitriolic acid raises a slight effervescence with it, but dissolves exceeding little. Vinegar likewise acts on it very weakly.

BEZOAR was formerly accounted a high alexipharmac; infomuch that the other medicines, possessed, or supposed to be possessed, of alexipharmac powers, have been denominated from it bezoardies. It appears, however, that this notion, adopted from the Arabian schools, has no just foundation; and that this calculous concrete, which lies inactive and indigestible in the stomach of the animal in which it is produced, is equally indiffoluble and inactive in the human stomach, unless where either a morbid acid is generated in the body, which is rarely the case in the acute difeafes wherein bezoar has been chiefly given, or acid liquors are taken along with it. Solutions of it in the nitrous or marine acids, given in doses of a few drops with proper diluents, the form in which it was used by Kæmpfer, may, doubtless, be of service in acute diseases, as antifeptic and antiphlogistic saline compounds; though not more so than folutions of the common testaceous earths. The bezoar in substance can have no other falutary operation than as an absorbent of acid humours; and appears, from experiment, to be the most weakly absorbent, or the most difficultly acted on by animal and vegetable acids, of all the earthy bodies commonly made use of in this intention. The London college nevertheless, after discarding it from the compound powder of crabsclaws, in which it was originally an ingredient, were in some measure necessitated to allow so much to ancient prejudice, as to direct, for the use of those who may still have some dependence on it, a separate officinal composition, distinguished by its name, consisting of one part of the prepared or levigated bezoar and twelve of the crabs-claw powder.

Pulv. bezoardicus Ph. Lond.

2. BEZOARD OCCIDENTALIS Pharm. Parif. Occidental bezoar: faid to be found in the stomach of an animal of the stag kind, a native

of Peru and some other parts of the Spanish West Indies. It is larger than the oriental, from the size of a walnut to that of a hen's egg or more (a): its surface is rough, and the colour less green, being often greyish or brownish without any greenness: it is likewise more brittle, of a looser texture, composed of thicker coats, and exhibits, when broken, a number of fine crystalline striæ curiously interwoven. It is less esteemed than the foregoing; though apparently not inferiour, so far as is known, in any respect that can influence its virtue as a medicine.

- 3. Lapis simiæ seu bezoard simiæ Pharm. Paris. Bezoar of the monkey: said to be found in the stomach of certain monkies; which are common in the Brasils, and in some parts of the East Indies, but which very rarely produce the admired stone. This species is about the size of a hazel nut, harder than the other bezoars, and of a very dark greenish colour almost black. Its great scarcity has rendered it of more value, and, among some, of more medicinal estimation, than the two foregoing, but prevented its having a place in the shops.
- 4. CALCULUS HUMANUS, bezoar microcosmicum quibusdam dictus. The calculus of the human bladder. This concrete is various in degree of hardness, as well as in appearance, figure, and size: the softer masses are for the most part pretty easily, the harder more difficultly, dissolved, in part at least, by acids, and corroded by soap leys and lime water: other menstrua for them are not known. Some have employed this stone as a succedaneum, and even in preference, to the foregoing costly bodies, and report that they have found it to act as an excellent sudorisic and diuretic (b); ascribing to the stony matter the effects of the theriaca, oil of amber, and oil of juniper berries, with which it was joined.
- 5. LAPIS PORCINUS Pharm. Parif. Lapis malacensis. Bezoar hystricis. Pedro del porco. Bezoar of the porcupine: said to be found in the

⁽a) There are accounts of occidental bezoars of much larger fizes: Mercatus (ubi supra, p. 174, 175) describes and figures a stone of this kind weighing no less than fifty-fix ounces, though part of the outer crusts had been removed.

⁽b) Bontius, Animadversiones in Garciam, lib. i. cap. 46.

gall-bladder of an Indian porcupine, particularly in the province of Malacca. It is of a roundish figure, of a pale purplish colour, of a soft substance, smooth and slippery to the touch (a). This concrete is of a very different nature from the four preceding: it has an intensely bitter taste, and, on being steeped in water for a very little time, impregnates the sluid with its bitterness, and with aperient, stomachic, and, as is supposed, with alexipharmac virtues. How far it differs in virtue from the similar concretions found in the gall-bladder of the ox and other animals, does not appear.

BISMUTHUM.

BISMUTHUM five marcasita Pharm. Paris. BISMUTH of TINGLASS: a bright whitish pulverable metal; near ten times specifically heavier than water; melting long before ignition, a little sooner than lead, and a little later than tin; sublimable, by a strong fire, into white slowers; calcining, by a continuance of a heat sufficient to keep it melted, into a greyish powder, which on raising the fire runs into a yellow very susple glass; dissolving with violence in the nitrous acid, and precipitating in form of a bright white powder on diluting the solution largely with pure water; very dissicultly acted on by the marine acid, and scarcely at all by the vitriolic; giving out very little to the vegetable acids, but impregnating them with a nauseous taste. It is extracted from an ore found hitherto chiefly in Saxony; by eliquation, or suspendent in a small heat without addition. The ore is generally very arsenical: whether the bismuth retains any of the arsenic, has not been sufficiently examined.

This metal remarkably promotes the facility and tenuity of the fusion of other metallic bodies: with lead and tin, it forms compounds which melt in so small a heat as to have been proposed by some for anatomical injections: the proportions, that have been found to compose the most suspections are, two parts of lead, three of tin, and sive of the bismuth. It likewise remarkably promotes the solution of lead in mercury, but has not been observed to produce a like effect on other metallic bodies.

Magiste-

The white flowers fublimed from this metal, and the white magistery Flor. bisprecipitated by water from its folution in aqua fortis, have been recommended externally against gleeting fores, and internally as diaphoretics rium biffimilar to the milder antimonial medicines. In the first intention, they appear to be greatly inferiour to some of the saturnine preparations: in the latter, it is not certain what their real effects are, or even whether they are fafe. At prefent, they are employed only as a fucus, nor is this use of them entirely innocent; for they gradually impair the natural complexion, and, as the college of Strafburg observes, occasion a thickness and defedation of the skin.

BISTORTA.

BISTORTA Pharm. Lond. & Edinb. Bistorta major radice minus intorta C. B. Polygonum Bistorta Linn. BISTORT OF SNAKEWEED: a plant with oval, pointed, wrinkled leaves, of a dark green colour above and blueish underneath, standing on long pedicles, and continued a little way down the pedicle, forming a narrow margin on each fide: among these arise round, slender jointed, unbranched stalks, furnished with fmaller and narrower leaves which have no pedicles; bearing on the top fpikes of imperfect five-leaved red flowers, which are followed by triangular feeds. It is perennial, grows wild in moist meadows in feveral parts of England, and flowers in May and June.

THE root of this plant is bent and jointed, commonly about the thickness of the finger, surrounded with bushy fibres, of a blackish brown colour on the outfide, and reddish within: it is distinguished from the roots of the other bistorts, by being less bent; that of the officinal fpecies having only one or two bendings, and those of the others three or more.

This root has a strong astringent taste, without any manifest smell or particular flavour. It is one of the strongest of the vegetable styptics, and frequently made use of as such, in disorders proceeding from a laxity and debility of the folids, for restraining alvine fluxes, after due evacuations, and other preternatural discharges both serous and sanguineous. It has been fometimes given in intermitting fevers; and fometimes, also, in small doses, as a corroborant and antiseptic, in acute malignant and colliquative colliquative fevers; in which intentions, Peruvian bark has now de-

fervedly superfeded both this and all the other astringents.

The common dose of bistort root, in substance, is fifteen or twenty grains: in urgent cases, it is extended to a dram. Its astringent matter is totally dissolved both by water and rectified spirit*(a); the root, after the action of a sufficient quantity of either menstruum, remaining insipid: the watery tinctures are of a dark brownish colour, the spirituous of a brownish red. On inspissating the tinctures, the water and spirit arise unslavoured, leaving extracts of intense stypticity.

BITUMEN JUDAICUM.

BITUMEN JUDAICUM Ph. Lond. Affibaltus. Jews PITCH: a folid, light bituminous substance; of a dusky colour on the outside, and a deep shining black within; of very little taste, and scarcely any smell, unless heated, in which circumstance it emits a strong pitchy one; not soluble in vinous spirits or in oils; difficultly and only imperfectly melting in the fire; and leaving, on being burnt, a large quantity of ashes. It is said to be found plentifully in the earth in several parts of Egypt, and sloating on the surface of the dead sea; at first soft, and growing hard by age.

ABUNDANCE of virtues are attributed to this bitumen; refolvent, discutient, sudorific, emmenagogue, and others. It has long, however, been disregarded in this country: the college of Edinburgh has now expunged it from the catalogue of officinals, and that of London retains it only as an ingredient in one of the compositions which complaisance to antiquity has preserved in the shops. Nor is it, among us, to be often met with; its place being generally supplied by different bituminous substances found in France, Germany, and Switzerland, sometimes by the caput mortuum remaining after the distillation of amber, and sometimes by common pitch. Its melting in the fire only partially or not at all, and the quantity of ashes it leaves in burning, distinguish it from these substances, and shew, at the same time, that in its most genuine and perfect state it is a very impure bitumen, mixed largely with

earthy matter. Distilled in a retort, it yields, according to Neumann, a little insipid phlegm, and about one eleventh its weight of oil, refembling the native petrolea, but of a somewhat more disagreeable empyreumatic smell.

BOLUS.

BOLE: a friable earthy substance, uniting with water into a smooth paste, adhering to the tongue, and dissolving as it were in the mouth: of the clayie kind, but more readily imbibing water than the clays strictly so called; when moistened, less viscous and cohesive; more easily diffusible through water by agitation; and more freely subsiding from it.

- 1. Bolus Armena Pharm. Lond. Armenian bole, or bole-armenic: of a pale but bright red colour, with a tinge of yellow; harder, and of a less glossy surface, than most of the other boles.
- 2. Bolus Gallica Pharm. Lond. & Edinb. French bole: of a pale red colour, variegated with irregular specks and veins of whitish and yellow.

Many other bolar earths have been recommended for medicinal uses, and were formerly ranked among the officinals; as, red boles from Armenia, Lemnos, Strigonium, Portugal, Tufcany, and Livonia; yellow boles from Armenia, Tockay, Silefia, Bohemia, and Blois; white boles from Armenia, Lemnos, Nocera, Eretria, Samos, Chio, Malta, Tuscany, and Goltberg. Several of these earths have been commonly made up into little cakes or flat masses, and stamped with certain impresfions; from whence they received the name of terræ figillatæ, or fealed earths. The Armenian and Lemnian have been generally supposed to be the best, but are rarely met with in the shops: the common French bole, and fome bolar earths found in our own country, and even white clay artificially coloured with ochre or colcothar of vitriol, have commonly fupplied the place both of those and of the other coloured boles. The fubflitution of the French to the Armenian, in the feveral compositions wherein that earth is directed as an ingredient, is now allowed by the London college: and indeed all these earthy bodies, however differing from one another in the degree or species of their colour, or in their

texture

texture and compactness, appear, in regard to their medicinal qualities, to be very nearly, if not entirely, alike.

ALL the boles have for their basis one and the same argillaceous earth; which is not dissoluble, by the heat of boiling water, in acids, in alkalies, or in any other known menstruum; which, in a strong heat, grows hard, contrary to all the other bodies of an earthy or stony nature, which receive from fire a greater or less degree of friability; and at the same time loses its property of imbibing water, and of being reduced thereby into a tenacious mass. The boles and clays, both in their natural state and when indurated by fire, become dissoluble in part, by strongly boiling them in the concentrated vitriolic acid, till the more phlegmatic parts of the liquor have exhaled, and the matter remains dry. The compound which the earth, by this process, forms with the acid, is of the same nature with alum: it dissolves in water, and may be crystallized into perfect alum, by adding a suitable quantity of any volatile or fixt alkaline salt to saturate the redundant acid, and after due evaporation setting the liquor to shoot.

The colours of the boles proceed from a flight admixture of a ferrugineous calx; which may be extracted by digestion in spirit of salt
or aqua regis, but is scarcely acted upon by any acid of the vegetable or
animal kingdom. Some of them contain a portion of calcareous earth,
which is extracted by all acids except the vitriolic, and discovers itself
by raising an effervescence on the affusion of the acid. The specimens
I examined of the bole of Blois gave out a considerable quantity of this
earth, those of the common French bole exceeding little, and the Armenian none: possibly, however, different masses of one kind of bole
may differ, in this respect, as much as different boles. All the boles
seem to participate also of vitriolic acid; which is so intimately blended
with the other matter, as not to be separable, or discoverable, without
violence of fire.

The ferrugineous calx and calcareous earth are likewise very intimately blended with the proper bolar matter; insomuch, that when the compound is diffused through water, it settles equally and uniformly without any separation of its parts. If the bole contains any sand or small stones, or has been artificially coloured, the sand, stones, and colouring ingredients, separate in the water, and being heavier than the bolar earth, subside before it. On this principle, the boles may be purified from the

gritty

gritty matter often intermixed among them, and the natural boles dif-

tinguished from artificial compositions.

The medical virtues of the boles appear to depend on the simple bolar or argillaceous earth. As this earth is not disfoluble by any sluid that can exist in the bodies of animals, it can act no otherwise than by imbibing, or giving a greater degree of consistence to thin sharp humours in the first passages, and in some measure defending the solids from their acrimony. In consequence of this virtue, the boles may be of some service in alvine sluxes, cardialgic complaints, and in some kinds of acute diseases; though they are not possessed, as they have been commonly supposed to be, of any truly aftringent, or absorbent, and much less of any alexipharmac powers. The sensation of astringency which they generally occasion, in some degree, in the mouth, seems to consist only in their adhering to and drying the part, by imbibing the fluids that moisten it. Their dose is from fifteen or twenty grains to a dram.

BONUS HENRICUS.

BONUS HENRICUS, five Lapathum unctuosum, Pharm. Edinb. (a) Chenopodium Pharm. Paris. Tota bona Dod, Lapathum unctuosum folio triangulo C.B. Chenopodium Bonus Henricus Linn. English mercury: a plant with triangular leaves, covered underneath with a whitish unctuous meal; and striated hollow stalks, partly erect, and partly procumbent, bearing on the top spikes of small imperfect flowers, each of which is followed by a small black seed inclosed in the cup. It is perennial, grows by road sides and in waste grounds, and flowers in August.

THE leaves of this plant, to the taste mucilaginous and somewhat unpleasantly subsaline, are accounted emollient; and in this intention have been made an ingredient in decoctions for glysters. They are applied by the common people to slesh wounds and sores, under the notion of drawing and healing. In some places, the young shoots are eaten in the spring as asparagus, and said to loosen the belly, and promote urine.

(a) Expunged.

BORAGO.

BORAGO Pharm. Edinb. (a) Borago bortensis Raii synops. Buglossum latifolium borrago store cæruleo C. B. Borago officinalis Linn.
BORAGE: a very hairy rough plant, with wrinkled blackish green leaves
approaching to an oval shape, and round hollow stalks, on which the
leaves are set alternately: on the tops of the branches come forth blue,
sometimes reddish or whitish, monopetalous slowers, each of which is
divided into sive sharp-pointed segments, and sollowed by sour wrinkled
blackish seeds lying naked in the enlarged cup. It is perennial, and
grows wild on waste grounds and on old walls.

THE leaves of this plant are very juicy, of no fmell, and of hardly any particular tafte: they feem nevertheless to contain substances of some medicinal activity, though in too small proportion to be sensible till separated from the herbaceous matter. Mr. Boulduc relates, that a decoction of borage leaves, evaporated to the confistence of a syrup, and set by for a few days, yielded faline crystals, partly in form of fine needles, and partly cubical: that the needled crystals were found to be perfect nitre, and the cubical fea falt: that by paffing the decoction through quicklime before the inspissation, both salts were obtained in greater purity and in larger quantity: that the substance of the leaves, remaining after the boiling, being dried and burnt, and the ashes elixated with water. the lye, properly evaporated and fet to shoot, yielded first a vitriolated tartar, and afterwards fea falt, the liquor, after the crystallization, proving fimply alkaline (b). From this analysis it may be presumed, that the aperient and refrigerating virtues, ascribed to borage leaves, are not wholly without foundation; though these virtues are undoubtedly very weak. Malouin remarks, that the juice of the leaves, which is not green, like that of most other herbs, but of a brown colour, added to bitter mixtures of the juices of creffes and chervil, diffipates their bitterness.

The flowers of the plant have been principally made use of, and are generally ranked among the cordials. Medicines may act as cordials, either by virtue of some warmth, pungency, or fragrance; or by a saline quality, abating immoderate heat: but borage flowers seem to have little

power of operating in either of these intentions. When fresh, they have a very flight smell, of the agreeable kind, which in drying is lost: to the taste, both the flowers in substance, and an extract made from them by water, are only mucilaginous and fweetish.

BORAX.

BORAX Pharm. Lond. Borax, Tincar, Ph. Edinb. BORAX: a crystalline salt; very difficultly soluble in cold water; swelling and bubbling up in the fire, and changing into a light white fpongy friable matter, which, foon fubfiding on a continuance of the fire, melts into a fubstance resembling glass, but which is still found to be dissoluble in water, though more difficultly than the borax at first.

It is brought from the East Indies in a very impure state; confisting partly of large hexaedral prismatic flatted crystals, but chiefly of smaller and more irregular ones, partly whitish and partly green, joined together as it were into one lump by a fetid greafy, or oily, yellow fubstance; intermingled with fand, small stones, and other impurities. Of its origin

and preparation we have no certain account.

This impure borax was formerly refined at Venice, afterwards in Holland only, and now by fome particular perfons in England also, into large irregular colourless masses, in appearance resembling alum. The falt is commonly called tincal in its rough state, and borax when thus purified or refined. The method of refining it is kept a fecret. Certain additional matters are fuspected to be employed; the common refined borax being different in some respects, particularly in its power of vitrefying earthy bodies, from the crystals unrefined or simply purified by folution.

THE purer crystals of tincal, or refined borax, disfolve, by boiling, in a finall quantity of water, fo as in cooling to concrete almost all into a folid mass, only a very little liquid remaining on the top: to keep them disfolved in the cold, more than fourteen times their weight of water is necessary. On boiling the impure tincal in water, the oil diffolves along with the falt into a foapy liquid; from whence it may be prefumed that the oily matter is not of a mineral, but of a vegetable or animal origin. From this folution it is very difficult to separate the oil,

Aa2

without additions which alter the quality of the falt: but if the rough tincal be previously heated in an iron ladle, or other convenient vessel, till it ceases to bubble and slame, the oil is destroyed or made indissoluble, and boiling water extracts from the black mass only the pure borax.

Pure borax has a fweetish somewhat pungent taste, leaving in the mouth an impression like that of alkaline salts, but far milder. Like alkaline falts also, it changes the colour of blue flowers to green, precipitates earthy and metallic bodies diffolved in acids, and renders vegetable and animal oils miscible with water: it does not, however, sensibly effervesce with acid any more than with alkaline liquors. It dissolves in acids more easily than in water, and promotes likewise the solution of fome vegetable acid falts of themselves difficultly dissoluble. A mixture of borax with twice its weight of tartar diffolves in about one fixth part of the quantity of water that would be necessary for their solution separately: the liquor tastes acid, like tartar by itself, and deposites a confiderable quantity of tartar in cooling. About equal parts of the two form a compound perfectly neutral, in taste more like borax than tartar, which is kept diffolved by five times its weight of water a little above freezing. On inspissation, a viscous tenacious mass is left, which does not crystallize, and which deliquiates in the air. Borax affords also glutinous compounds with all the other acids except the vitriolic: faturated folutions even of the borax by itself are confiderably fo.

This falt appears to confift of the mineral alkali or basis of sea falt, united with a smaller proportion of a peculiar saline subacid concrete. By all the mineral acids, and, as is said, by the acetous, its constituent parts are separable from one another; the acid uniting with the alkaline basis, and disjoining therefrom the subacid ingredient of the borax.

This analysis is most commodiously effected by the vitriolic acid. A mixture of nine parts of borax, three of oil of vitriol, and one of water, being urged, in a wide-necked retort, with a fire at first gentle and afterwards pretty hastily increased till the vessel becomes red hot; the subacid salt of the borax, called sedative salt, rises into the neck, and concretes into thin shining white plates. But as this salt proves volatile only while moist, a part of it remains behind, and may be sublimed, like the first, by pouring back on the residuum the liquor that distills, and renewing the operation. The same salt may be obtained more commodiously, though scarcely in so pure a state, by adding the oil of vitriol

Sal. fedativ. Phar. Parif.

to the borax diffolved in water, and, after due evaporation, fetting the mixture to shoot: the sedative salt crystallizes on the surface, much fooner than the other faline matter, into thin plates; which, uniting together, and growing heavier, fall to the bottom. The falt, which in either case remains after the separation of the sedative salt, is a combination of the vitriolic acid with the alkaline basis of the borax, and has not been observed to differ from the common combination of that acid with the alkaline basis of sea salt, that is, from the sal mirabile or cathartic falt of Glauber. The fedative falt, joined to the marine alkali, recomposes borax again.

The peculiar and characteristic ingredient of the borax, though called fubacid from its property of neutralizing alkalies, fcarcely difcovers any other mark of acidity. Its taste is bitterish, accompanied with a slight impression of coolness. It makes no change in the colour of blue flowers, and no effervescence with alkalies or with acids. It melts in a moderately ftrong fire, and affumes a perfect vitreous appearance; but this apparent glafs, as well as the falt itself, may be totally sublimed, if repeatedly moistened, by a less degree of heat; and totally, though difficultly, dis-

folved both by water and by rectified spirit.

It is observable, that the spirituous solution of the sedative falt, set on fire, burns with a green flame; and that borax itself, boiled in spirit, is partially diffolved, and tinges its flame of the same colour. Perhaps it is principally, or folely, this falt, that the spirit extracts from the borax; for spirit burnt on the alkali of borax, exhibits no greenness.

Borax is accounted an efficacious deobstruent, diuretic, emmenagogue, and promoter of delivery. Its virtues have not as yet been thoroughly afcertained by experience, and are by many questioned; the borax having generally been given in conjunction with other fubstances, to which the effects, experienced from the compound, may be, in part at leaft, attributed. Thus, in the powder for promoting delivery, half a dram of Pulvis ad the borax, the mean dose of this falt, is accompanied with eleven grains partum Ph. and a quarter of faffron, the same quantity of castor, a drop of oil of cinnamon, and three fourths of a drop of oil of amber. That the borax itself, however, has really some virtues of this kind, may be presumed from the effects it has been observed to produce when used in large quantity: Trioen relates, that an ounce and a half of borax having been taken by a young woman in mistake for creme of tartar, an uterine

hemorrhage succeeded, so profuse, that life was despaired of: the flux was got under by medicines; but the ill state of health, and almost universal ædema, which followed it, were lasting (a). Solution of borax has been found to be a powerful dissolvent of aphthous crusts in the mouth and sauces of children (b).

The peculiar faline concrete, extricated from borax by acids, is supposed to be antispassmodic and anodyne, whence its name sedative salt. It is said to calm the heat of the blood in burning severs, to prevent or remove delirious symptoms, and allay, for a time, melancholical, hypochondriacal, and hysterical complaints. It continues in some esteem in France, where it was first discovered by Mr. Homberg, but has never come into practice among us. Its dose is from three grains to a scruple.

BOTRYS.

BOTRYS: a low fomewhat hairy plant, full of branches, bearing numerous imperfect flowers: the flower stands in a five-leaved cup, which forms a covering to a small roundish seed.

- I. Botrys Pharm. Edinb.(c) Botrys sive ambrosia Pharm. Paris. Botrys ambrosioides vulgaris C.B. Chenopodium Botrys Linn. Jerusalem oak: with oblong pointed, deeply sinuated leaves like those of the oak tree, of a yellowish green colour on the upper side, purplish underneath, marked with large red veins, and placed alternately: the flowers stand in clusters, on divided pedicles, in the bosoms of the leaves.
- 2. BOTRYS MEXICANA Pharm. Parif. Botrys ambrosioides mexicana C. B. Chenopodium ambrosioides Linn. Mexico tea: with pale green triangular leaves, and undivided flower-stalks: greatly resembling the wild orach.

BOTH these plants are natives of the southern parts of Europe, and sown annually with us in gardens. The leaves and slowery heads have a pretty strong and not unpleasant smell, and a moderately aromatic somewhat bitterish taste: on much handling them, an unctuous resinous

⁽a) Observationum medico-chirurg. fasciculus, p. 18. (b) Bisset, Essay on the medical constitution of Great Britain, p. 203.

juice adheres, in considerable quantity, to the singers. The proper menstruum of their active matter is rectified spirit: they nevertheless give
out their more valuable parts to boiling water also, which they impregnate strongly with their smell, and considerably with their taste.
The insusions, which are not unpalatable, drank as tea, are said to be
of service in humoural asthmas and coughs, and other disorders of the
breast: they are supposed also to be antispasmodic and antihysteric.

BRASSICA.

BRASSICA Pharm. Edinb. (a) CABBAGE. Brassica capitata alba C.B. White cabbage and coleworts. Brassica capitata rubra C.B. Red cabbage. Brassica rubra C.B. Red coleworts. Brassica sabauda: brassica alba capite oblongo non penitus clauso, C.B. Savoy. Brassica caulissora C.B. Caulislower. These and the other forts of cabbages, raised in our culinary gardens, are supposed to be only varieties of the smaller kind, which, in some parts of England, about the sea coasts, is found wild: accordingly they are joined by Linnæus into one species, under the name of brassica (oleracea) radice caulescente tereti carnosa. They are all biennial.

The feveral forts of cabbages are commonly accounted hard of digestion, and of little nourishment, but perhaps not very justly. For as they have manifestly a strong tendency to putrefaction, running into this state sooner than almost any other vegetable, and emitting also during the putrefaction a more offensive smell, nearly approaching to the fetor of the animal kingdom; it does not seem irrational to presume, that of all the oleraceous herbs, cabbages may be the most easily resoluble in the stomach, the most nutritious, and the least remote from the nature of animal food. Thus much is certain, that they are, in general, not unwholesome; that they do not induce, or promote, a putrid disposition in the human body, but on the contrary prove a salubrious aliment in the true putrid scurvy*(b); that when taken freely, they tend to loosen

(a) Expunged.

^{*(}b) Cabbages cut in slices and packed up in a cask with salt and other additions, ferment, and acquire an acid taste, in which state they are much eaten in Germany under the name of faur kraut. This has lately been introduced as an article of diet in the British navy, and has been found to keep well in long voyages, and to prove a very useful antiscorbutic.

the belly and produce flatulencies; and that their laxative matter is ex-

tracted by long boiling in water.

Of all these herbs, the white cabbage is the most putrescible; and the red sweetest, and most emollient or laxative. If the stalks of red cabbage, towards the end of autumn, be cut longitudinally and set by for some time in a place not warm, a laxative juice, resembling honey or manna, exudes from the incisions (a). A decoction of this last kind has been greatly recommended in hoarsenesses and some disorders of the breast, for softening acrimonious humours, and promoting expectoration. Boerhaave tells us of very dangerous disorders of the chest cured by the use of a decoction of red cabbage with a little salt and orange juice (b).

BRASSICA MARINA.

BRASSICA MARINA, foldanella, Pharm. Edinb. (c) Soldanella & convolvulus maritimus Ph. Parif. Soldanella maritima minor C. B. Convolvulus Soldanella Linn. Soldanella: a species of convolvulus, with roundish or kidney-shaped leaves set on long pedicles; and large reddish-purple flowers standing solitary in the bosoms of the leaves. It grows wild about the sea coasts in the north of England, and slowers in June.

THE leaves of foldanella are faid to be a strong and a rough cathartic, generally occasioning gripes and disordering the stomach. Their virtue resides in an acrid, bitterish, milky juice, which exudes upon wounding them. In drying, they lose much of their taste, but seem to retain their purgative virtue: a decoction of from half a dram to two or three drams of the dried leaves is directed for a dose. In some places, it is customary among the common people to prepare a strongly purgative liquor by boiling a small handful of the fresh leaves in broth. But as their degree of strength is very little known, and as we have many other medicines, for the same intention, whose qualities have been ascertained by general experience, the soldanella, though retained in most catalogues of the materia medica, stands excluded from practice.

(a) Hoffman, De remediis domesticis, §. 14.

(b) Boerhaave, Hist. plant. bort. Lugd. Bat. p. 423.

(c) Expunged.

BRYONIA.

BRYONIA.

J. B. Bryonia aspera sive alba baccis rubris C. B. Bryony or wild vine: a perennial rough plant, growing wild in hedges, and climbing on the bushes with curled tendrils. The leaves are in shape somewhat like those of the vine, irregularly pentagonal, with a considerable indentation between every two angles, and the extreme segment longest: in their bosoms come forth clusters of greenish-white bell-shaped monopetalous slowers, divided into five roundish sections, adhering firmly to the cup; succeeded by red berries, containing an extremely viscid pulp with small seeds. The root is very large, sometimes as thick as a man's thigh, of a yellowish or brownish colour on the outside, and white and fungous within.

FRESH bryony root, taken up in the beginning of spring, abounds with a thin milky juice: if the upper part of the root be bared of earth, and the top cut over transversely, the juice continues to rise gradually to the surface, in notable quantity, for two or three days successively, and may be collected by forming a cavity in the middle to receive it. Both the root in substance, and the juice, have a disagreeable smell, and a nauseous, bitter biting taste: applied for some time to the skin, they inflame or even vesicate the part. On drying the one, or inspissating the other, they lose most of their acrimony and nearly the whole of their ill scent. In summer, the root proves much less juicy, and weaker both in smell and taste.

This root, taken in powder from a scruple to a dram, proves a strong cathartic. It was formerly given, both as a purge, in maniacal and hydropic cases; and, in smaller doses, as a resolvent and deobstruent, in uterine and asthmatic disorders, in which it is reported to have been of great efficacy. At present it is, in this country, very rarely made use of in either intention; on account, not entirely of the violence of its operation, for purgatives as violent as it are still retained in practice; but partly of its degree of activity, in different states and forms, being very variable, and less ascertained than that of other cathartics in more general use. It is said, that when fresh it operates, violently, upwards as well as downwards; and that when dry, it acts with less violence, and B b

chiefly by flool (a); that the juice, which issues from it spontaneously, purges mildly in doses of a spoonful (b); that the fresh root, bruised and freed from its juice by pressure, and afterwards dried, is still purgative (c); and that the expressed juice exsiccated, and the farinaceous fecula which subsides from it on standing, are of little activity (d): that an extract made from the fresh root, by boiling it in wine, and pressing out and inspissating the decoction, operates with sufficient mildness, in doses of from half a dram to a dram (e), both by stool and urine; and that infusions in water are chiefly diuretic. Burggrave relates, as from his own knowledge, a pretty remarkable account of the effects of the watery infusion and juice in this last intention. From a fresh root, as thick as can be procured and about a span long, he directs about an inch of the top to be cut off, and a large conical piece to be cut out to two thirds the depth of the root: into this cavity put two ounces or more of fugar-candy in powder, above which infert the cone properly detruncated, and fet the root upright in a warm place for twenty-four hours: the fugar being now diffolved by the native juice of the bryony, the excavated part of the root is to be cut off, and one, two, or three flices, from the lower folid part, infused in water. "Give, says the "author, to an hydropic person, one spoonful of the saccharine solution "in the morning, and repeat it every two hours till the patient begins "to make water profusely, for it will not purge: when great thirst is " complained of, give a draught of the infusion, which will likewise not " purge, but work still more by urine. Then carefully provide against " any ill effects ensuing from the inanition of the abdomen and collapsion " of the integuments (f)."

Externally, the fresh root has been employed in cataplasms, as a refolvent and discutient; against hard and ædematous tumours, stagnations

⁽a) Hermann, Cynosura m. m. edit. Boeeler. i. 141, &c. — Boulduc's opinion, of the dry root being strongest (mem. de l'acad. roy. des scienc. de Paris, pour l'ann. 1712) seems to have been deduced from a principle, which cannot be admitted, that the root suffers no other change in drying than the dissipation of its watery humidity.

⁽b) Stoffelius, apud Joan. Baubin. bift. plant. tom. ii. p. 143.

⁽c) Le Mort, Morley collect. chym. Leydenf. p. 120.

⁽d) Boerhaave, Hift. plant. Lugd. Bat. p. 497. Geoffroy, m. m. iii. 223.

⁽e) Geoffroy, m. m. iii. 223.

⁽f) Burggrave, Lexicon medicum, p. 1710.

and coagulations of blood from external injuries, and ischiadic and other rheumatic pains.

BUGLOSSUM.

BUGLOSSUM SATIVUM Pharm. Edinb.(a) Buglossim angustifolium majus C. B. Anchusa officinalis Linn. Bugloss: a rough plant,
greatly resembling borage, and differing from it chiefly in the leaves being
narrow, less prickly, not wrinkled, and in colour blueish green; and
in the segments of the flowers being obtuse. It grows wild, on waste
grounds, in the southern parts of Europe, is cultivated with us in
gardens, flowers from June to the end of summer, and in winter dies
to the ground, the roots abiding.

This plant appears to be nearly fimilar to borage, in its medicinal qualities as well as in its external form. The principal difference feems to confift, in the leaves being fomewhat lefs juicy, and the roots more mucilaginous. The roots, leaves, and flowers, are ranked among the articles of the materia medica, but are very rarely made use of.

BUGULA.

BUGULA, consolida media, Pharm. Edinb. (b) Prunella germanis Trag. Consolida media pratensis cærulea C. B. Ajuga reptans Linn. Bugle or MIDDLE CONSOUND: a low plant, with two kinds of stalks; round creeping ones, which strike root at the joints; and upright square ones, hairy on two of the opposite sides, alternately, from joint to joint, bearing loose spikes of blue labiated flowers, of which the upper lip is wanting: the leaves are somewhat oval, soft, slightly cut about the edges, and set in pairs at the joints. It is perennial, found wild in woods and moist meadows, and flowers in May.

THE leaves of bugle discover, on first chewing, a sweetish taste, which is followed by a considerable bitterishness and roughishness. Insusions of them, or the expressed juice, are recommended as vulneraries, or as mild astringents and corroborants, in sluxes and other disorders. Some

(a) Expunged.

(b) Expunged.

B b 2

have

have observed, that they do not bind the belly, like the other consolida, but that, on the contrary, decoctions of them are gently laxative, and of great use in phthises and internal ulcerations (a). The roots of the plant are considerably astringent, as appears both from their taste, and from their striking a black colour with solution of chalybeate vitriol.

BURSA PASTORIS.

BURSA PASTORIS Pharm. Parif. Bursa pastoris major folio sinuato C. B. Thlaspi Bursa Pastoris Linn. Shepherds purse: a plant with small tetrapetalous whitish flowers, along the upper part of the branches, followed each by a triangular seed vessel resembling a purse, whence its name: the lower leaves are for the most part deeply jagged like those of dandelion, and widen from the bottom forwards; those on the stalks are entire, and most of them broadest at the bottom, with a little ear on each side at the juncture with the stalk. It is annual, common in waste grounds, and flowers from April to the end of summer.

This herb has, when fresh, an unpleasant smell, which in drying is dissipated: its taste is almost merely herbaceous. An extract made from the dry leaves by water is somewhat ungratefully mucilaginous and subsaline: an extract made by rectified spirit has somewhat more of an unpleasant, though weak, taste. No pungency or astringency could be perceived either in the leaves themselves or in the extracts; nor did a decoction of them strike any degree of blackness with solution of chalybeate vitriol. There does not appear, therefore, to be any soundation for the strong styptic virtues, for which this herb has been generally recommended by writers on the materia medica; or for the acrid in-slammatory power, which some (misled probably by its botanic affinity with mustard and some other acrid vegetables) have ascribed to it.

BUXUS.

BUXUS Pharm. Lond. Buxus arborescens C. B. Buxus sempervirens Linn. Box TREE: a small evergreen tree, or shrub, with numerous branches, clothed with firm, shining, somewhat oval leaves: the

(a) La Poterie (Poterius) Pharm. spagyr. lib. i. sett. i. cap. 2.

wood is of a yellow colour, and more compact and ponderous than any of the other European woods: the flowers are imperfect; the fruit, which grows on a distinct part of the tree, is a green berry, divided into three cells, containing fix small seeds. It is found wild in some parts of England.

THE leaves of this tree have a faint unpleasant smell, which is in great part distipated in drying, and pretty strongly impregnates water in distillation: their taste is somewhat of the bitter kind, very strong, and very nauseous. It is said, that their effluvia are narcotic; and that the leaves in substance, and insusions or decoctions of them, are aperient and purgative.

The wood gives a bright yellow tincture to spirituous menstrua, and a paler yellow to water. Chewed, in substance, it scarcely discovers any taste: an extract made from it by rectified spirit, which amounts to little more than one sifteenth part of the weight of the wood, is weakly bitterish: by water was obtained nearly one thirtieth its weight, of a stronger tasted, ungratefully saline extract. From these experiments it may be presumed, that boxwood contains little active matter; and that this matter is not of the pungent resinous, but of the saline kind; and consequently that it differs greatly from guaiacum wood, to which it is by many writers supposed to be similar.

CACAO.

CACAO Pharm. Lond. CACAO or CHOCOLATE NUT: an oblong roundish nut, nearly of the shape of an almond, but larger: the shell is dark coloured, brittle, and thin: the kernel is both externally and internally brownish, divided into several unequal portions, which are joined firmly together. It is the produce of a small American tree, (Cacao Sloan. Jam. Amygdalis similis guatimalensis B. C. Theobroma Cacao Linn.) bearing a large red fruit shaped like a cucumber, which contains thirty or more of the nuts. There are several sorts of these nuts in the shops, distinguished by their size, and the places whence they are brought: the larger kind, from the province of Nicaragua in Mexico, is most esteemed.

CACAO nuts have a light agreeable smell, and an unctuous, bitterish. roughish, not ungrateful taste: those of Nicaragua and Caracco are the most agreeable; those of the French Antilles, and our own American islands, the most unctuous. All the forts, thoroughly comminuted and committed to the press, yield a considerable quantity of a fluid oil, of the same general qualities with those obtained from other feeds and kernels: boiled in water, they give out a large proportion, half their weight or more, of a febaceous matter, which gradually concretes upon the furface as the liquor cools. For obtaining this product to the best advantage, the faculty of Paris directs the nuts to be flightly roafted in an iron pan, cleared from the rind and germ, levigated on a hot stone, then diluted with a proper quantity of hot water, and kept in a waterbath till the oil rifes to the top; which, when concreted, looks brown, and by repeated liquefactions in hot water becomes white. This vegetable fevum is not liable to grow rancid in long keeping; and hence is recommended as a basis for odoriferous unguents, and the compositions called apoplectic balfams.

Oleum feu butyrum e nucleis cacao Pharm. Parif.

The principal use of these nuts is for the preparation of the dietetic liquor, chocolate; a mild unctuous fluid, supposed to be serviceable in consumptive disorders, emaciations, and an acrimonious state of the juices in the first passages.

CALAMINARIS.

LAPIS CALAMINARIS Ph. Lond. & Edinb. Cadmia fossilis; Cadmia lapidosa. Calamine or Calamy: a mineral substance, of a greyish, brownish, yellowish, or pale reddish colour, and sometimes of all these colours variously mixed; considerably heavy; and moderately hard, but never sufficiently so to strike fire with steel; when mixed with powdered charcoal, changing copper, by sustain, into a yellow metal, called brass. It is found plentifully in England, Germany, and other countries; either in distinct mines, or intermingled with the ores of lead or other metals.

The matter, which copper imbibes from this mineral in its conversion into brass, separates again from the brass on keeping it melted in an open vessel, and exhales in sumes; which condense, upon such adjacent bodies, as are less hot, into white slowers, the same with those into which

zinc

zinc is converted by fire. A mixture of calamine and powdered charcoal yields by itself, in open vessels, or if the air is admitted, the same flowers: in close vessels, the zinc is revived, and either runs off, or sublimes, in its proper metallic form, into that part of the vessel which is most remote from the action of the fire. The quantity of zinc is variable, as of other metals in their ores: Marggraf informs us, in the Berlin memoirs, that some of the foreign calamines yielded two sixteenths of their weight, an English calamine three sixteenths, and another English specimen from Holywell in Flintshire, seven sixteenths: from several parcels of the common calamine of the shops, I have gained nine sixteenths. The most exact way of determining the quantity of zinc appears to be, by mixture with a pretty large proportion of copper; by which the zinc, resolved into sume, is imbibed and detained.

This ore of zinc, employed principally for the making of brass, is for that purpose roasted or calcined; partly with a view to distipate some sulphureous matter, which the crude mineral is supposed to contain; but chiefly to render it friable, and more easily reducible into sine powder. It is with the ore thus calcined, that the shops are generally supplied. The roasted calamine is levigated into an impalpable powder.

Lap. calamin. præparatus Pharm. Lond.

In this state it proves, for external purposes, an excellent restringent, desiccative, and epulotic; of great use in collyria, against desluxions of thin acrid humours upon the eyes; and in unguents and cerates, for cutaneous ulcerations and excoriations. The officinal epulotic cerate, commonly called Turner's, is made by melting six ounces of yellow wax in a pint of oil olive, over a gentle fire, sprinkling in six ounces of levigated calamine assoon as the mixture begins to grow stiff, and keeping the whole stirring till grown quite cold. The college of Edinburgh uses only one part of the calamine to sive parts of a cerate composed of oil, wax, and spermaceti.

Cerat. epulotic. Ph. L.

Cerat. e lapide calaminari Ph. Edinb.

CALAMINTHA.

CALAMINT: a plant with fquare stalks: the leaves set in pairs; the slowers on branched pedicles, whereof two issue from one joint in the bosoms of the leaves: the upper lip of the flower is divided into two segments, the lower lip into three. It is perennial, and flowers in July and August.

I. CALA-

1. CALAMINTHA Pharm. Lond. Calamintha pulegii odore seu nepeta C. B. Melissa nepeta Linn. Calamint, field calamint: with reclining stalks; small, irregularly oval leaves, very slightly indented, without pedicles; and the flower-stalks longer than the leaves. It grows wild in dry grounds, and by the sides of fields.

This herb has a strong aromatic smell, approaching to that of pennyroyal; and a moderately pungent taste, somewhat like that of spearmint, but warmer. In virtue, it appears to be nearly similar to a mixture of those herbs: insusions of the leaves are drank as tea, in weaknesses of

the stomach, flatulent colics, and uterine obstructions.

Water extracts by infusion nearly all the virtues of the calamint; and carries off, in evaporation, the whole of its specific flavour. In distillation with water, there separates from the aqueous sluid a considerable quantity of essential oil, of a very pungent taste, and smelling strongly of the herb. The remaining decoction, thus divested of the aromatic part of the plant, is unpleasantly roughish, bitterish, and mucilaginous.

Rectified spirit extracts the virtues of the calamint more perfectly than water, and gains from it a deep green tincture. On gently distilling the filtered liquor, a part of the flavour of the herb rises with the spirit, and a part remains behind in the inspissated extract. Spirit manifestly brings over more from this plant than from spearmint, and less than from pennyroyal; its active matter being more volatile than that of the one, and less so than that of the other.

2. CALAMINTHA MONTANA Pharm. Edinb.(a) Calamintha vulgaris vel officinarum germaniæ C. B. Melissa Calamintha Linn. Common calamint, so called: with upright stalks; larger, short, serrated, pointed leaves, set on pedicles; and the slower-stalks of the length of the leaves. It is found wild about the sides of highways, but is less common, in this country, than the other.

The leaves of this species are in taste weaker than those of the preceding. Their smell is strong, not like that of pennyroyal, but rather approaching to that of the wild mints, though more agreeable. The essential oils of the two plants differ in slavour as the herbs themselves: in the spirituous extracts the difference is less considerable. They are

supposed to agree in virtue, and have been used indiscriminately; the shops being generally supplied with the species which is most easily procurable.

3. CALAMINTHA MAGNO FLORE Pharm. Parif. & C. B. Galamintha montana flore magno ex calyce longo J. B. Melissa grandistora Linn. Mountain calamint: with larger leaves and flowers than the two preceding, but smaller stalks; the leaves set on pedicles, pointed, acutely and deeply serrated like those of nettles; the flower-stalks shorter than the leaves, and of the length of the flowers themselves. It is a native of the southern parts of Europe, and raised with us in gardens.

This species has a moderately pungent taste, and a more agreeable aromatic smell than either of the other calamints. It appears to be

the most eligible of the three as a stomachic.

CALAMUS.

CALAMUS AROMATICUS Pharm. Lond. Acorus verus, Calamus aromaticus Pharm. Edinb. & C. B. Acorus Calamus Linn. CA-LAMUS OF SWEET-SCENTED FLAG: a plant with long, narrow, pointed leaves, like those of the narrow iris, of a bright green colour, divided by the longitudinal rib into two unequal portions, one of which is smooth, the other transversely wrinkled. The flowers are imperfect, and stand thick together, forming an elegant spike like the catkin of the hazel, which rifes in the bosom of one of the leaves about the middle of its height. The root, which spreads obliquely under the surface of the earth, is long, crooked, full of joints, about an inch thick, fomewhat flatted; externally of a greenish white colour, which changes, in drying, into a brownish yellow; internally white, and of a loose fungous texture. This plant grows plentifully, in rivulets and marshy places, about Norwich and in some other parts of this kindom, and, as is faid, in the canals of Holland: the flowers appear early in the fummer, the leaves die in the winter, the roots are perennial. The shops have been usually fupplied from the Levant with dried roots, not fuperiour, and fcarcely equal, to those of our own growth.

THE roots of calamus have a moderately strong aromatic smell, and a warm, pungent, bitterish taste. Their slavour, when fresh, is un-C c pleasant, pleafant, approaching in some degree to that of leeks or garlic: by drying, it is greatly improved, but does not become truly grateful. Some report them to be superiour in aromatic flavour to any other vegetables produced in these northern climates; but the specimens I examined fell

short, in this respect, of many of our common plants.

Infusions of calamus in water smell strongly of the root, and have a moderately warm and very bitter taste: spirit, applied after water, receives no fmell, and scarcely any taste. Tinctures of the root in rectified spirit are warmer and more pungent than the watery infusion, but much less bitter, and of very little smell: water, applied after spirit, gains a confiderable bitterness, but no smell. It appears, therefore, that water is the most perfect menstruum of the bitter matter, as rectified -spirit is of the aromatic, and that the smell of the calamus is covered or suppressed by spirit. The tinctures in both menstrua are of a yellow or brown colour, according as they are less or more saturated.

In distillation with water, there arises a small quantity of essential oil, amounting only to about two ounces from fixty-fix pounds of the root (a): both the oil and distilled water have a strong smell, somewhat less grateful than that of the root in substance: the remaining decoction, thus deprived of the aromatic matter, is nauseously bitter. On distilling the spirituous tincture, the distilled spirit has scarcely any smell or taste of the calamus: the extract, nevertheless, has very little smell, and much less taste than might have been expected in the extract of so warm a

root.

Calamus aromaticus was formerly held in confiderable esteem as a warm stomachic; and was commonly made an ingredient in bitter tinctures and infusions: among us, it has given place, in this intention, to bitters of a more grateful kind. The root, candied, is faid to be used at Constantinople as a preservative against contagion.

CALENDULA.

CALENDULA Pb. Edinb.(b) Caltha vulgaris C. B. Calendula SINGLE MARIGOLD: a plant, with oblong undivided officinalis Linn. leaves, joined close to the stalk, widening from thence to the extremity, juicy, and somewhat clammy to the touch; and moderately large, bright yellow or gold coloured flowers, composed of a number of indented petals standing round a middle disk, on which, after the flower has fallen, several rough crooked seeds lie naked. It is annual, common in gardens, propagates itself by seeds, and flowers from May to the end of autumn.

Marigold flowers have been recommended as aperients in uterine obstructions and icteric disorders; as sudorifies, alexipharmaes, and for promoting eruption in malignant and exanthematous severs. They appear, from their sensible qualities, to be of little activity: when fresh, they have a faint unpleasant smell, which is lost in drying: their taste is chiefly mucilaginous, with a slight bitterishness. They give a pale yellow tincture to water, and a deeper yellow to spirit: the watery insusion has the most smell, and the spirituous the most taste of the slowers. The extract obtained by inspissating the spirituous tincture, is bitterish and slightly roughish: the watery extract is a tenacious mucilage, of less taste than the other.

The leaves of the plant appear to be of greater virtue than the flowers. Chewed, they impress at first a viscid sweetness, which is followed by a penetrating pungency, very durable in the mouth, not of the hot or aromatic, but rather of the subsaline kind. Their expressed juice, which contains great part of the pungent matter of the herb, has been given, in doses of two or three ounces or more, as an aperient; and is said to loosen the belly, and promote the natural secretions in general.

CALX VIVA.

CALX VIVA Pharm. Lond. & Edinb. QUICKLIME: an acrimonious, friable fubstance; dissolving, very readily, in the nitrous, marine, and vegetable acids; uniting with the vitriolic acid into an indissoluble and nearly insipid concrete; producing heat on the affusion of water, partly dissolving in the water, and impregnating it with a strong taste.

Quicklime is prepared, about London, chiefly from chalk; in other parts of the kingdom, from different forts of stones called, from their use, limestones; by calcining them, in kilns made for this purpose, with a strong fire. All the native mineral earths and stones, that dissolve in acids, and all the sea shells that have been tried, are reduced by fire

Cc2

into quicklime; and fuffer, in the calcination, a great diminution of their weight. If the lime be exposed long to the atmosphere, it falls by degrees into powder, increases in weight, loses of its acrimony, and at last becomes similar in quality to what the earth was before calcination: it retains its acrimony much longer in a moist than a dry state.

THE earths and stones, from which quicklime is produced, contain a large quantity of air, which in calcination is expelled: hence strong quicklime raises no effervescence, or emits no air-bubbles (which the crude earths do in great abundance) during its diffolution in acids. A theory, which now begins to prevail, confiders the proper calcareous matter as a substance, which is in its pure state quicklime; which, by the simple coalition of air *(a) with it, loses its acrimony, solubility in water, and other distinguishing characters; and which, on the bare separation of this incorporated air, proves quicklime again (b). Thus much appears demonstrated, that either the qualities of the calcareous matter are affected by the air itself imbibed or expelled, or that both (in all the experiments hitherto known) are equally affected by some other cause: in either case, the discovery is valuable in regard to practical utility; the calcareous earths and stones becoming quicklime by all those means by which air is expelled from them, whether calcination by fire, or folution in acids and precipitation with substances void of air; and quicklime lofing its qualities by all those means by which air is introduced, whether direct exposure to the atmosphere, or commixture with certain other bodies from which it inftantaneously absorbs the aereal matter.

Quicklime is employed for increasing the activity of alkaline salts: if water, strongly impregnated with the lime, be gradually added to a solution of fixt alkali, the calcareous matter separates and subsides, saltiated with air, and no longer acrid or dissoluble in water; the alkali at the same time losing its air, so as to make no effervescence with acids, and proving in this state much more acrid than at first. Quicklime is sometimes used also in external applications as a depilatory; and has been sometimes made into an unguent with honey for rheumatic and other

obstinate

^{*(}a) Not atmospherical air, but the species of air termed fixed or fixable.

⁽b) See Dr. Black's Experiments on Magnesia, in the Edinburgh essays physical and literary, vol. ii. art. 8. and Dr. Macbride's Experimental essays.

obstinate fixt pains of the joints or limbs: this unquent is greatly commended by Fuller, who observes that it is almost caustic.

Solutions of the calcareous matter in water are given internally with fafety, and in many cases with advantage. For this purpose, a gallon and a half of water is poured by degrees upon a pound of fresh burnt Aqua calcis quicklime, the veffel shaken when the ebullition ceases, and then set by Pb. Lond. till the undiffolved lime has fettled; after which, the liquor is poured off, and passed through a filter. In the last Edinburgh dispensatory, Aqua calcis half a pound of lime is directed to be sprinkled with four ounces of water Ph. Edinb. in a close vessel, and when it is fallen to powder, twelve pounds more of water are to be added, and the whole agitated about ten times, keeping the veffel still close; and the liquor then filtered. Only a small portion of the lime is diffolved by the water, and the remainder gives a strong impregnation to large quantities of fresh water, though not so strong as to the first; great part remaining at last undissolved: this refiduum, calcined again, becomes quicklime as before: and by repetitions of this process nearly the whole may be dissolved.

The folution has a strong, styptic kind of taste. It changes the juices of blue flowers to a green colour; precipitates metallic bodies diffolved in acids; tinges filver of a coppery hue; and turns red wine to a dark colour: by these properties, the strength of its impregnation with the calcareous matter may be in some measure estimated. The specific gravity of the liquor is increased by the lime, in a much greater ratio than the fmall quantity taken up can effect by the apposition of its own weight (a); on account, perhaps, of the water being deprived of its air. In veffels quite filled with the lime-water, and exactly closed, it may be kept unchanged for many months: in open vessels, the calcareous matter soon separates from the aqueous fluid, and concretes upon the surface into a crust, insipid and indissoluble as the earth in its natural state, and again convertible into quicklime by a repetition of the calcination. As most kinds of liquids, and many other bodies, are impregnated, more or less, with the fubstance which lime greedily imbibes, and which renders it indiffoluble; lime-water suffers a separation of part of its lime in most mixtures, and probably also in the act of its dissolving bodies: hence, when this liquor is employed as a menstruum, it is advisable to add some

⁽a) Whytt, Edinb. eff. & obf. pbyf. & lit. vol. i. art. xiii. p. 383.

quicklime in substance, in order to continue the impregnation of the water with the lime.

Lime-water diffolves, by the affiftance of heat, mineral fulphur, vegetable oils and refins, and animal fats: it extracts, in the cold, the virtues of fundry refinous and oily vegetables, and diffolves thick phlegm or mucous matters, and the curd of milk; with which last it forms a white liquid, nearly similar in its appearance to milk in its natural state. It has lately been found to dissolve also the human calculus, particularly the lime-water prepared from calcined oyster-shells, which proves a more active menstruum for this concrete (and possibly for other substances also) than that made from the stone limes; the dissolving power of the oyster-shell lime-water seeming, from Dr. Whytt's experiments, to be more than double to that of the stone lime-waters (a). Taken internally, in considerable quantity, it impregnates the urine in some degree with its lithontriptic power, and in sundry calculous cases has happily given relief.

Lime-water, drank to the quantity of a quarter of a pint three or four times a day, has been found ferviceable in scrophulous complaints, fluxes, seminal weaknesses, and other disorders proceeding from an impurity of the fluids, or laxity and debility of the solids. It generally promotes urine; oftentimes the cuticular discharge; and where the stomach is oppressed with viscid phlegm, expectoration. It for the most part binds the belly, and sometimes occasions a troublesome costiveness, unless this effect be occasionally provided against by the interposition of proper laxatives. It answers best in cold, sluggish, phlegmatic, and corpulent habits; and is to be used more cautiously in hot bilious dispositions, and where the patient is greatly emaciated, or the appetite weak, and at the time of any critical or periodical evacuation.

It is customary to impregnate lime-water with different materials, partly for rendering it more acceptable to the palate and stomach, and partly for improving its medicinal efficacy against cutaneous defedations. The college of London directs, in this view, half an ounce of sassafras bark and one ounce of liquorice+, with the addition, in some cases, of four ounces of rasped guaiacum wood and three drams of coriander

Aqua calcis † minus composita Ph. Lond.

⁽a) Edinb. medical effays, vol. v. art. 69. See on this subject his Treatise on the virtues of lime-water.

feeds 1; to be macerated for four days in three quarts of lime-water, 1 magis and the liquor strained off for use. These infusions are taken in the composita Ph. Lond. fame quantities as the fimple lime-water, by themselves, or with the addition of milk.

CAMPHORA.

CAMPHORA Pharm. Lond. & Edinb. Caphura. CAMPHOR: a folid concrete, fomewhat unctuous to the touch: totally volatile in the heat of boiling water, and fubliming unaltered: melting in a less degree of heat into the appearance of oil; readily taking fire on a red-hot iron, and burning entirely away, with a bright white flame, and copious fumes, which condensing form foot; soluble in spirit of wine, and in oils, and in the nitrous and vitriolic acids, not in water, nor in vegetable acids. From the nitrous acid, diluted with a little water, it absorbs the stronger acid matter, and forms therewith a substance like oil, which floats on the furface of the more phlegmatic liquor: with the vitriolic, it mingles uniformly into a yellowish red fluid.

Camphor is extracted, by a process similar to that by which essential oils are obtained, from the wood and roots of a large tree of the bay kind, growing in Japan, called by Linnæus Laurus (campbora) foliis trinerviis lanceolato-ovatis: nervis supra basin unitis. A species of camphor is fometimes likewise found naturally concreted into little grains, in the medullary part of this and fome other trees: I have in my possession a piece of a reddish wood, which seems to be part of the trunk of a large tree, and which on being split in different places exhibits camphor

plentifully concreted in it.

* In the Adversaria of the learned Gaubius, is an account transmitted by a correspondent, of the production of camphor in the island of Sumatra, which is worthy of notice. There are, it feems, only two kinds of camphor known in commerce in the East Indies, the Sumatran and Japanese. The former is so much superior to the latter, that the Japanese themselves readily give a hundred pounds of their own, for one pound of the Sumatran. The reason of this probably is, both that the climate of Sumatra is much warmer, and that the camphor there is entirely prepared by nature. In the northern parts of this island the camphor tree grows to fuch a fize, that planks two feet in breadth are fawn out of its trunk. After it has stood for a certain number of years, its branches naturally crack, and an oily liquor exudes from the fissures. When the inhabitants observe this, they collect the oil in pieces of bamboo, and watch the time when they have learned by experience that the formation of the camphor is complete. They then, after some superstitious ceremonies, cut down the tree, and splitting the branches, which are found full of camphor, pick out the pieces, making separate parcels of the large and the small ones. They conclude with rasping the wood itself; and thus make three forts of camphor, which they bring to the Dutch sactory; selling, however, all three together by weight. The largest and finest pieces are called, in the Malay language, Copal, or head; the smaller, Poeroet, or belly; and the raspings of the wood, Cacki, or feet*(a).

As first sublimed or distilled from the wood, it appears brownish, and composed of semipellucid grains mixed with some impure matter. In this state, it is imported by the Dutch, and purified by a second sublimation, by which it becomes clear and white: this last process is so managed, that the head of the subliming glass is kept warm enough, to make the camphor run together into a mass of its own figure; in which form it is brought into the shops. It may likewise be purified by solution in spirit of wine; recovered from the spirit by distillation, the spirit rising before the camphor; and afterwards formed into loaves by suspense that a gentle heat, in a close vessel.

This concrete has a fragrant smell, somewhat approaching to that of rosemary, but much stronger; and a bitterish, aromatic, pungent taste, accompanied with an impression of coolness. It is looked upon as one of the principal diaphoretics and antiseptics, and as possessing some degree of an anodyne or antispassmodic power. It is apparently of great subtility and penetration, quickly diffusing itself through the habit in a very sensible manner: taken in any considerable quantity, it generally produces very uneasy sensations about the stomach and præcordia, and often in the remoter parts; though it does not heat the body near so much, as might be expected from the great pungency of its taste.

Hoffman

^{*(}a) The editor has been favoured with an account of the camphor tree, fent from Sumatra, which agrees perfectly with the above. The writer further fays, that the Sumatran camphor tree is certainly a new genus, and not at all refembling the tribe of Lauri.

Hoffman reports, that doses even of half a dram did not increase the pulse, or excite any immoderate heat, but occasioned rather a sense of coolness; and that on continuing the use of the camphor for some time, the blood became more sluid, and the quantity of watery serum, which the habit before abounded with, was notably diminished.

* A remarkable account of the effects of camphor in a large dose on the relator himself, Mr. Alexander, surgeon, in Edinburgh, is contained in the Philosophical Transactions, vol. lvii. part 1. After taking one scruple of this medicine, he found his pulse somewhat abated in frequency, but no other change in himself is remarked. He next took two scruples; the first effect of which was to fink the pulse from 77 to 70. In less than half an hour it returned to its former number; and at this time a giddiness came on, which gradually increased, till all consciousness of present, and memory of past, objects was obliterated. Violent efforts to vomit, with strong convulsions and a temporary mania, fucceeded. The pulse was now raised to 100. Some degree of recollection returned; accompanied with a fensation of violent heat, and tremors of the whole body. The exhibition of warm water now caused a rejection of great part of the camphor; and from this time its effects by degrees wore off. A great foreness and rigidity of the whole body were felt the next day and the day following. A fimilar account is related in a thefis de viribus camphoræ by Dr. Griffin, printed at Edinburgh in 1765.

In acute diseases, this medicine is given from a quarter of a grain to one or two grains, and sometimes more, in conjunction commonly with nitre, or other substances of the anti-inflammatory saline kind. Hossman observes (a), that it answers best on the approach of a criss, or in the decline: that it is to be used with caution during the increase, and when the fever is at the height, more especially where the internal heat is great, moisture desicient, and the skin dry: and that it is sparingly to be given also when nature is weak; where a tumidness and redness of the face, with vertiginous complaints, torpor, and sleepiness, shew the vessels of the head to be distended; as also in palsies, convulsions, and in plethoric and costive habits.

⁽a) Diff. de ufu campboræ securissimo & præstantissimo; & Med. rational. de febrib. passim.

In chronical diforders proceeding from a redundance of serous defluxions, or from an impurity of the humours, and as an affistant to mercurial alteratives, it is used more freely, and with less danger: in some cases a little opium is joined, which prevents the uneasiness which camphor of itself is apt to produce, and at the same time increases its operation by sweat, a mixture of camphor and opium being one of the most potent sudorifics. Some recommend camphor to be given in maniacal cases, to the quantity of half a dram every night or oftener; and instances have been produced (a) of this practice being attended with success.

It has been generally supposed that this concrete corrects the irritating power of cantharides; and Hossiman looks upon it as a corrector also of the stimulating cathartics and emetics. It apparently corrects, in a considerable degree, the more active mercurial preparations; that is, it determines their operation to the cuticular emunctories, and by promoting their diaphoretic, restrains their purgative or emetic virtue: but how far it varies the action of cantharides, and the stimulants, purgatives, and emetics of the vegetable kingdom, is not as yet certainly known.

Camphor may be dissolved in watery liquors, and thus fitted for being commodiously taken, by grinding it with sugar, almonds, or thick mucilages, and adding the water by degrees. A dram of camphor, rubbed with a few drops of rectified spirit of wine till it grows soft, requires about four drams of fine fugar: a pint of boiling water is poured on this mixture, the veffel closely covered, and the liquor, when grown cold, strained out for use. Vinegar also, by this treatment, dissolves the camphor equally with water, and is often preferred in acute diseases, whether putrid or inflammatory, as rendering the julep somewhat more grateful both to the palate and ftomach, and excellently coinciding with the medicinal intention. The whole of the camphor, however, is not disfolved by either; a part, and generally a confiderable one, remaining behind upon the strainer. Almonds or mucilages render it completely dissoluble into an emulfion or milky form. The above quantity of camphor requires about twelve almonds; to which mixture a pint of some suitable aqueous fluid, as the diffilled water of pennyroyal, is commonly added, and half an ounce of fine fugar diffolved in the strained liquor. In this

Julepum e camphora Ph. Lond.

(a) Philosophical Transactions, n. 400. Suenska vetenskaps acad. bandl. tom. v. ann. 1744.

form, vinegar or other acids can have no place, as they coagulate the emulfion, or at least render it incapable of keeping the camphor dissolved : but nitre may be added in any quantity that may be thought proper, this neutral falt mingling uniformly with the liquor, and producing no feparation of its parts. Emulfions made with mucilages admit both nitre and acids.

A folution of camphor in rectified spirit of wine, in the proportion of an ounce to a pint, is employed externally against rheumatic pains and paralytic numbneffes, for discussing tumours and inflammations, and restraining the progress of gangrenes. On diluting this solution with watery liquors, the mixture becomes milky, and on standing for some time greatest part of the camphor separates. It has been said, that with the spirit of fal ammoniac made by quicklime, and with saturated alkaline lixivia, it mingles without separation: but, on trial, it turned milky with the former, in the same manner as with water; and with the latter it did not mingle at all, the camphorated spirit swimming distinct upon the furface of the alkaline lye. It has been reported also, that a camphorated spirit, uniformly miscible with water, may be obtained, by grinding the camphor with fomewhat more than equal its weight of fixt alkaline falt, then adding a proper quantity of proof spirit, and drawing off one half by distillation. This spirit, however, does not answer expectation: the quantity of camphor that rifes with it is exceeding small, greatest part remaining behind in the distilling vessel: hence, though when the spirit is mixed with a large quantity of water, it occasions no fensible turbidness, yet when mixed with only a little water, it exhibits the same appearances as the common solution, differing no otherwise than in degree.

Camphor is used also in unguents, for burns, itchings, and serpiginous eruptions on the skin: for these purposes, a dram and a half of camphor, first ground with a few drops of oil, are mixed with a pound and a half Ung. album of the white ointment. It is mixed in a larger proportion, with cataplasms for the throat against inflammations of the uvula and tonsils; and disfolved, for rheumatic and other pains, in oil of olives, in the propor- Ol. camphotion of one part of camphor, to four of the oil. Hoffman reports, that a folution of camphor, in empyreumatic vegetable oils that have been rectified by distillation from quicklime, procures immediate relief in some kinds of violent pains (a).

Spiritus vini ratus Ph.

camphorat.

rat. Ph. Edin.

(a) In notis ad Poterium, p. 483.

CANCRORUM CHELÆ.

CHELÆ CANCRORUM Pharm. Lond. CRABS CLAWS: the black tips of the claws of the cancer marinus or common sea crab.

Chelæ canc. præparatæ Pb. Lond.

lis cancr. comp. Pb. Lond.

This testaceous matter, levigated into an impalpable powder, is made use of for absorbing acidities in the first passages, and makes the basis of the compound absorbent powders of the shops: to four parts of the Pulv. e che- prepared claws, the college of London joins one part of prepared red coral and one of pearl. All these three ingredients consist of the same calcareous animal earth: how far any of them is superiour, as a medicine, to the others, or any composition of them to any one of the three separately, does not appear. The distinguishing characters of this kind of earth are, its being convertible, by calcination with a strong fire, into quicklime; its not perfectly vitrefying with a moderate proportion of vitreous fluxes, but rendering the flux or glass opake and white; its being diffoluble in all acids except the vitriolic, and precipitable by this laft from the others.

CANCRORUM OCULI.

CANCRORUM OCULI dicti Pharm. Lond. & Edinb. lapides. CRABS-EYES fo called: stony concretions, found in the head, or rather stomach, of the astacus suviatilis or river craw fish; generally about the fize of peas, or larger; of a roundish shape, flatted on one fide; in colour white, fometimes with a reddish, and sometimes with a blueish cast; internally of a leafy texture. They are said to be brought to us chiefly from Holland: perhaps the greatest quantities are the produce of Muscovy, particularly of the river Don, where the craw fish, as I have been informed, are extremely plentiful, and have been commonly laid in heaps to putrefy, after which the stones are picked out.

These stones are said to be sometimes counterfeited with tobacco-pipe clay, or chalk mixed with glutinous materials. Compositions of this kind may be readily diftinguished from the genuine crabs-eye, by their texture being uniform and not leafy, and by their flicking to the tongue, and being foftened with water. They differ also in their habitude to

acids; either not dissolving at all, or dissolving in another manner.

If genuine crabs-eyes be put entire into strong vinegar, or into aqua fortis largely diluted with water; their earthy part is gradually extracted, and there remains a soft transparent gelatinous substance, of the same figure with the original concrete: such as were at first coloured, retain their colour after the action of the acid. The quantity of this gelatinous matter is much less than might be judged from the volume which it occupies; amounting, when the nitrous acid has been used, scarcely to one tenth part of the weight of the crabs-eyes: vinegar leaves a larger quantity, a part of the earth itself seeming to escape the action of this acid.

Crabs-eyes are used as an absorbent of acid humours, and are supposed, when combined with the acid, to be more aperient and resolvent than most of the other absorbent earths: in this intention they are commended by Hoffman, who looks upon a folution of them in vinegar as capable of refolving both stagnant thick humours and coagulated blood. * They have lately been employed, by an eminent physician at Leyden, with great fuccess, in the cure of the fluor albus. He gives to the quantity of half an ounce in a day; and remarks, that it is particularly ferviceable in that species of the disease in which the discharge is so acrimonious as to corrode the parts (a). Their earth differs remarkably from that of the preceding article, in not being convertible into quicklime; but the medical differences of their folutions in vinegar, or in other acids of the vegetable or animal kingdom, do not appear to be very great, the folutions of the two earths being in taste nearly alike. The earth of crabs-eyes, in regard to its chemical characters, is of the fame nature with that of hartshorn.

CANELLA.

CANELLA ALBA Pharm. Lond. & Edinb. Canella alba & Costus corticosus Ph. Paris. Cinamomum sive canella tubis minoribus alba C. B. Winterania Canella Linn. Canella Alba: the inner bark of a large bay-leaved tree, growing in the low lands of Jamaica and other American islands: brought over in the form of quills; of which some are large and thick, taken from the trunk of the tree; others slender and thinner,

from the branches; having generally pieces of a wrinkled brownish coat adhering to the outside; lined on the inside with a fine white membrane; breaking over with a close even surface, and appearing internally of an unequal, pale, brownish or yellowish white colour.

Canella alba has hitherto been rarely employed in medicine, unless as a substitute for winter's bark, which it pretty much resembles, and has been commonly mistaken for. The London college has now received it in two officinal compositions, for alleviating the ill flavour of aloes; and the Edinburgh, in their Tinctura Amara. It is a moderately warm aromatic; of an agreeable smell, somewhat resembling that of cloves, but far weaker; and of a pungent taste, accompanied with a considerable bitterishness.

Infusions of it in water are of a yellowish colour, and smell moderately of the canella, but in taste are rather bitter than aromatic. Tinctures made in rectified spirit are of a darker reddish yellow colour, and have more of the aromatic warmth of the bark, but very little of its smell. Tinctures in proof spirit are more agreeable than either; this menstruum dissolving the aromatic as well as the bitter matter of the canella, without

covering or suppressing its flavour like the pure spirit.

In distillation with water, it yields an essential oil, of a dark yellowish colour, of a thick tenacious consistence, difficultly separable from the aqueous sluid, in smell sufficiently grateful, though rather less so than the bark itself: the remaining decoction, inspissated, leaves an extract of great bitterness, in consistence not uniform, seemingly composed of a resinous and gummy matter imperfectly mixed. On inspissating the spirituous tincture, the spirit, which distils, has no great smell or taste of the canella, but is so far impregnated with its more volatile oil as to turn milky on the admixture of water: the remaining extract retains the bitterness of the bark, but has little more of its warmth or flavour than the extract made with water.

CANNABIS.

CANNABIS Pharm. Edinb. (a) & C. B. Cannabis fativa Linn. HEMP: a tall annual herb, with digitated leaves, cultivated in fields

on account of the mechanic uses of its tough rind. Some of the plants, called male, produce flowers; composed of yellowish stamina set in siveleaved cups. Others, called female, produce seeds; moderately large, covered with a shining dark grey-coloured shell; under which is lodged a white kernel.

This plant has a rank smell, of the narcotic kind, and is supposed to be prejudicial to health. It is said that the effluvia of the fresh herb weaken the eyes, and affect the head(a): and that the water, in which the herb has been steeped for facilitating the separation of the tough rind, is a violent and sudden poison(b). The deleterious power of this liquor may depend, however, not solely on the specific virtues of the hemp, but in great part on the strong putrid taint which the soluble matter of the herb contracts during the process: for flax, a plant not suspected of any hurtful qualities, is reckoned to give a like poisonous impregnation to the water in which it is long macerated; insomuch that the steeping of one, as well as of the other, in spring or running waters, or ponds in which cattle drink, is prohibited by law(b). The leaves of an oriental hemp, called bangue or bang, and by the Egyptians as a faid to be used, in the eastern countries, as a narcotic, and aphrodisac(c).

The feeds of hemp, when fresh, have a faint smell of the herb, which is dissipated in keeping: their taste is unctuous and somewhat sweetish, accompanied with a slight warmth. They yield upon expression a considerable quantity of insipid oil; and unite with water, by trituration, into an emulsion. Decoctions of them in milk, and the emulsions, have been recommended against coughs, heat of urine, $\mathcal{C}c$. in which cases they may be of service, as emollients and obtunders of acrimony: but the virtues attributed to them against incontinence of urine, and for restraining venereal appetites, appear to have less, if any foundation. They are said to be used in some places as food; and, when taken freely, to affect the head (d).

⁽a) Lindestolpe, De venenis, edit. Stentzel. cap. x. thef. xiii. p. 541.

⁽b) Ray, Hift. plant. i. 159.

⁽c) Kæmpfer, Amanitates exot. p. 645. Alpinus, De med. Ægypt. lib. iv. cap. 2.

⁽d) Ray, Hift. plant. p. 158.

CANTHARIDES.

CANTHARIDES Pharm. Lond. & Edinb. CANTHARIDES or Spanish flies: an infect of the beetle kind, (Meloe vesicatorius Linn.) generally about half an inch in length; on the upper side, of a shining green colour, variegated with more or less of a blue and a gold yellow; on the lower side, brownish. These infects are frequent in Spain, Italy, and the southern parts of France: they are collected from herbs and bushes, killed by the steam of strong vinegar, and afterwards dried in the sun. The largest and best are said to come from Italy. They should be chosen fresh coloured, entire, and free from dust: on long keeping, they are apt to lose of their colour, and become powdery, and in this state are to be rejected.

CANTHARIDES have little or no fmell, unless the quantity is large; in which case they yield a faint disagreeable one, Cautiously tasted, they impress a slight sense of acrimony: those who describe the taste as highly acrimonious and caustic, have probably judged, not from the direct sensation of taste, but from the consequential effects. Applied to the skin, they first inslame, and afterwards excoriate the part; raising a more perfect blister, and producing a more plentiful discharge of serum, than any of the vegetable acrids. Hence their common use as a vesicatory.

Veficatories are employed, either as a general stimulus, for raising the pulse and quickening the circulation, in low fevers, and in lethargic disorders; or for resolving topical obstructions. Fixt pains, whether external or internal, as in the rheumatism, sciatica, dysentery, pleurisses and peripneumonies, are frequently observed to yield to a blister upon the part; though frequently also the matter is lodged so deep, as to be beyond the reach of this as well as of other external medicines. Blisters are likewise applied to the head in epileptic and maniacal disorders, inveterate and periodic headachs, and obstinate defluxions on the eyes, in which cases they are not to be considered merely as topical remedies: Hossman relates, that in defluxions on the eyes, he has known a blister, applied to the nape of the neck, increase the pain; whilst one laid on the soles of the feet has procured relief as soon as the discharge from its operation began to take place. Blisters on the head give the least pain; on the legs the most.

The

The bliftering applications are generally composed of cantharides reduced into fine powder and mixed with plasters or other compositions of a due confistence. Three ounces of the powder are stirred into three ounces Emplast. each of hogs lard, yellow wax, and white refin, melted together +; or into fix ounces of the drawing plaster with the addition of a quarter of a pint of vinegar ‡. This last ingredient is supposed to promote or facilitate † Pb. Lond. the action of the cantharides: for in some cases, where the plaster without vinegar has failed of taking effect, on removing it and washing the part with vinegar, the same plaster, applied again, has blistered freely: it is probable, however, that this was owing, not fo much to any peculiar quality of the vinegar, as to its foftening and deterging the fkin; an effect which is not to be expected from it when mixed with the other ingredients of the plaster. Other stimulating ingredients are sometimes added, as pepper, mustard-feed, and verdigris; but it does not appear that these kinds of fubstances give any material affistance to the action of cantharides. The powdered flies spread on the surface of a common plaster operate as effectually as any of the compositions, and in this form they are often used.

In some cases, as in variolous eruptions or other inequalities of the skin, compositions of a softer consistence than plasters are required, that they may apply themselves to the depressed parts: for these purposes, equal quantities of finely powdered cantharides and wheat flour are mixed with Epithema vinegar into a paste. Where blisters are intended to be made perpetual, Ph. Lond. or continued, as a constant drain of serous humours, for a considerable time, fome cantharides are added occasionally in the dressings, to keep the ulcers open: ointments for this intention are prepared in the shops, by melting together equal parts of the bliftering-plafter and hogs lard, and keeping them stirring together till grown cold; or by melting seven parts of yellow bafilicon, and then adding one part of powdered cantharides.

Cantharides are applied also, in smaller quantity, sufficient to warm and stimulate the part considerably, but not to raise a blister, against some rheumatic pains, chilblains, and paralytic affections. In this intention, the bliftering compositions are diluted with other plasters, in such proportion, that the quantity of the fly may be about one twenty-fixth part Emp. caliof the whole compound.

The external use of cantharides, if the quantity be considerable, is often followed by a strangury and heat of urine; this infect being peculiarly

Ung. ad veficatoria Ph. Lond. Ung. epifpaf. e pulvereCantharidum Pb.

dum No focom. Edinb.

culiarly disposed to affect the urinary organs, though applied to the remotest parts. This inconvenience is prevented or remedied, by emulsions or mucilaginous liquors plentifully drank.

SMALL doses of cantharides are given internally in suppressions of urine, and for deterging ulcerations of the bladder. They have likewise been found remarkably serviceable in seminal weaknesses and old gleets; in which the balsamic medicines, generally recommended, are often ineffectual (a). In leprous cases also they have frequently had excellent effects, in virtue perhaps of their diuretic power; for so great is the consent of the kidneys with the skin, that the humours accumulated in the cutaneous glands may be discharged by urine; as the urinary liquor, when the kidneys fail in their office, sometimes transpires through the skin(b).

Great caution is requifite in the use of this highly stimulating medicine; a small excess in the dose producing not only a strangury, but a discharge of blood, with intense pains about the neck of the bladder: a grain, and even a quarter of a grain (c), has in some cases had this effect. The remedy for these symptoms, in good habits, and where the cantharides have not been greatly overdosed, consists in plentiful dilution with emollient liquors in which some nitre has been dissolved, with the interposition of moderate doses of opium. It is commonly supposed that camphor, given along with the fly, corrects in some degree its irritating power.

CANTHARIDES, digested in rectified spirit, impart to it a bright yellow tincture, and have their own colour improved: boiling water receives from them a muddy yellowish or brownish hue, and considerably impairs the colour of the sly. The active matter of the cantharides is completely taken up by both menstrua, and does not rise with either in distillation or evaporation: the substance of the sly remaining after digestion either in water or in spirit, does not in the least blister or inslame the skin; whereas both the watery and spirituous extracts blister freely.

(a) Mead, Monita & præcepta, p. 256. (b) Idem, Medica sacra, p. 24.

The fafest and most commodious form for taking cantharides internally, is the spirituous tincture; which, dropt into watery or vinous liquors, mingles uniformly, without precipitation or turbidness. Two drams of Tina. the cantharides, bruised a little, are commonly digested two or four days canth. in a pint and a half or two pints of proof spirit, with or without the Ph. Edinb. addition of half a dram or more of cochineal as a colouring ingredient. These tinctures are usually given from fifteen to thirty or more drops twice a day: the most certain method of obtaining, without danger, the full effect of the cantharides, is, to begin with the smaller dose, and increase it by two or three drops at a time, till a little uneasiness is perceived in making water; after which, the medicine being intermitted for a day or two, the dose is to be diminished a little, and continued just below the quantity which produced that effect.

A foft extract of cantharides is in many cases preferable, for external purposes, to the ointments and plasters made with the powdered fly, particularly for the dreffing of perpetual blifters; as it acts more uniformly than the compositions containing the fly in substance, and occasions less pain in the dressing. Hostman's mild blister which gives little pain, mentioned now and then in his works, feems to have been, or to have had for its basis, a preparation of this kind; and probably the empyrical perpetual blifter is no other. The college of Edinburgh have Unguentum now received a composition on the same principle: they direct an ounce of cantharides to be infused for a night in four ounces of boiling water, cantharidum the liquor to be strongly pressed and strained out, and boiled with two ounces of hogs lard till the humidity is wasted; after which, an ounce of white refin, an ounce of yellow wax, and two ounces of Venice turpentine, are to be added, and the whole well mixed fo as to form a fmooth ointment.

epispasticum ex infuso Ph. Edinb.

CARANNA.

CARANNA Pharm. Edinb.(a) CARANNA: a concrete refinous juice; exuding from a large tree, of which we have no particular account; brought from New Spain, and some other parts of America, in little masses, rolled up in leaves of flags; externally of a dark brownish colour,

(a) Expunged.

Ee2

internally

internally brown with a cast of red, variegated with irregular white streaks; somewhat soft and tenacious as it first comes over, but in length of time growing dry and friable.

This juice has an agreeable smell, especially when heated, and a bitterish and slightly pungent taste. Water dissolves about one fourth of it, and rectified spirit above three fourths: what is left by the one menstruum dissolves in the other, a small quantity of impurities excepted: both

folutions are of a bright yellow colour, the spirituous deepest.

The watery tincture smells agreeably of the caranna, and is in taste bitterish and somewhat warm. In distillation with water, there separates from the aqueous sluid a considerable quantity of an orange-coloured essential oil, of a very fragrant smell, and a moderately pungent taste: the remaining decoction, inspissated, leaves an extract of an ungrateful,

though weak, bitterishness.

The fpirituous tincture is both in smell and taste stronger and more agreeable than the watery. Inspissated, it yields a very tenacious adhesive resin, with an oily matter which separates and floats on the surface: the resin has very little smell, and scarcely makes any impression on the organs of taste: the oil is considerably aromatic, and moderately bitter, in which last respect it differs from the purer oil obtained by distillation with water.

Caranna has been chiefly employed as an ingredient in vulnerary balfams, corroborant and discutient plasters, and other external applications. It has very seldom been given internally, and is now, in this country, almost wholly in disuse.

*CARDAMINE.

CARDAMINE Pharm. Edinb. Nasturtium pratense magno store C. B. Cardamine pratensis Linn. Ladies-smock of Cuckow flower: a plant, of the class tetradynamia of Linnæus, in taste resembling cress. It has an erect stalk; and leaves set in pairs on a middle rib, with an odd one at the end. Its slower is white or purplish, and is succeeded by a bivalvular pod. It grows plentifully in moist meadows, and slowers early in the spring.

The

The virtue of the flowers of ladies-smock, in hysteric and epileptic cases, was first noticed by Ray; and their use has been revived by Sir George Baker, who has published some cases of their efficacy in the Medical Transactions, vol. 1. The flowers are given in powder, in doses of from thirty to ninety grains. They have little sensible effect, sit easy on the stomach, and increase the appetite. Their antispasmodic powers seem, from the cases related, to be considerable.

CARDAMOMUM.

CARDAMOM: a dried fruit or pod, brought from the East Indies; divided internally into three cells, in each of which are contained two rows of triangular seeds, of a brownish colour on the outside and white within.

1. CARDAMOMUM MINUS Pharm. Lond. & Edinb. Cardamomum fimpliciter in officinis dictum C. B. Cardamom, lesser cardamom: with short triangular husks, scarce half an inch in length; the produce of a plant with reed-like stalks, described in the Hortus malabaricus under the name of Elettari; the Amomum Cardamomum of Linnæus.

These seeds, freed from the husks, are an elegant and useful aromatic, of a grateful smell and flavour, very warm, yet not fiery, or subject, like the spices of the pepper kind, to produce immoderate heat. The husks should be separated only at the time of use; for the seeds soon lose a part of their flavour in being kept without this defence.

Their virtue is extracted, not only by rectified spirit, but almost completely by water also; with this difference, that the watery insusion is cloudy or turbid, the spirituous clear and transparent: the colour of both is a pale yellow. Scarcely any of the aromatic seeds give out so much of their warmth to watery menstrua, or abound so much with gummy matter, which appears to be the principle by which the aromatic part is made dissoluble in water: the insusion is so mucilaginous, even in a dilute state, as hardly to pass through a filter.

In distillation with water, a considerable quantity of essential oil separates from the watery sluid, of a pale yellowish colour, in smell exactly resembling the cardamoms, and of a very pungent taste: the remaining decoction is disagreeably bitterish and mucilaginous, retaining nothing

of the pungency or warmth, any more than of the peculiar flavour of the spice. On inspissating the tincture made in rectified spirit, a part of the flavour of the cardamoms arises with the spirit, but greatest part remains behind concentrated in the extract; which fmells moderately of the feeds, and has a pungent aromatic taste, very durable in the mouth, and rather

more grateful than that of the feeds in fubstance.

Tinctures of this spice both in rectified and proof spirit are more agreeable than the watery infusions; and proof spirit, impregnated with its flavour by distillation, more agreeable than the simple distilled water. A tincture of fix ounces of the feeds, in a quart + or two pints and a half t of proof spirit; and a spirituous water more lightly flavoured with them, by drawing off a gallon of proof spirit from four ounces; are kept in the shops, and occasionally made use of as pleasant warm cordials and for flavouring other medicines. I have not observed any of the aromatics to answer, in general, so well as the tincture of this fpice, for rendering mineral waters and other faline liquors acceptable to the stomach.

Tinet. cardamomi † Ph. Lond. ‡ Ph. Edinb. Aq. femin. cardam. Ph. Lond.

> 2. CARDAMOMUM MEDIUM Pharm. Parif. Cardamomum majus officinarum C. B. Greater cardamom: with thicker and tougher husks, an inch or more in length; the produce of a plant of the same kind with the preceding, but larger. There is some confusion in regard to the name, that of cardamomum majus being applied among us to this species, and in France to the grana paradifi, of which hereafter.

> The feeds of the greater cardamom are allowed by the faculty of Paris to be used indifferently with those of the lesser: the large kind, however, is much weaker than the other, both in fmell and tafte, and hence has in this country been long difregarded, and is now become a stranger to the shops. Both forts are nearly of the same nature, the difference being chiefly in degree.

CARDIACA.

CARDIACA seu Agripalma Pharm. Paris. Marrubium cardiaca dictum C. B. Leonurus Cardiaca Linn. Motherwort: a large plant, with fquare branched stalks, the leaves fet in pairs on long pedicles at the joints, and the flowers in clusters round the upper joints: the leaf is dark coloured, cut deeply into three sharp-pointed indented segments, of which the middle one is longest, and the two lateral ones commonly again deeply cut: the flower is purplish, labiated, with the upper lip long and arched, the lower short and cut into three sections. It is biennial, grows wild in waste grounds, and flowers in July.

This plant is faid to be useful in disorders of the stomach proceeding from thick phlegm; to loosen the belly; to promote perspiration, urine, and the uterine purgations. Such, in effect, are the virtues, which may be expected from its sensible qualities. The leaves and the tops have a moderately strong smell, not very agreeable; and a very bitter taste. In keeping for some time, or on boiling them in water, their smell is dissipated: the decoction, inspissated to the consistence of an extract, discovers to the taste a strong penetrating subsaline bitterness.

CARDUUS.

CARDUUS BENEDICTUS Pharm. Lond. & Edinb. Cnicus filvestris birsuitor sive carduus benedictus C.B. Centaurea benedicta Linn. Holy thistle: a plant with rough, narrow, jagged leaves, terminating in soft prickles; and large, hairy, branched stalks, leaning to the ground; on the tops of which grow large, scaly, prickly heads, including a number of yellow slosculi, which are followed by oblong striated seeds inclosed in down. It is a native of Spain and some of the islands of the Archipelago, and sown annually with us in gardens.

THE leaves of carduus have a penetrating bitter taste, not very strong, or very durable in the mouth; accompanied, in their recent state, with somewhat of an ungrateful slavour, which they soon lose in keeping. The herb, when thoroughly dried, should be hung up loosely in an airy place; being very subject, if pressed close, to rot or grow mouldy.

Cold water, poured on the dry leaves, extracts, in an hour or two, a light grateful bitterness: by standing long upon the plant, the liquor becomes disagreeable: a strong decoction is very nauseous and offensive to the stomach. A cold insusion and a decoction being separately inspissated, the same differences were observed between the extracts, as between the liquors in their dilute state; the extract obtained from the

infusion

infusion being a sufficiently agreeable bitter, and that from the decoction disgussful; a proof, that the differences of the liquors do not depend, as might be supposed, on their degree of saturation, but on their being

impregnated with matters of a different kind.

Rectified spirit also extracts, in a short time, the lighter bitter part of the carduus, but does not take up the nauseous near so easily as water: a spirituous tincture prepared by warm digestion for several hours, and the extract obtained by inspissating it, were more strongly but not unpleasantly bitter. The colour of the watery tinctures is a yellowish or greenish, inclining more or less to brown, according as they are more or less saturated; that of the spirituous, a deep green.

On keeping the foft watery extracts for some months, a considerable quantity of saline matter was found to have shot upon the surface, into small crystals, in shape approaching to those of nitre, in taste bitterish

with an impression of coolness.

The virtues of this plant feem to be little attended to in the present practice. The nauseous decoction is sometimes used to excite vomiting, and a strong insusion to promote the operation of other emetics: but this elegant bitter, when extracted from the offensive parts of the herb, may be advantageously applied to other purposes. I have frequently observed excellent effects from a light insusion of carduus, in weakness of appetite and indigestion, where the stomach was injured by irregularities and oppressed by viscid phlegm: nor have I found any one medicine of the bitter kind to sit so easily on weak stomachs, or to heat so little. These insusions, taken freely, promote the natural secretions. Drank warm in bed, they commonly increase perspiration or excite sweat; and as they act with great mildness, not heating or irritating considerably, they have been used, in this intention, in acute as well as chronical diseases.

The feeds of carduus are likewise considerably bitter, and have sometimes been used as sudorifies or diaphoretics, in the form of emulsion. Cartheuser observes, that they give the proper consistence of an emulsion to ten times their weight or more of water: and that they do not impart a perfect whiteness, but a greyish colour to the liquor.

CARICÆ.

CARICÆ Pharm. Lond. & Edinb. Figs: the dried fruit of the ficus communis C. B. Ficus Carica Linn. a tree of a middling fize, with large leaves cut into five fegments; remarkable for producing no flowers previous to the fruit; growing spontaneously in the warmer climates, and cultivated in our gardens.

Figs are accounted moderately nutrimental, grateful to the stomach, and easier of digestion than any of the other sweet fruits. Their principal medicinal use is as a lubricating emollient sweet; in which intention, they are commonly made an ingredient in pectoral decoctions, and in lenitive electraies. They are employed externally, in cataplasms, for promoting the suppuration of inflammatory tumours; for which purpose they appear to be equally adapted with other soft substances void of acrimony or irritation.

CARLINA.

CARLINA: a perennial plant, with long, narrow, deeply jagged, and very prickly leaves, lying on the ground; in the middle of which grows a large roundish head, without any stalk, encompassed with smaller leaves, full of sharp prickles: the flower issues from the middle of the head.

magno flore C. B. Carlina acaulis Linn. Cardopatium. CARLINE THISTLE: with the flower composed of a number of white petals set round a middle disk. It is a native of the mountainous parts of Italy and Germany, from whence the dried roots are sometimes brought to us. These are about an inch thick, externally of a rusty or reddish brown colour, internally of a pale yellowish or brownish, corroded as it were upon the surface, and perforated with small holes, so as to appear, when cut, as if worm-eaten.

THE roots of carline thiftle have a moderately strong, not agreeable smell; and a weak, bitterish, subacrid, somewhat aromatic taste. In-

(a) Expunged.

fusions of them in water have very little taste, and not much smell: distilled with water, they yield a two-hundredth part of their weight, or a little more, of a thick ponderous essential oil, which, on being rectified or redistilled, leaves a considerable proportion of resinous matter and becomes thin (a): the decoction, remaining after the separation of this most active principle of the root, is unpleasantly bitterish and subsaline, though only weakly so even when inspissated to an extract. A tincture and extract prepared with rectified spirit are stronger in taste than those made with water, but have little smell. Both the watery insusion and extract are of a brownish yellow colour, the spirituous of a deep gold yellow.

This root is supposed to be diaphoretic, antihysteric, and anthelmintic. It has been greatly esteemed by some foreign physicians in acute malignant as well as in chronical diseases; and given in substance from a scruple to a dram, and in insussion from one to two drams and more. It never came much into use among us, and is now rarely to be met with in the shops. Frederic Hossman the elder relates, that he has known a decoction of it in broth excite vomiting (b), but does not mention the quantity which produced this effect.

2. CARLINA GUMMIFERA; Carduus pinea; Ixine. Carlina acaulos gummifera C. B. Chamæleo albus diofcoridis Columnæ. Atractylis gummifera Linn. Pine thistle: with the flowers composed of purplish flosculi, like those of the common thistle. It is a native of Italy and the island of Candy.

THE roots of the pine thiftle are larger than those of the carline, and of a stronger smell. Wounded when fresh, they yield a viscous milky juice, which concretes into tenacious masses, at first whitish and refembling wax, when much handled growing black, supposed to be the ixion, ixia, and acanthina massiche of the ancients. The juice, in taste and smell not ungrateful, is said to have been formerly chewed for the same purposes as massich; and the root itself to be of the same virtue with that of the preceding species.

(a) Neumann, Chem. works, p. 406.

(b) Clavis Schrader. p. 431.

CARPOBALSAMUM.

CARPOBALSAMUM Pharm. Lond. The fruit of the tree that yields the balfam of Gilead. It is about the fize of a small pea, with a short pedicle; of a roundish or oval figure, pointed at the top; composed of a dark brown or reddish black, wrinkled bark, marked with four ribs from top to bottom, and a whitish or yellowish medullary substance (a).

This fruit, when in perfection, is faid to have a pleafant warm, bitterish taste, and a fragrant smell resembling that of the balsam itself. But such as is now and then met with in the shops, (for it is but rarely to be met with there) has almost wholly lost both its smell and taste. It is no otherwise made use of in this country than as an ingredient in mithridate and theriaca; in both which, its place is commonly supplied by materials of more efficacy than itself: some direct juniper berries, the London college cubebs, for its substitute.

CARTHAMUS.

CARTHAMUS Pharm. Parif. Cnicus sativus sive carthamus officinarum C. B. Carthamus tinctorius Linn. SAFFLOWER: a plant with
oval pointed leaves, somewhat prickly about the edges, joined close to
the stalk, which is round, sirm and branched: on the tops grow large
scaly heads, with saffron-coloured sistular slowers standing out from
them: these are followed by smooth white seeds, of an oblong roundish
shape, yet with four sensible corners, remarkably heavy so as to sink in
water. It is an annual plant, a native of Egypt, and cultivated in large
quantity in some parts of Germany on account of the uses of its slowers
in dying. It is sometimes raised among ourselves; but the seeds, which
are the part that has been chiefly made use of in medicine, seldom come
to perfection in this climate.

THE feeds of carthamus, freed from the shells, have an unctuous sweetish taste, which on chewing them for a little time becomes acrid

and disagreeable: they form an emulsion on trituration with water, and give out to spirit a little nauseous acrid matter. They have been celebrated as a gentle cathartic, in doses of a dram or two in substance and six or eight drams in emulsion: but as they operate very slowly, and are apt, especially when given in substance, to occasion nauseæ, slatulencies, and distensions of the stomach; their use has long been laid aside, and the colleges both of London and Edinburgh have now discarded them from their catalogues of officinals.

The flowers have been fometimes employed as a colouring drug for alimentary and medicinal fubstances; and when well cured, are not easily distinguishable by the eye from fasfron, though they have nothing of its smell or taste. They give a deep saffron tincture to rectified spirit, and a paler yellow to water. After the yellow matter has been extracted by water, the flower appears red, and communicates a deep red colour to spirit of wine or to alkaline lye.

CARUI.

CARUI Pharm. Lond. Carvi Pharm. Edinb. Cuminum pratense carui officinarum C. B. Carum Carvi Linn. CARAWAY: an umbelliferous plant, with striated branched stalks, two or three feet high; and finely divided leaves set in pairs along a channelled rib; every two of which ribs or pedicles cross one another at their origin on the stalk: the seeds are small, of a brownish or blackish colour, somewhat bent, striated, slat on one side, convex on the other. It is a native of the northern climates: in this kingdom it is rarely found wild, but commonly cultivated in gardens for culinary and consectionary as well as medicinal purposes. It is biennial.

CARAWAY feeds are an useful stomachic and carminative; of a sufficiently agreeable aromatic smell, and a moderately warm taste: they are given, in substance, from a scruple to a dram. The leaves have the same kind of slavour with the seeds, but are considerably weaker and less grateful. The roots have a sweetish taste, accompanied with a slight warmth, and very little smell.

The feeds give out the whole of their virtue, by moderate digestion, to rectified spirit; for after the action of this menstruum they prove

infipid and inodorous: the tincture taftes strongly of the caraways, but their smell is in great measure covered by the menstruum. The spirit, gently distilled off from the filtered liquor, brings over very little of the flavour of the caraways, leaving nearly all their active matter concentrated in the extract, which proves a very warm pungent aromatic. colour, both of the tincture and extract, is a yellowish verging to green.

Infusions of the seeds in water are stronger in smell than the spirituous tincture, but much weaker in taste: after repeated infusion in fresh portions of water, they still give a considerable taste to spirit. The colour of the watery infusions is a pale reddish brown. In distillation or evaporation, water elevates all the aromatic part of the caraways: the remaining extract is almost insipid, and thus discovers, that in caraways there is less, than in most of the other warm feeds of European growth, of a bitterish or ungrateful matter joined to the aromatic. Along with the aqueous fluid there arises in distillation a very considerable quantity, about one ounce from thirty, of effential oil, of a bright yellow colour, Ol. effent. fmelling strongly of the caraway, in taste hotter and more pungent than fem. carai those obtained from most of our other warm feeds *(a): this oil is given from one to five or fix drops, as a carminative; and is supposed also to be of peculiar efficacy for promoting urine, to which it communicates fome degree of its smell. The leaves of the plant afford likewise an oil, nearly fimilar, both in colour and quality, to that of the feeds, but in far less quantity: fixteen pounds of the herb in flower, stripped from the stalks, yielded scarcely an ounce. The essential oil of the seeds is directed as an officinal; as also a cordial water, pretty strongly flavoured Aqua semin. with them by drawing off a gallon + or nine pints tof proof spirit from carui half a pound of the caraways.

+ Pb. Lond. t Ph. Edinb.

CARYOPHYLLA.

CARYOPHYLLA AROMATICA Pharm. Lond. & Edinb. Caryophyllus aromaticus seu potius garyophyllus Pharm. Paris. the unripe fruit, or perhaps more properly the cups of the unopened flowers, of a bay-like tree growing in the East Indies; Caryophyllus aro-

*(a) M. Beaumé obtained from fix pounds of unbruifed caraway feeds, four ounces of effential oil, as colourless as water.

maticus

maticus Linn. In shape they somewhat resemble a short thick square nail, of a rusty colour inclining to black: in the inside of each clove are found a stylus, and stamina, with their apices: at the larger end shoot out, from the four angles, four little points like a star; in the middle of which is a round ball, of a lighter colour than the rest, composed of four small scales or leaves, which seem to be the unexpanded petala of the slower. The tree is one of those, whose slower is produced above the rudiments of the fruit: the ripe fruit, sometimes brought into Europe under the name of anthophyllus, is marked on the top with the remains of the slower; it is about the size and shape of an olive, and contains, under a thin blackish shell, a hard kernel of the same colour, which has a deep longitudinal seam on one side. The cloves are said to be cured by exposing them to smoke, and afterwards drying them in the sun.

THE clove has a strong agreeable smell, and a bitterish, hot, very pungent taste: it is one of the hottest and most acrid of the substances of the aromatic class, and as such is often used, not only internally, but as an external stimulant. The antophyllus has the same kind of slavour with the clove itself; but being far weaker, in smell as well as in taste, it is very rarely applied to any medicinal purposes, and is now scarcely ever to be met with in the shops.

The clove is remarkably disposed to imbibe humidity; and when robbed of its active parts by infusion in menstrua or distillation, and afterwards mixed with fresh cloves, it regains from them a considerable share both of taste and smelt. The Dutch, through whose hands this spice is brought to us, have often practised this abuse; which, however, may be easily discovered; for those cloves which have once lost their virtue, continue always not only weaker than the rest, but likewise of a much paler colour.

Tinctures of cloves in rectified spirit are of a dark reddish brown colour, of no great smell, but of a highly acrid taste: if the quantity of spirit be considerable, it leaves the clove deprived of all its virtue. On inspissating the filtered tincture, the spirit, which distils, is found to have very little impregnation from the spice: the remaining extract, nevertheless, does not discover so much smell as the clove in substance, but its taste is excessively pungent and fiery. The quantity of this burning extract amounts to about one third the weight of the clove.

Digested

Digested or insused in water, they impregnate the liquor more strongly with their smell than they do spirit, but not near so much with their taste: after repeated insusion in water, they impart still a considerable tincture to rectified spirit. In distillation with water, they give over, very slowly, near one sixth their weight (a) of essential oil; when carefully distilled, colourless; by age, changing to a yellow, and at length to a reddish brown colour; when drawn with a strong sire, proving often of this colour at first; smelling strongly of the cloves; but in taste only moderately pungent, very much less so than the spirituous extract. Neither the remaining clove nor decoction have any considerable taste; the pungency of this spice seeming to depend, not on the volatile or fixt parts separately, but on the combination of the two (b).

The oil of cloves commonly met with in the shops, and received from the Dutch, is indeed highly acrimonious: but this oil is plainly not the genuine distilled oil of the clove; for notwithstanding its being more pungent than that which cloves afford by the common process of distillation, it contains a large admixture, oftentimes half its weight or more, of an insipid expressed oil; as appears upon treating it with rectified spirit, which dissolves the pungent aromatic matter, and leaves the gross insipid oil. It is probably from an admixture of the resinous part of the clove, that this sophisticated oil receives both its acrimony and high colour. Fresh cloves are said to yield a high coloured, thick, fragrant oil upon expression: possibly the common oil of cloves, brought from the spice islands, is no other than this oil, diluted with insipid ones. The college of London seems to require both the common and the genuine oil to be kept in the shops, making essential oil of cloves both an article of the materia medica and an officinal preparation: perhaps the common

CARYOPHYLLATA.

oil, as being most pungent, is best adapted for some external purposes, as

CARYOPHYLLATA Pharm. Edinb. (c) Caryophyllata vulgaris C.B. Geum urbanum Linn. Avens, or Herb-Benit: a roughish plant, with

the genuine doubtless is for those of an internal aromatic.

⁽a) Hoffmann, Observationes physico-chymicæ, lib. i. obs. 3.

⁽b) Neumann, De caryoph. aromat. Chem. works, p. 413. Cartheuser, m. m. ii. 383.

⁽c) Expunged.

dark-coloured winged leaves, resembling those of agrimony; and pentapetalous yellow flowers, standing in ten-leaved cups on the tops of the branches, followed, each, by a round cluster of hairy seeds with hooked tails: the roots are slender, full of fibres, of a dark brownish colour on the outside, and reddish within. It is perennial, grows wild in woods and hedges, and is found in flower greatest part of the summer.

THE root of avens has been employed as a gentle flyptic, corroborant, and stomachic; and for these intentions continues, not undeservedly, of some esteem in foreign countries, though very little regarded among us. It has a mildly austere somewhat aromatic taste; and a very pleasant smell, somewhat of the clove kind, especially in the spring, and when produced in dry warm soils: such as is the growth of close, shady, moist places, has little and often nothing of this flavour.

This root gives out its astringent matter equally to watery and spirituous menstrua, its aromatic part most perfectly to the latter: the aqueous insusion is of a reddish brown colour, the spirituous of a deep yellow. In distillation with water, it yield a small quantity of a whitish concrete oily matter, of a very grateful fragrance: the remaining decoction, inspissated to the consistence of an extract, is moderately astringent. On committing to distillation the spirituous tincture, little or nothing comes over with the spirit; the aromatic part of the root, as well as the austere, remaining concentrated in the extract. The smell, which in the tincture is concealed or suppressed by the menstruum, discovers itself again when the spirit is drawn off.

CARYOPHYLLUS RUBER.

phyllus altilis major C. B. Dianthus Caryophyllus Linn. CLOVE-JULY-FLOWER, or GILLIFLOWER: a plant with many smooth round jointed stalks, and grass-like blueish-green leaves standing in pairs at the joints: the flower is composed of five petals, narrow at the bases, broad and jagged at top, set in an oblong cylindrical cup, which is covered at bottom with four short scales forming as it were a secondary cup: after the flower has fallen, the cup becomes a covering to a number of small, stat, wrinkled, black seeds. It is perennial, and said to be a native of Italy.

MANY

MANY species or varieties of these flowers are common in our gardens. Those employed for medicinal uses, to which the name of clove-julyflower is more particularly appropriated, are of a deep crimfon colour, and a pleasant aromatic smell somewhat akin to that of the clove spice: their taste is bitterish and subastringent. In drying, their taste becomes stronger, and their smell is not so soon dissipated as that of many other fragrant flowers.

Clove-july-flowers have been recommended as cardiacs and alexipharmacs. Simon Paulli relates, that he has often cured malignant fevers by the use of a decoction of them; which, he says, powerfully promotes fweat and urine without greatly irritating nature, and at the

fame time raifes the spirits, and abates thirst.

At present, these flowers are valued chiefly for their fine flavour; which is readily extracted by infusion in water, and distipated even by light coction. Three pounds of the fresh flowers clipt from the heels, Syr. carycommunicate, by infusion in a close vessel for a night, a grateful and oph rubromoderately strong smell, and a deep red colour, to five + and even to + Ph. Lond. twelve pints of water: these liquors, with a proper quantity of fine t Pb. Edinb. fugar, form very agreeable fyrups. On diffilling the fresh flowers with water, the distilled liquor proves considerably impregnated with their fragrance, but no effential oil feparates, though feveral pounds of the flowers be submitted to the operation. The remaining decoction is of a deep red colour, and yields, upon being inspissated, a dark purplish red extract, of little or no smell, and of a bitterish, austere, subsaline tafte.

Rectified spirit, digested on the flowers, receives a much paler tincture than watery liquors, but extracts the whole of their active matter. In distillation or evaporation, spirit elevates much less than water; the spirituous extract retaining a considerable share of the fine smell of the flowers, as well as their tafte: its colour is purplish, like that of the watery extract.

CASIA CARYOPHYLLATA.

CASIA CARYOPHYLLATA Pharm. Parif. Cortex caryophyl-CLOVE BARK: the bark of a tree of the clove kind, (caryophyllus aromaticus fructu rotundo, caryophyllon plinii C. B. Myrtus caryo-Gg phyllata

phyllata Linn.) brought from the island Cuba, Jamaica, and other parts of the West Indies; rolled up in quills; like cinnamon, but somewhat thinner, rougher on the outside, and of a darker rusty brown colour.

This bark is a warm aromatic, nearly of the same kind of smell and taste with the clove spice, but weaker, and with a little admixture as it were of the cinnamon flavour. It agrees nearly with cloves also in regard to the solubility and volatility of its active principles. Tinctures of it in rectified spirit smell and taste strongly of the bark: the watery infusions are considerably impregnated with its smell, but have very little of its taste. On inspissating the spirituous tincture, the spirit which distils has little or nothing of its slavour: the remaining extract smells lightly of the bark, and proves in taste very hot and pungent, though much less so than the spirituous extract of cloves. In distillation with water, it yields a very small portion of essential oil, nearly similar in slavour to the oil of cloves, but more pungent than the genuine oil of that spice: the remaining decoction is ungratefully austere and bitterish.

A bark of the same kind is sometimes brought from the East Indies under the name of culitlawan or culilawan; a Malaccan compound word, of which the Latin cortex caryophylloides or clove bark is said to be a translation. That distinguished in Europe by the name of culilawan is thicker than the other, and in colour approaches somewhat more to cinnamon, but scarcely differs in smell or taste.

The same with this appears likewise to be the carabaccium of Baglivi; which he describes as being in taste like cloves, but very temperate and grateful, and in colour having a great resemblance to cinnamon; and which, he says, he made use of, with great benefit in decoction, for correcting acrimony and scorbutic dissolution of the lymph, and for strengthening the stomach and promoting digestion (a).

Rumphius observes, that the outer and inner barks, and the barks of different parts of the tree, differ somewhat in colour and in taste from one another; (whence, probably, such differences as may have been observed in those brought under different names into Europe); and that

the bark of the root approaches both in appearance and in flavour to faffafras, to which it was, in Batavia, frequently substituted.

CASIA FISTULARIS.

CASIA FISTULA: a hard woody cylindrical pod, of a tree refembling the walnut, (cassia fistula alexandrina C. B. & Linn.) which grows spontaneously in Egypt and the warmer parts of the East Indies, and has been thence introduced into the West. The pods or canes are about an inch in diameter, and a foot or more in length; externally, of a dark brown colour, somewhat wrinkled, with a large seam running the whole length upon one side, and another less visible on the opposite one; internally, of a pale yellowish colour, divided by thin transverse woody plates into a number of little cells, containing each a flattish oval seed with a soft black pulp.

The pulp of casia has a sweetish taste, followed by more or less of an ungrateful kind of acrimony. The casia of the East Indies has a more agreeable sweetness, and less acrimony than that of the West; and hence the former is universally preferred: they may be distinguished from one another by the eye; the oriental canes being smaller, smoother, and thinner-rinded, and their pulp of a deeper shining black colour, than the occidental. The lighter canes of either fort, and those in which the seeds rattle on being shaken, are generally rejected: in these, indeed, the pulp has become dry, but it does not necessarily follow that it is damaged: it loses nothing in drying but its aqueous humidity, and by this loss it should seem to be effectually secured from growing mouldy or four, inconveniences to which in its moist state it is very subject.

The pulp of casia, whether moist or dry, dissolves both in water and in rectified spirit; readily in the former, slowly and dissicultly in the latter, and not totally in either: the part which remains undissolved appears to be of little or no activity. It is usually extracted by boiling the bruised pods in water, and evaporating the strained solution to a due consistence: the exhaling vapour carries off nothing considerable of the casia. As it is very apt to grow sour in keeping, only small quantities should be prepared at a time.

Gg2

Cafia,

Casia, in doses of a few drams, is a gentle laxative; of good use in costive habits, in inflammatory cases where purgatives of the more acrid or irritating kind can have no place, and, as Geoffroy observes, in the painful tension of the belly which sometimes follows the imprudent use of antimonials. It is rarely given in such doses as to have the full effect of a cathartic; the quantity necessary for this purpose, an ounce and a half or two ounces, being apt to nauseate the stomach, and produce flatulencies and gripes, especially if the casia is not of a very good kind: mild aromatics, and dilution with warm liquors, are the best correctives.

It is fometimes acuated with the stronger purgatives, or with the antimonial emetics; of which last it is said by some to diminish the activity so far, that four grains and a half of the tartarum emeticum may be taken, in a decoction of casia, by those who can bear but one grain and a half of the antimonial preparation by itself (a). It is often joined also as an auxiliary to the milder purgatives, as crystals of tartar, tamarinds, and manna; and of these, particularly of the latter, it is supposed to increase the cathartic virtue: a mixture of four drams of casia, and one and a half or two of manna, is said, by Vallisneri, to purge as much as twelve drams of casia or thirty-two of manna by themselves. * In the shops are kept electaries of this kind, composed of six ounces each of pulp of casia and syrup of pale roses; and two ounces; or one and a half; of manna; with one ounce; or one and a half; of pulp of tamarinds.

Elect. e Cafia † Pb. Lond. ‡ Pb. Edinb.

It is observable, that during the use of casia, the urine appears frequently of a green colour, and sometimes, where the quantities taken are considerable, of a dark brown or blackish.

CASIA LIGNEA.

CASIA LIGNEA Pharm. Edinb. & Parif. CASIA LIGNEA: the bark of a tree of the cinnamon kind, (cinnamomum feu canella malavarica & javenensis C. B. Laurus Cassia Linn.) brought from the East Indies*(b); exactly resembling cinnamon in appearance, but dis-

(a) Malouin, Chimie medicinale, part. iii. chap. 38.

*(b) Bergius in his Mat. Med. gives the Laurus Malabatrum as the tree producing the cassia lignea brought from the East Indies, and the Laurus Cassia as the West India cinnamon or cassia,

which he fays is stronger than the former.

tinguishable

tinguishable by its breaking short or smooth, while cinnamon breaks fibrous or shivery like wood.

This bark resembles cinnamon in aromatic flavour as well as in external appearance; but differs in being weaker, or containing less active matter, and in its abounding with a viscous mucilaginous substance. Chewed, it disfolves as it were in the mouth into a kind of slime: powdered and boiled in water, it renders a confiderable quantity of the fluid thick and glutinous, fo as to concrete on cooling into the confiftence

of a gelly.

Rectified spirit of wine, digested on the bark, dissolves and extracts its aromatic matter; the powder retaining its mucilage, fo as to form a gelly with water as at first. The aromatic part may be separated also by distillation with water; in which process, if a large quantity of casia is used, a small portion of effential oil may be collected. * M. Beaumé procured two drams and a half of oil, from twelve pounds and a half of casia. The spicy principle of the casia, thus freed from the mucilage, in the form of spirituous tincture, or spirituous extract, or distilled water, or effential oil, appears the same with that of cinnamon; provided, in regard to the distilled fluids, that they have not received an empyreumatic taint in the operation, an inconvenience to which they are very subject on account of the mucilaginous matter fwelling up and burning to the veffel. * The Edinburgh college have now directed a fimple water to Aq. Caffiæ be kept, ten pints of which are drawn from a pound and a half of caffia Edinb. lignea.

Cafia lignea was employed by the ancients as a fuccedaneum to cinnamon, of which it was reckoned equivalent to half its own quantity. At present, it is not unfrequently mixed with, or substituted to, that spice in the shops, but is scarcely ever made use of under its own name.

CASTOREUM.

CASTOREUM Pharm. Lond. & Edinb. CASTOR: the inguinal glands of the castor or beaver; a four-footed amphibious animal, frequent in feveral parts of Europe, and in North America. These glands are of different shapes and fizes, covered with a thick skin, including an unctuous liquid matter, which in keeping grows dry and hard: on cutting cutting the dry cods, as they are called, they are found full of a brittle friable substance, of a brownish red colour, interspersed with fine membranes and fibres exquifitely interwoven. The best castor comes from Russia in large, round, hard cods; an inferior fort, smaller and moister, from Dantzick; the worst of all from New England, in longish thin cods.

Russra caftor has a ftrong not agreeable fmell, and a biting bitterish nauseous taste: the other forts are weaker than that of Russia, yet more ungrateful. It is generally looked upon as one of the capital nervine, antifpafmodic and antihysteric medicines: its virtues have undoubtedly been much exaggerated; but though they are not near fo great as they have by most writers been represented, they appear nevertheless to be confiderable. The common dose is from two or three grains to a scruple; though it has been fometimes taken by drams, and these doses very often

repeated (a).

Rectified spirit, proof spirit, and water, extract, by the affistance of heat, nearly all the active matter of castor: rectified spirit dissolves most readily the finer and less ungrateful, and water the more nauseous bitter part: proof spirit acts equally, but difficultly, on both. Three ounces of powdered Russia castor, digested with a gentle heat, for three days, in a quart of rectified spirit, communicated a deeper reddish tincture, and a stronger though rather less ungrateful taste, than four ounces did to the fame quantity of proof spirit by maceration without heat for ten days. On digefting in the two spirits equal quantities of the powder, with equal degrees of heat, for ten or twelve hours, the tincture in proof spirit proved fenfibly most ungrateful, and that in rectified spirit possessed most perfectly the specific flavour of the castor: an infusion in boiling water was bitterer and more nauseous than either. The castor remaining after the action of water, retained a little of its flavour, but nothing of its nauseous bitterness: that left by rectified spirit retained a little of the latter, but nothing of the former.

Tinet. caf-Pb. Edinb.

Ph. Lond.

In distillation, it gives over to water the whole of its smell and Aq. castorei flavour: a quart of water, distilled from an ounce of Russia castor, receives a confiderably strong impregnation, but gradually loses greatest

part of it in being kept. It is faid, that on submitting to this operation large quantities of the caftor, a small portion of effential oil is obtained, which fmells exceeding strongly, and diffuses its ungrateful scent to a great distance (a). This odorous and most active principle of the castor is carried off by water in a very gentle heat; infusions or light decoctions, which are very naufeous, yielding, however flowly inspiffated, a brittle extract, which has nothing of the specific flavour of the drug, and proves in tafte but weakly though ungratefully bitterish. Rectified spirit on the other hand, distilled from the tincture made in that menstruum, brings over fearcely any fenfible impregnation; nearly all that it had extracted from the castor, remaining entire in the inspissated mass, which proves of an unctuous confistence; not easily reducible to dryness.

Caftor is commonly joined in prescription with the deobstruent fetid gums, volatile alkaline falts, the volatile oily spirits, and other materials of fimilar intention. The volatile oily spirits are well adapted also as menstrua for dissolving the active matter both of the castor and of the fetid gums; at the same time that they prove in many cases excellent additions to their virtue, as particularly in some hysteric disorders, and the feveral fymptoms which accompany them: in this view, an ounce of Russia castor, and half as much asafetida, are digested about six days, Tine. casin a close vessel, with a pint of the volatile spirit.

tor. comp. Ph. Edinb.

CASUMUNAR.

CASUMUNAR Pharm. Lond. Bingalle; Rifagon. CASMUNAR: the root of an East India plant, of which we have no certain account; brought over in irregular flices of various forms, fome cut transversely and others longitudinally: the cortical part is marked with circles, and of a dusky brownish colour: the internal part is paler, and unequally yellow.

This root was introduced fome time ago by Marloe, as a medicine of uncommon efficacy in hysteric, epileptic, paralytic, and other nervous disorders. At present it is sometimes employed as a stomachic, but its use is not yet become so general as it seems to deserve. It is an elegant

mild aromatic, moderately warm, lightly bitterish, in smell somewhat resembling ginger. Its virtues are extracted in perfection by rectified spirit, and, on drawing off the menstruum from the filtered tincture, remain concentrated in the inspissated mass; which smells very agreeably, and impresses on the organs of taste a grateful bitterishness, and a durable glowing warmth, not a fiery or pungent heat. Both the tincture and extract are of a deep saffron colour.

CEDRUS.

CEDRINUM LIGNUM Pharm. Parif. Cedrus conifera foliis laricis C. B. Pinus Cedrus Linn. CEDAR OF LIBANUS: a large evergreen coniferous tree, with very narrow stiff sharp-pointed leaves standing several together in tusts. It is a native of the bleak snowy mountains of Syria, and is not as yet become common in this kingdom.

THE cedar is one of the odorous refiniferous trees; in its general medicinal qualities fimilar to the fir, but in fome respects different. The refinous juice, extracted from incisions made in the trunk, has a stronger and more agreeable kind of fmell, and is much more disposed to concrete into a folid brittle mass, without losing much of its valuable parts in the exficcation. The wood, which is of a fine reddish colour and very light, is likewise more fragrant than the fir, and its odorous matter less volatile: a tincture of it in rectified spirit, which is reddish like the wood itself, being committed to distillation, the spirit brings over nothing of its virtue; all the active matter of the cedar remaining behind, concentrated into an elegant balfamic extract. Even boiling water does not eafily carry off its flavour: the watery extract smells considerably of the wood, and is in tafte bitterish and saline. Marggraf relates, that on keeping the extract for some time, small crystals shot upon the surface, which were found on trial to be common falt: and that on distilling the wood with water, it yielded about one fixty-fourth its own weight of a thick, yellowish, essential oil, which grew thicker in a moderate degree of cold, and quite confistent in a strong one (a). In the saline nature of the watery extract, this wood differs from all the refinous ones that

have been examined; and in the thickness, and congelability of its effential oil, from all but the lignum aloes and yellow faunders.

CENTAURIUM.

CENTAURIUM MINUS Pharm. Lond. & Edinb. & C.B. Gentiana Centaurium Linn. Lesser centaury: a small plant, with three-ribbed, somewhat oval leaves, set in pairs on the stalks; which divide, towards the top, into several branches, bearing umbel-like clusters of bright red, sunnel-shaped flowers, cut into sive acute segments, followed by little oblong capsules full of very small seeds. It is annual, grows wild in dry pasture grounds, and slowers in July.

THE leaves and tops of centaury are strong bitters, of scarcely any smell or particular flavour. The seeds also are very bitter; the petala of the flowers, and the roots, almost insipid. The flowery tops are generally made choice of, and are of considerable estimation in the present practice as corroborant stomachic bitters.

The active parts of this plant are diffolved readily both by water and rectified spirit, the herb, after insusion in sufficient quantities of either menstruum, remaining insipid: insusions of the leaves in water are of a paler or deeper brownish colour, according as they are less or more saturated; to rectified spirit, the fresh leaves give a green, the dry a dark brownish red tincture. All these liquors are sufficiently elegant bitters.

Water takes up, along with the bitter, a large quantity of an infipid mucilaginous substance, whereas rectified spirit seems to dissolve little more than the pure bitter part. Hence, on inspissating the two solutions to the same consistences, the watery extract proves much less bitter than the spirituous, and its quantity above four times greater: according to Cartheuser's experiments, an ounce of the herb yields with water above half an ounce of extract, but with spirit scarcely two scruples.

CENTAURIUM MAJUS.

CENTAURIUM MAJUS, sive Rhaponticum vulgare officinarum, Pharm. Paris. Centaurium majus folio in plures lacinias diviso C.B. Centaurea Centaureum Linn. GREAT CENTAURY: a large plant, with the leaves composed of oblong serrated segments set in pairs on a middle Hh

rib, which is edged, in the intermediate spaces, with a serrated margin: the stalk divides, towards the upper part, into several branches, bearing, on the tops, round soft scaly heads, from which come forth blueish slosculi, followed by down inclosing the seeds. It is perennial, a native of the southern parts of Europe, and raised with us in gardens.

The root of this plant, of a dark blackish colour on the outside, is internally reddish, and yields, when fresh, a juice of a deep red. It has a slight smell, not disagreeable; and in chewing discovers a viscous sweetishness and roughness, with some degree of acrimony. It is reckoned aperient and corroborant, and supposed to be particularly useful in alvine sluxes; in which intention it has by some been greatly recommended, though apparently much inferiour to the root whose place it was employed to supply, to wit, the true rhapontic. Among us it has long been discarded from practice, and is now dropt by the colleges both of London and Edinburgh.

CEPA.

CEPA Pharm. Lond. Cepa vulgaris C. B. Allium Cepa Linn. Onion: a plant with a fingle bulbous root composed of a number of coats; producing long fistular leaves, and a tall naked bellied stalk, bearing a large cluster of hexapetalous white flowers, each of which is followed by a roundish capsule, containing a number of black angular seeds. It is cultivated in gardens for culinary uses.

Onions are very acrid, stimulating, and of little nourishment. Taken freely in hot bilious dispositions, they produce statulencies, thirst, headachs, and febrile symptoms. In cold sluggish phlegmatic temperaments, they are of service; warming the habit, attenuating viscid humours, and promoting the natural excretions, particularly expectoration and urine. They are likewise powerfully antiseptic, and by virtue of this quality are recommended by some as a salubrious addition to the food in scorbutic cases. Externally, they are employed in cataplasms for suppurating hard tumours: some recommend them also to be rubbed on bald places for promoting the growth of hair. Frederic Hoffman reports, that suppressions of urine, in children, are speedily relieved, by the application of roasted onions to the region of the pubes.

The

The root, which is the most acrid part of the plant, loses greatly in drying, both of its smell and taste, together with near seven eighths of its weight. It gives out its virtue, by infusion, both to water and to rectified spirit, but not readily, nor completely, to either. In distillation, the whole of its acrimony and peculiar slavour arise with water, and a very considerable part with spirit: the distilled water smells exceeding strongly and offensively of the onion, but no essential oil could be obtained on submitting to the operation several pounds of the root. The watery decoction, inspissated to the consistence of an extract, is very mucilaginous, but of scarcely any particular taste; and the taste of the spirituous extract is very weak. The active matter of onions appears therefore to be of a much more volatile kind than that of garlic, with which, in other respects, it nearly agrees.

CERA.

BEES WAX: a folid concrete, collected from vegetables by the bee; and extracted from the combs, after the honey is got out, by heating and prefling them: lighter than water, heavier than proof spirit; soluble in rectified spirit, very sparingly, and not without the affishance of heat, into a gelatinous liquid; not dissoluble at all in watery liquors; melting, by a heat a little greater than that which the hand can support, into the appearance of oil, and in this state easily miscible with oils and liquid sats; readily inflammable, and burning totally away; almost totally arising in distillation, partly in form of a thick empyreumatic oil, and partly in that of a consistent butyraceous matter, which by repeated distillation becomes sluid and thin.

- 1. CERA FLAVA Pharm. Lond. & Edinb. Yellow wax; in the state wherein it is obtained from the combs. When new, it is of a lively yellow, somewhat tough, yet easy to break: by age, it loses its fine colour, and becomes harder and more brittle.
- 2. CERA ALBA Pharm. Lond. & Edinb. White wax: the yellow wax artificially bleached, by reducing it into thin flakes, exposing these for a length of time to the sun and open air, and sprinkling them occa-sionally with water: when sufficiently whitened, the wax is melted and cast into cakes.

YELLOW

Yellow wax, when in perfection, has an agreeable smell, somewhat resembling that of honey: by long keeping, and in the process by which it is whitened, its smell is in good measure distipated. Distilled with water, by a boiling heat, it impregnates the liquor slightly with its scent, but gives no appearance of any essential oil; nor is the whole of its odorous matter to be easily separated by this process. Chewed, it proves tenacious, does not mingle with the saliva, or discover any particular taste. The gelatinous solution, obtained by boiling it in spirit of wine, by mixture with a thick mucilage of gum-arabic, becomes soluble in water, so as to form therewith an emulsion or milky liquor: the wax itself is made in like manner soluble, without the intervention of spirit, by thoroughly mixing it with the gum in fine powder: when thus dissolved, it proves still insipid, and perfectly void of acrimony.

The chief medicinal use of wax is in plasters, unguents, and other like external applications; partly for giving the requisite consistence to other ingredients; and partly on account of its own emollient quality. The yellow sort, dissolved into an emulsion, or mixed with spermaceti, oil of almonds, conserve of roses, &c. into the form of an electary; or divided, by stirring into it, when melted over a gentle fire, as much, as it will take up, of powdery matters, as the compound crabs-claw powder; is given also internally, and often with great success, in diarrhæas and dysenteries, for obtunding the acrimony of the humours, supplying the natural mucus of the intestines, and healing their excoriations or erosions.

The empyreumatic oil, into which wax is refolved by distillation with a strong heat, is greatly recommended by Boerhaave and others, for healing chaps and roughness of the skin, for discussing chilblains, and, with the assistance of proper fomentations and exercise, against stiffness of the joints and contractions of the tendons. It is, doubtless, highly emollient, but does not appear to have any other quality by which it can act in external applications: it has nothing of the acrimony or pungency, which prevail in all the other known distilled vegetable oils; though in smell it is not a little disagreeable and empyreumatic, a circumstance which occasions it to be at present more rarely used than it has been heretofore. As the wax swells up greatly in the distillation, it is convenient to divide it, by melting it with twice its weight of sand, or putting the sand above it in the retort, that it may mingle with the wax when brought into suspenses. The oil, which is preceded by a small quantity

quantity of acid liquor, congeals in the neck of the retort, from whence it may be melted down by applying a live coal, and made fluid by rediffilling it two or three times without addition.

CETERACH.

ASPLENIUM five Ceterach Pb. Parif. Ceterach officinarum C. B. Scolopendria vera Tragi. Afplenium Ceterach Linn. Spleenwort or Miltwaste: a small bushy plant, growing in sissures of rocks and old walls; consisting of capillary blackish roots, and long narrow leaves, cut down to the rib, on each side, alternately, into a number of oblong obtuse narrow sections with broad bases. It has no stalk or slower: the seeds are a yellow powder produced on the backs of the leaves.

The leaves of ceterach have an herbaceous, mucilaginous, roughish taste, and no considerable smell: with solution of chalybeate vitriol, they strike a blackish colour. They stand recommended as a pectoral, similar to maidenhair, to which they have been frequently joined in insussions and apozems; and likewise as an aperient in obstructions of the viscera. Mr. Morand relates, that there has lately been discovered in them an excellent diuretic virtue; that they were used with great success by count D'Auteuil, a Spanish naval commander, against the gravel, with which he was violently tormented; that they have since come greatly into use at Paris, Verdun, and Grenoble; that from the observations made there, they appear to gently carry off sand, cleanse the kidneys, and allay pains in the urinary passages; and that the way of using them is, to drink insusions of them in the morning as tea, with the addition of such other medicines as particular cases may require.

CEVADILLA.

CEVADILLA hispanorum Ph. Paris. (i. e. hordeolum). Sebadilla. Sabadilla. Hordeum causticum C. B. Indian caustic barley: the seed-vessel of a Mexican plant, resembling in its form and structure a barley ear, but with smaller seeds, not above the size of linseed.

These seeds appear, from the accounts given of them by some authors, to be the strongest of the vegetable caustics. Monardes reports, that in putrid

putrid verminous ulcers, and gangrenes, they have the same effects as corrosive sublimate, or the actual cautery; and that the way of using them is, to sprinkle a little of the powdered seed upon the part, or, for the greater safety, to dilute it with watery liquors, and apply lint dipt in the mixture (a). In Linnæus's amænitates academicæ they are said to be the most effectual of all medicines for destroying cutaneous insects in children. In France, they are ranked among the officinals: in this country they are very rarely to be met with.

CHÆREFOLIUM.

CHEREFOLIUM Pharm. Edinb.(b) Chærophyllum fativum C. B. Gingidium. Scandix Chærefolium Linn. Chervil: an umbelliferous plant, with winged leaves somewhat like those of parsley, producing smooth longish seeds shaped like a bird's beak. It is a native of the southern parts of Europe, and sown annually with us in gardens.

CHERVIL is a falubrious culinary herb; fufficiently grateful both to the palate and stomach; slightly aromatic; gently aperient and diuretic. The expressed juice is recommended by Riverius and others in dropsies; and Geossfroy relates, that he has found it, from experience, of remarkable service in this disease, that it acts mildly and without irritation, and abates inflammatory symptoms. He directs three or four ounces of the depurated juice, with seven or eight grains of nitre, and half an ounce of a syrup of the sive opening roots, to be taken daily every four hours; though he intimates also that the chervil juice has succeeded without any affistance. He observes that it is to be used with caution where the patient is troubled with a cough or a spitting of blood, as being liable to aggravate these complaints, in consequence of a nitrous salt by virtue of which he supposes this juice to act.

The depurated juice, inspissated to the consistence of an extract, is manifestly saline to the taste, but not entirely of the nitrous kind: it is more pungent than nitre, and did not visibly deslagrate in the fire. Of the aromatic slavour of chervil, little or nothing accompanies the juice; though water, as well as spirit, extracts greatest part of it by insusion.

The aromatic matter of this herb is of a very volatile kind, being foon distipated in drying or boiling: in distillation with water, there separates from the aqueous sluid a small portion of essential oil, resembling in taste, as Hossman observes, the essential oil of sensel seeds.

CHAMEDRYS.

CHAMEDRYS Pharm. Lond. Chamædrys minor repens C. B. Chamædrys vulgo vera existimata J. B. Teucrium Chamædrys Linn. Germander: a low creeping shrubby plant; with square stalks; small stiff oval leaves, notched from about the middle to the extremity, like those of the oak tree, set in pairs at the joints; and purplish labiated slowers, set thick together, wanting the upper lip. It grows wild in some of the woods of France, Germany, and Switzerland: with us it is raised in gardens, and slowers in June and July.

THE leaves and tops of germander have a moderately bitter taste, accompanied with a weak aromatic flavour, which is diminished a little in drying, but not totally diffipated in keeping for feveral months. They stand recommended as mild aperients and corroborants, in uterine and other obstructions, intermitting fevers, and in the rheumatism and gout. They make a principal ingredient in the alterative antiarthritic compositions prescribed by the ancients; whose use has lately been revived, with little variation; and which are faid, when long perfifted in, by strengthening the habit, rendering the blood more fluid, and promoting perspiration, to prevent returns of the gouty paroxysms. In fome arthritic cases, these and other warm bitter medicines have been of confiderable fervice: in others, they have been continued for years without any apparent benefit: in others, particularly in hot dispositions, in persons of an advanced age, and who had long suffered the disease, the abatement they procured of the gouty paroxysms has been followed by fymptoms more alarming.

The tops of the plant, gathered when the feeds are formed, are generally preferred to the leaves. Their dose, in substance, dried and powdered, is from half a dram to a dram or more. They give out their virtues both to watery and spirituous menstrua; and tinge the former of a yellowish colour inclining more or less to brown, according to the

degree

degree of faturation; the latter of a deep green. Water feems to disfolve the bitter matter more perfectly than pure spirit; the watery extract being stronger in taste than the spirituous, though the quantity of both extracts, according to Cartheuser's experiments, is very nearly alike.

CHAMEMELUM.

CAMOMILE: a plant with finely divided leaves; and moderately large flowers, standing solitary on the tops of the stalks, upon long naked pedicles: the flower is composed of a number of white petala, set round a yellow convex disk.

1. CHAMÆMELUM Pharm. Lond. & Edinb. Chamæmelum nobile seu leucanthemum odoratius C. B. Anthemis nobilis Linn. Chamomilla. Trailing perennial camomile, called Roman. It is found wild in moist pasture grounds in several parts of England, but commonly cultivated in gardens, and flowers in June and July.

The leaves and flowers of this plant have a strong, not ungrateful smell; and a very bitter nauseous taste. The flowers are somewhat bitterer, and considerably more aromatic than the leaves; and the yellow disk of the flower is, in both respects, far stronger than the white petala. The smell, as well as the taste, is rather improved than weakened by careful drying, and does not soon suffer any considerable diminution in keeping.

This plant, besides its general virtues as a bitter, has been supposed to have some degree of a carminative, anodyne, and antispasmodic power, depending on its odorous matter. It is recommended in colics of disferent kinds, particularly such as arise from flatulencies or cold; in hysterical and hypochondriacal disorders, and nephritic pains; in the pains of childbed women, and deficiencies of the uterine purgations; and intermitting severs, where a viscidity of the humours, or obstructions of the viscera, render the Peruvian bark ineffectual or prejudicial. In this last intention, the camomile is generally assisted by fixt alkaline salts, sal ammoniac, or other aperients, and often, also, by corroborating materials: Baglivi's preparation of camomile, which he looks upon as the

most certain specific in obstinate intermittents (a), was probably rather a composition of this kind, than any particular preparation of the camomile alone. The dose of the dry flowers, in substance, is from ten or twelve grains to half a dram or more; in decoction or infusion, two drams.

Camomile flowers give out their virtues both to water and rectified spirit; infusions made in the former are of a yellowish brown colour, in the latter of a bright gold yellow: when the flowers have been dried fo as to be pulverable, the infusions prove more grateful than when they are fresh or but moderately dried. Distilled with water, they impregnate the aqueous fluid pretty strongly with their flavour: if the quantity of camomile, fubmitted to the operation, is large, a little effential oil Ol. effent. feparates and rifes to the furface of the water, in colour yellow with a mem. Pb. cast of greenish or brown, of a pungent taste, and a strong smell exactly Lond. refembling that of the camomile. Decoctions of the flowers, inspiffated, though with a very gentle heat, to the confistence of honey, lose almost all the peculiar flavour of the plant, retaining its bitterness entire: the extract discovered to the taste a slight saline kind of austerity joined to the bitter; and on keeping for fome months, threw off to the furface a number of minute faline crystals.

flor. cha-

Rectified spirit, drawn off from the spirituous tincture, brings over likewife a part of the flavour of the camomile, but leaves a confiderable part behind in the extract. The fmell is in great measure covered or fuppressed by the spirit, in all the spirituous preparations; but the taste, both in the spirituous tincture and extract, is considerably stronger than in the watery.

The leaves and flowers are frequently employed externally, in difcutient and antiseptic fomentations, and in emollient and carminative glysters. They appear, from Dr. Pringle's experiments, to stand very high in the scale of antiseptics; the soluble part of the flowers resisting the putrefaction of animal flesh, with a power at least one hundred and twenty times greater than fea falt. Some endeavour to impregnate oil olive with the active matter of the camomile, for external uses, by gently boiling the fresh herb and flowers bruised, in thrice their quantity of the oil, till they become crifp; and then straining and pressing out the fluid. A preparation of this kind might be obtained to better purpose,

by a process similar to that, whereby expressed oils are perfumed with the fragrance of the more odoriferous flowers (see Ben); or by infusing the flowers in the oil without heat; for the strong heat, necessary for making the fluid boil, impresses a disagreeable taint, and dissipates greatest part of the volatile matter of the camomile.

2. CHAMÆMELUM FLORE PLENO: Chamæmelum nobile flore multiplici C. B. Double camomile: a variety of the foregoing, produced by culture; differing in the flowers being double, or having feveral rows of

the white petala, and the disk proportionably smaller.

The fingle and double flowered camomiles have been often used indiscriminately, and are allowed to be so used by the faculty of Paris. The leaves of the two plants are indeed alike, in quality as well as in their external form: but with regard to the flowers, as their active matter is almost wholly confined to the yellow disk, and as the single have large disks, but the double very small ones, and when very double, scarcely any at all; it is plain that the latter cannot be equivalent to the former unless taken in much greater quantity; and that therefore the single or large-disked flowers alone ought to be employed for medicinal uses.

3. CHAMÆMELUM VULGARE, Anthemis. Chamæmelum vulgare leucanthemum dioscoridis C. B. Matricaria Chamomilla Linn. Common

camomile; upright, annual, growing wild in corn fields.

This species also is allowed by the faculty of Paris to be used indifferently with the Roman camomile. Both its leaves and flowers are much weaker than those of the Roman, and their smell of a less agreeable kind: sometimes they have scarce any smell at all. They yield in distillation considerably less oil: from eight pounds of the flowers of the Roman were obtained about five drams, or a very little more; from the same quantity of those of the common, scarcely three drams. The oils of the two plants are in smell and taste nearly alike, but in colour remarkably different, that of the common being of a beautiful deep blue: if the oil is carefully kept, it retains its fine colour for many years; but if the air is admitted to it, the blue degenerates in a short time to a yellow, like that which the oil of the other fort has on its first distillation.

4. COTULA FOETIDA Chamæmelum fætidum C. B. Anthemis Cotula Linn. Mayweed or stinking camomile: annual, more upright than the other camomiles, with finer leaves, the flowers thicker together and their disks more convex and protuberant. It grows in waste grounds, and among corn.

This species differs greatly in quality from the three preceding. Its smell is disagreeable: the flowers have little or no taste; the leaves a strong one, of the acrid bitterish kind. It has never been much in use, nor are its medicinal effects well known. Decoctions of it are said to have been sometimes employed as a bath or somentation, against hysteric suffocations, and hæmorrhoidal pains and swellings. Mr. Ray says, that a decoction of the herb has by some been given internally, with success, in scrophulous cases. Brown Langrish gives an account of a decoction of it throwing a person affected with rheumatism into a prosuse sweat, and curing him.

CHAMÆPITYS.

CHAMEPITYS Pharm. Lond. Chamæpitys lutea vulgaris five folio trifido C. B. Abiga & ajuga quibusdam. Teucrium Chamæpitys Linn. GROUNDPINE: a low, hairy, creeping plant, with square stalks; whitish clammy leaves, cut deeply into three long narrow segments like those of the pine tree, set in pairs at the joints; and yellow labiated flowers, without pedicles, and wanting the upper lip. It is annual, grows wild in sandy and chalky grounds in some parts of England, and slowers in July.

The leaves of groundpine are moderately bitter, and of a refinous, not difagreeable smell; approaching in this respect, as in their external form, to those of the pine tree. They are recommended as aperients, and corroborants of the nervous system; and said to be particularly serviceable in female obstructions, paralytic disorders, and when continued for a length of time, either by themselves or with the assistance of germander, in rheumatic, ischiadic, and gouty pains.

The leaves in substance, dried and powdered, are directed to be given from half a dram to a dram. Their virtues are extracted both by water and spirit, most perfectly by the latter: the aqueous tinctures are yellowish,

li2

tl

the spirituous green. In distillation, they weakly impregnate water with their resinous scent: on distilling large quantities of the herb, a little essential oil may be collected, in quality somewhat approaching to that of turpentine. The watery extract has, joined to its bitterness, a weak saline austerity; the spirituous, a slight sweetishness and warmth.

CHEIRI.

CHEIRI, five Leucoium luteum Pharm. Edinb. (a) Leucoium luteum vulgare C. B. Viola lutea Gerard. Cheiranthus Chieri Linn. WALL FLOWER: a plant with woody brittle stalks and branches; smooth, dark green, oblong, narrow, sharp-pointed leaves; and numerous, tetrapetalous, yellow flowers, opening successively, on the tops, followed by long slender pods containing reddish slat seeds. It grows wild upon old walls and among rubbish, and slowers in April and May.

The flowers of cheiri have a moderately strong pleasant smell, and a nauseous, bitterish, somewhat pungent taste; which seem to afford some foundation for the nervine and deobstruent virtues commonly ascribed to them. They give out their active matter, together with a deep yellow tincture, both to water and spirit; and impregnate water, in distillation, with their odoriferous principle separated from the other parts; but no oil is obtained, at least when only moderate quantities, as a pound or two of the slowers, are submitted to the operation at once. The decoction, after the dissipation of the aromatic matter, discovers, besides the strong taste manifest in the flowers themselves, a sensibly saline one.

CHELIDONIUM MAJUS.

CHELIDONIUM MAJUS Pharm. Edinb. (b) Chelidonium majus vulgare C. B. & Linn. CELANDINE: a plant with longish leaves, divided to the rib into roundish indented portions, of which those at the extremities are much larger than the others, of a bright green colour on the upper side, blueish green underneath, full of a gold coloured juice, as are likewise the stalks: from the bosoms of the leaves issue long

pedicles, bearing clusters of tetrapetalous yellow flowers, which are followed by brownish pods containing flattish shining black seeds: the root is pretty thick at top, with a number of fibres at bottom, externally brownish, internally of a deep yellowish red or saffron colour. It is perennial, grows wild in hedges and shady waste places, and flowers in May and June.

THE leaves and roots of celandine have a faint unpleasant smell, and a bitterish, very acrid, and very durable taste, which is considerably stronger in the roots than in the leaves. Both water and rectified spirit extract nearly the whole of their pungent matter: the leaves, notwithstanding the yellow juice which issues so plentifully from a slight wound, and in which their activity appears to refide, give to rectified spirit a green tincture: the roots, which yield a copious faffron red juice, tinge the same menstruum of a brownish yellow. The pungency of this plant is not of the volatile kind, little or nothing of it rifing in distillation, with water, any more than with spirit: it is nevertheless greatly abated by drying the plant itself, or by inspissating, with a gentle heat, the spirituous or watery infusions. The smell of the herb is wholly diffipated in drying.

This acrid plant stands recommended as a powerful aperient and attenuant, in obstinate jaundices when not accompanied with inflammatory fymptoms, in cachexies, chlorofes, dropfies, and other difeafes. Half a dram or a dram of the dry root in powder; or an infusion, in wine or water, of a dram or a dram and a half of the fresh root; or three or four drops of its faffron-coloured juice, in any convenient vehicle; are directed for a dose. Great caution is requisite in the internal use of a medicine so acrimonious and irritating; more particularly in acute distempers, in which infusions of it, made in vinegar, have by some been recommended as a fudorific. Among us, it is employed chiefly by the common people for fome external purposes; as the destroying of warts, cleansing foul fores, removing fome cutaneous defedations, and clouds and beginning fuffusions of the eyes: for this last intention, the juice is diluted largely with milk, being of itself much too sharp to be applied with safety to so tender an organ.

CHELIDONIUM MINUS.

CHELIDONIUM MINUS Pharm. Edinb.(a) Chelidonia rotundifolia minor C. B. Ranunculus Ficaria Linn. PILEWORT: a small plant,
with roundish, smooth, shining green leaves, set on long pedicles; and
slender procumbent stalks; bearing bright gold-coloured solitary flowers,
of eight or nine petala, which stand in three-leaved cups, and are sollowed by clusters of naked seeds: the root consists of slender sibres,
with a number of tubercles or little knobs among them. It is perennial,
grows wild in hedges and moist meadows, and flowers in April.

The leaves of pilewort are ranked among the antiscorbutics, but do not promise to be of much virtue: they have no smell, and only an herbaceous taste, which, on chewing considerable quantities of them for some time, is followed by a very slight pungency. The roots are celebrated as a specific against the piles: they have been sometimes given inwardly, but chiefly applied externally, in the form of a cataplasm, lotion, or unguent. Boerhaave relates, that he cured an atrabiliary person who was troubled with the piles, by using daily a decoction of two ounces of this root, after sundry other medicines had been tried in vain. Perhaps the pilewort root acts, in these cases, little otherwise than as a simple emollient: it has a soft sweetish taste, and yields with water a large proportion of a mucilaginous extract.

CHINA.

CHINA RADIX Pharm. Edinb. (a) CHINA ROOT: an oblong thick jointed root, full of irregular knobs, of a reddish brown colour on the outside, and a pale reddish within. Two sorts are common in the shops, an oriental and occidental: the first, which is accounted the best, is considerably paler coloured and harder than the other. Of either kind, such should be chosen, as is fresh and heavy, and which, when cut, exhibits a close, smooth, glossy surface.

The plant is a climber, with tendrils like those of the vine, producing clusters of small flowers, which are followed by pretty large berries.

The oriental species (Smilax China Linn.) has roundish prickly stalks and red berries, and is a native of China and Japan: the occidental (Smilax Pseudo-China Linn.) has rounder smooth stalks and black berries, grows wild in Jamaica and Virginia, and bears the colds of our own climate.

THESE roots have scarce any smell or particular taste: when fresh, they are said to be somewhat acrid, but as brought to us they discover, even when long chewed, no other than a slight unctuosity in the mouth. Boiled in water, they impart a reddish colour, and a kind of vapid softness: the decoction, inspissated, yields an unctuous, farinaceous, almost insipid mass, amounting to upwards of half the weight of the root. They give a gold yellow tincture to rectified spirit, but make no sensible alteration in its taste: on drawing off the spirit from the filtered liquor, there remains an orange-coloured extract, nearly as insipid as that obtained by water, but scarcely in half its quantity.

China root is generally supposed to promote perspiration and urine, and by its soft unctuous quality, to obtund acrimonious humours. It was first introduced into Europe about the year 1535, with the character of a specific against venereal disorders: the patient was kept warm, a weak decoction of china root used for common drink, and a stronger decoction taken twice a day in bed to promote a sweat. Such a regimen is doubtless a good auxiliary to mercurial alteratives: but whatever may be its effects in the warmer climates, it is found in this to be, of itself, greatly insufficient. At present, the china root is very rarely made use of, having for some time given place to sarsaparilla, which is supposed to be more effectual. Prosper Alpinus informs us, that this root is in great esteem among the Egyptian women for procuring fatness and plumpness.

CICHOREUM.

CICHOREUM Pharm. Edinb. (a) Cichoreum silvestre sive officinarum C. B. Cichorium Intybus Linn. WILD CICHORY: a plant with oblong, dark green, somewhat hairy leaves, deeply jagged, like those of dandelion,

but larger; in the bosoms of which, towards the tops of the branches, the flowers come forth in spikes, consisting, each, of a number of blue flat flosculi, set in a scaly cup, which afterwards becomes a covering to several short angular seeds: the root is long and slender, of a brown colour on the outside, and white within. It is biennial, grows in hedges and by road sides, and flowers in June and July.

WILD cichory abounds with a milky juice, of a penetrating bitterish taste, and of no remarkable smell or particular slavour: the roots are bitterer than the leaves or stalks, and these much more so than the flowers. By culture in gardens it loses its green colour, and in great measure its bitterness, and in this state is a common sallad herb. The darker coloured, and the more deeply jagged the leaves are, the bitterer is the taste of all the parts of the plant.

The roots and leaves of wild cichory are very useful aperients, acting mildly and without irritation, tending rather to abate than to increase heat, and which may therefore be given with safety in hectic and inflammatory cases. Taken freely, they keep the belly open, or procure a gentle diarrhæa; and when thus continued for some time, they have often proved salutary in beginning obstructions of the viscera, in jaundices, cachexies, hypochondriacal and other chronical disorders. Geoffroy relates, that he has known inveterate and stubborn intermittents cured by the daily use of wild cichory leaves, after many febrifuges had been tried in vain.

The virtues of the cichory refide in the milky juice; and may be extracted by expression, or by infusing or boiling the herb or root in water or in spirit. The plant seems to lose nothing of its taste in drying; or the juice or infusions in being gently inspissated to the consistence of an extract; though the plant in its recent state, or the liquors uninspissated, are supposed to be of most efficacy. The watery extract is somewhat larger in quantity than the spirituous, and this last is proportionably strongest in taste.

CICUTA.

CICUTA Pharm. Edinb. Cicuta major C. B. Conium maculatum Linn. Hemlock: a tall umbelliferous plant, with large leaves, of a blackish

blackish green colour on the upper side, and a whitish green underneath, divided into a number of small oblong somewhat oval segments, which stand in pairs on middle ribs: these segments are again deeply cut, but not quite divided, on both sides; and many of these ultimate sections have one or two slighter indentations. The stalk is round, smooth, hollow, irregularly variegated with spots and streaks of a red or blackish purple colour. The slowers are white; the seeds greenish, slat on one side, very convex and marked with five surrows on the other. The root is oblong, about the size of a middling parsnep, yellowish without, white and sungous within. The plant is annual or biennial; common about the sides of fields, under hedges, and in moist shady grounds; and slowers in June and July.

THE leaves of hemlock have a rank smell: the organs of taste, they affect but little. On expression, they give out their smell to the juice; which, on being directly inspissated with a gentle heat, to the consistence of an extract, retains great part of the scent, and discovers an unpleasant subacrid taste seemingly of the subsaline kind. If the juice be suffered to settle till it becomes clear, it loses nearly all the specific slavour of the hemlock, the odorous principle seeming to separate and subside along with the herbaceous seculencies: the proper menstruum of this matter is rectified spirit of wine; which completely extracts the smell, both of the leaves in substance and of the inspissated juice, and receives, from both, a green tincture. The saturated tincture, mixed with water, grows turbid, and deposites a green resin.

This herb is recommended externally, in cataplasms, somentations, and plasters, as a powerful resolvent and discutient. Taken internally, in no great quantity, it has occasioned disorders of the senses, sleep, convulsions, and in some instances death; and hence it is ranked among the poisonous plants: Boerhaave tells us, that by the effluvia of the herb bruised and strongly smelt to, he became vertiginous (a). It is said, that to certain brute animals, it is innocent (b); and that its ill qualities

Kk

⁽a) Historia plantar. Lugd. Bat. p. 94. Haller, Stirpes belveticæ, p. 434.

⁽b) Quippe videre licet, pinguescere sæpe cicuta

Barbigeras pecudes, homini quæ est acre venenum. Lucretius.

are corrected by vinegar or other vegetable acids (a). Of its effects in small doses, insufficient to do harm, in which it has been by some recommended, nothing material was known, till the happy experiments of Dr. Stærck, lately published (b), gave room to hope, not only that the virtues ascribed to it in external applications are better founded than practitioners in general seem now to suppose, but likewise that it is a

plant of very great importance as an internal medicine.

Dr. Stærck relates, that bags of the dry leaves, quilted together, boiled for a few minutes in water, (or in milk, where they could not otherwife be borne, on account of their fmell and the itching they produced) then fqueezed from the fuperfluous liquid, and applied warm, checked the progress of very bad gangrenes, and procured a separation of the corrupted parts: that the same application, in a person of fixty who had been gouty for feveral years, immediately abated the pains, foftened and discussed the tophaceous concretions, and occasioned the next fit to be milder and of shorter continuance: that its effects were likewise considerable in ædematous tumours, scirrhous strumæ, indurations of the glands of the breast, and in very bad cancers: that nevertheless some received from it no benefit, though no one harm: that in inflammations or hot ferous tumours, it was less proper than in the above cases, or had place only after evacuations: and that plasters, containing the juice of the hemlock, often refolve and discuss what resists all other applications. It is in the form of plaster that this herb, among us, has been chiefly made use of: the. recent juice is mixed with a folution of twice its quantity of gum ammoniacum made in vinegar of fquills, and the mixture boiled down to a due confistence.

(a) Cicuta, præsens illud venenum, si coquitur in aceto, sine noxa comedi potest, quod probavi aliquoties, experimenti ergo, Lugduni Batavorum, ubi in sossis extra urbem frequens crescit, Lindestolpe, de venenis, edit. Stentzel. p. 431.—Cicutæ caules, aceto macerati, impune comeduntur,

& ipfe edi. Id. p. 781.

A large spoonful of hemlock juice given to a cat, had no sensible effect: a second produced a visible embarras on the region of the reins: in a little time the animal staggered, but did not fall: on swallowing a third spoonful, she ran away, and was presently out of sight. A quarter of an hour after, she was found, stretched out, motionless, her paws rigid. Half a dram of theriaca, diluted with two large spoonfuls of wine, had no good effect. A large spoonful of fresh lemon juice was scarce swallowed, when the animal got up on her legs, appeared free from pain, as if nothing had happened—continues in persect health. Mr. Haram, apothecary at Chartres. Rozier Tableau, tom. i. 1773.

⁽b) Libellus quo demonstratur cicutam, &c. Vindobonæ, 1760.

For internal purposes, he directs the juice, while fresh, without fuffering it to fettle, to be inspissated in an earthen vessel over a very gentle fire, and kept continually stirring to prevent its burning, till it acquires the confiftence of a thick extract; which is to be mixed with fo much of the powdered leaves, as will reduce it into a mass fit for being formed into pills. This preparation, he fays, was given to a little dog, in the quantity of a scruple; taken by himself, in doses of one and two grains, every morning and evening, for feveral days; continued by persons in health, for a year or two; increased, in some cases, to a dram and a half in a day; without producing any ill confequence, or affecting any of the actions, fecretions, or excretions of the body. It nevertheless had very powerful and falutary effects in some reputed incurable diseases; acting, though flowly and infenfibly, as a high refolvent: he relates histories of inveterate scirrhuses, cancers, and the worst kinds of ulcers and fiftulæ, being completely cured by it; and fays it refolves also recent cataracts, or at least restrains their progress. He begins always with fmall doses; giving one pill, of two grains, first twice a day, then thrice a day, and gradually increasing the number to fix or more for a dose. The good effects of the medicine were sometimes visible in a few days; though the cure generally required feveral months.

The trials of this medicine, made among us, have not, as yet, so far as I can learn, been accompanied with so much success. I have been informed of some cases, in which it was apparently of great benefit; of some, in which it did no service; and, of some, in which it affected the speech and hearing. It has been suspected that the general inefficacy of the medicine in this country was owing either to our hemlock being different in quality from the German, or to the extract being less skilfully prepared; but some of the extract has been procured from Dr. Stærck himself, and sound to succeed no better than our own: though the medicine has seldom been accompanied with very happy effects, it had

activity enough to be productive of alarming symptoms.

In the third volume of the medical observations and inquiries published by a society of physicians in London, Dr. Fothergill and Dr. Rutty have given two papers on this medicine, drawn from extensive practice in England and Ireland; from which we have grounds to persuade ourselves, that though the cicuta is far from answering the expectations which Dr. Stærck had raised, it may nevertheless be an useful acquisition, and

Kk2

may affift in curing some disorders, and alleviating others, in which the common medicines are inadequate auxiliaries. In real cancers, whether ulcerated or occult, there is no instance of its effecting a complete cure; but it was found to retard the progress of the deplorable disease, to mitigate the pain for a time, and to change the thin, ichorous, fetid discharge to a state more approaching that of laudable pus. In different kinds of malignant ulcers, it in like manner mended the discharge, and disposed the ulcer to heal. Some scrophulous tumours were completely refolved by it, and the cure has stood for several seasons: in other cases of this kind, the patient has frequently fuffered a relapfe, especially in the fpring. Dr. Fothergill observes, that the success of this medicine depends on its being given in as large a dose as the patient can bear; for otherwise, though continued for a length of time, it seldom procures any benefit: that the hemlock is in greatest perfection when the flowers begin to fade, and the habit of the plant inclines to yellow; and that in making the extract, the less heat it undergoes, the better: that he has found twenty grains of one fort of extract equal in point of efficacy to near forty of another: that the dose is to be increased by degrees, till it produces certain effects, which feldom fail to arife from a dull dofe, and which for the most part are, either a giddiness affecting the head, and motions of the eyes, as if fomething pushed them outwards; or a flight fickness and trembling agitation of the body; or a laxative stool or two: that here we must stop till none of these effects are felt, and in three or four days advance a few grains more: that a greater quantity can commonly be borne at night than at noon, and at noon than in the morning: that the method he commonly follows is, to order two drams of the extract to be made into thirty pills, of which two are to be taken in a morning, two at noon, and three or four at night, and one pill added to each dose, according as the patient can bear it: that the extract, given in this manner, is apparently anodyne; promotes rest, and eases pain; feldom creates thirst, or that kind of morning headach which fucceeds an opiate; rarely occasions costiveness, but in most procures a laxative stool the day following: that in some habits, very small doses offend the stomach, excite spasmodic twitchings, heat, and thirst; and that in fuch cases, its use is immediately forbid.

* In a paper of Dr. Fothergill's in Vol. 5th of the London Med. Observ. and Inq. the efficacy of extract of hemlock in a particular painful affection

affection of the face, is related. This difease is a sudden violent pain attacking some part of the face, continuing a very short time, and returning at irregular intervals, the nature and cause of which is not distinctly known. The Doctor was led, from some circumstances, to consider it as owing to a cancerous acrimony. In most of the cases he met with, the cicuta, taken in sufficient quantity, and long enough persisted in, removed the complaint.

*This remedy has likewise been strongly recommended in the cure of the chincough, by Dr. Butter, in a treatise on that disease. He represents it as no less efficacious in this complaint, than bark in an ague; but the trials made by other practitioners do not seem to have confirmed

this opinion.

* The following directions for preparing in the most perfect manner the extract of hemlock, are given by Dr. Withering in his Botanical Arrangement of British Vegetables. "Let several people be employed to gather the plant; and as fast as it is cut, let others carry it in handbaskets to the press; but it must lie light and loosely in the baskets. Let the juice be immediately squeezed out; and as fast as it runs from the press, it must be put over the fire, and boiled till three parts out of four of the whole liquor is wasted. Then it must be put into a water bath, and evaporated to the consistence of honey. If it is now taken and spread thin upon a board or marble slab, and exposed to the sun and the air, it will soon be of a proper consistence to be formed into pills. From five to ten grains of this extract is a proper dose; sew constitutions will bear more without experiencing disagreeable effects (a)."

The root of hemlock is generally supposed to be, both in external applications and when taken internally, of more activity than the leaves. Stærck relates, that on being cut, it yields a bitter acrid milk, of which a drop or two, applied to the tip of the tongue, occasioned a rigidity, pain, and swelling of the part, so as to prevent speech; and that he was freed from this complaint, by washing and rubbing the tongue with citron juice. In drying, it seems to lose of its virulence: he says he has taken a grain or two of the powder without injury: there are instances of twenty and thirty grains being given, with advantage, in scirrhuses of the liver, $\mathcal{C}c.(b)$, in quartan agues on the approach of a fit, and even

⁽a) Vol. i. p. 163.

⁽b) Renealm, Observat, iii. & iv. Etmuller, Schrader dilucidat. par. i. sett. ii. p. 111.

in acute fevers (a). Nor does the fresh root appear to be at all times of equal virulence: I have seen it chewed freely, without any other effect being perceived, than an impression of sweetishness resembling that of parsley roots or carrots: there are instances of some drams, and even ounces, having been taken, without producing any ill consequence (b). So variable does this plant appear to be in its qualities; if really the subject of the several histories was precisely the same species of plant.

The feeds have been recommended by fome as demulcent, paregoric, and antaphrodifiac. Of their real qualities, little more is known with certainty, than that they are innocent to fome kinds of birds: Mr. Ray fays, he found the crop of a thrush full of hemlock feeds, even at the

feafon when corn was plentiful.

* In the Medical Commentaries (c), it is afferted, that the extract prepared from the feeds of hemlock has been observed to be much more powerful than from the leaves. And in the last Edinburgh pharmacopæia, this is the only extract directed to be kept as an officinal.

CIMOLIA.

TWO forts of argillaceous earth are directed under this name in catalogues of the materia medica: CIMOLIA ALBA feu Argilla alba Pharm. Edinb.(d) the pure white strong clay, called, from the use to which it is principally applied, Tobacco-pipe clay: and CIMOLIA PURPURASCENS Pharm. Edinb.(d) a compact bolar earth, commonly of a greyish brown colour, called, from its use, Fuller's earth.

THESE earths appear to be nearly of the same medical qualities with the boles formerly treated of; but are rarely or never employed, at least under their own names, for any medicinal purposes. If the virtue of these kinds of substances depends, as it most probably does, on their soft viscous quality, (for they do not appear to have any other, by which they can act in the bodies of animals) the white cimolia is obviously the most effectual of them all.

⁽a) Bowle, apud Raium, bist. plant. i. 451.

⁽b) Petiver, Philosoph. transact. Abr. Lowthorp. ii. 641. Henley, ibid. Jungius, Eph. nat. curios. dec. i. ann. iv. obs. 106. Trew, Commerc. lit. Norimberg. ann. 1740. hebd. 47.

⁽c) Vol. i. p. 326. (d) Expunged.

CINARA.

CINARA; Scolymus. Cinara bortensis Pharm. Edinb. Cinara bortensis foliis non aculeatis C. B. Cynara Scolymus Linn. ARTICHOKE: a rough plant, with large greyish leaves, divided almost to the rib into irregularly indented segments: among these arises a thick stalk, bearing a large scaly head, which, opening at top, sends out a number of purplish blue slosculi, followed by whitish seeds winged with down. It is perennial, a native of the southern parts of Europe, and cultivated in our culinary gardens.

The bottoms of the heads, and the fleshy part of the scales, are supposed to be of easy digestion; but gross, flatulent, and of little nourishment. The leaves are bitter, and give out their bitterness, along with their juice, on being bruised and pressed. The expressed juice is given in dropsies, and in some instances has proved successful after the medicines more commonly made use of in that disorder had failed: for this purpose, the juice, not depurated, or freed only from its grosser feculency by passing it through a coarse strainer, is mixed with equal its quantity of white wine, and three or four spoonfuls or more of the mixture taken every morning and evening. Its operation is chiefly by urine.

CINNABARIS.

CINNABARIS NATIVA. Minium Græcorum. NATIVE CINNABAR: a ponderous, red, fulphureous ore of mercury; found in Spain,
Hungary, and feveral other parts of the world. The finest is imported
from the East Indies: partly in pretty large irregular masses; partly in
smaller roundish ones, smooth without, and striated within; both externally and internally of an elegant deep red colour, which is greatly
improved by grinding the mass into fine powder.

Cinnabar confifts of quickfilver and common brimstone; in the proportion of not less than four (a), commonly six or seven (b) parts of the

⁽a) Lemery, Cours de chimie, part. i. chap. viii. operat. 2.

⁽b) Cramer, Elem. artis docimasticæ, tom. i. edit. ii. p. 287. Malouin, Chimie medicinale, part. iv. chap. 38.

mercury, to one of the fulphur: the finer its colour, the more mercury and the less sulphur it is found to hold. The native cinnabar generally contains also a quantity of earthy matter, from which it may be purified by fublimation. If this earth should be of the calcareous kind, or if calcareous earths, iron filings, or other fubstances that absorb fulphur more strongly than mercury does, be added; more or less of the sulphur, proportionably to the quantity of such absorbent addition, will be detained at the bottom of the fubliming veffel: on this principle, the coarfer cinnabars may be freed from their redundant fulphur as well as from their earthy matter, and thus rendered of a high colour: or the whole of the fulphur may be detained, by an increase of the absorbent material, and the pure mercury distilled off in its running form: one part of lime or iron filings is generally fufficient for extricating all the mercury from four parts of cinnabar. The humid menstrua, that diffolve either one or the other of the ingredients of cinnabar by themselves, have little effect upon the compound; the mercury being protected by the fulphur from the action of acids, and the fulphur by the mercury from that of alkaline liquors: alkalies indeed, even in the dry way of fublimation, do not so perfectly detain the fulphur as bodies of the earthy or metallic kind.

Native cinnabar has by many been preferred, as a medicine, to that which is made by art, but apparently on no good foundation. The only difference between them confifts in this; that the native is subject to an admixture of heterogeneous matters, which are not perhaps always innocent (a); and that the proportions of its constituent ingredients are more precarious than in the factitious. The native cinnabar is therefore deservedly rejected by the London and Edinburgh colleges.

CINNAMOMUM.

CINNAMOMUM Pharm. Lond. & Edinb. Cinnamomum five canella zeilanica C. B. Cassia cinamomea Hermann. hort. Lugd. Bat. Cin-NAMON: the bark of a tree of the bay kind, (Laurus Cinnamomum Linn.)

⁽a) Accidit nonnunquam quod . . . nauseas & vomitiones excitet, necnon etiam anxietates circa præcordia: quod & ipse bis terve observavi, licet cinnabaris pluribus lotionibus purgata fuisset. Geoffroy, mat. med. i. 246.

growing in the island Ceylon; freed from the outer green or greyish part, and cut into long flices, which curl up, in drying, into quills or canes, the form in which it is brought to us; very thin, light, of a reddish yellow or pale rusty iron colour, somewhat tough in breaking, and of a fibrous texture. It is frequently mixed with another bark, greatly resembling it in appearance, but much weaker in virtue, casia lignea: this last is distinguished by the close smooth surface which it exhibits on being broken, and by its remarkably flimy tafte.

This bark is one of the most grateful of the aromatics; of a very fragrant smell, and a moderately pungent, glowing, but not fiery taste, accompanied with confiderable fweetness, and some degree of astringency. It is faid, that the fine flavour refides, originally, only in the thin pellicle, which lines the interiour furface of the bark, and which abounds with vehicles of effential oil; the rest of the bark, while fresh, being merely fubaftringent, and receiving the flavour, which we find it to have, from the inner pellicle in drying (a). Accordingly the thinnest pieces are found to be strongest; as containing the largest proportion of this active part, and the least of the inert woody matter.

Cinnamon, infused in boiling water in a close vessel, gives out to the fluid greatest part of its virtue: together with a reddish brown tincture, deeper or paler, according to the proportion of cinnamon employed. Rectified and proof spirit extract its virtues more perfectly than water, and without the affistance of heat; three ounces of the powdered bark, Tindura by cold maceration for a few days, give a strong impregnation to a quart+

or two pints and a half t of proof spirit.

The aromatic principle of this spice is an essential oil; which, in distillation with water, rises slowly and difficultly, and renders the liquor fomewhat milky: the water continues to run milky, and gratefully impregnated with the fragrance of the cinnamon, till about a gallon Aq.cinnam. has been drawn off from a pound: when large quantities of the spice fimp. Ph. Lond. Edin. are submitted to the operation at once, a small portion of oil commonly feparates and finks to the bottom of the water; in colour gold yellow; OI. effent. of a delightful smell-like that of the cinnamon itself; and of a fiery cinnamom. pungency, fo as not to be fafely tafted or applied to the skin without & Edinb.

cinnamomi + Pb. Lond. 1 Ph. Edinb.

dilution; for, as Boerhaave observes, it burns the part to a gangrenous eschar: in doses of a drop or two, diluted by the means of sugar, mucilages, \mathfrak{Sc} . it is one of the most immediate cordials and restoratives, in languors, singultuses, and all debilities. If the milky distilled water be long kept, great part of the ponderous oil, suspended in it, separates and subsides: some, with a view to the perfection of the water, endeavour to prevent this separation, by adding a small proportion of sugar, which contributes to keep the oil dissolved: others, with a view only to the obtaining of the oil, endeavour to promote the separation, by setting the liquor in a very cold place, and perhaps by other means not commonly known. It is said, that from sixteen ounces of good cinnamon, a dram and a half or two drams of oil have been collected.

The watery decoction, remaining after the distillation, yields, on being inspissated, a mildly astringent mass, which has nothing of the sweetness, any more than of the peculiar slavour of the cinnamon. It is observable, that this extract is free from the nauseous relish which most of the other spices discover, in a greater or less degree, when divested by the same means of their proper aromatic matter.

Rectified spirit, distilled from cinnamon, brings over very little of its slavour. An extract, made by this menstruum, retains nearly all the valuable parts of the spice, the sweet aromatic matter as well as the restringent: it has a durable and very grateful warmth and pungency, not a siery heat like the spirituous extracts of many other spices; the heat and pungency of cinnamon residing in the pure essential oil. The quantity of this extract is about one sixth of that of the cinnamon employed.

On distilling proof spirit from this spice, the purely spirituous part, which comes over first, proves almost slavourless, but the watery part which follows brings with it the essential oil; and this oil being disfolved by means of the spirituous portion, the liquor proves limpid. A cordial water of this kind is commonly prepared in the shops, by drawing off a gallon or nine parts of proof spirit from a pound of cinnamon. A like preparation might be obtained rather more advantage-ously, and free from the foreign slavour which the common proof spirits are accompanied with, by adding to the simple water a suitable quantity of pure rectified spirit.

Aq. cinnam. fpiritu. + Pb. Lond. 1 Pb. Edinb.

Some other products of the cinnamon tree are used medicinally in the eastern countries, and have been sometimes, though very rarely, brought into Europe; to wit, an aromatic effential oil distilled from the roots, Ol.caphuræ. and a species of camphor which separates from this oil on redistilling it: an oil drawn from the leaves, fimilar in flavour to the genuine oil of cloves with a little admixture of that of the cinnamon bark; and a whitish sebaceous matter, said to resemble the expressed oil of nutmegs, obtained either by expression or by coction in water from the fruit (a).

Caphurabaros Indorum.

Oleum malabathri. Cera cinnamomi.

CITREA.

CITREA MALUS Pharm. Edinb.(b) Malus medica C. B. Citrus medica Linn. CITRON: a fmall evergreen tree, refembling the lemon, and differing from it chiefly in the fruit; which is much larger, less juicy, and contains, under the yellow rind, a thicker fungous white bark. It is a native of Asia, and cultivated in the southern parts of Europe.

This fruit has a near affinity with lemons in its medicinal qualities as well as in its external form: the principal difference lies, in the juice of the citron being fomewhat less acid, and the yellow rind being fomewhat hotter and accompanied with a confiderable bitterishness. The rind gives out its virtue, equally with that of lemons, both to watery and spirituous menstrua; but its flavour is not equally retained in the spirituous extract, the effential oil of citron peel being very light and volatile, fo as in great part to rife in distillation along with the highest rectified spirit. Several varieties of these trees are common in the warmer climates. Oils obtained from the fresh peels of the more odoriferous kinds, by rolling the fruit on a plane stuck full of points, are brought from Italy, and used as perfumes: these are somewhat more Essentia de grateful, and confiderably less pungent, than such as are drawn by cedro Gerdistillation with water. The oil prepared in either of these ways is very Ol. cedrifubject to lose of its fine flavour, and become thick and refinous in keeping: when distilled with rectified spirit, and afterwards separated from the spirit by dilution with a large portion of water, it retains much longer its odour, fluidity, and limpidness.

(a) Albertus Seba, Ada academiæ cæsareæ, ubi supra, p. 11, 12.

(b) Expunged.

COCCINELLA.

COCCINELLA Pharm. Lond. Cochinilla Pharm. Edinb. Coccus Casti Linn. Cochineal: little wrinkled grains, of an irregular figure, convex on one fide and flat or somewhat hollowed on the other, externally of a dark red colour generally sprinkled with a whitish clammy powder, internally of a deep bright red. This substance, brought from Mexico and New Spain, supposed formerly the feed of a plant, appears to be an insect of the scarabæus kind, found adhering to the leaves and branches of the opuntia or American prickly pear tree, and carefully preserved and cured by the natives. The male insects have wings, and are about the size of a slea: the semales have no wings, and are larger; when full of young, they become torpid, and swell so as on first sight to resemble berries, in which state they are swept off with a pencil: if lest till the young ones creep out, the parent dies, and its body becomes an empty husk.

The principal use of cochineal is as a colouring drug: it gives a fine deep durable red both to rectified and proof spirit, and a deep purplish crimson to water: neither the watery or spirituous insusions suffer any change of their colour on being inspissated to the consistence of an extract. Cochineal has been sometimes used also in a directly medicinal view, and supposed to act as a mild corroborant and diaphoretic. It has a faint musty kind of smell, and a very slight bitterish roughish taste; both which are taken up, along with the colouring matter, by watery and by spirituous menstrua, and, though scarcely perceptible in the dilute tinctures or insusions, are very sensible in the inspissated extracts, particularly in that made with spirit. Cartheuser observes, that the mucilaginous bitterish watery extract amounts to three fourths of the weight of the cochineal; and the balsamic bitter and moderately astringent spirituous extract, to nearly as much.

COCHLEARIA.

SCURVYGRASS: a low plant; with thick juicy leaves, fomewhat hollowed fo as to refemble a fpoon, those from the root standing on long pedicles, those on the stalk joined close to it without pedicles; producing,

ducing, towards the upper part of the stalks, small white tetrapetalous slowers, followed by roundish seed-vessels. It is annual; grows wild in several parts of England, particularly about the sea coasts and salt marshes; and slowers in May or sooner.

- folio subrotundo C. B. Cochlearia officinalis Linn. Garden or Dutch scurvygrass: with the radical leaves unevenly roundish, and those on the stalks oblong. It is commonly cultivated, for the use of the shops, in gardens; and does not appear, like many other maritime plants, to change its qualities with the soil.
- 2. COCHLEARIA MARINA feu britannica. Cochlearia folio finuato C.B. Cochlearia anglica Linn. English or sea scurvygrass: with all the leaves alike, oblong, pointed, deeply and irregularly indented or sinuated.

THE fresh leaves of these plants have an ungrateful kind of smell, and a penetrating acrid taste: the first fort is considerably the strongest, and hence has long superfeded the use of the other. The slowers and seeds also are pungent, but less so than the leaves.

Scurvygrass is a powerful antiseptic, attenuant, and aperient: it manifestly promotes the sensible excretions, particularly urine, without heating or irritating fo much as might be expected from its great pungency. It is one of the capital antifcorbutic herbs, and in this intention has been principally made use of, in conjunction, generally, with mild vegetable acids, or substances of less acrimony, as orange juice, forrel, becabunga, &c. It is of fervice also in paralytic and cachectic indispositions; and in the wandering rheumatic pains, of long continuance, unaccompanied with a fever, called by Sydenham the scorbutic rheumatism. I have had frequent experience of the efficacy of the composition which he prefcribes in this commonly obstinate distemper, and which, he says, if the public had not outweighed private advantage, he should have concealed; to wit, fixteen parts of fresh made conserve of garden scurvygrass, eight of conferve of wood forrel, and fix of the compound powder of arum root, made up with fyrup of orange peel into an electary, of which two drams are to be taken thrice a day for a month, along with fome ounces of a distilled water impregnated with scurvygrass, mint, nutmegs,

&c. Among different aromatic materials made trial of for covering the ill flavour of scurvygrass, nutmegs seemed to answer the best. Instead of the compound powder of arum in the above composition, an equal quantity or more of fresh arum root, mixed with as much powdered gum-arabic, may be advantageously substituted; and probably the virtues of the medicine depend as much upon this root as on the scurvygrass.

The active matter of this plant is extracted by maceration both in watery and in spirituous menstrua, and accompanies the juice obtained by expression. The most considerable part of it is of a very volatile kind; the peculiar penetrating pungency totally exhaling, in the exficcation of the herb, and in the evaporation of the liquors; and only a flight biting bitterishness remaining in the dried leaves, in the inspissated juice, and in the spirituous as well as in the watery extracts. The fresh leaves, beaten into a conferve with thrice their weight of fine fugar, may be kept in a close vessel, without much diminution of their virtue, for years. juice also, purified from its feculencies by fettling and straining, may be preferved for a confiderable time, though by no means equally with the conferve, by fetting it in a cool place, and covering the furface with a little oil to prevent the access of air. The orange or forrel juice, commonly added to that of the scurvygrass, seem to promote the depuration; for if the juices, feparately, are made moderately fine, they foon deposite, on being mixed together, a fresh feculence.

Conferv.fol. cochl. hort. Pb. Lond.

The principal virtue of this plant refides in an effential oil; feparable, in very small quantity, by distillation with water. The oil is so ponderous, as to sink in the aqueous sluid; but of great volatility, subtility, and penetration. One drop, dissolved in spirit, or received on sugar, communicates to a quart of wine or other liquors the smell and taste of scurvygrass. It rises in distillation with rectified spirit as well as with water: a pint of rectified spirit, drawn off, in the heat of a water bath, from two pounds of the fresh herb bruised, brings over nearly all the oil, and proves exceeding strongly impregnated with the volatile pungency of the scurvygrass. Both the oil and the spirit, particularly the former, require the bottles they are kept in to be very carefully secured; the subtile matter of the plant, when thus disengaged by distillation from the grosser parts, being extremely disposed to escape.

COFFEA.

what smaller than a common bean, convex on one side, flat on the other with a remarkable surrow. It is the produce of a tree of the jasmine kind, (jasminum arabicum castaneæ solio, slore albo odoratissimo, Commelin. bort. amst. Cossea arabica Linn.) growing in Arabia, and thence introduced into the West Indies: the fruit is a juicy berry, including two of the seeds, joined by the flat sides, and covered each with a thin shell.

Coffee feeds have a farinaceous, somewhat unctuous, bitterish taste, and little or no smell: the flavour, for which they are valued, is procured by gentle roasting; and some of our own seeds and kernels acquire, by that process, a flavour somewhat of the same kind. The roasted seeds, ground into powder, soon lose their flavour in the air, impart it to water and spirit by slight coction or digestion, and give over great part of it with water in distillation. An extract made from them by water is, not disagreeably, bitterish: an extract made by rectified spirit is stronger, and not a little nauseous.

The dietetic use of coffee is said to strengthen the stomach and promote the secretions; to be serviceable in phlegmatic corpulent habits; to be injurious in thin habits and bilious temperaments, in melancholic and hypochondriacal disorders, and to persons subject to hemorrhages.

* From some experiments related by Dr. Percival in Vol. II. of his Essays Medical and Experimental, it appears that coffee is slightly astringent and antiseptic; that it restrains fermentation; and has a powerful sedative action on the nervous system. Sir John Pringle, in a letter to the same author, recommends strong coffee as the best abator of the paroxysms of the periodic asthma with which he is acquainted. He directs for this purpose an ounce of best Mocha coffee to be made into a single dish, to be repeated fresh after the interval of a quarter or half an hour, and drunk without milk or sugar.

(a) Expunged.

COLCHICUM.

COLCHICUM: Colchicum autumnale Stærck. & Linn. Colchicum commune C. B. Colchicum anglicum purpureum & album Ger. & Park. MEADOW SAFFRON: a plant with a fleshy bulbous root, producing from its lower part a smaller bulb: from this last arises, in autumn, along a furrow in the fide of the old root, a flender hollow transparent pedicle, widening at top into a flower like those of the crocuses, divided into fix fegments, of a purplish or whitish colour, withering in two or three days: from the fame root, next fpring, come forth three or four upright leaves, like those of the lily; in the middle of which appear, on fhort pedicles, commonly three triangular pods, about the fize of small walnuts, divided into three cells full of roundish dark-coloured seeds. It grows wild in rich moift meadow grounds in the fouthern and western parts of England. The roots, freed from the outer blackish coat and the fibres at bottom, are while fresh of a white colour, and full of a milky juice; in drying they become wrinkled, and of a blackish or dark reddish brown.

This is one of those plants, whose violent and singular effects in the bodies of animals engaged the attention of Dr. Stærck; in hopes that by giving it in very small doses, or by due preparation, it might be converted into a medicine not only fafe, but capable of relieving diforders in which the common remedies prove ineffectual. He observes, that on cutting the fresh root into slices, the acrid particles emitted from it irritate the nostrils, fauces, and breast, and that the ends of the fingers with which it had been held become for a time benumbed: that applied for two minutes to the tip of the tongue, it rendered the part rigid and almost void of sensation for fix hours: that less than a grain, wrapt up in crumb of bread and taken internally, produced alarming fymptoms, a burning heat and pain in the stomach and bowels, strangury, tenesmus, thirst, total loss of appetite, &c. which were greatly relieved by an acidulous mixture with fyrup of poppies, and which on the fourth day went entirely off: that an infusion of three grains of the root in four ounces of wine, flowly fwallowed, occasioned a tickling in the larynx and short dry cough, foon after a heat in the urinary paffages and a copious discharge of pale urine, without sensibly affecting the other functions of

the body: that an ounce of the fliced juicy root being digested with a gentle heat in a pound of vinegar for forty-eight hours, and the bottle frequently shaken; the root became almost insipid, and the strained vinegar proved acrid in taste, irritated and constringed the fauces, and raised a short cough: that this vinegar, mixed with twice its quantity of honey, and gently boiled down to the confistence of honey (frequently Oxymel stirring the mixture with a wooden spoon) proved a sufficiently grateful oxymel, which taken in doses of a tea-spoonful, that is, a dram, promoted a copious discharge of urine, without inconvenience. After these experiments on himself, he made trial of this oxymel, in the hospital at Vienna, in desperate hydropic and other serous disorders, in which it was always found to act without disturbance, as a most potent diuretic, after the common medicines employed in that intention had failed. He begins with giving a dram twice a day in any fuitable vehicle, and gradually increases the dose to an ounce and sometimes an ounce and a half in a day: if this last quantity should prove ineffectual, he thinks there are little hopes of any benefit from this medicine. *The Edinburgh college have now received into their pharmacopæia a fyrup of colchicum, made with Syr. colchici the fame infusion of the root in vinegar as above described, in which are disfolved twenty-fix ounces of fine fugar.

COLOCYNTHIS.

COLOCYNTHIS Ph. Edinb. Colocynthidis medulla Ph. Lond. COLOQUINTIDA: the dried medullary or pulpy part of a species of gourd or cucumber, (Cucumis Colocynthis Linn.) brought from Aleppo. It is very light, white, of a fungous texture, composed as it were of membranous leaves, with a number of roundish feeds lodged in the cavities.

THE fungous medulla, freed from the feeds, (which are fomewhat unctuous and fweetish, like those of the common cucumber) has a naufeous, acrid, intenfely bitter tafte. It is a very strong irritating cathartic; commended by fome, not only as an efficacious purgative, but likewife as an alterative in obstinate chronical disorders; by others condemned, as a dangerous and deleterious drug. Thus much is certain, that when given by itself, in substance, in such doses as to purge effectually, as Mm eight

eight or twelve grains, it operates for the most part with great vehemence; disordering the constitution, occasioning violent gripes and sometimes bloody discharges. Its principal use is as a stimulus to other purgatives.

Colocynth, boiled in water, renders a large quantity of the liquor ropy and flimy: even a tincture of it made in proof spirit is so glutinous, as not to pass through a filter, and not easily through a common strainer. The watery decoctions inspissated, yield a large proportion, half the weight of the colocynth or more, of a mucilaginous extract; which purges strongly, but with much less irritation, and greater safety than the colocynth itself, and appears to be the best preparation obtainable from this drastic drug. The college of Edinburgh uses a soft extract of this kind, made from coloquintida and black hellebore, for mixing into pills with scammony and aloes (a); that of London directs a tincture of the coloquintida in proof spirit, in which the scammony and aloes are to be dissolved, and the whole inspissated together. Both these forms are greatly preferable to those in which the colocynth in substance is joined to the same materials.

Pil. Rudii.

Extract.

*COLUMBO.

COLOMBA radix Ph. Edinb. Calumba Redi Exper. nat. A root brought from Columbo, a town in the island of Ceylon, to which it was originally transplanted from the continent of India. It is called by the Portugueze Raijs de Mosambique. We are as yet unacquainted with the vegetable of which it is a part.

Columbo root comes to us in circular pieces, which are from half an inch to three inches in diameter, and from two inches to one quarter of an inch in length. The fides are covered with a thick wrinkled bark, of a dark brown colour externally, but of a light yellow within. The furfaces of the transverse sections appear very unequal, highest at the edges, with a concavity towards the center. On paring off this rough furface, the root is seen to consist of three lamina, the cortical, ligneous, and medullary. This last is much the softest, and when chewed seems very mucilaginous. A number of small sibres run longitudinally through it, and appear on the surface. The cortical and ligneous parts are divided by a circular black line. All the thicker pieces have small holes drilled through them, for the convenience of drying.

This root has an aromatic fmell, but is disagreeably bitter and pungent to the taste, somewhat resembling mustard-seed long kept.

From a number of pharmaceutical experiments on the columbo root, it appears to give out its virtues more completely to spirituous, than to watery menstrua. The watery insusion is more perishable than that of other bitters. A copious precipitation takes place in it in twenty-sour hours; and in two days it becomes ropy and musty. By the united action of water and rectified spirits, an extract weighing eight ounces and two drams was obtained from twelve ounces of the root. This extract is found to retain the entire flavour of the columbo, and to be equal, if not superiour, in virtue to the powder of the root. The antiseptic power of an insusion of columbo, upon animal sless, was found to be less than that of Peruvian bark; but its efficacy in correcting and preventing the putridity of the bile, appeared to be superiour to that of bark or chamomile flowers. Columbo considerably checks the progress of fermentation in alimentary mixtures; and neutralizes a large proportion of acid. It has little or no aftringency.

The columbo root has long been a medicine in repute among the natives of the countries which produce it, in diforders of the stomach and bowels. They carry it about with them, and take it, fliced or fcraped, in Madeira wine. Our practitioners in the East Indies adopted the use of it from them; and frequently found it of great service in the cholera morbus, so common and fatal in those hot climates. It was observed to stop the violent vomiting in this complaint, more speedily and effectually than any other remedy; an effect attributed to its property of correcting the putrid disposition of the bile. It was, however, little known or regarded in this country, till Dr. Percival, in his Essays Medical and Experimental, Vol. II. published observations and experiments on this root, (from which the fubstance of this article is taken) with cases of its efficacy in various diseases depending on the state of the bile; as the bilious colic, bilious fevers, diarrhœas, habitual vomitings, &c. The experience of other practitioners has confirmed its utility in these cases. The dose of the powder usually employed has been from one to two scruples.

CONESSI.

CONESSI Med. eff. edinb. Codago-pala Hort. malabar. Ph. Parif. Conessi: the bark of a small tree, (Nerium antidysentericum Linn.) growing in the island Ceylon and Malabar; of a dark blackish colour on the outside, covered more or less with a white moss or scurf: the bark of the small young branches, which has the least of this matter, is preferved. It has but lately been introduced into Europe, and is as yet little known in the shops.

This bark, to the tafte gratefully auftere and bitter, is recommended in the Hortus Malabaricus for the cure of diarrhœas; and its efficacy in this diforder has been confirmed in the Edinburgh medical effays. bark, freed from the fourf, is directed to be made into an electary with fyrup of oranges, and taken to the quantity of half a dram or more four times a day: it fenfibly loses of its roughness and its virtue if kept for two or three days, in the form either of powder or electary; and hence fresh quantities are to be prepared at least every other day. It is faid, that the first day of taking the medicine, it increases the number and quantity of the stools, without increasing the gripes; that on the second day, the colour of the stools is mended; and that on the third or fourth day, if it fucceeds at all, the confiftence generally comes near to a natural state: that in recent diarrhow, from irregularities, it seldom fails to cure, if a vomit, of ipecacoanha, is given immediately before its use: that the fame management fucceeds in persons of a lax habit, to whom diarrhee are familiar in moist weather: but that, in any case, if a fever is joined, the conessi has no place till the fever is removed. Mr. Monro informs us, in the Essays above mentioned, that he cured a dysentery of three years standing, which had refisted a great variety of other medicines, by giving this bark in the form above prescribed.

CONSOLIDA.

CONSOLIDA major sive Symphytum majus Ph. Edinb. (a) Symphytum consolida major C. B. Symphytum officinale Linn. Comfry: a rough plant; with large, somewhat oval, pointed leaves; producing,

on the tops of the branches, spikes of white or purplish, pendulous, nearly cylindrical flowers, followed each by four shining black seeds: the root is thick and sleshy, black on the outside and white within. It is perennial, grows wild in moist grounds, and flowers in May or June.

The roots of comfry abound with a viscid glutinous juice, of no particular taste or slavour. The dried root, boiled in water, renders a large proportion of the sluid slimy: the decoctions, inspissated, yield a strong slavourless mucilage, similar to that obtained from althea, but somewhat stronger-bodied or more tenacious, and in somewhat larger quantity, amounting to about three sourths the weight of the comfry. From this analysis it may be presumed, that the consolida, though rarely employed in practice, is rather superiour to the althea in the several intentions in which that root is made use of; the mucilaginous matter being, in both roots, the only medicinal principle.

CONTRAYERVA.

contrayerva: the root of a fmall plant (dorstenia sphondylii folio, dentariæ radice, Plum. gen. dorstenia contrayerva Linn.) growing in Peru and other parts of the Spanish West Indies. The root is an inch or two in length, and about half an inch thick; full of knots and irregular tubercles; surrounded on all sides with numerous long tough sibres, most of which are loaded with scaly knobs; of a reddish brown colour on the outside, and pale within. It was sirst brought into Europe, about the year 1581, by Sir Francis Drake, whence its name Drakena radix.

This root, freed from the fibres, which are much weaker than the tuberous part, has a light aromatic fmell; and a roughish, bitterish, penetrating taste, which, as Fuller observes, is not easily concealed by a large admixture of other substances. It is given, as a diaphoretic and antiseptic, in low and malignant severs, and appears to be one of the mildest and safest of the substances of the pungent kind commonly made use of in these intentions; not being liable to produce, though taken pretty freely, any considerable heat. The dose, in substance, is from five or six grains to half a dram and more; in decoction or insusion, from half a dram to two drams.

Contrayerva

Contrayerva root gives out its virtue, by the affistance of heat, both to water and rectified spirit; and tinges the former of a dark-brownish red, the latter of a brighter reddish colour: the watery decoction is very mucilaginous, so as not to pass through a filter. In distillation or evaporation, pure spirit elevates nothing, and water very little of its virtues; the active matter of this root being of the fixt kind, and remaining nearly entire in the watery as well as in the spirituous extracts. The extract made by rectified spirit tastes strongly of the contrayerva, and leaves in the mouth a durable, glowing, vibrating kind of pungency, like that of peppermint, but far milder: its quantity is about three drams from sixteen of the root. The quantity of watery extract is more than double to that of the spirituous, and its taste proportionably weaker.

COPAL.

COPAL Pharm. Edinb.(a) COPAL: supposed to be a resinous concrete, obtained from certain large trees growing in New Spain; more probably a mineral bitumen, analogous to amber. It is brought to us in irregular masses; some of which are transparent, and of all the intermediate shades of colour from a very light yellow to a gold colour and a deep brown; others are whitish and semitransparent. In the middle of some of the masses is found a cavity containing some drops of a clear saline liquor: in others, insects and vegetable matters are inclosed (b).

These concretes, however various in appearance, are in quality very nearly if not entirely alike. They have a more agreeable smell than frankingense, to which some have resembled them, and when laid on a red-hot iron, they do not melt so thin, or burn away so fast. They do not soften in the mouth on being chewed, like anime, with which they have been confounded by others. From those, and from all the other known resinous bodies, they differ more remarkably, in being exceeding difficultly dissoluble in rectified spirit of wine, and yielding in distillation an oil which like the mineral petrolea is also indissoluble in spirit; in being readily dissolved by the concentrated vitriolic acid,

⁽a) Expunged.

and effential oils, not by expressed oils, or volatile or fixt alkalies. Though pure spirit of wine, by moderate digestion, seems to have little action on copal; a boiling heat, or long agitation, enables it to take up a considerable portion: the undissolved part is tenacious and white: the filtered solution is of a gold yellow colour, in taste at first sweetish, afterwards agreeably aromatic, inclining to bitter. It is said that spirit of wine saturated with camphor dissolves the copal more easily than pure spirit. The medicinal effects of this bitumen are not much known, as it has never been much in use: it is recommended as a warm corroborant; and may be presumed to be similar to amber.

CORALLINA.

corallina officinarum C. B. Coralline: a marine production, common on rocks, shells, &c. resembling a small plant without leaves; consisting of numerous slender jointed branches, generally of a greyish or whitish colour, sometimes greenish, yellowish, reddish, or blackish; of a brittle stony substance, friable betwixt the singers, and crackling between the teeth. It has been commonly supposed a vegetable; but late observations afford grounds to believe, that it is of an animal origin: it is apparently the habitation, and probably the production, of marine polypi.

Powdered coralline has been celebrated in doses of from half a scruple to a dram, as an anthelmintic; probably on too slight foundation. As usually met with in the shops, it has no smell, and very little if any taste. It is almost wholly dissolved by aqua fortis, is precipitated from this acid by the admixture of the vitriolic, and by calcination in a strong fire becomes a true quicklime; proofs that it consists chiefly of an abforbent calcareous earth, and that it is of the same general nature with crabs-claws, oyster-shells, and other testaceous marine bodies.

CORALLIUM.

CORALLIUM rubrum Pharm. Lond. RED CORAL: a hard, brittle, branched fubstance, resembling the stalk of a plant; usually

about the thickness of a goose quill; full of knots; sometimes straight, and sometimes variously bent; both externally and internally of a deep bright red colour. It is sound adhering to rocks and other bodies in the sea, particularly in the Mediterranean; covered with a soft sungous bark, in which is a great number of cells curiously divided, containing a milky juice, with apertures on the surface: this cortical part is separated while fresh and soft. It has been generally referred to the vegetable kingdom; but is more probably, like the preceding, the work and the nest of little animals.

RED coral appears to have for its basis the same calcareous animal earth as the corallines and the shells of sea sistes: like them, it is changed by calcination into quicklime, dissolves in all acids except the vitriolic, and is precipitated by this last from its solutions made in the others. It is used also, like those productions, as an absorbent of acid humours in the first passages; and like them, when satiated with such acids as are generated in the bodies of animals, it forms therewith a restringent saline compound. That it has any virtues distinct from those of the other calcareous absorbents, or that it is superiour in absorbent power to the cheaper testaceous bodies, there are no grounds to suspect.

Corallium præparatum Pb. Lond. The levigated coral is fometimes counterfeited in the shops with the common testacea coloured with dragons blood or red bole. These abuses may be discovered, by shaking the powder with water and suffering it to settle, when the white and the coloured matter will separate from one another and appear in great part distinct: dragons blood may be known likewise by its giving a red tincture to spirit of wine; and bole, by its retaining its redness in the fire, whilst coral burns white.

CORIANDRUM.

CORIANDRUM Pharm. Lond. & Edinb. Coriandrum majus C. B. Coriandrum fativum Linn. Coriandrum sativum Linn. Coriandrum pale yellowish or brownish, striated, hemispherical seeds, which are joined, by the flat sides, two together. It is annual, a native of Italy, and cultivated in some parts of England.

THE leaves of coriander have a strong smell, somewhat of the aromatic kind, but not a little disagreeable. The seeds also have, when fresh, a very unpleasant slavour, which by drying is altered and becomes tolerably grateful: their taste, in this dry state, is moderately warm

and flightly pungent.

The dried feeds are fometimes employed as a stomachic and carminative, though less frequently than the other warm feeds. They give out their virtue totally to rectified spirit, but only partially to water: the spirituous tincture is of a pale bright yellow colour, the watery insusion of a deeper brownish. In distillation with water, they yield a small quantity of a yellowish essential oil, which smells strongly and pretty agreeably of the coriander. Pure spirit likewise carries off, in evaporation or distillation, great part of their slavour; the spirituous extract proving, in taste as well as in smell, considerably weaker than the tincture uninspissated.

CORNU CERVI.

CORNU CERVI Pharm. Lond. HARTSHORN: large branched horns of the hart or male red deer. The horns usually met with in the shops are those of the common or fallow deer.

Hartshorn, rasped or shaved, gives out to water, by boiling, a soft gelatinous matter, of scarcely any particular slavour, amounting, when inspissated, to about one fourth the weight of the horn. The decoction and gelly are sometimes directed in diarrhæas and other disorders, partly as affording a mild nutriment, and partly for obtunding and incrassating acrimonious thin humours. An elegant hartshorn gelly is prepared, by boiling half a pound of the shavings in three quarts of water till two parts are wasted, and adding to the strained liquor an ounce of Seville orange or of lemon juice, a quarter of a pint of mountain wine, and half a pound of fine sugar; and boiling down the mixture to a due consistence. Compositions of this kind are very grateful to many in acute diseases.

On distilling the horn with a red heat, it gives over a volatile salt and spirit, together with a fetid empyreumatic oil. The same products are obtainable, in greater or less quantity, from all animal substances,

N n though

though those prepared from hartshorn have been in most general use: see Sal alkalinus volatilis.

Cornu cervi philosophice præparatum.

Cornu cervi calcinatum
Ph. Lond.

The horns, so far freed from their gelatinous matter as to be pulverable, either by boiling them in water or by exposing them to its steam, have been used by some in the same intentions as the absorbent earths: by calcination in a strong sire, with the free admission of air, their earthy part may be obtained in a much purer state, in quantity about one half of the weight of the original horn. The calcination may be performed in a potter's surnace; or by stratifying the horns with charcoal in any common surnace or stove, and setting the whole on fire together: the horns, after the burning, retain their sigure, and a considerable degree of hardness, so as to be easily separable from the ashes into which the vegetable coal is reduced. The horns remaining after the distillation of the volatile salt are as proper for this use as fresh ones.

The pure earth of hartshorn differs from that of coral and the testacea, in not being convertible into quick lime; and agrees with them, in being diffoluble by the vegetable, nitrous, and marine acids, and in being precipitated from these acids on the addition of the vitriolic. The earth of the horns, and of the bones also, of other animals, appears to be of the same nature with that of hartshorn. How far this species of earth differs from the others in its medicinal effects, is little known. It is customary, in acute diseases accompanied with a looseness, to impregnate the common drink with the calcined hartshorn levigated into an impalpable powder, on a supposition of its acting as a mild restringent: solutions of it in vegetable acids are apparently restringent, as they discover a degree of aufterity to the tafte; but the pure earth is infipid, fo that probably it tends to restrain fluxes only in consequence of its uniting with acid humours in the first passages. Hoffman reports, that this earth, when combined with acids, is more disposed, than the other absorbents, to promote perspiration.

COSTUS.

COSTUS Ph. Lond. Costus dulcis officinarum C. B. Costus arabicus Linn. Costus: a root, brought from the East Indies, about the fize of the finger, of a pale greyish or whitish colour on the outside, and yellow within. An Arabian, a bitter, and a sweet costus were formerly distinguished

distinguished in the shops: whether they were, as some suspect, the roots of different plants, or, as others, of one and the same plant in different states, does not fully appear. At present, only one sort is met with, and this but rarely.

This root has been recommended as a stomachic, uterine, diaphoretic, and diuretic. The editor of the Wirtemberg pharmacopæia observes, that it impregnates the urine, like the turpentines, with a violet smell. The smell of the root itself approaches to that of violets or Florence orris: its taste is warm and bitterish. Both the smell and taste are confined chiefly to the brittle cortical part, the internal tough woody matter having very little of either.

Decoctions of costus in water are of a brownish colour, a bitter taste, and less grateful smell than the root in substance: in evaporation they diffuse a very disagreeable scent: the inspissated extract is moderately bitter, of scarcely any particular flavour, in quantity amounting to no less than two thirds the weight of the root; in keeping, it soon grows mouldy, and dusty on the surface. The spirituous tincture is of a bright yellow colour, a bitter aromatic taste, and a more agreeable smell than the watery decoction; inspissated, it yields a very warm pungent, bitter extract, of an aromatic slavour, less grateful than that of the costus itself, in quantity not exceeding one ninth the weight of the root.

CRETA.

CRETA Ph. Lond. & Edinb. CHALK: a pure white mineral calcareous earth, met with in most parts of the world, of various degrees of hardness. The softer masses, included in sea shells (which are commonly in chalk pits) called from their figure chalk eggs, are by many preferred to the others for medicinal use.

There are innumerable concretes in the mineral kingdom, of the same general nature with chalk; or which consist chiefly, or wholly, of the same earth, formed into masses, which differ from one another little otherwise than in their external appearance, compactness, or texture. Such are the limestones; the marbles; some of the marles; the sine earth called agaricus mineralis, medulla saxi, or lac lunæ: the transparent crystalline concretes called spars; most of the petrefactions; and most of the

N n 2 ftalactitæ,

stalactive, or the earthy matter, which, in its concretions from waters, incrustates the sides of caverns, or hangs in icicles from the tops. Many of these bodies were formerly introduced into medicine, from an erroneous supposition of their possessing distinct qualities: chalk, one of the purest of them, is the only one now retained in practice; nor would the art suffer any detriment, if a like reduction was made in the analogous bodies surnished by the animal kingdom.

The distinguishing characters of this earth, in all its forms, are, its not dissolving in the vitriolic acid, though readily dissolving in all the others; its being precipitated by the vitriolic acid from its solutions in the others, and being thus changed into a selenites; its being convertible, by calcination with a strong fire, into quicklime; and its melting easily, with certain vitrefactive sluxes, as borax, into a transparent glass. In this last property it differs from the calcareous animal earths; which appear to be unvitrescible, communicating, to a large proportion of vitreous matters, an opake milkiness.

Pure chalk is a very useful absorbent in cardialgic and other complaints from acidities in the first passages. For this purpose it is formed, in the shops, into troches, with sugar and a little nutmeg, and generally with the addition of some of the other absorbent earths, which add nothing to its virtue; and into juleps, by mixing the chalk, levigated into a subtile powder, with water, in the proportion of half an ounce to a pint; with the addition of two drams of sugar, and one dram of gumarabic, to give some consistence to the liquor, so as to enable it to keep the powder suspended. *The Edinburgh college have now directed a powder, consisting of sour ounces of prepared chalk, with a dram and a half of nutmeg and a dram of cinnamon, to supply the place of the cardialgic troches.

Julepum e creta Ph. Lond.

Pulv. Cretaceus Ph. Edinb.

When chalk is combined with fuch acids, as may be deemed most analogous to those which are preternaturally generated in the human stomach, as sour milk and sour vegetable liquors; the compounds prove somewhat more austere than those resulting from the coalition of the same acids with the animal absorbents: hence chalk, given in cases of acidities, is generally observed to bind the belly more; and thus to prove more injurious in costive habits, and more beneficial in suxes: two drams given in a dose, and repeated at proper intervals, have often effected a speedy

a speedy cure both in simple diarrhoe and in dysenteries. But that it has any astringent power, as many have supposed, independently of its combination with acids, or in disorders where there are no acid juices in the first passages to dissolve it, is not so clear: the sense of astringency, which the chalk in substance produces in the mouth, appears to proceed, like that of the bolar earths, only from its adhering to the part and imbibing its moisture.

CROCUS.

CROCUS Ph. Lond. & Edinb. Crocus fativus C. B. & Linn. Crocus autumnalis fativus Morifon. bift. SAFFRON: a bulbous-rooted plant, with narrow grafs-like leaves which have a white line running along the middle: the stalk is short and undivided, and bears on the top a purplish blue flower, deeply cut into six segments; in the middle of the flower arises, among the stamina, a whitish pistil, divided at the top into three stigmata or sleshy silaments, the lower part of which is slender and pale coloured, the upper broader, of a deep orange red, and very sinely indented about the sides: these silaments, carefully picked, moderately dried in kilns, and pressed together into cakes, are the saffron of the shops. The plant is perennial, and slowers in autumn: the common spring crocuses of our gardens are reckoned by Linnæus to be no other than varieties of it.

Saffron is cultivated in different parts of the world: that produced in our own country is greatly superiour to the forts brought from abroad, and may be distinguished from them by its blades being broader. It should be chosen fresh, not above a year old, in close cakes, neither dry nor yet very moist, tough and firm in tearing, of a high fiery colour, staining the hands on rubbing it, and of the same colour within as on the outside.

SAFFRON is a very elegant and useful aromatic; of a strong, penetrating; diffusive smell, and a warm, pungent, bitterish taste. It is supposed to have a considerable degree of anodyne power, depending on its subtile odorous principle; to be more cordial and more exhibitant than almost any of the other aromatics, so as, when taken too freely, to occasion even immoderate mirth (a); to be particularly serviceable in disorders of the

breast, in female obstructions, and hysteric depressions. It tinges the urine of a high colour. The dose is commonly from two or three grains to ten or twelve: Geoffroy says it may be extended with safety to a scruple and more.

Saffron gives out the whole of its virtue and colour to rectified spirit, proof spirit, wine, vinegar, and water: about three parts in sour of the saffron are taken up by each of these menstrua; and the matter which remains undissolved is inodorous, insipid, and of a pale clay colour. The officinal tinctures are made with sack to French brandy to in the proportion of an ounce to a pint. By dissolving in the vinous tincture a proper quantity of sine sugar, twenty-sive ounces to a pint, an elegant cordial syrup is obtained. Tinctures drawn with vinegar, or other liquors sensibly acid, soon lose of their rich colour in keeping: the colour of the vinous tinctures also sades a little, and a part of the dissolved saffron is apt to be in time thrown off: those made in proof spirit, and in rectified spirit, particularly the latter, may be kept in perfection for years.

In distillation, it impregnates water strongly with its flavour: if the quantity of saffron is large, a small portion of a fragrant and very pungent essential oil may be collected, amounting, as is said by Vogel, to about a dram and a half from sixteen ounces. The remaining decoction, inspissated, yields an extract of a high colour, in taste unpleasantly bitterish, without any thing of the distinguishing smell or slavour of the saffron.

Rectified spirit elevates also a considerable share of its slavour, but leaves much the greatest part concentrated in the extract. This extract, inspissated only to the consistence of oil, is recommended by Boerhaave as one of the highest cordials and exhibit extracts: the dose is a few drops, which may be taken in a glass of rich wine. It dissolves in wine and in water, as well as in spirit, and mingles also with oils both expressed and distilled; appearing to be a substance of a peculiar nature. The spirit, distilled from saffron, is said to have an advantage above most other cordial spirits, of disposing the patient to sweat.

† Vinum
croceum
Ph. Lond.

‡ Tinctura
croci Ph.
Edinb.

|| Syrupus
croci Ph.
Lond.

CRYSTALLUS.

CRYSTALLUS Pharm. Edinb. (a) ROCK CRYSTAL: a transparent colourless stone; of a regularly angular figure, which is generally that of an hexagonal column terminated by a pyramid of the same number of sides; hard, so as to strike fire freely with steel; becoming white, opake, and friable, by repeated ignition and extinction in water; not dissoluble by any acid, either in its natural state or when calcined; fusible, with vitrefactive sluxes, into a nearly colourless glass.

Rock crystal, and some other stones of the same general nature, introduced into medicine by the credulity of former times, not yet expunged from catalogues of the materia medica, and in some places still made ingredients in officinal compositions; appear, from their indissolubility in every known species of humid menstruum, to be incapable of exerting any action in the human body: unless that by the rigidity and hardness, which their particles retain, however finely levigated, they may offend the stomach and intestines; or that by virtue of the calcareous earth, which they abrade plentifully from the marble instruments with which they are levigated, the prepared powder may act as an absorbent.

The colours of the precious stones appear to depend on a principle distinct from the stony matter which makes their basis. It is said, that the sapphire, emerald, amethyst, and cornelian, on being urged with a strong fire, become colourless, and nearly similar to common crystal: that the emerald, in parting with its tinging matter in the fire, emits visible stames (b): that the hyacinth and garnet melt, in a vehement fire, into a vitreous mass, of a brownish or blackish colour like that which ferrugineous calces communicate (c). From these kinds of experiments it has by some been inferred, that the coloured precious stones, though their stony basis is confessedly inactive, may, nevertheless, have some medicinal powers depending on the tinging metallic impregnation (c). But surely this reasoning does not take off the impropriety, or rather absurdity, of using, as medicines, these costly concretes, from a possibility of their producing effects, which far cheaper substances are known to produce with certainty.

⁽a) Expunged. (b) Geoffroy, Trad. de mat. med. tom. i. p. 95.

⁽c) Pott, Chymische untersuchungen von der lithogeognosia, p. 45.

CUBEBÆ.

CUBEBÆ Pharm. Lond. & Edinb. Cubebæ vulgares C. B. Cubebæ: dried berries, of an ash brown colour, generally wrinkled, greatly resembling pepper, but surnished each with a slender stalk, whence they are called by some piper caudatum. They are the fruit of an East Indian tree, of which we have no particular account, said to resemble the apple tree, and to produce its berries in clusters.

Cubebs are a warm spice; of a pleasant smell, and a moderately pungent taste. Their heat and pungency are weaker than those of pepper, but of the same kind; and reside, like those of that spice, not in the volatile, but in the more fixt matter. In distillation with water, they yield a small quantity of a ponderous essential oil, of an agreeable and moderately strong smell, but in taste mild: the remaining decoction, inspissated to the consistence of an extract, retains a considerable share of the warmth and pungency of the cubebs. An extract made with rectified spirit possesses the whole of their slavour in perfection, for even the odorous principle does not exhale or distil with this menstruum: the taste of this extract is very hot and pungent, though not near so much so as that of the spirituous extract of pepper.

CUCUMIS.

CUCUMBER: an annual herb, with naked monopetalous flowers divided into five fegments, and a large juicy fruit produced under the flower. Thus far the characters of the cucumber agree with those of fome other plants, whose feeds have been commonly ranked among the officinals, and which may properly be placed together.

- 1. CUCUMIS HORTENSIS Ph. Edinb. (a) Cucumis fativus vulgaris C. B. Cucumis fativus Linn. Cucumber: with oblong fruit, often covered with little protuberances; and oblong white feeds.
- 2. CITRULLUS Pharm. Edinb. (a) Anguria citrullus dicta C. B. Cucurbita Citrullus Linn. Citrul: with very large, roundish, smooth hard rinded fruit; and oblong, broad, rhomboidal, blackish seeds.

- 3. CUCURBITA Pharm. Edinb. (a) Cucurbita lagenaria flore albo, folio molli C. B. Cucurbita lagenaria Linn. Gourd, bottle-gourd: with very large, thick woody rinded fruit, bellied like a bottle; and long whitish seeds, having two angles like horns at the top.
- 4. Pepo Pharm. Edinb. (a) Pepo vulgaris Raii hist. Cucurbita Pepo Linn. Common pumpion: with very large, roundish or oval fruit; and rhomboidal whitish seeds, having a rim or elevated line round the edges.

THE cucumber and citrul are esteemed cooling and relaxing; salubrious in hot bilious dispositions, and where there is a tendency to inflammation; prejudicial in the opposite circumstances; dissicult of digestion, and of very little nourishment.

The feeds of all these plants are similar in quality; and have been generally used promiscuously, and distinguished by the title of the greater cold seeds. They have a sweetish taste, accompanied with some unctuosity, and no smell or particular flavour: on expression, they yield a soft insipid oil, of the same general nature with that of almonds: on trituration with water, their oil, by the mediation of the mucilaginous and farinaceous matter of the seed, unites with the water into an emulsion or milky liquor. These emulsions have been used as diluents, refrigerants, and emollients, in the same cases as those prepared from sweet almonds; which last are now almost universally preferred. The seeds in substance have likewise been made ingredients in some officinal emollient powders; for which purposes they are not well adapted, as being liable to grow soon mouldy and rancid in keeping, especially in a powdery form: those of the cucumber seem to be the least subject to this inconvenience.

5. CUCUMIS AGRESTIS Ph. Lond. Cucumis filvestris assininus dictus C. B. Momordica Elaterium Linn. Wild cucumber: with warty, hairy, somewhat oval fruit, not above two inches in length: the fruit, when ripe, bursts on being touched, and throws out with violence its whitish juice and its black seeds. It is sown annually, as all the preceding, in gardens.

(a) Expunged.

MATERIA MEDICA.

ALL the parts of the wild cucumber are strongly purgative: the fruit appears to be fomewhat more fo than the root, and this than the leaves. The juice that iffues spontaneously, or by very light pressure, on slitting the fruit when almost ripe, has an unpleasant smell, and a very durable nauseous bitter taste: on standing for a few hours, it parts into a thick matter which fettles to the bottom, and a thin watery fluid, which floats above: this last may be commodiously drained off, after the clearer part is decanted, by means of strips of woollen cloth or skains of cotton laid over the fides of the veffel. The thick fecula, dried in the fun or any other gentle heat, is a very firong, irritating, but flow, cathartic; and often operates likewise upwards. It remarkably raises the pulse, and feems to kindle a degree of fever for a time: Lifter and Hoffman observe, that its effect in increasing the pulse is perceivable even in the extremities of the fingers. Its use therefore is in cold indolent phlegmatic cases; particularly in dropsies, in which it has some times been given with fuccess after medicines of a milder kind had proved ineffectual. Two or three grains are in general a fufficient dose: in some cases this quantity has acted violently: in others, five grains have procured plentiful evacuation, without much uneafiness or disturbance to the constitution. It is said, that in Holland, an extract made with wine from the roots of the plant is substituted to the elaterium, and has been found to be equally efficacious and fafe: though Boulduc (a) speaks of an extract of this root, made probably with water, which appears to have been much weaker; the dose being from twenty-four to thirty grains. In what kind of matter the purgative virtue of this plant refides, has not been fufficiently examined: according to Boulduc, spirit of wine has scarcely any action upon it: and that water is not its proper menstruum appears from its quitting the watery juice and fettling to the bottom.

Elaterium Pb. Lond. & Edinb.

CUPRUM.

CUPRUM Pharm. Lond. Cuprum five Venus Pharm. Edinb. Copper: a reddish metal, nearly nine times specifically heavier than water; requiring for its sussion a strong white heat, and calcining by a continuance of a weaker red heat into a dark reddish powder; contracting,

(a) Histoire de l'acad. roy. des scienc. de Paris, pour l'ann. 1719.

from long exposure to the air, a greenish rust; soluble in all acids and in volatile alkaline spirits, and exhibiting, when dissolved, a blue or green colour, or a colour composed of the two. Volatile spirits, in particular, receive from a small proportion of it a beautiful deep blue; and if added to solutions of it made in acids, when so far diluted as to appear almost or altogether colourless, change them immediately to the same sine colour. If a piece of bright iron be immersed in the acid solutions, the acid quits the copper to attack the iron; and the copper, in its separation from the menstruum, adheres to the iron, which soon appears covered with a cupreous coat. On these principles, very minute quantities of copper, dissolved in liquors, may be readily discovered.

There are confiderable mines of copper in England, Sweden, Germany, and many other parts of the world. The ores are often, wholly or in part, of beautiful blue or green colours like those of the solutions: all the mineral stones, tinctured with these colours, are supposed to receive them from this metal. Most of the ores abound with sulphur: which this metal very strongly retains, and which is difficultly separated by repeated calcinations and suspenses.

Pure copper, in its metallic state, or calcined by fire, appears to be indissoluble, and of no considerable effect, in the bodies of animals. There are instances of pieces of copper having been swallowed, and lying long in the bowels, without seeming to act any otherwise than by their bulk or figure.

Diffolved in the nitrous or marine acids, and crystallized or exsiccated by heat, it proves a strong caustic. These preparations, particularly that with the nitrous acid, were formerly sometimes made use of in this intention; but have long been laid aside, on account of their great disposition to liquefy.

Combined with the vitriolic acid, or with the acids of the vegetable kingdom, or corroded by the air, it acts, when externally applied, as an efficacious detergent and a gentle escharotic; when taken internally, as a virulent emetic and cathartic. Some have ventured on small doses of these preparations, as quick emetics, for procuring immediate evacuation where poisonous substances have been swallowed: but that end may surely be obtained by less dangerous means. A more particular

002

account

account of these preparations is given under their respective names, Ærugo and Vitriolum.

A faturated folution of the metal in volatile spirits is recommended by Boerhaave in diforders proceeding from an acrid, weak, cold, phlegmatic cause. He says, that if three drops be taken in the morning with a glass of mead, and the dose doubled every day to twenty-four drops, it proves attenuating, warming, and diuretic: that by this medicine he once cured a confirmed afcites, though in other cases of the same kind it failed: that it is the only preparation of copper which does not prove emetic; and that, as it does no harm, it may be tried with fafety(a). It is probable, that this preparation differs from the preceding only in containing less copper, the quantity which volatile spirits dissolve being extremely small; and that in considerable doses, it would exert the same virulent operation with the other folutions or foluble preparations of this metal. * A folid preparation of this kind made by rubbing together in a glass mortar two parts of blue vitriol and three of the volatile salt procured from fal ammoniac, till all effervescence has ceased, and then gently drying the concrete, is ordered in the last Edinburgh pharmacopæia, under the name of cuprum ammoniacum. It has frequently been given with fuccess in epileptic and convulsive disorders.

A tincture, differing little in its cupreous impregnation from that made with volatile spirits, is used in the shops as an external detergent, and for confuming specks or films of the eyes. This is prepared with a folution of fal ammoniac in lime-water, in the proportion of a dram+ Ph. Edinb. or four scruples; of the salt to a pint of the liquor; which solution is tinged of a fapphire blue colour by flanding for fome days with fome flips of copper or in a copper veffel+, or by the addition of eight grains of verdegris 1.

> Copper vessels, in certain circumstances, give a taint to almost all kinds of liquors, even to pure water; more especially if they have not been thoroughly cleanfed from the rust which they contract by lying exposed to a moist air: in certain circumstances, however, they appear to refift even liquors of confiderable acidity. Most of the vegetable acids, fo long as they are kept boiling in copper veffels, have little or no action on the metal; though in a gentle heat, or in the cold, they become in

Aq. fapphir. + Ph. Lond.

a short time impregnated with its ill taste and with its pernicious qualities, corroding it chiefly at the surface of the liquor: if the metal is only moistened and exposed to the air, it is corroded more speedily; and sooner still if exposed to the vapours of the acid. The most acid syrups are prepared by the confectioners, by boiling in copper vessels kept perfectly clean, without receiving any hurtful impregnation; whereas the far less acid liquor, that rises towards the end of long protracted distillations of simple waters, corrodes, in its passage through the copper head, in the form of vapour, so much of the metal as to prove emetic.

Brass is a combination of copper with the metallic part of calamine, that is, with zinc, a metallic body still easier of dissolution than the copper itself. The two metals, nevertheless, form by their coalition a new compound, which does not rust in the air, or dissolve in vegetable acids, or calcine in the fire, near so soon as either of them separately.

CURCUMA.

CURCUMA Pharm. Lond. & Edinb. Curcuma radice longa Herm. bort. lugd. bat. Maniella kua Hort. malabar. Crocus indicus; Terra merita; Cypira. Curcuma longa Linn. Turmeric: a small tuberous knotty root, brought from the East Indies; externally greyish, internally of a deep lively yellow or saffron colour, which by age changes towards a red. Two sorts are mentioned by authors, one longish, the other roundish: only the first is met with in the shops.

TURMERIC has a flight, aromatic, and not very agreeable smell; and a bitterish somewhat warm taste. It readily gives out its active matter both to aqueous and spirituous menstrua: to the former it communicates its own deep yellow, to the latter a fine yellowish red tincture. In distillation with water, it yields a small quantity of a gold coloured essential oil, of a moderately strong smell, and a pungent taste: the remaining decoction, inspissated, leaves a bitterish, considerably saline, mass. Rectified spirit elevates little or nothing of its virtue; all the active parts of the root being left behind in the inspissated extract, which is moderately warm, and bitter, and not a little nauseous.

This

This root is said to be in general use in the eastern countries, both tor the colouring and seasoning of food, and as a medicine: it is accounted one of the most effectual remedies in obstructions of the viscera and mesentery, which are there frequent; in uterine disorders, difficulties of urine, and affections of the kidnies (a). Among us it has been employed also as a deobstruent, and esteemed by some a specific in the jaundice: the dose in substance is from a scruple to a dram; in decoction or insusion, twice as much. It tinges the urine of a deep yellow colour.

CYANUS.

CYANUS Pharm. Parif. Cyanus fegetum C. B. Centaurea Cyanus Linn. Blue Bottle: a greyish green plant, with long narrow leaves, of which the lower are deeply jagged, the upper entire, those between furnished with one or two long perpendicular ears on each side towards the bottom; the stalk divides, near the top, into several branches, each of which is terminated by a large blue flower, consisting of tubulous indented flosculi set in a smooth scaly head; the outer flosculi are larger than the inner, and widened in the upper part like a funnel; the scales are serrated about the edges. It is annual, common in corn fields, and found in flower greatest part of the summer.

The flowers of cyanus, hastily dried, preserve their colour better than most of the other blue flowers: they agree with the others in giving no blue tinge to spirit, and differ from most of them in giving none to water. In substance, they discover very little smell, and scarcely any taste: an extract made from them by rectified spirit has a weak saline austerity mixed with a kind of sweetishness: an extract made by water is less austere and more manifestly saline. From hence it may be presumed, that among the various and opposite virtues ascribed to these flowers, the antiphlogistic, aperient, and diuretic ones have the best foundation; though even these they appear to possess only in a low degree.

The varieties of this plant, produced by culture in gardens, are not materially different in quality from the wild fort. Another species, of

oriental origin, cyanus orientalis major moschatus slore purpureo & albo Morif. bift. ox. Centaurea moschata Linn. commonly called sultan flower or fweet fultan, promises, by its musky fragrance, to have some claim to the cordial and antispasmodic virtues, which have been groundlessly afcribed to our indigenous cyanus.

CYDONIA.

CYDONIA MALUS Pharm. Lond. & Edinb. Cotonea malus J. B. Mala cotonea majora & minora C. B. Pyrus Cydonea Linn. a low tree, with uncut leaves, bearing a fruit like a pear; a native of the rocky banks of the Danube, and common in our gardens.

This fruit has a pleasant strong smell, and a very austere acid taste. Its expressed juice, taken in little quantities, as a spoonful or two, proves a mild, cooling, reftringent stomachic; of good service in nauseæ, vomitings, nidorous eructations, and some kinds of alvine fluxes. A grateful and lightly cordial restringent syrup is prepared, by digesting Syrup. eythree pints of the depurated juice with a dram of cinnamon, half a dram doniorum Ph. Lond. of ginger, and half a dram of cloves, on warm ashes, for fix hours, then adding a pint of red port, and diffolving in the strained liquor nine pounds of fugar. An useful restringent gelly or marmalade is made, by boiling the juice with fine fugar to a due confistence, in the proportion commonly of three pints to a pound. If the quinces, after they are gathered, be kept for some time in a dry airy place, their juice will become richer by the diffipation of a part of their aqueous humidity.

The feeds of quinces abound with a mucilaginous substance, which they readily give out to boiling water. A dram, boiled in fix ounces of water by measure, renders the liquor slimy, almost like the white of an egg: two drams make it quite thick. On inspissating the decoctions, Mucilago the quantity of dry extract amounts to about half the weight of the feeds. Ph. Lond. This mucilage has a flight agreeable smell, and a sweetish taste, more grateful than that of the other common mucilages. In keeping, in its foft state, it foon grows mouldy.

fem. cydoni,

CYMINUM.

CYMINUM.

CYMINUM Pharm. Lond. Cuminum Pharm. Edinb. Cuminum femine longiore C. B. Cuminum Cyminum Linn. Cummin: an umbelliferous plant, refembling fennel, but much smaller; producing longish, slender, plano-convex seeds, of a brownish colour with yellowish striæ. It is annual; a native of Egypt and Ethiopia; and cultivated in the islands of Sicily and Malta, from whence we are supplied with the seeds.

CUMMIN feeds have a bitterish warm taste, accompanied with an aromatic slavour, but not agreeable. They give out great part of their smell by insusion in water, but very little of their taste: in distillation with water, a pungent oil arises, of a strong ungrateful flavour like that of the seeds: the decoction, inspissated, leaves a weakly roughish bitterish extract. Rectified spirit takes up the whole virtues of the cummin by insusion, and leaves them nearly uninjured in evaporation: the inspissated mass is very warm, moderately pungent, and not a little nauseous.

These seeds are accounted good carminatives and stomachics; but have now, in great measure, given place, for these purposes, to medicines of a more grateful kind. Their principal use is in external applications, as a warm discutient and antiseptic: the college of London have given a compound plaster and cataplasm of cummin.

CYNOGLOSSUM.

CYNOGLOSSUM Pharm. Parif. Cynoglossum majus vulgare C. B. Lingua canina. Cynoglossum officinale Linn. Hounds Tongue: a biennial plant; producing, the first year, large, fost, tongue-shaped, long-pointed leaves; the second year, a thick, branched stalk, with narrower and shorter-pointed leaves joined to it without pedicles; bearing, on the tops of the branches, dark purplish slowers, each of which is divided into five segments, and followed by four stat rough seeds: the root is oblong, thick, of a dark brown or blackish colour on the outside, and white within. It is found wild in shady lanes and uncultivated grounds, and slowers in June.

THE roots of hounds tongue are very juicy, and liable to grow mouldy in drying. Such as are produced in moift grounds have, when fresh, a rank, though not very strong smell, like that of the narcotic plants, which in drying is in good meafure diffipated: those, which are the produce of dry grounds, have fcarcely any fmell(a). On the organs of tafte, they make no great impression.

The medical effects of these roots are somewhat doubtful. It has been generally supposed that they are narcotic; by some, that they are virulently fo. The argument for their innocence (b), from the frequent and fafe use of a pill, to which they still give name in foreign pharmacopæias, appears unfair; the quantity of the hounds tongue root, in a dose of that opiate pill, being only about a grain; whereas the root by itself is ordered, in decoction, to the quantity of an ounce, in catarrhs,

coughs, diarrhœas, dysenteries, and hemorrhagies.

The leaves of the plant are supposed to be of the same quality with the roots: to the fmell, they are stronger and more disagreeable. Fuller reports, that he has used a syrup of the juice a multitude of times, and could never find it to cause sleep, or to be in the least virulent; and that he had often experienced it to be a great remedy, fecond to none, against hot, sharp, thin, catarrhous humours, and a cough occasioned thereby. Neither the leaves nor the roots are ever made use of in practice; and the colleges, both of London and Edinburgh, have now rejected the plant from their catalogue of officinals; all the good effects, which the accounts of those who have recommended it afford grounds to expect from it, being obtainable, without suspicion of malignity, from the products of the poppy.

CYPERUS.

CYPERUS: a plant with grass-like leaves, and three-square stalks, branched at top, bearing tufts of small imperfect flowers, each of which is followed by a naked triangular feed. It is perennial.

1. CYPERUS LONGUS Pharm. Parif. & Linn. Cyperus odoratus radice longa five cyperus officinarum C. B. Cyperus or English galangal:

⁽a) Hermann, Cynofura mat. med. edit. Boecler. tom. i. p. 178.

⁽b) Geoffroy, Mat. med. iii. 395. Ray, Hift. plant. i. 490.

with a long slender root, crooked and full of knots, of a dark brown or blackish colour on the outside, and whitish within. It grows wild in marshy places in some parts of England: the shops have been usually supplied from Italy and France, with dried roots, not superiour to those which are produced in our own country.

CYPERUS root has a pleasant aromatic smell, and a warm bitterish taste; both which it imparts almost totally to watery and to spirituous menstrua: it tinges the former of a dark reddish brown colour, the latter of a bright reddish yellow. In distillation, it impregnates water with its grateful smell; but yields, at least when only moderate quantities are submitted to the operation, no separable oil: rectified spirit carries off likewise, in evaporation or distillation, a great part of its odorous matter. The watery extract is moderately bitter, slightly pungent, and subastringent: the spirituous is in taste bitterer and warmer; but has not much smell, any more than the watery extract. The quantity of both extracts is very nearly alike, amounting to about one fourth of the weight of the root.

2. CYPERUS ROTUNDUS Pharm. Parif. & Linn. Cyperus rotundus orientalis major C. B. Round cyperus: with feveral roundish roots, about the fize and shape of an olive, connected by fibres, rough and rusty-coloured on the outside, whitish or yellowish within. It is a native of the East Indies, from whence the roots are sometimes, but rarely, brought to us. It is said to grow wild also in France.

This root has the same kind of smell and taste, and nearly in the same degree, as the foregoing. It differs, according to Cartheuser's examination of it, in giving a gold yellow tincture to water, in yielding a little more extract, one third its weight, both with water and spirit; and in the spirituous extract being more tenacious or resinous.

DAUCUS.

CARROT: an umbelliferous plant, with finely divided leaves; producing pale coloured, hairy, striated, somewhat oval, plano-convex seeds:

feeds: the entire umbel, and each of its subdivisions, have a circle of little leaves at their origin: the petala are unequal and heart-shaped.

1. DAUCUS CRETICUS Pharm. Lond. & Edinb. Daucus foliis fæniculi tenuissimis C. B. Athamanta cretensis Linn. Candy carrot: with white flowers, flat umbels; and oblong seeds, swelled or bellied in the middle, and pointed at one end. It is perennial, a native of the Levant and the mountains of Switzerland, and cultivated in some of our gardens. The seeds have been generally brought to us from the isle of Candy.

THE feeds of the candy carrot have a light aromatic smell, and a moderately warm biting taste. They have been occasionally employed as carminatives, and supposed likewise to be diuretic and emmenagogue: at present they are little otherwise made use of than as ingredients in mithridate and theriaca.

Water, digested on the seeds, becomes impregnated with their smell, but takes up very little of their taste: in distillation or evaporation, it elevates the whole of their smell and aromatic warmth, leaving a weakly-bitterish mucilaginous extract: on distilling large quantities, a small portion of a yellowish essential oil is obtained, of a moderately pungent taste, and smelling strongly of the daucus. Rectified spirit takes up the whole of their virtue by digestion, and elevates little in distillation: the remaining extract smells weakly, and tastes strongly of the seeds. The colour both of the tincture and extract made with spirit, is a bright yellow; of those with water, brownish: the quantity of spirituous extract is about one half of the watery.

2. Daucus silvestris Pharm. Edinb. Staphylinus. Pastinaca silvestris tenuisolia dioscoridis vel daucus ossicinarum C. B. Daucus Carota Linn. Wild carrot, or birds-nest, so called from the appearance of the umbels, which close and form a roundish cavity in the middle after the flowers have fallen: one or more of the inner flowers are commonly of a deep red colour, and several of the others red in part: the seeds are smaller, shorter and rounder than those of the preceding. It is biennial, common in uncultivated grounds, and flowers in June.

THE feeds of the wild carrot are similar in smell and taste to those of the daucus creticus, but weaker. The essential oils obtained from the

Pp2

two

two are nearly alike in quality, but somewhat different in quantity, the wild yielding a little less than the other. The spirituous extract of the wild is somewhat less pungent than that of the candy sort. Malt liquors, fermented with these seeds, receive from them an agreeable flavour somewhat resembling that of lemon peel, and are supposed to become useful diuretic drinks in cachectic and scorbutic disorders. * Insusions of them in water, in the proportion of three spoonfuls of the seed to a pint of boiling water, are said to have done great service in calculous cases, and to give speedy relief in strangury (a). In the shops they have frequently supplied the place of the daucus creticus, and been themselves supplied by the seeds of the garden carrot, which are much weaker in aromatic warmth than either. The garden and wild carrot are reckoned by botanists the same species of plant, their differences proceeding only from culture.

* A poultice of the root of garden carrot has been fuccessfully used to cancerous and phagædenic ulcers, the fætor of which it has not failed very speedily to remove, and generally with a great amendment of the state of the sore. Some have been brought to cicatrise by its use solely. The method of making the cataplasm is, to grate the carrots, and mix them with as much water as is necessary. The application is to be renewed two or three times a day. It is found to be most efficacious when the carrots are fresh and juicy. This remedy was recommended by Mr. Soultzer in a letter in a Magazine; and some cases of its efficacy were afterwards published in the 4th Vol. of the London Medical Observations and Inquiries.

A marmalade of carrots has also been proposed as an addition to the stock of ship's provisions, for preventing the scurvy.

DENS LEONIS.

DENS LEONIS five Taraxacum Pharm. Edinb. Dens leonis latiore folio, & angustiore folio C.B. Leontodon Taraxacum Linn. Dandelion: a low plant, with long, narrow, deeply indented or jagged leaves, lying on the ground; among which arises a single, naked, hollow pedicle; bearing a large yellow sloculous flower, set in a double cup, the outermost of which consists of several little oblong leaves turned downwards:

the flower is followed by small seeds, covered with a tust of long down: the root is oblong, slender, yellowish or brownish on the outside, and white within. It is perennial, common in uncultivated grounds, and flowers from April to the end of summer.

The roots, leaves, and flower-stalks of dandelion abound with a bitterish milky juice, of no smell or particular flavour. They promise to be medicines of no inconsiderable efficacy in sundry chronical disorders, as mild detergents and aperients, similar to the cichoreum silvestre, but stronger. Boerhaave had a high opinion of this and the other lactescent plants; and esteems them capable, if duly continued, of resolving very obstinate coagulations and obstructions of the viscera. Their more immediate sensible operation is, to loosen the belly, promote the urinary discharge, and render the water high coloured, without exciting any preternatural heat. *Bergius affirms that he has frequently succeeded in resolving scirrhi of the liver, by the long continued use of a decoction of dandelion root and sorrel leaves in whey or water, with the addition of yolk of egg; at the same time giving cream of tartar. This method also succeeded in the stone of the gall-bladder, and ascites (a).

The expressed juice of the plant has been taken to the quantity of a quarter of a pint or more, three or four times a day: it seems to lose nothing of its virtue in being gently inspissated to the consistence of an extract; which is moderately and not unpleasantly bitter, with some degree of sweetishness. The dried roots, which are stronger in taste than the leaves, give out their virtues both to water and rectified spirit; and tinge the former of a brown, the latter of a yellow colour. The tinctures and insusions, gently inspissated, differ little from the inspissated juice; except that the watery extract is rather weaker; and that the spirituous, which is in smaller quantity, has a stronger bitter taste, and discovers also a slight astringency. Cartheuser says, that the watery extract amounts to one fourth, the spirituous only to one eighth the weight of the root.

Neither the plant in substance, nor its preparations, bear keeping well: after the dried root had lain about a twelvemonth, its bitterness was wholly lost, and only a slight sweetishness remained: an extract made from the fresh root, inspissated to dryness, and kept for the same length of time, suffered nearly the same change.

DICTAMNUS.

DICTAMNUS ALBUS Pharm. Edinb. & Linn. Dictamnus albus vulgo five fraxinella C.B. WHITE OF BASTARD DITTANY: a plant with oval acuminated leaves, like those of the ash tree, but smaller and more juicy, set in pairs on a middle rib, which is terminated by an odd one: on the tops of the stalks stand elegant long spikes of irregular white or purplish slowers, followed each by sive pods full of shining black seeds. It is perennial, and grows wild in the mountainous parts of France, Italy, and Germany; from whence the white cortical part of the root, freed from the sibres and pith, is sometimes brought to us, dried, and rolled up in the form of quills.

THE herb has a strong smell, of an unpleasant resinous or bituminous kind: on the tops of the stalks, and the flower cups, the microscope discovers innumerable little vesicles, filled with an essential oil, the source of its strong scent. It is said, that on the approach of a candle, in very hot dry weather, its essential vest and the second scenarios.

The root, when fresh, has a moderately strong, not disagreeable smell; as met with in the shops, it has scarcely any. To the taste it discovers a pretty strong, and very durable, bitterness; which is taken up both by watery and spirituous menstrua, and on inspissating the filtered tinctures, remains entire in the extracts: the aqueous extract is in much larger quantity than the spirituous, and proportionably weaker in taste. This root has been recommended as a stomachic, anthelmintic, and as an aperient in uterine obstructions; but is at present very rarely made use of. *Dr. Stærck has published some cases in which the root of white dittany, was given with good effect in epileptic sits, melancholy, intermittent severs, worms, and semale obstructions. It was given either in powder in the dose of a scruple or half a dram, or in tincture.

DICTAMNUS CRETICUS Pharm. Lond. & C. B. Origanum creticum latifolium tomentosum Tourn. Origanum Dictamnus Linn. Dittany of Crete: a small, shrubby, branched plant; with square stalks; roundish leaves about an inch in length, covered with a thick white down, set in pairs at the joints; and purplish labiated flowers, in loose scaly heads or spikes,

spikes, drooping downwards. It is perennial, a native of stony grounds in Greece and the island of Candy, and bears the ordinary winters of our own climate. The shops are generally supplied from Italy, with the leaves tied up in bundles, which are often damaged or decayed, and at best not superiour to those of our own growth: they have now and then pieces of the slowery heads among them, but oftener stalks, and different foreign matters.

THE leaves of dittany of Crete have been chiefly recommended as emmenagogue, alexipharmac, and vulnerary. They are apparently, when in perfection, a very warm aromatic; of an agreeable fmell, and a hot biting tafte, refembling that of the thymus citratus, but stronger and more pungent. They impart their virtues both to water and rectified spirit; and tinge the former of a yellowish, the latter of a green colour. Distilled with water, they give over a moderately strong impregnation to the aqueous fluid; from which, if the quantity of dittany be large, there feparates, as Neumann observes, a small portion of a yellowish essential oil, of a highly pungent aromatic tafte and smell, and which congeals in the cold into the appearance of camphor: the remaining decoction, inspissated, leaves a bitterish, disagreeable mass, totally divested of the warmth and flavour of the herb. Rectified spirit, distilled off from the tincture made in that menstruum, brings over little or nothing of the virtue of the dittany: the spirituous extract is a tolerably grateful and very hot pungent aromatic.

DIGITALIS.

DIGITALIS Ph. Parif. Digitalis purpurea Pharm. Edinb. & Linn. Digitalis purpurea folio aspero C. B. Foxglove: a somewhat hairy plant; with oblong, acuminated, serrated leaves; and a thick, angular, hollow stalk, on which numerous purple tubulous flowers, resembled to the singer of a glove, hang downwards, in a row along one side, each on a short pedicle: the flower is followed by an oblong pointed capsule, full of small angular seeds. It is biennial; grows wild in woods and on heaths; and puts forth in June or July its elegant flowers, which often continue a month or longer. It is observable of this plant, that it grows only

only on gravelly foils; rarely or never on those where there are strata of calcareous earths or stones underneath.

The leaves of foxglove have a bitterish very nauseous taste; which they communicate both to watery and spirituous menstrua. They have been strongly recommended in epileptic disorders: Parkinson relates, that after two or three sits had been suffered every month for twenty-six years, a cure was obtained by taking twice a week a decoction made in ale, of two handfuls of foxglove leaves with four ounces of polypody of the oak. The operation of this medicine, or of the foxglove by itself, is by stool and vomit; and appears, from the accounts given of it by authors, to be so violent, as to afford sufficient soundation for the present disuse of the plant (a). Externally, the leaves and slowers have been employed, with greater safety, and sometimes, as is said, with success, in cataplasms and unguents for ill-conditioned ulcers.

*DOLICHOS.

DOLICHOS Pharm. Edinb. Dolichos pruriens Linn. Cacara pruritus Rumph. Amb. Couhage, or Cow-itch. Cadjuet, Bengalis.

This is an herbaceous plant, of the papilionaceous tribe, growing in the East and West Indies. It bears pods, densely covered with sharp hairs, which have the property of penetrating the skin, and causing a most troublesome itching. This quality, with us, is only employed in performing mischievous tricks; but in the West Indies, the cow-itch is given internally as a very efficacious anthelmintic. The most particular account of the use of this remedy is contained in Mr. Bancrost's Hist. of Guiana, and it is confirmed by a letter in the Medical Comment. Vol. II. p. 82.

The

⁽a) Boerhaave judges it to be of a poisonous nature, and says it is so acrid as to exulcerate the mouth, fauces, essophagus and stomach. Hist. plant. Lugd. Bat. p. 308. & Haller Stirp. Helvet. p. 617. Dr. Alston, on the other hand, ranks it among those indigenous vegetables, "which, "though now disregarded, are medicines of great virtue, and scarcely inseriour to any that the "Indies afford." Index medicamentorum simplicium, prassat. p. 5. For my own part, I have had no experience of this plant used as a medicine; and, in regard to the taste, little can be judged, from that quality, of the virtues of substances of this kind: the taste of the digitalis is strong and nauseous, but not near so acrid or pungent as that of many vegetables which are taken with great safety.

The manner in which it is employed, is to mix the hairy matter scraped off from the pods, with syrup or melasses, into a thin electary, of which a tea-spoonful is given to a child two or three years old, and double the quantity to an adult. The dose is exhibited in the morning, fasting, for three successive days, after which a dose of rhubarb is given. Its effects are represented as remarkably powerful and certain, without the least dangerous consequence. The spiculæ seem by their mechanical action either to excite the peristaltic motion of the intestines, or to irritate and annoy the worms themselves. Neither a tincture nor decoction of the cow-itch were found to possess the least anthelmintic power.

A particular botanical description of the plant by Mr. Kerr is given in the Medical Comment. Vol. II. p. 202.

DORONICUM.

LEOPARDS BANE: a hairy plant, with uncut leaves, and yellow radiated discous flowers, which stand solitary upon long pedicles on the tops of the stalks and branches, and are followed by small seeds winged with down: the lower leaves have long pedicles, those which grow on the stalks have none. It is perennial.

1. DORONICUM sive Alisma & Arnica germanorum Pharm. Paris. Arnica Pharm. Edinb. Doronicum plantaginis folio alterum C. B. Arnica German leopards bane: with oval-pointed, ribbed montana Linn. leaves, like those of plantane, set in pairs upon the stalk; and oblong roots. It is a native of the mountainous parts of Germany, and flowers throughout the fummer.

THE leaves and flowers of this plant have a penetrating bitterish taste, and emit, when bruifed, a light pungent smell, which provokes sneezing. Both water and rectified spirit extract their virtues by infusion, and carry off a confiderable share of them in evaporation. The roots are more of an aromatic nature than the other parts, and their active matter fomewhat less volatile.

This plant has been greatly esteemed in different parts of Germany, as a specific for resolving coagulated blood occasioned by falls or bruises; from

Qq

from its efficacy in which intentions, it received the title of lapforum panacea: the dose is an infusion of one or two pugils of the leaves, or flowers, or both, and in some cases of the roots. It is said, that soon after the taking of this medicine, a great pain is felt about the affected part, and generally a cardialgic anxiety of the stomach, with nausea, reaching, and gripes, and fometimes extreme difficulty of breathing: that these symptoms are alleviated a little by walking about, and may be removed, if they should be very alarming, by venæsection: that in a short time, they terminate spontaneously in a copious discharge of urine, or in a profuse sweat, and sometimes in vomiting or purging (a). This herb is recommended likewise, in consequence of its supposed resolvent power, in fundry obstinate chronical disorders. It appears, however, to be much too violent in its operation for general use, unless repeated fmall doses should be accompanied with the good effects, without the disturbance, which a full dose is said to produce. Simon Paulli suspects, that it is made an ingredient in the malt liquors used in some places, by the common people, against bruises and other disorders.

* In a differtation on the virtues of the flowers of arnica, written by Dr. Collin of Vienna, a variety of cases are related of their good effects in paralytic disorders, amaurosis, and convulsive and spasmodic cases. An infusion or decoction of the flowers was used in the proportion of from one to four drams, to the pint of liquid. It generally occasioned pain in the affected parts, but no other disagreeable consequences of its use are mentioned.

2. Doronicum romanum: Doronicum radice scorpii C. B. Doronicum graphoy dictum J. B. Aconitum pardalianches minus quod falso doronicum vocant Matth. Doronicum pardalianches Linn. Roman leopards bane: with obtuse heart-shaped leaves set alternately on the stalk, and slender knotty roots, supposed to resemble the scorpion's tail. It is a native of the Alps, cultivated in some of our gardens, and slowers in June or July.

THE root of the Roman leopards bane has a sweetish somewhat astringent taste, accompanied with a weak aromatic slavour. Its medi-

cinal

⁽a) Tabernæmontanus, Herbar. lib. ii. seat. i. cap. xxii. p. 714. & seat. xi. p. 116. Fehr, Eph. nat. curios. dec. i. ann. ix & x. obs. 2. Gohl, Ad. med. Berolinens, vol. i. n. 4. p. 46. De la Marche, Dissert. de arnicæ veræ usu.

cinal qualities were formerly the subject of a considerable dispute; some affirming it to be poisonous, and others falutary. Geiner relates, that though it poisons dogs, he had himself many times eaten the herb as a grateful aromatic, and the roots also both fresh and dry, without perceiving any ill effect from them: that he had given the roots, with advantage, in vertiginous and epileptic cases: that nevertheless having once taken two drams of the powdered root, a fwelling of the belly and stomach succeeded in about eight hours, accompanied by a weakness of the whole body; that these symptoms continued for two days, and that he was cured by a warm bath(a). The Augustan college, after once expunging the doronicum from their pharmacopæia, have lately received it again, and affure us, that experience has declared it innocent. It does not however appear, its innocence admitted, to be poffessed of virtues fufficient to recommend it to practice. Among us, it has long been discarded, together with all the Arabian compositions, as an ingredient in which (from a prefumption, not perhaps very well grounded, of its being their duronegi) it was originally introduced.

DRACONTIUM.

DRACUNTIUM five Serpentaria Pharm. Parif. Dracunculus polyphyllus C. B. Arum Dracunculus Linn. DRAGONS: a plant with fmooth gloffy leaves, fet on long pedicles, divided into fix or feven or more long narrow fegments; and a fingle, thick, whitish stem, elegantly variegated with reddish or purplish streaks, composed as it were of membranes enveloping one another: on its top is a long sheath, greenish on the outside and purplish within, inclosing a dark-coloured pistil, like that of arum, but larger, succeeded by a cluster of red berries: the root is large, roundish, externally yellowish, internally white. It is perennial, a native of the southern parts of Europe, and cultivated in our gardens: it dies to the ground early in the autumn.

⁽a) Some have alledged, that the taking of this root occasioned his death; Casp. Hoffman de medicament. officinal. lib. ii. cap. iv. §. 8. Boerbaave, Hist. plant. Lugd. Bat. p. 151. but the epistle in which he mentions his being cured of the symptoms which the doronicum had occasioned, is dated above five months after; and Simler, his cotemporary, who gives a particular account of his death, informs us that he was carried off by an epidemic disease.

The dracontium appears to be fimilar, in medicinal virtues, as in botanic characters, to arum; the roots and leaves being, like those of that plant, extremely acrimonious, seeming, when slightly tasted, to burn or corrode the tongue, and continuing to painfully vellicate the part for many hours. The acrimonious matter is likewise of the same kind in regard to its pharmaceutical properties; being in great part dissipated or destroyed by exsiccation; not being dissoluble either by watery or spirituous menstrua; not rising with either menstruum in distillation, but being destroyed in the process; being extracted, by expression, along with the watery juice; but soon separating from the aqueous sluid, and being now found, though not a little weakened, in the secula or sediment. This plant might therefore be used in the same cases as arum, but general practice employs only the latter. So far as can be judged, between substances of such vehement pungency, the dracunculus is rather the strongest of the two.

ELATINE.

ELATINE Pharm. Lond. Elatine dioscoridis Lobel. adv. Linaria segetum nummulariæ solio villoso Tourn. Antirrhinum Elatine Linn. Flu-Ellin or Female speedwell: a low, procumbent plant; with oval, acuminated, downy leaves, set alternately on the stalks: from their bosoms issue long pedicles, bearing irregular, monopetalous, labiated, gaping flowers, with a crooked tail or spur behind, sollowed by roundish capsules full of small seeds: the upper lip is of a dark purplish colour, the lower lip and the spur yellow. It is annual, grows wild in corn fields, and slowers in July.

THE leaves of elatine have a roughish very bitter taste, and scarcely any smell. Both watery and spirituous menstrua extract their active matter by infusion, and leave it entire in evaporation. This herb was formerly accounted an excellent detergent and purifier of the blood. An extract made from it by water has been given in doses of a dram, and the expressed juice from three to five ounces, twice or thrice a day, and the juice applied also externally; against scorbutic disorders, and different kinds of old ulcers. A decoction of it has been used likewise in glisters for alvine fluxes. A combination of its active matter with honey, prepared

pared by boiling four pints of the depurated juice with four pounds of Mel elatines clarified honey, is fometimes kept in the shops; but neither this preparation, nor the herb in any form, are at present much made use of.

E L E M I.

GUMMI ELEMI Pharm. Lond. ELEMI: a concrete refinous juice, faid to be obtained from a large tree refembling an olive; (Amyris elemifera Linn.) brought from the Spanish West Indies, and sometimes from the East Indies, in oblong roundish cakes, generally wrapt up in flag leaves. The best fort is softish, somewhat transparent, of a pale whitish yellow colour, inclining a little to greenish. The faculty of Paris mentions a spurious elemi, or gummi chibou, which is not yet known among us.

ELEMI has a strong, tolerably pleasant smell; and a slight bitterish tafte. It gives out very little to aqueous menstrua, but almost totally diffolves in rectified spirit, tinging the fluid of a pale gold colour. Distilled with water, it yields a thin pale coloured essential oil, amounting to about one ounce from fixteen, of a moderately pungent tafte, and finelling strongly of the elemi: a friable inodorous refin remains behind in the still. On submitting to distillation the solution made in rectified spirit, a little of the fragrance of the elemi arises with the spirit, greatest part remaining in the inspissated mass, which has a considerable share of the smell, though it makes little impression on the organs of taste.

This refin is scarcely otherwise employed among us, than as an ingredient in digestive ointments; one of the best of the officinal digestives, commonly called the ointment or liniment of Arcæus, confifts of fix parts of the elemi, five or fix of turpentine, and twelve of lard or of a mixture of lard and fuet, melted together. This refin should nevertheless seem applicable to other purposes, and to be preferable, for internal use, to some refinous substances that have been held in greater esteem.

ELEUTHERIA.

ELEUTHERIÆ CORTEX Pharm. Lond. Cafcarilla Pharm. Edinb. Thus judæorum Park. Cortex thuris nonnullis dictus, vel thymiama, miama, vel thus judæorum Raii hift. ELEUTHERIA or CASCARILLA: the bark, probably of the shrub described and figured by Catesby under the name of ricinoides elæagni folio or ilathera, Croton Cascarilla Linn. which grows plentifully in most of the Bahama islands (a). From those islands, particularly, as it is said, from one of them called Elatheria, it is immediately brought to us; in curled pieces, or rolled up into short quills, about an inch in width: covered on the outside with a rough whitish matter, and brownish on the inner side; exhibiting, when broken, a smooth, close, blackish brown surface.

This bark, freed from the outer whitish coat which is insipid and inodorous, has a light agreeable smell, and a moderately bitter taste, accompanied with a considerable aromatic warmth. It is easily in-slammable, and yields, whilst burning, a very fragrant smell, somewhat resembling that of musk; a property which distinguishes the eleutheria from all other known barks (b).

Stiffer appears to have been the first who employed the cortex eleutheriæ as a medicine in Europe. He relates, that he received this aromatic bark from England; and that some time after, it was sold at Brunswick for Peruvian bark: that a tincture of it in alkalized vinous spirits, or in dulcified alkaline spirits, proved carminative and diuretic, and did considerable service in arthritic and scorbutic cases; and that if taken immediately after meals, it affected the head a little (c).

Eleutheria was foon after employed by Apinus, in an epidemic fever of the intermittent kind, which raged in some parts of Norway in 1694 and 1695. This disease, which at first had the appearance of an ordinary intermittent, was at length accompanied with petechial spots. The

⁽a) Essay towards a natural history of Carolina, Florida, and the Bahama islands.

⁽b) This property seems to confirm the above account; that eleutheria (not of the growth of the East Indies as some have supposed, nor of the Spanish West Indies as others) is really the produce of the Bahama ricinoides of Catesby; whose bark, he says, insused either in wine or water, gives a fine aromatic bitter, and being burnt yields a fine persume. Those, who imagine the eleutheria to be the bark of a Peruvian tree, seem to have been missed by the name cascarilla; which is applied by the Spaniards to the Peruvian bark strictly so called, and signifies no more than bark in general. See Hossman's Dissertatio de cascarilla, anno 1738. Operum omnium supplement. ii. par. i. p. 704.

⁽c) Acta laboratorii chymici, Specim. ii. cap. ix & x. edit. ann. 1693. & de febribus intermittentibus consult. nov. cap. xvi.

common alexipharmacs and sudorifics were found ineffectual: but the powder or extract of eleutheria, joined with them, proved successful, even after petechiæ had appeared: dysenteries, succeeding the sever, were removed by the same means. During the use of the eleutheria, the patient generally sweated plentifully, without loss of strength or other inconvenience: the belly was at the same time kept open, and those who did not sweat had commonly three or sour stools a day: where the menstrual or hæmorrhoidal sluxes were suppressed at the beginning of the disorder, they generally, upon the use of this medicine, reappeared (a).

The gentlemen of the French academy found this bark of excellent fervice against an epidemic dysentery in the year 1719; in which, ipecacoanha proved ineffectual. Mr. Boulduc observes, that this last left a lowness of the spirits, and weakness of the stomach, which continued for a long time: whereas the eleutheria soon raised the strength,

and promoted appetite (b).

At prefent, eleutheria is in great esteem among the Germans, as a warm stomachic and corroborant, in statulent colics, internal hæmor-rhagies, dysenteries, the diarrhææ of acute severs, and in common intermittents; in which last it is often joined to the Peruvian bark, and by many preferred to it, as being less subject to some inconveniences, which the other, by its great astringency, is apt to produce. Among us, it has but lately been received into practice; and its use is not yet become so general as it well deserves to be.

The virtues of eleutheria are partially extracted by water, and totally by rectified spirit: after the action of the former, it retains a considerable share of its slavour, after the latter it proves inodorous and insipid: the watery tinctures are of a reddish brown, the spirituous of a brownish orange colour. Distilled with water, it yields a greenish essential oil, of a very pungent taste, and of a fragrant penetrating smell, more grateful than that of the cascarilla itself, in quantity, according to Hossman's experiments, not exceeding one dram from sixteen ounces: the decoction, inspissated, leaves an extract of a moderate dull bitterness, much weaker than might have been expected from the strong taste of the bark in sub-

⁽a) Historica relatio febris epidemica, edita anno 1697.

⁽b) Histoire de l'acad. royale des sciences, pour l'ann. 1719.

stance. On inspissating the spirituous tincture, with a gentle heat, nothing considerable of the active matter of the cascarilla was found to arise with the menstruum: the remaining extract, nevertheless, was rather weaker in taste than the bark itself, and when thoroughly exsictated, scarcely discovered any taste at all, being almost a pure resin, not dissoluble by the saliva. It is probably the dry pulverable extract that Cartheuser means, when he says it has no taste; and the extract in its moist state that was examined by Boulduc, who says it is bitter, biting, and aromatic.

ELICHRYSUM.

ELICHRYSUM five Stæchas citrina Pharm. Parif. Elichrysum sive stæchas citrina angustifolia C. B. Gnaphalium Stæchas Linn. Goldiocks: a small, shrubby, downy plant; cloathed with long, very narrow leaves: producing, on the tops of the branches, several small round heads of bright yellow scaly flowers. It is a native of the southern parts of France, slowers in our gardens in May and June, and holds its leaves all the winter.

THE flowers or scaly heads of this plant, naturally dry and firm, retain their figure and glossy yellow colour for years. Both the flowers and leaves, rubbed a little, yield an agreeable and moderately strong smell: to the taste they discover a considerable warmth, pungency, and bitterishness: from whence it may be presumed, that the aperient and corroborant virtues, commonly ascribed to them, are not wholly without foundation.

ENDIVIA.

ENDIVIA Pharm. Edinb. (a) Intybus fativa latifolia five endivia vulgaris C. B. Cichorium endivia Linn. Endive: a common culinary plant, resembling cichory in the flowers and seeds; and differing from it chiefly in being annual, and in the leaves being shorter, broader, and only slightly crenated, not jagged, about the edges.

ENDIVE agrees with cichory in quality as in appearance; containing, like it, a milky juice, of a bitterish taste, which it loses on being blanched by culture: the greener the colour of the leaves, the greater is the bitterness of the whole plant. In its bitterest state, however, it is somewhat less so than the cichory.

ENULA CAMPANA.

ENULA CAMPANA Pharm. Lond. Enula campana seu belenium Pharm. Edinb. Inula Gesn. bort. Helenium vulgare C. B. Aster omnium maximus Tourn. Inula Helenium Linn. ELECAMPANE: a large plant, with long, wrinkled, oval, acuminated leaves, serrated about the edges, pale green above, and hoary underneath, joined close to the stalk, which divides towards the top into several branches, bearing large yellow flowers of the radiated discous kind, followed by oblong seeds winged with down: the roots are short and thick, somewhat unctuous to the touch, brown or blackish on the outside, and whitish within. It is perennial, grows wild in moist rich soils, and slowers in June.

The fresh roots of elecampane have a weak not very grateful smell; which, on thoroughly drying and keeping them for some time, is greatly improved, and approaches to that of Florence orris. Chewed, they discover at first a kind of rancid glutinous taste, quickly succeeded by an aromatic bitterness, which by degrees becomes considerably pungent.

This root stands recommended as a diaphoretic and stomachic, for promoting expectoration in humoural asthmas and coughs, and for attenuating viscid juices in general, and disposing them to excretion: taken freely, it is said to gently loosen the belly, and increase the urinary discharge. The dose of the dry root in substance is from a scruple to a dram or two.

It gives out its virtue partially to aqueous, totally to spirituous menstrua: the former it tinges of a muddy yellowish, the latter of a bright
pale yellow colour. In distillation with water, it gives over an essential
oil, which concretes into white slakes, partly swimming on the water,
and partly subsiding, in quantity about one dram from thirty ounces,
of no great heat or pungency, smelling at first pretty strongly of the
elecampane, but very apt to lose its smell in keeping. Great part of the

R r

Extractum enulæ camp. Ph. Lond. aromatic warmth and pungency, as well as the bitterness, reside in a matter of a more fixt kind, which does not easily exhale in the heat of boiling water, and is preserved in tolerable perfection in the watery extract. Rectified spirit elevates little or nothing from this root: the spirituous extract is considerably stronger than the watery, though its pungency is not near so great as might be expected considering the small-ness of its quantity: it scarcely exceeds one sisteenth the weight of the root, whereas the watery extract amounts to almost one half.

EPITHYMUM.

EPITHYMUM Ph. Parif. Epithymum five cuscuta minor C. B. Cuscuta Epithymum Linn. Dodder of theme: a plant without leaves, growing on theme, consisting of a number of slender juicy filaments, producing here and there small heads of white or reddish flowers, which are followed by roundish capsules sull of minute seeds. Dodder receives its nourishment from the vegetable on which it climbs, its own roots quickly perishing. A large kind, vulgarly called hellweed, is common in heaths, upon surzes, nettles, &c. and in fields of flax and other manured herbs. The smaller fort, sound upon theme, has been generally preferred for medicinal use, and imported to us from Turkey and Leghorn, intermixed with stalks and tops of theme. It is supposed by some, that the dodder partakes of the qualities of the plant by which it is supported.

Dodder of thyme has a pretty strong not disagreeable smell, and a peculiar kind of subtile pungent taste, very durable in the mouth, and sinking as it were into the tongue. Though it was early received into medicine, its medicinal qualities are not as yet known. The ancients accounted it cathartic, but when given by itself it is found to have very little purgative virtue. Some late writers recommend it rather as a de-obstruent, in melancholic and other disorders. It is in this country an entire stranger to practice; though the remarkable subtilty of its taste seems to promise some considerable medicinal power.

EQUISETUM.

CAUDA equina five Equifetum Ph. Edinb.(a) Equifetum palustre longioribus setis C. B. Hippuris vulgaris Linn. HORSETAIL: a plant, with a thick hollow straight stalk, full of joints; and long, slender, rough, stiff, jointed, rush-like leaves, standing several round every articulation in form of a star. It is perennial, and common in watery places.

The leaves of equifetum have been accounted powerfully astringent, and hence recommended in different fluxes and hemorrhagies: the dose commonly directed is a dram of the dry leaves in powder, or two ounces of the expressed juice. They appear, indeed, from their sensible qualities, to have some virtues of this kind, but in a very low degree: in the leaves themselves, the astringency is so weak, that the taste scarcely gives any notice of it, though solution of chalybeate vitriol renders it apparent by the inky blackness it receives from them: the astringent matter is extracted both by watery and spirituous menstrua, and when concentrated by inspissating the insusions, proves still of so weak a kind, and in so little quantity, that the plant can be ranked only among the milder restringent corroborants. In this intention, an insusion of the dried herb may be used as tea.

ERIGERUM.

ERIGERUM five Senecio Pharm. Edinb. (a) Senecio minor vulgaris C. B. Senecio vulgaris Linn. GROUNDSEL: a low, branched plant, with oblong narrow leaves, deeply cut into wing-like sections, joined to the stalk by broad bases: on the tops grow numerous yellow slowers, of the naked discous kind, set in large cups, and sollowed by small seeds winged with down. It is a common annual weed in dry grounds.

THE leaves of groundfel have an herbaceous fomewhat faline tafte, and no remarkable fmell. They stand recommended, externally, as a vulnerary and refrigerant, internally as a mild and safe emetic. The

(a) Expunged.

expressed juice, or an infusion or decoction of the herb, are by several writers directed in this last intention, but neither of them appear to have any considerable effect: perhaps those, who ascribe to them an emetic power, were imposed on by giving such large doses, as to nauseate the stomach by the quantity of sluid.

ERUCA.

ROCKET: a plant with smooth oblong narrow leaves deeply jagged about the edges, bearing on the tops of the stalks numerous tetrapetalous slowers, which are followed by angular pods, full of small roundish seeds flatted on one side. It is annual.

- I. ERUCA: eruca latifolia alba, sativa dioscoridis C. B. Brassica Eruca Linn. Garden or Roman rocket, or rocket gentle: with leaves like those of turneps, but much smaller; and whitish flowers variegated with black streaks. It is a native of Switzerland, and cultivated among us in gardens.
- 2. ERUCA SILVESTRIS: eruca sylvestris major lutea caule aspero C. B. Brassica Erucastrum Linn. Wild rocket: with leaves like those of dandelion, and yellow flowers; common on old walls, and among rubbish.

THE leaves of both the rockets have an acrid taste like that of cresses, and a rank disagreeable smell: the roots are as acrid as the leaves: the seeds much more so, approaching to the pungency of mustard. The wild fort is, in all its parts, considerably more acrid than the garden, though the faculty of Paris allows both forts to be taken indiscriminately. They are accounted good aperients and antiscorbutics, but are now rarely made use of on account of their ill slavour: to the aphrodisac virtues, commonly ascribed to them, they appear to have no other title than the rest of the pungent stimulating plants.

The active matter of the leaves is extracted by expression, by infusion in boiling water, and by digestion or maceration in rectified spirit; with this difference, that the infusions and tinctures retain the ill smell of the herb, which in expression is in great measure destroyed. On drying the herb itself, or inspissating the juice of the watery or spirituous tinctures,

the

the pungency, as well as the fmell, is almost totally distipated. In distillation with water, a very small quantity of a yellowish, very pungent, and very volatile essential oil is obtained.

The pungency of the feeds is of a less volatile kind; not exhaling in exficcation, and arising more difficultly with water in distillation: and though it appears to reside, as that of the leaves, in an essential oil, it is but partially extracted by rectified spirit. In these respects the seeds of rocket agree with those of mustard; to which they appear to be similar also, but inferiour, in medicinal virtue.

ERYNGIUM.

ERYNGIUM Pharm. Lond. Eryngium maritimum C. B. & Linn. Eryngo or Sea holly: a blueish, branched, umbelliserous plant, with mallow-like, thick, prickly leaves, angular or jagged about the edges: the flowers are white and set in prickly heads, under which a number of little oblong leaves stand in form of a star: the roots are slender, very long, with a sew knots, brownish on the outside, and white within. It is perennial, grows plentifully on some of our sandy and gravelly shores, and slowers in July.

ERYNGO roots have an agreeable sweetish taste, which on chewing them for some time is followed by a light aromatic warmth and pungency. They are accounted aperient, diuretic, and aphrodisiac: Boerhaave says they are the principal of the aperient diuretic roots, and that he constantly made them an ingredient in his prescriptions against scurvies: their virtues, however, appear to be but weak, and they are now scarcely otherwise used than as made into a sweetmeat.

ERYSIMUM.

ERYSIMUM Pharm. Edinb. (a) Erysimum vulgare C. B. Irio sive erysimum dioscoridis Lob. Erysimum officinale Linn. HEDGE MUSTARD: a hairy plant, with oblong narrow leaves, divided into wing-like sections, triangular at the extremity; and tough, branched stalks, bearing nu-

merous small, yellow, tetrapetalous flowers, which are followed by short roundish pods, standing close to the stalks, full of small reddish brown seeds. It is annual, common in waste places, and flowers in July.

The leaves of eryfimum are faid to be attenuant, expectorant, and diuretic; and stand particularly recommended against chronical coughs, and hoarseness, whether humoural, or occasioned by immoderate exertion of the voice. Lobel greatly commends for this purpose a compound syrup, which, as Geoffroy observes, is not superiour to a simple mixture of the expressed juice of the herb with honey; and indeed it is not very clear whether the virtue of the honey is much improved by the erysimum. The herb has no smell, and its taste, at least when moderately dried, is little other than herbaceous, with somewhat of a slight faline impregnation.

The feeds of eryfimum are confiderably pungent, and appear to be nearly of the same quality with those of mustard, but weaker. Their acrimony, like that of mustard-seed, is extracted totally by water, and partially by rectified spirit, and strongly impregnates water in distillation.

EUPATORIUM.

EUPATORIUM arabum Pharm. Parif. Eupatorium cannabinum C. B. & Linn. Hemp agrimony: a plant with oblong, acuminated, deeply indented leaves, set three on one pedicle, and the pedicles in pairs: the flowers, which stand in umbel-like clusters, consist of purplish flosculi set in scaly cups, followed by oblong seeds winged with down. It is perennial, grows wild by the sides of rivers and ditches, and flowers in July.

The leaves of eupatorium have a light agreeable smell, and a pungent very bitter taste. They are recommended as aperients, laxatives, and corroborants; in beginning dropsies, jaundices, intermitting severs, and other consequences of obstructions of the viscera, succeeding frequent relapses into acute, or a long continuance of chronical diseases. They are said to be the common medicine of the turf-diggers in Holland, against the scurvies, soul ulcers, and swellings of the feet, to which

they

they are subject. Insusions of the herb may be drank as tea, or the expressed juice taken in doses of one, two, or three ounces; in large quantity, it purges or vomits.

EUPHORBIUM.

EUPHORBIUM Pharm. Edinb. (a) EUPHORBIUM: a gummy refinous concrete juice; exuding from an oriental, prickly, lactescent shrub, of the same name, euphorbia (officinarum) aculeata nuda multangularis: aculeis geminatis Linn. spec. plant. The juice is brought immediately from Barbary, in drops or tears of an irregular form; some of which are found, on being broken, to contain little thorns, twigs, slowers, and other vegetable matters; others are hollow, without any thing in the cavity: the tears are in general easy to break, of a pale yellow and sometimes of a gold colour on the outside, and white within.

EUPHORBIUM, applied lightly to the tongue, discovers a sharp biting taste: held for some time in the mouth, it proves vehemently acrimonious, inflaming and exulcerating the parts. The finer dust which slies off in pulverization, unless great care is taken to avoid it, is apt to affect the operator's head and throat in a violent manner.

It consists of about equal parts of resinous and gummy matter. The acrimony resides chiefly in the former; the spirituous tincture being excessively siery, and when inspissated still more so; whereas the watery insusand extract are bitterish with only a slight, though a very durable acrimony. A single drop of a strong spirituous tincture produces in the mouth a sensation of burning, which, as Cartheuser observes, is scarcely to be obtunded by mucilages or oils in less than an hour. It gives over nothing in distillation either to water or spirit.

The extreme acrimony of this drug renders it absolutely unfit for any internal use: several correctors have indeed been proposed for abating its virulence, but the best of them are not to be trusted to. It is employed only, and that but seldom, for external purposes; in stimulating unguents and plasters for paralytic limbs, carious bones, &c. Some have ventured on a minute portion of it, mixed with other powders, as an errhine, in obstructions of the nostrils, and mucous disorders of the head; a practice

by no means adviseable, as we are in no want of medicines for these purposes, equally effectual, and far more safe. I have seen violent and dangerous inflammations produced by rashly using even very small quantities of it mixed with snuffs.

EUPHRASIA.

EUPHRASIA Pharm. Edinb. (a) Euphrafia officinarum C. B. & Linn. EYEBRIGHT: a small herb, with little, oval, serrated leaves, set in pairs, without pedicles: in their bosoms, towards the tops of the stalks, come forth labiated, monopetalous, whitish flowers, streaked internally with purple and yellow; followed each by a flattish capsule, full of small whitish seeds. It is annual, grows wild in uncultivated grounds, and flowers from July to September.

This plant has long been celebrated as an ophthalmic, both taken internally, and applied externally. Hildanus fays, he has known aged people, who had their fight impaired, recover it again by the use of this herb; but later practitioners have not been so happy as to meet with the like success. It may indeed, in some cases, be of service as a mild corroborant; for it discovers an astringent quality to the taste, and, in a more sensible manner, by striking a black colour with solution of chalybeate vitriol. The astringent matter is extracted both by water and spirit; and when concentrated by inspissating the tinctures, is still found to be very weak.

F E L.

FEL five Bilis. GALL or BILE: a bitter animal juice, secreted from the blood in the liver, and collected in a particular receptacle. The galls of the ox, the eel, and the pike fish, have been chiefly made use of in medicine.

This fluid mingles uniformly with water, spirit of wine, fixt alkaline lixivia, and volatile alkaline spirits, without change of its yellow colour. The concentrated mineral acids coagulate and render it whitish: diluted

acids, those at least of the vegetable and animal kingdom, change it green: the addition of alkalies to the green mixtures restores the natural vellow colour of the bile. Inspissated by heat, it dissolves almost totally in water, but is more sparingly acted upon by rectified spirit (a). It renders oily, unctuous, and refinous fubstances miscible with watery liquors; preferves milk from coagulating or turning four, and rediffolves it when already coagulated. Such at least are the properties of the gall of the ox: how far that of other animals agrees with or differs from it

in these respects, is not known (b).

This stimulating resolvent bitter has been given, and as is said with good fuccess, for opening obstructions of the viscera, promoting urine, the menses, and labour pains: in this last intention, the gall of the eel, which is faid to be one of the most acrid, has been chiefly recommended. Boerhaave relates, that he has cured pale ricketty children by pills made of the galls of the eel and the pike; that the medicine operated powerfully by urine; and that, by its use, the belly, before swelled, subsided furprizingly (c). In want of appetite and other complaints proceeding from a deficiency of bile in the first passages, this animal bitter may probably be of more fervice, than those of the vegetable kingdom usually directed in fuch intentions. Among us these fluids are employed only for external purposes: a mixture of ox gall with camphorated spirit of wine is faid to be an useful embrocation for sprains, bruises, rheumatic pains, &c.

FERRUM.

IRON: a greyish, hard metal, between seven and eight times specifically heavier than water: distinguished from all other metallic bodies, in its metallic state, by its attracting, or being attracted by, the loadstone; but losing this attractive power on being reduced, by fire or menstrua, to a calx: not fusible without an intense white heat, and calcining, by a continuance of a weaker heat, first into blackish scales, and afterwards into a dark reddish powder: corrosible by moist air into a reddish yellow rust, and soluble in all acids, from which it precipitates

⁽a) Cartheuser, Fundamenta mat. med. feet. ix. cap. i. Edit. Parif. tom. i. p. 519.

⁽b) Vide Baglivi, Differtationes variæ, diff. iii. de experiment. circa bilem. Opp. p. 428, fegg.

⁽c) Praxis medica, tom. i. p. 164.

all the common metallic bodies except zinc: forming with the marine acid a yellow, with the nitrous a dark red, and with the vitriolic a pale green folution; which is changed to an inky blackness by the addition of a little galls, and by most of the other vegetable astringents, and to a reddish or purple by a mixture of astringents with a minute proportion of any alkaline salt. All the solutions, by whatever acid effected, on the addition of a lixivium of alkaline salts that have been calcined and fully impregnated with animal coals (a), are changed to a deep blue, and on standing deposite the iron in form of a powder of the same colour. By these characters, a most minute portion of iron may be discovered in liquors.

Cæruleum berolinenfe; Pruffian blue.

Ores of iron, and minerals more flightly impregnated with it than those which are strictly called ores, are common in most parts of the world: the red and yellow earths and stones generally owe their colour to an admixture of this metal. The iron, extracted from the ore by fusion in large surnaces in mixture with the suel, is impure and brittle: being again laid on burning charcoal in a smaller surnace or forge, and melted down, a quantity of sulphureous scoria separates, the iron proves less suspenses tough enough to be forged into bars. The iron thus purified is employed as an article of the materia medica in two states.

- I. FERRUM Pharm. Lond. Ferrum five Mars Pharm. Edinb. Iron, or forged iron: iron in its foftest state; capable of being easily filed; acquiring little or no additional hardness on being made red hot and quenched in water; appearing, when broken, of a fibrous texture; exceeding difficult of susion, and perhaps not susible at all by common fire without the contact of the suel or other additions.
- 2. CHALYBS Pharm. Lond. Steel: iron in a hard state, so as to resist the file, or acquiring this hardness by heating and quenching it; when broken, of a fine granulated texture; much easier of susion, some-

⁽a) Volatile alkaline spirits may be completely saturated with the matter which tinges dissolved iron blue, by digesting them with the pigment called Prussian blue; from which they acquire a yellowish or greenish tincture, leaving the iron in form of ochre or a rusty coloured calx.

what more difficult of folution, and fomewhat less subject to rust in the air, than soft iron. Iron, cemented in close vessels, with vegetable or animal coals, becomes steel; and steel, kept red hot for some time in an open vessel, becomes soft iron again.

This metal, when dissolved, discovers a strong austere corrugating taste, and contracts and hardens all the vascular and soft fibrous parts of animals. To constringe and corroborate the animal solids appears to be its primary medical operation.

In weak, lax, pale habits, and in chronical diforders proceeding from languor and debility, cachectic, hypochondriacal, and others, this metal has generally good effects: strengthening the stomach, and chylopoietic organs, and the system in general; quickening the circulation and raising the pulse; rendering the blood more florid, and as it were expanding and rarefying the juices; promoting, when they are deficient, and restraining, when immoderate, the secretions that are made from the blood, as perspiration, urine, and the uterine purgations; but for the most part binding the belly, though this evacuation also, in some circumstances, it promotes.

By the same corroborating power, whereby it promotes deficient and restrains redundant discharges where the suppression or slux arise from debility and relaxation; it, contrariwise, increases sluxes and confirms obstructions when they proceed from tension, rigidity, or spasmodic strictures of the vessels. Where either the circulation is quick, or the habit plethoric; by increasing the blood's velocity, and all the plethoric symptoms, it produces heaviness, dulness, vague heats and slushings, or kindles more dangerous severs or inflammations, or bursts some of the over-distended small vessels.

In some constitutions, even where iron is proper and salutary, particularly in hysterical and hypochondriacal cases, and where the stomach is very weak, it is apt at first to occasion great sickness and perturbation: Sydenham observes, that these inconveniences may be, in some measure, prevented by beginning with very small doses, and giving it for a while only at bed-time, in conjunction with a slight opiate (a). In other circumstances, it is commonly taken in the morning and afternoon,

⁽a) Dissertatio epistolaris de variol. constuent. & affect. hysteric. Oper. p. 409.

and moderate exercise used to promote its action. The dose in all cases should be small, and rather repeated than enlarged: a grain or half a grain of the metal, disfolved or in a soluble state, is generally a sufficient dose. Nidorous eructations, and the alvine feces being tinged of a black

colour, are marks of its taking effect.

Iron is fometimes given in fubflance, reduced into fine filings; which answer, in many cases, as well as its most elaborate preparations; but their action is less certain, as depending upon their meeting, in the first passages, with juices capable of dissolving them: they are likewise the most subject, when they do act, to produce troublesome eructations, probably from the property of this metal of yielding copious fetid vapours during its diffolution. The dose of the filings is from two or three grains to a scruple and more: it is probable that the whole quantity taken does not prove operative, even when the stomach abounds most with acidities; for on digefting a scruple of the filings in a quarter of a pint or more of strong vinegar, a very considerable proportion remains undiffolved.

Ferri limatura purifi-cata Ph. Ed.

Iron filings, procured from the common work-shops, may be cleanfed from earthy matters or fragments of other metallic bodies, by means of a magnet, though not so perfectly as could be wished. When other metals have been previously melted with the iron, the filings of the compound cannot thus be separated from those of pure iron, the loadstone attracting both: regulus of antimony is the only exception, this metal being found, even in a fmall proportion, to destroy the magnetic power of the iron.

The filings are fometimes candied with fugar; a preparation which is very commodious for taking, but which requires a good deal of address in the operator, and is made chiefly by the confectioners. Two parts of fine fugar, diffolved in water and boiled down to a candy confiftence, are added, by little and little, to one part of the cleanfed filings, in a kettle suspended over a very gentle fire; and the vessel continually shaken, that the filings may be crusted over with the sugar: to render the matter less subject to run into lumps, a little starch is previously mixed with the fugar, in the proportion of a dram to a pound.

The filings, moistened with vinegar or water, and exposed to a moist air, or occasionally moistened afresh, soon change in great part into rust, which may be separated from the uncorroded part, by grinding and

washing

bigo, vulgo ferri limatura præpa-rata Pb. Ed.

washing over the finer powder with water. The rust is given in the Chalyis fame dose as the crude filings, and is perhaps rather easier of solution. Hoffman fays, he has often used it, with remarkable success, in obstinate Lond. chlorotic cases accompanied with excessive headachs and other alarming fymptoms; and that he usually joined with it pimpinella, arum root, and falt of tartar, with a little cinnamon and fugar.

A piece of steel, heated in a very fierce fire, as that of a smith's forge, to a strong white heat, and immediately applied to a roll of brimstone held over a vessel of water, is in part corroded by the sulphur, and, melting, falls down in brown coloured drops; which, picked out from Chalybs the yellow strings of sulphur, and levigated into an impalpable powder, phure præare given in the same doses as the filings and rust, and nearly with the Paratus Ph. fame effects.

As this process is somewhat troublesome and accompanied with offensive fumes, the shops have been generally supplied with a sulphurated preparation made more commodiously, by mixing iron filings with twice their weight of flowers of brimstone, and as much water as will make them into a paste, which on standing at rest for some hours swells up, and is then pulverized, put into a heated crucible to deflagrate, and kept constantly stirring with an iron spatula till it falls into a deep black powder.

This powder, urged longer in the fire, becomes red, and in this state has been usually distinguished in the shops by the name of aperient crocus: when further reverberated with a very intense heat, it is called aftringent crocus. This notion, of opposite virtues in the two preparations, does not appear to have any just foundation; chalybeate medicines in general acting by an aftringent power, though with different degrees The college of Edinburgh allows colcothar of vitriol as a Colcothar substitute both to the aperient and the astringent crocus; and indeed it appears to be at bottom, if duly prepared, the very fame thing with them: all the three are no other than iron, that has been corroded by the fulphureous or vitriolic acid, and afterwards by fire divested of greatest part of the acid, and reduced to a state of calx: the colcothar, however, as remaining after the distillation of the vitriol, commonly retains much more acid than the others, a circumstance to be attended to in the substitution of it. In all these kinds of preparations, only a small quantity of the metal is in a foluble or active state, more or less according to the proportion

proportion of acid: when iron is perfectly calcined, and no acid com-

bined with it, it has fearcely any fenfible operation.

Oil of vitriol, diluted with from equal to five or fix times its measure. or more, of water, and affifted by a gentle heat, acts readily on iron, and emits, during its action, a strong sulphureous vapour, which on the approach of any flaming body, catches fire and explodes, fo as fometimes to burst the vessel, especially if its mouth is narrow. The solution filtered, and evaporated till a pellicle appears on the furface, yields, on standing in the cold, green crystals, the same with the common green vitriol. To four parts of oil of vitriol fome direct three of the iron filings+; others, that the acid may prevail a little, and the crystals be

less subject to grow rusty coloured in the air, only two !.

+ Vitriolum martis, feu fal chalybis Pb. Edinb. t Sal martis Ph. Lond.

on macerating half a pound of iron filings in three pounds of spirit of falt till the acid ceases to act, a notable quantity remains at last undissolved. The folution is excessively styptic, far more so than the combinations of iron with any other acid: it has likewise this peculiarity, that it mingles equally with, and when inspissated dissolves in, rectified spirit of wine; on which foundation the spirituous chalybeate tinctures depend. Some inspissate the marine solution, made in the quantity above mentioned, to the weight of a pound, and then add three pints of rectified spirit: others dissolve three ounces of clean iron scales in a sufficient quantity of the

acid, and then add so much rectified spirit as to make the weight of the whole, two pounds and a half. Of these tinctures, a few drops are a

The marine acid diffolves much less of this metal than the vitriolic:

Tinet. martis in spiritu falis Ph. Lond.

Tinet. martis Ph. Edin.

fufficient dose.

Flores martiales vulgo Ens Veneris Pb. Edinb.

Flores martiales Pb. Lond.

On grinding iron filings, or washed colcothar of green vitriol, with equal or twice their weight of fal ammoniac, moistening the mixture with water, gently drying it, and repeating the pulverization, humectation, and exficcation, a few times; the iron is in a confiderable degree attenuated, and on fublimation with a quick fire, fo much of it arises with the falt as to communicate a deep yellow or orange colour. If the iron and fal ammoniac be only mixed together, and the fublimation performed with a flow fire, fuch as a glass retort will bear, the flowers prove at first pale, and require, in order to their being fufficiently tinged with the metal, to be ground with the refiduum, fublimed again, and this process repeated. These flowers have a very pungent austere taste, and are supposed to be more aperient and attenuating than the other chalybeates,

by virtue of the faline matter joined to the iron. They are most conveniently given in the form of a bolus, from three or four grains to twenty: they occasion pills to swell and crumble, except such as are composed of gummy refins: in a liquid form they are nauseous, except in spirituous tinctures. A tincture made by digesting four ounces of the Tinct. floflowers in a pint of proof spirit, is a sufficiently elegant chalybeate, and rum martimay be given in doses of a tea-spoonful.

The matter which remains after the fublimation of the flowers, exposed to a moift air, runs into a liquid, in tafte extremely flyptic, and greatly Lixivium refembling a faturated folution of iron made in spirit of falt; the marine martis Pb. acid and volatile alkali of the fal ammoniac being in part separated from one another in the process.

Solutions of iron in vegetable acids are much more mild, and lefs ungrateful both to the palate and stomach, than such as are made in the acids of the mineral kingdom. Vinegar, juices of oranges, lemons, apples, and other fruits, acidulous wines, and tartar, have been made use of for this purpose. A vinous tincture is prepared in the shops, by macerating three ounces of iron filings, for four weeks, in three pints Vin. chalyb. of Rhenish; to which a slight aromatic impregnation is given, by three drams of cinnamon and the same quantity of mace. The dose of the tincture is from a tea-spoonful to a common spoonful and upwards. For making these kinds of preparations, fine iron wire, cut in pieces, is more eligible than the filings, as we may always depend on the wire being pure iron, and as, by lying loofer, and exposing a larger surface to the fluid, it is more easily acted on.

Some direct folutions of iron made in wine, or other vegetable acids, to be inspissated to the consistence of an extract. These kinds of pre- Extract. parations are commodious for fome purposes, particularly for being made into pills; as being tenacious enough to give a due confiftence to a confiderable admixture of powdery matters. They are most of them very apt to grow mouldy in keeping; an extract made with the juice of golden rennets is faid by Neumann to be free from this inconvenience.

A combination of iron with the acid of tartar is most commodiously obtained, by grinding the filings with equal their weight of crystals of Mars solub. tartar, forming the mixture into a mass with water, then pulverizing, seu chalybs tartarizatus. and repeating the humectation and exficcation alternately, till the whole falls into an impalpable powder. This is a very elegant and useful chalybeate,

chalybeate, the tartar rendering almost all the iron dissoluble. It is given either in a solid or liquid form, from two or three grains, to ten or more. It has been usually distinguished in the shops by the name of its inventor Dr. Willis.

If the mixture of iron filings and tartar be calcined in a crucible for fome time with a red heat, and such part as cannot easily be reduced into fine powder, calcined again; the tartar will be converted into a fixt alkaline salt, and by this also the iron will be in part corroded and rendered soluble. There are several other methods of obtaining alkaline solutions of iron; but these kinds of combinations appear ill adapted for medicinal use, and are at present wholly neglected.

Mars folub. alkalizatus.

Some have made trial of the blue precipitate of iron called Prussian blue, and report that it seemed to act as a diaphoretic, and in some cases as an aperient (a). Of all the known preparations of iron, this promises the least activity: the perfect calces, almost if not wholly inert, are soluble in certain acids, particularly in the marine; but the Prussian blue is not acted upon by any kind of acid menstruum.

*FILIX.

FILIX MAS Pharm. Edinb. Filix non ramofa dentata C. B. Polypodium Filix mas Linn. MALE FERN: a plant of that class which has the fructification at the back of the leaves. This species has large doubly-pinnated unbranched leaves, with stalks rising singly from the root. The root consists of an oblong scaly body, terminating in numerous sibres. The plant grows commonly under hedges.

The efficacy of fern root against worms was known in the time of Dioscorides; and towards the beginning of this century, Messrs. Audry and Marchant published accounts of successful modes of exhibiting it in these cases. It was, however, fallen into neglect, till a few years ago, when it came again into notice, by being discovered to be the remedy which had become greatly celebrated in Switzerland as a specific in the cure of the tænia or tape-worm. The secret was purchased by the king of France, after its efficacy had been attested upon trial by some of the

(a) Geoffroy, Memoires de l'acad. roy. des sciences de Paris, pour l'ann. 1743.

principal

principal physicians at Paris. The following has been published as the mode of its exhibition. After the patient has been prepared by an emollient clyster and a supper of panada with butter and salt, he is directed to take in bed in the morning a dose of two or three drams of the powder of male fern root. The dose to infants is only one dram. The powder must be washed down with a draught of water, but nothing else must be taken till two hours after, when a bolus of calomel, joined with some of the stronger cathartics, is to be given. If this does not operate, it must be followed by a dose of purging salts. By this method, the worm is commonly expelled in a few hours. If the first trial does not succeed, the process must be repeated at due intervals (a).

*FLAMMULA JOVIS.

FLAMMULA JOVIS Stærck, Pharm. Edinb. Flammula recta C.B. Clematis recta Linn. Upright Virgin's Bower: this species of clematis, distinguished by its pinnated oval leaves, and erect stalk, grows wild in thickets in the southern parts of France and Germany. Its leaves and slowers are extremely acrid; the former, when fresh,

raifing blifters on the part to which they are applied.

The flammula jovis is one of the new medicines introduced by Dr. Stærck. He has published several cases of its efficacy in cancerous, venereal, and other malignant ulcers, obstinate pains of the head and bones, inveterate itch, and other diseases proceeding from peculiar acrimony. It was used internally, in infusion of the flowers or leaves, and extract of the plant; and the powder was sprinkled on the ulcers externally, where it was found to act as a most excellent escharotic and detergent. The medicine is said to have proved diuretic to some patients, and sudorishe to others, but rarely to have moved the belly. Small doses, of only half a grain of the exract, and half a dram of the dried leaves in infusion, were at first exhibited, which were gradually increased.

⁽a) Précis du traitement contre le Ténia ou Ver solitaire, pratiqué a Morat en Suisse, examiné & approuvé a Paris. Publié par ordre du Roi. Paris, 1775.

FOENICULUM.

FOENICULUM; Marathrum. Fennel: an umbelliferous plant, with dark green leaves divided into long capillary fegments: the umbels are fomewhat concave, and have no leaves or cup at their origin: the feeds oval, oblong, marked with prominent striæ: the root straight, white, about the thickness of the finger.

I. FOENICULUM DULCE Pharm. Lond. & Edinb. & C. B. Anethum Fæniculum Linn. Sweet fennel: with whitish or pale greenish yellow seeds, generally crooked. It is annual, a native of the warmer climates, and cultivated in our gardens. The shops are commonly supplied, from Germany, with seeds, superiour to those of our own growth.

SWEET fennel feeds are an useful stomachic and carminative; of an agreeable aromatic smell, and a moderately warm sweetish taste. They are sometimes given in powder, from a scruple to a dram; and sometimes candied.

Water extracts the virtue of these seeds very impersectly by insussion, but carries it off totally in evaporation: after repeated insussion, they retain part of their aromatic warmth, and the liquors are much less agreeable than the seeds in substance; after boiling for some time, the seeds prove entirely insipid, and the decoction, inspissated to the consistence of an extract, is very nearly so. By distillation, they impregnate water with their flavour: a gallon receives a strong impregnation from a pound of the seeds. A large proportion of essential oil separates in the distillation, and floats on the surface of the aqueous sluid; in colour yellowish, in smell moderately strong and diffusive, and exactly resembling the sennel, in taste mild and sweetish like the oil of aniseeds, and like it also congealing, by a slight cold, into a white butyraceous mass.

Aqua femin fœniculi Pb. Lond.

Ol.effentiale femin. fœnic dulcis Pb. Lond.

These seeds contain likewise a considerable quantity of a gross oil of the expressed kind, which, when freed from the essential oil, discovers no particular smell or taste. This oil is extracted, along with the aromatic matter of the sennel; by digestion in rectified spirit, but separates and rises to the surface upon inspissating the filtered tincture. The spirit, gently distilled off, has very little of the slavour of the seeds;

feeds; the oily matter retains a part both of their taste and smell; but much the greatest part remains concentrated in the extract.

2. FOENICULUM VULGARE Pharm. Edinb. Fæniculum vulgare germanicum C. B. Common fennel or finckle: with smaller, dark coloured almost blackish seeds. It is now reckoned only a variety of the
former.

THE feeds of this kind are warmer and more pungent, but less sweet, and of a less grateful flavour, than those of the preceding; and the same difference obtains in the distilled waters, distilled oils, and the spirituous extracts of the two kinds. The spirituous tinctures are somewhat different also, as the seeds themselves, in colour: those of the sweet fennel feeds being yellowish, of the common greenish.

The leaves of common fennel have the same kind of slavour with the seeds, and are in smell stronger, though in taste weaker and less agreeable. They impregnate water, by distillation, with a sufficiently grateful slavour, and yield a considerable portion of essential oil. An extract made from them by rectified spirit is likewise no inelegant aro-

matic: the colour of the spirituous tincture is a deep green.

The roots, taken up early in the spring, have a pleasant sweetish taste, with a flight aromatic warmth; but nothing of the peculiar strong flavour of the leaves and feeds. They are ranked among the aperient roots, and supposed by some to be equivalent in virtue to the celebrated ginseng of the Chinese, from which however they differ considerably in their fenfible qualities (fee Ginfeng). They give out their virtue, by infusion or slight coction to water, and by moderate digestion to rectified spirit: to the latter they communicate a pale amber colour, to the former a wheyish appearance. The aqueous infusions are in taste confiderably the strongest, but on being inspissated, they yield an extract of very little taste and in very small quantity; greatest part of the fweetish matter as well as the aromatic being diffipated in the evaporation. The spirituous extract is in larger quantity, about one twelfth the weight of the root, and of a moderately strong taste; agreeable, unctuous, fweetish, lightly aromatic, with some small admixture of bitterifhness.

FOENUM GRÆCUM.

FOENUM GRÆCUM Pharm. Lond. & Edinb. Fænum græcum fativum C. B. Fænu-græcum J. B. Trigonella Fænum-græcum Linn. Penugreek: a plant with flightly ferrated oblong or roundish leaves, set three on a pedicle; and whitish papilionaceous flowers, which are followed by long slender crooked flattish pods, containing yellowish rhomboidal seeds furrowed from angle to angle. It is sown annually in the southern parts of Europe, from whence the seeds are brought to us.

Fenugreek feeds have a strong disagreeable smell, and an unctuous farinaceous taste accompanied with a slight bitterishness. An ounce renders a pint of water thick and slimy: the decoction, inspissated, yields an unctuous mucilaginous bitterish extract, retaining a considerable share of the ill slavour of the seeds, and amounting to about three fourths their quantity. To rectified spirit they give out the whole of their distinguishing smell and taste; and afterwards to water a strong slavour-less mucilage.

The principal use of these seeds is in cataplasms and somentations, for softening, maturating, and discussing tumours; and in emollient and carminative glysters. They are an ingredient in the oleum e mucilaginibus of the shops, to which they communicate a considerable share of their smell.

FORMICA.

FORMICA Pharm. Parif. The ANT or PISMIRE: a fmall, oblong, reddish or blackish insect, furnished with a sting: the male has four wings, naked or uncovered; the female none.

This infect contains an acid juice; which it sheds on being irritated; with which, by agitation or boiling, it impregnates both water and rectified spirit; less volatile than pure spirit, so as to be concentrable from the spirituous insusion by drawing off a part of the menstruum; not quite so volatile as water, a considerable part of the water arising first with only a slight acidulous impregnation, and the strong acid coming

coming over with the remainder; differing in its properties from all the other known acids; and approaching nearest to those produced from vegetables by fermentation. The ant contains likewise a gross oil, separable by boiling in water, rising to the surface of the aqueous sluid, and similar in its general qualities to the expressed oils of vegetables; as also a subtile oil, which comes over in distillation both with rectified spirit and with water, analogous to the vegetable essential oils, but

wanting their pungent taste (a).

The medical qualities of this infect, and its remarkable productions, are not certainly known. It has been generally supposed, that the ants in substance, and insuspense and distilled waters of them, have an aphrodisac virtue; a virtue for which the above analysis does not appear to afford much foundation, though they are still retained in the aquæ magnanimitatis and other like compositions in foreign pharmacopæias. The insects in their chrysalis state, commonly called ants eggs, (which discover no marks of acidity) are said to be strongly diuretic and carminative: a decoction of a spoonful of them in butter-milk has been directed by some to be taken every morning in dropsies. The acid is recommended by Hossiman as one of the best menstrua of iron for medicinal uses.

FRANGULA:

FRANGULA five Alnus nigra Pharm. Edinb. (b) Alnus nigra baccifera C. B. Rhamnus Frangula Linn. BLACK ALDER: a small tree, or shrub, with slender slexible branches, and broad roundish leaves; bearing black berries, which contain a blue juice, with two seeds in each. It is common in moist woods in several parts of England.

The internal yellow bark of this shrub is a strong cathartic, and in this intention is sometimes made use of by the common people in dropsies and other disorders: it generally operates with great violence, occasioning nauseæ, sickness, gripes, and often vomiting. An infusion or decoction of it in water, inspissated to the consistence of an extract, acts with greater mildness than the bark itself. It gives a deep yellow tincture both to water and spirit.

⁽a) Vide Neumann, Chemical works, p. 497. Marggraf, Memoires de l'acad. roy. de scienc. de Berlin, pour l'ann. 1749.

(b) Expunged.

The berries also are strongly purgative; but are scarcely ever made use of, at least under their own name. In our markets, they are said to be sometimes substituted to those of buckthorn; which last may be distinguished by their green juice, and by their containing generally four seeds.

FRAXINUS.

FRAXINUS Pharm. Edinb. (a) Fraxinus excelsior C. B. & Linn. Ash: a tall tree common in woods and hedges; with a whitish bark, and oblong reddish brown seeds in shape somewhat resembling a bird's tongue, whence their names lingua avis, ornithoglossum, &c.

The bark of the ash tree, when fresh, has a moderately strong, bitterish, unpleasant taste, which in drying grows weaker. It has been given in substance from half a dram to a dram, and an extract made from it by water in smaller doses, as a resolvent and diuretic (b), and in intermitting fevers, in which it is faid to have often proved fuccefsful, especially when affisted by fixt alkaline salts. Vander Mye reports, that at the fiege of Breda, in defect of guaiacum, a decoction of this bark was made trial of in its place, and was found to be a potent fudorific; that in consequence of this discovery, it was given in pestilential cases, but that the decoction being difguftful by its quantity, a diffilled water was substituted, which in doses of two spoonfuls excited sweat freely, and was falutary to many (c). It must be observed, that this water was distilled in a fand heat, and is described as being ungrateful and smoky: from whence it appears to have been, not what is commonly called a distilled water, but an acid empyreumatic liquor, such as is forced out by fire from all vegetables: whatever might have been its virtues in malignant difeases, they apparently depended, not upon its being a preparation of ash bark, but on its being an acid.

Among us, this bark is regarded only on account of a phenomenon, of more curiofity than use, observed in its watery insusion, similar to that of the insusion of the lignum nephriticum. The liquor, if only slightly impregnated with the bark, on being held against the light, appears of a pale yellowish colour; looked down upon, or placed betwixt

⁽a) Expunged. (b) Duchesne, (Quercetanus) Pharm. dogmat. restitut. cap., 26.

⁽c) Fred. Vander Mye, de morbis Bredanis, p. 26.

the eye and an opake object, blue: the addition of an acid destroys the blueness, and alkalies recover it again. The spirituous tincture exhibits the same variability of colour; with this difference, that against the light, it appears of a much deeper gold or orange yellow.

The feeds of the ash tree have been given to the quantity of a dram, as diuretics, aphrodisiacs, and for reducing corpulent habits. They have a considerable taste, of a bitterish, aromatic, not very agreeable

kind.

FRUCTUS HORÆI.

FRUCTUS HORÆI Medicorum. Fraga, cerasa, ribesia, mora, fructus rubi idæi, &c. Summer fruits: strawberries, cherries, currants, mulberries, raspberries, &c.

THESE mild fweet subacid fruits are sometimes used medicinally, as refrigerants, antiseptics, relaxants, attenuants and aperients. Boerhaave looks upon their continued use as one of the principal remedies in cases of obstruction and viscidity, and in putrid disorders (a); and Hossman gives instances of some obstinate diseases being cured by them (b): they apparently promote the alvine and urinary excretions; and in some fevers, where watery liquors run off almost unchanged, these fruits or their juices render the urine coloured. As dietetic articles, they afford little nutriment, and are liable to produce flatulencies: to persons of a bilious temperament and rigid sibres, and where the habit is disposed, naturally or from extrinsic causes, to an inflammatory or putrescent state, their moderate, and even plentitum as salubrious: by those of a cold inactive disposition, where the vesses are lax, the circulation languid, and the digestion weak, they should be used very sparingly.

The juices, extracted from fruits by expression, contain the medicinal parts, freed from the grosser indigestible matter. On standing, they ferment, and change to a vinous or acetous state: by a proper addition of sugar, and by boiling, their fermentative power is suppressed, and their medicinal qualities preserved. The inspissated juices are found to be less statulent, and less disposed, when taken freely, to pro-

⁽a) Elementa chemiæ, process. iii. Praxis medica passim.

⁽b) Med. rational. de affectione phthisica, obs. i. Oper. tom. iii. p. 295.

duce gripes and fluxes, than an equivalent quantity of the fruits in fubftance or of the juices unboiled.

Syrup. mororum, & fructus rubi idæi Ph. L. These juices, purified from their seculencies by settling and straining, are made into syrups by a less proportion of sugar, than water or the common watery insusions require. For a quart of the depurated juices of mulberries, raspberries, &c. sifty ounces at most are sufficient; whereas the generality of vegetable insusions require sifty-eight. The more juicy berries give out their juice by heat without expression: if equal parts of picked currants and sugar be set over a gentle fire, the sugar dissolves in the juice of the fruit, and by boiling for a little time, an elegant jelly is formed, which may be freed from the skins by straining. These preparations may be occasionally dissolved in water, and used as diluents, resolvents, &c. in acute and other diseases.

The kernels of the stones of fruits, as of cherries, plums, &c. are of the same general nature with almonds. Those which have any bitterishness or particular slavour, receive these qualities from a subtile principle; which is extracted by maceration in vinous spirits; which rises in distillation with water; and which, when thus separated from the oily and farinaceous matter of the kernel, and combined with only a small quantity of the menstruum, appears to be, like the slavouring matter of bitter almonds, poisonous. Some physicians having found, that a distilled water very strongly impregnated with black cherry kernels, no more than two pints being distilled from sourteen pounds of the strong bruised, proves poisonous to brutes; the committee of the London college, appointed to reform their pharmacopæia, repeated the experiment with the same event.

FRUMENTUM.

FRIMENTA, Farinacea, Cerealia, Medicorum. Triticum ejusque amylum & furfur; oryza; avena; bordeum, &c. BREAD-CORN or GRAIN: wheat, with the starch made from it by maceration in water, and its bran; rice; oats, barley, &c.*(a)

THESE

^{*(}a) The nutrimental matter in grain and other vegetables is, 1. starch, which yields in analysis the same principles with honey, sugar, manna; viz. an acid, and an oil which sinks: 2. glutinous vegeto-animal matter; the dark coloured substance which in starch-making settles on the surface of the white, and is taken off for feeding hogs; of the same nature with the

These farinaceous feeds are less nutritious than the dietetic articles taken from the animal kingdom, but the nutriment they afford is milder and more benign: it is said that gouty and other chronical disorders, which are palliated or relieved by using milk for the only aliment, receive like relief from farinaceous aliments, provided the stomach is sufficient for their digestion (a). In all cases, they are a necessary addition to animal food; and are, perhaps, the best correctors of the putrid disposition which animal substances of themselves would produce. The alimentary animal substances, which of themselves run into high putrefaction, undergo, when mixed with the farinacea, a resolution of another kind; the mixture tending, not to a putrid, but to an acid state (b). These mixtures appear to be sooner resoluble than either the animal or vegetable matter separately: probably they are easier also of digestion in the stomach.

Among the common kinds of grain, rice is accounted the mildest and most nutritious, and supposed to be particularly serviceable in dysenteries and diarrheas. It is less viscous than wheat, or of less tenacity when boiled with water. It swells to a larger bulk in water than any other grain.

Wheat, whether in the form of flower or of starch, is perhaps the most glutinous of all the farinacea. The viscous substance which the flower forms with milk, is often a salubrious aliment in fluxes and catarrhs. The starch is used medicinally for the same intentions, in powders, mixtures, and other forms: a dram of starch, with three ounces of any agreeable simple water, and a little sugar, compose an elegant gelly, of which a spoonful may be taken every hour or two. These gelatinous mixtures are likewise an useful injection in some

cafeous part of milk; not foluble in water or spirit; yielding in analysis no acid, but a volatile alkaline spirit and salt, and an oil which swims. — Starch is the same substance in all vegetables containing it: feculæ of arum root, bryony, iris, dandelion, American manioc or yacca, freed from the poisonous juice, is an identical matter with starch of wheat. — The glutinous matter is likewise the same in all vegetables. From juices of herbs, heated, a green secula separates; from which spirit extracts green resin, of the same general properties with other resins, leaving behind a pure glutinous vegeto-animal substance. The secula which separates sirst, or with least heat, is richest in tinging resin; the next, in the glutinous matter. Repeated digestion in large quantities of spirit is necessary for complete separation. Rouelle, &c. Journ. de Med. 1773.

⁽a) Vide Malouin, Chimie medicinale, part. iii. chap. 2. tom. i. p. 234.

⁽b) See Dr. Pringle's experiments on this subject, in the appendix to his observations on the diseases of the army.

U u diarrheeas.

diarrhœas, particularly where the lower intestines have their natural mucus abraded by the flux, or are constantly irritated by the acrimony of the matter.

Oats are reckoned to be less viscous, and less nutrimental, than the two preceding.

Barley is less nutrimental, less glutinous, more cooling, more easily refoluble by fermentation, and probably in the stomach also, than either of the foregoing. Among the ancients, decoctions of it were the principal medicine, as well as aliment, in acute diseases; and from the common use of shelled barley, ptisana, in that form, other preparations of this kind, though made with different ingredients, have been often distinguished by the same name. The barley is freed from the shell in mills, and in this state called French or Scotch barley. A fort of shelled barley has been commonly brought from Holland in small round grains, called from their pearly whiteness, bordeum perlatum, or pearl barley *(a).

Decoctions of the farinacea in water, containing only their lighter and more agreeable parts, are very useful diluents in acute and other inflammatory difeases: however trivial preparations of this kind may appear to be, they are often, in these cases, medicines of primary importance. The most elegant and grateful beverage is prepared from pearl barley, by washing, for example, two ounces of it, from the mealy matter that adheres, with cold water; then boiling it with about half a pint of fresh water, which will acquire some tinge, and is to be thrown away; and afterwards putting the barley into two quarts of boiling water, and continuing the coction till one half is wasted.

drank plentifully, has effected a cure.

Bran, confifting of the hufks or shells of the grain, with a portion of its farinaceous matter, is supposed to have a laxative and detergent

An infusion or decoction of well toasted bread is likewise a very agreeable diluent, of the aftringent kind. In the cholera morbus, or bilious vomitings and purgings, it is often retained by the stomach when other liquors and medicines are rejected; and in feveral instances, by being

*(a) MAIZ, with water forms the most gluey viscid substance of any of the farinacea. We

Aq. hordeata Pb. Lond.

have never been able to make it undergo fuch a fermentation as to have the friability of our grains. Cullen, Mat. Med.

quality. Decoctions of it sweetened with sugar, are used by the common people, and fometimes with good fuccefs, against coughs.

* WORT. Dr. Macbride, in his very ingenious Experimental Essays, having laid down as a principle, "that the cure of the fcurvy depends on the fermentative quality in the remedies made use of," was led to enquire after a substance, capable of being preserved during a long sea voyage, and yet containing materials by which a fermentation might be occafionally excited in the bowels. Such an one appeared to him to be found in malt, which is well known to be the grain of barley, brought fuddenly to a germinating state by heat and moisture, and then dried, whereby its faccharine principle is developed, and rendered eafy of extraction by watery liquors. The fweet infusion of this, called wort, he proposed to give as a dietetic article to scorbutic persons, expecting that it would ferment in their bowels, and give out its fixed air, by the antiseptic powers of which the strong tendency to putrefaction in this difease might be corrected.

It was fome time before a fair trial of this proposed remedy could be obtained; and different reports were made concerning it. By fome cases, however, published in a postscript to the second edition of the Doctor's work, in 1767, it appears that fcorbutic complaints of the most dangerous kind have actually been cured at sea by the use of wort. Its general effects were to keep the patients open, and to prove highly nutritious and strengthening. It sometimes purged too much, but this effect was easily obviated by the tinctura thebaica. Other unquestionable cases of its success in this disease are to be seen in Vol. V.

of the London Med. Eff. and Ing.

The use of wort has hence been adopted in other cases where a strong putrid disposition in the fluids appeared to prevail, as in cancerous and phagædenic ulcers; and instances are published in the 4th Vol. of the work above-mentioned of its remarkable good effects in these cases.

As the efficacy of the malt infusion depends upon its producing changes in the whole mass of fluids, it is obvious that it must be taken in large quantities for a confiderable length of time; and rather as an article of diet than medicine. From one to four pints daily have generally been directed. The proportion recommended in preparing it, is one measure of ground malt to three equal measures of boiling water.

Uu 2

The

The mixture must be well stirred, and left to stand, covered, three or four hours. It should be made fresh every day.

FULIGO.

FULIGO LIGNI Pharm. Lond. & Edinb. WOOD SOOT: the smoke of burning wood condensed into a shining black concrete.

Wood foot has a disagreeable smell, and a pungent, bitter, nauseous taste: the more resinous the wood, the bitterer is the soot. On a chemical analysis, it is resolved into a volatile alkaline salt, an empyreumatic oil, a fixt alkali, and an insipid earth. The volatile salt and spirit are sometimes kept in the shops, and have in some cases been pre-

ferred to those of the animal kingdom.

Wood foot is directed in hysteric cases, and in different nervous disorders, as an antispasmodic and corroborant. It is used chiefly in the form of a spirituous tincture, in conjunction, commonly, with asafetida or other materials of similar intention: the officinal tincture is drawn from two ounces of soot and one of asafetida, with a quart of proof spirit, or a pint of proof and a pint of rectified spirit. The virtues of the soot are extracted, almost equally, by proof spirit, rectified spirit, and water; each of which, if the soot is of a good kind, dissolves about one fourth of it. The extracts, obtained by inspissating the filtered solutions, are excessively bitter: the spirituous extract retains most perfectly the peculiar slavour of the soot.

Tin&t. fulig. † Ph. Lond. 1Ph. Edinb.

FUMARIA.

FUMARIA Pharm. Edinb. Fumaria officinarum & dioscoridis C. B. Fumaria officinalis Linn. Fumitory: a plant with blueish green finely divided leaves; producing, towards the tops of the stalks, opposite to the leaves, spikes of irregular purplish flowers followed each by a fingle seed. It is an annual weed in shady grounds, and flowers in May and June.

THE leaves of fumitory are very juicy; of a bitter, somewhat saline taste; and no remarkable smell. The expressed juice, and a decoction of

of the leaves in water, inspissated to the consistence of extracts, are very bitter, and considerably saline: on standing for some time, they throw up to the surface copious saline efflorescences, in sigure somewhat resembling the crystals of nitre, to the taste bitterish and slightly pungent. A tincture of the dry leaves in rectified spirit yields, on inspissation, an extract, less in quantity, and bitterer in taste, than either the watery extract or inspissated juice: no saline matter separated from this extract, nor did it appear to the taste any other than simply bitter.

This herb is recommended as an aperient and resolvent, in obstructions of the viscera, in scorbutic and cutaneous maladies: Hossman has a high opinion of it as a purifier of the blood, and gives it the preference to all the other herbs made use of in that intention(a). It appears from the above analysis to be a plant of no inconsiderable virtue, though at present a stranger to practice: its sensible operation is by loosening the belly and promoting urine.

GALANGA.

GALANGA MINOR Pharm. Edinb. (b) & C. B. Wanhom Kæmpfer amænitat. exot. GALANGAL or Lesser GALANGAL: the root of a graffy-leaved plant, (Kæmpferia Galanga Linn.) brought from China and the East Indies, in pieces about an inch long and scarce half so thick, branched, full of knots and joints with several circular rings, of a reddish brown colour on the outside and a pale reddish within. We sometimes meet with a larger root of the same kind, an inch or more in thickness, under the name of galanga major: this is to be rejected, as being much weaker, yet more disagreeable, than the small.

GALANGAL root has an aromatic smell, not very grateful; and an unpleasant, bitterish, hot, biting taste. It was formerly in common use as a warm stomachic bitter, and generally made an ingredient in bitter insusions; but is now almost wholly laid aside, on account of its unpleasant slavour. Nor indeed does bitterness appear to be its proper medical character; the heat and pungency greatly prevailing. An extract made from it with rectified spirit is excessively fiery, setting the mouth

as it were in a flame: the watery extract is likewise very hot and pungent, though much less so than the spirituous, its quantity being about three times as large: neither one nor the other extract discovers any great bitterness. In distillation with water, there arises an essential oil, to the quantity of about a dram from sixteen ounces, of little smell, and of no great pungency. The pungent matter of the galangal appears from these experiments to be of the same nature with that of pepper; residing, not in the volatile oil, but in a more fixt matter.

GALBANUM.

GALBANUM Pharm. Lond. & Edinb. GALBANUM: the concrete gummy-refinous juice of an evergreen umbelliferous plant growing in Ethiopia, ferula africana galbanifera foliis & facie ligustici Herman. par. bat. Bubon Galbanum Linn. It is brought to us in pale coloured, semi-transparent, soft, tenacious masses: the better sorts of which, on being opened, appear composed of clear whitish tears, often intermixed with little stalks or seeds of the plant. By age it grows yellowish or brown.

This juice has a strong unpleasant smell, and a bitterish, warm, somewhat biting taste. In medical virtue, as in its sensible qualities, it is similar to ammoniacum; but is generally accounted less efficacious in asthmatic disorders, and more so on account of its stronger slavour, in hysteric cases.

Galbanum, like the other gummy refins, unites with water, by trituration, into a milky liquor; but does not perfectly diffolve, as fome have reported, in water, vinegar, or wine. Rectified spirit takes up much more than either of those menstrua, but not the whole: the tincture is of a bright golden colour. A mixture of two parts of rectified spirit and one of water dissolves all but the impurities, which are commonly in considerable quantity. It is best purified by including it in a bladder, and keeping it in boiling water till it melts or becomes soft enough to be strained, by pressure, through a hempen cloth: if this process be skilfully managed, the galbanum loses but little of the essential oil, in which great part of its virtue consists, and which appears to be carried off in evaporation both by water and spirit. In distillation with water, the oil separates and rises to the surface, in colour yellowish, in quantity

Galbanum purificatum Ph. Lond. about one twentieth of the weight of the galbanum: in this respect it differs from ammoniacum, which has not been observed to yield any essential oil. From that and the other common gummy resins it differs also in regard to the appearance of the empyreumatic oil obtained by distillation in a retort without addition; the empyreumatic oil of galbanum being, as Neumann observes, of a blue colour, which changes in the air to a purple.

GALLÆ.

GALLÆ Pharm. Lond. & Edinb. GALLS: hard roundish excrescences, found in the warmer countries, on the oak tree; produced from the puncture of an insect, and affording a lodgment for its young till they are capable of eating a passage through: those galls, which have no hole, are found to have the dead insect remaining in them. Two sorts are distinguished in the shops, one said to be brought from Aleppo, the other from some of the southern parts of Europe. The former are generally of a blueish colour, or of a greyish or blackish verging to blueness, unequal and warty on the surface, hard to break, and of a close compact texture: the others are of a pale brownish or whitish colour, smooth, round, easily broken, less compact, and of a much larger size. The two sorts differ in strength, but in other respects appear to be of the same quality: the Aleppo or blue galls are the strongest, two parts of these being equivalent to at least three of the others.

This excrescence is a strong astringent; one of the strongest of those of the vegetable kingdom. It has no smell, or particular flavour; simple astringency being its medical character. The cortical hard part of the gall appears considerably stronger than the interiour more brittle matter. The virtue of both is taken up by watery and by spirituous menstrua: on inspissating the tinctures, both the water and spirit rise unflavoured, leaving extracts of intense stypticity: the spirituous extract is in smaller quantity than the watery, and somewhat stronger in taste. The galls in substance have been given in small doses in different disorders proceeding from relaxation, and recommended by some in intermitting fevers, in doses of half a dram and a dram; but it is surely imprudent to venture on such large quantities, of so strong an astringent. Among

us they are employed chiefly as an external flyptic, in embrocations and injections.

G A L L I U M.

GALLIUM Pharm. Edinb. (a) Gallium luteum C. B. Galium verum Linn. LADIES BEDSTRAW: a plant with square stalks, and long narrow foft leaves, standing generally eight at a joint in form of a star: on the tops, and on short pedicles issuing from among the leaves, grow thick clusters of small yellow monopetalous bell-shaped flowers, divided, each, into four fegments, and followed by two feeds. It is perennial, common in dry waste grounds, and flowers in June and July.

THE flowers of this plant have a moderately strong, not disagreeable fmell; the leaves, little or none. They both discover to the taste a fenfible acidity; which they manifest also by changing the juices of blue flowers to a red, and by coagulating boiling milk: they are faid to be, in fome places, commonly made use of in this last intention, whence one of the common names of the plant, cheefe-rennet. Their acid matter appears to be, if Borrichius's experiment is to be depended on, of a more fubtile kind than that of forrel, and than the other native vegetable acids that have been examined; the flowery tops, committed to diffillation as foon as gathered, giving over a pretty strong acid liquor, in a moderate heat, wherein forrel yielded only an infipid phlegm (b). The restringent and refrigerating virtues, ascribed to this plant, appear from these experiments to have some foundation.

GAMBOGIA.

GAMBOGIA Pharm. Lond. & Edinb. Gummi gutta, gamandra, gamma, jemu, &c. Cambogia gutta Linn. GAMBOGE: the concrete gummy-refinous juice of certain trees growing in Cambogia or Cambodia and some other parts of the East Indies: brought over in large cakes or rolls, externally of a brownish yellow, internally of a deep reddish yellow or orange colour, changing to a pale bright yellow on being moistened.

(a) Expunged. (b) Alla medica & philosophica Hafniensia, vol. i. obs. 69.

This juice has no smell, and when first chewed makes but little impression on the organs of taste: kept in the mouth for some time, it discovers a considerable acrimony. Rectified spirit of wine, poured upon it, acquires immediately a deep gold colour, and dissolves about five parts out of six. Water, assisted by heat, takes up nearly as much; but the solution is turbid, and deposites, on cooling, a considerable quantity of resinous matter. Water, impregnated with fixt alkaline salt, totally dissolves it into a transparent blood red liquor, which passes through a filter without any separation of its parts, and deposites no sediment on standing. It is wholly taken up likewise, and in considerable quantity, by vinous spirits impregnated with volatile alkalies, or the dulcified spirit of sal ammoniac: this solution mingles uniformly both with water and rectified spirit, without precipitation or turbidness.

Gamboge is a strong and quick cathartic; producing copious evacuations, and usually finishing its operation soon. In such hydropic cases as require the brisker cathartics, and in other disorders accompanied with a redundance of serous humours, it is an useful and safe hydragogue: in hot, dry, bilious constitutions, it is never to be ventured on: in all cases, it is liable, on first using it, to vomit as well as purge. The dose is from three or sour grains to twelve, or at most fifteen.

This medicine is most disposed to act upwards, when given in the solid form of a bolus or pill: by joining to it mercurius dulcis, its emetic power is generally restrained. It is principally made use of in conjunction with that mercurial preparation and with other purgative materials. Solutions of gamboge in alkalized water, and in dulcified alkaline spirits, act only by stool and urine, and with much greater mildness than the juice in substance. The watery tincture is still milder, the menstruum dissolving only a part of the resin: the spirituous tincture operates with extreme irritation both upwards and downwards.

GENISTA.

GENISTA Pharm. Edinb. Genista angulosa & scoparia C. B. Spartium scoparium Linn. Broom: a shrubby plant; with numerous, slender, angular, tough twigs; small, somewhat oval leaves, set three on one pedicle; and deep yellow papilionaceous flowers, which are followed

Xx

by broad pods, containing hard brownish flat seeds. It is common on heaths and uncultivated sandy grounds, and slowers in May and June.

The leaves and stalks of broom have a nauseous bitter taste; which they give out, by insussion, both to water and rectified spirit; and which, on gently inspissating the filtered liquors, remains concentrated in the extracts: the watery tincture is of a yellowish green or brownish, the spirituous of a dark green colour. They are accounted laxative, aperient, and diuretic; and in this intention have been often used by the common people in dropsies and other serous disorders. Dr. Mead relates a case of an hydropic person, who, after the paracentesis had been thrice persormed, and sundry purgatives and diuretics had been tried without relief, was persectly cured, by taking, every morning and evening, half a pint of a decoction of green broom tops with a spoonful of whole mustard seed: by this medicine, the thirst was abated, the belly loosened, and the urinary discharge increased to the quantity of at least five or six pints a day (a).

Infusions of the ashes of the plant in acidulous wines have likewise been employed in the same intention, and often with good success. The virtue of this medicine does not depend, as some have supposed, on any of the peculiar qualities of the broom remaining in the ashes; but on the alkaline salt and earth, which are the same in the ashes of broom as in those of other vegetables, combined, wholly or in part, with the vinous acid. A solution even of the pure earthy part of vegetable ashes,

made in vegetable acids, proves notably purgative and diuretic.

Of the feeds and flowers, the medicinal qualities are not well known. It is faid that the feeds, in doses of a dram and a half in substance, and five or six drams in decoction or insusion, prove purgative or emetic. Some report that the flowers also operate in the same manner, but Lobel assures us, from his own observation, that they have been taken in quantity, without producing any such effect; and I have known insusions of the flowery tops drank freely in some asthmatic cases, without any other sensible operation than a salutary increase of urine and expectoration. The seeds, slightly roasted, are used in some places as coffee, and said to act as diuretics.

(a) Monita & præcepta medica, p. 138.

GENTIANA.

GENTIANA Pharm. Lond. & Edinb. Gentiana major lutea C. B. Gentiana lutea Linn. GENTIAN: a plant with an unbranched jointed stalk, and oblong acuminated ribbed leaves set in pairs at the joints upon broad pedicles: the flowers, which stand in clusters round the stalk in the bosoms of the upper leaves, are of a pale yellow colour, somewhat bell-shaped, deeply cut into five segments, followed by oblong capsules full of small seeds: the root is moderately long, slender, branched, brownish on the outside, of a reddish yellow or gold colour within. It is perennial, a native of the mountainous parts of Germany, &c. from whence the shops are generally supplied with the dried roots.

Among the gentian brought to London some years ago, a root of a different kind was mixed: the use of which occasioned violent disorders, and in some instances, as is said, proved fatal. This root is externally of a paler colour than gentian, and its longitudinal wrinkles finer and closer: on cutting the two roots, the difference is more remarkable, the poisonous root being white, without any degree of the yellow tincture which is deep in gentian; nor is its taste bitter, like that of gentian, but mucilaginous.

GENTIAN root is a strong flavourless bitter; in taste less exceptionable than most of the other common strong bitters, and hence among us most generally made use of. The flavour and aromatic warmth, wanting to render it grateful, and acceptable to the stomach, are supplied by additions. An ounce of the gentian root, with the same Infus. amar. quantity of fresh lemon peel, and a dram and a half of dried orange peel, infused for an hour or two in three quarters of a pint of boiling water, make a very elegant bitter. The lemon peel is an excellent addition in the watery infusions, but the perishableness of its flavour excludes it from spirituous tinctures designed for keeping. The Edinburgh college Infus. amar. have directed an addition of spirits to their bitter infusion, which is made of half an ounce of gentian root, one dram of dried orange peel, and half a dram of coriander feeds, infused in a quarter of a pint of proof spirit, and a pint of water. The bitter tinctures are commonly prepared, by macerating an ounce of the root, for some days, in a pint Tind.amar of proof spirit, with four drams of dried orange peel and two of leffer Pb. Lond.

X x 2

cardamom

Tinctura amara, five Elixir itomachic. Pb. Edinb.

Vinum amar. Ph. Lond.

Extractum gentianæ Ph. L. & Ed.

cardamom feeds; or in a pint and quarter of proof spirit, with the above quantity of orange peel, two drams of canella alba, and fifteen grains of cochineal. Wines and malt liquors are likewise impregnated with the same or similar materials, in different proportions: an ounce of the gentian root, the same quantity of fresh lemon peel, and two drams of long pepper, communicate, by maceration without heat, a grateful warmth and bitterness to a quart of mountain. The virtue of the root is extracted by all these menstrua, as also by rectified spirit; not totally, however, by any, in the above proportions; and not in so great a degree by water as by spirit. The tincture in rectified spirit is of an orange yellow colour: inspissated, it yields an intensely bitter extract, the spirit rising unflavoured. The watery insusions are of a dark brownish red; and leave, on being inspissated, an extract, in larger quantity, and less bitter, than the other. These extracts are made into pills, by themselves, or with aromatic additions.

The German ephemerides mention a root brought from America by the Portugueze, under the name of *Indian gentian*; of a pale yellowish colour, jointed, marked with various knots and circles like ipecacoanha; of a penetrating aromatic bitterness, not ungrateful, though far more intense than the bitterest of the officinal bitter drugs. This root is greatly commended in obstinate intermittents, and many other disorders: a scruple is said to do more than repeated half drams of bark (a).

*GEOFFRÆA.

GEOFFREA Jamaicensis inermis Dris. Wright. Geosfræa Pharm. Edinb. Cabbage bark, or Worm-bark tree. This is a tree growing abundantly in the low savannahs of Jamaica, of a considerable height, but no great thickness. It has a straight smooth trunk, and sends off its branches near the top. It bears dark green leaves, and rose-coloured flowers of the papilionaceous kind, set in purple flower-cups. These are succeeded by a green hard fruit, of the size of a small plumb, having a skin the thickness of a crown piece, and a nut within.

THE bark of this tree is externally of a grey colour, black and furrowed on the infide. To the taste it is mucilaginous and sweetish.

⁽a) Michael Schendo, Ada physico-medica acad. nat. curios. vol. i. anno 1727. Append. p. 112.

It fmells difagreeably, whence it has been called by fome the bulge-water tree. It has been long a celebrated anthelmintic in the West Indies,

and has lately been introduced into European practice.

The first account we meet with of its use is in a letter from Mr. Duguid, in Vol. II. of Essays Physical and Literary. Several subsequent accounts appeared in different numbers of the Medical Commentaries. But the sullest relation, together with an accurate botanical description of the plant, is given by Dr. William Wright in the Philosophical

Transactions, Vol. LXXVII. Part II.

The bark is used in the several forms of decoction, syrup, powder, and extract. For making the decoction, an ounce of fresh-dried bark is to be boiled slowly in a quart of water, till the liquor be of the colour of Madeira wine. This is to be strained off for use. The syrup is made by adding a sufficient quantity of sugar to this decoction. By evaporating the decoction, the extract is formed, which must be carefully stirred during the process, to prevent the resinous part from rising to the top. The decoction is generally preferred in Jamaica, and seems to be the most efficacious as an anthelmintic.

As this medicine is rather a violent one, it should always be exhibited at first in small doses. The most immediate effect of these is to produce naufea, which is fucceeded by brifk purging, especially when the powder is given. If cold water is drank during its operation, it is apt to occasion fickness, vomiting, fever, and delirium. These symptoms, whether occafioned by this cause, or by an over dose, are removed by washing the stomach with warm water, purging with castor oil, and giving plenty of drink acidulated with vegetable acid, which last seems a kind of specific against its deleterious effects. The manner in which Mr. Anderson, the writer of a paper concerning it in Vol. IV. of the Medical Commentaries, recommends its exhibition, is to give gradually augmented doses of the decoction for eight or nine mornings fuccessively, and then a dose of jalap and calomel, which feldom fails to bring away the worms, fome dead, and fome alive. This writer also remarks, that there are two kinds of the bark, one much paler than the other, which acts with greater violence.

GERANIUM.

GERANIUM Pharm. Parif. CRANESBILL: a plant, so called from the remarkable long beak of its seed-vessel, which consists of five capsules opening inwards, and containing each a single seed: the flowers are pentapetalous.

- 1. GERANIUM SANGUINARIUM seu hæmatodes. Geranium sanguineum maximo store C. B. Geranium sanguineum Linn. Bloody cranesbill: with solitary slowers, on their first appearance red, but soon changing to a blueish; the leaves roundish, but divided almost to the pedicle into sive segments, which are often subdivided at the extremities into three.
- 2. GERANIUM COLUMBINUM sive Pes columbinus. Geranium folio malvæ rotundo C. B. Geranium rotundifolium Linn. Dovesfoot: with purple flowers standing two on one pedicle; and mallow-shaped leaves on long footstalks.
- 3. GERANIUM BATRACHIOIDES five Batrachium. Geranium batrachioides gratia dei germanorum C. B. Geranium pratense Linn. Crowfoot cranesbill: with two blue (sometimes white) flowers on one pedicle: and large wrinkled leaves, divided into sive or seven segments, which are again deeply cut on the edges.
- 4. HERBA ROBERTI five Gratia dei. Geranium robertianum C. B. & Linn. Herb-robert: with two reddish or purplish flowers on one pedicle; the leaves divided quite to the footstalk into three segments; and these again deeply cut.
- 5. GERANIUM MOSCHATUM five Acus moschata. Geranium cicutæ folio moschatum C. B. Geranium moschatum Linn. Musk cranesbill: with a number of red flowers on one pedicle; and oval indented leaves, set in pairs along a middle rib, which is terminated by an odd one.

All these plants are found wild in different parts of this kingdom: the four first are common, the last rare. They flower in May, June, and

and July; the fourth earliest, the first latest. The second and fifth are annual, the fourth biennial, the others perennial.

THE above geraniums, formerly ranked among the officinals, and many other plants of the same genus, indigenous or commonly cultivated; discover to the taste a considerable astringency, and strike an inky blackness with solution of chalybeate vitriol: some of them are, apparently, of the stronger kind of vegetable styptics. The three first forts have no great fmell; the fourth, an unpleasant one, somewhat like that of the dead nettle, but stronger; the fifth has an agreeable musky fcent, which is destroyed by bruising the plant. The odoriferous principle is feparated by distillation with water, and gives a moderate impregnation to the distilled fluid; but no essential oil was obtained on submitting to the operation moderate quantities either of the fetid or the musky forts. The styptic matter is extracted both by water and rectified spirit; and on evaporating the filtered liquors, remains entire in the inspissated extracts: the watery infusions are yellowish or brownish, the spirituous tinctures of a deep green colour. The watery extracts, those at least of the second and fourth kinds, on standing for some weeks, throw off to the furface a confiderable quantity of small saline crystals, in shape somewhat like those of nitre, in taste austere and bitterish. From these experiments it may be presumed, that the geraniums have no ill claim to the vulnerary, that is, astringent virtues, commonly ascribed to them, as in alvine fluxes, hemorrhagies, defluxions on the breast, &c. *(a)

GINSENG.

GINSENG Pharm. Edinb. Aureliana canadensis, sinensibus gins-eng, iroquæis garent-oguen, Lasiteau memoir. sur le gins. Panax quinquesolium Linn. GINSENG: the root of a small plant; growing in China, Tartary, and likewise in some parts of North America, particularly Canada and Pensylvania, from whence considerable quantities have lately been brought over. It is two or three inches in length; taper; about

^{*(}a) Our host at Carlisle told us that he used to be troubled with the stone, and the best remedy he ever had experience of to give him ease was the decoction of geranium robertianum. Ray's remains, published by Scott, 1760.

the thickness of the little finger, or less, in the thickest part; often forked at bottom; elegantly striated with circular wrinkles; of a brownish or yellowish colour on the outside, and whitish or of a pale yellowish within: on the top are commonly one or more little knots or tubercles, which are the remains of the stalks of preceding years, and from the number of which, the age of the root is accordingly judged of.

On comparing the American roots with some specimens received from Nankin, no material difference could be observed between them, either in their external appearance or in their quality; except that the Chinese were in general somewhat paler coloured on the outside, and internally rather whiter. It is said that in China, the roots, taken up in spring or autumn and carefully cleaned from the fibres, are washed and soaked for a time in a decoction of rice or millet-seed, and afterwards exposed to the steam of the liquor; that by this means they acquire a greater firmness and clearness than they have in their natural state; that nevertheless the American roots were received and purchased as true ginseng in China itself, though without the supposed advantage of the Chinese preparation.

Ninzin or Nindsin has been commonly supposed a name synonymous to ginseng. It appears from later observations, that the ninzin is the root of a different plant (a) which is cured in the same manner, and very nearly resembles the ginseng, but is supposed to be of weaker virtue. This also is a native of America as well as China. It is called by Kæmpser, sistem montanum coræense, radice non tuberoso; by Linnæus, sium foliis serratis pinnatis ramis ternatis.

GINSENG root, a medicine of extraordinary esteem among the Chinese as a general restorative and corroborant, though undoubtedly very far unequal to the character that has been commonly given of it, promises nevertheless, from its sensible qualities, to be an useful addition to the officinal drugs. To the taste it discovers a mucilaginous sweetness, approaching to that of liquorice, accompanied with some degree of bitterishness, and a slight aromatic warmth; with little or no smell. It is far sweeter, and of a more grateful kind, than the roots of fennel, to which it has by some been supposed similar; and differs likewise

remarkably from those roots, in the nature and pharmaceutic properties of its active principles; the sweet matter of the ginseng being preserved entire in the watery as well as in the spirituous extract, whereas that of fennel roots is destroyed or dissipated in the inspissation of the watery tincture. The slight aromatic impregnation of the ginseng is likewise in good measure retained in the watery extract, and perfectly in the spirituous; which last is a very pleasant, bitterish, warm sweet.

GLYCYRRHIZA.

GLYCYRRHIZA Pharm. Lond. & Edinb. Glycyrrhiza filiquofa vel germanica C. B. Glycyrrhiza glabra Linn. LIQUORICE: a plant with oval leaves, fet in pairs along a middle rib; and fmall blueish papilionaceous flowers, standing in spikes, on naked pedicles, at the junctures of the ribs of the upper leaves with the stalk: the flower is followed by a fmooth pod containing flat kidney-shaped seeds: the root is very long, slender, flexible, of a brownish colour on the outside and yellow within. It is perennial, a native of the fouthern parts of Europe, and plentifully cultivated in England: the roots are fit for being taken up in the third year after the flips or offsets have been planted. The liquorice root of our own growth is nowife inferiour to that which is produced in its native climate. The root carefully dried and powdered, is of a richer and more agreeable taste than when fresh, and of a dull yellow colour with a cast of brown: the liquorice powder commonly fold is of a weaker taste, and a paler and bright yellow colour, from an admixture probably of other fubstances.

LIQUORICE, one of the principal fweets, is almost the only one of the common substances of that class which tends to abate thirst: this property was known to the Greeks, who hence distinguished it by the name adipson, and employed it, as Galen observes, in hydropic cases, for alleviating the desire of drinking. It is an useful emollient and incrassant in defluxions on the breast, and supposed to prove at the same time gently detergent. Insusions and extracts, made from it, afford likewise very commodious vehicles or intermedia for the exhibition of other medicines: the liquorice taste concealing that of unpalatable drugs more effectually than syrups or any of the sweets of the saccharine kind. It differs also from the

Yy

fweets

fweets of the faccharine or honey kind, from the fweet juices of fruits, and from the fweet matter afforded by the common forts of grain when beginning to vegetate; in being far less disposed to run into a fermentative state. The cortical part of the root is considerably sweeter than the more compact internal fubstance, but the fweet matter appears to be in both of the fame kind.

Extractum glycyrrhizæ Ph. Lond.

Liquorice root, lightly boiled in a little water, gives out nearly all its fweetness: the decoction, pressed through a strainer, and, after settling, carefully inspissated, with a gentle heat, till the matter will no longer flick to the fingers, affords an extract exceedingly fweet, more agreeable than that brought from abroad or prepared among ourselves in the way of business, of a pleasant smell, of a dark reddish brown colour in the mass, and when drawn out into strings, of a golden colour: its quantity amounts to near half the weight of the root. If the liquorice be long. boiled, its fweetness is greatly impaired, and the preparation contracts an ungrateful bitterness (a).

Rectified spirit takes up the sweet matter of the liquorice equally with water; and as it disfolves much less of the insipid mucilaginous substance of the root, the spirituous tinctures and extracts are proportionably fweeter than the watery: they are accompanied also with a slight, but very fenfible, pungency. The quantity of spirituous extract amounts only to about one half of the aqueous; and rectified spirit, digested on the aqueous extract, diffolves about one half of it, taking up nearly the

whole of its fweetness.

GRAMEN.

GRAMEN CANINUM Ph. Lond. Gramen caninum arvense seu gramen dioscoridis C.B. Triticum repens Linn. Dogs grass, Couch: a creeping grass, of a whitish green colour, with knotty stalks, bearing a spike of imperfect flowers somewhat resembling a wheat ear: the roots are whitish or pale yellowish, long, slender, jointed at distances, variously bent and interwoven. It is a perennial weed.

THE

^{*(}a) The extract from the first infusion is bronze yellow, exceedingly sweet, without any acrid or bitter relish : - from a second infusion, much deeper coloured, and far less agreeable : - afterwards, by coction, a black acrid extract is obtained, in which the tafte of the liquorice can hardly be perceived. Beaumé.

THE roots of this plant, to the taste agreeably sweetish, are recommended as mild aperients in obstructions of the viscera. Boerhaave directs the expressed juice to be taken, in this intention, to the quantity of some pints a day; and observes, that cattle are generally sound to have indurated livers in the winter, but that from fresh grass in the spring a diarrhoea ensues, and the obstruction is resolved.

GRANA PARADISI.

GRANA PARADISI Pharm. Edinb.(a) Cardamomum majus seu grana paradisi Pharm. Paris. Meleguetta & maniguetta & cardamomum piperatum quibusdam. Amomum Grana Paradisi Linn. GRAINS OF PARADISE, called by some GREATER CARDAMOMS: angular reddish brown seeds, smaller than pepper, in appearance much resembling cardamom seeds, brought from the East Indies. The grains of paradise and cardamom plants belong both to one botanic genus, the amomum of Linnæus.

In pharmaceutical properties, the grains of paradife differ greatly from cardamom feeds, and greatly refemble pepper. They have fomewhat of the flavour of the former, joined to the heat and pungency of the latter; which pungency refides, not like that of cardamoms in the volatile parts or effential oil, but like that of pepper in the refinous or more fixt matter. The diftilled oil of grains of paradife, in smell sufficiently agreeable, is in taste of the milder kind: the remaining decoction, inspissated, yields an extract of considerable pungency: an extract made by rectified spirit is highly siery. This spice is employed in some places for the same purposes as pepper; among us it is rarely directed for medicinal purposes.

GRANATA.

GRANATA malus Ph. Edinb. Malus punica fativa C. B. Punica granatum Linn. Pomegranate: a prickly tree or shrub; with long narrow leaves; deep red pentapetalous flowers set in bell-shaped cups of

(a) Expunged.

Y y 2

the

the same colour; producing fruit about the size of an orange, which consists of a thick tough rind, externally brownish and internally yellow, with a juicy pulp and numerous seeds in cells like a honeycomb. It is a native of the southern parts of Europe, and rarely brings its fruit to full persection in this climate.

The flowers of this tree are mild aftringents, fimilar to those of the wild pomegranate or balaustine, which last are preferred only on account of their being larger. The pulp of the fruit, when in perfection, is of a grateful subacid sweet taste, and of the same general qualities with the other summer fruits. The rind of the fruit is moderately astringent, and in this intention is now and then directed, under the names of cortex granati, malicorium, psidium, and sidium: it yields with water near half its own weight of a very austere extract, but gives out very little to rectified spirit, its astringent matter, like that of the fruit of the acacia tree, seeming to be indissoluble in spirituous menstrua: in this respect the astringency of the fruit differs from that of the balaustine or flower of the tree.

+ Ph. Lond.

GRATIOLA.

GRATIOLA Ph. Edinb. Gratiola centaurioides C. B. Gratia dei. Gratiola officinalis Linn. Hedge hyssop: a low plant, with oblong finely ferrated leaves fet in pairs on the stalks without pedicles: in their bosoms come forth solitary, whitish, tubulous, irregular flowers, followed by roundish pointed capsules full of small seeds: the root is slender, whitish, jointed, surrounded with sibres. It is perennial, a native of the southern parts of Europe, and raised in some of our gardens.

The leaves of gratiola have a nauseous bitter taste, and no remarkable smell. They are said to be a strong hydragogue purgative; to operate upwards as well as downwards; and in weak constitutions, to occasion oftentimes violent gripes or superpurgations. Geosfroy observes, that the dose is an insusion of half a handful of the fresh, or a dram of the dry herb, in wine or water; that a slight decoction of it in milk operates far more mildly; and that an extract made with wine may be given to half a dram or two scruples. The roots are less ungrateful in taste, and less

less violent in operation, than the leaves: given in substance, from a scruple to a dram, they are said to vomit and purge, without much inconvenience: in some parts of Germany, they are reported (a) to be

commonly employed in dyfenteries, as ipecacoanha among us.

*In a differtation on the medical virtues of this plant, published at Vienna by a Polish physician, Jac. Kostrzewski, several cases are related of its efficacy in maniacal disorders; and also in the venereal disease, accompanied with ulcerations, in tumours, and fluor albus. In mania, the powder of the root to the quantity of half a dram was given, which constantly excited vomiting, purging, and a copious flow of urine. In the other cases, pills were exhibited, composed of an extract of gratiola, with sugar and absorbent and aromatic powders. This preparation excited a nausea, but did not vomit. In one venereal case a salivation was induced by its use. On the whole, the author recommends the gratiola as a most effectual remedy in all disorders proceeding from a superabundance of serum.

GUAIACUM.

GUAIACUM Pharm. Lond. & Edinb. Guaiacum americanum primum fructu aceris sive legitimum Breyn. prodr. Guaiacum ossicinale Linn. Guaiacum: a large tree, with roundish box-like winged leaves, pentapetalous blue slowers in clusters, and a maple-like heart-shaped fruit including a single seed (b), a native of Jamaica, Hispaniola, and other warm parts of America; from whence the wood with its bark, and a concrete resinous juice exuding from incisions made in the trunk, are brought to us. The wood is called by some lignum vitæ and sanctum.

THE WOOD is very hard, compact, and so heavy as to fink in water: the outer part is of a pale yellowish colour, the heart of a dark blackish brown with a greater or less admixture of green. It scarcely discovers any smell, unless heated, or while rasping; in which circumstances it yields a light aromatic one: chewed, it impresses a mild acrimony, biting the palate and sauces. Its pungency resides in a resinous matter,

⁽a) Kramer, Tentamen botanicum emendat. & auct. Introduct. p. 18.

⁽b) Sloane, Catalogus plantarum in insula Jamaica, p. 186. Voyage to Jamaica, vol. ii. p. 133.

which is totally extracted by digestion in rectified spirit, and partially by boiling in water: the spirituous tinctures are of a deep brownish red colour, the watery decoctions of a dark yellowish brown. On inspiffating the liquors, nothing of the pungency of the guaiacum exhales or distils with either menstruum: the spirituous extract, nevertheless, discovers but little of the pungent tafte which prevailed in the tincture, proving a tenacious almost pure resin, not dissoluble in the mouth or miscible with the faliva: the watery extract, which contains likewife no small proportion of refinous matter, disfolves slowly, and then manifests a notable degree of pungency. During the inspissation of the watery decoction, the refinous part is apt to separate and subside, unless a little rum Ph. Lon. spirit be added towards the end of the process to keep it united with the gummy: this extract is kept in the shops in a foft and a hard form; the first of a proper consistence for making into pills, the latter for being reduced into powder. The quantity of folid extract obtained by rectified fpirit amounts to about one fourth the weight of the wood; with water scarcely one fixth is obtained. After a pound of the shavings of the wood has been boiled in a gallon of water till half the liquor is wasted, and the coction fuccessively repeated with five or fix fresh gallons of water, a confiderable portion of refin may still be extracted by moderate digestion in rectified spirit.

Extractum ligni guaiaci molle & du-

> The BARK of guaiacum is confiderably less hard, but not much lighter, than the wood: it is thin, fmooth, composed as it were of a number of fine plates joined closely together, externally of a blackish grey colour variegated for the most part with greenish or livid specks, internally of a whitish or pale yellow. In taste and smell, it is similar to the wood, but weaker: the watery and spirituous extracts are of the fame quality, but in less quantity.

> The refin, or GUM, fo called, is brought over in irregular maffes, usually friable, of a dusky greenish, and sometimes of a reddish hue; intermixed with small pieces of the wood; of a pungent taste, but of little or no fmell, unless heated. It contains more refin than the watery extract made from the wood; and more gummy matter than the spirituous extract. The refin, which is the only active part, is obtained pure both from the gummy substance and from the woody and other indissoluble impurities, by digefting the compound in rectified spirit, drawing off the spirit from the filtered solution till the matter begins to grow thick.

thick, and then adding a quantity of water, which will precipitate the pure refin, and keep diffolved fuch of the gummy parts as the spirit may have taken up. The quantity of refin, thus obtained, amounts commonly to about three fourths of the weight of the gum guaiacum.

GUAIACUM was first received in Europe as a remedy for the venereal difease; and is said, in the warmer climates, to have been sometimes fufficient for fubduing it. Though of itself greatly unequal among us to that distemper, it is a good assistant to mercurial alteratives, and a medicine of great use also in several other cases. To warm and stimulate the habit, to promote the excretions made from the blood, as perspiration and urine, and likewise the groffer evacuations from the intestinal canal, appear to be its primary virtues: in large doses, it operates as a purgative. Where the excretory glands are obstructed, the vessels lax and flaccid, and the habit replete with impure ferous humours; in fundry cutaneous and catarrhous diforders, some female weaknesses, and chronical rheumatisms; it frequently has good effects. In thin emaciated habits and an acrimonious state of the fluids, it often does harm: in such cases, it has converted a simple itching of the skin to ill-conditioned eruptions, or increased the itching to an almost insupportable degree: where this happens, nitre, whey, faline laxatives, and warm bathing, are commonly found most effectual for abating the complaints. Hoffman observes that it is, in general, less proper in an advanced age than in other circumstances.

A decoction of half an ounce of the wood or bark may be taken in a day, at proper intervals, the patient keeping warm to promote a diaphoresis. The gum, or extracts made from the wood, are given from a few grains to a scruple or half a dram, and sometimes two scruples; which last dose proves, for the most part, considerably purgative. The extract is recommended by Hoffman as an excellent errhine, which occasions a great discharge from the nose, and which he supposes, besides its stimulating power, to be endued with a corroborating one, very friendly to the nervous parts.

Solutions or tinctures of the gum guaiacum are made in the shops, both with rectified spirit of wine, and with the dulcified aromatised volatile alkaline spirits, vulgarly called fal volatile, which in many cases Tind. guaipromote its virtues. To fix ounces of the gum guaiacum are directed, by acina volatilis Ph. Lon.

Balfamum guaiac. Ph. Lond.

Elixir guaiacin. volat. Ph. Edinb.

Elixir guaiacin. Ph. Ed.

the London college, two pints and a quarter of the volatile spirit, and for a more faturated folution, one pint and a half of rectified spirit of wine, with the addition of a dram and a half of balfam of Peru. The Edinburgh college directs, for the volatile tincture, four ounces of the guaiacum, with two drams of balsam of Peru, and half a dram of effential oil of fassafras, to be dissolved in a pound and a half of dulcified spirit of fal ammoniac; and for the spirituous solution, one pound of gum guaiacum and three drams of balfam of Peru, with two pounds and a half of rectified spirit. All these are sufficiently elegant solutions of the guaiacum, and the additional articles coincide with its virtue: they may be given from twenty drops to a tea-spoonful or more in any convenient vehicle. The gum, or refinous extracts, may be diffolved also, by the mediation of thick mucilages, in watery liquors; and in this form are more commodiously taken than in spirituous solutions, the mucilage in great measure covering the pungency of the guaiacum: the mucilaginous folution, at first greyish or brownish, changes in a few hours to a fine blue or blueish green colour.

GUMMI.

GUM: a concrete vegetable juice; of no particular smell or taste; becoming viscous and tenacious when moistened with water; totally disfolving in water into a liquid more or less glutinous in proportion to the quantity of the gum; not dissolving in vinous spirits or in oils; burning in the fire to a black coal, without melting or catching slame; suffering no dissipation in the heat of boiling water.

I. Gummi Arabicum Pharm. Lond. & Edinb. Gummi acanthinum & thebaicum quibusdam. Gum-arabic: the gum, exuding from the Egyptian acacia tree, (Mimosa milotica Linn.) whose fruit affords the inspissated juice of that name; brought to us from Turkey in small irregular masses, of a clear whitish or very pale yellowish colour.

The medical character of gum-arabic is its glutinous quality; in confequence of which, it ferves to incraffate and obtund acrimonious thin humours, in tickling coughs, alvine fluxes, and other like diforders: Prosper Alpinus says, it is often used successfully by the Egyptians for restraining hemorrhagies. It is given chiefly in the form of powder,

from

from a scruple to a dram or two; and sometimes dissolved in water, in such proportion as not to make the liquor disagreeably slimy. An ounce renders a pint of water considerably glutinous: four ounces give a thick syrupy consistence. The solutions mingle equally with vegetable and with mineral acids, and with neutral saline mixtures; but on the addition of alkalies, fixt or volatile, the liquor grows turbid, and the dissolved gum separates.

Though the gum in its dry state is not affected by oily liquids, yet when softened with water into the consistence of a thick mucilage, it unites, by trituration, both with the sluid oils and the thicker balsams, so intimately, that the whole compound dissolves in water, without separation, into an emulsion or milky liquor; one part of gum-arabic, softened with an equal weight of water, is sufficient for rendering sour parts or more of oil or balsam dissoluble (a). The solid resins may in like manner be reduced into emulsions, by grinding them thoroughly with powdered gum, and adding the water by degrees. By these means, all resinous and oily bodies may be dissolved in watery liquors, and thus excellently fitted for being taken in a liquid form, without any alteration in their smell, taste, or virtue. These emulsions, like the solutions of the gum itself, mingle uniformly with acids and neutral salts; but on the mixture of any alkali, they suffer immediately a separation of their parts.

2. Gummi senegalense Pharm. Parif. Gummi senegal vel senica. Gum-senegal or senica: a gum brought from the island Senegal on the coast of Africa, said to be the produce of a tree of the same genus with that which affords the gum-arabic, acacia siliquis compressis Pharm. Parif. Mimosa senegal Linn. Greatest part of this gum is in larger and darker coloured masses than the arabic, and not smooth like it, but rough on the outside. In quality, the two sorts are scarcely different from one another, or from that which exudes from plum, cherry, and other trees among ourselves: in the shops, the clearer pieces of the gum-senegal generally supply the place of the more costly gum-arabic. It is supposed that the Senegal gum is the strongest and most substantial, and the Arabic the purest and finest.

⁽a) See on this subject the medical observations and enquiries published by a society of physicians in London, vol. i. art. 28. p. 358.

3. TRAGACANTHA Ph. Lond. Gummi tragacantha Ph. Edinb. Gummi tragacanthæ & dragacanthæ. Gum-tragacanth or dragant: the gum exuding from a prickly bush of the same name, (tragacantha C. B. goats thorn; tragacantha cretica incana flore parvo lineis purpureis striato Tour. Astragalus Tragacantha Linn.) which grows wild in the warmer climates, and endures the cold of our own, but does not here yield any gum. This commodity is brought chiefly from Turkey, in irregular lumps, or long vermicular pieces bent into a variety of shapes: the best sort is white, semitransparent, dry, yet somewhat soft to the touch.

Gum-tragacanth differs from all the other known gums, in giving a thick confishence to a much larger quantity of water; and in being much more difficultly disfoluble, or rather disfolving only imperfectly. Put into water, it slowly imbibes a great quantity of the liquid, swells into a large volume, and forms a soft but not fluid mucilage: if more water be added, a fluid solution may be obtained by agitation, but the liquor looks turbid and wheyish, and on standing the mucilage subsides, the limpid water on the surface retaining little of the gum. Nor does the admixture of the preceding more soluble gums promote its union with the water, or render its dissolution more durable: when gumtragacanth and gum-arabic are dissolved together in water, the tragacanth seems to separate from the mixture more speedily than when dissolved by itself.

Tragacanth is usually preferred to the other gums for making up troches, and other like purposes, and is supposed likewise to be the most effectual as a medicine; but on account of its impersect solubility is unfit for liquid forms. It is commonly given in powder with the addition of other materials of similar intention: thus to three parts of gum-tragacanth, are added three of gum-arabic, three of marshmallow root, one of liquorice, one of starch, and six of sugar.

Pulv. e trag. comp. Ph. L.

HEMATITES.

FIEMATITES LAPIS Pharm. Lond. BLOOD-STONE: an elegant iron ore, found either along with the other ores of that metal, or in distinct mines by itself; in irregular masses, convex on one side and angular on the other, generally of a dark reddish colour with more or less

less of a yellowish cast, very heavy, and of great hardness. Broken longitudinally, it exhibits a number of striæ converging to the smaller end: the transverse fracture appears of a granulated texture. Exposed to a moderately strong fire, it falls by degrees into scales; and in this state is attracted by the magnet, and gives out its iron to acids, both which have little action upon it in the mass or when barely reduced to powder.

This mineral, finely levigated, and freed from the groffer parts by Hæmat. lap. washing over with water, has long been recommended in hemorrhagies, Pharm.Lond. fluxes, uterine obstructions, &c. in doses of from one scruple to three or four. We presume, that it is not expected to act any otherwise than by virtue of its ferrugineous matter; and that pure iron itself, or its preparations, are preferable to a stony ore of variable and uncertain contents.

HEDERA

HEDERA ARBOREA Pharm. Edinb. (a) & C. B. Hedera communis major J. B. Hedera Helix Linn. Ivy: an evergreen plant, climbing and spreading on trees and old walls; with numerous slender twigs, and angular leaves. When grown old, the angles of the leaves disappear, the plant becomes erect, produces flowers, small and herbaceous, in autumn, and clusters of black berries in winter.

THE leaves of ivy have a very nauseous taste, and little or no smell. Haller fays, they are commended in Germany against the atrophy of children. Among us they are fometimes applied externally by the common people to running fores, and for keeping iffues open.

The berries were supposed by the ancients to have a purgative and emetic quality; and an extract made from them by water is called by Quercetanus extractum purgans. Later writers have recommended them in fmall doses, as alexipharmac and fudorific: it is faid, that in the London plague, the powder of them was given in vinegar or white wine with good fuccess. It is probable, however, that the virtue of this compound was rather owing to the vehicle than to the ivy-berries.

(a) Expunged.

From the stalks of this plant exudes, in the eastern countries, and sometimes in our own(a), a resinous juice, which has been directed as an officinal, under the name of gummi bederæ. This is in hard compact masses, externally of a reddish brown colour, internally of a bright brownish yellow with reddish specks or veins, of a vitreous glossiness, but not pellucid, of a light agreeable smell when rubbed or heated, and a resinous subastringent taste. Rectified spirit receives from it a deep brownish red tincture, and dissolves near three sourths: near one fourth remains undissolved after the successive action of water and spirit. It has been recommended as corroborant, and resolvent, in cachexies, and uterine obstructions; but has rarely been otherwise made use of than as an ingredient in plasters: nor does it appear to have any virtues that common resin does not posses in at least an equal degree.

HEDERA TERRESTRIS.

HEDERA TERRESTRIS Pharm. Lond. & Edinb. Hedera terrestris vulgaris C. B. Corona terræ Lobel. Chamæciss; Chamæclema. Glechoma hederacea Linn. GROUND-IVY: a low, somewhat hairy, creeping plant: with square stalks; roundish or kidney-shaped leaves set in pairs at the joints; in the bosoms of which come forth clusters of blue labiated flowers, whose upper lip is cloven and turned backwards. It is common in hedges and shady places, slowers from April to near the end of summer, and is generally found green all the winter.

This herb has a quick, bitterish, warm taste; and an aromatic but not very agreeable smell, which is in great measure dissipated by drying. It is supposed to be particularly serviceable in disorders of the breast, for cleansing and healing ulcerations in general, resolving coagulated juices, and purifying the blood. It has been customary to macerate the herb for a diet drink, in malt liquors; to which it readily communicates its virtue, and which it remarkably helps to fine down. It gives out its virtues also, together with a yellowish brown tincture, by infusion in water: on inspissating the filtered liquor, only the unpleasant smell of the herb exhales, its more valuable parts remaining concentrated in the

extract; which, on being tasted, impresses first a kind of sweetness, then a degree of bitterness, and soon after discovers a strong pungency. To rectified spirit of wine it yields its virtue only in part: the deep green spirituous tincture has but little of the subtile pungency of the watery insusant and the brownish yellow extract, obtained by inspissating the tincture, is much weaker in taste, as well as less in quantity, than the extract made with water.

HELLEBORUS ALBUS.

HELLEBORUS ALBUS Pharm. Lond. Veratrum Pharm. Edinb. Elleborum album Matth. Helleborus albus flore subviridi C. B. Veratrum flore subviridi Tourn. Veratrum album Linn. White Hellebore: a plant with large oval ribbed leaves, crumpled and plaited as it were, set alternately on a round firm stalk, and embracing it by a tubulous basis: in their bosoms, towards the top, appear clusters of hexapetalous greenish white flowers, followed each by three slat pods containing whitish triangular seeds: the root is short, commonly near an inch thick, with numerous sibres hanging from it, externally of a brownish colour, internally white. It is common in mountainous places in Germany, Switzerland, and some other parts of Europe.

WHITE hellebore root has, when fresh, a disagreeable smell; but as brought into the shops, scarcely any: its taste is nauseous, bitterish, acrid, very penetrating and durable. The juice of the fresh root, in taste extremely acrid, is said, when mixed with the blood, to act as a poison. The powder of the dry root is sometimes mixed with external applications for destroying cutaneous insects: snuffed up the nose, in small quantity, it proves a violent sternutatory, and in this intention is sometimes used in lethargic and other disorders.

This root, taken internally, in doses of ten or fifteen grains, operates with great violence both upwards and downwards, and has sometimes brought on convulsions and other terrible symptoms: Hoffman observes, that it peculiarly affects the fauces, producing a strangulation and danger of suffocation, with extreme anxiety. It has been chiefly employed, and that but seldom, in some kinds of maniacal cases, as a last resource; in which it is said to have taken place after the stronger of

the

the antimonial preparations had been given without effect. In minute doses, it has been sometimes used for acuating other purgatives and emetics; and sometimes also as an alterative or deobstruent in stubborn chronical distempers. In this last intention it is doubtless a medicine of great power, but its effects have not yet been sufficiently ascertained, to

entitle it to a place in general practice.

Infusions of white hellebore root in water, and the extracts obtained by inspissating them, in colour yellowish, in taste less acrid than the root itself, appear to operate with less violence. Hermann, who makes the dose of the hellebore in powder from ten grains to fifteen, directs an infusion of a dram; and of the extract he gives about as much as of the root in substance. In the shops, the active parts of the root, extracted by water, are thence transferred into honey: a pound of the root is macerated three days in four pints of water, then boiled a little, the decoction, pressed out and strained, mixed with three pounds of clarified honey, and the mixture boiled down till the water has exhaled and the honey appears of its original confistence. This preparation is used fometimes, but rarely, in glysters: a similar combination of the active matter of the hellebore with vinegar and honey, reduced to the confiftence of a syrup, is recommended by Gesner, in an express treatise on this plant, as a fafe internal medicine in phlegmatic diforders, particularly those of the breast, and faid to promote, without disturbance, all the natural excretions: preparations of this kind, however, have one great inconvenience, that they do not admit fo much precision, in regard to the strength, as is requisite in a medicine of so great activity.

Mel helleboratum Ph. L.

Tinctura veratri

+ Ph. Lond. † Ph. Edinb. A tincture of white hellebore made in proof spirit is likewise milder, both in taste and in operation, than the root in substance: a tincture drawn with a quart †, or two pounds and a half ‡ of proof spirit from eight ounces of the root, is kept in the shops, and given sometimes in doses of a few drops as an alterative, and one or two drams and upwards as a cathartic and emetic. On inspissating this tincture, the remaining extract is found to taste stronger than that made with water, though not quite so pungent as the root itself.

HELLEBORUS NIGER.

HELLEBORUS NIGER Pharm. Lond. & Linn. Melampodium & Helleborus niger Ph. Ed. Helleborus niger flore roseo C. B. Black Helleborus niger plore roseo C. B. Black externally stream the pedicles of the leaves and flowers, which are pedicles in the flower is large, indented from about the middle to the extremity; the flower is large, naked, pentapetalous, of a pale rose colour, with numerous stamina in the middle, which are followed by five or six pods full of shining blackish seeds, the petala continuing, and changing greenish: the root consists of numerous sibres, hanging generally from a knotty head, externally of a black colour, internally white. It is perennial, grows wild in the mountainous parts of Germany, and flowers in our gardens in January.

BLACK hellebore root, in doses of from ten grains to half a dram, proves a very strong, though not very violent cathartic. The hellebore of the ancients, which was never ventured on without extreme caution and as a last resource, appears to have been a different species from ours, much larger and of more violent operation, called by Tournesort belleborus niger orientalis, amplissimo folio, caule præalto, flore purpurascente, which is still said to be found in plenty about mount Olympus, and in the island of Anticyra, celebrated of old for the production of this reputed antimaniacal drug. In the present practice, this root or its preparations are used sometimes as a purgative in cases where the stronger cathartics are required; but oftener in small doses as an attenuant and deobstruent. It is found particularly serviceable against obstinate suppressions of the menstrual purgations, in plethoric habits and sanguine constitutions, where chalybeates are inessectual or injurious.

The taste of this root is bitter and pungent: chewed for a few minutes, it seems to benumb the tongue. The sibres are stronger in taste, and medicinally more active, than the tuberous head; and the cortical part of this, than the internal. It is said to prove purgative when applied only externally, in somentations, to the belly or seet. Water extracts by coction, and proof spirit by digestion, nearly all the virtue of the hellebore: rectified spirit takes up chiefly the irritating

refinous

refinous part. After due coction in water, it gives out little to spirit: but after repeated digestions in pure spirit, it is said still to yield to water a considerable proportion of a diuretic mucilaginous substance: the quantity of watery extract amounts to about one third of the root, the spirituous only to about one sifth.

Extractum helleb. nigri Pharm. Lond. & Edinb.

The extract made with water is the best and safest preparation of this root when defigned for a cathartic; as it contains both the purgative and diuretic parts of the hellebore, and as the irritating power of its active matter is confiderably abated by the boiling: it may be given from eight or ten grains to a scruple or more, but is used oftener in conjunction with other materials of fimilar intention, than by itself. * An extract of black hellebore, made in a very operofe way by macerating the root in rectified spirits, and wine, and then strongly expressing the liquor, which is afterwards repeatedly mixed with water, and evaporated to a due confistence, is the principal ingredient of a celebrated medicine for the dropfy in France, known by the name of Bacher's tonic pills. The other articles are an extract of myrrh, and powder of Carduus benedictus. These pills are said to produce a very copious evacuation both by stool and urine; and at the same time to brace and strengthen the solids. Their use is prohibited to persons of a tense fibre, or who are suspected to have internal suppurations. They are to be used with great caution also in hot climates. In dropfies with relaxation, they are faid to produce the happiest effects.

Tinctura melampod. Pb. Lond.† five helleb. nigri Pharm. Edinb.‡ A tincture made in proof spirit appears the most eligible preparation for the purposes of an alterative and deobstruent: four ounces of the root may be digested in a quart, or two pounds and a half of the spirit, with the addition of thirty; or forty; grains of cochineal to render the colour more sightly; and the filtered tincture given to the quantity of a tea-spoonful twice a day, in warm water or any other convenient vehicle. Dr. Mead informs us, that in menstrual obstructions the power of this medicine is so great, that when, from an ill conformation of the parts, or other causes, the expected discharge does not succeed upon the use of it, the blood is so forcibly propelled, as to make its way through other passages.

HEPATICA.

HEPATICA terrestris sive Lichen Ph. Edinb. (a) Lichen petræus latisolius sive hepatica fontana C. B. Hepatica terrestris vulgaris seu lichen ossicinarum Raii hist. i. 124. Jecoraria. Marchantia polymorpha Linn. Liver-wort: a species of moss; consisting of numerous obtusely laciniated leaves, lying over one another, marked on the surface with white tubercles reticularly disposed through the green substance of the leaf, shooting out underneath fine capillary roots: from the extremities of the leaves issue clear whitish pedicles, bearing globular bodies, which when ripe contain black minute seeds like dust. It is perennial, grows on moist stony places, and runs up to seed in March or April.

This moss is recommended as an aperient, resolvent, and purisher of the blood. From the penetrating though mild pungency and bitterishness of its taste, sinking as it were into the tongue, it promises to be a plant of no inconsiderable virtue, though in this country at present disregarded. It gives out its active matter both to watery and spirituous menstrua.

HEPATICA NOBILIS.

HEPATICA NOBILIS Pharm. Edinb.(a) Hepatica trifolia five berba trinitatis Pharm. Parif. Trifolium bepaticum flore fimplici C. B. Anemone Hepatica Linn. HEPATICA, HERB-TRINITY: a low plant, without any other stalk than the pedicles of the leaves and flowers: the leaf is cut, not very deep, into three lobes, entire about the edges: the flower is commonly blue, sometimes reddish or white, hexapetalous, set in a three-leaved cup, with numerous stamina in the middle, which are followed by a cluster of whitish seeds. It is perennial, grows wild on gravelly shady grounds in Germany and other parts of Europe, and flowers in our gardens in March or sooner.

This herb is a mild restringent and corroborant; in which intentions, infusions of it have been drank as tea, or the powder of the dry leaves

(a) Expunged.

given to the quantity of half a spoonful at a time. Its astringent matter is dissolved equally by water and spirit; and on inspissating the filtered tinctures, remains entire in the extracts: it is found however, even when thus concentrated, to be still but weak. The watery extract is larger in quantity, and proportionably weaker in taste, than the spirituous.

HERMODACTYLUS.

HERMODACTYLUS Pharm. Parif. HERMODACTYL: the root of a species of colchicum, brought from Turkey; of the shape of a heart, flatted on one side, with a surrow on the other; of a white colour; compact and solid, yet easy to cut or powder.

This root has a viscous, sweetish farinaceous taste, and no remarkable smell. It was ranked by the ancients among the cathartics, but such as we now meet with in the shops does not appear to have any purgative virtue. Alpinus relates, that hermodactyls, the same with what are sold in Europe, are eaten by the Egyptian women to the quantity of several roots at a time; that they do not move the belly, or produce any ill effect: that they are supposed to be very nutritious, and contribute to procure the fatness and plumpness there admired (a). Those, who report them to be purgative, have probably ascribed to the hermodactyls a part of the effects of the substances which were joined to them; for, being acknowledged flow in operation, they have been commonly acuated with aloes and other cathartics. They have long been an entire stranger to practice, and the colleges both of London and Edinburgh have now deservedly expunged them from their catalogues of officinals.

HIBERNICUS LAPIS.

HIBERNICUS LAPIS & Tegula bibernica & Ardesia bibernica Pharm. Edinb. (b) Hardesia Pharm Paris. IRISH SLATE: a kind of slate, or very soft stone, found in different parts of Ireland; in the mass of a blueish black colour and staining the hands; when powdered, pale or whitish at first, and in keeping growing black; in the fire, yielding

(a) De medicina Ægyptiorum, lib. iii. cap. 16. & lib. iv. cap. 1.

(b) Expunged.

fulphureous fumes, and acquiring a pale red colour with an additional hardness.

This mineral appears from Dr. Rutty's experiments in his fynopfis of mineral waters, to be a matrix of ferrugineous vitriol; which it discovered by its taste, and by the black colour which infusions of it struck with galls. The specimens I examined, procured from our shops, had no vitriolic taste, and infusions of them in water suffered no visible change from galls: but that they contained materials from which vitriol is producible, appeared from the fulphureous fumes they emitted in the fire, and from their giving out, when burnt, a calx of iron to aqua regis; the iron discovered itself by the tincture striking a blue colour with the lixivium described under the article ferrum: their burning hard shews their earthy matter to be of the argillaceous kind. It may therefore be prefumed, that this fosfil consists of argillaceous earth impregnated, like the pyritæ, with fulphur and iron; and that, like the pyritæ alfo, it is capable of becoming vitriolic by long exposure to the air and moisture. It has been sometimes taken by the common people, powdered and mixed with spruce beer, against inward bruises; but its medicinal use in any intention is not much to be commended, on account of the variability of its qualities.

HIPPOSELINUM.

HIPPOSELINUM five Smyrnium Ph. Edinb. (a) Hipposelinum theophrasti vel smyrnium dioscoridis C. B. Macerone & olus atrum & herba alexandrina quibusdam. Smyrnium Olusatrum Linn. ALEXANDERS: an umbelliserous plant, with leaves like those of smallage but larger: producing thick, roundish, striated, black seeds. It is biennial, grows wild about the sea coasts, and sides of rivers, and slowers in May and June.

This plant is nearly fimilar in quality to smallage, but somewhat stronger both in smell and taste: it was formerly blanched in our gardens for culinary use, but has now given place to celeri, which is more grateful. The seeds are bitterish and aromatic, and have been now

(a) Expunged.

and then employed, like the other warm feeds, as carminatives, sto-machics, and aperients: they give out their virtue imperfectly to water, completely to rectified spirit: in distillation with water, they yield a small quantity of essential oil, smelling agreeably of the seeds, and in taste moderately pungent. The roots are bitterer than the seeds, and stand recommended as resolvents, diuretics, and emmenagogues: they yield, on incision, a whitish juice, which is said, when inspissated, to approach in taste to myrrh; whence the plant has been called, from one of the names of that gummy-resin, smyrnium.

HORMINUM.

HORMINUM SATIVUM five Sclarea Pharm. Edinb.(a) Horminum sclarea dictum C. B. Gallitrichum sativum J. B. Salvia Sclarea Linn. Garden clary: a whitish green, slightly hairy plant, with square stalks, and large wrinkled oblong somewhat heart-shaped leaves: both the leaves, and the divisions and subdivisions of the branches, stand in pairs: on the tops grow long spikes of blueish labiated flowers, at the origin of which are little concave purplish leaves without pedicles: the upper lip of the flower is long and arched, the lower smaller and cut into three segments, the middlemost of which is hollowed like a spoon. It is biennial, a native of the warmer climates, cultivated with us in gardens, and slowers in July and August.

The leaves and feeds of clary are recommended as corroborants and antispasmodics; particularly in the fluor albus and other like weaknesses, and in hysterical complaints. They have a bitterish warm taste; and a strong smell, of the aromatic kind, but to many persons not agreeable. The leaves discover to the touch a large quantity of unctuous resinous matter, in which the virtue of the herb appears to reside, which is readily dissolved by rectified spirit, and which, on inspissating the sine green tincture, remains nearly entire in the dark brownish extract: this extract smells more agreeably than the herb in substance, and is in taste moderately warm, bitterish, and pungent. Water takes up likewise by insusion great part of the active matter of the clary, and carries off its

whole flavour in evaporation, leaving a weak, disagreeably bitterish, roughish extract. In distillation with water, there arises both from the leaves and seeds, a small quantity of essential oil, smelling strongly of the clary: from sixty-four ounces, or sive hundred and twelve drams, of the seeds, was obtained only about one dram of oil. The leaves or seeds, fermented with malt liquors, are said to remarkably increase their inebriative quality.

HYOSCYAMUS.

HYOSCYAMUS Pharm. Parif. HENBANE: a plant with foft, hairy, oblong, deeply notched leaves; and bell-shaped flowers cut into four segments, followed by irregular cup-like capsules, whose cover falling off discloses numerous small seeds in two cells.

- 1. HYOSCYAMUS Pharm. Edinb. Hyoscyamus niger vel vulgaris C. B. Hyoscyamus niger, sive apollinaris herba, altercum arabum Lob. Faba suilla. Hyoscyamus niger Linn. Black henbane: with large leaves joined close to the stalk; dusky coloured flowers; and greyish seeds. It is annual, grows wild in waste rich grounds, and flowers in June.
- 2. HYOSCYAMUS ALBUS Linn. Hyoscyamus albus major vel tertius dioscoridis & quartus plinii C. B. White henbane: with smaller and woollier leaves set on pedicles; white flowers; and whitish seeds. It is annual, and a native of the southern parts of Europe.

THESE plants have a fetid fmell, of the narcotic kind: on the organs of taste they make no considerable impression, the leaves being little other than herbaceous, the seeds somewhat mucilaginous, and the roots sweetish. The leaves, applied externally, in the form of cataplasm, fomentation, or unguent, are said to be refrigerant, discutient, and to abate not only instammatory but rheumatic pains. All the parts, when taken internally, though in no great quantity, prove highly narcotic, and occasion violent disorders of the senses, sometimes of long duration, and sometimes fatal. There are several instances also of the senses being strongly affected by the effluvia of the plant; and by the vapour that arises upon scorching it over the fire, which has by some been imprudently

dently directed to be received into the mouth against toothachs (a). The effect of small doses, insufficient to do harm, are not well known. The seeds have been given from two or three grains to a scruple, and said to have proved beneficial in spittings of blood and thin sharp defluxions on the lungs, and likewise in some convulsive disorders. The present practice, however, has deservedly rejected a medicine of such suspicious qualities, and which appears to be at best but a precarious succedaneum to the products of the poppy. It is said, that the seeds are the mildest part, and the root the strongest; that the black fort is in all its parts stronger than the white; and that the seeds are the more deleterious, the more they approach to blackness.

HYPERICUM.

HYPERICUM Pharm. Lond. Hypericum vulgare C. B. Androsæmum sive perforata Gesn. Hypericum perforatum Linn. St. John's wort: a plant with slender round woody reddish stalks, which have two fine ridges, or sharp edgings, opposite, alternately, from joint to joint; small oblong obtuse leaves, set in pairs without pedicles; and numerous gold-coloured pentapetalous flowers on the tops of the branches, sollowed by little rough blackish husks, each of which is divided into three cells full of minute seeds. It is perennial, grows wild in woods and uncultivated places, and slowers in June and July.

This plant has been recommended as a medicine of peculiar efficacy in hysterical and hypochondriacal disorders, and alienations of mind; from its supposed virtue in which cases, it received the name of fuga damonum. It promises to be of some use as a mild detergent and corroborant, discovering to the senses a resinous, bitterish, balsamic impregnation. The leaves, viewed against the light, exhibit numerous transparent points, which are found to be little vesicles full of essential oil or resinous matter; in distillation with water, the oil separates and rises to the surface, approaching in some degree to that of turpentine. About

⁽a) Vide Lindestolpe, De venenis, cap. x. thes. 20. edit. Stentzel. p. 559. Konig, Regnum vegetabile, sett. iv. p. 869. Hoyer, Atta physico-medica naturæ curiosorum, vol. v. obs. 69. p. 260. Hossman, Philosophia corporis humani morbosi, p. ii. sap. vii. §. 7. Oper. omn. i. 223. Haller, & auctores ab eo citatos, Stirp. helvet. p. 513.

the edges of the flower are observed black points, and on the feed veffels fmall tubercles, which appear to be fimilar oily veficles: the tops, when the feeds are formed, have the strongest terebinthinate smell, and yield in distillation the greatest quantity of oil. The flowery tops give a deep vellowish red tincture to rectified spirit, and a paler red to expressed oils: this colour does not appear to proceed from the substance of the leaves or flowers, but from the refinous juice in the veficles above mentioned. A tincture of the flowers in oil olive, made by macerating four ounces of Ol. hyperici the full blown flowers, fresh gathered and freed from the cups, in a quart of oil, till the oil is sufficiently coloured, is kept for external purposes in the shops, but very rarely made use of.

HYPOCISTIS.

HYPOCISTIDIS SUCCUS Pharm. Lond. The juice of HYPO-CISTIS: an inspissated juice, of a firm consistence and a bright black colour; prepared from a certain fleshy juicy vegetable production (bypocistis sub cisto C. B. Asarum Hypocistis Linn.) which, in the warmer climates, grows up from the root of different kinds of the ciftus or rockrose, three or four inches high, easy to break, cloathed with scales, bearing a number of little bell-shaped flowers on the top, but no leaves.

This juice is a mild aftringent, of no particular smell or flavour. It is looked upon as fimilar to the Egyptian acacia; from which, however, it differs remarkably in its pharmaceutical properties, the hypocistis being almost totally dissoluble in rectified spirit of wine, whereas acacia gives out little or nothing to that menstruum. It is at present scarcely otherwise made use of among us, than as an ingredient in some of the old compositions.

HYSSOPUS.

HYSSOPUS Pharm. Lond. & Edinb. Hyssopus officinarum carulea five spicata C. B. Hyssop officinalis Linn. Hyssop: a low shrubby plant; with brittle branched stalks, square when young, and round when they grow woody; oblong narrow dark green leaves fet in pairs; and loofe spikes of labiated blue flowers, whose upper lip is cloven and turned upwards, standing in rows, towards the tops of the stalks, generally rally all on one fide, in long striated cups. It is perennial, cultivated in gardens, and flowers in July and August.

The leaves of hysfop have an aromatic smell, and a bitterish moderately warm taste. They give out their active matter both to water and rectified spirit, to the last most perfectly: the watery insusions are of a brownish or greenish yellow, the spirituous tinctures of a dark blackish green colour. On inspissating the spirituous tincture, very little of the slavour of the herb exhales or distils with the menstruum: the remaining extract is bitterish and very warm, and discovers a penetrating pungency, somewhat like that of camphor. Water, distilled from the fresh herb, is found pretty strongly impregnated with its slavour: an essential oil separates and rises to the surface, to the quantity of about an ounce from six pounds of the leaves, in smell exactly resembling the hyssop, in taste very pungent, in colour, when newly distilled, yellowish with a slight cast of green, which by age changes to a brownish: the decoction, remaining after the distillation, is disagreeably roughish, bitterish, and subsaline.

This plant is accounted particularly ferviceable, as an attenuant, corroborant, and expectorant, in humoural asthmas, coughs, and other disorders of the breast and stomach unaccompanied with inflammatory symptoms: in these cases, insusions of the leaves, which are not unpalatable, may be sweetened with honey or sugar, and drank as tea. The distilled water, by some made choice of as a basis for pectoral mixtures and juleps, does not appear superiour or equal to the insusion.

JACOBÆA.

JACOBÆA Pharm. Parif. Jacobæa vulgaris laciniata C. B. Senecio major five flos fancti jacobi Matthiol. Senecio Jacobæa Linn. RAGWORT: a plant with a firm round stalk generally purplish; oblong dark green leaves, deeply jagged almost to the rib, and the sections jagged again and somewhat crumpled: on the tops of the stalks grow umbel-like clusters of yellow flowers, of the radiated discous kind, followed by small oblong seeds winged with down. It is perennial, common in uncultivated fields and by road sides, and slowers in July.

THE

The leaves of ragwort have a roughish bitterish subacrid taste, extremely nauseous, far different from that of the herbaceous groundsel to which they have by some been accounted similar. Simon Paulli relates, that they were found of great service in an epidemic camp dysentery, many soldiers having been cured of that disease by drinking a decoction of them; and expresses some concern, that a medicine of so much efficacy should be at the same time so very disgussful to the palate. This however is an inconvenience that may be easily palliated: the active matter of ragwort, whatever its virtue may be, is dissolved both by water and by rectified spirit, and on inspissating the filtered tinctures, remains concentrated in the extracts; which may be taken, without offence to the palate, in the form of a bolus or pill. The spirituous extract is in less quantity than the watery, and proportionably stronger in taste; though rather less nauseous than the herb in substance.

JALAPIUM.

JALAPIUM Pharm. Lond. Jalapa Pharm. Edinb. Mechoacanna nigra. JALAP: the dried root of the mirabilis peruviana or marvel-ofperu, mirabilia peruviana Gerard. solanum mexiocanum flore parvo C. B. Convolvulus Jalapa Linn.* (a) a plant with thick, fleshy radish-like roots; jointed stalks and branches; acuminated somewhat oval leaves set in pairs; and elegant, numerous, monopetalous, funnel-shaped flowers, purple, yellow, white, or diverfly variegated, standing in double cups, of which the innermost incloses the flower and the outer surrounds its basis: each flower is followed by a wrinkled, roundish, pentagonal umbilicated fruit, about the fize of a pepper corn, including a white kernel. It is perennial, a native of the West Indies, and cultivated in our gardens on account of the beauty and duration of its odoriferous flowers, which open only during the night, and of which it produces continual fuccessions from June or July till checked by frosts; at which time the roots, which do not endure our winter, are taken up, and preserved in fand till spring. Whether the roots produced here are equivalent in virtue to those which are brought from abroad, has not, that I know of, been tried.

^{* (}a) The later botanists are not perfectly agreed concerning the genus of the plant producing jalap. Linnæus first made it a mirabilis; and Bergius now gives it as the mirabilis dichotoma Linn.

The officinal jalap roots come from the province of Xalapa in New Spain; in thin transverse slices, solid, hard, weighty, of a blackish colour on the outside or cortical part, internally of a dark greyish with several black circular striæ: the hardest, darkest coloured, and those which have the most of these resinous veins, are the best. Slices of bryony root, which are said to be sometimes mixed with them, may be distinguished by their whiter colour and less compact texture.

This root has fearcely any smell, and very little taste upon the tongue: swallowed, it affects the throat with a slight kind of pungency and heat. Taken in doses of a scruple or half a dram, it proves an effectual and in general a safe purgative; very rarely occasioning any severe gripes or nausea, which too frequently accompany the other strong cathartics. Some have prohibited the use of this cathartic to children; probably on no very good foundation. Young children, from the laxity of their solids, and the soft lubricating quality of their food, generally bear these kinds of medicines better than adults, and adults of a spongy, lax, or weak habit, better than the rigid or robust. Few, if any, of the strong resinous purgatives are in either case more innocent than jalap.

Jalap root, digested in as much rectified spirit as will cover it to the height of about four singers, gives out greatest part of the resinous matter in which its activity resides, and tinges the menstruum of a yellowish brown colour. On inspissating the filtered tincture to about one half, and adding to the remainder a proper quantity of water, the liquor becomes milky, and on standing deposites the pure resin. This preparation, given by itself, irritates and gripes violently, without proving considerably purgative: thoroughly triturated with testaceous or other powders, or with soap; or ground with almonds or powdered gum-arabic, and made into an emulsion with water; or dissolved in rectified spirit, and mixed with a proper quantity of syrup, that the solution may bear being diluted with watery liquors without precipitation; it purges, in doses of eight or ten grains, as effectually, and for the most part as mildly, as the jalap in substance.

The jalap remaining after sufficient digestion with spirit, has no cathartic virtue: boiled in water, it gives out a mucilaginous substance, which operates only by urine. Water applied at first takes up a portion of the resin along with the gum, and hence the watery decoction and extract

extract prove weakly cathartic as well as diuretic: the root still retaining great part of its refin, so as to purge considerably. The refinous and gummy parts may be united into one extract, by first drawing a tincture from the powdered root with rectified spirit, then boiling the residuum in fresh quantities of water, evaporating the decoctions till they begin to grow thick, mixing in by degrees the tincture inspissated to a like thickness, and continuing a gentle heat till the whole is reduced to a due confistence. This extract may be taken by itself in doses of twelve grains Extractum or more: the gummy matter of the jalap being sufficient to divide the jalappæ Ph. refin and prevent its too violent irritation.

The proportion of active matter differs greatly in different parcels of jalap; fixteen ounces of fome forts yielding hardly two of refin, while the same quantity of others affords three or four. Hence the extracts of jalap appear preferable to the root in fubstance, not only on account of the dose being rendered smaller by the rejection of the woody parts, but likewise as being more uniform and certain in strength. Tinctures of jalap made in proof spirit are nearly similar in quality to the gummyrefinous extract, this menstruum taking up both the refinous and gummy parts of the root: these preparations, made from different kinds of jalap, will vary in strength somewhat more than the solid extract or refin, but not fo much as some have suspected, or as the roots in substance; for in the proportions usually employed, the proof spirit does not take up the whole of the virtue of any kind of jalap, and perhaps it does not extract much more from one kind than from another, provided the jalap be of Tind. jalap. moderate goodness. If three + or four tounces of jalap be digested in + Ph. Edinb. a pint of proof spirit, the residuum will still give out a portion of resinous matter to rectified spirit, and this refin will be in greater quantity in proportion as the root itself was the more refinous.

I Pb. Lond.

ICHTHYOCOLLA.

ICHTHYOCOLLA Pb. Edinb.(a) ISINGLAS OF FISH-GLUE: a folid glutinous substance, prepared from a fish of the sturgeon kind, caught in the rivers of Russia and Hungary. The skin, fins, &c. are boiled in water, the decoction inspissated to a due consistence, and then

(a) Expunged.

poured out so as to form thin cakes; which are either exsiccated in that form, or cut while soft into slices and rolled up into spiral, horse shoe, and other shapes. The best is in thin, clear, and almost transparent pieces.

* A different account of the formation of ifinglass is given by Mr. Jackson in *Philos. Trans.* Vol. LXIII. Part I. He afferts that the solution of animal substances of every kind gives glue, not ifinglass; that this last is nothing more than certain membranous parts of sishes, as the air-bladder, intestines, peritonæum, &c. in their entire state, only freed from their natural mucus and adhering matters, and rolled and twisted into the forms in which we get it.

ICHTHYOCOLLA is one of the purest and finest of the animal glues, of no particular smell or taste. Beaten into shreds, it dissolves pretty readily in boiling water or milk, and forms a gelatinous substance, which yields a mild nutriment, and proves useful medicinally in some disorders arising from a sharpness and colliquation of the humours. A solution of it in water, curiously spread, whilst hot, upon silk, affords an elegant sticking plaster for slight injuries of the skin, not easily separable from the part by water, and scarcely inferiour to the more compounded one sold under the name of the ladies black plaster, in which different balsams and resins are joined to the ichthyocolla.

ILLECEBRA.

ILLECEBRA, Vermicularis, Piper murale, Sedum minus. Sempervivum minus vermiculatum acre C. B. Sedum acre Linn. Wallpepper or Stonecrop: a small plant, having its stalks covered with little sleshy conical leaves set thick together in the manner of scales: on the tops appear pentapetalous yellow flowers, each of which is followed by several pods full of small seeds. It is annual, grows on old walls and dry stony grounds, and slowers in July.

This plant has a very acrid tafte, and no remarkable smell: applied externally, it vesicates the part: taken internally, in no great quantity, it proves strongly emetic. Its active matter appears, from the accounts given by authors, to be in great part forced out along with the watery juice

juice by expression; to dissolve both in water and sermented liquors by insussion; and not to be dissipated, or not soon, by boiling. It is said to have been used with success in sundry chronical disorders (a), but its durable acrimony, and the great vehemence of its operation, have prevented its being received in practice.

IMPERATORIA.

IMPERATORIA Ph. Edinb. Imperatoria major C. B. Imperatoria astrutium Lob. & Linn. Astrantia Dod. Smyrnion hortense Trag. Struthium hodie vocatum Cord. Masterwort: an umbelliferous plant, with large winged leaves divided into three indented segments; producing thick oblong, striated seeds surrounded with a narrow leasy margin: the roots are oblong, thick, knobby, jointed, with several lateral sibres, brown on the outside and whitish within. It is perennial, a native of the Alps and Pyreneans, from whence we are supplied with roots supposed to be superiour to those which are raised in our gardens.

The root of imperatoria is a very warm and moderately grateful aromatic, nearly of the same nature with that of angelica. Insused in water, or digested in rectified spirit, it impregnates both menstrua strongly with its fragrant smell; the former weakly, the latter strongly, with its warmth, pungency, and bitterishness; the former with a muddy brownish, the latter with a bright yellow colour. On inspissating the spirituous tincture, very little of its slavour exhales with the spirit: the remaining deep yellow extract smells moderately of the root, and impresses on the organs of taste a considerable bitterness and glowing pungency. Water carries off in evaporation nearly all the specific slavour of the masterwort, leaving, in the dark brown extract, a nauseous bitterness with a slight degree of warmth or acrimony.

IPECACOANHA.

IPECACOANHA Ph. Lond. Ipecacuanha Ph. Edinb. Hipecacuanna; Radix braziliensis. Psychotria emetica Linn.*(b) IPECACO-

⁽a) Below, Eph. nat. euriof. dec. i. ann. vi & vii. obf. 22. Boerhaave, Hift. plant. p. 369.

^{*(}b) This is the name given it in the fupplement to Linnaus: it was formerly reckoned by him a species of Lonicera.

ANHA: a flender root, brought from the Spanish West Indies, in short pieces, variously bent and contorted, full of wrinkles and deep circular fissures, which reach quite down to a small whitish woody fibre that runs in the middle of each piece: the cortical part is compact, brittle, and looks smooth and resinous on breaking. Two forts of this root are met with in the shops, one brought from Peru, the other from Brazil; usually denominated from their external colour, the first whitish, grey, or ash-coloured, the other brown ipecacoanha. The first is generally preferred, being found to operate with the greatest certainty and mildness.

A root has been brought over under the name of white ipecacoanha, which has little or nothing of the virtues of the two foregoing: this is readily distinguished by its yellowish white colour, woody texture, and having no fissures or wrinkles. More dangerous abuses have sometimes been committed, by the substitution or mixture of the roots of an American apocynum, which have been found to operate with great violence both upwards and downwards, and in some instances, as is said, to prove satal: these may be known by their being larger than the true ipecacoanha, the sissures more distant, the intermediate spaces smoother, and more particularly by the colour of the medullary sibre, which in the poisonous roots is a deep reddish yellow, in the true ipecacoanha a whitish or pale greyish.

IPECACOANHA has scarcely any smell, unless during its pulverization or insussion in liquors, in which circumstances it emits a faint nauseous one: in chewing, the wrinkled cortical part proves bitterish and subacrid, and covers the tongue as it were with a kind of mucilage; the medullary woody fibre is nearly insipid, and gives out to menstrua very little active matter. Geosffroy observes, that in pulverizing considerable quantities, the finer powder that slies off, unless great care be taken to avoid it, is apt to affect the operator with a difficulty of breathing, a spitting of blood, a bleeding at the nose, or a swelling and inflammation of the eyes and face, and sometimes of the throat; and that these symptoms go off in a few days, either spontaneously, or by the affishance of venæsection. *In the Philosophical Transactions, Vol. LXVI. Part I. is a remarkable case of violent asthmatic fits in a lady caused by the effluvia of powdered ipecacoanha.

THIS

This root is the mildest and safest emetic that has yet been discovered; and may be ventured on almost in the lowest circumstances where the stomach requires to be unloaded. The common dose is from ten grains to a scruple and upwards: in the medical observations and enquiries published by a society of physicians in London, a great number of cases are mentioned, in which two grains operated sufficiently: in constitutions which bore vomiting ill, and which were greatly russled by the usual doses, two or three grains operated with great ease. Where it fails of operating upwards, it commonly purges, and sometimes considerably: in this intention it may be employed, in several cases, to advantage, in conjunction with other purgatives, to determine its action downwards: I have found sisteen grains of jalap, with two or three of ipecacoanha, purge more than twice the quantity of jalap by itself.

The ipecacoanha was first introduced, about the middle of last century, as a specific in dysenteries; and repeated experience has confirmed its efficacy in this distemper, not only when used as an emetic, but likewife when given in fuch fmall doses as scarcely to affect the groffer emunctories. In common dysenteric fluxes, it frequently performs a cure in a very short space of time; not by its exerting an astringent power, as some have supposed, for it does not appear to have any real aftringency; nor by its mucilaginous substance covering the intestines and incraffating thin humours, as others, with more plaufibility, have inferred both from its mucilaginous taste, and from the ropiness and fliminess which it manifestly communicates to the contents of the stomach; but apparently by promoting perspiration, the freedom of which is in these cases of the utmost importance, and an increase of which, even in a state of health, is generally observed to diminish the evacuation by stool. In common dysenteries, the skin is for the most part dry and tense, and perspiration obstructed: and indeed this obstruction, and the conversion of the perspirable matter upon the intestines, is very frequently the immediate cause of the disease. Most of the common diaphoretics pass off, in these cases, without effect: but ipecacoanha, if the patient after a puke or two be covered up warm in bed, brings on a free diaphoresis or a plentiful sweat, by which I have often known the distemper terminated at once.

In putrid or malignant dysenteries, or where the patient breathes a tainted air, it has not been found equally successful: it requires here to

be continued for feveral days, or repeated as an evacuant, with the further affiftance of rhubarb, cordial antifeptics, and mild opiates or aftringents. Where plentiful evacuation is necessary, or the offending matter lodged deep, and the operation can be borne without inconvenience, the ipecacoanha, as Dr. Pringle observes, is most advantageously given in small quantities at a time, and repeated at proper intervals, till a vomiting or purging comes on.

*In the spasmodic asthma, Dr. Akenside remarks, that where nothing contraindicates repeated vomiting, he knows no medicine so effectual as ipecacoanha. In violent paroxysms, a scruple procures great and immediate relief. For habitual indisposition, from three to five grains every morning, or from five to ten every other morning, may be given for a month or six weeks. It is equally useful where it does not vomit, as where it does. The relief seems owing to its general antispasmodic or relaxing property, of which its emetic operation is probably a particular consequence (a).

*In the Stockholm acts 1770, are several cases of uterine hæmorrhagies cured by \(\frac{1}{3} \) or \(\frac{1}{2} \) grain, rubbed with sugar, given every four hours or oftener. In one case, the hæmorrhage returned on discontinuing the medicine, and ceased on repeating it. These small doses had good effects in catarrhal coughs, even in those which attend consumptions; and if not beneficial, are at least not hurtful, in bloody coughs, in which vomiting has several times been observed to come on, without any increase of the hæmorrhage. They may be useful in peripneumony and pleurisy, in which cough is often the most troublesome symptom, and in which Seneka root (which in increased doses proves also emetic) has been so much recommended.

The emetic virtue of ipecacoanha refides in its refinous parts. By digesting the root in fresh quantities of rectified spirit, and inspissating the filtered tinctures, a resinous extract is obtained, to the quantity of about three ounces from sixteen, which, by itself, vomits strongly, and with great irritation: the residuum yields to water nearly four ounces of a soft tenacious mucilage, which has scarcely any sensible operation. If only a part of the resin be extracted, by slight digestion in a little highly rectified spirit, the remaining root proves more gentle, and rather purgative than emetic: in this state it is recommended by some in

dysenteries accompanied with a considerable fever, where the root with its natural quantity of refin might irritate too much; but as small doses of the root itself operate with all the ease and gentleness that can be wished for, this precarious method of weakening it does not appear adviseable.

By boiling it in water, a part of the refin is taken up with the mucilage; the extract amounting to about fix ounces from fixteen, and proving mildly emetic. The best menstruum for extracting the entire virtue of the root appears to be a mixture of one part of pure spirit with two or rather three parts of water: after sufficient digestion in this menstruum, neither water nor spirit took up any thing considerable from the remainder. In the shops wine is employed: an ounce of the root Vinum ipeis macerated or digested in a pint of canary + or fifteen ounces of cacoanhæ + Pb. Lond. mountain 1; to which some add a quarter of an ounce of dried orange 1 Ph. Edinb. peel + to cover the unpleasant flavour of the ipecacoanha. tinctures, in doses of from half an ounce or less to an ounce and upwards, prove mildly emetic.

IRIS.

IRIS: a perennial plant with long narrow fword-like leaves, standing edgewife to the stalk; and large naked flowers, divided deeply into fix fegments, of which, alternately, one is erect and another arched downwards, with three smaller productions in the middle, inclosing the stamina and pistil: the roots are tuberous, irregular, and full of joints.

I. IRIS NOSTRAS PURPUREA Ph. Edinb. (a) Iris vulgaris germanica five filvestris C. B. Iris germanica Linn. Flower-de-luce, common iris or orrice: with blue flowers, whose arched segments are bearded with a yellowish matter, standing several on one stalk higher than the leaves. It is a native of the mountainous parts of Germany, common in our gardens, and flowers in June.

THE roots of this plant have, when fresh, a disagreeable smell, and an acrid nauseous taste. They are a strong irritating cathartic; in which intention, their expressed juice has been given in hydropic cases, from

(a) Expunged.

one or two drams to three or four ounces, diluted largely with watery or vinous liquors, to prevent its inflaming the throat. The remarkable differences in the dose, as directed by different practical writers, appear to have proceeded from hence; that some employed the juice in its recent turbid state, loaded with the acrimonious cathartic matter of the root; others, fuch as had been depurated by fettling, and which had deposited, along with the feculencies, a great share of the active parts. By gently inspissating the juice, it is rendered less violent in cathartic power, and less liable to irritate and inflame; but becomes at the fame time too precarious in strength to be depended on: by inspissation to perfect dryness, its purgative virtue is almost, if not altogether, destroyed. The root itself loses also, in drying, its offensive smell, and its naufeous acrimony, and along with thefe its cathartic quality: in this state, it discovers a slight and not disagreeable pungency and bitterishness, accompanied with a kind of aromatic flavour, nearly of the same kind with that of the following species, but weaker and less grateful.

The blueish expressed juice of the flowers changes on being inspissated, especially if a little lime-water is added, to a fine green; and in this form is directed, in foreign pharmacopæias; for tinging some of the unctuous compositions called odoriferous or apoplectic balsams.

2. IRIS Ph. Lond. Iris florentina Ph. Edinb. & Linn. Iris alba florentina C. B. Florence orrice; supposed to be only a variety of the foregoing occasioned by difference of climate; distinguishable from it in our gardens, by the flowers being white, and the leaves inclining more to blueish. The shops are supplied from Italy with dried roots superiour to those of our own growth; in oblong statish pieces freed from the sibres and brownish bark, externally of a whitish colour with brownish specks, internally inclining to yellowish, easily reducible into a farinaceous yellowish white powder.

This root, in its recent state, does not seem to differ much from the preceding; being, like it, nauseous, acrimonious, and purgative, though not quite in so great a degree; and losing these qualities on being dried. The dry root, as met with in the shops, has an unctuous, bitterish, pungent taste, not very strong, but very durable in the mouth: and a light agreeable fmell, approaching to that of violets. It is used in perfumes; in sternutatory powders; for communicating a grateful flavour, somewhat like that of raspberries, to wines and to spirits; and medicinally in disorders of the breast, for attenuating viscid phlegm, and promoting expectoration. Its smell and taste are extracted both by water and rectified spirit, most perfectly by the latter. In distillation, it gives over with water the whole of its peculiar slavour, its bitterness and a slight acrimony remaining in the inspissated extract: the distilled water smells very agreeably, but no essential oil is obtained though some pounds of the root be subjected to the operation at once. Rectified spirit brings over a part of its violet smell, but little or nothing of its warmth or taste: the inspissated extract is a pungent, bitterish, balsamic mass, glowing in the mouth like pepper; its quantity is about one sisteenth of the weight of the root.

3. GLADIOLUS LUTEUS Ph. Lond. Iris palustris Ph. Edinb. Iris palustris lutea Ger. Acorus vulgaris Pharm. Augustan. Acorus adulterinus C. B. Pseudoacorum Matth. Pseudoris Dod. Butomon Clus. Iris Pseud-Acorus Linn. Yellow water-flag, bastard acorus, sedge: with reddish roots, yellow unbearded flowers standing several on one stalk, and the middle ribs of the leaves prominent. It is common by the sides of rivers and marshes, and flowers in June.

The roots of this species are, when fresh, rather more acrid, and more strongly cathartic, than either of the preceding. The expressed juice, given to the quantity of eighty drops every hour or two, and occasionally increased, has, in some instances, produced plentisul evacuations, after jalap, gamboge, and mercurials had failed (a): but however successful it may have sometimes been as a drastic purgative, it is accompanied, like the other irises, with a capital inconvenience; its strength being so precarious, or so variable in different states, that it is by no means sit for general use. The juice, both of this and of the other kinds of iris, has been employed also externally for clearing the skin of serpiginous eruptions; and sometimes snuffed up the nose as a

strong errhine: even for these purposes it is to be used with caution, being subject, by its great acrimony, to instame or vesicate the parts.

The dry roots are much weaker and less agreeable than those of either of the preceding species of iris. They have scarcely any smell; and when chewed in substance, discover very little taste. An extract made from them by rectified spirit is likewise weaker and more nauseous, though its quantity is less, amounting only to one twenty-fourth of the weight of the root: it has nothing of the flavour or aromatic warmth of those of the other two, but an ungrateful austere bitterishness and a kind of saline pungency. It is the root in this dry state that the writers on medicines mean, when they speak of the yellow water-flag root as being aftringent and stomachic: it does not, however, appear to have any great claim to these virtues, and among us is no otherwise made use of than as an ingredient in the officinal arum powder, in which it is said to be employed only in deference to the original of Birckmann first published by Quercetanus.

JUJUBÆ.

JUJUBÆ Ph. Parif. Jujubæ majores oblongæ C. B. Zizyphus Dod. Rhamnus Zizyphus Linn. Jujubæs: a half-dried fruit of the plum kind, about the fize and shape of an olive: confisting of a pretty thick reddish yellow skin, a whitish fungous pulp, and a wrinkled stone pointed at both ends: the produce of a prickly tree, with three-ribbed leaves, and herbaceous or yellowish flowers, sometimes found wild, and commonly cultivated in the southern parts of Europe.

This fruit, when in perfection, has an agreeable sweet taste; and in those countries where it is common, makes an article of food in its recent state, and of medicine when half dried; decoctions of it being used, like other glutinous sweets, as incrassants, and demulcents in defluxions on the breast. Among us, it has long stood neglected, and is now become a stranger to the shops; the tree not producing fruit in this climate; and that, which we received from abroad, being commonly mouldy or carious.

ANOTHER fruit of the same kind, of a dark blackish hue, furnished with an ash-coloured cup at the bottom, from which it easily parts,

is sometimes brought from the eastern countries, under the names of sebesten, myxa, or myxaria. It is produced by the Cordia Myxa of Linnæus. It is more glutinous than the jujube; to which it has been commonly joined in pectoral decoctions; and along with which it is now discarded by the colleges both of London and Edinburgh.

JUNCUS ODORATUS.

JUNCUS ODORATUS Pharm. Lond. Juncus odoratus five aromaticus C. B. Schænanthus, squinanthum, sænum camelorum, & palea de mecha Quibusdam. Andropogon Schænanthus Linn. Sweet rush of Camel's hay: a dried herb, of the grass kind, brought from Turkey and Arabia, in bundles about a foot long; consisting of smooth stalks, in shape and colour somewhat resembling barley straws, full of a sungous pith like those of rushes; and leaves like those of wheat surrounding the stalk with several coats: towards the tops of the stalks are sometimes found short woolly spikes of impersect red slowers, set in double rows.

THE fweet rush, when in perfection, has an agreeable smell, and a warm, bitterish, not unpleasant taste. Distilled with water, it yields a small quantity of a yellowish, fragrant, and very pungent essential oil: the remaining decoction, thus divested of the aromatic matter of the plant, proves unpleasantly roughish, bitterish, and somewhat acrid. A tincture made in rectified spirit, in colour greenish yellow, yields, on being inspissated, a tolerably grateful, bitterish, aromatic extract. This plant, formerly employed as a warm stomachic and deobstruent, appears from the above experiments to be of no inconsiderable activity; but in this country, more common aromatic vegetables have now superfeded its use. It is kept in the shops only as an ingredient in the mithridate and theriaca; and the college of Edinburgh, having at last expunged those compositions, have dropt the juncus odoratus.

JUNIPERUS.

JUNIPERUS: juniperus vulgaris fruticosa C. B. Juniperus communis Linn. Juniper: an evergreen tree or bush, clothed with slender narrow stiff sharp leaves, like prickles, which stand generally three together: gether: the flowers are a kind of small scaly catkins growing on one plant; the fruit, round berries, growing on a different one, containing, each, three oblong irregular seeds. It is common on heaths in different parts of Europe; and is found, at all seasons of the year, both with

unripe green or red berries, and with ripe blueish black ones.

The BERRIES, baccæ juniperi Pharm. Lond. & Edinb. are brought chiefly from Holland and Italy: they should be chosen fresh, not much shrivelled, and free from mouldiness, which they are very subject to contract in keeping. They have a moderately strong not disagreeable smell, and a warm pungent sweetish taste, which if they are long chewed or previously well bruised, is followed by a considerable bitterness. The sweetness appears to reside in the juice or soft pulpy part of the berry; the bitterness, in the seeds; and the aromatic slavour, in oily vesicles, spread throughout the substance both of the pulp and of the seeds, and distinguishable even by the eye. The fresh berries yield, on expression, a rich, sweet, honey-like aromatic juice: if previously powdered, so as to thoroughly break the seeds, which is not done without difficulty, the juice proves tart and bitter. The same differences are observable also in tinctures and infusions made from the dry berries, according as the berry is taken entire or thoroughly bruised.

They give out nearly all their virtue both to water and rectified spirit, and tinge the former of a brownish yellow, the latter of a bright orange colour. Distilled with water, they yield a yellowish essential oil, very subtile and pungent, in smell greatly resembling the berries, in quantity (if they have been sufficiently bruised) about one ounce from forty: the decoction, inspissated to the consistence of a rob or extract, has a pleasant, balsamic, sweet taste, with a greater or less degree of bitterishness. A part of the flavour of the berries arises also in distillation with rectified spirit: the inspissated tincture consists of two distinct substances; one oily and

fweet: the other tenacious, refinous, and aromatic.

These berries are useful carminatives, detergents, and diuretics. The distilled oil is a very stimulating diuretic, approaching in quality to that of turpentine, like which, it impregnates the urine with a violet smell: the spirituous extract gives the same kind of smell; as does likewise the berry in substance, in a lower degree; but the watery extract or rob, as being divested of the oil, has no such effect. This last may be used with advantage in cases where the more stimulating preparations would be im-

Ol. e baccis juniperi Ph. Lond. & Edin.

proper;

proper; as in catarrhs, debilities of the stomach and intestines, and difficulties of the urinary excretions, in persons of an advanced age. Among the arcmatics that have been tried in composition with juniper berries, fweet fennel feeds and caraway feeds feem the best adapted to improve their flavour: a cordial water is prepared in the shops by drawing off a gallon of proof spirit from a pound of the berries and an ounce and a Aq. juniperi half of each of the feeds. The water is strongly impregnated with the volatile virtue of the berry; to which the more fixt ones may in many cases be usefully superadded, by mixing with it a proper quantity of the rob.

The WOOD, lignum juniperinum Ph. Edinb. (a) cedrinum lignum Ph. Parif. has been recommended as a fudorific, and by fome accounted fimilar to guaiacum or fassafras, to either of which it is greatly inferiour. It has a weak not unpleasant smell, and very little taste: decoctions and extracts, made from it with water, are difagreeably bitterish, subaftringent, and balfamic: the spirituous tinctures are weaker than the watery, and yield, on being inspissated, an almost insipid refin. The quantity of watery extract, according to Cartheufer's experiments, is about one twelfth the weight of the wood; of spirituous extract, one eighth.

In the warmer climates, particularly on the coasts of Africa, there exudes, from a larger species of juniper, a refinous juice, which concretes into femipellucid pale yellowish tears or glebes, resembling mastich, but larger; the fandaracha of the Arabians, and gummi juniperinum of the shops (Pb. Edinb.(a) called by some, from the use to which it has been principally applied, vernix. This refin has a light agreeable fmell, and no confiderable tafte: it diffolves in rectified spirit, and in oils both expressed and distilled, but gives out little or nothing to watery liquors, and thus discovers that it is nearly a pure refin. It is supposed to be fimilar in quality, as in appearance, to mastich; and has been fometimes given internally, against hemorrhagies, old fluxes, and ulcerations; but principally employed externally in corroborant, nervine, traumatic applications. Among us, it is scarcely ever made use of for any medicinal purposes; other refinous substances, more common in the shops, being apparently superiour to it.

KALI.

KALI Pharm. Edinb. (a) Kali majus cochleato semine C. B. Salsola quibusdam. Salsola Soda Linn. Snail seeded glasswort or saltwort: a plant with spreading, reddish, pretty thick branches; oblong, narrow, pointed, sleshy leaves like those of the houseleeks; and imperfect flowers in the bosoms of the leaves, followed each by one seed spirally curled and inclosed in the cup. It is annual, and grows wild on the sea coasts in the southern parts of Europe, particularly of the Mediterranean.

This herb is very juicy, in taste bitterish and remarkably saline. The expressed juice, and insuspense or decoctions of the leaves, are said to be powerfully aperient and diuretic, and in this intention have by some been greatly recommended in hydropic cases: half a dram of the juice is reckoned a sufficient dose. But the kali is principally regarded, on account of its yielding copiously, when burnt, the fixt alkaline salt called soda or soude: an impure soda is prepared from it about Montpelier, where the plant is said to be cultivated for this use in the salt marshes; and a purer kind at Alicant in Spain from a somewhat different species of kali (b). The salt called kelp, prepared among ourselves from different marine plants, contains an alkali of the same kind, but more impure.

The foda is much milder in taste than the common vegetable alkalies, and is in several other respects also very considerably different from them, being of the same nature with the mineral alkali or basis of sea salt (see Natron). It promises to be an useful article of the materia medica, and has for some time past been received in practice in this country, as it has long been among the French, both by itself, and combined with tartar into the neutral salt called sal rupellense. * The Edinburgh college have received a purified salt of this kind, under the title of sal alcalinus sixus sossilis purificatus.

(a) Expunged.

⁽b) Kali hispanicum supinum annuum sedi foliis brevibus, Mem. de l'acad. des scienc. de Paris, pour l'ann. 1717. & Pharm. Paris. p. lxiv. Salsola sativa Linn.

KERMES.

KERMES Pharm. Lond. Granum tinctorium & coccus bapbica quibusdam. KERMES: round reddish-brown grains, about the fize of peas: found in Spain, Italy, and the fouthern parts of France, adhering to the branches of the scarlet oak. These grains appear, when fresh, full of minute reddish ova or animalcules, of which they are the nidus, and which in long keeping change to a brownish red powdery substance. They are cured by fprinkling with vinegar before exficcation: this prevents the exclusion of the ova, and kills such of the animals as are already hatched; which would otherwise become winged insects, and leave the grain an empty husk.

FRESH kermes yield upon expression a red juice, of a light pleasant fmell, and a bitterish, subastringent, somewhat pungent taste: this juice, or a fyrup made from it, are brought from the fouth of France, and fometimes made use of as mild restringents and corroborants. An elegant cordial confection, for these intentions, is prepared in the shops, Confectio by dissolving, in the heat of a water bath, fix ounces of fine sugar in fix Pharm. Lond. ounces by measure of damask rose water, then adding three pounds of the juice of kermes warmed and strained, and after the whole has grown cold, mixing in half a fcruple of oil of cinnamon: this confection is taken from a scruple to a dram or more; either by itself, or in juleps, with which it mingles uniformly without injuring their transparency. The dried grains, if they have not been too long kept, give out, both to water, and to rectified spirit, the same deep red colour, and nearly the same kind of smell and taste, with those of the expressed juice. The watery tinctures lose nearly all their smell in evaporation: the spirituous retain nearly the whole of their fmell as well as of their tafte. The inspissated extracts are considerably bitter, astringent, and of a kind of mild balfamic pungency: the spirituous is stronger and in somewhat fmaller quantity than the watery, but the difference in strength is more confiderable than that of the quantity, spirit seeming to extract the active matter more completely than water.

KINO.

GUMMI rubrum astringens gambiense D. Fotbergill in med. obs. Lond. 1757. Kino, Pharm. Edinb. Red astringent gum from Gambia; supposed to exude from incisions made in the trunks of certain trees, called pau de sangue, growing in the inland parts of Africa.

It is very friable, so as to be crumbled in pieces by the hands; of an opake dark reddish or almost black colour in the mass, and when reduced to powder, of a deep brick red: small particles of it, viewed with a magnifying glass, appear of a semitransparent red like bits of garnet. In chewing, it first crumbles, then sticks together a little, and in a short time seems wholly to dissolve, impressing a very considerable astringency accompanied with a slight sweetishness. It has no smell.

To oils it gives little or no tincture. On a red-hot iron, it glows for a long time like a bit of burning charcoal, without shewing any disposition to melt: it yields, during a little while, a slight dull slame hovering about the surface, and leaves at length a large proportion of greyish ashes.

Both rectified spirit and water dissolve, each, about two thirds of it, the spirit somewhat more than the water. Both solutions, when made with the same quantities of the two menstrua, as twenty or thirty times the weight of the powdered gum, appear of the same deep bright red colour, the spirituous rather deepest: with solution of chalybeate vitriol, they both produce inky mixtures, from which the black matter speedily concretes and settles to the bottom, leaving the liquors colourless (a). The watery solution suffers no apparent change from the addition of alkalies fixt or volatile; but acids render it turbid, and occasion a copious precipitation.

The part, which water leaves undissolved, seems as dark-coloured as the gum at first: it gives the same deep red tincture to spirit, and this tincture strikes the same black with solution of vitriol. The part which

⁽a) The black matter in these kinds of mixtures appears to consist of the iron of the vitriol, disengaged from its acid solvent, and combined with the vegetable aftringent substance; the acid serving only as a necessary intermedium for procuring this union. The above black precipitates, after repeated ablutions with water, retained their blackness; and the clear liquors from which they had settled, being examined with alkaline salt on the principles to be mentioned hereafter under the article salkalini, seemed to contain as much acid as the quantity of vitriol employed in them.

spirit leaves undiffolved is much paler than the original gum, gives no tincture to water, and produces no change with the vitriolic folution.

It appears therefore that both the colouring and aftringent matter are more completely taken up by spirit than by water; though water extracts readily enough a great share of both.

ONLY a little quantity of this drug has hitherto been brought over. Dr. Fothergill, the first person, as far as I can find, who gave notice of it to the public, and who favoured me with the specimens on which the above experiments were made, informs us, that he had the first intimation of it from a physician, who had met with good effects from it in obstinate chronical diarrhoæ; and that a parcel was afterwards shewn to him, which had been received from a Guinea ship, and taken for a fine kind of dragons blood, which it pretty much refembles in appearance, though in quality effentially different. He observes, that from the trials which have been made, and from its fensible qualities, it promifes to be an article worth enquiring after, and to become in time a valuable addition to the materia medica. In diforders from laxity and acrimony, it may, doubtless, be of great advantage; nor do I recollect any other drug, that is so much of a gummy nature, and at the fame time fo aftringent. Terra japonica comes the nearest to it, but is manifestly less astringent. The terra japonica differs likewise, in its watery folutions fuffering no confiderable separation of their parts from the addition of acids; and in the black matter, which they produce with vitriol, being little disposed to concrete and precipitate. Whether the cause, on which these kinds of diversities depend, be sufficient to influence also their medicinal powers, our knowledge, both in the chemical composition of bodies, and in the operation of medicines, is as yet too imperfect to permit us to judge. * The Edinburgh college have now received this gum as an officinal, and have directed a tincture, Tinctura e Kino Pharm. in which two ounces of it are dissolved in a pound and a half of proof Edinb. spirit.

LABDANUM.

LABDANUM Pharm. Lond. Ladanum. LABDANUM: a refinous juice, exuding upon the leaves of a small shrub, ciftus ladanifera cretica flore purpureo Tourn. Ciftus creticus Linn. which grows plenti-3 D 2 fully fully in Candy and some of the other islands of the Archipelago, and bears the winters of our own climate. The juice is said to be collected, by lightly brushing the shrub, in the summer heats, with a kind of rake having several straps or thongs of leather fixed to it instead of teeth (a): the unctuous juice adheres to the thongs, and is afterwards scraped off with knives. The shrub is said to be very plentiful also in Spain (b), but it does not appear that any labdanum is brought from thence.

Two forts of labdanum are met with in the shops. The best, which is very rare, is in dark-coloured black masses, of the consistence of a soft plaster, growing still softer on being handled: the other is in long rolls coiled up, much harder than the preceding, and not so dark. The first has commonly a small, and the last a very large admixture of sine sand, which, in the labdanum examined by the French academy, amounted to three fourths of the mass. It is scarcely indeed to be collected pure, independently of designed abuses; the dust, blown on the plant by winds from the loose sands among which it grows, being retained by the tenacious juice.

LABDANUM has been fometimes exhibited as a refinous corroborant and restringent, but principally employed in external applications and persumes: the soft kind makes an useful ingredient in the cephalic and stomachic plasters of the shops. This sort has an agreeable smell, and a lightly pungent bitterish taste: the hard is much weaker, and the common means of purifying these kinds of substances, though they separate the sandy matter mixed with it, render it weaker still. Rectified spirit of wine dissolves nearly the whole of the pure labdanum into a gold-coloured liquor: on inspissating the siltered solution, the siner part of the labdanum rises with the spirit, and the remaining resin proves both weaker and less agreeable than the juice at first. On insufing the labdanum in water, it impregnates the liquor considerably with its smell and taste: in distillation with water, there comes over a fragrant essential oil; and there remains in the still a brittle almost insipid resin, with a pale coloured liquor, which, inspissated, yields a weakly bitterish extract.

⁽a) Belon, (Bellonius) Observations des choses memorables trouvées en Grece, &c. 1. i. c. vii.

⁽b) Clusius, Rariorum stirpium per Hispanias observatarum bistoria, l. i. c. v.

The specific flavour of this juice seems to be sooner diffipated by heat than that of almost any of the other officinal resins or gummy refins.

LAC.

LAC; lac asininum, caprinum, muliebre, ovillum, vaccinum. MILK: asses, goats, human, sheeps, and cows milk: a fluid prepared and secreted in the bodies of animals, but not completely elaborated into an animal nature. On a chemical analysis, it yields the same general principles with substances of the vegetable kingdom.

MILK is a mild nutritious balfamic fluid; when taken freely, an excellent obtunder of acrid and deleterious fubstances, and of over-doses of the stronger cathartics and emetics; one of the best restoratives in emaciated habits; a palliative, whilst its use is continued for the only aliment, in gouty cases not inveterate, and in some rheumatic pains; the medicine principally depended on in hectics and consumptions; prejudicial in acute diseases, bilious sluxes and dysenteries, swellings of the præcordia, and obstructions of the abdominal viscera.

It sometimes happens, that when the body stands most in need of this medicinal nutriment, the intestines are too slippery to retain it. In such cases it may be advantageously boiled with gentle astringents, as granate peel, balaustines, red roses; about an equal quantity of water being added, by a little at a time as the milk boils up, so as that all the water

may be wasted in the boiling (a).

It may be prefumed that milk thickens in a found stomach, before its digestion, nearly in the same manner as it is thickened by the runnet or infusion of the stomach of a calf; and that, where the gastric juices are too inert to produce this change, or so acid as to produce it in too great a degree and to separate a firm curd from the serous part; the milk will be difficult of digestion. Debilities of the stomach are endeavoured to be corrected by the medication above mentioned, or by the interposition of proper stomachies; acidities, by the absorbent earths. The absorbent earths, however, are in this intention commonly insufficient, unless assisted by stomachies; for as they absorb only the acid already generated,

and have no power of remedying the weakness or indisposition which tends to produce more, they afford only a temporary and palliative relief: and indeed it may be questioned, whether they are capable of so far destroying the force, even of the acid they are mixed with, as to prevent its curdling milk in the stomach.

Milk is curdled by all acids; by most, perhaps by all, of the combinations of acids with earthy and metallic bodies; by alkaline salts both fixt and volatile; by some vegetables that have no acidity or alkaline quality, as mustard seed; and by strong vinous spirits. The concentrated acids produce a strong curd immediately on mixture: most of the other substances scarcely have their sull effect without a boiling heat. The coagulum made by acids salls to the bottom of the serum: that made with alkalies swims on the surface, forming, especially if the alkali is of the volatile kind, a thick coriaceous skin. The serum, with alkalies, proves of a greenish hue: that made with the other substances is nearly of the same appearance with the whey which separates spontaneously.

The perfect neutral falts, or those compounded of an acid and an alkali, produce no coagulation, either with or without heat: some of them, particularly nitre and sal ammoniac, make the milk less coagulable, and, if added to the boiling mixture when already curdled by vegetable acids, render nearly the whole sluid again (a). Sugar retards the spontaneous coagulation, and impedes likewise the separation of the creme from milk, and of the butyraceous part from creme. Lime-water and animal gall redisfolve the coagula.

MILK, distilled with a gentle warmth, gives over a colourless and tasteless liquor, which seems to be mere water, but is found to differ from the simple element in growing sour upon keeping. The residuum is a grumous, unctuous, yellowish or brownish mass; which, on being boiled in water, partially dissolves. This solution contains the sweet substance of the milk, freed from the grosser unctuous caseous matter; and proves an elegant whey, more agreeable in taste, and which keeps better, than those prepared in the common manner. These sorts of liquors are very useful, cooling, diluent, aperients and detergents; in hypochondriacal complaints, impurities of the humours, acute diseases,

&c. They promote the natural excretions in general, and remarkably increase the action of the purgative sweets, casia and manna. The faline matter of these liquors may be obtained in a pure solid crystalline state, Saccharum by clarifying the whey with whites of eggs, and, after due evaporation, lactis P. Par. fetting it to shoot, in the same manner as other saline solutions.

Thus milk is refolved into a watery fluid; a gross substance indissoluble in water, which appears to contain the directly nutrimental part; and a fweet aperient falt. The milks of different animals differ remarkably in the proportions of these ingredients, and in the quality of the salt.

Breaft milk and affes milk are very nearly alike: twelve ounces leave on evaporation, according to Hoffman's experiments, eight drams of folid matter, of which boiling water disfolves fix drams: the folution, inspissated or crystallized, yields a salt of a rich honey-like or saccharine fweetness. The same quantity of cows milk leaves thirteen drams of folid matter, from which water extracts only about a dram and a half: the falt obtained from this folution is much lefs fweet, when purified is almost infipid, diffolves very difficultly, and feems to have little claim to the pectoral and antiphthifical virtues vulgarly afcribed to it. All the other milks that have been examined are of an intermediate nature between the two first and the last: goats milk approaches more to that of the ass than sheeps milk does, though both of them come nearer to that of the cow than of the afs.

There are confiderable differences in the milk of one and the same animal according to its different aliment. Dioscorides relates, that the milk of goats, which fed on the scammony plant and spurges, proved cathartic; and instances are given, in the Acta Hafniensia, of bitter milk from the animal having eaten wormwood. It is a common observation, that cathartics, spirituous liquors, &c. taken by a nurse, affect the child; that the milk of animals, feeding on green herbs, is more dilute than when they are fed on dry ones; and that many of the common plants, which are eaten by cattle, give a particular tafte to their milk. Hoffman is of opinion, that, on this principle, milk may be usefully impregnated with the virtues of different medicinal substances.

LACCA.

LACCA Pharm. Edinb. (a) LAC, STICK-LAC, improperly called Gum-LAC: a concrete brittle substance, of a dark red colour; brought from the East Indies incrustated on pieces of sticks; internally divided into several cells; said to be the resinous juice of certain trees, collected by winged red insects of the ant kind, impregnated with the tinging matter of the insects, and by them deposited either on the branches of the trees, or on sticks sastened in the earth for that purpose. In the cells are often observed small red bodies, which appear to be the young insects (b).

* A curious account by Mr. Kerr of the insect producing this gum, is contained in the Philos. Trans. Vol. LXXI. Part II. From this it appears, that these insects are inhabitants of four trees; the Ficus religiosa Linn. the Ficus Indica Linn. the Plaso Hort. Malabar. and the Rhamnus Jujuba Linn. The lac is however rarely found upon this last, and of an inferiour quality. The two species of Ficus yield a milky juice when wounded, which instantly coagulates into a viscid substance. The Plaso tree by incision gives out a red gum very similar to the lac. Hence the insect seems to have little trouble in animalizing the juices of these trees so as to make its cell, which is the stick-lac. It is found in very great quantities on the uncultivated mountains on both sides the Ganges; and is of great use to the natives in various works of art, as varnish, painting, dying, &c.

The tinging red animal matter of the stick-lac dissolves both in water and in rectified spirit, and appears to be of the same general nature with that of cochineal; like which it is made dull by alkalies, and brighter by acids, and turned to a scarlet by solution of tin. If the lac be broken in small pieces, or grains, and insused in warm water, till it ceases to give any tincture to the liquor; the remainder appears of a transparent yellowish or brownish colour, and, on raising the heat so as to make the water boil, melts and rises to the surface. The grains, or the plates formed from them by liquesaction, thus robbed of great part of the

Seed lac of the shops. Shell lac of

the Shops.

(a) Expunged.

(b) See the Memoires de l'acad. roy. des sciences de Paris, pour l'ann. 1714.

animal

animal tincture, feem to be of an intermediate nature between that of wax and refins, or to partake of the nature of both: they crumble on chewing, and do not foften or stick together again: laid on a red-hot iron, they instantly catch fire, and quickly burn off, with a strong and not disagreeable smell: distilled, they yield, like wax, an acid spirit and a butyraceous oil: alkaline lixivia, and volatile alkaline spirits, dissolve them into a purplish liquor: they dissolve also, by the assistance of heat, in rectified spirit of wine, and communicate to it a yellowish or brownish red colour, an agreeable smell, and a bitterish, subastringent, not unpleasant taste. The lac in substance, whether entire, or freed from so much of its colouring matter as boiling water is capable of extracting, has no manifest taste or smell.

A spirituous tincture of stick-lac has been sometimes given as a mild restringent and corroborant in semale weaknesses, and in rheumatic and scorbutic disorders. But the principal medicinal use of this concrete is as a topical corroborant and antiseptic, in laxities and scorbutic bleedings and exulcerations of the gums: some employ for this purpose a tincture of the lac in alum water; others, a tincture made in vinous spirits impregnated with the pungent antiscorbutics.

LACTUCA.

LETTUCE: a plant with slender but firm stalks, which yield, as do the leaves, a milky juice on being wounded: the flower consists of a number of flat flosculi set in a small scaly cup, followed by short flat seeds, which are pointed at both ends and winged with down.

I. LACTUCA Pharm. Edinb. (a) Lactuca sativa C. B. & Linn. Garden lettuce: with oblong, broad, rounded, uncut leaves; and numerous flowers standing on long pedicles in the form of an umbel. It is annual, and raised at different times of the year in culinary gardens.

THE young leaves of the several species or varieties of garden lettuce are emollient, cooling, in some small degree laxative and aperient, easy of digestion, but of little nourishment; salubrious in hot, bilious, in-

dispositions; less proper in cold phlegmatic temperaments. In some cases, they tend to procure sleep; not as being possessed of any strictly hypnotic power; but by virtue of their refrigerating and demulcent quality. When the plant is grown up, it proves considerably bitter, though less so than most of the others of the lactescent kind, to which it is similar in its general virtues.

The feeds, which in the common lettuce are of a grey or ash colour, in the cabbage lettuce black, unite with water, by trituration, into an emulsion or milky liquor, which has nothing of the aperient bitterness of the milky juice of the leaves, and is nearly similar to the emulsions made with almonds. The lettuce emulsions have been supposed to be more refrigerant than those of the almond, and hence have sometimes been preferred in heat of urine and other disorders from acrimony or irritation.

2. LACTUCA SILVESTRIS Medicorum. Lactuca filvestris & Scariola Pharm. Paris. Lactuca silvestris costa spinosa C. B. Lactuca Scariola Linn. Wild lettuce: with the leaves cut almost to the rib into indented triangular segments; and the stalks and the ribs prickly. It is biennial, grows wild in hedges, and slowers in June.

This species is considerably bitterer than the garden lettuces, and more aperient and laxative. It is nearly similar, in virtue as in taste, to endive unblanched.

3. LACTUCA GRAVEOLENS: lactuca filvestris odore viroso C. B. Lactuca virosa Linn. Strong scented lettuce, by some erroneously supposed to be the wild lettuce of medical writers: with the lower leaves entire, the upper jagged, the stalks and leaves prickly. It is biennial, found in hedges and by the sides of ditches, and slowers in June.

This species differs greatly in quality from the two preceding, though reckoned by botanists to be only a variety of the second. It smells strongly of opium, and appears to partake, in no small degree, of the virtues (a) of that narcotic drug. The opiate power of the lettuce, like that of the poppy-heads, resides in its milky juice, but whether the

milk of the lettuce is of equal fafety, or its virtue precifely of the fame kind, with that of the poppy, is not known.

*Dr. Collin of Vienna has written a tract recommending the use of this plant in the cure of dropsies. The preparation he employs, is an extract from the expressed juice, first sufficiently clarified, and evaporated by a very gentle heat. He begins with small doses; but in dropsies of long standing, originating from visceral obstructions, he rises to the quantity of from one to three drams in twenty-four hours. He has constantly found it a mild remedy, agreeing perfectly with the stomach. It usually kept the body open, but without exciting a purging. It feldom failed of proving powerfully diuretic, and at the same time mildly diaphoretic. The patients' thirst is said to have been totally extinguished by its use; but at the same time we are told that they were allowed to drink freely of diluting liquors during the course. Dr. Collin afferts, that out of twenty-four dropsical cases, all but one were cured by the use of this medicine; a degree of success that certainly entitles it to the further notice of the faculty.

LAMIUM.

LAMIUM ALBUM Pharm. Lond. & Linn. Lamium album non fætens folio oblongo C. B. Galeopsis & archangelica, & urtica mortua sive alba Quibusdam. White archangel or Dead nettle: a plant with square stalks; oblong indented acuminated leaves, like those of the stinging nettle, set in pairs at the joints; and clusters, in the bosoms of the leaves, of white labiated flowers, whose upper lip is entire, arched, and hairy, the lower lip cloven. It is perennial, common in hedges and about the borders of fields, and found in flower from April to near the end of summer.

Infusions of this plant, drank as tea, are faid to be beneficial in uterine hemorrhagies and the fluor albus: the flowers are supposed to be more efficacious than the leaves, and hence those only are directed by the college of London. The sensible qualities, either of the one or the other, afford little foundation to expect from them any considerable virtues. The flowers have only a slight mucilaginous sweetishness, without any remarkable smell or flavour: the leaves have a weak not 3 E 2 unpleasant

unpleasant smell, and a small degree of roughness, which may entitle them to a place among the milder corroborants.

LAMPSANA.

LAMPSANA, Lapfana, Napium, Papillaris herba. Soncho affinis lampfana domestica C. B. Lapfana communis Linn. Dock cresses, Nipplewort: a roughish plant; bearing small yellow sloseulous flowers, set in form of an umbel on the top of the stalk, followed by little crooked naked seeds: the lower leaves are deeply cut, towards the pedicle, into generally two or four opposite sections; the upper are oblong, narrow, undivided, and have no pedicles. It is annual, grows wild by road sides, and flowers greatest part of the summer.

This is one of the bitter lactescent plants, nearly similar in virtue to dandelion, endive, cichory, and the others of that class. It has been employed chiefly for external purposes, against wounds and ulcerations, particularly of the nipples, whence its names nipplewort and papillaris.

LAPATHUM.

DOCK: a perennial plant bearing numerous imperfect flowers fet in double cups: the outermost cup consists of three small green leaves; the inner of three larger reddish ones, which become a covering to a glossy triangular feed.

I. OXYLAPATHUM Pharm. Edinb. (a) Lapathum acutum folio plano C. B. Rumex acutus Linn. Sharp-pointed wild dock: with long acuminated leaves, not curled about the edges, growing gradually narrower from the bottom to the point; and the feed-covers indented and marked with little tubercles. The roots are of a brownish colour on the outside, and of a yellowish within, which grows deeper in drying.

THE roots of the sharp-pointed dock have a bitterish astringent taste; and no remarkable smell: the roots of the other common wild docks

are nearly of the same quality, equally discover their astringent matter both to the tafte and by striking an inky blackness with solution of chalybeate vitriol, and have been often substituted in our markets to those of the sharp-pointed kind; which last are generally, and, so far as can be judged from their taste, justly, accounted the most efficacious. They are supposed to have an aperient and laxative, as well as an astringent and corroborating virtue; approaching in this respect to rhubarb, but differing widely in degree, their stypticity being greater, and their purgative quality, if really they have any purgative quality at all, far less. They stand recommended in habitual costiveness, obstructions of the viscera, scorbutic and cutaneous maladies: in which last intention, fomentations, cataplasms, or unguents of the roots, have been commonly joined to their internal use: in many cases, the external application alone is faid to be fufficient. Their active matter is taken up both by water and rectified spirit, and, on inspissating the tinctures, remains in the extracts; both the watery and spirituous extracts are confiderably bitter and very auftere. A decoction of half an ounce or an ounce of the fresh roots, or of a dram or two of the dry roots, is commonly directed for a dofe.

2. HYDROLAPATHUM sive Herba britannica Pharm. Edinb. Lapathum aquaticum folio cubitali C. B. Rumex aquaticus Linn. Great wild water-dock: with very large leaves, two or three feet long; the feed-covers not indented. The roots are externally blackish, internally white with a faint reddish tinge, which, in drying, changes in some parts to a yellowish: the internal part of the fresh root, exposed to the air, or of the dry root moistened, soon changes superficially to a deep yellow or brown.

The roots of the water-dock strike a black colour with solution of chalybeate vitriol, like those of the preceding species, but have a much stronger and more acerb taste; which is diffused equally, so far as can be judged, through the whole substance of the root. They give out their active matter both to water and rectified spirit, and tinge both menstrua of a pale yellowish or reddish brown colour, though in chewing they render the saliva only milky.

The

The berba britannica of the ancients, celebrated as an antiscorbutic, and of which the knowledge was long loft, was proved by Muntingius, towards the end of last century, to be no other than this great waterdock. Muntingius endeavours to prove also, that its name britannica was not derived from that of our island, but from Teutonic words expressing its power of fastening loose teeth, or of curing the disease which makes them loofe. Later experience has shewn, that it is a medicine of very confiderable efficacy, both externally in lotions against putrid spongy gums and ulcerations, and as an internal antiscorbutic: Boerhaave affures us, that in these cases he has known many instances of its happy effects. It is supposed to be of service also in cutaneous defedations different from the true scurvy, in rheumatic pains, and in chronical diforders proceeding from obstructions of the viscera. It has been chiefly used in medicated wines and small ales, with the addition generally of some spicy materials, and sometimes of other antiscorbutic plants, as scurvygrass, buckbean, horse-radish, &c.

3. Rhabarbarum monachorum Pharm. Parif. Lapathum hortense latifolium C. B. Hippolapathum; Patientia. Rumex Patientia Linn. Monks rhubarb, garden patience: with large, broad, acuminated leaves; reddish, branched stalks; the leaves that cover the seeds unindented, and a tubercle on one of them: the root is of a yellow colour, with red veins, approaching in appearance to rhubarb.

This root is supposed to possess the virtues of rhubarb, in an inferiour degree. It is obviously more aftringent than rhubarb: but comes very far short of it in purgative virtue, though given, as usually directed, in double its dose; nauseating the stomach, without producing any considerable evacuation. It communicates a deep yellow tincture both to water and spirit.

LAVENDULA.

LAVENDER: a shrubby plant, with its leaves set in pairs, the stalks square when young, and round when grown woody; producing, on the tops of the branches, naked spikes of blue, sometimes white, labiated flowers, of which the upper lip is erect and cloven, the lower divided into three roundish segments.

I. LAVENDULA

1. LAVENDULA Pharm. Lond. & Edinb. Lavendula angustifolia C. B. Pseudonardus quæ lavendula vulgo J. B. Lavendula minor sive spica Ger. & Park. Lavandula Spica Linn. Lavender: with oblong, very narrow, somewhat hoary, undivided leaves; a native of dry gravelly soils in the fouthern parts of Europe, common in our gardens, and flowering in July.

THE flowers of lavender have a fragrant smell, to most people agreeable, and a bitterish, warm, somewhat pungent taste: the leaves are weaker and less grateful. They are often employed as a perfume; and medicinally, as mild stimulants and corroborants, in vertigoes, palfies, tremors, and other debilities of the nervous system, both internally and externally.

The flowers are fometimes taken in the form of conferve; into which Conferva flothey are reduced, by beating them, while fresh, with thrice their weight of double refined fugar. Their fragrance is less injured by beating or bruifing them than most of the other odoriferous flowers, but is nevertheless considerably diminished: the flavour of the leaves is of a much less destructible kind.

Water extracts by infusion nearly all the virtue both of the leaves and flowers. In distillation with water, the leaves yield a very small portion of effential oil; the flowers a much larger, amounting, in their most perfect state, when they are ready to fall off spontaneously and the seeds begin to shew themselves, to about one ounce from fixty. The oil is Ol.essentiale of a bright yellowish colour, a very pungent taste, and possesses, if carefully distilled, the fragrance of the lavender in perfection: it is given internally from one drop to five, and employed in external applications for stimulating paralytic limbs and for destroying cutaneous insects. The decoction, remaining after the distillation of the oil, is disagreeably bitterish and somewhat saline.

Rectified spirit extracts the virtue of lavender more completely than water. The spirit elevates also in distillation a considerable part of the odoriferous matter of the leaves, and greatest part of that of the flowers; leaving, in the inspiffated extracts, a moderate pungency and bitterishness with very little smell. A spirit prepared by pouring a gallon of proof Spiritus laspirit on a pound and a half of the fresh-gathered flowers, and drawing off vend. simpl. Pharm. Lond. five pints by the heat of a water bath; or by adding eight pounds of rectified spirit to two of the flowers, and drawing off seven pounds, is Pharm. Edin.

rum lavendulæ Ph. L.

Ph. L. & Ed.

Spiritus lavend. comp. Pharm.Lond.

Spirit, lavend comp. Pharm. Edin. richly impregnated with the fragrance of the flowers. More compounded fpirits of this kind, in which other aromatics are joined to the lavender, have been distinguished by the name of English or palfy drops: the college of London directs three pints of the simple spirit of lavender, and one pint of spirit of rosemary, to be digested on half an ounce of cinnamon, half an ounce of nutmegs, and three drams of red saunders as a colouring ingredient: the college of Edinburgh, to the same quantity of both spirits, orders one ounce of cinnamon, two drams of cloves, half an ounce of nutmegs, and three drams of red saunders. These preparations are taken internally, on sugar or in any convenient vehicle, from ten to an hundred drops, and used externally in embrocations, &c.

2. LAVENDULA LATIFOLIA C. P. Lavendula major seu spica Pharm. Paris. Pseudonardus quæ vulgo spica J. B. Lavendula major sive vulgaris Park. Broad lavender: with longer, broader, and hoarier leaves, less numerous on the stalks and branches; and much larger spikes, though smaller slowers; common in the southern parts of Europe, but rare among us. The name spike is applied by foreign writers to this species, by some of ours to the first. Linnæus makes this only a variety of the former.

THE broad-leaved lavender is stronger both in smell and taste than the narrow, and yields in distillation almost thrice as much essential oil, but the slavour both of the oil and of the plant itself, is much less grateful: the oil is likewise of a much darker colour, inclining to green. Watery and spirituous extracts, made from the two sorts of lavender, are very nearly alike; the difference seeming to reside only in the volatile parts.

LAUROCERASUS.

LAUROCERASUS Pharm. Parif. Cerasus folio laurino C. B. Prunus Lauro-cerasus Linn. LAUREL, CHERRY-BAY: an evergreen tree or shrub, with large, thick, oblong, glossy leaves, pointed at both ends, and slightly indented: towards the tops of the branches come forth pentapetalous slowers, in sive-leaved cups, followed by clusters of berries, like cherries or damsons. It is cultivated in gardens, slowers in May, and ripens its fruit in August or September.

THE

THE leaves of the laurocerafus have a bitter tafte, accompanied with a flavour refembling that of the kernels of certain fruits, as those of black cherries, apricots, bitter almonds, &c. Like those kernels, they communicate an agreeable flavour both to watery and spirituous liquors, by distillation and by infusion; and, like them also, they appear from fome late trials to be poisonous. A distilled water, strongly impregnated with their flavour, given in the quantity of four ounces to a large mastiff dog, occasioned in a few minutes terrible convulsions, and within an hour put an end to his life: dogs have been killed also, in a few minutes, by smaller quantities, of the distilled water, of an infusion of the leaves in water, and of their expressed juice, taken into the stomach, or injected by the anus; and there are some instances of liquors slavoured with the distilled water being poisonous to human subjects. The dissections of dogs killed by this poifon have shewn no other morbid appearances, or alterations, than fuch as may be reasonably supposed the immediate effect of the convulfions: when the distilled water, or the leaves in substance, were given in fuch fmall quantities as not to kill, and continued for fome time, the pulse became quicker, and the blood more fluid, and of a more florid red colour (a). It is faid that infusions of the leaves (made probably very weak) are commonly used in Holland in disorders of the lungs (b).

The kernel of the fruit is of the same nature with the leaves. The pulpy part discovers no ill quality to the palate, is coveted by birds, and appears to be innocent.

LAURUS.

LAURUS Pharm. Lond. & Edinb. Laurus vulgaris C. B. Laurus nobilis Linn. BAY: an evergreen tree or shrub, with oblong, stiff, smooth leaves, pointed at both ends, pale yellowish, monopetalous flowers, divided into four sections; and oblong dry berries, containing, under a thin black skin, a horny shell, within which are lodged two dark brownish seeds joined together. It is a native of the southern parts of Europe, and not uncommon in our gardens: it flowers in April and May, and

⁽a) See Dr. Langrish's experiments on brutes, and No. 418 and 420 of the philosophical transactions.

⁽b) Linnæi Amænitat. academic. iv. 40. i. 409.

ripens its berries in September. The shops have been commonly supplied with the berries from the Streights.

THE leaves of the bay have a light agreeable finell, and a weak aromatic roughish taste: in distillation with water they yield a small quantity of a very fragrant effential oil: with rectified spirit they afford a moderately warm pungent extract. The berries are stronger both in smell and taste than the leaves, and yield a larger quantity of effential oil: they difcover likewise a degree of unctuosity in the mouth, give out to the press an almost insipid fluid oil, and on being boiled in water a thicker butyraceous one, of a yellowish green colour, impregnated with the flavour Pharm. Edin. of the berry.

Oleum expressum baccarum lauri

> The leaves and berries of the bay are accounted stomachic, carminative, and uterine: in these intentions, infusions of the leaves are sometimes drank as tea; and the effential oil of the berries given, on fugar or diffolved by means of mucilages or in spirit of wine, from one to five or fix drops. The principal use of these simples in the present practice is external: they are made ingredients in carminative glyfters, warm cataplasms, and uterine baths; and the butyraceous oil of the berries serve as a basis for some nervine liniments, and mercurial and sulphureous unguents.

LAZULI LAPIS.

LAZULI LAPIS Pharm. Edinb. (a) Lapis cæruleus; Lapis cyanus; Cæruleum nativum. LAPIS LAZULI: a compact ponderous fossil; less hard than flint; of a deep blue colour, variegated commonly with gold or filver coloured points or veins; retaining its colour in a moderately strong fire, in a very strong one calcining to a brown, and at length melting into a dufky coloured glass with blueish clouds; losing its colour, and in great part diffolving, by digeftion in mineral acids; faid to be found in the mines of gold, filver, and copper, in the eastern countries and in some parts of Germany (b).

This stone, levigated into an impalpable powder, and freed from the groffer parts by washing with water, has been given in doses of half

(a) Expunged.

(b) See Cronstedt's mineral system, and Marggraf's chemical works.

a dram

a dram and a dram, and faid to operate strongly by stool and vomit. Some have recommended it in epilepfies and intermitting fevers: Dolæus tells us, that in this last disorder, the above doses, taken on the approach of a fit, with two or three spoonfuls of brandy, were with him a singular fecret. The ancients supposed, that it evacuates chiefly what they called melancholic humours or adust black bile, probably, as Geoffroy fuspects, on account of its tinging the feces black; a property, from which it may rather be prefumed that the mineral participates of iron. The constringing power, which is likewise ascribed to it, depends perhaps on this ingredient; but neither its real medical qualities, nor its chemical composition, are as yet known. Its ferrugineous impregnation is apparent, from its yielding yellow martial flowers on fublimation with fal ammoniac, and from folutions of it in mineral acids affording a blue precipitate with the tincture of Prussian blue described under the article ferrum. It has been generally supposed to participate pretty largely of copper; but pure lapis lazuli gives no mark of copper; and those, who speak of experiments discovering that metal in it (a), have probably taken for lapis lazuli some other blue stones, as the lapis armenus, which plainly contains copper, and which some celebrated naturalists have ranked as a species of the lazuli. The lapis armenus may be readily distinguished, by its being less hard than the lazuli, soon losing its blue colour in moderate fire, and raising an effervescence with acids, its basis seeming to be a calcareous earth.

LENTISCUS.

LENTISCUS Ph. Edinb. (b) Lentiscus vulgaris C.B. Lentiscus verus ex insula chio cortice & foliis suscis Commel. Pistachia Lentiscus Linn. Lentisk or Mastich tree: an evergreen tree or shrub, with soft slexible branches hanging downwards, and small stiff leaves, pointed at both ends, set in pairs on a surrowed rib, which terminates in a soft prickle: some trees produce reddish impersect flowers, in the bosoms of the leaves; others, clusters of black sirm berries including a whitish kernel. It is a native of the southern parts of Europe, and bears the ordinary winters of our own climate: large plantations of it are culti-

(a) Hoffman, in notis ad Poterium, p. 628,

(b) Expunged.

vated in the island of Chio, on account of the resinous juice, called mastich, obtained from incisions made in the trunk. The wood is sometimes brought to us from Marseilles, in thick knotty pieces, covered with a brownish bark, internally of a whitish or pale yellowish colour.

This wood is accounted a mild balfamic restringent: infusions and decoctions of it are greatly commended, in the German ephemerides, against catarrhs, nauseæ, weakness of the stomach, and in general as a corroborant and an alterative or fweetener(a). It may indeed be prefumed, from its fenfible qualities, to possess virtues of this kind, though in no very high degree. Its fmell and tafte are aromatic and refinous, but very weak: the fmall tough sprigs are stronger than the larger pieces, and the bark than the wood. It impregnates water with a red colour, and a light agreeable finell: to rectified spirit it gives a bright yellow tincture, and fcarce any fmell. On gently diffilling off the menstrua from the filtered liquors, the remaining extracts prove refinous, fubastringent, and slightly pungent: the watery extract discovers more of the flavour of the wood, and is in tafte rather stronger, though much larger in quantity, than the spirituous; the spirit covering or suppressing the fmell, and not taking up enough of the gummy or mucilaginous matter to render the refin diffoluble in the mouth. According to Cartheuser's experiments, the watery extract amounts to one eighth the weight of the wood, the spirituous to one twentieth or one fixteenth.

LEPIDIUM.

LEPIDIUM: a plant with undivided leaves, and small tetrapetalous white flowers on the tops of the stalks and branches, followed by little short heart-shaped sharp-pointed pods, which are divided longitudinally into two cells full of minute seeds.

1. LEPIDIUM Pharm. Edinb. (b) Lepidium latifolium C. B. & Linn. Piperitis. Raphanus sylvestris Belgis & Gallis. DITTANDER, PEPPERWORT, POOR-MANS-PEPPER: with oblong, broad, acuminated,

(a) Wenck, Alla nat. curiof. dec. iii. ann. ix & x. p. 254. (b) Expunged. ferrated

MATERIA MEDICA.

ferrated leaves. It is perennial, and found wild in some parts of England by the sides of rivers and in other moist shady places.

2. IBERIS, seu Cardamantica: Iberis latiore folio C. B. Lepidium gramineo folio sive iberis Tourn. Lepidium Iberis Linn. Sciatica cresses: with long narrow leaves, the lower on long pedicles and serrated; the upper entire, without pedicles. It is annual; a native of the southern parts of Europe; and raised in our gardens, as the preceding, for culinary use.

These herbs, when fresh, have a quick penetrating pungent taste; which is in great part dissipated or destroyed by exsiccation, retained in the expressed juice, extracted by water and by rectified spirit, and elevated by both menstrua in distillation or evaporation. They are recommended as antiseptics, stomachies, attenuants, and aperients; and appear to be of the same general nature with the cochleariæ, nasturtium, and other acrid antiscorbutics. The second fort has been supposed particularly serviceable, externally, against the sciatica; whence its common English name.

LEVISTICUM.

LEVISTICUM seu Ligusticum, Pharm. Edinb. Ligusticum vulgare C. B. Angelica montana perennis paludapii solio Tourn. Ligusticum Levisticum Linn. Lovage: a tall umbelliserous plant, with large leaves divided and subdivided into sections like those of smallage: the umbels stand on short pedicles, with several little leaves at the origin of each of the primary ones, and a few at the ramifications; the seeds are of a pale brown colour, oblong, plano-convex, marked with sive longitudinal ridges: the root thick, sleshy, juicy, branched, of a dark brownish colour on the outside, and a whitish or pale yellowish within. It is a native of the southern parts of Europe, and raised with us in gardens: it is perennial, slowers in June, and ripens its seeds in August.

ALL the parts of this plant are of the aromatic kind; of a strong slavour, somewhat like that of angelica, but less agreeable; supposed particularly useful in semale disorders. The leaves, which have been generally made choice of in this intention, have the most unpleasant smell, and suffer no great loss of it in keeping for some months; their taste

taste is moderately warm, and acrid, and very durable in the mouth and throat. The root, whose smell is nearly of the same kind with that of the leaves, though more approaching to gratefulness, discovers to the taste a considerable sweetness joined to its mild aromatic warmth; an extract made from it by water retains little more than the sweet matter; the slavour exhaling in the inspissation, and impregnating the distilled sluid, from which, if the quantity of the root subjected to the operation be large, a small portion of essential oil separates: an extract made by rectified spirit retains the aromatic part as well as the sweet, and proves moderately warm, but much less so than the extract of angelica: towards the end of the inspissation of the spirituous tincture, a thin unctuous matter appears upon the surface, in taste highly aromatic, and which seems to be the part that gives activity to the rest of the mass. The seeds of the plant have little of the sweetness of the roots, but are rather of more warmth and pungency, and of a more agreeable slavour.

LICHEN.

LIVERWORT: a kind of imperfect vegetable production, confifting of spreading leaves, of a leathery crustaceous matter. A fort of flowers both male and semale have been discovered in it, the latter producing innumerable seeds, like meal.

LICHEN CINEREUS TERRESTRIS Pharm. Lond. Lichen terrestris cinereus Raii. Lichen caninus Linn. Ash-coloured ground liverwort: consisting of roundish pretty thick leaves, divided about the edges into obtuse segments, shat above, of a reticular texture underneath, sastened to the earth by small sibres; when in perfection, of an ash grey colour; by age turning darker coloured or reddish. It grows on commons and open heaths, spreads quickly on the ground, and is to be met with at all times of the year, but is supposed to be in its greatest vigour about the end of autumn.

This herb is faid to be a warm diuretic. It is particularly celebrated as a preservative against the terrible consequences of the bite of a mad dog: an account of the remarkable efficacy, in this intention, of a powder composed of the dry leaves and black pepper, was communicated

to the royal fociety by Mr. Dampier, and published in No. 237 of the Philosophical Transactions. This powder was afterwards inserted, in the year 1721, into the London pharmacopæia, at the defire of Dr. Mead, who had large experience of its good effects, and who declares, that he had never known it to fail, where it had been used, with the affiftance of cold bathing, before the hydrophobia began. He directs a dram and a half of the powder to be taken in the morning fasting, in half a pint of cows milk warm, for four mornings fuccessively: previoully to these four doses, the patient is to be blooded nine or ten ounces; and after them, to be dipt in cold water every morning fasting for a month, and then dipt thrice a week for a fortnight longer (a). The powder was originally composed of equal parts of the lichen and pepper: Pulvis antibut this quantity of pepper rendering the medicine too hot, only one lyffus Pharm. part is now used to two of the lichen.

If cold bathing, bleeding, black pepper, and lichen, conjointly, be really of fufficient efficacy against the poison of the mad dog, it will not perhaps follow that any share of this efficacy belongs to the lichen: and indeed greater stress has been laid in general on the cold bath, than on this or the other parts of the prescription. The lichen does not promise to have any valuable medicinal power: to the organs of tafte or fmell it discovers no activity: taken by itself, in double the quantity above prescribed, it did not appear to have any sensible operation. Digested in rectified spirit, it tinged the menstruum of a deep yellowish green colour: on distilling off the spirit from the filtered tincture, the remaining grumous extract had very little tafte, and amounted only to twentyfix grains from two ounces, or about one thirty-feventh of the weight of the lichen. A decoction of the herb in water was brownish, and of a faint smell, somewhat like that of mushrooms: the extract, obtained by inspissating it, weighed one eighth of the lichen, and had some taste, but so little, that it is hard to say of what kind.

*LICHEN ISLANDICUS Linn. Lichen Pharm. Edinb. Lichen terrestris, foliis Eryngii, Buxb. Cent. II. Lichenoides rigidum, Eryngii folia referens, Raii & Dillen. Eryngo-leaved, or eatable Iceland, liverwort. This species of lichen consists of nearly erect leaves, stiff when dry, but foft and pliant when moift, irregularly divided into broad diftant fegments, smooth, and ciliated at the margins. It grows in the mountainous parts of this country, and in various other parts of Europe.

The Iceland lichen infused in water gives a bitterish liquor, which is reddened by a mixture of martial vitriol. A decoction of it is very thick and viscid; and on cooling concretes into a strong gelly. An ounce of the lichen boiled for a quarter of an hour in a pound of water, and afterwards strained, yielded seven ounces of a mucilage as thick as that procured by the solution of one part of gum-arabic in three of water.

The inhabitants of Iceland make great use of this lichen both as food and as physic. When fresh, according to Borrichius (a), it is employed as a purgative; but Olassen (b) decries that it has any more than a very gently opening quality. It is usually dried and ground into a meal, with which they make pottage and other preparations, adding either water or milk, and find it an agreeable and very nutritive article of food. It is best first to steep it for a sufficient time in water, in order to extract the bitterness.

The prepared lichen has been much used of late, particularly at Vienna, as a remedy for consumptive disorders. The celebrated Scopoli (c) has published some cases of its successful exhibition in the phthis; and other practitioners (d) have confirmed his account. It is used boiled in milk to a kind of pottage, of which the patient's diet is chiefly to consist. It is said to be antiseptic, easy of digestion, and remarkably nourishing. It is also recommended in other cases, where the stomach is so weak that common aliments are rejected. The Edinburgh college have received this lichen into their catalogue of simples.

LIGNUM ALOES.

LIGNUM ALOES, Xyloaloes, & Agallochum, Pharm. Parif. Lignum Calambac. AGALLOCHUM, CALAMBAC, or Aloes wood: a wood brought from China, and the inner parts of Tartary, in small pieces, compact and ponderous, of a yellowish or rusty brown colour, with black or purplish veins, sometimes purple with ash-coloured veins, and sometimes all over blackish. Of its origin, we have no very satis-

⁽a) Bartholini Ad. med. Hafn. 1671. (c) Annus 2. bistorico-natural, p. 114.

⁽b) Journey to Iceland.
(d) Bergius, mat. med. 858.

factory account: most of the writers, to whom we are indebted for information about the productions of those countries, report, that it is the internal part of certain trees; that a large tree affords only a very small quantity of this valuable part; and that there are several different forts of it, of which the best is never brought to us, being fold in China itself for twice or thrice its weight of filver.

THE best fort of agallochum wood brought into Europe, has a bitterish refinous taste, and a light aromatic smell. Set on fire, it seems to melt like wax, and emits, during the burning, an agreeable fragrance, which continues till the wood is wholly confumed. It is this fragrance in burning which makes the wood precious in the eastern countries for fumigations, and which affords the furest criterion of its genuineness and goodness. As this wood is apparently very refinous, rectified spirit takes up more from it than watery menstrua: according to Cartheuser's experiments, an ounce yields with spirit three drams of extract, and with water only two. The watery decoction and extract are moderately bitter and subacrid. The spirituous make less impression on the organs of taste, being less dissoluble in the mouth, or less miscible with the saliva: the Resina ligni pure refin, obtained by precipitation with water from the fomewhat in- aloes Pharm. spissated spirituous tincture, as directed by the faculty of Paris, is still weaker in taste. Hoffman observes, that in distillation with water, it yields an effential oil, of a whitish colour, of a thick confishence, of great fragrance, but in small quantity, not exceeding half an ounce from one hundred and fixty ounces of the wood: this oil, in which the more valuable parts of the agallochum are concentrated, he recommends, diffolved in spirit of wine, as one of the best cordials and corroborants, in weaknesses of the stomach and depressions of strength (a).

In our shops, we rarely meet with any agallochum that answers the above characters. In its place have been substituted woods of an inferiour kind, probably the aspalathus, lignum aquilæ, and calambour of authors; which are faid to be woods of the nature of agallochum, but, when in their greatest perfection, far weaker.

(a) Observ. physico-chym. lib. i. obs. 4. Not. ad Poterium, p. 487. De medicament. balfamic. §. 15.

LIGNUM CAMPECHENSE.

LIGNUM CAMPECHENSE Pharm. Edinb. Lignum tinetile campechense Pharm. Lond. Lignum campescanum & lignum indicum Mont. exot. Lignum campechianum, species quædam brasil Sloan. Lignum sappan quibusdam. Campeachy wood or Logwood: the wood of a prickly pod-bearing tree (Hæmatoxylum Campechianum Linn.) a native of Campeachy, in the bay of Honduras; from whence the wood is brought over in large compact hard logs of a red colour.

This wood, imported from America as a dying drug, has of late been introduced into medicine, and found to be a very useful restringent and corroborant, in diarrheas, dysenteries, and other disorders from a laxity of the solids. It has a sweetish subastringent taste, and no remarkable smell: extracts made from it, by water and spirit, have a great degree of sweetness, mixed with a mild grateful astringency. It gives a deep purplish red tincture both to watery and spirituous menstrua*(a); and frequently tinges the stools, and sometimes the urine, of the same colour: of this the patient ought to be apprised, that he may not be alarmed by judging the colour of the discharge to be owing to blood.

Watery menstrua readily extract part of the virtue of this wood, but are very difficultly made to take up the whole. To promote the extraction, the wood should previously be reduced into fine powder, which is to be strongly boiled in the water, in the proportion, for example, of a pound to a gallon, till half the liquor is wasted: the powder will still give a considerable impregnation to the same quantity of fresh water, and this repeatedly for four or five times or oftener: the extract obtained by inspissating the decoctions, of a dark blackish colour in the mass, tinges water of a sine red, like that of the liquors before inspissation, but does not totally dissolve: it is given in doses of from ten grains to a scruple and upwards. Rectified spirit takes up more from the logwood than watery menstrua. Some digest the powdered wood in four times its weight of spirit, and afterwards boil it in water: the matters taken up

Extract. lign.campechenf. Phar. Lond.

Phar. Edinb.

[&]quot;(a) Pure rain water acquires only a deep orange or mahogany colour from logwood; and rectified spirit a fine yellowish red. The purple hue seems to be communicated by some extraneous saline matter, as the selenitic or aluminous salts in hard spring water. A very small quantity of fixed alkali will also give it still more perceptibly.

by the two menstrua are then united into one extract, by inspissating the watery decoction to the confistence of honey, and then gradually stirring in the inspissated spirituous tincture.

LIGNUM RHODIUM.

LIGNUM RHODIUM Pharm Lond. RHODIUM OF ROSEWOOD: the wood or root of a tree of which we have no certain account; brought from the Canary islands, in long crooked pieces, full of knots, externally of a whitish colour, internally of a deep yellow, with a reddish cast. The largest, smoothest, straightest, heaviest, and deepest coloured pieces should be chosen; and the small, thin, pale, light ones rejected.

This wood has a flightly bitterish, somewhat pungent, balsamic tafte, and a fragrant smell, especially when scraped or rubbed, resembling that of roses. Digested in rectified spirit, it gives out pretty readily the whole of its active matter, and tinges the menstruum of a reddish yellow colour: on committing to distillation the filtered tincture, the spirit brings over little or nothing of its flavour; the fine smell, as well as the balfamic pungency, of the rhodium, remaining nearly entire in the inspiffated extract, which proves tenacious and adhesive like the turpentines. Infused in water, it gives out likewise great part of its fmell and tafte, together with a bright yellow colour: in evaporation, the water carries off the specific flavour of the wood, leaving in the extract only a flight pungency and bitterishness. Distilled with water, it gives over, fomewhat difficultly and flowly, a highly odoriferous effential oil, at first of a gold colour, by age turning reddish, amounting, Oleum ligni if the rhodium is of a good kind, to about one ounce from fifty: the distilled water is likewise agreeably impregnated with the fragrance of the rhodium, and resembles that of damask roses.

The effential oil is used as a perfume, for scenting pomatums, &c. and in this light only the rhodium wood is generally regarded. It promifes, however, to be applicable to more important purposes, and bids fair to prove a valuable cordial and corroborant.

LILIUM.

LILIUM ALBUM Pharm. Edinb. Lilium album flore erecto & vulgare C. B. Lilium candidum Linn. WHITE LILY: a plant with a fingle straight round stalk, clothed with oblong, acuminated, thick, smooth, pale green, ribbed leaves, which have no pedicles; bearing on the top several elegant, naked, white, upright, hexapetalous, bell-shaped flowers, which open successively, and are followed each by an oblong triangular capsule, divided into three cells sull of brownish seeds: the root is a single bulb, composed of sleshy scales, with several sibres at the bottom. It is perennial, a native of Syria and Palestine, common in our gardens, and flowers in June.

THE flowers of the white lily have a pleafant sweet smell, and a slightly mucilaginous taste. Their odorous matter is of a very volatile kind, being totally dissipated in drying, and totally carried off in evaporation by rectified spirit as well as water: both menstrua become agreeably impregnated with it by infusion or distillation, but no essential oil has been obtained, though many pounds of the flowers were submitted to the operation at once. The principal use of these flowers is for flavouring expressed oils; which, by insolation with fresh parcels of them continued about three days each time, are supposed to receive from them, along with their smell, an anodyne and nervine virtue. The distilled water has been sometimes employed as a cosmetic.

Oleum liliorum, fusinum, &c.

The roots also have been used chiefly for external purposes; as an ingredient in emollient and suppurating cataplasms: they abound with a strong mucilage, and do not seem to have much active matter besides. Gerard indeed relates, that several persons were cured of dropsies, by the constant use, for a month or six weeks, of bread made of barleymeal with the juice of white lily roots: but there are examples of similar cures being obtained by the use of common dry bread; and probably in one case, as well as in the other, abstinence from liquids was the remedy.

LILIUM CONVALLIUM.

LILIUM CONVALLIUM Pharm. Edinb. (a) Lilium convallium album C. B. Convallaria, Maianthemum. Convallaria maialis Linn. Lily of the valley, May Lily: a plant with two or three oblong, acuminated, ribbed leaves; in the bosoms of which arises a naked stalk bearing a number of small, naked, white, drooping, bell-shaped, monopetalous flowers, cut about the edges into six segments, and followed by red berries: the roots are long, slender, and white. It is perennial, grows wild in woods and shady places, and slowers in May.

THE flowers of this plant have a fragrant delightful smell, and a penetrating bitterish taste; both which they readily impart to watery and to spirituous menstrua. Their odorous matter, like that of the white lily, is very volatile; being totally dissipated in exsiccation, and elevated in distillation both by water and rectified spirit: there is no appearance of essential oil in either distillation; nor does the distilled spirit turn milky on the admixture of water, as those spirits do, which are impregnated with actual oil. The pungency and bitterness, on the other hand, reside in a fixt matter, which remains entire both in the watery and spirituous extracts, and which, in this concentrated state, approaches, as Cartheuser observes, to hepatic aloes.

It is principally from the volatile parts of these flowers, that medicinal virtues have been expected, in nervous and catarrhous disorders; but probably their fixt parts also have no small, perhaps the greatest, share in their esticacy. The flowers, dried and powdered, and thus divested of their odoriferous principle, prove strongly sternutatory. Watery or spirituous extracts made from them, given in doses of a scruple or half a dram, act as gentle stimulating aperients and laxatives; and seem to partake of the purgative virtue, as well as of the bitterness, of aloes.

The roots have nothing of the fine smell which is admired in the flowers, but discover to the taste a greater degree of penetrating bitterness. The bitter matter appears to be of the same kind in these as in the flowers; being equally extracted by water and spirit; remaining entire behind upon inspissating the tinctures or insusions; acting as a sternuta-

tory when fnuffed up the nose, and as a laxative or purgative when taken internally.

The leaves have the same kind of bitterness, in a lower degree, mixed with a considerable roughness, and a slight sweetishness.

LIMACES.

LIMACES terrestres sive Cochlææ terrestres Pharm. Edinb.(a) The SNAIL: an animal, lodged in a short thick turbinated shell, whose aperture is closed in the winter with a kind of cement. The large ash-coloured snail is said to be the species intended for medicinal use; but the smaller, dark-coloured, spotted, striped fort, more common in gardens, is taken indiscriminately, and appears to be not at all different in quality from the other.

This animal abounds with a viscid slimy juice, which it readily gives out, by boiling, to milk or water, so as to render them thick and glutinous. The decoctions in milk are apparently very nutritious and demulcent, and stand recommended in a thin acrimonious state of the humours, in consumptive cases, and emaciations.

LIMONES.

LIMONES Pharm. Lond. Limonia mala Pharm. Edinb. LEMONS: the fruit of the malus limonia fructu acido Pharm. Lond. malus limonia acida C.B. Citrus Limon Linn. a tree refembling the orange; from which it differs chiefly in the leaves having no appendages at the bottom; and in the fruit having a nipple-like production at the end: it is a native of Afia, and cultivated in the warmer parts of Europe, from whence we are supplied with the fruit. There are many varieties of this tree in regard to the fruit: by Linnæus, the several citrons, as well as lemons, are reckoned varieties of one species, which is distinguished from those of the orange kind, only by the pedicles of the leaves being naked. The terms citron and lemon have been often confounded together; what is commonly called citron by the French (b) and Germans (c) being our lemon, and their lemon our citron.

(c) Hoffmann, Differt. de citriis, Opera omnia, supplement. ii. par. i. p. 720.

⁽a) Expunged. (b) Codex medicamentarius facultatis Parifiensis, p. xxxviii. & lxx.

MATERIA

THE yellow rind of lemons is a grateful aromatic, of common use in stomachic tinctures and infusions, and for rendering other medicines acceptable to the palate and stomach: its flavour is one of those which is best adapted for accompanying medicines of the bitter kind. It is less hot than orange peel, and yields in distillation a less quantity of effential oil: the oil is extremely light, almost colourless, in smell nearly Ot. Billat. as agreeable as the fresh peel, and frequently employed as a perfume: it is generally brought to us from the fouthern parts of Europe, under the name of effence of lemons. The flavour of the lemon peel is more Effentia liperishable in keeping than that of orange peel, yet does not rise so easily in distillation with spirit of wine: for a spirituous extract, prepared from the rind of lemons, retains the aromatic taste and smell of the peel in a much greater degree than an extract prepared in the fame manner from that of oranges. After digestion in the spirit, lemon peel proves tough, that of oranges crifp.

The juice of lemons differs from that of oranges only in being more acid. Six drams of it faturate about half a dram of fixt alkaline falt: this mixture, with the addition of a fmall quantity of some grateful aromatic water or tincture, as fimple cinnamon water, is given in cases of nauseæ and reachings, and generally abates, in a little time, the severe vomitings that happen in fevers, when most other liquors and medicines are thrown up as foon as taken: it is used also as a saline aperient in icterical, hydropical, inflammatory and other diforders. A fyrup made Syrup.efucby dissolving fifty ounces of fine sugar in a quart + or two pounds and colimonum + Phar. Lond. a half ‡ of the depurated juice, is mixed occasionally with draughts and ‡ Phar. Edin. juleps as a mild antiphlogistic, and fometimes used in gargarisms for inflammations of the mouth and tonfils.

cort. limon. Pharm. Edin.

monum P.L.

LINARIA.

LINARIA Pharm. Edinb. (a) Linaria vulgaris lutea flore majore C. B. Ofyris, linaria, sive urinaria Lobel. Antirrbinum linaria Linn. TOADFLAX: a plant with fmooth round blueish stalks, and numerous, oblong, narrow, pointed leaves; greatly refembling the efula minor or pine spurge, so as scarcely to be otherwise distinguishable, before flowering, than by its wanting the milky juice with which the efula abounds :

on the tops of the stalks and branches appear spikes of yellow, irregular, monopetalous, gaping flowers, with a long tail behind, followed by roundish bicapsular seed-vessels: it is perennial, grows wild about the sides of dry fields, and flowers in June and July.

THE leaves of this herb have a bitterish somewhat saline taste: and when rubbed betwixt the fingers, yield a faint smell, resembling that of elder. Taken internally, they are said to be powerfully resolvent, diuretic, and purgative: their principal use, however, has been external, in unguents and cataplasms, for painful swellings of the hemorrhoidal vessels; against which they have been said to be particularly effectual.

LINGUA CERVINA.

LINGUA CERVINA seu Scolopendrium Pharm. Edinb. (a) Lingua cervina officinarum C. B. Phyllitis Gerard. Asplenium Scolopendrium Linn. Harts-tongue: a plant with long, uncut, narrow bright green leaves, set on long hairy pedicles, and nipt at the bottom: it has no stalks or manifest flowers; the seeds are a fine dust, lying in large, rough, brown, transverse streaks on the backs of the leaves. It is perennial, and found green at all times of the year, in moist, shady, strong places.

The leaves of harts-tongue stand recommended as aperients and corroborants, in obstructions of the hypochondriacal viscera, laxities of the intestines, and some disorders of the breast: they have been chiefly used in apozems and infusions, along with maidenhair, spleenwort, and other plants of the same kind, with which they appear to agree in virtue. To the taste they are slightly roughish and sweetish: with solution of chalybeate vitriol, they strike a blackish colour. When fresh, they yield, on being rubbed or bruised, a faint unpleasant smell, which in drying is in great part dissipated.

LINI SEMEN.

LINI SEMEN Pharm. Lond. Lini vulgaris semen Pharm Edinb. Lini sativi C. B. Lini usitatissimi Linn. Linseed: reddish-brown,

(a) Expunged.

gloffy, flippery, flat, pointed nearly oval feeds, of the common flax; an annual herb, cultivated in fields, on account of the mechanic uses of its tough filamentous rind.

THESE feeds have an unctuous, mucilaginous, fweetich tafte, and no remarkable fmell. On expression, they yield a large quantity of oil; Oleum exwhich, when carefully drawn, without the application of heat, has no preffum feparticular taste or slavour, though in some properties it differs consider- Phar. Edinb. ably from most of the other oils of this kind; not congealing in winter; not forming a folid foap with fixt alkaline falts (a); acting more powerfully, as a menstruum, on sulphureous bodies, than any other expressed oil that has been tried. The feeds, boiled in water, yield a large proportion of a strong flavourless mucilage: to rectified spirit they give out little or nothing.

Infusions of linseed, like other mucilaginous liquors, are used as emollients, incrassants, and obtunders of acrimony, in heat of urine, stranguries, thin defluxions on the lungs, and other like disorders: a spoonful of the seeds, unbruised, is sufficient for a quart of water, larger proportions rendering the liquor difagreeably flimy. The mucilage obtained by inspissating the insusions, or decoctions, is an excellent addition for reducing difguftful powders into the form of an electuary; occasioning the compound to pass the fauces freely, without sticking or discovering its taste in the mouth. The expressed oil is supposed to be more of a healing and balfamic nature than the other oils of this class; and has been particularly recommended in coughs, spitting of blood, colics, and conftipations of the belly. The feeds in fubftance, or the matter remaining after the expression of the oil, are employed externally, in emollient and maturating cataplasms. In some places, these seeds, in times of scarcity, have supplied the place of grain, but appeared to be an unwholesome, as well as an unpalatable food: Tragus relates, that those who fed upon them in Zealand, had the hypochondres in a fhort time distended, and the face and other parts swelled; and that not a few died of these complaints.

(a) Geoffroy, Memoires de l'acad. roy. des sciences de Paris, pour l'ann. 1741.

LINUM CATHARTICUM.

LINUM CATHARTICUM Pharm. Edinb. (a) & Linn. Linum pratense storibus exiguis C. B. Chamælinum. Purging flax or Mill-mountain: a small plant, with little oblong smooth leaves, having one vein or rib running along the middle, joined in pairs close to the stalks, which are round, slender, reddish, divided towards the upper part into fine branches, bearing on the tops white pentapetalous uncut flowers, followed each, as in the common slax, by a roundish, ribbed, acuminated capsule, containing ten flattish slippery seeds in as many cells. It is annual, and grows wild on chalky hills, and in dry pasture grounds.

This herb is faid to be an effectual and fafe cathartic: an infusion of an handful of the fresh leaves in whey or white wine, or a dram of the leaves in substance with a little creme of tartar and aniseeds, are directed for a dose. Linnæus recommends an infusion of two drams of the dry leaves as a mild laxative. Their taste is bitterish and disagreeable.

LIQUIDAMBRA.

LIQUIDAMBRA Pharm. Edinb. (a) Ambra liquida Pharm. Argent. Liquidamber: a refinous juice, of a yellow colour inclining to red, at first about the confistence of turpentine, by age hardening into a solid brittle resin; obtained from the tree that yields liquid storax, syrax aceris folio Raii, Liquidambar styracistua Linn. growing in Virginia, Mexico, and other parts of America, and bearing the colds of our own climate.

This juice has a moderately pungent, warm, balfamic tafte; and a very fragrant smell, not unlike that of storax calamita heightened with a little ambergris. It was formerly in common use as a persume, and might probably be applied to valuable medicinal purposes, but it is not at present much regarded, different artificial compositions having been often substituted to it in the shops.

LITHOSPERMUM.

LITHOSPERMUM seu Milium solis Pharm. Edinb. (a) Lithospermum majus erectum C. B. Lithospermum officinale Linn. Gromwell: a rough plant, with stiff branched stalks, oblong acuminated leaves set alternately without pedicles; and whitish monopetalous flowers, scarcely longer than the cup, divided into five obtuse sections, followed by little roundish, hard, pearl-like seeds inclosed in the cup. It is perennial, grows wild in dry fields and by road-sides, and flowers in May and June.

THE feeds of gromwell have been accounted notably diuretic; and recommended for cleanfing the kidneys and urinary passages from viscid mucous matters, and promoting the expulsion of gravel. Their virtues do not appear to be very considerable; they have no smell, and their taste is little other than farinaceous. They have long been discarded from practice.

* LOBELIA.

LOBELIA Pharm. Edinb. Rapunculus galeatus virginianus flore majore violaceo Morison. Lobelia siphilitica Linn. Blue Cardinal-flower: an herbaceous perennial plant, with an erect stalk three or four feet high, and ovate-lanceolate subservated leaves, bearing long spikes of labiated, irregular, blue flowers, each with five stamina having connate antheræ, succeeded by a bilocular capsule, containing many small seeds. The whole plant has a milky juice, and something of a rank smell. It grows in moist places in Virginia, and bears the winters of our climate.

The root of this plant confifts of white fibres, a line in thickness, and about two inches in length. It resembles tobacco in its taste, which dwells long on the tongue, and is apt to excite vomiting. It was long a famous secret among the North American Indians for the cure of the venereal disease. The secret was purchased by Sir William Johnson, and has been made public in the writings of Bartram, Kalm, and others.

(a) Expunged.

The dose and mode of administering this medicine are not exactly defined, but the following directions are given as the most accurate. A decoction is made of a handful of the roots in three measures of water. Of this, half a measure is taken in the morning fasting, and repeated in the evening; and the dose is gradually increased till its purgative effect becomes too violent, when the medicine is for a time to be intermitted, and then renewed, till a perfect cure is effected. One dose daily is fufficient during the latter part of the treatment; and the regimen during the whole process is to be equally strict with that observed in a course of mercurial falivation. From the third day, the ulcers are to be well washed twice daily with the decoction; and it is said that when they are very deep and foul, the Indians sprinkle them with powder of the internal bark of the spruce-tree. By this method we are assured that inveterate venereal complaints are cured without the aid of mercury; and the Edinburgh college feem to give credit to the efficacy of the lobelia, by receiving it into their latest catalogue of simples.

LUJULA.

LUJULA Pharm. Lond. Trifolium acetofum vulgare C. B. Oxys alba Gerard. Alleluja, oxytriphyllum, & panis cuculi Quorundam. Oxalis Acetosella Linn. WOOD SORREL: a plant, with the leaves and flowers iffuing on separate pedicles from the root: the leaves are broad, shaped fomewhat like a heart, and stand three together: the flowers are folitary, whitish, monopetalous, divided deeply into five segments, followed by angular capfules, which burft on being touched, and shed numerous fmall brownish feeds. It is perennial, grows wild in woods, and flowers in April.

THE leaves of the wood forrel are useful faline antiseptics and antiphlogistics; fimilar, both in taste and in medicinal virtue, to those of the acetofæ or common forrels, but somewhat more acid, and rather more grateful both to the palate and stomach. Beaten with thrice their weight Confervalu. of fine fugar, they form a grateful fubacid conferve. Their expressed juice, depurated, is a very agreeable acid: duly inspissated, and set to shoot, it yields a crystalline acid salt of the same nature with that of the sorrels:

julæ Ph. Lon.

the faline matter feems to amount to nearly one hundredth part of the weight of the fresh leaves.

LUMBRICUS.

LUMBRICUS TERRESTRIS Pharm. Edinb. (a) Vermes terrestres. Earth worms. These insects are supposed to have a diuretic
and an antispasmodic virtue. The faculty of Paris directs them to be
prepared for medicinal use, by washing and drying them with a moderate
heat. Moistened with wine, or vinous spirits, to prevent their putresying,
and set in a cellar, they are almost wholly resolved in a few days into
a slimy liquor, which is said, when mixed with alkaline salts, to yield
crystals of nitre.

LUPULUS.

LUPULUS Pharm. Parif. Lupulus mas & femina C. B. Lupulus falictarius Ger. Humulus Lupulus Linn. Hop: a rough plant, with very long, angular, climbing hollow stalks, and broad ferrated leaves, cut generally into three or five sharp-pointed sections, and set in pairs at the joints: on the tops grow loose scaly heads, with small slat seeds among them. It is found wild in hedges and at the bottoms of hills, in England and other parts of Europe, but commonly cultivated in large plantations. It is perennial, and ripens in August or September its leafy heads, which are cured by drying with a gentle heat on kilns made for that purpose.

Hops have a very bitter taste, less ungrateful than most of the other strong bitters, accompanied with some degree of warmth and aromatic slavour. They give out their virtue by maceration without heat, both to rectified and proof spirit; and, by warm insusion, to water: to cold water they impart little, though macerated in it for many hours. The extracts obtained both by watery and spirituous menstrua, particularly by the latter, are very elegant balsamic bitters, and promise to be applicable to valuable purposes in medicine; though the hop is at present scarcely regarded as a medicinal article, and scarcely otherwise used than

for the preserving of malt liquors; which, by the superaddition of this balsamic, aperient, diuretic bitter, become less mucilaginous, more detergent, more disposed to pass off by urine, and in general more falubrious.

LYCOPERDON.

LYCOPERDON five Crepitus lupi Pharm. Edinb. (a) Fungus rotundus orbicularis C.B. Bovista officinarum Dill. Lycoperdon Bovista Linn. Puffball: a round or egg-shaped whitish fungus, with a very short or scarcely any pedicle, growing in dry pasture grounds; when young, covered with tubercles on the outside, and pulpy within; by age becoming smooth without, and changing internally into a very fine, light, brownish dust.

THE dried fungous matter and the dust of lycoperdon have been long used among the common people, particularly in Germany, for restraining the bleeding of wounds, and immoderate hemorrhoidal fluxes, and drying up running ulcers. In some late trials, the dust has been found to produce the same effect, in stopping hemorrhages after amputation, as the celebrated agaric of the oak.

MACIS.

MACIS Pharm. Lond. Macis officinarum C. B. MACE: a pretty thick, tough, unctuous membrane, reticular or variously chapt, of a lively reddish yellow colour approaching to that of saffron, enveloping the shell of the fruit whose kernel is the nutmeg. The mace, when fresh, is of a blood-red colour, and acquires its yellow hue in drying: it is dried in the sun, upon hurdles fixed above one another, and then, as is said, sprinkled with sea-water, to prevent its crumbling in carriage.

MACE has a pleasant aromatic smell, and a warm, bitterish, moderately pungent taste. It is in common use as a grateful spice; and appears to be, in its general qualities, nearly similar to the nutmeg, both as the subject of medicine and of pharmacy. The principal difference consists

in the mace being much warmer, more bitterish, less unctuous, and sitting easier on weak stomachs; in its yielding by expression a more studied oil, and in distillation with water a more subtile volatile one. What is called in the shops expressed oil of mace is prepared, not from this spice, but from the nutmeg.

MAGNESIA.

MAGNESIA ALBA Pharm. Edinb. MAGNESIA: a fine white earth; foluble readily in all acids, the vitriolic as well as the others, into a bitter purgative liquor.

This earth has not hitherto been found naturally pure or in a separate state: it was for several years a celebrated secret in the hands of some particular persons abroad, till the preparation was made public by Lancisi in the year 1717 (a), and afterwards by Hossman in 1722 (b). It was then extracted from the mother-lye, or the liquor which remains after the crystallization, of rough nitre; either by precipitation with a solution of fixt alkaline salt; or by evaporating the liquor, and calcining the dry residuum, so as to dissipate the acids by which the earth had been made dissoluble.

The magnefia, in this mother-lye, appears to have proceeded from the vegetable ashes, which are either made ingredients in the compositions from which nitre is obtained, or else added in the elixation of the nitre: for the ashes of different woods, burnt to perfect whiteness, and freed from their alkaline salt, were found to be, in part, of the same nature with the true magnesia (c). But as quicklime also, in most of the German, French, and other European nitre-works, is commonly employed in large quantity, the earth obtained from the mother-lyes of those works is rather a calcareous earth than magnesia. What is now brought from abroad, under the name of magnesia, gives plain proofs of its calcareous nature, by its burning into quicklime, and forming a selenites with the vitriolic acid.

⁽a) Annot. in Mercati metallothec. vatican. Arm. ii. cap. x. p. 50.

⁽b) Observationes Physico-chymica, lib. ii. obs. 2.

⁽c) Of vegetable ashes, moderately or strongly calcined, only a part was found to dissolve in acids, and this part appeared to be perfect magnesia. It is probable that the remainder might be reduced to the same state by repeating the calcination.

The true magnesia is obtained in great purity, from a filtered solution of fal catharticus amarus, by adding a filtered solution of any alkaline salt. *This, by its superiour affinity with the vitriolic acid of the sal catharticus, precipitates its earthy basis, which is the magnesia. The method of conducting the process is thus directed in the last Edinburgh dispensatory. Dissolve separately equal weights of sal catharticus, and any pure fixed alkaline salt, in double their weight of water. Strain, and then mix them, and immediately add eight times the quantity of hot water. Let the liquor boil a while over the fire, and at the same time agitate it. Then let it stand till the heat be abated; and strain it through a linen cloth, on which the magnesia will be left. This is to be washed by affusions of pure water till it become perfectly tasteless.

*A method of preparing magnesia in the most perfect and convenient manner, was published by Mr. Henry in Vol. II. of the Medical Transactions. The same writer, likewise, in a publication of Experiments and Observations on various subjects, recommends the calcination of magnesia, as rendering it a fitter medicine in certain cases. Magnesia is found by experiment to contain above half its weight of fixed air. The evolution of this in the stomach may increase flatulency, and cause uneasiness in weak bowels. By strong calcination the air is expelled from magnesia, while its purgative virtue remains unimpaired; nor does it acquire any of the acrimony or causticity of lime. The Edinburgh college have now

received this preparation, under the title of magnefia ufta.

The magnesia is recommended by Hossiman as an useful antacid, a safe and inossensive laxative in doses of a dram or two, and a diaphoretic and diuretic, when given in small doses, as sisteen or twenty grains. Since this time, it has had a considerable place in the practice of foreign physicians, and has of late come into some esteem among us, particularly in heart-burns, and for preventing or removing the many disorders which children are thrown into from a redundance of acid humours in the first passages. It is preferred, on account of its laxative quality, to the testaceous and other absorbent earths, which, unless gentle purgatives are given occasionally to carry them off, are apt to lodge in the body, and occasion a costiveness very detrimental to infants. It must be observed, however, that it is not the magnesia itself which proves laxative, but the saline compound resulting from its coalition with acids: if there are no acid juices in the stomach to dissolve it, it has no sensible operation, and

in fuch cases it may be rendered purgative by drinking any kind of acidulous liquors after it. All the other known foluble earths yield with

acids, not purgative, but more or less aftringent compounds.

It may be proper to observe, that the name magnesia has been principally Magnesia, Manganese, applied to a substance of a very different kind; a native mineral, found in iron mines, and in the lead mines of Mendip hills, in Somersetshire, usually of a dark grey colour, sometimes bright and striated like antimony, fometimes dull, with only a few small striæ; remarkable for communicating, to a large proportion of glass in fusion, a purplish or red tinge, which disappears on a continuance of the fire, at the same time destroying the effect of many other colouring matters, and rendering foul or coloured glass clear: supposed to be an ore of iron, and recommended medicinally, when calcined by a strong fire, as an astringent; but yielding no iron, or marks of iron, on any of the common trials by which that metal is diftinguished in ores; and in its nature and composition as yet little known. Mr. Pott relates, that on being calcined with fulphur, and afterwards elixated with water, it yielded a large quantity of a white crystalline salt, of a bitterish astringent taste, followed by a kind of sweetness; and that the salt, after strong calcination, tasted like burnt alum, but more acid(a); from whence it may be prefumed, that this mineral confifts in great part of an earth analogous to that of alum, which, in combination with acids, makes one of the strongest styptics.

vitrariorum & minerologorum.

MAJORANA.

MAJORANA Pharm. Lond. & Edinb. Majorana vulgaris C. B. Origanum Majorana Linn. SWEET MARJORAM: a low plant, with flender, fquare, branched, woody stalks; and little, oval, somewhat downy leaves, fet in pairs: on the tops grow fealy heads of small whitish labiated flowers, whose upper lip is erect and cloven, the lower divided into three fegments. It is fown annually in gardens, for culinary as well as medicinal uses: the feeds, which rarely come to perfection in this country, are procured from the fouth of France, where the plant is faid to be indigenous.

MATERIA MEDICA.

THE leaves and tops of marjoram have a pleasant smell, and a moderately warm aromatic bitterish taste. Infusions of them in water, in colour brownish, smell pretty strongly, and taste weakly and unpleafantly of the herb: the blackish green tinctures, made in rectified spirit, have less smell, but a stronger and more agreeable taste. distillation with water, an essential oil is obtained, amounting, as Hoffman observes, to about one ounce from fixty-four of the leaves flightly dried; when carefully drawn, of a pale yellow colour; by age, or too hasty fire in the distillation, contracting a reddish hue; of a very hot penetrating tafte, and in smell not near so agreeable as the marjoram itself: the remaining decoction, thus divested of the volatile aromatic matter, is weakly, but unpleasantly bitterish and austere. Great part of the aromatic matter of the herb rifes also in the inspissation of the spirituous tincture, and impregnates the distilled spirit: the remaining extract is stronger in taste than that made with water, its quantity being less, but has not much of the warmth or flavour of the marjoram.

This plant has been chiefly recommended in diforders of the head and nerves, in uterine obstructions and mucous discharges proceeding from what is called a cold cause (that is, from a laxity and debility of the solids, and a sluggish state of the juices) and in the humoural asthmas and catarrhs of old people. The powder of the leaves, their distilled water, and the essential oil properly diluted, are agreeable errhines, and accounted particularly useful in pituitous obstructions of the nostrils, and disorders of the olfactory organs.

MALABATHRUM.

MALABATHRUM Pharm. Lond. Tamalapatra, Folium indum. Indian leaf: the leaf of the cinamomum five canella malavarica & javanensis C. B. Laurus Cassia Linn. or casia-lignea tree, brought from the East Indies. It is of a firm texture; of an oblong oval figure, pointed at both ends; smooth and glossy on one side, which is the upper, and less so on the lower; of a yellowish green colour on the former, and a pale brownish on the latter; furnished with three ribs, running its whole length, very protuberant on the lower side, and two smaller ones which bound the edges.

THESE

Ol. effent. majoranæ Pharm.Lond.

THESE leaves have a remarkable affinity, in one respect, with the casia or bark of the tree, both the leaves themselves and their pedicles being, like it, extremely mucilaginous: chewed, they render the faliva flimy and glutinous: infused in water, they give out a large proportion of a strong tenacious mucilage. But of the aromatic flavour, which is strong in the bark, the leaves, as brought to us, have very little: they fcarcely discover any warmth or pungency to the taste, and have little or no smell unless well rubbed, when they yield an agreeable, though weak, spicy odour. They are no otherwise made use of than as an ingredient in mithridate and theriaca; and are, when in their greatest perfection, far inferiour to the mace which our college directs as a fuccedaneum to them.

MALVA.

MALVA Pharm. Lond. & Edinb. Malva silvestris folio sinuato C. B. Malva sylvestris Linn. Common Mallow: a plant with firm branched stalks, and roundish, notched leaves, set alternately on long pedicles: in their bosoms come forth bell-shaped monopetalous flowers, deeply divided into five heart-shaped sections, of a pale purplish or whitish colour variegated with deeper streaks, followed by a number of capfules fet in form of a flat disk, and containing each a kidney-shaped feed: the root is long, flender, and whitish. It is perennial, common in uncultivated grounds, and found in flower throughout the fummer.

THE leaves and flowers of the mallow are in taste mucilaginous, and The leaves were formerly of some esteem, as of no remarkable fmell. an emollient or laxative dietetic article, in dry constipated habits in the warmer climates: at prefent, infusions or decoctions of the leaves and flowers, and a conferve made by beating the fresh flowers with thrice Conf. florum their weight of fine fugar, are fometimes directed in dyfuries, heat and sharpness of urine, and other like complaints; but the principal use of the herb is in emollient glysters, cataplasms, and fomentations. The roots have been recommended in diforders of the breast, and though now difregarded, may perhaps deserve some notice: they have a soft sweet taste, without any particular flavour, approaching in some degree to that of liquorice: an extract made from them by rectified spirit of wine is of great fweetness.

malvæ P. L.

MANNA.

MANNA Pharm. Lond. & Edinb. Manna feu Ros calabrinus Pharm. Parif. Manna: a sweet juice obtained from certain ash trees (a) in the southern parts of Europe, particularly in Calabria and Sicily, exuding from the leaves, branches, or trunk of the tree, and either naturally concreted, or exsiccated and purished by art.

* There are three ways in which manna is collected in Calabria. From the middle of June to the end of July, a very clear liquor exudes fpontaneously from the trunk and branches of the tree, which by the sun's heat concretes into whitish masses, which are scraped off the next morning with wooden knives, and dried in the sun. This is called Manna in the tear. At the beginning of August, when this ceases to slow, the peasants make incisions in the bark, whence a juice flows, which concretes in larger masses and of a redder colour. This is the sat or common Manna. Besides these forts, a third is procured by receiving the spontaneous exudations in June and July on straws or chips of wood fastened to the tree. This is the cannulated or slaky Manna, and is accounted the finest of the three.

Juices of the same nature are collected, in the eastern countries, from other trees and shrubs (b): and similar exudations are sometimes found on different kinds of trees in Europe, as particularly on the larch in the Briançonois in Dauphiny. How far the manna juices of different vegetables differ from one another, is not well known: but thus much is certain, that one and the same tree affords mannas very considerably different, in their colour, in their taste, and in their disposition to assume a solid concrete form; that is, in their purity, or the greater or less admixture of oily or resinous matter.

The best fort of the officinal or Calabrian manna is in oblong pieces or slakes, moderately dry, friable, very light, of a whitish or pale yellow colour, and in some degree transparent: the inferiour kinds are moist, unctuous, and brown. Both sorts are said to be sometimes counterfeited by compositions of sugar, honey, and purgative materials;

Manna brigantiaca.

compositions

⁽a) Ornus: Fraxinus rotundiore folio, & fraxinus humilior minore & tenuiore folio C. B. Fraxinus Ornus Linn.

⁽b) Vide Clusii exotic. lib. i. p. 164. Rauwolf itin. p. 74. Teixeira bift. Perf. p. 29.

compositions of this kind, in a solid or dry form, may be distinguished by their weight, compactness, and untransparency: both the dry and moist compositions may be distinguished by their taste, which is sensibly different from that of true manna, and with greater certainty by their habitude to menstrua.

This juice liquefies in a moift air, dissolves readily in water, and, by the assistance of heat, in rectified spirit also; the impurities only being left by both menstrua. On inspissating the watery solution, the manna is recovered of a much darker colour than at first. From the saturated spirituous solution, great part of it separates as the liquor cools, concreting into a slaky mass, of a snowy whiteness, and a very grateful sweetness: the liquor, remaining after the separation of this pure sweet part of the manna, leaves, on being inspissated, an unctuous, dark coloured, disagreeable matter, in greater or less quantity according as the manna made use of was less or more pure.

Manna, in doses of an ounce and upwards, proves a gentle laxative: it operates in general with great mildness, so as to be safely given in inflammatory or acute diftempers, where the stimulating purgatives have no place. It is particularly proper in stomachic coughs, or those which have their origin in the stomach; the manna, by its sweetness and unctuosity, contributing to obtund as well as to evacuate the offending humours: in this intention it is fometimes made into a linctus or lohoch, with equal quantities of oil of almonds and of fyrup of violets. fome constitutions, however, it acts unkindly, especially if given in confiderable quantity, occasioning flatulencies, gripes, and distensions of the belly; inconveniences which may be generally obviated by a fmall addition of fome grateful aromatic. It does not produce the full effect of a cathartic, unless taken in large doses, as two ounces or more, and hence is rarely employed in this intention by itself: it may be commodiously dissolved in the purging mineral waters, or acuated with the cathartic falts, or other purgatives: its efficacy is faid to be peculiarly promoted by cafia fiftularis, a mixture of the two purging more than both of them separately. See Casia.

MARGARITE.

MARGARITÆ Ph. Lond. Perlæ. Uniones. Pearls: small calculous concretions, of a bright semitransparent whiteness, sound on the inside of the shell of the concha margaritisera or mother-of-pearl sish, as also of certain oysters and other shell-sishes. The sinest pearls are brought from the East and West Indies: the oriental, which are most esteemed, have a more shining silver-like hue than the occidental, which last are generally somewhat milky: an inferiour sort is sometimes met with in the shell-sishes of our own seas, particularly on the coasts of Scotland. The coarse rough pearls, and the very small ones which are unsit for ornamental uses, called rag pearl and seed pearl, are those generally employed in medicine.

It is faid, that counterfeit pearls are often brought from China, made of pellets of clay coated with the white matter of oyster-shells. The clay may be distinguished by its acquiring an additional hardness in the fire, and resisting acids; whereas the true pearls calcine in the fire and become quicklime, and readily dissolve in acids; the vitriolic excepted, which precipitates them when previously dissolved by other acids.

These properties of the pearl, shew that it is an earth of the same kind with crabs-claws, oyster-shells, and the other calcareous animal absorbents. It has no other virtues than those of the other substances of this class, and does not possess those virtues in any greater degree than the common testacea.

MARRUBIUM.

MARRUBIUM Pharm. Lond. & Edinb. Marrubium album vulgare C. B. Marrubium vulgare Linn. WHITE HOREHOUND: a hoary plant, with square stalks, and roundish wrinkled indented leaves, set in pairs on long pedicles; in the bosoms of which come forth thick clusters of whitish labiated flowers, in striated cups, whose divisions terminate in sharp points or prickles. It is perennial, grows wild in uncultivated grounds, and flowers in June.

The leaves of horehound have a moderately strong smell, of the aromatic kind, but not agreeable, which by drying is improved, and in keeping

keeping for some months is in great part diffipated: their tafte is very bitter, penetrating, diffusive, and durable in the mouth. From these qualities, and their fensible operation, when taken in any considerable doses, of loosening the body, it may be presumed that this herb is a medicine of fome efficacy, and has no ill claim to the corroborant and aperient virtues, for which it is recommended, in humoural afthmas, and in menstrual suppressions, cachexies, and other chronical disorders proceeding from a viscidity of the fluids and obstructions of the viscera: a dram of the dry leaves in powder, or two or three ounces of the expressed juice, or an infusion of half a handful or a handful of the fresh leaves, are commonly directed for a dose. The dry herb gives out its virtue both to watery and spirituous menstrua, tinging the former of a brownish, the latter of a green colour: on inspissating the watery infusion, the smell of the horehound wholly exhales, and the remaining extract proves a strong and almost flavourless bitter: rectified spirit carries off likewise greatest part of the flavour of the herb, leaving an extract in less quantity than that obtained by water, and of a more penetrating bitterness.

MARUM.

MARUM SYRIACUM Pharm. Lond. Majorana syriaca vel cretical C. B. Marum cortusi J. B. Chamædrys maritima incana frutescens soliis lanceolatis Tourn. Origanum Syriacum Linn. Marum, Syrian herbemastich: a low shrubby plant, with small oval leaves, pointed at each end, set in pairs, without pedicles, of a dilute green colour above, and hoary underneath: in their bosoms appear solitary, purple, labiated slowers, wanting the upper lip; the lower lip is divided into five segments, the middlemost of which is larger than the rest, and hollowed like a spoon: each flower is followed by four roundish seeds inclosed in the cup. It is said to be a native of Syria, and of one of the Hieres islands, on the coast of Provence: in our climate it does not well bear severe winters without shelter.

THE leaves of marum have a bitterish, aromatic, very pungent taste; and when rubbed a little, yield a quick piercing smell, which provokes sneezing. They have been chiefly made use of as an ingredient in sternutatory powders, though, from their sensible qualities, they promise to

be applicable to more important purposes, and to have no ill title to the stimulating, attenuating, deobstruent, antiseptic virtues ascribed to them by Wedelius in a differtation on this plant: they seem particularly well adapted as an ingredient in the volatile oily aromatic spirits with which

their agreeable pungency in a great degree coincides.

The marum loses but little of its pungency on being dried, and in this respect it differs remarkably from many other acrid herbs, as those called antiscorbutic. It gives out its active matter partially to water, and completely to rectified spirit: the watery insusions, in colour yellow, though pretty strongly impregnated with the smell of the marum, have only a weak taste: the spirituous tinctures, in colour yellowish-green, are strongly impregnated with the taste, but have the smell in great measure covered by the menstruum. Distilled with water, it yields a highly pungent, subtile, volatile essential oil, similar to that of scurvygrass, but stronger, and of a less perishable pungency: the remaining decoction is little other than bitterish. Rectified spirit carries off likewise, in the inspissation of the spirituous tincture, a considerable share of the smell and pungency of the marum, but leaves much the greatest part concentrated in the extract; which, on being tasted, fills the mouth with a durable, penetrating, glowing warmth.

MARUM VULGARE.

MARUM VULGARE Pharm. Lond. Sampfucus sive marum mastichen redolens C. B. Thymbra hispanica majoranæ solio Tourn. Clinopodium quibusdam, mastichina gallorum J. B. Thymus mastichina Linn. Common herb-mastich: a low shrubby plant, with small oblong leaves, pointed at both ends, set in pairs, without pedicles: at the tops of the branches stand woolly heads, containing small white labiated slowers, whose upper lip is erect and cloven, the lower divided into three segments: each slower is sollowed by sour seeds inclosed in the cup. It grows spontaneously on dry gravelly grounds in Spain, and in the like soils it bears the ordinary winters of our own climate.

This plant is employed chiefly, like the foregoing, as an errhine. It is confiderably pungent, though far less so than the marum syriacum; and of a strong agreeable smell, somewhat resembling that of mastich.

MASTICHE.

MASTICHE.

MASTICHE Pharm. Lond. & Edinb. MASTICH: a concrete resin, obtained in the island Chio from the lentisk tree; brought over in small yellowish transparent brittle grains or tears. From transverse incisions made in the bark of the tree, about the beginning of August, the resin exudes in drops, which running down, and concreting on the ground, are thence swept up (a). The tree is raised also in several parts of Europe; but no resin has been observed to issue from it in these climates: nor do all the trees of this species, in the island Chio itself, afford this commodity.

This refin has a light agreeable fmell, especially when rubbed or heated: in chewing, it first crumbles, soon after sticks together, and becomes soft and white like wax, without impressing any considerable taste. It totally dissolves, except the earthy impurities, which are commonly in no great quantity, in rectified spirit of wine, and then discovers a degree of warmth and bitterness, and a stronger smell than that of the resin in substance: the colour of the solution is a pale yellow. Boiled in water, it impregnates the liquor with its smell, but gives out little or nothing of its substance: distilled with water, it yields a small proportion of a limpid essential oil, in smell very fragrant, and in taste moderately pungent. Rectified spirit brings over also, in distillation, the more volatile odorous matter of the mastich.

Mastich is recommended, in doses of from half a scruple to half a dram, as a mild corroborant and restringent, in old coughs, hemoptyses, diarrhæas, weakness of the stomach, &c. It is given either in substance, divided by other materials; or dissolved in spirit and mixed with syrups; or dissolved in water into an emulsion by the intervention of gum-arabic or almonds: the decoctions of it in water, which some have directed, have little or nothing of the virtue of the mastich. It is said that this resin is commonly employed as a masticatory, in Chio and among the Turkish women, for sweetening the breath, and strengthening the gums and teeth; and that when thus used, by procuring a copious excretion of saliva, it proves serviceable in catarrhous disorders.

MATRICARIA.

MATRICARIA Pharm. Lond. Matricaria vulgaris seu sativa C.B. Febrifuga Dorsten. Matricaria Parthenium Linn. Feverfew: a plant with firm branched stalks, and roughish leaves, each of which is composed of two or three pairs of indented oval segments set on a middle rib, with an odd one at the end, cut into three lobes: the flowers stand on the tops in the form of an umbel, consisting, each, of a number of short white petala, set round a yellow disk, which is followed by small striated seeds. It is biennial, or of longer duration; grows wild in hedges and uncultivated places, and slowers in June.

The leaves and flowers of feverfew have a strong, not agreeable smell, and a moderately bitter taste; both which they communicate, by warm insusion, to water and to rectified spirit. The watery insusions, inspissated, leave an extract of considerable bitterness, and which discovers also a saline matter, both to the taste, and in a more sensible manner, by throwing up to the surface small crystalline efflorescences in keeping: the peculiar slavour of the matricaria exhales in the evaporation, and impregnates the distilled water: on distilling large quantities of the herb, a yellowish strong-scented essential oil is found floating on the surface of the water. Rectified spirit carries off but little of the slavour of this plant in evaporation or distillation: the spirituous extract is far stronger in taste than that made with water, and more agreeable in smell than the herb itself. The quantity of spirituous extract, according to Cartheuser's experiments, is only about one sixth the weight of the dry leaves, whereas the watery extract amounts to near one half.

This herb is recommended as a warm, aperient, carminative bitter; and supposed to be particularly serviceable in semale disorders. It appears, from the above analysis, to be a medicine of no inconsiderable virtue, in some degree similar to camomile.

MECHOACANNA.

MECHOACANNA Pharm. Edinb. (a) Bryonia mechoacanna alba C. B. Convolvulus americanus mechoacan dictus Raii. Jalappa alba,

Rhabarbarum album Quibusdam. MECHOACAN: the root of an American convolvulus, (Convolvulus Mechoacanna Linn.) brought chiefly from a province in Mexico of the same name, in thin transverse slices, like jalap, but larger and whiter.

This root was first introduced, about the year 1524, and continued in esteem for a considerable time, as a mild cathartic, of very little taste or smell, not liable to offend the stomach, of slow operation, but essectual and safe: by degrees, it gave place to jalap, which has now, among us, almost wholly superseded its use. It seems to differ from jalap only in being weaker: the resins obtained from the two roots appear to be of the same qualities, but mechoacan scarcely yields one sixth part so much as jalap does, and hence requires to be given in much larger doses to produce the same effects. The dose of the mechoacan in substance is from one dram to two or more.

MEL.

MEL Pharm. Lond. Honey: a fweet vegetable juice; collected by the bee from the flowers of different plants, and deposited in the cells of the combs; from which it is extracted, either by spontaneous percolation through a sieve in a warm place, or by expression. That which runs spontaneously is purer than the expressed; a quantity of the waxy and other impurities being forced out along with it by the pressure, especially when the combs are previously heated. The best fort of honey is of a thick consistence, a whitish colour, an agreeable smell, and a very pleasant taste: both the colour and slavour are said to differ in some degree according to the plants which the bees collect it from.

Honey, exposed to a gentle heat, as that of a water bath, becomes thin, and throws up to the surface its waxy impurities, together with the meal or flower sometimes fraudulently mingled with it, which may thus be separated by despumation, so as to leave the honey pure. On continuing the heat, there rises a considerable quantity of aqueous sluid, impregnated with the sine smell of the honey: the inspissated residuum, like the honey at first, dissolves both in water and in rectified spirit, and promotes the union of oily and resinous substances with watery liquors. By treating the inspissated mass with moist clay, as practised

by the fugar-bakers for purifying fugar from its unctuous treacly matter, the unctuous parts of honey may in like manner be feparated, and its pure fweet matter obtained in the form of a folid, faline, white concrete.

This juice is an useful sweet, for medicinal as well as domestic purposes; more aperient and detergent than the simpler sweet prepared from the fugar cane; particularly ferviceable for promoting expectoration in disorders of the breast, and as an ingredient in cooling and detergent gargarisms. For these, and other similar intentions, it is sometimes mixed with vinegar, in the proportion of about two pounds to a pint, and the mixture boiled down to the confistence of a syrup; sometimes impregnated with the virtues of different vegetables, by boiling it in like manner with their juices or infusions, till the watery parts of the juice or infusion have exhaled and left the active matter incorporated with the honey. It excellently covers the tafte of purging falts and waters. The boiling of honey, though it diffipates great part of its odorous matter, and thus proves in some cases injurious to it, is in some cases also of advantage: there are particular constitutions with which honey remarkably difagrees, and in which even very fmall quantities occasion gripes, purging, and great diforder: by boiling, it lofes of that quality by which it produces these effects.

* The Edinburgh college feem at prefent of opinion that honey has no qualities which render it in any case preferable to sugar; since they have entirely expunged it, and all preparations in which it entered, from their last pharmacopæia.

MELILOTUS.

MELILOTUS Pharm. Edinb. (a) Melilotus officinarum germaniæ C. B. Lotus filvestris. Trifolium odoratum. Trifolium Melilotus officinalis Linn. Melilot: a plant with smooth oval striated leaves, standing three together, on slender pedicles; and round, striated, branched stalks, terminated by long spikes of papilionaceous flowers drooping downwards, which are followed by short thick wrinkled pods, containing, each, one or two roundish seeds. It is annual or biennial, and found in flower, in hedges and corn fields, greatest part of the summer.

(a) Expunged.

MELILOT

Oxymelfimplex Pb. Lon. MELILOT has been said to be resolvent, emollient, anodyne, and to participate of the virtues of camomile. In its sensible qualities, it differs very materially from that plant: its taste is unpleasant, subacrid, subsaline, but not bitter: when fresh, it has scarcely any smell; in drying, it acquires a pretty strong one, of the aromatic kind, but not agreeable. Linnæus observes, in the third volume of the Amænitates Academicæ, that distilled water of melilot, of little smell itself, remarkably heightens the fragrance of other substances. The principal use of this plant has been in glysters, somentations, and other external applications: it formerly gave name to one of the officinal plasters; which received from the melilot a green colour and an unpleasant smell, without any addition to its efficacy.

MELISSA.

MELISSA Pharm. Lond. & Edinb. Melissa hortensis C. B. Melyssophyllum, mellisolium, mellitis, citrago, citraria, cedronella, apiastrum. Melissa ossicinalis Linn. Balm: a plant with square stalks; and oblong, pointed, dark green, somewhat hairy leaves, set in pairs; in the bosoms of which come forth pale reddish labiated flowers, standing several together on one pedicle, with the upper lip roundish, erect, and cloven, and the lower divided into three segments. It is perennial; a native of mountainous places in the southern parts of Europe; and slowers in our gardens in June.

This plant, formerly celebrated for cephalic, cordial, stomachic, uterine, and other virtues, is now justly ranked among the milder corroborants. It has a pleasant smell, somewhat of the lemon kind, and a weak aromatic taste; of both which it loses a considerable part on being dried; a slight roughishness, which the fresh herb is accompanied with, becoming at the same time more sensible. Insusions of the leaves in water, in colour greenish or reddish brown according to the degree of saturation, smell agreeably of the herb, but discover no great taste, though, on being inspissated, they leave a considerable quantity of a bitterish and somewhat austere extract: the insusions are sometimes drank as tea in chronical disorders proceeding from debility and relaxation, and sometimes acidulated with lemon juice for a diluent in acute diseases. On distilling the fresh herb with water, it impregnates the first runnings

pretty

pretty strongly with its grateful flavour: when large quantities are subjected to the operation at once, there separates, and rises to the surface of the aqueous sluid, a small portion of essential oil, in colour yellowish, of a very fragrant smell, apparently of great medicinal activity, commended by Hossman as an excellent corroborant of the nervous system. Tinctures of the newly-dried leaves made in rectified spirit, in colour blackish green, discover less of the balm smell than the watery insusions, but have its taste in a greater degree: inspissated, they leave an extract in somewhat less quantity than that obtained by water, in taste stronger, and which retains a considerable share of the specific smell and slavour of the balm, but is less agreeable than the herb in substance.

MENTHA.

MINT: a perennial herb; with square stalks; serrated leaves set in pairs; and spikes of monopetalous slowers, each of which is cut into four sections, and sollowed by sour seeds inclosed in the cup.

I. MENTHA VULGARIS Pharm. Lond. Mentha fativa Pharm. Edinb. Mentha angustifolia spicata C. B. Mentha viridis Linn.*(a) Mint, hartmint, spearmint: with oblong, narrow, pointed leaves, joined close to the stalk; and small purplish flowers standing in long spikes on the tops. It is a native of the warmer climates, common in our gardens, and flowers in June and July.

This herb has a strong agreeable aromatic smell, and a bitterish, roughish, moderately warm taste. It is in general use as a restringent stomachic and carminative: in vomitings and weakness of the stomach, there are, perhaps, sew simples of equal essicacy. Some report that it prevents the coagulation of milk, and hence recommend it to be used along with milk diets, and even in cataplasms and somentations for resolving coagulated milk in the breasts: upon experiment, the curd of

^{*(}a) There is much confusion in the species of mints as described by different botanists. Though the viridis of Linnaus is given as our spearmint, it can scarcely be the same, from Bergius's account of it. The Mentha viridis, he says, is stronger and less agreeable than the splvestris, of a weaker taste than the piperita, and therefore erroneously substituted to it in the shops. Mat. Med. p. 516.

milk, digested in a strong infusion of mint, could not be perceived to be any otherwife affected than by common water, but milk, in which mint leaves were fet to macerate, did not coagulate near so soon as an

equal quantity of the same milk kept by itself.

The leaves are fometimes taken in substance, beaten with thrice their Conserv.fol. weight of fine fugar into a conferve. Moderately bruised, they yield menth.vulg. Pharm.Lond. upon expression about two thirds their weight of a turbid, browncoloured, fomewhat mucilaginous juice; which is commonly supposed to retain the full virtues of the mint, but which, though participating of the bitterness and subastringency of the herb, is found to have little or nothing of the peculiar aromatic flavour in which the principal virtue of this plant refides. The leaves lose in drying about three fourths of their weight, without fuffering much loss of their smell or taste; nor is the fmell foon diffipated by moderate warmth, or impaired in keeping.

Cold water, by maceration for fix or eight hours on the dry herb, and warm water in a shorter time, become richly impregnated with its flavour: if the maceration be long continued, the groffer parts of the mint are extracted, and the liquor proves less grateful: on boiling the mint in water till the aromatic matter is diffipated, the remaining darkbrown liquor is found nearly fimilar to the recent juice; unpleafant, bitterish, subaffringent, and mucilaginous. By distillation, a pound and a half of the dry leaves communicate a strong impregnation to a gallon of water +: the distilled water proves rather more elegant if drawn + Aq. menfrom the fresh plant in the proportion of ten pints from three pounds ! than from the dry plant, though the latter is frequently made use of Lond. as being procurable at all times of the year. Along with the aqueous fluid, an effential oil diftils, of a pale yellowish colour, changing by age to a reddish, and at length to a dark red, in quantity near an ounce from ten pounds of the fresh herb in flower, smelling and tasting strongly of the mint, but fomewhat less agreeable than the herb itself.

Dry mint, digested in rectified spirit, either in the cold, or with a gentle warmth, gives out readily its peculiar tafte and fmell, without imparting the groffer and more ungrateful matter, though the digestion be long continued. The tincture appears by day-light of a fine dark green, by candle-light of a bright red colour: a tincture extracted from the remaining mint by fresh spirit appears in both lights green: the colour of both tinctures changes, in keeping, to a brown. On gently inspissating

thæ vulg. fimpl. Phar. ‡ Aq. menthæ fativæ Phar. Edinb. Ol. menthæ essentiale Pb. Lond. & Edin.

inspissating the filtered tinctures, little or nothing of their flavour rises with the spirit: the remaining extract possesses the concentrated virtues of about ten times its weight of the dry herb; and differs from the products obtained by distillation with water, in this, that the bitterness and subaftringency of the mint, which are there separated from the

aromatic part, are here united with it.

Proof spirit extracts the smell and taste of mint, but not its green colour. The tincture is brown, like the watery infusions; and, like them also, it becomes ungrateful, if the digestion is long continued. On gentle distillation, the more spirituous portion, which rises at first, discovers little flavour of the mint; but as soon as the watery part begins to distil, the virtues of the mint come over plentifully with it. Hence the officinal spirituous water, prepared by drawing off a gallon of proof Aq. menthæ spirit from a pound and a half of the dried leaves, proves strongly im-rituosa P. L. pregnated with the mint.

After mint has been repeatedly infused in water, rectified spirit still extracts from it a green tincture, and a fenfible flavour of mint: on the other hand, fuch as has been first digested in spirit, gives out afterwards to water a brown colour, and a kind of nauseous mucilaginous taste very different from that which distinguishes mint. The spirituous tinctures mingle with watery liquors without precipitation or turbidness; but spirituous liquors, impregnated with its pure volatile parts by distillation, turn milky on the admixture of water.

- 2. MENTHA AQUATICA five Mentastrum Pharm. Paris. aquatica five sifymbrium J. B. Mentha rotundifolia palustris sive aquatica major C. B. Mentha aquatica Linn. Water-mint: with somewhat oval leaves fet on pedicles, and long stamina standing out from the flowers.
- 2. MENTASTRUM HIRSUTUM: Auricularia officinarum Dale: Mentha palustris folio oblongo C. B. Hairy water-mint: with long hairy leaves having no pedicles; and broad spikes of flowers.

BOTH these plants grow wild in moist meadows, marshes, and on the brinks of rivers, and flower towards the end of fummer. They are less agreeable in smell than the spearmint, and in taste bitterer and more pungent: pungent: the second fort approaches in some degree to the slavour of pennyroyal. They yield a much smaller proportion of essential oil: from twenty pounds of the water-mint were obtained scarcely three drams. With regard to their virtues, they appear to partake of those of spearmint; to which they are obviously far inferiour as stomachics. The hairy water-mint is supposed to be the auricularia, planta zeylanica, or earwort, celebrated by Marloe for the cure of deasness: though probably not more effectual against that complaint, than the other water-mint against nephritic ones, in which it is said to have been formerly an empyrical secret (a).

4. Mentha piperitis Pharm. Lond. & Edinb. Mentha spicis brevioribus & babitioribus, foliis menthæ fuscæ, sapore fervido piperis Raii synops. Mentha piperita Linn. Pepper-mint: with acuminated leaves on very short pedicles; and the flowers set in short thick spikes or heads. It is a native of this kingdom; and, so far as is known, of this kingdom only: it is much less common, however, than the other wild mints; but having been of late received in general practice as a medicine, it is now raised plentifully in gardens, and does not appear, like many of the other plants that grow naturally in watery places, to lose any thing of its virtue with this change of soil.

This species has a more penetrating smell than any of the other mints; and a much stronger and warmer taste, pungent and glowing like pepper, and sinking as it were into the tongue. It is a medicine of great importance in statulent colics, hysteric depressions, and other like complaints; exerting its activity as soon as taken into the stomach, and diffusing a glowing warmth through the whole system; yet not liable to heat the constitution near so much as might be expected from the great warmth and pungency of its taste.

By maceration or infusion, it readily and strongly impregnates both water and spirit with its virtue; tinging the former of a brownish colour, and the latter of a deeper green than the other mints. In

⁽a) In the first volume of Linnæus's Amænitates Academicæ, the auricularia is said to be not at all of the mint kind, but a stellated plant, akin to galium.

menthæ pi-peritidis Pb.

Ol.effentiale distillation with water, it yields a considerable quantity of essential oil. of a pale greenish yellow colour, growing darker coloured by age, very Lond. & Edin. light, fubtile, possessing in a high degree the specific smell and penetrating pungency of the pepper-mint *(a): the decoction remaining after the feparation of this active principle, is only bitterish and subastringent, like those of the other mints. Rectified spirit, drawn off with a gentle heat from the tincture made in that menstruum, brings over little of the virtue of the herb, nearly all its pungency and warmth remaining concentrated in the extract, the quantity of which amounts to about Aq. menthæ one fourth of the dried leaves. A simple and a spirituous distilled water, piperit. simp. Ph. L. & Ed. drawn in the same proportions as those of spearmint, and the essential Aq. menthæ oil, are kept in the shops.

piperit. fpirituofa Phar. Lond. & Edin.

MERCURIALIS.

HERB-MERCURY: a plant with oblong, acuminated, indented leaves, standing in pairs: in their bosoms come forth, either spikes of imperfect flowers, fet in three-leaved cups, falling off without any feeds; or little rough balls, joined two together, including each a fingle feed.

1. MERCURIALIS mas & femina Pharm. Edinb. (b) testiculata sive mas, & mercurialis spicata sive femina dioscoridis & plinii C. B. Mercurialis annua Linn. French mercury: with smooth glossy leaves, and branched stalks. The flowering plants, called female, and those which produce feeds, called male, are both annual, and grow wild together in shady uncultivated grounds.

The leaves of this plant have no remarkable fmell, and very little taste: when freed by exsiccation from the aqueous moisture, with which they abound, their prevailing principle appears to be of the mucilaginous kind, with a small admixture of saline matter. They are ranked among the emollient oleraceous herbs, and faid to gently loofen the belly: their principal use has been in glysters.

^{• (}a) Some particles of a true camphor were procured from dried pepper-mint by cohobation. Gaubii Adversar.

2. CYNOCRAMBE: Mercurialis montana testiculata & spicata C. B. Mercurialis perennis Linn. Dogs mercury, male and semale: with rough leaves and unbranched stalks. It is perennial, and grows wild in woods

and hedges.

This species has been said by some to be similar in quality to the foregoing, and to be more acceptable to the palate as an oleraceous herb: it has lately however been found to be possessed of noxious qualities, acting as a virulent narcotic. An instance is related in N°. 203 of the Philosophical Transactions, of its ill effects on a family, who eat at supper the herb boiled and fried: the children, who were most affected by it, vomited, purged, and fell sast assessment two slept about twenty-sour hours, then vomited and purged again, and recovered: the other could not be waked for sour days, and then opened her eyes and expired.

MEUM.

MEUM ATHAMANTICUM Pharm. Lond. Meum foliis anethic. B. Meu & athamanta & radix ursina quibusdam. Æthusa Meum Linn. Spignel, Bauldmony: an umbelliferous plant, with bushy leaves divided into slender segments, like those of fennel, but finer; producing large, oblong, striated seeds, flat on one side, and convex on the other: the root is long, variously branched, with generally a number of hairs or silaments at top, which are the remains of the stalks of former years, of a brownish colour on the outside, pale or whitish within, when dried of a sungous texture. It is perennial, grows wild in meadows in some of the mountainous parts of England, and slowers in June.

The root of spignel, recommended as a carminative, stomachic, and for attenuating viscid humours, appears to be nearly of the same nature with that of lovage; differing, in its smell being rather more agreeable, somewhat like that of parsneps, but stronger, and in its taste being less sweet and more warm or acrid. The difference betwixt the two roots is most considerable in the extracts made from them by water; the extract of spignel root being unpleasantly bitterish, with little or nothing of the sweetness of that of lovage roots. The spirituous extract of spignel, more aromatic than that of the lovage, is moderately warm, bitterish, and pungent.

MILLEFOLIUM.

MILLEFOLIUM Pharm. Edinb. Millefolium vulgare album, & millefolium purpureum C. B. Achillea, Myriophyllon, Chiliophyllon, Militaris berba, Stratiotes, Carpentaria, Lumbus veneris, & Supercilium veneris. Achillea Millefolium Linn. MILFOIL or YARROW: a plant with rough stiff leaves, divided into small segments, set in pairs along a middle rib like seathers: the little flowers stand thick together in form of an umbel on the top of the stiff stalk, and consist each of several whitish or pale purplish petala set round a kind of loose disk of the same colour, sollowed by small crooked seeds. It is perennial, grows plentifully by the sides of fields and on sandy commons, and is found in flower greatest part of the summer.

The leaves and flowers of milfoil are greatly recommended by fome of the German physicians (a) as mild corroborants, vulneraries, and antifpasmodics, in diarrhœas, hemorrhagies, hypochondriacal and other disorders. They promise, by their sensible qualities, to be of no inconsiderable activity. They have an agreeable though weak aromatic smell, and a bitterish, roughish, somewhat pungent taste. The leaves are chiefly directed for medicinal use, as having the greatest bitterishness and austerity: the flowers have the strongest and most subtile smell, are remarkably acrid, and promise to be of most efficacy, if the plant has really any such efficacy, as an anodyne or antispasmodic. Dr. Grew observes, that the young roots have a glowing warm taste, approaching to that of contrayerva, and thinks they might in some measure supply its place; but adds, that they lose much of their virtue in being dried (b), from whence it may be presumed that their active matter is of another kind.

The virtue of the leaves and flowers is extracted both by watery and spirituous menstrua; the astringency most perfectly by the former, their aromatic warmth and pungency by the latter, and both of them equally by a mixture of the two. The flowers, distilled with water, yield a penetrating essential oil, possessing the flavour of the milsoil in perfection,

⁽a) Stahl, Dissert. de therapia passionis hypochondriacé. Hoffman, De prestantia remed. domest. §. 18. (b) Idea of philos. hist. of plants, §. 29. Of diversities of tastes, chap. v. §. 2.

though rather less agreeable than the flowers themselves, in consistence fomewhat thick and tenacious, in colour remarkably variable, fometimes of a greenish yellow, sometimes of a deep green, sometimes of a blueish green, and sometimes of a fine blue: these differences seem to depend in great measure on the foil in which the plant is produced; the flowers gathered from moist rich grounds yielding generally a blue oil; whereas those, which are collected from dry commons, afford only, fo far as I have observed, a green one with a greater or less admixture of yellow: the decoction remaining after the separation of this volatile principle, leaves, on being inspissated, a dark brownish mass, ungratefully austere, bitterish, and somewhat saline. On inspissating the yellowish tincture made in rectified spirit, scarcely any thing of the flavour of milfoil exhales or diffils with the menstruum: the remaining deep yellow extract is more agreeable in fmell than the flowers themfelves, of a moderately warm penetrating tafte, somewhat like that of camphor, but much milder, accompanied with a flight bitterishness and fubastringency.

MILLEPEDÆ.

MILLEPEDÆ Pharm. Lond. & Edinb. Centipedes & onisci Quibusdam. Oniscus Asellus Linn. Millepedes, Wood-Lice: an oblong insect, with sourteen feet, and its body composed of sourteen rings, rolling itself up into a round ball on being touched; sound in cellars, and under stones and logs of wood in cold moist places; rarely met with in the warmer climates. Two sorts are commonly used indiscriminately; one large, of a dusky blueish-black or livid colour; the other smaller, slatter, thinner, of a pale brownish grey, and differing also from the former in the last division of the body being not annular, but pointed, and in the tail being forked. The first species is said to be the true officinal fort, though some have preferred the second: but there does not seem to be any material difference between them in quality.

MILLEPEDES have a faint disagreeable smell, and a somewhat brackish, sweetish, unpleasant taste. They are celebrated as resolvents, aperients, and diuretics; in jaundices, asthmas, scrophulous and other disorders; but that their virtues are so great as they are generally supposed to be,

may be justly questioned, at least when given in the customary doses. I have known two hundred taken every day for some time together, without producing any remarkable effect: in large doses, indeed, it is probable that their activity may be considerable; as they are said to have sometimes produced an universal heat and thirst with a pain in the region of the pubes (a), and sometimes a scalding of urine (b).

Millepedæ præparatæ Ph. L. & Ed.

These insects may be commodiously swallowed entire, as they spontaneously contract themselves, on being touched, into the form of a pill. In the shops they are commonly reduced into powder; for which purpose they are prepared, by inclosing them in a thin canvas cloth, and fuspending them over hot spirit of wine in a close vessel, till they are killed by the steam and rendered friable. Of the extraction of their active matter by menstrua, no direct experiments have been made: it is rather by expression, than on the principle of extraction or dissolution, that their virtues are commonly endeavoured to be obtained in a liquid form; though fome liquors are generally added previously to the expression, partly to improve their virtue for particular intentions, partly to preserve the animal juice from corruption, and partly to render it more completely separable. The college of Edinburgh directs two ounces of live millepedes to be flightly bruifed, and digested for a night in a pint of rhenish wine, after which the liquor is to be pressed out through a strainer.

Vinum millepedatum Pharm. Edin.

MOLDAVICA.

MOLDAVICA seu Melissa turcica: an Melissa americana trifolia odore gravi Tourn. inst.? Camphorosma Morison. hist. ox.? Dracoce-phalum canariense Linn. Turkey or rather Canary Balm, commonly called Balm-of-gilead: a plant with square stalks, and acuminated leaves, slightly and obtusely indented, set generally three on one pedicle: of each three, the end one is largest, and the other two are nipt at the bottom on the upper side, or do not reach so far down their middle ribs on that side as on the other: the pedicles stand in pairs at the joints, with similar sets of smaller leaves in their bosoms.

⁽a) Frid. Hoffman, De mat. med. regn. animal. cap. 18. Opera omnia, suppl. ii. par. iii. p. 157. (b) Fuller, Pharmacopæia extemporanea, sub Express. milleped. simp.

On the tops come forth thick spikes, or heads, of pretty large, reddish, labiated flowers; whereof both the upper and lower lip are cut into two parts, and the cup into five. It is perennial, a native of the Canary islands, and scarcely bears the winters of our climate without shelter.

THIS or some of the other species of the Turkey balm, of which there are several, is greatly commended by Hoffman, for strengthening the tone of the stomach, and the nervous system: in this country, it has not yet been, though it feems to have a good claim to be, received among the medicinal plants: infusions of it may be drank as tea, and are very grateful. The leaves and flowery tops have a fragrant fmell, fomewhat refembling that of balm, but far stronger, and approaching to that of the fine balfam from which the plant received its name. Their tafte is likewise agreeable, but so covered with the aromatic flavour, that its particular species is not easily determined: when the herb is infused in water, and the aromatic part distipated by inspissating the filtered infusion, the remaining extract impresses on the palate a moderately strong, though only momentary, pungency and bitterness. In Ol. syrize distillation with water, it yields a fragrant essential oil.

Germanis quibusdam.

MOSCHUS.

MOSCHUS Pharm. Lond. & Edinb. Musk: an odoriferous, grumous substance: found in a little bag, situated near the umbilical region of an oriental quadruped, which is faid by some to bear the greatest resemblance to the goat, by others to the stag kind. The best musk is brought from Tonquin in China, an inferiour fort from Agria and Bengal, and a still worse from Russia.

Fine musk comes over in round thin bladders, generally about the fize of pigeons eggs, covered with short brown hairs, well filled, and without any aperture or any appearance of their having been opened. The musk itself is dry, with a kind of unctuosity; of a dark reddish brown or rufty blackish colour; in small round grains, with very few hard black clots; perfectly free from any fandy or other visible foreign matter. Chewed, and rubbed with a knife on paper, it looks bright, yellowish, smooth and free from grittiness. Laid on a red-hot iron, it catches flame, and burns almost intirely away, leaving only an exceeding small quantity of light greyish ashes: if any earthy substances have been mixed with the musk, the quantity of the residuum will discover them.

THIS concrete has a bitterish subacrid taste; and a fragrant smell, agreeable at a distance, but so strong as to be disagreeable when smelt near to, unless weakened by a large admixture of other substances. A fmall quantity, macerated for a few days in rectified spirit of wine, imparts a deep colour, and a strong impregnation to the spirit: this tincture, of itself, discovers but little smell, the spirit covering or suppressing the smell; but on dilution it manifests the full fragrance of the musk, a drop or two communicating to a quart of wine or watery liquors a rich musky scent. The quantity of liquor which may thus be flavoured by a certain known proportion of musk, appears to be the best criterion of the genuineness and goodness of this commodity; a commodity, which is not only faid to vary in goodness according to the feason of its being taken from the animal (a), but which is oftentimes fo artfully fophisticated, that the abuses cannot be discovered by any external characters, or by any other known means than the degree of its specific smell and taste, which the above experiment affords the most commodious method of measuring. The rectified spirit takes up completely the active matter of the musk; watery liquors extract it only in part. The shops endeavour to procure an union of its virtues with water by the intervention of fugar: twelve grains of musk and a dram of fine fugar are thoroughly ground together, and fix ounces by meafure of damask rose water added by degrees: this intermedium, however, does not answer so effectually as could be wished; the mixture, at first turbid, becoming clear on standing for a little time, and depositing, along with the feculent matter, a great share of the medicinal parts of the musk: the most successful medium for uniting musk with water, appears to be mucilage of gum-arabic; but the most elegant of all the liquid preparations of this drug, is the tincture in rectified spirit, which may be occasionally diluted with any watery liquors, like the other spirituous tinctures. By distillation, water becomes strongly impregnated with the

Julepum e moscho P. L.

(a) Strahlenberg, Descript. Ruff. Siber. &c. p. 340.

fcent of the musk, and seems to elevate all its odoriferous matter; while rectified spirit, on the contrary, brings over little or nothing of it.

Musk, a medicine of great esteem in the eastern countries, has lately come into general use among us also, in some nervous disorders: though liable, by its strong impression on the organs of smell, to offend and disorder hysterical persons and constitutions of great sensibility, yet, when taken internally, it is found to abate fymptoms of that kind which its smell produces, and to be one of the principal medicines of the antifpafmodic class. Dr. Wall relates, that two persons labouring under a fubfultus tendinum, extreme anxiety, and want of fleep, occasioned by the bite of a mad dog, were perfectly relieved by two doses of musk of fixteen grains each: that convulfive hiccups, attended with the worst fymptoms, were removed by a dose or two of ten grains: but in some cases, where this medicine could not, on account of strong convulsions, be administered by the mouth, it proved of service when injected as a glyfter: that he never met with any person, how averse soever to perfumes, but could take it in the form of a bolus without inconvenience: that under the quantity of fix grains, he never found much effect from it, but that when given to ten grains and upwards, it produces a mild diaphoresis, without heating or giving any uneasiness, but on the contrary, abating pain and raifing the spirits; and that after the sweat has begun, a refreshing sleep generally succeeds(a). This medicine is now received in general practice, in different convultive diforders; and its dose has been increased, with advantage, to a scruple, and half a dram, every four or fix hours. It has been tried also in some maniacal cases; in which it seemed to procure a temporary relief.

M O X A.

MOXA five lanugo artemifiæ japonicæ Pharm. Parif. Moxa: a foft lanuginous substance, prepared in Japan, from the young leaves of a species of mugwort, by beating them, when thoroughly dried, and rubbing them betwixt the hands, till only the fine fibres are left. A like substance is said, in the German ephemerides, to have been obtained, by treating the leaves of our common mugwort in the same manner.

(a) Philosoph. Transact. No. 474-

Moxa is celebrated in the eastern countries, for preventing and curing many disorders, by being burnt on the skin: a little cone of moxa, laid on the part previously moistened, and set on fire at top, burns down with a temperate glowing heat, and produces a dark coloured spot, the exulceration of which is promoted by applying a little garlic, and the ulcer either healed up when the eschar separates, or kept running for a length of time, as different circumstances may require. A sungous substance, found in fissures of old birch trees, is said to be in common use among the Laplanders for the same purposes (a); and some have used cotton, impregnated with a solution of nitre, and afterwards dried, which answers the end as effectually as the moxa of the Japonese (b). It is obvious, that all these applications are no other than means of producing an exulceration of the skin, and its consequence a drain of humours.

MYROBALANI.

MYROBALANI Pharm. Parif. Myrobalans: dried fruits, of the plum kind, brought from the East Indies. Five forts, produced by different trees, have been distinguished in the shops.

- 1. Myrobalani Belliricæ: Myrobalani rotundæ belliricæ, arabibus belleregi, &c. C. B. Belliric myrobalans: of a yellowish grey colour, and an irregularly roundish or oblong figure, about an inch in length, and three quarters of an inch thick.
- 2. Myrobalani citrinæ: Myrobalani teretes citrini bilem purgantes C. B. Yellow myrobalans: somewhat longer than the preceding; with generally five large longitudinal ridges, and as many smaller between them; somewhat pointed at both ends.
- 3. Myrobalani chebulæ: Myrobalani maximi angulosi pituitam purgantes, arabibus quebolia, &c. C. B. Chebule myrobalans: refembling the yellow in figure and ridges, but larger, of a darker colour inclining to brown or blackish, and with a thicker pulp.

- 4. Myrobalani emblicæ in segmentis nucleum habentes angulosæ J. B. Emblic myrobalani: of a dark blackish grey colour, roundish, about half an inch thick, with six hexagonal faces opening from one another; the fruit of the Phyllanthus Emblica of Linnæus.
- 5. Myrobalani indicæ nigræ sine nucleis J. B. Indian or black myrobalans: of a deep black colour, oblong, octangular, differing from all the others, in having no stone, or only the rudiments of one; from whence they are supposed to have been gathered before maturity.

ALL the myrobalans have an unpleasant, bitterish, very austere taste; and strike an inky blackness with folution of chalybeate vitriol. They are faid to have a gently purgative, as well as an aftringent and corroborating virtue; and are directed to be given, in fubstance from half a dram to four drams, and in infusion or slight decoction from four to twelve drams. It is faid also, that the fruit in substance acts barely as a flyptic, without exerting its purgative quality; that this last is difcovered only in the infusions (a), and that by boiling it is diffipated or destroyed (b). A difference of this kind, between the fruit and its infusions, might be easily conceived, if the astringency of the myrobalans was not extracted by watery liquors, but the contrary of this was found on trial to be true; the infusions, decoctions, and the decoctions inspissated to the consistence of an extract, being strongly styptic. In this country, they have long been entire strangers to practice, and are now discarded, by the colleges both of London and of Edinburgh, from their catalogue of officinals.

MYRRHA.

MYRRHA Pharm. Lond. & Edinb. MYRRH: a gummy-refinous concrete juice, of an oriental tree, of which we have no certain account. * Mr. Bruce informs us that it grows spontaneously in the eastern part of Arabia felix, and in that part of Abyssinia which the Greeks named

⁽a) Geoffroy, Tract. de materia medica, tom. ii. p. 332.

⁽b) Benancius, Declaratio fraudum & errorum apud pharmacopæos, è museo Bartholini, p. 68.

Troglodytria; and that the Abyssinian myrrh, which is the least plentiful, is the best. The best kind is that which slows from deep incisions of the larger branches, hardening upon the tree. It continues to distil every year from the same wound; but the myrrh is of an inferiour quality after the first, being mixed with foreign impurities, and the decayed juices of the tree. The worst is that which comes from near the root, or the old trunks (a). It comes over in glebes or drops, of various colours and magnitudes: the best sort is somewhat transparent, friable, in some degree unctuous to the touch, of an uniform brownish or reddish yellow colour, often streaked internally with whitish semi-circular or irregular veins; of a moderately strong, not disagreeable smell; and a lightly pungent, very bitter taste, accompanied with an aromatic slavour, but not sufficient to prevent its being nauseous to the palate.

There are sometimes found among it hard shining pieces, of a pale yellowish colour, resembling gum-arabic, of no taste or smell: sometimes masses of bdellium, darker coloured, more opake, internally softer than the myrrh, and differing from it both in smell and taste: sometimes an unctuous gummy-resin, of a moderately strong somewhat ungrateful smell, and a bitterish very durable taste, obviously different both from those of bdellium and myrrh: sometimes likewise, as Cartheuser observes, hard compact dark coloured tears, less unctuous than myrrh, of an offensive smell, and a most ungrateful bitterness, so as, when kept for some time in the mouth, to provoke reaching, though so resinous, that little of them is dissolved by the saliva. Great care is therefore requisite in the choice of this drug.

This bitter aromatic gummy-refin is a warm corroborant, deobstruent, and antiseptic. It is given from a few grains to a scruple and upwards, in uterine obstructions, cachexies, putrid severs, &c. and often employed also as an external antiseptic and vulnerary. *In doses of half a dram, Dr. Cullen remarks that it heated the stomach, produced sweat, and agreed with the balsams in affecting the urinary passages. It has lately come more into use as a tonic in hectical cases, and is said to prove less heating than most other medicines of that class.

Myrrh disfolves almost totally in boiling water, but as the liquor cools, a portion of refinous matter subsides. The strained solution is of a dark vellowish colour, somewhat turbid, smells and tastes strongly of the myrrh, and retains both its tafte, and a confiderable share of its scent, on being inspiffated with a gentle heat to the consistence of an extract. By distillation with a boiling heat, the whole of its flavour arises, partly impregnating the diffilled water, partly collected and concentrated in the form of an effential oil; which is in fmell extremely fragrant, and rather more agreeable than the myrrh in substance, in taste remarkably mild, fo ponderous as to fink in the aqueous fluid, whereas the oils of most, perhaps of all, of the other gummy-refins swim: the quantity of oil, according to Hoffman's experiments, is about two drams from fixteen ounces, and when the myrrh is of a very good kind, near three drams.

Rectified spirit dissolves less of this concrete than water, but extracts more perfectly that part in which its bitterness, flavour, and virtues, refide: the refinous matter, which water leaves undiffolved, is very bitter; but the gummy matter, which spirit leaves undissolved, is insipid, the spirituous folution containing all the active parts of the myrrh. Tinctures of myrrh, made by digefting three ounces of the concrete in Tinc. myrr. two pounds and a half of rectified + or a quart of proof 1 spirit, are kept + Pb. Edinb. in the shops, and given sometimes internally from fifteen drops to a teaspoonful, but oftener used among us externally for cleanfing ulcers and promoting the exfoliation of carious bones: both tinctures are of a reddish yellow colour. In distillation, rectified spirit brings over little or nothing of the flavour of the myrrh: the extract, obtained by inspissating the tincture, is a fragrant, bitter, very tenacious resin, amounting to one third or more of the weight of the myrrh employed.

MYRTUS.

MYRTUS Pharm. Edinb.(a) Myrtus communis italica Cafp. Baubin. Myrtus communis Linn. MYRTLE: an evergreen shrub; with oblong leaves, pointed at both ends; in the bosoms of which spring solitary white pentapetalous flowers, followed by black oblong umbilicated

(a) Expunged.

berries

berries full of white crooked feeds. It is a native of the fouthern parts of Europe, from whence the shops have been usually supplied with the berries, called *myrtilli*, which rarely come to perfection in our climate; nor does the shrub bear our severe winters without shelter.

THE berries of the myrtle, recommended in alvine and uterine fluxes and other diforders from relaxation and debility, appear to be among the milder reftringents or corroborants: they have a roughish not unpleasant taste, accompanied with a degree of sweetishness and aromatic slavour. The leaves have likewise a manifest astringency, and yield, when rubbed, a pretty strong aromatic smell, agreeable to most people.

MYRTUS BRABANTICA.

MYRTHUS BRABANTICA Pharm. Parif. Rhus myrtifolia belgica C. B. Gale, frutex odoratus feptentrionalium, elæagnus Cordo, chamælæagnus Dødonæo J. B. Myrica Gale Linn. Gaule, Sweet willow, Dutch myrtle: a small shrub, much branched; with oblong, smooth, whitish green leaves, somewhat pointed or converging at each end; among which arise pedicles bearing flowery tusts, and separate pedicles bearing scaly cones which include the seeds, one little seed being lodged in each scale. It grows wild in waste watery places in several parts of England: in the isle of Ely it is said to be very plentiful. It slowers in May or June, ripens its seeds in August, and loses its leaves in winter.

THE leaves, flowers, and feeds of this plant, have a strong fragrant smell, and a bitter taste. They are said to be used among the common people, for destroying moths, and cutaneous infects, being accounted an enemy to insects of every kind; internally, in insusions, as a stomachic and vermisuge; and, as a substitute to hops, for preserving malt liquors, which they render more inebriating, and of consequence less salubrious (a): it is said that this quality is destroyed by boiling (b).

(a) Ray, Historia plantarum, tom. ii. p. 1707. (b) Linnai, Aman. Academic. iii. 96.

· NAPUS.

NAVEW: a plant of the turnep kind, with oblong roots growing slenderer from the top to the extremity. Two forts of it, ranked among the articles of the materia medica, are supposed by Linnæus to be only varieties, and are therefore joined into one species, under the name of brassica (napus) radice caulescente suffermi. They are both biennial.

- I. NAPUS; napus dulcis officinarum Pharm. Lond. Napus sativa C. B. Garden or fweet navew, or French turnep: cultivated for the culinary use of its roots, which are warmer and more grateful than those of the common turnep, and are faid to afford likewife, in their decoctions, a liquor beneficial in diforders of the breaft. The feeds, in figure roundish and in colour reddish, are the part principally directed for medicinal purpofes: they have a moderately pungent tafte, fomewhat approaching to that of mustard seed, of the virtues of which they appear to partake: with mustard feed they agree also in their pharmaceutic properties, their pungent matter being taken up completely by water, and only partially by rectified spirit, and being diffipated in the inspissation of the watery infusion, only an unpleasant bitterishness remaining in the extract. As the navew feeds, nearly fimilar in kind to those of mustard, are apparently much inferiour in degree, the college of Edinburgh has difcarded them, and that of London retains them only as an ingredient in theriaca.
- 2. Bunias Pharm. Parif. Napus silvestris C. B. Wild navew, or rape: growing on dry banks and among corn: with leaves somewhat different from those of the preceding, being more like those of cabbages than of turneps; the root smaller, and of a stronger unpleasant taste; and the seeds also rather more pungent, on which account they are preferred by the faculty of Paris. The seeds of both kinds yield upon expression a large quantity of oil: the oil called rape-oil is extracted from the seeds of the wild fort, which is cultivated in abundance, for that use, in some parts of England; the cake, remaining after the expression of the oil, retains, like that of mustard, the acrimony of the seeds.

NARDUS CELTICA.

NARDUS CELTICA Pharm. Lond. Nardus celtica dioscoridis C. B. Valeriana celtica Tourn. & Linn. Spica celtica & saliunca quibusdam. Celtic nard: a small species of valerian, with uncut, oblong, obtuse, somewhat oval leaves. It is a native of the Alps, from whence the shops have been generally supplied with the dried roots, consisting of a number of blackish sibres, with the lower parts of the stalks adhering; which last are covered with thin yellow scales, the remains of the withered leaves.

This root has been recommended as a stomachic, carminative, and diuretic: at present, it is scarcely otherwise made use of, in this country, than as an ingredient in mithridate and theriaca, though its sensible qualities promise some considerable medicinal powers. It has a moderately strong smell, of which it is extremely retentive (a), and a warm bitterish subacrid taste, somewhat resembling those of common wild valerian: an extract made from it by rectified spirit has a strong penetrating taste, and retains in good measure the particular slavour, as well as the bitterness and pungency of the root.

NARDUS INDICA.

NARDUS INDICA Pharm. Lond. Nardus indica quæ spica, spica nardi, & spica indica officinarum C. B. Indian nard, or spikenard: the bushy top of the root, or the remains of the withered stalks and ribs of the leaves, of an Indian grassy-leaved plant of which we have no particular description. The nard, as brought to us, is a congeries of small tough reddish brown fibres; cohering close together, but not interwoven, so as to form a bunch or spike about the size of a singer: sometimes two or three bunches issue from one head, and sometimes bits of leaves and stalks in substance are found among them.

THE Indian nard, now kept in the shops chiefly as an ingredient in the mithridate and theriaca, was formerly employed in the same intentions as

⁽a) Linnaus observes that this plant, in a dry herbal, has retained its fragrance above a century. Amanitat. Academic. vol. iii. p. 71.

the Celtic, and is faid to be used among the orientals as a spice. It is moderately warm and pungent, accompanied with a flavour not disagreeable.

NASTURTIUM AQUATICUM.

NASTURTIUM AQUATICUM Ph. Lond. & Edinb. Nasturtium aquaticum supinum C. B. Sisymbrium aquaticum Tourn. Cressio quibusdam. Sisymbrium Nasturtium aquaticum Linn. Water cresses: a juicy plant, with brownish, oblong, obtuse leaves, set nearly in pairs, without pedicles, on a middle rib, which is terminated by an odd one larger and longer-pointed than the rest: the stalks are hollow, pretty thick, channelled, and crooked: on the tops grow tusts of small tetrapetalous white slowers, followed by oblong pods, which bursting throw out a number of roundish seeds. It grows in rivulets and the clearer standing waters, and slowers in June: the leaves remain green all the winter, but are in greatest perfection in the spring.

The leaves of the water cresses have a moderately pungent taste; and, when rubbed betwixt the fingers, emit a quick penetrating smell, like that of mustard seed, but much weaker. Their pungent matter is taken up both by watery and spirituous menstrua, and accompanies the aqueous juice which issues copiously upon expression: it is very volatile, so as to arise, in great part, in distillation with rectified spirit as well as with water, and almost totally to exhale in drying the leaves, or inspissating by the gentless heat, to the consistence of an extract, either the expressed juice, or the watery, or spirituous tinctures: both the inspissated juice and the watery extract discover to the taste a faline impregnation, and in keeping throw up crystalline efflorescences to the surface. On distilling with water considerable quantities of the herb, a small proportion of a subtile, volatile, very pungent essential oil is obtained.

This herb is one of the milder acrid, aperient, antifcorbutics; of the fame general virtues with the cochlearia, but confiderably less pungent, and in great measure free from the peculiar flavour which accompanies that plant. Hoffman has a great opinion of it, and recommends it as of singular efficacy for strengthening the viscera, opening obstructions of the glands, promoting the fluid secretions, and purifying the blood

3 N

MATERIA MEDICA.

and humours: for these purposes, the herb may be used as a dietetic article, or the expressed juice taken in doses of from one to sour ounces twice or thrice a day.

NASTURTIUM HORTENSE.

NASTURTIUM HORTENSE Pharm. Edinb. (a) Nasturtium bortense vulgatum C.B. Lepidium sativum Linn. GARDEN CRESSES: a low plant, with variously cut winged leaves, bearing on the top of the round stalk and branches tusts of tetrapetalous white slowers, which are followed by roundish capsules, slatted on one side, full of reddish round seeds. It is annual, and raised in gardens.

THE garden cress is an useful dietetic herb in scorbutic habits, viscidities of the juices, obstructions of the viscera, and for promoting digestion; nearly of the same quality with water cress, but somewhat milder. The seeds are considerably more pungent than the leaves, and agree in their general qualities with those of mustard.

NATRON.

NATRON, Anatron, Soude blanche, Pharm. Parif. Nitrum antiquorum. Aphronitrum. Baurach. NATRON, or MINERAL FIXT ALKALINE SALT. This falt is contained in great abundance in the waters of the ocean, and makes the basis of the neutral salt so plentifully extracted from them for alimentary uses. It is likewise discoverable in sundry mineral springs, even of those which do not participate of sea salt. The celebrated Seltzer waters, in the archbishoprick of Treves, appear to be no other than a dilute solution of this salt mixed with a little earthy matter: twelve ounces of the water, according to Hossman's analysis, yield a scruple of the pure alkali. In some of the eastern countries it is sound in considerable quantities on the surface of the earth, sometimes pure, but more commonly blended with various heterogeneous matters, from which it is extracted by means of water. I have been favoured by Dr. Heberden with a sample of this salt in a very pure state, which was

taken up on the Pic of Teneriffe, and with which some parts of that mountain are covered. An account of this salt, as sound sossil in a crystalline state, in the country of Tripoli, is contained in the Phil. Trans. Vol. LXI. Part II. The alkali called soda, or barilla, prepared by incinerating the maritime plant kali or glasswort, contains a salt of the same kind.

The mineral alkali agrees in its general qualities with the common lixivial falts of vegetables. The differences which have been observed are, that it is milder and less acrid in taste: that it melts easier in the fire, and requires more water for its solution: that when dissolved in water it concretes, on evaporation, into crystalline masses: that when exposed to a moist air, though it grows somewhat moist on the surface, it does not run into a liquid form: that in a dry air, the crystals lose the water necessary for their crystalline form, and fall by degrees into a white powder: that the neutral salt, resulting from its coalition with the vitriolic acid, sal glauberi, is very easily dissoluble in water and susible in the fire: that with the nitrous acid it forms cubical crystals, nitrum cubicum; with the marine, perfect sea salt; with tartar, a salt which easily crystallizes, sal rupellense. Made caustic by lime, it proves greatly inferiour to the vegetable alkali in dissolving the urinary calculus. (a).

This falt appears to possess the same general virtues with the vegetable alkalies; but as it does not liquefy in the air, it is better adapted for an ingredient in powders; and as it is less acrimonious, it may be presumed to be less disposed to stimulate the first passages: some of the chemists have taken great pains, in the preparation of the common alkalies, to preserve in them a part of the oil of the plant, so as to reduce them to such a degree of mildness, as this alkali, with much greater uniformity and certainty, possesses in its pure state.

NEPETA.

NEPETA Pharm. Lond. Mentha cataria vulgaris, & major C. B. Cataria & Herba felis quibusdam. Nepeta Cataria Linn. NEP, or CATMINT, so called from its being often destroyed by cats: a hoary

(a) Med. Tranf. i. 124.

plant; with fquare stalks; heart-shaped, acuminated, serrated leaves, set in pairs on long pedicles; and whitish labiated flowers standing in spikes on the tops of the branches: the upper lip of the flower is divided into two, the lower into three sections. It is sometimes found wild in hedges and on dry banks, and flowers in June.

The leaves of catmint have a moderately pungent aromatic taste, and a strong smell, not ill resembling that of a mixture of spearmint and pennyroyal; of the virtues of which herbs, in weaknesses of the stomach, and more particularly in uterine disorders, they appear also to participate. Their active matter is extracted both by water and rectified spirit, most perfectly by the latter: the watery tinctures are of a greenish yellow or brownish colour, the spirituous of a deep green. In distillation with water, they yield a yellowish essential oil, smelling strongly of the catmint, but rather less agreeable than the herb itself: the remaining decoction is ungratefully bitterish and subastringent. Rectified spirit elevates likewise a part of the smell and aromatic warmth, but leaves the greatest share behind concentrated in the extract, which proves more grateful than the leaves in substance, having more of the mint and less of the pennyroyal slavour.

NEPHRITICUM LIGNUM.

NEPHRITICUM LIGNUM Pharm. Edinb. (a) Lignum peregrinum aquam cæruleam reddens C. B. NEPHRITIC wood: an Ameriean wood, brought to us in large compact ponderous pieces, without
knots: the outer part is of a whitish or pale yellowish colour, the medullary substance of a dark brownish or reddish. It is the product of
the Guilandina Moringa of Linnæus. This wood, macerated in water for
half an hour or an hour, imparts a deep tincture, appearing, when placed
betwixt the eye and the light, of a golden colour, in other situations
of a fine blue: a property in which it agrees with the bark of the ash
tree, and differs from all other known woods. Pieces of a different kind
of wood, are often mixed with it, which give only a yellow tincture
to water.

(a) Expunged.

Neptrettic wood has a flightly bitterish somewhat pungent taste; and in rasping or scraping emits a faint smell of the aromatic kind. The blue watery tincture has neither smell nor taste: but a strong insusion, which appears not blue, but of a dark brownish colour, is manifestly bitter, and smells pretty agreeably; inspissated, it leaves a blackish brown extract, in which the bitterness is more considerable, and accompanied with a slight astringency. A saturated tincture made in rectified spirit, is of a blackish red colour: the extract, obtained by inspissating it, is a tenacious resin, larger in quantity and weaker in taste than the watery extract. According to Cartheuser, the spirituous extract amounts to about one sifth, the watery only to one twelfth the weight of the wood. Both menstrua seem to extract the whole of the active matter; for if the wood remaining after the action of the one, be digested or boiled in the other, and the liquors inspissated, the extracts thus obtained have neither smell nor taste.

This wood stands greatly recommended in difficulties of urine, nephritic complaints, and all disorders of the kidneys and urinary passages; and is said to have this peculiar advantage, that it does not, like the warmer diuretics, heat or offend the parts: the blue aqueous tincture is directed to be used as common drink, and fresh water to be poured on the remaining wood so long as it communicates any blueness. For my own part, I have never known its being given medicinally, nor is it received in practice: Geosfroy says he has seen some instances of its being used without success; and indeed, whatever may be the virtues of strong insusions or extracts of the wood, the exceedingly dilute blue tincture cannot be expected to have much efficacy.

NICOTIANA.

TOBACCO: a plant with alternate leaves, and monopetalous tubulous flowers divided into five fections: the flower is followed by an oval capfule, which opening longitudinally, sheds numerous small seeds.

1. NICOTIANA Pharm. Lond. Nicotiana major latifolia C. B. Nicotiana Tabacum Linn. Tobacco: with large, sharp-pointed, pale green, soft leaves, about two feet in length, joined immediately to the stalk without pedicles. It was brought into Europe by M. Nicot, from

from the island Tobago in America, about the year 1560, and is now cultivated for medicinal use in our gardens. It is perennial, as is said, in America; and annual with us.

THE leaves of tobacco have a strong disagreeable smell, and a very acrid burning taste. They give out their acrid matter both to water and spirit, most perfectly to the latter: the aqueous insusions are of a yellow or brown colour, the spirituous of a deep green. They yield nothing considerable in distillation with either menstruum: nevertheless their acrimony is greatly abated in the inspissation of the tinctures, the watery extract being less pungent than the leaves themselves, and the spirituous not much more so. The several sorts of tobacco brought from abroad, are stronger in taste than that of our own growth, and the extracts made from them much more fiery, but in less quantity.

Tobacco taken internally, even in a small dose, or decoctions of it used as a glyster, prove virulently cathartic and emetic, occasioning extreme anxiety, vertigoes, stupors, and disorders of the senses: some have nevertheless ventured upon it both as an evacuant, and in minuter quantities as an aperient and alterant, in epilepsies and other obstinate chronical disorders; a practice which, though in some cases it may have been successful, appears much too hazardous to be followed, particularly in the more irritable, hot, dry, bilious constitutions. By long boiling in water, its deleterious power is abated, and at length destroyed: an extract made by long coction is recommended by Stahl and other German physicians, as the most effectual and safe aperient, detergent, expectorant, diuretic, &c. but the medicine must necessarily be precarious and uncertain in strength, and has never come into use among us.

The smoke of tobacco, received by the anus, is said to be of singular efficacy in obstinate constipations of the belly. Hossman observes, that horses have often been relieved by this remedy, but in human subjects it has been rarely tried; and says he has known some of the common people, who laboured under excruciating pains of the intestines, freed in an instant from all pain by swallowing the smoke. *Both the decoction and the smoke have not unfrequently been injected in cases of incarcerated herniæ, and often with success. The smoke thus applied is recommended as one of the principal means for the revival of persons

apparently

apparently dead from drowning or other fudden causes; but some suspect the narcotic powers of tobacco, as unfavourable in these cases.

Tobacco is fometimes employed externally in unguents and lotions, for cleanfing foul ulcers, destroying cutaneous infects, and other like purposes: it appears to be destructive to almost all kinds of infects, to those produced on vegetables as well as on animals. Beaten into a mash with vinegar or brandy, it has sometimes proved a serviceable application for hard tumours of the hypochondres (a). Some caution however is requisite even in these external uses of tobacco, particularly in solutions of continuity: there are instances of its being thus transmitted into the blood, so as to produce virulent effects. Of the common uses of the leaves brought from America, prepared in different forms, both the advantages and inconveniences are too well known to require being mentioned here.

2. NICOTIANA MINOR C. B. Priapeia quibusdam nicotiana minor J. B. Tobacco anglicum Park. Hyoscyamus luteus Ger. Nicotiana rustica Linn. English tobacco: with short, somewhat oval leaves, set on pedicles. It is annual, originally a native of America, but now propagates itself plentifully in England and other parts of Europe.

THE leaves of this species are said by some to be of the same quality with those of henbane; by others, to be similar to the preceding, but weaker, which, in point of taste, they manifestly are. They have been sometimes substituted, in our markets, to the true tobacco; from which they are readily distinguishable by their smallness, their oval shape, and their being furnished with pedicles.

NITRUM.

NITRUM Pharm. Lond. & Edinb. NITRE, or SALTPETRE: a neutral falt, formed by the coalition of the common vegetable fixt alkaline falt with a peculiar acid: of a sharp penetrating cooling taste: soluble in eight times its weight of very cold water, in less than thrice its weight of water temperately warm, and, as is said, in one third its weight of

boiling water: concreting from its faturated folutions, on evaporation of a part of the fluid or a gradual diminution of the heat that kept it diffolved, into colourless transparent crystals, which in figure are hexagonal prisms terminated by pyramids of the same number of sides: melting thin as water in a moderate heat: when heated to ignition, deflagrating, on the contact of any inflammable substance, with a bright flame and a considerable hissing noise; and leaving, after the detonation, its fixed alkaline salt, the acid being destroyed in the act of accension.

Nitrum fixum

The origin of nitre, or rather of the acid which makes the characteristic part of nitre, is unknown. Thus much only is known with certainty, that common waters, both atmospherical and subterraneous, often contain a little of this acid in combination with earthy or other bodies, so as to yield, by crystallization, on supplying the vegetable fixt alkali, a persect nitre: and that when animal and vegetable substances, mixed with porous absorbent earths, have lain exposed to the air till they are thoroughly rotted, they are found in like manner to contain a small portion of nitre or of nitrous acid, so as to give out a little nitre to water, either without addition, or on being supplied with the proper alkaline basis. On this foundation, some nitre is prepared in different parts of Europe: but the greatest quantities are the produce of the East Indies; the means by which it is there so plentifully obtained, or whether it is a natural or artificial production, have not yet, so far as I can learn, been revealed.

Nitrum purificatum Pharm. Lond. Nitre, as brought into the shops, has generally a greater or less admixture of sea salt; from which it is purified, by dissolving it in boiling water, and, after duly evaporating the filtered solution, setting it in a cold place to crystallize. The more impure brown nitre requires repeated dissolved in lime-water, or the solution suffered to percolate through quicklime or a mixture of quicklime and wood ashes. It is observable that nitrous solutions differ from those of most other salts in contracting no pellicle in evaporation: if a solution of rough nitre, containing sea salt, be boiled down till a pellicle appears, or till a part of the salt begins to concrete and salt to the bottom, all that thus separates is said to be sea salt, boiling water keeping far less of this salt dissolved than it does of nitre: but if the liquor be now poured off, though it should still retain a quantity of the sea salt, only the nitre will crystallize in cooling,

fea

fea falt continuing diffolved in nearly as little water when cold, as was . fufficient to keep it diffolved when boiling.

This falt is one of the principal medicines of the antiphlogistic class; of general use in disorders accompanied with inflammatory symptoms whether chronical or acute, and as a corrector of the inflammation or irritation produced by stimulating drugs. Hossman thinks it has an advantage above the refrigerants of the acid kind, in not being liable to coagulate the animal juices; solutions of it mingling with or dissolving recent thick blood, and in some degree preserving it from coagulation as well as corruption; at the same time changing its colour, when dark or blackish, to a crimson, an effect which it produces also, in a less degree, upon the sleshy parts of dead animals (a). It retards likewise the coagulation of milk, but seems, from Stahl's account, to increase the consistence of thin serous humours; for he observes, that when used in gargarisms for inflammations of the sauces in acute severs, it thickens the salival fluid into a mucus, which keeps the parts moist for a considerable time, whereas, when nitre is not added, adryness of the mouth presently ensues (b).

This medicine generally promotes urine, and often gives relief in stranguries and heat of urine whether simple or proceeding from a venereal taint. It sometimes loosens the belly, particularly in hot dispositions: in cold phlegmatic temperaments it rarely has this effect, though given in very large dose: the diarrheas of acute diseases, and fluxes in other circumstances from an acrimony of the bile or inflammation of the intestines, have been frequently restrained by it. In high fevers, it often promotes a diaphoresis or sweat; in malignant fevers, where the pulse is low and the strength greatly depressed, it impedes that salutary excretion and the eruption of the exanthemata; in consequence of its general power of diminishing inflammation and heat. It seems to be prejudicial in disorders of the lungs, though some (c) have ventured to prescribe it in hamoptyses.

The

⁽a) Hoffman, De falium mediorum virtute, §. 16. De medicamentis felectioribus, §. 13. De præftantissima nitri virtute, §. 5. — We cannot, however, conclude much, from these kinds of experiments, in regard to the medical powers of nitre, or its effects on the animal sluids whilst under
the laws of the vital occonomy.

⁽b) De usu nitri medico, Mensis martius, Opusc. p. 569.

⁽c) Riverius, Cent. i. obs. 83. Stahl, ubi supra, & Observ. chym. phys. med. curios. p. 464. Tralles, Virium terreis ascriptorum examen, p. 246. Dickson, Lond. med. obs. iv. 106. This last writer ventures to assert, that he can depend upon an electuary of conserve of roses and nitre in the cure of an hæmoptoë almost equally with bark in an intermittent.

The usual dose of nitre, among us, is from two or three grains to a scruple; though in many cases it may be given with great safety, and to better advantage, in larger quantities. It has been faid, that nitre loses, in being melted, half its weight of watery moisture, and recovers this weight again on being diffolved and crystallized (a); from whence it would follow, that one part of melted nitre is equivalent to two of the crystals: but there was probably some mistake in this experiment, for I have repeated it with different parcels of nitre, and never found the lofs to be fo much as one twentieth of its weight.

Trochifci e

Nitre may be commodiously taken in the form of troches; one part of the purified falt is commonly ground with three parts of fine fugar, nitro Pb. L. and the mixture made up with mucilage of gum tragacanth. In this and all other folid forms it is accompanied, however, with one inconvenience; being liable, especially when the dose is considerable, to occasion a pain or uneasiness at the stomach, which can be prevented only by plentiful dilution. A liquid form is therefore, in general, the most eligible, and may be eafily rendered grateful by a proper addition of fugar.

Sal prunellæ Crystallus mineralis.

Sal poly-chrest Ph. E.

The chemists have thought to improve the virtue of nitre, by deflagration with a small portion of sulphur: they melt the nitre, in a crucible, and gradually sprinkle on it one twenty-fourth its weight of flowers of fulphur: when the deflagration is over, they pour out the melted falt into clean, dry, warm brass moulds, so as to form it into little cakes. In this process, a part of the acid of the nitre, and the inflammable principle of the fulphur, detonating together, are both destroyed; while that part of the alkali of the nitre, which is thus forfaken by its acid, unites with the acid of the fulphur, which is the fame with that of vitriol, into a new neutral falt, the fame with vitriolated tartar; and the preparation is found to be no other than a mixture of unchanged nitre with a small portion of this vitriolated falt. If the nitre and fulphur be taken in equal quantities, the mixture injected by a little at a time into a red-hot crucible, and kept in till all detonation ceases, nearly the whole of the nitre will thus be changed; and the remaining falt, purified by folution in water, proves almost wholly the same with vitriolated tartar.

The fame falt is produced by pouring gradually on nitre the pure acid of vitriol or fulphur: this acid, uniting with the alkali, difengages the

acid of the nitre, which begins to exhale, immediately on mixture, in yellow or red fumes, and may be collected by distillation in a glass retort with a moderate fire. Two parts of nitre to one of vitriolic acid, is a Acidum niproper proportion for difengaging all the acid of the nitre; the remaining falt is nearly a pure vitriolated tartar. If three parts of nitre be used nitri glauberi Ph. E. to one of the vitriolic acid, a part of the nitre remains unchanged: on diffolving the whole refiduum in hot water, and fetting the filtered folution to crystallize, the vitriolated salt shoots first, greatest part of the nitre continuing dissolved.

The nitrous spirit is obtained also by distillation in a strong fire with vitriol in fubstance; the vitriol parting, when strongly heated, with its own acid, which then acts upon the nitre and extricates its acid in the fame manner as when the pure vitriolic acid is used. The spirit thus distilled, called aqua fortis, is more phlegmatic than the preceding, in proportion as the vitriol employed contains more phlegm than the oil of vitriol: it is likewife liable to an admixture of the vitriolic acid, more or less of which is generally forced over. The proportion directed by the London college is three parts of nitre, three of green vitriol uncalcined, and one and a half of the same vitriol calcined. The ingredients are well mixed together, the distillation performed in an earthen Aqua fortis retort or an iron pot fitted with an earthen head and a receiver, and continued fo long as any red vapours arife. * The college of Edinburgh has now discarded this kind of preparation, and directs a weaker nitrous Acidum niacid to be made by mixing equal parts of the strong acid and pure water.

The nitrous spirit, usually distilled from rough nitre, contains often an admixture of the marine acid as well as of the vitriolic. The first is discovered, and separated, by dropping in a little solution of filver, the latter by a folution of chalk or any other calcareous earth, made in the pure nitrous acid; the filver abforbing the marine acid, and the chalk the vitriolic, and forming with those acids, respectively, indissoluble concretes, which immediately render the liquor milky, and on standing settle to the bottom. The solutions are to be cautiously and flowly dropt in, fo long only as they continue to produce a milkiness: in case of an excess in their quantity, if the spirit is required perfectly pure, it is to be rectified by redistillation.

By the property on which the above method of purification depends, the nitrous spirit may be readily distinguished from the other two 302 mineral

trofum vulgo spiritus Spir. nitri glauberi Ph. Lond. Nitrum vitriolatum Ph. Lond.

Phar. Lond.

trofum tenue Ph. Ed. mineral acids. By the red or yellowish red colour of its sumes; by its forming with one fourth its weight of sal ammoniac, or with sea salt or its acid, a menstruum that perfectly dissolves gold; by its deflagrating on the contact of any inflammable matter, when heated to ignition, whatever other body it be previously combined with; it may with certainty be distinguished both from those and from every other known species of acid.

This acid has been fometimes given as a diuretic, from two or three to fifty drops, diluted largely with water; but its principal use is in combination with other bodies.

Nitrum cubicum. fo

Nitrum flammans, volatile, five ammoniacale. Combined with vegetable fixt alkalies, it reproduces common nitre. With the mineral fixt alkali, or foda, it composes a species of nitre in some respects different from the common, crystallizing not into a prismatic but a cubical figure; with volatile alkalies, a subtile pungent salt remarkable for its solubility in spirit of wine: *(a) of these two compounds, the medicinal qualities are little known, though they should seem to be well deserving of enquiry. The acid, in the most concentrated state in which it is commonly met with, saturates about five sixths its weight of vegetable fixt alkali (b).

*(a) NITRUM FLAMMANS, VOLATILE, AMMONIACALE. This falt diffolves readily in water, and becomes pappy or fluid in a moift air: by flow evaporation in gentle warmth it shoots into large crystals, much resembling those of common nitre. It disfolves in fix times its weight, or less, of rectified spirit of wine. In a heat equal to that of boiling water, it melts and looks like oil, without suffering any loss of its substance; on increasing the heat a very little beyond that degree, it begins to exhale, and in a little time is wholly dissipated: the sumes, caught in proper vessels, condense not into a concrete salt, but a sluid spirit; in which, however perfect the neutralization was at first, the acid appears now to prevail. The salt thrown into a red-hot crucible, without addition of any inflammable matter, emits bright slames, without detonation or noise: the slashes continue to play on the surface till the whole quantity of the salt is dissipated.

This falt is in taste similar to common nitre, but somewhat sharper or more penetrating. Taken in doses of from ten to twenty-five grains, it sensibly promotes urine; and if the patient is kept warm, perspiration or sweat. It is recommended by Kurella, preserably to the other neutral saline medicines, in inflammatory cases, in exanthematous severs, and as an attenuant and resolvent in obstructions of the viscera. He gives it either in powder, mixed with absorbents neutralized by lemon juice, or dissolved in well dulcissed spirits of vitriol or nitre, in which last form he finds it in some cases to answer best: from fisteen to twenty-sive drops of the saturated solution are given for a dose in any agreeable warm liquor. He recommends it likewise externally against inflammations, erysipelases, and gouty pains, dissolved in spirit of wine, either by itself, or with the addition of camphor and opium. M. S. of Dr. Lewis.

(b) Homberg, Memoires l'acad. roy. des scienc. de Paris, pour l'ann. 1699.

Solutions

Solutions of calcareous earths in this acid are in tafte bitterish and very pungent. They are difficultly made to assume a crystalline appearance; and when evaporated and exficcated by heat, the dry falt deli- Nitrum calquiates again in the air. This falt has not hitherto been employed rum. medicinally, nor is it as yet much known. It is a common ingredient in waters, which when its quantity is confiderable, it renders hard and indisposed to putrefy, apparently impeding putrefaction in a much greater degree than an equal quantity of fea falt. Alkaline falts, fixt or volatile, added to the folutions, precipitate the earthy basis; and uniting with the acid in its flead, compose therewith, according to the species of alkali employed, the common, cubical, or ammoniacal nitre mentioned in the preceding paragraph.

The nitrous spirit dissolves zinc, iron, copper, bismuth, lead, mercury, and filver, the most readily of all the acids: tin it disfolves imperfectly: regulus of antimony it only corrodes: fee the respective metals.

The concentrated acid, combined with a due proportion of rectified spirit of wine, loses its acidity; the coalition of the two producing a new compound, of a gratefully pungent tafte and colour, and which is given from a few drops to a tea-spoonful or more as mildly aperient, diuretic, antiphlogistic, in some degree anodyne and antispasmodic. On mixing the two spirits together, a great heat, ebullition, and noxious red vapours arise: this conflict is less violent when, cautiously and by little and little, the acid spirit is added to the vinous, than when the vinous is added to the acid. It is prudent also to place the bottle containing the spirit of wine, in a vessel of cold water. One part of the Spiritus niftrong acid spirit is commonly taken to three of the spirit of wine t, or half a pound to a quart +: the mixture, after standing for some time Acidum nithat the two liquors may in some degree unite, is fet to distil with a vulgo spirigentle fire, by which the union is completed, and the very volatile dulcified spirit separated from the more fixt acid that remains undulcified. 1 Ph. Edinb. The distillation is directed to be continued so long as the spirit that comes over raifes no effervescence with fixt alkaline salts+; it may be regulated more commodiously by performing the process in a water bath 1, for all that rifes in this heat will be found to be a pure dulcified spirit.

tri dulcis + Ph. Lond. tri vinofum, tus nitri dulcis

A fubtile

450

Nitrous ether. A fubtile ethereal fluid, fimilar in its general qualities to that defcribed under the head of vitriolic acid, is obtainable with the nitrous in a more compendious manner. If equal parts by measure of spirit of nitre and spirit of wine, of moderate strength, be mixed together, the bottle closely stopt, and set in a cool place, a large proportion of ether rises to the surface in a few days: it may be purified from the adhering acid, by shaking it with water in which some fixt alkaline salt has been dissolved, and then drawing off the ether by distillation. The medicinal qualities of this subtile sluid are not as yet much known.

NUMMULARIA.

NUMMULARIA Pharm. Parif. Nummularia major lutea C. B. Hirundinaria. Centimorbia. Lysimachia Nummularia Linn. Money-wort: a low creeping plant, with square stalks, and smooth little roundish or heart-shaped leaves set in pairs at the joints upon short pedicles: in their bosoms appear yellow solitary monopetalous flowers, each divided into sive oval segments, and sollowed by a small round capsule sull of minute seeds. It is perennial, grows wild in moist pasture grounds, and slowers from May to near the end of summer.

This herb is accounted restringent, antiscorbutic, and vulnerary. Boerhaave looks upon it as similar to a mixture of scurvygrass with sorrel: it appears indeed to have some degree both of pungency and acidity, but it is far weaker than those herbs, or than any mixture of the two.

NUX MOSCHATA.

NUX MOSCHATA Pharm. Lond. & Edinb. Nux myriftica fructu rotundo C. B. Nucifta. Myriftica officinalis Linn. Nutmeg: the aromatic kernel of a large nut, produced by a tree faid to refemble the pear tree, growing in the East Indies. The outer part of the fruit is a soft fleshy substance like that of the walnut, which spontaneously opens when ripe: under this lies a red membrane called mace, forming a kind of reticular covering, through the sissures of which is seen the hard woody shell that includes the nutmeg. Two sorts of this kernel are distinguished: one of an oblong figure, called male; the other roundish,

or of the shape of an olive, called female: this last is the officinal species, being preferred to the other on account of its stronger and more agreeable flavour, and its being, as is faid, less subject to become carious. The nutmegs are cured, according to Rumphius, by dipping them in a fomewhat thick mixture of lime and water, that they may be every where coated with the lime, which contributes to their preservation.

THE nutmeg is a moderately warm, grateful, unctuous spice; supposed to be particularly useful in weakness of appetite, and the nauseæ and vomitings accompanying pregnancy, and in fluxes; but liable, when taken too freely, to fit very uneasy on the stomach, and, as is said, to affect the head. Roasted with a gentle heat, till it becomes easily Nux mosch. friable, it proves less subject to these inconveniences, and is supposed likewise to be more useful in fluxes.

torrefacta Pharm. Lond.

Nutmegs, distilled with water, yield nearly one fixteenth (a) their weight of a limpid effential oil, very grateful, possessing the flavour of Ol. still. feu the spice in perfection, and which is faid to have some degree of an antispasmodic or hypnotic(b) power: on the surface of the remaining Pharm. Lond. decoction is found floating an unctuous concrete matter like tallow, of a white colour, nearly infipid, not eafily corruptible, and hence recommended as a basis for odoriferous balsams (c): the decoction, freed from this febaceous matter, and inspissated, leaves a weakly bitter fubastringent extract. Rectified spirit takes up, by maceration or digeftion, the whole smell and taste of the nutmegs, and receives from them a deep bright yellow colour: the spirit, drawn off by distillation from the filtered tincture, is very flightly impregnated with their flavour; greatest part of the specific smell, as well as the aromatic warmth, bitterishness and subastringency of the spice remaining concentrated in the extract. The effential oil, and an agreeable cordial water, lightly flavoured with the volatile parts of the nutmeg by drawing Aqua nucis off a gallon or nine pounds of proof spirit from two ounces of the spice, mosch. Phar.

effent. nucis moschatæ

mosch. Phar.

⁽a) Hoffman, Observationes physico-chymicæ, lib. i. obs. 1.

⁽b) Miscell. nat. curiosor. dec. III. ann. ii. obs. 120. Bontius, de medicina Indorum, p. 20.

^{* (}c) After a fluid essential oil had been procured from nutmegs by distillation, on repeating the process upon the residuum, an oil of a butyraceous consistence arose, which possessed the tafte and odour of the nutmeg, and was perfectly foluble in alcohol. Gaubii Adversar.

are kept in the shops. Both the oil, and the spirituous tincture and extracts, agree better with weak stomachs than the nutmegs in substance.

Nutmegs, heated, and strongly pressed, give out a sluid yellow oil, which concretes on growing cold into a febaceous confistence. Rumphius informs us, that in the spice islands, when the nuts are broken. those kernels which appear damaged, carious, or unripe, are separated for this use, and that seventeen pounds and a quarter of such kernels yield only one pound of oil, whereas, when the nutmeg is in perfection.

it is faid to afford near one third its own weight.

Ol. nucis mosch. expreffum, macis vulgo dictumPhar.

Two kinds of sebaceous matter, faid to be expressed from the nutmeg, are distinguished in the shops by the name of oil of mace: the best fort. brought from the East Indies in stone jars, is somewhat soft, of a yellow colour, and of a strong agreeable smell greatly resembling that of the Lon. & Edinb. nutmeg itself: the other comes from Holland in solid masses, generally flat and of a square figure, of a paler colour and much weaker smell. These oils are employed chiefly externally in stomach plasters, and in anodyne and nervine unguents and liniments. They appear to be a mixture of the gross sebaceous matter of the nutmeg with a little of the effential or aromatic oil; both which may be perfectly separated from one another by maceration or digestion in rectified spirit, or by distillation with water. The spirituous tincture, the distilled water, and the effential oil, are nearly fimilar to those drawn from the nutmeg itself, the pure white febaceous fubstance being left behind.

NUX PISTACIA.

NUX PISTACIA Pharm. Edinb. (a) PISTACHIO nut: an oblong, pointed nut, about the fize and shape of a filberd; including a kernel of a pale greenish colour, covered with a yellowish or reddish skin. It is the produce of a large tree, with winged leaves, refembling those of the ash, pistacia peregrina fructu racemoso sive terebinthus indica theophrasti C. B. Piftachia vera Linn. which grows spontaneously in the eastern countries, and bears the cold of our own.

PISTACHIO nuts have a pleafant sweetish unctuous taste, resembling that of fweet almonds: their principal difference from which confifts

in their having rather a greater degree of sweetness, accompanied with a light grateful flavour, and in being more oily, and hence somewhat more emollient, and perhaps more nutritious. They have been ranked among the principal analeptics, and greatly esteemed by some in certain weaknesses and emaciations. They are taken chiefly in substance, their greenish hue rendering them unsightly in the form of an emulsion. They are very liable to grow rancid in keeping.

NUX VOMICA.

NUX VOMICA Pharm. Parif. Nux metella. Vomic nut: a flat roundish seed or kernel, about an inch broad and near a quarter of an inch thick, with a prominence in the middle on both sides, of a grey colour, covered with a kind of woolly matter, internally hard and tough like horn. It is the produce of a large tree growing in the East Indies, called by Plukenet cucurbitisera malabariensis, ænopliæ soliis rotundis, fructu orbiculari rubro cujus grana sunt nuces vomicæ officinarum; by Linnæus, Strychnos Nux Vomica.

This feed discovers to the taste a considerable bitterness, but makes little or no impression on the organs of smell. It has been recommended in tertian and quartan severs, in virulent gonorrheas, and as an alexipharmac: Fallopius relates, that it was given with success in the plague; that in doses of from a scruple to half a dram, it procured a plentiful sweat; and that where this evacuation happened, the patient recovered (a). At present it is looked upon, and not without good foundation, as a deleterious drug; which, though like many other deleterious substances, capable, in certain doses and in certain circumstances, of producing happy effects, has its salutary and pernicious operations so nearly and so indeterminably allied, that common prudence forbids its being ventured on. Hossman tells us of a girl of ten years of age, to whom sifteen grains, given at twice, for the cure of an obstinate quartan, proved mortal (b). The principal symptoms it has been observed to produce, in human subjects and brutes (c), are, great anxieties,

⁽a) Tract. de tumoribus præternaturalibus, cap. 27.

⁽b) Philosophia corp. buman. morbosi, P. ii. cap. viii. §. 8.

⁽c) Vide Wepfer, De cicuta aquatica, cap. xiii. p. 194, & feq.

strong convulsions or epileptic fits, paralytic tremors and resolutions, a great increase of the motion of the heart and of respiration, and reachings and subversions of the stomach. Dissections of dogs killed by it have shewn no material injury of the grosser parts; from whence we may presume that it is the nervous system which it immediately offends. It is probable, that the active matter of this seed is of the same nature with that of bitter almonds, but more developed and in a more concentrated state.

* The nux vomica was lately used in Sweden in an epidemic dyfentery, as it is said, with remarkably good effects. A scruple of the powder was given to adults once a day in barley water, proper evacuants having been premised. Bergius (a), however, afferts, that though the flux was suppressed for twelve hours by this medicine, it never failed to return. He also mentions a case in which the above dose caused convulsive stretchings and vertigo; and after the cure of the dysentery by other medicines, a pain in the stomach and epigastric region remained for a long time. In the isle of Ceylon the nux vomica is said to be used internally as a specific against the bite of a species of water-snake.

THE wood or roots of the tree, or of other trees of the same genus, are sometimes brought from the East Indies under the name of lignum colubrinum (Pharm. Paris.) or snakewood, in pieces about the thickness of a man's arm, covered with a brownish or rusty coloured bark, internally of a yellowish colour with whitish streaks.

This wood, in rasping or scraping, emits a faint not disagreeable smell; and when chewed for some time discovers a very bitter taste. Cartheuser relates, that it gives a gold-coloured tincture both to water and spirit, and that the inspissated extracts are brownish; that the watery insusion has an agreeable smell like that of rhodium, the spirituous little or none; that the insussons and extracts made with both menstrua are very bitter; that the quantity of watery extract amounts to one sixth of the wood, and that of the spirituous to near one fourth; and that the wood remaining after the action of spirit, yields still, to water, a gold-coloured

tincture, and one eighth its weight of a bitter subacrid extract: from whence water appears to be the proper menstruum of its active matter.

The lignum colubrinum has been recommended, in small doses, not exceeding half a dram, as an anthelmintic, and in obstinate quartans, jaundices, cachexies, and other chronical disorders: it is said to operate most commonly by sweat, sometimes by stool, and sometimes by vomit. It appears however to be possessed of the same ill qualities with the nux vomica itself, though in a lower degree, having in sundry instances been productive of convulsions, tremors, stupors, and disorders of the senses.

THE faba indica Pharm. Parif. Faba sancti ignatii, or faba febrifuga, is the produce of a tree of the same kind, growing in the East Indies and in the Philippine islands, called by Plukenet cucurbitisera malabathri foliis scandens, catalongay & contara philippinis orientalibus dicta, cujus nuclei pepitas de besayas aut catbalogan & sancti ignatii ab hispanis, igasur & mananaog insulanis nuncupati; by Linnæus, Strychnos Ignatii. The seeds of the gourd-like fruit, improperly called beans, are of a roundish sigure, very irregular and uneven, about the size of a middling nutmeg, semitransparent, and of a hard horny texture.

These seeds have a very bitter taste, and no considerable smell: when fresh they are said to have somewhat of a musky scent. Neumann observes, that an extract made of them by rectified spirit impresses at first a very agreeable bitterness, somewhat like that of peach kernels, which going off leaves in the mouth a strong bitter; that an extract made with water is likewise bitter; that the watery extract is greenish and in quantity one half of the seeds, the spirituous yellowish and little more than one sifth; that the seeds remaining after the action of water scarcely gave out any thing to spirit, but that after spirit they yielded above one fourth of extract with water.

St. Ignatius's bean is faid by father Camelli to be employed by the common people in the Philippine islands against all diseases. The effects attributed to it are similar to those of the two foregoing substances: he observes, that it generally vomits, sometimes purges, and almost always produces in the Europeans, though not in the Indians, spasmodic motions; that the dose in substance, as an emetic, is ten or twelve grains, to be taken an hour after eating; and that in smaller doses it spasmodic formetimes

fometimes promotes a plentiful fweat (a). Neumann says he has known intermitting fevers cured by drinking, on the approach of a paroxysm, an insusion of some grains of the seed made in carduus water (b); and I have been informed, that two grains were found to have as much effect as a full dose of bark. This seed, nevertheless, as it apparently partakes of the qualities of the two preceding articles, seems much too hazardous for general use.

NYMPHÆA.

WATER-LILY: an aquatic plant, with thick firm roundish leaves, furnished with two obtuse ears at the pedicle, floating on the surface of the water: the flowers, which stand on separate pedicles, are large, composed of several petala with numerous stamina in the middle, followed by single capsules full of blackish shining seeds: the root is long, thick, internally white and sungous.

- 1. NYMPHÆA ALBA sive Nenuphar Pharm. Edinb. (c) Nymphæa alba major C. B. Leuconymphæa. Nymphæa alba Linn. White waterlily: with white flowers set in four-leaved cups: the seed vessels round, and the roots externally brownish or blackish.
- 2. NYMPHÆA LUTEA: Nymphæa major lutea C. B. Nymphæa lutea Linn. Yellow water-lily: with yellow flowers fet in large five-leaved cups, the feed vessels shaped like a pear, and the roots externally greenish.

BOTH these plants are found in rivers and large lakes; the yellow is most common: they are perennial, and flower usually in June. The roots and flowers have been employed, both internally and externally, as demulcent, anti-inflammatory, and in some degree anodyne. Their virtues, however, do not appear to be very great, as they have no smell, at least when dried, and but little taste: extracts made from them both by water and spirit are weakly bitterish, subastringent, and subsaline. Lindestolpe informs us, that in some parts of Sweden, the roots, which

⁽a) Philosophical transactions, numb. 250.

⁽b) Chymia medica, &c. i. 737. Chemical works, p. 347. (c) Expunged.

are the strongest part, were in times of scarcity used as food, and did not prove unwholesome.

OCHRA.

ochra five Minera ferri lutea vel rubra Pharm. Parif. Ochre: an argillaceous earth; less tenacious, when moistened, than the clays and the boles; impregnated with a calx of iron, and thereby tinged of a yellow or red colour. The dark red fort is called reddle or ruddle, rubrica fabrilis Pharm. Edinb. (a) the yellow fil; ochra plinio & latinis fil dicta Charleton. Those which are naturally yellow become red by burning. Both kinds are dug in several parts of England.

These earths discover their argillaceous nature, by burning hard in the fire; and their ferrugineous impregnation, by digestion in aqua regis, which extracts the iron, leaving the earth nearly white. To the taste they seem somewhat astringent, in consequence, not of the metallic, but of the earthy part, for the iron is in such a state as not to be acted on by any sluid that exists in the bodies of animals: it may therefore be presumed, that they do not differ materially, in virtue, from the boles; except in being less viscid, and therefore of less efficacy for obtunding acrid humours: see Bolus and Cimolia. Among us they are rarely or never used medicinally under their own name; though sometimes applied in the shops to the counterfeiting of earths that are less common.

OCIMUM.

OCIMUM, Basilicum, Herba regia. BASIL: a plant, with square stalks; oval leaves set in pairs; and long spikes of labiated flowers, whose upper lip is divided into sour parts, the lower entire: the cup also has two lips, one cut into sour sections, the other into two.

1. Ocimum Pharm. Edinb. (a) Ocimum vulgatius C. B. Ocymum medium citratum Ger. Ocimum Bafilicum Linn. Common or citron bafil: with most of the leaves indented, and the flower-cups edged with fine hairs.

2. OCIMUM CARYOPHYLLATUM: Ocimum minimum C. B. & Linn. Small or bush basil: with uncut leaves.

BOTH these plants are natives of the eastern countries, and sown annually in our gardens for culinary as well as medicinal uses. The seeds, which rarely come to perfection in this climate, especially those of the second sort, are brought from Italy and the south of France.

The leaves of basil are accounted mildly balsamic: insusions of them are sometimes drank as tea in catarrhous and uterine disorders, and the dry leaves in substance made an ingredient in cephalic and sternutatory powders. They are very juicy, of a weakly aromatic and very mucilaginous taste, and of a strong smell, which is somewhat disagreeable when the herbs are fresh, but is improved by drying: those of the first sort approach to the lemon scent, those of the second to that of cloves. Distilled with water, they yield a considerable quantity of essential oil, of a penetrating fragrance, commended by Hossman as a nervine, similar, but greatly superiour, to oil of marjoram (a).

* OE NANTHE.

OENANTHE: this is the botanical name of a genus of plants of the umbelliferous class, of which there are three species natives of Great Britain. One of these only is known by its effects on the human body, the

OENANTHE CROCATA Linn. Hemlock dropwort: this is a large umbelliferous plant, growing in ditches and other moist places; with pinnated leaves, resembling those of celery or chervil, and ribbed stalks. Its roots afford the easiest mark of distinction, which are white, thick, and short, and grow several together, forming a kind of bunch.

The hemlock dropwort has long been known as a most dangerous poison; the most virulent, perhaps, that this country produces. Its roots or leaves eaten by mistake, have frequently proved fatal, occasioning violent sickness and vomiting, rigors, convulsions, delirium, and other terrible affections of the nervous system. The head has been said to be

affected even by being in the same room with a quantity of the plant. Like so many other deleterious vegetables, it, however, is capable of being rendered a powerful remedy. A case is published by Dr. Pulteney in the Philos. Transact. Vol. LXII. in which this plant, used by mistake instead of the water parsnep, proved remarkably efficacious in removing an inveterate scorbutic complaint, which had resisted a variety of other remedies. The dose first given was a common spoonful of the juice of the root, which at the first exhibition produced very alarming effects. This was afterwards reduced to three tea-spoonfuls; which quantity was persisted in a considerable time, and then changed for a tea of the leaves. The medicine never proved purgative, but was diuretic. It always occasioned a degree of vertigo; accompanied, when the juice itself was taken, with nausea and sickness.

If this experiment be imitated, it is obvious that the greatest degree

of caution will be necessary.

OLEA.

OLEA sativa C. B. Olea europæa Linn. OLIVE: an evergreen tree, with oblong, narrow, willow-like leaves, and monopetalous whitish flowers, cut into four sections, followed by clusters of oval black fruit, containing, under a slesshy pulp, a hard rough stone. It is a native of the southern parts of Europe, and bears the ordinary winters of our own climate.

THE fruit of this tree (oliva) has a bitter, austere, very disagreeable taste: pickled, as brought from abroad, it proves less ungrateful, and is supposed to promote appetite and digestion, and attenuate viscid phlegm in the first passages: the Lucca olives, which are smaller than the others, have the weakest taste; and the Spanish, or larger, the strongest: those brought from Provence, which are of a middling size, are in general most esteemed. But the principal consumption of olives is in the preparation of the common sallad oil (oleum olivarum Pharm. Lond. & Edinb.) which is obtained by grinding and pressing them when thoroughly ripe: the finer and purer oil issues first by gentle pressure; and inferiour sorts, on heating the residuum and pressing it more strongly. All these oils contain a portion of watery moisture; and of the mucilaginous substance of

the

the fruit: to separate these, and thus prevent the oil from growing rancid, some sea salt is added, which not being dissoluble in the pure oil, imbibes the watery and mucilaginous parts, and sinks with them to the bottom. As this oil grows thick in a moderate degree of cold, a part of the salt, thrown up by shaking the vessel, is sometimes detained in it, so as to render the taste sensibly saline. In virtue, it does not differ materially from the other slavourless expressed oils: it is preferred to the others for dietetic uses, and in plasters and unguents, but is more rarely employed as an internal medicine.

OLIBANUM.

OLIBANUM Pharm. Lond. & Edinb. OLIBANUM: a gummy-refin brought from Turkey and the East Indies, usually in drops or tears like those of mastich, but larger, of a pale yellowish colour, which by age becomes reddish. It is the product of a tree of the juniper kind growing in Arabia; the juniperus lycia of Linnæus.

THIS gummy-refin has a moderately strong, not very agreeable smell, and a bitterish somewhat pungent taste: in chewing, it sticks to the teeth, becomes white, and renders the faliva milky. Laid on a red-hot iron, it readily catches flame, and burns with a strong, diffusive, not unpleafant fmell: it is supposed to have been the incense used by the ancients in their religious ceremonies, though it is not the substance now known by that name in the shops. On trituration with water, greatest part of it dissolves into a milky liquor, which on standing deposites a portion of refinous matter, and being now gently inspissated, leaves a yellow extract, which retains greatest part of the smell as well as the tafte of the olibanum; its odorous matter appearing to be of a less volatile kind than that of most other gummy-refins. Rectified spirit dissolves less than water, but takes up nearly all the active matter: the transparent yellowish solution, inspissated, yields a very tenacious refin, in which the active parts of the juice are fo enveloped and locked up, that they are scarcely to be discovered, either by the smell or taste.

Olibanum is recommended in disorders of the head and breast, in hæmoptoës, and in alvine and uterine sluxes: the dose is from a scruple to a dram or more.

ONONIS.

ONONIS & Anonis seu Aresta bovis Pharm. Edinb. (a) Anonis spinosa slore purpureo C.B. Resta bovis. Remora aratri. Ononis spinosa Linn. Rest-harrow: a plant with flexible branches terminating in sharp prickles; small oval indented leaves, standing generally three together, without pedicles; and purplish papilionaceous flowers, set in pairs, followed each by a short pod containing three unequal kidney-shaped seeds. It is perennial, grows wild in waste grounds and dry fields, and with its long tough spreading roots obstructs the plough or harrow.

THE roots of rest-harrow have a faint unpleasant smell, and a sweetish, bitterish, somewhat nauseous taste. Their active matter is confined to the cortical part; which has been sometimes given in powder, in doses of a dram, and made an ingredient in apozems or decoctions, as an aperient and diuretic. Its virtue is extracted both by water and spirit.

OPIUM.

OPIUM Pharm. Lond. & Edinb. OPIUM: a concrete gummy-refinous juice; fomewhat foft and tenacious, especially when much handled or warmed; of a dark reddish brown colour in the mass, and when reduced into powder yellow. It is brought from Egypt, Persia, and some other parts of Asia, in slat cakes or irregular masses, from four to about sixteen ounces in weight, covered with leaves to prevent their sticking together.

It is extracted from the heads of white poppies (see Papaver) which in those countries are cultivated in fields for this use. Kæmpser reports, that the heads, when almost ripe, are wounded with a five-edged instrument, by which as many parallel incisions are made at once from top to bottom; that the juice which exudes is next day scraped off, and the other sides of the heads wounded in like manner; and that the juice is afterwards worked with a little water, till it acquires the consistence, tenacity, and brightness of the finest pitch. The best opium was

formerly called Thebaic opium, from its being prepared about Thebes in Egypt: no distinction is now made in regard to the places of its production, though the epithet thebaic serves to distinguish some of its officinal preparations.

OPIUM has a faint disagreeable smell, and a bitterish, somewhat hot, biting taste. Watery tinctures of it strike a black colour with chalybeate solutions, and thus seem to discover some astringency. Mixed with the serum of blood, they thicken and render it whitish; and on blood itself, newly drawn, they have nearly a like effect: Mr. Eller observes, that on examining with a microscope blood thickened by a vinous tincture of opium, the nature of its globules seemed to be destroyed. But neither from these, nor any of the other known sensible properties of this drug, can its surprizing operation in the human body be deduced.

Taken in proper doses, it commonly procures sleep, and a temporary respite from pain, or the action of any stimulating power. The cause of the pain it in many cases confirms or augments; and in not a few, it fails even of giving palliating relief. The cases in which it is proper or improper will be best understood from a view of its general effects; which, so far as experience has hitherto discovered them, are the

following.

It renders the folids, while the operation of the opium continues, less sensible of every kind of irritation, whether proceeding from an internal cause, or from acrimonious medicines, as cantharides, and the more active mercurials, of which it is the best corrector - It relaxes the nerves; abating or removing cramps or spasms, even those of the more violent kind; and increasing paralytic disorders and debilities of the nervous system-It incrassates thin serous humours in the sauces and adjacent parts; by which means, it proves frequently a speedy cure for fimple catarrhs and tickling coughs; but in phthifical and peripneumonic cases, dangerously obstructs expectoration, unless this effect be provided against by suitable additions, as ammoniacum and squills-It produces a fullness and distension of the whole habit; and thus exasperates inflammations both internal and external, and all plethoric fymptoms - It promotes perspiration and sweat; but restrains all other evacuations, unless when they proceed from a relaxation and infensibility of the parts, as the colliquative diarrhoe in the advanced stage of hectic feversfevers—It promotes labour-pains and delivery (a) more effectually than the medicines of the stimulating kind usually recommended for that purpose; partly perhaps by increasing plenitude, and partly by relaxing the solids or taking off spasmodic strictures—And indeed all the preceding effects are perhaps consequences of one general power, being nearly allied to those which natural sleep produces (b).

The operation of opium is generally accompanied with a flow but strong and full pulse, and a slight redness, heat and itching of the skin: it is followed by a weak and languid pulse, lowness of the spirits, some difficulty of breathing or a sense of tightness about the breast, a slight giddiness of the head, dryness of the mouth and sauces, and some degree of nausea. Given on a full stomach, it commonly occasions a nausea from the beginning, which continues till the opium is rejected along with the contents of the stomach. Where the evacuation of acrid humours, accumulated in the first passages, is suppressed by it, great sickness and uneasiness are generally complained of, till the salutary discharge either takes place again spontaneously or is promoted by art.

An over dose occasions either immoderate mirth or stupidity, a redness of the face, swelling of the lips, relaxation of the joints, vertigo, deep sleep with turbulent dreams and startings, convulsions, and cold sweats. Geoffroy observes, that those who recover, are generally relieved by a diarrhæa, or by a profuse sweat, which is accompanied with a violent itching. The proper remedies, besides emetics, blisters, and bleeding, are acids and neutral mixtures: Dr. Mead says he has given, with extraordinary success, repeated doses of a mixture of salt of wormwood with lemon juice.

A long continued use of opium is productive of great relaxation and debility, sluggishness, heaviness, loss of appetite, dropsies, tremors, acrimony of the humours, frequent stimulus to urine, and propensity to venery. On leaving it off, after habitual use, an extreme lowness of the spirits, languor, and anxiety, succeed; which are relieved by having again recourse to opium, and in some measure by spirituous or vinous liquors.

With regard to the dose, one grain is generally a sufficient, and sometimes too large a one: maniacal persons, and those who labour under violent spasms, require oftentimes two, three, or more grains; though even in these cases, it is generally more advisable to repeat the dose at proper intervals, than to enlarge it. By frequent use, much greater quantities may be borne: the Turks, who habituate themselves to opium as a succedaneum to spirituous liquors, are said to take commonly a dram at a time, and Garcias says that he knew one who every day took ten drams.

Opium colat. vel Extract. thebaic. Ph. Lon. OPIUM appears to confift of about five parts in twelve of gummy matter, four of refinous matter, and three of earthy or other indiffoluble impurities (a). From these last it is purified, in the shops, by softening the opium with boiling water, in the proportion of a pint to a pound, into the consistence of a pulp, with care to prevent its burning; and whilst it remains quite hot, strongly pressing it from the seces through a linen cloth: the strained opium is then inspissated in a water-bath, or other gentle heat, to its original consistence. When thus softened with a small quantity of water, the gummy and resinous parts pass the strainer together; whereas, if dissolved by a larger quantity, they would separate from one another.

It has been disputed, whether it is in the gummy or in the refinous parts of opium, that its activity resides. From the experiments of Hossman(b) and Neumann(c), it seems to be neither in the direct gum, nor in the direct resin, but in a certain subtile part of the resinous matter, somewhat analogous to essential oils, but of a much less volatile kind: they report, that on boiling the opium in water, there arises to the surface a frothy, viscid, unctuous, strong-scented substance, to the quantity of two or three drams from sixteen ounces: that this substance, in the dose of a few grains, has killed dogs that could bear above a dram of crude opium; that in distillation with water, though it does not rise itself, it gives over, at least in part, the active principle of which it is the matrix; impregnating the distilled liquor with its scent and its soporific power; as essential oils exhale their odoriferous principle in the air, without being dissipated themselves. What this subtile and highly active

⁽a) Alfton, Edinburgh medical essays, vol. v. art. 12.

⁽b) Diss. de opii correctione genuina & usu, Oper. supplement. II. P. i. p. 645. Not. ad Poterium, p. 437.

⁽c) Chymia medica, vol. i. p. 996. Chemical works, p. 308.

principle really is, in effential oils, in odorous vegetables that yield no

oil, and in opium, is equally unknown.

Both water and rectified spirit extract, difficultly, by maceration or digestion, the active matter of opium, and receive from it a yellow or brownish tincture. The watery solution is found to contain great part of the refin along with the gum; and the spirituous, a smaller proportion of the gum along with the refin. Such part of the gum as is left by spirit, and such part of the resin as is left by water, seem to be equally

Tinctures of opium in water, wine, and proof spirit, have the same effects as the opium in substance; with this difference, that they exert themselves sooner in the body, and are less disposed to leave a nausea on the stomach. Tinctures made in rectified spirit are said to act with greater power than the others: Geoffroy relates, from his own observation, that while the watery and vinous tinctures occasioned quiet fleep, the spirituous brought on a phrenzy for a time. It is faid likewife, that alkaline falts diminish the soporific virtue of the opium; that fixt alkalies render it diuretic, whilst volatile ones determine its action to the cutaneous pores; and that acids almost entirely destroy its force.

The officinal tinctures of opium are made in wine or proof spirit. The college of London directs two ounces of strained opium to be ma- Tinetur.thecerated without heat for a week in a pint of mountain, with the addition of a dram of cinnamon and as much cloves to alleviate the ill fmell of the opium: the college of Edinburgh orders the same quantity of crude opium to be digested for four days in a pound and a half of spirituous cinnamon water: a mixture of wine and proof spirit has been sometimes num liquimade choice of, in order to prevent in some measure an inconvenience which both of them feparately, confidered as officinals, are liable to, being apt to throw off in long standing a part of the opium, which in wine falls to the bottom, and forms a crust on the surface of spirit. Of the first of the above tinctures twenty drops, and of the latter twenty-five drops, are reckoned to contain one grain of opium: but as these quantities of the menstrua do not easily dissolve all the active matter of so large a proportion of the opium, those doses are generally observed to have somewhat less effect than a grain of the drug in substance. Asdrops also, according to different circumstances, vary in quantity, though

baic. Ph. L.

Tinctura thebaica vulgo laudadum Pb. Ed.

in number the same, it were to be wished that the shops were furnished with a solution of this drug, made in a quantity of menstruum large enough not only for the complete extraction of the active parts, but to admit of the dose being exactly determined by weight or measure.

In a folid form, independently of fuch materials as may be subservient to the other indications of cure, it is sometimes mixed with saponaceous or gummy substances which promote its dissolution in the stomach, and sometimes with resinous ones which render its dissolution and operation more gradual and slow: to these is commonly superadded some aromatic ingredient, to prevent its occasioning a nausea. Thirty-two parts of almond soap, four of strained opium moistened with a little wine, and one of essence of lemons, beaten together; or one part of opium, sour of extract of liquorice, three of Spanish soap, and two of powdered Jamaica pepper, make an elegant pill of the first kind; and sixteen parts of strained storax, eight of saffron, and sive of strained opium, of the latter.

Pil. faponaceæ Pb. Lon.

Pil. thebaicæ vulgo pacificæ Pb. E.

Pil. e ftyrace Pharm. Lond.

Many have endeavoured to correct certain ill qualities, which they fuppose opium to be possessed of, by roasting it, by fermentation, by long continued digestions or boiling, by repeated dissolutions and distillations. These kinds of processes, though recommended by several late writers, do not promise any singular advantage. That they weaken the opium is indeed very probable; but this intention is answered as effectually, and with far greater certainty, by diminishing the dose of the opium itself: for the ill effects, which opium produces in certain circumstances, do not depend on any distinct property or principle, and appear to be no other than the necessary consequences of the same power, by which in other circumstances it proves so beneficial: the only rational way of improving or correcting this valuable drug seems to be, by joining or interposing such medicines, as may counteract or remove those particular effects of it, which in particular cases may be injurious.

OPOBALSAMUM.

OPOBALSAMUM Pharm. Lond. OPOBALSAMUM OF BALSAM OF GILEAD: a refinous juice, obtained from an evergreen tree or shrub (balfamum syriacum rutæ folio C. B.) said to grow in Arabia. The best sort, which naturally exudes from the plant, is scarce known in Europe;

and the inferiour kinds, said to be extracted by lightly boiling the branches and leaves in water, are very rarely seen among us.

The true opobalsam, according to Prosper Alpinus, is at first turbid and white, of a very strong pungent smell, like that of turpentine; but much sweeter and more fragrant, and of a bitter, acrid, astringent taste: on being kept for some time, it becomes thin, limpid, light, of a greenish hue, and then of a gold yellow, after which it grows thick like turpentine, and loses much of its fragrance (a). Some resemble the smell of this balsam to that of citrons, others to that of a mixture of rosemary and sage flowers. I have sometimes met with a curious balsam of this last kind of smell, exceedingly fragrant, limpid, and thin: dropt on water, it spread itself all over the surface, imparting to the liquor a considerable share of its taste and smell: the grosser part, that remained on the top of the water, was so tenacious, as to be easily taken up at once with the point of a needle, which is reckoned, by Alpinus and others, as a characteristic of the true balsam.

This precious balfam is of great esteem in the eastern countries, both as a medicine, and as an odoriferous unguent and cosmetic. Its great scarcity has prevented its coming into use among us: nor are its virtues, probably, superiour to those of some of the resinous juices more common in the shops; all these substances being in their general qualities alike, though differing in the degree of their gratefulness, pungency, and warmth.

OPOPANAX.

OPOPANAX Pharm. Lond. OPOPANAX: a concrete gummy-refinous juice, obtained from the roots of an umbelliferous plant, which grows fpontaneously in the warmer countries, and bears the colds of our own (Pastinaca Opopanax Linn.) The juice is brought from Turkey and the East Indies, sometimes in little round drops or tears, more commonly in irregular lumps, of a reddish yellow colour on the outside with specks of white, internally of a paler colour and frequently variegated with large white pieces.

THIS gummy-refin has a strong disagreeable smell, and a bitter, acrid, somewhat nauseous taste. It readily mingles with water, by triture, into a milky liquor, which on standing deposites a portion of resinous matter and becomes yellowish: to rectified spirit it yields a gold-coloured tincture, which tastes and smells strongly of the opopanax. Water distilled from it is impregnated with its smell, but no essential oil is obtained on committing moderate quantities to the operation.

Opopanax is an useful attenuant and deobstruent, and in considerable doses loosens the belly. It is given from a scruple to a dram, in the same intentions as ammoniacum or galbanum; and joined in smaller doses

as an auxiliary to those and the other deobstruent gums.

ORIGANUM.

ORIGANUM Pharm. Lond. Origanum filvestre, cunila bubula plinii C. B. Agrioriganum sive onitis major Lob. Origanum anglicum Ger. Origanum vulgare Linn. Origanum or Wild marjoram: a plant with firm round stalks, and oval, acuminated, uncut, somewhat hairy leaves, set in pairs upon short pedicles: on the tops grow scaly heads of pale red labiated flowers, whose upper lip is entire and the lower cut into three segments, set in form of a convex umbel, intermixed with roundish purplish leaves: each flower is followed by four minute seeds inclosed in the cup. It is perennial, grows wild on dry chalky hills and gravelly grounds, in several parts of England, and flowers in June. The flowers, or rather flowery tops, of a somewhat different species, origanum creticum, were formerly brought from Candy, but have long given place to those of our own growth, which are nearly of the same quality.

THE leaves and flowery tops of origanum have an agreeable aromatic fmell, and a pungent taste, warmer than that of the garden marjoram, and much resembling thyme; with which they appear to agree in medicinal virtue. Insusions of them are sometimes drank as tea, in weakness of the stomach, disorders of the breast, for promoting perspiration and the sluid secretions in general: they are sometimes used also in nervine and antirheumatic baths; and the powder of the dried herb

as an errhine. Distilled with water, they yield a moderate quantity of Ot. effent. a very acrid penetrating effential oil, fmelling strongly of the origanum, but less agreeable than the herb itself: this oil is applied on a little cotton for eafing the pains of carious teeth; and fometimes diluted and rubbed on the nostrils, or snuffed up the nose, for attenuating and evacuating mucous humours.

OSTEOCOLLA.

OSTEOCOLL A Pharm. Edinb. (a) Ofteocolla, aliis offifragus, ofteites, ammosteus, osteolithos, bolosteus, stelochites, Worm. mus. Osteo-COLLA OF BONE-BINDER: a fossil substance, found in some parts of Germany, particularly in the marchè of Brandenburgh, and in other countries. It is met with in loofe fandy grounds, spreading, from near the furface to a confiderable depth, into a number of ramifications like the roots of a tree: it is of a whitish colour, soft while under the earth, friable when dry, rough on the furface, for the most part either hollow within, or filled with folid wood, or with a powdery woody matter (b).

This earth has been celebrated for promoting the coalition of fractured bones and the formation of a callus; a virtue to which it does not feem to have any claim. It is found to be composed of two different earthy fubstances, which are nearly in equal proportions, and which may be separated from one another, by washing the powdered ofteocolla with water: the finer matter, which washes over, appears from its burning into quicklime, and its properties in other experiments, to be mere calcareous earth, not different in quality from chalk: the groffer matter that remains is no other than fand.

OSTREUM.

OSTREUM Pharm. Lond. The OYSTER; a common, bivalvous, marine shell fish.

(a) Expunged.

(b) A more particular account of this fossil may be feen in Neumann's chemical works, p. 11. and the Memoires de l'academie royale des sciences de Berlin, pour l'ann. 1748.

Testæ ostreorumprepar. Pharm. Lond. THE shell of the oyster, levigated into a subtile powder, is employed as an absorbent, in heart-burns and other like complaints arising from acidities in the first passages: the hollow shells are generally made choice of, as containing more, than the thinner slat ones, of the fine white earth, in proportion to the outer rough coat, which last is found to be considerably impregnated with sea salt. By calcination, they are converted into a strong quicklime, which imparts to water a greater degree of lithontriptic power than the mineral limes; see Calx viva.

O V U M.

OVUM gallinaceum Pharm. Lond. HENS EGG.

Eggs are accounted very nutritious, but difficult of digestion, especially if boiled hard. In medicine, the yolk has been employed as an intermedium for rendering resinous juices and balsams soluble in water: it answers this purpose less effectually, and less elegantly than vegetable gums, the solutions obtained by means of the animal matter being apt on standing to become putrid or rancid. The yolk, exsiccated by a gentle warmth, forms a friable concrete; the white, a firm semi-transparent one, in appearance resembling amber or gum-arabic, and soluble again in watery liquors. The boiled white, placed in a moist cellar, deliquiates spontaneously, and gummy-resinous substances, included in it, dissolve along with it: preparations of this kind have been directed for medicinal uses, but it does not appear that more of the gummy-resin is thus dissolved by the liquamen of the egg than by simple water.

Testæ ovorum præpar. Pharm. Lond. The shells of eggs, freed, after boiling, from the inner skin, and levigated into fine powder, are sometimes used as absorbents, and supposed, when combined with the acid humours in the first passages, to be less disposed to bind the belly than most of the other testaceous powders.

PÆONIA.

PÆONIA Pharm. Lond. Pæonia folio nigricante splendido quæ mas, & pæonia femina flore pleno rubro majore C.B. Pæonia officinalis Linn.

MALE

Male and Female Peony or Piony: a plant with large leaves, divided deeply into oblong fegments, or rather composed of a number of these fegments set on divided pedicles: on the tops of the branches grow large rose-like flowers, followed each by two or more horned pods, internally of a deep red colour, containing roundish shining red or black seeds. The male sort has dark green leaves, pale red single flowers, long thick roots, and the stalks and pedicles streaked with red: the semale has longer narrower and paler leaves, deep red double flowers, and irregular roots composed of several tuberous pieces hanging by tough filaments from one head. They are both sound wild in some parts of Europe, and cultivated with us in gardens: they are perennial, produce their flowers in May, and very soon shed them.

THE male peony has been generally preferred for medicinal use: but the female, which is the largest and most elegant, and for this reason the most common, is the species which the shops have been principally supplied with. In quality, there does not appear to be any material difference betwixt the two; and hence the college allow both sorts to be taken indiscriminately.

The roots and feeds of peony have, when fresh, a faint unpleasant smell, somewhat of the narcotic kind; and a mucilaginous subacrid taste, with a slight degree of bitterishness and astringency. In drying, they lose their smell, and part of their taste. Extracts made from them by water are almost insipid as well as inodorous; but extracts made by rectified spirit are manifestly bitterish and considerably astringent.

The leaves are nearly inodorous. To the taste, the leaves themselves discover a moderate degree of roughness, and their pedicles of sweetness; both which are preserved in great measure in the watery, but more perfectly in the spirituous extracts.

The flowers have rather more smell than any of the other parts of the plant, and a rough sweetish taste, which they impart, together with their colour, both to water and spirit: the watery insusion leaves, on being inspissated, a blackish red, austere, sweetish, and somewhat bitterish extract: the spirituous tincture yields an extract of a beautiful bright red, of an agreeable though weak smell, a moderate astringency, and an almost saccharine sweetness.

The roots, flowers, and feeds, are looked upon as lightly anodyne and corroborant; to the latter, at leaft, of which virtues, they appear from the above experiments to have fome claim. They have been principally recommended in spasmodic and epileptic complaints; in which, we are afraid, their effects are not very considerable.

PALMA.

PALM: a tall unbranched tree, with long reed-like leaves elegantly disposed on the top. Different species of it grow spontaneously in the eastern countries, and in the warmer parts of the West Indies.

THE palma major C. B. Phænix dactylifera Linn. is cultivated in fome of the fouthern parts of Europe. Its fruit, the dates of the shops, is of an oblong shape, like an acorn, but generally larger; and confists of a thick fleshy substance including, and freely parting from, an oblong hard stone, which has a remarkable furrow running its whole length The best dates come from Tunis: they should be upon one fide. chosen large, foftish, not much wrinkled, of a reddish yellow colour on the outfide, with a whitish membrane betwixt the flesh and the stone. They have an agreeable fweet tafte, accompanied with a flight aftringency; and hence stand recommended in tickling coughs and thin acrid defluxions on the lungs, and in alvine fluxes. Among the Egyptians and Africans, they are faid to be a principal article of food, and when used too freely, to be difficult of digestion, occasion headachs, sometimes gripes, and, in length of time, obstructions of the viscera, cachectic, and melancholic diforders.

THE palma oleosa (palma foliorum pediculis spinosis, fructu pruniformi luteo oleoso Sloan. jam.) is a native of the coast of Guinea (a) and the Cape Verd islands, from whence it has been introduced into Jamaica and Barbadoes. From its fruit is extracted an oil, which, as brought to us, is about the consistence of an ointment, of a strong, not difagreeable smell, and scarcely any particular taste: by long keeping it

Oleum expressum palmæ Pb. Ed.

loses

^{*(}a) According to Bergius, another species of the oil palm grows on the coast of Guinea and in Senegal, the palma altissima non spinosa, fructu pruniformi minore, racemo sparso Sloan. Jam. & Adanson.

loses its high colour, and becomes white, and in this state is to be rejected. The inhabitants of the Guinea coast are said to employ the palm oil for the same purposes as we do butter. With us, it is only used in some external applications, for pains and weakness of the nerves, cramps, sprains, and other like complaints. The common people sometimes apply it to chilblains; and, when used early, not without benefit. It is said to be peculiarly serviceable in hardness of the belly, both of adults and children (a).

THE medullary part of certain oriental palm trees (palma indica caudice in annulos protuberantes distincto, fructu pruniformi, Raii. Sagus, seu palma farinaria Rumph. Amb. (b) affords another article of food to the natives, and of the materia medica to us. The farinaceous medulla, freed from the filamentous matter with which it is inveloped, is beaten with water, and made into cakes, which are afterwards reduced into fmall grains, and dried. The cakes are faid to be the bread used by the Indians in fcarcity of rice: the grains are the fago or fagou of the shops. This substance, commonly recommended as a restorative in phthises and emaciations and for restraining fluxions, appears to be a light, moderately nutritious demulcent food; in which view it is by some directed (c) as a proper aliment for young children, in preference to the more tenacious and less digestible preparations of wheat flour. It disfolves in water into a viscid mucilage; is less acescent and flatulent than other farinæ; keeps longer in the grain, even for twenty years in a dry place, and also in its mucilaginous state a long time (d).

PAPAVE R.

POPPY: a plant with oblong leaves, and round stalks, divided into a few branches, each of which is terminated by a large tetrapetalous flower, set in a two-leaved cup that falls off as the flower opens: the flower itself likewise soon falls, leaving a smooth roundish head or

⁽a) Bergii Mat. Med. 882.

^{*(}b) The Cycas circinalis Linn. has been given as the fago plant, but, as Bergius supposes, erroneously.

⁽c) Albertus Seba, Thefaur. vol. i. p. 40. Act. nat. curiof. vol. i. Append.

⁽d) Cullen, Mat. Med.

capfule, covered with a radiated crown, and containing a number of fmooth roundish seeds. It is annual, and flowers from June to near the end of summer.

- 1. PAPAVER ALBUM Pharm. Lond. & Edinb. Papaver hortense semine albo C. B. Papaver somniferum Linn. White poppy: with smooth, slightly indented leaves; and whitish flowers and seeds.
- 2. PAPAVER NIGRUM: Papaver hortense nigro semine C. B. Black poppy: a variety of the former, with smooth, slightly indented leaves, purple flowers, and black seeds.

THESE plants are found wild in some parts of Europe; and several varieties of them, in regard to the flowers, are produced by culture in our gardens. The heads, stalk, and leaves, have an unpleasant smell, and a bitterish biting taste, of the same kind with those of opium. fmell and tafte is lodged in a milky juice; which abounds chiefly in the cortical part of the heads; which may be collected, in confiderable quantity, by flightly wounding them when almost ripe; and which, on being exposed for a little time to a warm air, thickens into a tenacious darkcoloured mass, similar to the opium brought from abroad, but stronger in fmell and taste. The juices thus obtained from the two forts of poppies, appear to be of the fame quality, the difference being only in the quantity afforded: the white poppy, which is the largest, is the fort cultivated by the preparers of opium in the eastern countries, and for medicinal uses in this. * The following extract from Mr. Kerr's account of the culture of this plant, and the preparation of opium, in the province of Bahar in the East Indies, may convey useful information.

"The feeds are fown in October or November. The plants are allowed to grow fix or eight inches distant from each other, and are plentifully supplied with water. When the young plants are fix or eight inches high, they are watered more sparingly. But the cultivator strews all over the areas a nutrient compost of ashes, human excrements, cow-dung, and a large portion of nitrous earth, scraped from the highways and old mud-walls. When the plants are nigh flowering, they are watered pro-

fusely to increase the juice.

When

"When the capsules are half grown, no more water is given, and they begin to collect the opium. At fun-fet they make two longitudinal double incisions upon each half-ripe capfule, passing from below upwards, and taking care not to penetrate the internal cavity of the capfule. incisions are repeated every evening, until each capsule has received fix or eight wounds; they are then allowed to ripen their feeds. The ripe capfules afford little or no juice. If the wound was made in the heat of the day, a cicatrix would be too foon formed.—The night-dews, by their moisture, favour the exstillation of the juice. Early in the morning old women, boys, and girls, collect the juice, by scraping it off the wounds with a small iron scoop, and deposite the whole in an earthen pot, where it is worked by the hand in the open fun-shine, until it becomes of a confiderable spissitude: it is then formed into cakes of a globular shape, and about four pounds in weight, and laid into little earthen basins to be further exficcated. These cakes are covered over with the poppy or tobacco leaves, and dried until they are fit for fale. Opium is frequently adulterated with cow-dung, the extract of the poppy-plant procured by boiling, and various other substances which they keep in secrecy." (a)

The collection of the pure milky juice of the poppy has not, among us, been as yet practifed in large, or with a view to the supplying of the common demand of opium. Instead of this troublesome process, we extract the narcotic matter by menstrua; the active parts of opium, as observed under that article, being completely dissoluble both by water and rectified spirit. A portion of the herbaceous inert substance of the plant is indeed, at the same time, taken up, at least when water is made use of, so as to render an enlargement of the dose necessary: but this addition to the bulk of a dose of opium would be of no inconvenience, if the compound was always of the same strength, or the narcotic and inert matter in the same proportions to one another; a point which cannot be attained with so much precision as could be wished, but which may nevertheless, by due care in the preparation, be adjusted as nearly as common practice in most cases requires.

The college of London directs the dried heads, cut and cleared from the feeds, to be boiled in water, in the proportion of three pounds and a half to fix gallons, and now and then stirred to prevent their burning, till only about one third part of the liquor remains, which will be almost Syrup. e meconio five diacodion P.L.

Syrupus papaveris albi, feu de meconio, vulgo diacodion Ph. Edinb.

Extract. capitum papaveris albi Phar. Edinb. entirely soaked up by the poppies: the decoction is to be strongly pressed out, boiled down to about four pints, then strained whilst hot, first through a sieve and afterwards through a sine woollen cloth, and set by for a night to settle: to the liquor, poured off clear, six pounds of double-resined sugar are to be added; and the mixture boiled till its weight comes to nine pounds or a little more, that it may become a syrup of a proper consistence. An ounce of this syrup is reckoned equivalent to about a grain of opium. *The Edinburgh college directs two pounds of poppy heads without the seeds to be macerated for a night in thirty pounds of boiling water, the liquor then boiled down till only a third part remains, which is to be strongly expressed and strained, then boiled again to the half, strained, and made into a syrup with a sufficient quantity of sugar. They also allow this syrup to be made by dissolving one dram of the extract of white poppy heads in two pounds and a half of simple syrup.

A decoction of poppy heads in water, strongly pressed out, depurated by settling, then clarified with whites of eggs, and inspissated, yields an extract amounting to one fifth or one fixth the weight of the heads: it is said, that two grains of this preparation are equivalent to one grain of opium, and that the extract is not liable to produce a nausea or giddiness which generally follow the use of pure opium (a): but the consequential effects which opiates produce, in different subjects, and in different circumstances, are so variable, that the trials which have been made of this preparation, however successful, do not appear sufficient for establishing this superiority. Of tinctures or extracts made with spirituous menstrua, no medicinal trials, so far as I can learn, have as yet been made: in smell and taste they approach more to opium than any other preparation of the poppy I have seen.

Many have supposed the seeds of the poppy to be, like the other parts of the herb, narcotic (b); missed, perhaps, by analogical reasoning from other plants. Though the seeds of many plants are more efficacious than the vessel in which they are lodged; those of the poppy have nothing of the narcotic juice which is diffused through their covering,

(a) Mr. Arnot, Edinburgh medical effays, vol. v. art. 11.

⁽b) Hermann, Cynosur. mat. med. Edit. Boecler, p. 436. Juncker, Conspectus therapiæ generalis, p. 279.

through the stalks, and more sparingly through the leaves. If emulfions of poppy feeds have been found ferviceable in coughs, catarrhs, heat of urine, and other like diforders; it is not to an anodyne, but an emollient quality, that this virtue is to be ascribed. The seeds in substance have a sweetish unctuous farinaceous taste, and yield upon expression a large quantity of insipid oil: both the seeds themselves and the oil are faid to be in some places common articles of food (a).

3. PAPAVER ERRATICUM Pharm. Lond. Papaver erraticum majus C.B. Papaver Rhaas Linn. Wild or red poppy, or corn-rose: with deep red flowers, dark-coloured feeds, hairy leaves and stalks, and the leaves cut almost, or quite, to the pedicle into indented segments. It is common in corn-fields; and is fometimes, like the others, made to vary its flowers by culture.

THE heads of this species appear to contain the same kind of narcotic juice with those of the two preceding, but in so much smaller quantity that they are wholly neglected. The only part made use of is the flowers, which are supposed to be likewise impregnated in some degree with the fame anodyne principle, and stand recommended in catarrhs, coughs, spitting of blood, and other disorders: they have a slight narcotic smell, and a very mucilaginous taste, accompanied with a sensible bitterishness. They are at present regarded rather on account of their colour, than for any great virtues expected from them: they yield upon expression a deep red juice, and impart the same colour to watery liquors, and a brighter though paler red to rectified spirit. A strong infusion of them is prepared in the shops, by pouring four pints and a half of boiling water upon four pounds of the fresh flowers, stirring them over the fire till the flowers are all immerged, and fetting them by to steep for a night: without the application of fire fo as to scald or shrink the flowers a little, they can scarcely be moistened with the water; if the heat is continued longer than this effect is produced, the liquor turns out quite flimy. This infusion, Syr. papav. pressed out and depurated by settling, is reduced, by a proper addition of fugar, into a deep red fyrup. The colouring matter of the red poppy

errat. Ph. L.

⁽a) Prosper Alpinus, De medicina Ægyptiorum, lib. iv. cap. 1. Geoffroy, Mat. med. tom. ii. p. 715. Linnæi, Amanitat. Academic. iii. 71.

MATERIA MEDICA.

differs from that of clove-gilly flowers, red roses, and other bright red flowers, in this; that on the admixture of alkaline liquors, it does not change, like them, to a green, but to a dark purple.

PARALYSIS.

PARALYSIS: a plant with oblong wrinkled leaves, hairy on the upper fides of the ribs; and naked stalks, bearing monopetalous flowers, each of which is divided about the edge into five segments, and set in a loose tubulous, ridged cup, which, after the slower has fallen, incloses a husk full of roundish seeds. It is perennial, and slowers early in the spring.

1. PARALYSIS Pharm. Lond. Verbasculum pratense odoratum C. B. Primula veris major Gerard. Primula veris officinalis Linn. Cowslip, paigil, or peagle: with several flowers set together on one stalk, of a deep yellow colour, drooping downwards. It grows wild in marshes and moist meadows.

Cowslip flowers have a moderately strong pleasant smell, and a somewhat roughish bitterish taste; both which they impart, together with a yellow tincture, to watery and to spirituous menstrua. Vinous liquors, impregnated with their flavour by maceration or fermentation, and strong insusions of them drank as tea, are supposed to be mildly corroborant, antispasmodic, and anodyne. An insusion of three pounds of the fresh flowers in five pints of boiling water is made in the shops into a syrup, of a fine yellow colour, and agreeably impregnated with the slavour of the cowslips.

Syr. e flor. paralyfis Pharm. Lond.

2. PRIMULA VERIS Pharm. Edinb.(a) Primula veris minor Ger. Verbasculum silvestre majus singulari slore C.B. Primula veris acaulis Linn. Primrose: with pale yellow solitary slowers. It grows wild in woods and hedges.

THE flowers of this species are much weaker and less agreeable in smell than those of the preceding. The leaves and the roots seem to partake

in some degree of the nature of those of asarum; acting as strong errhines or sternutatories, when snuffed up the nose, and as emetics (the roots at least) when taken internally. Gerard reports, as from the experience of a skilful practitioner, that "a dram and a half of the powder of the dried roots (taken up in autumn) purgeth by vomit very forcibly, but safely, in such manner as asarum doth."

PAREIRA.

PAREIRA BRAVA Pharm. Edinb.(a) Pareyra, Ambutua, Butua, Overo brutua, Zan. hift. Pharm. Parif. Pareira brazilianus flore octopetalo monococcos Raii hift. Cissampelos Pariera Linn.) brought from Brazil, generally in crooked pieces of different fizes, some no bigger than the finger, others as large as a child's arm: the outside is brownish and variously wrinkled; the internal substance of a pale dull yellowish hue, and interwoven as it were with woody fibres, so that on a transverse section, there appears a number of concentric circles, crossed with strike running from the center to the circumference.

This root is extolled by the Brasilians and Portuguese in a variety of diseases, particularly in suppressions of urine and in nephritic and calculous complaints. Geosfroy is of opinion, that its virtue consists in dissolving and attenuating tenacious juices; and reports, that in sundry disorders arising from their viscidity, it was found remarkably beneficial: that in nephritic pains and suppressions of urine, he has often given it with happy success: that he has sometimes seen the patient freed from pain almost in an instant, and a plentiful discharge of urine brought on: that in ulcers of the kidneys and bladder, where the urine was mucous and purulent, and could scarcely be voided, or not without great uncafiness, the symptoms were soon relieved by pareira, and the ulcer at length healed by joining to it balsam of copaiba: that in an asthmatic case, where the patient was almost suffocated by thick phlegm, an infusion of pareira, after many other medicines had been tried in vain, brought on a copious expectoration, which proved a solution of the

(a) Expunged.

disease: that a person who, from an acute pain under the liver, had become in a few hours icterical, had the pain relieved, after bleeding, by the third cup of the decoction, and all the symptoms removed by a continuance of it; and that the same disorder frequently returning, she always found relief from the same medicine: but that in another icterical case, where the liver was swelled, it did no good. He cautions against giving too large doses, which might, he observes, raise a heat, and perhaps an inflammation in the kidneys: of the root in substance he prescribes from twelve grains to half a dram, and in decoction or infusion two or three drams; this quantity of the root, bruised, he directs to be boiled in a pint and a half of water till only a pint remains, which is to be strained off, sweetened with a little sugar, divided into three portions, and drank as tea at intervals of half an hour.

The use of this root has not been in general accompanied with so much fuccefs: but though, like many other medicines, it has not been found to answer the character at first given of it, and has thence fallen into neglect, we may prefume, from its fensible qualities, that it is not destitute of medical virtue. It has no remarkable smell; but to the taste it manifests a notable sweetness, of the liquorice kind, together with a confiderable bitterness and a flight roughness covered by the sweet matter. It gives out great part both of the bitter and the fweet substance to watery and spirituous menstrua: in evaporating the watery decoction, a confiderable quantity of refinous matter separates, which does not mingle with the remaining extract or diffolve in water, but is readily taken up by spirit; whence spirit appears to be the most perfect diffolvent of its active parts. Both the spirituous tincture and extract are in taste stronger than the watery.

PARIETARIA.

Parietaria officinarum & PARIETARIA Pharm. Lond. & Edinb. dioscoridis C. B. Parietaria officinalis Linn. PELLITORY OF THE WALL: a plant with tender reddish stalks; rough, uncut, oblong leaves, pointed at both ends; and imperfect rough flowers, growing in clusters along the stalks, followed each by a small shining seed. It is perennial, common on old walls and among rubbish, and flowers in May.

THE

The leaves of pellitory of the wall have been used in cataplasms for discussing inflammatory swellings: decoctions of them, and their expressed juice, have been given as emollient diuretics in nephritic cases and ischuries, and are said, when long persisted in, to be useful aperients or sweeteners in cutaneous defedations. The plant appears to be of no great activity, being rather oleraceous than medicinal: to the taste, the leaves in substance and their juice, are little other than herbaceous and watery.

PASTINACA.

PASTINACA: an umbelliferous plant, with naked umbels, yellow flowers, and flat feeds furrounded with a leafy margin: the leaves are oblong, and stand in pairs on a middle rib, without pedicles.

- 1. PASTINACA: Pastinaca latifolia sativa C. B. Pastinaca sativa Linn. Garden parsnep: with pale-coloured smooth indented leaves, and a large sleshy root.
- 2. ELAPHOBOSCUM: Pastinaca silvestris latifolia C. B. Bancia & branca leonina quibusdam. Wild parsnep: with dark green rough indented leaves and slender woody roots; common about the sides of fields; flowering, as the other, in June and July, and ripening its seeds in September. The garden fort is supposed to be only a variety of this, and to owe its differences to culture.

THE roots of the garden parsnep, in taste considerably sweetish, are accounted a very nutritious aliment: they yield with rectified spirit a very sweet extract, and in distillation with water a small portion of essential oil possessing the specific slavour of the roots. It is said that by standing in the ground for some years, it contracts pernicious qualities, so as to occasion disorders of the senses (a).

The feeds of the garden fort are fomewhat aromatic; those of the wild a little more so; of considerable smell, but no great pungency or warmth. By infusion, they impregnate water moderately with their

⁽a) Ray, Historia plantarum, i. 420. Dan. Hostman, Ada acad. cæsar. nat. curiosor. vol. vi. anno 1742. obs. 128. p. 426.

fmell, but communicate very little taste: in distillation they give over a small quantity of a pale yellowish essential oil, in taste moderately pungent, and smelling strongly of the seeds: sive pounds of the seeds of the garden parsnep yielded little more than a dram. Rectified spirit takes up by digestion the whole of their active matter, and carries off little in the inspissation of the tincture: the extracts of both sorts have a moderate warmth and bitterishness, differing in degree as the seeds themselves. These seeds have been commended as diuretics, similar to those of daucus, but weaker, which, in their sensible qualities, they apparently are: Haller reports, that those of the wild species, made into pills, with extract of liquorice, were much used by Boerhaave against nephritic complaints and ulcerations of the bladder.

3. Panax: Panax beracleum Morison. Panax pastinacæ solio C. B. Sphondylio vel potius pastinacæ germanicæ assinis panax vel pseudocostus store luteo J. B. Laserpitium Chironium Linn. Hercules's allheal or woundwort: with uncut leaves, somewhat heart-shaped, but having one of the sides lower than the other: the middle ribs, bearing the several sets of leaves, stand in pairs along a larger rib. It is a native of the warmer climates, and bears the colds of our own.

BOTH the feeds and the roots of this species are considerably warmer than those of the two preceding. The roots and stalks have a strong smell and taste resembling those of opopanax; and Boerhaave relates, that on wounding the plant in summer, he obtained a yellow juice, which, being inspissated a little in the sun, agreed perfectly, in both respects, with that exotic gummy-resin.

PENTAPHYLLUM.

PENTAPHYLLUM Pharm. Lond. Quinquefolium majus repens C. B. Potentilla reptans Linn. CINQUEFOIL, or FIVE-LEAVED GRASS: a trailing plant, with oval ferrated leaves, fet five together on long pedicles, and pentapetalous yellow flowers standing solitary on like pedicles: the cup is divided into ten unequal segments, the five innermost of which form a covering to a button of seeds: the root is long and slender.

flender, dark coloured on the outfide, and reddish within. It is perennial, grows wild on open clayie grounds, and flowers in June.

THE roots of pentaphyllum are mild aftringents, and give out their aftringent matter both to water and spirit. They have been used in diarrhœas and other fluxes, in intermitting severs, sometimes as corroborants and antiseptics in low colliquative acute severs, in gargarisms for strengthening the gums, &c. Their virtue is confined chiefly to the red cortical part, the whitish woody fibre in the middle being nearly insipid.

PERSICA.

PERSICA MALUS Pharm. Edinb. (a) Perfica molli carne vulgaris viridis & alba C. B. Anygdalus Perfica Linn. Peach: a tree common in gardens; with oblong, narrow, pointed, ferrated leaves; pale reddish flowers, composed of five broad petala with numerous stamina in the middle, set in five-leaved reddish cups, adhering to the branches without pedicles; and a sleshy fruit covered with downy matter and including a furrowed stone.

The flowers of the peach tree have an agreeable but weak smell, and a bitterish taste: Boulduc observes, that when distilled without addition, by the heat of a water-bath, they yield one sixth their weight or more of a whitish liquor, which communicates, to a considerable quantity of other liquids, a flavour like that of the kernels of fruits. These flowers appear to be gently laxative: it is said, that an insusion in water of half an ounce of the fresh gathered flowers, or of a dram of them when dried, sweetened with sugar, proves, for children, an useful purgative and anthelmintic; and that the leaves, more unpalatable than the flowers, are somewhat more efficacious. The fruit is of the same quality with the other dulco-acid summer fruits: see Fructus boræi.

PERSICARIA.

ARSMART: an annual plant with oblong uncut leaves pointed at both ends, and imperfect flowers fet in spikes on the tops of the stalks:

the cup is thick and fleshy, divided into five oval segments, which, closing, form a cover to an angular glossy seed.

1. Persicaria mitis: Persicaria mitis maculosa C. B. Pharm. Paris. Plumbago. Polygonum Persicaria Linn. Spotted arsmart; so called from most of the leaves having a blackish spot in the middle. It grows wild in moist watery places, and slowers in July.

This plant is faid to be a good vulnerary and antiseptic; and decoctions of it in wine, to restrain the progress of gangrenes (a). It has a slightly acerb taste inclining to acidity, and no remarkable smell.

2. Persicaria urens Pharm. Edinb. (b) Persicaria urens sive bydropiper C. B. Polygonum Hydropiper Linn. Biting arsmart, lakeweed, water pepper: distinguished from the former by the spikes of slowers being slenderer; the leaves shorter, narrower, and without any spots; but more remarkably by its taste. In our markets, a plant of a different genus, the second of the ranunculi hereaster described, is sometimes sold for it.

THE leaves of this species have an acrid burning taste, and seem to be nearly of the same nature with those of arum; their acrimony not rising in distillation, and being destroyed in the process (c). They are commended as antiseptic, aperient, diuretic; in scurvies and cachexies, humoural asthmas, hypochondriacal and nephritic complaints, and in the wandering gout. The fresh leaves have been sometimes applied externally, in stimulating cataplasms, and for cleansing soul ulcers and consuming sungous sless; in which last intention they are said to be used by the farriers.

⁽a) Tournefort, Memoires de l'acad. des scienc. de Paris, pour l'ann. 1703. (b) Expunged.

⁽c) Rutty, Synopfis of mineral waters, p. 524. Dr. Cullen however fays, "its acrimony operates chiefly on the kidneys. What is remarkable, it gives out its diuretic virtue in distillation to water." Mat. Med. 308.

PERUVIANUS CORTEX.

PERUVIANUS CORTEX Pharm. Lond. & Edinb. PERUVIAN BARK: the bark of a middling-fized tree, growing in Peru, called by the Spaniards, from its efficacy against intermitting fevers, palo de calenturas, or the fever tree; by Linnæus, Cinchona officinalis. This virtue of the bark is faid to have been discovered by the Indians about the year 1500, but not revealed to their European masters till 140 years after; when a fignal cure having been performed by it on the Spanish viceroy's lady, the countess del Cinchon, it came into general use in those parts, and was distinguished by the appellations pulvis comitissa, cortex china china or chinchina, kina kina or kinkina, and quina quina or quinquina. In 1649, a jesuit brought a large quantity of it into Italy, which was distributed by the fathers of that order, at a great price, in different parts of Europe: about the same time a quantity was purchased by cardinal de Lugo for the use of the poor at Rome. From these it received the names of cortex or pulvis jesuiticus, pulvis patrum, and pulvis cardinalis de Lugo.

This bark is brought to us in pieces of different fizes, some rolled up into short thick quills, and others flat: the outside is brownish, and generally covered in part with a whitish moss: the inside is of a yellowish, reddish, or rusty iron colour. The best fort breaks close and smooth, and proves friable betwixt the teeth: the inferiour kinds appear when broken of a woody texture, and in chewing separate into fibres. The former pulverises more easily than the latter, and looks, when powdered, of a light brownish colour, resembling that of cinnamon, but somewhat paler.

A bark was some time ago brought from America under the name of the semale Peruvian bark. This was sound, from experience, to be less effectual as a medicine than the genuine sort, which it was frequently substituted to or mixed with in France, insomuch that its importation, as the editor of Geoffroy informs us, was prohibited by law. It is considerably thicker, whiter on the outside, redder within, and weaker in smell and taste than the true bark.

PERUVIAN bark has a flight fmell, approaching as it were to mustiness, yet so much of the aromatic kind as not to be disagreeable. Its 3 T taste take is confiderably bitter, aftringent, very durable in the mouth, and accompanied with some degree of aromatic warmth, but not sufficient

to prevent its being ungrateful.

The febrifuge virtue, for which alone this medicine was at first recommended, has now been established by the daily experience of about a century: and that, when judiciously and seasonably administered, it proves as fafe as it is effectual, is now also beyond dispute. An emetic, which is in most cases necessary, being taken towards the approach of a paroxysm, that its operation may be over before the fit comes on; the bark is begun at the end of the paroxysm, or even in the time of the hot fit, and repeated, in doses of half a dram or more, every third or fourth hour, during the intermission: after the fever has been removed, the medicine is continued for a time, but more sparingly, to prevent a return. During the use of the bark, the pulse, which betwixt the paroxysms is generally weak and slow, becomes stronger and quicker, the appetite mends, the patient grows more cheerful, and perspiration increases: these may be looked upon as sure presages of its fuccess. At first it frequently occasions a looseness, and this also is falutary; but if the purging runs on too long, as the fever rarely yields while this evacuation continues, it is usually checked by the addition of a little opium: if too great costiveness ensues, recourse is had to glysters. In gross impure habits, gentle purgatives are premised to the bark, or given for a time in conjunction with it: in agues of the inflammatory kind, or accompanied with great heat, a little nitre is joined or interposed: in lax spongy constitutions, and a thin watery state of the blood, the bark is affisted by bitters, snakeroot, camphor, and chalybeates: where obstructions of the abdominal viscera are apprehended, it is not ventured on without the addition of fixt alkaline falts, fal ammoniac, or other aperients. In all cases, moderate exercise, and the drinking of warm liquids, promote its effects. As the bark is hurtful in the inflammatory diathefis, it is not near fo effectual in vernal, as in fummer and autumnal intermittents (a).

In remitting fevers, this medicine is less successful than in those which have perfect intermissions: in hectics, or wherever pus is formed, or juices are extravasated, it does harm. In the decline of long nervous

fevers or after a remission, and in those of the low malignant kind where the blood is colliquated and the strength exhausted, it proves an excellent cordial, corroborant, and antiseptic.

Peruvian bark has likewise been sound serviceable in gangrenes and mortifications, and in soul obstinate ulcers and running sores of other kinds: in these cases, taken in large and repeated doses, it frequently brings on a laudable suppuration, which degenerates on discontinuing the use of the medicine, and again turns kindly upon resuming it. The like effects have been observed from it in variolous cases, where either the pustules did not duly suppurate, or petechiæ shewed a disposition to a gangrene: by the use of bark, the empty vesicles filled with matter, watery sanies changed into thick white pus, and the petechiæ became gradually paler and at length disappeared. The principal symptom in this disease that contraindicates this valuable suppurant and antiseptic, is great obstruction at the breast or dissiculty of breathing; which are always by this medicine increased, insomuch that small doses have in some cases endangered suffocation.

In tumours of the glands, the Peruvian bark appears to promote, not fuppuration, but refolution. In the medical observations and inquiries published by a society of physicians in London, there are several instances of its being given with fuccess in scrophulous complaints. Dr. Fothergill observes, that inveterate ophthalmiæ generally yield to it: that beginning glandular tumours are very frequently refolved and their farther progress stopt by it: that swelled lips, cutaneous blotches arising from a like cause, are healed, and the tendency to a strumous habit corrected: that it does not succeed in all cases, but that there are few in which a trial can be attended with much detriment: that he has never known it to avail where the bones were affected, or where the scrophulous tumour was fo fituated as to be attended with much pain, as in the joints or under the membranous covers of the muscles; for when it attacks these parts, the periosteum, and consequently the bone, seldom escape being injured; that here the bark, instead of lessening, adds to the fever which accompanies these circumstances, and if it does not increase the force of the mischief, seems at least to hasten its progress.

Peruvian bark has been applied likewise, in conjunction with other appropriated medicines, and often with good success, to the cure of periodic headachs, hysterical, hypochondriacal, vertiginous and epileptic 3 T 2 complaints,

complaints, and other disorders that have regular intermissions. By its bitterness, astringency, and mild aromatic warmth, it strengthens the whole system, and proves a medicine of great utility in weakness of the stomach, uterine sluxes, and sundry chronical diseases proceeding from a laxity and debility of the sibres. To strengthen the solids appears indeed, in all cases, to be its primary operation; and its salutary virtues in different diseases, to be no other than consequential effects of this general power. In all the distempers where bark is known to take place, other astringent and bitter medicines, singly or combined, have likewise been of service, though not equally with this natural combination of them*(a).

The virtues of this bark are very difficultly extracted by long coction in water, and part of what the liquor is by heat enabled to take up begins to separate as soon as it is cold. This resinous part, which is rather melted out by the boiling heat than dissolved by the water as a menstruum, seems to contain chiefly the astringency of the drug: the bitter matter appears to be perfectly dissoluble, though more difficult to be got completely out. *After repeated insusion in cold water, till the liquor came off colourless and suffered no change from solution of vitriol, warm water extracted a considerable colour, and vitriol produced with this insusion an opake black: after warm water would extract no more, very hot water received a deeper colour than that of the strongest cold insusion of fresh bark; and this likewise struck a deep black with vitriol: boiling water had the same effect, after very hot water had ceased to act (b).

On boiling a pound of finely powdered bark for an hour or two in five or fix quarts of water, the decoction whilst hot looks clear and reddish, but in cooling becomes turbid and of a pale yellowish or wheyish hue: in this state it is found to partake, in a great degree, both of the bitterness and astringency of the bark, but in proportion as it deposites the matter that made it turbid, it loses more and more of its stypticity, the bitterness seeming to continue undiminished. The remaining bark, boiled in fresh water, exhibits the same appearances for two or three

^{*(}a) Dr. Percival found, that on mixing infusion of bark with putrid ox gall, an instant coagulation ensued, and the sector was increased. Hence he accounts for the disagreement of this medicine in the bilious severs of the West Indies. Ess. Med. and Exper. Vol. II. p. 24.

MATERIA MEDICA.

times fuccessively; and when, at length, it ceases to render the water turbid, it imparts a bitterness without astringency (a), retaining still fome share of bitterness itself. The vapour which exhales in the first coction being caught in proper veffels, condenses into a limpid liquor which smells strongly of the bark; though no separable oil is obtained on submitting many pounds to the operation. The several decoctions, Extr. cort. strained and inspissated together, yield an extract, rather less bitter, and much less styptic, than the bark in substance: this extract is kept in

peruv. molle & durum Ph.

(a) In the above experiments, I judged of the aftringency only from the tafte: folution of chalybeate vitriol, fo useful on other occasions for discovering aftringent matter in vegetable decoctions or infusions, seemed here to fail; for having often mixed it, in different quantities, with even the first decoctions of bark, it produced, not a black, but a deep green. I have fince observed, that when the vitriolic solution is used in very small proportion, it strikes a black with the turbid decoctions of bark, as with other aftringents; and that even the green mixtures, refulting from a greater addition of the vitriol, on being largely diluted with water, become black or blueish like diluted ink. The refinous matter, which subsides on standing from the turbid decoctions, being diffolved in spirit of wine, gave likewise a black with vitriol. But when the bark had been boiled in fresh waters, till it no longer gave any turbidness to the liquor, the last transparent decoctions, though fill pretty firong in tafte, gave no blackness at all.

Some doubts having arisen with regard to this experiment, I have repeated it twice, and found the event both times the same as before. The last decoctions, on dropping in the chalybeate folution, contracted indeed a flight dusky hue, which in certain positions might be mistaken for a low degree of blackness; but the mixtures, held between the eye and the light, appeared only of a kind of olive yellowish or brownish colour, and, on standing for a little while, deposited, not a black, but an ochery precipitate; whereas the first infusions or decoctions, though so far diluted with water as fearcely to difcover any tafte, ftruck a blueish colour like that of diluted ink, and

what little precipitate could be separated was black.

After the boiling of the bark in water had been repeated till the filtered liquor no longer made any change with folution of vitriol, the remaining bark gave no tincture at all to rectified spirit.

But fresh bark, boiled in successive portions of rectified spirit, till it ceased to impart any colour to the menstruum, gave still a deep tincture to boiling water; and this decoction, on the addition of folution of vitriol, exhibited nearly the same appearances as the last decoctions above mentioned, only in a higher degree, the precipitate being much more copious, and its colour

Though by repeated boilings in water the bark may be fo exhausted as to give out nothing to fpirit, but after the repeated action of spirit still gives out something to water; yet spirit appears to be the most active menstruum of its medicinal parts. For all, that spirit can dissolve, is extracted by a far less quantity of spirit than of water; and what spirit leaves undissolved is of little taste. Equal quantities of bark being digested for the same length of time with equal quantities of water and rectified spirit, with or without heat; the spirituous tinctures proved always fironger in tafte than the watery, and left on evaporation a larger proportion of extract * ||.

^{• ||} This affertion feems contrary to the refult of fome experiments by Dr. Percival, related in his first Vol. of Esf. Med. and Exper. p. 91, in which a dram of bark infused seven days in three ounces each of rectified spirit, proof spirit, and water, loft in the first, fix grains; in the second, eight and a quarter; and in the third, eight. These accounts can be reconciled only upon the supposition that watery liquors do, indeed, extract more of the inert gummy matter of bark; but spirituous, more of the active matter.

the shops in a foft and a hard form; the one of a proper consistence for

making into pills; the other fit for being reduced into powder.

As this drug gives out its virtue fo difficultly and imperfectly to boiling water, it has not been suspected that cold water would have any considerable action on it: I have nevertheless found, that an infusion in cold water, though perfectly transparent, is rather stronger in taste than even the turbid decoction, though the latter has somewhat more of a kind of fulness in the mouth (a). It is by means of a gummy matter in vegetables, that the refinous parts become diffoluble in watery liquors; and it feems probable that, in boiling, part of the gummy principle of the bark is haftily diffolved and difunited from the refinous, whereas cold water, acting more gradually, extracts them both together. I have given the infusions in intermitting fevers as well as other disorders, with all the fuccess that could have been expected from any preparation of this valuable medicine: the proportions commonly followed were, one ounce of the bark in fine powder, and eight or twelve of water, which were macerated without heat for twenty-four hours (b), and the clear liquor given in doses of two or three ounces.

It

⁽a) I have endeavoured to compare the strength of the two preparations by characters that may be thought more fatisfactory than the tafte. A cold infusion and decoction were made with equal quantities of bark and water, and both liquors passed through a filter: the infusion ran through faft; the decoction exceeding flowly, and continued turbid and opake after filtration. The two liquors, examined hydrostatically, were found very nearly of the same specific gravity. Equal quantities of them being turned black with equal quantities of folution of vitriol, the quantity of water necessary for diluting the blackness of the mixtures to an imperceptible degree, was very nearly the same for both. These experiments were often repeated, and seemed to prove, that the infusion and decoction are not considerably different in the quantity of matter taken up from the bark, but that this matter is in the cold infusion transparently dissolved, whereas in the decoction great part of it is only diffused through the liquor in an undiffolved state.-In the infufion itself, however, the solution does not appear to be very intimate. The transparent liquor becomes in a day or two turbid, and on standing for some weeks (being now and then shaken to prevent its growing mouldy) deposites so much of the resinous part, that it is in taste simply bitter, and produces no blackness with vitriol. The refinous sediment gives to spirit of wine a dark-coloured aftringent tincture, which strikes a black with vitriol like the tincture of bark itself.

⁽b) Since the above account was written, this preparation has been received in general practice, and found to answer the character here given of it. The time of maceration has been diminished to twelve hours, and some late experiments shew, that it may be still further reduced, without any injury to the medicine. A mixture of one part of bark and eight of water being siltered after standing for one hour, the liquor appeared, from its taste, from its colour, from its specific gravity, and from the trial with solution of vitriol, to be very nearly, if not fully, as strong,

It is a common opinion, that bark in substance is more effectual than any preparation of it. Thus much is plain, that the infusions, as well as the decoctions, have not near fo much effect as the quantity of bark they were made from, as the menstruum does not in either case completely extract its active matter: but their effects are evidently the same in kind, and the difference in degree may be compensated by an increase in the quantity.

The turbid decoctions, on the addition of any of the concentrated mineral acids, in the proportion of one drop to about a quarter of an ounce, become transparent, of a bright pale yellow colour, and of a rougher or more acerb taste, but with the loss of their bitterness: the vegetable acids, added in proportionably larger quantity, render them likewise transparent and improve their roughness, without much diminishing their bitterness: all these mixtures deposite, on standing, a little powdery fediment. Alkalies, both fixt and volatile, occasion a more copious precipitation, and instead of making the turbid decoctions clear, make the clear turbid.

Rectified spirit of wine receives from bark a deep reddish brown colour, and takes up much more of its active matter than watery liquors * (a): by digesting the powder first in some rectified spirit, and then boiling it in water, nearly the whole of its virtue is pretty readily got out. On inspissating the filtered tincture, the spirit carries off nothing remarkable of its smell or taste: the remaining extract retains the peculiar flavour of the bark, as well as its aftringency and bitterness, and proves a very elegant preparation, preferable to the pure refin obtained by precipitation from the tincture by water, as containing a part of the gummy matter, which is a medicinal principle of the bark as well as the refin. The spirituous tincture, and the decoction of the residuum, may be united Extr. cort. into an extract, possessing this advantage in a greater degree, by inspif- peruv. P. E. fating them feparately to the confishence of a fyrup, then mixing them together, and continuing the evaporation with a gentle heat.

frong, as those which had stood 2, 4, 6, 8, 12, 24 hours. On doubling the quantity of bark, and shaking it with the water for only two or three minutes, the liquor proved rather stronger than any of the preceding; and being afterwards kept 24 hours on the same bark, it gained no fensible addition to its strength. So that a very strong infusion may be obtained in a very expeditious manner.

^{*(}a) See Dr. Percival's different opinion, at the note in page 489.

†Tinct. cort. peruv. P. L. ‡ Ph. Edinb. Proof spirit extracts less from bark than rectified spirit, but more than water. Four ounces of the powder, macerated for some hours without heat, in a quart + or two pounds and a half ‡ of proof spirit, impart a considerable degree both of bitterness and astringency: on applying heat +, the taste becomes stronger, the colour darker, and the liquor somewhat turbid; from whence it may be concluded, that the resinous part is not by this menstruum completely dissolved.

Tinct. cort. peruv. vol. Pharm. Lond. Spirit of fal ammoniac made with fixt alkaline falt, by maceration with powdered bark in the above proportion, receives from it very little taste or colour. The spirit prepared with quicklime, and the dulcified spirit, extract in a few hours a very deep colour, and become strongly impregnated with its virtue. Though the spirit made with quicklime is held too acrimonious to be given internally by itself, it is not liable to that objection here; its pungency being sheathed by the substance which it dissolves.

Among the several substances which I have tried for covering the taste of bark, to some persons offensive, liquorice seemed to answer the best. Aromatics alone leave the taste of the bark very sensible in the mouth; but liquorice appeared to cover it effectually, whether in draughts or electraries, with the bark in substance or its preparations: to this compound any proper aromatic material may be superadded, to give a grateful slavour. For liquid forms, an infusion of the liquorice, and for electraries the extract should be used: for making up the electraries, mucilages are more proper than syrups, as the former occasion the compound to pass down freely without sticking about the mouth and sauces.

*Peruvianus cortex ruber: Red Peruvian bark. In the year 1779, a Spanish ship from Lima was taken by an English frigate, and carried into Lisbon. Her cargo chiefly consisted of bark, part of which was afterwards brought to London, and purchased by several druggists. From its large coarse appearance, it was some time before practitioners could be prevailed on to use it. At length, it was tried in some of the hospitals, and sound to be so efficacious, that an opinion soon prevailed of its being of a much superiour quality to the best common bark. Trials were multiplied throughout the kingdom, in a year when intermittents were remarkably frequent and obstinate; and its reputation increased with every experiment. Chemical tests were equally savourable to it,

as they proved it to contain a much greater proportion of active matter, than the other forts. At length, Dr. Saunders, a physician in London, eminent for chemical knowledge, published a treatise, in which various experiments on this bark were related, and attestations of its great medical efficacy from several practitioners were annexed. From this pamphlet, together with the editor's own experiments, the following account is extracted.

The red bark, as it is called, is in much larger and thicker pieces than the common. Most of the pieces are concave, though not rolled together, like the quilled bark. They break short, like the best common bark; and appear evidently composed of three layers. The outer is thin, rugged, frequently covered with a mostly substance, and of a reddish brown colour. The middle is thicker, more compact, and of a darker colour: it is very brittle and resinous. The innermost layer is more woody and sibrous, and of a brighter red. In powdering this bark, the middle layer, which seems to contain the greatest proportion of resinous matter, does not break so readily as the rest; a circumstance to be attended to, lest the most active part should be lest out of the sine powder.

This red bark to the taste discovers all the peculiar flavour of the Peruvian bark, but much stronger than the common officinal fort. An infusion in cold water is intensely bitter; more so than the strongest decoction of common bark. Its astringency is in an equal degree greater than that of the infusion of common bark, as is shewn by the addition of martial vitriol. The spirituous tincture of the red bark is also proportionally stronger than that of the pale. The quantity of matter extracted by rectified spirit from the powder of the former, was to that from the latter, as 3 to 2 in one experiment, and as 229 to 130 in another. And yet, on infusing the two residuums of the first experiment in boiling water, that of the red bark gave a liquor considerably bitter, and which struck a black with martial vitriol; while that yielded by the other was nearly tasteless, and void of astringency.

With respect to medical properties, from numerous and repeated trials it appears, that the red bark possesses the same virtues with the common, but in a much higher degree. A single half ounce of this has radically cured an obstinate intermittent, where many ounces of the other kind had either had no effect, or merely a temporary one.

3 U

Upon the whole, there is the strongest reason to conclude, with Dr. Saunders, that the red bark is the true Peruvian bark, of the best quality, or in its highest perfection. It was probably the kind of bark first introduced into Europe, and which acquired so much reputation in the hands of Sydenham and Morton. It is the fort still preferred by the Spaniards for their own use; and they are surprized at our preference of an inferiour kind. Whether it be, as Dr. Saunders first imagined, the bark of the trunk of full grown trees, the branches or young trees of which yield the pale bark; or whether the trees be different species, or, at least, varieties, does not feem accurately determined. The latter opinion is, perhaps, rendered the most probable, by an observation in the third edition of Dr. Saunders's pamphlet. He fays, that "he has lately feen fome exceeding good red bark imported by a Spanish merchant, a confiderable part of which was as fmall as the quilled bark in common use, yet still preserved its redness in that form, approaching, however, to the colour of cinnamon. It was extremely refinous, and gave evident proofs of its being the quill of the larger red bark which was in the same chest." This idea feems to be confirmed by fome curious remarks on the natural history of the cinchona, communicated by Dr. Simmons from the papers of the late M. Justieu, and subjoined to the same edition.

This writer makes feveral different species of bark, which may, however, be reduced to two. The first includes the red, the yellow, and the knotty barks, all of which have very smooth leaves, purplish slowers, with a bark that is bitter to the taste, and more or less coloured. Of these, the red is held in the highest estimation, and was that first imported into Europe, but is now become exceeding scarce, so that its place has been supplied by the yellow and knotty kinds. The second species includes the white barks, of which there are four varieties. All these have broad hairy leaves, and red, very odoriserous flowers, surnished with hairs on their inside. In two of these varieties the inner layers of bark are of a reddish hue. These have a slightly bitter taste, and somewhat of a febrifuge quality, which, however, they soon lose. The bark of the other two is quite white and insipid.

There have been lately discovered in the province of Santa-Fe, four degrees and a half north of the equator, two kinds of cinchona, one of which appears to be the same with the red bark of Peru; the other, one of the white species. This is a fortunate discovery, as it points out a

new store of this most valuable medicine, when the ancient ones shall be exhausted. We shall see in the next article, that our own settlements are not unprovided with a plant of the same genus, and similar virtues.

* CINCHONA CARRIBÆA Linn. Cinchona Jamaicensis Dris. Wright, Phil. Trans. Vol. LXXVII. Part II. This is a species of the jesuit's bark, produced in Jamaica and the Carribee islands, of which an accurate description, with an account of its virtues, has been published by Dr. Wright in the volume of Philosophical Transactions above referred to; and fome additions are made to this, in a letter from the same physician

to Dr. Duncan. Med. Comment. Vol. V. p. 398.

This tree, called in Jamaica the fea-fide beech, grows to the height of from twenty to forty feet. The outer bark of the large trees is white, furrowed, and very thick. This is inert, and may be knocked off from the inner. This latter is of a dark brown colour. Its flavour is at first fweet, with a mixture of the taste of horseradish and of the eastern aromatics; but when swallowed, it has that very bitterness and astringency which characterize the Peruvian bark. It yields its virtues both to cold and warm water; and a decoction of half an ounce of it boiled in a quart of water to the confumption of a pint, proved as strong as a decoction of an ounce and a half of the true bark. With the addition of orange peel it makes an elegant and grateful bitter tincture.

Its medicinal powers have been frequently tried by Dr. Wright, and it was found very efficacious in the dangerous remittent fevers of the West Indies, and also in nervous fevers. It has been administered in London in an intermittent, and effected a cure as completely as the Peruvian bark. From these accounts, we may hope that it will prove an useful and efficacious substitute for the cinchona of Peru, if ever the supplies of

this medicine should fail.

PETASITES.

PETASITES Pharm. Edinb. (a) Petasites major & vulgaris C. B. Galerita & tuspilago major quibusdam. Tuspilago Petasites Linn. But-TERBUR or PESTILENTWORT: a perennial plant, found wild by the

(a) Expunged.

fides of ditches and in meadows; producing early in the spring a thick naked roundish stalk, with a spike of small naked purplish sloseulous slowers on the top: the slowers and stalks soon wither, and are succeeded about May, by very large, roundish or somewhat heart-shaped leaves, standing on long pedicles, somewhat hollowed in the middle so as to refemble a bonnet (petasos): the root is long, thick, of a dark brownish or black colour on the outside, and white within.

The roots of butterbur are recommended as aperient and alexipharmac; and promife, though now difregarded in practice, to be of confiderable activity. They have a strong smell, and a bitterish acrid taste, of the aromatic kind, but not agreeable, very durable and diffusive, scarcely to be concealed, as Fuller observes, by a large admixture of other substances. Their virtue appears to reside in a resinous matter; which is distinguishable by the eye in the dried root, and which is readily extracted by spirit of wine.

PETROLEUM.

PETROLEUM, Oleum petræ, Oleum terræ. Rock oil: a fluid bitumen or mineral oil; exuding from the clefts of rocks or from the earth, or found floating on the furface of waters, in different parts of Europe, and more plentifully in the warmer countries; fimilar, in its general properties, to the oils extracted by distillation from pitcoal, amber, and other solid bituminous bodies. The more fluid petrolea have been distinguished by the name of naphtha; and the thicker by those of pissasphaltum and pisselæum.

- I. Petroleum album. White petroleum: nearly colourless; almost as clear, sluid, and transparent, as water; of a strong penetrating smell, not disagreeable, somewhat resembling that of rectified oil of amber. The principal, or only, part of Europe, in which it is sound, is the dutchy of Modena in Italy.
- 2. Petrolæum flavum seu italicum Pharm. Paris. Yellow petroleum: of a clear yellow colour; somewhat less sluid than the former; in smell rather less penetrating, less agreeable, and more nearly allied

allied to that of oil of amber. This also is found chiefly in the dutchy of Modena, and does not appear to differ very materially from the white fort.

3. PETROLÆUM RUBRUM seu gabianum, sive Oleum gabianum Pharm. Parif. Red petroleum: of a blackish red colour; of a thicker confiftence, and a less penetrating and more disagreeable smell, than either of the foregoing; found in Italy, and about the village Gabian in Languedoc.

There are many variations of these oils in regard to colour, fluidity, fubtility, and the pungency of their fmell and taste: the most fluid are in general the most subtile and pungent. Among us, the finer kinds are rarely to be met with; and even the inferiour forts are rarely unfo-

phisticated.

Fine petroleum catches fire on the approach of a flaming body, even without the contact of its substance with the slame; and burns entirely away. The hafty affusion of concentrated mineral acids, which raises a violent ebullition with distilled vegetable oils, and generally sets them on fire, makes no great conflict with petroleum: its confistence becomes thicker by this admixture, and its smell more fragrant. By distillation, it loses much of its natural fcent, and becomes fomewhat more pellucid than at first; a small quantity of a brownish or yellowish matter, similar to amber (a), remaining behind. Dropt on water, it spreads to a great distance, forming a various-coloured film on the surface. It floats also on rectified spirit of wine, and appears to be indissoluble in this menstruum; but unites with the essential oils of vegetables (b).

The finer petrolea, more agreeable than oil of amber, and more mild than that of turpentine, partake of the virtues of both. They have been fometimes taken internally in nervous complaints and as a diuretic; but used chiefly as an external stimulant, against rheumatic pains, palsies, chilblains, &c. In these intentions, some mineral oils, procurable among ourselves, are used by the common people, and often with benefit: the empyrical medicine, called British oil, is of the same nature with the petrolea; the genuine fort being extracted by distillation from a hard bitumen, or a kind of stone-coal, found in Shropshire and other parts of

England.

⁽a) Borrichius, Ada medica & philosoph. Hafniensia, tom. i. obs. 57.

⁽b) See l'Histoire & les memoires de l'acad. roy. des scienc, de Paris, pour les années 1715 & 1726.

4. Petroleum Barbadense Pharm. Lond. & Edinb. Bitumen barbadense. Pisselæum indicum. Barbadoes tar: of a reddish black colour, and a thick consistence, approaching to that of treacle or common tar. It is found in several of our American islands, particularly, as is said, in that from which it receives its name.

This bitumen, greatly esteemed by the Americans as a sudorific, in disorders of the breast, and as an external discutient and antiparalytic, is in smell more disagreeable, and both in smell and taste less pungent, than the foregoing petrolea. It is likewise less inflammable, and leaves on being burnt a considerable quantity of ashes. In distillation, it yields an oil different, in regard to its colour, from those afforded by such of the other bitumens as have been examined; appearing, when placed betwixt the eye and the light, of an orange colour, in other positions blue; but losing this variability of aspect in long keeping, and then looking in all situations yellow. This oil, and a balsam prepared by boiling the petroleum itself with one fourth its weight of slowers of sulphur, are directed by the London college to be kept in the shops.

Ol. petrolei barbadenfis Phar. Lond.

Balf. fulph. barbadenfe Phar. Lond.

PETROSELINUM.

PETROSELINUM VULGARE Ph. Lond. Petrofelinum Ph. Edinb. Apium bortense seu petroselinum vulgo C. B. Apium Petroselinum Linn. Parsley: an umbelliserous plant, with deep green winged leaves, of which those that grow on the stalk are divided into sine oblong narrow segments: the seeds are small; somewhat crookedly planoconvex, of a dusky greenish colour, with sour yellow ridges along the convex side; the root long, whitish, about the thickness of the singer. It is biennial, a native of moist grounds in the southern parts of Europe, and common in our culinary grounds.

The roots of parsley are sometimes used in apozems, and supposed to be aperient and diuretic, but liable to produce statulencies. Their taste is sweetish, accompanied with a slight warmth or slavour, somewhat resembling that of a carrot. Rectified spirit takes up, by digestion, all their active matter, and on inspissating the tincture, leaves it entire in the extract; in which, the sweetness is very considerable, the warmth very weak. Distilled with water, they impregnate the first runnings pretty

pretty strongly with their flavour: when large quantities are distilled, there separates a small portion, two or three drams from two hundred pounds, of essential oil, which partly swims on the water, partly sinks, and partly concretes about the nose of the worm into a butyraceous matter.

The leaves of the plant have a greater warmth and less sweetness than the roots. In distillation with water, they yield a greater quantity of essential oil, about ten drams from two hundred pounds, smelling agree-

ably of the herb, and in tafte moderately pungent.

The feeds, faid to be carminative, resolvent, and diuretic, and commended in the German ephemerides for destroying cutaneous insects in children, are in taste warmer and more aromatic than any other part of the plant, and accompanied with a considerable bitterness. In distillation, three pounds yielded above an ounce of essential oil, great part of which sunk in the watery suid. They give out little by insusion to watery menstrua, but readily impart all their virtue to rectified spirit: the tincture loses nothing considerable in being gently inspissated to the consistence of an extract, which proves a moderately warm, pungent, bitterish, not very grateful, aromatic.

PETROSELINUM MACEDONICUM.

PETROSELINUM MACEDONICUM Pharm. Lond. Apium macedonicum C.B. Apium petræum & petrapium quibusdam. Bubon macedonicum Linn. Macedonian parsley: differing from the foregoing, in the upper and lower leaves being alike, the stalks hairy and much branched, the seeds dark coloured and covered with a rough hoariness. It is a native of stony soils in Macedonia, and cultivated in some of our gardens.

THE Macedonian parsley is similar in quality to the common fort, but, weaker and less grateful. The seeds are the only part made use of, and these only as ingredients in the mithridate and theriaca: hence the Edinburgh college, having now dropt those compositions, has dropt also the Macedonian parsley.

PEUCEDANUM.

PEUCEDANUM Pharm. Edinb. (a) Peucedanum germanicum C. B. Pinastellum, fæniculum porcinum, fæniculum silvestre, marathrum silvestre, marathrum silvestre, marathrum silvestre, marathrum officinale Linn. Hogs fennel, Horestrong, Sulphurwort: an umbelliserous plant, with large leaves divided and subdivided tripartitely into sine oblong narrow segments: the seed is somewhat oval, slattish, marked with three striæ, and surrounded with a leasy margin: the root long and thick, with a tust of silaments on the top, blackish on the outside and pale coloured within. It is perennial, grows wild by the sea shores and in moist shady grounds, and slowers in July.

The roots of fulphurwort have a strong setid smell, somewhat resembling that of sulphureous solutions; and an unctuous, subacrid, bitterish taste. Wounded when fresh, in the spring or autumn, particularly in the former season, in which they are most vigorous, they yield a considerable quantity of yellow juice, which soon dries into a solid gummy-resin, retaining the taste and the strong smell of the root. This gummy-resin stands recommended as an aperient, and antihysteric.

PILOSELLA.

PILOSELLA, Myosotis, seu Auricula muris. Pharm. Paris. Pilosella major repens birsuta C.B. Hieracium Pilosella Linn. Mouse-ear: a low creeping hairy plant; with oval leaves, in shape like those of the daisy, joined to the stalks without pedicles, green above and white underneath: the flowers, which stand solitary on upright naked stalks, are composed of a number of yellow sloscules, set in scaly cups, and followed by small black seeds, winged with down. It is perennial, grows wild in dry pasture grounds, and slowers in June and July.

PILOSELLA is one of the bitterish lactescent plants. Its leaves differ from those of dandelion, cichory, and the other herbs of that class, in being much less juicy, less bitter, accompanied with some aftringency

which feems to prevail above the bitter, and a flight fweetishness very durable in the mouth: in the extracts made from them, both by water and spirit, the astringency is more manifestly the prevailing principle, though even when thus concentrated it is not very strong. The roots are considerably bitterer than the leaves, and less, if at all, astringent.

PIMPINELLA.

PIMPINELLA SAXIFRAGA Pharm. Lond. & Edinb. & Linn. Burnet-saxifrage: a perennial umbelliferous plant; with naked umbels; the outermost flowers composed of unequal petals, the inner equal; the seeds small, oblong, somewhat pointed, flat on one side, convex and striated on the other; the lower leaves roundish, indented, set in pairs along a middle rib with an odd one at the end; the upper leaves oblong and very narrow; the roots long, slender, and whitish.

- 1. PIMPINELLA ALBA Germanorum: Pimpinella saxifraga major umbella candida C. B. Greater or white burnet-saxifrage: with some of the leaves pretty deeply cut, the odd one into three sections. It is not very common in this country, and therefore our markets have been generally supplied with the following.
- 2. PIMPINELLA SAXIFRAGA: Pimpinella faxifraga minor foliis fanguisorbæ Raii; Tragoselinum alterum majus Tourn. Smaller burnet-faxifrage; with uncut leaves. It grows wild in dry pasture grounds.
- 3. PIMPINELLA SAXIFRAGA MINOR C. B. Tragoselinum minus Tourn. Small burnet-saxifrage; with the upper leaves divided into oblong narrow segments; taller than the others, but with smaller leaves. This is the most common fort in the fields about London.

ALL these plants appear to be possessed of the same qualities, and to differ little otherwise than in external appearance: and even in this, their difference is so inconsiderable and inconstant, that Linnæus has joined them into one species, under the name of pimpinella foliis pinnatis, foliolis radicalibus subrotundis, summis linearibus: he says he has seen the second fort produced from the seeds of the first sown in a richer soil. Instead of

3 X

the first, which has been generally understood as the officinal kind, our college allows either of the others to be taken indifferently.

The roots of pimpinella have a hot, pungent, not very durable tafte; and emit, when fresh, an acrid halitus, of no particular smell, but affecting the eyes like that of horseradish or mustard seed, though in a lower degree. In drying, they lose this subtile matter, and in long keeping the pungency of their taste is diminished. Their virtue is extracted, partially by water, and completely by rectified spirit. In distillation with water, a part of their pungency arises and impregnates the distilled shuid, and a part remains behind in the decoction: when large quantities are distilled, there separates from the water a small portion of a yellowish essential oil extremely acrid and siery. On inspissating the spirituous tincture, little or nothing of the virtue of the pimpinella rises with the spirit: the remaining extract, small in quantity, is of great pungency and heat. The leaves and seeds of the plant have likewise a considerable acrimony; the leaves less than the seeds, and both less than the roots.

This pungent root is in great esteem among the Germans, as a warm stimulating resolvent, aperient, diaphoretic, &c. in weakness of the stomach from viscid phlegm, infarctions of the breast, tumours and obstructions of the glands, impurities of the blood, and in general wherever tenacious humours are to be attenuated, or the fluid secretions promoted. It is an useful ingredient in our officinal compound arum powder, supplying in good measure the pungency which the arum root loses in being reduced into that form. It is employed also as a masticatory for stimulating the salival glands; and in gargarisms for dissolving viscid mucus in the sauces.

PINGUEDO.

PINGUEDO sive adeps: Sevum ovillum & bircinum, Axungia porcina & viperina. ANIMAL FATS: sheeps suet, goats suet, hog's lard, and vipers sat.

THE medical use of these substances is wholly external, as the basis of ointments and other unctuous applications. In their effects, they do not seem to differ materially from one another; all of them having one common emollient virtue, suppling and relaxing the part to which they

are applied, and obstructing its perspiration. The principal difference to be confidered in them is that of their confistence, by which they are adapted to different forms, or for receiving different admixtures; the folid feva ferving to give the thick confiftence of an unguent to oils and the more fluid refinous juices, while the fofter axungiae procure a like confistence to folid refins and powders. The fat of the viper is commonly preferred to the others in affections of the eyes; but its fuperiority, in these cases, to other soft fats, does not appear to have been fufficiently determined by experience. Nor indeed does it appear, that animal fats, and flavourless vegetable oils, of similar confistences, are materially different, respectively, from one another, in their effects when used in external applications. Even in regard to qualities, more remote than those, by which they can act when applied to the external parts of the body, the difference between the vegetable and animal fats is, perhaps, less than might be expected, and apparently less than that which is observed between the other corresponding substances of the two kingdoms, as the gelatinous matters of the one and the gummy of the other: animal fats, in their refolution by fire, yield neither the peculiar stench, nor the volatile alkaline falt, which fubstances completely animalized

Lard and fuet are directed to be tried or purified, by chopping them into small pieces; melting them by a gentle heat, with the addition of a little water, which fecures them from any danger of burning or turning curatioP.L. black, this fluid not being susceptible of a degree of heat sufficient for that effect; and then straining them from the membranes. Vipers fat, Axungia vifeparated from the heart, liver, and other bloody parts, is ordered to be melted without addition, and then strained through a linen cloth; the quantity of this fat, usually purified at a time, being so small, that the heat may be eafily regulated, fo as to prevent burning, without water.

Tried lard is formed into an elegant ointment, commonly called pomatum, by beating it with rofewater, in the proportion of three ounces of the water to two pounds of the lard, till they are well mixed; then melting it over a very gentle fire, and after standing for a little while, that the watery part may fettle, pouring off the lard, and inceffantly stirring and beating it about till it grows cold, fo as to reduce it into a light yielding mass; and afterwards adding so much essence of lemons as will be fufficient to give a grateful smell. Some scent it with oil of rhodium;

Axungiæ porcinæ fe-vique ovilli

perinæ curatio Ph. Lond.

Unguentum fimplex P.L. Pomatum rubrum Ph. Parif. and previously digest the lard for ten days with common water, renewing the water every day, a process which does not appear to be of much use. These ointments may be tinged of a fine red colour, for lip-salves, by a proper addition of alkanet root: the faculty of Paris directs, for this purpose, twenty-four parts of the white pomatum, eight of oxes marrow, and eight of white wax cut in small pieces, to be melted together by the heat of a water-bath; one part of powdered alkanet root to be added; the mixture stirred at times till it appears tinged of a deep red colour, and then strained through a linen cloth.

Animal fats are not dissoluble by spirit of wine any more than by water: when scented with essential oils, the oil may be totally extracted by digestion in rectified spirit, so as to leave the fat inodorous. By the same menstruum, fats may be freed from their ill smell, and even those that have grown considerably rancid by keeping may be made sweet again as at first; the rancidity and smell seeming to consist in a part of the fat attenuated, or subtilized, into a state analogous to that of the oil into which fats are resolved by distillation, which oil is totally dissoluble in spirit.

PINGUICULA.

PINGUICULA: Sanicula montana flore calcari donato C. B. Pinguicula sive sanicula eboracensis Gerard. Viola palustris, liparis, cucullata, & dodecatheon plinii quibusdam. Pinguicula vulgaris Linn. Butterwort or Yorkshire sanicle: a small plant, with a sew, oblong, obtuse, uncut, pale, glossy, unctuous leaves, lying on the ground; among which rise naked pedicles, bearing, each, a purplish monopetalous flower divided into two lips (of which the upper is cut like a heart, the lower into three sections) with a slender cylindrical spur or tail at bottom: the slower is followed by a roundish capsule full of small seeds. It is perennial, grows wild in elevated marshy grounds, and slowers in the spring.

THE remarkable unctuosity of this plant, and of some others of the same genus, seems to entitle them to a surther examination than has yet been bestowed upon them (a). It is said, that the unctuous and glutinous

juice of the *pinguicula* is used in some places as a liniment for chaps (a), and as a pomatum for the hair (b): that new milk, poured upon the fresh leaves, on a strainer, and after quick colature, set by for a day or two, becomes thick, tenacious, very agreeable and salubrious, and throws off no whey unless long kept; and that a little of the milk, so thickened, serves for bringing fresh milk to the same state (c): that a syrup made from the juice, and decoctions of the leaves in broth, are used among the common people in Wales as cathartics: and that the herb is hurtful to cattle that feed upon it (d).

PIPER.

PEPPER: the small, round, aromatic fruit of a trailing plant growing in Sumatra, Java, and Malabar, (Piper nigrum Linn.) The pepper-corns adhere in clusters to the stalks, without pedicles: when ripe, they are firm, not juicy, of a red colour, which changes in drying to a black.

- 1. PIPER NIGRUM Pharm. Lond. & Edinb. Melanopiper. Common or black pepper: the fruit gathered, probably, before perfect maturity, and dried in the fun.
- 2. PIPER ALBUM Pharm. Lond. Leucopiper. White pepper: the ripe fruit decorticated by maceration in water. Some of the grains, as brought to us, have pieces of a dark-coloured skin still upon them.

Or these pungent hot spices, the black fort is the hottest and strongest, and most commonly made use of for medicinal as well as culinary purposes. They both seem to heat the constitution more than some other spices that are of equal pungency upon the palate; and from those spices they differ in this, that their pungency does not reside in the volatile parts or essential oil, but in a substance of a more fixt kind, which does not rise in the heat of boiling water.

Pepper, infused in water, impregnates the menstruum pretty strongly with its slavour, but weakly with its taste: by boiling for some time,

⁽a) Simon Paulli, Quadripartit. botanic.

⁽c) Gifsler, Suenska vetenskaps academiens handl. 1749.

⁽b) Ray, Historia plantarum, i. 752.

⁽d) Gerard, Herbal emaculated, p. 789.

a little more of its pungent matter is extracted, and its flavour diffipated. On collecting the fluid that exhales in the boiling, the water is found agreeably impregnated with the odour of the spice, but scarcely discovers any taste: the essential oil, which rises to the surface of the water, thin, light, and limpid, in smell strong and agreeable, is in taste mild; a drop or two impressing on the tongue only a moderate grateful warmth. On inspissating the decoction, a part of the pungency of the pepper is found in the mucilaginous extract, and a part is retained by the pepper itself *(a).

Rectified spirit extracts completely the active matter of the pepper. The tincture is extremely hot and fiery, a few drops setting the mouth as it were in a slame: inspissated, it leaves an extract still more fiery; the spirit carrying off in its exhalation a little of the slavour, but nothing of the heat or pungency of the spice. The quantity of extract is nearly the same from both kinds of pepper; the spirituous amounting to about one eighth, and the watery to near one half their weight: but those of the white, like the spice in substance, are weaker than those of the black fort.

PIPER LONGUM.

PIPER LONGUM Ph. Lond. & Edinb. Macropiper Ph. Parif. Piper longum orientale C. B. Long pepper: the fruit of an East Indian plant of the same genus with that which produces the black pepper, (Piper longum Linn.); of a cylindrical figure, about an inch and a half in length, having numerous minute grains disposed round it in a kind of spiral direction.

This spice is hotter and more pungent than either of the preceding kinds, and its spirituous extract is proportionably more siery. In pharmaceutic properties, it entirely agrees with them; its active matter being only partially dissoluble in watery menstrua, and its pungency not rising in the heat of boiling water. Decoctions of it are very mucilaginous, rather more so than those of the black or white.

^{*(}a) The acrid matter of pepper is fo strongly retained, that a quantity boiled successively in fresh parcels of water, had not lost all its taste till the forty-third boiling. Gaubii Adversar.

PIPER JAMAICENSE.

PIPER JAMAICENSE Pharm. Lond. Pimenta five piper jamaicense Pharm. Edinb. Amomum Pharm. Wirtemb. JAMAICA PEP-PER, PIMENTO, ALL-SPICE: the dried aromatic berry of a large tree growing in the mountainous parts of Jamaica, reckoned a species of myrtle, and called by Sir Hans Sloane myrtus arborea aromatica foliis laurinis, by Linnæus myrtus (pimenta) foliis alternis.

PIMENTO is a moderately warm spice, of an agreeable flavour, somewhat refembling that of a mixture of cloves, cinnamon, and nutmegs. Distilled with water, it yields an elegant essential oil, so ponderous as to fink in the aqueous fluid, in taste moderately pungent, in smell and flavour approaching to oil of cloves, or rather a mixture of those of cloves and nutmegs: the remaining decoction, inspissated, leaves an extract somewhat ungrateful but not pungent, and the berry itself is now found to be almost wholly deprived of its taste as well as flavour; the warmth of this spice residing rather in the volatile than in the fixt parts. To rectified spirit it imparts, by maceration or digestion, the whole of its virtue, together with a brownish green tincture: in distillation, it gives over nothing confiderable to this menstruum, nearly all its active matter remaining concentrated in the inspissated extract; which is very warm and pungent, but not of a fiery heat like those obtained from the foregoing forts of pepper.

This spice, at first brought over for dietetic uses, has been long employed in the shops as a succedaneum to the more costly oriental aromatics: from them it was introduced into our hospitals, and is now received both in the London and Edinburgh pharmacopæias. A simple water is directed to be distilled from it, in the proportion of a gallon or ten pounds from half a pound: this is strongly impregnated with the Aq. piperis flavour of the pimento, though it is less elegant than the spirituous water phar. Lond. which the shops have been accustomed to prepare, by drawing off two Phar. Edinb. or three gallons of proof spirit from the same quantity of the spice. The Aq. piper. Edinburgh college directs only nine pounds from this quantity. effential oil does not feem to be much known in practice; though Ol. effent. it promises to be a very useful one, and might, doubtless, on many pip. jamaic. occasions, supply the place of many of the dearer oils. The quantity

of oil afforded by the spice is very considerable: Cartheuser indeed says, that only about half a dram is to be got from sixteen ounces; a mistake, which probably has arisen from inadvertence in copying Neumann's proportion, of half a dram from an ounce, or one sixteenth: so large a proportion as this last cannot, however, be collected in its proper form, the oil that remains dissolved in the distilled water being here included.

PIPER INDICUM.

PIPER INDICUM Pharm. Edinb. Capficum Pharm. Parif. Piper indicum, brazilianum, guineense, calecuticum, hispanicum, & lustanicum, quibusdam. Capsicum siliquis longis propendentibus Tourn. Siliquastrum plinii J. Bauh. Capsicum annuum Linn. Capsicum or Guinea pepper: long, roundish, taper, bright red pods, divided into two or three cells sull of small whitish seeds: the fruit of an annual plant, with square stalks, oblong acuminated leaves, and white slowers growing in their bosoms divided into sive segments in form of a star; a native of the East and West Indies, and raised in some of our gardens.

This fruit, when fresh, discovers to the organs of smell, a penetrating acrid halitus, which in drying is dissipated: its taste, whether fresh or dry, is extremely pungent and acrimonious, setting the mouth as it were on fire, and producing a painful burning vellication of long continuance, like that occasioned by arum root, but more of the warm aromatic kind. It gives out its pungency to rectified spirit, together with a pale yellowish red tincture: the spirit, gently distilled off, has no considerable impregnation from the capsicum: the remaining extract is insupportably fiery.

Capficum is fometimes given, in minute quantities, as one of the highest stimulants, in cold sluggish phlegmatic temperaments, in some paralytic cases, in relaxations and insensibility of the stomach, and for promoting the efficacy of aloetic medicines and the deobstruent gums in uterine disorders. It is used principally at table: a species of it, called in the West Indies bird-pepper, is the basis of the powder brought from thence under the name of Cayenne pepper. It is observable that this fruit, perhaps the strongest of the aromatic stimulants, is used freely,

as is faid, by the natives even of the warm climates: possibly these pungent antiseptic kinds of substances may there be more salubrious than they are, in general, among us, as they seem qualified to resist or correct the putredinous colliquation of the humours which immoderate heat produces.

pleisung of PIX.

PIX LIQUIDA Pharm. Lond. & Edinb. TAR: a thick, black, refinous, very adhefive juice; melted out by fire from old pines and fir-trees. The trees, cut in pieces, are inclosed in a large oven, which being heated by a fire on the outside, or the wood itself kindled and smothered, the juice runs off by a canal at the bottom.

TAR differs from the turpentine or native refinous juice of the trees, in having received a difagreeable empyreumatic impression from the fire; and in containing, along with the pungent bitter terebinthinate matter, a portion of the acid which is extricated from the wood by the heat, and likewise of its gummy or mucilaginous matter. By the mediation of these principles, a part of the terebinthinate oil and resin becomes dissoluble in watery liquors, which extract little or nothing from the purer turpentines.

Water impregnated with the more foluble parts of tar has been recommended as a remedy for almost all diseases. The proportions that have been commonly followed are, two pounds of tar to a gallon of water; which are to be well stirred together, then suffered to settle for two days, and the clear liquor poured off for use. It is observed, that " tar-water, when right, is not paler than French, nor deeper coloured "than Spanish white wine, and full as clear: if there be not a spirit "very fenfibly perceived in drinking, the tar-water is not good. It " may be drank either cold or warm. As to the quantity, in common "chronical indispositions a pint a day may suffice, taken on an empty "flomach, at two or four times: more may be taken by ftrong " stomachs. But those who labour under great and inveterate maladies, "must drink a greater quantity, at least a quart every twenty-four "hours. In acute diseases, it must be drank in bed warm, and in great " quantity (the fever still enabling the patient to drink) perhaps a pint 3 Y " every "every hour." Though this medicine is undoubtedly very far inferiour to the character that has been given of it, it is apparently capable of answering important purposes, as a deobstruent balsamic solution, moderately warm and stimulating. It sensibly raises the pulse, and increases either perspiration or the grosser evacuations. I have been informed of some late instances of its good effects in disorders of the leprous kind.

Some have imagined the acid to be the principle that gives virtue to tar-water; and hence have endeavoured to introduce, instead of the infusion, an acid spirit extracted from tar by distillation. But the effects of this, as of other acids, are opposite to those experienced from tarwater; nor does the acid of tar differ from that which is extricated by fire from all kinds of recent wood. Tar-water, distilled, yields a liquor very considerably impregnated with its slavour, though more grateful than the insusion itself both in smell and taste: there remains a light, spongy, blackish substance, not acid but bitter, partially dissoluble again in water.

Pil. piceæ Nofocom. Ed.

Unguent. e pice + Ph. L.

t Pbar . Edin.

This juice is fometimes given also in substance, mixed with so much powdered liquorice, or other like powdery matters, as is sufficient to render it of a due consistence for being formed into pills. An ointment, made by melting it with an equal weight of mutton suet, and straining the mixture whilst hot +, or by melting together sive parts of tar and two of yellow wax +, is sometimes used as a digestive, and said to be particularly serviceable against scorbutic and other cutaneous eruptions.

On inspissating tar, or boiling it down to dryness without addition, it gives over an acid liquor in considerable quantity, and an ethereal oil of the same general nature with that of turpentine, but impregnated with the empyreumatic slavour of the tar. The solid residuum is the common pitch, pix arida Ph. Lond. Pix sicca, palimpissa dioscoridi C. B. This is less pungent, and less bitter than the liquid tar, and used only in some external applications, as a warm adhesive resinous substance. Neumann observes, that when melted with oils, resins, and fats, into ointments and plasters, the pitch is greatly disposed to separate and precipitate.

PLANTAGO.

PLANTANE: a small perennial herb, common in fields and by road fides; with the leaves lying on the ground; and naked unbranched stalks, bearing on the top a spike of small imperfect four-leaved flowers, followed by little capfules, which, opening horizontally, shed numerous crooked feeds.

- I. PLANTAGO Pharm. Edinb. Plantago latifolia finuata C. B. Plantago septinervia. Plantago major Linn. Common greater plantane: with oval leaves, having feven ribs, prominent on the lower fide, running from end to end; and long flender spikes.
- 2. PLANTAGO MINOR seu quinquenervia: Plantago major angustifolia C. B. Plantago lanceolata J. B. Plantago lanceolata Linn. Narrow-leaved plantane or ribwort: with oblong, five-ribbed leaves; and short thick spikes.

THE leaves and feeds of plantane, recommended as vulneraries, in phthifical complaints, spittings of blood, alvine fluxes, &c. appear to be of the milder kind of restringents or corroborants. The leaves, of no remarkable smell, are in taste slightly acerb: their expressed juice, depurated by fettling and colature, or clarified with white of eggs, and inspissated to the consistence of honey, discovers a considerable saline austerity. The two forts are not sensibly different in quality from one another, though the first has been generally directed for medicinal use in preference to the other. The leaves are, in some places, the usual application made by the common people to flight wounds.

PLUMBUM.

PLUMBUM Pharm. Lond. LEAD: a pale, livid, foft, very flexible metal: above eleven times specifically heavier than water; fusible in a small heat, somewhat less than that in which expressed oils begin to boil. Continued in fusion it contracts a various-coloured pellicle on the furface, and if kept stirring, fo as that fresh surfaces may be exposed to the air, it changes by degrees into a powdery dusky-coloured calx:

3 Y 2

+ Minium Pharm. Lond.

† Lithargy-

calx: this powder, calcined for some time in a stronger fire, in such a manner that the flame may reverberate all over it, becomes first yellow, and afterwards of a deep red colour +: all these calces, if the fire be haftily raised to a considerable degree, melt into the appearance of oil, and on cooling form a foft flaky pulverable substance called litharge t, of a pale yellowish or reddish colour, according as the lead has been less or more calcined: if the calces be urged with a pretty strong fire, they run into a yellowish glass, which, while in fusion, powerfully dissolves most kinds of earthy bodies, and corrodes the common crucibles till it has faturated itself with their earth.

The ores of lead, in colour commonly refembling lead itself, and of a cubical or parallelopipedal structure, are plentiful in England and other parts of the world. The metal, extracted from the ore by fusion, contains frequently a portion of filver, and fometimes of gold: on keeping the compound melted in a due degree of heat, the lead calcines and turns to litharge, which is raked or blown off till the noble metals remain pure; all the other common metallic bodies being scorified and carried off by the lead. From the works, wherein filver is thus extracted from lead in the large way, the shops are supplied with litharge; which, when pale coloured, is called litharge of filver; when high coloured, litharge of gold. The latter is to be preferred, not as containing any of the metal by whose name it is distinguished, but as being more thoroughly calcined than the pale fort: the pale may be freed from the uncalcined lead it holds, by melting it; the uncalcined part falling to the bottom during the fusion.

The nitrous acid, diluted with about an equal quantity of water, diffolves lead pretty readily into a gold-coloured liquor: by the vitriolic and marine acids it is very difficultly acted on; and when previously diffolved in the nitrous, it is by either of these precipitated. Vegetable acids, digested on lead in substance, dissolve it exceeding sparingly: by certain managements they may be made to act more vigorously, and

to fatiate themselves with the metal.

Thin plates of lead, suspended over vinegar in a proper vessel, and set to digeft in a gentle heat, as that of horsedung, that the acid vapour may rise and circulate round the plates, are found, in about twenty days, covered with a white powdery or flaky matter: this being scraped off, and the process repeated, the whole of the metal is thus corroded by degrees

Ceruffa Pb. Lon. & Edin.

degrees into ceruffe or white lead. This commodity, the preparation of which makes a confiderable trade, is frequently adulterated with a mixture of whiting: the entire flaky masses, called flake lead, should be chosen, as not being liable to abuse. The adulteration may be discovered by means of vinegar, which will effervesce with and disfolve the whiting or calcareous earth: the liquor being then poured off clear, or filtered, the addition of a little spirit of falt will precipitate such part of the lead as the vinegar may have taken up; after which the calcareous earth will manifest itself on adding a little vitriolic acid.

The calces of lead are much easier of solution in vegetable acids than lead in its metallic form. On digesting four ounces of litharge about three days in a fand heat with a pint of strong vinegar, and now and then shaking the vessel; the liquor, filtered, is found to have received a Acetum listrong impregnation from the litharge, and to have disfolved about one tenth of it, whereas, of the same quantity of lead in substance, scarcely one hundredth part would be diffolved. Lead even in its vitreous state, or in the glazing of the common earthen-ware veffels, is confiderably acted on by vegetable acids; which, by being boiled in those vessels, receive from them the peculiar tafte, and pernicious qualities of faturnine folutions.-Lead may be discovered in acid liquors by a reddish, brown, or blackifh colour being produced in them on adding a few drops of a folution of orpiment or common fulphur made in lime-water, and by the colour not being destroyed on the superaddition of a little spirit of falt (a): other metals, diffolved in vegetable acids, produce, as well as lead, a dark colour with the fulphureous folutions, but spirit of falt redissolves them, and totally discharges the colour.

Of all the faturnine calces, the ceruffe, on account of the corrofion it has previously undergone from the steam of vinegar, is the most easily diffoluble in fresh vinegar, and hence is made choice of where a saturated folution is required. The folution made in vinegar, inspissated to the Saccharum confistence of honey and set in the cold, shoots by proper management into crystals, called, from their taste, fugar of lead. All the solutions, Sal plumbi and soluble preparations of this metal, have a remarkably sweet taste, sulgo facch. mixed with a confiderable aufterity.

thargyrites.

faturni P. L.

⁽a) The brownish or reddish colour produced by alkalies in cyder impregnated with lead, is totally discharged by spirit of falt. M. S. of Dr. Lewis.

† Tinct. faturnina P.L. † Tinct. faturnina vulgo antiphthifica Phar. Edinb.

LEAD in its metallic form, or when calcined by fire, does not appear to have any medicinal operation: diffolved or rendered foluble by acids, it is one of the most powerful styptics, but at the same time, for internal uses, one of the most dangerous. A few grains of the sugar have been ventured on for checking obstinate hemorrhagies and other profuse evacuations: a tincture drawn with rectified spirit, by maceration without heat, from fugar of lead and green vitriol, in the proportion of two ounces of each of the falts reduced separately into powder+, or three ounces of the fugar and two of the vitriol 1, to a quart of spirit, has been given from fifteen to thirty drops, for reftraining the colliquative sweats attending phthises and hectic fevers. This practice has in some instances been fuccessful, but the hazard is very great: all the faturnine preparations that have any activity are in a peculiar manner injurious to the nervous fystem, and ought never to be ventured on but in desperate cases as a last resource. Obstinate constipations, violent colics, pains and contractions of the limbs, tremors and refolutions of the nerves, and flow wasting fevers, are the general consequences of saturnines taken in any confiderable quantities internally, and of the fumes to which the workmen are exposed in the fusion of the metal in the way of business (a). Externally, this metal and its preparations are of fufficient fafety and

Empl. commune + P.L. † Pb. Edinb.

Empeminio Pharm. Lond.

of great utility. The plaster, in general use for slight cutaneous injuries, and which makes the basis of several other plasters, is a solution of litharge in oil olive, in the proportion of sive pounds of the litharge, subtilely powdered, to eight pints to ten pounds to the oil. The union is effected by boiling them together over a gentle fire, with the addition of about a quart of water to prevent their burning, and keeping them continually stirring, till they incorporate and acquire a due consistence: if all the water should be consumed before this happens, some more water, previously made hot, is added. A red plaster is prepared in the same manner with minium instead of litharge, but as it does not stick so well as the other, it is more rarely used: it is likewise more difficult of preparation, the compound being very apt, though a considerable quantity of water be used, to burn and grow black in the boiling.

⁽a) Vide Hoffman, Philosophia corp. human. morbosi, P. II. cap. viii. §. 20. & seq. Hundert mark, Acta acad. casarea nat. curios. vol. vii. Append. p. 96.

The ceruffe and fugar, particularly the latter, are cooling, drying, and astrictive: the fugar is used in collyria for inflammations and defluxions of the eyes, and in injections for restraining simple gonorrheas; and both preparations in unguents and liniments, against cutaneous heats and excoriations, flight ferpiginous eruptions, and for anointing the lips of wounds or ulcers that itch much or tend to inflammation. Compositions Ung. e cefor these purposes are made in the shops, by mixing one part of cerusie with five of the fimple ointment made with oil and wax; by grinding two ounces of litharge, and adding, alternately and by little and little, two ounces of vinegar and fix of oil+; or by boiling and stirring, over a + Ung. nugentle fire, four ounces of the common plaster, with one of vinegar, and two of oil where a thick unguent is required t, or four of oil for a fofter t Ung. triliniment | : this last is a less troublesome method of uniting the litharge with the oil and vinegar, than trituration; and the composition proves likewise more smooth and uniform, and less liable to grow hard in keeping *(a). But the most elegant and effectual of all the saturnine unguents, are those made with the sugar; in the proportion of half an ounce § to a pint of oil and three ounces of white wax; or one part, to twenty parts of the fimple oil and wax ointment §§ *(b).

* Mr. Goulard, a furgeon of Montpellier, has been the means of greatly extending for fome years past the external use of lead. The basis of his preparations is what he calls the extract of lead, or a folution of litharge in strong vinegar, boiled down to almost a syrupy consistence. This, diluted in a large quantity of foft water, makes his vegeto-mineral water, which is employed as a lotion or fotus, or boiled with bread to make a cataplasm. The extract is likewise combined with unguentous matters into a variety of forms. These preparations have, in fact, been found of the greatest utility in various cases of inflammation, particularly of the eryfipelatous kind, and the confequences of burns and scalds. most liberal application has not, in the opinion of most practitioners, been observed to produce any of those affections of the nervous system, which characterize the poisonous effects of lead taken internally. At the

ruffa, vulgo album P. E.

tritum.

pharmacum Pharm.Lond. || Lin. tripharmacum Pharm. Lond.

Ung. faturni. & Ph. Lon.

§§ Ph. Edin.

^{*(}a) The ung. nutritum, made without heat, though now expunged from our dispensatories, is much the best of the above preparations, and a very excellent application in many cases. It should not be long kept, but made fresh as wanted.

^{*(}b) These are by no means the efficacious preparations here represented. The oil and wax so cover the metallic falt, that its action is prevented; or, if it acts at all, it proves highly stimulating from the undiffelwed flate in which it is applied.

fame time, the abuse of saturnine applications, on the ground of those salse and inconsistent ideas of their action which Mr. Goulard has supported, has not infrequently been attended with disagreeable consequences.

POLIUM.

POLIUM Pharm. Lond. POLEY-MOUNTAIN: a small shrubby plant, with square stalks, oblong woolly leaves set in pairs; and labiated slowers wanting the upper lip and having the lower divided into five segments.

- 1. Polium maritimum erectum monspeliacum C. B. Teucrium capitatum Linn. Poley-mountain of Montpellier; with the leaves indented towards the end and joined to the stalk without pedicles, the slowers white and set in roundish spikes on the tops of the branches.
- 2. Polium angustifolium creticum C. B. Teucrium frutescens, stæchados arabicæ folio & facie Tourn. Teucrium creticum Linn. Poley-mountain of Candy; with the leaves not indented and set on short pedicles, the slowers standing in loose clusters, each on a separate soot-stalk.

SEVERAL other species, or varieties, of polium, erect and procumbent, with white, yellow, and purplish flowers, have been received in the shops. The second above described has been commonly understood as the true officinal fort, and procured dry from the island Candy, of which it is a native: the first, which better bears the winters of our own climate, appears to be of the same quality; and hence the college allow either fort to be taken indifferently.

THE leaves and tops of poley-mountain have a moderately strong aromatic smell, and a disagreeable bitter taste: distilled with water, they yield a small quantity of a yellowish essential oil, of a pungent taste, in smell less agreeable than the herb itself; the remaining decoction, inspissated, leaves a strongly bitter extract. They stand recommended as corroborants, aperients, and antispasmodics; but are at present scarcely otherwise made use of than as an ingredient in mithridate and theriaca.

POLYGALA.

MILKWORT: a fmall perennial plant; with the leaves alternate, uncut, and those on the upper parts of the stalks larger than on the lower; the flowers irregular, tubulous, tripetalous, labiated, set in loose spikes on the tops; the cup composed of sive leaves, the two larger of which continue after the flower has fallen, and embrace, like wings, a flat bicellular seed-vessel.

1. Seneka Pharm. Edinb. Polygala (Senega) floribus imberbibus spicatis, caule erecto herbaceo simplicissimo, foliis lato-lanceolatis Linn. Seneka or Senegaw milkwort, rattlesnake-rooted milkwort: with oblong, somewhat oval, pointed leaves; upright unbranched stalks; white slowers; and a variously bent and divaricated jointed root, about the thickness of the little singer, with a membranous margin running its whole length on each side, externally of a yellowish or pale brownish colour, internally white. It is a native of Virginia, Pensylvania, and Maryland, and cultivated in some of our gardens.

THE root of this plant is faid to be the specific of the Senegaw Indians against the poison of the bite of the rattlesnake; and to be effectual, when used early, even in the middle of the summer heats, when the poison is in its highest vigour, and when all their other antidotes fail. The powder or a decoction of the root is taken internally; and either the powder, or cataplasms made with it, applied to the wound.

Dr. Tennent, observing that this poison produces symptoms refembling those of pleurisies and peripneumonies (a difficulty of breathing, cough, spitting of coagulated blood, and a strong quick pulse) conjectured that it might be serviceable in those distempers also: and from the trials made by the gentlemen of the French academy, as well as those mentioned by him, its virtues appear to be great. It made the sizy blood sluid, procured a plentiful spitting, increased perspiration and urine, and sometimes purged or vomited. The usual dose was thirty or thirty-sive grains of the powder; or three spoonfuls of a decoction prepared by boiling three ounces of the root in a quart of water till near half the liquor was consumed.

The

The feneka root has been tried likewise in hydropic cases, and sound in some instances to procure a copious evacuation by stool, urine, and perspiration, after the common purgatives and diuretics had failed. Mons. Bouvart observes, that though dropsies were thus removed by the seneka, the cure did not seem complete, a swelling and hardness of the spleen remaining, which sometimes occasioned a fresh extravasation: that the medicine sometimes acts by liquesying the blood and juices, without producing a due discharge; and that in these cases it does harm unless affisted by proper additions, but that so long as it proves cathartic, nothing is to be feared from it. It is said to have been found serviceable also in the rheumatism and gout.

This root, of no remarkable smell, has a peculiar kind of subtile pungent penetrating taste. Its virtue is extracted both by water and spirit, though the powder in substance is supposed to be more effectual than either the decoction or tincture. The watery decoction, on first tasting, seems not unpleasant, but the peculiar pungency of the root quickly discovers itself, spreading through the sauces, or exciting a copious discharge of saliva, and frequently, as Linnæus observes, a short cough: those to whom I have directed this medicine, have generally found a little Madeira most effectual for removing its taste from the mouth, and making it to sit easy on the stomach. A tincture of the root in rectified spirit is of a more fiery pungency, extremely durable in the mouth and throat, and apt to promote vomiting or reaching.

2. Polygala vulgaris C.B. & Linn. Flos ambarvalis. Common milkwort: with the stalks procumbent; the lower leaves roundish, the upper oblong, narrow, and pointed; the flowers blue, purplish or red, sometimes white, with a kind of fringed appendix on the lower lip; the roots slender and hard. It grows wild in dry pasture grounds.

THE roots of this species are somewhat similar in taste to those of the preceding, but far weaker: they have been sound likewise to produce the same effects in pleurisies, in a lower degree. The leaves of the plant are very bitter: Gesner, who from this quality gives it the name of amarella, relates, that an insussion of a handful of them in wine is a safe and gentle purgative.

POLYPODIUM.

POLYPODIUM.

POLYPODIUM Pharm. Edinb. (a) Polypodium vulgare C.B. & Linn. Polypody: a plant with long leaves issuing from the root, divided on both sides, down to the rib, into a number of oblong segments, broadest at the base: it has no stalk, or manifest flower; the seeds are a fine dust, lying on the backs of the leaves, in roundish specks, which are disposed in rows parallel to the rib: the roots are long and slender, of a reddish brown colour on the outside, greenish within, full of small tubercles, which are resembled to the feet of an insect, whence the name of the plant. It grows in the clefts of old walls, rocks, and decayed trees: that produced on the oak has been generally accounted the best, though not sensibly different from the others. It is found green at all seasons of the year.

The leaves of polypody have a weak ungrateful fmell, and a naufeous fweet taste, leaving a kind of roughness and slight acrimony in the mouth. They give out their smell and taste, together with a yellow colour, both to water and rectified spirit: the spirituous tincture is sweeter than the watery, but in inspissation its sweetness is in great part destroyed or covered by the other matter; the spirituous extract, as Cartheuser observes, being to the taste only subastringent and subacrid, with very little sweetness, while the watery extract seems to retain the full sweetness of the polypody. The root is supposed to be aperient, resolvent, and expectorant: it was formerly ranked among the purgatives, but operates so weakly, a decoction of an ounce or two scarcely moving the belly, that it has long been expunged from that class: the present practice pays very little regard to it in any intention.

POPULUS.

POPULUS NIGRA Pharm. Edinb. (a) & C. B. & Linn. Populus nigra five aigeiros J. B. BLACK POPLAR: a large tree; with dark green, fomewhat rhomboidal acuminated leaves; producing imperfect flowers, in catkins: in fome of the individuals, called male, the

(a) Expunged.

flowers are barren; in others, called female, they are followed by membranous pods, containing a number of feeds winged with down. It is indigenous in watery places, and quick of growth.

The young buds or rudiments of the leaves, which appear in the beginning of the spring, were formerly employed in an officinal ointment, which received its name from them. At present, they are almost entirely disregarded; though they should seem, from their sensible qualities, to be applicable to purposes of some importance. They abound with a yellow, unctuous, odorous, balsamic juice, which they readily impart, by maceration or digestion, to rectified spirit. The tincture, inspissated, yields a fragrant resin, superiour to many of those brought from abroad, and approach to the nature of storax.

*A species of poplar growing in Siberia and in North America, called by Linnæus *Populus balfamifera*, is said to be much more abundant in balfamic juice than the former, insomuch that the buds give it out on

mere expression (a).

PRUNELLA.

PRUNELLA sive Brunella Ph. Edinb. (b) Prunella major folio non dissecto C. B. Consolida minor. Symphitum minus. Prunella vulgaris Linn. Self-heal: a small plant; with square stalks; oval uncut leaves set in pairs on pedicles; and short thick spikes of purplish labiated slowers. It is perennial, grows wild in pasture grounds, and slowers in June and July.

This herb is recommended as a mild restringent and vulnerary, in spittings of blood, and other hemorrhagies and fluxes; and in gargarisms against aphthæ and inflammations of the sauces. Its virtues do not appear to be very great: to the taste it discovers a very slight austerity or bitterishness; which is more sensible in the flowery tops than in the leaves; though the latter are generally directed for medicinal use.

(a) Bergius, Mat. Med. 804.

(b) Expunged.

PRUNUS.

PRUNUS: a tree with pentapetalous white flowers; each of which is fucceeded by a roundish or oval fruit, standing on a long pedicle, composed of a sleshy pulp including a flat stone pointed at both ends.

T. PRUNUS HORTENSIS. Prunus domestica Linn. Garden plum tree; without prickles; bearing a sweet fruit. Three sorts of this fruit are ranked among the articles of the materia medica: they are all met with in our gardens, but the shops are supplied with them, moderately dried, from abroad. I. PRUNA BRIGNOLENSIA Pharm. Edinb. (a) Pruna ex slavo rusescentia mixti saporis gratissima C. B. The Brignole plum or prunelloe, brought from Brignole in Provence, of a reddish yellow colour, and a very grateful sweet subacid taste. 2. PRUNA GALLICA Ph. Lond. Edinb. Pruna parva dulcia atro-cærulea C.B. The common or French prunes, called by our gardeners the little black damask plum. 3. PRUNA DAMASCENA: Pruna magna dulcia atro-cærulea C. B. Damsons, the larger damask violet plum of Tours: this is seldom kept in the shops, and has been generally supplied by the common prune.

All these fruits possess the same general qualities with the other summer fruits. The prunelloes, in which the sweetness has a greater mixture of acidity than in the other sorts, are used as mild refrigerants in severs and other hot indispositions, and are sometimes kept in the mouth for alleviating thirst in hydropic cases. The French prunes and damsons are the most emollient, subricating and laxative: they are taken by themselves for gently loosening the belly in costive habits and where there is a tendency to inflammation: decostions of them afford an useful basis for laxative or purgative mixtures, and the pulp in substance for

electuaries.

2. PRUNUS SILVESTRIS Pharm. Lond. & Edinb. & C. B. Acacia Germanorum. Prunus spinosa Linn. Black thorn or sloe: a prickly bush, common in hedges, producing austere fruit, somewhat smaller than an ordinary cherry.

THE fruit of the floe bush is so harsh and austere, as not to be eatable till thoroughly mellowed by frosts. The juice expressed from it while unripe, or before it has been thus mellowed, inspissated by a gentle heat to dryness, is called German acacia, and has been usually fold in the shops for the Egyptian juice of that name; from which it differs in being harder, heavier, darker coloured, of a sharper or tarter taste, and more remarkably in this, that it gives out its aftringency in good meafure to rectified spirit as well as to water, whereas that of the Egyptian acacia is not at all dissoluble in spirit. A conserve of this fruit is likewise prepared in the shops, by mixing the pulp with thrice its weight of double-refined fugar: the floes being previously steeped in water, over the fire, with care that they do not burst, till they are sufficiently foftened to admit of the pulp being pressed out through a sieve. In some places, the unripe floes are dried in an oven, and then fermented with wines or malt liquors, for a restringent diet drink in alvine and uterine laxities.

Conferv. prunor fylvestrium Pb. Lond. & Ed.

The bark, both of the branches and of the roots, is faid to have been given with success in intermitting fevers, and by some stands recommended as equal to the Peruvian bark. It is apparently a strong styptic; and its styptic matter is of that kind which is not easily extracted by watery menstrua.

The flowers, in smell very agreeable, and in taste bitterish, appear to have a laxative virtue, like those of the peach tree or the damask rose. They impregnate water, by distillation, strongly with their fragrance; and give out their active matter, by infusion, both to water and spirit. The watery infusion, sweetened with sugar, or made into a syrup, is said to be a very useful purgative for children.

PSYLLIUM.

PSYLLIUM Pharm. Parif. Pulicaris berba Lugdun. Pfyllium majus erectum C. B. Plantago Pfyllium Linn. FLEAWORT: an herb of the plantane kind, differing from the common plantanes in being annual, and having its stalks branched, with leaves upon them, which are long, slender, and somewhat hairy. It grows wild in the warmer parts of Europe, and is sometimes raised in our gardens. The seeds have been usually brought from the south of France: they are small, smooth, slippery,

flippery, of a shining brown colour, of an oblong flattish figure supposed to resemble that of a flea, whence the name of the plant.

The feeds of fleawort have a naufeous mucilaginous tafte, and no remarkable fmell: a dram renders near a pint of water flimy and yellowish: the decoction, inspiffated, leaves a strong dark brown mucilage, which impresses on the palate an unpleasant, weak, but penetrating acrimony. This mucilage has been employed chiefly in emollient glysters, in gargarisms for hoarseness and asperity of the sauces, and in external applications for chaps of the lips and inflammations of the eyes. Prosper Alpinus relates, that among the Egyptians, the mucilage or an insusion of the seeds is given internally, in ardent severs; and that it generally either loosens the belly or promotes sweat. The particular virtue of these seeds, or whatever virtue they may have distinct from that of mucilaginous substances in general, appears to reside in the acrid matter, which may be separated from the mucilaginous by rectified spirit: the feeds, digested in rectified spirit, give out their acrimony and ill taste, and yield afterwards to water an almost insipid mucilage.

PTARMICA.

PTARMICA, Pseudopyrethrum, Pyrethrum silvestre, Draco silvestris, Tarchon silvestris, Sternutamentoria: Dracunculus pratensis serrato solio C.B. Achillea Ptarmica Linn. Sneezewort of Bastard pellitory: a plant with long narrow leaves finely serrated about the edges, and radiated discous white slowers set in form of umbels on the tops of the branches. It is perennial, grows wild on heaths and in moist shady grounds, and is sound in flower from June to the end of summer.

THE roots of this plant have a hot biting taste, approaching to that of pyrethrum, with which they nearly agree also in their pharmaceutic properties, and to which they have been sometimes substituted in the shops. They are by some recommended internally as a warm stimulant and attenuant; but their principal use is as a masticatory and sternutatory.

PULEGIUM.

PENNYROYAL: a plant of the mint kind; differing from the mints strictly fo called, in the flowers being disposed, not in spikes on the tops, but in thick clusters, at distances, round the joints of the stalks; and the upper fegment of the flower not being nipped at the extremity.

- I. PULEGIUM Pharm. Lond. & Edinb. Pulegium latifolium C. B. Mentha palustris sive pulegium Pharm. Paris. Pulegium regium Ger. emac. Common pennyroyal: with fomewhat oval Mentha pulegium Linn. obtuse leaves, and trailing stalks, striking root at the joints. It grows wild on moift commons and in watery places, and flowers in June.
 - 2. PULEGIUM ERECTUM: Pulegium erectum officinarum Dale: Pulegium mas Ger. emac. Upright pennyroyal; with the stamina standing out from the flowers; faid to be a native of Spain, common in our gardens, and usually substituted in our markets to the foregoing species.
 - 3. Pulegium cervinum: Pulegium angustifolium C. B. aquatica satureiæ folio Tourn. Mentha cervina Linn. Harts pennyroyal: with small oblong narrow leaves; said to grow wild about Montpellier.

ALL the pennyroyals are warm pungent herbs, fomewhat fimilar to mint, but more acrid and less agreeable both in smell and taste, less proper in common nauseæ and weakness of the stomach, more efficacious as warm carminatives and deobstruents in hysteric cases and disorders of the breast: the last species is the strongest, though least ungrateful, of the three. Their active principle is an effential oil; of a more volatile nature than that of mint, coming over hastily with water at the beginning of the distillation, and rising also in great part with highly-rectified spirit; in taste very pungent, and of a strong smell; when newly drawn, of a yellowish colour with a cast of green; by age turning brownish. The oil, and a simple + and spirituous t water strongly impregnated with it, Pharm. Lond. by drawing off a gallon of water or proof spirit from a pound and a half of the dry leaves, are kept in the shops.

Ol. pulegii effent. Ph.L.

Aq. pulegii + Pb. Edinb. fimplex t fpirituofa Pharm. Lond.

PULMONARIA.

maculosum sive pulmonaria latifolia C. B. Pulmonaria officinalis Linn. Spotted lungwort, Jerusalem cowslips, Jerusalem sage: a hairy scabrous plant, with the leaves of a dark brownish green colour on the upper side and spotted for the most part with white, underneath of a paler green, the lower oval and set on broad pedicles, those on the stalks narrower, long-pointed, set alternately, without pedicles: the slowers are monopetalous, cut into sive sections, of a purple or blue colour, and sometimes white, sollowed each by sour seeds inclosed in the cup. It is perennial, grows wild in several parts of Europe, and slowers in our gardens in April and May.

The leaves of pulmonaria, recommended in hemoptoës, tickling coughs, afperities of the fauces, &c. appear to be of little medicinal virtue. The dried leaves have hardly any smell; and their taste is just perceptibly mucilaginous, sweetish, and roughish. They seem to be nearly of the same nature with the adianthum and trichomanes.

*PULSATILLA.

PULSATILLA NIGRICANS Stærck, Pharm. Edinb. Pulsatilla flore minore nigricante C. B. Anemone pratensis Linn. A species of anemone, much resembling the pulsatilla vulgaris, or pasque flower, but its flower is less, and of a darker hue. It is a native of the south of Germany, and other neighbouring countries.

All the anemonies have a confiderable degree of acrimony; but this feems to possess the largest share. The whole plant when chewed impresses the tongue with a sharp, burning, durable taste. The root is milder than the other parts. On distilling the plant with water, the liquor which comes over is strongly impregnated with its virtues; and the remaining extract is also considerably active.

Dr. Stærck of Vienna, to whom the introduction of so many of the more powerful vegetables is owing, has likewise recommended this to the medical practitioner. From numerous trials, he celebrates its efficacy

in various chronic diseases of the eye; in venereal nodes and nocturnal pains; in soul ulcers with caries; in serpigo; and suppressed menses. He relates instances of its curing blindness of many years continuance, by dissipating and dissolving silms and obscurities of the cornea. In these cases, its good effects were first indicated by considerable pain excited in the eye. The sensible operation of the medicine was nausea and vomiting, particularly when the distilled water was used; an increased flow of urine; and sometimes gripes and looseness; with increased pain at first in the affected part. From all these circumstances, the pulsatilla seems to be endued with very active and penetrating powers, yet such as may be employed with perfect safety if proper caution be used. The dose of the distilled water to adults is about half an ounce, twice or thrice a day; of the extract, reduced to powder with the addition of sugar, sive or six grains. Bergius mentions having given the extract copiously, especially in diseases of the eyes, but without any effect (a).

Extract. folior. Pulfatillæ nigricantis Pb. E. The Edinburgh college had adopted the distilled water of pulsatilla, but has now changed it for the extract.

PYRETHRUM.

PYRETHRUM Pharm. Lond. & Edinb. Pyrethrum flore bellidis C. B. Chamæmelum specioso flore, radice longa fervida Shaw afr. Dentaria, berba salivaris, & pes alexandrinus quibusdam. Anthemis Pyrethrum Linn. Pellitory of Spain: a trailing perennial plant; with finely divided leaves somewhat like those of camomile or sennel; and naked thick stalks, bearing each a large flower, which consists of a yellow disk surrounded with petala of a pure white colour on the upper side and a sine purple underneath: the root, which sinks deep in the ground like a carrot, is of a brownish colour on the outside and whitish within. It is a native of the warmer climates, but bears the cold of our own, and often produces flowers in succession from January to May: the roots also, as Parkinson observes, grow larger with us, than those which the shops are supplied with from abroad.

Pellitory root has a very hot pungent taste, without any sensible smell. Its pungency resides in a resinous matter, of the more fixt kind;

being extracted completely by rectified spirit, and only in small part by water; and not being carried off, in evaporation or distillation, by either menstruum. The spirituous extract is extremely siery, but in small quantity, scarcely amounting to one twentieth of the weight of the root. The watery insusion is nauseous, but scarcely discovers any acrimony till concentrated by inspissation; when reduced to the consistence of an extract, it proves considerably pungent: the quantity of this extract is commonly five or six times as large as that of the spirituous. The root remaining, after the action of water, yields still with rectified spirit a very siery extract; whereas that, which has been digested in spirit, yields with water only an insipid mucilaginous substance.

The principal use of pyrethrum, in the present practice, is as a masticatory, for stimulating the salival glands, &c. and evacuating viscid humours from the head and parts adjacent: by this means it frequently relieves toothachs, some kinds of headachs, lethargic complaints, and paralyses of the tongue. It has sometimes likewise been given internally, from a few grains to a scruple, as a hot stimulant and attenuant, in paralytic and rheumatic disorders.

PYRITES.

MARCASSITA Pharm. Parif. Pyrites or Marcasite: a hard fossil; striking fire with steel, copiously, and in large sparks; becoming vitriolic, either by simple exposure to the air, or by calcination and subsequent exposure.

This mineral varies extremely in its appearances. It is found of a bright brass yellow, of a greenish, of a grey or whitish colour, and of different intermediate or mixt shades; in masses, rarely of any great size, globular, oblong and slattish, cubical, octoedral, dodecaedral; sometimes covered with a coat or crust, but oftener bare; internally sometimes striated, and sometimes of an even and simple structure (a). It is met with in different places of this kingdom, and in most parts of the world; on the surface of the earth, on the sea shores, in clay pits, embedded in earthy and stony bodies of various kinds.

The pyritæ confift, in general, of fulphur, iron, and unmetallic earth: in some, a little copper is joined to the iron; and in some, copper is the prevailing metal. In some, particularly the yellow kind, the quantity of sulphur is large: in others, particularly the white, both the

fulphur and metal are in fmall proportion.

If artificial mixtures of fulphur with iron or copper be gently calcined, the inflammable principle of the fulphur exhales, and its acid remains united with the metal, forming therewith a faline vitriolic compound: a mixture of iron filings and fulphur, moistened with water, fuffers a like change without external heat, and if the quantity is large, bursts spontaneously into fire. A resolution of the same kind happens in the natural pyritæ on exposing them to the air and rain; provided, where they are very fulphureous, a part of the fulphur be previously dislipated by calcination. On this exposure, they all become powdery and acquire a vitriolic tafte, the ferrugineous much more eafily than those which have any admixture of copper: fome shoot out efflorescences of vitriol upon the furface: from others, the faline matter, washed off by rain, is found to confift chiefly of the fulphureous or vitriolic acid. If the pyritæ, even fuch as have the least fulphureous and metallic impregnation, as those from which the English vitriol is made, be laid in large heaps, they grow hot, and take fire, and emit, during the burning, strong diffusive fulphureous vapours (a).

The pyritæ, in substance, are never used medicinally; but in their products they are very important. It is from these, that common sulphur is extracted, in Sweden and Saxony; that the native vitriols are produced in caverns of the earth or on its surface; that the greatest quantities of artificial vitriol are prepared; and that the chalybeate mineral waters are supposed to receive their impregnation: see the respec-

tive articles.

*2.UASSIA.

QUASSIA Pharm. Edinb. Lignum Quassiæ Amænit. Acad. Vol. VI. Bois de Coissi Fermin Surinam. Quassy Root: the woody root of a tree growing in Surinam, called by Linnæus, Quassia amara, of the class and order decandria monogynia in his system. This root is as thick as a

MATERIA MEDICA.

man's arm. Its wood is whitish, hard, solid, and tough, becoming yellowish on exposure to the air. It is covered by a thin, grey, fissured, and brittle bark.

Quassi root has no fensible odour. Its taste is that of a pure bitter, more intense and durable than that of almost any other known substance. Its watery insusions and decoctions, and its spirituous tinctures, are all almost equally bitter, of a pale yellowish hue, which is not blackened by the addition of martial vitriol. The watery extract is from a sixth to a ninth of the weight of the wood; the spirituous, about a twenty-fourth. The bark of the root is reckoned in Surinam more powerful than the wood. The flowers also are a strong bitter.

THE medical use of the quassi has been a considerable time known in Surinam. The flowers were long ago employed by the natives as an excellent stomachic. The root was a fecret remedy used by a negro, named Quaffi, in the fatal fevers of that country, from whom it was purchased by Dan. Rolander, a Swede, who returned from thence in 1756. Some specimens of the wood and of the fructification were, in 1761, presented by M. Dahlberg to the celebrated Linnæus; who drew up a botanical description of the plant, with an account of its virtues, and published it in the fixth vol. of the Amani. Acad. A confirmation of its medical powers appeared in a letter from Mr. Farley, a practitioner in Antigua, printed in the Phil. Transact. Vol. LVIII. He found it remarkably efficacious in suppressing vomitings, stopping a tendency to putrefaction, and removing fevers. It feemed capable of producing all the good effects of Peruvian bark, without heating. Some further experiments on the quassia are contained in a late medical thesis by Dr. Ebeling. He confirms the general account of its virtues, with this additional circumstance, that though its general antiseptic powers were inferiour to those of Peruvian bark, yet it preserved bile a longer time from putrefaction. In this circumstance it agrees with another pure bitter, the columbo root.

From these relations, the quassi appears to be a valuable addition to our tonic remedies, and has therefore obtained a place in the last Edinburgh and other pharmacopæias. It may be used either in infusion, or extract:

the latter, made into pills, on account of the intense bitterness of the drug, is preferable for delicate stomachs.

QUERCUS.

QUERCUS Ph. Edinb. Quercus cum longis pediculis C. B. Quercus Robur Linn. OAK: a large tree, with oblong leaves, widening from the bottom to the extremity, and finuated or bluntly indented about the edges: the fruit is an acorn, or kernel with a coriaceous covering, inclosed at bottom in a scaly cup. It is a common forest tree in most parts of Europe.

THE bark of the oak is a strong astringent, accompanied with a moderate bitterness, but no remarkable smell or particular slavour: with solution of chalybeate vitriol it strikes an inky blackness. It is said to have been employed with success, not only for restraining hemorrhagies, and other immoderate evacuations, but likewise in intermitting severs and in gleeting gangrenous wounds and ulcers; in which cases, an extract made from it is said by some to be equal to that of the Peruvian bark. A decoction of it used as a somentation is said to have cured a procidentia recti (a). It gives out its virtue both to water and rectified spirit.

QUERCUS MARINA.

QUERCUS MARINA sive fucus vesiculosus: Fucus maritimus sive quercus maritima vesiculas babens C.B. Fucus vesiculosus Linn. Sea wrack or Sea oak: a soft, very slippery, marine plant; common upon rocks that are left dry at the ebb tide; with the leaves somewhat resembling in shape those of the oak tree; the stalks running along the middle of the leaves, and terminated by warty bladders containing either air or a slippery sluid. The vesicles begin in March to fill with a thin juice; and about the end of July they burst, and discharge a matter as thick as honey.

DR. RUSSEL relates, that he found this plant an useful affistant to sea water in the cure of disorders of the glands: that he gave it in powder

to the quantity of a dram, and that in large doses it nauseated the stomach: that by burning in the open air it was reduced into a black faline powder; which feemed, as an internal medicine, greatly to excel Æthiops vethe officinal burnt sponge; which was used with benefit, as a dentifrice, getabilis D. Russel. for correcting laxities of the gums; and which shewed a notable degree of detergent virtue by its effect in cleaning the teeth: that the juice of the veficles, after standing to putrefy, yielded, on evaporation, an acrid pungent falt, amounting to above a scruple from two spoonfuls: that the putrefied juice, applied to the skin, finks in immediately, excites a flight fense of pungency, and deterges like a folution of soap: that one of the best applications for discussing hardness, particularly in the decline of glandular fwellings, is a mixture of two pounds of the juicy veficles, gathered in July, with a quart of fea water, kept in a glass veffel for ten or fifteen days, till the liquor comes near to the confiftence of very thin honey: the parts affected are to be rubbed with the strained liquor twice or thrice a day, and afterwards washed clean with sea water.

* RADIX LOPEZIANA.

RADIX INDICA LOPEZIANA Pharm. Edinb. Radix Indica a Joanne Lopez denominata Gaubii Adversar. Cap. vi. Rais di Juan Lopez Lusitanis. The root of an unknown tree, growing, as some affert, at Goa, as others suppose, in Malacca, from whence it is sometimes brought to Batavia. It is met with in pieces of different thickness, fome, at least, of two inches diameter. The woody part is whitish, and very light; fofter, more fpongy, and whiter next the bark, including a denfer fomewhat reddish medullary part. The bark is rough, wrinkled, brown, foft, and as it were woolly, pretty thick, covered with a thin paler cuticle.

NEITHER the woody nor cortical part has any remarkable smell or taste, nor any appearance of resinous matter. On boiling in water, no odour is emitted; and the strained liquor, which is of a vellow hue, is almost infipid, only impressing the tongue with a very light obscure bitterishness; and without viscidity. The extract obtained by evaporating the decoction is equally void of fensible activity. Rectified spirit is tinged by the root of a brown colour, but acquires no particular tafte.

After

After drawing off the spirit from the tincture, a matter remains refembling balfam, which bubbles and inflames in the fire, and has a

bitterish taste, like that of opium.

Though the preceding examination of this root is not favourable to the opinion of its medical powers, yet it is regarded in the East Indies as a medicine of extraordinary efficacy in diarrheas; and the learned Gaubius, in his Adversaria, has published an account of some experiments made with it, which in some degree confirm its reputation. From his own trials, and those of his friends, it appeared most remarkably effectual in stopping colliquative diarrheas which had resisted the usual remedies. Those attending the last stage of consumptions were particularly relieved by its use. It seemed to act not by any astringent power, but by a faculty of restraining and appeasing spasmodic and inordinate motions in the intestines. Gaubius compares its action to that of simarouba, but thinks it more efficacious than this medicine.

The mode of exhibiting it in India, is to levigate the root with water on a porphyry till reduced to a fine pulp. In Europe the powder of it has been given with any proper vehicle, in doses from fifteen to thirty grains, repeated three or four times a day: one practitioner found a tincture of it in common spirits equally effectual with the root in substance. Of this, a tea-spoonful was given thrice a day in red wine. The colleges of Edinburgh and Brunswick have received this root into their catalogues; but it is scarcely yet to be met with in the shops.

RANUNCULUS.

CROWFOOT: a plant with pentapetalous flowers fet in five-leaved cups; followed each by a round cluster of naked feeds. It is perennial.

1. RANUNCULUS. Ranunculus pratensis radice verticilli modo rotunda C.B. Ranunculus bulbosus Linn. Bulbous crowfoot, butter flower, gold cup: with a round tuberous root about the size of an olive; the leaves divided commonly into three segments, and these further subdivided; the stalks erect: the flowers of a bright glossy yellow, and their cups turned downwards. It is common in pasture grounds, and flowers in May.

2. FLAM-

2. FLAMMULA: Ranunculus longifolius palustris minor C. B. Ranunculus Flammula Linn. Smaller water crowfoot or spearwort: with shous roots, long narrow leaves acuminated at both ends, and leaning or procumbent stalks. It grows in watery places or moist meadows, and slowers in June.

The roots and leaves of these plants are of no considerable smell, but in taste highly acrid and siery. Taken internally, they appear to be deleterious, even when so far freed from the caustic matter, by boiling in water, as to discover no ill quality to the palate. The effluvia likewise even of the less acrid species or varieties cultivated in gardens, when freely inspired, have occasioned headachs, anxieties, vomitings, and spasmodic symptoms. The leaves and roots, applied externally, inspame and exulcerate or vesicate the part, and are liable to affect also the adjacent parts to a considerable extent (a): they have sometimes, particularly among empyrics and the common people, supplied the place of the far safer and not less effectual vesicatory, cantharides, for procuring an ulcer and discharge of serum, in sciaticas and some fixt pains of the head. Their pungency is diminished by drying, and by long keeping seems to be dissipated or destroyed.

RAPHANUS.

RAPHANUS RUSTICANUS Pharm. Lond. & Edinb. & C. B. Cochlearia folio cubitali Tourn. Cochlearia Armoracia Linn. Horse-radish: a plant refembling scurvygrass in the flowers and seeds, but differing in the leaves being very large and long, and indented about the edges. It is sometimes sound wild about the sides of ditches and rivulets, but for medicinal and culinary uses is cultivated in gardens. It is perennial, flowers in June, rarely perfects its seeds, and is propagated from transverse cuttings of the roots.

Horseradish root affects the organs both of taste and smell with a quick penetrating pungency: it nevertheless contains in certain vessels a sweet juice, which sometimes exudes in little drops upon the surface.

Its pungent matter is of a very volatile kind; being totally diffipated in drying, and carried off in evaporation or distillation both by water and rectified spirit: as the pungency exhales, the sweet matter of the root becomes more fenfible, though this also is in great measure dislipated or destroyed. It impregnates both water and spirit, by infusion or by distillation, very richly with its active matter: in distillation with water it yields a small quantity of effential oil exceedingly penetrating and This root appears therefore to agree with scurvygrass and creffes, and to differ from mustard feed to which it is by some resembled, in the volatility of its pungent matter, and its folubility in spirit.

Horseradish is a moderately stimulating, aperient, and antiseptic medicine: it fenfibly promotes perspiration, urine, and the expectoration of viscid phlegm, and excites appetite when the stomach is weakened or relaxed, without being fo liable to produce immoderate heat, or inflammatory symptoms, as the stimulants of the aromatic kind. It is principally used in paralytic and rheumatic complaints, in scurvies and fcorbutic impurities of the humours, in cachectic diforders, and in dropfies, particularly in those which follow intermitting fevers. Taken

in confiderable quantities, it provokes vomiting.

RAPUM.

RAPUM Pharm. Edinb. (a) Rapa sativa rotunda C. B. Brassica Rapa Linn. TURNEP: a plant with a round root; jagged leaves, rude to the touch; tetrapetalous flowers, commonly yellow; and fmall round smooth reddish or blackish seeds lodged in long pods. The garden turnep is supposed to be a variety produced by culture from the smaller fort which grows wild in fandy grounds in some parts of England. It is biennial.

TURNEPS are accounted a falubrious food; demulcent, detergent, fomewhat laxative and diuretic; but liable in weak stomachs to produce flatulencies, and prove difficult of digestion: the liquor pressed out from them after boiling, is fometimes taken medicinally, in coughs and disorders of the breast. The seeds have been accounted alexipharmac or diaphoretic: they have no smell, but discover to the taste a mild acrimony, feemingly of the same nature with that of mustard seed, though far weaker.

RHABARBARUM.

RHUBARB: a plant with large dock-like leaves, among which arises a single thick stalk bearing loose clusters of naked monopetalous bell-shaped slowers divided into six segments: each flower contains nine stamina (whereof the docks strictly so called have but six), and is followed by a triangular seed surrounded about the edges with a leasy margin.

1. Rhabarbarum Ph. Lond. Rheum Ph. Edinb. Lapathum orientale folio latissimo undulato & mucronato Mill. diet. Rheum undulatum foliis subvillosis, petiolis æqualibus Linn. Rhubarb: with the leaves somewhat heart-shaped, acuminated, and slightly hairy, and the pedicles planoconvex. It is a native of China and Siberia, and has lately been raised in some of our gardens, where it is found to grow with vigour in the open ground (a).

Two forts of rhubarb roots are met with in the shops. The first is imported from Turkey and Russia, in roundish pieces, freed from the bark, with a hole through the middle of each, externally of a yellow colour, internally variegated with lively reddish streaks. The other, which is less esteemed, comes immediately from the East Indies, in longish pieces, harder, heavier, and more compact than the foregoing. The first fort, unless kept very dry, is apt to grow mouldy and wormeaten: the second is less subject to these inconveniencies. Some of the more industrious artists are said to fill up the worm holes with certain mixtures, and to colour the outside of the damaged pieces with powder of the finer sorts of rhubarb, and sometimes with cheaper materials. The

⁽a) The plant above described is that which is generally reckoned the true rhubarb plant, having been produced from the seeds, sent from Russia, as those of the true rhubarb, to Justieu at Paris, Rand at Chelsea, and Linnæus at Upsal. Dr. Hope received lately rhubarb seeds from the same country, which being sown in the open ground at Edinburgh, produced a different species, Rheum palmatum Linnæi, with the leaves deeply cut into pointed segments. He observes that the root of this plant, though taken up too young, and at an improper season, viz. in July, agreed perfectly with the best foreign rhubarb, in colour, smell, taste, and purgative quality. See Philosoph. Transact. vol. lv. for the year 1765.—Perhaps the roots of both species may be of the same quality, and taken promiscuously.

marks of the goodness of rhubarb are, the liveliness of its colour when cut; its being firm and solid, but not flinty or hard; its being easily pulverable, and appearing when powdered of a fine bright yellow colour; its imparting to the spittle, on being chewed, a deep saffron tinge, and not proving slimy or mucilaginous in the mouth. Its taste is subacrid, bitterish, and somewhat styptic; the smell, lightly aromatic.

Rhubarb is a mild cathartic, and commonly looked upon as one of the fafest and most innocent of the substances of this class. Besides its purgative virtue, it has a mild astringent one, discoverable by the taste, and by its striking an inky blackness with chalybeate solutions: hence it is found to strengthen the tone of the stomach and intestines, to leave the belly costive, and to be one of the most useful purgatives in diarrheas, dysenteries, and all disorders proceeding from a debility and laxity of the fibres: it is frequently indeed given with a view rather to this stomachic and corroborating virtue, than to its producing any considerable evacuation. It tinges the urine of a high yellow colour.

Rhabarb. torrefactum Pharm. Lond. Rhubarb in substance purges more effectually than any preparation of it: the dose is from a scruple to a dram. By roasting it with a gentle heat, till it becomes easily friable, its cathartic power is diminished, and its astringency supposed to be increased.

In its habitude to menstrua, it differs remarkably from most of the other cathartic drugs, its purgative virtue being extracted far more perfectly by water than by rectified spirit: the root remaining after the action of water is almost, if not wholly, inactive; whereas, after repeated digeftion in spirit, it proves still very considerably purgative: the colour of both tinctures is a fine deep yellow, that of the spirituous palest; when the rhubarb has given out to spirit all that this menstruum can extract, it still imparts a deep colour, as well as a purgative impregnation, to water. The watery infusion, in being inspissated by a gentle heat, has its virtue so much diminished, that a dram of the extract is faid to have scarcely any greater effect than a scruple of the root in substance: the spirituous tincture looses less; half a dram of this extract proving moderately purgative, though scarcely more so than an equal quantity of the powder. The spirituous extract dissolves almost wholly in water; and hence the tincture does not, like the spirituous infusions of most other vegetables, turn milky on being mixed with aqueous liquors: of the watery extract, scarce above one fourth is dissolved by

rectified

rectified spirit, and the part that does not dissolve proves more purgative than that which does.

*A watery infusion is directed in the Edinburgh pharmacopæia, made Infus. rhei by infufing half an ounce of rhubarb for a night in eight ounces of boiling water, and adding to the strained liquor one ounce of spirituous cinnamon water.

Phar. Edinb.

Tinctures of this root are drawn in the shops with proof spirit and with mountain wine. The London college directs an ounce of rhubarb with two drams of cardamom feeds, and one of faffron (a), for each tincture+: that of Edinburgh, orders for the vinous tincture, two ounces of rhubarb and one dram of canella alba to be infused in fifteen ounces of mountain wine, and two of proof spirit; for the simple spirituous tincture, three ounces of rhubarb, and half an ounce of leffer cardamom feeds, to two pounds and a half of proof spirit |; in which, sometimes are diffolved, four ounces of fugar candy §; and a compound tincture, composed of two ounces of rhubarb, half an ounce of gentian, one dram of inakeroot, and two pounds and a half of proof spiritss. These preparations are used chiefly as mildly laxative corroborants, in weakness of the stomach, indigestion, diarrheas, colicky and other like complaints. The last tincture is, in many cases, an useful affistant to the Peruvian bark in the cure of intermittents.

+Tinct. rhabarb. vinof. & spirituosa Pharm. Lond. t Vinum rhei Phar. Edinb. ||Tinct.rhei. & Tinct. rhei dulcis.

§§ Tinct. rhei amara Phar. Edinb.

The Turkey rhubarb is, among us, univerfally preferred to the East India fort, though this last appears to be for some purposes at least equal to the other. It is manifestly more astringent, but has somewhat less of an aromatic flavour. Tinctures made from both with equal quantities of rectified spirit, have nearly the same taste: on drawing off the menstrua, the extract left by the tincture of the East India rhubarb proves in taste confiderably stronger than the other. Both forts appear to be the produce of the same climate, and the roots of the same species of plant, taken up probably at a different feafon, or cured in a different manner.

2. RHAPONTICUM Pharm. Parif. Rhabarbarum dioscoridis & anti-Rhaponticum folio lapathi majoris glabro C. B. Rheum quorum Tourn. Rhapontic: with fmooth roundish leaves, and Rhaponticum Linn.

fomewhat

⁽a) Saffron does not appear to be a very proper ingredient in these preparations, as it renders the taste rather more unpleasant; nor indeed does rhubarb seem, for general use, to want any aromatic addition.

fomewhat channelled pedicles. It grows wild on the mountain Rhodope in Thrace, from whence it was brought into Europe by Alpinus about

the year 1610: it bears the hardest winters of this climate.

The root of this plant, which appears to have been the true rhubarb of the ancients, is by some confounded with the modern rhubarb, though considerably different from that root in appearance as well as in quality. The rhapontic is of a dusky colour on the surface, and a loose spongy texture; more astringent than rhubarb, and less purgative: in this last intention two or three drams are required for a dose.

*RHODODENDRON.

RHODODENDRON CHRYSANTHEMUM. This plant, which is a new fpecies of the rhododendron of Linnæus, discovered by Professor Pallas, is a shrub growing near the tops of the high mountains named Sajanes, in the neighbourhood of the river Jenisea in Siberia.

It is called by the natives of the place chei, or tea, from their commonly drinking a weak infusion of it, as we do the Chinese plant of that name. A stronger preparation of it is, however, used by them as a powerful medicine in arthritic and rheumatic disorders. For this purpose, they take about two drams of the dried shrub, stalk and leaves together, and infuse it in nine or ten ounces of boiling water for a night, in the heat of an oven. This is drunk next morning for a dofe; which occasions heat, a degree of intoxication, with a fingular uneafy kind of fensation, and a fort of vermiculation in the affected parts. The patient is not permitted to quench the thirst this medicine occasions, as liquids, especially cold water, would produce vomiting, and diminish the effect of the remedy. In a few hours, all difagreeable fymptoms go off, commonly with two or three stools; and the patient finds his disease greatly relieved. A repetition of the dose twice or thrice generally completes the cure. This is the fubstance of the account given in a letter from Dr. Guthrie, of Petersburgh, to Dr. Duncan, Med. Comment. Vol. V. p. 434.

The rhododendron has been fince tried by Dr. Home in the Infirmary at Edinburgh; and the refult of his trials, as published in his Clinical Cases and Experiments, is, that it is a very powerful sedative, remarkably diminishing the frequency of the pulse; but that it was not peculiarly

efficacious in removing the acute rheumatifm.

RICINUS.

RICINUS.

RICINUS & ricinoides: large plants, with small flowers in clusters, and the fruit growing at a little distance from, or succeeding only a few of, the flowers: the fruit consists of three capsules, containing each a single seed, flatted on one side, generally about the size of a small bean, composed of a thin skin or shell including an oily kernel.

- 1. PALMA-CHRISTI Pharm. Parif. Ricinus, cataputia major, cherva major, kiki, & granum regium quibusdam: Ricinus vulgaris C. B. Ricinus communis Linn. Palma-christi, Mexico seed: with the fruit triangular, the seed furnished with a little knob at one end, externally variegated with blackish and whitish streaks, resembling both in shape and colour the insect ricinus or tick.
- 2. RICINOIDES, seu pineus purgans, vel pinhones indici Pharm. Paris. Carcas, nux barbadensis, & faba purgatrix quibusdam: Ricinus americanus major semine nigro C. B. Jatropha Curcas Linn. Barbadoes nut: with an oval walnut-like fruit, and oblong black seeds.
- 3. AVELLANA PURGATRIX C. B. Nuces purgantes Ger. Jatropha multifida Linn. Purging nut: with oval fruit, and roundish, somewhat triangular, pale brownish seeds.
- 4. TIGLIUM, grana tiglia. Pinus indica nucleo purgante, & lignum moluccense foliis malvæ, fructu avellanæ minore, cortice molliore & nigricante, pavana incolis C. B. Croton Tiglium Linn. Grana tilia: with roundish fruit, and dark greyish seeds in shape nearly like those of the first species.

THE first of these plants is said to be found wild in some of the southern parts of Europe: it is biennial. The others are middling sized trees, natives of America and the East Indies, from whence the seeds are sometimes brought to us.

THE two first of these seeds are sweetish, nauseous, and acrid: the third has scarcely any acrimony, and tastes nearly like almonds: the fourth is intensely hot and acrimonious. They are all strong evacuants, operating,

operating, in doses of a few grains, both upwards and downwards; the sweet species not excepted. The grana tilia are the most violent, too much so to be taken with any tolerable safety; and indeed they all appear too drastic to be ventured on in substance.

They yield upon expression a considerable quantity of oil, impregnated more or less with the taste and the purgative quality of the seeds: of the oil of the grana tilia, Geoffroy limits the dose to one grain, which is probably an error of the press for one dram: that of the Barbadoes nut is said to be taken in America in larger quantities, and to purge without much inconvenience.

The oil of the palma-christi, vulgarly called in America castor oil, has been often given from two to four spoonfuls, and found to act as a sufficiently mild laxative: it is said to be particularly useful in the dry-belly-ach, and in other disorders where irritating purgatives cannot be borne, and where the common laxatives, on account of the large dose in which they require to be given, are apt to be rejected by the stomach. From such trials as I have made of this medicine, it did not seem to have any peculiar good qualities, or to produce any other effects than may be equally obtained by combining the more common purgatives, as tincture or insusion of sena, with common oil.—It is said that some, or all, of the above oils act as purgatives, when applied externally to the umbilical region.

The wood and leaves of the plants are likewise strong cathartics: Hermann relates, that the wood of the tilia, called panava or pavana, operates violently, when fresh, in the dose of a scruple or half a dram: that when dried and long kept, it is given to the quantity of a whole dram as a purgative, and in smaller doses as a sudoristic. Among us, all these substances are entire strangers to practice (except that the oil of the first species has of late been sometimes made use of;) and, so far as can be judged from the accounts given of them, they have little claim to be received *(a).

^{*(}a) The oleum ricini has increased in reputation fince the above was written, and is now much used as an aperient in calculous, nephritic, and various other cases. It is however liable to the inconvenience of great uncertainty in its operation, owing, probably, to the different degree of genuineness of different specimens of the oil.

ROSA.

ROSE: a prickly bush; with oval serrated leaves, set in pairs along a middle rib, which is terminated by an odd one; producing large elegant flowers, whose cup is divided into five long segments, with a knob at the bottom, which becomes an umbilicated foft fruit full of hairy feeds.

I. ROSA DAMASCENA Pharm. Lond. Rosa pallida Pharm. Edinb. Rosa purpurea C. B. Rosa centifolia Linn. The damask rose: with double flowers, of the fine pale red called from them rofe-colour.

The pleasant smell of damask roses is of a less perishable kind than that of many other odoriferous flowers, not being much diminished in drying, nor foon diffipated in keeping. They impart their odorous matter to watery liquors both by infusion and distillation: fix pounds of the fresh roses impregnate, by distillation, a gallon or more of water Aq. rosar. strongly with their fine flavour. On distilling large quantities, there pharm. Lond. separates from the watery fluid a small portion of a fragrant butyraceous pallid. Ph.E. oil, which liquefies by heat and appears yellow, but concretes in the cold into a white mass: an hundred pounds of the flowers, according to the experiments of Tachenius and Hoffman, afford scarcely half an ounce of oil. The oil and water, used chiefly as perfumes and flavouring materials, are recommended by Hoffman as excellent cordials, for raifing the strength and spirits, and allaying pain. They appear to be of a very mild nature, and not liable to irritate or heat the constitution; even the effential oil discovering to the taste but little pungency.

These flowers contain likewise a bitterish substance; which is extracted by water along with the odoriferous principle; which, after this last has been separated by distillation or evaporation, is found entire in the remaining decoction; and which appears to be of a gently purgative nature. The decoction, or a strong infusion of the flowers, made into a fyrup with a proper quantity of fugar, proves an useful laxative for chil- Syrup.rofar. dren, in doses of a spoonful: of the extract obtained by inspissating the decoction, from a scruple to a dram is said to be sufficient for adults. Phar. Edinh. The college of London directs the fyrup to be made, by pressing out the liquor remaining after the distillation of fix pounds of damask roses, and boiling it down to three pints; then, after it has fettled for a night,

folutiv. P.L. rofar. pallid.

adding five pounds of fine fugar, and boiling the mixture to the weight of feven pounds and a half: a spoonful of this syrup appears to be equivalent to about three drams of the fresh flowers. The solutive matter of the flowers is combined also in the same manner, for the purposes of glysters, with brown sugar and honey: towards the end of the boiling down of the strained decoction, an ounce of cummin seeds, bruised a little and tied in a linen cloth, is added; and the liquor afterwards boiled with four pounds of brown sugar and two of honey.

Mel, folutiv. Pharm. Lond.

Rectified spirit extracts both the odoriferous and the purgative matter of the damask rose, equally with water, or rather more completely. The spirit, distilled off from the filtered tincture, proves lightly impregnated with the fragrance of the flowers, and the inspissated extract retains likewise a part of their flavour along with the bitterish matter. This extract, in quantity smaller, and in taste stronger, may be presumed to be more purgative, than that made with water.

2. Rosa Rubra Pharm. Lond. & Edinb. Rosa rubra multiplex C. B. Rosa gallica Linn. The red rose: with double flowers of a deep red colour.

The red rose has very little of the fine slavour admired in the pale fort: to the taste, it is bitterish and subastringent. The astringency is greatest before the slowers have opened, and, in this state, they are chosen for medicinal use as a mild corroborant: the full-blown flowers are probably as laxative as those of the foregoing species, for Poterius relates, that he found a dram of powdered red roses occasion three or four stools, and this not in a few instances, but constantly, in an extensive practice, for several years. The astringency of the buds is improved by hasty exsiccation in a gentle heat: by slow drying, both the astringency and the colour are impaired.

Confervarofarum Phar. Lond. & Edin. The fresh buds, clipt from the white heels, and beaten with thrice their weight of fine sugar, form an agreeable and useful conserve; which is given in doses of a dram or two, dissolved in warm milk, in weaknesses of the stomach, coughs, and phthisical complaints. Instances are mentioned in the German ephemerides, and in Riverius's praxis, of very dangerous phthisical disorders being cured by the continued use of this medicine: in one of these cases, twenty pounds of the conserve were taken in the space of a month, and in another upwards of thirty pounds.

Mixture

Mixtures of the roses with a larger proportion of sugar are made in the Saccharum shops into lozenges: one part of the buds clipt from the heels and rosaceumPh. haftily dried, and twelve parts of fine fugar, are separately reduced into powder, then mixed, and moistened with so much water as will render them of a due confistence for being formed: or the conserve is mixed with as much fresh sugar as is sufficient to bring it to a like consistence, that is, about thrice its own weight.

These flowers give out their virtue both to water and rectified spirit, and tinge the former of a fine red colour, but the latter of a very pale one: the extract obtained by inspissating the watery insusion, is moderately auftere, bitterish, and subsaline; the spirituous extract is confiderably stronger both in astringency and bitterness. In the shops, seven ounces of the dried rose-buds are infused in five pounds of boiling water; and the infusion made into a syrup with six pounds of fine sugart, or boiled to a fyrupy confiftence with feven pounds of clarified honey: the fyrup is valued chiefly for its gratefulness and fine red colour: the mixture with honey is used as a mild cooling detergent, particularly in gargarisms for inflammations and ulcerations of the mouth and tonfils. The infusions acidulated with a little vitriolic acid, and sweetened with fugar, make a grateful, cooling, restringent julep, which is sometimes directed in hectic cases and hemorrhagies, and along with boluses or electuaries of Peruvian bark, and sometimes is used as a gargarism: the college of London orders two pints and a half of boiling water, mixed with a scruple of oil of vitriol, to be poured on half an ounce of the fresh buds, and an ounce and a half of fine fugar to be diffolved in the strained infusion: that of Edinburgh orders two pounds and a half of water, and Infusium half a dram of the acid, to half an ounce of the dry buds and an ounce Tinet. rofaof fugar.

+ Syrup. e rofis ficcis Phar. Edinb. Mel rofaceum t Ph. Lon.

Tinctura rofarum Pb. L.

rum Ph. Ed.

3. CYNOSBATOS Pharm. Lond. Rosa silvestris vulgaris flore odorato incarnato C. B. Rosa silvestris inodora seu canina Park. Cynorrhodon. Rosa canina Linn. Dog-rose, wild briar, hipp-tree: with single pentapetalous flowers, of a whitish colour mixed with various shades of red. It is one of the largest plants of the rose kind; grows wild in hedges; and flowers, as the garden forts, in June.

The flowers of this species, of an agreeable but weak smell, and in taste bitterish and roughish, are said to have a greater degree of laxative

4 C 2

virtue

virtue than those of the damask rose, together with a mild corroborating or restringent quality. The fruit, the only part of the dog-rose
made use of in medicine among us, is agreeably dulco-acid, and stands
recommended as a cooling restringent, in bilious sluxes, sharpness of
urine, and hot indispositions of the stomach: the fresh pulp is made in
the shops into a conserve, by mixing three ounces of it with five of sine
sugar. The pulp should be separated with great care from the rough
prickly matter which incloses the seeds; a small quantity of which,
retained in the conserve, is apt to occasion an uneasiness at the stomach,
pruritus about the ahus, and sometimes vomiting.

Conf. fructus cynosbati Pharm. Lond.

ROSMARINUS.

ROSMARINUS Pharm. Lond. & Edinb. Rosmarinus bortensis angustiore folio C.B. Libanotis coronaria quorundam. Rosmarinus officinalis Linn. Rosemary: a large shrubby plant, clothed with long narrow stiff leaves, set in pairs, of a dark green colour above and hoary underneath; producing pale blueish labiated flowers, which stand in clusters round the stalk in the bosoms of the leaves. It is a native of the southern parts of Europe, common in our gardens, and seems to grow larger and more woody in this than in most other countries. It slowers in April and May, and sometimes again about the end of August.

Rosemary is a warm pungent aromatic; particularly useful in phlegmatic habits and debilities of the nervous system; of the same general nature with lavender, but with more of a camphorated kind of pungency, and of a stronger, and to most people less grateful, smell. The tender tops are the strongest both in smell and taste, and next to these the cups of the slowers; which last, though somewhat weaker than the leaves or tops, are nevertheless the most pleasant, and hence are generally preferred: it is chiefly, if not wholly, in the cup, that the active matter of the flower resides; for the blueish petalum, carefully separated, has very little smell or taste. The fragrance of these flowers is greatly diminished, or in great measure destroyed, by bruising or beating; and hence the officinal conserve, made by beating them with thrice their weight of sugar, has very little of the slavour of the rosemary.

Conferv.flor. rorifmarini Pharm. Lond.

The

The leaves and tops of rosemary give out their virtues completely to rectified spirit, but only partially to water: the spirituous tinctures are of a yellowish green colour, the aqueous of a dark greenish brown. Distilled with water, they yield a thin, light, pale coloured effential oil, Ol. rorismar. inclining a little to yellowish or greenish, of great fragrancy, though not lord for Flair. quite fo agreeable as the rosemary itself: from one hundred pounds of the herb in flower were obtained eight ounces of oil: the decoction, thus divested of the aromatic part of the plant, yields on being inspissated an unpleasant weakly bitterish extract. Rectified spirit likewise, distilled from rosemary leaves, becomes considerably impregnated with their fragrance, leaving however in the extract the greatest share both of their flavour and pungency. The active matter of the flowers is somewhat more volatile than that of the leaves, greatest part of it arising with spirit. The Hungary water, used as a perfume, and sometimes medicinally in nervous complaints, and which is faid to have received its name from its being first made public by an empress of that nation who was cured by its continued use of a paralytic disorder, is a strong spirit distilled from fresh rosemary flowers: the college of Edinburgh directs a gallon Sp.rorismar. of rectified spirit to be drawn over in the heat of a water bath from two pounds of the flowers as foon as they are gathered +: that of London takes the tops, and a spirit not quite so strong; putting a gallon of proof spirit to a pound and a half of the fresh tops, and drawing off in the heat of a water bath five pints ‡. The Hungary water brought from † Pb. Lond. France is more fragrant than fuch as is generally prepared among us.

Lond. & Edin.

RUBIA.

RUBIA TINCTORUM Pharm. Lond. & Linn. Rubia Pharm. Rubia tinctorum sativa C. B. Radix rubra, & erythrodanum MADDER: a rough procumbent plant, with fquare jointed stalks, and five or fix oblong pointed leaves set in form of a star at every joint: on the tops come forth greenish yellow monopetalous flowers, deeply divided into four, five, or fix fegments, followed by two black berries: the root is long, flender, juicy, of a red colour both externally and internally, with a whitish woody pith in the middle. It is perennial, and cultivated in different parts of Europe (in some of which it is said to be indigenous) for the use of the dyers: the roots have been brought

to us chiefly from Zealand; but those which have for some years past been raised in England, appear superiour to the foreign, both as a colouring and a medicinal drug.

The roots of madder have a bitterish, somewhat austere taste, and a slight smell, not of the agreeable kind. They impart to water a dark red tincture, to rectified spirit and distilled oils a bright red: both the watery and spirituous tinctures taste strongly of the madder. The root taken internally tinges the urine and milk red; and in the philosophical transactions, and the memoirs of the French academy of sciences, there are accounts of its producing a like effect upon the bones of animals with whose food it had been mixed: all the bones, particularly the more solid ones, were changed both externally and internally to a deep red, though neither the sleshy nor the cartilaginous parts suffered any alteration. The bones, so tinged, gave out nothing of their colour either to water or spirit of wine.

This root appears therefore to be possessed of great subtility of parts, which may possibly render its medical virtues more considerable than they are now in general supposed to be. It has been chiefly recommended as a resolvent and aperient, in obstructions of the viscera, particularly of the urinary organs, in coagulations of blood from falls or bruises, in jaundices, and in beginning dropsies. * It has lately come into reputation as an emmenagogue, and is said to be a very efficacious medicine of this class (a). From a scruple to half a dram of the powder, or two ounces of the decoction, may be given three or four times a day in this intention.

RUSCUS.

RUSCUS five Bruscus Pharm. Edinb. (b) Ruscus C. B. Ruscus myrtifolius aculeatus Tourn. Oxymyrsine, myrtacantha, myacantha & scopa regia quibusdam. Ruscus aculeatus Linn. Butchers-broom or Kneeholly: a low woody plant, with oblong stiff prickly leaves joined immediately to the stalks: from the middle ribs of the leaves, on the upper side, issue small yellowish slowers succeeded by red berries:

the root is pretty thick, knotty, furnished with long fibres matted together, of a pale brownish colour on the outside and white within. It grows wild in woods and on heaths, is perennial and evergreen, flowers in May, and ripens its berries in August.

THE root of butchers-broom has a sweetish taste, mixed with a slight bitterishness. It stands recommended as an aperient and diuretic, in urinary obstructions, nephritic cases, dropsies, &c. Riverius tells us of an hydropic person who was completely cured by using a decoction of butchers-broom for his only drink, and taking two purges of fena. The virtues of the root are extracted both by water and spirit, and on inspiffating the liquors, feem to remain entire behind: neither of the extracts is very strong in taste; the watery the least so.

R U T A.

RUTA Pharm. Lond. & Edinb. Ruta hortensis latifolia C. B. Ruta graveolens Linn. Rue: a small shrubby plant, with thick blueish green leaves divided into numerous roundish segments: on the tops of the branches come forth yellowish tetrapetalous (sometimes pentapetalous) flowers, followed each by a capfule, which is divided into four partitions full of small blackish rough seeds. It is cultivated in gardens, flowers in June, and holds its leaves all the winter. The mankets are frequently supplied with a narrow-leaved fort, which is cultivated in preference to the other, on account of its appearing variegated during the winter with white streaks.

This herb has a strong unpleasant smell, and a penetrating pungent bitterish taste: much handled, it is apt to inflame and exulcerate the skin. It is recommended as a powerful stimulant, aperient, antiseptic, and as possessing some degree of an antispasmodic power; in crudities and indigeftion, for preferving against contagious diseases and the ill effects of corrupted air, in uterine obstructions and hysteric complaints, and externally in discutient and antiseptic fomentations. Among the common people, the leaves are fometimes taken with treacle, on an empty stomach, as an anthelmintic. A conserve, made by beating the fresh Conserv. for.

leaves ruta Ph. Lon.

leaves with thrice their weight of fine fugar, is the most commodious form for the exhibition of the herb in substance.

The virtues of rue are extracted both by water and rectified spirit, most perfectly by the latter: the watery infusions are of a greenish yellow or brownish; the spirituous, made from the fresh leaves, of a deep green, from the dry of a dark yellowish brown colour: the leaves themselves, in drying, change their blueish green colour to a yellow. On inspissating the spirituous tincture, very little of its flavour rises with the menstruum; nearly all the active parts of the rue remaining concentrated in the extract, which impresses on the palate a very warm, subtile, durable pungency, and is in fmell rather less unpleasant than the herb in substance. Oleum rutæ In distillation with water, an essential oil separates; in colour yellowish or brownish, in taste moderately acrid, and of a very penetrating smell rather more unpleasant than that of the herb: a very considerable part of the virtue of the rue remains behind; the decoction, inspissated, yielding a moderately warm, pungent, bitterish extract. matter of this plant appears therefore to be chiefly of the more fixt kind : the effential oil itself is not very volatile, or at least is so strongly locked up by the other principles, as not to be readily elevated in distillation. The feeds and their capfules appear to contain more oil than the leaves: from twelve pounds of the leaves, gathered before the plant had flowered, only about three drams were obtained; whereas the fame quantity of the herb with the feeds almost ripe yielded above an ounce.

effentialePh. Lond.

Extractum rutæ Pharm. Lond. & Edin.

SABINA.

SABINA Pharm. Lond. & Edinb. Sabina folio tamarisci dioscoridis C. B. Savina quibusdam. Juniperus Sabina Linn. SAVIN: an evergreen shrub or small tree, clothed with very short narrow leaves so stiff as to be prickly; producing small imperfect flowers, and sometimes, when grown old, blueish black berries like those of juniper, of which the modern botany reckons it a species. It is a native of some of the fouthern parts of Europe, and raifed with us in gardens.

THE leaves and tops of favin have a moderately strong smell, of the difagreeable kind; and a hot, bitterish, acrid taste. They give out great part of their active matter to watery liquors, and the whole to rectified rectified spirit; tinging the former of a brownish, and the latter of a dull dark green colour. Distilled with water, they yield a large quantity O1. effent. of essential oil: Hoffman says, that from thirty-two ounces he obtained Lond. Edin. full five ounces of oil, and observes that there is no other known vegetable fubstance, except some of the refinous juices, as turpentine, that affords fo much. The oil fmells strongly, and tastes moderately of the favin: decoctions of the leaves, freed from this volatile principle Extractum by inspissation to the consistence of an extract, retain a considerable share of their pungency and warmth along with their bitterishness, and have likewise some degree of smell, but not resembling that of the plant itself. On inspiffating the spirituous tincture, there remains an extract confifting of two diffinct substances; one yellow, unctuous or oily, bitterish and very pungent; the other black, refinous, tenacious, less pungent, and subastringent.

fabinæ P. L.

Savin is a warm stimulant and aperient; supposed particularly serviceable in uterine obstructions, proceeding from a laxity or weakness of the vessels, or a cold fluggish indisposition of the juices. The distilled oil is accounted one of the most potent emmenagogues: it is likewise a strong diuretic, and, as Boerhaave observes, impregnates the urine with its fmell. * The powdered leaves have been recommended as a very effectual escharotic for confuming warty venereal excrescences (a).

SACCHARUM.

SUGAR: a sweet substance, of a saline nature; prepared from the juice of an elegant large cane or reed, arundo faccharifera C. B. which grows spontaneously in the East Indies and some of the warmer parts of the West, and is cultivated in large plantations in several of the American islands. The expressed juice of the cane is clarified with the addition of lime water, and boiled down to a fomewhat thick confistence: being then removed from the fire, the faccharine part concretes into brown coloured maffes, faccharum rubrum Pharm. Lond. & Edinb. leaving an unctuous liquid matter called melasses or treacle, from which a little more folid fugar, but of a coarfer kind, is obtainable by a repetition of the boiling and clarification. The brown fugar is purified in conical

(a) Med. Eff. Edinb. III. 395.

moulds, by spreading, on the upper broad surface, some moist clay; whose watery moisture, slowly percolating through the mass, carries with it a considerable part of the remains of the treacly matter. The clayed sugar, imported from America, is by our refiners dissolved in water, the solution clarified with whites of eggs, and after due inspissation, poured as before into conical moulds, where, as soon as the sugar has concreted, and the sluid part is drained off by an aperture at the bottom, the surface of the loaf is again covered with moist clay. The sugar, thus once refined, saccharum album Pharm. Edinb. becomes, by a repetition of the process, the double-refined sugar of the shops, saccharum purissimum Pharm. Lond. Solutions of the brown or white sugars, boiled down till they begin to grow thick, and then removed into a very hot room, shoot, upon sticks placed across the vessels for that purpose, into brown or white crystals or candy, saccharum crystallinum Pharm. Edinb.

SUGAR disfolves, by the affistance of heat, in rectified spirit; but greatest part of it separates again in the cold, and concretes into a crystalline form: On this foundation, faccharine concretions are obtained from faturated spirituous tinctures of several of the sweet plants of our own growth; the faccharine part feparating when the tincture is fet in the cold, while the refinous or other matter extracted from the plant, remains disfolved in the spirit. Solutions of sugar mingle uniformly with those of other saline substances, whether acid, alkaline, or neutral; and make no visible alteration in the infusions of the coloured flowers of vegetables, or other liquors, in which acids or alkalies produce a change of colour or a precipitation. This fweet faline fubstance appears on all trials completely neutral *(a); and unites with most kinds of humid bodies, without altering their native qualities: it ferves as an intermedium for uniting together fome bodies naturally repugnant, as distilled oils and water. On the same principle it impedes the coagulation of milk, and the feparation of its butyraceous part.

Sugar, in consequence of this property, is supposed to unite the unctuous part of the food with the animal juices. Hence some have

^{*(}a) An acid of a peculiar kind has been feparated from it in small proportion and by a laborious process, in which the nitrous acid is employed as the separating medium.

concluded, that it increases corpulence or fatness; others, that it has a contrary effect, by preventing the separation of the oily matter, which forms fat, from the blood; and others, that it renders the juices thicker and more sluggish, retards the circulation, obstructs the natural secretions, and thus occasions or aggravates scorbutic, cachectic, hypochondriacal and other disorders. General experience, however, has not shewn, that sugar produces any of these effects in any remarkable degree: its moderate use appears to be innocent; and perhaps, of all that have yet been discovered, it is the most universally innocent and inosfensive, as well as the most simple, sweet.

Sugar preserves both animal and vegetable substances from putrefaction, and appears to possess this power in a higher degree than the common alimentary salt: I have seen animal sless preserved by it untainted for upwards of three years. From this property it has been sometimes applied externally as a balsamic and antiseptic.

The impure brown fugars, by virtue of their oily or treacly matter, prove emollient and gently laxative. The crystals or candy are most difficult of solution, and hence are properest where this soft lubricating sweet is wanted to dissolve slowly in the mouth, as in tickling coughs and hoarseness. The uses of sugar in medicinal compositions, whether for their preservation, for procuring the intended form and consistence, or for reconciling to the stomach and palate substances of themselves disgustful, are too obvious to require being enlarged on.

SAGAPENUM.

SAGAPENUM: the concrete gummy-refinous juice of an oriental plant, of which we have no certain account, but which appears, from the feeds and pieces of stalks sometimes found among the juice as brought to us, to be of the ferulaceous or umbelliferous kind. The sagapenum comes immediately from Alexandria; either in distinct tears, or run together into large masses; outwardly of a yellow colour, internally somewhat paler and clear like horn; growing soft on being handled, so as to stick to the singers. It is sometimes supplied in the shops by the larger and darker coloured masses of bdellium broken in pieces; which greatly resemble

Sensition.

refemble it in appearance, but may be distinguished by their much weaker smell.

SAGAPENUM has a strong disagreeable smell, somewhat of the leek kind, or like that of a mixture of galbanum with a little asafetida; and a moderately hot biting taste. It is one of the strongest of the deobstruent gums, and frequently prescribed, either by itself, or in conjunction with ammoniacum or galbanum, in hysteric cases, uterine obstructions, asthmas, and other disorders. It may be commodiously taken in the form of pills, from two or three grains to a scruple or half a dram: in doses of a dram, it loosens the belly.

On boiling this gummy-resin in water, about three-fourths of it are resolved into a turbid yellowish white liquor, which smells and tastes weakly of the sagapenum. Rectified spirit scarcely takes up above one half, and receives very little colour: the solution smells weakly, and tastes pretty strongly. Both the watery and spirituous solutions lose much, in evaporation, of their taste as well as their smell; the watery loses most, the extract being very considerably weaker than the sagapenum in substance. It is probable that the more active parts are carried off by the watery vapour, but that in the spirituous extract they are only inviscated by the grosser resinous matter: for the water, collected by distillation, is notably impregnated with the flavour of the sagapenum, and discovers likewise a small portion of essential oil; whereas the distilled spirit is almost flavourless.

SALES ALKALINI.

ALKALIES, or ALKALINE SALTS: substances of a very pungent taste; dissoluble in cold water; changing the colours of the blue flowers of vegetables to a green; destroying the acidity of sour liquors, and forming with the acid a neutral compound; precipitating earthy bodies dissolved in acids (a); producing no precipitation or turbidness in so-

lutions

⁽a) To this character of alkalicity there is one exception or limitation. Volatile alkaline spirits made completely caustic by quicklime, on being mixed with a solution of calcareous earth in the nitrous or marine acids, occasion no precipitation or cloudiness. If the mixture be exposed for some time to the air, the alkaline spirit gradually loses its causticity; and then precipitates the earth: on blowing into it air from the lungs, through a glass pipe, the precipitation began immediately.

lutions of the lixivial falts of vegetables. These lixivial salts are themselves alkalies: and to mingle uniformly with these bodies of their
own kind, in a liquid state, is the most commodious and sure mark I
can recollect, for distinguishing alkalies, universally, from certain solutions of earthy bodies in acids; some of which have, in a greater or
less degree, all the common characters of alkalicity; but on being
examined by this criterion, readily betray their composition, by rendering the limpid lixivial liquor milky, and depositing their earth; the
acid, which before held the earth dissolved, being absorbed from it by
the lixivial salt.

I. SAL ALKALINUS FIXUS. Fixt alkaline or lixivial falt: obtained from the ashes of vegetables, by macerating or boiling them in water, and afterwards evaporating the lye till the salt remains dry. It is fixt and fusible in the fire *(a), deliquiates in a moist air, dissolves in equal its weight or less of water, and does not assume a crystalline form (b).

FIXT alkaline falts have an acrid fiery taste, and leave in the mouth a kind of urinous slavour. Saturated solutions of them in water corrode the solid parts of animals, dissolve fats and oils into saponaceous compounds, and liquefy almost all the animal humours, except perhaps only milk, which, when heated, they coagulate. Diluted largely with water, and drank warm in bed, they generally excite sweat: if that evacuation is not favoured by external warmth, they operate chiefly by urine, of which, in many cases, as in maniacal and hydropic ones, they frequently procure a copious and salutary discharge: they likewise loosen the belly, and in costive habits, where the direct purgatives or laxatives give only temporary relief, they render the benefit more lasting. They seem in general to act by stimulating and deterging the solids, and resolving the

^{*(}a) If in fusion a coal falls in, the alkali is resolved into dense white sumes, which act prodigiously on the brain and nervous system, rendering the head weak and benumbed, as in convalescence from some great disease, occasioning impatience and inquietude in every member. Beaumé.

⁽b) Though these salts, as commonly prepared, are never found to shoot into crystals, they do crystallize in part when solutions of them have been exposed for a length of time to the open air. The crystals are far milder in taste, and effervesce more strongly with acids, than the alkali in its common state.

viscidities of the humours; and by these means opening obstructions, or promoting secretion, in all the organs through which they pass. The dose is from two or three grains to sifteen or twenty; in some cases, it has been extended to a dram. That they may be given, and continued for some time, with safety, in very considerable doses, appears from the experience of those, who have taken the strong solution of them called soap-lyes for the relief of calculous complaints. In putrid disorders, and a colliquated state of the humours, these salts have been generally, and I think justly, condemned: for though they have lately been discovered to resist putrefaction both in the sluids and solids of dead animals, yet in living ones they apparently increase the colliquation, with which all putrid diseases are accompanied.

Fixt alkaline salts are obtainable, in greater or less quantity, from almost all vegetables; excepting perhaps only a few of the volatile acrid kind, as mustard seed. The salts of different plants, in the state wherein they are first extracted from the ashes, are found to differ in degree of strength, and in some other respects, from one another; many of them containing a portion, and some a very considerable one, of neutral salts of the vitriolic or marine kind (a). Purished by calcination, so as that all remains of the oil of the vegetable may be burnt out; and by deli-

(a) The readiest way of discovering neutral salt in the lixivial salts of vegetables is, by shaking a strong solution of them in a vial with about an equal quantity of rectified spirit of wine. If the salt is purely alkaline, the two liquors, on standing for a moment, will separate from one another; the spirit rising to the top, and the alkaline solution collecting itself at the bottom, both of them transparent as at sirst. If neutral salts are mixed with the alkali, though in very small proportion, the spirit produces instantly an opake milkiness in the lye; and on standing for a few minutes, a saline matter separates and falls to the bottom, in greater or less quantity, according as the alkali has a greater or less admixture of the neutral salt.

The exact quantity of pure alkali in any kind of lixivial falt or potash may be determined by means of acids. Some alkaline salt known to be pure, as good salt of tartar, is to be melted in an iron ladle, that all remains of watery moisture may be expelled: a certain quantity of this salt, as a dram, weighed out while warm, is to be dissolved in a little water, and saturated with any convenient acid, as diluted spirit of salt: the point of saturation is readily and accurately obtained by means of stained paper, as directed in page 19. For the greater facility in trials of this kind, a quantity of the spirit of salt may be so diluted, that sixteen drams of it for instance may exactly saturate the one dram of pure alkali. If then a solution of one dram of any given salt be saturated with the same acid liquor, so many drams or parts of a dram of the acid as are required for the saturation, so many sixteenths or parts of sixteenths of pure alkali does the given salt contain. This appears to be the most simple and commodious, as well as the most accurate way, that has yet been contrived, for determining the alkalicity, or degree of purity, of all kinds of lixivial salts.

quiation in the air, by which only the alkali diffolves; they are all, except those of some marine plants (see Natron), so much alike, as not to be distinguishable, by any known method of trial, from one another.

The falts of the leaves and other herbaceous parts of plants are more difficultly brought to a state of perfect purity than those of the more woody and compact; a portion of oily matter being tenaciously retained, minute indeed, yet sufficient to give a brownish tinge. A salt of this kind is generally prepared, or expected to be prepared, from wormwood +, +Salabsinth. fometimes from broom, and fometimes from bean stalks, all which are fufficiently well adapted to this use, their ashes yielding as large a proportion as most of the common herbaceous matters, and their falt seeming to be almost merely alkaline, or free from any considerable quantity of the other kinds of faline matter, of which the ashes of some vegetables contain more than they do of alkaline falt. About London, the shops are usually supplied from the country with the ashes of wormwood ready burnt; but that more of the oil may be confumed than the simple burning of the herb has diffipated, they are further calcined with a red heat, and occasionally stirred, for some hours: the white ashes are then boiled in water, and the filtered lye evaporated to dryness.

Some have endeavoured to retain in the falt as much as possible of the oil, by burning the plant with a close smothering heat, continued no longer than till it is reduced fully to ashes; that is, till the alkaline falt is generated, for these salts do not appear to exist naturally in vegetables. The alkalies thus prepared are of a dark brown colour, and supposed to be much milder and less acrimonious, and more of a saponaceous nature, than those which have been further divested of oil. But as we now have, in the foda or natron, an alkali as mild as can be wished for, this inelegant, precarious and unfrugal method of suppressing the acrimony of the common alkalies, becomes unnecessary.

Among all the known vegetables, or vegetable productions, there are none from which a pure alkaline falt is obtainable fo eafily, and in fo large a quantity, as from tartar. If red or white tartar be burnt with a moderately strong fire, either in a proper vessel, or wrapped up in wetted brown paper, to prevent the fmaller pieces from dropping down through the interstices of the coals on being first injected into the furnace, it foon turns to white ashes, which yield on the first elixation a

ftrong

Ph. L. & Ed.

+ Sal tartari strong fiery falt+, of a snowy whiteness, amounting to about one fourth the weight of the tartar. The strength of the salt is somewhat further increased, by keeping it melted for some hours in an intense fire; in which operation, if the crucible cracks or is left uncovered, fo as that the flame may have any access to the falt, or if a minute portion of any inflammable matter is introduced, it assumes, in part at least, a greenish or blue colour, which is commonly looked upon as a mark of its ftrength. -A pure and strong alkaline solution is obtained, by exposing to the air, in a moist place, either the salt, or the white ashes t of tartar: the alkali imbibes in a few days fo much of the aereal moisture, as to run wholly into a liquor, leaving, how highly foever the falt has been purified before, a confiderable quantity of earthy matter. If the liquor be inspissated to dryness, and the dry falt again deliquiated in the air or dissolved in pure water, an earthy matter is still left: and even if the filtered folution be kept for a length of time in a close stopt glass vessel, an earthy substance gradually separates and falls to the bottom.

1 Lixivium tartari Pb.L.

> Alkaline falts are prepared for common uses, in the way of trade, chiefly from wood; of which, in the forests of Germany, Russia, and Sweden, large piles are burnt on purpose. To save the trouble of boiling down the lye, the finer part of the ashes unelixated is in some places tempered with it into the confistence of mortar, which is afterwards stratified with some of the more inflammable kinds of wood, and burnt a fecond time: in others, the lye is foaked up in dry straw, and this drained and burnt. The impure faline masses, obtained by these or fimilar methods, are called Potasbes; the strongest of which has been generally reckoned that brought from Russia (Cineres russici Ph. Lond.) in dark-coloured hard masses, of a very pungent taste, yet containing so much earthy matter as not readily to liquefy or grow moist in the air. This potash is said to be prepared in the first of the ways above mentioned: but it appears from some late experiments, that another ingredient is made use of in the process; the masses, as brought to us, being found to contain more quicklime than alkaline falt (a), and on this depends

⁽a) See Dr. Home's Experiments on bleaching .- It has been suspected that the matter in Russia potash, which seemed from Dr. Home's experiments to be quicklime, is no other than the earth of the vegetable ashes themselves, which earth, by strong calcination, such as this kind of potash is faid to undergo, assumes some of the most striking characters of true quicklime. Since the establishment

depends the great strength and corrosiveness of the Russia potash. For a purer falt, the lye is boiled down in large iron vessels; and the darkcoloured dry falt, which concretes into a hard crust on the sides and bottom of the veffel, is beaten off with a mallet and chifel, and calcined in an oven, with a gradual fire, till it becomes white; in which state it is called, from its pearly appearance, pearl-ash. For some years past, we have been supplied chiefly from our American colonies, with compact alkaline masses, much more pure than the above pot-ashes, though less so than the pearl-ash; prepared by boiling down the lye to dryness, and then increasing the fire till the falt becomes red hot, and melts, so as to be conveniently laded out with iron ladles: the troublesome operation of getting off the indurated falt from the boiler is thus avoided; and the strong melting heat, though of short continuance, supplies in great measure the tedious calcination of the salt; for though the inflammable matter, on which the colour depends, is in fusion not confumed, it is burnt to an indiffoluble coaly state, so that lyes made from these melted potashes with water are nearly as colourless as those of the whitest pearlashes. * The college of Edinburgh, which has discarded the oily Salalkalinus alkalies of wormwood, broom, &c. now directs a pure incinerated alkali to be made from pearl-ashes, first burned with a red heat in a crucible, catus Ph. Ed. then diffolved in water, cleared by fubfidence, and evaporated to drynefs in an iron pot. This falt will diffolve in equal its weight in water, and the folution is analogous to the former oleum tartari per deliquium.

fixus vegetabilis purifi-

Quicklime remarkably increases the activity of all these salts; enabling them, in a liquid or dilute form, to diffolve oils, fats, &c. far more powerfully than either the lime or alkali by themselves; and in a solid or more concentrated one, to act as caustics. For these purposes, the

establishment of the American manufacture, the Russia fort has in this country fallen so much into difuse, that it is very difficult to procure a specimen that can be depended on as genuine. What has been fent to me as true Ruffia potash (and which indeed has greatly the appearance of what used to be fold under that name) on being elixated with water, leaves a large quantity of earthy matter, greatest part of which dissolves readily in aquafortis. This solution has exactly the same tafte with a folution of chalk made in the fame acid: on dropping into it a little vitriolic acid, the liquor grows inflantly milky, and a copious precipitation enfues. This precipitability by the vitriolic acid is one of the properties of calcareous earths, which the earth of vegetables has, not been found to acquire by any degree of calcination; and therefore we may conclude that in the making of this potash real quicklime is mixed, in very large proportion.

ponarium Pharm. Lond.

London college directs equal parts of quicklime and Russia potash (a) to be gradually sprinkled with so much water as will flake the lime: more Lixivium fa- water is then added, the whole stirred together, and suffered to stand for a day or two. The liquor, poured off clear from the undiffolved lime, is the common foap lye; and is directed to be made of fuch a strength, that an exact wine pint may weigh just fixteen ounces troy: if it is heavier, for every dram that it exceeds this weight, an ounce and a half of water by measure is to be added to each pint of the liquor: if lighter, it is to be boiled till the like quantity is wasted, or poured upon fresh lime and potash. The common lyes of our soft-soap makers are considerably stronger than this: Dr. Pemberton observes, that their lyes will be reduced to the strength here proposed, by diluting them with somewhat less than an equal measure of water. * In the Edinburgh pharmacopæia this preparation is thus directed. Eight ounces of fresh quicklime are put into an iron or earthen vessel with twenty-eight ounces of warm water. When the ebullition is over, fix ounces of pure vegetable fixt alkali are added, and after perfect mixture, the veffel is covered and fuffered to cool. The matter is then poured into a glass funnel, lined with a linen rag, and is fet to drain into a glass bottle as long as any liquor will run. Some more water is then to be poured to the matter in the funnel, which will drain through it; and this is to be repeated till three pounds of liquor are procured, which is to be shaken together, and kept in a well stopped phial.

Lixivium causticum Phar. Edinb.

> The dry falt obtained by evaporating these lyes is a strong and sudden caustic: for the greater convenience of using, it is urged in a crucible with a strong fire, till it flows like oil, then poured upon a flat plate made hot, and whilst the matter continues soft, cut into pieces of a proper fize and figure, which are kept in a glass vessel closely stopt. It deliquiates much fooner in the air, and disfolves more readily in watery liquors, than the milder alkalies, and in this confifts its principal inconvenience; being apt to liquefy fo much upon the part to which it is applied, as to spread beyond the limits in which it is intended to operate.

Causticum commune acerrimum Phar. Edinb.

⁽a) Pure alkaline falt requires commonly about twice its weight of quicklime to render it completely caustic. Complete causticity is known by the lye making no effervescence with acids. A redundance of lime is known, by the lye growing milky on dropping into it a little common alkaline lye, or on blowing into it with the breath through a glass pipe.

This inconvenience is avoided, by boiling down the foap lye only to one third+ or fourth t part, and then, while the liquor continues boiling, sprinkling in, by little and little, so much powdered quicklime as will absorb it so as to form a kind of paste: the addition of the lime in substance renders the preparation less apt to liquefy, and hence more easily Pharm. Lond. confinable within the intended limits, but at the same time proportionably more flow in its operation.

+ Causticum mitius P. E. 1 Causticum comm. fort.

2. SAL ALKALINUS VOLATILIS. Volatile alkaline falt: obtainable, by distillation with a strong fire, from all animal matters, from foot, and in fmall quantity from most vegetables: producible also in animal substances, very plentifully in urine, by putrefaction, and in this case feparable by distillation with a gentle heat. When the falt is once formed, whether by ignition or putrefaction, it gradually exhales in moderately warm air; and rifes fooner in distillation than highly-rectified vinous spirits, condensing about the sides of the recipient into crystalline concretions. It requires for its folution three or four times its weight of water.

These salts are in smell as well as taste very penetrating and pungent: they are the only concrete falts that in their pure state emit sensible effluvia. They dissolve oils, refins, fats, &c. more languidly than the fixt alkalies, on account perhaps of their not being susceptible of any confiderable heat, by which their menstrual power might be promoted. In the bodies of animals, they operate more powerfully than the fixt, both as refolvents and stimulants; are more disposed to direct their force to the cutaneous pores, and less to the groffer emunctories; and act more remarkably upon the nervous fystem. They are particularly useful in lethargic and apoplectic cases; in hysterical and hypochondriacal diforders, and the languors, headachs, inflations of the stomach, flatulent colics, and other fymptoms attending those distempers, especially in aged perfons and those of a phlegmatic habit: in languors and faintings, their stimulating smell gives oftentimes immediate relief. In some kinds of fevers, particularly those of the low kind, accompanied with a cough, hoarfeness, redundance of phlegm, and lentor of the blood, they are of great utility; liquefying the thick juices, raising the pulse, and exciting a falutary diaphoresis. In putrid fevers, scurvies, and wherever the mass of blood is thin and acrimonious, they are hurtful: for though they 4 E 2 powerfully

powerfully refift the putrefaction of animal substances, that are detached from the vital economy, yet, in living animals, one of their primary effects is a colliquation of the humours, which in its advanced state is very nearly allied to the advanced state of putrefactive colliquation: their immoderate use has brought on high scorbutic symptoms, resembling those of the true putrid scurvy (a). These salts are most commodiously taken in a liquid form, largely diluted; or in that of a bolus, which should be made up only as wanted, the salt soon slying off. The dose is from two or three grains to ten, twelve, or more.

The volatile alkalies obtained from different substances appear, like the fixt, to be, in their state of perfect purity, one and the same thing. But as first distilled from the subject, they are largely impregnated with its oil rendered setid or empyreumatic by the process in which the salt was generated; and as these oils differ from one another in degree of subtility and setidness, the salts partake of the same differences, till repeated distillations or other processes have either separated the adhering oils, or subtilized and purished them to the same degree. By repeated distillations, all animal oils become limpid as water, lose their setor, acquire a penetrating fragrant smell, and a gratefully pungent taste: thus rectified, they are said, by Dippelius, Hossman, and others, to act, in doses of half a drop, as diaphoretics, anodynes, and antispasmodics. The volatile alkalies used in medicine are expected to be either pure from oil, or to have their oil in this subtilized state.

Oleum animale vulgo.

The oily volatile alkalies have been chiefly prepared from hartshorn, by distillation in large iron pots, with a fire increased by degrees to a strong red heat. At first there arises an aqueous liquor, then the volatile salt, along with a yellowish and at length a dark reddish oil: if the aqueous liquor is not removed before the salt begins to come over, a part of the salt dissolves in it, and thus forms what is called spirit.

The oil, excepting so much of it as is incorporated with the alkali, may be separated from the spirit by filtration through wetted paper, which transmits the spirit and retains the oil. The salt and spirit are then distilled again together, with a very gentle heat, in a glass retort; and the distillation carefully repeated several times, till the salt becomes exceedingly white, and the spirit limpid as water, and of a grateful

finell. The falt becomes the fooner pure, if it be feparated from the fpi- Spiritus &fall rit, and fublimed first from an equal weight of pure chalk and afterwards from a little rectified spirit of wine. If the whole of the volatile salt is required in a folid form, it may be recovered from the spirit by sublimation in a tall narrow cucurbit, the falt rifing into the head, while the watery fluid remains behind. In all the distillations of the spirit, greatest part of the falt comes over before the phlegm; and the process should be continued no longer than till so much of the phlegm has followed as is nearly sufficient to dissolve it; that a part of the salt remaining undiffolved may be a criterion, to the purchaser, of the saturation or strength of the spirit. - A spirit, salt, and oil, are sometimes distilled in the same spirit. sal & manner from wood foot, but here more labour is required to render the falt and spirit pure.

Though the whiteness and limpidity which the falts and spirits of hartshorn, foot, and other like substances, acquire by the above methods of purification, feem to shew that they are divested of oil; they are nevertheless found to participate still of that principle in no small degree. long keeping they contract a yellow colour, and at length become again nauseous and fetid; the oil seeming to be more and more extricated, or to lose by degrees of the fubtility and gratefulness which it received from the rectification. The oftener the distillation is repeated, the more per-

manent is the fubtilization of the oil.

The most effectual purification of these salts is obtained, by combining them with mineral acids, and afterwards separating the acid. It is not needful to make fuch a combination on purpose: for such a one is produced more compendiously, in the way of trade, and called in the shops fal ammoniac: fee the following article.

If fal ammoniac be mingled with any fixt alkaline falt, either in the form of powder or folution, its acid will be absorbed by the fixt alkali; and the volatile alkali, thus fet at liberty again, will immediately discover itself by its pungent odour, and may be collected perfectly pure by distillation. Eighteen ounces of fixt alkali, and one pound of fal ammoniac, Spirit. fal. may be distilled with four pints of water in a gentle heat, till two pints are drawn off +; or fixteen ounces each of the two falts may be distilled + Pb. Lond. with two pounds of water, to dryness t. The volatile alkaline salt may t Ph. Edinbs. be extricated likewise by means of chalk, but with this difference, that the chalk does not begin to act upon the fal ammoniac, or abforb its acid,

Pharm. Lond.

ol. fuliginis Pharm. Lond.

ammon.P.L. Alcali volatile ex fale ammon. vulgo fal ammon. volat. Phar. Edinb.

Sal vol. fal till the mixture is confiderably heated: one part of the fal ammoniac may be mixed with two of chalk, and the mixture fet to sublime in a retort with a strong fire.

Spirit. volatil.causticus. vulgo fp. fal amm. cum calce viv.

Alcali volat. caust. Ph. Ed.

Quicklime, which heightens the pungency of fixt alkalies even to causticity, has a like effect upon the volatile: it renders the fixt more eafily liquefiable, and the volatile permanently liquid, preventing their concretion into a folid form: the volatile alkali, like the fixt, in having its activity thus increased by quicklime, loses its power of effervescing with acids; from whence it may be prefumed, that the lime acts, on one alkali as on the other, by abforbing their air (fee page 176). This pungent volatile spirit may be prepared, by flaking about a pound and a half of quicklime in four pints of water, and adding to the mixture, in a retort, a pound of powdered fal ammoniac: immediately adapt a recipient, (for the pungent vapours begin to arise on the first contact) and with a gentle heat draw off two pints. *Or two pounds of quicklime may be flaked with one pound of water, and covered till it falls to a powder, which is then to be put in a retort with fixteen ounces of fal ammoniac dissolved in four times its weight of water. The vessels are to be very carefully luted, and with a very gentle heat twenty ounces are to be drawn off. This spirit is held too acrimonious for internal use, and has therefore been chiefly employed in fmelling-bottles. It is an excellent menstruum for certain vegetable substances, as Peruvian bark, which the milder spirits extract little from; and when saturated with bodies of this kind, its pungency is fo far sheathed, that it may be taken inwardly with as great safety as tinctures made in the other spirits. In long keeping, unless the bottle is quite full and very closely secured, it gradually imbibes air, as appears from the effervescence which it raises with acids; and loses proportionably of its pungency.

Some have mixed a quantity of this caustic spirit with the officinal spirits both of sal ammoniac and of hartshorn; which thus become more pungent, fo as to bear an addition of a confiderable proportion of water without danger of discovery from the taste or smell. This abuse may be detected, by adding to the suspected spirit a quantity of spirit of wine; which, if the volatile spirit is genuine, will precipitate a part of its volatile falt, but has no fuch effect either on the caustic spirit itself, or on fuch as is sophisticated with it.

Some

Some have fubftituted to the spirit a folution of fal ammoniac and fixt alkaline falt: this liquor eludes the above method of trial, as it deposites a faline matter on the addition of spirit of wine, in the same manner as the genuine volatile spirit: it may be distinguished by the matter, thus deposited, being not volatile but fixt; or by a fixt falt being left upon evaporating a little of the liquor; or more compendiously, by adding a drop or two of folution of filver made in aquafortis, which immediately produces a milkiness in the counterfeit, but makes no apparent change in the genuine spirit.

The addition of spirit of wine to volatile alkaline spirit affords means of judging, in some degree, of their strength or saturation as well as of their purity. If the volatile spirit be fully saturated with salt, a quantity of highly-rectified spirit of wine, poured on it slowly down the sides of the glass, in a cool place, produces immediately an opake dense coagulum on the furface where the liquors touch: on shaking them together, the whole becomes a confiftent mass, which soon resolves by warmth into a fluid and a folid part. This is supposed by some to be a volatile soap, Offa alba composed of the alkaline salt of the one spirit and the oily principle of the other; though in effect it is no more than the alkaline falt itself diflodged, by the vinous spirit, from the watery fluid in which it was diffolved: the quantity of falt, thus separated, will be in proportion to the strength of the volatile spirit.

Though volatile alkalies, not caustic, appear from the above experiments to be averse to any union with vinous spirits; a solution of them even in rectified spirit is nevertheless obtainable, by adding it, along with water, in the process by which they are extricated from the sal ammoniac. For this purpose, three pints of proof spirit are put to four ounces of fal ammoniac, and fix (or less) of any fixt alkaline falt, and one half of the liquor drawn off with a gentle heat +. Or fixteen ounces of quicklime +Spirit. falis and eight of fal ammoniac are powdered together, and put into a retort ammon. with thirty-two ounces of rectified spirit, and the distillation continued 18piritus sal. till all the spirit is come over . This preparation has lately come into nofus Ph. Ed. esteem both as a medicine and a menstruum.

ammon. vi-

Mixtures of volatile and vinous spirits, flavoured with different aromatic oils and other like materials, have long been in general use under the name of fal-volatile. The college of London orders a quart of the spir. volat. above dulcified spirit, two drams of essential oil of nutmegs, the same aromatic. Pharm. Lond. quantity

aromat. vulgo Spir. vo-lat. oleos. & Spir. falinus aromat.P.E. Spir. volat. fætid. + Pb. Lond. 1 Pb. Edinb.

Spirit.volat. quantity of effence of lemons, and half a dram of oil of cloves, to be distilled together with a very gentle heat: that of Edinburgh directs a dram and a half of oil of rosemary, and a dram of essence of lemons, to be diffolved in eight ounces of the dulcified spirit. Volatile spirits are impregnated also in the shops with asafetida, in the proportion of four ounces to five pints+, or one ounce to fixteen ounces 1.

> These kinds of compositions may be made extemporaneously, by dropping any proper effential oils into the dulcified spirit of sal ammoniac, (as now directed above by the Edinburgh college) which will readily dissolve them without the assistance of distillation. By this method, a fal-volatile may be occasionally prepared, of any particular flavour, or adapted to particular purposes: thus, in hysterical disorders, where the uterine purgations are deficient, a preparation of this kind, made with the oils of rue or favin, proves an useful remedy: in weakness of the stomach, oil of mint may be used; and in flatulent cases, those of aniseeds or fweet fennel feeds: thefe last remarkably cover the pungency of the volatile spirit, and render it supportable to the palate. The dose of these compounds is from a few drops to fixty or more.

> The caustic spirit made with quicklime appears in some cases preferable, for these kinds of compositions, to the other volatile spirits; as being perfectly miscible with rectified spirit of wine, in any proportions, without any separation of its volatile alkaline part; and as being a more powerful menstruum for some difficultly soluble oils. The very penetrating pungent volatile spirit, which has lately come into vogue under the name of eau de luce, is faid to be made with this caustic spirit, and oil of amber that has been rectified or redistilled till it becomes limpid and loses its smell: thirty-six drops of the oil, so rectified, are directed to be dissolved in half an ounce of rectified spirit of wine, and fix ounces of the caustic spirit added gradually to this solution: the mixture generally appears milky, and if required limpid may be rendered fo by distillation: fome tinge it of a fine blue colour, when defigned only for fmelling to, by adding a drop or two of folution of copper (a).

> > (a) Malouin, Chimie medicinale, i. 202. ii. 431.

SAL AMMONIACUS.

SAL AMMONIACUS Pharm. Lond. & Edinb. SAL AMMONIAC: a neutral falt; volatile in a moderate heat, but not in that of boiling water; formed by the coalition of volatile alkaline falt with marine acid. On mixing it with a fixt alkaline falt or calcareous earth, and exposing the mixture to the fire, its ingredients are disunited: the volatile alkali exhales, and may be collected in proper vessels (see the foregoing article): and the acid remains combined with the fixt alkali or earth, forming therewith the same compounds as if the pure marine acid had been poured upon them. Hence, in the preceding operations, where the volatile alkali of the sal ammoniac is separated by the intervention of fixt alkalies, the residuum, dissolved and crystallized, is found to be the same with regenerated sea salt; and when chalk or lime is used for the intermedium, the residuum affords calcareous marine salt: see Sal communis.

The other mineral acids form likewise ammoniacal salts with volatile alkali; and it is said, that one made with the vitriolic acid is often substituted to the true officinal one with the marine.—The most obvious character of ammoniacal salts in general is, their yielding a pungent urinous odour on being ground with a little quicklime. The marine sal ammoniac may be distinguished from those made with the other acids, by its emitting white sumes on dropping upon it some oil of vitriol; and by a solution of it in purified aquasortis being able to dissolve gold leaf or a mark made with gold on a touchstone. The nitrous sal ammoniac is distinguished, by its deslagrating or slaming when thrown into a red-hot vessel, dissolving in spirit of wine, and yielding red sumes with oil of vitriol. The vitriolic is distinguished by a solution of it rendering solution of chalk in aquasortis milky; and by its not being acted upon by oil of vitriol.

SAL AMMONIAC has been hitherto prepared chiefly in Egypt: it is faid, that the earth abounds there with marine falt; that grass and other vegetables are sensibly impregnated with this salt; that the dung of graminivorous quadrupeds is used as suel, and the soot carefully collected; and that from this soot, sal ammoniac is extracted, by sublimation without addition. The salt is brought to us, sometimes in conical loaves, most commonly in large round cakes, convex on one side and concave on the

other, appearing when broken of a needled texture, or composed of strice running transversely and parallel to one another: the internal part is generally pure, and of an almost transparent whiteness; the outside for the most part foul and of a yellowish grey or blackish hue. It is purified, either by sublimation, with a gradual fire, in an earthen cucurbit having a blindhead adapted to it; or, perhaps more perfectly, by solution in water, filtration, and crystallization. It dissolves, in temperately warm weather, in about thrice its weight of water, and by the affistance of heat in a much smaller quantity; and crystallizes into long shining spicula, or thin sibrous plates like feathers. In sublimation, especially if the fire is hastily raised, it remarkably volatilizes many kinds of bodies, perhaps all those that are soluble by the marine acid.

† Sal amm. purificat. Phar. Lond.

> This falt has a very sharp penetrating taste. It is a powerful attenuant and deobstruent, feeming to liquefy the animal juices almost like alkaline falts: Boerhaave observes that its liberal and continued use renders the blood fo thin as to burst through the vessels, particularly those of the lungs and the urinary organs. In doses of half a dram or a dram, diffolved in water, if the patient is kept warm, it generally proves sudorific: by moderate exercise, or walking in the open air, its action is determined to the kidneys: in larger doses it loosens the belly. It has by some been held a fecret for the cure of intermittents; and is undoubtedly, in many cases, as an aperient, an excellent affistant to the Peruvian bark, where that aftringent drug by itself would produce dangerous obstructions, or aggravate those already formed. This falt is employed likewise externally as an antiseptic, and in lotions and fomentations for ædematous tumours; as also in gargarisms, for inflammations of the tonsils, and for attenuating and diffolving thick mucus in the mouth and fauces. Saturated folutions of it are faid to confume warts.

SAL CATHARTICUS.

PURGING SALT: a falt of a bitter tafte; foluble in twice its weight or less of water, and shooting into long prismatic crystals like those of nitre; liquefying and bubbling up in a moderate heat, emitting a large quantity of aqueous vapours, and changing to a white spongy mass, bitterer than the salt at first. It is of two kinds; one a combination of the vitriolic acid with the earth called magnesia; the other, a combination

bination of the same acid with the fixt alkali called natron. The most obvious criterion of their acid being the vitriolic is, their precipitating chalk dissolved in aquafortis or in other acids.

1. SAL CATHARTICUS AMARUS Pharm. Lond. & Edinb. Nitrum calcareum Listero & bydrologis quibusdam. Purging bitter salt: composed of the vitriolic acid and magnesia; distinguishable from that whose basis is an alkali, by solutions of it being turned milky, and depositing their earth, on the addition of any alkaline salt.

This is the falt with which the purging mineral waters are principally impregnated, and on which their purgative quality depends. It was first extracted from the Epsom waters, and has been commonly distinguished, both in this and other countries, by the name of Epsom salt: but those waters yielding the salt very sparingly, and their quantity being insufficient for its great demand, it was sought for elsewhere, and sound, in plenty, in the bitter liquor remaining after the crystallization of common salt from sea water; from which it is now generally prepared.

This falt is a gentle purgative, operating in general with ease and fasety, yet with sufficient efficacy, and quickly sinishing its operation: its passing off hastily, and not extending its action so far as most other purgatives, seems to be its principal impersection. For a sull dose, eight or ten drams may be dissolved in a proper quantity of common water, or four or five drams in a pint or quart of the purging waters; to which may be added a little tincture of cardamom seeds, or some other grateful aromatic, to render the liquor more acceptable to the stomach. These liquors, in smaller doses, pass further into the habit, promote the secretions in general, and prove excellent aperients in sundry chronical disorders.

2. SAL CATHARTICUS GLAUBERI, vulgo fal mirabile. Glauber's cathartic falt: composed of the vitriolic acid and the mineral alkali natron, and hence suffering no change from an admixture of fresh alkali.

This falt was discovered, by the chemist whose name it bears, in extracting the acid spirit of sea salt by means of the vitriolic acid. When oil of vitriol is poured on sea salt, the marine acid, thereby disengaged from its own alkaline basis, begins immediately to exhale, and by applying heat may be totally expelled; the vitriolic acid remaining combined

4 F 2 with

Sal cathart.
glauberi
† Ph. Lond.
Soda vitriolata vulgo fal
cath. glauberi † Pharm.
Edinb.

with the natron or marine alkali. This combination is still procured chiefly in the same manner: to the sea salt is added equal to rhalf this weight of oil of vitriol diluted with water, and the marine acid being distilled off, the residuum is dissolved and crystallized. The smallest of these proportions of oil of vitriol appears to be sufficient for expelling the acid and saturating the alkali of the sea salt; but the larger, or rather perhaps one between the two, is more eligible, as the Glauber's salt does not well crystallize unless the acid prevails in the solution.

This falt is nearly of the same medicinal qualities with the foregoing, which frequently supplies its place in the shops. The Glauber's salt, somewhat the least unpleasant to the taste, is supposed to be the mildest of the two, and to operate the most kindly.

SAL COMMUNIS.

COMMON or Culinary salt; called, from its most obvious fource, sea salt; though found also, in immense quantities, in the bowels of the earth. It is a persectly neutral salt, composed of a peculiar acid denominated from it the marine acid, and of the mineral alkali natron. It dissolves in less than thrice its weight of boiling water, and does not, like the other neutral salts, concrete again in the cold, so long as the evaporation of the sluid is prevented; cold water dissolving nearly as much of this salt as boiling water. By gentle continued evaporation it shoots into cubical crystals, several of which unite together into the form of hollow truncated pyramids. The crystals, exposed to the fire, burst and crackle; soon after melt, and appear thin and limpid as water: if the salt be melted along with other susible salts or with vitreous matters, it does not persectly unite with them, but flows in part distinct upon the surface. After suffering a considerable heat, it liqueses in the air.

\$ Sal decrepitatum.

1. SAL GEMMÆ Pharm. Lond. Sal gem, rock falt, fosfil common falt. This is met with in several parts of the world, but in greatest plenty in certain deep mines, of prodigious extent, near Cracow in Poland: some is likewise found in England, particularly in Cheshire. It is for the most part very hard; sometimes pure, transparent and colourless; more commonly mixed with earthy or stony matters, of an opake whiteness, or of a red, green, blue, or other colours. These

last forts are purified, for the common uses of salt, by solution and crystallization.

2. SAL MARINUS Pharm. Lond. Sal marinus hispanus Pharm. Edinb. The falt extracted from fea water and faline springs. Sea waters yield from one fiftieth to about one thirtieth their weight of pure falt: from feveral springs much larger quantities are obtained: those in our own country at Nantwich, Northwich, and Droitwich, afford from one fixth to one third their weight. Sea water contains, besides the common salt, a portion of purging bitter falt, and of another faline fubstance which remains diffolved after the crystallization of the latter, of a very pungent taste, scarce reducible into a crystalline form, composed of marine acid and calcareous earth: from both these salts the spring waters are usually free. There are two general methods of extracting the common falt from these natural solutions of it: the one, a hasty evaporation, continued till the falt concretes and falls in grains to the bottom of the pan, from whence it is afterwards raked out, and fet to drain from the bittern: the other, a flow and gradual evaporation, effected in the warmer climates by the fun's heat, by which the falt is formed, not into fmall grains, but into large crystals, called bay-falt. The falts obtained by these two processes differ in some respects from one another: that got by hafty evaporation, especially if a boiling heat, or one approaching to it, be continued during the time of the falts concreting, is apt to liquefy in a moist air; an inconvenience which the crystallized fort is not subject to: the crystals are found likewise to be stronger than the other, and to answer better for preserving provisions. Both forts prove impure and brown-coloured if the folutions are evaporated directly, but of perfect whiteness if previously clarified by boiling with a little ox blood, or other like fubstances, which concreting by the heat, inviscate the unctuous matter, and carry it to the furface in form of fcum. Both forts generally retain a portion of the bitter falt; whose basis being an earth, folutions of them deposite this earth on the addition of any alkali.

COMMON falt differs from other faline substances in occasioning drought, and tending, not to cool, but rather to heat the body. It prevents putrefaction less than most others, and in small quantities, such

as are taken with food, promotes it: by this quality it probably promotes also the resolution of the aliment in the stomach, at the same time that it proves a mild stimulus to that viscus itself. Salted animal foods are generally, perhaps juftly, accounted one of the principal causes of the scurvy at sea; not that the salt is of itself prejudicial, but on account of its being incapable of preferving the animal subjects, for a length of time, in a perfectly uncorrupted state. Pure sea falt, and sea water, are rather falubrious than hurtful, both in the true fcurvy, and in impurities of the blood and humours in general. In confiderable doses, they act as purgatives: Hoffman observes, that an ounce of the falt, diffolved in a proper quantity of water, occasions commonly fix stools or more, without uneafiness; that this falt checks the operation of emetics, and carries them off by stool; that in glysters it is more effectual, though used only in the quantity of a dram, than any of the purgatives; and that where other glysters fail of opening the belly, a folution of common falt takes place.

* A remarkable instance of the efficacy of common salt given in very large doses in a worm case, is related in the Medical Transactions of the London College, Vol. I. A person reduced to the utmost extremity with pain in the stomach, obstinate constipation, and contracted limbs, was advised, after many remedies had been used in vain, to drink salt and water. He drank within an hour two pounds of salt mixed in two quarts of water. It occasioned violent vomiting, which brought up a quantity of small worms, and its operation ended with purging and a profuse sweat. Great rawness and soreness of the gullet and stomach, with unquenchable thirst, and dysury, remained. These symptoms went off by free dilution; and he ventured the third day after to repeat the dose of salt, which had effects similar to the former. A persect cure was the consequence of this singular practice.

THE common forts of fea falt, contrary to other neutral ones, part with a little of their acid in the boiling down of folutions of them to drynefs (a). To this cause are attributed the weakness of the falt pre-

⁽a) This acid appears to proceed, not from the pure marine falt, but from the calcareous muriatic falt, or a combination of marine acid with earth, which all the common forts of fea falt are found to partake of. On dropping into a folution of common fea falt a little alkaline lye, the earth precipitates; and the acid, being thus faturated with alkali, is no longer disposed to evaporate on boiling down the liquor.

pared by that process, and its disposition to deliquiate in the air; both which imperfections are faid to be corrected by a finall addition of fresh acid when the falt begins to concrete. Hence also distilled sea water is manifestly impregnated with acid, fo as to be unfit for drinking or for the common purposes of life; unless a little chalk, vegetable ashes, or other like fubstances, be added in the distillation, to absorb and keep down the acid extricated by the heat *(a): by this means the distilled

fluid proves perfectly sweet.

The acid of fea falt is completely difengaged from its alkaline basis by the more powerful acid of vitriol; and may now be collected, in a concentrated state, by distillation; but as, in this concentrated state, its fumes very difficultly condense, a little water is commonly added to promote that effect. On two pounds of dry fea falt, the college of Spirit. falis London directs two pounds of oil of vitriol diluted with a pint of water, that of Edinburgh one pound of oil of vitriol diluted with equal its quantity of warm water, to be poured by little and little, under a go fpir. fall chimney, that the operator may not be incommoded by the noxious fumes: the retort is placed in fand, and the distillation performed with a fire gradually increased till nothing more will arise. The spirit may be freed from its superfluous water, by a second distillation in a glass cucurbit; the phlegmatic part rifing in a proper degree of heat, while the stronger acid remains behind. The distilled spirit proves nearly the fame, whether the larger or smaller of the above proportions of oil of vitriol are used, the difference affecting chiefly the residuum: see the foregoing article.

The marine acid is diftinguished from the others, by its rifing in white fumes; by its peculiar pungent fmell; by its enabling the nitrous acid to diffolve gold, preventing its diffolving filver, and precipitating filver previously disfolved, but producing no precipitation in folutions of calcareous earths. It is fometimes given, from ten to fixty or feventy drops, properly diluted, as an antiphlogistic, diuretic, and for promoting appetite, and applied externally to chilblains, which are faid by Linnæus to be radically cured by it in a short time, without fear of a return; but its

principal use is in combination with other bodies.

marini. Ph. Acidum muriaticum vulmarin. Ph.

^{*(}a) This affertion is contradicted by the fuccefs of fome late attempts to supply ships with fresh water by the distillation of sea water. In these, good sweet water was obtained merely by fitting an apparatus to the ships' boilers in which falt water was used for common purposes. See, particularly, Phipp's Voyage towards the North Pole. Cambined

Spirit. falis marini coag. Ph. Lond. & Wirtem. Sal marin. rege-

Sal ammon. fixum vulgo. Sal muriatic. calcareus.

Combined with volatile alkalies, it produces the officinal fal ammoniac. With the mineral fixt alkali, it regenerates common falt. With vegetable fixt alkalies, it forms a neutral falt of a sharper taste, and somewhat more difficult of fusion and folution, than common falt: this combination is prepared in the shops, by dropping into the marine spirit a lixivium of the fixt alkali till all effervescence ceases, and then evaporating the mixneratusvulgo. ture to dryness: the same salt may be obtained from the matter which remains after the distillation of spirit of sal ammoniac with fixt alkali.

With calcareous earths, it forms a very pungent faline compound, which difficultly assumes a crystalline form, deliquiates in the air, diffolves not only in water but in rectified spirit of wine, and changes the colour of blue flowers of vegetables to a green. This falt is contained, in confiderable quantity, in fea water, and remains fluid after the crystallization of its other faline matters: it is found also in fundry common waters, to which, like the calcareous nitre, it communicates, according to its quantity, a greater or less degree of hardness and indisposition to putrefy: it is far more antiseptic than the perfect marine salt. It is faid to be diuretic and lithontriptic: the medicine commonly fold as a lithontriptic under the name of liquid-shell, appears to be no other than a combination of this kind, confifting of calcined shells dissolved in marine acid. These combinations have been chiefly prepared, by mixing the calcareous earth with fal ammoniac, and urging the mixture with a gradual fire, till the volatile alkali of the fal ammoniac is either diffipated in the air or collected by distillation, and only its acid left incorporated with the earth: fo much of the earth, as is fatiated with the acid, may be separated from the rest by elixation with water.

This acid diffolves, among metallic bodies, zinc and iron pretty readily; copper and tin languidly; bifmuth and arfenic very difficultly and fparingly; lead, mercury, regulus of antimony, and filver, not at all, unless highly concentrated and applied in the form of fume: it diffolves, by digestion, all metallic bodies when reduced to a state of calx, gold not excepted. Though it difficultly unites with metals, it adheres to most of them more strongly than any other acid, and in part volatilizes them: it renders them likewise more fusible in the fire than the other acids do,

and more disposed to solution in spirit of wine.

Of itself it is nevertheless the most averse of all acids to a perfect union with vinous spirits. If poured gradually into thrice its quantity of rectified spirit of wine, and the mixture, after digestion for some days, submitted to distillation in a sand heat; the spirit that comes over, appears to be little other than the acid simply diluted with the vinous spirit; whereas, when the nitrous or vitriolic acids are treated in the same manner, a new compound is formed by the intimate coalition of the acid spirit with the inflammable (a). The dulcified marine acid has by some been held in great esteem against weakness of the stomach, indigestion, and other like complaints brought on by irregularities.

SALIX.

SALIX Pharm. Edinb. Salix vulgaris alba arborescens C.B. Salix Ger. Salix fragilis Linn. Common white willow: a pretty large tree, frequent in moist woods and hedges; producing loose spikes or catkins, either of imperfect barren flowers, or of seeds inclosed in down: it is the largest of the willows; and differs from the others, in the oblong pointed serrated leaves being hoary on both sides, though most so on the lower, and in the branches not being tough.

The bark of this tree has lately been found an useful medicine in agues, of which many persons have been cured by taking a dram of the powdered bark every four hours during the intermissions, though in some cases it was necessary to join to it a little Peruvian bark (see the Philosophical Transactions for the year 1763). To the taste this bark discovers a pretty strong bitterness and astringency: with solution of chalybeate vitriol, it strikes an inky blackness. These observations serve therefore to confirm what has been remarked under the head of the Peruvian bark, that in the distempers where that valuable medicine takes place, other bitters and astringents are likewise useful, though in an inferiour degree. The astringency of the willow bark is extracted both by water and spirit; and

⁽a) It is faid that the marine acid may be combined with vinous spirits as intimately as the others, and an ethercal fluid produced from the mixture, by applying the acid spirit to the vinous while both are resolved by fire into vapour; or more commodiously, by using the acid in a high degree of concentration, such as is obtained by distillation from a mixture of mercury sublimate with tin, commonly called the smoking spirit of Libavius, and proceeding with this spirit and spirit of wine, in the same manner as with the other acids. It is supposed that this acid, in distillation from metallic substances, takes up a portion of the inflammable principle of the metal, which promotes its union with the vinous spirit.

the black matter, produced by adding vitriol to the watery infusion, is not disposed to precipitate.

SALVIA.

SAGE: a low shrubby plant, with square stalks; obtuse wrinkled dry leaves set in pairs, and large blueish labiated slowers, in loose spikes on the tops of the branches: the upper lip of the flower is nipt at the extremity, the lower divided into three segments. It is a native of the southern parts of Europe, common in our gardens, and slowers in June.

- 1. SALVIA Pharm. Lond. & Edinb. Salvia major C. B. Salvia officinalis Linn. Common fage: with the leaves nearly oval, but acuminated, fometimes green and fometimes red: both the green and red forts rife from the feeds of one and the fame plant.
- 2. SALVIA HORTENSIS MINOR. Salvia minor aurita & non aurita C. B. Small fage, or fage of virtue: with narrower leaves, generally whitish, never red: most of them have at the bottom a piece standing out at each side in the form of ears. This is a variety of the former.

THE leaves and tops of fage are moderately aromatic and corroborant, and used in debilities and relaxations both of the nervous and vascular system. Their smell is pretty strong and not disagreeable; their taste somewhat warm, bitterish, and subastringent: with solution of chalybeate vitriol, they strike a deep black colour. The second fort is both in smell and taste the strongest, the first most agreeable. Of both kinds, the slowers are weaker and more grateful than the leaves; and the cup of the slower stronger, and obviously more resinous, than any other part.

The leaves of fage give out their virtue both to water and rectified spirit, most perfectly to the latter; to the former they impart a brownish, to the latter a dark green tincture. The watery insusion is often used as tea, and often acidulated with a little lemon juice for a diluent in febrile distempers: the spirituous tincture is in taste stronger than the watery, but the smell of the sage is by this menstruum covered or suppressed. The leaves and slowery tops, distilled with water, yield a small quantity

of effential oil, smelling strongly and agreeably of the herb, in taste very warm and pungent, when newly distilled of a fine greenish colour, by age turning yellow or brown: the remaining decoction, divested of this aromatic and most active principle of the sage, yields an extract weakly bitterish, subastringent, and subsaline. The spirituous extract, in smell weak and somewhat different in kind from that of the herb itself, discovers to the taste a considerable aromatic warmth and pungency, refembling that of camphor, but milder.

SAMBUCUS.

ELDER: a plant, with finely ferrated sharp-pointed leaves, set in pairs on a middle rib, with an odd one at the end; producing, on the tops of the branches, umbel-like clusters of small white flowers, followed each by a juicy berry, containing generally three seeds.

1. Sambucus Pharm. Lond. & Edinb. Sambucus fructu in umbella nigro C. B. Acte. Sambucus nigra Linn. Elder tree: with nearly oval leaves, of which five or seven stand on one rib. It is a small tree or shrub, covered with an ash-coloured chapt bark, under which lies a thinner green one, and under this a white: it grows wild in hedges, slowers in May, and ripens its black berries in September.

The bark of this tree is recommended as a strong hydragogue in hydropic cases. Sydenham directs three handfuls of the inner bark to be boiled in a quart of milk and water till only a pint remains, of which one half is to be taken in the morning, and the other at night, and this repeated every day: he observes, that this medicine operates both upwards and downwards; and that if it does not vomit or purge at all, or but gently, it does no service. Boerhaave says, that the expressed juice of the middle bark, given from a dram to half an ounce (some go as far as an ounce), is the best of hydragogues where the viscera are sound; and that it so powerfully dissolves the humours, and procures so plentiful watery discharges from all the emunctories, that the patient is ready to faint from the large and sudden inanition. The decoction and juice are recommended also, in smaller doses, as useful aperients and deobstruents in different chronical disorders. This bark has scarcely any smell, and very little taste: on first chewing, it impresses a kind of

4 G 2 fweetish-

fweetishness, which is followed by a very slight but very durable acrimony, in which its medical activity seems to reside, and which it imparts both to watery and spirituous menstrua.

The leaves, of a faint unpleasant smell, and a strong, very nauseous, bitter kind of taste, are said to be purgative and emetic like the bark. They are celebrated externally against burns and inflammations, and for these purposes an ointment has been prepared for them in the shops: four ounces of the leaves, and the same quantity of the inner bark, fresh, were thoroughly bruised, and boiled in a quart of linseed oil till the watery moisture was consumed and the oil tinged of a green colour: the oil was then pressed out, and brought to the consistence of an ointment by melting in it six ounces of white wax.

The flowers of elder have an agreeable flavour, which they give over in distillation with water, and impart by infusion both to water and rectified spirit: on distilling with water a large quantity of the flowers, a small portion of a butyraceous essential oil separates. Insusions made from them while fresh are gently laxative and aperient: when dry, they are said to promote chiefly the cuticular excretion, and to be particularly serviceable in erysipelatous and eruptive disorders. From these also an unguent is prepared, probably of equal efficacy with the other, and preferred by some as being more elegant, by melting three pounds of mutton sut that a pint of oil-olive, and boiling in this mixture four pounds of the full blown flowers till they are almost crisp.

Unguentum fambucinum Pharm. Lond.

The berries in take (weetilb and not unpleafar

Rob baccar. fambuci + Ph. Lond. 1 Ph. Edinb.

The berries, in taste sweetish and not unpleasant, yield on expression a fine purplish juice, which inspissated to the consistence of honey, either by itself to with the addition of half a pound of fine sugar to two pounds and a half to proves an useful aperient and resolvent in recent colds and sundry chronical disorders, gently loosening the belly, and promoting urine and perspiration.

2. EBULUS Pharm. Edinb. (a) Sambucus bumilis sive ebulus C. B. Chamæacte. Sambucus Ebulus Linn. Dwarf-elder or Danewort: an herbaceous plant, dying to the ground in winter; with longer leaves than those of the elder tree, and nine leaves on one rib. It grows wild in some parts of England, slowers in July, and produces ripe black berries in the beginning of September.

It is faid that this species has the same virtues with the preceding, but differs somewhat in degree: that the bark (that of the root has been chiefly used) and the berries, are respectively more efficacious, and the leaves less so: that the rob or inspissated juice of the berries, in doses of half an ounce to an ounce, acts as a strong hydragogue, and in smaller doses as a powerful resolvent and deobstruent.

SANGUIS DRACONIS.

SANGUIS DRACONIS Pharm. Lond. & Edinb. Cinnabaris Græcorum. DRAGONS-BLOOD, so called: a resin, obtained from certain large palm-like trees (Palmijuncus Draco Rumph. amb. Calamus Rotang Linn.) growing in the East Indies; brought over in oval drops wrapt up in flag-leaves, or in large and generally more impure masses composed of smaller tears; of a deep dark red colour, which changes, in pulverization, to a crimfon. Sundry artificial compositions, coloured with the true dragons-blood, or with brazil wood or other materials, have been fometimes fold in the room of this commodity: fome of these dissolve like gums in water, and others crackle in the fire without proving inflammable; whereas the genuine dragons-blood readily melts and catches flame, and is scarcely acted on by watery liquors. It dissolves almost totally, by the affistance of heat, in rectified spirit, and tinges a large quantity of the menstruum of a deep red colour: it is likewise foluble in expressed oils, and imparts to them a red tincture, less beautiful than that which anchusa communicates.

This refin, in fubstance, has no perceptible smell or taste: when dissolved, whether in vinous spirits or in oils, it discovers some degree of pungency and warmth. It is usually looked upon as a gentle incrassant, desiccative, and restringent; and sometimes prescribed in these intentions against alvine and uterine sluxes, and ulcerations both internal and external.

SANICULA.

SANICULA seu Diapensia Ph. Ed. (a) Sanicula officinarum C. B. Sanicula europæa Linn. SANICLE: an umbelliferous plant, with shining

dark green roundish leaves, cut into five segments, serrated about the edges; and rough seeds, which stick to the clothes. It is perennial and evergreen, grows wild in woods on hilly grounds, and flowers in May.

This herb, recommended both externally and internally as a vulnerary or mild restringent, and supposed to have received its name from the sanative virtues ascribed to it, discovers to the taste a kind of bitterishness and roughness, followed by an impression of acrimony which affects chiefly the throat: in the fresh leaves, the taste is very weak; in the dry leaves, considerable; in the extracts made from them, by water and spirit, moderately strong.

SANTALUM.

SAUNDERS. Three different woods are brought under this name from the East Indies, in large billets: they are said to be the produce, chiefly, of the island Timor in the Indian ocean.

- Yellow faunders: of a pale yellowish or brownish colour, and a close even grain. This wood has a pleasant smell, and a bitterish aromatic taste accompanied with an agreeable kind of pungency. Distilled with water, it yields a fragrant essential oil, which thickens in the cold into the consistence of a balsam, approaching in smell to ambergris, or a mixture of ambergris and roses: the remaining decoction, inspissated to the consistence of an extract, is bitterish and slightly pungent. Rectified spirit extracts, by digestion, considerably more than water: the colour of the tincture is a rich yellow. The spirit, distilled off, is lightly impregnated with the fine slavour of the wood: the remaining brownish extract has a weak smell, and a moderate balsamic pungency. This wood therefore, though at present among us disregarded, promises to have a good claim to the corroborant virtues ascribed to it by Hossman and others.
- 2. SANTALUM ALBUM. White faunders: of a close texture and straight fibres like the preceding, but of a paler whitish colour. This

species,

^{*(}a) The Santalum album, below, is faid not to be a wood of a different species, but the alburnum of the trunk of the same tree, the medullary part of which is the citrinum.

species, far weaker than the yellow, both in smell and taste, promises very little medicinal virtue: it has long been entirely neglected, and is now rarely to be met with in the shops.

3. Santalum rubrum Pharm. Lond. & Edinb. Pterocarpus Santolinus Linn. Suppl. Red faunders: of a dull red almost blackish colour on the outside, and a deep brighter red within: its sibres are now and then curled, as in knots. This also, recommended as an astringent and corroborant, appears to be of very little virtue, as it has no manifest smell, and little or no taste: even of extracts made from it, with water or with spirit, the taste is inconsiderable. Its principal use is as a colouring drug. To watery liquors it communicates only a yellowish tinge, but to rectified spirit a fine deep red: a small quantity of an extract made with this menstruum tinges a large one of fresh spirit of the same elegant colour; though it does not, like most other resinous bodies, dissolve in expressed oils, or communicate its colour to them: of distilled oils, there are some, as that of lavender, which receive a red tincture both from the wood itself and from the resinous extract, but the greater number does not.

Geoffroy and others take notice that the brazil woods are sometimes substituted to red saunders, and the college of Brussels doubts whether all that is sold among them for saunders is not really a wood of that kind. According to the account which they have given of their red saunders, it is plainly the brazil wood of the dyers; the distinguishing character of which is, that it imparts its colour to common water. Of the same kind also is the wood examined by Cartheuser under the name of red saunders, the watery insusion and extract of which were both of a dark red.

SANTONICUM.

SANTONICUM SEMEN Pharm. Edinb. Semen cinæ, semen sanctum, semen contra, sementina. Wormseed: a small light oval seed; composed as it were of a number of thin membranous coats; of a yellowish-greenish colour with a cast of brown; easily friable, by rubbing between the singers, into a fine chaffy kind of substance. The seeds have commonly mixed with them a considerable quantity of this chaffy matter, and small bits of stalks and leaves. They are brought from the Levant, and supposed to be the produce of a species of artemisia, resembling in its general appearance our fine-leaved mugwort, called by Linnæus Artemisia (Santonicum) foliis caulinis linearibus pinnato-multisidis, ramis indivisis, spicis secundis restexis; the Artemisia austriaca of Jacquin.

THESE feeds have a moderately strong, not agreeable smell, somewhat of the wormwood kind; and a very bitter subacrid taste. They have been chiefly recommended as anthelmintics; and commonly taken, in this intention, either along with melafies, or candied with fugar. They might be used also for other purposes; as they appear (at least the specimens which I examined) to be a not inelegant strong bitter. They give out their virtue both to water and spirit, together with a brownishhue, which in the watery tincture has an admixture of reddiff, in the spirituous of yellow: the spirituous is less ungrateful in taste, and discovers less also of the ill smell of the santonicum than the watery infusion. In evaporation, water carries off greatest part of the disagreeable flavour of the feeds, the inspissated extract being little other than simply bitter. An extract made by rectified spirit retains a considerable share of the flavour: this extract appears to be the most eligible preparation of the fantonicum for the purposes of an anthelmintic; and the watery extract, or a tincture drawn from it, for the more general intentions of bitter medicines.

SAPO.

SOAP: a composition of oils or fats with alkaline salts, incorporated so as to dissolve together in water into a milky semitransparent liquid.

Sapo amygdalinus P. L.

1. SAPO DURUS. Hard foap. The finest hard soap is prepared with fresh-drawn oil of almonds, by digesting it with thrice its measure of the soap-lyes formerly described (see Sales alcalini) in such a heat that they may just simmer. In a few hours they unite into a turbid sluid, which, on being boiled a little, becomes more transparent, and ropy, so that if a little be suffered to cool, it will concrete like gelly. Some sea salt is now thrown in, till the boiling liquor loses its ropiness; and the coction continued till, on receiving some drops upon a tile, the soap is found to coagulate, and the water to freely separate from it. The sire being then removed, the soap rises gradually to the surface; from whence

it is taken off before it grows cold, and put into a wooden mould, or frame, with a cloth bottom: being afterwards separated from the mould, it is fet by till it has acquired a due confistence. After the same manner a hard foap is made with oil-olive, which should be of the finest kind, Sapo ex oleo that the foap may prove as little ungrateful as possible either to the palate or stomach. By the same or similar processes this commodity is prepared Sapo durus for common uses in the way of trade. The finest of the common soaps is that called Spanish or Castile soap, which is made with oil-olive, and the alkaline falt called foda or barilla: our foap-boilers find that this alkali gives a better confistence, or greater hardness to the soap, than the mat. med. other potashes or common vegetable alkalies.

olivarum Ph. Phar. Lond. mat. med. Sapo albus hifpanus Pb. Edin.

Hard foap, triturated with vegetable refins and thick balfams, incorporates with them into a compound, foluble, like the foap itself, in watery liquors: hence it proves an ufeful ingredient in refinous pills, which of themselves are apt to pass entire through the intestines, but by the admixture of foap become diffoluble in the stomach. It renders unctuous and thick mucous animal matters dissoluble in like manner in aqueous fluids, and hence may be presumed to act as a menstruum for these kinds of substances in the body, that is to attenuate viscid juices and resolve obstructions: such, in effect, are the virtues which it appears to exert in cachectic, hydropic, and icteric cases, in which last, particularly, its aperient and resolvent powers have been often experienced. Solutions of it have been found likewise to dissolve certain animal concretions of the harder kind, as the filaments which are fometimes feen floating in the urine of rheumatic and arthritic persons, the matter secreted in gouty joints, and the more compact urinary calculus: on these substances (at least on the latter) though soap of itself acts more languidly than limewater, yet when joined to that menstruum it remarkably increases its activity, the diffolving power of a composition of the two being, according to Dr. Whytt's experiments, confiderably greater than that of the foap and lime-water unmixed: of the good effects of these medicines in calculous cases there are several instances; but what their effects may be in gouty and rheumatic ones, is not yet well known.

The usual dose of soap, as an aperient, is half a dram or a dram: as a lithontriptic, half an ounce, or an ounce, or more, are taken in a day at proper intervals. It is given in the form of a bolus or pills; or made into an electuary with some grateful syrup, as that of orange peel; or diffolved

4 H

dissolved in milk or other liquids. *It is excellently covered by chocolate: two drams in a pint are not in the least perceived; the chocolate is thought by some better than without it. A little soap is always added in the composition of chocolate, to make it froth.

In watery liquors it dissolves only imperfectly, the solution being always turbid. Rectified spirit, though it has no action on the alkaline salt or oil separately, dissolves the soap into a limpid liquor. Proof spirit, free from acidity, dissolves it as perfectly, and in much larger quantity; rectified spirit not taking up one tenth its own weight, but proof spirit one third or more. The spirituous solutions bear to be largely diluted with pure water, without suffering any turbidness or separation of their parts: but on the addition of any acid, or of any combination of acids with earthy or metallic bodies, as the salt catharticus amarus, &c. the soap is resolved into its constituent ingredients; its alkaline salt being absorbed by the acid, and the oil rising to the surface. The oil, thus extricated from soap by acids, dissolves like essential oils, in rectified spirit.

Soap is employed externally for discussing rheumatic pains, arthritic tumours, the humours stagnating after sprains, &c. Some pretend that the indurated tophaceous concretions in arthritic joints have been refolved by the external use of soapy cataplasms. Several compositions for external purposes are prepared in the shops. One part of Spanish soap, shaved or cut in thin slices, is stirred into fix parts of common plaster melted over the fire, and the mixture boiled till it acquires the confisence of a plaster; which is formed into rolls whilst hot, the soap disposing it to grow brittle as it cools +: fome endeavour to promote the refolvent virtue of the foap, by adding to four parts of the common plaster, two of gum plaster, with one of foap t. But foap acts to much better advantage in the form of a cataplasm or liniment, than in the stiff one of a plaster. The officinal saponaceous liniments are made, by digesting three ounces of Spanish soap in a pint of spirit of rosemary till the spirit is faturated, and diffolving in this folution an ounce of camphor || ; or by digesting two ounces of soap in a pint of rectified spirit of wine, and afterwards adding an ounce of camphor, and two drams of oil of rofemary §. Sometimes opium is joined, by which the compound is fupposed to be rendered more effectual for allaying violent pains: half an ounce of opium is digested with the soap in the last mentioned composition. This is given also internally, in nervous colics, jaundices, &c. 2. SAPO

Emplastrum † e sapone Phar. Lond. ‡ saponaceum Ph. Ed.

Linimentum faponaceum || Pb. Lond.

vulgo Balf.
faponaceum
§ Ph. Edin.
Linimentum
anodynum
vulgo Balf.
anodyn. Ph.
Edinb.

2. SAPO MOLLIS. Soft foap. The common foft foap used about Sapo mollis London, generally of a greenish hue with some white lumps, is prepared chiefly with tallow: a blackish fort more common in some other places, is faid to be made with whale oil. Both kinds are confiderably more acrid than the hard foaps, and are employed only for fome external purposes: a mixture of equal parts of our common foft soap and quick- + Caustic. lime is used as a mild caustic +.

com. mitius Pharm. Lond.

3. SAPO VOLATILIS. Volatile foap. Of this there are three kinds: one composed of fixt alkalies and volatile oils; another, of volatile alkalies, and oils of the groffer or more fixt kind; and the third, in which both the alkali and the oil are volatile.

Fixt alkalies are very difficultly made to unite with diffilled oils. The most commodious method of obtaining the combination appears to be, by throwing the falt red hot into a heated mortar, immediately reducing it into powder, then pouring on it, while it continues quite hot, by little at a time, an equal quantity or more of the oil, and continuing to grind them together, fo as to form a fmooth foft mass. Stahl reports that the union may foon be obtained also, by agitating the falt with a fmall proportion of the oil, and a quantity of phlegmatic vinous spirit; the spirit seeming to serve as a medium for joining them together. This Sapo philomedicine, prepared with oil of turpentine, was formerly celebrated as a diuretic, in nephritic complaints, and as a corrector of certain vegetables, particularly of opium: its virtues have not been fully determined by experience, nor does the present practice pay any regard to it. Beaumé observes, that it consists of only the resinous part of the oil united with the alkali; that the more fluid and well rectified the oil is, the less soap is obtained; and that by adding a little turpentine in fubstance, the preparation is considerably expedited.

Combinations of volatile alkalies with expressed oils, and with the oily balfamic juices, are obtained more readily. One ounce of spirit Liniment. of fal ammoniac, and four of oil of almonds, shaken together in a wide- volatile P.L. mouthed vial, unite perfectly, in a short time, into a white saponaceous liquid. A more confistent soapy mass is prepared, by gradually Epithema dropping the spirit into equal its weight of Venice turpentine, and volat. Ph. L. stirring them carefully together. Both these compositions are very

4 H 2

acrimonious.

acrimonious, and are used only externally, as stimulants, in rheumatic

and ischiadic pains.

Combinations of volatile alkalies with volatile oils, in a liquid form, have been already mentioned under the head of fal alkalinus volatilis: compositions of the same kind may be obtained in a solid state, by mixing the salt with the oil, and subliming them together. It may be observed, that in all these combinations made with volatile salts, though the pungency of the salt is more or less covered, it is never completely sheathed as that of the fixt alkalies is in the hard soaps; and that none of the compositions, in which either the alkali or the oil is volatile, are so perfectly saponaceous as those in which they are both of the more fixt kind.

SAPONARIA.

SAPONARIA Pharm. Edinb. (a) Saponaria major lævis C. B. Saponaria officinalis Linn. Soapwort or Brussewort: a smooth herb, with plantane-like three-ribbed leaves set in pairs on short broad pedicles; producing, on the tops of the stalks, umbel-like clusters of red, purple, or whitish flowers, cut deeply into five segments nipt at the ends, standing in long cups, followed by pear-shaped capsules full of small seeds: the root is long, slender, spreading to a great distance, so as scarce to be extirpated, of a brownish colour on the outside, internally white, with a yellowish sibre in the middle. It grows wild, but not very common, in moist grounds, and slowers in July.

The roots and leaves of saponaria discover to the taste a kind of glutinous softness or smoothness; accompanied, in the roots, with a sweetishness and slight pungency; in the leaves, with a degree of bitterness and roughness. The smoothness or soapiness, from which the plant received its name, is strongest in the leaves; which, on being agitated with water, raise a slippery froth, and are said to impart a detergent quality approaching to that of solutions of soap itself. This matter is dissolved also by rectified spirit as well as water, and hence appears evidently of a different nature from gummy or mucilaginous substances:

on inspissating the solutions, it remains entire in the extracts, and proves stronger in the spirituous extract than in the watery. This plant therefore, among us disregarded, may be presumed to have some considerable medicinal virtues: by the German physicians, the roots are used in venereal maladies, and supposed to be similar, but superiour, to those of sarsaparilla. *A physician in Paris is said lately to have given the inspissated juice of this plant to the quantity of half an ounce in a day, to persons labouring under a gonorrhæa, with success.

SAPONARIÆ NUCULÆ.

NUCULÆ saponariæ non edules C. B. Saponariæ sphærulæ arboris filicisoliæ J. B. Baccæ bermudenses Marloe. Soap-berries: a spherical fruit, about the size of a cherry; whose cortical part is yellow, glossy, and so transparent, as to shew the spherical black nut, which rattles within, and which includes a white kernel. It is the produce of a small tree, growing in Jamaica and other parts of the West Indies, called by Sir Hans Sloane prunifera racemosa, solio alato, costa media membranulis utrinque extantibus donata, fructu saponario; by Linnæus, Sapindus Saponaria.

It is faid that this fruit, at least its cortical part, has a very bitter taste, and no smell: that it raises a soapy froth with water, and has similar effects with soap in washing: that it is a medicine of singular and specific virtue in chloroses: and that a tincture or extract are preferable to the berry in substance, from whence it may be presumed that its soapy matter, like that of the saponaria, is dissoluble in spirit. Its medicinal virtue was first published by Marloe in a letter to Mr. Boyle; but the fruit having been concealed under the sictitious name of Bermudas berries, its use died with the author. That Marloe's Bermudas berries were the same with the soap-berries of America, had been suspected by some, and was confirmed by Dale in examining the Bermudas berries which Marloe had left under that title behind him. They are still, however, unknown in practice, and in the shops.

SARCOCOLLA.

SARCOCOLLA Pharm. Lond. A concrete gummy-refinous juice, brought from Persia and Arabia, in small, spongy, crumbly, whitish-yellow grains, with a few of a reddish and sometimes of a deep red colour mixed with them: the tears, when entire, are about the size of peas: the whitest tears or fragments are preferred, as being the freshest. The plant which produces this juice, and the place of its production, are unknown.

SARCOCOLLA has a bitterish subacrid taste, followed by a nauseous kind of sweetishness. It softens in the mouth, bubbles and catches slame from a candle, dissolves almost wholly in water, and greatest part of it in rectified spirit. Its medicinal qualities are not well known: it is said, when taken internally, to act as a slow and dangerous purgative; externally, to cleanse and promote the cicatrization of ulcers: dissolved in breast-milk, to be an useful collyrium for defluxions on the eyes.

SARSAPARILLA.

the root of a species of bindweed, smilax aspera peruviana sive sarsaparilla C. B. Smilax (sarsaparilla) caule aculeato angulato, soliis inermibus retuso-mucronatis Linn. growing in the Spanish West Indies, and scarcely bearing the winters of our climate without shelter. The root consists of a number of strings, of great length, about the thickness of a goose-quill or thicker, slexible, free from knots, composed of sibres running their whole length, so that they may be stript in pieces from one end to the other. They are covered with a thin, brownish, or yellowish ash-coloured skin, under which lies a thicker, white, friable substance, and in the middle runs a woody pith.

This root has a farinaceous fomewhat bitterish taste, and no smell. To water it communicates a reddish brown, to rectified spirit a yellowish red tincture, but gives no considerable taste to either menstruum. An extract, obtained by inspissating the spirituous tincture, has a weak, somewhat nauseous, balsamic bitterishness, which is followed by a slight but

but durable pungency: the watery extract is much weaker, and in larger

quantity.

Sarfaparilla was first brought into Europe by the Spaniards, about the year 1563, with the character of a specific for the cure of the lues venerea, which made its appearance a little before that time. Whatever good effects it might have produced in the warmer climates, it was found to be infufficient in this, infomuch that many have denied it to have any virtue at all, and supposed that it could do no more than, by its farinaceous foftness, to obtund the force of the gastric fluid, and thus weaken the appetite and digeftion. It appears however, from experience, that though greatly unequal to the character which it bore at first, yet, in many cases, strong decoctions of it, drank plentifully and duly continued, are of very confiderable fervice, for promoting perspiration, and what is called fweetening or purifying the blood and humours. In the medical observations published by a society of physicians in London, there are feveral instances of its efficacy in venereal maladies, as an affistant to mercury, or when mercury had preceded its use: it oftentimes answered, and that speedily, after mercurial unctions, and long continued courses of strong decoctions of guaiacum, had failed. Three ounces of the root are boiled in three quarts of river water, till the liquor when strained amounts to about one quart, which is taken at three or four doses, either warm or cold, every twenty-four hours. Dr. Harris fays, that infants who have received the infection from the nurse, though full of pustules and ulcers, and fometimes troubled with nocturnal pains, are cured by farfaparilla without mercurials: he directs the powder of the root to be mixed with their food,

SASSAFRAS.

SASSAFRAS Ph. Lond. & Edinb. The root of a large American tree of the bay kind, (laurus (sassafras) foliis integris trilobisque Linn. Arbor ex florida ficulneo folio C. B.) brought over in long straight pieces, very light and of a spongy texture, covered with a rough sungous bark, outwardly of an ash-colour, inwardly of the colour of rusty iron.

This root has a fragrant fmell, and a fweetish, subastringent, aromatic taste: the bark is much stronger than the internal woody part, and the small

Ol.effentiale rad. faffafr. Pb. L. & Ed. fmall twigs than the larger pieces. It gives out its virtues, together with a reddish colour, totally to spirit, less perfectly to water: the spirituous tincture smells weakly and tastes strongly, the watery smells stronger and tastes weaker of the root. Distilled with water, it yields a fragrant essential oil, of a penetrating pungent taste, so ponderous as to sink in water, limpid and colourless when newly distilled, by age growing yellowish and at length of a reddish brown colour: the remaining decoction, inspissated, yields a bitterish subastringent extract. Rectified spirit, distilled from the tincture made in that menstruum, brings over with it nothing considerable: the inspissated extract retains, along with the bitterness and subastringency, nearly all the aromatic matter of the root, though the smell is in great part suppressed in the extract as well as in the tincture.

Saffafras is used as a mild corroborant, diaphoretic, and sweetener, in scorbutic, venereal, cachectic, and catarrhal disorders. For these purposes, both the volatile and the fixt parts, the distilled oil and the watery extract, have been given with success: the spirituous tincture or extract, which contain both, appear to be the most elegant preparations. Insusions made in water, from the cortical or the woody part rasped or shaved, are commonly drank as tea: in some constitutions, these liquors, by their fragrance, are apt, on first taking them, to affect the head; an inconvenience, which is generally got the better of on continuing their use for a little time, and which neither the watery nor spirituous extracts are at all subject to.

SATUREIA.

SATUREIA Pharm. Edinb. (a) Satureia hortensis sive cunila sativa plinii C. B. Thymbra. Satureia hortensis Linn. Summer savoury: a low, shrubby, somewhat hairy plant: with small oblong narrow leaves, narrowest at the bottom, set in pairs; and small clusters, in the bosoms of the leaves, of pale purplish labiated flowers, whose upper lip is nipt at the extremity, the lower cut into three segments. It grows wild in some of the southern parts of Europe, and is sown annually in our culinary gardens.

The leaves of favoury are a warm aromatic; of a grateful fmell, like that of thyme but milder; and a penetrating pungent tafte. To rectified fpirit, they give out the whole of their active matter, together with a dark green tincture: water receives from them a reddish brown colour, and a considerable smell, but very little of their taste. In distillation with water, they yield a small quantity of a fragrant essential oil, very pungent, and of great subtility and volatility: the remaining decoction, inspissated, leaves a weakly bitterish, subastringent, ungrateful extract. Rectified spirit elevates in distillation much less than might be expected from the remarkable volatility of the oil: the extract smells agreeably, though weakly, of the savoury, and has a very warm, pungent, aromatic taste.

SATYRION.

SATYRION Pharm. Edinb. Orchis morio mas foliis maculatis C. B. Cynosorchis & testiculus caninus quibusdam. Orchis mascula Linn. Orchis: a plant with fix or seven long smooth narrow leaves, variegated with dark-coloured streaks or spots, issuing from the root; and one or two embracing the stalk, which is single, roundish, and striated: on its top appears a long loose spike of irregular, naked, purplish red slowers, consisting each of six petala; one of which is large, cut into three sections, hanging downwards; the others smaller, forming a kind of hood above it, with a tail behind: the root consists of two roundish whitish tubercles, about the size of nutmegs, one plump and juicy, the other sungous and somewhat shrivelled, with a few large sibres at the top. It is perennial, grows wild in shady grounds and moist meadows, and slowers in the beginning of May or sooner.

The plump roots or bulbs (the only part directed for medicinal use) have a faint somewhat unpleasant smell, and a viscid sweetish taste. They abound with a glutinous slimy juice, in virtue of which they have been found serviceable, like althea root and other mucilaginous vegetables, in a thin acrid state of the humours and erosions of the intestines. They have been celebrated also for aphrodisiac virtues, to which they appear to have little claim.

The substance brought from the eastern countries under the names of Salep, saleb, and serapias, and recommended, like our orchis root, in bilious

bilious dysenteries, defluxions on the breast, and as a restorative, appears to be no other than the prepared roots of some plants of the orchis kind. of which different species are said to be taken indiscriminately. The falep comes over in oval pieces, of a yellowish white colour, somewhat clear and pellucid, very hard and almost horny, of little or no smell, in taste like gum tragacanth. The common orchis root, boiled in water, freed from the skin, and afterwards suspended in the air to dry, gains exactly the same appearance: the roots thus prepared do not grow moift or mouldy in wet weather, which those, that have been barely dried, are very liable to: reduced into powder, they foften or diffolve as it were in boiling water into a kind of mucilage; which may be diluted, for use,

with a larger quantity of water, or with milk.

* The following process for the preparation of salep from the English orchifes, by Mr. Moult, of Rochdale, is published in the Philos. Transact. Vol. LIX. "The new root is to be washed in water, and the fine brown skin which covers it is to be separated by means of a small brush, or by dipping the root in hot water, and rubbing it with a coarse linen cloth. When a fufficient number of roots have been thus cleaned, they are to be fpread on a tin plate, and placed in an oven heated to the usual degree, where they are to remain fix or ten minutes, in which time they will have lost their milky whiteness, and acquired a transparency like horn, without any diminution of bulk. Being arrived at this state, they are to be removed, in order to dry and harden in the air, which will require feveral days to effect; or by using a very gentle heat, they may be finished in a few hours."

SAXIFRAGA.

SAXIFRAGA ALBA Pharm. Edinb. (a) Saxifraga rotundifolia Saxifraga granulata Linn. WHITE SAXIFRAGE: a plant alba C. B. with kidney-shaped crenated yellowish-green leaves, and round slender purplish branched stalks, on the tops of which grow short loose spikes of pentapetalous white flowers, followed each by a two-horned capfule full of small seeds: the root is composed of small fibres, with a number of little tubercles among them, about the fize of pepper-corns, containing,

under a chaffy covering, irregular whitish bodies somewhat brittle like the kernels of fruits. It is perennial, grows wild in sandy pasture-grounds, and slowers in May: the leaves and stalks wither soon after slowering, and by degrees the tubercles of the roots also disappear.

The leaves of this plant, of little or no smell, and of a weak unpleafant taste; and the tubercles of the roots, improperly called seeds, of no smell, and in taste sweetish with a very slight acrimony; are recommended as aperients and diuretics, in obstructions of the menses, stranguries, and nephritic cases. Among us, they have long been disused, and unknown in the shops; a more common plant, of the same name, but of a different genus, and of more activity, having generally supplied their place, viz.

SAXIFRAGA VULGARIS.

SAXIFRAGA vulgaris five anglica, Hippomarathrum anglicum, Fæniculum erraticum: Sefeli pratense, silaus forte plinio C. B. Angelica pratensis apii solio Tourn. Peucedanum Silaus Linn. English of Meadow saxifrage: an umbelliserous plant, with winged leaves subdivided into oblong narrow sharp-pointed segments: the slowers are of a yellowish white colour, the umbel naked, but its subdivisions have several little leaves at their origin; the seeds are short, brownish or reddish, plano-convex, with three deep surrows so as to appear winged: the root is long, about the thickness of the singer, brownish or blackish on the outside, and white within. It is perennial, common in meadows and pasture grounds, and slowers in June.

THE roots, leaves, and feeds of this plant have been commended as aperients, diuretics, and carminatives; and appear, from their aromatic fmell, and moderately warm pungent bitterish taste, to have a better claim to these virtues than the preceding saxifrage. They are rarely or never used.

SCAMMONIUM.

SCAMMONIUM Pharm. Lond. & Edinb. Diagrydium. SCAM-MONY: the concrete gummy-refinous juice of the roots of a species of convolvulus (convolvulus (scammonia) foliis sagittatis postice truncatis, pe-4 I 2 dunculis dunculis bifloris Linn.) distinguished by the leaves being shaped like an arrow-head and having two semicircular notches at the bottom on each side of the footstalk, the slowers being of a pale yellowish colour and standing two on one stem: it is a native of Syria, and has been lately found to bear the colds of our own climate. The scammony is extracted in Syria, by baring the upper part of the root in June, cutting off the top obliquely, and placing a shell or some other receptacle at the depending part to receive the milky juice, which on standing concretes into solid masses.

The best scammony is brought from Aleppo, in light spongy masses, easily friable, glossy, of different shades of colour from a grey or yellowish white almost to black, when reduced to powder of a brownish white colour. An inferiour sort comes from Smyrna, in compact hard ponderous pieces, full of sand and other impurities. Such should be chosen as crumbles the most easily betwixt the singers, grows instantly white on the contact of watery moisture, and leaves little or no seces on being dissolved. Its colour in the mass affords no criterion of its purity or goodness.

Scammony has a flight unpleafant smell, and a weak bitterish sub-acrid taste. It consists of about equal parts of resinous and gummy matter, and hence dissolves almost totally in a mixture of equal parts of rectified spirit and water, that is, in proof spirit. Rectified spirit takes up the resin, with some part of the gum: if the tincture be inspissated a little, and then mixed with water, the gum continues dissolved, and the pure resin precipitates. By trituration with water, or by bare maceration, the scammony is resolved into a milky liquor verging to greenish; which on standing deposites some portion of the resin, but retains its milkiness.

This gummy-refin is one of the strong stimulating cathartics; more kindly in operation, and hence in more general use, than most of the other substances of that class: the dose is from two or three grains to twelve. Sundry ill qualities have been ascribed to it, which it is not found to posses: and sundry correctors have been devised, which it does not appear to want. In cold indolent serous habits, scammony itself procures generally a plentiful evacuation with great ease and safety: in inflammatory cases, and the more irritable dispositions, it is indeed dangerous; but no otherwise so than the rest of the strong purgatives; and no otherwise than by virtue of that power on which its efficacy in the opposite circumstances depends.

By

By the smallness of the dose of this medicine, its easy solubility, and its having little tafte, it is fitted for being commodiously taken in almost any form. It is made in the shops into a powder, with the addition of three fourths its weight of calcined hartshorn to divide and disunite it +; Pulvis e or with equal its weight of crystals of tartar 1. A scammoniate electuary is composed of one ounce of scammony and sour of honey, aromatised Lond. ‡Ph. with half an ounce of cloves, half an ounce of ginger, and a scruple of the effential oil of caraway-feeds; of which composition, one dram and a half contain fifteen grains of the scammony +. Agreeable purging troches, for those who are not easily prevailed upon to take medicines of this kind in other forms, are prepared, by grinding together three drams of scammony, four drams of crystals of tartar, four drops of oil of cinnamon, and eight ounces of fine fugar, and moistening the mixture with fo much rose-water as will render it of a due confistence for being formed: each tablet is made to weigh about a dram t, and confequently contains two grains and a half of scammony. One of the most elegant liquid preparations is a folution of the scammony in a strong infusion or decoction of liquorice, poured off from the feces, and aromatifed with fome grateful distilled water or aromatic tincture; as those of cardamomfeeds.

fcammonio

+ Electar. e fcammonio Pharm. Lond.

t Morfuli purgantes Ph. Brandenburg.

The dried root of the plant, as well as its juice, may perhaps deserve fome notice. Dr. Ruffel, to whom the public is obliged for an accurate history of this drug, relates that a decoction of half an ounce of the root procured five stools, without gripes, fickness, or any manner of uneafiness, and, on repeating the trial several times, had the same effect: and that the decoctions are entirely without smell, and in taste rather fweetish than disagreeable. Neither the stalks, leaves, flowers, or seeds, feemed to have any purgative virtue (a).

SCILLA.

SCILLA Pharm. Lond. & Edinb. Scilla radice alba, & scilla vulgaris radice rubra C. B. Ornithogalum maritimum Tourn. Scilla maritima SQUILL or SEA-ONION: a plant with a large bulbous onion-like root; from which rife, first a naked stalk bearing several hexapetalous

white flowers, and afterwards large green lily-like leaves with a remarkable rib in the middle of each. It grows fpontaneously on fandy shores in Spain and in the Levant, from whence we are annually supplied with the roots. They should be chosen large, plump, fresh, and full of a clammy juice: some are of a reddish colour, and others white, but no difference is observed in the qualities of the two forts, and hence the college allows both to be taken promiscuously.

This root is to the tafte very nauseous, intensely bitter, and acrimonious: much handled, it exulcerates the skin. Taken internally, it acts as a powerful attenuant and aperient: in doses of a few grains it promotes expectoration and urine: in fomewhat larger ones, it proves emetic and fometimes purgative. It is one of the most certain diuretics in hydropic cases, and expectorants in asthmatic ones, where the lungs or stomach are oppressed by tenacious phlegm, or injured by the imprudent use of opiates.

Phar. Edinb.

This medicine, on account of its ungrateful tafte, is most commodiously taken in the form of pills; into which the dried root may be reduced, Pil.feillitica by beating it with thrice its weight each of ammoniacum and leffer cardamom-feeds in powder, and extract of liquorice, and a sufficient quantity of fimple fyrup. In whatever form fquills are given, unless when defigned to act as an emetic, the addition of some grateful aromatic material is of use, to prevent the nausea which of themselves they are very apt, even in fmall doses, to occasion.

The fresh root loses in drying about four fifths of its weight, without any confiderable loss of its tafte or virtue; the vapour which exhales appearing to be little other than merely aqueous. Hence four grains, which are the mean dose of the dry root in powder, are equivalent to Scilla exfic- near a scruple of the fresh squill. The most convenient way of drying cata Pharm.

Lond. & Ed. it is, after peeling off the outer skin, to cut the roots transversely into thin flices (not to separate the coats, as has been usually directed) and expose them to a gentle warmth.

The ancients, in order to abate the acrimony of the squill for certain purposes, inclosed it (after separating the skin, and the fibres at the bottom with the hard part from which they iffue) in a paste made of flour and water, and then baked it in an oven, till the paste became dry, and the fquill foft and tender throughout. The fquill, fo prepared, was

Scilla cocta Pharm. Lond.

beaten

beaten with two thirds its weight of flour, the mixture formed into Troch.esciltroches, and dried with a gentle heat. These troches were supposed to be alexipharmac, and in this light were made an ingredient in theriaca, for which purpose they are still retained.

Water, wine, proof spirit, and rectified spirit, extract the virtues both of the fresh and the dry root. Nothing rifes in distillation with any of these menstrua, the entire bitterness and pungency of the squill remaining concentrated in the inspissated extracts: the spirituous extract is in fmaller quantity than the watery, and of a proportionably stronger almost fiery tafte.

Alkalies confiderably abate both the bitterness and acrimony of the fquill: vegetable acids make little alteration in either, though the admixture of the acid taste renders that of the squill more supportable. These acids extract its virtue equally with watery or spirituous menstrua; and, as an expectorant in diforders of the breaft, excellently coincide with it. The college of London directs an acetous tincture to be Acetum scilprepared, by macerating a pound of the dry roots in fix pints of vinegar, liticum Ph. with a gentle heat: to the liquor pressed out, and after settling poured off from the feces, one twelfth its quantity of proof spirit is added, to prevent its growing foon foul. The college of Edinburgh for the same preparation directs the proportions of two ounces of dried fquills, two pounds and a half of distilled vinegar, and three ounces of rectified spirit. A scillitic oxymel is obtained by boiling a quart of the acetous tincture Oxym. scilwith three pounds of clarified honey, till the mixture acquires the confiftence of a fyrup: and a fyrup of fquills, by diffolving three pounds Syr. feillit. and a half of fine fugar in two pounds of the vinegar. These preparations are used, as expectorants, in doses of one, two, or three drams, along with cinnamon or some other grateful water: where the first passages are overloaded with viscid phlegm, an ounce or more is given at once, to procure a more speedy and effectual evacuation by vomit.

lit. Ph. Lon.

Phar. Edin.

SCINCUS.

SCINCUS Pharm. Lond. Scincus seu crocodilus terrestris Raii. THE SKINK: a fmall amphibious animal, of the lizard kind, clothed with greyish scales, caught about the Nile, &c. and thence brought, dried, to us, remarkably smooth and glossy as if varnished. The flesh of this animal, particularly of the belly, has been faid to be diuretic, alexipharmac, aphrodifiac, and useful in leprous disorders. Whatever virtues it may have when used fresh, as food, it is not expected to be of any importance as it comes to us, and serves only to increase the number of the articles of which mithridate is composed.

SCORDIUM.

SCORDIUM Pharm. Lond. & Edinb. C. B. Chamædrys palustris & trissago palustris quibusdam Teucrium Scordium Linn. WATER-GERMANDER: a trailing plant, with oblong, oval, indented, soft, hoary leaves, set in pairs, without pedicles: in their bosoms issue purplish monopetalous flowers, not above four or five together, each cut into five segments and followed by four small seeds lodged in the cup. It is sometimes found wild in watery places, but the shops are supplied chiefly from gardens: it is perennial, and flowers in June.

The leaves of scordium, rubbed betwixt the fingers, yield a moderately strong smell, somewhat of the garlick kind: to the taste they discover a considerable bitterness and some pungency; but the astringent power, which some ascribe to them, could not be distinguished, either by the taste, or by solution of chalybeate vitriol. They are recommended as alexipharmacs and corroborants, in malignant and putrid disorders, and in laxities of the intestines: they enter several officinal compositions in those intentions, and are sometimes employed externally in antiseptic cataplasms and somentations.

On keeping the dry herb for some months, its smell is dissipated; and the bitterness, thus divested of the slavouring matter, proves considerably less ungrateful than at first. The leaves, moderately and newly dried, give out their smell and taste both to water and to rectified spirit; and tinge the former of a brownish, the latter of a deep green colour. In distillation, their peculiar slavour arises with water; but the impregnation of the distilled sluid is not strong, nor could any essential oil be obtained on submitting to the operation several pounds of the herb: the remaining decoction, inspissated, leaves a very bitter mucilaginous extract. Rectified spirit brings over little or nothing: the inspissated extract partakes in a considerable degree of the slavour of the scordium, and proves in bitterness also far stronger than the watery.

SCOR-

SCORZONERA.

SCORZONERA Pharm. Edinb. (a) Scorzonera latifolia sinuata C. B. Viperaria & serpentaria hispanica quibusdam. Scorzonera hispanica Linn. Vipers-grass: a plant with large sharp-pointed leaves, slightly sinuated about the edges, having a large prominent rib in the middle, joined to the stalks without pedicles: on the tops of the branches grow yellow sloculous flowers, set in scaly cups, followed by oblong roundish striated seeds winged with down: the root is long, single, from the size of a goose-quill to that of the little singer, of a dark colour on the outside and white within. It is perennial, a native of Spain, and common in our culinary gardens.

The roots of fcorzonera have been employed medicinally as alexipharmacs, and in hypochondriacal diforders and obstructions of the viscera; but at present are more properly looked upon as alimentary articles, in general salubrious, and moderately nutritious. They abound with a milky juice, of a soft sweetish taste, but which in drying contracts a slight bitterness. Extracts made from them by water are considerably sweet and mucilaginous: extracts made by rectified spirit have a less degree of sweetishness, accompanied with a slight grateful warmth. In Cartheuser's experiments, the spirituous extract amounted to one third the weight of the root, and the watery to above one half: as his watery extract, though in larger quantity than the spirituous, was nevertheless, like mine, sweeter, it should seem that the sweet matter of scorzonera is somewhat different, in regard to its solubility, from that of most of the other vegetable sweets that have been examined, the spirituous extracts having generally much the greatest sweetness.

SCROPHULARIA.

FIGWORT: a plant with square stalks; the leaves set in pairs, at distances, in opposite directions; the branches terminated by loose spikes of irregular, purple, helmet-shaped slowers; each of which is followed

by a roundish pointed capsule, containing numerous small seeds in two cells. It is perennial.

1. SCROPHULARIA VULGARIS Pharm. Edinb. (a) Scropbularia nodosa fætida C. B. Millemorbia quibusdam. Scropbularia nodosa Linn. Common figwort or kernelwort: with the leaves somewhat heart-shaped and serrated about the edges; the roots long, thick, and full of knots and tubercles. It grows wild in woods and hedges, and flowers in July.

The roots and leaves of this plant have been celebrated both internally and externally, against inflammations, the piles, scrophulous tumours, and old ulcers. Their sensible qualities are, a rank smell somewhat like that of elder leaves but stronger, and a disagreeable bitterish taste. The anodyne and anti-inflammatory virtues, which they are reckoned to exert in external applications, are attributed in great part to the odorous matter, which is supposed to be somewhat of the narcotic kind: the root, which has less of this smell than the leaves, has been generally preferred for internal use. At present, they are both among us disregarded.

2. Scrophularia aquatica major G. B. Scrophularia aquatica Pharm. Edinb. (a) Scrophularia aquatica major C. B. Scrophularia aquatica Linn. Greater water figwort, water betony: with the leaves oblong, nearly oval, crenated about the edges; the stalks winged at the angles; the root composed of numerous white strings issuing from one head. It grows in watery places, and slowers in July.

The leaves of this species are recommended for the same purposes as those of the preceding, to which they have by some been preferred: in taste and smell, they are similar, but weaker. Mr. Marchant reports, in the memoires of the French academy, that this plant is the same with the iquetaia of the Brazilians, celebrated as a specific corrector of the ill slavour of sena: on his authority, the Edinburgh college, in their common insusion of that drug, directed two thirds its weight of the water signort leaves to be joined; but as they have now discarded this ingredient, we may presume that it was not found to be of much use.

S E D U M.

SEDUM MAJUS seu Sempervivum majus Pharm. Edinb. (a) Sedum majus vulgare C. B. Aizoon & barba jovis quibusdam. Sempervivum Houseleek or Sengreen: a plant with numerous, tectorum Linn. thick, stiff, sleshy, pointed leaves, lying over one another in form of a roundish cluster; in the middle of which rises a stiff stalk, covered with fmaller leaves, divided at the top into feveral branches, bearing purplish flowers with twelve petala, which are followed by the fame number of capfules full of small feeds. It is perennial and evergreen, grows on old walls and the tops of houses, and flowers in June.

THE leaves of houseleek, of no remarkable smell, discover to the taste a mild fubacid aufterity: their expressed juice, of a pale yellowish hue when filtered, yields on infpiffation a deep yellow, tenacious, mucilaginous mass, considerably acidulous and acerb: from whence it may be prefumed, that this herb has fome claim to the refrigerant and restringent virtues that have been ascribed to it. It is observable that the filtered juice, on the addition of an equal quantity of rectified spirit of wine, forms a light white coagulum, like creme or fine pomatum, of a weak but penetrating taste: this, freed from the fluid part, and exposed to the air, almost totally exhales. From this experiment it is concluded by fome that houseleek contains a volatile alkaline falt (b): but the juice coagulates in the fame manner with volatile alkalies themselves, as also with fixt alkalies: acids produce no coagulation.

SELENITES.

SELENITES: an earthy or stony concrete; not disfoluble in acids; calcining in a gentle heat into a foft powder+, which forms a + Plaster of tenacious paste with water: composed of calcareous earth and vitriolic acid.

THE vitriolic acid, poured on crude calcareous earths, as chalk, limestone, marble, does not dissolve or unite with them, at least in any

(b) Burghart, Medicorum Silefiacorum fatyræ, specim. IV. obf. ii. p. 11. (a) Expunged.

confiderable degree: but if the earth be previously disfolved in any other acid, the vitriolic acid, superadded to this solution, absorbs the dissolved earth, and forms with it a concrete no longer foluble, which of course renders the liquor milky, and on standing settles to the bottom, either in a powdery or crystalline form, according as the liquor was less or more diluted with water. Native mineral concretes of this kind, when pellucid and crystalline, are called felenites; when composed of a number of thin transparent coats or leaves, lapis specularis, Muscovy glass, or isinglass; when in large stony masses, of a granulated texture, gypsum; and when the masses are of a fibrous texture, striated gypsum or English talc. All these substances are made to discover their composition, by strongly calcining them in contact with the burning fuel: the inflammable principle of the coals absorbs their vitriolic acid, from which combination is produced common fulphur, greatest part of which exhales; and the remaining calcined earth, thus deprived of the acid, is found to be a perfect quicklime.

This concrete, in its different forms, has been recommended as an aftringent in fluxes and hemorrhagies; a virtue which agrees but ill with its indiffolubility and want of taste. It is often met with in the residua of waters, both of the common and medicinal springs.

SENA.

SENA Ph. Lond. Senna Ph. Edinb. Folium orientale. Sena: the leaf of an annual, woody, pod-bearing plant (fenna alexandrina five foliis acutis C. B. Cassia (senna) foliolis trijugatis quadrijugatisque Linn.) brought dry from Alexandria in Egypt. It is of a lively yellowish green colour, an oblong somewhat oval figure, sharp-pointed at the ends, about a quarter of an inch broad, and not a full inch in length. Some inferiour sorts are brought from Tripoli and other places: these may be distinguished, by their being either narrower, longer, and sharper pointed; or larger, broader, and round pointed, with small prominent veins; or large, obtuse, and of a fresh green colour without any yellow cast.

SENA is a moderately strong, and in general a safe cathartic: Geoffroy specifies hemorrhagies, inflammations of all kinds, and disorders of the breast,

breast, as being almost the only exceptions to its use. The dose in substance is from a scruple to a dram; in infusion, from one dram to three or four. It gives out its virtue both to watery and spirituous menstrua: to water and proof spirit it communicates a brownish colour, more or less deep according to the proportions; to rectified spirit, a fine green. There are two inconveniences often complained of in this medicine, its being liable, in most constitutions, to occasion gripes; and its being accompanied with an ill flavour, which is apt to nauseate both the stomach and the palate. The first may be greatly obviated by dilution, the latter by aromatic and other additions; feveral compositions of this kind are prepared in the shops, both sufficiently palatable, and which operate for the most part with ease and mildness.

Six drams of tamarinds, one of crystals of tartar, half a dram of cori- Infusum taander feeds, half an ounce of brown fugar, and one, two or three drams cum fenna of fena are infused in eight ounces of boiling water, in an unglazed Phar. Edinb. earthen vessel, for four hours, and then strained. Or three drams of Infus. senze fena are infused in a quarter of a pint of boiling water, till the liquor has commune Pharm. Lond. grown cold, with half a dram of leffer cardamom feeds hufked, and three fourths of a dram of crystals of tartar, which last are previously boiled in the water till dissolved; or with two drams of fresh lemon peel, and two Infus. sense drams by measure of lemon juice. The committee of the London limoniatum college observe, that this last is the most agreeable form, they have been able to contrive, for the exhibition of fena to those who are more than ordinarily offended with its flavour; and that though acids are generally supposed to impede the action of water on vegetables, yet the infusions of fena made with acids are found, from experience, not to fail in their intention. Indeed if the acids really weaken the diffolving power of the . water, which it is probable they do in some degree, it should seem to be, on this account, rather of advantage than otherwise; for, as the committee further observe, in a medicine very nauseous to many, it is of primary consequence that only the lighter and least disgustful parts be extracted. On this principle, some macerate the sena for a night in cold water, which becomes fufficiently impregnated with its purgative virtue, without extracting fo much, as boiling water does, of the naufeous matter: if the liquor, poured off from the sena, be boiled a little by itself, great part of its ill flavour will be distipated; and the remains of its offensiveness may be covered by infusing in it some bohea tea. coction

marindorum

coction is continued for any confiderable time, the purgative virtue of the fena will be diminished; for the inspissated watery extracts are scarcely found to purge so much, as one fourth of the insusion or decoction they were made from, or so much as an equal weight of the leaves in substance.

† Tinctura fenæ Pb. Lon. † Tinct. fennæ comp. vulgo elixir falutis Pb. E.

The officinal spirituous tinctures of sena are prepared by digestion for fome days, with + or without t heat, in proof spirit. The proportions, in the London pharmacopæia, are three ounces of sena to a quart of the spirit, to which are added four ounces of stoned raisins, three drams of caraway feeds, and one dram of leffer cardamom feeds hufked: in the Edinburgh, two ounces of fena to three pounds and a half of the spirit, with the addition of one ounce of jalap, half an ounce of coriander feeds, and four ounces of white fugar-candy in powder, which last is directed to be dissolved in the tincture after straining it from the other ingredients. Both these tinctures are agreeable and useful carminative purgatives, especially to those who have accustomed themselves to spirituous liquors: the ill flavour of the sena is in great measure covered, and its offending the stomach or producing gripes prevented, by the warm feeds and the fweets. Several compositions of this kind have been offered to the public, under different names: the two above are inferiour to none; and superiour to most of them.

SERPENTARIA.

SERPENTARIA VIRGINIANA Pharm. Lond. & Edinb. Serpentaria virginiana & Viperina & Colubrina virginiana Pharm. Parif. VIRGINIAN SNAKEROOT: the root of a species of aristolochia growing in Virginia and Carolina, aristolochia (serpentaria) foliis cordatis oblongis planis, caulibus insirmis flexuosis teretibus, sloribus solitariis Linn. The root is small, light, bushy, composed of a number of strings or sibres issuing from one head and matted together, of a brownish colour on the outside, and paler or yellowish within.

SNAKEROOT has an aromatic smell, approaching to that of valerian, but more agreeable, and a warm bitterish pungent taste, which is not easily concealed or overpowered by a large admixture of other materials. It gives out its active matter both to water and rectified spirit, and tinges

the former of a deep brown, the latter of an orange colour. Greatest part of its smell and flavour is carried off in evaporation or distillation by both menstrua: along with water there arises, if the quantity of the root submitted to the operation be large, a small portion of a pale-coloured essential oil, of a considerable smell, but no very strong taste, greatest part of the camphorated pungency, as well as bitterishness of the root, remaining in the inspissated extract. The spirituous extract is stronger than the watery; not so much from its having lost less in the evaporation, as from its containing the active parts of the root concentrated into a fmaller volume; its quantity amounting only to about one half of that of the other.

This root is a warm diaphoretic and diuretic. It is reckoned one of the principal medicines of the alexipharmac kind; and as fuch is in general use, in low malignant fevers and epidemic diseases, for raising the pulse, promoting a diaphoresis, and correcting a putrid disposition of the humours. It is given, in fubstance, from a few grains to a scruple or half a dram; in decoction or infusion, to a dram and upwards. Tinctures of it are prepared in the shops, by macerating three ounces Tinctura of the root in a quart of proof spirit+, or two ounces, with one dram of cochineal, in two pounds and a half of the fame spirit ‡.

ferpentarize + Ph. Lond. 1 Pb. Edinb.

SESELI.

SESELI Pharm. Lond. Ligusticum quod seseli officinarum C. B. Laferpitium Siler Linn. HARTWORT OF SERMOUNTAIN: a tall umbelliferous plant, with large leaves, composed of oblong pointed sections set in pairs or three together: the entire umbel, and its fubdivisions, have a circle of little leaves at their origin: the feeds are large, of a pale brown colour, oblong, flat on one fide, convex and striated on the other, and edged with a leafy margin: the root is large, thick, and branched. It is perennial, grows wild in some of the southern parts of Europe, is raifed with us in gardens, and flowers in June.

BOTH the feeds of this plant, which are the part directed in our pharmacopæias, and the roots, appear to be useful aromatics, though not regarded in practice; of an agreeable smell, and a warm glowing sweetish tafte. The roots have the greatest warmth and pungency: the feeds, the greatest greatest sweetness and the most pleasant flavour. A spirituous extract of the seeds is a very elegant aromatic sweet.

SESELI MASSILIENSE.

SESELI MASSILIENSE Pharm. Edinb. (a) Sefeli massiliense fæniculi solio C. B. Fæniculum tortuosum J. B. Seseli tortuosum Linn. HARTWORT OF MARSEILLES: a large spreading branched umbelliserous plant; with the stalk and branches sirm, woody, knotty, and variously bent; the leaves sinely divided, like those of sennel, but somewhat thicker, shorter, stiffer, and more distant from one another; the seeds also in shape like those of sennel, and of a pale grey colour. It is perennial, and a native of the southern parts of Europe, from whence the seeds are sometimes brought to us.

THE feeds of this plant have an agreeable aromatic smell, and a very warm biting taste: they are more pungent than those of the foregoing sessel, but want their sweetness.

SIGILLUM SALOMONIS.

GONVALLARIA feu Sigillum Salomonis Pharm. Edinb. (b) Polygonatum latifolium vulgare C.B. Convallaria multiflora Linn. Solomonsseal: a plant with unbranched stalks, bearing oval narrow leaves ribbed like those of plantane, generally all on one side: on the other side hang oblong monopetalous white flowers, two or more together, on long pedicles, followed each by a black berry: the root is white, thick, sleshy, with several joints, and some flat circular depressions supposed to resemble the stamp of a seal. It is perennial, grows wild in woods, and slowers in May.

THE roots of Solomons-seal are recommended externally as restringents; and internally as incrassants and mild corroborants. They have little or no smell; to the taste they discover a considerable sweetness and viscidity, followed by a very slight impression of bitterishness and acri-

⁽a) Expunged.

^{*(}b) The Edinburgh college give the Convallaria Polygonatum of Linnaus as their species.

mony, which is diffipated by boiling. It is faid that they have been used with success in the hamorrhoids (a). The flowers, berries, and leaves, are acrid and poisonous (b).

SIMAROUBA.

SIMAROUBA Pharm. Edinb. the bark of the Quassia Simarouba, Linn. suppl. Quassia dioica, Pharm. Suec. Evonymus fructu nigro tetragono Barrer. Equin. It is brought from Guiana, in long pieces, of a yellowish white colour, light, tough, and of a sibrous texture.

Mr. de Justieu reports, that this bark is of common use in Guiana, against dysenteric fluxes, and was brought from thence into Europe in the year 1713: that the fluxes which, in France, fucceeded the exceffively hot fummer of 1718, and which not only refifted, but were aggravated by, purgatives, aftringents, and ipecacoanha, happily yielded to fimaruba: that decoctions of an ounce or half an ounce in a small quantity of water, the dose used by the natives of Guiana, occasioned often vomiting, almost always uneasy sweats, and sometimes an increase of the bloody and ferous discharges by stool; but that a decoction of two drams in a quart of water, boiled to the confumption of one third, divided into four doses, and taken warm at intervals of three hours, abated the pain in one day, and when continued for a short time completed the cure, without producing any nausea or disturbance: that it is not accompanied with the ill effects of aftringents: that it abates spasmodic and hysteric symptoms: that it answers best in fluxes of the seroso-bilious, bloody and mucous kind, supported by a convulsive motion of the intestines, where there is no fever, where the functions of the stomach are unhurt, and in tenesmi (c). Dr. Degner likewise made use of this bark in the above form, with good fuccess, after proper evacuations, in an epidemic putrid dysentery, which raged at Nimeguen during the summer and autumn of 1736: he fays it acted mildly and almost infensibly, and that its effects were speedier in bloody than in bilious discharges: he takes notice also that the barks procured under the name of simaruba, in different parts of Holland, from Leipfick, and from Paris, differed

⁽a) Cullen, Mat. Med. (b) Id.

⁽c) Mem. de l'acad. des scienc. de Paris, 1729. Geoffroy, mat. med. ii. 211.

MATERIA MEDICA.

greatly in quality from one another; but does not mention what the differences were, nor the qualities of the genuine or best fort (a).

The simaruba, which I have met with in our shops, has a moderately strong, durable, not very ungrateful bitter taste, without smell, and without any manifest astringency. Macerated in water, or in rectified spirit, it quickly impregnates both menstrua with its bitterness, and with a yellow tincture. It seems to give out its virtue more perfectly to cold than to boiling water; the cold infusion being rather stronger in taste than the decoction; which last, of a transparent yellow colour whilst hot, grows turbid and reddish brown as it cools. The milky appearance, which Justieu says it communicates to boiling water, I have not observed in the decoction of any of the specimens I examined.

SINAPI.

sinapi Pharm. Lond. Sinapi album Pharm. Edinb. Sinapi rapi folio C. B. Sinapis nigra Linn.*(b) Mustard: an annual plant; with long rough leaves divided to the rib into irregular fegments, of which the extreme one is largeft; producing, at the tops of the branches, tetrapetalous yellow flowers, followed, each, by a short, smooth, quadrangular pod, divided longitudinally by a membrane which projects at the ends, containing small, roundish, reddish-brown or dark-coloured feeds. It is a native of England, but commonly cultivated for medicinal and dietetic use.

Mustard feed is one of the strongest of the pungent, stimulating, diuretic medicines that operate without exciting much heat. It is sometimes taken, unbruised, to the quantity of a spoonful at a time; in paralytic, cachectic, and serous disorders. In this manner of exhibition it generally opens the body; whereas the powder is apt to occasion vomiting, in which intention it is sometimes given diffused in warm water, of which repeated draughts must be drunk, to continue the effect. It is applied also, as an external stimulant, to benumbed or paralytic limbs;

⁽a) Hist. dysenteriæ bilioso-contagiosæ, in Append. ad act. nat. curios. vol. v.

^{*(}b) The Sinapis alba Linn. is the Edinburgh species, which differs little from the black, or common, except in being less pungent and bitter. It should seem therefore to be less proper for external use, at least.

to parts affected with fixt rheumatic pains; and to the foles of the feet, in the low stage of acute diseases, for raising the pulse: in this intention, a mixture of equal parts of the powdered feeds and crumb of bread, with the addition, fometimes, of a little bruifed garlick, are made into a cataplasm with a sufficient quantity of vinegar.

Mustard seed yields upon expression a considerable quantity of oil, oil. semin. which is by fome recommended externally against rheumatisms and palfies, though it has nothing of that quality by which the feeds themfelves prove useful in those disorders; the oil being mild and insipid as that of olives, and the pungency of the feed remaining entire in the cake left after the expression. Nor is any considerable part of the pungent matter extracted by rectified spirit; the tincture, which is of a pale amber colour, having very little tafte; and the extract, obtained by inspiffating it, being only bitterish and oily: the quantity of extract is about one fixteenth the weight of the feeds. The bruifed feeds give out readily to water nearly the whole of their active matter: added to boiling milk, they curdle it, and communicate their pungency to the whey. Distilled with water, they yield a limpid effential oil, extremely pungent and penetrating both in fmell and tafte, and fo ponderous as to fink in the aqueous fluid: the remaining decoction, thus divested of the principle in which alone the acrimony of the mustard resides, leaves on being inspiffated a fweetish mucilaginous extract.

SOLANUM.

NIGHTSHADE: a plant with a monopetalous flower, divided into five fegments, having its cup divided in the same manner, with the fame number of stamina in the middle, and followed by a juicy berry.

1. Belladona Pharm. Edinb. & Parif. Solanum melanocerafos C. B. Solanum lethale. Atropa Belladonna Linn. Deadly nightshade or dwale: with the leaves oval, pointed, somewhat hairy; the flowers folitary in the bosoms of the leaves, of a dull purplish colour, tubulous, flightly cut, with the stamina separate from one another; the berries of a gloffy black. It is perennial, grows wild in some shady waste grounds, and flowers in July.

2. Solanum Pharm. Parif. Solanum officinarum C. B. Solanum nigrum Linn. Garden nightshade: with the leaves oval, pointed, having generally some irregular indentations; the flowers in clusters, white, not tubulous, deeply cut, the segments spread out, and the tips of the stamina united into one button; the berries black. It is annual, grows spontaneously in cultivated grounds, and slowers in August.

The leaves of these plants have a faint smell, somewhat of the narcotic kind, which in drying is diffipated: on the organs of tafte, whether fresh or dry, they make scarcely any impression. Their effects are nevertheless very powerful: in external applications, they are faid to act as refrigerants, refolvents, and discutients: taken internally, in the quantity of not many grains, they are highly deleterious, the first somewhat the most so. In very small doses, as an infusion in boiling water of half a grain or a grain of the dried leaves, they occasion a warmth over the whole body; which is often followed by a fweat, or an increase of the urinary discharge, or some loose stools, or a sickness and vomiting; and often by a headach, giddiness, dimness of the fight, and other paralytic symptoms (a). In some cancerous, ulcerous, and hydropic cases, these infufions have been repeated, at bed-time, every two or three nights or oftener, and the quantity of the leaves in each dose increased gradually to five or fix grains or more, with apparent benefit: but they are fo variable and irregular in their operation, and so liable, not only to fail of giving relief, but to be productive of very alarming fymptoms by ftrongly affecting the nervous fystem, that their use is deservedly laid aside. Their good effects, when they happen to prove medicinal, feem to depend, not on any alterative or peculiarly deobstruent power, but merely on the evacuations they produce: where they do not act as evacuants, they generally aggravate the complaints (b).

* In the Med. Comment. Vol. I. p. 419. is a remarkable case of the efficacy of an external application of belladonna in discussing a scirrhous

⁽a) Mr. Ray gives an account, from his own knowledge, of a pretty remarkable effect of a fmall piece (particula) of a fresh leaf of belladonna applied externally to a little ulcer, supposed cancerous, below the eye: the uvea became in one night so relaxed, that it lost all power of contracting the pupil, which, though exposed to the strongest light, continued dilated to four times its natural size, till the leaf being removed the parts gradually recovered their tone. The application was repeated three several times, and produced always the same effect. Hist. plant. 680.

⁽b) See Mr. Gataker's observations (and the supplement thereto) on the internal use of the nightshade, and Mr. Bromfeild's account of the English nightshades and their effects.

Maria.

tumour in the rectum, near the anus, which almost totally blocked up the passage. The mode of application was a poultice of the root boiled in milk. This was applied to the anus and perinæum, and renewed morning and evening. In the space of a month it entirely dissolved the tumour, without any suppuration, or discharge of matter. The writer fays that he could add more inflances of the good effects of this plant externally applied.

The roots and berries appear to partake of the deleterious qualities of the leaves, though probably in different degrees: the berries in particular feem to be of much less activity. It is faid that three or four of the berries of the deadly nightshade, which are reckoned more virulent than those of the other fort, have been sometimes eaten without injury: Gesner reports that their expressed juice, boiled with a little sugar to the confiftence of a fyrup, proves, in doses of a tea-spoonful, an effectual and safe anodyne, but gives a particular caution not to exceed this dose. The Succus spissed Edinburgh college has directed the inspissated juice of these berries to be fatus belladonna P. E. kept as an officinal.

3. DULCAMARA Pharm. Edinb. Solanum scandens seu dulcamara C. B. Amaradulcis & glycypicros quibusdam. Solanum Dulcamara Linn. Woody nightshade or Bittersweet: with several of the leaves, particularly the upper ones, cut deeply into three fections, or rather furnished with two fmaller appendages at the bottom; the flowers in clusters, of a blue colour, with the fegments spread out and the stamina united as in the fecond species; the berries red. It grows by the sides of ditches and in moist hedges, climbing upon the bushes, with winding, woody, but brittle stalks. It is perennial, and flowers in June and July.

The roots and stalks of this species impress, on first chewing them, a confiderable bitterness, which is foon followed by an almost honey-like fweetness. They have been commended in different disorders, as high refolvents and deobstruents: their sensible operation is by sweat, urine, and stool; the dose from four to fix ounces of a tincture made by digesting four ounces of the twigs in a quart of white wine. Experience has shewn, that they are by no means equally deleterious with the two preceding nightshades; that they act more regularly and uniformly; and that, without producing nervous complaints, they produce more con-

fiderable

fiderable evacuations, especially by stool; but their virtues in particular

cases have not yet been sufficiently ascertained.

* In a medical differtation on this plant, printed at Upfal, a light decoction and infusion of the stalks is the preparation recommended, and is said to have been frequently employed with success in violent ischiadic and rheumatic pains. The efficacy of the dulcamara in the jaundice, source, suppressed menses, and the lues venerea, is also mentioned from other authors.

SPERMA CETI.

SPERMA CETI Pharm. Lond. & Edinb. Spermaceti, improperly so called: a species of fat; found in certain whales, particularly in their heads; artificially purified, by boiling with alkaline lye, to a snowy whiteness; and afterwards broken into slakes. It differs from the other animal fats, in not being dissoluble by alkalies or combinable with them into soap; and in rising almost totally in distillation, not in form of a fluid oil, but in that of a butyraceous matter resembling, both in consistence and smell, the butter of wax. In long keeping, it is apt to turn yellow and rancid: the matter, very small in quantity, which has suffered this change, and which taints the rest, is found to have lost the discriminating characters of the spermaceti; being dissoluble both by alkaline lye and by vinous spirits, so as to leave the remainder white and sweet as at first.

This concrete, of a foft butyraceous taste and no remarkable smell, is given with advantage in tickling coughs, in dysenteric pains and erosions of the intestines, and in such cases in general as require the solids to be softened and relaxed, or acrimonious humours to be obtunded. It readily dissolves in oils, and unites by liquesaction with wax and resins; and in these forms is applied externally. For internal use, it may be dissolved in aqueous liquors into the form of an emulsion, by trituration with almonds, the yolk or white of an egg, and more elegantly by mucilages; or made into a lohoch with proper additions.

* SPIGELIA.

SPIGELIA Pharm. Edinb. Anthelmia Dris. Lining. Spigelia marilandica Linn. Periclymeni virginiani flore coccineo planta marilandica, spica erecta, foliis conjugatis Catesby carol. Indian Pink: this plant has a perennial fibrous root, whence rise single stems, beset with opposite oval-lanceolate, entire leaves, and crowned with a spike of tubular monopetalous red flowers, with five stamina and one pistil. Each flower is succeeded by two round united bivalvular capsules, containing several small seeds. It grows spontaneously in South Carolina, and other southern provinces of North America.

The use of the root of this plant as an anthelmintic was communicated from the native Indians to the colonists, and it has since been much employed in that country. The first account of its virtues is to be met with in a paper of Dr. Lining's, in Vol. I. of the Essays Physical and Literary; and Dr. Garden has confirmed it in Vol. III. of the same publication, and has given a figure and particular description of

the plant.

The root is given both in powder and infusion; but the powder is esteemed most efficacious. The dose is not accurately ascertained, but extends from twelve to sixty or seventy grains of the powder. It is found to be most efficacious when it purges, which it does not always do without some additions. The exhibition of a vomit previous to the use of the Indian pink has proved very serviceable. It sometimes produces disagreeable effects on the nervous system, such as giddiness, dimness of the sight, and convulsive motions of the muscles of the eye. These, according to Dr. Garden, are more likely to happen from a small dose than a large one, the latter more certainly proving purgative or emetic. Dr. Lining, on the other hand, represents these effects as consequent upon too large a dose. It is said to act powerfully as a sedative in abating the exacerbations of low remittent worm-fevers.

SPINA CERVINA.

SPINA CERVINA Pharm. Lond. Rhamnus catharticus sive Spina cervina Pharm. Edinb. Rhamnus catharticus C. B. & Linn. Spina infectoria et cervispina quibusdam. Buckthorn: a prickly bush, or low

low tree, common in hedges: with oval pointed leaves; producing in June small greenish flowers; and about the beginning of October ripening its black berries, which contain a dark green juice, with four seeds in each. The berries of the black alder and dogberry tree, which are frequently, in our markets, mixed with or substituted for those of buckthorn, may be distinguished, by their juice having no greenness, and by their containing only one or two seeds.

BUCKTHORN berries have a faint unpleasant smell, and a bitterish, acrid, nauseous taste. They operate briskly by stool; and occasion, at the same time, a thirst and dryness of the mouth and throat, and not unfrequently fevere gripes, especially if water-gruel or other foft diluents are not freely drank foon after taking them. The dose is faid to be, about twenty of the fresh berries in substance; twice or thrice that number in decoction; a dram, or a dram and a half, of the dried berries; an ounce of the expressed juice; or half an ounce of the rob or extract obtained by inspissating the juice. Among us they have been employed only in the form of a fyrup, in which they feem to operate less unkindly than in any other, and which is given by itself in doses of three or four spoonfuls, or mixed in smaller quantities with other cathartics. The college of Edinburgh directs the fyrup to be prepared by boiling the depurated juice with fugar to a due confistence: that of London adds a little cinnamon, ginger, and nutmegs, with a view to cover in some degree the ill flavour of the buckthorn: but notwithstanding this improvement of the medicine, it is still so unpleasant and so churlish, that it has now almost fallen into disuse.

Syrupus e fpina cervina.

SPIRITUS VINOSUS.

VINOUS SPIRIT: an inflammable fluid, obtained by distillation from wines or other fermented liquors. As first distilled, it partakes both of the phlegm or watery part, and of the oil, of the fermented liquor; which oil, in the liquors commonly used for this purpose, is nauseous and fetid.

i. Spiritus vinosus rectificatus *Pharm. Lond. & Edinb.* Rectified spirit of wine: a vinous spirit purified as much as possible both from its phlegm and ill smell.

Spirits

Spirits drawn from wine, fuch as French brandy, may be in great measure purified by simple distillation, in tall vessels, with a gentle heat, the pure spirituous part rising before the phlegm: if French brandy be thus distilled to one half, the distilled spirit proves tolerably pure. But wine or brandy being in this country too dear an article for distillation; and all vinous spirits, when perfectly purified, being one and the same thing; this purification is chiefly practifed, among us, on the cheaper spirits of melasses and malt liquors. These spirits, when freed by distillation from greatest part of their phlegm, are still found, particularly the latter, to abound with a very offensive oil. To separate this, they are mixed with equal their quantity of spring water, and the spirit gently drawn off again: a considerable portion of the oil is thus left behind in the water, which now proves turbid, and milky, and very nauseous both in smell and taste. By repeating this ablution with fresh quantities of water, the foulest and most offensive spirits may be purified from all ill flavour.

Though spirits, by this treatment, may be divested of their oil, they cannot be freed wholly from phlegm; the gentlest heat, in which they can be distilled, being sufficient to raise a little watery vapour. To complete the purification, therefore, a little fixt alkaline falt, thoroughly dried and powdered, is added; which, imbibing the phlegm, is thereby diffolved into a ponderous liquid, that does not mingle with the spirit, but fettles at the bottom. If the spirit is very phlegmatic, four pints will require a pound of the alkali: if the distillation has been performed with due care, half this quantity, or less, will be sufficient: in either case, if all the falt disfolves, the spirit is to be digested with a little more, till at least a part remains undiffolved. The spirit, now poured off, is to be again distilled, in order to separate from it a portion of the salt, which has united with it, and which, though extremely minute, is fufficient to vary, in some respects, its qualities. As some particles of the alkali are apt to be carried up with it even in the distillation, so as to communicate an ill flavour, it is adviseable to previously add a small portion of calcined vitriol or burnt alum, which will completely abforb the alkali, without giving any new impregnation to the spirit. In this manner was prepared the spirit used in the experiments of the present work under the name of rectified or pure spirit of wine.

4 M

Vinous

Vinous spirits, thus rectified, have a very hot pungent taste, without any particular slavour. They readily take fire, and burn totally away, without leaving any mark of an aqueous moisture behind; though on catching the vapour that exhales from the slame, a considerable quantity of mere water is collected. On distilling them with the gentlest heat, the last runnings prove as colourless, slavourless, and inflammable, as the first. They dissolve distilled vegetable and animal oils, and all the pure resins, into an uniform transparent sluid. They are the lightest of almost all known liquids: expressed oils, which swim on water, sink freely in these to the bottom: a measure which holds ten ounces by weight of water, will contain little more than eight and a quarter of pure spirit.

+ Ph. Lond.

2. Spiritus vinosus tenuior Ph. Lond. & Edinb. Proof spirit: the same spirit containing an admixture of an equal quantity [by measure] of water. "The best proof spirit is that distilled from French wine; "but for common uses, may be employed the spirit drawn from the sprupy matter which separates in the purification of sugar, commonly called melasses."—The spirits usually met with, under the name of proof, are those distilled from different sermented liquors, freed from their phlegm and their slavour only to a certain degree. Their purity, with regard to slavour, may be judged by the taste, especially if the spirit be first duly diluted: of their strength, or the proportion of phlegm contained in them, the least uncertain criterion seems to be their gravity, which is estimated most commodiously by the hydrometer. For the nicer purposes, a pure slavourless proof spirit may be obtained by mixing the foregoing rectified spirit with an equal measure of pure water.

RECTIFIED spirit coagulates all the fluids of animal bodies, that have been tried, except bile and urine. It hardens the solid or consistent parts, and preserves them from corruption. Applied externally to living animals, it strengthens the vessels, contracts the extremities of the nerves, and deprives them of sensibility: hence its power of restraining hemorrhagies, abating superficial pains, &c. Received into the stomach, undiluted, it produces the like effects; thickening the sluid, and contracting all the solid parts which it touches, and destroying, at least for a time, their use and office: if the quantity taken is considerable, a palfy, or apoplexy, follows, and speedily proves mortal.

Proof

Proof spirits, and such as are diluted below the proof strength, have the same effects in a lower degree. Externally they are of use in corroborant, anodyne, and antifeptic fomentations. Taken inwardly, in fmall quantity, they strengthen lax fibres, incrassate thin sluids, and warm the habit: in larger quantity, they diforder the fenses, destroy voluntary motion, and produce, like the rectified spirit, a mortal apoplexy or palfy. -Vinous spirits, therefore, in small quantity and properly diluted, may be applied to useful purposes in the relieving of some disorders; whilst in larger ones, or imprudently continued, they act as a poison of a particular kind. Their moderate use is the most serviceable to those, who are exposed to heat and moisture, to corrupted air, or other causes of colliquative and putrid diseases; the most pernicious in the opposite circumstances, and to those who are afflicted with hysterical or hypochondriacal complaints; for whatever temporary relief these spirituous cordials may afford in the lownesses to which hysterical and hypochondriacal perfons are fubject, we entirely agree with Dr. Pemberton, that there are none who feel fo foon the ill effects arifing from their habitual ufe.

SPONGIA.

SPONGIA Pharm. Lond. & Edinb. SPONGE: a foft, light, very porous and compressible substance, readily imbibing water; found in the fea, adhering to rocks, particularly in the Mediterranean, about the islands of the Archipelago. It has been commonly supposed a vegetable production, but is more probably, like the corallines, of animal origin. Chemically analysed, it yields, like animal substances, a volatile alkaline falt, and this even in larger quantity than I have obtained from any of the other animal matters except the bags of the filk-worm: the caput mortuum, incinerated, yields also a large proportion of fixt falt, not however an alkaline one like that of vegetables, but chiefly of the marine kind: a like falt is obtainable by boiling the sponge in water without burning.

Dry sponge, from its property of imbibing and swelling by moisture, is fometimes used as a tent for dilating wounds and ulcers: for this purpose, after being carefully freed from the small stones generally lodged in it, it is dipt in melted wax, and the wax squeezed out from it in a press-+. * It has also been found to be the most efficacious of all those fubstances

† Spongia præparata Ph. Parif.

fubstances which have been employed to suppress hemorrhagies on the ground of their strong adhesion to the mouths of the wounded vessels; such as agaric, puff ball, \mathfrak{Sc} . For this purpose, a very dry and solid piece, of a cubical or conical form, should be applied in close contact with the vessel, and retained by proper compression. It soon adheres with great force; and indeed its difficult removal is one of its chief inconveniencies. Very large arteries have been prevented from bleeding by this application (a).

† Spongia usta Pharm. Lond. & Ed. Burnt in a close earthen vessel, till it becomes black and friable+, it has been given in doses of a scruple against scrophulous complaints and cutaneous desedations; in which it has sometimes been of service, in virtue, probably, of its saline matter, the proportion of which, after the great reduction which the other matter of the sponge has suffered in the burning, is very large. By virtue of this saline matter also, the preparation, if ground in a brass mortar, corrodes so much of the metal, as to contract a disagreeable taint and sometimes an emetic quality: hence the college expressly orders it to be powdered in a mortar of glass or marble. *The burned sponge is a principal article in what is called the Coventry method of cure in the bronchocele, and also in that published by Mr. Prosser(b).

STANNUM.

metal, not liable to rust, but losing its brightness in the air; easily slexible, and making a crackling noise in being bent; little more than seven times specifically heavier than water; sussible in a heat far below ignition, and somewhat less than that in which lead melts. Heated till almost ready to melt, it proves extremely brittle, so as to fall in pieces from a blow. Melted, and nimbly agitated at the time of its beginning to congeal (as by shaking in a wooden box rubbed on the inside with chalk) it is reduced partly, and by repetitions of the process totally, into powder. Continued in sussion for some time, and kept stirring with an iron rod, it changes into a dusky calx; which, urged longer in the fire, gains a perfect whiteness, a mark of the purity of the tin. It is corroded

Stannum pulveratum Pharm. Lond. Stanni pulvis Ph. Edin.

⁽a) See White's Cases in Surgery.

⁽b) See Wilmer's Cases in Surgery, and Proster on the Bronchocele.

by vegetable acids, and renders them turbid and whitish: the nitrous acid pretty readily diffolves it, but foon deposites a part in form of a thick mucilage, especially if the acid has any admixture of the vitriolic: the vitriolic and marine acids are very difficultly made to act upon it: its most perfect menstruum is a mixture of the marine and nitrous.

THE principal use of this metal in the present practice is as an anthelmintic: even the flat worms, which too often elude the force of other medicines, are faid to be effectually destroyed by powdered tin. The common dose of the powder is from a scruple to a dram, but Dr. Alston affirms, in the Edinburgh medical esfays, that its fuccess depends chiefly on its being given in much larger quantities, as half an ounce or an ounce. It is possible, that the anthelmintic virtues of tin may proceed, not so much from the pure metal, as from a certain substance of a different nature, which there are grounds to suspect that the purest forts of tin usually met with, participate of; filings of tin, held in the flame of a candle, emit a thick fume fmelling like garlick: Mr. Marggraf reports (a), that by gentle diffolution in aqua regis and flow evaporation, he obtained crystals, which on being exposed to the fire, with the addition of some fixt alkaline falt to absorb their acid, sublimed into a white concrete; and that this exhaled in the fire in fumes of a strong garlick smell, formed with sulphur yellow and red compounds, and whitened copper (see Arsenicum). It must be observed, however, that notwithstanding these strong presumptions, not to say proofs, of an arfenical impregnation in tin, the metal taken in substance has not been observed to be noxious, though the fumes which it emits in a red heat are undoubtedly fo.

A sparkling gold-coloured preparation of tin, called mosaic gold, is prepared by adding fix ounces of quickfilver to twelve of melted tin, Aurur. mapulverizing the mass when grown cold, mixing with it seven ounces of flowers of fulphur and fix of fal ammoniac, and fubliming in a matras: the mosaic gold is found under the sublimed matter, with some dross at This preparation is chiefly valued for its beautiful appearance: as a medicine it is at prefent little regarded, though formerly held in confiderable efteem against hysterical and hypochondriacal com-

plaints, malignant fevers, and venereal diforders. It appeared, upon experiment, to be little more than a calx of tin: tin, calcined by itself, gains nearly as much in weight, as it does by being made into mosaic gold; and the mosaic gold, melted with inflammable fluxes, is revived into tin again without suffering much more loss than the simple calx. The volatile ingredients, sal ammoniac, sulphur, and quicksilver, sublime in the process, partly escaping, and partly forming the scoriæ: great part of the sulphur and mercury are found united together into the form of cinnabar.

Sal Jovis.

A falt of tin is directed to be prepared, from twelve ounces of calx of tin and four of aqua regia diluted with twenty-four of water: after digestion for two days, the vessel is to be shaken, the more ponderous part of the undissolved calx suffered to settle, the turbid liquor poured off and evaporated nearly to dryness, and the mass further exsiccated on brown paper: to the remaining calx, half the quantity of fresh menstruum is to be added, and the process repeated. Of the virtues of this salt I can say nothing from experience, except that it is in taste very sharp and almost corrosive. Nor do I apprehend the use of calcining the metal, as tin uncalcined dissolves much more easily and more plentifully: the solution is in both cases the same, the fire in the calcination dissipating only the inflammable principle of the tin, which the acid equally does in the solution and evaporation. Hossman says, that solution of tin is a strong purgative.

STAPHISAGRIA.

STAPHISAGRIA Pharm. Edinb. (a) & C. B. Staphys, pedicularia, & herba pedicularis quibusdam. Delphinium Staphysagria Linn. Stavesacre: a plant with large leaves, set on long pedicles, deeply divided into several segments; producing irregular blue flowers with a tail behind like those of larkspur, followed by pods containing large, rough, triangular, dark-coloured seeds. It is annual, a native of the southern parts of Europe, from whence the shops have been generally supplied with the seeds.

THE feeds of stavesacre have a disagreeable smell, and a very nauseous bitterish burning taste. They were formerly employed sometimes as a cathartic, in doses of from twelve grains to a scruple: but they operate with so much violence both upwards and downwards, and are so liable not only to disorder the bowels, but likewise to instame the throat, that their internal use has been long laid aside. They are now used only in external applications, for some kinds of cutaneous eruptions, and for destroying insects. Their acrimony is extracted partially by water, totally by rectified spirit, and not elevated in distillation by either.

STOECHAS.

STOECHAS Pharm. Lond. Stæchas purpuræa C. B. Lavandula Stæchas Linn. French lavender: a low shrubby plant, with small oblong narrow leaves, bearing on the tops of the branches short thick spikes or scaly heads, from which issue several small purple labiated slowers, followed each by four seeds inclosed in the cup. It is a native of the southern parts of Europe, common in our gardens, and slowers in May or June. The shops have been generally supplied, from Italy and the south of France, with the slowery tops, often mouldy, and never equal to those of our own growth.

THE best stechas which we receive from abroad has no great smell or taste; Pomet affirms, that such as is to be met with in the shops of Paris is entirely destitute of both; whereas ours, both whilst fresh and when carefully dried, has a pretty strong aromatic smell, and a moderately warm pungent bitterish taste. Distilled with water, it yields a considerable quantity of a pale-coloured fragrant essential oil: the remaining decoction is unpleasantly bitterish and subastringent. With rectified spirit, it gives over nothing considerable, greatest part of the active matter of the stechas being left in the extract. Both the herb itself and its preparations are much less grateful than lavender, with which it is supposed to have some agreement in virtue.

* STRAMONIUM.

STRAMONIUM Pharm. Edinb. Solanum fætidum, pomo spinoso oblongo C. B. Datura Stramonium Linn. THORN-APPLE: an herbaceous

baceous plant, with a thick branched stalk, two or three feet high, large sinuated indented leaves, and long tubular white or purplish flowers, succeeded by large, prickly, green, sleshy seed-vessels, which open at the end in four divisions, and disclose numerous black seeds. It is sown in gardens, and sometimes found wild among rubbish. It slowers in July.

This plant, which has been long known as a narcotic poison, has been introduced into the catalogue of medicines by Dr. Stærck. An extract made from the expressed juice of the leaves is acrid and saline to the taste, and yields crystals of nitre on standing. This preparation given in doses of from one to five grains twice or thrice a day, is said to be a very powerful remedy in various convulsive and spasmodic disorders, epilepsy and mania. The accounts of other practitioners have confirmed that of the first introducer, and it has been received into some pharmacopæias.

An abridged account of its medicinal properties, with some instances of its efficacy, from a treatise printed at Upsal by Dr. Wedenberg, is to be met with in the *Med. Comment*. Vol. III. p. 18. An ointment prepared from the leaves has been found to give ease in external inflammations and hamorrhoids.

Several instances are recorded of the bad effects of inadvertently eating the seeds of thorn-apple. Emetics and purgatives give the speediest relief in these cases, which it is sometimes necessary frequently to repeat, as some of the seeds have been found to lodge a considerable time in the stomach.

STYRAX.

SOLID STORAX: an odoriferous refin, exuding in the warmer climates from a middling-fized tree (flyrax folio mali cotonei C. B. Styrax officinale Linn) with leaves like those of the quince, flowers like those of the orange tree, and fruit like filberds; a native of Asia, and, as is said, of Italy. Two forts of this refin have been commonly distinguished in the shops.

1. Storax in the tear: not in separate tears, or exceeding rarely, but in masses, sometimes composed of whitish and pale reddish brown tears, and sometimes of an uniform reddish-yellow or brownish appearance; unctuous

unctuous and foft like wax, and free from visible impurities. This is supposed to be the fort which the ancients received from Pamphylia in reeds or canes, and which was thence named calamita.

2. Common storax: in large masses, considerably lighter and less compact than the foregoing, and having a large admixture of woody matter like faw-duft. This appears to be the kind intended by the London college, as they direct their flyrax calamita to be purified, for medicinal Styrax colause, by softening it with boiling water, and pressing it out from the feces betwixt warm iron plates; a process which the first fort does not stand in need of. And indeed there is rarely any other than this impure storax to be met with in the shops.

tus Ph. Lond.

The writers on the materia medica in general prefer the pure storax in the tear, and reject that which is mixed with woody matter. It appears however, upon comparison, that this last, notwithstanding its large proportion of impurities, is the most fragrant of the two: nor is it difficult to affign a reason for this superiority, as the pure juice must have required, for its inspissation to a firm consistence, a longer exposure to the fun and air, and confequently loft more of its volatile parts, than when absorbed and thickened by the woody substance.

Common storax, infused in water, imparts to the menstruum a gold yellow colour, some share of its smell, and a slight balsamic taste. It gives a confiderable impregnation to water by distillation, and strongly diffuses its fragrance when heated, though it scarcely yields any effential Hence, in the purification of it by straining, it is apt to suffer a confiderable loss of its finer matter, which is partly diffipated by the heat, and partly kept dissolved by the water: a part of the storax is likewise defended by the woody substance from the action of the press, and left behind among the feces. It may be purified rather more elegantly by means of rectified spirit, which readily dissolves the fine refin, leaving only the impurities and a little inert gummy matter: the spirit gently distilled off from the filtered reddish-yellow solution, brings over with it very little of the fragrance of the storax; and the remaining refin is more fragrant than the finest storax in the tear which I have The pure refin, distilled without addition, yields, along with an empyreumatic oil, a portion of faline matter fimilar to the flowers

4 N

of benzoine; I have fometimes also extracted from it a substance of the same nature by coction in water.

Storax is one of the most agreeable of the odoriferous resins, of a mild taste, of no great heat or pungency, nearly similar, in its medical as in its pharmaceutic qualities, to benzoine and balsam of Tolu. It is not, however, much used in common practice, unless as an ingredient in some of the old compositions, and in an opiate pill to which it gives name in the shops.

Pil. e styrace Pharm. Lond.

STYRAX LIQUIDA.

STYRAX LIQUIDA Pharm. Edinb. (a) LIQUID STORAX: a refinous juice; obtained from a large tree with angular leaves like those of the maple, and a round fruit composed of a number of pointed seed-vessels, called by Ray syrax aceris folio, by Linnæus Liquidambar styracistua, a native of Virginia and Mexico, and lately naturalized to our own climate. The juice called liquidambar is said to exude from incisions made in the trunk of this tree, and the liquid storax to be obtained by boiling the bark or branches in water.

Two forts of liquid storax are distinguished by authors: one, the purer part of the resinous matter that rises to the surface in boiling, separated by a strainer, of the consistence of honey, tenacious like turpentine, of a reddish or ash brown colour, moderately transparent, of an acrid unctuous taste, and a fragrant smell, faintly resembling that of the solid storax, but somewhat disagreeable: the other, the more impure part, which remains on the strainer, untransparent, in smell and taste much weaker, and containing a considerable proportion of the substance of the bark. What is most commonly met with under this name in the shops, is of a weak smell, and a grey colour, and is supposed to be an artificial composition.

Liquid storax has been employed chiefly in external applications. Among us, it is at present almost wholly in disuse.

(a) Expunged.

SUCCINUM.

SUCCINUM Pharm. Lond. & Edinb. Ambarum citrinum & electrum AMBER: a folid, brittle, bituminous substance, dug out of the earth or found upon the sea shores, most plentifully in Polish Prussia and Pomerania; of a white, yellow, or brown colour, sometimes opake, and fometimes very clear and transparent; of very little tafte; and fearcely any fmell, unless heated or briskly rubbed, in which circumstances it yields a pretty strong one, to most people agreeable.

Boiled in water, it neither foftens, nor undergoes any fenfible alteration. Digested in rectified spirit, it imparts a yellowish or brownish colour, a fragrant smell, and a bitterish aromatic taste: by repetitions of the process with fresh quantities of spirit, a considerable part of the amber by degrees diffolves. The spirit, distilled off from the tinctures, is strongly impregnated with their fmell; nevertheless the remaining balfam, or soft

extract, is found to be very strong both in smell and taste.

By alkalies, fixt and volatile, the vegetable, nitrous, and marine acids, it is fearcely at all acted upon: the vitriolic acid diffolves it into a deep purple liquor, from which the amber is precipitated on the mixture of

any other acid, or of water, or spirit of wine (a).

The spirituous tincture and balfam are medicines of great efficacy in hysterical disorders, cachexies, the fluor albus, some rheumatic pains, and in debilities and relaxations in general: in fome cases of this kind they have taken place, after bark and other corroborants of the vegetable kingdom had been given with little effect. The spirit, which distils in concentrating the tincture, may be referved for extracting a fresh tincture, either from another parcel of amber, or from that which remained after the former extraction. It is faid that if a little vitriolic acid be previously combined with the spirit, it will dissolve more of the amber than pure vinous spirits. The amber is sometimes given also in substance, levigated Succinum into an impalpable powder, but does not appear to act with fo much ad- preparatum Phar. Lond. vantage in this form as in a disfolved state.

This concrete, exposed to the fire in open vessels, melts into a black mass, takes flame, emits a copious smoke, with a smell like that which arises from the finer kinds of pitcoal, and burns almost entirely away. † Spiritus fuccini Ph. Lond. Distilled in a retort, it yields first an acidulous phlegm intermingled with a thin limpid oil, which grows thicker and deeper coloured as the fire is increased: at length a brownish faline matter arises into the neck of the retort, succeeded by a grosser oil, and at last, in a great heat, by a black thick pitchy matter. About the time that the first oil begins to rise, the amber melts in the retort, and, unless the heat be cautiously regulated, is apt to boil over into the receiver: to prevent this accident, some previously mix with the amber an equal quantity of clean sand, which does not appear, however, to be of much use, for with due care the process succeeds equally without as with it.

† Sal fuccini Ph. L. & Ed.

The falt is purified from its adhering oil, either by fublimation, or by repeated folution, filtration, and crystallization 1. When perfectly pure, it is of a white colour, and of a penetrating gratefully acid subastringent tafte. It disfolves in rectified spirit sparingly by the assistance of heat, not at all in the cold. Of cold water, in the common temperate state of the atmosphere, it requires for its folution above twenty times its own weight; of boiling water, only about twice its weight: in flow cooling, it shoots into triangular prismatic crystals, with the points obliquely truncated. In the heat of boiling water, it does not exhale, or fuffer any visible alteration: in a greater one, it first melts, then rises in white fumes, and concretes again in the upper part of the glass into fine white flakes; leaving behind a fmall quantity of a dark coaly matter. It effervesces with alkalies and absorbent earths, and forms with them compound falts fomewhat resembling those made with vegetable acids, its acid matter feeming to have a confiderable analogy to the acids of the vegetable kingdom, and being effentially distinct from the three called mineral acids (a): mixed with acids, it makes no fensible commotion. By these characters this salt may be distinguished from all the other matters that have been mixed with or vended for it. With regard to its virtues, it is accounted aperient, diuretic, and antihysteric: its great price has prevented its coming much into use, and probably its real virtues, though doubtless considerable, fall greatly short of the opinion that has been generally entertained of them.

Ol. fuccini | Ph. Edinb. 6 Ph. Lond.

The oil ||, distilled again by itself, is divided into a thinner oil which arises \(\), and a thicker part, which remains behind, called balsam of

amber †: fome distil it from brine of sea salt, or from plain water ‡, by which it becomes purer than when distilled without addition. This oil has a strong bituminous smell, and a hot pungent taste; and approaches more to the nature of the mineral petrolea than of the vegetable or animal distilled oils, being very difficultly, if at all, dissoluble in vinous spirits. It is sometimes given internally, in doses of ten or twelve drops, as an antihysteric and emmenagogue; and sometimes employed externally in antihysteric, paralytic, and rheumatic liniments or unguents.

+ Balfamum fuccini P. L. ‡ Ol.fuccini rectificatum Phar. Edin.

SULPHUR.

SULPHUR Pharm. Lond. BRIMSTONE: a yellow concrete, of no taste, and scarcely any smell; melting in a small degree of heat into a viscous red fluid, and totally exhaling on an increase of the heat; readily inflammable, and burning with a blue slame and a suffocating acid sume.

It consists of the vitriolic acid combined with a small proportion of inflammable matter. If a combination of pure vitriolic acid with a pure fixt alkaline salt be melted in a close vessel with the addition of a little powdered charcoal, a true sulphur will be produced, and the compound will be the same (excepting for the earthy part of the charcoal) as if the alkali had been melted with common brimstone. And contrariwise, if a combination of alkaline salt with common brimstone be reduced into powder, and roasted with a gentle heat, the inflammable principle exhales, and the remainder proves the same as if the alkali had been combined with the pure vitriolic acid: the diminution of weight, resulting from this avolation of the inflammable principle, does not exceed two drams upon sixteen ounces of the sulphur (a).

Greatest part of the sulphur met with in the shops is either extracted from certain ores by a kind of distillation (b); or prepared from minerals abounding with vitriolic acid, by stratifying them with wood, which being set on fire, the sulphur is collected in cavities made in the upper part of the pile (c). The largest quantities are brought from Saxony, in irregular masses, which are afterwards melted and cast into cylindrical rolls. Sulphur is found likewise native in the earth; sometimes in

⁽a) Vide Stahlii Mensis Julius, Experimenta & animadversiones ccc, &c.

⁽b) Leopold, Relatio de itinere suo Suecico.

⁽c) Hoffman, Observationes physico-chemicæ, lib. iii. obs. 9.

transparent pieces, of a greenish or bright yellow colour; more commonly in opake grey ones with only fome streaks of yellow: this last is the fort which is understood by the name fulpbur vivum, though what is fold under this name in the shops is no other than the dross which remains after the fublimation of fulphur. The native fulphurs should never be employed for any internal use without purification: they almost always participate of arfenic, which is discoverable in some by their having naturally more or less of a red colour, and in the others by their exhibiting this colour after a part of the pure fulphur has been separated by fublimation.

Flor. fulph. Phar. Edin.

Sublimation is the most effectual method of purifying sulphur from arfenical as well as earthy admixtures; and by the same process it is reduced into a fine powder, somewhat of a softer kind than that obtained by triture. Those who prepare the flowers in the way of trade, use for the fubliming veffel a large iron pot, capable of holding two or three hundred weight: this is placed under an arched chamber, lined with glazed tiles, which ferves for the recipient. Some fmall portion of fulphur that rifes first, especially when the vessels are very large, or the air not fufficiently excluded, is apt to take fire, and give out its acid, which adhering to the flowers that fublime afterwards, communicates to them a fensible acidity or roughness; in consequence of which, they are fometimes found to coagulate milk, when taken internally to produce gripes, and to receive from some metalline vessels a disagreeable taint: hence the London college directs fuch of the flowers, as may happen to concrete or melt together from the vicinity of the receiver to the fire, to be reduced to powder, not with metalline instruments, but either in + Flor. fulph. a wooden mill, or in a marble mortar with a wooden peftle+. From this extraneous or superficial acid they are freed, by boiling them in water, t Flor. fulph. and afterwards carefully washing them with cold water !.

Pharm. Lond.

loti Pb. L.

Pure fulphur, taken in doses of from ten grains to a dram or more, gently loofens the belly, and promotes perspiration. It seems to pass through the whole habit; and manifestly transpires through the skin, as appears from the fulphureous fmell of persons who have taken it, and filver being stained in their pockets to a blackish hue as by the vapour of fulphureous folutions. In confequence of these properties, and of this fubtility of parts, it promifes to be of great medicinal powers;

but

but what its particular virtues are, experience has not as yet clearly determined. It is principally recommended against the piles, in disorders of the breast, and in cutaneous eruptions: in the itch indeed it is a certain remedy, whether internally or externally used, but in other kinds of eruptions it has not equally succeeded, and perhaps its efficacy against the first depends not so much on its purifying the blood, as on its fumes being destructive to the cuticular animalcules to which the present theory ascribes that distemper. It remarkably corrects or restrains the power of certain mineral substances of the more active kind: by the admixture of fulphur, mercury becomes inert, the virulent antimonial regulus mild, and arfenic itself almost innocent: hence though sulphur should contain a fmall proportion of arfenic, it possibly may not receive from that poisonous ingredient any very hurtful quality.

This concrete is not acted on by water, by acids, or by vinous spirits; but disfolves, by the assistance of heat, in oils both expressed and distilled, and in the mineral petrolea: when diffolved, it yields a very offenfive fmell, and difcovers to the tafte a naufeous pungency and heat. Expreffed oils and petroleum dissolve it more readily and more plentifully than the distilled, taking up so much as to become thick and almost consistent: the college of London directs one part of flowers of fulphur and four of oil-olive +, and the same proportions of the flowers and of the + simplex. petroleum called Barbadoes tart, and that of Edinburgh one ounce of tbarbadense. the flowers and eight ounces of oil-olive |, to be boiled together till || craffum. they unite into the confistence of a balfam. Essential oils do not load themselves so much with the sulphur as to become thick. As soon as the fulphur begins to be ftrongly acted on either by expressed or essential oils, which happens nearly about the point of ebullition, or in fuch a heat as the fulphur by itself would melt in, the matter rarefies and fwells up greatly, fo as to require the veffel to be very large and occasionally removed from the heat; and at the same time throws out impetuously great quantities of an elastic vapour, which, if a free exit is not allowed it, produces violent explosions. The volatile flavour of the effential oils is in great measure diffipated in this process by the great heat requisite for effecting the folution: a more elegant composition of this kind might be obtained by adding to the effential oil a proper quantity of the balfam made with expressed oils, which will unite with it by a gentle warmth. The balfams of fulphur have been employed externally for cleanfing and healing

healing foul running ulcers. They are recommended internally in some cachectic and hydropic cases; as also in coughs and consumptions, in which they promise, by their manifest heat and acrimony, to be oftener injurious than beneficial: they have been frequently observed to hurt the appetite, and excite febrile symptoms.—It may be observed, that in these solutions the component parts of the sulphur are in some measure disunited from one another; insomuch that a considerable quantity of sulphureous acid, but no actual sulphur, separates in distillation.

Hepar fulphuris.

Tinctura fulphuris.

Syrupus fulphuris. Fixt alkaline falt, stirred by little and little into twice [or rather half] its weight of sulphur in suspense with it into a red mass called from its colour liver of sulphur. This compound has a fetid smell, and a nauseous taste: it dissolves in water, and deliquiates in the air, into a yellow shuid: thrown, whilst hot from the fire, into rectified spirit of wine, in the proportion of about sour ounces to a pint, and digested about twenty-four hours, it communicates a rich gold colour, a particular not ungrateful smell, and a hot somewhat aromatic taste. Solutions of the liver in water, made with sugar into a syrup; and a few drops of the tincture mixed with a glass of canary or other rich wine, to which it communicates a milky hue; have been sometimes given in the same intentions as the balsams, and seem to be accompanied with the same inconveniencies.

Flowers of fulphur may be dissolved in water by boiling them with thrice their weight of quicklime, though not so readily as by alkaline salts. If the filtered solution be long exposed to the atmosphere, or if air from the lungs be blown into it for a short time through a glass pipe, the lime gradually separates, as it does from common lime-water; and the sulphur, which was dissolved by means of the lime, separates and precipitates along with it. Common alkalies, fixt or volatile, added to the solution of sulphur in lime-water, occasion a precipitation of the lime, the sulphur continuing dissolved: caustic alkalies make no precipitation.

On adding to the fulphureous folution, whether made by lime or by alkalies, some of the weak spirit of vitriol, (or any other acid) the liquor becomes milky, an extremely setid and diffusive inflammable vapour arises, and on standing for some time the sulphur settles to the bottom in form of a white powder, which, when washed with fresh quantities of water, becomes insipid and inodorous, and is vulgarly called sac or milk of sulphur: the liquor after the precipitation retains still a sulphureous impregnation,

impregnation, which further additions of acid will not precipitate. The method of preparing the lac with fixt alkalies is the most expeditious and least troublesome, provided the sulphur has been thoroughly united with a fufficient quantity of the alkali; and on the other hand, quicklime + gives the preparation a more faleable whiteness. The medicine + Sulphur proves in either case nearly the same: it would be exactly the same if præcipitat. the precipitation was made with any other acid than the vitriolic; which forms with the dissolved lime a selenitic concrete, not separable from the lac by any ablution, but with the alkali a neutral falt, which by hot water may be totally diffolved and washed off; whereas the combinations of all the other acids, with lime as well as with alkalies, are eafily diffoluble even in cold water. The pure lac is not different in quality from pure fulphur itself; to which it is preferred, in external applications, only on account of its colour. The whiteness does not proceed from the sulphur having loft any of its parts in the operation, nor from any new matter superadded: on being melted with a gentle fire, it resumes its yellow hue.

A folution of fulphur in volatile alkaline spirits may be obtained, by Tinet. fulph. boiling half a pound of flowers of fulphur with a pound of quicklime, in a gallon of water, till half the liquor is wasted; then putting the remainder into a retort, with eight ounces of powdered fal ammoniac, and distilling with a gradual fire. The spirit comes over loaded with the fulphur, and has a strong offensive smell, somewhat resembling that which rifes in the precipitation of the lac. Hoffman fays, a mixture of it with thrice its quantity of spirit of wine, given in doses of thirty or forty drops, proves a powerful diaphoretic; and that applied externally as a fomentation, with the addition of camphor, it alleviates gouty pains.

The flowers of fulphur in fubstance seem to be preferable for internal use to any of the preparations: they are certainly more safe, and perhaps not less effectual; as they do not heat or irritate the first passages, and yet are evidently diffolved in the body and carried through the habit. They are most commodiously taken in the form of troches: the college Trochisci of London directs for this purpose two ounces of the washed flowers, and four of double refined fugar, to be beaten together, and made up with mucilage of quince feeds; that of Edinburgh, one ounce of the flowers of fulphur, ten grains of flowers of benzoine, fifteen grains of factitious Trochifei e cinnabar, and two ounces of fine fugar, to be formed with mucilage of gum tragacanth: by the addition of the flowers of benzoine in this last Phar. Edinb.

e fulphure

volatilis

vulgo.

fulphure five diafulphuris

prescription, the medicine is supposed to be rendered more efficacious in some disorders of the breast.

Unguentum e fulphure † Ph. Lond. five antipforicum ‡ Ph. Edin.

A fulphureous ointment, for the itch, is prepared, by mixing two ounces of the unwashed flowers, with fix + or eight tounces of the simple ointment called pomatum+, or of hogs-lard‡, and a scruple of effence of lemons. Half this quantity is, in most cases, sufficient for a cure; though it may be proper to renew the application, and touch the parts most affected, for some nights longer, till the whole quantity is exhausted. Some have been of opinion, that this external use of sulphur is unsafe; that as fulphur taken inwardly promotes the expulsion of impure humours and the eruption of cutaneous efflorescences, it must act, when outwardly applied, by repressing them. This consequence, however, does not follow; nor is it by affecting the humours that it performs the cure: for it equally removes the itch, whether used internally or externally, by its vapours diffused through the skin. All the danger, that is to be apprehended from fulphureous unquents, is that which may arise from the obstruction of the cutaneous pores by the unctuous matter; and to prevent any diforders from this cause, only a part of the body is to be anointed at one time.

SUMACH.

SUMACH five Rhus obsoniorum Pharm. Edinb. (a) Rhus folio ulmi C. B. Rhus coriaria Linn. Sumach: a shrub or low tree; with oval, pointed, serrated, downy leaves, having each a red rib running along the middle, set in pairs without pedicles; producing clusters of small yellowish or greenish flowers, each of which is followed by a small, red, flattish berry, including a roundish reddish-brown seed. It is a native of the southern parts of Europe, and cultivated in some of our gardens.

THE berries of sumach have an acid austere taste: they were formerly used for restraining bilious fluxes, and hemorrhagies, and colliquative hectic sweats: some direct an infusion of half an ounce of the berries, and others two or three drams of an extract made from them by water, for a dose. The leaves and young twigs are strong astringents, and have been directed in the same intentions.

TACAMAHACA.

TACAMAHACA Pharm. Edinb. (a) A refin; obtained from a tree, resembling the poplar, (Populus balfamifera Linn.) bearing, at the extremities of the branches, small roundish fruits including a feed like a peach-kernel; a native of the temperate parts of the continent of America, and in a sheltered situation enduring the winters of our own country.

Two forts of this refin are fometimes to be met with. The best, called, from its being collected in a kind of gourd-shells, tacamahaca in shells, is somewhat unctuous and soft, of a pale yellowish or greenish colour, a bitterish aromatic taste, and a fragrant delightful smell approaching to that of lavender and ambergris. This fort is very rare. That commonly found in the shops is in semitransparent grains or glebes, of a whitish, yellowish, brownish, or greenish colour, and of a less grateful fmell than the foregoing. The first is said to exude from the fruit of the tree, the other from incisions made in the trunk. The tree, as raised among us, affords in its young buds, or the rudiments of the leaves, a refinous juice of the same kind of fragrance (b).

Tacamahaca is used chiefly as an ingredient in warm nervine plasters; though the fragrance and taste of the finer fort points out its being applicable to other purposes, as an internal balfamic corroborant. Both kinds diffolve in rectified spirit into a gold-coloured liquor, only a small quantity of impurities being left: they impregnate water also considerably with their smell and taste, but give out very little of their substance to this menstruum.

TALCUM.

TALC: an earthy concrete; of a fibrous or leafy texture; more or less pellucid, bright or glittering; smooth and slippery to the touch; in some degree flexible and elastic, so as scarce to be pulverable; soft, fo as to be easily cut; fuffering no change in an intense fire, or no other than a diminution of its brightness, flexibility, and unctuosity; not acted upon by acids, either in its crude state, or after vehement calcination. There are feveral different appearances of this earth; among which, the

(a) Expunged.

(b) See article Populus.

greenish

greenish foliaceous Venice talc has been selected for medicinal use; though it does not appear capable of answering any medicinal intention, as not being dissolved, or sensibly affected, by any known humid menstruum: on account of its unctuous softness, and the silver hue which it exhibits when reduced by rasping or otherwise into powder, it has been employed externally as a cosmetic. The sibrous slexible amianthus or asbestos, and the more rigid sibrous alumen plumosum, seem to approach to the nature of the talcs, and made an equally insignificant addition to the articles of the materia medica in our former pharmacopæias.

TAMARINDUS.

TAMARINDUS Ph. Lond. & Ed. Oxyphænicon. TAMARIND: the fruit of a pretty large tree, (filiqua arabica quæ tamarindus C. B. Tamarindus indica Linn.) growing in Arabia and in the East and West Indies. The fruit is a pod, somewhat resembling a bean-cod, including several hard seeds, together with a dark-coloured viscid pulp: the East India tamarinds are longer than those of the West, the former containing six or seven seeds each, the latter rarely above three or four: they nevertheless seem both to be the produce of one species of plant. The pulp, with the seeds, connected together by numerous tough strings or sibres, are brought to us freed from the outer shell: the oriental fort is dryer and darker coloured than the occidental, and has more pulp: the former is sometimes preserved without addition, the latter has always an admixture of sugar.

THE pulp of tamarinds is an agreeable laxative acid; of common use in inflammatory and putrid disorders, for abating thirst and heat, correcting putrefaction, and loosening the belly. The dose, as a laxative, is two or three drams: an ounce or two prove moderately cathartic. It is an useful addition to the purgative sweets, cassia and manna, increasing their action, and rendering them less liable to produce flatulencies: the resinous cathartics are said to be somewhat weakened by it. Tournesfort relates that an essential salt may be obtained from tamarinds, by dissolving the pulp in water, and setting the filtered solution, with some oil upon the surface, in a cellar for several months; that the salt is of a sourish taste, and difficultly dissoluble in water; and that a like

falt is fometimes found also naturally concreted on the branches of the tree. The salt, as Beaumé observes, may be obtained more expeditiously, by clarifying the decoction of the tamarinds with whites of eggs, then filtering and evaporating it to a proper confishence, and setting it to cool: the salt shoots into crystals, of a brown colour, and very acid taste; but in dissolving and crystallizing them again, or barely washing them with water, they lose almost all their acidity, the acid principle of the tamarinds seeming not to be truly crystallizable.

TANACETUM.

TANACETUM Pharm. Lond. & Edinb. Tanacetum vulgare luteum C.B. Tanasia, athanasia, & parthenium mas quibusdam. Tanacetum vulgare Linn. Tansy: a plant with large leaves, divided to the rib, on both sides, into oblong deeply indented segments; producing, on the tops of the stalks, several gold-coloured discous flowers, in umbel-like clusters, followed by small oblong blackish seeds. It is perennial, grows wild by road-sides and about the borders of fields, and flowers in June and July.

The leaves and flowers of tanfy have a strong, not very disagreeable smell, and a bitter somewhat aromatic taste: the flowers are stronger though rather less unpleasant than the leaves. They give out their virtue both to water and spirit, most perfectly to the latter: the tincture, made from the leaves, is of a fine green, from the flowers of a bright pale yellow colour. Distilled with water, they yield a greenish-yellow esfential oil, smelling strongly of the herb: the remaining decoction, inspissated, affords a strong bitter subsaline extract. The spirituous tinctures give over also, in inspissation, a considerable part of their flavour *(a); a part of it remaining, along with the bitter matter, in the extract.

This plant is used as a warm deobstruent bitter, in weakness of the stomach and in cachectic and hysteric disorders; and likewise as an anthelmintic. The seeds have been chiefly recommended in this last

^{*(}a) Alcohol, distilled from tansy, proved, after standing for upwards of sisteen years, richly impregnated with the slavour of the plant, and sufficiently grateful. M. S. of Dr. Lewis.

intention, and supposed by some to be the fantonicum of the shops, from which they differ not a little in quality as well as in appearance, being much less bitter, and of a more aromatic flavour.

TARTARUM.

TARTARUM vini albi vel rubri Pharm. Lond. TARTAR: an acid concrete falt thrown off from wines, after complete fermentation, to the fides and bottoms of the casks; of a red or white colour, and more or less drossy, according to the colour and quality of the wine. The white is generally purer than the red: both kinds, when purified, are exactly the same.

THIS falt is one of those which are most difficultly dissoluble in water, being scarcely affected by it in the cold, and requiring ten or twelve times its own weight when affifted by a boiling heat. From this faturated folution the tartar begins to separate almost as soon as the boiling ceases: if the quantity of water is greater, as about twenty times the weight of the falt, it continues long enough suspended to be passed, with due care, through a woollen strainer or a filter. The filtered liquor appears nearly colourless, whether the tartar made use of was red or white: if hastily cooled, the falt separates in small grains like fand, but if the veffel is closely covered, and the heat very leifurely diminished, it shoots into semitransparent whitish crystals: if the filtered liquor be kept boiling, a thick skin forms on the surface, which, being taken off with a perforated wooden skimmer, is succeeded by fresh cuticles, till the whole of the falt is thus formed into what is called creme of tartar. The refining of tartar is practifed, in the way of trade, chiefly about Montpellier, from whence the shops are generally supplied both with the crystals and creme; the process being so troublesome, and requiring so large conveniencies, that it is scarcely ever attempted here. A certain earth, of the argillaceous kind, is added in the process, the chief use of which seems to be, to promote the separation of the colouring matter; for the falt extracted from the coloured tartars by water only is feldom of perfect whiteness. It is faid that the earth generally contains fome small portion of chalky matter, soluble in acids, which of consequence will be taken up by the tartar; I have sometimes observed

Crystalli tartari Phar. Lon. & Edin.

Cremor tartari. observed solutions of the crystals to deposite an earthy precipitate on adding alkaline lye. The purer fort of white tartar, unrefined, especially that of Rhenish wine, is, for many purposes, particularly for combinations with other bodies, not inferiour either to the creme or crystals.

Pure tartar, in doses of half a dram or a dram, is a mild cooling aperient: two or three drams gently loofen the belly; and fix or eight prove moderately cathartic. Its acidity and laxative power are its medical characters.

* The difficult folubility of creme of tartar being an objection to its medical use, some experiments were made by Dr. Peter Jonas Berg, for rendering it more foluble by certain additions, without altering its medicinal qualities. Borax was found to answer best for this purpose. To four parts of creme of tartar, one of borax was added. diffolved in a fufficient quantity of water, and the liquor strained. About a fixteenth part of impurities were left behind. The pure folution evaporated yielded an acid and extremely foluble white falt (a).

Tartar, disfolved in water, effervesces with fixt alkaline salts, and faturates, of the vegetable alkalies, near one third its own weight. The compound falt, refulting from their union, is a neutral one, more purgative than the tartar itself, and far easier of solution, whence its name foluble tartar. This falt is prepared, either by boiling the refined tartar in a fufficient quantity of water till it is diffolved, and then dropping in strong alkaline lye; or by dissolving the alkali in boiling water in the proportion of a pound to a gallon+, or to fifteen pounds I, and Tart. folubil. then adding the tartar, till a fresh addition occasions no further effervescence; which generally happens before triple the weight of the alkali vegetabile is thrown in: the liquor is then filtered while hot, and either crystallized or evaporated to drynefs. As this falt difficultly crystallizes, inspissation to dryness is the most convenient method; and in this case, to secure the neutralization of the falt, the tartar may be made to prevail at first, and the liquor fuffered to cool a little before filtration, that the redundant tartar may concrete and separate from it; or the neutralization may be more perfectly obtained by means of stained papers, as mentioned at the end of the article Acetum.

+ Ph. Lond. Alcali fixum tartarifatum vulgo tart. folub. † Ph. Edinb. Sal vegetabilis quibufd.

⁽a) Nova Acta Physico-Medica Academiæ Cæsareæ Leopoldino-Carolinæ Naturæ Curiosorum, tom. quart.

Soda tartarifata vulgo fal rupellenfis Ph. Edinb. Sel de Seignette, Rochelle falt. Of the mineral fixt alkali or foda, this acid faturates, according to the faculty of Paris, four fifths its own weight. The neutral falt refulting from its coalition with this alkali, is fomewhat lefs diffoluble than that with the vegetable; and shoots much more easily, into pretty large, hard, multangular crystals, some columnar and flattish, others more irregular. It is milder in taste, and said to be less purgative, requiring to be given to the quantity of an ounce or an ounce and a half to purge effectually: eight drams are reckoned by some to be equivalent, in cathartic power, to six of the soluble tartar.

Tartar forms likewise soluble compounds with all the absorbent earths, and with some metallic bodies, but with these last it is difficultly made to satiate itself completely, the part that is first saturated seeming to impede the action of the rest; for after long boiling, a very considerable

part of the tartar separates on crystallization unchanged.

It is observable, that if any of these combinations of tartar, with alkalies, with earths, or with metals, be dissolved in water, and any other acid added, the pure tartar separates and falls to the bottom, as acid, and as difficult of solution, as at first; the substance, that was combined with it, being absorbed by the acid superadded. As the acids of the vegetable kingdom, whether native or fermented, vinegar, lemon juice, &c. have this effect of disuniting tartar from all the bodies that are combinable with it, equally with those of the mineral kingdom; it sollows, that the tartareous acid is of a kind essentially different from all the other known vegetable ones, and that no acid, unless it be tartar itself, can be joined in prescription to the tartarum solubile, the sel de Seignette, or the combinations of tartar with earths or metals.

* In the Swedish Transactions, Part III. for the year 1770, was published an analysis of creme of tartar by Mr. Scheele. By this it appears, that creme of tartar is not a pure acid, but a compound salt, containing the fixed vegetable alkali united with a superabundance of the tartareous acid. It differs, therefore, from soluble tartar, only in the proportion

of acid it contains.

TELEPHIUM.

TELEPHIUM: a plant with unbranched stalks, clothed with thick sleshy oval leaves, but producing no leaves immediately from the root:

root: the flowers stand in form of umbels on the top of the stalk, and are followed, each, by from three to fix pods full of small seeds: the root is irregular and knobby. It is indigenous in England, and perennial.

- 1. CRASSULA five Telephium Pharm. Edinb. (a) Telephium vulgare C. B. Anacampseros, fabaria, & faba crassa quibusdam. Sedum Telephium Linn. Orpine: with the leaves very slightly or not at all serrated: growing in hedges and moist shady grounds, and producing reddish or whitish pentapetalous flowers in June. The leaves have been supposed to be possessed of an anti-instammatory power; but their virtues appear to be very inconsiderable, as they have no smell, and only an herbaceous mucilaginous taste.
- 2. Rhodiola five rosea: Rhodia radix C. B. Rosewort: with serrated leaves; growing in mountainous places, and producing yellow tetrapetalous flowers in the spring. The root of this species, of little smell when fresh, has when dry a very pleasant one, resembling, when the root is in persection, that of the damask rose: in this odorous matter consists the medical virtue of the rhodiola, and its principal medical difference from the preceding species. Linnæus observes, that when raised in gardens, it has not one hundredth part of the smell or virtue of that which is produced on its native mountains.

TEREBINTHINÆ.

TURPENTINES: the native balfams or refinous juices of certain trees. Four kinds are distinguished by medical writers.

1. TEREBINTHINA CHIA Pharm. Lond. Chio or Cyprus turpentine: generally about the confishence of thick honey, very tenacious, clear and almost transparent, of a white colour with a cast of yellow and frequently of blue, of a warm pungent bitterish taste, and a fragrant smell more agreeable than that of any of the other turpentines. It is the produce of the common terebinth (terebinthus vulgaris C.B. Pistachia terebinthus Linn.), an evergreen bacciferous tree or shrub, growing spon-

taneously in the eastern countries and in some of the southern parts of Europe. The turpentine brought to us is extracted in the islands whose name it bears, by wounding the trunk and branches a little after the buds have come forth: the juice issues thin and clear as water, and by degrees thickens into the consistence in which we meet with it. A like juice, exuding from this tree in the east, inspissated by a slow sire, is said by Kæmpser to be used as a massicatory by the Turkish women, for preserving the teeth, sweetening the breath, and promoting appetite.

- 2. TEREBINTHINA VENETA Pharm. Edinb. Venice turpentine: usually thinner than any of the other forts, of a clear whitish or pale yellowish colour, a hot pungent bitterish disagreeable taste, and a strong smell, without any thing of the fine aromatic flavour of the Chian kind. The true Venice turpentine is said to be obtained from the larch (larix C. B. Pinus larix Linn.), a coniferous tree, with small cones, and short leaves standing in tusts, which fall off in the winter, growing in great abundance on the Alps and Pyreneans, and not uncommon in the English gardens. Though this kind of turpentine bears the name of Venice, it is not the produce of the Venetian territories: it is brought from some parts of Germany, and one greatly resembling it, as is said, from New England. In the shops this turpentine is often supplied by a composition of rosin and the distilled oil of common turpentine.
- 3. TEREBINTHINA ARGENTORATENSIS Pharm. Lond. Strasburg turpentine: generally of a middle consistence between the two foregoing, more transparent and less tenacious than either, in colour yellowish brown, in smell more agreeable than any of the other turpentines, except the Chian, in taste the bitterest yet least acrid. This juice is extracted, in different parts of Germany, from the silver and red fir, by cutting out, successively, narrow strips of the bark, from as high as a man can reach to within two feet of the ground. In some places, a resinous juice is collected from certain knots under the bark: this, called lacryma abiegna and oleum abietinum, is accounted superiour to the turpentine. Neither this turpentine, nor any thing under its name, is at present common in the shops.
- 4. TEREBINTHINA COMMUNIS Pharm. Lond. Common turpentine: about the confistence of honey, of an opake brownish white colour, the coarsest

coarfest, heaviest, in smell and taste the most disagreeable, of all the kinds of turpentine. It is obtained from the wild pine (pinus filvestris C. B. & Linn.), a low coniferous tree, with the leaves longer than those of the firs and issuing two together from one tubercle, growing wild in the different parts of Europe. This tree is extremely refinous, infomuch that, if not evacuated of its juice, it often swells and bursts. The juice, as it issues from the tree, is received in trenches made in the earth, and afterwards freed from its groffer impurities by colature through The cones of the tree appear to contain a refinous wicker baskets. matter, of a more grateful kind than that of the trunk : distilled while fresh, they are said to yield a fine effential oil greatly superiour to that Carpathicum of the turpentines.

oleum Germanis.

All these juices dissolve totally in rectified spirit, but give out little to watery menstrua: they become miscible with water, into a milky liquor, by the mediation of the yolk or white of an egg, and more elegantly by mucilages. Distilled with water, they yield a notable quantity of a subtile penetrating effential oil + vulgarly called spirit; a yellow t or blackish refin remaining in the still: this is the common rofin of the shops. It is fupposed that the officinal Burgundy pitch ||, which is brought from Saxony, is a preparation of the same kind, only less divested of the oil, made by boiling the common turpentine till it acquires a due confistence. The effential oil, rediffilled by itself in a retort, with a very gentle heat, becomes more fubtile, and in this state is called ethereal §; a thick matter remaining behind, called balfam of turpentine ¶. A like balfam is obtained also by distilling, with a stronger fire, the common resin; from which there arises, first a thin yellow oil, and afterwards the thicker dark-reddish balsam, a blackish refin+ remaining in the retort.

+ O1. terebinth. Phar. Lond. & Ed. t Refina flava Ph. Lond. alba Ph. Ed. || Pix Burgundica Ph. Lond & Ed.

§ Ol. terebinth. æthereum P. L. Balf. terebinth. P. L.

+ Refina nigra feu colo-phonia P. L.

All the turpentines are hot stimulating corroborants and detergents. They are given, where inflammatory symptoms do not forbid their use, from half a scruple to half a dram and upwards, for cleaning the urinary passages and internal ulcerations in general, and in laxities of the seminal and uterine vessels. They seem to act in a peculiar manner on the urinary organs, impregnating the water with a violet fmell, even when applied externally, particularly the Venice fort. This last is accounted the most powerful as a diuretic and detergent, and the Chio and Strafburgh as corroborants: they all loofen the belly, the Venice most; and on this account they are supposed by Riverius and others to be less hurtful than fuch irritating diuretics, as are not accompanied with that advantage. Terebinthinate glysters, in obstinate costiveness, are said to be much preferable to saline, as being more certain and durable (a). The common turpentine, as being the most offensive, is rarely given internally: its principal use is in some external applications, among the farriers, and for the distillation of the oil.

The oil is a most potent stimulating detergent diuretic. It is sometimes given, in doses of a few drops, in rheumatisms and fixt pains of the joints; and fome have ventured on much larger quantities. Cheyne recommends, as a perfect cure for sciaticas, though of many years standing, from one to four drams of the ethereal oil, to be taken with thrice its quantity of honey, in a morning fasting, with large draughts of fack whey after it, and an opiate at bed-time: this medicine is to be repeated, with the intermission of a day now and then, if daily repetitions cannot be borne, for four or five days, or eight at furthest (b). It appears, however, highly imprudent, to venture on fuch large doses at once of a medicine fo very hot and stimulating. Boerhaave, after recounting, not without fome exaggeration, its flyptic, anodyne, healing, antiseptic, and discutient virtues when applied hot externally, and its aperient, warming, fudorific and diuretic qualities when taken internally, adds, that it must be used with great caution; that when taken too freely, it affects the head, excites heat and pain therein, and, violently urging a diabetes, brings on a flux of the femen and of the liquor of the proftates; and that in venereal runnings, in which it has by fome been commended, it tends to inflame the parts and increase the disorder.

The balfam and the inspissated refins are used chiefly externally: the balfam is less pungent than the oil, and the refins much less so than the turpentines in substance. The common yellow refin, in taste considerably bitter, is sometimes given as an internal corroborant, in preference to the turpentines themselves, as being divested of the stimulating oil.

TERRA JAPONICA.

TERRA JAPONICA Ph. Lond. Terra japonica dieta Ph. Ed. JAPAN EARTH, improperly so called. *The plant which yields the terra japonica, grows in the East Indies, and is called coira, or caira, by

the natives of Bahar province. It appears to be the same with that mentioned by Cleyerus and Herbert de Jagur, from which the natives of Pegu prepare this extract: they name the tree Kheir or Khadira.

It is a species of the genus mimosa of Linnæus, and called by him mimosa catechu. Its stem grows to about a foot in thickness, and from three to five feet high. It branches out into a thick spreading top, seldom above twelve feet high. The bark is thick and rough. The wood is extremely hard and heavy; its interiour part varies from a pale brown to a dark red, sometimes approaching to black, but always covered with one or two inches thick of white wood. The leaves are doubly pinnated, and have two prickles at the base. From the axillæ of the leaves, arise dense spikes of small flowers, succeeded by pods.

The terra japonica is an extract of the wood of this plant, prepared in the following manner. After the trees are felled, the exteriour white wood is carefully cut off, and the interiour coloured part is cut into chips, with which narrow-mouthed unglazed earthen pots are filled, and water poured upon them till it appears among the upper chips. When this is half evaporated by boiling, the decoction, without straining, is poured into a flat earthen pot, and boiled to one third part; this is set in a cool place for one day, and afterwards evaporated by the heat of the sun, stirring it several times in the day; when it is reduced to a considerable thickness, it is spread upon a mat or cloth which has previously been covered with the ashes of cow-dung; this mass is divided into square or quadrangular pieces by a string, and completely dried by turning them frequently in the sun.

This extract is called by the natives cutt; by the English cutch. In making it, the pale brown wood is preferred, as it produces the fine whitish extract. The darker the wood is, the blacker the extract, and of less value. From the flovenly manner in which the preparation is made, it generally contains a considerable quantity of earth, besides what may be designedly put in it for the purpose of adulteration (a).

This concrete is a mild aftringent, more agreeable in tafte than most of the other substances of that class, being accompanied with a considerable degree of sweetness. It is often suffered to dissolve leisurely in the mouth; both as a topical restringent for laxities and exulcerations of

Trochifci e terra japon. Pharm.Lond.

Electuar. japonic. vulgo confect. japon. Ph. Ed. the gums; and in alvine and uterine fluxes, and catarrhal coughs and hoarseness; medicines of this kind acting in general to much better advantage when thus gradually swallowed, than when taken in full doses at once. With this view the terra japonica is made in the shops into troches; beaten with equal its weight of gum-arabic, and four times the weight of both of sugar of roses, and so much water to be dropt in as will reduce them into a mass of a due consistence for being formed. The Edinburgh college directs a compound electuary, of which terra japonica is the basis, joined with other astringents and aromatics, and a small proportion of opium, which is a very elegant and efficacious medicine of the kind.

Japan-earth dissolves almost totally in water, excepting the impurities; which are usually of the sandy kind, and in considerable quantity, amounting, in the specimens I examined, to about one eighth of the mass. Of the pure matter, rectified spirit dissolves about seven eighths, into a deep red liquor: the part, which it leaves undissolved, is an almost insipid mucilaginous substance. In the shops a solution of it is made in proof spirit, with the addition of cinnamon, a spice the best adapted of any to the intention of this medicine: three ounces of the japan-earth and two of cinnamon are digested in a quart † or two pounds and a half ‡ of the spirit, and the strained tincture given commonly in doses of two or three tea-spoonfuls. It dissolves also in volatile alkaline spirits, in alkaline lye, in the mineral acids, partially and more difficultly in the vegetable acids, and not at all in oils: all the solutions are of a red or purplish colour.

Tinctura japonica † Ph. Lond. ‡ Ph. Edinb.

* By the natives where this extract is made, it is employed medicinally as a cooler in the diseases considered by them as of a hot nature. It is said, when profusely used, to destroy the venereal appetite. It is given at the rate of two ounces a day to tame vicious horses. It is a principal ingredient in one of their ointments of great repute, composed of blue vitriol four drams, Japan-earth four ounces, alum nine drams, white resin four ounces; these are reduced to a fine powder, and mixed by the hand with ten ounces of oil-olive, and water enough to give the mass a proper consistence. This ointment is used in every fore, from a fresh wound to a venereal ulcer; and has been found remarkably serviceable by European practitioners (a).

(a) Kerr's Account, above quoted.

TERREA

TERREA ABSORBENTIA.

ABSORBENT EARTHS: distinguishable from other earthy and stony substances by their solubility in acids. Such are, the mineral calcareous earths, as chalk: the animal calcareous earths, as crabs-claws, oyster-shells, egg-shells, pearl, coral, coralline: animal earths not calcareous, as crabs-eyes and burnt hartshorn. See the respective substances; which have been separately treated of, so far as concerned each in particular; and whose general and common qualities were reserved for this article.

THE obvious and immediate virtue of these bodies is, to obtund acid humours in the first passages, and thus to relieve the cardialgic and other complaints occasioned by them: the relief, however, which they afford, is oftentimes only temporary; from their acting only upon the acid already generated, without correcting the indisposition which tends to produce more. If no acid humours are contained in the first passages, these earthy bodies, not soluble by any other kind of fluid, can have no falutary operation; and, by concreting with the vifcous contents of the stomach into indigestible masses, may prove injurious in a high degree (a).

Absorbents are of more general use in infancy than in adult age; acidities being very familiar to young children, being often in that tender age productive of alarming fymptoms, and having a greater or less share in most of their diseases; whereas, in adults, they are much less frequent, accompanying chiefly hypochondriacal affections, cardialgiæ, and fuch diforders as happen in the first passages from the immoderate use of acid and fermentable food. .

An hypothesis formerly obtained, which ascribed the acute diseases of adults to a morbific acid: against which the absorbent earths were introduced as the most direct alexipharmacs. This theory is now justly exploded; these diseases, instead of being produced, being in general most fuccessfully controlled, by acids. The use of absorbents, in different kinds of fevers, is nevertheless still continued, and sometimes perhaps with advantage: for, though the earths of themselves are apparently rather injurious than beneficial, yet as acids are often given freely at the fame time, the folution of the earth in the acid may prove a medicine more ferviceable in particular cases than the acid unobtunded. It is however, doubtless, more adviseable, to use the earth previously dissolved in the acid, than to give them separately.

The college of Berlin, sensible of the advantage of having the earths, in these cases, previously dissolved, or reduced to a soluble saline form; as well as of the absurdity, retained in other German pharmacopæias, of precipitating them from their solutions by fixt alkaline salts, and thus rendering them wholly inert; directs them to be digested in distilled vinegar, with a gentle heat, till the menstruum ceases to act, and the filtered solution to be inspissated to dryness. This preparation is greatly preferable to the simple imbibition with vinegar or lemon juice recommended by some; as by this last management the earth is made soluble only in part, and in an undeterminable proportion.

Magisterium folubile, coralliorum, perlarum,&c. P. Brandenb.

Solutions of these earths in vegetable acids are in taste somewhat austere. The different earths differ somewhat from one another, both in the degree and in the species of the taste, and probably also in the medical effects, of the solutions: but whether these differences are such, that some of them, as crabs-claws, pearl, coral, and bezoar, are most disposed to promote a diaphoresis in severs, while others, as egg-shells and oyster-shells, act rather by promoting urine, as seems to be generally supposed, has not been determined by fair experience, the earths having rarely been given in a dissolved or in a soluble state. It is most probable that they all act, when dissolved, as mild cooling restringents; for when given in substance, as absorbents, in cases of acidities, they all tend to restrain sluxes of the belly, or to bring on costiveness, an effect to which regard ought to be had in the use of them.

There are two foluble earths, not commonly ranked among the abforbents, whose effects, when combined with acids, are known with more certainty, as they have been used oftener, so combined, than otherwise; to wit, the aluminous earth and magnesia; of which the one is strongly styptic, and the other moderately purgative.

Combinations of the absorbent earths with the nitrous and marine acids are bitterish and of great pungency, particularly those with the marine: the medical effects of these solutions are little known. The vitriolic acid does not dissolve them into a liquid form, but precipitates

them

them from all the others, and is thus combined with them into concretes nearly infipid.

Experiments have been made for determining the comparative strength of different absorbents, or the quantities of acid they are capable of fatiating. Langius reports, that ten grains of crabs-claws destroyed the acidity of forty drops of spirit of falt; that egg-shells, crabs-eyes, and mother of pearl, taken in the same quantity, saturated fifty drops each; red coral, white coral, and fixt alkaline falt, fixty drops each; volatile alkaline falt and pearl, eighty drops each; chalk, an hundred drops; oyfter-shells, an hundred and twenty; and some lime stones no less than an hundred and fixty (a). These experiments however (admitting their accuracy, and the acid to have been equally neutralized in all, which may be reasonably questioned) do not answer the end so perfectly as could be wished; for, to different acids, the earths have different habitudes: from a fet of experiments made by Homberg, it appears that oystershells, for example, require for their solution more of the marine acid than coral does; whereas of the nitrous acid, contrariwife, the coral requires more than the oyster-shells (b). Neither the nitrous nor the marine acids are those which absorbents are destined to satiate in the human stomach, and by which their strength should be examined: the acids of the vegetable kingdom, and the acid of milk, may be prefumed to be the most analogous to such as are generated in the bodies of animals. On trying, with these, the several substances enumerated at the beginning of this article, the differences in their absorbent powers appeared not to be very great: they all faturated pretty nearly the fame quantities of the acids; and there remained, from all, quantities very confiderable, but not very greatly different, of a matter which further additions of the acid would not diffolve.

THEA.

THEA Pharm. Edinb. (c) TEA: the leaf of a Chinese shrub, evonymo affinis arbor orientalis nucifera flore roseo Pluk. alm. Thea bohea & viridis Linn. The leaves, carefully picked, are dried hastily on warm iron plates; whereby they are said to lose in great measure some noxious

⁽a) Vide Langii Opera omnia medica, Lipfiæ 1704, p. 452 & feq.

⁽b) Vide Memoires de l'acad. roy. des sciences de Paris, pour l'ann. 1700.

⁽c) Expunged.

qualities which they have when fresh, and to preserve their admired slavour which by slow exsiccation would be lost. The several sorts of tea brought to us are supposed to be the leaves of the same plant, collected at different times, and cured in a somewhat different manner: Neumann suspects that the brown colour, and the slavour, of the bohea forts, are introduced by art.

BOTH the green and bohea teas have an agreeable smell, and a lightly bitterish subastringent taste: with solution of chalybeate vitriol, they strike an inky blackness. They give out their smell and taste both to watery and to spirituous menstrua: to water, the green sorts communicate their own green tincture, and the bohea their brown: to rectified spirit they both impart a fine deep green. On gently drawing off the menstrua from the filtered tinctures, the water is found to elevate nearly all the peculiar slavour of the tea, while rectified spirit brings over little or nothing, leaving the smell as well as the taste concentrated in the extract: both extracts are very considerably astringent, and not a little ungrateful; the spirituous most so.

Infusions of tea, as dietetic articles, have been extravagantly commended by some and condemned by others; and notwithstanding the frequency of their use, their real effects are scarcely as yet clear. They seem, when moderately used, to be for the most part innocent: in some cases, they seem to be salutary: in some, they are apparently prejudicial. They dilute thick juices and quench thirst more effectually, and pass off by the natural emunctories more freely, than mere watery fluids: they refresh the spirits in heaviness and sleepiness, and seem to counteract the operation of inebriating liquors. From their manifest astringency, they have been supposed to strengthen and brace up the solids, but this effect experience does not countenance: it is in disorders, and in constitutions, wherein corroborants are most serviceable, that the immoderate use of tea is peculiarly hurtful; in cold indolent habits, cachexies, chloroses, dropsies, and debilities of the nervous system.

THLASPI.

THLASPI Pb. Lond. a plant with oblong narrow undivided leaves joined immediately to the stalks, on the tops of which grow numerous tetrapetalous

tetrapetalous flowers, each of which is followed by a short flat seed-vessel divided transversely into two cells.

- 1. This pi arvense siliquis latis C. B. This pi arvense Linn. Treacle-mustard: with roundish-pointed jagged leaves, and broad capsules containing about four seeds in each cell. It is annual, and grows wild in corn-fields.
- 2. Thlaspi arvense, vaccariæ incano solio majus C. B. Thlaspi campestre Linn. Mithridate mustard: with hoary sharp-pointed leaves shaped like an arrow-head; and only one seed in each cell. It is biennial, and grows in fields and open clayie grounds.

The feeds of these plants have an acrid biting taste, approaching to that of the common mustard; with which they agree nearly in their pharmaceutic properties, their pungent matter being totally extracted by water, only partially by rectified spirit, and being elevated by water in distillation. They have, joined to their acrimony, an unpleasant slavour, somewhat of the garlick or onion kind; and this they give out to spirituous as well as watery menstrua. They are rarely made use of any otherwise than as ingredients in the compositions whose names they bear: though some recommend them in different disorders, preferably to the common mustard.

THUS.

THUS Ph. Lond. Frankincense: a solid brittle resin, brought to us in little glebes or masses, of a brownish or yellowish colour on the outside, internally whitish or variegated with whitish specks. It is supposed to be the produce of the pine that yields the common turpentine, and to concrete upon the surface of the terebinthinate juice soon after it has issued from the tree.

This refin has a bitterish acrid unpleasant taste, and no considerable smell: it dissolves totally in rectified spirit, but is scarcely acted upon by watery menstrua. It may be looked upon as a mild corroborant; though at present it is little otherwise made use of than as an ingredient in theriaca, and externally in plasters.

THYMELEA.

THYMELÆA Pharm. Parif. A shrubby plant; with smooth uncut leaves; and monopetalous flowers set thick together: each flower is cut into sour acute sections, and followed by an oblong, red, yellow, or black berry, containing one seed, which resembles a hemp-seed.

- 1. THYMELÆA: Thymelæa foliis lini C.B. Daphne Gnidium Linn. Spurge-flax: with the stalks and branches clothed with evergreen leaves like those of flax; and white flowers in clusters on the tops.
- 2. LAUREOLA seu Chamælæa: Laureola sempervirens flore viridi, quibus dam laureola mas C. B. Daphne laureola Linn. Spurge laurel: with evergreen shining bay-like leaves, standing several together, only at the tops of the branches; and greenish flowers on pedicles in their bosoms.
- 3. MEZEREUM: Mezereon Pharm. Edinb. Laureola folio deciduo, flore purpureo, officinis laureola femina C. B. Daphne Mezereum Linn. Spurge-olive, widow-wail: with pale purplish or white flowers clothing the branches; on the tops of which appear, after the flowers have fallen, bay-shaped leaves not shining.

The first of these plants grows on mountainous places in the southern parts of Europe: the second in moist woods in some parts of England: the third, a native of Germany (a), is cultivated in our gardens, on account of the elegance and earliness of its slowers, which sometimes appear in the end of January: the berries of all the sorts ripen in August or September.

The leaves of these plants have little or no smell, but a nauseous, acrid, very durable taste: taken internally, in small doses, as ten or twelve grains, they are said to operate with violence, by stool, and sometimes by vomit, so as not to be ventured on with safety unless their virulence be previously abated by long boiling, and even then they are much too precarious to be trusted to.

⁽a) The mezereon has of late been observed to be a native of England also, being found plentifully in some woods near Andover in Hampshire.

The flowers are of a different nature, being in taste little other than mucilaginous and sweetish, and of a light pleasant smell.

The pulpy part of the berries appears also to be harmless; but the feeds, called coccognidia or grana cnidia, are as acrid, and as virulently

purgative as the leaves.

* The bark of the spurge laurel, macerated in water, has of late been much employed in France as a topical application to the skin, for the purpose of excoriating and exciting a discharge. That of the mezereon

has been recommended for the same purpose.

The root of the mezereon has lately been used with success in cases of venereal nodes. Dr. Ruffel, to whom the public is obliged for the communication of its efficacy in this frequently obstinate complaint, observes, in the medical inquiries above mentioned, that the cortical part of the root, on first chewing, is not pungent, but after a little time proves greatly fo; and that the difagreeable stimulus in the fauces lasts for many hours: that a decoction of an ounce of the fresh cortical part in a gallon and a half of water (the boiling being continued till half a gallon is wasted, and an ounce of sliced liquorice added towards the end) may be taken to the quantity of half a pint four times a day, is not nauseous to the taste, and has not been found to disagree with any stomach or constitution, or to remarkably increase any of the secretions; but that on doubling the quantity of the mezereon, the decoction proved fo pungent, that no stomach could bear it. He recommends the above decoction principally in those venereal nodes that proceed from a thickening of the membrane of the bones, which appears to be the cause of greatest part of these tumours, at least when recent: when there is an exostosis, nothing is to be hoped for from this medicine; and when the bone is carious, no cure is to be expected without an exfoliation, though even here it sometimes disperses the tumour, as appears from some of the cases which he relates. In a thickening of the periosteum from other causes, it has likewise had good effects.

THYMIAMA.

THYMIAMATIS cortex Officinarum Germaniæ: Thus judæorum Quorundam. A bark, in small brownish-grey pieces, intermixed with bits of leaves, seeming as if the bark and leaves had been bruised and pressed

pressed together; brought from Syria, Cilicia, &c. and supposed to be the produce of the liquid-storax tree.

This bark has an agreeable balfamic smell, approaching to that of liquid-storax, and a subacrid bitterish taste accompanied with some slight aftringency. Insusions of it in water are of an orange colour, in taste and smell ungratefully balfamic: inspissated, they leave a dark reddish brown extract, retaining some of the smell of the bark, in taste austere, slightly bitter, and of a mild aromatic acrimony. To rectified spirit it communicates a dark colour like that of a solution of balfam of Peru: the spirit, distilled off from this tincture, is highly fragrant, insomuch that a dram communicates an agreeable odour to some quarts of water: the remaining extract is likewise of a pleasant smell, and amounts to at least one eighth of the weight of the bark.

This bark, said to be common in the German shops, is in this country very rarely to be met with. Cartheuser and Hossman, from whom the above account is extracted, report, that it affords an excellent sumigation for ædemas, rheumatisms, and catarrhs; and that the spirituous tincture and extract, and the distilled spirit, are useful anodynes or antispasmodics in convulsive coughs and other disorders.

THYMUS.

THYME: a low shrubby plant; consisting of numerous slender tough stalks, with little roundish leaves in pairs, and loose spikes, on the tops, of purplish or whitish labiated flowers, whose upper lip is nipt at the extremity, the lower divided into three nearly equal segments.

I. THYMUS Pharm. Edinb. Thymum vulgare folio tenuiore C. B. Thymus vulgaris Linn. Common thyme: with upright stalks, and dark brownish green somewhat pointed leaves; a native of the southern parts of Europe, common in our gardens, and slowering in June and July.

This herb is a moderately warm pungent aromatic. To water it imparts, by infusion, its agreeable smell, but only a weak taste, with a yellowish or brown colour: in distillation, it gives over an essential oil, in quantity about an ounce from thirty pounds of the herb in slower, of a gold yellow colour if distilled by a gentle fire, of a deep brownish red

if by a strong one, of a penetrating smell resembling that of the thyme itself, but less grateful, in taste excessively hot and siery: the remaining decoction, inspissated, leaves a bitterish, roughish, subsaline extract. The active matter, which by water is only partially dissolved, is by rectified spirit dissolved completely; though the tincture, in colour blackish-green, discovers less of the smell of the thyme than the watery insusion: the spirit brings over in distillation a part of its slavour, leaving an extract of a weak smell and of a penetrating camphorated pungency.

- 2. SERPYLLUM Pb. Ed. Serpyllum vulgare minus C. B. Thymus Serpyllum Linn. Mother-of-thyme: with trailing stalks, and obtuse leaves; growing wild on heaths and dry pasture grounds. This also is an elegant aromatic plant, similar to the foregoing species, but milder, and in flavour rather more grateful. Its essential oil is both in smaller quantity and less acrid, and its spirituous extract comes greatly short of the penetrating warmth and pungency of that of the other. It is said to afford an agreeable distilled water, more durable, but less active and penetrating than pepper-mint (a). Both the leaves themselves, and their spirituous tincture, are of a bright green colour, without any thing of the brown or blackish hue of those of common thyme.
- 3. Thymus citratus Pb. Lond. Serpyllum foliis citri odore C. B. Lemon-thyme: in appearance differing little from the second sort, of which Linnæus makes it a variety, except that it is more upright and more bushy; a native of dry mountainous places, common in gardens, and flowering as the others in July. This species is less pungent than the first sort, more so than the second, and much more grateful than either: its smell in particular is remarkably different, approaching to that of lemons. Distilled with water, it yields a larger quantity than the other sorts, of a yellowish very fragrant oil of the lemon flavour, containing nearly all the medicinal parts of the plant, for the remaining decoction is almost insipid as well as inodorous. It gives over also with rectified spirit its finer odorous matter; a less agreeable flavour, and a moderate warmth, remaining in the spirituous extract.

TILIA.

TILIA Pharm. Lond. Tilia femina folio majore C.B. Tilia europæa Linn. Lime or Linden: a tall spreading-branched tree, with large heart-shaped, serrated, soft, somewhat hairy leaves: in the bosoms of these rise long narrow leafy productions, from the middle rib of which issue one or three pedicles bearing three slowers apiece, or one pedicle bearing nine: the flower is whitish, pentapetalous, and sollowed by a kind of dry berry about the size of a filberd. It is a native of England, flowers in July, and begins to lose its leaves in August.

THE flowers of the lime-tree are supposed to have an anodyne and antispassmodic virtue: Hossman seems to entertain a great opinion of them in these intentions, and as his theory deduces most diseases from spassms and spassmodic strictures, they are accordingly very frequent in his prescriptions: he says he knew a chronical epilepsy cured by the use of an insussion of them drank as tea. The fresh flowers have a moderately strong smell, in which their virtue (whatever it may be) seems to consist, and which in keeping is soon dissipated: when divested of this odorous principle, they discover to the taste only a strong mucilage, from which may be extracted, by rectified spirit, a slightly bitterish subastringent matter.

TITHYMALUS.

SPURGE: a plant with small smooth leaves, round stalks full of a milky juice, and umbel-like clusters of tetrapetalous flowers, whose cups are divided into sour segments set alternately with the petala: the flower is followed by a roundish or three-square capsule containing three seeds.

1. TITHYMALUS PARALIOS: Tithymalus maritimus C. B. Euphorbia Paralias Linn. Sea spurge: with oblong narrow flax-like leaves, broadest in the middle, clothing the stalks, and lying over one another in an upward direction, like scales; and two roundish, heart-shaped, or kidney-shaped leaves encompassing each of the subdivisions of the umbel: found wild on sandy shores, and slowering in June. All the parts of this

this plant are extremely acrid irritating cathartics; apt to inflame the mouth, fauces, and stomach; operating with so great violence, that though some may perhaps have borne their operation without much injury to the constitution, yet common prudence forbids their being ever ventured on. Several correctors have been employed for them, but none with commendable success: maceration of the middle bark of the root in vinegar, directed by the faculty of Paris, renders it indeed less virulent, but of precarious operation: digestion of the milky juice with alkaline salts, recommended by others, leaves it still too acrid. For alleviating inflammatory symptoms produced by imprudently swallowing or tasting these acrid substances, milk, plentifully drank, seems the most effectual remedy. Gerard relates, that on taking but one drop of the milk of the sea spurge, it did so swell and inflame in his throat, that he hardly escaped with his life, and that on drinking milk, the extremity of the heat ceased.

2. TITHYMALUS CYPARISSIUS C. B. Euphorbia Cyparistias Linn. Cypress spurge: with numerous oblong slender leaves, not wider in the middle than at the ends; the umbel divided into numerous ramifications, each of which is divided and subdivided into two; the divisions perforating as it were the two roundish leaves which encompass them; a native of Germany, Switzerland, and some other parts of Europe. This species, though allowed by the faculty of Paris to be used indiscriminately with the preceding, is in all its parts less acrimonious. Poterius says he has found half a dram or a dram of the powdered root to act as a mild cathartic; and that the juice obtained from the bruised herb and root, depurated and exsiccated in the sun, is of the same operation with scammony (a).

SEVERAL other spurges are enumerated in catalogues of the materia medica, under the names of efula, pityufa, cataputia, lathyrus, alypum, peplus, apios, &c. among which there does not appear to be any one more virulent than the first above described, or less virulent than the second. None of them are among us ventured on for any internal use: the milky juice of the wild spurges is sometimes applied externally by the common people for consuming warts.

4 R

TORMENTILLA.

TORMENTILLA Pharm. Lond. & Edinb. Tormentilla filvestris C. B. Heptaphyllum. Tormentilla erecta Linn. Tormentilla or Septeoil: a plant with slender, weak, upright stalks; oblong leaves, indented towards the extremity, and converging from the indented part to their juncture with the stalk, standing generally seven at a joint; and small yellow tetrapetalous flowers on the tops of the branches, sollowed by naked seeds: the root is generally crooked and knotty, of a dark brown or blackish colour on the outside, and reddish within. It is perennial, grows wild in woods and on commons, and slowers in June.

Tormentil root is a strong and almost flavourless astringent, and gives out its astringency both to water and rectified spirit, most perfectly to the latter: the watery decoction, of a transparent brownish red colour whilst hot, becomes turbid in cooling like that of the Peruvian bark, and deposites a portion of resinous matter: the spirituous tincture, of a brighter reddish colour, retains its pellucidity. The extracts, obtained by inspissation, are intensely styptic, the spirituous most so. It is generally given in decoction: an ounce and a half of the powdered root may be boiled in three pints of water to a quart, adding, towards the end of the boiling, a dram of cinnamon: of the strained liquor, sweetened with an ounce of any agreeable syrup, two ounces or more may be taken four or five times a day.

TRICHOMANES.

TRICHOMANES Pharm. Lond. Polytrichum sive Trichomanes Pharm. Edinb. & C. B. Callitrichum. Asplenium Trichomanes Linn. English maidenhair: a small plant, without stalks: the leaves are long, narrow, composed of little roundish dark-green segments set in pairs along a shining black rib: the seeds are a fine dust lying on the backs of the leaves. It is perennial, and grows wild on shady grounds and old walls.

This herb has a mucilaginous somewhat sweetish and roughish taste, and little or no particular slavour. It is accounted serviceable in disorders

orders of the breast, particularly in tickling coughs and hoarseness from thin acrid defluxions, and in these intentions has been long substituted among us to the adianthum, from which it appears to be very little, if at all, different in quality. It is usually directed in infusion or decoction, with the addition of a little liquorice: a pectoral fyrup is prepared in the Syrup. pec-shops, from an infusion of five ounces of the dry leaves and four of toralis P. L. liquorice root in five pints of boiling water.

TRIFOLIUM PALUDOSUM.

TRIFOLIUM PALUDOSUM Ph. Lond. Menyanthes Ph. Ed. Trifolium palustre C. B. Menyanthes trifoliata Linn. BUCKBEAN: a plant with large oval leaves, pointed at each end like those of the garden bean, fet three together on long pedicles, which embrace the stalk to fome height, and there parting leave it naked to near the top, where issues a short spike of pretty large reddish white monopetalous flowers, each of which is cut into five fegments, hairy on the infide, and followed by an oval feed-veffel. It is perennial, grows wild in marshy places, and flowers in May.

THE leaves of buckbean have a bitter penetrating taste, which they impart both to watery and spirituous menstrua; without any remarkable fmell or flavour. They have of late years come into common use, as an alterative and aperient, in impurities of the humours, and some hydropic and rheumatic cases. They are usually taken in the form of infusion, with the addition of some of the acrid antiscorbutic herbs, which in most cases improve their virtue, and of orange-peel or some other grateful aromatic to alleviate their ill tafte: they are fometimes, among the common people, fermented with malt liquors, for an antifcorbutic dietdrink. Their fenfible operation is by promoting urine and fomewhat loofening the belly.

TURPETHUM.

TURPETHUM five Turbith Ph. Ed. (a) TURBITH: the cortical part of the root of a species of convolvulus (Convolvulus Turpethum

(a) Expunged.

4 R 2

Linn.),

Linn.), brought from the East Indies, in oblong pieces, of a brown or ash colour on the outside and whitish within: the best is ponderous, not wrinkled, easy to break, and discovers to the eye a large quantity of resinous matter.

This root, on the organs of taste, makes at first an impression of sweetness; but when chewed for some time, betrays a nauseous acrimony. It is accounted a moderately strong cathartic, but does not appear to be of the safest or most certain kind; the resinous matter, in which its virtue resides, being very unequally distributed; insomuch that, as is said, some pieces, taken from a scruple to a dram, purge violently, whilst others, in larger doses, have very little effect.

TUSSILAGO.

TUSSILAGO sive Farfara Pharm. Edinb. Tussilago vulgaris C. B. Bechium & ungula caballina quibusdam. Tussilago Farfara Linn. Coltsfoot: a low plant, producing early in the spring single stalks, each of which bears a yellow sloculous flower followed by several seeds winged with down: the leaves, which succeed the slowers, are short, broad, somewhat angular, slightly indented, green above, and hoary underneath. It is perennial, and grows wild in moist grounds.

The leaves and flowers of coltsfoot, in taste somewhat mucilaginous, bitterish, and roughish, and of no remarkable smell, are ranked among the principal pectoral herbs. Insusions of them, with a little liquorice or with the other herbs of similar intention, are drank as tea, and sometimes with considerable benefit, in catarrhous disorders and coughs threatening consumptions. They have been found serviceable in hectics and colliquative diarrhous (a).

TUTIA.

TUTIA Pharm. Lond. & Edinb. Tutia alexandrina. TUTTY: an argillaceous ore of zinc, found in Persia; formed on cylindrical moulds into tubulous pieces like the bark of a tree, and baked to a moderate

MATERIA MEDICA.

hardness (a); generally of a brownish colour and full of small protuberances on the outside, smooth and yellowish within, sometimes whitish, and sometimes with a blueish cast. Like other argillaceous bodies, it becomes harder in a strong sire; and after the zinc has been revived and dissipated by inflammable additions, or extracted by acids, the remaining earthy matter affords, with oil of vitriol, an aluminous salt (see Bolus and Calaminaris).

TUTTY, levigated into an impalpable powder, is, like the lapis calaminaris and calces of zinc, an useful ophthalmic, and frequently used as such in ointments and collyria. Ointments for this intention are prepared in the shops, by mixing the levigated tutty with so much vipers fat as is sufficient to reduce it to a due consistence; or by adding one part to sive parts of a simple liniment made of oil and wax;

Tutia præparata Phar. Lon. & Edin.

Ung. tutiæ † Pb. Lond. † Pb. Edinb.

VALERIANA.

WALERIANA filvestris Ph. Lond. & Edinb. Valeriana silvestris major montana C. B. Valeriana officinalis Linn. WILD VALERIAN: a plant with channelled stalks; the leaves in pairs; each leaf composed of a number of long narrow sharp-pointed segments, indented about the edges, of a dull green colour, set along a middle rib, which is terminated by an odd one; producing, on the tops of the stalks, umbel-like clusters of small monopetalous slowers, each of which is divided into five segments, set in a very little cup, and followed by a single naked seed winged with down: the root consists of tough strings with numerous smaller threads, matted together, issuing from one head, of a dusky brownish colour approaching to olive. It is perennial, and grows wild in dry mountainous places.

Another species, or variety, of wild valerian, is met with in moist watery grounds, distinguishable by the leaves being broader and of a deep

(a) The above account of the origin of tutty is supported by the authority of Teixeira and Douglas, and by its chemical properties. That the common opinion, of its being a sublimate produced in the European sounderies where zinc is melted with other metals, is erroneous, appears from hence; that tutty is not found, upon strict enquiry, to be known at those sounderies; and by its confisting in great part of an earth not capable of rising in sublimation. Thus much, however, is probable, that sublimates or the common ores of zinc are often mixed with argillaceous earths and baked hard, in imitation of the genuine oriental tutty.

glossy green colour. Both forts have been used indiscriminately; but the mountain sort is by far the most efficacious, and is therefore expressly ordered for the officinal species by the London college.

THE mountain valerian root has a strong not agreeable smell, and an unpleafant warm bitterish subacrid taste: the strength of the smell and tafte is the only mark to be depended on of its genuineness and goodness. It is a medicine of great esteem in the present practice against obstinate hemicrania, hysterical, and the different kinds of nervous disorders, and is commonly looked upon as one of the principal antispassmodics. Columna reports, that he was cured by it of an inveterate epilepfy after many other medicines had been used in vain: on more extensive trials it has been found, in some epileptic cases, to effect a cure, in several to abate the violence or frequency of the fits, and in many to prove entirely ineffectual: oftentimes, it either purges, or operates by fweat or by urine, or brings away worms, before it prevents a fit. The dose of the root in powder is from a scruple to a dram or two, which may be repeated, if the stomach will bear it, two or three times a day. * A remarkable instance of its efficacy in a catalepsy is given by Mr. Mudge(a): doses of half an ounce of the powder were exhibited twice a day, and a less quantity was found ineffectual.

The powdered root, infused in water or digested in rectified spirit, impregnates both menstrua strongly with its smell and taste, and tinges the former of a dark brown, the latter of a brownish red colour. Water distilled from it smells considerably of the root, but no essential oil separates, though several pounds be submitted to the operation at once: the extract obtained by inspissating the watery insusion, has a pretty strong taste, disagreeably sweetish and somewhat bitterish: the spirituous extract is less disagreeable, and more perfectly resembles the root itself: the quantity of watery extract is about one sourth the weight of the root; of the spirituous, about one eighth. Tinctures of it are prepared in the shops, by digesting sour ounces of the powdered valerian in a quart of proof spirit; in the same quantity of the volatile aromatic spirit; or of the dulcified spirit of sal ammoniac. The root in substance, however, is generally sound to be more effectual than any preparation

Tinct. valer. † fimp. P. L. ‡ volat. P. L. || Pb. Edinb. of it. Among the materials I have made trial of for covering its flavour, mace feemed to answer the best.

VANILLA.

VANILLA seu Banilia Pharm. Paris. Aracus aromaticus. VA-NELLOE: the fruit of a climbing plant (volubilis siliquosa mexicana foliis plantaginis Raii bist. Epidendrum Vanilla Linn.) growing in the Spanish West Indies. It is a long flattish pod, containing, under a wrinkled brittle shell, a reddish brown pulp, with small shining black seeds.

Vanelloes have an unctuous aromatic taste, and a fragrant smell like that of some of the finer balsams heightened with musk. They are used chiefly in perfumes; scarcely ever, among us at least, in any medical intention; though they should seem to deserve a place among the principal medicines of the nervous class. By distillation, they impregnate water strongly with their fragrance, but give over little or nothing with pure spirit: by digestion, spirit totally extracts their smell and taste, and in great measure covers or suppresses the smell.

VERBASCUM.

TAPSUS BARBATUS five Verbascum Pharm. Edinb. Verbascum mas latifolium luteum C. B. Candelaria & lanaria quibusdam. Verbascum Thapsus Linn. Mullein: a large plant, all over white and woolly; with a single woody stalk, clothed with oblong oval leaves joined to it without pedicles, bearing on the top a long spike of large yellow monopetalous slowers cut into sive segments, and followed by conical seed-vessels. It is biennial, grows wild by road-sides, and slowers in July.

THE leaves of mullein, recommended as mild aftringents, have a roughish drying kind of taste, and very little smell. The flowers have also a slight roughishness, with a considerable sweetness. A decoction of the leaves has lately been used with some success in diarrheas (a).

(a) Home's Clin. Caf. and Exp.

VERBENA.

VERBENA Pharm. Edinb.(a) Verbena communis flore cæruleo C. B. Hierobotane, herba facra, herba cephalalgica & peristerium quibustam. Verbena officinalis Linn. Vervain: a plant with wrinkled, oblong, obtuse leaves, deeply jagged and indented, set in pairs on the stalks, the upper ones divided into three segments: on the tops of the branches appear irregularly labiated blue flowers, in long spikes, without any leaves among them, followed each by four seeds inclosed in the cup. It is annual, grows wild in uncultivated places, and slowers in July or August.

This herb has been celebrated for abundance of virtues, for which its fensible qualities afford little or no foundation. It has no remarkable smell, and hardly any taste.

VERONICA.

VERONICA: a low, fomewhat hairy, trailing plant, with firm leaves fet in pairs: from the joints arise slender pedicles, bearing spikes of blue monopetalous flowers, each of which is divided, as is the cup, into four segments, and followed by a flat bicellular capsule, which opens at the upper broad part and sheds small brown seeds.

1. VERONICA MAS five Betonica pauli Ph. Edinb. (a) Veronica mas fupina & vulgatissima C. B. Thea Germanica quibusdam. Veronica officinalis Linn. Male speedwell: with crenated leaves of a roundish oval figure; those on the flowering twigs, long, narrow, and not crenated. It is perennial, and grows wild on sandy grounds and dry commons.

The leaves of veronica have a weak not disagreeable smell, which in drying is dissipated, and which they give over in distillation with water, but without yielding any separable oil. To the taste they are bitterish and roughish: an extract made from them by rectified spirit is moderately bitter and subastringent: the watery extract is weaker, though the quantity of both is nearly the same; whence spirit seems to extract their

virtue more completely than water. This herb is of great efteem among the Germans; in diforders of the breast both catarrhous and ulcerous, and for purifying the blood and humours: infusions of the leaves, which are not unpalatable, are drank as tea, and are found to operate fenfibly by urine.

2. TEUCRIUM Act. med. berolinens. Chamædrys spuria major angustifolia C. B. Veronica Teucrium Linn. Mountain speedwell: with sharply serrated leaves of a long oval figure; the lower embracing the stalk by a broad basis. It is a native of Germany.

The leaves and flowers of this species have been greatly commended for dietetic infusions; and said to promote perspiration and urine, to be in general falubrious, and medicinal in feveral diforders (a). Cartheufer observes, that they impart to boiling water a greenish colour, a pleasant balfamic fmell, and a much more agreeable tafte than the preceding veronica. Among us they have not yet been introduced, nor is the plant common: what has usually been called teucrium is a plant of another genus, a large species of germander.

VINCETOXICUM.

VINCETOXICUM, Asclepias, Hirundinaria, Pharm. Edinb. (b) Asclepias albo flore C. B. Asclepias Vincetoxicum Linn. Swallow-wort, tame-poifon: a plant with unbranched stalks; smooth oblong acuminated leaves fet in pairs, and clusters of white monopetalous flowers, each of which is divided into five fections, and followed by two long pods full of a white cottony matter with small brownish feeds: the root is large, composed of a great number of slender strings hanging from a transverse head, externally brownish, internally white. It is perennial, grows wild in gravelly grounds in fome parts of England, and flowers in July.

THE root of vincetoxicum has, when fresh, a moderately strong not agreeable fmell, approaching to that of wild valerian, which in drying is in great part diffipated; chewed, it impresses first a considerable sweetness, which is soon succeeded by an unpleasant subacrid bitterishness: an extract made from it by water, is moderately sweetish, balsamic, and bitterish; the spirituous extract is stronger in taste, proportionably smaller in quantity, and retains a part of the specific slavour of the root. It is recommended as resolvent, sudorisic, and diuretic; in catarrhal, cachectic, and scrophulous disorders, and in uterine obstructions; in doses of from a scruple to a dram or more in substance, and three or four drams in insusion. It has been employed by some of the Germans as an alexipharmac, and hence received the name of contrayerva Germanorum. Some have however suspected it to possess noxious qualities, and observe that when fresh it excites vomiting. Among us it is scarcely ever made use of in any intention.

VINUM.

WINE: the fermented juice of the grape. It differs in colour, flavour, and strength, partly from differences in the grape itself, but chiefly from different managements or additions. Five forts are employed in the shops as menstrua for medicinal substances: Vinum album, Mountain: Vinum album gallicum, French white wine: Vinum canarinum, Canary or sack: Vinum rbenanum, Rhenish: Vinum rubrum, Red Port.

ALL wines confift of an inflammable spirit, and water, separable by distillation; an unctuous viscid substance, which abounds particularly in the sweet wines, as Canary, and impedes their dissolving power; and an acid, obvious in some to the taste, as in Rhenish, which hence becomes an useful menstruum for some bodies of the metallic kind, particularly iron and the antimonial regulus. In distillation, after the inflammable spirit has arisen, they all yield more or less of a peculiar grateful acid; a grosser tartareous acid remaining in the still, along with the unctuous and mucilaginous matter. In long keeping, a part of the tartar is thrown off from the wine, and incrustates the sides of the cask.

Media fubflantia vini Beccheri.

Wine, confidered as a medicine, is a valuable cordial in languors and debilities; more grateful and reviving than the common aromatic infufions and distilled waters, particularly useful in the low stage of malignant or other fevers, for raising the pulse and supporting the vis vitæ, promoting a diaphoresis, and resisting putrefaction. Dietetically, its moderate

moderate use is of service to the aged, the weak, and the relaxed, and to those who are exposed to warm and moist, or to corrupted air: in the opposite circumstances, it is less proper, or prejudicial. Externally, it is used as a corroborant, antiseptic, and antiphlogistic somentation.

The acid obtained from wine by distillation, apparently of a different nature from the acetous as well as from the native vegetable acids, seems to deserve some regard, both as a medicine, and as a more elegant menstruum, for iron and some other bodies, than the common acids.

With regard to the medical differences of wines, it may be observed, that the effects of the full-bodied are much more durable than those of the thinner: that all sweet wines are in some degree nutritious; the others not at all, or only accidentally so, by promoting appetite and strengthening the organs concerned in digestion: that sweet wines in general do not pass freely by urine, and that they heat the constitution more than an equal quantity of any other, though containing sull as much spirit: that those which are manifestly acid pass freely by the kidneys, and gently loosen the belly; and that most of the red ones are subastringent, and tend to restrain immoderate excretions.

VIOLA.

VIOLA Pharm. Lond. & Edinb. Viola martia purpurea flore simplici odoro C. B. Violaria. Viola odorata Linn. VIOLET: a low creeping plant, without any other stalk than the pedicles of the leaves and flowers: the leaves are roundish, somewhat heart-shaped, obtusely crenated about the edges: the flower consists of five irregular petala, of the deep purplish blue called, from the name of the plant, violet colour: the fruit is a little capsule divided into three cells, full of small roundish seeds. It is perennial, grows wild in hedges and shady places, and flowers in March.

The flowers of a different species, greatly inferiour to the above, are frequently substituted in our markets. This fort may be readily distinguished; the herb, by its having stalks, which trail on the ground, and bear both leaves and flowers, and by the young leaves being hairy; the flower, by the three lower petala being spotted with white, and by their want of smell.

Syr. violar. † Ph. Lond. † Ph. Edinb.

THE officinal violet flowers have a very agreeable fmell, and a weak mucilaginous bitterish taste. Taken to the quantity of a dram or two, they are faid to be gently laxative or purgative; and the feeds, which have more taste than the flowers, to be more purgative, and sometimes emetic. The flowers give out to water both their virtue and their fine colour, but fearcely impart any tincture to rectified spirit, though they impregnate the spirit with their fine flavour, and probably also with their purgative quality. An infusion of two pounds of the fresh flowers in five + or eight + pints of boiling water, passed through a fine linen cloth without pressure, is made in the shops into a syrup, which proves an agreeable laxative for children. Both the flowers themselves and the fyrup lose their colour in being long kept: acids change them instantly into a red; alkalies, and fundry combinations of acids with earthy and metallic bodies, to a green: perfect neutral falts, or those compounded of an acid and alkali, make no alteration. Some have been accustomed to communicate to fyrups a violet colour with materials of greater durability than the violet itself, or than any other blue flower: these sophisticated preparations may be distinguished by their colour withstanding alkalies and acids, or being affected by them in a different manner.

* VIOLA TRICOLOR Linn. Pansies, or heartsease: this well known plant has lately been recommended by a German physician, Dr. Strack, as a specific in the crusta lastea of children. He directs a handful of the fresh, or half a dram of the dried leaves to be boiled in half a pint of milk, which is to be strained for use. This dose is repeated morning and evening. He observes, that when it has been administered eight days, the eruption usually increases considerably, and the patient's urine acquires a smell like that of cats. When the medicine has been taken a fortnight, the scurf begins to fall off in large scales, leaving the skin clean. The remedy is to be persisted in, till the skin has resumed its natural appearance, and the urine ceases to have any particular smell.

VIPERA.

VIPERA Pharm. Lond. & Edinb. Coluber Berus Linn. THE VIPER or ADDER, a viviparous reptile, about an inch or less in thickness, and twenty or thirty in length, with a small sharp-pointed tail. It is found

in the heat of fummer, under hedges in unfrequented places; and in winter retires into holes in the earth.

The poison of this serpent is confined to its mouth. At the basis of the phangs, or long teeth which it wounds with, is lodged a little bag containing the poisonous liquid; a very minute portion of which, if mixed immediately with the blood, proves fatal; though it does not appear to be pernicious when fwallowed, provided there is no folution of continuity in the parts which it comes in contact with (a). Our viper-catchers are faid to prevent the mischiefs otherwise following from

the bite, by rubbing oil-olive warm upon the part.

The flesh of the viper is perfectly innocent, and has been greatly commended as a medicine in fundry diforders. It appears to be very nutritious, and hence an useful restorative in some kinds of weaknesses and emaciated habits: but in fcrophulous, leprous, and other like distempers, the good effects, which have been ascribed to it, are more uncertain: I have known a viper taken every day for above a month, in diforders of the leprous kind, without any apparent benefit. The form in which they are used to best advantage, is that of broth, of which an elegant preparation is directed by the London college+; that the + Jus viperiwines (made commonly by macerating for a week, with a gentle heat, two ounces of the dried flesh in three pints of mountain 1) have any great 1 Vinum vivirtue, cannot perhaps be affirmed from fair experience.

The fat of the viper is accounted particularly useful in disorders of the eyes; but what advantages it has above other foft fats, is by no means clear: see Pinguedo. It was formerly supposed to have some specific power of refisting the poison of the viper's bite, by being rubbed immediately on the part; but experience has now shewn that common

oil is in this intention of equal efficacy.

VIRGA AUREA.

VIRGA AUREA Pharm. Edinb. (b) Virga aurea angustifolia minus ferrata C. B. Herba doria & Consolida saracenica quibusdam. Solidago Virga aurea Linn. GOLDEN ROD: a plant with long fomewhat oval leaves, pointed at both ends, flightly or not at all indented; and upright

perinumP.L.

spikes, along the stalks, of small yellow flowers, composed of several flosculi set in scaly cups, followed by small seeds winged with down. It is perennial, grows wild in woods and on heaths, and flowers in August.

THE leaves and flowers of golden rod are recommended as corroborants and aperients; in urinary obstructions, nephritic cases, ulcerations of the bladder, cachexies, and beginning dropsies. Their sensible qualities promise considerable medical activity: their taste, which they readily impart both to water and rectified spirit, and which remains entire in the inspissated extracts, is of a subtile penetrating durable kind, not very ungrateful, weak in the herb in substance, strong in the watery extract, and stronger in the spirituous.

VISCUS.

VISCUS QUERNUS Ph. Edinb. (a) Viscum baccis albis C. B. Viscum album Linn. Misseltoe: a bushy evergreen plant, with woody branches variously interwoven; firm narrow leaves, narrowest at the bottom, set in pairs; and imperfect white flowers in their bosoms, followed each by a transparent white berry containing a single seed. It grows only on the trunks and branches of trees, and may be propagated by rubbing the glutinous berries on the bark that the seeds may adhere.

The leaves and branches of miffeltoe, formerly recommended as fpecifics in convultive and other nervous diforders, and now fallen into general neglect, do not appear to have any confiderable medicinal power. Instances have indeed been produced of their seeming to prove beneficial: but as there are, perhaps, no disorders, whose nature is so little understood, whose causes are so various, and whose mitigations and exasperations have less dependence upon sensible things; there are none in which medicines operate more precariously, and in which the observer is more liable to deception.

Half a dram or a dram of the wood or leaves in substance, or an infusion of half an ounce, the doses commonly directed, have no sensible effect. Both the leaves and branches have very little smell, and a very

MATERIA MEDICA.

weak taste, of the nauseous kind. In distillation they impregnate water with their faint unpleasant smell, but yield no essential oil. Extracts made from them by water are bitterish, roughish, and subsaline: the spirituous extracts, in quantity smaller than the watery, are in taste stronger, nauseous, bitterish, and subaustere: the spirituous extract of the wood has the greatest austerity, and that of the leaves the greatest bitterishness. The berries abound with an extremely tenacious, not ungrateful, fweet mucilage.

VITRIOLUM.

VITRIOLUM & Calcanthum Ph. Parif. VITRIOL: a faline crystalline concrete, composed of metal united with a certain acid called the vitriolic acid. There are three metals with which this acid is found naturally combined, zinc, copper, and iron: with the first it forms a white, with the fecond a blue, and with the third a green falt.

1. VITRIOLUM ALBUM Ph. Lond. Vitriolum album sive Zinci Ph. Ed. White vitriol, or vitriol of zinc; found in the mines of Goslar, fometimes in transparent pieces, more commonly in white efflorescences; which are diffolved in water, and crystallized into large irregular masses fomewhat refembling fine fugar; in taste sweetish, nauseous, and styptic.

The common white vitriol of the shops contains a quantity of ferrugineous matter; of which, in keeping, a part is extricated from the acid, in an ochery form, so as to tinge the mass of a yellow hue. On disfolving the whitest pieces in water, a considerable portion of ochre immediately feparates: the filtered folution, transparent and colourless, becomes again turbid and yellow on being made to boil, and deposites a fresh ochery fediment; and a like feparation happens, though much more flowly, on standing without heat. Hence, when the folution is evaporated to the usual pitch, and set to crystallize, the crystals generally prove foul; unless some fresh acid be added (as an ounce of the strong spirit or oil of vitriol to a pound of the falt +) to keep the ferrugineous matter dissolved: + Sal vitrioli this addition both fecures the whiteness of the crystals, and prevents their growing foon yellow in the air. White vitriol generally contains also a small portion of copper, distinguishable by the cupreous stain which it communicates to polished iron immersed in solutions of it, or rubbed

Pharm. Lond.

with

with it in a moift state. The quantity of this metal is so exceedingly minute, that it is not, perhaps, of any inconvenience in the intentions for which white vitriol is commonly employed: the separation, if it should be thought necessary, may be effected, by boiling the solution for fome time, along with bright pieces of iron, which will extricate all the copper: by continued or repeated coction, greatest part of the ferru-

gineous matter also may be separated.

White vitriol is fometimes given, from five or fix grains to half a dram and more, as an emetic; and appears to be one of the quickest in operation of those that can be employed with safety. Its chief use is for external purposes, as a cooling restringent and desiccative: a dilute solution of it, as fixteen grains in eight ounces of water, with the addition of fixteen drops of weak vitriolic acid, is an excellent collyrium in defluxions and flight inflammations of the eyes, and, after bleeding and purging, in the more violent ones: a folution of it with alum, in the proportion of two drams of each to a pint of water, is used as a repellent fomentation for some cutaneous eruptions, for cleanling foul ulcers, and as an injection in the fluor albus and gonorrhæa when not accompanied with virulence. This vitriol is fometimes likewife employed as an errhine, and faid to be a very effectual dissolvent of mucous matters; in which intention it is recommended, in the German ephemerides, against obstructions of the nostrils in new-born infants.

Aqua vitrio-lica Ph. Ed.

Aqua alumi-nofa bateana Pharm. Lond.

> 2. VITRIOLUM CÆRULEUM Ph. Lond. Vitriolum cæruleum sive cupri Blue vitriol, or vitriol of copper, commonly called Roman or Cyprian vitriol, or blue-stone. This kind of vitriol is in many places produced from fulphureous ores of copper: the acid of fulphur is no other than the vitriolic; and the inflammable principle of the fulphur being diffipated either by fire or by a spontaneous resolution of the mineral, the acid remains combined with the copper (fee Pyrites): the vitriol, now formed, is either extracted by the application of water, or washed out by rain or subterraneous waters: hence in some copper mines are found blue waters, which are true vitriolic folutions of copper, and which deposite that metal on the addition of iron or of any other substance which the acid more strongly attracts. The greatest part of the blue vitriol, now met with in the shops, is prepared in England, by artificially combining copper with fulphur or its acid.

The

The vitriol of copper is of an elegant fapphire blue colour; hard, compact, and semitransparent; when perfectly crystallized, of a slattish, rhomboidal, decahedral figure; in tafte extremely nauseous, styptic, and acrid. Exposed to a gentle heat, it first turns white, and then of a yellowish red or orange colour: on increasing the fire, it parts, difficultly, with its acid, and changes at length to a very dark red calx, reducible, by fusion with inflammable fluxes, into copper.

This falt, like the other preparations of copper, acts, in doses of a few grains, as a most virulent emetic. Its use is chiefly external, as a detergent, escharotic, and for restraining hemorrhagies: for which last intention, a strong styptic liquor is prepared in the shops, by dissolving three ounces of blue vitriol and two+ or three t of alum in a pint and + Aqua via half + or two pounds to of water, then adding two ounces + or one and rulea Ph. L.

a half t of oil of vitriol, and filtering the mixture for use.

‡ Aqua ftypticaPh. Edin.

* Blue vitriol has of late been confiderably employed as an emetic by fome practitioners; and is faid to be by no means an unfafe one, as it operates the inftant it reaches the stomach, before it has time to injure by its corrofive quality. The peculiar advantage in using it is reprefented to be, that it has no tendency to become also purgative, and that its astringent power prevents the tone of the stomach from being impaired after vomiting with it. It is much recommended in the early state of tubercles in the lungs; and the following method of exhibition is directed (a). Let the patient first swallow about half a pint of water, and immediately afterwards, the vitriol dissolved in a cup-full of water. The dose may be varied according to age, constitution, &c. from two grains to ten, or even twenty; always taking care to begin with fmall ones. After the emetic is rejected, another half pint of water is to be drunk, which is likewife fpeedily thrown up, and this is commonly fufficient to remove the nausea.

In still smaller doses, the blue vitriol has been much used by some as a tonic in intermittents, and other difeases.

3. VITRIOLUM VIRIDE Pharm. Lond. Vitriolum viride, sive ferri Pharm. Edinb. Green vitriol, or vitriol of iron; commonly called English vitriol or copperas; the Roman vitriol of the Italian writers.

This fort of vitriol is produced from fulphureo-ferrugineous pyritæ, as the blue from fulphureo-cupreous ones; and as the ferrugineous minerals are much easier of resolution than the others, the ferrugineous vitriol is much oftener found native. In this native state, neither fort is free from an admixture of the other; the native green vitriols having always more or less of a blueish cast, and the blue of a greenish. The common green vitriol is prepared in large quantity at Deptsord and Blackwall near London, and at Newcastle, by boiling iron with the acid liquor, which runs from certain pyritæ after long exposure to the air: this vitriol appears to be purely martial, for if it should receive any cupreous particles from the mineral, the superadded iron would precipitate them. All vitriols may be freed perfectly from copper by adding iron to solutions of them: those, which contain even a small portion of that metal, readily discover it by staining the iron of a copper hue.

Pure vitriol of iron is confiderably transparent, of a fine bright, though not very deep, grass green colour; of a nauseous astringent taste accompanied with a kind of fweetishness. Dissolved, and set to crystallize, it shoots into thick rhomboidal masses; a part generally rising at the same time in efflorescences about the sides of the vessel. The solution deposites in standing a considerable quantity, and in boiling a much larger one, of the metallic basis of the vitriol, in form of a rusty calx or ochre: iron feems to be the only metallic body that thus feparates fpontaneously, in any confiderable quantity, from the vitriolic acid. On exposing the vitriol itself to a moist air, a similar resolution happens on its surface; which, fooner or later, according as the acid is more or less faturated with the metal, changes its green to a rufty hue. In a warm dry air, it loses a part of the phlegm or water necessary to its crystalline form, and falls by degrees into a white powder. Exposed to a gentle fire, it liquefies and boils up; but foon changes, on the exhalation of the watery part that rendered it fluid, to a folid, opake, whitish or grey mass: this, pulverized and urged with a stronger fire, continues to emit sumes, becomes † Vitriotum yellow+, afterwards red, and at length, having parted with most of its acid as well as its phlegm, turns to a deep purplish-red calx t, revivable t Colcothar by inflammable fubstances into iron.

† Vitriolum calcinatum
Ph. L. & Ed.
† Colcothar
vitrioliPhar.
Lond. & Edin.
Chalcitis
factitiaFhar.
Parif.

Pure green vitriol is in no respect different from the artificial fal martis. It is one of the most certain of the chalybeate medicines, scarcely

MATERIA MEDICA.

scarcely ever failing to take effect where the calces and other indisfoluble preparations pass inactive through the intestinal tube. It may be conveniently given in a liquid form, largely diluted with aqueous fluids: two or three grains or more, dissolved in a pint or a quart of water (which from this quantity receives no disagreeable taste) may be taken in a day, divided into different doses. This vitriol is used also, especially when calcined, as an external styptic: the styptic of Helvetius, and as is faid that of Eaton, is no other than French brandy very flightly impregnated with the calcined vitriol: a dram of the vitriol is commonly directed to a quart of the spirit, but only a minute portion of the dram dissolves in Tinatura it. As French brandy has generally an aftringent impregnation from the oaken casks in which it has been kept, the vitriol changes it, as it does the watery infusions of vegetable aftringents, to a black colour; but makes no fuch change in spirituous liquors that have not received some astringent tincture.

ftyptica P.L.

IT is from the green vitriol that the acid called vitriolic has been generally extracted; by distilling the calcined vitriol in earthen longnecks, with a strong fire continued for two days or longer. The distilled spirit appears of a dark blackish colour; and contains a quantity of phlegm, greater or less according as the vitriol has been less or more calcined. On committing it a fecond time to distillation, in a glass retort placed in a fand-heat, the phlegmatic parts rife first, together with a portion of the acid, and are kept apart under the name of spirit or weak spirit of vitriol +: at the same time the remaining strong spirit, or oil as + Spir.vitriit is called, loses its black colour and becomes clear; and this is the Pharm. Lond. usual mark for discontinuing the rectification. The college of Edinburgh now directs a weak vitriolic acid of more certain strength, made by Pharm. Edin. mixing one part of the strong acid with seven parts of water ||.

The strong acid or oil of vitriol is the most ponderous of unmetallic fluids, and the most fixed of saline ones, yielding no smell in the greatest heat of the atmosphere, and requiring, to make it boil or distil, a heat confiderably greater than that in which lead melts. Exposed to the air, Pharm. Edin. it imbibes its humidity, fo as to gain by degrees an increase of about twice its own weight. Mixed directly with water, it produces a heat fo great as to render the veffel insupportable to the hand: glass veffels are apt to crack from the suddenness of the heat, unless the commixture is

‡Acidum vitriolicum Spir. vitrioli fortis Pb. L. || Acidum vitriolicum tenue, vulgo Spiritus vitrioli tenuis very flowly performed. The most ready method of distinguishing it, in a dilute state, or when mixed with other acids, is by adding a solution of some calcareous earth, as chalk, made in any kind of acid liquor: this solution is by a minute portion of the vitriolic acid rendered milky, but suffers no change from any other species of acid; see Selenites.

Spir. vitrioli volat. Stabl. If the long-neck, in the extrication of the acid from vitriol, happens to crack in the fire, the acid that rifes after this period is found remarkably changed. It emits in the air suffocating vapours like the sumes of burning brimstone, and rifes in distillation with a heat not much greater than that which the hand can bear: to the taste it discovers little corrosiveness or acidity. Combined with alkaline salts, it loses its pungent odour; but on the addition of any other acid, it is disengaged from the alkali, so as to rise again in distillation as volatile and suffocating as before. It destroys or whitens the blue and red colours of the flowers of plants; whereas, in its fixt state, like the other acids, it changes the blue to red, and heightens those which are naturally red. This volatile spirit loses its suffocating odour, and resumes its corrosiveness, fixedness, and other qualities, by exposure to the air, which seems to carry off the inflammable principle whereon its volatility depended.

Aqua fulphurata Ph. Lond. Gas fulphuris vulgo.

Spir. fulph. per campanam Ph. L.

The fumes of burning brimftone are no other than the vitriolic acid in its volatile state; see Sulphur. If a little burning sulphur be suspended over some water in a close vessel till the sumes subside, and this repeated with fresh portions of sulphur, till about half a pound has been used to a quart of water, the liquor will be found strongly impregnated with the volatile suffocating acid, and in keeping for some time, if the vessel is not closely stopt, it will become exactly similar to water acidulated with the fixt acid. If a very large glass, open at bottom, be hung over the burning fulphur, in a damp place screened from wind, a part of the fumes will condense upon the fides of the glass, and run down in drops, which may be collected by placing a glass dish underneath: the acid thus obtained is called, from the shape of the vessel that has been generally used for condensing the sumes, spirit or oil of sulphur by the bell. The quantity of acid collected by this process is very small, greatest part of the sumes escaping: fixteen ounces of fulphur, in the most favourable circumstances, yield fearcely one ounce of phlegmatic spirit; though it is certain, that out of this quantity of fulphur, more than fifteen ounces are pure acid, of fuch strength, as to require being diluted with above an equal quantity

of water to reduce it to the pitch of common spirit of sulphur; so that if fulphur could be burnt without the loss of any of its fumes, we might obtain double its weight of an acid of the ordinary strength. The process has lately been improved, by fome particular perfons, though not perhaps to this degree, yet fo far as to afford at a very low price almost all the acid now fold under the name of oil of vitriol. The improvement confifts chiefly in burning the fulphur in very large glass vessels, in the bottoms of which some warm water is placed, whose steam serves to collect and condense the fumes.

THE acid of vitriol or fulphur, largely diluted fo as to be supportable or but gratefully tart to the palate, is the most falubrious of all the mineral acids. It is mixed with watery infusions, spirituous tinctures and other liquids, as an antiphlogistic; as a restringent in hemorrhagies; and as a stomachic and corroborant in weaknesses, loss of appetite, and decays of constitution, accompanied with slow febrile symptoms, brought on by irregularities, or fucceeding the suppression of intermittents by Peruvian bark. In feveral cases of this kind, after bitters and aromatics of themselves had availed nothing, a mixture of them with the vitriolic acid has happily taken place: the form commonly made use of is that of a spirituous tincture: fix ounces of oil of vitriol are dropt by degrees into a quart of rectified spirit of wine, the mixture digested for three days in a very gentle heat, and afterwards digested for three days longer with an ounce and a half of cinnamon, and an ounce of ginger +; or a pint + Elixir viof an aromatic tincture drawn with proof spirit is mixed with four ounces of the strong acid : these liquors are given from ten to thirty or forty t Elix.vitridrops, in any convenient vehicle, at fuch times as the stomach is most empty. A mixture of oil of vitriol with spirit of wine alone, in the proportion of one part of the former to three of the latter, digested together for some time, is used in France as a restringent in gonorrheas, female fluors, and spittings of blood.

* This acid, diluted with water, has been given internally with great fuccess in the itch. It was first used for this purpose in the Prussian army in 1756, and has fince been much employed in feveral parts of Germany. The dose recommended is from an eighth to a fourth of a dram of the pure acid twice or thrice a day. It is faid to fucceed equally in the dry

trioli Pb. E.

oli acidum Phar. Lond.

Aqua rabelliana vulgo Eau de Rabel. P. Parif.

and

and moist itch; and when given to nurses, to cure both themselves and their children.

When oil of vitriol and rectified spirit of wine are long digested to-

gether or distilled, a part of the acid unites with the vinous spirit into a new compound, very volatile and inflammable, of no perceptible acidity, of a strong and very fragrant smell, and an aromatic kind of taste: this dulcified part, more volatile than the rest, separates and rises first in distillation, and may thus be collected by itself. The college of London directs a pound of oil of vitriol and a pint of rectified spirit of wine to be cautiously and gradually mixed (a great conflict and heat ensuing if they are mixed hastily) and set to distil with a very gentle heat: that of Edinburgh orders the same quantity of the oil of vitriol to be dropt into four times as much of the vinous spirit, and the mixture to be digested' in a close vessel, for eight days, previously to the distillation, with a view to promote the coalition of the two ingredients. The different proportions of the acid spirit to the vinous, in these prescriptions, make no material variation in the qualities of the product, provided the distillation is duly conducted; for the smallest of the above proportions of acid is much more than the vinous spirit can dulcify, and all the redundant acid remains in either case behind. The true dulcified spirit rises in thin fubtile vapours, which condense upon the fides of the recipient in straight striæ: these are succeeded by white sumes, which form either irregular striæ or large round drops like oil; on the first appearance of which, the process is either to be stopt, or the receiver changed. The spirit which these fumes afford, very different from the dulcified one, has a pungent acid smell like the fumes of burning sulphur: on its surface is found a small quantity of oil, of a strong penetrating and very agreeable fmell, readily diffoluble in spirit of wine, to a large proportion of which it communicates the fmell and tafte of the aromatic or dulcified spirit. The college of Edinburgh, in order to fecure against any acidity in the dulcified spirit, order it to be rectified, by mixing it with an equal measure of water, in every pint of which a dram of salt of tartar has been diffolved, and drawing off the spirit again by a gentle heat *(a).

Spiritus vitrioli dulcis.

O1. vitrioli dulce Hoffm.

This

[&]quot;(a) The Edinburgh college, in their last pharmacopoxia, have manifestly shewn how little they conceive the acid to enter as a constituent part of this preparation, and at the same time have directed an effectual method of preventing its presence in it. They order the acidum vitriolicum vinosum, vulgo spiritus vitrioli dulcis, to be made by simply mixing one part of vitriolic ether with two of rectified spirit.

This spirit, taken from ten to eighty or ninety drops, strengthens the stomach and digestive powers, relieves statulencies, promotes urine, and in many cases abates spasmodic strictures, and procures rest. It is not effentially different from the celebrated mineral anodyne liquor of Hoffman; to which it is frequently, by the author himself, directed as a substitute. It is evident, from Hoffman's writings, that his anodyne Liquor anowas composed of the dulcified spirit and the aromatic oil which comes over after it, but the particular proportions of the two he has no where specified: the faculty of Paris directs, under the title of his preparation, twelve drops of the oil to be dissolved in two ounces of the spirit; the college of Wirtemberg feems to think, that all the oil, and all the spirit, obtained in one operation, were mixed together, without regard to the precise quantities.

dynus mineralis Hoffm.

The dulcified spirit is sometimes used as a menstruum for certain resinous and bituminous bodies, which are more difficultly and languidly acted upon by pure vinous spirits. It is often mixed with aromatic and stomachic tinctures, in cases where the stomach is too weak to bear the acid elixirs above mentioned: eight ounces are commonly added to a pint Elix. vitrioli of the officinal aromatic tincture+, in which it does not, like the acid dulce + P. L. undulcified, occasion any precipitation; or the ingredients of the aromatic tincture are infused in the dulcified acid, instead of common rectified spirit . A medicine of this kind was formerly in great esteem under the 1 Pb. Edinb. name of Vigani's volatile elixir of vitriol, the preparation of which was long kept a fecret, and first made public in the pharmacopæia reformata: it is prepared by macerating, in some dulcified spirit of vitriol free from acidity, a small quantity of mint leaves curiously dried, till the spirit has acquired a fine green colour: to prevent the necessity of filtration, during which the more volatile parts would exhale, the mint may be suspended. in the spirit in a fine linen cloth.

If the dulcified spirit, rectified as above prescribed from a solution of fixt alkaline falt, be shaken with equal its quantity of a like solution, and the mixture fuffered to rest; an ethereal fluid rifes to the furface, and great part of the dulcified spirit may be recovered again from the remainder by distillation. I am informed by Dr. Hadley, that he has observed the largest proportion of ether to be obtained, by using the strongest vitriolic acid of the shops with equal its quantity by measure of spirit of wine, and distilling immediately by a heat sufficient to make the mixture boil; and that by this management, from three pints of oil of vitriol, and fix pints of rectified spirit of wine, he obtained two pints and a half of the ether. * The following is the method prescribed for making ether, in the last Edinburgh pharmacopæia. To thirty-two ounces of rectified spirit of wine in a glass retort, add at once an equal weight of strong spirit of vitriol. Mix them gradually by gentle agitation, and immediately set them to distil in sand previously heated, so that the mixture may be brought as soon as possible to boil, in which heat it is to be continued till sixteen ounces are come over. The receiver must be cooled by water or snow. To this distilled liquor, two drams of the strong alkaline caustic are to be added, and the distillation repeated in a very high retort, with a very gentle heat, till ten ounces are come over. To the residuum after the first distillation may be added sixteen ounces of fresh rectified spirit, when more ether will be procured; and this may be several times repeated.

Liquor æthereus vitriolicus Pb. Ed.

> The ether or ethereal spirit is the lightest, most subtile, volatile, and inflammable, of all known liquids: it quickly exhales in the air, diffufing an odour of great fragrance: it does not mingle with water, with acid liquors, with alkaline liquors, or with vinous spirits, at least not in any confiderable quantity, only a fmall portion of the ether being imbibed by them: it unites with oils in all proportions, diffolves balfams, and refins, and extracts the oily and refinous parts of vegetables. It has been hitherto regarded chiefly as a matter of curiofity, nor are its medicinal qualities as yet much known *(a). Malouin looks upon it as one of the most perfect tonics, friendly to the nerves, cordial and anodyne; and fays he has found it to be a good remedy in rheums, for abating coughs, especially those of the convulsive kind. Its great volatility renders the taking of it very incommodious: the author above mentioned orders, as the most convenient form, from three to twelve drops to be dropt on fugar or powdered liquorice, a little warm water or fome warm infusion to be immediately added, and the whole swallowed directly. It has been reported to give immediate ease in violent headachs, by being rubbed on the temples.

^{* (}a) It has fince come more into use in flatulent and spasmodic complaints, the gout in the stomach, nervous assumas, and the like. Though it will not mix with water, it may be disfused in a sufficient quantity of it so as to be taken without much difficulty.

The vitriolic acid faturates a larger quantity of fixt alkaline falts than any of the other acids, and diflodges therefrom fuch other acids as have been previously combined with them: of the strong spirit or oil of vitriol, about five parts are fufficient for eight of the common vegetable fixt alkalies. The neutral falt refulting from its coalition with this Tartarum kind of alkali, is of a bitterish taste, very difficultly soluble in water, and scarcely fusible in the fire: in small doses, as a scruple or half a Sal enixum dram, it is an ufeful aperient; in larger ones, as four or five drams, a mild cathartic, which does not pass off so hastily as the sal catharticus, quibusdam. and feems to perform its office more thoroughly. This falt has been commonly prepared with the alkali obtained from tartar, and is hence called vitriolated tartar: fome dilute the oil of vitriol with fix times the Ph. Edinb. quantity of warm water, and drop into it a folution of the alkaline falt till a fresh addition occasions no further effervescence: others, instead of Ph. Lond. the pure acid, use vitriol in substance, which being dissolved in boiling water, any alkaline falt, gradually superadded till the effervescence ceases, absorbs the pure acid, and throws down the metallic basis of the vitriol; one part of the alkali is nearly fufficient for two of the vitriol. This last method, more troublesome than the other in regard to the preparation, more effectually secures the medicine from a redundance of acidity. The wholefale dealers substitute for this falt that which remains after the extraction of the acid of nitre by means of oil of vitriol, which, when duly prepared, is exactly the same (see Nitrum): but as they commonly employ in that process an over-proportion of the vitriolic acid, and vend the remaining falt uncrystallized, we often meet with it so very acid as to be utterly unfit for use.

With the mineral fixt alkali, and the earth called magnefia, this acid forms compound falts of a bitterer tafte, fomewhat less purgative, and much easier of folution, than that with vegetable alkalies: with volatile alkalies a very pungent ammoniacal falt, whose medicinal effects are not well known. The strong acid, boiled on argillaceous earths to dryness, corrodes a portion of them, and concretes therewith into an austere styptic falt. Calcareous earths it does not dissolve into a liquid state, but may be combined with them, by precipitation from other acids, into an indiffoluble concrete feemingly of no medicinal activity. Among metallic bodies, it dissolves zinc and iron readily; copper, silver, quickfilver, lead, and tin, very difficultly: it is fitted for acting on the two 4 U first

vitriolatum.

& Arcanum duplicatum

first by dilution with three or four times its quantity of water: the others require the undiluted acid, and a heat sufficient to make it boil; when, the more phlegmatic parts exhaling, so much of the pure acid matter remains combined with the metals, as to render them, in part at least, dissoluble in water; see the respective metals.

The medical qualities of the acid in its volatile state are very little known, and those of the combinations thereof with alkalies not at all, though they should seem to deserve enquiry. The volatile acid of burning brimstone may be commodiously transferred into fixt alkalies, by dipping linen clothes in a strong folution of the alkali and suspending them over the fumes, of which they will quickly imbibe fo much as to neutralize the alkali: this neutral falt being rubbed off, the clothes may be again moistened with the alkaline lye, exposed to the acid fumes, and these processes alternately repeated (a). The neutral falt thus obtained differs greatly in its tafte and other properties, and doubtlefs also in its medical virtues, from that which is produced by the coalition of the fixt acid with the same alkali, that is, from vitriolated tartar. It dissolves more eafily in water, and shoots, not into octangular crystals, but into fmall flender ones like short needles. On adding to it the fixt vitriolic acid (or even the weaker acids of nitre or fea-falt) the volatile acid is difengaged from the alkali; and though, in the compound falt, its pungent fmell was wholly suppressed, it now rifes in distillation as pungent and fuffocating as the original fumes of the brimstone. The neutral falt, in a dry form, may be kept unchanged for years: disfolved in water, and exposed for some time to the air, or if roasted with a gentle heat, it becomes the same with vitriolated tartar.

ULMARIA.

ULMARIA five Regina prati Pharm. Edinb. (b) Barba capræ floribus compactis C. B. Spiræa Ulmaria Linn. Meadowsweet or Queen-of-the meadows: a plant with tall, smooth, reddish, brittle stalks; and oval, sharp-pointed, indented leaves, set in pairs along a middle rib, with smaller pieces between, and at the end a larger odd one divided into three sections, wrinkled and green above, white underneath: on the tops come forth large thick clusters of little whitish flowers, followed each by several crooked seeds set in a roundish head. It is perennial, common in moist meadows, and slowers in June.

The leaves of ulmaria recommended as mild aftringents, discover to the taste or smell very little soundation for any medical virtues. The slowers have a strong and pleasant smell, in virtue of which they are supposed to be antispasmodic and diaphoretic, and which in keeping is soon dissipated, leaving in the slowers only an insipid mucilaginous matter. As these slowers are more rarely used in medicine than their fragrant smell might rationally persuade, Linnæus suspects that the neglect of them has arisen from the plant being possessed of some noxious qualities, which it seemed to betray by its being left untouched by cattle: it may be observed, however, that the cattle, which refused the ulmaria, resused also angelica, and other herbs, whose innocence is apparent from daily experience.

ULMUS.

ULMUS Pharm. Edinb. Ulmus campestris & theophrasti C. B. Ulmus campestris Linn. Elm: a tall common tree; covered with a rough, chapt, brownish, brittle bark, under which lies a white, smooth, tough, coriaceous one; producing in the spring, before the leaves appear, imperfect flowers, followed by slat roundish capsules, containing each a single seed.

THE inner tough bark of the elm tree, of no manifest smell, discovers, on being chewed, a copious slimy mucilage, of no particular taste: the outer brittle bark is much less slimy, but equally void of smell and taste. It may therefore be presumed, that if elm bark has been found of use in nephritic cases, in which it is recommended by authors; or externally against burns, for which it is applied by the common people; it was of use no otherwise than as a simple emollient. Neither the purgative virtue ascribed to it by some, nor the astringent by others, appear to have any foundation.

*A decoction of the inner bark of elm has been employed in cutaneous diseases in some of our hospitals; and an account of its efficacy has been published by Dr. Lysons in Vol. II. of *Medical Transactions*, and since, in a separate work. In making this decoction, four ounces of the bark 4 U 2

fresh from the tree are boiled in two quarts of water to one. It is of a beautiful light purple colour, when the elm is in flower; but browner at other times. Its taste is mildly aftringent; and an extract from it is very austere. It has no purgative effects, as some have alledged, but rather the contrary. Where it succeeds, it generally at first increases the efflorescence. Patients are usually directed to drink half a pint twice a day, and to perfift in the use of it some months.

URINA.

URINA Ph. Paris. URINE. The recent urine of healthful subjects is naufeoufly bitter, very faline, scarce manifestly alkaline or acid. As foon as it begins to putrefy, it emits volatile alkaline vapours; and if distilled, when moderately putrefied, by a gentle heat, it yields a concrete volatile alkaline falt: as volatile alkalies have a strong antiseptic power, the vapours of putrefied urine are not observed, like those of cadaverous animal substances, to be productive of putrid diseases. A pungent caustic volatile spirit may likewise be obtained from recent urine, by inspissating, and then distilling it with the addition of quicklime.

If the putrefied urine be flowly inspissated, in glass or stone-ware veffels, to the confiftence of a thin fyrup, and fet for fome weeks in a cold place, brown crystals will shoot from it, confisting partly of marine falt, and partly of a falt of a peculiar kind, which shoots before the marine, and which, by repeated folutions, filtrations, and crystallizations, may be purified both from that falt and from the adhering oil. In this + Sal micro- state +, it appears perfectly neutral, and impresses on the tongue a sense of coolness with a slight bitterishness: laid on a red-hot iron, it bubbles, emits volatile alkaline vapours, and runs into a colourless pellucid substance resembling fine glass: this apparent glass is manifestly acid, though but weakly fo, diffolves in water, neutralizes alkaline falts, and with volatile alkalies regenerates the original neutral falt. One of its most distinguishing characters is, that a mixture of it with inflammable matters, as foot or powdered charcoal, on being heated to ignition in an open veffel, emits flashes like lightening, and, on being distilled in a retort with a moderately ftrong fire, yields the highly inflammable concrete called phosphorus.

cosmicum, five fal effentiale urinæ.

Urine

Urine is fometimes applied externally, boiled with bran, as a refolvent and discutient, in which intentions it is said to be very efficacious. Recent cows urine has been drank in the spring, to the quantity of a pint or more every morning, for several days, as an attenuant and deobstruent in different disorders: the nauseous draught purges plentifully by stool, and sometimes vomits. The peculiar salt of urine is but of late discovery, and its medicinal qualities are as yet unknown.

URTICA.

URTICA Pharm. Edinb. Urtica urens maxima C. B. Urtica dioica Linn. Common stinging nettle. Infusions and decoctions of this herb, or its expressed juice, are recommended in different disorders as aperients, and said to loosen the belly: the juice, depurated and gently inspissated, discovers a considerable taste, of the subsaline kind.

UVÆ PASSÆ.

RAISINS: rich sweet grapes, dried by the sun's heat in the warmer parts of Europe. Two sorts are directed for medicinal use. 1. Uvæ Passæ majores Pharm. Lond. Passæ majores Pharm. Edinb. Raisins of the sun; the fruit of the vitis damascena dried upon the tree; the stem of each cluster, when the grapes are ripe, being cut almost through, so as to prevent the afflux of any fresh juice. 2. Uvæ Passæ minores seu corinthiaca. Currants; the fruit of the vitis corinthiaca picked from the stalks.

THESE fruits are used as agreeable lubricating acescent sweets, in pectoral decoctions, and for obtunding the acrimony of other medicines and rendering them acceptable to the palate and stomach: the first fort inclines most to acidity, the sweetness of the latter being more of the mucilaginous kind. They both give out their sweetness and their pleasant flavour to water and spirit: the stones or seeds are supposed to communicate a disagreeable relish, and hence are generally directed to be taken out; but it did not appear on trial that they give any taste at all to water, proof spirit, or rectified spirit.

UVA URSI.

UVA URSI: Vitis idea foliis carnosis & veluti punctatis, sive idea radix dioscoridis C.B. Arbutus (Uva Ursi) cauliculis procumbentibus, soliis integerrimis Linn. BEARS WHORTLEBERRY: an evergreen trailing shrubby plant; with numerous small oblong oval leaves; monopetalous white slowers with a sless-coloured edge cut into five sections; and red berries. It greatly resembles the common red whort-bush; from which it may be distinguished, by the leaves being more oblong, and by the slower having ten stamina, and the berry five seeds; whereas the slower of the common whort has only eight stamina, and the berry often twenty seeds. It is found on the snowy hills of Austria and Styria, but more plentifully on the Swedish hills. It is also a native of the Highlands of Scotland, and is now cultivated in some of our gardens.

THE leaves of this plant have a bitterish astringent taste; without any remarkable smell, at least in the dry state in which they have been brought to us from Germany. Insusions of them in water strike a deep black colour with solution of chalybeate vitriol, but soon deposite the black matter, and become clear: I do not recollect any other astringent insusion, from which the blackness, produced by vitriol, separates so very speedily.

The leaves of uva urfi have of late been greatly celebrated in calculous and nephritic complaints, and other diforders of the urinary organs: the dose is half a dram of the powder of the leaves, every morning, or two or three times a day. De Haen relates, after large experience of this medicine in the hospital of Vienna, that suppurations, though obstinate and of long continuance, in the kidneys, ureters, bladder, urethra, fcrotum and perinæum, where there was no venereal taint or evident marks of a calculus, were in general completely cured by it: that of those who had a manifest calculus, several found permanent relief, so that, long after the medicine had been left off, they continued free from pain or inconvenience in making water, though the catheter shewed that the calculus still remained: that others, who seemed to be cured, relapsed on leaving off the medicine, were again relieved on repeating its use, and this for feveral times fuccessively; while others obtained from it only temporary and precarious relief, the complaints being often as fevere during

during the continuance of the medicine as when it was not used. It may be observed, that in several cases which he relates, paregorics were joined to the uva ursi; and that other mild astringent plants have been recommended for the same intentions; from some of which De Haen himself expects the same good effects. The trials of the uva ursi, made in this country, have by no means answered expectation: in all the cases that have come to my knowledge, it produced great sickness and uneasiness, without any apparent benefit, though continued for a month.

*WINTERANUS CORTEX.

WINTERANUS CORTEX, Cortex magellanicus. WINTER'S BARK. The tree producing the Winter's bark (Winterana aromatica Soland.) is one of the largest forest trees on Terra del Fuego. Its leaves are ever-green, smooth, oval, and entire. Its flowers consist of seven petals, with from sifteen to thirty stamina, and from three to six germina, terminating in as many stigmata. Each germen becomes a seed-vessel, containing several seeds. The bark of the trunk of the tree is externally grey, and very little wrinkled.

The Winter's bark, which takes its name from Capt. Winter, who discovered it on the coast of Magellan in 1577, is brought to us in pieces of different degrees of thickness, from a quarter to three quarters of an inch. It is of a dark brown cinnamon colour, with an aromatic smell when rubbed, and of a pungent, hot, spicy taste, which is lasting on the palate, though imparted slowly. A watery infusion of it struck a black colour with a solution of green vitriol. From an infusion of two ounces of the bark, coarsely powdered, was obtained on evaporation an extract weighing two drams and twenty-sour grains. The same quantity, treated with rectified spirit, yielded two drams of extract. A pound of the bark was insused in a proper quantity of water, and the liquor submitted to distillation. The distilled water was clear, of a pleasant taste, and somewhat of the cinnamon slavour. There was no appearance of effential oil. The residuum afforded six ounces of a soft extract, of a grateful aromatic taste.

A mixture of this bark feemed very effectually to cover and correct the difagreeable taste and smell of certain drugs; a property common to it with the canella alba.

Almost

Almost the only use hitherto made of the Winter's bark has been by the crews of ships navigating the streights of Magellan, as a preservative from the scurvy. It has been confounded in the shops with the canella alba, from which it is totally different.

An exact description of the plant, with a figure, is contained in a paper published in the Medical Obs. and Ing. Vol. V. from whence this account is extracted.

ZEDOARIA.

ZEDOARIA Pharm. Lond. & Edinb. Zedoaria longa & Zedoaria ZEDOARY: the root of an Indian plant, (Amomum rotunda C. B. scapo nudo, spica laxa truncata, Berg. Mat. Med.) brought over in oblong pieces, about the thickness of the little finger and two or three inches in length; or in roundish ones + about an inch in diameter; of an ash-colour on the outside, and white within. The long fort is said by fome to be the strongest, but the difference, if any, is very inconfiderable, and hence the college allows both to be used indiscriminately.

† Zerumb-eth P. Parif.

This root has an agreeable fmell, and a bitterish aromatic taste. It impregnates water with its fmell, a flight bitterness, a considerable warmth and pungency, and a yellowish brown colour: the reddishyellow spirituous tincture is in taste stronger, and in smell weaker, than the watery. In distillation with water, it yields a thick ponderous effential oil, fmelling strongly of the zedoary, in taste very hot and pungent: the decoction, thus deprived of the aromatic matter, and concentrated by inspissation, proves weakly and disagreeably bitter and subacrid. A part of its odorous matter rifes also in the inspissation of the spirituous tincture: the remaining extract is a very warm, not fiery, moderately bitter aromatic, in flavour more grateful than the zedoary in fubstance.

Zedoary root is a very useful warm stomachic. It was employed by fome as a fuccedaneum to gentian root; at a time when a poisonous article, mixed with the gentian brought from abroad, rendered its use hazardous: but from the above analysis it appears to be not entirely fimilar to that fimple bitter; its warm aromatic part being the prevailing principle, in virtue of which, its spirituous extract (the most elegant preparation of it) is made an ingredient in the cordial confection of the London pharmacopæia.

ZIBETHUM.

ZIBETHUM.

ZIBETHUM Pb. Edinb. (a) Civetta. CIVET: a foft unctuous odoriferous substance, about the consistence of honey or butter; of a whitish, yellowish, or brownish colour, and sometimes blackish; brought from the Brazils, the coast of Guinea, and the East Indies; found in certain bags situated in the lower part of the belly of an animal of the cat kind*(b). The bag has an aperture externally, by which the civet is shed or extracted.

This substance has a very fragrant smell, so strong as, when undiluted, to be disagreeable; and an unctuous subscrid taste. It is used chiefly in perfumes, rarely or never for medicinal purposes, though the singular effects which musk has been lately found to produce may serve as an inducement to the trial. It unites with oils, both expressed and distilled, and with animal sats: in watery or spirituous liquors it does not dissolve, but both menstrua may be strongly impregnated with its odoriferous matter, water by distillation, and rectified spirit by digestion: by trituration with mucilages, it becomes soluble in water.

ZINCUM.

ZINCUM Pharm. Edinb. ZINC, or TUTENAG: a blueish white metal; crackling, in being bent, like tin, and quickly breaking; about seven times specifically heavier than water; beginning to melt in a moderate red heat, and very slowly calcining on a continuance of the fire; in a moderate white heat flowing thin, burning, sulgurating, with a bright deep green or blueish green flame, and subliming into light white flowers, which concrete about the upper part of the vessel, or on the bodies adjacent, into thin crusts, or soft loose filaments like down or cobwebs. In its metallic form, and in that of a calx or flowers, it dissolves readily in all acids, and precipitates from them almost all the other metallic bodies.

The calces or flowers of zinc are difficultly revived into their metallic form. Though perfectly fixed in the fire so long as they continue in a

(a) Expunged.

*(b) Or rather of the weafel kind.

state of calx; yet, as calces in general require for their revival a greater heat than that in which the metal itself melts, and as a full melting heat is the greatest that zinc can support; the instant they are revived, they burn and calcine again in open veffels, and escape through the pores of close ones. Hence some ores and preparations of this metal have been long kept in the shops, and even chemically examined, without being discovered to be such. The revival may be effected, by using compact veffels, of fuch a structure, that the zinc, in proportion as it is restored to its metallic form by the charcoal powder or other inflammable additions commonly made use of for those purposes, may be suffered to fublime or run off from the heat without being exposed to the outward air; or by adding some other metallic substance to detain it, as copper, which is thus changed into brafs.

This metal has but lately been received into the shops in its own form; in which it deferves a place, as affording preparations superiour to the ores or productions of it now made use of. A white vitriol made from pure zinc, by diffolution in the diluted vitriolic acid and crystalli-+ Vitriolum zation +, is doubtless preferable for medicinal use to the common impure white vitriol; and the white flowers, into which it is changed by det Calx zinci flagration t, to the very impure calamine and tutty. Moderately pure white flowers, sublimed from it in the brass or other furnaces, wherein zinc or its ores are melted with other metals, were formerly kept in the shops, and distinguished by the names of pompholix and nihil album.

* The flowers of zinc were first used as an internal medicine by the celebrated chemist Glauber, but were little known in practice till Dr. Gaubius, of Leyden, gave an account of their virtues in his Adverfaria. They have fince been much employed in convulfive and spasmodic difeases, and sometimes with good effects. Even obstinate epilepsies have been rendered much less violent by their use. Like all other medicines, however, in diseases of this class, their good effects are often only temporary, and they often fail altogether. When the flowers are genuine, a grain or two generally at first excites nausea or sickness, but by degrees a considerable dose may be taken with little or no sensible effect. As they are liable to be adulterated, it may be proper to mention, as tests of their purity, that they make no effervescence with acids; and that, when exposed to a strong heat, they become yellow, but on cooling, turn white again. An application for external use, made by mixing one part of flowers

album P.Ed.

vulgo flores zinci P. Ed.

flowers of zinc with fix of the fimple liniment of wax and oil, is directed Unguent. e in the Edinburgh pharmacopæia.

calce zinci Phar. Edin.

ZINGIBER.

ZINGIBER Pharm. Lond. & Edinb. & C. B. GINGER: the root of a reed-like plant (Amomum Zingiber Linn.), growing spontaneously in the East Indies, and cultivated in some parts of the West; brought over in knotty branched flattish pieces, freed from the outer bark, of a pale colour and fibrous texture: that which is least fibrous is accounted the best.

This warm aromatic root; of common use as a spice in flatulent colics, &c. appears to be much less liable to heat the constitution than might be expected from the penetrating heat and pungency of its tafte, and from the fixedness of its active principles. It gives out the whole of its virtue to rectified spirit, and great part of it to water, tinging the former of a deep, the latter of a pale yellow colour: the spirituous tincture, inspissated, yields a fiery extract, smelling moderately of the ginger: the watery infusion, boiled down to a thick consistence, dissolved afresh in a large quantity of water and strongly boiled down again, retains still the heat and pungency of the root, though little or nothing of its fmell: there does not feem to be any of the common spices whose pungency is of fo fixed a kind. In the shops are kept a syrup made from an infusion of three + or four t ounces of the root in four + or three t pints of boiling water, which is agreeably impregnated with its warmth and flavour; and the candied ginger ||, brought from abroad, which is likewife moderately || Zingiber aromatic.

Syr. zingib. + Ph. Edinb. 1 Ph. Lond.

conditum Pb. Edinb.

*POSTSCRIPT.

Some progress having been made in printing this edition before the publication of the Edinburgh Pharmacopæia of the present year (1783), the following alterations and additions could not be noticed in their proper places.

WEIGHTS AND MEASURES. In order to avoid the confusion arising from the promiscuous use of terms signifying both weight and measure, the college have thought proper entirely to abolish liquid measure, and to reduce every thing, as well fluid as solid, to troy weight.

PILULÆ ALOETICÆ. The fal polychrest in these is omitted.

TINCTURA SACRA. Leffer cardamom feeds are fubstituted to Jamaica pepper; and two pounds of wine is directed instead of one pint and a half.

AMBRAGRISEA expunged.

ANGELICA SYLVESTRIS added.

TARTARUS EMETICUS five ANTIMONIALIS, is now directed to be made in the following manner. Infuse butter of antimony in water containing as much fixed alkali as will precipitate all the antimony.

Wash

Wash and dry this precipitate; and boil nine drams of it with two ounces and a half of finely powdered crystals of tartar, till the powders are dissolved. Strain the solution, evaporate to a pellicle, and set it to crystallize.

VINUM E TARTARO ANTIMONIALI made by diffolving tartar emetic in white wine, in the proportion of twenty-four grains to a pound.

EMPLASTRUM E HYDRARGYRO, five CÆRULEUM (in place of Empl. Mercuriale) made of common plaster six parts, quicksilver three parts, oil-olive and white resin each one part.

Pulvis Mercurii cinereus. Dissolve quicksilver in equal its weight of weak nitrous acid; dilute the solution with water, and add spirit of sal ammoniac enough to precipitate all the mercury. Wash and dry the precipitate.

TINCTURA FOETIDA. Two ounces of asafætida dissolved in a pound of dulcified spirit of sal ammoniac.

Hyoscyamus. The inspissated juice of it is directed as an officinal.

TINCTURA Moschi. Two drams of musk dissolved in a pound of rectified spirit.

THE three following articles of the Edinburgh catalogue were omitted in their proper places; the first through mistake, the two last for want of sufficient information concerning them, which the editor at length obtained by the favour of Dr. Duncan.

ACONITUM.

ACONITUM Pharm. Edinb. Aconitum Napellus Linn. Blue Wolfsbane. This is a perennial plant, having many stalks three feet high or upwards arising from one root; alternate petiolated leaves divided into five parts, each portion cut into linear segments; and terminal bunches of irregular blue flowers with five petals, many stamina, and three.

three pistils, succeeded by three capsules containing several seeds. It is a native of various parts of Europe, and of Virginia; and is planted in gardens with us.

Blue wolfsbane when fresh has a strong odour, but no peculiar taste. The leaves powdered are said by Stærck to impress the tongue with a durable acrimony; but Bergius (a) afferts that he did not find this to be the case; and from this circumstance, and the sigure of the plant given by Stærck, he concludes that the species with which he made his experiments was not the Napellus, but the Aconitum Cammarum of Linnæus, which much resembles it (b). The expressed juice has an ungrateful smell, and an acrid, slightly styptic taste. On inspissation it yields an extract, of similar smell and taste, and subsaline.

Succus fpiffatus Aconiti Pharm. Edin.

The fresh plant has long been known as one of the most virulent of the vegetable poisons. It occasions convulsions, giddiness, infanity, violent purgings both upwards and downwards, faintings, cold sweats, and death itself. Dr. Stærck was the first who ventured to introduce it into medicine. He found that the extract given to the quantity of ten, twenty, and even thirty grains, excited a sweat without inconvenience; and by persisting in the use of it, great relief was obtained in fixed rheumatic and arthritic pains, scirrhous glandular tumours, venereal nodes, anchyloses, amaurosis, and other similar complaints. Other practitioners have experienced the same good effects in some degree, and the Edinburgh college has received the extract as an officinal. In this, as in all the other medicines of suspicious and dangerous properties, it is proper to begin with very small doses, and increase them as they can be borne.

CURSUTA.

CURSUTÆ radix Pharm. Edinb. This is a foreign root, which has been used by some practitioners at Edinburgh for more than forty years. It is a strong bitter, and has very much the appearance and taste of gentian. Dr. Home, in his list of the materia medica, stiles it Gentiana lutea sylvestris; while he terms the common gentian, Gentiana lutea sativa. No botanical author, however, makes this distinction; nor can the name of cursuta be met with in any writer that the editor has

(a) Mat. Med. 483.

(b) Haller (Stirp. Helv.) makes the fame observation.

confulted.


consulted. The Edinburgh college received it on Dr. Home's recommendation; but it is little used there, and is not in general kept in the shops.

HIPPOCASTANUM.

HIPPOCASTANUM Ph. Edinb. Æsculus Hippocastanum Linn. Horse-chestnut. The fruit of this tree, which is a trilocular capsule, containing two seeds in each cell, has been given as a food to sheep; and steeped in water, so as to extract its bitterness, is said to fatten poultry. It falls spontaneously into a saponaceous gluten, which has been used instead of soap for washing linen. No writer mentions its medical application; but the Edinburgh college have admitted it on the recommendation of Dr. Gardiner, who says, that three or sour grains of the powder snuffed up the nostrils in the evening, operates next morning as an excellent sternutatory, and thereby proves very beneficial in obstinate inflammations of the eyes.

The bark of the horse-chestnut has been proposed in Italy, according to Haller, as a substitute to Peruvian bark in the cure of intermittents; and the experiment has proved successful (a).

(a) Stirp. Helvet. I. 442.



Λ			Page	00		Page
ABELMOSCH	The same	- Street	137	Aconitum Anthora		- 64
Abies	A Comment	2	I	Napellus	-	- 689
balfamea -	15-1	THE PARTY OF	3	pardalianch	es - ,	- 278
Canadenfis -	-		3	Acopa, V. Trifol. pale	id	- 655
Abiga -	- II- SIUYO	-	223	Acorus adulterinus	a pille	- 359
Abrotanum femina	- 15 1492		5	Calamus -	1	- 173
mas -	and Minney	-	4	verus -	Augustina Paris	- 173
Abfinthium -	- 5. 6		6	vulgaris -	113	- 357
Alpinum	- BULL DE LOUIS	-	9	Acte -	Ca Tembra em	- 575
maritimum		-	8	Acus mofchata -	-	- 322
minus	-	-	9	Adeps -		- 502
Ponticum	-	- 4	9	Adianthum Canadenfe	51 1	- 20
Romanum	-	-	8	verum	+ Symptoness	- 19
Seriphium	-	A TOP	9	vulgare, /	. Trichomanes	
Valefiacum		200	9	Adipfon -	Meine	- 325
Acacia Germanica			521	Aer fixus -		21, 81
vera -	-	-	10	Ærugo -	200 N	- 23
Acajou -	-	-	55	Æther -	- 450	, 573, 675
Acanthina mastiche	- 1	75	198	Æthiops antimonialis		- 110
Acanthus -	-		11	mineralis		- 101
Acer		-	11	vegetabilis	THE THE PARTY OF THE PARTY	- 531
Acetabulum -	-	The state of	57	Æthufa Meum -		- 423
Acetofa -	-	-	12	Agallochum -	STONESTERN !	- 388
Acetofella -	-	-	400	Agaricus -	A PROPERTY OF	- 24
Acetum -	The state of the s	Total Control	.13	mineralis	- Staglar	- 255
Achillea -	-	The state of	424	quercinus	ter amount of	- 26
Ageratum		21 61	131	Ageratum -	100000	- 131
Ptarmica -		100	523	Agnus caftus -	112	- 27
			4	Y	1000	Agrimonia
						The second secon

I N D E X

		Pag			THE PARTY
Agrimonia -		- 28			Page
odorata	the real line	- 20			54
Agrioriganum -		- 46			55
Agripalma -		- 194			139
Aigeiros -	Valletine.	- 519			438
Ajuga -		- 22		-	56
reptans -		16;			167
Aizoon		- 599			361
Alcanna -		- 50			57
Alchimilla -		- 20		- 1	346
Alexandrina herba	THE DESIGNATION	- 34		-	341
Alifma -		- 27			525
Alkekengi -		- 3			57
Alkermes -		- 36		7	302
Alleluja -		- 40		-	58
Alliaria -		- 3		4	591
Allium -	DE VERY	- 3		A.	60
Alnus -		1.70			
nigra -		- 3		1	61
Aloe		30		100	63
Aloes lignum -		38			69
Alfine -				1	675
Altercum -		- 39		1	461
Althæa -	alongart m	34		AUTET	91
Alumen -	STATE OF THE PARTY	- 4			306
plumofum	anailabana in	- 63		WHITE !	611
antiquo	rum	- 4		RI NEW	222
Alypum -	Carnison Lubu	65		1000	223
Amaracus, V. Majorana	Celamous	- 40		magi	220
Amaradulcis -	50137	- 600		Tank Sal	526
Amaranthus luteus, V. Se	mechas	- 28		ST OFFIE	III TO UBLICATE
Amarella -		- 51			544
Ambarvalis -	- bytilbho	- 51			280
Ambarum -		- 4			395
citrinum	catalogica (CD mass)	- 62	A TOP CONTROL OF THE PROPERTY		64
Ambra grifea -	mintak	- 4	The state of the s		201
liquida -	Assigna.	- 39			77
Ambrofia -		- 16		P	438
Ambrofioides -		- 16		45	417
Ambutua -		- 47		THE PARTY	653
Amianthus -		- 63		175-18	77
Ammi -	antaraoattana 1	- 4	OCCUPANT OF THE PROPERTY OF TH	Mant	78
Ammoniacum gummi	The state of the s	- 4		ange.	79
Ammosteus -	our ansakan	- 46		-	498
Amomum Germanorum	OF THE PARTY OF	- 50		THE NO	499
verum -	-	- 4		1	499
vulgare	Harry Wall	- 5		PRESIDE	498
Amygdalæ amaræ	STAN TORKAL	- 5		03	345
dulces	the many	- 5		11 12	447
Amylum -	-	- 30		1	448
Anacampferos -	-	- 63		-	80
					Aquæ

			Dage				Page
Aquæ medicinales			Page 84	Afparagus		1/-	123
alkalinæ			84	Afphaltus -			154
catharticæ		THE REAL PROPERTY.	85	Afplenium -			217
chalybeata		The state of	87	Scolopendrium	_	HELICAN	396
cupreæ			89	Trichomanes			654
marinæ			86	Affis -		-	187
fulphureæ			90	Aftacus -			184
Aquila alba -			109	After Atticus, V. Eryngium		- 4	289
Aquilæ lignum -			389	Aftrantia -			353
Aquilegia -			90	Athamanta -	-		423
Aquilina -	2 . 66		90	cretenfis			271
Arabicum gummi		-	332	Athanafia -	_	10	633
Aracus aromaticus	His		659	Atractylis gummifera			198
Aranearum telæ			91	Atriplex olida			124
Arantia -			124	Atropa Belladonna			607
Arbutus Uva Urfi			682	Avellana purgatrix			539
Arcanum corallinum			105	Avena -			308
duplicatum	0		677	Aurantia Curaslavensia			127
tartari			18	Hispalensia -		-	125
Archangelica -				Sinenfia		- 4	127
Arcium -		THE PERSON	375	Aureliana Canadensis		E	323
Arctium Lappa			137	Auricula muris			500
Ardefia -			137	Auricularia -		1000	420
Arefta bovis -			342 461	Auripigmentum			116
Argentina -		-		Aurum -			128
THE RESERVE OF THE PARTY OF THE		M. Marie	91	fulminans			
Argentum -			92	mufivum	100	Mary In The	617
Argilla alba -		MAIN OF	93			MET A TOTAL	
Ariftolochia -	- S. J.	HI HE	234	Axungia -	1500	AND THE PARTY	502
Armoracia -	5,50	OMES!	111	В.			
Arnica -	•		533	Baccæ Bermudenses			-0-
Arfenicum -		- A - A	277	Badian -	1		585
Artemifia -		Water o	113		3		63
Abrotanum	1	17.54	116	Balanus myrepfica Balauftia	*		141
Abfinthium	F 191	- 4 1	4		-	A PART	130
glacialis	-	1000		Balfamea -	-		3
pontica	-	1,70	9	Balfamita femina		STORES IN	131
Santonicum	5 11	11174	9	mas Balfamum -	21911	alphay 5	131
Artemisiæ lanugo		13 5 2	579	Americanum	Tip !		466
Arthanita -	2 97	47712	429	Brazilienfe		A CONTRACT	133
Arthritica, V. Chamæpitys	Man	71415	117	Canadenfe	MA	And the state of	132
Arum -		alai Ei	223	Canadenie	1000	THE WAR	3
Dracunculus	7.00	THE PARTY	118		-	PHO PARTY	132
Afa dulcis -	17 kmp	THE PARTY OF	279	Gileadenfe Indicum	3.4		466
fœtida -	5113	- 1	141		504		133
	7 24	aller -	120	Mexicanum	-	47	133
Afarum -			121	Peruvianum	-	Topin Ti	133
Asbestos -	1	50	632	album		1515	135
Afclepias Vincetoxicum	2		661	nigrur	n	the ledges	134
Afelli, V. Millepedes	7	1 1	425	Tolutanum	-	LE BOOK	135
Afpalathus -	4	100 700	389	Bamia mofchata	3	-	137
			4	Y 2			ncia

I N D E X

AND THE RESERVE OF THE PARTY OF		Page	经营业技术 是从1960年的			
Bancia -		- 481	Brassica Eruca -	The Park		Page
Bangue -		- 187	Erucastrum			288
Banilia -		- 659	marina -		-	288
Baptificula, V. Cyanus	AND AS A	- 266	Napus -		-	164
Barba capræ -		- 678	D		-	435
Jovis -			Britannica -			534
Bardana -		227	Brunella -		•	377
Bafilicum .	NEW YORK	-31	Brufcus -		1	520
Batrachium -		13/	Brutua -			546
Baurach -		- 322	Bryonia -			479
Bdellium	Billion .	- 438	Bubon macedonicum		-	165
D1	The state of	- 138				499
D . 1 .	-	- 139	Bugloffum -	-		167
Belladona -		- 656	Bugula -	-	-	167
D 111		- 607	Bumelia, V. Fraxinus			280
	-	- 140	Bunias -	-	-	435
Ben -	-	- 140	Buphthalmum -	-	-	140
Benzoinum -	*	- 141	Burfa paftoris -	2 3 1 1 1 1 1	120	168
		- 141	Butomon -	P. Colon		359
Berberis -	-	- 144	Butua -	- 240	-	479
Beta -		- 145	Butyrum antimonii	-	-	73
Betonica -	-	- 146	Buxus -			168
aquatica	- 100	- 598				
Pauli -	- 1915	- 660	C.			
Betula -		- 147	Cabureiba -	-	-	133
Alnus -		- 34	Cacao -	-	-	169
Bezoar microcofmicum	- mania	- 151	Cacara -		-	276
minerale	- ALDAN	- 73	Cadmia fossilis -	-	-	170
occidentale		- 150	lapidofa	+010	17-019	170
orientale	-	- 149	metallica	-	6211	113
porcinum		- 151	Cadjuct -	-	1494	276
fimiæ -	-	- 151	Cæruleum Berolinense	-		294
Bilis -	4 .	- 292	nativum	-	-3194	382
Bingalle -	- Livingar	- 211	Caffe -			243
Bismuthum -	-	- 152	Cajous -	- STATE TON	1	55
Bistorta -	4	- 153	Calambac -		-	388
Bitumen Barbadense	- miles	- 498	Calambour -	To America	-	389
Judaicum		- 154	Calaminaris -	- 1922		170
Blitum fætidum		- 124	Calamintha -	42574 1915	-	171
Bois de Coiffi -		- 528	magno flore	9111/	WAR	173
Boletus igniarius -		- 26	montana	-	2000	172
Bolus Armena & aliæ	al makein V	- 155	Calamus aromaticus	med Q Th	1	173
Bonus Henricus	- udikas	- 157	Calcanthum -	2	14	667
Borago -	viga sireni(- 158	Calculus humanus	2030170	102 (55	151
Borax -	- masth	- 159	Calendula -	6	120	174
Botrys -	(Checker)	- 162	Alpina, V. Arnic	a	1000	277
Mexicana	muncipality	- 162		2 4, 99	100	654
Bovista -		- 402	Calomelas -	1411/1-1	13.99	109
Branca leonina		- 481	Caltha -	plantoni	1 3516	174
urfina -	myampile"	- 11	Calumba -	12-14(74)	THE P	246
Braffica -	nie!	- 163	Calx viva	188 -11	The state	175
		103	Can title		Camb	
					Camil	514

			Page				Page
Cambogia Gutta	-	-	316	Cataria -	3-	THE HOSPITA	439
Camphora -	1 2 10 1		179	Catechu -	-	-	640
Camphorofma -	-	-	426	Cauda equina -	4	-	287
Cancrorum chelæ		-	184	porcina -	2444		500
oculi		-	184	Causticum antimoniale	-		73
Candelaria -	2	-	659	commune fortiu	S	- 558,	559
Canella alba -		-	185	mitius	Die.	559,	
malavarica	-	19100	208	lunare	-		92
Cannabina aquatica, V. E	upatorium	-	290	Cedrinum lignum	200	a topical	363
Cannabis -	4	1291	186	oleum	1200		239
Cantharides -		-10	188	Cedronella -	-		417
Caphura -	2	-	179	Cedrus -	-		212
Caphura baros -	112000		239	Celeri -	-	1000	79
Capillus Veneris	- Chilomat	mun i	. 19	Centaurea benedicta	-	. 1	195
Capficum -	LINE !	44	508	Centaureum	12		213
Carabaccium -		-	206	Cyanus		A CONTRACTOR	266
Carabe, V. Succinum	HE DE SELLE	1	623	mofchata			267
Caranna -		-	191	Centaurium majus	-	14.9	213
Carcas -	R. District 16.	-	539	minus		Contract to	213
Cardamantica -		1	385	Centimorbia -	-	V race &	450
Cardamine -	The state of	-	192	Centipedes -	-	1121000	425
Cardamomum majus	almin de	-	194	Cepa -	310	A	214
majus Gall	orum	-	327	Cera alba -			215
medium	L. combined	-	194	di cardo, V. Carlina		124	198
minus	-	-	193	cinnamomi			239
Siberiense	1 2 31		63	flava -		THE WAR	215
Cardiaca -	2000	-	194	Cerafa -		1	307
Cardopatium -	a good all the second		197	Cerealia -		CH20	308
Carduus benedictus	= 4		195	Cerefolium -			218
pineus		210	198	Cervifpina -	1.10	1410 20 7	611
Caricæ -	-	4	197	Ceruffa -	-		512
Carlina -	-	-	197	antimonii	-		67
gummifera	-	100	198	Ceterach -	- 11		217
Carpathicum oleum	-	10	639	Cevadilla -		Shau to	217
Carpentaria -	Man allerdi		424	Chacarilla -	-		281
Carpobalfamum	Market Street	-	199	Chærefolium -		AN PARTY	218
Carthamus -		-	199	Chalcitis -			670
Carum -	- 1000	10	200	Chalybs -			294
Caryophylla aromatica	Lu trust	1	201	Chamæacte -	-	nul and	576
Caryophyllata -	- 1000000	-	203	Chamæciffus -			336
Caryophylloides cortex	1 1 1 1 1 1 1	-	205	Chamæclema -			336
Caryophyllus ruber	CLOSE OF	Maria.	204	Chamæcypariffus		To Live	
Cafcarilla -	an agilen	-	281	Chamædrys -	-		219
Cafia caryophyllata	P.U.	-	205	palustris		Action 1	596
fiftularis -	- Volening	1	207	fpuria	-	1	661
lignea -	2 1/2	1000	208	Chamælæa -	-	1111111111	648
Castoreum -	Given Re	120	209	Chamælæagnus			434
Cafumunar -	'sameyeth	2	211	Chamæleon -	_		198
Cataputia -	-	230	653	Chamælinum -		19/,	398
major	and albert	2.1	539	Chamæmelum -			220
			333			Chamæme	

I N D E X

		Page				Page
Chamæmelum flore pleno -		222	Coccus Cacti			240
fætidum -		223	Cochlea -		1911	
Chamæpitys			Cochlearia -			394
Cheiranthus Chieri -		223	armoracia	STATISTICS.	1 1 14	240
01.::		224	marina		119	533
		224		T. EHAD	- 10	241
Chelæ cancrorum		184	Codago-pala -	-		248
Chelidonium majus -		224	Coffea -	5	1	243
minus -		226	Colchicum -	Can month		244
Chenopodium ambrofioides	en einli	162	Colcothar -	Shir large	- del	670
Bonus Henricus		157	.Colocynthis -		-	245
Botrys -	1 31/3	162	Colophonia -		-	639
fætidum -	312 7	124	Coluber Berus	-	-	664
Chermes	W. T.	365	Colubrina -	-	-	602
Cherva		539	Colubrinum lignum	-	-	454
Chibou gummi		281	Columbinus pes		-	322
Chiliophyllon		424	Columbo -			246
China		226	Coma aurea, V. Elichryfum			284
China chinæ		485	Conessi -	The state of		248
Chryfanthemum Leucanthemum		140	Conium maculatum	Control (Print)	S. Alley	228
			Confolida major	STATE OF THE		
Chrysocome, V. Elichrysum		284		- 389	5	248
Cichoreum		227	media		140,	
Endivia -	-	284	minima	510000000	FIRST	140
Cicla		145	minor	To stand	-	520
Cicuta	OF SHIP	228	faracenica	CONTRACTOR OF	-	665
aquatica -	1	78	Contrayerva -	*	-	249
Cimolia alba -	-	234	Germanica	-	-	662
purpurafcens -		234	Convallaria -		-	393
Cinæ femen	- 3	579	multiflora		-	604
Cinara	10.	235	Copal -	- 10	180,	250
Cinchona Carribæa -		495	Corallina -	-	-	251
Jamaicenfis -		495	Corallium -		- 14	251
officinalis -		485	Coriandrum -			252
Cineres clavellati -		556	Cornu cervi -		2011	253
			Corona terræ -			336
Ruffici -		556	Cortex cardinalis de Lugo			485
Cinnabaris antimonii		74	caryophylloides			100-200
factitia -	Dia-1990	102		1949	1010	205
Græcorum -	- 7	577	coneffi -		-	248
nativa -	- 141	235	culilawan	1	-	206
Cinnamomum -	- 141	236	eleutheriæ	A CONTRACTOR	1	281
Citrago	W.	417	granatorum	-	15.54	328
Citrea		239	Jefuiticus	1700 10016	-	485
Citrullus	Miles - Isk	260	Magellanicus		1-1630	683
Civetta	New-bull	685	Peruvianus	-	- 100	485
Clematis recta	Cr 10 2	301	ruber	· him	-	492
Clinopodium	81 - 2	412	fimarouba	-	-	605
Cnicus	194,		thuris -	- 10	-	281
Cobaltum	-	113	thymiamatis	-	-	649
Coccagnidia		649	Winteranus	-	-	683
Coccinella		240	Cortufa -	-	-	411
	77 (-132 14 1m)	365	Coftus corticofus			185
Coccus baphica -		202	College College		C	oftus
					TO THE PARTY OF TH	

14 0 14	111011	Page	OTHORI WORD			Done
Coftus hortorum	- miles	- 131	D.			Page _
minor	No. of the last	- 131				472
orientalis		- 254			L	648
Cotonea -		- 267				648
Cotula fœtida -	S. Linia	- 223		200	2.5	648
Cotyledon marina	The state of	- 57			-	619
Courbaril -		- 60				271
Craffula -	de modern	- 637		4 - 10	0	271
Crepitus lupi -	m. Vijisal	- 402				618
Crespinus -	2 2 1 1 1 1	- 144			- IA	272
Creflio -		- 437				526
Creta -		- 255		-	2 4	591
Crocus -	7100	- 257			12000	204
antimonii	- contin	- 69		-	-	577
Indicus -		- 265				274
martis -		72, 297				274
metallorum		- 69				275
Crystallus -		- 259				504
mineralis		- 446			a du	276
Cubebæ -		- 260		120000		665
Cucullata -	4 411	- 504				277
Cucumis -	3-16	- 260		3 14 15		278
agreftis		- 261				249
Colocynthis		- 245		-		523
Cucurbita -		- 261		13 713		426
Citrullus	1	- 260				279
Pepo		- 261				
Culilawan		- 206				523
Cuminum -		- 268		A STATE OF THE STA		334
pratenfe	The state of	- 200				609
Cunila bubula -	The Later	- 468				009
fativa -		- 588				6
Cuprum -		- 262				576
Curcuma -	The state	- 265			737	262
Curfuta -			Elatine -			280
Cufcuta -		- 286			72	
Cyanus -		- 266		-	T. C.	434 623
lapis -	- Word	- 382				281
mofchatus	- Contract	- 267			40	
Cyclamen -	enter de	- 117			49,	78
Cydonia -	2 3 30	- 267				281
Cyminum -	Valore in	- 268			12.	284
Cynocrambe -	net Steader	- 423			- 2	The second second
Cynogloffum -	- malau	- 268				337 284
Cynorrhodon -		- 543		1031		285
Cynosbatos -	- 300	- 543			1	286
Cynoforchis -		- 589				287
Cyperus longus	- 1956	- 269		-		287
rotundus	1 - 168 651	- 270			1	288
Cypira -		- 265		-		289
STATE OF STA		,			Eryfi	
					22711	-44 66 341

		Dane				
Eryfimum -		- 289	Fucus veficulofus			Page
Alliaria		100,000	Fuga dæmonum			530
Erythrodanum -		- 31	Fuligo -	, 100-0	-	346
Efula -	7	- 545	Fungo -			312
Eupatorium Avicennæ		- 653				312
cannabinum		- 290	Fungus arboreus		1116	26
Mefues		- 290	igniarius	- mitte	o de la	26
	-	- 131	laricis		-	25
odoratum		- 29	petræus mar		-448	57
verum	- date att	- 28	terrestris orb	icularis	101-12	402
Euphorbia Cyparissias		- 653	vinofus			27
Paralias		- 652	Furfur -	-	- 4	308
Euphorbium -	-	- 291				
Euphrafia -	de do Les A s	- 292		G.		
			Gabianum oleum	- lings	200	497
F.		Margarett .	Galanga -	100	iline s	313
Faba crassa -	manage of	- 637	Galbanum -	-	ma-	314
febrifuga -	-	455	Gale -	-	nie.	434
Indica -	- 1000	455	Galeopfis -	1 4 TH 1975	11-15	375
purgatrix -	-	- 539	Galerita -	18 com.	-	495
fancti Ignatii	·	- 455	Gallæ -	-	-	315
fuilla -	1-19-19/2 1110	345	Gallitrichum -	450 - 450	1.	344
Fabaria -	a service .	- 637	Gallium -	-	-	316
Farfara -	-	- 656	Aparine	- 1999	10-	77
Farinacea -	- 200	308	Gamandra -	11 - 1	0-5	316
Febrifuga -	- recultify	- 414	Gambogia -	-	-	316
Febrifugum Craanii	-	- 70	Garofmum -		-	124
Fel		292	Garyophyllus -	- 000		201
Ferrum -	- Tartin	- 293	Genista -		-	317
Ficaria, V. Chelidonium m	inus .	- 226	Gentiana	-		319
Ficus	- 11	- 197	Centaurium	-24	-	213
Filix -	-6	300	Indica		1	320
Flammula -		- 532	Geoffræa -		-	320
Jovis	- + 1	- 301	Geranium -		-	322
Fæniculum dulce	-	302	Geum urbanum		-	203
erraticum		591	Gingiberi -		-	687
porcinum	-	- 500	Gingidium -			218
Sinense	-	63	Ginfeng -	1881-189	27-	323
tortuofum	-	604	Gladiolus luteus			359
vulgare		303	Glans unguentaria			141
Fœnum camelorum		361	Glechoma hederacea	Sel Vinter		336
Græcum		304	Glycypicros		AUT CAN	609
Folium Indum		406	Glycyrrhiza		V. British	
orientale		600	Gnaphalium Steechas			3 ² 5 284
Formica -			Gramen caninum			
Fraga -		304	Grana cnidia	REINSTELL		326
Frangula -		307				649
Fraxinella -		305	paradifi		1000	327
Fraxinella -	1	274	regia -		The state of	539
Fructus horai		306	tiglia -	The state of the s	11/5/1/	539
		307	tinctoria	THE PARTY OF	12 5	365
Frumenta -		- 308	Granata -		-	327
				13 20 13 13 13 13 13	Grai	natus

			P.		Page				Page
Granatus filvestris		-	4-		130	Herba cephalalgica			660
Granum mofchi		-		-	137	doria	-		665
Graphoy -				-	278	felis		- H	439
Gratia Dei -		-		322,		fancti Jacobi	-	- 1	348
Gratiola -		-		-	328	julia	-		131
Guaiacum -		-		-	329	fanctæ Kunigu	ndis,	V. Eupatorium	290
Gummi acanthinum		* 13		-	332	militaris	-		424
ammoniacum		-Hilling		-	47	papillaris	-	HALL WE'TH' LAND	376
anime		-		-	61	pedicularis	-		618
arabicum		7		-	332	pulicaris	-		522
chibou		- 445		- 18	281	regia	-	STATE TO SE	457
courbaril		-		-	60	facra	-	A STATE OF THE PARTY OF THE PAR	660
elemi		-		-	281	falivaris	100	JIN AND THE	526
Gambiense		-			366	Roberti			322
guaiaci				To	329	trinitatis	-		341
gutta		-			316	Hermodactylus	-		342
hederæ		To but		7 +	336	Hibernicus lapis			342
juniperinum		-			363	Hiera picra	-		38
lacca		7			372	Hieracium Pilofella		,	500
rubrum aftrir	igens				366	Hierobotane	- 3	The section of	660
Senegalense		- 1			333	Hippocastanum		Union mortuain	691
Thebaicum		-			332	Hippolapathum	15		378
tragacanthæ					334	Hippomarathrum			591
Gutta gamba		1			316	Hippofelinum			343
Gypfum -				-	000	Hippuris vulgaris Hirundinaria	-		287
	н.					Hoitziloxitl	-	450,	
Hæmatites	11.				221	Holofteus		建设建筑场内的	133
Hæmatodes	177				334	Hordeolum			469
Halicacabum		and and			322	Hordeum		NEW PROPERTY.	217
Hardefia					342	causticum		A CONTRACT OF THE PARTY OF THE	308
Hedera arborea					335	perlatum			217
terrestris					336	Horminum			310
Helenium			-		285	Humulus Lupulus		Alleria	344 401
Heliochryfum, V. Sto	chas		-		284	Hydrargyrus		THE PROPERTY OF THE PARTY OF	
Helleborus albus	-		-		337	Hydrolapathum			93
niger			-		339	Hydropiper			484
Helvetii pulvis	-		-		43	Hyofcyamus	-		345
Helxine, V. Parietari	a		-		480	luteus	-		443
Hepar antimonii	-		-		69	Hypericum	-		346
fulphuris	-		-		628	Hypociftis	0-1		347
Hepatica nobilis			-		341	Hysfopus	-		347
terrestris	4-1		-		341	Hyftricis lapis	3-	A STATE OF THE PARTY	151
Hepatorium, V. Eupa	torium		-		290			Service Control of the Control of th	1.50
Heptaphyllum	-		-		654		I.		
Herba alba	-				9	Iberis	1 -		385
Alexandrina			-		343	Ibifcus	-	CONTRACTOR OF THE PARTY OF THE	40
Apollinaris	-		-		345	Ichthyocolla			351
bafilica			-		457	Igafur	-	-	455
Britannica	1500		-		377	Illecebra	-	visa de zama	352
and the second			-		4 2			Illi	cium

I N D E X

		-	-			
THE			Page			Page
Illicium anifatum	-	-		Lactucæ -	-	373
Imperatoria	-	-	353	Ladanum -	-	367
Inguinalis, V. Eryngiu	m	-	289	Lamium album -		375
Intybus	- 1 1995	-	284	Lampfana -	-	376
Inula Helenium	-	-	285	Lanaria -	-	659
Ipecacoanha	THE PERSON NAMED IN	-	353	Lapathum acetofum	-	12
Iquetaia	-	-	598	acutum -	-	376
Iringus, V. Eryngium		-	289	aquaticum		377
Irio -	-	-	289	hortenfe -	_	378
Iris Florentina	-	5 4	358	unctuofum		157
nostras purpurea	-		357	Lapides cancrorum -	1	184
palustris lutea	2	. 17 %	359	Lapis bezoar	21 17	148
Iva arthritica, V. Char	mænitys		223	cæruleus -		382
Ixine -			198	calaminaris -		
Ixion -		THE D	198	cyanus -		170
TATOII -			190	hæmatites -	-	382
	T			Hibernicus -	-	334
	J.				-	342
Jacobæa	-		348	hystricis -	-	151
Jalapium		-	349	lazuli -	-	382
Jalappa alba	-	-	414	Malacenfis -	-	151
Japonica terra	-	-	640	porcinus -	-	151
Jecoraria		-	341	fepticus -	-	558
Jemu -	-		316	fimiæ -		151
Jesuiticus cortex			485	fpecularis -	-	600
Jujuba	-	- 8	360	Lappa	-	. 137
Julia herba			131	Lapfana -	-	376
Juneus odoratus	-	100	361	Larix	-	638
Juniperus	-	-	361	Lafer		120
Sabina	-	200	548	Laserpitium -	-	120
Jupiter, V. Stannum		-	616	Chironium	-	482
		-		Siler -	-	603
The state of the s	K.			Lathyrus -		653
Kali -		grante	364	Laudanum -		465
Keiri -	La Maria	A Stoler	224	Lavendula angustifolia	Sain	Service Control of the Control of th
Kermes -		- Hier	365	latifolia -		379 380
mineralis			71	Stœchas -		
Ketmia				Laver Germanicum, V. Becabunga		619
Kiki -			137	Laureola -		139
			539	Laurocerafus -	-	648
Kina kina			485			380
Kino -	T.		366	Laurus -	-	381
Kunigundis herba, V.	Eupator.	-	290	Lazuli lapis -	-	382
				Leontodon Taraxacum	-	272
	L.			Leonurus Cardiaca -	-	194
Labdanum	=		367	Lentifcus -	-	383
Lac -	-	-	369	Lepidium -	-	384
lunæ	-	-	255	Iberis -	-	385
fulphuris		-	628	fativum -	-	438
virginis	-	- 5	143	Leucanthemum Diofcoridis	-	222
Lacca	-		372	odoratius	-	220
Lachryma abiegna	-	4	638	vulgare	-	140
			THE PAR		Le	ucoium

	10000		Dane			Done
Leucoium luteum			Page 224	Lopeziana radix -		Page
Leuconymphæa		a family	456	Lotus filvestris		531
Leucopiper				Lujula	The state of the s	4.600
Levisticum		n all all all	505	Lumbrici terrestres		400
Libanotis			385	Lumbus Veneris -		401
Lichen cinereus terres	1	A SECTION AND	544			424
	tris	Tolars	386	Luna		92
iflandicus	7		387	Lupulus -	- E-100	401
petræus		the state of the s	341	Lycoperdon -	No. 1 THE RES	402
Lignum aloes			388	Lyfimachia Nummularia		450
aquilæ		1000	389	M.		
calambac	300		388			
Campechense		1 4	390	Macerone -		343
cedrinum		-	212	Macis	F41 10 15 C	402
fpur	ium		363	Macropiper -	P. Maria Color	506
colubrinum			454	Magellanicus cortex -	100	683
guaiacum	-	- 1011	329	Magistrantia, V. Imperatoria	Bert Tay	353
Indicum	-	100	390	Magnefia alba	And the same	403
juniperinum	-	7 411	363	vitrariorum	7.5	405
lentifcinum		all ally	383	Maianthemum -	37137	393
Moluccenfe	-	15 A P 40	539	Majorana -	Line Par	405
nephriticum	-	Windson .	440	Syriaca -	idm to the	411
pavanum	-	moint Dies	539	Malabathri oleum -	BUT INTERNAL	239
Quaffiæ	-	ability.	528	Malabathrum -	Managar.	406
rhodium	-	-	391	Malicorium -		328
fanctum	2000	Section 1	329	Malva	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	407
fantalum	- 610	SPACE TO SERVICE STATE OF THE	578	Malvaviscus, V. Althæa	Anniety	40
fappan	- 813	The Table	390	Maniella kua -		265
faffafras	A- 14	-	587	Maniguetta -	15 to 18 to	327
vitæ	7		329	Manna	1100101	408
Ligusticum	-	-	385	Brigantiaca -	-	408
Lilium album		distanting	392	Marathrum -	10007	302
convallium	-	ALERO III	393	Marathrophyllum -	7	500
Limaces	-	-	394	Marcafita -	-	152
Limones	130	of the	394	Marcaffita -		527
Linaria	-	- 1	395	Marchantia polymorpha	Add to the first	341
fegetum	-	- mi	280	Margaritæ , -	The state of	410
Lingua avis	-	- 010	306	Marrubium -	-	410
canina	-		268	Cardiaca -	1.	194
cervina	- 101	PERSONAL PROPERTY.	396	Mars	-	294
Lini femen	-	non-	396	Marum Syriacum -	-	411
Linum catharticum	- Dell	-	398	vulgare -	Halane -	412
Liparis	-	19 H 1= 10	504	Maftiche -	September 1	413
Liquidambra	-	a statute of	398	acanthina -	-	198
Liquiritia	- 34	Mule I	325	Mastichina -	- 1	412
Lithargyrus	-	THE PARTY OF THE P	512	Materia perlata -	-	68
Lithospermam	17	TANK - ATT	399	Matricaria -		414
Lixivium causticum	-	The Contract of the Contract o	558	Chamomilla	-	222
faponarium		San Lies	558	Matrifylva, V. Ulmaria	1000	678
tartari	- 1	in the land	556	Mechoacanna -		414
Lobelia -			399	nigra -		349
				4 Z 2	Meco	

			Page				Page
Meconium	-	MATERIAL STREET	476	Myacantha	40	PE.	546
Medica malus	34	Name of the	239	Myofotis			
Medulla faxi	-	4.	255	Myrica Gale	William Co.		500
Mel -	-	milestal big	415	Myriophyllon			434
Melampodium		min new car	339	Myriftica nux			424
Melanopiper		10116	505	Myrobalani			450
Meleguetta		The last section	327	Myrrha		1	430
Melilotus		and dates	416	Myrtacantha		100	431
Meliffa	Deberter	a story rivers	No. of Concession, Name of Street, or other Publisher, Name of Street, or other Publisher, Name of Street, or other Publisher, Name of Street, Original Publisher, Name of Street, Name of Str	Myrtilli	14 H. (15)		546
Calamintha			417	Myrtus -		100	434
grandiflora	13/ 115		172	Brabantica		1	433
Nepeta			173			-	434
Turcica		ELECTRICAL DESIGNATION OF THE PERSON OF THE	172	Myxa -	-		360
	IN CASE	The Real	426	Myxaria		-	360
Meliffophyllon Mellitis -	1.20	the law selli	417		N T		is Page
Meloe veficatorius	Consult.		417	NT 11	N.		
	-	a stone about	188	Napellus	-	-	689
Mentastrum	-		420	Napha -			125
Mentha aquatica	-	-	420	Naphtha	-	-	496
cataria	-	-	439	Napium	-		376
cervina	-	-	524	Napus dulcis			435
corymbifera			131	filveftris	- 100000	-	435
palustris	-	Mr. Alex Disaver	524	Nardus Celtica		-	436
piperitis	-	-	421	Indica	- 1	-	436
Pulegium	-	-	524	ruftica		-	121
Saracenica	-		131	Nasturtium aquaticum		-	437
vulgaris	-	A PARTY OF	418	hortense	- 1 - 100	-	438
Menyanthes	-	10. 10.00	655	pratenfe	-	-	192
Mercurialis	-	L. L. Lindson	422	Natron -	200		438
montana	-		423	Nenuphar	-		456
Mercurius	-	SHEET THE	93	Nepeta -			172, 439
vitæ		4 5 5 5 6	73	Nephriticum lignum			440
Metella nux	-	14.00	453	Neroli essentia	THE RESERVE	1	125
Meum -	-	11.00	423	Nicotiana			441
Mezereon	-	Le ne	648	minor			443
Militaris herba	and the	avier wheels	424	Nihil -	MATE TO SERVICE		686
Milium folis		Control of	399	Ninzin -	Barrier State		
Millefolium		The state of the s	424	Nitrum -	The No. of London	12.00	324
Millemorbia		MENON CO.	598	ammoniacale		ATE.	443
Millepedæ				antiquorum			448
Minium	A SOLIN	1000	425	calcareum			438
Græcorum			512		-1		449
Mirabilis Peruviana			235		ologorum	-	567
			349	causticum		-	72
Moldavica	-	-	426	cubicum		-	448
Momordica Elaterium		-	261	vitriolatum	-	-	447
Mora -	-		307	volatile	STATE OF THE PARTY	-	448
Morfus gallinæ, V. Al	line	1	39	Nucista -	-	-	450
Mofchata nux		-	450	Nuculæ faponariæ	-	-	585
Mofchus	Sections.	4	427	Nammularia	- 10	-	450
Moxa -	-	TO LEGATE	429	Nux Barbadensis	10		539
Muscus maritimus	-	38.4	251	ben -		-	141
		A THE PARTY OF THE	18 36 1				Nux

			D			Dane
			Page	Ofyris -		Page
Nux metella			453	Ovum -		395
- mofchata	5	5	450	Oxalis -		470
pistacia	34	19109	452	Acetofella -		
purgans		Town I was	539			400
vomica			453	Oxyacantha -	760	144
Nymphæa			456	Oxylapathum -		376
				Oxmyrfine -		546
	0.			Oxyphænicon -		632
Ochra -	-		457	Oxys		400
Ocularia, V. Euphrafia		-	292	Oxytriphyllon		400
Oculi cancrorum	-		184	P		60
Oculus bovis	-	C 44 14	140	Pæonia -	-	470
Ocymum			457	Palea de Mecha -	-	361
Enanthe	-	-	458	Palimpiffa -		510
Offa alba		-	563	Palma -	-	472
Olea -	-		459	Palma Christi -		539
Oleum abietinum	-	-	638	Palum fanctum, V. Guaiacum	-	329
animale		-	560	Panava -	-	540
Carpathicum		- 1	639	Panax -	-	48z
Gabianum	-	-	497	quinquefolium -		323
laurinum	-	-	381	Panis cuculi -		400
Neroli	*	-	125	porcinus, V. Arthanita	-	117
olivarum	-	-	459	Papaver	46	1, 473
palmæ	- STREET	4	472	erraticum -		477
petræ	-	-	496	Papillaris -	-	376
fufinum	-	-	392	Paradifi grana -	1	327
Syriæ	-	u l	427	Paralyfis -	40.50	478
templinum	-	-	2	Pareira brava -	-	. 479
terræ	- 10	CIENT IN	496	Parietaria -	-	480
vitrioli	4	-	671	Parthenium -	-	414
Olibanum	-	-	460	mas -	12	633
Olivæ	-	-	459	Passerina, V. Ornithoglossum	12.39	306
Olufatrum		- 1715	343	Paffulæ -	-	681
Onisci	-	-	425	Pastinaca	27	1, 481
Onitis	-		468	Patientia -	-	378
Ononis	- DAMES	-	461	Patrum pulvis -	1000	485
Opium		-5/19	461	Pavana -	200	539
Opobalfamum	-	-	466	Pedicularia -	1	618
Opopanax		2010	467	Pedro del porco -		151
Orchis	Harry Holdy	2	589	Pentaphylloides -		91
Origanum			468	Pentaphyllum -	1	482
Dictamnus			274	Peplus -	2479	653
Majorana		S EN	405	Pepo -	-	251
Syriacum			411	Perforata -	-	346
Ornithogalum		-	593	Peristerium -	-	660
Ornithogloffum		125414	306	Perlæ -	-	410
Oryza		To the said	308	Perfica -	1	483
Ofteocolla	72 - Yal		469	Perficaria -	1000	484
Oftreum		-	469	Perfonata -	-	137
Oftrutium		-	353	Peruvianus cortex -	110	485
A COLOR OF THE PARTY OF THE PAR	A STATE OF THE PARTY OF THE PAR	HE LEE	333		Per	ivianus
					2 02 1	

			Page				Page
Peruvianus cortex rub	er		492	Plantago Pfyllium			522
Pes alexandrinus		-	526	Plumbago			484
columbinus			322	Plumbum			511
leonis			29	candidum, V.	Stannum		616
Petafites			495	Polium montanum	-	1	516
Petrapium			499	Polygala			14 10 20 20 20 20 20 20 20 20 20 20 20 20 20
Petroleum			496	Polygonatum			517 604
Barbadenfe			498	Polygonum Bistorta		-	
Petrofelinum			498	Hydropiper			153
Macedonie	Cum.			Perficaria			484
Petum	cum		499	Polypodium		-	484
Peucedanum			441	Filix mas		-	519
Silaus			500	Polytrichum		1-71-	300
Philanthropus			591		•	- 10	654
Phu, V. Valeriana		- with	77	Pompholyx		-	
		-	657	Populus balfamifera		7.	519
Phyllitis Phyfalis Alkekengi		-	396	Potentilla -		5	20, 631
Picea			30			-	91
Pilofella		-	1	reptans		-	482
		Park.	500	Priapeia		-	443
Pimenta	中级 保险	-	507	Primula veris	· Land	-	478
Pimpinella		17	501	Pruna	- 3400		521
Pinastellum		Table 9	500	Prunella	-	-	520
Pineus purgans	-	7 33	539	Germanorum		-	167
Pinguedo	-	-	502	Prunus Laurocerafus			380
Pinguicula	-	-	504	filveftris	-	-	521
Pinhones Indici	-	-	539	Pfeudoacorus		15	359
Pinus Abies		-	1	Pseudocostus	-/	1781	. 482
balfamea	-	-	3	Pfeudoiris		5	359
Cedrus	-	-	212	Pfeudonardus		-	379
Larix	-	-	638	Pfeudopyrethrum		-	523
Picea	-		I	Pfidium		-	328
fylvestris	•	-	639	Pfyllium	-	-	522
Piper album	-	-	505	Ptarmica	-	7	523
caudatum	-	-	260	Pulegium	-	-	524
Jamaicense	-	-	507	Pulicaris	-	-	522
Indicum	-	-	508	Pulmonaria maculofa		-	525
longum	-	-	506	Pulfatilla	-	-	525
murale	-	-	352	Punica	5	- 1	30, 327
nigrum	-	-	505	Pyrethrum	-	-	526
· Piperitis	The state of	-	384	filvestre	5 1 2 1	-	523
Pissasphaltum	-	- 10	496	Pyrites		-	527
Piffelæum	-	-	496		The second second		
Indicum	-	-	498		Q.		
Pistacia	-	-	452	Quaffia -	-	-	528
Pistolochia	1- 4	1	111	Quercula, V. Chamæd	rys	-	219
Pityufa	17	-	653	Quercus -	-		530
Pix Burgandica	-	-	639	marina		-	530
liquida	-	-	509	Quinquefolium	- 450100		482
ficca	-		510	Quinquina	-	- 1	485
Plantago	-	100	511				D. Jin
							Radix

140	MITHOL						Dago
	D		Page	Rumex -			Page 12
n !! n c!! c	R.						376
Radix Brafilienfis		-	353	acutus			
dulcis, V. Glyo		70 00	325	aquaticus Patientia			377 378
indica Lopezian	la	7 1	531				546
rofea.	-	-	637	Rufcus -			
rubra	-	4	545	Ruta -		1	547
urfina	-	-	423	c			
Raiis di Juan Lopez	- HERMAN	-	246				
di Mofambique	- dimension	-	531	Sabadilla		-	217
Ranunculus		-	532	Sabina -			548
Ficaria	-		226	Saccharum			549
Raphanus rusticanus	-	-	533	Canadense			12
filvestris	-	- 4	384	lactis			371
Rapum -	-	-	534	Saturni		1 54	513
Rapunculus virginianu	s -	-	399	Sagapenum	•	-	551
Realgar -	-	-	116	Sago -		-	473
Regina prati	-	- die	678	Sal alkalinus fixus	-	-	553
Regulus antimonii	-	-	66	fosilis	-		364, 438
mart		-	72	volatilis		-	559
ftella	itus	-	72	Sal ammoniacus	- 119/03	-	565
Remora aratri	-	-	461	fixus	-	-	572
Refina flava & nigra	-	-	639	catharticus amarus	1		567
Resta bovis	-	-	461	Glauberi		-	567
Rhabarbarum	-	-	535	communis	- 1	-	568
album	-	-	414	diureticus	· Daniel		18
antiquoru	ım	-	537	enixus	-	-	677
monachor	rum	-	378	gemmæ	- 1	-	568
Rhamnus catharticus		-	611	marinus	-	-	569
Frangula		- 03	305	microcofmicus	-	-	680
Zizyphus	-	-	360	mirabilis	-	-	567
Rhaponticum	1-19	-	537	muriaticus calcareus		-	572
vulgare	- 10 - 10 - 10	-	213	petræ	-	10	443
Rheum -		-	535	prunellæ		-	446
Rhodia radix		- 000	637	polychreftus	-	-	446
Rhodium lignum	- 100	-	391	Rupellenfis		-	636
Rhododendron	-	-	538	fedativus	21 1441	-	160
Rhus -	*_		630	vegetabilis	_	-	635
belgica	Estimated	-	434	Salep	-3	-	589
Ribefia -	-	-	307	Salivaris herba		-	526
Ricinus -		-	539	Saliunca			436
Rifagon -		-	211	Salix		1 23	573
Rifigal -			116	amerina			27
Ros Calabrinus		24	408	Salfola Kali			364
Rofæ -		2700	541	Salvia			574
Rofa filvestris			543	Sclarea			
Rofea radix			638	Sambucus			344
Rofmarinus			544	Sampfuchus			575 412
Rubia -			545	Sandaracha Arabum		MAN N	363
Rubrica -	1. 100			Græcorum	W. Lak	1111	116
Rubi Idæi fructus		2 6	457 307	Sanguis draconis		1139	
TIMOL ZUIGE IL MOUND	El de la company		30/	ounguis diaconis			577 Sanicula.
The state of the s							Onnicula.

			Page				Page
Sanicula		_	577	Seneka -		De la	Page 517
Eboracenfis			504	Serapias -		- 14	589
Santalum lignum			578	Serapinum	-		551
Santolina	. /		5	Seriola, V. Endivia	Die Familie Ce 1		284
Santonicum			579	Seriphium		-	A DESCRIPTION OF
Sapo			580	Seris, V. Endivia	La visita de	-	284
volatilis			583	Serpentaria Gallorum			
Saponaria			584	Hifpanica	1 4 2 2		279 597
nucula	100	- 1114	585	Virginiana			602
terra, V. Ci	molia	. 12	234	Serpyllum			651
Sappan lignum -	_		390	citratum		-	651
Sarcocolla			586	Sefeli -			603
Sarfaparilla			586	Massiliense			604
Saffafras	7.00		587	pratenfe		-	
Satureia		Daniel's	588	tortuofum	-		591 604
Saturnus			511	Sevum		2.12	502
Satyrion			589	Sicula			145
Savina -			548	Sidium	The Paris of the Control of the Cont		328
Saxifraga alba			590	Sigillum Salomonis			604
vulgaris		Arm was	591	Sil -	The state of the s		457
Scammonium			591	Silaus		ALL ALL	591
Scandix Chærefolium		-	218	Siler montanum, V. Se	efeli		603
Scariola, V. Endivia	E TARID	-	284	Siliquaftrum		- Contract	508
Gallorum			374	Simarouba	S Sant	*	605
Schenanthus			361	Simiæ lapis	1		151
Scilla -		Sec. of	593	Sinapi			606
Scincus -		- Links	595	Sifarum montanum		-	324
Sclarea -			344	Sifon	1 1 1 1 1 1		50
Scolopendria			217	Ammi	- Vanis		
Scolopendrium			396	Sifymbrium		10	47
Scolymus	The state of the s		235	aquaticum		-1100	437
Scopa regia			546	Sium aromaticum, V.	Amomum		50
Scordium			596	Smectis, V. Cimolia pr			234
Scorzonera	Maria Maria	-	597	Smyrnion	-	1	353
Scrophularia		-	597	Smyrnium Olufatrum			343
	Chelid. min.		226	Soda -		1-	364
Sebadilla	-		217	Sol -			128
Sebesten -	0	-	360	Solanum			607
Sedum -	The state of the s	1	599	fætidum		. 10	609
acre			352	lignofum		-	609
Telephium			637	nigrum			608
Selenites -			599	veficarium		-	30
Semen contra		2 6 0	579	Soldanella		-	164
fanctum -	S. T. R. L. A.	-	579	Solidago Virga aurea		_	665
Sementina			579	Spartium fcoparium	-	Lasage	317
Semperviyum	1	-	599	Specularis lapis	ELDER HELD	-	600
acre		Lesin	352	Sperma ceti	- Alberta	-	610
Sena -		200	600	Spica -	STATE OF STA	370	, 380
Senecio -		4	287	Celtica		- 3/3	436
Jacobæa	-	15711	348	nardi	- 1 6 11 1	12	436
			34-			Sp	igelia
							19 6385

	O MI I IV	- WI -	•	. HONT MOROM		
0.3			Page	Tr.		Page
Spigelia			611	Taraxacum -		272
Spina acida	-		144	Tarchon filvestris -	-	523
cervina			611	Tartarum -		634
infectoria			611	emeticum -		74
Spiræa Ulmaria		-	678	regeneratum		18
Spiritus vinosus recti		-	612	folubile -	-	635
teni	lior	, -, -	614	Tegula Hibernica -	-	342
Spongia		THE THE	615	Telephium -		636
Squilla	-		593	Templinum oleum	Se Hall	6
Squinanthus			361	Terebinthinæ -	-	637
Stalactitæ			255	Terra foliata tartari -		18
Stannum		-	616	Japonica - merita -		640
Staphifagria			618			265
Staphylinus			271	faponaria, V. Cimolia purp.	1 3	234
Stellaria, V. Alchim	iiiia		29	figillata -	-	155
Stelochites			469	Terræ oleum		496
Sternutamentoria		- in	523	Terrea abforbentia -	-	643
Stibium	-		65	Tefticulus caninus -	-	589
Stechas	7 7 7		619	Teucrium -	-	661
citrina	10.5		284	capitatum -	-	516
Stramonium	-		619	Chamædrys	-	219
Stratiotes	S- Anti-		424	Chamæpitys	-	223
Struthium			353	creticum	-	516
Styrax alba			135	Scordium -	-	596
calamita	775	1	620	Thea -	-	645
liquida			622	Germanica -		660
Succinum .			623	Theriaca rusticorum -	-	32
cinereum			44	Thermæ	-	90
grifeum	7	mat ris	44	Thlaspi -		646
Sulphur		76710	625	Burfa pastoris	-	168
Sumach			630	Thuris cortex	281,	649
Supercilium Veneris		-	424	Thus -	-	647
Sufinum oleum		-	392	Judæorum	281,	649
Sylphium	1		120	masculum, V. Olibanum	-	460
Symphytum			248	Thymbra -	-	588
maculof	um	AL STATE OF	525	Hispanica -		412
minus		The state of the s	520	Thymelæa -	-	648
Syriæ oleum			427	Thymiama	281,	649
	-			Thymus -	1	650
mah	T.	A Control of		citratus -	•	651
Tabacum	A COLUMN		441	maftichina -	-	412
Tacamahaca			631	Serpyllum -		651
Talcum	Estate III		631	Tiglia grana		539
Tamalapatra		1	406	Tilia -	To the last	652
Tamarindus	-	100	632	Tincar -	-	159
Tanacetum		1500	633	Tithymali -	-	652
Balfamit	a -	What she	13.1	Tormentilla -	-	654
hortense	MONTH STATE		131	Tota bona -		157
Tanafia			633	Tragacantha -	1	334
Tapíus barbatus	A STATE OF THE STA	Die State	659	Tragofelinum -	m · ·	501
		-	5	•	Trichom	anes

INDEX NOMINUM, &c.

	Trichomanes			Page	Veronica Becabunga			Page
	Trifolium acetofum			654	femina, V.	Clasina .	A	139
	aquaticum		1 -3 av	400	Teucrium		United By	281
	hepaticum		100	655	Vetonica	-		661
	Melilotus	-		341	Vincetoxicum	-		146
	paludofum	- Thursday		416	Vinum	1	the Wi	661
	Trigonella Fænum-gra		Will be	655	Viola	With the same	7101-20 10	662
	Trinitatis herba	- Cum		304	lutea			663
	Triffago, V. Chamædr			341	palustris		1 1 1 1 1 1 1	224
	paluftris		10 - 301	219	tricolor			504
	Triticum .	- 2	19-17-1	596	Violaria		100	664
		1	FI - IN	308			AL ASSESSMENT	663
	Turnethum		316-32	326	Vipera	15, 21		664
	Turpethum minerale	•	1.00	655	Viperaria	-	A-100M	597
		0.7	CALL SALE	105	Viperina	10	A STATE OF	602
	Tuffilago Petafites		Den alla	656	Virga aurea	- Lunca	172 - 1244	665
			-	495	Viride æris	407	- Capan	23
	Tutia	100	100	656	Vifcus		分为於 無額	666
		U.			Vitex Agnus castus . Vitis alba	-	1	27
	Ulmaria			6-0			1	165
	Ulmus	- Amai	The same	678	Vitriolum album		110000	667
		. Same	-	679	cæruleum		HE MENT	668
	Umbilicus marinus	- syligem	-	57	Romanum	-	- 34	668
	Ungula caballina	- 1, 4000	102	656	viride	-	A DESCRIPTION	669
	Uniones Urina	214.4		680	Vitrum antimonii	-		68
	Urinaria	2000				atum	41.85	76
		101111111111111111111111111111111111111	100	395	Vomica nux	-	aleased,	453
	Urfina radix	(19710)	NAME OF STREET	423	Vulgago		- 1 4 T 1 M 1	121
	Urtica		- 1	681	Vulvaria	1	213082	124
	mortua		- 1	375		377		
	Uvæ paffæ		-	681	W	W.		MANUEL .
	Uva urfi		1002031	682	Wanhom	-		313
2		37			Winteranus cortex	119	Charles Total	683
	Valeriene	v.		-		x.		
	Valeriana Celtica	F-0-18	Duti-Fill	657	Vulaslass	Δ.		-00
	Vanilla	-	F10	436	Xyloaloes			388
		45400	THE T	659		Z.		
	Venus Veratrum album	-	3 3 3 4	262	Zagarilla, V. Eleuth			-0-
			- Ellin	337	Zarza Zarza	eria	Will the state of	281
	nigrum	-	15 TH	339		-		586
	Verbafculum	- 2 64.30	10000	478	Zedoaria	MAN		684
	Verbafcum	a noid	oula dia	659	Zerumbeth	17.74	T STATE	684
	Verbena	-	6dans	660	Zibethum		165 165	685
	femina, V. E.	ryumum	SATELY	289	Zincum	15 15 10 10	1	685
	Vermicularis			352	Zingi	1 30	1 4 5 - Q.E.	63
	Vermis terrestris	100		401	Zingiber	FIRE S	1 2 20 3 1 5 1	687
*	Vernix		JING F FINE	363	Zizyphus	W. Mar	S. C. S. C. S.	360
	Veronica	- 5	NAT DE	660	TER SERVICE	and and an		

O F

ENGLISH NAMES.

Λ		-	Page			Page
ABSORBENT	earths	-	643	Amber -		623
Acacia -	-	- 200	10	Ambergris -		44
German	-	-	521	Amethyst -		259
Aconite	12	- 6	64, 689	Amianthus	A Thinks	632
Acorus, baftard		- 10	359	Ammoniac falt -	liste -	565
Adder	-	-	664	Ammoniacum gum -	-	47
Agallochum	-	(20th	388	Amomum -	-	49
Agaric		-	24	common -	-	50
of the oak	-	-	. 26	Anacardium -	-	54
Agnus-castus	-	- 100	27	Angelica -	-1	. 58
Agrimony	-	10114	28	Anime -	-	60
Hemp	-	- 10	290	Anife -	-	61
fweet-scented	1_	- 501	29	starry-beaded -	13 () ()	63
Water	-	211	290	Anodyne liquor of Hoffman	Contract to	675
Air, fixed	-	1-12	21	Antimony -	-	65
Alder	-	-	34	Ants		304
black	-	-	305	Aqua-fortis -		447
Alecost -	-	-	131	Aqua-regis -		448
Alehoof	-	-	336	Arabic gum -	THE STATE OF	332
Alexanders	-	-	343	Archangel -	1	375
Alkanet	-	270_23	56	Armenian bole -	-	155
Allheal	-	Accept - Second	482	Arrach, Stinking -	149	124
Allfpice	3-		507	Arfenic	100	113
Almonds	-	- Lotte No	51	Arfmart -	Friends 1	483
Aloes		DEATH PIN	35	biting -		484
Aloes-wood	-		388	Artichoke -		235
Alum	-	COUNTY OF BE	41	Asafetida -		120
plumous	-	S 12 20	632	Afarabacca -	State of the state of	121
independent of the life			5	A z		Ashtree

		1	Page				Page
Ashtree			306	Bezoar mineral			73
Ashes, Russia	-	-	556	occidental			150
Ashes of vegetables		-	403	oriental	-		149
Afparagus	-		123	of the ape		-	151
Avens	-	-	203	of the porcupi	ne	- 1	151
				Bile -	-		292
	B.			Birch			147
Bacher's pills	- 100	-	340	Birdfneft	-	-4-	271
Balaustines	-	-	130	Birdstongue		SUBSTITUTE OF THE	306
Baldmoney, V. Gentia	ın	-	319	Birthworts	-	-	111
Balm	-		417	Bishopsweed	-		47
of Gilead, herb		-	426	Bifmuth	-	1	152
fir		-	3	Biftort	-		153
Turkey		-	426	Bittergourd, V. Coloc	ynth		245
Balfam of Canada		-	3	Bitterfweet	-		609
of Copaiba	-		132	Blifters	-		189
of Gilead	- 100	-	466	Bloodstone	1 - 1 1 M		334
of Peru	•		133	Blue-stone	-	-	668
white of Peru		•	135	Bluebottle			266
of Tolu	-		135	Blue, Prussian	-		294
Barbadoes nut	-	-	539	Bois de Coiffi	*	12 N-111	528
tar	-	-	498	Bolar earths	-	-	155
Barberry		-	144	Bones		H11:08	254
Bark, Carribæan	-	-	495	Bonebinder	-	-	469
Jamaica		- 9 41	495	Borage	-	N 1 10	158
Peruvian	-	- 00000	485	Borax	-	+	159
red	- 1	A to Ministra	492	Box tree	-	The state of	168
Barley	a Vinda	to liver	310	Bran	-		310
caustic	-	- Peloine	217	Brankurfine		and the	11
Bafil	A- 17 100	120	457	Brafs	-		265
Bauldmony	-	- million	423	Bread	-		308
Bay			381	Briar, wild	-	Minister.	543
Cherry	-	-	380	Brignole plum	-	- VOIA	521
Bdellium	-	-	138	Brimstone	-	the man	625
Bean, Malaca	The State of the S	d-week	-54	Brooklime	- 100	A Marie	139
St. Ignatius's	A-Brill total	- 90	455	Broom -	-		317
Bears-breech	-	- (10.11)	11	Bruisewort	-	- 2.4	584
Bears-whortleberry		-	682	Bryony -	-		165
Bedftraw, Ladies	-	-	316	Buckbean	-	-	655
Beech, Sea-fide		-55014	495	Buckthorn	-		611
Bees	-	-devole	77	Bugle -	-	-	167
Beets	-	-150,000	145	Buglofs -	-	-	167
Ben nuts		ast anime	141	Bulgewater tree		- 19	320
Benit herb	- 1	-	203	Bullfist, V. Bovista	-		402
Benzoine		- 0100	141	Burdock	-		137
Bermudas berries			585	Burgundy pitch			639
Betony		-	146	Burnet-faxifrage			501
Paul's	-	-5470	660	Butchersbroom	-	- Bosse	546
Water	-	2 1800	598	Butter of antimony	-	-	73
Bezoar microcofmic	-	-tisonifes	151	Butterbur		All to the	495
		-	111			Butter	flower

OF ENGLISH NAMES.

			Dana				Desc
Butterflower			Page	Celandine			Page 224
			532 504	Celeri			
Butterwort			304	Centaury		138	79
	c.			greater			213
Cabbages			163	Ceruffe		1775	512
Cabbage bark			320	of antimon	v -		67
Cacao nut			169	Ceterach			217
Calamine	GARAGA .	8 .	170	Chalk			255
Calamints		77 197	171	Chamemel			220
Calambac	130		388	Chafte-tree			27
Calamus			173	Cheeferennet		-	316
Calamy	4.57	0.20	170	Cherries	2 2	MALE.	307
Calomel			109	Winter			30
Camels-hay			361	Cherry-bay			380
Cammock, V. Ononis			461	Chervil			218
Camomile		-	220	Chibou Gum	-nwavia	*	281
Rinking			223	Chickweed		-	39
Campeachy wood			390	China root		-	226
Camphor	-		179	Chocolate		-	169
of cinnamon		1011-01	239	Christmas-flower,	V. Black, Hell	eb	339
of peppermint	and the same	G 11 - 1	422	Cichory			227
Canella -	Park to		185	Cinnabar of antim	onv -	nelso.	74
Cantharides			188	factitions			102
Capficum		-	508	native			235
Caranna	-	1 1 1 and	191	Cinnamon			236
Caraway		Mills of	200	Cinquefoil	S G Live		482
Cardamom feeds			193	Citrons		3200	239
greater		-	194, 327	Citrul	A STATE OF THE STA		260
Cardinal-flower, blue			399	Civet		-	685
Carduus	-	-	195	Clary			344
Carline thiftle	-	2001	197	Clay			234
Carpobalfam		ebeletia.	199	Cleavers	100000	2	77
Carrot of Candy	Te rivers	No likh	271	Clotbur		213	137
wild	Waster !	W71 -	271	Clove bark		1	205
Cafcarilla	-	ICEO SE	281	Clove gilliflower		-	204
Cashew nut	-		55	Clove spice		-	201
Cafia fistula	would	WHEN THE	207	Cobalt	-	-	113
lignea	-	H-1	208	Cobweb	1	1	91
Cassidony, V. Steechas	-	14	284	Cochineal		1 12	240
Caftor	-	M-19	209	Coffee	-	Marie Du	243
Caftor oil	*	100	540	Colcothar	-	7-11	670
Cafumunar			211	Cold feeds		-23	261
Catmint	-	-	439	Caleworts		Mice	163
Cauliflower	-	1000	163		Soldanella	1-1	164
Caustic alkalies	-	-	558	Colophony		She is	639
barley	-	1	217	Coloquintida	-	- 1	245
Antimonial	-	-	73	Coltsfoot	-	-	656
Lunar	-	-0	92	Columbine	S	100	90
Cayenne pepper	-	-	508	Columbo	-		246
Cedar	-	-	212	Comfry			248
							Conessi

		D					
Coneffi bark			age 48		D.		Page
Confound, greater, V.	Comfry		48	Daify -			
middle	Cominy		67	Ox-eye		12 3 51	140
fmall				Damfons	5		140
Saracens			65	Dandelion			521
Contrayerva root				Danewort			172
Copaiba balfam			49	Dates -			576
Copal gum			32	Devils-dung, V. Afai	Catida		472
Copper			51 62	Dill -	etiua		120
				Dittander			57
Copper waters			89				384
Copperas Coral				Dittany of Crete		history	274
Coralline			51	white D1-			274
	114 115		51	Dock -		- 10	376
Coriander		The state of the s	52	four	-		12
Cornelian			59	Water	- GIRLING	- 2000	377
Cornflower, V. Cyanus			66	Dock-cresses		1	376
Cornrole	-		77	Dodder	-		286
Cottmary			31	Dog-rofe	-	The same	543
Costus root			54	Dogs-grafs		- TSS	326
Couch-grafs	- 0/1/10 423 410		26	Dogs-mercury	- asimi	DEE -10-	423
Couhage	-	- 11002	76	Dogstones, V. Orchis	-100010	adult-10 1	589
Counterpoison, V. Con			49	Dovesfoot		- 1	322
Monksh	ood, V. Ant	hora	64	Dragant gum	-	-tim	334
Cowitch	- 1000	- 2	76	Dragons		1.90	279
Cowflips	-	- 4	78	Dragons blood	-		577
of Jerufalem	-	- 5	25	Dropwort Hemlock		- 40	458
Crabs-claws	- 11	- 1	84	Dwale	**	His - chu	607
Crabs-eyes	-	- 1	84				
Cranefbill	-	- 3	22		E.		
Creme of tartar	- 1	- 6	34	Earths	-		80
Creffes	-	- 4	38	abforbent	-	010-11-0	643
Bank, V. Eryfim	um		89	aluminous	-	-	41
Dock			76	animal calcare	ous	ber a file	184
Sciatica	La Contract		85	not cale	careous	The same	254
Water	- in	ALCOHOL: NO	37	argillaceous	-	15	5, 234
Crowfoot			32	bolar		-1150	155
Cryftal			59	mineral calcare	eous	- 01100	255
Crystal mineral			46	vegetable		W. mai	403
Cubebs	N. Warrish		60	Earth Fullers	2000	2 1	234
Cuckow-flower			92	Japan	100		640
			18	Earths fealed		-313	155
Cuckowpint			60	Earthworms		1 3 3 1 1	401
Cucumber			61	Earwort	100		420
wild	Contractor of the		68	Eatons flyptic			671
Cummin				Eau de luce		Salar Sun S	564
Curaffoa apples			27	Eau de Rabel			673
Currents -		307,6			1		
Cutch			41	Eggs		La la constitución de la constit	470 262
Cyperus root		- 110 111 2	69	Elaterium		3	Elder
							Liuci

OF ENGLISH NAMES.

			Page		Page
Elder		-	575	G.	
Dwarf			576	Galangal -	- 313
Elecampane			285	English -	- 269
Elemi Gum		-	281	Galbanum -	- 314
Eleutheria bark			281	Gall of animals -	- 292
Elm			679	Gall-flones -	- 151
Emerald			259	Galls -	- 315
Emulsions			52	Gamboge -	- 316
Endive	CONTRACTOR		284	Garlic -	
Epfom falt			566	Garnet-stone -	- 32
Eryngo			289	Gas, fixed -	259
Eternal-flower, V. Sta	chae		284	Gaule -	21
Ether ,	chas	100	1000	Gelly of fruits -	- 434
nitrous		157	675		267, 308
	EG9199	PO YOU	450	horns	- 253
marine		1	573	Gentian	- 319
Ethiops antimonial	-	1	110	Indian -	- 320
mineral		15,00	101	Germander -	- 219
vegetable		Tisal	531	Water -	- 596
Euphorbium	- 100	1000	291	Gill-go-by-ground, V. Groundivy	336
Eyebright		1	292	Gilliflower -	- 204
				Ginger -	- 687
A Waller of the Control of the Contr	F.			Ginfeng root -	- 323
Fats	- 1	-	502	Glass of antimony -	- 68
Felwort, V. Gentian	- /	-	319	cerated	- 76
Fennel	-	-	302	Glaffwort -	- 364
Hogs	-	-	500	Glaubers falt -	- 567
Fenugreek	- 1000	-	304	Gold -	- 128
Fern, male	- 1300		300	Mofaic -	- 617
Feverfew	THE PERSON NAMED IN	-	414	Goldcup -	- 532
Figs		-	197	Golden rod -	- 665
Figwort		- 1	597	Goldilocks -	- 284
Finckle	-		303	Goofegrafs -	- 77
Fir tree	4 4 6 1 3 1 5	-	1	Gourd -	- 261
Balm-of-Gilead		-	3	bitter, V. Colocynth	- 245
Canada			3	Grains of paradife -	- 327
Fishglue		145	351	Grana-tilia -	
Fistick-nut, V. Pistaci	a		452	Grafs, Dogs -	- 539 - 326
Five-leaved grafs		-	482	Gromwell -	
Flag, Sweet	The same	200	173	Groundivy -	399
Flax, purging		10.07	398	Groundpine -	336
Fleawort	4	In s	522	Groundfel -	- 223
Flies, Spanish			188	Guaiacum -	- 287
Flower-de-luce		264		Gum ammoniacum -	- 329
Fluellin			357 280		- 47
Foxglove	THE CONTRACTOR	1		anime -	- 61
Frankincenfe		Merca	275	elemi -	332
Fruits, Summer		411.9	647		- 281
Fumitory		O(COS)	307	gambia -	- 366
Fullers earth		(Light	312	guaiacum -	- 329
Furze, Ground, V. On	onic	- Trail	234	hederæ -	- 336
raize, Grouna, v. On	OHIS	-5-49=	461	juniper -	- 363
					Gum

I N D E X

			Page				Page
Gum lac		- 1240	372	Houfeleek			599
red aftringent	-	_ Inger	366	Hartfickle, V. Cyanus	s	Que	266
fenica	200	1	333	Hyacinth stone		2000	259
tragacanth	-	-9200	334	Hypociftis			347
Gypfum	7-	DP39	600	Hyffop			347
				RIGHT STATE OF THE			A STATE OF
	H.				I.		
Hartshorn		- ayard	253	Jack-by-the-hedge	-	1,016	31
Hartstongue	44		396	Jalap	-	-	349
Hartwort		0208-19	603	Jamaica bark		SIA	495
of Marseille	S	- 1770	604	pepper		-	507
Hay Camels	-	- 71 9	361	James's wort, St. V.	Ragwort	The L	348
Heartseafe	- ati	12 30 V	664	Japan-earth		-	640
Hedgehyssop	07	10.1	328	Jerusalem cowslips		12(17)	525
Hedgemustard	- 1	- nau	289	oak	128 866	12/11/0	162
Hellebore black	17 3 11 3 1 A	natural second	339	fage	and Import	11200	525
white	The second second	200.000	337	Jefuits bark			485
Hellweed	-	-	286	Jews-pitch	Sour Singer		154
Helmet-flower	D. M. Manor	14-78	64	Ignatius's bean		Aure	455
Helvetius's powder	- 1	THE WOOD	43	Ilathera bark	-	-	281
flyptic		- 19	671	Indian-leaf		-	406
Hemlock		SCH SIL	228	pink		-	611
Dropwort	-42) - X4200	THE MAN	458	John's wort, St.	-	-	346
Water	AND THE	-	79	Ipecacoanha	Continu S	-	353
Hemp		12 FOREN	186	Irish slate			342
Water		of steel	290	Iron	-	12374	293
Hemp-Agrimony	THE WORLD	-	290	Ifinglass		-	351
Henbane		- Tarrella	345	mineral	T 4 (2) (6) (5)		600
Hepatica		- 45	341	Jujubes	-	-	360
Herb-benit	-	47.79	203	July flowers			204
Herb-of-Grace, V. G	ratiola	- Isoliy	328	Juniper	*-		361
Herb-mastich	-	1010	412	Ivy	-	-	335
Syrian	-	-	411				100
Herb-Robert	turisbury, sa	Spring	322	A SHARE THE SHARE	K.		
Herb-trinity	slinan	3-10 cm	341	Kelp		-	364
Hercules's allheal	-	all trees	482	Kermes	•		365
Hermodactyl	-	AFILT OF	342	mineral			71
Hipps	-	5-190E	543	Kernels of fruits		-	308
Hogs-fennel	5-	- 175	500	Kernelwort			598
Holly Sea .	- 219	still day	289	Keyfer's pills	The MICHELLINE	000	110
Holy Thiftle		The state of	195	Kino	50-0 H. H.		366
Honey	-	- minst	415	Kneeholly	-		546
Hops	- PARTIES	agmuna.	401				
Horehound	-	Separate .	410		L.		
Horestrong	-	910	500	Labdanum	7-		367
Horns	-	-	253	Lac Gum	Series of the se		372
Horsechestnut		AL THE	691	Ladies bedstraw	To the second		316
Horferadish		-	533	mantle			29
Horfetail		-	287	fmock	Towns of	-	192
Houndstongue	-	A PHIS	268	Lakeweed	-		484
							Larch

OF ENGLISH NAMES.

			Page				Page
Larch tree	1 2 3	silplin u. hip	638	Maidenhair, Canada	-	-	20
Lard	Marine .	The Contraction	502	English		1000	654
Laudanum		The state of	465	Malaca-bean	-	PALL SPECIAL	54
Lavender -	14 1	Station Laborate	378	Mallow	2 7 10	STATE OF THE PARTY	407
French		-12	619	Manganese	gol No		405
Lavender Cotton			5	Manna		Yair lane	408
Laurel		Acriedan 3	380	of Briançon	-	Tell Line	408
Spurge			648	Maple			11
Lazule-stone		ALL THE MAN	382	Marble	_	the areas	255
Lead			511	Marcafite	_	250W	527
Leaf, Indian	-		406	Marigold	-	WHILE WA	174
Lemnian Earth	0.0		155	Marjoram			405
Lemons			394	quild		A COLUMN	468
Lemon-thyme		Combined 18	651	Marle	-	die	255
Lentisk wood		ALC: WE	383	Marshmallow	_	. 14	40
Leopards bane	Sec.		277	Marshtrefoil or buckbea	n	D. 13.23	655
Lettuces		Barrier St.	373	Marvel of Peru		all play with the	349
Lignaloes		AR AND	388	Marum			411
Lily of the valley	of the party of	20084	393	Masterwort		1 littless	
May	Marie Land	A STATE OF THE PARTY OF THE PAR	393	Mastich		LABORA	353
Water	-	and the second	456	Maftich-wood			413
nubite -		. Land		Herb			383
Lime			392	Syrian Herb	T. ASS.	The state of	412
Lime stone	1	NE SECTION AND ADDRESS OF THE PARTY OF THE P	174 255	Maudlin		and the later of	411
Lime tree			652	May lily		N SELECT	131
Linden tree		The Paris	652	Mayweed		easts	393
Linden tree				Meadow-faffron			223
Liquidambar			396 398	Meadowsweet			244
Liquid-shell				Mechoacan		Percept 1	678
Liquorice		The Party	572	Melilot			414
Litharge			325 512	Mercury	1		416
Liverwort	-	and the second	The second second	Dogs			93
Ground		A Part Control	341	English Herb		and the	423
Iceland			387	French Herb	1		- 157
noble		THE ASSESSMENT	The second second	Mexico-feed		CHARLES TO	422
Logwood	BUILD	The Salles of Loc	341	Mexico tea		The 187 (1)	539
Lopez root	TATRO		390	Mezereon	00	(1,500 0 1)	162
Lovage		HE GO TO THE	531 385	Microcosmic bezoar			648
Lunar caustic			THE REAL PROPERTY.	falt		Total Sand	151
pills			92	Milfoil	1	19 Se 100 NO	680
			93	Milk	•	Tomas Trans	424
Lungwort Lye of tartar		STATE OF THE STATE	525	Milkwort		Santa Maria	369
Lyes, Soap			556	Millepedes	- Billian	1874 T	517
Lycs, bomp		1914 A	558	Millmountain			425
	M.	A Change		Miltwafte		1	398
Mace			100				217
Oil of	131		402	Mindererus's spirit		osiswamine	18
Madder	1		452	Mineral anodyne liquor Mineral waters		SHATTE TAB	675
Magnefia	THE ST	STATE OF THE STATE	545	Mint Waters	Selfie.	THE STATE OF	84
Maidenhair			403			Jesidon	418
2vialdennali	118		19	Pepper .		1980	421
			19	5 B			Mint,

			Page				Page
Mint, Water -	25.10			Nitre fixt or alkalized		1200	444
Miffeltoe -			666	volatile			448
Mithridate-mustard -				Nitrous acid		- 10 16	447
Mollipuff, V. Lycoperdon	n -		402	dulcified			449
Monkshood, V. Anthora			64	Nitrous ether		-	450
Monk's rhubarb -			378	Nutmeg	100		450
Moneywort -			450	Nut, Barbadoes	TO LAND		539
Mofaic gold -			617	Pistachio	120 15 113		452
Mother-of-thyme -			651	purging		Section 1	539
Motherwort -		200	194	Vomic	To the state of		453
Moufear -	TO THE REAL PROPERTY.		500				733
Moxa -			429		0.		
Mugwort .			116	Oak			530
Mulberries			307	of Jerufalem		The B	162
Mullein			659	Sea			530
Mufcovy-glafs -		- COLUMN	600	Oats		200	310
Mushroom, dusty, V. Ly	conerdon		402	Ochre		200	457
Muík			427	Oil animal			560
Musk-cranesbill			322	of bays		1	381
Musk-cyanus		214.95	267	of mace	MINIS		452
Muskseed			137	olive			459
Mustard		The state of the s	606	British		THE STATE OF	497
Hedge			289	Caftor		-	540
Mithridate		300	647	Palm		1988	472
~ ,			647	Rock			496
7. 11			430	Olibanum	11070178	1	460
Myrrh		1,000	431	Olives		- 45	459
The second secon	The state of the s	THE CAME	433	Olive, Spurge		ALL SE	648
Dutch		3419	434	Onion		44 2 1	214
Daile			тэт	- Sea	350		593
	N.			Opium	10000	74 123	461
Nard, Celtic		907	436	Opobalfamum	THE PARTY	- 1	466
Indian	THE STATE OF		436	Opopanax	11.		467
Navew	THE PARTY NAMED IN		435	Oppodeldoch	THE PROPERTY OF		582
qwild	THE SH	E S	435	Orache, flinking			124
Nep		-	439			A STATE OF	125
Nephritic wood		-	440				127
Nettle		3 97	681	Orchis	All years		589
dead	The same of	100	375	Origany,	STATE OF		468
Nightshade		2/14	607		-	110-00	116
deadly			607	THE RESERVE OF THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER.			637
Garden			608				357
Woody			609			SHAPE.	469
Ninzin		-	324				140
Nipplewort	E		376				469
Nitre		- 94	443	A STATE OF THE PARTY OF THE PAR			- California
ammoniacal		13278	448		P.		
calcareous	The latest states	-	449	Contract to the contract of th			478
	ly fo called	121711	567				539
cubical	-		448	With the state of			472
	The second						Panfies

OF ENGLISH NAMES.

			-				
D C			Page	D1-4		-	Page
Panfies Panfies	AUTO AND THE		The state of the s	Plaster-of-Paris		-	599
Pareira brava	-1220 (0.51 76		479	Plums		-	521
Parfley	-	- 1800	498	Poley-mountain	The state of the s	-	516
Macedonian		- 11 5	499	Polypody	-	-	519
Parinep		- 991 6 63	481	Pomegranate		-	327
Pafque flower	- A Republic	+ Town	525	Poor-man's pepper	-	-7.110	384
Patience		- 41003	378	Poplar	-	-	519
Paul's betony	-	-明明日	660	Рорру		46	1, 474
Peach		• 10000	483	wild	-	-	477
Peagle	-	- 10000	478	Potafh		-	556
Pearl	- 150	-	410	Primrofe		-	478
Pearl-ashes	-	-	557	Prunelloes	*	-	521
Pearl-barley	-	- 1000	310	Prunes	-	-	521
Pedro del porco	-	-	151	Pruffian blue		29	294
Pellitory of Spain	-	-	526	Puffball	-	-	402
of the wall	I Stillen	-	480	Pumpion			261
bastard	- The section		523	Purging-flax		-	398
Pennyroyal		170-10	524	nuts	200		539
Peony			470	falt			566
		Palain		waters			
Pepper Guinea			505	Pyrites			85
			508	Tyrries			527
Jamaica		- Milbred	507		•		
long		-	506	0	Q.		
Poor-man's		- 200	384	Quaffy		-44	528
Wall	· 16 15 107	-	352	Queen of the meadow	5 -	-	678
Water	-	- 101/19	484	Quickgrass		-	326
aubite	-	- Intellige	505	Quicklime	-	-	175
Peppermint	The Park	- Die	421	Quickfilver	(A)	- 10	93
Pepperwort		-	384	Quince	-	-	267
Peruvian bark	-	-111100	485		2 00		
red	-		492		R.		
Pestilentwort	- 10000 1 10	- 10	495	Ragwort	5	-	348
Petrefactions		- 14 83	255	Raifins	-		681
Petroleum	10000000	- 400	496	Rape		- 4	435
Petty-whin, V. Ononis		-	461	Rafpherries	Carrier St.		307
Pilewort	-Conducted S		226	Rattlefnake root	The Transport		517
Pimento			507	Realgar	200		116
Pimpernel			55	Reddle	The state of the s	30.18	
Water		1 12 1		Regulus of antimony		1	457 66
		-	139		martial	. Days	
Pine tree			639		Rellated		72
Pine-thiftle		THE RES	198		itenated		72
Pink, Indian	(a) helputytil		611	Refin		-	639
Piony		- (1757)	470	Reftharrow		-	461
Pismire	-	-	304	Rhapontic	A STATE OF THE STA	-	537
Piffabed, V. Dandelion		#8449B	272	Rhodium wood		-	391
Pistachio nut	- saling	30000	452	Rhubarb		- 140	535
Pitch	TO USE OF SEC	-2100	510	Monks		- 18	378
Burgundy		-	639	Ribwort	4- 11- 8	-	511
Jews	-	-016/19	154	Rice	Eps	-tgin	309
tree	- Mys	- mongo	1	Rifigal	- NEEDIN	- 11	116
Plantane	-	. 31	511	Rochelle falt		- 016	636
			5 B		8	I	Rocket
				MARKET BUT THE REAL PROPERTY.			

		Page			Page
Rocket -	12 10	288	Sandarach of the Arabians	4	363
Rock-oil -	- /	496	of the Greeks	07219	116
Rose, Damask -	-	541	Sanicle -	2	577
Dog -	- 3	543	Yorksbire -	Wall.	504
red -	Wat he	542	Sapphire -	14	259
Rofemary -	2000	544	Saracens confound -	- 4-1	665
Roferoot -	-	637	Sarcocol -	1.100	586
Rofewood -	-	391	Sarfaparilla -	1160	586
Rofin -	Vision.	639	Saffafras -	-	587
Ruddle -	-	437	Satyrion -	-	589
Rue -	-	547	Sauce-alone -	-	31
Rush, fweet -	-	361	Savin -	- dela	548
Ruffia ashes -	-	556	Savoury -	22/6/9	588
			Savoy		163
S.			Saunders woods -	12 00	578
Safflower -		199	Saxifrage, Burnet -		501
Saffron -		257	English -	114	591
baftard, V. Safflower	-	199	Meadow -	1	591
Meadow -	5.	244	white -		590
Sagapenum -		551	Scammony -		591
Sage -		574	Sciatica-creffes -	Man /	385
of Bethlehem -		525	Scordium -	1012	596
of Jerufalem -	-	525	Scorzonera -	-	
of virtue		574	Scurvygrafs -	LIEN P	597
Sago - Milan	ore in		Scotch, V. Soldanella	11.2 1	164
St. James's wort -		473	Sea-holly -	1	289
St. Ignatius's bean		348	Sea-mofs -		and the second second
St. John's wort -	1	455	Sea-oak -		251
Salep -		346 589	Sea-onion -		530
Sal-ammoniac -			Sea-water -		593 86
	1	565	Sea-water -		
Sal-gem - Sal-volatile -	-		Sealed earths -		530
Salts in water		563 81	Sebestens -		155
	1		Sedative falt		360
alkaline fixt - mineral	- The	553		-	160
		438	Sedge -		359
vegetable	-	553	Seggrum, V. Jacobæa		348
cauftic	Mag !	558	Sel de Seignette -		636
alkaline volatile		559	Selenite -	-	599
cauftic	- 4	562	Self heal	-	520
ammoniacal -		565	Sena -	-	600
cathartic -	1	566	Seneca gum	-	333
common -		568	Senegaw rattlesnake root		517
diuretic -	-	18	Sengreen -	150 0	599
microcofmic -	- 14	680	Septfoil -		654
muriatic calcareous	-	572	Sermountain -		603
Rochelle	5 56	636	Shepherd's purfe -	1 .	168
fea .	-	569	Silefian earth	-	155
fedative -	1	160	Silver -	-	92
Seignette's	-	636	Silverweed -	12 37	91
Saltpetre	-//	443	Simaruba bark -	4.4916	605
Saltwort		364	Skink -	- 70	595
					Slate,

OF ENGLISH NAMES.

			Page				Page
Slate, Irish	-	Was lift	342	Stonecrop	-		352
Slaters, V. Millepedes		75	425	Stone-parfley, baftare	d		. 50
Sloes	CHARLES OF PARTY	-	521	Stones, precious	7	-	259
Smallage	de la	STATE OF A STATE OF	79	Storax		10 1 - 5	620
Snails	4-194	-	394	Liquid		19 19 20	622
Snakeroot	L. Penny	- 11	602	white			135
Snakeweed		-	153	Strawberries		-	307
Snakewood		-	454	Sublimate corrofive	-	1500	106
Sneezewort	-	15.070	523	Succory		Delivery	227
Soap	Linchashas	-	580	Suet	-	0 114	502
Soaps, Volatile	-	CARLES	583	Sugar			549
Soap-lyes	- 23		558	of milk	1753370	DO TO	371
Soapherries		2	585	of lead	1	a supplied to	513
Soapwort		Samuel Samuel	584	Maple		S Spanis	11
Soldanella		-	164	Sulphur	-		625
Solomon's feal		1000	604	Sulphurwort		4 1926	500
Soot		-	312	Sultan-flower		To Amelia	267
Sorrel	Law Own	ALEN A	12	Sumach	to the fi	1 1 1 1	630
Wood	Transfer of	1000000	400	Swallowwort			661
Soude		1000	364	Sweet-rush			361
Soude blanche		The state of	438	Sweet-fultan			267
Sour-dock			12	Sweet willow		THE STREET	
Southernwood				Sycamore	The state of		434
Sowbread	BEET STELL	A CHILL	41	Sycamore		Delta Series	11
	TO SELECT		117		T		Town fell
Spanish slies		-		Theometica	T.		4.
Spar	15 7 7 1 1 1		255	Tacamahaca	000		631
Spearmint	THE TEN	1	418	Talc			631
Spearwort F.	THE REAL PROPERTY.	1	533	English	STATE OF		600
Speedwell, Female	THE REAL PROPERTY.		280	Tamarind			632
Male		To the second	660	Tamepoifon			661
Mountain	S N LA		661	Tanfy		9	633
Spermaceti		100	610	wild		Wan a	91
Spignel			423	Tar		44	509
Spike		-	380	Barbadoes			498
Spikenard		THE YELL	436	Tartar			634
Spirits, Vinous			612	emetic			74
Volatile			563	regenerated			18
Spleenwort	- 17116015	-	217	foluble	-	-	635
Sponge			615	vitriolated	-	- N-	677
Spunk	-	1	26	Tea	-	-	645
Spurges	-	-	652	German			660
Spurge-flax	-	-	648	Mexico		-	162
Spurge-laurel	-	-	648	Thistle, Carline	-	-	197
Spurge-olive	-	-,	648	holy	-	-	195
Squill	-		593	Pine		-	198
Squinanth		-	361	Thorn-apple	-	-	619
Starch		-	309	black		-	521
Stavefacre		-	618	Thyme	-	100	650
Stechas	-	- 1.4	619	Lemon	-		651
yellow	-	-	284	Mother of	-	-	651
Steel	-	4 6	294	Tin	-	-	616
							Tincal

INDEX OF ENGLISH NAMES.

		Page			Page
Tincal	-	- 159	Waters, common -	Desar.	80
Tinglass	-	- 152	Mineral -	1 Es	84
Toadflax		- 395	alkaline	- 14	84
Tobacco	-	- 441	chalybeate		87
Tobacco-pipe clay	-	- 234	cupreous	-	89
Tormentil		- 654	purging	-	85
Touchwood		- 26	fea	-	86
Tragacanth	-	- 334	fulphureous		90
Treacle-mustard		- 647	Water-creffes -	Acres 14	437
Trefoil, Marsh	-	- 655	Water-germander -	-	596
Trinity-herb	-	- 341	Water-flag		359
Tunhoof, V. Ground	ivy	- 336	Water-hemp -		290
Turbith mineral		- 105	Water-lily -	-000000	456
Turbith root	-	- 655	Water-pepper -	- 110	484
Turmerick	-	- 265	Wax -		215
Turnep		- `534	Wheat -	1.00	310
Turpentines		- 637	Whey -		371
Tutenag		- 685	Whin, Petty, V. Ononis		461
Tutty		- 656	Whortleberry, Bears	4 2 34	682
		Day May 5	Widow-wail -	-	648
	v.		Willow -		573
Valerian		- 657	faveet -	-	434
Vanelloes		- 659	Wine -		662
Verdegris		- 23	Winter-cherries -		30
Vermilion		- 102	Winter's bark -		683
Vervain		- 660	Wolfsbane -		64
Vine, wild	100	- 165	blue -		689
Vinegar		- 13	Woodlice -		425
Violets		- 663	Wood-forrel - 4		400
Viper		- 664	Wormbark -		320
Vipers-grafs		- 597	Wormfeed -		579
Virgins-bower, uprigi	ht	397	white, V. Coralline		251
Virgins-milk		- 143	Wormwood -		6
Vitriol, blue		- 668	Mountain -		9
green		- 669	of Valais		9
Roman		- 668	Roman -		9
white		- 667	Sea -	THE STREET	8
Vitriolic acid	120	- 671	Wort -	THE STATE	311
volatile		- 672	Woundwort -	100	482
dulcifie		- 674	Wrack, Sea	7112	530
Vitriolated falts	u	- 677	Track, oes		330
Vomic nut	1		Υ.		
y omic nut	200	453	Yarrow	No. of the	424
Urine	FILE	- 680			17
Office	TO S	000	Z.	Series à	
	w.	MEL	Zarnich	137	116
Wakerobin		- 118	Zedoary	PART DA	684
Wallflower	350	- 224	Zerumbeth -		684
		THE RESERVE THE PARTY OF THE PA	Zinc	100	685
Wallpepper	H-USD-	352		The state of the	1100

BOOKS in ANATOMY, MEDICINE, SURGERY, and NATURAL PHILOSOPHY, printed for J. JOHNSON, No. 72, St. Paul's Church-Yard.

- AIKIN on the Use and various Preparations of Lead, with Remarks on Goulard, 2d edit. 1s 6d Thoughts on Hospitals, 1s 6d - Heads of Chemistry, 1s 6d Translation of Beaume's Manual of Chemistry, with Notes, 3s 6d sewed, 4s bound Oeconomiæ Animalis delineatio; in usum Juventutis Medicæ, 15 6d Sketch of the Animal Œconomy, 1s Biographical Memoirs of Medicine in Great Britain, from the Revival of Literature to the Time of Harvey, 4s fewed 8 Alanfon on Amputation, and the After-Treatment, with an Account of the Amputation above the Ancle, with a Flap, 2d edition, greatly enlarged, 6s bound 9 Curry on the Nature of Fevers; on the Caufe of their becoming fo frequently Mortal; and on the Means to prevent it, 1s 6d 10 Dawfon's Cases in the Acute Rheumatism and Gout, with the Method of Treatment, 6s bound 11 - Account of a fafe and efficacious Medicine in Sore Eyes and Eye-lids, 1s 22 Else on the Hydrocele, and other Works; with some new Cases of the Hydrocele, and a Comparison of the different Methods of treating it by Causlic and Seton. By G. Vaux, 2s. 6d. fewed 13 Elliot's Physiological Observations, 15 6d - Medical Pocket-Book; containing a short Account of the Symptoms, Causes, and Methods of Cure, of the Diseases incident to the Human Body, 2s 6d sewed 15 - Account of the Medicinal Virtues of the principal Mineral Waters in Great Britain and Ireland, and on the Continent, with the Method of Impregnating Water with Fixed Air, invented by Dr. Priestley, and improved by others, 3s fewed 16 —— Elements of the Branches of Natural Philosophy connected with Medicine, viz. Chemistry, Optics, Sound, Hydroftatics, Electricity and Physiology. With Bergman's Tables of Elective Attractions, 5s fewed 17 ---- Philosophical Observations on Vision and Hearing, &c. 3s 6d sewed 18 Fothergill's Works, with Memoirs of his Life, by Dr. Elliot, 6s fewed 19 Fordyce's (Geo.) Practice of Physic, 5s fewed 20 - Elements of Agriculture, a Syllabus of his Chemical Lectures, 2s 6d fewed 21 Falconer's Observations and Experiments on the Poison of Copper, 2s sewed 22 Hunter's (Dr. W.) 34 Plates of the Gravid Uterus, with Explanations in Latin and English. The few Copies of this fuperb Work that remained unfold at the Author's Death, may be had at 31 138 6d which is 21 128 6d under the Original Price 23 - Medical Commentaries, 4to. 6s boards 24 Hunter's (John) Natural History, Anatomy, and Diseases of the Human Teeth, with 16 copperplates, 4to. 11 16 bound 25 Hewson's Experimental Enquiries, in Two Parts. Part 1. On the Blood. Part 2. On the Lymphatics, 2 vols. 8s in boards. - The First Part, on the Blood, may be had alone, 3s fewed 26 Hoffman's Practice of Physic, with a great Number of Cases, translated by Dr. W. Lewis, and revised by Dr. Duncan, 2 vols. 8vo. 12s in boards 27 Henry's Experiments and Observations on the Preparation, Calcination, and Medicinal Uses of Magnefia Alba: On the Solvent Powers of Quicklime: On Absorbents: On the Antifeptic Powers of Vegetable Infusions prepared with Lime, &c. On the sweetening
- 28 Henry's Medicinal Virtues of Magnefia, 6d 29 — Method of preserving Water at Sea, 28

Properties of fixed Air, &c. 2s 6d fewed

BOOKS printed for J. JOHNSON.

- 30 Henry's Memoirs of the Life of Haller, 2s 6d fewed
- 31 Hippocrates upon Air, Water, and Situation; upon Epidemical Diseases and Prognostics, translated by Clifton, 5s bound
- 32 London Medical Journal for 1781, 1782, 3 vol. 18s -- N.B. This Work is continued in Numbers, Quarterly, price 18 6d each
- 33 Lavoisier's Essays: containing a History of Discoveries relating to Air, &c. with Original Experiments, translated from the French by T. Henry, F. R. S. with Notes, 7s bound
- 34 Lewis's Abridgment of the Medical Essays, published by a Society in Edinburgh, 2 vol. 8vo. 10s 6d bound
- 35 Monro's (Alex. jun.) Observations on the Structure and Functions of the Nervous System, with 55 copper-plate Tables, folio, 21 12s 6d in boards
- 36 Motherby's New Medical Dictionary, or General Repository of Physic: containing an Explanation of the Terms, and an Account of the Discoveries and Improvements in Anatomy, Physiology, Physic, Surgery, Materia Medica, Pharmacy, &c. with copper-plates, folio
- 37 Moss on the Management of Children and Lying-in Women, 8vo. 6s bound
- 38 Mills on the Management and Diseases of Cattle, 7s bound
- 39 Priestley's History of Electricity, 4to. 4th edit. 11 1s bound
- 40 Introduction to the Study of Electricity, 2s 6d
- 41 Experiments and Observations on different Kinds of Air, 5 vol. 8vo. 11 10s in boards
- 42 Pott's Works, complete, 3 vol. 8vo. 11 4s bound
- Aftringents and Bitters. 4 On the Use and Operation of Blisters. 5 On Water. 6 On Inoculation. 7 On the Columbo Root. 8 On the Orchis Root. 9 On the Waters of Buxton and Matlock in Derbyshire. 10 On the Medicinal Uses of Fixed Air. 11 On the antiseptic and sweetening Powers, and on the Varieties of factitions Air. 12 On the Noxious Vapours of Charcoal. 13 On the Atrabilis. 14 On Sea Salt. 15 On Coffee. 16 On the proportional Mortality of the Small-Pox and Measles. 17 On the Solution of Stones of the Urinary and Gall Bladder, by Water impregnated with Fixed Air. 18 On the internal Regulation of Hospitals. 19 On the Action of different Manures. 20 On the Properties of different Absorbents. To which are added, select Histories of Diseases with Remarks, and Proposals for establishing more accurate and comprehensive Bills of Mortality, 3 vol. 8vo. 14s sewed.—N. B. The third Volume may be had alone, by those who are already possessed.
- 44 Percival's Observations and Experiments on the Poison of Lead, 2s
- 45 Park's Account of a new Method of treating Diseases of the Joints of the Knee and Elbow. In a Letter to Mr. Pott, 1s 6d
- 46 Sims's Observations on Epidemic Disorders, with Remarks on Nervous and Malignant Fevers, second edit. 5s bound
- 47 Sims's Discourse on the best Method of prosecuting Medical Enquiries, read before the Medical Society, second edit. 2s
- 48 Saunders on the superior Efficacy of the Red Peruvian Bark in the Cure of Agues and other Fevers, 4th edit. 2s 6d
- 49 Scheele's Experiments on Air and Fire, with an Introduction by Bergman: translated by Dr. Forster, with Notes by Mr. Kirwan, and a Letter from Dr. Priestley, 3s 6d sewed
- 50 Theobald's Dispensatory, compiled for the Use of the Army, 3s bound
- 51 Withers's Observations on the Abuse of Medicine: in Seven Chapters. 1 On Blood-letting.
 2 Emetics and Purgatives. 3 Sudorifics. 4 Blisters. 5 Stimulants. 6 Sedatives.
 7 Tonics, or Strengtheners. Dedicated to Dr. Cullen, 5s bound
- Wooffendale's Practical Observations on the Human Teeth, wherein the various Causes of their Diseases, Decay, and Loss, are enumerated, and the Methods of preventing, and of relieving them, are pointed out, 3s 6d boards

