

Observations on the superior efficacy of the red Peruvian bark, in the cure of agues and other fevers : Interspersed with occasional remarks on the treatment of other diseases by the same remedy.

Contributors

Saunders, William, 1743-1817.
University of Leeds. Library

Publication/Creation

London : J. Johnson; [etc., etc.], 1782.

Persistent URL

<https://wellcomecollection.org/works/cfxj2e9a>

Provider

Leeds University Archive

License and attribution


This material has been provided by This material has been provided by The University of Leeds Library. The original may be consulted at The University of Leeds Library. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



Leeds-Infirmary

MEDICAL-LIBRARY.

Entered *Jan^y 8th* 1783

Allowed for reading } Weeks Days
the first Year. } 1

—After the first Year

Forfeiture per Day for keeping } d. 2
it beyond the Time. }

LEEDS UNIVERSITY LIBRARY

Classmark:

Special Collections

Health Sciences Historical Collection

SC2 SAU



30106016200809


*The University Library
Leeds*



*The Library of the
School of Medicine*








E
89

OBSERVATIONS

ON THE
SUPERIOR EFFICACY
OF THE
RED PERUVIAN BARK,
IN THE
CURE of AGUES and other FEVERS.

INTERSPERSED WITH
OCCASIONAL REMARKS on the TREATMENT of
other DISEASES by the same REMEDY.

SECOND EDITION,
WITH CONSIDERABLE ADDITIONS.



By WILLIAM SAUNDERS, M. D.
Member of the Royal College of Physicians in London,
and Physician to GUY'S Hospital.

ÆQUE PAUPERIBUS PRODEST, LOCUPLETIBUS ÆQUE.
HORATIUS.

L O N D O N :

Printed for J. JOHNSON, St. Paul's Church Yard,
J. MURRAY, Fleet-street; J. SEWELL, Cornhill,
and HAWKINS, High Street, Southwark.

M D C C L X X X I I.

276.22

ROBERT THOMSON, M.D.

JAMES HERVEY, M.D.

Fellow of the Royal College of Physicians
of Guy's Hospital

Mr. JOSEPH W. ... F.R.S.

Mr. JAMES ...

Mr. WILLIAM ...

Secretary of the Hospital

GENTLEMEN,

PERMIT me to dedicate these Observations to you, as a Mark of my Devotion to you, and high Opinion of your Professional Character.

I have the Honour to be

Your most obedient

Humble Servant,

W.

SAUNDERS

Physician

T O

ROBERT THOMLINSON, M. D.

JAMES HERVEY, M. D.

FELLOWS of the ROYAL COLLEGE, and PHYSICIANS
of GUY'S HOSPITAL.

Mr. JOSEPH WARNER, F. R. S.

Mr. JAMES FRANK,

Mr. WILLIAM LUCAS,

SURGEONS of GUY'S HOSPITAL.

GENTLEMEN,

PERMIT me to dedicate these Observations to you, as a Mark of my sincere Esteem, and high Opinion of your PROFESSIONAL CHARACTERS.

I have the Honour to be,

GENTLEMEN,

Your most obedient,

Humble Servant,

W. SAUNDERS.

Jeffries Square, July 20, 1782.

MEDICAL LECTURES at GUY'S HOSPITAL.

IN the beginning of *October*, Dr. SAUNDERS, PHYSICIAN to GUY'S HOSPITAL, will begin his LECTURES, as usual, on the THEORY and PRACTICE of MEDICINE, in which the chief Objects of Discussion will be the *History, Diagnosticks, Nature, and Cure of Diseases.*

At the same Place, and about the same Time, Dr. KEIR, PHYSICIAN to St. THOMAS'S HOSPITAL, will begin his LECTURES on CHEMISTRY, in which, while the Principles of that Science are taught by Reasoning and Experiment, the Practice of it may be learned from the numerous Processes carried on in the Laboratory of the Hospital.

Dr. KEIR will also begin his LECTURES on the MATERIA MEDICA, about the same Time.

CLINICAL LECTURES are likewise delivered at GUY'S HOSPITAL, both by Dr. SAUNDERS, and Dr. KEIR, on the Cases of their Patients in GUY'S and St. THOMAS'S HOSPITALS; by Means of which the Student may enjoy every Advantage that the Practice of both Hospitals can afford, in the Study of his Profession.

Each of these Courses continue *Four Months*; they are all carried on at the same Time, and are repeated *Three Times* in the Year, beginning in *October, February, and June.*

To accommodate these Lectures to the Gentlemen who attend the Practice of the Hospitals, Care is taken that they shall not encroach on the Hours appropriated to the Transactions of the Hospital Business.

INTRODUCTION.

I AM encouraged to undertake a Second Edition of this Treatise on the Red Bark, from the favourable manner in which the first was received by the Public, and from the rapid Sale of a large impression in a few weeks. This Edition contains still more ample testimony, in favour of the doctrines and opinions I had laid down in the former. Such is now the evidence of the infallibility of the Red Bark in Intermittent Fevers, as to make it a subject of the first importance to the interests and happiness of mankind.

The

The Reader will observe, that I have added considerably to this Edition, especially on the subjects of Complicated Intermittents, and the Acute Rheumatism, from repeated and attentive observation.

I am happy in being able to assure my Readers, that they may be liberally supplied with the Red Bark, and I hope its advanced price, will not discourage them from its use. It is a cheap article, when we consider that it need only be employed in half the quantity generally used of the Pale Bark.

*There is some danger, from the avarice of Dealers, of its being adulterated, more especially, however, in the form of powder, a circumstance which may bring it into discredit, I have therefore taken some pains in collecting specimens of it, and arranging them in the order of their goodness, chiefly for the benefit of the young Gentlemen who attend my Lectures. The many applications to me on
this*

this subject, rendered this plan necessary, the Red Bark being hitherto little known by Apothecaries in the country. The efficacy of this remedy is so well established, that it is a matter of very little consequence, if Botanists should suppose that there are not yet Data sufficient to determine whether it be from the same species or not, with the Cinchona Officinalis of LINNÆUS. My friend, Dr. HOPE, Professor of Botany, in the University of Edinburgh, in a Letter to me, says,

*“ If this Red Bark, which seems to be more
 “ efficacious than the Common Bark, can be
 “ had; that is the great point, and whe-
 “ ther it be, or be not, the Cinchona Offi-
 “ cinalis of LINNÆUS, is but a secondary,
 “ and much less important consideration;
 “ however, I think your opinion the most
 “ probable; your Treatise will, I think,
 “ have the effect of making the public attend
 “ to this matter, and of rendering the use
 “ of it more general.”*

Since

Since the following pages went to the press, I have seen some exceeding good Red Bark, imported by a Spanish Merchant, a considerable part of which was as small as the Quilled Bark in common use, but it still preserved its Redness in that form, approaching, however, more to the colour of Cinnamon; it was evidently more compact and heavy than common Quill Bark, and appeared extremely resinous, its exterior coat, thin, whitish, and rough; it broke brittle, and gave evident proofs of its being the Quill of the larger Red Bark, which was in the same Chest: this, however, does not render my opinion less probable, that it may be only a variety of the *Cinchona Officinalis*, depending on AGE, SOIL or EXPOSURE.

OF THE
GENERAL HISTORY
OF THE
RED BARK.

I HAD long suspected that the Peruvian Bark in common use, was very inferior in power and efficacy to that recommended by the early writers on the subject; but more especially by our countrymen, MORTON and SYDENHAM, in whose works the medical virtues of this drug, in Intermittent and other Fevers are extolled as little short of infallibility. In their time the Quill Bark, (at least such as is now in use), was not mentioned;

B

their

their cotemporary writers on the *Materia Medica*, evidently describe the Peruvian Bark of that period, as of a larger kind, of more compact pieces, and of the colour of the rust of iron, which marks are very expressive of the Red Bark; the innermost coat of which has an ochrey appearance, and its resinous or middle layer resembles very much the *Lapis Hæmatitis*.* M. LA CONDAMINE expressed his surprise, when he was told by Mr. THOMAS BLACHYNDEN, Director of the English South Sea Company, at Panama, that the writers on Pharmacy and *Materia Medica* in England had preferred the Small and Quill Bark, while the inhabitants of New Spain, held the

* Cortex crassitie mediocri, foras scaber, paululum canescens, aliquando musco obductus, intus lævis, coloris ferruginei, saporis acris et intensè amari. Autumno colligitur et cortex circumcirca delibratur, tam trunci quam ramorum.

Dalei Pharmacologia, p. 201. Anno 1692.

the larger Bark in higher estimation. * I can only explain such a preference from this circumstance, that the larger pieces of Bark, which were imported into this country along with the smaller Bark, were either of a fibrous, spongy, or ligneous texture; or probably, damaged by moisture, and taken from the decayed trees.

B 2 Druggists

* They commonly reckon three species of the *Quina*, tho' some make four; the white, the yellow, and the red: I was told at *Loxa*, that these three kinds differ in their virtue only; the white having scarce any virtue, and the red excelling the yellow.

The trees from whence the first Bark was taken, which were very large, are all dead, having been entirely stripped, which infallibly kills them when they come to be old. Experience has shewn, that stripping kills some of the young ones also, but the greatest part escape. For this operation they use a common knife, which they hold in both hands; the barker sticks it into the Bark as high as he can reach, and so draws it downwards as low as he can. It does not appear that the

Druggists have sometimes suggested this as an argument against the use of the Red Bark, and many probably endeavour to oppose its introduction, until the quantity in the markets be greater, and more equally divided among them, or until they have been able to dispose of the large quantity of common Bark they have in their warehouses.

The

the trees which grow where the old ones stood, have less virtue than they, the situation and soil being the same; the difference, if there be any, may arise, perhaps, from the different ages of the trees. Few but young ones are now to be met with: I do not remember to have seen any much thicker than my arm, or above 12 or 15 feet high; those which are cut young, shoot forth new branches from the stumps.

I was informed at *Loxa*, that heretofore they preferred the coarsest Bark, and laid it by as a rarity, but now the finest is most esteemed: The merchants may possibly find their account in it, as it takes up less room in packing. But a Director of the English South Sea Company at *Panama*, through which all
the

The taste and flavour of the Red Bark is more difficultly evolved, and is therefore at first not so obvious from the closeness of its texture, and from the resinous coat being so well defended and inclosed between two other layers. It is evidently heavier than any other kind of Bark, and seems to have been prepared and dried with greater attention, its original appearance and form being better preserved.

B 3

I think

the *Quinquina* that comes to Europe must pass, assured me, that the preference given at present to the fine Bark, is in consequence of several chymical analyses and experiments which have been made on both sorts in England. It seems probable, that the difficulty of thoroughly drying the large coarse Bark, and the humidity it is naturally apt to contract and retain, has helped to bring it into disrepute. Vulgar prejudice will have it, that to lose nothing of its virtue, the tree should be barked in the moon's decrease, and on the east side. These circumstances, as also its being gathered on the mountain of *Cajanuma*, were certified by a Notary in 1735, where the Marquis de *Castellafuerte* procured a quantity of *Quinquina* from *Loxa*, to carry to Spain on his return.

Memoir de l'Acad. d. S. 1738.

I think it probable from a more attentive view of the subject, that it may be the Bark of the trunk, or larger branches of the tree, and I am the more confirmed in this opinion by the ideas of my friends Dr. WITHERING and Dr. FOTHERGILL, conveyed in their letters to me, in which they observe, that the essential and active parts of the Oak Bark are more intire, and in larger quantity in the trunk and larger branches, than in the twigs or smaller branches, which are comparatively of an imperfect growth; perhaps the Small and Quilled Bark may be procured from younger trees, not yet arrived at their full maturity, and therefore yielding a Bark of a weaker quality.*

I am

* Mr. ARROT, a Scotch Surgeon, who had gathered the Bark in the place where it grows, says, that the small curled Bark so much esteemed in England, is the Bark of younger trees, which frequently recover the
Barking

I am justified very much by the analogy of other Barks, and by the influence of soil and exposure, in changing the appearances, and even of affecting the virtues of the same species of vegetables, to conclude, that the diversity both in size and colour of the Red Bark from the common Peruvian Bark, may depend either on the largeness of the branches, or on other circumstances not necessarily implying a specific difference in the tree.—On comparing the larger Bark of the Oak with its twig Bark even collected from the same tree, I find the former of a reddish colour, while the latter

Bark of the Oak with its twig Bark even collected from the same tree, I find the former of a reddish colour, while the latter

Barking, while the older trees never do. This affords a strong proof, that the early Bark introduced into Europe was of the larger kind, and from the older trees, while the difficulty of procuring it, has been the means of introducing a small and younger Bark. Mr. ARROT, and all other Travellers agree in preferring the Red Bark to any other.

Encyclopædia. Cortex. Vol. I.

ter is much paler, the roughness of the coat of the larger Oak Bark and its general appearance may be compared to that of the Quill Oak Bark, as the Red Bark is to the common Peruvian Bark; and I have not only learned from consulting Tanners on this subject, that the larger Bark is superior in its powers as an astringent; but I have found by comparing infusions of both, and submitting them to the most decisive experiments regarding their astringency, by adding to them solutions of iron, that the precipitates were of a blacker colour and in greater quantity from the larger and more compact pieces of Bark, than from the smaller twig Bark. —I have convinced many medical gentlemen, who have seen my specimens of the Oak Bark, that they tend very much to illustrate and confirm the opinion that the Red Bark is the *Cinchona Officinalis* of *Linnaeus*.

The

The following description of the Peruvian Bark taken from Pomet is likewise a farther confirmation of this doctrine : —“ The *Kinquina* is the bark of a tree
 “ that grows in *Peru*, in the province
 “ of *Quitto*, upon the mountains near
 “ the city of *Loxa*. This tree is almost
 “ the size of a Cherry-tree, the leaves
 “ round and indented; it bears a long
 “ reddish flower, from whence arises a
 “ kind of pod, in which is found a ker-
 “ nel like an almond, flat and white,
 “ cloathed with a slight rind; that Bark
 “ which comes from the trees at the bot-
 “ tom of the mountains is thicker, be-
 “ cause it receives in more nourishment
 “ from the earth; it is smooth, of a
 “ whitish yellow without, and of a pale
 “ brown within. That which comes
 “ from trees on the top of the mountains
 “ is abundantly more delicate; it is un-
 “ even, browner without, and of an higher
 “ colour

“ colour within; but the trees which
 “ grow on the middle of the mountains,
 “ have a Bark yet browner than the other
 “ and more rugged. All these Barks are
 “ bitter, but that from the trees at the
 “ bottom of the mountains, less than the
 “ others.

“ It follows from hence, that the Bark
 “ of the least virtue, is that which grows
 “ in the lowest places, because it abounds
 “ more with earthy and watry parts, than
 “ that which grows high, and which for
 “ the contrary reason is better; but the
 “ best of all is that which grows in the
 “ MIDDLE of the MOUNTAINS, because
 “ it has not too much or too little nou-
 “ rishment.—There is another kind of
 “ Bark which comes from the mountains
 “ of *Potosi*, and is browner, more aroma-
 “ tic, and bitterer than the former, but
 “ abundantly

“ abundantly scarcer than any of the
 “ rest.

“ The conditions or qualities we ought
 “ to observe in the Bark, are, that it be
 “ HEAVY, of a FIRM SUBSTANCE, SOUND
 “ and DRY. Beware of such as is rotten,
 “ and will sink in water presently, and
 “ that flies into duft in breaking, or is
 “ dirty and unclean, as it happens to be
 “ sometimes; but make choice of such
 “ as is in little thin pieces, dark and
 “ blackish without, with a little white
 “ moss, or some small fern-leaves stick-
 “ ing to it, REDDISH within, of a bitter
 “ and disagreeable taste; and refuse that
 “ which is full of light chives when broke
 “ and of a ruffet colour, and take care
 “ that there be not several pieces of wood
 “ mixed with it, which you have more
 “ of sometimes than the Bark.

“ This

“ This was brought first into France in
 “ the year 1650, by the Cardinal Lago, a
 “ Jesuit, who having brought this from
 “ Peru, it was had in such vogue in
 “ France, as to be sold weight for weight
 “ at the price of gold.”

I have since the publication of the first edition of this treatise, extended my enquiries into other countries, from which I am convinced that there cannot a doubt be entertained of its being the *Cinchona Officinalis*. I have a specimen of the Red Bark which was given me by Mr. BABINGTON, the Apothecary of Guy's Hospital—it contains in it a branch of Quill Bark, exactly as it was imported.

I have seen some specimens of Red Bark so very large, that they contained a great proportion of woody part, and therefore less fit for use than those of a more moderate

derate size; indeed from having made such frequent experiments on this subject, I am able with great precision to ascertain the comparative quantity of resin in any two pieces of Bark from their external appearance.

It has been suggested by some, that the Red Bark resembled much the mahogany Bark; but having examined that Bark, and having conversed with persons to whom it is extremely familiar, I am persuaded that there is no foundation for the opinion of its being the same. If future naturalists, by having better opportunities of investigating this part of our subject, should be persuaded that it is the Bark of a tree of a different *genus*, or *species*, from the *Cinchona Officinalis*, such a discovery cannot invalidate the proofs of its superior efficacy; and I should have been happy had it been found to be the
production

production of any of our colonies, instead of its being as yet known in Europe only as a native of South America. Several very intelligent men, who were disposed to think it the Bark of a different tree, immediately changed their opinion, from examining it in tincture, decoction, or infusion, in which forms it conveys the genuine taste and flavour of the common Peruvian Bark, under the appearance of a much stronger impregnation. I suspect that we have been long in error by judging chiefly of the goodness of Peruvian Bark, from the colour of its external coat. I have seen some specimens of Red Bark extremely rich in resinous parts with a very white coat, but whose inner layers were compact, and of a dark red or ochry colour. I have examined twenty chests of this Red Bark in the very state in which it was imported, and there is always found a very considerable proportion of Quill Bark amongst it.

If the execution of this work was equal to its importance, it would challenge the attention of the public, in a degree far above most medical subjects. This will be unquestionably admitted by those who have been eye witnesses to the malignancy and fatality of intermittent and remittent Fevers, in every part of the globe, but more especially in warm climates; this fatality is by no means owing to the ignorance or unskilfulness of the practitioners in those countries, but to the inefficacy of the common Bark in general use.

The numbers who fall a sacrifice to the epidemic and seasoning Fevers of warm climates, are admitted infinitely to exceed those who are destroyed by the enemy. In almost all the dangerous Fevers which occur in our East and West India settlements, the Bark is a principal remedy. I think it therefore an object of the greatest national

national importance, that our fleets and armies should be liberally supplied with this Bark, which will seldom or ever disappoint them.

I have been told by a druggist, that its great activity rendered it a dangerous remedy; my answer was, that wine mixed with water was much safer in the hands of an unskilful practitioner, than wine alone; but that did not prove that wine was not a better cordial than water. The same reasoning may be applied to prove, that weak and decayed remedies, by being much milder in their operations, are therefore preferable to such as are more perfect of their kind. I believe the general, and best founded complaint is against the want of power and efficacy of Bark, and not that it is too powerful and active. In proof of this, I refer my readers to the letters annexed, which demonstrate, that such
was

was the stubbornness and obstinacy of the intermittent Fevers of the present year, even in this country, that they resisted common Bark, and only gave way to the Red Bark.

Notwithstanding I formed very early a favourable opinion of this Bark, yet it fell far short of that which I am justified in maintaining, from the collected evidence of so many gentlemen in distant and remote parts of the country. Many letters which I have received are written with such zeal in favour of its superior efficacy, that they could only be dictated by the strongest conviction, arising from extensive and diligent observations.

I have persuaded many of my medical friends to use the Red Bark in our foreign settlements, and I shall take pleasure in communicating to the public the result of

C

their

their observations as soon as I am favoured with them. A more powerful Bark is particularly desirable in those countries, where the violence and danger of the paroxysm is so great. In the following observations, I have confined myself very much to the use of the Red Bark in febrile diseases, but I am in possession of many facts in proof of its superior powers in other diseases, in which the common Peruvian Bark has been found useful.

The introduction of the Red Bark into this country was the effect of chance. In the year 1779, a Spanish Ship from Lima, bound to Cadiz, was taken by the Hussar frigate, and carried into Lisbon; her cargo consisted chiefly of this Bark, some part of which was immediately imported into this country, and a considerable quantity was bought at a very low price at Ostend,

Ostend, by some of our London Druggists. The Boxes in which it was brought to Europe were of the same kind as those in which the common Peruvian Bark was contained, and all sold by the general title of QUINQUINA. The Druggists in whose hands the Red Bark at first was, found it difficult to dispose of it, its appearance was so very unlike that of common Bark; at last they offered it by way of trial to such Apothecaries as reside in counties where agues are frequent; the success attending its use soon convinced them of its superior efficacy. It was early introduced into the hospitals, and its greater powers became universally acknowledged. It has continued ever since in general use in the Hospitals of St. Bartholomew, St. Thomas, Guy, and the London. The reputation therefore of the Red Bark stands better established, and is supported by the concurring testimony of more Physicians,

than that of any other article of the *Materia Medica*. I am assured by every Druggist with whom I have conversed on the subject, that the demand for it in this country is pressing and general. I am likewise informed, that the markets may be well supplied with it; and as it is no longer in the hands of a few dealers, the prejudices of the Druggists have subsided, and I have lately heard nothing of its destructive qualities, which were said to have arisen from its superior powers.

Being highly sensible of the difficulty of establishing such facts, either on the effects of remedies, or on any branch of medicine which regards the animal oeconomy, I have solicited the opinion of many ingenious and attentive practitioners, who from their situation have had frequent opportunities of trying the Red Bark. This caution appeared the more necessary, because

cause I am well persuaded that the love of novelty, and too great a credulity in admitting facts on very doubtful authorities, have corrupted medicine more than any other science, and proved more injurious than the most absurd and fanciful theories, the errors of which are easily detected.

OF ITS SENSIBLE QUALITIES.

THE Red Bark is in much larger and thicker pieces than the common Peruvian Bark. It evidently consists of three distinct layers. The external, thin, rugged, and frequently covered with a mossy substance, and of a reddish brown colour.* The middle, thicker, more compact, and

C 3

of

* I have lately seen some very good Red Bark whose external coat had a white appearance, though its internal surface is of a deep red colour, extremely resinous, compact, and heavy:

of a darker colour. In this appears chiefly to reside its resinous part, being extremely brittle, and evidently containing a larger quantity of inflammable matter than any other kind of Bark.

The innermost has a more woody and fibrous appearance, and is of a brighter red than the former.

The intire piece breaks in that brittle manner described by writers on the *Materia Medica*, as a proof of the superior excellence of the Bark.

In reducing it to powder, the middle layer, which seems to contain the greatest proportion of resin, will not give way to the pestle so easily as the other layers; this should be particularly attended to when it is used in fine powder. Its flavour is chiefly discoverable either in powder or solution, is evidently more aromatic, and
has

has a greater degree of bitterness than the common Bark.

OF ITS CHYMICAL AND PHARMACEUTICAL HISTORY.

EXPERIMENT I.

TO an ounce of Red Bark, reduced to a fine powder, were added sixteen ounces of distilled water; and after remaining together twenty-four hours in a Florence flask, the liquid was carefully filtered. The same experiment was made with the Peruvian Bark commonly in use.

The colour of the two infusions was very different, that made with the Red Bark being much deeper. The taste and flavour of the infusion of the Red Bark were considerably more powerful than of the other. In the opinion of many gen-

tlemen who tasted the infusions, the cold infusion of the Red Bark was more sensibly impregnated than even the strongest decoction of the common Bark.

EXPERIMENT II.

TO two ounces of the cold infusion of the Red Bark, were added *twenty* drops of the *Tinctura Florum Martialium*. It immediately became of a darker colour, soon lost its transparency, and after a short time precipitated a black powder.

EXPERIMENT III.

TO two ounces of the cold infusion of the common Bark were added *twenty* drops of the *Tinctura Florum Martialium* in the same manner as to the other. It retained its transparency some time, and afterwards became of a dark colour, but there was no precipitation from it as from the last.

E X P E-

E X P E R I M E N T IV.

To an ounce of red Bark, reduced to a coarse powder, were added *sixteen* ounces of distilled water, and after boiling until one half was evaporated, the liquid while hot was strained through a piece of linen. The same experiment, under similar circumstances, was made with the common Bark. The superior taste and flavour of the decoction of the Red Bark was equally observable with that of the infusion. The decoction of the Red Bark, in cooling, precipitated a larger quantity of resinous matter than the decoction of the common Bark. The difference of colour was likewise very distinguishable.

E X P E R I M E N T V:

To an ounce of Red Bark, reduced to a coarse powder, were added *eight* ounces of
proof

proof spirit, and, after standing a week together, the Tincture was filtered.

The same experiment, under similar circumstances, was made with the common Bark. The Tincture of the Red Bark, both when tasted by itself and under precipitation by water, had more flavour and taste than that of the common Bark.

The Tincture from the Red Bark is of a much deeper colour than the other.

EXPERIMENT VI.

To each *residuum* of the above Tinctures were added *eight* ounces of proof spirit, which were infused in a moderate sand heat for the space of *twenty-four hours*, and afterwards allowed to remain together a week, occasionally agitating them.

them. The Tinctures were then poured off, that of the Red Bark evidently appearing to be the strongest.

The Tinctures both of Experiments V. and VI. were by a gentle heat evaporated to the consistence of a resinous extract.

The extract from the Tincture of the Red Bark was of a smooth, homogeneous appearance, not unlike the Balsam of Peru, when thickened: The flavour and taste of the original Tincture were intirely preserved in it.

The extract from the Common Bark had a very different appearance. It seemed coarse and gritty, and by no means so characteristic of its original Tincture.

The quantity of extract procured from the Red Bark was considerably greater than
from

from the same quantity of common Bark ; but, as the *residuum* of neither was rendered entirely inert, the absolute quantity could not be ascertained. *

E X P E.

* To 26 lb. of red Bark were added 26 gallons of proof spirit, after remaining together for some time the Tincture was poured off, and submitted to a distillation in a water bath, the quantity of spirituous extract obtained was 12 lb. and a half ; a quantity of water being poured on the *residuum* of the Tincture, the watery extract obtained was 4 lb.

In another experiment with 30 lb. of Red Bark, of an inferior quality, treated in the same manner as the former, only 11 lb. and a half of spirituous extract was procured, and 4 lb. and half of watery extract.

The same quantity of the best Peruvian Bark hitherto in use gives from 6 lb. and a half to 7 lb. and a half of spirituous extract.

It may be proper to observe, that the facts here mentioned are on the authority of a very eminent druggist, who had accurately marked the quantity of extract obtained by the usual process from a given quantity of red Bark at two different trials ; the reader will likewise observe, that although the spirit employed for making the Tinctures may have been saturated with

EXPERIMENT VII.

A tea spoonful of each of the Tinctures, prepared by Experiment V. was added to *two* ounces of water; the resinous precipitation from the Red Bark was not only more copious, but fell more quickly to the bottom of the glass than that from the other, and yet what remained still dissolved in the water, was infinitely more in the Red Bark than in the common Bark, so far as we could judge from the taste and flavour of both.

with resin, yet a fresh quantity was not poured on the residuum, which by extracting the whole resin would have yielded a larger proportion of resinous extract, and consequently left little or nothing for the watery extract. Though these experiments were not made with any view to a philosophical purpose, yet I am sufficiently convinced of their accuracy; they are more conclusive than experiments conducted on a much smaller scale.

E X P E-

EXPERIMENT VIII.

In imitation of the experiments of my ingenious friend Dr. PERCIVAL, I added to *two* ounces of the watery infusion of each Bark a few drops of the *Sp. Vitriol, ten*. The acid lost its taste more in the infusion of the Red, than in the common Bark, so that there were more obvious appearances of its being neutralized.

EXPERIMENT IX.

A decoction of both Red and Common Peruvian Bark was prepared by taking an ounce of each and boiling them in a pint and a half of water, to *one* pint; the former had greatly the superiority in strength and power, as mentioned in a preceding Experiment. A pint of fresh water was added to each decoction; the boiling still continued till that quantity was

was evaporated. The decoction of the common Peruvian Bark seemed gradually to lose its sensible qualities, while that of the Red Bark still retained its own.

The same quantity of water was added as before to each, and the decoction repeated until a gallon of water was exhausted; at the expiration of which time, the common Peruvian Bark was rendered almost tasteless; the Red Bark still retained nearly its former sensible qualities. This experiment proves that the common practice of boiling the Bark is hurtful to its powers.

By my desire Mr. SKETE, a very ingenious and attentive young gentleman from Barbadoes, and a student of medicine in Guy's Hospital, made several Experiments in order to ascertain the comparative antiseptic

tiseptic power of Red Bark, with the common Peruvian Bark; and he found that the infusion of Red Bark preserved animal matter much better, and for a longer time, than the infusion, or even decoction of the common Bark; indeed, the decoction of common Bark, after its powdery part had subsided, was less bitter, and preserved animal matter for a shorter time than the infusion of the same Bark. His experiments were conducted with great accuracy, and the result of them were submitted to the examination of many gentlemen at Guy's Hospital.

The conclusions to which the above experiments evidently lead, are,

First, That the Red Bark is more soluble than the Peruvian Bark, both in water and spirit.

Secondly,

Secondly, That it contains a much larger proportion of active and resinous parts.

Thirdly, That its active parts, even when greatly diluted, retain their sensible qualities in a higher degree than the most saturated solutions of common Bark.

Fourthly, That it does not undergo the same decomposition of its parts by boiling as the common Peruvian Bark.

Fifthly, That the Red Bark is more astringent than the common Peruvian Bark.

Sixthly, That its antiseptic powers are greater; as an additional proof of this it may be proper to observe here, that both its cold infusion and decoction preserved entire their bitter and other medicated powers in the month of June, in the Ela-

D

boratory

boratory of Guy's Hospital for five weeks, and perhaps for a much longer time, while a decoction of common Bark gave evident marks of a change in a few days. In the decoction of Red Bark, the powder, which is separated during the cooling of it, remains intimately diffused through the liquor, which therefore continues loaded and turbid when at rest. In the decoction of common Bark, the powder quickly subsides to the bottom, the Red Bark therefore contains in it a large proportion of mucilaginous parts, such as have been proposed by the late Dr. FOTHERGILL, to be added to the decoction of the common Peruvian Bark, in order that it may remain turbid when at rest, and thereby that its resinous parts be more perfectly suspended in the body of the liquor. It is obvious that this circumstance will favour exceedingly the action of the stomach upon it.

The

The advantages therefore to be expected from the Red Bark cannot be obtained from any quantity of common Bark. The best common Bark, compared with the Red Bark, appears inert and effete.

All the above experiments were executed in the presence of several Gentlemen.

I was led more particularly to prosecute this subject, from an opinion that the Red Bark might so impregnate cold water by infusion, as to cure Intermittent Fevers with more certainty than could be done even by the decoction or powder of common Bark. The sensible qualities which appear from the above Experiments, being so much greater, in the cold infusion of the one than in the decoction of the other.

It cannot I think be denied, that the Experiments above related, and which have been executed and frequently repeated with great accuracy, sufficiently prove that the Red Peruvian Bark, exceeds the other in its sensible qualities, and that it contains a much larger proportion of those resinous and active parts on which the power and efficacy of Bark have been by all writers on the practice of medicine and *Materia Medica* believed to depend.

OF THE GENERAL OPERATION
OF BARK ON THE HUMAN
BODY.

THE following remarks are intended to apply to the Peruvian Bark, generally in use; but I am certain that the effects enumerated are found to be produced in a much higher degree by the Red Bark.

The

The cold infusion of Bark seems evidently to promote both appetite and digestion, it increases the tone and action of the stomach, by which the gastric liquor, the great *Menstruum* of our solid aliment, is more perfectly prepared.

In most cases, the Bark rather promotes costiveness, the common effect of strong and vigorous intestines. In very large doses however it generally proves purgative, but this effect ceases after a short time.

It renders the pulse stronger and fuller in health, and in most diseases unaccompanied with Fever.

In Low and Malignant Fevers, and more especially under remission, it renders the pulse stronger and even slower.

In particular circumstances of Fever marked with debility and a tendency to remission, though of a very irregular type, it diminishes febrile heat,

It encreases the animal heat and aggravates every symptom, in Fevers accompanied either with much local inflammation, or a general inflammatory diathesis, which is strongly indicated by the pulse, the manner of the original attack, the want of due freedom in the secretions, and the painful and oppressive exercise of every function.

It checks profuse and colliquative discharges, especially those by the skin, while it does not seem to diminish insensible perspiration, or other natural evacuations.

It checks every tendency to putrefaction or gangrene, occurring under circumstances of debility, but it may promote both, if injudiciously employed, while the action of the system is too violent, or the inflammatory diathesis is too prevalent. It seems more reasonable to refer its action, as an antiseptic, to its tonic power on the moving system, than to any primary action on the animal fluids.

Perhaps this doctrine will apply in explaining the history of remedies used in the Scurvy, a disease invited and favoured by every means which can induce debility, and evidently preceded by symptoms of a diminished *vis vitæ* which necessarily lead to others that in a secondary manner only take place in the animal fluids.*

D 4

It

* An Enquiry into the Source from whence the Symptoms of Scurvy and of putrid Diseases arise, by Dr. MILMAN.

It promotes under many circumstances a favourable suppuration, and improves the nature of sanious and ichorous discharges.

Its action here can only be explained from its general tonic power, for either general or local debility retards suppuration, and favours the obstinacy of ill-conditioned ulcers.

Upon the same principle its power of promoting the generation of true *pus* in the Small Pox may be explained.

The period of debility is that only in which the Bark should be employed.

I have seen patients under the most confluent Small Pox require Bark in the progress of suppuration, and yet, in the more advanced state of the disease, the
same

same persons have been saved by the seasonable and repeated use of the Lancet in the secondary Fever, which attacked with fresh rigors and inflammatory symptoms of a true Peripneumony.

In delicate and irritable habits, which seem more especially to favour scrophulous affections, and which produce inflammation of a peculiar nature, that gives way sooner to tonics than evacuants, the Bark has been found the best remedy: this may probably admit of the explanation already given on the subject of Scurvy.

OF ITS USE IN THE CURE OF INTERMITTENT FEVERS.

WHAT I have to offer on this subject is the result of careful, and diligent observation

tion, totally unconnected with prejudice of any kind, in favour of any particular theory, or a blind attachment to system.

Notwithstanding the cautious and timid practitioner has very generally forbid the use of Bark until evacuations should have been made; I am very well persuaded from observation, that in Intermittent Fevers, such as rage and are endemic, particularly in low and marshy situations, and such as frequently occur on the banks of the Thames, and the lower parts of this metropolis, the Bark cannot be given too early; the use of either emetics or purgatives, as preparatory, is not only unnecessary, but in some cases productive of greater debility, and therefore to be avoided,

The doctrine of concoction, however just it may be in continued Fevers, and in some cases of inflammation, does not apply
in

in Intermittents produced by the *Miasmata* of low and swampy grounds, and which afford the principal source of agues in the environs of this city.

In this opinion I am confirmed by the testimony of Dr. CLEGHORN and others; who frequently found it necessary to give it on the first accession of the disease, in order effectually to obviate, or weaken the return of a *second* paroxysm, which in many cases would inevitably have proved fatal; and Dr. LIND has very properly observed that such symptoms which have been attributed to Bark, are rather the effects of the paroxysm being allowed to return, from the neglect of that medicine.

There are many symptoms which would forbid the use of Bark, did they occur distinct and independent of Intermittent
Fever,

Fever, such as Cough, difficulty of breathing and pain in the side: they are frequently brought on by the paroxysm of the Intermittent, and only give way to the use of the Bark by which that paroxysm may be prevented. Such symptoms do not admit of a distinct and separate treatment, but are always aggravated by the use of evacuants, more especially bleeding, the most probable means of relief in common Depuratory Fevers.

I have sometimes found a complication of Intermittent and Hæctic Fever in the same person and could distinguish between the paroxysms of each; the Bark, while it cured the Intermittent, has even moderated the Hæctic; though Hæctic Fever of itself, especially as a symptom of *Phtisis Pulmonalis* does not appear to give way at any time under the use of the Bark; I think it probable, therefore, that
 although

although in the treatment of intermit-
tents, complicated with other diseases,
our chief attention should be first directed
to the cure of the intermittent, yet it
may be necessary to pursue an indication
that may have in view the other disor-
ders, not incompatible with the treat-
ment of the Intermittent. In Dropsies,
which accompany Intermittents, I have
found more benefit from the use of Bark,
joined to the neutral salts, and other mild
diuretics, than from active purgatives,
which always tend to protract the Inter-
mittent. Every returning paroxysm of
an ague confirms more and more the cause
of that disease which it has produced.

It is not intended to insinuate that no
cases do occur, in which it may not be
prudent to administer a vomit, especially
to persons subject to bilious accumula-
tions in the stomach; but this is more
with

with a view of removing an obstacle to the operation of Bark, than as necessary to render it safe; and I have frequently seen *nausea* and vomiting so much a symptom of the paroxysm, that they gave way only to the free use of Bark itself.

A practice more absurd than that of preceding evacuations has been adopted and recommended, *viz.* That of evacuating by purgatives after the cure had been compleated by the Bark, this seldom fails of bringing back the Intermittent, as one cause of inducing debility, the most favourable state of the body for the attack of Intermittent Fevers.

In persons who are rendered costive by the use of the Bark, I would recommend the common practice of giving a few grains of Rhubarb, or *Pilul. Rufi.* If on the other hand it should prove purgative,

tive, a few drops of the *Thebaic Tincture* is the usual and best means of checking that operation.

Intermittent Fevers are frequently so very anomalous in their appearances, and assume so much the character of other diseases, that an unskilful, or inattentive practitioner may be deceived; they, however, under any form or any type, generally give way to the Bark; sometimes aided and assisted by other means.

Much experience is required to detect the paroxysm of an Intermittent, which is frequently disguised under different forms, and although the Bark is our best remedy, yet the violence of some symptoms which accompany the disease, and interrupt its natural and usual form, render it necessary that other remedies be occasionally employed, as adapted to the particular

particular circumstances of the case; not always so much with a view of rendering the Bark a safe remedy, as of rendering it an effectual one.

Such anomalous appearances are greatly influenced by the nature of the prevailing epidemic of the season, they more particularly interrupt the progress of the cold fit, and are generally much aggravated during the paroxysm of the Intermittent.

The disorders which I have seen complicated with Intermittent Fevers have been chiefly those of a bilious kind, occurring in the autumnal season, such as violent and excessive vomiting, Diarrhæa, with pain in the bowels, Cholera Morbus, periodical Head-Achs, Pain in the Side, and frequent inflammatory disorders, such as Pleurisy, Peripneumony, and the
acute

acute Rheumatism, and even sometimes spasmodic diseases, terminating in apoplexy and death.

Such diseases have been supposed to be only varieties of the Intermittent Paroxysm, and have been believed to give way to the same remedy which cures an ague.

I think, however, we have no analogy in nature to support this doctrine, and it seems highly unreasonable to suppose, that the same cause can produce such a diversity of appearances. Indeed the influence of the prevailing epidemic diseases on sporadic complaints evidently shew, that the human body is subject to be acted upon by more than one cause at a time, and that the remedies to be employed should have a view to such a diversity of circumstances. It therefore becomes

E

the

the object of the physician to know which of two or more diseases are most deserving of his early attention, which the human body may labour under at the same time. He will generally find, that as the paroxysm of an Intermittent Fever excites such violent action in the system, and generally aggravates the symptoms of other diseases, it ought to be early removed.

In some cases however I have seen such active appearances of inflammation prevailing in a person labouring under an Intermittent Fever, and so much encreased in the hot fit, that unless a quantity of blood had been taken, which was always fizy, the patient most probably would have died. Such symptoms of inflammation retard the cure of the Intermittent, and therefore are in the first instance to be removed.

This

This may probably explain the reason why inflammatory Fevers in their decline often assume the appearance of Intermittent diseases.

In the same manner it may be necessary to remove Symptomatic Vomiting, Cholera Morbus, and the like, by remedies peculiarly adapted to these diseases, before the stomach will so far favour the action of Bark as to enable it to cure the Intermittent Fever.

Peruvian Bark is chiefly adapted to the cure of genuine and idiopathic Intermittent Fevers, and not those of a symptomatic nature, which frequently require remedies of a different kind.

It is impossible in a treatise of this nature to point out more minutely the cir-

cumstances which ought to regulate and direct the conduct of practitioners in the treatment of complicated Intermittents, and the condition of the habit, which may render necessary the previous use of other remedies, or the combination of them with Bark, in order to render its operation either safe or effectual.

It appears to me, that the advantages at any time derived from the use of other remedies, depend upon their having removed some other disease, which may have protracted the ague, or interrupted the action of Bark in the cure of the Intermittent, and not on their obviating future effects which have been falsely attributed to Bark, while they are chiefly produced by the obstinacy and the imperfect treatment of the Intermittent Fever.

Both the Vernal and Autumnal Intermittents of Dr. SYDENHAM yield to it, the latter however sometimes with more obstinacy than the former.

In several cases I have experienced the efficacy of the Red Bark in removing *Tertians* and *Quartans* which had resisted the common Bark, this however is not to be wondered at, when we consider the diversity in the power even of different kinds of the common Bark in general use.

I hope I shall be excused in digressing so far, as to mention the useful effects I have frequently experienced from the exhibition of Opium in Intermittent Fevers. We are chiefly indebted for this practice to Dr. LIND. It moderates so effectually the force of the paroxysm by shortening the duration of the cold fit, as well as by di-

minishing the violence of the hot fit, that I had often flattered myself it was capable of curing Intermittents. In this however I was disappointed..

From the experience of it in many hundred cases, I conclude with Dr. LIND,
 “ That an Opiate given soon after the
 “ commencement of the hot fit, by abating the violence and lessening the duration of the Fever, preserves the constitution so entirely uninjured, that
 “ since I used Opium in Agues, neither a
 “ Dropsy nor Jaundice has attacked any of
 “ my patients in these diseases.” The manner in which I employ it, is either by giving a grain of the *Thebaic* extract upon the accession of the cold fit, or *twenty* drops of the *Thebaic* Tincture upon the accession of the hot fit, the action of the former being later from its slower solubility.

The

The Red Bark is so much warmer than the other, that it would seem to answer all the purposes derived from the union of Cordials, Aromatics, Serpentaria, and the like, so much recommended in the obstinate *Quartan* Intermittents of elderly people.

Some difference in opinion has prevailed regarding the manner of giving the Bark. Most practitioners concur in thinking, that it cures Intermittents more readily when taken in substance than in any other form. In this state, it is both a bulky and nauseous dose in the quantity necessary to cure an Intermittent; at any rate, it ought rather to be diffused in some liquid, than given in the form of an electuary or pills, which are sometimes difficultly soluble.

I have found milk cover the taste of Bark, and make it more acceptable to children than any other vehicle. The extract of Liquorice dissolved in water, may be likewise employed to cover the taste of Bark. Its taste is also corrected by wine, especially by Old Hock.

It would appear from the general preference given to Bark in substance, that its decoction, infusion, or tincture, are found too weak in any quantity for the purpose of curing Intermittents, otherwise as they are much lighter to the stomach and act more quickly, they should be preferred.

I hope to make it appear, that in this respect the Red Bark has the advantage of any other kind now in use, since either its infusion or decoction will cure Intermittents,

mittents, and its powder in a much smaller dose than that of common Bark will produce similar effects.

The best time for giving the Bark is in the intermission between the paroxysms and when the stomach is empty. In *Quartan Fevers*, where there are *two* days of *Apyrexia*, we should be particularly desirous of getting down a large quantity on the day immediately preceding the approaching paroxysm, and in other Intermittents as near the period of the returning paroxysm as the stomach will bear it.

A very prevailing argument in favour of the Red Bark has been suggested to me both by apothecaries and their patients, viz. that it will cure when taken in half the quantity which has been found necessary of other Bark. I am likewise persuaded from a great variety of trials, that
while

while other Bark only gradually weakens the force of the Intermittent Fever, the Red Peruvian Bark will frequently obviate the return of a second paroxysm. It is seldom I have found it necessary to give more than half a dram every two hours in the interval of the fit, and in no one Intermittent, even of a *Quartan* type have I found it necessary to give more than six drams between the paroxysms. I have frequently known double that quantity of common Bark fail to produce the desired effect.

It is however unnecessary to limit the dose. One dram may be given every hour, if the stomach will retain it, and will perhaps in some cases remove the disease more quickly than a smaller quantity given at longer periods.*

The

* I have likewise frequently adopted the practice recommended by Dr. HOME, of giving the Bark as soon

The following facts will best determine how far I am justified in favouring the opinion of the superior excellence of the Red Bark.

EDWARD VIRGOE, aged 21, had laboured under an Intermittent Fever *five* months. It was first a *Tertian* and afterwards became a regular *Quotidian*, accompanied with Cough, Dyspnœa, and Hoarseness, particularly in the paroxysm. The common Peruvian Bark, given in the dose of one dram every hour, prevented the return of the paroxysm for a few days; the patient however relapsed notwithstanding the Bark was continued, he was at last cured by taking one dram of the Red Bark every second hour for the space of ten days.

It

soon as the sweating fit of the Fever has sufficiently carried off the hot fit; this is particularly proper when the interval is short between the paroxysms.

It appears from the above case of *Edward Virgoe*, and several others which have occurred to me, that Intermittent Fevers which had resisted common Bark and other remedies, have yielded to the Red Bark, even under very complicated and unfavourable appearances.

In this opinion I am likewise justified by the experience and testimony of many eminent practitioners, and so decided are they in its favour, that the demand for it every day increases, especially in some of the neighbouring counties where Intermittents are not only more frequent, but more obstinate.

I began now to suspect that its powers were even sufficient in cold infusion in most cases to cure Intermittent Fevers, and in all other cases to answer every purpose which might be expected from common
Peruvian

Petuvian Bark, in any form in which it had hitherto been employed.

The following facts are sufficient to authorise this opinion.

JAMES YOUNGMAN, aged sixteen, had laboured under a *Tertian* intermittent many months; it was accompanied with a severe cough, and his strength was considerably impaired. He was ordered to take *four* ounces of the cold infusion of the Red Bark every *third* hour; after taking it for *two* days the paroxysm did not return. Its use was persevered in *fourteen* days, and he continued perfectly well.

THOMAS PUGH, aged twenty-two, from *Woolwich* in *Kent*, had an Intermittent Fever of *five* weeks standing of the *Tertian*

tian type, accompanied with a considerable degree of cough, occurring especially in the cold fit; he took the cold infusion, as recommended in the former case, after which time he had only one slight return of the paroxysm, his cough is likewise totally removed.

JOHN WELDING, aged twenty-five, had an Intermittent Fever at *Sheerness* for near a year, he took large quantities of common Peruvian Bark, without effect. It was of the *Quotidian* type. After taking a cold infusion of the Red Bark in the quantity of a quart in *twenty-four* hours for *three* days, the paroxysm disappeared and has never since returned. It proved at first purgative, but soon lost that effect. His strength and appetite were greatly improved under the use of the infusion.

Since

Since the first edition of this work, I have had frequent experience of the efficacy of the cold infusion in the cure of Intermittent Fevers, but it requires to be continued for a greater length of time than is necessary, when the Bark is taken in substance with wine. I have however seen cases where the stomach had rejected the Bark in substance, which yielded readily to the cold infusion when taken in the dose of four ounces, every two hours in the interval of the paroxysm. I have likewise met with cases of Intermittent Fevers so complicated with other disorders, as to make it necessary to act more gradually and cautiously on the constitution, by the use of the cold infusion, than by giving the Bark in substance in the doses necessary for instantly removing the Intermittent Fever.

As

As I consider a perseverance in the use of the Bark proper for some time after the paroxysm has been removed, I think that it is only necessary to have recourse to the cold infusion for that purpose, and it will be found a much more agreeable preparation than any other. I have found great benefit from continuing its use in the convalescent state of persons after Intermittent and other Fevers.

I could here enumerate a great variety of cases which have occurred to me, both in public and private practice, in confirmation of the general doctrines I have now laid down, but I think it better to mention the general result of a successful practice.

The cold infusion employed in the above cases was prepared by pouring a
quart

quart of cold water on two ounces of the Red Bark in fine powder, frequently agitating them for the space of twenty-four hours. *

OF ITS USE IN THE CURE OF OTHER FEVERS.

Remittent Fevers are frequently equally steady in their periods of remission and accession with those of the intermittent kind, but the *Apyrexia* being less perfect and complete has given rise to many doubts regarding the safety of giving Bark. As remittent Fevers are more particularly

* Though the cases above mentioned, evidently prove that the cold infusion will cure Agues, yet they are not intended to divert the attention from other more effectual means of giving this medicine.

F

marked

marked by appearances indicating the prevalence of bile in the stomach; the propriety of giving an *emetic*, prior to the use of the Bark, seems well founded, and the remission is frequently rendered more complete by such a practice.

In the remitting fevers, however, of warm climates, the accession of the paroxysm is so extremely violent, and the strength of the patient so quickly exhausted, that it becomes absolutely necessary to catch the first opportunity of the most trifling remission, and to give the Bark with the same freedom that you would do in common Intermittents.

In Remitting Fevers, the concomitant symptoms are more to be regarded than in Intermittents, because they more especially disturb, interrupt, and shorten the periods of remission; even in the warmest climates,

climates, and under the severest prejudices, it has been found necessary to take away a few ounces of blood to relieve pain in the head, oppression in breathing, an intense dry heat on the skin, and other symptoms protracting the paroxysm; by such means the remission has been brought on, and the Bark given with greater effect.

The Fevers of this country seldom have regular remissions, until they have been properly treated by evacuations; the inflammatory by bleeding, and the bilious by vomiting and purging.

When Fevers are brought into a state of obvious remission; that is, when the pulse becomes from ten to twenty slower at some particular time in the twenty-four hours; when the restlessness, anxiety, and tendency to delirium abate; when the mouth and fauces are moist; when the

organs of secretion, and especially the skin are more open and pervious, such symptoms of remission admit the use of Bark with the same freedom as in Intermitting Fevers.

The ACUTE RHEUMATISM, notwithstanding its inflammatory attack, and the appearance of the blood, and although the joints often continue inflamed several weeks, very early assumes the form of a Remitting Fever.

Under such circumstances, a perseverance in the *Antiphlogistic* plan is generally found to be ineffectual. I have, in several cases of this kind, employed a cold infusion of the Red Bark, and the disease seemed to give way only to this treatment.

The ACUTE RHEUMATISM in its remissions assumes the form of a double *Tertian*, and the patient is frequently
greatly

greatly exhausted by the profuse sweatings which terminate the paroxysm.

It is in such cases that I would particularly recommend the use of Bark. I have found this practice more successful, and it must be allowed to be more rational, than the use of *Volatiles* and *Guaiac*.

The tediousness, as well as the inefficacy of the antiphlogistic practice in the cure of the Acute Rheumatism, has frequently directed my attention to that disease in a particular manner, and after being repeatedly disappointed and dissatisfied by pursuing the usual mode of treatment recommended by the best writers, either antient or modern, on the subject, I was determined to adopt other means, which I think have proved more successful. The Rheumatic Fever appears to me, notwithstanding the violence of its inflammatory symptoms, to be an Intermittent Fever in a state of disguise, and its periods

are evidently, as I have already mentioned, those of a double *Tertian*. Its inflammatory symptoms, however, ought in a certain degree to be reduced by moderate bleeding, occasional purging, and great dilution, before it can be treated as an Intermittent Fever; it does not however appear to me incompatible with the ideas of its being inflammatory, to have recourse to the moderate use of the Bark, to obviate the weakness which may be induced by the necessary use of the lancet, nor does it appear repugnant to the idea of its being intermittent, that the inflammatory symptoms which render the Intermittent anomalous and irregular, should be moderated and checked by an antiphlogistic treatment accompanying the use of Bark. I have found in many cases by this practice, the Rheumatic Fever greatly shortened, and the debility and torpor in the joints, which is frequently the effect of that disease, together with the disposition to the
Chronic

Chronic Rheumatism, generally prevented. The Acute Rheumatism appears from the state of the pulse, the tendency to profuse sweating, the deposition in the urine, the frequent accession of chilly paroxysm to be an Intermittent Fever, which is probably prevented from assuming the more usual and natural form of that disease by the inflammatory action on the joints, which I have sometimes seen merely local, (*i. e.*) unaccompanied with any general Inflammatory Fever in the habit: in such cases, while leeches have been applied to the joints, and the hemorrhage from them encouraged by fomentations, I have given Bark freely, which I have never known to increase the inflammatory symptoms while the usual means of promoting inflammation were guarded against. General and vague maxims, applied in reasoning on the action of Bark, and its tendency in all cases to promote inflammation are therefore ill found-

ed, and had they not been corrected by experience and observation, would have deprived us of the advantages we have derived from the use of this medicine in the cure of Rheumatic Fevers, Scrophulous Inflammation, and perhaps a great variety of other diseases. I have seen in delicate and irritable habits Rheumatic Inflammation on the joints, accompanied with a low Nervous Fever, which gave way only to Bark and Sedatives.

In the Rheumatic Fever I generally begin about the seventh day from the attack with the cold infusion of the Red Bark, in the dose of three ounces every two or three hours, until the evening paroxysm comes on; nor am I, by this practice, in any degree, diverted either from, general or local bleeding, or evacuations by stool, when the circumstances occur which may render them necessary.

In

In that *low Fever*, which HUXHAM has so well described under the title of nervous, I have prescribed the cold infusion of the Red Bark with advantage, where the skin hath been soft, and the pulse under one hundred and ten.

In such Fevers, it chiefly acts as a Cordial in supporting the *Vis Vitæ*, and for which reason I think it may be given with advantage in the decline of all Fevers, even where the symptoms on the attack of the disease were evidently inflammatory.

Almost every Fever remits in its decline.

Fevers originally putrid and malignant, as arising from *Miasmata* and putrid vapour, very seldom occur in this City, they are most generally to be found in Fleets and in Camps, and in situations where
air

air stagnates or where animal bodies are confined in a close place.

In such Fevers, the cure is effected by vomiting and warm Cordials; of the last is the Red Bark in an eminent degree.

It may be infused in wine, which will render its operation more antiseptic.

It intimately unites with the several acids, from which a very favourable operation in such cases may be expected.

In the Putrid Fever, attended with a gangrenous Sore Throat, I have in many instances experienced the efficacy of Bark, but care should be taken not to confound this disease with the *Angina Mucosa* of Dr. HUXHAM, or the *Angina Erysipelatosa* of Dr. GRANT, diseases, though contagious, and accompanied with *Eruptions* on the skin, evidently of the most inflammatory

matory nature, and requiring the use of
Evacuations.

In *Petechial Fevers*, with symptoms of great prostration of strength, I have frequently seen the pulse not much quickened, and the animal heat very little increased. In such cases I would recommend the use of the Red Bark infused in Old Hock.

I have seen a *Jail Fever* with no other distinguishing symptoms than Petechiæ and Debility; the tongue clean, the pulse moderate though small, and the secretions apparently not disturbed. In that case, blisters on the extremities, Bark and Wine, are the best remedies.

In general, we find that Fevers marked with symptoms of debility are chiefly found to remit, and therefore they admit of the use of Bark; by increasing the tone
and

and vigor of the system, it opposes a returning paroxysm.

Fevers of more violent action, such as we denominate inflammatory, do not remit at least so obviously, until that action is moderated by Evacuations, so that they chiefly remit only in their decline.

The primary attack of most Fevers in this country, is attended with violent action, which is best moderated sometimes by the prudent use of the lancet, but most frequently, by the *Tart. Emetic* with the infusion of *Sena* and some of the neutral salts. All Fevers beginning with a violent *Rigor*, and followed with great heat, require early evacuations, while such as creep on slowly and imperceptibly in the beginning, with giddiness in the head, rather than acute pain, much anxiety and watchfulness, tremor and debility, give way to Opiates, Bark, *Serpentaria* and Wine.

Blisters applied to the arms are extremely useful in keeping up the pulse, especially if the skin be soft, the tongue and fauces sufficiently moist; much more may be learned by attending to this last appearance than is generally known.

It is necessary, however, to distinguish between the dryness of the tongue and mouth, as a symptom of the Fever, and as arising from the circumstance of a patient sleeping with his mouth open.

I hope the observations here offered to the public, on the subject of this invaluable remedy, will remove all prejudices against a Peruvian Bark of a large and coarser appearance, than is generally employed.

It is at present in very great demand, the difficulty of procuring it will not, I hope, instigate Druggists and Dealers in
the

the article, to substitute at any time a spurious kind in its room.

Extract of a Letter from Mr. EDWARD JACOB, jun. an eminent Surgeon, at Faversham, in Kent.

I H A V E had such repeated opportunities of trying the Red Bark, that I hope to be able to give you every satisfaction you can wish, of its superior efficacy over the Bark in common use.

Our situation being in a country not far distant from the marshes, renders the inhabitants more exposed to intermitten complaints than those of cities and more inland counties; and I assure you, before the use of the Red Bark was known, the Ague, from its particular stubbornness (as we then thought, or what we have since observed from the want of efficacy in the other Bark) might be truly called
the

the *opprobrium Medicorum*; but now I think that stain is entirely removed, for I have not met with one single case, where (when I could persuade the patient to adhere steadily to my advice) I have ever found the Red Bark to fail.

The Peruvian Bark, with which I was formerly provided, was of the best kind, and always had in the quill; yet, it was even much inferior in its effects to what ought reasonably to be expected from it.

The first knowledge that ever we had of the Red Bark was in March, 1781, when a few pounds were sent us to try its effects; our Druggist informed us, that it was then in use at St. Bartholomew's Hospital, and was there found effectual. It remained in our house till May following, when I was attacked with an Ague; I did not

at

at first think of trying the *new* Bark, for I guessed, by taking of the *old* in great quantity, which would not be disagreeable to me, that my Ague would soon leave me; but, to my great disappointment, it succeeded fit, without shewing the least good effect of the remedy applied. I was then determined to try the *new* Bark; but finding my Ague stubborn, I emptied the *Primæ Viæ* by an emetic and carthatic, and immediately, on the Fever going off, I took one dram of the Red Bark in fine powder, mixed with three ounces of the decoction, and a small quantity of the tincture, which being a draught well loaded with Bark, at first seemed to sit uneasy on the stomach (which I have several times on being first taken, found the case with some of my patients, but which never took from its effect) but, after resting for some time, I found myself in a state to continue its use. My Ague from that time kept off;

off; till, from omitting the Bark too soon, and finding myself quite well, in a few weeks after, slight symptoms appeared again; but which never formed a direct fit; the Bark being again repeated, eradicated the disorder.

My father, now in the 70th year of his age, has since that time been attacked with an Ague; but, from his having been before seized with a chilliness without subsequent Fever, suffered himself to have three fits without trying any remedy; being then convinced of the reality of the Ague, applied to this Bark in decoction, with some tincture made of the same; he found it so effectual, that after taking it, the fit did not once return; he omitted the Bark too soon, and had one slight fit; but, after repeating the same remedy a few days, it has not since returned, and he is now perfectly well.

G

The

The patients who have received immediate relief from the new Bark, are so numerous, that I cannot, with any degree of certainty, guess the number.

The quantity that we have used from July, 1781, to the present time, is upwards of sixty pounds. The manner in which we have used it, with a view to prevent the return of an approaching paroxysm, is by giving half a dram or one dram of the powder, mixed with two ounces of the decoction, and a small quantity of the tincture. The decoction we have used, has been prepared by boiling three ounces of the gross powder boiled in two quarts of water to one quart.

When the patient has been of a more delicate frame, or when it has been used as a corroborant, we have given the decoction and tincture without the powder. When the Ague has been but recently contracted,

contracted, we have seldom trusted to any thing but the Bark, but when stubborn, evacuants have been first given; half an ounce of the powder has very frequently cured an Ague without evacuants, or more Bark, when the fit was a recent one; we have now entirely left off giving the *old* Bark in any form since we have found so good effects from the Red Bark, and we used the Resin of Red Bark with good success in many Cases, where the patient could take it only in the form of pills. I heartily hope the above account will prove satisfactory to you, if not I shall think myself very happy in answering at any time any future inquiries.

I am,

Sir,

with great respect,

your most obedient

and humble servant,

EDWARD JACOB, Junr.

Feversham, June 23, 1782.

*A Letter from Mr. BOYS, an eminent
Surgeon and Apothecary, at Sandwich in
Kent.*

S I R

I AM sorry it is not in my power to send you particular cases of the effects of the Red Bark: having made no memorandums, I can only say, in general, that it is a much better kind of Bark, than any I have been able to procure hitherto. Whether the Intermittents have been worse than common, or the Grey Bark has been of inferior quality, I will not take upon me to determine; but we were unusually foiled in our attempts to cure that complaint, till we were furnished with the *new Bark.*

I can have no inducement to give a higher character of this drug, than it deserves: but I do assure you, that since I began to use it, it has not once failed me,
when

when the patient has taken it in a proper manner. Hence, I have a firm persuasion of its superior efficacy, and I am the more confirmed in my opinion, by knowing that my sentiments correspond with yours, and with those of all my medical acquaintance, in this neighbourhood, who have made trial of it. My best wishes attend your publication; not only because I am persuaded it will operate to the advantage of mankind, by extending the knowledge and use of this valuable medicine, but, likewise, because I am in hopes it will hasten a fresh importation of the article, of which I have very little left, and I know not where to go for a supply.

I have the honor to be,

Sir,

Your most obedient servant,

W. BOYS,

Sandwich, June 19, 1782.

G 3 *In*

In addition to the character given of the Red Bark, by several practitioners in the country, is a Letter I received a few days ago from Sir WILLIAM BISHOP, an eminent Surgeon, at Maidstone, in Kent, dated June 16th.

IN which he shews, by a variety of cases, that in the cure of Intermittents, in that part of the world, the Red Bark had not only been infinitely more effectual than the common Peruvian Bark or any other remedy usually employed, but that it had radically cured where Cold Bathing, Emetics, Opiates, Bitters, and Friction failed, and where the best Pale Bark, both by itself, and with a variety of other medicines, was exhibited without effect, even to the quantity of sixteen ounces. The form he gives it in, is that of an electuary, with the addition of a small

small quantity of the essential Oil of Pepper-mint and Carraway-seed; and he seldom has occasion, he says, to use more than from four to six ounces. One case he mentions, where a Sphacelus had taken place in a Malignant Fever, accompanied with delirium and every other bad symptom, the patient was recovered by the use of the Red Bark, Anodynes, and Wine; the Sphacelus separating kindly, the Fever and every bad symptom speedily gave way.

I am likewise favoured with another Letter from Sir WILLIAM BISHOP, dated June 23, which contains many sensible and judicious hints upon this subject.

HE recommends the grinding Red Bark into the finest powder by a mill, and afterwards sifting it through the finest cypress sieve, in order that it may fit lightly on the stomach, and that its parts

may be uniformly blended together: he observes, that if, after breaking the Bark, you examine the broken pieces, by means of a glass in the sun-shine, you will see the resin like spangles of gold between the internal woody fibres and the outward grey coat; by comparing this appearance with the best Pale Bark, you will discover that the Red Bark contains a much larger quantity of resinous parts.

Two scruples of Bark, as coarsely powdered as is commonly sold by the apothecaries, will be as disagreeable to take, as a dram or four scruples finely ground. One dram is the dose he generally gave to an adult, although he sometimes met with patients, who would choose to take two or three drams at a dose, and those doses, at longer intervals; and from experience he found that, when this last practice agreed with the stomach, it was the most effectual way of curing the Intermittent,

mittent. One patient took three ounces
in twelve hours, and had no return of
an obstinate *Quartan*.

Many Intermittents, which could not
be cured by small doses, were removed by
larger doses, more frequently employed.

He mentions that a period of *five* or *six*
hours, before the expected return of a
Quotidian or *Tertian* Intermittent, is all
that is necessary for the taking a sufficient
quantity of the Red Bark, in order to ob-
viate the approaching paroxysm. Some
have been cured by taking *two* ounces, as
quick as possible after the fit, but such as
persevered in the use of it, until *four* or
six ounces were taken, acted most pru-
dently, their health was soonest and most
perfectly restored. He likewise observes,
that such persons ought to take the great-
est quantity of Bark, whose blood is in
the most dissolved state, and where the
fibres

fibres have been relaxed by the most profuse sweating, which frequently takes place in obstinate *Quartans*.

Quotidians require but a small quantity, *Tertians* will yield to fewer doses than our *Kentish Quartans*.

He found it better to get down *five* or *six* ounces of Bark, in eight or ten days, than to allow a larger time for the same quantity. Labouring people, who work out of doors in cold foggy mornings, either in marshy grounds or wet lands, from sun-rise till the dew of the evening, and who sweat profusely, and suffer their wet linen to dry on their backs, will not be safe from a return of the *Quartan* Fever, with less than five ounces taken after the last fit, as the good effects of all they had taken before, are very probably carried off and dissipated in the ensuing paroxysm.

He likewise observes, that the Bark in large doses will not cure the irregular fits of an ague and fever, which so often afflict persons, when matter is forming in the lungs, in the *Phtbisis Pulmonalis*.

The letters from Sir William Bishop, which convey these observations, likewise contain much valuable information on other medical subjects, and I hope the ingenious author will, on some future occasion, favour the public with them.

A Letter from Dr. WITHERING, an eminent Physician, at Birmingham, June 29, 1782.

DEAR SIR,

I AM much pleased that you have undertaken to give us an account of the Red Peruvian Bark. A publication upon that subject, cannot fail to produce good effects, by removing the prejudices of
some

some, the ignorance of others, and ultimately by exciting our merchants to obtain liberal supplies of a medicine so truly valuable. The universal prevalence of Intermittent Fevers this spring gave us a large experience of its effects. We have not many Intermittents in the town of Birmingham; but in other parts of the county of Warwick, in Staffordshire, Shropshire, Worcestershire, and Oxfordshire, so far as my rides extend, they have been more general than ever known before in the memory of the oldest practitioners I have conversed with.

After taking pains to recommend the Red Bark to all the Apothecaries I met with, and consequently not less anxious to learn their observations relative to its effects, I can say that they unanimously concur in asserting “ that they have never been disappointed in their expectations when they exhibited it to patients labouring under Intermittents.”

Now

Now the Intermittents which I have seen, have pretty generally assumed the *Tertian* Type in light soils, and the *Quartan* Type in clayey countries. Of the latter, I have had the care of some which were from six to eighteen months duration, originating in Kent and Essex. I have known two instances only of *Quartans* in which the Red Bark did not prevent a recurrence of the Fits; I saw one of these patients afterwards; he laboured under hepatic and anasarous symptoms, these were removed in a fortnight by the usual methods, and then the Bark effected a cure. The other case was at a considerable distance from this place, and I have not yet learnt its termination.

It will require some farther experience to ascertain the necessary doses. I know some practitioners who have given one or two drams every four hours betwixt the fits, but I have never had occasion to give

more

more than thirty or forty grains at similar intervals of time.

As to its preparations I can say but little, preferring always in my own practice the simple powder; but, I am told, that it makes a very rich tincture, and I have seen decoctions of it very high coloured and turbid.

But it may be asked—Would not the common or Quill Bark, have produced similar good effects in the Intermittents of the present year? From the reports of other practitioners, I believe it would not. From my own experience, I can give no other answer to such a question, than, by saying, that after frequent and almost continual disappointments, from the use of common Quill Bark, I have not tried to cure a single Intermittent with it for more than seven years past; relying entirely upon the use of evacuants, opium,
and

and metallic salts. To render this last assertion reconcilable to the daily experience of others, it may be necessary to observe, that with us, a Physician is seldom consulted in Fevers of the intermittent kind, unless some unusual appearance, or some uncommon obstinacy in the recurrence of the attack, alarms the patient or his friends.

Whether the medicine in question be the product of the same tree from which the Quill Bark is derived, may be difficult to determine, but I am persuaded it is the medicine that was used by MORTON and SYDENHAM, or its efficacy could never have been so proverbial. I know not what could first induce the writers upon the *Materia Medica*, to prefer the Quill Bark, but I know if you were to ask an English Tanner, whether the Bark from the trunk, or that from the twigs of the

Oak

Oak is the strongest, he would laugh at
your ignorance.

I communicated the contents of your
letter to my worthy friend and colleague
Dr. Ash, together with my opinion upon
the subject; he authorises me to say, that
the result of his experience perfectly co-
incides with mine.

I remain,

with the truest esteem,

Yours,

W. WITHERING.

P. S. Thus, my good friend, have I
freely communicated my sentiments and
observations upon the subject you have
in hand.

You are at liberty to make what use
you please of them, and I shall think
myself

myself happy in having contributed a little towards the extensive usefulness of your design. I have seen no bad effects from it, notwithstanding the reports of some of the London Druggists, but their motives were too evident to need a comment.

I feel, with you, that distance alone has interrupted our communications; but I feel too, that distance can never abate the regard with which I once more subscribe myself,

Affectionately,

Your's,

W. W.

To Dr. Saunders, Jefferies Square.

H

A Let-

A Letter from Mr. SHERWIN, an ingenious Surgeon at Enfield.

S I R,

S I N C E we have used the Red Peruvian Bark, we have had great success in curing Agues and Intermittent Fevers. These disorders returned so frequently after the use of the best Peruvian Bark, which we could formerly procure, that our credit, as well as that of the medicine, began to fail very much, and numbers of our patients got into the hands of persons unacquainted with physic. They seemed willing to trust rather to nostrums and charms, than to a medicine which they have seen so repeatedly fail, and to the effects of which they very ingeniously ascribe every ache and pain which continue after Agues, or which are the consequences of Agues when imperfectly cured.

I could

I could give a very ample detail of the inefficacy of the common Bark; but as the complaint is general, it would be unnecessary. I shall, therefore, only take the liberty to present you with the more agreeable history of a few cases that have immediately yielded to the Red Peruvian Bark, and that even when given in very moderate doses, after having obstinately resisted large quantities of the other.

RICHARD PARSLEY, a hard working young man, about twenty-eight years of age, was seized with an Ague last autumn, which continued either as a *Quotidian*, *Tertian* or *Quartan*, with very little intermission, till the 20th of April last, when I saw him accidentally. He gave me the following account :

That he had taken every thing that had been recommended to him; and that

his master, a worthy and respectable gentleman in this neighbourhood, had procured him the advice of different practitioners of physic, who had prescribed Bark for him in large quantities, but without benefit, as the disorder always returned with greater violence after it had been checked a few days. He added, that the last medicine which he had been advised to take, was half an ounce of Allum boiled in a pint of Ale, to half a pint, which he was ordered to repeat three different times, as soon as he felt the Ague approach. He had in this way, he said, taken seven half ounces upon the approach of as many different fits of the Ague.

I suspected, that so large a quantity of Allum, taken at one dose, must have produced some dangerous effects, and doubted whether he had swallowed the whole of it; but I found, upon more particular

particular enquiry, that he really swallowed half an ounce each time, as he says; that he stirred it up from the bottom and felt it gritty in his teeth. It gave him great pain in his stomach. So large a quantity of Allum being taken at one dose, without materially injuring the system, is a fact which I am persuaded you will think worthy of notice.

I took compassion upon this poor fellow, and sent him eight papers of the Red Peruvian Bark, containing a dram in each, which he finished in two intermediate days, the disorder being then a *Quartan*. It is now seven weeks since he took this medicine, and he has had no return of the complaint, but, to use an expression of his own, "hath felt ever since as if he had a new inside."

ANNE FIGOT, a poor girl, fourteen years old, has been afflicted with an Ague

since Autumn last in the same family with PARSLEY, and has also tried various medicines without effect. I prescribed small doses of the *Vitriolum Cærul.* fourteen days without gaining any advantage. I gave her also three or four doses of a hot aromatic powder, consisting of Bay Berries and Caian Pepper, which I have frequently seen to cure, but she found no benefit from it. I cannot say whether this girl had given the common Bark a fair trial. She begged to have some of the same medicine that cured her relation, and about a month since had three papers, containing only one scruple in each, and I assure you, she has been perfectly free from the complaint ever since.

Mr. BARNES, a very strong, hard working man, was seized last Autumn with a very violent Intermittent Fever, attended with delirium, and apparently much danger.

ger. After three or four paroxysms, I removed the disorder by giving him one dram of the common Peruvian Bark every hour, and continuing the same quantity twice a day afterwards; but at the end of fourteen days, the Fever returned with the same violence as before, and was again stopped by the same quantity of Bark. From Autumn, till the beginning of last *April*, it returned in this manner (at the end of about fourteen days from the time it was stopped) five or six times, and always with uncommon violence. The common Peruvian Bark, newly powdered, was repeatedly given,

In the beginning of *April*, I prescribed six doses of the Red Peruvian Bark, of one dram each, which he took upon the going off of the paroxysm, and I have the pleasure to inform you, that he has had no return since that time; though I no

longer used the precaution of continuing the daily use of the same medicine.

It is not worth while to mention common cases, where the Red Bark hath cured without the other having had the trial. A few have occurred, and I have not yet had any instance of a relapse.

I was lately called to a very severe case, like that of Mr. BARNES, where the intermission was not longer than six or eight hours. I prescribed, and the patient took six drams of the Red Bark, without interrupting the paroxysm, which came at the expected hour; and during the sickness and horripilation the Bark seemed to be entirely rejected by vomiting; however, the succeeding paroxysm abated something of its fury. My patient was extremely averse to Bark in every shape; but my deceiving him, during the following intermission, I got him to swallow two
drams

drams at once, which rendered him so entirely averse to the medicine, that he would take no more afterwards. However, to his great joy and my surprize, the Fever left him. He ascribes his cure to three or four draughts of Camphorated Julep, which were prescribed as a *placebo* upon his refusing to persevere in the use of the Bark. But it may with more justice be ascribed to the two drams of Red Bark, taken at one dose; as I think the first six were in a great measure lost.

I consider the Red Peruvian Bark as a valuable acquisition to the *Materia Medica*: or, perhaps, it may be only a restoration of what was used in the days of SYDENHAM, and some time after, when it was common for medical writers to say, that this, or that medicine would cure, with as much certainty as Bark would an Ague. An Eulogium to which the Bark
in

in this country has not been of late years entitled.

Before I became acquainted with the superior efficacy of the Red Peruvian Bark, I had tried a variety of *tonics*, with very indifferent success in general, but now and then with very happy effects, such as the *Cuprum Ammoniacum*, *Vitriolum Cæruleum*, *Sal Vitrioli*, *Sal Martis*, &c.

WILLIAM KING, an athletic young man, applied to me last Autumn, on account of a tedious irregular Intermittent, complicated with acute fixed pain in the side, and a full strong pulse, much cough and hoarseness. Under these circumstances, I thought it imprudent to administer the Bark in any Form, and had recourse to a plentiful bleeding, applying blisters to the part affected, and giving the *Decoct. Taraxac.* with *Tart. Solubile*, and Honey in large quantities. The disorder

order was clearly an Ague; but attended with symptoms that threatened a pulmonary consumption. His blood was remarkably fizy, which induced me to make use of *venesection* four times, which, with the above medicines, and a very strict *antiphlogistic* regimen, removed the cough and the pain in the side, though the Ague still continued. I ventured at last to give him eight doses of common Bark, one dram in each, which removed the Ague several weeks, and mended his habit much. A continuance of the Bark, twice a day, for some time after the Ague stopped, brought on a frequent bleeding at the nose.

The Ague returned twice during the winter, and gave way soon to the same Bark, and he kept tolerably well till the setting-in of cold north-east winds in spring, when his Ague again returned, and

and brought with it the pain in the side, a bloated icteric countenance, much hoarseness and cough. After one more bleeding, and the use of the *Apozem* for a month, I gave him six drams of the Red Peruvian Bark in April last, and have had the pleasure to see him continue free from the Ague since that time, though he never repeated the dose after it stopped.

I was unwilling to persist in the use of the Bark after the Ague stopped, lest it should encrease the circulation too much, and again excite a bleeding at the nose.

I have now indeed almost entirely laid aside the custom of persevering in the use of the Bark, after the Intermittent is stopped. I have for some time suspected that it answers no good purpose, and that it may possibly now and then be
the

the reason why large quantities of good Bark have been given in vain.

It is a well known fact, that every species of intermittent complaint frequently returns in fourteen days after being interrupted by means of Bark, notwithstanding that medicine is continued every day. May not the daily use of Bark so habituate the constitution to its effect, as to render it useless when the Intermittent returns?

I am now satisfied when I have put a stop to the expected paroxysm, and prescribe an anodyne to be in readiness if ever the cold fit returns, advising my patient to have recourse to the same quantity of Bark which he found necessary at first, and to take it as soon as the succeeding Fever abates. I adopted this practice a little time before I became acquainted with the superior

superior efficacy of the Red Bark, and thought it an improvement, but since that time I have succeeded so well by giving six or eight drams between the paroxysms, that I have seldom had occasion to repeat a single dose.

I am,

Sir,

with great respect,

your obliged and

most obedient Servant,

JOHN SHERWIN.

Enfield, June 23, 1782.

A Letter

*A Letter from Dr. FOTHERGILL, an
eminent Physician, in Harpur-street.*

DEAR SIR,

I N answer to your obliging request, I shall now proceed to lay before you the best information I can concerning the New Bark, lately introduced into practice, under the name of CORTEX RUBER, or RED BARK. As it was found on board a Spanish prize, intermixed with a small quantity of common Peruvian Bark, it would seem to be no other than the produce of the trunk or large branches of the same tree. It seems to be possessed of the same sensible qualities, only in a much higher degree, hence it yields a much larger proportion of resinous extract, and gives a more saturated tincture and decoction than the common Bark. Hence too I have found (according to what you
lately

lately observed) that the decoction may be exposed a long time to the open air, without contracting the degree of acidity, which manifestly takes place in that of the common Bark in a few days.

It has been remarked for many years past, that the Peruvian Bark has often disappointed the expectation of the public; having fallen greatly short of that transcendent degree of efficacy, for which writers of the last century have so highly extolled it. Few Physicians of extensive practice, but must sometimes have had the mortification to observe their favourite specific entirely baffled by a regular Intermittent, without being able to assign any probable cause for the defeat, except the *ungenuineness* of the medicine; whence it would seem to follow, either that the Intermittents of late years have been more *obstinate* in their nature, or that the Bark has been of an *inferior quality*

lity to that which was used by Dr. SYDENHAM, and many of his respectable contemporaries, of whose veracity we can entertain no reasonable doubt. The latter appears to me to be the more probable, especially when it is considered, that of late years, it has been customary (for what reason I cannot imagine) to select the Bark of the small branches, under the denomination of *Quill Bark*, in preference to that of the trunks. Should this preference hereafter be discovered to have been groundless (agreeable to what I have long suspected) it may perhaps contribute to explain the principal cause of our disappointments.

In the art of tanning, experience has long determined in favour of large Oak Bark, as being greatly preferable to that of the smaller twigs. Why the reverse of this should take place in a medicinal view, is by no means evident, especially

cially if the virtue of the Peruvian Bark keeps pace in any degree with its astringency.

Spanish practitioners, as I have been lately informed, are so well convinced of this, that they always prefer the large Peruvian Bark; separating it from the smaller sort, for all important purposes, which the late capture indeed seems to render probable.* Future observations may probably discover that this was the real genuine Bark, with which our ancestors cured all the various kinds of Intermittents, with a degree of certainty, which now astonishes their successors.

* This must not however be confounded with those coarse woody flakes, which constitute a large proportion of the worst kind of Bark, now in use. For these, having been already stripped of the outer rind, together with the resinous cells, consist of a mere ligneous substance, divested of medicinal virtue.

If

If Bark taken from the large branches should be found to be more efficacious than that of the small, is it not reasonable to believe, that, that of the trunk or rather of the root, might surpass both in virtue? Political reasons, however, respecting the preservation of the trees, which produce so important an article of commerce, will, it is to be apprehended, effectually deter the inhabitants from every experiment of this nature.

A very eminent Druggist, who purchased a large share of the above cargo, assures me, that many of the Apothecaries whom he has supplied with the Red Bark, and who have carefully compared its effects with those of the common *cortex*, make no scruple of deciding in its favour. He further adds, that the demand for it has increased so rapidly

of late, that the stock in hand is already nearly exhausted.

It is to be regretted, that the passion for interlarding this simple febrifuge with other bitters and astringents (after the usual complex mode) should still too much prevail in practice. Under this false idea of adding to its efficacy, its virtue may often be greatly diminished, and at all events, the result of the trial must be rendered extremely equivocal. In justice to the remedy, and to obviate this uncertainty concerning its effects, I have embraced every opportunity (which so short a space would admit) of administering it in its simple state, without intermixing it with other medicines. The cases in which I have chiefly tried it, have been low putrid Fevers, attended with extreme prostration of strength, delirium during the evening exacerbations, with short and obscure remissions in the day time.

For

For the sake of brevity I will but mention one instance. WILLIAM HENTON, a Silk Weaver, laboured under a Fever of this kind, accompanied with *Petechiæ*, and a *profuse nasal Hemorrhage*, which last symptom generally returned with the exacerbations. To these were added cold sweats, muttering delirium, involuntary twitchings, &c. Before I saw him, common Peruvian Bark had been prescribed, and his case pronounced *desperate*. In this very critical situation, however, I was not deterred from recommending the *Red Bark*, in proper diluents, acidulated with *Spiritus Vitrioli*. Accordingly he took from two scruples to a dram every two hours, except during the midnight exacerbation, when it was ordered to be cautiously avoided; because I have long been convinced by experience, and attentive observation, that the febrile symptoms are generally exasperated when a Bark remedy is given at that period. The

medicine agreed, the Hemorrhage abated, and the Fever soon subsided. Some weeks have now elapsed, and I have the satisfaction to add that lately, on entering his room, he assured me with a joyful countenance, that since he had taken the Red Bark (or as might now, perhaps, with more propriety be said, the *Real Bark*) he had suffered no symptoms of relapse, and that he remained free from all complaints, except a little weakness of sight, to which he had been formerly liable. Should this medicine be hereafter found to answer as completely in the other various types of Fevers, and that in smaller doses than the common Bark, it will become highly interesting to the patients, and no less pleasing to the Physician, especially when he shall be enabled beforehand, thus to pronounce with real confidence concerning the event.

“ *Hi motus, atque hæc certamina tanta,*

“ *Pulveris exigui jactu compressa quiescent.*”

VIRGIL.

From

From its success in this and *several other instances*, which have fallen under my own observation, I am strongly inclined to believe, that it will soon become an object worthy of attention, but must decline giving a decisive opinion concerning its superiority, till I know the result of a variety of cases, in which it is now under trial, and also receive additional confirmation from such correspondents, as are less prepossessed in its favour than myself, because they may be supposed to prove in reality more impartial judges. Therefore in the interim, I shall wait with impatience for your intended publication, hoping, that by thus collecting the sentiments and observations of several practitioners in different parts of the kingdom, we may at length be enabled to ascertain the true comparative merit of the

Red Bark, which cannot but afford great satisfaction to the public at large, as well as to,

Sir,

Your most obedient servant,

A. FOTHERGILL,

Harpur Street, July 6, 1782.

A Letter from Mr. EDWARD RIGBY, an eminent Surgeon, at Norwich.

DEAR SIR,

I am very much obliged to you for the honor you have done me, in sending me your Treatise on the Red Peruvian Bark, and I feel myself particularly flattered by the manner in which you have solicited an answer to your letter.

I am

I am happy that it is in my power to bear the most ample testimony to the great and certain efficacy of this Bark; I have had the fullest opportunity of trying it in every species of the Intermittent, and have given it to patients under the greatest variety of circumstances respecting age, constitution, date of the disease, &c. and out of more than a hundred and fifty persons, who have taken it under my direction, *it has not failed in a single case, stopping the return of the paroxysm in the very first instance of its application.*

I was fortunate enough to receive a small parcel of this Bark so early as October, 1781, owing to the friendly communication of Mr. TALBOT, Surgeon, at Wymondham, in this county, to whom Mr. HOPKINS had sent a few ounces as a specimen for trial, half of which he immediately sent to me; Intermittents being at that time very frequent in Norwich,

wich, and its neighbourhood, I had an opportunity of trying it the day after I received it, and I chose the two worst cases, which then occurred to me, the one was a *Quartan*, of many months standing, the patient, a boy about eleven years of age, very much worn down by a *Quotidian*, of nearly as long a date; the patient, a young woman of about eighteen years of age, whose constitution was also much shattered by the long continuance of the complaint:—to both of them I gave half a grain of Tartar Emetic, at the approach of the next fit, which brought off a good deal of Bile from the stomach, and when the Fever terminated, they began to take the Bark, an ounce of which was divided into twelve doses, all which were taken by each of them before the times of the expected returns of the complaint; they both lost the Fits, and though they took no more of it, for they had taken all which was sent me,

they

they had no return of them until many weeks after, when the weather proved very wet, and then they went away without taking any more of this, or the common Bark, for they were both at this time removed to such a distance in the country as not to be able to send to me.

My success in these cases, induced me to procure a considerable quantity of it, and as soon as I received it, I made use of it in several *Tertians* then under my care, in all which it immediately answered the intended purpose. About this time I had a patient, a gentleman about fifty years of age, who had had a *Quartan* more than three months, he had been endeavouring to cure himself by a variety of popular receipts, amongst which was one which contained a considerable quantity of the common Bark; when I first saw him it was the day after he had had a Fit; but he was then much
indisposed,

indisposed, his pulse was too quick, he had no appetite, and was much reduced, he took a saline Fever medicine, and an opening draught before the next fit, at the approach of which I gave him a grain of Emetic Tartar, which vomited him considerably, and he repeated half a grain of it every three hours, until the Fever terminated, which, though it lasted a long while, went off with a greater and more general perspiration than was usual with him.

I now thought it right to begin giving him the Bark, but his intermission not being quite so perfect as I wished it, and moreover my not having yet had a sufficient number of cases, in which I had given the Red Bark, fully to establish its reputation with me, I thought it most prudent to give the common Bark, of which he took two scruples at such intervals during the intermission, that he
got

got down more than an ounce and a half of it before the time when the fit was expected, but it did not succeed, and he had another fit fully as severe as his former one; I then resolved to make the next trial with the Red Bark, an ounce of which was taken in the course of the succeeding intermission, and in doses of two scruples, and this stopped the fit; I prevailed upon him to continue its use in the same dose three times a day for a little time, which he did till he had taken two ounces more, but he has taken none since; and though his employment exposes him very much to the weather, and he was soon after frequently wet, yet he had not the slightest return.

From this time I confidently gave it in every case which came under my care, and its use was invariably attended with the same immediate success. In the number of those which were cured, were several
whose

whose legs were much swelled, and their bodies hard, and who appeared to be very rapidly hastening into a Dropsy; more than twenty of them were children, two were infants, not a year old; and one, whose case I shall relate, was a striking instance of the truth of your remark in the Treatise, that the Bark given in considerable quantity as near as may be to the time of the approach of the fit, is particularly efficacious.

My patient was a gentleman about twenty-five years of age, robust, and of a full habit; he had had an irregular Intermittent about a fortnight, it began as a *Quartan*, of which he had three fits, it then became a *Quotidian*, and he had three or four more fits; I saw him about the time when it began to come every day, and I found the Fever was very considerable; the usual treatment, which I
 need

need not particularize, was made use of, until he appeared to be in a state to take Bark; the fit after which he was to begin to take it, terminated about ten o'clock at night, and its return was expected the next day between twelve and one at noon. The time of the intermission being short he began to take it in doses of a dram: I had a message from him in the night, that it disagreed with him, and that he could not keep it down; I desired him to persevere, but to consider what came up as if he had not taken it, and to supply its place with another dose: between five and six o'clock in the morning he sent to me again, and desired me to go and see him, I found him much fatigued for want of sleep, which he had been entirely hindered from getting, by his repeated endeavours to take the Bark, every dose of which, except the first, had been rejected, and he seemed convinced that his stomach would not retain it. I prevailed
upon

upon him, however, to take half a dose; and this kept down; I staid with him some time, and half an hour after taking the first half dram, I repeated the same quantity, which likewise kept down; I then left him, desiring him to continue it every half hour in the half doses. Between eight and nine o'clock I was sent for again, and he gave me the same account of its coming up again as before; notwithstanding this, I ventured to give him another dose, which he immediately threw out of his mouth, before he had swallowed any of it; observing this, I was more particular in enquiring in what manner he had vomitted up, as it was called, the former doses; and from the attendant's account I was convinced, that what he imagined to have been vomited up, had never been swallowed; by this unlucky mismanagement of himself, he had taken but two drams and a half instead

stead of five drams, and there remained five drams and a half to complete the ounce, which was to be taken in little more than three hours; however, I was determined he should persevere, and I immediately gave him a dram, as I was no longer under any apprehension of its coming up, being fully satisfied that all had been retained which had been actually taken into the stomach; this kept down, and by strictly attending him with a dose every half hour, or at longest every three quarters of an hour, the whole was gotten down before the time of the expected return of the fit, which happily prevented its coming, and he has continued well ever since. The night having been entirely without sleep, and the patient having been excessively fatigued and fretted by his many fruitless endeavours to take his medicine, were very unfavourable circumstances in his situation, and

K

prevented

prevented his Fever from going off so perfectly as it did in the former intermission, for he was much hotter, and his pulse quicker than was to be wished, when taking Bark; however, the event fully justified my perseverance, and strongly proved the efficacy of the medicine.

Many of my patients having been poor and ignorant people, and some of them living at some distance from Norwich, to whom I could only give general directions, as it was impossible for me to attend them, you may easily imagine that most of that class of them took it carelessly; some of them I know did not take the quantity prescribed; and I recollected one person, a strong country girl, about seventeen years of age, who took an ounce of it at two doses, yet all of them were immediately cured. One patient of this class was a child about eight years old, who had had a *Quotidian* more than

two

two months, and was directed to take half an ounce of the Cortex between the fit, which was to terminate on a Monday noon, and was expected to return on the Tuesday, about the same time; on the following Saturday the child's father called to inform me that he was cured. Upon enquiry how the child had taken it, I was surpris'd to find that he had not then finished the half ounce, for when I told him that I meant the child should have gotten the whole quantity down in one day, he said he had understood before that I directed it to be taken between the Monday and the Saturday night, so that it was evident the child could not, the first day, have taken more than one dram of the medicine, which it is clear proved sufficient to stop the fit.

From the foregoing, and from some other cases which I have had, as also from those communicated to you by my friend

Mr. Sherwin of Enfield, one may certainly very strongly presume, that a much less quantity of this Bark than what I have usually given would answer the purpose; it would be a very easy matter to ascertain this by more trials, but at present I am not willing to give it in a smaller quantity, nor would I recommend it to other practitioners to do so, until the reputation of the Bark has been fully and universally established, for whilst the prejudices of some practitioners, who are averse to new medicines, and the interest of some Druggists, who will be probably sufferers, by having large stocks of the common Bark by them, may in the least degree tend to oppose its general use, it is to be wished that the testimonies in its favour should not only be strong and clear, but that its success should be as uniform and invariable as the nature of the medicine admits of, and therefore I would not yet venture to prescribe a
quantity

quantity so small, as to run a possible
 risque of its failure, when the quantity
 I have hitherto used, which is an ounce
 to an adult, and a proportionable less
 quantity to children, has not, with me, in
 a single instance been fallible.

In relating the few cases above, I did
 not think it necessary to mention the
 names of the patients, but as the circum-
 stances which tend to recommend a new
 medicine cannot have too great notoriety,
 I will, as they occur to my memory,
 give you a list of some of the persons
 who have been cured of Intermittents by
 this Bark, and whose situations in this
 neighbourhood render them well known.

Mr. Thomas Smith—Mr. Garland—
 Mr. Carter, Jun. twice—Mr. Money—
 Master Money, and Miss Money, of
 Trowse—two children of Mr. Barham of
 Kirby—Mr. Oliver—Mrs. Taylor—Mr.

Kiddell's daughter of Colney—Mr. How-
 lat of Earlham—Servant of James Crowe,
 Esq.—Servant of John Gay, Esq.—Ser-
 vant of Mr. Bloom, of Trowse—Mr. and
 Mrs. Clift—Miss Clift—Son of Rev. Mr.
 Abdell—Mr. Dixon—Master Webb—
 Mrs. Kett—Master Bunn—Mrs. Denny of
 Shottisham—Mr. Wright of Brecondale
 —Mrs. Glover of Kirby—Mr. Smith of
 Burlingham—Servant of Sir Lambert
 Blackwell, Bart.

With regard to the common Peruvian
 Bark, notwithstanding the complaints of
 its inefficacy have been great and general,
 my experience for a few years past has
 given me no reason to be so much dis-
 satisfied with it as I find many others are.
 I will acknowledge, indeed, that for some
 time past, I have found it necessary to
 give a much larger quantity of it than
 usual, and that even when I have given
 from an ounce and a half to two ounces of

it in substance, it has not always succeeded in stopping the fit in the first instance of its application, but when I have been able to prevail upon my patients to persevere in its use, in the same quantity, I have never known it fail to stop the fit after the second intermission, in which it has been taken. I should imagine there is no reason to believe that the common Bark, which has been used for some years past, is not the same with what was formerly in use, or with what was perhaps, originally introduced; as far as can be judged by its taste, and its appearance, either in the lump, in powder, in decoction, or in any of the other preparations of it, it seems to me, at least, to be precisely the same as I have always seen it; I have, therefore, never once suspected that, as a natural production, it has degenerated, much less have I apprehended that any artful means have been used by Druggists, to render it more saleable, or to increase their profit upon it, by which

its medical quality has been diminished: Intermittents having been more general in this country for two years past, than, perhaps, was ever remembered by any Practitioners now living, probably the same cause which has made them so frequent, has made them of a worse kind, and consequently more difficult to remove; and to this cause, rather than to any change in the quality of the Bark, is, in my opinion, to be attributed the late general want of success in the treatment of this disease. Upon the whole then, Sir, from the experience I have had in using the two kinds of Bark, which has not been inconsiderable, it is evident to me, that they possess [the same medicinal quality, but that the Red Bark has it in a degree greatly superior to the pale, which strongly favours your supposition, that they are both the produce of the same tree, the Pale or Quill being the Bark of the smaller branches, and the

Red,

Red, that of the larger branches, or the trunk of the tree. Having myself found such singular satisfaction in the use of this Bark, I sincerely wish it may be universally introduced, and I am persuaded that every Practitioner, who will give it a fair trial, will immediately prefer it to the Pale. Nothing can be more agreeable in the administration of a medicine, than to be able to speak of and foretell its effects with confidence, as it must be very encouraging to the patient; this circumstance, and its answering the purpose in a much smaller quantity than the Pale, are very peculiar advantages which the Red Bark possesses; for in the use of the Pale, though, as I before observed, I make no doubt, but by perseverance, and taking it in large doses, it will for the most part succeed, yet I have more than once been awkwardly situated with patients, who have taken a large quantity of it without its having answered the intention in the first instance,

instance, I mean in stopping the fit, after the first intermission, in which it has been given, it not being always an easy matter to persuade persons, under such a disappointment, to persevere taking down a large quantity of a nauseous medicine, more especially, when there still remain popular prejudices against the Bark; and it is a notion received by some, that when it does not immediately succeed it must be hurtful.

I have just received a letter from Mr. TALBOT, the gentleman whom I mentioned before, as having first sent me a specimen of the Red Bark, with an extract from which, relative to the subject, I shall conclude this already too long letter; he informs me, that from the time he first made trial of it, he has used no other in Intermittents, that he has given it to more than fifty persons, and he has not failed removing the disease in a single instance

instance though before that time he had been very unsuccessful in the use of the Pale Bark. He mentions a case, in which four ounces of the common Bark had been given without effect, and that an ounce and half of the Red immediately put a stop to the Fit:—And he further says, that he lately made enquiries about it amongst some Practitioners in his neighbourhood, to whom he recommended it, and that Mr. SWALLOW, of Watton, Mr. BRINGLOE, of Hingham, Mr. GIBBS, of Buckenham, and one or two more, have been equally successful in its use as himself.

I am, Sir,

With the utmost respect,

Your obliged and humble Servant,

EDWARD RIGBY.

Norwich, Sept. 8, 1782.

A Letter

*A Letter from Dr. JAMES MADDOCKS,
Physician to the London Hospital.*

DEAR SIR,

AGREEABLY to your request, I here send you an answer to the questions you proposed to me, relating to the large and Red Peruvian Bark, lately introduced into use in England.

Your first question related to my opinion of the medicinal efficacy of this Bark, with that of the paler, smaller, and quilled Bark, which for a long series of years has been considered in this country as superior to every other species.

In answering this question, the short notice you have given me, and the little time I have at present to spare, will not admit either of my taking notice of many different

different kinds of diseases, in which I have had opportunities of observing its superior efficacy, or of describing particular cases of the few disorders I am to mention; on which I shall content myself with giving you the general result of my observations.

The cases which have afforded me the most frequent opportunities of observing, and of drawing the most satisfactory conclusions relative to the superior efficacy of the Red Bark, are those of the Intermittent Fever.

To the best of my recollection, it was about the month of October, 1781, when we first began to make use of the Red Bark, at the London Hospital. Immediately after its introduction, the difference, in point of efficacy, between this and the common Bark became very remarkable; infomuch, that my learned
 Colleague,

Colleague, Dr. DICKSON, and myself, recommended to the Committee of the Hospital, at one of their weekly meetings, to purchase of the Druggist, who had furnished the first specimen, the whole of his stock of the medicine, however great its quantity might be: upon which measure the Committee, without any hesitation, immediately resolved.

The London Hospital is, perhaps, never without a very considerable number of patients under Intermittent Fevers; to which its comparative vicinity to the county of Essex not a little contributes.—When, before the introduction of the Red Bark, we were in the use of employing the common Bark upon all occasions, we had found it, in Intermittents, to fall exceedingly short of that high character for efficacy, which is assigned to it by Dr. SYDENHAM, and his cotemporaries.

As

As to myself, I can truly assert, that in the cases of patients under Intermittent Fevers in the Hospital, very seldom indeed was the return of the fit prevented, or even the violence of it much diminished at the first attempt to stop it, by any quantity of the medicine given in the interval. On the contrary, portions of the Bark for a considerable length of time, and in very large quantities, were generally necessary to stop the progress of the disorder, or even to abate its violence; and on many occasions, from a total want of success, I have judged it proper to desist from its farther use, and to have recourse to other means of cure.

On the other hand, by the use of the Red Bark, I have frequently seen the return of the fit intirely prevented upon the first trial of the medicine given in the interval; where this is not the case, the
subsequent

subsequent fit is generally less violent, and in almost every case the disorder generally disappears in a short time.

Several of the cases in which I have observed the abovementioned good effects of the Red Bark, were cases in which the common Bark had previously been employed, and continued for a longer or shorter time, without success.

With respect to the out-patients, or such as do not reside in, but occasionally come to the Hospital for advice and medicines, these are much more numerous than the in-patients; among which there occur a great variety of Intermit-
tents, in all its different types. The superior efficacy of the Red, compared with the common Bark, I have found to be as remarkable in these cases as in those of the in-patients.

Whilst

Whilst I was in the practice of employing the common Bark, very large doses of it were generally necessary to the cure, and the patients used to return to the Hospital again and again, for repetitions of their medicine; but since I have used the Red Bark, many of these patients have not returned a second time. Of such as have returned, some have informed me that the dose prescribed to be taken during the first interval had intirely prevented the return of the fit; others, that the subsequent fits had been very moderate. And where the cases have been otherwise, and I have been informed of the event, the disorder has given way in a short time.

I shall trouble you with only one other tribe of diseases, in which I have had occasion to observe the superior efficacy of the Red Bark. These are pe-

L riodic

riodic pains, of which diseases, the periodic Head Ach is the most common, and most generally known.

But I meet pretty frequently with cases of a similar disorder affecting various other parts of the body than the head. In some of these, the seat of the pain seems, to the feelings of the patient, to be some part of the parietes of the abdominal, or thoracic cavity, but more frequently of the former; in others it seems to be some or other of the contents of one of these cavities, but more frequently of the abdominal.

With respect to the cases in which the disorder seems to be seated in some of the contents of the abdominal cavity, I have met with many of them, and with some that, during the paroxysm, have been accompanied with a sharp

sharp pyrexia, a most acute pain, and most, or all of the essential, or characteristic symptoms of an inflammation of the viscus; which, from the place of the pain, seemed to be the seat of the disorder.

These cases, however, differ from inflammations in this, that their paroxysms are succeeded by intermissions, and return at intervals, generally of the *Quotidian*, sometimes of the *Tertian*, at other times of less regular types; and blood taken from the patient during the utmost violence of the fit, is without the smallest appearance of the fize.

In many cases of these disorders, where our view is to prevent the returns of the fits, by remedies employed in their intervals, tonics, undoubtedly, are not always proper remedies; some of them, as in particular the periodic Head Ach,

when it occurs in young and plethoric subjects, may sometimes require the use of evacuants; but in those cases in which I have judged tonic remedies to be indicated, and have employed the Red Bark, I have found its effects, compared with those of the common Bark, to be similar to those I have above described, respecting Intermittent Fevers.

In support of my opinion of the superior efficacy of the Red Bark in the diseases I have above specified, I have been led from the nature of the subject, to draw my arguments principally from cases of hospital patients, as these, on account of their superior number, afford the better opportunities of comparison; but the observations I have made in private practice upon these, not to mention other diseases, correspond with and tend to support the same conclusion.

You

You desire to have my opinion relating to the nature of the Red Bark, from what tree it is taken, whether from that which affords the small Quilled Bark commonly in use, or from a different one. In my opinion there can be very little doubt, but that both are taken from the same tree, and that their difference in appearance depends on this circumstance only, that the small or Quilled Bark, is taken either from very young trees, of which all the parts are yet small, or if ever taken from large, is the produce of their twigs or very small branches; and that on the other hand the Red Bark is taken from well-grown trees, and from their trunks or larger branches.

Certain arguments which I find in your publication on the Red Bark, some of which are suggested by yourself, and others, by some of your correspondents, are, I think, sufficient to warrant our

resting in this conclusion. The principal reasons from which I have been led to adopt it are the following:—

First, Because the Red Bark agrees in its sensible and other qualities, with the small Quilled Bark, possessing however these qualities in a much higher degree.

Second, Because it is very well known, that the peculiar sensible qualities and powers of most vegetables, are comparatively little observable in the young plants, or tender shoots.

Third, Because the Tanners know very well from experience, that the Oak Bark which is taken from the trunk or larger branches of the tree, possesses much stronger powers than that taken from the smaller branches; and therefore always prefer this in the business of tanning.

Lastly,

Lastly, What appeared to me upon examining the specimens, you lately shewed me, of Oak Bark, which afforded me an opportunity of comparing the Bark of the trunk, or larger branches of the Oak, with that taken from the smaller branches, where the Bark of the larger kind appeared of a red hue, and expanded, that of the smaller, pale and quilled; a difference exactly similar to that we observe between the two different species of Peruvian Bark.

The only other observation I shall make relating to the Red Bark, is, that when we reflect on the very extraordinary virtues ascribed to the Peruvian Bark, by Dr. SYDENHAM and Dr. MORTON, and particularly on the degree of certainty with which it is by them affirmed to have cured Intermittent Fevers; of which virtues, the small and Quilled Bark is allowed by all Practitioners, to fall so very

far short; and consider further, that the descriptions given by the *Materia Medica* writers, cotemporary with the eminent persons now mentioned, of the Peruvian Bark then in use, does not apply to the small and Quilled Bark, but does exactly so to the Red Bark; and lastly, that the inhabitants of New Spain, and, if I am rightly informed by a gentleman lately arrived from that country, of Old Spain also, actually hold the Red Bark in higher estimation, it must appear highly reasonable to conclude, that the Red Peruvian Bark is truly the species of this medicine, the virtues of which are so much extolled by Dr. SYDENHAM and Dr. MORTON, and which was in common use with them and their cotemporaries.

To conclude, Sir, I consider the work in which you are engaged as highly commendable. I consider it as one not only
of

greater public utility, inasmuch as it tends to fix the reputation, and extend the use of a most efficacious and important medicine; but also as a necessary one to counteract the endeavours of prejudiced or interested persons to oppose its deserved reputation, and discourage its general use.

I am, Dear Sir,

Your sincere Friend,

And humble Servant,

JAMES MADDOCKS.

London, Capel Court, Sept. 20, 1782.

DR. KEIR, Physician to St. THOMAS'S HOSPITAL, informs me, that in that Hospital about 150lb. of the Red Bark have been used, and he thinks with more success than is usually experienced from the common Peruvian Bark.

The

The employment of it in his practice has not been confined to Intermittents; he has also used it in Mortifications, in Phagedænic Ulcers, in the Convalescence of Fevers, and in every other complaint that occurred, where the common Bark would have been deemed a proper remedy.

In opposition to the objection stated, and refuted by me, Dr. KEIR observes, that during the whole of this extensive and miscellaneous use of the Red Bark, no case occurred in which there was reason to believe any bad effects to have been produced by it.

Extract of a Letter from Mr. SHIREFF, an eminent Surgeon and Apothecary at Deptford in Kent.

After observing that the situation of Deptford, and its environs, renders the inhabitants

inhabitants of that village extremely subject to Intermittent Fevers of a very obstinate nature, some of which he found more difficult to cure, than even such as he had seen on the west coast of Sumatra, where they put on a more formidable appearance than in Europe. He proceeds as follows:

“ From the several patients whom I
 “ attended, I selected the three following
 “ cases to try the effects of the Red Bark;
 “ not to enumerate every particular, I
 “ shall only observe, that in all of them
 “ the common Bark had failed in a sin-
 “ gular manner. Each of them had suf-
 “ fered frequent relapses, the first case
 “ especially; she had scarce any respite
 “ for the space of nine months in Lin-
 “ colnshire; she had removed to this
 “ place to try the effects of a different
 “ air; before I saw her she had been here
 “ for three months, without finding any
 “ benefit.

C A S E

C A S E I.

A gentlewoman of a weak constitution, and naturally of a nervous habit of body, had been seized last autumn with a simple *Tertian* in Lincolnshire; upon her removing to this place it had assumed the *Quartan* type, and was of three months standing; having found very little relief from Bark and other remedies, she had declined calling in any assistance; but her husband, alarmed at her extreme weakness during a paroxysm, sent for me: I was informed of the above particulars, and found her with an icteric countenance, swelled ancles, and other marks of great debility: after some difficulty, I prevailed upon her to take medicines; I sent her immediately several doses of Red Bark, each containing only one scruple, on account of her nauseating every thing that was presented her, desiring

firing her to begin after a general moisture had come on, with an abatement of thirst and Head Ach, and to be repeated every four or six hours; she continued the medicine in this manner, for four or five days, and as the subsequent paroxysm had been more mild than the one preceding it; I could not prevail upon her to take the Bark so frequently; she however continued its use for fourteen days longer, each day taking four scruples, which entirely removed her complaints—she is now in perfect health.

C A S E II.

A young gentleman, naturally of a robust and healthy habit of body, had suffered several paroxysms of a double *Tertian* to attack him, without using any method to prevent them; it was his determination to trust to nature for a cure, rather than take such large, and frequent
doses

doses of the Bark, as he had some months before done, without any permanent effects. But a delirium seizing him in one of the paroxysms, his relations sent for me at midnight: I found him sensible, his body covered with a most profuse sweat, and loaded with bed-cloaths, the curtains drawn close, and the external air carefully excluded from the room; having removed every obstacle to the free admission of air, and his body wiped with a dry cloath, and in place of hot drinks, made still hotter with spice, I ordered toast and water, acidulated with lemon, and other diluting liquors, to be given him almost cold; I procured his consent to make one more trial of medicine; accordingly, half a dram of Red Bark was given him immediately, and repeated every four hours; he missed the next period, and after continuing his medicine three days longer, only three times in the day; he left it entirely off;

—he has suffered no relapse, and is now in health.

C A S E III.

A gentleman, after exposing himself to a damp evening, was seized with the common symptoms of Fever; an emetic was given immediately, and followed by a laxative, not apprehending that he was attacked again with an Intermittent, he continued to go to London; at the expected period, however, he was again taken ill, and the paroxysm was rendered very severe, by his imprudently walking home after it had commenced. When the febrile symptoms abated, and the skin became moist, I gave him half a dram of the Red Bark, with orders to repeat it every three hours; he suffered no return, and now remains well; being much exposed to the weather,

ther, I have advised him to continue small doses of it twice in the day.

I am,

Dear Sir,

Your obliged and humble Servant,

Deptford, J. L. SHIRREFF,
Sept. 14, 1782.

To Dr. Saunders.

A Second Letter from Dr. FOTHERGILL to Dr. SAUNDERS.

DEAR SIR,

As medical attention has of late been deservedly engaged on the subject of the Red Peruvian Bark, and as the public are not a little interested in the result, I take the liberty to submit to your consideration

deration, a few more cursory observations, which have occurred since my last.

The testimony which you have already produced from so many respectable Practitioners, who could be under no temptation, either to conceal its failures, or to exaggerate its virtues, renders it unnecessary to add any fresh evidence of its superiority; otherwise I might mention some late instances of its success, in certain inveterate Agues, which had entirely baffled the ordinary Bark. Therefore, whatever doubts or difficulties may be now raised concerning its identity with the Bark formerly used by MORTON and SYDENHAM, can by no means invalidate the facts which have been advanced in support of its real efficacy. That it was, however, in actual use about the beginning of the present century seems demonstrable.

In the year 1702, the cargo of Bark which was captured on board a Spanish galleon, a parcel of which fell into the possession of Mr. Pearson, an eminent apothecary in the city only four years ago, appears, from every circumstance, to have been no other than the drug now under consideration. But what seems still more worthy our attention, is, that after the space of about 78 years, it should still afford a much stronger decoction than that of the common Bark, and also surpass it in the cure of fevers, and other diseases: an evident proof that this Bark retains its medicinal powers much longer than could have been imagined. In further confirmation of this singular property, and also of its early use in this country, allow me to transcribe a remarkable passage from Dr. Lister, who mentions some of its most characteristic marks: “ Propria experientia testor, me
 “ ante 20 annos CORTICE TRUNCI saepe
 “ usum

“ usum esse ad CRASSITIEM, & LATITU-
 “ DINEM VOLÆ MANUS, MAGNIS, &
 “ PROFUNDIS SULCIS, & FISSURIS CON-
 “ SPICUO, velut in VETUSTO ARBORE,
 “ imo eundem aliquando cariosum; &
 “ olim, & nunc, vix unquam frustravit
 “ eventu optimo, & desiderato, maxime
 “ si ejus modus, et tempus exhibitionis
 “ ritè observantur*.” Add to this, a still
 further proof with which I have been lately
 favoured by Dr. Smith, a very ancient phy-
 sician near Andover, who assures me, that
 having obtained a sample of the Red Bark,
 he immediately recognized it, “ both by
 “ the smell, taste, and colour, to be the
 “ SAME that was COMMONLY USED FIF-
 “ TY YEARS AGO.” To which he subjoins
 some recent instances of its success in
 Intermittents, which had resisted the ordi-
 nary Bark.

* De *Hydr.* p. 56.

From the year 1640, that the Peruvian Bark was first imported into Spain, its reputation increased till the old unpeeled trees becoming scarce, the inhabitants of Loxa, mixed other Barks with it, which being detected, it fell into such discredit, that, in the year 1690, several chests of it lay in the warehouses at Piura, and nobody to purchase it. From this circumstance, and from the insignificant doses in which it was administered, it disappointed the public expectation so much, as to be generally discarded, till Tabor, an adventurous English practitioner, by giving more adequate doses of the genuine drug, revived its reputation; when its fame spread so rapidly, that the Spanish merchants, at length, found it difficult to supply the demand of their customers for full grown Bark, and therefore partly through necessity, and partly through political œconomy, substituted the *small* Bark with which they have long furnished the European markets.

Hence

Hence may be explained, why they now affect to extol the Quill Bark, which is more easily prepared, and more readily obtained, in almost any quantity, and that without destroying the trees. M. CONDAMINE, who visited Loxa, about fifty years ago, assures us, that the Red Bark was allowed to surpass the other sorts, but was grown, even then, extremely scarce, on account of the reason already assigned *

Of late years Peruvian Bark has become such an important article of commerce, that our merchants are glad to procure such as is offered; but no candid Spanish Practitioner, who has tried the different sorts, will, it is presumed, be at a loss in determining to which the real preference ought to be given.

* Mem. de l'Acad. des Sc. 1738.

As the present stock of genuine Red Bark cannot but be extremely disproportionate to the demand, it only remains, that we earnestly admonish younger practitioners not to be too precipitate in drawing unfavourable conclusions from the result of their present trials, but to suspend their judgment, till a fresh supply shall enable them to pronounce with more certainty, concerning its comparative powers.

As it hitherto promises to be much superior to the common Bark, in the speedy cure of Intermittents, it will also behove them to use the greater circumspection in ascertaining the true nature, and tendency of the disease, *viz.* whether it is a primary, or only a secondary affection, whether certain obstacles are not previously to be removed; whether the cortex is not contraindicated; and, lastly, whether the sudden suppression of periodical

Edical motions may not prove productive of some more dangerous derangement in the system.

From the present indiscriminate use of the Peruvian Bark, in diseases so diametrically opposite in their nature, I cannot help thinking, that the inactivity of this universal Catholicon, so generally lamented of late, has been rather a fortunate circumstance, and that the inertness of the remedy has often prevented a series of evils, which must have ensued from such a preposterous abuse of the genuine drug.

In Intermittents which are purely idiopathic, and proceed from an epidemic constitution of the atmosphere, without any concomitant disease, or internal inflammation, the Bark may generally, without hesitation, be freely exhibited; and in highly urgent cases of this kind, which

prevail in marshy countries, and sultry climates, wherein the remissions are very short, this medicine can scarcely be administered too soon, or too liberally. Under such hazardous circumstances time is too precious to admit of preparatory evacuations, and I moreover concur with you in considering them as frequently unnecessary, if not injurious.

On the contrary, it must be allowed, that Agues are sometimes merely symptomatic of some other more dangerous affection, and ought to be considered by the attentive Practitioner, as *remedies*, rather than diseases. Thus in the gout, the pain, inflammation, and tumefaction of the toe, is not the principal disease, but a critical *metastasis*, in order to its cure; so febrile paroxysms are, in certain cases, to be considered as the salutary efforts of nature, to subdue some morbid cause, or to remove some confirmed disease

fease, of a more fatal tendency. In such cases, nature is to be assisted, not disarmed of these useful weapons, by which she sometimes combats Palsies, Epilepsies, and other Herculean maladies, which all the artillery of medicine could otherwise never have subdued.

This being accomplished, the febrile paroxysms either subside spontaneously, or may be now safely removed by this powerful febrifuge.

On the other hand, there are not wanting instances, where the supervening Ague is so far from removing the former disease, that it serves but to exasperate its symptoms, and if suffered to continue, to produce still other dangerous affections. To determine with accuracy and precision in these different situations, demands a degree of medical discernment

discernment and deliberation, which but too seldom occur in the hurry of modern practice.

With respect to the general operation of the Peruvian Bark, I entirely coincide with what you have so satisfactorily advanced, and shall only presume to subjoin the following reflections.

Physicians, in attacking Putrid Fevers and other obstinate diseases with the Bark, seem extremely solicitous to impregnate the whole mass of fluids with its specific virtue, yet excellent as it is, when applied to the nervous surface of the alimentary canal, nature seems to me, never to have intended that it should enter the blood, and has therefore wisely placed firm barriers to prevent its admission into the interior parts of the machine. A substance which is capable of undergoing repeated macerations, and decoctions in water for many months,

months, without being wholly divested of its bitterness and astringency, could not easily be subdued in the blood-vessels; but would probably prove (at least in its native state) utterly incompatible with the laws of the system. Dr. FRIEND accordingly informs us, that no sooner had he injected two ounces of a decoction of this medicine into the jugular vein of a dog, than it produced severe palpitations, convulsions, and death.* The Provident Guardian of the human frame, thus kindly checks the wild career of aspiring mortals, when, through the mists of boasted science, they blindly pursue devious paths that often lead to dangerous errors. Not that our researches into the laws of the œconomy, and the operation of medicines can be too deep, or prosecuted with too much ardour, so long as we follow the clue of accurate observation, and draw no conclusions but what are fairly

* Emmenalog. c. xiv.

fairly deducible from the phenomena; but unfortunately, from the little we know of these matters, we often presume a great deal concerning the major part which remains unknown, and the misconceptions which thence ensue in theory, are transferred into practice.

In the late German war, the French army on their return from Bohemia, were seized with *Tertian* Agues of the putrid kind, which at length terminated in critical abscesses, which formed behind the ears, and in the arm-pits. When these abscesses were fully matured, they were opened according to the usual mode of practice in similar cases. But no sooner was this unfortunate operation performed, than the symptoms recurred, accompanied with extreme prostration of strength, under which the patients generally sunk in a few days. But when the sick were left to Nature's own management, without

any

any attempt to promote, or retard sup-
puration, or to open the abcesses, the pe-
rulent matter was spontaneously discharged
by the intestinal canal, or some of the
other emunctories. The practitioners
were now led to acquiesce in the mode of
cure pointed out by Nature, and from
this time, almost all who were affected
with the disease recovered.*

Thus art often boldly usurps the province
of nature, and undertakes to regulate the
inordinate motions of a complicated ma-
chine, and not unfrequently by very im-
proper, or very inadequate means. From
this source proceed innumerable errors in
the treatment of diseases, and endless mis-
takes concerning the effects of medicines.
Intermittent Fevers, and the Bark, the
subjects now under discussion, afford preg-
nant examples of both. Nor can any re-
formation be expected till more attention

i2

* Mem. de l'Acad. royale des Scien. de Stockholm.

is paid to that excellent maxim of the Illustrious VERULAM:

“ Non fingendum, aut excogitandum, sed invenendum quid NATURA faciat, aut ferat.”

I remain, Dear Sir,

Your most obedient Servant,

A. FOTHERGILL.

London, Sept. 20, 1782.

I hope I shall not be considered as presumptuous, in concluding, that the foregoing observations are sufficient for determining the superior efficacy of the Red Peruvian Bark. And it seems reasonable to hope, that the introduction of this kind of Bark may be attended with the happiest effects, and enable us to oppose more successfully those malignant and remittent fevers of warm climates, and unfavourable situations so destructive to our fleets and armies.

I cannot, however, finish, without returning my best thanks to the gentlemen

men who have favoured me with their correspondence; and I think it necessary to express my obligations to many gentlemen whose letters were too late for publication.

They all concur in recommending the Red Peruvian Bark, as more efficacious and powerful than any other kind.

From the numerous trials I have made with it, in Intermittent Fevers, and other diseases, I am disposed to conclude, that it need be employed only in half the quantity we generally recommend of other Bark.

I have likewise the satisfaction of assuring my readers, that it is now in general use in all the large Hospitals in London; and such is the preference given to it, that the demand is difficultly supplied.

Be

Be careful in the choice of it, by attending to the characters which distinguish it from the large Bark, hitherto rejected by our Druggists.

I shall continue to be diligent in my enquiries on the subject, and I most earnestly request the favour of my friends, that they will persevere in supplying me with accurate observations, so as to determine, with precision, in what other diseases this valuable remedy may be used with safety and advantage.

The operations and effects of remedies can only be ascertained by the united industry and experience of intelligent men; who, by being aware of the difficulty of making observations, are sufficiently guarded against the fallacy to which they are unavoidably exposed.

F I N I S.



