Aphorisms on the application and use of the forceps and vectis: on preternatural labours, on labours attended with hemmorhage, and with convulsions.

#### **Contributors**

Denman, Thomas, 1733-1815. University of Leeds. Library

#### **Publication/Creation**

London: E. Cox, 1815.

#### **Persistent URL**

https://wellcomecollection.org/works/q9hpbxyk

#### **Provider**

Leeds University Archive

#### License and attribution

This material has been provided by This material has been provided by The University of Leeds Library. The original may be consulted at The University of Leeds Library. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org

# APHORISMS

ON THE

## APPLICATION AND USE

OF THE

# FORCEPS AND VECTIS;

ON PRETERNATURAL LABOURS;

ON LABOURS ATTENDED WITH HEMORRHAGE,

AND WITH CONVULSIONS.

FIFTH EDITION.

# BY THOMAS DENMAN, M.D.

CIANS, LONDON; AND HONORARY MEMBER OF THE ROYAL MEDICAL SOCIETY AT EDINBURGH.

## London:

PRINTED FOR E. COX AND SON, ST. THOMAS'S STREET, BOROUGH.

1815

# APHORISMS

THE RO

APPLICATION AND USE

23 5 50

# PURCEES AND VECTIS:

PRUDENT ALBUTTANTERIANT MOURS

STANSONSH BUTTH ON SUBSTIL ESTONAL NO

SHOP WHEN COMPANY THE COMP

MOTHER RESERVE

HILE WATER TO BAME IN THE

CHARL TO RESTRUCT AND MEMORARS MEMBERS OF THE COLUMN TO A THE

: 15,0

CANNOT STATE AND ADDRESS OF HOS OFFICE AND ADDRESS OF THE PARTY OF

## ARRANGEMENT

OF

# LABOURS.

# FOUR CLASSES.

- I. NATURAL.
- II. DIFFICULT.
- III. PRETERNATURAL.
- IV. ANOMALOUS, or COMPLEX.

# CLASS I. NATURAL LABOURS.

CHARACTER. Every labour in which the process is completed within twenty-four hours, the head of the child presenting, and no adventitious assistance being required.

## VARIETIES.

- 1. The face inclined towards the sacrum.
- 2. The face inclined towards the ossa pubis.

- 3. The head presenting with one or both arms.
- 4. The face presenting.

That part of a child which descends lowest into the pelvis, is to be esteemed the presenting part.

Circumstances attending Labours.

- 1. Anxiety.
- 2. Rigours.
- 3. Strangury.
- 4. Diarrhœa.
- 5. Mucous discharge, with or without a mixture of blood.
- 6. Pain.

# Causes of pain.

- 1. Expulsatory action of the uterus.
- 2. Resistance made to the effect of that action.

# Distinctions of pain.

- 1. True.
- 2. False.

Causes and signs of false pain. Means of removing them. Digitized by the Internet Archive in 2015

Means by which true pains are supposed to be regulated, and their effect promoted.

Note. The pains attending labour are subsequent to the action of the uterus, though in common language the word pain, and the action of the uterus, are used synonymously.

# Progress of natural labours.

Three periods or stages.

1st period.

Dilatation of the os uteri.

Rupture of the membranes.

Discharge of the waters.

2d period.

Descent of the child.

Dilatation of the external parts.

Expulsion of the child.

3d period.

Separation of the placenta.

Expulsion or extraction of the placenta.

Note. It very often happens that the membranes do not break till the head of the child is on the point of being expelled. This

los io ena dia galacoreg krod

4. The face precenting.

Thet part of a child which decreads lowers into the present-

Committee of the street Laboury.

. Annuna .

Migount

d. Strategury

eministici .

in interest discharge, with or without a mixture of bisod.

G. Pain.

Carriery by prain.

I. Sispulsatory action of the afferen.

C. Resistance made to the thest of that

Distinctions of quia

Jone'l Jirae.

c. False.

Causes and signs of felor pain.

Means by which true pains are supposed to be regulated, and their effect promoted.

NOTE. The pains attending labour are subsequent to the action of the uterus, though in common language the word pain, and the action of the uterus, are used synonymously.

Progress of natural labours.

Three periods or stages.

1st period.

Dilatation of the os uteri.

Rupture of the membranes.

Discharge of the waters.

2d period.

Descent of the child.

Dilatation of the external parts.

Expulsion of the child.

3d period.

Separation of the placenta.

Expulsion or extraction of the placenta.

NOTE. It very often happens that the membranes do not break till the head of the child is on the point of being expelled. This

is the natural and most desirable progress of a labour, and it is a negative proof that the labour has been well conducted; that is, not interrupted. But the description given above, will answer the purpose of impressing a clear, general idea of labours.

The two circumstances which principally require attention in natural labours are, to guard the perinæum and to extract the placenta with discretion.

### CLASS II. DIFFICULT LABOURS.

CHARACTER. Every labour in which the process is prolonged beyond twenty-four hours, the head of the child presenting.

Note. Some objections may be made to this definition taken from time, but it will be found to apply to practical uses better than if it was taken from circumstances.

It would often be extremely difficult to say with precision when a labour actually begins, because

And the second of the second o

The time established formers in moderate established to the state of the polynomial formers and to the polynomial to the

Mitana, day

CHARACTERS INVESTIGATE IN A SOURCE OF STREET COMPANY STREET STREET STREET STREET STREET STREET STREET STREET STREET STREET

house, the board of the child proventing.

If a real and some objections may be seed to the little of the desired to the seed of the seed

It would often be extranely lifficult to any with precision when a bassur asterily legins, because

because of the number of concurrent changes. But in general some progress must be made before we can allow a labour to be commenced.

### FOUR ORDERS.

#### ORDER I.

Labours rendered difficult from the inert or irregular action of the uterus.

#### CAUSES.

- 1. Too great distention of the uterus.
- 2. Partial action of the uterus.
- 3. Rigidity of the membranes.
- 4. Imperfect discharge or dribbling of the waters.
- 5. Shortness of the funis umbilicalis.
- 6. Weakness of the constitution.
- 7. Fever.
- 8. Want of a due degree of irritability.
- 9. Passions of the mind.
- 10. General deformity.

segment in a reduction of the segment of the segmen

AREST ORDERS.

I REVED

The trans of the same of the s

Too peak detention of the steems.

C. Period extend of the plane.

E. English vi the mothermen.

d. Laperior double pe or well-bling of

in the state of the facilitation.

F. Ware of a deep degree of instability.

130,00

#### ORDER II.

# Labours rendered difficult by the rigidity of the parts to be dilated.

- 1. First child.
- 2. Advancement in age.
- 3. Too early rupture of the membranes.
- 4. Oblique position of the os uteri.
- 5. Fever or local inflammation.
- 6. Extreme rigidity of the os uteri.
- 7. Uncommon rigidity of the external parts.

#### ORDER III.

Labours rendered difficult from disproportion between the dimensions of the cavity of the pelvis and the head of the child.

- 1. Original smallness of the pelvis.
- 2. Distortion of the pelvis.
- 3. Head of the child unusually large, or too much ossified.
- 4. Head of the child enlarged by disease.
- 5. Face inclined towards the ossa pubis.

6. Presentation

onning II.

Labour vandayd shifted by the rigidity of the

dible teris .

e. Advancement in ago.

S. Too saily rupture of the members.

Culling position of the or stee

Teres or local infammation.

de Pentrate tiglifig et die et aler

the plant state out to abbinit an annual to

API MATERO

Labours readered eightailt from slight sparties hetween the dissensions of the carily of 'the

1. Original smalligers of the period.

a, these of the child southently large, of

.bahino dance on

A. Hoed of the child enlarged by uncorn.

G. Presentation

- 6. Presentation of the face.
- 7. Head presenting with one or both arms.

#### ORDER IV.

Labours rendered difficult by diseases of the soft parts.

- 1. Suppression of urine.
- 2. Stone in the bladder.
- 3. Excrescences of the os uteri.
- 4. Cicatrices in the vagina.
- 5. Adhesion of the vagina.
- 6. Steatomatose tumours.
- 7. Enlargement of the ovaria.
- 8. Rupture of the uterus.

Note. The disturbance of the natural progress of labours, more especially the premature rupture of the membranes, is the most general cause of difficulties in parturition.

Women are to be relieved in difficult labours,

- 1. By time and patience.
- 2. By encouragement to hope for a happy event.

Consider they discontinued in the contract Land of

- 3. By regulating their general conduct.
- 4. By lessening or removing the obstacles to the effects which should be produced by the pains.
- 5. By the assistance of instruments.

Intentions in the use of instruments.

- 1. To preserve the lives both of the mother and child.
- 2. To preserve the life of the mother.
- 3. To preserve the life of the child.

Instruments contrived to answer the first intention.

1. Fillets. 2. Forceps. 3. Vectis.

Three things are to be considered with respect to the Forceps or Vectis, and to the use of instruments in general.

- 1. To make an accurate distinction of those cases which require their use.
- 2. Of those cases which allow their use.
- 3. Of the manner in which they ought to be used.

We are in the first place to speak of the application and use of the forceps.

Directions

(8)

a. By regulating their general conduct.

A. the leavening or removing the obstacles

to the effects which should be pro-

c. By the pointance of instruments.

Jetentions in the use of her remedie.

I. To preserve the lives both of the mellion and the mellion

To preserve the his of the mother. To preserve the life of the child.

Latenments contribed to grant the first intension.

Three things are to be confidered with reof cet to the Forega or Fatte, and to the me
of instruments in general.

1. To make an accurate distinction of these
2. Cases which require their tax.
2. Of those came which allow their use.
3. Of the summer in which they ought to but
3. Tested

We are in the first placesto speak of the application and use of the favores.

Directions

Directions for, and admonitions in, the application and use of the Forceps.

#### SECTION I.

- 1. It has long been established as a general rule, that instruments are never to be used in the practice of midwifery; the cases in which they are used are therefore to be considered merely as exceptions to this rule.
- 2. But such cases can very seldom occur in the practice of any one person; and when they do happen, neither the *forceps* nor any other instrument is ever to be used in a clandestine manner.
- 3. The first stage of a labour must be completed, that is, the os uteri must be dilated and the membranes broken, before we think of applying the forceps.
- 4. The intention in the use of the forceps is, to preserve the lives both of the mother and child, but the necessity for using them must be decided by the circumstances of the mother only.

at beaut all at the he of a Warmen I mile play call by constanting out with his chart the

5. It is meant, when the forceps are used, to supply with them the insufficiency or want of labour pains; but so long as the pains continue, we have reason to hope they will produce their effect, and shall be justified in waiting.

6. Nor doth the cessation of the pains always prove the necessity of using the forceps, as there may be a total or a temporary cessa-

tion of the pains.

7. In the former, the pulse, the countenance, and the general appearances of the patient indicate extreme debility, and resemble those of a person worn out with disease or fatigue.

8. But in the latter there are no alarming symptoms, and the patient often enjoys short

intervals of refreshing sleep.

9. A rule for the time of applying the forceps has been formed from the following circumstance; that, after the cessation of the pains, the head of the child should have rested for six hours in such a situation as to allow the use of the forceps before they are used.

A. It is meant, when the prompt on and the stands of a second stands of the second stands of

of Mor thick the constion of the prime of the foreign.

The sale of the process of the particle of the particular country that indicate extrane debilling and rescalds there are the those of a person wars out with discount or in-

B. But in the latter there, are no slaveling symptoms, and the parlient offen or juye short istured to form of the parlies of the solution of

countracte; that he class of applying the for-

10. 'But

- 10. But this and every other rule intended to prevent the rash and unnecessary use of the forceps, must be subject to the judgment of the person who may have the management of any individual case.
- 11. Care is also to be taken that we do not, through an aversion to the use of instruments too long delay that assistance we have the power of affording with them.
- 12. The difficulties which attend the application and the use of the forceps are far less than those of deciding upon the proper time when, and the cases in which, they ought to be applied.
- 13. The lower the head of the child has descended, and the longer the use of the forceps is deferred, the easier will in general their application be, the success of the operation more certain, and the hazard of doing mischief less.
- 14. The forceps should always be applied over the ears of the child; it must therefore be improper to apply them when we cannot feel an ear.

can but to see all amond the property of

- 15. But when an ear can be felt by a common examination, the case is always manageable with the *forceps*, if the circumstances of the mother require their use.
- 16. The ear of the child which can be felt, will be found towards the ossa pubis, or under one of the rami of the ischia.
- 17. The ears are not turned to the sides of the pelvis till part of the hind head has emerged under the arch of the ossa pubis, when the use of the forceps can very seldom be required.
- 18. When you have determined on using the forceps, and explained the necessity of using them to the patient and her friends, she is to be placed in the usual position on her left side, near to the edge of the bed; and the instruments, warmed in water and smeared with some unctuous application, are to be laid conveniently by you.

Note. Women, impelled by their fears and their sufferings in difficult labours, will very frequently implore you to deliver them with instruments long before you will be convinced of the necessity of using them. In

the our of the child which our be fell. bringer of molific gray posterior be required. the places of the rotal words to washing aderror bind od of are caretorilyga montonit outseries by the parties of the series there when the property berg he and supported the conmany cases I have found it expedient and encouraging to them to fix upon some distant time when they should be delivered, if the child were not before born; six, or eight, or twelve hours, for instance. In some cases of great apprehension I have also shewn them, upon one of my knees, all that I intended to do with the forceps.

The following rules are given on the presumption that the head of the child presents with the face inclined or verging towards the hollow of the sacrum, and that the common short forceps are intended to be used; but if any other kind of forceps should be preferred, the rules must be adapted to the instrument.

#### SECTION II.

- 1. Carry the fore finger of the right hand to the ear of the child.
- 2. Then take the blade of the forceps to be first introduced, by the handle in the left hand, and conduct it between the head of the child

C

( 01

couraging to them to fix upon some distant time when they should be delivered, if the child were not before born; six, or eight, or twelve hours, for instance. In some cases of great apprehension I have also shown them, upon our of the times, and there also shown them, upon our of the times, and there also shown them,

For many that the head of the child presents with the face head of the child presents with the face has inched or veryon towards the hollow of the restaur, and that the common thout foreign the pathod to be need; but is the common that it is a start of the preferred, and other that of foreign should be preferred, the rules must be adapted to the instrument.

Werry at

to the car of the chills.

<sup>2.</sup> I non take the blade of the forego to be not introduced, by the handle in the left hand, and conduct it between the hand of the child

and the finger already introduced, till the point reaches the ear.

- 3. The farther introduction must be made with a motion resembling a slight degree of semi-rotation, and the point of the blade must be kept close to the head of the child, by gently raising the handle as the instrument is advanced.
- 4. The blade of the forceps must be carried up till the lock reaches the external parts, near the inferior edge of the ossa pubis.
- 5. Should any difficulty occur in the introduction of either of the blades, we must withdraw them a little, to discover the obstacle, and never strive to overcome it with violence.
- 6. When the first blade is introduced, it must be held steadily in its situation, as it will be a guide in the introduction and application of the second blade.
- 7. The second blade of the forceps must be conducted upon the fore finger of the left hand, passed between the head of the child and the perinæum, in the same cautious man-

C 2

F

ner as the first, till the lock reaches the perinæum, or even presses it a little backward.

- 8. When the second blade is properly introduced, its situation should be opposite to the first.
- 9. In order to lock the forceps, the handles of which are at a considerable distance from each other, the blade first introduced must be brought down and carried so far back that it will lock with the second blade, held in its first position.
- 10. Care should be taken that nothing be entangled in the lock of the forceps, by carrying the finger round it.
- 11. It is convenient to tie the handles of the forceps together, when locked, with force sufficient to keep them from sliding or shifting their position.
- 12. If the blades of the forceps were introduced so as not to be opposite to each other, they could not be locked.
- 13. Should the handles of the forceps when applied come close together, probably the bulk of the head is not included between them,

C 3

and

the near up the lock resulted the periserious promes it a main backward.

odi ol encogno si klanda ngularne sa te cho

and others are as a commission of the fields to an about the same failes to an about the same failes and commission of the same failes.

ed quidlon, sair reder to binon soir .01

the terripe tiggeness when located, with force

duced as us not to be opened to the cake, other,

and were it as to be adjusted to be been a single and a s

and therefore when we acted with them they would slip.

14. If the handles when locked are at a great distance from each other, they are not well applied, and will probably slip.

15. But in these estimations allowance is to be made for the different dimensions of the heads of children.

16. The forceps will never slip if judiciously applied, if the case be proper for their use, and we act circumspectly with them.

Note. The difficulties in the application of the forceps arise, from attempting to apply them too soon; from passing them in a hurry, or in a wrong direction; or from entangling the soft parts of the mother between the instrument and the head of the child. Of course, we are always to be guarded against these circumstances.

the part of a part of the part

### SECTION III.

- 1. There is no occasion, and it would be hurtful to attempt to change the position of the head, when the *forceps* are applied, before we began to extract.
- 2. For if the action with the forceps be slow, the head of the child will turn in the same manner, and for the same reasons, as in a natural labour.
- 3. Therefore the forceps being fixed upon the head must also change their position according to its descent, and the handles be gradually turned from the ossa pubis and sacrum, where they were first placed, to the sides of the pelvis.
- 4. The handles of the forceps likewise, though originally placed far back towards the sacrum, that is, in the direction of the cavity of the pelvis, will be gradually turned, as the child advances, more and more towards the pubes, that is, in the direction of the vagina.

- 5. The first action with the forceps must be to bring the handles, firmly grasped in one or both hands, slowly towards the pubes, till they come to a full rest.
- 6. After waiting till the pains return, or an imaginary interval if there should be a total want of pain, the handles are to be carried back in the same slow and cautious manner till the lock reaches the *perinæum*, using at the same time a certain degree of extracting force.
- 7. The subsequent actions must be from handle to handle, or occasionally by simple traction; but the action of that blade which was towards the *pubes*, must be stronger and more extensive throughout the operation, than the action with the other blade, which has no fulcrum to support it.
- 8. By a repetition of these actions, always directed according to the position of the handles, with their force increased, diminished, or continued, according to the exigence of the case, we shall in a short time perceive the head of the child descending.

DIVIDED OF THE PROPERTY OF THE

After the necessing to the position of the euroalter the necessing to the position of the eurodies, which there increased, to make both and is before the continue of the state of the saigneest with the saigneest with the saigneest of the continue of the saigneest of the sai

- 9. When the head begins to descend, the force of the action with the forceps must be abated, and as that advances, the direction of the handles must change by degrees more and more to each side, and towards the pubes.
- 10. The lower the head of the child descends, the more gently we must proceed, in order to prevent any injury or laceration of the perinæum or external parts, which are likewise to be supported in the same manner as in a natural labour.
- 11. In some cases, the mere excitement occasioned by the application of the forceps, or the very expectation of their being applied, will bring on a return or an increase of the pains sufficient to expel the child without their assistance.
- 12. In other cases we are obliged to exert very considerable force, and to continue it for a long time; so that one operation may be safely and easily finished in twenty minutes, or even a less time, and another may require more than an hour for its completion, and

the main means the dring makes all to serve the many of the main and the server to make t

TO A STORY THE BOOK OF THE STORY OF THE STOR

the ray expectation of their particular of the property of their particular of their p

The state of the s

the repeated exertions of very considerable force.

- 13. In some cases it happens also, that the obstable to the delivery exists at one particular part of the *pelvis*, and when that is surmounted, the remainder of the operation is easy; but in other cases there is some difficulty through the whole course of the *pelvis*.
- 14. Before the exertion of much force, we are always to be convinced that a small or a moderate degree of force is not equal to our purpose.
- 15. In every case in which the forceps have been applied, they are not to be moved before the head is extracted, even though we might have little or no occasion for them.
- 16. When the head of the child is born the forceps are to be removed, and the remaining circumstances are to be managed as if the labour had been natural.

Note. The general arguments against the use of instruments have been drawn from their abuse: it appears, however, that necessity will,

the repeated energing of ways capidaralite

13. In some eases it happens also, that the observable to the delivery exists at one presidenter part of the period, and when that is an amount of the remainder of the remainde

n'n' armid stant to mainten ail quotati et .

n'n to thoma adade beomismo si es esani sur

rou or tanpa tur of constitut de segui-standam

the state of plied, they are fast to be moved be forest the forest the face at the land is extracted, even thought and might have little or to occasion for them.

forces are to be removed, and the remaining

Louis and hed tood

Norse: it appears, however, that necessity will, abuse: it appears, however, that necessity will,

in some instances, justify the use of the forceps; that when such necessity exists, their use is not only justifiable, but often highly advantageous; that delay to apply them, and slowness in their application and use, will secure, as far as is possible, both the mother and child from untoward accidents; but that mischief cannot be prevented if they are applied too soon, or the operation with them be performed in a hurry.

It would be a very desirable thing that every student should have an opportunity of seeing the operation with the forceps performed before he goes into practice; but that is not always possible. Yet if he has been properly instructed in the principles of the application and use of the forceps, reflects seriously before he determines on performing the operation, and proceeds slowly but not timidly in it, he can hardly fail to succeed. Hurry, in any operation, is a very common sign both of want of information and of fear; and attention is to be paid to the order of the rule in Celsus, 1. tutò, 2. citò, 3. jucundè.

### SECT. IV.

# On the application and use of the Vectis

- 1. We shall have a just idea of the vectis by considering it as one blade of the forceps, a little lengthened and enlarged, with the handle placed in a direct line with the blade, that is, without any lateral curvature.
- 2. The general condition and circumstances of labours before stated, as requiring and allowing the use of the *forceps*, will hold equally good when the *vectis* is intended to be used.
- 3. In the application of the *vectis* two fingers, or the fore finger of the right hand is to be passed to the ear of the child.
- 4. Then taking the *vectis* by the handle, or with the blade shortened, in the left hand, conduct it slowly till the point of the *vectis* reaches the ear, however that may be situated.
- 5. The instrument is then to be advanced, as was advised with the forceps, till according to your judgement the extremity of the blade reaches

YI, YOUR

On the applications and use of the fronts

i., Wa shall bayen just idea of the regts by coundering it as out thate of the streets, a little longitiesed and volument, with the bandle placed in a direct line regt the blade, that is, eithout any larged convenience.

In the animal an interest of the second streets and the second streets are second streets and the second streets are second streets and the second streets are second streets and the s

or at the stage of the range out off to any

4. I now telling the week by the handle, or with the blands, convicts the bland short in the left hand, convicts the bland the transfer duct it shows to be able to the converted that can be able to the converted that can be able to the converted.

philosophic and or med to knimerick and the observed guidence of the philosophic contraction of the believe of

reaches as far, or a little beyond, the chin of the child.

- 6. Then grasping the handle of the instrument firmly in the right hand, wait for the accession of a pain.
- 7. During the continuance of the pain raise the handle of the instrument gently but firmly towards the *pubes*, drawing at the same time with some degree of extracting force.
- 8. When the pain ceases let the instrument rest, and on its return repeat the same kind of action, alternately resting and acting in imitation of the manner of the pains.
- 9. By a repetition of this kind and manner of action the head of the child is usually advanced, and the face turning gradually towards the hollow of the sacrum, the position of the handle of the vectis will be altered, and the direction of the action with it of course should be changed.
- 10. When the head is perceived to descend, we must proceed more slowly and carefully, according to the degree of descent, in order to prevent any injury to the external parts, which

( 02 )

sactors as into of a hittle beyond, the cuin of

of Then providing the deadle of the instru-

the blands of the continuence of the paint for the third of the blands o

The content of the point consecrate she instrument to be been from the part of the part of

to present any injury to the restored in descend,
or present any injury to the extense parts,

which is to be prevented, as was directed when the forceps are used.

- 11. But if by the continuance of the moderate force before recommended, the head should not descend, it must be gradually and cautiously increased till it becomes sufficient to bring down the head.
- 12. In the action with the vectis the back part of the instrument must rest upon the symphysis of the ossa pubis, or upon the ramus of the ischium, according to its position, as upon a fulcrum, for its support.
- 13. By passing the flat part of the hand to the back of the blade of the instrument when in action, we shall be occasionally able to lessen or take off this pressure, which must otherways be made upon the parts of the mother.
- 14. Some have recommended the vectis to be used when the head of the child was higher up in the pelvis than is before stated, as justifying the use either of this instrument or the forceps.
- 15. They have also recommended the vectis when the head of the child was firmly locked

which is to be prevented; as was directed when

The flow of the commendate of the modernate to the beat that the second rate the second rate to the second rate the second rate the second rate that the sec

12. In the parties with the retain the back part of the free instrument mast use upon the synchronic as point, or upon the raines of the sisters, according to its position, to

about the start part part of the hond to about the bond to about the start when it were that the start which the start when the start which the start which the start when the start which the start when the start which the

ways be made upon the pasts of the mother.

14. Since helpering calculation is the decide to
be unce when the final calculations calculations singles
up in the salesting final calculations stated, earloadreing the uncesting of this instrument of the

the They have also recommended the rectal when the head of the child was finally lecked in

in the pelvis, and have asserted that by its use there is often obtained a very good chance of preserving the life of a child, which must otherwise be inevitably lost.

- 16. Others have by frequent use acquired such dexterity, as to be able to extract the head of a child, in the situation first stated, with a single sweep of the instrument.
- 17. Some have also advised the introduction of the *vectis* between the *sacrum*, or sacrosciatic ligaments, and the head of the child, from a belief that it could be equally or more advantageously used in this position than in that first stated.
- 18. But having ever considered the use of all instruments as a thing to be lamented, and when I did use them, esteemed the safety of using them as my principal object, I cannot deviate from these principles, or enter upon a discussion of points of practice, of which, as far as I am competent to judge, I cannot approve.

Note. Before, and immediately after the publication of my second Essay on Difficult

not the gooding and their services and arrive out the COULD BE SEED OF THE PROPERTY OF THE PROPERTY OF appear from the shopping and species and aben I did per them, extremel the melegral human them on any principal rate of mail gains society from these presciption or easter upon a

.. Norm. Indiana and impositional survey.

212 Sp. 1

Labours, several gentlemen, with whom I converse, and to whom I ought to pay great respect, reprehended in very decided terms what I have advanced with regard to the forceps and vectis. Some maintained that the forceps is an instrument far superior to the vectis, of which I was accused of speaking too favourably. Others, of equal respectability, accused me of speaking with timidity, or restraint, of those advantages which, they asserted, the vectis had over the forceps. This very strong evidence could only be invalidated by its contradiction, but the very respect which I bear to the witnesses, compelled me to pass over their evidence, and to rely upon my own experience and judgment.

I did not speak of the mechanism of the instruments, or of the operation performed when we had applied, and acted with them, as these have hitherto been very imperfectly and often erroneously explained. The subject came under consideration in the ordinary course of the work, and having frequently used both the instruments, I stated

D 2

the

remember of the appendict performed to the class of these infer historia been very importantly of often expansions of the came mader consideration in the the came caute of the work and harring in-

the matter equitably, according to the best of my abilities, and in such a way that, I thought, students, who were principally concerned in the discussion, being left with the choice of either instrument, according to the doctrines of the particular professors whom they might attend, could not be misled. It is not to be expected that men versed in practice should change their opinions or alter their practice, or, in short, pay much regard to disputes about instruments, if any were disposed to raise them.

It then was, and yet remains my opinion, founded, as I before observed, on my experience with both instruments, that the superior excellence which has been attributed to each of these instruments, ought chiefly to be ascribed to the dexterity which may be acquired by the habit of using either of them. It is also my opinion that we may, in general, either with the forceps or vectis, effectually and conveniently give that assistance which is required in cases of difficult parturition, allowing and justifying their use. In particular

D 3

reside vilage

it then was, and pet requires our apprises, standard, as I before obsciverit, on any expenses with both instruments, that the superior with both instruments, that the superior cought existly to be accept of these instruments, ought existly to be aterior of the obsciring which may be accepted by the habit of using aiting as them, as the my equired by the holds we may, in general, disher with the formation that we may, in general, edition with the formation that we may, in general, conveniently and the means of their particular and although the particular and although their man. In particular they and a their man. In particular they

cases it may perhaps be proved that one instrument is more commodious than another.

But if the vectis be depreciated by those who have never used it, and are not expert in its use, because they prefer the forceps; or if the known properties of the forceps be not allowed by those who do not use them, because they prefer the vectis, the proper inference would not be, that either of the instruments ought to be condemned; but that we are in possession of two instruments well adapted to answer the same purpose, if they are prudently used; or, that neither of them ought to be used.

In those cases, in which the face is turned towards the *pubes*, or in which the face of the child is the presenting part, it is generally more convenient to deliver with the *vectis*, or with one blade of the *forceps*, than with both blades.

come is purposed by proved that configured in the prove communitions than appreciated by these let it it it a prove to despreciated by these site is not because the despreciated by these sees they and the state of the sees of the sees of the sees they appreciate the sees of the sees of the sees of they appear they appear they appear they appear the sees of the see

In these cases, in which the face is turned towards the pyles, is in which the face of the cities in the presenting path, it is gonerally more convenient to deliver with the course, or while one hinds of the players with the course in the deliver with the course in the deliver with their back.

CLASS III. PRETERNATURAL LABOURS.

CHARACTER.—Labours in which any part of the child presents, except the head.

## TWO ORDERS.

### ORDER I.

Presentations of the Breech, or inferior Extre-

#### ORDER II.

Presentations of the Shoulder, or superior Extremities.

### SECTION I.

1. The presentation of children at the time of birth may be of three kinds. 1. With the head.

CLASS III. PRESENCATURAL LARBORS

CHARLOTER, well-bought in which any part of the child promits, except the bond.

ARMONIO ONLY

A RESTRICTION

Presentations of the Bleech, or inforget shifted

AL ERUNO

Correspondent of the Shoulder, or supprint due

### a noissan

1. In a presentation of children at the fone of birth may be exclude kinds. (1.3 With the leads.

- head. 2. With the breech, or inferior extremities. 3. With the shoulder, or superior extremities.
- 2. Presentations of the first kind are called natural, those of the second and third kind, preternatural.
- 3. Preternatural presentations have been subdivided into a much greater variety, but without any practical advantage.
- 4. The presumptive signs of the preternatural presentation of children are very uncertain, nor can it ever be determined what the presentation is, till we are able to feel the presenting part.
- 5. When any part of a child can be felt, we may form our judgment of the presenting part by the following marks.
- 6. The head may be distinguished by its roundness, its firmness, and its bulk.
- 7. The breech may be known by its bulk, by the cleft between the buttocks, by the parts of generation, and by the discharge of meconium.
  - 8. The foot may be distinguished by its length,

head. c. With the birecth, or leferies avermides. c. With the shoulder, or superiors avtromities.

2. Presentations of the first hind according

'S. Preservational preservations these bean enbelopided into a much granter variety, but without now practical estemateign.

the The principles signs of the presentatoral grammings of children are very uncertain, nor can it over he descrimed what the presentation is silver are able to feel the presenting part.

ney form our judgment of the presenting part by the following marks.

o. The head may be distinguished by its

The brunch may be known by its built, by the cleft between the buttocks, by the parts of generation, and by the discharge of meconium.

a, The foot may be distinguished by its length,

length, by the heel, by the shortness of the toes, and the want of a thumb; and the hand by its flatness, by the thumb, and the length of the fingers.

### SECTION II.

On the first Order of Preternatural Presentations.

- 1. In this kind of presentation the breech, one hip, the knees, and one or both legs, are to be included.
- 2. In these presentations it was formerly supposed necessary, as soon as they were discovered, to introduce the hand to bring down the feet, and to extract the child with expedition.
- 3. But, according to the present practice, such labours are not to be interrupted, but allowed to proceed as if the presentation were natural; unless the necessity of giving assistance should arise from some circumstance independent of the presentation.

( 10

isograph by the head, by the shortesses of the term to the head the head the head the head by its flatment, by the themb, and the longill of the flagmen.

# OF MOISONS

On the first Order of Preservatural Presidentium.

t. In this kind of presentation the breach, one hip, the braces, and one or both days, are to be included.

2. To these presentations it was formerly employed necessary, as show as they envadinged verificated, to introduce the hand to bring down the feet, and to extract the child with expedition.

3. But, according to the present practice, and inhours are not to be interrupted, but allowed to proceed as if the presentation were natural; unless the necessity of giving assistance about arise from some circumstance independent of the presentation.

- 4. By acting on this principle, when the breech of the child is expelled by the pains, the parts are sufficiently distended to allow the body and head to follow without any danger from delay.
- 5. But if the feet of the child were to be brought down in the beginning of labour, the difficulty with which it would be expelled or could be extracted, increasing as it advanced, the child would probably die before the woman was delivered, and she would be in danger of suffering mischief.
- 6. In cases of this kind there is also equal reason, when the breech is on the point of being excluded, for our guarding the *perinæum* from the hazard of laceration as in presentations of the head.
- 7. In first labours, the child, unless it be small, will not unfrequently be born dead when the breech, or inferior extremities, present; but in subsequent labours they will usually be born living, if there be no other impediment than that which is occasioned by the presentation.

4. By sexual on this principle, when the breach of the child is sapplied by the pains, the pains, the pains are dedicated to allow the body and head to follow willows any dancers are deleased to follow willows any dancers.

of or ease blists and the beginning of fabour, the confidence of fabour, the british down in the beginning of fabour, the billion of the supplied as a specific for the british of the confidence of the british down the state of the british down man was delivered, and what what was the british down man of the british and the british down the state was dear

for the course of this kind there is also equal teams, when the investi of on the paint of her ing uncluded, for our granding the permanent from the handred of lucuration as in presenta-

7. in first labouts, the child, onless it he small, will not unfrequently be horn dead when the breed when the breed, or inferior extraolties, establish protest he has in subsequent labours they will be united her he had been been other impedienced than that which is exceptantly by the presents.

- 8. The injuries which the presenting part of the child, especially the *penis* and *scrotum*, may sustain will often be alarming, and appear dangerous, but by soothing and gentle treatment, they are soon recovered.
- 9. Should there be reason to think the child dead, or the powers of the mother insufficient to expel it, we must then give such assistance as may be required.
- 10. This assistance must be given with the hand or with a blunt hook or crotchet, hitched in the groin of the child; or, which I prefer, by passing a ligature round the bent part of the child at the groin, with which we can hardly fail to extract it.
- 11. But every assistance of this kind must be given with discretion, and we must first be convinced of the necessity before we interfere.
- 12. Should a child presenting with the breech advance, though slowly, it is better to be satisfied with this slow progress; or, we might break, without much force, the neck of the thigh bone, or separate the bones of the

of the table and which the presenting pairs of the table that the common has and another that the common man and appearance to the common has been and appearance to the common the common that the common that the common the common that the com

O. Should them be reason to think the child iced, or the powers of the monthly insufficient to days it, we must then give such assistance as more be required.

10. This consumers must be given with the least or with the least or with a blunt book or createlet, hitchest in the groin of the chief; or, which I perfect by presing a liquide round, the bend, part of the chief at the grain, with which we can hardly fail to extract it.

is given with discretion, and we must first be convinced of the necessity before we sales.

12. Should a child presenting with the breach breach advance, though rlowly, it is better to be satisfied with this clow progress; or, we haught break, without much force, the neck of the third bone, or separate the bones of the pairs

pelvis of the child, by either of which accidents future lameness would be occasioned.

### SECTION III.

Of the second Order of Preternatural Presentations.

- 1. In this kind of presentation are included the shoulders, the elbows, and one, or both arms.
- 2. In all these presentations we shall be under the necessity of turning the child, but as they may be attended with circumstances widely different, it is necessary to make the following distinctions.
- 3.—I. When the os uteri is fully dilated, the membranes unbroken, or the waters lately discharged, a superior extremity being perceived to present, before the uterus is contracted.
- 4.—II. When the membranes break in the beginning of labour, the os uteri being little dilated.

Annual state of bloom resummed stated only

THE WOLFSTAR

The second tector of Presentings Streets

the shoulders, the elbows, and one, or both

2. In all three presentations we shall be under the certain the child, but under the certain the child, but at they may be actualled with circumstances which with circumstances who wildered, it is percentary to make the collection of the collecti

The encontrained patenties of the fully dilately the encontraint lately dilately and exception of the words lately dilately form of the contraint to progent, before the name in ton-

dependency of labours, the columns hands in the

5.—III. When the os uteri has been fully dilated, the membranes broken, and the waters long discharged, the uterus being at the same time strongly contracted, and the body of the child jammed at the superior aperture of the pelvis.

6.—IV. When, together with any of these circumstances, there is a great disproportion between the size of the head of the child, and the dimensions of the cavity of the pelvis.

### SECTION IV.

On the Cases which come under the first Distinction.

- 1. Whenever there is a necessity of turning a child, the patient is to be placed upon her left side, near the edge of the bed; or sometimes, when we expect or find much difficulty, in a prone position, resting upon her elbows and knees.
  - 2. All the advantage to be gained from any particular

Seed II. Vibra the or after the bean filly distributed, and new designation of the new distribution of the new distribution of the child jointant at the sequence of the child jointant at the sequence of the child jointant at the sequence of the cold.

of carty, types, regeries with any of these circumstances, there is a great dispreparties between the size of the head of the childrania the consessions of the cavity of the privile

# VE MOISSONS

On the Cases which casts earlier the first Di-

I. Whenever there is a necessity of tunding a whild, the propert is to be placed upon her left side, near the adge of the brd; or sometimes, when we expect or independ difficulty; in a prope position, resting squar her elbows and knows.

2. All the advantage to be gained from any

particular position of the patient is, to allow us the free and dextrous use of our hands; the situation of the child not being altered by the position of the patient.

- 3. The os externum is then to be dilated with the fingers reduced into a conical form, acting with a semi-rotatory motion of the hand.
- 4. The artificial dilatation of all parts must be made slowly, in imitation of the manner of natural dilatation.
- 5. The os externum should be amply distended before the hand is carried farther, or its contraction round the wrist will be an impediment in the subsequent part of the operation.
- 6. When the hand is passed through the os externum, it must be slowly conducted to the os uteri, which being wholly or sufficiently dilated, we must break the membranes by perforating them with a finger, or by grasping them firmly in the hand.
- 7. The hand must then be passed along the sides, thighs, and legs of the child, till we come to the feet.

while of all finisher and the soling interior of the standard of the first soling and the soling

hardin oil et mest ai reprinter se mill de arreit hardin et mill de bander sergen et dans de arreit hardin et mest de la grico e transfer et mest de la grico e transfer et mest de la grico e de la g

A. The or assesses about to neeply allelended before the head is curious sarriage, or in executations erwest, the partial, will be an impediment in the subsequencement of the ope-

distributed the bead of passed through the expressured it make be able to conducted to the

or stori, which being whally or sufficiently
dilated, we must break throughnesses by payforesting through with a fidger, or by grashing

F. The hand must their be possed along the sides, thighs, and less of the child, till we come to the fast.

- 8. If both the feet lie together we must grasp them firmly in our hand; but if they are distant from each other, and we cannot conveniently lay hold of both feet, we may deliver by one foot without much additional difficulty.
- 9. Before we begin to extract we must be assured that we do not mistake a hand for a foot.
- 10. The feet must be brought down, with a slow waving motion, into the pelvis; when we are to rest and wait till the uterus begins to contract, still retaining them in our hand.
- 11. When the action of the uterus comes on, the feet are to be brought lower at each return of pain, till they are extracted through the external orifice, and the labour may then be finished, partly by the efforts of the mother, and partly by art.
- 12. If the toes are turned towards the pubes, the back of the child is towards the back of the mother, which is an unfavourable position.
  - 13. But if the toes are towards the sacrum,

the continues the feet his topping by and a property of the common of th

the first we short man might sur metall All with the said that we have the said that t

distributed into the brought down, with a sign of the state of the sta

construct the state of the stat

12. If the topic are topic defined the fine the lasts when the fact the lasts of the fact in an end-content the lasts of the factors in an end-content the fact.

13. Dot if the form next concesse the minute.

of the mother, which is proper; and all other positions of the child must be gradually turned to this as the body is extracting.

- 14. Yet this position of the child is only advantageous when the head comes to be extracted.
- 15. When the feet of the child have passed through the os externum, wrap them in a cloth, and holding them firm, wait till there is a contraction of the uterus, or a pain, during the continuance of which gently draw down the feet.
- 16. When the pain ceases we must rest, and proceed in this manner through the delivery, assisting the efforts of the patient, but not making the delivery wholly artificial.
- 17. When the breech comes to the os exernum, the child must be extracted very slowy through it, and in the proper direction, or here will be danger of lacerating the perinæum.
  - 18. When the child is brought so low that he funis reaches the os externum, a small por-

E 2

tion of it is to be drawn out, to slacken it to lessen the chance of compression, or to prevent the separation of it from the body of the child, or of the placenta from the uterus; and from this time the operation should be finished as speedily as it can with safety.

- 19. But if the circulation in the funis be undisturbed there is no occasion for haste, as the child, we are then assured, is in safety.
- 20. The child may be extracted without much difficulty if we act alternately from side to side, by making a lever of its body, and sometimes by pressing it from the ossa pubis with the fingers.
- 21. If the child should stick at the shoulders, the arms must be successively brought down.
- 22. This is to be done by raising the body the opposite way, and by successively bending them at the elbow very slowly, lest they should be broken, and the hand must be cleared toward the *pubes*.
- 23. When both the arms are brought down, the body of the child must be supported upon our left hand placed under the breast, the

in or it is to be drawn out, to sinchen it to
each the cimuco of compression, or to preeat the appropriate of the firm the body of the
falls, or of the placesta from the body of the
interior of the placesta from the body of the

19. But if the circulation in the rings be made in colors.

to a company of the first property of the body and

with the fregies.

fingers on each side of the neck, and the body supported upon our left arm.

- 24. Then placing the right hand over the shoulders, and pressing with our fingers the head towards the sacrum, we must ease the head along, gradually turning the body of the child as it advances toward the abdomen of the mother.
- 25. If the head should not come easily away, we must introduce the fore finger of the left hand into the mouth of the child, by which the position of the head will be rendered more convenient.
- 26. When the head begins to enter the os externum, we must proceed very slowly, and support the perinæum, by spreading the fingers of the left hand over it.
- 27. In some cases there may be a necessity of speedily extracting the child in order to preserve its life, but we must also recollect, that the child is often lost by endeavouring to extract it too hastily.
- 28. When a child has been extracted by the feet, the placenta usually separates very soon

fingers on each side of the neck, and the hody

shoulders, and pressing with our fingers the shoulders, and pressing with our fingers the band fewards the aurrows, we must core the bead fewards transming the bady of the band of the distance the children of the distance of

pilite ample the Bleich band all it let an apple of the production of the production

The same of the last hand the same the or content the same of the

man gray manage glass trees of all

and very easily; but in the management of this we are to be guided by the general rules.

#### SECTION V.

On the Cases which come under the second Distinction.

- 1. We are first to ascertain the presenting part, and if, together with the arm, the head is perceived by a common examination, there may be no occasion to turn the child, such case only constituting the third variety of natural labour.
- 2. But if the case should be such as to require the child to be turned, it might be doubted whether it were proper to dilate the os uteri by art, or to wait for its spontaneous dilatation.
- 3. Perhaps neither of the methods can be constantly followed, but we may generally say, that there is under these circumstances neither danger nor increase of difficulty, from waiting

and very smally; but in the spennyement of

NECTEDR V.

en all Care white come ander the percent like

A series fint to most his presenting and the presenting series that a series that the bead series and the series that the series that the series are the child, englished and early constituting the third their their cities of the

equire the child to be tormed, it inight be equired the child to be tormed, it inight be equired to the child to the equire proper is distincted to the contract of the end of t

S. Parkays believe of the methods can be constituted for the lower properties of the control of

for the spontaneous dilatation, which is therefore in general to be preferred.

- 4. But if more speedy dilatation should be required, whatever is done by art should be done slowly, and in imitation of nature.
- 5. The os uteri is always to be considered as completely dilated when we judge it will allow of the easy introduction of the hand.
- 6. When we have fixed upon the proper time and begin the operation, the os externum must be dilated in the manner before advised.
- 7. The hand must always be introduced into the uterus, on that side of the pelvis where it will pass most conveniently; and there is usually most room at that part which will lead to the feet.
- 8. It is generally most convenient to pass the hand between the body of the child and the ossa pubis, the feet being most commonly found lying toward the belly of the mother.
- 9. In cases which come under this distinction the uterus is seldom contracted very strongly upon the body of the child, but always in some degree.

the billy all la your not sport I bond out

10. But the difficulties which may occur in the operation of turning the child, in these cases, will be fully explained under the following distinction.

#### SECTION VI.

On the Cases which come under the third Distinction.

- 1. The difficulty in the management of these cases depends upon the degree of contraction of the uterus, and upon the distance or awkward position of the feet of the child, but chiefly upon the former circumstance.
- 2. The uterus is in some cases contracted in a globular, and in others in a longitudinal form.
- 3. It is always easier with an equal degree of contraction to turn the child when the uterus is contracted in a globular, than in a longitudinal form.
- 4. When we are called to a case of this kind it is better not to form, or to give a hasty opinion,

betractions some contracted in at anything 2. Lis adways caller with an count degree or contraction to tem the child when the . torol landost qual opinion, nor to attempt to deliver the patient immediately, but to deliberate upon it, and then to make a second examination.

- 5. If the second examination should confirm our first opinion, we may prepare for the operation.
- 6. We shall be able to judge in what part of the uterus the feet of the child lie, if we consider whether it be the right or left hand which presents, which may be known by the direction of the thumb and of the palm of the hand.
- 7. But the contraction of the uterus is the principal difficulty to be surmounted, and the danger in turning the child is in proportion to the difficulty.
- 8. The danger in turning a child when there is a strong contraction of the *uterus*, is a fingle danger, that of rupturing the *uterus*.
- 9. The contraction of the uterus is of two kinds; first, the permanent contraction, in consequence of the waters having been long drained off, which may occur when there has been little or no pain.

- 10. Second, the extraordinary contraction arising from the action of the uterus, returning at intervals, and always attended with pain.
- 11. The hand must be introduced with a degree of force sufficient gradually to overcome the permanent contraction of the uterus, or the operation could never be performed.
- 12. But if we were to attempt to overcome the extraordinary contraction, it must follow, that we can, or cannot overcome it.
- 13. In the first instance we should be in danger of rupturing the *uterus*, and in the second the hand would be cramped, and we should be unable to proceed with the operation.
- 14. The deduction is therefore clear, that we ought not to proceed in our attempts to turn the child while the *uterus* is acting with violence.
- 15. The action of the uterus is rendered more frequent and strong by the generally increased irritability of the patient.
  - 16. Before we attempt to deliver, it will be prudent

blug from the action of the serves retninthe operation which never be principalized. hould be analyte to educated with the operaprudent to endeavour to lessen this irritability, in many cases by bleeding, by clysters, and by an opiate, which, to answer this purpose, should be given in two or three times the usual quantity.

- 17. When the opiate takes effect, and the patient becomes disposed to sleep, we must consider this state as extremely favourable, and proceed without loss of time to the delivery.
- 18. There never can be occasion to separate the arm which presents from the body of the child, and when this has been done, instead of facilitating, it has impeded the operation.
- 19. Without regarding the arm, the right or left hand, as may be most convenient to ourselves, must be introduced in the manner before directed, and conducted slowly into the uterus if there be sufficient room.
- 20. But if the child be jammed at the superior aperture of the *pelvis*, the hand cannot be introduced.
  - 21. We must then fix our forefinger and thumb

product to endouvous to be seen this irrivability, in analy cases the blooding, by circles, and by an equation which, to answer this purpose, thought be given in two or three times the neual

17. When the opiate takes effect, and the patient sheeps we must outsider talk states as extratagely involutible, and proceed without these at time to the de-

to a the same the variety of the contents of t

The legit sair large out anti-copy recount of the large of a share of recommon recommon and continued find the recommon will sair be supposed to be a faithful for the sair and the sair an

Albertalista beresanj aktifika alikurasi .00

kikananiali binanialis pirka mindo beresani ilanga ana

Sala regulared two till code salar ave ale.

thumb in the form of a crutch in the armpit of the child, and pushing the shoulders towards the head, and towards the fundus of the uterus, we must by degrees raise the body of the child till there be room for the introduction of the hand.

- 22. If while we are introducing our hand we perceive the action of the uterus come on, we must not proceed till that ceases or is abated.
- 23. The hand, when introduced, is also to be laid flat during the continuance of the action of the uterus, lest the uterus be injured by its own action on the knuckles.
- 24. When the action ceases or is abated, we must renew our attempts to carry up our hand to the feet of the child.
- 25. In this manner we are to proceed, alternately resting and exerting ourselves, till we can lay hold of one or both feet.
- 26. There is sometimes much difficulty in getting to the feet, and sometimes in extracting them, especially when the uterus is contracted in a longitudinal form.

shoots at the tagen, of a cratch in the athense

- 27. In such cases it is often convenient, when we can reach the knees, to bend them cautiously, and to bring down the legs and feet together.
- 28. But before we begin to extract we should examine the parts we hold, and be assured they are the feet; and we must extract slowly and steadily.
- 29. If we hurry to bring down the feet they may slip from us, and return to the place from which they were brought.
- 30. We must then carry up the hand again, and grasping the foot or feet more firmly, bring them down in the cautious manner before advised.
- 31. When the feet are brought down, if there be difficulty in extracting them, we must endeavour to slide a noose, first formed upon our wrist, over the hand to secure the feet, by which the hazard of their return will be prevented, and the succeeding part of the operation much facilitated.
  - 32. When the noose is fixed over the ancles, we must pull by both ends of it with one hand,

gon metch freedmard,

hand, and grasp the feet with the other, but we must not attempt to proceed hastily.

- 33. When there is afterward much difficulty in extracting the child, it is probably owing to the body of the child being jammed across the superior aperture of the pelvis.
- 34. It will then be proper to pass the finger and thumb as directed at 21, to raise the shoulders and body of the child toward the fundus of the uterus, with one hand, and with the other extract at the same time with the noose.
- 35. When the breech of the child has entered the pelvis, we must proceed with deliberation, but there will be little farther difficulty, except from the smallness of the pelvis, of which we shall speak in the next section.

Commercy paried blints will be glood out of egutions but there will be bride forther diffi-

# SECTION VII.

On those Cases which come under the fourth Distinction.

- 1. The disproportion between the head of the child and the dimensions of the pelvis, may be added to any of the circumstances mentioned under the preceding distinctions.
- 2. But as the management of these has been already directed, there is now occasion to speak only of the peculiar difficulties arising from that cause.
- 3. The degree of difficulty in these cases is greater or less according to the degree of disproportion; but the difficulty of extracting any part of the body of the child is little, compared with that which attends the extraction of the head.
- 4. We will therefore suppose the body of the child to be brought down, but that the head cannot be extracted by any of the methods before recommended.
  - 5. The force with which we endeavour to F 2 extract

of cause month of globaldth 75 worden and . about the examined by any or thousand having extract must then be increased, till it is sufficient to overcome the difficulty or resistance.

- 6. But as the necessity of using great force can only be known by the failure of a less degree to produce the desired effect, we must begin our attempts with moderation, and gradually increase our efforts according to the exigence of the case.
- 7. The force exerted should also be uniform, controuled or commanded, and exerted by intervals, in the manner of the natural pains.
- 8. If the head should not descend with the force which we judge can be safely exerted, we must rest, and give it time to collapse.
- 9. We may then renew our attempts, extracting from side to side, or backwards and forwards, as may best conduce to ease the head through the distorted *pelvis*, alternately resting and endeavouring to extract.
- 10. But if the head should descend in ever so small a degree, the force is not to be increased with the view of finishing the delivery

F 3 expeditiously,

extract must agen be topropied, till it is audit--any has automorphist the committee the de-Divisors Cont. Althoughts on Estimation and s. We may then treatly carr attempts, exexpeditiously, but we must be satisfied with our success, and proceed circumspectly.

- 11. When the head once begins to descend there is seldom much subsequent difficulty in finishing the delivery, as the cause of the difficulty usually exists at one particular part of the pelvis.
- 12. But should the head rest in this situation for several hours, no additional inconvenience would thence arise to the mother, and the longer it rested the greater advantage we should probably gain when we renewed our attempts to extract it.
- 13. It may be presumed when the head of the child has been wedged for a long time in the position we are supposing, and great force has been used to extract it, that there is little reason to expect the child should be born alive; yet instances of this are said to have occurred in practice.
- 14. When we can hook a finger on the lower jaw of the child, the direction of the head may be changed to one more favourable, and the delivery thereby facilitated.

( 12

this believes of room any too related with

out rejected, and proceed circumspacely.

It. When the hard once begins to descend
there is solden males subsequent difficulty in
finishing the distribute, as the court of the difficulty unually asists at one paralicular part of

then for several hours, no administration of the several hours, no administration of the several hours, no administration of the several the mother advantage as the several our schemings are several our several our several our several our

15. It oney be presented when the head of the child has been reciped for a long sime in the position or are supposing, and great force that here here has been used to extract for, that there is little transon to oxipper the child chould be here align; yet instances of this are said to have

14. When we can hash a flagor on the lover jow of the head in the direction of the head near the head near be changed to one wow favourable, and the delivery thereby theiltened.

- 15. But we must not extract with so much force as to incur the hazard of breaking or tearing away the jaw.
- 16. Pressing the head of the child from the ossa pubis to the sacrum, with the fingers or thumbs carried up as high as we can reach, will often be of great use in these cases.
- 17. If the difficulty of extracting the head arises from its enormous size, occasioned by some disease, as the hydrocephalus, &c. these methods steadily pursued will answer our intention, as by a prudent use of the force in our power, the integuments will burst, or even the bones be broken.
- 18. Cases of this kind, in which it might be necessary or expedient to use one or both blades of the *forceps*, or to lessen the head, very seldom occur.
- 19. But if such cases should occur, the latter operation is preferable to the use of the forceps, and the utmost care must be taken that we do no injury to the mother.
- 20. Under these circumstances should it be absolutely necessary to lessen the head of the child,

( 88 .)

to provide as as incurred the bodies of brooking or

16. Previous the best of the child from

many the min op as bigle as on cash reach.

17. If the difficulty of extracting the bend arises from the energones atto, organizated by some dispose, as the hydrographilm, the these methods sendily puriod will anisote during tention, to by a prodest me of the roses in tention, to by a prodest me of the roses in tention, to by a prodest me of the roses in

19. Hat if such cases should occur, the light or opension is preferable to the use of the form cap, and the standards care must be taken that

ed at blands around markin seeds abould in he should at the absolutely successary to idaga viscolately successary to idaga.

child, the perforation may be conveniently made behind either of the ears, or in any part where we can most conveniently fix the point of the *perforator*, and the general rules of the operation must be followed.

- 21. By the force used, should the neck of the child have given way, we are not to separate the body from the head altogether, but we must rest longer and act more moderately.
- 22. But should the body be separated from the head by the force we have used, or should we be called to a case of this kind, there will be no occasion, for this reason alone, to act hastily or rashly, as the head may even then be expelled by the pains.
- 23. But if this should be impossible, or if it be absolutely necessary to extract the head speedily, on account of the state of the mother;
- 24. Then the general rules for lessening the head must be accommodated to the exigencies of this particular case, and the head may be confined to a proper situation by compressing

child, the periodicien may be conveniently made parts where the point absolute we can make conveniently by the point of the perference and the general roles of the

21. By the force mont, should the ment of the seemthe eight inner given when we are not to seemone the hooly from the treat attogether, but
we much rest langer and occurs made materials.

22. This should the hools the should be administrated
the bead by the first mode to administrate and
the bead by the first we had the kind, that, there will
be no caracted to a rate of the kind, that, there will
be no caracted to a rate of the kind, then, there will
be no caracted to a rate of the kind, then the see

the administration of the short of the beautiful of the short of the party of the short of th

Then the general artes for bireening the biree originalist for the exigencing the broad and to the exigencing for the finite passenger case, out the best may be confirmed to a proper situation by rempressing the finite form.

the abdomen with a napkin passed across it, or by the hands of an assistant.

### SECTION VIII.

# Miscellaneous Observations.

- 1. It sometimes happens that no part of the child can be perceived before the membranes break, though the os uteri be fully dilated.
- 2. In such cases we should not be absent when the membranes break, lest it should prove a preternatural presentation, requiring the child to be turned.
- 3. In some cases even when the os uteri is dilated, the membranes broken, and the waters discharged, no part of the child can be felt.
- 4. It will then be prudent, in the cautious manner before directed, to introduce the hand far enough into the *uterus*, to discover the part which does present.

prove a prefermieral presentation, requiring Abornat ed or bilde and ed and hims out to flee on seen through .

5. If the head be found to present we should withdraw our hand, and suffer the labour to proceed in a natural way.

6. If the inferior extremities should present, we may bring down the feet, and then suffer the labour to go on uninterruptedly.

7. But if the shoulder or superior extremities should present, we must proceed to the feet, and turn the child as was before directed.

8. By this conduct we shall guard against the danger of turning a child in a contracted uterus.

9. If we should be called to a case in which the arm presented, and much force had been used to extract the child in that position, the arm having perhaps been mistaken for a leg, and the pains being at the same time violent, it may be impossible, without giving much pain, and incurring some danger of rupturing the uterus, to turn the child, or even to introduce the hand into the uterus, the shoulder of the child being pushed low down into the pelvis.

10. Under such circumstances, it is improper to attempt to introduce the hand into the

( 50 )

5. If the fiend herfound to present we should withdraw dur hand, and suffer the labour to

o. If the inferior outseeming the theuld present,

To the is the choulder or appeared to the second of the chould present, we must spreaded to the second of the chould as inic testore directed.

In By this conduct we shall guard against the day that continue we shall guard against the day this conduct we shall guard against

It we show if he called to a rese in which the store had near the store parential, and makin here had near that the prostance the a folder to their position, the later having periods then then along the later the paid and the single that the paid of the paid that the improvement that the single that the improvement that the single the single that t

to autompt to introduce the bond into the

uterus, or to turn the child, as it will generally be expelled by the efforts of the mother; or it may be extracted by methods less painful and hazardous to the mother.

- 11. Yet in these cases the body of the child does not come doubled, but the breech is the first part delivered, and the head the last, the body turning, as it were, on its own axis.
- 12. Nor is this observation made with regard to a small child coming prematurely, as it will apply to a child of a common size, and when a woman is at her full time, provided the pelvis be well formed.
- 13. This fact, of the possibility of a child being expelled in this position, though originally contradicted with great confidence, is now confirmed in the most satisfactory manner by many cases which have been recorded, in some of which the children have even been born living.
- 14. From these it might be inferred, that a woman in a state of nature, or in perfect health, would not die undelivered, though the arm of

\*Liserror How on sistant

to man in a state of nettern or in perfect health, the second of the sec

the child might present, supposing that she was not assisted by art.

- 15. Yet it is always requisite and proper to turn children when the superior extremities present, if the operation can be performed without the hazard of injuring the mother, and we have a better chance of lessening the sufferings of the mother, and of preserving the child.
- 16. But when there is no chance of preserving the child, and yet it cannot be turned without the greatest danger to the mother, knowing the possibility of its being expelled in this position, it is necessary to consider the propriety of the operation before we perform it.
- 17. It remains, however, to be proved by future experience, how far, and in what cases, the preceding observation ought to be a guide in practice.
- 18. In cases of presentation of the superior extremities, in which the difficulty of turning the child would be very dangerous, and great

he shift might present, supposing blast shi

1.5. Yet it is always requirits and proper to teen children when the suppositor extremition properties the the operation can be performed without the housening the injuring the mother, and we have a house above of lessoning the singerings of the implies, and of-preserving the

16. But when then it was a chance of preserving the shild, and yet it enamed to turned
without the president slanger to the mother,
knowing the passibility of its being expelled
in this position, is is necessary to consider the
propriety; at the operation before we per-

17. It remains, however, to be proved by funce experience; bow far, and in whit cases, the proceding observation ought to be a guide

18. In cases of presentation of the superior extremely, in which the difficulty of turning the cold mount be very dangerous, and greater or

or insurmountable, another method has been recommended.

- 19. But of this method, which has been practised by one gentleman to whose know-ledge and experience I pay great respect, I am not a competent judge, having never tried it.
- 20. I therefore refer to the annexed note for an explanation and history of the method to which I allude.

Note. Hoorneus, sæpe laudatus, adhuc peculiarem, novum eumque breviorem modum, fætum mortuum cum brachio arctissimè in vagina uteri hærente extrahendi, invenit atque descripsit, qui in eo consistit, ut quando ad pedes pervenire nequit, collum, utpote quod in fætibus valdè adhuc tenerum est, vel scalpello a reliquo trunco resecet, vel unco idoneo quam cautissimè auferat. Hoc enim facto, vel sponte mox prorumpit ex utero fætus, vel tamen, dum brachium propendens attrahitur, quod medico loco habenæ inservit, quam facillimè excutitur. Caput vero deinde seorsim mox vel manu, vel aliis

-month compared parameters. Our del parameter du convenier venunt college, algore cue, et mente d'acconstructe per les altre obtants applier s toto nghena imervit igana facilityo anequitur. live smeath list from mismals obmon our targe? aliis propositis artificiis, si manus parum esset, ejiciendum.

HEISTER. cap. cliii. sect. ix.

The latter part of this description is further explained in the next section.

### SECTION IX.

I am induced to reprint the following, as they were the very cases which first gave me an opportunity of observing the spontaneous evolution.

### CASE I.

In the year 1772, I was called to a poor woman in Oxford Street, who had been in labour all the preceding night, under the care of a midwife. Mr. Kingston, now living in Charlotte Street, and Mr. Goodwin, surgeon, at Wirksworth, in Derbyshire, who were at that

er ni mont had a present being of minera of a policyle. . 3ff King-on, now living in

Songram administration of the constants of the file

that time students in midwifery, had been sent for some hours before I was called. The arm of the child presenting, they attempted to turn and extract it by the feet, but the pains were so strong as to prevent the introduction of the hand into the uterus. I found the arm much swelled, and pushed through the external parts in such a manner that the shoulder nearly reached the perinaum. The woman struggled vehemently with her pains, and, during their continuance, I perceived the shoulder of the child to descend. Concluding that the child was small, and would pass, doubled, through the pelvis, I desired one of the gentlemen to sit down to receive it, but the friends of the woman would not permit me to move. I remained by the bed-side till the child was expelled, and I was very much surprised to find, that the breech and inferior extremities were expelled before the head, as if the case had originally been a presentation of the inferior extremities.

The child was dead, but the mother reco-G vered of turn and o ... I by the less, but the country and tomorrow the street of the street enter pending of the hand that the meres. I found

vered as soon, and as well, as she could have done after the most natural labour.

#### CASE II.

In the year 1773, I was called to a woman in Castle Street, Oxford Market, who was attended by a midwife. Many hours after, it was discovered that the arm of the child presented. Mr. Burosse, surgeon, in Poland Street, was sent for, and I was called into consultation. When I examined, I found the shoulder of the child pressed into the superior aperture of the pelvis. The pains were strong, and returned at short intervals. Having agreed upon the necessity of turning the child, and extracting it by the feet, I sat down and made repeated attempts to raise the shoulder, with all the force which I thought could be safely used; but the action of the uterus was so powerful that I was obliged to desist. I then called to mind the circumstances of the case before related, mentioned them to Mr. Burosse,

G 2

and

and the could be settle as the could be been and the could be coul

ATT BEZO

seried. Alcolings, suggest, fallologi bland, spalments of the Louis and Lane and Lane and Lane and the priest . The police were strong out? - Make self white to be the the block of both of the with all the force which I thought qualit how make shore related, the control december str. Bureses, and proposed that we should wait for the effect, which a continuance of the pains might produce, or till they were abated, when the child might be turned with less difficulty. No further attempts were made to turn the child. Then every pain propelled it lower into the pelvis, and in a little more than one hour the child was born, the breech being expelled, as in the first case.

This child was also dead, but the mother recovered in the most favourable manner.

Having been prepared for observing the progress of this labour, I understood it more clearly, and attempted to explain both in my lecture on the subject, and in the aphorisms which were printed for the use of the students, my opinion of the manner in which the body of the child turned, as it were, upon its own axis. I also pointed out the circumstances, in which, I supposed, the knowledge of the fact might be rendered useful in practice; but with great circumspection.

soud proposed that we should wait for the soffeet, which a configurates of the pains might crosses, which a configurates were abstra, when the child solds he turned with ign difficulty. No latility attached with ign difficulty. No latility attached is breed into the latin and in a hittle more than one hour the child was here. Into the child, and in a hittle more than one hour the child was here. Into the child, and in a hittle more than one hour the child was here. Into the child, and in a hittle more than one hour the child was here. Into the child, and here here the heing copelities, and a single capalities, and the the single copelities, and the the single capalities and the child was here and the single capalities, and the the single capalities and the child was here and the single capalities, and the single capalities and the child was here and the single capalities and the child was here and the single capalities and the child was here and the single capalities and the child was here.

response our that come calls over aller and

Finning over prepared for alterring the finning over prepared for alterring the finning over this labour, I understood it more drawly, and luterapped to explaid outh in any casely, and luterapped to explaid outh in any casely, and in the total apportune of the subject, and in the the apportune of the subject of the subject of the subject out its make, the which the body of the child function of the which the core matters in unice, upon its own its ow

#### CASE III.

January the 2d, 1774, I was called to Mrs. Davis, who keeps a toy-shop, in Crown Court, Windmill Street. She had been a long time in labour, and the arm of the child presented.

The late Mr. Eustace had been called on the preceding evening, and had made attempts to turn the child, which he had continued for several hours without success. I was sent for about one o'clock in the morning, and on examination found the arm pushed through the external parts, the shoulder pressing firmly upon the perinæum. The exertions of the mother were wonderfully strong. I sat down while she had two pains; by the latter of which, the child was doubled, and the breech expelled. I extracted the shoulders and head, and left the child in the bed. Mr. Eustace expressed great astonishment at the sudden change; but I assured him that I could claim no other merit on account of this delivery, except that I had not impeded an effect which was wholly produced by the pains.

## CARR LIL.

to bein the chart which are one continued for and the same of th extensive parts, the should? proming freally ther were wenderfully strains. I sat down the child was nougled, and the breech expelled. the ban Jugal Dan trableuds out belowskin I trees accompliment at the sudden change; not produced by the pains.

This child was also dead, but the mother recovered in the most favourable manner.

In all these cases, the women were at the full period of utero-gestation, and the children were of the usual size.

Many other cases of the same kind have occurred to me; and with the histories of several, varying in the time or manner in which the evolution of the child was made, I have lately been favoured by gentlemen of eminence in the profession, and many others have been published in different countries. But these are sufficient to prove the fact, that in cases in which children present with the arm, women would not necessarily die undelivered, though they were not assisted by art.

With respect to the benefit we can, in practice, derive from the knowledge of this fact, I may be permitted to repeat, that the custom of turning and delivering by the feet in presentations of the arm, will remain necessary and proper, in all cases in which the operation can be performed with safety to the mother, or give a chance of preserving the life of the child.

avanuari oldanio rat team dit en process ti serioù af the sorgenistion, noch the children Canoril Servetti ar ita calling a change of party and a string to child. But when the child is dead, and when we have no other view but merely to extract the child, to remove the danger thence arising to the mother, it is of great importance to know the child may be turned spontaneously, by the action of the uterus. If we avail ourselves of that knowledge, the pain and danger which sometimes attend the operation of turning a child may be avoided. Nor would any person, fixing upon a case of preternatural presentation, in which he might expect the child to be turned spontaneously, be involved in difficulty, if, from a defect of the pains, or any other cause, he should be disappointed in his expectations. Nor would the suffering, or chance of danger to the patient, be increased by such proceeding, as the usual methods of extracting the child could, under any such circumstances, be safely and successfully practised.

be safely and successfully practised.

# CLASS IV. Anomalous or Complex Labours.

# FOUR ORDERS.

ORDER I.

Labours attended with Hemorrhage.

ORDER II.

Labours attended with Convulsions.

ORDER III.

Labours with two or more Children.

ORDER IV.

Labours in which the Funis Umbilicalis presents before the Child.

CAMPS BY: Assessment of Congress

BELLEVILLE STREET

Sa andao

Spring satisfact to the Planta with the

HI REILES

Labour Manded with Constitutes

ORDER III.

Labour with new or more Children.

TI SHIRE

Labour in takich the Funds Umbilicalls presents

400re.the Child.

On Labours attended with Hemorrhage.

HEMORRHAGE. A discharge of blood from the uterus, inordinate with respect to time or quantity.

## VARIETIES.

- 1. In abortions.
- 2. At the full period of utero-gestation.
- 3. After the birth of the child.
- 4. After the expulsion of the placenta.

Note. No general description or character can be given to Anomalous Labours as a class, because the different orders bear no resemblance to each other. They are brought together merely to prevent the multiplication of classes.

On Judowe wanded with Memoriane,

the second in the second to a second to the second transfer transfer to the second transfer tran

# Variation

The first feet following the state of charges of charges of the shift.

The first one birth or the shift.

North No general description or character can be gived to Anomalous Lalaces as a class, because the discrept arders been no resemblative to each other. They are breacht begriffer marrily to prevent the multiplecation

# ON ABORTIONS.

### SECTION I.

- 1. With respect to the time of pregnancy, all expulsions of the fatus may be reduced under two distinctions.
- 2. In the first will be included all those which occur before the uterus is sufficiently distended to allow of any manual operation; and these may be properly called abortions.
- 3. In the second may be classed all those which allow of manual assistance, if required; and which are therefore to be esteemed as labours, premature or at the full time.
- 4. But no precise period of pregnancy can be fixed as a line for these distinctions.
- 5. We may, however, in general say, that all expulsions of the fætus, before the end of the sixth month, are to be considered as abortions.

# ON ABORTIONS.

## SECTION 1.

1. With respect to the time of pregnancy, all expulsions of the faces may be reduced under two distinctions.

2. In the first will be included all those which occur before the stores is sufficiently distended to allow of any manual operation; and these may be properly called abortions.

3. In the second may be closed all these which allow of manual assistance, if required; and which are therefore to be externed as labours, premature as the fall time.

4. But no precise period of pregnancy can

so fixed as a little for these distinctions.

for We make however, in general say, that
all expelsions of the firms hafore the and an
the vixth month here to be considered as abortions, at their

- 6. But all expulsions of the fatus, after the expiration of the sixth month, are to be esteemed as labours, and, if attended with the same circumstances, should be managed upon the same principles.
- 7. Yet expulsions of the fatus sometimes happen so critically, as to make it doubtful to which distinction they should be ascribed, especially in cases in which there are two or more children.
- 8. When manual assistance is thought needful, the longer the time wanting to complete the full period of pregnancy, the more difficult must be any operation.

## SECTION II.

# On the Causes of Abortions.

- 1. The predisposing causes of abortion are, 1st, general indisposition of the constitution; 2d, infirmity of the uterus.
  - 2. The general state of women who are disposed

6. But all expulsions of the flow, after the expiration of the sixth month, are to be esteemed as labours, and, if attended with the same circumstances, should be managed upon the same principles.

To Yet expulsions of the distance sometimes happen as cratically, as to make it doubtful to which distinction they should be ascribed, superially in cases in which there are two or more children.

20. When rememble against over is thought needful, the longer the time wanting to complete the full pariod of pregnancy, the more difficult tours on any operation.

in Rolling in

the shall name of Alloratories

1. Piet pridisposing timera of abortion are, 24, agrees indisposition of the constitutions, 24, 24, agrees of the state of the constant are 22.

disposed to abortion is very different, some being weak and reduced, and others plethoric.

- 3. Weakly women become more liable to abortion, because they are susceptible of violent impressions from slight external causes.
- 4. Plethoric women are more liable to abortion, from the peculiar disposition which the vessels of the *uterus* have, from structure and habit, to discharge their contents.
- 5. Every action in common life has been assigned as a cause of abortion.
- 6. But it is to the excess of these actions that we are to attribute their effects, for women in health seldom abort, unless from violent external causes.

### SECTION III.

# On the Prevention of Abortion.

1. As every disease to which women are liable may dispose to abortion, the method instituted to prevent it must be accommodated

( 77 )

dispond to abstrict is very different, some being week and reduced, and editors pleathering.

3. Weakly wranin become more liable to viaabstrict, because they are anscaptible of vialett impressors rises they are anscaptible of viadett impressors rises their extensal causes.

4. Postporic wedgen are more liable to abore
their facts the peculiar disposition which the

a. Every serion in common life has been

the limiter is to attribute their edects, for sothat we has to attribute their edects, for somen in ligable seldons above, valous from via-

ALL WOLFSON

On the Presenting of Aportlan.

I. As every circuss to which women are liable may dispose to abortion, the method in-

to the disease, or to the state of the constitu-

- 2. In some constitutions, abortions may be prevented by repeated bleeding in small quantities, by antiphlogistic medicines, and sometimes by warm bathing.
- 3. In others, abortion may be prevented by nourishing and invigorating diet and medicines, by bark, by the liberal use of wine, especially claret, and often by cold bathing.
- 4. But it will be proper, in every case, to avoid all violent exercise, to keep the mind composed, and to rest frequently in an horizontal position.
- 5. Women seldom abort while they have the vomiting which usually attends early pregnancy.
- 6. In women who have no spontaneous vomiting, this may be excited, with safety and advantage, by frequently giving small doses of *Ipecacuanha*.
- 7. Pregnant women are usually costive, and abortions have been very often occasioned by too great assiduity to remove this costiveness, which

ed your constitutions, abortique may be -many lines at passend halvager yet between ares, by anapaleglade facticines, and some-

to seech Hams priving givening doesn't desert of

which is a natural and proper state, in the early part of pregnancy.

### SECTION IV.

## On the Signs of Abortion.

- 1. The signs of abortion are, frequent micturition, tenesmus, pains in the back, abdomen, and groins, with a sense of weight in the region of the uterus.
- 2. But the most certain sign is, a discharge of blood, which proves that some part of the ovum is separated from the uterus.
- 3. It has been supposed when this last sign appears, that there is scarcely a possibility of the patient proceeding in her pregnancy.
- 4. But I have met with an infinite number of cases in practice, in which, notwithstanding this appearance, once or oftener, to a considerable degree, the discharge has ceased, and no ill consequences have followed.
- 5. We are therefore to persevere in the use of those means of prevention which are thought

H

or production of the product of the state of the mun ataltur on drew redressant tola . reasonable and proper, till the abortion has actually happened.

6. It is not always prudent to give a decided opinion of the probable event of those cases in which abortion is threatened, as their termination is often different from what might have been expected from the symptoms.

### SECTION V.

On the Treatment of Women at the Time of Abortion.

- 1. The treatment must vary according to the nature and degree of the symptoms.
- 2. There is an endless variety in the manner in which abortion takes place. Some women abort with sharp and long continued pains, others with little or no pain; some with a profuse and alarming hemorrhage, others with very little discharge. In some the ovum has been soon and perfectly expelled, in others after a long time, in small portions, or very

H 2

the treatment that wary document and and become all arms al. squadash all the much decayed; but the only alarming symptom is the hemorrhage.

- 3. The hemorrhage in abortions is not always in proportion to the period of pregnancy, this being in some advanced cases very small; and in others, though very early, abundant.
- 4. The hemorrhage usually depends upon the difficulty with which the *ovum* may be expelled, and upon the state of the constitution of the patient naturally prone to hemorrhage.
- 5. The general principles which should guide us in the treatment of hemorrhages, from any other part of the body, are applicable to those of the uterus, regard being had to the structure of the uterus.
- 6. If the patient be plethoric, some blood should be taken from the arm at the commencement of the hemorrhage, and the saline draughts with nitre, or acids of any kind, may be given in as large a quantity, and as often, as the stomach will bear.
  - 7. These may also be given during its con-H 3 tinuance

mark decayed r bardin only marming symptom.

S. The homoralings in abartions is not garged to the period of party. State bunglin some advanced cases very early, and in cinery through very early,

to The namericage washly depends upon the construction of the cons

If it is the treatment of hemorrhages, from may a in the treatment of hemorrhages, from may after part of the body, are applicable to the struct of the struct being had to the struct

of the passent by plethoric, some blood about the comsence sheet of the hemoritage, and the saling design with nitre, or acids of any kind, may given in as large a quantity, and as often, as

There, may also be given during its con-

tinuance, and cloths wet with cold vinegar may be applied to the abdomen and loins, and renewed as they become warm. The patient should be exposed to, and suffered to breathe, the cold air.

- 8. Every application or medicine, actually or potentially cold, may be used. A large draught of cold water or ice may be given with great propriety; and it is the custom in Italy to sprinkle ice over the body of the patient if the danger of the case be imminent.
- 9. Every medicine or application which has the power of slackening the circulation of the blood, eventually becomes an astringent; but astringents, properly so called, can have no power in stopping hemorrhages from the uterus.
- 10. Hemorrhages are stayed by the contraction of the coats of the blood vessels, or by the formation of *coagula*, plugging up the orifices of the open blood vessels.
- 11. Both these effects are produced more favourably during a state of faintness, which, though occasioned by the loss of blood, becomes a remedy in stopping hemorrhages.

12. Cordials

- 12. Cordials are not therefore to be hastily given to those who are faint from loss of blood; unless the faintness should continue so long as to make us apprehensive for the immediate safety of the patient.
- 13. The introduction of lint, a piece of sponge, or any other soft substance, into the vagina, has been recommended, and sometimes used with advantage, by favouring the formation of coagula.
- 14. Cold or astringent injections into the vagina, or even a piece of ice introduced into the vagina, have also been recommended.
- 15. Opiates have been advised in abortions attended with profuse discharges, and they may sometimes be proper to ease pain, or to quiet the patient, especially when there is a chance of preventing the abortion, or after the accident has happened.
- 16. But when there is no hope of preventing the abortion, the degree of pain proving the degree of action of the *uterus*, and the action of the *uterus* producing and favouring the contractile power of the blood vessels, if by opiates

121 Condide dee not therefore to be hastily from to those who are thin from loss of blood; and the from the blood openions should continue so long as to teste in apprehensive for the immediate

13. The introduction of Bot, a piece of gentle, or may other soft substance, into the marries, but been teconomended, and sometimes

Sulmura to aids

all bini molitochi injectiona inte the

The Opinion have been advised in absortions aftended with profuse discharges, and they have sometimes for proper to case pain, or to their the plaient, repectable when there is a transfer of preparating the abortion, or after the transfer of preparating the abortion, or after the

ignificant has happened.

16. Dut when there is no hope of preventing the chartes, the degree of pain proving the degree of action of the narrow, and the action of the narrow, and the action of the structure predicting and in conting the continuent predicting and in continuents of the blood restrict, if her opinion the blood restrict, if her opinion the

the action of the uterus should be prevented or checked, they may contribute to the continuance of the hemorrhage.

- 17. Hemorrhages in abortions, independent of other complaints, though sometimes very alarming, are not dangerous.
- 18. But if women abort in consequence of acute diseases, or if they are attended with violent spasms, there will be real and great danger.
- 19. For they abort because they are already in great danger, and the danger is increased and accelerated by the abortion.
- 20. The *ovum* has been sometimes retained in the *uterus* for many months after the symptoms of abortion had appeared, and when it had lost the principle of increasing.
- 21. But it is not now thought necessary or proper in abortions, to use any means for bringing away the *ovum*, or any portion of it which may be retained, with instruments or manual assistance.

To binespent to Stabile termin wit to native and decised, the state is and indicate.

indepolyant and in abortions, independent

to surseparate to the property of the property

beninter equilibring pool and surrand 122 copyrige and realth editional young not upon add to
the spring has been appropriate the spring to another

the partition of the partition of the same of the same

MOTOTA II

#### SECTION VI.

On Hemorrhages at the full Period of Uterogestation.

- 1. Under this section will be included all those hemorrhages which may happen in the three last months of pregnancy.
- 2. These are occasioned first by the attachment of the placenta over the os uteri; secondly, by the separation of a part, or of the whole placenta, which had been attached to some other part of the uterus.
- 3. Hemorrhages arising from the first cause are more dangerous than from the second; but those from the second have sometimes proved fatal.
- 4. The danger of attending hemorrhages is to be estimated from a consideration of the general state of the patient, of their cause, of the quantity of blood discharged, and of the effect of the loss of blood, which will vary in different constitutions.

5. Hemorrhages

### AND ROPES

the Hemorrhages at the full Period of Chera-

1. Under this section will he included all these home things which may pappen in the face last months of premiency.

2. These are occisioned first by the attachment of the planeau over the m steet; secondly, by the separation of a part, or of the while glacesta, which had been attached to some other part of the merus.

3. Hopsordiages arising from the first cause are more dangerous than from the second; but those from the second have sometimes proved fetal.

de. The danger of nitending heatershapes is to be estimated from a consideration of the general state of the patient, of their cause, of the quantity of blood discharged, and of the chart of the blood discharged, and of the chart of the blood, which will-wary in

6. Hemorringes

- 5. Hemorrhages are infinitely more dangerous with sudden than with slow discharges of blood, even though the quantity lost may be equal.
- 6. The danger arising from hemorrhages is indicated by the weakness or quickness of the pulse, or by its becoming imperceptible, by the paleness of the lips, and a ghastly countenance, by inquietude, by continued fainting, by a high and laborious respiration, and by convulsions.
- 7. The two last symptoms are usually mortal, though when women are extremely reduced, they are liable to hysteric affections of a similar kind, that are not dangerous.
- 8. The vomiting, which generally follows violent hemorrhages, indicates the injury which the constitution has sustained by the loss of blood, but by the action of vomiting the patient is always relieved, and it contributes to the suppression of hemorrhages.
- 9. Near the full period of utero-gestation, women are always in greater danger in those hemorrhages

of blood, even though the quantity less may be seemed.

of The danger arising from hemorrhages is is dicated by the versioness or quickness of the pulse, or by its incoming imperceptible, by the palences of the lips, and a glandly counternance by inquictude, by continued fainting, by a high and laborious respiration, and by converse.

7. The two last symptoms are usually mortal, though when women are extremely reduced, they are inclie to bysteric affections of

8. The vomiting, which generally follows violent bemorrhages, indicates the injury which the constitution has sustained by the loss of blood, but by the action of vomiting the partient is always religious, and it contributes to the suppression of hemorrhages.

9. Pearthe full period of utero-gestation, seemen are always in greater clarger in those homorrhanes

hemorrhages which are not accompanied with pain.

10. For the pain proving the contraction of the uterus, and this proving that the strength of the constitution is not exhausted, the danger in hemorrhages may often be estimated by the absence or degree of pain.

### SECTION VII.

On those Hemorrhages which are occasioned by the Attachment of the Placenta over the Os Uteri.

- 1. Though the placenta, which may easily be distinguished from the membranes, or from coagulated blood, as soon as the os uteri is a little opened, be attached over the os uteri, the woman usually passes through the early part of pregnancy without any inconvenience, or symptom which denotes the circumstance.
- 2. But before or when the changes previous to labour come on, there must be an hemorrhage, because a separation of a part of

drive braining or and agreempanied with

The recent, and this proving that the arreagely
the recent, and this proving that the arreagely
at the resolitation is the arbumined, the dealget in bearinging may allow be referrated by
the character or degrees of pain.

## HAY MOLYGER

the displaced of the Pharmin car the Os.

distinguished from the resemblance, or from congulated blood, or and the or ward is desirable opened, be attached over the or ward is desirable opened, be attached over the or ward is desirable opened, be attached over the carrier and the carry part of the carry part.

2. Due la chenges previous de la la la company de la la company de la la company de la

the placenta is thereby occasioned, and as the disposition to labour advanceth, the hemor-rhage is generally, though not universally, increased.

- 3. With this circumstance very slight external causes are also apt to occasion hemorrhage.
- 4. When an hemorrhage from this cause has once come on, the patient is never free from danger till she is delivered.
- 5. The powers of the constitution are undermined by hemorrhages profuse or often returning, so that no efforts, or only very feeble and insufficient ones, are commonly made for the expulsion of the child.
- 6. We are therefore often obliged to free the patient from the imminent danger she is in by artificial delivery.
- 7. Of the propriety of this delivery, in cases of dangerous hemorrhage, there is no doubt, or can be any dispute, except as to the precise time when the patient ought to be delivered.
  - 8. On the first appearance of the hemor-rhage,

the placests is thereby necessioned, and an the imposition to Iniough advancetts, the hemor-these is generally, though not universally, in-

S. With this circumstance very slight ex-

de Nineman bennouchage from this cause has once come com rio patrent is never free frem danner till also a delivered.

derwised by heaptrobagus profuse or often rederwised by heaptrobagus profuse or often returning, so that no offers, or only very fields
and installicient ones, are continuedy made for
the exculsion of the while

6. We are therefore when obliged to free

2. Of the propriety of this delivery, in cases of dangerous hemorrhage, there is no doubt, or can be any dispute, reacept as to the precise time when the patient aught to be delivered.

8. On the first appearance of the hemor-

rhage, unless it be prodigious in quantity, or unusually terrifying in its effect, it is seldom either requisite or proper to attempt to deliver by art.

- 9. Nor does it often happen that a second or a third return of the discharge compel us to the delivery by art.
- 10. But as a patient with this circumstance cannot be secured till she is delivered, and as the delivery is seldom completed by the natural efforts, and as the artificial delivery, though performed before it is absolutely necessary, is not dangerous, if performed with care, we must be on our guard not to delay the delivery too long.
- 11. In some cases in which it might be thought eligible to deliver on account of the hemorrhage, the parts are so unyielding as not to allow of the operation itself without some hazard.
- 12. Yet when the parts requiring dilatation make no resistance to the passage of the hand, the event of the operation is always more precarious,

them, makes it to produce in quantity, or understanded to it is seldent to modely territying sucide execut, it is seldent extent to require the contract to altern the deliver

or a third return of the discharge compet us

council to fecured will the circumstance council of delivered, and as the delivery is added coinciled by the nateral effects, and as the artificial delivery, though performed before it is absolutely necessary, is not dangered; if performed with care, we not congress on our goard not to delay the delivery too long.

It. In some cases in which it might be thought eligible to deliver out account of the hemorrhage, the parts are so unyielding as not to allow of the operation itself without some harmed

12. Yet when the pasts requiring dilatation sanks and resistance to the passage of the hand, the exent of the operation is always more pre-

carious, the operation having been deferred too long.

- 13. But though it may be proper in some cases to determine on immediate delivery, the operation must always be performed with the utmost deliberation.
- 14. The first part of the operation has been described under preternatural presentations.
- 15. When the hand is carried to the placenta attached over the os uteri, it is of little consequence whether we perforate the placenta with our fingers, or separate it on one side till we come to the edge, though the latter is generally preferable.
- 16. If the hand be passed through the placenta, we shall come directly to the part of the child which presents.
- 17. But if we separate the placenta to the edge, the hand will be on the outside of the membranes, which must be ruptured before we lay hold of the feet of the child.
  - 18. No regard is to be paid to the part of the

( 48 ) was To obtain out on ad him bank bulk, who ting out of blay of or of bring all ay I - I the child which may present, as it must always be delivered by the feet.

- 19. The feet of the child being brought slowly into the pelvis, we must wait till the uterus is contracted to the body of the child, which will be indicated by pain, and known by the application of our hand to the abdomen.
- 20. The delivery must then be finished very slowly, to give the uterus time to contract as the child is withdrawn from its cavity; but this part of the operation has likewise been described under preternatural presentations.
- 21. An assistant should make a moderate pressure upon the *abdomen* during the operation, to aid the contraction of the *uterus*, and to prevent ill consequences from the sudden emptying of the *abdomen*.
- \*22. When the child is born, the hemorrhage will be generally stayed, if the operation has been performed slowly.
  - 23. But if the hemorrhage should continue

the child which may present, as it must al-

19. The feet of the child being brought the lawly into the privite we must wait till the lawly in the child the body of the child thick will be indicated, by policy and known with the application of our hand to the se-

feed and apply the give the where the bear to conthe child is wishdrawn from his reand as the child is wishdrawn from his like reand the child is wishdrawn from his likeand the continuous properation has likeand the continuous properation from his like-

The state of the contraction of the state of the opening the state of the opening the state of the contraction of the state of the stat

ege will be generally stayed, if the operation been performed slewly.

if the hemorrhage should continue

or return, the *placenta* is to be managed as will be afterwards directed.

- 24. Should no uncommon difficulty attend the delivery, children will be often born living in cases of hemorrhage which are attended with the utmost danger to the mother; or, as it has sometimes happened, after the death of the mother.
- 25. Before, during, or after delivery in cases of hemorrhage, the means and applications before recommended, may be occasionally used with advantage.

### SECTION VII.

On those Hemorrhages which are occasioned by the Separation of a Part, or of the whole Placenta, before or in the Time of Labour.

- 1. Hemorrhages arising from this cause are seldom so alarming or dangerous as the preceding.
  - 2. But if the separation of the placenta be I 2 sudden

Her as finishment of an in meaning ods , angles of

the delivery, chieffing will be offen berit living the delivery, chieffing will be offen berit living the case of the chieff of the case o

the more against the mount and applications in the same

Lagalancia de V

## THEY WOLLDON

the close Hericardages which are established by

the properties and paints and the passent of the pa

all shorted and the separation of the printents the

sudden and extensive, the danger may be equal, and the same mode of proceeding required.

- 3. Our conduct must be guided by a consideration of the degree and effect of the hemorrhage, and of the state of the labour when it occurs.
- 4. Should the hemorrhage from this cause occur in the first period of labour, the action of the *uterus* will be weakened, but it may be sufficient to dilate the *os uteri*.
- 5. If the quantity of blood lost in these cases be very considerable when the os uteri is sufficiently dilated, the greater the degree the better, if the case will allow us to wait so long, the membranes containing the waters may be ruptured.
- 6. By the discharge of the waters the distention of the *uterus* will be lessened, and by the consequent contraction, the size of the vessels being diminished, the hemorrhage will of course be abated or removed.
- 7. After the abatement or suppression of the hemorrhage, the action of the uterus will

equal, and the same pieds of proceeding re-

dutted.

25. Our conduct must be guided by a conelisamica of the e-gree and affect of the hemorrange, and of the class of the labour when

de Chicult the bemorringe from this camer occur in the first period of tabour, the metion of the exemptal be weakened, and it may be

of the seas considerable when the in-contract the state of the season of the season of the state of the season of

for the decirate will be tenues, and by denter of the sines of the interest will be tenues, and by denter of the sparsparint time size of the stands being distintished, the bencerings, will

To delicate the setting of the series of the

become stronger, so that the delivery will, in general, be then completed without further assistance.

- 8. But if the hemorrhage should continue after the discharge of the waters in such a degree as to threaten danger; or if it should commence in the second period of the labour, the interposition on our part must vary according to the circumstances, and chiefly according to the situation of the child.
- 9. It may in some cases be necessary to deliver by art as in the preceding section, and in others to deliver with the *forceps* or *vectis*, if the hemorrhage be profuse, and we despair of the child being expelled by the natural efforts.
- 10. The proper management of all such cases may be collected from what will be generally said on the subject, being always on our guard to distinguish between fear and real danger.

b come stronger, so that the delivery will, in general, he then completed without further

S. Her if the hemorrhage should continue, after the discharge of the waters in apole a degree on to the catenders or if it should commence in the second period of the labour, the interposition on our part must vary or grading to the circumstances, and charge according to the circumstances, and charge ac-

9. It may in some cases be mounterly to defiver by art as in the proceeding section, and it others to-deliver with the /earge or users, if the hemographs prefuse, and we despois the the child herog expelied sty the natural

the proper management of all such cores may be collected from what will be generally said on the subject, today stones on our guard to distinguish between har and real danger, see the grant of the subject.

machana

高·

#### SECTION VIII.

- On those Hemorrhages which occur when the Placenta is retained after the Birth of the Child.
- 1. The placenta is generally expelled by the spontaneous action of the uterus in a short time after the birth of the child.
- 2. But sometimes the placenta is retained, 1st, from the inaction or insufficient action of the uterus; 2d, by the irregular action of the uterus; 3d, by the scirrhous adhesion of the placenta to the uterus.
- 3. Sometimes there is a profuse discharge of blood, when no action is exerted by the uterus to expel the placenta, and this is found in practice to be far the most common cause of hemorrhage at the time of delivery.
- 4. Whenever there is a hemorrhage, the whole or a portion of the placenta must have been previously separated, and the hemorrhage usually continues, or returns till the placenta is expelled or extracted out of the cavity of the uterus.

#### SECTION IX.

On the Retention of the Placenta from the Inaction or insufficient Action of the Uterus.

- 1. Though the placenta be retained after the birth of the child, if there be no hemorrhage, we are to wait, without any interposition on our part, in expectation of the action of the uterus.
- 2. The time which it may be proper and expedient to wait, will depend upon the state of the patient, and the state of the patient generally depends upon the previous circumstances of the labour; so that it may not be proper to wait in one case for any length of time, and in another we may safely wait four, six, or even twelve hours.
- 3. But no patient ought to be left before the *placenta* is brought away, because though there may be no existing hemorrhage, a dangerous one may at any time come on.
- 4. When the patient complains of pain, the expulsion of the *placenta* may be safely forwarded, by aiding the contraction of the *uterus*

minuted the most at aspects with produced the soamong the six by measter in the state of a man and a dynast to
any tests be at any of any of the state of a decay
are to be at any of the state of any of the state of a decay
are to the state of any of the state of a decay
are made which if the state of the state of any of the state of the st

Barrangen and the state of the

the state of his physical desiration, on the last of the state of the

by moderate pressure with the hand upon the abdomen, and by pulling very gently by the funis.

- 5. But if the first pains, with the aid we think it prudent to give, should not bring down the placenta, we are to wait for a return of the pains, proceeding in the same cautious manner.
- 6. When that part of the placenta into which the funis is inserted can be felt, little danger or difficulty is to be apprehended; but we are either to wait longer, or to extract it very slowly.
- 7. But if a hemorrhage were to come on, the placenta being retained, it would be equally necessary to extract the placenta as it would be to extract the child, provided the degree of hemorrhage was equally profuse or sudden.
- 8. After the birth of the child, the extraction of the *placenta* is therefore to be considered as the only method by which an apprehended or present dangerous hemorrhage is to be prevented or avoided.
  - 9. Yet all discharges of blood do not require

of and and by pulling way goody by the

district of prodess to give, should not bring almost think of prodess to give, should not bring almost in district a which of the placesta, we are to wall for a which of the paint, exerciseding in the same described

which the jags is muched the be fell, firther that the first hinted the supposite of the first is to be supposited their in the supposite of t

The struct the child, provided the character of the countries of the provided the child, provided the character of the character of the child, provided the character of the child, provided the character of the

O. After the hirth of the child, the extraction of the placema is therefore to be countdened as the only method by which an appreson had or present dangerous hereovelops in to

con ran oh hould to especialist 84 to 1. Q

quire a speedy extraction of the *placenta*, but such only as by their violence or continuance, or frequent returns, threaten danger.

- 10. If much force be used in pulling by the funis, there will be danger; 1st, of tearing it from the placenta; 2d, of inverting the uterus; 3d, of injuring the uterus by the violence; 4th, of increasing the hemorrhage.
- 11. The danger of these consequences is greater when force is used to extract the placenta by the funis, than by the prudent introduction of the hand into the uterus for that purpose.
- 12. In cases in which the uterus acts insufficiently, by attending to the respiration you will sometimes be able to bring down the placenta, just using so much force, in pulling by the funis, as will prevent the retrocession of it in the act of inspiration.
- 13. But in whatever manner the placenta may be brought into the pelvis, it should be suffered to remain there till the action of the uterus comes on, or so long as there is reason to fear a return of the hemorrhage, and it

armanulised as cumber and we at the doe Appetito and the basement in must then be carefully withdrawn, or until it drop away.

### SECTION X.

On the Retention of the Placenta from the irregular Action of the Uterus.

- 1. When all the parts of the uterus act with equivalent force, and at the same time, the combined power will contribute to the expulsion of whatever is contained in its cavity.
- 2. But if the uterus should act irregularly, the contrary effect might be produced.
- 3. If the fundus uteri should not act when the other parts are in action, the longitudinal contraction of the uterus would be produced; but if the central parts should only act, the uterus would then be contracted in the form of an hour-glass.
- 4. As the placenta cannot be excluded when the uterus acts in this irregular manner, it must be extracted by introducing the hand into the uterus, provided the state of the hemorrhage

should

( 80 )

must then be carefully withdrawn, or charl in

AN ROLLSES

on the Meterician of the Placents from the dire-

I. When all the purts of the stores not with equivalent force, and at the same time, the contact of power will established to the less pointed of whatever is contained in its cavity.

A. Dat if the same should act irregularity.

the other pairs are in action, the hondered.

The other pairs are in action, the hongracing the other pairs are in action, the hongracing contracted to produced; the hard the produced; the hard action of the central bear should only not, the honer would then be sentracted in the deep of

de de la company campot de marigoral ad de la company de de la company de de la company de la compan

should require it; and when it cannot be extracted by using the means before mentioned.

- 5. The hand ought never to be introduced into the uterus except in cases of real necessity, and then with the utmost circumspection and care; and the hand when introduced should not be withdrawn until the placenta is detached and brought into the pelvis.
- 6. If the whole placenta be loosened, this is easily effected; but if a portion of it should be found adhering, this must be separated by bending it back from the uterus, or by passing gently the fingers between it and the uterus.
- 7. When the uterus is found contracted in the form of an hour-glass, and this is the most common cause of the retention of the placenta, the contracted part must be dilated in the manner recommended for the dilatation of the os uteri, and it must be amply dilated, or it will immediately contract again round the wrist.
- 8. We must then proceed as is before advised.

· ( We )

tracted by using the means, before mentioned.

2. The head ought envier to be introduced into the read ought envier to be introduced into the read ought envier to be introduced into the read out of the their with the utmost circumspection sity, and then with the utmost circumspection and then with the placetion introduced about out he seithermy until the placetie is

of side, heremost of mesoning along settle of an Simula is to noting a li rad plant of the visual by the superior of the state of the spinished by particular the superior of the state of the spinished by particular the superior of the state of the spinished by particular the superior of the state of the state of the state of the spinished spinished the spinished spinished the spinished spinished the spinished spi

If the form of an binarylain, and this is the most
deciment of an binarylain, and this is the most
common cause of the retriction of the planting
the commercial partiment he cilqued in the manare recommended for the dilutation of the exmer recommended for the dilutation of the exmer recommended for the dilutation of the exmer recommended for the dilutation of the ex-

somediately contract again round the wrist.

S. We must then proceed as in before ad-

### SECTION XI.

On the Retention of the Placenta from the scirrhous Adhesion of it to the Uterus.

- 1. Should there be a degree of hemorrhage sufficient to make it necessary to introduce the hand to extract the placenta, a part of it must be separated, though there may be a scirrhous adhesion of the remainder to the uterus.
- 2. Then the method advised in the last section must be put in practice, and the firmer we find the adhesion, the slower the separation ought to be made.
- 3. But if there should be no hemorrhage of importance, and merely a retention of the placenta beyond its due time, we may say, for example, more than four hours, and the means before recommended are insufficient to bring down the placenta;
- 4. It may then be necessary to introduce the hand carefully to separate and extract the placenta,

ATTENNA NI.

On the Method of the Placents of the soir of the star of the soir of the soir University of the soil o

1. Should those is degree at hemorrhage collicient as make it necessary to introduce the hand to extract the planning, a part of it must be reparated, though their range for it reinflows asky for it from their remainder to the many for its remainder to the

2. Then the method advised in the last section must be part in practice, and the firmer we had the shape at the slower the acquisition could se the make

If there is no be no benerous of the participant of the proportion of the place of the place of the place of the place of the recommended are insufficient to bring flows the recommended are insufficient to bring slows the place of the plac

4. It may then be necessary to introduce the hand carried the the hand carried the

placenta, and the difficulty will not be increased by the delay.

- 5. Following the navel string as our guide, we must then pass the hand to the placenta; and if it should be found almost wholly adhering, we must begin with great caution to separate at the edge, and gradually proceed as before directed until the separation is completed.
- 6. Then grasping the placenta, we must slowly withdraw our hand, that the uterus may contract accordingly, and the chance of a subsequent hemorrhage be prevented.
- 7. The irritation made by the introduction of the hand, will often occasion a return of the action of the uterus, before dormant, that will greatly facilitate the separation.
- 8. Yet it is possible that a portion of the placenta may adhere so firmly as to make it unsafe to separate it with our fingers.
- 9. Should this circumstance occur notwithstanding the most deliberate and firm proceeding, it may sometimes be more justifiable to leave the adhering part remaining, than to use violence in separating it.

placement and the difficulty will not be interested

S. Following the name of the or the placents; our must then passedimental to the placents; and if tradequid of found almost wholly adhered if tradequid of found almost schools achoever the must begin with great causion in soparates at the edge, and gradually proceed as before directed and the experience is completed.

6. From gradual, the experience we may interstantly withers and the separation is completed.

slowly withers we can beauty and the enters of any or the enters.

of the head, beginning the prevented.

7. The instantion made by the introduction of the head, will eiten secusion a return of the action of the party, before definant, that

all to accurate that a persion of the

S. Should this circulations occur notwith tradian the most deliberate and from yearcording, itsmuy sometimes he more justificable to leave the adhering part remaining than to use violence in separating it. 10. But though hemorrhages are stayed when the greater portion of placenta is brought away, it is always a desirable thing to bring away the placenta and membranes in a perfect state; and if these are slowly extracted, any coagula formed in the uterus will usually be enveloped in them.

## SECTION XII.

On those Hemorrhages which follow the Expulsion or Extraction of the Placenta.

- 1. The hemorrhage in these cases may be either a continuation of that which existed before the exclusion of the *placenta*, or it may only follow the exclusion of the *placenta*.
- 2. When it is of the former kind, we may presume that it was not within our power to prevent it; but the latter kind may often be attributed to the violence or hurry with which the placenta has been extracted.
  - 3. This is not so dangerous as either of K

\* PLANTER OF THE PROPERTY OF THE PARTY OF TH of the last two will be the same and to find the last to the varieties of hemorrhage of which we have last spoken, though with imprudent management, or under particular circumstances, it has sometimes proved fatal.

- 4. All the cautions given with respect to the general management of the *placenta*, relate to the prevention of this kind of hemorrhage.
- 5. When the strength of women is much reduced by any cause which existed previous to labour, or when they have gone through much fatigue in the course of it, there is usually great heat and a rapid circulation of the blood at the time of delivery.
- 6. While they are in this situation, if the placenta were to be brought away hastily, an extraordinary quantity of blood must of necessity be discharged.
- 7. The interval of time which passeth between the birth of the child and the expulsion of the *placenta*, should therefore be employed in cooling the patient, and recovering her from her fatigue.
  - 8. Even when the placenta is excluded out K 2 of

the anagodisessed to emed

an appropriate free manus materials of the community of t

donor of receive an approve oil rath.

the fight along the first the make good of the

of the cavity of the *uterus*, it should be suffered to remain till all tumult is quieted, and then, with the membranes, slowly extracted.

- 9. The quantity of blood discharged in consequence of the separation of the placenta will vary in different women, or in the same women at different labours, independently of the manner in which the placenta may come away.
- 10. The less the quantity of blood discharged, the better women in general recover, provided there be no morbid cause of its diminution.
- 11. Some women are always prone to a great discharge of blood after the separation of the *placenta*, whatever care may be taken in extracting it.
- 12. This may often be prevented by keeping the patient out of bed till the membranes are broken and the waters discharged to the very moment of the child being born, rather retarding than forwarding its expulsion.
- 13. In all cases of dangerous hemorrhage, after the extraction of the placenta, it is first

K 3

necessary

of the cavity of the medical deployed he suffered in cemein till ed turnellt is quieted, and then,

only the membranes, durity excinated,

9. The quantity of those discharged in
consequence of the separation of the placests
will vary in enterent weamen, or in the same
women at different behours, independently of
the manner in which the placeme may come

10. The less the quantity of blood discharged, the better women in general recover, provided their be no moride cause of in direlation.

11: Some women my always prone to a great-discharge of blood, effect the separation of the placeton, whatever care may be taken in

12. This may often be prevented by keeping the patient out of bed till the membranes are broken and the waters discharged to the very moment of the child being born, rather

13. In all cases of dangerous hemorrhage, after the extraction of the placeson, it is first K. S. heckenney

necessary that we should be assured, by an examination per vaginam, that the uterus is not inverted.

- 14. Should there be an alarming hemorrhage after the separation and exclusion of the placenta, notwithstanding all the care which can be taken according to the methods before mentioned,
- 15. The doctrine of hemorrhages before given, and the general treatment already recommended, will enable you to fix upon the line of conduct it will be expedient to pursue, and to restrain or suppress them as far as they are under the influence of art.
- 16. In cases of hemorrhage so very profuse as to occasion frightful faintings, continuing so long as to raise great solicitude for the immediate safety of the patient, it was generally said, that cordials ought not to be given.
- 17. But this requires explanation. When the patient has continued faint so long as to give time, according to our judgement, for the vessels of the *uterus* to contract, then cordials

and

· See tri lunes forwards

the said planting of the planting of the said beauty of the said beaut

entrand paintrale du ed stadt blustê de adrine amendant han reprintipa ade tont again Ende again the line pallementalization palatinis states and the pallementalization palatinis

The ductions of handrings and the regreen, and the general mentioned about a regreen, and the general mentioned in the duction discommonded, will english and him to provide
and so restrain or approvale man for safety

16. In ruses of heimorrange of very proinse as the societies frequent lapstings, evenalaming to long as to rute prent salication
for the immediate salety of the partent, it
was greatfally said, that cardials ought not to
be given.

17. But this requires explanation. When the patient has configured him to long as to give time, eccording to our jud canent, for the remain of the therm to contract, then confinished and

# ON LABOURS

# ATTENDED WITH CONVULSIONS.

- 1. The convulsions which occur in pregnancy very much resemble the epilepsy; but to the symptoms, which these have in common, may be added, the peculiar hisping noise which women almost universally make with their lips during the convulsions.
- 2. When convulsions happen to women with child, they are generally, but not constantly, accompanied or followed with symptoms of labour; but though the convulsions may be removed, the child is most frequently afterwards born dead.
- 3. These convulsions are indicated by a piercing pain in the head, by giddiness and other vertiginous complaints, by blindness, by vacillation of the mind or a slight delirium, by violent cramp or pain at the stomach, by a fulness or apparent strangulation of the neck and fauces, and other affections of the vascular and nervous system.

4. The

# BHIZORAL MO

ATTENDED TO STATE AND STREET, STREET,

de l'un come much reseaulte de demony; les en comen new voys much reseaulte de demony; les en comto-the resugnime, which elma have m comto-the resugnime, the peculiar hisping none tripe is wobern almost eneverable mike with

The danks of an amplied proise age of the A to a standard of the standard of t

And pall shows

2. These convincions are judicated by a picture, and the control of the control o

ANTEN A

and nourishment in small quantities, very often repeated, are really needful.

- 18. Other means are also to be used for the purpose of recovering women from this long continued fainting; and one of the most effectual is, sprinkling the face freely with cold water.
- 19. After a profuse hemorrhage, the patient will frequently have a disposition to sleep, which has generally been considered as dangerous.
- 20. But short sleeps are very refreshing; though long ones, in a very weak state, are, under every circumstance, found to be injurious.
- 21. When there has been a dangerous hemorrhage, the patient should remain for many hours undisturbed, and in an horizontal position; and our attention must be continued as long as any danger is to be apprehended.

and monishment in small quantities, very often

18. Other means are also to be used for the paypose of recovering restace from this long continued minutes; and one of the most effectual is, sprobiling the face freely with cold tester.

19. After a profuse hemorehage, the patient nith frequently have a disposition to sleep, which has generally been equalished as dan-

20. But chort sleeps are very refreshing; though love ones, in a very weak state, are, under every circumstance, found to be in-

21. When there has been a staggerous he morrhage, the patientshould remain for many hears undisturbed, and in an horizontal position; and our attention must be continued as long as any danger is to be apprehended.

- 4. The means to be used for the prevention or cure of convulsions when threatened or existing, must be regulated according to the constitution of the patient and the violence of the symptoms.
- 5. It will always be necessary to take away some blood, and commonly to repeat the bleeding; and it has been found particularly serviceable to open the jugular vein; or to take away blood by cupping; and by applying leeches to the temples. Emetics, when they could be given, have been useful, as has sometimes also the warm bath. Clysters may be frequently exhibited. Opiates, joined with nervous medicines, may be given; and the patient is, by all the means in our power, to be soothed and restrained from violent exertions.
- 6. During the convulsions, the means by which contrary irritations may be excited are to be used; and of these the most powerful is, the dashing of cold water in the face, which has been known to prevent, or even to cure, convulsions.
  - · 7. Some writers have recommended the speedy

desired on comments to be used for the preventer conting, ment he regulated according to the consting, ment he regulated according to

trappetity exhibited. Opinion, idented with stated continue instantions may be explicit and and been Literal to provent, or even to chee,

To come writers have recompressed the

speedy delivery of the patient, as the most eligible, and only effectual method of removing puerperal convulsions; but others have insisted that the labour should be uninterrupted.

- 8. From the histories of all the cases of puerperal convulsions which have been hitherto recorded, it appears, that a greater number have died of those who were delivered by art, than when the labours were resigned to nature.
- 9. As far as my experience enables me to judge, we ought not to attempt to deliver women with convulsions before some progress is made in the labour.
- 10. But when the os uteri becomes dilated sufficiently, or to a certain degree, the patient safely may, and ought to be delivered by art, if from the urgency of the convulsions, and the general danger of the case, delivery should appear necessary.
- 11. The manner of delivering women in these cases, whether the operation be performed with the *forceps* or *vectis*, or by turning and extracting the child by the feet, has already been fully explained.

restrict of the partition of the partition; he the plant partition of the plant partition of the partition o

A from the histories of all the ones of an from the ones of passeporal convenience which has elected birtheres reconsist, it expenses, that a greater, muchber force died of chose who were delivered by any, have died of chose who were delivered by any, has when the labour were resigned to nature. It she filt as my expensions combine me to this were the taken of the compit not to extend to the very weare and a state of the compitation of the state of the sta

10. Hat when the as also becomes dilated a continue, or to a certain degree, the patient safety and ought to be delivered by act, if from the sugency of the convolutions, and the general danger of the case, delivery should

in The manner of delivering wantend in their cases of the cheration was part that cheration to the manner with the foreign or resting or by manner and extracting the child be the feet, has thready been failly explained.

and on

- 12. The event of the operation, both to the mother and child, will also very much depend upon the skill and circumspection with which it may be performed.
- 13. When dangerous convulsions come on in the early part of pregnancy, it is often clear that they arise from excessive uterine irritation.
- 14. It will then be justifiable and proper to forward the exclusion of the fætus, by puncturing the membranes as soon as it can be done with safety.

THE END.

( 001

without the amount the character to the section of the section of

no omos sacializados acestas como en m im entila perios presentente, il inclica cluca m im entila perios ligar excessiva attenino invita-

In the state of the property of the state of

aut in

# Published by Dr. DENMAN.

- 1. An Introduction to the Practice of Midwifery. 2 vols. 8vo. 4th edition. Price 17s. in boards.
- 2. The same, in 4to. 5th edition. Price 2l. 2s. in boards.

Printed for J. Johnson, No. 72, St. Paul's Church Yard.

Printed by S. Hamilton, Weybridge, Surrey.

THE PRINCIPLE OF THE PARTY OF Printed for J. Jonaton C. Marie Santa